

ADVANCES IN
IMMUNOLOGY

VOLUME 67

Cumulative Subject Index
Volumes 37-65



ACADEMIC PRESS

ADVANCES IN
Immunology

CUMULATIVE SUBJECT INDEX
VOLUMES 37-65

This Page Intentionally Left Blank

ADVANCES IN
Immunology
CUMULATIVE SUBJECT INDEX
VOLUMES 37-65

EDITED BY

FRANK J. DIXON

*The Scripps Research Institute
La Jolla, California*

ASSOCIATE EDITORS

Frederick Alt
K. Frank Austen
Tadamitsu Kishimoto
Fritz Melchers
Jonathan W. Uhr

VOLUME 67



ACADEMIC PRESS

San Diego London Boston
New York Sydney Tokyo Toronto

This book is printed on acid-free paper. ☺

Copyright © 1998 by ACADEMIC PRESS

All Rights Reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without permission in writing from the Publisher.

The appearance of the code at the bottom of the first page of a chapter in this book indicates the Publisher's consent that copies of the chapter may be made for personal or internal use of specific clients. This consent is given on the condition, however, that the copier pay the stated per copy fee through the Copyright Clearance Center, Inc. (222 Rosewood Drive, Danvers, Massachusetts 01923), for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale. Copy fees for pre-1998 chapters are as shown on the title pages. If no fee code appears on the title page, the copy fee is the same as for current chapters.
0065-2776/98 \$25.00

Academic Press

a division of Harcourt Brace & Company

525 B Street, Suite 1900, San Diego, California 92101-4495, USA

<http://www.apnet.com>

Academic Press Limited

24-28 Oval Road, London NW1 7DX, UK

<http://www.hbuk.co.uk/ap/>

International Standard Book Number: 0-12-022467-4

PRINTED IN THE UNITED STATES OF AMERICA

97 98 99 00 01 02 QW 9 8 7 6 5 4 3 2 1

CONTENTS

CONTENTS OF VOLUMES 37–65	vii
CUMULATIVE SUBJECT INDEX	1
CONTRIBUTOR INDEX	321

This Page Intentionally Left Blank

CONTENTS OF VOLUMES 37-65

Volume 37

Structure, Function, and Genetics of Human Class III Molecules	1
ROBERT C. GILES AND J. DONALD CAPRA	
The Complexity of Virus—Cell Interactions in Abelson Virus Infection of Lymphoid and Other Hematopoietic Cells	73
CHERYL A. WHITLOCK AND OWEN N. WITTE	
Epstein-Barr Virus Infection and Immunoregulation in Man	99
GIOVANNA TOSATO AND R. MICHAEL BLAESE	
The Classical Complement Pathway: Activation and Regulation of the First Complement Component	151
NEIL R. COOPER	
Membrane Complement Receptors Specific for Bound Fragments of C3	217
GORDON D. ROSS AND M. EDWARD MEDOF	
Murine Models of Systemic Lupus Erythematosus	269
ARGYRIOS N. THEOFILOPOULOS AND FRANK J. DIXON	

Volume 38

The Antigen-Specific, Major Histocompatibility Complex-Restricted Receptor on T Cells	1
PHILIPPA MARRACK AND JOHN KAPPLER	

Immune Response (<i>Ir</i>) Genes of the Murine Major Histocompatibility Complex	31
RONALD H. SCHWARTZ	
The Molecular Genetics of Components of Complement	203
R. D. CAMPBELL, M. C. CARROLL, AND R. R. PORTER	
Molecular Genetics of Human B Cell Neoplasia	245
CARLO M. CROCE AND PETER C. NOWELL	
Human Lymphocyte Hybridomas and Monoclonal Antibodies	275
DENNIS A. CARSON AND BRUCE D. FREIMARK	
Maternally Transmitted Antigen	313
JOHN R. RODGERS, ROGER SMITH III, MARILYN M. HUSTON, AND ROBERT R. RICH	
Phagocytosis of Particulate Activators of the Alternative Complement Pathway: Effects of Fibronectin	361
JOYCE K. CZOP	

Volume 39

Immunological Regulation of Hematopoietic/Lymphoid Stem Cell Differentiation by Interleukin 3	1
JAMES N. IHLE AND YACOB WEINSTEIN	
Antigen Presentation by B Cells and Its Significance in T-B Interactions	51
ROBERT W. CHESNUT AND HOWARD M. GREY	
Ligand-Receptor Dynamics and Signal Amplification in the Neutrophil	95
LARRY A. SKLAR	
Arachidonic Acid Metabolism by the 5-Lipoxygenase Pathway, and the Effects of Alternative Dietary Fatty Acids	145
TAK H. LEE AND K. FRANK AUSTEN	
The Eosinophilic Leukocyte: Structure and Function	177
GERALD J. GLEICH AND CHERYL B. ADOLPHSON	

CONTENTS OF VOLUMES 37-65

ix

Idiotypic Interactions in the Treatment of Human Diseases	255
RAIF S. GEHA	
Neuroimmunology	299
DONALD G. PAYAN, JOSEPH P. MCGILLIS, AND EDWARD J. GOETZL	

Volume 40

Regulation of Human B Lymphocyte Activation, Proliferation, and Differentiation	1
DIANE F. JELINEK AND PETER E. LIPSKY	
Biological Activities Residing in the Fc Region of Immunoglobulin	61
EDWARD L. MORGAN AND WILLIAM O. WEIGLE	
Immunoglobulin-Specific Suppressor T Cells	135
RICHARD G. LYNCH	
Immunoglobulin A (IgA): Molecular and Cellular Interactions Involved in IgA Biosynthesis and Immune Response	153
JIRI MESTECKY AND JERRY R. MCGHEE	

The Arrangement of Immunoglobulin and T-Cell Receptor Genes in Human Lymphoproliferative Disorders	247
THOMAS A. WALDMANN	

Human Tumor Antigens	323
RALPH A. REISFELD AND DAVID A. CHERESH	
Human Marrow Transplantation: An Immunological Perspective	379
PAUL J. MARTIN, JOHN A. HANSEN, RAINER STORB, AND E. DONNALL THOMAS	

Volume 41

Cell Surface Molecules and Early Events Involved in Human T Lymphocyte Activation	1
ARTHUR WEISS AND JOHN B. IMBODEN	

Function and Specificity of T Cell Subsets in the Mouse JONATHAN SPRENT AND SUSAN R. WEBB	39
Determinants on Major Histocompatibility Complex Class I Molecules Recognized by Cytotoxic T Lymphocytes JAMES FORMAN	135
Experimental Models for Understanding B Lymphocyte Formation PAUL W. KINCade	181
Cellular and Humoral Mechanisms of Cytotoxicity: Structural and Functional Analogies JOHN DING-E YOUNG AND ZANVIL A. COHN	269
Biology and Genetics of Hybrid Resistance MICHAEL BENNETT	333

Volume 42

The Clonotype Repertoire of B-Cell Subpopulations NORMAN R. KLINMAN AND PHYLLIS-JEAN LINTON	1
The Molecular Genetics of the Arsonate Idiotypic System of A/J Mice GARY RATHBUN, INAKI SANZ, KATHERYN MEEK, PHILIP TUCKER, AND J. DONALD CAPRA	95
The Interleukin 2 Receptor KENDALL A. SMITH	165
Characterization of Functional Surface Structures on Human Natural Killer Cells JEROME RITZ, REINHOLD E. SCHMIDT, JEAN MICHON, THIERRY HERCEND, AND STUART E. SCHLOSSMAN	181
The Common Mediator of Shock, Cachexia, and Tumor Necrosis B. BEUTLER AND A. CERAMI	213
Myasthenia Gravis JON LINDSTROM, DIANE SHELTON, AND YOSHITAKA FUJII	233

Alterations of the Immune System in Ulcerative Colitis and Crohn's Disease	285
RICHARD P. MACDERMOTT AND WILLIAM F. STENSON	

Volume 43

The Chemistry and Mechanism of Antibody Binding to Protein Antigens	1
ELIZABETH D. GETZOFF, JOHN A. TAINER, RICHARD A. LERNER, AND H. MARIO GEYSEN	
Structure of Antibody–Antigen Complexes: Implications for Immune Recognition	99
P. M. COLMAN	
The $\gamma\delta$ T-Cell Receptor	133
MICHAEL B. BRENNER, JACK L. STROMINGER, AND MICHAEL S. KRANGEL	
Specificity of the T-Cell Receptor for Antigen	193
STEPHEN M. HEDRICK	
Transcriptional Controlling Elements in the Immunoglobulin and T-Cell Receptor Loci	235
KATHRYN CALAME AND SUZANNE EATON	
Molecular Aspects of Receptors and Binding Factors for IgE	277
HENRY METZGER	

Volume 44

Diversity of the Immunoglobulin Gene Superfamily	1
TIM HUNKAPILLER AND LEROY HOOD	
Genetically Engineered Antibody Molecules	65
SHERIE L. MORRISON AND VERNON T. OT	
Antinuclear Antibodies: Diagnostic Markers for Autoimmune Diseases and Probes for Cell Biology	93
ENG M. TAN	

Interleukin-1 and Its Biologically Related Cytokines	153
CHARLES A. DINARELLO	
Molecular and Cellular Events of T-Cell Development	207
B. J. FOWLKES AND DREW M. PARDOLL	
Molecular Biology and Function of CD4 and CD8	265
JANE R. PARNES	
Lymphocyte Homing	313
TED A. YEDNOCK AND STEVEN D. ROSEN	

Volume 45

Cellular Interactions in the Humoral Immune Response	1
ELLEN S. VITETTA, RAFAEL FERNANDEZ-BOTRAN, CHRISTOPHER D. MYERS, AND VIRGINIA M. SANDERS	
MHC-Antigen Interaction: What Does the T-Cell Receptor See?	107
PHILIPPE KOURILSKY AND JEAN-MICHEL CLAVERIE	
Synthetic T- and B-Cell Recognition Sites: Implications for Vaccine Development	195
DAVID R. MILICH	
Rationale for the Development of an Engineered Sporozoite Malaria Vaccine	283
VICTOR NUSSENZWEIG AND RUTH S. NUSSENZWEIG	
Virus-Induced Immunosuppression: Infections with Measles Virus and Human Immunodeficiency Virus	335
MICHAEL B. MCCHESNEY AND MICHAEL B. A. OLDSSTONE	
The Regulators of Complement Activation (RCA) Gene Cluster	381
DENNIS HOURCADE, V. MICHAEL HOLERS, AND JOHN P. ATKINSON	
Origin and Significance of Autoreactive T Cells	417
MAURICE ZAUDERER	

Volume 46

Physical Maps of the Mouse and Human Immunoglobulin-like Loci	1
ERIC LAI, RICHARD K. WILSON, AND LEROY E. HOOD	
Molecular Genetics of Murine Lupus Models	61
ARCYRIOS N. THEOFILOPOULOS, REINHARD KOFLER, PAUL A. SINGER, AND FRANK J. DIXON	
Heterogeneity of Cytokine Secretion Patterns and Functions of Helper T Cells	111
TIM R. MOSMANN AND ROBERT L. COFFMAN	
The Leukocyte Integrins	149
TAKASHI K. KISHIMOTO, RICHARD S. LARSON, ANGEL L. CORBI, MICHAEL L. DUSTIN, DONALD E. STAUNTON, AND TIMOTHY A. SPRINGER	
Structure and Function of the Complement Receptors, CR1 (CD35) and CR2 (CD21)	183
JOSEPH M. AHEARN AND DOUGLAS T. FEARON	
The Cellular and Subcellular Bases of Immunosenescence	221
MARILYN L. THOMAN AND WILLIAM O. WEIGLE	
Immune Mechanisms in Autoimmune Thyroiditis	263
JEANNINE CHARREIRE	

Volume 47

Regulation of Immunoglobulin E Biosynthesis	1
KIMISHIGE ISHIZAKA	
Control of the Immune Response at the Level of Antigen-Presenting Cells: A Comparison of the Function of Dendritic Cells and B Lymphocytes	45
JOSHUA P. METLAY, ELLEN PURÉ, AND RALPH M. STEINMAN	
The CD5 B Cell	117
THOMAS J. KIPPS	

Biology of Natural Killer Cells	187
GIORGIO TRINCHIERI	
The Immunopathogenesis of HIV Infection	377
ZEDA F. ROSENBERG AND ANTHONY S. FAUCI	
The Obese Strain of Chickens: An Animal Model with Spontaneous Autoimmune Thyroiditis	433
GEORGE WICK, HANS PETER BREZINSCHEK, KAREL HÁLA, HERMANN DIETRICH, HUGO WOLF, AND GUIDO KROEMER	

Volume 48

Internal Movements in Immunoglobulin Molecules	1
ROALD NEZLIN	
Somatic Diversification of the Chicken Immunoglobulin Light-Chain Gene	41
WAYNE T. McCORMACK AND CRAIG B. THOMPSON	
T Lymphocyte-Derived Colony-Stimulating Factors	69
ANNE KELSO AND DONALD METCALF	

The Molecular Basis of Human Leukocyte Antigen Class II Disease Associations	107
DOMINIQUE CHARRON	

Neuroimmunology	161
E. J. GOETZL, D. C. ADELMAN, AND S. P. SREEDHARAN	
Immune Privilege and Immune Regulation in the Eye	191
JERRY Y. NIEDERKORN	
Molecular Events Mediating T-Cell Activation	227
AMNON ALTMAN, K. MARK COGGESHALL, AND TOMAS MUSTELIN	

Volume 49

Human Immunoglobulin Heavy-Chain Variable Region Genes: Organization, Polymorphism, and Expression	1
VIRGINIA PASCUAL AND J. DONALD CAPRA	

Surface Antigens of Human Leucocytes	75
V. HOŘEJŠÍ	
Expression, Structure, and Function of the CD23 Antigen	149
G. DELESPESSE, U. SUTER, D. MOSSALAYI, B. BETTLER, M. SARFATI, H. HOFSTETTER, E. KILCHERR, P. DEBRE, AND A. DALLOUL	
Immunology and Clinical Importance of Antiphospholipid Antibodies	193
H. PATRICK MCNEIL, COLIN N. CHESTERMAN, AND STEVEN A. KRILIS	
Adoptive T-Cell Therapy of Tumors: Mechanisms Operative in the Recognition and Elimination of Tumor Cells	281
PHILIP D. GREENBERG	
The Development of Rational Strategies for Selective Immunotherapy against Autoimmune Demyelinating Disease	357
LAWRENCE STEINMAN	
The Biology of Bone Marrow Transplantation for Severe Combined Immune Deficiency	381
ROBERTSON PARKMAN	

Volume 50

Selective Elements for the Vβ Region of the T Cell Receptor: Mls and the Bacterial Toxic Mitogens	1
CHARLES A. JANEWAY, JR.	
Programmed Cell Death in the Immune System	55
J. JOHN COHEN	
Avian T-Cell Ontogeny	87
MAX D. COOPER, CHEN-LO H. CHEN, R. PAT BUCY, AND CRAIG B. THOMPSON	
Structural and Functional Chimerism Results from Chromosomal Translocation in Lymphoid Tumors	119
T. H. RABBITS AND T. BOEHM	
Interleukin-2, Autotolerance, and Autoimmunity	147
GUIDO KROEMER, JOSÉ LUIS ANDREU, JOSÉ ANGEL GONZALO, JOSÉ C. GUTIERREZ-RAMOS, AND CARLOS MARTÍNEZ-A.	

Histamine Releasing Factors and Cytokine-Dependent Activation of Basophils and Mast Cells	237
ALLEN P. KAPLAN, SESHA REDDIGARI, MARIA BAEZA, AND PIOTR KUNA	
Immunologic Interactions of T Lymphocytes with Vascular Endothelium	261
JORDAN S. POBER AND RAMZI S. COTRAN	
Adoptive Transfer of Human Lymphoid Cells to Severely Immunodeficient Mice: Models for Normal Human Immune Function, Autoimmunity, Lymphomagenesis, and AIDS	303
DONALD E. MOSIER	

Volume 51

Human Antibody Effector Function	1
DENNIS R. BURTON AND JENNY M. WOOF	
The Development of Functionally Responsive T Cells	85
ELLEN V. ROTHENBERG	
Role of Perforin in Lymphocyte-Mediated Cytolysis	215
HIDEO YAGITA, MOTOMI NAKATA, AKEMI KAWASAKI, YOICHI SHINKAI, AND KO OKUMURA	
The Central Role of Follicular Dendritic Cells in Lymphoid Tissues	243
FOLKE SCHRIEVER AND LEE MARSHALL NADLER	
The Murine Autoimmune Diabetes Model: NOD and Related Strains	285
HITOSHI KIKUTANI AND SUSUMU MAKINO	
The Pathobiology of Bronchial Asthma	323
JONATHAN P. ARM AND TAK H. LEE	

Volume 52

Cell Biology of Antigen Processing and Presentation to Major Histocompatibility Complex Class I Molecule-Restricted T Lymphocytes	1
JONATHAN W. YEWDELL AND JACK R. BENNINK	
Human B Lymphocytes: Phenotype, Proliferation, and Differentiation	125
JACQUES BANCHEREAU AND FRANÇOISE ROUSSET	

CONTENTS OF VOLUMES 37-65

xvii

Cytokine Gene Regulation: Regulatory <i>cis</i> -Elements and DNA Binding Factors Involved in the Interferon System	263
NOBUYUKI TANAKA AND TADATSUGU TANIGUCHI	
Cellular and Molecular Mechanisms of B Lymphocyte Tolerance	283
G. J. V. NOSSAL	
Cell Surface Structures on Human Basophils and Mast Cells: Biochemical and Functional Characterization	333
PETER VALENT AND PETER BETTELHEIM	
Animal Models for Acquired Immunodeficiency Syndrome	425
THOMAS J. KINDT, VANESSA M. HIRSCH, PHILIP B. JOHNSON, AND SANSANA SAWASDIKOSOL	

Volume 53

Lymphokine and Cytokine Production by Fc ϵ RI $^+$ Cells	1
WILLIAM E. PAUL, ROBERT A. SEDER, AND MARSHALL PLAUT	
The Leukemia Inhibitory Factor and Its Receptor	31
DAVID P. GEARING	
The Role of CD4 and CD8 in T-Cell Activation and Differentiation	59
M. CARRIE MICELI AND JANE R. PARNES	
B Lymphopoiesis in the Mouse	123
ANTONIUS ROLINK AND FRITZ MELCHERS	
Compartmentalization of the Peripheral Immune System	157
GUIDO KROEMER, EDUARDO CUENDE, AND CARLOS MARTINEZ-A.	
Immunological Memory	217
CHARLES R. MACKAY	
Recognition of Bacterial Endotoxins by Receptor-Dependent Mechanisms	267
RICHARD J. ULEVITCH	

Cell Adhesion Molecules as Targets of Autoantibodies in Pemphigus and Pemphigoid, Bullous Diseases Due to Defective Epidermal Cell Adhesion	291
JOHN R. STANLEY	

Volume 54

Interleukin-6 in Biology and Medicine	1
SHIZUO AKIRA, TETSUYA TAGA, AND TADAMITSU KICHIMOTO	
Interleukin-9	79
J.-C. RENAUD, F. HOUSSIAU, J. LOUAHED, A. VINK, J. VAN SNICK, AND C. UYTTEHOVE	
Superantigens and Their Potential Role in Human Disease	99
BRIAN KOTZIN, DONALD Y. M. LEUNG, JOHN KAPPLER, AND PHILIPPA MARRACK	
Interleukin-1 Receptor Antagonist	167
WILLIAM P. AREND	
The Mechanism and Regulation of Immunoglobulin Isotype Switching	229
ROBERT L. COFFMAN, DEBORAH A. LEBMAN, AND PAUL ROTHRMAN	
CD44 and Its Interaction with Extracellular Matrix	271
JAYNE LESLEY, ROBERT HYMAN, AND PAUL W. KINCADE	
Immunoglobulin Receptor-Associated Molecules	337
NOBUO SAKAGUCHI, TATSUYA MATSUO, JUN NOMURA, KAZUHIKO KUWAHARA, HIDEYA IGARASHI, AND SEIJI INUI	

Analysis of B Cell Tolerance <i>in Vitro</i>	393
DAVID W. SCOTT	

Volume 55

The kit Ligand, Stem Cell Factor	1
STEPHEN J. GALLI, KRISZTINA M. ZSEBO, AND EDWIN N. GEISSLER	
Interleukin-8 and Related Chemotactic Cytokines—CXC and CC Chemokines	97
MARCO BAGGIOLINI, BEATRICE DEWALD, AND BERNHARD MOSER	

Receptors for Transforming Growth Factor- β	181
KOHEI MIYAZONO, PETER TEN DIJKE, HIDENORI ICHIJO, AND CARL-HENRIK HELDIN	
Biochemistry of B Lymphocyte Activation	221
MICHAEL R. GOLD AND ANTHONY L. DEFRAZCO	
CD5 B Cells, a Fetal B-Cell Lineage	297
RICHARD R. HARDY AND KYOKO HAYAKAWA	
Human Natural Killer Cells: Origin, Clonality, Specificity, and Receptors	341
LORENZO MORETTA, ERMANNO CICCONE, MARIA CRISTINA MINGARI, ROBERTO BIASSONI, AND ALESSANDRO MORETTA	
MHC Class I-Deficient Mice	381
DAVID H. RAULET	
The Immune System of Mice Lacking Conventional MHC Class II Molecules	424
SUSANNA CARDELL, MATTHIAS MERKENSCHLAGER, HELEN BODMER, SUSAN CHAN, DOMINIC COSGROVE, CHRISTOPHE BENOIST, AND DIANE MATHIS	

Volume 56

Properties and Functions of Interleukin-10	1
TIM R. MOSMANN	
The Mechanism of V(D)J Joining: Lessons from Molecular, Immunological, and Comparative Analyses	27
SUSANNA M. LEWIS	
Involvement of the Protein Tyrosine Kinase p56 ^{lck} in T-Cell Signaling and Thymocyte Development	151
STEVEN J. ANDERSON, STEVEN D. LEVIN, AND ROGER M. PERLMUTTER	
Generating the Antibody Repertoire in Rabbit	179
KATHERINE L. KNIGHT AND MARY A. CRANE	

Immunotherapeutic Strategies Directed at the Trimolecular Complex AMITABH GAUR AND C. GARRISON FATHMAN	219
Therapeutic Regulation of the Complement System in Acute Injury States FRANCIS D. MOORE, JR.	267
Chemoimmunoconjugates for the Treatment of Cancer GEOFFREY A. PIETERSZ, APRIL ROWLAND, MARK J. SMYTH, AND IAN F. C. MCKENZIE	301
The Molecular Basis of Susceptibility to Rheumatoid Arthritis ROBERT WINCHESTER	389
Retrovirus-Induced B Cell Neoplasia in the Bursa of Fabricius PAUL E. NEIMAN	467

Volume 57

Molecular Basis of Fc Receptor Function MARK D. HULETT AND P. MARK HOGARTH	1
Fas and Fas Ligand: A Death Factor and Its Receptor SHIGEKAZU NAGATA	129
Interleukin-5 and Its Receptor System: Implications in the Immune System and Inflammation KIYOSHI TAKATSU, SATOSHI TAKAKI, AND YASUMICHI HITOSHI	145
Human Antibodies from Combinatorial Libraries DENNIS R. BURTON AND CARLOS F. BARBAS III	191
Immune Response against Tumors CLAUDE ROTH, CHRISTOPH ROCHLITZ, AND PHILIPPE KOURILSKY	281
Formation of the Chicken B-Cell Repertoire: Ontogenesis, Regulation of Ig Gene Rearrangement, and Diversification by Gene Conversion CLAUDE-AGNÈS REYNAUD, BARBARA BERTOCCI, AURIEL DAHAN, AND JEAN-CLAUDE WEILL	353

Volume 58

NF- κ B and Rel Proteins in Innate Immunity	1
ELIZABETH B. KOPP AND SANKAR GHOSH	
V(D)J Recombination and Double-Strand Break Repair	29
DAVID T. WEAVER	
Development and Selection of T Cells: Facts and Puzzles	87
PAWEŁ KISIELOW AND HARALD VON BOEHMER	
The Pharmacology of T-Cell Apoptosis	211
GUIDO KROEMER	
Intraepithelial Lymphocytes and the Immune System	297
GEK-KEE SIM	
Leukocyte Migration and Adhesion	345
BEAT A. IMHOF AND DOMINIQUE DUNON	
Gene Transfer as Cancer Therapy	417
GLENN DRANOFF AND RICHARD C. MULLIGAN	

Volume 59

The CD1 Family: A Third Lineage of Antigen-Presenting Molecules	1
STEVEN A. PORCELLI	
Positive Selection of Thymocytes	99
PAMELA J. FINK AND MICHAEL J. BEVAN	
Molecular and Cellular Aspects of X-Linked Agammaglobulinemia	135
PASCHALIS SIDERAS AND C. I. EDWARD SMITH	
The Common δ -Chain for Multiple Cytokine Receptors	225
KAZUO SUGAMURA, HIRONOBU ASAO, MOTONARI KONDO, NOBUYUKI TANAKA, NAOTO ISHII, MASATAKA NAKAMURA, AND TOSHIKAZU NAKAMURA	

Self-Tolerance Checkpoints in B Lymphocyte Development	279
CHRISTOPHER C. GOODNOW, JASON G. CYSTER, SUZANNE B. HARTLEY, SARAH E. BELL, MICHAEL P. COOKE, JAMES L. HEALY, SRINIVAS AKKARAJU, JEFFREY C. RATHMELL, SARAH L. POGUE, AND KEVAN P. SHOKAT	
The Regulation of Pulmonary Immunity	369
MARY F. LIPSCOMB, DAVID E. RICE, C. RICHARD LYONS, MARK R. SCHUYLER, AND DAVID WILKES	

Volume 60

The Janus Protein Tyrosine Kinase Family and Its Role in Cytokine Signaling	1
JAMES N. IHLE	
X-Linked Agammaglobulinemia and Immunoglobulin Deficiency with Normal or Elevated IgM: Immunodeficiencies of B-Cell Development and Differentiation	37
RAMSAY FULEIHAN, NARAYANASWAMY RAMESH, AND RAIF S. GEHA	
Defective Glycosyl Phosphatidylinositol Anchor Synthesis and Paroxysmal Nocturnal Hemoglobinuria	57
TAROH KINOSHITA, NORIMITSU INOUE, AND JUNJI TAKEDA	
The Use of Multiple Antigen Peptides in the Analysis and Induction of Protective Immune Responses against Infectious Diseases	105
ELIZABETH H. NARDIN, GIANE A. OLIVEIRA, J. MAURICIO CALVO-CALLE, AND RUTH S. NUSSENZWEIG	
Eosinophils: Biology and Role in Disease	151
ANDREW J. WARDLAW, REDWAN MOQBEL, AND A. BARRY KAY	
In Situ Studies of the Germinal Center Reaction	267
GARNETT KELSOE	
Cytotoxic T Lymphocytes: The Newly Identified Fas (CD95)-Mediated Killing Mechanism and a Novel Aspect of Their Biological Functions	289
HAJIME TAKAYAMA, HIDEFUMI KOJIMA, AND NOBUKATA SHINOHARA	
The Role of Nitric Oxide in Inflammation	323
C. RICK LYONS	

Volume 61

CD40-CD40 Ligand: A Multifunctional Receptor-Ligand Pair	1
CEES VAN KOOTEN AND JACQUES BANCHEREAU	
 Antibody Class Switching	 79
JANET STAVNEZER	
 Interleukin-2 Receptor Signaling Mechanisms	 147
LARRY M. KARNITZ AND ROBERT T. ABRAHAM	
 Control of the Complement System	 201
M. KATHRYN LISZEWSKI, TIMOTHY C. FARRIES, DOUGLAS M. LUBLIN, ISABELLE A. ROONEY, AND JOHN P. ATKINSON	
 V(D)J Recombination Pathology	 285
KLAUS SCHWARZ AND CLAUS R. BARTRAM	
 Major Histocompatibility Complex Class II Deficiency: A Disease of Gene Regulation	 327
VIKTOR STEIMLE, WALTER REITH, AND BERNARD MACH	
 TH1-TH2 Cells in Allergic Responses: At the Limits of a Concept	 341
IWAN AEBISCHER AND BEADA M. STADLER	

Volume 62

Organization of the Human Immunoglobulin Heavy-Chain Locus	1
FUMIHIKO MATSUDA AND TASUKU HONJO	
 Analysis of Gene Function in Lymphocytes by RAG-2-Deficient Blastocyst Complementation	 31
JIANZHU CHEN	
 Interferon-γ: Biology and Role in Pathogenesis	 61
ALFONS BILLIAU	
 Role of the CD28-B7 Costimulatory Pathways in T-Cell-Dependent B-Cell Responses	 131
KAREN S. HATHCOCK AND RICHARD J. HODES	

Prostaglandin Endoperoxide H Synthases-1 and -2 WILLIAM L. SMITH AND DAVID L. DEWITT	167
Human Tumor Antigens Are Ready to Fly ROBERT A. HENDERSON AND OLIVERA J. FINN	217
Inflammatory Mediators, Cytokines, and Adhesion Molecules in Pulmonary Inflammation and Injury NICHOLAS W. LUKACS AND PETER A. WARD	257

Volume 63

Surrogate Light Chain in B-Cell Development HAJIME KARASUYAMA, ANTONIUS ROLINK, AND FRITZ MELCHERS	1
CD40 and Its Ligand LISA B. CLARK, TERESA M. FOY, AND RANDOLPH J. NOELLE	43
Human Immunodeficiency Virus Infection of Human Cells Transplanted to Severe Combined Immunodeficient Mice DONALD E. MOSIER	79
Lessons from Immunological, Biochemical, and Molecular Pathways of the Activation Mediated by IL-2 and IL-4 ANGELITA REBOLLO, JAVIER GÓMEZ, AND CARLOS MARTÍNEZ-A.	127
B Lymphocyte Development and Transcription Regulation <i>in Vivo</i> DAVINA OPSTELTEN	197
Soluble Cytokine Receptors: Their Roles in Immunoregulation, Disease, and Therapy RAFAEL FERNANDEZ-BOTRAN, PAULA M. CHILTON, AND YUHE MA	269
Cytokine Expression and Cell Activation in Inflammatory Arthritis LIONEL B. IVASHKIV	337
Prolactin, Growth Hormone, and Insulin-like Growth Factor I in the Immune System RON KOOIJMAN, ELISABETH L. HOOGHE-PETERS, AND ROBERT HOOGHE	377

Volume 64

Proteasomes and Antigen Processing	1
KEIJI TANAKA, NOBUYUKI TANAHASHI, CHIZUKO TSURUMI, KIN-YA YOKOTA, AND NAOKI SHIMBARA	
Recent Advances in Understanding V(D)J Recombination	39
MARTIN GELLERT	
The Role of Ets Transcription Factors in the Development and Function of the Mammalian Immune System	65
ALEXANDER G. BASSUK AND JEFFREY M. LEIDEN	
Mechanism of Class I Assembly with β^2Microglobin and Loading with Peptide	105
TED H. HANSEN AND DAVID R. LEE	
How Do Lymphocytes Know Where to Go: Current Concepts and Enigmas of Lymphocyte Homing	139
MARKO SALMI AND SIRPA JALKANEN	
Plasma Cell Dyscrasias	219
NORIHIRO NISHIMOTO, SACHIKO SUEMATSU, AND TADAMITSU KISHIMOTO	
Anti-Tumor Necrosis Factor-α	283
MARC FELDMANN, MICHAEL J. ELLIOTT, JAMES N. WOODY, AND RAVINDER N. MAINI	

Volume 65

NF-IL6 and NF-κB in Cytokine Gene Regulation	1
SHIZUO AKIRA AND TADAMITSU KISHIMOTO	
Transporter Associated with Antigen Processing	47
TIM ELLIOTT	
NF-κB as a Frequent Target for Immunosuppressive and Anti- Inflammatory Molecules	111
PATRICK A. BAEUERLE AND VIJAY R. BAICHWAL	

Mouse Mammary Tumor Virus: Immunological Interplays between Virus and Host	139
SANJIV A. LUTHER AND HANS ACHA-ORBEA	
IgA Deficiency	245
PETER D. BURROWS AND MAX D. COOPER	
Role of Cellular Immunity in Protection against HIV Infection	277
SARAH ROWLAND-JONES, RUSUNG TAN, AND ANDREW McMICHAEL	
High Endothelial Venules: Lymphocyte Traffic Control and Controlled Traffic	347
GEORG KRAAL AND REINA E. MEBIUS	

SUBJECT INDEX

Boldface numerals indicate volume number

A

- Aarskog-Scott syndrome, **63**:149
Ab2s, **42**:149-157
ABC transporters, **65**:55, 56-57, 59
Abelson disease
 anti-A-MuLv antibodies in study of *c-abl*-encoded protein, **37**:77-78
 c-abl expression pattern, normal and leukemic cells, **37**:78-79
 complexity
 genetics of susceptibility, **37**:88-89
 helper virus role in neoplastic transformation, **37**:87-88
 proliferating cells as transformation targets *in vitro*, **37**:88
 transformation of cells from genetically resistant mice, **37**:89-90
 use of site-directed antibodies, **37**:90, 91
 variations in disease process induced by mutants, **37**:90
complexity of transformation *in vitro*
 associated cellular changes, **37**:93-94
 cultured B cell lines in study of, **37**:94-95
 early effects of *v-abl* expression, **37**:91
 preneoplastic growth properties of infected pre-B cells, **37**:91-92
 tumorigenicity progression, **37**:92-93
history of, **37**:75-76
neoplastic transformation
 NIH 3T3 fibroblasts, **37**:83-84
 pre-B cells, **37**:79-80
 rapid induction of thymomas, **37**:83
target cells in monocyte-macrophage lineage, **37**:82-83
techniques for, mature B and plasma cells, **37**:80-81
transformed cell lines with differentiation potential, **37**:82
non-neoplastic changes induced, **37**:84-85
 agar colony formation from fetal liver erythroblasts, **37**:85
 lethality, **37**:86-87
 resistance of GM-CFC to leukemia-associated inhibition activity, **37**:85
overview, **37**:74-75
properties of *v-abl* and tyrosine kinase transforming protein, **37**:76
Abelson-murine leukemia virus
 fetal liver cell infection, **39**:40-41
 IL-3-regulated growth and, **39**:40-41
 helper T cell infection, **39**:41
 IL-2-regulated growth and, **39**:41
 plasmacytogenesis, **64**:230
 transfected cells, rearrangement by V(D)J joining of antigen receptor genes, **56**:48-49
 transformed mast cell lines, lymphokine production by, **53**:4-6
Abelson virus
 B cell formation and, **41**:195, 209, 217, 219, 225
 hybrid resistance and, **41**:397
abl, tumor antigens derived from, **57**:296-297
ABL-MYC retrovirus, plasmacytogenesis, **64**:231-238

- Accessibility model, **61**:84, 86
 germline transcription, **54**:241
- Accessory cells
 B cell formation and, **41**:220
 hybrid resistance and, **41**:350, 373
 immune reconstitution and, **40**:416–417
 isotype-specific responses and, **40**:219
 T cell activation and, **41**:7–14, 21
 T cell subsets and, **41**:63–66, 69
- Accessory genes, HIV-1 disease, **63**:85–86, 94, 105
- Accessory molecules
 CD4 or CD8, role in immune response, **54**:124–125
 CD40-dependent B cell activation, **63**:56–57
 cytotoxic T cells and, **41**:150–152, 158, 168–170
 T cell receptor and, **45**:108–110
- Acetylcholine receptor, *see also* Experimental autoimmune myasthenia gravis; Myasthenia gravis
 antigenic structure of, **42**:240–248
 discovery of, **42**:233–236
 Ig heavy-chain variable region genes and, **49**:31, 46, 50–51
 main immunogenic region, Abs to, **42**:240–241, 246, 262
 muscle, **42**:239
 autoantibody effects on, **42**:260–263
 neuromuscular transmission and, **42**:236–237
 neuronal, **42**:239
 polypeptide chain, **42**:238
 skeletal muscle, **42**:238
 structure of, **42**:237–240
 suppressor T cells and, **42**:269
 T cell epitopes and, **42**:241–245
- Acetylcholinesterase
 autoimmune thyroiditis and, **46**:268
 human erythrocyte, ethanolaminephosphates in, **60**:62
- N-Acetylcysteine
 effect on apoptosis, **58**:250, 262–263
 inhibition of HIV lymphocyte death, **58**:279
- N-Acetylglucosamine
 and glucosamine, in GPI-anchored proteins, **60**:60–72
- transfer to PI from UDP-GlcNAc, **60**:64–70
- Acetyltransferase, antinuclear antibodies and, **44**:109, 110
- AChR, *see* Acetylcholine receptor
- α 1-Acid glycoprotein, **65**:22
- Acid hydrolysis, hybrid resistance and, **41**:398
- Acid phosphatase
 B cell formation and, **41**:212, 213
 cytolysis and, **41**:281
- cis-Aconitic anhydride, in chemoimmunoconjugation, **56**:315–316
- Acquired agammaglobulinemia, **59**:138–139, 146
- Acquired C1 esterase inhibitor deficiency, B cell lymphoma reaction to anti-Id antibodies, **39**:288–291
 idiotypic network and, **39**:289, 291
- Acquired immune deficiency syndrome, *see AIDS; AIDS, animal models*
- Acromegaly, **63**:409
- ACTH, *see Adrenocorticotrophic hormone*
- Actin
 cytolysis and, **41**:275
 hybrid resistance and, **41**:378
 immunosenescence and, **46**:224
 leukocyte integrins and, **46**:151, 169
 polymerization, **63**:149
- Activated partial thromboplastin time, aPL antibodies and, **49**:200–202, 244
- Activated protein C, aPL antibodies and, **49**:253–254, 257
- Activating peptide receptors, **52**:366–375
 for formyl-methionine peptides and related compounds, **52**:371–372
 for IL-8 and related integrins, **52**:367–371
 for substance P and other neuropeptides, **52**:372–375
- Activation
 antigen-dependent, **52**:199
 antigen receptor-dependent, *see Antigen receptor-dependent activation*
 complement components and, **38**:214, 234
 contact, *see Contact activation*
 maternally transmitted antigen gene, **38**:342
 polyclonal, B cell, *see Polyclonal B cell activation*
 SCID and, **49**:386
 sequential, of Ig genes, **40**:256–257

- surface antigens of human leukocytes and, **49:125**
- T cell, *see* T cell activation
- T cell receptor $\gamma\delta$ lymphocytes, **43:171–172**
- Activation antigen, **48:248–250**
accessory, **48:241–242**
glycosyl-phosphatidylinositol, **48:237–241**
surface antigens of human leukocytes and, **49:115, 117**
- Activation-induced death, hybridomas, **50:67**
- Active immunotherapy, **62:219**
- Activin type II receptors
amino acid sequences, **55:190–192**
cDNA cloning, **55:188–192**
structure, **55:190–193**
- Acute asthmatic response, and airway tone, **51:350–351**
- Acute lymphoblastic leukemia, **46:45; 63:244**
B-lineage, **61:11–12**
- Acute lymphocytic leukemia
CD1 proteins, **59:60**
with 4:11 chromosomal translocation, **40:293**
T and B cell lineage-specific genes and, **40:284–290**
- Acute myelomonocytic leukemia, **46:160**
- Acute negative selection, **59:106–107**
- Acute phase proteins, **65:6**
NF- κ B-dependent, **58:9–10**
synthesis, **54:25–27, 216**
- Acute phase response factor, **65:19**
- Acute respiratory distress syndrome, **62:260–265, 267, 270, 271, 286**
IL-8, **66:140**
MIF, **66:214–215**
- Acyl salicylates, PGHS, **62:187**
- Adapter molecules, **63:134, 141–143**
- Adaptive immunity, NK cells and, **47:291–295**
- Addressin cell adhesion molecule-1, ligand for $\alpha 4\beta 7$ receptor, **60:170–171**
- Addressins, **62:257**
lymphocyte–HEV interaction, **65:361–363, 377**
mucosal, MadCAM-1, **58:360**
mucosal vascular, lymphocyte–HEV interaction, **65:362–363**
- PLN-specific vascular, *see* Peripheral node vascular addressins
- surface antigens of human leukocytes and, **49:88**
vascular, role of GlyCAM-1, **58:359**
- Adenocarcinoma, rat pancreatic, **54:324**
- Adenoma, pituitary hormones, **63:380**
- Adenosine deaminase, **64:69**
- Adenosine deaminase deficiency, **59:151, 246**
SCID and, **49:382–383, 390, 396**
- Adenosine receptors, **52:396–398**
- Adenovirus
cytotoxic T cells and, **65:279**
E3/19K glycoprotein, effects on antigen processing and presentation, **52:44–47**
hybrid resistance and, **41:368**
- Adenylate cyclase, immunosenescence and, **46:226**
- Adhesins, antigen-presenting cells and, **47:61**
- Adhesion, *see* Cell adhesion
- Adhesion molecules, *see* Cell adhesion molecules
- Adipocytes
B cell formation and, **41:210**
IFN- γ and, **62:91**
- Adipose tissue, cachectin and, **42:215, 222–223**
- Adjuvant, *see also* Freund's complete adjuvant
antigen concentration and, **40:91–92**
Freund's, **60:111–112, 116–117, 131**
T cell subsets and, **41:70, 71**
- incomplete Freund's, autoimmune thyroiditis and, **46:277**
- lipopeptide moiety, **60:141**
- liposome–lipid A/alum, **60:112**
- MAP/alum, **60:117–118**
- RIBI, **60:131–132**
type, and differential results, **60:132**
- Adjuvant arthritis, **64:289**
- Adoptive immunotherapy, **58:437–438**
- Adoptive T cell therapy, tumors, *see* Tumors, adoptive T cell therapy
- Adrenal corticosteroids, MG therapy and, **42:270–271**
- β -Adrenergic mediator, **48:162**
- Adrenocortotropic hormone, **48:170–172; 54:26–27**
in vitro IgE synthesis, **61:370–371**
NK cells and, **47:268, 269**
production by leukocytes, **39:315, 316**

- Adrenocorticotrophic hormone (*continued*)
 spontaneous autoimmune thyroiditis and, **47:477, 479, 493**
 thymic factor-induced release, **39:315**
- Adriamycin
 antitumor activity, **56:333–334**
 monoclonal antibody conjugation procedure, **56:334–336, 338**
- Adsorption
 experiments, IL-2 and, **42:166**
 hybrid resistance and, **41:363**
 T cells, **38:155–157**
- Adult respiratory distress syndrome, therapeutic regulation of complement system in
 clinical studies, **56:278–279**
 inhibition using soluble CRI, **56:284–286**
- Advanced glycosylation end products, **63:390–391**
- AES, *see* Anti-eosinophil serum
- Afferent blockade
 anterior chamber, **48:203–204**
 corneal graft survival, **48:195–196**
- Afferent lymph dendritic cells, CD1 protein expression, **59:63–64**
- Affinity chromatography, leukemia inhibitory factor, **53:32**
- Affinity maturation, memory B cells, **53:233–234**
- Affinity purification, aPL antibodies and, **49:228–231, 257, 259**
 specificity, **49:242–243, 251**
- Affinity threshold
 receptor-ligand interaction, **42:48–51**
 tolerance induction B cells, **42:77–78**
 variable region gene complexes and, **42:73**
- African swine fever virus, **65:127–128**
- Agammaglobulinemia, **38:291; 59:139–141**, *see also* X-linked agammaglobulinemia congenital, **59:138–139**
- Agarose gel electrophoresis, complement components and, **38:207, 208**
- Age
 antiidiotypic regulation and, **42:57–58**
 associated replacement of Vγ6 T cells by Vγ4, **58:303–304**
 and B cell genesis, **63:215–216**
 CD5 B cells and, **47:122, 123**
 related extrathymic expansion of Vδ2 T cells, **58:309**
 V gene segment utilization and, **42:30–32**
- Agglutination
 epitope, **48:23**
 hybrid resistance and, **41:342, 359**
 segmental flexibility, **48:23–24**
- Agglutination test, aPL antibodies and, **49:235**
- Agretopes
 immune response genes and
 antigen binding, **38:107**
 competitive inhibition, **38:108**
 cytotoxic T cells, **38:165**
 tolerance, **38:130**
 synthetic T and B cell sites and, **45:201, 202, 210**
- T cell receptor and
 antigen processing, **45:122, 123**
 structure-function relationships, **45:135, 139**
 T cell repertoire, **45:166**
- T cell subsets and, **41:53, 54**
- AIDS, *see also* AIDS, animal models
 anti-TNF-α therapy, **64:311**
 aPL antibodies and, **49:210**
 associated lymphopenia, and lymphocyte PCD, **58:214**
- CD4 molecules and, **44:290**
- CTL suppression in, **65:305**
 epidemic, **52:425–426**
 exacerbation, **58:16**
- HIV-1, **63:87–89**
 chronic infection, **63:89**
 immunity, **63:95–98**
 infection, **63:84**
 neuropathology, **63:95**
 pathogenesis, **63:90–95, 100**
 primary infection, **63:88**
 transmission, **63:88**
 treatment, **63:94–95**
 xenograft models, **63:79, 107**
 deletion mutant studies, **63:94, 105**
 hu-PBL-SCID model, **63:89–98**
 pediatric infection, **63:95**
 SCID-hu thy/liv model, **63:89–90, 99–107**
- HIV infection and, **47:377, 413, 414**
 activation, **47:398, 400, 403**
 etiological agent, **47:378, 379, 384**
 immune response, **47:405–409**
 immunopathogenic mechanism, **47:388–391, 394, 395, 397**

- neuropsychiatric manifestations, 47:403–405
- HIV progression to, 65:285, 299, 302, 304
- humoral immune response and, 45:78
- hybrid resistance and, 41:353
- and IL-6, 54:37–39
- HIV infection, 54:37–38
- Kaposi's sarcoma, 54:38–39
- membrane cofactor protein, 61:219–220
- NK cells and, 47:285, 302, 303
- synthetic T and B cell sites and, 45:219, 264
- virus-induced immunosuppression and, 45:335, 336, 351–357, 360, 363, 365
- AIDS, animal models, 52:427
- animal infection with HIV-1, 52:454–466
- chimpanzee, infection of, 52:455–461
- model for testing HIV-1 vaccines, 52:457–459
- pathogenesis of HIV-1 infection in, 52:455–457
- use to test therapies based on CD4, 52:459–461
- conclusions, 52:466
- mouse, SCID-hu, 52:463–466
- rabbit, infection with HIV-1, 52:461–463
- epidemic of AIDS, 52:425–426
- feline immunodeficiency virus, 52:437–441
- use to study antiviral therapy and vaccines, 52:440–441
- feline leukemia model, 52:434–436
- antiviral therapy, 52:436–437
- pathogenic mechanism of FeLV-FAIDS, 52:436
- models using viruses distantly related to HIV-1, 52:430–432
- murine leukemia virus, 52:432–434
- ungulate lentiviruses, 52:430–432
- murine retroviral models, 58:233–237
- relationships among viruses used, 52:426–430
- simian immunodeficiency virus
- infection of macaques, 52:442–452
- pathogenesis of simian genetic drift of SIV molecular clones *in vivo*, 52:451–452
- immunodeficiency virus infection of macaques, 52:443–452
- natural history of SIV infection of macaques, 52:443–448
- use of SIV molecular clones to define viral determinants of pathogenesis, 52:448–451
- phylogeny of nonhuman primate lentiviruses, 52:442–443
- SIV/macaque model for testing vaccines, 52:452–454
- AIDS-dementia complex, 47:377, 403
- AIDS-related complex
- immune response, 47:408
- immunopathogenic mechanism, 47:388, 395, 397, 403
- neuropsychiatric manifestations, 47:404
- Airway hyperresponsiveness, 51:351–352
- and leukotrienes, 51:344–346
- Airway hypersensitivity, guinea pig, IL-5 role, 57:173–174
- Airway inflammation, *see* Pulmonary inflammation
- Airway mucosa, in bronchial asthma, 51:326
- Airway tone, and acute asthmatic response, 51:350–351
- Alamethicin, cytolysis and, 41:315
- AL-amyloidosis, 64:220–221
- Alanine
- antinuclear antibodies and, 44:119
- genetically engineered antibody molecules and, 44:84
- IL-1 and, 44:158
- Albumin
- hybrid resistance and, 41:392, 394
- IL-1 and, 44:165, 170, 186
- membrane attack complex of complement and, 41:300
- ALDC, *see* Afferent lymph dendritic cells
- Algorithms, T cell epitope prediction, 66:78–84
- Alkaline phosphatase
- B cell formation and, 41:186, 213
- hybrid resistance and, 41:380
- Alkylating agent–antibody conjugates, as cancer treatment
- chlorambucil, 56:328–330
- cis-platinum, 56:329, 332
- melphalan, 56:329–331
- mitomycin C, 56:329, 332–333
- phenylenediamine mustard, 56:329, 331
- properties, 56:328–329
- structure, 56:328–329
- trenimon, 56:329, 331–332

- Alleles
 adoptive T cell therapy of tumors and, **49:297**
 aPL antibodies and, **49:255**
 complement receptor 1 and, **46:188, 189, 191**
 complement receptor 2 and, **46:202**
 Ig gene superfamily and, **44:27–30, 44, 46**
 Ig heavy-chain variable region genes and, **49:9, 14, 22–28**
 surface antigens of human leukocytes and, **49:81**
- Allele specificity, cytotoxic T cells and, **38:162, 165, 167**
- Allele-specific oligonucleotide, **48:118–119**
 hybridization, **48:118**
 polymerase chain reaction, **48:118**
- Allelic exclusion
 B cell development, **63:2, 19–21**
 chicken, **57:372–374**
 T cell receptor β -chain, in control of thymocyte by p56 lt , **56:164–166**
- Allergen challenge
 in fiberoptic bronchoscopy studies, **60:218–219**
 and increased bronchial hyperresponsiveness, **60:168–171**
- Allergens, **61:363–364**
 antibodies from combinatorial libraries, **57:228**
- CD23 antigen and, **49:152**
- Allergic asthma
 CD23 antigen and, **49:160**
 IL-4 storage by lung mast cells, **53:15**
- Allergic bronchopulmonary aspergillosis complication of asthma, **60:216**
 in pulmonary eosinophilia, **60:226**
- Allergic diseases, **60:215–216**
 CD23 antigen and, **49:174**
 contact activation in, **66:258–259**
 IgE biosynthesis and, **47:1**
 antibody response, **47:7, 29, 30, 38**
 binding factors, **47:13**
- Allergic pulmonary inflammation, **62:283–285**
- Allergic response, **61:341, 374–375**
 APCs, **61:361–364**
 conditioning, **61:368–369**
 hybrid resistance and, **41:351**
- IgE
 regulation, **61:369–374**
 synthesis, **61:342–344, 370–374**
- IL-4, **63:310**
 inflammation, *see* Inflammation, allergic
 mast cells, **61:360–361**
 nervous system role, **61:364–368, 374**
 neuroendocrine factors, **61:366–367, 369–374**
 prostaglandin E₂, **61:98**
 psychological factors, **61:368–369, 374–375**
 T cell regulatory potential, **61:344–360**
 T helper responses, **61:344–347**
 APCs, **61:361–363**
 CD8 $^+$, **61:358–360**
 cytokines, **61:347–355**
 extrinsic factors, **61:357–358**
 surface molecules, **61:355–357**
- Allergic rhinitis, **60:221**
- Allergic thyroiditis, murine lupus models and, **46:79**
- Allergy, human
 anti-Id autoantibodies and, **39:275–276**
 idiotypic network and, **39:275–276**
- Alloantibodies, hybrid resistance and, **41:356**
- Alloantigens
 B cell formation and, **41:190, 193, 194, 197, 198**
 cytotoxic T cells and, **41:135, 136, 169, 170**
 β 2-microglobulin, **41:156**
 amino acid, **41:159, 163**
 hybrid resistance and, **41:333**
 antibodies, **41:376**
 bone marrow cells, **41:336–340**
 leukemia/lymphoma cells, **41:359, 361**
 lymphoid cells, **41:354, 356**
 immunosenescence and, **46:221, 245, 252**
Ir genes and, **38:114, 115**
 presentation by cultured endothelial cells, **50:266–267**
 response to, EC costimulator in, **50:271–272**
- T cell subsets and
 cell surface molecules, **41:44**
 H-2 molecules in thymus, **41:99, 107–109**
 H-2-restricting antigen recognition, **41:63, 70, 74**
- T cell specificity, **41:111**
- Allogeneic cells, immunosenescence and, **46:250**

- Allogeneic inhibition, hybrid resistance and, **41**:334, 358–361, 369
- Allogeneic lymphocyte cytotoxicity, hybrid resistance and, **41**:356, 357
- Allografts
- autoimmune thyroiditis and, **46**:286
 - corneal, *see* Corneal allograft
 - cytotoxicity and, **41**:270
 - cytotoxic T cells and, **41**:135
 - hybrid resistance and
 - antibodies, **41**:379
 - antigen expression, **41**:404, 405, 408–410
 - bone marrow cells, **41**:348, 350
 - effector mechanisms, **41**:370
 - leukemia/lymphoma cells, **41**:366
 - lymphoid cells, **41**:355
 - macrophages, **41**:371, 372
 - marrow engraftment, **41**:387
 - marrow microenvironment, **41**:391
 - NK cells, **41**:372–375, 377
 - syngeneic stem cells, **41**:389, 390
 - T cells, **41**:383
 - rejection
 - CD28–B7 costimulation, **62**:142
 - IFN- γ and, **62**:101–103
 - T cell subsets and, **41**:43, 92–95
 - tumor, *see* Tumor allografts
- Alloimmune effector, anterior chamber, **48**:204
- Alloimmunization, human
- anti-HLA antibodies and, **39**:281–283
 - idiotypic network and, **39**:281–283
- Allophenic mice, *Ir* genes and, **38**:62, 63
- Alloreactivity
- autoreactive T cells and, **45**:420, 421, 425
 - B cell formation and, **41**:190
 - cytotoxic T cells and, **41**:135–137, 169, 170
 - β_2 -microglobulin, **41**:156
 - amino acid, **41**:159, 162, 163, 165
 - carbohydrate moieties, **41**:153
 - exon shuffling *41*, **41**:142, 144–148
 - monoclonal antibodies, **41**:156–158
 - somatic cell class I variants, **41**:167
 - hybrid resistance and, **41**:357, 372, 380, 381
- Immoral immune response and, **45**:40
- Ir* genes and, **38**:121–123
- synthetic T and B cell sites and, **45**:234
- T cell receptor and, **45**:148–157
 - homogeneity, **45**:128, 130
- MHC molecules, **45**:116
- peptides, **45**:138
- polymorphic residues, **45**:158, 159, 162
- structure-function relationships, **45**:136, 140
- T cell repertoire, **45**:165
- T cell subsets and, **41**:40
- cell surface molecules, **41**:44
 - H-2 alloantigen recognition, **41**:78–83
 - H-2 molecules in thymus, **41**:100–102
 - H-2-restricted antigen recognition, **41**:68
- Allospecificities, NK-defined, genetic analysis, **55**:348–349
- Allotype
- complement receptor 1 and, **46**:185, 187–190, 195, 196
 - immunosenescence and, **46**:248
 - $\alpha\beta$ heterodimer receptor structure, **42**:168
 - α chain, **43**:282, 284–287
 - free, retention of, **52**:63–64
 - IgA, structure and function, **40**:155–158
 - IL-2
 - cell division and, **42**:173
 - growth signal, **42**:172–173
 - α enhancer, T cell receptor gene, **43**:268
 - Alpha helix, amphipathic, **43**:204
 - Alternate antigen pathway, and trapping of immune complexes, **51**:265–269
 - Alternative complement pathway
 - antibodies as activators, **51**:28–29
 - particulate activators, **38**:384–386, 392
 - Alum adjuvant
 - adsorbed MAP, **60**:141
 - HIV V3 loop in, **60**:131
 - with liposome–lipid A, MAP in, **60**:112 (T1B), MAP in, **60**:117–118
 - Alum/*Borderella pertussis* (ABP), **46**:137
 - Alveolitis
 - CD23 antigen and, **49**:160
 - extrinsic allergic, **62**:267
 - Alzheimer's disease
 - causes, **66**:260–261
 - and IL-6, **54**:42–43
 - Amino acids
 - antibody diversity and, **42**:97
 - antigenic equivalence, **43**:68–72
 - antigen-presenting cells and, **47**:60, 66, 89
 - antinuclear antibodies and, **44**:106, 108, 125, 129–131

- Amino acids (*continued*)
 autoimmune demyelinating disease and, 49:358, 363, 366, 368
 autoimmune thyroiditis and, 46:317
 antigens, 46:268, 272
 genetic control, 46:284
 humoral responses, 46:302
 prevention, 46:315, 316
 B cell formation and, 41:197, 198
 CD1 proteins, 59:11–22
 CD4 molecules and, 44:285, 286, 289, 292, 295, 296
 CD5 B cell and, 47:145, 146
 CD8 molecules and, 44:272–276, 278
 CD23 antigen and, 49:156, 162–166
 complement receptor 1 and, 46:184, 186, 187, 194
 complement receptor 2 and, 46:202, 205, 210
 conservation, 43:115, 221–222
 contact residue, 43:119
 critical residue, 43:67–68
 critical to binding energy, 43:24
 cytology and, 41:285
 cytotoxic T cells and, 41:135, 170
 β_2 -microglobulin, 41:154
 exon shuffling, 41:142, 147, 148
 H-2 mutant strains, 41:158–163
 HLA class I subtypes, 41:163–165
 HLA subtypes, 41:163–165
 monoclonal antibodies, 41:157
 somatic cell class I variants, 41:166, 167
 energy penalty of voids, 43:72
 epitope, 43:65–67
 genetically engineered Abs and
 antigen-combining sites, 44:84
 biological properties, 44:83
 cloning, 44:73, 75
 expression, 44:78, 79
 production, 44:68
 HIV infection and, 47:386, 405–408
 human T cell activation and, 41:3, 5, 10, 13
 humoral immune response and, 45:18, 20
 hybrid resistance and, 41:394
 IgE biosynthesis and, 47:15–17, 20, 39
 Ig gene superfamily and
 evolution, 44:49
 homology unit, 44:3, 6, 7
 nonimmune receptor members, 44:40
 receptors, 44:11, 18, 21, 22
 Ig heavy-chain variable region genes and
 D segments, 49:28, 31
 J_H segments, 49:34
 V_H families, 49:7, 14, 18
 V_H gene expression, 49:35, 43, 51
 IL-1 and, 44:153, 154
 biological effects, 44:167, 168
 receptor, 44:179, 181, 182
 structure, 44:155–158
 TNF, 44:185, 188
 intrinsic hydrophilicity parameters, 43:49
 leukocyte integrins and, 46:154–156, 159, 174, 176
 lymphocyte homing and, 44:342, 355
 maps of Ig-like loci and, 46:2
 structure, 46:7, 8, 10, 12, 13, 15, 21, 25
 T cell receptor, 46:35, 38, 41
 membrane attack complex of complement and, 41:307
 MHC contact, 43:210
 murine lupus models and, 46:69
 pore formers and, 41:313–318
 propensity factors, sequential epitopes, 43:66–67
 regulators of complement activation and, 45:381, 411
 genes, 45:405, 406
 protein expression, 45:399, 400, 402
 RCA-like protein utilization, 45:409, 410
 short consensus repeat, 45:391, 392, 394
 replaceability, 43:69–71, 77
 sequential epitope role, 43:39
 sporozoite malaria vaccine and
 antibodies, 45:300
 CS proteins, 45:292, 294–300
 CS-specific T cells, 45:311–313
 human trials, 45:315
 Plasmodium vivax, 45:317
 surface antigens of human leukocytes and, 49:81, 92
 synthetic T and B cell sites and, 45:196, 255, 256, 258, 262
 antigens, 45:232, 233
 bacterial antigens, 45:230, 231
 candidate synthetic peptide vaccines, 45:253, 255, 256, 258
 globular protein antigens, 45:210, 216
 immunological considerations, 45:200
 prediction, 45:251
 viral antigens, 45:219–221, 227, 228

- T cell receptor, **43**:157–159
 T cell receptor and
 alloreactivity, **45**:151–153
 antigen processing, **45**:120–123
 epitopes, **45**:143, 145, 148
 peptides, **45**:138
 polymorphic residues, **45**:159
 T cell repertoire, **45**:165
 T cell subsets and, **41**:85
 activated T cells and hybridomas,
 41:54, 56
 cell surface molecules, **41**:46, 47
 H-2 alloantigen recognition, **41**:83
 virus-induced immunosuppression and,
 45:350, 362, 364
 VL and VH domains, **43**:109, 125
 Amino acid sequence, **48**:108
 activin type II receptor, **55**:190–192
 amino-terminal, peptides, **54**:350
 antibodies from combinatorial libraries
 analysis, **57**:233–238
 RGD, **57**:261–263
 Ars A response and, **42**:95, 97, 111–112
 heavy chains and, **42**:99
 J_{H2} segment and, **42**:102–108
 light chains and, **42**:139–141
 chemokine, **55**:98–99
 chemokine receptors, **55**:129, 132–133
 desmoglein, **53**:314
 ELR motif of CXC chemokines,
 55:136–138
 IL-8 receptor, **55**:129–131
 interferon regulatory factors 1 and 2,
 mouse, **52**:268
 LIF, human, **53**:40
 oncostatin M, **53**:40
 PV antigen, **53**:314
 TCR γ genes, human
 C_γ, **43**:143
 J_γ, **43**:142
 V_γ, **43**:141
 TGF-β receptor II, **55**:190–192
 V_{B1} and V_{B2} gene segments, chicken,
 50:110
 Aminoacetylation, antinuclear antibodies and,
 44:94, 119, 127, 134, 135, 138
 3-Aminobenzamide, effect on cytolysis,
 58:252–253
 Aminopterin, in cancer treatment, monoclonal
 antibody conjugation procedure, **56**:342
 5-Aminosalicylic acid, *see also* Sulfasalazine
 prostaglandin synthesis and, **42**:312–313,
 317, 318–329
 AML1, **65**:24–25
 Amoebas, pore-forming proteins of, **41**:312,
 313
 Amphiphaticity, T cell receptor, **43**:204
 Amphiphilic nature, C5b-9, membrane attack
 complex of complement and, **41**:299–304
 Amyloidosis, **64**:219, 220–221
 Anaphylatoxins, **61**:204, 255
 Anaphylaxis, IFN-γ and, **62**:103
 Anatomical sequestration, lens, **48**:220
 Anchor, GPI, defective synthesis, **60**:57–74
 Anchor residues, **62**:231
 Androgens, spontaneous autoimmune
 thyroiditis and, **47**:480, 481
 Anemia, **59**:146–147, 343
 aPL antibodies and, **49**:218, 225, 250–251,
 255, 257
 aplastic, PNH associated with, **60**:91
 erythropoiesis and, **42**:224
 hemolytic, in PNH, **60**:74–75
 Anemic cells, hybrid resistance and, **41**:346,
 348, 388
 Anemic mouse, B cell formation and, **41**:224
 Anergy
 B cells, **59**:321–322
 γδ T cells, **58**:313
 and peripheral tolerance to cancer, **58**:421
 resulting from tolerization processes,
 58:271–272
 surviving T cells, **58**:260
 T cell, cyclosporin A effect, **58**:262
 Angioedema, hereditary, **61**:206
 Angiogenesis, in rheumatoid arthritis, **64**:295
 Angloimmunoblastic lymphadenopathy, gene
 rearrangement and, **40**:294
 Angiotensin II, synthetic T and B cell sites
 and, **45**:232, 233
 Anhydrides
 cis-aconitic, in conjugation procedure for
 chemoimmunoconjugates in cancer
 treatment, **56**:318–319
 mixed, in conjugation procedure for
 chemoimmunoconjugates in cancer
 treatment, **56**:319
 Anions, cytolysis and, **41**:282, 312
 ANK repeat, **65**:14
 Ankylosing spondylitis, HLA-B27 and, **65**:94

- Ankyrin repeats, interaction with NF- κ B rel domain, **58**:3–4
- Anopheles*, sporozoite malaria vaccine and, **45**:283, 284
- Anorexia, cachectin and, **42**:213, 219
- Anterior chamber, **48**:191, 192
afferent blockade, **48**:203–204
alloimmune effector, **48**:204
alloimmunity down-regulation, **48**:204
antigen-processing cells, **48**:206, 207
DBA/2 mastocytoma cell, **48**:204
immunological privilege, **48**:203–212
intraocular HSV-1 infection, **48**:210
 contralateral eye, **48**:210
 DTH suppression, **48**:210
mastocytoma, **48**:213–215
physiochemical environment, **48**:206, 207
systemic cell-mediated immunity, **48**:204
tumor allograft, **48**:204
tumorigenesis, **48**:213–215
UV5C25, **48**:213–215
- Anterior chamber-associated immune deviation, **48**:205–211, 206
- cytotoxic T cell, **48**:208, 209
- DBA/2 mastocytoma model, **48**:207
- delayed-type hypersensitivity, **48**:219
 antigen-specific down-regulation, **48**:208, 209
 antigen-specific suppression, **48**:211
 anterior chamber, **48**:211
 efferent mode, **48**:211
 hapten model, **48**:211
 suppressor T cell, **48**:211
 intracellular tumor allograft, **48**:208, 209
- effector mechanism, **48**:211–212
- expression, **48**:207–210
- eye, **48**:206
induction, **48**:205–207
ocular HSV infection, **48**:208–210
S antigen, **48**:219
spleen, **48**:206, 211
tumor allograft, intraocular, **48**:207–208
tumorigenesis, **48**:213
vascular route, **48**:205
- Anthracycline–antibody conjugates, as cancer treatment
- direct conjugation
 C-14 methyl group, **56**:336
 hydroxyl group, **56**:336–337
 keto group, **56**:337
- sugar amino group, **56**:334–336
sugar moiety, **56**:334
- intermediary carriers
 dextran, **56**:337–338
 polyglutamatic acid, **56**:338
- properties, **56**:333–334
structure, **56**:333–334
- Antihadhesion therapy, **64**:195–197
- Antibodies, **48**:16–18
 adoptive T cell therapy of tumors and, **49**:331, 333
 mechanisms, **49**:312–315
 principles, **49**:289, 297
affinity threshold of, **42**:69, 70–71
- anti-BrMRBC/anti-PC, secreting CD5 B cells, **55**:315–316
- antidansyl
 genetically engineered antibody molecules and, **44**:78, 80, 81
 segmental flexibility, **48**:13, 14
- anti-DNA, **38**:297, 298
- antigen binding, **43**:105
- antigen-presenting cells and
 antigen presentation, **47**:68, 73, 75
 antigen processing, **47**:92
 APC-T cell binding, **47**:95, 97, 99
 cell surface, **47**:57–59, 61, 62
 immunogeneity, **47**:102, 103
 interaction with T cells, **47**:93
 T cell-dependent antibody responses, **47**:81–86
- T cell growth, **47**:100, 101
- tissue distribution, **47**:49, 50, 53, 57
- anti-Id, experimental autoimmune encephalomyelitis, **48**:164, 166
- anti-Id
 autoimmune thyroiditis and, **46**:318
 genetic control, **46**:287
 humoral responses, **46**:301, 304–309
 prevention, **46**:308, 312, 314
 Ig heavy-chain variable region genes and, **49**:37, 40, 46
 immunosenescence and, **46**:247, 248, 252, 253
 MG therapy and, **42**:268, 272
 specific immunoregulation, **42**:55–57
- anti-Id, animal, as vaccines, protection from bacterial antigens, **39**:283
parasite antigens, **39**:284
viral antigens, **39**:283

- anti-Id, human therapeutic
 B cell tumors and, **39**:285–291
solid tumors and, **39**:291–292
 T cell leukemias and, **39**:288
- anti-Id, rabbit
 to human anti-TT antibodies
 binding to B cells, **39**:264–266
 binding to Id, **39**:261–263
 binding to T cells, **39**:265–267
 human immune responses and,
 39:267–269
 production, **39**:261
 to human IgM-RF, **39**:263
- anti-Ig, **52**:185–188
 anti-TNF- α therapy, **64**:323–325
Branhamella catarrhalis and others,
52:188
 induced proliferation of normal human B
 cells, **54**:363
 inhibitory effects, **52**:186–187
 inhibitory effects of anti-Ig antibodies,
52:186–187
Staphylococcus aureus Strain Cowan,
52:187–188
 stimulatory effects, **52**:185–186
 as surrogate antigens, differential effects
 on mature and immature cells,
52:295–297
- anti-IgE, **43**:279–281
- anti-IgM
 growth arrest and apoptosis induced by,
54:410–411
 induced unresponsiveness in adult B
 cells, **54**:404–405
 role of apoptosis in neonatal B cell
 tolerance, **54**:403–404
 surrogate systems, **54**:402–410
 tolerance in B cell subsets, **54**:405–407
- anti-IgM and anti-CD40, costimulatory
 effect on B cell proliferation, **52**:190
- antinuclear, *see* Antinuclear antibodies
- antipeptide, **43**:61
 binding, **43**:118
 design, **43**:59–65
 affinities of monoclonal anti-MH_r
 antibodies, **43**:61–62
 local mobility and antigenicity, **43**:61
 antiphospholipid, *See* Antiphospholipid
 antibodies
- antiprotein, binding, **43**:42
- antiprothrombin, **49**:247, 256
- antisporozoite, **60**:112–124
- antithrombin III, **49**:256
- autogeneic, *see* Autoantibodies
- autoimmune demyelinating disease and,
49:363, 368–374
- autoimmune thyroiditis and, **46**:263,
 317–319
- antigens, **46**:272, 273
- cellular immune responses, **46**:287, 289,
 290, 293–295
- experimental models, **46**:276–278, 281,
 282
- genetic control, **46**:283, 285–287
- humoral responses, **46**:298–307, 309
 prevention, **46**:308, 310, 312, 314, 315
- autoreactive T cells and, **45**:427
- B cell formation and, **41**:182, 183
 B cell precursors, **41**:189, 191, 192, 194,
 201, 202
- bone marrow cultures, **41**:213
- Ig genes, **41**:203, 204
- lymphohemopoietic tissue organization,
41:187
- C1q interaction, **51**:9–21
- catalysis, **43**:1–2
- CD1, **59**:39–43
- CD2, role in T cell death, **58**:226
- CD3, IGF-I-R, **63**:403
- CD4 molecules and, **44**:297, 300
- CD5 B cell and, **47**:161, 162
 Ig gene expression, **47**:150–154, 157,
 158
- lineage, **47**:129
- marker for activation, **47**:127
- physiology, **47**:130, 131, 133, 135–137,
 139, 140
- primordial immune network, **47**:159–161
- surface antigen, **47**:143, 144
- CD23 antigen and, **49**:149, 176
 biological activity, **49**:168–169, 172
 cellular expression, **49**:152–153, 155
 expression regulation, **49**:159
 Fc ϵ RII, **49**:165
- CD40, **52**:188–191
 and anti-IgM, costimulatory effect on B
 cell proliferation, **52**:190
- CD59, in venous thrombosis, **60**:91–92
- cell mediation, **41**:271, 272

- Antibodies (*continued*)
 cold-reactive antilymphocyte, IgG levels in, **42:292**
 combining site, structure variability, **43:14–15**
 complement receptor 1 and, **46:193, 196, 200, 201**
 complement receptor 2 and, **46:206**
 complex with antigens, trapping by follicular dendritic cells, **60:274**
 conformation, **43:17, 124–125**
 crystallographic studies, I diversity, **43:133**
 cytolysis, **41:275, 278, 283**
 cytotoxic T cells and
 β_2 -microglobulin, **41:156**
 exon shuffling, **41:141, 144–146**
 HLA class I antigens, **41:150, 151**
 monoclonal antibodies, **41:156–158**
 δ dextran
 class switching, **61:91–92**
 Ig production, **63:65**
 Dob structure, **43:8**
 to double-stranded cDNA, cA2 treatment, **64:325**
 electrostatic recognition with antigen, **43:12–13**
 enzymatic activity, **43:81**
 F23.1, induction of thymic cell depletion, **58:274**
 Fab Kol, crystal packing, **43:19–20**
 Fab region, **43:8–9**
 Fab structures, **43:17**
 Fc region, **43:8–9**
 genetically engineered, *see* Genetically engineered antibodies
 glycoprotein, phase I clinical trials, **40:366–370**
 helper T cell cytokines and
 cross-regulation of differentiation, **46:131, 132**
 differential induction, **46:127, 129, 130**
 functions, **46:116, 118–120**
 immune responses, **46:133, 137, 138**
 precursors of differentiation states, **46:124**
 HIV-1 immunity, **63:97–98**
 HIV infection and
 immune response, **47:405–410**
 immunopathogenic mechanism, **47:388, 389, 393, 395, 398**
- neuropsychiatric manifestations, **47:404**
 human, from combinatorial libraries
 affinity, strategy for improvement, **57:263–266**
 allergens, **57:228**
 amino acid sequences
 analysis, **57:233–238**
 RGD, **57:261–263**
 bacteria, **57:228**
 cloning strategies, **57:203–206**
 design, **57:261–263**
 Epstein–Barr virus-transformed cell line—
 derived, **57:229–230**
 expression of antibody fragments, **57:207–208**
 to Graves' ophthalmopathy self-antigens
 from human donors, **57:245**
 to HIV-1 self-antigens from human
 donors, **57:246–250**
 HuSCID mouse-derived, **57:230–231**
 hybridomas, **57:229–230**
 immune donors, **57:208–242**
 naive repertoire approach, **57:250–252**
 overview, **57:191–192**
 phage display
 antibody fragments, **57:195–198**
 vector systems, **57:199–203**
 to primary biliary cirrhosis self-antigens
 from human donors, **57:244–245**
 primate-derived, **57:231–232**
 principles of combinatorial approach, **57:192–195, 266–267**
 selection strategies, **57:206–207, 261–263**
 staphylococcal protein A, **57:229**
 study of responses, **57:232–242**
 synthetic repertoire approach, **57:250, 252–260**
 to thyroid disease self-antigens from
 human donors, **57:242–244**
 viruses, **57:209–228**
 cytomegalovirus, **57:220–221**
 hepatitis B virus, **57:220**
 herpes simplex virus type 1, **57:222–227**
 herpes simplex virus type 2, **57:222–227**
 HIV-1, **57:211–217**
 measles, **57:227–228**

- respiratory syncytial virus, **57**:217–219
- varicella Zoster virus, **57**:221–222
- whole antibody molecules, **57**:208
- human antichimeric, **64**:323–324, 330
- human anti-mouse, **64**:323
 - immune response, chemotherapy limited by, **56**:304, 358–360
- humoral immune response and, **45**:1–6
 - antigens, **45**:10, 12
 - cellular interactions *in vivo*, **45**:78, 81–83, 85
 - helper T cells, **45**:24–30, 32
 - interleukins, **45**:62, 67, 72, 74
 - physical interaction, **45**:38, 40–54
- hybrid resistance and
 - antigen expression, **41**:400, 402–405
 - bone marrow cells, **41**:336, 338, 341, 349, 350
 - effector mechanisms, **41**:370, 376–380
 - leukemia/lymphoma cells, **41**:359, 365, 368
 - lymphoid cells, **41**:352, 356, 357
 - macrophages, **41**:371, 372
 - marrow engraftment, **41**:386, 387
 - NK cells, **41**:374
 - T cells, **41**:383
- IBD and, **42**:290–291, 294–295, 298
- IFN- γ , **62**:63, 87–89
- IgE biosynthesis and, **47**:1
 - binding factors, **47**:12, 14, 15, 17, 19, 21–23, 25, 28
 - response, **47**:1–12
 - response suppression, **47**:28–39
- Ig gene superfamily and, **44**:4, 9, 38, 44
- Ig heavy-chain variable region genes and, **49**:1–3, 61
 - D segments, **49**:31–33
 - J_H segments, **49**:34
 - organization, **49**:4, 6
 - V_H families, **49**:9, 14, 16
 - V_H gene expression, **49**:40–45
- IL-1, **44**:189, 190, 192; **66**:140
 - gene expression, **44**:163
 - immunocompetent cells, **44**:173, 177, 178
 - receptor, **44**:182
 - structure, **44**:157, 158
 - systemic effects, **44**:169
- TNF, **44**:187
- IL-6, **66**:132
 - clinical applications, **54**:50
- immunosenescence and, **46**:221, 253
 - lymphocyte activation, **46**:222
 - mucosal immunity, **46**:252
 - regulatory changes, **46**:247, 248, 250, 251
- stem cells, **46**:246
- induced complementarity, **43**:16–20
 - immunological, biochemical and crystallographic data, **43**:16–18
 - mobility data, **43**:20
 - variable domain structure superposition, **43**:18–20
- to Leu7, **42**:191–192
- leukocyte integrins and, **46**:164
- lung, **59**:393–394, 411–412
- lymphocyte homing and
 - high endothelial venules, **44**:317, 318
 - immune response, **44**:330–332, 335
 - molecules, **44**:342–344, 346, 350, 352
 - regional specificity, **44**:326, 328, 338, 341
- lymphotoxin and, **41**:298
- membrane attack complex of complement and, **41**:301, 302, 307, 309
- MG and, **42**:262
- monoclonal, *see* Monoclonal antibodies
- murine lupus models and, **46**:99
 - Ig germline, **46**:65, 66, 72, 74, 76, 77
 - lupus strains, **46**:64
 - T cell antigen receptor, **46**:79, 88, 90, 98
- neutralizing, **48**:95–96
 - HIV infection and, **47**:407, 408, 410, 413
- NK cells and, **47**:189
 - adaptive immunity, **47**:292–294
 - alteration, **47**:300
 - antimicrobial activity, **47**:286–288
 - cell-mediated cytotoxicity, **47**:190, 196
 - congenital defects, **47**:224
 - cytotoxicity, **47**:249–251, 254, 257–259, 261
 - differentiation, **47**:231, 233
 - effector mechanisms, **47**:235, 237, 238, 240, 241, 243–248
 - hematopoiesis, **47**:274, 275, 277, 281
 - identification, **47**:196, 198
 - malignant expansion, **47**:227–229
 - surface phenotype, **47**:201, 202, 204–213
 - tissue distribution, **47**:221

- Antibodies (*continued*)
 PGHS-1 or PGHS-2, **62**:178
 phosphorylcholine, **42**:10
 PM-1, reshaped, **64**:261
 polyclonal, phagocytosis and, **38**:376
 pore-forming protein, **41**:291, 296
 to rabbit molecules, **56**:182–183
 raised against native protein, **43**:3
 receptor structure, *see* Antibodies, structure
 regulators of complement activation and
 complement pathways, **45**:381
 protein expression, **45**:396, 397
 protein roles, **45**:389, 390
 reshaped, human PM-1, construction,
 64:261
 SCID and, **49**:397, 401
 secondary B cell lineage and, **42**:69–71
 secretory, **65**:249, 250
 sequence variation, **43**:13–16
 serum, **42**:291–292
 serum, IBD and, **42**:291–292
 SLE and, **42**:291–292
 spontaneous autoimmune thyroiditis and,
 47:433, 492
 altered thyroid function, **47**:460
 disturbed immunoregulation, **47**:479
 histopathology, **47**:442
 humoral immune reactions, **47**:444, 445,
 447, 479
 potential effector mechanisms, **47**:468
 sporozoite malaria vaccine and, **45**:320, 321
 CS proteins, **45**:292, 299, 300
 CS-specific T cells, **45**:306–313
 endemic areas, **45**:315–317
 human trials, **45**:313–315
 immunity, **45**:286–292
 interferon- γ , **45**:301, 304, 306
 Plasmodium vivax, **45**:317, 318
 role, **45**:300, 301
 structure
 conformational changes, **43**:113–114
 domain-domain interactions, **43**:110
 domain structure, **43**:107–109
 Fab fragments, **43**:107
 H and L chain CDR, **43**:108–109
 module structure, **43**:110–113
 polypeptide loops, **43**:107
 sheet-sheet interactions, **43**:110
 V and C domains, **43**:109
 variable domains, **43**:109
 VL–VH interface, **43**:110, 112, 124
 VL–VL pairing, **43**:112
 V module interface, **43**:111
 surface antigens of human leukocytes and,
 49:113, 117
 synthetic T and B cell sites and, **45**:196,
 260–264
 antigens, **45**:233, 236
 bacterial antigens, **45**:231
 candidate synthetic peptide vaccines,
 45:253, 255–259
 globular protein antigens, **45**:215, 216
 immunological considerations, **45**:197
 parasitic antigens, **45**:228, 229
 peptides, **45**:237–239
 prediction, **45**:251
 viral antigens, **45**:220–223, 225, 227
 Tac, **42**:167, 169, 171, 172
 T cell activation and, **41**:6, 7, 9, 11, 13
 T cell development and, **44**:207, 255
 ontogeny, **44**:218, 221
 thymocyte subpopulation, **44**:226, 237,
 239, 241
 T cell receptor and
 accessory molecules, **45**:108, 109
 alloreactivity, **45**:152, 154
 homogeneity, **45**:128, 130
 peptides, **45**:138
 polymorphic residues, **45**:158, 160
 structure-function relationships, **45**:136,
 140
 T cell repertoire, **45**:164
 T cell subsets and, **41**:39, 40
 activated T cells and hybridomas,
 41:54, 56
 cell surface molecules, **41**:49
 effector phase, **41**:76
 H-2 alloantigen recognition, **41**:83–86,
 89–93
 H-2 molecules in thymus, **41**:97, 98, 105
 H-2 restricted recognition of antigen,
 41:50, 52
 T accessory molecules, **41**:61, 62
 T cell receptor, **41**:40
 unprimed and resting T cells, **41**:64, 65,
 70, 72
 to TCR1, TCR2, and TCR3, embryonic
 treatment with, **50**:103–105
 TCR antibody, **48**:90
 TNF- α , *see* Anti-TNF- α therapy

- to TNF, treatment with, **66**:124, 132, 150
TNKTR, **42**:201–202
trimolecular complex
 anti- α/β T cell receptor antibody, **56**:228–229
 anti-CD3 antibody, **56**:220–223
 anti-CD4 antibody, **56**:223–228
 anti-T cell receptor V β antibody, **56**:229–233
virus-induced immunosuppression and HIV, **45**:351–353, 356, 357, 363, 365, 368
 measles, **45**:340, 341, 343, 345–347
X-ray analysis, **43**:121
Antibody–antigen interaction, **43**:99–128, *see also* Antigen recognition
 affinities to small molecules, **43**:82
 analysis of structure, **43**:22
 buried surface, **43**:82
 chemistry and binding energy, **43**:72–84
 concerted shifts, **43**:22
 contact residue, **43**:26
 criteria for systematic studies, **43**:5–8
 critical residues, **43**:27
 crystallographic approaches, **43**:4
 degree of fidelity, **43**:60
 dominant epitopes and collective response, **43**:46–49
 electrostatic attraction, **43**:80
 evaluation of predictive methods
 criteria for assessing prediction success, **43**:50–59
 sequence data, **43**:49–50
 immunological cross-reactivity, **43**:64
 interface adaptor hypothesis, **43**:125–128
 local antibody conformation effect, **43**:17–18
 lock-and-key fit, **43**:78, 128
MHr, **43**:42
 mobile structural elements, **43**:80
 nature of immune response, **43**:83
 recognition and binding components, **43**:79
 reconciling crystallographic and peptide mapping data, **43**:23–27
 surface mobility role, **43**:16–17
 synthesis of structural description, **43**:26
 three-dimensional structures, **43**:119–120
 complementarity, **43**:122
 conformational changes, **43**:123–125
 epitope and paratope size, **43**:120–121
 epitope character, **43**:121–122
 X-ray crystallography, **43**:8–11
 X-ray structures, **43**:73
Antibody class switching, **61**:79–81
 heavy chain isotype expression, **61**:82–84
 problems in other studies, **61**:81–82
 regulation of isotype specificity, **61**:84–89
 switch recombination, **61**:99–130
Antibody complexes, crystallographic structures, **43**:4
Antibody-dependent cellular cytotoxicity, **41**:269, 273; **46**:114, 122; **58**:223
 adoptive T cell therapy of tumors and, **49**:312–313
 cytolysis and, **41**:275, 277–281, 283
 eosinophil, for helminthic parasites, **60**:211–213
 Fc γ R-mediated, **60**:180
 HIV infection and, **47**:384, 388, 405, 410, 413
 hybrid resistance and
 bone marrow cells, **41**:336, 338
 effector mechanisms, **41**:376–378
 lymphoid cells, **41**:351
 macrophages, **41**:371
 T cells, **41**:383
 mechanisms
 IBD and, **42**:294–295
 NK cells and, **42**:191, 192, 203–204
 membrane attack complex of complement, **41**:307
 NK cells and, **47**:189, 190
 antimicrobial activity, **47**:285
 cytotoxicity, **47**:251, 253, 258
 effector mechanisms, **47**:241–243
 genetic control, **47**:222
 malignant expansion, **47**:227, 228
 reproduction, **47**:271, 272
 surface phenotype, **47**:204, 212, 213
 spontaneous autoimmune thyroiditis and, **47**:451, 452, 466, 469
Antibody diversity, *see Arsonate idiotypic system, A/J mouse*
Antibody effector function, human
 bacterial Fc receptors, **51**:56–60
 catabolism of antibodies, **51**:54–55
 complement activation by, **51**:7–29
 antibodies as activators of alternative pathway, **51**:28–29

- Antibody effector function (*continued*)**
- C1 activation, **51:21–25**
 - C3 activation, **51:25–28**
 - C4 activation, **51:25–28**
 - cell lysis, **51:25–28**
 - interaction of C1q and antibody, **51:8–21**
 - conclusions, **51:62–64**
 - human leukocyte Fc Receptors, **51:29–54**
 - molecular explanation, **51:1–2**
 - structure, considerations, **51:2–7**
 - Fc, **51:2–4**
 - IgG, conformation and flexibility, **51:4–7**
 - other antibody classes, **51:7**
- Antibody–hapten complexes, 43:119**
- Antibody isotype switching, 48:83–84**
- Antibody-mediated complement-dependent cytotoxicity, HIV infection and, 47:409, 410, 413**
- Antibody repertoire, generation in rabbit**
- B cell development
 - B lymphopoiesis, **56:180–186**
 - immune response ontogeny, **56:186–187, 209**
 - development of repertoire
 - contributing factors, **56:194–195**
 - somatic gene conversion, **56:198–202**
 - somatic mutation, **56:201, 203**
 - V_H allotype, **56:195**
 - V_H usage, **56:195–198**
- gut-associated lymphoid tissue**
- as bursal equivalent, **56:204**
 - follicular structure, **56:203**
 - germfree rabbits, **56:204–206**
 - model, **56:206–209**
- Ig genes**
- C_H, **56:191–193**
 - D, **56:189–191**
 - J_H, **56:188–191**
 - κ light-chain, **56:193–194**
 - λ light-chain, **56:194**
 - organization, **56:187–188**
 - V_H, **56:188–191, 195–198**
 - mechanism, **56:179–180, 209**
- Antibody response, 45:4**
- B cell proliferation and, **40:6–12**
 - Ir* genes and
 - gene dosage, **38:31, 80**
 - insulin, **38:142, 143**
 - T cell suppression, **38:135, 147**
- T cell-T cell interactions, **38:154, 156, 157, 160**
 - tolerance, **38:128, 129**
 - model of, **40:1–2, 10, 45**
 - suppression by CTLs, **60:306–308**
 - T cell dependent, **40:3–4**
- Antibody secretion**
- IBD and, **42:301–307**
 - inhibition by specific T cells, **40:139–140, 141**
- Anti-bromelaine-treated red blood cells, 46:72, 74**
- Anticardiolipin antibodies, 49:193, 259**
- affinity purification, **49:230–232**
 - antibody subsets, **49:233**
 - antimitochondrial antibodies, **49:249–250**
 - binding to cell membrane, **49:250–252**
 - clinical aspects, **49:203–212, 215–218, 220–224**
 - history, **49:199**
 - isotype, **49:224–227**
 - LA antibodies, **49:241, 243–247**
 - pathogenic potential, **49:254–255, 257**
 - reactivity, **49:248–249**
 - specificity, **49:234–240**
- Anticholinesterase drugs, MG and, 42:269–270, 271**
- Anticytokine therapy, 66:151**
- Anti-DNA transgenic model, 52:314–315**
- Antiendothelial antibodies, 49:252**
- Anti-eosinophil serum, parasite-infected animals and, 39:209**
- Antiergotypic T cell, 48:166**
- Antigen-antibody binding, first stage, 43:79**
- Antigen binding**
- Ars A response and, **42:109, 139–141**
 - assay, *Ir* genes and, **38:35**
 - cachectin and, **42:225**
 - α chain, **42:168–172**
 - Ir* genes and
 - competitive inhibition, **38:108–110**
 - determinant selection, **38:98, 99**
 - Ia molecules, pigeon cytochrome c and, **38:99–104**
 - macrophages, **38:111–115**
 - T cell activation, **38:104–108**
 - T cell-T cell interactions, **38:155, 156,**

- Antigen-binding lymphocytes, **38:285, 286**
Antigen bridging, T cell suppression and, **38:142, 143, 151**
Antigen catabolism, *Ir* genes and, **38:95, 96**
Antigen-dependent activation, **52:199**
Antigenic determinants, *see* Epitopes
Antigenicity
 chemistry, **43:2**
 epitope prediction, **43:58**
 flexibility, **43:105, 118**
 relationship with local mobility, **43:61**
 secondary structure, **43:76**
 in terms of recognition frequency, **43:29–32**
 value, **43:50**
Antigenic microassemblies, **43:42**
Antigenic peptide, KAVYNFATC, induced positive selection of thymocytes
 Antigenic peptide, **59:126**
Antigenic signaling, constraints to biochemical study of, **52:294–295**
Antigenic stimulation, **42:19–22**
 memory generation by, **53:221–223**
 sensitivity of naive and memory T cells to, **53:228**
 T cell differentiation after, **53:229–230**
Antigenic variation, Delbrück model of, **38:340–344**
Antigen-independent recruitment, T cells into tissues by endothelial cells, **50:274–287**
 expression of endothelial leukocyte adhesion molecules *in vivo*, **50:283–284**
 extravasation of T cells, **50:285–287**
 leukocyte adhesion molecules, **50:279–281**
 migration of memory T cells to inflammatory sites, **50:278–284**
 migration of naive T cells to lymph nodes, **50:275–278**
 migration of pre-T cells to thymus, **50:275**
 role of endothelial leukocyte adhesion molecules in lymphocyte binding *in vitro*, **50:281–283**
 tissue-specific homing of lymphocytes, **50:284–285**
Antigen-induced arthritis, **64:289**
Antigen–MHC determinant, formation
 antigen competition, **43:208–211**
 binding of antigen and MHC molecules in solution, **43:211–213**
 immune response gene control, **43:206–208**
Antigen–MHC interactions, **43:219**
 binding rate constants, **43:212**
 correlation with immune-response phenotype, **43:212**
Antigen pathway, alternate, and trapping of immune complexes, **51:265–269**
Antigen polymerization, and B cell tolerance, **52:287–290**
Felton's paralysis and other early results with polymers, **52:287–288**
polymeric antigens and IgE responses, **52:289–290**
size-fractionated polymers: Dintzis model, **52:288–289**
Antigen presentation, *see also* Antigen processing and presentation
 by B cells, tolerance, **52:321–322**
 Ir genes and, **38:189**
 antigen binding, **38:98–115**
 antigen processing, **38:93–97**
 blocking, **38:69–71**
 complementation, **38:74–79**
 cytotoxic T cells, **38:164, 165, 186**
 expression, **38:55–60, 188**
 gene dosage, **38:81**
 H-Y antigen, **38:176, 177, 179**
 Ia molecules, **38:83, 84**
 I region mutations, **38:88–90**
 MHC restriction, **38:64, 66**
 T cells, **38:153, 157, 158, 160**
 proliferation, inhibition of, **38:68, 69**
 selection, **38:131, 133, 134**
 suppression, **38:151, 152**
 tolerance, **38:120, 122, 123, 125, 127, 128, 130**
 islet beta cell, **48:142**
 protein antigen, by cultured endothelial cells, **50:265–266**
to T cells by
 B cell lymphomas, **39:54, 63–66, 86**
 B cells, **39:52–54**
 comparison with non-B cells, **39:85–86**.
 Ig role, **39:52–53, 55–56, 58–59**

- Antigen-presenting cells, **47:45–47**, 105; **48:232**; **62:231**
 adoptive T cell therapy of tumors and, **49:283**, 291, 334
 expression, **49:324–325**, 328–329, 331
 recognition, **49:319–321**, 323
 antigen presentation, **47:64**
 allogeneic MHC products, **47:73–76**
 H-Y antigens, **47:80**
 Mls stimulation, **47:76–79**
 protein, **47:65–73**
 TNP-modified syngeneic cells, **47:79**, 80
 viral antigens, **47:80**, 81
 antigen processing
 B cells, **47:89**, 90
 dendritic cells, **47:90**, 91
 MHC-peptide complexes, **47:87–89**
 MHC products, **47:92**
 for antitumor immunity, **58:436–437**
 APC-T cell binding, **47:95–99**
 autoimmune demyelinating disease and, **49:367**
 autoimmune thyroiditis and, **46:266**, 317
 cellular immune responses, **46:289**, 294–297
 experimental models, **46:281**
 autoimmunity, **47:104**, 105
 autoreactive T cells and
 activation, **45:421–427**
 origin, **45:419**
 regulation, **45:432**
 B cell formation and, **41:225**
 CD23 antigen and, **49:167–168**
 cell surface, **47:57**
 CD5 leukocyte common antigen, **47:60**, 61
 integrins, **47:61**
 MHC products, **47:57–60**
 receptors, **47:62–64**
 eosinophils as, **60:200**
 helper T cell cytokines and, **46:112**, 128–130
 human T cell activation and, **41:1**, 30
 accessory molecules, **41:14**
 cell surface molecules, **41:2**
 T cell antigen receptor, **41:2**, 5, 7
 humoral immune response and, **45:5–8**
 B cells, **45:10–12**
 cellular interactions *in vivo*, **45:79–81**, 84
 class II molecules, **45:18–20**
 helper T cells, **45:24**, 30, 31
 interleukins, **45:55**, 62–64, 71, 73
 physical interaction, **45:35**, 36, 41, 42, 45, 53
 processing, **45:8**, 9, 12, 13
 types, **45:9**, 10
 in vitro, **45:21**
 in vivo, **45:21–24**
 IgE biosynthesis and, **47:27**, 28, 33, 37, 39
 immunogeneity, **47:101–104**
 interaction with T cells, **47:92–94**
 modification, **58:438**
 SCID and, **49:384**, 387, 393
 sporozoite malaria vaccine and, **45:287**, 305, 307, 316
 surface antigens of human leukocytes and, **49:77–78**, 80, 82–83, 87
 synthetic T and B cell sites and, **45:197**, 201, 202
 antigens, **45:232**, 237
 globular protein antigens, **45:215**
 prediction, **45:252**
 viral antigens, **45:220**, 225
 T cell-dependent antibody responses
 B cells, **47:81**, 82, 84–86
 dendritic cells, **47:83**
 helper T cell subsets, **47:83**, 84
 T cell growth, **47:99–101**
 T cell receptor and, **45:107**
 accessory molecules, **45:108**, 109
 alloreactivity, **45:153**
 antigen processing, **45:118**, 120, 123, 124, 127
 epitopes, **45:143**, 144
 homogeneity, **45:129**
 MHC molecules, **45:116**
 peptides, **45:132**, 138
 T cell repertoire, **45:165**
 T cell subsets and
 H-2 alloantigen recognition, **41:80**, 83, 88, 92
 H-2 molecules in thymus, **41:100–102**, 105, 107
 H-2-restricted antigen recognition, **41:53**, 56, 58, 60–62, 68, 70, 74, 75
 T cell specificity, **41:110**, 111
 T cell triggering, **41:111**, 112
 T cell surface molecules in communication with, **58:112–139**

- TH response, **61**:361–364
 tissue distribution
 circulation, **47**:54–57
 delayed-type hypersensitivity, **47**:57
 life span, **47**:49, 50
 lymphoid organs, **47**:50–52
 nonlymphoid organs, **47**:52–54
 origin, **47**:47–49
 virus-induced immunosuppression and, **45**:366
- Antigen processing, **52**:11–78; **64**:1–3
 by B cell lymphomas, **39**:63–66
 by B cells, **39**:63–66, 87
 entry of proteins into class I processing pathway, **52**:13–14
Ir genes and, **38**:93–97, 152, 172, 188
 by macrophages, **39**:62–66
 MHC class I-associated peptides, generation, **64**:16–31
 preliminaries, **52**:11–13
 proteasome system, **64**:8–32
 activator protein, **64**:21–24
 cleavage properties, **64**:24–26
 inhibitors, **64**:11–12
 LMP and IFN- γ regulation, **64**:12–14
 molecular adaptation, **64**:19–21
 structural and catalytic features, **64**:8–11
 subunits X/Y and immunoproteasomes, **64**:14–16
 subunits Z and MECL1, **64**:16–18
 threonine residues, **64**:18–19
 proteolytic degradation, **39**:63–64, 66
 regulation of antigen processing, **52**:76–78
 TAP system, **64**:3–5, 30–31
 T cells, **43**:196–197
 translocation, **52**:35–42
 ubiquitin system, **64**:3, 5–8
 unfolding, **52**:22
 unfolding and proteolysis, **52**:22–78
 assembly of class I molecules, **52**:42–76
 control of class I export to cell surface, **52**:62–76
 proteolysis, **52**:22–35
 Antigen processing and presentation
 class Ib molecules, **52**:92–104
 characteristics of individual class Ib products, **52**:94–104
 general aspects of structure and function, **52**:92–94
- clinical relevance: immune, autoimmune and immunodeficiency states, **52**:106–108
 autoimmunity, **52**:107
 novel immunodeficiency states, **52**:107–108
 tumor immunology, **52**:104–106
 vaccines, **52**:106–107
 immune system evolution, **52**:1–11
 history of discovery of MHC restriction, **52**:4–6
 overview of T cell recognition and function, **52**:1–4
 properties of cell surface class I molecules, **52**:78–92
 association with exogenous β 2m, **52**:86–87
 interaction with exogenous peptides, **52**:78–86
 internalization, **52**:87–92
 evidence for internalization, **52**:87–89
 internalization signals, **52**:89–92
 T_{CD8+}, what it recognizes, **52**:104–111
 clinical relevance of antigen processing and presentation, **52**:104–108
- Antigen receptor
 genes, V(D)J joining, *see* V(D)J joining
 antigen receptor genes
 homology motif, signal transduction role, **57**:81–83
 loci, V–D–J recombination, **61**:304–306
 modulation, **59**:322–324
 structure, **43**:105–107
- Antigen receptor-dependent activation
 cytokines and B cell differentiation, **52**:204–206
 IL-2, **52**:204–205
 IL-4, **52**:205–206
 other stimulatory cytokines, **52**:206
 cytokines and B cell growth, **52**:195–200
 antigen-dependent activation, **52**:199
 polyclonal activators, **52**:195–199
 IL-2, **52**:195–196
 IL-4, **52**:196–198
 other stimulatory cytokines, **52**:198–199
 transforming growth-factor, **13**, **52**:199
- Antigen recognition, **43**:60, 223–225
 altered self model, **43**:202
 antigen dose-response curve, **43**:202

- Antigen recognition (*continued*)
 chemistry, **43:6**
 amino acid residues, **43:27**
 antigenicity in terms of recognition frequency, **43:29–32**
 reactive site stereochemistry, **43:32–39**
 systematic data base, **43:27–29**
 class II MHC molecules, **43:203–205**
 difference between T and B cells, **43:134**
 dual recognition, **43:202**
 hydrogen bonds and electrostatic forces, **43:83**
 hypothesis for origin, **43:224**
 mechanisms, **43:39**
 critical residues, **43:39–42**
 induced complementarity, **43:42–46**
 MHC-restricted, **43:202–203**
 T cell, **43:194**
 T cell receptor As, **43:176**
- Antigens, **43:117–119**
 ABC transporters, **65:55, 56–57, 59**
 adoptive T cell therapy of tumors and, **49:281–285, 334–335**
 expression of antitumor responses, **49:324, 327, 329–332**
 macrophages, **49:307–308, 311–312**
 mechanisms, **49:299–301, 304, 313, 318**
 principles, **49:287, 289–290, 295–296, 298**
 recognition, **49:318–324**
 and antigen recognition, **58:318–324**
 antinuclear antibodies and, **44:93, 94, 120**
 autoantibodies, **44:126, 127, 137, 138**
 autoantigen, **44:127–131**
 autoimmune response, **44:131–134, 136**
 dermatomyositis, **44:118–120**
 drug-induced autoimmunity, **44:109**
 scleroderma, **44:114, 121–125**
 Sjögren's syndrome, **44:111–113, 118**
 SLE, **44:95, 100–107**
 aPL antibodies and, **49:193–194, 256, 258–259**
 affinity purification, **49:228**
 clinical aspects, **49:200, 205**
 history, **49:198–199**
 LA antibodies, **49:241, 243**
 specificity, **49:234–239, 248–251**
Ascaris, inhalation challenge, **60:168–169**
 autogeneic, *see* Autoantigens
- autoimmune demyelinating disease and, **49:368–370, 372–374**
 autoimmune thyroiditis and, **46:263, 265–273, 317–319**
 cellular immune responses, **46:288, 289, 291, 293–295, 297, 298**
 experimental models, **46:273, 276, 281**
 genetic control, **46:284, 285**
 humoral responses, **46:298, 299, 302–305, 307**
 prevention, **46:310, 312, 314–316**
 autoreactive T cells and, **45:417, 418, 433**
 activation, **45:421–426**
 origin, **45:418–421**
 physiology, **45:427, 428, 430, 431**
 regulation, **45:431, 432**
 bacterial, IBD and, **42:290**
 based vaccination strategies, **58:435–437**
 B cell, **52:173–174, see also** B cell surface antigens
 Bgp 95, **52:173**
 CK226 antigen, **52:173**
 IgM-binding protein (Fc μ receptor), **52:173**
 B cell anergy, **59:321–322**
 B cell elimination, **59:306–308**
 B cell populations and
 assessment, **42:19–22**
 secondary, **42:75**
 binding by B cells, activation and, **39:71**
 BrMRBC, autoantibodies, secretion by
 CD5 B cells, **55:315–316**
 CD1-restricted, **59:78–87**
 CD2, role in T cell death, **58:226**
 CD3, expression during NK cell maturation, **55:346**
 CD4 and CD8, **44:265–270, 299–302**
 CD4 molecular biology, **44:293, 295**
 CD8 molecular biology, **44:270, 272, 282, 284**
 on IELs, **58:306–307**
 CD4 $^{-}$ CD8 $^{+}$, majority of $\gamma\delta$ T cells as, **58:102**
 CD4/CD8 coreceptor, role in negative selection, **58:171–172**
 CD5 B cell and, **47:117, 118, 120, 161**
 bone marrow transplantation, **47:129**
 genetic influence, **47:148**
 Ig gene expression, **47:150, 151, 155, 156, 158**

- lineage, **47:129**
 malignancies, **47:123, 124**
 marker for activation, **47:125, 126**
 ontogeny, **47:122**
 physiology, **47:131, 133–136, 138–143**
 primordial immune network, **47:159–161**
 surface, **47:143–147**
- CDw60, on CD4⁺ T cells, **58:108–109**
 changes in amino acid side chain, **43:24**
 clustered, *see* Clustered antigens
 competition, **43:208–211**
 complementarity-determining regions, **43:9–10**
 complementarity induced by, **43:20–23**
 complement receptor 1 and, **46:200**
 complement receptor 2 and, **46:205,**
 207–209, 211
 concentration, effect of Fc fragments and, **40:91**
 conformational changes, **43:123–124**
 conformational hierarchy levels, **43:74**
 differentiation in avian T cells, **50:88–89**
 electrostatic recognition with antibody, **43:12–13**
 environmental, influences on GALT, **40:207–209**
 eosinophil, recognition by antisera, **39:183–184**
 epithelial, **62:238–239**
 epitopes, *see* Epitopes
 equivalence, **43:68–72**
 evolution, **52:1–11**
 history of discovery of MHC restriction, **52:4–6**
 overview of T cell recognition and function, **52:1–4**
 T_{CD8⁺}, what it recognizes, **52:6–11**
 nature of antigens bound to class I molecules, **52:8–10**
 what T cell antigen receptor sees, **52:10–11**
 expression, hybrid resistance and genetics of, **41:397–401**
 hemopoietic cell grafts, **41:404–407**
 Hh-1, **41:401–403**
 marrow allograft reactivity, **41:408–410**
 transplanted cells, **41:403, 404**
- Fas, **57:129–131**
 flexibility, **43:118–119**
- follicular dendritic cell expression of, **51:257–261**
 foreign, CD5 B cell response, **55:318–319**
 fragment properties, **43:204**
 ganglioside, *see* Ganglioside antigens
 genetically engineered Abs and, **44:65,**
 66, 89
 antigen-combining sites, **44:84, 85**
 chimeric antibodies, **44:79, 80, 87, 88**
 cloning, **44:75**
 expression, **44:76–78**
 fusion proteins, **44:85**
 production, **44:66, 67**
 global structure, binding process, **43:22**
 glycoprotein, *see* Glycoprotein antigens
 heat-stable, for B and T cells, **53:235–236**
 helper T cell cytokines and, **46:111**
 cross-regulation of differentiation, **46:130–132**
 differential induction, **46:128–130**
 functions, **46:114, 115, 117–122**
 immune responses, **46:138**
 precursors of differentiation states, **46:122–127**
 secretion patterns, **46:112, 113**
 HIV infection and
 activation, **47:398, 400, 402**
 etiological agent, **47:378, 380, 383, 384**
 immune response, **47:408–410**
 immunopathogenic mechanism, **47:388,**
 390–392, 394, 397
 neuropsychiatric manifestations, **47:404**
 homing of plasma cell precursors and, **40:205**
 humoral immune response and, **45:2, 5–7**
 cell surface molecules, **45:41–46**
 cellular interactions *in vivo*, **45:79, 80,**
 82–85
 helper T cells, **45:24–33, 35**
 lymphokines, **45:46–54**
 physical interaction, **45:35–41**
 presentation, **45:9–12, 15–24**
 processing, **45:8, 9, 12–15, 17, 18**
- Ia, *see* Ia molecules
 IBD and
 Crohn's disease, **42:289**
 cytotoxic effector cells, **42:297**
 DR, **42:288, 292–294**
 lymphocytes, **42:290**
 somatic mutations and, **42:34**

- Antigens (*continued*)
 IFN- γ and, 62:80
 IgE biosynthesis and, 47:1
 antibody response, 47:3–6, 8, 10, 11
 antibody response suppression, 47:28,
 29, 31–39
 binding factors, 47:12, 14, 15, 21, 23–28
 Ig gene superfamily and, 44:1
 evolution, 44:5, 46:!!
 homology unit, 44:4
 MHC complex, 44:24–26, 28, 29
 nonimmune receptor members,
 44:29–31, 34–36, 38, 39, 41
 receptors, 44:9–12, 17, 18, 20, 23, 24
 Ig heavy-chain variable region genes and,
 49:26, 28, 34, 54
 V_H gene expression, 49:36, 39, 41–43,
 49, 51–52
 IL-1 and, 44:154, 189, 192
 biological effects, 44:166
 immunocompetent cells, 44:173–175,
 177
 structure, 44:156
 systemic effects, 44:169, 171
 immunosenescence and, 46:253
 lymphocyte activation, 46:222–224, 228,
 231, 235
 lymphocyte subsets, 46:237
 mucosal immunity, 46:253
 regulatory changes, 46:249–251
 stem cells, 46:247
 interaction energy, 43:23–24
 intestinal, IBD and, 42:290
Ir genes and
 blocking, 38:70, 71
 competitive inhibition, 38:108–110
 cytotoxic T cells, 38:161, 163–165,
 170–174
 expression
 in B cells, 38:57, 58
 in lymphoid tissue, 38:162, 190
 in T cells, 38:62, 190
 gene dosage, 38:80
 gene function, 38:92
 genetic control, 38:34–36
 H-Y, *see* H-Y antigen
 immunodominance, 38:181, 184, 185
 macrophages, 38:111–115
 MHC, 38:44, 46, 188
 MHC restriction, 38:64–67
 mutations, I region, 38:84–90
 pigeon cytochrome c, 38:77–79
 positive T cell selection, 38:130–134
 response to polypeptides, 38:46, 49
 suppression, 38:134, 135, 137–142, 144,
 148, 150, 152
 T cells, 38:40, 41, 43, 154, 160
 activation, 38:105–108
 proliferation, inhibition of, 38:68
 repertoire, 38:117, 118
 tolerance, 38:120, 123, 125–129
 leukocyte integrins and, 46:149, 153,
 161–163, 169
 lymphocyte homing and, 44:313, 314
 carbohydrate, 44:360–362, 371
 high endothelial venules, 44:321
 inflammation, 44:340, 341
 molecules, 44:342, 343, 345, 347–352
 regional specificity, 44:326, 327,
 330–338
 lymphocyte hybridomas and, 38:275, 276
 donor lymphocytes, 38:283
 EB nuclear, 38:277
 human-human, 38:297
 human-murine, 38:290
 maps of Ig-like loci and, 46:1, 7, 29, 31,
 32, 34
 MHC, *see* Major histocompatibility
 complex antigens
 microsomal, autoimmune thyroiditis and,
 46:266, 273, 298, 299
 multiple antigen peptides as, 60:137–139
 murine lupus models arid, 46:65, 99
 Ig germline, 46:72, 74, 76–78
 T cell antigen receptor, 46:81, 90, 91
 mutants, 43:117–118
 mycobacterial, recognition, 58:321–322
 neuroimmunology and, 39:299–300
 NK cell, *see* Natural killer cells, surface
 antigens
 NK cells and, 42:182, 183; 47:187–190
 adaptive immunity, 47:292, 293, 295
 alterations, 47:302
 antimicrobial activity, 47:288, 289
 CNS, 47:266
 congenital defects, 47:226
 differentiation, 47:232, 233
 effector mechanisms, 47:235, 237–240,
 247, 248
 genetic control, 47:222

- hematopoiesis, **47**:273, 275, 281, 282
 lymphokine production, **47**:264
 malignant expansion, **47**:227
 regulation, **47**:257, 259, 261
 reproduction, **47**:271, 272
 surface phenotype, **47**:200–212
 tissue distribution, **47**:221
 nonclustered, B cell, *see* Nonclustered antigens, B cell
 non-self, *see* Non-self antigens
 nonspecific killing, **60**:302
 parasitic, synthetic T and B cell sites and, **45**:228–230
 peptide, *see* Peptide antigens
 plasmacyte, **52**:174
 other antigens of plasmocytes, **52**:174
 PC-1, a plasma cell threonine-specific protein kinase, **52**:174
 polymeric, and IgE responses, **52**:289–290
 position-specific, leukocyte integrins and, **46**:153
 pulmonary immunity, **59**:387–388
 regulators of complement activation and, **45**:381, 389
 replaceability, amino acids, **43**:69–70
 residues, definition, **43**:23
 responses to, enhancement by V β selective elements, **50**:36–37
 schistosome
 Sm23, **60**:126–127
 Sm28-CST, **60**:127–128
 TPI, **60**:125–126
 SCID and, **49**:381–388
 graft-versus-host disease, **49**:398
 lack of stem cell engraftment, **49**:397
 posttransplant immunocompetence, **49**:400, 402
 stem cell engraftment, **49**:391
 tolerance, **49**:398–399
 transplantation, **49**:393–395
 self-, *see* Self-antigens
 sites, hydrophilicity, **43**:76
 specific activation, molecular basis, **54**:337–338
 specific killing, CTL-mediated, **60**:292–303
 sporozoite malaria vaccine and, **45**:283, 319–322
 CS proteins, **45**:299
 CS-specific T cells, **45**:307–311
 endemic areas, **45**:317
 human trials, **45**:314
 immunity, **45**:287, 291, 292
 interferon- γ , **45**:304–306
Plasmodium vivax, **45**:318
 staphylococcal protein A, antibodies to, **57**:229
 structural characteristics, antibody-binding sites, **43**:118
 superantigens, *see* Superantigens
 surface
 accessibility, **43**:119
 NK cell, *see* Natural killer cells, surface antigens
 surface, human leukocyte, **49**:75–76, 116–126
 adhesion molecules, **49**:82–89
 antigen-specific receptors, **49**:76–80
 expression, **49**:113, 116–117, 119–124
 membrane enzymes, **49**:111–114
 MHC glycoproteins, **49**:80–82
 receptors
 chemotactic, **49**:111
 for complement components, **49**:91–94
 for growth factors, **49**:99–101
 for IgGs, **49**:89–91
 for interleukins, **94**, **49**:98–99
 membrane proteins, **49**:93–97
 for neurotransmitters, **49**:101, 111
 T cell molecules, **49**:101–110
 transport proteins, **49**:114–116
 synthetic T and B cell sites and, **45**:195, 197, 260–264
 bacterial antigens, **45**:230–232
 candidate synthetic peptide vaccines, **45**:253–259
 globular protein antigens, **45**:203–216
 immunological considerations, **45**:197–202
 parasitic antigens, **45**:230
 peptides, **45**:203, 232–250
 prediction, **45**:251
 viral antigens, **45**:216–228
 tac, **59**:228
 tachyzoite, **65**:208
 T and B cell sites and, **45**:197, 263
 antigens, **45**:209, 212–214, 234–236
 immunology, **45**:198–200, 202
 prediction, **45**:252
 viral antigens, **45**:217, 218, 221, 225, 226

- Antigens (*continued*)
- T cell development and, **44**:207, 208, 255
 - antigen recognition, **44**:209, 210, 212–214
 - cellular selection within thymus, **44**:245–254
 - ontogeny, **44**:215, 219, 220, 221
 - recognition, **44**:209–214
 - thymocyte subpopulation, **44**:228, 239, 241–243
 - T cell-independent, examples, **40**:2–3
 - T cell receptor and, **38**:2
 - protein properties, **38**:3–5
 - specific binding, **38**:19, 20, 23
 - target recognition, **38**:22, 23
 - three-dimensional structures, **43**:117
 - thymus-dependent and -independent, **60**:272–277; **63**:43
 - transplantation, tumor-specific, characteristics, **57**:285
 - tumor, *see* Tumor antigens
 - tumor immunogenicity affected by low level of expression, **57**:303–305
 - uptake by B cells, **39**:59–62
 - comparison with macrophages, **39**:60–62
 - Ig role, **39**:60–62
 - valency, **59**:302–305, 308
 - virus-induced immunosuppression and HIV, **45**:351–353, 355–358, 360–362, 365–367
 - measles, **45**:339, 344, 346, 347
- Antigen specificity
- Ir* genes and
 - alloreactivity, **38**:121–123
 - antigen binding, **38**:101, 106, 107
 - antigen processing, **38**:96
 - B cell response, **38**:50
 - cytotoxic T cells, **38**:162, 164, 167, 171–174
 - expression, **38**:53, 55, 57, 60
 - gene function, **38**:37, 38, 93
 - genetic regulation, **38**:36, 37, 188
 - immunodominance, **38**:181, 183, 184
 - macrophages, **38**:111
 - MHC, **38**:46
 - “Schlepper” experiment, **38**:39, 40
 - T cells, **38**:41–43
 - interactions, **38**:153–158, 160
 - receptor, **38**:115–117
- selection, **38**:131
 - suppression, **38**:135, 136, 140, 149, 151, 152
 - lymphocytes hybridomas and
 - B cells **286**, **38**:306
 - human-human hybridomas, **38**:296
 - Antigen-specific recognition, T cell cultures for, **42**:165–166
 - Anti-H-2K^k transgenic model, **52**:311–313
 - Antiidiotypes, *see* Antibodies, anti-Id
 - Antiidiotypic immunoregulation, B cell repertoire expression and, **42**:35, 82
 - Antiidiotypic recognition, B cell repertoire expression and, **42**:51–62
 - Antiinflammatory cytokines
 - IL-6 effect, **54**:27–28
 - during placental/fetal development, **54**:32
 - role in embryonic development, **54**:31–32
 - rheumatoid arthritis and, **64**:291, 295–296
 - Antiinflammatory drugs
 - mode of action, **64**:172, 195–196
 - receptors, **52**:402–404
 - receptors for, **52**:402–404
 - Antiinflammatory and immunosuppressive molecules, NF-κB as target, **65**:128–132
 - antioxidants and, **65**:122–123
 - anti-TNF-α antibodies, **65**:123–124
 - cAMP and, **65**:124–125
 - cyclosporin A and FK506, **65**:120
 - deoxyspergualin and, **65**:125–126
 - gliotoxin and, **65**:126–127
 - glucocorticoids and, **65**:118–120
 - gold compounds and, **65**:123–124
 - rapamycin and, **65**:121
 - reactive oxygen intermediate-generating molecules and, **65**:122–123
 - rheumatoid arthritis drugs and, **65**:123–124
 - salicylates and, **65**:121–122
 - spergualin, **65**:125–126
 - steroids and, **65**:118–120
 - Antimicrobial activity, NK cells and, **47**:282–291
 - Antimitochondrial antibody, **49**:219–220, 233, 249–250
 - Antinuclear antibodies, **44**:93, 94; **49**:210, 212, 219
 - autoantibodies
 - biology, **44**:136–138
 - rare occurrence, **44**:125–127

- autoantigen
function, **44**:127, 128
molecular cloning, **44**:128–131
- autoimmune response, **44**:131, 132
antigen driven, **44**:132
epitopes, **44**:134–136
immunogens, **44**:132–134
- dermatomyositis, **44**:113, 118–120
- drug-induced autoimmunity, **44**:109–111
- localization, **44**:120
autoantibodies, **44**:125, 127
autoimmune response, **44**:133
scleroderma, **44**:121, 122, 125
SLE, **44**:106, 107
- mixed connective tissue disease, **44**:111, 112
- polymyositis, **44**:113, 118–120
- scleroderma, **44**:114, 120, 121
centromere antigens, **44**:121, 122
nuclear antigens, **44**:122–125
- Sjögren's syndrome, **44**:111–113, 115–118
- SLE, **44**:94–97
DNA, **44**:97, 98
histones, **44**:98–100
ku, **44**:104, 105
PCNA, **44**:106–109
ribosomal RNP, **44**:105, 106
RNP, **44**:100–102
subcellular particles, **44**:102–104
- Antioxidants, NF- κ B and, **65**:122–123
- Antipain, T cell receptor and, **45**:124
- Antiphospholipid antibodies, **49**:193–194, 257–259
- biochemistry, **49**:194–197
- clinical aspects
autoimmune disorders, **49**:207–210
detection methods, **49**:200–206
drug-induced antibodies, **49**:212–213
genetic studies, **49**:221–222
infectious disease, **49**:210–212
normal populations, **49**:206
syndromes, **49**:213–221
treatment, **49**:222–224
- history, **49**:198–199
- immunology
aCL antibodies, **49**:236–240
affinity purification, **49**:228–231
antimitochondrial antibodies, **49**:249–250
binding, **49**:250–252
isotype, **49**:224–228
- LA antibodies, **49**:240–248
reactivity, **49**:248–249
reagin, **49**:234–236
specificity, **49**:233–234
subsets, **49**:231–233
- pathogenic potential, **49**:252–257
- Antiphospholipid syndrome, **49**:218–221, 235, 243
- Antiproliferative effect, IFN- γ , **62**:76–77
- Antisense *myc*, blocking of negative signaling, **54**:413–415
- Antisense oligonucleotides, **58**:249–250
- Antisera, anti-MHr, polyclonal, **43**:29
- Antithrombin III, **49**:256; **66**:238
- Antithrombotic treatment, **49**:223, 258
- Anti-TNF- α therapy, **64**:333–334, *see also*
cA2 therapy
anti-CD4 with, **64**:332
combination therapy, **64**:331–333
Crohn's disease, **64**:307–309
drug modalities, **64**:330–331
HIV and AIDS, **64**:311
mode of action, **64**:311–323, 330
problems, **64**:323–330
anti-double-stranded cDNA antibodies, **64**:325
anti-Ig antibodies, **64**:323–325
malignancy risk, **64**:329–330
susceptibility to infection, **64**:325–329
- rheumatoid arthritis, **65**:124
animal models, **64**:298
cA2, **64**:299–305, 311, 312–315
clinical studies, **64**:299–307
- sepsis, **64**:310–311
- synergies with other treatments, **64**:331–333
ulcerative colitis, **64**:309–310
- Anti-transgenic models, with absent or partial
B cell tolerance, **52**:315–317
- α_1 -Antitrypsin, **62**:266
- Antitumor activity, NK cells and, **47**:295–300
- Antiviral activity, NK cells and, **47**:282–288
- Antiviral therapy
for FeLV, **52**:436–437
and vaccines, use to study, **52**:440–441
- AP-1, interaction with NF-IL6, **65**:18–19
- AP1-Ets interactions, **64**:81
- AP-1 family, **65**:29
- Apamin, synthetic T and B cell sites and, **45**:232

- Apigenin, action of, **65:123**
- aPL antibodies, *see* Antiphospholipid antibodies
- Aplasia, hybrid resistance and, **41:349, 372, 376, 388**
- Aplastic anemia, **41:236, 350**
graft rejection by patients, **40:383–385**
- Apo-1/Fas membrane-spanning proteins
induction of PCD, **58:135–136**
and negative selection, **58:172–173**
role in T cell death, **58:226**
- Apolipoprotein J, **61:249–251**
- Apo-MHr, monoclonal anti-MHr antibody
affinity, **43:61–62**
- Apoptosis, **52:157; 59:177; 61:9, 17, see also Cell death**
abnormal regulation, **58:214–215**
adoptive T cell therapy of tumors and, **49:312**
Bcl-2 effects, **58:137**
biochemical pathways, **50:60–61**
CD4⁺8⁻ cells, linomide effects, **58:263–265**
CD23 antigen and, **49:155, 171**
c-myc gene, **63:162**
control during intrathymic development, **58:168–174**
cytotoxic T cells, TCR-driven, **58:243**
different types, classification, **58:265–267**
due to cell damaging agents, **58:237–240**
eosinophil
aspect of senescence, **60:162**
in culture, **60:160**
TGF β -induced, **60:184**
- Fas antigen/APO-1 role, **53:241–242**
- Fas ligation, **61:37, 39**
- Fas-mediated, **57:129, 133–134, 140**
germinal center B cells, **60:282–283**
and growth arrest induced by anti-IgM, **54:410–411**
- HIV-induced, **58:236–237**
- HIV-specific CTLs, **65:306**
- hypothetical steps, **58:241**
in immune system, **50:63–70**
in B cells, **50:63–66**
clonal abortion, **50:64**
definitions, **50:55–56**
faulty recombination, **50:63–64**
growth arrest of WE III-231 B cell line, **50:64–65**
- mechanisms, **50:56–63**
apoptosis, morphology, **50:56–57**
biochemical pathways, **50:60–61**
DNA damage, **50:57–60**
phagocytosis, surface changes resulting in, **50:62–63**
somatic mutation and terminal differentiation, **50:65–66**
in T cells, **50:66–69**
CTL targets, death, **50:69**
deprived of growth factors, **50:68–69**
hybridomas, activation-induced death, **50:67**
interphase death of T and B cells and nonspecific damage, **50:69–70**
non-selected thymocyte, death, **50:67–68**
splenic B cells, **50:65**
thymus, negative selection, **50:66**
triggering and regulation, **50:71–73**
induction activation of death genes, **50:71–73**
release of death program, **50:75–76**
transduction, **50:73–75**
- inducing signals, **58:89**
- induction by Apo-1/Fas, **58:135–136**
- induction via TCR/CD3 complex, **58:216–219**
- inhibition of, **51:272–274**
- leukocyte, **58:280–281**
- as mechanism of T cell negative selection, **58:166–168**
- modulation, functional consequences, **58:272–280**
- morphology, **50:56–57**
- and nuclear DNA fragmentation, **58:212–213**
- and positive and negative selection of T cells, **58:89**
- Ras protein, **63:146–147**
- regulation, multiple pathways, **58:265–268**
- role in neonatal B cell tolerance, **54:403–404**
- role of reactive oxygen species, **58:250**
- and signal transduction pathways, **58:268–269**
- surface antigens of human leukocytes and, **49:117**
- susceptibility of murine lymphocytes, **58:234**

- target cell, Ca^{2+} -independent, **58:226**
- T cell
and IL-2, **50:166–167**
immunopharmacological manipulation *in vivo*, **58:254–255**
pharmacology, **58:211–296**
role of p53, **58:138**
and T cell persistence, **58:273–275**
thymocytes, **58:166–170**
TNF and, **66:114**
- Appendix, germinal centers, in rabbit,
rearrangements of VDJ genes, **56:204**
- Arachidonate
cyclooxygenase reaction, **62:178–183**, 198
metabolism
cachectin and, **42:219**
IBD and, **42:314**
- Arachidonic acid, **42:311**
corticosteroids and, **42:316**, 318
cyclooxygenase pathway, **39:147**
inhibition by EPA and docosahexaenoic acid, **39:160**, 164–165
formation from N-3 and N-6 fatty acids, **39:159–160**
hybrid resistance and, **41:360**
5-lipoxygenase pathway, **39:147–151**, 153, 160–161
metabolism
in eosinophils, human, **39:204**, 207–208
in neutrophils, human
dietary EPA and, **39:166–167**
nonesterified and esterified EPA and, **39:167–168**
- Arachidonic acid metabolite receptors, **52:398–402**
other arachidonic acid derivatives, **52:401–402**
prostaglandin binding sites, **52:399–401**
- Arachidonic acid metabolites
Fe fragments and, **40:107**
immune complexes and, **40:74**
receptors, **52:398–402**
suppressive properties of, **40:100–106**
- ARDS, *see* Adult respiratory distress syndrome
- Arg120, **62:175**
- Arg277, **62:174**
- Arginine, **42:97**, 141–144
- Arginine analogs, as NOS inhibitors, **60:325**
- Arsonate, genetically engineered antibody molecules and, **44:79**, 84
- Arsonate idiotypic system, A/J mouse
general considerations in, **42:95–96**, 157–158
- J558* V_{H} gene family and, **42:112–115**
deletion mapping analysis, **42:133–137**
evolution, **42:123**, 125–127
expressed and germline sequences, **42:127–128**
structural features, **42:115–117**
subfamilies, **42:117–123**, 124
- light chain
arginine, **42:141–144**
hybridomas, **42:139–141**
K locus, **42:138**
Vκ10, **42:144–148**
- methodology for, **42:96–99**
- molecular genetics, heavy chain cross-reactive idiotype-positive molecules, **42:109–110**
- DFL16.1* gene segment, **42:105–107**
- J_H* gene segment, **42:107–108**
kinetics, **42:108–109**
repetitive substitutions, **42:110–111**
serine, **42:111–112**
single V_{H} gene segment, **42:99–105**
- murine V_{H} complex and
deletion mapping analysis, **42:133–137**
idiotypic V_{H} map, **42:129–130**
inbred strains, **42:137**
preferential utilization, **42:131–133**
southern filter hybridization analysis, **42:130**
- serologic and structural analysis of Ab2 antibodies and, **42:149–150**
- CR-A*, **42:150–151**
- D_H* segments, **42:154–157**
heterogeneity, **42:151–154**
- V_{H} complex organization and, **42:128–129**
- Arthritis, **59:146**, 343; **61:46**, 252, 367, *see also* Juvenile Rheumatoid arthritis; Rheumatoid arthritis
- adjuvant-induced, **64:289**
antiadhesion therapy, **64:197**
antigen-induced, **64:289**
autoimmune, **62:99–100**, 107
autoimmune thyroiditis and, **46:284**, 310,

- Arthritis (continued)**
- collagen-induced, autoimmune thyroiditis
 - and, 46:312
 - and IL-1ra
 - in animal models, 54:210
 - in patients with arthritis, 54:210–211
 - in vitro* studies, 54:209–210
 - IL-8 role in, 55:143–144
 - induction, 64:289
 - inflammatory, IL-1ra in, 54:209–211
 - animal models, 54:210
 - patients with, 54:210–211
 - in vitro* studies, 54:209–210
 - joint erosion in, 64:287–288
 - lymphocyte migration into, 64:191
 - murine lupus models and, 46:61, 75, 79, 90
 - NO-induced, 60:343–344
 - pituitary hormones, 63:421–422
 - transgenic model, 64:298
- Arylsulfatase, in eosinophils, 39:218
- ASFV, see African swine fever virus**
- ASLC, 48:132**
- Asn410, 62:178**
- Asn580, 62:174**
- Asp³⁷**
- IDDM, 48:135–137
 - location, 48:139
- Asparagine**
- cytotoxic T cells and, 41:152; 153
 - genetically engineered antibody molecules and, 44:69, 84
- Aspergillosis, allergic bronchopulmonary, *see also* Allergic bronchopulmonary aspergillosis**
- Aspergillus infections, gliotoxin, 65:126–127**
- Aspirin**
- antiplatelet cardiovascular therapy, 62:200
 - aPL antibodies and, 49:223
 - colon cancer, 62:190
 - inhibition of NF- κ B activation at high doses, 58:19–20
 - PGHS, 62:187
- Aspirin-induced asthma, 51:353**
- Aspirin-sensitive asthma, 51:353–358**
- conclusion, 51:358
 - mechanism, 51:353–358
 - abnormal arachidonate metabolism at rest, 51:355
 - cellular source, 51:355–356
 - cyclooxygenase inhibition, 51:353
 - leukotrienes and AIA, 51:353–354
 - target organ sensitivity, 51:356–358
- Assembly deficient cells, 52:47–49**
- Association rate constants, IL-2 receptor and, 42:169–172**
- Asthma, 59:415–417; 62:283–285**
- acute response, and airway tone, 51:350–351
 - allergic
 - CD23 antigen and, 49:160
 - IL-4 storage by lung mast cells, 53:15
 - aspirin-sensitive, 51:353–358
 - bronchial, *see Bronchial asthma*, pathobiology of
 - CD23 antigen and, 49:160
 - eosinophil role, fiberoptic bronchoscopy studies, 60:218–223
 - human, *see also Bronchial asthma*
 - anti-Id antibody as mediator of, 39:279
 - bronchoconstrictive response to LTC₄ and LTD₄, 39:155
 - leukotrienes in neutrophils, dietary EPA and, 39:166–167
 - IL-5 role, 57:174–176
 - monkey model, 60:219
 - pathophysiology, *see Eicosanoids, and asthma pathophysiology*
 - related deaths, pathology, 60:217
 - release of leukotrienes in, 51:346
- Asthmatic mucosa, pathology of, 51:324–340**
- Aspirin, *see also* Eicosanoids, and eosinophils, 51:331–336**
- eosinophils and adhesion mechanisms, 51:333–336
 - epithelial cells and basement membrane, 51:324–327
 - lymphocytes, 51:339–340
 - macrophages and monocytes, 51:337–339
 - mast cells 327–331, mast cells and cytokines, 51:328–331
 - neutrophils, 51:336
- Astrocytes, 62:84**
- CD4 molecules and, 44:290
 - IL-1 and, 44:177
 - immunological mediator, 48:168–169
- Astrocytes, autoimmune thyroiditis and, 46:293**
- Ataxia telangiectasia, 58:67–69; 61:191, 315**
- T cell receptors and, 38:17
- ATF-2, Ets transcription factors and, 64:81**
- Atk, 59:169**

- Atomic solvation parameter, **43**:101
- ATP
- cytolysis and, **41**:277
 - as effector molecule in CTL cytotoxicity, **60**:301–302
 - effect on target cell killing, **60**:310
 - in IL-3-receptor interactions, **39**:39
 - TAP peptide binding and, **65**:84–85, 87–88
 - TAP peptide transport and, **65**:76–77
 - T cell activation and, **41**:16, 18
- ATPase, antigen-presenting cells and, **47**:53, 54
- ATP binding cassette, **52**:36
- Aurin tricarboxylic acid
- effect on thymic cellularity, **58**:262
 - prevention of T cell PCD, **58**:251
- Autoantibodies, **59**:330–331, 342
- AChR
 - cellular immunology, **42**:256–260
 - detection, **42**:255
 - anti-Id, human
 - in abnormal immune responses, **39**:274–275
 - in allergic diseases, **39**:275–276
 - to anti-TT antibodies, immune response and, **39**:269–274
 - natural occurrence, **39**:274 - anti-IgG, **38**:278
 - human-human hybridomas, **38**:297–299
 - monoclonal, **38**:298 - antinuclear, *see* Antinuclear antibodies
 - antithymocyte, CD5 B cell production, **55**:316–318
 - aPL antibodies and, **49**:212, 226, 257–258
 - pathogenic potential, **49**:252–253, 257
 - specificity, **49**:238, 248–249 - autoimmune thyroiditis and, **46**:298, 317
 - B cell formation and, **41**:227, 228, 231
 - CD5 B cell and, **47**:133–135
 - Ig gene expression, **47**:150, 155, 157, 158
 - physiology, **47**:130–136
 - primordial immune network, **47**:159 - CD23 antigen and, **49**:150
 - germline origin, **62**:19
 - helper T cell cytokines and, **46**:120
 - hybrid resistance and, **41**:379
 - Ig heavy-chain variable region genes and, **49**:2, 23, 33–34, 63
 - V_H gene expression, **49**:41–51

- in LEMS, **42**:252
 - murine lupus models and, **46**:99
 - Ig germline, **46**:65–67, 71–77
 - lupus strains, **46**:63, 64 - spontaneous autoimmune thyroiditis and, **47**:491, 492
 - altered thyroid function, **47**:456, 457, 459, 462
 - cellular immune reactions, **47**:449
 - clinical symptoms, **47**:439
 - humoral immune reactions, **47**:444, 446–448
 - potential effector mechanisms, **47**:466, 468, 469 - surface antigens of human leukocytes and, **49**:101
 - synthetic T and B cell sites and, **45**:247, 248
 - targeted adhesion molecules in pemphigus and pemphigoid, 8; **53**:309, 314
 - V_H segments, **62**:19
- Autoantigens, **51**:305–306
- antinuclear antibodies and, **44**:93, 94
 - autoantibodies, **44**:136
 - autoantigen, **44**:127–131
 - autoimmune response, **44**:131–134
 - scleroderma, **44**:125
 - SLE, **44**:100, 105
 - cDNA cloning
 - BP antigen, **53**:306–310
 - PF antigen, **53**:310–312
 - PV antigen, **53**:312–314 - hybridomas and, **38**:302
 - Ku, and V(D)J recombination and DNA repair, **58**:55–60
 - neural, dominant epitope, **48**:164–165
 - spontaneous autoimmune thyroiditis and, **47**:491, 492
 - cellular immune reactions, **47**:456, 459
 - disturbed immunoregulation, **47**:470
 - humoral immune reactions, **47**:445, 446, 448 - valency, **59**:302–305, 308
- Autochthonous tumor cells, adoptive T cell therapy of tumors and, **49**:284–285, 333
- Autocrine aspects, B cell growth and differentiation, **52**:219–221
- studies with B cell lines and tumors, **52**:219–220
 - studies with normal B cells, **52**:220–221

- Autocrine effects, IL-1, **54**:201–202
- Autocytotoxic cells, hybrid resistance and, **41**:394
- Autoimmune, immune, and
immunodeficiency states, antigen
processing and presentation, **52**:106–108
- autoimmunity, **52**:107
- novel immunodeficiency states, **52**:107–108
- tumor immunology, **52**:104–106
- vaccines, **52**:106–107
- Autoimmune attack, IL-2 and IL-2R at site
of, **50**:171–172
- Autoimmune cells, or normal human cells,
transfer to SCID mice, **50**:304–310
- Autoimmune demyelinating disease, **49**:357
multiple sclerosis
HLA antibodies, **49**:368–371
- TCR V region, antibodies to molecules,
49:372–374
- TCR V region, vaccination to,
49:374–375
- myelin basic protein, **49**:357–363
- TCR usage restriction, **49**:363
EAE in rat, **49**:366–368
- encephalitogenic C terminus, **49**:366
- encephalitogenic N terminus,
49:363–366
- Autoimmune diabetes, murine model
autoimmunity, **51**:294–306
autoantigens, **51**:305–306
diabetogenic T cells, repertoire of,
51:298–301
- macrophages and B cells, involvement
of, **51**:303–305
- suppressor mechanism controlling
development, **51**:301–303
- T cell-dependent autoimmunity,
51:294–296
- T cell subsets involved in insulitis and
overt diabetes, **51**:296–298
- clinical and histopathological
characteristics, **51**:289–290
- conclusions, **51**:310–311
- genetics, **51**:290–294
MHC-linked diabetogenic genes,
51:290–293
- non-MHC-linked genes, **51**:293–294
- IDDM, **51**:285–286
- intervention, **51**:306–310
- class II MHC molecule expression,
51:308–310
- cytokines, **51**:307
- immunopotentiators, **51**:306
- immunosuppressants, **51**:307
- virus infection, **51**:308
- NOD strain, development, **51**:287–289
- Autoimmune diseases, **61**:367
antiadhesion therapy, **64**:197
- anticolon antibodies and, **42**:291
- aPL antibodies and, **49**:193, 198, 258
clinical aspects, **49**:208–209, 216
- immunology, **49**:225, 235–236, 240,
250
- bcl-2* gene, **63**:163
- CD5 B cell and, **47**:148–150
- CD5⁺ B cells in, **53**:182–183
- CD28 costimulation, **62**:143
- cytokines and, **64**:290
- disrupted T cell therapy for, **42**:269
- elevated serum levels of soluble IL-2Ra in,
50:183–186
- Fas role, **57**:138, 140
- genetic defects in B lineage cells,
53:144–147
- HLA and, **66**:67, 76, 84–85, 93
- insulin-dependent diabetes mellitus,
66:91
- methimazole-induced insulin
autoimmune syndrome, **66**:87–88
- multiple sclerosis, **66**:88–91
- pemphigus vulgaris, **66**:87
- rheumatoid arthritis, **66**:85, 87
- IFN- γ and, **62**:78, 93, 98–101, 105, 107
and IL-1 α
- Alzheimer's disease, **54**:42–43
- cachexia, **54**:43
- cardiac myxoma, **54**:39–40
- Castleman's disease, **54**:40–41
- and IL-6, **54**:39–43
- mesangial proliferative
glomerulonephritis, **54**:41
- psoriasis, **54**:41
- rheumatoid arthritis, **54**:40
- IL-2, **63**:128, 132, 294, 297
- immunointervention, **66**:91–93
- induction, prevention by apoptosis-inhibitory drugs, **58**:278
- lymphocyte dependence, **64**:192
- MG and, **42**:260

- NF- κ B and, **65:117–118**
 pathogenesis, **64:331–332**
 pituitary hormones, **63:420, 425–426, 433**
 diabetes mellitus, **63:423**
 experimental allergic encephalitis,
 63:424
 multiple sclerosis, **63:425**
 rheumatoid arthritis, **63:421–422**
 SLE, **63:420–421**
 thyroiditis, **63:424**
 uveitis, **63:425**
 radioactive antigen suicide and, **42:268**
 SLE-like, **54:141–142**
 B/WF1 mouse, **53:144–147**
 spontaneous autoimmune thyroiditis, obese
 strain chicken, **47:433–435**
 altered thyroid function, **47:462, 464**
 cellular immune reactions, **47:449, 455**
 disturbed immunoregulation, **47:481**
 histopathology, **47:444**
 humoral immune reactions, **47:448**
 potential effector mechanisms, **47:468**
 streptococcal infection-associated,
 54:140–141
 SLE, **54:41–42**
 superantigens and, **65:207**
 TAP and, **65:92–94**
 treatment, **58:273–275**
 V_H polymorphisms, **62:9**
- Autoimmune manifestations
 lack of, in mice transgenic for human IL-2
 or IL-2R components, **50:192–193**
 after *in vivo* applications of recombinant
 IL-2, **50:189–191**
- Autoimmune response
 retinal antigen, **48:217**
 rheumatoid arthritis, **64:288**
- Autoimmune thyroiditis, **46:263–266,**
 317–319, *see also* Experimental
 autoimmune thyroiditis; Spontaneous
 autoimmune thyroiditis
 antigens, **46:266**
 thyroglobulin, **46:267–271**
 thyroid peroxidase 272, **46:273**
 cellular immune responses, **46:287**
 lymphoid cells, **46:287–291**
 target thyroid cells, **46:291–298**
 genetic control, **46:282**
 experimental induction, **46:284–287**
 SAT, **46:282–284**
- humoral responses, **46:298**
 antiidiotype autoantibodies, **46:305–309**
 autoantibodies, **46:298–306**
 prevention, **46:308, 310–316**
- Autoimmune uveitis
 experimental, **62:98–99**
 immune regulation, **48:215–216**
- Autoimmunity, **43:26; 52:107**
 antigen-presenting cells and, **47:104, 105**
 B cell repertoire expression and, **42:49**
 B cell tolerance, **59:342–344**
 CD23 antigen and, **49:150, 174–175**
 and diabetes, **51:294–306**
 autoantigens, **51:305–306**
 diabetogenic T cells, repertoire of,
 51:298–301
 macrophage and B cell involvement,
 51:303–305
 suppressor mechanism controlling
 development of diabetes,
 51:301–303
 T cell-dependent autoimmunity,
 51:294–296
 T cell subsets involved in insulitis and
 overt diabetes, **51:296–298**
 Ig heavy-chain variable region genes and,
 49:2–3, 61, 63
 polymorphism of V_H gene segments,
 49:21, 26
 regulation, **49:53–54**
 V_H gene expression, **49:39, 41, 43,**
 46–47, **50–51**
- and IL-2, **50:171–187**
 abnormalities in IL-2 expression and
 responsiveness on circulating
 lymphocytes, **50:186–187**
 elevated serum levels of soluble IL-2Ra
 in autoimmune disease, **50:183–186**
 at site of autoimmune attack,
 50:171–172
in vitro IL-2 production in autoimmune
 disease, **50:172–178**
 autoantibodies, **50:176**
 decrease of IL-2 producing cells,
 50:176
 low IL-2, **50:175 –176**
 signal transduction, **50:177–178**
 suppressor macrophages, **50:176**
 suppressor T cells, **50:176–177**

- Autoimmunity (*continued*)
in vivo IL-2 production in autoimmune disease, **50**:178–183
 immunological memory and, **53**:251–252
lpr mouse, **60**:308–309
Lyl B cells and, **42**:64
 murine, CD5 B cell effect on, **55**:313–315
 role of IL-2-induced cytokines in, **50**:210–213
 IFN- γ and, **50**:212–213
 IL-6 and, **50**:210–211
 TNF- α and, **50**:211–212
 X-linked agammaglobulinemia, **59**:146–147
- Autologous mixed lymphocyte responses, autoreactive T cells and, **45**:418, 425, 426
- Autologous plaque-forming B cell response, NK cells and, **47**:300, 301
- Autonomous model, interaction of C1q and IgG, **51**:19–20
- Autoradiography
 B cell formation and, **41**:187
 hybrid resistance and, **41**:356
- Autoreactive T cells, **45**:417, 418, 433
 activation
 specificity, **45**:425–427
 thymic stromal cells, **45**:421–425
 deletion, **58**:272–275
 origin, **45**:418, 419
 antigen-specific T cells, **45**:420
 antigen-stimulated precursors, **45**:419, 420
 immature precursors, **45**:421
 physiology
 autoimmune response, **45**:430, 431
 nonspecific helper function, **45**:427–429
 tumor resistance, **45**:428, 430
 regulation, **45**:431
 stimulators, **45**:432, 433
 suppressor T cells, **45**:431, 432
- Autoreactivity, CD1 proteins, **59**:76–78
- Autosomes, murine lupus models and, **46**:63, 94
- Autotolerance, and IL-2, **50**:165–171
 effect *in vivo*, **50**:165–166
 interference with clonal deletion, **50**:166–167
 nonspecific killing induced by IL-2, **50**:170–171
 the system, **50**:167–170
- Auxiliary signaling receptors, **51**:157–160
 CD2, **51**:157–159
 CD2S, **51**:159–160
- Avian leukosis virus
 B cell neoplasia induced by
 myb gene role, **56**:477–478
 myc gene role, **56**:472–473
 resistance determinants, **56**:470–471
 susceptibility determinants, **56**:469–471
 in chicken B cell DT40 induction, **57**:361–362
 pathogenesis of bursal lymphomas induced by, **56**:471–472
 mechanism, **56**:469
- Azathioprine, MG and, **42**:271, 272
- AZH, DR2 cellular subtype, **48**:135
- Azidodeoxythymidine, *see* AZT
- Azodisalicylate, **42**:318
- Azophenylarsonate, **42**:96–99, *see also* Arsonate idiotypic system, AJ mouse
- AZT, **63**:99
- B**
- B₁ receptors, **66**:239, 240
- B₂ receptors, **66**:240
- B7
 and CD28, costimulation, Td activation of B cells, **62**:132, 142–144, 148–157
 costimulatory effects on immunogenicity, **57**:305–307
 IFN- γ and, **62**:78
 interaction with CD28, **58**:133–134
- B7-1, **62**:132, 135–138, 156–157
- B7-1/B7-2 ligand family, **62**:135–138, 156
- B7-2, **62**:132, 135–138, 156–157
- B7/BB1, **52**:168–169
- B7 receptors, **59**:377
- B8
 DQw2, **48**:134
 DQw3, **48**:134
 DQw3.1, **48**:134
 DQw3.2, **48**:134
 DQw3.3, **48**:134
 DQw7, **48**:134
 DQw8, **48**:134
 DQw9, **48**:134
 HLA-DQ, **48**:134
 IDDM, **48**:137

- B15
 DQw2, **48**:134
 DQw3, **48**:134
 DQw3.1, **48**:134
 DQw3.2, **48**:134
 DQw3.3, **48**:134
 DQw7, **48**:134
 DQw8, **48**:134
 DQw9, **48**:134
 HLA-DQ, **48**:134
 IDDM, **48**:137
 1 B 236/myelin-associated glycoprotein, **48**:161–162
 Bacillus Calmette–Guerin, **42**:215
 Bacillus Calmette–Guerin factor, low molecular-weight, **52**:185
 Bacteria, *see also specific bacteria*
 adoptive T cell therapy of tumors and, **49**:308
 antibodies from combinatorial libraries, **57**:228
 antigen-presenting cells and, **47**:87
 aPL antibodies and, **49**:211, 248–249
 CD5 B cell and, **47**:160
 complement receptor 2 and, **46**:205
 cytotoxicity and, **41**:269, 270
 cytolytic proteins, **41**:317, 318
 membrane attack complex of complement, **41**:300
 genetically engineered antibody molecules and, **44**:66–70, 72
 helper T cell cytokines and, **46**:133
 HIV infection and, **47**:402
 hybrid resistance and, **41**:357, 378
 IL-1 and, **44**:163, 165–167, 194, 195
 infection
 IL-6 effects, **54**:36–37
 pneumonia, **59**:395–402
 responses in MHC class I-deficient mouse, **55**:412–413
 X-linked agammaglobulinemia, **59**:135, 136, 143–145, 150
 leukocyte integrins arid, **46**:151, 161
 lymphocyte homing and, **44**:354, 366
 maps of Ig-like loci and, **46**:1
 NK cells and, **47**:216, 224, 235, 263, 289
 regulators of complement activation and, **45**:388, 409
 SCID and, **49**:382
 spontaneous autoimmune thyroiditis and, **47**:462
 superantigens, **54**:113–115; **65**:14, 208–210
 Clostridium and *Pseudomonas* toxins, **54**:114–115
 Mycoplasma arthritides mitogen, **54**:113
 streptococcal M proteins, **54**:114–115
 surface antigens of human leukocytes and, **49**:81, 111, 115
 T cell receptor and, **45**:125
 toxic mitogens, **50**:15–16
 and murine V β gene segments, **50**:15
 retroviral sequences bearing no homology to, **50**:41
 Bacterial antigens
 humoral immune response and, **45**:82, 83
 IBD and, **42**:290
 synthetic T and B cell sites and, **45**:230–232, 245
 Bacterial disease, **60**:128–136
 multiple antigen peptides as immunogens, **60**:134–136
 BAF/B03 cell line, IL-2, **61**:156–157
 BAGE, **62**:229, 233
 BAL, *see* Bronchoalveolar lavage
 BALB/c mouse, splenectomized, **48**:208
 BALB invariant delta chain, **58**:304, 310, 312, 315, 320
 BALT, *see* Bronchus-associated lymphoid tissue
 BALU, *see* Bronchus-associated lymphoid unit
 Bare lymphocyte syndrome, **59**:246–247; **61**:327–338
 biochemical and genetic heterogeneity, **61**:329–331
 cell typing, **48**:117
 CIITA, **61**:331–334, 336–338
 RFX, **61**:334–336
 SCID and, **49**:387, 391
 Basement membrane
 B cell formation and, **41**:186
 lymphocyte homing and, **44**:316, 320
 Basophils, **48**:176, 177; **61**:15, 20, 360–361
 CD23 antigen and, **49**:165, 170
 characterization, **53**:2–3
 chemokine effects
 CC, **55**:116–117
 CXC, **55**:116
 differentiation, **60**:163–164

- Basophils (continued)**
- genetically engineered antibody molecules and, 44:68, 80
 - histamine release
 - major basic protein and, 39:191
 - somatostatin and, 39:305–306
 - IgE receptor/Fc ϵ RI numbers and binding constants, 52:386–387
 - IL-2 and, 44:164, 187
 - and IL-4 production in response to FcR crosslinkage, 53:13
 - IL-5-mediated activity, 57:156
 - IL-8 effects, 55:116
 - major basic protein, detection, 39:186–187
 - and mast cells, human, cell surface structure on
 - adhesion receptors and recognition molecules, 52:357–366
 - integrins, 52:358–361
 - other recognition molecules, 52:366
 - recognition molecules of Ig supergene family, 52:361–364
 - selectins and related recognition molecules, 52:364–366
 - cells, 52:333–335
 - cell surface typing with mAbs, 52:335–339
 - complementing binder sites, 52:375–377
 - conclusions, 52:404
 - IgE/Fc ϵ RI, 52:378–381
 - assembly, 52:381–382
 - cell surface structures functionally associated with Fc ϵ RI molecules, 52:391–396
 - control of synthesis and expression of Fc ϵ RI molecules, 52:384–386
 - functional characterization of receptor, 52:387–391
 - sequence and structural homologies, 52:383–384
 - topology of, 52:378
 - Ig receptors, 52:378–396
 - negative regulators of growth and differentiations of human basophils and mast cells, 52:355–357
 - interferons, 52:355–356
 - transforming growth factors, 52:356–357
 - receptors for activating peptides, 52:366–375
 - receptors for formyl-methionine peptides and related compounds, 52:371–372
 - receptors for IL-8 and related integrins, 52:367–371
 - receptors for substance P and other neuropeptides, 52:372–375
 - receptors for growth and differentiating factors, 52:339–355
 - hemopoietic growth factor receptor superfamily, HRS, 52:340–351
 - receptors for low-molecular-weight regulators and pharmacological compounds, 52:396–404
 - receptors for adenosine, 52:396–398
 - receptors for antiinflammatory drugs, 52:402–404
 - receptors for arachidonic acid metabolites, 52:398–402
 - RTK family, c-kit, and related oncogenes, 52:351–355
 - neuropeptide production, 39:316–317
 - phenotypic properties, 53:1–2
 - precursors, 53:2
 - properties, 53:1–2
 - role in inflammatory reactions, 53:2
 - secretions, 53:2
 - BB10 antibody, 43:306–307
 - bcl-2* oncogene, immunological memory and, 53:241
 - bcl-2* transgenic mice, pre-B cell differentiation into B cells, 53:143–144
 - B cell activation factor 1, humoral immune response and, 45:67
 - B cell activators, isotype specificity, 61:90–92
 - B cell antigen receptor, 52:136–138, 185–188; 55:282
 - anti-Ig in study of, 52:185–188
 - Branhamella catarrhalis* and others, 52:188
 - inhibitory effects of anti-Ig antibodies, 52:186–187
 - Staphylococcus aureus* Strain Cowan, 52:187–188
 - stimulatory effects, 52:185–186
 - cytoskeletal attachment, 55:262
 - Ig- α , tyrosine phosphorylation, 55:246–248
 - Ig- β , tyrosine phosphorylation, 55:246–248

- Ig complex, **52**:136–138
 ion movement, changes in, **55**:265–266
 mIg-induced tyrosine phosphorylation targets, **55**:234–249
 murine lupus models and, **46**:61, 65
 proteins associated with Ig complex, **52**:138
 protein tyrosine phosphorylation,
55:231–233
 signaling, **55**:273
 signaling pathway, CD40 expression,
61:37–38
 structure, **55**:226–230
 tyrosine kinase activation, **55**:248–256
 coreceptors, **55**:259–261
 signal initiation model, **55**:256–259
 tyrosine phosphorylation targets,
55:233–234
- B cell-associated nuclease, as truncation factor in V(D)J joining, **56**:80
- B cell differentiation factor, properties of, **40**:31–34
- B cell growth factor, **40**:35
 CD5 B cell and, **47**:119, 140, 141
 function of, **40**:14
 low-molecular-weight, CD23 antigen and,
49:160, 169
 NK cells and, **47**:265
 properties of, **40**:29–31
- B cell growth factor I, humoral immune response and, **45**:65, 66
- B cell growth factor II, **48**:78
 B cell formation and, **41**:235, 236
 T cell subsets and, **41**:76
- B cell growth factors
 CD23 antigen and, **49**:161, 171–172
 surface antigens of human leukocytes and,
49:99
- B cell immunopoiesis, **61**:47–54
- B cell lineage specific activator protein, **63**:228–229
- B cell lines
 AChR-specific, **42**:260–263
 Ars A response and, **42**:136
 cloned, and biochemistry of negative signaling, **52**:298–302
 deletion mapping of, **42**:133–135
 V_H gene segments and, **42**:131–132
- B cell lymphoma, **38**:248, 272; **54**:144–145;
61:121
A20-2J, **43**:196–197
- as animal models for tolerance, **54**:410–415
 antigen presentation to T cells, **39**:54
 cell line differences, **39**:86
 high activity of, **39**:66
 Ia structure and, **39**:76–78
 antigen processing
 chloroquine-inhibited, **39**:63–64
 comparison with macrophages, **39**:63–66
 heterogeneity of time for, **39**:64–65
 kinetics, **39**:63
 prefixation-inhibited, **39**:64–65
 anti-Id antibody therapy
 lack of response to, **39**:285
 acquired C1 esterase inhibitor deficiency and, **39**:288–291
 Id variants and, **39**:287–288
 remission, **39**:285–286
- CD5 B cell and
 Ig gene expression, **47**:150–152, 154
 malignancies, **47**:123–125
 chromosomal translocations and,
40:273–275
- cytotoxic T cells and, **41**:149
 follicular, human, **50**:73
 and GM-CSF, expression, **58**:435
 human B cell neoplasia and, **38**:248
 stimulation modes, **54**:253–254
 stimulatory factor 2, **54**:1–3
 superantigen expression, **54**:144–145
 T cell functions and, **39**:84
 T cell receptor β genes and, **40**:272
- B cell neoplasia
 human
 Burkitt lymphoma, *c-myc* oncogene in deregulation, **38**:254–257
 translocation, **38**:248–253
 variant chromosome translocation,
38:257, 259
 chromosomal translocation, **38**:246–248
c-myc activation, **38**:259–264
 t(14;18) chromosome translocation,
38:267, 271
 t(11;14) chromosome translocation,
38:264, 267
 retrovirus-induced, in bursa of fabricius, see
 Bursa of fabricius
 B cell peptide epitope vaccine, **43**:78

- B cell repertoire expression, *see also* Variable region, gene expression**
- B cell subpopulations and**
- adult primary B cell subpopulation, 42:67–68**
 - general considerations, 42:62–64, 79–83**
 - Ly1 B cells, 42:64–65**
 - neonatal, 42:65–67**
 - secondary B cell lineage, 42:68–79**
 - environment and, 42:34–35**
 - down-regulation, 42:42–43**
 - antiidiotypic recognition, 42:51–62**
 - tolerance to self-antigens, 42:43–51**
 - predominant clonotype expression, 42:39–42**
 - primary B cell repertoire diversification, 42:36–39**
 - sIg bone marrow precursor cells, 42:35–36**
 - neonatal sequential development of, 42:29**
- B cell response**
- Ir* gene influence on, 38:50–52**
 - to MMTV**
 - endogenous Mtv, 65:196–208**
 - neonatal, 65:171–175**
 - T–B cell interaction, 65:180–188**
 - T cell-dependent differentiation, 65:178–180**
 - T cell-independent activation, 65:168–171**
 - T cell suppression, 38:136** - B cells, 43:133**
 - AChR and, 42:257–258**
 - activation**
 - accessory molecules, 63:56–57**
 - early events, 63:52–53**
 - effector phase, 63:44**
 - extrafollicular, 52:221–224**
 - for proliferation and differentiation, 54:345–346**
 - and proliferation and differentiation, factors affecting, 40:29–50**
 - T-dependent, 63:43**
 - triggered by T_b membranes, 63:46** - activation biochemistry, 55:221–222, 282**
 - antigen contact, early cellular events following, 55:262–265**
 - antigen uptake, 55:264–265**
 - capping, 55:263–264**
 - contact-dependent T cell help, mechanism, 55:269–273**
 - cytokine receptor structure, 55:274**
 - cytoskeletal attachment, 55:262–264**
 - gene expression induction, 55:266–269**
 - IL-2 receptor, signaling by, 55:274–278**
 - IL-4 receptor, signaling by, 55:278–279**
 - IL-6 receptor, signaling by, 55:279–280**
 - ion movement changes, 55:265–266**
 - models, 55:222–225**
 - survival, 55:280–282**
 - adoptive T cell therapy of tumors and, 49:289, 306, 312–315, 330–332**
 - adult, abortive signaling and second signals, 54:407–410**
 - adult primary subpopulation, repertoire expression, 42:67–68**
 - aging and, 63:215–216**
 - allelic exclusion, 38:18**
 - anergy, 59:321–322**
 - antibody class switching, 61:84–85**
 - antibody isotype switching, 48:83–84**
 - antibody repertoire generation in rabbit GALT, 56:205–209**
 - Ig gene rearrangements, 56:189–192**
 - immune response ontogeny, 56:186–187, 209**
 - lymphopoiesis, 56:180–186**
 - somatic mutation, 56:203**
 - antibody response to β-galactosidase, 39:88–90**
 - helper T cells and, 39:88–90**
 - antigen presentation to T cells**
 - antigen specificity and, 39:57–59**
 - comparison with non-B cells, 39:85–86**
 - fractionation and, 39:68–69**
 - gamma radiation and, 39:67–72**
 - increase by activation, 39:66–72**
 - antigen binding and, 39:72–73**
 - Ia expression in membrane and, 39:74–76**
 - Ia structure and, 39:73–74, 76–78**
 - IL-1-like activity and, 39:78–79**
 - non-Ia molecules and, 39:78** - LPS and, 39:67, 69, 71**
 - NRGG as antigen, 39:55–56**
 - RAM1G as antigen, 39:52–53, 55–56**

- antigen-presenting cells and, **47:45**, 46, 47, 87
antigen presentation, **47:66–70**, 72–76, 78, 79
 antigen processing, **47:88–90**, 92–94
 APC-T cell binding, **47:95**, 97–99
 cell surface, **47:57–64**, 59, 61–64
 immunogenicity, **47:101**, 103, 104
 T cell-dependent antibody responses, **47:81–86**
 T cell growth, **47:99–101**
 tissue distribution, **47:47–57**, 48–51, 53–57
 antigen-reactive, specific killing, **60:305–306**
 antigen recognition, **43:223**
 antigens and, **38:2**
 antigen uptake, **39:59–62**
 comparison with macrophages, **39:60–62**
 Ig role, **39:60–62**
 aPL antibodies and, **49:249**
 autoocrine aspects of cell growth and differentiation, **52:220–221**
 autoimmune disease, **59:342–344**
 autoimmune thyroiditis and, **46:263**, 318
 antigens, **46:270**, 271
 cellular immune responses, **46:288**, 290, 295, 297
 experimental models, **46:275–277**, 280, 282
 humoral responses, **46:299–304**, 306, 307
 prevention, **46:314**, 315
 B1 cells, **59:285**; **63:207**
 B2 cells, **63:207**
 BALU, **59:386**
 at birth, **63:215**
 bone marrow
 neonatal, **42:67**
 repertoire expression in, **42:32–34**
 secondary B cells and, **42:76**
 in bone marrow, **63:208–215**
 CD1 distribution, **59:67–68**
 CD4 molecules and, **44:290**, 291
 CD5, **47:117–120**, 161; **55:297–299**, 330–333
 anti-BrMRC specificity, **55:315–316**
 anticarbohydrate specificity, **55:318**
 antiphosphorylcholine specificity, **55:318**
 antithymocyte autoantibody specificity, **55:316–318**
 in autoimmune disease, **53:182–183**
 autoreactivity, **55:328–329**
 bone marrow transplantation, **47:129**, 130
 clonal expansions, **55:322–323**
 in development, **55:327–328**
 in diseases, **55:327–328**
 expression on B cells, initial reports, **55:298**
 from fetal B progenitors, **55:300–304**
 generation, model for, **55:304–306**
 human population, homologous, **55:326–330**
 Ig gene expression, **47:150**, 152
 lineage, **47:128**, 129
 malignancies, **47:125**
 marker for activation, **47:125–128**
 mutation, **55:330**
 neoplasias, **55:322–323**
 ontogeny, **47:121**, 122
 peritoneal cavity localization, **55:299–300**
 physiology, **47:131**, 133, 134, 136–138
 primordial immune network, **47:160**
 response to foreign antigens, **55:318–319**
 specificity, biases in, **55:312–319**
 surface antigen, **47:143**, 147
 V genes
 human, **55:329–330**
 rearrangements, **55:323–324**
 usage, biases in, **55:319–326**
 CD5, murine
 autoimmune mouse, **55:313–315**
 cytokine production, **55:310–311**
 cytokine regulation, **55:310–311**
 differentiation, **55:309**
 expression on malignancies, **55:311–312**
 isotype expression, **55:309–310**
 properties, **55:306–312**
 strain variation, **55:308–309**
 surface phenotype, **55:306**, 308–309
 CD5⁺/CD5⁻ dichotomy, **53:178–183**
 CD23 antigen and, **49:149–151**, 176
 biochemical structure, **49:156**, 158
 biological activity, **49:167–173**, 169, 171
 cellular expression, **49:150–151**, 153
 expression in clinical conditions, **49:174–175**

- B cells (*continued*)
 expression regulation, **49**:158–160,
 158–161
FcεRII, **49**:162, 164–167, 165
 CD40 expression, **61**:2, 11–12, 32–35
 CD45 and, **66**:22–31, 36–37
 monoclonal antibody studies and,
66:12–15
 CD45-deficient, **66**:22–26
 cell-mediated killing and, **41**:297
 censoring, **59**:302–304, 308–309, 311–312,
 331–341
 chicken, repertoire formation in,
57:353–375
 development
 allelic exclusion, **57**:372–374
 antisilencer regulation, **57**:372–374
 bursa role, **57**:364–367
 embryonic, **57**:367–369, 374
 Ig sequence selection, **57**:369–372
 silencer regulation, **57**:372–374
 gene conversion
 avian leukosis virus induction of DT40
 cell line, **57**:361–362
 DT40, **57**:361–362
 Holliday junction, possible formation,
57:362–365
 hyperconversion mechanism,
57:358–361
 recombination models, **57**:362–365
 Ig gene organization
 amino acid composition, **57**:356, 358
 D elements, **57**:356–358
 heavy chain loci, **57**:354–356
 light chain loci, **57**:354–356
 chronic lymphocytic leukemia, **54**:50–51
 circulating, functions, **53**:201–203
 classes of, **53**:178–179
 clonal deletion, **58**:257
c-myc activation, **38**:261
 complement receptor 1 and, **46**:183, 193,
 200, 201
 complement receptor 2 and, **46**:203–211,
 213
Cμ-only transcription, **43**:237–238
 cytolysis and, **41**:273
 delineation of responsive subsets, **40**:4–5
 development, **59**:151–162; **63**:1–3,
 203–216
 bursa of Fabricius, **48**:42–44
- differential expression of lineage-related
 markers, **53**:125–127
 and gene rearrangements, **60**:38–39
 heavy chain and κ gene rearrangement
 and expression, **43**:236–237
μ heavy chains, **63**:1–4, 11–15, 16–29
 Ig, gene expression, **43**:236–240
 IL-2Rγ mutation and, **59**:252
 in MHC class I-deficient mouse,
55:393–394
 in mouse, **64**:221
 pituitary hormones, **63**:412–414
 preimmune repertoire, **59**:281–286
 subpopulation proliferation and
 differentiation, **53**:127–130
 surrogate light chain, **63**:2–4, 15–25,
 29–31
 terminally differentiated plasma cells,
43:239
 transduction, **63**:25–29
 XLA as defect, **60**:39–42
 development and differentiation, **60**:
 37–49
 differential expression of lineage related
 markers, **53**:125–127
 differentiation
 in CBA/N immunodeficient mice,
53:142–143
 cytokines and, **52**:204–218
 activated T cells, **52**:212–213
 antigen receptor-dependent activation,
52:204–206
 CD40-dependent activation,
52:206–212
 IL-2 and autoimmune system, **52**:215
 isotype switching, **52**:215–218
in vivo activated plasmablasts,
52:213–215
 heavy chain isotype expression, **61**:82–84
*HIGM*X-1 as defect, **60**:42–48
 isotype switching during, **60**:38–39
 surrogate L chain functions in,
53:136–143
 differentiation and expression, **54**:338–
 343
 directly responsive to T cell factors, **40**:19
 Epstein-Barr virus-transformed, **48**:125
 Fc receptors on, **40**:66–67, 68
 FOE receptors, **43**:298–300
 follicular B cells, **59**:283–285

- formation, **41**:181–185, 235–239
B cell precursors and, **41**:188, 189
cell size changes, **41**:202, 203
cell surface markers, **41**:189
distinctions, **41**:198, 199
functional assays, **41**:199–202
Ly-5 family of glycoproteins,
41:189–193
markers, **41**:193–195
phosphatidylinositol-linked lymphocyte
antigens, **41**:197, 198
technical considerations, **41**:196, 197
tumor cell lines, **41**:195, 196
bone marrow cultures and, **41**:208, 209
CBA/N mice, **41**:226
characteristics, **41**:216–218
inducible cell line, **41**:223
lymphocyte adhesion to stromal cells,
41:214–216
recent innovations, **41**:219, 220
structural organization, **41**:209–214
W/W anemic mice, **41**:224
genetically determined defects, **41**:223, 224
C3H/HeJ mice, **41**:230, 231
CBA/N mice, **41**:226, 227
cyclic neutropenia, **41**:228, 229
moth-eaten mice, **41**:231, 232
NOB mice, **41**:227–229
SCID mice, **41**:224–226
W/W anemic mice, **41**:224
Ig genes, **41**:203–205
inducible cell line, **41**:220–225
lymphohemopoietic tissue organization,
41:185–188
population dynamics, **41**:205–208
soluble mediators, **41**:232–235
frequency of, **42**:19–22
fusion, **38**:282, 283, 285, 286
Fyn expression, **63**:135
gene rearrangement, **38**:8, 12, 14
genesis, transcription factors, **63**:
216–250
genetically engineered antibodies, **38**:304,
305; **44**:88, 89
genetic defects leading to autoimmune
disease, **53**:144–147
genomic DNA structure or *mb-1* gene,
54:355
germinal center, maintenance of self-
tolerance, **60**:280–283
growth, cytokines and, **52**:195–204
activated T cells, **52**:203
antigen receptor dependent activation,
52:195–200
CD40-dependent activation, **52**:200–203
IL-6, **52**:203–204
growth and differentiation
autocrine aspects, **52**:219–221
B cell lines and tumors, **52**:219–220
normal B cells, **52**:220–221
complement role, **52**:218–219
cytokines involved, **52**:174–185
IL-2, **52**:177–178
IL-4, **52**:178–181
IL-6, **52**:182–184
IL-10, **52**:181–182
low molecular-weight BCGF-12 kDa,
52:185
TH1 and TH2 CD4+ helper T cells,
52:175–177
in vivo aspects, **52**:221–230
cytokines and B cells, a synthesis,
52:227–230
extrafollicular B cell activation,
52:221–224
follicular dendritic cells, **52**:225–227
germinal centers, development,
52:224–225
hapten-specific, T cell hybrid stimulation,
39:56–57
helper T cell cytokines and
cross-regulation of differentiation, **46**:131
differential induction, **46**:128–130
functions, **46**:114, 116–122
immune responses, **46**:133, 136
precursors of differentiation states,
46:125, 126
HIV infection and, **47**:390, 397
human
in Id recognition by T cells, **39**:259–260
interaction with rabbit anti-Id antibodies
binding to, **39**:264–266
suppression by, **39**:267–268
human counterpart of IgR-associated
proteins, **54**:356–357
human-human hybridomas, **38**:297
human-mouse heterohybridomas, **38**:301
human-mouse hybridomas, **38**:291
human T cell activation and, **41**:29
hybridization cell lines, **38**:279–282

- B cells (*continued*)
 hybrid resistance and
 antibodies, 41:378, 379
 antigen expression, 41:397, 408
 leukemia/lymphoma cells, 41:366
 lymphoid cells, 41:355, 357
 marrow engraftment, 41:387
 marrow microenvironment, 41:392
 NK cells, 41:366
 IBD and, 42:297
 6-MP therapy, 42:300
 idiotypic-specific inhibition by suppressor T cells, 40:136–143
 IFN- γ , 62:87
 IFN- γ and IL-4, 63:278
 Ig, enhancer preference, 43:242
 Ig class switching, 60:270–271
 Ig diversity, 48:52
 IgE-binding factor, 43:300–301
 IgE biosynthesis and
 antibody response, 47:5–12
 antibody response suppression, 47:28
 binding factors, 47:13, 19–23
 Ig gene superfamily and
 evolution, 44:47
 nonimmune receptor members, 44:37, 42
 receptors, 44:9, 10, 12, 21, 22, 24
 somatic diversification, 44:14, 16–18
 Ig heavy-chain variable region genes and, 49:2, 3, 21, 33, 53, 56
 regulation, 49:53–56
 tumors, 49:9, 11
 V_H gene expression, 49:35–38, 41, 43, 49–52
 IL-1 and, 44:189, 190, 194, 195
 biological effects, 44:164
 immunocompetent cells, 44:172, 173, 177, 178
 receptor, 44:182, 183, 185
 structure, 44:156
 IL-1-like activity, 39:78–79
 IL-5 effects
 regulation of development, 57:154–155
 signaling in X chromosome-linked immunodeficient mouse, 57:171–172
 IL-6 effects, 54:21
 IL-10 biological effects, 56:10–11
 immature, 59:285–286
 immune response results, 52:125
 immunosenescence and, 46:253
 lymphocyte activation, 46:231–234, 236
 mucosal immunity, 46:251, 253
 regulatory changes, 46:248, 251
 stem cells, 46:240, 242, 243, 246, 247
 induction of responses by, 40:12–16
 intermitotic life span of, 53:237–241
 Ir genes and, 38:50, 73, 153, 156, 188, 189
 antigen binding, 38:115
 expression, 38:53–55, 57, 58, 63
 helper T cells, 38:60
 Ia molecules, 38:68, 83, 84
 I region mutations, 38:85, 89
 MHC restriction, 38:64, 65
 T cells, 38:41, 42
 interactions, 38:155–158
 repertoire, 38:117, 121, 133, 134
 suppression, 38:142
 isotype-specific inhibition by suppressor T cells, 40:145–148
 leukocyte integrins and, 46:151, 154, 166, 171
 lineage expression of CD44, 54:293
 Ly1, 42:64–65
 splenic B cell and, 42:68
 subpopulation, 42:82–83
 in lymph follicles, 53:185–187
 lymphocyte homing and
 high endothelial venules, 44:315
 immune response, 44:330, 334, 335
 regional specificity, 44:325, 326, 329, 339
 lymphocytotoxic antibodies and, 42:291–292
 malignancies, 61:12
 Ig heavy-chain variable region genes and, 49:39–40
 mature, Ig gene rearrangements in, 40:270–276
 malignant, 61:38–39
 CD40 expression, 61:11–12, 38–39
 T cell-mediated suppression of, 40:144–145
 maps of Ig-like loci and, 46:1
 marginal zone, 59:284–285
 marginal zone B cells, 59:284–285
 maternally transmitted antigen, 38:313, 334
 maturity and tolerance, 52:290–294
 clonal abortion and clonal anergy, 52:291–294

- memory, *see* Memory B cells
 MG and, **42**:242–245
 in MHC class II-deficient mouse,
55:429–430
 MHC glycoproteins, **49**:81–82
 MLR stimulation, **39**:79–83, *see also*
 Mixed-lymphocyte reaction
 monoclonal antibodies, **38**:306
 murine lupus models and, **46**:64–66, 76, 98
 murine peritoneal, **54**:377–378
myc gene product in, **38**:254
 naïve
 adhesion molecules affecting,
53:248–249
 homing, **58**:350
 migration
 adhesion molecules and, **53**:248–249
 basic pattern, **53**:247–248
 migration pattern, **53**:247–248
 natural history, **52**:126–127
 neoplasia, and IL-6, **54**:33–36
 NK cells and, **47**:188, 199, 225, 232
 adaptive immunity, **47**:291
 alterations, **47**:300
 congenital defects, **47**:225, 226
 differentiation, **47**:230
 effector mechanisms, **47**:255, 264
 hematopoiesis, **47**:281
 malignant expansion, **47**:228
 surface phenotype, **47**:201, 209, 211
 PC-responsive, **42**:65
 in peripheral compartments, **53**:177–186
 in peritoneum, **53**:183–185
 in Peyer's patches, **40**:194, 195–197, 206,
 208
 phenotype of, **52**:129–174
 nonclustered antigens, **52**:164–174
 polyclonal activation, **52**:185–195
 activated T cells inducing resting B cells
 to proliferate and differentiate,
52:191–193
 anti-CD40, **52**:188–191
 antigen receptor, **52**:185–188
 pools, **53**:135–136
 pre-B cell receptor, **59**:155–156; **63**:20–21,
 25–26
 pre-B cells, **59**:286
 70Z/3 murine, crosslinking of ^{125}I -labeled
 ASD-Re595 LPS, **53**:278
 contacts with stromal cells, **53**:133–134
 defined, **63**:6
 differentiation into B cells
 in *bcl-2* transgenic mice, **53**:143–144
 in vitro, **53**:140–141
 in vivo, **53**:141–142
 exogenous recombination substrates,
43:237
 expression D μ protein, **63**:22–23
 long-term proliferating, **53**:131–136
 nonproliferating on stromal cells,
53:134–135
 RAG genes, **63**:20
 surrogate light chain, expression, **63**:6–7,
 11–12, 13
 precursors, **61**:11, 38
 B cell formation and, **41**:188, 189
 cell size changes, **41**:202, 203
 cell surface markers, **41**:189
 distinctions, **41**:198, 199
 functional assays, **41**:199–202
 Ly-5 family of glycoproteins,
41:189–193
 markers, **41**:193–195
 phosphatidylinositol-linked lymphocyte
 antigens, **41**:197, 198
 technical considerations, **41**:196, 197
 tumor cell lines, **41**:195, 196
 prenatal B cell genesis, **63**:207–208, 212
 pre-pre B cells, **43**:237
 pro-B cells, **59**:286, 298; **63**:23, 25
 long-term proliferating, **53**:131–136
 programmed cell death, **50**:63–66
 clonal abortion, **50**:64
 faulty recombination, **50**:63–64
 growth arrest of WE III-231 B cell line,
50:64–65
 somatic mutation and terminal
 differentiation, **50**:65–66
 proliferation, **63**:412–413
 CD40 ligands, **63**:58, 60–61
 CD40 signaling, **63**:53–58
 proliferation and antibody responses,
40:6–12
 pulmonary immunity, **59**:391–394
 receptor, **43**:133
 receptors, **49**:99, 101
 recognition, **43**:224
 regulatory elements of *mb-1* gene
 expression in restricted stage of B cell
 differentiation, **54**:355–356

- B cells (*continued*)
 role in V β selective element action, **50:31**
 SCID and, **49:381–383, 387–390, 393, 400–401**
 self-tolerance checkpoints, **59:279–281, 342–344**
 immune repertoire, **59:319–342**
 preimmune repertoire, **59:281–319**
 signaling, role of Btk, **60:5–6**
 signal transduction
 CD40, **61:32–35**
 MB-1, **54:351–355**
 sites of origin, **63:204–207**
 size, subsets and, **40:18**
 specific genes involved in IgR complex, **54:348–351**
 specific proteins in IgR complex identified by molecular cDNA cloning, **54:348–357**
 splenic, **50:65**
 spontaneous autoimmune thyroiditis and, **47:434, 467, 469, 481, 492**
 cellular immune reactions, **47:449, 450, 452, 455**
 histopathology, **47:440, 444**
 humoral immune reactions, **47:446–448**
 steady state production, **53:135–136**
 stimulatory factor 2, **54:1–3**
 subpopulations
 proliferation and differentiation capacities, **53:127–130**
 responsiveness of, **40:17–29**
 surface antigens, major, expression, **52:130**
 surface antigens of human leukocytes and, **49:76, 78, 92, 101, 115–117, 125**
 adhesion molecules, **49:86**
 antigen-specific receptors, **49:76–78**
 Ig's, **49:89, 91**
 switch recombination, **61:83**
 T cell activation and, **41:22**
 T cell-dependent responses, **62:131**
 B7-1/B7-2 ligand family, **62:135–138, 140–142**
 CD28-B7 costimulation, **62:142–144, 148–157**
 CD28/CTLA-4 receptor family, **62:132–135, 138–140**
 cellular events, **62:144–148**
 costimulation, **62:138–144, 148–154**
 model, **62:131–132**
- T cell development and, **44:209, 243, 252**
 –T cell interactions, **39:57–58, 83–85, 91–92; 63:44, 46, 71–72**
 epitopic specificity and, **39:87–90**
 and T cells, interphase death and nonspecific damage, **50:69–70**
 T cell subsets and, **41:39, 40**
 cell surface molecules, **41:42, 45, 46**
 H-2 alloantigen recognition, **41:89–91**
 H-2 molecules in thymus, **41:99, 100, 102**
 H-2-restricted recognition of antigen, **41:51, 54, 68–71, 76, 77**
 T cell specificity, **41:112**
 T-dependent and -independent, **42:63–65**
 V-DJ joining, **43:239**
 V(D)J joining of antigen receptor genes, rearrangement, **56:47, 49**
 pre-B cells, **56:47, 49**
 VH-only transcription, **43:236–238**
 Xid mutation, **59:317–318**
 B cell stimulating factor-1
 B cell formation and, **41:236**
 B cell precursors, **41:193, 194, 197**
 genetically determined defects, **41:231**
 inducible cell line, **41:222**
 soluble mediators, **41:234, 235**
 T cell subsets and, **41:66, 76, 77, 95**
 B cell surface antigens, **40:17–18**
 major, expression, **52:130**
 non-Ig, **52:138–174**
 non-Ig, plasmacyte, **52:174**
 other antigens of plasmacytes, **52:174**
 PC-1, a plasma cell threonine-specific protein kinase, **52:174**
 non-Ig clustered, **52:140–164**
 CD5, **52:140–144**
 CD9, **52:144**
 CD10 or CALLA, **52:144–145**
 CD20, **52:149–150**
 CD22, **52:150–151**
 CD23/FeeRII, **52:151–153**
 CD24, **52:153**
 CD37, **52:154–155**
 CD38, **52:155**
 CD40, **52:155–158**
 CD44, **52:158–159**
 CD45, **52:159–160**
 CD48, **52:160**
 CD72, **52:160**

- CD73, **52**:160–161
 CD74, **52**:161
 CD76, **52**:162
 CD77, **52**:162–163
 CD19, **52**:145–148
 CDw32/Fc γ RII, **52**:153–154
 CDw75, **52**:161–162
 complement receptors: CR2, CD21, and
 CD35, **52**:148–149
 other clustered antigens, **52**:163–164
 CD1c, **52**:163
 CD27, **52**:163
 CD39, **52**:163
 CD69, **52**:163
 CD78, **52**:163–164
 non-Ig nonclustered, **52**:164–174
 adhesion molecules, **52**:166–168
 activation of, **52**:168
 Ig superfamily, **52**:167
 integrin family, **52**:166
 selection family, **52**:167–168
 B7/BB1, **52**:168–169
 cytokine receptors, **52**:169–173
 IL-1, **52**:169–171
 IL-2, **52**:171–172
 IL-4, **52**:172
 IL-6, **52**:172
 other cytokines, **52**:173
 TNF- α , **52**:172–173
 transforming growth factor β , **52**:173
 HLA class II, **52**:164–166
 Bgp 95, **52**:173
 CK226 antigen, **52**:173
 IgM-binding protein (Fc μ receptor),
 52:173
 other antigens of B cells, **52**:173–174
 B cell tolerance, **52**:283–284, 325–326
 antigen polymerization and, **52**:287–290
 Felton's paralysis and other early results
 with polymers, **52**:287–288
 polymeric antigens and IgE responses,
 52:289–290
 size-fractionated polymers: Dintzis
 model, **52**:288–289
 biochemical analysis of signal transduction
 pathways, **54**:415–420
 direct analysis of signal transduction,
 54:418
 identifying Ig complex and associated
 second messengers, **54**:419–420
 use of inhibitors to block *in vitro*
 tolerance, **54**:415–418
 biochemical basis, **52**:294–302
 anti-Igs as surrogate antigens,
 52:295–297
 cloned lines and biochemistry of negative
 signaling, **52**:298–302
 constraints to study of antigenic
 signaling, **52**:294–295
 role of surface IgM and IgD,
 52:297–298
 early studies affecting antibody formation,
 52:284–287
 dissection of T cell vs B cell tolerance,
 52:285–287
 tolerance and antibody formation before
 T and B cell era, **52**:284–285
 model systems of analysis, **54**:394–415
 anti-IgM surrogate systems, **54**:402–410
 B cell lymphoma model, **54**:410–415
 hapten-specific, **54**:395–402
 history, **54**:394–395
 preliminary studies with transgenic
 spleen cells *in vitro*, **54**:414–415
 perspectives including antigen presentation
 by B cells, **52**:321–322
 rationale, **54**:393
 definition, **54**:394
 value of *in vitro* studies, **54**:394
 resolution of apparently conflicting models,
 52:322–325
 in secondary repertoire, **52**:317–321
 transgenic approaches, **52**:303–317
 advantages and disadvantages,
 52:303–305
 anti-DNA transgenic model, **52**:314–315
 anti-H-2Kk transgenic model,
 52:311–313
 anti-transgenic models with B cell
 tolerance absent or partial,
 52:315–317
 hen egg lysozyme model, clonal anergy
 vindicated, **52**:305–311
 B cell tumors, *see* B cell lymphoma
 BCG, **42**:215
 Bcl-2, **59**:344; **62**:43, 53; **63**:162–163
 downregulation, **58**:151
 effect on apoptosis, **58**:137
 highly expressed in pro-B cells, **58**:
 256–258

- bcl-2*, **64**:235, 251
 antiapoptotic, **58**:233
 apoptosis-regulatory, manipulation, **58**:253–258
 in CD40 antiapoptotic signal, **60**:46
 characterization, **58**:136–137
 defective, and T cell resistance to PCD, **58**:215
 inhibition of apoptosis, **58**:250
 overexpression, **58**:168, 268, 270
 regulation of apoptotic cell death in bursa of fabricius, **56**:479, 481
- bcl-3* gene, **65**:14
Bcl-x, **62**:43, 53
bcl-x, **63**:163
 regulation of apoptotic cell death in bursa of fabricius, **56**:479–480
- BCR, **61**:50, 53
bcr, tumor antigens derived from, **57**:296–297
 BCR triggering, CD40 signaling differs from, **61**:29
- Bence-Jones proteins, structural analyses, **43**:112
 Benign monoclonal gammopathy, **64**:229
 Beta-adrenergic mediator, **48**:162
 β bulges, **43**:111
 Beta cells, islet, *see* Islet beta cells
 β chain, **52**:379–380, *see also* T cell receptor excess of, **42**:176–177
 IL-2 receptor, **42**:167–168
 binding characteristics, **42**:168–172
 mast cell-specific receptor, gene cloning studies, **43**:287–289
 primary structure, **43**:220
 β sheet, **43**:102, 104, 111
 β strand, **43**:108–110
 BFP-STS, *see* Biological false positive serological test for syphilis
- Bg/II RFLP fragment., 4-kDa, celiac disease, **48**:145
 Bgp 95, **52**:173
bic, retrovirus-induced B cell neoplasia in bursa of fabricius, role in, **56**:477
- Bimolecular reactions, simultaneous, **58**:390
- Binding, via cell-cell and cell-matrix adhesion molecules, **51**:269–271
- Binding constants, basophils/mast cells, **52**:386–387
- Biochemical typing, HLA class II, **48**:114–116
 interisotypic association, **48**:116
 intraisotypic hybrid molecule, **48**:116
 trans-association, **48**:116
- Biological false positive serological test for syphilis, **49**:198–199, 210
 isotype, **49**:227–228
 specificity, **49**:234–236, 241
- Biological filtration, thymocytes, **59**:106–107
- Biological fluids, measurements of prostanooids in, **51**:349
- Biosynthetic radiolabeling methods, for IL-2, **42**:166–167
- Birbeck granules, **59**:61, 63
- Bis Q, **42**:249–250
- BIV, *see* Bovine immunodeficiency-like virus
- Black patient
 DQw9 Asp⁵⁷ positive, **48**:140
 DR7 haplotype, **48**:140
 DR9-DQ β , **48**:140
 DR9 haplotype, **48**:140
- Bladder infection, effector cell role, **60**:346
- Blast cells, localization to gut, **58**:348–349
- Blastocysts
 LIF regulatory role, **53**:34
 RAG-2-deficient blastocyst complimentation, **62**:31–41
- Bleomycin, pulmonary fibrosis, **59**:418–419
- BLIMP, **59**:342
- Blocking, *Ir* genes and, **38**:69–72
- Blood
 fetal, containing T cell precursors, **58**:140
 IL-2 receptor subunits, **59**:234–235
 peripheral, eosinophil counts, **60**:216–218
- Blood clotting, contact activation, *see* Contact activation
- Blood group antigens, **62**:220–221
- Blood stage, *P. falciparum*, antigen Pf322, **60**:120–121
- Blood vessels, IL-6 effects, **54**:29–30
- Bloom's syndrome, **61**:316
 and DNA ligase deficiency, **58**:65–66
- BLS, *see* Bare lymphocyte syndrome
- BLV, *see* Bovine leukemia virus
- B lymphoblasts
 EBV-transformed, **38**:277, 278
 IM-9, human, **48**:180
 NK cells and, **47**:292
- bmi-1, B cell genesis, **63**:220–221, 241–243

- Bmx* gene, **59**:184
Boc chemistry
 with benzyl, **60**:128–130
 and Finoc, HIV-1, **60**:134
in synthesis of multiple antigen peptides, **60**:106–107
- Bone
 metabolism
 IL-6 effects, **54**:28
 LIF role, **53**:36–37
 resorption, cachectin and, **42**:224
- Bone marrow
 adoptive T cell therapy of tumors and, **49**:288, 300
 antigen-presenting cells and, **47**:45, 47, 48, 81
 autoimmune thyroiditis and, **46**:301
 B cell, **42**:74
 neonatal, **42**:67
 repertoire expression in, **42**:32–34
 secondary B cells and, **42**:76
 B cell elimination, **59**:288–309, 318, 319
 B cell formation and, **41**:183, 184, 235–238
 B cell precursors, **41**:191, 193–199, 201, 202
 C3HeJ mice, **41**:231
 cyclic neutropenia, **41**:229, 230
 Ig genes, **41**:204, 205
 lymphohemopoietic tissue organization, **41**:186, 187
 moth-eaten mice, **41**:231
 NZB mice, **41**:227–229
 population dynamics, **41**:205–208
 SCID mice, **41**:225, 226
 soluble mediators, **41**:233, 234
 B cell genesis, **63**:208–215
 CD5 B cell and
 aging, **47**:122
 anatomic localization, **47**:120, 121
 lineage, **47**:128
 malignancies, **47**:124
 physiology, **47**:132, 142
 primordial immune network, **47**:161
 transplantation, **47**:129, 130
 CD23 antigen and, **49**:155
 cells from, transplantation in mouse, **54**:46–47
 complement receptor 1 and, **46**:203, 204
 cytotoxicity and, **41**:270
 dendritic cells, originating site, **59**:380
 derived B cells, clonal deletion, **58**:257
 derived cells, and deletion of thymocytes, **58**:170
 Fc ϵ RI $^+$ cells, lymphokine production by (murine), **53**:9–14
 forbidden α/β T cells, **53**:166
 growth hormone, **63**:386
 helper T cell cytokines and, **46**:115, 117
 HIV infection and, **47**:395–397
 human, CD44 expression, **54**:293–294
 hybrid resistance and, **41**:333, 334
 antibodies, **41**:376–378
 antigen expression, **41**:397–399, 401–410
 effector mechanisms, **41**:370
 leukemia/lymphoma cells, **41**:362, 365, 367, 369
 lymphoid cells, **41**:353, 355, 356
 marrow engraftment, **41**:384–388
 marrow microenvironment, **41**:391
 NK cells, **41**:372, 373, 375
 normal hemopoietic cells, **41**:335–351
 syngeneic stem cell functions, **41**:388, 390
 T cells, **41**:380–383
 *in vitro*assays, **41**:393, 395, 396
 IGF-I, **63**:392–393
 Ig heavy-chain variable region genes and, **49**:33, 36, 49
 immunosenescence and
 regulatory changes, **46**:248, 249, 251
 stem cells, **46**:240–247
 leukocyte integrins and, **46**:160, 166
 lymphocyte homing to, **64**:177
 mast cells derived from, **53**:3
 murine lupus models and, **46**:64, 80, 81
 NK cells and, **47**:188, 201
 adaptive immunity, **47**:292
 antimicrobial activity, **47**:289, 290
 antitumor activity, **47**:298
 differentiation, **47**:229–233
 effector mechanisms, **47**:234, 235, 263, 265
 hematopoiesis, **47**:272–283
 reproduction, **47**:270, 271
 tissue distribution, **47**:219, 221
 prolactin, **63**:383
 rabbit, rearrangements of VDJ genes, **56**:184
 spontaneous autoimmune thyroiditis and, **47**:446, 450

- Bone marrow (*continued*)
 surface antigens of human leukocytes and, **49:88, 117**
 T cells, characterization, **53:171–173**
 transplantation for severe combined immunodeficiency, *see* Severe combined immunodeficiency
 X-linked agammaglobulinemia, **59:158**
- Bone marrow cells
 IL-3-dependent lines from, **39:34–35**
 20 α SDH activity, **39:35**
 IL-3 effects
 colony formation and, **39:20–21**
 natural cytotoxicity and, **39:21**
 Thy-1 $^+$ induction and, **39:21–22, 24, 26–29**
 transplantation
 graft-versus-host disease and, **39:280–281**
 T cell role, **39:280–281**
- Bone marrow chimeras, *see* Chimeras
- Bone marrow cultures, B cell formation and, **41:208, 209**
 CBA/N mice, **41:226**
 characteristics, **41:216–218**
 inducible cell line, **41:223**
 lymphocyte adhesion to stromal cells, **41:214–216**
 recent innovations, **41:219, 220**
 structural organization, **41:209–214**
- Bone marrow grafting, MG and, **42:252**
- Bone marrow transplantation
 cyclophosphamide therapy and, **42:264**
 histocompatible, **49:392**
 MG and, **42:252**
- Borrelia burgdorferi*, **54:207**
- Bovine immunodeficiency-like virus (BIV), **52:432**
- Bovine leukemia virus, **52:432**
- Bovine serum albumin, methylated, aPL antibodies and, **49:234–235**
- Bowel disease, inflammatory, *see* Inflammatory bowel disease
- Box1/box2 motifs
 role in signal transduction, **60:11–12**
 sequence homology, **60:1**
- Boyden chamber assay, **60:175–176**
- BP-1 marker, B cell formation and, **41:193, 217, 225**
- Bpk, **59:169**
- Bradykinin
 formation, **66:225, 226, 233–239**
 function, **66:239**
 IgE biosynthesis and, **47:18, 25**
- Bradykinin receptors, **66:239–240**
- Brain
 autoreactive T cells and, **45:430**
 lesions, autoimmune demyelinating disease and, **49:368**
 virus-induced immunosuppression and
 HIV, **45:349, 351, 360, 365**
 measles, **45:346**
- Brain-derived neurotrophic factor, **61:5**
- Branhamella catarrhalis* and others, **52:188**
- Breast
 genetically engineered antibody molecules and, **44:87**
 Ig gene superfamily and, **44:39**
 lymphocyte homing and, **44:326, 327, 336, 338**
- Breast cancer, etiology, stem cell factor role, **55:78**
- Breast tumors, CD44 expression, **54:323**
- Brefeldin A, **64:105**
 effects on antigen processing and presentation, **52:42–44**
- Bromocriptine, **63:380, 409, 420, 422**
- Bromodeoxyuridine, T cell subsets and, **41:101, 102**
- Bronchial asthma, corticosteroid resistant, **51:358–361**
- Bronchial asthma, eosinophil role
 bronchial epithelium damage by major basic protein, **39:219–223**
 bronchoalveolar eosinophilia, **39:223**
 bronchospasm induction by LTC₄, **39:223**
 degranulation in tissues, **39:223–224**
 ECP in serum and, **39:224**
 mechanism of, **39:224–225**
- Bronchial asthma, pathobiology of
 aspirin-sensitive asthma, **51:353–358**
 conclusion, **51:358**
 mechanism, **51:353–358**
 corticosteroid resistant bronchial asthma, **51:358–361**
- eicosanoids and pathophysiology of asthma, **51:340–353**
 leukotrienes, **51:341–347**
 platelet-activating factor, **51:352**

- prostaglandins and thromboxane, **51**:347–352
epidemiologic background, **51**:323
pathology of asthmatic mucosa, **51**:324–340
 eosinophils, **51**:331–336
 neutrophils, **51**:336
 summary, **51**:361–363
Bronchial hyperresponsiveness, increase after allergen challenge, **60**:168–171
Bronchial lymph nodes, immunosenescence and, **46**:236, 237, 251
Bronchial mucosa, *in vitro* mucus secretion
 LTC₄ and LTD₄ effects, **39**:155
Bronchiolitis obliterans, **59**:421
Bronchoalveolar lavage
 from asthmatics, **60**:203–204
 eosinophil recruitment into, **60**:171–172
 obtained by fiberoptic bronchoscopy, **60**:218–223
 prechallenge, **60**:168–169
Bronchoalveolar lavage fluid, **59**:373, 374, 382
 assay *in vivo*, **60**:177–178
Bronchoconstrictive response
 indirect to LTB₄, **39**:156
 to LTC₄ and LTD₄
 in asthmatic subjects, **39**:155, 223
 in guinea pig, **39**:154–155
Bronchopulmonary aspergillosis, allergic, *see* Allergic bronchopulmonary aspergillosis
Bronchopulmonary mucomycosis, **62**:283
Bronchus-associated lymphoid tissue, **59**:382, 385–387
Bronchus-associated lymphoid unit, **59**:385–386
Brucella abortus, **54**:257, 397
 helper T cell cytokines and, **46**:127, 137
Bruton's agammaglobulinemia, *see* X-linked agammaglobulinemia
Bruton's disease, *see* X-linked agammaglobulinemia
Bruton's tyrosine kinase, *see* Btk
BSAP, **63**:228–229
Btk, **59**:136, 169–176, 178–196, 317; **62**:42, 47
 in B cell signaling, **60**:5–6
 role in XLA pathogenesis, **60**:40–42
 signal transduction, **59**:184–185, 317
 through CD40, **63**:51
 in XLA pathogenesis, **60**:40–42
BTK family, **59**:178–185
Btk gene, **59**:170–176, 184, 196
 B cell development, **63**:28–29
 mutations, **59**:185–192, 195, 317
Bullous pemphigoid
 antigen
 180-kDa, uniqueness of, **53**:305–310
 230-kDa, homology to desmoplakin I, **53**:305–310
 cDNA cloning, **53**:305–310
 hemidesmosome localization, **53**:297–300
 structural and amino acid homologies
 with desmoplakin I, **53**:307
clinical presentation, **53**:292–295
histology, **53**:292–295
immunopathology, **53**:292–295
α-Bungarotoxin, MG and, **42**:237–239
Burkitt's lymphoma, **63**:388
CD23 antigen and, **49**:166
chromosomal translocation, **38**:246–248, 271
 c-myc activation, **38**:259–264
 c-myc oncogene
 deregulation of, **38**:254–257
 translocation of, **38**:248–253, 271, 272
gene translocations and, **40**:272–273
human-human hybridomas, **38**:293
Ig heavy-chain variable region genes and, **49**:3
translocations, *c-myc* in, **50**:124–128
variant chromosome translocation, **38**:259, 264
Burnet's clonal selection theory, Ars A
 response and, **42**:109
Bursa of Fabricius
 B cell clonal expansion, **48**:54
 B cell development, **48**:42–44
 B cell formation and, **41**:181, 237
 development, **48**:42–44
 early role, **48**:49–51
 embryogenesis, **48**:47
 hybrid resistance and, **41**:372
 Ig gene rearrangement, **48**:49
 Ig+ precursor selection, **48**:49–51
 retrovirus-induced B cell neoplasia
 analytical views, **56**:480
 apoptotic cell death, role of, **56**:478–481
 avian leukosis virus-induced lymphoma,
 pathogenesis, **56**:471–472

Bursa of Fabricius (*continued*)
bic role, 56:477
myb role, 56:477–478
myc role, 56:472–478, 481
rel role, 56:477–478
research origins, 56:468–469
resistance and susceptibility
determinants, 56:469–471
somatic diversification, 48:54
structure, 56:467–468

Bursal anlage, 48:49

Bursal follicle
epithelial components, 48:42–43
lymphoid components, 48:42–43

Bursal lymphocyte
diversity, 48:52–54
V gene segment, 48:52–54

Bursal stem cell, 48:51–52
decline, 48:52
embryogenesis, 48:52
frequency, 48:51
germ-line gene segment, 48:51–52

Burst-forming units-erythroid, hybrid
resistance and, 41:396

t-Butyloxycarbonyl, *see* Boc chemistry

Bystander cells, cytotoxicity and, 41:280, 296

Bystander help, T cell subsets and, 41:76–77

C

C1, activated
actions of, 37:202–204
regulation and fate of
C1q receptor interaction, 37:206
dissociation of activated Clr2Cls2 and
C1, 37:204–206
inhibition of activated C1 by C1-In,
37:204

C1 activation, 51:21–25
C1 activation requirements and process,
37:195–200
C1 complexing agents and activators,
37:194–195
regulation of, 37:200–201

C1 activation unit
complexes of
C1, 37:185–194
Clr2Cls2, 37:183–185

proteins of
C1 inhibitor, 37:178–183
C1q, 37:155–165
Clr and Cls, 37:165–178

C1 INH deficiency, in hereditary
angioedema, 66:257–258, 259

C1-Inh gene, 61:205

C1 inhibitor, 61:204–206; 62:80

C1q
and antibody, interaction of, 51:9–21
C1q-associated IgG at the molecular
level, 51:17–21
C1q-IgG interaction at molecular level,
51:11–17
C1q-IgM interaction at the molecular
level, 51:16–17
complement C1q, 51:8
thermodynamics of interaction, 51:
8–11

complement components and, 38:
231–233

C2, 61:97, 206
complement components and, 38:203, 204,
206, 236
C4bp, 38:234
cDNA, 38:209–211
expression, 38:217, 218
Factor B, 38:207–209, 228
genes, cloning of, 38:212–214
MHC class III region, 38:227
restriction fragment length
polymorphism, 38:215, 216

CHITA, 61:331–334, 336, 338

C3
activation, 51:25–28
background, 37:217–218
definition of receptors, 37:221
functions of receptors erythrocyte,
37:242–251
history, 37:219–221
kidney podocyte, 37:260–261
lymphocyte, 37:257–260
mast cell, 37:261
monocyte macrophage, 37:255–257
neutrophil, 37:251–255
in complement activation pathways,
58:9–10

complement components and, 38:204, 206,
223, 233, 236
convertase, 38:205, 206, 213, 219, 222

- eosinophil-mediated parasite killing and, 39:210
generation of ligands for receptors, 37:221–222
activation of C3 by classical and alternative pathway, 37:223–225
covalent binding of C3b, 37:222–223
degradation of C4b and C3b by cleavage with factor 1, 37:226–230
factors controlling C3 activation, 37:225–226
structure and binding site characteristics of receptors
 type four, 37:240–242
 type one, 37:230–235
 type three, 37:238–240
 type two, 37:235–238
C3a, 61:255; 62:261, 262, 275, 286
C3b
 cleavage, 61:213
 complement receptors, 61:228–229
 complement system, 61:201–204
 degradation, 61:209–210
 properdin, 61:243–244
C3-binding protein
 complement receptor 1 and, 46:185, 189, 195, 196
 complement receptor 2 and, 46:202
C3/C5 convertase, 61:206–210
 complement receptors, 61:228–230
 decay-accelerating factor, 61:222–228
 factor I, 61:210–214
fluid-phase
 C4BP, 61:235–241
 factor H, 61:230–235
 nephritic factors, 61:245–247
 properdin, 61:207, 241–245
inhibition by membrane cofactor protein, 60:75–76
membrane cofactor protein, 61:214–221
C4, complement components and, 38:203, 204, 206, 219
allotypes
 functional activity, 38:221–223
 molecular basis, 38:225–227
cDNA cloning, 38:223
deficiency, 38:208
gene expression, 38:228–231
isotypes, 38:223–225
MHC
 class III region, 38:227–229
 linkage, 38:220, 221
 protein structure, 38:219, 220
C4, murine, molecular genetics, 38:223, 224, 227, 228
C4a, 61:255
C4 activation, 51:25–28
C4b
 cleavage, 61:213–214
 complement receptors, 61:228–229
 degradation, 61:209–210
C4b-binding protein, complement components and, 38:205, 211, 221, 222, 234–236
C4-binding protein, 61:203, 204, 235–236
complement receptor 1 and, 46:184, 185, 189, 195, 197
complement receptor 2 and, 46:202
function, 61:238–241
structure, 61:236–238
VCP, 61:241
C5, complement components and, 38:204, 205, 234
 convertases, 38:205, 206, 213, 222, 236
C5a, 61:255; 62:261, 269, 275, 286
 attracting myeloid cells, 58:380
 as neutrophil receptor, 39:96–97
C6, complement components and, 38:204, 205
C7, complement components and, 38:204, 205
C8, complement components and, 38:205
C9, complement components and, 38:205, 234
cA2 therapy, 64:312–315
clinical studies
 rheumatoid arthritis therapy, 64:299–305
 sepsis, 64:311
problems
 anti-double-stranded cDNA, 64:325
 anti-Ig antibodies, 64:323–325
 malignancy risk, 64:329–330
 susceptibility to infection, 64:325–329
rheumatoid arthritis
 anemia and, 64:321–322
 cardiovascular risk factors, 64:322
 cell trafficking, 64:317–319
 clinical studies, 64:299–305

- cA2 therapy (*continued*)
 cytokine expression, 64:315–317
 neovascularization, 64:320
 systemic effects
 cardiovascular risk, 64:322
 hemopoiesis, 64:321–322
 immune response, 64:322–323
- Cabergoline, 63:380
- Cachectin, 61:5; 62:96; 66:105, *see also*
 Tumor necrosis factor α
 acute-phase reactants from hepatocytes
 and, 42:225
 amino-terminal sequence of, 42:216, 217
 biosynthesis of
 cell of origin, 42:220–221
 control, 42:221
 kinetics, 42:221–222
 discovery of, 42:213–216
 effects of
 benefits, 42:226
 metabolism, 42:222–223
 neutrophils, 42:223
 other, 42:224–225
 receptor distribution, 42:222
 toxic, 42:218–220
 hybrid resistance and, 41:400
 IL-2 and, 44:185
 lymphotoxin and, 42:216–217, 224
 murine, 42:216–217
 and muscle cells, 42:223
 structure of, 42:216–217
- Cachectin receptor, 42:225–226
- Cachexia, 62:96
 helper T cell cytokines and, 46:114
 and IL-6, 54:43
 LIF role, 53:37–38
- Cadherin gene superfamily, pemphigus,
 foliaceous antigen in, 53:310–312
- Caenorhabditis elegans*, 50:72
- CAF, 65:282
 HIV-1 replication, 66:281, 282–283, 287
- Calcineurin, 62:42, 46–47; 63:166–167
 cyclosporin A, action of, 65:120
- Calcitonin gene-related peptide, 48:169
- Calcium, 48:228
 adoptive T cell therapy of tumors and,
 49:307
- aPL antibodies and, 49:197, 243, 245–246
- CD4 and CD8 molecules and, 44:268, 270,
 300
- CD5 B cell and, 47:127, 144, 145
- CD23 antigen and, 49:151, 163
- cell mediated killing and, 41:288–290, 299,
 292–297
- cell mediation, nature of, 41:271–273
- complement receptor 1 and, 46:199
- complement receptor 2 and, 46:207, 208
- cytolysis, 41:274, 278, 281, 284
- cytolytic proteins, 41:317, 318
- cytoplasmic, 43:293–298
 influx of extracellular, 43:295–298
 influx pathway, 43:296–297
 intracellular stores mobilization,
 43:293–295
- ionophore dose effect, 43:297
- lag times, 43:294
- phosphoinositide hydrolysis, 43:291
- extracellular, in CTL function, 60:296–298
- genetically engineered antibody molecules
 and, 44:85, 89
- HIV infection and, 47:387
- human T cell activation and, 41:31
 cell surface molecules, 41:2, 12
 gene regulation, 41:27–29
 receptor-mediated signal transduction,
 41:19–26
- synergy, phorbol esters and, 41:15–19
- IgE biosynthesis and, 47:26
- IL-2 and, 44:175, 185
- immunosenescence and, 46:222–226, 229,
 235, 253
- influx, and cNOS phosphorylation, 60:324,
 326–327
- ion channel, 48:285–287
- leukocyte integrins and, 46:156, 157, 159
- lymphocyte homing and
 carbohydrate, 44:358, 360–362
 molecules, 44:346–349
 regional specificity, 44:329
- mobilization, 48:260–262
- mobilization response
 in B lineage cells through MB-1 model,
 54:354
 role in signal transduction, 54:418–419
 to signals through IgR, 54:367–368
- in neutrophils
 protein kinase C and, 39:128–135
 release during activation
 amplification and, 39:132–135

- inositolphosphate generations and, **39:124–127**
 phospholipase G and, **39:125**
 receptors and, **39:126**
- NK cells and, **47:291**
 cytotoxicity, **47:249–251, 253, 254, 261**
 effector mechanisms, **47:241, 244, 245**
 lymphokines, **47:265, 266**
- pore-forming proteins and, **41:317, 318**
 role in apoptosis, **50:61**
- SCID and, **49:385–386**
 spontaneous autoimmune thyroiditis and, **47:471**
 surface antigens of human leukocytes and, **49:90, 114–115**
- T cell development and
 antigen recognition, **44:214**
 thymocyte subpopulation, **44:226, 227, 229, 236, 241, 243**
- T cell subsets and, **41:65**
- Calcium channels, **48:285–287**
 LEMS and, **42:234–235**
- Calcium ionophore
 and factor-dependent mast cell lymphokine production, **53:6**
 T cell activation, **48:258–260**
 and transformed mast cell lymphokine production, **53:5**
- Calcium ionophore A23187
 5-lipoxygenase activation, **39:154**
 eosinophil degranulation and, **39:207–208**
- Calf serum
 B cell formation and, **41:209, 216, 218, 219, 222, 223**
 hybrid resistance and, **41:395**
- CALLA, *see* CD10
- Calmodulin, NK cells and, **47:249**
- Calnexin, **64:107–108**
 calnexin-negative cells, phenotype, **64:111–112**
 class I molecules, interaction site, **64:110–111**
 endoplasmic reticulum retention of assembly intermediates of class I molecules, **64:109–110**
 H-chain assembly, **64:108–109**
 MHC class I assembly and, **65:89, 90**
- 3' Cα enhancer, germline switch transcripts and class switching, **61:121–123**
- Calreticulin, **65:90, 91**
 MHC class I peptide loading, **64:128**
- CAMPATH-1H, **64:323**
- cAMP response element modulator τ, **61:189**
- cAMP response elements, **62:194**
- Cancer, **59:147–148**
 adoptive T cell therapy of tumors and, **49:283–285, 333**
 anemia following IL-6 administration, **64:321–322**
 antitumor activity of IL-6, **54:48–50**
 IFN-γ and, **62:94–96**
 IL-1ra in, **54:213–214**
 immunotherapy, potential of antigens, **58:437**
 malignant B cells, CD40 expression, **61:11–12, 38–39**
 maps of Ig-like loci and, **46:1**
 NK cells and, **47:297–300**
 p53 malignant transformation, **62:226–227**
 PGSH-2, **62:190, 201**
 rheumatoid arthritis and, **64:329–330**
 TAP dysfunction, **65:95**
 therapy, **54:49–50**
 gene transfer as, **58:417–454**
 lymphoid tumors, **58:282**
- TIL therapy, **64:197**
 treatment with chemoimmunoconjugates, *see* Chemoimmunoconjugates
- Cancer antigens, **62:240**
- Candida albicans*, cachectin and, **42:226**
- Capping
 B cell antigen receptor, **55:263–264**
 and resynthesis of receptors, **54:402–403**
- Caprine arthritis encephalitis virus (CAEV), virus-induced immunosuppression and, **45:349, 351, 354, 365**
- Carbodiimides, in crosslinkage procedure for chemoimmunoconjugates in cancer treatment, **56:316–318**
- Carbohydrate antigens, **62:219–221**
- Carbohydrate component, mobility, **48:26–32**
- Carbohydrates
 aPL antibodies and, **49:239**
 autoimmune thyroiditis and, **46:271**
 B cell formation and, **41:197**
 CD8 molecules and, **44:271**
 CD23 antigen and, **49:155–156, 163–165,** 176

- Carbohydrates (*continued*)**
- complement receptor 2 and, **46:204**
 - cytotoxic T cells and, **41:138, 165, 170, 152–154**
 - epitopes, tumor-associated alterations, **58:420**
 - genetically engineered antibody molecules and, **44:68, 69, 84**
 - IgE biosynthesis and, **47:15–17**
 - IL-2 and, **44:194**
 - leukocyte integrins and, **46:173, 174**
 - lymphocyte homing and, **44:314**
 - cell adhesion, **44:353–355**
 - high endothelial venules, **44:315**
 - homing receptors, **44:369, 370**
 - inhibition, **44:355–357**
 - ligands, **44:363–371**
 - molecules, **44:342**
 - PPME-binding receptor, **44:357–363**
 - NK cells and, **47:207, 241**
 - as selectin ligands, **58:358**
 - spontaneous autoimmune thyroiditis and, **47:460**
 - sulfated complex, in eosinophils, **39:204**
 - surface antigens of human leukocytes and adhesion molecules, **49:88–89**
 - Igs, **49:91**
 - membrane enzymes, **49:113–114**
 - receptors, **49:94, 98**
- Carboxyfluorescein, cell-mediated killing and, 41:290**
- Carboxypeptidase-N, 61:204**
- Carboxyterminal signal sequence, in precursor peptide to be GPI anchored, 60:62–63**
- Carcinoembryonic antigen, 62:221–222**
- genetically engineered antibody molecules and, **44:67**
 - Ig gene superfamily and, **44:39, 40, 43–45**
 - mAbs in identification, **56:305**
 - monoclonal antibody recognition, **56:305**
- Carcinogenesis, inflammation-induced, NO role, 60:345–346**
- Carcinomas**
- CD40 antigen, **61:14, 39**
 - hybrid resistance and, **41:358, 363, 369**
- Cardiolipin, *see also* Anticardiolipin antibodies**
- aPL antibodies and, **49:193, 258–259**
 - affinity purification, **49:229–231**
 - antibody subsets, **49:232**
 - antimitochondrial antibodies, **49:250**
 - binding to cell membrane, **49:251–252**
 - biochemistry, **49:194, 197**
 - clinical aspects, **49:200, 204–205**
 - history, **49:198**
 - LA antibodies, **49:240, 242–247**
 - pathogenic potential, **49:255**
 - reactivity, **49:248–249**
 - specificity, **49:234–240**
- Ig heavy-chain variable region genes and, 49:42, 46**
- Cardiopulmonary bypass, therapeutic regulation of complement system in, 56:279–280**
- Cardiotrophin-1, *see* CT-1**
- Cardiovascular system, chemokine role in, 55:146**
- Caries, dental, subunit vaccines, 60:135–136**
- Carp IgM antibody, 48:23–24**
- Carrageenan, hybrid resistance and, 41:366, 371–373, 395**
- Cartilage matrix protein, leukocyte integrins and, 46:158**
- Casein, cell-mediated killing and, 41:297**
- β -Casein gene, 65:20**
- Castleman's disease, and IL-6, 54:40–41**
- Cat, mesenteric microvascular models, 60:329–330**
- Catabolic cascades, inhibition at effector level, 58:247–253**
- Catabolic effects, IL-2 and, 44:168**
- Catabolin, IL-2 and, 44:154, 168**
- Catabolism, antigen processing and, 38:95**
- Catalysis**
- antinuclear antibodies and, **44:127, 134–136**
 - CD4 molecules, **44:265–270, 297–303**
 - cDNA structure, **44:285**
 - expression, **44:288–291**
 - function, **44:291–295**
 - gene structure, **44:287, 288**
 - HIV-1 receptor, **44:295–297**
 - Ig gene superfamily and, **44:31–33, 36, 41, 42**
 - IL-2 and, **44:189**
 - lymphocyte homing and, **44:325, 340**
 - molecular biology, **44:284**
 - polypeptide structure, **44:285–287**
- CD8 molecules, 44:265–270, 297–303**
- Ig gene superfamily and, 44:31–33, 35, 41, 47**

- lymphocyte homing and, **44**:325, 350
 molecular biology
 CD8 β , **44**:276–278
 cDNA structure, **44**:272–274
 expression, **44**:279, 280
 function, **44**:280–284
 gene organization, **44**:274–276
 polypeptide structure, **44**:270–272
 genetically engineered antibody molecules
 and, **44**:85, 86
- Cataract, immune-mediated, **48**:219–221
- Cathepsin G, **62**:266
- Cathepsins, **64**:2
- Cations
 B cell formation and, **41**:214, 215
 cytosis and, **41**:280, 311
- Caucasian patient
 DQw9 Asp⁵⁷ positive, **48**:140
 DR7 haplotype, **48**:140
 DR9-DQ β , **48**:140
 DR9 haplotype, **48**:140
- CBA/N mice, B cell differentiation in, **53**:142–143
- c-bcl-2* oncogene, IL-2 signal transduction, **63**:162–164
- CBF family, **65**:24
- CBF α proteins, **65**:24
- CD1, **59**:10–11, 90–91
 biochemistry, **59**:44–45
 CD23 antigen and, **49**:154, 156
 cellular expression, **59**:54–55
 family, **59**:2, 17–37, 90–91
 biochemical studies, **59**:44–54
 cellular expression, **59**:10–11, 54–74
 genetics, **59**:2–10
 immunology, **59**:74–90
 molecular biology, **59**:2–10
 serology, **59**:37–44
 structure, **59**:10–37
 gene, **59**:3–10
 gene cloning, **59**:3–4
 group 1, **59**:22–25
 cellular expression, **59**:55–69
 by B cells, **59**:67–68
 by dendritic cells, **59**:60–64
 by macrophages, **59**:66–67
 by monocytes, **59**:64–66
 by neoplastic cells, **59**:68–69
 in thymus, **59**:55–60
 immunological function, **59**:74–87
 group 2, **59**:22–25
 cellular expression, **59**:56, 69–74
 function, **59**:88–89
 human leukocytes and, **49**:78, 80–83
 serology, **59**:37–44
 structural features, **59**:10–37
 CD1a, **59**:39, 40, 42, 45, 46, 48, 50–53,
 82–85
 cellular expression, **59**:55–69
 CD1b, **59**:39, 41–43, 45, 48, 50–53
 cellular expression, **59**:55–69
 mycolic acid restricted by, **59**:82–85
 CD1c, **52**:163; **59**:39, 41, 42, 45, 46, 50–54
 cellular expression, **59**:55–69
 CD1d, **59**:38, 39, 40, 49–52
 cellular expression, **59**:69–74
 tissue distribution, **59**:56
- CD1E, gene, **59**:22
- CD2, **48**:235–237; **51**:157–159; **64**:69
 Abs, **48**:235
 B cell genesis, **63**:239
 CD45 and, **66**:7–8, 33, 35–36
 expression, **48**:235
 human leukocytes and, **49**:87, 117
 structure, **48**:235
- CD2S, **51**:159–160
- CD3, **43**:135; **48**:162
 CD45 and, **66**:9–12, 34
 CD59 as ligand, **61**:254
 complex, human leukocytes and, **49**:76–79,
 81, 87, 90
 complex with TCR, *see* CD3/TCR complex
 ϵ , δ , γ , and ζ/η invariant chains, in T cell
 development and selection,
 58:127–128
 ϵ subunit, **64**:69
 expression, **43**:166, 169
 SCID and, **49**:384, 386–387, 396–397
 T cell lines expressing, **58**:122–124
- CD3 complex, components, **48**:229, 230
- CD3–Jurkat mutant, anti-CD2, mAb,
 48:236
- CD3/TCR complex
 ligation, **58**:248
 mediated peripheral T cell deletion,
 58:219–224
 monoclonal antibodies to, **58**:300
 second messenger triggered by,
 58:230–232

- CD4, **48**:242–245, 304–306; **59**:119, 382; **63**:84, 403
 as accessory molecule, **53**:60
 adoptive T cell therapy of tumors and, **49**:333–335
 antigen recognition, **49**:318, 320–321, 324
 expression of antitumor responses, **49**:324–330, 332
 mechanisms, **49**:299–306
 principles, **49**:289–291
 autoimmune demyelinating disease and, **49**:364, 366–367, 374
 and CD8, **52**:363
 and CD8, coreceptors, **58**:129–131
 CD23 antigen and, **49**:152
 coreceptor effects on selechon, **53**:103–104
 coreceptor/MHC binding sites, **53**:68–70
 coreceptor model, **53**:65–66
 cytoplasmic tail
 protein tyrosine kinase p56^{ck} and, **53**:61–64
 role in mediating coreceptor signal, **53**:73–74
 differential signaling, **53**:91–94
 function in TCR-mediated recognition and activation, **53**:59–61
 GTP-binding protein association, **53**:64–65
 human leukocytes and, **49**:77–78, 81–83, 90, 113
 independent signal transduction, **48**:243
 localization, **44**:295, 297
 mechanism of function during thymocyte selection, **53**:104–108
 MHC class I interactions, **53**:60
 and negative selection, **53**:101–102
 and negative signaling, **53**:61
 and positive selechon, **53**:97–98
 pulmonary immunity, **59**:402–406
 regulatory functions, **48**:242–243
 SCID and, **49**:384, 385, 396
 signaling role in thymocytes, **53**:82
 T cell-specific tyrosine protein kinase, **48**:243–244
 TCR-coreceptor molecule association, **53**:66–68
 and TCR-mediated recognition and activation, **53**:59–61
 and thymocyte development and selection, **53**:94–97
- CD4 accessory molecule, **43**:218
 CD4 receptor, expression by eosinophils, **60**:186–187
 CD5, **48**:246; **52**:140–144; **63**:55–56, 62, 69
 antigen, **49**:151
 human leukocytes and, **49**:92, 101
 Ig heavy-chain variable region genes and, **49**:42
 CD6, **48**:247
 CD7, **48**:247
 CD45 and, **66**:34–35
 CD8, **48**:242–245, 304–306; **59**:52–54, 119, 382
 as accessory molecule, **53**:60
 adoptive T cell therapy of tumors and 283, **49**:288, 290–291, 333–335
 antigen recognition, **49**:319–321, 324
 expression of antitumor responses, **49**:324–330, 332
 mechanisms, **49**:299–306, 316, 318
 autoimmune demyelinating disease and, **49**:374
 and CD4, **52**:363
 coreceptors, **58**:129–131
 CD8 β role in T cell recognition and activation, **53**:84–85
 CD23 antigen and, **49**:152–153
 coreceptor effects on selection, **53**:103–104
 coreceptor/MHC binding sites, **53**:68–70
 coreceptor model, **53**:65–66
 cytoplasmic tail, protein tyrosine kinase p56^{ck} and, **53**:61–64
 differential signaling, **53**:91–94
 down-regulation, versus CD4 down-regulation, **51**:185–186
 function in TCR-mediated recognition and activation, **53**:59–61
 GTP-binding protein association, **53**:64–65
 human leukocytes and, **49**:77–78, 81–83, 101, 113
 IGF-1-R, **63**:403
 independent signal transduction, **48**:243
 mechanism of function during thymocyte selection, **53**:104–108
 MHC class I interactions, **53**:60
 and negative selection, **53**:102–103
 and negative signaling, **53**:61
 and positive selection, development models; **53**:98–100
 T cells, 108–113
 pulmonary immunity, **59**:403–405

- regulatory functions, **48**:242–243
TCR-coreceptor molecule association, **53**:66–68
and thymocyte development and selection, **53**:94–97
CD8 α , **64**:69
CD8 accessory molecule, **43**:218
CD8 $^+$ antiviral factor, **65**:282
 HIV-1 replication, **66**:281, 282–283, 287
CD9, **52**:144
CD10, **52**:144–145; **64**:245
CD11, **62**:258, 272, 276, 279
 human leukocytes and, **49**:86, 92
CD11a/CD18, **64**:146
CD14, role in cellular recognition of LPS, **53**:281–286
CD16, **59**:382
 eosinophils expressing, **60**:205
 human leukocytes and, **49**:77, 90
 neutrophils and NK cells expressing, **60**:179–181
CD18, **62**:258, 272, 276, 279; **64**:169, 187
 human leukocytes and, **49**:86, 92
 mAb, effect on vascular leakage, **60**:329–330
CD19, **52**:145–148; **62**:225
 and CR2 and CR1, molecular complexes on surface of human B cells, **52**:146
CD20, **52**:149–150; **62**:225
CD21, **62**:42, 43–44
 human leukocytes and, **49**:92
CD22, **52**:150–151; **63**:198
 B cell AgR-induced tyrosine phosphorylation of, **55**:245–246
 CD45 and, **66**:50–51
CD23, **49**:149–150, 176–177
 biochemical structure, **49**:155–158
 biological activity, **49**:167, 169
 expression in clinical conditions, **49**:174–175
 membrane CD23, **49**:167–169
 soluble CD23, **49**:169–173
cellular expression, **49**:150–155, 155
 B cells, **49**:150–151
 eosinophils, **49**:154
 Langerhans cells, **49**:154–155
 macrophages, **49**:153–154
 monocytes, **49**:153–154
 T cells, **49**:151–153
cleavage regulation, **49**:158
expression in various clinical conditions, **49**:174–176
expression regulation, **49**:159–161
 B cells, **49**:158–160
 EBV-induced expression, **49**:161
 macrophages, **49**:161
 monocytes, **49**:160–161
Fc ϵ RII
 cloning of cDNA, **49**:162–163
 genomic structure, **49**:165–167
 lectins, **49**:163–164
 structural analysis, **49**:164–165
human leukocytes and, **49**:90–91
IgE receptor corresponding to, **60**:181
ontogeny, **49**:150
transcriptional regulation, **49**:165–167
CD23/Fc ϵ RII, **52**:151–153
 expression, **49**:162, 166–167
CD24, **52**:153
CD25, **59**:379
 eosinophil-bound, **60**:187
 gp39 stimulatory effect, **63**:53
 subunit of IL-2 receptor, **60**:185
CD27, **52**:163; **61**:7, 8, 17
CD27-L, **61**:23, 24
CD28, **48**:244, 245–246, 304; **61**:36–37, 148; **62**:132–135, 156
 costimulatory effects on immunogenicity, **57**:305–307
 human leukocytes and, **49**:101
 and negative selection, **58**:172
 and T cell costimulation, **58**:133–134
CD28-B7 costimulation, Td activation of B cells, **62**:132, 142–144, 148–157
CD28-CD80/CD86 signaling pathway, CD40 expression, **61**:36–37
CD30, **61**:7, 8–9, 12, 17
CD30-L, **61**:23, 24, 25
CD31
 leukocyte transmigration, **65**:369
 lymphocyte homing, **64**:159–160
 properties, **64**:147
 redistribution to cell border, **58**:395
 shedding, **64**:171
 six-Ig domain molecule, **58**:375
CD34
 B cell genesis, **63**:239, 240
 ligand for L-selectin, **60**:163
 lymphocyte–HEV interaction, **65**:362, 377
 properties, **64**:147

- CD34+ cells, **61:12**
 CD37, **52:154–155**
 CD38, **52:155**
 lymphocyte homing, **64:163**
 CD39, **52:163**
 CD40, **52:155–158; 61:1, 2, 7, 54; 62:41–43, 146, 153–154; 63:43–44**
 B cell proliferation, **63:53–58**
 cloning, **61:2, 4**
 expression, **61:32–41; 63:34**
 basophils, **61:15**
 B cell precursors, **61:38**
 B cell receptor pathway, **61:37–38**
 B cells, **61:2, 11–12, 32–35**
 CD28–CD80/CD86 signaling pathway, **61:36–37**
 dendritic cells, **61:13, 40–41**
 endothelial cells, **61:14, 41**
 epithelial cells, **61:14, 41**
 Fas–Fas-L signaling pathway, **61:37**
 fibroblasts, **61:15, 41**
 follicular dendritic cells, **61:14, 41**
 hematopoietic progenitors, **61:12, 39**
 malignant B cells, **61:11–12, 38–39**
 monocytes, **61:13, 40**
 Reed Sternberg cells, **61:12, 39**
 synoviocytes, **61:15, 41**
 T cells, **61:13, 39–40**
 gene, **61:2, 4, 10–11; 63:45**
 gp39 interactions, **63:48, 52–53, 73**
 Ig production, **63:58–70**
 IgA, **63:64–65**
 IgE, **63:63–64**
 IgG, **63:65, 69–70**
 IgM, **63:59, 62–63**
 signaling, **63:58–59**
 intracellular signal transduction, **63:50–52**
 lymphocyte homing, **64:163**
 memory cell formation, **63:70–73**
 molecular structure, **63:45–46**
 signal transduction, **61:27–31, 92–98**
 cAMP, **61:98**
 CD58, **61:97–98**
 IgA, **61:94–95, 96–97**
 IgE, **61:93–94, 95–96**
 IgG, **61:94**
 IgG1, **61:95–96**
 soluble, **63:46**
 structure, **61:2, 4**
 CD40-binding proteins, **63:50–51**
 CD40–CD40-L interactions, **61:54, 55**
 B cell immunopoiesis, **61:47–55**
 in vivo role
 animal models, **61:44–45**
 arthritis, **61:46**
 glomerulonephritis, **61:46**
 immune response, **61:45–47**
 X-linked hyper-IgM syndrome, **61:1, 18, 41–44, 92–93**
 CD40-dependent activation, **52:206–212**
 CD40–gp39 interactions, **63:44, 48, 52–53, 73**
 CD40 ligand, **61:1, 24, 54**
 association with follicular dendritic cells and PALS, **60:269–271**
 B cell proliferation, **63:58, 60–61**
 cloning, **61:17–18; 63:46–48**
 expression, **61:18, 20–22; 63:48**
 gene, **61:18**
 and germinal center reaction, **60:273–276**
 identification, **63:46**
 in IgE isotype switching, **60:38–39**
 molecular structure, **63:48–49**
 in pathogenesis of HIGMX-1, **60:43–48**
 soluble form, **61:26**
 structure, **61:17–18, 19**
 trimerization, **61:26**
 X-linked hyper-IgM syndrome, **61:1, 18, 41–44, 92–93**
 CD40 receptor associated factor 1, **61:29–31**
 CD42, human leukocytes and, **49:85**
 CD43, **48:248**
 human leukocytes and, **49:117**
 CD44, **52:158–159; 59:384**
 and extracellular matrix, **54:282–291**
 human, schematic representation, **54:275, 276**
 human leukocytes and, **49:85–86, 88**
 isoforms, **54:275–281**
 ligand recognition, **54:271–272**
 lymphocyte–HEV interaction, **65:359–360**
 lymphocyte homing, **64:160–161, 165, 177**
 to thymus and bone marrow, **64:177**
 and lymphocytic homing, **54:291–302**
 molecular isoforms and posttranslational modifications, **54:272–282**
 properties, **64:146**
 regulation of interaction with extracellular matrix, **54:302–318**
 role in lymphocyte development, **54:271**

- and T cell development, **58:134**
 and tumor cell migration, role of CD44 in metastasis, **54:318–325**
- CD44** gene, **64:167**
CD45, **48:283; 52:159–160; 63:164–165**, 198
 B cell studies, **66:12–14**, 22–37
 CD2 and, **66:7–8**, 33, 35–36
 CD3 and, **66:9–12**, 34
 CD4⁺ cells and, **66:18**, 26, 35–36
 CD7 and, **66:34–35**
 CD8⁺ cells and, **66:18**, 26, 35–36
 CD22 and, **66:50–51**
 CD100 and, **66:16–17**
 characterization, **58:131–132; 66:1–5**, 55–56
 deficient cell lines
 B cells, **66:22–26**
 T cells, **66:18–22**
 functions, **66:55**
 CD45-deficient cell line studies, **66:18–26**
 gene-targeted mouse studies, **66:26–31**
 lymphocyte development and activation, **66:4**, 5–17
 molecular mechanisms, **66:31**
 regulation, **66:45–49**, 56
 Src family kinases, regulation, **66:4**, 23, 39–45
 structure-function analysis, **66:46–49**
 Fyn, **63:135**
 human leukocytes and, **49:87**, 113–114
 Lck, **63:136**
 lymphocyte adhesion, **66:15–17**
 p53/56^{kn} and, **66:24**, 37, 39, 44
 p56^{lk} and, **66:11**, 12, 19–21, 26, 37–39, 39–44, 53
 p59^{kn} and, **66:19–21**, 37, 38, 39–43
 p70^{ap} and, **66:21–22**, 26
 p72^{sk} and, **66:21**, 23–24, 26, 45
 phenotypic identification of memory T cells with, **53:225–227**
 phosphorylation, **66:53–54**
 post-translational modification, **66:4**, 53–55
 proteolysis, **66:46–47**
 regulation, **66:49–53**, 56
 role in negative selection, **58:172**
 structure
 domain structure, **66:2**
 isoforms, **66:2–3**, 5
 structure–function analysis, **66:46–49**
 substrates, **66:31–39**, 49–53, 55
 T cell studies, **66:6–12**, 18–22, 32–36
 ZAP70, **63:138**
CD45RA, **66:7**, 15, 16, 36
CD45RB, **66:35**
CD45RO, **66:7**, 15, 16, 35
CD46, *see* Membrane cofactor protein
CD48, **52:160**
CD49d, multiple myeloma, **64:246**, 247
CD49d/CD29, properties, **64:146**
CD50, lymphocyte homing, **64:158**
CD55, *see* Decay-accelerating factor
CD56, multiple myeloma, **64:247**
CD57, **61:51**
CD58, **59:377–378**
 CD40 expression, **61:97**
 expression, **64:160**
 multiple myeloma, **64:247**
CD59, **48:240–241; 61:204**, 251–254
 deficiencies in PNH, **60:75–78**, 89
 expression on all blood cells, **60:78**
CD62E, **65:354**
 properties, **64:147**
CD62L, **65:354**
CD62P, **65:354**
 properties, **64:147**
CD69, **48:248–249; 52:163**
 expression on lung eosinophils, **60:186**
CD72, **52:160**
CD73, **48:241; 52:160–161**
 human leukocytes and, **49:112**
 lymphocyte homing, **64:162–163**
 properties, **64:146**
 shedding, **64:171**
CD74, **52:161**
CD76, **52:162**
CD77, **52:162–163**
CD78, **52:163–164**
CD80, **62:132**
CD86, **59:325–326**, 378; **62:132**
CD95, **59:329–331**; **61:9**
 in myeloma cells, **64:259**
CD100, CD45 and, **66:16–17**
CD102, properties, **64:147**
CD106, properties, **64:147**
CD62I, properties, **64:146**
CD antigens, expression on basophils and mast cells, **52:336–337**
cdk, rapamycin, **61:183–187**
cDNA, *see* Complementary DNA

- CDP571**
 rheumatoid arthritis, **64:305, 306**
 ulcerative colitis, **64:310**
- CDR**, *see* Complementarity determining region
- CDw32/Fc_yRII**, **52:153–154**
- CDw75**, **52:161–162**
- C/EBP**, **65:3–7, 3–8**
 knockout mice studies, **65:30**
 structure of, **65:1**
 transcriptional regulation, **65:3–8**
- C/EBP γ** , **65:18–19**
- Celiac disease**
 4-kDa *Bg*/II RFLP fragment, **48:145**
 4-kDa *RSa*I class II fragment, **48:145**
 DP allele, **48:145**
 DP α allele, **48:145**
 DP β allele, **48:145**
 DP α -DQw1 β , **48:146–147**
 DP α polymorphism, **48:145**
 DP β polymorphism, **48:145**
 DQw2 α chain, **48:146**
 DR3, **48:144, 147**
 DR3-DQw2 haplotype, **48:144**
 DR5, **48:144, 147**
 DR7, **48:144, 147**
 DR α -DQw1 β , **48:146–147**
 DRw8, **48:147**
 gluten-derived peptide, **48:147–148**
 HLA class II, **48:144**
 HLA class II antigen, **48:147**
 hybrid HLA-DQ molecule, **48:147**
 with malignant histiocytosis, gene rearrangements and, **40:292–293**
 and $\gamma\delta$ TCR expression, **58:329**
 triggering factor, **48:144**
- Cell adhesion**
 B cell formation and, **41:185, 235, 237, 239**
 B cell precursors, **41:194, 195, 198, 200, 201**
 bone marrow cultures, **41:209, 212, 213, 219, 220**
 genetically determined defects, **41:228, 231**
 lymphohemopoietic tissue organization, **41:186–188**
 population dynamics, **41:208**
 soluble mediators, **41:234**
 stromal cells, **41:214–216**
 cascade, **58:381–389; 64:181–184**
- CD23 antigen and, **49:163–165, 169, 176**
 cytotoxicity and, **41:271, 272**
 deficiencies, **64:193–195**
 eosinophils, **60:164–178**
 flowing leukocytes to platelets, **58:355**
 hybrid resistance and, **41:368, 393**
 leukocyte–endothelial, **58:376–378**
 lymphocytes, **66:15–17**
 in high endothelial venules, **65:367–372**
 mechanisms, and eosinophils, **51:333–336**
 neutrophil, IL-8 effects on, **55:114**
 by stem cell factor and ligand interaction, **55:37**
 strong, role of integrins, **58:383–388**
 tight, pro-T cells to endothelium, **58:393**
 vascular, and emigration of inflammatory cells, **60:330–332**
- Cell adhesion, receptors and recognition molecules**, **52:357–366**
 integrins, **52:358–361**
 β_1 integrins, **52:358–359**
 β_2 integrins, **52:359–361**
 β_3 integrins, **52:361**
 other recognition molecules, **52:366**
 recognition molecules
 CD4 and CD8, **52:363**
 ICAM-1/CD54, **52:361–363**
 Ig supergene family, **52:361–364**
 LFA-2/CD2 and LFA-3/CD58, **52:361**
 MHC class I and class II, **52:364**
 other, Ig superfamily, **52:364**
 stem cell factor receptor/c-kit, **52:363–364**
 VCAM-1, **52:363**
 selectins and related molecules, **52:364–366**
- Cell adhesion molecules**, **52:166–168; 59:383**, *see also specific molecules*
 activation of, **52:168**
 affecting migration of naive and memory cells, **53:248–249**
 clustering, **64:173**
 cocapping, **64:173**
 endothelial, and lymphocytes, **58:352**
 endothelial binding, **64:146**
 expression and function regulation, **64:139–140, 146, 147, 163–173**
 Ig superfamily, **52:167**
 integrin family, **52:166**
 lymphocyte binding, **64:147**

- multistep adhesion cascade, **64**:181–184
in myeloma cells, **64**:246–249
parallels with regulatory mechanisms,
 54:308–310
 LFA-1, **54**:309–310
 L-selectin, **54**:310
 Mac-1/CR3, **54**:310
 platelet gpIIb-IIIa integrin, **54**:309
production during inflammation, **58**:10
pulmonary inflammation, **62**:267, 288
regulation
 alternative splicing, **64**:167
 conformational events, **64**:169–170
 internalization, **64**:170–172
 oligomerization, **64**:172–173
 ontogenetic, **64**:164–165
 posttranslational modifications,
 64:168–169
 secretion, **64**:170–171
 shedding, **64**:170–171
 upregulation and induction, **64**:
 165–167
SCID and, **49**:384, 397
selection family, **52**:167–168
surface antigens of human leukocytes and,
 49:79, 82–89, 86, 88, 114, 116
 antigen-specific receptors, **49**:78–80
 complement components, **49**:92–93
 as targets of autoantibodies in pemphigus
 and pemphigoid, **53**:309, 312, 314
- Cell cycle
control, IL-2-dependent G₁-phase
 progression, **61**:179–187
control mechanisms in tolerance,
 54:411–414
 blocking negative signaling with
 antisense myc, **54**:413–414
 myc, TGF- β and pRB, **54**:411–413
immunosenescence and, **46**:221–223, 227,
 234, 235, 253
interventions, **58**:248–250
regulation of V(D)J recombination and
 DSB repair, **58**:70–74
regulatory kinase p34^{cdk}, blockade, **58**:267
switch recombination, **61**:116–117
- Cell cycle checkpoints
G₀–G₁, and induction of apoptosis,
 58:249–250
mechanisms, **58**:66–69, 71–74
mutations in, **58**:29
- Cell death, *see also* Apoptosis
ataxia telangiectasia cells, **58**:69
B cells, **59**:296–300, 336–339
deficient regulation, and disease states,
 58:212
Fas-mediated, **57**:129, 133–134, 140
in immune system, *see* Apoptosis, in
 immune system
intrathymic, **58**:166–168
T cell
 induction, **58**:224–240, 238
 inhibition, **58**:240–265
 by IL-2, **58**:269
 in vitro, **58**:242–253
 in vivo, **58**:253–265
 TCR-mediated, **58**:215–224
- Cell degradation, redundant effector
pathways, **58**:269–270
- Cell density, IL-2 activity and, **42**:166
- Cell lines, *see also* specific cell lines
CD44 expression and hyaluronan binding,
 54:303
- CD44-positive, **54**:304
- IL-1ra production, **54**:190
- murine, perforin-deficient, **60**:293–294
mutant, and GPI anchor synthesis,
 60:65–67
- PNH, **60**:79–80
- T cell, Trac, **60**:43
verification of perforin in, **51**:224–227
- Cell lysis, **51**:25–28
- Cell-mediated cytotoxicity
HIV infection and, **47**:410, 411
NK cells and, **47**:189–196
- Cell-mediated killing, **41**:319, 320
granule proteins and
 cell lines, **41**:286, 287
 cytoplasmic granules, **41**:287–291
 lymphotoxin, **41**:298
 membrane attack complex, **41**:310
 proteoglycans, **41**:298, 299
 serine esterases, **41**:297, 298
 TNF-related polypeptides, **41**:298
- pore-forming proteins and
amoebas, **41**:312, 313
biochemical properties, **41**:292–295
eosinophil cationic protein, **41**:311, 312
membrane binding, **41**:295, 296
purification, **41**:291, 292

- Cell-mediated lympholysis
hybrid resistance and, **41**:355, 364,
394–396
T cell subsets and, **41**:43, 45, 51, 57,
108
- Cell-mediated lysis, hybrid resistance and,
41:407
- Cell receptor binding, segmental flexibility,
48:24–26
- Cell surface molecules
T cell activation and, **41**:1, 2
accessory molecules, **41**:14, 15
IL-1 receptor, **41**:13, 14
T1, **41**:13
T11, **41**:8–10
T cell antigen receptor, **41**:2–8
Thy-1, **41**:10, 11
Tp44, **41**:11–13
T cell subsets and, **41**:40, 43–48
accessory molecules, **41**:49, 50
T cell receptor, **41**:40–43
TH responses, **61**:355–357
- Cell surface structures, functionally associated
with Fc ϵ RI molecules, **52**:391–396
- cell surface glycolipids, **52**:393–394
- Fc γ receptors, **52**:394–396
- G63, **52**:394
mast cell chromolyn binding sites,
52:392–393
- ME491 (CD63), **52**:393
- Cell surface typing, with mAbs: human mast
cells and basophils, **52**:335–339
- Cell survival, role of stem cell factor
receptor-ligand interaction, **55**:
38–39
- Cell tropism, HIV infection, **63**:84, 90
- Cellular control mechanisms, summary,
52:73–74
- Cellular distribution, receptors for Fc region
of IgG3–64
basophils and mast cells, **40**:68
B cells, **40**:66–67
macrophages and monocytes, **40**:64–66
neutrophils, **40**:68
platelets, **40**:68
T cells, **40**:67–68
- Cellular immune response
effector phase, **60**:332–346
regulation of, **40**:77–79
and Th1 cells, **60**:348–349
- Cellular immunity, **64**:3; **65**:277–278,
322–323
adenovirus, **65**:279
cytomegalovirus infection, **65**:278–279
Epstein–Barr virus (EBV), **65**:279
evolution of, **52**:108–111
antigen processing and presentation,
52:108–110
T cell antigen receptor, **52**:110–111
herpes simplex virus, **65**:279
HIV-1 disease, **63**:97, 98
HIV infection, **65**:280
adoptive immunotherapy, **65**:317–320
CTL-mediated lysis, **65**:280–281
CTLs and, **65**:280–322
HLA and, **65**:284–286
replication suppression, **65**:281–284
in seronegatives, **65**:287, 307–309
vaccines, **65**:320–322, 323
HLA class I system, **65**:279
hybrid resistance and, **41**:370
influenza virus, **65**:278
pituitary hormones, **63**:407
- Cellular switches, self-tolerance, **59**:281
- Cellulose acetate, hybrid resistance and,
41:372
- Censoring, B cells, **59**:302–304, 308–309,
311–312, 331–341
- Central nervous system
allergic reactions, **61**:365–367
autoimmune demyelinating disease and,
49:357, 369
autoimmunity in, **62**:98–99
 β -endorphin receptors, distinction from
those on lymphocytes, **39**:312
HIV infection and, **47**:378, 413
immunopathogenic mechanism, **47**:393
neuropsychiatric manifestations, **47**:403,
404
IFN- γ and, **62**:91–92
lesions, immunological effects, **39**:299
NK cells and, **47**:266–269
- Centrocytes, **61**:51, 53
CD23 antigen and, **49**:171
- Centromeres
antineuclear antibodies and, **44**:121, 122,
130, 131, 133
maps of Ig-like loci and, **46**:26, 30, 33, 44
- Cerebrovascular disease, aPL antibodies and,
49:220

- Cervical cancer, antigens, **62:240**
Cetirizine, modulated eosinophil migration, **60:174**
c-fms, and *c-kit*, evolutionary relationship, **55:9–10**
c-Fos
 B cell genesis, **63:220–221**, 248–250
 and *c-Jun*, **58:138–139**
 T cell activation and, **41:27**, 28
c-fos
 IL-2 signal transduction, **63:161**, 171
 SIE, **63:141**
 T cell activation, **48:291–292**
CH3 domain
 Fc, **48:20–21**
 IgG, **48:20–21**
Channel formation
 cytolytic protein and, **41:318**, 319
 membrane attack complex of complement, **41:301–303**, 306, 307, 311
Charcot-Leyden crystal
 in basophils, **39:183**
 in eosinophils, **39:181–183**
 lysophospholipase activity, **39:182–183**
Charcot-Leyden crystal protein, **60:156**, 160, 164, 201
Chediak-Higashi syndrome, NK cells and, **47:224**, 225, 299
C-helix, peptide homologs, **43:63**
Chemiluminescence, NK cells and, **47:263**, 266
Chemoattractants
 eosinophil, **60:174–178**
 lymphocyte homing, **64:152–154**
Chemoimmunoconjugates, as cancer treatment, *see also* Alkylating agent–antibody conjugates; Anthracycline–antibody conjugates
barriers to antibody-targeted chemotherapy
 biodistribution, **56:348**
 effects of, **56:347–348**
 route of administration, **56:348–350**
 tumor antigen-heterogeneity, **56:352–354**
 tumor perfusion, **56:350–352**
 tumor vasculature, **56:350–352**
chemistry
 cis-aconitic anhydride, **56:318–319**
 carbodiimides, **56:316–318**
 conjugation procedure, **56:312–314**, 360
conjugation strategies, **56:314–323**
cyanuric chloride, **56:321**
design strategy, **56:324–325**
diazo reaction, **56:319**
ester linkage, **56:323**
glutaraldehyde, **56:315**
hydrazone linkage, **56:322–323**
imidoesters, **56:320–321**
intermediary carriers, **56:323–324**
mixed anhydrides, **56:319**
noncovalent bonds, **56:321–322**
periodate oxidation, **56:315–316**
photoactivation, **56:323**
N-succinimidyl 3-(2-pyridyl)dithio) propionate, **56:319–320**
thioether linkage, **56:321–322**
clinical trials, **56:354–361**
drug–monoclonal antibody conjugates
 alkylating agent–antibody, **56:328–333**
 anthracycline–antibody, **56:333–338**
 antineoplastic drugs, **56:313**
 cytotoxic drugs, **56:342–343**
 folic acid antagonists, **56:338–340**
 functional studies, **56:344–347**
 mode of action, **56:344–347**
 morphological studies, **56:347**
 preclinical testing, **56:326–328**, 343
 vinca alkaloids, **56:340–342**
monoclonal antibodies as carriers
 conjugate size as determinant of permeability, **56:310–311**
development, **56:302–304**
heterogeneity, **56:306–308**
immunogenicity, **56:311–312**
internalization, **56:308–310**
localization, **56:306–308**
modulation, **56:308–310**
specificity, **56:305–306**
targets, **56:305–306**
toxicity, **56:311–312**
studies, **56:360–361**
Chemokine receptors
 C5a, gene mapping, **55:135**
CC
 amino acid sequences, **55:129**
 biochemical studies, **55:125–128**
 on erythrocytes, **55:128**
cloning, **55:131**, 134
CXC
 amino acid sequences, **55:129**

- Chemokine receptors (*continued*)
 binding, 55:136
 biochemical studies, 55:122–125
 ELR motif, 55:136–138
 on erythrocytes, 55:128
 genes, 55:135
 G-protein coupling, 55:138–139
 LESTR52, cloning of, 55:131, 134
 lymphocytes in high endothelial venules, 65:367–368
 MCP-1, binding studies of, 55:125–126
 MDCR15, cloning of, 55:131, 134
 MIP-1 α
 cloning, 55:131
 gene cloning, 55:135
 RANTES, binding studies, 55:126–128
 regulation, 55:139–140
 structure, 55:132–135
 7-TM, G-protein coupling to, 55:138
- Chemokines, 62:80, *see also* Cytokines
 allergic airway inflammation, 62:285
 arthritis role, 55:143–144, 147
 biological effects, 55:115–122
 cardiovascular system role, 55:146
 CC
 biological effects, 55:120
 on basophils, 55:116–117
 cellular sources, 55:105
 chemical properties, 55:98, 100–102
 inactivation, 55:102–103
 molecular structure, 55:98, 100–102
 proteolytic processing, 55:102–103
- cellular effects
 basophils, 55:116–117
 eosinophils, 55:117
 neutrophils, 55:115–116
- cellular sources, 55:105–112
 leukocytes, 55:105–107
 macrophages, 55:106
 monocytes, 55:105–106
 tissue cells, 55:107, 111
- connective tissue-activating peptide III
 biological effects on neutrophils, 55:115
 chemical properties, 55:97, 99
- CXC
 biological effects, 55:115, 118–120
 on basophils, 55:116
 on lymphocytes, 55:119
 on monocytes, 55:118
 on neutrophils, 55:115–116
- cellular sources, 55:105
 chemical properties, 55:97–101
 inactivation, 55:102–103
 molecular structure, 55:97–101
 proteolytic processing, 55:102–103
 in eosinophil migration, 60:177–178
- genes
 chromosomal localization, 55:103–104
 expression regulation, 55:104–105
 organization, 55:103
- granulomatous lesions, 62:282
- GRO α , chemical properties, 55:100
- HIV-1 antiviral effect, 66:281–282
- in integrin activation, 58:378–379
- in vivo* effects, 55:121–122
- lung disease role, 55:144–147
- lymphocyte adhesion and, 64:152–154
- lymphocytes in high endothelial venules, 65:367–368
- MCP-1
 biological effects on monocytes, 55:118
 chemical properties, 55:101
 production by monocytes, 55:106
- MIP-1 β
 biological properties, 55:101
- NAP-2, chemical properties, 55:99–100
- in neutrophil recruitment, 60:339
- pathology role, 55:142–148
- platelet factor 4
 biological effects on neutrophils, 55:115
 chemical properties, 55:97, 99–100
- pulmonary immunity, 59:373
- pulmonary inflammation, 62:259, 268–270
- RANTES, biological effects on eosinophils, 55:117
- rheumatoid arthritis and, 64:294, 316, 317
- sepsis and
 IL-8, 66:135–141
 MCP-1, 66:141–142
- signal transduction, 55:140–142
- skin disease role, 55:142–143, 147
- Chemotactic receptor, surface antigens of
 human leukocytes and, 49:93, 111
- Chemotaxis
 adoptive T cell therapy of tumors and, 49:328, 335
 complement receptor 1 and, 46:192, 199

- eosinophils, **60**:174–178
IBD and, **42**:307, 308
IL-2 and, **44**:166, 169
leukocyte integrins and, **46**:151, 160, 161,
 163, 164, 166, 167
lymphocyte homing and, **44**:320
stem cell factor–ligand interaction in,
 55:37–38
- Chemotherapeutic agents, MG and,
 42:265–268, *see also* specific agents
- Chemotherapy
 adoptive T cell therapy of tumors and,
 49:287, 292–296, 311
 hybrid resistance and, **41**:350
 SCID and, **49**:388, 390, 396–397
 targeted, concept of, **56**:301–303
- CH gene, **43**:239
- Chicken
 B cells, repertoire formation in, *see* B cells
 Ig_l gene, **48**:41–64
 genomic organization, **48**:44–46
 germ-line allele, **48**:49–51
 Ig_l allele, **48**:49–51
 JL gene segment, **48**:44–46
 rearrangement, **48**:46–49
 somatic diversification, **48**:41–64
 structure, **48**:44–46
 V gene segment, **48**:44–46
 V segment, **48**:52–54
- ovalbumin, T cell subsets and, **41**:53
- retrovirus-induced B cell neoplasia in bursa
 of fabricius
 research, **56**:468–469
 resistance, **56**:470
 susceptibility, **56**:469–470
 transformed follicles in lymphoma-
 resistant strains, **56**:471
- Chimeras, *Ir* genes and, **38**:58–61, 70
 allogeneic, **38**:62, 64
 allophonic, **38**:62, 63
 antigen binding, **38**:114
 complementation, **38**:73, 74
 cross-reactive lysis, **38**:186
 II-Y antigen, **38**:176–179
 immunodominance, **38**:182–184
T cells, **38**:64
 cytotoxic, **38**:165–167, 172
 interactions, **38**:158–160
 selection, **38**:131, 133, 134
 tolerance, **38**:120, 123, 126–129
- Chimeras, radiation, **59**:106, 109–111
Chimeric antibodies, **44**:79–82; **48**:2; **64**:261,
 263, 323
 therapeutic, **44**:87, 88
- Chimeric genes, genetically engineered
 antibody molecules and, **44**:76, 77,
 83, 84
- Chimeric mouse, RAG-2-deficient blastocyst
 complementation, **62**:36
- Chimerism, **59**:106, 109–111, *see also*
 Genetic chimerism
 functional, involving DNA binding and
 protein dimerization, **50**:133–138
- IL-2 receptors, **59**:258–259
- Chimpanzee, HIV-1 infection, **52**:455–461
 model for testing HIV-1 vaccines,
 52:457–459
 pathogenesis of HIV-1 infection in,
 52:455–457
 use to test CD4-based therapies,
 52:459–461
- Chlamydia trachomatis*, vaccine design,
 60:136
- C_{II} locus, **62**:1
- Chlorambucil, chemotherapeutic use in
 cancer treatment, **56**:328–330, 344–345
- Chloroleukemia, cell line, rat, observation,
 50:73
- Chloroquine, **41**:56, 57, 281; **42**:176
 antigen processing and, **39**:63–64
 humoral immune response and
 antigens, **45**:13–15, 17, 20
 physical interaction, **45**:38, 51
 synthetic T and B cell sites and, **45**:223
- Chlorpromazine, aPL antibodies and, **49**:203,
 212–213, 221, 224, 257
- Cholecystokinin receptor, **48**:181
- Cholera toxin, **61**:98
- Cholestasis, intrahepatic severe, **50**:190–191
- Cholesterol
 aPL antibodies and
 affinity purification, **49**:229–230
 LA antibodies, **49**:244–245
 specificity, **49**:234–236, 248
 cytotoxicity and, **41**:279, 300, 302, 317, 318
 spontaneous autoimmune thyroiditis and,
 47:439
- surface antigens of human leukocytes and,
 49:116

- Cholinergic nerve differentiation factor, *see also*
 Leukemia inhibitory factor
- Chondroitin sulfate A, cell-mediated killing
 and, 41:298, 299
- CHOP-10, 65:3–6
- Chromatids
 maps of Ig-like loci and, 46:3
 murine lupus models and, 46:83
- Chromatin
 antinuclear antibodies and, 44:130
 B cell formation and, 41:204, 221
 configuration during V(D)J joining of
 antigen receptor genes, 56:37, 41–46
- cytolysis and, 41:283
- Ig gene superfamily and, 44:12
- Ig heavy-chain variable region genes and,
 49:37, 56
- immunosenescence and, 46:234
- NK cells and, 47:214
- structure, transcription regulation,
 63:200–201
- T cell development and, 44:217
- Chromosomal abnormalities
 on adjacent oncogenes, consequences,
 50:124–132
c-myc in Burkitt's lymphoma
 translocations, 50:124–128
 gene fusion resulting from chromosome
 translocation, 50:128–129
 helix-loop-helix oncogenes in
 chromosome translocacions,
 50:129–130
 oncogene location after translocation,
 50:131–132
 transcriptional disruption of LIM domain
 oncogenes by translocation,
 50:130–131
- lymphoid cells, effect, 50:132–140
 differentiation-related translocation
 oncogenes, 50:139
- functional chimerism involving DNA
 binding and protein dimerization,
 50:133–138
- other unusual proteins affected by
 chromosome translocacions,
 50:139–140
- Chromosomal mapping
 B cell surface antigens, 52:129, 137
 T cell receptor and, 38:16, 17
- Chromosomal translocation, *see also*
 Translocation
- Ars A response and, 42:129–130
- Burkitt's lymphoma, *c-myc* in, 50:124–128
c-myc gene, 64:227–229
 gene fusion resulting, 50:128–129
 helix-loop-helix oncogenes in, 50:129–130
 human B cell neoplasia and, 38:245–248,
 272
 in Burkitt lymphoma, 38:253–259, 271
c-myc activation, 38:261, 263
c-myc oncogene, 38:248, 253–259, 271
 t(8;14), 38:249–251, 254, 260
 t(11;14), 38:264–267
 t(14;18), 38:267–271
 and inversion, in lymphoid tumors,
 mechanism, 50:121–122
 other unusual proteins affected by,
 50:139–140
- t(8;14)
c-myc activation, 38:260, 264
 human B cell neoplasia, 38:249–251,
 254, 269
 t(11;14) chromosome translocation,
 38:264–267, 272
 t(14;8) chromosome translocation,
 38:264
 t(14;18) chromosome translocation,
 38:267–272
 T cell receptor δ genes, 43:160–161
 and timing of inversion, 50:122–124
- Chromosome 1
 CD1 locus, 58:319
 selectins on, 58:353
- Chromosome 2, radiation hybrid with *xrs*
 cells, 58:57
- Chromosome 4q, IL-2, gene, human, 39:5
- Chromosome 5, transfer into *XR-1* cells,
 58:52
- Chromosome 8, complementation of *scid*
 defects, 58:50–51
- Chromosome 11
 GM-CSF gene, murine 9, 39:18
 IL 3-gene, murine 7–8, 39:18
- Chromosome 14
 q region shared by ECP and eosinophil-
 derived neurotoxin, 60:193
 rearrangement in ataxia telangiectasia,
 58:68
- V_H locus mapping, 62:4–7, 23, 24

- Chromosome 15
orphan V_H and D_H loci, **62**:10–12, 19, 20, 23
translocation, **64**:228
- Chromosome 16, orphan V_H and D_H loci, **62**:10–12, 19, 20, 23
- Chromosome 19, integration of adeno-associated virus, **58**:424–425
- Chromosome 22, location of p70^{Ku}, **58**:57
- Chromosomes
antinuclear antibodies and, **44**:99, 105, 121, 122, 125
autoimmune thyroiditis and, **46**:268, 272
CD5 B cell and, **47**:132, 148, 157
CD8 molecules and, **44**:271, 276–278
chemokine gene, localization, **55**:103–104
c-kit location, **55**:7–9
complement receptor 1 and, **46**:190
complement receptor 2 and, **46**:202
errors, **58**:72–74
GPI synthesis genes, location, **60**:71
HIV infection and, **47**:380
humoral immune response and, **45**:32
Ig gene superfamily and, **44**:9, 34, 47
Ig heavy-chain variable region genes and, **49**:1, 3, 6, 55
IL-1 and, **44**:156
IL-1ra genes, localization, **54**:177–180
interchromosomal recombination during V(D)J joining of antigen receptor genes, **56**:120–121
leukocyte integrins and, **46**:159, 160, 170
maps of Ig-like loci and, **46**:2, 3, 5–7
MHC, **46**:28, 31, 33
structure, **46**:7, 9–13, 18, 19, 21, 23, 25–27
T cell receptor, **46**:36, 39, 42, 43, 45
murine and human Jak's, location, **60**:3–4
murine lupus models and, **46**:63, 64
NK cells and, **47**:227, 228, 231
regulators of complement activation and, **45**:381, 403, 404, 411
stem cell factor location, **55**:13–15
surface antigens of human leukocytes and, **49**:91–92
T cell receptor and, **45**:110, 115
- Chronic graft-versus-host disease
cellular mechanisms, **40**:408–409
clinical manifestations, **40**:406–407
pathologic manifestations, **40**:407–408
risk factors, **40**:409–410
treatment and prevention, **40**:410
- Chronic granulomatous disease, X-linked, **59**:248
- Chronic inflammatory disease, *see* Inflammatory bowel disease
- Chronic lymphocytic leukemia
CD5 B cell and
aging, **47**:123
autoimmune diseases, **47**:150
Ig gene expression, **47**:153–159
malignancies, **47**:123, 124
physiology, **47**:136, 140, 141
surface antigen, **47**:146, 147
NK cells and, **47**:298
- Chronic myelogenous leukemia
circulating IL-4 levels, **53**:14–15
leukocyte integrins and, **46**:160
lymphoid blast crisis of, gene rearrangements and, **40**:290–291
- Chronic myeloid leukemia, B cell formation and, **41**:188
- Chronic obstructive pulmonary disease, **62**:261, 286
- Churg–Strauss syndrome
complication of asthma, **60**:216
in pulmonary eosinophilia, **60**:226
- Ciliary neurotrophic factor, **65**:20
CNTF/LIF/OSM/IL-6 subfamily, **61**:151–152
- Ciliary neurotrophic factor receptor and high-affinity LIF receptor, **53**:43–44, 46
pleiotropic nature of, **53**:46–47
cim, **65**:48–49, 78
- Circumsporozoite proteins
derived T cell epitopes, **60**:108–123
sporozoite malaria vaccine and, **45**:292–294, 319–322
antibodies, **45**:300
cloning, **45**:294–296
endemic areas, **45**:315–317
human trials, **45**:315
immunity, **45**:291, 292
interferon-γ, **45**:306
Plasmodium vivax, **45**:317, 318
repeats, **45**:296–300
synthetic T and B cell sites and, **45**:228, 258

- Circumsporozoite reaction, sporozoite malaria vaccine and, 45:288, 290
- Cirrhosis, primary biliary, and antibodies to self-antigens from human donors, 57:244–245
- Cisplatin and apoptosis, 58:237–240
in cancer treatment, 56:329, 332
- cis-regulatory elements, 62:37, 42–43, 49–52
- c-Jun, 62:42, 48
and c-Fos, 58:138–139
- c-jun oncogene, IL-2 signal transduction, 63:160–161, 171
- CK-2 sequence, 48:81–82
- CK226 antigen, 52:173
- CK-I sequence, 48:81–82
- c-kit, 60:26
- c-kit
allelism with W, 55:6–7
and c-fms, evolutionary relationship, 55:9–10
characteristics, 55:3–6
chromosomal localization, 55:7–9
genomic organization, 55:7–9
and PDGFR, evolutionary relationship, 55:9–10
- c-kit, stem cell factor receptor, 52:352–354
- C3 complement component degradation fragments, 52:147
third component of, 52:148
- CLA, *see* Cutaneous lymphocyte antigen
- Classic CD1 proteins, *see* CD1, group 1
- Class II inactivator (CIITA), 62:106
- Class I modifiers, 65:48–49, 78
- Class switching, 61:79–81
B cell differentiation, 61:82–84
heavy chain isotype expression, 61:82–84
isotype specificity regulation, 61:84–99
B cell activators, 61:90–92
CD40 signaling, 61:92–98
cytokines, 61:84–89
problems in other studies, 61:81–82
switch recombination, 61:129–130
3' Cα enhancer, 61:121–123
control during cell cycle, 61:116–117
fine sequence specificity, 61:107–108
germline transcripts, 61:111–116
illegitimate priming model, 61:108–111
intrachromosomal deletion, 61:99–100
mutations, 61:108
protein-binding sites, 61:117–122
RNA splicing, 61:101–102
sequential switching, 61:105–107
sites, 61:102–105
substrates, 61:123–129
transchromosomal switching, 61:100–101
- Clathrin-coated pits, complement receptor 1 and, 46:199
- CLC, *see* Charcot–Leyden crystal
- CLC protein, *see* Charcot–Leyden crystal protein
- Cleavage
CD23 antigen and, 49:156, 158, 160, 173
coupled, V(D)J recombination, 64:53–54
cytolytic protein and, 41:318
membrane attack complex of complement and, 41:299, 306
model, V(D)J recombination, 58:38–39
RAG, 64:57–58
reactions, in V(D)J recombination pathway, 58:30, 33–35
V(D)J recombination and, 64:52–54
- Clinical disorders, and IL-6, 54:44, 45
- Clonal abortion, 50:64; 52:312
and clonal anergy, 52:291–294
- Clonal anergy, 48:217, 218
- Clonal deletion, *see also* Deletion
intrathymic, 51:165–173
cell biology of negative selection, 51:165–170
search for mechanism, 51:170–173
- lymphocyte, physiology, 58:270–272
- MTV superantigen-reactive T cells, 65:203–204, 210
- nascent bone marrow B cells, 58:257
and peripheral tolerance to cancer, 58:421
putative interference of IL-2, VVith, 50:166–167
- Clonal dominance, PIG-A mutant cells, 60:90–91
- Clonal expansion, and V gene expression, 42:32–34
- Clonal inactivation, by Vβ selective elements, 50:35–36
- Clonality, definition based on Ig and T cell receptor gene rearrangements, 40:295–305
- Clonal origin, PNH cells, 60:80–81
- Clonal suppression, 48:218

Clones

adoptive T cell therapy of tumors and, **49**:284, 289–290, 298, 334
antigen recognition, **49**:321, 324
expression of antitumor responses, **49**:325–326, 330
mechanisms, **49**:299, 301, 303–305, 311
antigen-presenting cells and, **47**:45
antigen presentation, **47**:65, 66, 68–71, 73
antigen processing, **47**:91, 92
APC-T cell binding, **47**:95, 99
cell surface, **47**:61, 62
immunogenicity, **47**:102, 103
interaction with T cells, **47**:94
T cell-dependent antibody responses, **47**:83
T cell growth, **47**:99
tissue distribution, **47**:48, 49
antinuclear antibodies and
autoantibodies, **44**:127, 137
autoantigen, **44**:128–131
autoimmune response, **44**:132
scleroderma, **44**:122
SLE, **44**:101, 108
aPL antibodies and, **49**:238–239, 246, 249
autoimmune demyelinating disease and, **49**:357
multiple sclerosis, **49**:370, 372, 374
myelin basic protein, **49**:358–361
TCR usage restriction, **49**:363–368
autoimmune thyroiditis and, **46**:263, 317, 318
antigens, **46**:270, 272
cellular immune responses, **46**:288, 289, 296
experimental models, **46**:276, 279, 281, 282
humoral responses, **46**:298, 301, 306, 308
prevention, **46**:312, 313
autoreactive T cell, **45**:418, 433
activation, **45**:421–426
origin, **45**:418–420
physiology, **45**:427, 428, 430, 431
regulation, **45**:431, 432
B cell formation and, **41**:182, 184, 185, 238
B cell precursors, **41**:191, 196, 200, 201
bone marrow cultures, **41**:213, 216, 217, 219, 220

C3H/HeJ mice, **41**:231
Ig genes, **41**:203, 205
inducible cell line, **41**:220–222
lymphohemopoietic tissue organization, **41**:187
NZB mice, **41**:226, 229
population dynamics, **41**:207
SCID mice, **41**:225, 226
W/W anemic mice, **41**:224
CD4 and CD8 molecules and, **44**:266–269, 298, 299, 301
CD4 molecular biology, **44**:285, 292, 293
CD5 B cell and
aging, **47**:122
Ig gene expression, **47**:151, 152, 154, 156, 158
physiology, **47**:133, 134, 136, 140, 142
surface antigen, **47**:146
CD8 molecular biology, **44**:271, 273, 274, 276–280, 282
CD23 antigen and, **49**:151–152, 154, 162–165
complement receptor 1 and, **46**:188, 199
complement receptor 2 and, **46**:201, 202, 211
cytolysis and, **41**:274, 276, 280, 281, 285
cytotoxicity and, **41**:319, 320
granule proteins, **41**:286, 287, 291, 297
nature of mediation, **41**:271, 273
pore farmers, **41**:317, 319
cytotoxic T cells and, **41**:167
 β_2 -microglobulin, **41**:156
amino acid changes, **41**:162
carbohydrate moieties, **41**:154
exon shuffling, **41**:138, 142, 144–146, 149
HLA class I antigens, **41**:149–151
mAbs and, **41**:162, 166, 167
genetically engineered antibody molecules and, **44**:73–75, 78–80
helper T cell cytokines and, **46**:111, 115, 116, 118, 121, 122
cross-regulation of differentiation, **46**:130, 131
differential induction, **46**:130
functions, **46**:114, 117, 119, 120
human subsets, **46**:139
immune responses, **46**:136–138

- Clones (*continued*)**
- precursors of differentiation states, **46:123**
 - secretion patterns, **46:112**, 113
 - HIV infection and, **47:400**, 411
 - etiological agent, **47:380**, 385
 - immunopathogenic mechanism, **47:388**, 390, 394
 - humoral immune response and
 - antigens, **45:16**, 17, 24
 - cellular interactions *in vivo*, **45:80**, 83
 - helper T cells, **45:25–28**, 30–32
 - interleukins, **45:62**, 63, 66, 69, 74, 75
 - physical interaction, **45:35–37**
 - hybrid resistance and antibodies, **41:377**
 - antigen expression, **41:405**
 - marrow engraftment, **41:387**
 - NK cells, **41:375**
 - T cells, **41:383**
 - in vitro* assays, **41:394**
 - IgE biosynthesis and
 - antibody response, **47:6**, 7, 10–12
 - antibody response suppression, **47:33**, 39
 - binding factors, **47:13**, 15–18, 20, 21, 27
 - Ig gene superfamily and
 - evolution, **44:46**
 - MHC complex, **44:29**
 - receptors, **44:17**, 18, 22–24
 - Ig heavy-chain variable region genes and,
 - 49:5**, 8, 22, 32
 - regulation, **49:55**
 - V_H gene expression, **49:41–42**, 47–48
 - IL-1 and, **44:155**, 179, 189
 - immunosenescence and, **46:237**, 249
 - leukocyte integrins and, **46:156**, 164, 171, 172
 - maps of Ig-like loci and, **46:3–5**
 - MHC, **46:29**, 32, 33
 - structure, **46:11**, 13, 15, 17, 18, 21, 23–27
 - T cell receptor, **46:36**, 37, 45
 - murine lupus models and, **46:99**, 100
 - Ig germline, **46:66**, 67, 71, 76–78
 - lupus strains, **46:63**
 - T cell antigen receptor, **46:79**, 82, 83, 88, 90, 91, 93, 95–97
 - NK cells and, **47:188**, 199, 291
 - cytotoxicity, **47:252**
 - differentiation, **47:231**, 234
 - effector mechanisms, **47:240**, 243, 245–248, 254, 264, 265
 - hematopoiesis, **47:273**, 274, 277, 279
 - malignant expansion, **47:228**
 - morphology, **47:217**
 - surface phenotype, **47:202**, 204–207, 209, 210, 212
 - regulators of complement activation and, **45:391**, 396, 398, 399
 - SCID and, **49:393**, 398, 402
 - self-reactive, **58:213**
 - spontaneous autoimmune thyroiditis and, **47:464**, 477
 - sporozoite malaria vaccine and
 - antibodies, **45:300**
 - CS proteins, **45:292**, 294–296, 299
 - CS-specific T cells, **45:310–312**
 - surface antigens of human leukocytes and, **49:79**, 91, 98, 111, 114, 116
 - synthetic T and B cell sites and, **45:261**, 262
 - bacterial antigens, **45:230–232**
 - globular protein antigens, **45:211**, 214
 - parasitic antigens, **45:229**
 - viral antigens, **45:218–220**, 222–224
 - T cell activation and, **41:1**
 - gene regulation, **41:27**
 - IL-1 receptor, **41:13**
 - receptor-mediated signal transduction, **41:20–22**, 24
 - synergy, **41:17**
 - T11, **41:9**
 - T cell antigen receptor, **41:2**, 3, 6, 7
 - Thy-1, **41:10**
 - T cell development and, **44:207**, 255
 - antigen recognition, **44:209**, 210, 213
 - cellular selection within thymus, **44:245–247**, 250, 251, 253
 - ontogeny, **44:215**
 - thymocyte subpopulation, **44:233**, 242
 - T cell receptor and
 - accessory molecules, **45:111**
 - alloreactivity, **45:149**, 151–156
 - antigen processing, **45:119**, 120, 125, 126
 - epitopes, **45:143**
 - experimental systems, **45:131**
 - polymorphic residues, **45:159–162**
 - structure-function relationships, **45:135**, 140, 141

- T cell subsets and
cell surface molecules, **41**:40, 49, 50
H-2 alloantigen recognition, **41**:79,
87, 94
H-2-restricted antigen recognition,
41:52, 53, 57, 59–61, 64, 66, 71, 72,
74, 75
virus-induced immunosuppression and
HIV, **45**:349–351, 366
measles, **45**:346
- Cloning
activin type II receptor cDNA, **55**:188–192
antibodies from combinatorial libraries,
57:203–206
B cells, **54**:399–400
B cell specific proteins in IgR complex,
54:348–357
CD1 genes, **59**:3–4
complement components and, **38**:209–214
cytokine genes, **58**:425
directional, **38**:205
expansion, V gene usage in, **55**:322–323
IL-2R genes, **59**:226, 227
IL-9, from human and mouse, **54**:80–82
LIF, **53**:31–32
LIF receptor, **53**:38–40
NK cells, **55**:354–357
positional, X-linked agammaglobulinemia
gene, **59**:165–169
XLA genes, **59**:165–176
- Clonotypes
murine lupus models and, **46**:78, 99
predominant, expression
in B cell subpopulations, **42**:22–26
environment and, **42**:39–42
repertoire of B cell subpopulations, *see* B
cell repertoire expression
- Clostridium* toxin, and *Pseudomonas* toxin,
54:114–115
- Clotting, aPL antibodies and
affinity purification, **49**:232
clinical aspects, **49**:200–202
pathogenic potential, **49**:253, 256–257
specificity, **49**:241, 244–245, 247
- Clustered antigens, B cell, **52**:140–164
CD5, **52**:140–144
CD9, **52**:144
CD10 or CALLA, **52**:144–145
CD20, **52**:149–150
CD22, **52**:150–151
CD23/Fc ϵ RII, **52**:151–153
CD24, **52**:153
CD37, **52**:154–155
CD38, **52**:155
CD40, **52**:155–158
CD44, **52**:158–159
CD45, **52**:159–160
CD48, **52**:160
CD72, **52**:160
CD73, **52**:160–161
CD74, **52**:161
CD76, **52**:162
CD77, **52**:162–163
CDl9, **52**:145–148
CDw32/Fc γ RII, **52**:153–154
CDw75, **52**:161–162
complement receptors: CR2, CD21, and
CD35, **52**:148–149
other clustered antigens, **52**:163–164
CD27, **52**:163
CD39, **52**:163
CD69, **52**:163
CD78, **52**:163–164
CDlc, **52**:163
Clusterin, **61**:204, 249–251
c-myb, **62**:227
B cell genesis, **63**:238–239
and *c-myc*, **62**:227
T cell activation, **48**:293–294
c-Myc, **63**:202
and thymocyte cell death, **58**:138
c-myc, **62**:227; **63**:147; **64**:232
apoptosis, **63**:162
in Burkitt lymphoma, **38**:271, 272
activation, **38**:263, 264, 271
deregulation, **38**:254–257, 260
t(14;18) chromosomal translocation,
38:269
translocation, **38**:248–253
variant chromosome translocations,
38:257–259
in Burkitt's lymphoma translocations,
50:124–128
chromosomal translocation, **64**:227–229
and *c-myb*, **62**:227
deregulated expression of, **64**:230
human B cell neoplasia and, **38**:259
activation, **38**:259–264
expression, **38**:259
translocation, **38**:271

- c-myc* (*continued*)
 IL-2 signal transduction, **63**:161–162
 immunosenescence and, **46**:223, 227, 228
 overexpression and cell death, **58**:249
 role in thymocyte apoptosis, **58**:169
 T cell activation, **48**:292–293
 T cell activation and, **41**:27, 28
- c-myc/v-Ha-ras* retrovirus,
 plasmacytogenesis, **64**:230–231
- CNTF, **65**:20
- CNTF/LIF/OSM/IL-6 subfamily, **61**:151–152
- CNTFR, **61**:151
 and high-affinity LIF receptor, **53**:43–44, 46
 pleiotropic nature of, **53**:46–47
- Coagulation
 aPL antibodies and, **49**:200, 258
 pathogenic potential, **49**:256
 specificity, **49**:239, 241–244, 246
 disseminated intravascular, endotoxic shock, **66**:260
- Coagulation/kinin-forming cascade, intrinsic,
see Intrinsic coagulation/kinin-forming cascade
- Cobra toxin, MG and, **42**:237
- Cobra venom factor, **61**:207; **62**:287–288
- Coding ends
 with P nucleotides, **58**:37–38
 and RSS ends, passage through DSB intermediates, **58**:33–36
 TCRδ, hairpin accumulation at, **58**:48
- Coding flanks, **64**:55–56
- Coding joints
 formation, microheterogeneity, **58**:39–40
 microhomology, **58**:40–42
 terminal deoxynucleotidyl transferase, **64**:51
 V-3 cells, **58**:51–52
 V(D)J recombination, **64**:40, 41–42
- Coding junctions
 and RSS junctions, **58**:42
 and *scid* mutant features, **58**:46–48
- Codominance, *Ir* genes and
 antigen processing, **38**:97
 blocking, **38**:71
 complementation, **38**:72
 gene dosage, **38**:80, 81
 genetic control, **38**:36
 immunodominance, **38**:181
 T cells 131, **38**:164
 tolerance, **38**:119, 120
- Cofactor proteins, **61**:210
- Cognate help, T cell subsets and, **41**:77
- Cognate target lysis, by CD4⁺ CTLs, **60**:304
- Coimmunoprecipitation, T cell activation and, **41**:4, 5
- Coincidental transmission, **38**:327
- Colchicine
 cytolysis and, **41**:282
 immunosenescence and, **46**:224
- Cold agglutinins, Ig heavy-chain variable region genes and, **49**:46–47, 51–52
- Cold-reactive antilymphocyte antibodies, IBD and, **42**:292
- Cold target cell competitors, hybrid resistance and, **41**:396, 402
- Coley's broth, **42**:215
- Colicins, cytolysis and, **41**:318, 319
- Colitis, *see* Ulcerative colitis
- Colitis colon-bound antibody, IgG levels in, **42**:298
- Collagen
 aPL antibodies and, **49**:197
 complement components, **38**:232, 233
 helper T cell cytokines and, **46**:128
 IL-1 and, **44**:168, 169, 186, 195
Ir genes and, **38**:61, 86
 leukocyte integrins and, **46**:153, 159
 murine lupus models and, **46**:79, 90
 surface antigens of human leukocytes and, **49**:85–86, 88
- Collagen arthritis, autoimmune thyroiditis and, **46**:312
- Collagenase, **62**:266, 288
 antigen-presenting cells and, **47**:52
 in eosinophils, **39**:203
- Colloid osmotic killing
 membrane attack complex of complement, **41**:300
 membrane damage, **41**:280
 pore formers, **41**:315, 318
 pore-forming protein, **41**:295, 312
- Colon
 genetically engineered antibody molecules and, **44**:87
 lymphocyte homing and, **44**:330, 332, 333, 338
- Colon cancer, **62**:190, 201
 properties of associated gangliosides, **40**:364–366
- Colon tumors, CD44 expression, **54**:323

- Colony-forming B cells, V_{H} utilization and, **42:133**
- Colony-forming cells
B cells and, **41:182, 191, 200, 230, 237**
hybrid resistance and
 bone marrow cells, **41:336, 344**
 leukemia/lymphoma cells, **41:359**
 NK cells, **41:376**
 T cells, **41:381, 382**
- Colony-forming unit, hybrid resistance and, **41:336, 396**
- Colony-stimulating factor α , eosinophil activation, **39:215**
- Colony-stimulating factor, murine, **48:73**
- Colony-stimulating factor 1, **48:71**
- Colony-stimulating factor gene, expression
posttranscriptional regulation, **48:83**
transcriptional regulation, **48:81–83**
- Colony-stimulating factor receptor
biochemical nature, **48:72–74**
distribution, **48:74**
- Colony-stimulating factors, **46:116; 48:69–77**,
see also Granulocyte colony-stimulating factor; Granulocyte-macrophage colony-stimulating factor
B cell formation and, **41:219, 234, 236, 237, 228–230**
biochemical nature, **48:72–74**
biological actions, **48:74–76**
cell survival, **48:75**
cellular sources, **48:76–77**
characterization, **48:70–72**
chronic deregulated production, **48:93–94**
colony-stimulating activity, **48:94**
detection in vivo, **48:94–96**
differentiation commitment, **48:75**
discovery, **48:70–72**
eosinophil, **48:92**
functional activation, **48:76**
granulocyte, **39:2–3; 48:75**
hemopoietic recovery, **48:93**
high-affinity membrane receptors, **48:73–74**
hybrid resistance and, **41:382**
macrophage, **48:75, 92**
maturation induction, **48:75**
monocyte, **39:2–3; 48:92**
multi-CSF, *see* Interleukin-3
myelosuppression, **48:93**
neutralizing antibody, **48:95–96**
- neutrophil, **48:92**
- proliferative stimulation, **48:74–75**
 target cell selectivity, **48:74–75**
- secondary structure, **48:73**
- sequence homology, **48:73**
- serum half-lives, **48:93**
- single unique gene encoding, **48:73**
- surface antigens of human leukocytes and, **49:99, 113**
- synthesis
 induction, **48:78–81**
 pathways, **48:78–81**
- T cell
 studies in vitro, **48:77–78**
 studies in vivo, **48:92–98**
- T cell synthesis, **48:96–98**
 antigen-presenting cell, **48:97**
 local polarization secretion, **48:97**
 local response, **48:96**
 target cells, **48:97**
therapeutic effects, **48:93**
- Combinatorial libraries, human antibodies
 from, *see* Antibodies, human, from
 combinatorial libraries
- Common acute lymphoblastic leukemia-associated antigen, *see* CD10
- Common γ -chain, multiple cytokine receptors, **59:239**
- Common variable immunodeficiency, **59:138, 143, 147, 150; 65:256–260**
- CD23 antigen and, **49:174**
and SCID mice, **50:308**
- Competition, autoimmune demyelinating disease and, **49:370–371**
- Competitive elimination, B cells, **59:312–315**
- Competitive inhibition, *Ir* genes and, **38:108–110, 182, 183**
- Complement, **61:201–204**
 alternative pathway
 antibodies as activators, **51:28–29**
 particulate activators, **38:384–386; 39:2**
 anaphylatoxins, **61:255**
 cascade, pulmonary inflammation, **62:260–262, 271, 286–288**
 classical pathway, history of, **37:153–154**
 membrane attack complex, **61:203, 247–255**
- molecular genetics, **38:203–207**
C1q, **38:231–233**
C2 and Factor B, **38:207–209, 228**

- Complement (*continued*)
 C3, 38:233–234
 C4, 38:219–231
 C4b-binding protein (C4bp), 38:234–236
 C9, 38:234
 cDNA, 38:209–211
 cloning, 38:212–214
 expression, 38:217, 218
 restriction fragment length polymorphism, 38:215, 216
 pathway, products, in IBD, 42:308–310
 role in B cell growth and differentiation, 52:218–219
 therapeutic inhibition in acute injury states, soluble CR1
 ARDS, 56:284–286
 ischemia, 56:282–284
 local injuries, 56:286
 range of use, 56:289
 reperfusion, 56:282–284
 tissue transplantation, 56:286–287
 therapeutic regulation in acute injury states
 activation
 clinical assessment, 56:270–271
 clinical injury, 56:278–281
 intrinsic regulation, 56:271–274
 mechanisms, 56:267–270
 ARDS, 56:278–279
 C1 esterase inhibitor, 56:272
 cardiopulmonary bypass, 56:279–280
 clinical injury, 56:278–281
 deficiency states, 56:277–278
 hemodialysis, 56:278–279
 leukocyte interaction, 56:274–277
 sepsis, 56:281–282
 septic shock, 56:281–282
 study approaches, 56:267
 therapeutic inhibition
 fungal products, 56:288–289
 heparin, 56:287–288
 thermal injury, 56:280–281
- Complement activation, *see also* Regulation of complement activation; Regulators of complement activation
 by antibody effector function, 51:7–29
 antibodies as activators of alternative pathway, 51:28–29
 C1 activation, 51:21–25
 C3 activation, 51:25–28
 C4 activation, 51:25–28
- cell lysis, 51:25–28
 interaction of C1q and antibody, 51:8–21
 segmental flexibility, 48:24–26
- Complementarity
 antibody-antigen complexes, 43:122
 induced
 antigen, 43:20–23
 differential contributions, 43:73
 evidence, 43:42–46
 protein folding, 43:83
 protein-protein interactions, 43:101–103
 structural, protein-protein interactions, 43:104–105
- Complementarity-determining region, 43:9–10, 107
 Ars A response and, 42:113, 115, 126–128
 nucleotide sequence of, 42:119
 buried surface, 43:111
 conformational changes, 43:18–20, 124
 diversity, 43:222
 forming antigen binding site, 43:13
 framework residues, 43:16
 genetically engineered antibody molecules and, 44:75, 84, 85
 Ig heavy-chain variable region genes and 34, 49:51, 61, 63
 D segments, 49:28, 31
 V_H families, 49:9, 14–15, 18
 length, 43:222
 loop
 flexibility, 43:107
 structures, 43:20
 maps of Ig-like loci and, 46:7
 mobility, 43:20
 models, 43:10
 modulating VL–VH pairing, 43:125
 movement from canonical positions, 43:127
 murine lupus models and, 46:72, 78
 sequences, 43:125
 size, 43:120–121
 spatial arrangement, 43:109
 structural variability, 43:73
 structures, 43:107–108
 T cell subsets and, 41:52
 variable domain, 43:79
 V module interface modification, 43:113, 126

- Complementary DNA
activin type II receptor, cloning, **55**:188–192
 α chain, **43**:284–286
antinuclear antibodies and
autoantibodies, **44**:127
autoantigen, **44**:128–131
autoimmune response, **44**:132
scleroderma, **44**:122
SLE, **44**:101, 108
biosynthetically engineered peptides based
constructs, **43**:281
CD4 and CD8 molecules and, **44**:297,
298
CD4 molecular biology, **44**:290, 292,
293, 296
CD8 molecular biology, **44**:272–274,
276–278
CD5 B cell and, **47**:145–147, 154
CD23 antigen and
biological activity, **49**:169, 172
cellular expression, **49**:152, 154
Fc ϵ RII, **49**:162–163, 165
CD44 constructs, transfection and
expression, **54**:286–287
clones
characterization, **43**:148
for IL-10, **56**:3–5
T cell receptor δ genes, **43**:154–155
T cell-specific, **43**:135
cloning of autoantigens
BP antigen, **53**:306–310
PF antigen, **53**:310–312
PV antigen, **53**:312–314
coding, **43**:287, 304
complement components and
C1q, **38**:232, 233
C2, **38**:209, 211, 212
C3, **38**:233, 236
C4, **38**:223–226, 228, 236
C4bp, **38**:234
C5, **38**:234
C9, **38**:234
expression, **38**:217
Factor B, **38**:209, 211, 212
from liver, **38**:209, 234
restriction fragment length
polymorphism, **38**:215, 216
complement receptor 1 and, **46**:187, 188,
191, 193, 194, 199
complement receptor 2 and, **46**:211
cytotoxicity and, **41**:319, 320
cell-mediated killing, **41**:297
hydrolytic enzymes, **41**:276
lymphotoxin-like molecules, **41**:284,
285
pore farmers, **41**:317
cytotoxic T cells and, **41**:138, 148
genetically engineered antibody molecules
and, **44**:67, 68, 73–75
for GM-CSF, **39**:9–10
helper T cell cytokines and, **46**:112, 115,
116
human LIF receptor, **53**:38–40
human-murine hybridomas, **38**:291
human T cell antigen and, **41**:3, 10, 13
humoral immune response and, **45**:63, 66,
72, 74, 75
IgE biosynthesis and, **47**:16–18, 21
Ig gene superfamily and, **44**:33–35, 39, 40
Ig heavy-chain variable region genes and,
49:19, 22, 33
 V_H gene expression, **49**:35, 37–38, 44–45
IL-1ra clones, purification and expression,
54:171–176
IL-2 and, **44**:155
for IL-3, **39**:5, 7–8
IL-5, organization, **57**:148–149
IL-10, **56**:3–5
Ir genes, **38**:116, 117, 148
Ir genes and, **38**:117
leukocyte integrins and, **46**:154–158, 160,
174, 176
maps of Ig-like loci and, **46**:34, 43
maternally transmitted antigen, **38**:348
murine lupus models and, **46**:83, 87, 95
NK cells and, **47**:209
normal and mutant in PNH, **60**:83–84
regulators of complement activation and
genes, **45**:403, 405
protein expression, **45**:396, 398, 399
short consensus repeat, **45**:391
spontaneous autoimmune thyroiditis and,
47:462, 483
surface antigens of human leukocytes and,
49:79, 114, 118
T cell receptor 7, **43**:145
T cell receptor and
in peripheral T cells, **38**:13
protein, **38**:4, 6, 8, 9

- Complementary DNA (*continued*)
 receptor-like genes, **38:10**
T3, 38:18
 in thymus, **38:12**
- T cells, **38:10**
 chromosomal mapping, **38:16**
 peripheral, **38:13, 14**
 rearrangement, **38:11, 12**
 receptor proteins, **38:5, 6, 8, 9**
 V region genes, **38:15, 16**
- T cell-specific, **43:135, 220**
- T cell subsets and, **41:41, 50**
- transfection experiments, **42:168**
- virus-induced immunosuppression and, **45:349–351**
- X-linked agammaglobulinemia, **59:165–167**
- Complementation, MHC-linked *Ir* genes and, **38:72–84**
 H-Y antigen, **38:177, 178**
 I region mutations, **38:85, 88**
- T cells
 interactions, **38:160**
 repertoire, **38:133**
 suppression, **38:140, 145, 148, 149**
- Complementation classes, mutants, biochemical defects, **60:66–67**
- Complement binding activity
 hinge region structure, **48:13, 15, 22**
 segmental flexibility, **48:13, 15, 22**
- Complement control, **61:255–257**
 C3/C5 convertases, **61:206–247**
 initiation, **61:201–206**
- Complement control protein repeats, **61:207–208**
- Complement deficiency alleles, aPL antibodies and, **49:222**
- Complement fixation, by Fc region of Ig, **40:70–71**
- Complementing binder sites, **52:375–377**
- Complement lysis sensitivity test, **60:75**
- Complement-mediated cytotoxicity, *see* Cytotoxicity, lymphocyte and complement-mediated
- Complement receptors
 C3b and C4 on eosinophils, **39:179–180**
 C5a, gene mapping, **55:135**
CR1, 46:183, 208, 209; 60:182; 61:228–230
 allotypes, **46:187–190**
 aPL antibodies and, **49:255**
- and CR2 and CD19, molecular complexes on surface of human B cells, **52:146**
- function
 endocytosis, **46:198–200**
 erythrocytes, **46:198**
 immunoregulation, **46:200, 201**
 ligand binding sites, **46:193–196**
 regulation, **46:196, 197**
- soluble, in therapeutic complement inhibition, **56:284–286**
 ischemia, **56:282–284**
 local injuries, **56:286**
 range of use, **56:289**
 reperfusion, **56:282–284**
 tissue transplantation, **56:286–287**
- structure, **46:183–187**
- in therapeutic complement activation, **56:274–277**
- tissue distribution, **46:190–193**
- CR2, **46:183; 61:228, 230**
 and CR1 and CD19, molecular complexes on surface of human B cells, **52:146**
- function, **46:205**
 EBV tropism, **46:210–213**
 immunoregulation, **46:205–210**
- structure, **46:201–203**
 tissue distribution, **46:203–205**
- CR2, CD21, and CD35, **52:148–149**
- CR3, **46:209, 210; 60:182–183**
 leukocyte integrins and, **46:150, 163, 164**
 leukocyte interactions, **56:276–277**
- regulators, **61:228–230**
- surface antigens of human leukocytes and, **49:91–97**
- Complement regulatory proteins, **61:201–204, 202, 203–204**
- Complete Freund's adjuvant, *see* Freund's complete adjuvant
- Conalbumin
 humoral immune response and, **45:37**
 IgE biosynthesis and, **47:11**
- Concanavalin A, **48:90, 233**
 activated spleen cells, T cell subsets and, **41:64, 65, 70**
- antigen-presenting cells and, **47:84, 100**
 autoimmune thyroiditis and, **46:281**
 autoreactive T cells and, **45:421**
- B cell formation and, **41:198, 229**

- B cell proliferation, **63:57**
 CD5 B cell and, **47:121**
 humoral immune response and, **45:45**
 hybrid resistance and, **41:375, 409**
 IBD and, **42:299**
 IgE biosynthesis and, **47:15, 18–20**
 IL-1 and, **44:173, 174, 182, 189**
 immunosenescence and
 lymphocyte activation, **46:223, 225, 226, 228–230**
 mucosal immunity, **46:252**
 stem cells, **46:245**
 murine lupus models and, **46:98**
 pore-forming protein and, **41:313**
 spontaneous autoimmune thyroiditis and,
 47:492
 cellular immune reactions, **47:452–455**
 disturbed immunoregulation,
 47:470–473, 475, 477–479
 genetics, **47:487, 488, 490**
 potential effector mechanisms, **47:467**
 T cell activation and, **41:8, 28**
 T cell development and, **44:243**
 T cell subsets and, **41:63–65**
 virus-induced immunosuppression and,
 45:357, 362
- Conditioning, allergic reactions, **61:368–369**
 Congenic strain, cross-reactive idiotype
 expression and, **42:141**
 Congenital agammaglobulinemia, **59:138–139**
 Congenital myasthenic syndrome,
 42:252–253, *see also* Myasthenia gravis
 Conjugate formation
 cell mediation and, **41:272**
 cytosis and, **41:274–276, 278, 282, 283**
 Connective tissue
 cachectin and, **42:225**
 IFN- γ , **62:90–91**
 Connective tissue-activating peptide III,
 50:240–243
 biological effects on neutrophils, **55:115**
 chemical properties of, **55:97, 99**
 comparison with histamine releasing factor,
 50:242
 conversion of to NAP-2, **50:242–243**
 in T cell development in MHC class I-
 deficient mouse, **55:399–401**
 Consensus sequences
 CD23 antigen and, **49:162**
- Ig heavy-chain variable region genes and,
 49:30
 Constant region, heavy chain, gene order and
 class switching, **40:257–258**
 Constant region complex, Ig heavy-chain
 variable region genes and, **49:3–4, 6**
 Contact activation, **66:225, 233–239**
 allergic diseases and, **66:258–259**
 bradykinin, **66:225, 226, 233–240**
 defined, **66:225**
 disease states and
 allergic diseases, **66:258–259**
 Alzheimer's disease, **66:260–261**
 endotoxic shock, **66:260**
 hereditary angioedema, **66:257–258**
 pancreatitis, **66:260**
 rheumatoid arthritis, **66:260**
 factor XI, **66:225, 226, 230, 237, 244**
 factor XII, **66:225, 226–229, 234–236, 244, 246, 249–257**
 high-molecular-weight kininogen,
 66:225, 226, 230–233, 245, 247–257
 interaction with cells and, **66:258–259**
 kallikrein, **66:225, 238, 243–244**
 low-molecular weight kininogen, **66:233**
 mechanism, **66:225–226**
 prekallikrein, **66:225, 226, 229–230, 236**
 regulation, **66:235–239**
 Contactin, **48:161–162**
 Contact residue, **43:26, 72, 119**
 Converter protein, **42:168**
 Coreceptor model, of CD4 and CD8
 and adhesiveness during TCR triggering,
 53:83–84
 coreceptor/MHC binding sites, **53:68–70**
 cytoplasmic tail role in signal mediation,
 53:73–74
 description, **53:65–68**
 effects on positive vs. negative selection,
 53:103–104
 function, gene transfer for, **53:70–78**
 p56^{lck} role in TCR-mediated signaling,
 53:74
 TCR-coreceptor simultaneous binding,
 53:66–68
- Coreceptors
 T cell, recognition/auxiliary, **51:91–92**
 TCR/CD3, signalling through, **51:92–98**
 Cornea, MHC antigen, **48:193–194**

- Corneal allograft
graft bed
 afferent blockade theory, **48**:195–196
 avascular, **48**:195
 graft location, **48**:195
 host cellular component, **48**:195
 orthotopically placed, **48**:195
 immunological privilege, **48**:192–203
- Corneal endothelial graft, cytotoxic T cell, **48**:194
- Corneal epithelial cell, accessory antigen-processing cell, **48**:202
- Corneal epithelial cell-derived T cell-activating factor, **48**:201–202
- Corneal epithelium, Langerhans cell, **48**:197
- Corneal graft
 Ia antigen, **48**:199
 MHC class II antigen
 cell-surface expression, **48**:199
 expression, **48**:199
 expression timing, **48**:199–200
 MHC class II loci, **48**:199–200
 latex bead treatment, **48**:200
 rejection, donor Langerhans cells, **48**:196–200
- Corneal transplantation, **48**:191
 HLA matching, **48**:193
 immunological recognition, **48**:193–196
 immunosuppressive drug, **48**:193
 success, **48**:193
- Coronary artery bypass grafting, aPL
 antibodies and, **49**:220, 222–223
- Coronary heart disease, dietary fish oil and, **39**:145–146
- Coronavirus, neutrotrophic, **48**:141
- Cortical cells, loss of responsiveness in, **51**:153–156
- Corticosteroid-binding globulin, **47**:492, 494
 genetics, **47**:488–490
 immunoregulation, **47**:476, 479, 481
- Corticosteroid resistant bronchial asthma, **51**:358–361
- Corticosteroids, **42**:311
 adrenal, MG therapy and, **42**:270–271
 B cell formation and, **41**:197
 and estrogens, **54**:50
 IBD and, **42**:316–319
 IL-1 and, **44**:160, 161, 182, 186, 189, 190
 MG therapy and, **42**:270–271
- mode of action, **64**:195–196
prostaglandin synthesis and, **42**:312–313
spontaneous autoimmune thyroiditis and, **47**:477
- Corticosterone, spontaneous autoimmune thyroiditis and, **47**:493
disturbed immunoregulation, **47**:476–481
genetics, **47**:488, 489
- Corticotropin-releasing factor
 in vitro IgE synthesis, **61**:370–372
 lymphocyte-produced, **39**:316
 NK cells and, **47**:268, 269
 spontaneous autoimmune thyroiditis and, **47**:493
- Corticotropin-releasing hormone, **48**:170–172
- Cortisone, hybrid resistance and, **41**:338, 359, 360
- Corynebacterium parvum*
 B cell formation and, **41**:208
 hybrid resistance and
 bone marrows, **41**:340, 347
 macrophages, **41**:370
 marrow microenvironment, **41**:391, 392
 syngeneic stem cell functions, **41**:389
- Cosignals, inactivation of death programs, **58**:243–245
- Cosmid clones
 complement components and, **38**:213
 C2, **38**:218
 C4, **38**:228
 Factor B, **38**:212, 218
 Ig heavy-chain variable region genes and, **49**:5–6, 8
- Cosmids, maps of Ig-like loci and, **46**:4, 5
 MHC, **46**:29, 32, 33
 structure, **46**:17, 18, 27
 T cell receptor, **46**:36, 37
- Costimulation, **62**:132
 B7-1 and B7-2, **62**:140–142
 CD28 and CTLA-4, **62**:138–139
 CD28-B7, **62**:132
 Td B cell response, **62**:148–157
 in vivo immune response, **62**:142–144
- Costimulator activities, vascular endothelial cells, **50**:267–274
- EC costimulators
 in alloantigen responses, **50**:271–272
 in polyclonal CD4+ T cell activation, **50**:268–270

- in polyclonal CD8+ T cell activation, **50**:270–271
soluble, EC production of, **50**:272–274
Costimulators, soluble, EC production of, **50**:272–274
Costimulus, **59**:280, 325–326
Coupled cleavage, V(D)J recombination, **64**:53–54
Coupling proteins, *see also G proteins*
G_i-receptor interaction in neutrophils, **39**:114–116
amplification during transduction and, **39**:116–117
model of, **39**:115–116
termination of, **39**:117–118
G_s
–catecholamine interaction, **39**:112–113
guanyl nucleotide binding, **39**:112
neutrophil activation and, **39**:113–114, 118
in retina, rhodopsin and, **39**:112
stimulatory and inhibitory, **39**:112–113
subunit activities, **39**:112
CpG islands, **65**:50, 51
CRAF1 (CD40 receptor associated factor 1), **61**:29–31
C-reactive protein, surface antigens of human leukocytes and, **49**:111
CREB, interaction with NF-IL6, **65**:19
C-REL knockout mice, cytokine induction, **65**:31–32
CREST, antinuclear antibodies and, **44**:120, 122, 131
CRF, *see Corticotropin-releasing factor*
Critical residue, Ab binding to protein antigens, **43**:27, 72
amino acids, **43**:67–68
antigen binding, **43**:61, 63
antiprotein antibodies, **43**:42
buried, **43**:42–43
identification, **43**:39–42
peptide mapping, **43**:74
properties, **43**:40–41
sequential epitope, **43**:64
stereochemical relationships, **43**:43
Crohn's disease, **59**:146, *see also Inflammatory bowel disease*
6-MP therapy and, **42**:300
anti-TNF- α therapy, **64**:307–310
granulomas in, **42**:297
helper T cell function and, **42**:300, 301
IgG response to infectious agent in, **42**:305
infections with cA2 treatment, **64**:326–328
intestinal MNC and, **42**:303–304
K cell function and, **42**:295
lymphocytes 290, number, **42**:297
skin test antigens and, **42**:289
submucosa of specimens of, **42**:298
Cromolyn-binding protein, **43**:297
Cross-hybridization
CD23 antigen and, **49**:162
Ig heavy-chain variable region genes and, **49**:1, 8
Crosslinking
CD23 antigen and, **49**:156, 168–169, 172–173
cell mediation and, **41**:272
FC ϵ RI
and factor-dependent mast cell lymphokine production, **53**:6
and IL-3 regulation of lymphokine production, **53**:21–22
and IL-4 production by splenic and bone marrow Fc ϵ RI⁺ cells, **53**:10
and signaling of IL-4 production, **53**:19–22
and transformed mast cell lymphokine production, **53**:5
photochemical, for lipid A receptor identification, **53**:277–279
surface antigens of human leukocytes and, **49**:89–90, 101, 111, 125
T cell activation and, **41**:4, 5, 9, 11, 12, 25
T cell subsets and, **41**:62, 111
Cross-priming, T cell subsets and, **41**:74
Cross-reactive idiotype, **42**:11
Ars A response and, **42**:97–99, 109–110, 139–141
autoimmune thyroiditis and, **46**:303–305, 308
CD5 B cdl and, **47**:150, 151, 153–158
Ig heavy-chain variable region genes and, **49**:43, 46, 48–49, 51
secondary B cell lineage and, **42**:70–71
Cross-reactivity
aPL antibodies and, **49**:209, 211, 236, 243, 248–249
Cytotoxic T cells and, **41**:144, 147, 162, 170
hybrid resistance and, **41**:357

- Cross-reactivity (*continued*)
 Ig heavy-chain variable region genes and, **49:47**
 membrane attack complex of complement and, **41:307, 309**
 T cell subsets and, **41:79, 88, 112**
- Cryptococcus neoformans*, pneumonia, **59:402–408**
- Crystallographic temperature factors, mobility data from, **43:20**
- Crystallography, reconciling data, **43:23–27**
- Crystal packing, Fab Kol, **43:19–20**
- Crystal studies, two-dimensional, major features, **51:18**
- Csa effects, development of autoimmune phenomena, **50:209–210**
- CSF, *see* Colony-stimulating factors
- Csk, **62:42, 45**
- CT-1, **64:256**
- CTL, *see* Cytotoxic T cells
- CTLA-4, **62:132–135**
 costimulatory effects on immunogenicity, **57:305–307**
 T cell receptor for B7, **58:134**
- CTLA4Ig, **62:138, 141, 143, 149, 150**
- Culture supernatants, CD23 antigen and, **49:157, 159**
- Curare, MG and, **42:237**
- Curcumin, action of, **65:123**
- Cutaneous diseases
 eosinophil degradation and, **39:225–226**
 major basic protein extracellular deposition, **39:225–226**
- Cutaneous lymphocyte antigen, **64:146, 151**
 chronic inflammation and, **64:190**
 properties, **64:146**
- CVID, *see* Common variable immunodeficiency
- CX5a, **62:261**
- C-X-C family, **62:269–270**
- CXCR4, **65:283; 66:282**
- Cyanuric chloride, linkage procedure for chemoimmunoconjugates in cancer treatment, **56:321**
- Cyclic AMP
 in activated neutrophils, **39:113–116, 118**
 autoimmune thyroiditis and, **46:268, 272**
 CD40 expression, **61:98**
 IL-1 and, **44:160, 161, 195**
- immunosenescence and, **46:223, 226, 227, 232**
 immunosuppressive activity of, **65:124–125**
 intracellular levels, elevation, **54:379**
 NK cells and, **47:249, 250**
- Cyclic AMP response element modulator τ , **61:189**
- Cyclic AMP response elements, **62:194**
- Cyclic GMP, immunosenescence and, **46:223**
- Cyclic neutropenia, B cell formation and, **41:222, 228–230, 234, 237**
- Cyclic nucleotides, immunosenescence and, **46:226, 227, 232**
- Cyclin-cdk complexes, **61:183, 184**
- Cycloheximide, **42:174**
 antigen-presenting cells and, **47:59**
 effect on apoptosis induction, **58:265–267**
- Cyclooxygenase
 active site, PGHS-2, **62:174–176**
 cachectin and, **42:219**
 IBD and, **42:317**
 IL-1 and, **44:160, 168, 195**
 isozymes
 catalysis by PGHS, **62:178–188**
 cellular and physiologic action of PGHS, **62:196–200**
 future work, **62:201–202**
 pathophysiologies, **62:200–201**
 regulation of gene expression, **62:188–196**
 structure-function relationships, **62:169–178**
- NO role, **60:349–350**
- NSAIDs and, **42:313**
- Cyclophosphamide
 adoptive T cell therapy of tumors and, **49:293, 295–297, 302–303, 317**
 diabetes induced by, **62:100–101**
 hybrid resistance and
 bone marrow cells, **41:347, 348**
 effector mechanisms, **41:370, 388, 390, 393**
 leukemia/lymphoma cells, **41:366, 368**
 lymphoid cells, **41:354**
- MG therapy and, **42:263–264, 271**
- Cyclosporin, **41:60, 350; 59:424**
 autoimmune thyroiditis and, **46:294**
 effect on thymocyte death, **58:173**
 mAbs and, **38:277, 286**
 treatment, **50:104–105**

- Cyclosporin A
autoimmune side effects, **58:278–279**
and autoimmunity, **50:208–210**
blockade of CD4⁺8⁺ thymocytes, **58:267**
CD40-L expression, **61:21**
effect on CD40L expression, **60:47–48**
effect on PCD induction, **58:261–262**
GM-CSF in T cells and, **39:14–15**
humoral immune response and, **45:47**
IL-2 in T cells and, **39:14–15**
IL-3 in T cells and, **39:14–15**
inhibition of B cells, **54:408**
for MG, **42:264**
NF-κB, **65:120**
suppressor T cells and, **42:258–259**
T cell activation and, **41:29, 30**
and thymic T cell maturation, **58:314–315**
- CYP2D5 gene, **65:23**
- Cysteine
autoimmune thyroiditis and, **46:316**
CD4 molecules and, **44:285–287**
CD8 molecules and, **44:273, 274, 276, 277**
CD23 antigen and, **49:163–164**
α chain of IgA, **40:155, 163**
complement receptor 1 and, **46:184, 194**
complement receptor 2 and, **46:202**
cytotoxic T cells and, **41:148, 163**
genetically engineered antibody molecules
and, **44:81, 82**
IgE biosynthesis and, **47:17**
J chain of IgA, **40:160**
leukocyte integrins and, **46:154, 155**
maps of Ig-like loci and, **46:2, 13**
membrane attack complex of complement
and, **41:307, 309**
relationship between loops, Ig domains,
43:285
residues, **43:137, 156**
secretory component of IgA, **40:162–163**
surface antigens of human leukocytes and,
49:116
- Cytocalasin B, cytolysis and, **41:282**
- Cytochrome, secondary B cells and, **42:71–72**
- Cytochrome c
acetimidylated, **43:200**
as antigen, **43:198–200**
autoreactive T cells and, **45:419**
beef, **43:199**
cross-reactive, **43:202**
immune response, **43:224**
- Ir* genes and, **38:77–79, 82, 89**
antigen binding, **38:98–107**
antigen processing, **38:94, 96**
competitive inhibition, **38:108**
gene dosage, **38:82**
T cells
activation, **38:104**
repertoires, **38:116, 119, 132–134**
- moth, **43:203**
- mouse, **43:199**
- pigeon, **43:199–200**
- secondary structure, **43:204**
- sequence differences in, **43:4**
- synthetic T and B cell sites and
bacterial antigens, **45:240**
globular protein antigens, **45:203, 204,**
210, 211, 214
immunological considerations, **45:201**
- T cell receptor and
antigen processing, **45:121, 127**
epitopes, **45:143**
experimental systems, **45:131, 132**
structure-function relationships, **45:135,**
136
- T cell subsets and, **41:53**
- Cytochrome P450, **65:23**
- Cytofluorographic analysis, two-color, T cell
receptor γδ proteins, **43:146–147**
- Cytofluorometry, IBD and, **42:295**
- Cytokine–endotoxin score, **66:123**
- Cytokine–hemopoietin receptors,
63:393–394, 397
- Cytokine-modulating therapy, **66:151**
- Cytokine receptors, **63:269–270, 273, see also**
specific receptors
- B cell, **52:169–173**
IL-2, **52:171–172**
IL-4, **52:172**
IL-6, **52:172**
IL-1, **52:169–171**
other cytokines, **52:173**
TGF-β, **52:173**
TNF-α, **52:172–173**
- common γ-chain for multiple receptors,
59:225–267
- disease and, **63:310–314**
- disease models, **63:312–314**
- IL-2R, **63:296–299**
- IL-6R, **63:302**
- TNF-R, **63:306**

- Cytokine receptors (*continued*)
- family, **52**:170
 - HLA class II molecules, **52**:164–166
 - human, **52**:171
 - human basophils, mast cells, KU-812 cells and HMC-1 cells, **52**:340, 341
 - IL-1R type I, **63**:312–313
 - membrane-bound form, **63**:273–274, 279–280
 - membrane proximal domains, role in Jak association, **60**:11–12
 - nonreceptor proteins, **63**:277–278
 - other antigens of B cells, **52**: 173–174
 - Bgp 95, **52**:173
 - CK226 antigen, **52**:173
 - IgM-binding protein (Fell receptor, **52**:173
 - receptor antagonists, **63**:277
 - soluble forms, **63**:275–276, 279–280, 314–315
 - GM-CSF-R, **63**:308
 - IFN, **63**:307–308
 - IFN- γ -R, **63**:307, 309–310, 314
 - IL-1R, **63**:290–292
 - IL-1R type II, **63**:310
 - IL-2R, **63**:292–294, 296–298
 - IL-4R, **63**:281–282, 284, 285, 294–295, 299, 313
 - IL-5R, **63**:275, 299–300
 - IL-6R, **63**:275, 280, 282, 300–303
 - IL-7R, **63**:303
 - immunoregulatory function, **63**: 285–290
 - production, **63**:280–285, 314–315
 - TNF-R, **63**:303–307, 313–314
 - TNF-R type II, **63**:309
 - viral, **63**:308–310
 - structure, **55**:274
 - TGF- β regulation, **55**:187
 - as therapeutic agents, **63**:311–312
- Cytokine release syndromes, IFN- γ and, **62**:103–104
- Cytokine response element, **65**:20
- Cytokines, **42**:165; **48**:70, 71; **62**:195; **63**:127, 269, *see also* Chemokines
- activation of Jak and Stat isoforms, **61**:174
 - acute effects, **48**:174–175
- adoptive T cell therapy of tumors and, **49**:285, 289–290, 308
- expression of antitumor responses, **49**:324, 326, 330, 332
- mechanisms, **49**:309–310
- alarm, **58**:394–395
- antigen-presenting cells and, **47**:47, 104
- antigen presentation, **47**:72
 - cell surface, **47**:63
 - immunogenicity, **47**:103
 - T cell growth, **47**:100
 - tissue distribution, **47**:53
- antiinflammatory, *see* Antiinflammatory cytokines
- aPL antibodies and, **49**:257
- and autoimmune diabetes, **51**:307
- autoimmune thyroiditis and, **46**:275, 298, 307, 317
- B cell differentiation, **52**:204–218
- activated T cells, **52**:212–213
 - antigen receptor-dependent activation, **52**:204–206
 - IL-2, **52**:204–205
 - IL-4, **52**:205–206
 - other stimulatory cytokines, **52**:206
- CD40-dependent activation, **52**:206–212
- general considerations on, **52**:215–217
- IL-4 and IgE switching, **52**:217–218
- IL-6 and autoimmune system, **52**:215
- in vivo-activated* plasmablasts, **52**:213–215
 - isotype switching of human B cells, **52**:215–218
- B cell growth, **52**:195–204
- activated T cells, **52**:203
 - antigen receptor dependent activation, **52**:195–200
 - antigen-dependent activation, **52**:199
 - polyclonal activators, **52**:195–196, 195–199, 196–198, 198–199, 199
- CD40-dependent activation, **52**:200–203
- IL-6 and B cell growth, **52**:203–204
- B cell growth and differentiation, **52**:174–185
- IL-2, **52**:177–178
 - IL-4, **52**:178–181
 - pleiotropic effects of, **52**:179–181
 - sources and structure of and its receptor, **52**:178–179

- IL-6, **52**:182–184
 pleiotropic effects of transforming growth factor β , **52**:184
 sources and structure of transforming growth factor β and its receptor, **52**:183–184
- IL-10, **52**:181–182
 pleiotropic effects of, **52**:181–182
 sources and structure of, **52**:181
- low molecular-weight bacillus Calmette Guerin factor (BCGG-12 kDa), **52**:185
- notions of TH1 and TH2 CD4+ helper T cells, **52**:175–177
- B cell production, **61**:33, 47
- B cell proliferation, **63**:54–55
and B cells, a synthesis, **52**:227–230
- binding, coupled to tyrosine phosphorylation, **60**:4–9
- binding activity, **63**:285–290
- biological activity, **59**:254
- CD1 expression, **59**:64–65
- CD5 B cell and, **47**:120
 marker for activation, **47**:127
 physiology, **47**:139, 142
 surface antigen, **47**:145
- CD23 antigen and, **49**:159, 169–170, 172
and chemicals inhibiting IL-6 synthesis, **54**:50–51
- circulating levels, measurement, **66**:104–105
- complement receptor 1 and, **46**:192, 199
- disease and, **63**:270, 310–311
- downregulation, **64**:315–316
effects on IL-1 α production by monocytes, **54**:182–184
- enhancement and inhibition of iNOS, **60**:333–339
and eosinophils, **60**:203–205
- fibrosis, **59**:418
- function, **66**:103–104
- functional effects, **48**:167–179
- gene regulation
 description of, **52**:263
 interferon system, **52**:264–265
 by NF-IL6 and NF- κ B, **65**:1–33
 regulatory DNA sequences in type I interferon genes, **52**:265–267
- transcriptional regulation of IFN- γ
 inducible genes, and role of IRF-1 and IRF-2, **52**:272–275
- transcription factors binding to regulatory elements of type I interferon genes
- IFN regulatory factors 1 and 2, **52**:267–271
- NF- κ B, **52**:271–272
other factors, **52**:272
- understanding, **52**:263–264
- genes
 cloning and expression, **58**:425–427
 in tumor cell engineering, **57**:321–323
 upregulated by NF- κ B, **58**:10–12
- germline transcripts, **61**:84–89
- granulomatous lesions, **62**:281
- growth hormone secretion, **63**:385
- helper T cell, *see* T helper cells, cytokines and histamine release, **50**:247–248
- HIV infection and, **47**:401–404
- human, properties, **52**:175
- humoral immune response and, **45**:1, 55–61, 73
- Ig production, **63**:58
induced by NF- κ B, **58**:10–12
- induction of, **51**:274
- inflammatory process, **65**:111
- inflammatory response, **63**:310
- inhibition of viral replication, **66**:288, 290
- Jak isoforms, **61**:174
- Jak kinases, **63**:138–139
- leukocyte integrins and, **46**:170, 172, 176
- LPS-induced release, CD14 role, **53**:284
- lung transplant rejection, **59**:424, 426
- lymphocyte–HEV interaction, **65**:374
and mast cells, **51**:328–331
- membrane receptors and, **63**:274–275
- modulation of histamine release by, **50**:248–250
- mRNA expression, Th2-like profile, **60**:222
- multiple myeloma pathogenesis and, **64**:252–260
- network, position of IL-2 in, **50**:160
- neural generation, **48**:168–169
- NK cells and, **47**:241, 263, 265, 266
- persistent effects, **48**:173–174
- pleiotropy of, **53**:47
- produced by different vector systems, **58**:429–430
- production, **48**:167–179
 by CD5 B cells, **55**:310–311
 by human CD4+ T cells, **61**:347–355
 TGF- β regulation, **55**:187

- Cytokines (*continued*)
 proinflammatory, **64**:291–294
 IL-6 effects, **54**:27–28
 pulmonary immunity, **59**:373
 pulmonary inflammation, **62**:267–271, 280,
 288
 receptor antagonists, **63**:277
 redundancy of, **53**:47
 regulation, **63**:270–275, 310–311
 regulation of apoptosis, **58**:229–230
 regulatory effects, **64**:165–166, 315–317
 rheumatoid arthritis, **64**:290–298, 316;
 65:123–124
 neovascularization, **64**:320
 role in peripheral clonal deletion,
 58:271–272
 role in regulating committed progenitors,
 60:163–164
 SCID and, **49**:386, 393
 sepsis and, **66**:101–102, 104
 G-CSF, **66**:149–150
 GM-CSF, **66**:150
 IFN- γ and, **66**:145–146
 IL-1, **66**:104, 117–127, 150
 IL-4, **66**:144–145
 IL-6, **66**:130–135
 IL-8, **66**:135–141
 IL-10, **66**:142–145
 IL-12, **66**:146–147
 LIF, **66**:147–148
 MCP-1, **66**:104, 135, 141–142
 MIF, **66**:147–148
 TNF, **66**:104, 117–127, 150
 signaling, **58**:369
 negative regulation, **60**:25–26
 role of Stats, **60**:19–22
 shared signals through GP130,
 54:15–19
 signal transduction, **63**:127–176
 adapter molecules, **63**:134, 141–143
 calcineurin, **63**:166–167
 CD45, **63**:164–165
 Jak–Stat pathway, **63**:138–141
 MAPK family, **63**:148–149
 nuclear transcription factor NF- κ B,
 63:157–160
 PAC-1, **63**:167
 phosphatidylinositol 3-kinase,
 63:149–154
- phosphoprotein phosphatases,
 63:164–167
 protein kinase C, **63**:154–157
 protein tyrosine kinases, **63**:133–143
 PTP-1C, **63**:165–166
 Raf kinase, **63**:147–148
 Ras, **63**:143–147
 Rho protein, **63**:149
 Src family, **63**:133–137
 SYK family, **63**:134, 137–138
 spontaneous autoimmune thyroiditis and,
 47:494
 disturbed immunoregulation, **47**:471,
 477, 478
 genetics, **47**:488, 489
 Stat protein activation, **61**:174–175
 stimulatory, **52**:198–199
 surface antigens of human leukocytes and,
 49:98–100, 111
 synthesis and secretion, **63**:271, 273
 synthesis coordination by NF- κ B, **58**:1–2
 TH2, **60**:177
 functional similarities to IL-10, **56**:12–13
 TH responses, **61**:347–348
 Cytokine synthesis inhibitory factor, **46**:131,
 133, 135; **62**:85, *see also* Interleukin-10
 discovery, **56**:1–2
 Cytokinoplasts, L-NMMA effect, **60**:352
 Cytolysis, cell-mediated killing and, **41**:291
 Cytolysis
 adoptive T cell therapy of tumors and
 antigen recognition, **49**:319
 expression of antitumor responses,
 49:325, 328, 332
 mechanisms, **49**:299–303, 305–308,
 312–313, 316
 principles, **49**:290, 295
 af27 polarity, **41**:273–275
 cell mediation, **41**:272
 cognate target, by CD4 $^{+}$ CTLs, **60**:304
 colloid osmotic killing, **41**:280
 cytoplasmic granules, **41**:287, 288, 291
 granule exocytosis, **41**:281, 282
 HIV-1-specific, **66**:283–284, 287–298
 HIV-infected cells, **65**:280–281
 hydrolytic enzymes, **41**:273–275
 intracellular damage, **41**:282–284
 leukoregulin, **41**:286
 lymphotoxin-like molecules, **41**:284–286

- membrane attack complex of complement, **41**:299, 300, 310
membrane damage, **41**:277–281
NK cell-mediated, p58 in, **55**:354–357, 362–363
perforin role, **51**:215–216, 224–232
perforin-independent mechanisms, **51**:231–232
verification by genetic approach, **51**:227–231
verification in cell lines, **51**:224–227
polypeptide toxins, **41**:316–319
pore-forming proteins amoebas, **41**:312, 313
cytolytic proteins, **41**:316–319
eosinophil cationic protein, **41**:311, 312
polypeptide toxins, **41**:316–319
small peptides, **41**:313–316
reactive, inhibition by CD59, **60**:76–77
reactive oxygen metabolism intermediates, **41**:286
TCR $\gamma\delta$ cells, **43**:172–174
Cytolysis inhibitor, **61**:204, 249–251
Cytolytic effector cells, NK cells and, **42**:204–205
Cytolytic T cells, **41**:365, 367, 368; **42**:165
CD5 B cell and, **47**:144, 145
clones, **41**:9, 12, 14
CTLL, **42**:165
CTLL-2, **61**:155
IL-2 activation, **61**:176–179
rapamycin, **61**:182
hybrid resistance and, **41**:364, 380, 395, 396, 402, 404
Cytomegalovirus
cytotoxic T cells and, **65**:278–279
HIV infection and immune response, **47**:409
immunopathogenic mechanism, **47**:390, 392, 400
neuropsychiatric manifestations, **47**:392, 403, 405
human, antibodies from combinatorial libraries, **57**:220–221
Cytoplasm
adoptive T cell therapy of tumors and, **49**:283
antigen-presenting cells and
 antigen presentation, **47**:81
 antigen processing, **47**:87
cell surface, **47**:60
T cell growth, **47**:101
tissue distribution, **47**:48, 51
autoimmune thyroiditis and, **46**:295
CD5 B cell and, **47**:144–146
CD23 antigen and, **49**:162, 166
complement receptor 1 and, **46**:184, 188–190
complement receptor 2 and, **46**:202, 209
HIV infection and, **47**:393, 396, 397
IgE biosynthesis and, **47**:4, 20, 21
leukocyte integrins and, **46**:154–156, 166, 171
NK cells and, **47**:197
congenital defects, **47**:225
effector mechanisms, **47**:249, 253, 262
morphology, **47**:214
SCID and, **49**:386
surface antigens of human leukocytes and, **49**:87, 89, 117
membrane enzymes, **49**:113–114
receptors, **49**:98–100
Cytoplasmic domains, **54**:281–282
IL-2R β , **59**:255–258
IL-2R γ , **59**:254–255, 258
LIF receptor, **53**:39–40
regulation through, **54**:311–313
role of phosphorylation, **54**:311–313
Cytoplasmic granules, cytotoxicity and, **41**:281, 287–291
Cytoplasmic tails, CD1 proteins, **59**:17
Cytoskeletal interactions, **54**:346
Cytoskeleton
attachment to B cell antigen receptor, **55**:262–264
autoimmune thyroiditis and, **46**:298
complement receptor 1 and, **46**:186, 198
complement receptor 2 and, **46**:210, 213
immunosenescence and, **46**:224, 235
leukocyte integrins and, **46**:163, 169, 171, 176
surface antigens of human leukocytes and, **49**:85–86, 92–93, 113
Cytosol
IL-1 and, **44**:161, 163, 175, 176, 179, 184
TAP system, **64**:5–6
T cell activation and, **41**:17, 22
Cytotoxic effector cells
IBD and, **42**:296–297
immune reconstitution and, **40**:416–417

- Cytotoxicity, **43**:173
 ADCC, *see* Antibody-dependent cellular cytotoxicity
 allogeneic lymphocyte, hybrid resistance and, **41**:356; 357
 CTL-mediated
 Fas/APO-1 pathway, **60**:294–296
 perforin pathway, **60**:292–294
 role of effector molecules, **60**:300–302
 and eosinophil mediator release, **60**:205–208
 hybrid resistance and, **41**:359, 361, 362, 366, 372, 393, 394
 lectin-mediated, **60**:300
 lymphocyte and complement-mediated, **41**:269, 270, 319, 320
 af27 polarity, **41**:273–275
 colloid osmotic killing, **41**:280
 CTL cells, **41**:270–273
 granule exocytosis, **41**:281, 282
 hydrolytic enzymes, **41**:275–277
 intracellular damage, **41**:282–284
 leukoregulin, **41**:286
 lymphotoxin-like molecules, **41**:284–286
 membrane attack complex of complement, **41**:299–307
 membrane damage, **41**:277–281
 NK cells, **41**:270–273
 pore-forming proteins, **41**:311–319
 reactive oxygen metabolism intermediates, **41**:286
 SCID and, **49**:395
 Cytotoxic T cell differentiation factor, T cell subsets and, **41**:64, 76
 Cytotoxic T cell recognition, maternally transmitted antigen and, **38**:313, 315, 316
 class I antigens, **38**:330, 331, 334–337
 generation, **38**:348–350
 maternally transmitted factor, **38**:328
 polymorphisms, **38**:320
 Cytotoxic T cells, **38**:2, 10, 86, 87, 105, 121, 122, 126, 127, 133; **41**:269, 270; **43**:195; **62**:87, 228, 231; **64**:2; **65**:277–278
 adenovirus, **65**:279
 adoptive T cell therapy of tumors and, **49**:283, 290–291
 antigen recognition, **49**:319–321, 324
 mechanisms, **49**:300, 303–305, 316, 318
 anterior chamber-associated immune deviation, **48**:208, 209
 antigen-presenting cells and, **47**:73, 79–81
 antigen processing, **47**:92
 immunogenicity, **47**:104
 antiviral therapy, **66**:295–298
 autoimmune demyelinating disease and, **49**:363
 autoimmune thyroiditis and, **46**:288
 autoreactive T cells and, **45**:425
 CD4⁺, **60**:303–310
 CD4⁺ and CD8⁺, **60**:289–292
 CD8⁺
 class II-restricted, **60**:309–310
 and HIV-1 immunopathology, **66**:273–274, 276, 277, 281
 in HIV-1 infection, **60**:133–134
 cell-mediated killing and, **41**:287–289, 291, 297
 class I antigens, **65**:279–280
 clones, **38**:20
 CTL epitope, *P. yoelii* circumsporozoite protein, **60**:112
 cytolysis and, **41**:273
 cytomegalovirus infection, **65**:278–279
 differentiation of, **40**:97–98
 effector T cell, **48**:197
 epitopes, **66**:276–280
 Epstein–Barr virus, **65**:279
 Fas expression in, **57**:138
 gene expression, **38**:13, 14
 granule exocytosis, **41**:281
 granule proteins, **41**:286, 287
 cytoplasmic granules, **41**:287
 pore-forming protein, **41**:291, 296, 297
 serine esterase, **41**:297, 298
 helper T cell cytokines and, **46**:111
 cross-regulation of differentiation, **46**:132
 functions, **46**:114–117
 immune responses, **46**:133, 135
 precursors of differentiation states, **46**:125
 secretion patterns, **46**:112, 113
 herpes simplex virus, **65**:279
 HIV and, **45**:356, 358, 365, 366
 HIV infection, **47**:390, 410, 411, 413; **65**:280
 adoptive immunotherapy, **65**:317–320
 CTL-mediated lysis, **65**:280–281
 CTLs and, **65**:280–322

- HLA and, **65:284–286**
 replication suppression, **65:281–284**
 in seronegatives, **65:287**, 307–309
 vaccines, **65:320–322**, 323
- HIV-1 persistence, **66:287**, 292–294
- HLA class I system, **65:279**
- HLA-DR antigens and, **42:293**
- hybrid resistance and, **41:385**
- hydrolytic enzymes, **41:275–277**
- immunosenescence and, **46:244**, 245, 249, 250
- influenza virus, **65:278**
- intracellular damage, **41:283**
- Ir* gene control of responses
- allogeneic cross-reactive lysis, **38:185–187**
 - class I molecules, **38:171–174**
 - H-Y antigen, **38:175–180**
 - immunodominance, **38:180–185**
 - I region genes, **38:161–171**
- Ir* genes and, **38:87**, 116, 174
- Langerhans cell, **48:197–198**
- leukocyte integrins and, **46:150**
- inflammation, **46:161**, 163, 164
 - ligand molecules, **46:172**
- lymphotoxin-like molecules, **41:284**, 285
- measles and, **45:341**, 345, 346
- mediated antigen-specific killing, **60:292–296**
- mediated cytotoxicity, **60:300–302**
- mediation, **41:270–273**
- MHC recognition, **41:135–138**; **55:364**
- carbohydrate moieties, **41:152–154**
 - exon shuffling, **41:138–149**
 - H-2 mutant strains, **41:158–163**
 - HLA class I antigens, **41:149–152**
 - HLA subtypes, **41:163–165**
 - β-microglobulin, **41:154–156**
 - monoclonal antibodies, **41:156–158**
 - somatic cell class I variants, **41:165–167**
- MHC restriction, **43:214–215**
- monoclonal antibodies, **38:277**
- mononuclear cells and, **42:294**
- mRNA, **38:21**
- NK cells and, **47:187**, 188, 196, 283, 295, 302
- congenital defects, **47:225**
 - cytotoxicity, **47:252–254**
 - differentiation, **47:229**, 232
 - effector mechanisms, **47:241**, 245, 246
- hematopoiesis, **47:282**
- reproduction, **47:272**
- surface phenotype, **47:208**, 211
- ocular HSV infection, **48:210**
- polarity, **41:273–275**
- response to foreign antigens, **43:205**
- response to virus infection, **43:205**
- SCID and, **49:396**, 398–399
- self-MHC class I recognition by effector cells, **57:314–315**
- sporozoite malaria vaccine and, **45:312**
- synthetic T and B cell sites and, **45:262**, 264
- antigens, **45:234**, 235
 - immunological considerations, **45:199**
 - viral antigens, **45:218–220**, 226, 227
- target recognition, **38:22**
- targets, death of, **50:69**
- T cell receptor and
- accessory molecules, **45:110**
 - alloreactivity, **45:151–157**
 - antigen processing, **45:118–120**, 125–127
- experimental systems, **45:137**
- MHC molecules, **45:116**
- peptides, **45:137**, 138
- polymorphic residues, **45:159**, 162
- structure-function relationships, **45:139–141**
- T cell repertoire, **45:166**
- T cell subsets and, **41:39**, 51
- activated T cells and hybridomas, **41:54**, 56, 57
 - alloreactivity, **41:78**, 81, 82
 - effector phase, **41:75**, 92, 94, 95
 - H-2 molecules in thymus, **41:97**, 99–103, 108
 - resting T cell subsets, **41:83–88**
 - T accessory molecules, **41:60**
 - unprimed and resting T cells, **41:65**, 71–75
- transplant, **48:197–198**
- transporters, need for, **65:47–48**
- tumor immunity and, **65:94**, 95
- Cytotoxins, **41:269**, 270

D

- D1.3, **43:20–21**, 222
- DAG, *see* Diacylglycerol

- Daudi cells, stimulation with CD40, **63:51**
- Daunomycin
in cancer treatment
antitumor activity, **56:333–334**
functional studies, **56:345**
monoclonal antibody conjugation, **56:334–336**, 338
- MC and, **42:265–268**
MG and, **42:265–268**
- DBA/2 mastocytoma cell, anterior chamber, **48:204**
- Death domain, **61:9**, 27, 31
- Death factor, Fas as, **57:129–140**
- Death genes, induction activation, **50:71–73**
- Death program, release, **50:75–76**
- Decay-accelerating factor, **48:241**; **61:203**, 204, 222–228
complement receptor 1 and, **46:189**, 197
deficiencies in PNH, **60:75–78**
expression on all blood cells, **60:78**
function, **61:219**, 224–228
gene, **61:222–223**
in GPI-anchored proteins, **60:60–66**
regulators of complement activation and genes, **45:403**, 408
protein expression, **45:395–402**
protein interactions, **45:383**
protein roles, **45:385–388**, 390, 391
structure, **61:207**, 222–224
surface antigens of human leukocytes and, **49:93**
therapeutic complement activation role, **56:273–274**
- Deep venous thrombosis, aPL antibodies and, **49:219**
- Degradation, *Ir* genes and, **38:93**, 95–97
- Degranulation
cytotoxicity and, **41:282**, 296, 297, 312
eosinophils
and $\beta 2$ integrins, **60:207**
morphology, **60:160–162**
mast cells, NO effect, **60:332**
- Dehydroepiandrosterone, **61:357**
- Delayed-type hypersensitivity
adoptive T cell therapy of tumors and, **49:289**, 327
- anterior chamber-associated immune deviation, **48:219**
antigen-specific down-regulation, **48:208**, 209
- intraocular tumor allograft, **48:208**, 209
- antigen-presenting cells and, **47:57**, 104
effector T cell, **48:197**
helper T cell cytokines and
cross-regulation of differentiation, **46:131**
differential induction, **46:127–130**
functions, **46:117**
human subsets, **46:139**
immune responses, **46:133**, 137, 138
precursors of differentiation states, **46:125**
- humoral immune response and, **45:24**, 27, 79, 85
- hybrid resistance and, **41:383**
IFN- γ and, **62:96–98**
- immunosenescence and, **46:221**, 237
- Ir* genes and
expression, **38:55**, 57, 60, 61
genetic control, **38:34–36**
H-Y antigen, **38:180**
I region mutations, **38:86**
MHC restriction, **38:65**
“Schlepper” experiment, **38:39**, 40
T cells, **38:41**, 42
interaction, **38:157**, 158
suppression, **38:135**
- Langerhans cell, **48:197–198**
- leukocyte integrins and, **46:162**
- NK cells and, **47:226**
- ocular HSV infection, **48:210**
- spleen, **48:211–212**
- spontaneous autoimmune thyroiditis and, **47:450**
- T cell subsets and, **41:76**, 77, 94
- transplant, **48:197**
tumor allograft, **48:208**, 209
UV5C25, **48:214**
- Delbrück model, antigenic variation, **38:340–344**
- Deletion, *see also* Clonal deletion
autoreactive T cells, **58:272–275**
- B cell formation and, **41:203**
 $CD4^+8^+$ thymocytes, **58:166–168**, 170–171, 259
complement components, **38:229**
cytotoxic T cells and, **41:139**, 167
 $\gamma\delta$ T cells, **58:313**
Ig heavy-chain variable region genes and, **49:9**, 14, 25, 31

- intrachromosomal, switch recombination, **61**:99
- Ir* genes, **38**:189
 antigen-specific clones, **38**:121
 clonal, **38**:92, 125
 macrophages, **38**:114, 115
 T cell selection, **38**:131, 132
 tolerance, **38**:127, 130
- peripheral, *in vitro* models, **58**:215–219
 peripheral T cell, CD3/TCR-mediated, *in vitro* models, **58**:219–224
- reduced level in G₁₀ or C₁₀ coding ends, **58**:40
- RSS junctional, **58**:54
 single and multiple base, in PNH, **60**:84–88
- Deletional switch recombination, in B cells, **60**:38–39
- Deletion mapping analysis, **42**:133–137
- Delta-kaolin clotting time, aPL antibodies and, **49**:201, 206
- Demyelinating disease, autoimmune, *see* Autoimmune demyelinating disease
- Dendritic cells
 adoptive T cell therapy of tumors and, **49**:330
 afferent lymph, CD1 protein expression, **59**:63–64
 antigen-presenting cells and, **47**:45–47, 87, 104, 105
 antigen presentation, **47**:71, 72, 74–76, 78–81
 antigen processing, **47**:88–92
 APC-T cell binding, **47**:95–97, 99
 cell surface, **47**:57–64
 immunogeneity, **47**:101–104
 interaction with T cells, **47**:93, 94
 T cell-dependent antibody responses, **47**:83, 84, 86
 T cell growth, **47**:99–101
 tissue distribution, **47**:47–57
- autoimmune thyroiditis and, **46**:280, 281, 296, 297, 300
- CD1 protein distribution, **59**:60–64
- CD23 antigen and, **49**:155, 171
- CD40 expression, **61**:13, 40–41
- complement receptor 1 and, **46**:190
- complement receptor 2 and, **46**:209
- distinction of FDCs from other, **51**:244–247
- helper T cell cytokines and, **46**:128
 IBD and, **42**:293, 296
 lung transplants, **59**:423
 pulmonary, **59**:376–381, 412
 role in V β selective element action, **50**:31
 SCID and, **49**:402
 as target for vaccination strategies, **58**:435–436
- T cell subsets and
 H-2 alloantigen recognition, **41**:89–92
 H-2 molecules in thymus, **41**:103–106, 108–110
 H-2-restricted antigen recognition, **41**:74, 75, 77
- Dengue virus, IFN- γ and infection, **62**:77
- Density, alterations during eosinophil activation, **60**:202–203
- Dental caries, subunit vaccines, **60**:135–136
- Deoxyguanosine, T cell subsets and, **41**:104, 106, 108
- Deoxynucleotidyltransferase, B cell formation and, **41**:194, 203, 204, 220, 231
- Deoxyspergualin, **65**:125–126
- Depletion
Ir genes
 complementation, **38**:74
 immunodominance, **38**:182, 184
 macrophages, **38**:113, 115
 tolerance, **38**:127
 SCID and, **49**:381, 394–397, 399, 401
- Depolarization
 cell-mediated killing and, **41**:273
 pore-forming protein and, **41**:293, 294
- Dermatitis
 CD23 antigen and, **49**:154
 SCID and, **49**:398
- Dermatomyositis, antinuclear antibodies and, **44**:111–113, 118–120, 137
- Desetope, *Ir* genes and, **38**:107
- Desmoglein
 amino acid sequence homologies and
 conserved sites with PV antigen, **53**:314
 characterization, **53**:300–303
 comparison with PV antigen and cadherins, **53**:311
- Desmoplakin I
 homology with 230-kDa pemphigoid antigen, **53**:305–310

- Desmoplakin I (*continued*)
 structural and amino acid homologies with
 BP antigen, **53**:307
- Desmosomes
 pemphigus foliaceus antigen localization,
53:300–303
 ultrastructure diagram, **53**:297
- Desotope, T cell subsets and, **41**:53, 58
- Determinant selection
Ir genes and, **38**:98, 99
 model, MHC restriction, **43**:207
- Development
 fetal, acquisition of clonotype repertoire during, **42**:26–30
 IL-6 role
 embryo, **54**:31–32
 fetus, **54**:32
 placenta, **54**:32
 intrathymic, *see* T cells, intrathymic development
 LIF role in, **53**:33–37
 neonatal, *see* Neonatal development
 regulation of IgR functional components, **54**:375–378
- DEX, *see* Dextran
- Dexamethasone, **62**:192, 195
 GM-CSF mRNA in T cells and, **39**:14
 hybrid resistance and, **41**:387
 IL-2 mRNA in T cells and, **39**:14
 IL-3 mRNA in T cells and, **39**:14
 induced T cell death, **58**:262–265
 NF- κ B and, **65**:119
- Dexter's culture system, B cell formation and bone marrow, **41**:208–210, 213, 215–217, 219, 220
- W/W anemic mice, **41**:224
- Dextran
 Ars A response and, **42**:131
 B cell repertoire expression and, **42**:58–62
 as intermediate carrier in
 anthracycline–antibody conjugation, **56**:337–338
 secondary B cell lineage and, **42**:70
- DFL16.1 gene segment, Ars A antibodies and, **42**:105–107
- DHA, **62**:186
- DH segments, **62**:1, 23
 Ig heavy-chain variable region genes and, **49**:2–3
 mouse and human, **62**:17–19
 organization, **62**:9–10
- Diabetes, *see also* Autoimmune diabetes, murine model
 autoimmune thyroiditis and, **46**:269, 284, 298
 insulin-dependent, *see* Insulin-dependent diabetes mellitus
 murine lupus models and, **46**:79
 NK cells and, **47**:301
 overt, T cell subsets involved in insulitis and, **51**:296–298
 patients, IL-2 deficiency of, **50**:173–174
 spontaneous autoimmune thyroiditis and, **47**:448, 468, 492
 suppressor mechanism controlling development of, **51**:301–303
- Diabetes mellitus
 antiadhesion therapy, **64**:197
 human-human hybridomas and, **38**:299
 IFN- γ and, **62**:100–101
 insulin-dependent, *see* Insulin-dependent diabetes mellitus
 mucosa-derived effector cells, **64**:192
 NO role, **60**:344–345
 pituitary hormones, **63**:423
 role of IL-1 α , **54**:214
- Diabetogenic genes, MHC-linked, **51**:290–293
- Diabetogenic T cells, repertoire of, **51**:298–301
- Diacylglycerol, **48**:228, 309
 IL-1 and, **44**:175, 176, 185
 immunosenescence and, **46**:223, 224
 in neutrophils
 amplification and, **39**:131–135
 protein kinase C and, **39**:130–131
- T cell activation and, **41**:15–19, 30
- Diamphotxin, cytolysis and, **41**:318
- Diapedesis, **65**:369
- Diazo reaction, as procedure for
 chemoimmunoconjugation, **56**:319
- Diutyryl cyclic AMP, autoimmune thyroiditis and, **46**:294
- Dicetyl phosphate, aPL antibodies and, **49**:229–230
- Diclofenac, **62**:184, 186
- Dif, in rel family, characterization, **58**:14
- Differentiation
 antigen-presenting cells and, **47**:48, 83, 84
 autoimmune thyroiditis and, **46**:266

- B cell
in CBA/N immunogenic mice, **53**:142–143
and growth, *in vivo* aspects, **52**:221–230
HICMX-1 as defect, **60**:42–48
isotype switching during, **60**:38–39
murine CD5 B cells, **55**:309–310
- B cell formation and, **41**:181, 235–237, 239
B cell precursors, **41**:188, 189, 191–192, 194, 195, 197, 199–201
bone marrow cultures, **41**:210, 212, 214–218, 220
genetically determined defects, **41**:223–226, 230, 231
inducible cell line, **41**:220, 221, 223
lymphohemopoietic tissues, **41**:185, 187
population dynamics, **41**:206, 207
soluble mediators, **41**:232, 235
- CD5 B cell and, **47**:117, 118, 123, 125
physiology, **47**:139, 141, 142
- complement receptor 2 and, **46**:204, 206
- cytolysis and, **41**:270
- eosinophils, **60**:162–164
- helper T cell cytokines and
cross-regulation, **46**:130–133
human subsets, **46**:139
immune responses, **46**:137
induction, **46**:127–130
precursors, **46**:122–127
- HIV infection and, **47**:379, 393
- hybrid resistance and
antigen expression, **41**:400
bone marrow cells, **41**:336
leukemia/lymphoma cells, **41**:369
marrow microenvironment, **41**:390, 392
- NK cells, **41**:372, 373
- syngeneic stem cells, **41**:388
- T cells, **41**:381
- IgE biosynthesis and
antibody response, **47**:8, 9, 11
binding factors, **47**:12, 13, 21, 23
- immunosenescence and, **46**:221, 253
lymphocyte activation, **46**:231, 232
stem cells, **46**:240, 242–247
- NK cells and, **47**:188, 198
adaptive immunity, **47**:291, 292, 294
congenital defects, **47**:225
effector mechanisms, **47**:235, 239
hematopoiesis, **47**:272, 276
morphology, **47**:216
- surface phenotype, **47**:201, 205, 208
in vitro, **47**:232–234
in vivo, **47**:229–232
- pre-B cells into B cells
in bc1-2 transgenic mice, **53**:143–144
in vitro, **53**:140–141
in vivo, **53**:141–142
- related translocation oncogenes, **50**:139
- spontaneous autoimmune thyroiditis and, **47**:447
- stem cell factor–ligand interaction in, **55**:40–42
- subpopulations, **53**:127–130
- surrogate L chain functions in, **53**:136–143
- T cell activation and, **41**:22
- T cells, CD4 and CD8 roles, **53**:94–113
negative selection, **53**:101–103
positive selection, **53**:97–98
thymocyte development and selection, **53**:94–97
- T cell subsets and
H-2 molecules in thymus, **41**:98–102, 104, 105, 107, 108
H-2-restricted antigen recognition, **41**:52, 64, 66, 76, 77
- terminal, and somatic mutation, apoptosis and, **50**:65–66
- Differentiation-inhibiting factor/differentiation-retarding factor, *see* Leukemia inhibitory factor
- Diffuse large cell lymphoma, gene rearrangements and, **40**:292
- DiGeorge syndrome, **59**:247
SCID and, **49**:383–384, 391
- Dilute Russell viper venom time, aPL antibodies and, **49**:201, 242
- Dimerization, through SH2 domain, **60**:19
- Dimerization/oligomerization, Epo receptor, **60**:9–11
- Dimethyl sulfoxide, for MG, **42**:264
- Dinitrophenyl
IgE biosynthesis and, **47**:28
synthetic T and B cell sites and, **45**:199
- Dinitrophenyl-Asc, IgE biosynthesis and, **47**:3–5
- Dinitrophenyl-KLH, IgE biosynthesis and antibody response, **47**:5, 6
antibody response suppression, **47**:29, 31, 33, 34, 37

- Dinitrophenyl-OVA, IgE biosynthesis and antibody response, **47:5**
 antibody response suppression, **47:28–30**,
32–34, **37**
 binding factors, **47:12**, **14**
- Dinitrophenyl-Rag, IgE biosynthesis and, **47:4**
- Dintzis model, size fractionated polymers, **52:288–289**
- Diphtheria, **38:32**, **289**, **295**
- Direct cDNA selection, **59:165**
- Direct cellular cytotoxicity, spontaneous autoimmune thyroiditis and, **47:451**, **452**, **469**
- Directional cloning, **38:205**
- Direct two-color immunofluorescence, for NK cell surface antigens, **42:200**
- DIR segments, Ig heavy-chain variable region genes and, **49:30–31**, **33**
- Disease, *see also* Xenograft models
 clinical
 fiberoptic bronchoscopy studies, **60:219–223**
 and IL-6, **54:44**, **45**
 cytokines, **63:270**, **310–314**
 IL-2R, **63:294**, **296–298**
 and IL-6, **54:39–43**, **44–45**
 IL-6, **63:301**, **302**
 immunity, role of IELs, **58:**
 326–329
 role of eosinophils, **60:151–228**
 TNF, **63:306**
 virulence factors, **63:308–309**
- Disseminated intravascular coagulation, endotoxic shock, **66:260**
- Dissociation rate constants, **42:168–172**
- Disulfide
 CD4 molecules and, **44:285**, **286**
 CD8 molecules and, **44:270–273**
 genetically engineered antibody molecules and
 biological properties, **44:83**
 chimeric antibodies, **44:81**, **82**
 expression, **44:76**
 production, **44:67**, **69**
- Ig gene superfamily and
 homology unit, **44:3**
 nonimmune receptor members, **44:31**, **34**, **35**, **38**, **41**, **42**
 receptors, **44:9**
- lymphocyte homing and, **44:342**
 T cell development and, **44:207**, **213**, **233**
- Disulfide bridge, **48:13**
- Dithiocarbamates, action of, **65:123**
- Divergence, functional, and positive selection
 T cell mechanism, **51:186–189**
- Diversification
 germinal centers, **60:283–284**
 somatic
 Ig gene superfamily and, **44:12–20**
 Ig VL gene, **48:60–62**
- Diversity
 combinatorial, in leukocyte–endothelial cell recognition, **58:389–390**
 N region, **58:301–303**
- DNA, *see also* Complementary DNA;
 Mitochondrial DNA; Recombinant DNA
 antigen-presenting cells and, **47:75**
 antigen processing, **47:92**
 APC-T cell binding, **47:95–97**
 T cell growth, **47:99–101**
- antinuclear antibodies and, **44:93**
 autoantibodies, **44:125**, **127**, **137**, **138**
 autoantigen, **44:127**, **128**, **131**
 drug-induced autoimmunity, **44:109**
 scleroderma, **44:121**, **122**
 SLE, **44:95–100**, **105–107**
- aPL antibodies and, **49:209**, **226**, **240**, **248–249**, **251**, **256**
- autoimmune thyroiditis and, **46:267**, **303**
- B cell formation and, **41:183**, **184**, **203**, **217**, **235**
- breakage
 immunodeficiency syndromes, **61:314–316**
 KU 70/86, **61:301–303**
 scid mutation, **61:299–301**, **317**
 V–D–J recombination, **61:298–303**
- broken, in V(D)J recombination, **64:43–45**
- CD4 molecules and, **44:286–288**
- CD5 B cell and, **47:161**
 Ig gene expression, **47:152**, **154–156**
 marker for activation, **47:127**
 physiology, **47:131**, **133**, **134**, **136**, **137**
- CD8 molecules and, **44:272**, **275**, **277**
- CD23 antigen and, **49:167**
- cell-mediated killing and, **41:298**
 cloning, **44:74**
 expression, **44:70**, **71**, **77**, **78**
 fusion proteins, **44:85**, **86**

- complement receptor 1 and, **46**:188
complement receptor 2 and, **46**:206
cytolysis and, **41**:277, 283–285
damage, *see* DNA damage
degradation, target cells, by CTLs, **60**:300–301
deletion and looping out, **54**:234
double-strand break repair, **64**:49, 51
model, **61**:111
relevance of joining activity, **58**:45
and V(D)J recombination, **58**:29–85; **64**:49–51
cell cycle regulation, **58**:70–74
and Ku autoantigen, **58**:55–60
double-strand breaks, role in V(D)J
recombination, **58**:32–36
Ets transcription factor binding, **64**:66–68
fragmentation, **58**:246–248, 251–253,
262–264, 270–271
genetically engineered antibody molecules
and, **44**:66, 89
genomic rearrangement, **38**:9
HIV infection and
etiological agent, **47**:380, 382
immunopathogenic mechanism, **47**:387,
401, 402
neuropsychiatric manifestations, **47**:403
humoral immune response and, **45**:6,
32, 35
hybrid resistance and, **41**:343, 377, 378
IgE biosynthesis and, **47**:17
Ig gene superfamily and
evolution, **44**:46
homology unit, **44**:8
receptors, **44**:12, 14, 17, 20
Ig heavy-chain variable region genes and,
49:1, 3
D segments, **49**:29, 31–33
J_H segments, **49**:33
polymorphism of V_H gene segments,
49:23–24
regulation, **49**:53, 55, 61
V_H families, **49**:8, 9, 11
V_H gene expression, **49**:35, 39–40,
42–43, 46, 48–50
IL-1 and, **44**:165, 177
immunosenescence and, **46**:230, 234, 253
Ir genes and
cytotoxic T cells, **38**:169
recombination, **38**:86
T cell repertoire, **38**:116, 145
T cell suppression, **38**:148, 150
–Ku complex, **58**:55–57, 61–62
leukocyte integrins and, **46**:160, 174, 175
maps of Ig-like loci and, **46**:3–6, 46
structure, **46**:11–13, 15, 18, 25, 27
T cell receptor, **46**:36, 37, 39, 41–43, 45
MHC class I molecule-encoding, **58**:433
mitochondrial, *see* Mitochondrial DNA
murine lupus models and
Ig germline, **46**:66, 69, 71, 72, 76–78
T cell antigen receptor, **46**:83, 84, 87
naked
and cytotoxic T cell responses,
58:436–437
linkage of cell-surface protein, **58**:423
NK cells and, **47**:209, 235, 253, 254
nuclear, fragmentation, and PCD,
58:212–213
PIG-A, analysis, **60**:84
promoters and enhancers, **43**:239
–protein intermediates, **58**:38–39
and rabbit antibody repertoire
B cell rearrangements, **56**:190–192
probes to molecules, **56**:182–183
repair, plasmacytoma development,
64:240–241
repair defects, **61**:310–311
repair syndromes, **58**:62–69
repair synthesis, across double-strand gap,
58:44
replication, role in immune response,
54:237–238
RNA-duplex switch region complex,
61:114–116
SCID patients, **61**:310–311
spontaneous autoimmune thyroiditis and,
47:435, 493
altered thyroid function, **47**:458, 464
disturbed immunoregulation, **47**:470
genetics, **47**:483, 484
sporozoite malaria vaccine and
antibodies, **45**:301
CS proteins, **45**:297
immunity, **45**:291
interferon- γ , **45**:302
Plasmodium vivax, **45**:317
strand polarity, **60**:279
surface antigens of human leukocytes and,
49:116

- DNA (*continued*)
 synthetic T and B cell sites and, **45:231**
 T cell activation and, **41:29**
 T cell receptor and, **45:109, 165**
 T cell subsets and, **41:41, 44, 46–48, 78**
 transcription inhibition, **58:240**
 transfection, **38:280, 305, 306**
 expression of IgR on cell surface, **54:360–361**
 as substrate introduction method in V(D)J joining, **56:50–53**
 virus-induced immunosuppression and HIV, **45:354, 355**
 measles, **45:344, 348**
- DNA binding
 effect of Stats, **60:28**
 Stat homology blocks in, **60:18–19**
- DNA damage
 and p53 accumulation, **58:258**
 in programmed cell death, **50:57–60**
 fragmentation, **50:57–58**
 methods, **50:59–60**
 observations in rodent thymocyte, **50:58–59**
 pattern of fragments, **50:58**
- DNA-dependent protein kinase, **61:300–301, 63:154**
 association with Ku, **58:60–62**
 DNA-PK_{cs}, **64:49–50, 60**
- DNA duplication, **38:211, 212**
- DNA flap, **61:304**
- DNA ligase, deficiency in Bloom syndrome, **58:65–66**
- DNA-PK, *see* DNA-dependent protein kinase
- DNA polymerase
 antinuclear antibodies and, **44:94, 108, 135**
 genetically engineered antibody molecules and, **44:85**
- DNA rearrangement, T cell receptor and, **38:5, 9**
- DNA RFLP typing strategy, **48:117–118**
- DNase I
 sensitivity as indicator of gene recombination, **56:43–44**
- T cell-specific hypersensitive site, **43:268**
- DNA sequence
 complement components, **38:212**
 human B cell neoplasia, **38:266**
 maternally transmitted antigen and, **38:330**
 T cell receptor and, **38:7, 15**
- DNA topoisomerase 1, antinuclear antibodies and, **44:94, 121, 131, 133, 134, 137**
- Dob, Fc, **48:21**
- Dob antibody structure, **43:8**
- Docosahexaenoic acid
 arachidonic acid metabolism and, **39:159–160, 164**
 in human neutrophils, **39:165–166**
 formation from N-3 and N-6 fatty acids, **39:159–160**
 metabolism, 5-lipoxygenase and, **39:160–161**
- Dog leukocyte antigen, hybrid resistance and, **41:349, 350**
- Domain, defined, **63:201**
- Domain-domain interactions, **43:110**
- Domain organization, CD1 proteins, **59:35–36**
- Domain structure
 genetically engineered antibody, **48:14, 16, 22**
 segmental flexibility, **48:14, 16, 22**
- Dominant resistance, IDDM, **48:143–144**
- Double-homologous recombination, **48:56**
- Down-regulation
 in clonotype repertoire expression, *see* Environment, B cell repertoire expression and signal transduction through B cell Ig receptor, **54:374**
- Down syndrome, leukocyte integrins and, **46:160**
- DP, **48:109, 110–111**
- DP allele, celiac disease, **48:145**
- DP α allele, celiac disease, **48:145**
- DP β allele, celiac disease, **48:145**
- DP α -DQ β , multiple sclerosis, **48:149**
- DP α -DQw1 β , celiac disease, **48:146–147**
- DPM1, **60:73–74**
- DP α polymorphism, celiac disease, **48:145**
- DP β polymorphism, celiac disease, **48:145**
- D μ protein, **63:7, 22–23**
- D-proximal V μ genes, **49:5**
 V μ families, **49:18, 20–21**
 V μ gene expression, **49:36–39**
- DPw4, multiple sclerosis, **48:149–150**
- DQ, **48:109, 110–111**
 allele, linkage disequilibrium, **48:121–122**
 heterodimer, DQ2-DQ3 cell, **48:120–121**

- linkage disequilibrium phenomenon, **48:136–137**
- DQ α
 - cis*-complementation, **48:139–140**
 - functional dimer, **48:123**
 - gene polymorphism, multiple sclerosis, **48:149**
 - polymorphism, **48:121**
 - structural requirement, **48:138–139**
 - trans*-complementation, **48:139–140**
- DQ β
 - amino acid 57, location, **48:139**
 - cis*-complementation, **48:139–140**
 - functional dimer, **48:123**
 - IDDM, **48:134–136**
 - position 57 aspartic acid, **48:135–136, 137**
 - polymorphism, **48:121**
 - structural requirement, **48:138–139**
 - trans*-complementation, **48:139–140**
- DQ α -DQ β , multiple sclerosis, **48:149**
- DQ6, **65:286**
- DQR2-6, multiple sclerosis, **48:148–149**
- DQw1, multiple sclerosis, **48:148**
- DQw1a, multiple sclerosis, **48:148**
- DQw1b, multiple sclerosis, **48:148**
- DQw2
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw2 α chain, celiac disease, **48:146**
- DQw3
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw3.1
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw3.2
 - B8, **48:134**
 - B15, **48:134**
 - β chain, **48:136**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw3.3
 - B8, **48:134**
- B15, **48:134**
- DR3, **48:134**
- DR4, **48:134**
- DQw7
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw8
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw9
 - B8, **48:134**
 - B15, **48:134**
 - DR3, **48:134**
 - DR4, **48:134**
- DQw9 Asp⁵⁷ positive
 - black patient, **48:140**
 - caucasian patient, **48:140**
- DR, **48:109, 110–111**
 - linkage disequilibrium phenomenon, **48:136–137**
- DR1, **48:127**
- DR2, **48:112; 65:286**
 - IDDM, **48:136, 143**
 - AZH, **48:136**
 - multiple sclerosis, **48:148–150**
- DR3
 - celiac disease, **48:144, 147**
- DQw2, **48:134**
- DQw3, **48:134**
- DQw3.1, **48:134**
- DQw3.2, **48:134**
- DQw3.3, **48:134**
- DQw7, **48:134**
- DQw8, **48:134**
- DQw9, **48:134**
- DR4, **48:112**
 - DQw2, **48:134**
 - DQw3, **48:134**
 - DQw3.1, **48:134**
 - DQw3.2, **48:134**
 - DQw3.3, **48:134**
 - DQw7, **48:134**
 - DQw8, **48:134**
 - DQw9, **48:134**
- HLA-DQ, **48:134**
- juvenile rheumatoid arthritis, **48:131**

- DR4 Dw haplotype, **48:118**
 DR4 haplotype, **48:127**
 DR5
 celiac disease, **48:144, 147**
 juvenile rheumatoid arthritis, **48:131**
 DR7, celiac disease, **48:144, 147**
 DR7 haplotype
 black patient, **48:140**
 caucasian patient, **48:140**
 DR9-DQ β
 black patient, **48:140**
 caucasian patient, **48:140**
 DR9 haplotype
 black patient, **48:140**
 caucasian patient, **48:140**
 DR allele, linkage disequilibrium, **48:121–122**
 DR antigens, IBD and, **42:288, 292–294**
 DRB1, RA susceptibility associated with,
 56:410–416
 DR α -DQ β , multiple sclerosis, **48:149**
 DR α -DQw2 β , **48:125**
 DR α -DQw3 β , **48:125**
 DR α -DQw I β , **48:125**
 celiac disease, **48:146–147**
 DR α gene, **48:124**
 expression modulation, **48:125**
 DR β gene, rheumatoid arthritis, **48:131**
Drosophila
 antinuclear antibodies and, **44:129**
 dorsal gene, **52:271**
 identification of Jak homolog, **60:3–4**
 leukocyte integrins and, **46:153**
 maternally transmitted antigen and, **38:322**
 DR α protein, **48:124**
 Drug development, HIV-1 infection,
 63:95–99, 107
 Drug-induced autoimmunity, antinuclear
 antibodies and, **44:109–111**
 autoantibodies, **44:136**
 autoimmune response, **44:131**
 scleroderma, **44:124**
 Sjögren's syndrome, **44:112**
 Drug-induced syndromes, aPL antibodies
 and, **49:193, 209, 212–213, 250**
 Drug-monomoclonal antibody conjugates, as
 cancer treatment, *see*
 Chemoimmunoconjugates
 Drugs
 cytotoxic, and apoptosis, **58:237–240**
 therapeutic, and leukocyte homing,
 58:395–396
- DRw8
 celiac disease, **48:147**
 juvenile rheumatoid arthritis, **48:131**
 DRw9, IDDM, **48:137**
 DRw15, IDDM, **48:143**
 DRw16, IDDM, **48:143**
 DRw52, **48:112**
 DRw53, **48:112**
 DRw53 molecule, rheumatoid arthritis,
 48:128
 DSB repair, *see* DNA, double-strand break
 repair
 D segments, Ig heavy-chain variable region
 genes and, **49:18, 28–34, 36, 53**
 organization, **49:3–6**
 DuP697, **62:186**
 Dw1O, **48:127**
 rheumatoid arthritis, **48:126–127**
 Dw4
 position 67, **48:128–129**
 position 70, **48:128–129**
 position 71, **48:128–129**
 rheumatoid arthritis, **48:126–127**
 DW14, rheumatoid arthritis, **48:126–127**
 DX α polymorphism, **48:139**
 Dyscrasias, *see* Plasma cell dyscrasias
- E**
- E α , **62:69**
 E β , **62:43, 51–52**
 E γ , **62:69**
 E1A, **65:11**
 E1B 19K, NF- κ B activation and, **65:128**
 E2A gene, B cell genesis, **63:218–219,**
 232–235
 E3/19K glycoprotein, adenovirus, effects on
 antigen processing and presentation,
 52:44–47
 α E β 7, lymphocyte homing and, **64:156**
 α E β 7 Integrin, **65:359**
 E47/E12, binding site, switch recombination,
 61:117, 118–119
 EA-1, **48:248–249**
 EAF, *see* Eosinophil-activating factor
 EAMG, *see* Experimental autoimmune
 myasthenia gravis

- Early activation antigen-1, **48**:248–249
 Early response cytokines, **62**:267, 276, 280
EBF, **63**:250
EBP, **43**:307–308
EBP γ , **65**:18
EBV, *see* Epstein–Barr virus
EC, *see* Endothelial cells
ECF-A, *see* Eosinophil chemotactic factor of anaphylaxis
E chain, **43**:278
 biosynthetically engineered peptides based cDNA constructs, **43**:281
 digestion, **43**:282
 peptide immunogens, **43**:279–280
 sites protected by receptor interaction, **43**:280
 stabilization, **43**:282
Echoviral infections, X-linked
 agammaglobulinemia, **59**:145
ECP, *see* Eosinophil cationic protein
ECP1, **61**:10
Ecto-5'-nucleotidase, **48**:241
Edema, paw, NO role, **60**:328–329
Edema, tissue, in IBD, **42**:310
EDN, *see* Eosinophil-derived neurotoxin
Effector cells
 activated in immune response, **58**:298
 adoptive T cell therapy of tumors and, **49**:284, 333–334
 antigen recognition, **49**:318–319, 324
 expression of antitumor responses, **49**:324–325, 327, 329, 332
 mechanisms, **49**:299, 301–302, 306, 308–313, 315
 NK cells, **49**:316–318
 principles, **49**:286, 288–290, 294–295, 297
 cell-mediated killing and, **41**:296
 cytosis and, **41**:274, 275
 granule exocytosis, **41**:281
 hydrolytic enzymes, **41**:276
 intracellular damage, **41**:283, 284
 membrane damage, **41**:277
 cytolytic, generation by CD4 $^{-}$ 8 $^{+}$ $\alpha\beta$ T cells, **58**:108
 cytotoxicity and, **41**:269, 270, 272
 cytotoxic T cells and, **41**:148, 150, 170
 hybrid resistance and, **41**:334, 370
 antibodies, **41**:376, 378, 379
 antigen expression, **41**:402, 403
 bone marrow cells, **41**:346
 leukemia/lymphoma cells, **41**:359, 366, 368
 lymphoid cells, **41**:353
 macrophages, **41**:371
 marrow microenvironment, **41**:390
 NK cells, **41**:372, 373
 syngeneic stem cells, **41**:390
 T cells, **41**:382–384
in vitro assays, **41**:393–396
 lymphocyte homing and, **44**:332
 NK cells and, **47**:196, 302
 antimicrobial activity, **47**:286, 287
 hematopoiesis, **47**:273, 274, 277, 280
 trafficking to vascularized tumors, **58**:432
 variance with introduced gene, **58**:427
Effector cell–target cell interaction, mediated by leukocyte Fc receptors, **51**:51–54
Effector mechanisms
 hybrid resistance, **41**:369, 370
 antibodies, **41**:376–380
 macrophages, **41**:370–372
 marrow engraftment, **41**:384–388
 marrow microenvironment, **41**:390–393
 NK cells, **41**:372–376
 syngeneic stem cell functions, **41**:388–390
 T cells, **41**:380–384
in vitro assays, **41**:393–396
 in IBD, **42**:321
 NK cells and, **47**:234, 235
 cytotoxicity, **47**:248–254
 lymphokines, **47**:262–266
 receptors, **47**:242–248
 regulation, **47**:254–262
 target cells, **47**:235–242
 spontaneous autoimmune thyroiditis and, **47**:446, 449, 465–469
Effector molecules
 ATP, in CTL cytotoxicity, **60**:301–302
 downregulation during inflammation, **60**:338–339
 macrophage, in cell-mediated immune response, **60**:333–341
 NO, during inflammation, **60**:332–346
 perforin, in cytotoxicity, **60**:290–292
Effector phase
 B cell activation, **63**:44
 T cell subsets and, **41**:75–77, 92–95

- Effector T cells
 cytotoxic T cell, **48:197**
 delayed-type hypersensitivity, **48:197**
 functional and phenotypic properties, **53:225–227**
 functional differences with memory and naive cells, **53:227–228**
 maturation into, **53:229–230**
 migration
 basic pattern, **53:242–244**
 tissue topic subsets, **53:244–247**
 primary and secondary, **53:227–228**
 EGF, *see* Epidermal growth factor
 Eicosanoids, and asthma pathophysiology, **51:340–353**
 leukotrienes, **51:341–347**
 biological activities, **51:343**
 leukotriene antagonists and inhibitors, **51:346–347**
 leukotrienes and airway
 hyperresponsiveness, **51:344–346**
 potency, **51:343**
 release of leukotrienes in asthma, **51:346**
 synthesis and metabolism, **51:341–343**
 platelet-activating factor, **51:352**
 prostaglandins and thromboxane, **51:347–352**
 biological activities, **51:347**
 measurements of prostanoids in biological fluids, **51:349**
 pharmacological modulation of airway hyperresponsiveness, **51:351–352**
 airway tone and acute asthmatic response, **51:350–351**
 prostanoid action, **51:350–352**
 potency, **51:347–349**
 prostanoids and airway
 hyperresponsiveness, **51:349**
 synthesis and metabolism, **51:347**
 Eicosanoids, as eosinophil-derived lipid mediators, **60:188–190**
 Eicosapentaenoic acid
 cyclooxygenase pathway inhibition, **39:160, 164–165**
 dietary, animal studies leukotriene production and, **39:162–163**
 dietary, human studies
 arachidonic acid in neutrophils and, **39:166**
 in asthma patients, **39:166–167**
 T cell subpopulations and, **39:167**
 dietary N-3 fatty acids and, **39:160**
 esterified and nonesterified, arachidonic acid in human neutrophils and, **39:167–168**
 formation from N-3 and N-6 fatty acids, **39:159–160**
 metabolism
 5-lipoxygenase pathway, **39:160–161**
 leukotriene production, **39:160–162**
 SLE and, **39:164**
 ELAM-1, **59:414**
 Elastase, IL-1 and, **44:163**
 Elastin, **62:266**
 Electron microscopy
 antigen-presenting cells and
 antigen processing, **47:91**
 interaction with T cells, **47:94**
 tissue distribution, **47:51, 52, 57**
 antinuclear antibodies and, **44:124**
 autoimmune thyroiditis and, **46:267, 274, 291**
 B cell formation and, **41:212**
 cell-mediated killing and, **41:292, 294, 295**
 complement receptor 1 and, **46:184**
 cytolysis and, **41:282**
 humoral immune response and, **45:37**
 IL-1 and, **44:184**
 leukocyte integrins and, **46:167**
 lymphocyte homing and, **44:319**
 membrane attack complex of complement and, **41:300, 306**
 NK cells and, **47:218**
 regulators of complement activation and, **45:392, 393**
 spontaneous autoimmune thyroiditis and, **440, 47:462**
 T cell subsets and, **41:106**
 Electron-spin resonance, membrane attack complex of complement and, **41:303**
 Electrophoresis
 complement receptor 1 and, **46:189, 190, 192**
 Ig heavy-chain variable region genes and, **49:4–5, 11**
 immunosenescence and, **46:230**

- leukocyte integrins and, **46:154**, 157
 maps of Ig-like loci and, **46:5**
 MHC, **46:28**, 30, 33
 structure, **46:15**, 18, 27
 T cell receptor, **46:36**, 37, 44
- Electrophoretic mobility shift assays, **54:255**
- Electrostatic complementarity, **43:12–13**
- Electrostatic forces, binding role, **43:13**
- Electrostatic potential, MHR, **43:35**
- Electrostatic recognition, between antigen and antibody, **43:12–13**
- Elf-1
 binding sites, **64:76–77**
 domain, **64:76–77**
 expression, **64:91**
 gene targeting, **64:92**
 structure, **64:78**
 structure–function studies, **64:91–92**
- ELISA, aPL antibodies and, **49:205**, 235–236, 242–244, 247
- Embryo
 anti-TCR1, -TCR2, -TCR3 antibody treatment, **50:103–105**
 thymectomy, **50:102**
 B cell formation, **41:181**, 193, 199, 201, 224, 227
 chicken, B cell, early commitment in, **57:367–369**, 374
 development, role of IL-6, **54:31–32**
- Embryogenesis
 bursar stem cell, **48:52**
 maternally transmitted antigen and, **38:321**, 330, 333
 V-J joint, **48:49**, 50
- Embryonic stem cells
 B cells from, **53:124–130**
 LIF effects, mouse, **53:33–34**
 mutant, **62:32**, 36–37, 40
- Emphysema, **62:188**, 266
- Encephalitis, experimental allergic, *see* Experimental allergic encephalitis
- Encephalomyelitis, experimental autoimmune, *see* Experimental autoimmune encephalomyelitis
- End-joining
 and nonhomologous recombination, **58:43–45**
 and *scid* mutation, **58:50**
 in *xrs* group mutants, **58:54**
- Endocrine cells, IFN- γ and, **62:93–94**
- Endocytosis
 antigen-presenting cells and
 antigen presentation, **47:64**, 67, 68, 71, 81
 antigen processing, **47:87–91**
 immunogenicity, **47:102**
 tissue distribution, **47:51**
 CD4 molecules and, **44:296**
 cell-mediated killing and, **41:298**
 complement receptor 1 and, **46:193**, 198–200
 complement receptor 2 and, **46:210**
 HIV infection and, **47:379**
 Ig gene superfamily and, **44:31**
 IL-1 and, **44:181**
 and processing of membrane IgR/antigen complex, **54:348**
 surface antigens of human leukocytes and, **49:89**, 98, 115, 125
- Endogenous pyrogen, cachectin as, **42:224**
- Endogenous retroviral insertions, inseparable from Vbse, **50:38–39**
- Endoglin, *see* Transforming growth factor β receptor
- Endoglycosidase F, autoimmune thyroiditis and, **46:270**, 271
- Endoglycosidase H
 autoimmune thyroiditis and, **46:271**
 leukocyte integrins and, **46:154**, 174
- Endonuclease
 in apoptotic cell degradation, **58:269–270**
 blockade, in apoptosis inhibition, **58:251**
- Endoplasmic reticulum, **64:105**
 adoptive T cell therapy of tumors and, **49:283**, 320, 323
 antinuclear antibodies and, **44:120**, 128
 binding of PIG-1, **60:67**
 cytoplasmic face, **60:65**
 CPI anchoring in, **60:60**
 MHC class I molecules
 peptide anchoring, **64:124–127**
 peptide binding, **64:114–115**, 118, 127–128
 peptide loading, **64:105–128**
 regulators of complement activation and, **45:400**
- T cell receptor and, **45:123**, 125
- α -Endorphin, immunosuppressive, **39:309–310**
- β -Endorphin, **48:170–172**
 effects on lymphocytes
 chemotaxis and, **39:307–308**

- β -Endorphin (continued)**
- immune responses and, **39:310**
 - natural killing activity and, **39:310**
 - production by leukocytes, **39:315–316**
- β -Endorphin receptors**
- in CNS, **39:312**
 - on lymphocytes, **39:312**
- Endosteum, B cell genesis**, **63:214**
- Endothelial cell costimulators**
- in alloantigen responses, **50:271–272**
 - in polyclonal CD4+ T cell activation, **50:268–270**
 - in polyclonal CD8+ T cell activation, **50:270–271**
 - soluble, production, **50:272–274**
- Endothelial cells**, **54:29–30**, *see also Human umbilical vein endothelial cells; Vascular endothelial cells*
- adhesion molecules, regulation, **64:139–140**, **146**, **147**, **163–173**
 - adhesion to leukocytes, **58:376–378**, **381–389**
 - antigen-independent recruitment of T cells into tissues by, **50:274–287**
 - extravasation of T cells, **50:285–287**
 - migration of memory T cells to inflammatory sites, **50:278–288**
 - endothelial leukocyte adhesion molecules, **50:279–281**
 - migration of naive T cells to lymph nodes, **50:275–278**
 - migration of pre-T cells to thymus, **50:275**
 - tissue-specific homing of lymphocytes, **50:284–285**
- B cell formation and**, **41:186**, **235**, **236**, **222–225**
- cachectin and, **42:223–224**
- CD40 expression, **61:14**, **41**
- cultured, alloantigen presentation by, **50:266–267**
- high endothelial, lymphocyte homing and, **44:315**, **316**, **318**, **320**
- hybrid resistance and, **41:400**
- IFN- γ , **62:81**
- IL-1 and, **44:153**, **190**
 - biological effects, **44:166–168**
 - gene expression, **44:159**, **160**
 - human disease, **44:194**, **195**
- immunocompetent cells, **44:177**
- receptor, **44:181**
- structure, **44:158**
- systemic effects, **44:169**, **171**
- TNF, **44:185**, **186**
- leukocyte integrins and
- inflammation, **46:162**, **164**, **165**, **168**
 - ligand molecules, **46:170–172**
- lymphocyte homing and, *see High endothelial cells; High endothelial venules*
- lymphocyte interaction with, **64:140–143**
- migration of memory T cells to
- inflammatory sites, endothelial leukocyte adhesion molecules expression *in vitro*, **50:283–284**
 - role in lymphocyte binding *in vitro*, **50:281–283**
- production of soluble costimulators, **50:272–274**
- as source of chemokines, **55:107**
 - T cell subsets and, **41:46**, **90**
- Endothelial-derived relaxant factor**, **54:29–30**
- effects via NO, **60:326**
- Endothelialitis**, **59:420**
- Endothelial leukocyte adhesion molecules**, **50:279–281**
- expression *in vivo*, **50:283–284**
 - role in lymphocyte binding *in vitro*, **50:281–283**
 - in vitro* binding in lymphocyte, **50:281–283**
 - in vitro* expression, **50:283–284**
 - in vivo* expression of, **50:283–284**
- Endothelial venules, L-selectin association**
- with lymphocyte adhesion, **62:257**
- Endothelium**
- aPL antibodies and, **49:193**, **250**, **252–254**
 - autoimmune thyroiditis and, **46:275**
 - integrity alterations, NO role, **60:327–330**
 - leukocyte transmigration, **58:388–389**
 - surface antigens of human leukocytes and, **49:86–88**
 - tumor vasculature, **64:180**
 - vascular, injury caused by TNF, **56:351**
- Endotoxemia**, *see also Sepsis*
- human experimental
 - IL-1, **66:116–117**
 - IL-8, **66:137–138**

- model, **66**:102
TNF, **66**:114–116
- IL-1, **66**:116–117
sepsis and, **66**:130–131
TNF and, **66**:114–116
- Endotoxic shock, contact activation proteins, **66**:260
- Endotoxin
B cell formation and, **41**:224, 230
cachectin and, **42**:214–216
and IL-1, **54**:31
IL-1 and
biological effects, **44**:166, 167
gene expression, **44**:159–161, 163, 164
human disease, **44**:192, 194, 195
TNF, **44**:185–188
lymphotoxin and, **42**:216
- Endplates, in MC patient, **42**:261–262
- Enhancers, Ig heavy-chain variable region genes and, **49**:52–62
sequences, **49**:39
- Enkephalin, helper T cell cytokines and, **46**:116
- Enkephalin heptapeptide, in pulmonary airways, **39**:304
- Enterotoxin, *see Staphylococcal enterotoxin A; Staphylococcal enterotoxin B*
- Enteroviral infections, X-linked
agammaglobulinemia, **59**:145–146, 150
env, adoptive T cell therapy of tumors and, **49**:320–321
- Environment, B cell repertoire expression and, **42**:35
assessment of, **42**:35–36
diversification and, **42**:36–39
down-regulation in clonotype repertoire expression and, **42**:42–43
antiidiotypic recognition, **42**:51–62
self antigens, **42**:43–51
predominant clonotype expression and, **42**:39–42
up-regulation, **42**:22–23
- Environmental agents, aPL antibodies and, **49**:221–222
- Enzymes
antineuclear antibodies and, **44**:122, 127, 133
aPL antibodies and, **49**:203, 239
CD23 antigen and, **49**:156–157, 172–173
- genetically engineered antibody molecules and, **44**:70, 75, 85
- granule-stored, **60**:201
- Ig gene superfamily and, **44**:12, 17, 47
- Ig heavy-chain variable region genes and, **49**:17
- IL-1 and, **44**:160, 170
lymphocyte homing and, **44**:317, 320, 360, 365, 369
- metal-requiring, **60**:340
- mitochondrial, action of, **65**:122
- myeloid-specific, **60**:25–26
- SCID and, **49**:382–383, 398
surface antigens of human leukocytes and, **49**:100, 111–114
- Eosinopenia, ascribed to treatment with glucocorticoids, **60**:227
- Eosinophil-activating factor, from monocytes, **39**:215–216
- Eosinophil cationic protein, **41**:311, 312
amino acid sequence, **39**:192–195
homology to pancreatic RNase, **39**:195
BAL, **60**:220–222
bronchial asthma and, **39**:224
coagulation and, **39**:196
fibrinolysis and, **39**:196
in granule matrix, **39**:195
induced damage to schistosomula, **60**:212–213
lymphocyte proliferation inhibition, **39**:196–197
neurotoxicity, **39**:196
parasite killing, **39**:196, 213
purification, **39**:191–194
release from eosinophils, **60**:204–205
sharing chromosome 14q with eosinophil-derived neurotoxin, **60**:193
- Eosinophil chemotactic factor of anaphylaxis
eosinophil activation, **39**:214–215
production by mast cells, **39**:214
- Eosinophil-derived neurotoxin, **41**:311
antihelminthic activity, **60**:212
Gordon phenomenon induction, **39**:198–199
in granule matrix, **39**:198
purification and properties, **39**:197–198
ribonuclease activity, **60**:194
sharing chromosome 14q with ECP, **60**:193
- Eosinophil differentiation factor, **48**:78

- Eosinophilia**
- experimental, IL-5 role, **57:173**
 - helper T cell cytokines and, **46:133**, **137**
 - IL-5** role
 - asthmatic patients, **57:174–176**
 - experimental, **57:173**
 - guinea pig, **57:173–174**
 - mRNA expression in patients, **57:158–159**
 - parasite infection association, **57:172**
 - tumors associated with, **57:177–178**
 - pulmonary, including Churg–Strauss syndrome, **60:226**
 - Eosinophilia myalgia syndrome**, from ingestion of L-tryptophan, **60:224–225**
 - Eosinophil mediators**
 - cytokines, **60:195–201**
 - granule proteins, **60:191–194**
 - lipid, **60:188–191**
 - release, and cytotoxicity, **60:205–208**
 - Eosinophil myelocytes**, morphology, **60:155–158**
 - Eosinophil peroxidase**
 - antihelminthic activity, **60:212**
 - binding to
 - effector cells, **39:202**
 - mast cell granules, **39:201–202**
 - concentration in BAL, **60:171**
 - deficiency, **60:227**
 - in cat family, **39:200**
 - in granule matrix, **39:200**
 - in humans, Yemenite Jews, **39:200–201**
 - IgE effects, **60:207**
 - leukotriene inactivation, **39:202–203**
 - oxidation of halides, **60:194**
 - parasite killing and, **39:201**
 - purification and properties, **39:199–200**
 - tumor cell killing and, **39:202**
 - Eosinophils**, **51:331–336**
 - activation, **60:201–210**
 - morphology, **60:158–160, 202–203**
 - activation by
 - α -CSF, **39:215**
 - EAF from monocytes, **39:215–216**
 - ECF-A from mast cells, **39:214–215**
 - ESP from lymph node cells, **39:214**
 - LTB₄, **39:215**
 - M-ECEF from monocytes, **39:216**
 - T cell-produced factor, **39:217**
 - TNF**, **39:216**
 - worm-produced factors, **39:217**
 - adhesion and migration, **60:164–178**
 - and adhesion mechanisms, **51:333–336**
 - as antigen-presenting cells, **60:200**
 - asthma, **62:283, 285**
 - autofluorescence, **39:205**
 - biology, overview, **60:151–155**
 - cachectin and, **42:224**
 - in cardiac disease, **39:226–227**
 - CD23 antigen and, **49:154, 161, 167**
 - chemokine action on, **55:117**
 - complement receptor 1 and, **46:190, 191**
 - in cutaneous diseases, **39:225–226**
 - and cytokines, **60:203–205**
 - cytotoxicity and, **41:269, 270, 291, 312**
 - degranulation
 - arachidonic acid metabolism and, **39:207**
 - in diseased tissues, **39:208, 223–226**
 - induction, methods of, **39:206–207**
 - differentiation and maturation, **60:162–164**
 - discovery, **39:177**
 - and disease, **60:216–227**
 - eicosanoid production, **39:204, 218**
 - enzymes
 - arylsulfatase, **39:218**
 - collagenase, **39:203**
 - EPO, **39:199–203**
 - variety of, **39:203–204**
 - granule proteins
 - ECP, **39:191–197**
 - EDN, **39:197–199**
 - major basic protein, **39:185–189**
 - high-density
 - activation, **39:205–206**
 - eosinophilia and, **39:205**
 - partial degranulation, **39:206**
 - in human reproduction
 - cyclic eosinopenia, ovulation and, **39:228**
 - major basic protein production by placenta, **39:228**
 - IgE, **43:306–307**
 - IL-1 and, **44:164, 188**
 - IL-5-mediated production, **57:156**
 - in immediate hypersensitivity, **39:217–219**
 - histamine release and, **39:217–218**
 - leukotrienes and, **39:218**
 - major basic protein and, **39:219**
 - mature
 - granule populations, **60:153–154**
 - morphology, **60:155–158**

- membrane proteins
 antigens, **39**:183–184
 CLC, **39**:181–183
 lysophospholipase activity, **39**:182–183
 complement receptors, **39**:179–180
 Ig receptors, **39**:179–181
 receptor for sheep erythrocytes, **39**:181
 PAF production, **39**:205
 parasite killing
 AES effects, animals, **39**:209
 cationic protein release and, **39**:213–214
 by moving into tissues and degranulation, **39**:212–213
 in vitro, mediated by
 C3, **39**:210
 IgE, **39**:210–211
 IgG, **39**:209–211
 IgG2a, **39**:210–212
 and parasites, **60**:210–216
 phagocytosis, **39**:208
 proliferation, GM-CSF, **48**:75
 recruitment, **58**:431
 in rodent uterus
 estrogen-induced eosinophilia and
 edema, **39**:227–228
 infiltration during estrus cycle, **39**:227
 role in asthma, fiberoptic bronchoscopy
 studies, **60**:218–223
 signal transduction, **60**:208–210
 sulfated complex carbohydrates, **39**:204
 toxicity to various cells, **39**:219
 ultrastructure, **39**:177–179
- Eosinophil stimulation promoter
 eosinophil activation, **39**:214
 production by lymph node cells, **39**:214
- EPA, *see* Eicosapentaenoic acid
- Epidermal cells
 dendritic
 expressing Ly-5, **58**:299
 reactivity, **58**:320
 Thy-1⁺, **58**:301–303
 V γ 5V δ 1, colonization of skin, **58**:317
- Ig gene superfamily and, **44**:31
- Epidermal growth factor
 autoimmune thyroiditis and, **46**:272, 294, 295
 complement receptor 1 and, **46**:187
 connection with NO production, **60**:350–351
 IL-1 and, **44**:168, 169, 184
 leukocyte integrins and, **46**:159
- Epidermal growth factor domain, PGHS, **62**:169, 171
- Epidermal growth factor receptor
 activation of Stats and Jaks, **60**:27
 in chimeric receptors, **60**:10
- Episalin, **62**:223
- Episomal vectors, **61**:124–128
- Epistasis, cytotoxic allogeneic T cells and
 allogeneic cross-reactive lysis, **38**:185–187
 H-Y antigen, **38**:175–180
 immunodominance, **38**:180–185
- Epithelial antigens, **62**:238–239
- Epithelial barriers, IFN- γ , **62**:90
- Epithelial cells
 and basement membrane, **51**:324–327
 CD1 protein distribution, **59**:56, 68, 72
 CD40 expression, **61**:14, 41
 corneal
 accessory antigen-processing cell, **48**:202
 derived T cell-activating factor, **48**:201–202
 hybrid resistance and, **41**:335
 IBD and, **42**:292–294, 319–320, 321
 interaction with IgA, **40**:174–177
 positive selection, **59**:120–121
 respiratory tract, **59**:372–373, 382
 as source of chemokines, **55**:107
 T cell activation and, **41**:1, 10
 T cell subsets and
 cell surface molecules, **41**:46
 restricted T cells, **41**:103–106
 T cell development, **41**:96–98
 T cell specificity, **41**:113
 tolerance induction, **41**:107–110
 thymic, induction of positive selection, **58**:164–165
 and thymocyte deletion, **58**:170
 thyroid, *see* Thyroid epithelial cells
- Epithelial tumors, antigens, **62**:220–221, 223–224
- Epithelium
 antigen-presenting cells and, **47**:54
 autoimmune thyroiditis and, **46**:266, 287, 296, 297
 CD23 antigen and, **49**:150, 155, 161, 170–171
 complement receptor 2 and, **46**:204, 210, 213
 corneal, Langerhans cell, **48**:197

- Epithelium (*continued*)**
- development in MHC class I-deficient mouse
 - intestinal, 55:395–396
 - thymic, 55:396–398
 - intestinal, derived T cell lineages, 58:110–112
 - leukocyte integrins and, 46:164, 172
 - SCID and, 49:384, 393, 402
- Epitopes**, 42:71–72; 43:106, *see also*
- Sequential epitope in AChR
 - B cells, 42:258
 - T cells, 42:257
 - adoptive T cell therapy of tumors and, 49:334
 - antigen recognition, 49:321–324
 - expression of antitumor responses, 49:329–330
 - mechanisms, 49:312–313, 315
 - agglutination, 48:23
 - amino acids, 43:65–67
 - antigen-presenting cells and, 47:60, 62, 63
 - anticellular antibodies and, 44:118, 119, 127
 - autoantigen, 44:128–130
 - autoimmune response, 44:132, 134–136
 - SLE, 44:97, 100, 101, 108
- aPL antibodies and, 49:193, 234–235, 239–240, 259
- affinity purification, 49:228
 - antibody subsets, 49:232–233
 - binding to cell membranes, 49:252
 - clinical aspects, 49:213
 - LA antibodies, 49:241, 246–248
 - pathogenic potential, 49:256–257
 - specificity, 49:234–235, 239–240, 248, 250
- autoimmune demyelinating disease and multiple sclerosis, 49:369–370, 372, 374
- myelin basic protein, 49:357–363
 - TCR usage restriction, 49:363–368
- autoimmune thyroiditis and, 46:318, 319
- antigens, 46:270, 271, 273
 - cellular immune responses, 46:291
 - experimental models, 46:278, 282
 - humoral responses, 46:302–305, 307
 - prevention, 46:312, 314, 315
 - autoreactive T cells and, 45:420
- B cell, 43:78
- multiple antigen peptides containing, 60:109–124
- B cell formation and, 41:191, 192
- binding energy contribution, 43:24
- CD4 and CD8 molecules and, 44:268, 295
- CD5 B cell and, 47:155
- CD23 antigen and, 49:157
- cell surface-expressed, 48:114
- character, 43:121–122
- complement receptor 1 and, 46:196
- complement receptor 2 and, 46:206, 207
- conformational, 48:114
- sites mapped by peptides cluster into, 43:38–39
- crystallographically defined, 43:24–25
- cytotoxic T cells and, 41:138, 169, 170
- $\beta 2$ -microglobulin, 41:155
 - amino acid changes, 41:159, 162, 163
 - carbohydrate moieties, 41:152, 154
 - exon shuffling, 41:139–142, 144, 145, 147
 - HLA class 1 antigens, 41:150, 151
 - monoclonal antibodies, 41:156–158, 165–167
- definition, 43:25–26, 72
- dominant, 43:46–49
- epicenter, 43:27
- flexibility, 43:63, 119
- helper T cell cytokines and, 46:124
- HIV CTL, 66:276–280
- HIV infection and, 47:384, 386
- immune response, 47:407, 410, 411
- HLA class II, 48:112
- humoral immune response and
- antigens, 45:8, 19
 - cellular interactions *in vivo*, 45:80
 - helper T cells, 45:28
 - physical interaction, 45:44
- Ig gene superfamily and, 44:11, 32, 33
- Ir* genes and
- antigen binding, 38:107
 - competitive inhibition, 38:108
- leukocyte integrins and, 46:168, 173
- lymphocyte homing and, 44:321, 342, 343, 348, 368
- methods, 43:75–76
- mobility, 43:121
- murine lupus models and, 46:74, 76

- NK cells and, surface antigens of, **42**:187; **47**:221, 245, 247, *see also* Natural killer cells
observed and predicted, **43**:52–55
overlapping, fine specificity studies, **43**:25
precipitation, **48**:23
prediction, **43**:56–58
sequence variability, **43**:77
sequential, *see* Sequential epitope
shared, RA susceptibility associated with concept, **56**:403–404, 416–418
DRB1 genes, **56**:410–416
ethnic differences, **56**:408–410
haplotype encoding, **56**:438–440
size, **43**:120–121
spontaneous autoimmune thyroiditis and, **47**:462
sporozoite malaria vaccine and, **45**:283, 319–322
CS proteins, **45**:292, 299
CS-specific T cells, **45**:307, 308, 310–313
endemic areas, **45**:315
human trials, **45**:313
interferon- γ , **45**:306
Plasmodium vivax, **45**:317, 318
surface antigens of human leukocytes and, **49**:87, 125
synthetic T and B cell sites and, **45**:195–197, 260–264
antigens, **45**:236
bacterial antigens, **45**:230–232
globular protein antigens, **45**:210–214
immunological considerations, **45**:201, 202
parasitic antigens, **45**:228–230
peptides, **45**:250
vaccines, **45**:252, 255–258
viral antigens, **45**:217–220, 222, 223
T cell, *see* T cell epitopes
T cell activation and, **41**:1, 9, 11, 20
T cell receptor and, **45**:142
amphipathic character, **45**:144, 145
antigen processing, **45**:121, 122
conformation, **45**:145, 146
experimental systems, **45**:132
helical conformation, **45**:142–144
location, **45**:146–148
peptides, **45**:137
structure-function relationships, **45**:135
T cell repertoire, **45**:164–166
T cell subsets and
cell surface molecules, **41**:44, 47
H-2 alloantigen recognition, **41**:79, 80, 83
H-2-restricted antigen recognition, **41**:53, 54, 56, 62
T cell specificity, **41**:111, 112
T helper, pathogen-derived, **60**:140–141
three-dimensional structure, **43**:76–77
as vaccines, **43**:59–60
virus-induced immunosuppression and, **45**:363–365, 367
EPO, *see* Eosinophil peroxidase
Epo
relation to tyrosine phosphorylation, **60**:5–6
role in Stat phosphorylation, **60**:22
Epo receptors
association with Jak2, **60**:11–12
cytoplasmic domain, **60**:13–14
induction of dimerization/oligomerization, **60**:9–11
Epstein–Barr virus, **52**:193–195, 464; **54**:143–144
abnormal infection, associated disorders, **37**:129–138
adoptive T cell therapy of tumors and, **49**:283
antibodies from transformed cell lines, **57**:229–230
antigen-presenting cells and, **47**:63, 99
associated lymphoproliferative disease in hu-PBL-SCID mice, **50**:310–313
autoimmune thyroiditis and, **46**:300, 306
CD4 molecules and, **44**:290
CD5 B cell and, **47**:136
CD23 antigen and, **49**:149, 161–162, 167, 171
complement receptor 1 and, **46**:192
complement receptor 2 and, **46**:204, 205, 208, 210–213
cytotoxic T cells and, **65**:279
cytotoxic T cells and, **41**:144, 165, 167
and herpesvirus saimiri, **54**:115–116
HIV infection and, **47**:397, 400
HLA and, **65**:286
human B cell neoplasia and, **38**:271
Burkitt lymphoma, **38**:246, 247, 254

- Epstein-Barr virus (*continued*)
 transformed lymphoblastoid cell line, **38:256**
 humoral immune response and, **45:16, 17**
 Ig heavy-chain variable region genes and, **49:2, 33, 35, 38, 42, 49-50**
 Ig production induced by, and
 immortalization, **37:110-112**
 IL-1 and, **44:156, 190**
 induced disease, **54:143-144**
 infectious mononucleosis and, **37:112-115**
 leukocyte integrins and, **46:173**
 mAbs and, **38:278, 279**
 B cell hybridomas, **38:280**
 donor lymphocytes, **38:284, 285**
 genetically engineered antibodies, **38:304**
 human-human hybridomas, **38:292, 293, 300**
 human-mouse heterohybridomas, **38:301**
 NK cells and, **47:224, 227, 267, 283, 292**
 persistent infection in normal individuals, **37:122-129**
 polyclonal B cell activation by, **37:102-110**
 QKRAA
 Dw4 peptide, **48:130**
 gp 110 peptide, **48:130**
 regulators of complement activation and, **45:388, 409**
 SCID and, **49:398**
 surface antigens of human leukocytes and, **49:92**
 synthetic T and B cell sites and, **45:226, 240, 241**
 T cell receptor and, **45:138**
 transformation mediated by
 genetically engineered antibodies, **38:305**
 human-human hybridomas, **38:293, 294**
 human-murine hybridomas, **38:289**
 monoclonal antibody production, **38:277-279, 301**
 transformed B cells, NK clones and, **42:182-184**
 tumor antigen induction, **57:292**
 virus-induced immunosuppression and, **45:351, 352, 356, 358**
 Equilibrium dissociation constant, **42:168-172; 43:101**
 Equine infectious anemia virus, **52:431**
 Erg, **64:65**
 Erythroblasts, hybrid resistance and, **41:376**
 Erythrocytes, *see also* Sheep red blood cells
 with abnormalities in complement interaction, **60:74-75**
 aPL antibodies and, **49:231, 239, 250-252, 254-255**
 B cell formation and, **41:189, 208**
 CD23 antigen and, **49:157**
 CD59 deficiency, **60:76-77**
 cell-mediated killing and, **41:289, 290, 293, 295**
 complement receptor 1 and, **46:190, 191, 194, 196, 198, 199**
 complement receptor 2 and, **46:203, 206, 210**
 cytosis and, **41:277, 278, 280**
 equine, T cell subsets and, **41:68**
 GPI anchor-deficient, **60:93**
 humoral immune response and, **45:17, 63, 65**
 hybrid resistance and, **41:337, 342, 349, 351, 352, 359, 376, 386, 388, 400**
 IgE biosynthesis and, **47:8, 12, 19**
 IL-8 binding to, **55:128**
 interaction with IgA, **40:170-174**
 leukocyte integrins and, **46:163**
 lysis by complement, protection from, **60:77-78**
 membrane, LPS-binding proteins, **53:279**
 membrane attack complex of complement and, **41:299-301, 303, 304, 306**
 pore formers and, **41:314, 316, 317, 319**
 protein roles, **45:388, 389**
 regulators of complement activation and
 protein expression, **45:395, 396, 399, 400, 402**
 SCID and, **49:383, 388**
 surface antigens of human leukocytes and, **49:114-115, 117**
 types I and II, **60:89**
 Erythroid cells, B cell formation and, **41:186, 213**
 Erythroleukemia, hybrid resistance and, **41:336, 372, 378**
 Erythropoiesis
 hybrid resistance and, **41:336, 337, 339, 348**
 IFN- γ and, **62:93**
 Erythropoietin
 hybrid resistance and, **41:348**
 properties, **39:2-3**

- Escherichia coli*, **41**:302, 318; **50**:162; **54**:174, 236
 expression library, immunoscreening, **54**:13
 genetically engineered antibodies and, **38**:304
 expression, **44**:72, 78
 fusion proteins, **44**:85
 production, **44**:66–69
 human-mouse heterohybridomas, **38**:300, 301
 IBD and, **42**:290–291, 294
 leukocyte integrins and, **46**:163
 lipopolysaccharide, structure, **53**:269–270
 maps of Ig-like loci and, **46**:5
 maternally transmitted antigen, **38**:344
 sporozoite malaria vaccine and, **45**:315
 synthetic T and B cell sites and, **45**:257
 T cell receptor and, **45**:126, 165
E-Selectin, **59**:383; **62**:259, 267, 277; **64**:144, 145; **65**:354, 355, 363, 367
 crystal structure, **64**:172
 downregulation, **64**:172
 downregulation by degradation in lysosomes, **58**:357–358
 genetic studies, **64**:194
 ligands, **64**:150, 151
 CLA, 250-kDa, and SSEA-1, **58**:361–362
 CLA⁺ memory T cells, **58**:392
 properties, **64**:147
 soluble form, **64**:171, 317
 upregulation, **64**:165
ESL-1, **64**:146, 151, 173
ESP, *see* Eosinophil stimulation promoter
Esterase
 B cell formation and, **41**:212, 213
 cell-mediated killing and, **41**:287
Esterase inhibitors, MC and, **42**:237
Ester linkage, in chemoimmunoconjugation, **56**:323
Estradiol
 depletion of CD4⁺8⁺ thymocytes, **58**:227–229
 hybrid resistance and, **41**:374, 389–392
Ethanolaminephosphate, in GPI-anchored proteins, **60**:60–67
Ethnicity, *see also* Race
 RA susceptibility associated with, **56**:408–410
N-Ethylcarboxyamido-adenosine, **52**:397
Etodolac, **62**:186
Ets-1, **62**:42, 47–48; **63**:250
 binding sites, **64**:70–73, 76–77
 domains, **64**:76–77
 expression, **64**:74–77
 gene, **64**:75
 gene targeting, **64**:81–82, 84–85
 oncogenic potential, **64**:65
 structure, **64**:79
 structure-function studies, **64**:79–81, 83–84
Ets-2
 binding sites, **64**:70–73, 76–77
 building sites, **64**:76–77
 domains, **64**:76–77
 expression, **64**:74–77, 82–83
 gene, **64**:84
 building sites, **64**:76–77
 gene targeting, **64**:81–82, 84–85
 oncogenic potential, **64**:65
 structure, **64**:79
 structure-function, **64**:79–81, 83–84
Ets binding sites, **64**:68–74
Ets domain, *PU.1P* domain, **63**:217, 222
Ets-Jun interaction, **64**:86
Ets transcription factors, **64**:65–66, 94–96
 binding sites, **64**:66, 68–74
 Elf-1 protein, **64**:76–77, 91–92
 Ets-1 protein, **64**:65, 74–82
 Ets-2 protein, **64**:76–77, 82–85
 Fli-1 protein, **64**:65, 76–77, 89–90
 GABP α protein, **64**:66, 76–77, 94
 PU.1 protein, **64**:65, 66, 76–77, 85–87
 Spi-B protein, **64**:76–77, 87–88
 ternary complex factors, **64**:76–77, 93–94
Eukaryotes, maternally transmitted antigen and, **38**:319, 320
Eu protein sequence, Ig heavy-chain variable region genes and, **49**:6, 8
Evolution
 CD23 antigen and, **49**:165
 Ig heavy-chain variable region genes and, **49**:14, 16, 26, 55, 61
J558 V μ family, **42**:123–127
 surface antigens of human leukocytes and, **49**:79, 117
V μ loci, **62**:19–26
Exocytosis
 autoimmune thyroiditis and, **46**:271
 IL-1 and, **44**:163

- Exocytosis (*continued*)**
- lymphocyte homing and, **44**:349
 - in neutrophils, IL-8-dependent, **55**:113
- Exoerythrocytic forms, sporozoite malaria**
- vaccine and, **45**:284, 319
 - antibodies, **45**:301
 - CS-specific T cells, **45**:307, 312
 - human trials, **45**:314
 - immunity, **45**:285, 286, 289–291
 - interferon- γ , **45**:301–303, 305, 306
- Exogenous help, T cell subsets and**, **41**:84, 85, 87, 112
- Exons**
- CD1 genes, **59**:7–8
 - CD4 molecules and, **44**:287–289
 - CD8 molecules and, **44**:274, 276, 278
 - complement components and, **38**:211, 213, 214, 236
 - genetically engineered antibody molecules and
 - biological properties, **44**:82, 83
 - cloning, **44**:73, 75
 - fusion proteins, **44**:86
 - humoral immune response and, **45**:32
 - Ig gene superfamily and
 - evolution, **44**:43–45, 47, 48
 - homology unit, **44**:8, 9
 - nonimmune receptor members, **44**:31, 34, 39
 - IL-1 and, **44**:157
 - regulators of complement activation and, **45**:404–406, 411
 - T cell receptor and, **45**:136
- Exon sequences, T cell receptor protein and**, **38**:6, 7
- Exon shuffling, CTL and**, **41**:144–148, 169
- altered cytoplasmic regions, **41**:148, 149
 - class I gene transfection, **41**:138, 139
 - CTL recognition, **41**:141–144
 - mAbs, **41**:156, 165
 - serology, **41**:139–141
- Experimental allergic autoimmune**
- encephalomyelitis, synthetic T and B cell sites and, **45**:235, 236
- Experimental allergic encephalitis, pituitary hormones**, **63**:424
- Experimental allergic encephalomyelitis**, **46**:79, **49**:357
- antibodies and, **49**:369–375
- myelin basic protein and**, **49**:357–358, 360–362
- TCR usage restriction and**, **49**:363, 366–368
- Experimental autoimmune encephalitis**, myelin basic and proteolipid proteins and, **66**:88, 90
- Experimental autoimmune encephalomyelitis**, **46**:291, 293, 312; **47**:433, 449; **48**:163–167; **62**:98–99, 105, 107
- anti-CD4 antibody, **48**:164, 166
 - anti-Ia antibody, **48**:164, 166
 - immunomodulation, **48**:166–167
 - inflammatory demyelination, **48**:166–167
 - T cell receptor V gene, **48**:166
 - TH cell receptor, **48**:166
- Experimental autoimmune myasthenia gravis**, *see also Acetylcholine receptor*
- Myasthenia gravis**
 - acute phase response in, **42**:260–261
 - development of, **42**:232
 - diagnosis of, **42**:254
 - etiology of, **42**:248–250
 - penicillamine and, **42**:235
 - T cell proliferation and, **42**:256
 - therapy for, **42**:263–269
- Experimental autoimmune thyroiditis**, **46**:266, 273, 317, 318; **47**:433, 440, 442, 447
- antigens, **46**:267, 270
 - cellular immune responses, **46**:287, 290, 296–298
 - genetic control, **46**:284–287
 - humoral responses, **46**:301, 302, 306, 308
 - induction, **46**:276–282
 - models, **46**:273–275
 - prevention, **46**:308, 310–315
 - SAT, **46**:273–276
- Experimental autoimmune uveitis**, **62**:98–99
- immune privilege, **48**:216–219
 - induction, **48**:215–216
 - pathogenesis, **48**:215–216
 - S antigen, **48**:219
 - self-tolerance, **48**:216–219
 - T cell, **48**:216
- Experimental eosinophilia, IL-5 role**, **57**:173
- Extracellular domain**
- LIF receptor, **53**:39
 - modification, **54**:313–315
- Extracellular matrix**
- and CD44, **54**:282–291

- CD44 as receptor for hyaluronan, **54:284–287**
 eosinophil adhesion, **60:173–174**
 features of CD44 molecule mediating interaction with hyaluronan, **54:287–290**
 general features, **54:282–284**
 leukocyte integrins and, **46:149, 153**
 ligand molecules, **46:171**
 structure, **46:156–159**
 and metastasis, **54:318–322**
 production, TGF- β activation, **55:186**
 protein interaction with CD44, **54:290–291**
 regulation, and interaction with CD44, **54:302–318**
 mechanisms for regulating CD44 receptor function, **54:310–318**
 regulated hyaluronan binding, **54:302–308**
 and regulatory mechanisms of other adhesion molecules, **54:308–310**
 surface antigens of human leukocytes and, **49:82–86**
- Extracellular processing, synthetic peptides, **52:84–85**
- Extravasation
 activated T cells in skin, **58:392**
 lymphocytes, **64:142; 65:360, 365–367**
 lymphocytes in high endothelial venules, **58:347**
 T cells, and endothelium, **50:285–287**
- Extrinsic allergic alveolitis, **62:267**
- Extrinsic factors, TH responses, **61:357–358**
- Extrinsic pathway inhibition, aPL antibodies and, **49:256**
- Eye, *see also* Anterior chamber; Cornea
 anatomy, **48:192**
 autoimmune diseases, **48:216–219**
 HSV-1 infection, anterior chamber, **48:210**
 contralateral eye, **48:210**
 delayed-type hypersensitivity suppression, **48:210**
 HSV infection
 anterior chamber-associated immune deviation, **48:208–210**
 cytotoxic T cell, **48:210**
 delayed-type hypersensitivity, **48:210**
 immune privilege, **48:191–221**
 immune regulation, **48:191–221**
- self-tolerance, strategies, **48:217**
 trabecular meshwork, **48:206–207**
 transplantation, **48:191**
- F**
- Fab
 constant domain, **48:19**
 C portion, **48:20**
 elbow bend, **48:19**
 heavy chain, **48:19**
 hinge region structure, **48:21–23**
 internal motions, **48:19–21**
 light chain, **48:19**
 protein antigen antibodies, **48:20**
 switch region flexibility, **48:19**
 V portion, **48:20**
- Fab-lysozyme complexes, **43:121**
- Fab subunit, rotation modes, **48:33**
- Faciogenital dysplasia, **63:149**
- Factor B, complement components and, **38:203, 206, 236**
 C2 and, **38:207–209, 228**
 C4b-binding protein and, **38:235**
 cDNA, **38:209–211**
 cloning, **38:212–214**
 deficiency, **38:208**
 expression, **38:217, 218**
 restriction fragment length polymorphism, **38:215, 216**
- Factor H, **61:203, 204, 209, 221, 230–235**
 complement components and, **38:204, 211, 235, 236**
 complement receptor 1 and, **46:189, 197**
- Factor I, **61:210–214, 239–240**
 complement components and, **38:235**
 in therapeutic complement activation, **56:272–273**
- Factor IX, complement components and, **38:236**
- Factor X
 aPL antibodies and, **49:242–243, 256**
 complement components and, **38:236**
- Factor XI, **66:225**
 activation, **66:230, 237**
 platelet receptors, **66:244**
 properties, **66:226**
 structure, **66:230**

- Factor XIa
 inhibition, **66:239**
 platelet receptors, **66:244**
- Factor XII, **66:225**
 activation, **66:225**, 227, 234, 235–236
 autoactivation, **66:234–235**, 236, 245, 248
 endothelial cell receptor, **66:249–257**
 function, **66:241**
 interaction with cells, **66:244**, 246
 properties, **66:226**
 structure, **66:226–229**
- Factor XIIa, **66:225**, 228, 229
 inactivation, **66:238**
- Factor XII_I, **66:227–228**, 229
 function, **66:241**
 inactivation, **66:238**
- FADD, **61:31**
- Family studies, RA susceptibility, **56:433–435**
- Fanconi's anemia, **61:315**
- Farnesyltransferase inhibitors, in cancer therapy, **58:417**
- Fas, **57:129–131**, 129–140; **61:9**, 17, 30, 31
 apoptosis role, **57:129**, 133–134, 140
in vitro, **57:133–134**
in vivo, **57:133–134**
 autoimmune diseases associated with, **57:138**, 140
 dependent cell lysis by CD4⁺ CTLs, **60:304–305**
 dependent MHC class II-restricted CTLs, **60:303–310**
 expressing B cells, **60:292**
 expression, **57:130–132**
 gene, **61:9**, 10–11
 mutation in *lpr* mouse, **57:132–133**
 loss of function mutation, **57:139**
 mutation in *lpr* mouse, **57:132–133**
 physiologic roles, **57:137–139**
 signal transduction role, **57:134–135**, 139
 T cell development role, **57:137**
 transmission of death signal, **60:299**
- Fas/APO-1
 immunological memory and, **53:241**
 in myeloma cells, **64:259–260**
 in target cell killing, **60:294–296**
- Fasciitis, eosinophilic, **60:227**
- Fas–Fas-L signaling pathway, **61:32**, 37
- Fas–Fc chimeric protein, **60:295–296**
- Fas ligand, **57:130**, 135–139; **61:24**, 25, 26
 characteristics, **57:135–137**
- expression, **60:295–298**, 303
 expression in cytotoxic T cells, **57:138**
 mediated killing, **60:308–309**
- Fatty acids
 aPL antibodies and, **49:194–197**, 238, 251
 IL-2 and, **44:161**, 176
 N-3 polyunsaturated
 dietary
 coronary heart disease and, **39:145–146**
 hypolipidemic effects, **39:145–147**
 in mammalian phospholipids, dietary fish oil and, **39:160**
 metabolism, **39:159–160**
 N-6 polyunsaturated, metabolism, **39:159–160**
 NK cells and, **47:249**
- FBL, adoptive T cell therapy of tumors and antigen recognition, **49:320–321**, 324
 expression of antitumor responses, **49:325–329**, 331–332
 mechanisms, **49:300**, 302–304, 308–310, 313–315
 NK cells, **49:317–318**
 principles, **49:292**, 295–296, 298
- Fc
 Asn²⁹⁷ residue, carbohydrate unit, **48:27**
 CH2 domain lateral contact, **48:20**
 CH3 domain, **48:20–21**
 Dob, **48:21**
 hinge region structure, **48:21–23**
 internal motions, **48:19–21**
 Mcg, **48:21**
 rotational properties, **48:20**
 structure, considerations, **51:2–4**
 three-dimensional structure, **48:20**
- Fc-binding factors, **43:277**
- Fc carbohydrate, **48:27**
- Fc fragment, active sequence, **40:116–118**
- Fc oligosaccharide, protein surface, **48:28–29**
- Fc receptors, **43:277**, 305; **51:29–54**
 adoptive T cell therapy of tumors and, **49:312–313**
 bacterial, **51:56–60**, 56–61
 B cell formation and, **41:194**
 binding sites, localization, **40:68–69**
 cells bearing, IBD and, **42:297**
 cellular distribution, **40:63–68**
 characteristics, **57:1**
 characterization of, **40:68–70**

- complement activation by, **51:7–29**
 antibodies as activators of alternative pathway, **51:28–29**
 C1 activation, **51:21–25**
 C3 activation, **51:25–28**
 C4 activation, **51:25–28**
 cell lysis, **51:25–28**
 interaction of C1q and antibody, **51:8–21**
- complement receptor 1 and, **46:198**
 cytotoxicity and, **41:269, 271, 279, 296**
- Fc α R, **51:48–50**
 characteristics, **57:38–39**
 Fc α -IgA interaction at molecular level, **51:49–50**
 positive cells, IBD and, **42:290**
- Fc γ R, **51:29–41; 52:394–396**
 biological function, **57:69–70**
 characteristics, **57:2–3**
 signal transduction role
 mechanisms, **57:72–77**
 phosphorylation, **57:74–77**
 second messenger interactions, **57:72–74**
 structural factors, **57:82, 84–89**
- Fc δ R, **51:51**
 characteristics, **57:43**
- Fc ϵ R, **51:41–48; 62:77**
 characteristics, **57:30**
 IgE biosynthesis and
 antibody response, **47:8, 9**
 binding factors, **47:12, 15, 17, 19–23**
- Fc μ R, **51:50–51; 52:173**
 characteristics, **57:41–42**
- FcRI
 bacterial, **51:56–58**
 expressed by monocytes, **60:179–181**
- Fc α RI
 biochemical structure, **57:39–40**
 cell distribution, **57:40–41**
 characteristics, **57:39**
 ligand properties, **57:40**
 mAbs, **57:9, 40–41**
 molecular structure, **57:39–40**
 polymorphisms, **57:41**
- Fc γ RI, **51:30; 62:77**
 biochemical structure, **57:3–7**
 biological function, **57:69**
 cell distribution, **57:8**
 characteristics, **57:2–4, 4**
- gene structure, **57:4–5, 7**
 IgG interaction at molecular level, **51:30–34**
- Ig interactions, molecular basis, **57:48–54**
- ligand properties, **57:7–8**
 molecular structure, **57:3–7**
 monoclonal antibodies, **57:8–9**
 polymorphisms, **57:8–10**
 signal transduction role
 phosphorylation, **57:76–77**
 structural factors, **57:84**
 structure, **57:5, 7**
- Fc ϵ RI, **51:41–43**
 biochemical structure, **57:31–36**
 biological function, **57:70–71**
 cell distribution, **57:36–37**
 α chain, **52:379; 57:32–33**
 molecules related to, **52:383–384**
 β chain, related molecules, **52:**
 379–380, 383–384; **57:33–34**
- γ chain, **52:380–381; 57:34–36**
 characteristics, **57:30**
 control of synthesis and expression, **52:384–386**
 dependence on IgE, **52:385**
 half-life, internalization, degradation
 and reexpression, **52:385–386**
 synthesis during cell cycle, **52:385**
 synthesis during differentiation of
 basophils/mast cells from
 hemopoietic precursor cells, **52:384–385**
- crosslinking
 and basophil ILL production, **53:13**
 and factor-dependent mast cell
 lymphokine production (murine), **53:6**
 and ILL production by splenic and bone marrow Fc ϵ RI $^+$ cells, **53:10**
 and signaling of IL-4 production, **53:19–22**
 and transformed mast cell lymphokine production, **53:5**
- functional characterization, **52:**
 387–391
 binding to IgE, **52:387–388**

- Fc receptors (*continued*)
- effector mechanisms and functional changes of mast cells/basophils following activation through IgE binding sites, **52**:391
 - signal transduction and distal events, **52**:388–391
 - gene structure, **57**:31–34
 - IgG interaction at molecular level, **51**:43–46
 - Ig interactions, molecular basis, **57**:49–50, 61–68
 - ligand properties, **57**:36
 - molecular structure, **57**:31–36
 - monoclonal antibodies, **57**:9, 36–37
 - polymorphisms, **57**:37
 - polypeptide components α , β , and γ , **53**:21
 - positive cells
 - bone marrow, lymphokine production by (murine), **53**:9–14
 - human, lymphokine production by, **53**:14–15
 - signal transduction role
 - mechanisms, **57**:77–81
 - phosphorylation in, **57**:79–81
 - second messenger interactions, **57**:77–79
 - structural factors, **57**:89–90
 - structure, **57**:31–34
 - transcripts, **57**:31
- FcRII
- bacterial, **51**:58–59
 - expressed by resting eosinophils, **60**:179–181
- FcγRII, **51**:34–36; **62**:77
- biochemical structure, **57**:10–16
 - biological function, **57**:70
 - cell distribution, **57**:17–19
 - characteristics, **57**:2–3, 11
 - gene structure, **57**:11–13, 15–16
 - IgG interaction at molecular level, **51**:36–38
 - Ig interactions, molecular basis, **57**:49–50, 54–59
 - ligand properties, **57**:16–17
 - molecular structure, **57**:10–16
 - monoclonal antibodies, **57**:9, 17–19
 - polymorphisms, **57**:19–21
- signal transduction role
- phosphorylation, **57**:75–77
 - structural factors, **57**:82, 84–87
 - structure, **57**:12–13, 15–16
 - transcripts, **57**:13
- FcεRI, **51**:46–47
- biochemical structure, **57**:37–38
 - CD23 antigen and, **49**:149, 162–167
 - cell distribution, **57**:38
 - characteristics, **57**:30
 - gene structure, **57**:37
 - IgE interaction at molecular level, **51**:47–48
 - ligand properties, **57**:38
 - molecular structure, **57**:37–38
 - monoclonal antibodies, **57**:38
 - structure, **57**:37
- FcRIII
- bacterial, **51**:59–60
 - eosinophils expressing, **60**:205
 - neutrophils and NK cells expressing, **60**:179–181
- FcγRIII, **51**:38–40; **62**:77
- biochemical structure, **57**:21–22
 - biological function, **57**:70
 - cell distribution, **57**:28–29
 - characteristics, **57**:2–3, 22
 - gene structure, **57**:22–26
 - IgG interaction at molecular level, **51**:40–41
 - Ig interactions, molecular basis, **57**:49–50, 59–61
 - ligand properties, **57**:27–28
 - molecular structure, **57**:21–27
 - monoclonal antibodies, **57**:9, 28–29
 - polymorphisms, **57**:29–30
 - signal transduction role
 - phosphorylation, **57**:76–77
 - structural factors, **57**:84, 87–88
 - structure, **57**:23–26
 - transcripts, **57**:23
- FcRIV, bacterial, **51**:61
- FcRV, bacterial, **51**:61
- FcRVI, bacterial, **51**:61
- FcRn, characteristics, **57**:46–47
- function
 - biological, **57**:69–71
- FcγR, **57**:69–70
- FcεRI, **57**:70–71
- signal transduction
 - mechanism, **57**:71–81
 - structural basis, **57**:81–90

- genetically engineered antibodies and, **44**:68, 80, 83, 87
human leukocyte, **51**:29–54
hybrid resistance and, **41**:371, 376–378, 394
Ig gene superfamily and, **44**:39, 43, 45
Ig interactions, molecular basis
 Fc γ RI, **57**:48–54
 Fc ϵ RI, **57**:49–50, 61–68
 Fc γ RII, **57**:49–50, 54–59
 Fc γ RIII, **57**:49–50, 59–61
 overview, **57**:48–50
leukocyte, effector cell-target cell
 interaction mediated by, **51**:51–54
lung dendritic cells, **59**:379–380
phagocytosis and
 IgG, **38**:362, 367, 392
 nonelicited murine macrophages, **38**:377, 379
 zymosan, **38**:370, 371, 393
polymeric IgR, characteristics, **57**:43–46
research directions, **57**:91
signal transduction, mechanism,
 57:71–81
 phosphorylation role, **57**:74–77, 79–81
 protein kinase role, **57**:74–77, 79–81
 second messenger interactions,
 57:72–74, 77–79
signal transduction, structural basis,
 57:81–90
 antigen receptor homology motif role,
 57:81–83
 Fc γ R role, **57**:82, 84–89
surface antigens of human leukocytes and,
 49:80, 83, 89–91, 111
T cell, role for isotype response,
 40:212–216
T cell activation and, **41**:9
Fc region, biological properties of
 activity of peptide fragments of Fc,
 40:111–119
complement fixation, **40**:70–71
enhancement of *in vitro* specificity of
 humoral immune response,
 40:90–95
enhancement of specific *in vitro* T cell-
 mediated immune response,
 40:95–100
nonspecific lymphocyte activation by Fc,
 40:79–90
regulation of cell-mediated immune
 response, **40**:77–79
regulation of specific humoral immune
 response, **40**:71–77
stimulation of macrophages by Fc region,
 40:100–111
FDC, *see* Follicular dendritic cells
Feedback, spontaneous autoimmune
 thyroiditis and, **47**:476, 478
Feedback loop, NF- κ B induction by cytokines
 and vice versa, **58**:7
Feline immunodeficiency virus, **52**:
 437–441
 LTR, **65**:10
 in study antiviral therapy and vaccines,
 52:440–441
Feline leukemia virus, model, **52**:
 434–436
 antiviral therapy, **52**:436–437
 pathogenic mechanism of FeLVFAIDS,
 52:436
Felton's paralysis, and other early results with
 polymers, **52**:287–288
FeLV, *see* Feline leukemia virus
FELV-FAIDS, pathogenic mechanism of,
 52:436
Ferritin, complement receptor 1 and,
 46:200
Fetal calf serum, autoreactive T cells and,
 45:426
Fetal development
 acquisition of clonotype repertoire during,
 42:26–30
 role of IL-6, **54**:32
Fetal loss, aPL antibodies and, **49**:252–253,
 257–258
 clinical aspects, **49**:213–216, 218–224
 isotype, **49**:225–226
Fetal tissues, transplantation with, SCID and,
 49:392–393
Fever
 cachectin and, **42**:224
 IL-1 and, **44**:154, 155, 190, 194
 biological effects, **44**:166
 immunocompetent cells, **44**:178
 systemic effects, **44**:170, 171
 TNF, **44**:185–187
Fiberoptic bronchoscopy, for obtaining BAL
 fluid, **60**:218–223
Fibrillarin, antinuclear antibodies and,
 44:122–124

- Fibrinogen
 IL-1 and, **44**:164, 189
Ir genes and, **38**:97
- Fibrinolysis, aPL antibodies and, **49**:254–255
- Fibrinolytic cascade, **66**:240–254
- Fibropeptide B
 helper T cell cytokines and, **46**:128
 human, synthetic T and B cell sites and, **45**:233
- Fibroblast growth factor, IL-1 and, **44**:153, 154
 biological effects, **44**:164, 167–169
 structure, **44**:158
- Fibroblasts
 B cell formation and, **41**:235, 236, 238
 genetically determined defects, **41**:230
 Ig genes, **41**:205
 lymphohemopoietic tissue organization, **41**:186, 188
- CD4 and CD8 molecules and, **44**:272, 302
- CD40 expression, **61**:15, 41
- CTL and, **41**:145, 150–153
- growth factor receptors, **54**:365
- hybrid resistance and
 antigen expression, **41**:400
 leukemia/lymphoma cells, **41**:359, 361
 NK cells, **41**:372
 in vitro assays, **41**:393, 394
- IFNs, **62**:61
- Ig gene superfamily and, **44**:36
- IL-1 and, **44**:153, 190
 biological effects, **44**:165, 166, 168, 169
 human disease, **44**:195
 receptor, **44**:179, 181, 183
 structure, **44**:158
- TNF, **44**:186, 188
- IL-1ra production, **54**:188–189
- induction of positive selection, **58**:164–165
- leukocyte integrins and, **46**:162, 171, 172
- maps of Ig-like loci and, **46**:6
- regulators of complement activation and
 protein expression, **45**:395, 397, 399
 protein roles, **45**:387
- RCA-like protein utilization, **45**:410
- as source of chemokines, **55**:107
- synthetic T and B cell sites and, **45**:217
- T cell activation and, **41**:10
- T cell receptor and, **45**:121
- T cell subsets and, **41**:46, 89–91, 94
- trans-acting negative factors, **43**:248
- Fibronectin, **38**:362
 antigen-presenting cells and, **47**:61
 B cell formation and, **41**:215
 eosinophil adherence to, **60**:173–174
 leukocyte integrins and, **46**:149, 153
 inflammation, **46**:168, 169
 leukocyte adhesion deficiency disease, **46**:176
 structure, **46**:157, 158
- lymphocyte homing, **64**:191
- multiple myeloma, **64**:248–249
- opsonic effects, **38**:388–391, 393
- structure, **38**:386–388
- Fibrosarcoma
 CTL and, **41**:45
 hybrid resistance and, **41**:359, 361, 363
- Fibrosis, cytokines, **59**:418
- Filariasis, major basic protein in blood, human, **39**:213
- Fine sequence specificity, switch recombination, **61**:107–108
- Fine specificity, **43**:3
 HyHEL5, **43**:26
 overlapping epitopes, **43**:25
- Fish oil, dietary
 arachidonic acid metabolism and, **39**:168–169
- coronary heart disease and, **39**:145–146
- guinea pig pulmonary response to antigen and, **39**:164–165
- plasma LTB₄ and, **39**:164
- murine SLE and, **39**:163–164
- rat rheumatoid arthritis and, **39**:163
- FIV, *see* Feline immunodeficiency virus
- FK506, **61**:180–187
 antiapoptotic effects, **58**:267
 effect on eosinophil survival, **60**:197–198
 inhibition of negative selection, **58**:261–262
- NF- κ B, **65**:120
- FK506-binding proteins, **61**:180–181
- FKBP12, **61**:181, 187
- FKBP12-rapamycin complex, **61**:181–190
- FLA-3, multiple myeloma and, **64**:247
- Flanking regions
 Ig heavy-chain variable region genes and, **49**:26, 39
 probes, **49**:23
- J*558 V_H subfamilies, **42**:122–123
- Flap endonuclease-1, **61**:304

- Flexibility
antigenicity, **43**:105, 118
epitope, **43**:57, 63, 119
loop, CDR, **43**:107
main-chain parameters, **43**:50
- Fli-1*, **64**:90
- Fli-1*
binding sites, **64**:76–77
domain, **64**:76–77
expression, **64**:89
gene targeting, **64**:90
oncogenic potential, **64**:65
structure, **64**:78
structure–function studies, **64**:89–90
- Florid glomerulonephritis, **59**:343
- Flow cytometry, histograms of antigen expression, **52**:131–135
- Fluorenyl-methoxy-carbonyl, *see* Fmoc chemistry
- Fluorescein, specific neonatal B cells, **54**:399
- Fluorescein isothiocyanate, CTL and, **41**:166, 167
- Fluorescence
antigen-presenting cells and, **47**:86, 94
antinear antibodies and, **44**:105
B cell formation and, **41**:196
CD5 B cell and, **47**:118, 127
cell-mediated killing and, **41**:298
complement receptor 2 and, **46**:204, 207, 211
genetically engineered antibody molecules and, **44**:65
immunosenescence and, **46**:229, 231, 232, 242
lymphocyte homing and, **44**:318, 323, 358, 359
NK cells and, **47**:198, 213, 219
spontaneous autoimmune thyroiditis and, **47**:454
T cell development and, **44**:237
- Fluorescence-activated cell sorter, T cell subsets and, **41**:84
- Fluorescence microscopy, CD5 B cell and, **47**:149
- Fluorescence polarization
IgA, rotational correlation times, **48**:16–17
IgE, rotational correlation times, **48**:16–17
IgM, rotational correlation times, **48**:16–17
- 5-Fluoro-2-deoxyuridine, use in cancer treatment, monoclonal antibody conjugation procedure, **56**:344–345
- Flurbiprofen, **62**:184, 186
IBD and, **42**:313
- FMLP, *see* Formyl-methionyl-leucyl-phenylalanine
- Fmoc chemistry
and Boc chemistry, HIV-1, **60**:134
MAP synthesis, **60**:106–107
Pf322 blood-stage antigen synthesis, **60**:120–121
with *tert*-butyl, **60**:128–130
TPI epitopes constructed with, **60**:125–126
- FOE receptors, **43**:282
degradation, **43**:302
human B cells, **43**:298–300
murine B cells, **43**:302
T cells, **43**:301, 303
- Folic acid antagonists, as cancer treatment, **56**:338–340
- Follicles, transformed, avian leukosis virus infection, response to, **56**:471–472
- Follicular B cell lymphoma, human, **50**:73
- Follicular B cells, **59**:283–285
- Follicular dendritic cells, **52**:225–227; **61**:33
antigen complex binding, **60**:269
CD40 expression, **61**:14, 41
cellular origin, **51**:261–263
central role of in lymphoid tissues
antigen expression of, **51**:257–261
cellular origin of, **51**:261–263
conclusions, **51**:278–280
distinction of from other dendritic cells, **51**:244–247
expression of in malignant lymphomas, **51**:274–275
process, **51**:243–244
regulation of normal germinal center by, **51**:263–274
apoptosis, inhibition of, **51**:272–274
in germinal center light zone, **60**:271
humoral immune response and, **45**:9, 24, 79, 80
role in lymphoid tissues
regulation of normal germinal center by binding via cell-cell and cell-matrix adhesion molecules, **51**:269–271
cytokines, induction of, **51**:274
requirements of FDC's for ontogeny of germinal center, **51**:263–265

- Follicular dendritic cells (*continued*)
 secondary immune response,
 regulation of, **51:271–272**
 trapping of immune complexes and
 alternative antigen, **51:265–269**
 as targets of HIV, **51:275–278**
 technical difficulties, solution of,
51:246–257
 stored antigen on, **53:222, 231**
- Follicular exclusion, autoreactive B cells,
59:310–319
- Folliculostellate cells, IFN- γ effects,
62:94
- Food poisoning, **54:128**
- Foot and mouth disease virus
 MAP vaccines, **60:130**
 synthetic T and B cell sites and, **45:227,**
253, 254
- Footprinting studies, **43:2–3**
- Forbidden T cells, in peripheral immune system, **53:164–166**
- N-Formyl-kynurenone, **62:71**
- Formyl-methionine peptides, and related compounds, receptors for, **52:371–372**
- Formyl-methionyl-leucyl-phenylalanine, **42:315; 64:154**
 5-lipoxygenase activation, **39:154**
- Formyl peptides
 comparison with leukotrienes in neutrophil activation, **39:135**
 as neutrophil receptor ligand, *see* Neutrophil receptors
- Forskolin, action of, **65:124–125**
- fos* gene family, **63:248**
- 4-1BB, **61:7, 10, 17**
 4-1BB-L, **61:24, 25**
- Four-way developmental choice, molecular basis of, **51:180–186**
 CD4 vs CD8 down-regulation, **51:185–186**
 positive selection vs two kinds of death, **51:175–177**
 positive vs negative selection, **51:180–183**
 signaling pathways controlling positive selection, **51:183–185**
- Fractionation
 antinuclear antibodies and, **44:102**
 lymphocyte homing and, **44:344**
 T cell development and, **44:238**
- Fragment cultures, specific, and limiting dilution, **54:395–398**
 importance of affinity and valence, **54:396–398**
 role of B cell maturational stage, **54:395–397**
- Fragmentins
 induced DNA degradation, **60:301**
 or granzymes, cytotoxic T cells containing, **58:240**
 peptide hydrolysis by, **58:243**
- FRAP, **63:154**
- FRAP1, **61:181**
- Free energy, protein-protdn interactions, **43:100**
- Free radicals
 formation or action, inhibition, **58:250**
 NO, scavenging, **60:340–341**
- Freund's adjuvant
 HIV V3 loop in, **60:131**
 multiple antigen peptides emulsified in, **60:111–112, 116–117**
- Freund's complete adjuvant
 autoimmune thyroiditis and experimental models, **46:276–282**
 genetic control, **46:286, 287**
 humoral responses, **46:301, 302, 307**
 prevention, **46:309–312, 314**
 helper T cell cytokines and, **46:137**
 IgE biosynthesis and, **47:2–5, 14, 24**
 T cell subsets and, **41:70, 71**
- Friend erythroleukemia, hybrid resistance and, **41:389**
- Friend leukemia virus, hybrid resistance and antigen expression, **41:400, 403**
 bone marrow cells, **41:345**
 lymphoid cells, **41:354**
 marrow engraftment, **41:387**
 NK cells, **41:376**
- Friend murine leukemia virus, adoptive T cell therapy of tumors and, **49:320–321, 324**
- Friend spleen focus-forming virus, hybrid resistance and, **41:389**
- Friend virus, **41:168, 378, 389**
- FS cells, IFN- γ effects, **62:94**
- Fucoidin, lymphocyte homing and carbohydrate, **44:356–358, 360–362**
 high endothelial venules, **44:320**
- Fucosyltransferases, **64:168, 169**
- Fuc-TVII, **65:364–365, 373**

- Functional divergence, and positive selection
T cell mechanism, **51**:186–189
- Functional redundancy, defined, **63**:173
- Fungal pneumonia, **59**:397–406
Cryptococcus neoformans, **59**:402–405
- Fungal products, in therapeutic complement inhibition, **56**:288–289
- Fungi, cytotoxicity and, **41**:269, 270
- Fusion
 antinuclear antibodies and, **44**:130, 131
 CD4 molecules and, **44**:295, 296
 donor lymphocyte preparation, **38**:282–286
 genetically engineered antibody molecules
 and
 expression, **44**:70, 71, 77
 production, **44**:67
 proteins, **44**:85, 86
 human-human hybridomas, **38**:297, 300
 human-mouse heterohybridomas, **38**:300,
 301
 hybridoma propagation, **38**:322
 maternally transmitted antigen, **38**:314,
 324, 341, 343
 phagocytosis, **38**:363
 somatic, B cell hybridization and, **38**:279
 T cell development and, **44**:215
- Fyn, **63**:135, 165, 399
 expression, **51**:144–145
 IL-2R β coupling, **61**:155
- G**
- G₁-phase progression
 IL-2 dependent, **61**:179–180
 rapamycin, **61**:182–183
- G63, **52**:394
- GABP α
 binding sites, **64**:76–77
 domains, **64**:76–77
 expression, **64**:94
 gene targeting, **64**:94
 structure, **64**:78
 structure-function studies, **64**:94
- gag, adoptive T cell therapy of tumors and,
49:320–321, 323–324
- GAGE, **62**:229, 233
- Galactans, phagocytosis and, **38**:368, 381
- Galactose, surface antigens of human
 leukocytes and, **49**:88–89, 91
- β -Galactosidase, antibody response in B cells
to, **39**:88–90
- helper T cells and, **39**:88–90
- GALT, *see* Gut-associated lymphoid tissue,
rabbit
- Gamma-activated factor, **62**:68
- Gamma chains
 γ_c -common chain, **61**:150, 151–152
 mast cell-specific receptor, gene cloning
 studies, **43**:288–289
- Gamma globulin
 agammaglobulinemia, history, **59**:137–141
 aPL antibodies and, **49**:222–223
 normal rabbit, presentation by B cells,
 39:55–56
 replacement therapy, **59**:150
- XLA, *see* X-linked agammaglobulinemia
- Gamma radiation, B cells
 antigen presentation and, **39**:67–72
 dose effects, **39**:67, 71–72
 LPS activation and, **39**:71
 MLR stimulation and, **39**:80–81
- Ganglionic nicotinic AChR, **42**:239, *see also*
 Acetylcholine receptor
- Ganglioside antigens, tumor antigens,
40:351–352
- cell-associated gangliosides, properties,
40:353–355
- tumor cell-associated, properties
 colon cancer, **40**:364–366
 melanoma, **40**:357–364
 neuroblastoma, **40**:355–357
- Gangliosides, **62**:219–220
 abnormal expression in PNH, **60**:92
- Gaps, **63**:144
- GAS-binding proteins, cytokine activation,
63:141
- GAS elements
 DNA-binding activity, **60**:21–22
 Stat dimer affinities for, **60**:24–25
- Gastrin receptor, **48**:181
- Gastrointestinal side effects,
 anticholinesterase drugs, **42**:270
- GATA-2, B cell genesis, **63**:217, 218–219
- GBP-1, **62**:76
- gC1aR, as endothelial cell receptor,
66:249–257
- G-CSF, *see* Granulocyte colony-stimulating
factor
- GD2, **62**:219

- GD3, **62**:219
GDP β S, **43**:292
 Gelatinase, **62**:288
 Gel electrophoresis
 Ir genes and, **38**:78, 147
 two-dimensional, **38**:225
 Gelonin, MG and, **42**:265, 266
 Gender, RA susceptibility studies, **56**:434
 Gene amplification, human B cell neoplasia
 and, **38**:254
 Gene conversion, **62**:24–26
 Ars A response and, **42**:120
 avian Ig gene, **48**:60–62
 chicken B cell genes, **57**:358–365
 complement components and, **38**:227
 Ig heavy-chain variable region genes and,
 49:15, 52, 61
 D segments, **49**:28–29, 32
 polymorphism of V_H gene segments,
 49:24, 26
 Ig VL gene, **48**:60–62
 Ir genes and, **38**:86
 somatic, and antibody repertoire generation
 in rabbit, **56**:198–202
 ψ-V gene segment, **48**:58
 VL gene segment, **48**:56–60
 Gene dosage
 hybrid resistance and, **41**:357
 Ir genes and, **38**:79–83, 139
 Gene duplication
 Ig heavy-chain variable region genes,
 49:25, 29
 in *J558* V_H family, **42**:123–127
 Gene expression
 B cell hybridomas, **38**:279
 donor lymphocytes, **38**:285
 genetically engineered antibodies, **38**:304
 induction via B cell antigen receptor,
 55:266–269
 maternally transmitted antigen and,
 38:314
 antigenic variation, **38**:342, 343
 MHC, **38**:329
 Gene fusion, resulting from chromosome
 translocation, **50**:128–129
 Gene knockouts, **59**:116–118, 124, 128
 Genealogy, *J558* V_H gene segments, **42**:124
 Generalized lymphoproliferative disease,
 murine lupus models and, **46**:63, 64,
 94–97
- Gene rearrangements
 cis-acting elements for, **58**:30–31
 DNA sequencing, **58**:33–36
 endogenous, **58**:37
 lymphoid-restricted, **58**:43
 TCR, **38**:8, 10, 11
 allelic exclusion, **38**:17, 18
 developmentally programmed,
 58:308–309
 peripheral T cells, **38**:13–15
 thymus, **38**:13
 TCR_y, **58**:142
 V α TCR, **58**:112
 V γ , developmentally regulated program,
 58:301
 V γ δ , **58**:311
 V γ γ , **58**:314
 Gene regulation, by cytokines, NF-IL6 and
 NF-κB, **65**:1–33
 Genes
 chicken B cell, conversion
 DT40, **57**:361–362
 Holliday junction, possible formation of,
 57:362–365
 hyperconversion mechanism, **57**:358–361
 recombination models, **57**:362–365
 cytokine
 cloning and expression, **58**:425–427
 in tumor cell engineering, **57**:321–323
 upregulated by NF-κB, **58**:10–12
 in DSB repair and V(D)J recombination,
 58:45–55
 in GPI anchor synthesis, **60**:67–74
 heavy chain, assembly of, **40**:252–253
 in hematopoietic cells, **60**:23–25
 of human κ light chains, **40**:249–251
 IL-2, localization, human, **39**:5, 18
 IL-3, localization, murine, **39**:7–8, 18
 λ light chain, **40**:251–252
 lymphocyte, function, **62**:31–53
 mb-1, genomic structure, **54**:355
 MHC, *see* Major histocompatibility
 complex
 natural and artificial mutations, **58**:90–101
 NF-κB-responsive, **58**:5–6
 non-MHC-linked, **51**:293–294
 oncogenes, *see* Oncogene
 perforin, **60**:291–294
 RA susceptibility-associated, alleles,
 56:441–446

- suicide, **58**:428–429
 transgenes, *see* Transgene
- Gene silencer, TCR γ , **58**:120
- Gene splicing, adhesion molecules, **64**:167
- Gene targeting
 Elf-1 protein, **64**:92
 Ets-1 protein, **64**:81–82
 Ets protein, **64**:84–85
 Fli-1 protein, **64**:90
 GABP α protein, **64**:94
 PU.1 protein, **64**:86–87
 Spi-B protein, **64**:88
 ternary complex factor, **64**:94
- Gene therapy, X-linked agammaglobulinemia, **59**:150–151
- Genetically engineered antibodies, **38**:304–306; **44**:65, 66, 89
 antigen-combining sites, **44**:84, 85
 biological properties, **44**:82, 83
 chimeric antibodies, **44**:79–82, 87, 88
 cloning, **44**:73–75
 domain structure, **48**:14, 16, 22
 expression, **44**:75–79
 expression vectors, **44**:70–72
 fusion proteins, **44**:85, 86
 lymphoid mammalian cells, **44**:69, 70
 production
 in bacteria, **44**:66–69
 in yeast, **44**:69
 segmental flexibility, **48**:14, 16, 22
 surface Ig, **44**:89
- Genetic chimerism, **59**:106, 109–111, *see also* Chimerism
- Genetic defects, in B lineage cells leading to autoimmune disease, **53**:144–147
- Genetic drift, *Ir* genes, **38**:87, 148
- Genetic mapping
 autoimmune demyelinating disease and, **49**:369
 CD1 loci, **59**:4–5
 CD23 antigen and, **49**:156, 165–166
 Ig heavy-chain variable region genes and, **49**:1, 29, 55
 organization, **49**:3–4
 V_H families, **49**:18, 20–21
- IL-2R genes, **59**:226, 227
- V_H locus, **62**:4–7
- Xid mutation, **59**:175
- XLA gene, **59**:163–165, 194–195
- Genetic maps, Ig-like loci, **46**:1–3
 chromosomal locations, **46**:6, 7
 mechanisms, **46**:45–47
 methods, **46**:3–6
 MHC, **46**:28
 human HLA, **46**:30–34
 mouse H-2, **46**:28–30
 structure, **46**:7
 human κ chain, **46**:15–19
 human λ chain, **46**:10–12
 human heavy chain, **46**:24–28
 mouse κ chain, **46**:12–15
 mouse λ chain, **46**:8–10
 mouse heavy chain, **46**:19–24
- T cell receptor, **46**:34, 35
 α -chain, **46**:41–45
 β -chain, **46**:35–37
 γ -chain, **46**:38–41
 δ chain, **46**:42, 43, 45
- Genetic mosaicism, X-chromosome, **59**:159, 195
- Genetic recombination, *see* Recombination
- Genetics
 adoptive T cell therapy of tumors and, **49**:283, 297
 aPL antibodies and, **49**:222, 255
 B cell progenitors, **59**:152
Btk gene, **59**:170–176, 184, 196
 mutations, **59**:185–192, 195, 317
 CD1 family, **59**:2–10
 CD23 antigen and, **49**:162
 and diabetes, **51**:290–294
 MHC-linked diabetogenic genes, **51**:290–293
 non-MNC-linked genes, **51**:293–294
 gene inactivation, **59**:108
 gene knockouts, **59**:116–118, 124, 128
 reconstitution, IL-2 receptors, **59**:231–234, 254
- rheumatoid arthritis susceptibility studies
 family, **56**:433–435
 female sex, **56**:434
 number of genes, **56**:434–438
 twins, **56**:433–435, 446–449
- somatic cell
 complement components and, **38**:204, 233
 human B cell neoplasia and, **38**:247

- Genetics (*continued*)**
- maternally transmitted antigen and, **38:320**
 - spontaneous autoimmune thyroiditis and, **47:481–491**
 - transgenes, **59:107–108, 113–116, 124–127**
 - X-linked agammaglobulinemia, **59:161–176, 187, 193–194**
- Gene transfer**
- as cancer therapy, **58:417–454**
 - for CD4/CD8 coreceptor function, **53:70–78**
 - retroviral-mediated, **58:437–438**
 - techniques, **58:422–425**
- Genome**
- c-kit*, organization of, **55:7–9**
 - stem cell factor, organization of, **55:13–15**
- Genomic mapping**, *see* Genetic mapping
- Genomic organization**
- T cell receptor protein and, **38:7**
 - TCR δ gene segments, **43:155–156**
- Genomic structure**, CD23 antigen and, **49:165–167**
- Genotype**, autoimmune demyelinating disease and, **49:364, 366**
- Geographic distribution**
- PIG-A* mutations, **60:81, 87**
 - Plasmodium vivax* malaria, **60:122**
 - PNH, **60:74**
- Germ cells**, development, stem cell factor receptor effect on, **55:66–69**
- Germinal center cells**, **62:146–148, 150**
- Germinal center reaction**, responses to thymus-dependent antigens, **60:272–276**
- Germinal centers**
- B cell censoring, **59:331–341**
 - development, **52:224–225**
 - diversification, **60:283–284**
 - formation, CD40 and, **63:72**
 - and memory B cells, **53:231–233**
 - normal, regulation of by FDCs, **51:263–274**
 - apoptosis, inhibition of, **51:272–274**
 - binding via cell-cell and cell-matrix adhesion molecules, **51:269–271**
 - cytokines, induction of, **51:274**
 - formation of, **51:243–244**
 - requirements of FDC's for ontogeny of germinal center, **51:263–265**
 - secondary immune response, regulation of, **51:271–272**
 - trapping of immune complexes and alternative antigen pathway, **51:265–269**
- ontogeny**, requirement of FDCs for, **51:263–264**
- pathways of cell differentiation**, **53:232–233**
- phenotypic selection**, **60:280**
- polarization**, **60:271–272**
- requirements of FDC's for ontogeny of**, **51:263–265**
- somatic hypermutation** in, **53:233–234**
- Germline**
- C_{H} transcription, **54:240–245**
 - counterparts, Ig heavy-chain variable region genes and, **49:25**
 - genes
 - V_{H} segment, **42:113–115**
 - $V_{\kappa}10$ family, **42:144–148**
 - sequences
 - Ig heavy-chain variable region genes, **49:27**
 - $J558 V_{\text{H}}$ gene, **42:127–128**
 - V_{H} segments, **62:1–26**
 - VL, sequence comparison, **48:55**
- Germline mutational approach**, **62:31, 34–36**
- Germline transcripts**
- CD40 signaling, **61:92–98**
 - class switching, cytokines, **61:84–85**
 - I_{μ} transcripts, **61:106–107**
 - regulation of transcription, **61:89–90**
 - structure and function, **61:89**
 - switch recombination, **61:111–116**
- GF109203X**, **63:155**
- GFA**, *see* Granulocyte functional antigen
- GH-N gene**, **63:384–385**
- Glia**, cells, retinal, T helper lymphocyte, **48:218**
- Glia growth-promoting factor**, **48:173**
- Glioblastoma cells**, immunological mediator, **48:168–169**
- Gliotoxin**, **65:126–127**
- Globular protein antigens**, synthetic T and B cell sites and, **45:203–216, 240**
- Glomerulonephritis**, **61:46, 245**
- florid, **59:343**
 - membranoproliferative, **61:245**
 - mesangial proliferative, and IL-6, **54:41**

- Glucagon, spontaneous autoimmune thyroiditis and, **47:440**
- β**-Glucan receptor, phagocytosis and alternative complement pathway activators, **38:285**, 386
- fibronectin, **38:390**
- particulate activators, **38:381**, 382, 393
- zymosan
- by human monocytes, **38:362**, 363, 365–372, 374–376, 392
 - by nonelicited murine macrophages, **38:377**, 379, 380
- α**-Glucans
- alternative complement pathway, **38:384**–386
 - phagocytosis, **38:366**–368, 381
- Glucocorticoid-inducing factors, spontaneous autoimmune thyroiditis and, **47:477**–479, 488
- Glucocorticoid receptors
- and antigen-driven T cell deletion, **58:258**–260
 - interaction with NF-IL6, **65:21**–22, 23
 - linomide effects, **58:264**–265
 - NF-κB and, **65:118**–120
 - and T cell fate determination, **58:137**–138
- Glucocorticoid response element, **65:22**
- Glucocorticoids, **62:259**; **65:21**, 22, 23
- cachectin and, **42:221**
 - CD5 B cell and, **47:145**
 - CD23 antigen and, **49:154**, 160–161
 - effect on CD4⁺8⁺ thymocytes, **58:166**–170
 - IgE biosynthesis and, **47:24**
 - immune system and, **66:197**, 218
 - induced thymocyte apoptosis, **58:245**–246
 - mode of action, **64:172**
 - NF-κB and, **65:118**–120
 - spontaneous autoimmune thyroiditis and, **47:493**, 494
 - disturbed immunoregulation, **47:476**, 477–479, 480
 - genetics, **47:488**, 489
- T cell death and, **58:227**–228, 269
- treatment with
- asthma and hypereosinophilic syndrome, **60:222**–224
 - eosinopenia ascribed to, **60:227**
 - in vitro* IgE synthesis, **61:370**–371
- Glucosamines, and GlcNAc, in GPI-anchored proteins, **60:60**–72
- Glucose, membrane attack complex of complement and, **41:300**
- γ-Glutamyl transpeptidase, LTC₄ conversion to LTD₄, **39:152**–153
- Glutaraldehyde
- in crosslinkage procedure for chemoimmunoconjugates in cancer treatment, **56:315**
 - T cell subsets and, **41:53**
- Gluten-derived peptide, celiac disease, **48:147**–148
- Gluten enteropathy, *see* Celiac disease
- GlyCAM-1, **64:149**, 168
- lymphocyte–HEV interaction, **65:361**–362, 363, 364, 373, 377
 - as selectin ligand, **58:359**
- Glycans, CTL and, **41:152**, 153
- Glycogen, cell-mediated killing and, **41:287**
- Glycolipid anchor, GPI, **60:57**–74
- Glycolipids
- aPL antibodies and, **49:228**–230, 239, 255
 - cell surface, **52:393**–394
 - membrane attack complex of complement and, **41:301**
 - surface antigens of human leukocytes and, **49:111**
- β**-Glycoprotein, complement receptor 1 and, **46:184**, 185
- β**-2-Glycoprotein-I, aPL antibodies and, **49:240**, 247, 252, 256–257, 259
- Glycoprotein antibodies, phase I clinical trials, **40:366**–370
- Glycoprotein antigens
- biosynthesis and intracellular transport, **40:337**–341
 - immunochemical and molecular profiles, **40:328**–337
- immunochemical profile, **40:326**–327
- immunological characterization, **40:324**–328
- preclinical models for immunotherapy
- in vitro* studies, **40:341**–345
 - in vivo* reactions in animal model systems, **40:345**–351
- Glycoproteins
- adoptive T cell therapy of tumors and, **49:320**
 - antigen-presenting cells and, **47:92**
 - aPL antibodies and, **49:239**–240, 255–256, 259

- Glycoproteins (*continued*)
 autoimmune thyroiditis and, **46:267**, 270
 B cell formation and, **41:189–193**, 227
 CD4 molecules and, **44:265**, 285, 295, 297
 CD5 B cell and, **47:118**, 146
 CD23 antigen, **49:149**, 155, 158
 complement receptor 2 and, **46:203**, 205, 210
 CTL and, **41:152**
 HIV and SIV, envelope variation, **52:451**
 HIV infection and, **47:378**, 395
 immune response, **47:405**, 407–409
 human T cell activation and, **41:2**, 4, 8
 humoral immune response and, **45:42**
 hybrid resistance and, **41:378**, 403
 Ig gene superfamily and, **44:39**, 40, 42
 IL-1 and, **44:165**, 166
 leukocyte integrins and, **46:149**, 151, 153
 ligand molecules, **46:171**, 172
 structure, **46:157**, 159
 lymphocyte homing and, **44:317**, 341, 350, 365, 369
 maps of Ig-like loci and, **46:34**
 myelin-associated, *see* Myelin-associated glycoprotein
 NK cells and, **47:298**
 antimicrobial activity, **47:286**, 287
 effector mechanisms, **47:237**, 248
 morphology, **47:218**
 surface phenotype, **47:207**
 regulators of complement activation and, **45:399**, 400, 410
 SCID and, **49:395**
 surface antigens of human leukocytes and, **49:112–114**, 118, 125
 adhesion molecules, **49:82–88**
 antigen-specific receptors, **49:76–78**
 complement components, **49:93**
 Igs, **49:90–91**
 MHC glycoproteins, **49:80–82**
 receptors, **49:94**, 98, 100
 synthetic T and B cell sites and, **45:225**, 226, 255
 T cell subsets and, **41:43**, 44, 46, 49
 virus-induced immunosuppression and, **45:350**
 HIV, **45:361**, 364–366
 measles, **45:339**, 346
 Glycosylation, **52:59–61**
 aPL antibodies and, **49:256**
 autoimmune thyroiditis and, **46:267**, 271
 B cells and, **41:190**, 192, 215
 CD4 molecules and, **44:285**
 CD8 molecules and, **44:271**, 273, 274, 277, 279
 CD23 antigen and, **49:156–158**, 162–164
 complement components and, **38:218**, 220
 complement receptor 1 and, **46:192**
 complement receptor 2 and, **46:202**, 204, 205
 CTL and, **41:135**, 138, 152, 153
 cytotoxicity and, **41:272**
 genetically engineered antibodies and, **38:304**; **44:69**, 76, 79, 80, 84
 humoral immune response and, **45:42**
 IgE biosynthesis and, **47:16**, 17, 20, 23, 25, 26
 Ig gene superfamily and, **44:35**, 39
 IL-1 and, **44:179**, 183, 194
 leukocyte integrins and, **46:154**, 170, 174
 lymphocyte homing and, **44:342**
 maternally transmitted antigen and, **38:331**
 patterns, cell-type specific, **54:314**
 regulators of complement activation and, **45:400–403**
 surface antigens of human leukocytes and, **49:93**, 99, 113
 T cell development and, **44:233**
 T cell receptor proteins, **38:4**, 5, 10, 18
 T cell subsets and, **41:41**, 50
 Glycosylation-enhancing factor, IgE
 biosynthesis and, **47:32**, 36, 38
 binding factors, **47:23**, 25, 26, 27
 Glycosylation-inhibiting factor, IgE
 biosynthesis and, **47:32–39**
 binding factors, **47:23–27**
 N-Glycosylation sites, PGHS, **62:173–174**
 Glycosylphosphatidylinositol anchor
 core backbone, **60:60–61**
 post-translational modification by, **60:65**
 anchored proteins, **60:57–74**, 79
 anchor precursor, biosynthesis, **60:63–65**
 anchor synthesis
 defective, **60:78–80**
 genes, **60:67–74**
 hydrolysis, signal transduction, **59:264–265**
 signal transduction, **59:264–265**
 surface antigens of human leukocytes and, **49:75**, 93–94, 112, 116, 125

- Glycosylphosphatidylinositol activation antigen, **48**:237–241
- Glycosyltransferase, *Ir* genes and, **38**:94
- Glycyrrhizin, **64**:196
- GM2, **62**:219
- GM3, **62**:219
- GM-CSF, *see* Granulocyte-macrophage colony-stimulating factor
- Goitrous thyroiditis, **46**:284
- Gold compounds, action of as rheumatoid arthritis treatment, **65**:124
- Colgi apparatus
- cytosis and, **41**:275, 281, 282
 - lymphocyte homing and, **44**:317, 320, 360
 - NK cells and 214, **47**:216, 303
 - T cell subsets and, **41**:57
- Gordon phenomenon, induction by EDN, animal, **39**:198–199
- Gout, inflammation in, **66**:260
- Gp, activation, **43**:290
- gp33
- covalent association of TCR β protein, **58**:150
 - gene, cloning, **58**:145
- gp39
- B cell proliferation, **63**:54
 - CD40-dependent B cell activation, **63**:56–57
 - cloning, **63**:46–48
 - expression, **63**:44, 48
 - gene, hyper-IgM syndrome, **63**:49–50
 - identification, **63**:46
 - molecular structure, **63**:48–49
 - structure, in hyper-IgM syndrome, **63**:49–50
- gp39-CD8 fusion protein, **63**:65
- gp48, **64**:116–117, 128
- gp75, **62**:229, 236–237
- gp96, **62**:244
- gp100, **62**:229, 235–236
- gp120
- binding by eosinophils, **60**:187
 - CD4-binding domain, **60**:138
 - HIV infection, **63**:84
 - induction of NO production, **60**:353–354
 - V3 loop, **60**:131–134
- gp130, **66**:127–128
- and high-affinity OSM/LIF receptor, **53**:43
 - IL-6-related cytokines and, **64**:256
- GPI, *see* Glycosylphosphatidylinositol
- GPI-H*, **60**:72–73
- CPL, aPL antibodies and, **49**:203–204
- G proteins, *see also* Coupling proteins
- in chemokine activation of integrin, **58**:379
 - coupling to chemokine receptors, **55**:138–139
- CTL and, **41**:154
- phosphoinositide-specific phospholipase, **48**:266
- role in IgR-mediated signaling, **54**:
 - 364–365
 - role in phosphoinositide hydrolysis, **43**:291–292
- SCID and, **49**:386
- surface antigens of human leukocytes and, **49**:76, 98, 111
- T cell, **48**:267–269
- T cell antigen receptor-CD3 complex, **48**:306–307
- Graft bed, corneal allograft
- afferent blockade theory, **48**:195–196
 - avascular, **48**:195
 - graft location, **48**:195
 - host cellular component, **48**:195
 - orthotopically placed, **48**:195
- Graft rejection
- adoptive T cell therapy of tumors and, **49**:301, 305, 328
 - antiadhesion therapy, **64**:197
 - hybrid resistance and
 - antigen expression, **41**:404–407
 - bone marrow cells, **41**:344, 350
 - effector mechanisms, **41**:370, 371
 - antibodies, **41**:376, 378
 - marrow engraftment, **41**:385, 388
 - NK cells, **41**:375
 - syngeneic stem cells, **41**:390
- IL-5 role, **57**:177
- lymphocyte dependence, **64**:192
- SCID and, **49**:397
- Grafts, *see also* specific type
- autoimmune thyroiditis and, **46**:286, 287
 - hybrid resistance and
 - antigen expression, **41**:397, 399, 401, 402, 404–407, 410
 - bone marrow cells, **41**:336–351
 - effector mechanisms antibodies, **41**:376, 377
 - macrophages, **41**:371
 - marrow, **41**:384–388, 390

- Grafts (*continued*)
 - NK cells, 41:373–375
 - in vitro* assays, 41:393, 395
 - leukemia/lymphoma cells, 41:362, 364–367
 - lymphoid cells, 41:352–355
 - syngeneic stem cell functions, 41:388, 390
 - liver cell, hybrid resistance and, 41:333
 - in MHC class I-deficient mouse
 - hematopoietic cell, 55:388
 - liver cell, 55:391
 - pancreatic islet, 55:390
 - skin, 55:388–390
- Graft-versus-host disease, 59:420–426
 - acute
 - clinical manifestations, 40:397
 - pathologic manifestations and pathogenesis, 40:397–399
 - prevention, 40:403–406
 - prognostic factors, 40:399–402
 - treatment, 40:402–403
 - autoreactive T cells and, 45:432
 - cachectin and, 42:220
 - CD5 B cell and, 47:129, 130
 - CD40-L expression, 61:47
 - hematopoietic engraftment and
 - acute, 40:397–406
 - chronic, 40:406–410
 - pathogenesis, 40:395–396
 - hybrid resistance and
 - antigen expression, 41:403
 - bone marrow cells, 41:335, 336, 340, 350
 - leukemia/lymphoma cells, 41:360, 367
 - lymphoid cells, 41:351–356
 - marrow engraftment, 41:385–387
 - T cells, 41:380
 - in vitro* assays, 41:394
 - IFN- γ , 62:101–103, 105
 - IL-1ra effects, 54:215
 - NK cells and, 47:231, 280–282
 - rat, in bone marrow transplantation, 39:280–281
 - idiotypic network and, 39:281
 - SCID and, 49:388, 391–395, 397–399
 - spontaneous autoimmune thyroiditis and, 47:455, 474, 482, 483
 - T cell subsets and
 - cell surface molecules, 41:43
- H-2 alloantigen recognition, 41:83, 84, 86, 88, 92, 95
- H-2-restricted antigen recognition, 41:71, 72
- Graft-versus-leukemia, 40:419–422
- Gramicidin A, cytolysis and, 41:316
- Granule proteins
 - cell-mediated killing and
 - cell lines, 41:286, 287
 - cytoplasmic granules, 41:287–291
 - lymphotoxin, 41:298
 - membrane attack complex of complement, 41:310
 - pore-forming protein, 41:291–297
 - proteoglycans, 41:298, 299
 - serine esterases, 41:297, 298
 - TNF-related polypeptides, 41:298
 - eosinophil, 60:191–194, 205–208, 221
 - role in DNA fragmentation, 60:300–301
- Granules
 - eosinophil, 60:155–162, 191–194
 - exocytosis
 - cytotoxicity and, 41:273, 281, 282
 - model, 60:290–291
 - Granule-stored enzymes, 60:201
 - Granulocyte colony-stimulating factor, 48:72; 66:149
 - activation of Jak, 60:7
 - gene regulation, 65:28–29
 - and GM-CSF, in immune response, 58:10–11
 - sepsis and, 66:104, 149–150
 - single receptor chain, 60:10
 - Granulocyte functional antigen, GFA-2 on eosinophils, parasite killing and, 39:184
 - Granulocyte-macrophage colony-stimulating factor, 42:221; 46:112, 115, 117, 122, 133; 48:71, 169; 54:174
 - and antitumor immunity, 58:429–432
 - cellular sources of, 39:2, 3, 10–13
 - and CR3 expression, 60:182–183
 - effect on genetically modified cells, 58:427
 - eosinophil-derived, 60:196–197
 - eosinophil proliferation, 48:75
 - expressing tumor cells, 58:438
 - factor-dependent mast cell production of, 53:8–9
 - functions, 39:2–3
 - and G-CSF, in immune response, 58:10–11

- GM-CSF/IL-3/IL-5 subfamily, **61**:151, 166–167
GMS/BCG vaccine, **62**:220
IL-5, **48**:75
production by T cells
 cyclosporin A-inhibited, **39**:13–14
 dexamethasone-inhibited, **39**:14
 mRNA accumulation and, **39**:14
purification and properties, murine, **39**:9
rheumatoid arthritis, **64**:319
role in cytokine signaling, **60**:5
sepsis and, **66**:150
single gene-coded, human, murine, **39**:9–10
transformed mast cell production of, **53**:4–6
- Granulocyte–macrophage colony-stimulating factor receptors, **52**:351; **63**:275, 276, 308
expression by eosinophils, **60**:183–185
- Granulocyte–macrophage response, sequential nature, **48**:70
- Granulocytes, **48**:69
 B cells and, **41**:238
 B cell precursors, **41**:191, 195
 bone marrow cultures, **41**:209, 210, 219
 genetically determined defects, **41**:229, 230
 cell-mediated killing and, **41**:312
 colony-stimulating factor, **48**:75
 helper T cell cytokines and, **46**:115, 117
 hybrid resistance and, **41**:336, 370, 380
 IL-4, **48**:75
 IL-6, **48**:75
 leukocyte integrins and, **46**:151, 161–163, 166
 leukocyte surface antigens and, **49**:88, 90, 93, 111–112, 116
 nonrecirculating, **58**:350–351
 opioid receptor, human, **48**:180
 pituitary hormones, **63**:411–412
 PNH, clonal origin, **60**:81
 with two PNH clones, **60**:89
- Granuloma
 in Crohn's disease, **42**:297
 oil, plasmacytogenesis and, **64**:224–225
 in pulmonary inflammation, **62**:280–283
- Granulopoiesis, hybrid resistance and, **41**:336
- Granzymes, **64**:1
 and apoptosis, **58**:226
- as effector molecules in CTL cytotoxicity, **60**:300–301
or fragmentins, cytotoxic T cells containing, **58**:240
- Graves' disease, **62**:19
 autoimmune thyroiditis and, **46**:263
- Graves' ophthalmopathy
 antibodies to self-antigens from human donors, **57**:245
 autoimmune thyroiditis and, **46**:268
- Grb2, **63**:143, 144, 146
 signal transduction, **59**:262
- GRB2, recruitment of ras exchange factor, **60**:13
- G reactive protein, NK cells and, **47**:246, 247
- Grooves, MHC molecules
 mutations, **58**:161
 peptide antigens bound to, **58**:330
- Growth, B cells, and differentiation, *in vivo* aspects, **52**:221–230
- Growth arrest, and apoptosis induced by anti-IgM, **54**:410–411
- Growth factor receptor, **48**:253
 Ig gene superfamily and, **44**:30, 39
 IL-1 and, **44**:180
- Growth hormone, **48**:172; **63**:379–380
 expression, **63**:386–388
 gene regulation, **63**:400
 gene regulation by, **65**:20
 immune system, **63**:377–378
 leukemia, **63**:428
 lymphohemopoietic system, **63**:386–388, 431–432
 macrophage, **48**:178
 regulation, **63**:385–388
 structure and physiology, **63**:384–385
- Growth hormone deficiency
 in humans, **63**:408–409, 432–433
 X-linked agammaglobulinemia, **59**:149, 194
- Growth hormone receptor, **63**:396–401
- Growth hormone-releasing hormone, **63**:385
- GRs, *see* Glucocorticoid receptors
- GTP
 binding of ras, **60**:209
 neutrophil activation and, **39**:99, 100, 116, 120, 121, 132
- GTPase
 G proteins and, **39**:112
 in neutrophil activation, **39**:113–115

GTP-binding protein, **43:290**
 association with CD4-p56^{lk} and CD8^{lt},
53:64–65
 T cell signal transduction, **48:265–269**

GTP γ S, **43:292**

Guanylyl cyclase, **62:74**

Gut, lymphokine production by isolated T cells, **53:167**

Gut-associated lymphoid tissue
 immunosenescence and, **46:252, 253**
 rabbit
 antibody repertoire generation role, **56:180, 209**
 as bursal equivalent, **56:180, 204**
 follicular structure, **56:203**
 germfree rabbits, **56:204–206**
 model, **56:206–209**

Gut-associated tissues, homing to, **58:391–392**

GVHD, *see* Graft-versus-host disease

H

2H1, **48:249–250**

H-2 molecules
 alloantigen recognition and, **41:78**
 alloreactivity, **41:78–83**
 antigen-presenting cells, **41:88–92**
 effector phase, **41:92–95**
 resting T cell subsets, **41:83–88**

antigen recognition and, **41:51, 52**
 effector phase, **41:75–77**

T accessory molecule function, **41:59–62**
 triggering of activated T cells and
 hybridomas, **41:52–59**

triggering of unprimed and resting T cells, **41:62–75**

B cell formation and, **41:213**

CTL and, **41:136, 170**

β_2 -microglobulin, **41:155, 156**

carbohydrate moieties, **41:152–154**

exon shuffling, **41:138–148**

HLA class I antigens, **41:149, 152**

monoclonal antibodies, **41:156, 157, 166, 167**
 mutant strains, **41:158–163, 165**

cytotoxicity and, **41:269, 279–281**

hybrid resistance and
 antibodies, **41:376–378**

antigen expression, **41:397–404**
 bone marrow cells, **41:336–348**
 leukemia/lymphoma cells, **41:358–369**
 lymphoid cells, **41:352–354**
 marrow engraftment, **41:386–388**
 NK cells, **41:375, 376**
 syngeneic stem cells, **41:390**
 T cells, **41:381–383**
in vitro assays, **41:395, 396**

T cell subsets and, **41:40, 43–48**

thymus, and, **41:95, 96**
 development, **41:96–99**
 restricted T cells, **41:99–107**
 tolerance induction, **41:107–110**

Haemophilus influenzae
 pulmonary infection, **59:396–397**
 type B, **38:289, 295, 296, 302**

Hageman factor, *see* Factor XII

Hairpin
 defined, **64:45**
 formation, **64:55, 56–59**
 opening end, **64:60**

Hairpin resolution model, for P nucleotide formation, **58:36–38**

Hairy cell leukemia
 gene rearrangements and, **40:291–292**
 leukocyte integrins and, **46:151**

Hairy leukemia, **54:50–51**

Ham1/2, **64:4**

HAM genes, **65:49**

Haplotype
 aPL antibodies and, **49:222**
 autoimmune demyelinating disease and,
49:365, 368
 autoimmune thyroiditis and, **46:275, 279, 284–286, 319**
 complement components and, **38:236**
C4, 38:220, 229–231
MHC, 38:208
 restriction fragment length
 polymorphism, **38:215, 216**

CTL and, **41:159**

hybrid resistance and
 antigen expression, **41:397–399, 409**
 bone marrow cells, **41:341, 343**
 leukemia/lymphoma cells, **41:364**
 lymphoid cells, **41:353**

Ig heavy-chain variable region genes and,
49:23

- Ir* genes and
 antigen binding, **38**:114
 antigen specificity, **38**:121, 122
 complementation, **38**:72, 73, 79
 gene function, **38**:91
 Ig molecules, **38**:70–72, 83
 I region mutations, **38**:84, 90
 polypeptides, response to, **38**:46, 49
 T cell, cytotoxic, **38**:161, 166, 185
 T cell interactions, **38**:153, 157–159
 T cell selection, **38**:131
 T cell suppression, **38**:135, 138–140,
 142–145, 149
 tolerance, **38**:126, 128
 maps of Ig-like loci and, **46**:32
 maternally transmitted antigen and, **38**:318
 class 1 antigens, **38**:333, 335, 336, 351
 strain combinations, **38**:349
 MHC, RA susceptibility associated with,
56:438–446
 murine lupus models and, **46**:99
 Ig germline, **46**:66, 67, 69, 71, 72
 T cell antigen receptor, **46**:84–86, 88, 90
 T cell subsets and, **41**:47, 48, 78, 79, 102
 Haplotype *b*, **48**:123
 Haplotype *d*, **48**:123
 Haplotype *k*, **48**:123
 Hapten
 augmentation, immunosenescence and,
46:247, 248, 252
 binding, **43**:17
 CD8 molecules and, **44**:280, 282
 derived T suppressor-inducer cell, **48**:212
 genetically engineered antibody molecules
 and
 antigen-combining sites, **44**:84, 85
 chimeric antibodies, **44**:79–81
 expression, **44**:79
 fusion proteins, **44**:85
 production, **44**:66
 helper T cell cytokines and, **46**:120, 121
 immunosenescence and, **46**:247
Ir genes and
 antigen binding, **38**:103
 B cell responses, **38**:51
 cytotoxic T cells, **38**:165
 “Schlepper” experiment, **38**:39, 40
 T cell activation, **38**:106, 107
 T cells, **38**:41
 pulmonary hypersensitivity, **59**:417–418
 specific B cell tolerance, **54**:395–402
 analysis with affinity-purified B cells,
54:398–401
 differential signaling via IgM and IgD,
54:401–402
 specific fragment cultures and limiting
 dilution, **54**:395–398
 Hapten-carrier effect, humoral immune
 response and, **45**:3, 4, 6, 7
 Haptotaxis, stem cell factor receptor-ligand
 interaction in, **55**:37–38
 Hashimoto’s thyroiditis, **46**:263, 266, 317
 antigens, **46**:268, 269, 271–275
 cellular immune responses, **46**:287–291,
 293
 genetic control, **46**:284
 humoral 298–301, **46**:303, 305, 306
 Hay fever, IgE biosynthesis and, **47**:1, 28
 H-CAM, multiple myeloma and, **64**:247,
 248
 Heart, myxoma, and IL-6, **54**:39–40
 Heart disease
 coronary, dietary fish oil and, **39**:145–146
 eosinophil degranulation and, **39**:226
 eosinophil localization and, **39**:226–227
 Heat-shock proteins, **62**:78, 244
 anticellular antibodies and, **44**:90, 96, 109,
 132
 hsp70, **62**:244
 MHC class I peptide loading, **64**:128
 mycobacterial, recognition, **58**:321–322
 surface antigens of human leukocytes and,
49:79, 82
 Heat-stable antigen, for B and T cells,
53:235–236
 Heavy chain
 allotypes, Crohn’s disease and, **42**:288
 Ars A response and, *see Arsonate idiotypic*
 system, A/J mouse, molecular genetics
 constant region gene order and class
 switching, **40**:257–258
 genes, **48**:41
 assembly of, **40**:252–253
 expression, sequences upstream of Ig
 promoter, **43**:253–254
 hypermutations, **60**:277–278
 locus, **62**:1–26
 mu heavy chain
 allelic exclusion, **63**:2, 19–21

- Heavy chain (*continued*)
 B cell development, **63:1–4**, 11–15,
 16–25, 31
 ligand for PreB cell receptor, **63:25–26**
 signal transduction through PreB cell
 receptors, **63:26–29**
 surrogate, **63:8**, 25
- Heavy chain diseases, **64:219**, 220
 Ig gene defects in, **40:305–306**
- Heavy chain variable region, *see also* Variable
 region, gene expression
 Ars A response and, **42:99–105**, *see also*
 Arsonate idiotypic system, A/J mouse
 cross-reactive idioype-positive
 molecules, **42:109–110**
 kinetics, **42:108–109**
 repetitive substitutions, **42:110–111**
 serine, **42:111–112**
- Heavy chain variable region, genes,
43:237–238; **49:1–3**, 61, 63
 and antibody repertoire generation in
 rabbit
 allotype, **56:195**
 organization, **56:188–191**
 preferential rearrangement, **56:197–198**
 preferential selection, **56:197–198**
 preferential use, **56:195–196**
- D-proximal genes, **49:20–21**
- D segments, **49:28–33**
 expression, *see* Variable region, gene
 expression
- J_H segments, **49:33–35**
 ontogeny, **49:2**, 33, 36–39, 41, 57
 organization, **49:3–6**
 polymorphism of V_H gene segments,
49:21–28
- promoter, **43:238**
 activation, **43:243**
 conserved octanucleotide and activity,
43:252–253
 enhancer independence and
 dependence, **43:265–266**
- regulation, **49:52–54**
 enhancer elements, **49:56–62**
 promoter elements, **49:54–56**
- V_H families, **49:6–8**
 V_H gene expression, **49:35–36**, 51–52
 autoantibodies, **49:41–51**
 B cell malignancies, **49:39–40**
 ontogeny, **49:36–39**
- V_HI genes, **49:5–9**, 11, 14–15, 61
 expression, **49:37**, 40, 42–43, 47, 49
 polymorphism, **49:22–23**, 27
- V_HII genes, **49:6–9**, 11, 61
 expression, **49:37**, 40, 42, 50
 polymorphism, **49:22–23**, 27
- V_HIII genes, **49:5**, 7–9, 11–15, 61
 expression, **49:37–38**, 40, 42–43, 48
 myasthenia gravis, **49:50–51**
 polymorphism, **49:22–27**
 regulation, **49:58–61**
- V_HIV genes, **49:5**, 11, 14–17, 61, 63
 expression, **49:38**, 40, 42–43, 47–51
 polymorphism, **49:26–27**
- V_HV genes, **49:4**, 17–19, 61
 expression, **49:40**, 50
 polymorphism, **49:26–27**
- V_HVI genes, **49:4–5**, 19–20, 61
 expression, **49:37–38**, 40, 42
 polymorphism, **49:26**, 28
 YH families, **49:6–11**, 14–20
- Heligmosomoides polygyrus*, **54:94**
- Helix, movement, **43:104**
- Helix-loop-helix oncogenes, in chromosome
 translocations, **50:129–130**
- Helminth infections
 eosinophil role, **60:210–216**, 227–228
 IL-5 role, **57:176–177**
 multiple antigen peptides as immunogens,
60:124–128
- Helper T cells, *see* T helper cells
- Hemagglutinin
 lymphocyte homing and, **44:354**, 355, 363
 spontaneous autoimmune thyroiditis and,
47:435, 445, 488, 489
- sporozoite malaria vaccine and, **45:307**
 synthetic T and B cell sites and
 peptides, **45:241**, 242
 vaccine, **45:253–255**
 viral antigens, **45:216**, 219
- T cell receptor and, **45:122**, 125, 133, 135
 T cell subsets and, **41:57**
 virus-induced immunosuppression and,
45:346
- Hematopoiesis, **48:83–84**
- B cell formation and, **41:181**, 182, 235–238
 B cell precursors, **41:188**, 191, 196, 197,
 200, 201
 bone marrow cultures, **41:208–210**, 212,
 214, 215, 217, 219, 220

- genetically determined defects, **41**:224, 225, 228, 229, 231, 232
 inducible cell line, **41**:220, 221, 223
 lymphohemopoietic tissues, **41**:185–188
 soluble mediators, **41**:233
 B cell genesis, **63**:203–211
 cachectin and, **42**:224
 CD44 activation states, **54**:306
 by CD44-specific antibodies *in vitro*, inhibition, **54**:299–300
 IFN- γ and, **62**:92–93
 IL-6 effects, **54**:22–25
 LIF role, **53**:35–36
 NK cells and, **42**:182, 183
 physiological experiments implying role for CD44, **54**:296–302
 pituitary hormones, **63**:404–405
 stem cell factor receptor effect on, **55**:43–48
 transcription factors, **63**:216–250
 vav, **63**:142
- Hematopoietic cells
 CD23 antigen and, **49**:150, 169
 CD44 expression, **54**:291–296
 during hematopoietic development in mouse and human, **54**:291–294
 on memory T cells and activated T cells, **54**:294–296
 cell lines, IL-2 α and β , **61**:150
 gene regulation by Stat proteins, **60**:23–25
 grafts, in MHC class I-deficient mouse, **55**:388
 helper T cell cytokines and, **46**:115–117
 HIV infection and, **47**:395
 hybrid resistance and
 antigen expression, **41**:400, 403, 405–407, 410
 bone marrow cells, **41**:335–351
 effector mechanisms, **41**:369, 370
 macrophages, **41**:371, 372
 marrow engraftment, **41**:385–388
 marrow microenvironment, **41**:391, 392
 NK cells, **41**:373, 374, 376
 syngeneic stem cell functions, **41**:388–390
 T cells, **41**:380
 in vitro assays, **41**:396
 leukemia/lymphoma cells, **41**:361, 363
 Ig gene superfamily and, **44**:30
- induction of positive selection, **58**:164–165
 leukocyte integrins and, **46**:160, 166
 NK cells and, **47**:187, 198, 303
 differentiation, **47**:233
 effector mechanisms, **47**:235
 malignant expansion, **47**:228
 surface phenotype, **47**:201, 212
 normal, **54**:304–305
 phosphatase, **63**:165
 role in cytokine signaling, **60**:25–26
 T cell development and, **44**:215, 226, 250, 254
 TGF- β effect on, **55**:186–187
 Hematopoietic engraftment, **40**:382–383
 engraftment of HLA-incompatible marrow, **40**:386–387
 experimental studies of transfusion effects, **40**:386
 facilitation of engraftment by T cells, **40**:391–392
 graft rejection in patients receiving T cell-depleted marrow, **40**:389–391
 graft rejection in patients transplanted for aplastic anemia, **40**:383–385
 marrow graft resistance, **40**:387–389
 reconsideration of marrow graft failure in humans, **40**:392–395
- Hematopoietic growth factor receptor superfamily, **52**:340–351
 granulocyte-macrophage colony-stimulating factor receptor, **52**:351
 IL-2 receptor, **52**:342–343
 IL-3 receptor, **52**:343–349
 IL-4 receptor, **52**:349–350
 IL-5 receptor, **52**:350–351
- Hematopoietic progenitors
 CD40 expression, **61**:12, 39
 IL-6 effects, **54**:22–23
- Hematopoietic stem cells
 immunosenescence and, **46**:240–247
 murine lupus models and, **46**:64, 65
 mutants, **60**:90–91
 SCID and, **49**:382, 385
 engraftment, **49**:388–391
 lack of engraftment, **49**:396–397
 transplantation with T cell-depleted bone marrow, **49**:394
- Hematopoietic system, **54**:83–84
 receptor superfamily, **54**:82–83

- Hematopoietic tumors, stem cell factor role
in etiology of, **55:72–74**
- Hemidesmosomes
pemphigoid antigen localization,
53:297–300
ultrastructure diagram, **53:297**
- Hemodialysis, therapeutic regulation of
complement system in, **56:278–279**
- Hemolysis, cell-mediated
cytolytic proteins and, **41:317, 318**
cytoplasmic granules, **41:288–290**
membrane attack complex of complement
and, **41:299, 305, 306, 309**
membrane damage and, **41:279**
pore-forming proteins and, **41:291, 293,**
295, 296
proteoglycans and, **41:299**
- Hemolytic anemia, **59:146–147, 343**
in PNH, **60:74–75**
- Hemolytic overlay, **38:208**
- Hemophiliacs, HIV-infected, **65:291–292**
- Hemopoiesis, *see* Hematopoiesis
- Hemopoietic growth factor receptor
granulocyte-macrophage colony-stimulating
factor receptor, **52:351**
superfamily, HRS, **52:340–351**
- Hemopoietin I, B cell formation and, **41:234**
- Hen egg lysozyme, synthetic T and B cell
sites and
globular protein antigens, **45:204, 211–214**
peptides, **45:240**
- Hen egg lysozyme model, clonal anergy,
vindicated, **52:305–311**
- Heparin
aPL antibodies and, **49:223, 240, 256**
cell-mediated killing and, **41:295, 299**
hybrid resistance and, **41:371**
leukocyte integrins and, **46:159**
SCID and, **49:395**
surface antigens of human leukocytes and,
49:88
in therapeutic complement inhibition,
56:287–288
- Hepatic growth factor, **64:153**
- Hepatitis, **59:68, 150**
chronic infection, effector cell role,
60:345–346
- Hepatitis B virus
antibodies from combinatorial libraries,
57:220
- C/EBP and, **65:10**
MAP immunogenicity, **60:128–130**
rabbit, anti-Id antibodies as vaccines
against, **39:283**
synthetic T and B cell sites and, **45:196,**
261, 263
bacterial antigens, **45:230**
globular protein antigens, **45:206, 212,**
214
immunological considerations, **45:201,**
202
peptides, **45:202, 239, 242, 243**
vaccine, **45:254–256**
viral antigens, **45:220, 221, 224**
- Hepatitis virus, helper T cell cytokines and,
46:129
- Hepatocyte growth factor, effects on
lymphocytes, **58:379–380**
- Hepatocytes
acute-phase protein synthesis, **54:25–27**
grafts in MHC class I-deficient mouse,
55:391
helper T cell cytokines and, **46:116**
IL-1 α production, **54:188**
iNOS production, **60:351–352**
interaction with IgA, **40:171, 177–184**
leukocyte integrins and, **46:162**
T cell activation and, **41:16**
- Hepatocyte-stimulating factor
IL-1 and, **44:165**
and IL-6, **54:4**
- Hepatocyte-stimulating factor III, *see*
Leukemia inhibitory factor
- Hepatoma, adoptive T cell therapy of tumors
and, **49:301, 305**
- HER-2/neu, **62:223–224, 237–238**
- Heredity angioedema, **61:206**
contact activation and, **66:239**
kinin formation in, **66:257–258, 259**
- Heredity primary immunodeficiency
disorders, gene defects in Ig and T cell
receptor, **40:306–309**
- HERMES-3, **65:360**
- Hermes antigen, mediating PLN adhesion,
54:296–298
- Herpes simplex virus, **41:153, 375, 376**
cytotoxic T cells and, **65:279**
ocular infection, **48:208–210**
regulators of complement activation and,
45:409, 410

- synthetic T and B cell sites and, **45**:225, 243, 256
- T cell receptor and, **45**:132
- thymidine kinase gene, **58**:434
- virus-induced immunosuppression and, **45**:367
- Herpes simplex virus type 1, **48**:210
antibodies from combinatorial libraries, **57**:222–227
intraocular infection, anterior chamber, **48**:210
multiple antigen peptides, **60**:130
- Herpes simplex virus type 2, antibodies from combinatorial libraries, **57**:222–227
- Herpesvirus, IL-10 gene acquisition, **56**:11–12
- Herpesvirus saimiri, and Epstein–Barr virus, **54**:115–116
- Herpesvirus type 6, human, and HIV-1, exacerbation of AIDS disease progression, **58**:16
- 5-HETE, *see* 5-Hydroxyeicosatetraenoic acid
- Heterogeneity
autoimmune thyroiditis and, **46**:274, 281, 300
B cell formation and, **41**:196, 216, 217
CTL and, **41**:138, 139, 152
cytotoxicity and
cell-mediated killing, **41**:312
mediation, **41**:271
membrane attack complex of complement, **41**:305, 306, 309, 310
helper T cell cytokines and, **46**:125, 126
hybrid resistance and, **41**:372
leukocyte integrins and, **46**:151, 154, 174, 175
T cell activation and, **41**:27
T cell subsets and, **41**:43, 79, 98
- Heterohybridoma antibodies, **38**:288–290, 296
- Heterohybridomas, human–mouse, **38**:300, 301
- Heterokaryons, hybridomas and, **38**:287, 291, 302
- Heterozygosity, Ig gene superfamily and, **44**:26, 28, 29
- High endothelial binding factors, lymphocyte homing and
carbohydrate, **44**:364, 370
molecules, **44**:344–346, 349, 351
- High endothelial cells, lymphocyte homing and, **44**:315, 316, 318, 320
- High endothelial venules, **59**:385, 386; **62**:257; **64**:142; **65**:379–380
binding assay, **64**:185–186
flattening, **65**:373
history, **65**:347–348
lymph node, **58**:353, 359, 390–391
lymphocyte binding to, role of CD44, **58**:376–377
- lymphocyte extravasation in, **58**:347
- lymphocyte homing and, **44**:314
cell adhesion, **44**:353–355
inhibition, **44**:355–357
interaction, **44**:318–322
ligands, **44**:363–371
molecule attachment, **44**:342–353
receptors, **44**:357–363, 369, 370
regional Specificity, **44**:322–330, 333–341
structure, **44**:315–318
- lymphocyte migration and, **65**:350–351
- lymphocytes and, **64**:142; **65**:352
addressins, **65**:361–363, 377–379
adhesion, **65**:365–372
CD34, **65**:362–377, 377
CD44, **65**:359–360
extravasation, **65**:360, 365–367
GlyCAM-1, **65**:361–362, 363, 364, 373, 377
homing receptor ligands, **65**:360–365
integrins, **65**:352, 358–359, 368–369, 370
MAdCAM-1, **65**:352, 359, 362–363, 367, 370, 377, 379
migration, **65**:350–352
mucins, **65**:360–361
selectins, **65**:352, 354–357, 363–365, 367
Sgp200, **65**:362, 377
sugars, **65**:360–361, 363–365
VAP-1, **65**:365
in vitro binding assay, **65**:351–352
ontogenesis, **64**:164–165
regulation, **65**:372–375
differentiation, **65**:377–379
immune response and, **65**:375–377
rolling, **65**:367, 368
structure of, **65**:348–350
tethering, **65**:367

- High-molecular-weight kininogen
 contact activation and, **66**:236, 237
 endothelial cell receptor, **66**:249–257
 inactivation, **66**:245
 properties, **66**:226, 256
 structure, **66**:230–233
- High-performance liquid chromatography,
 complement receptor 1, **46**:187
- Hinge region
 IgA molecules, **40**:158
 structure, **48**:11
 complement binding activity, **48**:13,
 15, 22
 Fab, **48**:21–23
 Fc, **48**:21–23
 segmental flexibility, **48**:13, 15, 22
- upper part
 complement activation, **48**:25
 Fab motion, **48**:25
 length, **48**:25
 segmental flexibility, **48**:25
- Histamine**
 genetically engineered antibody molecules
 and, **44**:80
 IL-1 and, **44**:164, 187
- Histamine receptors**, **52**:402
- Histamine release**
basophil
 major basic protein and, **39**:191
 somatostatin and, **39**:305–306
- basophil/mast cell**
 clinical considerations, **50**:250–254
 cytokines and histamine release,
50:247–248
 factors, **50**:237
 modulation of histamine release by
 cytokines, **50**:248–250
- mast cell**
 major basic protein and, **39**:191
 SP and, **39**:305–306
- Histamine releasing factor**, **43**:307
 cell sources, **50**:243–244
 mechanism of action, **50**:244–247
 purification, **50**:238–243
- Histidine**, IL-1 and, **44**:158, 181–182
- Histiocytes**, hybrid resistance and, **41**:335
- Histiocytosis**, malignant, complicating celiac
 disease, gene rearrangements and,
40:292–293
- Histocompatibility antigen modifier gene**,
65:49
- Histocompatibility antigens**, **38**:382
 H-2 alloantigen recognition and, **41**:82, 87
 H-2 molecules in thymus, and, **41**:99, 100
 H-2-restricted antigen recognition and,
41:51, 57, 71–75
 hybrid resistance and, **41**:333, 334, 350
- Histocompatible bone marrow**
 transplantation, **49**:392
- Histones**, antinuclear antibodies and
 autoantibodies, **44**:125, 127
 autoantigen, **44**:128, 130
 drug-induced autoimmunity, **44**:109, 110
 SLE, **44**:95, 96, 98–100
- Histoplasma capsulatum***, leukocyte integrins
 and, **46**:163
- Histotope**
Ir genes and, **38**:107
 T cell subsets and, **41**:53, 58
- HIV**, *see* Human immunodeficiency virus
- HIVEN86A**, **48**:298
- HLA**
 autoimmune demyelinating disease and,
49:367
 class II molecules, **49**:357, 362–363,
 368–371, 374
- CTL** and, **41**:168, 170
 β_2 -microglobulin, **41**:155, 156
 amino acid, **41**:163–165
 carbohydrate moieties, **41**:153
 exon shuffling, **41**:138, 139, 141, 142,
 144
 monoclonal antibodies, **41**:158, 165, 167
 transfected cells, **41**:149–152
- cytotoxicity and, **41**:270
 human leukocyte antigen, oligosaccharide
 mobility, **48**:31
- hybrid resistance and, **41**:350, 369,
 400–403
- idiotypic network in responses to,
39:281–283
- matching, corneal transplantation, **48**:193
- NK cells and, **47**:198, 223
- polymorphisms, RA susceptibility associated
 with
 remission during pregnancy, **56**:431
 testing, **56**:432–433
 toxicity, **56**:431–432
- population distribution, **48**:126

- rheumatoid arthritis, **48**:126
 rheumatoid arthritis susceptibility
 associated with
 risk ratios, **56**:439
 studies, **56**:436–438
 rotational correlation times, **48**:28–29
 SCID and, **49**:392–393, 398
 type, IBD and, **42**:288
 HLA A1-B8-DR3, HIV progression to AIDS
 and, **65**:285
 HLA-A2
 allele
 composition, **64**:127
 MHC class I molecule peptide binding, **64**:126–127
 structure, **43**:215
 synthetic T and B cell sites and, **45**:219,
 223, 234, 235
 T cell receptor and
 alloreactivity, **45**:150, 151, 153, 155, 156
 epitopes, **45**:142, 144
 homogeneity, **45**:129
 MHC molecules, **45**:111, 112, 115, 116
 peptides, **45**:137, 138
 polymorphic residues, **45**:158, 160, 161
 structure-function relationships,
 45:139–141
 HLA-A2.1, **62**:234
 HLA-A3, multiple sclerosis, **48**:148–150
 HLA-B7, multiple sclerosis, **48**:148–150
 HLA-B8, HIV and, **65**:285
 HLA-B27
 HIV and, **65**:285
 spondyloarthritic diseases, **65**:94
 HLA-B35, HIV progression to AIDS and,
 65:285
 HLA-B37, **65**:286
 HLA-B49, **65**:286
 HLA-B57, HIV and, **65**:285
 HLA class I
 crystal structure, **48**:111
 system
 cytotoxic T cells and, **65**:279
 HIV infection and, **65**:284–286
 HLA class II
 allelic variability amino acid stretches,
 48:111–112
 amino acid sequence, **48**:108
 antigen recognition site, **48**:111
 autoimmune disease associations, **66**:67, 76,
 84–85, 93
 immune intervention, **66**:91–93
 insulin-dependent diabetes mellitus,
 66:91
 methimazole-induced insulin
 autoimmune syndrome, **66**:87–88
 multiple sclerosis, **66**:88–91
 pemphigus vulgaris, **66**:87
 rheumatoid arthritis, **66**:85, 87
 biochemical typing, **48**:114–116
 interisotypic association, **48**:116
 intraisotypic hybrid molecule, **48**:116
 trans-association, **48**:116
 celiac disease, **48**:144, 147
 disease associations, **48**:107–153
 disease susceptibility, **48**:108
 DP, **48**:109, 110–111
 DQ, **48**:109, 110–111
 DQ gene, **48**:120–121
 DQ haplotype, **48**:120–121
 DQ molecule, **48**:120–121
 DR, **48**:109, 110–111
 epitope, **48**:112
 molecular localization, **48**:152–153
 gene location, **48**:109
 genes, **66**:67
 gene structure, **48**:109–112
 homologous, **48**:109
 hybrid molecule, **48**:119–126
 cell surface expression, **48**:119
 stable heterodimer, **48**:119
 ligands
 class II-binding groove, **66**:68–69
 class II-binding motifs, **66**:69–77
 computational identification, **66**:77–84
 HLA-DP and -DQ, **66**:76–77
 HLA-DR, **66**:70, 72–76
 pocket specificity profiles, **66**:74–76
 quantitative matrices, **66**:72–76
 molecules, **52**:164–166
 interisotypic hybrid, **48**:124–126
 transfected genes, **48**:124
 other antigens of B cells, **52**:173–174
 Bgp 95, **52**:173
 CK226 antigen, **52**:173
 IgM-binding protein (*Fcμ* receptor),
 52:173
 structure, **48**:109–112
 synovial cell, **48**:132

- HLA class II (continued)**
- one-dimensional isoelectrofocusing, **48:114–116**
 - peptide binding motifs, **66:69–77**
 - structure, **66:68–69**
 - polymorphism, **48:109, 112–119**
 - serology, **48:112–114**
 - two-dimensional polyacrylamide gel electrophoresis, **48:114–116**
 - silver staining, **48:116**
 - theoretical limitations, **48:116**
 - typing, **48:112–119**
 - Western blotting, **48:116**
- HLA-D**
- deletion mutants, **37:15–18**
 - function of, **37:63–65**
 - genes, **37:53–60**
 - invariant (γ) chain, **37:50–53**
 - restriction fragment length polymorphism, **37:60–63**
 - supertypic specificity localization, **37:47–50**
- HLA-DP**
- biochemistry, **37:43–47**
 - serology/HTC, **37:14–15**
- HLA-DQ**
- B8, **48:134**
 - B15, **48:134**
 - biochemistry, **37:31–43**
 - DR4, **48:134**
 - homologous hybrid, **48:137**
 - serology, **37:9–14**
 - HLA-DQ hybrid molecule, **48:119–124**
 - celiac disease, **48:147**
 - cell surface expression, **48:122**
 - disease susceptibility, **48:122**
 - trans*-complementation, **48:119–120**
 - transectants, **48:122**
 - HLA-DR
- aPL antibodies and, **49:222**
 - biochemistry, **37:23–31**
 - cachectin and, **42:224**
 - CD23 antigen and, **49:156, 170**
 - IBD and, **42:292–293**
 - serology/HTC, **37:5–9**
 - HLADR3-DQ2, **65:286**
 - HLA-DR5, **65:286**
 - HLA-DR13, **65:286**
 - HLA-D region, **48:108**
 - HLA-D region products, monoclonal antibodies to, **37:18–21**
 - HLA-D region specificity, **48:113–114**
- HLA DR specificity, HLA D typing,** **48:113–114**
- HLA-Dw14, T cell clone reactivity,** **48:129**
- HLA factor, IDDM,** **48:133–144**
- HLA markers, MG and,** **42:251**
- hlx,** **63:240–241**
- hlx, B cell genesis,** **63:239–241**
- Hmt, maternally transmitted antigen and,** **38:315, 316**
- antigenic variation, **38:342, 344**
 - class I antigens, **38:333, 336**
 - mitochondrial transmission, **38:347, 348**
 - phylogenetic considerations, **38:338**
- Hmt,101–104**
- nature of, **52:102**
 - nature of antigen presented by, **52:102–104**
- HNK-1, NK cells and,** **47:291, 292**
- CNS, **47:266**
 - differentiation, **47:234**
 - effector mechanisms, **47:257**
 - genetic control, **47:222**
 - hematopoiesis, **47:275**
 - malignant expansion, **47:226**
 - morphology, **47:218**
 - surface phenotype, **47:206, 207**
 - tissue distribution, **47:221**
- Hodgkin's disease,** **38:299–301; 61:8, 12, 39**
- gene rearrangements and, **40:294**
 - IL-5 mRNA expression in patients, **57:158**
- Holoricin-AChR conjugates, MG and,** **42:265**
- Homeobox domain, Ig heavy-chain variable region genes and,** **49:55, 57**
- Homeodomain proteins**
- hlx, **63:239–241**
 - hox11, **63:244–245**
 - oct-2, **63:243–244**
- Homeostasis, B cell formation and,** **41:185, 208, 236**
- Homing**
- to gut-associated tissues, **58:391–392**
 - leukocytes
 - integrins in, **58:362**
 - molecular basis, **58:389–390**
 - therapeutic drug development and, **58:395–396**
 - naive B cells, **58:350**
 - PLN, **58:390–391**
 - preferences, avian, **50:96–97**
 - skin, **58:392**

- tissue-specific, lymphocytes, endothelium and, **50**:284–285
- Homing receptors
surface antigens of human leukocytes and, **49**:87–88
- tissue-specific, **58**:316–318
- Homodimerization, β cytoplasmic domains, **60**:10
- Homogeneity
B cell formation and, **41**:221, 228
cytotoxicity and, **41**:270, 284, 287, 300
hybrid resistance and, **41**:338, 339
T cell activation and, **41**:13
T cell receptor and, **45**:128–131
- Homologous recombination, **54**:242
gene inactivation, **59**:108
switch recombination, **61**:110–111
- Homologous restriction factor, **61**:254–255, 257
- NK cells and, **47**:252, 254
- Homology
antinuclear antibodies and, **44**:129
autoimmunedemyelinating disease and, **49**:367
CD4 molecules and, **44**:285, 286, 288, 295, 302
CD8 molecules and, **44**:272, 276, 278, 302
CD23 antigen and, **49**:162–164, 166
CTL and, **41**:136
 β_2 -microglobulin, **41**:154, 155
exon shuffling, **41**:140–142, 144–146
monoclonal antibodies, **41**:156, 158
cytotoxicity and, **41**:285, 307, 309, 314, 316
genetically engineered antibody molecules and, **44**:81
human T cell activation and, **41**:3
accessory molecules, **41**:14
gene regulation, **41**:29
synergy, **41**:17
T cell antigen receptor, **41**:4
Thy-1, **41**:10
hybrid resistance and, **41**:335
Ig heavy-chain variable region genes and D segments, **49**:29, 32–33
polymorphism of V_H gene segments, **49**:24, 26
 V_H families, **49**:8, 15, 17
 V_H gene expression, **49**:35, 39, 44–45, 48–51
IL-1 and, **44**:155–158, 167
- lymphocyte homing and, **44**:346
sequence and structural
conservation during evolution, **52**:383
conversation during evolution, **52**:383
Fc ϵ RI α chain related molecules, **52**:383
Fc ϵ RI β chain related molecules, **52**:383–384
Fc ϵ RI γ chain related molecules, **52**:384
surface antigens of human leukocytes and, **49**:76, 88, 92–93, 99, 114
T cell subsets and, **41**:41, 46, 49, 50
- Homology blocks
in Jaks, **60**:4
in Stats, **60**:18–19
- Homology unit, Ig gene superfamily and, **44**:1, 3–9
evolution, **44**:43–46, 48
MHC complex, **44**:24, 25, 27–29
nonimmune receptor members, **44**:31–35, 37–42
receptors, **44**:9–11
- Hormone radioreceptor assay, IL-2, **42**:166–167
- Hormones
autoimmune thyroiditis and
antigens, **46**:267, 272
cellular immune responses, **46**:298
experimental models, **46**:274, 278, 280
genetic control, **46**:283
B cell formation and, **41**:196, 229
comparison with cytokines, **63**:271
hybrid resistance and, **41**:402
IL-1 and, **44**:164, 171
immunosenescence and, **46**:244, 246, 247
murine lupus models and, **46**:63
myasthenia gravis and, **42**:263
NK cells and, **47**:299
reproduction, **47**:269, 270
pore formation and, **41**:314
spontaneous autoimmune thyroiditis and, **47**:491, 493
altered thyroid function, **47**:456, 461
cellular immune reactions, **47**:450
clinical symptoms, **47**:439
disturbed immunoregulation, **47**:476, 480, 481
potential effector mechanisms, **47**:465
sporozoite malaria vaccine and, **45**:305, 312
surface antigens of human leukocytes and, **49**:94, 101

- Hormones (*continued*)
 synthetic T and B cell sites and, **45**:232,
 246, 247, 259
 T cell development and, **44**:236
- Horror autotoxicus*, **54**:394
- Host red blood cells, T cell subsets and,
41:68
- Horse serum, B cell formation and, **41**:209,
 219
- Host defense mechanisms, **66**:101
- Host response, sequential nature, **48**:70
- Household contacts, IBD and, **42**:291
- Housekeeping, immunosenescence and,
46:234
- Hox11, **63**:244–245
hox11, **63**:244
- HPA axis, **54**:30–31; **61**:367–368, 375
- 5-HPETE, *see* 5-Hydroperoxy-5,8,11,13-eicosatetraenoic acid
- HS1, B cell development, **63**:29
- 8HS-20 gene, **63**:29
- HSP, *see* Heat shock proteins
hst-1, **65**:155
- HSV, *see* Herpes simplex virus
- HT-2 cells, IL-2 activation, **61**:176–179
- HTLV-1, *see* Human T lymphotropic virus type I
- Human
 antibodies from combinatorial libraries, *see* Antibodies, human, from combinatorial libraries
 antibody effector function, *see* Antibody effector function, human
 common mucosal systems in, **40**:199–204
 κ light chain genes of, **40**:249–251
 lymphoid neoplasms, Ig and T cell receptor gene rearrangements and, **40**:264–270
 and mouse, heavy chain locus organization, **54**:231–233
 mutations affecting stem cell factor or SCF receptor, **55**:23–24
- Human chorionic gonadotropin, synthetic T and B cell sites and, **45**:259
- Human cytomegalovirus, antibodies from combinatorial libraries, **57**:220–221
- Human herpesvirus type 6, and HIV-1, exacerbation of AIDS disease progression, **58**:16
- Human immunodeficiency syndromes, and DNA repair syndromes, **58**:62–69
- Human immunodeficiency virus, **47**:377, 378, 411–414; **52**:192–193, *see also* Human immunodeficiency virus type 1
 anti-TNF-α therapy, **64**:311
 aPL antibodies and
 clinical aspects, **49**:202, 205, 210–211
 isotype, **49**:224, 226–227
 CD8⁺ antiviral factor, **65**:282–284
 chemokine receptor family, **65**:283
 cytotoxic T cells, **65**:286–287
 acute infection, **65**:293–296
 adoptive immunotherapy, **65**:317–320
 adverse effects of CTL activity, **65**:308–311
 apoptosis, **65**:306
 asymptomatic period of infection, **65**:296–297
 CTL decline in late disease, **65**:300
 CTL exhaustion, **65**:304–305
 disease progression, **65**:297–298
 epitopes, **65**:291–293, 301, 312, 313
 escape mutation, **65**:311–317, 322–323
 infection, **65**:290–307
 long-term nonprogressors, **65**:298–300
 lysis of infected cells, **65**:280–281
 measurement of HIV-specific CTLs, **65**:287–290
 replication suppression, **65**:281–284
 in seronegatives, **65**:287, 307–309
 Th1 to Th2 switch, **65**:302–304
 therapeutic implications, **65**:317–322
 vaccines, **65**:320–322, 323
 enhanced apoptotic decay, **58**:233–237
 etiological agent, **47**:378
 genome, **47**:380–384
 HIV-2, **47**:384, 385
 life cycle, **47**:379, 380
 FDCs as targets of, **51**:275–278
 HLA class I system, **65**:284–286
 humoral immune response and, **45**:78
 IgE biosynthesis and, **47**:22
 Ig heavy-chain variable region genes and, **49**:57
 immune response, **47**:405
 cellular response, **47**:409–411
 humoral response, **47**:405–409
 immunopathogenic mechanism, **47**:397
 activation, **47**:398–403
 CD4 cell depletion, **47**:386–389
 CD4-HIV interaction, **47**:385, 386

- macrophages, **47**:392–395
 monocytes, **47**:392–395
 precursor cells, **47**:395–397
 T4 cells, **47**:389–392
 induced immunosuppression, **45**:335, 336, 355–363
 history of infection, **45**:352–355
 immune response, **45**:363–368
 virology, **45**:347–352
 infected CD4⁺ eosinophil precursors, **60**:187
 infection, **47**:377
 CTL-mediated lysis, **65**:280–281
 etiological agent, **47**:378, 380, 383, 384
 hu-PBL-SCID mouse, **50**:314–317
 other tumor models in SCID mice, **50**:313–314
 and IL-6, **54**:37–38
 immune response, **47**:406, 407, 409
 immunopathogenic mechanism, **47**:391
 in seronegatives, **65**:287, 307–309
 inhibition of lymphocyte death, **58**:279
 integrases of, **64**:57
 lymphocytic tendency to undergo PCD, **58**:214
 lysis of infected cells, **65**:280–281
 and mitosis, **58**:424
 neuropsychiatric manifestations
 brain, **47**:403, 404
 mechanisms, **47**:404, 405
 NK cells and, **47**:302, 303
 progression to AIDS, **65**:285, 299, 302, 304
 pseudotypes, in HIV-infected hu-PBL-SCID mice, biosafety issues and, **50**:317–319
 replication suppression, **65**:281–284
 sporozoite malaria vaccine and, **45**:308–310
 synthetic T and B cell sites and, **45**:196, 263
 globular protein antigens, **45**:207
 peptides, **45**:243
 viral antigens, **45**:219, 220
 variants, **65**:306–307
 Human immunodeficiency virus type 1, **63**:83–87
 animal infection, **52**:454–466
 chimpanzee, infection of, **52**:455–461
 model for testing HIV-1 vaccines, **52**:457–459
 pathogenesis of HIV-1 infection in, **52**:455–457
 use to test therapies based on CD4, **52**:459–461
 mouse, SCID-hu, **52**:463–466
 rabbit, infection with HIV-1, **52**:461–463
 antibodies
 from combinatorial libraries, **57**:211–217
 to self-antigens from human donors, **57**:246–250
 CD4 molecules and, **44**:290, 295–297
 CD8⁺ cells and, **66**:285–291, 297–298
 antiviral activity, **66**:274–275
 antiviral therapy, **66**:295–299
 lytic activity, HIV-1-specific, **66**:275–277
 dissemination, **63**:88
 genomic organization, **63**:84–85
 infection in rabbits, **52**:461–463
 inhibition, **66**:280–291
 CD8⁺ cell-derived antiviral factor, **66**:282–283
 chemokines and, **66**:281–282
 IL-16, **66**:283
 LTR and, **65**:10
 MAP immunogens designed for, **60**:130–134
 models using viruses distantly related to, **52**:430–432
 murine leukemia virus, **52**:432–434
 and NF-κB, **58**:14–17
 pathogenesis, **63**:90–95, 100
 progression, **66**:283
 persistence, **66**:291–295
 replication, **63**:85, 89–90
 targets for therapy, **66**:295–297
 CTLs, **66**:283, 286, 292–293
 cytolysis, **66**:284–286
 epitopes, **66**:276–280
 transmission, **63**:88
 ungulate lentiviruses, **52**:430–432
 V3 loop, **60**:137–138
 Human leukocyte antigen, *see* HLA
 Human papillomavirus, **62**:229, 239–240, *see also* Papillomavirus
 C/EBP and, **65**:10
 Human T lymphotropic virus
 HIV infection and
 etiological agent, **47**:378
 immunopathogenic mechanism, **47**:388, 391, 393, 400, 402

- Human T lymphotropic virus (*continued*)**
- IgE biosynthesis and, **47:21**
 - NK cells and, **47:229**
 - Human T lymphotropic virus type I, 50:180,**
 - 182; **52:429; 63:285, 310**
 - tax protein, **58:17–18**
 - Human umbilical vein endothelial cells**
 - CD40 antigen, **61:14**
 - cytokines, regulatory effects, **64:165–166**
 - as endothelial cell model, **64:186**
 - eosinophil adhesion to, **60:167–170**
 - lymphocytes and, **64:142**
 - Humoral immune response, 45:1; 64:2, *see also* Antibody response**
 - antigen-bridging model, **45:7, 8**
 - antigen presentation, **45:9–12, 22**
 - B cells, **45:15–17**
 - class II molecules, **45:18–21**
 - in vitro*, **45:21**
 - in vivo*, **45:21, 23, 24**
 - antigen processing, **45:12, 22**
 - B cells, **45:17, 18**
 - intracellular events, **45:12–14**
 - macrophages, **45:14, 15**
 - requirements, **45:8, 9**
 - antigen-specific receptors, **45:6, 7**
 - cellular interactions *in vivo*, **45:78, 79, 85**
 - bacterial antigens, **45:82, 83**
 - lymphoid organ, **45:79–82**
 - parasites, **45:84, 85**
 - viral antigens, **45:83, 84**
 - hapten-carrier effect, **45:3, 4**
 - helper T cells, **45:24, 25**
 - activation, **45:29–31**
 - clones, **45:29**
 - differences, **45:27, 28**
 - heterogeneity, **45:25–27**
 - isotype switching in B cells, **45:32–35**
 - subtype surface markers, **45:28, 29**
 - in vivo*, **45:31, 32**
 - HIV infection and, **47:405–410**
 - immune reconstitution and, **40:414–416**
 - to infection, IBD and, **42:299–300**
 - interleukins, **45:54–62, 75–77**
 - IL-1, **45:72, 73**
 - IL-2, **45:67–70**
 - IL-3, **45:73, 74**
 - IL-4, **45:62–65**
 - IL-5, **45:65–67**
 - IL-6, **45:74, 75**
 - interferon- γ , **45:70, 71**
 - mechanisms, spontaneous autoimmune thyroiditis and, **47:465–470**
 - against MMTV, **65:188–192**
 - models, **45:1–3**
 - physical interaction, **45:35, 36**
 - biochemical events, **45:38, 39**
 - cell surface Ig, **45:40, 41**
 - cell surface molecules, **45:41–46**
 - functional outcome, **45:38**
 - intracellular events, **45:39**
 - lymphokines, **45:46–54**
 - model systems, **45:36–38**
 - monogamous nature, **45:39, 40**
 - pituitary hormones, **63:407**
 - spontaneous autoimmune thyroiditis and, **47:444–449, 491**
 - suppression, **45:5**
 - Humoral memory, germinal center B cells, 60:283**
 - Hu-PBL-SCID mouse**
 - EBV-associated lymphoproliferative disease in, **50:310–313**
 - HIV infection of, **50:314–317**
 - Hyaluronan**
 - binding regulation, evidence, **54:302–308**
 - CD44 as receptor for, **54:284–287**
 - evidence, **54:284–287**
 - features, **54:287–290**
 - sequence of CD44 in hyaluronan recognition, **54:284**
 - CD44-specific antibody binding, inhibition, **54:284–286**
 - Hyaluronate, IGF-I, 63:390**
 - Hyaluronic acid**
 - CD44 binding to, **58:376–377**
 - Ig gene superfamily and, **44:40, 41, 44**
 - H-Y antigen, cytotoxic T cells and, 38:175–180**
 - cross-reactive lysis, **38:186**
 - immunodominance, **38:181**
 - Hybrid histocompatibility antigens**
 - antigen expression, **41:397–399, 401, 409, 410, 405–407**
 - trans gene model, **41:401–403**
 - bone marrow cells, **41:340–343, 345, 347**
 - genetics of expression, **41:397**
 - leukemia/lymphoma cells, **41:360–367**
 - lymphoid cells, **41:352, 353**
 - marrow engraftment, **41:385, 386**
 - in vitro* assays, **41:394–396**

- Hybrid hyperreactivity, **41**:360, 379
Hybridization, *see also* Somatic cell hybridization
 antigen-presenting cells and, **47**:65–67, 70, 71, 87
 autoreactive T cells and, **45**:420, 424
 CD5 B cell and, **47**:134
 CD23 antigen and, **49**:152, 162, 170
 complement components and
 C4 genes, **38**:228, 231
 cloning, **38**:209, 212
 restriction fragment length polymorphism, **38**:215
 complement receptor 1 and, **46**:188–190
 cytotoxicity and, **41**:276, 288
 helper T cell cytokines and, **46**:112
 human B cell neoplasia and, **38**:257
 Burkitt lymphoma cells, **38**:260
 chromosome translocation, **38**:266, 268
 IgE biosynthesis and, **47**:17
 Ig heavy-chain variable region genes and
 polymorphism of V_H gene segments, **49**:23, 27
 V_H families, **49**:8–9, 15, 17
 V_H gene expression, **49**:37–38
 leukocyte integrins and, **46**:159, 160, 173
 maps of Ig-like loci and, **46**:2–4
 MHC, **46**:30, 33
 structure, **46**:10, 12, 13, 15, 17, 18, 21, 25, 27
 maternal transmitted antigen and, **38**:324, 326
 murine lupus models and, **46**:74, 83, 85, 87
 NK cells and
 genetic control, **47**:223
 hematopoiesis, **47**:273, 274, 280
 surface phenotype, **47**:204, 209
 regulators of complement activation and, **45**:405
 SCID and, **49**:397
 spontaneous autoimmune thyroiditis and
 altered thyroid function, **47**:462
 genetics, **47**:483, 484, 486–491
 subtractive, **43**:135
 T cell receptor and, **45**:135, 139, 140
 T cell receptors and, **38**:15
 T cell subsets and, **41**:41, 100, 101
 virus-induced immunosuppression and, **45**:354
Hybrid joints, V(D)J recombination, **64**:41
Hybridoma/plasmacytoma growth factor, **54**:3
Hybridoma reagents, **38**:276
Hybridomas, **42**:1, 2
 activation-induced death, **50**:67
 adoptive T cell therapy of tumors and, **49**:301
 A/J mice, **42**:131–132, 135
 antiarsonate, **42**:97–98, 99–101, 128–129
 antibodies from, **57**:229–230
 antigen-presenting cells and, **47**:65, 68–70, 77
 antigen processing, **47**:87
 T cells, **47**:86, 93
 antinuclear antibodies and, **44**:98
 aPL antibodies and, **49**:246, 249
 Ars A, **42**:102, 108–109
 light chains, **42**:139–141
 autoimmune demyelinating disease and, **49**:358–359, 365, 367
 autoimmune thyroiditis and, **46**:289, 296, 304, 312
 autoreactive T cells and, **45**:420, 424–426
 B cell formation and, **41**:223, 225, 237
 CD4 and CD8 molecules and, **44**:268, 270, 299, 302
 CD4 molecular biology, **44**:291–293, 297
 CD8 molecular biology, **44**:280, 282, 284
 CD5 B cell and, **47**:159
 Ig gene expression, **47**:151, 152, 156
 physiology, **47**:133–135, 138
 CTL and, **41**:150, 168, 169
 fetal liver, **42**:131–132
 genetically engineered antibody molecules and, **44**:65, 66
 chimeric antibodies, **44**:79–81, 87
 cloning, **44**:73, 75
 expression, **44**:70, 71, 75, 77, 78
 fusion proteins, **44**:86
 lymphoid mammalian cells, **44**:69
 production, **44**:67
 helper T cell cytokines and, **46**:116
 HIV infection and, **47**:391
 human–human
 advantages, **38**:300
 antigens, **38**:294–297, 299
 autoantibodies, **38**:297–299
 fusion, **38**:292, 293
 propagation, **38**:302–304
 human–murine
 advantages, **38**:291, 292

- Hybridomas (*continued*)**
- heterohybridoma antibodies, **38**:288–290
 - Ig secretion, **38**:287, 288
 - propagation, **38**:302
 - humoral immune response and
 - antigens, **45**:15
 - interleukins, **45**:63, 65, 74, 75
 - physical interaction, **45**:36–39, 44, 50
 - IgE biosynthesis and
 - antibody response, **47**:8, 32, 33, 35–39
 - binding factors, **47**:13, 15, 17, 18, 21, 25–27
 - Ig heavy-chain variable region genes and, **49**:2, 35, 37, 41
 - IL-1 and, **44**:178, 189
 - immunosenescence and, **46**:249
 - mAbs and, **38**:279, 302, 306
 - murine lupus models and, **46**:72, 74, 78
 - neonatal development and, **42**:28–29
 - A/J mice, **42**:131–132
 - somatic mutation, **42**:66–67
 - synthetic T and B cell sites and
 - antigens, **45**:232, 234, 237
 - globular protein antigens, **45**:213, 215, 216
 - immunological considerations, **45**:200
 - viral antigens, **45**:217, 225, 228
 - T cell, *see* T cell hybridomas
 - T cell activation and, **41**:1, 23, 25
 - T cell development and, **44**:215, 216, 218
 - T cell receptor and, **45**:151
 - cDNA, **38**:6
 - gene expression, **38**:13, 14
 - mice, **38**:5, 14
 - protein properties, **38**:3, 4
 - rearrangement, **38**:11, 12
 - specific binding, **38**:19–21
 - target recognition, **38**:122, 123
 - V region genes, **38**:15, 16
 - T cell subsets and
 - H-2 alloantigen recognition, **41**:79, 82, 91
 - H-2-restricted antigen recognition, **41**:62, 70, 52–60
 - T cell receptor, **41**:41, 42
 - T cell triggering, **41**:111
 - technology, **43**:1
- Hybrid resistance, **41**:333, 334**
- antigen expression genetics, **41**:397–401
 - hemopoietic cell grafts, **41**:404–407
- Hh-1, **41**:401–403**
- marrow allograft reactivity, **41**:408–410
 - transplanted cells, **41**:403, 404
 - effector mechanisms, **41**:369, 370
 - antibodies, **41**:376–380
 - macrophages, **41**:370–372
 - marrow engraftment, **41**:384–388
 - marrow microenvironment, **41**:390–393
 - NK cells, **41**:372–376
 - syngeneic stem cell functions, **41**:388–390
 - T cells, **41**:380–384
 - in vitro* assays, **41**:393–396
 - hemopoietic cells
 - bone marrow cells, **41**:335–351
 - lymphoid cells, **41**:351–358
 - leukemia/lymphoma cells, **41**:358–369
- Hybrids**
- antinuclear antibodies and, **44**:108, 129
 - B cell formation and, l9O, **41**:226
 - CD4 molecules and, **44**:285, 296
 - CD8 molecules and, **44**:272, 273, 282
 - CTL and, **41**:169
 - β_2 -microglobulin, **41**:156
 - exon shuffling, **41**:139, 142, 144, 145, 147
 - HLA class I antigens, **41**:150, 151
 - mAbs, **41**:158
 - genetically engineered antibody molecules and, **44**:77, 78, 80
 - Ig gene superfamily and, **44**:12, 27, 31, 40, 47
 - IL-1 and, **44**:167
 - Ir* genes
 - competitive inhibition, **38**:109
 - complementation, **38**:72
 - cross-reactive lysis, **38**:186
 - cytotoxic T cells, **38**:163, 164, 169
 - gene dosage, **38**:80, 82, 83
 - H-Y antigen, **38**:176
 - I region mutations, **38**:85, 90
 - T cell repertoire, **38**:116, 117, 144, 145
 - T cell-T cell interactions, **38**:155, 160
 - lymphocyte homing and, **44**:329, 336, 337, 353
 - resistance phenomenon, NK cells in, **55**:366–367
 - T cell development and, **44**:217, 218, 227
- Hydrazone, *in linkage procedure for***
- chemoimmunoconjugates in cancer treatment, **56**:322–323

- Hydrocortisone
 hybrid resistance and, **41**:387, 390
 in vitro IgE synthesis, **61**:370–371
- Hydrolysis
 in signal transduction, **54**:418–419
 T cell activation and, **41**:15, 16, 23, 25, 30
- Hydrolytic enzymes, cytolysis and, **41**:275–277, 281
- Hydropathicity, mast cell-specific receptor, β subunit profile, **43**:288
- 5-Hydroperoxy-5,8,11,13-eicosatetraenoic acid, dehydration to LTA₄, **39**:150–151
- Hydrophilicity, **43**:76
 epitope prediction, **43**:56
 intrinsic parameters, **43**:49
 MHR, **43**:35
 profiles, **43**:14
- Hydrophobic domains
 human T cell activation and, **41**:5, 13, 25
 T cell subsets and, **41**:45, 54
- Hydrophobicity
 B cell formation and, **41**:197, 198
 CD1 proteins, **59**:37, 38
 CD4 molecules and, **44**:285
 CD8 molecules and, **44**:273
 cytotoxicity and
 cell-mediated killing, **41**:295
 cytolysis, **41**:279, 280
 membrane attack complex of complement, **41**:299–302
 pore formers, **41**:314–317
 genetically engineered antibody molecules and, **44**:67
 Ig gene superfamily and, **44**:3, 6, 35, 46
 IL-1 and, **44**:158, 181
- Hydrophobic regions, **43**:76
- 5-Hydroxyeicosatetraenoic acid
 in neutrophils, human, **39**:153–154, 166–167
 production by eosinophils, **39**:204
- 21-Hydroxylase, maps of Ig-like loci and, **46**:28, 32, 33
- 4-Hydroxy-3-nitrophenyl acetyl, genetically engineered antibody molecules and, **44**:66, 69, 80, 85, 86
- 20 α -Hydroxysteroid dehydrogenase
 in IL-3-dependent cell lines, **39**:35, 36
 IL-3-induced in T cells, **39**:2, 5, 18–19, 23–31
- N-Hydroxsuccinimide, in
 chemoimmunoconjugate crosslinking in cancer treatment, **56**:316–318
- Hydroxyurea
 Ig-secreting cells and, **40**:9
 NK cells and, **47**:230, 231, 283
- HyHEL5
 binding, **43**:20–21
 blind peptide mapping study, **43**:26
 fine specificity study, **43**:26
 lysozyme binding, **43**:12–13, 122
 model, **43**:121
- HyHEL10, blind peptide mapping study, **43**:26
- Hypereosinophilia, reactive, and neoplasms, **60**:225
- Hypereosinophilic syndrome
 clinical picture, **60**:223–224
 eosinophils
 cocultured with cytokines, **60**:185
 synthesis of cytokines, **60**:199–200
 high eosinophil counts, **60**:217
 small granules in, **60**:160
- Hypergammaglobulinemia, murine lupus models and, **46**:63, 64
- Hyper IgE recurrent infection, **62**:89
- Hyper IgM, HIGMX-1
 defect of B cell differentiation, **60**:42–48
 and XLA, immunodeficiency diseases, **60**:37
- Hyper IgM syndrome
 gp39 structure in, **63**:49–50
 X-linked, **59**:248
- Hyperimmunization, secondary B cells and, **42**:72
- Hyperlipidemia, spontaneous autoimmune thyroiditis and, **47**:440
- Hypermutation
 in germinal centers, **53**:233–234
 Ig gene superfamily and, **44**:18, 20, 22, 23
 Ig somatic, **60**:277–278
 locus-specific, **60**:278–280
 somatic
 in germinal centers, **53**:233–234
 T cell subsets and, **41**:42, 88, 106
 V(D)J, **60**:284
- Hyperprolactinemia, **63**:409–410, 430
- Hypersensitivity
 antigen-presenting cells and, **47**:104, 105
 B cell formation and, **41**:183

- Hypersensitivity (*continued*)
 delayed-type, *see* Delayed-type hypersensitivity
 immediate
 eosinophil effects, 39:217–219
 histamine secretion and, 39:217, 218, 304–306
 neuropeptide effects, 39:304–307
- Hypersensitivity lung diseases, 59:387, 412–420
- Hypersensitivity pneumonitis, 59:419–420
- Hypertriglyceridemia, cachectin and, 42:214, 218
- Hypogammaglobulinemia, hybridomas and, 38:287
- Hypogammaglobulinemia, 59:141
 CD23 antigen and, 49:174
 Crohn's disease and, 42:300
 Ig heavy-chain variable region genes and, 49:3
- Hypophysis, IFN- γ effect on, 62:93
- Hypothalamic–pituitary–adrenal axis, 54:30–31; 61:367–368, 375
- Hypothalamus, NK cells and, 47:266
- Hypothyroidism, 46:263, 269, 274, 275, 278
 spontaneous autoimmune thyroiditis and
 altered thyroid function, 47:456
 breeding, 47:435–437
 clinical symptoms, 47:438
 disturbed immunoregulation, 47:470
 histopathology, 47:442, 444
- Hypoxanthine-aminopterin-thymidine,
 lymphocyte hybridomas and, 38:280, 286
 heterohybridomas, 38:288, 289
 human-human hybridomas, 38:293–296, 300
 human-mouse heterohybridomas, 38:300, 301
- Hypoxanthine-guanine phosphoribosyl-transferase
 B cell hybridomas and, 38:279, 280
 human-murine hybridomas, 38:289
- Hypoxanthyl ribosyl transferase gene, 61:113–114
- Hypoxia, IL-6 expression, 65:27
- I
- I ϵ , 62:43, 49–50
- I γ 2b, 62:43, 49–50
- Ia-like molecules, expression by T cells 156, 38:157, 159, 160
- Ia molecules
 on activated and resting B cells
 B cell antigen-presenting capacity and, 39:73–78
 expression, 39:74–76
 MLR and, 39:81–82
 structure, 39:73–74, 76–78
- allogeneic, Ir genes and, 38:115, 126, 132
- corneal graft, 48:199
- IgE biosynthesis and, 47:9, 23, 27
- Ir genes and, *see also* Ia molecules,
 allogeneic; Self-antigens, self-Ia
 antigen binding, 38:98–100, 103
 antigen processing, 38:96, 97
 antigen-specific clones, 38:121
 blocking, 38:69–72
 competitive inhibition, 38:108, 110
 complementation, 38:75–77
 cytochrome c, pigeon, 38:77–79
 discovery of, 38:67, 68
 gene dosage, 38:79–83
 gene function, 38:91–93
 H-Y antigen, 38:179
 immunodominance, 38:184
 I region mutations, 38:84–91
 macrophage factors, 38:111–114
- T cells
 activation, 38:104–107
 cytotoxic, 38:161, 163, 165, 167
 proliferation, inhibition of, 38:68, 69
 selection, 38:131, 132, 134
 suppression, 38:131, 135, 136, 138, 139, 142, 143, 149–152
 T cell-T cell interactions, 38:153, 156, 159, 160
 tolerance, 38:122, 123, 125, 126
 xid gene, 38:83, 84
- MG and, 42:256–257
 positive cell, 48:197
- I-A molecules, role in V β selective element action, 50:20–21
- I κ B, 63:157–159; 65:117
 deficiency effects, 65:32
 description, 58:2–4
 family, 65:14–15
 knockout mice, cytokine induction, 65:32
 phosphorylation by TNF α , 58:8

- rapid degradation and release of NF- κ B, 58:4–5
- I κ B α** , 65:14
gene, 65:23
- I κ B family**, transcriptional regulation, 65:14–15
- IBD**, *see* Inflammatory bowel disease
- Ibuprofen**, 62:184, 186
- ICAM**, *see* Intercellular adhesion molecule
- Icocomes**, humoral immune response and, 45:24
- ICP47**, 65:279
- Id**, *see* Idiotype
- IDDM**, *see* Insulin-dependent diabetes mellitus
- Idiopathic pulmonary fibrosis**, 62:261, 267, 270, 286
- Idiotope-determining region**, 43:14, 16
- Idiotype**
autoimmune thyroiditis and, 46:318, 319
humoral responses, 46:302, 304–308
prevention, 46:315
binding to rabbit anti-Id antibodies, 39:261–263
cross-reactivity, 39:263–265
network theory, 39:255–257
recognition by suppressor T cells, 40:142
recognition by T cells, 39:257–261
B cells and, 39:259–260
- Idiotypic determinants**, Ars A response and, 42:138
- Idiotypic interactions**, human
allergic responses and, 39:175–176
antiself antibodies and, 39:278–279
autoimmune diseases and, 39:277–278
anti-Id antibodies as mediators of, 39:279–280
- B cell tumor therapy** with anti-Id antibodies and, 39:289, 291
immune responses to HLA and, 39:281–283
during pregnancy, 39:281–282
self-tolerance and, 39:276–277
- Idiotypic map**, 42:129–130
- Idiotypic regulation**, immunosenescence and, 46:247–249
- IELs**, *see* Intraepithelial lymphocytes
- I-E molecules**, role in V β selective element action, 50:18–19
- Ig- α** , B cell AgR, tyrosine phosphorylation, 55:246–248
- Ig- β** , B cell AgR, tyrosine phosphorylation, 55:246–248
- IgA**, 61:79, 80, 94–97
antibody concentration on mucosal surfaces, 58:298
antigen-presenting cells and, 47:52, 53
aPL antibodies and, 49:204, 224–226, 232, 238
biosynthesis, 40:184–226; 47:7, 19
cellular and molecular aspects, 40:184–190
common mucosal immune system in human and animal models, 40:197–207
factors regulating S-IgA immune response, 40:207–226
IgA and, induction of, 40:190–192
induction of IgA immune response, 40:190–192
isotype-specific, accessory cells and, 40:219
mucosa-associated lymphoid tissue, 40:192–197
specific humoral, regulation by Fc region of Ig, 40:71–77
specific *in vitro* humoral, enhancement by Fc portion of Ig, 40:90–95
specific *in vitro* T cell-mediated, enhancement by Fc fragments of Ig, 40:95–100
- CD5 B cell** and, 47:138
- distribution of cells producing, 40:189–190
in eosinophil degranulation, 60:182
fluorescence polarization, rotational correlation times, 48:16–17
- function, 65:251
- heavy chain class switch of, 40:219–223
- IBD** and, 42:298, 301–303, 306–307, 321
interaction with nonlymphoid cells, 40:168–184
- epithelial cells, 40:174–177
- erythrocytes, 40:174
- hepatocytes, 40:177–184
- phagocytic cells, 40:168–174
- production, 65:248–251
- CD40 signaling, 63:64–65, 66–67
- respiratory immunity, 59:373, 387, 396, 411, 414

- IgA (*continued*)
 secretory, interaction of component polypeptide chains, **40**:163–168
 secretory component
 binding of, **40**:165–166, 167, 186
 epithelial cells and, **40**:175–177
 hepatocytes and, **40**:179–180, 182
 structure and function, **40**:160–163
 segmental flexibility, **48**:14–18
 of serum and secretory, structure and function of component polypeptide chains, **40**:154–168
 structure, **65**:248
 surface antigens of human leukocytes and, **49**:91
 transforming, expression, **54**:262
- IgA₁, wide extreme, **48**:30
- IgA₂, rotation freedom, **48**:31
- IgA deficiency, **65**:251–256, 263
 clinical manifestations, **65**:246–248, 256
 common variable immunodeficiency (CVID) and, **65**:256
 genetic susceptibility, **65**:256–260
 pathogenesis, **65**:260–263
 treatment, **65**:263
- Ig complex, **52**:136–138
- IgD, **61**:82–83
 antigen-presenting cells and, **47**:48
 B cell expression, **59**:322–324
 biosynthesis and, **47**:7, 19
 CD5 B cell and, **47**:121, 122
 CD23 antigen and, **49**:150, 168, 175
 expression by B cells, **40**:19–26
 and IgM, differential signaling, **54**:401–402
 surface, and surface IgM, respective roles, **52**:297–298
 surface antigens of human leukocytes and, **49**:76, 91
- IgE, **61**:91
 60-kDa factor, **43**:304–305
 allergic airway inflammation, **62**:283
 allergic response, **61**:374–375
 α chain, **43**:282
 anti-dinitrophenol, **43**:294
 antigen-presenting cells and, **47**:63
 anti-ovalbumin, **43**:294
 binding site localization, **43**:279–280
 biosynthesis, **43**:303; **47**:1
 antibody response, **47**:1–3
 B cells, **47**:7–12
- T cells, **47**:8, 10–12
 in vitro, **47**:3–7
 antibody response suppression, **47**:28
 antigens, **47**:28–30
 anti-IL-4, **47**:31
 GIF, **47**:32–35
 interferon-γ, **47**:31, 32
 T cell hybridomas, **47**:36–39
- binding factors
 biological activities, **47**:12–15
 FcεRII structure, **47**:19–22
 formation, **47**:22–28
 physicochemical properties, **47**:15–19
- CD23 antigen and, **49**:149–150, 174, 176
 biochemical structure, **49**:155–157
 biological activity, **49**:167–173
 cellular expression, **49**:150–152, 154–155
 cleavage regulation, **49**:158
 expression regulation, **49**:161
 FcεRII, **49**:162–163, 165
- CD40 expression, **61**:93–94, 95–96
 dependent granule protein stimulation, **60**:207
- eosinophils, **43**:306–307
- ε chain, **43**:278
- Fcε molecules dependence on, **52**:385
- Fc receptors, **43**:277
 fluorescence polarization, rotational correlation times, **48**:16–17
 functional characteristics, **43**:305
 functions, **48**:18
 high-affinity receptor, **43**:281
 histamine-releasing factors, **43**:307
 IL-4 stimulated production, **54**:247–248
 and IL-4 switching, **52**:217–218
 interaction with eosinophils, **60**:181
 isotype switching, **60**:38–39
 macrophage receptor, **43**:306
 monoclonal anti-β antibody, **43**:283
 monocytes, **43**:305–306
 myeloma, ESR spectra, **48**:30–31
 plate-bound, IL-4 production in response to, **53**:20–21
- platelets, **43**:306–307
 production, CD40 signaling, **63**:62–64, 66–67
- protein-binding sites, **43**:281–282
 receptor binding to, **52**:387–388
 regulation, **61**:369–370
 role, **43**:277

- segmental flexibility, **48**:18–19
 sequential switching, **61**:105
 sites interacting with receptors, **43**:278–279
 anti-IgE antibody studies, **43**:279–281
 inhibition studies, **43**:281–282
 structural properties
 comparative studies on rodent cells,
43:302–305
 Fc ϵ receptors, **43**:298–301, 303
 IgE-binding factor, **43**:300–302
 surface antigens of human leukocytes and,
49:77, 81, 90–91
 synthesis, **43**:277; **61**:342–344, 370–374
 wide extreme, **48**:30
- IgE-binding factors
 CD23 antigen and, **49**:149
 cDNA coding, **43**:304
 functional characteristics, **43**:305
 human B cells, **43**:300–301
 retroviral protein, **43**:304
 T cells, **43**:301–303
- IgE-binding proteins, **43**:307–308
- Ig/EBP, **65**:3–5
- IgE/Fc ϵ RI, γ dimer, **52**:380–381
 IgE/Fc ϵ RI receptor, high-affinity,
52:378–381
 assembly, **52**:381–382
 cell surface structures functionally
 associated with Fc ϵ RI molecules,
52:391–396
 cell surface glycolipids, **52**:393–394
 Fcy receptors, **52**:394–396
 G63, **52**:394
 mast cell chromolyn binding sites,
52:392–393
 ME491 (CD63), **52**:393
 chromosomal location, genomic
 organizations, and mRNA species,
52:382–383
 control of synthesis and expression of
 dependence on IgE, **52**:385
 Fc ϵ RI molecules, **52**:384–386
 half-life, internalization, degradation and
 reexpression, **52**:385–386
 synthesis during cell cycle, **52**:385
 synthesis during differentiation of
 basophils/mast cells from
 hemopoietic precursor cells,
52:384–385
- functional characterization of receptor,
52:387–391
 binding to IgE, **52**:387–388
 effector mechanisms and functional
 changes of mast cells/basophils
 following activation through IgE
 binding sites, **52**:391
 signal transduction and distal events,
52:388–391
- numbers and binding constants on
 basophils/mast cells, **52**:386–387
 sequence and structural homologies,
52:383–384
 conversation during evolution, **52**:383
 molecules related to Fc ϵ RI α chain,
52:383
 molecules related to Fc ϵ RI β chain,
52:383–384
 molecules related to Fc ϵ RI γ chain,
52:384
- topology of, **52**:378
 α chain, **52**:379
 β chain, **52**:379–380
 γ dimer, **52**:380–381
 nomenclature, **52**:378
- IgE-potentiating factor, **47**:5
 IgE receptors, eosinophil, and killing of
 schistosomula, **39**:180–181
- Ig fold, **43**:221
- IgG, **61**:80, 94
 adoptive T cell therapy of tumors and,
49:296
 antigen-presenting cells and, **47**:62, 67, 85
 aPL antibodies and, **49**:199, 242, 251, 255,
 258
 affinity purification, **49**:229
 clinical aspects, **49**:204–206, 210, 212,
 222, 224
 isotype, **49**:224–228
 pathogenic potential, **49**:253, 255
 specificity, **49**:238, 242–243
 syndromes, **49**:220
- B cell formation and, **41**:223, 237
 biosynthesis and
 antibody response, **47**:2–11, 31, 32
 binding factors, **47**:12, 19, 27
 CD5 B cell and, **47**:132, 134, 136, 144
 CD23 antigen and, **49**:157
 CD40 signaling, **63**:63, 65–66, 69–70
 CH3 domain, **48**:20–21

- IgG (*continued*)
 complex and associated second messengers, identification, 54:419–420
 conformation and flexibility, 51:4–7
 cytotoxicity and, 41:271
 eosinophil-bound, parasite killing, 39:209–211
 hybrid resistance and, 41:336, 355, 357
 effector 376–378, 41:394
 IBD and, 42:297–298
 antibody expression in, 42:321
 subclass expression in, 42:305
 Ig heavy-chain variable region genes and, 49:2, 6, 49–51
 IL-4-stimulated production, 54:247–248
 induced monocytes, 54:172, 173
 NK cells and, 47:189, 293
 cell-mediated cytotoxicity, 47:190, 196
 cytotoxicity, 47:249, 256
 effector mechanisms, 47:237, 243–245, 247, 248
 malignant expansion, 47:227
 surface phenotype, 47:201, 202, 212
 physicochemical characterization of Fc receptor for, 40:69–70
 segmental flexibility, 48:5–19
 early studies, 48:5–6
 fluorescence polarization, 48:6–10
 hinge region structure, 48:21–23
 IgG subclasses, 48:11–14
 spin-label method, 48:10–11, 12
 X-ray crystallography, 48:11
 surface antigens of human leukocytes and, 49:83, 89–90
- IgG1, 48:13
 CD40 expression, 61:95–96
 heavy chain, hinge region structure, 48:21–22
 oligosaccharide, ¹³C nuclear magnetic resonance, 48:29
- IgG2, 48:13
- IgG2a, eosinophil-bound, parasite killing, 39:210–212
 inhibition by IgG2C, 39:211–212
- IgG-Fc_{RIII} interaction at molecular level, 51:40–41
- IgG receptor, isotypes, 54:343–344
- IgH
 B cell formation and, 41:198
 CD5 B cell and, 47:129, 161
- Igh haplotypes, *J*558 and, 42:135
 Ig recombinant strains, *V_H* gene segment and, 42:6
- Igκ
 gene 3' enhancer, 62:43, 49
 gene enhancers, 62:43, 50–51
- Ig_l gene
 diversification, potential mechanisms, 48:58
 somatic diversification, 48:52, 53
- IgM, 42:13; 48:16–18; 61:79, 80, 82, 93
 adoptive T cell therapy of tumors and, 49:314, 331
 antigen-presenting cells and, 47:48, 85
 aPL antibodies and
 affinity purification, 49:228–229
 clinical aspects, 49:204, 206, 209–210, 212, 222
 B cell expression, 59:322–324
 B cell formation and, 41:183
 B cell precursors, 41:194, 200
 inducible cell line, 41:221, 222
 population dynamics, 41:206
 soluble mediators, 41:234
 B cells secreting, 40:22–23
 biosynthesis and, 47:4, 6, 7, 9–11
 CD5 B cell and, 47:129
 physiology, 47:130–133, 135, 136
 CD23 antigen and, 49:159, 167–168, 172, 175
 CD40 signaling, 63:59, 62–63, 66–67
 fluorescence polarization, rotational correlation times, 48:16–17
 human myeloma, 48:16–18
 hybrid resistance and, 41:336, 377, 378
 IBD and, 42:299, 300
 and IgD, differential signaling, 54:401–402
 Ig heavy-chain variable region genes and, 49:38, 41–43, 48–51
 myeloma, ESR spectra, 48:30–31
 NK cells and, 47:207, 292, 293
 normal or elevated, 60:37–49
 secondary B cells and, 42:76
 secretion in NZB spleen cells, 55:313–314
 segmental flexibility, 48:14–18
 surface, and surface IgD, respective roles, 52:297–298
 surface antigens of human leukocytes and, 49:76, 91, 101
 X-linked hyper IgM syndrome, 59:248
- IgM-binding protein, 52:173

- IgM hybridoma, somatic mutation and, **42:67**
 IgM receptors, on eosinophils, human, **39:180**
 Ig oligosaccharide, spin-labeling, **48:27–28**
 Ig+ precursor selection, bursa of Fabricius, **48:49–51**
 IGUP I-5111, **64:21, 23**
 Ig VL gene
 gene conversion, **48:60–62**
 somatic diversification, **48:60–62**
 Ikaros
 B cell genesis, **63:218–219, 224–226**
 gene expression, **63:224–226**
 IL-1ra, *see* Interleukin-1 receptor antagonist
¹²⁵I-labeled antigen therapy, for MG, **42:268**
¹²⁵I-labeled IL-2, **42:172**
 Ileum, lymphocyte homing and, **44:330**
 Illegitimate priming model, switch
 recombination, **61:108–111**
 Imidoesters, in chemoimmunoconjugation, **56:320–321**
 Immune complex disease, pulmonary, rat model, **60:345**
 Immune complexes
 and alternative antigen pathway, trapping, **51:265–269**
 cell-mediated immune responses and, **40:77–79**
 lung inflammation, **62:277–283**
 regulation of immune responses and, **40:71–72, 73–76**
 regulators of complement activation and, **45:388, 389, 410, 411**
 Immune function, animal models
 EBV-associated lymphoproliferative disease in hu-PBL-SCID mice, **50:310–313**
 HIV-infection of hu-PBL-SCID mice, **50:314–317**
 other tumor models in SCID mice, **50:313–314**
 search for better models, **50:303–304**
 transfer of normal or autoimmune human cells to SCID mice, **50:304–310**
 viral pseudotypes in HIV-infected hu-PBL-SCID mice, biosafety issues and, **50:317–319**
 Immune interferon, *see* Interferon- γ
 Immune intervention, IL-2R-targeted, **50:187–189**
 Immune privilege
 experimental autoimmune uveitis, **48:216–219**
 eye, **48:191–221**
 Immune reconstitution, marrow grafts and cytotoxic cells, accessory cells and neutrophils, **40:416–417**
 effects of T cell depletion, **40:417–419**
 humoral immunity, **40:414–416**
 phenotype and function of peripheral T cells, **40:412–414**
 repopulation of lymph nodes and thymus, **40:411–412**
 Immune regulation
 autoimmune uveitis, **48:215–216**
 eye, **48:191–221**
 intraocular tumor rejection, **48:212–215**
 neoplasm, **48:212**
 spontaneous neoplasm, **48:212**
 Immune repertoire, self-tolerance
 checkpoints, **59:319–342**
 Immune response, **52:125; 65:111, see also**
 Antibody response; Cellular immune response; Cellular immunity; Humoral immune response
 affinity maturation, **43:126–127**
 antigens, **64:1–3**
 assessment, **43:6**
 cellular dynamics, **60:267–268**
 diversity in, antigen receptors, **43:106**
 environmental antigen influences, **40:207–209**
 factors regulating, accessory cells and isotype-specific response, **40:219**
 gene control, **43:206–208**
 heavy chain class switch, **40:219–223**
 to HIV, **65:290–293**
 acute infection, **65:293–296**
 adverse effects of CTL activity, **65:309–311**
 apoptosis, **65:306**
 asymptomatic period of infection, **65:296–297**
 CTL decline in late disease, **65:300**
 CTL exhaustion, **65:304–305**
 disease progression, **65:297–298**
 escape mutation, **65:311–317**
 long-term nonprogressors, **65:298–300**
 in seronegatives, **65:287, 307–309**
 Th1 to Th2 switch, **65:302–304**
 therapeutic implications, **65:317–322**

- Immune response (*continued*)
 to MMTV, **65**:167–168, 212
 adult T cell response, **65**:175–178
 cellular response, **65**:192–194
 humoral response, **65**:188–192
 neonatal response, **65**:171–175, 203
 receptors for, **65**:194–196
 superantigen and T cell-independent, **65**:168–171
 superantigen dependent, **65**:171–194
 T–B interaction, **65**:180–188
 T cell-dependent B cell differentiation, **65**:178–180
 negative regulation, role of CTLs, **60**:303–310
 ontogeny, in rabbits, **56**:186–187, 209
 oral tolerance, **40**:223–226
 phenotype, correlation with antigen-MHC interactions, **43**:212
 plasmacytogenesis, **64**:225–226
 primary humoral, histology, **60**:269–272
 regulation, IL-6 effects, **54**:21–22
 B cells, **54**:21
 T cells, **54**:21–22
 role of T cell Fc receptor for isotype response, **40**:212–216
 to superantigens, **54**:117–127
 binding to major histocompatibility complex, **54**:117–120
 direct non-T cell effects, **54**:127
 recognition by T cell receptor, **54**:121–124
 role of accessory CD4 or CD8 molecules, **54**:124–125
 T cell deletion and anergy in injected animals, **54**:125–127
 T cell signaling in response to superantigens vs conventional antigens, **54**:127
 T cell networks in secretory immune system, **40**:217–218
 T cell regulation of IgA response, **40**:209–212
 Td B cell response, **62**:131–157
 tumor antigens, **62**:217–245
 against tumors. *see* Tumors, immune response
 Immune response genes, **38**:32, 33; **43**:207, 212
 antigen binding, **38**:98–115
 antigen processing, **38**:93–97
 antigen specificity, **38**:36, 37
 complementation, **38**:72–83
 cytotoxic T cells
 classical type, **38**:161–174
 epistatic effects, **38**:175–187
 expression
 in antigen-presenting cells, **38**:55–57
 B cell responses, **38**:50, 52
 in B cells, **38**:57, 58
 codominance, **38**:36, 71, 80, 81, 97
 complementation, **38**:73, 79, 82, 83
 H-Y antigen, **38**:175, 177
 Ia molecules, **38**:83, 84
 I region mutations, **38**:84, 88, 90, 91
 in lymphoid tissue, **38**:52–55
 MHC restriction 65, **38**:67
 T cells, **38**:40–43, 58–63
 cytotoxic, **38**:165, 167
 interaction, **38**:156, 157, 159, 160
 positive selection, **38**:130, 134
 repertoire, **38**:116, 117
 tolerance, **38**:127
 function, theories of, **38**:91–93
 gene function models, **38**:37–39
 genetic locus, **38**:33–37
 humoral immune response and, **45**:5
 hybrid resistance and, **41**:361
 Ia molecules, **38**:67–72
 I region mutations, **38**:84–91
 MHC, **38**:43–46
 MHC restriction, **38**:63–67
 polypeptides, responses to, **38**:46–50
 “Schlepper” experiment, **38**:39, 40, 42
 sporozoite malaria vaccine and, **45**:308–310
 synthetic T and B cell sites and antigens, **45**:232
 globular protein antigens, **45**:210, 211, 214
 immunological considerations, **45**:198–201
 viral antigens, **45**:221
 T cells
 germline repertoire limitations, **38**:115–118
 interactions, **38**:153–160
 positive selection, **38**:130–134
 receptors, **38**:2
 suppression, **38**:134–152

- T cell subsets and, **41**:44, 51, 54, 55
tolerance, **38**:119–130
- Immune states, and autoimmune and immunodeficiency states, antigen processing and presentation, **52**:106–108
autoimmunity, **52**:107
novel immunodeficiency states, **52**:107–108
tumor immunology, **52**:104–106
vaccines, **52**:106–107
- Immune surveillance theory, **48**:212
- Immune system, **63**:269
B cell, and memory
affinity maturation, **53**:233–234
bcl-2 oncogene role, **53**:241
Fas antigen/APO-1 role, **53**:241–242
germinal centers role, **53**:231–233
intermitotic life span and, **53**:237–241
memory cells
functional properties, **53**:236
generation of, **53**:236–238
phenotype of, **53**:234–236
peripheral, compartmentalization
anatomic localization of colonizing lymphocytes, **53**:159
CD5⁺/CD5 cell dichotomy, **53**:178–183
generation of compartments, **53**:187–194
lymph follicle B cells, **53**:185–186
peritoneal B cells, **53**:183–185
significance of, **53**:199–203
site-dependent tolerance, **53**:201–203
specialized functions of, **53**:199–201
- T cells
intestinal intraepithelial, **53**:163–167
 γ/δ invariant intraepithelial cells, **53**:167–169
in liver, **53**:169–171
in lymph nodes, **53**:161–163
in peritoneum, **53**:173–177
in spleen, **53**:161–163
- common mucosal, **40**:197–199
in humans, **40**:199–204
mechanism of homing of plasma cells to mucosal tissues and secretory glands, **40**:204–207
- complement system, **61**:201–257
- Ets transcription factors, **64**:65–94
- evolution of antigens, T_{CD8⁺}, what it recognizes, structure of class I molecules, **52**:6–8
- glucocorticoids and, **66**:197, 218
high endothelial venules, **65**:373, 375–377
hyperprolactinemia, **63**:409–410
IL-1ra effects, **54**:202–204
lymphocyte migration, HEVs and, **65**:350–372
nonspecific, pituitary hormones, **63**:407–408
peripheral compartments
B cell, **55**:429–430
T cell, **55**:424–429
- pituitary gland, **63**:377, 407–408
- programmed cell death in, *see* apoptosis, in immune system
- secretory, T cell networks in, **40**:217–218
- soluble cytokine receptors, **63**:285–290
- T cell, and memory
bcl-2 oncogene role, **53**:241
differentiation after antigenic stimulation, **53**:229–230
effector cells, **53**:227–230
Fas antigen/APO-1 role, **53**:241–242
intermitotic life span and, **53**:237–241
memory cells, **53**:227–230
naive cells, **53**:227–230
phenotypic identification of memory T cells, **53**:224–227
TCR and signal transduction, **53**:238–239
- V(D)J recombination, **64**:39–61
- Immunity, *see also* Cellular immune response; Cellular immunity; Humoral immune response
adaptive, NK cells and, **47**:291–295
antitumor, antigen-presenting cells for, **58**:436–437
epithelial and mucosal, **58**:297–298
HIV-1 disease, **63**:95–98
IL-2, **63**:132
to infections and diseases, role of IELs, **58**:326–329
- innate
and gene products regulated by NF- κ B, **58**:20–21
role of NF- κ B and rel proteins, **58**:1–27
role of IELs, **58**:326–329
- systemic
enhancement, **58**:428–429
and MHC class II molecules, **58**:433–434
- tumor-specific, **58**:418–419

- Immunity/tolerance phenomena, **59:279–281**
- Immunization
- antigen-presenting cells and, **47:67, 72, 86, 94**
 - with anti-Id antibodies, animal, **39:283–284**
 - HIV infection and, **47:407–410**
- IgE biosynthesis and
- antibody response, **47:2–6, 8, 9**
 - antibody response suppression, **47:30–32, 36**
 - binding factors, **47:14, 15, 23**
 - NK cells and, **47:282, 292, 293**
 - spontaneous autoimmune thyroiditis and, **47:433**
 - with TT, human responses to, *see* *Tetanus toxoid*
- Immunoadhesins, **64:310**
- Immunocytes, immature, **52:285**
- Immunodeficiencies, **54:142–143; 61:41, 311–312, 317**
- adenosine deaminase deficiency, **59:151, 246**
 - angioedema, **61:206**
 - ataxia telangiectasia, **61:191, 315**
 - autoimmune diseases, **59:146–147; 61:367, 342–344**
 - bare lymphocyte syndrome, **59:246–247; 61:327–338**
 - B cell formation and, **41:199, 200, 207, 216**
 - genetically determined defects, **41:224–226, 231**
 - Bloom's syndrome, **61:316**
 - Bruton's agammaglobulinemia, **61:312–314**
 - common variable immunodeficiency (CVID), **59:138, 143, 147, 150**
 - DiGeorge syndrome, **59:247**
 - DNA breakage syndromes, **61:314–316**
 - factor I, **61:213**
 - Fanconi's anemia, **61:315**
 - idiotypic network and, **39:274–275**
 - IgA deficiency, **65:245–263**
 - Lesch–syndrome, **59:159**
 - major histocompatibility complex, **61:327–338**
 - prevention by apoptosis-inhibitory drugs, **58:278–280**
 - purine nucleotide phosphorylase deficiency, **59:246**
 - selective T cell defect, **59:247**
- severe combined, *see* Severe combined immunodeficiency
- Wiskott–Aldrich syndrome, **59:159, 248**
- xeroderma pigmentosum, **61:315**
- X-linked, **59:248, 317**
- murine, Btk implicated in, **60:41–42**
 - severe combined immunodeficiency, *see* Severe combined immunodeficiency
 - X-linked agammaglobulinemia, *see* X-linked agammaglobulinemia
- Immunodeficiency states, **52:106–108**
- and immune and autoimmune states, **52:106–108**
 - autoimmunity, **52:107**
 - novel immunodeficiency states, **52:107–108**
 - tumor immunology, **52:104–106**
 - vaccines, **52:106–107**
 - novel, **52:107–108**
 - tumor immunology, **52:104–106**
 - vaccines, **52:106–107**
- Immunodeficient mouse
- HuSCID-derived antibodies, **57:230–231**
 - IL-5 production in, **57:171–172**
- Immunodepression
- hybrid resistance and, **41:385**
 - systemic, and T cell PCD, **58:214**
- Immunodiffusion, antinuclear antibodies and, **44:118, 120, 121**
- Immunodominance, cytotoxic T cells and, **38:180–185**
- Immunofixation, **38:207**
- Immunofluorescence
- anticomplement, **38:293**
 - antinuclear antibodies and, **44:109, 113, 120, 125**
 - autoimmune response, **44:133, 136**
 - scleroderma, **44:121, 122, 124, 125**
 - SLE, **44:104–106**
 - autoimmune thyroiditis and, **46:293**
 - B cell formation and, **41:188, 221**
 - CD5 B cell and, **47:118**
 - Ig gene expression, **47:153, 157**
 - lineage, **47:129**
 - marker for activation, **47:127**
 - complement receptor 1 and, **46:193**
 - complement receptor 2 and, **46:204, 210**
 - cytotoxicity and, **41:275, 298**

- direct two-color, for NK cell surface antigens, **42**:200
- Epstein-Barr virus transformation, **38**:277
- human-human hybridomas, **38**:295, 296, 299
- human-murine hybridomas, **38**:290
- IgE biosynthesis and, **47**:4, 8, 19, 21
- immune response (*Ir*) gene complementation, **38**:78
- indirect, in studies of CD antigen expression on basophils and mast cells, **52**:336–337
- leukocyte integrins and, **46**:158, 171
- murine lupus models and, **46**:88, 95
- NK cells and, **47**:211
- spontaneous autoimmune thyroiditis and, **47**:445, 446, 462
- two-color, NKHI antigen analysis, **42**:189
- Immunofluorescent cell sorting, NK clones and, **42**:182–184
- Immunogenicity, **43**:75
- antigen-presenting cells and, **47**:101–105
- multiple antigen peptides based on schistosome antigens, **60**:125–128
- multiple-branched multiple antigen peptides, **60**:140–141
- (T1B)₄ MAP construct, **60**:114–120
- Immunogens
(BT)₄, **60**:109
- multiple antigen peptides in helminthic diseases, **60**:124–128
- in protozoan diseases, **60**:108–124
- for viral and bacterial diseases, **60**:128–136
- Immunoglobulin, **42**:95; **43**:235, *see also* specific IgS
- adoptive T cell therapy of tumors and, **49**:312, 314, 330–331
- in antigen presentation by B cells, **39**:52–53, 55–56, 58–59
- antigen-presenting cells and
- antigen presentation, **47**:66–70
 - antigen processing, **47**:89, 90
 - cell surface, **47**:62–64
 - immunogeneity, **47**:103
 - interaction with T cells, **47**:93
 - tissue distribution, **47**:47
- in antigen uptake by B cells, **39**:60–62
- antinuclear antibodies and, **44**:104, 110, 111, 120, 124
- aPL antibodies and, **49**:194, 199, 259
- affinity purification, **49**:228, 230
- antibody subsets, **49**:232
- binding to cell membranes, **49**:250, 252
- clinical aspects, **49**:200, 203, 211
- isotype, **49**:224
- pathogenic potential, **49**:253–254, 257
- specificity, **49**:237, 241, 243–244, 248
- autoimmune thyroiditis and, **46**:318
- cellular immune responses, **46**:288, 297
- experimental models, **46**:274
- humoral responses, **46**:300, 303–308
- prevention, **46**:315, 316
- autoreactive T cells and, **45**:427
- avian, V_L gene diversification by gene conversion, **48**:60–62
- B cell formation and, **41**:182, 183, 237, 238
- B cell precursors, **41**:188, 189, 191, 193–195, 199–202
- bone marrow cultures, **41**:209, 217, 218
- CBA/N mice, **41**:226
- inducible cell line, **41**:221, 223
- lymphohemopoietic tissue organization, **41**:187
- population dynamics, **41**:206, 207
- SCID mice, **41**:225
- carbohydrate component, mobility, **48**:26–32
- CD4 and CD8 molecules and, **44**:268, 269, 297, 302
- CD4 molecular biology, **44**:286–288, 296
- CD8 molecular biology, **44**:272, 274–276
- CD5 B cell and, **47**:161, 162
- gene expression, **47**:150–159
 - genetic influence, **47**:148
 - lineage, **47**:128
 - malignancies, **47**:123
 - physiology, **47**:137, 138, 140–142
 - surface antigen, **47**:143, 146
- CD23 antigen and, **49**:150–151, 156–158
- cell-mediated killing and, **41**:291
- characterization, **43**:7
- CH switching rearrangement, **43**:239
- class switching by B cells, **60**:270–271
- complement receptor 1 and, **46**:193, 198, 200
- complement receptor 2 and, **46**:203, 205, 207, 208, 211
- CTL and, **41**:168

- Immunoglobulin (*continued*)
 definition of clonality based on, **40**:295–305
 distribution in body, **40**:153
 diversification
 evolutionary mechanism, **48**:62
 mechanisms, **48**:62
 domains, cysteine loop relationships, **43**:285
 effector functions, I flexible, **48**:1
 Fc receptors for, *see* Fc receptors
 fold, **43**:108
 genetically engineered antibody molecules
 and, **44**:66, 89
 antigen-combining sites, **44**:84
 biological properties, **44**:82, 83
 chimeric antibodies, **44**:80–82, 87, 88
 cloning, **44**:73–75
 expression, **44**:71, 72, 75–79
 fusion proteins, **44**:85, 86
 lymphoid mammalian cells, **44**:69, 70
 production, **44**:67–69
 surface Ig, **44**:89
 heavy chain, *see* Heavy chain
 heavy chain variable region genes, *see*
 Heavy chain variable region, genes
 helper T cell cytokines and
 cross-regulation of differentiation, **46**:132
 differential induction, **46**:127, 128, 130
 functions, **46**:114, 116, 118–122
 human subsets, **46**:139
 immune responses, **46**:133, 135, 137,
 138
 precursors of differentiation states, **46**:126
 secretion patterns, **46**:113
 HIV and, **45**:351, 353, 356, 362, 365
 HIV infection and, **47**:390, 406, 408
 human, rotational correlation times, **48**:28–29
 humoral immune response and, **45**:4, 6, 8
 antigens, **45**:10, 12, 13, 16–18, 21
 cellular interactions *in vivo*, **45**:80,
 82–84
 helper T cells, **45**:26, 27, 29, 32–35
 interleukins, **45**:55, 63–72, 76
 physical interaction, **45**:40, 43, 44, 49,
 50, 52
 hybrid resistance and, **41**:357, 365, 366,
 378, 379
- IBD and, **42**:301–307
 IL-1 and, **44**:177–179, 183, 190
 IL-5-regulated production, **57**:151–154
 IgA, **57**:152–153
 IgE, **57**:153
 IgG, **57**:153
 immunosenescence and
 lymphocyte activation, **46**:223, 224, 231,
 232, 234, 236
 lymphocyte subsets, **46**:237, 239
 mucosal immunity, **46**:251
 regulatory changes, **46**:248, 249, 251
 stem cells, **46**:240, 242, 243
 internal movement, **48**:1–34
 intravenous, replacement therapy with,
 60:40
 isotype, B cell subsets and, **40**:19
 isotype switching
 history, **54**:229–231
 in humans, **54**:260–262
 IL-4 role in regulation, **54**:260–262
 TGF- β and IgA expression, **54**:262
 IFN- γ regulation, **54**:256–257
 mechanism of switch recombination,
 54:234–240
 organization of heavy chain locus in
 mouse and human, **54**:231–233
 switch regions, **54**:233–234
 IL-4-induced regulation
 enhancement of switching to IgE and
 IgC, **54**:248–249
 molecular mechanisms, **54**:252–256
 role in T cell-dependent switching,
 54:249–252
 stimulation of IgE and IgG
 production, **54**:247–248
 molecular mechanism, **54**:231–246
 directed switching and germline C_{H}
 transcription, **54**:240–245
 expression of downstream isotypes
 without switch recombination,
 54:245–246
 TGF- β -induced, **54**:257–259
 leukocyte integrins and, **46**:149, 150, 168,
 171
 ligand, **48**:1
 light chain, *see* Light chain
 lymphocyte homing, **44**:326, 327, 335, 336;
 64:147, 157–160
 measles and, **45**:340, 341

- membrane, induced tyrosine phosphorylation, targets, **55**:248–249
- CD22, **55**:245–246
- Ig- α , **55**:246–248
- Ig- β , **55**:246–248
- mitogen-activated protein kinase, **55**:244–245
- p21^{ras}, **55**:238–240
- p95^{cav}, **55**:243–244
- phosphatidylinositol 3-kinase, **55**:241–243
- phospholipase, **55**:234–238
- ras*GAP, **55**:239–240
- ras*GAP-associated proteins, **55**:240–241
- Vap-1, **55**:243
- mRNA, **43**:239
- μ chains, **43**:238–239
- NK cells and, **47**:203, 292–294
- production
- Cd40 and, **63**:58–70
 - switch recombination signals, **63**:58–59
- promoter elements, **43**:269
- B cell preference, **43**:258
 - enhancer interactions, **43**:263–265
 - heavy chain, common factors with enhancers, **43**:256–257
 - interaction between multiple proteins and conserved octanucleotide, **43**:254–255
 - light chain, sequence conservation, **43**:257
 - nuclear factor interaction with conserved heptamer, **43**:255
 - sequence conservation, heavy chain V gene promoters, **43**:252
 - sequences required for heavy chain gene expression, **43**:253–254
 - tissue and stage specificity, **43**:251, 255–256
 - VH promoter activity, **43**:252–253
 - κ activity, required octamer, **43**:257–258
 - κ V gene promoters, **43**:258
- rabbit anti-mouse
- presentation by B cells, **39**:52–53, 55–56
 - B cell activation and, **39**:70
 - uptake by B cells and macrophages, **39**:60–62
- recognition function, **48**:1
- regulating transcription, **43**:240
- regulators of complement activation and, **45**:382, 390, 398
- replacement therapy, **59**:150
- rotational correlation times, **48**:28–29
- SCID and, **49**:388, 400–401
- secretion
- human-murine hybridomas, **38**:287, 288
 - hybridoma propagation, **38**:302
 - requirements for, **40**:83–90
 - segmental flexibility, **48**:5–19
 - electron microscopy, **48**:2–3
 - fluorescence polarization, **48**:3–5
 - hydrodynamic study methods, **48**:2
 - nanosecond, **48**:4–5
 - spin-label approach, **48**:5
 - steady state, **48**:4
 - study methods, **48**:1–34
 - X-ray crystallography, **48**:3
- sequence selection in chicken B cell development, **57**:369–372
- sequence variability, **43**:11, 14
- somatic rearrangement of gene elements to create functional antibody gene
- γ light chain genes, **40**:251–252
 - heavy chain gene assembly, **40**:252–253
 - heavy chain gene order and class switching, **40**:257–258
 - human Ig light chain genes, **40**:249–251
 - recombinational mechanisms in joining of segments of variable region, **40**:253–256
 - sequential activation of Ig genes, **40**:256–257
- spontaneous autoimmune thyroiditis and altered thyroid function, **47**:458
- cellular immune reactions, **47**:452
- histopathology, **47**:440
- humoral immune reactions, **47**:445, 446, 448
- potential effector mechanisms, **47**:469
 - sporozoite malaria vaccine and, **45**:286, 307
 - structure of, **40**:61–63
- superfamily molecules, **58**:372–376
- superfamily of adhesion molecules, **52**:167
- supergene family, recognition molecules of, **52**:361–364
- CD4 and CD8, **52**:363
 - ICAM-1/CD54, **52**:361–363
 - LFA-2/CD2 and LFA-3/CD58, **52**:361
 - MHC class I and class II molecules, **52**:364

- Immunoglobulin (*continued*)
 other recognition molecules of Ig superfamily, **52**:364
 stem cell factor receptor (SCF-R)/c-kit, **52**:363–364
 VCAM-1, **52**:363
 surface antigens of human leukocytes and, **49**:81–82, 98, 118
 antigen-specific receptors, **49**:76–79
 receptors, **49**:89–91
 surface Ig, B cell proliferation, **63**:56
 synthesis, hybridomas and, **38**:293, 299
 T cell development and, **44**:209, 217, 243
 T cell receptor and, **38**:1, 2
 accessory molecules, **45**:109
 antigen processing, **45**:118
 experimental systems, **45**:131
 homogeneity, **45**:129
 MHC molecules, **45**:111
 protein properties, **38**:3–5
 and T cell receptor gene rearrangements as lineage markers
 acute lymphoid leukemia with 4:11
 chromosomal translocation, **40**:293
 diffuse large cell lymphoma, **40**:292
 hairy cell leukemia, **40**:291–292
 Hodgkin's disease, Lennert's lymphoma and angioimmunoblastic lymphadenopathy, **40**:294
 lymphoid blast crisis of chronic myelogenous leukemia, **40**:290–291
 malignant histiocytosis complicating celiac disease, **40**:292–293
 pseudo T cell lymphoma, **40**:294–295
 transcriptional enhancer elements, *see also* Protein-binding sites
 activation and V-DJ joining, **43**:241–242
 B cell preference, **43**:258
 common factors with heavy chain promoters, **43**:256–257
 containing bind sites, **43**:264
 dependent transcription, **43**:246
EcoRI site, **43**:249
 IgH enhancer-binding proteus, **43**:247–248
 increase local concentration of factors, **43**:264
in vivo protein-binding sites, **43**:243–244
 κ enhancer, **43**:249
 light chain enhancer elements, **43**:249–250
 location, **43**:242
 mechanism, **43**:240–241
 negative regions, **43**:248–249
 promoter interactions, **43**:263–265
 tissue specificity, **43**:242
 transcription initiation, **43**:242–243
 transcriptional regulation mechanisms, **43**:258–259
 enhancer independence and dependence of VH promoter, **43**:265–266
 lymphoid-specific octamer factor, **43**:261–262
 octamer as tissue-specific element, **43**:260–263
 octamer-binding protein forms, **43**:261
 promo/a-enhancer interactions, **43**:263–265
 tissue and stage specificity, **43**:259–260
 ubiquitous factor, **43**:263
 V-D-J joining, **43**:237–239, 241–242
 transgenic mouse, CD5 B cells in, **55**:324–326
 V(D)J joining of antigen receptor genes at loci
 endogenous substrate, **56**:34–37
 gene assembly, **56**:28–32
 Immunoglobulin genes, **59**:332, *see also* Heavy-chain variable region, genes and antibody repertoire generation in rabbit
 C_H, **56**:191–193
 D, **56**:189–191
 IgA, **56**:192
 IgD, **56**:192
 J_H, **56**:188–191
 κ light-chain, **56**:193–194
 λ light-chain, **56**:194
 organization, **56**:187–188
 B cell development, **63**:20–21
 chicken B cell
 conversion
 allelic exclusion, **57**:372, 374
 avian leukosis virus induction of DT40 cell line, **57**:361–362
 rearrangement in bursa, **57**:365–367
 recombination models, **57**:362–365
 organization
 D elements, **57**:356–358

- heavy chain loci, **57**:354–356
light chain loci, **57**:354–356
- defects
heavy chain disease and, **40**:305–306
hereditary primary immunodeficiency disorders, **40**:306–309
- expression
during B cell development, **43**:
236–240
B cell-specific, **43**:253
heavy chain and κ chain, **43**:236–237
tissue specific, **43**:235
- germline, murine lupus models and,
46:65, 66
autoantibodies, **46**:71–76
inducing agent, **46**:78
murine lupus, **46**:66–71
somatic mutation, **46**:76, 77
T cell antigen receptor, **46**:98
- H- and L-chain, **60**:277–278
- human T cell antigen and, **41**:3, 10
- rearrangements, **58**:31, 37–41
aberrant, **58**:63
applications to clinical medicine,
40:264–270
regulation, **54**:375
- superfamily, **44**:1–3
evolution
diversity, **44**:44–46
DNA, **44**:46, 47
homology unit, **44**:43, 44
implications, **44**:47–49
homology unit, **44**:3–9
MHC complex, **44**:24–29
nonimmune receptor members,
44:29, 30
growth factor, **44**:39
Ig-binding molecules, **44**:38, 39
kinase, **44**:39
 β 2-microglobulin, **44**:30, 31
miscellaneous members, **44**:39–41
nervous system molecules, **44**:35–38
T cell-associated molecules, **44**:31–36
uncertain members, **44**:41, 42
- receptors, **44**:9–12
diversification strategies, **44**:20–24
organization, **44**:12
somatic diversification, **44**:12–20
- T cell subsets and, **41**:40, 41, 46, 52
- Immunoglobulin receptors, **52**:378–396
B cell, signal transduction through,
downregulation, **54**:374
B cell differentiation and expression,
54:338–343
B cell-specific proteins in IgR complex
identified by molecular cDNA cloning,
54:348–357
Fc α , **60**:182
Fc γ , **60**:179–181
Fc ϵ , **60**:181
functions, **54**:345–348
activation of B cells for proliferation and
differentiation, **54**:345–346
cytoskeletal interactions, **54**:346
endocytosis and processing of membrane
IgR/antigen complex, **54**:348
transportation of assembled IgR
structure retaining functions,
54:347
- future perspectives, **54**:379–380
- high affinity receptor for IgE/Fc ϵ RI,
52:378–381
assembly, **52**:381–382
cell surface structures functionally
associated with Fc ϵ RI molecules,
52:391–396
- cell surface glycolipids, **52**:393–394
Fc γ receptors, **52**:394–396
G63, **52**:394
mast cell chromolyn binding sites,
52:392–393
ME491 (CD63), **52**:393
- chromosomal location, genomic
organizations, and mRNA species,
52:382–383
- control of synthesis and expression of
dependence on IgE, **52**:385
Fc ϵ RI molecules, **52**:384–386
half-life, internalization, degradation
and reexpression, **52**:385–386
synthesis during cell cycle,
52:385
- synthesis during differentiation of
basophils/mast cells from
hemopoietic precursor cells,
52:384–385
- functional characterization of receptor,
52:387–391
binding to IgE, **52**:387–388

- Immunoglobulin receptors (*continued*)
 effector mechanisms and functional
 changes of mast cells/basophils
 following activation through IgE
 binding sites, **52:391**
 signal transduction and distal events,
52:388–391
 numbers and binding constants on
 basophils/mast cells, **52:386–387**
 sequence and structural homologies,
52:383–384
 conversation during evolution, **52:383**
 molecules related to Fc ϵ RI α chain,
52:383
 molecules related to Fc ϵ RI β chain,
52:383–384
 molecules related to Fc ϵ RI γ chain,
52:384
 topology of, **52:378**
 α chain, **52:379**
 β chain, **52:379–380**
 γ dimer, **52:380–381**
 nomenclature, **52:378**
- IgM-RF, **64:288**
 isotypes of IgG receptor, **54:343–344**
 membrane, **63:43**
 membrane and secreted, **54:342–343**
 membrane IgM receptor induction by Ig
 gene transfection, **54:357–359**
 polymeric, **65:249**
 regulation by interaction of CD44
 extracellular domain, **54:315–317**
 signal transduction and pathway through
 IgR transmembrane signaling,
54:359–378
- T cell subsets and, **41:77**
- Immunointervention
 in autoimmune diseases, **66:91–93**
 in sepsis, **66:150–151**
- Immunological antigen receptors, **43:133–136**
- Immunological cell, neuropeptide mediation,
48:176, 177
- Immunological factor
 low-molecular-weight mediators, **48:175**
 neural effects, **48:173–174**
- Immunological mediator
 astrocyte, **48:168–169**
 glioblastoma cell, **48:168–169**
 microglial cell, immune synthetic
 capabilities, **48:168**
 neural sources, **48:168–169**
- Immunological memory
 and autoimmunity, **53:251–252**
bcl-2 oncogene and, **53:241**
 and B cell immune system, germinal
 centers role, **53:231–233**
 Fas antigen/APO-1 and, **53:241**
 generation, theories of, **53:219–220**
 life span of, **53:218–219**
 lymphocyte migration and, **53:242**
 maintenance
 by constant antigenic stimulation,
53:221–222
 by long- and short-lived cells,
53:223–224
 by long-lived clonally expanded cells,
53:220–221
 and self-tolerance, **53:249–251**
 and T cell immune system
 differentiation after antigenic stimulation,
53:229–230
 effector T cell function, **53:227–228**
 memory T cell functions, **53:227–228**
 naive T cell function, **53:227–228**
 TCR and signal transduction,
53:228–229
- Immunological privilege
 anterior chamber, **48:203–212**
 corneal allograft, **48:192–203**
 haptens model, **48:211–212**
- Immunological systems, programmed cell
 death in, **50:63–70**
 in B cells, **50:63–66**
 clonal abortion, **50:64**
 faulty recombination, **50:63–64**
 growth arrest of WE III-231 B cell line,
50:64–65
 somatic mutation and terminal
 differentiation, **50:65–66**
- interphase death of T and B cells and
 nonspecific damage, **50:69–70**
 splenic B cells, **50:65**
 in T cells, **50:66–69**
 CTL targets, death, **50:69**
 deprived of growth factors, **50:68–69**
 hybridomas, activation-induced death,
50:67
 non-selected thymocyte, death, **50:**
67–68
 thymus, negative selection, **50:66**

- Immunomodulation, experimental
autoimmune encephalomyelitis,
48:166–167
- Immunopathology, IFN- γ and, **62**:96–104,
106–107
- Immunopoiesis, T cell-dependent B cell,
antigen-induced, in secondary lymphoid
organs, schematic representation,
52:222–223
- Immunopotentiators, and autoimmune
diabetes, **51**:306
- Immunoprecipitation
antinuclear antibodies and, **44**:111, 120,
129
autoimmune response, **44**:135, 136
scleroderma, **44**:123, 124
Sjögren's syndrome, **44**:113, 115
SLE, **44**:104, 105
B cell formation and, **41**:221
CD1 proteins, **59**:45–46
CD4 and CD8 molecules and, **44**:271, 285
IL-1 and, **44**:163
T cell activation and, **41**:19
T cell development and, **44**:218, 233, 243
- Immunoprophylaxis, role of MAP
immunogens, **60**:105–106
- Immunoproteasomes, **64**:16, 17
- Immunoregulation
antiidiotypic, B cell repertoire expression
and, **42**:35, 82
complement receptor 1 and, **46**:200, 201
complement receptor 2 and, **46**:205–210
immunosenescence and, **46**:247
NO role, **60**:348–349
- Immunoregulation disorders, IBD and,
42:286
- Immunosenescence, **46**:221, 253
lymphocyte activation, **46**:221, 222,
234–236
cell cycle, **46**:222, 223
chromatin, **46**:234
early events, **46**:223–227
housekeeping, **46**:234
protein, **46**:227–230
secondary signal, **46**:230–234
lymphocyte subsets, **46**:240
human, **46**:238, 239
murine, **46**:236–238
marker density, **46**:239
mucosal immunity, **46**:251–253
- regulatory changes, **46**:247
idiotypic, **46**:247–249
suppressor cells, **46**:249–251
- stem cells, **46**:246, 247
lymphoid, **46**:242–244
pluripotent, **46**:240, 241
thymus, **46**:244–246
- Immunostimulation, by apoptosis-inhibitory
drugs, **58**:278
- Immunosuppressants, and autoimmune
diabetes, **51**:307
- Immunosuppression
aPL antibodies and, **49**:223, 258
hybrid resistance and, **41**:340, 353, 354,
386–388
IFN- γ as mediator, **62**:89–90, 105–107
virus-induced, *see* Virus-induced
immunosuppression
- Immunosuppressive drug, corneal
transplantation, **48**:193
- Immunosuppressive factors, tumor
immunogenicity affected by, **57**:307–309
- Immunosuppressive molecules, *see*
Antiinflammatory and
immunosuppressive molecules
- Immunosuppressive therapy, in PNH, **60**:91
- Immunotherapy
active, **62**:219
adoptive, **58**:437–438
for EAMG, **42**:263–269
HIV, **65**:317–320
for MG, **42**:235, 269–272
NF- κ B inhibition, **58**:19–20
pulmonary inflammation, **62**:259, 285–290
tumors, **62**:218–219
- Immunotoxins, MG and, **42**:264–265
- Importin- α family, **64**:48
- Incomplete Freund's adjuvant, autoimmune
thyroiditis and, **46**:277
- Indirect immune precipitation assay, for MG,
42:254–255
- Indoleamine-2,3-dioxygenase, **62**:71
- Indomethacin, **62**:184, 186, 201
IBD and, **42**:313
Ig secretion and, **40**:103, 106
mode of action, **64**:172
- Inducing agents, murine lupus models and,
46:78
- Induction, mechanism of action of death
genes, **50**:71–73

- Infection, *see also* Viral infection
 cachectin and, **42**:213, 219
 gut microflora and, **40**:208–209
 IBD and, **42**:319
 IgG response, **42**:305
 IFN- γ and, **62**:94–96
 and IL-1ra administration *in vivo*,
54:204–209
 in animal models of sepsis, **54**:204–207
 in human infections and sepsis,
54:207–209
 IL-5 role
 helminth, **57**:176–177
 parasite, **57**:172–173
 immunity, role of IELs, **58**:326–329
 lung, immunity, **59**:394–412
 MHC class I-deficient mouse responses
 bacterial infections, **55**:412–413
 filarial parasites, human, **55**:415
 Leishmania major, **55**:414–415
 pathogens, **55**:404–415
 Trypanosoma cruzi, **55**:414–415
 viral infections, **55**:406–412
 models, NO role, **60**:341–343
 retroviral, as substrate introduction method
 in V(D)J joining, **56**:50–51
 rheumatoid arthritis, **64**:289–290
 susceptibility, cA2 therapy, **64**:325–329
 X-linked agammaglobulinemia, **59**:
 135–136, 143–146, 150
 Infections, mycoplasmic, X-linked
 agammaglobulinemia, **59**:146
 Infectious disease
 aPL antibodies and, **49**:193, 210–212, 221,
 257
 protective immune responses to, role of
 multiple antigen peptides, **60**:105–141
 role of multiple antigen peptides,
60:105–141
 Infectious mononucleosis
 acute, immunoregulatory cell functions in,
37:115–122
 associated suppressor T cells activity,
 reversal by aD D-mannose and
 saccharides, **37**:138–142
 Epstein-Barr virus and, **37**:112–115
 NK cells and, **47**:224
 Infectivity, SIVmac and SIVsm molecular
 clones, **52**:448
- Inflammation
 adoptive T cell therapy of tumors and,
49:290–291, 311, 328
 allergic
 chronic, **60**:164–165
 and eosinophils, **60**:217–223
 expression of selectins and integrins,
60:169–173
 generation of SRS-A leukotrienes,
60:190
 CD23 antigen and, **49**:161
 chemokines and, **66**:135
 clinical stages, **60**:325–326
 complement receptor 2 and, **46**:205
 cytokines, **63**:310
 eosinophilic, **60**:159–162
 helminth-induced, **60**:214
 helper T cell cytokines and, **46**:115–117
 IFN- γ and, **62**:103–104, 105, 281, 288–289
 IL-5-associated, in asthmatic patients,
57:174–176
 IL-6 and, **66**:129
 induced carcinogenesis, **60**:345–346
 intrinsic coagulation/kinin-forming cascade
 and, **66**:225–261
 leukocyte integrins and, **46**:149–151, 160,
 161, 177
 animal models, **46**:165
 expression, **46**:166–169
 function, **46**:161–165
 ligand molecules, **46**:170
 mAbs, **46**:165, 166
 leukocyte recruitment during, **58**:394–395
 LIF role, **53**:37–38
 lymphocyte homing, **64**:185–187, 188–193
 lymphocyte homing and
 high endothelial venules, **44**:315, 317
 molecules, **44**:344, 349, 352
 regional specificity, **44**:338–341
 mediators of, **42**:286, *see also* Cachectin
 in IBD, **42**:310–315
 MIF, **66**:206–207
 and NF- κ B, **58**:9–14
 nonspecific, IFN- γ and, **62**:103–104
 PGHS-2, **62**:201
 pulmonary, *see* Pulmonary inflammation
 role of NO, **60**:323–354
 sepsis
 contact cascade and, **66**:235
 cytokines and, **66**:103–150
 MIF and, **66**:210–212

- surface antigens of human leukocytes and, **49:87**, 111
and T helper cell activation, **58:106**
- Inflammatory arthritis, IL-1ra in, **54:209–211**
animal models, **54:210**
patients with, **54:210–211**
in vitro studies, **54:209–210**
- Inflammatory bowel disease, *see also* Crohn's disease; Ulcerative colitis
antibody secretion in, **42:301–307**
anti-TNF- α therapy, **64:307–310**
cell-mediated cytotoxicity in, **42:294–297**
clinical characteristics of, **42:285**
complement pathway products in, **42:308–310**
cytotoxicity as cause of, **42:295–297**
etiology of, **42:285**
general considerations in, **42:285–286**, 287
genetic defect in, **42:319**
genetic markers in, **42:288**
granulocyte function in, **42:307–308**
household contacts and, **42:291**
IL-1ra in, **54:214–215**
immunopathology of, **42:297–298**
immunoregulatory alterations in, **42:299–301**
inflammatory cells in lesions of, **42:297**
lipid extract of mucosa in, **42:315**
mediators of inflammation in, **42:310–315**
pathogenesis of, **42:319–322**
peripheral blood lymphocyte function in, **42:288–290**
pitfalls in study of, **42:286**
sequence of immunologic events in, **42:287**
serum antibodies
 DR antigens, **42:292–294**
 Escherichia coli, **42:291–292**
 lymphocytotoxic antibodies, **42:291–294**
therapy for, **42:316–319**
- Inflammatory conditions, complement components and, **38:233**
- Inflammatory demyelination, experimental autoimmune encephalomyelitis, **48:166–167**
- Inflammatory diseases
and IL-6, **54:39–43**
MIF, **66:210**
 acute respiratory distress syndrome, **66:214–215**
- arthritis, **66:213–214**
delayed-type hypersensitivity, **66:212**
glomerulonephritis, **66:212–213**
malaria, **66:215–216**
septic shock, **66:210–212**
models, **60:343–345**
NF- κ B and, **65:117–118**
- Inflammatory joint diseases, cachectin and, **42:225**
- Inflammatory mediators, **66:101**
- Inflammatory reactions, **65:11**, 122–123
- Inflammatory response
autoreactive T cells and, **45:428–430**, 433
B cell formation and, **41:208**, 237
lymphoid cells and, **41:354**
therapy, NF- κ B inhibition, **58:19–20**
- Influenza
sporozoite malaria vaccine and, **45:307**
synthetic T and B cell sites and
 antigens, **45:234**
 vaccine, **45:253–255**
 viral antigens, **45:216–219**
- T cell receptor and
 antigen processing, **45:119**, 122, 125, 126
 epitopes, **45:143**
 homogeneity, **45:128**
 peptides, **45:133**
 structure-function relationships, **45:135**, 141
- virus-induced immunosuppression and, **45:356**
- Influenza virus
cachectin and, **42:220**
CTL and, **41:136**, 168
 amino acid, **41:165**
 exon shuffling, **41:144**, 146
 HLA class I antigens, **41:151**, 152
 mAbs, **41:151**, 166, 167
cytotoxic T cells and, **38:165**, 172; **65:278**
 type A, **38:165**
hybrid resistance and, **41:375**
and memory $\gamma\delta$ T cells, **58:328–329**
pneumonia, **59:408–409**
responses in MHC class I-deficient mouse, **55:406–407**
- Inheritance
Mendelian, maternally transmitted antigen and, **38:338**
- RA susceptibility, **56:441–446**

- Inhibition, *see also* Allogeneic inhibition
 adoptive T cell therapy of tumors and, **49**:285, 320, 327
 mechanisms, **49**:307, 313
 principles, **49**:287–289
- antigen-presenting cells and, **47**:69, 77
- aPL antibodies and, **49**:193, 259
 clinical aspects, **49**:200–202
 history, **49**:198–199
 pathogenic potential, **49**:253–254, 256–257
 specificity, **49**:236, 239, 241–244, 247, 252
- autoimmune demyelinating disease and, **49**:358, 370–371
- B cell formation and, **41**:236
 B cell precursors, **41**:91, 92
 bone marrow cultures, **41**:216
 inducible cell line, **41**:222, 223
- CD5 B cell and, **47**:161
- CD23 antigen and
 biochemical structure, **49**:156–157
 biological activity, **49**:168–170, 172–173
 cellular expression, **49**:152, 154
 expression regulation, **49**:160–161
 $\text{Fc}\epsilon\text{RII}$, **49**:164
- CTL and, **41**:168, 169
 amino acid, **41**:162
 carbohydrate moieties, **41**:152–154
 exon shuffling, **41**:139
 HLA class I antigens, **41**:149–151
 mAbs, **41**:157, 158
- cytotoxicity and
 cell-mediated killing, **41**:295, 296
 cytolysis, **41**:276, 281, 282
 cytolytic proteins, **41**:317
 membrane attack complex of complement, **41**:309
- HIV infection and
 etiological agent, **47**:379
 immune response, **47**:406, 408
 immunopathogenic mechanism, **47**:386, 390–392, 394, 395
- hybrid resistance and
 antibodies, **41**:378
 antigen expression, **41**:397, 402, 404, 409
 bone marrow cells, **41**:337–340, 347, 348
 effector mechanisms, **41**:370
 leukemia/lymphoma cells, **41**:358, 364, 367
- lymphoid cells, **41**:351, 352, 354, 356
 macrophages, **41**:371, 372
 marrow engraftment, **41**:387
 marrow microenvironment, **41**:390–392
 NK cells, **41**:374–376
 syngeneic stem cells, **41**:388, 390
- T cells, **41**:381, 382
in vitro assays, **41**:394–396
- IgE, **43**:281–282
- IgE biosynthesis and
 antibody response, **47**:7, 9, 33, 39
 binding factors, **47**:12, 19, 23–25
- Ig heavy-chain variable region genes and, **49**:46
- NK cells and, **47**:298
 adaptive immunity, **47**:291–294
 antimicrobial activity, **47**:284, 287–290
 CNS, **47**:268, 269
 cytotoxicity, **47**:249, 253, 254, 257, 259
 effector mechanisms, **47**:237, 238, 240, 242, 243, 245–247
- hematopoiesis, **47**:273, 274, 276–280
 morphology, **47**:218
- SCID and, **49**:383
- spontaneous autoimmune thyroiditis and, **47**:477
- surface antigens of human leukocytes and, **49**:89, 92–93, 111–112, 114
- T cell activation and, **41**:9, 22, 26
- T cell subsets and
 H-2 alloantigen recognition, **41**:79, 82, 85, 86, 90, 91
 H-2 molecules in thymus, **41**:109, 110
 H-2-restricted antigen recognition, **41**:60, 61, 74, 76
- Inhibitors, use to block *in vitro* tolerance, **54**:415–418
- Inhibitory proteins, $\text{I}\kappa\text{B}$, **63**:157–159
 description, **58**:2–4
 phosphorylation by $\text{TNF}\alpha$, **58**:8
 rapid degradation and release of $\text{NF}-\kappa\text{B}$, **58**:4–5
- Injury
 acute states, therapeutic regulation of complement system in, *see* Complement
- thermal, therapeutic regulation of complement system in, **56**:280–281
- vascular endothelial, by TNF , **56**:351

- Inner core matrix, branched, in multiple antigen peptides, **60**:106–108
- myo*-Inositol, structure, **48**:255
- Inositol 1-monophosphate, phosphoinositide-specific phospholipase, **48**:253
- Insitol phosphates
in cells undergoing PI hydrolysis, **43**:292
isomers, dephosphorylation pathways, **48**:255
- Inositol triphosphate, T cell activation and, **41**:15, 16, 21, 24, 26, 30
- Inositol 1,4,5-trisphosphate, **43**:295; **48**:228
 Ca^{2+} release by neutrophils and, **39**:124–127
PIP₂ hydrolysis in neutrophils and, **39**:124
- Insertions, endogenous retroviral, inseparable from Vbse, **50**:38–39
- Insulin
autoimmune thyroiditis and, **46**:269
autoreactive T cells and, **45**:420
B cell formation and, **41**:197
bovine
 B chain, processing by B cells, **39**:87
 intact, defective processing by B cells, **39**:87
CD4 molecules and, **44**:293
gene, transcriptional factor, **48**:142
hexamers, **43**:103–104
hybrid resistance and, **41**:402
IL-1 and, **44**:169, 184, 188
immune response genes and, **38**:32
 I region mutations, **38**:86–89
 MHC restriction, **38**:66, 84
 T cell suppression, **38**:142–144
 tolerance, **38**:125, 126
murine lupus models and, **46**:79
spontaneous autoimmune thyroiditis and, **47**:440
synthetic T and B cell sites and, **45**:202, 209, 214, 237
T cell receptor and, **45**:117, 124, 131
- Insulin autoimmune syndrome, **66**:88
methimazole-induced, HLA class II peptide binding and, **66**:87–88
- Insulin-dependent diabetes mellitus, **51**:285–286
aberrant HLA class II expression, **48**:141–142
Asp⁵⁷, **48**:135–137
B8, **48**:137
- B15, **48**:137
dominant resistance, **48**:143–144
- DQ β chain, **48**:134–136
 position 57 aspartic acid, **48**:135–136, 137
- DR2, **48**:136, 143
AZH, **48**:136
- DRw9, **48**:137
- DRw15, **48**:143
- DRw16, **48**:143
- HLA factor, **48**:133–144
suppression phenomenon, **48**:143–144
- HLA class II peptide binding and, **66**:91
TAP and, **65**:92–93
- Insulin-like growth factor-binding proteins, **63**:419
- Insulin-like growth factor I, **63**:379–380
expression, **63**:389–393
immune system, **63**:377–378
leukemia, **63**:427
lymphohemopoietic system, **63**:389–393, 431–432
regulation, **63**:389–393
structure and physiology, **63**:388–389, 401
- Insulin-like growth factor I receptor, **63**:401, 427
- Insulin-like growth factor II, **63**:417–419
- Insulin-like growth factor II receptor, **63**:401, 418
- Insulin-like growth factor receptors, **63**:401–404
- Insulin promoter gene, **48**:141–142
- Insulin receptor, **63**:146, 168
surface antigens of human leukocytes and, **49**:100, 113
- Insulin receptor substrate-1, phosphorylation, **60**:15–16
- Insulitis
autoimmune, **62**:100–101
and diabetes, development of, **51**:292
and overt diabetes, T cell subsets involved in, **51**:296–298
- Integral membrane proteins
leukocyte integrins and, **46**:154, 156
types, **52**:144; 145
- Integrases, HIV, **64**:57

- Integrins, **52**:166, 358–361; **59**:383, *see also*
- Leukocyte integrins
 - $\alpha 4\beta 1$, **58**:371; **65**:358, 368
 - conformational effects, **64**:169
 - lymphocyte homing, **64**:146, 156
 - $\alpha 4\beta 7$, **58**:371; **64**:154, 173
 - lymphocyte–HEV interactions and, **65**:352, 358–359, 368
 - lymphocyte homing, **64**:146, 156, 192
 - $\alpha 6\beta 1$, **58**:371
 - $\alpha 6\beta 7$, lymphocyte homing and, **64**:157
 - $\alpha E\beta 7$, **58**:372
 - defined by monoclonal antibody HML-1, **58**:317–318
 - expression on intraepithelial lymphocytes, **58**:392
 - $\alpha v\beta 3$, lymphocyte homing and, **64**:157
 - $\beta 1$, **52**:358–359; **58**:370
 - $\beta 1\alpha 4$, **62**:258
 - $\beta 2$, **52**:359–361; **58**:370; **62**:279
 - and degranulation, **60**:207
 - $\beta 3$, **52**:361
 - activation, chemotactic molecules in, **58**:378–381
 - antigen-presenting cells and, **47**:61, 97
 - CD23 antigen and, **49**:163
 - characterization, **58**:362–370
 - conformational effects, **64**:169
 - expression in allergic inflammation, **60**:172–173
 - expression on human mast cells and
 - basophils, assessment by mAbs and indirect immunofluorescence, **52**:359
 - family of adhesion receptors and ligands, **58**:366–367
 - interaction with eosinophils, **60**:169–172
 - lymphocyte–HEV interactions and, **65**:352, 358–359, 368–369, 370
 - lymphocyte homing, **64**:154–157
 - properties, **64**:146
 - subunits, biochemical characterization, **58**:364–365
 - surface antigens of human leukocytes and, **49**:83–86, 92
- Intercellular adhesion molecule, **59**:383, 414
- Intercellular adhesion molecule-1, **52**:361–363; **59**:377–378; **62**:77–78; **64**:147, 154
- adhesion molecule, **62**:77–78
- as anti-adhesive, **64**:171
- CD23 antigen and, **49**:161
- central nervous system cells, **62**:91
- downregulation, **64**:172
- expression, **64**:179
- expression in allergic inflammation, **60**:172–173
- gene, NF- κ B and, **65**:119
- humoral immune response and, **45**:43
- IFN- γ , **62**:92
- IL-1 and, **44**:166
- interaction with LFA-1 in negative selection, **58**:172
- leukocyte integrins and, **46**:149, 150, 176, 177
- inflammation, **46**:163, 166, 167
- ligand molecules, **46**:169–172
- structure, **46**:158
- lymphocyte homing and, **44**:350; **64**:147, 157
- mAbs, **60**:168–170
- Mac-1 receptor, **58**:370
- modifications, **64**:168
- multiple myeloma, **64**:246, 247
- multiple myeloma and, **64**:246, 247
- properties, **64**:147
- pulmonary inflammation, **62**:267, 272, 276, 277, 281, 288
- recruitment of leukocytes to inflammation site, **58**:395
- rheumatoid arthritis, cell trafficking, **64**:317–319
- rolling cells containing, **58**:381–382
- synergistic activation of, **65**:21
- T cell activation, **63**:44
- upregulation, **64**:165, 166
- Intercellular adhesion molecule-2 expression, **64**:179
- and ICAM-1, affinity for LFA-1, **58**:373–374
- lymphocyte homing and, **64**:157, 158
- properties, **64**:147
- Intercellular adhesion molecule-3 five-Ig domain molecule, **58**:373–374
- lymphocyte homing and, **64**:158
- Interdigitating cell antigen, **59**:379
- Interdigitating cells
- antigen-presenting cells and, **47**:50–52
 - hybrid resistance and, **41**:356, 357
 - within PALS, **60**:269

- Interdigitating dendritic cells, **51**:245–246
 staining of, **51**:248–249, 250–251, 259
- Interdigititation, *J558 V_H* subfamily, **42**:122
- Interface adaptor hypothesis, **43**:125–128
- Interferon, *see also* specific interferon
 adoptive T cell therapy of tumors and, **49**:289–290, 307–308, 311, 316
 autoimmune demyelinating disease and, **49**:361
 autoimmune thyroiditis and, **46**:265
 antigens, **46**:272
 cellular immune responses, **46**:289, 293–298
 B cell formation and, **41**:228, 234, 265
 cachectin and, **42**:221
 CD23 antigen and, **49**:152, 154, 158, 161, 174–175
 helper T cell cytokines and, **46**:112
 cross-regulation, **46**:131
 functions, **46**:114, 117, 120–122
 immune responses, **46**:133, 137, 138
 precursors of differentiation states, **46**:122, 123, 125
 HIV infection and, **47**:394, 402
 hybrid resistance and, **41**:368, 377
 antigen expression, **41**:405, 406
 bone marrow cells, **41**:345
 macrophages, **41**:371, 372
 marrow microenvironment, **41**:392
 syngeneic stem cells, **41**:388–390
- IBD and, **42**:295
- IL-1 and, **44**:189, 195
 gene expression, **44**:161
 immunocompetent cells, **44**:178
 structure, **44**:157
 systemic effects, **44**:171
- immunosenescence and, **46**:231, 232, 237
- inducible genes, transcriptional regulation and role of IRF-1 and IRF-2, **52**:272–275
- leukocyte integrins and, **46**:168, 170
- lymphocyte homing and, **44**:321
- NK cells and, **42**:183
 adaptive immunity, **47**:291, 293–295
 alterations, **47**:300, 301, 303
 antimicrobial activity, **47**:283–290
 antitumor activity, **47**:295, 298
 cell-mediated cytotoxicity, **47**:195, 196
 CNS, **47**:269
- congenital defects, **47**:225, 226
 cytotoxicity, **47**:250, 251, 254–260, 262
 differentiation, **47**:229, 233, 234
 effector mechanisms, **47**:235, 237, 238, 241–244
 genetic control, **47**:222–224
 hematopoiesis, **47**:274, 277–280, 283
 identification, **47**:198
 lymphokines, **47**:264–266
 reproduction, **47**:271
 surface phenotype, **47**:201
 tissue distribution, **47**:219
- SCID and, **49**:385–386
- spontaneous autoimmune thyroiditis and, **47**:493
 altered thyroid function, **47**:461–463
 disturbed immunoregulation, **47**:472, 477
- surface antigens of human leukocytes and, **49**:99
- T cell development and, **44**:219
- T cell receptor and, **45**:117
- type I, *see* Interferon- α ; Interferon- β
 type II, *see* Interferon- γ
 virus-induced immunosuppression and, **45**:343, 361
- Interferon- α , **48**:168, 169; **52**:264–275; **61**:353; **62**:61, 84–85; **63**:275
 effect on L-selectin cell surface density, **58**:353–354
 IgE biosynthesis and, **47**:9
 Jak kinases, **63**:140
 multiple myeloma, **64**:258–259
 regulators of complement activation and, **45**:397
 soluble form, **63**:276, 307
 sporozoite malaria vaccine and, **45**:302
 TH responses, **61**:353
 virus-induced immunosuppression and, **45**:339, 344
- Interferon- α/β
 genes
 regulatory DNA sequences in, **52**:265–267
 transcriptional regulation of IFN- α
 inducible genes and role of IRF-1 and IRF-2, **52**:272–275

- Interferon- α/β (continued)**
- transcription factors binding to regulatory elements of: interferon regulatory factors 1 and 2, **52**:267–271
 - NF- κ B, **52**:271–272
 - other factors, **52**:272
 - induction of ISGF3, **60**:17
 - role in cytokine signaling, **60**:7–9
- Interferon- β , 48**:168, 169, **52**:264–275; **62**:61, 84–85, **63**:275
- regulators of complement activation and, **45**:397
 - released by macrophages, **60**:334–335
 - soluble form, **63**:276, 307
 - sporozoite malaria vaccine and, **45**:302
- Interferon- $\beta/26$ -kDa, 54**:3
- Interferon- γ , 48**:70, 71, 78, 132, 162; **52**:264–275; **62**:61–62, 105–107; **63**:275; **64**:106, **65**:282; **66**:145
- antagonists, **62**:82–83, 105
 - antigen-presenting cells and, **47**:59, 83, 97
 - and autoimmunity, **50**:213
 - autoreactive T cells and, **45**:427, 430
 - B7 expression, **62**:136
 - B cell formation and, **41**:197, 222, 230
 - B cell proliferation, **40**:37–38, 42–44; **63**:55
 - B cells, **63**:278
 - biochemistry
 - cellular, **62**:71–76
 - signal transduction, **62**:67–71
 - biological effects
 - adipocytes and, **62**:91
 - antibody formation, **62**:87–89
 - antigen presentation, **62**:80
 - antiproliferative effect, **62**:76–77
 - central nervous system cells and, **62**:91–92
 - connective tissue and, **62**:90–91
 - endocrine cells and, **62**:93–94
 - endothelial cells, **62**:81
 - epithelial barriers and, **62**:90
 - in hematopoiesis, **62**:92–93
 - immune suppression mediator, **62**:89–90, 105
 - lymphocytes and, **62**:86–87
 - membrane protein expression, **62**:77–78
 - mononuclear phagocytes, **62**:78–80
 - natural IFN- γ antagonists, **62**:82–86
 - synergy between IFN- γ and TNF, **62**:81–82
 - CD1 expression, **59**:65
 - CD28 costimulation, **62**:139
 - class switching, **61**:98
 - cytotoxicity and, **41**:272, 284
 - factor H, **61**:232
 - function, **66**:145
 - genes, transcriptional regulation, **52**:272–275
 - germline transcripts, **61**:86, 88–90
 - human T cell antigen and, **41**:6, 11, 27–29
 - humoral immune response and, **45**:61, 62, 68, 70–72, 76
 - cellular interactions *in vivo*, **45**:82, 84, 85
 - helper T cells, **45**:26–30, 33
 - hybrid resistance and, **41**:371
 - IgE biosynthesis and
 - antibody response, **47**:9, 10, 12, 31, 32
 - binding factors, **47**:22–24
 - immunopathology
 - allograft rejection, **62**:101–103
 - autoimmune disease, **62**:98–101
 - cytokine release syndrome, **62**:103–104
 - delayed-type hypersensitivity, **62**:96–98
 - granulomatous lesions, **62**:281
 - nonspecific inflammation, **62**:103–104
 - infection and cancer, **62**:94–96
 - Jak kinases, **63**:140
 - MHC class I gene expression, **61**:328–329
 - monocytes, **62**:136
 - multiple myeloma, **64**:259
 - in NO production, **60**:334–337
 - production, **62**:63–66, 105
 - proteasomes, **64**:106
 - regulation, **64**:12–32
 - pulmonary inflammation, **62**:281, 288–289
 - recent research, **62**:106–107
 - regulation of isotype switching, **54**:256–257
 - regulators of complement activation and, **45**:397
 - secreting IEL, **58**:329
 - sepsis and, **66**:104, 145–146
 - soluble form, **63**:276, 307
 - sporozoite malaria vaccine and, **45**:319
 - CS-specific T cells, **45**:307, 312
 - liver stages, **45**:301–306
 - structure–function relationship, **62**:62–63
 - T cell receptor and, **45**:117, 119, 129

- T cell subsets and
 H-2 alloantigen recognition, **41**:89, 90, 94, 95
 H-2 antigen recognition, **41**:16, 64, 66, 69, 77
 H-2 molecules in thymus, **41**:106
 TH responses, **61**:350, 358
 upregulation of adhesion molecules, **64**:165
 upregulation of FcRIII, **60**:180–181
 virus-induced immunosuppression and, **45**:356, 357, 360, 361, 363, 366
- Interferon- γ receptor
 biochemistry, **62**:66–67, 105
 IFN- γ R α , **62**:66
 IFN- γ R β , **62**:66
 soluble forms, **63**:307, 309–310, 311, 314
- Interferon receptor, soluble forms, **63**:307–308
- Interferon regulatory factor, *see* IRF
- Interferon response factor-1, *see* IRF-1
- Interferons, **52**:355–356; **66**:145
- Interferon-sensitive response elements, **62**:69–70
- Interferon-stimulated response element, **61**:174
- Interferon system, **52**:264–265
- Intergenic probes, **62**:19–20
- Interleukin, **48**:70, 71
 adoptive T cell therapy of tumors and, **49**:289, 312
 and CD1 expression, **59**:64–65
 CD23 antigen and, **49**:151, 154–155, 176
 enhancement of antibody responses and, **40**:107–110
 humoral immune response and, **45**:54–62, 75–77
 neuroimmunology and, **39**:301
 surface antigens of human leukocytes and, **49**:94, 98, 101
- T cell development and
 ontogeny, **44**:219
 thymocyte subpopulation, **44**:225–233, 236, 241
- Interleukin-1, **44**:153–155; **48**:168, 169, 52:169–171; **61**:232, 353, 354; **63**:290–291
 adhesion molecules, **62**:81
 adoptive T cell therapy of tumors and, **49**:287, 289, 325–329, 334
- antigen-presenting cells and, **47**:64, 99–101, 103
 anti-IL-1, **66**:140
 autocrine or paracrine effects, **54**:34–35, 201–202
 autoimmune arthritis, **62**:99–100
 autoimmune thyroiditis, **46**:295
 autoreactive T cells and, **45**:423, 424
 B cell formation and, **41**:192, 208, 222, 236, 237
 genetically determined defects, **41**:228, 230
 soluble mediators, **41**:234, 235
- B cell responses and, **40**:13–14, 35–36
- biochemistry, **66**:105–107
- biological effects, **44**:164, 165
 catabolic effects, **44**:168
 endothelial cells, **44**:166–168
 fibroblasts, **44**:168, 169
 hepatic protein synthesis, **44**:165, 166
 cachectin and, **42**:224
 cardiomyocytes and, **66**:114
 CD5 B cell and, **47**:142, 143, 145, 146
 CD23 antigen and, **49**:169–171
 CD28 costimulation, **62**:139
 cell-mediated killing and, **41**:288
 circulating levels, **66**:104–107
 in endotoxemia, **66**:116–117
 gene expression, **44**:159–164
 gene regulation, **65**:25–26
 granulomatous lesions, **62**:281
 growth hormone secretion, **63**:385
 helper T cell cytokines and, **46**:130
 HIV infection and, **47**:394, 404
 human disease, **44**:191–195
 humoral immune response and, **45**:56, 68, 70–75, 80
 IFN- γ , **62**:80
 IgE biosynthesis and, **47**:22
 IL-6 and, **44**:189–192
 immunocompetent cells
 B cell activation, **44**:177, 178
 diacylglycerol, **44**:175, 176
 IL-2 production, **44**:174, 175
 NK cells, **44**:178
 peripheral blood T cells, **44**:177
 T cell activation, **44**:172, 173
 thymocytes, **44**:173, 174
 immunosenescence and, **46**:228

- Interleukin-1 (*continued*)**
- in induction of acute-phase proteins by liver, **58**:10–12
 - induction of NF- κ B, **58**:7–9
 - inhibition of iNOS, **60**:338
 - in iNOS expression, **60**:351–352
 - in vitro* effects, **66**:110–112
 - in vivo* effects, **66**:112–114
 - leukocyte integrins and, **46**:170
 - local effects, **44**:171, 172
 - localization, **44**:162, 163, 167
 - membrane-bound, T cell subsets and, **41**:90
 - multiple myeloma pathogenesis, **64**:258
 - neuropeptide, **48**:178
 - neutrophils and, **66**:111
 - NF- κ B-induced, **58**:10–12
 - NK cells and, **47**:233, 262, 266
 - NO production in response to, **60**:337
 - prolactin secretion, **63**:382
 - pulmonary hypersensitivity, **59**:418, 419
 - pulmonary inflammation, **62**:259, 267, 273, 276, 277, 279, 280, 288–289
 - regulation, **66**:108–110
 - regulators of complement activation and, **45**:397
 - SCID and, **49**:396
 - deficiency, **49**:384–385, 390
 - receptor deficiency, **49**:385
 - sepsis and, **66**:104, 126–127, 150
 - animal models, **66**:117–121
 - clinical, **66**:121–124
 - treatment, **66**:124–126, 134
 - signaling events, **65**:117
 - spontaneous autoimmune thyroiditis and, **47**:477, 478, 493
 - structure, **44**:155–158
 - systemic effects, **44**:169–171
 - systemic inflammatory response syndrome, **63**:310
 - T cell activation and
 - cell surface molecules, **41**:2, 8, 13, 14
 - receptor-mediated signal transduction, **41**:20, 22, 23, 26
 - T cell subsets and, **41**:63–66, 84, 85, 107, 111
 - TNF, **44**:185–189
 - TNF- α , **64**:293, 316
 - virus-induced immunosuppression and, **45**:357
- Interleukin-1 α** , **66**:106, 107, 117
- binding to type I receptors, **54**:193
 - synthesis by eosinophils, **60**:195
- Interleukin-1 β** , **66**:106–107, 116, 117, 122
- Interleukin-1 inhibitors**, IL-1ra, **54**:168–169
- Interleukin-1-like activity**, in B cells, **39**:78–79
- Interleukin-1 receptor**, **44**:178–180; **63**:275, 290–292; **66**:108
 - binding, **44**:181, 182
 - expression regulation, **44**:182
 - IL2, **44**:180, 181
 - post-receptor-binding events, **44**:184, 185
 - structure, **44**:182, 183
 - type I, **63**:276, 291–292, 312–313
 - type II, **63**:274, 276, 291–292, 310
- Interleukin-1 receptor antagonist**, **54**:168–169; **63**:277; **66**:104, 116–117, 120, 121
 - binding
 - soluble IL-1 receptors, **54**:195–197
 - type II IL-1 receptors, **54**:194–195
 - biological relevance, **54**:217–219
 - cDNA clones, purification and expression, **54**:171–176
 - chromosomal localization and gene structure, **54**:177–180
 - chromosomal location and structure, **54**:177–180
 - in diabetes mellitus, **54**:214
 - discovery, **54**:169–171
 - effects on immune system, **54**:202–204
 - function as receptor antagonist, **54**:197–200
 - identification, **54**:167–168
 - IL-6 and, **66**:132
 - in infection and sepsis *in vivo*, **54**:204–209, 215–217
 - in inflammatory arthritis, **54**:209–211
 - in inflammatory bowel disease, **54**:214–215
 - inhibitors, **54**:168–169
 - in vitro* and *in vivo* effects, **54**:200–202
 - in malignancies, **54**:213–214
 - murine, cloning and expression, **54**:175–176
 - in nervous system, **54**:211–213
 - receptor binding of IL-1ra, **54**:192–200
 - regulation of IL-1ra production, **54**:180–192
 - structural variants, **54**:176–177
 - treatment with, **66**:124–126

- Interleukin-2, **44**:189; **48**:77; **52**:171–172, 177–178, 195–196, 204–205; **59**:225, 254; **61**:147, 149, 190, 351–352; **62**:88; **63**:292; **64**:69
adoptive T cell therapy of tumors and, **49**:284, 289–291, 293, 334
expression of antitumor responses, **49**:324–329, 332
mechanisms, **49**:300, 302–305, 307, 309–310, 315–316
allograft rejection, **62**:102, 103
antigen-presenting cells and, **47**:75
cell surface, **47**:63–65
T cells, **47**:84, 86, 100
tissue distribution, **47**:53, 55
autoimmune demyelinating disease and, **49**:361
autoimmune disease, **63**:128, 132, 294, 297
autoimmune thyroiditis and, **46**:275, 289, 310
autoimmunity, **50**:171–187
abnormalities in IL-2 expression and responsiveness on circulating lymphocytes, **50**:186–187
elevated serum levels of soluble IL-2Ra in autoimmune disease, **50**:183–186
in vitro IL-2 production in autoimmune disease, **50**:172–178
autoantibodies, **50**:176
decrease of IL-2 producing cells, **50**:176
low IL-2, **50**:175–176
signal transduction, **50**:177–178
suppressor macrophages, **50**:176
suppressor T cells, **50**:176–177
in vivo IL-2 production in autoimmune disease, **50**:178–183
at site of autoimmune attack, **50**:171–172
autotolerance, **50**:165–171
effect *in vivo*, **50**:165–166
interference with clonal deletion, **50**:166–167
nonspecific killing induced by IL-2, **50**:170–171
the system, **50**:167–170
autoreactive T cells and, **45**:425
B7 expression, **62**:136
B cell activation and, **40**:36–37, 41–50
B cell formation and, **41**:193, 228, 237
B cell proliferation, **63**:54, 55
- β chain interaction, **42**:177
binding constants, kinetic and equilibrium, **42**:169–172
bioavailability, **50**:164
cachectin and, **42**:221
CD4 and CD8 molecules and, **44**:298, 299
CD4 molecular biology, **44**:291–293
CD5 B cell and, **47**:144, 145, 161
CD23 antigen and, **49**:159, 168–169, 175
CD28 costimulation, **62**:139
cell-mediated killing and, **41**:287, 288
cells producing or expressing, detection and distribution, **50**:152–154
chain specific internalization of, **42**:175–176
complement receptor 1 and, **46**:185
cytotoxic T cells and, **38**:175
dependent cells, rescue, **58**:244
dependent T cell line, **39**:41
v-abl effect, **39**:41
eosinophil granule-associated, **60**:197
expression, **63**:44
functions, **39**:1, 3
gene, transcriptional regulation, **48**:294–297
gene regulation, case study, **51**:103–105
helper T cell cytokines and, **46**:112
cross-regulation of differentiation, **46**:130, 131
functions, **46**:114, 120–122
immune responses, **46**:136, 138
precursors of differentiation states, **46**:122–126
high-affinity binding sites, **42**:174
history of, **50**:147–149
HIV and, **65**:282
HIV infection and, **47**:390, 392, 398, 399, 411
human T cell activation and, **41**:1, 6, 11–14
gene regulation, **41**:27–300
hybrid resistance and, **41**:351, 380, 382, 396
receptor-mediated signal transduction, **41**:22, 25
synergy, **41**:15, 19
T11, **41**:9, 10
humoral immune response and, **45**:57, 62, 66–71, 73, 76
cellular interactions *in vivo*, **45**:84

- Interleukin-2 (*continued*)
 helper T cells, **45**:26, 28–33
 physical interaction, **45**:44
 IBD and, **42**:296
IFN- γ , **62**:80
IgE biosynthesis and
 antibody response, **47**:10, 11, 36, 37
 binding factors, **47**:22
Ig production, **63**:60, 62
 and IL-2R, gene regulation, **50**:154–157
 and IL-2R, physiology, **50**:149–165
 compartmentalization reduces pleiotropy,
50:163–165
 IL-2 bioavailability, **50**:164
 IL-2 production, **50**:163–164
 IL-2 responsiveness, **50**:164–165
 detection and distribution of cells
 producing or expressing,
50:152–154
 effects on peripheral immune system and
 toxicity, **50**:160–163
 gene regulation, **50**:154–157
 mechanics of IL-2/IL-2 system,
50:149–152
 and thymocyte differentiations,
50:158–160
 –IL-2R system, **63**:127
 cytokine signal transduction, **63**:127–176
 signal transduction, **63**:127, 173–176
 signaling pathways, **63**:169–173
 signal molecules, **63**:167–169
 target genes, **63**:160–164
 three-channel model, **63**:171–173
 immunocompetent cells, **44**:173–176,
 178
 immunosenescence and
 lymphocyte activation, **46**:222, 226,
 228–230, 232
 lymphocyte subsets, **46**:237, 238
 mucosal immunity, **46**:252
 regulatory changes, **46**:250
 induced cytokines, role in autoimmunity,
50:210–213
IFN- γ and autoimmunity, **50**:212–213
IL-6 and autoimmunity, **50**:210–211
TNF- α and autoimmunity, **50**:211–212
 inhibition of T cell death, **58**:269
in vitro production in autoimmune disease,
50:172–178
 autoantibodies, **50**:176
 decrease of IL-2 producing cells, **50**:176
 low IL-2, **50**:175–176
 signal transduction, **50**:177–178
 suppressor macrophages, **50**:176
 suppressor T cells, **50**:176–177
in vitro production in response to T cell
 mitogens ConA or PHA, **50**:173
in vivo antitolerance effect, **50**:165
in vivo interventions in autoimmunity,
50:187–210
 cyclosporin A and autoimmunity,
50:208–210
 effects of *in vivo* applications of rIL-2,
50:189–191
 IL-2R-targeted immune intervention,
50:187–189
 induction of manifest autoimmunity in
 athymic mice by IL-2/vaccinia virus,
50:203–206
 effect of IL-2.VV on neonatally
 thymectomized mice, **50**:206–208
 effect of IL-2.VV on nu/nu mice,
50:203–205
 lack of autoimmune manifestations in
 mice transgenic for human IL-2 or
 IL-2R components, **50**:192–193
 recombinant IL-2/vaccinia virus construct
 effect on SLE of MRL/Mp-lpr/lpr
 mice, **50**:193–203
 effect of IL-2.VV on life expectancy
 and autoimmune manifestations,
50:195–196
 effect of IL-2.VV on T cell
 compartments, **50**:196–200
 mechanism of beneficial effect of IL-2
 on lymphoproliferation and
 autoimmunity, **50**:200–203
 Jak kinases, **63**:140
 lung transplant rejection, **59**:424, 426
 lymphocyte proliferation, **63**:127
 NK cells and, **42**:184; **47**:189, 292
 adaptive immunity, **47**:294, 295
 antitumor activity, **47**:295, 296, 298
 cell-mediated cytotoxicity, **47**:196
 congenital defects, **47**:226
 cytotoxicity, **47**:250, 252, 253, 257–262
 differentiation, **47**:231–234
 effector mechanisms, **47**:235, 237, 238,
 242–244, 247
 genetic control, **47**:222, 223

- hematopoiesis, **47**:280
 identification, **47**:199
lymphokines, **47**:264, 265
 malignant expansion, **47**:228
 surface phenotype, **47**:206, 207, 209
 tissue distribution, **47**:221
 nuclear responses induced by, **63**:158
 pituitary hormones, **63**:385
 production, **50**:163–164
 production by
 LBRM-33 cells, **39**:17
 T cells, **39**:1, 3, 10–13
 cyclosporin A and, **39**:14–15
 dexamethasone and, **39**:14
 mRNA and, **39**:13–14
 production by activated T cells, **58**:18–19
 prolactin secretion, **63**:382
 protein kinase C, **63**:155
 purification and properties, **39**:4–5
 Ras proteins, **63**:145
 recombinant, autoimmune manifestations
 after *in vivo* application of,
50:189–191
 recombinant vaccine virus infection,
50:205
 regulators of complement activation and,
45:394, 402, 404
 responsiveness, **50**:164–165
 SCID and, **49**:384, 386–387
 deficiency, **49**:385, 390
 lack of stem cell engraftment,
49:396
 posttransplant immunocompetence,
49:400
 receptor deficiency, **49**:385–386
 stem cell engraftment, **49**:390–391
 tolerance, **49**:399
 transplantation with fetal tissue,
49:393
 sepsis and, **66**:104
 signaling, **48**:299–304
 signal transduction, **59**:253–259, 265–266;
63:174
bcl-2, **63**:162–164
c-fos, **63**:161, 171
c-jun, **63**:160–161, 171
c-myc, **63**:161–162
 Jak/Stat pathway, **63**:171–172
Lck, **63**:171
PI3K, **63**:171
 Ras, **63**:171
Rho, **63**:171
 single gene-coded, human, **39**:5
 spontaneous autoimmune thyroiditis and,
47:492
 altered thyroid function, **47**:461
 cellular immune reactions, **47**:454–456
 disturbed immunoregulation,
47:471–474, 479
 genetics, **47**:488
 histopathology, **47**:442
 humoral immune reactions, **47**:446
 Stat proteins, **63**:141
 stimulation of NK cells, **58**:426–429
 structure, **48**:299–300
 sulfasalazine and, **42**:317
 surface antigens of human leukocytes and,
49:94, 98
 T cell subsets and
 H-2 alloantigen recognition, **41**:87,
 93–95
 H-2 molecules in thymus, **41**:98, 99, 107
 H-2-restricted antigen recognition,
41:52, 53, 56, 62–66, 72–74, 76, 77
 T cell triggering, **41**:111, 112
 Th cells, **62**:146
 virus-induced immunosuppression and
 HIV, **45**:356, 358, 360, 362, 363, 366,
 367
 measles, **45**:342, 344
 in vivo production in autoimmune disease,
50:178–183
 Interleukin-2 β , Jak kinases, **63**:139
 Interleukin-2 γ , Jak kinases, **63**:139
 Interleukin-2 receptor, **44**:180, 181;
52:342–343; **59**:226; **61**:148–153;
63:128–132
 activation, **61**:148–149
 $\alpha\beta$ heterodimer receptor structure, **42**:168
 $\alpha\beta$ interaction, **42**:176
 assays for, **42**:166–167
 binding characteristics of, **42**:168–172, 176
 cachectin and, **42**:224
 γ -chain, Jak2 associated with, **60**:12
 cell division and, **42**:167, 178
 cellular expression, **59**:234–237
 chimerism, **59**:258–259
 c-myc expression, **63**:162
 expressions, **42**:173–176
 function, **48**:300–302

- Interleukin-2 receptor (*continued*)
 gene cloning, **59**:226, 227
 and IL-2, gene regulation, **50**:154–157
 induction by IL-5, **57**:155
 membrane form, **63**:285
 membrane proximal region, **60**:14
 noncovalent interaction of, **42**:176–177
 production, **58**:19
 reconstitution, **59**:231–234
 signaling, **61**:153–191, 191
 in B cells, **55**:274–278
 G_i-phase progression, **61**:179–187
 Jaks, **61**:160–161, 173–176
 mTOR, **61**:188–190
 PI3-K pathway, **61**:167–173
 proximal signaling events
 initiation, **61**:160–161
 Jak PTK's, **61**:158–159
 Src family kinases, **61**:153–158
 syk, **61**:159–160
 ZAP-70, **61**:160
 rapamycin, **61**:180–183
 cdk activities, **61**:183–187
 FKBP12-rapamycin cycle,
 61:187–190
 Ras/MAP kinase pathway, **61**:161–167
 Stats, **61**:162, 176–179
 soluble form, **63**:276, 279, 284, 292–294,
 296–298
 structure, **42**:167–168; **55**:274; **61**:149–153
 structure and function, **59**:227–238
 X-linked severe combined
 immunodeficiency, **59**:248–253
 Interleukin-2 receptor α , **42**:167–168;
 59:227–228, 231–234; **61**:149–150, 152;
 63:128–130, 274, 275, 276, 279, 284,
 293–294
 binding characteristics and, **42**:168–172
 gene, transcriptional regulation,
 48:294–295, 297–299
 T cell development and selection,
 58:134–135
 Interleukin-2 receptor β , **59**:228–229,
 231–234, 254; **61**:150, 152–153, 155;
 63:129, 130–131, 137–138, 274, 275,
 276, 293
 chimeric molecules, **59**:258
 cytoplasmic domain, **59**:255–258
 T cell development and selection,
 58:134–135
- Interleukin-2 receptor γ , **59**:229–234, 241,
 243, 254; **63**:129, 131, 133, 274, 275
 cellular expression, **59**:237–238
 chimeric molecules, **59**:258
 cytoplasmic domain, **59**:254–255
 gene, **59**:227, 254–255
 mutations, **59**:246–253
 sharing among multiple cytokine receptors,
 59:238–246
 T cell development and selection,
 58:134–135
 X-linked severe combined
 immunodeficiency and, **59**:246–253
 Interleukin-2 receptor chain, gene, NF- κ B,
 63:160
 Interleukin-2 receptor complex
 signal transduction, **59**:253–259
 T cell development and selection,
 58:134–135
 Interleukin-2/vaccinia virus
 effect of on neonatally thymectomized
 mice, **50**:206–208
 effect of on *nulnu* mice, **50**:204–206
 induction of manifest autoimmunity in
 athymic mice by, **50**:203–208
 Interleukin-3, **48**:71, 72, 168, 169; **63**:55, 139
 B cell formation and, **41**:228–230,
 232–234, 236, 237
 CD23 antigen and, **49**:170
 eosinophil proliferation, **48**:75
 helper T cell cytokines and, **46**:116, 117
 humoral immune response and, **45**:26, 57,
 65, 73–75
 hybrid resistance and, **41**:389
 IgE biosynthesis and, **47**:10, 22
 multiple myeloma pathogenesis, **64**:256,
 258
 murine
 20 α SDH induction in T cells, **39**:2, 5,
 18–19, 23–31
 activities *in vivo*
 hematopoiesis and, **39**:30
 during immune responses, **39**:31–34
 amino acid sequence, **39**:5–7
 in bone marrow cell culture
 CSF activity, **39**:20–21
 natural cytotoxicity increase, **39**:21
 Thy-1 $^{+}$ induction, **39**:21–22, 24,
 26–31
 burst-promoting activity, **39**:20

- cDNA library for, 39:5, 7
-dependent cells
 c-myc expression, 39:40, 42
 IL-3 receptor activity, 39:38–39,
 42–43
 properties, 39:34–37
 protein kinase C redistribution, 39:39
 v-abl effect, 39:40–41
 v-myc effect, 39:40
functions, 39:2–3
growth regulation, mechanism of,
 39:37–42
 future research, 39:43–45
hematopoietic growth and, 39:19
hemopoietin-2 activity, 39:20
mast cell growth and, 39:19–20
production by
 mast cells, 39:17
 T cells, 39:2–3, 10–13
 WEHI-3 cells, 39:13, 15–17, 19
purification and properties, 39:5
single gene-coded, 39:7–8
NK cells and, 47:232, 233, 266, 277, 278
produced by eosinophils, 60:197–198
production
 by crosslinkage of Fc ϵ R on peritoneal
 mast cells, 53:9
 by factor-dependent mast cell lines,
 53:6–8
 by transformed mast cells, 53:4–6
regulation of lymphokine production by
 Fc ϵ RI $^+$ cells, 53:21–22
signaling, role of protein tyrosine kinases,
 60:5–9
Interleukin-3 receptor, 52:343–349; 63:139
expression by eosinophils, 60:183–185
murine, IL-3 interaction with, 39:38–39,
 42, 43
ATP role, 39:39
Interleukin-3 receptor α , 63:275
Interleukin-3 receptor β , 63:275
Interleukin-4, 44:172, 174, 177; 48:71, 72;
 52:172, 178–181, 196–198, 205–206;
 59:235–236, 254; 62:83–84, 85; 63:168,
 295, 299
adoptive T cell therapy of tumors and,
 49:290, 307, 327–328, 334
allergic reactions, 63:310
allograft rejection, 62:102
antibody response, 62:88
antigen-presenting cells and, 47:75, 84, 100
 cell surface, 47:59, 63, 64
 immunogenicity, 47:103
asthma, 62:283, 285
autoreactive T cells and, 45:427
B7 expression, 62:136
B cell production, 61:34
B cells, 63:278
 proliferation, 63:54, 55
biological activity, 59:238–240
CD5 B cell and, 47:120
CD23 antigen and biochemical structure,
 49:157
 biological activity, 49:169–170, 172–173
 cellular expression, 49:150–155
 cleavage regulation, 49:158
 expression, 49:159–161, 174–175
 Fc ϵ RII, 49:162, 166–168
CD28 costimulation, 62:139
CD40, 61:38
class switching, 61:98
effect on nitrite production, 60:353
enhancement of isotype switching to IgE
 and IgC, 54:248–249
eosinophil granule-associated, 60:198
expression, 63:44
functional similarities with IL 10,
 56:12–13
gene promoter, 65:29
germline transcripts, 61:86–90
granulocyte, 48:75
granulomatous lesions, 62:281, 290
helper T cell cytokines and, 46:113
 cross-regulation of differentiation,
 46:130–131
 functions, 46:114–116, 119–122
 immune responses, 46:133, 136–138
 precursors of differentiation states,
 46:122–125
humoral immune response and, 45:58,
 62–67, 69–71, 75, 76
 cellular interactions *in vivo*, 45:82–84
 helper T cells, 45:26–34, 42, 44
IFN- γ , 62:105
IgE biosynthesis and
 antibody response, 47:6–12, 31
 binding factors, 47:13, 21–23
 and IgE switching, 52:217–218
Ig production, 63:58–67, 69, 79–80

- Interleukin-4 (*continued*)
 and IL-10, in CD40 system, growth-promoting activity of, **52:202**
 immunosenescence and, **46:231, 232, 234, 237, 238**
 inhibition of iNOS, **60:338–339**
 Jak kinases, **63:140**
 lung transplant rejection, **59:424**
 macrophage, **48:75**
 molecular mechanisms of IL-4-induced switching, **54:252–256**
 NK cells and, **47:233, 262**
 pleiotropic effects of, **52:179–181**
 preferential rescue of Th2 cells, **58:244**
 production
 FceR⁺ cells in infected/immunized mice, **53:15–17**
 FceRI⁺ bone marrow cells, **53:10**
 FceRI⁺ cells, and lymphokine producing phenotypes of CD4⁺ T cells, **53:17–19**
 FceRI⁺ spleen cells, **53:9–10**
 human FceR⁺ cells, **53:14–15**
 N. brasiliensis-infected mice, **53:15–17**
 response to plate-bound IgE, **53:20–21**
 signaling mechanisms, **53:17–22**
 and T cell lymphokine-producing phenotype, **53:18**
 molecular regulation, **53:22**
 murine, infected or immunized, **53:15–17**
 peritoneal, **53:9–14**
 role in disease process, **53:17**
 signaling processing in response to FeR crosslinkage, **53:20–21**
 splenic, **53:9–14**
 by transformed mast cells, **53:4–6**
 pulmonary inflammation, **62:260, 290**
 role of Stat6 in response to, **60:21–22**
 role in T cell-dependent isotype switching, **54:249–252**
 SCID and, **49:386**
 selective induction of VCAM-1 on
 HUVEC, **60:170**
 sepsis and, **66:144–145**
 signaling molecules for, **63:167–168**
 sources and structure, **52:178–179**
 Stat protein activation, **61:176–179**
 stimulation of IgE and IgG production, **54:247–248**
 surface antigens of human leukocytes and, **49:89, 91**
 TH responses, **61:348–350, 350, 354, 358, 374**
 transcription induced by, **65:19**
 upregulation of adhesion molecules, **64:165**
 Interleukin-4 receptor, **52:349–350; 59:240–241; 63:132**
 activation, **63:168**
 production, **63:281–282, 284, 285**
 Ras proteins, **63:145**
 signaling, in B cells, **55:278–279**
 soluble form, **63:276, 281–282, 284, 294–295, 299, 311, 313**
 sources and structure, **52:178–179**
 Interleukin-4 receptor α , **63:275**
 association with 4PS, **60:16**
 Interleukin-4 receptor antagonist, **63:277**
 Interleukin-4 receptor complex, **59:238–241, 254**
 Interleukin-4 receptor F, Ras activation, **61:161**
 Interleukin-5, **48:71, 72, 78; 61:98, 352–353, 358; 63:299**
 allergic airway inflammation, **62:283–285**
 animal models of production
 airway hypersensitivity, **57:173–174**
 eosinophilia in guinea pigs, **57:173–174**
 experimental eosinophilia, **57:173**
 parasite infection, **57:172–173**
 transgenic mouse, **57:169–170**
 tumor rejection, **57:174**
 X chromosome-linked immunodeficient mouse, **57:171–172**
 antigen-presenting cells and, **47:63, 64, 103**
 B cell proliferation, **63:55**
 characteristics, **57:145–146**
 expression, **63:44**
 functional properties
 basophil regulation, **57:156**
 B cell development regulation, **57:154–155**
 eosinophil production, **57:156**
 Ig production regulation, **57:151–154**
 IL-2 receptor induction, **57:155**
 future perspectives, **57:178–179**
 and gene, molecular structure, **57:148–149**
 GM-CSF, **48:75**

- helper T cell cytokines and, **46**:113
 functions, **46**:116, 119, 120
 immune responses, **46**:133, 136, 137
 precursors of differentiation states,
46:122, 123, 125
- histological background, **57**:146–148
 in human disease
 asthma, **57**:174–176
 graft rejection, **57**:177
 helminth infections, **57**:176–177
 tumors, **57**:177–178
- humoral immune response and
 cellular interactions in vivo, **45**:82–84
 helper T cells, **45**:26, 28, 32, 33
 physical interaction, **45**:48
- IgE biosynthesis and, **47**:9, 10
- Ig production, **63**:60–62
- immunosenescence and, **46**:231, 232, 237,
 238
- inhalation at time of challenge, **60**:176
- Jak kinases, **63**:139
- mAbs, **60**:215
 mediated eosinophil signal transduction,
60:209–210
- messenger RNA expression, **57**:156–159
- molecular structure
 biological activity, **57**:149
 cDNA organization, **57**:148–149
 gene organization, **57**:148–149
 polypeptides, **57**:149–151
- mRNA, **60**:198–199, 222
- production by factordependent mast cell
 lines, **53**:6–8
- recombinant human, **60**:204–205
- Th cells, **62**:146
- Interleukin-5 receptor, **52**:350–351;
63:299–300
 expression
 aberrations in mouse, **57**:170–172
 analysis, **57**:168–169
 by eosinophils, **60**:183–185
 function, **57**:165–166
 signaling pathway, **57**:166–168
 signal transduction, **57**:146
 structure, **57**:159–165
- Interleukin-5 receptor α , **63**:275, 276
 gene structure, **57**:162–163
 soluble forms, **57**:163–164
 structure, **57**:160–164, 162–163
- Interleukin-5 receptor β , structure,
57:164–165
- Interleukin-6, **44**:153–155; **48**:71, 72, 168,
169; **52**:172, 182–184; **61**:33, 353, 354,
370; **62**:86; **63**:300; **66**:127
 and acute-phase protein synthesis in
 hepatocytes, **54**:25–27
 antibody response, **62**:88
 antigen-presenting cells and, **47**:64, 100,
 103
 anti-IL-6, **66**:132
 and autoimmune system, **52**:215
 and autoimmunity, **50**:210–211
 and B cell growth, **52**:203–204
 biochemistry, **66**:127–128
 biological function, **54**:20–33
 in blood vessels, **54**:29–30
 and bone metabolism, **54**:28
 effect on skin, **54**:28–29
 growth regulatory functions, **54**:20
 during hematopoiesis, **54**:22–25
 immune regulation, **54**:21–22
 in neuronal cells, **54**:30–31
 in placental/fetal development, **54**:32
 proinflammatory and antiinflammatory
 cytokine, **54**:27–28
 role in embryonic development,
54:31–32
- CD23 antigen and, **49**:160
- CD28 costimulation, **62**:139
- cell sources and inducers, **54**:6
 clinical applications, **54**:48–50
 common signaling pathways, **53**:44–46
 comparison to IL-1 186, **44**:189–191
 description, **54**:1–2
 as diagnostic marker, **54**:47–48
 and disease, **54**:33–47
- AIDS, **54**:37–39
 bacterial and parasite infection,
54:36–37
- B cell neoplasia, **54**:33–36
- IL-6 transgenic mouse, **54**:45–47
 inflammatory or autoimmune diseases,
54:39–43
- viral infection, **54**:37
- downregulation, **64**:315, 316
- eosinophil-derived, **60**:199
- expression regulation, **54**:8–10
- NF- κ B, **54**:10
- NF IL-6, **54**:9–10

- Interleukin-6 (*continued*)
- gene
 - NF- κ B and, **65:119**
 - regulation 23, **65:25**, 26–28
 - gene promoter, **65:26**
 - granulocyte, **48:75**
 - helper T cell cytokines and, **46:116**, 133
 - hemopoiesis, **64:321–322**
 - historical overview, **54:1–4**
 - HIV infection and, **47:403**, 404
 - human disease, **44:192**, 194
 - humoral immune response and, **45:60**, 74, 75, 82
 - Ig production, **63:64**
 - immunocompetent cells, **44:172**, 177, 178
 - immunosenescence and, **46:231**
 - inflammatory cytokine, interaction with NF- κ B, **58:12–14**
 - inhibitors, clinical applications, **54:50–51**
 - in vitro* effects, **66:128–129**
 - in vivo* effects, **66:129–130**
 - Jak kinases, **63:139**, 140
 - Jaks interdependence in response to, **60:8–9**
 - and LIF, **53:44–47**
 - multiple myeloma pathogenesis, **64:252–256**
 - pituitary hormones, **63:385**
 - plasmacytogenesis, **64:234–235**
 - pristane-induced, **64:226–227**
 - pleiotropic effects of transforming growth factor B, **52:184**
 - pleiotropic nature of, **53:44**
 - production
 - by crosslinkage of Fc ϵ R on peritoneal mast cells, **53:9**
 - by factor-dependent mast cell lines, **53:6–8**
 - prolactin secretion, **63:382**
 - pulmonary hypersensitivity, **59:418**, 419
 - pulmonary inflammation, **62:289**
 - schematic ternary structure, **54:5**
 - sepsis and, **66:104**, 127, 130–135
 - signal transduction
 - all-trans* retinoic acid, **64:261–262**
 - reshaped human PM-1, **64:261**
 - therapies using, **64:260–263**
 - structure and expression, **54:4–11**, 12
 - inducers and producers, **54:6–8**
 - regulation of IL-6 expression, **54:8–10**
 - repression of IL-6 expression, **54:10–11**
 - structure, **54:4–5**
 - transcriptional regulatory elements, **54:9**
- Interleukin-6 receptor, **54:11–19**; **61:151**;
 - 63:275**, 276, 280, 282, 300–303
 - signaling, in B cells, **55:279–280**
 - structure, **66:127**
- Interleukin-6 receptor α , **63:275**, 276, 280, 300–301
- Interleukin-6 receptor β , **63:275**, 276
- Interleukin-7, **59:235–236**, 254; **61:178**; **63:140**, 303
 - B cell development, **59:298**
 - biological activity, **59:241–243**
 - cytotoxic T cells, **65:306**
 - Jak protein activation, **61:178**
 - pre-B cells nonproliferating on stromal cells, **53:134–135**
 - pro- and pre-B cells reactive to, **53:131–136**
 - signal induction, **59:259**
 - T cell development, **59:251**
 - and TCR β regulation, **58:142**
- Interleukin-7 receptor, **59:243–244**; **63:139**, 303
 - T cell activation, **48:250–253**
 - signal transduction, **48:251–252**
- Interleukin-7 receptor α , **63:275**, 276
- Interleukin-7 receptor complex, **59:241–244**, 254
- Interleukin-8, **55:97**
 - biochemistry, **66:135–136**
 - biological effects
 - basophils, **55:116–117**
 - eosinophils, **55:117**
 - lymphocytes, **55:118–119**
 - monocytes, **55:117–118**
 - neutrophils, **55:113–115**
 - in vivo*, **55:118–119**
 - cellular sources, **55:108**
 - endothelial cells, **55:107**
 - epithelial cells, **55:107**
 - fibroblasts, **55:107**
 - keratinocytes, **55:111**
 - leukocytes, **55:105–107**
 - macrophages, **55:106**
 - mesangial cells, **55:111**
 - monocytes, **55:105–106**
 - production, regulation, **55:111–112**
 - synovial cells, **55:111**
 - tissue cells, **55:107**, 111

- chemical properties, **55:97–101**
chemotactic for primed eosinophils, **60:177–178**
gene regulation, **65:23, 28**
genes
 chromosomal localization, **55:103–104**
 expression, regulation, **55:104–105**
 organization, **55:103**
IFN- γ , **62:80**
immunoreactivity, **60:199**
inactivation, **55:102–103**
inflammatory cytokine, transcription, **58:12**
molecular structure, **55:97–101**
pathology role, **55:147–148**
 arthritis, **55:143–144, 147**
 lung disease, **55:144–147**
 skin diseases, **55:142–143, 147**
production by factor-dependent mast cell
 lines, **53:8**
proteolytic processing, **55:102–103**
pulmonary inflammation, **62:260, 270, 273,**
 274
regulation, **66:136**
and related integrins, receptors for,
 52:367–371
sepsis and, **66:104, 137–141**
signal transduction, **55:140–142**
TNF- α , **64:294, 295, 316**
- Interleukin-8 receptor, **66:136**
binding, **55:136**
biochemical studies, **55:122–125**
cloning, **55:128–134**
ELR motif, **55:136–138**
on erythrocytes, **55:128**
expression, **55:135**
genes, **55:135**
G-protein coupling, **55:138–139**
mapping, **55:135**
regulation, **55:139–140**
signal transduction, **55:141**
- Interleukin-9, **59:244, 254; 63:140, 167,**
 168
in hematopoietic system, **54:83–84**
in human and mouse, cloning and
 characterization, **54:80–82**
and human T cells, **54:85–88**
and mast cells, **54:93–95**
and mouse T cells, tumorigenesis,
 54:88–93
studies with, **54:79–80**
- Interleukin-9 receptor, **54:82–83;**
 59:244–245; 63:139
Interleukin-9 receptor complex, **59:244–245,**
 254
Interleukin-10, **52:181–182; 62:85; 63:54, 55,**
 60–67, 69, 139, 140
allograft rejection, **62:102**
antibody response, **62:88**
B cell production, **61:34**
biological effects
 B cells, **56:10–11**
 macrophages, **56:6–8**
 mast cells, **56:10**
 NK cells, **56:10**
 T cells, **56:8–10**
CD40, **61:38**
characterization, **56:1**
class switching, **61:98–99**
complementary DNA clones, **56:3–5**
discovery
 cytokine synthesis inhibitory factor,
 56:1–2
 TH1/TH2 dichotomy, **56:1–2**
functions, **56:7, 18**
gene acquisition by herpesvirus, **56:**
 11–12
gene structure, **56:4**
granulomatous lesions, **62:281**
herpesvirus acquisition of, **56:11–12**
IFN- γ , **62:105**
and IL-4, in CD40 system, growth-
 promoting activity of, **52:202**
levels, *in vivo* manipulation, **56:16–17**
multiple myeloma pathogenesis, **64:258**
PGHS-2, **62:192, 195**
physical properties, **56:3**
pleotropic effects of, **52:181–182**
as potent B cell differentiation factor,
 52:207
in pregnancy, **56:17**
production, **56:4–6**
properties, **56:3**
pulmonary inflammation, **62:259–260,**
 274–275, 279, 289
related proteins, properties, **56:3**
removal, **56:16–17**
sepsis and, **66:104, 142–145**
sources and structure of, **52:181**
structure, **56:4**
Th cell, **62:146**

- Interleukin-10 (*continued*)
 Th2 cell
 functional similarities of cytokines,
 56:12–13
 responses correlated with expression,
 56:13–15
 Th responses, **61**:351, 358
 treatment, **56**:16
 tumor immunogenicity reduced by,
 57:307–309
- Interleukin-10 receptor, characteristics, **56**:18
- Interleukin-11, **63**:139
 biological function, **54**:16–17
 multiple myeloma pathogenesis, **64**:256
 relationship to LIF, OSM, CNTF, and IL-6, **53**:47
- Interleukin-12, **61**:350–351; **63**:139, 140;
 66:146–147
 granulomatous lesions, **62**:281
 IFN- γ , **62**:105, 107
 sepsis and, **66**:104, 147
 and Stat4 phosphorylation, **60**:21
 Th cells, **62**:146
- Interleukin-13, **61**:34, 354, 374; **62**:85–86,
 260
 B cell proliferation, **63**:55
 Ig production, **63**:64, 69
 Jak kinases, **63**:139, 140
 signaling molecules for, **63**:169
 upregulation of adhesion molecules, **64**:165
- Interleukin-13 receptor, **63**:133
- Interleukin-15, **59**:245, 254; **61**:178;
 62:106–107; **63**:140, 173
- Interlenkin-15 receptor, **59**:245; **63**:139
- Interleukin-15 receptor complex, **59**:245–246,
 254
- Interleukin-16, **62**:136, 146; **65**:282
 HIV-1, **66**:281, 283
- Interleukin hybridoma plasmacytoma 1,
 64:226
- Interleukin receptor antagonist, **62**:289
- Intermitotic life span, of B and T cells,
 53:237–241
- Internalization
 CD4 molecules and, **44**:296
 evidence for, **52**:87–89
 IL-1 and, **44**:181, 184, 185
 signals, **52**:89–92
 T cell development and, **44**:226, 228, 229
- Interphotoreceptor retinoid-binding protein,
 48:215–216
- Interstitial macrophages, **59**:375
- Interstitial nephritis, induced by IL-2,
 50:200–201
- Interstitial T cell surface glycoprotein, avian,
 50:97–101
- Intestinal antigens, IBD and, **42**:290
- Intestinal mononuclear cells
 antibody secretion and, **42**:302–307
 IBD and, **42**:295–296
- Intestine
 infections and diseases, immunity, **58**:329
 inflammation, lymphocyte homing, **64**:145
 intraepithelial lymphocytes, **53**:163–167
 majority as CD8 $^{+}$, **58**:307
 lymphocyte homing and
 immune response, **44**:331, 332, 334–336
 molecules, **44**:352
 regional specificity, **44**:323, 327, 329,
 339
- int* genes, **65**:154, 155
- Intrachromosomal deletion, switch
 recombination, **61**:99
- Intracisternal A particle core protein,
 polyclonal antibody, **43**:304
- Intraepithelial lymphocytes
 CD4 and CD8 on, **58**:306–307
 effector potential, **58**:324–326
 and immune system, **58**:297–343
 immunosenescence and, **46**:237
 intestinal, differentiation, **53**:163–167
 invariant γ/δ T cells (skin, female
 reproductive tract), **53**:167–169
- origin, **58**:308–311
- role in immunity
 intestine, **58**:329
 lung, **58**:328–329
 skin, **58**:326–328
- T cell lineages, **58**:110–112
 as T cells, **58**:298–300
 as T cells bearing $\gamma\delta$ TCR, **58**:301
- Intrahepatic cholestasis, severe, **50**:190–191
- Intraocular tumor rejection
 basic patterns, **48**:215
 cytolytic T cell, **48**:215
 hemorrhagic necrosis, **48**:215
 immune regulation, **48**:212–215
 immune rejection, **48**:215

- Intrathymic clonal deletion, **51**:165–173
 cell biology of negative selection,
51:165–170
 search for mechanism, **51**:170–173
- Intrathymic defects, SCID and, **49**:383–384
- Intrathymic development, *see* T cells,
 intrathymic development
- Intrathymic induction, clonal anergy,
 examples, **50**:169
- Intrauterine influences, maternally
 transmitted antigen and, **38**:321, 322,
 338
- Intrinsic affinity, **59**:305
- Intrinsic coagulation/kinin-forming cascade,
66:240–243
 bradykinin
 formation, **66**:225, 226, 233–239
 functions, **66**:239
 receptors, **66**:239–240
 contact activation, **66**:225, 233–239
 allergic diseases, **66**:258–259
 Alzheimer's disease, **66**:260
 disease states and, hereditary
 angioedema, **66**:257–258
 endotoxic shock, **66**:260
 interaction with cells, **66**:243–248
 pancreatitis, **66**:260
 rheumatoid arthritis, **66**:260
 endothelial cells and, **66**:245
 factor XI, **66**:225, 226, 230, 237, 244
 factor XII, **66**:225, 226–229, 234–236, 244,
 246, 249–257
 high-molecular-weight kininogen, **66**:225,
 226, 230–233, 245, 247–257
 kallikrein, **66**:225, 238, 243–244
 low-molecular-weight kininogen, **66**:233
 mechanism, **66**:225–226
 platelets and, **66**:244–245
 prekallikrein, **66**:225, 226, 229–230, 236
- Intron/exon mapping, CD1 genes, **59**:7–8
- Inulin, cytotoxicity and, **41**:304, 306
- Inversion
 T cell receptor protein and, **38**:8
 and translocation, in lymphoid tumors,
 mechanism, **50**:121–122
- Iodine
 autoimmune thyroiditis and
 antigens, **46**:267, 269–272
 cellular immune responses, **46**:294
 experimental models, **46**:278
 humoral responses, **46**:303
- spontaneous autoimmune thyroiditis and
 altered thyroid function, **47**:456, 459,
 460
 genetics, **47**:487–490
- Iodo-2'-deoxyuridine, hybrid resistance and
 antigen expression, **41**:397, 404
 bone marrow cells, **41**:340, 342, 345,
 347
 effector mechanisms, **41**:370
 leukemia/lymphoma cells, **41**:364–368
 lymphoid cells, **41**:352, 354
 T cells, **41**:380
- Ion channels, T cell, **48**:285–288
 regulation, **48**:285, 286
- Ion-exchange chromatography, leukemia
 inhibitory factor, **53**:32
- Ionizing radiation
 and *scid* defect, **58**:49–51
 sensitivity of ataxia telangiectasia cells,
58:68–69
- Ionomycin, **48**:80, 90
 effects
 factor-dependent mast cell lymphokine
 production, **53**:6
 IL-3 production by mast cells, **53**:5
 immunosenescence and, **46**:228
 T cell subsets and, **41**:65, 98
- Ionotypic regulation, murine lupus models
 and, **46**:65
- IP₃, *see* Inositol 1,4,5-trisphosphate
- IP-10, **62**:76, 80
- Ipr* gene, **61**:9
- IRAK, **63**:159; **65**:117
- IRF, **62**:69–70
 model for, **52**:276
- IRF-1, **62**:69–70
 B cell genesis, **63**:235–236
 and IRF-2, **52**:267–271
 knockout mouse, **60**:335
 requirement for NOS expression,
60:17
- IRF-1, **60**:24–25
- IRF-2, **62**:69–70; **63**:218–219, 235–238
 B cell genesis, **63**:218–219, 235–238
 and IRF-1, **52**:267–271
- IRF-2, **63**:235–236
- IRF-binding sites, in IFN-inducible genes,
52:273
- IRG-47, **62**:76
- Ir* genes, *see* Immune response genes

- Irradiation
 and apoptosis, **58**:237–240
 hybrid resistance and
 antigen expression, **41**:397, 403, 404,
 408–410
 bone marrow cells, **41**:339, 344,
 346–348, 350
 effector mechanisms, **41**:369, 370
 antibodies, **41**:376, 377, 379
 marrow, **41**:384–386, 388, 390
 NK cells, **41**:373–375
 syngeneic stem cells, **41**:389, 390
 T cells, **41**:380–384
in vitro assays, **41**:394, 396
 leukemia/lymphoma cells, **41**:358, 359,
 361, 363–367
 lymphoid cells, **41**:352, 354–357
 NK cells and, **47**:291, 295
 differentiation, **47**:229–232
 effector mechanisms, **47**:260, 261
 hematopoiesis, **47**:273, 274, 276
- IRS-1, **63**:146, 168–169
 IRS-2, **63**:169
- Ischemia
 aPL antibodies and, **49**:218–219, 258
 leukocyte integrins and, **46**:165, 166, 177
 therapeutic complement inhibition using
 soluble CR1, **56**:282–284
- Ischemia/reperfusion
 model, role of NO, **60**:330–332, 346–348
 pulmonary inflammation, **62**:275–277
- ISGF3 transcription complex, induction by
 IFN- α/β , **60**:17
- Islet beta cells, **48**:140–141
 antigen presentation, **48**:142
 dysfunction, NO role, **60**:345
 lymphokine, **48**:141
 T cell cytotoxicity, **48**:142
- Islets of Langerhans, IFN- γ and, **62**:100–101
- Isoelectric focusing, complement components
 and, **38**:207, 208, 215
- Isoelectrofocusing, one-dimensional, HLA
 class II, **48**:114–116
- Isotype-mismatched molecule, **48**:125
- Isotypes
 IgG receptor, **54**:343–344
 production, secondary B cells and,
42:75–76
 response, role of T cell Fc receptors and,
40:212–216
- Isotype specificity, regulation, **61**:84–89,
 90–92
- Isotype switching, **65**:253–254
 during B cell differentiation, **60**:38–39
 B cells, human, **52**:215–218
 general considerations on, **52**:215–217
 IL-4 and IgE switching, **52**:217–218
 defect in HIGMX-1, **60**:42–48
 directed, and germline C_H transcription,
54:240–245
 dual isotype expression without
 recombination, **54**:246
 expression of downstream isotypes without
 switch recombination, **54**:245–246
 in humans, **54**:260–262
 IL-4 regulation, **54**:260–262
 TGF- β and IgA expression, **54**:262
 to IgE and IgC, IL-4 enhancing,
54:248–249
 nondeletional, **54**:245–246
 sequential, **54**:238–239
 by TGF- β , regulation, **54**:257–259
- Itk, **59**:265–266
 signal transduction, **59**:265–266
- J**
- J*558 V_H gene family, *see also* Arsonate idiotypic system, A/J mouse, *J*558 V_H gene family and
- J*774.1 cell line, specific binding of LPS,
53:277–278
- Jak*, and *JAK*, **60**:3–4
- JAK-1, **62**:68–69, 106; **63**:139, 168
- JAK-1/2, IL-2, **61**:173–174
- JAK-2, **62**:68, 69, 106; **63**:139
 activation of, **65**:20–21
- JAK-3, **63**:139
- JAK-STAT-GAS pathway, pituitary hormones, **63**:399–400
- JAK-STAT mechanism, IFN- γ binding,
62:67–71
- Jak-Stat pathway, **61**:179, 190; **63**:138–141,
 171–172, *see also* Janus kinase family
- Janus kinase family, **62**:68–69, 106; **63**:134,
 138–141
 activation
 by EGF receptor, **60**:27

- role of receptor dimerization/
oligomerization, **60**:9–11
- association with cytokine receptors,
60:11–12
- coupling of cytokine binding to tyrosine
phosphorylation, **60**:4–9
- origins and structure, **60**:2–4
- PTK's, IL-2R coupling, **61**:158–161, 190
- Stat1 as substrate, **60**:19
- Tyk2, **62**:68
- in IFN- α/β response, **60**:7–9
- Jarisch–Herrheimer reaction, release of IL-6,
66:11–34
- JC_v, B cell development, **63**:29–30
- J chain, **65**:248
- cells producing, **40**:186, 188–189
- of IgA, structure and function of,
40:158–160
- IgA polymerization and, **40**:187–188
- Jejunum, lymphocyte homing and, **44**:330,
332, 333
- Jerne hypothesis, **59**:99–100
- Jerne model, T cell repertoire and,
38:130–132
- J gene segments, Ig heavy-chain variable
region genes and, **49**:4–6, 28
- J_H gene segments, **62**:1
- Ars A antibodies, **42**:107–108
- deletion, **62**:32–33, 42
- Ig heavy-chain variable region genes and,
49:2–3, 29, 31, 33–35, 53
- in mouse and human, **62**:17–19
- organization, **62**:9–10
- Joining, V(D)J, antigen receptor genes, *see*
V(D)J joining, antigen receptor genes
- Joints
- morphology of, **64**:287–288
 - NO-induced destruction, **60**:344
 - open-and-shut, V(D)J recombination,
64:42
 - rheumatoid arthritis, **64**:287–288, 291, 315
 - joint destruction mechanism, **64**:
320–321
 - neovascularization, **64**:319–320
- jsd*, **60**:20–21
- Junction, diversity of, variable region and,
40:255–256
- jun* genes, **63**:160, 248
- Jun kinases, **63**:148
- Ets transcription factors and, **64**:81
- Jurkat leukemic T cells, **48**:182
- IL-1 and, **44**:174–176, 185
- Juvenile rheumatoid arthritis
- DR4, **48**:131
 - DR5, **48**:131
 - DRw8, **48**:131
- K**
- K562 cells, as target cells for NK activity,
42:184
- Kallikrein, **66**:225
- in allergic reactions, **66**:259
 - IgE biosynthesis and, **47**:25
 - inhibition, **66**:238
 - interactions with cells, **66**:243–244
- Kaolin clotting time, aPL antibodies and,
49:201–202, 241
- Kaposi's sarcoma, and IL-6, **54**:38–39
- κ enhancer, **43**:249–251
- κ gene, **43**:238, 240
- κ light chain, **63**:5, 10, 13, 24
- rearrangement, **61**:305
 - variability, **42**:4
- κ light chain promoter, **43**:257
- κ locus, murine, Ars A response and,
42:138
- κ promoter, **43**:257–258
- Karyotype, SCID and, **49**:388, 392
- KAVYNFATC, induced positive selection of
thymocytes, **59**:126
- Kawasaki syndrome, **54**:133–137
- analysis of T cell receptors, **54**:135
 - diagnostic criteria for, **54**:133
- KC, **62**:80
- K cell function, Crohn's disease and,
42:295
- KEKE motif, **64**:30
- Keratinocytes
- B cell formation and, **41**:234
 - expression of dendritic epidermal cell
ligand, **58**:320
 - helper T cell cytokines and, **46**:115
 - IFN- γ effect on, **62**:76
 - IL-1 and, **44**:153, 160, 177, 185
 - IL-1ra production, **54**:190–192
 - and NO production, **60**:350–351
 - as source of chemokines, **55**:111
- Ketorolac tromethamine, **62**:186

- Keyhole limpet hemocyanin, **61**:369; **62**:149, 220
 antigen-presenting cells and, **47**:66, 69, 70
 autoreactive T cells and, **45**:419, 421
 HIV infection and, **47**:400
 humoral immune response and, **45**:10, 11, 15, 33
 IgE biosynthesis and, **47**:6, 14, 34, 35
 sporozoite malaria vaccine and, **45**:301
 synthetic T and B cell sites and, **45**:227, 255, 256
 T cells, **47**:86, 97, 99
 T cell subsets and, **41**:70, 71
 TNP-KLH-sensitized B cells, **60**:305–308
 virus-induced immunosuppression and, **45**:361, 362
- Ki antigen, **64**:22–23
- Kidney
 antigen-presenting cells and, **47**:76
 CD5 B cell and, **47**:134
 CTL and, **41**:149
 hybrid resistance and, **41**:338, 348, 356, 404
 IgE biosynthesis and, **47**:16
 NK cells and, **47**:187, 282
 regulators of complement activation and, **45**:389
 spontaneous autoimmune thyroiditis and, **47**:462, 493
 transplant, SCID and, **49**:399
 virus-induced immunosuppression and, **45**:337
- Kinase inhibitor protein-1, **61**:186–187
- Kinases, **59**:169, 178–179
 Ig gene superfamily and, **44**:30, 39, 44
 in signal transduction, **59**:261–266
- Kinetics
¹²⁵I-labeled IL-2 internalization, **42**:175
 Ars A response, **42**:108–109
- Kinin
 in hereditary angioedema, **66**:257–258, 259
 inactivation, **66**:240
- Kininogen
 genes, **66**:233
 high-molecular-weight
 contact activation and, **66**:236, 237
 endothelial cell receptor, **66**:249–257
 inactivation, **66**:245
 properties, **66**:226, 256
 structure, **66**:230–233
 low-molecular-weight, **66**:233
- Kit ligand, *see* Stem cell factor; Stem cell factor receptor
- Klebsiella pneumoniae*, **50**:162
- Knockout animals
 adhesion studies, **64**:187–188
 lymphocyte homing, **64**:194–195
- Knockout experiments
 cytokine induction, **65**:29–32
 with perforin gene, **60**:291–292
- Knockout mouse, LIF knockout, **53**:33, 48
- KOL, crystal packing, **43**:113
- KSH 4.13.6
 changes in after deprivation of RCS, **51**:226
 characterization of CTL activity of, **51**:225
- Ku, **64**:49, 60
 antinuclear antibodies and, **44**:96, 97, 104, 105
- KU 70/86, **61**:301–303, 317
- Kupffer cells, CD1 proteins, **59**:67
- Kynurenone, **62**:71

L

- L-β, **61**:22
- L1 cell adhesion molecule, 200-kDa glycoprotein, **58**:375–376
- L3T4, humoral immune response and, **45**:25, 26, 44, 45
- Labeling, CD1 proteins, **59**:45–46
- Lactate dehydrogenase B, suppressor T cells and, **38**:149–152, 184
- Lactation, lymphocyte homing and, **44**:321, 326, 327, 335
- LAK cells, *see* Lymphokine-activated killer cells
- LAM-1, surface antigens of human leukocytes and, **49**:87–88
- λ₅, **63**:7–8
- λ₅ gene, **63**:2
 expression, **63**:5–15
 μ heavy chains, **63**:2–3
 regulation, **63**:4–5
 structure, **63**:4
- λ light chain, **63**:4, 13
- λ light chain promoter, **43**:257
- λ phage repressor protein, **43**:209–210

- Lambert Eaton myasthenic syndrome, **42**:234, *see also* Myasthenia gravis
etiology, **42**:251, 252
- Lamina propria, **59**:372
lymphocytes, immunosenescence and, **46**:237
- MNC, IBD and, **42**:297
- T cells, IBD and, **42**:300–301
- Laminarin, phagocytosis of zymosan and, **38**:366, 377
alternative complement pathway, **38**:384–386
ingestion inhibition, **38**:381
- Laminin
complement receptor 1 and, **46**:199
hybrid resistance and, **41**:403
leukocyte integrins and, **46**:153
NK cells and, **47**:246
- Laminine, multiple myeloma and, **64**:249
- Langerhans' cells, **59**:60–62, 377, 379; **61**:13
antigen-presenting cells and, **47**:71–73, 75
antigen processing, **47**:90–92
APC-T cell binding, **47**:96, 97
cell surface, **47**:57, 59, 62, 63
immunogeneity, **47**:102
interaction with T cells, **47**:94
T cell growth, **47**:100
tissue distribution, **47**:50, 51, 53–57
- CD23 antigen and, **49**:154–155, 161, 167, 171
- corneal epithelium, **48**:197
- corneal graft rejection, **48**:196–200
- cytotoxic T cell, **48**:197–198
- delayed-type hypersensitivity, **48**:197–198
- dynamic distribution, **48**:200–202
- epithelial distribution, **48**:201
- Fc ϵ RI expression, **53**:1–2
Ia+, **48**:197
- IBD and, **42**:296
- immunoregulatory effects, **48**:203
- male vs. female tissue, **48**:198–199
- migration, **48**:201
- orthotopic graft, **48**:198
- regulatory process, **48**:202
- T cell subsets and, **41**:69, 92, 94
- LAP, **65**:2
- Large granular lymphocytes, **47**:188, 222, 283, 295
adoptive T cell therapy of tumors and, **49**:315
- CD23 antigen and, **49**:155
congenital defects, **47**:226
cytotoxicity, **47**:250, 253, 257
cytotoxicity and, **41**:271, 287, 289, 291, 309
differentiation, **47**:231, 234
effector mechanisms, **47**:248
hematopoiesis, **47**:275, 276, 278
hybrid resistance and, **41**:376, 380
identification, **47**:197–199
leukocyte integrins and, **46**:151
lymphocytosis, **47**:227–229, 275, 276, 279
lymphokines, **47**:262, 263
malignant expansion, **47**:226–228
morphology, **47**:214, 216–218
NK cells and, surface antigens of, **42**:182,
see also Natural killer cells
- reproduction, **47**:271
- surface phenotype, **47**:204, 205, 208, 211, 213
- tissue distribution, **47**:219–221
- Laron dwarfism, **63**:397
- Latent membrane protein, synthetic T and B cell sites and, **45**:226
- Latent membrane protein 1, CD23 antigen and, **49**:161
- L cells
B cell formation and, **41**:198
CTL
and amino acid, **41**:163
exon shuffling 138–140, **41**:144, 147, 148
HLA class I antigens, **41**:149–152
- Lck, **63**:135–137, 165, 171
IL-2R β coupling, **61**:155
- lok, expression regulation
restricted accumulation of transcripts, **56**:151–152
transcription, **56**:152–153
- LCL, *see* Large granular lymphocytes
- Lectin
antigen-presenting cells and, **47**:99, 100
autoimmune thyroiditis and, **46**:294
B cell formation and, **41**:201, 210
CD4 and CD8 molecules and, **44**:267, 268, 300, 302
CD23 antigen and, **49**:163–165, 176
CTL and, **41**:150
human T cell activation and, **41**:7–9, 12, 28
IgE biosynthesis and, **47**:15–20
immunosenescence and, **46**:221, 223, 245

- Lectin (continued)**
- lymphocyte homing and, **44**:354, 363, 367, 368
 - NK cells and, **47**:241, 250, 261, 286
 - spontaneous autoimmune thyroiditis and, **47**:470, 471
 - surface antigens of human leukocytes and, **49**:88–91, 94, 100
 - T cell subsets and, **41**:60, 76
- Lectins**, mitogenic, IBD and, **42**:289
- Lederberg's theory**, tolerance induction, **42**:48
- Legionella pneumophila*, **59**:398
- Leishmania*
- complement receptor 1 and, **46**:200
 - leukocyte integrins and, **46**:163
- Leishmania donovani*, and diabetes, **51**:294
- Leishmania major*
- helper T cell cytokines and, **46**:128, 129, 137, 138
 - NO expression during infection, **60**:342–343
 - responses in MHC class I-deficient mouse to infection, **55**:414–415
- Leishmaniasis, and $\gamma\delta$ T cell accumulation, **58**:326–328
- LEMS**, *see* Lambert Eaton myasthenic syndrome
- Lennert's lymphoma, gene rearrangement and, **40**:294
- Lens**
- anatomical sequestration, **48**:220
 - autoantigens, **48**:220
 - crystallin, **48**:219
 - α , **48**:220
 - antigen, **48**:191
 - γ , **48**:220
 - self-tolerance, **48**:220
- Lentiviruses**
- HIV infection and, **47**:395, 411
 - etiological agent, **47**:378, 384, 385
 - nonhuman primate, phylogeny, **52**:442–443
 - primate, phylogenetic tree, **52**:444
 - ungulate, **52**:430–432
 - virus-induced immunosuppression and, **45**:347–349, 351, 355
- Leprosy**
- CD1 proteins in lesions, **59**:66
 - Ig heavy-chain variable region genes and, **49**:46, 48
 - and $\gamma\delta$ T cell accumulation, **58**:326–328
- Lesch-Nyhan syndrome**, **59**:159
- Lesions**, cytotoxicity and, **41**:288–291, 318, 305–307
- LESTR**, **66**:282
- Leu-1**, **48**:246
- Leu-7**
- antigen, NK cells and, **47**:275, 291, 292
 - CNS, **47**:266
 - congenital defects, **47**:225
 - differentiation, **47**:234
 - genetic control, **47**:222
 - malignant expansion, **47**:226
 - morphology, **47**:218
 - surface phenotype, **47**:206, 207
 - tissue distribution, **47**:220, 221
- Leu-19**
- antigen, NK cells and, **47**:289, 293, 294, 302
 - CNS, **47**:266, 267
 - differentiation, **47**:233, 234
 - reproduction, **47**:271
 - surface phenotype, **47**:205, 206
- Leucine**
- antinuclear antibodies and, **44**:115
 - CTL and, **41**:166
- Leukemia**
- adoptive T cell therapy of tumors and mechanisms, **49**:300–305, 308, 311, 313, 317
 - principles, **49**:286, 291–293, 295–296
- B cell formation and, **41**:185, 191, 193, 205, 220, 233
- B-lineage, **61**:11–12
- CD1 proteins, **59**:60
- CD5 B cell and, **47**:123, 124
 - bone marrow transplantation, **47**:129, 130
- Ig gene expression, **47**:153, 154, 156, 157
- physiology, **47**:136, 140, 141, 143
 - surface antigen, **47**:147
- CD23 antigen and, **49**:150–151, 175
- diagnosis of, **40**:247–248, 249
- eosinophilic, **60**:223–224
- graft-versus, **40**:419–422
- human B cell neoplasia and, **38**:245, 271
 - acute B cell, **38**:268
 - acute lymphocytic, **38**:247, 264, 269
 - chromosome translocation, **38**:266
 - chronic lymphocytic, **38**:264, 265

- chronic myelogenous, **38:245**
human promyelocytic, **38:254**
- hybrid resistance and, **41:358–369**
antibodies, **41:378**, 380
antigen expression, **41:401**, 403
bone marrow cells, **41:346**, 348
- IgE biosynthesis and, **47:20**
- Ig heavy-chain variable region genes and,
49:17, 39–40
- IL-1 and, **44:194**
- lymphocyte hybridomas and
chronic lymphatic, **38:282**, 288, 291, 299,
302
human-human, **38:292**
Immunoglobulin secretion, **38:288**
- maps of Ig-like loci and, **46:7**
- mature T cell, receptor gene
rearrangements in, **40:276–284**
- myeloid
20αSDH activity, **39:36**
c-mpb locus rearrangement, **39:36–37**
IL-3-dependent lines from, **39:35–37**
- NK cells and, **47:190**, 199, 298
congenital defects, **47:227**
differentiation, **47:231**
effector mechanisms, **47:238**, 262
hem atop oie si s, **47:276**, 277
malignant expansion, **47:226**
surface phenotype, **47:205**
- pituitary hormones, **63:426**, 427
growth hormone, **63:409**, 428
prolactin receptor, **63:396**
- in PNH patients, **60:92**
- risk in rheumatoid arthritis patient,
64:329–330
- SCID and, **49:397**
- surface antigens of human leukocytes and,
49:92, 114, 117–118
- T cell, *see* T cell leukemia
- Leukemia inhibitory factor, **54:15–16**, 24–25;
61:151; **64:256**; **66:147–148**
and blastocyst regulation, **53:34**
in cachexia, **53:37–38**
in development, **53:33–37**
effects on mouse embryonic stem cells,
53:33–34
in hematopoiesis, **53:35–36**
human, amino acid sequence, **53:40**
and IL-6, **53:44–47**
immobilized form, **53:34**
- in inflammation, **53:37–38**
LIF knockout mouse, **53:48–49**
local action of, **53:49**
matrix-associated form, **53:34**
molecular cloning, **53:31–32**
pleiotropic nature of, **53:44**
polyfunctionality in vitro, **53:32–33**
pseudonyms for, **53:32–33**
purification, **53:31–32**
sepsis and, **66:104**, 148
soluble form, **53:34**
as stem cell factor, **53:35**
- Leukemia inhibitory factor receptor
alternatively spliced versions (murine),
53:39
cloned, **53:38–39**
cytoplasmic domain, **53:39–40**
extracellular domain, **53:39**
genes, chromosome locations, **53:39**
high affinity
characterization, **53:38**
and CNTF receptor, **53:43–44**, 46
oncostatin M binding, **53:41–43**
human, **53:38–39**
low affinity, **53:38**
transfected, **53:41**
types, **53:38**
- Leukemia inhibitory factor receptor β , **61:151**
- Leukemia virus, cytotoxic T cells and, **38:173**,
174
- Leukemic cell lines, **42:168**, 169, 172, 175
- Leukemic cells
cell typing, **48:117**
depletion, **58:272–275**
- Leukocyte adhesion deficiency, types I and II,
64:193
- Leukocyte adhesion deficiency disease,
46:151, 173–176
inflammation, **46:161**, 162, 165, 168
ligand molecules, **46:170**
- Leukocyte adhesion molecules, endothelial,
see Endothelial leukocyte adhesion molecules
- Leukocyte common antigen, antigen-presenting cells and, **47:60**, 61
- Leukocyte elastase, *see* Neutrophil elastase
- Leukocyte–endothelial cell recognition, models,
58:381–389
- Leukocyte function antigen, *see* LFA

- Leukocyte integrins, **46**:149–153, *see also* Integrins
 chromosomal localization, **46**:159, 160
 future research, **46**:176, 177
 inflammation, **46**:160, 161
 adhesion deficiency, **46**:161
 animal models, **46**:165
 expression, **46**:166–169
 function, **46**:161–164
 functional redundancy, **46**:164, 165
 mAbs, **46**:165, 166
 leukocyte adhesion deficiency disease, **46**:173–176
 ligand molecules, **46**:169–173
 structure, **46**:153, 154
 biosynthesis, **46**:154
 ligand binding, **46**:158, 159
 primary, **46**:154–158
- Leukocytes
 adhesion to endothelial cells, **58**:376–378, 381–389
 antigen-presenting cells and, **47**:76
 APC-T cell binding, **47**:97
 cell surface, **47**:61, 63
 tissue distribution, **47**:48, 51, 53
 B cell formation and, **41**:192, 194, 198, 229
 cachectin and, **42**:222
 CD5 B cell and, **47**:145
 as cellular source of chemokines, **55**:105–107
 cytotoxicity and, **41**:270, 312
 growth hormone, **63**:386–388
 helper T cell cytokines and, **46**:117
 HIV infection and, **47**:400
 hybrid resistance and, **41**:349, 350, 356
 IFNs, **62**:61
 IGF-I, **63**:389–391
 interaction with complement in acute injury states, **56**:274–277
 lymphocyte homing and
 carbohydrate, **44**:369
 molecules, **44**:342, 350
 regional specificity, **44**:339, 341
 migration, **58**:345–351
 NK cells and, **47**:188, 189, 295
 effector mechanisms, **47**:243, 263
 genetic control, **47**:222
 surface phenotype, **47**:201, 208
 perforin expression, **51**:222–223
 prolactin, **63**:382–384, 383
 recruitment during inflammation, **58**:394–395
 SCID and, **49**:383–384, 398
 spontaneous autoimmune thyroiditis and, **47**:492
 surface antigens, *see* HLA
 T cell subsets and, **41**:108
 and transendothelial migration, **58**:388–389
- Leukopenia, hybrid resistance and, **41**:388
 Leukophoresis, donor lymphocytes and, **38**:284
 Leukoregulin, cytolysis and, **41**:286
 Leukotriene A₄
 5-HPETE conversion to, **39**:150–151
 LTC₄ formation from, **39**:151–152
 Leukotriene A₄ hydrolase, **39**:152
 Leukotriene antagonists and inhibitors, **51**:346–347
 Leukotriene B₄, **62**:269
 attracting myeloid cells, **58**:381
 biological effects, **39**:156–157
 receptor-mediated, **39**:158
 CD23 antigen and, **49**:154, 159, 161
 corticosteroids and, **42**:316, 318–319
 eosinophil receptors for, **60**:185–186
 eosinophil response to, **60**:174–176
 IBD and, **42**:314–315
 immunoreactive, dietary fish oil and, **39**:164
 inactivation by EPO, **39**:202–203
 neutrophil activation by, **39**:135
 produced by neutrophils, **60**:189
 Leukotriene B₄ 20-hydroxylase, **39**:152
 Leukotriene B₄ receptors, neutrophil characteristics, **39**:96–98
 heterogeneity, **39**:99
 Leukotriene B₅
 biological effects, **39**:161–162
 EPA conversion to, **39**:160–161
 Leukotriene C₄
 5-HPETE conversion to, **39**:151–152
 biological effects, **39**:154–156
 receptor-mediated, **39**:157–158
 bronchial asthma and, **39**:223
 and eosinophil antihelminthic activity, **60**:212–214
 inactivation by EPO, **39**:202–203
 LTD₄ formation from, **39**:152–153
 Leukotriene C₄ synthetase, **39**:151–152

- Leukotriene C₅**
 biological effects, **39:162**
 EPA conversion to, **39:160–161**
- Leukotriene D₄**
 biological effects, **39:154–156**
 receptor-mediated, **39:157–158**
 inactivation by EPO, **39:202–203**
 LTC₄ conversion to, **39:152–153**
- Leukotriene E, IBD and**, **42:315**
- Leukotriene E₄, biological effects**,
39:154–155
 receptor-mediated, **39:157–158**
- Leukotrienes**, **51:341–347**
 arachidonic acid conversion to, **39:**
147–149
 5-lipoxygenase and, **39:147–150**
 cell specificity, **39:153**
 biological activities, **51:343**
 EPA-derived
 5-lipoxygenase pathway, **39:160–161**
 biological properties, **39:161–162**
 in human neutrophils, dietary EPA and,
39:166
 in asthmatic subjects, **39:166–167**
 hybrid resistance and, **41:360**
IBD and, **42:310–315**
 immediate hypersensitivity and,
39:218
 leukotriene antagonists and inhibitors,
51:346–347
 leukotrienes and airway
 hyperresponsiveness, **51:344–346**
 metabolism
 5-HPETEdehydration to LTA₄,
39:150–151
 γ-glutamyl transpeptidase and,
39:152–153
 LTA₄ hydrolase and, **39:152**
 LTB₄ 20-hydroxylase and, **39:152**
 LTC₄ synthetase and, **39:151–152**
 as neutrophil activators, **39:135–136**
 potency, **51:343**
 production by
 eosinophils, **39:204**
 marine mastoeytoma, dietary EPA and,
39:162
 rat peritoneal cavity
 EPA-enriched diet and, **39:163**
 fish oil-enriched diet and, **39:**
162–163
 release of leukotrienes in asthma, **51:346**
- sulfidopeptide
 binding sites for, **39:157–158**
 biological effects, **39:154–156**
 synthesis and metabolism, **51:341–343**
- Leupeptin**
 synthetic T and B cell sites and, **45:223**
 T cell receptor and, **45:124**
- LFA**
 CTL and, **41:136, 139, 150–152, 154**
 cytotoxicity and, **41:271, 272**
 IL-1 and, **44:166**
- LFA-1**, **48:247; 54:309–310**
 activation by chemokine, **58:381**
 affinity of ICAM-1 and ICAM-2,
58:373–374
- B cell formation and, **41:194**
- CD23 antigen and**, **49:161**
 conformational effects, **64:169–170**
 humoral immune response and, **45:39, 41,**
43, 45, 51, 52, 54
- IFN-γ and**, **62:78, 81**
- integrins**, **58:369–370**
 interaction with ICAM-1 in negative
 selection, **58:172**
- leukocyte integrins and**, **46:149–153**
 inflammation, **46:160–164, 166–169**
 leukocyte adhesion deficiency disease,
46:173, 174
 ligand molecules, **46:169–172**
 structure, **46:153, 154, 156, 158**
- lymphocyte–HEV interactions and**, **65:352,**
358, 368, 370
- lymphocyte homing and**, **44:350, 351, 361;**
64:146, 155, 157
- multiple myeloma and**, **64:247**
- surface antigens of human leukocytes and**,
49:86, 88, 92
- LFA-2/CD2 and LFA-3/CD58**, **52:361**
- LFA-3**, **48:236; 64:160**
 CD23 antigen and, **49:161**
 surface antigens of human leukocytes and,
49:87
- LH**, *see* Luteinizing hormone
- LIF**, *see* Leukemia inhibitory factor
- Life expectancy, and autoimmune**
 manifestations, effect of IL-2.VV,
50:195–296
- Life span**
 of immunological memory, **53:218–219**
 intermitotic, of B and T cells, **53:237–241**

- Li-Fraumeni syndrome, and mutant p53 allele, **58:66–67**
- Ligand binding
association with hematopoietic cell phosphatase, **60:26**
cellular consequences, **60:1–2**
induced receptor aggregation, **60:8**
- Ligand blotting, for LPS receptor identification, **53:279–280**
- Ligand–receptor interactions, **38:363; 51:148**
T cell activation and, **41:1, 26**
- Ligand recognition, and CD44, **54:271–272**
- Ligands
antigen-presenting cells and, 0; **47:64, 65, 68, 87**
antigen processing, **47:88, 89, 92**
cell surface, **47:61**
T cells, **47:94–96, 101**
aPL antibodies and, **49:228–231**
CD4 and CD8 molecules and, **44:267, 282, 298, 300, 302**
CD5 B cell and, **47:146**
CD23 antigen and, **49:158, 165, 167**
complement receptor 1 and, **46:183, 192–196, 199, 200**
complement receptor 2 and, **46:205, 211**
CTL and, **41:168, 169**
carbohydrate moieties, **41:154**
exon shuffling, **41:139, 142, 147**
HLA class I antigens, **41:149–152**
mAbs, **41:158**
- cytotoxicity and, **41:307**
- HIV infection and, **47:391**
- Ig gene superfamily and evolution, **44:43, 44**
- IL-1 and, **44:180, 181, 184, 185**
- immunosenescence and, **46:224**
- Ir* genes and, **38:105, 173**
- leukocyte integrins and, **46:150, 153, 169–173, 176, 177**
inflammation, **46:163, 167**
structure, **46:156, 158, 159**
- lymphocyte homing and, **44:314**
carbohydrate, **44:355, 360, 362–371**
high endothelial venules, **44:319–322**
molecules, **44:344, 349, 350, 352, 353**
regional specificity, **44:323, 324, 326, 335, 336, 340**
- murine lupus models and, **46:94**
- NK cells and, **47:221, 247, 265**
- nonimmune receptor members, **44:32, 33, 35, 39**
- phagocytosis and, **38:367–369, 371**
- positive selection, **58:160–164; 59:122–128**
- preB cell receptors, **63:25–26**
and receptors, in positive selection, **51:178–180**
- SCID and, **49:385**
- specificity, **38:381, 385**
- surface antigens of human leukocytes and, **49:113, 115, 117**
adhesion molecules, **49:81, 86–87**
antigen-specific receptors, **49:76**
Igs, **49:89**
receptors, **49:99, 101, 111**
- T cell activation and, **41:30**
cell surface molecules, **41:2, 7, 8, 10–13**
gene regulation, **41:28, 30**
receptor-mediated signal transduction, **41:20, 22**
- synergy, **41:17**
- T cell development and
antigen recognition, **44:213, 214**
cellular selection within thymus, **44:245, 246**
- thymocyte subpopulation, **44:228, 229**
- T cell subsets and, **41:40, 50–53**
- TCR, inducing deletion of thymocytes, **58:170–171**
- Light chain, *see also* κ light chain; λ light chain
- B cell development, **63:1–2**
- genes, **48:41–64; 63:4**
diversity, **48:41**
gene conversion, **48:41–42**
hypermutations, **60:277–278**
independent genes, **48:41**
joining events, **48:41**
- surrogate
B cell development, **63:1–31**
expression, **63:5–20**
genes, **63:4–5**
transduction in complex, **63:25–29**
variable region gene expression, **42:3–5**
- Light microscopy, antigen-presenting cells and, **47:51, 52**
- LIM domain, oncogenes by translocation, transcriptional disruption of, **50:130–131**

- Limiting dilution analysis
NK clones and, **42**:182–184
SCID and, **49**:394, 400
- Limnea*, maternally transmitted antigen and, **38**:322
- Lineage fidelity, B cell formation and, **41**:195
- Lineage-related markers, differential expression during development (murine B cells), **53**:125–127
- Linear tracking, Ig heavy-chain variable region genes and, **49**:33, 38
- Linkage disequilibrium phenomenon
DQ, **48**:136–137
DR, **48**:136–137
- Linked substitution, Ig heavy-chain variable region genes and, **49**:9, 14
- Linomide
autoimmune side effects, **58**:278–279
reduction of DNA fragmentation, **58**:263–265
- LIP, **65**:2
- Lipid
antigen-presenting cells and, **47**:65
aPL antibodies and, **49**:252, 256–257, 259
affinity purification, **49**:228–230
antibody subsets, **49**:232
biochemistry, **49**:195, 197
clinical aspects, **49**:205
specificity, **49**:234, 238–240, 243, 246–248
- B cell formation and, **41**:197, 213
- cell-mediated killing and, **41**:290, 291, 312, 313, 294–296
- complement receptor 2 and, **46**:210
- CTL and, **41**:169
- cytolysis and, **41**:275, 278–280
- helper T cell cytokines and, **46**:114
- HIV infection and, **47**:387
- IL-8 bioactivation in neutrophils, **55**:114
- leukocyte integrins and, **46**:170
- membrane attack complex of complement and, **41**:299–303, 305, 306
- NK cells and, **47**:217
- pore formers, **41**:315, 317, 318
- spontaneous autoimmune thyroiditis and, **47**:439, 491
- T cell receptor and, **45**:120, 124
- Lipid A
antagonists, **53**:272–274
binding proteins, **53**:274–281
- chemical structure, **53**:270
from enterobacteria, **53**:269
from nonenterobacteria, **53**:269–270
receptor identification, **53**:274–281
recognition by receptor-mediated mechanisms, **53**:271–272
- structure, **53**:269
structure-function relationships, **53**:272–274
- Lipid bodies, eosinophil, characterization, **60**:155–156
- Lipid IVa
32P-labeled, for lipid A binding site characterization, **53**:278–279
recognition by scavenger receptor, **53**:280
- Lipid mediators, eosinophil-derived, **60**:188–191
- Lipoarabinomannan, **59**:85, 90; **65**:26
- Lipocortin, IgE biosynthesis and, **47**:24, 25, 38
- Lipodystrophy, partial, **61**:245
- Lipomodulin, **42**:312
IBD and, **42**:316, 318–319
- Lipopolsaccharide, **54**:373–374; **65**:27
adoptive T cell therapy of tumors and, **49**:307–310, 314
antigen-presenting cells and, **47**:68, 70, 76, 80
antigen processing, **47**:89, 90, 92
cell surface, **47**:57, 59, 61–63
immunogenicity, **47**:104
tissue distribution, **47**:49
- aPL antibodies and, **49**:211
- autoimmune thyroiditis and
cellular immune responses, **46**:294
experimental models, **46**:276, 277, 279
genetic control, **46**:286
prevention, **46**:310, 311
- autoreactive T cells and, **45**:428
- bacterial, structure, **53**:268–271
- B cell activation
antigen presentation and, **39**:67, 69
radiosensitivity and, **39**:71
- B cell formation and, **41**:196, 200, 221, 222, 231
- cachectin and, **42**:215, 218, 220, 223
- CD5 B cell and, **47**:159
anatomic localization, **47**:121

- Lipopolysaccharide (*continued*)
 marker for activation, **47:126**
 physiology, **47:133–135, 142, 143**
 CD23 antigen and, **49:150–151**
 cellular recognition, role of LBP/CD4 pathway, **53:281–286**
 complement receptor 2 and, **46:205**
 enhancement of iNOS, **60:333–337**
 helper T cell cytokines and, **46:116, 119, 121**
 hepatotoxicity, **60:347**
 heterodimeric receptor models, **53:285–286**
 HIV infection and, **47:402**
 humoral immune response and
 helper T cells, **45:27, 33, 34**
 interleukins, **45:63, 64, 66, 68, 70**
 physical interaction, **45:44, 46, 53**
 IgE biosynthesis and, **47:6, 8, 9, 11**
 immunosenescence and, **46:223, 234, 242**
 leukocyte integrins and, **46:171**
 lipid A
 antagonists, **53:272–274**
 binding proteins, **53:274–281**
 chemical structure, **53:270**
 from enterobacteria, **53:269**
 from nonenterobacteria, **53:269–270**
 receptor identification, **53:274–281**
 recognition by receptor-mediated mechanisms, **53:271–272**
 structure, **53:269**
 structure-function relationships, **53:272–274**
 murine lupus models and, **46:63, 72**
 NK cells and, **47:289**
 pore-forming protein and, **41:313**
 regulators of complement activation and, **45:397**
 SCID and, **49:384**
 spontaneous autoimmune thyroiditis and, **47:474**
 stimulated splenic B cells, IL-4 effects, **54:247–248**
 treatment with, **50:65**
 virus-induced immunosuppression and, **45:357**
 Lipopolysaccharide-binding protein, in LPS-induced macrophage stimulation, **53:281–286**
 Lipopolysaccharide–cytokine score, **66:134**
- Lipoprotein, aPL antibodies and, **49:213, 256**
 Lipoprotein lipase
 aPL antibodies and, **49:257**
 cachectin and, **42:214–216**
 Liposomes
 aPL antibodies and affinity purification, **49:228–230**
 LA antibodies, **49:242, 244–246**
 specificity, **49:234–236, 239, 250**
 CTL and, **41:154**
 cytotoxicity and
 cytolytic proteins, **41:319**
 membrane attack complex of complement, **41:300, 302, 305, 309**
 pore-forming proteins, **41:312, 313**
 with Ia from normal and lymphoma B cells, antigen-presenting capacity, **39:76–77**
- Lipoxygenase
 action of, **65:122**
 products, IBD and, **42:313–314**
- 5-Lipoxygenase
 activation by
 calcium ionophore A23187, **39:154**
 FMLP, **39:154**
 leukotriene biosynthesis and, **39:147–151, 153, 160–161**
 5-HETE dehydration to LTA₄, **39:150–151**
 properties, **39:150–151**
- Listeria monocytogenes*, **54:207; 59:398**
 and IFN- γ -secreting IEL, **58:329**
- Liver
 acute phase protein synthesis, **54:25**
 antinuclear antibodies and, **44:110, 123, 125, 129, 130**
 B cell formation and, **41:181, 182, 224, 225**
 B cell precursors, **41:191, 193**
 bone marrow cultures, **41:216**
 Ig genes, **41:204**
 lymphohemopoietic tissue organization, **41:187**
 population dynamics, **41:207**
 CD1 protein distribution, **59:73**
 CD5 B cell and, **47:141, 158**
 complement components and
 cDNA, **38:234**
 Factor B synthesis, **38:217**
 mRNA, **38:226**
 complement receptor 1 and, **46:198**

- complement receptor 2 and, **46:204**
 CTL and, **41:149**
 hybrid resistance and, **41:335**, 351, 356,
 363, 366, 373, 386
 Ig gene superfamily and, **44:37**
 IL-1 and, **44:166**, 168, 178, 189
 lymphocyte homing and, **44:328**
 lymphocyte homing to, **64:178–179**
 NK cells and, **47:283**, 296, 297
 genetic control, **47:222**
 hematopoiesis, **47:276**
 tissue distribution, **47:220**, 221
 protein synthesis
 cachectin and, **42:225**
 IL-1 and, **44:165**, 166
 regulators of complement activation and,
 45:410
 protein expression, **45:397**, 399
 protein interactions, **45:384**
 protein roles, **45:388**, 389
 SCID and, **49:392–393**, 401
 sporozoite malaria vaccine and, **45:319**
 antibodies, **45:301**
 CS proteins, **45:300**
 CS-specific T cells, **45:312**
 immunity, **45:287**, 289, 290
 interferon- γ , **45:301–306**
 Plasmodium vivax, **45:317**
 synthetic T and B cell sites and, **45:228**
 T cell receptor and, **45:129**
 T cell subsets and, **41:67**, 69
 T lymphopoiesis in, **53:169–171**
 Liver cell grafts, hybrid resistance and,
 41:333
 Liver diseases, aPL antibodies and, **49:250**
 Liver-enriched transcriptional activator
 protein, **65:2**
 Liver inhibitory protein, **65:2**
 L-IVS regions, V_H gene segments and,
 42:116, 117–119
 LM34 antibody, **63:14–15**
 LMP1-associated protein 1, **61:29**
 LMP-2, **64:12**, 14, 106, 107
LMP-2, **64:12**, 14, 19–20, 106
 LMP-7, **64:12**, 14, 106, 107
LMP-7, **64:12**, 14, 19–20, 106
 LMP-9, **64:106**
 LMP-10, **64:107**
LMP-10, **64:106**
 LMP-17, **64:106**
 LMP-19, **64:106**
 LNGFR, *see* Low-affinity nerve growth factor
 receptor
 Lock-and-key fit, **43:78**
 Loefflers syndrome, in pulmonary
 eosinophilia, **60:226**
 Long homologous repeat
 complement receptor 1 and, **46:185**,
 187–189, 191, 194–196
 complement receptor 2 and, **46:203**
 Long terminal repeat enhancer, viral infection
 and, **65:9–10**
 Long terminal repeats
 HIV-1, **58:14–17**
 HIV infection and
 etiological agent, **47:380**, 382, 385
 immunopathogenic mechanism,
 47:399–402
 spontaneous autoimmune thyroiditis and,
 47:464
 Long-term nonprogressors, HIV, **65:298–300**
 Low-affinity nerve growth factor receptor,
 61:1, 2, 7
 Low-density lipoprotein
 antigen-presenting cells and, **47:89**
 B cell formation and, **41:212**, 213
 cytotoxicity and, **41:295**, 307, 309
 regulators of complement activation and,
 45:402
 Lower respiratory tract, pulmonary immunity
 regulation and, **59:372–373**
 Low-molecular-weight B cell growth factor,
 CD23 antigen and, **49:160**, 169
 Low-molecular-weight kininogen, **66:233**
 LPAM-1, **65:359**
 LPL, cachectin and, **42:214–216**
lpr mouse, Fas gene mutation in, **57:132–133**
lpr phenomenon, and IL-2, **50:202**
 LPS, *see* Lipopolysaccharide
 LR1, binding site, **61:120**
 L-Selectin, **54:310**; **59:384**; **62:257–258**;
 64:144–145, 173; **65:352**, 354, 367
 activation and signaling via, **65:357**
 downregulation, **64:172**
 genetic studies, **64:194**
 lymphocyte–HEV interaction, **65:355**, 357,
 358, 361
 in PLNs, **64:174**
 properties, **64:146**
 as pro-T cell homing molecule, **58:393**

- L-Selectin (*continued*)
 regulation of, 65:356–357
 shedding, 64:170
 shedding of, 65:356
 soluble form, 64:171
 structure and function of, 65:354–356
 sugar structure of, 65:363–365
 T cells positive and negative for,
 58:390–391
 upregulation, 64:165
 widespread distribution on leukocytes,
 58:353–354
- LT- β , 61:22, 23, 24
- Ltk, 59:169
- Lung
Cryptococcus neoformans clearance, 59:403
 expansion of resident $\gamma\delta$ T cells, 58:321
 genetically engineered antibody molecules
 and, 44:68, 87
 hybrid resistance and, 41:356, 366, 373
 Ig gene superfamily and, 44:39
 inflammation, *see* Pulmonary inflammation
 invariant $\gamma\delta$ T cell receptors of murine
 IEL, 58:303–304
 lymphocyte homing, 64:178–179
 carbohydrate, 44:363
 regional specificity, 44:328–330, 334,
 336–338
 lymphocyte occurrence, 58:299
 mast cells, IL-4 storage by, 53:15
 pulmonary immunity, 59:369–427
 sensitized guinea pig, hypersensitivity,
 dietary fish oil and, 39:164–165
 structure, 59:370–373
 BALT, 59:385–387
 dendritic cells, 59:376–381
 lymphocytes, 59:381–385
 macrophages, 59:373–376
 transplantation, 59:420–426
 graft-versus-host disease, 59:420–426
 Lung-associated lymph nodes, 59:372, 376,
 379, 381, 388–389, 391
- Lung cancer, etiology, stem cell factor role,
 55:77–78
- Lung disease
 chemokine role, 55:144–146
 graft-versus-host disease, 59:420–426
 hypersensitivity diseases, 59:387, 412–420
 IL-1ra effects, 54:215–216
 IL-8 role, 55:144–146
- immunity, 58:328–329
 pneumonia, 59:379, 383, 395–408
- Lupus, *see* Systemic lupus erythematosus
- Lupus anticoagulant, aPL antibodies and,
 49:193, 258–259
 affinity purification, 49:229–231
 antibody subsets, 49:231–233
 clinical aspects, 49:200–210, 212, 221, 224
 history, 49:198–199
 isotype, 49:227
 pathogenic potential, 49:253–256
 specificity, 49:234, 239–248, 251–252
 syndromes, 49:215–218
- Lupus mouse, *see also* Systemic lupus
 erythematosus, murine
 cellular abnormalities
 defects in tolerance, 37:330–332
 functional abnormalities of B cells, T
 cells, macrophages and related
 interleukins, 37:308–330
 surface characteristics and numerical
 abnormalities of T and B cells,
 37:303–308
 thymic defects, 37:332–336
 transfer of autoimmune disease,
 37:301–303
- derivation of, 37:271–273
 inbred strains, 50:193
 natural history and pathology of,
 37:273–274
 morphologic manifestations, 37:276–284
 serologic manifestations, 37:284–300
 survival and body weight, 37:274–276
- Luteinizing hormone
 synthetic T and B cell sites and, 45:259
 thymic factor-induced release, 39:315
- L-VAP-2, and VAP-1, 58:377–378
- Ly1 B cell, 42:64–65
 splenic B cell and, 42:68
 subpopulation, 42:82–83
- Ly1 lineage, CD23 antigen and, 49:150–151
- Ly-6, 48:240–241
- Lymphadenopathy
 and autoimmune symptoms in MRL/Mp-
 lpr/lpr mice, amelioration, 50:199
 cardinal feature of HIV-related, 51:277
 murine lupus models and, 46:63
- Lymph follicles
 B cell populations, 53:186–187
 zonal organization, 53:185–186

- Lymph node cells
 ESP production, **39:214**
 IgE biosynthesis and
 antibody response, **47:3–5**
 binding factors, **47:12, 14, 16, 17**
- Lymph nodes
 adoptive T cell therapy of tumors and, **49:284, 331**
 antigen-presenting cells and, **47:65, 71–73, 76**
 immunogenicity, **47:102**
 T cells, **47:85, 94**
 tissue distribution, **47:50–52, 54, 55**
 autoimmune demyelinating disease and, **49:358–359, 364, 372**
 autoimmune thyroiditis and, **46:280, 296**
 autoreactive T cells and, **45:419**
 B cell formation and, **41:191, 223**
 CD4 molecules and, **44:288**
 CD8 molecules and, **44:271**
 complement receptor 2 and, **46:204, 209**
 forbidden α/β T cells, **53:166**
 helper T cell cytokines and, **46:129, 137, 138**
 high endothelial venules, **65:373–377**
 HIV infection and, **47:410**
 humoral immune response and
 antigens, **45:23, 24**
 cellular interactions *in vivo*, **45:79, 81**
 physical interaction, **45:54**
 hybrid resistance and, **41:359, 384, 393**
 bone marrow cells, **41:337, 338**
 lymphoid cells, **41:352, 353, 355, 356**
 NK cells, **41:372**
 IL-1 and, **44:194**
 immunosenescence and, **46:236, 245, 250, 251**
 lymphocyte homing and
 carbohydrate, **44:363, 365**
 high endothelial venules, **44:315, 317, 318, 321**
 mesenteric, *see* Mesenteric lymph nodes
 molecules, **44:344**
 mucosal immune system, **44:328, 329**
 peripheral, *see* Peripheral lymph nodes
 regional specificity, **44:322, 323, 325, 331, 336, 337**
 murine lupus models and, **46:95, 96, 98**
 NK cells and, **47:211, 219, 220, 228, 298**
 peripheral, *see* Peripheral lymph nodes
- prolactin, **63:382–383**
 repopulation, immune reconstitution and, **40:411–412**
 reticular conduit system, **65:375**
 sporozoite malaria vaccine and, **45:305**
 T cell development and, **44:243, 244**
 T cell populations, **53:161–163**
 T cell receptor and, **45:134**
 T cells, **41:381**
 T cell subsets and
 H-2 alloantigen recognition, **41:78, 92**
 H-2 molecules in thymus, **41:99**
 H-2-restricted antigen recognition, **41:67–71**
 virus-induced immunosuppression and, **45:350, 353, 354, 360**
- Lymphoblast foci, PALS-associated, **60:270–271**
- Lymphoblastoid cells
 complement receptor 1 and, **46:194**
 complement receptor 2 and, **46:201–204, 207, 209, 210, 213**
 human B cell neoplasia and, **38:254, 257**
 c-myc activation, **38:260, 261, 263**
 IM-9, SP receptor recovery from, **39:313**
 leukocyte integrins and, **46:172**
- Lymphoblasts, *see also* B lymphoblasts; T lymphoblasts
 antigen-presenting cells and, **47:94, 96, 103**
 B cell hybridomas, **38:282**
 CTL and, **41:139**
 Epstein-Barr virus-transformed, **38:280**
 germinal center, **53:186**
 human-human hybridomas, **38:293**
 leukocyte integrins and, **46:162**
 lymphocyte homing and, **44:314**
 high endothelial venules, **44:317**
 regional specificity, **44:322–324, 328, 330**
 NK cells and, **47:231**
 spontaneous autoimmune thyroiditis and, **47:469**
- Lymphochoriomeningitis virus, *see*
 Lymphocytic choriomeningitis virus
- Lymphocyte activating determinants, **38:88**
- Lymphocyte-activating factor, IL-1 and, **44:154, 172, 173, 189**
- Lymphocyte function-associated antigen, *see*
 LFA

- Lymphocyte homing, **44**:313, 314; **64**:139–140, 173–174, 198
 carbohydrate
 cell adhesion, **44**:353–355
 homing receptors, **44**:369, 370
 inhibition, **44**:355–357
 ligands, **44**:363–371
 PPME-binding receptor, **44**:357–363
 CD44 role *in vivo*, **54**:298–299
 expression of CD44 on hematopoietic cells, **54**:291–296
 and hematopoiesis
 inhibition *in vitro* by CD44-specific antibodies, **54**:299–300
 physiological experiments implying role for CD44, **54**:296–302
 hermes antigen mediating PLN adhesion, **54**:296–298
 high endothelial venules
 HEV-associated molecules, **44**:352, 353
 interaction, **44**:318–322
 lymphocyte associated molecules, **44**:342–352
 structure, **44**:315–318
 inflammation and, **64**:188–193, 195–197
 liver and lung, **64**:178–179
 migration, role of CD44 *in vivo*, **54**:298–299
 mucosa, **64**:175–177
 PLN, **64**:174–175
 regional specificity
 blast cell migration, **44**:322, 323
 breast, **44**:326, 327
 immune response, **44**:330–336
 inflammation, **44**:339–341
 mucosal immune system, **44**:327–330
 regulation, **44**:336–339
 selective homing model, **44**:323–326
 regulation, **64**:173–188
 signaling through CD44-ligand interactions, **54**:300–302
 thymocyte progenitors, migration inhibition, **54**:299
 thymus and bone marrow, **64**:177
 Lymphocyte hybridomas
 cell lines for, **38**:279–282
 fusion, **38**:286, 287
 mAbs, **38**:275–277
 Lymphocyte receptors, **62**:41–44
 neuromediator, **48**:179–184
- Lymphocytes, **51**:339–340
 adhesion, **66**:15–17
 adhesion deficiencies, **64**:193–195
 adhesion molecules
 multistep adhesion cascade, **64**:181–184
 regulation, **64**:139–140, 146, 147, 163–173
 anatomic compartments colonized by, **53**:159
 antiadhesion and proadhesion therapy, **64**:195–197, 198–199
 antigen-presenting cells and, **47**:65, 91, 95, 105
 antinuclear antibodies and
 autoimmune resopns, **44**:132
 scleroderma, **44**:120
 SLE, **44**:95, 102, 107, 108
 aPL antibodies and, **49**:239, 249
 autoimmune demyelinating disease and, **49**:368–369
 autoimmune thyroiditis and, **46**:263, 318
 antigens, **46**:269
 cellular immune responses, **46**:287–289, 293, 294, 296–298
 experimental models, **46**:273, 274, 276, 278, 279, 281, 282
 genetic control, **46**:285, 286
 humoral responses, **46**:299–301, 303, 306
 prevention, **46**:310, 312, 315
 B, *see* B cells
 biology, LIF role, **53**:36
 CD4 and CD8 molecules and, **44**:265, 288
 CD5 B cell and, **47**:117, 118, 149, 150, 153, 157, 161
 bone marrow transplantation, **47**:130
 lineage, **47**:128, 129
 malignancies, **47**:123
 marker for activation, **47**:125, 126
 ontogeny, **47**:121, 122
 physiology, **47**:135, 137–139
 surface antigen, **47**:143–145
 CD23 antigen and, **49**:164, 173, 175
 CD45 and, **66**:4, 5–31
 B cells, **66**:12–15, 36–37
 lymphocyte adhesion, **66**:15–17
 monoclonal antibody studies, **66**:1, 3, 5–17
 T cells, **66**:6–12, 32–36

- chemotaxis effects of
 opioid peptides, 39:307–308
 SP, 39:308
- circulating, abnormalities in IL-2 and IL-2R expression and responsiveness, 50:186–187
- circulating, functions of, 53:201–203
- compartments
 generation
 by different lineages, 53:188–189
 by local selection, 53:192–199
 by selective migration and
 extravasation, 53:189–192
 site-dependent tolerance, 53:201–203
 specialized functions of, 53:199–201
- complement-mediated cytotoxicity and, *see*
 Cytotoxicity, lymphocyte and
 complement-mediated
- complement receptor 1 and, 46:190, 191,
 193, 198–201
- complement receptor 2 and, 46:203, 205,
 207, 209, 210, 213
- CRF production, 39:316
- cytokines produced, 48:71
- cytotoxic T, *see* Cytotoxic T cells
- deficient development, 55:393–394
 epithelium, intestinal, 55:395–396
 lymphoid organs, 55:394
 models, instructive, 55:398–399
 models, selective, 55:398–399
 thymus, 55:394
- development
 CD44 involvement, 54:271
 self-tolerance checkpoints, 59:279–344
- β -endorphin receptors, 39:312
- endothelial cell interactions, 64:140–143,
 163, 185–186
 model, 64:182
- function, neuroendocrine mediator,
 48:177–179
- gene function
 Bcl-2 and Bcl-x, 62:43, 53
 cis-regulatory elements, 62:42–43, 49–52
 perforin, 62:43, 52–53
 signal transduction, 62:42, 44–47
 TdT, 62:43, 52
 transcription, 62:42, 47–49
- genetically engineered antibody molecules
 and, 44:76
- helper T, *see* T helper cells
- helper T cell cytokines and, 46:123, 124
- hemopoietic regulators produced, 48:71
- hepatocyte growth factor effects,
 58:379–380
- HIV, and programmed cell death, 58:214
- HIV infection and
 immune response, 47:410, 411
 immunopathogenic mechanism, 47:390,
 392, 393, 397, 398, 400
- homing, *see* Lymphocyte homing
- homing receptors, 62:257; 64:143, 146, 147
- CD40, 64:163
- CD73, 64:162–163
- chemoattractants, 64:152–154
- Ig superfamily, 64:147, 157–160
- integrins, 64:146, 154–157
- MEL-14, 64:142–143
- multiple myeloma, 64:246
- proteoglycans, 64:146, 160–161
- selectins, 64:144–145, 146, 147, 148–152
- sialomucins, 64:144, 146, 147, 148–152
- vascular adhesion protein, 64:161–1662
- hybrid resistance and
 antigen expression, 41:400
 bone marrow cells, 41:348, 350
 leukemia/lymphoma cells, 41:361, 364
 lymphoid cells, 41:353, 355–357
 marrow, 41:387, 390
 T cells, 41:380
- IBD and, 42:295–296, 297, 321, *see also*
 Peripheral blood lymphocytes
- antibodies and, 42:291
- Crohn's disease and, 42:297
- IFN- γ and, 62:86–87
- IgE biosynthesis and
 antibody response, 47:3–8, 11
 antibody response suppression, 47:33, 38
 binding factors, 47:13, 14, 19–22, 24, 25
- IL-1, 44:153
 biological effects, 44:166
 human disease, 44:192
 immunocompetent cells, 44:173
 receptor, 44:181
 structure, 44:155
 TNF, 44:185, 186
- IL-8 effects on, 55:118–119
- immune responses, neuropeptides and,
 39:309–311
- immunosenescence and, 46:221, 253
 activation, 46:221–234

- Lymphocytes (*continued*)
 mucosal immunity, **46**:251, 252
 regulatory changes, **46**:249
 stem cells, **46**:240, 246, 247
 subsets, **46**:236–240
 interaction with IgA, **40**:171
 intraepithelial, *see* Intraepithelial lymphocytes
Ir genes and, **38**:37, 56, 62
 H-Y antigen, **38**:177
 macrophages, **38**:113
 positive T cell selection, **38**:130
 T cell suppression, **38**:147
 lamina propria, immunosenescence and, **46**:237
 leukocyte integrins and, **46**:149, 151
 inflammation, **46**:162–164, 169
 leukocyte adhesion deficiency disease, **46**:173
 ligand molecules, **46**:170–172
 mAbs and, **38**:276
 Epstein-Barr virus, **38**:277
 fusion, **38**:282–286
 human-human hybridomas, **38**:293, 294, 296–298
 human-mouse heterohybridomas, **38**:301
 human-murine hybridomas, **38**:290–292
 maps of Ig-like loci and, **46**:1
 mediated cytolysis, perforin role expression, **51**:219–224
 in human leukocyte, **51**:222–223
 regulation of, **51**:223–224
 perspectives, **51**:232–233
 role of in cytolysis, **51**:215–216, 224–232
 perforin-independent mechanisms, **51**:231–232
 verification by genetic approach, **51**:227–231
 verification in cell lines, **51**:224–227
 structure of, **51**:216–219
 memory and effector, **58**:347–350
 migration
 diapedesis, **65**:369
 extravasation, **65**:360, 365–367
 GlyCAM-1, **65**:361–362, 363, 364, 373, 377
 CD34, **65**:362, 377
 Sgp200, **65**:362, 377
 high endothelial venules and, **65**:350–351
 addressins, **65**:361–363, 377–379
 adhesion, **65**:365–372
 CD44, **65**:369–370
 extravasation, **65**:360, 365–367
 GlyCAM-1, **65**:361–362, 363, 364, 373, 377
 homing receptor ligands, **65**:360–365
 integrins, **65**:352, 358–359, 368–369, 370
 MAdCAM-1, **65**:352, 359, 362–363, 367, 370, 377, 379
 mucins, **65**:360–361
 selectins, **65**:352, 354–357, 363–365, 367
 sugars, **65**:360–361, 363–365
 VAP-1, **65**:365
in vitro binding assay, **65**:351–352
 organ specificity, **65**:370–372
 recirculation pathways, **65**:371–372
 mixed, met-enkephalin receptors, **39**:312
 naive, **58**:389; **64**:180–181
 neuropeptide, **48**:177
 differentiation, **48**:178
 immunological functions, **48**:178
 tissue cycles, **48**:178
 tissue homing, **48**:178
 neuropeptide receptor, **48**:180
 NK cells and, **47**:187–189
 adaptive immunity, **47**:291
 alterations, **47**:302
 antimicrobial activity, **47**:289
 antitumor activity, **47**:297, 299
 cell-mediated cytotoxicity, **47**:190, 194–196
 congenital defects, **47**:224–226
 cytotoxicity, **47**:252–255, 257, 258, 260, 261
 differentiation, **47**:230, 231, 234
 effector mechanisms, **47**:243, 246
 genetic control, **47**:222, 223
 hematopoiesis, **47**:272–274, 276, 280, 282
 identification, **47**:197, 199
 malignant expansion, **47**:226
 morphology, **47**:214, 216
 reproduction, **47**:270
 surface phenotype, **47**:201, 205–207, 209–213
 tissue distribution, **47**:219, 220

- nonspecific, activation by Fc fragment of Ig, **40**:79–90
- opioid receptor, **48**:180
- of peripheral blood, IgA and, **40**:202–203
- phagocytosis and, **38**:364
- pituitary hormones, **63**:391–392, 405–407
- pulmonary, **59**:381–385
- RAG-2-deficient blastocyst complementation, **62**:31–41
- recirculation, **64**:139–143, 198
- genetic studies, **64**:184–188
- naive and memory lymphocytes, **64**:180–181
- through spleen, **64**:177–178
- regulation, **59**:176–178, 383–385
- resident pulmonary, CD4⁺8⁻ $\alpha\beta$ T cells, **58**:319
- responses, maternally transmitted antigen and, **38**:313
- retinal glial cell, **48**:218
- SCID and, **49**:386–388
- shedding, **64**:170
- as source of chemokines, **55**:106
- spontaneous autoimmune thyroiditis and altered thyroid function, **47**:456, 462
- cellular immune reactions, **47**:451, 453, 455
- disturbed immunoregulation, **47**:470–472, 474, 477, 481
- genetics, **47**:487
- potential effector mechanisms, **47**:466, 467, 469
- SP receptors, **39**:313
- stem cell factor and SCF receptor effects, **55**:48–51
- surface antigens of human leukocytes and, **49**:94, 113, 115
- T, *see* T cells
- T9 positive, in Crohn's disease, **42**:290
- T cell development and, **44**:213, 229, 236
- T cell subsets and, **41**:70, 81, 110
- TGF- β regulation, **55**:187
- thyroid-infiltrating, **47**:492
- effector mechanisms, **47**:466, 467, 469
- immunoregulation, **47**:473, 475
- tumor-infiltrating, *see* Tumor-infiltrating lymphocytes
- turnover and clonal deletion, physiology, **58**:270–272
- VIP receptors, **39**:312–313
- Lymphocyte tumors, **64**:179
- Lymphocytic choriomeningitis virus, **52**:316
- hybrid resistance and, **41**:355, 390
- infection, cytotoxic T cells and, **65**:278
- NK cells and, **47**:244, 245, 283, 284
- responses in MHC class I-deficient mouse, **55**:409–410
- clearance, **55**:411–412
- immunopathology, **55**:410–411
- synthetic T and B cell sites and, **45**:226
- Lymphohematopoiesis, stem cell factor and
- SCF receptor effects
- hematopoiesis, **55**:43–48
- lymphocytes, **55**:48–51
- NK cells, **55**:48–51
- Lymphohematopoietic cells, T cell subsets and, **41**:107–109
- Lymphohemopoietic tissue, organization, B cell formation and, **41**:185–188
- Lymphoid blast crisis, chronic myelogenous leukemia, **40**:290–291
- Lymphoid cells, **48**:69
- adoptive T cell therapy of tumors and, **49**:302
- autoimmune thyroiditis and cellular immune responses, **46**:287–291, 293, 295
- experimental models, **46**:274, 275, 279, 280, 282
- B cell formation and, **41**:196, 213, 220, 225, 226
- CD1 protein distribution, **59**:56
- CD5 B cell and, **47**:125
- effect of chromosomal abnormalities, **50**:132–140
- differentiation-related translocation oncogenes, **50**:139
- functional chimerism involving DNA binding and protein dimerization, **50**:133–138
- proteins affected by chromosome translocations, **50**:139–140
- genetically engineered antibody molecules and, **44**:71, 87
- HIV infection and, **47**:382, 388
- hybrid resistance and, **41**:333
- antibodies, **41**:376, 380
- antigen expression, **41**:401, 405

- Lymphoid cells (*continued*)
 bone marrow cells, **41**:338, 339
 leukemia/lymphoma cells, **41**:359, 369
 marrow graftment, **41**:386
 NK cells, **41**:374, 375
 normal hemopoietic cells, **41**:351–358
 T cells, **41**:380, 383
in vitro assays, **41**:393, 394
- IgE biosynthesis and, **47**:11
 Ig heavy-chain variable region genes and,
49:52–53, 55
 immunosenescence and, **46**:242–244, 246,
 247
 leukocyte integrins and, **46**:162, 168, 171
 mammalian, genetically engineered
 antibody molecules and, **44**:69, 70
 NK cells and, **47**:188, 221
 precursors, maturation, role of CD44,
58:377
 SCID and, **49**:393
 spontaneous autoimmune thyroiditis and,
47:435, 491
 cellular immune reactions, **47**:454
 disturbed immunoregulation, **47**:475,
 476, 481
 histopathology, **47**:440, 442, 443
 T cell subsets and, **41**:44, 78, 90, 96, 101,
 109
 Lymphoid hyperplasia, murine lupus models
 and, **46**:63
 Lymphoid organs
 antigen-presenting cells and, **47**:104
 APC-T cell binding, **47**:95
 immunogeneity, **47**:102, 103
 interaction with T cells, **47**:82–84
 tissue distribution, **47**:50–52
 CD5 B cell and, **47**:122
 deficient development of lymphocytes,
55:394
 humoral immune response and, **45**:79–82
 NK cells and, **47**:211, 266
 spontaneous autoimmune thyroiditis and
 cellular immune reactions, **47**:449, 450
 disturbed immunoregulation, **47**:477
 histopathology, **47**:444
 humoral immune reactions, **47**:446, 447
 potential effector mechanisms, **47**:469
 Lymphoid sheath, periaxial, *see*
 Periaxial lymphoid sheath
- Lymphoid stem cells, SCID and, **49**:384, 389,
 400–401
 absence, **49**:382, 390–391, 393
 Lymphoid tissues
 central role of FDCs in,
 antigen expression of, **51**:257–261
 cellular origin of, **51**:261–263
 conclusions, **51**:278–280
 distinction of from other dendritic cells,
 51:244–247
 expression of in malignant lymphomas,
 51:274–275
 process, **51**:243–244
 regulation of normal germinal center by,
 51:263–274
 as targets of HIV, **51**:275–278
 technical difficulties, solution of,
 51:246–257
 human, cell types and antigenic profiles,
52:129, 136
 immunosenescence and, **46**:236, 252, 253
Ir gene expression in, **38**:52–55
 mucosa-associated
 functional anatomy of Peyer's patches,
40:192–195
 studies on disassociated Peyer's patches
 cells, **40**:195–197
 tyrosine protein kinase, **48**:277–280
 Lymphoid tumors
 chromosomal translocation in
 development of T cell leukemia,
50:141–142
 effect of chromosomal abnormalities on
 adjacent oncogenes, **50**:124–132
 c-myc in Burkitt's lymphoma
 translocations, **50**:124–128
 gene fusion resulting from
 chromosome translocation,
50:128–129
 helix-loop-helix oncogenes in
 chromosome translocations,
50:129–130
 oncogene location after translocation,
50:131–132
 transcriptional disruption of LIM
 domain oncogenes by
 translocation, **50**:130–131
 effect of chromosomal abnormalities on
 lymphoid cells, **50**:132–140

- developmentally regulated translocation oncogenes, **50**:139
- functional chimerism involving DNA binding and protein dimerization, **50**:133–138
- other unusual proteins affected by chromosome translocations, **50**:139–140
- mechanism of translocation and inversion, **50**:121–122
- presence of abnormalities, **50**:119–121
- timing of chromosome translocation and inversion, **50**:122–124
- therapy, **58**:282
- Lymphokine-activated killer cells, **42**:182, 190; **47**:188, 259, 260, 296, *see also* Natural killer cells
- adoptive T cell therapy of tumors and, **49**:284–285, 302, 306
- antigen-nonspecific killing, **60**:302–303
- humoral immune response and, **45**:68, 70
- IBD and, **42**:296
- Lymphokine receptors, **59**:177
- Lymphokines, **42**:165–166
- adoptive T cell therapy of tumors and antigen recognition, **49**:319
- expression of antitumor responses, **49**:324, 328–329
- mechanisms, **49**:299–301, 307–309, 313, 316
- principles, **49**:289–290
- antigen-presenting cells and, **47**:75, 80, 87
- cell surface, **47**:63
- immunogeneity, **47**:103, 104
- T cells, **47**:83, 84, 99–101
- autoimmune demyelinating disease and, **49**:360
- autoimmune thyroiditis and, **46**:283, 289, 291, 294, 298
- autoreactive T cells and, **45**:430, 431, 433
- B cell formation and, **41**:199, 222, 231
- B cell growth and, **40**:16
- CD4 and CD8 molecules and, **44**:299
- cytolysis and, **41**:284, 286
- fresh tissues, **48**:95
- human T cell antigen and, **41**:1, 6, 20
- gene regulation, **41**:27, 29
- synergy, **41**:15
- humoral immune response and, **45**:5
- cellular interactions *in vivo*, **45**:85
- helper T cells, **45**:24–35
- interleukins, **45**:55, 62, 65, 67, 69, 71–76
- physical interaction, **45**:36, 39, 41, 46–54
- hybrid resistance and, **41**:407
- IgE biosynthesis and
- antibody response, **47**:7–11
- antibody response suppression, **47**:32, 33, 39
- binding factors, **47**:12, 22–28
- immunosenescence and, **46**:253
- lymphocyte activation, **46**:230–236
- lymphocyte subsets, **46**:237
- regulatory changes, **46**:250
- islet beta cell, **48**:141
- NK cells and, **42**:183; **47**:189, 289, 290, 293
- differentiation, **47**:233
- effector mechanisms, **47**:235, 262–266
- morphology, **47**:216
- reproduction, **47**:271
- tissue distribution, **47**:220
- production
- AB-Mul V-transformed mast cells, **53**:4–6
- factor-dependent mast cell lines, **53**:6–9
- human Fc ϵ R $^+$ cells, **53**:14–15
- murine Fc ϵ RI $^+$ cells, **53**:9–14
- regulators of complement activation and, **45**:389
- role in B cell activation, **40**:38–39
- spontaneous autoimmune thyroiditis and, **47**:472, 473, 477
- sporozoite malaria vaccine and, **45**:303, 304, 306, 312
- as substitutes for T cells, **40**:92
- surface antigens of human leukocytes and, **49**:101
- synthesis, **48**:78–81
- IL-2-induced, **48**:81
- transcription, **48**:81
- synthetic T and B cell sites and, **45**:219, 248
- T11, **41**:9, 10
- T cell development and, **44**:219, 221, 229
- T cell subsets and
- H-2 alloantigen recognition, **41**:94
- H-2 molecules in thymus, **41**:107
- H-2-restricted antigen recognition, **41**:52, 62–66, 76, 77
- T cell trigger, **41**:112

- Lympholysis, hybrid resistance and, **41:388**
- Lymphoma*, *see also specific types*
- adoptive T cell therapy of tumors and, **49:283, 286, 291–292, 295, 305**
 - avian leukosis virus-induced
 - myc* gene role, **56:472–476**
 - pathogenesis, **56:471–472**
 - resistance, **56:470**
 - susceptibility, **56:471**
 - B cell, *see* B cell lymphoma
 - B cell formation and, **41:202**
 - BSAP binding, **63:229**
 - Burkitt's, *see* Burkitt's lymphoma
 - CD1 proteins, **59:60**
 - CD5 B cell and, **47:123–125**
 - Ig gene expression, **47:150–152, 155, 156**
 - CD23 antigen and, **49:161, 164, 175**
 - CD40, **61:39**
 - cell-mediated killing and, **41:293**
 - chromosomal translocations, **38:247, 266**
 - c-myc* activation, **38:261**
 - development, and *bcl-2* gene
 - overexpression, **58:270**
 - follicular, **38:264, 267, 268, 271, 272**
 - Hodgkin's disease, **61:8, 12, 39**
 - human B cell neoplasia and, **38:245**
 - hybrid resistance and, **41:334, 358–369**
 - antigen expression, **41:401, 403**
 - NK cells, **41:373, 374**
 - Ig heavy-chain variable region genes and, **49:40, 51**
 - induction by MMTV, **65:155–156**
 - malignant, differential expression of FDCs in, **51:274**
 - NK cells and, **47:190, 228**
 - pituitary hormones, **63:388, 400, 427**
 - prolactin, **63:400**
 - prolactin receptor, **63:396**
 - risk in rheumatoid arthritis patient, **64:329–330**
 - SCID and, **49:393**
 - surface antigens of human leukocytes and, **49:114, 117**
 - T cell
 - depletion, **58:272–275**
 - murine, **54:289**
 - Lymphoma cells, lymphocyte homing and carbohydrate, **44:359, 361**
 - molecules, **44:343**
 - regional specificity, **44:324, 338**
- Lymphopenia
- chronic AIDS-associated, and lymphocyte PCD, **58:214**
 - and viral infection, **58:233–237**
- Lymphopoiesis
- B cell, in fetal and neonatal rabbit, **56:180–186**
 - B cell formation and, **41:213, 237, 238, 229–231**
 - hybrid resistance and, **41:339**
 - T cell, in liver, **53:169–171**
- Lymphoproliferation
- and autoimmunity, mechanism of beneficial effect of IL-2 on, **50:200–203**
 - B cell formation and, **41:193, 208**
 - defective mouse, CD5 B cell analysis, **55:314–315**
 - treatment, **58:273–275**
- Lymphoproliferation gene, murine lupus models and, **46:63, 94–98**
- Lymphoproliferative disease, **54:143–145**
- B cell lymphoma, **54:144–145**
 - CD23 antigen and, **49:175**
 - Epstein–Barr virus, **54:143–144**
 - in hu-PBL-SCID mice, EBV-associated, **50:310–313**
- Lymphoproliferative disorder, post-transplant, **54:35–36**
- Lymphoproliferative syndrome, X-linked, **59:248**
- Lymphosarcoma cells, hybrid resistance and, **41:358**
- Lymphotoxin, **42:216–217**
- adoptive T cell therapy of tumors and, **49:308**
 - autoimmune demyelinating disease and, **49:360**
 - B cell formation and, **41:236**
 - cytotoxicity and, **41:284–286, 298**
 - helper T cell cytokines and, **46:112**
 - functions, **46:114, 115, 117, 122**
 - immune responses, **46:133**
 - hybrid resistance and, **41:400, 407**
 - IgE biosynthesis and, **47:10**
 - IL-2 and, **44:153, 154, 185**
 - maps of Ig-like loci and, **46:30, 33, 34**
 - multiple myeloma and, **42:224**
 - NK cells and, **47:278, 279**
- Lymphotoxin- α , **61:5, 22, 24, 26**
- Lymphotoxin- β , **61:22, 23, 24**
- Lymphotoxin-like molecules, cytology and, **41:284–286**

- Lyn, **59**:324
 IL-2R β coupling, **61**:155
 in myeloid cells, **60**:209
- lyn*, in CD40-induced signaling, **60**:45–46
- Lysine, **42**:97
 cell-mediated killing and, **41**:297
 coupling to Boc, **60**:107
 Ig gene superfamily and, **44**:33
- Lyo-phosphatidylethanolamine, **49**:246, 248
- Lysophospholipase, CLC identification with, **39**:182–183
- Lysosomes
 antigen-presenting cells and, **47**:52
 autoimmune thyroiditis and, **46**:267, 268
 CAL and, **41**:168
 complement receptor 1 and, **46**:200
 cytotoxicity and, **41**:276, 281, 288
 humoral immune response and
 antigens, **45**:12–14, 20, 21
 cellular interactions *in vivo*, **45**:79
 IL-1 and, **44**:163, 184
 lymphocyte homing and, **44**:360
 NK cells and, **47**:214, 218, 225, 238
 synthetic T and B cell sites and, **45**:215
- Lysozyme
 antibody complex, **43**:22–23
 antigenic structure, **43**:118
 hen egg, T cell response, **43**:201
 H-Y antigen and, **38**:179
 HyHEL5 binding, **43**:12–13, 122
 mobility, **43**:21
 side-chain movement, **43**:22
 synthetic T and B cell sites and, **45**:204,
 211, 213, 214
- T cell receptor and
 antigen processing, **45**:122, 123
 experimental systems, **45**:131
 homogeneity, **45**:129
 peptides, **45**:133, 134
 structure-function relationships, **45**:135, 136
- T cell suppression and, **38**:139–142
 T cell-T cell interactions and, **38**:159, 160
- Lyt-1, **48**:246
- M**
- mAbs, *see* Monoclonal antibodies
- Mac-1, leukocyte integrins and, **46**:149–153,
 176
 chromosomal localization, **46**:160
 inflammation, **46**:163, 164, 166–168
- ligand molecules, **46**:173
 structure, **46**:153, 154, 156, 158, 159
- Mac-1/CR3, **54**:310
- Macaque, SIV pathogenesis
 genetic drift of SIV molecular clones *in vivo*, **52**:451–452
 immunodeficiency virus infection of,
52:443–452
 natural history of SIV infection of,
52:443–448
 use of SIV molecular clones to define viral determinants of pathogenesis,
52:448–451
- α_2 -Macroglobulin, **63**:278–279
 complement component and, **38**:234, 236
- Macrophage-activating factor, **48**:78;
62:61–62, 96
 adoptive T cell therapy of tumors and,
49:307, 309–311
- Macrophage colony-stimulating factor, **48**:71
- Macrophage inflammatory protein, **46**:115
- Macrophage inflammatory protein-1, **62**:260,
 270, 280, 285
- Macrophage inflammatory protein-1 α , mRNA expression in hypereosinophilic syndrome, **60**:200
- Macrophage inflammatory protein-2, **62**:274
- Macrophage migration inhibitory factor, **66**:197
 background, **66**:197–199
 history, **66**:198
 in IBD, **42**:290
 immune system and, **66**:218–219
 localization, **66**:202–206
 MIF gene, **66**:199–202, 206
 properties, **66**:199, 206–210
 sepsis and, **66**:104, 148–149
 structure-function studies, **66**:216–218
 surface antigens of human leukocytes and,
49:111
- Macrophages, **48**:69
 activation, **62**:78–80, 86, 277
 adoptive T cell therapy of tumors and,
49:287, 289–290, 333
 antigen recognition, **49**:319
 expression of antitumor responses,
49:325–328, 330
 mechanisms, **49**:300, 302, 306–313, 316
 alveolar, **59**:370, 374–375, 381
 human, iNOS in, **60**:353–354
 IL-1ra production, **54**:185

- Macrophages (*continued*)**
- antigen-presenting cells and, **47:68, 80, 104**
 - antigen processing, **47:89, 90**
 - cell surface, **47:61, 62**
 - immunogeneity, **47:103, 104**
 - interaction with T cells, **47:93, 94**
 - T cell, **47:86, 95, 97, 99**
 - tissue distribution, **47:48, 51, 54, 55**
 - antinuclear antibodies and, **44:134**
 - autoimmune thyroiditis and, **46:265, 287, 307**
 - B cell formation and, **41:186, 208, 234–236, 238**
 - B cell precursors, **41:191, 192, 194, 195**
 - bone marrow cultures, **41:212, 213, 218**
 - cytotoxicity and, **41:269, 285, 293**
 - genetically determined defects, **41:230–232**
 - hybrid resistance and, **41:340, 356, 366**
 - antigen expression, **41:402, 405**
 - effector mechanisms, **41:370–372, 374, 387, 392, 393**
 - and B cells, involvement of, **51:303–305**
 - cachectin and, **42:214, 215, 220, 221**
 - CD1 protein distribution, **59:66–67**
 - CD4 molecules and, **44:288, 290**
 - CD5 B cell and, **47:120–122, 132**
 - CD23 antigen and, **49:157, 160, 166**
 - biological activity, **49:167, 171**
 - cellular expression, **49:153–154**
 - expression in clinical conditions, **49:176**
 - colony-stimulating factor, **48:75**
 - comparison with B cells
 - antigen processing, **39:62–66**
 - antigen uptake, **39:60–62**
 - complement components and
 - bronchoalveolar, **38:218**
 - C2, **38:218**
 - C4, **38:220**
 - Factor B, **38:217, 218**
 - complement receptor I and, **46:190, 192, 200**
 - cosignals from, **58:243–245**
 - differentiation, IL-6 effects, **54:24–25**
 - effector cells in cell-mediated immune response, **60:333–341**
 - Fc receptors on, **40:64–66**
 - fibronectin and, **38:388**
 - GM-CSF production, **39:2, 3, 11**
 - growth hormone, **48:178**
 - helper T cell cytokines and
 - differential induction, **46:128–130**
 - functions, **46:114–117, 122**
 - immune responses, **46:133**
 - HIV infection and, **47:413**
 - activation, **47:400, 402**
 - immunopathogenic mechanism, **47:392–397**
 - neuropsychiatric manifestations, **47:404**
 - hybridomas and
 - B cells, **38:283**
 - genetically engineered antibodies, **38:306**
 - heterohybridoma antibodies, **38:289**
 - IBD and, **42:285, 287O, 292, 297**
 - IgE biosynthesis and
 - antibody response suppression, **47:30, 33, 36, 37**
 - binding factors, **47:20, 21, 27**
 - IGF-I, **63:389–391**
 - Ig gene superfamily and, **44:39**
 - IL-1 and, **44:153, 190**
 - biological effects, **44:169**
 - gene expression, **44:159–161, 164**
 - human disease, **44:192, 195**
 - immunocompetent cells, **44:173, 177**
 - structure, **44:155**
 - systemic effects, **44:171**
 - TNF, **44:185, 187**
 - IL-4, **48:75**
 - IL-10 biological effects, **56:6–8**
 - immunosenescence and, **46:228**
 - interaction with IgA, **40:170**
 - interstitial, **59:375**
 - in vitro*-derived, IL-1ra production, **54:185–186**
 - Ir* genes and
 - antigen binding, **38:98, 109, 111–115**
 - antigen processing, **38:93–97**
 - expression in
 - antigen-presenting cells, **38:55–57**
 - B cells, **38:57, 58, 189**
 - in lymphoid tissue, **38:55**
 - in T cells, **38:59–63**
 - H-Y antigen, **38:179**
 - Ia molecules, **38:68, 70, 83, 84**
 - MHC restriction, **38:64–67**
 - T cells
 - cytotoxic, **38:161, 162, 164**
 - interactions, **38:155 –157**

- selection, **38**:133, 134
 suppression, **38**:136–139
 tolerance, **38**:125, 126
 leukocyte integrins and, **46**:151, 154, 163,
 164, 167
 lymphocyte homing and, **44**:368
 lymphotoxin and, **42**:216
 and monocytes, **51**:337–339
 neuropeptide, **48**:177
 NK cells and **189**, **47**:288–290
 congenital defects, **47**:224
 differentiation, **47**:229
 effector mechanisms, **47**:235, 237, 262,
 263, 265, 266
 hematopoiesis, **47**:274, 278
 identification, **47**:296
 malignant expansion, **47**:228
 reproduction, **47**:270
 surface phenotype, **47**:201, 202, 211–213
 tissue distribution, **47**:220
 peritoneal, IL-1ra production, **54**:188
 PGHS-2, **62**:191–192
 phagocytosis and, **38**:361, 362, 369
 human alveolar, **38**:365
 noneelicited murine, **38**:376–380, 383,
 384
 rat alveolar, **38**:374
 pituitary hormones, **63**:411–412
 present at vaccinating site, **58**:431
 pulmonary, **59**:373–376
 interstitial, **59**:375
 lung transplants, **59**:423
 response to Fc fragments, **40**:80–83
 SCID and, **49**:384–385, 389–391, 401–402
 soluble factors, antibody responses and,
 40:35
 as source of chemokines, **55**:106
 spontaneous autoimmune thyroiditis and,
 47:492, 494
 disturbed immunoregulation, **47**:475, 476
 genetics, **47**:488
 histopathology, **47**:440, 442
 potential effector mechanisms, **47**:467,
 468
 stimulation by Fc fragments of Ig,
 40:100–111
 suppressor, **50**:176
 surface antigens of human leukocytes and,
 49:89, 91–92, 98–99, 116
 synovial, IL-1ra production, **54**:187–188
 T cell development and, **44**:248
 T cell stimulation, **39**:84
 T cell subsets and
 cell surface molecules, **41**:46
 H-2 alloantigen recognition, **41**:89,
 90, 94
 H-2 molecules in thymus, **41**:103–106,
 108–110
 H-2-restricted antigen recognition,
 41:51, 69–71, 74, 75, 77
 TGF- β regulation, **55**:187
 Macrophages/dendritic cells, antigen
 presentation, **39**:85–86
 MAdCAM-1, **62**:101, 257; **64**:188
 lymphocyte–HEV interaction, **65**:352, 359,
 362–363, 367, 370, 377, 379
 lymphocyte homing, **64**:147, 158–159
 to thymus and bone marrow, **64**:177
 in mucosal lymphocyte homing, **64**:176
 ontogenesis, **64**:164
 Maedi-visna virus, **52**:431
 MAGE, **62**:228, 229, 231–232
 MAGE-3, **62**:232
 Magnesium
 cytotoxicity and, **41**:271, 294
 genetically engineered antibody molecules
 and, **44**:85
 leukocyte integrins and, **46**:150, 156, 157,
 163
 lymphocyte homing and, **44**:351, 360, 361
 NK cells and, **47**:249, 250
 MAH cell line, LIF/CNTF receptors,
 53:43–44
 MAIDS, *see* Murine acquired
 immunodeficiency syndrome
 Main-chain flexibility parameters, Ab binding
 to protein antigens, **43**:50
 Main immunogenic region, AChR, antibodies
 to, **42**:240–241, 262
 specificity and, **42**:246
 Major basic protein
 antihelminthic activity, **60**:212–213
 basophil, **39**:186–187
 cell-mediated killing and, **41**:311, 312
 cytotoxicity for mammalian cells,
 60:191–193
 eosinophil
 biochemical properties, **39**:185
 bronchial epithelium damage by,
 39:219–223

- Major basic protein (*continued*)
granule core, **39**:185–186
histamine release stimulation, **39**:191
parasite killing, **39**:188–190, 209,
 212–213
toxicity to various cells, **39**:190–191, 219
- HL-60 promyelocyte, **39**:187–188
- hypereosinophilic syndrome, **60**:160
- placenta, **39**:188, 228
- tissue from asthma deaths, **60**:217
- Major histocompatibility complex, **48**:78, 107
- ABC protein
 function, **52**:38–42
 genes, **52**:35–38
 restriction, history of discovery of,
 52:4–6
- adoptive T cell therapy of tumors and,
 49:281, 283, 294–295, 334
- antigen recognition, **49**:319, 322–323
- expression, **49**:324–325, 327–330, 332
- mechanisms, **49**:299, 315
- allele, **43**:219
- antigen-presenting cells and, **47**:45, 47, 87,
 104, 105
- antigen presentation, **47**:64, 66–69,
 71–73, 75–80
- antigen processing, **47**:87–89, 91, 92
- APC-T cell binding, **47**:95–97, 99
- cell surface, **47**:57–60
- immunogeneity, **47**:103
- T cells, **47**:82, 83, 85, 101
- tissue distribution, **47**:51–54
- autoimmune demyelinating disease and,
 49:357–358
- multiple sclerosis, **49**:369–371, 374–375
- TCR usage restriction, **49**:363, 365–367
- autoimmune thyroiditis and, **46**:263, 265,
 266, 317, 319
- antigens, **46**:268
- cellular immune responses, **46**:289, 290,
 295
- experimental models, **46**:275, 279, 280
- genetic control, **46**:282–286
- humoral responses, **46**:299, 301, 305,
 307
- prevention, **46**:310, 315
- autoreactive T cells and, **45**:417, 418, 433
- activation, **45**:421, 423, 427
- origin, **45**:418–420
- physiology, **45**:428, 430
- regulation, **45**:432
- B cell formation and, **41**:194, 197
- CD4 and CD8 molecules and, **44**:265–267,
 297, 299, 300, 302, 303
- CD4 molecular biology, **44**:291–293,
 296
- CD8 molecular biology, **44**:271, 272,
 282, 284
- CD5 B cell and, **47**:137, 142, 160
- CD23 antigen and, **49**:150–151, 169
- class I, **43**:213–214
- binding process, **43**:214
- determinant nature, **43**:205–206
- class II alleles, RA susceptibility associated
 with
- ethnic differences, **56**:409–410
- penetrance influenced by, **56**:440–441
- properties, **56**:444–446
- risk for severe disease, **56**:441–444
- class II α chain, **48**:141–142
- class II β chain, **48**:141–142
- class II loci, corneal graft, **48**:199–200
- latex bead treatment, **48**:200
- complement components and, **38**:203, 207,
 236
- C2 and, **38**:208
- C4 deficiency, **38**:229
- C4 linkage to, **38**:220, 221
- class III region, **38**:227–229
- Factor B, **38**:209
- complement receptor 1 and, **46**:189
- contact amino acids, **43**:210
- control of anticolon type II response,
 48:130–131
- cytotoxicity and, **41**:270
- electron density in cleft, **43**:214
- helper T cell cytokines and, **46**:112, 117,
 118
- HIV and, **45**:349, 361, 365–367
- HIV infection and, **47**:380, 411
- immunopathogenic mechanism, **47**:386,
 389, 391, 392
- human T cell activation and, **41**:2, 3, 5, 14
- humoral immune response and, **45**:5–8, 80
- antigens, **45**:8, 10, 19, 20
- helper T cells, **45**:25, 26
- interleukins, **45**:62, 66, 67
- physical interaction, **45**:37, 41, 43,
 46–48, 52–54
- hybrid resistance and, **41**:348, 357, 361,
 369, 378, 403

- IgE biosynthesis and
 antibody response, **47:2**, 29, 33
 binding factors, **47:25**, 27
- Ig gene superfamily and
 gene organization, **44:24**–29
 receptors, **44:9**–12
- immunosenescence and, **46:239**
- Ir* genes and, **38:33**, 43–46, 188, 190
 adoptive transfer, **38:52**, 53
 antigen binding, **38:98**, 99, 105, 114
 antigen processing, **38:94**, 97
 B cell response, **38:51**
 complementation, **38:72**–83
 cross-reactive lysis, **38:186**
 expression, **38:57**, 59, 63
 gene function, **38:91**
 H-Y antigen, **38:175**, 180
 Ia molecules, **38:170**, 171
 immunodominance, **38:180**, 181, 185
 I region mutations, **38:84**, 85, 87
 restriction, *see* Major histocompatibility
 complex restriction
 self, *see* Self-antigens, self-MHC
- T cells
 cytotoxic, **38:161**, 163, 164, 167–169,
 173
 interaction, **38:153**, 154, 157, 159, 160
 positive selection, **38:130**, 131
 repertoire, **38:118**, 144, 145
 suppression, **38:138**, 147, 151
 tolerance, **38:121**, 122, 124, 125, 127,
 129
- linked diabetogenic genes, **51:290**–293
- locus, MMTV-induced tumors and, **65:156**
- lymphocyte homing and, **44:340**, 350
- maps of Ig-like loci and, **46:1**, 28–34, 46
 structure, **46:19**
- T cell receptor, **46:34**
- maternally transmitted antigen and, **38:313**,
 315, 318, 351, 352
 class I genes, **38:329**, 330
 mitochondrial transmission, **38:347**
- measles and, **45:345**, 346
- membrane proximal and distal domains,
43:214
- murine lupus models and, **46:65**, 79, 81,
 82, 85, 86, 90, 91, 94
- NK cells and, **47:187**, 188, 292
 cytotoxicity, **47:190**
- effector mechanisms, **47:239**, 240, 242,
 257, 259, 261
 surface phenotype, **47:201**, 211
- recognition, **43:194**–196, 224
- regulators of complement activation and,
45:404
- restricted antigen recognition, AChR and,
42:256–259
- restriction, *see* Major histocompatibility
 complex restriction
- SCID and, **49:384**, 386–387, 391
- self-encoded, **43:213**
- spontaneous autoimmune thyroiditis and,
47:435, 491, 493, 494
 breeding, **47:436**–438
 effector mechanisms, **47:467**, 469
 function, **47:459**, 461, 462, 464
 genetics, **47:481**–491
 immune reactions, **47:450**, 455
 immunoregulation, **47:470**–472, 476,
 477
- sporozoite malaria vaccine and
 CS-specific T cells, **45:307**, 310
 immunity, **45:287**
 interferon- γ , **45:304**, 305
 structure, **66:68**–69
- surface antigens of human leukocytes and,
49:78–79
- T cell development and, **44:207**
 antigen recognition, **44:209**, 210,
 212–214
- cellular selection within thymus,
44:245–254
- ontogeny, **44:221**
- T cell receptor and, *see* T cell receptor
- T cell subsets and, **41:39**, 71, 78, 81, 101,
 108
 cell surface molecules, **41:43**, 44, 50
 transgenic, transcriptional factor, **48:142**
- Major histocompatibility complex antigens,
59:1–2, 99–100, 112, 120; **61:327**–338
- alloantigens, B cell repertoire expression
 and, **42:78**–79
- binding in solution, **43:211**–213
- blocking by analogs in trimolecular
 complex, **56:244**–247
- CD4-CD8 coreceptor/MHC binding sites,
53:68–70
- class I, **64:2**, 31; **65:48**–49
 assembly, **52:42**–76

- Major histocompatibility complex antigens
(continued)
- assembly, subcellular location, **52:42–50**
 - assembly deficient cells, **52:47–49**
 - class I molecules associated with antigen, place of, **52:49–50**
 - effects of adenovirus E3/19K glycoprotein on antigen processing and presentation, **52:44–47**
 - effects of brefeldin A on antigen processing and presentation, **52:42–44**
 - assembly and transport, posttranslational modifications of, **52:59–62**
 - glycosylation, **52:59–61**
 - palmitylation, **52:61–62**
 - phosphorylation, **52:62**
 - calnexin, **64:107–112, 119–121**
 - cell surface-associated, mechanism of peptide binding to, **52:81–84**
 - cytotoxic T cells and, *see* Cytotoxic T cells
 - deficient mouse, *see* Mouse
 - entry into processing pathways, **52:13–14**
 - exogenous β_2m association with, **52:86–87**
 - exogenous peptide association with, site of, **52:79–81**
 - export to cell surface, control of, **52:62–76**
 - general considerations, **52:62–63**
 - induction of class I transport by exogenous peptides or hypothermia, **52:69–73**
 - retention of β_2m – α chain complexes, **52:65–68**
 - retention of free α chains, **52:63–64**
 - summary of cellular control mechanisms, **52:73–74**
 - viral subterfuges, **52:74–76**
 - folding and assembling of, **52:50–52**
 - generation, model, **64:26**
 - induction of transport by exogenous peptides or hypothermia, **52:69–73**
 - interaction of exogenous peptides with, **52:78–86**
 - internalization, **52:87–92**
 - evidence for internalization, **52:87–89**
 - internalization signals, **52:89–92**
 - phosphorylation involvement, **52:90–92**
 - region of proteins, **52:89–90**
 - Ir* genes and, **38:163, 164**
 - cross-reactive specificity, **38:185, 186**
 - cytotoxic T cells, **38:168–171, 173, 174**
 - epistasis, **38:176, 180**
 - immunodominance, **38:185**
 - maternally transmitted antigen and, **38:330–337, 348, 350**
 - membrane proximal domains, **43:214**
 - β_2 -microglobulin, **58:380**
 - molecular structure, **64:24**
 - murine NK cell receptors for, **55:364, 366**
 - nature of antigens bound to, **52:8–10**
 - NK cell receptors for, **55:356, 358–362, 372**
 - NK cell recognition, **55:363, 365**
 - in NK cell-specific functions, **55:349–353**
 - peptide anchoring, **64:124–127**
 - peptide building, **64:114–115, 118, 127–128**
 - peptide loading, **64:105–128**
 - properties, **52:78–92**
 - association of exogenous β_2M with class I molecules, **52:86–87**
 - extracellular processing of synthetic peptides, **52:84–85**
 - interaction of exogenous peptides with class I molecules, **52:78–86**
 - mechanism of peptide binding to cell surface-associated class I molecules, **52:81–84**
 - physiological relevance of exogenous peptide association, **52:85–86**
 - site of exogenous peptide association with class I molecules, **52:79–81**
 - proteasomes, **64:3, 105–107**
 - in proteolysis, role of, **52:29–30**
 - restoration by single peptides, **58:162**
 - role in proteolysis, **52:29–30**
 - self-, **57:314–320**
 - metastatic phenotype affected by, **57:318–320**
 - nonimmunological effects on tumor cells, **57:317–318**
 - recognition by cytotoxic T cell effector cells, **57:314–315**

- recognition by natural killer effector cells, **57**:315–317
TAP and, **65**:87–92
TAP system, **64**:3–5, 30–31, 112–121
T cell recognition, **55**:364
tumor immunogenicity affected by low level of expression, **57**:303–305
class Ia, as trimolecular complex target, **56**:247–250
class I and class II, **52**:364
and $\alpha\beta$ T cells, **58**:107–112
deficient mouse, **58**:156
and $\gamma\delta$ T cell function, **58**:103–104
and $\gamma\delta$ T cell reactivities, **58**:323–324
interaction with CD4 and CD8 coreceptors, **58**:129–131
ligation of $\alpha\beta$ TCR and coreceptor, **58**:159
peptide binding for T cell recognition, **58**:88
as potential tumor antigens, **58**:420–421
class Ib, **52**:92–104
characteristics of individual class Ib products, **52**:94–104
Hmt, **52**:101–104
Q10, **52**:94–95
Qa-2, **52**:95–98
TL region gene products, **52**:98–101
general aspects of structure and function, **52**:92–94
class II, **62**:80; **64**:2
appropriate, expression, **51**:308–310
on B cells, MLR and, **39**:81–83
corneal graft
cell-surface expression, **48**:199
expression, **48**:199
expression timing, **48**:199–200
deficiency, **61**:327–328
CIITA, **61**:331–334, 336–338
RFX, **61**:334–336
deficient mouse, *see* Mouse
HLA types, HIV and, **65**:286
incompetent, **50**:22
Ir genes and, **38**:168–171, 189
MMTV superantigens and, **65**:164–165
MUC-1 mucin, **62**:229, 238–239
mucins, **62**:222–223, 238–239
MUM-1, **62**:229, 237
mutated antigens, **62**:236–237
mutated oncogene products, **62**:226–227
oncogenic proteins, **62**:237–238
p53, **62**:226–227, 237–238
Pmel17, **62**:229, 235
prostate-specific antigen, **62**:224–225
protein antigens, **62**:221–225
RA susceptibility associated with alleles, **56**:429–430
DR4, **56**:404–408, 420–426
Dw determinants, **56**:407–408
serologic identification, **56**:402–404
restricted antigen presentation, **58**:319
restricted CTLs, **60**:303–310
restricted soluble, **60**:306–308
self-, tumor immune response to increased expression, **57**:313
transducing signals, **48**:142
tyrosinase, **62**:233–234
viral antigens, **62**:239–240
class II-containing vesicle, **64**:2
conformation, **43**:203–205
cornea, **48**:193–194
determinants, polymorphic, **58**:304
expression on vascular endothelium, regulation, **50**:262–265
gene expression, **61**:328–329
in graft-versus-host disease, rat, **39**:280–281
T cell reaction, **39**:280–281
and immunological interactions of T cells with vascular endothelium, **50**:261–266
and positive selection, **58**:164
presentation, **58**:330–331
RA susceptibility associated with concepts, **56**:390–397
determinants, **56**:400–402
DR1, **56**:420–426
DR6, **56**:419–420, 426
DR10, **56**:424–427, 429
function influenced by structure, **56**:451
function of molecules, **56**:449–450
genetic studies, **56**:436–438
mapping region, **56**:418–419
polymorphism, **56**:398–399
terminology, **56**:390–397
role in $V\beta$ selective element action, **50**:17–22
I-A molecules, **50**:20–21
I-E molecules, **50**:18–19

- Major histocompatibility complex antigens (*continued*)
 incompetent MHC class II molecules, **50:22**
 superantigen binding, **54:117–120**
 T cell receptor and, **61:148**
 three-dimensional structure, **43:213–215**
 tumor immune response to increased expression of allo-MHC, **57:312–313**
- Major histocompatibility complex glycoprotein, surface antigens of human leukocytes and, **49:80–82, 125**
 adhesion, **49:82–83**
 receptors, **49:77–78, 94, 100**
- Major histocompatibility complex restriction determinant selection model, **43:207**
- Ir* genes and
 antigen binding, **38:98**
 epistasis, **38:175**
 gene control, **38:63–67**
 gene function, **38:91, 92**
 Ia molecules, **38:71, 76**
 I region mutations, **38:88–90**
 macrophages, **38:112**
 T cells
 interactions, **38:157–160**
 selection, **38:131, 133**
 tolerance, **38:120, 125–129**
- Malaria
 cachectin and, **42:220, 226**
 human, **60:112–124**
 regulators of complement activation and, **45:389**
 rodent, **60:108–112**
 synthetic T and B cell sites and
 globular protein antigens, **45:207**
 parasitic antigens, **45:228, 229**
 vaccine, **45:254**
 viral antigens, **45:222**
- T cell receptor and, **45:132, 138**
 vaccine, *see* Sporozoite malaria vaccine
- Malignancy, cachectin and, **42:213, see also Cachectin**
- Malignant histiocytosis, complicating celiac disease, gene rearrangements and, **40:292–293**
- Malignant lymphomas, expression of FDCs in, **51:274–275**
- Malnutrition, IBD and, **42:289**
- Maloney sarcoma virus, **38:163**
- MALT, **65:251**
- Mammalian TOR, **61:181, 188–190**
- Mammary gland, lymphocyte homing and high endothelial venules, **44:321**
 molecules, **44:353**
 regional specificity, **44:326, 327, 335**
- Mammary gland factor, Stat5, **60:21**
- Mammary tumors, in mice, *see* Mouse mammary tumor virus
- Mammary tumor virus
 in B cells with Vbse expression, **50:39**
 superantigens, **54:102**
 transcripts expressed in activated B cells, **50:40**
- Mangano-heme PGHS-1, **62:188**
- α -Mannan, phagocytosis of zymosan by human monocytes, **38:365, 367, 368, 374–376**
 by nonelicited murine macrophages, **38:377**
- Mannose
 CD23 antigen and, **49:163, 165**
 complement receptor 2 and, **46:202, 204**
 in GPI-anchored proteins, **60:60–74**
 leukocyte integrins and, **46:154**
 surface antigens of human leukocytes and, **49:98**
- MAPK, *see* MAP kinase
- MAP kinase, **63:148–149, 166**
 activation, **60:13–14**
 in B cell activation, **55:244–245**
 signal transduction, **59:262–263, 341**
- MARE-I, **43:279**
- Marek's disease virus, induced tumors, **50:105–107**
- Marginal zone B cells, **59:284–285**
- Marker density, immunosenescence and, **46:237, 239**
- Markers, lineage-related, differential expression during development (murine B cells), **53:125–127**
- Marrow
 HLA-incompatible, engraftment of, **40:386–387**
 microenvironment, hybrid resistance and, **41:390–393**
 T cell-depleted, rejection by patients, **40:389–391**
- Marrow grafts
 hybrid resistance and
 hemopoietic cells, **41:386–388**
 specific unresponsiveness, **41:384–386**

- immune reconstitution and, **40**:410–411
 cytotoxic effector cells, accessory cells
 and neutrophils, **40**:416–417
 effects of T cell depletion, **40**:417–419
 humoral immunity, **40**:414–416
 phenotype and function of peripheral T
 cells, **40**:412–414
 repopulation of lymph nodes and
 thymus, **40**:411–412
 reconsideration of failure in humans,
40:392–395
- MART-1, **62**:229, 235
- Masking, in regulation of CD44 receptors,
54:317–318
- Mast cell growth enhancing activity, **54**:93
- Mast cell leukemia, etiology, stem cell factor
 role, **55**:74
- Mast cells, **48**:176, 177; **51**:327–331
 AB-Mul V-transformed, lymphokine
 production (murine), **53**:4–6
 and basophils, human, cell surface
 structure on
 adhesion receptors and recognition
 molecules, **52**:357–366
 integrins, **52**:358–361
 other recognition molecules, **52**:
 366
 recognition molecules of Ig supergene
 family, **52**:361–364
 selectins and related recognition
 molecules, **52**:364–366
- cells, **52**:333–335
 cell surface typing with mAbs,
52:335–339
- complementing binder sites, **52**:
 375–377
- conclusions, **52**:404
- IgE/Fc ϵ RI, **52**:378–381
 assembly, **52**:381–382
 cell surface structures functionally
 associated with Fc ϵ RI molecules,
52:391–396
 control of synthesis and expression of
 Fc ϵ RI molecules, **52**:384–386
 functional characterization of receptor,
52:387–391
 sequence and structural homologies,
52:383–384
 topology of, **52**:378
- Ig receptors, **52**:378–396
- negative regulators of growth and
 differentiations of human basophils
 and mast cells, **52**:355–357
 interferons, **52**:355–356
 transforming growth factors,
52:356–357
- receptors for activating peptides,
52:366–375
 formyl-methionine peptides and
 related compounds, **52**:371–372
 IL-8 and related integrins, **52**:367–371
 substance P and other neuropeptides,
52:372–375
- receptors for growth and differentiating
 factors, **52**:339–355
 hemopoietic growth factor receptor
 superfamily, HRS, **52**:340–351
- receptors for low-molecular-weight
 regulators and pharmacological
 compounds, **52**:396–404
 adenosine, **52**:396–398
 antiinflammatory drugs, **52**:402–404
 arachidonic acid metabolites,
52:398–402
- RTK family, c-kit, and related
 oncogenes, **52**:351–355
- bone marrow-derived, **53**:3
- cachectin, **42**:220
- CD40-L expression, **61**:20
- characterization, **53**:2–3
- chromolyn binding sites, **52**:392–393
- connective type, **53**:2–3
- and cytokines, **51**:328–331
- differentiation/dedifferentiation, **53**:3
- ECF-A production, **39**:214
- factor-dependent cell lines, lymphokine
 production (murine), **53**:6–9
- Fc receptors, **40**:68
- granules, EPO binding to, **39**:201–202
- helper T cell cytokines and, **46**:113, 114,
 116, 117
- histamine release
 major basic protein and, **39**:191
 SP effect, **39**:305–306
- IgE receptor/Fc ϵ RI, numbers and binding
 constants on primary cells,
52:386–387
- IL-3 effects, **39**:19–20
- IL-3 production, **39**:17
- and IL-9, **54**:93–95

- Mast cells (*continued*)
 IL-10 biological effects, **56:10**
 lung, IL-4 storage by, **53:15**
 mucosal, **53:2–3**
 neuropeptide production, **39:316–317**
 phenotypic properties, **53:1–2**
 precursors, **53:2**
 properties, **53:1–2**
 role in inflammatory reactions, **53:2**
 secretions, **53:2**
 stem cell factor effects
 development, **55:52–60**
 function, **55:60–64**
 secretion, **55:43**
 Substance P, **61:372**
 TH2 type cells, **61:360–361**
 triggered by oxidative events, **60:331–332**
 Mast cell-specific receptor, **43:282–283**
 β subunit, hydrophobicity profile, **43:288**
 characteristics, **43:283**
 gene cloning studies, **43:283–289**
 α chain, **43:284–287**
 β and γ chains, **43:287–289**
 mechanisms of action, **43:289–298**
 aggregation of receptors, **43:289–291**
 cytoplasmic calcium rise, **43:293–298**
 intrinsic and extrinsic, **43:289**
 phosphoinositide hydrolysis, **43:290–293**
 Mast cell tumors, hybrid resistance and, **41:361**
 Mastocytoma, anterior chamber, **48:213–215**
 Mastocytosis, *Nipponstrongylus*-induced, **54:95**
 Maternal effect, maternally transmitted antigen and, **38:321, 323, 339, 343**
 Maternally transmitted antigen, **38:313–317, 351, 352**
 class I antigens and, **38:330, 331, 336, 337**
 CTL generation, **38:348–350**
 expression, **38:331, 336, 344, 345, 351, 352**
 gene activation, **38:342**
 genetic mechanism models, **38:338–340**
 Delbrück model of antigenic variation, **38:340–344**
 mitochondrial transmission, **38:344–348**
 MHC, class I genes and, **38:329, 330**
 phylogenetic considerations, **38:337, 338**
 polymorphisms, **38:314, 318–329, 336, 337**
 repressors, **38:341, 342**
 Maternally transmitted factor, **38:321, 323, 327, 329, 351, 352**
 class I antigens, **38:337**
 CTL responses, **38:328**
 genetic mechanism models, **38:339, 340, 345–348**
 hybridization, **38:324**
 phylogenetic, **38:338**
 Maternal transmission, maternally transmitted antigen and, **38:321, 322, 326, 343, 352**
 Matrix metalloproteinases, rheumatoid arthritis, **64:297, 320–321**
 Maturation
 eosinophils, **60:162–164**
 functional, vs commitment to T cell lineage, **51:162–164**
 phenotypic, sequence, **58:141**
 post-thymic, T cells, **51:160–162**
 stem cell factor receptor-ligand interaction in, **55:40–42**
 Maturation arrest, **52:312**
 Max, **63:161**
 MB-1 molecule
 induction of calcium mobilization in early B lineage cells, **54:354**
 signal transduction through, **54:351–355**
 MC1, rheumatoid arthritis, **48:128**
 M cells, **65:251**
 Meg, and Ig flexibility, **48:21**
 MCP, *see* Membrane cofactor protein
 McPC603
 binding site, **43:9, 19**
 electrostatic complementarity with phosphorycholine, **43:10**
 variable domains, **43:18–19**
 ME491 (CD63), **52:393**
 Mean fluorescence intensity, immunosenescence and, **46:222**
 Measles
 antibodies from combinatorial libraries, **57:227–228**
 infection, membrane cofactor protein, **61:220–221**
 T cell receptor and, **45:124, 127**
 virus-induced immunosuppression and, **45:335–347, 358**
 Measles virus, hybridomas against, **38:294**
 MECA79, **64:149, 174; 65:361–363, 365**
 MECA367, **65:361, 362**

- M-ECEF, *see* Monocyte-derived eosinophil cytotoxicity enhancing factor
- MECLI*, **64**:16, 20
- MECL1, proteasome subunits, **64**:16–18
- Meclofenamate, **62**:184, 186
- Medulloblastoma, *pax5*, **63**:229–230
- Megakaryocytes, IL-6 effects, **54**:23–24
- Megakaryopoiesis, LIF effects, **53**:36
- MEK kinase, **63**:149
- MEL-14, **64**:142–143; **65**:352, 354
- lymphocyte homing and carbohydrate, **44**:360–362, 370
 - molecules, **44**:342–345, 347, 348, 350–352
 - regional specificity, **44**:341
- MELAN-A, **62**:229, 235
- Melanocytes, stem cell factor effects, **55**:64–66
- Melanocytic differentiation antigens, **62**:233–236
- Melanoma
- adoptive T cell therapy of tumors and, **49**:285–286, 288, 290, 305
 - etiology, stem cell factor role, **55**:74–76
 - structural and functional properties of associated gangliosides, **40**:357–364
- Melanoma antigens, **62**:219–220, 228, 230, 231–236
- Melanoma-derived lipoprotein lipase inhibitor, *see* Leukemia inhibitory factor
- α -Melanotropin, *in vitro* IgE synthesis, **61**:371–372
- Melittin, pore formation and, **41**:313–316
- Melphalan, chemotherapeutic use in cancer treatment, **56**:329–331
- Membrane attack complex
- amphiphilic nature of C5b-9, **41**:299–304
 - analogues, **41**:307–311
 - and C3 and C5 convertases, **60**:75–76
 - complement activation role, **56**:269–270
 - deposited CD59, **61**:251–254
 - fluid phase
 - clusterin, **61**:249–251
 - S protein, **61**:247–249
 - homologous restriction factor, **61**:254–255, 257
 - regulation, **61**:204
 - subunit composition, **41**:304–307
- Membrane-binding domain, PGHS, **62**:172–173
- Membrane cofactor protein, **61**:214–221
- biosynthesis and tissue distribution, **61**:216–219
 - complement receptor 1 and, **46**:189, 197
 - function, **61**:218–221
 - AIDS, **61**:221
 - measles, **61**:220–221
 - reproduction, **61**:219–220
 - Streptococcus*, **61**:221
 - transplantation, **61**:220
 - inhibition of C3 and C5 convertases, **60**:75–76
 - precursor, **61**:216
 - regulators of complement activation and genes, **45**:403, 404, 408
 - protein expression, **45**:395–398, 400–402
 - protein interactions, **45**:383
 - protein roles, **45**:386, 387, 390
 - RCA-like protein utilization, **45**:410
 - short consensus repeat, **45**:391
 - structure, **61**:214–216
- Membrane cofactor protein 1, **62**:80, 282, **64**:154, 318
- Membrane damage
- cytolysis and, **41**:277–281
 - pore formation and cytolytic proteins, **41**:316–319
 - small peptides, **41**:313–316
- Membrane enzymes, surface antigens of human leukocytes and, **49**:111–114
- Membrane proteins
- aPL antibodies and, **49**:197
 - Apo-1/Fas, membrane-spanning induction of PCD, **58**:135–136
 - and negative selection, **58**:172–173
 - role in T cell death, **58**:226
 - CD23 antigen and, **49**:164
 - IFN- γ , **62**:66, 77–78
 - surface antigens of human leukocytes and, **49**:93–97, 111, 114, 117
 - T3, **43**:116
- Membranes
- IgM receptor expression by Ig gene transfection, **54**:357–359
 - IgR/antigen complex processing, **54**:348
 - Ig receptors, **63**:43
 - integral proteins of, **52**:144, 145
 - and secreted Ig molecules, **54**:342–343
- Membranoproliferative glomerulonephritis, **61**:245

- Memory, *see also* Immunological memory
 humoral, germinal center B cells, **60:283**
- Memory B cells, **59:340–341, 393, 412**
 adhesion molecules affecting migration,
53:248–249
 affinity maturation, **53:233–234**
 functional properties, **53:236**
 generation, alternative models for,
53:236–238
 generation, alterativemodels for,
53:236–238
 germinal centers and, **53:231–233**
 life span, **53:240–241**
 migration, **60:274–277**
 adhesion molecules and, **53:248–249**
 basic pattern, **53:247–248**
 migration pattern, **53:247–248**
 pathways of differentiation, **53:232–233**
 phenotype of, **53:234–235**
 Ig isotypes, **53:235**
 surface markers, **53:235–236**
 switched, **59:340–341**
- Memory cells
 formation, CD40 and, **63:70–72, 73**
 immunosenescence and, **46:246, 247**
 lymphocyte homing and, **44:314, 325,**
330–334, 337
- Memory phenotype, peripheral T cells
 expressing, **58:216**
- Memory T cells
 CD44 expression, **54:294–295**
 differentiation after antigenic stimulation,
53:229–230
 functional differences with naive and
 effector cells, **53:227–228**
 maturation into, **53:229–230**
 migration
 adhesion molecules affecting,
53:248–249
 basic pattern, **53:242–244**
 to inflammatory sites, endothelium and,
50:278–284
 tissue-specific, **53:244–247**
 phenotypic identification with CD45,
53:224–227
 TCR and signal transduction, **53:228–229**
- Mendelian inheritance, maternally
 transmitted antigen and, **38:338**
- 6-Mercaptopurine, NK activity and, **42:300**
- Mesangial cells, as source of chemokines,
55:111
- Mesangial proliferative glomerulonephritis,
 and IL-6, **54:41**
- Mesenteric lymph node cells, IgE
 biosynthesis and
 antibody response, **47:3–5**
 binding factors, **47:12, 14, 16, 17**
- Mesenteric lymph nodes, lymphocyte homing
 and
 carbohydrate, **44:351, 353**
 regional specificity, **44:323, 326–328, 333,**
335–338
- Mesenteric microvascular models, anti-CD18
 effect, **60:329–330**
- Messenger RNA
 antinuclear antibodies and, **44:94**
 autoantibodies, **44:137**
 autoantigen, **44:129**
 autoimmune response, **44:131, 134**
 scleroderma, **44:122**
 SLE, **44:101**
 autoimmune demyelinating disease and,
49:360
 autoimmune thyroiditis and, **46:273, 294,**
295, 297
- cachectin and, **42:215, 221–222, 223**
- CD4 and CD8 molecules and, **44:298**
 CD4 molecular biology, **44:288–291,**
293, 296, 297
 CD8 molecular biology, **44:271, 272,**
274–278, 280
- CD5 B cell and, **47:147**
- CD23 antigen and
 biological activity, **49:170**
 cellular expression, **49:152–155**
 cleavage regulation, **49:158**
 expression in clinical conditions,
49:174–175
 expression regulation, **49:158–160**
 Fc ϵ RII, **49:162, 166**
- CD40L, **60:48**
- complement components and
 C1q, **38:232**
 C2, **38:212, 217**
 C4, **38:223, 226**
 Factor B, **38:209, 211, 212, 218**
- complement receptor 1 and 184, **46:193**
- complement receptor 2 and, **46:202**
- CTL and, **41:148**

- cytokines, **60**:222
genetically engineered antibody molecules and, **44**:73
GM-CSF, induction in T cells, **39**:14
helper T cell cytokines and, **46**:112, 115, 138
HIV infection and, **47**:380, 381, 392, 397
humoral immune response and, **45**:40, 62
Ig, **43**:239
IgE biosynthesis and, **47**:16
Ig gene superfamily and, **44**:18
Ig heavy-chain variable region genes and, **49**:2, 55
IL-1 and, **44**:174, 195
 biological effects, **44**:165, 167
 gene expression, **44**:159–161, 163, 164
 structure, **44**:155, 156
IL-2, induction in T cells, **39**:13–14
IL-3, induction in T cells, 14
IL-5, **57**:156–159; **60**:198–199
IL-5 receptor, **60**:184
immunosenescence and, **46**:222, 228, 229
leukocyte integrins and, **46**:174, 175
lymphocyte hybridomas and, **38**:279, 291
major basic protein, **60**:192
maps of Ig-like loci and, **46**:3, 12, 32
maternally transmitted antigen and, **38**:344, 345
murine lupus models and, **46**:72, 86–88, 90, 91, 96
NK cells and, **47**:210, 246, 254, 263, 265, 266
PIG-A, **60**:82–84
regulators of complement activation and, **45**:397–400
SCID and, **49**:386
sporozoite malaria vaccine and, **45**:294
surface antigens of human leukocytes and, **49**:83, 89, 101, 113
T cell activation and, **41**:28
T cell development and, **44**:207
 ontogeny, **44**:216, 217
 thymocyte subpopulation, **44**:226–229, 231, 233
T cell receptor and, **45**:165
T cell receptor γ , **43**:145
T cells and, **38**:2
 cDNA, **38**:8
 clones, **38**:5
 cytotoxic, **38**:21
expression, **38**:13, 14
Ir genes, **38**:159, 189
rearrangement, **38**:12, 13
transcription, **38**:2, 10, 11
V region genes, **38**:15, 21
T cell subsets and
 cell surface molecules, **41**:41–43, 45, 47, 50
 H-2 molecules in thymus, **41**:97, 99
TGF α , **60**:195
virus-induced immunosuppression and, **45**:344, 361
Metalloproteases, **62**:266–267, 288
Metal-requiring enzymes, **60**:340
Metaphase, B cell formation and, **41**:206
Metastasis
 adoptive T cell therapy of tumors and, **49**:286, 297, 305
 and leukocyte homing, **58**:396
Met-enkephalin, **48**:172
lymphocyte ehemotaxis and, **39**:307–308
Met-enkephalin receptors
 on mixed lymphocytes, **39**:312
 on T cells, **39**:312
Methimazole, induced insulin autoimmune syndrome, HLA class II peptide binding and, **66**:87–88
Methionine
 antinuclear antibodies and, **44**:118, 120
 CD4 molecules and, **44**:289, 290
Methotrexate, in cancer treatment
 functional studies, **56**:346–347
 monoclonal antibody conjugation procedure, **56**:338–340
Methotrexate serum, hybrid resistance and, **41**:350
Methylated bovine serum albumin, aPL antibodies and, **49**:234–235
Methylation
 status in *cis*, role in gene rearrangements, **58**:30–31
 V(D)J joining, of antigen receptor genes, endogenous substrate, **56**:44–46
MG, *see* Myasthenia gravis
Mg21, **62**:76
MGF/STAT5, **65**:20
MGUS, **64**:219, 220
MHC, *see* Major histocompatibility complex;
 Major histocompatibility complex antigens

- MHr, *see* Myohemerythrin
- Micelles, cytotoxicity and, **41**:303
- Microfilaments, complement receptor 1 and, **46**:199
- β_2 -Microglobulin
- T cell receptor and
 - alloreactivity, **45**:155, 156
 - MHC molecules, **45**:111, 115
 - structure-function relationships, **45**:140
- virus-induced immunosuppression and, **45**:353
- β_2 -Microglobulin, **64**:115–116
- α chain complexes, retention, **52**:65–68
 - B cell formation and, **41**:198
 - CD1 heavy chains and, **59**:43, 47–50
 - CD8 molecules and, **44**:271, 272
 - chemotactic for pro-T cells, **58**:380
 - CTL and, **41**:135, 138, 165, 170
 - exon shuffling, **41**:138, 139, 147
 - HLA class I antigens, **41**:149, 150
 - T cell recognition, **41**:154–156
 - deficient mouse, *see* Mouse, MHC class I-deficient
 - exogenous, association with class I molecules, **52**:86–87
 - Ig gene superfamily and, **44**:24, 25, 30, 31
 - maps of Ig-like loci and, **46**:2
 - T cell subsets and, **41**:44, 75
- Microsomal antigens, autoimmune thyroiditis and, **46**:266, 273, 298, 299
- Microtubule organizing center
- cytolysis and, **41**:275, 282
 - humoral immune response and, **45**:37, 38, 42
- Microtubules, immunosenescence and, **46**:224
- MIF, *see* Macrophage migration inhibitory factor
- Migration, *see also* Transmigration
- CD23 antigen and, **49**:164, 169
 - defect, in IBD, **42**:307–308
 - effector T cells
 - basic pattern, **53**:242–244
 - tissue-specific, **53**:244–247
 - eosinophils, **60**:164–178
 - inflammatory cells, NO role, **60**:330–332
 - leukocytes, **58**:345–351
 - lymphocyte, and immunological memory, **53**:242
 - memory B cells, **60**:274–276
 - adhesion molecules and, **53**:248–249
 - basic pattern, **53**:247–248
- memory T cells
- adhesion molecules and, **53**:248–249
 - basic pattern, **53**:242–244
 - tissue tropic subsets, **53**:244–247
- naive B cells
- adhesion molecules and, **53**:248–249
 - basic pattern, **53**:247–248
- naive T cells
- adhesion molecules and, **53**:248–249
 - basic pattern, **53**:242–244
 - tissue-specific, **53**:244–247
- plasma cell precursors, **40**:198–199
- plasma cells, **53**:247–248
- Migration inhibition factor, *see* Macrophage migration inhibitory factor
- Milk, S-IgA antibodies in, **40**:201
- mim-1* gene, **65**:22
- Minor histocompatibility antigens, maternally transmitted antigen and, **38**:333, 334
- MIP-1 α , HIV and, **65**:282
- MIP-1 β , HIV and, **65**:282
- MIP-1 α receptor, cloning, **55**:135
- MIR, antibodies to, *see* Main immunogenic region, AChR, antibodies to
- Mitochondria, *see also* Antimitochondrial antibody
- aPL antibodies and, **49**:197, 239, 249–250, 256
 - autoimmune thyroiditis and, **46**:272
 - cell-mediated killing and, **41**:288
 - defects, maternally transmitted antigen and, **38**:333
 - lymphocyte homing and, **44**:317
 - maternally transmitted antigen and, **38**:337, 352
 - genetic mechanism models, **38**:340
 - maternal transmission, **38**:322, 323
 - polymorphisms, **38**:319, 320, 324, 326
 - strain combinations, **38**:349
 - transfer via cell fusion, **38**:314
 - natural killer cells and, **47**:214
 - protein export, maternally transmitted antigen and, **38**:315
 - spontaneous autoimmune thyroiditis and, **47**:442, 448
 - surface receptor, maternally transmitted antigen and, **38**:341
- T cell receptor and, **45**:123

- Mitochondrial DNA
 damage, **38:319**
 genetic mechanism models, **38:339**
 maternally transmitted antigen, **38:316**,
 317, **329**, **351**
 polymorphism, **38:323–327**
 rearrangement, **38:320**
 restriction enzyme polymorphism, **38:337**
 restriction patterns, **38:327**, **328**
- Mitochondrial enzymes, action of, **65:122**
- Mitochondrial transmission, maternally
 transmitted antigen and, **38:351**
 of class I-like antigen, **38:348**
 of conventional antigen, **38:346–348**
 of enzymatic activity, **38:346**
 induced nuclear response, **38:344–346**
- Mitogen-activated protein kinase, *see* MAP
 kinase
- Mitogenic lectins, IBD and, **42:289**
- Mitogenic signaling, Ig, **59:326–328**
- Mitogens
 adoptive T cell therapy of tumors and,
 49:314
 antigen-presenting cells and, **47:71**, **73**, **100**
 antinuclear antibodies and, **44:107**
 autoimmune thyroiditis and, **46:289**, **310**
 autoreactive T cells and, **45:426**
 B cell formation and, **41:192**, **200**, **228**, **231**
 CD5 B cell and, **47:140**, **142**, **147**
 CD23 antigen and, **49:170**
 complement receptor 2 and, **46:206**, **208**
 cytotoxicity and, **41:272**, **284**, **285**
 donor lymphocytes and, **38:283**, **284**
 helper T cell cytokines and, **46:122**
 HIV infection and, **47:390**, **391**, **398**, **399**
 human T cell activation and, **41:7**, **8**, **11**,
 12, **26**, **27**
 humoral immune response and, **45:67**,
 70, **72**
 antigens, **45:23**
 helper T cells, **45:26**, **29**, **33–35**
 interleukins, **45:64**, **73**, **75**
 physical interaction, **45:43**
 hybrid resistance and, **41:350**, **393**, **395**
 IL-1 and, **44:154**, **168**, **189**
 human disease, **44:192**, **194**
 immunocompetent cells, **44:173–177**
 immunosenescence and, **46:221**
 lymphocyte activation, **46:223–225**, **228**,
 234
- mucosal immunity, **46:252**
 regulatory changes, **46:250**
 stem cells, **46:244**, **245**
- leukocyte integrins and, **46:173**
Mycoplasma arthritides, **54:113**, **141**
 NK cells and, **47:206**, **261**, **264**
 potential effector mechanisms, **47:467**
 SCID and, **49:382–385**, **387**, **398**, **400**
 spontaneous autoimmune thyroiditis and,
 47:491, **493**
 cellular immune reactions, **47:452**, **456**
 disturbed immunoregulation,
 47:470–474, **476**
 genetics, **47:488**
- T cell development and cellular selection
 within thymus, **44:248**
 ontogeny, **44:221**
 thymocyte subpopulation, **44:229**, **236**,
 241
- T cell subsets and, **41:60**, **63–66**, **111**
 tissue distribution, **47:54**
- virus-induced immunosuppression and
 HIV, **45:355**, **356**, **358**, **359**, **361**, **362**
 measles, **45:338**, **339**, **342**, **344**
- Mitomycin C, in cancer treatment, **56:329**,
 332–333
- Mitosis
 B cell formation and, **41:184**
 HIV and, **58:424**
 immunosenescence and, **46:241**, **246**
- Mixed connective tissue disease, antinuclear
 antibodies and, **44:93**, **111**
 autoantibodies, **44:136**, **138**
 autoantigen, **44:130**
 autoimmune response, **44:131**
 scleroderma, **44:124**
 SLE, **44:100**
- Mixed-isotype dimer, formation, **48:125**
- Mixed leukocyte cultures, NK cells and,
 47:295
- Mixed-leukocyte reaction, *see* Mixed-
 lymphocyte reaction
- Mixed-lymphocyte reaction, **59:376**
 antigen-presenting cells and, **47:65**, **71–73**,
 75, **76**, **78–80**
 APC-T cell binding, **47:95**, **97**, **99**
 cell surface, **47:59**, **61**
 T cell, **47:83**, **84**, **99**
 tissue distribution, **47:48**, **49**, **53**, **54**
- Ia role, **39:81**

- Mixed-lymphocyte reaction (*continued*)
 immunosenescence and, **46**:243, 244
 MHC class II-mediated, B cells and,
39:81, 83
 Mls-mediated, B cells and, **39**:81–83
 primary, activated B cells and, **39**:80–81
 secondary, resting B cells and, **39**:80
 T cell subsets and
 cell surface molecules, **41**:43
 H-2 alloantigen recognition, **41**:81,
 83–87
 APC, **41**:88–92
 H-2 molecules in thymus, **41**:107, 108
 virus-induced immunosuppression and,
45:341
- Mixed-lymphocyte response, *see* Mixed-lymphocyte reaction
- MLA-144 leukemic cell line, **42**:172
- MLR, *see* Mixed-lymphocyte reaction
- mls
 identification as product of MMTV,
54:100–102
 in vitro response to, **54**:124–125
- Mls antigen, **50**:7–10, **65**:139, 196
 antigen-presenting cells and, **47**:47
 antigen presentation, **47**:64, 76–79
 APC-T cell binding, **47**:95, 99
 and bacterial toxic mitogen action, working
 model, **50**:4, 5–6
 B cell
 MLR and, **39**:81–83
 during ontogeny, **39**:81
 definition, **50**:7–9
 multiple loci, **50**:9–10
 Mls loci, multiple, **50**:9–10
 current nomenclature for, **50**:11
- MMTV, *see* Mouse mammary tumor virus
- 6-MNA, **62**:186
- Mobility, **43**:33–35
 crystallographic temperature factors, **43**:20
 functional significance, **43**:73–74
 interacting structural elements, **43**:80
 local, relationship with antigenicity, **43**:61
 secondary structure, **43**:76
 T cell receptor $\gamma\delta$ proteins, **43**:150
- Molecular adaptation, **64**:20
- Molecular complexes, CD19, CR2, and CR1
 on surface of human B cells, **52**:146
- Moloney leukemia virus, **41**:364, 400
- Moloney sarcoma–leukemia virus, **41**:153
- Moloney sarcoma virus, **41**:378
- Monensin, **41**:281
- Monkey
 Ascaris antigen challenge, **60**:168–169
 asthma model, **60**:219
 complement components and, **38**:207–209
 (T1B)₄ MAP immunization, **60**:116–117
- Monoclonal antibodies, **53**:5D3, 276–277,
64:323
 5D3, LPS mimesis, **53**:276–277
 109d6, RA susceptibility associated with,
56:422–425
 109 D6, rheumatoid arthritis, **48**:128
 AChR and, **42**:240–241
 adoptive T cell therapy of tumors and,
49:283, 320, 327
 mechanisms, **49**:308, 312–314, 317
 principles, **49**:290, 296–297
 anti- β , **43**:283
 antigen-presenting cells and, **47**:70, 76,
 77, 96
 cell surface, **47**:58, 60–62
 tissue distribution, **47**:50, 51, 53
 anti-Id, **42**:149–157
 antinuclear, **44**:98, 108, 118, 134–136
 antiphospholipid, **49**:239, 242, 245,
 248–249, 251
 anti-Tac, **42**:167
 Ars A response and, **42**:98
 autoimmune demyelinating disease, **49**:357,
 363, 366
 autoimmune thyroiditis
 antigens, **46**:267, 269, 272, 273
 cellular immune responses, **46**:287, 295
 experimental models, **46**:274, 275, 277,
 280
 humoral responses, **46**:300, 302–306,
 308
 prevention, **46**:310, 314
 autoreactive T cells and, **45**:420
 B cell formation, **41**:181, 182, 235, 238
 B cell precursors, **41**:188, 189
 bone marrow cultures, **41**:213
 cell size changes, **41**:202
 functional assays, **41**:200, 201
 Ig genes, **41**:204
 Ly-5 family of glycoproteins, **41**:189, 192
 lymphomohemopoietic tissues, **41**:187
 markers, **41**:193, 194
 NZB mice, **41**:227

- PI-linked lymphocyte antigens, **41**:198
SCID mice, **41**:225
tumor cell lines, **41**:195
CD1 proteins, **59**:39–43
CD2, **48**:235–236
 CD3- Jurkat mutant, **48**:236
 CD3- thymocyte, **48**:236
 NK cell, **48**:236
CD3, effect on IL-9 expression in T cells, **54**:85
CD4 and CD8, **44**:298–303
 CD4 molecular biology, **44**:284, 285, 290, 291, 293, 295–297
 CD8 molecular biology, **44**:271, 278, 280, 282
CD5 B cell and, **47**:117
 genetic influence, **47**:143–147
 Ig gene expression, **47**:153, 155
 physiology, **47**:133, 134
 primordial immune network, **47**:160
CD23, **49**:177; **60**:181
 biochemical structure, **49**:155, 157
 biological activity, **49**:171–172
 cellular expression, **49**:152–153, 155
 Fc ϵ RII, **49**:162, 164–165
CD28, **48**:245
 mediated T cell activation, mechanism, **48**:246
CD40, **60**:45–49
CD44, *in vitro* synthesis, **54**:301
CD45, studies, **66**:1, 3, 5–17
in chemoimmunoconjugates for cancer
 treatment
agents, **56**:302
alkylating agent–antibody conjugates, **56**:328–333
anthracycline–antibody conjugates, **56**:333–338
antigen heterogeneity affected by, **56**:352
barriers to chemotherapy, **56**:347–354
clinical trials, **56**:354–356, 358–359
conjugation strategies, **56**:312–314, 318–319, 323
cytotoxic drugs, **56**:342–343
design strategy, **56**:324–325
development, **56**:302–304
folic acid antagonists, **56**:338–340
functional studies, **56**:344–347
immunogenicity, **56**:311–312
intermediary carriers, **56**:323–324
internalization, **56**:308–310
localization, **56**:306–308
mode of action, **56**:344–347
modulation, **56**:308–310
morphological studies, **56**:347
preclinical studies, **56**:326–328, 343
radiolabeling, **56**:354
size effects, **56**:310–311
targets, **56**:305–306
toxicity, **56**:311–312
vinca alkaloids, **56**:340–342
complement receptor 1 and, **46**:196, 200, 201
complement receptor 2 and, **46**:203, 204, 206, 211
CTL and, **41**:169, 170
 β_2 -microglobulin, **41**:155, 156
 amino acid, **41**:163
 blocking, **41**:156–158
 exon shuffling, **41**:140, 141, 145, 148
 HLA class I antigens, **41**:150
 somatic cell class I variants, **41**:165–167
cytotoxicity and, **41**:285, 291, 307, 309
EG1, **60**:193
EG2, **60**:193, 220
Fc receptors
 Fc ϵ R, **43**:299–300
 Fc α RI, **57**:9, 40–41
 Fc γ RI, **57**:8–9
 Fc γ RII, **57**:9, 17–19
 Fc γ RIII, **57**:9, 29–30
 Fc ϵ RI, **57**:9, 36–37
 Fc ϵ RII, **57**:38
fibronectin and, **38**:387–389
helper T cell cytokines and, **46**:112, 119, 123–125
HIV infection and, **47**:386
human B cell neoplasia and, **38**:269
hybrid resistance and, **41**:336, 374, 377, 378, 386
IBD and, **42**:289
ICAM-1 and VCAM-1, **60**:168–170
IgE biosynthesis and
 antibody response, **47**:8, 31, 33–35
 binding factors, **47**:16, 18–22, 24, 27
Ig gene superfamily and, **44**:33, 41
Ig heavy-chain variable region genes and, **49**:46, 48, 51
IL-1, **44**:174

- Monoclonal antibodies (*continued*)
 IL-2Ra, treatment with, **50**:188–189
 IL-5, **60**:215
 immunorescence and, **46**:249
Ir genes and
 antigen binding, **38**:99, 107
 Ia molecules, **38**:68–72, 75, 77, 78
 mutations, **38**:88, 90
 T cell repertoire, **38**:116
 T cell suppression, **38**:150, 151
 tolerance, **38**:125
J11D, **42**:74, 76
 leukocyte integrins and, **46**:150, 151, 177
 inflammation, **46**:162–166, 168
 leukocyte adhesion deficiency disease, **46**:173
 ligand molecules, **46**:170, 172, 173
 structure, **46**:153
 lymphocyte homing and
 carbohydrate, **44**:354, 361, 364, 371
 high endothelial venules, **44**:317, 318
 molecules, **44**:342–346, 348, 350–352
 regional specificity, **44**:327, 329, 338
 lymphocyte hybridomas and, **38**:275–277, 306
 antitetanus toxoid, **38**:303
 B cell hybridization, **38**:279, 280
 donor lymphocytes, **38**:282, 286
 EBV-transformed B cells, production by, **38**:277–279
 genetic engineering, **38**:304, 305
 human, **38**:301, 302
 human-human hybridomas, **38**:292, 293, 296, 299
 human-mouse heterohybridomas, **38**:301
 human-murine hybridomas, **38**:289, 290
 maternally transmitted antigen and, **38**:331, 351
 MC therapy and, **42**:272
 to MHC, protein and peptide-induced, **43**:61–62
 mouse, segmental flexibility, **48**:13, 14
 multiple myeloma, **64**:242–243, 260–263
 multiple sclerosis and, **49**:369, 372–375
 murine lupus models and, **46**:95
 neoplastic plasma cells as immunogens, **64**:243–244
 NK cells and, **42**:182; **47**:198, 296
 CD16, **42**:188, 190–191
 congenital defects, **47**:224, 225
 differentiation, **47**:233
 effector mechanisms, **47**:243, 258
 hematopoiesis, **47**:278, 279
 HNK-1, **42**:191–192
 NKH1 antigen, **42**:188–190
 phenotype, **42**:186
 surface phenotype, **47**:200, 201, 205, 207, 208, 210, 211
 tissue distribution, **47**:219, 220
 phagocytosis of zymosan, **38**:369, 370
 reactivity with isolated FDCs, **51**:259
 negative, **51**:250–251
 positive, **51**:248–249
 regulators of complement activation and, **45**:390, 396, 397, 410
 reshaped human PM-1, construction, **64**:261
 SCID and, **49**:394–395, 397
 spontaneous autoimmune thyroiditis and, **47**:435
 altered thyroid function, **47**:460–462
 cellular immune reactions, **47**:454, 455
 disturbed immunoregulation, **47**:472, 479
 histopathology, **47**:442
 potential effector mechanisms, **47**:466
 sporozoite malaria vaccine and, **45**:292, 293, 306, 312
 in studies of CD antigen expression on basophils/mast cells, **52**:336–337
 surface antigens of human leukocytes and, **49**:87, 117, 125–126
 receptors, **49**:98, 100–101, 111
 synthetic T and B cell sites and, **45**:238, 239
 T cell activation and, **41**:30
 accessory molecules, **41**:14
 cell surface molecules, **41**:1, 2
 gene regulation, **41**:28
 IL-1 receptor, **41**:14
 receptor-mediated signal transduction, **41**:19–22, 24–26
 T1, **41**:13
 T11, **41**:9, 10
 T cell antigen receptor, **41**:2, 4–8
 Thy-1, **41**:11
 Tp44, **41**:11
 T cell development and, **44**:209
 cellular selection within thymus, **44**:245–247, 254, 255
 ontogeny, **44**:218, 220, 224

- thymocyte subpopulation, **44**:227, 228, 232, 239, 243
- T cell receptors and, **45**:130, 141, 153, 156
protein properties, **38**:3–5
target interaction, **38**:21
thymus, **38**:12
- T cell subsets and
H-2 alloantigen recognition, **41**:79, 84, 89, 93
H-2-restricted antigen recognition, **41**:53
T cell receptor, **41**:40, 43
- therapeutic use, **64**:196
- virus-induced immunosuppression and, **45**:346, 357, 362
- to VSAG, structure–function studies, **54**:105–106
- Monoclonal gammopathies, undetermined significance, *see* MGUS
- Monocyte chemoattractant protein-1, **66**:135
sepsis and, **66**:104, 141–142
- Monocyte-derived eosinophil cytotoxicity enhancing factor, **39**:216
- Monocyte/macrophages
LIF role in, **53**:35–36
LPS receptors, **53**:274–281
- Monocytes
adherent IgG effects on inductor of IL-1 α production, **54**:182
adoptive T cell therapy of tumors and, **49**:328
antinuclear antibodies and, **44**:95
autoimmune thyroiditis and, **46**:294
CD1 protein distribution, **59**:64–66
CD5 B cell and, **47**:142
CD23 antigen and, **49**:153, 158, 174, 176
biological activity, **49**:167, 169, 171, 173
expression regulation, **49**:160–161
Fc ϵ RII, **49**:166
CD40 expression, **61**:13, 40
complement receptor 1 and, **46**:190, 199–201
complement receptor 2 and, **46**:207
differential regulation of IL-1 β and IL-1 α production, **54**:181–184
effects on cytokines in IL-1 α production by, **54**:182–184
Fc receptors on, **40**:64–66
fibronectin and, **38**:390
HIV infection and
activation, **47**:400, 402, 403
- immunopathogenic mechanism, **47**:392–397
neuropsychiatric manifestations, **47**:404
- human, phagocytosis
alternative complement pathway, **38**:385
particulate activators, **38**:380, 382
zymosan, **38**:363–376
- IFN- γ and, **62**:80, 136
- IgE, **43**:305–306
- IGF-I, **63**:389–391
- IL-1 and, **44**:153, 190
biological effects, **44**:166, 167, 169
gene expression, **44**:159–161, 163, 164
human disease, **44**:193, 194
immunocompetent cells, **44**:178
structure, **44**:155, 157
- TNF, **44**:185, 186
- IL-8 effects, **55**:117–118
- immunosenescence and, **46**:228
- leukocyte integrins and, **46**:161, 163–167
- lymphocyte homing and, **44**:339, 352
- opioid receptor, **48**:180
- PGHS-2, **62**:191–192
- pituitary hormones, **63**:411–412
- purification of IL-1 α from IgG-induced supernatants, **54**:172
- SCID and, **49**:384–385, 389–391, 396, 401
- as source of chemokines, **55**:105–106
- spontaneous autoimmune thyroiditis and, **47**:475
- surface antigens of human leukocytes and, **49**:111, 116
adhesion molecules, **49**:86, 88
Igs, **49**:89, 91
MHC glycoproteins, **49**:82
TGF- β regulation, **55**:187
- Monocybrids, CD5 B cell and, **47**:141–143
- Monocytopenia, hybrid resistance and, **41**:392
- N*^G-Monomethyl-L-arginine, **62**:72, 75, 99
in arthritis and diabetes, **60**:344–345
in conjunction with lipopolysaccharide, **60**:347
- effect on
cytokinoplasts, **60**:352
infection, **60**:341, 343
- inhibition of F-BSA extravasation, **60**:328

- Mononuclear cells
 IBD and, **42**:289, 294–297
 antibody secretion and, **42**:301–307
 intestinal
 antibody secretion and, **42**:302–307
 IBD and, **42**:295–296
 lamina propria, IBD and, **42**:297
- Mononuclear phagocytes, IFN- γ and, **62**:78–80
- Mononucleosis, virus-induced
 immunosuppression and, **45**:352
- MOPC-315, antigen-specific inhibition by suppressor T cells, **40**:143–144
- Morphine, NK cells and, **47**:268
- Morphology
 B cell formation and, **41**:182, 186, 204, 213, 219, 238
 cytotoxicity and, **41**:271
 cytosis, **41**:274, 276, 281–283
 cytolytic proteins, **41**:317
 granule proteins, **41**:287, 291, 294, 295
 membrane attack complex of complement, **41**:300, 304, 305, 309, 311
 degranulating eosinophils, **60**:160–162
 eosinophil activation, **60**:158–160
 HIV infection and, **47**:378
- NK cells and, **47**:214–218, 292
 antimicrobial activity, **47**:283, 290
 cytotoxicity, **47**:249, 250, 253, 256, 258
 differentiation, **47**:234
 hematopoiesis, **47**:275, 281
 identification, **47**:197, 198
 malignant expansion, **47**:226, 227
 reproduction, **47**:271
 surface phenotype, **47**:210, 213
 tissue distribution, **47**:219, 220
- normal mature eosinophils and eosinophil myelocytes, **60**:155–158
- spontaneous autoimmune thyroiditis and, **47**:444, 467, 472
- MORT-1, **61**:31
- Mosaicism, X-chromosome, **59**:159, 195
- Mouse
 acquired immunodeficiency disease syndrome, **54**:115
 A/J Inbred strains, V_H genes in, **42**:137
 athymic nude, **58**:309–311
 B cell repertoire expression, *see* B cell repertoire expression
- bearing male-specific transgenic T cell receptor, effect of IL-2 on, **50**:170–171
- CD2-deficient, **58**:172
- class A gene *Pig-a*, **60**:72, 93
- EAE-susceptible, MBP epitope identification, **48**:164–165
- erythrocytes, rosette formation with human B cells, **40**:18
- genome, non-Mls elements in, **50**:11–14 and human
 heavy chain locus organization, **54**:231–233
 IL-9 characterization and cloning, **54**:80–82
- HuSCID, antibodies derived from, **57**:230–231
- IELs, invariant $\gamma\delta$ T cell receptors, **58**:301–304
- Ig-transgenic, CD5 B cells in, **55**:324–326
- IL-1 receptor antagonist, cloning and expression, **54**:175–176
- IL-6 transgenic, **54**:45–47
- IL-9 receptor cDNA, **54**:82, 84
- immunodeficient, IL-5 production, **57**:171–172
- IRF-1 knockout, **60**:335
- lpr*
 etiology of autoimmunity, **60**:308–309
 Fas gene mutation in, **57**:132–133
lpr/lpr, accumulation of CD4⁻⁸⁻ $\alpha\beta$ T cells, **58**:112
- malaria model, **60**:108–112
- MHC class I-deficient, **55**:381–383, 415–417; **58**:313
 autoimmune disease, **55**:415
 bacterial infection, response to, **55**:412–413
- cell surface expression, **55**:385–387
 development, **55**:384–385
 functional class I molecules on β_2 m-deficient cells, **55**:386–387
- hematopoietic cell grafts, **55**:388
- human filarial parasites, response to, **55**:415
- impaired NK cell activity, **55**:404–405
Leishmania major infection, response to, **55**:414–415
- liver cell grafts, **55**:391

- lymphocytes, deficient development of, **55:393–404**
- NK cells, target cell susceptibility, **55:392–393**
- pancreatic islet grafts, **55:390**
- pathogens, responses to, **55:405–406**
- production, **55:383–384**
- serological analysis, **55:385–386**
- skin grafts, **55:388–390**
- transplantation studies, **55:387–391**
- Trypanosoma cruzi* infection, response to, **55:413–414**
- viral infections, response to, **55:406–412**
- MHC class II-deficient, **55:423–424**
- B cells, **55:429–430**
- CD4⁺ T cells, **55:424–427**
- CD8⁺ T cells, **55:427–428**
- double deficient mouse, **55:435–436**
- E-only mouse, **55:434–435**
- immune system, peripheral compartment, **55:424–430**
- $\gamma\delta$ T cells, **55:428–429**
- thymus, **55:433**
- reconstituted, **55:433–434**
 - single-positive compartment, **55:430–433**
- motheaten, CD5 B cell expression in, **55:314**
- MRL/Mp-lpr/lpr, recombinant IL-2/vaccinia virus construct on, **50:193–203**
- Mtv-7⁺ and Mtv-7⁻, **58:222**
- mutations affected by stem cell factor or SCF receptor, **55:17–23**
- neonatally thymectomized, effect of IL-2.VV, **50:205–208**
- Nu/nu*, effect of IL-2.VV on, **50:204–206**
- p53-deficient, **58:169**
- perforin gene mutation, **60:292–296**
- retention of Vbse sequences, **50:41–42**
- scid*, **60:341–343**
- and DSB repair and V(D)J recombination, **58:45–51**
 - genetic reconstitution experiments, **58:142–143**
 - lymphocytes, **58:317**
 - rearrangement-deficient, **58:155–156**
 - thymocytes, **58:37–38**
- SEB shock model, **60:348–349**
- self-reactive T cells, **58:274–277**
- with severe combined immune deficiency, *see* Severe combined immunodeficiency, SCID mouse
- T cell activation, **41:1**
- cell surface molecules, **41:10, 11, 13, 14**
- receptor-mediated signal transduction, **41:23, 25**
- T cell development stages in, **51:118–119**
- T cells, tumorigenesis, **54:88–93**
- transgenic
- bcl2/scid*, **58:48–49**
 - for given Mtv, **50:40**
 - H-2^b and H-2^d, **58:313–316**
 - IL-5 production models, **57:169–170**
 - with knockout mutations, **58:30–33, 41, 89**
 - RAG^{-/-}, **58:143–145**
 - TAP^{1-/-}, **58:162–163**
 - TCR- α , - β , - γ , and - δ loci, **58:118–119**
- transgenic spleen cells, *in vitro* studies with, **54:414–415**
- Vbse not stimulating primary T cell response, **50:11–13**
- X-linked immunodeficient, **60:41–42**
- IL-5 production models, **57:171–172**
- Mouse mammary tumors, **65:139**
- induction of, **65:153–155**
- Mouse mammary tumor virus, **50:39; 65:139–141, 211–212**
- endogenous
- immune stimulation by superantigens, **65:196–203**
 - superantigen expression, **65:152–153**
 - tolerance induction, superantigens and, **65:203–208**
 - virology, **65:141–146**
- exogenous
- adult T cell response, **65:175–178**
 - neonatal response, **65:171–175, 203**
 - T cell-dependent B cell differentiation and, **65:178–180**
 - virology, **65:141–146**
- immune response to, **65:167–168, 212**
- adult T cell response, **65:175–178**
 - cellular response, **65:192–194**
 - humoral response, **65:188–192**
 - neonatal response, **65:171–175, 203**
 - receptors for, **65:194–196**
 - superantigen-dependent, **65:171–194**

- Mouse mammary tumor virus (*continued*)
superantigens and T cell-independent, **65**:168–171
T–B interaction, **65**:180–188
T cell-dependent B cell differentiation, **65**:178–180
superantigens, **65**:140, 152–153
cellular expression of antigens, **54**:106–107
expression, **65**:152–153
immune stimulation by, **65**:140, 141, 171–194, 196–203
MHC class II molecules, interaction with, **65**:164–165
Mls identification as product, **54**:100–102
protein structure, **65**:157–164
role in life cycle of virus and host, **54**:107–108
structure–function studies, **54**:103–106
TCR V β , interaction with, **65**:165–167
tumor formation and, **65**:152, 155
virology, **65**:141–145
amplification and spread, **65**:180–188
infection and transmission, **65**:148–151
life cycle, **65**:211
structure, **65**:146–148
tissue distribution, **65**:151–152
transcriptional regulation, **65**:148
- Mouse mammary tumor virus receptors, **65**:194–196
- MPL units, aPL antibodies and, **49**:203–204
- mRNA, *See* Messenger RNA
- α -MSH, *in vitro* IgE synthesis, **61**:371–372
- mTOR, **61**:181, 188–190
- ntp* genes, **65**:49
- MUC-1, **62**:223, 229, 238–239
- μ chains, **43**:238–239
- Mucins, **62**:222–223, 238–239
lymphocyte–HEV interaction, **65**:360–361
tumor antigen recognition of, **57**:302–303
- Mucosa
addressin, **62**:257
associated lymphoid tissue
functional anatomy of Peyer's patches, **40**:192–195
studies on disassociated Peyer's patches cells, **40**:195–197
and immunity, **58**:297–298; **59**:372
lymphocyte homing into, **64**:175–177
- Mucosal immunity, immunosenescence and, **46**:221, 251–253
- Mucosal inflammatory infiltrate, in IBD, **42**:298
- Mucosal organs, lymphocyte homing and, **44**:323, 327–330, 334–336
- Mucosal tissues, mechanisms of homing of plasma cell precursors to, **40**:204–207
- Mucosal vascular addressin, lymphocyte–HEV interaction, **65**:362–363
- Mu heavy chain
allelic exclusion, **63**:2, 19–21
B cell development, **63**:1–4, 11–15, 16–25, 31
ligand for PreB cell receptor, **63**:25–26
signal transduction through PreB cell receptors, **63**:26–29
- Muller cells, **48**:218–219
- Multicatalytic protease complex, **52**:26
- Multichain immune recognition receptors, **61**:154
- Multi-CSF, *see* Interleukin-3
- Multidrug resistance gene, mammalian, **65**:51
- Multimeric–Synthetic Peptide Combinatorial Library, **60**:139
- Multiple antigen peptides
as antigens, **60**:137–139
based on schistosome antigens, **60**:125–128
definition, **60**:105–106
design and synthesis, **60**:106–108
in helminthic diseases, **60**:124–128
mono-, di-, and triepitope, **60**:120–123
[(NAAG)₆] MAP, immunized mouse, for *P. malariae*, **60**:123
in protozoan diseases, **60**:108–124
sporozoite malaria vaccine and, **45**:321
(T1B)₄ MAP construct, immunogenicity, **60**:114–120
for viral and bacterial diseases, **60**:128–136
- Multiple myeloma, **42**:224; **64**:219, *see also* Myeloma
adhesion molecules, **64**:246–249
CD40, **61**:12, 39
cell identification, **64**:242–246
chromosome and oncogene abnormalities in, **64**:250–251
cytokines, **64**:252–260
IL-6 signal transduction therapies, **64**:260–263

- immunophenotype, **64:242**
oncogenic transformation, **64:249–251**
- Multiple sclerosis, **49:357**
antibodies, **49:368–374**
aPL antibodies and, **49:209**
DP α -DQ β , **48:149**
DPw4, **48:149–150**
DQ α -DQ β , **48:149**
DQ α gene polymorphism, **48:149**
DQR2-6, **48:148–149**
DQwl, **48:148**
DQwl_a, **48:148**
DQwl_b, **48:148**
DR2, **48:148–150**
DR α -DQ β , **48:149**
Dw2 subtype, **48:148–150**
HLA-A3, **48:148–150**
HLA-B7, **48:148–150**
HLA class II peptide binding and, **66:88–91**
IFN- γ and, **62:75**, 91
myelin basic protein and, **49:362–363**
NK cells and, **47:301**, 302
pituitary hormones, **63:425**
T cell response, **48:150**
TCR usage restriction and, **49:367–368**
vaccination to TCR V regions, **49:374–375**
- MUM-1, **62:229**, 237
- Murine acquired immunodeficiency syndrome, **52:432**
- Murine colony-stimulating factor, **48:73**
- Murine cytomegalovirus, hybrid resistance and, **41:389**
- Murine encephalomyocarditis virus, resistance to infection, **65:31**
- Murine hepatitis virus, hybrid resistance and, **41:375**, 376
- Murine leukemia virus, **52:432–434**
 antineuclear antibodies and, **44:130**
 plasmacytogenesis, **64:230**
- Murine mammary tumor virus, *see* Mouse mammary tumor virus
- Muscle, AChR, **42:239**
 autoantibody effects on, **42:260–263**
- Muscle cells, cachectin and, **42:223**
- Mutagenesis
 CD4 molecules and, **44:295**
 genetically engineered antibody molecules and antigen-combining sites, **44:84**
- biological properties, **44:83**
cloning, **44:75**
expression, **44:77**, 79
production, **44:68**
HIV infection and, **47:386**
spontaneous autoimmune thyroiditis and, **47:464**
- Stat4, **60:20**
- Mutants
 GPI anchor-deficient, **60:65–67**
 PIG-A, clonal dominance, **60:90–91**
 V(D)J recombination/DSB repair and cell cycle arrest, **58:71–72**
 leading to chromosome errors, **58:72–74**
- Mutations, *see also* Deletion; Hypermutation; Somatic mutation
adoptive T cell therapy of tumors and, **49:320**, 327, 332, 334
affected by stem cell factor or SCF receptor
 human, **55:23–24**
 mouse, **55:17–23**
 rat, **55:23**
antigen-presenting cells and, **47:80**, 103
autoimmune thyroiditis and, **46:263**, 286
- autoreactive T cells and, **45:417**, 420, 424, 427
- B cell formation and, **41:185**
B cell precursors, **41:195**, 198
genetically determined defects, **41:224**, 226, 228, 231, 232
- B cells, **59:300**
box1 motif, **60:12**
Btk genes, **59:185–192**, 195
- CD4 molecules and, **44:291**, 296
- CD5 B cell and, **47:132**, 146, 154, 155
- CD5⁺ B cell lines, **55:330**
- CD23 antigen and, **49:163–164**
- cell cycle checkpoints, **58:29**
- complement receptor 1 and, **46:193**, 194, 196
- complement receptor 2 and, **46:211**
- CTL and, **41:138**, 169, 170
- β_2 -microglobulin, **41:155**
- amino acid, **41:158–163**, 165
- carbohydrate moieties, **41:153**, 154
- exon shuffling, **41:141**, 142, 145, 148
- mAbs, **41:156**, 157, 165, 166

- Mutations (*continued*)
- Fas gene
 - loss of function, **57**:139
 - lpr* mouse, **57**:132–133
 - genetically engineered antibody molecules and, **44**:79, 84
 - grooves of MHC molecules, **58**:161
 - HIV genome, **65**:311–317
 - HIV infection and
 - etiological agent, **47**:381, 383
 - immune response, **47**:408
 - immunopathogenic mechanism, **47**:388, 400
 - hopscotch*, **60**:3
 - hybrid resistance and, **41**:343, 368, 374, 397
 - IgE biosynthesis and, **47**:16, 38
 - Ig gene superfamily and, **44**:18, 27, 28
 - Ig heavy-chain variable region genes and, **49**:6, 22, 32, 43, 53
 - IL-1 and, **44**:158, 182
 - IL-2R γ , **59**:246–253
 - immunosenescence and, **46**:222, 234
 - I region, *Ir* genes and, **38**:84–91
 - cytotoxic T cells, **38**:163, 164
 - gene function, **38**:92, 93
 - H-Y antigens, **38**:176
 - immunodominance, **38**:180
 - T cells, **38**:130, 131, 144
 - tolerance, **38**:126
 - kinase domain of XLA patients, **60**:40–41
 - leukocyte integrins and, **46**:160, 171, 172, 174–176
 - lpr*, **60**:294
 - maps of Ig-like loci and, **46**:3, 12
 - murine lupus models and, **46**:99
 - Ig germline, **46**:65, 66, 69, 76–78
 - lupus strains, **46**:63, 64
 - T cell antigen receptor, **46**:83, 94, 98
 - natural and artificial, **58**:90–101
 - NK cells and, **47**:225
 - p53 protein-derived tumor antigens, **57**:295–296
 - scid*, **58**:45–51; **61**:299–301, 317
 - SCID and, **49**:383
 - sporozoite malaria vaccine and, **45**:298, 311, 312
 - switch recombination, **61**:108
 - sxil*, **58**:54–55
- synthetic T and B cell sites and, **45**:217, 228, 229, 261
- T cell activation and, **41**:5, 8, 12, 20
- T cell receptor and
 - alloreactivity, **45**:150–152, 154, 156, 157
 - homogeneity, **45**:128, 129
 - structure-function relationships, **45**:135, 139–142
- T cell repertoire, **45**:164, 165
- T cell subsets and, **41**:42, 46
 - H-2 alloantigen recognition, **41**:81, 82, 85, 86, 88, 93
 - and transgenes, **58**:147–149
- V-3, **58**:51–52
- virus-induced immunosuppression and, **45**:349–351
- W, allelism with *c-kit*, **55**:6–7
- Xid mutation, **59**:174–176, 317–318, 328
- XR-1, **58**:52–53
- xrs, **58**:53–54
- MW, *see* Maedi-visna virus
- Myasthenia gravis, *see also* Acetylcholine receptor; Experimental autoimmune myasthenia gravis
 - anti-Id antibody as mediator of, **39**:279
 - aPL antibodies and, **49**:209
 - diagnosis of, **42**:254–256, 272
 - discovery of autoimmune nature of, **42**:233–234
 - etiology of, **42**:250–253
 - experimental approaches to, **42**:233–235
 - fatigue and, **42**:237
 - future prospects for, **42**:272–274
 - genetic factors in, **42**:259
 - Ig heavy-chain variable region genes and, **49**:46, 50–51
 - incidence of, **42**:255–256
 - murine lupus models and, **46**:79
 - pathology of, **42**:272
 - autoantibody effects, **42**:260–263
 - autoantibody production, **42**:256–260
 - penicillamine and, **42**:235
 - therapy for, **42**:235, 269–274
 - transient neonatal, **42**:252, *see also* Myasthenia gravis
- Myb
- B cell genesis, **63**:220–221, 238–239
 - interaction with NF-IL6, **65**:22

- myb*, **65:22**
 retrovirus-induced B cell neoplasia in bursa
 of fabricius, role in, **56:477–478**
- myc*, **63:161, 162**
 antisense, blocking of negative signaling, **54:413–415**
 retrovirus-induced B cell neoplasia in bursa
 of fabricius
 deregulation of expression, **56:480**
 role in, **56:472–478, 481**
 TGF- β and pRB, **54:411–413**
- Mycobacteria
 CD1-restricted antigens, **59:82–85, 89**
 reactivity, **58:321–322**
- Mycobacterium avium*, **50:162; 54:36**
- Mycobacterium bovis*, and diabetes, **51:294**
- Mycobacterium paratuberculosis*, IBD and, **42:319**
- Mycobacterium tuberculosis*, **54:36**
 infection, **59:398–402**
 pneumonia, **59:379, 383, 398–402**
 responses in MHC class I-deficient mouse, **55:412–413**
- Mycolic acid, **59:82–85, 90**
- Mycophenolic acid, genetically engineered antibody molecules and, **44:70–72**
- Mycoplasma, infections, X-linked agammaglobulinemia, **59:146**
- Mycoplasma arthritidis*, mitogens, **54:113, 141**
- Myelin, Ig gene superfamily and, **44:37, 38**
 Myelin-associated glycoprotein, **46:171, 172**
 Ig gene superfamily and, **44:37, 38, 44, 45, 48**
- Myelin basic protein, **48:162, 163–166**
 α -acetylated amino-terminal peptide, **48:167**
 autoimmune demyelinating disease and, **49:357–363**
 multiple sclerosis, **49:369–370, 372–374**
 TCR usage restriction, **49:363–366**
- autoimmune thyroiditis and, **46:289**
 dominant epitope, **48:164–165**
 multiple sclerosis and, **66:88, 90**
 murine lupus models and, **46:79**
 specific tolerance induction, **48:167**
 synthetic T and B cell sites and, **45:209, 235–237**
- Myeloablative therapy, multiple myeloma, **64:250**
- Myeloblasts, leukocyte integrins and, **46:166**
- Myelocytes, eosinophil, morphology, **60:155–158**
- Myeloid cells
 B cell formation and, **41:237**
 B cell precursors, **41:189–191**
 bone marrow cultures, **41:209, 210, 212, 213, 219, 220**
 cyclic neutropenia, **41:229, 230**
 inducible cell line, **41:220**
 lymphohemopoietic tissue organization, **41:87**
 NZB mice, **41:229**
 SCID mice, **41:225**
 soluble mediators, **41:232, 233**
 W/W anemic mice, **41:224**
- CD1 protein distribution, **59:56**
 chemotactic molecules attracting, **58:380–381**
- cytolysis and, **41:285**
 helper T cell cytokines and, **46:114, 115, 117**
 hybrid resistance and, **41:373, 376**
 leukocyte integrins and, **46:150, 163, 165**
 surface antigens of human leukocytes and, **49:92, 111–112, 116**
- Myeloid-specific enzymes, **60:25–26**
- Myeloid zinc finger, **63:1, 239**
- Myeloma
 CD23 antigen and, **49:168**
 genetically engineered antibody molecules and, **44:89**
 antigen-combining sites, **44:84, 85**
 chimeric antibodies, **44:79, 80, 82, 87**
 expression, **44:70, 71, 75–79**
 fusion proteins, **44:85**
 lymphoid mammalian cells, **44:69, 70**
 production, **44:68, 69**
 human, **48:16–18**
 IgM, **48:16–18**
 human B cell neoplasia and
 chromosomal translocations, **38:247, 248, 254, 257, 259**
 c-myc gene expression, **38:260**
 t(11;14) chromosomal translocation, **38:264, 272**
- hybridomas
 genetically engineered antibodies and, **38:305**

- Myeloma (continued)**
- human-human, 38:292, 293, 296, 297, 300
 - human-mouse heterohybridomas, 38:300, 301
 - human-murine, 38:290-292
 - mAbs and, 38:306
 - IgE, ESR spectra, 48:30-31
 - Ig heavy-chain variable region genes and, 49:1-2
 - D segments, 49:33
 - V_H families, 49:6-7
 - V_H gene expression, 49:39-40, 46, 48-49
 - IgM, ESR spectra, 48:30-31
 - surface antigens of human leukocytes and, 49:98
 - Myeloma cells**
 - adhesion molecules in, 64:246-249
 - autocrine and/or paracrine growth, 64:257
 - B cell formation and, 41:220
 - FAS/APO-1 in, 64:259-260
 - hybrid resistance and, 41:378
 - identification of, 64:242-246
 - IL-6 production, 64:252-253
 - precursors, 64:249-250
 - tumors in idiotype immune mice and, 40:136-137
 - Myeloma protein, 48:2**
 - Ir* genes and, 38:45, 68
 - Myelopathy, HTLV-I associated, 50:182**
 - Myelopoiesis, 62:92-93**
 - B cell formation and, 41:186, 187, 213, 219
 - pituitary hormones, 63:404-405, 410-411
 - Myelosuppression**
 - colony-stimulating factor, 48:93
 - IL-6 treatment, 54:48
 - Myocardial infarction, aPL antibodies and, 49:219-220**
 - Myoglobin**
 - antigenic structure, 43:118
 - epitopes, 43:51-52, 55
 - humoral immune response and, 45:19
 - synthetic T and B cell sites and
 - globular protein antigens, 45:204, 211, 212, 215
 - peptides, 45:203, 240, 250
 - T cell receptor and, 45:123, 131, 134
 - Myohemerythrin**
 - antibody-antigen interaction, 43:42
 - antigenic response, 43:29-31, 47
 - antigenic sites, 43:80
 - critical residues, stereochemical relationships, 43:43
 - crystallographic structure, 43:21
 - electrostatic potential, 43:35
 - epitopes, 43:51, 53
 - features, 43:28
 - hexapeptide homologs, 43:32
 - hydrophilicity, 43:35
 - immune response, 43:28-29, 44-46, 75
 - mobility, 43:33-35
 - molecular surface chemistry and immunological reactivity, 43:33-34
 - monoclonal anti-MHr antibody affinity, 43:61-62
 - packing density, 43:33-35
 - peptides, monoclonal anti-MHr antibody affinity, 43:61-62
 - protein fold, 43:28
 - reactive sites, 43:29
 - secondary structure, 43:38
 - sequential epitopes, 43:36, 38-40
 - shape accessibility and exposed surface area, 43:35-36
 - side-chain contributions, 43:36-38
 - site 4-9, 43:37, 40, 43, 45
 - site 90-95, 43:45-46
 - Myosin**
 - cytolysis and, 41:275
 - hybrid resistance and, 41:378
 - Myositis, antinuclear antibodies and, 44:118, 119**
 - MZ2-E, 62:228, 231-232

N

- [(NAAG)₆] MAP, immunized mouse, for *P. malariae*, 60:123
- Na⁺/H⁺ antiporter, intracellular pH, 48:288
- L-NAME, see *N^G-Nitro-L-arginine methyl ester*
- NAP-2, conversion of CTAP III to, 50:242-243
- Naproxen, 62:184, 186
- Nasal polyps, endothelium, eosinophil adhesion to, 60:167-168
- Natural cytotoxic cells, 47:213, 286
 - hybrid resistance and, 41:373, 374

- Natural killer cell colony inhibiting activity, cachectin and, **42**:220
Natural killer cell cytotoxic factor, **47**:241, 250, 251, 253, 262, 302
Natural killer cell receptors activation, mediation, **55**:367–369 hybrid resistance, **55**:367 for MHC class I molecules, **55**:356, 358–360, 372 murine, **55**:364, 366–367 p58 dimers, **55**:358, 361–362 murine, **55**:364, 366–367
Natural killer cells, **47**:187–189 adaptive immunity B cell response, **47**:291–294 T cell response, **47**:294, 295 adoptive T cell therapy of tumors and, **49**:302, 315–319, 333 alterations, **47**:300–303 anti-CD2 mAb, **48**:236 antimicrobial activity antiviral activity, **47**:282–288 infection, **47**:288–291 antitumor activity cancer patients, **47**:297–300 experimental animals, **47**:295, 296 apoptosis, **58**:227 autoimmune thyroiditis and, **46**:279, 288, 289, 295 B cell formation and, **41**:192, 199, 225, 231, 236, 237 CD23 antigen and, **49**:155 cell-mediated cytotoxicity, **47**:189–196 clonal human cell lines of, **42**:184–186 CNS, **47**:266–269 congenital defects, **47**:224–226 cytotoxicity, **41**:269–273; **60**:289–291 cytolysis, **41**:275–277, 280–286 granule proteins, **41**:286, 287, 289, 291, 297, 298 membrane attack complex of complement and, **41**:309 defined allospecificities, genetic analysis, **55**:348–349 differentiation, **47**:229–234 effector mechanisms, **47**:234, 235 cytotoxicity, **47**:248–254 lymphocyte production, **47**:262–266 receptors, **47**:242–248 regulation, **47**:254–262 target cells, **47**:235–242 effects of MHC class I-deficiency, **55**:392–393 functions of, **42**:181–182 genetic control, **47**:222–224 granules in, **42**:183 helper T cell cytokines and, **46**:114, 116 hematopoiesis, **47**:272, 273 graft-versus-host reaction, **47**:280–282 inhibition, **47**:276, 277 regulation, **47**:273–276 soluble factors, **47**:277–280 heterogeneity of, **42**:182–183, 186, 188 HIV infection and, **47**:411 human, **55**:341–342, 370–373 allospecificity, genetic analysis of, **55**:348–349 CD3 gene expression during maturation, **55**:346 class I molecules, **55**:349–353 cytolysis, p58 in, **55**:354–357 MHC recognition by, **55**:363–365 ontogeny of, **55**:342–346 p58 modulation, **55**:362–363 repertoire, **55**:347–348 specificity, **55**:347 human clonal cell lines, **42**:184–186 humoral immune response and, **45**:68, 70, 83, 84 hybrid resistance and, **41**:334 antibodies, **41**:376–380 antigen expression, **41**:402–410 bone marrow cells, **41**:336, 338, 340, 351 effector mechanisms, **41**:372–376 leukemia/lymphoma cells, **41**:361, 363–369 lymphoid cells, **41**:354, 357 macrophages, **41**:371, 372 marrow engraftment, **41**:386–388 marrow microenvironment, **41**:390–393 syngeneic stem cell functions, **41**:388–390 T cells, **41**:382 IBD and, **42**:294–296 identification, **47**:196–199 IFN- γ production, **62**:63–66, 105, 106–107 IL-1 and, **44**:164, 166, 178 IL-10 biological effects, **56**:10

- Natural killer cells (*continued*)
 impaired activity in MHC class I-deficient mouse, **55**:404–405
 leukocyte integrins and, **46**:161, 163, 169
 lineage derivation of, **42**:186–188
 lymphocyte homing and, **44**:349, 350
 malignant expansion, **47**:226–229
 morphology, **47**:214–218
 murine splenic, perforin expression, **51**:220, 221
 pituitary hormones, **63**:416–417
 prolactin receptor, **48**:180
 regulators of complement activation and, **45**:395, 396
 reproduction, **47**:269–272
 SCID and, **49**:382, 396–397
 self-MHC class I recognition in tumor cells, **57**:315–317
 specificity, **42**:181
 specificity of, **42**:181
 stem cell factor receptor
 lymphohematopoietic effect, **55**:48–51
 stimulation by IL-2, **58**:426–429
 sulfasalazine and, **42**:317–318
 surface antigens, **42**:187, 204–206
 CD2, **42**:187, 194–198, 205
 CD3, **42**:187, 198–201, 205–206
 CD11a, **42**:187, 192–194
 CD16, **42**:187, 190–191
 HNK-1, **42**:187, 191–192
 NKH1 antigen, **42**:187, 188–190
 phenotype, **42**:185–186, 188
 TCR_{TAR}, **42**:198–204, 205
 TNK_{TAR}, **42**:201–202
 surface antigens of human leukocytes and, **49**:90, 94, 114–115
 adhesion molecules, **49**:83, 88
 antigen-specific receptors, **49**:79
 MHC glycoproteins, **49**:81–82
 surface phenotype, **47**:199–201, 207, 208
 experimental animals, **47**:210–214
 FcR antigen, **47**:201–205
 HNK-1, **47**:206, 207
 NKH-I antigen, **47**:205, 206
 T cell-associated antigens, **47**:208–210
 T cell activation and, **41**:9, 10, 24
 T cell subsets and, **41**:49
 tissue distribution, **47**:219–221
 virus-induced immunosuppression and, **45**:341, 357, 358, 360, 363
- Natural killer cell stimulating factor, **62**:105
 Natural killer cytotoxic factor, **41**:285, 296
 Natural resistance, hybrid resistance and, **41**:334
 NBD, *see* Nucleotide-binding domain
 NC41, **43**:21, 123–125
 N-CAM, *see* Neural cell adhesion molecule
 NDP52, **63**:202
 Necrosis, vs apoptosis, **58**:240
nef gene, HIV-1 disease, **63**:85–86, 94, 105–106
 Negative regulators, growth and differentiations of human basophils and mast cells, **52**:355–357
 interferons, **52**:355–356
 transforming growth factors, **52**:356–357
 Negative regulatory element, HIV infection and, **47**:381, 382
 Negative selection
 inhibition by FK-506, **58**:261–262
 T cells, **51**:165–175; **58**:166–168
 intrathymic clonal deletion, **51**:165–173
 cell biology of, **51**:165–170
 search for mechanism, **51**:170–173
 other mechanisms, **51**:173–175
 vs positive selection, **51**:180–183
 thymocytes, **58**:165–174, 171–174; **59**:105–107
 transgenic T cells, **58**:316
 Negative signaling
 blocking with antisense myc, **54**:413–414
 CD4 and, **53**:61
 CD8 and, **53**:61
Neisseria meningitidis, MAP vaccines, **60**:134–135
neo, **62**:36–38
 Neolactosylceramides, sulfoglucuronyl-containing, **58**:358
 Neonatal development
 anti-Id antibodies and, **42**:52–54
 B cells and, **42**:76
 clonotype repertoire acquisition and, **42**:26–30
 MG and, **42**:263
 preferential utilization of gene segments in mice and, **42**:132–133
 Neoplasia
 B cell, retrovirus-induced, in avian bursa of fabricius, **56**:467–481
 CD5⁺ B cell, V gene usage, **55**:323

- etiology, stem cell factor role, **55**:71–78
hybrid resistance and, **41**:387
immune regulation, **48**:212
and reactive hypereosinophilia, **60**:225
spontaneous, immune regulation, **48**:212
- Neoplastic cells, CD1 proteins, **59**:69
- Neostigmine, for MG, **42**:269–270, 271
- Nephritic factors, **61**:245–247
- Nephritis, *see also* Interstitial nephritis
 murine lupus models and, **46**:64
 regulators of complement activation and, **45**:389
virus-induced immunosuppression and, **45**:338
- Nerve growth factor, **48**:173–174; **61**:5
- Nerve growth factor receptor, family members, **52**:156
- Nervous system
 allergic reactions, **61**:364–368, 375
 autoimmune reaction, T cell component, **48**:163–167
 IL-1ra in, **54**:211–213
 stem cell factor receptor in, **55**:69–71
- Neural autoantigen, dominant epitope, **48**:164–165
- Neural cell adhesion molecule, **46**:171
 CD4 molecules and, **44**:288
 Ig gene superfamily and, **44**:36–38, 43–45, 48
 lymphocyte homing and, **44**:365
 multiple myeloma, **64**:247
 surface antigens of human leukocytes and, **49**:88, 114
- Neural cells
 Ig gene superfamily and, **44**:30, 35–38, 49
 immunological mediation, **48**:173–174
- Neuraminidase
 B cell antigen-presenting capacity and, **39**:74
 humoral immune response and, **45**:42, 44
 hybrid resistance and, **41**:400
 la structure and, **39**:74
 T cell subsets and, **41**:64, 91
- Neuroblastoma, structure and functional properties of associated gangliosides, **40**:355–357
- Neuroendocrine mediators, *see also* Neuromediators
 acute inflammation, **48**:175–176
 chronic inflammation, **48**:177–179
 immediate hypersensitivity, **48**:175–176
 immunological effects, **39**:301; **48**:175–179
 immunological generation, **48**:169–173
 lymphocyte function, **48**:177–179
- Neuroendocrine peptides, *in vitro* IgE synthesis, **61**:371–374
- Neuroendocrine system, **54**:30–31
- Neuroimmunology, **48**:161–184
 antigens and, **39**:299–300
 interleukins and, **39**:301
 neuroendocrine mediators and, **39**:300–301
 neuropeptides and, *see* Neuropeptides
- Neurological abnormalities, HIV infection and, **47**:377, 413
- Neuromediators, **48**:162, *see also* Neuroendocrine mediators
 cellular source, **48**:170–172
 identification method, **48**:170–172
 immunological sources, **48**:170–172
 lymphocyte receptors, **48**:167, 179–184
 structure, **48**:170–172
- Neuromuscular transmission, MG and, **42**:233–234, 236–237, 263, *see also* Acetylcholine receptor
- Neuronal cells
 IL-6 effect, **54**:30–31
 TE671 cell line, human, **42**:239
- Neuropeptidase
 cellular localization, **48**:167
 specificity, **48**:167
- Neuropeptide receptors
 immune cell, **39**:311–313
 lymphocyte, **48**:180
- Neuropeptides
 effects on nonneuronal tissues, **39**:301–304
 future research, **39**:318–319
 human lymphocyte receptor, **48**:180
 IL-1, **48**:178
 immediate hypersensitivity and, **39**:304–307
 immunopathogenic role, **39**:317–318
 inflammation and, **39**:307
 leukocyte chemotaxis and, **39**:307–308
 leukocyte-derived, **39**:315–317
 lymphocyte, **48**:177
 differentiation, **48**:178
 immunological functions, **48**:178
 tissue cycles, **48**:178
 tissue homing, **48**:178

- Neuropeptides (*continued*)
 lymphocyte functions and, **39**:309–311
 macrophage, **48**:177
 mediation by, immunological cell, **48**:176,
 177
 similarity to thymic factors, **39**:314–315
 structure, **48**:167
 tissue repair and, **39**:308–309
- Neuropsychiatric manifestations, HIV
 infection, **47**:395, 403–405
- Neurotoxin, eosinophil-derived, *see*
 Eosinophil-derived neurotoxin
- Neurotransmitter receptors, surface antigens
 of human leukocytes and, **49**:101,
 111
- Neurotrophic virus, **48**:165
- Neurotrophin-3, LNGFR recruitment of,
61:5
- Neurotrophin-4, LNGFR recruitment of,
61:5
- Neutralization
 aPL antibodies and, **49**:244
 apoptosis-inducing stimuli, **58**:242–243
 virus-induced immunosuppression and,
45:365
- Neutralizing antibodies
 colony-stimulating factor, **48**:95–96
 HIV infection and, **47**:407, 408, 410, 413
- Neutropenia
 agammaglobulinemia, **59**:146
 cyclic, B cell formation and, **41**:222;
 228–230; 234; 237
 hybrid resistance and, **41**:376
- Neutrophil-activating factor, **46**:115
- Neutrophil-activating peptide-1, **66**:135
- Neutrophil elastase, **62**:266
- Neutrophil receptors
 characteristics, **39**:96–98
 ligand interaction
 dynamics at 4°C, **39**:99–101
 dynamics at 15–25°C, **39**:101–102
 free radical production, **39**:105
 in permeabilized cells, **39**:109–110, 115
 pulse response analysis at 37°C
 antibody to ligand and, **39**:103–104
 occupancy response, **39**:102–104, 106
 static model of, **39**:98–99
 transiently active
 Ca²⁺ release and, **39**:126, 132–135
 cell signal generation and, **39**:110–111
- G protein interaction, **39**:113–116
 amplification during transduction and,
39:116–117
 cAMP elevation and, **39**:113–116, 118
 model of, **39**:115–116
 termination of, **39**:117–118
 potential functions, **39**:110–112
- Neutrophils, **51**:336
 adoptive T cell therapy of tumors and,
49:328
 arachidonic acid metabolism, human
 dietary EPA and, **39**:166–167
 esterified and nonesterified EPA and,
39:167–168
 B cell formation and, **41**:229, 230
 cachectin and, **42**:223
 chemotaxis, SP and, **39**:308
 complement receptor 1 and, **46**:190–193,
 198–200
 CXC proteins as activators, **55**:115–116
 cytotoxicity and, **41**:269
 decay-accelerating factor-deficient, **60**:79
 Fc receptors on, **40**:68
 FcRIII expression, **60**:179–181
 function, IBD and, **42**:307–308
 IBD and, **42**:285, 286, 287, 307, 308
 complement pathway products, **42**:309
 LTB₄, **42**:314–315
 IL-1_{ra} production, **54**:189–190
 IL-8 as agonist, **55**:112–113
 adhesion, **55**:114
 exocytosis, **55**:113
 lipid formation, **55**:114
 receptor up-regulation, **55**:113–114
 respiratory burst, **55**:114–115
 shape change, **55**:113
 immune reconstitution and, **40**:416–417
 interactions with endothelium, **60**:331–332
 interaction with IgA, **40**:169
 leukocyte integrins and, **46**:151, 177
 inflammation, **46**:160, 161, 163–169
 ligand molecules, **46**:173
 structure, **46**:154
 LTA hydrolase-containing, **60**:189
 neuropeptide production, **39**:317
 NK cells and, **47**:189, 190, 194, 201–204,
 260
 NO generation, **60**:352
 PNH, *PIG-A* abnormalities, **60**:81

- polymorphonuclear
interactions with endothelium,
60:331–332
as source of iNOS, **60**:329
recruitment by chemokines, **60**:339
regulators of complement activation and,
45:389, 390, 403, 410
rolling, selectin-mediated, **60**:165–167
senescent, **58**:270
as source of chemokines, **55**:106–107
as source of iNOS, **60**:329
surface antigens of human leukocytes and,
49:86, 88, 93, 112
TNF and IL-1 and, **66**:111
upregulation of CR3 expression, **60**:182
- Neutrophils, activation
amplification
Ca²⁺ release and, **39**:132–135
ligand binding rate and, **39**:133–134
oxidase activation, **39**:132
phospholipase C lifetime and, **39**:124,
132–134
sequence of events, **39**:131
termination steps, **39**:133–134
transient responses, **39**:132–133
branchpoints, of, **39**:134–135
Ca²⁺ release, **39**:124–127
future research, **39**:136
G proteins, linkage to
phosphatidylinositol turnover, **39**:119
phosphoinositide metabolism,
39:119–121
phospholipase C, **39**:119–120, 122
GTP and, **39**:99–100, 116, 120–121, 132
leukotriene low effects, **39**:135–136
phospholipase C role, **39**:119–120, 122,
124–125, 132–134
protein kinase C role, **39**:127–131
transduction sequence, **39**:111, 120–121
- Neurotrophic coronavirus, **48**:141
- Newcastle disease virus, **41**:166, 167, 389
lymphocyte homing and, **44**:366
- NF-AT, **63**:166–167
role in CD40L gene expression, **60**:47–48
- NFAT family, **65**:120
- NF-IL6, **58**:12–14; **65**:3–8
activation of, **65**:2–3
CYP2DG gene and, **65**:23
gene, **65**:2
- gene regulation, **65**:8–9
AP-1, interaction with, **65**:18–19
CREB, interaction with, **65**:19
cytokine induction, **65**:29–30
glucocorticoid receptors, interaction with,
65:21–22
Myb, interaction with, **65**:22
NF-κB, interaction with, **65**:16–18
PU.1, interaction with, **65**:21
STAT family, interaction with, **65**:19–21
knockout mice studies, **65**:29–30
structure and function of, **65**:1–2
transcriptional regulation, **65**:1–3, 8–11,
16–18
in viral infection, **65**:9–11
- NF-IL6β, **65**:3–6
- NF-IL-6/C/EBβ response elements, **62**:194
- NF-κB, **43**:250–251; **48**:298; **52**:271–272;
63:157–160; **65**:1, 111
activation, **58**:4–6; **65**:16–18, 130
inappropriate, **58**:14–18
by lipopolysaccharide, **60**:335
role of Fas, **60**:300
- antiinflammatory/immunosuppressive
molecule target, **65**:128–132
antioxidants, **65**:122–123
anti-TNF-α antibodies, **65**:123–124
cAMP, **65**:124–125
cyclosporin A and FK506, **65**:120
deoxyspergualin, **65**:125–126
gliotoxin, **65**:126
glucocorticoids, **65**:118–120
gold compounds, **65**:123–124
rapamycin, **65**:121
rheumatoid arthritis drugs, **65**:123–124
ROI-generating molecules, **65**:122–123
salicylates, **65**:121–122
spergualin, **65**:125–126
steroids, **65**:118–120
- description, **58**:2–4
disease and, autoimmune and
inflammatory, **65**:117–118
family, transcriptional regulation, **65**:11–12,
15–22, 30–32
- gene regulation, **65**:1, 26, 32, 111
cytokine induction, **65**:30–32
interaction with NF-IL6, **65**:16–18
glucocorticoid-mediated repression, **65**:23
induction, **65**:111
endogenous, **65**:114
exogenous, **65**:112–113

- NF- κ B (*continued*)
 as inflammatory mediator, 58:14
 inflammatory process, 65:111–118
 inhibition, 58:19–20
 in innate immunity in vertebrates, 58:1–27
 knockout mouse studies, 65:30–32
 physiologic inducers, 58:6–9
 structure and function of, 65:11–12
 target genes for, 65:115–116
 viruses and, 65:112, 127–128
- NF- κ B box, TAP genes, 65:60
- NF- κ B/p50, binding site, switch recombination, 61:117
- NF-M, 65:2–3, 7
- NF-S μ , binding site, 61:119
- Nicotinic AChR, *see* Acetylcholine receptor
- Nijmegen breakage syndrome, chromosome 7 and 14 translocations, 58:69
- Nippostrongylus brasiliensis*, 54:94–95
 helper T cell cytokines arc, 46:137
 infections, IL-4 production by Fc ϵ R⁺ cells in, 53:15–17
- Nitrate, urine
 L-NMMA effect, 60:343
 production in septic trauma patients, 60:351
- Nitric oxide
 in biological systems, 62:72–76, 99, 105, 264–265
 in immunoregulation, 60:348–349
 in inflammation
 effector molecule, 60:332–346
 protective molecule, 60:346–348
 in leukocyte adhesion and migration, 60:330–332
 in pain mediation, 60:349–350
 in vascular leak syndromes, 60:327–330
 in wound healing, 60:350–351
- Nitric-oxide synthase, 62:72–73
 constitutive, 62:72–73
 expression, role of IRF-1, 60:17
 inducible
 in human macrophages, 60:353–354
 regulation, 60:333–341
 isoforms
 in edema formation, 60:328–329
 expression in human tissues, 60:351–354
 in inflammation, 60:323–326
- Nitrite
 inhibition by L-NMMA, 60:353
 production by RAW 264.7 cells, 60:349–350
- N^G-Nitro-L-arginine methyl ester
 in conjunction with SEB, 60:348–349
 effect on aortic ring reactivity, 60:336–337
 infusion, and lymph flux increase, 60:329–330
 inhibition of F-BSA extravasation, 60:328
- Nitrogen, lymphocyte homing and, 44:354
- NK cells, *see* Natural killer cells
- NK-CIA, cachectin and, 42:220
- NKH-1, 42:188–190
 NK cells and, 47:188, 293, 294, 302
 CNS, 47:266, 267
 differentiation, 47:233, 234
 effector mechanisms, 47:261
 reproduction, 47:271
 surface phenotype, 47:205, 206
 tissue distribution, 47:219, 220
- NLDC-145, 59:379
- L-NMMA, *see* N^G-Monomethyl-L-arginine
- N-myc, 64:232–233
- NOD mouse, *see* Nonobese diabetic mouse
- Nonclustered antigens, B cell, 52:164–174
 adhesion molecules, 52:166–168
 activation of, 52:168
 Ig superfamily, 52:167
 integrin family, 52:166
 selection family, 52:167–168
- B7/BB1, 52:168–169
 cytokine receptors on B cells, 52:169–173
 IL-1, 52:169–171
 IL-2, 52:171–172
 IL-4, 52:172
 IL-6, 52:172
 other cytokines, 52:173
 TNF- α , 52:172–173
 transforming growth factor β , 52:173
- HLA class II molecules, 52:164–166
 other antigens of B cells, 52:173–174
 Bgp 95, 52:173
 CK226 antigen, 52:173
 IgM-binding protein (Fc μ receptor), 52:173
- Noncovalent bimolecular IL-2 receptor structure, 42:176
- Nongerminal elements, as V(D)J recombination products, 56:54–56
- Nonlymphoid cells, interaction with IgA, 40:168–184

- Nonobese diabetic mouse, **51**:286, *see also*
 Autoimmune diabetes, murine model
IDDM-prone, **48**:139
NOD.SCID, **63**:80
 strain development, **51**:287–289
Nonobese nondiabetic mouse, **51**:288
Non-self antigens, **64**:1, 2
 antibodies
 allergens, **57**:228
 bacteria, **57**:228
 Epstein–Barr-virus-transformed cell
 lines, **57**:229–230
 HuSCID mouse, **57**:230–231
 hybridomas, **57**:229–230
 overview, **57**:208–209
 primates, **57**:231–232
 staphylococcal protein A, **57**:229
 study of responses, **57**:232–242
 viruses, **57**:209–228
Nonspecific immune system, pituitary
 hormones, **63**:407–408
Nonspecific inflammation, IFN- γ and,
62:103–104
Nonsteroidal antiinflammatory drugs
 IBD and, **42**:313
 PGHS and, **62**:167, 184–187
 class I NSAIDs, **62**:184, 185–186,
 201
 class II NSAIDs, **62**:184, 185, 186
 class III NSAIDs, **62**:185
Normal rabbit gamma globulin, presentation
 by B cells, **39**:55–56
Northern blot analysis
 Ig gene rearrangements, **40**:269
 T cell receptor genes, **38**:11
NOS, *see* Nitric-oxide synthase
NP κ b, secondary B cell lineage and, **42**:70
N region diversity, T cell receptor,
58:301–303
NRGG, *see* Normal rabbit gamma globulin
NRI, concentration, **43**:265
NS398, **62**:186
NSAIDs, *see* Nonsteroidal antiinflammatory
 drugs
Nuclear factors, protein-binding sites, **43**:255
Nuclear matrix, **63**:201
Nuclease
 B cell-associated, as truncation factor in
 V(D)J joining, **56**:80
 sequence, **43**:209
Nucleic acids
 antinear antibodies and, **44**:119, 132,
 134
 murine lupus models and, **46**:76
Nucleolus, antinear antibodies and,
44:122–125, 133
Nucleoprotein
 synthetic T and B cell sites and, **45**:261,
 263
 antigens, **45**:234
 bacterial antigens, **45**:230
 viral antigens, **45**:217, 218
T cell receptor and
 antigen processing, **45**:119, 125
 homogeneity, **45**:128
 peptides, **45**:137
 structure-function relationships,
 45:139
Nucleosomes, antinear antibodies and,
44:98–100
Nucleotide binding, TAP, **65**:86
Nucleotide-binding domain, **65**:56
Nucleotides
 antinear antibodies and, **44**:98, 119, 130,
 135
 autoimmune demyelinating disease and,
 49:386
 B cell formation and, **41**:191, 194, 197,
 203
 CD4 molecules and, **44**:285, 288
 CD5 B cell and, **47**:144
 CD8 molecules and, **44**:274, 276
 complement receptor 2 and, **46**:201
 conjugate, **58**:322
 CTL and, **41**:159
 in gene rearrangements, **58**:32
 genetically engineered antibody molecules
 and, **44**:70, 73, 79
 HIV infection and, **47**:383, 384
 IgE biosynthesis and, **47**:16, 20
 Ig gene superfamily and, **44**:16–18, 48
 Ig heavy-chain variable region genes and
 D segments, **49**:28–29, 31, 33
 J μ segments, **49**:34
 polymorphism of V μ gene segments,
 49:22–27
 regulation, **49**:57
 V μ families, **49**:7–9, 16, 19
 V μ gene expression, **49**:39, 43, 48–49, 51
 immunosenescence and, **46**:223, 226, 227

- Nucleotides (*continued*)
 maps of Ig-like loci and MHC, **46**:31, 32
 structure, **46**:13, 15, 17, 20, 21, 25,
 27, 28
 T cell receptor, **46**:36, 39
 mismatched, **58**:44
 murine lupus models and, **46**:67
 NK cells and, **47**:202, 252
 nontemplated, **58**:308
 P (palindromic), **58**:36–38
 as V(D)J recombination product,
56:56–58
 SCID and, **49**:383
 T cell activation and, **41**:18, 26
 T cell receptor, **45**:109
 γ V-J junction sequences, **43**:145
 T cell subsets and, **41**:42, 43, 46, 47
 two, overlap internal to coding end, **58**:42
- Nucleotide sequence
 A/J *FLI6.1* *D_H* gene segment, **42**:105–106
 Ars A response and, **42**:101, 143
 Ig heavy-chain variable region genes
 D_H genes, **49**:28–33, 30
 J_H gene segments, **49**:35
 V_HI germline gene sequences, **49**:10–11
 V_HIII germline gene sequences,
 49:12–13
 V_HIV gene segments, **49**:16–17
 V_HV gene segments, **49**:18
 V_HVI gene segments, **49**:20
J558 *V_H* gene segments, **42**:112–113, 115
 surface antigens of human leukocytes and,
49:77, 90
V_H gene segments, **42**:102–105
 Nucleus, signal transduction, **48**:311
nur-77, encoding orphan steroid receptor,
58:229
Nur-77, nuclear hormone receptor, **58**:137
 Nylon fibers, IBD and, **42**:309
- O**
- Obese strain chicken, autoimmune
 cellular immune responses, **46**:298
 experimental models, **46**:273–276, 279
 genetic control, **46**:282–284
 humoral responses, **46**:301, 303
 prevention, **46**:310
 thyroiditis and antigens, **46**:269, 270
- Obese strain chicken, spontaneous
 autoimmune thyroiditis, *see* Spontaneous
 autoimmune thyroiditis
oct-1, **63**:243
oct-2, B cell genesis, **63**:220–221, 243–244
 Octameric sequence, *V_H* gene, **42**:113
 Oil granulomas, **64**:224–225
 Oligomeric proteins, buried surfaces, **43**:101
 Oligomerization, cytotoxicity and, **41**:306,
 315, 317, 318
 Oligonucleotide capture, in V(D)J joining of
 antigen receptor genes, **56**:62–63
 Oligonucleotides
 antinuclear antibodies and, **44**:108, 119,
 130
 antisense, **58**:249–250
 CD23 antigen and, **49**:162
 Ig heavy-chain variable region genes and,
 49:23, 27
 in ligation reactions, **58**:33
 synthesis, complement components and
 C1q B chain, **38**:232
 C4 genes, **38**:228
 Factor B and C2 cDNA, **38**:209
 Oligosaccharide rotation, **48**:26–27
 Oligosaccharides
 complement receptor 1 and, **46**:187, 192
 complement receptor 2 and, **46**:202, 204
 CTL and, **41**:135, 152, 153
 IgE biosynthesis and, **47**:16, 18, 19, 23
 leukocyte integrins and, **46**:154
 Omenn's syndrome, with V(D)J
 recombination defects, **58**:63–64
Onchocerca volvulus, major basic protein in
 patient tissues and, **39**:213
 Onchocerciasis, human, eosinophilic infiltrate,
60:213–214
- Oncogenes
 adjacent, effects of chromosomal
 abnormalities, **50**:124–132
 c-myc in Burkitt's lymphoma
 translocations, **50**:124–128
 gene fusion resulting from chromosome
 translocation, **50**:128–129
 helix-loop-helix oncogenes in
 chromosome translocations,
 50:129–130
 oncogene location after translocation,
 50:131–132

- transcriptional disruption of LIM domain oncogenes by translocation, **50**:130–131
- cellular
- c-myb*, rearrangement in myeloid leukemia, **39**:36–37
 - c-myc*, expression in IL-3-dependent cells, **39**:40, 42
 - complement receptor 1 and, **46**:187
 - complement receptor 2 and, **46**:210
 - derived tumor antigens, **57**:296–299
 - Ig gene superfamily and, **44**:39
 - LIM domain, by translocation, transcriptional disruption of, **50**:130–131
 - location after translocation, **50**:131–132
 - regulators of complement activation and, **45**:395
 - synthetic T and B cell sites and, **45**:248, 249
 - translocation, differentiation-related, **50**:139
 - viral
 - v-abl*, cell response to IL-2 and, **39**:41
 - IL-3 and, **39**:40–41
 - v-myc*, cell response to IL-3 and, **39**:40
 - virus-induced immunosuppression and, **45**:348
- Oncogenic proteins, **62**:237–238; **64**:65
- recognition by T cells, **58**:420–421
- Oncostatin M, **54**:15–16; **61**:151; **64**:256
- amino acid sequence, **53**:40
 - binding to high-affinity LIF receptor, **53**:41–43
 - and CNTF, **53**:44–47
 - and LIF, **53**:44–47
 - pleiotropic nature of, **53**:46
- One-dimensional isoelectrofocusing, HLA class II, **48**:114–116
- Ophthalmia, sympathetic, **48**:216, 217
- Ophthalmopathy, antibodies to self-antigens from human donors, **57**:245
- Opioid receptor
- human granulocyte, **48**:180
 - human lymphocyte, **48**:180
 - human monocyte, **48**:180
 - monocyte, **48**:180
- Opioids, NK cells and, **47**:268, 269
- Opsonic effects, fibronectin fragment, **38**:389–391, 393
- intact, **38**:388, 389, 393
- Opsonins, phagocytosis and, **38**:362, 392
- Oral tolerance, IgA and, **40**:223–226
- Organ transplantation, graft-versus-host disease, **59**:420–426
- Ornithine decarboxylase
- half-life, **48**:289
 - polyamine metabolism, **48**:289–290
- Orphans, **62**:10–12, 20, 22–24
- Orthogonal field electrophoresis, **38**:205
- OSM, *see* Oncostatin M
- Osteoarthritis, **54**:187
- cartilage destruction in, **64**:297
- Osteoblasts, LIF effects, **53**:36–37
- Osteoclasts, IL-1 and, **44**:154, 168, 195
- Osteopetrosis
- B cell formation and, **41**:232
 - hybrid resistance and, **41**:374, 390, 392, 393
- Ouabain, hybridomas and
- human-human, **38**:293–295
 - human-mouse heterohybridomas, **38**:300
 - human-murine, **38**:289
- Ouabain resistance, hybridomas and, **38**:280, 286, 301
- Outer membrane protein, *Chlamydia*, **60**:136
- Ovalbumin
- 17 amino acid peptide, binding to B cells, **39**:72–73
 - CTL and, **41**:167, 168
 - humoral immune response and, **45**:11
 - IgE biosynthesis and
 - antibody response, **47**:8
 - antibody response suppression, **47**:29, 30, 32–37, 39 - binding factors, **47**:27
- Ir* genes and, **38**:49, 55, 86
- antigen processing, **38**:95
 - competitive inhibition, **38**:109
- synthetic T and B cell sites and
- globular protein antigens, **45**:204, 210, 214–216
 - immunological considerations, **45**:200
 - viral antigens, **45**:228
- T cell receptor and, **38**:49, 55, 86
- alloreactivity, **45**:153
 - antigen processing, **45**:123, 125

- Ovalbumin (*continued*)

 experimental systems, **45**:131

 peptides, **45**:133
- T cell subsets and, **41**:55, 70

 urea-denatured, IgE biosynthesis and, **47**:29, 30, 33
- Ox40, **61**:7, 9

 OX40 gene, **61**:9

 Ox40-L, **61**:24–26

 OX62, **59**:380
- Oxidant formation, **62**:262–267

 Oxidation, periodate, as strategy for chemoimmunoconjugation, **56**:315–316
- Oxidative burst, surface antigens of human leukocytes and, **49**:111

 Oxygen radicals, **62**:72, 262–264
- P**
- P3C–MAP construct, HIV, **60**:133–134

 p21^{as}, B cell antigen receptor-induced activation, **55**:238–240

 p50, **65**:15

 B cell genesis, **63**:220–221, 245–246

 and p52, **58**:3

 p50 knockout mouse, cytokine induction, **65**:30–31

 p53

 association with apoptosis, **58**:138

 BCL-2, **63**:164

 dependent pathways in immunodeficiency syndromes, **58**:66–67

 role in PCD regulation, **58**:258

 p53, **62**:226–227, 237–238; **64**:251

 regulation, **65**:25

 p53/p56^{as}, CD45 and, **66**:24, 37, 39, 44

 p55, transcriptional regulation, **48**:294–295, 297–299

 p56^{as}, **64**:69

 binding to CD4 and CD8 cytoplasmic tails, **53**:61–64

 CD45 and, **66**:11, 12, 19–21, 26, 37–39, 39–44, 53

 interaction with CD4 and CD8 coreceptors, **58**:129–131

 phosphorylation of TAM, **58**:127–128, 130

 role in signaling through pre-TCR, **58**:150

 role in T cell activation, **53**:78–82

 structure, **53**:62–64
- T cell signaling by

 CD4 association, **56**:160–161

 CD8 association, **56**:160–161

 cell-surface receptor interactions, **56**:157–159

 cellular targets, **56**:159–160

 function, **56**:151

 lck expression regulation

 restricted accumulation of transcripts, **56**:151–152

 transcription, **56**:152–153

 regulation

 post-translational modification, **56**:156–157

 structure as factor, **56**:153–156

 T cell receptor interactions, **56**:161–162

 thymocyte development control

 allelic exclusion, **56**:164–166

 differentiation, **56**:162–163

 maturation, **56**:164–166

 modes of action, **56**:167–168

 T cell receptor β-chain, **56**:163–164

 and TCR-mediated signaling, **53**:85–91

 p58

 dimers, **55**:358, 361–362

 in modulation of NK-mediated cytosis, **55**:354–357, 362–363

 as NK receptor for MHC class I molecules, **55**:356, 358–360, 372

 p59^{as}, CD45 and, **66**:19–21, 37, 38, 39–43

 p70^{as}, located on human chromosome 22, **58**:57

 p70^{asK}, mTOR and, **61**:188–190

 p70^{asP}, CD45 and, **66**:21–22, 26

 p72^{asK}, CD45 and, **66**:21, 23–24, 26, 45

 p75 LNCFR, **61**:2, 4–5

 p95^{as}, B cell antigen receptor-induced tyrosine phosphorylation, **55**:243–244

 p105, **65**:15–16

 p150,95, leukocyte integrins and, **46**:149–153

 chromosomal localization, **46**:160

 inflammation, **46**:164, 167

 ligand molecules, **46**:173

 structure, **46**:153, 156, 158, 159

 PA28, **64**:21

 PA28α, **64**:21–24, 30

 PA28β, **64**:21–24

 PAC-1, **63**:167

 Packaging cells, development, **58**:424

 Packing density, **43**:33–35

- PAF, *see* Platelet-activating factor
 Pain, mediation with NO, 60:349–350
 Palmitylation, 52:61–62
 PALS, *see* Periarteriolar lymphoid sheath pan, 63:232
 Pancreas
 IFN- γ and, 62:100–101
 rat adenocarcinoma cell line, 54:324
 Pancreatic islets, grafts in MHC class I-deficient mouse, 55:390
 Pancreatitis, autoimmune, 62:98
 Pannus, 64:287–288
 Papillomavirus, *see also* Human papillomavirus
 tumor antigen induction, 57:292–293
 Paracrine effects, IL-1, 54:201–202
 Parallel tubular arrays, NK cells and, 47:216, 218, 256, 281
 Paralysis, autoimmune demyelinating disease and, 49:371–372
Paramecium
 antigenic variation in, 38:340
 maternally transmitted antigen and, 38:315
 Paraprotein
 aPL antibodies and, 49:242
 Ig heavy-chain variable region genes and, 49:35, 43, 46–48
 Parasites
 cytotoxicity and, 41:269
 and eosinophils, 60:210–216
 helper T cell cytokines and, 46:137–139
 human filarial, responses in MHC class I-deficient mouse, 55:415
 humoral immune response and, 45:84, 85
 infection
 IL-5 production in animal models, 57:172–173
 and IL-6, 54:36–37
 Parasitic antigens, synthetic T and B cell sites and, 45:228–230, 245
 Paratope, *see also* Complementarity-determining region
 Ir genes and, 38:107
 Paroxysmal nocturnal hemoglobinuria, 61:201, 253
 decay-accelerating factor and CD59 deficiencies, 60:75–78
 defective GPI synthesis, 60:78–80
 genetic basis, 60:80–91
 monoclonal or oligoclonal, 60:89
 regulators of complement activation and, 45:387, 400
 Partial lipodystrophy, 61:245
 Partial thromboplastin time, aPL antibodies and, 49:200
 Partial thromboplastin time with kaolin, aPL antibodies and, 49:201
 Particulate activators
 alternative complement pathway, 38:384–386
 monocyte ingestion, 38:366
 phagocytosis, 38:361, 362, 392, 393
 by human monocytes, 38:380–382
 by nonelicited murine macrophages, 38:383, 384
 Passive transfer
 AChR and
 B cell line, 42:260
 T cell clones, 42:258
 with Ig, 42:262
 MG, 42:234
 Pathogenesis
 aPL antibodies and, 49:193, 227, 238, 252–257
 autoimmune demyelinating disease and, 49:36O, 37O, 357–358, 368, 374
 CD23 antigen and, 49:174
 Ig heavy-chain variable region genes and, 49:49–50
 surface antigens of human leukocytes and, 49:81
 Pathogenicity, SIVmac and SIVsm molecular clones, 52:448
 Pathogens, responses in MHC class I-deficient mouse, 55:405–415
 Paw edema model, NO role, 60:328–329
 pax5, B cell genesis, 63:218–219, 228–232
 Pax-5/BSPAP, binding site, 61:120–121
 PBMC, *see* Peripheral blood mononuclear cells
 PC-1, a plasma cell threonine-specific protein kinase, 52:174
 PCD, *see* Apoptosis
 Peanut agglutinin
 antigen-presenting cells and, 47:86
 IgE biosynthesis and, 47:15, 18
 T cell subsets and, 41:97
 PEBP2 family, 65:24
 PEL-CTLs, Perforin expression in murine splenic, 51:220, 221

- Pemphigus foliaceus
antigen
 cadherin gene superfamily, **53**:310–312
 cDNA cloning, **53**:310–312
 desmosomal localization, **53**:300–303
 relationship to PV antigen, **53**:312–314
clinical presentation, **53**:295–297
histology, **53**:295–297
immunopathology, **53**:295–297
- Pemphigus vulgaris
antigen
 characterization, **53**:303–305
 identification as PF antigen-related
 cadherin, **53**:312–314
 sequence homologies and conserved sites
 with desmoglein, **53**:314
clinical presentation, **53**:295–297
histology, **53**:295–297
HLA class II peptide binding and, **66**:87
immunopathology, **53**:295–297
- Penicillamine, MG and, **42**:235, 251–252,
 253, 272
- Pentanoylsalicylate, **62**:187
- Pentoxifylline, sepsis and, **66**:130
- Peptide
 B cell formation and, **41**:215
 CTL and, **41**:136, 137, 167–169
 amino acid, **41**:162
 exon shuffling, **41**:148
 HLA antigens, **41**:152
 cytotoxicity and, **41**:279, 280, 309, 313
 pore formation and, **41**:313–319
 epitope vaccines, **43**:77–78
 α -helical conformation, **43**:203–204
 MHC interaction, rate constants for
 binding reaction, **43**:211
 monoclonal anti-MHr antibody affinity,
 43:61–62
 synthesis, **43**:77
 T cell activation and, **41**:3
 T cell subsets and, **41**:41, 47, 110
 H-2-restricted antigen recognition,
 41:52–55, 57, 58, 74, 75, 77
- Peptide add-back experiments, **59**:123
- Peptide antigens
 T cell specificity for, **43**:197–206
 in trimolecular complex
 features, **56**:239, 250–251
 MHC blockers, **56**:244–247
- peptide determinant as tolerogen,
 56:240–244
- T cell antagonists, **56**:244–247
- Peptide antigen X, RA susceptibility role,
 56:455
- Peptide binding
 to cell surface-associated class I molecules,
 mechanism, **52**:81–84
 to TAP, **65**:84–86
- Peptide-binding motifs, HLA class II
 molecules, **66**:69–71
- Peptide mapping, **43**:3
 blind study, **43**:26
 critical residue, **43**:74
 reconciling data, **43**:23–27
 sites mapped into conformational epitopes,
 43:38–39
- T cell receptor $\gamma\delta$ subunits, **43**:150
- Peptide–MHC complexes, **64**:2
- Peptide receptors, functionally active,
 expression on basophils and mast cells,
 52:367
- Peptides
 adoptive T cell therapy of tumors and,
 49:283, 319, 321–323, 334
 antigen-presenting cells and, **47**:87, 104
 antigen presentation, **47**:64, 66, 71, 74,
 76, 77, 79, 81
 antigen processing, **47**:87, 88, 90–92
 T cells, **47**:82, 96, 97, 99, 101
 antineuronal antibodies and, **44**:106, 129,
 134, 135
 autoimmune demyelinating disease and,
 49:357
 multiple sclerosis, **49**:369–371, 373–375
 myelin basic protein, **49**:357–361
 TCR usage restriction, **49**:363, 366
 autoreactive T cells and, **45**:425, 426
 bacterial formyl, attracting myeloid cells,
 58:381
 biologically active, from Fc region of Ig,
 40:111–119
 branched, synthesized on polylysine core,
 60:105–108
- CD4 and CD8 molecules and, **44**:269
 CD4 molecular biology, **44**:285, 286,
 289, 295, 296
 CD8 molecular biology, **44**:272, 274–276
- CD23 antigen and, **49**:156

- exogenous, association with class I molecules, **52**:78–86
 physiological relevance, **52**:85–86
 site of, **52**:79–81
- fMLP
 eosinophil response to, **60**:185–186
 induced eosinophil degranulation, **60**:173
- fusogenic, **58**:423
- genetically engineered antibody molecules and, **44**:67
- HIV infection and
 immune response, **47**:406, 410
 immunopathogenic mechanism, **47**:389, 391, 392
 neuropsychiatric manifestations, **47**:404
 humoral immune response and, **45**:7, 18–21
- IgE biosynthesis and
 antibody response, **47**:39
 binding factors, **47**:15–17, 20, 23, 25
- Ig gene superfamily and
 MHC complex, **44**:24, 26, 29
 nonimmune receptor members, **44**:32–34, 36, 42
- IL-1 and, **44**:164, 173, 190
 gene expression, **44**:160, 163
 receptor, **44**:179, 181
 structure, **44**:155, 157, 158
- immunogenic, conformation, **43**:63
- MHC class I molecules
 anchoring, **64**:124–127
 binding, **64**:114–115, 118, 127–128
 loading, **64**:105–128
- Multimeric-Synthetic Peptide
 Combinatorial Library, **60**:139
- multiple-antigen, *see* Multiple antigen peptides
- neuroendocrine, *in vitro* IgE synthesis, **61**:371–374
in vitro IgE synthesis, **61**:371–374
- NK cells and, **47**:202, 204, 268, 303
 P2 and P30, **60**:110–111, 121–122
 P12 and P14, **60**:127
- P30, purified from *T. gondii*, **60**:124
- positive selection, **59**:122–128
- precursor, to be GPI anchored, **60**:61–63
 production in cytosol, functional evidence, **52**:28–29
- regulators of complement activation and, **45**:389, 394, 399, 400
 role in positive selection, **58**:161–164
 SCID and, **49**:387
 sporozoite malaria vaccine and, **45**:320–322
 antibodies, **45**:304, 305
 CS proteins, **45**:296
 CS-specific T cells, **45**:307, 308, 310, 311
 human trials, **45**:314, 315
 immunity, **45**:287
 surface antigens of human leukocytes and, **49**:80–82, 112
 switch, **43**:113
- synthetic
 extracellular processing, **52**:84–85
 recognition sites defined by, *see*
 Recognition sites, synthetic technology, applications, **43**:59
- T cell development and, **44**:209, 218, 233, 253
- T cell receptor and, **45**:107, 108
 alloreactivity, **45**:148–157
 antigen processing, **45**:118–123, 125, 127, 128
 binding, **45**:132–134
 epitopes, **45**:142–148
 experimental systems, **45**:131, 132, 137
 homogeneity, **45**:129–131
 MHC molecules, **45**:112, 115, 116
 polymorphic residues, **45**:158–161
 selection, **45**:137–139
 structure-function relationships, **45**:135, 139–142
 T cell repertoire, **45**:162, 163, 165–167
- tumor-specific, T cell recognition, **57**:288–290
- virus-induced immunosuppression and
 human immunodeficiency virus, **45**:350, 351, 362, 363, 366–368
 measles, **45**:337
- Peptide supply factor gene, **65**:50
- Peptidoleukotrienes,, and asthma, **51**:341
- Peptides, or proteins, in entry of proteins into class I processing pathway, **52**:13–19
- Percoll density gradients, NK cells and, **42**:182, 184
- Perforin, **62**:43, 52–53; **64**:1, 69
 in cytolysis, **51**:215–216, 224–232
 perforin-independent mechanisms, **51**:231–232

- Perforin (*continued*)**
- verification in cell lines, **51:224–227**
 - verification by genetic approach, **51:227–231**
 - cytotoxicity and, **41:269, 270, 280, 281, 291, 305, 307–309, 319**
 - cell-mediated killing, **41:291–299**
 - cytolytic, **41:311–319**
 - membrane attack complex of complement, **41:299, 305, 307–309, 311**
 - effector molecule in cytotoxicity, **60:290–292**
 - in lymphocyte-mediated cytolysis
 - expression of, **51:219–224**
 - in human leukocyte, **51:222–223**
 - regulation of, **51:223–224**
 - perspectives, **51:232–233**
 - structure of, **51:216–219**
 - NK cells and, **47:251–254, 262, 291**
 - pathway of CTL-mediated cytotoxicity, **60:292–296**
 - verification by genetic approach, **51:227–231**
- Perforin-independent mechanisms, 51:231–232**
- Periarteriolar lymphoid sheath**
- associated foci, **60:270–271**
 - B cell activation in, **60:276**
 - containing interdigitating cells, **60:268–269**
 - T cells, **63:48, 71**
- Periodate oxidation, as strategy for chemoimmunoconjugation, 56:315–316**
- Peripheral blood**
- CD5 B cell and, **47:122, 131**
 - autoimmune diseases, **47:150**
 - marker for activation, **47:127**
 - surface antigen, **47:147**
 - eosinophil counts, **60:216–218**
 - HIV infection and
 - immune response, **47:411**
 - immunopathogenic mechanism, **47:394, 397, 400, 402, 403**
 - NK cells and, **47:289, 293, 298, 300, 303**
 - CNS, **47:266**
 - differentiation, **47:232–234**
 - effector mechanisms, **47:252, 254, 258, 264**
 - hematopoiesis, **47:275, 277, 279**
 - identification, **47:199**
- malignant expansion, 47:226**
- morphology, 47:216**
- surface phenotype, 47:201, 205, 206**
- spontaneous autoimmune thyroiditis and, 47:447**
- cellular immune reactions, **47:449, 451**
 - disturbed immunoregulation, **47:471, 473, 474**
- Peripheral blood lymphocytes**
- adoptive T cell therapy of tumors and, **49:16**
 - aPL antibodies and, **49:245**
 - autoimmune thyroiditis and, **46:287, 289, 299, 300**
 - CD5 B cell and, **47:144, 149**
 - CD23 antigen and, **49:166**
 - HIV infection and, **47:388**
 - in IBD, IBD and; **42:288–290, see also Lymphocytes**
- immunosenescence and**
- lymphocyte activation, **46:224, 226–230**
 - lymphocyte subsets, **46:238, 239**
 - stem cells, **46:239, 240**
- in MG, 42:259**
- NK cells and, 47:187, 294, 297, 301, 302**
- antimicrobial activity, **47:286–289**
 - cell-mediated cytotoxicity, **47:195**
 - CNS, **47:267**
 - cytotoxicity, **47:254, 258–261**
 - differentiation, **47:234**
 - effector mechanisms, **47:234, 240, 243, 246**
 - genetic control, **47:222**
 - hematopoiesis, **47:275, 276**
 - lymphokines, **47:264, 265**
 - morphology, **47:214, 217, 218**
 - reproduction, **47:270**
 - surface phenotype, **47:205–209**
 - tissue distribution, **47:219**
- SCID and, 49:385**
- spontaneous autoimmune thyroiditis and, 47:492**
- cellular immune reactions, **47:452, 453, 456**
 - disturbed immunoregulation, **47:471, 472, 475, 479**
 - effector mechanisms, **47:469**
 - genetics, **47:487, 488**
 - humoral immune reactions, **47:447, 449**

- Peripheral blood mononuclear cells, **61**:214
 CD23 antigen and, **49**:152, 157, 174
 HIV infection and
 activation, **47**:398, 400, 403
 immune response, **47**:411
 immunopathogenic mechanism,
 47:390–392
 human immunodeficiency virus and,
 45:350, 352, 354–362, 367
 IBD and, **42**:289, 294–297, 302
 IgE biosynthesis and
 antibody response, **47**:6, 7, 9, 12, 38
 binding factors, **47**:21, 22
 measles and, **45**:338, 340, 342, 346
 NK cells and, **42**:181–184
 PGE, IBD and, **42**:311–312
 PGE₂, corticosteroids and, **42**:316, 318–319
 phagocytic invasion, MG and, **42**:261
 phagocytosis, IBD and, **42**:307
 C3b and, **42**:308
- Peripheral immune system
 compartmentalization
 anatomic localization of colonizing
 lymphocytes, **53**:159
 B cells
 CD5⁺/CD5⁻ dichotomy, **53**:178–183
 in lymph follicles, **53**:185–186
 in peritoneum, **53**:183–185
 generation of compartments, **53**:187–194
 significance of, **53**:199–203
 site-dependent tolerance, **53**:201–203
 specialized functions of, **53**:199–201
 T cells
 intestinal intraepithelial, **53**:163–167
 γ/δ invariant intraepithelial cells,
 53:167–169
 in liver, **53**:169–171
 in lymph nodes, **53**:161–163
 in peritoneum, **53**:173–177
 in spleen, **53**:161–163
 IL-2 effects and toxicity, **50**:160–163
- Peripheral lymph nodes
 lymphocyte homing, **58**:390–391
 carbohydrate, **44**:354–359, 362–367,
 369, 371
 molecules, **44**:342–348, 350, 351, 353
 regional specificity, **44**:327, 328, 331,
 334–341
 selective homing model, **44**:323–325
 lymphocyte migration into, **64**:174–175
 lymphocyte recirculation, **64**:141, 142
- Peripheral node vascular addressins
 expression, **64**:149
 lymphocyte–HEV interaction, **65**:361–363
 ontogenesis, **64**:164
 in PLNs, **64**:174
 properties, **64**:147
- Peripheral T cells, *see* T cells, peripheral
- Peripolesis, spontaneous autoimmune
 thyroiditis and, **47**:442, 447, 469
- Peritoneal cavity, localization of CD5 B cells,
55:299–300
- Peritoneum
 B cells, **53**:183–185
 forbidden α/β T cells, **53**:166
 IL-1ra injection, **54**:212
 IL-1ra production by macrophages, **54**:188
 lymphokine production by Fc ϵ RI⁺ cells
 (murine), **53**:9–14
 T cells, characterization, **53**:173–177
- Peroxidase, eosinophil, *see* Eosinophil peroxidase
- Peroxidase active site, PHGS-2, **62**:176–177,
 187–188
- Pertussis toxin, **43**:291–292
 autoimmune side effects, **58**:278
 and G protein-mediated regulation,
 58:260–261
 suppression of cell death, **58**:246–248
- PERUN (software), T cell epitope prediction,
66:82
- Peyer's patches
 disassociated cells, studies with,
 40:195–197
 functional anatomy of, **40**:192–195
 immunosenescence and, **46**:236, 237, 251,
 252
 lymphocyte homing
 carbohydrate, **44**:362–371
 high endothelial venules, **44**:315, 317,
 318
 molecules, **44**:342–347, 353
 into mucosa, **64**:175–177
 regional specificity, **44**:322, 324–329,
 334–338, 341
- PF, *see* Pemphigus foliaceus
- Pf322, *P. falciparum* blood stage, **60**:120–121
- PGHS, *see* Prostaglandin endoperoxide H synthase
- P-glycoprotein, surface antigens of human leukocytes and, **49**:114–115

pH

- antigen-presenting cells and, **47:88**
- IgE biosynthesis and
 - antibody response, **47:8, 33, 34, 36**
 - binding factors, **47:12**
- Na^+/H^+ antiporter, **48:288**
- NK cells and, **47:263**
- spontaneous autoimmune thyroiditis and, **47:446**

Phagocytes, mononuclear, IFN- γ and, **62:78–80**

Phagocytic cells, interaction with IgA, **40:168–174**

Phagocytosis, **38:361, 392, 393**
adoptive T cell therapy of tumors and, **49:319, 323, 328**

B cell formation and, **41:212, 213**

CD23 antigen and, **49:167**

complement receptor 1 and, **46:183, 186, 198, 199**

by eosinophils, **39:208**

hybrid resistance and, **41:340, 357, 370**

Ir genes and, **38:94**

leukocyte integrins and, **46:151, 161, 163, 168**

by mononuclear cells, **38:363**

of particulate activators

- by human monocytes, **38:380–382, 392**

- by nonelicited murine macrophages, **38:383, 384**

pore-forming proteins and, **41:313**

surface antigens of human leukocytes and, **49:89**

surface changes resulting in, **50:62–63**

zymosan

- by human monocytes, **38:363–376**

- by nonelicited murine macrophages, **38:376–380**

Phenothiazine, aPL antibodies and, **49:212, 227**

Phenotype

adoptive T cell therapy of tumors and, **49:325, 333**

mechanisms, **49:299–300, 316**

principles, **49:290, 297**

antinuclear antibodies and, **44:109, 110**

autoimmune demyelinating disease and, **49:362–363**

autoimmune thyroiditis and

cellular immune responses, **46:287, 288, 290**

experimental models, **46:274, 275**

genetic control, **46:282**

prevention, **46:311**

B cell, developmentally regulated generation, **55:301**

B cell formation and

B cell precursors, **41:192, 201**

Ig genes, **41:204, 205**

lymphohemopoietic tissues, **41:187**

NZB mice, **41:227, 229**

W/W anemic mice, **41:2a4**

CD4 and CD8 molecules and, **44:266, 277, 279**

CD5 B cell and, **47:120, 122**

malignancies, **47:124**

physiology, **47:132, 135, 140, 142**

cytotoxicity and, **41:273, 287**

genetically engineered antibody molecules and, **44:70, 72**

helper T cell cytokines and, **46:111, 123, 126–128, 139**

HIV infection and, **47:384**

human B cell, **52:129–174**

B cell antigen receptor, **52:136–138**

Ig complex, **52:136–138**

proteins associated with Ig complex, **52:138**

non-Ig B cell surface antigens, **52:138–174**

antigens of plasmacytes, **52:174**

clustered antigens of B cells, **52:140–164**

non-Ig nonclustered antigens, **52:164–174**

adhesion molecules, **52:166–168**

B7/BB1, **52:168–169**

cytokine receptors on B cells, **52:169–173**

HLA class II molecules, **52:164–166**

plasmacyte, **52:174**

non-Ig nonclustered antigens, other, **52:173–174**

Bgp 95, **52:173**

CK226 antigen, **52:173**

IgM-binding protein (Fc μ receptor), **52:173**

- hybrid resistance and, **41**:338, 374, 382, 400, 401
IgE biosynthesis and, **47**:10, 12
Ig gene superfamily and, **44**:24, 29, 48
Ig heavy-chain variable region genes and, **49**:24
immunosenescence and, **46**:252
Ir genes and, **38**:61–64
antigen binding, **38**:115
cross-reactive lysis, **38**:186, 187
immunodominance, **38**:181
T cells
cytotoxic, **38**:163, 165, 167, 178
interaction, **38**:159
repertoire suppression, **38**:151, 152
tolerance, **38**:125
leukocyte integrins and, **46**:174–176
lymphocyte homing and, **44**:321
murine lupus models and, **46**:64, 82, 94
NK cells and, **47**:292, 298, 302
cell-mediated cytotoxicity, **47**:195
differentiation, **47**:229, 231–233
effector mechanisms, **47**:246, 260, 264
genetic control, **47**:223
hematopoiesis, **47**:274–277
identification, **47**:197–199
malignant expansion, **47**:226–228
reproduction, **47**:271
surface, **47**:199–214
tissue distribution, **47**:220
SCID and, **49**:381–385, 387, 391
lack of stem cell engraftment, **49**:396–397
posttransplant immunocompetence, **49**:400
T cell-depleted bone marrow, **49**:394–395
spontaneous autoimmune thyroiditis and, **47**:435–437
disturbed immunoregulation, **47**:470
genetics, **47**:484, 488
surface, murine CD5 B cell, **55**:306, 308–309
surface antigens of human leukocytes and, **49**:101
T cell activation and, **41**:1, 7, 28
T cell development and
cellular selection within thymus, **44**:254
ontogeny, **44**:220, 221
thymocyte subpopulation, **44**:225, 229–232, 235, 237–244
T cell subsets and
cell surface molecules, **41**:49, 50
H-2 alloantigen recognition, **41**:84, 87, 88
H-2 molecules in thymus, **41**:96
H-2-restricted antigen recognition, **41**:60, 69, 73
Phenotypic selection, germinal centers, **60**:280
Phenylalanine, CTL and, **41**:162, 166
Phenylbutazone, **62**:186
Phenylenediamine mustard, in cancer treatment, **56**:329, 331
2-Phenyl-4,4,5,5-tetramethylimidazoline-3-oxide-1-oxyl, interaction with NO, **60**:340
Philadelphia chromosome, **38**:245
Phorbol esters
adoptive T cell therapy of tumors and, **49**:307
B cell formation and, **41**:22a
CD4 molecules and, **44**:296
CD23 antigen and, **49**:151, 172
cell-mediated killing and, **41**:14–19, 24, 26
complement receptor 1 and, **46**:199
complement receptor 2 and, **46**:208
Ig gene superfamily and, **44**:34, 41
IL-1 and, **44**:174–176
leukocyte integrins and, **46**:162, 163, 165–169
T cell activation, **48**:258–260
T cell development and, **44**:236, 243
Phorbol myristate acetate, **48**:234
autoimmune thyroiditis and, **46**:296
CD4 and CD8 molecules and, **44**:268, 269, 298, 301
CD5 B cell and, **47**:125, 126, 141
immunosenescence and, **46**:224, 226, 228, 230
leukocyte integrins and, **46**:168
T cell activation and, **41**:8, 9, 11, 12, 14, 19
gene regulation, **41**:28, 30
receptor-mediated signal transduction, **41**:20, 22, 23, 25
T cell development and, **44**:226, 227, 229, 237, 241
Phosphatases
hematopoietic cell, role in cytokine signaling, **60**:25–26
T cell, signal transduction, **48**:284

- Phosphates
 CTL and, **41**:153
 T cell activation and, **41**:16, 24
- Phosphatidic acid, aPL antibodies and, **49**:194, 236–237, 242, 248
- Phosphatidylcholine
 aPL antibodies and, **49**:259
 biochemistry, **49**:194, 196–197
 LA antibodies, **49**:242–246
 specificity, **49**:234–237, 248, 251
 cytotoxicity and, **41**:275, 276
 IL-1 and, **44**:175, 176, 185
- Phosphatidylethanolamine, aPL antibodies and
 biochemistry, **49**:194, 196–197
 clinical aspects, **49**:202
 LA antibodies, **49**:242–248
 specificity, **49**:236–237, 251–252
- Phosphatidylinositol
 aPL antibodies and, **49**:194, 229, 251
 LA antibodies, **49**:242–243, 246
 specificity, **49**:236–237
 B cell formation and, **41**:197, 198, 202, 222, 239
 bone marrow cultures, **41**:215, 216
 leukocyte integrins and, **46**:171
 neutrophil
 cell activation and, **39**:122–124, 130
 G proteins and, **39**:119–120
 transfer of GlcNAc from UDP-GlcNAc, **60**:64–70
- Phosphatidylinositol biphosphate, T cell activation and, **41**:15, 16, 23, 25
- Phosphatidylinositol 4,5-biphosphate, **48**:228
 neutrophil
 amplification and, **39**:123–124
 metabolism, **39**:122–123
 phospholipase C and, **39**:122, 124–125
 T cell antigen receptor–CD3 complex
 Ca²⁺ response, **48**:262–265
 hydrolysis, **48**:262–265
- Phosphatidylinositol-4,5-biphosphate kinase, **48**:253
- Phosphatidylinositol 3-kinase, **54**:368–369; **61**:167; **63**:144, 149–154, 171
 activation, **61**:168–171
 B cell antigen receptor-induced activation, **55**:241–243
 cytokine-induced activity, **60**:15
 downstream effectors, **61**:171–173
- pathway, IR-2R-induced activation, **61**:167–173
- phosphorylation, **61**:28
- signal transduction, **59**:263
- Phosphatidylinositol 4-kinase, as p56^{ck} target in T cell signaling, **56**:160
- Phosphatidylinositol phospholipase C bacterial, **60**:60
 resistance of GPI anchor, **60**:61
- Phosphatidylinositol-specific phospholipase, B cell formation and, **41**:198, 214, 215
- Phosphatidylinositol-specific phospholipase C bacterial, **60**:60
 regulators of complement activation and, **45**:399
 resistance of GPI anchor, **60**:61
- Phosphatidylserine, aPL antibodies and, **49**:194, 259
 affinity purification, **49**:229–231
 antibody subsets, **49**:232
 clinical aspects, **49**:201, 204
 specificity, **49**:236, 240, 242–248, 251
- Phosphodiesterase type IV inhibitors, **64**:331
- Phosphodiester groups, aPL antibodies and, **49**:235, 237, 240, 248
- Phosphoinositide, **48**:228
 hydrolysis, **48**:253–254
 mast cell-specific receptor, **43**:290–293
 G protein role, **43**:291–292
 inositol phosphate role, **43**:292
 protein kinase role, **43**:292–293
 role in signal transduction, **54**:418–419
 turnover, T cell, **48**:258–265
 direct measurements, **48**:260
 secondary messenger formation, **48**:253–265
- Phosphoinositide cycle, **48**:253–255
- Phosphoinositide 3-kinase, **62**:133–135
- Phosphoinositide kinase type I, **48**:253
- Phosphoinositide kinase type II, **48**:253
- Phosphoinositide-specific phospholipase, **48**:256–258
 function, **48**:256–257
 G protein, **48**:266
 inositol 1-monophosphate, **48**:253
 lymphocyte, **48**:256
 multiple forms, **48**:256
 regulation, **48**:257–258
 structure, **48**:256–257

- Phospholipase
action of, **65:122**
aPL antibodies and, **49:235, 239, 246, 251**
B cell antigen receptor-induced activation of, **55:234–238**
B cell formation and, **41:198, 216, 239**
cytotoxicity and, **41:300, 314**
IgE biosynthesis and
 antibody response, **47:39**
 binding factors, **47:18, 24–26, 28**
NK cells and, **47:249**
SCID and, **49:386**
surface antigens of human leukocytes and, **49:111**
Phospholipase A, cytolysis and, **41:275, 276**
Phospholipase A2, pore formers and, **41:314–316**
Phospholipase C
activation, **54:365–367**
IgR complex to, **54:366**
immunosenescence and, **46:223**
leukocyte integrins and, **46:171**
neutrophil
 amplification and, **39:124, 132–134**
 G proteins and, **39:119–120, 122**
 phosphoinositide turnover and, **39:124**
 PIP₂ content and, **39:122, 124–125**
Phospholipase C_{y1}, as p56^{lck} target in T cell signaling, **56:159**
Phospholipase C_{y2}, phosphorylation, **61:28**
Phospholipids
aPL antibodies and, *see* Antiphospholipid antibodies
cytotoxicity and, **41:276, 300, 302, 303**
 pore formation, **41:314, 315, 318**
IgE biosynthesis and, **47:26**
immunosenescence and, **46:223**
T cell activation and, **41:15–19**
Phosphomannan polysaccharide, **65:356**
Phosphoprotein phosphatases, **63:164–167**
Phosphorylation, **52:62**, *see also* Tyrosine phosphorylation
 antigen-presenting cells and, **47:101**
 antinear nuclear antibodies and, **44:115, 120, 124**
 autoimmune thyroiditis and, **46:267**
 CD4 and CD8 molecules and, **44:269, 296, 300**
 CD23 antigen and, **49:156**
 CD45, **66:53–54**
 complement receptor 1 and, **46:187, 199**
 complement receptor 2 and, **46:202, 208, 209**
 CTL and, **41:135, 148**
 cytoplasmic tail of CD44, **54:281**
 cytotoxicity and, **41:273**
 in Fc receptor-mediated signal transduction
 Fc_YR, **57:74–77**
 Fc_εRI, **57:79–81**
 IκB by TNF α , **58:8**
 IgE biosynthesis and, **47:24**
 IL-1 and, **44:163, 184**
 immunosenescence and, **46:222**
 involvement, **52:90–92**
 leukocyte integrins and, **46:155, 169**
 lymphocyte homing and, **44:356–358**
 NF-κB, **65:15–16**
 NF-IL6 activation, **65:2–3**
 possible role, **54:311–313**
 protein kinases, in Fas ligand expression, **60:298–299**
 protein tyrosine, B cell antigen receptor-induced, **55:231–233**
 SCID and, **49:383**
 serine, **48:270–277**
 Src family kinases, **66:40–41**
 surface antigens of human leukocytes and, **49:117**
 adhesion molecules, **49:86**
 antigen-specific receptors, **49:76–77**
 complement components, **49:92**
 receptors, **49:98–100**
 TAM, by Src family protein tyrosine kinases, **58:127–128, 130**
T cell activation, **41:4, 16, 18, 19, 23, 24; 48:269–285**
threonine, **48:270–277**
transcription factors by DNA-PK, **58:60–62**
Phosphorylcholine
 electrostatic complementarity with antibody McPC603, **43:10**
 immunosenescence and, **46:248**
Phosphorylcholine response, **42:131**
 B cells
 BALB/c mice, **42:31**
 level, **42:65**
 secondary B cell lineage, **42:70**
 bone marrow isolates, **42:23–26**
Phosphotyrosine-binding domain, **61:161–162**
Ras/MAP kinase cascade, **61:161–162**

- Photoactivation, linkage procedure for chemoimmunoconjugates in cancer treatment, **56**:323
- Photochemical crosslinking, for lipid A receptor identification, **53**:274–279
- PhOX, B cell repertoire and, **42**:71
- Physical mapping, V_{II} locus, **62**:4–7
- Physiological abnormalities, IL-1 α blocking, **54**:205
- Physiological experiments, implying role for CD44, **54**:296–302
- in hematopoiesis *in vitro*, inhibition by CD44-specific Abs, **54**:299–300
 - inhibition of thymocyte progenitor migration, **54**:299
 - in lymphocytic migration *in vivo*, **54**:298–299
 - relationship to hermes antigen mediating PLN adhesion, **54**:296–298
 - signaling through CD44-ligand interactions, **54**:300–302
- Phytohemagglutinin, **48**:233; **54**:85
- autoimmune thyroiditis and, **46**:293, 310
 - B cell formation and, **41**:198
 - CD23 antigen and, **49**:150, 171
 - CTL and, **41**:50
 - HIV infection and, **47**:390, 398, 399, 401
 - hybrid resistance and, **41**:359, 361
 - polyclonal growth of human T cells, **42**:165
 - SCID and, **49**:386
 - spontaneous autoimmune thyroiditis and, **47**:492
 - cellular immune reactions, **47**:452, 455
 - disturbed immunoregulation, **47**:470, 471, 473
 - genetics, **47**:487
 - potential effector mechanisms, **47**:467
- T cell activation and, **41**:8, 14, 22, 26–28
- T cell subsets and, **41**:60, 63
- virus-induced immunosuppression and human immunodeficiency virus, **45**:356–358, 361–363
- measles, **45**:338
- PI, *see* Phosphatidylinositol
- PID, Ras/MAP kinase cascade, **61**:161
- PIG-A, **60**:67–71
- PIG-A*, **60**:57, 67–72, 80–91
- human class A gene, **60**:57, 67–72
 - mutant cells, clonal dominance, **60**:90–91
- and *Pig-a*, **60**:67–72, 93
- responsible for PNH, **60**:80–82
- somatic mutations, **60**:82–89
- PIG-B*, **60**:73
- Pigeon cytochrome c, *see* Cytochrome c
- PIG-F*, **60**:73
- pin-1*, and PCD, **58**:258
- Pinocytosis, B cell formation and, **41**:212, 213
- PIP₂, *see* Phosphatidylinositol 4,5-biphosphate
- Piroxicam, **62**:184, 186, 201
- Pit-1, **63**:381
- Pituitary adenoma, **63**:380
- Pituitary cells, thymic factor effects on secretion of
- ACTH, **39**:315
 - LH, **39**:315
- Pituitary gland
- anatomy, **63**:379
 - immune system, **63**:377, 433
- Pituitary hormone receptors, **63**:393–404, 426–427, 431–432
- Pituitary hormones, **63**:379–380, 459, *see also specific hormones*
- assays, **63**:429–430
 - autoimmune disease, **63**:420, 425–426, 433
 - depletion, chemical, **63**:430
 - hyperprolactinemia, **63**:409–410, 430
 - immune system, **63**:377–378, 407–408, 433
 - in vitro* effects, **63**:410–417
 - in vivo* effects, **63**:404–410
 - knockout animals, **63**:430–431
 - lymphohemopoietic system
 - growth hormone, **63**:386–388, 431–432
 - growth hormone receptor, **63**:396–397
 - IGF-I, **63**:389–393, 402–404, 431–432
 - IGF-II, **63**:417–419
 - as lymphohemopoietic growth factors, **63**:431–432
 - prolactin, **63**:382–384, 431–432
 - prolactin receptor, **63**:394–396 - lymphoproliferative diseases, **63**:426–429
 - mutant animals, **63**:430–431
 - production, **63**:430
 - secretion, **63**:382
 - as therapeutic agents, **63**:408, 433
 - transgenic animals, **63**:430–431
- Placenta
- development, role of IL-6, **54**:32
 - major basic protein production, human, **39**:188, 228

- Placental lactogens, **63:379**
- Plakoglobin, 130-kDa glycoprotein complexed with, **53:303–305**
- Plaque-forming cells
anti-BMRBC, **55:315–316**
autoimmune thyroiditis and, **46:299, 301**
CD5 B cell and
physiology, **47:130–132, 137, 139**
surface antigen, **47:144**
hybrid resistance and, **41:352**
immunosenescence and, **46:247, 248, 251**
spontaneous autoimmune thyroiditis and, **47:446**
- Plasma
antigen-presenting cells and, **47:45, 52, 53**
aPL antibodies, **49:259**
antibody subsets, **49:232**
binding to cell membrane, **49:252**
clinical aspects, **49:200–202, 206, 224**
isotype, **49:227**
pathogenic potential, **49:253–254, 257**
specificity, **49:240–242, 244–245, 247–248**
CD5 B cell and, **47:132**
genetically engineered antibodies, **44:76**
HIV infection and, **47:408**
IgE biosynthesis and, **47:4–6, 9, 13**
NK cells and, **47:267, 268**
regulators of complement activation, **45:387, 388, 397**
spontaneous autoimmune thyroiditis and
clinical symptoms, **47:440**
disturbed immunoregulation, **47:476, 479–481**
genetics, **47:488**
humoral immune reactions, **47:446, 447**
potential effector mechanisms, **47:467, 469**
- Plasmablasts, *In vitro* activated, **52:213–215**
- Plasma cell dyscrasias
amyloidosis, **64:219, 220–221**
defined, **64:219**
heavy chain diseases, **64:219, 220**
MGUS, **64:219, 220**
multiple myeloma, **64:219, 242**
adhesion molecules, **64:246–249**
cell identification, **64:242–246**
cytokines, **64:252–260**
IL-6 signal transduction therapies, **64:260–263**
oncogenic transformation, **64:249–251**
- plasmacytogenesis, **64:221–242**
chromosomal translocation, **64:227–229**
pristane-induced, **64:222–227**
spontaneous, **64:229–230**
susceptibility and resistance to, **64:238–242**
transgenic studies, **64:232–238**
virus infection induced by, **64:230–232**
- Waldenström's macroglobulinemia, **64:219–220**
- Plasma cells
distribution, **40:184–185**
humoral immune response and, **45:81–83**
lymphocyte homing and, **44:326, 327, 335, 336**
migration, **53:247–248**
precursors, mechanisms of homing to
mucosal tissues and secretory glands, **40:204–207**
surface antigens of human leukocytes and, **49:77, 98, 116**
- Plasmacyte foci, PALS-associated, **60:270–271**
- Plasmacytes, antigens, **52:174**
PC-1: threonine-specific protein kinase, **52:174**
- Plasmacytoma, **64:219**
adoptive T cell therapy of tumors and, **49:301, 305**
CD23 antigen and, **49:153**
cell line, dysregulated expression, **54:15**
helper T cell cytokines and, **46:116**
humoral immune response and, **45:74, 75**
hybrid resistance and, **41:367, 368**
IL-1 and, **44:178, 189**
plasmacytoma/hybridoma growth factor, **54:3**
- Plasmacytogenesis, **64:221–242**
chromosomal translocation, **64:227–229**
pristane-induced, **64:222–227**
spontaneous, **64:229–230**
susceptibility and resistance to, **64:238–242**
transgenic studies, **64:232–238**
virus infection induced by, **64:230–232**
- Plasmacytosis, polyclonal, Castleman's disease patients, **54:40–41**
- Plasma membrane
antigen-presenting cells and, **47:51, 68**
antineuclear antibodies and, **44:109**

- Plasma membrane (*continued*)
 aPL antibodies and, **49**:193, 197, 204,
 212–213
 CD8 molecules and, **44**:273
 complement receptor 1 and, **46**:191, 192,
 197, 199
 complement receptor 2 and, **46**:204, 209
 HIV infection and, **47**:393
 IL-1 and, **44**:163, 166, 175, 182, 184, 185
 immunosenescence and, **46**:224, 226, 235
 lymphocyte homing and, **44**:319, 368
 NK cells and, **47**:253
 P-selectin translocation, **58**:356
- Plasma membrane glycoprotein, **48**:161–162
- Plasmapheresis, for MG, **42**:271
- Plasmids, maps of Ig-like loci and, **46**:4
- Plasmin, lymphocyte homing and, **44**:347,
 348, 358
- Plasminogen, activation, **66**:241
- Plasminogen activator
 NK cells and, **47**:254
 tissue PA, **46**:166
- Plasmodium*, sporozoite malaria vaccine and,
45:296, 297, 299
- Plasmodium berghei*, **54**:36
 rodent malaria model, **60**:108–111
 sporozoite malaria vaccine and
 CS-specific T cells, **45**:308, 309
 immunity, **45**:285–290
 interferon-γ, **45**:302, 306
- Plasmodium falciparum*, **54**:36
 CS-specific T cells, **45**:308–311
 endemic areas, **45**:315, 316
 human trials, **45**:313
 immunity, **45**:285, 286, 288, 290, 292
 interferon-γ, **45**:306
 MAP candidate vaccines, **60**:112–122
 sporozoite malaria vaccine and, **45**:283,
 317, 320–322
 synthetic T and B cell sites and, **45**:229,
 258
 T cell receptor and, **45**:143
- Plasmodium malariae*
 response to MAP [(NAAG)₆]_n, **60**:123
 sporozoite malaria vaccine and, **45**:283,
 293, 294, 297
- Plasmodium vivax*
 circumsporozoite protein-based vaccines,
60:122–123
 sporozoite malaria vaccine and, **45**:283
 CS proteins, **45**:297–299, 317, 318
- CS-specific T cells, **45**:308–310, 313
 immunity, **45**:286, 288, 292
 interferon-γ, **45**:302
- Plasmodium yoelii*, **54**:36
 MAP immunogenicity, **60**:111–112
- Plasmacytosis, polyclonal, in patients with
 Castleman's disease, **54**:40–41
- Platelet-activating factor, **51**:352
 attracting myeloid cells, **58**:381
 CD23 antigen and, **49**:154, 159–161
 as eosinophil-derived lipid mediator,
60:190–191
 eosinophil receptors for, **60**:185–186
 IL-1 and, **44**:166, 171, 187
 induced neutrophil infiltration, **60**:176
 inducing granule protein release, **60**:207
 as neutrophil receptor, **39**:96–98
- Platelet-derived growth factor, **48**:308; **62**:268
 Ig gene superfamily and, **44**:39, 44, 45
 IL-1 and, **44**:153, 158, 164, 168, 169
 IL-6 production and, **66**:128
- Platelet-derived growth factor receptor,
 genes, and c-kit, evolutionary
 relationship, **55**:9–10
- Platelet factor 4
 biological effects on neutrophils, **55**:115
 chemical properties, **55**:97, 99–100
- Platelet neutralization procedure, aPL
 antibodies and, **49**:202, 244
- Platelets
 aggregation, PGHS-1, **62**:200–201
 aPL antibodies and, **49**:193, 197
 affinity purification, **49**:231
 binding to cell membranes, **49**:251–252
 clinical aspects, **49**:200, 202–203, 213
 pathogenic potential, **49**:254–257
 specificity, **49**:239, 241–242, 244–246
 CD23 antigen and, **49**:161, 167
 contact activation system, **66**:244
 Fc receptors on, **40**:68
 gpIIb-IIIa integrin, **54**:309
 IgE, **43**:306–307
 IL-1 and, **44**:153, 167
 SCID and, **49**:382
 surface antigens of human leukocytes and,
49:85–86, 88
- cis*-Platinum, *see* Cisplatin
- Platinum compounds, in cancer treatment,
56:329, 332

- Pleckstrin homology domain, Btk, **59:182–183, 185, 187–189, 191**
- Pleiotropic effects, TGF- β
 IL-4, **52:179–181**
 IL-6, **52:184**
 IL-10, **52:181–182**
- Pleiotropy
 LIF, OSM, CNTF, and IL-6, **53:44–47**
 reduction by compartmentalization, **50:163–165**
 IL-2 bioavailability, **50:164**
 IL-2 production, **50:163–164**
 IL-2 responsiveness, **50:164–165**
- PLNs, *see* Peripheral lymph nodes
- Pmel17, **62:229, 235**
- PML domains, **63:202**
- pml gene, **63:202, 203**
- PMN leukocytes, *see* Neutrophils
- Pneumococcus* polysaccharide, **48:28**
- Pneumonia, **59:397–398**
 bacterial
Legionella pneumophila, **59:398**
Listeria monocytogenes, **59:398**
Mycobacterium tuberculosis, **59:379, 383, 398–402**
Streptococcus pneumoniae, **59:395–396**
- fungal, **59:397–406**
Cryptococcus neoformans, **59:402–405**
- streptococcal, **59:395–396**
- viral
 influenza virus, **59:406–408**
 respiratory syncytial virus, **59:406–408**
- PNH, *see* Paroxysmal nocturnal hemoglobinuria
- P nucleotides, **58:36–38**
 as V(D)J recombination product, **56:56–58**
- Pocket specificity profiles, quantitative matrices and, **66:74–76**
- Pokeweed mitogen
 activation of B cells and, **40:13**
 B cells responsive to, **40:20, 23–29**
 complement receptor 1 and, **46:200, 201**
 hybridomas and, **38:297, 298**
 virus-induced immunosuppression and
 HIV, **45:356, 362**
 measles, **45:340–342, 346**
- pol β* , **61:303–304**
- Polarity
 cytolytic mechanisms and, **41:273–275, 282**
 T cell subsets and, **41:41**
- Poliovirus type II, mouse infection, anti-Ig antibodies as vaccines against, **39:283**
- Polyacrylamide gel electrophoresis, two-dimensional, HLA class II, **48:114–116**
 silver staining, **48:116**
 theoretical limitations, **48:116**
- Polyadenylation
 complement receptor 1 and, **46:184**
 regulators of complement activation and, **45:407, 408**
- Poly(ADP-ribose) polymerase, switch recombination, **61:110**
- Polyamine metabolism, **48:288–290**
 ornithine decarboxylase, **48:289–290**
- Polyarthritis, agammaglobulinemia, **59:146**
- Polyclonal activators, **52:195–199**
 IL-2, **52:195–196, 196–198**
 other stimulatory cytokines, **52:198–199**
 transforming growth-factor α , **52:199**
- Polyclonal B cell activation, **52:185–195**
 activated T cell induction of, **52:191–193**
 anti-CD40, **52:188–191**
 antigen receptor, **52:185–188**
 anti-Igs, **52:185–188**
 inhibitory effects, **52:186–187**
 stimulatory effects, **52:185–186**
Branhamella catarrhalis and others, **52:188**
Staphylococcus aureus Strain Cowan, **52:187–188**
- Epstein-Barr virus, **52:193–195**
 virus-induced immunosuppression and, **45:356, 357, 362**
- Polyclonal plasmacytosis, in patients with Castleman's disease, **54:40–41**
- Polyclonal stimulation, **42:19–22**
- Polycomb* group, **63:241**
- Polyglutamatic acid, anthracycline–antibody conjugation role in cancer treatment, **56:338**
- Polyinosinic:polycytidylc acid, hybrid resistance and, **41:371–372, 388–390**
- Poly(L-lysine), synthetic T and B cell sites and, **45:198, 199**
- Polymerase chain reaction
 autoimmune demyelinating disease and, **49:357, 368**
 Ig heavy-chain variable region genes and, **49:27, 35–36**
- leukocyte integrins and, **46:175, 176**
 quantitative techniques, **54:137**

- Polymeric antigens, and IgE responses, **52**:289–290
- Polymeric Ig receptor, **65**:249
- Polymerization
- aPL antibodies and, **49**:230
 - CD23 antigen and, **49**:168
 - cell-mediated killing and, **41**:290, 292–295, 298
 - complement components and, **38**:234
 - genetically engineered antibody molecules and, **44**:79
 - IL-1 and, **44**:179, 183
 - Ir* genes and, **38**:35, 37
 - membrane attack complex of complement and, **41**:304, 305, 309, 311
- Polymers, size-fractionated: Dintzis model, **52**:288–289
- Polymorphism
- autoimmune demyelinating disease and, **49**:365, 368–369
 - B cell formation and, **41**:193, 197, 223, 224, 229
 - CD1 proteins, **59**:46–47
 - complement components and, **38**:206, 236
 - of C2, **38**:208, 209
 - of C4, **38**:223, 225, 229–231
 - of C4b-binding protein, **38**:204
 - of Factor B, **38**:207–209, 227
 - restriction fragment length, **38**:215, 216
 - CTL and, **41**:135–137, 169
 - amino acid, **41**:158–163, 165
 - exon shuffling, **41**:141, 142
 - HLA class I antigens, **41**:151
 - mAbs, **41**:158, 165
 - hybrid resistance and, **41**:367, 400, 401, 405
 - Ig heavy-chain variable region genes and, **49**:1–3
 - J_H* Segments, **49**:34
 - V_H* families, **49**:9, 17–18, 20–21
 - V_H* gene expression, **49**:41, 48
 - V_H* gene segments, **49**:21–28
 - Ir* genes and, **38**:190
 - cytotoxic T cells, **38**:169
 - gene function, **38**:93
 - Ia molecules, **38**:68
 - T cell repertoire, **38**:145
 - maternally transmitted antigen, **38**:336, 347
- regulators of complement activation and, **45**:388, 396, 407
- sporozoite malaria vaccine and, **45**:311–313
- surface antigens of human leukocytes and, **49**:80–81, 92
- synthetic T and B cell sites and, **45**:229
- TAP, **65**:71–75
- T cell receptor, **38**:16; **45**:108
 - alloreactivity, **45**:149, 154, 156
 - homogeneity, **45**:129
 - MHC molecules, **45**:112, 115–117
 - peptides, **45**:132
 - residues, **45**:157–162
 - structure-function relationships, **45**:140
 - T cell repertoire, **45**:162, 167
- T cell subsets and
- cell surface molecules, **41**:41, 44–47
 - H-2 alloantigen recognition, **41**:182, 183
 - H-2-restricted antigen recognition, **41**:58, 59, 62
 - V_H* gene segments, **49**:22–27
 - V_H* locus, **62**:8–9
- Polymorphism analysis, **48**:114
- Polymorphonuclear leukocytes, *see* Neutrophils
- Polymyositis, antinuclear antibodies and, **44**:118–120
- autoantibodies, **44**:137
- autoimmune response, **44**:131, 132
- mixed connective tissue disease, **44**:111
- scleroderma, **44**:124
- Sjögren's syndrome, **44**:112, 113
- SLE, **44**:97, 109
- Polypeptide chain, AChR, **42**:238
- Polypeptides
- antigen-presenting cells and, **47**:82, 88
 - antinuclear antibodies and, **44**:120
 - autoantibodies, **44**:127
 - autoantigen, **44**:129, 131
 - autoimmune response, **44**:135
 - scleroderma, **44**:121, 123, 124
 - SLE, **44**:101
 - autoimmune thyroiditis and, **46**:272
 - B cell formation and, **41**:190
 - CD4 and CD8 molecules and, **44**:298
 - CD4 molecular biology, **44**:284–287
 - CD8 molecular biology, **44**:270–275, 277–280, 284
 - CD5 B cell and, **47**:142

- cell-mediated killing and, **41**:291, 292, 298, 313
complement receptor 1 and, **46**:184
complement receptor 2 and, **46**:204
CTL and, **41**:168
cytolysis and, **41**:276, 279, 285
cytotoxicity and, **41**:309, 319
genetically engineered antibody molecules and
 cloning, **44**:73
 expression, **44**:79
 fusion proteins, **44**:85, 86
 production, **44**:67–69
helper T cell cytokines and, **46**:124
IgE biosynthesis and, **47**:17–19, 34
IL-1 and, **44**:153
 human disease, **44**:193
 receptor, **44**:180, 183
 structure, **44**:155
TNF, **44**:185, 186
IL-5, molecular structure, **57**:149–151
immunosenescence and, **46**:229, 230
Ir gene control and, **38**:46–50
leukocyte integrins and, **46**:153, 156
lymphocyte homing and, **44**:342, 371
maps of Ig-like loci and, **46**:2, 46
 structure, **46**:7, 12, 20
T cell receptor, **46**:35, 41–43
NK cells and, **47**:202, 203, 208, 213
pore formers and, **41**:316, 317
regulators of complement activation and, **45**:392, 405
sporozoite malaria vaccine and
 CS proteins, **45**:294, 295, 300
 human trials, **45**:315
Plasmodium vivax, **45**:318
surface antigens of human leukocytes and, **49**:92, 94, 101, 112–113, 115
synthetic T and B cell sites and
 bacterial antigens, **45**:238
 prediction, **45**:251
 vaccine, **45**:253, 255
 viral antigens, **45**:221, 224
T cell activation and, **41**:10, 17, 23
T cell receptor and
 accessory molecules, **45**:109
 antigen processing, **45**:120
 homogeneity, **45**:130
MHC molecules, **45**:112
T cell repertoire, **45**:165
T cell subsets and, **41**:41, 43, 50, 51, 75
Polyperforin 1, cytotoxicity and, **41**:280, 289
Polyperforin 2, cytotoxicity and, **41**:280
Polyphosphomonoester, lymphocyte homing and, **44**:356–363, 370
Polyps, nasal, endothelial adhesion of eosinophils, **60**:167–168
Polyribosomes, B cell formation and, **41**:183
Polysaccharides
 CD23 antigen and, **49**:162
 lymphocyte homing and, **44**:320, 356, 357, 362
 surface antigens of human leukocytes and, **49**:87
Poly-2-vinylpyridine-N-oxide, hybrid resistance and, **41**:371
POMC, **61**:371
Population dynamics, B cell formation and, **41**:205–208
Population studies, RA susceptibility, **56**:425–427
Pore formation, cytotoxicity and, **41**:313–320
Pore-forming protein, *see* Perforin
Positional cloning, X-linked
 agammaglobulinemia gene, **59**:165–169
Position-specific antigens, leukocyte integrins and, **46**:153
Positive selection
 CD4⁺ cells, **58**:158–159
 definition, **59**:105, 110
 discovery, **59**:109–113
 epithelial cells, **59**:120–121
 genetically manipulated animals, **59**:113–118
 inducing cells, **58**:164–165
 ligand for, **59**:122–128
 ligands, **58**:160–164
 mechanism, **51**:175–189
 functional divergence, **51**:186–189
 ligands and receptors in positive selection, **51**:178–180
 molecular basis of four-way CD4 vs CD8 down-regulation, **51**:185–186
 developmental choice, **51**:180–186
 positive vs negative selection, **51**:180–183
 signaling pathways that control positive selection, **51**:183–185
 vs two kinds of death, **51**:175–177

- Positive selection (*continued*)
 thymocytes, **59**:99–129
 vs negative selection, **51**:180–183
 vs two kinds of death, **51**:175–177
- Post-translational modification
 antinuclear antibodies and, **44**:131
 lymphocyte homing and, **44**:370
 role in class I assembly and transport, **52**:59–62
 glycosylation, **52**:59–61
 palmitylation, **52**:61–62
 phosphorylation, **52**:62
- T cell development and, **44**:241
- Post-translational processing event, **48**:123
- Post-transplant lymphoproliferative disorder, **54**:35–36
- Potassium, cytotoxicity and, **41**:294, 318
- Potassium channels, **48**:287–288
- Poxviruses
 homologs, **61**:10
 virulence factors, **63**:308–309
- Praomys*, autoimmune thyroiditis and, **46**:273, 275
- Prasnitz-Kustner reaction, genetically engineered antibody molecules and, **44**:68
- PRE2*, **64**:20
- PRE3*, **64**:20
- Pre-B cells, *see* B cells, pre-B cells
- Precipitation
 epitope, **48**:23
 segmental flexibility, **48**:23–24
- Precursor cells, HIV infection and, **47**:395–397
- Precursor peptides, to be GPI anchored, **60**:61–63
- Prednisolone, IBD and, **42**:318–319
- Prednisone
 aPL antibodies and, **49**:223
 MG and, **42**:272
 prostaglandin synthesis and, **42**:312–313
- Pregnancy
 alloimmunization, human, anti-HLA antibodies and, **39**:281–282
 idiotype network, **39**:281–282
 and IL-10, **56**:17
 and preterm delivery, IL-1ra effects, **54**:216–217
- rheumatoid arthritis remission during, **56**:431
- Pregnancy zone protein, **38**:234, 236
- PRE-I, **65**:18–19
- Preimmune repertoire, self-tolerance checkpoints, **59**:281–319
- Prekallikrein, **66**:225
 activation, **66**:249
 function, **66**:236
 properties, **66**:226, 229
 structure, **66**:229
- Pre-pre B cells, **43**:237
- Preproenkephalin, helper T cell cytokines and, **46**:112, 116
- Pre-T cell receptor α , **62**:42, 44
- Pre-T cells, endothelium and migration of to thymus, **50**:275
- Preterm delivery, IL-1ra effects, **54**:216–217
- Primary amyloidosis, **64**:219, 220–221
- Primary antiphospholipid syndrome, **49**:257–259
 clinical aspects, **49**:209, 219, 221–222
 isotype, **49**:225–226
- Primary B cell repertoire, antiidiotypic recognition and, **42**:51–62, *see also* B cell repertoire expression
- Primary biliary cirrhosis, antibodies to self-antigens from human donors, **57**:244–245
- Primate lentiviruses, phylogenetic tree, **52**:444
- Primates, antibodies derived from, **57**:231–232
- Priming
 adoptive T cell therapy of tumors and, **49**:308–310, 321, 325, 329–332
 antigen-presenting cells and, **47**:65, 68, 71–73, 75, 80
 T cells, **47**:82, 83, 85, 86, 94
 tissue distribution, **47**:56
 autoimmune demyelinating disease and, **49**:370
 CD5 B cell and, **47**:137, 139
 hybrid resistance and, **41**:370
 IgE biosynthesis and
 antibody response, **47**:6, 29, 30, 32, 33, 36–39
 binding factors, **47**:14, 22–24, 26, 28
 T cell subsets and, **41**:72, 74, 100
- Primordial germ cells, LIF effects, **53**:34–35

- Pristane
 arthritis induction by, **64:289**
 plasmacytoma induction by, **64:222–227**
- Pro-B cells, *see* B cells, pro-B cells
- Probes, of T cell proliferations, **40:268–269**
- Procainamide, antinuclear antibodies and, **44:109, 111**
- Procarbazine serum, hybrid resistance and, **41:350**
- Programmed cell death, *see* Apoptosis
- Proinflammatory cytokines, **64:291–294**
 IL-6 effects, **54:27–28**
- Prokaryotes, maternally transmitted antigen and, **38:348**
- Prolactin, **63:379–380**
 activation of Jak2, **60:7**
 expression, **63:381–384**
 gene regulation, **63:400**
 hyperprolactinemia, **63:380, 430**
 immune system, **63:377–378**
 lymphohemopoietic system, **63:382–384, 431–432**
 regulation, **63:381–384**
 secretion, **63:382**
 structure and physiology, **63:380–381**
 in women, **63:380, 400, 409**
- Prolactinoma, **63:380**
- Prolactin receptor, **48:180–181; 63:394–396, 396–401**
 NK cell, **48:180**
- Proliferating cell nuclear antigen, antinuclear antibodies and, **44:94**
 autoantigen, **44:130**
 autoimmune response, **44:131, 134, 135**
 SLE, **44:95–97, 106–108**
- Proliferation
 adoptive T cell therapy of tumors and, **49:283, 334–335**
 antigen recognition, **49:319, 321**
 expression of antitumor responses, **49:325–330**
 mechanisms, **49:300, 302, 304**
 principles, **49:289–290, 294, 298**
 autoimmune demyelinating disease and, **49:359–360, 362, 364–366, 370**
- B cell formation and, **41:183, 184, 186, 187**
 B cell precursors, **41:192, 200, 202**
 bone marrow cultures, **41:210, 212**
 genetically determined defects, **41:225, 229, 231, 232, 234**
- inducible cell line, **41:220**
 population dynamics, **41:205–207**
- B cells, **40:6**
 antibody response and, **40:7–12**
 B cell subpopulations, **53:127–130**
 cytotoxicity and, **41:269**
 hybrid resistance and, **41:384, 390, 391, 397**
 bone marrow cells, **41:335, 336, 338, 340, 349**
 lymphoid cells, **41:352, 353**
 long-term, pro- and pre-B cells, **53:131–136**
 SCID and, **49:384–387, 390–391, 400**
 stem cell factor receptor-ligand interaction in, **55:39–40**
- T cell activation and, **41:1, 6, 7, 9, 10, 12–15, 26**
 T cell clone, **58:126–127**
 T cells, **51:106–110**
 anergy, **51:109–110**
 growth, **51:106–109**
- T cell subsets
 H-2 alloantigen recognition, **41:85, 86, 88, 92**
 H-2-restricted recognition of antigen, **41:51, 52, 60, 62–66, 70, 72**
- Pro-MCP, **61:216**
- Promonocytes, hybrid resistance and, **41:371**
- Promoters
 CD23 antigen and, **49:166–167**
 Ig heavy-chain variable region genes and, **49:53–56, 61**
 OTF2, **49:55**
 PGHS-1 and PGHS-2, **62:192–196**
- Promyelocytic leukemia cells, hybrid resistance and, **41:348**
- Proopiomelanocortin, **48:172–173**
- Properdin, **61:207, 241–245**
 sporozoite malaria vaccine and, **45:296**
- Propionylsalicylate, **62:187**
- Prostacyclin
 aPL antibodies and, **49:253, 255**
 IL-1 and, **44:160**
- Prostaglandin binding sites, **52:399–401**
- Prostaglandin E, release, Fc fragments and, **40:103–106**
- Prostaglandin E₂, **59:376; 62:198; 63:57–58**
 action of, **65:124–125**
 CD40 expression, **61:98**

- Prostaglandin E₂ (*continued*)
 production by RAW 264.7 cells, **60**:349–350
- Prostaglandin endoperoxide H synthase, **62**:192
 cellular and physiologic action of PGHS, **62**:196–200
 future work, **62**:201–202
 membrane-binding domain, **62**:172–173
 NSAIDs, **62**:184–187
 pathophysiologies, **62**:200–201
 regulation of gene expression, **62**:188–196
 structure–function relationships, **62**:169–178
- Prostaglandin endoperoxide H synthase-1
 catalysis by, **62**:178–188
 cellular and physiologic action, **62**:196–200
 future work, **62**:201–202
 gene structure, **62**:192
 mangano-heme PGHS-1, **62**:188
 NSAIDs, **62**:184–187
 pathophysiology, **62**:200–201
 platelet aggregation, **62**:200–201
 promoters, **62**:192–196
 prostanoid production, **62**:196–199
 regulation of gene expression, **62**:188–189
 structure–function relationship, **62**:169–178
- Prostaglandin endoperoxide H synthase-2
 catalysis by, **62**:178–188
 cellular and physiologic action, **62**:196–200
 cyclooxygenase active site, **62**:174–176
 future work, **62**:201–202
 gene structure, **62**:192
 inflammation, **62**:201
 NSAIDs, **62**:184–187
 pathophysiology, **62**:200, 201
 peroxidase active site, **62**:176–177, 187–188
 promoters, **62**:192–196
 prostanoid production, **62**:199–200
 regulation of gene expression, **62**:188–196
 structure–function relationship, **62**:169–178
 trypsin cleavage sites, **62**:174
- Prostaglandin E synthase, **62**:198
- Prostaglandin G, **62**:174
- Prostaglandin H₂, **62**:198
- Prostaglandin I₂ synthase, **62**:200
- Prostaglandins
 adoptive T cell therapy of tumors and, **49**:287
 B cell formation and, **41**:208, 236
 CD23 antigen and, **49**:160
 hybrid resistance and, **41**:360, 387
 IBD and, **42**:310–315
 IL-1 and
 biological effects, **44**:165, 166, 168
 gene expression, **44**:160, 161
 human disease, **44**:193, 195
 TNF, **44**:186–188
 NK cells and, **47**:250, 257
 production by IL-1ra, **54**:201
 SCID and, **49**:384
 and thromboxane, **51**:347–352
 biological activities, **51**:347
 measurements of prostanoids in biological fluids, **51**:349
 pharmacological modulation
 airway hyperresponsiveness, **51**:351–352
 airway tone and acute asthmatic response, **51**:350–351
 prostanoid action, **51**:350–352
 potency, **51**:347–349
 prostanoids and airway hyperresponsiveness, **51**:349
 synthesis and metabolism, **51**:347
- Prostanoids, **62**:167, 168, 196–200
 action, pharmacological modulation, **51**:350–352
 airway hyperresponsiveness, **51**:351–352
 airway tone and acute asthmatic response, **51**:350–351
 and airway hyper-responsiveness, **51**:349
 measurements of in biological fluids, **51**:349
- Prostate-specific antigen, **62**:224–225
- Protease inhibitors, prevention of DNA fragmentation, **58**:252
- Proteases, cytotoxicity and, **41**:282, 298
- Proteasome inhibitors, **64**:11–12
- Proteasomes, **64**:1–3, 8; **65**:15
 20S, **64**:9, 11, 12, 20, 24–25, 30, 105
 26S, **64**:9–11, 21, 25–26, 30, 105–107
 activator protein, **64**:21–24
 cleavage properties, **64**:24–26
 in cytosolic degradation pathwaypathway, **52**:25–28

- genes, evolution, **64**:19–21
 inhibitors, **64**:11–12
 LMP and IFN- γ regulation, **64**:12–14
 molecular adaptation, **64**:19–21
 structural and catalytic features, **64**:8–11
 subunits X/Y, **64**:14–16
 subunits Z and MECL1, **64**:16–18
 threonine residues, **64**:18–19
- Protective immunity, HIV-1 disease, **63**:95–98
- Protein antigen presentation, by cultured endothelial cells, **50**:265–266
- Protein antigens, **62**:221–225
- Proteinases, pulmonary inflammation, **62**:266–267
- Protein-binding sites
 B cell-specific, **43**:256
 enhancers containing, **43**:264
 fibroblast-specific site, **43**:256
 heavy and κ chain enhancers, **43**:243–244
 IgE, **43**:281–282
 multiple
 κ enhancer, **43**:250–251
 required, **43**:245–247
 in vitro mapping, **43**:244–245
 NF- κ B, **43**:250–251
 nuclear factors, **43**:255
 protein purification, **43**:247
 sequences, **43**:246
 site A, **43**:248
 site B/ μ E1, **43**:247
 site D, **43**:248
 site E, **43**:247
 switch recombination, **61**:117, 120
 E47/E12, **61**:117
 LR1, **61**:120
 NF- κ B/p50, **61**:117, 118–119
 NF- κ B/ μ , **61**:119
 Pax5, **61**:120–122
 S γ 3 regions, **61**:117–118
 S μ p-2, **61**:119
 switch regions, at or near, **61**:117–122
in vivo, **43**:243–244
 V β promoter, **43**:268–269
- Protein C, aPL antibodies and, **49**:253–256
- Protein-DNA complexes, **58**:56–57
- Protein electrophoresis, complement components and, **38**:225
- Protein export, mitochondrial, maternally transmitted antigen and, **38**:315
- Protein interaction domain, Ras/MAP kinase cascade, **61**:161
- Protein kinase, **48**:228
 antigen-presenting cells and, **47**:101
 Ca²⁺-dependent, **48**:274–275
 CaM-dependent, **48**:274–275
 cAMP-dependent, VIP effect, in lymphocytes, **39**:311
 CD4 molecules and, **44**:296
 CD5 B cell and, **47**:125
 cyclic nucleotide-dependent, **48**:275–277
 DNA-dependent, *see* DNA-dependent protein kinase
 in Fc receptor-mediated signal transduction
 Fc γ R, **57**:74–77
 Fc ϵ RI, **57**:79–81
 HIV infection and, **47**:382
 IgE biosynthesis and, **47**:26
 IL-1 and, **44**:175, 176, 179, 185
 mitogen-activated, *see* MAP kinase
 NK cells and, **47**:249, 271
 p34^{cdc}, cell cycle regulatory, blockade, **58**:267
 stress-activated, **63**:148
 surface antigens of human leukocytes and, **49**:77, 81, 83, 113
 T cell, signal transduction, **48**:284
 T cell development and, **44**:214, 226
- Protein kinase I α , interaction with CD3/TCR complex, **58**:232
- Protein kinase A, inhibition, **63**:51
- Protein kinase B, **63**:153–154
- Protein kinase C, **48**:79, 270–274;
63:154–157
 autoimmune thyroiditis and, **46**:296
 B cell formation and, **41**:222
 CD45, **63**:165
 cell-mediated killing and, **41**:296, 297
 complement receptor 1 and, **46**:187, 199
 complement receptor 2 and, **46**:202, 208,
 211
 growth hormone, **63**:399
 humoral immune response and, **45**:43
 IL-2 signaling, **63**:171, 172
 in IL-3-dependent cells, **39**:39
 immunosenescence and, **46**:222–227, 230,
 235
 isozymes, **54**:371
 leukocyte integrins and, **46**:169

- Protein kinase C (*continued*)**
- neutrophil
 - activation
 - Ca²⁺ and, **39**:130–135
 - DAG and, **39**:130–135
 - termination of, **39**:131
 - metabolism activation by, **39**:128–129
 - protein targets of, **39**:129
 - translocation, **39**:127–128
 - Ca²⁺ role, **39**:128–130
- plasma membrane kinase, **54**:370
- role in phosphoinositide hydrolysis, **43**:292–293
- role in thymocyte apoptosis, **58**:245–246
- signal transduction, **59**:263–264
- T cell activation and, **41**:8, 31
- Ca²⁺, **41**:15–19
 - gene regulation, **41**:28–30
 - receptor-mediated signal transduction, **41**:19, 20, 23, 25, 26
- T cell subsets and, **41**:65
- tyrosine kinase, role in IgR-associated transmembrane signaling, **54**:359–364
- Protein phosphatase, **48**:283–285**
- Protein–protein interactions, **43**:81, 265**
- buried surfaces, **43**:100–101
 - complementarity, **43**:101–103
 - conformational changes, **43**:103–105
 - free energy, **43**:100
 - structural complementarity, **43**:104–105
 - thermodynamics, **43**:100
- Protein S, **61**:237**
- Proteins**
- adoptive T cell therapy of tumors and, **49**:283, 334–335
 - antigen recognition, **49**:319–324
 - expression of antitumor responses, **49**:324, 331–332
 - antigen-presenting cells and, **47**:46, 64–73, 77
 - antigen processing, **47**:87, 88, 90, 91
 - cell surface, **47**:60
 - immunogeneity, **47**:104
 - T cells, **47**:86, 94, 99, 101
 - antinuclear antibodies and, **44**:93, 94
 - autoantibodies, **44**:125, 127, 137, 138
 - autoantigen, **44**:127–131
 - autoimmune response, **44**:137, 138
 - dermatomyositis, **44**:118–120
 - mixed connective tissue disease, **44**:111
 - scleroderma, **44**:121, 122, 124, 125
 - Sjögren's syndrome, **44**:112, 113, 115
 - SLE, **44**:98, 100–106, 108, 109
 - aPL antibodies and, **49**:195, 259
 - affinity purification, **49**:229–230
 - clinical aspects, **49**:200, 203, 213
 - pathogenic potential, **49**:253–254, 256–257
 - specificity, **49**:239–240, 247–248, 250–252
 - autoimmune demyelinating disease and, **49**:369
 - autoimmune thyroiditis and, **46**:265, 266, 315, 316
 - antigens, **46**:268, 272
 - cellular immune responses, **46**:294
 - experimental models, **46**:276
 - B cell formation and, **41**:205, 216, 239
 - B cell precursors, **41**:191, 197, 198
 - inducible cell lines, **41**:221, 222
 - biosynthesized, targeting to class I processing pathway, **52**:19–21
 - CD4 and CD8 molecules and, **44**:265, 266, 297, 298, 303**
 - CD4 molecular biology, **44**:285, 288–291, 297
 - CD8 molecular biology, **44**:271–274, 276–280, 284
 - CD5 B cell and, **47**:126**
 - Ig gene expression, **47**:150, 153, 154, 157
 - physiology, **47**:131, 134
 - primordial immune network, **47**:160
 - surface antigen, **47**:146
 - CD23 antigen and, **49**:176**
 - biological activity, **49**:169, 171
 - cellular expression, **49**:152–154
 - cleavage regulation, **49**:158
 - expression regulation, **49**:160
 - Fc ϵ RII, **49**:162–165, 167
 - complement receptor 1 and, **46**:185, 186, 190, 194, 197, 199
 - complement receptor 2 and, **46**:201–203
 - CTL and, **41**:135, 148, 153, 154, 159, 165, 168
 - cytotoxicity and, **41**:269, 270, 320
 - cell-mediated killing, **41**:289–295, 297, 311, 312
 - cytolysis, **41**:277–281, 316–319
 - mediation, **41**:270–273

- membrane attack of complement and, **41**:299, 301, 303, 307, 309–311
small peptides, **41**:314
degradation mediated by ubiquitin-proteasome system, **64**:6–8
entry into class I processing pathway, **52**:13–14
epitopes, **43**:2–4
extracellular matrix, interaction of CD44, **54**:290–291
folding, **43**:83, 100
fusion
expressing B cell lymphoma and
GM-CSF, **58**:435
L-selectin–IgG, **58**:353, 358–359
genetically engineered antibody molecules
and, **44**:89
antigen-combining sites, **44**:84, 85
chimeric antibodies, **44**:79, 80, 82, 87
cloning, **44**:75
expression, **44**:76, 78
fusion, **44**:85, 86
lymphoid mammalian cells, **44**:70
production, **44**:66–68
CPI-anchored, **60**:57–74, 79
granule
eosinophil, **60**:191–194, 205–208, 221
role in DNA fragmentation, **60**:300–301
helper T cell cytokines and, **46**:115, 116, 123
HIV infection and
activation, **47**:398–402
etiological agent, **47**:378–382, 384, 385
immune response, **47**:405–407, 409, 411
immunopathogenic mechanism, **47**:387, 392, 395
human counterpart of IgR-associated, **54**:356–357
human T cell activation and, **41**:1, 4, 8, 18, 23
gene regulation, **41**:29, 30
IL-1 receptor, **41**:13, 14
humoral immune response and
antigens, **45**:10, 14, 15, 17, 18
helper T cells, **45**:35
interleukins, **45**:72, 74, 75
hybrid resistance and, **41**:393
Ig complex-associated, **52**:138
IgE biosynthesis and, I
antibody response, **47**:2, 3, 11
antibody response suppression, **47**:31, 32, 34, 39
binding factors, **47**:16–18, 20, 22, 24–27
Ig gene superfamily and
evolution, **44**:43, 44
homology unit, **44**:5, 6, 8
MHC complex, **44**:24, 26, 28
nonimmune receptor members, **44**:30, 31, 34–37, 39–41
receptors, **44**:10, 11, 14, 16, 17, 21
Ig heavy-chain variable region genes and, **49**:1–3
D segments, **49**:33
regulation, **49**:53–57
 V_H families, **49**:6–9, 11, 15, 18
 V_H gene expression, **49**:39, 41, 43, 46, 51
IL-1 and, **44**:154, 155
biological effects, **44**:164–166
gene expression, **44**:159–161, 163
human disease, **44**:192, 194
immunocompetent cells, **44**:173, 177, 178
receptor, **44**:179, 182–184
IL-10-related, properties, **56**:3
immunogenicity, **43**:2
immunosenescence and, **46**:222, 223, 227–230, 234
interaction with Ig conserved
octanucleotide, **43**:254–255
isoform-specific TGF- β -binding, **55**:199
 κ B-dependent, inappropriate expression, **58**:17–18
leukocyte integrins and
inflammation, **46**:164, 165, 169
leukocyte adhesion deficiency disease, **46**:174, 175
ligand molecules, **46**:171
structure, **46**:153, 154, 158, 159
lymphocyte homing and
carbohydrate, **44**:354, 356, 364, 369, 370
molecules, **44**:342, 345–349, 351, 353
maps of Ig-like loci and
MHC, **46**:28–30, 32
structure, **46**:8, 10, 12, 13, 15, 17, 19
T cell receptor, **46**:45
molecule, **43**:118
murine lupus models and, **46**:69

- Proteins (continued)**
- naturally occurring, T cell specificity for linear sequences, **43**:198–201
 - NK cells and, **47**:284, 287, 291
 - cytotoxicity, **47**:252, 255
 - effector mechanisms, **47**:241, 242, 244–246
 - lymphokines, **47**:263, 265
 - surface phenotype, **47**:203, 205
 - octanucleotide, **43**:254–255
 - oligomeric, buried surfaces, **43**:101
 - oncogenic, **62**:237–238; **64**:65
 - recognition by T cells, **58**:420–421
 - outer membrane, *Chlamydia*, **60**:136
 - posttranslational modification, **64**:169
 - regulators of complement activation and, **45**:381–383, 410, 411
 - expression, **45**:394–403
 - genes, **45**:403, 405, 408
 - interactions, **45**:383, 384
 - roles, **45**:384–390
 - short consensus repeat, **45**:391–394
 - utilization, **45**:409, 410
 - retroviral, IgE-binding factor, **43**:304
 - SCID and, **49**:387
 - soluble TGF- β -binding, **55**:199–200
 - spontaneous autoimmune thyroiditis and, **47**:462, 476, 479
 - sporozoite malaria vaccine and, **45**:319–322
 - antibodies, **45**:300
 - CS proteins, **45**:292–300
 - CS-specific T cells, **45**:306–313
 - immunity, **45**:291, 292
 - interferon- γ , **45**:306
 - Plasmodium vivax*, **45**:317, 318
 - surface antigens of human leukocytes and, **49**:85, 90, 115–117, 125
 - antigen-specific receptors, **49**:76–77, 79
 - MHC glycoproteins, **49**:80, 82
 - in switch recombination, **54**:239–240
 - synthetic T and B cell sites and, **45**:195, 197, 259–264
 - antigens, **45**:234
 - globular protein antigens, **45**:203–216
 - immunological considerations, **45**:201, 202
 - parasitic antigens, **45**:228–230
 - peptides, **45**:203, 230–232, 237–239
 - prediction, **45**:250–252
 - vaccine, **45**:255–258
 - viral antigens, **45**:218, 219, 221, 223, 227, 228
 - T cell development and, **44**:232, 234, 241
 - T cell receptor and, **45**:107
 - accessory molecules, **45**:110
 - alloreactivity, **45**:149
 - antigen processing, **45**:118, 119, 124, 125, 127
 - epitopes, **45**:143–148
 - experimental systems, **45**:131, 132, 137
 - homogeneity, **45**:128, 129
 - MHC molecules, **45**:117
 - peptides, **45**:138, 139
 - polymorphic residues, **45**:160
 - structure-function relationships, **45**:139
 - T cell repertoire, **45**:165, 166
 - T cell subsets and, **41**:90, 97, 99
 - cell surface molecules, **41**:45–47, 50
 - H-2-restriction antigen recognition, **41**:52, 54, 56–58, 62, 70, 74, 75
 - T cell receptor, **41**:41–43
 - tumor antigens derived from
 - abl oncogenes, **57**:296–297
 - ber oncogenes, **57**:296–297
 - mutated p53, **57**:295–296
 - overexpression role, **57**:299–302
 - ras oncogenes, **57**:297–299
 - tum⁺, **57**:293–295
 - tyrosine phosphorylation, B cell antigen receptor-induced, **55**:231–233
 - as V(D)J joining factors
 - nonamer-binding, **56**:69–70
 - RBP-Jk, **56**:70–71
 - recognition protein, **56**:71
 - signal-binding, **56**:69
 - V(D)J joining protein, **56**:72
 - virus-induced immunosuppression and
 - HIV, **45**:348, 349, 361–363, 365–367
 - measles, **45**:337–339, 346
 - virus-produced, **58**:16–17
 - ζ and ϵ
 - invariant chains, **58**:127–128
 - and structure and assembly of TCR complexes, **58**:121–124
 - Protein synthesis
 - hepatic, cachectin and, **42**:225
 - maternally transmitted antigen and, **38**:348

- Protein-tyrosine kinase, **59**:178–185, 259–261, 265–266; **63**:51
activation by B cell antigen receptor components, **55**:248, 250–253
coreceptors, **55**:259–261
functions, **55**:255–256
signal initiation model, **55**:256–259
src family, regulation, **55**:253–255
adapter molecules, **63**:134, 141–143
CD23 antigen and, **49**:169
complement receptor 1 and, **46**:202
cytoplasmic, **59**:176–185
developmental expression, **54**:362
IL-2R activation, **61**:153–159
lymphoid tissue, **48**:277–280
phosphorylation, in Fas ligand expression, **60**:298–299
protein expression, **54**:416
regulate lymphocyte function, **59**:176–178
signal transduction, **59**:259–261
surface antigens of human leukocytes and, **49**:78, 99–100, 113
T cell, **48**:277–280
substrates, **48**:281–283
transforming protein, properties of *v-2bl* and, **37**:76–77
- Protein-tyrosine-phosphatase, **63**:164, 198
calcineurin, **63**:166–167
CD45, **63**:164–165
CD45R(0), and T cell apoptosis, **58**:246
negative regulation of cytokine signaling, **60**:25–26
PAC-1, **63**:167
SHPTP2, in cytokine signaling, **60**:26
- Protein-tyrosine-phosphatase 1C, **59**:299–300, 308–309; **63**:165–166
- Proteoglycans
cachectin and, **42**:224
CD44
binding to β -chemokine, **58**:383
role in leukocyte–endothelial adhesion, **58**:376–377
cell-mediated killing and, **41**:298
- Proteolipid protein
autoimmune demyelinating disease and, **49**:362, 374
multiple sclerosis and, **66**:88, 90
- Proteoliposomes, cytotoxicity and, **41**:274, 278, 303
- Proteolysis, **52**:22–35; **64**:3
additional factors in selectivity, **52**:34–35
antigen-presenting cells and, **47**:66, 87, 88
in antigen processing, functional evidence, **52**:24–25
antinear antibodies and, **44**:132, 135
background information, **52**:22–24
CD5 B cell and, **47**:130, 161
CD23 antigen and, **49**:156–158, 173
chemokines, **55**:102–103
cytotoxicity and, **41**:275, 276, 301, 305
functional evidence for peptide production in cytosol, **52**:28–29
functional evidence for proteolysis in antigen processing, **52**:24–25
genetically engineered antibody molecules and, **44**:66, 86
IgE biosynthesis and, **47**:16, 17, 20
IL-1 and, **44**:158, 162, 164, 188, 192
IL-8, **55**:102–103
lymphocyte homing and, **44**:320, 354
NK cells and, **47**:203, 250, 254
proteasomes in pathway, **52**:25–28
proteolytic machinery demonstrating selectivity, **52**:30–34
rasGAP, B cell antigen receptor-induced activation, **55**:239–240
role of class I molecules, **52**:29–30
spontaneous autoimmune thyroiditis and, **47**:461
surface antigens of human leukocytes and, **49**:90, 93, 112
T cell activation and, **41**:18
T cell subsets and, **41**:53
- Prothrombin, aPL antibodies and, **49**:231, 242–243, 248
- Prothrombin activator complex, aPL antibodies and, **49**:199, 241
- Prothrombin time, aPL antibodies and, **49**:200
- Protooncogene, T cell activation, **48**:291–294
- Prototypic sequences, Ig heavy-chain variable region genes and, **49**:6–7
- Protozoan diseases, **60**:108–124
multiple antigen peptides as immunogens, **60**:108–124
- Protozoan infections, X-linked agammaglobulinemia, **59**:146
- Provirus, **65**:143
Mtv, **65**:142–146

- Provoked immunity, **62:244**
- Proximal tubular epithelial cells, CD40 expression, **61:14**
- 4PS**, **63:168**
phosphorylation, **60:15–16**
- Psc** (posterior sex combs), **63:241**
- P-Selectin, **59:383; 62:276; 64:144, 145; 65:354, 355, 363, 367**
cooperative ligand binding site, **58:354–357**
expression, **64:148, 167**
genetic studies, **64:194**
ligands, **58:360; 64:150–152**
120-kDa, **58:361**
PSGL-1, **58:360–361**
properties, **64:147**
redistribution, **64:166**
shedding, **64:171**
synthesis, induction by alarm cytokines, **58:394**
upregulation, **64:165, 166**
- Pseudoallelism, Ig heavy-chain variable region genes and, **49:14**
- Pseudogenes, **42:138; 62:16–17, 24–26**
conserved, **62:16, 24**
diverged, **62:16**
- Ig gene superfamily and, **44:20, 30**
Ig heavy-chain variable region genes and, **49:57, 61**
D segments, **49:29**
 J_{H1} segments, **49:33–34**
organization, **49:3, 6, 8–9, 14–15, 17**
 V_{H1} gene expression, **49:39, 52**
 $J558 V_{H1}$, **42:115**
- Pseudogout, inflammation in, **66:260**
- Pseudo-light chain, **63:2**
- Pseudomonas* exotoxin A, **54:114**
- Pseudo T cell lymphoma, gene rearrangement and, **40:294–295**
- PSF1/2**, **64:4**
- psf1* gene, **65:50, 51**
- PSGL-1, **64:146, 150–151, 173**
- Psoriasis, and IL-6, **54:41**
- P/STEL sequences, PHGS-2, **62:177**
- Psychological factors, allergic reactions, **61:368–369, 374–375**
- pT α , **62:42, 44**
- PTB domain, Ras/MAP kinase cascade, **61:161–162**
- Pteridin metabolism, IFN- γ , **62:72**
- PU.1**, **64:86**
- PU.1
B cell genesis, **63:217, 218–219, 222–224**
binding sites, **64:76–77**
domains, **64:76–77**
expression, **64:85, 89**
gene targeting, **64:86–87**
interaction with NF-IL6, **65:21**
oncogenic potential, **64:65**
structure, **64:79**
structure–function studies, **64:85–86**
- PU.1P**, **63:217, 222**
- Pulmonary disease, *see* Lung disease
- Pulmonary embolism, aPL antibodies and, **49:219**
- Pulmonary eosinophilia, including Churg–Strauss syndrome, **60:226**
- Pulmonary fibrosis, bleomycin, **59:418–419**
- Pulmonary hypersensitivity, **59:387, 412–420**
- Pulmonary immunity, **59:369–427**
hypersensitivity lung diseases, **59:387, 412–420**
- transplantation, **59:420–426**
vaccination, **59:409–412**
- Pulmonary inflammation, **62:257–260, 290**
adhesion molecules, **62:267, 288**
allergic, **62:283–285**
animal models, **62:271–285**
complement cascade, **62:260–262, 271, 286–288**
- cytokines, **62:267–271, 280, 288**
and eosinophil increase, **60:219–220**
granuloma formation, **62:280–283, 290**
immune-complex mediated, **62:277–279**
ischemia/reperfusion, **62:275–277**
oxidant formation, **62:262–267**
proteinases, **62:266–267**
sepsis, **62:271–275**
therapeutic interventions, **62:259, 285–290**
- Pulps, white and red, splenic, **60:268–269**
- Pulsed-field gel analysis, Ig heavy-chain variable region genes and, **49:4–5**
- PUP1** gene, **64:20**
- Purified protein derivative
autoreactive T cells and, **45:419**
humoral immune response and, **45:16**
virus-induced immunosuppression and, **45:355, 356**
- Purine nucleotide phosphorylase deficiency, **59:246**

- Putrescine, **48:288**
PV, see Pemphigus vulgaris
 Pyridostigmine, for MG, **42:269–270, 271**
 Pyrogen, endogenous, cachectin as, **42:224**
 Pyrrolidine dithiocarbamate, **65:123**
- Q**
- Q10, **52:94–95**
 Qa-2, **52:95–98**
 QKRAA, Epstein-Barr virus
 Dw4 peptide, **48:130**
 gp 110 peptide, **48:130**
 Q region gene products, **52:94**
 Q10, **52:94–95**
 Qa-2, **52:95–98**
 Quantitative matrices
 algorithms based on, **66:81–82**
 peptide binding motifs by, **66:72–76**
 Quantitative regulatory mechanism, **48:123**
- R**
- RA, *see* Rheumatoid arthritis
 Rabbit, infection with HIV-1, **52:461–463**
 Rabbit anti-mouse Ig, *see* Immunoglobulin, rabbit anti-mouse
 Rabies, synthetic T and B cell sites and, **45:226, 244**
 Rabies virus
 AChR and, **42:272**
 nucleocapsid, **54:116–117**
 Race, *see also* Ethnicity
 as RA susceptibility factor, **56:427–429**
 Radiation chimeras, **59:106, 109–111**
 Radiautoautography, B cell formation and, **41:205**
 Radioactive antigen suicide, **42:268**
 Radioactivity, hybrid resistance and, **41:349**
 Radioimmunoassay
 IgE biosynthesis, **47:1, 20**
 IL-1 and, **44:160, 163, 192**
 Radioreceptor assay, IL-2, **42:166–167**
 Radioresistance, hybrid resistance and, **41:365, 376, 380, 383, 390, 393, 409**
 Radiosensitivity, hybrid resistance and, **41:350, 368**
 Raf, **63:144, 147, 399**
- Raf-1 kinase, signal transduction, **59:262–263**
 RAFT1, **61:181**
 RAG, **64:46**
 B cell development, **63:20–21**
 RAG1, **64:46, 48**
 RAG1
 aging and, **63:215**
 in rabbit B lymphopoiesis, **56:184, 186**
 as V(D)J joining agent, **56:64–67**
 V(D)J recombination, **64:45–47, 52–54, 59**
 RAG1/2, **61:123**
 active in immature lymphocytes, **58:311**
 mutations, **61:293–296, 317**
 SCID, **61:308–310**
 and signal joint product formation, **58:48**
 transcription by CD44⁺25⁺ cells, **58:141**
 transfection into XR-1 G12 cells, **58:53**
 transient transfection
 into Bloom syndrome cells, **58:65**
 into nonlymphoid cells, **58:51**
 and V(D)J recombination, **58:32–33**
 V-D-J recombination, **61:291–296**
 RAG2, **64:46, 48**
 RAG2
 deficient blastocyst complementation, **62:31–41**
 in rabbit B lymphopoiesis, **56:184, 186**
 as V(D)J joining agent, **56:66–67**
 V(D)J recombination, **64:46–48, 52–54, 59**
 Raji cells, complement receptor 2 and, **46:201–203, 206, 207, 209, 210**
 RalGDS, **63:144**
 RAMIG, *see* Immunoglobulin, rabbit anti-mouse
 RANTES, **62:285; 64:154**
 composed of basophil chemotaxins, **60:177–178**
 HIV and, **65:282**
 Rapamycin, **61:180–187, 182–183; 63:154**
 effect on eosinophil survival, **60:197–198**
 NF- κ B and, **65:121**
 RAPT1, **61:181**
 Ras, **62:42, 46; 63:143–147**
 GTP-bound, **60:209**
 IL-2 signaling, **63:171, 172**
 mutant, **58:417**
 prolactin signaling, **63:399**
 signaling pathway, activation by cytokine receptors, **60:12–14**
 signal transduction, **59:261–262**

- ras*, tumor antigens derived from, **57:297–299**
- ras*-GAP, as p56^{ck} target in T cell signaling, **56:159–160**
- ras*GAP-associated proteins, B cell antigen receptor-induced activation, **55:240–241**
- Ras-GTP, **63:147**
- Ras/MAP kinase, pathway, IL-2R initiation, **61:161–167**, **190**
- Rat
- arthritis model, **60:343–344**
 - chloroleukemia cell line, observation, **50:73**
 - IL-6 administration *in vitro*, **54:25–26**
 - mutations affecting stem cell factor or SCF receptor, **55:23**
 - paw edema model, **60:328–329**
 - RT-6 alloantigen, **48:241**
 - wound healing model, **60:350–351**
- Rat basophilic leukemia tumor, **43:282–283**
- RAW 264.7 cells
- lipid A binding sites, **53:278–279**
 - LPS-stimulated, **60:349–350**
- RB, **65:25**
- Rb, **62:42**, **48–49**
- Rb gene, **62:42**, **48–49**
- Rch1, **64:48**
- Reactive hypereosinophilia, and neoplasms, **60:225**
- Reactive oxygen intermediates
- cytosis and, **41:286**
 - NF-κB and, **65:122–123**
- Reactive oxygen species
- generation, inhibition by Bcl-2, **58:256**
 - IFN-γ, **62:72**
 - induction of NF-κB, **58:6–7**
 - pulmonary inflammation, **62:262**, **263**
 - role in PCD, **58:250**
 - scavenging by acute-phase proteins, **58:9**
- Reactive site
- association with electrostatic regions and surface grooves, **43:79**
- MHr, **43:29**
- stereochemistry, **43:32–39**
- electrostatic potential, **43:35**
 - hydrophilicity, **43:35**
 - mobility and packing density, **43:33–35**
 - secondary structure, **43:38**
 - shape accessibility and exposed surface area, **43:35–36**
 - side-chain contributions, **43:36–38**
- sites mapped by peptides cluster into conformational epitopes, **43:38–39**
- superassemblies, **43:74–75**
- Reading frames, preB cells expressing D_μ protein, **63:22**
- Reaggregate cultures, thymocyte maturation, **59:109**
- Reagin, aPL antibodies and, **49:258**
- biochemistry, **49:198**
 - specificity, **49:234–236**, **240**, **249–250**
- Rearrangement
- human B cell neoplasia and
 - Burkitt lymphoma cells, **38:257**
 - chromosome translocation, **38:267–269**,
 - 271**
 - c-myc* gene, **38:249**
 - c-myc* oncogene, **38:264**
 - Ir* genes and, **38:96**
 - maternally transmitted antigen and, **38:331**, **339**
 - V_α11, **60:278–280**
 - V(D)J region, **60:38**, **268**
 - V_{H6} gene, **60:47**
- Receptor affinity, B cell development and, **59:305–306**
- Receptor antagonists
- cytokines, **63:277**
 - function of IL-1ra as, **54:197–200**
- Receptor desensitization, **59:324–328**
- Receptor editing, B cells, **59:300–302**
- Receptor-mediated signal transduction, T cell activation and, **41:19–26**
- Receptors
- for activating peptides, **52:366–375**
 - formyl-methionine peptides and related compounds, **52:371–372**
 - IL-8 and related intercrines, **52:367–371**
 - substance P and other neuropeptides, **52:372–375**
- auxiliary signaling, **51:157–160**
- for growth and differentiating factors, **52:339–355**
- and ligands/counterreceptors, promiscuity in interactions between, **52:138**
- for low-molecular-weight regulators and pharmacological compounds, **52:396–404**
- adenosine, **52:396–398**
 - antiinflammatory drugs, **52:402–404**

- arachidonic acid metabolites, **52**:398–402
 other arachidonic acid derivatives,
52:401–402
 prostaglandin binding sites,
52:399–401
 histamine, **52**:402
 Receptor shedding, **63**:280
 Receptor subunit oligomerization, ligand-triggered, **61**:151
 Receptor tyrosine kinases, **52**:351–355
 other receptors of RTK family, **52**:354–355
 stem cell factor receptor: c-kit, **52**:352–354
- Recognition, **58**:124–128
 antigen-specific, T cell cultures for,
42:165–166
 antiidiotypic, B cell repertoire expression
 and, **42**:51–62
 leukocyte–endothelial cell, models,
58:381–389
 target, TCRs and, **38**:22–24
 Recognition molecules, Ig superfamily,
52:364
 Recognition phase, T cell activation,
63:43–44
 Recognition sites, synthetic, T and B cell,
45:159–164, 195–197
 B cell sites, synthetic peptides and,
45:237–250
 candidate synthetic peptide vaccines,
45:252–259
 immunological considerations, **45**:197–202
 prediction
 B cell, **45**:250, 251
 T cell, **45**:251, 252
 T cell sites, synthetic peptides and, **45**:202,
 203
 antigens, **45**:232–237
 bacterial antigens, **45**:230–232
 globular protein antigens, **45**:203–216
 parasitic antigens, **45**:228–230
 viral antigens, **45**:216–228
 Recombinant DNA
 and carcinogenesis, **58**:417
 mAbs and, **38**:304
 Recombinase activating gene, *see* RAG
 Recombination
 complement components and, **38**:229,
 230
 deletional switch, in B cells, **60**:38–39
 double-homologous, **48**:56
 experiments, Ars A response and, **42**:141
 frequency, V_{H} complex and, **42**:137
 gene inactivation, **59**:108
 homologous, *see* Homologous
 recombination
 Ig heavy-chain variable region genes and,
49:3, 6, 9
 D segments, **49**:28, 30–32
 polymorphism of V_{H} gene segments,
49:22, 24, 26
Ir genes and
 antigen binding, **38**:98
 complementation, **38**:79, 80
 cytotoxic T cells, **38**:166
 immunodominance, **38**:180
 I region mutations, **38**:86
 MHC, **38**:45, 73
 T cell suppression, **38**:145, 148
 maps of Ig-like loci and, **46**:3, 4, 47
 structure, **46**:19–21, 23
 mechanisms, joining of segments of Ig
 variable region and, **40**:253–256
 murine lupus models and, **46**:87
 nonhomologous, and end-joining,
58:43–45
 switch, *see* Switch recombination
 tolerance, **38**:129
 V(D)J, *see* V(D)J joining, antigen receptor
 genes; V(D)J recombination
 XLA, **59**:161–162
 X-linked agammaglobulinemia, **59**:161–162
 Recombination recognition sites, in V(D)
 joining of antigen receptor genes, **56**:86
 Recombination signal sequences, **64**:39
 junctions, and coding junctions, **58**:42, 46,
 53–55
 in V(D)J recombination, **58**:31–36
 Reconstitution, *see also* Immune
 reconstitution
 IL-2 receptors, **59**:231–234, 254
 Recruitment
 neutrophils, by chemokines, **60**:339
 ras exchange factor, by GRB2, **60**:13
 Stats, to cytokine receptor complexes,
60:22–23
 Rectal dialysis, lipoxygenase products and,
42:314–315
 Rectal mucosa, estimate of prostaglandin
 syntheses by, **42**:311–312
 Red blood cells, *see* Erythrocytes

- Redundancy
 functional, defined, **63:173**
 killing mechanisms, **60:303**, 311
 LIF, OSM, CNTF, and IL6, **53:44–47**
- Reed Sternberg cells, CD40 expression, **61:12**, 39
- Regulation of complement activation
 complement receptor 1 and, **46:189**
 complement receptor 2 and, **46:202**
- Regulators of complement activation, **45:381**, 410, 411; **61:207**
 complement pathways, **45:381–383**
- genes
 divergent function, **45:408**, 409
 exon structure, **45:404**, 405
 gene duplication, **45:405**
 intragenic duplication, **45:406**, 407
 organization, **45:403**, 404
 polyadenylation, **45:407**, 408
- protein expression
 alternate RNA processing, **45:398**, 399
 anchoring systems, **45:399**, 400
 differential glycosylation, **45:400–403**
 tissue-specific, **45:394–398**
- protein interactions, **45:383**, 384
- protein roles
 autologous tissue, **45:386**, 387
 history, **45:384**, 385
 immune complexes, **45:388**, 389
 plasma, **45:387**, 388
 transmembrane functions, **45:389**, 390
- RCA-like protein utilization, **45:409**, 410
- short consensus repeat
 common structural motif, **45:391**, 392
 localization, **45:392–394**
- Regulatory DNA sequences, in type I
 interferon genes, **52:265–267**
- Reiter's syndrome, HLA-B27 and, **65:94**
- REKS, **63:144**
- rel*, retrovirus-induced B cell neoplasia in
 bursa of fabricius, role in, **56:477–478**
- Rel
 and ankyrin repeats, **58:3–4**
 B cell genesis, **63:245–248**
 in innate immunity in vertebrates, **58:1–27**
 subunits, and promotion of HIV-1
 transcription, **58:15**
- relA*, **63:159–160**
- RelA/p65 knockout mouse, cytokine
 induction, **65:31**
- Rel-B
 B cell genesis, **63:220–221**, 246–248
 deficiency, **63:159**
- RelB knockout mouse, cytokine induction, **65:31–322**
- REL family, **65:12–13**
 transcriptional regulation, **65:12–13**
- Repair, double-strand break, *see* DNA,
 double-strand break repair
- Repair synthesis, across double-strand gap, **58:44**
- Reperfusion
 injury due to, antiadhesion therapy, **64:196**
 leukocyte integrins and, **46:165**, 166, 177
 pulmonary inflammation, **62:275–277**
 therapeutic complement inhibition using
 soluble CR1, **56:282–284**
- Repetitive amino acid interchanges, Ars A
 response and, **42:110–111**
- Replacement therapy, intravenous Ig, **60:40**
- Replication
 B cell formation and, **41:183–185**, 236
 B cell precursors, **41:200**
 bone marrow cultures, **41:210**, 213, 216,
 217
 genetically determined defects, **41:228**,
 231
 population dynamics, **41:206**, 207
 soluble mediators, **41:232–234**
- genetically engineered antibody molecules
 and, **44:72**
- maps of Ig-like loci and, **46:4**
 and V(D)J joining of antigen receptor
 genes, **56:80–81**
- Reproduction
 clusterin, **61:250–251**
 membrane cofactor protein, **61:219–220**
 NK cells and, **47:269–272**
 PGHS-1, **62:191**
- Reproductive tract, female
 invariant intraepithelial γ/δ T cells,
53:167–169
- invariant γ/δ T cell receptors, **58:303**
- Rescue
 CD4 $^+$ 8 $^-$ cells, **58:158**
 cells in thymic cortex, **58:154–155**
 IL-2-dependent cells, **58:244**
scid coding junction products, **58:48–49**
 signal, **58:241**
 thymocytes, from PCD, **58:165**

- Reshaped antibody, human PM-1, construction, **64:261**
- Resistance, to retrovirus-induced B cell neoplasia, **56:470–471**
- Respiratory burst, induction by IL-8 in neutrophils, **55:116–117**
- Respiratory syncytial virus, **59:406–408** antibodies from combinatorial libraries, **57:217–219**
- Respiratory tract
 immunity mechanisms, **59:371–373**
 lung diseases
 bacterial pneumonia, **59:379, 383, 395–402**
 fungal pneumonia, **59:402–406**
 graft-versus-host disease, **59:420–426**
 hypersensitivity diseases, **59:387, 412–420**
 viral pneumonia, **59:406–409**
 lung transplantation, **59:420–426**
- Responding cells, **50:29–31**
- Response genes, **51:98–105, 146–157** characteristics of, as defined by cloned lines, **51:100**
combinatorial regulation of, **51:105**
IL-2 gene regulation, case study, **51:103–105**
inducibility, **51:151–157**
 cortical loss of responsiveness in, **51:153–156**
 emergence from paralysis to functional maturity, **51:156–157**
 immature thymocytes, early inducibility of responsive genes in, **51:151–153**
proliferation, **51:146–151**
T cell subsets, **51:98–103**
- Restitope, *Ir* genes and, **38:107**
- Restriction enzymes, maps of Ig-like loci and, **46:3, 4**
- Restriction fragment length polymorphism, **48:117–119**
 allele-specific oligonucleotide, **48:117–119**
analysis, pitfalls, **48:117**
- Ars A response and, **42:107**
- autoimmune thyroiditis and, **46:284**
- complement receptor 1 and, **46:191**
- Ig heavy-chain variable region genes and, **49:4, 22–23, 25, 27, 43**
- leukocyte integrins and, **46:160**
- maps of Ig-like loci and, **46:23**
- murine lupus models and, **46:66, 67**
TaqI fragment, **48:139**
- Restriction mapping, Ars A response and, **42:106**
A/J *J558 V_H* gene segments and, **42:101–102**
- Reticular cells, lymphocyte homing and, **44:316**
- Reticulo genesis, SCID and, **49:382, 389**
- Reticuloendothelial hyperplasia, cachectin and, **42:222**
- Reticuloendotheliosis virus-induced tumors, **50:107–108**
- Retina, organ-resident nonlymphoid cell, **48:218**
- Retinal antigen
 autoimmuneresponse, **48:217**
 specific T cell clone, deletion, **48:217, 218**
- Retinal glial cell, T helper lymphocyte, **48:218**
- Retinoblastoma gene, **62:42, 48–49**
- 9-cis-Retinoic acid
 PCD inhibition, **58:245**
 targeted to steroid receptor family, **58:259**
- all-trans-Retinoic acid, IL-6 signal transduction modulation, **64:261–262**
- Retinol, *see* Vitamin A
- Retroviral protein, IgE-binding factor, **43:304**
- Retroviral switch recombination vector, **61:123–124**
- Retroviruses, **50:40–41**
 adoptive T cell therapy of tumors and, **49:282, 320, 335**
 mechanisms, **49:313–314**
 principles, **49:292, 295**
- antinuclear antibodies and, **44:130**
- autoimmune thyroiditis and, **46:275**
- B cell formation and, **41:195, 233**
- CD4 and CD8 molecules and, **44:269, 284, 290, 291**
- endogenous, insertions, inseparable from Vbse, **50:38–39**
- endogenous or extrinsic, **50:37–42**
- endogenous retroviral insertions
 inseparable from Vbse, **50:38–39**
 expressions of Mtv in B cells with Vbse expression, **50:39**
- mice transgenic for given Mtv, **50:40**
- mouse mammary tumor virus (MMTV), **50:39**

- Retroviruses (*continued*)
 mouse retention of Vbse sequences, **50:41–42**
 Mtv transcripts expressed in activated B cells, **50:40**
 other retroviruses, **50:40–41**
 retroviral sequences bearing no homology to bacterial toxic mitogen, **50:41**
 genetically engineered antibody molecules and, **44:68**
 hybrid resistance and, **41:378, 400, 403**
 IgE biosynthesis and, **47:17**
 induction of B cell neoplasia in avian bursa of fabricius, *see* Bursa of fabricius
 infection, as substrate introduction method in V(D)J joining, **56:50–51**
 integration of viral DNA, **65:168**
 NK cells and, **47:303**
 origin of endogenous superantigens, **54:101–102**
 SCID and, **49:400**
 surface antigens of human leukocytes and, **49:99**
 transformed tumors, adoptive T cell therapy of tumors and, **49:281, 295**
 virus-induced immunosuppression and
 HIV, **45:348, 349, 361, 365**
 measles, **45:344**
 Reverse-phase high-performance liquid chromatography, leukemia inhibitory factor, **53:32**
 Reverse transcriptase, virus-induced immunosuppression and, **45:358, 360, 361, 366**
 RF, *see* Rheumatoid factor
 RFX, **61:334–336**
 RGD sequence, leukocyte integrins and, **46:153**
 ligand molecules, **46:171–173**
 structure, **46:158, 159**
 RGL, **63:144**
 Rhe, **43:112–113**
 Rhesus monkey, complement components and, **38:207–209**
 Rheumatoid arthritis, **38:284, 298; 54:137–140, 187; 59:146, 343; 61:367**
 109 D6 mAb, **48:128**
 anemia in, **64:321–322**
 animal models, **64:288, 298**
 antinuclear antibodies and, **44:99, 111**
 anti-TNF- α antibodies and gold compounds and, **65:123–124**
 anti-TNF- α therapy, **64:289, 293, 333–334**
 animal models, **64:298**
 clinical studies, **64:299–307**
 aPL antibodies and, **49:208–210**
 autoimmune response in, **64:288**
 future prospects, **64:330–333**
 mode of action, **64:311–323**
 problems in, **64:323–330**
 cardiovascular disease and, **64:322**
 CD5 B cell and, **47:136, 147, 148–150**
 CD23 antigen and, **49:174**
 conformational epitope three-dimensional structure, **48:129**
 cytokines in, **64:290–298**
 DR β gene, **48:131**
 DRw53 molecule, **48:128**
 Dw4, **48:126–127**
 DW14, **48:126–127**
 Dw1O, **48:126–127**
 etiology, **48:126–132; 64:283**
 HLA, **48:126**
 HLA-B27 and, **65:94**
 HLA class II peptide binding and, **66:85, 87**
 Ig heavy-chain variable region genes and, **49:43, 46–47**
 IL-1 and, **44:191, 193–195**
 and IL-6, **54:40**
 infections, **64:288–289**
 susceptibility and outcome, **64:326**
 juvenile
 DR4, **48:131**
 DR5, **48:131**
 DRw8, **48:131**
 kallikrein in, **66:260**
 lymphocyte homing and, **44:315, 338, 344**
 MCI, **48:128**
 molecular basis of susceptibility alleles
 properties as factor in determination, **56:444–446**
 risk of severe disease associated with, **56:441–444**
 blacks, incidence of disease in, **56:427–429**
 characteristics of disease process, **56:389–390**

- epitope, shared
 concept, **56**:403–404, 416–418
 haplotype encoding, **56**:438–440
 identification by mAbs, **56**:422–425
 mapping, **56**:410–416
 ethnic differences, **56**:408–410
 genetic studies
 early knowledge, **56**:400–401
 family, **56**:433–435
 female sex, **56**:434
 number of genes, **56**:433–438
 twins, **56**:433–435, 446–449
- HLA polymorphism, actions of
 remission during pregnancy, **56**:431
 testing, **56**:432–433
 toxicity, **56**:431–432
- immune response, **56**:455–456
 incidence, **56**:399–400
 inheritance mode, **56**:440–441
 mapping
 in region facing antigen-binding
 groove, **56**:418–419
 into sequence motif, **56**:410–416
- MHC associated with
 class II molecules, **56**:429–430
 concepts, **56**:390–397
 determinants, **56**:400–402
 DRI, **56**:420–422, 426
 DR4, **56**:404–408, 422–426
 DR6, **56**:419–420
 DR10, **56**:424–427, 429
 Dw determinants, **56**:407–408
 function of molecules, **56**:449–450
 haplotype interactions, **56**:438–446
 polymorphism, **56**:398–399
 terminology, **56**:390–397
- MHC class II molecules, serologic
 identification, **56**:402–404
 monoclonal antibody 109d6 associated
 with, **56**:422–425
 peptide antigen X role, **56**:455
 population studies, **56**:425–427
 serologic association, **56**:406–407
 structural determinants, **56**:455–456
- T cell repertoire
 exogenous events, **56**:452–455
 formation, **56**:450–452
- therapeutic directions, **56**:456
- mortality, **64**:286
- NK cells and, **47**:300
- pathogenesis, **64**:283–288
 pituitary hormones, **63**:421–422
 rat, dietary fish oil and, **39**:163, 165
 TNF- α , **64**:293, 296–298, 315, 316
 cell trafficking, **64**:317–319
 joint destruction mechanism,
 64:320–321
 neovascularization, **64**:319–320
- Rheumatoid factor, **38**:278, 284; **39**:263
 murine lupus models and, **46**:72, 75, 77, 78
- Rheumatoid factors
 aPL antibodies and, **49**:210
 Ig heavy-chain variable region genes and,
 49:31, 43, 46–47, 52
- Rheumatoid synovial fibroblast-like cells,
 54:193
- Rheumatoid synovial tissue, **54**:170
- Rhinitis
 allergic, **60**:221
 seasonal allergic and perennial, **60**:172–173
- Rho, **63**:149, 171
- RIBI adjuvant, with MAP, **60**:131–132
- Ribonucleic acid, in suppressed cells,
 40:140–141
- Ribonucleoprotein, antinuclear antibodies
 and, **44**:93, 94
 autoantibodies, **44**:138
 autoantigen, **44**:129–131
 autoimmune response, **44**:131–134, 136
 mixed connective tissue disease, **44**:111
 SLE, **44**:96, 97, 100–102, 104–106
- Ribosomes
 antinuclear antibodies and, **44**:120, 128
 autoimmune response, **44**:132, 133
 SLE, **44**:96, 97, 105, 106
 genetically engineered antibody molecules
 and, **44**:70
 lymphocyte homing and, **44**:317
- Ricin A chain, MG and, **42**:265
- Rigin, **40**:115
 potency of, **40**:112
- Rin1, **63**:144
- RING4, **65**:50, 51
- RING4/11, **64**:4
- RING11 gene, **65**:50, 51
- RIP, **61**:31; **65**:117
- RNA, *see also* Messenger RNA
 antinuclear antibodies and, **44**:120
 autoantigen, **44**:129, 130
 autoimmune response, **44**:132, 133

- RNA (*continued*)
 mixed connective tissue disease, **44:111**
 scleroderma, **44:124**, 125
Sjögren's syndrome, **44:113**, 118
SLE, **44:100**, 101, 104
 B cell formation and, **41:191**, 198, 221
 double-stranded, induction of NF- κ B, **58:6**, 11–12
 HIV infection and, **47:403**
 etiological agent, **47:378**, 380
 immunopathogenic mechanism, **47:387**, 395
 hybrid resistance and, **41:80**, 377
IgE biosynthesis and, **47:21**
Ig gene superfamily and, **44:10**, 12, 34, 39, 44, 46
Ig heavy-chain variable region genes and, **49:33**, 38, 55
Ig-like loci, maps, **46:8**, 13, 25
IL-1 and, **44:156**, 159–161
 immunosenescence and, **46:227**
 leukocyte integrins and, **46:174**–176
 murine lupus models and
Ig germline, **46:74**
T cell antigen receptor, **46:86**–88, 90–92, 95
 negative strand transcript, **58:15**–16
NK cells and, **47:203**, 241, 242, 284
 regulators of complement activation and, **45:396**, 398, 399, 407
SCID and, **49:383**, 385
 sporozoite malaria vaccine and, **45:302**
T cell activation and, **41:27**
T cell development and, **44:217**
T cell receptor and, **45:116**
 translation inhibition, **58:240**
 virus-induced immunosuppression and
HIV, **45:348**, 354, 355, 366
measles, **45:336**, 344, 346
RNA colony blot hybridizations, Ars A response and, **42:132**–133
RNA–duplex DNA complex, **61:114**–116
RNA polymerases, antinuclear antibodies and, **44:93**, 94
 autoantigen, **44:127**, 129
 autoimmune response, **44:133**, 134
sclerodama, **44:122**–125
Sjögren's syndrome, **44:113**
SLE, **44:104**
- RNase
 complement receptor 2 and, **46:202**
 murine lupus models and, **46:86**–88, 91
RNA splicing, switch recombination, **61:101**–102
RNA viruses, in IBD, **42:291**
Rolipram, **64:331**
Rolling
 eosinophils and neutrophils, **60:165**–167
 high endothelial venules, **65:367**, 368
 leukocytes, **58:382**–383
RPMI 8866, **43:298**–300
RSal class II fragment, 4-kDa, celiac disease, **48:145**
RSS, *see* Recombination signal sequences
RT-6, **48:241**
RU-38486, effect on CD4 $^+$ 8 $^+$ thymocyte deletion, **58:259**
Runt domain, *AML1*, **65:24**, 25
- S**
- Sy3* regions, binding site, switch recombination, **61:117**–118
S6 kinase, signal transduction, **59:264**
Saccharomyces cerevisiae, cytolysis, **41:319**
SAGs, *see* Superantigens
Salicylates
NF- κ B and, **65:121**–122
PGHS, **62:187**
Salicylic acid, **62:184**–187
Saliva, S-IgA antibodies in, **40:201**–202
Salmonella adelaide, **54:395**
Salmonella paratyphi, flagella, and human antibody effector function, **51:16**–17
Salmonella typhimurium, secondary B cells and, **42:71**–72
Salt, cytotoxicity and, **41:290**, 301
S antigen, **48:215**–216
 anterior chamber-associated immune deviation, **48:219**
 experimental autoimmune uveitis, **48:219**
Saporin, MG and, **42:265**
Sarcoidosis, **62:267**, 281
 and TCR junctional regions, **58:328**–329
Sarcoma
 adoptive T cell therapy of tumors and, **49:286**, 291, 297, 301, 305
 hybrid resistance and, **41:358**, 359, 363

- SASD, identification of lipid A-binding membrane proteins, **53**:274–275
- SC58125, **62**:186
- Scatter factor, *see* Hepatocyte growth factor
- Schistosoma mansoni*, **54**:37
- anti-Id antibodies as vaccines against, rat, **39**:284
 - eosinophil-activating substance production, **39**:217
 - eosinophil-mediated killing of
 - AES and, **39**:209
 - C3 and, **39**:210
 - ECP and, **39**:106, 213
 - F(ab') of anti-GFA-2 and, **39**:184
 - IgE receptor and, **39**:180–181
 - IgG antibodies and, **39**:209–212
 - major basic protein and, **39**:188–190, 209, 212–213
 - infections, lymphokine production in, **53**:16
- Schistosomiasis, immunogen selection, **60**:124–125
- Schizosaccharomyces pombe*, maps of Ig-like loci and, **46**:5
- Schlepper experiment, *Ir* genes and, **38**:39, 40, 42, 51
- SCID, *see* Severe combined immunodeficiency
- Scleroderma
- antinuclear antibodies and, **44**:94, 114, 119–121
 - autoantibodies, **44**:125, 136, 137
 - autoimmune response, **44**:131–133
 - centromere antigens, **44**:121, 122
 - mixed connective tissue disease, **44**:111
 - nuclear antigens, **44**:122–125
 - Sjögren's syndrome, **44**:112
 - SLE, **44**:97, 105
 - IL-1 and, **44**:193
- Scleroderma–polymyositis overlap syndrome, **61**:301–302
- Sclerosing panencephalitis, virus-induced immunosuppression and, **45**:346
- SCMC assays, IBD and, **42**:295
- SDF-1, **65**:283
- 20 α SDH, *see* 20 α -Hydroxysteroid dehydrogenase
- SDS-PAGE, **42**:166–168, 167–172, 172, 188
- T cell receptor, **43**:150
- SEB, *see* Staphylococcal enterotoxin B
- Secondary B cell lineage, **42**:83, *see also* B cell repertoire expression
- environment and, **42**:68–79
- Secondary B cell repertoire expression, somatic mutation and, **42**:9–12, *see also* B cell repertoire expression
- Secondary immune response, regulation, **51**:271–272
- Secondary structure
- antigenicity and, **43**:76
 - antigenic recognition and, **43**:38
- Second messengers
- B cell formation and, **41**:197
 - in Fc receptor-mediated signal transduction
 - Fc γ R, **57**:72–74
 - Fc ϵ RI, **57**:77–79
 - T cell antigen receptor and, **41**:2, 26, 27
- Secretion, helper T cell cytokines, **46**:111–114, 127, 136
- Secretory antibodies, **65**:249, 250
- Secretory component, IgA binding of, **40**:165–166, 167, 186
- epithelial cells and, **40**:175–177
 - hepatocytes and, **40**:179–180, 182
 - structure and function, **40**:160–163
- Secretory glands, mechanisms of homing of plasma cell precursors to, **40**:204–207
- Secretory leukoprotease inhibitor, **62**:287–288
- Segmental flexibility
- agglutination, **48**:23–24
 - antidansyl antibody, **48**:13, 14
 - cell receptor binding, **48**:24–26
 - complement activation, **48**:24–26
 - complement binding activity, **48**:13, 15, 22
 - domain structure, **48**:14, 16, 22
 - evolution, **48**:26
 - functional implications, **48**:23–26
 - genetically engineered antibody, **48**:14, 16, 22
 - hinge region structure, **48**:13, 15, 22
 - Ig, **48**:5–19
 - electron microscopy, **48**:2–3
 - fluorescence polarization, **48**:3–5
 - hydrodynamic study methods, **48**:2
 - nanosecond, **48**:4–5
 - spin-label approach, **48**:5
 - steady state, **48**:4
 - study methods, **48**:1–34
 - X-ray crystallography, **48**:3

- Segmental flexibility (*continued*)
 IgA, 48:14–18
 IgE, 48:18–19
 IgG, 48:5–19
 early studies, 48:5–6
 fluorescence polarization, 48:6–10
 hinge region structure, 48:21–23
 IgG subclasses, 48:11–14
 spin-label method, 48:10–11, 12
 X-ray crystallography, 48:11
 IgM, 48:14–18
 mouse monoclonal antibody, 48:13, 14
 precipitation, 48:23–24
 SEK1, 63:148
 Selectin ligands, 64:168, 172, 194
 Selectins, 52:167–168; 59:383–385;
 62:257–258, 259, 267, 276, 277; 64:144,
 148; 65:352; 367, *see also* E-Selectin;
 L-Selectin; P-Selectin
 CD23 antigen and, 49:163
 characterization, 60:165–167
 genetic studies, 64:193–195
 interactions with eosinophils, 60:167–169
 monospecific ligands, 58:362
 multistep adhesion cascades, 64:183–184
 and related recognition molecules,
 52:364–366
 surface antigens of human leukocytes and,
 49:88
 Selection, *see also* Negative selection; Positive selection
 extrathymic, 58:315–316
 Ig heavy-chain variable region genes and,
 49:22, 25, 40
 intracellular, 58:120–121
 intrathymic, T cells, 58:139–174
 phenotypic, in germinal centers, 60:280
 T cell, processes, 51:164–189
 negative selection, 51:165–175
 intrathymic clonal deletion,
 51:165–173
 positive selection, 51:175–189
 functional divergence, 51:186–189
 ligands and receptors in positive
 selection, 51:178–180
 vs two kinds of death, 51:175–177
 positive selection, molecular basis of
 four-way
 CD4 vs CD8 down-regulation,
 51:185–186
- developmental choice, 51:180–186
 positive vs negative selection,
 51:180–183
 signaling pathways that control positive
 selection, 51:183–185
 thymic, 58:311–315
 Self-antigens
 autoimmune thyroiditis and, 46:263, 265,
 305, 307, 319
 B cell repertoire and, 42:43–51
 and foreign, 58:297–298
 from human donors, antibodies to
 Graves' ophthalmopathy, 57:245
 HIV-1, 57:246–250
 primary biliary cirrhosis, 57:244–245
 thyroid disease, 57:242–244
 Ir genes and
 cytotoxic T cells, 38:162
 T cell suppression, 38:152
 tolerance, 38:120, 123–125, 127, 128
 murine lupus models and, 46:65
 recognition, 58:319–320
 self-Ia, 38:120, 123, 127, 132, 134
 self-MHC, 38:119, 120
 sequestration, 48:217
 Self-nonself discrimination, 59:280
 in complement system, 60:75–76
 Self-peptides, positive selection, 59:122–128
 Self-reactive cells, elimination, 59:288–295
 Self-reactive clones, 58:213
 Self-reactive immune cells, clonal deletion,
 48:216, 217
 Self-tolerance
 experimental autoimmune uveitis,
 48:216–219
 eye, strategies, 48:217
 germinal center B cells, maintenance,
 60:280–283
 idiotypic network and, 39:276–277
 immunological memory and, 53:249–251
 lens, 48:220
 Self-tolerance checkpoints, 59:279–281,
 342–344
 immune repertoire, 59:319–342
 preimmune repertoire, 59:281–319
 Sendai virus
 cachectin, 42:220
 cytotoxic T cells and, 38:163, 164, 167
 responses in MHC class I-deficient mouse,
 55:408–409

- Sepsis
 anticytokine therapies, **66**:151
 anti-TNF- α therapy, **64**:310–311, 326
 chemokines and, **66**:104
 IL-8, **66**:104, 135–141
 MCP-1, **66**:141–142
 cytokines and, **66**:101–102, 103–104, 104
 G-CSF, **66**:149–150
 GM-CSF, **66**:150
 IFN- γ and, **66**:145–146
 IL-2, **66**:104, 150–151
 IL-4, **66**:144–145
 IL-10, **66**:142–145
 IL-12, **66**:146–147
 LIF, **66**:147–148
 measurement, **66**:104–105
 MIF, **66**:147–148
 TNF and IL-1, **66**:104, 117–127, 150
 defined, **66**:101
 and IL-1ra
 animal models, **54**:204–207
 in human infections, **54**:207–209
immuno干预, **66**:150–151
models, **66**:102–103, 117–121
pathogenesis, **66**:150–151
 chemokines, **66**:135–142
 cytokines, **66**:104, 105–135, 142–151
pulmonary inflammation, **62**:271–275
therapeutic regulation of complement
 system in, **56**:281–282
- Septic shock
 complement cascade, **62**:271
 contact cascade in, **66**:235
 experimental treatment, **58**:280–281
 IFN- γ and, **62**:103, 104
 NO role, **60**:337
 therapeutic regulation of complement
 system in, **56**:281–282
- Sequential epitope, **43**:25, 72
amino acid, **43**:39, 66–67
collective responses of different species,
 43:48
critical residue, **43**:64
DFLEKI, **43**:63–64
EVVPH, **43**:63–64
individual side chain role, **43**:40
replaceability matrix, **43**:77–78
superassemblies, **43**:36, 38
- Sequential switching, switch recombination,
61:105–107
- Ser530, cyclooxygenase active site, **62**:175
Serine
 antinuclear antibodies and, **44**:115
 Ars A response and, **42**:111–112, 155
 CD4 and CD8 molecules and, **44**:269
 CTL and, **41**:135, 148
 genetically engineered antibody molecules
 and, **44**:84
 IL-1 and, **44**:184
 phosphorylation, **48**:270–277
Serine esterases, cytotoxicity and, **41**:267,
 288, 297, 298, 310
Serine proteinase inhibitor, **61**:202
Serine proteinases, **64**:297
 complement components and
 C2, **38**:213, 214
 Factor B, **38**:209, 211, 214
 cytotoxicity and, **41**:277, 297, 310
 IL-1 and, **44**:157, 158, 163
Serine/threonine kinase inhibitors, **54**:18–19
Serine/threonine kinase receptors, in TGF- β
 signal transduction, **55**:188, 194–195
 binding of isoforms, **55**:194
 cDNA cloning, **55**:188, 192
 structure, **55**:190–193
Serine/threonine phosphatases, **63**:166
 as p56^{lck} targets in T cell signaling, **56**:160
SeroLOGY
 in analysis of MHC class I-deficient cells,
 55:385–386
 CD1, **59**:37–44
 CTL and, **41**:139–141
 HLA class II, **48**:112–114
Serpins, **61**:202, 205
Serratia, tumor necrosis and, **42**:215
Serum
 antibodies, IBD and, **42**:291–292
 antithymocyte, hybrid resistance and,
 41:350
 IL-6 levels, **54**:30
Serum amyloid A, lymphocyte adhesion and,

- Severe combined immunodeficiency
(continued)
 graft-versus-host disease, **49**:397–398
 histocompatible bone marrow
 transplantation, **49**:392
 hybrid resistance and, **41**:351, 365, 366,
 384, 379–381
 antigen expression, **41**:404, 405, 408, 409
 ontogeny, **49**:398–400
 posttransplant immunocompetence,
 49:399–402
 RAG1/2 mutations, **61**:308–310, 311
 SCID-Beige mouse, **63**:80
 SCID defect, V(D)J recombination and,
 64:49–50
 SCID-hu mouse, HIV infection in,
 52:463–466
 SCID-hu thy/liv xenograft model, **63**:79
 HIV-1 infection, **63**:89–90, 99–107
 T cells, **63**:82–83
 SCID mouse, **50**:303
 EBV-associated lymphoproliferative
 disease in hu-PBL-SCID mice,
 50:310–313
 expression of defective B lineage cells,
 53:146
 HIV-infection of hu-PBL-SCID mice,
 50:314–317
 other tumor models in SCID mice,
 50:313–314
 transfer of normal or autoimmune
 human cells to, **50**:304–310
 scid mutation, **61**:299–301, 317
 Scid syndromes, with V(D)J recombination
 defects, **58**:62–63
 stem cell engraftment, **49**:388–392
 lack of, **49**:396–397
 tolerance, **49**:398–399
 transplantation
 with fetal tissue, **49**:392–393
 with T cell-depleted bone marrow,
 49:393–396
 and V(D)J joining of antigen receptor
 genes, **56**:73–78
 Sex-limited protein, **46**:29
 Sex steroids, spontaneous autoimmune
 thyroiditis and, **47**:480, 481
 SGP2, **61**:204, 249–251
 SH1 domain, BTK, **59**:179, 185–186,
 188–191
- SH2 domain
 BTK, **59**:179, 182, 186, 188–190
 lacking in Jaks, **60**:4
 in ras pathway activation, **60**:13–14
 Stat activation specificity in, **60**:23
 Stat proteins, **60**:18–20
- SH2 and IRS1 NPXY (SAIN)-binding
 domain, **61**:161
- SH2-PTP, **63**:165–166
- SH3 domain, Btk, **59**:179, 182, 184–187
- Shape accessibility, MHR, **43**:35–36
- Shared epitope, **64**:283–285
- Shc, **63**:143, 146
 IL-2-induced Ras activation, **61**:161–163,
 164–166
 signal transduction, **59**:262
- SHC, tyrosine phosphorylation, **60**:13–14
- Shedding
 lymphocytes, **64**:170
 in regulation of CD44 receptors,
 54:317–318
- Sheep red blood cells
 cytotoxicity and, **41**:293
 humoral immune response and, **45**:1, 2, 68,
 70, 72
 hybrid resistance and, **41**:352
 T cell subsets and, **41**:67–71, 77, 90, 92
- Sheet–sheet interactions, **43**:102–103, 110
- Sheet–sheet packing, schematic, **43**:102–103
- Shock, **42**:213, *see also* Cachectin
- Shope fibroma virus, T2 ORF product
 resemblance to TNF-R domain, **61**:10
- Short consensus repeats
 complement receptor 1 and, **46**:184–186,
 188–190, 194, 195
 complement receptor 2 and, **46**:202, 203,
 211
 regulators of complement activation and
 common structural motif, **45**:391, 392
 genes, **45**:404–408
 localization, **45**:392–394
 protein expression, **45**:399
 protein utilization, **45**:409, 410
- SHPTP2, in cytokine signaling, **60**:26
- Sialic acid, **64**:162, 168
 CD23 antigen and, **49**:155–156
 lymphocyte homing and, **44**:354, 355, 363,
 367–369, 371
- Sialidases, lymphocyte homing and,
 44:364–368, 371

- Sialomucins, **64**:144, 146–152
CD34, and selectin ligand, **58**:359–360
- Sialophorin, **48**:248
- Sialyl Lewis X, selectins and, **60**:166–167; **65**:363
- Sialytransferases, **64**:168
- Sicca-CD8⁺ lymphocytosis syndrome, **65**:286
- Side chain
contribution to solvent-exposed surface area, **43**:36–38
hydrophobic, **43**:42, 80–81
sequential epitope role, **43**:40
- Signal binding proteins, recombination, **61**:303–304
- Signaling, *see also* Negative signaling
abortive, and second signals in adult B cells, **54**:407–410
antigenic, constraints to biochemical study of, **52**:294–295
cytokines in, **58**:369
differential, between CD4 and CD8, **53**:91–94
in IL-4 production, mechanisms, **53**: 17–22
mitogenic, Ig, **59**:326–328
in naive and memory T cells, **53**:228–229
pathways, controlling positive selection, **51**:183–185
- TCR-mediated
and CD4, **53**:85–91
and CD8, **53**:85–91
and p56^k, **53**:85–91
transmembrane, **48**:227
- Signal joints, V(D)J recombination, **64**:40, 41
- Signal sequence recognition, V(D)J recombination, **64**:54
- Signal transducers and activators of transcription, **59**:261; **60**:17–25, *see also* Stat
activation by EGF, **60**:27
in cytokine signaling, **60**:19–22
in IFN signaling, **60**:17–19
phosphorylation by Jak2, **60**:210
recruitment, **60**:22–23
- Signal transduction, **59**:176–177
adoptive T cell therapy of tumors and, **49**:287, 289, 302, 313, 316
B7-1 and B7-2, **62**:138
Btk, **59**:184–185; **60**:42
cachectin and, **42**:226
- CD4 and CD8 molecules and, **44**:267, 270, 299, 300, 302
- CD23 antigen and, **49**:156
- CD28, **62**:134
- CD40, **60**:45–46; **61**:32–38; **63**:50–52
B cell proliferation, **63**:53–54
B cell receptor pathway, **61**:37–38
B cells, **61**:32–35
CD28-CD80/CD86 pathway, **61**:36–37
Fas-Fas-L pathway, **61**:37
Ig production, **63**:58–70
- CD44-ligand interactions, **54**:300–302
- CD45 and, **66**:1–56
and cytokine receptor superfamily, **60**:1–2
cytokines
adapter molecules, **63**:134, 141–143
calcineurin, **63**:166–167
CD45, **63**:164–165
Jak-Stat pathway, **63**:138–141
MAPK family, **63**:148–149
negative regulation, **60**:25–26
nuclear transcription factor NF- κ B, **63**:157–160
Pac-1, **63**:167
phosphatidylinositol 3-kinase, **63**:149–154
phosphoprotein phosphatases, **63**:164–167
protein kinase C, **63**:154–157
protein-tyrosine kinases, **63**:133–143
protein-tyrosine-phosphatase 1C, **63**:165–166
Raf kinase, **63**:147–148
Ras, **63**:143–147
Rho protein, **63**:149
Src family, **63**:133–137
Stats role, **60**:19–22
Syk family, **63**:134, 137–138
eosinophil, **60**:208–210
Fas-mediated, **57**:134–135, 139
Fc receptor-mediated
antigen receptor homology motif, **57**:81–83
Fc γ R, mechanisms of activity, **57**:72–77
Fc γ R, structural factors, **57**:82, 84–89
Fc ϵ RI, **57**:77–81, 89–90
mechanism, **57**:71–81
phosphorylation, **57**:74–77, 79–81
protein kinase involvement, **57**:74–77, 79–81

- Signal transduction (*continued*)
 second messenger interactions, **57**:72–74, 77–79
 structural factors, **57**:81–90
 GPI hydrolysis, **59**:264–265
Grb2, **59**:262
IFN- γ binding, **62**:67–71, 106
Ig gene superfamily and, **44**:31, 36
IgR-mediated cascade
 functional molecules, **54**:351, 352
 stimulation, **54**:378–379
IL-1 and, **44**:183
IL-2/IL-2R system, **63**:127, 173–176
 signaling pathways, **63**:169–173
 signal molecules, **63**:167–169
 target genes, **63**:160–164
 three-channel model, **63**:171–173
IL-2-induced, **59**:253–259, 265–266
IL-2R, **42**:178
IL-2R γ , **59**:227
IL-2R-triggered proximal signaling events, **61**:153–161
IL-6R complex chain, **54**:14, 18
IL-9 receptors and, **54**:83
lymphocyte, **62**:42, 44–47
lymphokine-mediated, **48**:310
MAP kinase, **59**:262–263
MB-1 molecule and, **54**:351–355
 mechanism of *IL-2R*-mediated, **50**:152
 in neutrophil regulation by *IL-8*, **55**:140–142
 new molecules, **61**:29–32
NF- κB , **58**:5
nucleus, **48**:310–311
PI3-K, **59**:263
pituitary hormones, **63**:397–399, 401–402
preB cell receptor and, **63**:26–29
pre-TCR, **58**:145–146
protein kinase C, **59**:263–264
Raf-1 kinase, **59**:262–263
Ras, **59**:261–262; **60**:12–14
 regulators of complement activation and, **45**:384, 410
S6 kinase, **59**:264
SCID and, **49**:386
Shc, **59**:262
SOS, **59**:262
 surface antigens of human leukocytes and, **49**:113, 125
 adhesion molecules, **49**:87
 antigen-specific receptors, **49**:76–78
MHC glycoproteins, **49**:81
 receptors, **49**:98, 100
T cell development and, **44**:212
T cell receptor and, **45**:110
 $\alpha\beta$ TCR complex, **58**:124–128
TGF- β
 endoglin functions, **55**:203–204
 formation of complexes, **55**:200–202
 intracellular, **55**:204–205
 type III receptor functions, **55**:203–204
TNF-R superfamily, *CD40*, **61**:26
 tyrosine kinases, **59**:259–261
 Signal transduction pathways
 artificial inducers, **58**:269
 artificial triggering, **58**:230–232
Bcl-2 effects, **58**:256
 biochemical analysis, **54**:415–420
 direct analysis, **54**:418
 identification of IgG complex and associated second messengers, **54**:419–420
 use of inhibitors to block *in vitro* tolerance, **54**:415–418
DNA damage, **58**:29
 via *IgR* transmembrane signaling, **54**:359–378
 and intervention of apoptosis, **58**:245–247
 leading to NF- κB activation by *TNF α* , **58**:7–9
 and p53, **58**:67
 TCR-mediated, relation of Apo-1/Fas, **58**:136
SIINFEKL, **59**:122, 124
SIIRFEKL, **59**:125
Silica, hybrid resistance and, **41**:350, 368, 395, 371–373
Simian immunodeficiency virus, **63**:79
 HIV infection and, **47**:378, 385, 404
 infection of macaques, **52**:442–452
 pathogenesis, **52**:443–452
 genetic drift of SIV molecular clones *in vivo*, **52**:451–452
 natural history, **52**:443–448
 viral determinants, analysis with SIV molecular clones, **52**:448–451
 phylogeny of nonhuman primate lentiviruses, **52**:442–443
 for vaccine testing, **52**:452–454
 vaccines, **65**:321–322

- Simian immunodeficiency virus type 1, **63**:79, 85
 Simian virus 40
 early promoter, **43**:268
 Ig heavy-chain variable region genes and, **49**:56–57
- Sindbis virus, cytotoxic T cells and, **38**:162
 Single strand annealing model, **61**:111
 sis-inducible element, in *c-fos* gene, **60**:24
 Sister chromatid exchange, **54**:234–235
 Site-specific tolerance, peripheral
 lymphocytes, **53**:201–203
 Sjögren's syndrome, **62**:76–77
 antineur nuclear antibodies and, **44**:111–113,
 115–118
 autoantibodies, **44**:127, 136
 SLE, **44**:93, 95, 102, 104
 aPL antibodies and, **49**:209–210
- Skin
 allergic reaction, contact activation, **66**:259
 homing, **58**:392
 IL-6 effects, **54**:28–29
 infections, immunity, **58**:326–328
 invariant intraepithelial γ/δ T cells,
53:167–169
 invariant γδ T cell receptors of murine
 IEL, **58**:301–303
 suppressor antigen-presenting cell, **48**:206
 Skin disease, IL-8 role, **55**:142–143
 Skin grafts
 allografts, T cell subsets and, **41**:86, 88, 93
 B cell formation and, **41**:225
 CTL and, **41**:158
 hybrid resistance and, **41**:369, 376, 383
 in MHC class I-deficient mouse,
55:388–390
 rejection
 CTL and, **41**:159, 165
 T cell subsets and, **41**:71, 88, 93, 94
 Skin test antigens, Crohn's disease and, **42**:289
 Skin window chamber technique, **42**:307
 SLE, *see* Systemic lupus erythematosus
Sl locus, kit ligand encoded at, **55**:10–12
 Smoking, related lung disease, **60**:220–221
 Smooth muscle cells, cachectin and, **42**:220
Sμbp-2, binding site, switch recombination,
61:119
 Snake venom toxin, MG and, **42**:237
 Sodium azodisalicylate, **42**:318
 Sodium dodecyl sulfate, denatured AChR
 subunits, **42**:257, 258
 Sodium dodecyl sulfate–polyacrylamide gel
 electrophoresis, *see* SDS-PAGE
 Sodium salicylate, inhibition of NF-κB
 activation at high doses, **58**:19–20
 Solid-phase immunoassay, aPL antibodies
 and, **49**:193, 199, 258–259
 affinity purification, **49**:230–231
 antibody subsets, **49**:232
 clinical aspects, **49**:200, 203–206
 isotype, **49**:227
 specificity, **49**:235, 243, 249
 Solid-phase synthesis, multiple antigen
 peptides, **60**:106–108
 Soluble costimulators, EC production of,
50:272–274
 Soluble mediators, in IBD, **42**:310–315, 322
 Solvation energy, estimating, **43**:101
 SOM28, **48**:172
 Somatic cell genetics
 complement components and, **38**:204, 233
 human B cell neoplasia and, **38**:247
 maternally transmitted antigen and, **38**:320
 Somatic cell hybridization
 human B cell neoplasia and
 Burkitt lymphoma cells, **38**:257, 259
 chromosomal translocation, **38**:248, 265
c-myc activation, **38**:263
c-myc oncogene, **38**:248, 249, 259
 lymphoblastoid cells, **38**:260
 lymphocyte hybridomas and, **38**:275
 B cell immortalization, **38**:306
 genetically engineered antibodies, **38**:305
 human-human, **38**:293, 297, 298
 human-mouse, **38**:287–289
 mAbs, **38**:277, 282, 302
 Somatic diversification
 Ig gene superfamily and, **44**:12–20
 Ig VL gene, **48**:60–62
 Somatic fusion, B cell hybridization and,
38:279
 Somatic hypermutation
 in germinal centers, **53**:233–234
 T cell subsets and, **41**:42, 88, 106
 Somatic mutation, **42**:96, 138
 adoptive T cell therapy of tumors and,
49:282
 and antibody repertoire generation in
 rabbit, **56**:201, 203

- Somatic mutation (*continued*)
 antigen and, **42:34**
 Ars A response and, **42:108**
 B cell repertoire expression and
 junctional diversity, **42:15–18**
 primary, **42:12–15**
 random vs nonrandom V region segment
 selection, **42:18–19**
 age factor, **42:30–32**
 antigen responsive cell populations,
 42:19–22
 fetal and neonatal development,
 42:26–30
 predominant clonotype expression,
 42:22–26
 secondary, **42:9–12, 69–71**
 Ig heavy-chain variable region genes and,
 49:2, 7, 26
 D segments, **49:28, 32**
 V_H gene expression, **49:37, 39, 41, 46,**
 49, 52
 neonatal hybridoma and, **42:67**
PIG-A gene, **60:82–89**
 secondary B cells and, **42:79**
 and terminal differentiation, apoptosis and,
 50:65–66
- Somatic variation, T cell receptors and, **38:16**
- Somatostatin, **48:169; 63:385, 386–387**
 cells of origin, **39:317**
 histamine release by mast cells and,
 39:305–306
 immunosuppressive properties, **39:310–311**
 secretion regulation by, **39:302–303**
- Sos, **63:143, 144, 146**
- SOS, signal transduction, **59:262**
- Southern blot analysis
 autoimmune demyelinating disease and,
 49:363
 clonally rearranged genes, **40:267–268**
 complement components, **38:212, 215, 229,**
 237
 human B cell neoplasia, **38:248, 266, 267**
 hybridomas, **38:299**
Ir genes, **38:117**
 SCID and, **49:385–386**
 T cell receptors, **38:11, 13, 15, 16**
- Southern filter hybridization analysis
 Ars A V_H gene segment and, **42:101**
 of BALB/c liver, **42:121**
 of V_H complex, **42:130**
 V_x10 family of germline genes and, **42:144**
- sox-4*, B cell genesis, **63:218–219, 226–228**
- Soybean agglutinin, SCID and, **49:394–396**
- SP, *see* Substance P
 SP-40,40, **61:204, 249–251**
 Sp100/ND, **63:202**
 Spacer sequences, Ars A response and,
 42:116
- Specific unresponsiveness, hybrid resistance
 and, **41:384–386**
- Spectrotypes, autoimmune thyroiditis and,
 46:301, 304, 306
- Spergualin, **65:125–126**
- Spermidine, **48:288**
- Spermine, **48:288**
 depletion, induction of DNA
 fragmentation, **58:252**
- Sphingomyelin, turnover, role of Fas,
 60:299–300
- Spi-B
 binding sites, **64:76–77**
 domain, **64:76–77**
 expression, **64:87–88**
 gene targeting, **64:88**
 structure, **64:79**
 structure-function studies, **64:88**
- Spleen
 adoptive T cell therapy of tumors and
 antigen recognition, **49:321**
 expression of antitumor responses,
 49:331–332
 mechanisms, **49:300, 309**
 principles, **49:293, 295–298**
 anterior chamber-associated immune
 deviation, **48:206, 211**
 antigen-presenting cells and, **47:69–72,**
 74–76, 78–81
 antigen processing, **47:92**
 immunogenicity, **47:102**
 T cells, **47:83, 85**
 tissue distribution, **47:47, 50–52, 56**
 antinuclear antibodies and, **44:102**
 architecture, **60:268–269**
 autoimmune thyroiditis and, **46:280, 281,**
 306, 310–312
- autoreactive T cells and
 activation, **45:421, 422**
 physiology, **45:428**
 regulation, **45:432, 433**
- B cell formation and, **41:181, 182, 184, 238**
 B cell precursors, **41:198, 200–202**

- genetically determined defects, **41**:229
 inducible cell line, **41**:223
 population dynamics, **41**:207, 208
B cells, **42**:74, 75
 adult, **42**:68
 CD5⁺ and CD5⁻, **53**:180–181
 neonatal, **42**:65–67
CD4 molecules and, **44**:288
CD5 B cell and, **47**:120–123
 genetic influence, **47**:147
 Ig gene expression, **47**:152, 157
 lineage, **47**:128
 physiology, **47**:130, 132, 133
 primordial immune network, **47**:159
CD8 molecules and, **44**:271, 282
 complement receptor 2 and, **46**:204–206
CTL and, **41**:138
 delayed-type hypersensitivity, **48**:211–212
FcεRI⁺ cells, lymphokine production by, **53**:9–14
 forbidden α/β T cells, **53**:166
 helper T cell cytokines and, **46**:122, 123,
 128, 129, 137, 138
 histology, **60**:268–272
 humoral response and, **45**:2
 antigens, **45**:11, 16, 17
 helper T cells, **45**:27, 34
 physical interaction, **45**:40, 53
 hybrid resistance and, **41**:334
 antigen expression, **41**:397, 402, 404, 408
 bone marrow cells, **41**:336, 338–340,
 342, 345, 347, 348
 effector mechanisms, **41**:370
 leukemia/lymphoma cells, **41**:358,
 361–368
 lymphoid cells, **41**:351–356
 marrow, **41**:384–388, 391, 393
 NK cells, **41**:372, 373, 375
 syngeneic stem cells, **41**:388–390
 T cells, **41**:380–384
in vitro assays, **41**:394–396
IgE biosynthesis and
 antibody response, **47**:5, 8, 11
 antibody response suppression, **47**:29,
 32–34, 36, 37
 binding factors, **47**:13–15, 22–26, 28
IGF-I, **63**:391–392
 immunosenescence and
 lymphocyte activation, **46**:226, 228, 232
 lymphocyte subsets, **46**:237, 238
 marker density, **46**:239
 mucosal immunity, **46**:251, 253
 regulatory changes, **46**:247–251
 stem cells, **46**:240, 241, 244, 245
lymphocyte homing
 high endothelial venules, **44**:315
 molecules, **44**:344, 351, 353
 regional specificity, **44**:328, 335
 lymphocyte migration, **58**:346–347
 lymphocyte recirculation, **58**:350;
64:177–178
 murine lupus models and, **46**:64, 91, 92, 98
NK cells and, **47**:187, 283, 289, 290, 292,
 300
 CNS, **47**:268
 congenital defects, **47**:226
 differentiation, **47**:229, 230
 effector mechanisms, **47**:234, 260, 262
 genetic control, **47**:223
 hematopoiesis, **47**:274, 276
 identification, **47**:199
 malignant expansion, **47**:228
 surface phenotype, **47**:210, 211, 213
 tissue distribution, **47**:220, 221
 regulators of complement activation and,
45:384, 388, 389, 410
 spontaneous autoimmune thyroiditis and
 cellular immune reactions, **47**:450, 452,
 454, 456
 disturbed immunoregulation, **47**:471,
 473–476, 481
 humoral immune reactions, **47**:446, 447
 sporozoite malaria vaccine and, **45**:287,
 305, 306
 synthetic T and B cell sites and, **45**:199
 T cell development and, **44**:250, 251
 T cell populations, **53**:161–163
 T cell receptor and, **45**:134
 T cell subsets and
 H-2 alloantigen recognition, **41**:78, 86,
 89, 91, 92
 H-2 molecules in thymus, **41**:95, 102
 H-2-restricted antigen recognition,
41:67, 69, 72
 virus-induced immunosuppression and,
45:354
Spleen cells
 NZB, IgM secretion, **55**:313–314
 transgenic, preliminary *in vitro* studies,
54:414–415

- Spleen colony formation, hybrid resistance and
 antigen expression, **41**:402, 404
 bone marrow cells, **41**:337, 347
 syngeneic stem cells, **41**:388–390
 T cells, **41**:382
- Spliceosomes, antinuclear antibodies and, **44**:102, 131
- Splicing
 CD1 genes, **59**:9–10
 events, abnormal, in *PIG-A* gene, **60**:88–89
- Splicing domains, **63**:201
- Spondyloarthritic diseases, TAP and, **65**:93
- Spontaneous autoimmune thyroiditis
 antigens, **46**:269, 270
 cellular immune responses, **46**:287
 experimental models, **46**:273–276, 279
 genetic control, **46**:282–284
 humoral responses, **46**:298, 305, 306
 obese strain chicken, **47**:433–435, 491–494
 altered thyroid function, **47**:456–465
 breeding, **47**:435–438
 cellular immune reactions, **47**:449–456
 clinical symptoms, **47**:438–440
 disturbed immunoregulation, **47**:470, 476, 477, 479
 genetics, **47**:484
 dysfunctions, **47**:487–491
 MHC, **47**:481–486
 target organ, **47**:485, 487
 histopathology, **47**:440–444
 humoral immune reactions, **47**:444–449
 immunoregulation, **47**:469
 extrinsic, **47**:476–481
 intrinsic, **47**:470–476
 potential effector mechanisms, **47**:465–469
 prevention, **46**:310, 311
- Spontaneous cell-mediated cytotoxicity, assays, IBD and, **42**:295
- Spontaneous cellular cytotoxicity, **47**:451, 452
- Spontaneous neoplasm, immune regulation, **48**:212
- Sporozoite malaria vaccine, **45**:283, 284
 antibodies, **45**:300, 301
 CS proteins, **45**:292–294
 cloning, **45**:294–296
 evolution of repeats, **45**:296–299
 function of repeats, **45**:299, 300
- CS-specific T cells, **45**:306–308
 epitopes, **45**:310–313
 Ir gene control, **45**:308–310
 endemic areas, **45**:315–317
 human trials, **45**:313–315
 immunity, **45**:285, 286
 specificity, **45**:287–292
 viability, **45**:286, 287
- interferon- γ , **45**:301–306
 perspectives, **45**:319–322
Plasmodium vivax, **45**:317, 318
- Sporozoites, *Plasmodium*, antibodies, **60**:112–124
- S protein, **61**:204, 247–249
- 11S protein, **64**:21, 25
- SPT14*, **60**:72
- Src family, **55**:253–255; **61**:153–158; **63**:133–137
 in cytokine signaling, **60**:5–9
 src homology 2, *see* SH2 domain
 Src tyrosine kinases, **63**:133–137
 activation by B cell antigen receptor, **55**:253–255
 in cytokine signaling, **60**:5–9
 IL-2R coupling, **61**:153–158
 regulation by CD45, **66**:4, 23, 39–45
- S region, **54**:236
- SRP1, **64**:48
- Stamper–Woodruff assay, **60**:167–168
- Staphylococcal α -toxin, cytolysis and, **41**:317
- Staphylococcal δ -toxin, cytolysis and, **41**:315, 316
- Staphylococcal enterotoxin A
 action on TCR, **58**:323
 deletion of $V\beta 3^+$ and $V\beta 11^+R$ lymphocytes, **58**:222
- Staphylococcal enterotoxin B
 beneficial effect, **58**:274–275
 induction
 cytokines, **58**:271–272
 thymocyte apoptosis, **58**:216
 $V\beta 8^+$ thymocyte depletion, **58**:259–260, 262
 with L-NAME, **60**:348–349
 pleiotropic effects *in vivo*, **58**:219–224
- Staphylococcal enterotoxins, **52**:165
- Staphylococcal protein A, antibodies from combinatorial libraries, **57**:229

- Staphylococcus aureus*, **52**:160; **54**:108, 109
 antineclear antibodies and, **44**:115, 118,
 135
 autoimmune thyroiditis and, **46**:299
Cowan I strain, CD23 antigen and, **49**:151
Cowan strain, **52**:187–188
 and anti-CD40 results in T-independent
 B cell differentiation, **52**:191
and food poisoning, **54**:128
genetically engineered antibody molecules
 and, **44**:78, 85
infection-associated autoimmune disease,
 54:140–141
 pulmonary infection, **59**:397
Staphylococcus aureus B
 B cells responsive to, **40**:20, 21
cell response and, **40**:14–15, 16,
 41–44
Stat1, **59**:261; **62**:69, **63**:141
Stat2, **59**:261; **62**:69
STAT3, **65**:19
Stat3, **59**:261; **61**:177; **63**:141, 399
STAT5, **65**:20
Stat5, **63**:141, 399
 IL-2 activation, **61**:177, 178
STAT6, **65**:20
Stat6, **61**:177
Stat48, **62**:69
Stat91, **62**:68
STAT family, interaction with NF-IL6,
 65:19–21
Stat proteins, **59**:261; **62**:68, 69; **63**:140, 398,
 see also Signal transducers and activators
 of transcription
 IL-2R coupling, **61**:162, 174–179
Stem cell factor, **55**:1–2, 81–84; **62**:189
 chromosomal location, **55**:13–15
 effect on secretion in SCFR⁺ cells, **55**:43
 expression regulation, **55**:31–33
 genomic organization, **55**:13–15
 germ cell development, **55**:66–69
 identification, **55**:10–12
 kit ligand, identification, **55**:10–12
 lymphohematopoiesis effects
 hematopoiesis, **55**:43–48
 lymphocytes, **55**:48–51
 NK cells, **55**:48–51
 in mast cell biology
 development, **55**:52–60
 function, **55**:60–64
 in melanocyte biology, **55**:64–66
 mutations affecting
 human, **55**:23–24
 mouse, **55**:17–23
 rat, **55**:23
neoplasia, **55**:71–72
 brain tumors, **55**:77
 breast cancer, **55**:78
 gonadal tumors, **55**:76–77
 hematopoietic tumors, **55**:72–74
 lung cancer, **55**:77–78
 mast cell leukemia, **55**:74
 melanoma, **55**:74–76
nervous system, **55**:69–71
as therapeutic agent, **55**:78–81
tissue distribution, **55**:15–17
Stem cell factor receptor
 c-kit gene encoding, **55**:3–10
 allelism with W, **55**:6–7
 and c-fms, evolutionary relationship,
 55:9–10
 chromosomal location, **55**:7–9
 genomic organization, **55**:7–9
 and PDGFR genes, evolutionary
 relationship, **55**:9–10
expression regulation, **55**:24–31
germ cell development, **55**:66–69
interaction with ligand, consequences
 adhesion, **55**:37
 chemotaxis, **55**:37–38
 differentiation, **55**:40–42
 haptotaxis, **55**:37–38
 maturation, **55**:40–42
 proliferation, **55**:39–40
 secretion, **55**:43
 survival, **55**:38–39
lymphohematopoiesis effects
 hematopoiesis, **55**:43–48
 lymphocytes, **55**:48–51
 NK cells, **55**:48–51
mast cells bearing, secretion, SCF effects,
 55:43
melanocyte biology, **55**:64–66
mutations
 human, **55**:23–24
 mouse, **55**:17–23
 rat, **55**:23
neoplasia, **55**:71–78
signal transduction through, **55**:33–36
tissue distribution, **55**:15–17

- Stem cell factor receptor/c-kit, **52**:352–354,
363–364
- Stem cells
B cell formation and, **41**:181, 182, 184,
187, 206, 237
B cell precursors, **41**:188, 191, 195, 199,
200
bone marrow cultures, **41**:209, 217, 219
genetically determined defects, **41**:220,
224, 225
soluble mediators, **41**:232–234
- embryonic, *see* Embryonic stem cells
- hematopoietic, *see* Hematopoietic stem cells
- hybrid resistance and, **41**:352, 362, 367
antigen expression, **41**:402–407
bone marrow cells, **41**:336–340, 344,
346
effector mechanisms, **41**:370–373,
377–380, 383, 386, 388–391
- lymphoid, *see* Lymphoid stem cells
- self-renewal, **58**:422
- switching, developmentally programmed,
58:308–309
- syngeneic stem cell functions, hybrid
resistance and, **41**:388–390
- Steroids
aPL antibodies and, **49**:223, 232
B cell formation and, **41**:209, 219, 236, 238
hybrid resistance and, **41**:348, 387
NF- κ B and, **65**:118–120
spontaneous autoimmune thyroiditis and,
47:480, 481
virus-induced immunosuppression and,
45:338
- Steroid therapy, IBD and, **42**:286, 289,
316–319
- Stimulating cells, **50**:31–32
B cells, role in V β selective element action,
50:31
dendritic cells, role in V β selective element
action, **50**:31
transfer of Vbse among cells, **50**:32
- Stochastic processes, maternally transmitted
antigen and, **38**:324, 326, 327, 341
- Stomach cancer, X-linked
agammaglobulinemia, **59**:147
- Streptococcal M proteins, **54**:114–115
- Streptococcal pneumonia, **59**:395–396
- Streptococcal toxins, **54**:108–113
- Streptococcus*
infection, membrane cofactor protein,
61:221
tumor necrosis and, **42**:215
- Streptococcus mutans*, subunit vaccines,
60:135–136
- Streptococcus pneumoniae*, mouse infection
anti-Id antibodies as vaccines against,
39:283
- Streptococcus pyogenes*, **54**:109, 114
 $\nu\beta$ specificity for, **54**:109, 110–112
- Streptococcus zooepidemicus*, and Fc of IgG,
51:61
- Streptolysin O, 317, **41**:318
- Streptomyces tsukubaensis*, **52**:403
- Streptozotocin, diabetes induced by, **62**:100
- Stress-activated protein kinases, **63**:148
- Stroke, aPL antibodies and, **49**:218, 221, 258
- Stroma, hybrid resistance and, **41**:336, 404
- Stromal cells
B cell formation and, **41**:184, 185, 237, 238
B cell precursors, **41**:198, 200, 203
bone marrow cultures, **41**:209, 210,
212–216, 219, 220
genetically determined defects, **41**:223,
227, 229, 231, 232
lymphohemopoietic tissue organization,
41:186–188
soluble mediators, **41**:234
- contacts with pre-B cells, **53**:133–134
- immunosenescence and, **46**:241
- pre-B cell nonproliferation on, **53**:134–135
- pro- and pre-B cells reactive to,
53:131–136
- thymic, autoreactive T cells and,
45:421–425
- uterine, IL-1ra production, **54**:188
- Stromelysins, **62**:288; **64**:297
- Structural complementarity, protein-protein
interactions, **43**:104–105
- Structure-function studies, MMTV
superantigens, **54**:103–106
- Subclones
B cell formation and, **41**:210
CTL and, **41**:139
- Subcloning, hybridomas and, **38**:291
- Submucosa, Crohn's disease specimens,
42:298
- Substance P, **61**:372–373
arthritis and, **39**:317–318

- bronchospasm induction and, **39:318**
effects on
 secretory functions, **39:302–303**
 smooth muscle and vasculature,
 39:302–303
- histamine release by mast cells and,
39:305–306
- immunostimulatory properties,
39:311
- leukocyte chemotaxis and, **39:308**
- lymphocyte adhesion and, **64:153**
in nasal secretion, **39:304**
and other neuropeptides, receptors for,
52:372–375
- tissue repair and, **39:308–309**
- Substance P receptors
 lymphocyte, **39:313**
 recovery from IM-9 cells, **39:313**
 T cell, **39:313**
- Subtilisin, site-directed mutants, **43:23**
- Subtractive hybridization, **43:135**
- N*-Succinimidyl 3-(2-pyridyl)dithio) propionate,
 in crosslinking for chemoimmuno-
 conjugates in cancer treatment,
56:319–320
- Sucrose, cytotoxicity and, **41:290, 304**
- Sugars
 immunosenescence and, **46:234**
 lymphocyte–HEV interaction, **65:360–361,**
 363–365
 as selectin ligands, chemical structure,
 58:355
- Sulfapyridine, **42:317, 318, see also**
 Sulfasalazine
- Sulfasalazine, **42:311**
 IBD and, **42:289, 296, 316–318**
 prostaglandin synthesis and, **42:312–313**
- Sulfated glycoprotein 2, **61:204, 249–251**
- Sulfidoleukotrienes, IBD and, **42:314, 315**
- Sulfosuccinimidyl-2(*p*-azidosalicyamino)-1,3'-
 dithiopropionate identification of lipid A-
 binding membrane proteins, **53:274–275**
- Sulfotransferases, **64:168**
- Sulindac, **62:184, 186, 201**
- SUN1*, **64:20, 21**
- Superantigens, *see also* Staphylococcal
 enterotoxin A; Staphylococcal
 enterotoxin B
bacterial, **65:14, 208–210**
classes, **54:99–100**
- endogenous murine, **54:100–108**
cellular expression of MMTV antigens,
54:106–107
- identification of Mls antigens as MMTV
products, **54:100–102**
- role in life cycle of MMTV and host,
54:107–108
- structure-function studies, **54:103–106**
- exogenous, **54:108–117**
- bacterial products with related function
 as superantigen, **54:113–115**
produced by infectious viruses,
54:115–117
- staphylococcal and streptococcal toxins,
54:108–113
- and human disease, **54:128–145**
- autoimmune disease associated with
 streptococcal infection, **54:140–141**
- food poisoning, **54:128**
- immunodeficiency disease, **54:142–143**
- Kawasaki syndrome, **54:133–137**
- lymphoproliferative diseases, **54:143–145**
- rheumatoid arthritis, **54:137–140**
- systemic lupus-like autoimmune disease,
54:141–142
- toxic shock syndrome, **54:128–133**
- immune responses
 binding to major histocompatibility
 complex, **54:117–120**
 direct non-T cell effects, **54:127**
 involvement of accessory CD4 or CD8
 molecules, **54:124–125**
 recognition by T cell receptors,
 54:121–124
- T cell deletion and anergy in adult
 animals injected with superantigens,
54:125–127
- T cell signaling in response to
 superantigens vs conventional
 antigens, **54:127**
- induced deletion and anergy, differential
 regulation, **58:272**
- MMTV, **65:140, 152–153**
 expression, **65:152–153**
 immune stimulation by, **65:140, 141,**
 171–194, 196–203
- MHC class II molecules, interaction
 with, **65:164–165**
- protein structure, **65:157–164**
- TCR V β , interaction with, **65:165–167**
- tumor formation and, **65:155, 156**

- Superantigens (*continued*)
 mycobacterial, **58**:321–323
 T cells stimulated by, **58**:260–262
- Supergenes, cytotoxicity and, **41**:310
- Supernatants, humoral immune response and,
45:5
 antigens, **45**:15
 interleukins, **45**:62, 65, 67, 69, 70
 physical interaction, **45**:54
- Superoxide
 interaction with NO, **60**:324–325
 relation to P-selectin, **58**:355
- Suppression
 allotypic and anti-idiotypic, Fe region and,
40:73
 hybrid resistance and, **41**:380, 381, 384,
 387, 389, 395
 IDDM, **48**:143–144
in vitro, induced by V β selective elements,
50:37
- T cell-mediated
Ir genes and, **38**:188
 GAT, **38**:134–141
 GT, **38**:144–149
 H-Y antigen, **38**:175, 179
 immunodominance, **38**:182–184
 insulin, **38**:142–144
 lactate dehydrogenase B, **38**:149–
 152
 lysozymes, **38**:140–142
 tolerance, **38**:129
 malignant B cell proliferation,
40:144–145
- Suppressor antigen-presenting cell, **48**:206
- Suppressor cells, *see also* Suppression, T cell-mediated; Suppressor T cells
 hybrid resistance and, **41**:340, 368, 410
 marrow, **41**:385–387, 391, 392
 in IBD, **42**:299
 immunosenescence and, **46**:249–251
 in MG, **42**:260
- Suppressor macrophages, **50**:176
- Suppressor T cells, **43**:266; **50**:176–177, *see also* Suppression, T cell-mediated
 AChR-specific, **42**:258–259
 adoptive T cell therapy of tumors and,
49:287–289, 293
 antigen-specific inhibition of MOPC-315
 by, **40**:143–144
 autoreactive T cells and, **45**:431, 432
- covert, in Crohn's disease, **42**:300, 301
 historical background, **40**:135–136
 hybrid resistance and, **41**:379, 386, 410
 idiotype-specific inhibition of B cells by,
40:136–143
 immunosenescence and, **46**:251, 252
 isotype-specific inhibition of B cells by,
40:145–148
 lymphocytotoxic antibodies and,
42:291–292
 MG therapy and, **42**:269
 T cell subsets and, **41**:102, 107, 110
- Surface area, exposed, MHR, **43**:35–36
- Surface markers, for memory B cells,
53:235
- Surface receptors, alternative, death induction
 by, **58**:224–227
- Surface variability analysis, **43**:11, 14
- Surrogate heavy chain, **63**:8, 25
- Surrogate light chain
 B cell development, **63**:2–4, 15–25, 29–31
 expression, **63**:5–12, 13–15
 genes, **63**:4–5
 transduction in complex, **63**:25–29
- Switched memory B cells, **59**:340–341
- Switch peptides, **43**:113
- Switch recombination
 B cells, **61**:83
 CD40 signaling, **61**:92–98
 comparisons between VJD and switch
 recombination, **54**:235–237
 control during cell cycle, **61**:116–117
 in DNA replication, **54**:237–238
 experimental systems for studying, **54**:237
 fine sequence specificity, **61**:107–108
 germline transcripts, **61**:111–116
 homeologous recombination similarity,
61:110–111
 Ig production, **63**:58–59
 illegitimate priming model, **61**:108–111
 intrachromosomal deletion, **61**:99
 looping out and deletion, **54**:234
 mutations, **61**:108
 poly(ADP-ribose) polymerase, **61**:110
 protein-binding sites, **61**:117, 120
E47/E12, **61**:117
LR1, **61**:120
NF- κ B/p50, **61**:117, 118–119
NF-S μ , **61**:119
Pax5, **61**:120–122

- sy3 regions, **61**:117–118
s_{ub}bp-2, **61**:119
- proteins involved in switch recombination, **54**:239–240
- RNA splicing, **61**:101–102
- sequential switching, **54**:238–239; **61**:105–107
- sister chromatoid exchange, **54**:234–235
- sites, **61**:102–105
- substrates, **61**:123–129
- transchromosomal switching, **61**:100–102
- vectors, **61**:123–129
- Syk, **59**:261, 324
- IL-2R coupling, **61**:159–160, 190
- Syk kinase, **59**:261, 324
- Syk tyrosine kinase, **63**:28, 134, 137–138
- Sympathetic ophthalmia, **48**:216, 217
- Syntium formation, virus-induced immunosuppression and, **45**:349, 350, 358, 359
- Synergy
- anti-TNF- α therapy, **64**:331–333
 - B cell formation and, **41**:222, 234
 - cytotoxicity and, **41**:284, 296
 - hybrid resistance and, **41**:406
 - T cell activation and, **41**:20, 23, 28, 30
 - Ca2+ ionophores, **41**:15–19
 - cell surface molecules, **41**:2, 12, 14
- Syngeneic mixed lymphocyte responses, autoreactive T cells and activation, **45**:426
- origin, **45**:418–420
- regulation, **45**:432, 433
- Syngeneic preference, hybrid resistance and, **41**:334, 358
- Syngeneic stem cell functions, hybrid resistance and, **41**:388–390
- Synovial cells
- HLA class II molecule, **48**:132
 - as source of chemokines, **55**:111
- Synoviocytes, **64**:288
- CD40 expression, **61**:15, 41
- Synovitis, lymphocytes, **64**:191
- Synovium
- IL-1ra production in macrophages, **54**:187–188
 - lymphocyte homing and, **44**:344
 - normal, **64**:287
 - rheumatoid, **64**:287–288, 291, 315
 - joint destruction mechanism, **64**:320–321
 - neovascularization, **64**:319–320
- Synthetic peptides
- defined recognition sites, *see* Recognition sites, synthetic
 - extracellular processing, **52**:84–85
 - technology, applications, **43**:59
- Syphilis, *see also* Biological false positive serological test for syphilis
- aPL antibodies and antibody subsets, **49**:232
- history, **49**:198–199
- specificity, **49**:234–236, 240, 249–250
- Systemic inflammatory response syndrome, **63**:310
- criteria for, **66**:101, 123
- Systemic lupus erythematosus, **46**:196, 200
- antinuclear antibodies and, **44**:94–97
 - autoantibodies, **44**:125, 136
 - autoimmune response, **44**:131, 132
 - DNA, **44**:97, 98
 - drug-induced autoimmunity, **44**:109
 - histones, **44**:98–100
 - ku, **44**:104, 105
 - mixed connective tissue disease, **44**:111
 - PCNA, **44**:106–109
 - ribosomal RIP, **44**:105, 106
 - RNP, **44**:100–102
 - scleroderma, **44**:120, 124
 - Sjögren's syndrome, **44**:112
 - subcellular particles, **44**:102–104
- aPL antibodies and, **49**:193, 198, 209, 212, 219, 250, 257, 259
- affinity purification, **49**:229
- anybody subsets, **49**:231
- binding, **49**:251–252
- clinical aspects, **49**:206–209, 212
- isotype, **49**:225–227
- LA antibodies, **49**:245–246
- pathogenic potential, **49**:252–255, 257
- specificity, **49**:234–236, 238–239, 248–249
- syndromes, **49**:213–216, 218–222
- CD23 antigen and, **49**:174–175
- human idiotypic network and, **39**:277–278
- Ig heavy-chain variable region genes and, **49**:46–50
- IL-1 and, **44**:193
- and IL-6, **54**:41–42
- lymphocytotoxic antibodies and, **42**:291–292
- NK cells and, **47**:300, 301

- Systemic lupus erythematosus (*continued*)
 overexpression of Fas variant, **58:274**
 patients, low IL-1 production in, **50:175–176**, **186–187**
 pituitary hormones, **63:420–421**
 transfer to hu-PBL-SCID mouse, **50:307**
- Systemic lupus erythematosus, murine, **46:61**, **62**, **65**, **71**, **99**, **100**, *see also* Lupus mouse
 aPL antibodies and, **49:219**, **248**
 EPA and docosahexaenoic acid in diet and, **39:163–164**
 fish oil in diet and, **39:163–165**
 functional abnormalities in
 B cells, **37:309–316**
 macrophage defects, **37:324–326**
 natural killer cells, **37:327**
 other cellular and humoral abnormalities, **37:327–330**
 T cells, **37:316–324**
 genetics, **37:336–337**
 complementarity of genetic backgrounds among lupus mice and role of accelerator and other autosomal genes, **37:341–345**
 inheritance of autoimmune traits, **37:337–338**
 relationship among autoimmune traits and their association with disease, **37:338–341**
 relationship between the traits, H-2 and Ig genes, **37:345–346**
- Ig germline, **46:65**, **66**
 autoantibodies, **46:71–76**
 inducing agent, **46:78**
 murine lupus, **46:66–71**
 somatic mutation, **46:76**, **77**
- influence of sex and sex hormones on pathogenesis of, **37:346–348**
- lupus strains, **46:61–65**
- morphologic manifestations
 arthritis, **37:282–283**
 glomerulonephritis, **37:276–278**
 lymphoid hyperplasia, **37:279–282**
 neoplasia, **37:282**
 other significant histopathologic characteristics, **37:283–284**
 thymic atrophy, **37:278**
 vascular disease and myocardial infarction, **37:278–279**
- recombinant IL-2/vaccinia virus construct on, **50:193–203**
- serologic manifestations
 antierythrocyte antibodies, **37:295–297**
 antiretroviral gp70 autoantibodies, **37:293–295**
 autoantibodies to IgG, **37:298–300**
 autoantibodies to nuclear antigens, **37:286–293**
 complement levels, **37:295**
 cryoprecipitates, **37:295**
 Igs, **37:284–286**
 natural thymocytotoxic antibody, **37:297–298**
- T cell antigen receptor, **46:78**, **79**
 autoimmune strains, **46:83–86**
 expression, **46:86–99**
 gene diversity, **46:79–83**
 treatment of, **37:350–355**
 viruses in, **37:348–349**
- Systemic lupus erythematosus-like autoimmune disease, **54:141–142**
- B/WF1 mouse, **53:144–147**

T

- T3, T cell receptor and, **38:18**, **22**
 T3 membrane proteins, **43:116**
 T9, positive lymphocytes, in Crohn's disease, **42:290**
- T11, *see* CD2
- T15' B cells in mice, **42:72**
- Tac antigen, **59:228**
- Tachykinin receptor, **48:181**
- Tachyzoite antigen, **65:208**
- Talin, leukocyte integrins and, **46:169**
- T-ALL cell line (YT), **42:171**
- TAM, *see* Tyrosine-containing sequence motifs
- TAP, **64:4**
- TAP, **65:47–96**, **96**
 background, **65:47–56**
 disease and
 autoimmune disease and, **65:92–94**
 dysfunction in tumors, **65:94–95**
 polymorphism, **65:92–94**
 viral inhibitors, **65:95**

- gene structure, **65:58**, 59–61
intracellular localization, **65:62**–63
MHC class I assembly and, **65:87**–92
polymorphism, **65:71**–75, 92–94
protein structure
as heterodimer, **65:61**–62, 79, 81
intracellular localization, **65:62**–63
subcellular localization, **65:63**, 65
topology, **65:64**, 65–71
subcellular localization, **65:63**, 65
- TAPI, **64:4**; **65:50**, 85
- TAPI, structure, **65:58**, 60–61
- TAPI/TAP2, **64:4**, 112
function, **64:112**
MHC class I molecules and, **64:113**–115, 117–119
peptide binding, **64:114**–115
peptide transport, **64:112**–113
- TAP2, **64:4**; **65:51**, 85
- TAP2, structure, **65:58**, 60–61
- Tapasin, **65:91**
- TAP complex, **65:75**–76
length specificity, **65:78**
peptide and nucleotide binding, **65:84**–86
sequence specificity, **65:78**–84
transport assays, **65:76**–78
transport model, **65:86**–87
- TAP system, **64:3**–5, 30–31
- Target-binding cells, NK cells and, **47:230**, 269
- Target cell–effector cell interaction, mediated by leukocyte Fc receptors, **51:51**–54
- Target cells
CTL and, **41:137**, 158, 162, 168
carbohydrate moieties, **41:153**, 154
exon shuffling 142, **41:144**, 149
HLA class I antigens, **41:149**–152
cytotoxicity and, **41:269**–272, 300, 319, 320
cell-mediated killing 294, **41:296**–298, 312
cytolysis, **41:273**–284
cytotoxic T cells and, **38:173**
HIV infection and
etiological agent, **47:378**, 384
immune response, **47:409**, 411
immunopathogenic mechanism, **47:400**
hybrid resistance and, **41:363**, 369, 372, 373, 383
antibodies, **41:376**–378
antigen expression, **41:400**, 404
- marrow, **41:387**, 391, 392
in vitro assays, **41:393**–396
- NK cells and, **47:292**, 294
antimicrobial activity, **47:285**–288
cytotoxicity, **47:249**–256, 259
effector mechanisms, **47:234**–242
hematopoiesis, **47:280**
lymphokines, **47:262**–264
receptors, **47:242**–248
spontaneous autoimmune thyroiditis and, **47:451**
- T cell activation and, **41:1**, 14, 30
T cell subsets and, **41:76**, 94, 95
- Targeted chemotherapy, concept of, **56:301**–303
- Target organ, spontaneous autoimmune thyroiditis and
defect, **47:456**–465
genetics, **47:482**, 485, 487
- Target recognition, T cell receptors and, **38:22**–24
- TATA box, TAP genes, **65:60**
- Tax, **65:29**
HTLV-1, activation of NF- κ B, **58:17**–18
- TBAM, **63:47**
- T cell-activating protein, **48:240**–241
- T cell activation, **41:1**, 4, 8, 10, 17, 19, 21, 22, 24, 25, 30, 31; **58:18**–19; **61:148**–149; **62:131**
- accessory activation molecule, **48:304**–306
alternative activation molecule, **48:304**–306
alternative pathway, **48:235**
calcium ionophore, **48:258**–260
CD8a role, **53:84**–85
cell cycle progression, **48:290**
cell surface molecules, **41:1**–15
c-fos, **48:291**–292
c-myb, **48:293**–294
c-myc, **48:292**–293
de novo-induced genes, **48:290**
gene products, **48:290**
gene regulation, **41:26**–30
IL-1 receptor, **48:250**–253
intracellular signals, **41:26**
in vivo, **63:71**
- Ir* genes and, **38:106**, 107, 189
antigen binding, **38:104**–108
competitive inhibition, **38:110**
complementation, **38:79**
cytotoxic T cells, **38:165**, 171

- T cell activation (*continued*)
- expression, **38:58**
 - gene function, **38:91**
 - H-Y antigen, **38:177**
 - I region mutations, **38:91**
 - selection, **38:132**
 - suppression, **38:135, 152**
 - molecular events mediating, **48:227–311**
 - murine system, **41:1**
 - cell surface molecules, **41:10, 11, 13, 14**
 - receptor-mediated signal transduction, **41:23, 25**
 - nuclear events, **48:290–299**
 - phorbol ester, **48:258–260**
 - protooncogene, **48:291–294**
 - Ras proteins, **63:144–145**
 - receptor-mediated signal transduction, **41:19–26**
 - recognition phase, **63:43–44**
 - signal transduction, **48:251–252**
 - synergy, **41:15–19**
 - T cell antigen receptor-CD3 complex, **48:232–235**
 - two stages of, **51:107**
 - tyrosine kinase p56^{lck} role, **53:78–82**
 - T cell antigen receptor, *see* T cell receptor
 - T cell-B cell collaboration, antigens and, **42:75**
 - T cell-B cell interaction, **60:282–283; 63:44, 46, 71–72, 71–72**
 - T cell clones, **38:3, 22; 43:134; 48:78**
 - activation, **43:216**
 - alloreactive, **38:3**
 - antigen binding, **38:99, 105, 106**
 - antigen dose-response curves, **43:202**
 - colony-stimulating factor gene differential expression, **48:89–92**
 - competitive inhibition, **38:109, 110**
 - cross-reactive lysis, **38:187**
 - cytotoxic T cells 165, **38:170**
 - gene dosage, **38:81**
 - gene function, **38:92**
 - germline repertoire, **38:116–118**
 - GM-CSF preferential synthesis, **48:91**
 - immunizing, **38:3**
 - immunodominance, **38:184, 185**
 - lines, **43:196**
 - lymphokine mRNA expression, **48:87–88**
 - macrophages, **38:111–114**
 - reactivity, HLA-Dw14, **48:129**
 - suppression, **38:134, 143, 152**
 - T cell selection, **38:131, 134**
 - T cell-T cell interactions, **38:160**
 - tolerance, **38:119–121, 124, 126, 129**
 - T cell cytokines, *see* T helper cells, cytokines
 - T cell development, **44:207, 255; 58:90–101; 63:1**
 - antigen recognition, **44:209–214**
 - avian, experimental manipulation, **50:102–105**
 - cyclosporin treatment, **50:104–105**
 - embryonic treatment with antibodies, **50:103**
 - thymectomy, **50:102**
 - CD4 vs. CD8, **53:108–113**
 - cellular selection within thymus, **44:245**
 - negative selection, **44:245–252**
 - positive selection, **44:252–255**
 - four-way developmental choice, **51:180–186**
 - CD4 vs CD8 down-regulation, **51:185–186**
 - positive selection vs two kinds of death, **51:175–177**
 - positive vs negative selection, **51:180–183**
 - signaling pathways that control positive selection, **51:183–185**
 - functionally responsible intrathymic transitions, **51:117–139**
 - ontogeny of major thymic lineages, **51:133–138**
 - ontogeny of minor thymic lineages, **51:138–139**
 - summary, **51:139**
 - TCRa,B lineage, **51:118–124**
 - TCRγδ lineage, **51:124–133**
 - mature T cells, properties, **51:85–110**
 - proliferative response, **51:106–110**
 - recognition/auxiliary coreceptors, **51:91–92**
 - recognition structures, **51:86–91**
 - response genes, **51:98–105**
 - signalling through TCR/CD3 and coreceptors, **51:92–98**
 - selection, and mechanism, **51:164–189**
 - negative selection, **51:165–175**
 - positive selection, **51:175–189**
 - summary, **51:189–194**

- thymocytes, functional maturation, **51**:139–164
 auxiliary signaling receptors, **51**:157–160
 functional maturation vs commitment to T cell lineage, **51**:162–164
 mature vs immature characteristics, history, **51**:139–140
 postthymic maturation, **51**:160–162
 response genes, **51**:146–157
 TCR/CD3 complexes and signaling mediators, **51**:140–146
 thymus and its seeding, **51**:110–117
 ontogeny, **44**:214, 215, 221–225
 accessory molecules, **44**:219–221
 function, **44**:221
 receptor expression, **44**:218, 219
 thymocyte, **44**:215–218
 thymocyte subpopulation, **44**:225–245
 in thymus, **51**:112
 T cell epitopes, **43**:4, 60, 73, 78
 autoimmune diseases and, **66**:88–91
 circumsporozoite protein-derived, **60**:108–120
 computational identification, **66**:77–84
 nonmalarial protein-derived, **60**:110–112
 T cell growth factor, **46**:112, 114; **61**:147
 T cell hybridomas, *Ir* genes and, **38**:40
 antigen binding, **38**:99, 107
 antigen processing, **38**:95
 competitive inhibition, **38**:109, 110
 germline repertoire, **38**:116
 suppression, **38**:151
 T cell-T cell interactions, **38**:156, 157, 160
 T cell hybrids
 alloreactive responses to
 B cells, **39**:85–87
 macrophages/dendritic cells, **39**:85–86
 haptens-specific B cells and, **39**:56–57
 T cell leukemias, **38**:18
 anti-Id antibody therapy, **39**:288
 development, **50**:141–142
 mature, receptor gene rearrangements in, **40**:276–284
 T cell lymphocytopenia, steroid therapy and, **42**:289
 T cell lymphoma
 depletion, **58**:272–275
 murine, **54**:289
 T cell receptor, **41**:2–8; **43**:105, 114–117; **45**:107, 108, 131; **48**:78–81; **58**:119–120; **63**:43–44, *see also* Antigen-MHC determinant; T cell development
 α , **64**:69
 and β and γ and δ
 generation of diversity, **58**:115–116
 genomic organization, **58**:114–115
 rearrangement and expression, **58**:116–119
 rearrangement, **58**:150–152; **61**:305–306
 transcription inhibition, **58**:243
 α enhancer, **43**:268
 $\alpha\beta$, **43**:115; **59**:100–105
 antigen receptor dimer, **43**:116–117
 CD3 complex, **43**:135
 chain alignment, **43**:221
 compared to T cell receptor Ad, **43**:175
 developmental control, **58**:157
 expression, **43**:146–147, 165
 genes, **43**:136
 heterodimers, **43**:221, 266
 induction of CD8 $^{+}$ T cell apoptosis, **58**:223
 loci, **43**:140
 lymphocytes, ontogeny, **43**:169
 molecules, **43**:134
 mRNA, **43**:146–147
 protein complex, **43**:134–135
 role in signal transduction, **58**:125–127
 specificity, **58**:124–125, 174
 transcripts, **43**:168
 variable region sequences, **43**:221
 $\alpha\beta$ and $\gamma\delta$
 complexes, **58**:113–128
 diversified
 in human, **58**:305–306
 in mouse, **58**:304–305
 $\alpha\beta$ lineage, **51**:118–124
 β , **43**:136
 germ-line genes, **43**:267
 induction of development, **58**:142–144
 primary structure, **43**:220
 rearrangement, **61**:305–306
 signal mediation, and p56 ck control, in
 thymocyte development, **56**:163–164
 transcription, **43**:268
 β enhancer, **62**:43, 51–52

- T cell receptor (*continued*)
 epitopes, 45:142
 conformation, 45:145, 146
 helical conformation, 45:142–144
 location, 45:146–148
- $\gamma\delta$ gene
 disulfide-linked form, 43:148
 human, 43:140–143
 IDP2 transcript organization, 43:154
 mRNA transcripts, 43:167
 murine genes, 43:136–140
 puzzle, 43:144–145
 rearrangement and diversity, 43:143–144
 role in fetal life, 43:144
- $\gamma\delta$, 43:175–176
 expression by fetal CD44⁺CD25⁺ cells, 58:141
 heterodimers, 43:266
 importance, 43:177–178
 invariant, 58:301–304
 lung, 58:303–304
 skin, 58:301–303
 tongue and female reproductive tract, 58:303
 ontogeny, 43:165–171, 177
 gene expression, 43:266
 protein structure, 43:174–175
 thymic ontogeny, 43:177
- V3, 64:69
- $\gamma\gamma$ homodimers, 43:150
- $\gamma\delta$ lineage, 51:124–133
- $\gamma\delta$ lymphocytes
 activation, 43:171–172
 allogenic target cell recognition, 43:173–174
 anti-CD3 mAb, 43:172
 cell surface expression, 43:162
 cytolysis, 43:172–174
 functional studies, 43:174
 peripheral blood and thymic, 43:161–164
 skin, 43:164–165
- $\gamma\delta$ proteins, 43:146
 CD3 polypeptides, 43:150–152
 cytofluorographic analysis, 43:146–147
 forms, 43:148–149
 human subunit structure, 43:148–152
 identification, 43:146–148
 mobility, 43:150
 murine subunit structure, 43:152–153
 peptide mapping, 43:150
- SDS-PAGE, 43:150
 two-color cytofluorographic analysis, 43:146–147
- δ , 43:150
 hairpin accumulation at, 58:48
- δ gene
 breakpoints, 43:160
 C δ -C α locus organization, 43:155–156
 C6 gene segment, 43:156–157
 chromosomal translocation, 43:160–161
 genomic organization, 43:155–156
 identification, 43:153–155
 IDP2 transcript organization, 43:154
 rearrangement and diversity, 43:158–160, 167
 accessory molecules, 45:108, 109
 CD3 complex, 45:109, 110
 adoptive T cell therapy of tumors and, 49:299
 allelic exclusion, 38:17, 18
 alloreactivity, 45:148–157
 amino acid conservation, 43:221–222
 amphipathicity, 43:204
 antigen, 38:2
 conformation and recognition, 43:194, 223–225
 MHC interaction model, 43:201
 processing, 43:196–197
 specificity, 43:193–194
 antigen binding, 38:98, 99, 104–107
 antigen processing
 class II molecules, 45:117, 118
 class I molecules, 45:118, 119
 exceptions, 45:119, 120
 factors, 45:122–124
 molecular mechanisms, 45:127, 128
 pathways, 45:124–127
 peptides, 45:120, 121
 recycling, 45:121, 122
 antigen-specific, 38:115–117
 autoimmune demyelinating disease and, 49:357
 myelin basic protein, 49:358–360
 usage restriction, 49:363–366
 autoimmune thyroiditis and, 46:265, 318, 319
 cellular immune responses, 46:291
 humoral responses, 46:307
 prevention, 46:312, 314–316
 autoreactive T cells and, 45:417, 419

- B cell formation and, **41**:194, 199, 225, 226, 237
C_a domain, **43**:221
 CDR, diversity, **43**:222
 chromosomal mapping, **38**:16, 17
 class I and class II-specific, **43**:205
C_δ locus deletion, **43**:170
 combining site
 antigens bound to MHC, **43**:215–220
 selections based on receptor specificity, **43**:218–219
 shape, **43**:222
 T cell receptor structure, **43**:220–222
 complex, **58**:128–136
 CTL and, **41**:136, 137, 139, 152, 167, 168, 170
 cytotoxic, **38**:170, 173
 cytotoxicity and, **41**:271, 272, 274
 defined, **64**:1
 encoding, **38**:5–9
 evolution, **52**:110–111
 experimental systems, **45**:131, 132, 137
 expression, peripheral T cells, **38**:13–15
 expression at different anatomical sites, **58**:300–306
 gene defects, in hereditary primary immunodeficiency disorders, **40**:306–309
 gene expression, **43**:135
 peripheral T cells, **38**:13–15
 putative transcription signals, **43**:267
 during T cell ontogeny, **43**:266–267
 tissue specific, **43**:235
 trans-acting negative regulation, **43**:267
 transcriptional regulatory elements, **43**:267–269
 gene organization, **40**:259–262
 gene rearrangements, **58**:31, 36–41, 63, 68–69
 applications to clinical medicine, **40**:264–270
 definition of clonality based on, **40**:295–305
 developmentally programmed, **58**:308–309
 and expression, thymus, **38**:10–13
 genes
 allelic exclusion, **38**:17–18
 α enhancer, **43**:268
 avian, **50**:108–112
 conserved structural features, **50**:111–112
 genomic organization, **50**:108–109
 TCR β diversity, **50**:109–111
 hypermutation, **60**:278–280
 germline repertoire, **38**:118
 homogeneity
 peptidic self model, **45**:128, 129
 self-component, **45**:128
 self-peptides, **45**:129–131
 humoral immune response and, **45**:5, 6
 antigens, **45**:18, 19
 physical interaction, **45**:36, 38, 41, 44–46, 52–54
 Ia molecules, **38**:68
 Ig fold, **43**:221
 Ig heavy-chain variable region genes and, **49**:5
 IL-2 and, **42**:173–174
 immediate responses, mediators for, **51**:143–146
 interaction with antigen bound to MHC, **43**:222
 intracellular selection, **58**:120–121
 J gene segment, **43**:115
 ligands inducing deletion of thymocytes, **58**:170–171
 mAb, SFI, **43**:161
 male-specific, CD4 $^+$ 8 $^+$ thymocytes
 expressing, **58**:261
 maps of Ig-like loci and, **46**:1, 5, 34–45
 chromosomal locations, **46**:6
 mechanisms, **46**:46, 47
 mediated induction of T cell death, **58**:215–224
 mediated signaling
 and CD4, **53**:85–91
 and CD8, **53**:85–91
 in naive and memory T cells, **53**:228–229
 and p56 lk , **53**:85–91
 membrane proximal domains, **43**:221
 methodological breakthroughs in discovery, **58**:88–89
 MHC molecules, **45**:110, 111
 antigen presentation, **45**:115–117
 class II molecules, **45**:112–115
 HLA-A2, **45**:111
 models, **45**:111, 112

- T cell receptor (*continued*)
- MHC-restricted, **38:2**
 - peripheral T cells, **38:14**, 15
 - protein properties, **38:3–5**
 - specific binding, **38:19–21**, 23, 24
 - target recognition, **38:22**, 23
 - thymus, **38:10**
 - V region genes, **38:15**
 - molecule recognition, **43:194–196**
 - murine lupus models and, **46:61**, 65, 78, 79
 - autoimmune strains, **46:83–86**
 - expression, **46:86–99**
 - gene diversity, **46:79–83**
 - NK cells and, **42:198–204**, 205
 - oligoclonal expansion, **58:327**
 - p56^{lck} interactions, **56:161–162**
 - peptides
 - binding, **45:132–134**
 - selection, **45:137–139**
 - peptide vaccination in trimolecular complex, **56:236–239**
 - peripheral T cells, **38:13–15**
 - polymorphic residues, **45:157–160**
 - modeling, **45:160**, 161
 - recognition, **45:161**, 162
 - polymorphism recognition, **43:214–215**
 - precursors, structure and function, **58:144–151**
 - pre-T cell receptor α , **62:42**, 44
 - properties, **38:3–5**
 - recognition of superantigens, **54:121–124**
 - models, **54:122–123**
 - modifications of $\nu\beta$ rule, **54:123–124**
 - restricted recognition, **43:202–203**, 216
 - restriction, **43:195–196**
 - SCID and, **49:384**, 386–388, 396, 400
 - signaling in CD4 $^+$ thymocytes, **58:173–174**
 - signalling through TCR/CD3 and coreceptors, **51:92–98**
 - specific, **43:193**
 - specific binding, **38:19–21**, 23, 24
 - structure, **40:258–259**; **43:220–222**
 - structure–function relationships, **45:135**, 136, 139–142
 - surface antigens of human leukocytes and, **49:77–79**, 81–83, 86–87
 - synthetic T and B cell sites and antigens, **45:233**, 234, 236
 - globular protein antigens, **45:210**, 211, 213, 216
 - immunological considerations, **45:200**, 201
 - prediction, **45:252**
 - viral antigens, **45:219**
 - T3, **38:18**, 19
 - T cell repertoire, **45:162**
 - positive selection in thymus, **45:163–167**
 - recognition, **45:167**, 168
 - T cell subsets and, **41:40–43**, 46, 111
 - H-2 alloantigen recognition, **41:79–81**, 88
 - H-2 molecules in thymus, **41:97**, 99, 106, 107
 - H-2-restricted antigen recognition, **41:51–54**, 56, 60–65
 - TCR1, functional capabilities, **50:101–102**
 - antibodies, embryonic treatment, **50:103–105**
 - migration of subpopulations, **50:96**
 - TCR2, functional capabilities, **50:101–102**
 - antibodies, embryonic treatment, **50:103–105**
 - migration of subpopulations, **50:96**
 - TCR3, functional capabilities, **50:101–102**
 - antibodies, embryonic treatment, **50:103–105**
 - migration of subpopulations, **50:96**
 - TCR/CD3 complexes and signaling mediators, **51:140–146**
 - TCR $\gamma\delta$ lineages, **51:124–133**
 - TH cell
 - experimental autoimmune encephalomyelitis, **48:166**
 - T cell receptor $\tilde{\nu}$ gene, **48:166**
 - thymus, **38:10–13**
 - tolerance, **38:127**
 - transgenes, **59:107–108**, 113–116, 124–127
 - transgenic, **58:154–156**, 159, 166–167
 - triggered CTL-mediated killing, **60:291**
 - triggering, coreceptor molecules
 - modulation of adhesiveness during, **53:83–84**
 - tumor antigen interactions
 - costimulatory pathway-determined immogenicity, **57:288–290**
 - early evidence of recognition, **57:285–288**
 - tumor-specific peptide recognition, **57:288–290**
 - tumor antigens defined by, **62:227–231**

- tyrosine protein kinase, **48**:277–280
V β , interaction with MMTV superantigen, **65**:165–167
V β 2, **64**:69
V β 8⁺, SEB effects, **58**:275
V β selective elements, **50**:2–3, 6–16
 bacterial toxic mitogens, **50**:15–16
 cellular basis of action, **50**:29–32
 B cell role, **50**:31
 dendritic cell role, **50**:31
 responding cells, **50**:29–31
 stimulating cells, **50**:31–32
 transfer of Vbse between cells, **50**:32
 in intrathymic development, **50**:33
 negative selection, **50**:34–35
 positive selection, **50**:34
Mls, **50**:7–10
 definition, **50**:7–9
 multiple loci, **50**:9–10
 molecular basis of action, **50**:16–22
 non-Mls, in murine genome, **50**:11–14
 general nomenclature, **50**:13–14
 not stimulating primary T cell response, **50**:11–13
 null alleles, **50**:14
 questions, **50**:42–47
 biological function, **50**:45
 distribution, **50**:44–45
 molecular nature, **50**:44
 validity, **50**:42–44
V γ 5 and **V γ 6** subsets, **58**:103–104
variable region gene recombination
 diversity and, **40**:262–264
variable regions
 autoimmune demyelinating disease and, **49**:357
 usage restriction, **49**:363–364, 366–367
 multiple sclerosis and, **49**:372–375
V and **C** domains, **43**:115
V(D)J joining of antigen receptor genes
 endogenous substrate, **56**:34–37
 gene assembly, **56**:28–32
 resulting in rearrangement, **56**:47, 49
V gene
 experimental autoimmune encephalomyelitis, **48**:166
TH cell receptor, **48**:166
- V region repertoire and variability, **38**:15–16
retroviruses, endogenous or extrinsic, **50**:37–42
 endogenous retroviral insertions
 inseparable from Vbse, **50**:38–39
 expressions of Mtv in B cells with Vbse expression, **50**:39
 mice transgenic for given Mtv, **50**:40
 mouse mammary tumor virus, **50**:39
 mouse retention of Vbse sequences, **50**:41–42
 Mtv transcripts expressed in activated B cells, **50**:40
 other retroviruses, **50**:40–41
 retroviral sequences bearing no homology to bacterial toxic mitogen, **50**:41
role of MHC molecules, **50**:17–22
 I-A molecules, **50**:20–21
 I-E molecules, **50**:18–19
 incompetent class II, **50**:22
role of T cell receptors and coreceptors, **50**:23–28
 coreceptors, **50**:27–28
 mapping relevant sites on region, **50**:25–27
 T cell receptor, **50**:23–24
 V β region of TCR, **50**:24–25
T cell development and function, **50**:32–37, 35–37
T cell development and function in periphery, **50**:35–37
 clonal inactivation, **50**:35–36
 enhancement of responses to antigen, **50**:36–37
 suppression induced *in vitro*, **50**:37
working model, **50**:4–6
working model, evidence, **50**:28–29
V β -specific antibodies, **58**:275
V-V interface edge strands, **43**:115–116
V-V pairings, **43**:116
what it sees, **52**:10–11
X-linked severe combined immunodeficiency, **59**:248–252
X phage repressor protein response, **43**:209–210
zone for, **59**:312–315, 338–339

- T cell receptor-CD3 complex, **43**:171
 chains, **48**:229-232
 and coreceptors, signaling through, **51**:92-98
 G protein, **48**:306-307
 heterodimers, **48**:229
 phosphatidylinositol 4,5-biphosphate² Ca²⁺ response, **48**:262-265
 hydrolysis, **48**:262-265
 polymorphic proteins, **48**:229
 and signaling mediators, **51**:140-146
 mediators for immediate TCR responses, **51**:143-146
 TCR/CD3 complex, **51**:140-143
 structure, **48**:229-232
 synthesis, **48**:232
 T cell activation, **48**:232-235
- T cell receptor I, **43**:169
- T cell replacement factors, humoral immune response and, **45**:65, 66, 68
- T cell-replacing factor, interfering and, **40**:38
- T cells, **43**:134; **48**:70, 71
- αβ**
 - CD4⁻8⁻, expression of V β 8 TCR chains, **58**:109-110
 - CD4⁺⁸⁻, heterogeneity, function, and specificity, **58**:105-108
 - CD4⁺⁸⁺, heterogeneity, function, and specificity, **58**:108-109
 - extrathymically derived, **58**:111-112
 - negative and positive selection, **58**:311
 - reactivities, **58**:318-319
 - αβ** and **γδ**
 - differences, **58**:300
 - lineage divergence, **58**:141
 - γδ**
 - extrathymically derived, **58**:110-111
 - function, **58**:104-105
 - heterogeneity, **58**:102-103
 - murine, properties of classes of, **51**:128
 - reactivities, **58**:319-324
 - self-tolerance, **58**:313
 - specificity, **58**:103-104
- 20 α SDH expression
- differentiation and, **39**:26-31
 - IL-3-i induced 2, **39**:5, 18-19, 23-24
 - in Thy-1⁺, **39**:21-22, 24, 26-29
- AChR and, **42**:256-260
- activated
- CD44 expression, **54**:294-296
- cytokine and B cell differentiation, **52**:212-213
- cytokines and B cell growth, **52**:203
- defective CD40L expression in, **60**:44-48
- inducing resting B cells to proliferate and differentiate, **52**:191-193
- T cell subsets and, **41**:59-59
- activation, *see* T cell activation
- adjuvant action and, **40**:92-94
- allogeneic, *Ir* genes and, **38**:125, 126
- alternative transmembrane signaling pathways, **48**:307-309
- antagonists, in trimolecular complex, **56**:244-247
- antigen-independent recruitment into tissues by endothelial cells, **50**:274-287
- endothelium and tissue-specific homing of lymphocytes, **50**:284-285
- extravasation of T cells, **50**:285-287
- migration of memory T cells to inflammatory sites, **50**:278-284
- endothelial leukocyte adhesion molecules, **50**:279-281
- expression of endothelial leukocyte adhesion molecules *in vivo*, **50**:283-284
- role of endothelial leukocyte adhesion molecules in lymphocyte binding *in vitro*, **50**:281-283
- migration of naive T cells to lymph nodes, **50**:275-278
- migration of pre-T cells to thymus, **50**:275
- antigen-presenting cells and, **47**:47, 64-74, 76, 77
- antibody responses, **47**:81-86
- antigen presentation, **47**:78-81
- antigen processing, **47**:87, 90-92
- APC-T cell binding, **47**:95-99
- cell surface, **47**:57, 59, 61-63
- immunogenicity, **47**:102-104, 103
- T cell growth, **47**:99-101
- T cells, **47**:86, 96, 99
- tissue distribution, **47**:48, 49, 50, 52-57, 54
- antigen recognition, **43**:134, 223
- antigens, **38**:2
- conformation, **43**:203

- antinuclear antibodies and, **44**:134
apoptosis, pharmacology, **58**:211–296
autoimmune demyelinating disease and
 multiple sclerosis, **49**:369–375
 myelin basic protein, **49**:358, 360–362
 TCR usage restriction, **49**:363–366
autoimmune thyroiditis and, **46**:263, 265,
 266, 317–319
 antigens, **46**:270
 cellular immune responses, **46**:287–291,
 293–298
 experimental models, **46**:273, 275–282
 genetic control, **46**:283
 humoral responses, **46**:299, 302–304,
 306–308
 prevention, **46**:308, 310–312, 314–316
autoimmunity, theories, **56**:453
autoreactive, *see* Autoreactive T cells
BALU, **59**:386
B cell activation, **40**:2–4, 12–13; **63**:43, 46
B cell differentiation and, **40**:31–35
B cell formation and, **41**:205, 212, 225,
 236
B cell precursors, **41**:189, 191, 192, 198,
 199
genetically determined defects, **41**:226,
 227, 230, 231
Ig genes, **41**:204
soluble mediators, **41**:233–235
–B cell interactions
 antigen bridging, **39**:51
 antigen presentation by B cells, *see* B
 cells
and B cells, interphase death and
 nonspecific damage, **50**:69–70
bone marrow, characterization, **53**:171–173
cachectin and, **42**:220, 222
CD1 proteins, **59**:76–90
CD3, T cell receptor and, **45**:109, 110
CD4⁺, **48**:84–86, 198; **60**:38–39, 114,
 271–272, 294–296
 antigen-presenting cells and
 antigen presentation, **47**:64, 75, 80
 cell surface, **47**:57
 immunogeneity, **47**:102
 interaction with T cells, **47**:93, 94
 T cell-dependent antibody responses,
 47:83, 84, 86
 tissue distribution, **47**:50
 CD5 antigen, NK cells and, **47**:227
CD5 B cell, **47**:117, 118, 161, 162
 aging, **47**:122, 123
 anatomic localization, **47**:120, 121
 autoimmune diseases, **47**:148–150
 bone marrow transplantation, **47**:129,
 130
 definition, **47**:118–120
 genetic influence, **47**:147, 148
CD5 cells, antigen-presenting cells and,
 47:47, 55, 63
CD8⁺ antiviral effect, **66**:274–275,
 282–283, 292–294
CD8 cells, antigen-presenting cells and
 antigen presentation, **47**:64, 75, 80, 81
 antigen processing, **47**:92
 immunogeneity, **47**:102
CD11 antigen, NK cells and, **47**:291
 congenital defects, **47**:224
 differentiation, **47**:234
 effector mechanisms, **47**:245, 246
 hematopoiesis, **47**:281
 malignant expansion, **47**:226
 surface phenotype, **47**:207, 208
CD16 antigen, NK cells and, **47**:188,
 189
 adaptive immunity, **47**:291, 295
 alterations, **47**:302
 antimicrobial activity, **47**:288, 289
 antitumor activity, **47**:298
 CNS, **47**:267
 congenital defects, **47**:224
 cytotoxicity, **47**:249, 257, 258, 261
 differentiation, **47**:233, 234
 effector mechanisms, **47**:237, 242,
 244, 247
 genetic control, **47**:222, 223
 hematopoiesis, **47**:275, 279
 lymphokines, **47**:265
 morphology, **47**:218
 reproduction, **47**:271
 surface phenotype, **47**:201–206, 208
 tissue distribution, **47**:219, 220
CD18 antigen, NK cells and, **47**:207,
 224
CD45 and, **66**:18, 26, 35–36
clone, **48**:197
development models, **53**:108–113
HIV infection and
 depletion, **47**:386–389

- T cells (continued)**
- etiological agent, **47**:379, 382, 384, 385
 - immune response, **47**:405, 408
 - immunopathogenic mechanism, **47**:385, 386, 389–392, 394
 - neuropsychiatric manifestations, **47**:405
 - humoral immune response and, **45**:20, 44, 45
 - Ig gene expression
 - lymphomas, **47**:150–152
 - malignancies, **47**:152–159
 - lineage, **47**:128, 129
 - lymphokine-producing phenotypes, **53**:17–19
 - lymphokine production patterns, **48**:85–86
 - malignancies, **47**:123–125
 - marker for activation, **47**:125–128
 - ontogeny, **47**:121, 122
 - physiology
 - autoantibody production, **47**:130–136
 - helper B cells, **47**:137–141
 - monocytoid features, **47**:141–143
 - primordial immune network, **47**:159–161
 - role in V β selective element action, **50**:30
 - sporozoite malaria vaccine and, **45**:303, 304, 319
 - surface antigen, **47**:143–147
 - T cell receptor and, **45**:109, 110, 127
 - T helper, **48**:163
 - virus-induced immunosuppression and, **45**:349–358, 360–363, 366
- CD4 and CD8 molecules and, **44**:265–270, 298–303**
- CD4 molecular biology, **44**:285, 286, 288–293, 295, 297**
 - CD8 molecular biology, **44**:271–273, 278–280, 282, 284**
- CD5 B cell and, **47**:118, 126, 162**
- bone marrow transplantation, **47**:130
 - lineage, **47**:129
 - marker for activation, **47**:125, 126
 - physiology, **47**:131, 137, 140–142
 - primordial immune network, **47**:160, 161
 - surface antigen, **47**:143–147, 144, 147
- CD8 $^{+}$, **48**:84–86, 197, 198**
- CD45 and, **66**:18, 26, 35–36**
- development models, **53**:108–113
- in HIV-1 infection, **66**:283–291, 297–298
- antiviral activity, **66**:274–275
- antiviral therapy, **66**:295–297
- HIV-1-specific lytic activity, **66**:275–277
- viral inhibition, mechanism, **66**:280–291
- humoral immune response and, **45**:20
- lymphokine production patterns, **48**:85–86
- role in V β selective element action, **50**:30–31
- sporozoite malaria vaccine and, **45**:319
- CS-specific T cells, **45**:312
 - interferon- γ , **45**:303–306
 - T cell receptor and, **45**:109, 110
 - TH responses, **61**:358–360
 - virus-induced immunosuppression and HIV, **45**:352, 353, 356–358, 366
 - measles, **45**:339
 - what it recognizes, **52**:6–11
- CD23 antigen and, **49**:149, 151–153, 176**
- biological activity, **49**:167, 169–170
 - cellular expression, **49**:151–153, 155
 - cleavage regulation, **49**:158
 - expression regulation, **49**:159
 - Fc ϵ RII, **49**:164
- CD40 expression, **61**:13, 39–40**
- CD40-L expression, **61**:20**
- CD45 and, **66**:32–36**
- monoclonal antibody studies, **66**:6–12
 - CD45-deficient, **66**:18–22
 - CD45R $^{-}$, life span, **53**:240
 - CD45RA $^{+}$, functional and phenotypic properties, **53**:225–227
 - CD45RO $^{+}$, functional and phenotypic properties, **53**:225–227
- circulating, functions, **53**:199–201
- clonal proliferation, **63**:127
- clones, *see* T cell clones
- colony-stimulating factor
- studies *in vitro*, **48**:77–78
 - studies *in vivo*, **48**:92–98
 - synthesis, **48**:96–98
 - antigen-presenting cell, **48**:97
 - local polarization secretion, **48**:97
 - local response, **48**:96
 - target cells, **48**:97

- complement receptor 1 and, **46**:183, 200, 201
 complement receptor 2 and, **46**:203, 205–207, 213
 costimulation, molecules involved in, **58**:132–134
 costimulator activities of vascular endothelial cells, **50**:267–274
 EC costimulator in polyclonal CD4+ T cell activation, **50**:268–270
 EC costimulator in polyclonal CD8+ T cell activation, **50**:270–271
 EC costimulators in alloantigen responses, **50**:271–272
 EC production of soluble costimulators, **50**:272–274
 cross-reactivity, **43**:200
 CTL and, **41**:165
 cytotoxic, *see* Cytotoxic T cells
 cytotoxicity, islet beta cell, **48**:142
 cytotoxicity and, **41**:271–273, 277, 297
 dependent autoimmunity, **51**:294–296
 dependent B cell responses, **62**:131–157, 144–148
 B7-1/B7-2 ligand family, **62**:135–138, 140–142
 CD28–B7 costimulation, **62**:142–144, 148–157
 CD28/CTLA-4 receptor family, **62**:132–135, 138–140
 costimulation, **62**:138–144, 148–154
 two-signal model, **62**:131–132
 dependent isotype switching, role of IL-4, **54**:249–252
 depleted haploidentical bone marrow, SCID and, **49**:393–396
 depletion, immune reconstitution and, **40**:417–419
 derived colony-stimulating factor, **48**:69–98
 derived lymphokine, *in vivo* studies, **48**:94–95
 development, *see* T cell development
 differentiation, CD4 and CD8 roles, **53**:94–113
 in negative selection, **53**:101–103
 in positive selection, **53**:97–100
 DR antigens and, **42**:293, 294
 effector
 functional differences with memory and naive cells, **53**:227–228
 maturation into, **53**:229–230
 migration pattern, **53**:242–244
 primary and secondary, **53**:227–228
 tissue-specific migration, **53**:244–247
 epithelial, clonotype dominance, **58**:298
 epitopes, *see* T cell epitopes
 experimental autoimmune uveitis, **48**:216
 expressing Epo receptor, **60**:14
 facilitation of hematopoietic engraftment by, **40**:391–392
 factors, Ig-secreting cell differentiation and, **40**:11–12
 Fas effects
 development role, **57**:137
 expression in cytotoxic T cells, **57**:138
 fate determination, **58**:136–139
 Fc ϵ receptors, **43**:301, 303
 Fc receptors on, **40**:67–68
 isotype response and, **40**:212–216
 forbidden, in peripheral immune system, **53**:164–166
 fusion, **38**:283, 286
 genetically engineered antibody molecules and, **44**:86
 G protein, **48**:267–269
 in graft-versus-host diseases, rat, **39**:280–281
 idiotypic network and, **39**:281
 and growth requirements, **53**:167
 helper, *see* T helper cells
 heterogeneity, **48**:83–92
 high-affinity self-reactive, deletion, **43**:219
 HIV-1 disease, **63**:87–89, 90
 SCID-hu thy/liv xenograft model, **63**:99–107
 HIV infection and, **47**:379
 activation, **47**:398–400, 402
 etiological agent, **47**:378, 379, 382
 immune response, **47**:405, 409, 410
 immunopathogenic mechanism, **47**:385, 386, 387–392, 388, 390, 391, 394, 397
 neuropsychiatric manifestations, **47**:405
 human
 interactions with rabbit anti-Id antibodies
 binding to, **39**:265–267
 suppression by, **39**:267–269
 met-enkephalin receptors, **39**:312
 SP receptors, **39**:313
 T8 $^{+}$, Id determinants, **39**:266–267

- T cells (*continued*)
 human, IL-9
 characterization and cloning, 54:80–82
 expression, 54:85
 growth-simulating activity, 54:85–88
 response to, 54:86, 87
 human leukemia, 38:5, 8
 hybridomas, *see* T cell hybridomas
 hybrid resistance and, 41:337, 353,
 364–366, 368
 antigen expression, 41:402–405, 408–410
 effector mechanisms, 41:372–374, 392,
 395, 376–388
 hybrids, 48:78
 hyperprolactinemia, 63:409–410
 IBD and, 42:297, 300–301
 subsets, 42:298
 IELs as, 58:298–300
 IFN- γ production, 62:63–66, 105
 IgE-binding factors, 43:301–303
 IgE biosynthesis and, 47:1, 28–30, 32–39
 antibody response, 47:4–12
 binding factors, 47:12, 13, 15–24, 26–28
 IGF-I, 63:406
 IGF-I-R, 63:402
 Ig gene superfamily and
 evolution, 44:43, 47
 MHC complex, 44:24, 26, 28, 29
 nonimmune receptor members,
 44:31–37, 41, 42
 receptors, 44:9–12, 14, 16, 17, 20–24
 Ig heavy-chain variable region genes and,
 49:52, 54
 IL-1 and, 44:154, 156, 164, 190
 human disease, 44:193
 immunocompetent cells, 44:172, 173,
 175–178
 receptor, 44:179, 181–183
 TNF, 44:185, 186
 IL-2 receptor and, 42:165–166, 177, *see also* Interleukin-2 receptor
 IL-6 effects, 54:21–22
 IL-10 biological effects, 56:8–10
 immune, transfer of idiotypic-suppression
 by, 40:139
 immunosenescence and, 46:221
 lymphocyte activation, 46:221, 222,
 224–232, 234, 236
 lymphocyte subsets, 46:236–239
 regulatory changes, 46:250–252
 stem cells, 46:240, 243–247
 intermitotic life span of, 53:237–241
 intestinal intraepithelial lymphocytes,
 53:163–167
 intrathymic development, 58:139–174
 control of apoptosis during, 58:168–174
 pre-T cell receptor control, 58:142–151
 role of V β selective elements in, 50:33
 negative selection, 50:34–35
 positive selection, 50:34
 and selection, 58:139–174
 $\alpha\beta$ TCR control, 58:151–165
 intrathymic transitions, 51:117–139
 ontogeny of major thymic lineages,
 51:133–138
 ontogeny of minor thymic lineages,
 51:138–139
 summary, 51:139
 TCR $\alpha\beta$ lineage, 51:118–124
 TCR $\gamma\delta$ lineage, 51:124–133
 invariant intraepithelial γ/δ T cells (skin,
 female reproductive tract),
 53:167–169
 ion channel, 48:285–288
Ir genes and
 antigen binding, 38:98–103
 antigen processing, 38:93, 94, 95, 96, 97
 competitive inhibition, 38:109
 complementation, 38:74–78
 expression, 38:40–43, 46, 50–63,
 188–190
 gene function, 38:91–93
 gene function H-Y antigen, 38:178
 Ia molecules, 38:68, 69, 70–72, 89
 I region mutations, 38:86, 89
 macrophages, 38:111–115
 MHC restriction, 38:64–67
 T cell-T cell interactions, 38:158
 tolerance, 38:124
 lamina propria, IBD and, 42:300–301
 leukocyte integrins and, 46:154, 169, 172
 lineages, 58:87–88
 extrathymically derived, 58:110–112
 thymus-derived, 58:102–110
 in lymph nodes, 53:161–163
 lymphocyte homing and
 carbohydrate, 44:362
 high endothelial venules, 44:315
 molecules, 44:349–351
 regional specificity, 44:325, 329, 330,
 334, 335, 339

- lymphocytes and, **38**:283, 286
 lymphogenesis in liver, **53**:169–171
 from lymphoid compartment, origins, **53**:164
 and lymphokine production, **53**:167
 maps of Ig-like loci and, **46**:1, 28, 29, 31
 maternally transmitted antigen and, **38**:313,
 329, 334
 mature, properties, **51**:85–110
 proliferative response, **51**:106–110
 anergy, **51**:109–110
 growth, **51**:106–109
 recognition/auxiliary coreceptors, **51**:91–92
 recognition structures, **51**:86–91
 response genes, **51**:98–105
 combinatorial regulation of, **51**:105
 IL-2 gene regulation, case study,
 51:103–105
 T cell subset characteristics, **51**:100
 T cell subsets, **51**:98–103
 signaling via TCR/CD3 and coreceptors,
 51:92–98
 mediated suppression, *see* Suppression, T
 cell-mediated
 memory, *see* Memory T cells
 MG and, **42**:241–245, 273
 MHC class I-deficient mouse
 CD4⁺, **55**:393–394
 TCRαβ⁺ CD4[−] CD8[−] T cells,
 55:401–403
 TCRαβ⁺ CD8⁺ CD4[−] T cells,
 55:394–401
 TCRγδ⁺ T cells, **55**:403–404
 MHC class II-deficient mouse
 CD4⁺, **55**:424–427
 single-positive compartment,
 55:430–433
 CD8⁺, **55**:427–428
 γδ, **55**:428–429
 MHC class I-restricted, antigen processing
 and presentation to, *see* Antigen
 processing and presentation
 MHC class II-restricted, **60**:305–306
 MHC recognition, **43**:224; **55**:364
 MHC-restricted recognition, **59**:100
 migration to periphery, avian, **50**:96–101
 homing preferences, **50**:96–97
 interstitial T cell surface glycoprotein,
 50:97–101
 migration of TCR1, TCR2 and TCR3
 sub-populations, **50**:96
 murine, **54**:88–93
 deletion and anergy in adult animals
 injected with superantigens,
 54:125–127
 murine lupus models and, **46**:64, 66, 67,
 99, 100
 naive
 adhesion molecules affecting migration,
 53:248–249
 endothelium and migration of to lymph
 nodes, **50**:275–278
 functional differences with memory and
 effector cells, **53**:227–228
 migration
 adhesion molecules and, **53**:248–
 249
 basic pattern, **53**:242–244
 tissue specific subsets, **53**:244–247
 migration pattern, **53**:242–244
 TCR and signal transduction,
 53:228–229
 negative selection, **58**:166–168
 networks in secretory immune system,
 40:217–218
 NK cells and, **47**:188, 190, 197, 294
 adaptive immunity, **47**:291–296
 antimicrobial activity, **47**:289
 CNS, **47**:266
 congenital defects, **47**:225, 226
 cytotoxicity, **47**:248, 255, 257–262
 differentiation, **47**:232
 effector mechanisms, **47**:240, 244, 245,
 247, 248, 257, 265
 genetic control, **47**:222
 hematopoiesis, **47**:272–275,
 277–281
 identification, **47**:196, 198, 199
 lymphokines, **47**:264, 265
 malignant expansion, **47**:227, 228
 morphology, **47**:218
 reproduction, **47**:271
 surface phenotype, **47**:200, 204–213
 tissue distribution, **47**:219–221
 ontogeny, avian
 development, interest, **50**:87
 differentiation antigens, **50**:88–89
 T cell development, experimental
 manipulation, **50**:102–105

- T cells (*continued*)
- embryonic treatment with anti-TCR_I, -TCR₂, -TCR₃ antibodies, **50:103–105**
 - thymectomy, **50:102**
 - T cell migration to periphery, **50:96–101**
 - homing preferences, **50:96–97**
 - interstitial T cell surface glycoprotein, **50:97–101**
 - migration of TCR₁, TCR₂ and TCR₃ sub-populations, **50:96**
 - T cell tumors, **50:105–108**
 - Marek's disease (MDV) virus-induced tumors, **50:105–107**
 - reticuloendotheliosis virus-induced tumors, **50:107–108**
 - TCR₁, TCR₂, TCR₃ cells, functional capabilities, **50:101–102**
 - TCR genes, **50:108–112**
 - conserved structural features, **50:111–112**
 - genomic organization, **50:108–109**
 - TCR β diversity, **50:109–111**
 - thymic attraction and origin, **50:89–90**
 - thymus, diversification in, **50:90–96**
 - embryonic waves of thymocyte development, **50:95–96**
 - intrathymic clonal selection, **50:94**
 - ontogeny of TCR₁, TCR₂, and TCR₃, **50:91–94**
 - ontogeny, T cell receptor gene expression, **43:266**
 - oral tolerance and, **40:224–226**
 - p56^{lck}, signaling role, *see* p56^{lck}
 - paracortical area, **58:431**
 - perforin-negative, **60:293, 303**
 - peripheral
 - classification, **53:159–161**
 - gene expression in, **38:13–15**
 - phenotype and function, immune reconstitution and, **40:412–414**
 - target recognition, **38:22** - V region genes, **38:16**
 - in Peyer's patches, **40:194, 195–197, 206, 210–212**
 - phosphatase, signal transduction, **48:284**
 - phosphoinositide turnover, **48:258–265**
 - direct measurements, **48:260**
 - secondary messenger formation, **48:253–265**
 - phosphorylation, activation-associated, **48:269–285**
 - pituitary hormones, **63:414–416**
 - plasmacytoma development, **64:240**
 - polyclonal CD4+, activation, EC costimulator in, **50:268–270**
 - polyclonal CD8+, activation, EC costimulator in, **50:270–271**
 - potential effector mechanisms, **47:465–469**
 - precursors, migration to thymus, **58:347, 392–394**
 - pre-T cells, endothelium and migration of to thymus, **50:275**
 - production of
 - EAF, **39:216–217**
 - GM-CSF, **39:2, 3, 10–15**
 - IL-2, **39:1, 3, 10–15**
 - IL-3 2-3, **39:10–15** - programmed cell death, **50:66–69**
 - CTL targets, death, **50:69**
 - deprived of growth factors, **50:68–69**
 - hybridomas, activation-induced death, **50:67**
 - non-selected thymocyte, death, **50:67–68**
 - thymus, negative selection, **50:66**
 - proliferation
 - IL-2-independent, **48:302–304**
 - immune complexes and, **40:77–78** - Ir* genes and
 - antigen binding, **38:103**
 - antigen processing, **38:94, 97**
 - competitive inhibition, **38:110**
 - complementation, **38:73, 79, 80**
 - cytotoxic T cells, **38:172**
 - H-Y antigen, **38:176**
 - immunodominance, **38:184, 185**
 - inhibition, **38:68, 69**
 - I region mutations, **38:86, 88**
 - macrophages, **38:111, 113, 114**
 - suppression, **38:136, 138, 141, 147, 149–152**
 - tolerance, **38:125, 126** - protein kinase, signal transduction, **48:284**
 - pulmonary, resident
 - CD4 and CD8 antigens, **58:306**
 - $\gamma\delta$ lineage, **58:303–304**
 - pulmonary hypersensitivity, **59:417–420**
 - pulmonary immunity, **59:389–391**
 - Ras proteins, **63:144–145**
 - receptor-like genes and, **38:10**

- recognition, antigen processing for, **58:420**
 recognition and function, overview, **52:1–4**
 regulation, **48:285, 286**
 regulation of IgA response and,
40:209–212
 regulatory potential, **61:344–360**
 CD8*, **61:358–360**
 cytokines, **61:347–355**
 extrinsic factors, **61:357–358**
 surface molecules, **61:355–357**
 TH-1–TH-2 concept, **61:346–347**
 repertoire, constraints
 germline limitations, **38:115–118**
 positive T cell selection, **38:130–134**
 suppression, **38:134–152**
 tolerance, **38:119–130**
 repertoire, and RA susceptibility
 exogenous events, **56:452–455**
 formation, **56:450–452**
 response
 cytotoxic T cells and, **65:279–280**
 to MMTV
 adult, **65:175–178**
 endogenous Mtv, **65:196–208**
 neonatal, **65:171–175**
 T–B cell interaction, **65:180–188**
 resting, accessory signal, **48:247–248**
 SCID and, **49:381–388**
 graft-versus-host disease, **49:397–398**
 lack of stem cell engraftment, **49:396**
 posttransplant immunocompetence,
49:399–402
 stem cell engraftment, **49:388–391**
 tolerance, **49:398–399**
 transplantation, **49:393–396**
 SCID-hu thy/liv xenograft model, **63:82–83**
 secondary B cell lineage and, **42:73**
 selection, **43:217; 51:164–189**
 negative selection, **51:165–175**
 intrathymic clonal deletion,
51:165–173
 positive selection, **51:175–189**
 functional divergence, **51:186–189**
 ligands and receptors in positive
 selection, **51:178–180**
 vs two kinds of death, **51:175–177**
 positive selection, four-way
 CD4 vs CD8 down-regulation,
51:185–186
 developmental choice, **51:180–186**
 positive vs negative selection,
51:180–183
 signaling pathways controlling,
51:183–185
 selective defect, **59:247**
 self-MHC class I recognition by cytotoxic T
 cell effector cells, **57:314–315**
 self- vs nonself-discrimination, **43:223**
 signaling response to superantigens and
 conventional antigens, **54:127**
 signal transduction
 calcineurin, **63:166–167**
 CTP-binding protein, **48:265–269**
 PTPs, **63:166**
 site-dependent tolerance, **53:201–203**
 soluble factors, B cell responses and,
40:13–14, 15, 26–29
 specific antigen presentation by vascular
 EC, **50:261–267**
 alloantigen presentation, **50:266–267**
 protein antigen presentation, **50:265–266**
 regulation of MHC molecule expression,
50:262–265
 specific for foreign antigens, **43:219–220**
 specificity for peptide antigens, **43:197–206**
 approaches, **43:197–198**
 cytochrome c, **43:198–200**
 hen egg lysozyme, **43:201**
 linear sequences, naturally occurring
 proteins, **43:198–201**
 minimum fragment size, **43:197**
 specific tyrosine protein kinase, CD4,
48:243–244
 splenic, **53:161–163**
 spontaneous autoimmune thyroiditis and,
47:434, 491, 492, 494
 altered thyroid function, **47:461, 462**
 cellular immune reactions, **47:449–452,**
454–456, 455
 disturbed immunoregulation,
47:470–472, 472–474, 475, 477, 481
 genetics, **47:488–490**
 histopathology, **47:442, 444**
 humoral immune reactions, **47:445,**
446
 potential effector mechanisms,
47:466–469
 stimulation by
 B cells, **39:83–85**
 B cell tumors, **39:84**

- T cells (*continued*)
 macrophages, 39:84
 TT-specific B cells, 39:57–58
 subsets, 41:44; 51:98–103
 characteristics defined by cloned lines, 51:100
 dietary EPA and, 39:167
 insulitis and overt diabetes, 51:296–298
 MHC molecules, 59:75–76
 subsets in mouse, 41:39, 110–113
 cell surface molecules, 41:40–50
 H-2 alloantigen recognition, 41:78
 alloreactivity, 41:78–83
 antigen-presenting cells, 41:88–92
 effector phase, 41:92–95
 resting T cells subsets, 41:83–88
 H-2 molecules in thymus, 41:95, 96
 development, 41:96–99
 restricted T cells, 41:99–107
 tolerance induction, 41:107–110
 H-2-restricted antigen recognition, 41:51, 52
 effector phase, 41:75–77
 T accessory molecule function, 41:59–62
 triggering of activated T cells and hybridomas, 41:52–59
 triggering of unprimed and resting T cells, 41:62–75
 substrates, 48:281–283
 supernatants, 54:247
 suppressor, *see* Suppressor T cells
 surface activation molecule, 48:228–253
 surface antigens of human leukocytes and, 49:86, 98, 112, 114, 116–117, 125
 adhesion molecules, 49:83, 85, 87, 89
 antigen-specific receptors, 49:76, 78–79
 Igs, 49:91
 membrane enzymes, 49:112–114
 MHC glycoproteins, 49:80–81
 receptors, 49:98, 101–110
 surface molecules, in communication with APCs, 58:112–139
 target, specific binding and, 38:20
 target interaction, 38:21–23
 as target molecules in trimolecular complex
 anti-CD3 antibody, 56:220–223
 anti-CD4 antibody, 56:223–228
 anti- α/β T cell receptor antibody, 56:228–229
 anti-T cell receptor V β antibody, 56:229–233
 perspective, 56:219
 receptor peptide vaccination, 56:236–239
 vaccination, 56:233–236
 therapy of tumors, *See* Tumors, adoptive T cell therapy
- T cell–T cell interactions, *Ir* genes and gene control, 38:189
 helper factors, 38:153–156
 Ia-like molecule expression, 38:156, 157
 MHC restriction, 38:157–160
- T cell tumors, avian, 50:105–108
 Marek's disease (MDV) virus-induced tumors, 50:105–107
 reticuloendotheliosis virus-induced tumors, 50:107–108
- TCR, *see* T cell receptor
- T-dependent and T-independent B cells, 42:63–65
- TdT, *see* Terminal deoxynucleotidyl-transferase
- Tec, association with c-kit, 60:5–6
- Telencephalin, 64:157
- Tel protein, 64:65
- Temperature, and apoptosis, 58:237–240
- TEMPO-amine spin label, 48:27–28
- TEPITOPE (software), T cell epitope prediction, 66:82–84, 86, 90–91
- Terminal deoxynucleotidyltransferase, 58:117; 62:43, 52; 64:51
 in cortical thymocytes, 39:23, 25
 Ig heavy-chain variable region genes and, 49:31
 V–D–J recombination, 61:296–298
 as V(D)J recombination product
 molecular genetics, 56:68–69
 structure, 56:54–56
- Ternary complex factors
 binding sites, 64:76–77
 domain, 64:76–77
 expression, 64:93
 gene targeting, 64:94
 structure, 64:78
 structure–function studies, 64:93–94
- Testosterone
 autoimmune thyroiditis and, 46:276, 310

- depletion of CD4⁺8⁺ thymocytes, 58:227–229
- spontaneous autoimmune thyroiditis and, 47:480, 481
- Tetanus toxoid
- B cell specificity for, 39:57–58
 - antigen-presenting capacity and, 39:57–58
 - immunization, human
 - auto-anti-Id antibodies and, 39:269–275
 - B- and T cell role, 39:264–269
 - cross-reactive Id and, 39:263–264
 - IgG response, 39:261–262
 - as carrier for (NANP)₃ vaccine, 60:112–113
 - derived epitopes, 60:121–122
 - donor lymphocytes and, 38:283–285
 - heterohybridoma antibodies, 38:288, 289
 - human-human hybridomas, 38:293, 294, 303
 - humoral immune response and, 45:13
 - sporozoite malaria vaccine and
 - antibodies, 45:300, 301
 - human trials, 45:313–315
 - synthetic T and B cell sites and, 45:229, 232, 255, 258
 - virus-induced immunosuppression and, 45:355, 357, 361, 362
- Tethering
- high endothelial venules, 65:367
 - leukocytes, 58:382–383
- Tetrahydrobiopterin, IFN- γ , 62:72
- Tetrahymena*, maternally transmitted antigen and, 38:315
- TFIIBA, concentration, 43:266
- TGF, *see* Transforming growth factor
- TH cell receptor
- experimental autoimmune
 - encephalomyelitis, 48:166
 - T cell receptor V gene, 48:166
- TH domain, Btk, 59:183, 187, 190
- Theiler's murine encephalomyelitis virus, 48:165
- Theiler's virus, responses in MHC class I-deficient mouse, 55:409
- T helper cell responses, 61:374–375
- APCs, 61:361–364
 - CD8⁺ T cells, 61:358–360
 - cell-surface molecules, 61:355–357
- cytokines, 61:347–348
- extrinsic factors, 61:357–358
- T helper cells, 59:319–321, 343
- adoptive T cell therapy of tumors and, 49:289–291
 - expression of antitumor responses, 49:327–328, 332
 - mechanisms, 49:299, 312
- B cell antigen receptor-activated, 55:269–273
- CD23 antigen and, 49:159
- clones, 54:89
- CTL and, 41:136, 137
- cytokines, 46:111
- cross-regulation of differentiation, 46:130–133
 - differential induction, 46:127–130
 - functions, 46:114–117
 - DTH, 46:117
 - helper, 46:117–120
 - human clones, 46:121, 122
 - isotype regulation, 46:120, 121
 - macrophage activation, 46:122
 - human subsets, 46:138, 139
 - immune responses, 46:133–138
 - precursors of differentiation states
 - antigen expression, 46:123–125
 - bulk cultures, 46:122, 123
 - clones, 46:123
 - heterogeneity, 46:125, 126
 - phenotypes, 46:126, 127
 - secretion patterns, 46:111–114
- cytolysis and, 41:276
- cytotoxicity and, 41:274, 297
- function, Crohn's disease and, 42:300, 301
- in β -galactosidase-induced antibody response, 39:88–90
- Id recognizing, 39:257–262
- B cells and, 39:259–260
- IL-2-dependent, *v-abl* effect, 39:41
- Ir* genes and, 38:188
- antigen binding, 38:105–107
 - cross-reactive lysis, 38:187
 - cytotoxic T cells, 38:165, 170–172
 - expression, 38:58–60
 - germline repertoire, 38:118
 - H-Y antigen, 38:175–179
 - immunodominance, 38:183
 - macrophage factors, 38:111

- T helper cells (*continued*)
 suppression, **38**:137–139, 141–144
 T cell-T cell interactions, **38**:154,
 157–160
 tolerance, **38**:124
 leukocyte integrins and, **46**:163
 surface antigens of human leukocytes and,
49:77
 T cell subsets and, **41**:40
 H-2 molecules in thymus, **41**:101–103,
 105
 H-2-restricted antigen recognition,
41:67, 70–72, 75, 77
 TH0, TH1, and TH2, characterization,
58:106–107
 TH1, **48**:86–89
 and IL-10 discovery, **56**:1–2
 TH1 and TH2
 CD4⁺, **52**:175–177
 function roles and cross-regulation,
52:176
 TH1–TH2, allergic responses, **61**:341,
 374–375
 APCs, **61**:361–364
 CD8⁺, **61**:358–360
 cytokines, **61**:347–355
 extrinsic factors, **61**:357–358
 IgE, **61**:342–344, 369–374
 mast cells, **61**:360–361
 nervous system, **61**:364–368
 neuroendocrine factors, **61**:366–367,
 369–374
 psychological factors, **61**:368–369
 surface molecules, **61**:355–357
 T cell regulatory potential, **61**:344–360
 TH1–TH2 concept, **61**:346, 374–375
 TH2, **48**:86–89, **62**:146
 and IL-10
 discovery, **56**:1–2
 functional similarities to cytokines,
56:12–13
in vitro stimulations of primed cells,
56:13–14
in vivo expression correlated to TH2
 responses, **56**:14–15
 PCD induction, inhibition,
58:243
 T helper epitopes, pathogen-derived,
60:140–141
 Theophylline, IL-1 and, **44**:160
- Therapy
 for EAMG, **42**:263–269
 for IBD, **42**:316–319
 for MG, **42**:235, 269–272
 Thermal injury, therapeutic regulation of
 complement system in, **56**:280–281
 Thermodynamics, protein-protein
 interactions, **43**:100
Thermoplasma acidophilum, proteasome
 from, **64**:105–106
 Thioether, linkage procedure for
 chemoimmunoconjugates in cancer
 treatment, **56**:321–322
 Thiol-activated Lysins, **41**:317, 318
 Thoracic duct lymphocytes, hybrid resistance
 and, **41**:356–358
 Threonine
 antinuclear antibodies and, **44**:115
 genetically engineered antibody molecules
 and, **44**:84
 phosphorylation, **48**:270–277
 Threonine phosphatases, as p56^{ck} targets in T
 cell signaling, **56**:160
 Threonine protease, proteasome, **64**:18–19
 Threonine/tyrosine phosphatase, PAC-1,
63:167
 Thrombin, aPL antibodies and, **49**:197
 Thrombocytopenia
 aPL antibodies and, **49**:199, 257–258
 clinical aspects, **49**:209, 222
 isotype, **49**:225–226
 pathogenic potential, **49**:252, 254
 specificity, **49**:250, 252
 syndromes, **49**:213, 215–216, 218, 221
 IL-6 treatment, **54**:48
 Thrombomodulin
 aPL antibodies and, **49**:254
 cachectin and, **42**:224
 downregulation, **66**:111
 Thromboplastin, aPL antibodies and
 antibody subsets, **49**:232
 clinical aspects, **49**:200–201, 205
 specificity, **49**:243–245
 Thromboplastin inhibition test, aPL
 antibodies and, **49**:202–203
 Thrombosis
 aPL antibodies and, **49**:194, 199, 257–258
 antibody subsets, **49**:233
 clinical aspects, **49**:204, 211–212
 isotype, **49**:225–227

- pathogenic potential, **49**:252–254, 256–257
 specificity, **49**:239, 250
 syndromes, **49**:213–221, 223
 venous, CD59-negative platelets in, **60**:91–92
- Thrombospondin, surface antigens of human leukocytes and, **49**:86
- Thrombospondin repeats, **61**:242
- Thromboxane
 aPL antibodies and, **49**:254
 and prostaglandins, **51**:347–352
- Thromboxane A₂, **62**:198
- Thromboxane-A synthase, **62**:198, 200
- Thy-1, **48**:238–239
 activation element, **48**:238
 amino acid sequencing, **48**:238
 dendritic epidermal cells, **43**:164–165
 humoral immune response and, **45**:46, 74
 independent signal-transducing molecule, **48**:239
 isolation, **48**:238
 surface antigens of human leukocytes and, **49**:117
- Thymectomy
 adoptive T cell therapy of tumors and, **49**:288, 300
 avian, **50**:102
 helper T cell cytokines and, **46**:125
 MG and, **42**:253, 259–260, 270–272
 neonatal, autoimmune thyroiditis and, **46**:275, 301, 302
- Thymic attraction, thymocyte precursors, **50**:89–90
- Thymic factors
 pituitary secretion induction, **39**:314–315
 similarity to neuropeptides, **39**:314
- Thymic fragments, human postnatal, and SCID mice, **50**:309
- Thymic hormones, SCID and, **49**:384, 391
- Thymic hyperplasia, MG and, **42**:235, 259–260, 270
- Thymic lineages
 major, ontogeny of, **51**:133–138
 minor, **51**:138–139
- Thymic lymphomas, **54**:92
- Thymic nurse cells, spontaneous autoimmune, thyroiditis and, **47**:455, 456, 475, 491
- Thymic stroma, SCID and, **49**:384, 392–393
- Thymic stromal cells, autoreactive T cells and, **45**:421–425
- Thymidine, **38**:280
 immunosenescence and, **46**:231–234, 243
 T cell development and, **44**:236
- Thymidine kinase, expressing cells, killing, **58**:434
- Thymocyte-activating molecule, **48**:250
- Thymocytes
 accumulation of TCR δ rearrangements, **58**:33
 antinuclear antibodies and, **44**:108, 130
 autoimmune thyroiditis and, **46**:274, 279
 B cell formation and, **41**:198, 199
 CD1 protein distribution, **59**:55–60
 CD3 $^{-}$, anti-CD2 mAb, **48**:236
 CD4 $^{+}$ 8 $^{+}$, **59**:101, 103
 commitment to CD4 or CD8 lineages, **58**:156–160
 deletion, blockade by cyclosporin A, **58**:267
 depletion by testosterone and estradiol, **58**:227–229
 generation and turnover, **58**:151–153
 reduction in ζ -deficient mouse, **58**:146–149
 TCR signaling in, **58**:173–174
 CD4 $^{+}$ 8 $^{+}$, **59**:101, 103
 CD4 $^{+}$ 8 $^{+}$, mature, expressing male-specific TCR, **58**:261
 CD4 $^{+}$ 8 $^{+}$ and CD4 $^{+}$ 8 $^{+}$, generation from CD4 $^{+}$ 8 $^{+}$ precursors, **58**:153–165
 CD4 and CD8 molecules and, **44**:265, 298
 CD8 molecular biology, **44**:270, 271, 278, 279
 CD4 signaling role, **53**:82
 cortical
 TdT expression, **39**:23, 25
 unaffected by IL-2 and IL-3, **39**:24–25
 cytolysis and, **41**:276, 279
 development and selection, CD4/CD8 roles, **53**:94–97
 differentiation, and IL-2, **50**:158–160
 functional maturation, **51**:139–164
 auxiliary signaling receptors, **51**:157–160
 CD2, **51**:157–159
 CD2S, **51**:159–160
 vs commitment to T cell lineage, **51**:162–164
 mature vs immature characteristics, history, **51**:139–140

- Thymocytes (*continued*)**
- postthymic maturation, **51:160–162**
 - response gene inducibility, **51:151–157**
 - cortical cells, loss of responsiveness in, **51:153–156**
 - early, in immature thymocytes, **51:151–153**
 - emergence from paralysis to functional maturity, **51:156–157**
 - response genes, **51:146–157**
 - proliferation, **51:146–151**
 - TCR/CD3 complexes and signaling mediators, **51:140–146**
 - mediators for immediate TCR responses, **51:143–146**
 - TCR/CD3 complex, **51:140–143**
 - helper T cell cytokines and, **46:116**
 - hybrid resistance and, **41:337, 376, 409**
 - Ig gene superfamily and, **44:22, 30, 31, 35, 36, 41**
 - IL-1 and, **44:173, 174, 189, 190**
 - immature, early inducibility of responsive genes in, **51:151–153**
 - Ir* genes and
 - expression, **38:54**
 - T cell suppression, **38:135**
 - T cell-T cell interactions, **38:154, 155, 158**
 - irradiated, biochemical events in, **50:70**
 - lymphocyte homing and, **44:345, 351, 352**
 - maturity, **59:101–105, 108–109**
 - mature vs immature characteristics, history, **51:139–140**
 - mechanism of selection by CD4 and CD8, **53:104–108**
 - medullary, **39:20erSDH expression, 23**
 - murine lupus models and, **46:82, 94–100**
 - negative selection, **58:165–174**
 - non-selected, death, **50:67–68**
 - p56^{lk} control of development
 - allelic exclusion, **56:164–166**
 - differentiation, **56:162–163**
 - maturity, **56:164–166**
 - modes of action, **56:167–168**
 - T cell receptor β -chain signaling, mediation, **56:163–164**
 - phagocytosis and, **38:383, 384**
 - positive selection, **59:99–129**
 - precursor, **43:166**
 - progenitors, inhibition of migration, **54:299**
 - rodent, and DNA damage, **50:58–59**
 - scid*
 - with broken DNA coding ends, **58:37–38**
 - hairpin accumulation at TCR δ coding ends, **58:48**
 - single-positive, **59:101, 103**
 - T cell development and, **44:208, 255**
 - antigen recognition, **44:213**
 - cellular selection, **44:245, 247–249, 252–255**
 - ontogeny, **44:214–221, 224, 225**
 - subpopulation, **44:225–245**
 - T cell receptor and
 - gene expression, **38:14**
 - target recognition, **38:22**
 - V region genes, **38:15**
 - T cell subsets and, **41:96–98, 105–108, 112 V88⁺, SEB-induced depletion, 58:259–260**
 - Thymoma, MG and, **42:235, 256, 259, 270, 272**
 - Thymoma cells
 - CD4 and CD8 molecules and, **44:269**
 - IL-1 and, **44:179, 181, 194**
 - T cell development and, **44:215**
 - Thymopoietin, **42:250**
 - Thymus
 - antigen-presenting cells and, **47:47, 48, 81**
 - antinuclear antibodies and, **44:106, 107, 134, 136**
 - autoimmune thyroiditis and, **46:276, 286**
 - auto-reactive T cells and, **45:417, 421, 423, 433**
 - B cell formation and, **41:191, 223, 236–238**
 - bone marrow cultures, **41:212, 214**
 - genetically determined defects, **41:231, 225–227**
 - population dynamics, **41:207, 208**
 - CD1 proteins, **59:67–72**
 - CD4 and CD8 molecules and, **44:265**
 - CD4 molecular biology, **44:288**
 - CD8 molecular biology, **44:272, 278, 279**
 - CD5 B cell and, **47:120, 121, 129**
 - CD23 antigen and, **49:169–171**
 - colonizing cells, and TCR β cells, **58:139–142**
 - cortex, related cells, and rescue, **58:154–155**

- deletion of superantigen-reactive T cells, **65**:203–204
 dependent antigens, **60**:272–277; **63**:43
 derived T cell lineages, **58**:102–110
 in development, waves of precursors
 populating, **51**:137
 diversification in, **50**:90–96
 embryonic waves of thymocyte
 development, **50**:95–96
 intrathymic clonal selection, **50**:94
 ontogeny of TCR1, TCR2, and TCR3,
 50:91–94
 endothelium and migration of pre-T cells
 to, **50**:275
 fetal, and synchronized thymocyte
 ontogeny, **50**:158–159
 HIV-1 infection, **63**:88
 humoral immune response and, **45**:2, 3
 cellular interactions *in vivo*, **45**:83
 helper T cells, **45**:32
 interleukins, **45**:64
 hybrid resistance and, **41**:337, 352
 effector mechanisms, **41**:372–374, 384,
 390–392
 IGF-I, **63**:391–392
 Ig gene superfamily and, **44**:9, 18, 22
 Ig heavy-chain variable region genes and,
 49:50
 IL-2 receptor subunits, **59**:235–236
 immunosenescence and, **46**:243–247
 independent antigens, **60**:272–277; **63**:43
Ir genes and
 cross-reactive lysis, **38**:187
 cytotoxic T cells, **38**:165, 167, 172
 expression in T cells, **38**:60, 61, 63
 H-Y antigen, **38**:177
 T cell repertoire, **38**:117, 127, 130–134,
 152
 T cell suppression T cell-T cell
 interactions, **38**:166
 lymphocyte homing, **64**:177
 MG and, **42**:250–251, 253, 259–260
 MHC class I-deficient mouse, TCR $\alpha\beta^+$
 CD8 $^+$ CD4 $^-$ T cell development,
 55:394
 MHC class II-deficient mouse, **55**:430–434
 negative selection, **50**:66
 NK cells and, **42**:200–201; **47**:187, 188,
 273, 276
 effector mechanisms, **47**:239
 genetic control, **47**:222
 malignant expansion, **47**:228
 prolactin, **63**:382–383
 prolactin receptor, **63**:395
 pro-T cell homing to, **58**:392–394
 repopulation, immune reconstitution and,
 40:411–412
 SCID and, **49**:383–384, 392–393, 398, 400,
 402
 and seeding, **51**:110–117
 origins and alternatives, **51**:115–117
 thymic environment, **51**:110–114
 spontaneous autoimmune thyroiditis and,
 47:492
 cellular immune reactions, **47**:449–451,
 454
 disturbed immunoregulation, **47**:474–476
 histopathology, **47**:434, 444
 humoral immune reactions, **47**:445, 447
 potential effector mechanisms, **47**:466,
 468
 surface antigens of human leukocytes and,
 49:78–79, 81, 83, 114–117
 synthetic T and B cell sites and, **45**:197
 T cell development and, **44**:207, 208
 antigen recognition, **44**:213
 negative selection, **44**:245–252
 ontogeny, **44**:215, 216, 220, 221, 224
 positive selection, **44**:252–255
 thymocyte subpopulation, **44**:225,
 228–238, 241, 243
 T cell development and programming,
 58:88
 T cell receptor and, **45**:115, 162–168
 T cell receptor genes in, **38**:11–13
 T cell subsets and, **41**:43, 44, 112
 H-2 molecules, **41**:95–110
 H-2-restricted antigen recognition,
 41:51, 59, 69
 V region genes, **38**:16
 Thyroglobulin, **46**:263, 317–319; **47**:491
 altered thyroid function, **47**:456, 459–461
 antigens, **46**:266–272
 autoantibodies, **47**:491, 492
 altered thyroid function, **47**:457, 459,
 460, 462
 cellular immune reactions, **47**:449,
 451
 disturbed immunoregulation, **47**:481
 effector mechanisms, **47**:466–469

- Thyroglobulin (*continued*)
 genetics, 47:482, 484–487
 humoral immune reactions, 47:444–448
 cellular immune responses, 46:287, 288,
 291, 297, 298; 47:450, 451
 experimental models, 46:274–278,
 281–283, 285, 287
 genetics, 47:485
 humoral responses, 46:298, 299, 302–307,
 309; 47:446, 447
 MG and, 42:265
 potential effector mechanisms, 47:466, 469
 prevention, 46:308, 310–312, 314, 315
- Thyroid
 antigen-presenting cells and, 47:76
 IFN- γ effect on, 62:93
 spontaneous autoimmune thyroiditis and,
 see Spontaneous autoimmune
 thyroiditis
- Thyroid cells
 autoimmune thyroiditis and, 46:293–295,
 297
 target, 46:291–298
- Thyroid disease, antibodies to self-antigens
 from human donors, 57:242–244
- Thyroid epithelial cells, spontaneous
 autoimmune thyroiditis and, 47:493
 altered thyroid function, 47:458, 459,
 461–463
 clinical symptoms, 47:440
 disturbed immunoregulation, 47:472
 effector mechanisms, 47:469
 genetics, 47:497
 histopathology, 47:442
- Thyroid-infiltrating lymphocytes, 47:492
 effector mechanisms, 47:466, 467, 469
 immunoregulation, 47:473, 475
- Thyroiditis, *see also* Autoimmune thyroiditis;
 Experimental autoimmune thyroiditis;
 Spontaneous autoimmune thyroiditis
 allergic, murine lupus models and, 46:79
 goitrous, 46:284
 Hashimoto's, 46:317
 antigens, 46:268, 269, 271–275
 cellular immune responses, 46:287–291,
 293
 genetic control, 46:284
 humoral 298–301, 46:303, 305, 306
 pituitary hormones, 63:424
 Thyroid peroxidase, 46:272, 273, 298, 317
- Thyroid-stimulating hormone, *see* Thyrotropin
- Thyrotropin, 47:457, 458, 461; 48:170–172
 antigens, 46:267–269, 272
 experimental models, 46:275, 278, 279, 294
 prevention, 46:311
- Thyrotropin-releasing hormone, 46:279, 311
- Thyroxine, 46:269, 278; 47:437, 439, 457, 470
- TIA-1 granule proteins, polyadenylate binding
 motif, 60:301
- Tissue edema, in IBD, 42:310
- Tissue factor pathway inhibitor, 66:132
- Tissue inhibitor of metalloproteinase-1,
 62:267, 288; 64:297–298
- Tissue inhibitor of metalloproteinase-2,
 62:267, 288
- Tissue plasminogen activator, 46:166
- Tissues, stem cell factor and SCF receptor
 distribution, 55:15–17
- Tissue-specific activator protein, B cell
 genesis, 63:228
- Tissue specificity
 Ig heavy-chain variable region genes and,
 49:3, 52–56
 regulators of complement activation and,
 45:394–398
- Tissue transplantation, therapeutic
 complement inhibition using soluble
 CR1, 56:286–287
- tk*, 62:36
- T killer cells
 cytolysis and, 41:280
 hybrid resistance and, 41:382, 383, 400
 T cell subsets and, 41:40, 54
- TL region, gene products, 52:98–101
- T lymphoblasts
 antigen-presenting cells and, 47:76, 84, 102
 spontaneous autoimmune thyroiditis and,
 47:467, 473
- TMV coat protein, 43:54, 76
- TNF, *see* Tumor necrosis factor
- TNF- α , *see* Cachectin; Tumor necrosis
 factor α
- Tobacco mosaic virus protein
 secondary B cell lineage and, 42:68–69
 synthetic T and B cell sites and, 45:227,
 244
 T cell receptor and, 45:132
- Tocopherol, action of, 65:122
- Tolerance, 59:279–281, 342–344
 B cell, *see* B cell tolerance

- early studies affecting antibody formation, **52**:284–287
 dissection of T cell vs B cell tolerance, **52**:285–287
 tolerance and antibody formation before T and B cell era, **52**:284–285
 immunological memory and, **53**:249–251
Ir genes and, **38**:190
 antigen-specific clones, **38**:121–123
 constraints, **38**:127–130
 cytotoxic T cells, **38**:172
 immunodominance, **38**:184
 induction, **38**:125–127, 188
 MHC restriction, **38**:120
 neonatal, **38**:183
 non-MHC-linked, **38**:123–125
 self MHC, **38**:119, 120
 T cell repertoire, **38**:132, 133
 monoclonal repertoires, **59**:310–311
 perspectives including antigen presentation by B cells, **52**:321–322
 SCID and, **49**:398–399
 self-tolerance checkpoints, **59**:281–342
 site-specific, peripheral lymphocytes, **53**:201–203
- Tolerance induction, **42**:48
 B cell repertoire expression and, **42**:35, 82
 B cell subpopulations and, **42**:77–78
 hybrid resistance and, **41**:356, 361, 409
 Mtv superantigens in, **65**:203–208
 murine lupus models and, **46**:65
 T cell subsets and, **41**:107–110
 TOMS-1, **59**:65
- Tongue, invariant $\gamma\delta$ T cell receptors of murine IEL, **58**:303
- Tonsils, human, isolation of highly purified FDC from, **51**:245, 246
- TOR1, **61**:187
- TOR2, **61**:187
- Tor2, **63**:154
- Torpedo*, **42**:249, 252, 257, 259, 260
- Total body irradiation, SCID and, **49**:397
- Total lymphoid irradiation, hybrid resistance and, **41**:386, 387
- Toxic oil syndrome, with endothelial cell proliferation, **60**:225
- Toxic shock syndrome, **54**:128–133
 clinical criteria used in classification, **54**:128, 129
 fatal, **54**:132–133
- IFN- γ and, **62**:103
Staphylococcus aureus toxins, **54**:129
 TSST-1, **54**:129, 131
- Toxoplasmosis, P30-induced IgG antibody, **60**:123–124
- Tp44, **48**:244, 245–246
- Tp67, **48**:246
- Tp90, **48**:248
- Tpl35–45, **48**:250
- TplO3, **48**:244, 249
- Trabecular meshwork, **48**:206–207
- Trachea, lymphocyte homing and, **44**:330, 332, 333
- TRADD, **61**:31
- TRAF1, **61**:29, **63**:50
- TRAF2, **61**:29, **63**:50
- Transchromosomal switching, switch recombination, **61**:100–102
- Trans-complementation, HLA-DQ hybrid molecule, **48**:119–120
- Transcription, **42**:176
 activation, Ig enhancers, **43**:241–242
 antinuclear antibodies and, **44**:94
 autoantibodies, **44**:137
 autoantigen, **44**:127–129
 scleroderma, **44**:124, 125
 SLE, **44**:104
 autoimmune demyelinating disease and, **49**:368
 autoimmune thyroiditis and, **46**:268, 295
 B cell formation and, **41**:192, 198, 204, 205, 221, 222, 226
 CD1 genes, **59**:7, 9–10
 CD4 molecules and, **44**:288–291
 CD8 molecules and, **44**:275, 277, 279
 CD23 antigen and, **49**:165–167
 CD44 gene, multiple products, **54**:278
 cytotoxicity and, **41**:271, 276, 277, 297
 endogenous substrate activation in V(D)J joining of antigen receptor genes, **56**:41–43
 genetically engineered Ab molecules and, **44**:89
 cloning, **44**:73–75
 expression, **44**:77
 lymphoid mammalian cells, **44**:70
 production, **44**:66
 germline, *see* Germline, C_{II} transcription; Germline transcripts

- Transcription (*continued*)
 HIV infection and, 47:380–382, 399, 401
 human B cell neoplasia and, 38:257, 260, 263, 264
 hybrid resistance and, 41:380, 403
 IgE biosynthesis and, 47:21
 Ig gene superfamily and, 44:43, 46
 Ig heavy-chain variable region genes and, 49:15, 33, 37, 53–56
 IL-1 and, 44:155, 174, 195
 gene expression, 44:159–161
 receptor, 44:179, 181
 immunosenescence and, 46:223, 228
Ir genes and, 38:82
 κB-dependent, 58:3
 in lymphocytes, 62:42, 47–49
 lymphokine synthesis, 48:81
 maps of Ig-like loci and
 MHC, 46:30, 34
 structure, 46:18, 19, 21, 27
 T cell receptor, 46:43, 44
 maternally transmitted antigen and, 38:340, 342
 mRNA, 38:217, 218
 murine lupus models and, 46:80, 95
 NK cells and, 47:265
 SCID and, 49:387
 signal transducers and activators, *see* Signal transducers and activators of transcription
 spontaneous autoimmune thyroiditis and, 47:477
 T cell activation and, 41:3, 27–31
 T cell development and
 ontogeny, 44:216–218
 thymocyte subpopulation, 44:229, 233, 241
 T cell subsets and, 41:41, 43, 48, 97
 Transcriptional factor
 insulin gene, 48:142
 transgenic MHC, 48:142
 Transcriptional regulation, 65:1
 C/EBP, 65:3–8
 chromatin structure, 63:200–201
 IFN-inducible genes, and potential role of
 IRF-1 and IRF-2, 52:272–275
 IκB family, 65:14–15
 IL-2 gene, 48:294–297
 IL-2R α gene, 48:294–295, 297–299
in vivo, 63:198–203
 lymphokine production by Fc ϵ RI $^+$ cells, 53:22
 NF-κB family, 65:11–12, 15–22, 30–32
 NF-IL6, 65:1–3, 8–11, 16–18
 nuclear scaffold, 63:201
 p55, 48:294–295, 297–299
 regulator proteins in nucleus, 63:201–203
 REL family, 65:12–13
 Transcription complex, ISGF3, induction by IFN- α/β , 60:17
 Transcription factors, 63:199–200, *see also specific transcription factors*
 assembly, role in gene rearrangements, 58:30–31
 B cell development, 63:5
 B cell development and function
 c-fos, 63:220–221, 248–250
 EBF, 63:250
 EtS, 63:250
 hox-11, 63:220–221, 244–245
 oct-2, 63:220–221, 243–244
 p50, 63:220–221, 246–248
 Rel proteins, 63:245–248
 B cell genesis, 63:216, 218–221
 bmi-1, 63:220–221, 241–243
 c-myb, 63:220–221, 238–239
 E2A gene, 63:218–219, 232–235
 GATA-2, 63:217, 218–219
 hlx, 63:239–241
 homeodomain proteins, 63:239–241, 243–245
 Ikaros proteins, 63:218–219, 224–226
 pax-5 gene, 63:218–219, 228–232
 PU.1, 63:217, 218–219, 222–224
 sox-4 gene, 63:218–219, 226–228
 binding to regulatory elements of type I IFN genes
 interferon regulatory factors 1 and 2, 52:267–271
 NF-κB, 52:271–272
 other factors, 52:272
 calcineurin-dependent, 60:47–48
 cloning, 43:269
 Dif, in rel family, characterization, 58:14
 phosphorylation, activation by DNA-PK, 58:60–62
 topography, 63:202

- Transduction, in programmed cell death, **50**:73–75
- Transfectants, adhesion studies, **64**:186
- Transfection
- adoptive T cell therapy of tumors and, **49**:320–321, 329
 - B cell formation and, **41**:198, 205
 - CD4 and CD8 molecules and, **44**:266
 - CD4 molecular biology, **44**:292, 293
 - CD8 molecular biology, **44**:272–278, 280, 282, 283
 - CD23 antigen and, **49**:161, 169
 - cell lines, positive and negative regulation of hyaluronan binding, **54**:306, 307
 - CTL and, **41**:138, 139, 144, 149–152, 163
 - cytotoxicity and, **41**:272
 - and expression of CD44 cDNA constructs, **54**:286–287
 - genetically engineered Ab molecules and, **44**:89
 - antigen-combining sites, **44**:84
 - chimeric antibodies, **44**:81, 82
 - expression, **44**:70, 71, 75–77, 79
 - fusion proteins, **44**:85, 86
 - lymphoid mammalian cells, **44**:69, 70
 - Ig heavy-chain variable region genes and, **49**:53
 - IL-1 and, **44**:165
 - Ir genes and, **38**:91, 169
 - κ V gene promoter, **43**:258
 - T cell activation and, **41**:3, 5, 8, 11, 25
 - T cell development and, **44**:210, 212
 - T cell subsets and, **41**:58, 59, 62, 80, 83
- Transferrin
- antigen-presenting cells and, **47**:89
 - B cell formation and, **41**:193, 212
 - immunosenescence and, **46**:222, 227
 - NK cells and, **47**:209, 240
 - SCID and, **49**:385–386
 - surface antigens of human leukocytes and, **49**:115
- T cell receptor and, **45**:124
- virus-induced immunosuppression and, **45**:344, 356
- Transfer RNA, antinuclear antibodies and, **44**:119, 127, 131, 134–138
- Transformation, malignant, p53, **62**:226–227
- Transformed producer cell line, **48**:69
- Transforming growth factor, IL-1 and, **44**:153
 - biological effects, **44**:164, 168, 169
 - human disease, **44**:194
 - structure, **44**:155, 158
- Transforming growth factor α , mRNA, expression by eosinophils, **60**:195
- Transforming growth factor β , **48**:169;
 - 52**:173, 199; **55**:181–183, 205–206; **59**:235, 418, 419; **61**:86, 87–90, 353, 374; **62**:84–85, 90, 105, 269
 - adoptive T cell therapy of tumors and, **49**:287
 - B cell growth and differentiation, **52**:182–184
- bioactivity effects
- bifunctional, **55**:185–186
 - cytokine receptors, **55**:187
 - cytokines, **55**:187
 - extracellular matrix production, **55**:186
 - hematopoietic cells, **55**:186–187
 - immune functions *in vivo*, **55**:188
 - lymphocytes, **55**:187
 - macrophages, **55**:187
 - monocytes, **55**:187
- CD23 antigen and, **49**:160–161
- and IgA expression, **54**:259, 262
- IgA synthesis, **63**:54
- induced eosinophil apoptosis, **60**:184
- inhibition of iNOS, **60**:338
- and its receptor, sources and structure of, **52**:183–184
- production by endothelial cells, **58**:358
- signal transduction
 - formation of complexes, **55**:200–202
 - intracellular, **55**:204–205
- sources and structure, **52**:183–184
- structure, **55**:183–184
- superfamily, **55**:184–185
- synthesis by eosinophils, **60**:196
- tumor immunogenicity affected by, **57**:307–309
- Transforming growth factor $\beta 1$, **61**:86, 88–90, 353, 374
- Transforming growth factor β receptor, **55**:182–183, 195
 - complex formation for intracellular signaling, **55**:200–202
 - endoglin, **55**:198
 - functions, **55**:203–204
 - serine/threonine kinase type, **55**:188, 194–195
 - binding of isoforms, **55**:194

- Transforming growth factor β receptor (*continued*)
 cDNA cloning, 55:188–189, 192
 structure, 55:190–193
 soluble TGF- β -binding proteins,
 55:199–200
 sources and structure, 52:183–184
 type I, 55:195–196, 206
 type II
 binding of isoforms, 55:194
 cDNA cloning, 55:190–192, 205
 signal transduction, 55:206
 structural comparisons with actin
 receptor II, 55:192–193
 type III, 55:196–198
 function, 55:203–204
 type IV, 55:198
 type V, 55:194–195
 Transforming retroviral protein, 48:253
 Transfusion effects, hematopoietic
 engraftment and, 40:386
 Transgene
 and mutations, effect on T cell
 development, 58:147–149
 T cell receptors, 59:107–108, 113–116,
 124–127
 Transgenic approaches, to B cell tolerance,
 52:303–317
 advantages and disadvantages, 52:303–305
 anti-DNA transgenic model, 52:314–315
 anti-H-2K k transgenic model, 52:311–313
 anti-transgenic models in which B cell
 tolerance is absent or only partial,
 52:315–317
 hen egg Lysozyme model, clonal anergy
 vindicated, 52:305–311
 Transgenic MHC, transcriptional factor,
 48:142
 Transgenic spleen cells, preliminary *in vitro*
 studies, 54:414–415
 Transglutaminase, and inhibition of apoptosis,
 58:253
 Translation
 antinuclear antibodies and, 44:128
 CD4 molecules and, 44:289, 290
 CD8 molecules and, 44:275, 277, 279
 IL-1 and, 44:155, 159–162, 164, 195
 IL-2 receptor and, 42:176
 T cell development and, 44:241
 Translocation, 52:35–42
 adoptive T cell therapy of tumors and,
 49:283, 320, 323
 antinuclear antibodies and, 44:120, 128
 chromosomal, *see* Chromosomal
 translocation
 c-myc gene, 64:227–229
 Ig heavy-chain variable region genes and,
 49:18
 immunosenescence and, 46:224, 225, 235
 leukocyte integrins and, 46:160
 maps of Ig-like loci and, 46:3, 26, 32, 45
 MHC ABC protein function, 52:38–42
 MHC ABC protein genes, 52:35–38
 surface antigens of human leukocytes and,
 49:98
 T cell activation and, 41:17–19, 22, 23
 V_H and D_H segments, 62:11, 19
 Transmembrane channel, cell-mediated
 killing and, 41:298
 Transmembrane domain, 65:56
 CTL and, 41:135, 146–148
 Transmembrane signaling, 48:227
 Transmembrane spanning segments, TAP
 protein, 65:66–67
 Transmigration, leukocyte–endothelial,
 58:388–389
 Transplantation, *see also specific type*
 antigen-presenting cells and, 47:75, 76, 80
 bone marrow, for SCID, *See* Severe
 combined immunodeficiency
 CD5 B cell and, 47:128–130
 graft-versus-host disease, 61:47
 haploidentical, SCID and, 49:393–397,
 399, 401
 histocompatible bone marrow, 49:392
 lung, graft-versus-host disease, 59:420–426
 membrane cofactor protein, 61:220
 in MHC class I-deficient mouse,
 55:387–388
 hematopoietic cell grafts, 55:388
 liver cell grafts, 55:391
 pancreatic islet grafts, 55:390
 skin grafts, 55:388–390
 murine lupus models and, 46:64
 NK cells and, 47:187, 293, 302
 differentiation, 47:231, 232
 hematopoiesis, 47:273, 280–282
 organ, graft-versus-host disease,
 59:420–426
 spontaneous autoimmune thyroiditis and,
 47:457, 458
 tissue, therapeutic complement inhibition
 using soluble CR1, 56:286–287

- Transplantation antigens, adoptive T cell therapy of tumors and, **49:281**
- Transplants
cytotoxic T cell, **48:197–198**
delayed-type hypersensitivity, **48:197**
rejection, graft-versus-host disease,
59:420–426
- Transporter associated with antigen processing, *see* TAP
- Transporters, **65:47–56**
ABC transporters, **65:55, 56–57, 59**
antigen processing, *see* TAP
need for, **65:47–49**
- Transport proteins, surface antigens of human leukocytes and, **49:114–116**
- Transposase, V(D)J recombination, **64:57**
- Transverse myelitis, aPL antibodies and, **49:209**
- Transversions, Ars A response and, **42:125, 126**
- Tremimon, in cancer treatment, **56:329, 331–332**
- Trichinella spiralis*, **54:94**
killing
by eosinophils, **39:209–211**
by major basic protein, **39:190**
- Trichuris muris*, **54:94**
- Triggering, leukocytes, **58:383**
- Triglycerides, cachectin and, **42:214–215**
- Trimolecular complex
creation, **52:52–59**
immunotherapeutic strategies
components, **56:219–220**
MHC Ia molecule as target, **56:247–250**
peptide antigens as targets
analog as MHC blockers, **56:244–247**
analog as T cell antagonists,
56:244–247
features, **56:239, 250–251**
peptide determinant as tolerogen,
56:240–244
- T cell as target
anti- α/β T cell receptor antibody,
56:228–229
anti-CD3 antibody, **56:220–223**
anti-CD4 antibody, **56:223–228**
anti-T cell receptor V β antibody,
56:229–233
perspective, **56:219**
receptor peptide vaccination,
56:236–239
vaccination, **56:233–236**
- Trinitrophenyl
antigen-presenting cells and, **47:68, 69, 79, 80**
T cells, **47:86, 97**
tissue distribution, **47:49**
CD5 B cell and, **47:160, 161**
genetically engineered Abs and, **44:80, 81, 89**
- Trinitrophenyl-antigen-binding cells (TNP-ABCs)
helper T cells, **45:27, 33**
interleukins, **45:62–64, 66**
physical interaction, **45:37–40, 42, 44, 47, 50, 51, 53, 54**
- Trinitrophenylated polyacrylamide beads, **52:199**
- Trinitrophenyl-derived T cell, **48:212**
- Trinitrophenyl-keyhole limpet hemocyanin, sensitized B cells, **60:305–308**
- Trinitrophenyl-memory-antigen-binding cells (TNP-MABCs), **45:50, 51**
- trk* receptors, **61:5**
- tRNA, antinuclear antibodies and, **44:119, 127, 131, 134–138**
- Trypanosoma cruzi*
infection, responses in MHC class I-deficient mouse, **55:413–414**
killing by major basic protein,
39:190
- Trypanosoma rhodensiense*, mouse infection
anti-Id antibodies as vaccines against,
39:284
- Trypanosomiasis, cachectin and, **42:214**
- Trypsin, **41:210, 227, 297, 298**
adoptive T cell therapy of tumors and,
49:301, 306
aPL antibodies and, **49:239, 251**
autoimmune thyroiditis and, **46:271, 273, 277, 278, 302**
- IgE biosynthesis and, **47:25**
- IL-1 and, **44:157, 158**
- lymphocyte homing and
carbohydrate, **44:358, 362**
molecules, **44:345, 348**
regional specificity, **44:329**
site-directed mutants, **43:23**
- Trypsin cleavage sites, PGHS-2, **62:174**
- Tryptophan
Ig gene superfamily and, **44:34, 35, 41**
metabolism, IFN- γ , **62:71–72**
- L-Tryptophan, ingestion-induced eosinophilia myalgia syndrome, **60:224–225**

- TS1, **63**:174–176
 TSG-6, **65**:18; **66**:111–112
 TSH, *see* Thyrotropin
 TSICSLYQLE (peptide), **66**:88
 T suppressor cells, *see* Suppressor T cells
 TT, *see* Tetanus toxoid
 Tuberculosis, virus-induced
 immunosuppression and, **45**:335, 336,
 338, 339
 Tubulin, **41**:275, 378
 NK cells and, **47**:302
 Tuftsin, biological activities of, **40**:111–112,
 115
 Tumor allografts
 anterior chamber, **48**:204
 anterior chamber-associated immune
 deviation, intraocular, **48**:207–208
 delayed-type hypersensitivity, **48**:208, 209
 Tumor antigens, **58**:418–422; **62**:217–228,
 240–245
 B7, *see* B7
 BAGE, **62**:229, 233
 blood group antigens, **62**:220–221
 carbohydrate antigens, **62**:219–221
 carcinoembryonic antigen, **62**:221–222
 monoclonal antibody recognition,
 56:305
 CD28, *see* CD28
 c-myb, **62**:227
 c-myc, *see* c-myc
 costimulatory pathway, **57**:305–307
 CTLA-4, *see* CTLA-4
 defined by antibodies, **62**:218–229
 defined by T cells, **62**:227–231
 epithelial antigens, **62**:238–239
 GAGE, **62**:229, 233
 gangliosides, **62**:219–220
 glycoproteins and
 biosynthesis and intracellular transport,
 40:337–341
 immunochemical and molecular profiles,
 40:328–337
 immunological characterization,
 40:324–328
 preclinical models for immunotherapy
 in vitro studies, **40**:341–345
 in vivo reactions in animal model
 systems, **40**:345–351
 gp100, **62**:229, 235–236
 HER-2/neu, **62**:223–224, 237–238
- heterogeneity as barrier to antibody-targeted chemotherapy, **56**:
 352–354
 HPV, **62**:229, 239–240
 idiotypes, **62**:225
 MACE, **62**:228, 229, 231–232
 MART-1, **62**:229, 235
 MELAN-A, **62**:229, 235
 melanocytic differentiation antigens,
 62:233–236
 monoclonal antibody use in identification,
 56:305–306
 mucin recognition, **57**:302–303
 protein-derived
 abl oncogenes, **57**:296–297
 bcr oncogenes, **57**:296–297
 mutated p53, **57**:295–296
 overexpression resulting in, **57**:
 299–302
 ras oncogenes, **57**:297–299
 tum⁺, **57**:293–295
 serological detection, **57**:283–284
 T cell recognition
 early evidence, **57**:285–288
 tumor-specific peptides, **57**:288–290
 virus-induced tumors, **57**:290–293
- Tumor cells
 genetic modification, **58**:425–435
 lysis, **43**:173
 transfected, effect on preexisting tumor,
 58:432–433
- Tumor growth, NK cells and, **42**:182,
 183
- Tumorigenesis
 anterior chamber, **48**:213–215
 anterior chamber-associated immune
 deviation, **48**:213
 pax family, **63**:229–230
 role of IL-9 and mouse T cells,
 54:88–93
 X-linked agammaglobulinemia,
 59:147–148
- Tumor immunology, **52**:104–106
- Tumor immunotherapy, preclinical model
 in vitro studies, **40**:341–345
 in vivo reactions in animal model systems,
 40:345–351
- Tumor-infiltrating lymphocytes, **58**:437–438;
 64:180, 197
 adoptive T cell therapy and, **49**:284, 293

- Tumor necrosis factor, **41**:235–237, 285, 298, 374, 400, 401; **48**:70, 71
adoptive T cell therapy of tumors and 301, **49**:306–309
anti-TNF, **66**:124, 132, 150
antitumor effects in preclinical trials, **56**:354
autoimmune demyelinating disease and, **49**:360
autoimmune thyroiditis and, **46**:294, 295
biochemistry, **66**:105–107
cardiomyocytes and, **66**:114
circulating levels, **66**:104–107
complement receptor 1 and, **46**:192
in endotoxemia, **66**:114–116
eosinophil activation, **39**:216
helper T cell cytokines and, **46**:112, 114, 115
HIV infection and, **47**:394, 402–404
IL-1 and, **44**:153–155, 185–188
biological effects, **44**:165, 166, 168, 169
gene expression, **44**:161
human disease, **44**:192, 195
human joint fluid, **44**:191, 192
immunocompetent cells, **44**:178
structure, **44**:155, 157, 158
synergism, **44**:188, 189
systemic effects, **44**:170
in vitro effects, **66**:110–112
in vivo effects, **66**:112–114
leukocyte integrins and, **46**:170, 172
maps of Ig-like loci and, **46**:30, 33
mediated bystander killing, **60**:304
multiple myeloma pathogenesis, **64**:258
neutrophils and, **66**:111
NK cells and, **47**:286, 289, 290
differentiation, **47**:233
effector mechanisms, **47**:251, 262, 265, 266
hematopoiesis, **47**:278, 279, 281
reproduction, **47**:271
surface phenotype, **47**:213
regulation, **66**:108–110
regulators of complement activation and, **45**:397
sepsis and, **66**:104, 126–127, 150
animal models, **66**:117–121
clinical, **66**:121–124
treatment, **66**:124–126
signaling events, **65**:117
surface antigens of human leukocytes and, **49**:98
synergy between IFN- γ and, **62**:81–82
vascular endothelial injury caused by, **56**:351
Tumor necrosis factor α , **48**:169; **52**:172–173; **61**:5, 22, 24, 25–26; **62**:81, 86, 139; **63**:303–304, *see also* Cachectin
allergic airway inflammation, **62**:285
allergic inflammation, **60**:169–170
antibody pretreatment, **60**:336–337
anti-TNF- α therapy, *see* Anti-TNF- α therapy
and autoimmunity, **50**:211–212
B cell proliferation, **63**:55
cardiovascular risk factors and, **64**:322
cytokine regulation of, **64**:315, 316
granulomatous lesions, **62**:281, 282
IGF-I, **63**:390
IL-1, **64**:293, 316
IL-8, **64**:294, 295, 316
mRNA, eosinophils expressing, **60**:200
NF- κ B induction, **58**:7–9
production
by crosslinkage of Fc ϵ R on peritoneal mast cells, **53**:9
by factor-dependent mast cell lines, **53**:8–9
by Fc ϵ RI $^+$ cells, **53**:9
by human mast cells, **53**:14–15
pulmonary inflammation, **62**:259, 260, 267, 273, 288–289
in hypersensitivity granuloma formation, **62**:280–281
regulation in IC-mediated inflammation, **62**:279
upregulation of vascular E-selectin and ICAM-1, **62**:276, 277
rheumatoid arthritis, **64**:293, 296–298, 316
cell trafficking, **64**:317–319
joint destruction mechanisms, **64**:320–321
neovascularization, **64**:319–320
systemic inflammatory response syndrome, **63**:310
TH responses, **61**:353
trimerization, **61**:26
Tumor necrosis factor β , **61**:5, 353; **63**:303–304
Tumor necrosis factor inhibitors, **63**:305

- Tumor necrosis factor receptor, **63:50**, 276, 284, 303–307, 313–314; **66:107**–108, 116
 genes, **61:25**
 p55, death domain, **58:135**
 superfamily, **61:54**; **63:45**, 49
 CD40 antigen, **61:2**–4
 CD40-L, **61:17**–22
 other members, **61:4**–11, 22–24
 signal transduction, **61:26**–27
 soluble forms, **61:15**–17, 25–26
 viral homologs, **61:10**
- TNF-R1, **61:5**, 7, 17; **63:276**, 305
 gene, **61:8**
- TNF-R2, **61:5**, 7, 8, 17; **63:276**, 305, 309
 gene, **61:8**
- Tumor necrosis factor α receptor, **42:225**–226
- Tumor necrosis factor receptor related protein, **61:7**, 8
 gene, **61:8**
- Tumor necrosis serum, **41:406**
- Tumor rejection
 adoptive T cell therapy of tumors and, **49:286**, 296, 318, 333
 B cells, **49:312**–315
 macrophages, **49:306**–312
 mechanisms, **49:299**–306
 NK cells, **49:315**–318
 intraocular, *see* Intraocular tumor rejection
- Tumors, *see also* Cancer
 adoptive T cell therapy, **49:281**–286, 332–335
 antigen recognition, **49:318**–324
 expression of antitumor responses, **49:324**–325
 B cells, **49:330**–332
 proliferation, **49:325** –330
 mechanisms of tumor rejection, **49:299**
 B cells, **49:312**–315
 macrophages, **49:306**–312
 NK cells, **49:315**–318
 T cell subsets, **49:299**–306
 principles, **49:286**
 requirement, **49:294**–299
 tumor burden, **49:286**–294
 antigen-presenting cells and, **47:66**, 67, 100
 autoreactive T cells and, **45:428**–430
 barriers to antibody-targeted chemotherapy
 perfusion, **56:350**–352
- tumor antigen-heterogeneity, **56:352**–354
 vasculature, **56:350**–352
- B cell formation and, **41:186**, 204, 230, 233, 237
 B cell precursors, **41:188**, 193, 194, 196
 bone marrow cultures, **41:214**, 217
 inducible cell line, **41:220**, 221
 lineage fidelity, **41:195**, 196
- CD5 B cell and, **47:118**, 125
 Ig gene expression, **47:150**, 151, 156
 marker for activation, **47:126**
- CD8 molecules and, **44:271**, 273, 276
- CD44 expression and metastasis, **54:322**–325
 cell migration, role of CD44 in metastasis, **54:318**–325
- chimeric, and IL-6, **54:51**
- CTL and, **41:135**, 138, 139, 154
- cytotoxicity and, **41:270**, 284–286, 293, 313
- defined by antibodies, **62:218**–227
- defined by T cells, **62:227**–240
- etiology, stem cell factor role
 brain, **55:77**
 gonadal, **55:74**–76
 hematopoietic, **55:72**–74
 lung cancer, **55:77**–78
 mast cell leukemia, **55:74**
 melanoma, **55:74**–76
- ganglioside antigens and, **40:351**–352
 properties of cell-associated gangliosides, **40:353**–355
- properties of ganglioside antigens
 associated with tumor cells,
 colon cancer, **40:364**–366
 melanoma, **40:357**–364
 neuroblastoma, **40:355**–357
- genetically engineered antibody molecules and, **44:87**, 88
- hematopoietic, stem cell factor role in
 etiology of, **55:72**–74
- humoral immune response and, **45:7**, 24
- hybrid resistance and, **41:338**
 antigen expression, **41:397**, 402–404
 effector mechanisms, **41:375**, 378, 380, 382, 387, 393, 394, 396
- leukemia/lymphoma cells, **41:358**–369
- IFN- γ and, **62:96**, 101–103
- Ig gene superfamily and, **44:39**

- Ig heavy-chain variable region genes and, **49**:9, 11
IL-1 and, **44**:166, 168, 178, 193
IL-5 effects, **57**:177–178
 rejection, **57**:174
IL-5 role, **57**:177–178
immune response, **57**:281–326
 antigens, *see* Tumor antigens
 B7-transfected cells, **57**:320–321
 cytokine genes in engineering of tumor cells, **57**:321–323
 escape mechanisms, **57**:324–325
 immunogenicity, factors contributing to low level
 antigen expression, low, **57**:303–305
 costimulatory molecules, lack of, **57**:305–307
 IL-10, **57**:307–309
 immune suppressive factors, **57**:307–309
 MHC class I expression, low, **57**:303–305
 TGF- β , **57**:307–309
 tumor environment modification, **57**:309–312
MHC expression increased by transfection
 allo-MHC, **57**:312–313
 self-MHC class I, **57**:314–320
 self-MHC class II, **57**:313
 tumor antigens, *see* Tumor antigens
induction by MMTV, **65**:153–156
leukocyte integrins and, **46**:150
maps of Ig-like loci and, **46**:8
metastasis and extracellular matrix, **54**:318–322
NK cells and, **47**:187–189, 295–300
 CNS, **47**:267
 differentiation, **47**:231, 234
 effector mechanisms, **47**:235, 238, 257, 259, 262
nonimmunogenic, **58**:419
regulators of complement activation and, **45**:395, 397, 402
reticuloendotheliosis virus-induced, **50**:107–108
solid
 adoptive T cell therapy and, **49**:284, 293
 anti-Id antibody therapy, **39**:291–292
spontaneous autoimmune thyroiditis and, **47**:450
studies with B cell lines and, **52**:219–220
surface antigens of human leukocytes and, **49**:115
TAP dysfunction in, **65**:94–95
T cell activation and, **41**:6
T cell development and, **44**:209
T cell receptor and, **45**:117, 118
T cell subsets and, **41**:74, 91
UV-induced, adoptive T cell therapy of tumors and, **49**:281, 283–284
Tumor-specific transplantation antigens, characteristics, **57**:285
Tumor transplantation, **49**:281–282, 286
TUNEL assay, **60**:279
 HIV-1 studies, **63**:102, 104–105
Tunicamycin
 CD23 antigen and, **49**:158, 162
 complement receptor 1 and, **46**:187
 complement receptor 2 and, **46**:201
 IgE biosynthesis and, **47**:18, 19
Turner's syndrome, **63**:409
12/23 Rule, V(D)J recombination, **64**:45, 53, 59
Twins studies, RA susceptibility
 discordance in identical twins, **56**:448–449
 formal genetics, **56**:433–435
 somatic nonidentity, **56**:446–449
Two-color cytofluorographic analysis, T cell receptor $\gamma\delta$ proteins, **43**:146–147
Two-color immunofluorescence analysis, NK1 antigen and, **42**:189
Two-signal model, T cell activation, **62**:131–132
Txk gene, **59**:184
Tyk2 Janus kinase, **62**:68
 in IFN- α/β response, **60**:7–9
Tyr385, **62**:175, 178, 180
Tyrosinase, **62**:233–234
Tyrosine, **42**:97
 CTL and, **41**:135, 148, 162
 phosphorylation, **48**:277–283
 surface antigens of human leukocytes and, **49**:77
Tyrosine-containing sequence motifs, **58**:127–128, 130
Tyrosine kinase, *see* Protein-tyrosine kinase

Tyrosine phosphorylation, **48**:277–283
 B cell antigen receptor-induced, targets, **55**:233–234
 coupled to cytokine binding, **60**:4–9
 mIg-induced, targets, **55**:234–249
 protein substrates, and lymphokine production, **53**:5–6
 protein tyrosine, B cell antigen receptor-induced, **55**:231–233
 role in signal transduction, **54**:418–419
 Vav, **60**:16

U

U937 cells, PGHS-2 expression regulation, **62**:194–195
 U2666 myeloma B cell, **48**:182
 Ubiquitin conjugation, **65**:117
 lymphocyte homing and, **44**:348
 Ubiquitination, **64**:6, 107
 Ubiquitin–proteasome system, **64**:3, 5–8
 Ubiquitynylation, **63**:396
 Ulcerative colitis, *see also* Inflammatory bowel disease
 antibody secretions and, **42**:304
 anti-TNF- α therapy, **64**:309–310
 IgA levels in, **42**:298
 Ultrastructure, purified tonsillar B cells cultured in CD40 system, **52**:210–211
 Ultraviolet light, induced tumors, adoptive T cell therapy of tumors and, **49**:281, 283–284
 Unfolding, **52**:2
 and proteolysis of antigens, **52**:22–78
 Ungulate lentiviruses, **52**:430–432
 bovine immunodeficiency-like virus, **52**:432
 bovine leukemia virus, **52**:432
 equine infectious anemia virus, **52**:431
 Maedi-visna virus, **52**:431
 Unresponsiveness, specific, hybrid resistance and, **41**:384–386
 Upper respiratory tract, pulmonary immunity regulation and, **59**:372–373
 Upstream induction sequence, **65**:25
 Urea-denatured OVA, IgE biosynthesis and, **47**:29, 30, 33
 Urine nitrate
 -NMMA effect, **60**:343

production in septic trauma patients, **60**:351
 Uromodulin, **63**:277
 U snRNA gene promoters, **43**:260, 262
 Uterus rodent
 edema, estrogen-induced eosinophilia and, **39**:227–228
 estrus cycle, eosinophil infiltration and, **39**:227
 stromal cells, IL-1ra production, **54**:188
 UV5C25 anterior chamber, **48**:213–215
 delayed-type hypersensitivity, **48**:214
 Uveitis autoimmune, *see also* Experimental autoimmune uveitis immune regulation, **48**:215–216
 pituitary hormones, **63**:425

V

V1-2 segment, **62**:18
 V1-3 segment, **62**:18
 V1-8 segment, **62**:18
 V1-12P pseudogene, **62**:23
 V1-18 segment, **62**:24
 V1-24P segment, **62**:16
 V1-45 segment, **62**:17
 V1-58P segment, **62**:16, 17
 V1-69 segment, **62**:18, 19
 V2-5 segment, **62**:18
 V3-7 segment, **62**:18, 19
 V3-9 segment, **62**:18, 20
 V3-11 segment, **62**:18
 V3-13 segment, **62**:20
 V3-15 segment, **62**:18, 19
 V3-16P segment, **62**:16
 V3-20 segment, **62**:17, 20
 V3-21 segment, **62**:20
 V3-23 segment, **62**:19
 V3-30 segment, **62**:18–19, 20
 V3-33 segment, **62**:20
 V3-35 segment, **62**:17
 V3-38P segment, **62**:16
 V3-43 segment, **62**:20, 25
 V3-47P segment, **62**:20
 V3-48 segment, **62**:19, 20
 V3-53 segment, **62**:18

- V3-54P segment, **62:16**
 V3-60P segment, **62:25**
 V3-62P segment, **62:25**
 V3-64 segment, **62:17**
 V3-72 segment, **62:17**
 V3 loop, HIV
 multiple antigen peptides with, **60:131–134**
 in octameric multiple antigen peptides,
 60:137–138
 V3 region, **62:24**
 V γ 4, expressed in lung, **58:304**
 V4-4 segment, **62:18, 25**
 V4-28 segment, **62:17, 20, 25**
 V4-31, **62:20**
 V4-39, **62:18, 19**
 V4-59, **62:18, 19**
 V4-61, **62:19**
 V5-51, **62:19**
 V6-1 segment, **62:18**
 V7-81 segment, **62:17**
 V13C segment, **62:24**
 V54 segment, **62:24**
 Vaccination
 antigen-based strategies, **58:435–437**
 cancer, strategies, **58:427, 433**
 efficacy of different clones, **58:430–431**
 GMS/BCG vaccine, **62:220**
 HIV infection and, **47:408, 411, 413**
 MMTV infection, **65:190–191**
 multiple sclerosis and, **49:374–375**
 pulmonary diseases, **59:409–412**
 and tumor-specific antigens, **58:418–420**
 Vaccines, **52:106–107; 65:323**
 HIV, **65:320–322**
 immune, autoimmune and
 immunodeficiency states, **52:106–107**
 (NANP) $_n$ -TT, for *P. falciparum*,
 60:112–113
 and *Neisseria* strain specificity, **60:134–135**
 neutralizing multiple *Chlamydia* serovars,
 60:136
 pneumococcal, mutation to, **60:277**
 SIV, **65:321–322**
 SIV as model for, **52:451, 452**
 sporozoite malaria, *see* Sporozoite malaria
 vaccine
 subunit, for dental caries, **60:135–136**
 synthetic T and B cell sites and, **45:195,**
 197, 260–264
 bacterial antigens, **45:231, 232**
- candidate synthetic peptide, **45:252–259**
 globular protein antigens, **45:213**
 parasitic antigens, **45:228, 230**
 peptides, **45:239**
 prediction, **45:251, 252**
 viral antigens, **45:217–219, 222, 223,**
 225, 226
- TT—multiple antigen peptides, **60:121–122**
 virus-induced immunosuppression and,
 45:336, 339, 340
- Vaccinia virus, *see also* Interleukin-2/vaccinia
 virus
 cytotoxic T cells and, **38:162–166,**
 180–185
 regulators of complement activation and,
 45:409
 responses in MHC class I-deficient mouse,
 55:407–408
- Vaccinia virus complement-control protein,
 61:241
- Valency, antigens, **59:302–305, 308**
- Valerylsalicylate, **62:187**
- VAP-1, **64:147, 161–162**
 and L-VAP-2, **58:377–378**
 lymphocyte–HEV interaction, **65:365**
- Vap-1, B cell, tyrosine phosphorylation after
 anti-Ig stimulation, **55:243**
- Variable domain structures, superposition,
 43:18–20
- Variable region, *see also* Heavy chain variable
 region
 diversity, **42:95–96, see also Arsonate**
 idiotypic system, A/J mouse
- gene expression, *see also* B cell repertoire
 expression
 evolutionary selection vs random somatic
 events in, **42:8–9**
 age factor, **42:30–32**
 antigen responsive cell populations,
 42:19–22
 clonal expansion, **42:32–34**
 fetal and neonatal development,
 42:26–30
 junctional diversity, **42:15–18**
 predominant clonotype expression,
 42:22–26
 segment selection, **42:18–19**
 somatic mutations, **42:9–15**
 mechanics of, **42:2–5**
 secondary B cell lineage and, **42:69–71**

- Variable region (*continued*)**
- secondary B cells and, **42:72–73**
 - unresolved issues in, **42:5–8**
 - Ig heavy-chain, *see* Heavy chain variable region, genes
 - recombinational mechanisms involved in joining segments of, **40:253–256**
 - T cell receptor, recombination creates diversity, **40:262–264**
- Variant surface glycoprotein, containing diacylglycerol**, **60:63–64**
- Variation, somatic, T cell receptors and**, **38:16**
- Variation pattern**
- germline sequences and, **42:127–128**
 - J558 V_H* gene and, **42:117, 125–127**
- Varicella Zoster virus, antibodies from combinatorial libraries**, **57:221–222**
- Vascular addressins**
- lymphocyte–HEV interaction, **65:361–363, 377**
 - lymphocyte homing and, **44:322, 353**
 - mucosal, lymphocyte–HEV interaction, **65:362–363**
- Vascular adhesion protein-1, *see* VAP-1**
- Vascular cell adhesion molecule-1, *see* VCAM-1**
- Vascular endothelial cells, specific antigen presentation by**, **50:261–267**
- alloantigen presentation**, **50:266–267**
- protein antigen presentation**, **50:265–266**
- regulation of MHC molecule expression**, **50:262–265**
- Vascular endothelial growth factor**, **64:319–320**
- Vascular endothelial injury, TNF as cause**, **56:351**
- Vascular leak syndromes, NO role**, **60:327–330, 347**
- Vasoactive intestinal peptide**, **48:170–172; 61:95**
- cAMP in lymphocytes and, **39:311**
 - in cerebral artery walls, **39:304**
 - gene, **65:20**
 - Ig production by Peyer's patches and, **39:311**
 - lymphocyte adhesion and, **64:153**
 - production by neutrophils and mast cells, **39:316**
- smooth muscle relaxation**, **39:303, 306**
- immediate hypersensitivity and**, **39:306**
- vascular dilatation**, **39:302–303, 306**
- immediate hypersensitivity and**, **39:306**
- Vasoactive intestinal peptide receptors**
- kinetics and properties, **39:312–313**
 - on lymphocytes, **39:312**
- Vasodilation**
- early event in inflammation, **60:326–327**
 - NO-induced, **60:347–348**
- Vav**, **60:16**
- Vav**, **62:42, 45–46**
- tyrosine phosphorylation, **60:16**
- Vbse, *see* V β selective elements**
- VCAM-1**, **52:363; 62:258, 267; 64:154; 65:359**
- cell trafficking, **64:317–319**
 - downregulation, **64:172**
 - expression in allergic inflammation, **60:172–173**
 - ligand for $\alpha 4\beta 1$ integrin, **58:374–375**
 - lymphocyte homing and, **64:158, 188**
 - to mucosa, **64:176–177**
- mAbs**, **60:168–171**
- pregnancy and**, **64:194–195**
- properties**, **64:147**
- recruitment of leukocytes to inflammation site**, **58:395**
- shedding**, **64:171**
- upregulation**, **64:165**
- V(D)J joining, antigen receptor genes**
- 12/23 rule, **56:32**
 - agents
 - biochemically defined, **56:69–73**
 - cutting, **56:71–72**
 - genetically defined, **56:73–79**
 - hairpin-nicking activity, **56:79**
 - joining signals, binding of, **56:69–71**
 - ligation, **56:72–73**
 - molecular genetics, **56:64–69**
 - properties, **56:64**
 - RAG-1, **56:64–67**
 - RAG-2, **56:66–67**
 - replication role, **56:80–81**
 - severe combined immunodeficiency, **56:73–78**
 - terminal deoxynucleotidyl transferase, **56:68–69**
 - truncation factors, **56:79–80**
 - end donation, **56:125–126**

- endogenous substrate
 accessibility, **56:37**, 41–42
 active cell lines, **56:38**–40
 chromatin configuration, **56:37**, 41–46
 DNase I sensitivity, **56:43**–44
 Ig loci, **56:34**–37
 methylation, **56:44**–46
 T cell receptor loci, **56:34**–37
 transcription, **56:41**–43
 fidelity, **56:119**–127
 gene assembly
 for IgS, **56:28**–32
 for T cell receptors, **56:28**–32
 mechanism
 analysis of process, **56:27**–28, 132
 cleavage, **56:129**–131
 end exchange, **56:130**–131
 hypothesis, **56:127**–131
 ligation, interim, **56:131**
 modification, interim, **56:131**
 synapsis, **56:127**–129
 model systems, **58:42**–43
 cell rearrangement, **56:47**–50
 introduced substrates, **56:50**–53
 in vivo-generated functions, **56:46**–47
 nonstandard products, **56:32**–34
 hybrid joint, **56:32**–33
 open-and-shut joint, **56:32**–33
 order, origins of
 12/23 rule, **56:82**–86
 distance effects, **56:96**–98
 joining signals, **56:86**–90
 locus deletion, **56:92**–94
 mechanisms, **56:81**–82, 118–119
 orientation, **56:101**–104
 pseudo-normal joining, **56:94**–95
 rearrangement
 cis, **56:98**–99
 successive, **56:95**–96
 trans, **56:98**–101
 recombination outcomes, **56:90**–95
 recombination recognition sites, **56:86**
 specificity of end exchange, **56:104**–111
 three-dimensional signals, **56:111**–118
 V gene replacement, **56:90**–92
 pathogenesis, **56:119**–127
 recombinant structure
 crossover site location, **56:53**–54
 germline joining, **56:63**–64
 homology, **56:60**–62
 junctional inserts, **56:54**–58
 N regions, **56:54**–55
 oligonucleotide capture, **56:62**–63
 P nucleotides, **56:56**–58
 terminal deoxynucleotidyl transferase, **56:54**–56
 truncation, **56:57**–60
 recombination
 cryptic site, **56:121**–127
 interchromosomal, **56:120**–121
 outcomes, **56:90**–95
 targets, **56:32**
 terminology, **56:34**
 VDJ joints, isolated from chicken thymus, diversity, **50:111**
 V(D)J junctions
 encoding CDR3 region, **58:125**
 and generation of TCR gene diversity, **58:116**–117
 V(D)J rearrangement, **60:38**, 268
 precursor B cells, **63:16**–19
 rabbit
 antibody repertoire development, **56:195**–198
 appendix germinal centers, **56:204**
 bone marrow, **56:184**
 GALT model, **56:206**–209
 mechanism, **56:179**
 organization, **56:190**–192
 somatic conversion, **56:198**–202
 somatic mutation, **56:201**, 203
 V(D)J recombinase, **64:46**
 V(D)J recombination, **61:123**, 285, 287–291; **62:1**
 12/23 Rule, **64:45**, 53, 59
 assay, **61:124**
 biochemistry, **64:51**
 biological consequences, **64:59**–61
 cleavage at an RSS, **64:51**–53
 coupled cleavage, **64:53**–54
 hairpin formation, **64:56**–59
 signal sequence recognition, **64:54**–56
 broken DNA molecules, **64:43**–45
 cleavage, **64:52**–54
 cleavage model, **58:38**–39
 coding joints, **64:40**, 41–42
 and DNA repair, and Ku autoantigen, **58:55**–60
 DSB repair, **58:29**–85; **64:49**–51
 cell cycle regulation, **58:70**–74

- V(D)J recombination (*continued*)
 hybrid joints, **64:42**
 initiation, **58:30–32**
 joining model, **58:42–43**
 mechanism, **61:109–110**
 antigen receptor loci, **61:304–306**
 DNA double-strand breaks, **61:298–303**
 lymphocyte-specific proteins, **61:291–298**
 RAG-1 and RAG-2, **61:291–296**
 signal binding proteins, **61:303–304**
 TdT, **61:296–298**
 open-and-shut joints, **64:42**
 pathology, **61:285–286, 316–317**
 ataxia telangiectasia, **61:191, 315**
 Bloom's syndrome, **61:316**
 DNA breakage syndromes, **61:314–316**
 Fanconi's anemia, **61:315**
 severe combined immunodeficiency, **61:306–311**
 xeroderma pigmentosum, **61:315–316**
 X-linked agammaglobulinemia, **61:312–314**
 properties, **64:39–42**
 RAG proteins, **64:45–48**
 relevance of joining activity, **58:45**
 SCID defect and, **64:49–50**
 signal joints, **64:40, 41**
 signal sequence recognition, **64:54–56**
 stages, **64:40–41**
 and switch recombination, comparison, **54:235–237**
 and TCR gene expression, **58:116–117**
 terminal deoxynucleotidyl transferase, **64:51**
 V(D)J region
 B cells, shared with germinal centers, **60:272**
 hypermutation in germinal centers, **60:278–280, 284**
 VDRL antigen, aPL antibodies and, **49:227–228, 234–238**
 Vectors
 episomal, **61:124–128**
 viral, and gene transfer techniques, **58:422–425**
 VEGF, **64:319–320**
 Veiled cells, **59:60**
 IBD and, **42:296**
 Venous sinuses, B cell formation and, **41:187, 214, 236**
 Venous thrombosis, CD59-negative platelets in, **60:91–92**
 Venules, endothelial, L-selectin association with lymphocyte adhesion, **62:257**
 Vesicles
 antigen-presenting cells and, **47:68, 87, 89**
 HIV infection and, **47:393, 396**
 NK cells and, **47:214, 216, 217, 221**
 Vesicular stomatitis virus, **41:142, 148, 153, 154, 163, 168**
 NK cells and, **47:286, 287**
 VF1-12P pseudogene, **62:23**
 V gene, **43:252, 255**
 V_β gene, **43:268–269**
 V gene replacement
 IgH locus, **64:54**
 outcome of V(D)J recombination, **56:90–92**
 V gene segment, bursar lymphocyte, **48:52–54**
 ψ-V gene segment
 diversification, **48:58–60**
 diversification sequences, **48:54–56**
 gene conversion, **48:58**
 VL, recombination event, **48:56**
 V gene usage
 biases in CD5 B cells, **55:319–320**
 clonal expansions, **55:322–323**
 human populations, **55:329**
 neoplasias, **55:322–323**
 normal cells, **55:320–322**
 chicken B cell
 diversification, **57:358–361**
 recombination models, **57:362–365**
 V_H1, in rabbit B lymphopoiesis, **56:184**
 V_H2 family, **62:13**
 V_H3 family, **62:18**
 V_H4 family, **62:13, 25**
 V_H5 family, **62:13, 17**
 V_H6, **60:47**
 V_H6 family, **62:13, 17**
 V_H7 family, **62:14, 15**
 V_H genes, *see* Heavy chain variable region, genes
 V_H locus
 evolution, **62:19–26**
 physical mapping, **62:4–7**
 polymorphisms, **62:8–9**
 repetitive sequences, **62:20, 22**
 V_H pseudogenes, **62:16–17**

- V_H segments, **62:1–2**, 7–8, 12–19
 germline, **62:1–26**
- vif* gene, HIV-1 infectivity, **63:106**
- Vinca alkaloids, in cancer treatment, mAb
 conjugation procedure, **56:340–342**
 vinblastine, **56:341–342**
 vincristine, **56:340–341**
 vindesine, **56:340–341**
- VIP, *see* Vasoactive intestinal peptide
- Viral antigens, **62:239–240**
 antigen-presenting cells and, **47:47**, 80, 81
 synthetic T and B cell sites and,
45:216–228, 240, 241
- Viral cytokine receptors, **63:308–310**
- Viral determinants, pathogenesis of HIV,
 analysis using SIV molecular clones,
52:448–451
- Viral diseases, **60:128–136**
 multiple antigen peptides as immunogens,
60:128–134
- Viral genomes, viruses used as animal models
 for AIDS, organization, **52:429**
- Viral infection
 and autoimmune diabetes, **51:308**
 and bacterial, induction of NF-κB, **58:6**
 cachectin and, **42:220**, 226
 cytotoxic T cells and
 adenovirus, **65:279**
 cytomegalovirus, **65:278–279**
 Epstein–Barr virus, **65:279**
 herpes simplex virus, **65:279**
 HIV, **65:280–322**
 influenza virus, **65:278**
 IFN-γ and, **62:77**, 95
 IL-6 and, **54:37**
 and lymphopenia, **58:233–237**
 MG and, **42:272**
 NF-IL6 in, **65:9–11**
 NK cells and, **42:182**, 183
 plasmacytogenesis-induced, **64:230–232**
 pneumonia, **59:406–409**
 responses in MHC class I-deficient mouse,
55:406
 influenza virus, **55:406–407**
 lymphocytic choriomeningitis virus,
55:409–412
 Sendai virus, **55:408–409**
 Theiler's virus, **55:409**
 vaccinia virus, **55:407–408**
- switch from latency to productivity, role of
 NF-κB, **58:16–17**
- X-linked agammaglobulinemia, **59:145**, 150
- Viral inhibition
 CD8⁺ cells, **66:273–275**, 281
 cell-mediated effects, **66:286–287**
 cytolysis, **66:283–286**, 287–291
 HIV-1-specific lytic activity, **66:275–277**,
 283–286
 mechanism, **66:280–291**
 chemokines and, **66:281–282**
 IL-16, **66:283**
- Viral replication, HIV, CD8⁺ and,
65:281–282
- Viral subterfuges, **52:74–76**
- Viral vectors, and gene transfer techniques,
58:422–425
- Virulence factors, **63:308–309**
- Virus-dependent cellular cytotoxicity, NK
 cells and, **47:286**
- Viruses, *see also specific viruses*
 adoptive T cell therapy of tumors and,
49:282–283, 323, 334
 mechanisms, **49:311**, 313–314, 318
 animal models for AIDS, relationships
 among, **52:426–430**
 antibodies, from combinatorial libraries
 cytomegalovirus, human, **57:220–221**
 hepatitis B virus, **57:220**
 herpes simplex virus type 1, **57:222–227**
 herpes simplex virus type 2, **57:222–227**
 HIV-1, **57:211–217**
 measles, **57:227–228**
 respiratory syncytial virus, **57:217–219**
 varicella Zoster virus, **57:221–222**
 autoimmune thyroiditis and
 cellular immune responses, **46:297**, 298
 experimental models, **46:274–276**
 genetic control, **46:283**
 autoreactive T cells and, **45:431**
 avian leukosis, in chicken B cell DT40
 induction, **57:361–362**
 B cell formation and, **41:209**
 complement receptor 2 and, **46:210**
 CTL and, **41:135**, 142, 153, 154, 165
 cytotoxicity and, **41:269**, 270
 helper T cell cytokines and, **46:127**, 129,
 133
 and host, role in life cycle, **54:107–108**

- Viruses (*continued*)**
- humoral immune response and, **45:24**, 78, 82, 83
 - hybrid resistance and, **41:375**, 378, 390, 400, 402, 403
 - Ig heavy-chain variable region genes and, **49:56**
 - infectious
 - exogenous superantigens produced by, **54:115–117**
 - production of exogenous superantigens, **54:115–117**
 - maps of Ig-like loci and, **46:1**
 - NF- κ B, strategies to control, **65:127–128**
 - regulators of complement activation and, **45:395**, 409
 - superantigens of, **65:208**
 - surface antigens of human leukocytes and, **49:80**, 118
 - synthetic T and B cell sites and, **45:195**, 219, 260, 262
 - T cell receptor and
 - alloreactivity, **45:156**
 - antigen processing, **45:119**, 125, 127
 - experimental systems, **45:131**, 137
 - structure-function relationships, **45:141**
 - T cell subsets and, **41:54**, 56, 57, 74–76, 101
 - tumor antigen induction by, **57:290–293**
 - Epstein-Barr virus, **57:292**
 - papillomavirus, **57:292–293**
 - virus-induced immunosuppression, **45:335**, 336, 368, 369
 - HIV, **45:355–363**
 - history of infection, **45:352–355**
 - immune response, **45:363–368**
 - virology, **45:347–352**
 - measles, **45:338–347**
 - virology, **45:336–338**
 - Vitamin A
 - binding to sex steroids, **58:229**
 - proautoimmune potential, **58:278**
 - Vitamin D₃, binding to sex steroids, **58:229**
 - Vitamin E
 - action of, **65:122**, 123
 - inhibition of HIV lymphocyte death, **58:279**
 - Vitronectin, **61:204**, 247–249
 - leukocyte integrins and, **46:153**, 157
- V-J joining**, **48:46–47**
 - circular episome, **48:47**
 - diversity, **48:47**
 - random nature, **48:47**
- V-J jointing**, embryogenesis, **48:49**, 50
- V_k promoters**, **43:238**
- v-kit**, characteristics, **55:3–6**
- VLA-4**, **62:258**, **64:154**; **65:359**
 - multiple myeloma, **64:246–248**
- VLA-5**, multiple myeloma, **64:248**
- VLA-6**, multiple myeloma, **64:248**
- VLA antigens**
 - leukocyte integrins and, **46:153**, 157
 - surface antigens of human leukocytes and, **49:83–85**, 88
- VL gene**, diversification sequences, **48:54–56**
- VL gene segment**, **48:41**
 - diversification, **48:56–60**
 - gene conversion, **48:56–60**
 - sequence substitution position, **48:56**, 57
 - sequence substitution size, **48:56**, 57
 - ψ -V gene, recombination event, **48:56**
- VL pseudogene 5**, sequence comparison, **48:55**
- VL pseudogene 7**, sequence comparison, **48:55**
- V_L-V_H interface**, **43:103**, 110, 112
 - CDR, **43:125**
- V_L-V_L pairing**, **43:112**
- von Willebrand factor**, leukocyte integrins and, **46:158**, 159
- vpr**, HIV-1 infectivity, **63:86–87**
- V_{preB}**, **63:7–8**, 29
 - expression, **63:5–15**
 - gene regulation, **63:4–5**
 - μ heavy chains, **63:2–3**
 - structure, **63:4**
- vpu**, HIV-1 infectivity, **63:87**, 106
- V region**, **62:1**
- V region genes**, T cell receptors and, **38:15**, 16
- V-rel oncogene**, **48:58**
- V β rule**, modifications, **54:123–124**
 - specificities of some exogenous bacterial superantigens, **54:109**
- V β selective elements**, *see also* T cell receptor, V β selective elements
- cellular basis of action**, **50:29–32**
 - responding cells, **50:29–31**
 - CD4 T cells, **50:30**
 - CD8 T cells, **50:30–31**

stimulating cells, **50**:31–32
 B cells, **50**:31
 dendritic cells, **50**:31
 transfer of Vbse among cells, **50**:32
 general nomenclature, **50**:13–14
in vitro suppression induced by, **50**:37
 murine, not stimulating primary T cell response, **50**:11–13
 null alleles, **50**:14
 role in T cell development and function, **50**:32–37
 in intrathymic development, **50**:33
 negative selection, **50**:34–35
 positive selection, **50**:34
 in periphery, **50**:35–37
 clonal inactivation by V β selective elements, **50**:35–36
 enhancement of responses to antigen by V β selective elements, **50**:36–37
 suppression induced by V β selective elements *in vitro*, **50**:37

W

Waldenström's macroglobulinemia, **64**:219–220
 aPL antibodies and, **49**:227–228
 Ig heavy-chain variable region genes and, **49**:40
 W alleles, and c-kit alleles, relationship between, **55**:6–7
 WEHI-13 cells
 chloroleukemia-derived, **39**:16
 IL-3 production, **39**:13, 15–17, 19
 rearranged allele and, **39**:15–16
 WEHI-3B, **48**:77
 WEHI-231 B cell line, growth arrest, **50**:64–65
 Weibel-Palade bodies, P-selectin stored in, **58**:356–357; **60**:166, 168
 Western blotting, HLA class II, **48**:116
 Whitlock-Witte cultures, B cell formation and, **41**:209, 216–220, 231
 Wiskott-Aldrich syndrome, **59**:159, 248
 Wnt-1 genes, **65**:154, 155
 Woodchuck, chronic hepatitis infection model, **60**:346

Worm burden
 eosinophil effect, **60**:215
 reduction, **60**:127–128
 Wound healing
 eosinophil role, **60**:228
 NO role, **60**:350–351
 WT31, CD3, **43**:161–163

X

Xanthine, genetically engineered antibody molecules and, **44**:70
 X-chromosome, mosaicism, **59**:159, 195
 Xenograft models
 HIV-1 disease, **63**:79, 89–90, 107
 hu-PBL-SCID model, **63**:89–98
 pathogenesis, **63**:90–95, 100
 pediatric infection, **63**:95
 protective immunity, **63**:95–98
 SCID-hu thy/liv model, **63**:89–90, 99–107
 hu-PBL-SCID model, **63**:79–82
 SCID-hu thy/liv model, **63**:82–83
 Xenopus laevis, antinuclear antibodies and, **44**:124
 Xenotransplantation, **61**:219, 220
 models, human-to-mouse, **50**:319
 Xeroderma pigmentosum, **61**:315
 Xid mutation, **59**:174–176
 B cells, **59**:317–318
 mitogenic signaling, **59**:328
 XLA, *see* X-linked agammaglobulinemia
 X-linked agammaglobulinemia, **59**:135–136,
 135–197, 317–318, 318; **61**:312–314
 B cell development, **59**:151–162, 317–318
 Btk, **59**:136, 169–176, 178–196
 carrier detection, **59**:194–195
 clinical manifestations, **59**:143–148
 defect of B cell development, **60**:39–42
 diagnosis, prenatal, **59**:194–195
 in females, **59**:149–150, 194
 genes
 cloning, **59**:165–169, 165–176
 mapping, **59**:163–165, 194–195
 genetic approach, **59**:162–176, 187,
 193–194
 with growth hormone deficiency, **59**:149
 and HIGM-X-1, immunodeficiency diseases,
 60:37

X-linked agammaglobulinemia (*continued*)
 history, **59**:136–142
 incidence and prognosis, **59**:142
 and SCID mice, **50**:308
 treatment, **59**:150–151

X-linked hyper-IgM syndrome, **59**:248
 CD40-L role, **61**:1, 18, 41–44, 92–93

X-linked immunodeficiency, *see also* Xid
 mutation
 diseases, **59**:248, 317
 immunology, **63**:51
 murine, Btk implicated in, **60**:41–42

X-linked lymphoproliferative disorder, NK cells and, **47**:224, 283, 299

X-linked severe combined immunodeficiency, **59**:227, 235, 246–247, 318; **61**:152
 atypical, **59**:252–253
 IL-2R γ mutations, **59**:248–250

X-ray analysis, antibody, **43**:121

X-ray crystallography, **43**:2–3, 8–11

X-ray structures, antibody-antigen interaction, **43**:73

XRCC4, V(D)J recombination, **64**:60

XRCC5, **64**:49

Y

Yeast
 antinuclear antibodies and, **44**:106
 cytotoxicity and, **41**:269, 270, 319
 genetically engineered antibody molecules and, **44**:66, 69
SPT14 gene, homolog of *PIG-A*, **60**:72

Yeast artificial chromosomes, maps of Ig-like loci and, **46**:4, 5

Yeast artificial chromosome system, **62**:3

Yeast infections, pneumonia, **59**:402–406
 YT cell line, **42**:171, 172

Z

ZAP70, **63**:138
 IL-2R coupling, **61**:160

Zinc, cell-mediated killing and, **41**:293, 294

Zinc finger, myeloid, **63**:1, 239

Zinc-finger proteins, BLIMP, **59**:342

Zonal organization, lymph follicles, **53**:185–186

Zymogens, **61**:202
 complement components and, **38**:206, 209, 235

Zymosan
 alternative pathway activation, **38**:386
 fibronectin, **38**:390
 β -glucan constituent of, **38**:362
 IBD and, **42**:309
 ingestion inhibition, **38**:381
 phagocytosis, human monocyte, **38**:392
 cellular responses, **38**:373–376, 393
 composition, **38**:363, 364
 receptor characterization, **38**:365, 366, 368–373

phagocytosis, nonelicited murine macrophages
 cellular responses, **38**:377–380
 receptor characterization, **38**:376, 377
 serum-treated, addition to eosinophils, **60**:183

CONTRIBUTOR INDEX

Boldface numerals indicate volume number

A

- Abraham, Robert T., **61**:147
Acha-Orbea, Hans, **65**:139
Adelman, D. C., **48**:161
Adolphson, Cheryl R., **39**:177
Aebischer, Iwan, **61**:341
Ahearn, Joseph M., **46**:183
Akira, Shizuo, **54**:1; **65**:1
Akkaraju, Srinivas, **59**:279
Altman, Amnon, **48**:227
Anderson, Steven J., **56**:151
Andreu, Jose Luis, **50**:147
Arend, William P., **54**:167
Arm, Jonathan P., **51**:323
Asao, Hironobu, **59**:225
Atkinson, John P., **45**:381; **61**:201
Austen, K. Frank, **39**:145

B

- Baeuerle, Patrick A., **65**:111
Baeza, Maria, **50**:237
Baggiolini, Marco, **55**:97
Baichwal, Vijay R., **65**:111
Banchereau, Jacques, **52**:125; **61**:1
Barbas III, Carlos F., **57**:191
Bartram, Claus R., **61**:285
Bassuk, Alexander G., **64**:65
Bell, Sarah E., **59**:279
Bennett, Michael, **41**:333
Bennink, Jack R., **52**:1
Benoist, Christophe, **55**:424
Bertocci, Barbara, **57**:353

- Bettelheim, Peter, **52**:333
Beutler, B., **42**:213; **49**:149
Bevan, Michael J., **59**:99
Biassoni, Roberto, **55**:341
Billiau, Alfons, **62**:61
Bodmer, Helen, **55**:424
Boehm, T., **50**:119
Brenner, Michael B., **43**:133
Brezinschek, Hans Peter, **47**:433
Bucy, R. Pat, **50**:87
Burrows, Peter D., **65**:245
Burton, Dennis R., **51**:1; **57**:191

C

- Calame, Kathryn, **43**:235
Calvo-Calle, J. Mauricio, **60**:105
Campbell, R. D., **38**:203
Capra, J. Donald, **42**:95; **49**:1
Cardell, Susanna, **55**:424
Carroll, M. C., **38**:203
Carson, Dennis A., **38**:275
Cerami, A., **42**:213
Chan, Susan, **55**:424
Charreire, Jeannine, **46**:263
Charron, Dominique, **48**:107
Chen, Chen-Lo H., **50**:87
Chen, Jianzhu, **62**:31
Cheresh, David A., **40**:323
Chesnut, Robert W., **39**:51
Chesterman, Colin N., **49**:193
Chilton, Paula M., **63**:269
Ciccone, Ermanno, **55**:341
Clark, Lisa B., **63**:43

Claverie, Jean-Michel, **45**:107
 Coffman, Robert L., **46**:111; **54**:229
 Coggeshall, K. Mark, **48**:227
 Cohen, J. John, **50**:55
 Cohn, Zanvil A., **41**:269
 Colman, P. M., **43**:99
 Cooke, Michael P., **59**:279
 Cooper, Max D., **50**:87; **65**:245
 Corbi, Angel L., **46**:149
 Cosgrove, Dominic, **55**:424
 Cotran, Ramzi S., **50**:261
 Crane, Mary A., **56**:179
 Croce, Carlo M., **38**:245
 Cuende, Eduardo, **53**:157
 Cyster, Jason G., **59**:279
 Czop, Joyce K., **38**:361

D

Dahan, Auriel, **57**:353
 Dalloul, A., **49**:149
 Debre, P., **49**:149
 DeFranco, Anthony L., **55**:221
 Delespesse, G., **49**:149
 Dewald, Beatrice, **55**:97
 DeWitt, David L., **62**:167
 Dietrich, Hermann, **47**:433
 Dinarello, Charles A., **44**:153
 Dixon, Frank J., **46**:61
 Dranoff, Glenn, **58**:417
 Dunon, Dominique, **58**:345
 Dustin, Michael L., **46**:149

E

Eaton, Suzanne, **43**:235
 Elliott, Michael J., **64**:283
 Elliott, Tim, **65**:47

F

Farries, Timothy C., **61**:201
 Fathman, C. Garrison, **56**:219
 Fauci, Anthony S., **47**:377
 Fearon, Douglas T., **46**:183
 Feldmann, Marc, **64**:283
 Fernandez-Botran, Rafael, **45**:1; **63**:269

Fink, Pamela J., **59**:99
 Finn, Olivera J., **62**:217
 Forman, James, **41**:135
 Fowlkes, B. J., **44**:207
 Foy, Teresa M., **63**:43
 Freimark, Bruce D., **38**:275
 Fujii, Yoshitaka, **42**:233
 Fuleihan, Ramsay, **60**:37

G

Galli, Stephen J., **55**:1
 Gaur, Amitabh, **56**:219
 Gearing, David P., **53**:31
 Geha, Raif S., **39**:255; **60**:37
 Geissler, Edwin N., **55**:1
 Gellert, Martin, **64**:39
 Getzoff, D. Elizabeth, **43**:1
 Geysen, H. Mario, **43**:1
 Ghosh, Sankar, **58**:1
 Gleich, Gerald J., **39**:177
 Goetzl, E. J., **48**:161
 Goetzl, Edward J., **39**:299
 Gold, Michael R., **55**:221
 Gomez, Javier, **63**:127
 Gonzalo, Jose Angel, **50**:147
 Goodnow, Christopher C., **59**:279
 Greenberg, Philip D., **49**:281
 Grey, Howard M., **39**:51
 Gutierrez-Ramos, Jose C., **50**:147

H

Hala, Karel, **47**:433
 Hansen, John A., **40**:379
 Hansen, Ted H., **64**:105
 Hardy, Richard R., **55**:297
 Hartley, Suzanne B., **59**:279
 Hathcock, Karen S., **62**:131
 Hayakawa, Kyoko, **55**:297
 Healy, James L., **59**:279
 Hedrick, Stephen M., **43**:193
 Heldin, Carl-Henrik, **55**:181
 Henderson, Robert A., **62**:217
 Hercend, Thierry, **42**:181
 Hirsch, Vanessa M., **52**:425
 Hitoshi, Yasumichi, **57**:145
 Hodes, Richard J., **62**:131

- Hofstetter, H., **49**:149
 Hogarth, P. Mark, **57**:1
 Holers, V. Michael, **45**:381
 Honjo, Tasuku, **62**:1
 Hood, Leroy, **44**:1; **46**:1
 Hooghe, Robert, **63**:377
 Hooghe-Peters, Elisabeth L., **63**:377
 Horejsi, V., **49**:75
 Hourcade, Dennis, **45**:381
 Houssiau, F., **54**:79
 Hulett, Mark D., **57**:1
 Hunkapiller, Tim, **44**:1
 Huston, Marilyn M., **38**:313
 Hyman, Robert, **54**:271

I

- Ichijo, Hidenori, **55**:181
 Igarashi, Hideya, **54**:337
 Ihle, James N., **39**:1; **60**:1
 Imboden, John B., **41**:1
 Imhof, Beat A., **58**:345
 Inoue, Norimitsu, **60**:57
 Inui, Seiji, **54**:337
 Ishii, Naoto, **59**:225
 Ishizaka, Kimishige, **47**:1
 Ivashkiv, Lionel B., **63**:337

J

- Jalkanen, Sirpa, **64**:139
 Janeway, Jr., Charles A., **50**:1
 Jelinek, Diane F., **40**:1
 Johnson, Philip R., **52**:425

K

- Kaplan, Allen P., **50**:237
 Kappler, John, **38**:1; **54**:99
 Karasuyama, Hajime, **63**:1
 Karnitz, Larry M., **61**:147
 Kawasaki, Akemi, **51**:215
 Kay, A. Barry, **60**:151
 Kelso, Anne, **48**:69
 Kelsoe, Garnett, **60**:267
 Kikutani, Hitoshi, **51**:285
 Kilcherr, E., **49**:149
- Kincade, Paul W., **41**:181; **54**:271
 Kindt, Thomas J., **52**:425
 Kinoshita, Taroh, **60**:57
 Kipps, Thomas J., **47**:117
 Kishiimoto, Tadamitsu, **54**:1; **64**:219; **65**:1
 Kishimoto, Takashi, **46**:149
 Kisielow, Pawel, **58**:87
 Klinman, Norman R., **42**:1
 Knight, Katherine L., **56**:179
 Kofer, Reinhard, **46**:61
 Kojima, Hidefumi, **60**:289
 Kondo, Motonari, **59**:225
 Kooijman, Ron, **63**:377
 Kopp, Elizabeth B., **58**:1
 Kotzin, Brian, **54**:99
 Kourilsky, Philippe, **45**:107
 Kourilsky, Philippe, **57**:281
 Kraal, Georg, **65**:347
 Krangel, Michael S., **43**:133
 Krilis, Steven A., **49**:193
 Kroemer, Guido, **47**:433; **50**:147; **53**:157; **58**:211
 Kuna, Piotr, **50**:237
 Kuwahara, Kazuhiko, **54**:337

L

- Lai, Eric, **46**:1
 Larson, Richard S., **46**:149
 Lebman, Deborah A., **54**:229
 Lee, David R., **64**:105
 Lee, Tak H., **39**:145; **51**:323
 Leiden, Jeffrey M., **64**:65
 Lerner, Richard A., **43**:1
 Lesley, Jane, **54**:271
 Leung, Donald Y. M., **54**:99
 Levin, Steven D., **56**:151
 Lewis, Susanna M., **56**:27
 Lindstrom, Jon, **42**:233
 Linton, Phyllis-Jean, **42**:1
 Lipscomb, Mary F., **59**:369
 Lipsky, Peter E., **40**:1
 Liszewski, M. Kathryn, **61**:201
 Louahed, J., **54**:79
 Lublin, Douglas M., **61**:201
 Lukacs, Nicholas W., **62**:257
 Luther, Sanjiv A., **65**:139
 Lynch, Richard G., **40**:135

Lyons, C. Richard, **59**:369
 Lyons, C. Rick, **60**:323

Mustelin, Tomas, **48**:227
 Myers, Christopher D., **45**:1

M

Ma, Yuhe, **63**:269
 MacDermott, Richard P., **42**:285
 Mach, Bernard, **61**:327
 Mackay, Charles R., **53**:217
 Maini, Ravinder N., **64**:283
 Makino, Susumu, **51**:285
 Marrack, Philippa, **38**:1
 Marrack, Philippa, **54**:99
 Martin, Paul J., **40**:379
 Martinez-A., Carlos, **50**:147; **53**:157; **63**:127
 Mathis, Diane, **55**:424
 Matsuda, Fumihiro, **62**:1
 Matsuo, Tatsuya, **54**:337
 McChesney, Michael B., **45**:335
 McCormack, Wayne T., **48**:41
 McGhee, Jerry R., **40**:153
 McGillis, Joseph P., **39**:299
 McKenzie, Ian F. C., **56**:301
 McMichael, Andrew, **65**:277
 McNeil, H. Patrick, **49**:193
 Mebius, Reina E., **65**:347
 Meek, Katheryn, **42**:95
 Melchers, Fritz, **53**:123; **63**:1
 Merkenschlager, Matthias, **55**:424
 Mestecky, Jiri, **40**:153
 Metcalf, Donald, **48**:69
 Metlay, Josua P., **47**:45
 Metzger, Henry, **43**:277
 Miceli, M. Carrie, **53**:59
 Michon, Jean, **42**:181
 Milich, David R., **45**:195
 Mingari, Maria Christina, **55**:341
 Miyazono, Kohei, **55**:181
 Moore, Jr., Francis D., **56**:267
 Moqbel, Redwan, **60**:151
 Moretta, Alessandro, **55**:341
 Moretta, Lorenzo, **55**:341
 Morgan, Edward L., **40**:61
 Morrison, Sherie L., **44**:65
 Moser, Bernhard, **55**:97
 Mosier, Donald E., **50**:303; **63**:79
 Mosmann, Tim R., **46**:111; **56**:1
 Mossalayi, D., **49**:149
 Mulligan, Richard C., **58**:417

N

Nadler, Lee Marshall, **51**:243
 Nagata, Shigekazu, **57**:129
 Nakamura, Masataka, **59**:225
 Nakamura, Toshikazu, **59**:225
 Nakata, Motomi, **51**:215
 Nardin, Elizabeth H., **60**:105
 Neiman, Paul E., **56**:467
 Nezlin, Roald, **48**:1
 Niederkorn, Jerry Y., **48**:191
 Nishimoto, Norihiro, **64**:219
 Noelle, Randolph J., **63**:43
 Nomura, Jun, **54**:337
 Nossal, G. J. V., **52**:283
 Nowell, Peter C., **38**:245
 Nussenzweig, Ruth S., **45**:283; **60**:105
 Nussenzweig, Victor, **45**:283

O

Oi, Vernon T., **44**:65
 Okumura, Ko, **51**:215
 Oldstone, Michael B. A., **45**:335
 Oliveira, Giane A., **60**:105
 Opstelten, Davina, **63**:197

P

Pardoll, Drew M., **44**:207
 Parkman, Robertson, **49**:381
 Parnes, Jane R., **44**:265; **53**:59
 Pascual, Virginia, **49**:1
 Paul, William E., **53**:1
 Payan, Donald G., **39**:299
 Perlmutter, Roger M., **56**:151
 Pietersz, Geoffrey A., **56**:301
 Plaut, Marshall, **53**:1
 Pober, Jordan S., **50**:261
 Pogue, Sarah L., **59**:279
 Porcelli, Steven A., **59**:1
 Porter, R. R., **38**:203
 Pure, Ellen, **47**:45

R

- Rabbitts, T. H., **50**:119
 Ramesh, Narayanaswamy, **60**:37
 Rathbun, Gary, **42**:95
 Rathmell, Jeffrey C., **59**:279
 Raulet, David H., **55**:381
 Rebollo, Angelita, **63**:127
 Reddigari, Sesha, **50**:237
 Reisfeld, Ralph A., **40**:323
 Reith, Walter, **61**:327
 Renauld, J.-C., **54**:79
 Reynaud, Claude-Agnes, **57**:353
 Rice, David E., **59**:369
 Rich, Robert R., **38**:313
 Ritz, Jerome, **42**:181
 Rochlitz, Christoph, **57**:281
 Rodgers, John R., **38**:313
 Rolink, Antonius, **53**:123; **63**:1
 Rooney, Isabelle A., **61**:201
 Rosen, Steven D., **44**:313
 Rosenberg, Zeda F., **47**:377
 Roth, Claude, **57**:281
 Rothenberg, Ellen V., **51**:85
 Rothman, Paul, **54**:229
 Rousset, Francoise, **52**:125
 Rowland, April, **56**:301
 Rowland-Jones, Sarah, **65**:277

- Shokat, Kevan P., **59**:279
 Sideras, Paschal, **59**:135
 Sim, Gek-Kee, **58**:297
 Singer, Paul A., **46**:61
 Sklar, Larry A., **39**:95
 Smith III, Roger, **38**:313
 Smith, C. I. Edvard, **59**:135
 Smith, Kendall A., **42**:165
 Smith, William L., **62**:167
 Smyth, Mark J., **56**:301
 Sprent, Jonathan, **41**:39
 Springer, Timothy A., **46**:149
 Sreedharan, S. P., **48**:161
 Stadler, Beda M., **61**:341
 Stanley, John R., **53**:291
 Staunton, Donald E., **46**:149
 Stavnezer, Janet, **61**:79
 Steimle, Viktor, **61**:327
 Steinman, Lawrence, **49**:357
 Steinman, Ralph M., **47**:45
 Stenson, William F., **42**:285
 Storb, Rainer, **40**:379
 Strominger, Jack L., **43**:133
 Suematsu, Sachiko, **64**:219
 Sugamura, Kazuo, **59**:225
 Suter, U., **49**:149

T**S**

- Sakaguchi, Nobuo, **54**:337
 Salmi, Marko, **64**:139
 Sanders, Virginia M., **45**:1
 Sanz, Inaki, **42**:95
 Sarfati, M., **49**:149
 Sawasdikosol, Sansana, **52**:425
 Schlossman, Stuart F., **42**:181
 Schmidt, Reinhold E., **42**:181
 Schriever, Folke, **51**:243
 Schuyler, Mark R., **59**:369
 Schwartz, Ronald H., **38**:31
 Schwarz, Klaus, **61**:285
 Scott, David W., **54**:393
 Seder, Robert A., **53**:1
 Shelton, Diane, **42**:233
 Shimbara, Naoki, **64**:1
 Shinkai, Yoichi, **51**:215
 Shinohara, Nobukata, **60**:289

- Taga, Tetsuya, **54**:1
 Tainer, John A., **43**:1
 Takaki, Satoshi, **57**:145
 Takatsu, Kiyoshi, **57**:145
 Takayama, Hajime, **60**:289
 Takeda, Junji, **60**:57
 Tan, Eng M., **44**:93
 Tan, Rusung, **65**:277
 Tanahashi, Nobuyuki, **64**:1
 Tanaka, Keiji, **64**:1
 Tanaka, Nobuyuki, **52**:263
 Tanaka, Nobuyuki, **59**:225
 Taniguchi, Tadatsugu, **52**:263
 ten Dijke, Peter, **55**:181
 Theofilopoulos, Argyrios N., **46**:61
 Thoman, Marilyn L., **46**:221
 Thomas, E. Donnall, **40**:379
 Thompson, Craig B., **48**:41; **50**:87
 Trinchieri, Giorgio, **47**:187

Tsurumi, Chizuko, **64:1**
 Tucker, Philip, **42:95**

U

Ulevitch, Richard J., **53:267**
 Uyttenhove, C., **54:79**

V

Valent, Peter, **52:333**
 van Kooten, Cees, **61:1**
 Van Snick, J., **54:79**
 Vink, A., **54:79**
 Vitetta, Ellen S., **45:1**
 von Boehmer, Harald, **58:87**

W

Waldmann, Thomas A., **40:247**
 Ward, Peter A., **62:257**
 Wardlaw, Andrew J., **60:151**
 Weaver, David T., **58:29**

Webb, Susan R., **41:39**
 Weigle, William O., **40:61; 46:221**
 Weill, Jean-Claude, **57:353**
 Weinstein, Yacob, **39:1**
 Weiss, Arthur, **41:1**
 Wick, George, **47:433**
 Wilkes, David, **59:369**
 Wilson, Richard K., **46:1**
 Winchester, Robert, **56:389**
 Wolf, Hugo, **47:433**
 Woody, James N., **64:283**
 Woof, Jenny M., **51:1**


Y

Yagita, Hideo, **51:215**
 Yednock, Ted A., **44:313**
 Yewdell, Jonathan W., **52:1**
 Yokota, Kin-ya, **64:1**
 Young, John Ding-E, **41:269**

Z

Zauderer, Maurice, **45:417**
 Zsebo, Krisztina M., **55:1**

ISBN 0-12-022467-4



9 780120 224678