

# Discrete Dynamical Systems


with an Introduction to Discrete Optimization Problems

Arild Wikan



Arild Wikan

# Discrete Dynamical Systems with an Introduction to Discrete Optimization Problems



Discrete Dynamical Systems with an Introduction to Discrete Optimization Problems

1<sup>st</sup> edition

© 2013 Arild Wikan & [bookboon.com](http://bookboon.com)

ISBN 978-87-403-0327-8


# Contents

<b>Acknowledgements</b>	<b>6</b>
<b>Introduction</b>	<b>7</b>
<b>Part 1 One-dimensional maps <math>f : \mathbb{R} \rightarrow \mathbb{R} \quad x \rightarrow f(x)</math></b>	<b>11</b>
1.1 Preliminaries and definitions	12
1.2 One-parameter family of maps	16
1.3 Fixed points and periodic points of the quadratic map	19
1.4 Stability	24
1.5 Bifurcations	30
1.6 The flip bifurcation sequence	35
1.7 Period 3 implies chaos. Sarkovskii's theorem	38
1.8 The Schwarzian derivative	42
1.9 Symbolic dynamics I	45
1.10 Symbolic dynamics II	50
1.11 Chaos	60
1.12 Superstable orbits and a summary of the dynamics of the quadratic map	64

## Free eBook on Learning & Development

By the Chief Learning Officer of McKinsey

[Download Now](#)





	<b>Part II <math>n</math>-dimensional maps <math>f : \mathbb{R}^n \rightarrow \mathbb{R}^n \quad \mathbf{x} \rightarrow f(\mathbf{x})</math></b>	<b>68</b>
2.1	Higher order difference equations	69
2.2	Systems of linear difference equations. Linear maps from $\mathbb{R}^n$ to $\mathbb{R}^n$	86
2.3	The Leslie matrix	97
2.4	Fixed points and stability of nonlinear systems	106
2.5	The Hopf bifurcation	115
2.6	Symbolic dynamics III (The Horseshoe map)	132
2.7	The center manifold theorem	138
2.8	Beyond the Hopf bifurcation, possible routes to chaos	147
2.9	Difference-Delay equations	173
	<b>Part III Discrete Time Optimization Problems</b>	<b>187</b>
3.1	The fundamental equation of discrete dynamic programming	188
3.2	The maximum principle (Discrete version)	198
3.3	Infinite horizon problems	206
3.4	Discrete stochastic optimization problems	218
	<b>Appendix (Parameter Estimation)</b>	<b>234</b>
	<b>References</b>	<b>247</b>

www.sylvania.com

**We do not reinvent  
the wheel we reinvent  
light.**

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

**OSRAM  
SYLVANIA**



# Acknowledgements

My special thanks goes to Einar Mjølhus who introduced me to the fascinating world of discrete dynamical systems. Responses from B. Davidsen, A. Eide, O. Flaaten, A. Seierstad, A. Strøm, and K. Sydsæter are also gratefully acknowledged.

I also want to thank Liv Larssen for her excellent typing of this manuscript and Ø. Kristensen for his assistance regarding the figures.

Financial support from Harstad University College is also gratefully acknowledged.

Finally I would like to thank my family for bearing over with me throughout the writing process.

Autumn 2012

Arild Wikan

# Introduction

In most textbooks on dynamical systems, focus is on continuous systems which leads to the study of differential equations rather than on discrete systems which results in the study of maps or difference equations. This fact has in many respects an obvious historical explanation. If we go back to the time of Newton (1642–1727), Leibniz (1646–1716), and some years later to Euler (1709–1783), many important aspects of the theory of continuous dynamical systems were established. Newton was interested in problems within celestial mechanics, in particular problems concerning the computations of planet motions, and the study of such kind of problems lead to differential equations which he solved mainly by use of power series method. Leibniz discovered in 1691 how to solve separable differential equations, and three years later he established a solution method for first order linear equations as well. Euler (1739) showed how to solve higher order differential equations with constant coefficients. Later on, in fields such as fluid mechanics, relativity, quantum mechanics, but also in other scientific branches like ecology, biology and economy, it became clear that important problems could be formulated in an elegant and often simple way in terms of differential equations. However, to solve these (nonlinear) equations proved to be very difficult. Therefore, throughout the years, a rich and vast literature on continuous dynamical systems has been established.

Regarding discrete systems (maps or difference equations), the pioneers made important contributions here too. Indeed, Newton designed a numerical algorithm, known as Newton's method, for computing zeros of equations and Euler developed a discrete method, Euler's method (which often is referred to as a first order Runge–Kutta method), which was applied in order to solve differential equations numerically.

Modern dynamical system theory (both continuous and discrete) is not that old. It began in the last part of the nineteenth century, mainly due to the work of Poincaré who (among lots of other topics) introduced the Poincaré return map as a powerful tool in his qualitative approach towards the study of differential equations. Later in the twentieth century Birkhoff (1927) too made important contributions to the field by showing how discrete maps could be used in order to understand the global behaviour of differential equation systems. Julia considered complex maps and the outstanding works of Russian mathematicians like Andronov, Liapunov and Arnold really developed the modern theory further.

In this book we shall concentrate on discrete dynamical systems. There are several reasons for such a choice. As already mentioned, there is a rich and vast literature on continuous dynamical systems, but there are only a few textbooks which treat discrete systems exclusively.

Secondly, while many textbooks take examples from physics, we shall here illustrate large parts of the theory we present by problems from biology and ecology, in fact, most examples are taken from problems which arise in population dynamical studies. Regarding such studies, there is a growing understanding in biological and ecological communities that species which exhibit birth pulse fertilities (species that reproduce in a short time interval during a year) should be modelled by use of difference equations (or maps) rather than differential equations, cf. the discussion in Cushing (1998) and Caswell (2001). Therefore, such studies provide an excellent ground for illuminating important ideas and concepts from discrete dynamical system theory.

Another important aspect which we also want to stress is the fact that in case of “low-dimensional problems” (problems with only one or two state variables) the possible dynamics found in nonlinear discrete models is much richer than in their continuous counterparts. Indeed, let us briefly illustrate this aspect through the following example:

Let  $N = N(t)$  be the size of a population at time  $t$ . In 1837 Verhulst suggested that the change of  $N$  could be described by the differential equation (later known as the Verhulst equation)

$$\dot{N} = rN \left( 1 - \frac{N}{K} \right) \quad (I1)$$

where the parameter  $r$  ( $r > 0$ ) is the intrinsic growth rate at low densities and  $K$  is the carrying capacity. Now, define  $x = N/K$ . Then (I1) may be rewritten as

$$\dot{x} = rx(1 - x) \quad (I2)$$

which (as (I1) too) is nothing but a separable equation. Hence, it is straightforward to show that its solution becomes

$$x(t) = \frac{1}{1 - \frac{x_0 - 1}{x_0} e^{-rt}} \quad (I3)$$

where we also have used the initial condition  $x(0) = x_0 > 0$ . From (I3) we conclude that  $x(t) \rightarrow 1$  as  $t \rightarrow \infty$  which means that  $x^* = 1$  is a stable fixed point of (I2). Moreover, as is true for (I1) we have proved that the population  $N$  will settle at its carrying capacity  $K$ .

Next, let us turn to the discrete analogue of (I2). From (I2) it follows that

$$\frac{x_{t+1} - x_t}{\Delta t} \approx rx_t(1 - x_t) \quad (I4)$$

which implies

$$x_{t+1} = x_t + r\Delta t x_t - r\Delta t x_t^2 = (1 + r\Delta t)x_t \left(1 - \frac{r\Delta t}{1 + r\Delta t} x_t\right) \quad (\text{I5})$$

and through the definition  $y = r\Delta t(1 + r\Delta t)^{-1}x$  we easily obtain

$$y_{t+1} = \mu y_t(1 - y_t) \quad (\text{I6})$$

where  $\mu = 1 + r\Delta t$ .

The “sweet and innocent-looking” equation (I6) is often referred to as the quadratic or the logistic map. Its possible dynamical outcomes were presented by Sir Robert May in an influential review article called “Simple mathematical models with very complicated dynamics” in *Nature* (1976). There, he showed, depending on the value of the parameter  $\mu$ , that the asymptotic behaviour of (I6) could be a stable fixed point (just as in (I2)), but also periodic solutions of both even and odd periods as well as chaotic behaviour. Thus the dynamic outcome of (I6) is richer and much more complicated than the behaviour of the continuous counterpart (I2).

Hence, instead of considering continuous systems where the number of state variables is at least 3 (the minimum number of state variables for a continuous system to exhibit chaotic behaviour), we find it much more convenient to concentrate on discrete systems so that we can introduce and discuss important definitions, ideas and concepts without having to consider more complicated (continuous) models than necessary.

—

The book is divided into three parts. In *Part I*, we will develop the necessary qualitative theory which will enable us to understand the complex nature of one-dimensional maps. Definitions, theorems and proofs shall be given in a general context, but most examples are taken from biology and ecology. Equation (I6) will on many occasions serve as a running example throughout the text. In *Part II* the theory will be extended to  $n$ -dimensional maps (or systems of difference equations). A couple of sections where we present various solution methods of higher order and systems of linear difference equations are also included. As in Part I, the theory will be illustrated and exemplified by use of population models from biology and ecology. In particular, Leslie matrix models and their relatives, stage structured models shall frequently serve as examples. In *Part III* we focus on various aspects of discrete time optimization problems which include both dynamic programming as well as discrete time control theory. Solution methods of finite and infinite horizon problems are presented and the problems at hand may be of both deterministic and stochastic nature. We have also included an Appendix where we briefly discuss how parameters in models like those presented in Part I and Part II may be estimated by use of time series. The motivation for this is that several of our population models may or have been applied on concrete species which brings forward the question of estimation. Hence, instead of referring to the literature we supply the necessary material here.

—



Finally, we want to repeat and emphasize that although we have used lots of examples and problems taken from biology and ecology this is a Mathematics text so in order to be well prepared, the potential reader should have a background from a calculus course and also a prerequisite of topics from linear algebra, especially some knowledge of real and complex eigenvalues and associated eigenvectors. Regarding section 2.5 where the Hopf bifurcation is presented, the reader would also benefit from a somewhat deeper comprehension of complex numbers. This is all that is necessary really in order to establish the machinery we need in order to study the fascinating behaviour of nonlinear maps.



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more

# Part 1

## One-dimensional maps

$$f : \mathbb{R} \rightarrow \mathbb{R} \quad x \rightarrow f(x)$$

## 1.1 Preliminaries and definitions

Let  $I \subset \mathbb{R}$  and  $J \subset \mathbb{R}$  be two intervals. If  $f$  is a map from  $I$  to  $J$  we will express that as  $f : I \rightarrow J$ ,  $x \rightarrow f(x)$ . Sometimes we will also express the map as a difference equation  $x_{t+1} = f(x_t)$ . If the map  $f$  depends on a parameter  $u$  we write  $f_u(x)$  and say that  $f$  is a one-parameter family of maps.

For a given  $x_0$ , successive iterations of map  $f$  (or the difference equation  $x_{t+1} = f(x_t)$ ) give:  $x_1 = f(x_0)$ ,  $x_2 = f(x_1) = f(f(x_0)) = f^2(x_0)$ ,  $x_3 = f(x_2) = f(f^2(x_0)) = f^3(x_0) \dots$ , so after  $n$  iterations  $x_{n+1} = f^n(x_0)$ . Thus, the orbit of a map is a sequence of points  $\{x_0, f(x_0), \dots, f^n(x_0)\}$  which we for simplicity will write as  $\{f^n(x_0)\}$ . This is in contrast to the continuous case (differential equation) where the orbit is a curve.

As is true for differential equations it is a well-known fact that most classes of equations may not be solved explicitly. The same is certainly true for maps. However, the map  $x \rightarrow f(x) = ax + b$  where  $a$  and  $b$  are constants is solvable.

**Theorem 1.1.1.** The difference equation

$$x_{t+1} = ax_t + b \tag{1.1.1}$$

has the solution

$$x_t = a^t \left( x_0 - \frac{b}{1-a} \right) + \frac{b}{1-a}, \quad a \neq 1 \tag{1.1.2a}$$

$$x_t = x_0 + bt, \quad a = 1 \tag{1.1.2b}$$

where  $x_0$  is the initial value.

**Proof.** From (1.1.1) we have  $x_1 = ax_0 + b \Rightarrow x_2 = ax_1 + b = a(ax_0 + b) + b = a^2x_0 + (a+1)b \Rightarrow x_3 = ax_2 + b = \dots = a^3x_0 + (a^2 + a + 1)b$ . Thus assume  $x_k = a^kx_0 + (a^{k-1} + a^{k-2} + \dots + a + 1)b$ . Then by induction:  $x_{k+1} = ax_k + b = a[a^kx_0 + (a^{k-1} + a^{k-2} + \dots + a + 1)b] + b = a^{k+1}x_0 + (a^k + a^{k-1} + \dots + a + 1)b$ . If  $a \neq 1$ :  $1 + a + \dots + a^k = (1 - a^{t+1})(1 - a)^{-1}$  so the solution becomes

$$x_t = a^t x_0 + \frac{1 - a^{t+1}}{1 - a} b = a^t \left( x_0 - \frac{b}{1-a} \right) + \frac{b}{1-a}$$

If  $a = 1$ :  $1 + a + \dots + a^{t-1} = t \cdot 1 = t$

$$x_t = x_0 + bt$$

□

Regarding the asymptotic behaviour (long-time behaviour) we have from Theorem 1.1.1: If  $|a| < 1$   $\lim_{t \rightarrow \infty} x_t = b/(1 - a)$ . (If  $x_0 = b/(1 - a)$  this is true for any  $a \neq 1$ .) If  $a > 1$  and  $x_0 \neq b/(1 - a)$  the result is exponential growth or decay, and finally, if  $a < -1$  divergent oscillations is the outcome.

If  $b = 0$ , (1.1.1) becomes

$$x_{t+1} = ax_t \tag{1.1.2}$$

which we will refer to as the linear difference equation. The solution is

$$x_t = a^t x_0 \tag{1.1.3}$$

Hence, whenever  $|a| < 1$ ,  $x_t \rightarrow 0$  asymptotically (as a convergent oscillation if  $-1 < a < 0$ ).  $a > 1$  or  $a < -1$  gives exponential growth or divergent oscillations respectively.

**Exercise 1.1.1.** Solve and describe the asymptotic behaviour of the equations:

a)  $x_{t+1} = 2x_t + 4$ ,  $x_0 = 1$ ,

b)  $3x_{t+1} = x_t + 2$ ,  $x_0 = 2$ . □

**Exercise 1.1.2.** Denote  $x^* = b/(1 - a)$  where  $a \neq 1$  and describe the asymptotic behaviour of equation (1.1.1) in the following cases:

a)  $0 < a < 1$  and  $x_0 < x^*$ ,

b)  $-1 < a < 0$  and  $x_0 < x^*$ ,

c)  $a > 1$  and  $x_0 > x^*$ . □

Equations of the form  $x_{t+1} + ax_t = f(t)$ , for example  $x_{t+1} - 2x_t = t^2 + 1$ , may be regarded as special cases of the more general situation

$$x_{t+n} + a_1 x_{t+n-1} + a_2 x_{t+n-2} + \dots + a_n x_t = f(t), \quad n = 1, 2, \dots$$

Such equations are treated in Section 2.1 (cf. Theorem 2.1.6, see also examples following equation (2.1.6) and Exercise 2.1.5).

—

When the map  $x \rightarrow f(x)$  is nonlinear (for example  $x \rightarrow 2x(1 - x)$ ) there are no solution methods so information of the asymptotic behaviour must be obtained by use of qualitative theory.

**Definition 1.1.1.** A fixed point  $x^*$  for the map  $x \rightarrow f(x)$  is a point which satisfies the equation  $x^* = f(x^*)$ . □

Fixed points are of great importance to us and the following theorem will be very useful.

**Theorem 1.1.2.**

- a) Let  $I = [a, b]$  be an interval and let  $f : I \rightarrow I$  be continuous. Then  $f$  has at least one fixed point in  $I$ .
- b) Suppose in addition that  $|f'(x)| < 1$  for all  $x \in I$ . Then there exists a unique fixed point for  $f$  in  $I$ , and moreover

$$|f(x) - f(y)| < |x - y|$$


□

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more



**Proof.**

a) Define  $g(x) = f(x) - x$ . Clearly,  $g(x)$  too is continuous. Suppose  $f(a) > a$  and  $f(b) < b$ . (If  $f(a) = a$  or  $f(b) = b$  then  $a$  and  $b$  are fixed points.) Then  $g(a) > 0$  and  $g(b) < 0$  so the intermediate value theorem from elementary calculus directly gives the existence of  $c$  such that  $g(c) = 0$ . Hence,  $c = f(c)$ .

b) From a) we know that there is at least one fixed point. Suppose that both  $x$  and  $y$  ( $x \neq y$ ) are fixed points. Then according to the mean value theorem from elementary calculus there exists  $c$  between  $x$  and  $y$  such that  $f(x) - f(y) = f'(c)(x - y)$ . This yields (since  $x = f(x)$ ,  $y = f(y)$ ) that

$$f'(c) = \frac{f(x) - f(y)}{x - y} = 1$$

This contradicts  $|f'(x)| < 1$ . Thus  $x = y$  so the fixed point is unique. Further from the mean value theorem:

$$|f(x) - f(y)| = |f'(c)| |x - y| < |x - y|$$

□

**Definition 1.1.2.** Consider the map  $x \rightarrow f(x)$ . The point  $p$  is called a periodic point of period  $n$  if  $p = f^n(p)$ . The least  $n > 0$  for which  $p = f^n(p)$  is referred to as the prime period of  $p$ .

Note that a fixed point may be regarded as a periodic point of period one. □

**Exercise 1.1.3.** Find the fixed points and the period two points of  $f(x) = x^3$ . □

**Definition 1.1.3.** If  $f'(c) = 0$ ,  $c$  is called a critical point of  $f$ .  $c$  is nondegenerate if  $f''(c) \neq 0$ , degenerate if  $f''(c) = 0$ . □

The derivative of the  $n$ -th iterate  $f^n(x)$  is easy to compute by use of the chain rule. Observe that  $f^n(x) = f(f^{n-1}(x))$ ,  $f^{n-1}(x) = f(f^{n-2}(x)) \dots$ ,  $f^2(x) = f(f(x))$ . Consequently:

$$f^{n'}(x) = f'(f^{n-1}(x))f'(f^{n-2}(x)) \dots f'(x) \tag{1.1.5}$$

(1.1.5) enables us to compute the derivative at points on a periodic orbit in an elegant way. Indeed, suppose the three cycle  $\{p_0, p_1, p_2\}$  where  $p_1 = f(p_0)$ ,  $p_2 = f(p_1) = f^2(p_0)$  and  $f^3(p_0) = p_0 \dots$ . Then

$$f^{3'}(p_0) = f'(p_2)f'(p_1)f'(p_0) \tag{1.1.6}$$

Obviously, if we have  $n$  periodic points  $\{p_0, \dots, p_{n-1}\}$  the corresponding formulae is

$$f^{n'}(p_0) = \prod_{i=0}^{n-1} f'(p_i) \quad (1.1.7)$$

(Later on we shall use the derivative in order to decide whether a periodic orbit is stable or not. (1.1.7) implies that all points on the orbit is stable (unstable) simultaneously.)

We will now proceed by introducing some maps (difference equations) that have been frequently applied in population dynamics. Examples that show how to compute fixed points, periodic points, etc., will be taken from these maps. Some computations are performed in the next section, others are postponed to Section 1.3.

## 1.2 One-parameter family of maps

Here we shall briefly present some one-parameter family of maps which have often been applied in population dynamical studies. Since  $x$  is supposed to be the size of a population,  $x \geq 0$ .

The map

$$x \rightarrow f_\mu(x) = \mu x(1 - x) \quad (1.2.1)$$

is often referred to as the quadratic or the logistic map. The parameter  $\mu$  is called the intrinsic growth rate. Clearly  $x \in [0, 1]$ , otherwise  $x_t > 1 \Rightarrow x_{t+1} < 0$ . If  $\mu \in [0, 4]$  any iterate of  $f_\mu$  will remain in  $[0, 1]$ . Further we may notice that  $f_\mu(0) = f_\mu(1) = 0$  and  $x = c = 1/2$  is the only critical point.

**Definition 1.2.1.** A map  $f : I \rightarrow I$  is said to be unimodal if a)  $f(0) = f(1) = 0$ , and b)  $f$  has a unique critical point  $c$  which satisfies  $0 < c < 1$ . □

Hence (1.2.1) is a unimodal map on the unit interval. Note that unimodal maps are increasing on the interval  $[0, c)$  and are decreasing on  $(c, 1]$ .

The map

$$x \rightarrow f_r(x) = x e^{r(1-x)} \quad (1.2.2)$$

is called the Ricker map. Unlike the quadratic map,  $x \in [0, \infty)$ . The parameter  $r$  is positive.

**Exercise 1.2.1.** Show that the fixed points of (1.2.2) are 0 and 1 and that the critical point is  $1/r$ . □

The property that  $x \in [0, \infty)$  makes (1.2.2) much more preferable to biologists than (1.2.1). Indeed, let  $\mu > 4$  in (1.2.1). Then most points contained in  $[0, 1]$  will leave  $[0, 1]$  after a finite number of iterations (the point  $x_0 = 1/2$  will leave the unit interval after only one iteration), and once  $x_t > 1$ ,  $x_{t+1} < 0$  which, of course, is unacceptable from a biological point of view. Such problems do not arise by use of (1.2.2).

The map

$$x \rightarrow f_{a,b}(x) = \frac{ax}{(1+x)^b} \quad (1.2.3)$$


where  $a > 1$ ,  $b > 1$  is a two-parameter family of maps and is called the Hassel family.

**Exercise 1.2.2.** Show that  $x = 0$  and  $x = a^{1/b} - 1$  are the fixed points of (1.2.3) and that  $c = 1/(b - 1)$  is the only critical point for  $x > 0$ .  $\square$

The map

$$x \rightarrow T_a(x) = \begin{cases} ax & 0 \leq x \leq 1/2 \\ a(1-x) & 1/2 < x \leq 1 \end{cases} \quad (1.2.4)$$


Cynthia | AXA Graduate



AXA Global  
Graduate Program


Find out more and apply

redefining / standards





where  $a > 0$  is called the tent map for obvious reasons. We will pay special attention to the case  $a = 2$ . Note that  $T_a(x)$  attains its maximum at  $x = 1/2$  but that  $T'(1/2)$  does not exist. Since  $T_a(0) = T_a(1) = 0$  the map is unimodal on the unit interval.



**Figure 1:** The graphs of the functions: (a)  $f(x) = 4x(1 - x)$  (cf. (1.2.1)), and (b) the tent function (cf. (1.2.4) where  $a = 2$ ).

All functions defined in (1.2.1)–(1.2.4) have one critical point only. Such functions are often referred to as one-humped functions. In Figure 1a we show the graph of the quadratic functions (1.2.1) ( $\mu = 4$ ) and in Figure 1b the “tent” function (1.2.4) ( $a = 2$ ). In both figures we have also drawn the line  $y = x$  and we have marked the fixed points of the maps with dots.

As we have seen, maps (1.2.1)–(1.2.4) share much of the same properties. Our next goal is to explore this fact further.

**Definition 1.2.2.** Let  $f : U \rightarrow U$  and  $g : V \rightarrow V$  be two maps. If there exists a homeomorphism  $h : U \rightarrow V$  such that  $h \circ f = g \circ h$ , then  $f$  and  $g$  are said to be topological equivalent.  $\square$

**Remark 1.2.1.** A function  $h$  is a homeomorphism if it is one-to-one, onto and continuous and that  $h^{-1}$  is also continuous.  $\square$

The important property of topological equivalent maps is that their dynamics is equivalent. Indeed, suppose that  $x = f(x)$ . Then from the definition,  $h(f(x)) = h(x) = g(h(x))$ , so if  $x$  is a fixed point of  $f$ ,  $h(x)$  is a fixed point for  $g$ . In a similar way, if  $p$  is a periodic point of  $f$  of prime period  $n$  (i.e.  $f^n(p) = p$ ) we have from Definition 1.2.2 that  $f = h^{-1} \circ g \circ h \Rightarrow f^2 = (h^{-1} \circ g \circ h) \circ (h^{-1} \circ g \circ h) = h^{-1} \circ g^2 \circ h$  so clearly  $f^n = h^{-1} \circ g^n \circ h$ . Consequently,  $h(f^n(p)) = h(p) = g^n(h(p))$  so  $h(p)$  is a periodic point of prime period  $n$  for  $g$ .

**Proposition 1.2.1.** The quadratic map  $f : [0, 1] \rightarrow [0, 1] \ x \rightarrow f(x) = 4x(1 - x)$  is topological equivalent to the tent map

$$T : [0, 1] \rightarrow [0, 1] \quad x \rightarrow T(x) = \begin{cases} 2x & 0 \leq x \leq 1/2 \\ 2(1 - x) & 1/2 < x \leq 1 \end{cases}$$

□

**Proof.** We must find a function  $h$  such that  $h \circ f = T \circ h$ . Note that this implies that we also have  $f \circ h^{-1} = h^{-1} \circ T$  where  $h^{-1}$  is the inverse of  $h$ .

Now, define  $h^{-1}(x) = \sin^2(\pi x)/2$ . Then

$$\begin{aligned} f \circ h^{-1} &= f\left(\sin^2 \frac{\pi x}{2}\right) = 4 \sin^2 \frac{\pi x}{2} \left(1 - \sin^2 \frac{\pi x}{2}\right) \\ &= 4 \sin^2 \frac{\pi x}{2} \cos^2 \frac{\pi x}{2} = \left(2 \sin \frac{\pi x}{2} \cos \frac{\pi x}{2}\right)^2 = \sin^2 \pi x \end{aligned}$$

$$h^{-1} \circ T = h^{-1}(2x) = \sin^2 \pi x \quad 0 \leq x \leq \frac{1}{2}$$

$$h^{-1} \circ T = h^{-1}(2(1 - x)) = \sin^2(\pi - \pi x) = \sin^2 \pi x \quad \frac{1}{2} < x \leq 1$$

Thus,  $f \circ h^{-1} = h^{-1} \circ T$  which implies  $h \circ f = T \circ h$  so  $f$  and  $T$  are topological equivalent. □

### 1.3 Fixed points and periodic points of the quadratic map

Most of the theory that we shall develop in the next sections will be illustrated by use of the quadratic map (1.2.1). In many respects (1.2.1) will serve as a running example. Therefore, in order to prepare the ground we are here going to list some main properties.

The fixed points are obtained from  $x = \mu x(1 - x)$ . Thus the fixed points are  $x^* = 0$  (the trivial fixed point) and  $x^* = (\mu - 1)/\mu$  (the nontrivial fixed point). Note that the nontrivial fixed point is positive whenever  $\mu > 1$ . Assuming that (1.2.1) has periodic points of period two they must be found from  $p = f_\mu^2(p)$  and since

$$f_\mu^2(p) = f(\mu p(1 - p)) = \mu^2 p[1 - (\mu + 1)p + 2\mu p^2 - \mu p^3]$$

the two nontrivial periodic points must satisfy the cubic equation

$$\mu^3 p^3 - 2\mu^3 p^2 + \mu^2(\mu + 1)p + 1 - \mu^2 = 0 \tag{1.3.1}$$



Since every periodic point of prime period 1 is also a periodic point of period 2 we know that  $p = (\mu - 1)/\mu$  is a solution of (1.3.1). Therefore, by use of polynomial division we have

$$\mu^2 p^2 - (\mu^2 + \mu)p + \mu + 1 = 0 \quad (1.3.2)$$

Thus, the periodic points are

$$p_{1,2} = \frac{\mu + 1 \pm \sqrt{(\mu + 1)(\mu - 3)}}{2\mu} \quad (1.3.3)$$

where  $\mu > 3$  is a necessary condition for real solutions.

Period three points are obtained from  $p = f_\mu^3(p)$  and must be found by means of numerical methods. (It is possible to show after a somewhat cumbersome calculation that the three periodic points do not exist unless  $\mu > 1 + \sqrt{8}$ .)

In general, it is a hopeless task to compute periodic points of period  $n$  for a given map when  $n$  becomes large. Therefore, it is in many respects a remarkable fact that it is possible when  $\mu = 4$  in the quadratic map. We shall now demonstrate how such a calculation may be carried out, and in doing so, we find it convenient to express (1.2.1) as a difference equation rather than a map.

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



Real work  
International opportunities  
Three work placements



**Month 16**  
I was a construction  
supervisor in  
the North Sea  
advising and  
helping foremen  
solve problems





Thus consider

$$x_{t+1} = 4x_t(1 - x_t) \tag{1.3.4}$$

Let  $x_t = \sin^2 \varphi_t$ . Then from (1.3.4):

$$\sin^2 \varphi_{t+1} = 4 \sin^2 \varphi_t \cos^2 \varphi_t = \sin^2 2\varphi_t$$

Further:

$$\begin{aligned} \sin^2 \varphi_{t+2} &= 4 \sin^2 \varphi_{t+1} (1 - \sin^2 \varphi_{t+1}) \\ &= 4 \sin^2 2\varphi_t \cos^2 2\varphi_t = \sin^2 2^2 \varphi_t \end{aligned}$$

Thus, after  $n$  iterations

$$\sin^2 \varphi_{t+n} = \sin^2 2^n \varphi_t$$

which implies:

$$\varphi_{t+n} = \pm 2^n \varphi_t + l\pi$$

Now, if we have a period  $n$  orbit ( $x_{t+n} = x_t$ )

$$\sin^2 \varphi_{t+n} = \sin^2 \varphi_t$$

Hence:

$$\begin{aligned} \varphi_{t+n} = \pm \varphi_t + m\pi &\Leftrightarrow \pm 2^n \varphi_t + l\pi = \pm \varphi_t + m\pi \\ &\Leftrightarrow (2^n \pm 1)\varphi_t = (m - l)\pi \end{aligned}$$

so

$$\varphi_t = \frac{k\pi}{2^n \pm 1}$$

where  $k = m - l$ . Consequently, the periodic points are given by

$$p_i = \sin^2 \frac{k\pi}{2^n \pm 1} \tag{1.3.5}$$

**Example 1.3.1.** Compute all the period 1, period 2 and period 3 points of  $f(x) = 4x(1 - x)$ . The period 1 points (which of course are the same as the fixed points) are

$$\sin^2 \frac{\pi}{2 - 1} = 0 \quad \text{and} \quad \sin^2 \frac{\pi}{2 + 1} = 0.75$$

The period 2 points are the period 1 points (which do not have prime period 2) plus the prime period 2 points.

$$\sin^2 \frac{\pi}{5} = 0.34549 \quad \text{and} \quad \sin^2 \frac{2\pi}{5} = 0.904508$$

(The latter points may of course also be obtained from (1.3.3).)

There are six points of prime period 3. The points

$$\sin^2 \frac{\pi}{7} = 0.188255, \quad \sin^2 \frac{2\pi}{7} = 0.611260 \quad \text{and} \quad \sin^2 \frac{4\pi}{7} = 0.950484$$

are the periodic points in one 3-cycle, while the points

$$\sin^2 \frac{\pi}{9} = 0.116977, \quad \sin^2 \frac{2\pi}{9} = 0.4131759 \quad \text{and} \quad \sin^2 \frac{4\pi}{9} = 0.969846$$

are the periodic points on another orbit. (The reason why it is one 2-cycle but two 3-cycles is strongly related to how they are created.)  $\square$

**Exercise 1.3.1.** Use (1.3.5) to find all the period 4 points of  $f(x) = 4x(1 - x)$ . How many periodic points are there?  $\square$

Since  $f(x) = 4x(1 - x)$  is topological equivalent to the tent map we may use (1.3.5) together with Proposition 1.2.1 to find the periodic points of the tent map. Indeed, since  $h^{-1}(x) = \sin^2(\pi x/2) \Rightarrow h(x) = (2/\pi) \arcsin \sqrt{x}$  (cf. the proof of Proposition 1.2.1) the periodic points  $p$  of  $T(x)$  may be found from  $T(h(p)) = T((2/\pi) \arcsin \sqrt{p})$ . Thus the fixed points of the tent map are

$$\begin{aligned} T\left(\frac{2}{\pi} \arcsin \sqrt{0}\right) &= \frac{4}{\pi} \arcsin 0 = 0 \\ T\left(\frac{2}{\pi} \arcsin \sqrt{\frac{3}{4}}\right) &= 2 \left(1 - \frac{2}{\pi} \arcsin \sqrt{\frac{3}{4}}\right) = 0.6666 \end{aligned}$$

**Exercise 1.3.2.** Find the period 2 points of the tent map ( $a = 2$ ).  $\square$

We shall close this section by computing numerically some orbits of the quadratic map for different values of the parameter  $\mu$ :

$$\begin{aligned} &\mu = 1.8 \text{ and } x_0 = 0.8 \text{ gives the orbit} \\ &\{0.8 \ 0.2880 \ 0.3691 \ 0.4192 \ 0.4382 \ 0.4431 \ 0.4442 \ 0.4444 \ 0.4444 \ \dots\} \end{aligned}$$

Thus the orbit converges towards the point 0.4444 which is nothing but the fixed point  $(\mu - 1)/\mu$ . In this case the fixed point is said to be locally asymptotic stable. (A precise definition will be given in the next section.)

$$\mu = 3.2 \text{ and } x_0 = 0.6 \text{ gives:}$$

$$\{0.6 \ 0.7680 \ 0.5702 \ 0.7842 \ 0.5415 \ 0.7945 \ 0.5225 \ 0.7984 \ 0.5151$$

$$0.7993 \ 0.5134 \ 0.7994 \ 0.5131 \ 0.7995 \ 0.5130 \ 0.7995 \ 0.5130 \ \dots\}$$

Thus in this case the orbit does not converge towards the fixed point. Instead we find that the asymptotic behaviour is a stable periodic orbit of prime period 2. The points in the two-cycle are given by (1.3.3).

$$\mu = 4.0 \text{ and } x_0 = 0.30 \text{ gives}$$

$$\{0.30 \ 0.84 \ 0.5376 \ 0.9943 \ 0.02249 \ 0.0879 \ 0.3208 \ 0.8716 \ 0.4476 \ 0.9890 \ \dots\}$$

Although care should be taken by drawing a conclusion after a few iterations only, the last example suggests that there are no stable periodic orbit when  $\mu = 4$ . (A formal proof of this fact will be given later.)

**ie** business school

**93%**  
OF MIM STUDENTS ARE  
WORKING IN THEIR SECTOR 3 MONTHS  
FOLLOWING GRADUATION

**MASTER IN MANAGEMENT**

- STUDY IN THE CENTER OF MADRID AND TAKE ADVANTAGE OF THE UNIQUE OPPORTUNITIES THAT THE CAPITAL OF SPAIN OFFERS
- PROPEL YOUR EDUCATION BY EARNING A DOUBLE DEGREE THAT BEST SUITS YOUR PROFESSIONAL GOALS
- STUDY A SEMESTER ABROAD AND BECOME A GLOBAL CITIZEN WITH THE BEYOND BORDERS EXPERIENCE

Length: 10 MONTHS  
Av. Experience: 1 YEAR  
Language: ENGLISH / SPANISH  
Format: FULL-TIME  
Intakes: SEPT / FEB

**5 SPECIALIZATIONS**  
PERSONALIZE YOUR PROGRAM

**#10 WORLDWIDE**  
MASTER IN MANAGEMENT  
FINANCIAL TIMES

**55 NATIONALITIES**  
IN CLASS

[www.ie.edu/master-management](http://www.ie.edu/master-management) | [mim.admissions@ie.edu](mailto:mim.admissions@ie.edu) | [f](#) [t](#) [y](#) Follow us on IE MIM Experience



**Exercise 1.3.3.** Use a calculator or a computer to repeat the calculations above but use the initial values 0.6, 0.7 and 0.32 instead of 0.8, 0.6 and 0.3, respectively. Establish the fact that the long-time behaviour of the map when  $\mu = 1.8$  or  $\mu = 3.2$  is not sensitive to a slightly change of the initial conditions but that there is a strong sensitivity in the last case.  $\square$

## 1.4 Stability

Referring to the last example of the previous section we found that the equation  $x_{t+1} = 1.8x_t(1 - x_t)$  apparently possessed a stable fixed point and that the equation  $x_{t+1} = 3.2x_t(1 - x_t)$  did not. Both these equations are special cases of the quadratic family (1.2.1) so what the example suggests is that by increasing the parameter  $\mu$  in (1.2.1) there exists a threshold value  $\mu_0$  where the fixed point of (1.2.1) loses its stability.

Now, consider the general first order nonlinear equation

$$x_{t+1} = f_\mu(x_t) \tag{1.4.1}$$

where  $\mu$  is a parameter. The fixed point  $x^*$  satisfies  $x^* = f_\mu(x^*)$ .

In order to study the system close to  $x^*$  we write  $x_t = x^* + h_t$  and expand  $f_\mu$  in its Taylor series around  $x^*$  taking only the linear term. Thus:

$$x^* + h_{t+1} \approx f_\mu(x^*) + \frac{df}{dx}(x^*)h_t \tag{1.4.2}$$

which gives

$$h_{t+1} = \frac{df}{dx}(x^*)h_t \tag{1.4.3}$$

We call (1.4.3) the linearization of (1.4.1). The solution of (1.4.3) is given by (1.1.4). Hence, if  $|(df/dx)(x^*)| < 1$ ,  $\lim_{t \rightarrow \infty} h_t = 0$  which means that  $x_t$  will converge towards the fixed point  $x^*$ .

Now, we make the following definitions:

**Definition 1.4.1.** Let  $x^*$  be a fixed point of equation (1.4.1). If  $|\lambda| = |(df/dx)(x^*)| \neq 1$  then  $x^*$  is called a hyperbolic fixed point.  $\lambda$  is called the eigenvalue.  $\square$

**Definition 1.4.2.** Let  $x^*$  be a hyperbolic fixed point. If  $|\lambda| < 1$  then  $x^*$  is called a locally asymptotic stable hyperbolic fixed point.  $\square$

**Example 1.4.1.** Assume that  $\mu > 1$  and find the parameter interval where the fixed point  $x^* = (\mu - 1)/\mu$  of the quadratic map is stable.



**Solution:**  $f_\mu(x) = \mu x(1 - x)$  implies that  $f'(x) = \mu(1 - 2x) \Rightarrow |\lambda| = |f'(x^*)| = |2 - \mu|$ . Hence from Definition 1.4.2,  $1 < \mu < 3$  ensures that  $x^*$  is a locally asymptotic stable fixed point (which is consistent with our finding in the last example in the previous section).  $\square$

It is clear from Definition 1.4.2 that  $x^*$  is a *locally* stable fixed point. A formal argument that there exists an open interval  $U$  around  $x^*$  so that whenever  $|f'(x^*)| < 1$  and  $x \in U$  and that  $\lim_{n \rightarrow \infty} f^n(x) = x^*$  goes like this:

By the continuity of  $f$  ( $f$  is  $C'$ ) there exists an  $\varepsilon > 0$  such that  $|f'(x)| < K < 1$  for  $x \in [x^* - \varepsilon, x^* + \varepsilon]$ . Successive use of the mean value theorem then implies

$$\begin{aligned} |f^n(x) - x^*| &= |f^n(x) - f^n(x^*)| = |f(f^{n-1}(x)) - f(f^{n-1}(x^*))| \\ &\leq K|f^{n-1}(x) - f^{n-1}(x^*)| \leq K^2|f^{n-2}(x) - f^{n-2}(x^*)| \\ &\leq \dots \leq K^n|x - x^*| < |x - x^*| < \varepsilon \end{aligned}$$

so  $f^n(x) \rightarrow x^*$  as  $n \rightarrow \infty$ .

Motivated by the preceding argument we define:

**Definition 1.4.3.** Let  $x^*$  be a hyperbolic fixed point. We define the local stable and unstable manifolds of  $x^*$ ,  $W_{\text{loc}}^s(x^*)$ ,  $W_{\text{loc}}^u(x^*)$  as

$$W_{\text{loc}}^s(x^*) = \{x \in U \mid f^n(x) \rightarrow x^* \text{ as } n \rightarrow \infty \text{ and } f^n(x) \in U \text{ for all } n \geq 0\}$$

$$W_{\text{loc}}^u(x^*) = \{x \in U \mid f^n(x) \rightarrow x^* \text{ as } n \rightarrow -\infty \text{ and } f^n(x) \in U \text{ for all } n \leq 0\}$$

where  $U$  is a neighbourhood of the fixed point  $x^*$ .  $\square$

The definition of a hyperbolic stable fixed point is easily extended to periodic points.

**Definition 1.4.4.** Let  $p$  be a periodic point of (prime) period  $n$  so that  $|f^{n'}(p)| < 1$ . Then  $p$  is called an attracting periodic point.  $\square$

**Example 1.4.2.** Show that the periodic points 0.5130 and 0.7995 of  $x_{t+1} = 3.2x_t(1 - x_t)$  are stable and thereby proving that the difference equation has a stable 2-periodic attractor.

**Solution:** Since  $f(x) = 3.2x(1 - x) \Rightarrow f'(x) = 3.2(1 - 2x)$  we have from the chain rule (1.1.7) that  $f^{2'}(0.5130) = f'(0.7995)f'(0.5130) = -0.0615$ . Consequently, according to Definition 1.4.4, the periodic points are stable.  $\square$

**Exercise 1.4.1.** Use formulae (1.3.3) and compute the two-periodic points of the quadratic map in case of  $\mu = 3.8$ . Is the corresponding two-periodic orbit stable or unstable?  $\square$

**Exercise 1.4.2.** When  $\mu = 3.839$  the quadratic map has two 3-cycles. One of the cycles consists of the points 0.14989, 0.48917 and 0.9593 while the other consists of the points 0.16904, 0.53925 and 0.95384. Show that one of the 3-cycles is stable and that the other one is unstable.  $\square$



"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"  
Jane, Chinese architect


ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



An alternative way of investigating the behaviour of a one-dimensional map  $x \rightarrow f(x)$  is to use graphical analysis. The method is illustrated in Figures 2a,b, where we have drawn the graphs of a)  $f(x) = 2.7x(1 - x)$ , and b)  $f(x) = 3.2x(1 - x)$  together with the diagonal(s)  $y = x$ . Now, considering Figure 2a, let  $x_0 (= 0.2)$  be an initial value. A vertical line from  $x_0$  to the diagonal gives the point  $(x_0, x_0)$ , and if we extend the line to the graph of  $f$  we arrive at the point  $(x_0, f(x_0))$ . Then a horizontal line from the latter point to the diagonal gives the point  $(f(x_0), f(x_0))$ . Hence, by first drawing a vertical line from the diagonal to the graph of  $f$  and then a horizontal line back to the diagonal we actually find the image of a point  $x_0$  under  $f$  on the diagonal. Continuing in this fashion by drawing line segments vertically from the diagonal to the graph of  $f$  and then horizontally from the graph to the diagonal generate points  $(x_0, x_0), (f(x_0), f(x_0)), (f^2(x_0), f^2(x_0)), \dots, (f^n(x_0), f^n(x_0))$  on the diagonal which is nothing but a geometrical visualization of the orbit of the map  $x \rightarrow f(x)$ . Referring to Figure 2a we clearly see that the orbit converges towards a stable fixed point (cf. Example 1.4.1). On the other hand, in Figure 2b our graphical analysis shows that the fixed point is a repeller (cf. Exercise 1.3.2), and if we continue to iterate the map the result is a stable period 2 orbit, which is in accordance with Example 1.4.2. In Figure 2c all initial transitions have been removed and only the period 2 orbit is plotted.

**Exercise 1.4.3.** Let  $x \in [0, 1]$  and perform graphical analyses of the maps  $x \rightarrow 1.8x(1 - x)$ ,  $x \rightarrow 2.5x(1 - x)$  and  $x \rightarrow 4x(1 - x)$ . In the latter map use both a)  $x_0 = 0.2$ , and b)  $x_0 = 0.5$  as initial values. □



**Figure 2:** Graphical analyses of a)  $x \rightarrow 2.7x(1 - x)$  and b), c)  $x \rightarrow 3.2x(1 - x)$ .

**Exercise 1.4.4.** Consider the map  $f : \mathbb{R} \rightarrow \mathbb{R} \quad x \rightarrow x^3$ .

- a) The map has three fixed points. Find these.
- b) Use Definition 1.4.2 and discuss their stability properties.
- c) Verify the results in a) and b) by performing a graphical analysis of  $f$ . □

Let us close this section by discussing the concept structural stability. Roughly speaking, a map  $f$  is said to be structurally stable if a map  $g$  which is obtained through a small perturbation of  $f$  has essentially the same dynamics as  $f$ , so intuitively this means that the distance between  $f$  and  $g$  and the distance between their derivatives should be small.

**Definition 1.4.5.** The  $C^1$  distance between a map  $f$  and another map  $g$  is given by

$$\sup_{x \in \mathbb{R}} (|f(x) - g(x)|, |f'(x) - g'(x)|) \tag{1.4.4}$$

□

By use of Definition 1.4.5 we may now define structural stability in the following way:

**Definition 1.4.6.** The map  $f$  is said to be  $C^1$  structurally stable on an interval  $I$  if there exists  $\varepsilon > 0$  such that whenever (1.4.4)  $< \varepsilon$  on  $I$ ,  $f$  is topological equivalent to  $g$ . □

To prove that a given map is structurally stable may be difficult, especially in higher dimensional systems. However, our main interest is to focus on cases where a map is not structurally stable. In many respects maps with nonhyperbolic fixed points are standard examples of such maps as we now will demonstrate.

**Example 1.4.3.** When  $\mu = 1$  the quadratic map is not structurally stable.

Indeed, consider  $x \rightarrow f(x) = x(1 - x)$  and the perturbation  $x \rightarrow g(x) = x(1 - x) + \varepsilon$ . Obviously,  $x^* = 0$  is the fixed point of  $f$  and since  $|\lambda| = |f'(0)| = 1$ ,  $x^*$  is a nonhyperbolic fixed point. Moreover, the  $C^1$  distance between  $f$  and  $g$  is  $|\varepsilon|$ . Regarding  $g$ , the fixed points are easily found to be  $\bar{x} = \pm\sqrt{\varepsilon}$ . Hence, for  $\varepsilon > 0$  there are two fixed points and  $\varepsilon < 0$  gives no fixed points. Consequently,  $f$  is not structurally stable. □

**Example 1.4.4.** When  $\mu = 3$  the quadratic map is not structurally stable.

Let  $x \rightarrow f(x) = 3x(1 - x)$  and  $x \rightarrow g(x) = 3x(1 - x) + \varepsilon$  and again we notice that their  $C^1$  distance is  $\varepsilon$ . Regarding  $f$ , the fixed points are  $x_1^* = 0$  and  $x_2^* = 2/3$ . Further,  $|\lambda_1| = |f'(0)| = 3$ ,  $|\lambda_2| = |f'(2/3)| = 1$ . Thus  $x_1^*$  is a repelling hyperbolic fixed point while  $x_2^*$  is nonhyperbolic. Considering  $g$ , the fixed points are  $\bar{x}_1 = (1/3)(1 - \sqrt{1 + 3\varepsilon})$  and  $\bar{x}_2 = (1/3)(1 + \sqrt{1 + 3\varepsilon})$ . Note that  $\varepsilon = 0 \Rightarrow \bar{x}_1 = x_1^*$ ,  $\bar{x}_2 = x_2^*$ .) Further,  $|\sigma_1| = |g'(\bar{x}_1)| = |1 + 2\sqrt{1 + 3\varepsilon}|$  and  $|\sigma_2| = |g'(\bar{x}_2)| = |1 - 2\sqrt{1 + 3\varepsilon}|$ . Whatever the sign of  $\varepsilon$ ,  $\bar{x}_1$  is clearly a repelling fixed point (just as  $x_1^*$ ) since  $\sigma_1 > 1$ . Regarding  $\bar{x}_2$  it is stable in case of  $\varepsilon < 0$  and unstable if  $\varepsilon > 0$ .

The equation  $x = g^2(x)$  may be expressed as

$$-27x^4 + 54x^3 + (18\varepsilon - 36)x^2 + (8 - 18\varepsilon)x + 4\varepsilon - 3\varepsilon^2 = 0 \quad (1.4.5)$$

and since  $\bar{x}_1$  and  $\bar{x}_2$  are solutions of (1.4.5) we may use polynomial division to obtain

$$9x^2 - 12x - 3\varepsilon + 4 = 0 \quad (1.4.6)$$

which has the solutions  $x_{1,2} = (2/3)(1 \pm \sqrt{3\varepsilon})$ . Thus there exists a two- periodic orbit in case of  $\varepsilon > 0$ .

Excellent Economics and Business programmes at:



university of  
 groningen




“The perfect start  
 of a successful,  
 international career.”

**CLICK HERE**  
 to discover why both socially  
 and academically the University  
 of Groningen is one of the best  
 places for a student to be

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)



Moreover, cf. (1.1.7)  $g^{2'} = g'(x_1)g'(x_2) = 9(1 - 2x_1)(1 - 2x_2) = 1 - 48\varepsilon$  which implies that the two-periodic orbit is stable in case of  $\varepsilon > 0$ ,  $\varepsilon$  small. Consequently, when  $\varepsilon > 0$  there is a fundamental structural difference between  $f$  and  $g$  so  $f$  cannot be structurally stable. (Note that the problem is the nonhyperbolic fixed point, not the hyperbolic one.)  $\square$

As suggested by the previous examples a major reason why a map may fail to be structurally stable is the presence of the nonhyperbolic fixed point. Therefore it is in many respects natural to introduce the following definition:

**Definition 1.4.7.** Let  $x^*$  be a hyperbolic fixed point of a map  $f : \mathbb{R} \rightarrow \mathbb{R}$ . If there exists a neighbourhood  $U$  around  $x^*$  and an  $\varepsilon > 0$  such that a map  $g$  is  $C^1 - \varepsilon$  close to  $f$  on  $U$  and  $f$  is topological equivalent to  $g$  whenever  $(1.4.4) < \varepsilon$  on this neighbourhood, then  $f$  is said to be  $C^1$  locally structurally stable.  $\square$

There is a major general result on topological equivalent maps known under the name Hartman and Grobman's theorem. The "one-dimensional" formulation of this theorem (cf. Devaney, 1989) is:

**Theorem 1.4.1.** Let  $x^*$  be a hyperbolic fixed point of a map  $f : \mathbb{R} \rightarrow \mathbb{R}$  and suppose that  $\lambda = f'(x^*)$  such that  $|\lambda| \neq 0, 1$ . Then there is a neighbourhood  $U$  around  $x^*$  and a neighbourhood  $V$  of  $0 \in \mathbb{R}$  and a homeomorphism  $h : U \rightarrow \mathbb{R}$  which conjugates  $f$  on  $U$  to the linear map  $l(x) = \lambda x$  on  $V$ .  $\square$

For a proof, cf. Hartman (1964).

**Example 1.4.5.** Consider  $x \rightarrow f(x) = (5/2)x(1 - x)$ . The fixed point is  $x^* = 3/5$  and is clearly hyperbolic since  $\lambda = f'(x^*) = -1/2$ . Therefore, according to Theorem 1.4.1,  $f(x)$  on a neighbourhood about  $3/5$  is topological equivalent to  $l(x) = -(1/2)x$  on a neighbourhood about 0.  $\square$

## 1.5 Bifurcations

As we have seen, the map  $x \rightarrow f_\mu(x) = \mu x(1 - x)$  has a stable hyperbolic fixed point  $x^* = (\mu - 1)/\mu$  provided  $1 < \mu < 3$ . If  $\mu = 3$ ,  $\lambda = f'(x^*) = -1$ , hence  $x^*$  is no longer hyperbolic. If  $\mu = 3.2$  we have shown that there exists a stable 2- periodic orbit. Thus  $x^*$  experiences a fundamental change of structure when it fails to be hyperbolic which in our running example occurs when  $\mu = 3$ . Such a point will from now on be referred to as a bifurcation point. When  $\lambda = -1$ , as in our example, the bifurcation is called a flip or a period doubling bifurcation. If  $\lambda = 1$  it is called a saddle-node bifurcation. Generally, we will refer to a flip bifurcation as supercritical if the eigenvalue  $\lambda$  crosses the value  $-1$  outwards and that the 2-periodic orbit just beyond the bifurcation point is stable. Otherwise the bifurcation is classified as subcritical.

**Theorem 1.5.1.** Let  $f_\mu : \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \rightarrow f_\mu(x)$  be a one-parameter family of maps and assume that there is a fixed point  $(x^*, \mu_0)$  where the eigenvalue equals  $-1$ . Assume

$$a = \left( \frac{\partial f_\mu}{\partial \mu} \frac{\partial^2 f_\mu}{\partial x^2} + 2 \frac{\partial^2 f_\mu}{\partial x \partial \mu} \right) = \frac{\partial f_\mu}{\partial \mu} \frac{\partial^2 f_\mu}{\partial x^2} - \left( \frac{\partial f_\mu}{\partial x} - 1 \right) \frac{\partial^2 f}{\partial x \partial \mu} \neq 0 \text{ at } (x^*, \mu_0)$$

and

$$b = \left( \frac{1}{2} \left( \frac{\partial^2 f_\mu}{\partial x^2} \right)^2 + \frac{1}{3} \left( \frac{\partial^3 f_\mu}{\partial x^3} \right) \right) \neq 0 \text{ at } (x^*, \mu_0)$$

Then there is a smooth curve of fixed points of  $f_\mu$  which is passing through  $(x^*, \mu_0)$  and which changes stability at  $(x^*, \mu_0)$ . There is also a curve consisting of hyperbolic period-2 points passing through  $(x^*, \mu_0)$ . If  $b > 0$  the hyperbolic period-2 points are stable, i.e. the bifurcation is supercritical.  $\square$

**Proof.** Through a coordinate transformation it suffices to consider  $f_\mu$  so that for  $\mu = \mu_0 = 0$  we have  $f(x^*, 0) = x^*$  and  $f'(x^*, 0) = -1$ .

First we show that one without loss of generality may assume that  $x^* = 0$  in some neighbourhood of  $\mu = 0$ . To this end, define  $F(x, \mu) = f(x, \mu) - x$ . Then  $F'(x^*, \mu) = -2 \neq 0$  and by use of the implicit function theorem there exists a solution  $\bar{x}(\mu)$  of  $F(x, \mu) = 0$ . Next, define  $g(y, \mu) = f(y + \bar{x}(\mu), \mu) - \bar{x}(\mu)$ . Clearly,  $g(0, \mu) = 0$  for all  $\mu$ . Consequently,  $y = 0$  is a fixed point so in the following it suffices to consider  $x \rightarrow f(x)$  where  $x^*(\mu) = 0$  and  $f'(0, 0) = -1$ .

The Taylor expansion around  $(x^*, \mu) = (0, 0)$  of the latter map is

$$\begin{aligned} g_\eta(\xi) &= -\xi + \frac{\partial f}{\partial \mu} \eta + \frac{1}{2} \left( \frac{\partial^2 f}{\partial x^2} \xi^2 + 2 \frac{\partial^2 f}{\partial x \partial \mu} \xi \eta \right) + \frac{1}{6} \frac{\partial^3 f}{\partial x^3} \xi^3 + \text{higher order} \\ &= -\xi + \alpha \eta + \beta \xi^2 + c \eta \xi + d \xi^3 + \text{higher order} \end{aligned}$$

where the parameter  $\eta$  has the same weight as  $\xi^2$ . The composite  $(g \circ g)(\xi)$  may be expressed as

$$g_\eta^2(\xi) = \xi + \alpha \eta \xi + \beta \xi^3 + \text{higher order}$$

Thus, in order to have a system to study we must assume  $\alpha, \beta \neq 0$  which is equivalent to

$$\begin{aligned} \alpha &= -(2c + 2ab) = - \left( 2 \frac{\partial^2 f}{\partial x \partial \mu} + 2 \frac{\partial f}{\partial \mu} \cdot \frac{1}{2} \frac{\partial^2 f}{\partial x^2} \right) \neq 0 \\ \beta &= -(2d + 2b^2) = - \left( 2 \cdot \frac{1}{6} \frac{\partial^3 f}{\partial x^3} + 2 \left( \frac{1}{2} \frac{\partial^2 f}{\partial x^2} \right)^2 \right) \neq 0 \end{aligned}$$






Figure 3: The possible configurations of  $\xi^2 \rightarrow h(\xi) = \xi + \alpha\eta\xi + \beta\xi^3$ .

and we recognize the derivative formulae as nothing but what is stated in the theorem.

## American online

# LIGS University

is currently enrolling in the  
Interactive Online **BBA, MBA, MSc,**  
**DBA and PhD** programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive **Online** education
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](http://www.ligsuniversity.com).





Next, consider the truncated map

$$\xi^2 \rightarrow h(\xi) = \xi + \alpha\eta\xi + \beta\xi^3$$

Clearly, the fixed points are

$$\bar{\xi}_1 = 0, \quad \bar{\xi}_{2,3} = \pm \sqrt{-\frac{\alpha}{\beta}\eta}$$

Further,  $h'(\xi) = 1 + \alpha\eta + 3\beta\xi^2$  so  $h'(\bar{\xi}_1) = 1 + \alpha\eta$  and  $h'(\bar{\xi}_{2,3}) = 1 - 2\alpha\eta$ . Thus we have the following configurations (see Figure 3), and we may conclude that the stable period-2 orbits corresponds to  $\beta < 0$ , i.e.

$$\frac{1}{2} \left( \frac{\partial^2 f}{\partial x^2} \right)^2 + \frac{1}{3} \frac{\partial^3 f}{\partial x^3} > 0$$

□

**Example 1.5.1.** Show that the fixed point of the quadratic map undergoes a supercritical flip bifurcation at the threshold  $\mu = 3$ .

**Solution:** From the previous section we know that  $x^* = 2/3$  and  $f'(x^*) = -1$  when  $\mu = 3$ . We must show that the quantities  $a$  and  $b$  in Theorem 1.5.1 are different from zero and larger than zero respectively. By computing the various derivatives at  $(x^*, \mu_0) = (2/3, 3)$  we obtain:

$$a = \frac{2}{9}(-6) + 2 \left( -\frac{1}{3} \right) = -2 \neq 0 \quad \text{and} \quad b = \frac{1}{2}(-6)^2 + \frac{1}{3} \cdot 0 = 18 > 0$$

Thus the flip bifurcation is supercritical. When  $x^*$  fails to be stable, a stable period-2 orbit is established. □

**Exercise 1.5.1.** Show that the Ricker map  $x \rightarrow x \exp[r(1-x)]$ , cf. (1.2.2), undergoes a supercritical flip bifurcation at  $(x^*, r) = (1, 2)$ . □


**Exercise 1.5.2.** Consider the two parameter family of maps  $x \rightarrow -(1-\mu)x - x^2 + \alpha x^3$ . Show that the map may undergo both a sub- and supercritical flip bifurcation. □

As is clear from Definition 1.4.1 a fixed point will also lose its hyperbolicity if the eigenvalue  $\lambda$  equals 1. The general case then is that  $x^*$  will undergo a saddle-node bifurcation at the threshold where hyperbolicity fails. We shall now describe the saddle-node bifurcation.

Consider the map

$$x \rightarrow f_\mu(x) = x + \mu - x^2 \tag{1.5.1}$$

whose fixed points are  $x_{1,2}^* = \pm\sqrt{\mu}$ . Hence, when  $\mu > 0$  there are two fixed points which equals when  $\mu = 0$ . If  $\mu < 0$  there are no fixed points. In case of  $\mu > 0$ ,  $\mu$  small, we have  $f'_\mu(x_1^* = \sqrt{\mu}) = 1 - 2\sqrt{\mu} < 1$ , hence  $x_1^* = \sqrt{\mu}$  is stable. On the other hand:  $f'_\mu(x_2^* = -\sqrt{\mu}) = 1 + 2\sqrt{\mu} > 1$ , consequently  $x_2^*$  is unstable. Thus a saddle-node bifurcation is characterized by that there is no fixed point when the parameter  $\mu$  falls below a certain threshold  $\mu_0$ . When  $\mu$  is increased to  $\mu_0$ ,  $\lambda = 1$ , and two branches of fixed points are born, one stable and one unstable as displayed in the bifurcation diagram, see Figure 4a.



**Figure 4:** (a) The bifurcation diagram (saddle node) for the map  $x \rightarrow x + \mu - x^2$ .  
(b) The bifurcation diagram (transcritical) for the map  $x \rightarrow \mu x(1 - x)$ .

The other possibilities at  $\lambda = 1$  are the pitchfork and the transcritical bifurcations. The various configurations for the pitchfork are given at the end of the proof of Theorem 1.5.1 (see Figure 3). A typical configuration in the transcritical case is shown in Figure 4b as a result of considering the quadratic map at  $(x^*, \mu_0) = (0, 1)$ .

**Exercise 1.5.3.** Do the necessary calculations which leads to Figure 4b. □

**Exercise 1.5.4.**

- a) Show that the map  $x \rightarrow \mu - x^2$  undergoes a supercritical flip bifurcation at  $(x^*, \mu_0) = (1/2, 3/4)$ .
- b) Perform a graphical analysis of the map in the cases  $\mu = 1/2$  and  $\mu = 1$ . □

**Exercise 1.5.5.** Find possible bifurcation points of the map  $x \rightarrow \mu + x^2$ . If you detect a flip bifurcation decide whether it is sub- or supercritical. □

**Exercise 1.5.6.** Analyze the map  $x \rightarrow \mu x - x^3$ . □

## 1.6 The flip bifurcation sequence

We shall now return to the flip bifurcation. First we consider the quadratic map. In the previous section we used Theorem 1.5.1 to prove that the quadratic map  $x \rightarrow \mu x(1 - x)$  undergoes a supercritical flip bifurcation at the threshold  $\mu = \mu_0 = 3$ . This means that in case of  $\mu > \mu_0$ ,  $|\mu - \mu_0|$  small, there exists a stable 2-periodic orbit and according to our findings in Section 1.3 the periodic points are given by (1.3.3), namely

$$p_{1,2} = \frac{\mu + 1 \pm \sqrt{(\mu + 1)(\mu - 3)}}{2\mu}$$

The period 2 orbit will remain stable as long as

$$|f'(p_1)f'(p_2)| < 1$$

cf. Section 1.4. Thus, in our example,

$$|\mu(1 - 2p_1)\mu(1 - 2p_2)| < 1$$

i.e.

$$|1 - (\mu + 1)(\mu - 3)| < 1 \tag{1.6.1}$$

**DON'T EAT YELLOW SNOW**

What will your advice be?

Some advice just states the obvious. But to give the kind of advice that's going to make a real difference to your clients you've got to listen critically, dig beneath the surface, challenge assumptions and be credible and confident enough to make suggestions right from day one. At Grant Thornton you've got to be ready to kick start a career right at the heart of business.

Sound like you? Here's our advice: visit [GrantThornton.ca/careers/students](https://www.grantthornton.ca/careers/students)

Scan here to learn more about a career with Grant Thornton.

**Grant Thornton**  
An instinct for growth™


© Grant Thornton LLP. A Canadian Member of Grant Thornton International Ltd



from which we conclude that the 2-periodic orbit is stable as long as

$$3 < \mu < 1 + \sqrt{6} \tag{1.6.2}$$

Since  $\lambda = f^2{}' = f'(p_1)f'(p_2) = -1$  when  $\mu_1 = 1 + \sqrt{6}$  there is a new flip bifurcation taking place at  $\mu_1$  which in turn leads to a 4-periodic orbit. We also notice that while the fixed point  $x^* = (\mu - 1)/\mu$  is stable in the open interval  $I = (2, 3)$ , the length of the interval where the 2-periodic orbit is stable is roughly  $(1/2)I$ . In Figure 5a we show the graphs of the quadratic map in the cases  $\mu = 2.7$  (curve a) and  $\mu = 3.4$  (curve b) respectively, together with the straight line  $x_{t+1} = x_t$ .  $\mu = 2.7$  gives a stable fixed point  $x^*$  while  $\mu = 3.4$  gives an unstable fixed point. These facts are emphasized in the figure by drawing the slopes (indicated by dashed lines). The steepness of the slope at the fixed point of curve a is less than  $-45^\circ$ ,  $|\lambda| < 1$ , while  $\lambda < -1$  at the unstable fixed point located on curve b. In general, if  $f_\mu(x)$  is a single hump function (just as the quadratic map displayed in Figure 5a) the second iterate  $f_\mu^2(x)$  will be a two-hump function. In Figures 5b and 5c we show the relation between  $x_{t+2}$  and  $x_t$ . Figure 5b corresponds to  $\mu = 2.7$ , Figure 5c corresponds to  $\mu = 3.4$ . Regarding 5b the steepness of the slope is still less than  $45^\circ$  so the fixed point is stable. However, in 5c the slope at the fixed point is steeper than  $45^\circ$ , the fixed point is unstable and we see two new solutions of period 2.



**Figure 5:** (a) The quadratic map in the cases  $\mu = 2.7$  and  $\mu = 3.4$ . (b) and (c) The second iterate of the quadratic map in the cases  $\mu = 2.7$  and  $\mu = 3.4$ , respectively.

Let us now explore this mechanism analytically: Suppose that we have an  $n$ -periodic orbit consisting of the points  $p_0, p_1 \dots p_{n-1}$  such that

$$p_i = f_\mu^n(p_i) \tag{1.6.3}$$

Then by the chain rule (cf. (1.1.7))

$$f_{\mu}^{n'}(p_0) = \prod_{i=0}^{n-1} f_{\mu}'(p_i) = \lambda^n(p_0) \quad (1.6.4)$$

Hence, if  $|\lambda^n(p_0)| < 1$  the  $n$ -periodic orbit is stable, if  $|\lambda^n(p_0)| > 1$  the orbit is unstable.

Next, consider the  $2n$ -periodic orbit

$$p_i = f_{\mu}^{2n}(p_i) = f_{\mu}^n(f_{\mu}^n(p_i))$$

By appealing once more to the chain rule we obtain

$$f_{\mu}^{2n'}(p_0) = \left( \prod_{i=0}^{n-1} f_{\mu}'(p_i) \right)^2 = \lambda^{2n}(p_0) \quad (1.6.5)$$

This allows us to conclude that if the  $n$ -point cycle is stable (i.e.  $|\lambda^n| < 1$ ) then  $\lambda^{2n} < 1$  too. On the other hand, when the  $n$ -cycle becomes unstable (i.e.  $|\lambda^n| > 1$ ) then  $\lambda^{2n} > 1$  too. So what this argument shows is that when a periodic point of prime period  $n$  becomes unstable it bifurcates into two *new* points which are initially stable points of period  $2n$  and obviously there are  $2n$  such points. This is the situation displayed in Figure 5c. So what the argument presented above really says is that as the parameter  $\mu$  of the map  $x \rightarrow f_{\mu}(x)$  is increased periodic orbits of period  $2, 2^2, 2^3, \dots$  and so on are created through successive flip bifurcations. This is often referred to as the flip bifurcation sequence. Initially, all the  $2^k$  cycles are stable but they become unstable as  $\mu$  is further increased.

—

As already mentioned, if  $f_{\mu}(x)$  is a single-hump function, then  $f_{\mu}^2(x)$  is a two-hump function. In the same way,  $f_{\mu}^3(x)$  is a four-hump function and in general  $f_{\mu}^p$  will have  $2^{p-1}$  humps. This means that the parameter range where the period  $2^p$  cycles are stable shrinks through further increase of  $\mu$ . Indeed, the  $\mu$  values at successive bifurcation points act more or less as terms in a geometric series. In fact, Feigenbaum (1978) demonstrated the existence of a universal constant  $\delta$  (known as the Feigenbaum number or the Feigenbaum geometric ratio) such that

$$\lim_{n \rightarrow \infty} \frac{\mu_{n+1} - \mu_n}{\mu_{n+2} - \mu_{n+1}} = \delta = 4.66920 \quad (1.6.6)$$

where  $\mu_n, \mu_{n+1}$  and  $\mu_{n+2}$  are the parameter values at three consecutive flip bifurcations. From this we may conclude that there must exist an accumulation value  $\mu_a$  where the series of flip bifurcations converge. (Geometrically, this may happen as a “valley” of some iterate of  $f_{\mu}$  deepens and eventually touches the  $45^\circ$  line (cf. Figure 5c), then a saddle-node bifurcation ( $\lambda = 1$ ) will occur.)



As is true for our running example  $x \rightarrow \mu x(1 - x)$  we have proved that the first flip bifurcation occurs at  $\mu = 3$  and the second at  $\mu = 1 + \sqrt{6}$ . The point of accumulation for the flip bifurcations  $\mu_a$  is found to be  $\mu_a = 3.56994$ .

**Exercise 1.6.1.** Identify numerically the flip bifurcation sequence for the Ricker map (1.2.2).  $\square$

In the next sections we will describe the dynamics beyond the point of accumulation  $\mu_a$  for the flip bifurcations.

### 1.7 Period 3 implies chaos. Sarkovskii's theorem

Referring to our running example (1.2.1),  $x \rightarrow \mu x(1 - x)$  we found in the previous section that the point of accumulation for the flip bifurcation sequence  $\mu_a \approx 3.56994$ . We urge the reader to use a computer or a calculator to identify numerically some of the findings presented below.  $\mu \in [\mu_a, 4]$ .

.....Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



Click on the ad to read more



When  $\mu > \mu_a$ ,  $\mu - \mu_a$  small, there are periodic orbits of even period as well as aperiodic orbits. Regarding the periodic orbits, the periods may be very large, sometimes several thousands which make them indistinguishable from aperiodic orbits. Through further increase of  $\mu$  odd period cycles are detected too. The first odd cycle is established at  $\mu = 3.6786$ . At first these cycles have long periods but eventually a cycle of period 3 appears. In case of (1.2.1) the period-3 cycle occurs for the first time at  $\mu = 3.8284$ . This is displayed in Figure 6. (The point marked with a cross is the initially fixed point  $x^* = (\mu - 1)/\mu$  which became unstable at  $\mu = 3$ . It is also clear from the figure that the 3-cycle is established as the third iterate of (1.2.1) undergoes a saddle-node bifurcation.





Figure 6: A 3-cycle generated by the quadratic map.

An excellent way in order to present the dynamics of a map is to draw a bifurcation diagram. In such a diagram one plots the asymptotic behaviour of the map as a function of the bifurcation parameter. If we consider the quadratic map one plots the asymptotic behaviour as a function of  $\mu$ . If a map contains several parameters we fix all of them except one and use it as bifurcation parameter. In somewhat more detail a bifurcation diagram is generated in the following way: (A) Let  $\mu$  be the bifurcation parameter. Specify consecutive parameter values  $\mu_1, \mu_2, \dots, \mu_n$  where the distance  $|\mu_i - \mu_{i+1}|$  should be very small. (B) Starting with  $\mu_1$ , iterate the map from an initial condition  $x_0$  until the orbit of the map is close to the attractor and then remove initial transients. (C) Proceed the iteration and save many points of the orbit on the attractor. (D) Plot the orbit over  $\mu_1$  in the diagram. (E) Repeat the procedure for  $\mu_2, \mu_3, \dots, \mu_n$ .

Now, if the attractor is an equilibrium point for a given bifurcation value  $\mu_i$  there will be one point only over  $\mu_i$  in the bifurcation diagram. If the attractor is a two-period orbit there will be two points over  $\mu_i$ , and if the attractor is a  $k$  period orbit there are  $k$  points over  $\mu_i$ . Later on we shall see that an attractor may be an invariant curve as well as being chaotic. On such attractors there are quasiperiodic orbits and if either of these two types of attractors exist we will recognize them as line segments provided a sufficiently number of iteration points. The same is also true for periodic orbits when the period  $k$  becomes large. (In this context one may in fact think of quasiperiodic and chaotic orbits as periodic orbits where  $k \rightarrow \infty$ .) Hence, it may be a hopeless task to distinguish these types of attractors from each other by use of the bifurcation diagram alone.



**Figure 7:** The bifurcation diagram of the quadratic map in the parameter range  $2.9 \leq \mu \leq 4$

In the bifurcation diagram, Figure 7, we display the dynamics of the quadratic map in the interval  $2.9 \leq \mu \leq 4$ . The stable fixed point ( $\mu < 3$ ) as well as the flip bifurcation sequence is clearly identified. Also the period-3 “window” is clearly visible. Our goal in this and in the next sections is to give a thorough description of the dynamics beyond  $\mu_a$ .

We start by presenting the Li and Yorke theorem (Li and Yorke, 1975).

**Theorem 1.7.1.** Let  $f_\mu : \mathbb{R} \rightarrow \mathbb{R}$ ,  $x \rightarrow f_\mu(x)$  be continuous. Suppose that  $f_\mu$  has a periodic point of period 3. Then  $f_\mu$  has periodic points of all other periods.  $\square$

**Remark 1.7.1:** Theorem 1.7.1 was first proved in 1975 by Li and Yorke under the title “Period three implies chaos”. Since there is no unique definition of the concept chaos many authors today prefer to use the concept “Li and Yorke chaos” when they refer to Theorem 1.7.1. The essence of Theorem 1.7.1 is that once a period-3 orbit is established it implies periodic orbits of all other periods. Note, however, that Theorem 1.7.1 does not address the question of stability. We shall deal with that in the next section.  $\square$

We will now prove Theorem 1.7.1. Our proof is based upon the proof in Devaney (1989), not so much upon the original proof by Li and Yorke (1975).

**Proof.** First, note that (1): If  $I$  and  $J$  are two compact intervals so that  $I \subset J$  and  $J \subset f_\mu(I)$  then  $f_\mu$  has a fixed point in  $I$ . (2): Suppose that  $A_0, A_1, \dots, A_n$  are closed intervals and that  $A_{i+1} \subset f_\mu(A_i)$  for  $i = 0, \dots, n-1$ . Then there is at least one subinterval  $J_0$  of  $A_0$  which is mapped onto  $A_1$ . There is also a similar subinterval in  $A_1$  which is mapped onto  $A_2$  so consequently there is a  $J_1 \subset J_0$  so that  $f(J_1) \subset A_1$  and  $f_\mu^2(J_1) \subset A_2$ . Continuing in this fashion we find a nested sequence of intervals which map into the various  $A_i$  in order. Therefore there exists a point  $x \in A_0$  such that  $f_\mu^i(x) \in A_i$  for each  $i$ . We say that  $f_\mu(A_i)$  covers  $A_{i+1}$ .

Now, let  $a, b$  and  $c \in \mathbb{R}$  and suppose  $f_\mu(a) = b$ ,  $f_\mu(b) = c$  and  $f_\mu(c) = a$ . We further assume that  $a < b < c$ . Let  $I_0 = [a, b]$  and  $I_1 = [b, c]$ , cf. Figure 6. Then from our assumptions  $I_1 \subset f(I_0)$  and  $I_0 \cup I_1 \subset f(I_1)$ . The graph of  $f_\mu$ , cf. Figure 6, shows that there must be a fixed point of  $f_\mu$  between  $b$  and  $c$ . Similarly,  $f_\mu^2$  must have fixed points between  $a$  and  $b$  and at least one of them must have period 2. Therefore we let  $n \geq 2$ . Our goal is to produce a periodic point of prime period  $n > 3$ . Inductively, we define a nested sequence of intervals  $A_0, A_1, \dots, A_{n-2} \subset I_1$  as follows. Let  $A_0 = I_1$ . Since  $I_1 \subset f(I_1)$  there is a subinterval  $A_1 \subset A_0$  such that  $f_\mu(A_1) = A_0 = I_1$ . Then there is also a subinterval  $A_2 \subset A_1$  such that  $f_\mu(A_2) = A_1$  which implies  $f_\mu^2(A_2) = f_\mu(f_\mu(A_2)) = f_\mu(A_1) = A_0 = I_1$ . Continuing in this way there exists  $A_{n-2} \subset A_{n-3}$  such that  $f_\mu(A_{n-2}) = f_\mu(A_{n-3})$  so according to (2), if  $x \in A_{n-2}$  then  $f_\mu(x), f_\mu^2(x), \dots, f_\mu^{n-1}(x) \in A_0$  and indeed  $f_\mu^{n-2}(A_{n-2}) = A_0 = I_1$ .



Join the best at  
the Maastricht University  
School of Business and  
Economics!

Top master's programmes

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht University is the best specialist university in the Netherlands (Elsevier)

Visit us and find out why we are the best!  
Master's Open Day: 22 February 2014

[www.mastersopenday.nl](http://www.mastersopenday.nl)



Click on the ad to read more

Now, since  $I_0 \subset f_\mu(I_1)$  there exists a subinterval  $A_{n-1} \subset A_{n-2}$  such that  $f_\mu^{n-1}(A_{n-1}) = I_0$ . Finally, since  $I_1 \subset f_\mu(I_0)$  we have  $I_1 \subset f_\mu^n(A_{n-1})$  so that  $f_\mu^n(A_{n-1})$  covers  $A_{n-1}$ . Therefore, according to (1)  $f_\mu^n$  has a fixed point  $p$  in  $A_{n-1}$ .

Finally, we claim that  $p$  has prime period  $n$ . Indeed, the first  $n - 2$  iterations of  $p$  is in  $I_1$ , the  $(n - 1)$ st lies in  $I_0$  and the  $n$ -th is  $p$  again. If  $f_\mu^{n-1}(p)$  lies in the interior of  $I_0$  it follows that  $p$  has prime period  $n$ . If  $f_\mu^{n-1}(p)$  lies on the boundary, then  $n = 2$  or  $3$  and again we are done.  $\square$

Theorem 1.7.1 is a special case of Sarkovskii's theorem which came in 1964. However, it was written in Russian and published in an Ukrainian mathematical journal so it was not discovered and recognized in Western Europe and the U.S. prior to the work of Li and Yorke. We now state Sarkovskii's theorem:

**Theorem 1.7.2.** We order the positive integers as follows:

$$1 \triangleleft 2 \triangleleft 2^2 \triangleleft \dots \triangleleft 2^m \triangleleft 2^k(2n + 1) \triangleleft \dots \triangleleft 2^k \cdot 3 \triangleleft \dots \triangleleft 2 \cdot 3 \triangleleft 2n - 1 \triangleleft \dots \triangleleft 9 \triangleleft 7 \triangleleft 5 \triangleleft 3$$

Let  $f_\mu : I \rightarrow I$  be a continuous map of the compact interval  $I$  into itself. If  $f_\mu$  has a periodic point of prime period  $p$ , then it also has periodic points for any prime period  $q \triangleleft p$ .  $\square$

**Proof.** Cf. Devaney (1989) or Katok and Hasselblatt (1995).  $\square$

Clearly, Theorem 1.7.1 is a special case of Theorem 1.7.2. Also note that the first part in the Sarkovskii ordering ( $1 \triangleleft 2 \triangleleft 2^2 \dots \triangleleft 2^m$ ) corresponds to the flip bifurcation sequence as demonstrated through our treatment of the quadratic map. As the parameter  $\mu$  in (1.2.1) is increased beyond the point of accumulation for the flip bifurcations, Sarkovskii's theorem says that we approach a situation where there are an infinite number of periodic orbits.

## 1.8 The Schwarzian derivative

In the previous section we established through Theorems 1.7.1 and 1.7.2 that a map may have an infinite number of periodic orbits. Our goal in this section is to prove that in fact only a few of them are attracting (or stable) periodic orbits.

**Definition 1.8.1.** Let  $f : I \rightarrow I$  be a  $C^3$  function. The Schwarzian derivative  $Sf$  of  $f$  is defined as

$$Sf(x) = \frac{f'''(x)}{f'(x)} - \frac{3}{2} \left( \frac{f''(x)}{f'(x)} \right)^2 \tag{1.8.1}$$

$\square$

Considering  $f_\mu(x) = \mu x(1-x)$  we easily find that  $Sf_\mu(x) = -6/(1-2x)^2$ . Note that  $Sf_\mu < 0$  everywhere except at the critical point  $c = 1/2$ . (However, we may define  $Sf_\mu(1/2) = -\infty$ .)

The main result in this section is the following theorem which is due to Singer (1978):

**Theorem 1.8.1.** Let  $f$  be a  $C^3$  function with negative Schwarzian derivative. Suppose that  $f$  has one critical point  $c$ . Then  $f$  has at most three attracting periodic orbits.  $\square$

**Proof.** The proof consists of three steps.

(1) First we prove that if  $f$  has negative Schwarzian derivative then all  $f^n$  iterates also have negative Schwarzian derivatives.

To this end, assume  $Sf < 0$  and  $Sg < 0$ . Our goal is to show that  $S(f \circ g) < 0$ . Successive use of the chain rule gives:

$$\begin{aligned}(f \circ g)'(x) &= f'(g(x))g'(x) \\(f \circ g)''(x) &= f''(g(x))(g'(x))^2 + f'(g(x))g''(x) \\(f \circ g)'''(x) &= f'''(g(x))(g'(x))^3 + 3f''(g(x))g'(x)g''(x) + f'(g(x))g'''(x)\end{aligned}$$

Then (omitting function arguments) Definition 1.8.1 gives

$$S(f \circ g) = \frac{f'''g'^3 + 3f''g'g'' + f'g'''}{f'g'} - \frac{3}{2} \left( \frac{f''g'^2 + f'g''}{f'g'} \right)^2$$

which after some rearrangements may be written as

$$\left( \frac{f'''}{f'} - \frac{3}{2} \left( \frac{f''}{f'} \right)^2 \right) g'^2 + \frac{g'''}{g'} - \frac{3}{2} \left( \frac{g''}{g'} \right)^2 = Sf(g(x))(g'(x))^2 + Sg(x)$$

Thus  $S(f \circ g)(x) < 0$  which again implies  $Sf^n < 0$ .

(2) Next we show that if  $Sf < 0$  then  $f'(x)$  cannot have a positive local minimum.

To this end, assume that  $d$  is a critical point of  $f'(x)$ . Then  $f''(d) = 0$ , and since  $Sf < 0$  it follows from Definition 1.8.1 that  $f'''/f' < 0$  so  $f'''(d)$  and  $f'(d)$  have opposite signs. Graphically, it is then obvious that  $f'(x)$  cannot have a positive local minimum, and in the same way it is also clear that  $f'(x)$  cannot have a negative local maximum. Consequently, between any two consecutive critical points  $d_1$  and  $d_2$  of  $f'$  there must be a critical point  $c$  of  $f$  such that  $f'(c) = 0$  and moreover, (1) and (2) together imply that between any two consecutive critical points of  $f^{n'}$  there must be a critical point of  $f^n$ .

(3) By considering  $f^{n'}(x) = 0$  it follows directly from the chain rule that if  $f(x)$  has a critical point then  $f^n(x)$  will have a critical point too. Finally, let  $p$  be a point of period  $k$  on the attracting orbit and let  $I = (a, b)$  be the largest open interval around  $p$  where all points approach  $p$  asymptotically. Then  $f(I) \subset I$  and  $f^k(I) \subset I$ . Regarding the end points  $a$  and  $b$  we have: If  $f(a) = f(b)$  then of course there exists a critical point. If  $f(a) = a$  and  $f(b) = b$  (i.e. that the end points are fixed points) it is easy to see graphically that there exist points  $u$  and  $v$  such that  $a < u < p < v < b$  with properties  $f'(u) = f'(v) = 1$ . Then from (2) and the fact that  $f'(p) < 1$  there must be a critical point in  $(u, v)$ . In the last case  $f(a) = b$  and  $f(b) = a$  we arrive at the same conclusion by considering the second iterate  $f^2$ . Thus in the neighbourhood of any stable periodic point there must be either a pre-image of a critical point or an end point of the interval and we are done.  $\square$



**Empowering People. Improving Business.**

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

**BI NORWEGIAN BUSINESS SCHOOL**

EFMD EQUIS ACCREDITED

[www.bi.edu/master](http://www.bi.edu/master)





**Example 1.8.1.** Assume  $x \in [0, 1]$  and let us apply Theorem 1.8.1 on the quadratic map  $x \rightarrow f_\mu(x) = \mu x(1 - x)$ . For a fixed  $\mu \in (1, 3)$  the fixed point  $x^* = (\mu - 1)/\mu$  is stable, and since  $f_\mu(0) = f_\mu(1) = 0$  and the fact that 0 is repelling there is one periodic attractor, namely the period-1 attractor  $x^*$  which attracts the critical point  $c = 1/2$ .

When  $\mu \in [3, 4]$  both  $x^*$  and 0 are unstable fixed points. Thus according to Theorem 1.8.1 there is at most one attracting periodic orbit in this case. (Prior to  $\mu_a$  there is exactly one periodic attractor.) When  $\mu = 4$  the critical point is mapped on the origin through two iterations so there are no attracting periodic orbits in the case.  $\square$

**Example 1.8.2.** Let us close this section by giving an example which shows that Theorem 1.8.1 fails if the Schwarzian derivative is not negative. The following example is due to Singer (1978). Consider the map

$$x \rightarrow g(x) = -13.30x^4 + 28.75x^3 - 23.31x^2 + 7.86x \quad (1.8.2)$$

The map has one fixed point  $x^* = 0.7263986$ , and by considering  $g^2(x) = x$  there is also one 2-periodic orbit which consists of the points  $p_1 = 0.3217591$  and  $p_2 = 0.9309168$ .

Moreover:  $\lambda_1 = g'(x^*) = -0.8854$  and  $\sigma = g'(p_1)g'(p_2) = -0.06236$ . Thus both the fixed point and the 2-periodic orbit are attracting.

The critical point of  $g$  is  $c = 0.3239799$  and is attracted to the period-2 orbit so it does not belong to  $W_{loc}^s(x^*)$ , cf. Definition 1.4.3. The reason that  $x^*$  is not attracting  $c$  is that  $Sg(x^*) = 8.56 > 0$  thus the assumption  $Sg(x) < 0$  in Theorem 1.8.1 is violated.  $\square$

**Exercise 1.8.1.** Compute the Schwarzian derivative when  $f(x) = x^n$ .  $\square$

**Exercise 1.8.2.** Show that  $Sf(x) < 0$  when  $f$  is given by (1.2.2) (the Ricker case).  $\square$

## 1.9 Symbolic dynamics I

Up to this point we have mainly been concerned with fixed points and periodic orbits. The main goal of this section is to introduce a useful tool called symbolic dynamics which will help us to describe and understand dynamics of other types than we have discussed previously. To be more concrete, we shall in this section analyse the quadratic map  $x \rightarrow \mu x(1 - x)$  where  $\mu > 2 + \sqrt{5}$  on the interval  $I = [0, 1]$ , and as it will become clear, although almost all points in  $I$  eventually will escape  $I$ , there exists an invariant set  $\Lambda$  of points which will remain in  $I$ . We shall use symbolic dynamics to describe the behaviour of these points.



First we need some definitions. Consider  $x \rightarrow f(x)$ . Suppose that  $f(x)$  can take its values on two disconnected intervals  $I_1$  and  $I_2$  only. Define an infinite forward-going sequence of 0's and 1's  $\{a_k\}_{k=0}^{\infty}$  so that

$$a_k = 0 \quad \text{if} \quad f^k(x_0) \in I_1 \quad (1.9.1a)$$

$$a_k = 1 \quad \text{if} \quad f^k(x_0) \in I_2 \quad (1.9.1b)$$

Thus what we really do here is to represent an orbit of a map by an infinite sequence of 0's and 1's.

**Definition 1.9.1.**

$$\Sigma_2 = \{\bar{a} = (a_0 a_1 a_2 \dots) \mid a_k = 0 \text{ or } 1\} \quad (1.9.2)$$

□

We shall refer to  $\Sigma_2$  as the sequence space.

**Definition 1.9.2.** The itinerary of  $x$  is a sequence  $\phi(x) = a_0 a_1 \dots$  where  $a_k$  is given by (1.9.1). □

We now define one of the cornerstones of the theory of symbolic dynamics.

**Definition 1.9.3.** The shift map  $\sigma : \Sigma_2 \rightarrow \Sigma_2$  is given by

$$\sigma(a_0 a_1 a_2 a_3 \dots) = a_1 a_2 a_3 \dots \quad (1.9.3)$$

□

Hence the shift map deletes the first entry in a sequence and moves all the other entries one place to the left.


**Example 1.9.1.**  $\bar{a} = (1111 \dots)$  represents a fixed point under  $\sigma$  since  $\sigma(\bar{a}) = \sigma^n(\bar{a}) = (111 \dots)$ . Suppose  $\bar{a} = (001, 001, 001, \dots)$ . Then  $\sigma(\bar{a}) = (010, 010, 010, \dots)$ ,  $\sigma^2(\bar{a}) = (100, 100, 100, \dots)$  and  $\sigma^3(\bar{a}) = (001, 001, 001, \dots) = \bar{a}$ . Thus  $\bar{a} = (001, 001, 001, \dots)$  represents a periodic point of period 3 under the shift map. □

The previous example may obviously be generalized. Indeed, if  $\bar{a} = (a_0 a_1 \dots a_{n-1}, a_0 a_1 \dots a_{n-1}, \dots)$  there are  $2^n$  periodic points of period  $n$  under the shift map since each entry in the sequence may have two entries 0 or 1.

**Definition 1.9.4.** Let  $U$  be a subset of a set  $S$ .  $U$  is dense in  $S$  if the closure  $\bar{U} = S$ . □

**Definition 1.9.5.** If a set  $S$  is closed, contains no intervals and no isolated points it is called a Cantor set. □

**Proposition 1.9.1.** The set of all periodic orbits  $P_{\text{er}}(\sigma) = 2^n$  is dense in  $\Sigma_2$ . □



**Figure 8:** The quadratic map in the case  $\mu > 2 + \sqrt{5}$ . Note the subintervals  $I_1$  and  $I_2$  where  $f_\mu(x) = \mu x(1-x) \leq 1$

**Proof.** Let  $\bar{a} = (a_0 a_1 a_2 \dots)$  be in  $\Sigma_2$  and suppose that  $\bar{b} = (a_0 \dots a_{n-1}, a_0 \dots a_{n-1} \dots)$  represent the  $2^n$  periodic points. Our goal is to prove that  $\bar{b}$  converges to  $\bar{a}$ . By use of the usual distance function in a sequence space,  $d[\bar{a}, \bar{b}] = \sum(|a_i - b_i|/2^i)$  we easily find that  $d[\bar{a}, \bar{b}] < 1/2^n$ . Hence  $\bar{b} \rightarrow \bar{a}$ . □

## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info

**Helpmyassignment**

We now have the necessary machinery we need in order to analyse the quadratic map in case of  $\mu > 2 + \sqrt{5}$ .

Let  $x \rightarrow f(x) = \mu x(1-x)$  where  $\mu > 2 + \sqrt{5}$ . From the equation  $\mu x(1-x) = 1$  we find  $x = 1/2 + 1/2\sqrt{1-4\mu}$ . Hence in the intervals  $I_1 = [0, 1/2 - 1/2\sqrt{1-4\mu}]$  and  $I_2 = [1/2 + 1/2\sqrt{1-4\mu}, 1]$ ,  $f \leq 1$ , cf. Figure 8. Moreover,  $|f'(x)| = |\mu - 2\mu x|$  and whenever  $\mu > 2 + \sqrt{5}$  we find that  $|f'(x)| \geq \lambda > 1$ .

Denote  $I = [0, 1]$ . Then  $I \cap f^{-1}(I) = I_1 \cup I_2$  so if  $x \in I - (I \cap f^{-1}(I))$  we have  $f > 1$  (cf. Figure 8) which implies  $f^2 < 0$  and consequently  $f^n \rightarrow -\infty$ . All the other points will remain in  $I$  after one iteration. The second observation is that  $f(I_1) = f(I_2) = I$  so there must be a pair of open intervals, one in  $I_1$  and one in  $I_2$ , which is mapped into  $I - (I_1 \cup I_2)$  such that all points in these two intervals will leave  $I$  after two iterations. Continuing in this way by removing pairs of open intervals (i.e. first the interval  $I - (I_1 \cup I_2)$ , then two intervals, one in  $I_1$  ( $J_1$ ) and one in  $I_2$  ( $J_2$ ), then  $2^2$  open intervals, two from  $I_1 - J_1$ , two from  $I_2 - J_2 \dots$  and finally  $2^n$  intervals) from closed intervals we are left with a closed set  $\Lambda$  which is  $I$  minus the union of all the  $2^{n+1} - 1$  open sets. Hence  $\Lambda$  consists of the points that remain in  $I$  after  $n$  iterations,  $\Lambda \subset I \cap f^{-1}(I)$  and  $\Lambda$  consists of  $2^{n+1}$  closed intervals.

Now, associate to each  $x \in \Lambda$  a symbol sequence  $\{a_i\}_{i=1}^{\infty}$  of 0's and 1's such that  $a_k = 0$  if  $f^k(x) \in I_1$  and  $a_k = 1$  if  $f^k(x) \in I_2$ .

Next, define

$$I_{a_0 \dots a_n} = \{x \in I / x \in I_{a_0}, f(x) \in I_{a_1} \dots f^n(x) \in I_{a_n}\} \quad (1.9.4)$$

as one of the  $2^{n+1}$  closed subintervals in  $\Lambda$ . Our first goal is to show that  $I_{a_0 \dots a_n}$  is non-empty when  $n \rightarrow \infty$ . Indeed,

$$\begin{aligned} I_{a_0 \dots a_n} &= I_{a_0} \cap f^{-1}(I_{a_1}) \cap \dots \cap f^{-n}(I_{a_n}) \\ &= I_{a_0} \cap f^{-1}(I_{a_1 \dots a_n}) \end{aligned} \quad (1.9.5)$$

$I_{a_1}$  is nonempty. Then by induction  $I_{a_1 \dots a_n}$  is non-empty, and moreover, since  $f^{-1}(I_{a_1 \dots a_n})$  consists of two closed subintervals it follows that  $I_{a_0} \cap f^{-1}(I_{a_1 \dots a_n})$  consists of one closed interval. A final observation is that

$$\begin{aligned} I_{a_0 \dots a_n} &= I_{a_0} \cap \dots \cap f^{-(n-1)}(I_{a_{n-1}}) \cap f^{-n}(I_{a_n}) \\ &= I_{a_0 \dots a_{n-1}} \cap f^{-n}(I_{a_n}) \subset I_{a_0 \dots a_{n-1}} \end{aligned}$$

Consequently,  $I_{a_0 \dots a_n}$  is non-empty. Clearly the length of all sets  $I_{a_0 \dots a_n}$  approaches zero as  $n \rightarrow \infty$  which allows us to conclude that the itinerary  $\phi(x) = a_0 a_1 \dots$  is unique.

We now proceed by showing that  $\Lambda$  is a Cantor set. Assume that  $\Lambda$  contains an interval  $[a, b]$  where  $a \neq b$ . For  $x \in [a, b]$  we have  $|f'(x)| > \lambda > 1$  and by the chain rule  $|f^{n'}(x)| > \lambda^n$ . Let  $n$  be so large that  $\lambda^n|b - a| > 1$ . Then from the mean value theorem  $|f^n(b) - f^n(a)| \geq \lambda^n|b - a| > 1$  which means that  $f^n(b)$  or  $f^n(a)$  (or both) are located outside  $I$ . This is of course a contradiction so  $\Lambda$  contains no intervals.

To see that  $\Lambda$  contains no isolated points it suffices to note that any end point of the  $2^{n+1} - 1$  open intervals eventually goes to 0 and since  $0 \in \Lambda$  these end points are in  $\Lambda$  too. Now, if  $y \in \Lambda$  is isolated all points in a neighbourhood of  $y$  eventually will leave  $I$  which means that they must be elements of one of the  $2^{n+1} - 1$  open sets which are removed from  $I$ . Therefore, the only possibility such that  $y \in \Lambda$  is that there is a sequence of end points converging towards  $y$  so  $y$  cannot be isolated.

From the discussion above we conclude that the quadratic map where  $\mu > 2 + \sqrt{5}$  possesses an invariant set  $\Lambda$ , a Cantor set, of points that never leave  $I$  under iteration.  $\Lambda$  is a repelling set. Our final goal is to show that the shift map  $\sigma$  defined on  $\Sigma_2$  is topological equivalent to  $f$  defined on  $\Lambda$ .

Let  $f : \Lambda \rightarrow \Lambda$ ,  $f(x) = \mu x(1 - x)$ ,  $\sigma : \Sigma_2 \rightarrow \Sigma_2$ ,  $\sigma(a_0 a_1 a_2 \dots) = a_1 a_2 \dots$  and  $\phi : \Lambda \rightarrow \Sigma_2$ ,  $\phi(x) = a_0 a_1 a_2 \dots$ . We want to prove that  $\phi \circ f = \sigma \circ \phi$ .

Observe that

$$\phi(x) = a_0 a_1 a_2 \dots = \bigcap_{n \geq 0} I_{a_0 a_1 a_2 \dots a_n \dots}$$

Further

$$I_{a_0 a_1 \dots a_n} = I_{a_0} \cap f^{-1}(I_{a_1}) \cap \dots \cap f^{-n}(I_{a_n})$$

so

$$f(I_{a_0 a_1 \dots a_n}) = f(I_{a_0}) \cap (I_{a_1}) \cap \dots \cap f^{-n+1}(I_{a_n}) = I_{a_1} \cap \dots \cap f^{-n+1}(I_{a_n}) = I_{a_1 \dots a_n}$$

This implies that

$$\phi(f(x)) = \phi \left( f \left( \bigcap_{n \geq 0} I_{a_0 \dots a_n} \right) \right) = \phi \left( \bigcap_{n \geq 1} I_{a_1 \dots a_n} \right) = \sigma(\phi(x))$$

Thus,  $f$  and  $\sigma$  are topological equivalent maps.

## 1.10 Symbolic dynamics II

In Section 1.8 we proved that if a map  $f : I \rightarrow I$  with negative Schwarzian derivative possessed an attracting periodic orbit then there was a trajectory from the critical point  $c$  to the periodic orbit. Our goal here is to extend the theory of symbolic dynamics by assigning a symbol sequence to  $c$  or more precisely to  $f(c)$ . We will assume that  $f$  is unimodal. The theory will mainly be applied on periodic orbits.

Note, however, that the purpose of this section is somewhat different than the others so readers who are not too interested in symbolic dynamics may skip this section and proceed directly to the next where chaos is treated.

**Definition 1.10.1.** Let  $x \in I$ . Define the itinerary of  $x$  as  $\phi(x) = a_0 a_1 a_2 \dots$  where

$$a_j = \begin{cases} 0 & \text{if } f^j(x) < c \\ 1 & \text{if } f^j(x) > c \\ C & \text{if } f^j(x) = c \end{cases} \quad (1.10.1)$$

□

What is new here really is that we associate a symbol  $C$  to the critical point  $c$ . Also note that we may define two intervals  $I_0 = [0, c)$  and  $I_1 = (c, 1]$  such that  $f$  is increasing on  $I_0$  and decreasing on  $I_1$ .

**Brain power**

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**



Click on the ad to read more

**Definition 1.10.2.** The kneading sequence is defined as the itinerary of  $f(c)$ , i.e.

$$K(f) = \phi(f(c)) \tag{1.10.2}$$

□

**Example 1.10.1.**

- 1) Suppose that  $x \rightarrow f(x) = 2x(1 - x)$ . Then  $c = 1/2$  and  $f(c) = 1/2, f^2(c) = 1/2 \dots f^j(c) = 1/2$  so the kneading sequence becomes  $K(f) = (C C C C \dots)$  which also may be written as  $(C C \overline{C} \dots)$  where the bar refers to repetition.
- 2) Suppose that  $x \rightarrow f(x) = 4x(1 - x)$ .  $c = 1/2, f(c) = 1, f^2(c) \dots = f^j(c) = 0$  so  $K(f) = (1 0 0 \overline{0} \dots)$ . □

An unimodal map may of course have several itineraries.

**Example 1.10.2.** By use of a calculator we easily find that the possible itineraries of  $x \rightarrow 2x(1 - x)$  are

$$(00 \dots 0 C C \overline{C} \dots) (C C \overline{C} \dots) (10 \dots 0 C C \overline{C} \dots) (00 \overline{0} \dots) (100 \overline{0} \dots)$$

(The last two itineraries correspond to the orbits of  $x_0 = 0$  and  $x_0 = 1$  respectively. Note that the critical point is the same as the stable fixed point  $x^*$  in this example.

In case of  $x \rightarrow 3x(1 - x)$  we obtain the sequences

$$\begin{aligned} &(00 \dots 0 1 1 \overline{1} \dots) (C 1 1 \overline{1} \dots) (1 1 \overline{1} \dots) \\ &(10 \dots 0 1 1 \overline{1} \dots) (00 \overline{0} \dots) (100 \overline{0} \dots) \\ &(0 C 1 1 \overline{1} \dots) (1 C 1 1 \overline{1} \dots) \end{aligned}$$

where the last two itineraries correspond to the orbits of  $x_0 = (1/6)(3 - \sqrt{3})$  and  $x_0 = (1/6)(3 + \sqrt{3})$  respectively. □

The reader should also have in mind that periodic orbits with different periods may share the same itinerary.

Indeed, consider  $x \rightarrow 3.1x(1-x)$ . Then  $x^* = 0.6774 > c = 1/2$  so the itinerary of the fixed point becomes  $\phi(x^*) = (1\ 1\ \bar{1}\ \dots)$ . However, there is also a two-periodic orbit whose periodic points are (cf. formulae (1.3.3))  $p_1 = 0.7645$ ,  $p_2 = 0.5581$ . Again we observe that  $p_i > c$  so the itinerary of any of the two-periodic points is also  $(1\ 1\ \bar{1}\ \dots)$ . (When  $\mu$  becomes larger than 3.1 one of the periodic points eventually will become smaller than  $c$  which results in the itinerary  $(1\ 0\ 1\ 0\ 1\ 0\ \dots)$  or  $(0\ 1\ 0\ 1\ 0\ 1\ \dots)$ .)

Our next goal is to establish an ordering principle of the possible itineraries of a given map. Let  $\bar{a} = (a_0 a_1 a_2 \dots)$  and  $\bar{b} = (b_0 b_1 b_2 \dots)$ . If  $a_i = b_i$  for  $0 \leq i < n$  and  $a_n \neq b_n$  we say that the sequences have discrepancy  $n$ . Let  $S_n(\bar{a})$  be the number of 1's among  $a_0 a_1 \dots a_n$  and assume  $0 < C < 1$ .

**Definition 1.10.3.** Suppose that  $\bar{a}$  and  $\bar{b}$  have discrepancy  $n$ . We say that  $\bar{a} \prec \bar{b}$  if

$$S_{n-1}(\bar{a}) \text{ is even and } a_n < b_n \tag{1.10.3a}$$

$$S_{n-1}(\bar{a}) \text{ is odd and } a_n > b_n \tag{1.10.3b}$$

□

**Example 1.10.3.** Due to a) we have the following order:

$$(1\ 1\ 0\ \dots) \prec (1\ 1\ C\ \dots) \prec (1\ 1\ 1\ \dots)$$

Due to b) we have

$$(1\ 1\ 0\ \dots) \prec (1\ 0\ 1\ \dots) \prec (1\ 0\ 0\ \dots)$$

□

Also note that any two sequences with discrepancy 0 are ordered such that the sequence which has 0 as the first entry is of lower order than the one with  $C$  or 1 as the first entry. Thus:

$$(0\ 1\ \dots) \prec (C\ 1\ \dots) \prec (1\ 1\ \dots)$$

**Exercise 1.10.1.** Let  $\bar{a} = (0\ 1\ 1\ 0\ 1\ 1\ \dots)$  be a repeating sequence. Compute  $\sigma(\bar{a})$  and  $\sigma^2(\bar{a})$  and verify the ordering  $\bar{a} \prec \sigma(\bar{a}) \prec \sigma^2(\bar{a})$ . □

The following theorem (due to Milner and Thurston) relates the ordering of two symbol sequences to the values of two points in an interval.



**Theorem 1.10.1.** Let  $x, y \in I$

- a) If  $\phi(x) \prec \phi(y)$  then  $x < y$
- b) If  $x < y$  then  $\phi(x) \preceq \phi(y)$

□

**Proof.** Suppose that  $\phi(x) = (a_0a_1a_2 \dots)$  and  $\phi(y) = (b_0b_1b_2 \dots)$  and let  $n$  be the discrepancy of  $\phi(x)$  and  $\phi(y)$ . First, suppose  $n = 0$ . Then  $x < y$  since  $0 < C < 1$ . Next, suppose that a) is true with discrepancy  $n - 1$ . Our goal is to show that a) also is true with discrepancy  $n$ . By use of the shift we have  $\phi(f(x)) = (a_1a_2a_3 \dots)$  and  $\phi(f(y)) = (b_1b_2b_3 \dots)$ . Suppose  $a_0 = 0$ . Then  $\phi(f(x)) \prec \phi(f(y))$  since the number of 1's before the discrepancy is as before. Therefore  $f(x) < f(y)$  but since  $f$  is increasing on  $[0, c)$  it follows that  $x < y$ .

Next, assume  $a_0 = 1$ . Then  $\phi(f(x)) \succ \phi(f(y))$  since the number of 1's among the  $a_i$ 's ( $i \geq 1$ ) has been reduced by one. Therefore  $f(x) > f(y)$  which implies that  $x < y$  since  $f$  decreases on  $(c, 1]$ . If  $a_0 = C$  we have  $x = y = c$ .

## TURN TO THE EXPERTS FOR SUBSCRIPTION CONSULTANCY

Subscribe is one of the leading companies in Europe when it comes to innovation and business development within subscription businesses.

We innovate new subscription business models or improve existing ones. We do business reviews of existing subscription businesses and we develop acquisition and retention strategies.

Learn more at [linkedin.com/company/subscribe](https://www.linkedin.com/company/subscribe) or contact  
Managing Director Morten Suhr Hansen at [mha@subscribe.dk](mailto:mha@subscribe.dk)

**SUBSCRIB**✓**BE** - to the future



Regarding b) suppose  $x < y$  and assume that  $\phi(x)$  and  $\phi(y)$  has discrepancy  $n$ . First, note that if  $x < c < y$  we have directly  $\phi(x) < \phi(y)$ . Otherwise (i.e.  $x < y < c$  or  $c < x < y$ ) note that  $f^i$  is monotone in  $[x, y]$  for  $i \leq n$ . Since the number of 1's (cf. the chain rule) directly says if  $f^n$  is increasing or decreasing it is easily verified that  $\phi(x) \leq \phi(y)$ .  $\square$

**Theorem 1.10.2.** Let  $x = \varphi(\bar{a}) = a_0 a_1 a_2 \dots$  and suppose that  $x \rightarrow f(x)$  unimodal. Then  $\phi(\sigma^n \varphi(\bar{a})) \preceq K(f(c))$  for  $n \geq 1$ .  $\square$

**Proof.** Since the maximum of  $f$  is  $f(c)$  we have  $f(x) < f(c)$  and  $f^n(x) \leq f(c)$ . Moreover,  $\sigma x = \sigma(\varphi(\bar{a})) = a_1 a_2 \dots = \varphi(f(x))$  so inductively  $\sigma^n x = \varphi(f^n(x))$ . Therefore, according to Theorem 1.10.1

$$\phi(\sigma^n \varphi(\bar{a})) \preceq \phi(f(c)) = K(f(c))$$

$\square$

The essence of Theorem 1.10.2 is that any sequence  $\bar{a}$  such that  $\phi(x) = \bar{a}$  has lower order than the kneading sequence.

Now, consider periodic orbits. In order to simplify notation, repeating sequences (corresponding to periodic points) of the form  $\bar{a} = (a_0 a_1 \dots a_n a_0 a_1 \dots a_n a_0 a_1 \dots a_n \dots) = (a_0 a_1 \dots a_n \overline{a_1 a_1 \dots a_n})$  will from now on be written as  $\bar{a} = (a_0 a_1 \dots a_n)$ .

We also define a sequence  $\hat{a} = (a_0 \dots a_{n-1} \hat{a}_n)$  where  $\hat{a}_n = 1$  if  $a_n = 0$  or  $\bar{\hat{a}}_n = 0$  if  $a_n = 1$ . If  $\bar{b} = (b_0 b_1 \dots b_m)$ ,  $\bar{a} \cdot \bar{b} = (a_0 a_1 \dots a_n b_0 b_1 \dots b_m)$ .

Suppose that there exists a parameter value  $\mu$  such that there are two periodic orbits  $\gamma_1$  and  $\gamma_2$  of the same prime period. We say that the orbit  $\gamma_1$  is larger than the orbit  $\gamma_2$  if  $\gamma_1$  contains a point  $p_m$  which is larger than all the points of  $\gamma_2$ . Note that, according to Theorem 1.10.1, the itinerary of  $p_m$  satisfies  $\phi(p_i) \preceq \phi(p_m)$  where  $p_i$  are any of the other periodic points contained in  $\gamma_1$ .

Our main interest is the ordering of itineraries of periodic points  $p$  which satisfy:

- (A) The periodic point  $p$  shall be the largest point contained in the orbit.
- (B) Every other periodic orbit of the same prime period must have a periodic point which is larger than  $p$ .

Before we continue the discussion of (A) and (B) let us state a useful lemma.

**Lemma 1.10.1.** Given two symbol sequences  $\bar{a} = (a_0 a_1 a_2 \dots)$  and  $\bar{b} = (b_0 b_1 b_2 \dots)$ .

Suppose that  $a_0 = b_0 = 1$  and  $a_1 = b_1 = 0$ .  $a_j = b_j = 1$  for  $2 \leq j \leq l$ ,  $a_l = 0$ ,  $b_l = 1$ .

If  $l$  is even then  $\bar{b} \prec \bar{a}$ . If  $l$  is odd then  $\bar{a} \prec \bar{b}$ . □

**Proof.** Assume  $l$  even. Then the number of 1's before the discrepancy is odd and since  $b_l > a_l$  Definition 1.10.3 gives that  $\bar{b} \prec \bar{a}$ .

If  $l$  is odd the number of 1's before the discrepancy is even and since  $a_l = 0 < b_l = 1$ ,  $\bar{a} \prec \bar{b}$  according to the definition. □

A consequence of this theorem is that sequences that begin with 1 0 are of larger order than sequences which begin with 1 1. In the same way, a sequence which first entries are 1 0 0 is larger than one which begins with 1 0 1.

Now, consider the quadratic map  $x \rightarrow \mu x(1 - x)$ . Whenever  $\mu > 2$  the fixed point  $x^* = (\mu - 1)/\mu > c = 1/2$  so the (repeating) itinerary becomes  $\phi(x^*) = (1)$ . When  $x^*$  bifurcates at the threshold  $\mu = 3$ , the largest point  $p_1$  contained in the 2-cycle is always larger than  $c$ , hence the itinerary of  $p_1$  starts with 1 in the first entry. Therefore, when  $\mu > 3$ , there may be two possible itineraries (1 0) and (1 1) and clearly (1 1)  $\prec$  (1 0). We are interested in (1 0). Considering the 4-cycle which is created through another flip bifurcation the itinerary of the largest point contained in the cycle which we seek is (1 0 1 1) which is of larger order than the other alternatives.

Turning to odd periodic orbits, remember that they are established through saddle-node bifurcations, thus two periodic orbits, one stable and one unstable, are established at the bifurcation. Considering the stable 3-cycle at  $\mu = 3.839$  (see Exercise 1.4.2 or the bifurcation diagram, Figure 7) two of the points in the cycle 0.14989 and 0.48917 are smaller than  $c$  while the third one 0.95943 is larger. Hence the itinerary of largest order of 0.95493 is (1 0 0). Referring to Exercise 1.4.2 the largest point contained in the unstable 3-cycle is 0.95384 and the other points are 0.16904 and 0.53392. Hence the itinerary of 0.95384 of largest order is (1 0 1) and according to (A) and (B) this is the itinerary we are looking for, not the itinerary (1 0 0).

Therefore, the itineraries we seek are the ones that satisfy (A) and (B) and correspond to periodic points which are established through flip or saddle-node bifurcations as the parameter in the actual family is increased. (A final observation is that sequences which contain the symbol  $C$  are out of interest since they violate (B).)

Now, cf. our previous discussion, define the repeating sequences:

$$S_0 = (1) \quad S_1 = (10) \quad S_2 = (1011) \quad S_3 = (10111010)$$

and

$$S_{j+1} = S_j \cdot \hat{S}_j \tag{1.10.4}$$

Clearly, the sequence  $S_j$  has prime period  $2^j$  so it represents a periodic point with the same prime period.

Another important property is that  $S_j$  has an odd number of 1's. To see this, note that  $S_0 = (1)$  has an odd number of 1's. Next, assume that  $S_k = (S_0 \dots S_{k-1} 1)$  has an odd number of 1's. Then  $\hat{S}_k = (S_0 \dots S_{k-1} 0)$  has an even number of 1's so the concatenation  $S_{k+1} = S_k \cdot \hat{S}_k$  clearly has an odd number of 1's. (If  $S_k$  has a 0 at entry  $S_k$  we arrive at the same conclusion.) We have also that

$$\hat{S}_{j+1} = S_j \cdot S_j = S_j \tag{1.10.5}$$

Indeed, suppose  $S_k = (S_0 \dots S_k)$ . Then  $S_{k+1} = S_k \cdot \hat{S}_k = (S_0 \dots S_k S_0 \dots \hat{S}_k)$  so  $\hat{S}_{k+1} = (S_0 \dots S_k S_0 \dots S_k) = S_k \cdot S_k = S_k$ .

What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



**Lemma 1.10.2.** The sequences defined through (1.10.4) have the ordering

$$S_0 \prec S_1 \prec S_2 \prec S_3 \prec \dots$$

□

**Proof.** Assume that  $S_j = (S_0 \dots S_{j-1} S_j)$ . If  $S_j = 1$  there must be an even number of 1's among  $(S_0 \dots S_{j-1})$  so according to Definition 1.10.3a  $\hat{S}_j \prec S_j$ . If  $S_j = 0$  there is an odd number of 1's among  $(S_0 \dots S_{j-1})$  so according to Definition 1.10.3b  $\hat{S}_j \prec S_j$  also here. Therefore, by use of (1.10.5), we have  $S_j \succ \hat{S}_j = S_{j-1} \cdot S_{j-1} = S_{j-1}$ . □

Let us now turn to periodic orbits of odd period. The following lemma is due to Guckenheimer.

**Lemma 1.10.3.** The largest point  $p_m$  in the smallest periodic orbit of odd period  $n$  has itinerary  $\phi(p_m) = \bar{a}$  such that  $a_i = 0$  if  $i = 1(\bmod n)$  and  $a_i = 1$  otherwise. □

**Example 1.10.4.** If  $n = 3$ ,  $\phi(p_m) = (101101101\dots) = (101)$  which is in accordance with our previous discussion of 3-cycles. □

**Proof.** Suppose that we have a sequence  $\bar{a}$  and that there exists a number  $k$  such that  $a_k = 1$  and  $a_{k+1} = a_{k+2} = 0$ . Then by applying the shift map  $k$  times we arrive at  $\sigma^k(\bar{a}) = (100\dots)$  which according to Lemma 1.10.1 has larger order than any sequence with isolated 0's. Hence the sequence  $\sigma^k(\bar{a})$  violates (A) and (B).

Therefore, the argument above shows that the sequence we are looking for in this lemma must satisfy that if  $a_k = 0$  then both  $a_{k-1}$  and  $a_{k+1}$  must equal 1. Consequently there are blocks in  $\bar{a}$  of even length where the first and last entry of the blocks consist of 0 and the intermediate elements of 1's. As a consequence of Lemma 1.10.1 the longer these blocks are the smaller is the order of the sequence. Note that the blocks in this lemma have maximum length  $n + 1$  for a periodic sequence of period  $n$ . □

**Example 1.10.5.**  $(1 \underbrace{0110} 1101)$  is a 3-cycle where the length of the block is 4.

$(1 \underbrace{011110} 111)$  is a 5-cycle where the length of the block is 6. Clearly, the order of the 5-cycle is smaller than the order of the 3-cycle. □

**Lemma 1.10.4.** Let  $n > 1$  be an odd number. Then there is a periodic orbit of period  $n + 2$  which is smaller than all periodic orbits of period  $n$ . □

**Proof.** The lemma is an immediate consequence of how the itinerary in Lemma 1.10.3 is defined combined with the results of Lemma 1.10.1.  $\square$

We now turn to orbits of even period where the period is  $2^n \cdot m$  where  $m > 1$  is an odd number. The fundamental observation regarding the associated symbol sequences is that they may be written as  $S_{j+1}S_j \dots S_j$  or  $S_j\hat{S}_jS_j \dots S_j$  where the number of  $S_j$  blocks following  $S_{j+1}$  (or  $\hat{S}_j$ ) is  $m - 2$ . (See Guckenheimer (1977) for further details.)

**Example 1.10.6.** If  $n = 2$  (cf. 1.10.4) and  $m = 3$  we have the sequence (101110101011) and if  $n = 1$  and  $m = 5$  we arrive at (1011101010).  $\square$

**Lemma 1.10.5.** Let  $P$  be a periodic orbit of odd period  $k$ . Then there exists a periodic orbit of even period  $l = 2^n \cdot m$  where  $m > 1$  is odd which is smaller than any odd period orbit.  $\square$

**Proof.** From Lemma 1.10.4 we have that the longer the odd period is the smaller is the ordering of the associated symbol sequence. From Lemma 1.10.3 it follows that such a symbol sequence may be written as (101111...1110111...). Therefore by comparing an even period sequence with the odd one above it is clear that the even period sequence has 0 as entry at the discrepancy. If the even period is 2 it is two 1's before the discrepancy. If the even period is larger there are three consecutive 1's just prior to the 0 and since the first entry of the sequence is 1 there is an even number of 1's before the discrepancy also here and the result of the lemma follows.  $\square$

We need one more lemma which deals with periodic orbits of even period.

**Lemma 1.10.6.** Let  $u = 2^n \cdot l$ ,  $v = 2^n \cdot k$  and  $w = 2^m \cdot r$  where  $l$ ,  $k$  and  $r$  are odd numbers.

- a) Provided  $1 < k < l$  there are repeating symbol sequences of period  $u$  which has smaller order than any repeating symbol sequence of period  $v$ .
- b) Provided  $m > n$  there are repeating symbol sequences of period  $w$  which has smaller order than any repeating symbol sequence of period  $v$ .  $\square$

**Sketch of proof.** Regarding a) consider  $S_j$  such that  $j$  is odd. Then by carefully examining the various sequences we find that the discrepancy occurs at entry  $2^j(k + 2)$  in the repeating sequence of the  $2^n \cdot k$  periodic point and it happens as the last entry of the  $\hat{S}_j$  block (which of course is 1 since  $j$  is odd) differs from the same entry in the  $2^n \cdot l$  sequence. Now, since  $S_j\hat{S}_j$  has an odd number of 1's the number of 1's before the discrepancy is even, so according to Definition 1.10.3a we have that sequences of period  $2^n \cdot l$  are smaller than any sequence of period  $2^n \cdot k$ . (The case that  $j$  is even is left to the reader.)



Turning to b) and scrutinizing sequences  $\bar{a}$  of period  $2^m \cdot k$  it is clear that all of them have 1 0 1 1 as the first entries and that  $a_i = 1$  if  $i$  is even and  $a_i = 0$  if  $i = 1 \pmod{4}$ . Moreover, assuming  $k > r$  whenever  $m > n$  we find that at discrepancy the sequence of period  $w$  has 1 as its element and in fact it is the last 1 in 1 0 1 1. Now, since  $S_j \hat{S}_j S_j \dots S_j$  has an even number of 1's the observation above implies that the sequence of period  $2^n \cdot k$  must have an even number of 1's before the discrepancy so the result follows.  $\square$

Now at last, combining the results from Lemmas 1.10.1–1.10.6 we have established the following ordering for the itineraries of periodic points that satisfy (A) and (B):

$$2 \prec 2^2 \prec 2^3 \prec \dots \prec 2^n \prec 2^n(2l+1) \prec 2^n(2l-1) \prec \dots \prec 2^n \cdot 5 \prec 2^n \cdot 3 \prec 2^{n-1}(2l+1) \prec \dots \prec 2^{n-1} \cdot 3 \prec \dots \prec (2l+1) \prec (2l-1) \dots \prec 5 \prec 3$$

which is nothing but the ordering we find in Sarkovskii's theorem.

We do not claim that we actually have proved the theorem in all its details, our main purpose here have been to show that symbolic dynamics is a powerful tool when dealing with periodic orbits. For further reading, also of other aspects of symbolic dynamics we refer to Guckenheimer and Holmes (1990), Devaney (1989) and Collet and Eckmann (1980).

**gaiteye**  
Challenge the way we run

**EXPERIENCE THE POWER OF FULL ENGAGEMENT...**

**RUN FASTER.  
RUN LONGER..  
RUN EASIER...**

**READ MORE & PRE-ORDER TODAY  
WWW.GAITEYE.COM**



## 1.11 Chaos

As we have seen, the dynamics of  $x \rightarrow \mu x(1 - x)$  differs substantially depending on the value of the parameter  $\mu$ . For  $2 < \mu < 3$  there is a stable nontrivial fixed point, and in case of larger values of  $\mu$  we have detected periodic orbits both of even and odd period. If  $\mu > 2 + \sqrt{5}$  the dynamics is aperiodic and irregular and occurs on a Cantor set  $\Lambda$  and points  $x \in (I \setminus \Lambda)$  approaches  $-\infty$ . ( $I$  is the unit interval.)

In this section we shall deal with the concept chaos. Chaos may and has been defined in several ways. We have already used the concept when we stated “Period three implies chaos”.

Referring to the examples and exercises at the end of Section 1.3 we found that whenever the long-time behaviour of a system was a stable fixed point or a stable periodic orbit there was no sensitive dependence on the initial condition  $x_0$ . However, when  $x \rightarrow f(x) = 4x(1 - x)$  we have proved that there is no stable periodic orbit and moreover, we found a strong sensitivity on the initial condition. Assuming  $x \in [0, 1]$  and that  $x_0 = 0.30$  is one initial condition and  $x_{00} = 0.32$  is another we have  $|x_0 - x_{00}| = 0.02$  but most terms  $|f^k(x_0) - f^k(x_{00})| > 0.02$  and for some  $k$  ( $k = 9$ )  $|f^k(x_0) - f^k(x_{00})| \approx 1^-$  which indeed shows a strong sensitivity.

Motivated by the example above, if an orbit of a map  $f : I \rightarrow I$  shall be denoted as chaotic it is natural to include that  $f$  has sensitive dependence on the initial condition in the definition. It is also natural to claim that there is no convergence to any periodic orbit which is equivalent to, say, that periodic orbits must be dense in  $I$ . Our goal is to establish a precise definition of the concept chaos but before we do that let us first illustrate what we have discussed above by two examples.

**Example 1.11.1.** This is a “standard” example which may be found in many textbooks. Consider the map  $h : S' \rightarrow S'$ ,  $\theta \rightarrow h(\theta) = 2\theta$ . ( $h$  is a map from the circle to the circle.) Clearly,  $h$  is sensitive to initial conditions since the arc length between nearby points is doubled under  $h$ . Regarding the dense property, observe that  $h^n(\theta) = 2^n\theta$  so any periodic points must be obtained from the relation  $2^n\theta = \theta + 2k\pi$  or  $\theta = 2k\pi/(2^n - 1)$  where the integer  $k$  satisfies  $0 \leq k \leq 2^n$ . Hence in any neighbourhood of a point in  $S$  there is a periodic point so the periodic points are dense so  $h$  does not converge to any stable periodic orbit. Consequently,  $h$  is chaotic on  $S'$ . □

**Example 1.11.2.** Consider  $x \rightarrow f(x) = \mu x(1 - x)$  where  $\mu > 2 + \sqrt{5}$ . We claim that  $f$  is chaotic on the Cantor set  $\Lambda$ . In order to show sensitive dependence on the initial condition let  $\delta$  be less than the distance between the intervals  $I_0$  and  $I_1$  (cf. Figure 7). Next, assume  $x, y \in \Lambda$  where  $x \neq y$ . Then the itineraries  $\phi(x) \neq \phi(y)$  so after, say,  $k$  iterations  $f^k(x)$  is in  $I_0$  ( $I_1$ ) and  $f^k(y)$  is in  $I_1$  ( $I_0$ ). Thus  $|f^k(x) - f^k(y)| > \delta$  which establishes the sensitive dependence.

Since  $f : \Lambda \rightarrow \Lambda$  is topological equivalent to the shift map  $\sigma : \Sigma_2 \rightarrow \Sigma_2$  it suffices to show that the periodic points of  $\sigma$  are dense in  $\Sigma_2$ . Let  $\bar{a} = (a_1 \dots a_n)$  be a repeating sequence of a periodic point and let  $\bar{b} = (a_1 a_2 a_3 \dots)$  be the sequence of an arbitrary point and note that  $\sigma^n(\bar{a}) = \bar{a}$ . By use of the distance  $d$  between two symbol sequences one easily obtains  $d[\bar{a}, \bar{b}] < 1/2^n$  so in any neighbourhood of an arbitrary sequence (point) there is a periodic sequence (periodic point). Hence periodic points of  $f$  are dense (and unstable).  $\square$

In our work towards a definition of chaos we will now focus on the sensitive dependence on the initial condition.

If a map  $f : \mathbb{R} \rightarrow \mathbb{R}$  has a fixed point we know from Section 1.4 that if the eigenvalue  $\lambda$  of the linearized system satisfies  $-1 < \lambda < 1$  the fixed point is stable and not sensitive to changes of the initial condition. If  $|\lambda| > 1$  one may measure the degree of sensitivity by the size of  $|\lambda|$ . We may use the same argument if we deal with periodic orbits of period  $k$  except that we on this occasion consider the eigenvalue of every periodic point contained on the orbit. If a system is chaotic it is natural to consider the case  $k \rightarrow \infty$  since we may think of a chaotic orbit as one having an infinite period. Therefore, define

$$\eta = \lim_{k \rightarrow \infty} \left| \frac{d}{dx} f^k(x)_{x=x_0} \right|^{1/k} \tag{1.11.1}$$

where we have used the  $k$ 'th root in order to avoid problems in order to obtain a well defined limit. If  $x_0$  is a fixed point  $\lambda = |(df/dx)(x = x_0)|$ . For a general orbit starting at  $x_0$  we may think of  $\eta$  as an average measure of sensitivity (or insensitivity) over the whole orbit. Let  $L = \ln \eta$ , that is

$$L = \lim_{k \rightarrow \infty} \ln \left| \frac{d}{dx} f^k(x_0) \right|^{1/k} = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{n=0}^{k-1} \ln |f'(x = x_n)| \tag{1.11.2}$$

The number  $L$  is called the Lyapunov exponent and if  $L > 0$  (which is equivalent to  $|\lambda| > 1$ ) we have sensitive dependence on the initial condition. By use of  $L$  we may now define chaos.

**Definition 1.11.1.** The orbit of a map  $x \rightarrow f(x)$  is called chaotic if

- 1) It possesses a positive Lyapunov exponent, and
- 2) it does not converge to a periodic orbit (that is, there does not exist a periodic orbit

$$y_t = y_{t+T} \text{ such that } \lim_{t \rightarrow \infty} |x_t - y_t| = 0.) \tag{1.11.3} \quad \square$$

Note that 2) is equivalent to, say, that periodic orbits are dense.

In most cases the Lyapunov exponent must be computed numerically and in cases where  $L$  is slightly larger than zero such computations have to be performed by some care due to accumulation effects of round-off errors. Note, however, that there exists a theorem saying that  $L$  is stable under small perturbations of an orbit.

**Example 1.11.3.** Compute  $L$  for the map  $h : S^1 \rightarrow S^1$ ,  $h(\theta) = 2\theta$ . In this case  $h' = 2$  for all points on the orbit so

$$L = \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{n=0}^{k-1} \ln |h'(x = x_n)| = \lim_{k \rightarrow \infty} \frac{1}{k} \cdot k \ln 2 = \ln 2 > 0$$

and since the periodic orbits are dense  $h$  is chaotic. □

**Example 1.11.4.** Compute  $L$  for the two periodic orbit of  $x \rightarrow f(x) = \mu x(1 - x)$  where  $3 < \mu < 1 + \sqrt{6}$ . Referring to formulae (1.3.3) the periodic points are

$$p_{1,2} = \frac{\mu + 1 \pm \sqrt{(\mu + 1)(\mu - 3)}}{2\mu}$$

Thus,

$$\begin{aligned} L &= \lim_{k \rightarrow \infty} \frac{1}{k} \{ \ln |f'(x = p_1)| + \ln |f'(x = p_2)| + \ln |f'(x = p_1)| + \dots + \ln |f'(x = p_2)| \} \\ &= \lim_{k \rightarrow \infty} \frac{1}{k} \left\{ \frac{k}{2} \ln |f'(x = p_1)| + \frac{k}{2} \ln |f'(x = p_2)| \right\} \\ &= \frac{1}{2} \ln |f'(x = p_1)f'(x = p_2)| \end{aligned}$$

This e-book  
is made with  
**SetaPDF**





**SETASIGN**

PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)



Since

$$f'(x = p_1)f'(x = p_2) = \mu(1 - 2p_1)\mu(1 - 2p_2) = 1 - (\mu + 1)(\mu - 3)$$

it follows that  $L = (1/2) \ln |1 - (\mu + 1)(\mu - 3)|$  and as expected  $L < 0$  whenever  $3 < \mu < 1 + \sqrt{6}$ . (Note that if  $\mu > 1 + \sqrt{6}$  then  $L > 0$  but the map is of course not chaotic since there in this case (provided  $|\mu - (1 + \sqrt{6})|$  small) exists a stable 4-periodic orbit with negative  $L$ .)  $\square$

**Example 1.11.5.** Show that the Lyapunov exponents of almost all orbits of the map  $f : [0, 1] \rightarrow [0, 1]$ ,  $x \rightarrow f(x) = 4x(1 - x)$  is  $\ln 2$ .

**Solution:** From Proposition 1.2.1 we know that  $f(x)$  is topological equivalent to the tent map  $T(x)$ . The “nice” property of  $T(x)$  which we shall use is that  $T'(x) = 2$  for all  $x \neq c = 1/2$ . Moreover,  $h \circ f = T \circ h$  implies that  $h'(f(x))f'(x) = T'(h(x))h'(x)$  so

$$f'(x) = \frac{T'(h(x))h'(x)}{h'(f(x))}$$

We are now ready to compute the Lyapunov exponent:

$$\begin{aligned} L &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \ln |f'(x = x_i)| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \ln \left| \frac{T'(h(x_i))h'(x_i)}{h'(f(x_i))} \right| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \ln |T'(h(x_i))| + \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \{\ln |h'(x_i)| - \ln |h'(f(x_i))|\} \end{aligned}$$

Since  $x_{i+1} = f(x_i)$  the latter sum may be written as

$$\lim_{n \rightarrow \infty} \frac{1}{n} \{\ln |h'(x_0)| - \ln |h'(x_n)|\}$$

which is equal to zero for almost all orbits. Thus, for almost all orbits:

$$L = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \ln |T'(h(x_i))| = \lim_{n \rightarrow \infty} \frac{1}{n} \cdot n \ln 2 = \ln 2$$

$\square$

For comparison reasons we have also computed  $L$  numerically with initial value  $x_0 = 0.30$  in the example above. Denoting the Lyapunov exponent of  $n$  iterations for  $L_n$  we find  $L_{100} = 0.67547$ ,  $L_{1000} = 0.69227$  and  $L_{5000} = 0.69308$  so in this example we do not need too many terms in order to show that  $L > 0$ .

A final comment is that since we have proved earlier (cf. Example 1.8.1) that the quadratic map does not possess any stable orbits in case of  $\mu = 4$ , Definition 1.11.1 directly gives that almost all orbits of the map are chaotic. Other properties of Lyapunov exponents may be obtained in the literature. See for example Tsujii (1993) and Thieullen (1994).

## 1.12 Superstable orbits and a summary of the dynamics of the quadratic map

The quadratic map has two fixed points. One is the trivial one  $x^* = 0$  which is stable if  $\mu < 1$  and unstable if  $\mu > 1$ . If  $\mu > 1$  the nontrivial fixed point is  $x^* = (\mu - 1)/\mu$  and as we have shown this fixed point is stable whenever  $1 < \mu < 3$ . Whenever  $\mu > 2$  the fixed point is larger than the critical point  $c$ . At  $\mu = 3$  the map undergoes a supercritical flip bifurcation and in the interval  $3 < \mu < 1 + \sqrt{6}$  the quadratic map possesses a stable period-2 orbit which has a negative Lyapunov exponent. The periodic points are given by formulae (1.3.3).

At the threshold  $\mu = 1 + \sqrt{6}$  there is a new (supercritical) flip bifurcation which creates a stable orbit of period  $2^2$  and through further increase of  $\mu$  stable orbits of period  $2^k$  are established. However, the parameter intervals where the period  $2^k$  cycles are stable shrinks as  $\mu$  is enlarged so the  $\mu$  values at the bifurcation points act more or less as terms in a geometric series. By use of the Feigenbaum geometric ratio one can argue that there exists an accumulation value  $\mu_a$  for the series of flip bifurcations. Regarding the quadratic map,  $\mu_a = 3.56994$ . In the parameter interval  $\mu_a < \mu \leq 4$  we have seen that the dynamics is much more complicated.


Still considering periodic orbits, Sarkovskii's theorem tells us that periodic orbits occur in a definite order so beyond  $\mu_a$  there are periodic orbits of periods given by Theorem 1.7.2 (see also Section 1.10). Even in cases where such orbits are stable they may be difficult to distinguish from non-periodic orbits due to the long period. In many respects the ultimate event occurs at the threshold  $\mu = 1 + \sqrt{8}$  where a 3-periodic orbit is created because period 3 implies orbits of all other periods which is the content both in Li and Yorke and in Sarkovskii's theorem.

Chaotic orbits may be captured by use of Lyapunov exponents. In Figure 9 we show the value of the Lyapunov exponent  $L$  for  $\mu \in [\mu_a, 4]$ .  $L < 0$  corresponds to stable periodic orbits,  $L > 0$  corresponds to chaotic orbits. (Figure 9 should be compared to the bifurcation diagram, Figure 7.) The regions where we have periodic orbits are often referred to as windows. The largest window found in Figure 7 (or 9) is the period 3 window. The periodic orbits in the interval  $3 < \mu < \mu_a$  are created through a series of flip bifurcations. However, the period-3 orbit is created through a saddle-node bifurcation. In fact, every window of periodic orbits beyond  $\mu_a$  is created in this way so just beyond the bifurcation value there is one stable and one unstable orbit of the same period. (If  $\mu$  is slightly larger than  $1 + \sqrt{8}$  there is one stable and one unstable orbit of period 3.) Within a window there may be flip bifurcations before chaos is established again, cf. Figure 7. Since the quadratic map has negative Schwarzian derivative there is at most one stable periodic orbit for each value of  $\mu$ .


# Free eBook on Learning & Development

By the Chief Learning Officer of McKinsey

[Download Now](#)



[Click on the ad to read more](#)



**Figure 9:** The value of the Lyapunov exponent for  $\mu \in [\mu_a, 4]$ .  $L < 0$  corresponds to stable periodic orbits.  
 $L > 0$  corresponds to chaotic orbits.

There is a way to locate the periodic windows. The vital observation is that at the critical point  $c$ ,  $f'(c) = 0$ , so accordingly  $\ln |f'(c)| = -\infty$  which implies  $L < 0$  and consequently a stable periodic orbit. Also, confer Singer's theorem (Theorem 1.8.1).

**Definition 1.12.1.** Given a map  $f : I \rightarrow I$  with one critical point  $c$ . Any periodic orbit  $\pi$  passing through  $c$  is called a superstable orbit.  $\square$

Hence, by searching for superstable orbits one may obtain a representative value of the location of a periodic window. Indeed, any superstable orbit of period  $n$  must satisfy the equation

$$f_\mu^n(c) = c \tag{1.12.1}$$

**Example 1.12.1.** Consider the quadratic map and let us find the value of  $\mu$  such that  $f_\mu^3(1/2) = 1/2$ .

We have

$$\begin{aligned} c = \frac{1}{2} \Rightarrow f_\mu(c) &= \frac{1}{4}\mu \Rightarrow f_\mu^2(c) = \frac{1}{4}\mu^2 - \frac{1}{16}\mu^3 \\ \Rightarrow f_\mu^3(c) &= \left(\frac{1}{4}\mu^3 - \frac{1}{16}\mu^4\right) \left\{1 - \left(\frac{1}{4}\mu^2 - \frac{1}{16}\mu^3\right)\right\} \end{aligned}$$

Hence, the equation  $f_\mu^3(1/2) = 1/2$  becomes

$$\mu^7 - 8\mu^6 + 16\mu^5 + 16\mu^4 - 64\mu^3 + 128 = 0 \tag{1.12.2}$$



By inspection,  $\mu = 2$  is a solution of (1.12.2) so after dividing by  $\mu - 2$  we arrive at

$$\mu^6 - 6\mu^5 + 4\mu^4 + 24\mu^3 - 16\mu^2 - 32\mu - 64 = 0 \quad (1.12.3)$$

This equation may be solved numerically by use of Newton's method and if we do that we find that the only solution in the interval  $\mu_a \leq \mu \leq 4$  is  $\mu = 3.83187$ . Therefore, there is only one period-3 window and the location clearly agrees both with the bifurcation diagram, Figure 7 and Figure 9. In the same way, by solving  $f_\mu^4(1/2) = 1/2$  one finds that the only solution which satisfies  $\mu_a < \mu < 4$  is  $\mu = 3.963$  which shows that there is also only one period-4 window. However, if one solves  $f_\mu^5(1/2) = 1/2$  one obtains three values which means that there exists three period-5 windows. The first one occurs around  $\mu_1 = 3.739$  and is visible in the bifurcation diagram, Figure 7. The others have almost no widths, the values that correspond to the superstable orbits are  $\mu_2 = 3.9057$  and  $\mu_3 = 3.9903$ .  $\square$

Referring to the numerical examples given at the end of Section 1.3 where  $\mu < \mu_a$  we observed a rapid convergence towards the 2-period orbit independent on the choice of initial value. Within a periodic window in the interval  $[\mu_a, 4]$  the dynamics may be much more complicated. Indeed, still considering the period-3 window, we have according to the Li and Yorke theorem that there are also periodic orbits of any period, although invisible to a computer. (The latter is a consequence of Singer's theorem.) If we consider an initial point which is not on the 3-periodic orbit we may see that it behaves irregularly through lots of iterations before it starts to converge, and moreover, if we change the initial point somewhat it may happen that it is necessary to perform an even larger amount of iterations before we are able to detect any convergence towards the 3-cycle. Hence, the dynamics within a periodic window in the interval  $[\mu_a, 4]$  is in general much more complex than in the case of periodic orbits in the interval  $[3, \mu_a]$  due to the presence of an (infinite) number of unstable periodic points.

By carefully scrutinizing the periodic windows one may find numerically that the sum of the widths of all the windows is roughly 10% of the length of the interval  $[\mu_a, 4]$ . In the remaining part of the interval the dynamics is chaotic. If we want to give a thorough description of chaotic orbits we may use symbolic dynamics in much of a similar way as we did in Sections 1.9 and 1.10. Here we shall give a more heuristic approach only. If  $\mu$  is not close to a periodic window, orbits are irregular and there is almost no sign of periodicity. However, if  $\mu$  is close to a window, for example, if  $\mu$  is smaller but close to  $1 + \sqrt{8}$  (the threshold value for the period-3 window) one finds that an orbit seems to consist of two parts, one part which appears to be almost 3-periodic and another irregular part where the point  $x$  may take almost any value in  $(0, 1)$ . The almost 3-periodic part of the orbit is established when the orbit becomes close to the diagonal line  $x_{t+1} = x_t$ . Then, since  $\mu$  is close to  $1 + \sqrt{8}$  the orbit may stay close to the diagonal for several iterations before it moves away. Therefore, a typical orbit close to a periodic window consists of an irregular part which after a finite number of iterations becomes almost periodic and again turns irregular in a repeating fashion. For further reading on this topic we refer to Nagashima and Baba (1999), Thunberg (2001), and Jost (2005). We also recommend the books by Iooss (1979), Bergé et al. (1984), Barnsley (1988), Devaney (1989), Saber et al. (1998), and Iooss and Adelmeyer (1999).

## Part II

# $n$ -dimensional maps

$$f : \mathbb{R}^n \rightarrow \mathbb{R}^n \quad \mathbf{x} \rightarrow f(\mathbf{x})$$

## 2.1 Higher order difference equations

Consider the second order difference equation

$$x_{t+2} + a_t x_{t+1} + b_t x_t = f(t) \quad (2.1.1)$$

If  $f(t) \neq 0$ , (2.1.1) is called a nonhomogeneous difference equation. If  $f(t) = 0$ , that is

$$x_{t+2} + a_t x_{t+1} + b_t x_t = 0 \quad (2.1.2)$$

we have the associated homogeneous equation.

**Theorem 2.1.1.** The homogeneous equation (2.1.2) has the general solution

$$x_t = C_1 u_t + C_2 v_t$$

where  $u_t$  and  $v_t$  are two linear independent solutions and  $C_1, C_2$  arbitrary constants.

**Proof.** Let  $x_t = C_1 u_t + C_2 v_t$ . Then  $x_{t+1} = C_1 u_{t+1} + C_2 v_{t+1}$  and  $x_{t+2} = C_1 u_{t+2} + C_2 v_{t+2}$  and if we substitute into (2.1.2) we obtain

$$C_1(u_{t+2} + a_t u_{t+1} + b_t u_t) + C_2(v_{t+2} + a_t v_{t+1} + b_t v_t) = 0$$

which clearly is correct since  $u_t$  and  $v_t$  are linear independent solutions. □

Regarding (2.1.1) we obviously have:

**Theorem 2.1.2.** The nonhomogeneous equation (2.1.1) has the general solution

$$x_t = C_1 u_t + C_2 v_t + u_t^*$$

where  $C_1 u_t + C_2 v_t$  is the general solution of the associated homogeneous equation (2.1.2) and  $u_t^*$  is any particular solution of (2.1.1).

Just as in case of differential equations there is no general method of how to find two linear independent solutions of a second order difference equation. However, if the coefficients  $a_t$  and  $b_t$  are constants then it is possible.

Indeed, consider

$$x_{t+2} + a x_{t+1} + b x_t = 0 \quad (2.1.3)$$

where  $a$  and  $b$  are constants. Suppose that there exists a solution of the form  $x_t = m^t$  where  $m \neq 0$ . Then  $x_{t+1} = m^{t+1} = mm^t$  and  $x_{t+2} = m^2m^t$  so (2.1.3) may be expressed as

$$(m^2 + am + b)m^t = 0$$

which again implies that

$$m^2 + am + b = 0 \tag{2.1.4}$$

(2.1.4) is called the characteristic equation and its solution is easily found to be

$$m_{1,2} = -\frac{a}{2} \pm \sqrt{\frac{a^2}{4} - b} \tag{2.1.5}$$

Now we have the following result regarding the solution of (2.1.3) which we state as a theorem:

**Theorem 2.1.3.**

- 1) If  $(a^2/4) - b > 0$ , the characteristic equation have two real solutions  $m_1$  and  $m_2$ . Moreover,  $m_1^t$  and  $m_2^t$  are linear independent so according to Theorem 2.1.1 the general solution of (2.1.3) is

$$x_t = C_1m_1^t + C_2m_2^t \quad \text{where} \quad m_{1,2} = -\frac{a}{2} \pm \sqrt{\frac{a^2}{4} - b}$$

- 2) The case  $(a^2/4) - b = 0$  implies that  $m = -a/2$ . Then  $m^t$  and  $tm^t$  are two linear independent solutions of (2.1.3) so the general solution becomes:

$$x_t = C_1m^t + C_2tm^t = (C_1 + C_2t)m^t \quad \text{where} \quad m = -a/2$$

(In order to see that  $tm^t$  really is a solution of (2.1.3) note that if  $a^2/4 = b$ , then (2.1.3) may be expressed as (\*)  $x_{t+2} + ax_{t+1} + (a^2/4)x_t = 0$ . Now, assuming that  $x_t = t(-a/2)^t$  we have  $x_{t+1} = -(a/2)(t+1)(-a/2)^t$ ,  $x_{t+2} = (a^2/4)(t+2)(-a/2)^t$  and by inserting into (\*) we obtain  $(a^2/4)[t+2 - 2(t+1) + t](-a/2)^t = 0$  which proves what we want.)

- 3) Finally, if  $(a^2/4) - b < 0$  we have

$$m = -\frac{a}{2} \pm \sqrt{-(b - (a^2/4))} = -\frac{a}{2} \pm \sqrt{b - (a^2/4)}i = \alpha + \beta i$$

From the theory of complex numbers we know that

$$\alpha + \beta i = r(\cos \theta + i \sin \theta)$$

where

$$r = \sqrt{\alpha^2 + \beta^2} = \sqrt{(-a/2)^2 + \sqrt{b - (a^2/4)}^2} = \sqrt{b}$$

and

$$\cos \theta = \frac{-a/2}{\sqrt{b}} \quad \sin \theta = \frac{\sqrt{b - (a^2/4)}}{\sqrt{b}}$$

which implies that

$$m^t = [r(\cos \theta + i \sin \theta)]^t = r^t(\cos \theta + i \sin \theta)^t = r^t(\cos \theta t + i \sin \theta t)$$

where we have used Moivre's formulae (cf. Exercise 2.1.2) in the last step. Since the real and imaginary parts of  $m^t$  are linear independent functions we express the general solution of (2.1.3) as

$$x_t = C_1 r^t \cos \theta t + C_2 r^t \sin \theta t$$

□

www.sylvania.com

**We do not reinvent  
the wheel we reinvent  
light.**

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM **OSRAM SYLVANIA**



Click on the ad to read more

**Example 2.1.1.** Find the general solution of the following equations:

- a)  $x_{t+2} - 7x_{t+1} + 12x_t = 0$ ,
- b)  $x_{t+2} - 6x_{t+1} + 9x_t = 0$ ,
- c)  $x_{t+2} - x_{t+1} + x_t = 0$ .

**Solutions:**

- a) Assuming  $x_t = m^t$  the characteristic equation becomes  $m^2 - 7m + 12 = 0 \Leftrightarrow m_1 = 4$ ,  
 $m_2 = 3$  so according to Theorem 2.1.3 the general solution is  $x_t = C_1 \cdot 4^t + C_2 \cdot 3^t$ .
- b) The characteristic equation is  $m^2 - 6m + 9 = 0 \Leftrightarrow m_1 = m_2 = 3$ . Thus  
 $x_t = C_1 \cdot 3^t + C_2 t \cdot 3^t = (C_1 + C_2 t)3^t$ .
- c) The characteristic equation becomes  $m^2 - m + 1 = 0 \Leftrightarrow m = (1 \pm \sqrt{-3})/2 = \frac{1}{2} \pm \frac{1}{2}\sqrt{3}i$ .

Further

$$r = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\sqrt{3}\right)^2} = 1$$

$$\cos \theta = \frac{\frac{1}{2}}{1} = \frac{1}{2} \quad \sin \theta = \frac{\frac{1}{2}\sqrt{3}}{1} = \frac{1}{2}\sqrt{3} \Rightarrow \theta = \frac{\pi}{3}$$

Thus

$$x_t = C_1 1^t \cos \frac{\pi}{3}t + C_2 1^t \sin \frac{\pi}{3}t = C_1 \cos \frac{\pi}{3}t + C_2 \sin \frac{\pi}{3}t$$

□

**Exercise 2.1.1.** Find the general solution of the homogeneous equations:

- a)  $x_{t+2} - 12x_{t+1} + 36x_t = 0$ ,
- b)  $x_{t+2} + x_t = 0$ ,
- c)  $x_{t+2} + 6x_{t+1} - 16x_t = 0$ .

□

**Exercise 2.1.2.** Prove Moivre's formulae:  $(\cos \theta + i \sin \theta)^t = \cos \theta t + i \sin \theta t$ .

(Hint: Use induction and trigonometric identities.)

□

**Definition 2.1.1.** The equation  $x_{t+2} + ax_{t+1} + bx_t = 0$  is said to be globally asymptotic stable if the solution  $x_t$  satisfies  $\lim_{t \rightarrow \infty} x_t = 0$ . □

Referring to Example 2.1.1 it is clear that none of the equations considered there are globally asymptotic stable. The solutions of the equations (a) and (b) tend to infinity as  $t \rightarrow \infty$  and the solution of (c) does not tend to zero either.

However, consider the equation  $x_{t+2} - (1/6)x_{t+1} - (1/6)x_t = 0$ . The characteristic equation is  $m^2 - (1/6)m - (1/6)m = 0 \Leftrightarrow m_1 = 1/2, m_2 = -(1/3)$  so the general solution becomes  $x_t = C_1(1/2)^t + C_2(-1/3)^t$ .

Here, we obviously have  $\lim_{t \rightarrow \infty} x_t = 0$  so according to Definition 2.1.1 the equation  $x_{t+2} + (1/6)x_{t+1} - (1/6)x_t = 0$  is globally asymptotic stable.

**Theorem 2.1.4.** The equation  $x_{t+2} + ax_{t+1} + bx_t = 0$  with associated characteristic equation  $m^2 + am + b = 0$  is globally asymptotic stable if and only if all the roots of the characteristic equation have moduli strictly less than 1. □

**Proof.** Referring to Theorem 2.1.3, the cases (1) and (3) are clear (remember  $|m| = r$  in (3)).

Considering (2): If  $|m| < 1$

$$\lim_{t \rightarrow \infty} tm^t = \lim_{t \rightarrow \infty} \frac{t}{s^t}$$

where  $s = 1/|m|$  and  $s > 1$ . Then by L'hospital's rule

$$\lim_{t \rightarrow \infty} \frac{t}{s^t} = \lim_{t \rightarrow \infty} \frac{1}{s^t \ln s} \rightarrow 0$$

and the results of Theorem 2.1.4 follows.

As we shall see later on, Theorem 2.1.4 will be useful for us when we discuss stability of nonlinear systems.

—

We close this section by considering the nonhomogeneous equation

$$x_{t+2} + ax_{t+1} + bx_t = f(t) \tag{2.1.6}$$



According to Theorem 2.1.2 the general solution of (2.1.6) is the sum of the general solution of the homogeneous equation (2.1.3) and a particular solution  $u_t^*$  of (2.1.6).

If  $f(t)$  is a polynomial, say  $f(t) = 2t^2 + 4t$  it is natural to assume a particular solution of the form  $u_t^* = At^2 + Bt + C$ .

If  $f(t)$  is a trigonometric function, for example  $f(t) = \cos ut$  we assume that  $u_t^* = A \cos ut + B \sin ut$ .

If  $f_t = c^t$ , assume  $u_t^* = Ac^t$  (but see the comment following (2.1.7)).

**Example 2.1.2.** Solve the following equations:

- a)  $x_{t+2} + x_{t+1} + 2x_t = t^2$ ,
- b)  $x_{t+2} - 2x_{t+1} + x_t = 2 \sin(\pi/2)t$ ,



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more

**Solutions:**

a) The characteristic equation of the homogeneous equation becomes

$m^2 - m - 2 = 0 \Leftrightarrow m_1 = 2$  and  $m_2 = -1$  so the general solution of the homogeneous equation is  $x_t = C_1 \cdot 2^t + C_2(-1)^t$ . Assume  $u_t^* = At^2 + Bt + C$ . Then

$u_{t+1}^* = A(t+1)^2 + B(t+1) + C$ ,  $u_{t+2}^* = A(t+2)^2 + B(t+2) + C$  which inserted into the original equation gives

$$A(t+2)^2 + B(t+2) + C - [A(t+1)^2 + B(t+1) + C] - 2[At^2 + Bt + C] = t^2$$

$$\Leftrightarrow$$

$$-2At^2 + (2A - 2B)t + (3A + B - 2C) = t^2 + 0t + 0$$

and by equating terms of equal powers of  $t$  we have (1)  $-2A = 1$ , (2)  $2A - 2B = 0$ , and (3)

$3A + B - 2C = 0$  from which we easily obtain  $A = -1/2$ ,  $B = -1/2$  and  $C = -1$ . Thus

$u_t^* = -(1/2)t^2 - (1/2)t - 1$  and the general solution is  $x_t = C_1 2^t + C_2(-1)^t - (1/2)t^2 - (1/2)t - 1$ .

b) The solution of the characteristic equation becomes  $m_1 = m_2 = 1 \Rightarrow$  homogeneous

solution  $(C_1 + C_2 t)1^t = C_1 + C_2 t$ . Assume  $u_t^* = A \cos(\pi/2)t + B \sin(\pi/2)t$ . Then,

$u_{t+1}^* = A \cos[(\pi/2)(t+1)] + B \sin[(\pi/2)(t+1)] = A[\cos(\pi/2)t \cos(\pi/2) - \sin(\pi/2)t \sin(\pi/2)] + B[\sin(\pi/2)t \cos(\pi/2) + \cos(\pi/2)t \sin(\pi/2)] = -A \sin(\pi/2)t + B \cos(\pi/2)t$ . In the same way,  $u_{t+2}^* = -A \cos(\pi/2)t - B \sin(\pi/2)t$  so after inserting

$u_{t+2}^*$ ,  $u_{t+1}^*$  and  $u_t^*$  into the original equation we arrive at

$$-2B \cos \frac{\pi}{2}t + 2A \sin \frac{\pi}{2}t = 0 \cos \frac{\pi}{2}t + 2 \sin \frac{\pi}{2}t$$

Thus  $-2B = 0$  and  $2A = 2 \Leftrightarrow A = 1$  and  $B = 0$  so  $u_t^* = \cos(\pi/2)t$ . Hence, the general

solution is  $x_t = C_1 + C_2 t + \cos(\pi/2)t$ . □

Finally, if  $x_{t+2} + ax_{t+1} + bx_t = c^t$  we assume a particular solution of the form  $u_t^* = Ac^t$ . Then

$u_{t+1}^* = Acc^t$  and  $u_{t+2}^* = Ac^2c^t$  which inserted into the original equation yields

$$A(c^2 + ac + b)c^t = c^t$$

Thus, whenever  $c^2 + ac + b \neq 0$  the particular solution becomes

$$u^* = \frac{1}{c^2 + ac + b} c^t \quad (2.1.7)$$

Note, however, that if  $c$  is a simple root of the characteristic equation, i.e.  $c^2 + ac + b = 0$ , then we try a solution of the form  $u_t^* = Btc^t$  and if  $c$  is a double root, assume  $u_t^* = Dt^2c^t$ .

**Example 2.1.3.** Solve the equations:

- a)  $x_{t+2} - 4x_t = 3^t$ ,
- b)  $x_{t+2} - 4x_t = 2^t$ ,

**Solutions:**

a) The characteristic equation is  $m^2 - 4 = 0 \Leftrightarrow m_1 = 2, m_2 = -2$  thus the homogeneous solution is  $C_1 2^t + C_2 (-2)^t$ . Since 3 is not a root of  $m^2 - 4 = 0$  we have directly from (2.1.7) that  $u_t^* = (1/5)3^t$  so the general solution becomes  $x_t = C_1 2^t + C_2 (-2)^t + (1/5)3^t$ .

b) The homogeneous solution is of course  $C_1 2^t + C_2 (-2)^t$  but since 2 is a simple root of  $m^2 - 4 = 0$  we try a particular solution of the form  $u_t^* = Bt2^t$ . Then  $u_{t+2}^* = 4B(t+2)2^t$  and by inserting into the original equation we arrive at

$$4B(t+2)2^t - 4Bt \cdot 2^t = 2^t$$

which gives  $B = 1/8$ . Thus  $x_t = C_1 2^t + C_2 (-2)^t + (1/8)t \cdot 2^t$ . □

**Exercise 2.1.3.** Solve the problems:

- a)  $x_{t+2} + 2x_{t+1} - 3x_t = 2t + 5$ ,
- b)  $x_{t+2} - 10x_{t+1} + 25x_t = 5^t$ ,
- c)  $x_{t+2} - x_{t+1} + x_t = 2^t$ ,
- d)  $x_{t+2} + 9x_t = 2^t$ ,
- e)  $x_{t+2} - 5x_{t+1} - 6x_t = t \cdot 2^t$ .

(Hint: Assume a particular solution of the form  $(At + B) \cdot 2^t$ .) □

In the examples and exercises presented above we found a particular solution  $u_t^*$  of the nonhomogeneous equation in a way which at best may be called heuristic. We shall now focus on a general method (sometimes referred to as variation of parameters) which enables us to find  $u_t^*$  of any nonhomogeneous equation provided the general solution of the associated homogeneous equation is known.

**Theorem 2.1.5 (Variation of parameters).** Let  $x_{1,t}$  and  $x_{2,t}$  be two linear independent solutions of (2.1.3) and let

$$w_t = \begin{vmatrix} x_{1,t} & x_{2,t} \\ x_{1,t-1} & x_{2,t-1} \end{vmatrix}$$

Then a particular solution  $u_t^*$  of the nonhomogeneous equation

$$x_{t+2} + ax_{t+1} + bx_t = f_t$$

may be calculated through

$$u_t^* = \sum_{m=0}^t \frac{\begin{vmatrix} x_{1,t} & x_{2,t} \\ x_{1,m-1} & x_{2,m-1} \end{vmatrix}}{w_m} f_{m-2} \quad t \geq 0$$


□

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more

**Proof.** The elements  $u_t^*$  must be linear functions of the preceding elements of the sequence  $\{f_t\}$ .

Hence,

$$u_t^* = \sum_{m=0}^t d_{t,m} f_{m-2}$$

which inserted into the nonhomogeneous equation gives

$$\sum_{m=0}^t (d_{t+2,m} + ad_{t+1,m} + bd_{t,m}) f_{m-2} + (d_{t+2,t+1} + ad_{t+1,t+1}) f_{t-1} + d_{t+2,t+2} f_t = f_t$$

The equation above must hold for any  $t > 0$ . Consequently, for each  $m$ , the coefficients of  $f_m$  on both sides of the equation must be equal. Therefore,

$$\begin{aligned} d_{t+2,m} + ad_{t+1,m} + bd_{t,m} &= 0 & t > m - 1 \\ d_{t+2,t+1} + ad_{t+1,t+1} &= 0 \\ d_{t+2,t+2} &= 1 \end{aligned}$$

The first of the three equations above expresses that the sequence  $\{d_{t,m}\}$  is a solution of the homogeneous equation in case of  $t > m - 1$ . Moreover, by imposing the initial condition  $d_{m,m-1} = 0$  the second equation may be replaced by the first if  $t \geq m - 1$  and we have the initial conditions  $d_{m,m-1} = 0$ ,  $d_{m,m} = 1$ . Now, since  $x_{1,t}$  and  $x_{2,t}$  are two linear independent solutions of the homogeneous equations there are constants  $c$  such that

$$d_{t,m} = c_{1,m} x_{1,t} + c_{2,m} x_{2,t}$$

and the initial conditions are satisfied whenever

$$\begin{aligned} c_{1,m} x_{1,m} + c_{2,m} x_{2,m} &= 1 \\ c_{1,m} x_{1,m-1} + c_{2,m} x_{2,m-1} &= 0 \end{aligned}$$

from which we easily obtain

$$c_{1,m} = \frac{x_{2,m-1}}{w_m} \quad c_{2,m} = -\frac{x_{1,m-1}}{w_m}$$

Consequently,

$$d_{t,m} = \frac{x_{1,t} x_{2,m-1} - x_{2,t} x_{1,m-1}}{w_m}$$

and the formulae in the theorem follows. □

**Example 2.1.4.** Use Theorem 2.1.5 and find a particular solution of

$$x_{t+2} - 5x_{t+1} + 6x_t = 2^t$$

**Solution.** Clearly, two linear independent solutions of the associated homogeneous equation are  $x_{1,t} = 2^t$  and  $x_{2,t} = 3^t$ . Moreover,

$$w_t = \begin{vmatrix} 2^t & 3^t \\ 2^{t-1} & 3^{t-1} \end{vmatrix} = -6^{t-1} \quad \begin{vmatrix} x_{1,t} & x_{2,t} \\ x_{1,m-1} & x_{2,m-1} \end{vmatrix} = 6^{m-1} [2^{t-(m-1)} - 3^{t-(m-1)}]$$

Thus

$$\begin{aligned} u_t^* &= \sum_{m=0}^t \frac{6^{m-1} [2^{t-(m-1)} - 3^{t-(m-1)}]}{-6^{m-1}} 2^{m-2} \\ &= - \sum_{m=0}^t \left[ 2^{t-1} - \frac{1}{4} 3^{t+1} \left( \frac{2}{3} \right)^m \right] = - \left\{ (t+1) 2^{t-1} - \frac{3}{4} (3^{t+1} - 2^{t+1}) \right\} \\ &= \frac{9}{4} 3^t - \frac{(t+4)}{2} 2^t \end{aligned}$$

Note that the particular solution found here is not the same as  $u_t^* = -\frac{1}{2}t \cdot 2^t$  which would be the result by use of a heuristic method (see Example 2.1.3b). However, the general solutions match. Indeed,

$$\begin{aligned} x_t &= C_1 x_{1,t} + C_2 x_{2,t} + u_t^* = C_1 \cdot 2^t + C_2 \cdot 3^t + \frac{9}{4} 3^t - \left( \frac{t+4}{2} \right) 2^t \\ &= (C_1 - 2) 2^t + \left( C_2 + \frac{9}{4} \right) 3^t - \frac{1}{2} t \cdot 2^t = D_1 \cdot 2^t + D_2 \cdot 3^t - \frac{1}{2} t \cdot 2^t \end{aligned}$$

which is in accordance with the heuristic method. □

**Example 2.1.5.** Find the general solution of

$$x_{t+2} - 5x_{t+1} + 6x_t = \ln(t+3)$$

**Solution.** By use of the findings from the previous example:

$$\begin{aligned} u_t^* &= - \sum_{m=0}^t [2^{t-(m-1)} - 3^{t-(m-1)}] \ln(m+1) \\ &= 3^{t+1} \sum_{m=0}^t 3^{-m} \ln(m+1) - 2^{t+1} \sum_{m=0}^t 2^{-m} \ln(m+1) \end{aligned}$$

Hence, the general solution becomes

$$x_t = C_1 \cdot 2^t + C_2 \cdot 3^t + 3^{t+1} \sum_{m=0}^t 3^{-m} \ln(m+1) - 2^{t+1} \sum_{m=0}^t 2^{-m} \ln(m+1)$$

Note that the solution above (in contrast to all our previous examples and exercises) contains sums which may not be expressed in any simple forms. However, in a somewhat more cumbersome way, we have obtained the general solution for any  $t \geq 0$ . Moreover, the constants  $C_1$  and  $C_2$  may be determined in the usual way if we know the initial conditions. Indeed, assuming  $x_0 = 0$  and  $x_1 = 1$  we arrive at the equations (A)  $C_1 + C_2 = 0$  and (B)  $2C_1 + 3C_2 + \ln 2 = 1$  from which we obtain  $C_1 = \ln 2 - 1$  and  $C_2 = 1 - \ln 2$ .  $\square$

**Exercise 2.1.4.** Use Theorem 2.1.5 and find the general solution of the equations

- a)  $x_{t+2} - 7x_{t+1} + 10x_t = 5^t$
- b)  $x_{t+2} - (a + b)x_{t+1} + abx_t = a^t$

(Hint: distinguish between the cases  $a \neq b$  and  $a = b$ .)  $\square$

**Exercise 2.1.5.** Consider the equation  $x_{t+2} = x_{t+1} + x_t$  with initial conditions  $x_0 = 0, x_1 = 1$ .

- a) Solve the equation.
- b) Use a) and induction to prove that  $x_t \cdot x_{t+2} - x_{t+1}^2 = (-1)^{t+1}, t = 0, 1, 2, \dots$   $\square$

Cynthia | AXA Graduate

**AXA Global Graduate Program**

Find out more and apply

redefining / standards AXA





Let us now turn to equations of order  $n$ , i.e. equations of the form

$$x_{t+n} + a_1(t)x_{t+n-1} + a_2(t)x_{t+n-2} + \cdots + a_{n-1}(t)x_{t+1} + a_n(t)x_t = f(t) \quad (2.1.8)$$

In the homogeneous case we have the following result:

**Theorem 2.1.6.** Assuming  $a_n(t) \neq 0$ , the general solution of

$$x_{t+n} + a_1(t)x_{t+n-1} + \cdots + a_n(t)x_t = 0 \quad (2.1.9)$$

is  $x_t = C_1u_{1,t} + \cdots + C_nu_{n,t}$  where  $u_{1,t} \dots u_{n,t}$  are linear independent solutions of the equation and  $C_1 \dots C_n$  arbitrary constants.  $\square$

**Proof.** Easy extension of the proof of Theorem 2.1.1. We leave the details to the reader.  $\square$

Regarding the nonhomogeneous equation (2.1.8) we have

**Theorem 2.1.7.** The solution of the nonhomogeneous equation (2.1.8) is

$$x_t = C_1u_{1,t} + \cdots + C_nu_{n,t} + u_t^*$$

where  $u_t^*$  is a particular solution of (2.1.8) and  $C_1u_{1,t} + \cdots + C_nu_{n,t}$  is the general solution of (2.1.9).  $\square$

If  $a_1(t) = a_1, \dots, a_n(t) = a_n$  constants we arrive at

$$x_{t+n} + a_1x_{t+n-1} + \cdots + a_nx_t = f(t) \quad (2.1.10)$$

and as in the second order case we may assume a solution  $x_t = m^t$  of the homogeneous equation. This yields the  $n$ -th order characteristic equation

$$m^n + a_1m^{n-1} + \cdots + a_{n-1}m + a_n = 0 \quad (2.1.11)$$

Appealing to the fundamental theorem of algebra we know that (2.1.11) has  $n$  roots. If a root is real with multiplicity 1 or complex we form linear independent solutions in exactly the same way as explained in Theorem 2.1.3. In case of real roots with multiplicity  $p$ , linear independent solutions are  $m^t, tm^t, \dots, t^{p-1}m^t$ .

**Example 2.1.5.** Solve the equations:

- a)  $x_{t+3} - 2x_{t+2} + x_{t+1} - 2x_t = 2t - 4$ ,  
 b)  $x_{t+3} - 6x_{t+2} + 12x_{t+1} - 8x_t = 0$ .

**Solutions:**

- a) The characteristic equation is  $m^3 - 2m^2 + m - 2 = 0$ . Clearly,  $m_1 = 2$  is a solution and  $m^3 - 2m^2 + m - 2 = (m - 2)(m^2 + 1) = 0$ . Hence the other roots are complex,  $m_{2,3} = \pm i$ . Following Theorem 2.1.3  $r = \sqrt{0^2 + 1^2} = 1$ ,  $\cos \theta = 0/1 = 0$ ,  $\sin \theta = 1/1 = 1 \Rightarrow \theta = \pi/2$  which implies the homogeneous solution  $C_1 \cdot 2^t + C_2 \cos(\pi/2)t + C_3 \sin(\pi/2)t$ . Assuming a particular solution  $u_t^* = At + B$  we find after inserting into the original equation,  $-2At - 2B = 2t - 4$  so  $A = -1$  and  $B = 2$ . Consequently, according to Theorem 2.1.7, the general solution is  $x_t = C_1 \cdot 2^t + C_2 \cos(\pi/2)t + C_3 \sin(\pi/2)t - t + 2$ .

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



Real work  
International opportunities  
Three work placements



**Month 16**  
I was a construction  
supervisor in  
the North Sea  
advising and  
helping foremen  
solve problems





b) The characteristic equation becomes  $m^3 - 6m^2 + 12m - 8 = 0 \Leftrightarrow (m - 2)^3 = 0$ .

Hence, there is only one root,  $m = 2$ , with multiplicity 3. Consequently,

$$x_t = C_1 \cdot 2^t + C_2 t \cdot 2^t + C_3 t^2 \cdot 2^t. \quad \square$$

**Exercise 2.1.6.** Find the general solution of the equations:

a)  $x_{t+3} - 2x_{t+2} - 5x_{t+1} + 6x_t = 0$     c)  $x_{t+1} - 2x_t = 1 + t^2$

b)  $x_{t+4} - x_t = 2^t$     d)  $x_{t+1} - 2x_t = 2^t + 3^t$

□

**Definition 2.1.2.** The equation  $x_{t+n} + a_1 x_{t+n-1} + \dots + a_n x_t = 0$  is said to be globally asymptotic stable if the solution  $x_t$  satisfies  $\lim_{t \rightarrow \infty} x_t = 0$ . □

**Theorem 2.1.8.** The equation  $x_{t+n} + a_1 x_{t+n-1} + \dots + a_n x_t = 0$  is globally asymptotic stable if all solutions of the characteristic equation (2.1.11) have moduli less than 1. □

It may be a difficult task to decide whether all roots of a given polynomial equation have moduli less than unity or not. However, there are methods and one of the most frequently used is the Jury criteria which we now describe.

Let

$$P(x) = x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_n \quad (2.1.12)$$

be a polynomial with real coefficients  $a_1 \dots a_n$ . Define

$$b_n = 1 - a_n^2, \quad b_{n-1} = a_1 - a_n a_{n-1}, \dots, \quad b_{n-j} = a_j - a_n a_{n-j}, \quad b_1 = a_{n-1} - a_n a_1$$

$$c_n = b_n^2 - b_1^2, \quad c_{n-1} = b_n b_{n-1} - b_1 b_2, \dots, \quad c_{n-j} = b_n b_{n-j} - b_1 b_{j+1}, \quad c_2 = b_n b_2 - b_1 b_{n-1}$$

$$d_n = c_n^2 - c_2^2, \dots, \quad d_{n-j} = c_n c_{n-j} - c_2 c_{j+2} \dots, \quad d_3 = c_n c_3 - c_2 c_{n-1}$$

and proceed in this way until we have only three elements of the type

$$w_n = v_n^2 - v_{n-3}^2, \quad w_{n-1} = v_n v_{n-1} - v_{n-3} v_{n-2}, \quad w_{n-2} = v_n v_{n-2} - v_{n-3} v_{n-1}$$

**Theorem 2.1.9 (The Jury criteria).** All roots of the polynomial equation  $P(x) = 0$  where  $P(x)$  is defined through (2.1.12) have moduli less than 1 provided:

$$P(1) > 0 \quad (-1)^n P(-1) > 0$$

$$|a_n| < 1, |b_n| > |b_1|, |c_n| > |c_2|, |d_n| > |d_3|, \dots, |w_n| > |w_{n-2}|. \quad \square$$

**Remark 2.1.1.** Instead of saying that all roots have moduli less than 1, an alternative formulation is to say that all roots are located inside the unit circle in the complex plane.  $\square$

Regarding the second order equation

$$x^2 + a_1x + a_2 = 0 \quad (2.1.13)$$

the Jury criteria become

$$\begin{aligned} 1 + a_1 + a_2 &> 0 \\ 1 - a_1 + a_2 &> 0 \\ 1 - |a_2| &> 0 \end{aligned} \quad (2.1.14)$$

If we have a polynomial equation of order 3

$$x^3 + a_1x^2 + a_2x + a_3 = 0 \quad (2.15)$$

the Jury criteria may be cast in the form

$$\begin{aligned} 1 + a_1 + a_2 + a_3 &> 0 \\ 1 - a_1 + a_2 - a_3 &> 0 \\ 1 - |a_3| &> 0 \\ |1 - a_3^2| - |a_2 - a_3a_1| &> 0 \end{aligned} \quad (2.16)$$

Evidently, the higher the order, the more complicated are the Jury criteria. Therefore, unless the coefficients are very simple or on a special form the method does not work if the order of the polynomial becomes large.

Later, when we shall focus on stability problems of nonlinear maps (which often leads to a study of polynomial equations), we will also face the fact that the coefficients  $a_1 \dots a_n$  do not consist of numbers only but a mixture of numbers and parameters. In such cases, even (2.1.16) may be difficult to apply.

However, let us give one simple example of how the Jury criteria works.

**Example 2.1.6.** Show that  $x_{t+3} - (2/3)x_{t+2} + (1/4)x_{t+1} - (1/6)x_t = 0$  is globally asymptotic stable.

**Solution:** According to Theorem 2.1.8 we must show that the roots of the associated characteristic equation  $m^3 - (2/3)m^2 + (1/4)m - (1/6) = 0$  are located inside the unit circle. Defining  $a_1 = -(2/3)$ ,  $a_2 = 1/4$ ,  $a_3 = -(1/6)$  the four left-hand sides of (2.1.16) become  $1/12$ ,  $25/12$ ,  $5/6$  and  $5/6$ , respectively. Consequently, all the roots are located inside the unit circle so the difference equation is globally asymptotic stable.  $\square$

Another theorem (from complex function theory) that may be useful and which applies not only to polynomial equations is Rouché's theorem. (In the theorem below,  $z = \alpha + \beta i$  is a complex number.)

**Theorem 2.1.10** (Rouché's theorem). If  $f(z)$  and  $g(z)$  are analytic inside and on a simple closed curve  $C$  and if  $|g(z)| < |f(z)|$  on  $C$  then  $f(z) + g(z)$  and  $f(z)$  and the same number of zeros inside  $C$ .  $\square$

**Remark 2.1.2.** If we take the simple closed curve  $C$  to be the unit circle  $|z| = 1$ , then we may use Theorem 2.1.10 in order to decide if all the roots of a given equation have moduli less than one or not.  $\square$

**Example 2.1.7.** Suppose that  $a > e$  and show that the equation  $az^n - e^z = 0$  has  $n$  roots located inside the unit circle  $|z| = 1$ .

**Solution:** Define  $f(z) = az^n$ ,  $g(z) = -e^z$  and consider  $f(z) + g(z) = 0$ . Clearly, the equation  $f(z) = 0$  has  $n$  roots located inside the unit circle. On the boundary of the unit circle we have  $|g(z)| = |-e^z| \leq e < a = |f(z)|$ . Thus, according to Theorem 2.1.10,  $f(z)$  and  $f(z) + g(z)$  have the same number of zeros inside the unit circle, i.e.  $n$  zeros.  $\square$

## 2.2 Systems of linear difference equations. Linear maps from $\mathbb{R}^n$ to $\mathbb{R}^n$

In this section our purpose is to analyse linear systems. There are several alternatives when one tries to find the general solution of such systems. One possible method is to transform a system into one higher order equation and use the theory that we developed in the previous section. Other methods are based upon topics from linear algebra, and of particular relevance is the theory of eigenvalues and eigenvectors. Later when we turn to nonlinear systems and stability problems it will be useful for us to have a broad knowledge of linear systems so therefore we shall deal with several possible solution methods in this section.

Consider the system

$$\begin{aligned}x_{1,t+1} &= a_{11}x_{1,t} + a_{12}x_{2,t} + \cdots + a_{1n}x_{n,t} + b_1(t) \\x_{2,t+1} &= a_{21}x_{1,t} + a_{22}x_{2,t} + \cdots + a_{2n}x_{n,t} + b_2(t) \\&\vdots \\x_{n,t+1} &= a_{n1}x_{1,t} + a_{n2}x_{2,t} + \cdots + a_{nn}x_{n,t} + b_n(t)\end{aligned}\tag{2.2.1}$$

Here, all coefficients  $a_{11} \dots a_{nn}$  are constants and if  $b_i(t) = 0$  for all  $1 \leq i \leq n$  we call (2.2.1) a linear autonomous system.

**ie** business school

**93%**  
OF MIM STUDENTS ARE  
WORKING IN THEIR SECTOR 3 MONTHS  
FOLLOWING GRADUATION

**MASTER IN MANAGEMENT**

- STUDY IN THE CENTER OF MADRID AND TAKE ADVANTAGE OF THE UNIQUE OPPORTUNITIES THAT THE CAPITAL OF SPAIN OFFERS
- PROPEL YOUR EDUCATION BY EARNING A DOUBLE DEGREE THAT BEST SUITS YOUR PROFESSIONAL GOALS
- STUDY A SEMESTER ABROAD AND BECOME A GLOBAL CITIZEN WITH THE BEYOND BORDERS EXPERIENCE

Length: 10 MONTHS  
Av. Experience: 1 YEAR  
Language: ENGLISH / SPANISH  
Format: FULL-TIME  
Intakes: SEPT / FEB

**5 SPECIALIZATIONS**  
PERSONALIZE YOUR PROGRAM

**#10 WORLDWIDE**  
MASTER IN MANAGEMENT  
FINANCIAL TIMES

**55 NATIONALITIES**  
IN CLASS

[www.ie.edu/master-management](http://www.ie.edu/master-management) | [mim.admissions@ie.edu](mailto:mim.admissions@ie.edu) | [f](#) [t](#) [in](#) Follow us on IE MIM Experience



It is often convenient to express (2.2.1) in terms of vectors and matrices. Indeed, let  $\mathbf{x} = (x_1, \dots, x_n)^T$ ,  $\mathbf{b} = (b_1, \dots, b_n)^T$  and

$$A = \begin{pmatrix} a_{11} & \cdots & a_{1n} \\ a_{21} & \cdots & a_{2n} \\ \vdots & & \vdots \\ a_{n1} & \cdots & a_{nn} \end{pmatrix} \quad (2.2.2)$$

Then, (2.2.1) may be written as

$$\mathbf{x}_{t+1} = A\mathbf{x}_t + \mathbf{b}_t \quad (2.2.3)$$

or in map notation

$$\mathbf{x} \rightarrow A\mathbf{x} + \mathbf{b} \quad (2.2.4)$$

First, let us show how one may solve a system by use of the theory from the previous section.

**Example 2.2.1.** Solve the system

$$(1) \quad x_{t+1} = 2y_t + t$$

$$(2) \quad y_{t+1} = x_t + y_t$$

Replacing  $t$  by  $t + 1$  in (1) gives

$$x_{t+2} = 2y_{t+1} + t + 1 \stackrel{(2)}{=} 2(x_t + y_t) + t + 1 = 2x_t + 2y_t + t + 1$$

Further, from (1):  $2y_t = x_{t+1} - t$ . Hence

$$x_{t+2} - x_{t+1} - 2x_t = 1$$

Thus, we have transformed a system of two first order equations into one second order equation, and by use of the theory from the previous section the general solution of the latter equation is easily found to be

$$x_t = C_1 \cdot 2^t + C_2(-1)^t - 1/2$$



$y_t$  may be obtained from (1):

$$\begin{aligned} y_t &= \frac{1}{2}(x_{t+1} - t) = \frac{1}{2} \left( C_1 2^{t+1} + C_2 (-1)^{t+1} - \frac{1}{2} - t \right) \\ &= C_1 2^t - \frac{1}{2} C_2 (-1)^t - \frac{1}{2} t - \frac{1}{4} \end{aligned}$$

The constants  $C_1$  and  $C_2$  may be determined if we know the initial values  $x_0$  and  $y_0$ . For example, if  $x_0 = y_0 = 1$  we have from the general solution above that

$$\begin{aligned} 1 &= C_1 + C_2 - 1/2 \\ 1 &= C_1 - \frac{1}{2} C_2 - 1/4 \end{aligned}$$

which implies that  $C_1 = 4/3$  and  $C_2 = 1/6$  so the solution becomes

$$x_t = \frac{4}{3} \cdot 2^t + \frac{1}{6} (-1)^t - \frac{1}{2} \quad y_t = \frac{4}{3} \cdot 2^t + \frac{1}{12} (-1)^t - \frac{1}{2} t - \frac{1}{4}$$

□

**Exercise 2.2.1.** Find the general solution of the systems

$$\begin{aligned} \text{a) } x_{t+1} &= 2y_t + t & \text{b) } x_{t+1} &= x_t + 2y_t \\ y_{t+1} &= -x_t + 3y_t & y_{t+1} &= 3x_t \end{aligned}$$

□

Another way to find the solution of a system is to use the matrix formulation (2.2.3). Indeed, suppose that the initial vector  $\mathbf{x}_0$  is known. Then:

$$\begin{aligned} \mathbf{x}_1 &= A\mathbf{x}_0 + \mathbf{b}(0) \\ \mathbf{x}_2 &= A\mathbf{x}_1 + \mathbf{b}(1) = A(A\mathbf{x}_0 + \mathbf{b}(0)) + \mathbf{b}(1) = A^2\mathbf{x}_0 + A\mathbf{b}(0) + \mathbf{b}(1) \end{aligned}$$

and by induction (we leave the details to the reader)

$$\mathbf{x}_t = A^t \mathbf{x}_0 + A^{t-1} \mathbf{b}(0) + A^{t-2} \mathbf{b}(1) + \cdots + \mathbf{b}(t-1) \quad (2.2.5)$$

In the important special case  $\mathbf{b} = \mathbf{0}$  we have the result:

$$\mathbf{x}_{t+1} = A\mathbf{x}_t \Leftrightarrow \mathbf{x}_t = A^t \mathbf{x}_0 \quad (2.2.6)$$

where  $A^0$  is equal to the identity matrix  $I$ .

**Exercise 2.2.2.** Consider the matrix

$$A = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

- Compute  $A^2$  and  $A^3$ .
- Let  $t$  be a positive integer and use induction to find a formulae for  $A^t$ .
- Let  $\mathbf{x} = (x_1, x_2)^T$  and solve the difference equation  $\mathbf{x}_{t+1} = A\mathbf{x}_t$  where  $\mathbf{x}_0 = (a, b)^T$ .

□

Our next goal is to solve the linear system

$$\mathbf{x}_{t+1} = A\mathbf{x}_t \tag{2.2.7}$$

"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"  
Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



in terms of eigenvalues and (generalized) eigenvectors. Recall that if there exists a scalar  $\lambda$  such that  $A\mathbf{u} = \lambda\mathbf{u}$ ,  $\mathbf{u} \neq \mathbf{0}$ ,  $\lambda$  is said to be an eigenvalue of  $A$ , and  $\mathbf{u}$  is called the associated eigenvector. Moreover, we call a vector  $\mathbf{v}$  satisfying  $(A - \lambda I)\mathbf{v} = \mathbf{u}$  a generalized eigenvector of  $A$ . (Note that the definitions above imply  $(A - \lambda I)\mathbf{u} = \mathbf{0}$  and  $(A - \lambda I)^2\mathbf{v} = \mathbf{0}$ .) Thus, consider (2.2.7) and assume a solution of the form  $\mathbf{x}_t = \lambda^t\mathbf{u}$  where  $\lambda \neq 0$ . Then

$$\begin{aligned}\lambda^{t+1}\mathbf{u} - A\lambda^t\mathbf{u} &= 0 \\ \Leftrightarrow \\ (A - \lambda I)\mathbf{u} &= 0\end{aligned}\tag{2.2.8}$$

so  $\lambda$  is nothing but an eigenvalue belonging to  $A$  and  $\mathbf{u}$  is the associated eigenvector. As is well known, the eigenvalues may be computed from the relation

$$|A - \lambda I| = 0\tag{2.2.9}$$

There are two cases to consider.

(A) If the  $n \times n$  matrix  $A$  is diagonalizable over the complex numbers, then  $A$  has  $n$  distinct eigenvalues  $\lambda_1, \dots, \lambda_n$  and moreover, the associated eigenvectors  $\mathbf{u}_1, \dots, \mathbf{u}_n$  are linear independent. Consequently, the general solution of the linear system (2.2.7) may be cast in the form

$$\mathbf{x}_t = C_1\lambda_1^t\mathbf{u}_1 + C_2\lambda_2^t\mathbf{u}_2 + \dots + C_n\lambda_n^t\mathbf{u}_n\tag{2.2.10}$$

(B) If  $A$  is not diagonalizable (which may occur when  $A$  has multiple eigenvalues) we may proceed in much of the same way as in the corresponding theory for continuous systems, see Grimshaw (1990) and express the general solution in terms of eigenvalues and (generalized) eigenvectors. Suppose that  $\lambda$  is an eigenvalue with multiplicity  $m$  and let  $\mathbf{u}_1, \dots, \mathbf{u}_p$  be a basis for the eigenspace of  $\lambda$ . If  $p = m$  we are done. If  $p < m$  we seek a solution of the form  $\mathbf{x}_t = \lambda^t(\mathbf{v} + t\mathbf{u})$  where  $\mathbf{u}$  is one of the  $\mathbf{u}_i$ 's.

Then from (2.2.7) one easily obtains

$$(A - \lambda I)\mathbf{v} = \lambda\mathbf{u}\tag{2.2.11a}$$

$$(A - \lambda I)\mathbf{u} = 0\tag{2.2.11b}$$

and after multiplying (2.2.11a) with  $(A - \lambda I)$  from the left we arrive at

$$(A - \lambda I)^2 \mathbf{v} = 0 \quad (2.2.12)$$

Now suppose that we can find  $\mathbf{v}_1, \dots, \mathbf{v}_q$  such that  $\mathbf{v}_1, \dots, \mathbf{v}_q, \mathbf{u}_1, \dots, \mathbf{u}_p$  are linear independent.

Now, if  $p + q = m$  we are done. If  $p + q < m$  we continue in the same fashion by seeking a solution of the form  $\mathbf{x}_t = \lambda^t(\mathbf{w} + t\mathbf{v} + (1/2!)t^2\mathbf{u})$ . In this case (2.2.7) implies

$$(A - \lambda I)\mathbf{w} = \lambda \left( \mathbf{v} + \frac{1}{2!}\mathbf{u} \right) \quad (2.2.13a)$$

$$(A - \lambda I)\mathbf{v} = \lambda\mathbf{u} \quad (2.2.13b)$$

$$(A - \lambda I)\mathbf{u} = 0 \quad (2.2.13c)$$

which again leads to

$$(A - \lambda I)^3 \mathbf{w} = 0 \quad (2.2.14)$$

and we proceed in the same way as before. Either we are done or we keep on seeking solutions where cubic terms of  $t$  are included. Sooner or later we will obtain the necessary number of linear independent eigenvectors, cf. Meyer (2000).  $\square$

**Exercise 2.2.3.**

- a) Referring to the procedure outlined above suppose a cubic solution of the form  $\mathbf{x}_t = \lambda^t(\mathbf{y} + t\mathbf{w} + (1/2!)t^2\mathbf{v} + (1/3!)t^3\mathbf{u})$ . Use (2.2.7) and deduce the following relations:  $(A - \lambda I)\mathbf{y} = \lambda(\mathbf{w} + (1/2!)\mathbf{v} + (1/3!)\mathbf{u})$ ,  $(A - \lambda I)\mathbf{w} = \lambda(\mathbf{v} + (1/2!)\mathbf{u})$ ,  $(A - \lambda I)\mathbf{v} = \lambda\mathbf{u}$ ,  $(A - \lambda I)\mathbf{u} = \mathbf{0}$ , and moreover that  $(A - \lambda I)^4\mathbf{y} = \mathbf{0}$ .

- b) In general, assume a solution of degree  $m - 1$  on the form

$$\mathbf{x}_t = \lambda^t \sum_{i=1}^m \frac{1}{(m-i)!} t^{m-i} \mathbf{v}_i$$

and show that  $\mathbf{v}_i$  may be obtained from

$$(A - \lambda I)\mathbf{v}_1 = \mathbf{0}$$

and

$$(A - \lambda I)\mathbf{v}_{i+1} = \lambda \sum_{k=1}^i \frac{1}{(i - (k - 1))!} \mathbf{v}_k, \quad i = 1, 2, \dots, m - 1.$$

□

**Remark 2.2.1.** A complete treatment of case (B) should include a proof of linear independence of the set of eigenvectors and generalized eigenvectors. However, such a proof requires a somewhat deeper insight of linear algebra than assumed here and is therefore omitted. □

Let us now illustrate the theory presented above through three examples. In Example 2.2.2 we deal with the easiest case where the coefficient matrix  $A$  has distinct real eigenvalues. In Example 2.2.3 we consider eigenvalues with multiplicity larger than one, and finally, in Example 2.2.4, we analyse the case where the eigenvalues are complex conjugated.

Excellent Economics and Business programmes at:



university of  
 groningen




“The perfect start  
of a successful,  
international career.”

**CLICK HERE**  
to discover why both socially  
and academically the University  
of Groningen is one of the best  
places for a student to be

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)



**Example 2.2.2.** Let

$$\mathbf{x} = (x_1, x_2)^T, \quad A = \begin{pmatrix} 2 & 1 \\ -3 & 6 \end{pmatrix}$$

and solve  $\mathbf{x}_{t+1} = A\mathbf{x}_t$ .

Assuming  $\mathbf{x} = \lambda^t \mathbf{u}$  the eigenvalue equation (2.2.9) becomes

$$\begin{vmatrix} 2 - \lambda & 1 \\ -3 & 6 - \lambda \end{vmatrix} = 0 \Leftrightarrow \lambda^2 - 8\lambda + 15 = 0 \Leftrightarrow \lambda_1 = 5, \quad \lambda_2 = 3$$

The eigenvector  $\mathbf{u}_1 = (u_1, u_2)^T$  belonging to  $\lambda_1 = 5$  satisfies (cf. (2.2.8))

$$\begin{pmatrix} 2 - 5 & 1 \\ -3 & 6 - 5 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Hence, we choose  $\mathbf{u}_1 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}$ .

In the same way, the eigenvector  $\mathbf{u}_2 = (u_1, u_2)^T$  belonging to  $\lambda_2 = 3$  satisfies

$$\begin{pmatrix} -1 & 1 \\ -3 & 3 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

Thus  $\mathbf{u}_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ . Therefore, according to (2.2.10), the general solution is

$$\mathbf{x}_t = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}_t = C_1 5^t \begin{pmatrix} 1 \\ 3 \end{pmatrix} + C_2 3^t \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

□

**Example 2.2.3.** Let

$$\mathbf{x} = (x_1, x_2, x_3)^T, \quad A = \begin{pmatrix} 2 & 1 & 1 \\ 0 & 2 & 2 \\ 0 & 0 & 2 \end{pmatrix}$$

and solve  $\mathbf{x}_{t+1} = A\mathbf{x}_t$ .

Assuming  $\mathbf{x}_t = \lambda^t \mathbf{u}$ , we arrive at the eigenvalue equation

$$\begin{vmatrix} 2 - \lambda & 1 & 1 \\ 0 & 2 - \lambda & 2 \\ 0 & 0 & 2 - \lambda \end{vmatrix} = 0 \Leftrightarrow (2 - \lambda)^3 = 0$$

so we conclude that  $\lambda = 2$  is the only eigenvalue and that it has multiplicity 3. Therefore, according to (B) the general solution of the problem is

$$\mathbf{x}_t = C_1 \lambda^t \mathbf{u} + C_2 \lambda^t (\mathbf{v} + t\mathbf{u}) + C_3 \lambda^t \left( \mathbf{w} + t\mathbf{v} + \frac{1}{2} t^2 \mathbf{u} \right)$$

where  $\lambda = 2$  and  $\mathbf{u}$ ,  $\mathbf{v}$  and  $\mathbf{w}$  must be found from (2.2.13a,b,c). Let  $\mathbf{u} = (u_1, u_2, u_3)^T$ ,  $\mathbf{v} = (v_1, v_2, v_3)^T$  and  $\mathbf{w} = (w_1, w_2, w_3)^T$ . (2.2.13c) implies

$$\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \\ u_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \Leftrightarrow \begin{cases} u_2 + u_3 = 0 \\ 2u_3 = 0 \end{cases}$$

so  $u_3 = 0 \Rightarrow u_2 = 0$  and  $u_1$  is arbitrary so let  $u_1 = 1$ . Therefore  $\mathbf{u} = (1, 0, 0)^T$ . (2.2.13b)

implies

$$\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = 2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \Leftrightarrow \begin{cases} v_2 + v_3 = 2 \\ 2v_3 = 0 \end{cases}$$

thus,  $v_3 = 0$ ,  $v_2 = 2$  and  $v_1$  may be chosen arbitrary so we let  $v_1 = 0$ . This yields  $\mathbf{v} = (0, 2, 0)^T$ .

Finally, from (2.2.13a):

$$\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix} = 2 \left( \mathbf{v} + \frac{1}{2} \mathbf{u} \right) = \begin{pmatrix} 1 \\ 4 \\ 0 \end{pmatrix} \Leftrightarrow \begin{cases} w_2 + w_3 = 1 \\ 2w_3 = 4 \end{cases}$$

Hence,  $w_3 = 2$ ,  $w_2 = -1$  and we may choose  $w_1 = 0$  so  $\mathbf{w} = (0, -1, 2)^T$ . Consequently, the general solution may be written as

$$\begin{aligned} \mathbf{x}_t &= \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}_t = C_1 2^t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + C_2 2^t \left( \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} + t \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) \\ &+ C_3 2^t \left( \begin{pmatrix} 0 \\ -1 \\ 2 \end{pmatrix} + t \begin{pmatrix} 0 \\ 2 \\ 0 \end{pmatrix} + \frac{1}{2} t^2 \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \right) \end{aligned}$$

□



**Example 2.2.4.** Let

$$\mathbf{x} = (x_1, x_2)^T, \quad A = \begin{pmatrix} -2 & 1 \\ -1 & -2 \end{pmatrix}$$

and solve  $\mathbf{x}_{t+1} = A\mathbf{x}_t$ .

Suppose  $\mathbf{x}_t = \lambda^t \mathbf{v}$ . (2.2.9) implies

$$\begin{vmatrix} -2 - \lambda & 1 \\ -1 & -2 - \lambda \end{vmatrix} = 0 \Leftrightarrow \lambda^2 + 4\lambda + 5 = 0$$

$\Leftrightarrow \lambda_1 = -2 + i, \lambda_2 = -2 - i$  (distinct complex eigenvalues).

Further:  $|\lambda_1| = \sqrt{(-2)^2 + 1^2} = \sqrt{5}$   $\cos \theta = (-2)/\sqrt{5}$   $\sin \theta = 1/\sqrt{5}$  so

$$\lambda_1 = \sqrt{5}(\cos \theta + i \sin \theta).$$

The eigenvector  $\mathbf{u} = (u_1, u_2)^T$  corresponding to  $\lambda_1$  may be found from

$$\begin{pmatrix} -2 - (-2 + i) & 1 \\ -1 & -2 - (-2 + i) \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \Leftrightarrow \begin{cases} -iu_1 + u_2 = 0 \\ -u_1 - iu_2 = 0 \end{cases}$$

## American online

## LIGS University

is currently enrolling in the  
Interactive Online **BBA, MBA, MSc,**  
**DBA and PhD** programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive **Online** education
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to  
find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](#).





Let  $u_2 = t$ ,  $u_1 = -it$  so  $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = t \begin{pmatrix} -i \\ 1 \end{pmatrix}$ , so we choose  $\begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} -i \\ 1 \end{pmatrix}$  as eigenvector. Therefore (by use of Moivre's formulae), the solution in complex form becomes

$$\mathbf{x}_t = \sqrt{5}^t (\cos \theta t + i \sin \theta t) \begin{pmatrix} -i \\ 1 \end{pmatrix} = \begin{pmatrix} \sqrt{5}^t \{-i \cos \theta t + \sin \theta t\} \\ \sqrt{5}^t \{\cos \theta t + i \sin \theta t\} \end{pmatrix}$$

Two linear independent real solutions are found by taking the real and imaginary parts respectively:

$$\begin{aligned} \text{Real part } \begin{pmatrix} x_{1r} \\ x_{2r} \end{pmatrix}_t &= \sqrt{5}^t \begin{pmatrix} \sin \theta t \\ \cos \theta t \end{pmatrix} \\ \text{Imaginary part } \begin{pmatrix} x_{1i} \\ x_{2i} \end{pmatrix}_t &= \sqrt{5}^t \begin{pmatrix} -\cos \theta t \\ \sin \theta t \end{pmatrix} \end{aligned}$$

Thus, the general solution may be written as

$$\mathbf{x}_t = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}_t = C_1 \begin{pmatrix} x_{1r} \\ x_{2r} \end{pmatrix}_t + C_2 \begin{pmatrix} x_{1i} \\ x_{2i} \end{pmatrix}_t = \begin{pmatrix} \sqrt{5}^t \{C_1 \sin \theta t - C_2 \cos \theta t\} \\ \sqrt{5}^t \{C_1 \cos \theta t + C_2 \sin \theta t\} \end{pmatrix}$$

□

**Exercise 2.2.4.** Let  $\mathbf{x} = (x_1, x_2)^T$ ,  $A = \begin{pmatrix} 1 & 2 \\ 3 & 2 \end{pmatrix}$ ,  $B = \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}$  and find the general solution of

a)  $\mathbf{x}_{t+1} = A\mathbf{x}_t$ ,

b)  $\mathbf{x}_{t+1} = B\mathbf{x}_t$ ,

c) Let  $\mathbf{x} = (x_1, x_2, x_3)^T$  and find the general solution of  $\mathbf{x}_{t+1} = C\mathbf{x}_t$  where

$$C = \begin{pmatrix} -3 & 1 & -1 \\ -7 & 5 & -1 \\ -6 & 6 & -2 \end{pmatrix}$$

□

We close this section by a definition and an important theorem about stability of linear systems.

**Definition 2.2.1.** The linear system (2.2.7) is globally asymptotic stable if  $\lim_{t \rightarrow \infty} \mathbf{x}_t = 0$ . □

**Theorem 2.2.1.** The linear system (2.2.7) is globally asymptotic stable if and only if all the eigenvalues  $\lambda$  of  $A$  are located inside the unit circle  $|z| = 1$  in the complex plane.

**Proof:** In case of distinct eigenvalues the result follows immediately from (2.2.10).

Eigenvalues with multiplicity  $m$  lead according to our previous discussion to terms in the solution of form  $t^q \lambda^t$  where  $q \leq m - 1$ .

Now, if  $|\lambda| < 1$ , let  $|\lambda| = 1/s$  where  $s > 1$ . Then by L'Hopital's rule:  $\lim_{t \rightarrow \infty} (t^q/s^t) = 0$  so the result follows here too.  $\square$

### 2.3 The Leslie matrix

In Part I of this book we illustrated many aspects of the theory which we established by use of the quadratic map. Here in Part II we will use Leslie matrix models which are nothing but maps on the form  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  or  $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ .

Leslie matrix models are age-structured population models. They were independently developed in the 1940s by Bernardelli (1941), Lewis (1942) and Leslie (1945, 1948) but were not widely adopted by human demographers until the late 1960s and by ecologists until the 1970s. Some frequently quoted papers where the use of such models plays an important role are: Guckenheimer et al. (1977), Levin and Goodyear (1980), Silva and Hallam (1993), Wikan and Mjølhus (1996), Behncke (2000), Davydova et al. (2003), Mjølhus et al. (2005), and Kon (2005). The ultimate book on matrix population models which we refer to is "Matrix population models" by Hal Caswell (2001). Here we will deal with only a limited number of aspects of these models.

Let  $\mathbf{x}_t = (x_{0,t}, \dots, x_{n,t})^T$  be a population with  $n + 1$  nonoverlapping age classes at time  $t$ .

$x = x_0 + \dots + x_n$  is the total population.

Next, introduce the Leslie matrix

$$A = \begin{pmatrix} f_0 & f_1 & \cdots & f_n \\ p_0 & 0 & \cdots & 0 \\ 0 & \ddots & & \vdots \\ \vdots & & & \\ 0 & \cdots & p_{n-1} & 0 \end{pmatrix} \tag{2.3.1}$$

The meaning of the entries in (2.3.1) is as follows:  $f_i$  is the average fecundity (the average number of daughters born per female) of a member located in the  $i$ 'th age class.  $p_i$  may be interpreted as the survival probability from age class  $i$  to age class  $i + 1$  and clearly  $0 \leq p_i \leq 1$ . The relation between  $\mathbf{x}$  at two consecutive time steps (years) may then be expressed as

$$\mathbf{x}_{t+1} = A\mathbf{x}_t \tag{2.3.2}$$

or in map notation

$$h : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}, \quad \mathbf{x} \rightarrow A\mathbf{x} \tag{2.3.3}$$

Hence, what (2.3.2) really says is that all individuals  $x_i$  ( $i > 1$ ) in age class  $i$  at time  $t + 1$  are the survivors of the members of the previous age class  $x_{i-1}$  at time  $t$  (i.e.  $x_{i,t+1} = p_{i-1}x_{i-1,t}$ ), and since the individuals in the lowest age class cannot be survivors of any other age class they must have originated from reproduction (i.e.  $x_{0,t+1} = f_0x_{0,t} + \dots + f_nx_{n,t}$ ).

**DON'T EAT YELLOW SNOW**

What will your advice be?

Some advice just states the obvious. But to give the kind of advice that's going to make a real difference to your clients you've got to listen critically, dig beneath the surface, challenge assumptions and be credible and confident enough to make suggestions right from day one. At Grant Thornton you've got to be ready to kick start a career right at the heart of business.

Sound like you? Here's our advice: visit [GrantThornton.ca/careers/students](http://GrantThornton.ca/careers/students)

Scan here to learn more about a career with Grant Thornton.

**Grant Thornton**  
An instinct for growth™

© Grant Thornton LLP. A Canadian Member of Grant Thornton International Ltd



Depending on the species under consideration, nonlinearities may show up on different entries in the matrix. For example, in fishery models it is often assumed that density effects occur mainly through the first year of life so one may assume  $f_i = f_i(x)$ . It is also customary to write  $f_i(x)$  as a product of a density independent part  $F_i$  and a density dependent part  $\hat{f}_i(x)$  so  $f_i(x) = F_i \hat{f}_i(x)$ . In the following we shall assume that every fertile age class has the same fecundity. Thus, we may drop the subscript  $i$  and write  $f_i(x) = f(x)$ . Frequently used fecundity functions are:

$$f(x) = F e^{-\alpha x} \tag{2.3.4}$$

which is often referred to as the overcompensatory Ricker relation and

$$f(x) = \frac{F}{1 + \alpha x} \tag{2.3.5}$$

the compensatory Beverton and Holt relation.

Instead of assuming  $f = f(x)$  one may alternatively suppose  $f = f(y)$  where  $y = \alpha_0 x_0 + \dots + \alpha_n x_n$  is the weighted sum of the age classes. If only one age class, say  $x_i$ , contributes to density effects one writes  $f = f(x_i)$ . In the case where an age class  $x_i$  is not fertile we simply write  $F_i = 0$ . (Species where most age classes are fertile are called iteroparous. Species where fecundity is restricted to the last age class only are called semelparous.)

The survival probabilities may of course also be density dependent so in such cases we adopt the same strategy as in the fecundity case and write  $p(\cdot) = P \hat{p}(\cdot)$  where  $P$  is a constant.

A final but important comment is that one in most biological relevant situations supposes  $p'(\cdot) \leq 0$  and  $f'(\cdot) \leq 0$ . The standard counter example is when the Allé effect (cf. Caswell, 2001) is modelled. Then one may use  $f'(x) \geq 0$  and/or  $p'(x) \geq 0$  in case of small populations  $x$ . (Allé effects will not be considered here.)

In the subsequent sections we shall analyse nonlinear maps and as already mentioned the theory will be illustrated by use of (2.3.2), (2.3.3). However, if both  $f_i = F_i$  and  $p_i = P_i$  the Leslie matrix is linear and we let

$$M = \begin{pmatrix} F_0 & \dots & & & F_n \\ P_0 & 0 & \dots & & 0 \\ 0 & & & & \\ \vdots & & \ddots & & \vdots \\ 0 & \dots & 0 & P_{n-1} & 0 \end{pmatrix} \tag{2.3.6}$$

We close this section by a study of the linear case

$$h : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}, \quad \mathbf{x} \rightarrow M\mathbf{x} \quad (2.3.7)$$

The eigenvalues of  $M$  may be obtained from  $|M - \lambda I| = 0$ .

**Exercise 2.3.1.**

a) Assume that  $M$  is  $3 \times 3$  and show that the eigenvalue equation becomes

$$\lambda^3 - F_0\lambda^2 - P_0F_1\lambda - P_0P_1F_2 = 0$$

b) Generalize and show that if  $M$  is a  $(n + 1) \times (n + 1)$  matrix then the eigenvalue equation may be written

$$\lambda^{n+1} - F_0\lambda^n - P_0F_1\lambda^{n-1} - \dots - P_0P_1 \dots P_{n-1}F_n = 0 \quad (2.3.8)$$

□

Next, we need some definitions:

**Definition 2.3.1.** A matrix  $A$  is nonnegative if all its elements are greater or equal to zero. It is positive if all elements are positive.

Clearly, the Leslie matrix is nonnegative.

□

**Definition 2.3.2.** Let  $N_0, \dots, N_n$  be nodes representing the  $n + 1$  age classes in a population model. Draw a directed path from  $N_i$  to  $N_j$  if individuals in age class  $i$  at time  $t$  contribute to individuals of age  $j$  at time  $t + 1$  including the case that a path may go from  $N_i$  to itself. A diagram where all such nodes and paths are drawn is called a life cycle graph.

□

**Definition 2.3.3.** A nonnegative matrix  $A$  and its associated life cycle graph is irreducible if its life cycle graph is strongly connected (i.e. if between every pair of distinct nodes  $N_i, N_j$  in the graph there is a directed path of finite length that begins at  $N_i$  and ends at  $N_j$ ).

□

**Definition 2.3.4.** A reducible life cycle graph contains at least one age group that cannot contribute by any developmental path to some other age group.


□

Examples of two irreducible Leslie matrices and one reducible one with associated life cycle graphs are given in Figure 10.




**Exercise 2.3.2.** Referring to Figure 10 write down the matrix and associated life cycle graph in the case of four age classes where only the two in the middle are fertile. □


$$\begin{pmatrix} F_0 & F_1 & F_2 \\ P_0 & 0 & 0 \\ 0 & P_1 & 0 \end{pmatrix}$$



$$\begin{pmatrix} 0 & 0 & F_2 \\ P_0 & 0 & 0 \\ 0 & P_1 & 0 \end{pmatrix}$$



$$\begin{pmatrix} F_0 & F_1 & 0 \\ P_0 & 0 & 0 \\ 0 & P_1 & 0 \end{pmatrix}$$



**Figure 10:** Two irreducible and one reducible matrices with corresponding life cycle graphs.

.....Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".





**Definition 2.3.5.** An irreducible matrix  $A$  is said to be primitive if it becomes positive when raised to sufficiently high powers. Otherwise  $A$  is imprimitive (cyclic) with index of imprimitivity equal to the greatest common divisor of the loop lengths in the life cycle graph.  $\square$

**Exercise 2.3.3.** Show by direct calculation that the first irreducible Leslie matrix in Figure 10 is primitive and that the second one is imprimitive (cyclic) with index of imprimitivity equal to 3.  $\square$

Regarding nonnegative matrices the main results may be summarized in the following theorem which is often referred to as the Perron-Frobenius theorem.

**Theorem 2.3.1 (Perron-Frobenius).**

- 1) If  $A$  is positive or nonnegative and primitive, then there exists a real eigenvalue  $\lambda_0 > 0$  which is a simple root of the characteristic equation  $|A - \lambda I| = 0$ . Moreover, the eigenvalue is strictly greater than the magnitude of any other eigenvalue,  $\lambda_0 > |\lambda_i|$  for  $i \neq 0$ . The eigenvector  $\mathbf{u}_0$  corresponding to  $\lambda_0$  is real and strictly positive.  $\lambda_0$  may not be the only positive eigenvalue but if there are others they do not have nonnegative eigenvectors.
- 2) If  $A$  is irreducible but imprimitive (cyclic) with index of imprimitivity  $d + 1$  there exists a real eigenvalue  $\lambda_0 > 0$  which is a simple root of  $|A - \lambda I| = 0$  with associated eigenvector  $\mathbf{u}_0 > \mathbf{0}$ . The eigenvalues  $\lambda_i$  satisfy  $\lambda_0 \geq |\lambda_i|$  for  $i \neq 0$  but there are  $d$  complex eigenvalues equal in magnitude to  $\lambda_0$  whose values are  $\lambda_0 \exp(2k\pi i/(d + 1))$ ,  $k = 1, 2, \dots, d$ .

For a general proof of Theorem 2.3.1 we refer to the literature. See for example Horn and Johnson (1985).

Concerning the Leslie matrix  $M$  (2.3.6) we shall study two cases in somewhat more detail: (I) the case where all fecundities  $F_i > 0$ , and (II) the semelparous case where  $F_i = 0$ ,  $i = 0, \dots, n - 1$  but  $F_n > 0$ . In both cases it is assumed that  $0 < P_i \leq 1$  for all  $i$ .

Let us prove Theorem 2.3.1 assuming (I):

Since  $F_n > 0$  and  $0 < P_i \leq 1$  it follows directly from (2.3.8) that  $\lambda = 0$  is impossible. Therefore, we may divide (2.3.8) by  $\lambda^{n+1}$  to obtain

$$f(\lambda) = \frac{F_0}{\lambda} + \frac{P_0 F_1}{\lambda^2} + \dots + \frac{P_0 P_1 \dots P_{n-1} F_n}{\lambda^{n+1}} = 1 \tag{2.3.9}$$

Clearly,  $\lim_{\lambda \rightarrow 0} f(\lambda) = \infty$ ,  $\lim_{\lambda \rightarrow \infty} f(\lambda) = 0$ , and since  $f'(\lambda) < 0$  for  $\lambda > 0$  it follows that there exists a unique positive  $\lambda_0$  which satisfies  $f(\lambda_0) = 1$ . Therefore, assume  $\lambda_0^{-1} = e^\gamma$  and rewrite (2.3.9) as

$$f(\lambda) = F_0 e^\gamma + P_0 F_1 e^{2\gamma} + \dots + P_0 P_1 \dots P_{n-1} F_n e^{(n+1)\gamma} = 1 \quad (2.43)$$

Next, let  $\lambda_j^{-1} = \exp(\alpha + \beta i) = e^\alpha (\cos \beta + i \sin \beta)$  for  $j = 1, \dots, n$  and since  $\lambda_0$  is unique and positive we may assume  $\beta$  real and positive and  $\beta \neq 2k\pi$ ,  $k = 1, 2, \dots$ . Then  $\lambda_j^{-p} = e^{\alpha p} (\cos p\beta + i \sin p\beta)$  which inserted into  $f(\lambda)$ , considering the real part only, gives

$$F_0 e^\alpha \cos \beta + P_0 F_1 e^{2\alpha} \cos 2\beta + \dots + P_0 P_1 \dots P_{n-1} F_n e^{(n+1)\alpha} \cos(n+1)\beta = 1 \quad (2.3.11)$$

Now, since  $\beta$  is not a multiple of  $2\pi$  it follows that  $\cos j\beta$  and  $\cos(j+1)\beta$  cannot both be equal to unity. Consequently, by comparing (2.3.10) and (2.3.11), we have  $e^\alpha > e^\gamma \Leftrightarrow |\lambda_j| < \lambda_0$  for  $j = 1, \dots, n$ .

Finally, in order to see that the eigenvector  $\mathbf{u}_0$  corresponding to  $\lambda_0$  has only positive elements, recall that  $\mathbf{u}_0$  must be computed from  $M\mathbf{u}_0 = \lambda_0\mathbf{u}_0$ , and in order to avoid  $\mathbf{u}_0 = \mathbf{0}$  we must choose one of the components of  $\mathbf{u}_0 = (u_{00}, \dots, u_{n0})^T$  free, so let  $u_{00} = 1$ . Then from  $M\mathbf{u}_0 = \lambda_0\mathbf{u}_0$ :  $P_0 \cdot 1 = \lambda_0 u_{10}$ ,  $P_1 u_{10} = \lambda_0 u_{20}$ ,  $\dots$ ,  $P_{n-1} u_{n-10} = \lambda_0 u_{n0}$  which implies

$$u_{10} = \frac{P_0}{\lambda_0}, \quad u_{20} = \frac{P_1 u_{10}}{\lambda_0} = \frac{P_0 P_1}{\lambda_0^2} \dots u_{n0} = \frac{P_0 \dots P_{n-1}}{\lambda_0^n}$$

which proves what we want. □

(This proof is based upon Frauenthal (1986).) The proof of Theorem 2.3.1 under the assumption (II) is left to the reader.

—

Let us now turn to the asymptotic behaviour of the linear map (2.3.7) in light of the results of Theorem 2.3.1.

In the case where all  $F_i > 0$  we may express the solution of (2.3.7) (cf. (2.2.10)) as

$$\mathbf{x}_t = c_0 \lambda_0^t \mathbf{u}_0 + c_1 \lambda_1^t \mathbf{u}_1 + \dots + c_n \lambda_n^t \mathbf{u}_n \quad (2.3.12)$$

where  $\lambda_i$  (real or complex,  $\lambda_0$  real) are the eigenvalues of  $M$  numbered in order of decreasing magnitude and  $\mathbf{u}_i$  are the corresponding eigenvectors. Further,

$$\frac{\mathbf{x}_t}{\lambda_0^t} = c_0 \mathbf{u}_0 + c_1 \left(\frac{\lambda_1}{\lambda_0}\right)^t \mathbf{u}_1 + \dots + c_n \left(\frac{\lambda_n}{\lambda_0}\right)^t \mathbf{u}_n$$

and since  $\lambda_0 > |\lambda_i|, i \neq 0$

$$\lim_{t \rightarrow \infty} \frac{\mathbf{x}_t}{\lambda_0^t} = c_0 \mathbf{u}_0 \tag{2.3.13}$$

Consequently, if  $M$  is nonnegative and primitive, the long term dynamics of the population are described by the growth rate  $\lambda_0$  and the stable population structure  $\mathbf{u}_0$ . Thus  $\lambda_0 > 1$  implies an exponential increasing population,  $0 < \lambda_0 < 1$  an exponential decreasing population, where we in all cases have the stable age distribution  $\mathbf{u}_0$ .

If  $M$  is irreducible but imprimitive with index of imprimitivity  $d + 1$  it follows from part 2 of the Perron-Frobenius theorem that the limit (2.3.13) may be expressed as

$$\lim_{t \rightarrow \infty} \frac{\mathbf{x}_t}{\lambda_0^t} = c_0 \mathbf{u}_0 + \sum_{k=1}^d c_k e^{(2k\pi/(d+1))it} \mathbf{u}_i \tag{2.3.14}$$



## Join the best at the Maastricht University School of Business and Economics!

### Top master's programmes

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht  
University is  
the best specialist  
university in the  
Netherlands  
(Elsevier)


Visit us and find out why we are the best!

Master's Open Day: 22 February 2014

www.mastersopenday.nl



As opposed to the dynamical consequences of 1) in the Perron-Frobenius theorem we now conclude from (2.3.14) that  $\mathbf{u}_0$  is not stable in the sense that an initial population not proportional to  $\mathbf{u}_0$  will converge to it. Instead, the limit (2.3.14) is periodic with period  $d + 1$ .



**Figure 11:** The hypothetical "beetle" population of Bernardelli as function of time.  $\Delta$  is the total population  $\square$ , + and  $\diamond$  correspond to the zeroth, first and second age classes respectively. Clearly, there is no stable age distribution.

**Example 2.3.1 (Bernardelli 1941).** The first paper where the matrix  $M$  was considered came in 1941. There, Bernardelli considered a hypothetical beetle population obeying the equation

$$\mathbf{x}_{t+1} = B\mathbf{x}_t \quad \text{where} \quad B = \begin{pmatrix} 0 & 0 & 6 \\ 1/2 & 0 & 0 \\ 0 & 1/3 & 0 \end{pmatrix}$$

Clearly,  $B$  is irreducible and imprimitive with index of imprimitivity equal to 3 (cf. Exercise 2.3.2). Moreover, the eigenvalues of  $B$  are easily found to be  $\lambda_1 = 1$  and  $\lambda_{2,3} = \exp(\pm 2\pi i/3)$  and it is straightforward to show that  $B^3 = I$  so each initial age distribution will repeat itself in a regular manner every third year as predicted by (2.3.14). In Figure 11 we show the total hypothetical beetle population together with the three age classes as function of time, and clearly there is no stable age distribution.  $\square$

## 2.4 Fixed points and stability of nonlinear systems

In this section we turn to the nonlinear case  $\mathbf{x} \rightarrow f(\mathbf{x})$  which in difference equation notation may be cast in the form

$$\begin{aligned} x_{1,t+1} &= f_1(x_{1,t}, \dots, x_{n,t}) \\ &\vdots \\ x_{n,t+1} &= f_n(x_{1,t}, \dots, x_{n,t}) \end{aligned} \tag{2.4.1}$$

**Definition 2.4.1.** A point  $\mathbf{x}^* = (x_1^*, \dots, x_n^*)$  which satisfies  $\mathbf{x}^* = f(\mathbf{x}^*)$  is called a fixed point for (2.4.1).  $\square$

**Example 2.4.1.** Assume that  $F_0 + P_0F_1 > 1$ ,  $x = x_0 + x_1$  and find the nontrivial fixed point  $(x_0^*, x_1^*)$  of the two-dimensional Leslie matrix model (the Ricker model)

$$\begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \longrightarrow \begin{pmatrix} F_0e^{-x} & F_1e^{-x} \\ P_0 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \tag{2.4.2}$$

According to Definition 2.4.1 the fixed point satisfies

$$x_0^* = F_0e^{-x^*}x_0^* + F_1e^{-x^*}x_1^* \tag{2.4.3a}$$

$$x_1^* = P_0x_0^* \tag{2.4.3b}$$

and if we insert (2.4.3b) into (2.4.3a) we obtain  $1 = e^{-x^*}(F_0 + P_0F_1)$ , hence the total equilibrium population becomes  $x^* = \ln(F_0 + P_0F_1)$ . Further, since  $x^* = x_0^* + x_1^*$  and  $x_1^* = P_0x_0^*$  we easily find

$$(x_0^*, x_1^*) = \left( \frac{1}{1 + P_0} x^*, \frac{P_0}{1 + P_0} x^* \right) \tag{2.4.4}$$

(Note that  $F_0 + P_0F_1 > 1$  is necessary in order to obtain a biological acceptable solution.)  $\square$

**Exercise 2.4.1.** Still assuming  $F_0 + P_0F_1 > 1$ , show that the fixed point  $(x_0^*, x_1^*)$  of the two-dimensional Beverton and Holt model

$$\begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \longrightarrow \begin{pmatrix} \frac{F_0}{1+x} & \frac{F_1}{1+x} \\ P_0 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} \tag{2.4.5}$$

becomes

$$(x_0^*, x_1^*) = \left( \frac{1}{1 + P_0} x^*, \frac{P_0}{1 + P_0} x^* \right) \tag{2.4.6}$$

where  $x^* = F_0 + P_0 F_1 - 1$ . □

**Example 2.4.2.** Find the nontrivial fixed point of the general Ricker model:

$$\begin{pmatrix} x_0 \\ \vdots \\ x_1 \end{pmatrix} \rightarrow \begin{pmatrix} F_0 e^{-x} & \cdots & & F_n e^{-x} \\ P_0 & 0 & \cdots & 0 \\ \vdots & & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & P_{n-1} & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ \vdots \\ x_n \end{pmatrix} \tag{2.4.7}$$

The fixed point  $\mathbf{x}^* = (x_0^*, \dots, x_n^*)$  obeys

$$\begin{aligned} x_0^* &= e^{-x^*} (F_0 x_0^* + \cdots + F_n x_n^*) \\ x_1^* &= P_0 x_0^* \\ &\vdots \\ x_n^* &= P_{n-1} x_{n-1}^* \end{aligned}$$

**Empowering People. Improving Business.**

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

**BI NORWEGIAN BUSINESS SCHOOL**

EFMD **EQUIS ACCREDITED**

[www.bi.edu/master](http://www.bi.edu/master)



From the last  $n$  equations we have  $x_1^* = P_0 x_0^*$ ,  $x_2^* = P_1 x_1^* = P_0 P_1 x_0^*$ ,  $x_n^* = P_0 \cdots P_{n-1} x_0^*$  which inserted into the first equation give

$$1 = e^{-x^*} (F_0 + P_0 F_1 + P_0 P_1 F_2 + \cdots + P_0 \cdots P_{n-1} F_n) \quad (2.4.8)$$

Hence,

$$x^* = \ln(F_0 + P_0 F_1 + \cdots + P_0 \cdots P_{n-1} F_n) = \ln \left( \sum_{i=0}^n F_i L_i \right)$$

where  $L_i = P_0 P_1 \cdots P_{i-1}$  and by convention  $L_0 = 1$ . From  $\sum x_i^* = x^*$  and  $x_1^* = P_0 x_0^* = L_1 x_0^*$ ,  $x_2^* = P_0 P_1 x_0^* = L_2 x_0^*$  and  $x_i^* = L_i x_0^*$  we obtain

$$(x_0^*, \dots, x_n^*) = \left( \frac{L_1}{\sum_{i=0}^n L_i} x^*, \dots, \frac{L_i}{\sum_{i=0}^n L_i} x^*, \dots, \frac{L_n}{\sum_{i=0}^n L_i} x^* \right) \quad (2.4.9)$$

Again,  $\sum_{i=0}^n F_i L_i > 1$  is required in order to have an acceptable biological equilibrium.  $\square$

**Exercise 2.4.2.** Generalize Exercise 2.4.1 in the same way as in Example 2.4.2 and obtain a formulae for the fixed point of the  $n + 1$  dimensional Beverton and Holt model. A detailed analysis of the Beverton and Holt model may be obtained in Silva and Hallam (1992).  $\square$

In order to reveal the stability properties of the fixed point  $\mathbf{x}^*$  of (2.4.1) we follow the same pattern as we did in Section 1.4. Let  $\mathbf{x} = \mathbf{x}^* + \boldsymbol{\xi}$ , then expand  $f_i(\mathbf{x})$  in its Taylor series about  $x^*$ , taking the linear terms only in order to obtain

$$\begin{aligned} x_{1,t+1}^* + \xi_{1,t+1} &\approx f_1(\mathbf{x}_t^*) + \frac{\partial f_1}{\partial x_1} \xi_{1,t} + \cdots + \frac{\partial f_1}{\partial x_n} \xi_{n,t} \\ &\vdots \\ x_{n,t+1}^* + \xi_{n,t+1} &\approx f_n(\mathbf{x}_t^*) + \frac{\partial f_n}{\partial x_1} \xi_{1,t} + \cdots + \frac{\partial f_n}{\partial x_n} \xi_{n,t} \end{aligned}$$

where all derivatives are evaluated at  $\mathbf{x}^*$ . Moreover,  $x_{i,t+1}^* = f_i(\mathbf{x}_t^*)$ . Consequently, the linearized map (or linearization) of (2.4.1) becomes

$$\begin{pmatrix} \xi_1 \\ \vdots \\ \xi_n \end{pmatrix} \longrightarrow \begin{pmatrix} \frac{\partial f_1}{\partial x_1}(x^*) & \cdots & \frac{\partial f_1}{\partial x_n}(x^*) \\ \vdots & & \vdots \\ \frac{\partial f_n}{\partial x_1}(x^*) & \cdots & \frac{\partial f_n}{\partial x_n}(x^*) \end{pmatrix} \begin{pmatrix} \xi_1 \\ \vdots \\ \xi_n \end{pmatrix} \quad (2.4.10)$$

where the matrix is called the Jacobian.



If the fixed point  $\mathbf{x}^*$  of (2.4.1) shall be locally asymptotic stable we clearly must have

$$\lim_{t \rightarrow \infty} \xi_t \rightarrow 0 \quad (2.59)$$

and according to Theorem 2.2.1 this is equivalent to say:

**Theorem 2.4.1.** The fixed point  $\mathbf{x}^*$  of the nonlinear system (2.4.1) is locally asymptotic stable if and only if all the eigenvalues  $\lambda$  of the Jacobian matrix are located inside the unit circle  $|z| = 1$  in the complex plane.  $\square$

**Example 2.4.3.**

a) Define  $\hat{F}\hat{x} = F_0x_0^* + F_1x_1^*$  and show that the fixed point (2.4.4) of the Ricker map (2.4.2) is locally asymptotic stable provided

$$\hat{F}\hat{x}(1 + P_0) > 0 \quad (2.4.12a)$$

$$2F_0 + \hat{F}\hat{x}(P_0 - 1) > 0 \quad (2.4.12b)$$

$$2P_0F_1 + F_0 - P_0\hat{F}\hat{x} > 0 \quad (2.4.12c)$$

b) Assume that  $F_0 = F_1 = F$  (same fecundity in both age classes) and show that (2.4.12b), (2.4.12c) may be expressed as

$$F < \frac{1}{1 + P_0} e^{2/(1-P_0)} \quad (2.4.13b)$$

$$F < \frac{1}{1 + P_0} e^{(1+2P_0)/P_0} \quad (2.4.13c)$$

**Solution:**

a) Rewrite (2.4.2) as

$$x_0 \rightarrow f_1(x_0, x_1) = F_0e^{-x}x_0 + F_1e^{-x}x_1$$

$$x_1 \rightarrow f_2(x_0, x_1) = P_0x_0$$

Then the Jacobian becomes

$$J = \begin{pmatrix} e^{-x^*}(F_0 - \hat{F}\hat{x}) & e^{-x^*}(F_1 - \hat{F}\hat{x}) \\ P_0 & 0 \end{pmatrix} \quad (2.4.14)$$

and the eigenvalue equation  $|J - \lambda I| = 0$  may be cast in the form

$$\lambda^2 - \frac{F_0 - \hat{F}\hat{x}}{F_0 + P_0F_1} \lambda - P_0 \frac{F_1 - \hat{F}\hat{x}}{F_0 + P_0F_1} = 0 \quad (2.4.15)$$

where we have used  $e^{-x^*} = (F_0 + P_0F_1)^{-1}$ .

(2.4.15) is a second order polynomial and  $|\lambda| < 1$  if the corresponding Jury criteria (2.1.14) are satisfied. Therefore, by defining

$$a_1 = -\frac{F_0 - \hat{F}\hat{x}}{F_0 + P_0F_1} \quad a_2 = -P_0 \frac{F_1 - \hat{F}\hat{x}}{F_0 + P_0F_1}$$

we easily obtain from (2.1.14) that the fixed point is locally asymptotic stable provided the inequalities (2.4.12a)-(2.4.12c) hold.

## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info

 **Helpmyassignment**

**Remark 2.4.1:** Scrutinizing the criteria, it is obvious that (2.4.12a) holds for any (positive) equilibrium population  $x^*$ . It is also clear that in case of  $\hat{F}\hat{x}$  sufficiently small the same is true for both (2.4.12b,c) as well which allow us to conclude that  $(x_0^*, x_1^*)$  is stable in case of “small” equilibrium population  $x^*$ . However, if  $\hat{F}\hat{x}$  becomes large, both (2.4.12b) and (2.4.12c) contain a large negative term so evidently there are regions in parameter space where (2.4.12b) or (2.4.12c) or both are violated and consequently regions where  $(x_0^*, x_1^*)$  is no longer stable.

b) If  $F_0 = F_1 = F$ , then  $\hat{F}\hat{x} = Fx^*$ , thus (2.4.15) may be expressed as

$$\lambda^2 - \frac{1 - x^*}{1 + P_0} \lambda - P_0 \frac{1 - x^*}{1 + P_0} = 0 \quad (2.4.16)$$

and the criteria (2.4.12b), (2.4.12c) simplify to

$$2 + x^*(P_0 - 1) > 0$$

$$2P_0 + 1 - Px^* > 0$$

(2.4.13b) and (2.4.13c) are now established by use of  $x^* = \ln[F(1 + P_0)]$ .

A final but important observation is that whenever  $0 < P_0 < 1/2$ , (2.4.13b) will be violated prior to (2.4.13c) if  $F$  is increased. On the other hand, if  $1/2 < P_0 \leq 1$ , (2.4.13c) will be violated first through an increase of  $F$ . (As we shall see later, this fact has a crucial impact of the possible dynamics in the unstable parameter region.)  $\square$

**Example 2.4.4 (Example 2.4.2 continued).** Let the fecundities be equal (i.e.  $F_0 = \dots = F_n = F$ ) in the general  $n + 1$  dimensional Ricker model that we considered in Example 2.4.2. Then,  $x^* = \ln(FD)$ ,  $D = \sum_{i=0}^n L_i$  and the fixed point  $\mathbf{x}^*$  may be written as  $\mathbf{x}^* = (x_0^*, \dots, x_i^*, \dots, x_n^*)$  where  $x_i^* = (L_i/D)x^*$ .

The eigenvalue equation (cf. (2.4.16)) may be cast in the form

$$\lambda^{n+1} - \frac{1}{D}(1 - x^*) \sum_{i=0}^n L_i \lambda^{n-i} = 0 \quad (2.4.17)$$

Our goal is to show that the fixed point  $\mathbf{x}^*$  is locally asymptotic stable whenever  $x^* < 2$  (i.e. that all the eigenvalues  $\lambda$  of (2.4.17) are located inside the unit circle.)

In contrast to Example 2.4.3, Theorem 2.1.9 obviously does not work here so instead we appeal to Theorem 2.1.10 (Rouché's theorem). Therefore, assume  $|1 - x^*| < 1$ , let  $f(\lambda) = \lambda^{n+1}$ ,  $g(\lambda) = -(1/D)(1 - x^*) \sum_{i=0}^n L_i \lambda^{n-i}$  and rewrite (2.4.17) as  $f(\lambda) + g(\lambda) = 0$ . Clearly,  $f$  and  $g$  are analytic functions on and inside the unit circle  $C$  and the equation  $f(\lambda) = 0$  has  $n + 1$  roots inside  $C$ .

On the boundary we have

$$\begin{aligned} |g(\lambda)| &= \left| -\frac{1}{D}(1 - x^*) \sum_{i=0}^n L_i \lambda^{n-i} \right| \\ &\leq \left| \frac{L_0}{D}(1 - x^*) \lambda^n \right| + \left| \frac{L_1}{D}(1 - x^*) \lambda^{n-1} \right| + \dots + \left| \frac{L_n}{D}(1 - x^*) \right| \\ &\leq |1 - x^*| < |f(\lambda)| \end{aligned}$$

Thus, according to Theorem 2.1.10,  $f(\lambda) + g(\lambda)$  and  $f(\lambda)$  have the same number of zeros inside  $C$ , hence (2.4.17) has  $n + 1$  zeros inside the unit circle which proves that  $x^* < 2$  is sufficient to guarantee a stable fixed point. Other properties of the Ricker model (2.4.7) may be obtained in Wikan and Mjølhus (1996).  $\square$

**Exercise 2.4.2 (Exercise 2.4.1 continued).**

a) Consider the two-dimensional Beverton and Holt model (see Exercise 2.4.1) and show that the fixed point  $(x_0^*, x_1^*)$  is always stable. ( $F_0 = F_1 = F$ .)

b) Generalize to  $n + 1$  age classes. ( $F_0 = \dots = F_n = F$ .)  $\square$

**Exercise 2.4.3:** Assume  $P_0 < 1$  and consider the two-dimensional semelparous Ricker model:

$$\begin{aligned} x_{0,t+1} &= F_1 e^{-x_t} x_1 \\ x_{1,t+1} &= P_0 x_0 \end{aligned} \tag{2.4.18}$$

a) Compute the nontrivial fixed point  $(x_0^*, x_1^*)$ .

b) Show that the eigenvalue equation may be written as

$$\lambda^2 + \frac{x_1^*}{P_0} \lambda - (1 - x_1^*) = 0$$

and use the Jury criteria to conclude that  $(x_0^*, x_1^*)$  is always unstable.

c) Show that

$$x_{0,t+2} = (P_0 F_1) e^{-x_{t+1}} x_{0,t} \quad (2.4.19)$$

$$x_{1,t+2} = (P_0 F_1) e^{-x_t} x_{1,t}$$

d) Assume that there exists a two-cycle where the points in the cycle are on the form  $(A, 0)$ ,  $(0, B)$  and show that the cycle is  $((1/P_0) \ln(P_0 F_1), 0)$ ,  $(0, \ln(P_0 F_1))$ .

e) Show that the two cycle in d) is stable provided  $0 < P_0 F_1 < e^2$ . □

—

Next, consider the general system (2.4.1) and its linearization (2.4.10) and let  $\lambda$  be the eigenvalues of the Jacobian. We now define the following decompositions of  $\mathbb{R}^n$ .

**Brain power**

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**

 [Click on the ad to read more](#)

**Definition 2.4.2.**

$E^s$  is the subspace which is spanned by the (generalized) eigenvectors whose eigenvalues satisfy  $|\lambda| < 1$ .

$E^c$  is the subspace which is spanned by the (generalized) eigenvectors whose eigenvalues satisfy  $|\lambda| = 1$ .

$E^u$  is the subspace which is spanned by the (generalized) eigenvectors whose corresponding eigenvalues satisfy  $|\lambda| > 1$ .

$\mathbb{R}^n = E^s \oplus E^c \oplus E^u$  and the subspaces  $E^s$ ,  $E^c$  and  $E^u$  are called the stable, the center and the unstable subspace respectively.  $\square$

By use of the definition above, the stability result stated in Theorem 2.4.1 may be reformulated as follows:

$\mathbf{x}^* = (x_0^*, \dots, x_n^*)$  is locally asymptotic stable if  $E^u = \{\mathbf{0}\}$  and  $E^c = \{\mathbf{0}\}$ .

$\mathbf{x}^*$  is unstable if  $E^u \neq \{\mathbf{0}\}$ .

$\mathbf{x}^* = (x_0^*, \dots, x_n^*)$  is called a hyperbolic fixed point if  $E^c = \{\mathbf{0}\}$  (cf. Section 1.4). ( $\mathbf{x}^*$  is attracting if  $|\lambda| < 1$ , repelling if  $|\lambda| > 1$ .)

We close this section by stating two general theorems which link the nonlinear behaviour close to a fixed point to the linear behaviour.

**Theorem 2.4.2 (Hartman-Grobman).** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a  $C^1$  diffeomorphism with a hyperbolic fixed point  $x^*$  and let  $Df$  be the linearization. Then there exists a homeomorphism  $h$  defined on some neighbourhood  $U$  on  $x^*$  such that

$$(h \circ f)(\xi) = Df(x^*) \circ h(\xi) \tag{2.4.20}$$

for  $\xi \in U$ .  $\square$

**Theorem 2.4.3.** There exists a stable manifold  $W_{\text{loc}}^s(\mathbf{x}^*)$  and an unstable manifold  $W_{\text{loc}}^u(\mathbf{x}^*)$  which are a) invariant, and b) is tangent to  $E^s$  and  $E^u$  at  $\mathbf{x}^*$  and have the same dimension as  $E^s$  and  $E^u$ .  $\square$

## 2.5 The Hopf bifurcation

There are three ways in which the fixed point  $\mathbf{x}^* = (x_0^*, \dots, x_1^*)$  of a nonlinear map,  $f_\mu : \mathbb{R}^n \rightarrow \mathbb{R}^n$  may fail to be hyperbolic. One way is that an eigenvalue  $\lambda$  of the linearization crosses the unit circle (sphere) through 1. Then, in the generic case, a saddle-node bifurcation occurs. Another possibility is that  $\lambda$  crosses the unit circle at  $-1$  which in turn leads generically to a flip bifurcation. The third possibility is that a pair of complex eigenvalues  $\lambda, \bar{\lambda}$  cross the unit circle. In this case the fixed point will undergo a Hopf bifurcation which we will now describe. Note that the saddle-node and the flip bifurcations may occur in one-dimensional maps,  $f_\mu : \mathbb{R} \rightarrow \mathbb{R}$ . The Hopf bifurcation may take place when the dimension  $n$  of the map is equal or larger than two. In this section we will restrict the analysis to the case  $n = 2$  only. Later on in section 2.7 we will show how both the flip and the Hopf bifurcation may be analysed in case of  $n > 2$ .

**Theorem 2.5.1.** Let  $f_\mu : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a  $C^3$  two-dimensional one-parameter family of maps whose fixed point is  $\mathbf{x}^* = (x_0^*, x_1^*)$ . Moreover, assume that the eigenvalues  $\lambda(\mu), \bar{\lambda}(\mu)$  of the linearization are complex conjugates. Suppose that

$$|\lambda(\mu_0)| = 1 \quad \text{but } \lambda^i(\mu_0) \neq 1 \text{ for } i = 1, 2, 3, 4 \quad (2.5.1)$$

and

$$\frac{d|\lambda(\mu_0)|}{d\mu} = d \neq 0 \quad (2.5.2)$$

Then, there is a sequence of near identity transformations  $h$  such that  $hf_\mu h^{-1}$  in polar coordinates may be written as

$$hf_\mu h^{-1}(r, \varphi) = ((1 + d\mu)r + ar^3, \varphi + c + br^2) + \text{higher order terms} \quad (2.5.3)$$

Moreover, if  $a \neq 0$  there is an  $\varepsilon > 0$  and a closed curve  $\xi_\mu$  of the form  $r = r_\mu(\varphi)$  for  $0 < \mu < \varepsilon$  which is invariant under  $f_\mu$ .  $\square$

Before we sketch a proof of the theorem let us give a few remarks.

**Remark 2.5.1.** Performing near identity transformations as stated in the theorem is also called normal form calculations. Hence, formulae (2.5.3) is nothing but the original map written in normal form.



**Remark 2.5.2.** If  $d > 0$  (cf. (2.5.2)) then the complex conjugated eigenvalues cross the unit circle outwards which of course means that  $(x_0^*, x_1^*)$  loses its stability at bifurcation threshold  $\mu = \mu_0$ . If  $d < 0$  the eigenvalues move inside the unit circle.  $\square$

**Remark 2.5.3.**  $\lambda(\mu_0) = 1$  or  $\lambda^2(\mu_0) = 1$  (cf. 2.5.1)) correspond to the well known saddle-node or flip bifurcations respectively.  $\lambda^3(\mu_0) = 1$  and  $\lambda^4(\mu_0) = 1$  are special and are referred to as the strong resonant cases. If  $\lambda$  is third or fourth root of unity there will be additional resonant terms in formulae (2.5.3).  $\square$

**Remark 2.5.4.** As is well known, if a saddle node bifurcation occurs at  $\mu = \mu_0$  it means that in case of  $\mu < \mu_0$  there are no fixed points but when  $\mu$  passes through  $\mu_0$  two branches of fixed points are born, one branch of stable points, one branch of unstable points.

If the fixed point undergoes a flip bifurcation at  $\mu = \mu_0$  we have (in the supercritical case) that the fixed point loses its stability at  $\mu = \mu_0$  and that a stable period 2 orbit is created.

Theorem 2.5.1 says that when  $(x_0^*, x_1^*)$  undergoes a Hopf bifurcation at  $\mu = \mu_0$  a closed invariant curve surrounding  $(x_0^*, x_1^*)$  is established whenever  $\mu > \mu_0$ ,  $|\mu - \mu_0|$  small.  $\square$

## TURN TO THE EXPERTS FOR SUBSCRIPTION CONSULTANCY

Subscribe is one of the leading companies in Europe when it comes to innovation and business development within subscription businesses.

We innovate new subscription business models or improve existing ones. We do business reviews of existing subscription businesses and we develop acquisition and retention strategies.

Learn more at [linkedin.com/company/subscribe](https://www.linkedin.com/company/subscribe) or contact  
Managing Director Morten Suhr Hansen at [mha@subscribe.dk](mailto:mha@subscribe.dk)

**SUBSCRIBE** - to the future



**Remark 2.5.5.** Much of the theory of Hopf bifurcations for maps have been established by Neimark and Sacker, cf. Sacker (1964, 1965) and Neimark and Landa (1992). Therefore, following Kuznetsov (2004), the Hopf bifurcation is often referred to as the Neimark-Sacker bifurcation, see for example Van Dooren and Metz (1998), King and Schaffer (1999), Kuznetsov (2004), Zhang and Tian (2008), and Moore (2008).  $\square$

**Sketch of proof, Theorem 2.5.1.** Let  $(x_0^*, x_1^*)$  be the fixed point of the two-dimensional map  $\mathbf{x} \rightarrow f(\mathbf{x})$  ( $\mathbf{x} = (x_0, x_1)^T$ ) and assume that the eigenvalues of the Jacobian  $Df(x_0^*, x_1^*)$  are  $\lambda, \bar{\lambda} = a_1 \pm a_2 i$ . Next, define the  $2 \times 2$  matrix  $T$  which columns are the real and imaginary parts of the eigenvectors corresponding to the eigenvalues at the bifurcation. Then, after expanding the right-hand side of the map in a Taylor series, applying the change of coordinates  $(\hat{x}_0, \hat{x}_1) = (x_0 - x_0^*, x_1 - x_1^*)$  (in order to bring the bifurcation to the origin) together with the transformations

$$\begin{pmatrix} \hat{x}_0 \\ \hat{x}_1 \end{pmatrix} = T \begin{pmatrix} x \\ y \end{pmatrix} \quad \begin{pmatrix} x \\ y \end{pmatrix} = T^{-1} \begin{pmatrix} \hat{x}_0 \\ \hat{x}_1 \end{pmatrix}$$

our original map may be cast into standard form at the bifurcation as

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} \cos 2\pi\theta & -\sin 2\pi\theta \\ \sin 2\pi\theta & \cos 2\pi\theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} R_1(x, y) \\ R_2(x, y) \end{pmatrix} \quad (2.5.4)$$

where  $\lambda, \bar{\lambda}$  equal  $\exp(2\pi i\theta), \exp(-2\pi i\theta)$  respectively, and  $\theta = \arctan(a_2/a_1)$ . Our next goal is to simplify the higher order terms  $R_1$  and  $R_2$ . This will be done by use of normal form calculations (near identity transformations). The calculations are simplified if they first are complexified. Thus we introduce

$$\begin{aligned} x' &= \cos 2\pi\theta x - \sin 2\pi\theta y + R_1(x, y) \\ y' &= \sin 2\pi\theta x + \cos 2\pi\theta y + R_2(x, y) \\ z &= x + yi \quad z' = x' + y'i \quad R = R_1 + R_2 i \end{aligned}$$

and rewrite (2.5.4) as

$$f : \mathbb{C} \rightarrow \mathbb{C}, \quad z \rightarrow f(z, \bar{z}) = e^{2\pi\theta i} z + R(z, \bar{z}) \quad (2.5.5)$$

where the remainder is on the form

$$R(z, \bar{z}) = R^{(k)}(z, \bar{z}) + O(|z|^{k+1})$$

Here,  $R^{(k)} = r_1^{(k)} z^k + r_2^{(k)} z^{k-1} \bar{z} + \dots + r_{k+1}^{(k)} \bar{z}^k$ .

Next, define

$$z = Z(w) \quad w = W(z) = Z^{-1}(z) \quad (2.5.6)$$

Then

$$z' = f(z) = f(Z(w)) \quad (2.5.7)$$

which in turn implies

$$w' = \hat{f}(w) = Z^{-1}(z') = (Z^{-1} \circ f \circ Z)(w) \quad (2.5.8)$$

Now, we introduce the near identity transformation

$$z = Z(w) = w + P^{(k)}(w) \quad (2.5.9)$$

and claim that

$$w = z - P^{(k)}(z) + O(|z|^{k+1}) = W(z) \quad (2.5.10)$$

This is nothing but a consequence of (2.5.9). Indeed we have

$$\begin{aligned} w &= z - P^{(k)}(w) = z - P^{(k)}(W(z)) \\ &= w + P^{(k)}(w) - P^{(k)}(w + P^{(k)}(w)) \\ &= w + \text{terms of order higher than } k \end{aligned}$$

Thus, we may now by use of the relations

$$\begin{aligned} f(z) &= e^{2\pi\theta i} z + R^{(k)}(z) + \text{h.o.} \\ Z(w) &= w + P^{(k)}(w) \\ Z^{-1}(z') &= z' - P^{(k)}(z') + \text{h.o.} \end{aligned}$$

(where h.o. means higher order) compute  $\hat{f}(w)$ . This is done in two steps.

First,

$$z' = (f \circ Z)(w) = e^{2\pi\theta i} w + e^{2\pi\theta i} P^{(k)}(w) + R^{(k)}(w + \dots)$$

Then

$$\begin{aligned}\hat{f}(w) &= (Z^{-1} \circ f \circ Z)(w) = z' - P^{(k)}(z') + \text{h.o.} \\ &= e^{2\pi\theta i} w + e^{2\pi\theta i} P^{(k)}(w) + R^{(k)}(w + \dots) - P^{(k)}(e^{2\pi\theta i} w) + \text{h.o.}\end{aligned}\tag{2.5.11}$$

Next, we want to choose constants in order to remove as many terms in  $R^{(k)}(w)$  as possible. To this end let  $H^k$  be polynomials of homogeneous degree  $k$  in  $w, \bar{w}$  and consider the map

$$K : H^k \rightarrow H^k \quad K(P) = e^{2\pi\theta i} P(w) - P(e^{2\pi\theta i} w)\tag{2.5.12}$$

Clearly,  $w^l \bar{w}^{k-l}$  is a basis for  $H^k$  and we have

$$\begin{aligned}K(w^l \bar{w}^{k-l}) &= e^{2\pi\theta i} w^l \bar{w}^{k-l} - e^{2\pi\theta i l} w^l e^{-2\pi\theta i(k-l)} \bar{w}^{k-l} \\ &= [e^{2\pi\theta i} - e^{2\pi\theta i(2l-k)}] w^l \bar{w}^{k-l} \\ &= \lambda w^l \bar{w}^{k-l}\end{aligned}$$

where  $k = 2, 3, 4, \dots, 0 \leq l \leq k$ .

What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



From this we conclude that terms in  $R^{(k)}(w)$  of the form  $w^l \bar{w}^{k-l}$  such that  $\lambda(\theta, k, l) = 0$  cannot be removed by near identity transformations. There are two cases to consider: (A)  $\theta$  irrational, and (B)  $\theta$  rational.

(A) Assume  $\theta$  irrational. Then  $\lambda = 0 \Leftrightarrow 2l = k + 1$  thus  $k$  is an odd number. Here  $k = 1$  corresponds to the linear term and the next unremoval terms are proportional to  $w^2 \bar{w}$  and  $w|w|^4$  (i.e. third and fifth order terms).

(B) Suppose  $\theta = \mu/r$  rational,  $\mu, r \in \mathbb{N}$ ,  $\mu/r$ . Then  $\lambda = 0 \Leftrightarrow (2l - (k + 1))\mu/r = m$  where  $m \in \mathbb{Z}$ . This implies  $(2l - (k + 1))\mu = mr$ . Therefore  $r$  must be a factor in  $(2l - (k + 1))$ . Thus the smallest  $k$  ( $l = 0$ ), equals  $r - 1$  which means that the first unremoval terms are proportional to  $\bar{w}^{r-1}$ . When  $r = 2$  the flip occurs. The cases  $r = 3, 4$  which corresponds to eigenvalues of third and fourth root of unity respectively are special (cf. Remark 2.5.3 after Theorem 2.5.1.)

Now, considering the generic case,  $\theta$  irrational, we may through normal form calculations remove all terms in  $R^{(k)}$  except from those which are proportional to  $w^2 \bar{w}$  and  $w|\bar{w}|^4$ , hence (2.5.5) may be cast into normal form as

$$z' = f(z) = e^{2\pi\theta i} z(1 + \alpha\mu + \beta|z|^2) + \mathcal{O}(5) \quad (2.5.13)$$

where  $\alpha$  and  $\beta$  are given complex numbers. Now introducing polar coordinates  $(r, \varphi)$ , (2.5.13) may after first neglecting terms of  $\mathcal{O}(5)$  and higher and then neglecting terms of  $\mathcal{O}(\mu^2, \mu r^2, r^4)$  be expressed as


$$r' = r(1 + d\mu + ar^2) \quad (2.5.14a)$$

$$\varphi' = \varphi + c + br^2 \quad (2.5.14b)$$

which is nothing but formulae (2.5.3) in the theorem.

Finally, observe that the fixed point  $r^*$  of (2.5.14a) is

$$r^* = \sqrt{-\frac{d\mu}{a}} \quad (2.5.15)$$



**Figure 12:** The outcome of a supercritical Hopf bifurcation. A point close to the unstable fixed point  $x$  moves away from  $x$  and approaches the attracting curve (indicated by a solid line). In the same way an initial point located outside the curve is also attracted.

Thus, if  $a$  and  $d$  have opposite signs we obtain an invariant curve for  $\mu > 0$ . In case of equal signs the curve exists for  $\mu < 0$ . Hence, the truncated map (2.5.14a) possesses an invariant curve. Moreover, the eigenvalue of the linearization of (2.5.14a) is  $\sigma = 1 - 2d\mu$ . Consequently, whenever  $a < 0, \mu > 0, d > 0$  and  $d\mu$  small,  $r^*$  is an attracting curve which corresponds to a supercritical bifurcation. This is displayed in Figure 12.  $\square$

**Remark 2.5.6.** To complete the proof of Theorem 2.5.1 we must show that the full system (2.5.13) possesses an invariant closed curve too. The basic idea here is to set up a graph transform of any closed curve (containing higher order terms) near  $r^*$  and show that this graph transform has a fixed graph close to  $r^*$ . However, in this procedure there are technical difficulties involved which are beyond the scope of this book, cf. the original work by Sacker (1964).  $\square$

Referring to section 1.5 where we treated the flip bifurcation we stated and proved a theorem (Theorem 1.5.1) where we gave conditions for the flip to be supercritical. Regarding the Hopf bifurcation there exists a similar theorem which was first proved by Wan (1978).

**Theorem 2.5.2 (Wan).** Consider the  $C^3$  map  $K : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  on standard form

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} \quad (2.5.16)$$

with eigenvalues  $\lambda, \bar{\lambda} = e^{\pm i\theta}$ . Then the Hopf bifurcation is supercritical whenever the quantity  $d$  (cf. (2.5.2)) in Theorem 2.5.2 is positive and the quantity  $a$  (cf. (2.5.14a)) is negative.  $a$  may be expressed as

$$a = -Re \left[ \frac{(1 - 2\lambda)\bar{\lambda}^2}{1 - \lambda} \xi_{11}\xi_{20} \right] - \frac{1}{2} |\xi_{11}|^2 - |\xi_{02}|^2 + Re(\bar{\lambda}\xi_{21}) \quad (2.5.17)$$

where

$$\xi_{20} = \frac{1}{8} [(f_{xx} - f_{yy} + 2g_{xy}) + i(g_{xx} - g_{yy} - 2f_{xy})]$$

$$\xi_{11} = \frac{1}{4} [(f_{xx} + f_{yy}) + i(g_{xx} + g_{yy})]$$

$$\xi_{02} = \frac{1}{8} [(f_{xx} - f_{yy} - 2g_{xy}) + i(g_{xx} - g_{yy} + 2f_{xy})]$$

$$\xi_{21} = \frac{1}{16} [(f_{xxx} + f_{xyy} + g_{xxy} + g_{yyx}) + i(g_{xxx} + g_{xyy} - f_{xxy} - f_{yyx})]$$

□

For a formal proof we refer to Wan's original paper (Wan, 1978).

(The idea of the proof is simple enough: we start with the original map, write it on standard form (i.e. (2.5.16)) and for each of the near identity transformations we then perform we express the new variables in terms of the original ones, thereby obtaining  $a$  in (2.5.14a) expressed in terms of the original quantities. The problem of course is that the calculations involved are indeed cumbersome and time-consuming as formulae (2.5.17) suggests.)

**gaiTEYE**  
Challenge the way we run

**EXPERIENCE THE POWER OF  
FULL ENGAGEMENT...**

**RUN FASTER.  
RUN LONGER..  
RUN EASIER...**

**READ MORE & PRE-ORDER TODAY  
WWW.GAITEYE.COM**



**Example 2.5.1.** Consider the stage-structured cod model proposed by Wikan and Eide (2004).

$$x_{1,t+1} = F e^{-\beta x_{2,t}} x_{2,t} + (1 - \mu_1) x_{1,t} \quad (2.5.18)$$

$$x_{2,t+1} = P x_{1,t} + (1 - \mu_2) x_{2,t}$$

Here the cod stock  $x$  is split into one immature part  $x_1$  and one mature part  $x_2$ .  $F$  is the density independent fecundity of the mature part while  $\beta$  measures the “strength” of cannibalism from the mature population upon the immature population.  $P$  is the survival probability from the immature stage to the mature stage and  $\mu_1, \mu_2$  are natural death rates. We further assume:  $0 < P \leq 1, 0 < \mu_1, \mu_2 < 1, \beta > 0, F > 0$  and  $FP > \mu_1 \mu_2$ .

Assuming  $x_1^* = x_{1,t+1} = x_{1,t}$  and  $x_2^* = x_{2,t+1} = x_{2,t}$  the fixed point of (2.5.18) is found to be

$$(x_1^*, x_2^*) = \left[ \frac{\mu_2}{\beta P} \ln \left( \frac{FP}{\mu_1 \mu_2} \right), \frac{1}{\beta} \ln \left( \frac{FP}{\mu_1 \mu_2} \right) \right] \quad (2.5.19)$$

The eigenvalue equation of the linearized map becomes (we urge the reader to work through the details)

$$\lambda^2 - (2 - \mu_1 - \mu_2)\lambda + (1 - \mu_1)(1 - \mu_2) - \mu_1 \mu_2 (1 - \beta x_2^*) = 0 \quad (2.5.20)$$

Now, defining  $a_1 = -(2 - \mu_1 - \mu_2), a_2 = (1 - \mu_1)(1 - \mu_2) - \mu_1 \mu_2 (1 - \beta x_2^*)$  and appealing to the Jury criteria (2.1.14) it is straightforward to show that the fixed point is stable as long as the inequalities

$$\beta \mu_1 \mu_2 x_2^* > 0 \quad (2.5.21a)$$

$$2(2 - \mu_1 - \mu_2) + \mu_1 \mu_2 \beta x_2^* > 0 \quad (2.5.21b)$$

$$\mu_1 + \mu_2 - \beta \mu_1 \mu_2 x_2^* > 0 \quad (2.5.21c)$$

hold. Clearly, (2.5.21a) and (2.5.21b) hold for any positive  $x_2^*$ . Thus, there will never be a transfer from stability to instability through a saddle-node or a flip bifurcation. (2.5.21c) is valid in case of  $x_2^*$  sufficiently small. Hence, the fixed point is stable in case of small equilibrium populations. However, if  $x_2^*$  is increased, as a result of increasing  $F$  which we from now on will use as our bifurcation parameter, it is clear that  $(x_1^*, x_2^*)$  will lose its stability at the threshold

$$x_2^* = \frac{\mu_1 + \mu_2}{\beta \mu_1 \mu_2} \quad (2.5.22a)$$

or alternatively when

$$F = \frac{\mu_1 \mu_2}{P} e^{(\mu_1 + \mu_2)/\mu_1 \mu_2} \quad (2.5.22b)$$

Consequently, the fixed point will undergo a Hopf bifurcation at instability threshold and the complex modulus 1 eigenvalues become

$$\lambda, \bar{\lambda} = \frac{2 - \mu_1 - \mu_2}{2} \pm \frac{b}{2} i \quad (2.5.23)$$

where  $b = \sqrt{4(\mu_1 + \mu_2) - (\mu_1 + \mu_2)^2}$ .

In order to show that the Hopf bifurcation is supercritical we have to compute  $d$  (defined through (2.5.2)) and  $a$  (defined through (2.5.17)) and verify that  $d > 0$  and  $a < 0$ .

By first computing  $\lambda$  from (2.5.20) we find

$$|\lambda| = \sqrt{(1 - \mu_1)(1 - \mu_2) - \mu_1 \mu_2 (1 - \beta x_2^*)} \quad (2.5.24)$$

which implies

$$\frac{d}{dF} |\lambda| = \frac{1}{2\sqrt{(1 - \mu_1)(1 - \mu_2) - \mu_1 \mu_2 (1 - \beta x_2^*)}} \cdot \frac{\mu_1 \mu_2}{F}$$

and since the square root is equal to 1 at bifurcation and  $F$  is given by (2.5.22b) we obtain

$$\frac{d}{dF} |\lambda| = \frac{1}{2} P e^{-(\mu_1 + \mu_2)/\mu_1 \mu_2} = d > 0 \quad (2.5.25)$$

which proves that the eigenvalues leave the unit circle through an enlargement of the bifurcation parameter  $F$ .

In order to compute  $a$  we first have to express (2.5.18) on standard form (2.5.16). At bifurcation the Jacobian may be written as

$$J = \begin{pmatrix} 1 - \mu_1 & \frac{1}{P} [(\mu_1 \mu_2 - (\mu_1 + \mu_2))] \\ P & 1 - \mu_1 \end{pmatrix} \quad (2.5.26)$$

so by use of standard techniques the eigenvector  $(z_1, z_2)^T$  belonging to  $\lambda$  is found to be

$$(z_1, z_2)^T = \left( \frac{\mu_1 - \mu_2}{2P} + \frac{b}{2P} i, 1 + 0i \right)^T \quad (2.5.27)$$

and the transformation matrix  $T$  and its inverse may be cast in the form

$$T = \begin{pmatrix} \frac{\mu_2 - \mu_1}{2P} & -\frac{b}{2P} \\ 1 & 0 \end{pmatrix} \quad T^{-1} = \begin{pmatrix} 0 & 1 \\ -\frac{2P}{b} & \frac{\mu_2 - \mu_1}{b} \end{pmatrix} \quad (2.5.28)$$

The next step is to expand  $f(x_2) = Fe^{-\beta x_2}$  up to third order. Then (2.5.18) becomes

$$\begin{aligned} x_{1,t+1} &= \left\{ f(x_2^*) + f'(x_2^*)(x_{2,t} - x_2^*) + \frac{1}{2}f''(x_2^*)(x_{2,t} - x_2^*)^2 \right. \\ &\quad \left. + \frac{1}{6}f'''(x_2^*)(x_{2,t} - x_2^*)^3 \right\} x_{2,t} + (1 - \mu_1)x_{1,t} \\ x_{2,t+1} &= P_{x,t} + (1 - \mu_2)x_{2,t} \end{aligned}$$

and by introducing the change of coordinates  $(\hat{x}_1, \hat{x}_2) = (x_1 - x_1^*, x_2 - x_2^*)$ , in order to bring the bifurcation to the origin, the result is

$$\begin{aligned} \hat{x}_{1,t+1} &= (1 - \mu_1)\hat{x}_{1,t} + \frac{1}{P}\mu_1\mu_2(1 - \beta x_2^*)\hat{x}_{2,t} - \frac{\beta}{P}\mu_1\mu_2\left(1 - \frac{\beta}{2}x_2^*\right)\hat{x}_{2,t}^2 \\ &\quad + \frac{\beta^2}{P}\mu_1\mu_2\left(\frac{1}{2} - \frac{\beta}{6}x_2^*\right)\hat{x}_{2,t}^3 \end{aligned} \quad (2.5.29a)$$

$$\hat{x}_{2,t+1} = P\hat{x}_{1,t} + (1 - \mu_2)\hat{x}_{2,t} \quad (2.5.29b)$$

where all terms of higher order than three have been neglected.

This e-book  
is made with  
**SetaPDF**



**SETASIGN**



PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)



Finally, by applying the transformations

$$\begin{pmatrix} \hat{x}_1 \\ \hat{x}_2 \end{pmatrix} = T \begin{pmatrix} u \\ v \end{pmatrix} \quad \begin{pmatrix} u \\ v \end{pmatrix} = T^{-1} \begin{pmatrix} \hat{x}_1 \\ \hat{x}_2 \end{pmatrix} \quad (2.5.30)$$

we obtain after some algebra that the original map (2.5.18) may be cast into standard form as

$$\begin{aligned} u_{t+1} &= \frac{2 - \mu_1 - \mu_2}{2} u_t - \frac{b}{2} v_t \\ u_{t+1} &= \frac{b}{2} u_t + \frac{2 - \mu_1 - \mu_2}{2} v_t + g(u_t, v_t) \end{aligned} \quad (2.5.31)$$

where

$$g(u, v) = \frac{2\beta}{b} \mu_1 \mu_2 \left(1 - \frac{\beta}{2} x_2^*\right) u^2 - \frac{2\beta^2}{b} \mu_1 \mu_2 \left(\frac{1}{2} - \frac{\beta}{6} x_2^*\right) u^3$$

Now at last, we are ready to compute the terms in formulae (2.5.17)

$$g_{uu} = \frac{4\beta}{b} \mu_1 \mu_2 A \quad g_{uuu} = -\frac{12\beta^2}{b} \mu_1 \mu_2 B$$

where  $A = 1 - (\beta/2)x_2^*$ ,  $B = (1/2) - (\beta/6)x_2^*$ . This yields:

$$\xi_{20} = \frac{1}{8} i g_{uu} \quad \xi_{11} = \frac{1}{4} i g_{uu} \quad \xi_{02} = \frac{1}{8} i g_{uu} \quad \xi_{21} = \frac{1}{16} i g_{uuu}$$

and

$$\begin{aligned} \operatorname{Re} \left[ \frac{(1 - 2\lambda)\bar{\lambda}^2}{1 - \lambda} \xi_{11}\xi_{20} \right] &= -\frac{g_{uu}^2}{256(\mu_1 + \mu_2)} \times \\ &\quad [3(\mu_1 + \mu_2) [(2 - u_1 - u_2)^2 - b^2] - 2(2 - \mu_1 - \mu_2)b^2] \end{aligned}$$

so finally, by computing  $|\xi_{11}|^2 = (1/16)g_{uu}^2$ ,  $|\xi_{02}|^2 = (1/64)g_{uu}^2$ ,  $\operatorname{Re}(\bar{\lambda}\xi_{21}) = (1/32)bg_{uuu}$  and inserting into (2.5.17) we eventually arrive at

$$a = -\frac{\beta^2}{16(\mu_1 + \mu_2)} \left\{ (2\mu_1\mu_2)^2 + (\mu_1 + \mu_2) [(2\mu_1\mu_2 - (\mu_1 + \mu_2))^2 - \mu_1\mu_2] \right\} \quad (2.5.32)$$

which is negative for all  $0 < \mu_1, \mu_2 < 1$ . Consequently, the fixed point (2.5.19) undergoes a supercritical Hopf bifurcation at the threshold (2.5.22a,b) (i.e. when  $(x_1^*, x_2^*)$  fails to be stable through an increase of  $F$ , a closed invariant attracting curve surrounding  $(x_1^*, x_2^*)$  is established). For further analysis of (2.5.18) we refer to the original paper by Wikan and Eide (2004) but also confer Govaerts and Ghaziani (2006) where a numerical study of the model may be obtained.  $\square$

In the next exercise most of the cumbersome and time-consuming calculations we had to perform in Example 1.5.1 are avoided.

**Exercise 2.5.1.** Assume that the parameter  $\mu > 1$  and consider the map

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{pmatrix} y \\ \mu y(1-x) \end{pmatrix} \quad (2.5.33)$$

a) Show that the nontrivial fixed point

$$(x^*, y^*) = \left( \frac{\mu-1}{\mu}, \frac{\mu-1}{\mu} \right)$$

b) Compute the Jacobian and show that the eigenvalue equation may be expressed as

$$\lambda^2 - \lambda + \mu - 1 = 0$$

c) Use the Jury criteria (2.1.14) and show that the fixed point is stable whenever  $1 < \mu < 2$  and that a Hopf bifurcation occurs at the threshold  $\mu = 2$ .

d) Show that  $|\lambda| = \sqrt{\mu-1}$  and moreover that

$$\frac{d}{d\mu} |\lambda|_{\mu=2} > 0$$

which proves that the eigenvalues leave the unit circle at bifurcation threshold.

e) Assuming  $\mu = 2$ , apply the change of coordinates  $(\hat{x}, \hat{y}) = (x - (1/2), y - (1/2))$  together with the transformations

$$\begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix} = T \begin{pmatrix} u \\ v \end{pmatrix} \quad \begin{pmatrix} u \\ v \end{pmatrix} = T^{-1} \begin{pmatrix} \hat{x} \\ \hat{y} \end{pmatrix}$$

where

$$T = \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ 1 & 0 \end{pmatrix}$$

(verify that the columns in  $T$  are the real and imaginary parts of the eigenvectors belonging to the eigenvalues of the Jacobian respectively) and show that (2.5.33) may be written on standard form at bifurcation threshold as

$$\begin{pmatrix} u \\ v \end{pmatrix} \rightarrow \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} f(u, v) \\ g(u, v) \end{pmatrix} \quad (2.110)$$


where  $f(u, v) = -u^2 - \sqrt{3}uv$  and  $g(u, v) = (1/\sqrt{3})u^2 + uv$ .

- f) Referring to Theorem 2.5.2 show that the quantity  $a$  defined in (2.5.17) is negative, hence that in case of  $\mu > 2$ ,  $|\mu - 2|$  small, there exists an attracting curve surrounding the unstable fixed point  $(x^*, y^*)$ . □

# Free eBook on Learning & Development

By the Chief Learning Officer of McKinsey

[Download Now](#)



[Click on the ad to read more](#)

**Exercise 2.5.2 (Strong resonant case I).** Consider the two-age structured population model

$$(x_1, x_2) \rightarrow (F_2 x_2, P e^{-x_1} x_1) \quad (2.5.35)$$

where  $0 < P \leq 1$ ,  $F_2 > 0$  and  $PF_2 > 1$ .

- Show that the fixed point  $(x_1^*, x_2^*) = (\ln(PF_2), (1/F_2) \ln(PF_2))$ .
- Show that the eigenvalue equation may be cast in the form  $\lambda^2 + x_1^* - 1 = 0$  and further that a Hopf bifurcation takes place at the threshold  $x_1^* = 2$  (or equivalently when  $F_2 = (1/P) \exp(2)$ ).
- Show that  $\lambda$  equals fourth root of unity at bifurcation threshold.

Note that the result obtained in c) violates assumption (2.5.1) in Theorem 2.5.1 which of course means that neither Theorem 2.5.1 nor Theorem 2.5.2 applies to map (2.5.35). We urge the reader to perform numerical experiments where  $F_2 > (1/P) \exp(2)$  in order to show that when  $(x_1^*, x_2^*)$  fails to be stable, an exact 4-periodic orbit with small amplitude is established. (For further reading, cf. Wikan (1997).) □

**Exercise 2.5.3 (Strong resonant case II).** Repeat the analysis from the previous exercise on the map

$$(x_1, x_2) \rightarrow (F e^{-(x_1+x_2)}(x_1 + x_2), x_1) \quad (2.5.36)$$

Hint:  $\lambda$  equals third root of unity at bifurcation threshold. □

As is shown in the sketch of proof of Theorem 2.5.1 most terms in (2.5.5) may be removed by a series of near identity transformations. In the next exercise the reader is actually asked to perform such transformations.

**Exercise 2.5.4.** Let  $\lambda^j \neq 1$ ,  $j = 1, 2, 3, 4, 5$  and consider

$$(i) \quad z_{t+1} = \lambda z_t + \alpha_1 z_t^2 + \alpha_2 z_t \bar{z}_t + \alpha_3 \bar{z}_t^2 + \mathcal{O}(3)$$

- Apply the near identity transformation (cf. (2.5.9))

$$z = w + \beta_1 w^2 + \beta_2 w \bar{w} + \beta_3 \bar{w}^2$$



together with (cf. (2.5.10))

$$w = z - (\beta_1 z^2 + \beta_2 z \bar{z} + \beta_3 \bar{z}^2)$$

and show that (i) may be written as

$$(ii) \quad w_{t+1} = \lambda w_t + (\lambda \beta_1 + \alpha_1 - \beta_1 \lambda^2) w_t^2 \\ + (\lambda \beta_2 + \alpha_2 - \beta_2 \lambda \bar{\lambda}) w_t \bar{w}_t + (\lambda \beta_3 + \alpha_3 - \beta_3 \bar{\lambda}^2) \bar{w}_t^2 + \mathcal{O}(3)$$

b) Show that if we choose

$$\beta_1 = -\frac{\alpha_1}{\lambda(1-\lambda)} \quad \beta_2 = -\frac{\alpha_2}{\lambda(1-\bar{\lambda})} \quad \beta_3 = -\frac{\alpha_3}{\lambda-\bar{\lambda}^2}$$

then all second order terms in (ii) will disappear. Thus, after one near identity transformation we have a system on the form (where we for notational convenience still use  $z$  as variable)

$$(iii) \quad z_{t+1} = \lambda z_t + \beta_1 z_t^3 + \beta_2 z_t^2 \bar{z}_t + \beta_3 z_t \bar{z}_t^2 + \beta_4 \bar{z}_t^3 + \mathcal{O}(4)$$

c) Apply

$$z = w + a_1 w^3 + a_2 w^2 \bar{w} + a_3 w \bar{w}^2 + a_4 \bar{w}^3$$

$$w = z - (a_1 z^3 + a_2 z^2 \bar{z} + a_3 z \bar{z}^2 + a_4 \bar{z}^3)$$

on (iii) and show that if we choose

$$a_1 = -\frac{\beta_1}{\lambda(1-\lambda^2)} \quad a_3 = -\frac{\beta_2}{\lambda(1-\bar{\lambda}^2)} \quad a_4 = -\frac{\beta_4}{\lambda-\bar{\lambda}^3}$$

then the  $w^3$ ,  $w\bar{w}^2$  and  $\bar{w}^3$  terms will disappear. Note that we cannot use

$$a_2 = -\frac{\beta_2}{\lambda(1-\lambda\bar{\lambda})}$$

because  $1 - \lambda\bar{\lambda} = 0$  for any  $\lambda$  located on the boundary of the unit circle.

d) After two near identity transformations our system is on the form

$$(iv) \quad z_{t+1} = \lambda z_t + \beta_2 z_t^2 \bar{z}_t + \mathcal{O}(4)$$

Write out all fourth order elements and perform a new near identity transformation in the same way as in a) and c) and show that all fourth order terms may be removed, hence that our system may be cast in the form (normal form!)

$$(v) \quad z_{t+1} = \lambda z_t + \beta_2 z_t^2 \bar{z}_t + \mathcal{O}(5)$$

□

**Remark 2.5.7.** Note that Exercise 2.5.4 in many respects offers an equivalent way of establishing the normal form (2.5.13). Moreover, if  $\lambda^3 = 1$ , the denominator in the expression for  $\beta_3$  becomes zero, hence the terms  $\bar{w}_t^2$  in (ii) is not removable. Consequently, there will be an additional resonant term on the form  $\alpha \bar{z}_t^2$  in (v). In case of  $\lambda^4 = 1$  or  $\lambda^5 = 1$  the additional terms are  $\gamma \bar{z}_t^3$  and  $\delta \bar{z}_t^4$  respectively. For further reading we refer to Kuznetsov (2004) and Kuznetsov and Meijer (2005). □

We close this section by once again emphasizing that the outcome of a supercritical Hopf bifurcation is that when the fixed point fails to be stable an attracting invariant curve which surrounds the fixed point is established. In section 2.8 we shall focus on the nonstationary dynamics on such a curve as well as possible routes to chaos. However, before we turn to those questions we shall in section 2.6 present an analysis of the Horseshoe map where we once again invoke symbolic dynamics and in section 2.7 we shall explain how we may analyse the nature of bifurcations in higher dimensional problems.

www.sylvania.com

We do not reinvent the wheel we reinvent light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.


Light is OSRAM

OSRAM SYLVANIA

Click on the ad to read more

## 2.6 Symbolic dynamics III (The Horseshoe map)

As we have seen maps may possess both fixed points and periodic points, and through Theorem 2.5.1 we have established that the dynamics may be restricted to invariant curves as well. However, in Part I our analysis also revealed other types of invariant hyperbolic sets. To be more concrete we showed in Section 1.9 that whenever  $\mu > 2 + \sqrt{5}$  the quadratic map possessed an invariant set of points  $\Lambda$  (a Cantor set) that never left the unit interval through iterations. Our next goal is to discuss a similar phenomenon in case of a two-dimensional map, the Horseshoe map, which is due to Smale (1963, 1967). There are several ways of visualizing the Horseshoe. We prefer the way presented in Guckenheimer and Holmes (1990),



**Figure 13:** a) The Horseshoe map  $f$ . b) The inverse  $f^{-1}$ . c) The image of four thin horizontal strips under  $f^2$ .

**Example 2.6.1 (The Horseshoe map).** Consider the unit square  $H = [0, 1] \times [0, 1]$ , see Figure 13a, and assume that we perform two operations on  $H$ : (1) a linear expansion of  $H$  by a factor  $\alpha$ ,  $\alpha > 2$ , in the vertical direction and a horizontal contraction by a factor  $\beta$ ,  $0 < \beta < 1/2$ . (2) A folding in such a way that the folding part falls outside  $H$ . The whole process is displayed in Figure 13a. We call this a map  $f : H \rightarrow \mathbb{R}^2$  and restricted to  $H$  we may express the two vertical strips as  $f(H) \cap H$ .

If we reverse the process (folding, stretching and contracting) we see from Figure 13b that we obtain two horizontal strips  $H_0$  and  $H_1$  and each of them has thickness  $\alpha^{-1}$ . Also note that the inverse image may be expressed as  $f^{-1}(f(H) \cap H) = f^{-1}(H) \cap H$ . Thus we conclude that on each of the horizontal strips  $H_0$  and  $H_1$ ,  $f$  stretches by a factor  $\alpha$  in the vertical direction and contracts by a factor  $\beta$  in the horizontal direction.

As is clear from Figures 13a,b and the text above, when  $f$  is iterated most points will leave  $H$  after a finite number of iterations. However, as we shall see (just as we did in the corresponding “one-dimensional example” in Section 1.9) there is a set

$$\Lambda = \{x \mid f^i(x) \in H\} \quad i \in \mathbb{Z}$$

which never leaves  $H$ . Now, let us describe the structure of  $\Lambda$ . First, observe that  $f$  stretches both  $H_0$  and  $H_1$  vertically by  $\alpha$  so that  $f(H_0)$  and  $f(H_1)$  both intersect  $H_0$  and  $H_1$  (Figure 13b). Therefore, points in  $H_0$  must have been mapped into  $H_0$  by  $f$  from two thinner strips, each of width  $\alpha^{-2}$  contained in  $H_0$ . The same is of course true for points in  $H_1$ , so after applying  $f$  twice on the four horizontal strips of widths  $\alpha^{-2}$  in Figure 13c the result is four thin vertical strips each of width  $\beta^2$  as displayed in Figure 13c. (Note, that after only one iteration of  $f$  the result is four rectangles, each of height  $\alpha^{-1}$  and width  $\beta$ .) Moreover, since  $H_0 \cup H_1 = f^{-1}(H \cap f(H))$  the union of the four thinner strips may be written as  $f^{-2}(H \cap f(H) \cap f^2(H))$ , and proceeding in this way  $f^{-n}(H \cap f(H) \cap \dots \cap f^n(H))$  must be the union of  $2^n$  such strips where each strip has a thickness of  $\alpha^{-n}$ . Since  $\alpha > 2$  the thickness of each of the  $2^n$  strips goes to zero when  $n \rightarrow \infty$ . Now, consider one of the  $2^n$  horizontal strips. Each time  $f$  is applied on the strip it is stretched by  $\alpha$  in the vertical direction and contracted by  $\beta$  in the horizontal direction so the image under  $f^n$  must be a strip of length 1 in the vertical direction and width  $\beta^n$  in the horizontal direction, and since  $0 < \beta < 1/2$  the latter tends to zero as  $n \rightarrow \infty$ . Thus, the  $2^n$  horizontal strips are mapped into  $2^n$  vertical strips. The points that will remain in  $H$  forever are those points which are located both in the horizontal and the vertical strips, hence  $\Lambda$  is nothing but the intersection of the horizontal and vertical strips. Moreover,  $\Lambda$  is a Cantor set. Indeed, when  $n \rightarrow \infty$ ,  $\Lambda$  contains just points (no intervals) and these points are not isolated but they are accumulation points of  $\Lambda$  (cf. Definition 1.9.5).

In order to describe the dynamics on  $\Lambda$  let us invoke symbolic dynamics in much of the same way as we did in Section 1.9 and assign a sequence  $\bar{a} = \{a_i\}_{i=-\infty}^{\infty}$  to every point  $x \in \Lambda$ . We also define another sequence  $\bar{b}$  through  $b_i = a_{i+1}$ . Thus  $\sigma : \Sigma_2 \rightarrow \Sigma_2$ .  $\sigma(\bar{a}) = \bar{b}$  is the shift map. The itinerary of  $x$ ,  $\phi : \Lambda \rightarrow \Sigma_2$  is defined as  $\phi(x) = \dots a_{-2}a_{-1}a_0a_1a_2\dots$  and we let  $a_i = 0$  if  $f^i(x) \in H_0$  and  $a_i = 1$  if  $f^i(x) \in H_1$  which means that  $x \in \Lambda$  if and only if  $f^i(x) \in H_{a_i}$  for every  $i$ .

First, observe that since  $f^{i+1}(x) = f^i(f(x))$  then  $\phi(f(x)) = \bar{b}$  which proves that  $\phi$  acts in the same way as  $\sigma$ . Consequently, if we are able to prove that  $\phi$  is a homeomorphism then (according to Definition 1.2.1),  $f$  and  $\sigma$  are topological equivalent maps on  $\Lambda$ . The 1 – 1 and continuity properties of  $\phi$  may be proved along the following line. Let  $S_V = S_V(b_{-m}, b_{-m+1}, \dots, b_{-1})$  be the central set of  $x$ 's such that  $f^i(x)$  is contained in one of the  $2^m$  vertical strips in  $H \cap f(H) \cap \dots \cap f^m(H)$  and let  $S_H(b_0, \dots, b_n)$  be the central set of  $x$ 's contained in a horizontal strip. Then  $S = S(b_{-m}, \dots, b_0, \dots, b_n) = S_V \cap S_H$  is the set of  $x$ 's such that  $f^i(x) \in H_{b_i}$  and clearly  $S$  must be a rectangle with height  $\alpha^{-(n+1)}$  and width  $\beta^m$ . When  $n, m \rightarrow \infty$  all areas  $\rightarrow 0$ . Consequently,  $\phi$  is both continuous and 1 – 1.

Regarding the onto property, following Guckenheimer and Holmes (1990), it suffices to prove that  $S$  is nonempty. To see this, observe that  $f^{n+1}(S_H(b_0, \dots, b_n))$  fills the entire  $S$  in the vertical direction. In particular it intersects both  $S_0$  and  $S_1$  so that  $S_H(b_0, \dots, b_n, b_{n+1})$  must be a nonempty horizontal strip. Moreover, every vertical strip  $S_V$  intersects every  $S_H$  which immediately implies that  $S = S_V \cap S_H$  is nonempty. Consequently,

$$\phi \circ f = \sigma \circ \phi \tag{2.6.1}$$

whenever  $f$  is restricted to  $\Lambda$ . □

Before we leave the Horseshoe map let us emphasize and comment on a few more topics. First, note the difference between the symbol sequence  $\{a_i\}_{i=-\infty}^{\infty}$  defined for the horseshoe and the sequence  $\{a_i\}_{i=0}^{\infty}$  we used in our study of the quadratic map in case of  $\mu > 2 + \sqrt{5}$  (see Section 1.9). Unlike the quadratic map (1.2.1), the two-dimensional horseshoe map is invertible (Figure 13b) so it makes sense to consider backward iteration. Therefore we may use negative indices in order to say which vertical strip  $f(x)$  is located in and positive indices in order to say which horizontal strip  $f(x)$  is contained in. If we glue together  $\{a_i\}_{i=-\infty}^{-1}$  and  $\{a_i\}_{i=0}^{\infty}$  we have a description of the whole orbit.

The shift map  $\sigma$  which in this context often is referred to as the two-sided shift, may be defined as in Example 2.6.1 or in the usual manner as

$$\sigma(\dots a_{-2} a_{-1} \cdot a_0 a_1 a_2 \dots) = (\dots a_{-2} a_{-1} a_0 \cdot a_1 a_2 \dots) \tag{2.6.2}$$

(cf. Definition 1.9.3). The inverse is defined through

$$\sigma^{-1}(\dots a_{-2}a_{-1}a_0 \cdot a_1a_2\dots) = (\dots a_{-2}a_{-1} \cdot a_0a_1a_2\dots) \quad (2.6.3)$$

Periodic points of period  $N$  for  $\sigma$  may be expressed as before. For example, a 3-period orbit may be expressed by the sequence  $\bar{c} = \{\dots 010010010\dots\}$  and  $\sigma^3(\bar{c}) = \bar{c}$ . Moreover, since each element of  $\{a_i\}$  may take two values (0 or 1) a period  $n$  orbit for  $\sigma$  corresponds to  $2^n$  periodic points. From this we may conclude that if  $\sigma^n$  has  $2^n$  periodic points in  $\Sigma_2$ , then from (3.6.1)  $f^n = \phi^{-1} \circ \sigma^n \circ \phi$  has  $2^n$  periodic points in  $\Lambda$ . Actually, these periodic points are unstable points of the saddle type. In order to see this, observe that segments contained in  $H_0$  and  $H_1$  are compressed horizontally by  $\beta$  ( $0 < \beta < 1/2$ ) and stretched by  $\alpha$  ( $\alpha > 2$ ) in the vertical direction. This means that  $f$  restricted to  $H \cap f^{-1}(H)$  is linear so the Jacobian becomes  $J = \text{diag}(\beta, \alpha)$  and if we apply  $f^n$  on one of the  $2^n$  horizontal strips described in Example 2.5.2 the resulting Jacobian may be expressed as  $Df^n = \text{diag}(\beta^n, \alpha^n)$ . Consequently, the eigenvalues are  $\lambda_1 = \beta$  and  $\lambda_2 = \alpha$ , and since  $\lambda_{1,2}$  are real and  $\lambda_1$  is located on the inside of the unit circle and  $\lambda_2$  on the outside the periodic points are saddle points.



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more



The distance function (cf. Proposition 1.9.1) between two sequences  $\bar{a}$  and  $\bar{b}$  in  $\Sigma_2$  is defined as

$$d[\bar{a}, \bar{b}] = \sum_{i=-\infty}^{\infty} \frac{|a_i - b_i|}{2^{|i|}} \quad (2.6.4)$$

where  $|a_i - b_i| = 0$  if  $a_i = b_i$  and  $|a_i - b_i| = 1$  if  $a_i \neq b_i$ . The fact that periodic points for  $\sigma$  are dense in  $\Sigma_2$  may be obtained from (2.6.4) and by use of the same method as in the proof of Proposition 1.9.1. We leave the details to the reader. There are also nonperiodic points for  $\sigma$  in  $\Sigma_2$  which are dense in  $\Sigma_2$ . In order to show this we must prove that the orbit of such a point comes arbitrarily close to any given sequence in  $\Sigma_2$ . Thus, let  $\bar{a} = (\dots a_{-k} \dots a_0 \dots a_k \dots)$  be a given sequence and let  $\bar{b}$  be a sequence whose central block equals the central block of  $\bar{a}$  (i.e.  $a_{-k} = b_{-k}, \dots, a_0 = b_0, \dots, a_k = b_k$ ). Then, from (2.6.4):

$$d[a, b] = \sum_{i=-\infty}^{\infty} \frac{|a_i - b_i|}{2^{|i|}} = \sum_{i=-\infty}^{-k-1} \frac{|a_i - b_i|}{2^{|i|}} + \sum_{i=k+1}^{\infty} \frac{|a_i - b_i|}{2^i} \leq \frac{1}{2^k} + \frac{1}{2^k} = 2^{1-k}$$

Hence, when  $k$  becomes large,  $\bar{b} \rightarrow \bar{a}$  so according to Definition 1.9.4,  $\bar{b}$  represents a dense orbit in  $\Sigma_2$ .

Finally, let us give a few comments on stable and unstable sets of points in  $\Lambda$ . In general, two points  $x_1$  and  $x_2$  are said to be forward asymptotic in a set  $S$  if  $f^n(x_1) \in S, f^n(x_2) \in S$  for all  $n$  and

$$\lim_{n \rightarrow \infty} |f^n(x_1) - f^n(x_2)| = 0 \quad (2.6.5a)$$

If  $f^{-n}(x_1) \in S, f^{-n}(x_2) \in S$  for all  $n$  and

$$\lim_{n \rightarrow \infty} |f^{-n}(x_1) - f^{-n}(x_2)| = 0 \quad (2.6.5b)$$

then  $x_1, x_2$  are said to be backward asymptotic in  $S$ . By use of (2.6.5a,b) we may define the stable set of a point  $x$  in  $S$  as

$$W^S(x) = \{y \mid |f^n(x) - f^n(y)| \rightarrow 0 \text{ as } n \rightarrow \infty\} \quad (2.6.6a)$$

and the unstable set as

$$W^U(x) = \{z \mid |f^{-n}(x) - f^{-n}(z)| \rightarrow 0 \text{ as } n \rightarrow \infty\} \quad (2.6.6b)$$

The shift map makes it easy to describe  $W^S(x)$  and  $W^U(x)$ . For example, if  $x^*$  is a fixed point of  $f$  and  $\phi(x^*) = (\dots a_{-2}^* a_{-1}^* a_0^* a_1^* a_2^* \dots)$  then any point  $y$  whose itinerary is the same as the itinerary of  $x^*$  to the right of an entry  $a_i^*$  is contained in  $W^S(x^*)$ . (2.6.6a) allows us to describe the stable set of points in  $\Lambda$ . Indeed, let  $x^*$  be a fixed point of  $f$  which lies in  $H_0$ . Then  $\phi(x^*) = \{\dots 0000\dots\}$ . Then, since  $f$  contracts in the horizontal direction, any point which is located in a horizontal segment through  $x^*$  must be in  $W^S(x^*)$ . But there are also additional points in  $W^S(x^*)$ . In fact, any point  $p$  which eventually is mapped into the horizontal segment through  $x^*$  after a finite number of iterations  $k$  is also contained in  $W^S(x^*)$  because  $|f^{k+n}(p) - x^*| < \beta^n$ . This implies that the union of all horizontal intervals given by  $f^{-n}(l)$ ,  $n = 1, 2, 3, \dots$ , (where  $l$  is a horizontal segment) lies in  $W^S(x^*)$ . We leave to the reader to describe the set  $W^U(x^*)$ .

—

## 2.7 The center manifold theorem

Recall that in our treatment of the flip bifurcation (cf. section 1.5) we considered one-dimensional maps of the form  $f : \mathbb{R} \rightarrow \mathbb{R}$  and when we studied the Hopf bifurcation in section 2.5 the main theorems were stated for two-dimensional maps  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ . Let us now turn to higher-dimensional maps,  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ . Of course,  $|\lambda| = 1$  at bifurcation in these cases too but how do we determine the nature of the bifurcation involved when the fixed point fails to be hyperbolic?

The main conclusion is that there exists a method which applied to a map on the form  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  reduces the bifurcation problem to a study of a map  $g : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  (Hopf), or  $g : \mathbb{R} \rightarrow \mathbb{R}$  (flip). The cornerstone in the theory which allows this conclusion is the center manifold theorem for maps which we now state.

**Theorem 2.7.1 (Center manifold theorem).** Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be a  $C^k$ ,  $k \geq 2$  map and assume that the Jacobian  $Df(0)$  has a modulus 1 eigenvalue and, moreover, that all eigenvalues of  $Df(0)$  splits into two parts  $\alpha_c, \alpha_s$  such that

$$|\lambda| = \begin{cases} 1 & \text{if } \lambda \in \alpha_c \\ < 1 & \text{if } \lambda \in \alpha_s \end{cases}$$

Further, let  $E_c$  be the (generalized) eigenspace of  $\alpha_c$ ,  $\dim E_c = d < \infty$ . Then there exists a domain  $V$  about 0 in  $\mathbb{R}^n$  and a  $C^k$  submanifold  $W^c$  of  $V$  of dimension  $d$  passing through 0 which is tangent to  $E_c$  at 0 which satisfies:

I) If  $x \in W^c$  and  $f(x) \in V$  then  $f(x) \in W^c$ .

II) If  $f^{(n)}(x) \in V$  for all  $n = 0, 1, 2, \dots$  then the distance from  $f^{(n)}(x)$  to  $W^c$  approaches zero as  $n \rightarrow \infty$ . □

For a proof of Theorem 2.7.1, cf. Marsden and McCracken (1976, p. 28  $\rightarrow$  43). Also cf. the book by Iooss (1979) and the paper by Vanderbauwhede (1987).


When  $E_c$  has dimension two, as it does for the Neimark-Sacker case at criticality, the essence of Theorem 2.7.1 is that there exists an invariant manifold of dimension  $2 \subset \mathbb{R}^n$  which has the eigenspace belonging to the complex eigenvalues as tangent space at the bifurcating nonhyperbolic fixed points. In case of flip bifurcation problems,  $\dim W^C = 1$ . Thus close to the bifurcation, our goal is to restrict the original map to the invariant center manifold  $W^C$  and then proceed with the analysis by using the results in Theorems 2.5.1 and 2.5.2 in case of Hopf bifurcation problems and Theorem 1.5.1 in the flip case.

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more

Let us now in general terms describe how such a restriction may be carried out. To this end, consider our discrete system written in the form

$$\begin{aligned} \mathbf{x}_{t+1} &= A\mathbf{x}_t + F(\mathbf{x}_t, \mathbf{y}_t) \\ \mathbf{y}_{t+1} &= B\mathbf{y}_t + G(\mathbf{x}_t, \mathbf{y}_t) \end{aligned} \quad (2.7.1)$$

where all the eigenvalues of  $A$  are on the boundary of the unit circle and those of  $B$  within the unit circle. (If the system we want to study is not on the form as in (2.7.1) we first apply the procedure in Example 2.5.1, see also the proof of Theorem 2.5.1.)

Now, since the center manifold  $W^C$  is tangent to the (generalized) eigenspace  $E_c$ , we may represent it as a local graph

$$W^C = \{(\mathbf{x}, \mathbf{y})/\mathbf{y} = h(\mathbf{x})\} \quad h(0) = Dh(0) = 0 \quad (2.7.2)$$

and by substituting (2.7.2) into (2.7.1) we have

$$\mathbf{y}_{t+1} = h(\mathbf{x}_{t+1}) = h(A\mathbf{x}_t + F(\mathbf{x}_t, h(\mathbf{x}_t))) = Bh(\mathbf{x}_t) + G(\mathbf{x}_t, h(\mathbf{x}_t))$$

or equivalently

$$h(A\mathbf{x} + F(\mathbf{x}, h(\mathbf{x}))) - Bh(\mathbf{x}) - G(\mathbf{x}, h(\mathbf{x})) = 0 \quad (2.7.3)$$

An explicit expression of  $h(\mathbf{x})$  is out of reach in most cases, but one can approximate  $h$  by its Taylor series at the bifurcation as

$$h(x) = ax^2 + bx^3 + O(x^4) \quad (2.7.4)$$

where the coefficients  $a, b$  are determined through (2.7.3), and finally the restricted map is obtained by inserting the series of  $h$  into (2.7.1).

**Example 2.7.1.** Consider the Leslie matrix model

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \rightarrow \begin{pmatrix} F(1 - \gamma x)^{1/\gamma} & F(1 - \gamma x)^{1/\gamma} \\ P & 0 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \quad (2.125)$$

where  $x = x_1 + x_2$  is the total population. □

(2.7.5) is often referred to as the Deriso-Schnute population model. Note that if  $\gamma \rightarrow 0$ , (2.7.5) is nothing but the Ricker model (see (2.3.4) and Examples 2.4.1 and 2.4.3). If  $\gamma = -1$  we are left with the Beverton and Holt model (see (2.3.5) and Exercise 2.4.1).

Our goal is to show that under the assumptions  $F(1 + P) > 1$ ,  $0 < P < 1/2$ ,  $\gamma > -(1 - P)/2$  the fixed point  $(x_1^*, x_2^*)$  of (2.7.5) will undergo a supercritical flip bifurcation at instability threshold.

We urge the reader to verify the following properties:

$$(x_1^*, x_2^*) = \left( \frac{1}{1 + P} x^*, \frac{1}{1 + P} x^* \right) \quad (2.7.6)$$

where  $x^* = (1/\gamma)[1 - (P + PF)^{-\gamma}]$ . Defining  $f(x) = F(1 - \gamma x)^{1/\gamma}$  the Jacobian becomes

$$\begin{pmatrix} f'x^* + f & f'x^* + f \\ P & 0 \end{pmatrix}$$

where  $f = f(x^*) = 1/(1 + P)$  and  $f' = f'(x^*)$ .

—

Show by use of the Jury criteria (2.1.14) that whenever  $0 < P < 1/2$ ,  $\gamma > -(1 - P)/2$  the fixed point (2.7.6) will undergo a flip bifurcation when  $f'x^* = -2/(1 - P^2)$  and that the Jacobian at bifurcation threshold equals

$$\begin{pmatrix} -\frac{1}{1-P} & -\frac{1}{1-P} \\ P & 0 \end{pmatrix} \quad (2.7.7)$$

and moreover, that the eigenvalues of (2.7.7) are  $\lambda_1 = -1$  and  $\lambda_2 = -P/(1 - P)$ .

Now, in order to show that the flip bifurcation is of supercritical nature we must appeal to Theorem 1.5.1 but since that theorem deals with one-dimensional maps, we first have to express (2.7.5) on the appropriate form (2.7.1) and then perform a center manifold restriction as explained through (2.7.2)-(2.7.4).

The form (2.7.1) is achieved by performing the same kind of calculations as in Example 2.5.1. The eigenvectors belonging to  $\lambda_1$  and  $\lambda_2$  are easily found to be  $(-1/P, 1)^T$  and  $(-1/(1 - P), 1)^T$  respectively so the transformation matrix  $T$  and its inverse become

$$T = \begin{pmatrix} -\frac{1}{P} & -\frac{1}{1-P} \\ 1 & 1 \end{pmatrix} \quad T^{-1} = \begin{pmatrix} \frac{P(1-P)}{2P-1} & \frac{P}{2P-1} \\ -\frac{P(1-P)}{2P-1} & -\frac{1-P}{2P-1} \end{pmatrix} \quad (2.7.8)$$

Further, by expanding  $f$  up to third order, i.e.

$$f(x) \approx f(x^*) + f'(x^*)(x - x^*) + \frac{1}{2}f''(x^*)(x - x^*)^2 + \frac{1}{6}f'''(x^*)(x - x^*)^3$$

and applying the change of coordinates  $(\hat{x}_1, \hat{x}_2) = (x_1 - x_1^*, x_2 - x_2^*)$ , using the fact that  $f'(x^*) = -2/(1 - P^2)$  at bifurcation threshold gives

$$\hat{x}_{1,t+1} = -\frac{1}{1-P}\hat{x}_{1,t} - \frac{1}{1-P}\hat{x}_{2,t} + \{1\}\hat{x}_t^2 + \{2\}\hat{x}_t^3 \quad (2.7.9)$$

$$\hat{x}_{2,t+1} = P\hat{x}_{1,t}$$

where all terms of higher order than 3 have been neglected and  $\{1\}$  and  $\{2\}$  are defined through

$$\{1\} = f' + \frac{1}{2}f''x^* \quad \{2\} = \frac{1}{2}f'' + \frac{1}{6}f'''x^*$$

Now, performing the transformations

$$\begin{pmatrix} \hat{x}_1 \\ \hat{x}_2 \end{pmatrix} = T \begin{pmatrix} u \\ v \end{pmatrix} \quad \begin{pmatrix} u \\ v \end{pmatrix} = T^{-1} \begin{pmatrix} \hat{x}_1 \\ \hat{x}_2 \end{pmatrix}$$

Cynthia | AXA Graduate

**AXA Global Graduate Program**

Find out more and apply

redefining / standards AXA



on (2.7.9) we arrive at

$$\begin{aligned} u_{t+1} &= -u_t + g(u_t, v_t) \\ v_{t+1} &= -\frac{P}{1-P} v_t - g(u_t, v_t) \end{aligned} \quad (2.7.10)$$

where  $g(u, v) = A[(1-P)^2u + P^2v]^2 + B[(1-P)^2 + P^2v]^3$

$$A = \frac{1}{P(2P-1)(1-P)} \{1\} \quad B = -\frac{1}{P^2(2P-1)(1-P)^2} \{2\}$$

and we observe that (2.7.10) is nothing but the original map (2.7.5) written on the desired form (2.7.1).

The next step is to restrict (2.7.10) to the center manifold. Thus, assume

$$v = i(u) = Ku^2 + Lu^3 \quad (2.7.11)$$

By use of (2.7.3) we now have

$$i(-u_t + g(u_t, v_t)) + \frac{P}{1-P} i(u_t) + g(u_t, i(u_t)) = 0$$

which is equivalent to

$$\begin{aligned} &\left[ K + \frac{PK}{1-P} + (1-P)^4A \right] u^2 + \\ &\left[ \frac{PL}{1-P} - 2KA(1-P)^4 - L + 2AP^2(1-P)^2K + B(1-P)^6 \right] u^3 = 0 \end{aligned}$$

from which we obtain

$$K = -(1-P)^5A \quad L = (1-P)^7[B + 2A^2(1-P)(1-2P)]$$

Finally, by inserting  $v = Ku^2 + Lu^3$  into the first component of (2.7.10) the restricted map may be cast in the form

$$\begin{aligned} u_{t+1} &= h(u_t) = -u_t + A(1-P)^4u_t^2 \\ &+ (1-P)^6[B - 2A^2P^2(1-P)]u_t^3 + O(u^4) \end{aligned} \quad (2.7.12)$$



Since  $u \rightarrow h(u)$  is a one-dimensional map we may now proceed by using Theorem 1.5.1 in order to show that the flip bifurcation is supercritical. A time-consuming but straightforward calculation now yields that the quantity  $b$  defined in Theorem 1.5.1 becomes

$$\begin{aligned} b &= \frac{1}{2} \left( \frac{\partial^2 h}{\partial u^2} \right)^2 + \frac{1}{3} \frac{\partial^3 h}{\partial u^3} \\ &= \left[ \frac{2\gamma}{1-P} + 1 \right]^2 \frac{2(1-P)^3}{P^2(1+P)(1-2P)} \left\{ (P-\gamma)^2 + \frac{1}{6}(1-\gamma)(4\gamma-3P+1) \right\} \quad (2.7.13) \end{aligned}$$

at bifurcation. Here we may observe that  $W(\gamma) = \{ \}$  attains its minimum when  $\gamma = (9/4)P - 3/4$  and that  $W((9/4)P - 3/4) > 0$  whenever  $0 < P < 1/2$ . Hence  $b > 0$ .

Regarding the nondegeneracy condition  $a$  defined in Theorem 1.5.1, it may be expressed as

$$a = \frac{\partial h}{\partial F} \frac{\partial^2 h}{\partial u^2} - \left( \frac{\partial h}{\partial u} - 1 \right) \frac{\partial^2 h}{\partial u \partial F} \neq 0 \text{ at } (u, v) = (0, 0)$$

Now, since the bifurcation is transformed to the origin it follows that  $\partial h / \partial u = -1$  and  $\partial h / \partial F = 0$ .

Therefore the condition  $a \neq 0$  simplifies to

$$a = 2 \frac{\partial^2 h}{\partial u \partial F} \neq 0 \Leftrightarrow 2 \frac{\partial \lambda}{\partial F} \neq 0$$

since in general  $\partial h / \partial u = \lambda$ . From the Jacobian:

$$\lambda = \frac{1}{2} \left( w - \sqrt{w^2 + 4Pw} \right)$$

where

$$w = f'x^* + f = \frac{1}{1+P} \left\{ -\frac{1}{\gamma} [(F + FP)^\gamma - 1] + 1 \right\}$$

it follows that

$$\begin{aligned} 2 \frac{\partial \lambda}{\partial F} &= \frac{dw}{dF} - \frac{1}{2\sqrt{w^2 + 4Pw}} \left( 2w \frac{dw}{dF} + 4P \frac{dw}{dF} \right) \\ &= \frac{dw}{dF} \left[ 1 - \frac{1}{\sqrt{w^2 + 4Pw}} (w + 2P) \right] \end{aligned}$$

At bifurcation,  $w = -(1 - P)^{-1}$  which inserted into the expression above gives

$$2 \frac{\partial \lambda}{\partial F} = -2 \left[ \frac{2\gamma}{1-P} + 1 \right]^{1-(1/\gamma)} \frac{(1-P)^2}{1-2P} \quad (2.7.14)$$

and clearly, (2.7.14) is nonzero whenever  $0 < P < 1/2$ . Consequently, the flip bifurcation is supercritical, which means that when the fixed point fails to be stable, a stable two-periodic orbit is established.

—

We close this section by showing the dynamics beyond the flip bifurcation threshold for the Ricker map

$$(x_0, x_1) \rightarrow (F e^{-x}(x_0 + x_1), P x_0) \quad (2.7.15)$$

which is a special case of map (2.7.5) (the case  $\gamma \rightarrow 0$ ). Assuming  $F(1 + P) > 1$  the nontrivial fixed point of (2.7.15) is

$$(x_0^*, x_1^*) = \left( \frac{1}{1+P} \ln(F(1+P)), \frac{P}{1+P} \ln(F(1+P)) \right)$$

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



**Month 16**


I was a construction supervisor in the North Sea advising and helping foremen solve problems

Real work  
International opportunities  
Three work placements










**Figure 14:** The bifurcation diagram of map (2.7.14) in the case  $P = 0.2$ . For small  $F$  values we see the stable fixed point of (2.7.14) which undergoes a supercritical flip bifurcation when  $F = 10.152$ . Through further increase of  $F$  stable orbits of period  $2^k$  are created until an accumulation value  $F_a$  for the flip bifurcations is reached. Beyond  $F_a$  the dynamics is chaotic.

and whenever  $0 < P < 1/2$  we have according to the preceding example that the fixed point undergoes a supercritical flip bifurcation at the threshold  $F = (1/(1 + P)) \exp(2/(1 - P))$ .

Now, consider the value  $P = 0.2$ . Under this choice the fixed point is stable in the  $F$  interval  $0.834 < F < 10.152$  and in Figure 14 we have plotted the bifurcation diagram of (2.7.15) in the range  $5 < F < 80$ . We clearly identify the supercritical flip at the threshold  $F = 10.152$  and beyond that stable periodic orbits of period  $2^k$  are established through further increase of  $F$  so what we recognize is essentially the same kind of dynamical behaviour as we found when we considered one-dimensional maps. Beyond the point of accumulation for the flip bifurcation sequence the dynamics becomes chaotic as displayed in Figure 15. Note that the chaotic attractor consists of 4 disjoint subsets (branches) that are visited once every fourth iteration so a certain kind of four periodicity is preserved in the chaotic regime. In case of higher  $F$  values the branches merge together.



**Figure 15:** The chaotic attractor consisting of four separate branches just beyond the point of accumulation for the ip bifurcations in the case  $P = 0.2, F = 58.5$ . The dynamics goes in the direction  $A \rightarrow B \rightarrow C \rightarrow D$

## 2.8 Beyond the Hopf bifurcation, possible routes to chaos

As we proved in section 2.5, the outcome of a supercritical Hopf bifurcation is that when the fixed point of a discrete map fails to be stable, an attracting invariant curve which surrounds the fixed point is created. Our goal in this section is to describe the dynamics on such an invariant curve. We will also discuss possible routes to chaos and as it will become clear, the dynamics may be much richer than in the one-dimensional cases discussed in Part I.

In general terms, the dynamics on an invariant curve (circle) created by a Hopf bifurcation may be analysed by use of equation (2.5.14b). Indeed, if we substitute the fixed point  $r^*$  of (2.5.14a) into (2.5.14b) we arrive at

$$\varphi \rightarrow \varphi + c - \frac{bd}{a} \mu = \varphi + \sigma(\mu) \tag{2.8.1}$$

where  $c = \arg \lambda$ . Also recall that when we derived (2.5.14a,b) we first transformed the bifurcation to the origin. If the Hopf bifurcation occurs at a threshold  $\mu_0 \neq 0$ ,  $\sigma(\mu) = c + (bd/a)(\mu_0 - \mu)$ .

Now, the essential feature is that successive iterations of (2.8.1) simply “move” or rotate points from one location to another on the invariant curve. Hence, the original map  $f_\mu : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  may be regarded as being topological equivalent to a circle map  $g : S^1 \rightarrow S^1$  once the invariant curve is established. Moreover, considering  $g$ , one may define its rotation number as the average amount that points are rotated by an iteration of the map. Therefore, we may (to leading order, recall that (2.8.1) is a truncated map) regard (2.8.1) as a circle map with rotation number  $\sigma(\mu)$ .

**Remark 2.8.1.** A more precise definition of the rotation number may be achieved along the following line: Given a circle map  $g : S \rightarrow S$  we first “lift”  $g$  to the real line  $\mathbb{R}$  by use of  $\pi : \mathbb{R} \rightarrow S$ ,  $\pi(x) = \cos(2\pi x) + i \sin(2\pi x)$  and then define the lift  $F$  as  $F : \mathbb{R} \rightarrow \mathbb{R}$ ,  $\pi \circ F = g \circ \pi$ . Next, let  $\sigma_0(F) = \lim_{n \rightarrow \infty} F^n(x)/x$  and finally define the rotation number of  $g$ ,  $\sigma(g)$ , as the unique number in  $[0, 1)$  such that  $\sigma_0(F) - \sigma(g)$  is an integer. In Devaney’s book there is an excellent introduction to circle maps.  $\square$

**ie** business school

**93%**  
OF MIM STUDENTS ARE  
WORKING IN THEIR SECTOR 3 MONTHS  
FOLLOWING GRADUATION

**MASTER IN MANAGEMENT**

- STUDY IN THE CENTER OF MADRID AND TAKE ADVANTAGE OF THE UNIQUE OPPORTUNITIES THAT THE CAPITAL OF SPAIN OFFERS
- PROPEL YOUR EDUCATION BY EARNING A DOUBLE DEGREE THAT BEST SUITS YOUR PROFESSIONAL GOALS
- STUDY A SEMESTER ABROAD AND BECOME A GLOBAL CITIZEN WITH THE BEYOND BORDERS EXPERIENCE

Length: 10 MONTHS  
Av. Experience: 1 YEAR  
Language: ENGLISH / SPANISH  
Format: FULL-TIME  
Intakes: SEPT / FEB

**5 SPECIALIZATIONS**  
PERSONALIZE YOUR PROGRAM

**#10 WORLDWIDE**  
MASTER IN MANAGEMENT  
FINANCIAL TIMES

**55 NATIONALITIES**  
IN CLASS

[www.ie.edu/master-management](http://www.ie.edu/master-management) | [mim.admissions@ie.edu](mailto:mim.admissions@ie.edu) | [f](#) [t](#) [in](#) Follow us on IE MIM Experience



Returning to map (2.8.1) the rotation number may be irrational or rational. In the former case this means that as the number of iterations of the map tends to infinity, the invariant curve will be filled with points. Whenever  $\sigma$  irrational, an orbit of a point is often referred to as a quasistationary orbit. If  $\sigma = 1/n$ , rational, the dynamic outcome is an  $n$ -period orbit. It is of great importance to realize that whenever the rotation number is rational for a given parameter value  $\mu = \mu_r$ , it follows from the implicit function theorem that there exists an open interval about  $\mu_r$  where the periodicity is maintained. This phenomenon is known as frequency locking of periodic orbits. Consequently, periodic dynamics will occur in parameter regions, not at isolated parameter values only. As we shall see, such regions (or intervals) may in fact be large. So in order to summarize: beyond the Hopf bifurcation (and outside the strongly resonant cases where  $\lambda$  is third or fourth root of unity) there are quasistationary orbits restricted to an invariant curve and there may also be orbits of finite period established through frequency locking as the value of the parameter  $\mu$  in the model is increased.

Our next goal is by way of examples to study in more detail the interplay between these cases as well as studying possible routes to chaos. We start by scrutinizing a population model first presented in Wikan and Mjølhus (1995).

—

**Example 2.8.1.** First, consider the two-age class population model

$$(x_0, x_1) \rightarrow (Fx_1, P e^{-\alpha x} x_0) \quad (2.8.2)$$

which is a semelparous species model where the fecundity  $F$  is constant while the survival probability  $p(x) = P \exp(-\alpha x)$  is density dependent.  $\alpha$  is a positive number (scaling constant) and we assume that  $PF > 1$ .

It is easy to verify that (2.8.2) possesses the following properties: The fixed point may be expressed as


$$(x_0^*, x_1^*) = \left( \frac{F}{1+F} x^*, \frac{1}{1+F} x^* \right) \quad (2.8.3)$$

where  $x^* = x_0^* + x_1^* = \alpha^{-1} \ln(PF)$ . Moreover, the eigenvalue equation may be cast in the form

$$\lambda^2 + \frac{\ln(PF)}{1+F} \lambda + \frac{F \ln(PF)}{1+F} - 1 = 0 \quad (2.8.4)$$

and from the Jury criteria one obtains that the fixed point is stable in case of  $PF$  small but undergoes a Hopf bifurcation at the threshold

$$e) \quad P = P_c = \frac{1}{F} e^{2(1+F)/F} \quad (2.8.5)$$



**Figure 16:** The dynamics of map (2.8.2) (a quasistationary orbit), just beyond the Hopfbifurcation threshold.

Note that  $\alpha$  drops out of (2.8.4), (2.8.5) which simply means that stability properties are independent of  $\alpha$ . At bifurcation threshold (2.8.5) the solution of the eigenvalue equation becomes

$$\lambda = -\frac{1}{F} \pm \sqrt{1 - \frac{1}{F^2}} i \quad (2.8.6)$$

A final observation is that by rewriting (2.8.2) on standard form (as in Example 2.5.1) and then apply Theorem 2.5.2, it is possible to prove that the bifurcation is supercritical.

Now, let us scrutinize a numerical example somewhat closer. Assume  $P = 0.6$ . Then from (2.8.5) the  $F$  value at bifurcation threshold is numerically found to be  $F = F_c = 14.1805$ . We want to investigate the dynamics when  $F > F_c$ . In Figure 16 we show the dynamics just beyond the instability threshold in the case  $(\alpha, P, F) = (0.02, 0.6, 15)$ . From an initial state  $(x_{00}, x_{10})$  500 iterations have been computed and the last 20 together with the (unstable!) fixed point are plotted. The invariant curve is indicated by the dashed line so clearly the original map (2.8.2) does nothing but rotate points around that curve, i.e. (2.8.2) acts as a circle map.






Figure 17: A 4-periodic orbit generated by map (2.8.2).

Moreover, Figure 16 demonstrates a clear tendency towards 4-periodic dynamics. This is as expected due to the location of the eigenvalues. Indeed, when  $F_c = 14.1805$  it follows from (2.7.6) that the eigenvalues are located very close to the imaginary axis ( $\lambda_{1,2} = -0.0750 \pm \sqrt{0.9975} i$ ), and since the rotation number (up to leading order!) has the form  $\sigma(F) = c + (bd/a)(F_c - F)$  where  $c = \arg \lambda$  it follows that  $\sigma$  must be close to  $1/4$  in case of  $F$  close to  $F_c$ .

“I studied English for 16 years but...  
...I finally learned to speak it in just six lessons”  
Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download

[Click on the ad to read more](#)

If we increase  $F$  beyond 15 we observe (due to frequency locking!) that an exact 4-periodic orbit is established. This is shown in Figure 17 in the case  $(\alpha, P, F) = (0.02, 0.6, 20)$  and further, it is possible to verify numerically that the exact 4-periodicity is maintained as long as  $F$  does not exceed the value 21.190.

At  $F = 21.190$  the fourth iterate of (2.8.2) undergoes a flip bifurcation, thus an 8-periodic orbit is established, and through further enlargement of  $F$  we find that new flip bifurcations take place at the parameter values 24.232 and 24.883 which again result in orbits of period 16 and 32 respectively. Hence we observe nothing but the flip bifurcation sequence which we discussed in Part I. The point of accumulation for the flip bifurcation is found to be  $F_a \approx 25.07$  and in case of  $F > F_a$  the dynamics becomes chaotic.

These findings are shown in Figures 18, 19 and 20. In Figures 18 and 19 periodic orbits of period 8 and 32 are displayed. In Figure 20 we show the chaotic attractor. Note that the attractor is divided in 4 disjoint subsets and that each of the subsets are visited once every fourth iteration so there is a kind of 4-periodicity preserved, even in the chaotic regime.  $\square$




Figure 18: An 8-periodic orbit generated by map (2.8.2).




Figure 19: A 32-periodic orbit generated by map (2.8.2).




Figure 20: Map (2.8.2) in the chaotic regime.

**Example 2.8.2.** The next example (which rests upon the findings in Wikan (1997)) is basically the same as the previous one but the dimension of the map has been extended by 1 and we consider a general survival probability  $p(x)$ ,  $0 < p(x) \leq 1$ ,  $p'(x) \leq 0$ , instead of  $p(x) = P \exp(-x)$ . Hence we consider the problem

$$(x_1, x_2, x_3) \rightarrow (F_3 x_3, p(x) x_1, p(x) x_2) \quad (2.8.7)$$

Skipping computational details (which are much more cumbersome here than in our previous example) we find that the nontrivial fixed point is

$$(x_1^*, x_2^*, x_3^*) = \left( \frac{x^*}{K}, p(x^*) \frac{x^*}{K}, p^2(x^*) \frac{x^*}{K} \right) \quad (2.8.8)$$

where  $K = \sum_{i=1}^3 p^{i-1}(x^*)$  and  $x^* = p^{-1}(F_3^{-1/(n-1)})$ . ( $p^{-1}$  denotes the inverse of  $p$ .)

Moreover, by first computing the Jacobian and then use the Jury criteria, it is possible to show that (2.8.8) is stable as long as

$$-p'(x^*) \frac{x^*}{K} < p(x^*) \frac{1 + p(x^*) - 2p^2(x^*)}{(1 + p(x^*))(1 - p^2(x^*))} \quad (2.8.9)$$

(2.8.8) becomes unstable when  $F_3$  is increased to a level  $F_{H1}$  where (2.8.9) becomes an equality.

At that level a (supercritical) Hopf bifurcation occurs and the complex modulus 1 eigenvalues may be expressed as

$$\lambda_{1,2} = -\frac{p^2(x^*)}{1 + p(x^*)} \pm \sqrt{1 - \frac{p^4(x^*)}{(1 + p(x^*))^2}} i \quad (2.8.10)$$

Excellent Economics and Business programmes at:



university of  
 groningen




“The perfect start  
 of a successful,  
 international career.”

**CLICK HERE**  
 to discover why both socially  
 and academically the University  
 of Groningen is one of the best  
 places for a student to be

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)



Now, for comparison reasons, assume that  $p(x) = P \exp(-x)$  just as in Example 2.8.1. Then it easily follows that  $F_3$  is a “large” number at bifurcation threshold  $F_{H1}$  and further that  $p(x^*) \ll 1$ . Consequently,  $\lambda_{1,2}$  are located very close to the imaginary axis, in fact even closer than the eigenvalues from Example 2.8.1. When we increase  $F_3$  beyond  $F_{H1}$  we observe the following dynamics: In case of  $F_3 - F_{H1}$  small we find an almost 4-periodic orbit restricted on an invariant curve and through further enlargement of  $F_3$  we once again find (through frequency locking) that an exact 4-periodic orbit is the outcome. Thus the dynamics is qualitatively similar to what we found in Example 2.8.1. However, if we continue to increase  $F_3$  we do not experience the flip bifurcation sequence. Instead we find that the fourth iterate of map (2.8.7) undergoes a (supercritical) Hopf bifurcation at a threshold  $F_3 = F_{H2}$ . Therefore, beyond that threshold, and in case of  $F_3 - F_{H2}$  small, the dynamics is restricted on 4 disjoint invariant attracting curves which are visited once every fourth iteration. This is displayed in Figure 21. At an even higher value,  $F_3 = F_s$ , map (2.8.7) undergoes a subcritical bifurcation which implies that whenever  $F_3 > F_s$  there is no attractor at all so in this part of parameter space we simply find that points  $(x_1, x_2, x_3)$  are randomly distributed in state space.  $\square$

So far we have demonstrated that although the dynamics is a quasistationary orbit just beyond the original Hopf bifurcation threshold, the dynamical outcome may be a periodic orbit as we penetrate deeper into the unstable parameter region. Such a phenomenon may happen when  $|\arg \lambda|$  is close to  $\pi/4$  at bifurcation threshold (4-periodicity). Another possibility (among others!) is that  $|\arg \lambda|$  is close to  $2\pi/3$  (3-periodicity).




Figure 21: Map (2.8.7) after the secondary Hopf bifurcation.

Note, however, that if  $\arg \lambda$  is close to a “critical” value, say  $\pi/2$ , at bifurcation it does not necessarily imply that a periodic orbit is created when we continue to increase the bifurcation parameter. In fact, when the parameter is enlarged the location of the eigenvalues may move away from the imaginary axis, hence the periodicity will be less pronounced as the bifurcation parameter grows. In our next example there is no periodicity at all.


**Example 2.8.3.** Consider the two-dimensional population map

$$(x_1, x_2) \rightarrow (F e^{-x_2} x_1 + F e^{-x_2} x_2, P x_1) \tag{2.8.11}$$

Hence, only the second age class  $x_2$  contributes to density effects. As before,  $F > 0$ ,  $0 < P \leq 1$  and  $F(1 + P) > 1$ .

We urge the reader to verify that the fixed point  $(x_1^*, x_2^*)$  may be written as

$$(x_1^*, x_2^*) = \left( \frac{1}{P} x_2^*, \ln[(1 + P)F] \right) \tag{2.8.12}$$



**Figure 22:** Dynamics generated by map (2.8.11). Parameter values: (a)  $(P, F) = (0.6, 2.5)$ ; (b)  $(P, F) = (0.6, 5.0)$ .

and further that a (supercritical) Hopf bifurcation occurs at the threshold

$$F = F_H = \frac{1}{1 + P} e^{(1+2P)/(1+P)} \tag{2.8.13}$$



and finally that the solution of the eigenvalue equation at threshold (2.7.13) becomes

$$\lambda = \frac{1}{2(1+P)} \left\{ 1 \pm \sqrt{4(1+P)^2 - 1} i \right\} \quad (2.8.14)$$

Now, assume that  $P$  is not close to zero. Then, the location of  $\lambda$  clearly suggests that frequency locking into an orbit of finite period will not take place. In Figure 22a we show the invariant curve just beyond the bifurcation threshold  $(P, F) = (0.6, 2.5)$  and on that curve we find no tendency towards periodic dynamics.

As we continue to increase  $F$  ( $P$  fixed) the “radius” of the invariant curve becomes larger. Eventually, the invariant curve becomes kinked and signals that the attractor is not topological equivalent to a circle anymore and finally the curve breaks up and a chaotic attractor is born. This is exemplified in Figure 22b. □

In our final example (cf. Wikan and Mjølhus (1996) or Wikan (2012b)) all bifurcations that we have previously discussed are present.

## American online LIGS University

is currently enrolling in the  
Interactive Online **BBA, MBA, MSc,**  
**DBA and PhD** programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive **Online** education
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to  
find out more!

Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](#).







**Example 2.8.4.** Referring to section 2.4, Examples 2.4.1 and 2.4.3 we showed that the fixed point  $(x_0^*, x_1^*)$  of map (2.4.2), i.e.


$$(x_0, x_1) \rightarrow (F_0 e^{-\alpha x} x_0 + F_1 e^{-\alpha x} x_1, P_0 x_0)$$

is stable in case of small equilibrium populations  $x^* = x_0^* + x_1^*$  but eventually will undergo a supercritical Hopf bifurcation at the threshold

$$F = F_H = \frac{1}{1 + P_0} e^{(1+2P_0)/P_0}$$

provided  $1/2 < P_0 < 1$  and equal fecundities  $F_0 = F_1 = F$ . In Figure 23 we have generated the bifurcation of the map in the case  $P_0 = 0.9$ ,  $\alpha = 0.01$ . The bifurcation parameter  $F$  is along the horizontal axis, the total population  $x$  along the vertical. Omitting computational details (which may be obtained in Wikan and Mjølhus (1996)) we shall now use Figure 23 in order to reveal the dynamics of (2.4.2).

In case of  $5.263 < F < 10.036$  there is one attractor, namely the stable fixed point  $(x_0^*, x_1^*)$ . (The lower limit 5.263 is a result of the requirement  $F(1 + P) > 1$ .) At the threshold  $F_s = 10.036$  a 3-cyclic attractor with large amplitude is created. Thus beyond  $F_s$  there exists a parameter ( $F$ ) interval where there are two coexisting attractors and the ultimate fate of an orbit depends on the initial condition. It is a well known fact that multiple attractors indeed may occur in nonlinear systems. What happens in our case is that the third iterate of the original map (2.4.2) undergoes a saddle-node bifurcation at  $F_s$ .



**Figure 23:** The bifurcation diagram generated by map (2.4.2).

This may be verified numerically by computing the Jacobian of the third iterate and show that the dominant eigenvalue of the Jacobian equals unity. Moreover (referring to section 1.5, see also Exercise 1.4.2 in section 1.4), a 3-cycle consisting of unstable points is also created through the saddle node at threshold  $F_s$ . This repelling 3-cycle is of course invisible to the computer.

In the interval  $10.036 < F < 11.81$  the large amplitude 3-cycle and the fixed point are coexisting attractors. At  $F_H = 11.81$  the fixed point undergoes a supercritical Hopf bifurcation (for a proof, cf. Wikan and Mjølhus (1996)), thus in case of  $F > F_H$ ,  $F - F_H$  small, there is coexistence between the 3-cyclic attractor and a quasistationary orbit restricted to an invariant curve. The coexistence takes place in the interval  $11.81 < F < 12.20$ . In somewhat more detail we also find that since  $\arg \lambda$  (where  $\lambda$  is the eigenvalue of the Jacobian of (2.4.2)) is close to  $2\pi/3$  at  $F_H$  there is a clear tendency towards 3-periodic dynamics on the invariant curve but there is no frequency locking into an exact 3-periodic orbit.

At  $F_K = 12.20$  the invariant curve disappears. Consequently, in case of  $F > F_K$ , there is again only one attractor, namely the attracting 3-cycle. The reason that the invariant curve disappears at threshold  $F_K$  is that it is “hit” by the three branches of the repelling 3-cycle. This phenomenon is somewhat akin to what is called a crisis in the chaos literature.

As we continue to increase  $F$  successive flip bifurcations occur, creating orbits of period  $3 \cdot 2^k$ ,  $k = 1, 2, \dots$ , in much of the same way as we have seen in earlier examples. Eventually an accumulation value  $F_a$  for the flip bifurcations is reached, and beyond that value the dynamics becomes chaotic. At first the chaotic attractor consists of three separate branches which are visited once every third iteration. When  $F$  is even more increased the branches merge together.  $\square$

Through our previous examples, which all share the common feature that the original (first) bifurcation is a Hopf bifurcation, we have experienced that the nonstationary dynamics beyond the instability threshold may indeed be different from map to map. In the following exercises even more possible dynamical outcomes are demonstrated.

**Exercise 2.8.1.** Consider the map (cf. Wikan (1998))

$$(x_0, x_1) \rightarrow (F_1 x_1, P_0(1 - \gamma \beta x)^{1/\gamma} x_0)$$

where  $\beta > 0$ ,  $\gamma \leq 0$ .

- a) Compute the nontrivial fixed point  $(x_0^*, x_1^*)$ .
- b) Assume that  $\gamma > \gamma_c = -(F_1/2(1 + F_1))$  and show that the fixed point undergoes a Hopf bifurcation at the threshold

$$P_0 = \frac{1}{F_1} \left[ 1 + \gamma \frac{2(1 + F_1)}{F_1} \right]^{1/\gamma}$$

- c) Assume that  $\gamma > \gamma_c$  but  $\gamma - \gamma_c$  small. Investigate numerically the dynamical outcomes when  $P_0$  is fixed and  $F_1$  is increased beyond the bifurcation threshold.
- d) (difficult!) Show that the Hopf bifurcation is supercritical. □

**Exercise 2.8.2.** Consider the semelparous population model

$$\begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix}_{t+1} = \begin{pmatrix} 0 & 0 & F_2 e^{-x} \\ P_0 & 0 & 0 \\ 0 & P_1 & 0 \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix}_t$$

**DON'T EAT YELLOW SNOW**

What will your advice be?

Some advice just states the obvious. But to give the kind of advice that's going to make a real difference to your clients you've got to listen critically, dig beneath the surface, challenge assumptions and be credible and confident enough to make suggestions right from day one. At Grant Thornton you've got to be ready to kick start a career right at the heart of business.

Sound like you? Here's our advice: visit [GrantThornton.ca/careers/students](https://www.grantthornton.ca/careers/students)

Scan here to learn more about a career with Grant Thornton.

**Grant Thornton**  
An instinct for growth™

© Grant Thornton LLP. A Canadian Member of Grant Thornton International Ltd



a) Show that the fixed point is

$$(x_0^*, x_1^*, x_2^*) = \left( \frac{1}{1 + P_0 + P_0 P_1} x^*, \frac{P_0}{1 + P_0 + P_0 P_1} x^*, \frac{P_0 P_1}{1 + P_0 + P_0 P_1} x^* \right)$$

where  $x^* = \ln(P_0 P_1 F_2)$ .

b) Compute the Jacobian and show that the eigenvalue equation may be cast in the form

$$\lambda^3 + \varepsilon \lambda^2 + P_0 \varepsilon \lambda + P_0 P_1 \varepsilon - 1 = 0$$

where  $\varepsilon = x^*/(1 + P_0 + P_0 P_1)$ .

c) Use the Jury criteria (2.1.16) and show that the fixed point is stable whenever

$$\varepsilon_4 < \varepsilon < \varepsilon_2$$

where

$$\varepsilon_4 = \frac{1 + P_0 - 2P_0 P_1}{P_0 P_1 (1 - P_0 P_1)} \quad \text{and} \quad \varepsilon_2 = \frac{2}{1 - P_0 + P_0 P_1}$$

d) Use the result in c) and show that the fixed point is stable provided

$$\frac{1}{2} < P_0 < 1 \quad P_1 > \frac{1 + P_0}{3P_0}$$

e) The results from c) and d) are special in the sense that they imply that the fixed point is unstable in case of  $x^*$  (or  $F_2$ ) small, becomes stable for larger values of  $x^*$  (or  $F_2$ ) and then becomes unstable again through further enlargement of  $x^*$  (or  $F_2$ ). Note that  $\varepsilon_4$  and  $\varepsilon_2$  are Hopf and flip bifurcation thresholds respectively. Investigate (numerically the dynamics in case of  $\varepsilon < \varepsilon_4$  (i.e.  $x^*$  small) and  $\varepsilon < \varepsilon_2$  (i.e.  $x^*$  large). (Hint: cf. Exercise 2.4.3.) Other properties of this model as well as properties of more general semelparous population models may be obtained in Mjølhus et al. (2005).  $\square$

**Exercise 2.8.3 (Coexistence of age classes).** Consider the two age class map (Wikan 2012a)

$$(x_1, x_2) \rightarrow (F e^{-\alpha x} x_2, P e^{-\beta x} x_1)$$

cf. (2.3.1) where  $x = x_1 + x_2$ ,  $0 < P \leq 1$ ,  $F > 0$  and  $\alpha, \beta > 0$ .

a) Show that the nontrivial fixed point of the map is

$$(x_1^*, x_2^*) = \left( \frac{1}{1 + aP} x^*, \frac{aP}{1 + aP} x^* \right)$$

where  $a = R_0^{-(\beta/(\alpha+\beta))}$ ,  $x^* = (\alpha + \beta)^{-1} \ln R_0$  and  $R_0 = PF > 1$ .

b) Use the Jury criteria and show that if  $\beta > \alpha$  then there exists a parameter region where  $(x_1^*, x_2^*)$  is stable and, moreover, that when  $R_0$  increases there will occur a Hopf bifurcation at the threshold

$$R_0 = \exp \left[ \frac{2(\alpha + \beta)(1 + aP)}{\beta + \alpha aP} \right]$$

c) Investigate numerically the behaviour of the map beyond instability threshold. (Hint: the cases  $\beta - \alpha$  small,  $\beta - \alpha$  large should be treated separately.)

d) The parameters  $\alpha$  and  $\beta$  may be interpreted as “strength” of density dependence. Show that if the strength of density dependence in the fecundity  $\alpha$  is equal or larger than the strength of density dependence in the survival  $\beta$  then  $(x_1^*, x_2^*)$  will always be unstable.

e) What kind of dynamic outcome do you find in the case  $\beta < \alpha$ ? □

**Exercise 2.8.4 (Permanence in stage-structured models).** In Example 2.5.1 we analysed a stage-structured cod model. A slightly more general form of such a model is

$$(i) \quad \mathbf{x}_{t+1} = \mathbf{A}_x \mathbf{x}_t$$

where  $\mathbf{x} = (x_1, x_2)^T$  and

$$\mathbf{A}_x = \begin{pmatrix} (1 - \mu_1)S(x) & f(x) \\ p(x) & (1 - \mu_2) \end{pmatrix}$$

Here,  $x_{1,t}$  and  $x_{2,t}$  are the immature and mature part of the population respectively and just as in the age-structured case  $f(x)$  is the fecundity.  $p(x)$  is the fraction of the immature population that survives to become mature, and  $\mu_1$  and  $\mu_2$  are (natural) death rates. Finally, it is also assumed that the remaining part of the immature population  $(1 - \mu_1)x_1$  is reduced by a nonlinear factor  $s(x)$ .



Further, let  $s(x) = S\hat{s}(x)$ ,  $f(x) = F\hat{f}(x)$ ,  $p(x) = P\hat{p}(x)$  where  $0 \leq S \leq 1$ ,  $0 < P \leq 1$ ,  $F > 0$ ,  $0 \leq \mu_1, \mu_2 < 1$ ,  $0 < \hat{s}(x), \hat{p}(x), \hat{f}(x) \leq 1$ ,  $\hat{s}(0) = \hat{p}(0) = \hat{f}(0) = 1$ . A final but important restriction in such models is  $(1 - \mu_1)S + P \leq 1$ . Otherwise, the fraction of juveniles that survives to become adults plus the fraction that survives but remain juveniles may be larger than 1 even in case of zero fecundity which of course is unacceptable from a biological point of view.

Definition. Let  $x_t = x_{1,t} + x_{2,t}$  be the total population at time  $t$ . Model (i) is said to be permanent if there exists  $\delta > 0$  and  $D > 0$  such that

$$\delta < \liminf_{t \rightarrow \infty} x_t \leq \limsup_{t \rightarrow \infty} x_t < D$$

□

Thus, if a population model is permanent, the total population density neither explodes nor goes to zero (see Kon et al. (2004)). Define the net reproductive number  $R_0$  as

$$R_0 = \frac{PF}{\mu_2[1 - (1 - \mu_1)S]}$$

.....Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



Click on the ad to read more

Our goal is to prove the following theorem:

**Theorem:** Suppose that model (i) is continuous and that one of  $\hat{p}(x)x_1$  or  $\hat{f}(x)x_2$  is bounded from above. Further assume that the matrix  $\mathbf{A}_0$  is irreducible and  $\mathbb{R}_+^2 \setminus \{0\}$  forward invariant (i.e. that  $\mathbf{A}_x \mathbf{x} \in \mathbb{R}_+^2 \setminus \{0\}$  for all  $\mathbf{x} \in \mathbb{R}_+^2 \setminus \{0\}$ ). Then model (i) is permanent provided  $R_0 > 1$ .  $\square$

a) Clearly,  $(\tilde{x}_1, \tilde{x}_2) = (0, 0)$  is a fixed point of (i). Use the Jury criteria and show that  $(0, 0)$  is unstable provided  $R_0 > 1$ .

b) Explain why  $\mathbf{A}_0$  is irreducible and  $\mathbb{R}_+^2 \setminus \{0\}$  forward invariant.

—

It remains to prove that the population density does not explode, i.e. that (i) is a dissipative model. From Kon et al. (2004), see also Cushing (1998), we apply the following definition of dissipativeness:

**Definition:** Model (i) is said to be dissipative if there exists a compact set  $X \subset \mathbb{R}_+^2$  such that for all  $\mathbf{x}_t \in \mathbb{R}_+^2$  there exists a  $t^M = t^M(\mathbf{x}_0)$  satisfying  $\mathbf{x}_t \in X$  for all  $t \geq t^M$ .  $\square$

c) Assume  $\hat{p}(x)x_1 \leq K_0$  where  $K_0$  is a constant. Use (i) and induction to establish the relations

$$x_{2,t+1} \leq PK_0 + (1 - \mu_2)x_{2,t}$$

and

$$x_{2,t} \leq (1 - \mu_2)^t x_{2,0} + \frac{PK_0}{\mu_2}$$

d) Use c) to conclude that there exists  $t^A = t^A(x_{2,0})$  such that for  $t > t^A$

$$x_{2,t} \leq \frac{2PK_0}{\mu_2} = K_1$$

e) Use the previous result together with (i) and induction to show that

$$x_{1,t+1} \leq (1 - \mu_1)Sx_{1,t} + FK_1$$

$$x_{1,t} \leq (1 - \mu_1)^t S^t x_{1,0} + \frac{FK_1}{1 - (1 - \mu_1)S}$$

$\square$



f) Show that there exists  $t^B = t^B(x_{1,0})$  such that for  $t > t^B(x_{1,0})$

$$x_{1,t} \leq \frac{2FK_1}{1 - (1 - \mu_1)S} = K_2$$

g) Take  $t^M = \max\{t^A, t^B\}$  and  $K = \max\{K_1, K_2\}$  and conclude that  $x_{1,t} \leq K$  and  $x_{2,t} \leq K$ , hence (i) is dissipative if  $\hat{p}(x)x_1$  is bounded from above.

h) Assume  $\hat{f}(x)x_2 \leq K_0$  and show in a similar manner that (i) is dissipative in this case too.

□

**Remark 2.8.2.** In Leslie matrix models nonoverlapping age classes are assumed. This is not the case in the stage-structured model from the previous exercise (or the model presented in Example 2.5.1). Moreover, while Leslie matrix models are maps from  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  (or  $\mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ ) where  $n$  may be a large integer, stage-structured models are mainly maps from  $\mathbb{R}^2 \rightarrow \mathbb{R}^2$  where we do not have the possibility to study the dynamic behaviour of age classes in detail. Some stage-structured models are maps from  $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ . Typically, they are insect models where the population is divided into three stages: larvae (L), pupae (P), and adult insects (A). In fact, such models are fully capable of describing and even predicting nonstationary and chaotic behaviour in laboratory insect populations, see Cushing et al. (1996), Costantino et al. (1997), Dennis et al. (1997), and Cushing et al. (1998). □

**Exercise 2.8.5 (Prey-Predator systems).** In 1920 Lotka introduced a system of differential equations which described the interaction between a prey species  $x$  and a predator species  $y$ . These equations were rediscovered by Volterra in 1926 and today they are often referred to as the Lotka-Volterra equations. A discrete version of the equations (written as a map) is

$$(i) \quad (x, y) \rightarrow [(1 + r) - ay)x, (-c + bx)y]$$

The first component of the map expresses that the growth rate of the prey is a constant  $(1 + r)$  due to the species itself minus a term proportional to the number of predators. In the same way, the growth rate of the predator is proportional to the number of prey minus a term  $c$  which is due to the predator species itself. All constants are assumed to be positive.

a) Find the nontrivial fixed point of the map and show that it is always unstable.

b) Consider the prey-predator map

$$(ii) \quad (x, y) \rightarrow (f(y)x, g(x)y)$$

where  $\partial f/\partial y < 0$  and  $\partial g/\partial x > 0$ . Show that  $|\lambda| > 1$  where  $\lambda$  is the solution of the eigenvalue equation. (See Maynard Smith (1979) for computational details.) What is the qualitative dynamic behaviour of maps like (i) and (ii)?

c) Next, consider the two parameter family prey-predator maps

$$(iii) \quad (x, y) \rightarrow [((1+r) - rx - ay)x, axy]$$

where  $r > 0$ ,  $a > 0$  (Maynard Smith, 1968). Show that (iii) has three fixed points,

$$(\hat{x}, \hat{y}) = (0, 0), (\tilde{x}, \tilde{y}) = (1, 0) \text{ and } (x^*, y^*) = (1/a, r(a-1)/a^2).$$

d) Following Neubert and Kot (1992) who perform a detailed analysis of (iii) show that 1)  $(\hat{x}, \hat{y})$  is always unstable, 2)  $(\tilde{x}, \tilde{y})$  is stable whenever  $0 < r < 1$  and  $0 < a < 1$ , and 3)  $(x^*, y^*)$  is stable provided  $1 < a < 2$  and  $0 < r < 4a/(3-a)$ .

e) Still referring to Neubert and Kot (1992), show that (iii) undergoes a transcritical bifurcation when  $a = 1$  and draw a bifurcation diagram similar to Figure 4b in Section 1.5.



Join the best at  
the Maastricht University  
School of Business and  
Economics!

Top master's programmes

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht University is the best specialist university in the Netherlands (Elsevier)

Visit us and find out why we are the best!  
Master's Open Day: 22 February 2014

[www.mastersopenday.nl](http://www.mastersopenday.nl)



Click on the ad to read more

((iii) has several other interesting properties. It should be easy for the reader to verify that in case of  $1 < a < 2$ ,  $r = 4a/(3 - a)$  gives birth to a flip bifurcation, but unlike most of the cases treated so far (however, see Exercise 1.5.2), this bifurcation is of the subcritical type and the predator goes extinct at instability threshold. (Formally, this may be proved by using the same procedure as in Example 2.7.1.) Moreover, when  $a = 2$  and  $r \neq 4, r \neq 6$  a Hopf bifurcation occurs and whenever  $a > 2$ ,  $|a - 2|$  small the dynamics is restricted on an invariant curve. In the strong resonant cases  $r = 4, r = 6$  we find the same qualitative picture as we did in Exercises 2.5.1 and 2.5.2. For further reading of this fascinating map we refer to the original paper by Neubert and Kot (1992).)

f) Finally, consider the age-structured prey-predator map

$$(x_1, x_2, y_1, y_2) \rightarrow \left( F_2 x_2, P e^{-(x+\beta_1 y)} x_1, G_2 x_2, \frac{Q}{1+y} \frac{\beta_2 x}{1+\beta_2 x} y_1 \right)$$

where  $F_2$  and  $G_2$  are the fecundities of the second age classes of the prey and predator respectively.  $P$  and  $Q$  are survival probabilities from the first to the second age classes,  $\beta_1$  and  $\beta_2$  are positive interaction parameters and  $x = x_1 + x_2, y = y_1 + y_2$ . Find the nontrivial fixed point  $(x_1^*, x_2^*, y_1^*, y_2^*)$  and show that it may not undergo a saddle node or a flip bifurcation at instability threshold. Thus stability or dynamics governed by Hopf bifurcations are the only possible dynamic outcomes.

g) If  $P = 0.6$  and  $F_2 = 25$  then the prey in absence of the predator exhibits chaotic oscillations. Now, suppose  $Q = 0.6, G_2 = 12$  and assume  $\beta = \beta_1 = \beta_2$ . Investigate numerically how the prey-predator system behaves in the following cases:  $\beta \in [0.1, 0.22]$  (weak interaction),  $\beta \in [0.4, 0.6]$  („normal” interaction),  $\beta \in [0.85, 1.00]$  (strong interaction) (see Wikan (2001)). □

**Exercise 2.8.6 (Host-Parasitoid models).** Following Kot (2001), see also the original work by Nicholson (1933), Nicholson and Bailey (1935), the books of Hassel (1978), and Edelstein-Keshet (1988), most host-parasitoid models are on the form

$$\begin{aligned} x_{t+1} &= a f(x_t, y_t) x_t \\ y_{t+1} &= c [1 - f(x_t, y_t)] x_t \end{aligned}$$

Here  $x_t$  and  $y_t$  are the number of hosts and parasitoids at time  $t$  respectively.  $f(x, y)$  is the fraction of hosts that avoids parasitoids at time  $t$  and  $a$  is the net reproductive rate of hosts.  $c$  may be interpreted as the product of the number of eggs laid per female which survive to pupate times the probability that a pupae will survive the winter and give rise to an adult next year (Maynard Smith, 1979). Kot (2001) simply refers to  $c$  as the clutch size of parasitoids.

- a) Assume that  $f(x, y) = f(y) = e^{-\beta y}$  where  $\beta > 0$  and find the nontrivial fixed point of the map. Use the Jury criteria and discuss its stability properties. What are the possible dynamic outcomes of this model?
- b) A slightly modified version of the Nicholson and Bailey model in a) which also contains a self-regulatory prey term was proposed by Beddington et al. (1975)

$$x_{1,t+1} = e^{r(1-x_t)-\beta y_t} x_t$$

$$y_{t+1} = c[1 - e^{-\beta y_t}]x_t$$

Denoting the nontrivial fixed point for  $(x^*, y^*)$ , show that

$$x^* = 1 - \frac{\beta}{r} y^* \quad 0 < y^* < \frac{r}{\beta}$$

and that  $y^*$  is the unique solution of

$$\frac{r y^*}{r - \beta y^*} - c[1 - e^{-\beta y^*}] = 0$$

Moreover, show (numerically) that there exists a parameter region where  $(x^*, y^*)$  is stable.

□

**Remark 2.8.3.** As is clear from Exercises 2.8.5a,b and 2.8.6a, if prey-predator models or host-parasitoid models shall possess a stable nontrivial equilibrium where both species exist we may not assume that one of the species is a function of only the other species. Thus, the function  $f$  in the exercises above should be on the form  $f = f(x, y)$  with properties  $\partial f / \partial x < 0$ ,  $\partial f / \partial y < 0$ . In prey-predator systems self-limitational effects are often assumed to be crowding or cannibalistic effects (the latter is typically the case in fish populations). However, what the self-regulatory effects in parasitoid species are, is far from obvious, cf. the discussion in Beddington et al. (1975), Hassel (1978), Edelstein-Keshet (1988), Murdoch (1994), and Mills and Getz (1996).

□

**Exercise 2.8.7 (Competition models).** Suppose that two species  $x$  and  $y$  compete on the same resource. From a biological point of view the competitive interaction between the two species would be that an increase of one of the species should reduce the growth of the other and vice versa. Hence, in a model of the form

$$(i) \quad \begin{aligned} x_{t+1} &= \alpha(x_t, y_t)x_t \\ y_{t+1} &= \beta(x_t, y_t)y_t \end{aligned}$$

where also self-regulatory effects are included, we should regard all partial derivatives of the functions  $\alpha$  and  $\beta$  as negative. (Note that these sign restrictions differ from the prey-predator models we studied in Exercise 2.8.5.)

a) Consider the competition model

$$(ii) \quad \begin{aligned} x_{t+1} &= (a - bx_t - c_1y_t)x_t \\ y_{t+1} &= (d - ey_t - c_2x_t)y_t \end{aligned}$$

where all constants are positive and  $a > 1, d > 1$ . Find all the fixed points of (ii). (There are four of them.)



**Empowering People. Improving Business.**

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

**BI NORWEGIAN BUSINESS SCHOOL**

EFMD **EQUIS ACCREDITED**

[www.bi.edu/master](http://www.bi.edu/master)



- b)  $(\tilde{x}, \tilde{y}) = ((a - 1)/b, 0)$  is one of the fixed points. Use the Jury criteria and find conditions for  $(\tilde{x}, \tilde{y})$  to be stable.
- c) (i) has a nontrivial fixed point  $(x^*, y^*)$ ,  $(x^* > 0, y^* > 0)$  which is a solution of the equations  $\alpha(x^*, y^*) = 1$  and  $\beta(x^*, y^*) = 1$ . Show that the solutions  $\lambda_{1,2}$  of the linearization of (i) may be expressed as (Maynard Smith, 1979)

$$\lambda_{1,2} = \frac{1}{2} \left\{ 2 - (a + d) \pm \sqrt{(a + d)^2 - 4(ad - bc)} \right\}$$

where

$$a = -x^* \frac{\partial \alpha}{\partial x} \quad d = -y^* \frac{\partial \beta}{\partial y}$$

and

$$ad - bc = x^* y^* \left( \frac{\partial \alpha}{\partial x} \frac{\partial \beta}{\partial y} - \frac{\partial \alpha}{\partial y} \frac{\partial \beta}{\partial x} \right)$$

Note that since all partial derivatives are supposed to be negative,  $a$ ,  $b$ ,  $c$  and  $d$  are positive.

- d) Explain that  $(\partial \alpha / \partial x)(\partial \beta / \partial y) > (\partial \alpha / \partial y)(\partial \beta / \partial x)$  (i.e. that the product of changes in  $\alpha$  and  $\beta$  due to self-regulatory effects are larger than the product of changes in  $\alpha$  and  $\beta$  due to the competitive species) is necessary in order for  $(x^*, y^*)$  to be stable.
- e) Discuss the possibility of having oscillatory behaviour in model (i).

For further reading of discrete competition models we refer to Adler (1990). □

**Exercise 2.8.8 (The Hénon map).** Consider the two parameter family of maps (the Hénon map)

$$H_{a,b} : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad (x, y) \rightarrow (y, 1 + bx - ay^2)$$

where  $0 < b < 1$ .

$H_{a,b}$  (in a slightly different version), was constructed and analysed by Hénon (1976), and is one of the first two-dimensional maps where there was found numerical evidence of a chaotic attractor. (Hénon's paper may also be obtained in Cvitanović (1996) where several classical papers on dynamical systems are collected.)

- a) Let  $V$  be a region elongated along the  $y$ -axis in the  $\mathbb{R}^2$  plane and consider the following maps:

$$h_1 : (x, y) \rightarrow (bx, y) \quad (\text{Contraction of } V \text{ along the } y\text{-axis})$$

$$h_2 : (x, y) \rightarrow (1 + x - ay^2, y) \quad (\text{Folding along the } x\text{-axis})$$

$$h_3 : (x, y) \rightarrow (y, x) \quad (\text{Change of orientation})$$

Show that  $H_{a,b} = h_3 \circ h_2 \circ h_1$ .

- b) Let  $a_0 = -((1-b)/2)^2$  and show that  $H_{a,b}$  has two fixed points if  $a > a_0$ , one fixed point if  $a = a_0$  and no fixed points if  $a < a_0$ .
- c) Show that  $H_{a,b}$  undergoes a saddle node bifurcation at the threshold  $a = a_0$ .
- d) Let  $a_1 = -3a_0$  and show that in the interval  $a_0 < a < a_1$  there is one stable fixed point  $(x_+^*, y_+^*)$  and one unstable fixed point  $(x_-^*, y_-^*)$ .
- e) Show by use of the Jury criteria that  $(x_+^*, y_+^*)$  undergoes a flip bifurcation at  $a = a_1$ .
- f) Show that the second iterate of  $H_{a,b}$  may be written as

$$x_{t+2} = 1 + bx_t - ay_t^2$$

$$y_{t+2} = 1 + by_t - a(1 + bx_t - ay_t^2)^2$$

and verify that whenever  $a > a_1$  there is a two-period orbit where the points are

$$(\tilde{x}_1, \tilde{y}_1) = \left( \frac{1 - a\tilde{y}_1^2}{1 - b}, \frac{1 - b + \sqrt{4a - 3(1-b)^2}}{2a} \right)$$

$$(\tilde{x}_2, \tilde{y}_2) = \left( \frac{1 - a\tilde{y}_2^2}{1 - b}, \frac{1 - b - \sqrt{4a - 3(1-b)^2}}{2a} \right)$$

- g) Show that

$$\lim_{a \rightarrow a_1} (\tilde{x}_1, \tilde{y}_1) = \lim_{a \rightarrow a_1} (\tilde{x}_2, \tilde{y}_2) = \lim_{a \rightarrow a_1} (x_+^*, y_+^*) = \left( \frac{2}{3(1-b)}, \frac{2}{3(1-b)} \right)$$



- h) Assume  $b = 1/2$  and let  $a > a_1$ . Investigate numerically if  $H_{a,1/2}$  possesses a chaotic attractor.
- i) Still assuming  $b = 1/2$ , generate a bifurcation diagram in case of  $a > a_0$ .
- j) Show that  $H_{a,b}$  has an inverse and compute  $H_{a,b}^{-1}$ .

—

Next, let  $b = 0$ . Then  $H_{a,0}$  contracts the entire  $\mathbb{R}^2$  plane onto the curve  $f_a(y) = 1 - ay^2$  and since the value of  $H_{a,0}$  is independent of the  $x$  coordinate we may study the dynamics through the one-dimensional map

$$y \rightarrow f_a(y) = 1 - ay^2$$

- k) Show that the map undergoes a saddle node bifurcation when  $a = -1/4$  and find a parameter interval where the map possesses a unique nontrivial fixed point.

## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info



1) Show that  $f_a(y)$  is topologically equivalent to the quadratic map  $g_\mu(y) = \mu y(1 - y)$ .

(Hint: Use Definition 1.2.2 and assume that  $h$  is a linear function of  $y$ . Moreover, show that the relation between  $a$  and  $\mu$  is given through  $\mu^2 - 2\mu = 4a$  and  $\mu > 1$ .)

(The case  $b = 1$  will be considered in the next exercise.) □

**Exercise 2.8.9** (Area preserving maps). Consider the map  $(x, y) \rightarrow f(x, y)$ . If the area of a region in  $\mathbb{R}^2$  is preserved under  $f$  we say that  $f$  is an area preserving map. In order to decide whether a map is area preserving or not we may apply the following theorem:

**Theorem.** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a two-dimensional map.  $f$  is area preserving if and only if  $|J| = 1$  where  $J$  is the Jacobian corresponding to  $f$ . □

A formal proof may be obtained in Stuart and Humphries (1998).

- a) Let  $b = 1$  in the Hénon map (cf. Exercise 2.8.8), and show that  $H_{a,1}$  is area preserving.
- b) Show that the map  $(x, y) \rightarrow (-xy, \ln x)$  is area preserving too.
- c) Compute all nontrivial fixed points of the maps in a) and b) and decide whether the fixed points are hyperbolic or not.
- d) In general, what can you say about the eigenvalues of the linearization of an area preserving map? □

## 2.9 Difference-Delay equations

Difference-Delay equations are equations of the form

$$x_{t+1} = f(x_t, x_{t-T}) \tag{2.9.1}$$

where  $T$  is called the delay.

Referring to population dynamical studies, equation (2.9.1) is often used when one considers species where there is a substantial time  $T$  from birth to sexual maturity. Hence, instead of using a detailed Leslie matrix model where the fecundities  $F_i = 0$  for several age classes, the more aggregated form (2.9.1) is often preferred.

One frequently quoted example is Colin Clark's Baleen whale model (Clark, 1976)

$$x_{t+1} = ux_t + F(x_{t-T}) \quad (2.9.2)$$

where  $x_t$  is the adult breeding population.  $u$  ( $0 \leq u \leq 1$ ) may be interpreted as a survival coefficient and the term  $F(x_{t-T})$  is the recruitment which takes place with a delay of  $T$  years. In case of the Baleen whale,  $5 \leq T \leq 10$ .

A slightly modified version of (2.9.2) was presented by the International Whaling Commission (IWC) as

$$x_{t+1} = (1 - u)x_t + R(x_{t-T}) \quad (2.9.3)$$

Here (just as in (2.9.2)),  $(1 - u)x_t$ ,  $0 < u < 1$ , is the fraction of the adult whales that survives at time  $t$  and enters the population one time step later.

$$R(x_{t-T}) = \frac{1}{2}(1 - u)^T x_{t-T} \left\{ P + Q \left[ 1 - \left( \frac{x_{t-T}}{K} \right)^z \right] \right\} \quad (2.9.4)$$

and regarding the parameters in (2.9.4) we refer to IWC report no. 29, Cambridge (1979). Other models where a variety of different species are considered may be obtained in Botsford (1986), Tuljapurkar et al. (1994), Higgins et al. (1997), see also Kot (2001) and references therein.

—

Now, returning to the general nonlinear equation (2.9.1), the fixed point  $x^*$  is found by letting  $x_{t+1} = x_t = x_{t-T} = x^*$ . The stability analysis follows the same pattern as in section 2.4. Let  $x_t = x^* + \xi_t$  where  $|\xi_t| \ll 1$ . Then from (2.9.1)

$$x^* + \xi_{t+1} \approx f(x^*, x^*) + \frac{\partial f}{\partial x_t}(x^*)\xi_t + \frac{\partial f}{\partial x_{t-T}}(x^*)\xi_{t-T} \quad (2.9.5)$$

Thus the linearization becomes

$$\xi_{t+1} = a\xi_t + b\xi_{t-T} \quad (2.9.6)$$

where  $a$  and  $b$  are  $\partial f/\partial x_t$ ,  $\partial f/\partial x_{t-T}$  evaluated at equilibrium respectively.

The solution of (2.9.6) is found by letting  $\xi_t = \lambda^t$  which after some rearrangements result in the eigenvalue equation

$$\lambda^{T+1} - a\lambda^T - b = 0 \quad (2.9.7)$$

which we recognize as a polynomial equation of degree  $T + 1$ . As before,  $|\lambda| < 1$  guarantees that  $x^*$  is locally asymptotic stable. The transfer from stability to instability occurs when  $x^*$  fails to be hyperbolic which means that  $\lambda$  crosses the unit circle through 1, through  $-1$  or crosses the unit circle at the location  $\exp(i\theta)$ .

**Brain power**

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**



**Example 2.9.1.** Compute the nontrivial fixed point  $x^*$  and derive the eigenvalue equation of the model

$$x_{t+1} = x_t \exp \left[ r \left( 1 - \frac{x_{t-T}}{K} \right) \right] \quad (2.9.8)$$

where  $r$  and  $K$  both are positive. ((2.9.8) is often called the delayed Ricker model and the parameters may be interpreted as the intrinsic growth rate ( $r$ ) and the carrying capacity ( $K$ )).

The fixed point obeys

$$x^* = x^* \exp \left[ r \left( 1 - \frac{x^*}{K} \right) \right]$$

so clearly,  $x^* = K$ .

The coefficients  $a$  and  $b$  in (2.9.7) become

$$\begin{aligned} a &= \frac{\partial f}{\partial x_t}(x^*) = 1 \cdot \exp \left[ r \left( 1 - \frac{K}{K} \right) \right] = 1 \\ b &= \frac{\partial f}{\partial x_{t-T}}(x^*) = K \left( -\frac{r}{K} \right) \exp \left[ r \left( 1 - \frac{K}{K} \right) \right] = -r \end{aligned}$$

Hence, the eigenvalue equation may be cast in the form

$$\lambda^{T+1} - \lambda^T + r = 0 \quad (2.9.9)$$

□

**Exercise 2.9.1.** Consider the difference-delay equation

$$x_{t+1} = x_t \left[ 1 + r \left( 1 - \frac{x_{t-T}}{K} \right) \right] \quad (2.9.10)$$

and repeat the calculations from the previous example.

□

**Exercise 2.9.2.** Repeat the calculations in Exercise 2.9.1 for the equation

$$x_{t+1} = \frac{\alpha x_t}{1 + \beta x_{t-T}} \quad (2.9.11)$$

□

—

Let us now turn back to the general eigenvalue equation (2.9.7). Although it is a polynomial equation of degree  $T + 1$ , its structure is simpler than most of the equations which we studied in Part II. Therefore, unless the delay  $T$  becomes too large, the Jury criteria work excellent when one tries to reveal stability properties. (The “Baleen whale equations” (2.9.4), (2.9.5) were analyzed by use of Theorem 2.1.9.) It is also possible to use (2.9.7) in order to give a thorough description of the dynamics in parameter regions where the fixed point is stable.

Our next goal is to demonstrate this by use of the difference-delay equation (2.9.8) and its associated eigenvalue equation (2.9.9).

As a prelude to the general situation, suppose that  $T = 0$  (no delay) in (2.9.8), (2.9.9). Then, from (2.9.9),  $\lambda = 1 - r$  from which we may draw the following conclusions: (i) If  $0 < r < 1$ , then  $0 < \lambda < 1$ , hence from a given initial condition we will experience a monotonic damping towards the fixed point  $x^* = K$ . (ii)  $1 < r < 2$  implies that  $-1 < \lambda < 0$ , thus in this case there will be oscillatory damping towards  $x^*$ . (iii) At instability threshold  $r = 2$  it follows that  $\lambda = -1$  and a supercritical flip bifurcation occurs (cf. Exercise 1.5.1). Consequently, in case of  $r > 2$  but  $|r - 2|$  small, the dynamics is a stable period-2 orbit.

Next, consider the small delay  $T = 1$ . Then (2.9.9) becomes  $\lambda^2 - \lambda + r = 0$  and by use of (2.1.14) stability of  $x^* = K$  is ensured whenever the inequalities  $r > 0$ ,  $r + 2 > 0$  and  $r < 1$  are satisfied. Hence, at instability threshold  $r = 1$  but in contrast to the case  $T = 0$  it also follows from (2.1.14) that  $\lambda$  is a complex number at bifurcation threshold.

If  $T = 2$  the eigenvalue equation may be written as  $\lambda^3 - \lambda^2 + r = 0$  and the four Jury criteria (2.1.16) simplify to  $r > 0$ ,  $2 - r > 0$ ,  $r < 1$  and  $r < (1/2)(\sqrt{5} - 1) \approx 0.6180$  respectively. Clearly,  $r = (1/2)(\sqrt{5} - 1)$  at bifurcation threshold and again we observe that  $\lambda$  is a complex number.

Now, consider the general case  $T \geq 1$ . From our findings above it is natural to assume that  $\lambda = \exp(i\theta)$  when the fixed point  $x^* = K$  loses its hyperbolicity. Moreover, the value of  $r$  at instability threshold becomes smaller as  $T$  increases which suggests that an increase of  $T$  acts as a destabilizing effect. Substituting  $\lambda = \exp(i\theta)$  into (2.9.9) gives

$$e^{i(T+1)\theta} = e^{iT\theta} - r \tag{2.9.12}$$

which after multiplication by  $\exp(-i(T + 1)\theta)$  may be written as

$$1 = e^{-i\theta} - r e^{-i(T+1)\theta} \quad (2.9.13)$$

Therefore

$$1 = \cos \theta - i \sin \theta - r \cos(T + 1)\theta + ir \sin(T + 1)\theta$$

and by separating into real and imaginary parts we arrive at

$$1 = \cos \theta - r \cos(T + 1)\theta \quad (2.9.14a)$$

$$0 = -\sin \theta + r \sin(T + 1)\theta \quad (2.9.14b)$$

Finally, by squaring both equations (2.9.14) and then add we obtain the relation between  $r$  and  $\theta$  as

$$r = 2 [\cos \theta \cos(T + 1)\theta + \sin \theta \sin(T + 1)\theta] = 2 \cos T\theta \quad (2.9.15)$$

## TURN TO THE EXPERTS FOR SUBSCRIPTION CONSULTANCY

**Subscribe is one of the leading companies in Europe when it comes to innovation and business development within subscription businesses.**

**We innovate new subscription business models or improve existing ones. We do business reviews of existing subscription businesses and we develop acquisition and retention strategies.**

**Learn more at [linkedin.com/company/subscribe](https://www.linkedin.com/company/subscribe) or contact  
Managing Director Morten Suhr Hansen at [mha@subscribe.dk](mailto:mha@subscribe.dk)**

**SUBSCR**✓**BE** - to the future



Substituting back into (2.9.13) then implies

$$1 = e^{-i\theta} - 2 \left( \frac{e^{iT\theta} + e^{-iT\theta}}{2} \right) e^{-i(T+1)\theta}$$

and through multiplication by  $\exp(i\theta)$  we get

$$e^{i\theta} = -e^{-i2T\theta} = e^{i(\pi-2T\theta)} \quad (2.9.16)$$

Thus

$$\theta = \pi - 2T\theta + 2k\pi \quad (2.9.17)$$

Hence

$$\theta = \frac{(2k+1)\pi}{2T+1} \quad (2.9.18)$$

From this we may draw the following conclusion. Since  $r = 2 \cos T\theta$  there are several values of  $r$  which result in modulus 1 solutions of the eigenvalue equation (2.9.9). The smallest  $r$  which results in a modulus 1 solution is clearly when  $k = 0$ , i.e.

$$r_2 = 2 \cos \frac{T\pi}{2T+1} \quad (2.9.19)$$

Let us now focus on possible real solutions of the eigenvalue equation (2.9.9). Assume  $\lambda = R$  ( $R$  - real). Then from (2.9.9):

$$r = R^T - R^{T+1} \quad (2.9.20)$$

and since  $r > 0$ ,  $T > 0$  it follows that  $R < 1$ . Moreover,

$$\frac{dr}{dR} = R^{T-1} [T - (T+1)R]$$


such that the maximum value of  $r$  occurs when

$$R = \frac{T}{T+1} \quad (2.9.21)$$

Hence,  $R$  is a positive number and the corresponding maximum value of the intrinsic growth rate  $r$  is

$$r_1 = \frac{T^T}{(T+1)^{T+1}} \quad (2.9.22)$$

**Exercise 2.9.3.** Show that  $\lim_{T \rightarrow \infty} r_1 = 1/Te$ . □



**Figure 24:** The graph of  $r(T) = R^T - R^{T+1}$  ( $T = 2$ ).

In Figure 24 we have drawn the graph of (2.9.20) in the case  $T = 2$ . (The graph has a similar form for other  $T \geq 1$  values.) Thus, when  $R$  is increasing from 0 to  $T/(T + 1)$ ,  $r$  will increase from 0 to  $r_1$  and when  $R$  increases from  $T/(T + 1)$  to 1,  $r$  will decrease from  $r_1$  to 0. Clearly, if  $0 < r < r_1$ , (2.9.20) has two positive roots. If  $r > r_1$ , there are no positive roots.

Following Levin and May (1976) we now have

$$r_1 = \frac{T^T}{(T + 1)^{T+1}} \leq \frac{1}{2} r_2 = \cos \frac{T\pi}{2T + 1} \quad (2.9.23)$$

Indeed, first observe that


$$\cos \frac{T\pi}{2T + 1} = -\sin \left( \frac{T\pi}{2T + 1} - \frac{\pi}{2} \right) = \sin \frac{\pi}{2(2T + 1)} > \frac{2}{\pi} \cdot \frac{\pi}{2(2T + 1)} = \frac{1}{2T + 1}$$

Next, by rewriting  $r_1$ :

$$r_1 = \frac{1}{(T + 1)(1 + \frac{1}{T})^T} \leq \frac{1}{(T + 1)2} < \frac{1}{2T + 1}$$

which establishes (2.9.23).

—



**Figure 25:** 30 iterations of  $x_{t+1} = x_t \exp[r(1 - x_{t-1})]$ . Monotonic orbit,  $r = 0.24$ . Oscillatory orbit,  $r = 0.90$ .

**What do you want to do?**


No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



Now, considering an orbit starting from an initial value  $x_0 \neq K$ , we may from the findings above conclude that in case of  $0 < r < r_1$  the orbit may approach  $x^* = K$  monotonically. If  $r_1 < r < r_2$  the orbit will always approach  $x^*$  as a convergent oscillation. If  $r > r_2$ ,  $x^*$  is unstable and an orbit will act as a divergent oscillation towards a limit cycle (provided the bifurcation is supercritical). These cases are demonstrated in Figure 25 and Figure 26. In Figure 25 we show the behaviour of the map  $x_{t+1} = x_t \exp[r(1 - x_{t-1})]$  (i.e.  $K = T = 1$  in (2.9.8)) in case of  $r = 0.24$  ( $< r_1$ ) and  $r = 0.90$  ( $r_1 < r < r_2$ ) respectively and clearly, one orbit ( $r = 0.24$ ) approaches the fixed point  $x^* = 1$  monotonically while the other orbit ( $r = 0.90$ ) approaches  $x^*$  in an oscillatory way. In Figure 26  $r = 1.02$  and  $r = 1.10$  ( $r > r_2$ ) and there is no convergence towards  $x^*$ . Note that the orbit with small amplitude ( $r = 1.02$ ) is almost 6-periodic.



**Figure 26:** 30 iterations of  $x_{t+1} = x_t \exp[r(1 - x_{t-1})]$ . Small amplitude orbit,  $r = 1.02$ . Large amplitude orbit,  $r = 1.10$ .

**Remark 2.9.1.** Note that in the case  $0 < r < r_1$  we have not actually proved that an orbit must approach  $x^*$  monotonically. After all (2.9.9) may have complex solutions with magnitudes larger than  $\lambda = R = T/(T + 1)$ . However, this is not the case as is proved in Levin and May (1976). (The proof is not difficult, it involves the same kind of computations as we did when (2.9.19) was derived.) □

—

Let us now comment on possible periodic dynamics. Referring to section 2.8 “Beyond the Hopf bifurcation” we learned that although the dynamics was a quasistationary orbit just beyond the Hopf bifurcation threshold, the dynamics could be periodic (exact or approximate) as we penetrated deeper into the unstable parameter region. Periodic phenomena may of course also occur in difference-delay equations. Indeed, consider

$$x_{t+1} = x_t \exp[1 - r(1 - x_{t-1})] \quad (2.9.24)$$

which is nothing but (2.9.8) where  $T = K = 1$ . At bifurcation threshold the dominant eigenvalue becomes (see (2.9.18))

$$\lambda_D = \exp(i\theta) = \exp\left(i\frac{\pi}{3}\right) \quad (2.9.25)$$

and since  $\lambda_D^6 = \exp(2\pi i) = 1$ ,  $\lambda_D$  is equal to 6th root of unity at bifurcation threshold. Therefore, in case of  $\lambda > \lambda_D$  but  $|\lambda - \lambda_D|$  small,  $\arg \lambda$  is still close to  $\pi/3$  which definitely signals 6-periodic dynamics. That the dynamics is almost 6-periodic is clearly demonstrated in Figure 26 ( $r = 1.02$ ). Through an enlargement of  $r$  ( $r = 1.10$ ) the periodicity is not so profound as the other orbit in Figure 26 shows. More about periodic phenomena in difference delay equations may be obtained in Diekmann and Gils (2000).

—

In one way the results presented above are somewhat special in the sense that we were able to find the complex eigenvalues at bifurcation threshold on closed form (cf. (2.9.18)). Typically, this is not the case. However, the method we used may still be fruitful in order to bring equations where it is difficult to locate modulus 1 solutions numerically to a form where it is much more simple. This fact will now be demonstrated through one example and one exercise.

**Example 2.9.2.** In Example 2.4.4 (section 2.4) we studied a  $(n \times 1) \times (n \times 1)$  Leslie matrix model with equal fecundities  $F$ . If we in addition assume that the year to year survival probabilities are equal, i.e.  $P_0 = P_1 = \dots = P_{n-1} = P$ ,  $0 < P < 1$ , the eigenvalue equation (2.4.17) may be cast in the form

$$\lambda^{n+1} - \frac{1}{D}(1 - x^*) \sum_{i=0}^n P^i \lambda^i = 0 \quad (2.9.26)$$

where

$$x^* = \ln(FD) \quad \text{and} \quad D = 1 + P + P^2 + \dots + P^n = \frac{1 - P^{n+1}}{1 - P}$$

Our goal is to locate complex modulus 1 solutions of (2.9.26) for given values of  $P$ . Using the fact that  $\sum P^i \lambda^i$  is nothing but a geometric series, it is straightforward to rewrite (2.9.26) as

$$\lambda^{n+2} + A\lambda^{n+1} - B = 0 \quad (2.9.27)$$

where

$$A = \frac{(1-P)(x^* - 1)}{1 - P^{n+1}} - P \quad \text{and} \quad B = \frac{(1-P)(x^* - 1)}{1 - P^{n+1}} P^{n+1}$$

By inspection, (2.9.27) has a root  $\lambda = P$  which is located inside the unit circle. The  $n + 1$  other roots of (2.9.27) are the same roots as of (2.9.26). Hence, assume that  $\lambda = \exp(i\theta)$  in (2.9.27). Then (we urge the reader to perform the necessary calculations), by using the same method as we did when we derived (2.9.14) from (2.9.12) we find that

$$\sin \theta = -B \sin(n + 1)\theta \quad (2.9.28a)$$

$$\cos \theta = \frac{B^2 - A^2 - 1}{2A} \quad (2.9.28b)$$

**gaiteye**  
Challenge the way we run

**EXPERIENCE THE POWER OF  
FULL ENGAGEMENT...**

**RUN FASTER.  
RUN LONGER..  
RUN EASIER...**

**READ MORE & PRE-ORDER TODAY  
WWW.GAITEYE.COM**

We know from Example 2.4.4 that the fixed point of (2.4.7) is stable in case of small equilibrium populations  $x^*$ . Therefore, numerically it is now easy to find the solutions of (2.9.27) (and (2.9.26)) at bifurcation threshold for given values of  $n$  and  $P$  by simply increasing  $F$  which means that  $x^*$  is increased too and compute  $B$  up to the point where (2.9.28a) is satisfied. Then we compute the corresponding value of  $A$  and finally  $\theta$  through (2.9.28b) as

$$\theta = \arccos\left(\frac{B^2 - A^2 - 1}{2A}\right) \quad (2.9.29)$$

□

**Exercise 2.9.4.** Consider the eigenvalue equation

$$\lambda^T - (1 + a)b\lambda^{T-1} + ab^2\lambda^{T-2} = D \quad (2.9.30)$$

where  $D$  is real,  $0 < a \leq 1$ ,  $a < 1/b$ .

a) Show that (2.9.30) may be written as

$$\lambda^{T-2}(\lambda - ab)(\lambda - b) = D$$

b) Assume that  $D = 0$  and conclude that the dominant root of the eigenvalue equation is  $\lambda = b$  if  $0 < a < 1$  or  $\lambda = ab$  if  $1 < a < 1/b$ .

c) Suppose  $D \neq 0$  and assume that  $\lambda = R$  is real and positive. Show that the maximum value of  $D$  is

$$R^T - (1 + a)bR^{T-1} + ab^2R^{T-2}$$

where

$$R = \frac{(1 + a)b(T - 1)}{2T} + \sqrt{\frac{(1 + a)^2b^2(T - 1)^2}{4T^2} - \frac{ab^2(T - 2)}{T}}$$

d) Assume that  $\lambda = \exp(i\theta)$  and separate (2.9.30) in its real and imaginary parts respectively.

Explain how  $\theta$  and  $D$  may be found numerically in case of given values of  $a$ ,  $b$  and  $T$ .

(Equation (2.9.30) arises in an analysis of the general Deriso-Schute model. A thorough discussion of the model may be obtained in Bergh and Getz (1988).)

□



In this section we have used a variety of different techniques in order to find the roots of polynomial equations. We close by stating Descartes' rule of signs which is a theorem that also may give valuable insight of location of the roots.

**Theorem 2.9.1** (Descartes' rule of signs). Consider the polynomial equation

$$a_0\lambda^n + a_1\lambda^{n-1} + a_2\lambda^{n-2} + \dots + a_{n-1}\lambda + a_n = 0$$

where  $a_n > 0$ .

Let  $k$  be the number of sign changes between the coefficients  $a_n, a_{n-1}, \dots, a_0$  disregarding any which are zero. Then there are at most  $k$  roots which are real and positive and, moreover, there are either  $k$  or  $k - 2$  or  $k - 4$  ... real positive roots.  $\square$

**Example 2.9.3.** Consider

$$\lambda^{T+1} - \lambda^T + r = 0$$

where  $r > 0$ . Here  $k = 2$ , hence there are at most 2 real positive roots and, moreover, there are either 2 or 0 such roots.

Next, suppose that  $\lambda = -\sigma$ .

- 1) If  $T$  is an even number, the equation may be written as  $-\sigma^{T+1} - \sigma^T + r = 0$ . Thus there is only one change of sign, consequently there is exactly 1 negative root  $\lambda$  of  $\lambda^{T+1} - \lambda^T + r = 0$ . (From our previous analysis of (2.9.9) this means that if  $0 < r < r_1$ , there are 2 positive roots, 1 negative root and  $T - 2$  complex roots. If  $r_1 < r$ , there are  $T$  complex roots and 1 negative root.)
- 2) If  $T$  is an odd number, the equation may be cast in the form  $\sigma^{T+1} + \sigma^T + r = 0$ . Hence, there are no sign changes so there are no negative roots  $\lambda$ . (Thus  $0 < r < r_1$  implies 2 positive roots and  $T - 1$  complex roots. If  $r_1 < r$ , all  $T + 1$  roots are complex.)  $\square$

# Part III

## Discrete Time Optimization Problems

### 3.1 The fundamental equation of discrete dynamic programming

In the following sections we shall give a brief introduction to discrete dynamic optimization. When one wants to solve problems within this field there are mainly two methods (together with several numerical alternatives which we will not treat here) available. Here, in section 3.1, we shall state and prove the fundamental equation of discrete dynamic programming which perhaps is the most frequently used method. In section 3.2 we shall solve optimization problems by use of a discrete version of the maximum principle.

Dynamic optimization is widely used within several scientific branches like economy, physics and biology. As an introduction to the kind of problems that we want to study, let us consider the following example:

**Example 3.1.1.** Let  $x_t$  be the size of a population at time  $t$ . Further, assume that  $x$  is a species of commercial interest so let  $h_t \in [0, 1]$  be the fraction of the population that we harvest at each time. Therefore, instead of expressing the relation between  $x$  at two consecutive time steps as  $x_{t+1} = f(x_t)$  or (if the system is nonautonomous)  $x_{t+1} = f(t, x_t)$ , we shall from now on assume that

$$x_{t+1} = f(t, x_t, h_t) \tag{3.1.1}$$

If the function  $f$  is the quadratic or the Ricker function which we studied in Part I, (3.1.1) may be written as

$$x_{t+1} = r(1 - h_t)x_t[1 - (1 - h_t)x_t] \tag{3.1.2}$$

or

$$x_{t+1} = (1 - h_t)x_t \exp[r(1 - (1 - h_t)x_t)] \tag{3.1.3}$$

respectively. In case of an age-structured population model (cf. the various examples treated in part II) the equation  $\mathbf{x}_{t+1} = f(t, \mathbf{x}_t, \mathbf{h}_t)$  may be expressed as

$$x_{1,t+1} = F_1 e^{-x_t} x_{1,t} (1 - h_{1,t}) + F_2 e^{-x_t} x_{2,t} (1 - h_{1,t}) \tag{3.1.4}$$

$$x_{2,t+1} = P x_{1,t} (1 - h_{2,t})$$

(For simplicity, it is often assumed that  $h_t = h$  and  $h_{i,t} = h_i$  which means that the population or the age classes are exposed to harvest with constant harvest rate(s).)

Now, returning to equation (3.1.1), assume that  $\pi_t = f_0(t, x_t, h_t)$  is the profit we can make of the harvested part of the population at time  $t$ . Our ultimate goal is to maximize the profit over a time period from  $t = 0$  to  $t = T$ , i.e. we want to maximize the sum of the profits at times  $t = 0, 1, \dots, T$ . This leads to the problem

$$\text{maximize}_{h_0, h_1, \dots, h_T} \sum_{t=0}^T f_0(t, x_t, h_t) \quad (3.1.5)$$

subject to equation (3.1.1) given the initial condition  $x_0$  and  $h_t \in [0, 1]$ .

To be somewhat more precise, we have arrived at the following situation: Suppose that we at time  $t = 0$  apply the harvest rate  $h_0$ . Then, according to (3.1.1)  $x_1 = f(0, x_0, h_0)$  is known at time  $t = 1$ . Further, assume that we at time  $t = 1$  choose the harvest  $h_1$ . Then  $x_2 = f(1, x_1, h_1)$  is known and continuing in this fashion, applying (different) harvest rates  $h_t$  at each time we also know the value of  $x_t$  at each time. Consequently, we also know the profit  $\pi_t = f_0(t, x_t, h_t)$  at each time. As stated in (3.1.5) our goal is to choose  $h_0, h_1, \dots, h_T$  in such a way that  $\sum_{t=0}^T f_0(t, x_t, h_t)$  is maximized.  $\square$

—

Let us now formulate the situation described in Example 3.1.1 in a more general context. Suppose that the *state* variable  $x$  evolves according to the equation  $x_{t+1} = f(t, x_t, u_t)$  where  $x_0$  is known. At each time  $t$  the path that  $x$  follows depends on discrete *control* variables  $u_0, u_1, \dots, u_T$ . (In Example 3.1.1 we used harvest rates as control variables.) We assume that  $u_t \in U$  where  $U$  is called the control region. The sum  $\sum_{t=0}^T f_0(t, x_t, u_t)$  where  $f_0$  is the quantity we wish to maximize is called the objective function.

**Definition 3.1.1.** Suppose that  $x_s = x$ . Then we define the value function as

$$J_s(x) = \text{maximize}_{u_s, u_{s+1}, \dots, u_T} \sum_{t=s}^T f_0(t, x_t, u_t) \quad (3.1.6)$$

$\square$

Hence, a more general formulation of the problem we considered in Example 3.1.1 is: maximize  $J_s(x)$  subject to  $x_{t+1} = f(t, x_t, u_t)$ ,  $x_s = x$  and  $u_t \in U$ .

We now turn to the question of how to solve the problem.

Suppose that we know the optimal control (optimal with respect to maximizing (3.1.6))  $u_s^*$  at  $s = 0$ . Then, according to the findings presented in Example 3.1.1, we find the corresponding  $x_1^*$  as  $x_1^* = f(0, x_0, u_0^*(x_0))$  and if we succeed in finding the optimal control  $u_1^*(x_1^*)$  at time  $t = 1$  we have  $x_2^* = f(1, x_1^*, u_1^*(x_1^*))$  and so on. Thus, suppose that  $x_s = x$  at time  $t = s$ , how shall we choose  $u_s$  in the best optimal way? Clearly, if we choose  $u_s = u$  as the optimal control we achieve the immediate benefit  $f_0(s, x, u)$  and also  $x_{s+1} = f(s, x, u)$ . This consideration simply means that the highest total benefit which is possible to get from time  $s + 1$  to  $T$  is  $J_{s+1}(x_{s+1}) = J_{s+1}(f(s, x, u))$ . Hence, the best choice of  $u_s = u$  at time  $s$  is the one that maximizes  $f_0(s, x, u) + J_{s+1}(f(s, x, u))$ . Consequently, we have the following theorem:

**Theorem 3.1.1.** Let  $J_s(x)$  defined through (3.1.6) be the value function for the problem

$$\underset{u}{\text{maximize}} \sum_{t=0}^T f_0(t, x_t, u_t) \text{ subject to } x_{t+1} = f(t, x_t, u_t)$$

where  $u_t \in U$  and  $x_0$  are given. Then

$$J_s(x) = \max_{u \in U} [f_0(s, x, u) + J_{s+1}(f(s, x, u))] , \quad s = 0, 1, \dots, T - 1 \quad (3.1.7)$$

$$J_T(x) = \max_{u \in U} f_0(T, x, u) \quad (3.1.8)$$

□

Theorem 3.1.1 is often referred to as the fundamental equation(s) of dynamical programming and serves as one of the basic tools for solving the kind of problems that we considered in Example 3.1.1. As we shall demonstrate through several examples, the theorem works “backwards” in the sense that we start to find  $u_T^*(x)$  and  $J_T(x)$  from (3.1.8). Then we use (3.1.7) in order to find  $J_{T-1}(x)$  together with  $u_{T-1}^*(x)$  and so on. Hence, all value functions and optimal controls are found recursively.

**Example 3.1.2.**

$$\underset{u}{\text{maximize}} \sum_{t=0}^T (x_t + u_t) \text{ subject to } x_{t+1} = x_t - 2u_t, \quad u_t \in [0, 1], \quad x_0 \text{ given}$$

**Solution:** From (3.1.8),  $J_T(x) = \max_u (x + u)$  so clearly, the optimal value of  $u$  is  $u = 1$ .

Hence at time  $t = T$ ,  $J_T(x) = x + 1$  and  $u_T^*(x) = 1$ .

Further, from (3.1.7):

$$J_{T-1}(x) = \max_u [x + u + J_T(x - 2u)] = \max_u [x + u + (x - 2u + 1)] = \max_u [2x - u + 1].$$

Consequently,  $u = 0$  is the optimal choice, thus at  $t = T - 1$  we have  $J_{T-1}(x) = 2x + 1$  and  $u_{T-1}^*(x) = 0$ .

This implies:  $J_{T-2}(x) = \max_u [x + u + J_{T-1}(x - 2u)] = \max_u [3x - 3u + 1]$  so again  $u = 0$  is the best choice and  $J_{T-2}(x) = 3x + 1$  and  $u_{T-2}^*(x) = 0$ .

From the findings above it is natural to suspect that in general

$$J_{T-k}(x) = (k + 1)x + 1, \quad u_{T-k}^*(x) = 0, \quad k = 1, 2, \dots, T$$

The formulae is obviously correct in case of  $k = 1$  and by induction we have from (3.1.7) that

$$\begin{aligned} J_{T-(k+1)} &= \max_u [x + u + J_{T-k}(x - 2u)] \\ &= \max_u [x + u + (k + 1)(x - 2u) + 1] = \max_u [(k + 2)x - 2(k + 1)u + 1] \\ &= (k + 2)x + 1 = [(k + 1) + 1]x + 1 \end{aligned}$$

hence the formulae is correct at time  $T - (k + 1)$  as well. Therefore

$$\begin{aligned} J_{T-k}(x) &= (k + 1)x + 1, & u_{T-k}^*(x) &= 0, & k &= 1, 2, \dots, T \\ J_T(x) &= x + 1 & u_T^*(x) &= 1 \end{aligned}$$

□

**Example 3.1.3.**

maximize  $\sum_{t=0}^T (-u_t^2 + u_t - x_t)$  subject to  $x_{t+1} = x_t + u_t, u_t \in \langle -\infty, \infty \rangle, x_0$  given

**Solution:** From (3.1.8),  $J_T(x) = \max_u (-u^2 - x + u)$  and since the function

$h(u) = -u^2 - x + u$  clearly is concave in  $u$  the optimal choice of  $u$  must be the solution of

$h'(u) = 0$ , i.e.  $u = 1/2$ . Hence, at time  $t = T$ ,  $u_T^*(x) = 1/2$  and

$$J_T(x) = -(1/4) - x + (1/2) = -x + (1/4).$$

Further, (3.1.7) gives

$J_{T-1}(x) = \max_u[-u^2 - x + u + J_T(x + u)] = \max_u[-u^2 - x + u - (x + u) + (1/4)] = \max_u[-u^2 - 2x + (1/4)]$  and again since  $h_1(u) = -u^2 - 2x + (1/4)$  is concave in  $u$  we find that  $u = 0$  is the optimal choice. Thus  $J_{T-1}(x) = -2x + (1/4)$  and  $u_{T-1}^*(x) = 0$ .

Proceeding in the same way (we urge the reader to work through the details) we find that  $J_{T-2}(x) = -3x + (1/2)$ ,  $u_{T-2}^*(x) = -(1/2)$  and  $J_{T-3}(x) = -4x + (3/2)$ ,  $u_{T-3}^*(x) = -1$ .

Therefore, it is natural to suppose that

$$J_{T-k}(x) = -(k + 1)x + b_k$$

where  $b_0 = 1/4$  and  $u_{T-k}^*(x) = -\frac{k-1}{2}$ ,  $k = 1, 2, \dots, T$ . The formulae is obviously correct when  $k = 0$  and by induction

$$\begin{aligned} J_{T-(k+1)} &= \max_u[-u^2 + u - x + J_{T-k}(x + u)] \\ &= \max_u[-(k + 2)x - u^2 - ku + b_k] \end{aligned}$$

Again, we observe that the function inside the bracket is concave in  $u$  so its maximum occurs at  $u = -(k/2)$  which means that the corresponding value function becomes

$$J_{T-(k+1)}(x) = -[(k + 1) + 1]x + b_k + k^2/4 = -[(k + 1) + 1]x + b_{k+1}$$

It remains to find  $b_k$ . The equation  $b_{k+1} - b_k = k^2/4$  has the homogeneous solution  $C \cdot 1^k = C$ . Referring to the remark following Example 3.1.4 we assume a particular solution of the form  $p_k = (A + Bk + Dk^2)k$ . Hence, after inserting into the equation and equating terms of equal power of  $k$  we find that  $A = 1/24$ ,  $B = -(1/8)$  and  $D = 1/12$  so the general solution becomes  $b_k = C + (1/24)k - (1/8)k^2 + (1/12)k^3$ . Finally, using the fact that  $b_0 = 1/4$  which implies that  $C = 1/4$ , we obtain

$$J_{T-k}(x) = -(k + 1)x + \frac{1}{24}(6 + k - 3k^2 + 2k^3) \text{ and } u_{T-k}^*(x) = -\frac{k-1}{2}$$

□



**Example 3.1.4 (Exam exercise, UiO).**

$$\text{maximize}_u \sum_{t=0}^T (x_t - u_t) \text{ subject to } x_{t+1} = u_t x_t, \quad u_t \in [0, 2], \quad x_0 \text{ given}$$

**Solution:**  $J_T(x) = \max_u (x - u)$ . Clearly,  $u = 0$  is the optimal choice so  $J_T(x) = x$  and  $u_T^*(x) = 0$ ,  $J_{T-1}(x) = \max_u [x - u + J_T(ux)] = \max_u [x + (x - 1)u]$ . Thus, if  $x \geq 1$  we choose  $u = 2$  and if  $x < 1$  we choose  $u = 0$ . Consequently,

$$J_{T-1}(x) = \begin{cases} x + (x - 1)2 = 3x - 2 & \text{if } x \geq 1 \text{ and } u_{T-1}^*(x) = 2 \\ x + (x - 1)0 = x & \text{if } x < 1 \text{ and } u_{T-1}^*(x) = 0 \end{cases}$$

(Note that  $J_{T-1}(x)$  is a convex function which is continuous at  $x = 1$ .)

In order to compute  $J_{T-2}(x)$  we must consider the cases  $J_{T-1}(x) = 3x - 2$  and  $J_{T-1}(x) = x$  separately.

This e-book  
is made with  
**SetaPDF**





**SETASIGN**

PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)



Assuming  $J_{T-1}(x) = 3x - 2$  we obtain

$$J_{T-2}(x) = \max_u [x - u + 3ux - 2] = \max_u [x + (3x - 1)u - 2]$$




Figure 27:  $J_{T-2}(x)$  possibilities.

so if  $x \geq 1/3$  our optimal choice is  $u = 2$  and if  $x < 1/3$  we choose  $u = 0$ .

In the same way, using  $J_{T-1}(x) = x$ , we find

$$J_{T-2}(x) = \max_u [x - u + ux] = \max_u [x + (x - 1)u]$$

so whenever  $x \geq 1$ ,  $u = 2$  and if  $x < 1$  our best choice is  $u = 0$ .

Hence, the possibilities are

$$J_{T-2}(x) = \begin{cases} x + (3x - 1) \cdot 2 - 2 = h_1(x) = 7x - 4 & \text{if } x \geq 1/3 \\ x + (3x - 1) \cdot 0 - 2 = h_2(x) = x - 2 & \text{if } x < 1/3 \\ x + (x - 1) \cdot 2 = h_3(x) = 3x - 2 & \text{if } x \geq 1 \\ x + (x - 1) \cdot 0 = h_4(x) = x & \text{if } x < 1 \end{cases}$$

In Figure 27 we have drawn the graphs of the  $h_i$  functions in their respective domains. The point of intersection between  $h_1(x)$  and  $h_4(x)$  is  $x = 2/3$  so clearly, if  $x \geq 2/3$ ,  $h_1(x)$  is the largest function. If  $x < 2/3$ ,  $h_4(x)$  is the largest function.

Consequently, we conclude that

$$J_{T-2}(x) = \begin{cases} 7x - 4 & \text{if } x \geq 2/3 \text{ and } u_{T-2}^*(x) = 2 \\ x & \text{if } x < 2/3 \text{ and } u_{T-2}^*(x) = 0 \end{cases}$$

and again we notice that  $J_{T-2}(x)$  is a convex function which is continuous at  $x = 2/3$ .

Now at last, let us try to find the general expression  $J_{T-k}(x)$ . The formulae for  $J_{T-1}$  and  $J_{T-2}$  suggest that our best assumption is

$$J_{T-k}(x) = \begin{cases} a_k x + b_k & x \geq \frac{b_k}{1-a_k} = c \\ x & x < \frac{b_k}{1-a_k} = c \end{cases}$$

$k = 1, 2, \dots, T$  and that  $u_{T-k}^*(x) = 2$  if  $x \geq c$  and  $u_{T-k}^*(x) = 0$  if  $x < c$ .

The formulae is certainly correct in case of  $k = 1$ . Further, by using the same kind of considerations as in the computation of  $J_{T-2}(x)$  and induction there are two separate cases.

$$J_{T-(k+1)}(x) = \max_u [x - u + a_k u x + b_k] = \max_u [x + (a_k x - 1)u + b_k]$$

Hence  $x \geq 1/a_k \Rightarrow u = 2$  and  $x < 1/a_k \Rightarrow u = 0$ , and

$$J_{T-(k+1)}(x) = \max_u [x - u + ux] = \max_u [x + (x - 1)u]$$

Thus  $x \geq 1 \Rightarrow u = 2$  and  $x < 1 \Rightarrow u = 0$ .

This yields (just as in the  $J_{T-2}(x)$  case) the following

$$J_{T-(k+1)}(x) = \begin{cases} (2a_k + 1)x + b_k - 2 = a_{k+1}x + b_{k+1} = g_1(x) & x \geq 1/a_k \\ x + b_k = g_2(x) & x < 1/a_k \\ 3x - 2 = g_3(x) & x \geq 1 \\ x = g_4(x) & x < 1 \end{cases}$$

and we recognize that the forms of  $g_1(x)$  and  $g_4(x)$  are in accordance with our assumption and moreover that the point of intersection between  $g_1(x)$  and  $g_4(x)$  is  $b_k(1 - a_k)^{-1}$  which also is consistent with the assumption.

Further,  $a_k$  obeys the difference equation  $a_{k+1} = 2a_k + 1$ . Therefore, the general solution is  $a_k = D \cdot 2^k - 1$  and since  $a_1 = 3 \Rightarrow D = 2$  we have  $a_k = 2^{k+1} - 1$ . In the same way,  $b_{k+1} = b_k - 2$  (see the remark following this example, see also (1.1.2b)) has the general solution  $b_k = K - 2k$  and since  $b_1 = -2 \Rightarrow K = 0$  we obtain  $b_k = -2k$ .

Finally, since (1)  $g_1(1) \geq g_3(1)$  and  $a_{k+1} > 3$ . (2)  $g_4(x) > g_2(x)$  and (3)  $g_1(x) > g_4(x)$  when  $x > b_{k+1}(1 - a_{k+1})^{-1}$  (recall that  $a_{k+1} > 3$ ) we obtain the general solution


$$J_{T-k}(x) = \begin{cases} (2^{k+1} - 1)x - 2k & x \geq \frac{k}{2^{k-1}} & u_{T-k}^* = 2 \\ x & x < \frac{k}{2^{k-1}} & u_{T-k}^* = 0 \end{cases}$$

□

# Free eBook on Learning & Development

By the Chief Learning Officer of McKinsey

[Download Now](#)



[Click on the ad to read more](#)

**Remark 3.1.1:** Referring to section 2.1, Exercise 2.1.3, the difference equation  $x_{t+2} - 5x_{t+1} - 6x_t = t \cdot 2^t$  has the homogeneous solution  $C_1(-1)^t + C_26^t$  and since the exponential function  $2^t$  on the right-hand side of the equation is different from both exponential functions contained in the homogeneous solution it suffices to assume a particular solution of the form  $(At + B)2^t$  in this case. In Example 3.1.3 we had to solve an equation of the form  $x_{t+1} - x_t = at^2$ . The homogeneous solution is  $C \cdot 1^t = C$  but since  $at^2 = at^2 \cdot 1^t$  we have the same exponential function on both sides of the equation. Therefore, we must in this case assume a particular solution of the form  $(A + Bt + Dt^2)t$ . In the same way, if  $x_{t+1} - x_t = bt$  we assume a particular solution  $(A + Bt)t$  and finally, in the case  $x_{t+1} - x_t = K$ , assume a particular solution  $A + Bt$  (cf. (1.1.2b)). □

**Exercise 3.1.1.** Let  $a$  be a positive constant and solve the problem

$$\max_u \sum_{t=0}^T (x_t + u_t) \text{ subject to } x_{t+1} = x_t - au_t, \quad u_t \in [0, 1], \quad x_0 \text{ given}$$

□

**Exercise 3.1.2.** Solve the problem (Exam Exercise, UiO):

$$\max_u \sum_{t=0}^T (x_t - u_t) \text{ subject to } x_{t+1} = x_t + u_t, \quad u_t \in [0, 1], \quad x_0 \text{ given}$$

(Hint: Use Remark 3.1.1.) □

**Exercise 3.1.3.** Solve the problem:

$$\max_u \sum_{t=0}^T (x_t + 1) \text{ subject to } x_{t+1} = u_t x_t, \quad u_t \in [0, 1], \quad x_0 \text{ given}$$

□

### 3.2 The maximum principle (Discrete version)

When  $t$  is a continuous variable, most optimization problems are formulated and solved by use of the maximum principle which was developed by Russian mathematicians about 60 years ago. The maximum principle, sometimes referred to as Pontryagin's maximum principle, is the cornerstone in the discipline called optimal control theory which may be regarded as an extension of the classical calculus of variation. An excellent treatment of various aspects of control theory may be found in Seierstad and Sydsæter (1987), see also Sydsæter et al. (2005). In this section we shall briefly discuss a discrete version of the maximum principle which offers an alternative way of dealing with the kind of problems presented in section 3.1.

Consider the problem

$$\begin{aligned} & \text{maximize } \sum_{t=0}^T f_0(t, x_t, u_t), \quad u_t \in U, \quad U \text{ convex} \\ & \text{subject to } x_{t+1} = f(t, x_t, u_t), \quad t = 0, 1, \dots, T-1, \quad x_0 \text{ given.} \end{aligned} \quad (3.2.1)$$

together with one of the following terminal conditions

$$\text{a) } x_T \text{ free,} \quad \text{b) } x_T \geq X_T, \quad \text{c) } x_T = X_T \quad (3.2.2)$$

Thus, the problem that we consider here is somewhat more general than the one presented in section 3.1 due to the terminal conditions (3.2.2b,c).

Next, define the Hamiltonian by

$$H(t, x, u, p) = \begin{cases} f_0(t, x, u) + pf(t, x, u) & t < T \\ f_0(t, x, u) & t = T \end{cases} \quad (3.2.3)$$

where  $p$  is called the adjoint function.

Then we have the following:

**Theorem 3.2.1 (The maximum principle, discrete version).** Suppose that  $(x_t^*, u_t^*)$  is an optimal sequence for problem (3.2.1), (3.2.2). Then there are numbers  $p_0, \dots, p_T$  such that

$$u_t^* \text{ maximizes } H'_u(t, x_t^*, u_t^*, p_t)u \text{ for } u \in U \quad (3.2.4)$$

Moreover,

$$p_{t-1} = H'_x(t, x_t^*, u_t^*, p_t), \quad t = 1, \dots, T-1 \quad (3.2.5a)$$

$$p_{T-1} = f'_{0x}(T, x_T^*, u_T^*) + p_T \quad (3.2.5b)$$

and to each of the terminal conditions (3.2.2) we have the following transversal conditions

a)  $p_T = 0.$  (3.2.6a)

b)  $p_T \geq 0$  ( $= 0$  if  $x_T^* > X_T$ ). (3.2.6b)

c)  $p_T$  no condition. (3.2.6c)

□

Theorem 3.2.1 gives necessary conditions for optimality. Regarding sufficient conditions we have:

**Theorem 3.2.2.** Suppose that  $(x_t^*, u_t^*)$  satisfies all the conditions in Theorem 3.2.1 and in addition that  $H(t, x, u, p)$  is concave in  $(x, u)$  for every  $t$ . Then  $(x_t^*, u_t^*)$  is optimal. □

**Proof.** Our goal is to show that

$$K = \sum_{t=0}^T f_0(t, x_t^*, u_t^*) - \sum_{t=0}^T f_0(t, x_t, u_t) \geq 0$$

www.sylvania.com

We do not reinvent the wheel we reinvent light.

Fascinating lighting offers an infinite spectrum of possibilities: Innovative technologies and new markets provide both opportunities and challenges. An environment in which your expertise is in high demand. Enjoy the supportive working atmosphere within our global group and benefit from international career paths. Implement sustainable ideas in close cooperation with other specialists and contribute to influencing our future. Come and join us in reinventing light every day.

Light is OSRAM

OSRAM SYLVANIA



Click on the ad to read more



Introducing the notation  $f_0 = f_0(t, x, u)$ ,  $f_0^* = f_0(t, x^*, u^*)$  and so on, it follows from (3.2.3) that

$$K = \sum_{t=0}^T (H_t^* - H_t) + \sum_{t=0}^T p_t (f_t - f_t^*)$$

Now, since  $H$  is concave in  $(x, u)$  we also have that  $H - H^* \leq H'_x(x - x^*) + H'_u(u - u^*)$ . Thus

$$K \geq \sum_{t=0}^T H'_u(u_t^* - u_t) + \sum_{t=0}^T H'_x(x_t^* - x_t) + \sum_{t=0}^{T-1} p_t (f_t - f_t^*)$$

Due to (3.2.4) and the concavity of  $H$  the first of the three sums above are equal or larger than zero. Indeed, suppose  $u_t \in [u_0, u_1]$ . If  $u_t^* \in (u_0, u_1)$  then  $H'_u(u_t^*) = 0$ . If  $u_t^* = u_0$ , then  $H'_u(u_t^*) \leq 0$  and  $u_t^* - u_t \leq 0$  and finally, if  $u_t^* = u_1$ ,  $H'_u(u_t^*) \geq 0$  and  $u_t^* - u_t \geq 0$ , hence in all cases  $H'_u(u_t^* - u_t) \geq 0$ .

Regarding the second and the third sum they may by use of (3.2.5a), (3.2.5b) and (3.2.1) be written as

$$\begin{aligned} & \sum_{t=0}^{T-1} p_{t-1} (x_t^* - x_t) + (p_{T-1} - p_T)(x_T^* - x_T) + \sum_{t=0}^{T-1} p_t (x_{t+1} - x_{t+1}^*) \\ & = p_T (x_T - x_T^*) = K1 \end{aligned}$$

Next, assume  $x_T$  free. Then from (3.2.6a),  $p_T = 0$  which implies  $K1 = 0$ . If  $x_T \geq X_T$ , (3.2.6b) gives  $p_T \geq 0$  and since  $x_T \geq X_T$  we must have  $K1 \geq 0$  if  $x_T^* = X_T$ . If  $x_T^* > X_T$ ,  $p_T = 0$ , thus in either case  $K1 \geq 0$ . Finally, if  $x_T = X_T$ ,  $K1 = 0$ . Therefore, whatever terminal condition (3.2.2),  $K1 \geq 0$  which implies  $K \geq 0$  so we are done.  $\square$

**Example 3.2.1.** Solve the problem given in Example 3.1.2 by use of Theorems 3.2.1 and 3.2.2.

**Solution:** From (3.2.3) it follows

$$H(t, x, u, p) = \begin{cases} x + u + p(x - 2u) & t < T \\ x + u & t = T \end{cases}$$

Consequently, whenever  $t < T$ ,  $H'_x = 1 + p$  and  $H'_u = -2p$  and if  $t = T$ ,  $H'_x = H'_u = 1$ .

By use of the results above, (3.2.5a,b) gives

$$p_{t-1} = 1 + p_t \quad t < T, \quad p_{T-1} = 1 + p_T$$

and since  $x_T$  is free, (3.2.6a) implies that  $p_T = 0$  so  $p_{T-1} = 1$ .

The equation  $p_{t-1} = 1 + p_t$  may be rewritten as  $p_{t+1} - p_t = -1$  and its general solution is easily found to be  $p_t = C - t$ . Further, since  $p_{T-1} = 1$  it follows that  $1 = C - (T - 1)$ . Thus  $C = T$  so  $p_t = T - t$  and we observe that  $p_t > 0$  for every  $t < T$ .

From the preceding findings, (3.2.4) may be formulated as

$$\begin{aligned} u = u_t^* & \text{ shall maximize } -2(T-t)u & t < T \\ u = u_T^* & \text{ shall maximize } 1u & t = T \end{aligned}$$

Accordingly, we make the following choices: If  $t = T$ , choose  $u_T^* = 1$ . If  $t < T$  (recall that  $-2(T-t) < 0$ ), choose  $u_t^* = 0$  for every  $t$ . Hence, we have arrived at the same conclusion as we did in Example 3.1.2.

A final observation is that the Hamiltonian is linear in  $(x, u)$  so  $H$  is also concave in  $(x, u)$ . Consequently,  $(x_t^*, u_t^*)$  solves the problem ( $x_t^*$  is found at each  $t$  from the equation  $x_{t+1}^* = x_t^* - 2u_t^*$  and  $x_0$  is given).  $\square$

**Example 3.2.2.** Solve the problem

$$\text{maximize}_u \sum_{t=0}^T (x_t - u_t) \text{ subject to } x_{t+1} = x_t + u_t$$

$$x_0 = 1, \quad x_T = X_T, \quad 1 < X_T < T + 1, \quad u_t \in [0, 1].$$

**Solution:**

$$H(t, x, u, p) = \begin{cases} x - u + p(x + u) & t < T \\ x - u & t = T \end{cases}$$

Therefore, whenever  $t < T$ ,  $H'_x = 1 + p$ ,  $H'_u = -1 + p$  and if  $t = T$ ,  $H'_x = 1$  and  $H'_u = -1$ .

Further, (3.2.5b) gives  $p_{T-1} = 1 + p_T$  and (3.2.5a) gives  $p_{t-1} = 1 + p_t$  if  $t < T$ . Clearly (cf. our previous example), the latter difference equation has the general solution  $p_t = C - t$  so  $p_t$  is a decreasing sequence of points.

From (3.2.4) it follows

$$\begin{aligned} u = u_t^* & \text{ shall maximize } (-1 + p_t)u & t < T \\ u = u_T^* & \text{ shall maximize } -1u & t = T \end{aligned}$$

Thus at  $t = T$  the optimal control is  $u_T^* = 0$ . In the case  $t < T$  we have that if  $p_t - 1 \geq 0$ , then  $u = u_t^* = 1$  and if  $p_t - 1 < 0$ , we choose  $u_t^* = 0$ .

First, assume  $p_t - 1 \geq 0$  for all  $t < T$ . Then  $u_t^* = 1$  and  $x_{t+1}^* = x_t^* + 1$  which has the general solution  $x_t^* = K + t$ .  $x_0^* = 1 \Rightarrow K = 1$ , which means that  $x_t^* = t + 1$ . This implies that  $x_T^* = T + 1$  but this is a contradiction since  $X_T < T + 1$ . Next, assume  $p_t - 1 < 0$  for all  $t \leq T$ . Then  $u_t^* = 0$ . Thus,  $x_{t+1}^* = x_t^*$  which has the constant solution  $x_t^* = M$ . Again we have reached a contradiction since  $1 < X_T$ .



Discover the truth at [www.deloitte.ca/careers](http://www.deloitte.ca/careers)

**Deloitte.**

© Deloitte & Touche LLP and affiliated entities.



Click on the ad to read more

Finally, let us suppose that there exists a time  $t_c$  such that whenever  $t \leq t_c$ , then  $p_t - 1 \geq 0$  and in case of  $t_c < t \leq T$ ,  $p_t - 1 < 0$ .

First, consider the case  $t \leq t_c$ . Then  $x_{t+1}^* = x_t^* + 1$  so  $x_t^* = K + t$ .  $x_0 = 1 \Rightarrow K = 1$ , hence  $x_t^* = t + 1$ . If  $t > t_c$  we have  $x_{t+1}^* = x_t^*$ . Hence,  $x_t^*$  is a constant, say  $x_t^* = M$ , and since  $x_t^* = X_T$  it follows that  $x_t^* = X_T$ .

Thus,

$$\begin{aligned} t \leq t_c, & \quad p_t - 1 = C - t - 1 \geq 0 & \quad x_t^* = t + 1 & \quad u_t^* = 1 \\ t > t_c, & \quad p_t - 1 = C - t - 1 < 0 & \quad x_t^* = X_T & \quad u_t^* = 0 \end{aligned}$$

It remains to determine  $t_c$  and the constant  $C$ . At time  $t_c$ ,  $C - t_c - 1 = 0$  so  $C = t_c + 1$ . Therefore,  $p_t = t_c - t$ . Further, from  $x_{t_c+1} = x_{t_c} + u_{t_c}$  we obtain  $X_T = t_c + 1 + 1$  so  $t_c = X_T - 2$ . Consequently, by use of the conditions in the maximum principle we have arrived at

$$\begin{aligned} x_t^* = t + 1 & \quad u_t^* = 1 & \quad 0 \leq t \leq X_T - 2 \\ x_t^* = X_T & \quad u_t^* = 0 & \quad X_T - 2 < t \leq T \end{aligned}$$

and  $p_t = X_T - 2 - t$  for every  $t$ . Finally, since  $H$  is linear and concave in  $(x, u)$  it follows from Theorem 3.2.2 that we have obtained the solution. □

—

We close this section by looking at one extension only.

If we have a problem which involves several state variables  $x_1, \dots, x_n$  and several controls  $u_1, \dots, u_m$  we may organize them in vectors, say  $\mathbf{x} = (x_1, \dots, x_n)$ ,  $\mathbf{u} = (u_1, \dots, u_m)$  and reformulate problem (3.2.1), (3.2.2) as

$$\text{maximize } \sum_{t=0}^T f_0(t, \mathbf{x}_t, \mathbf{u}_t) \tag{3.2.7}$$

subject to  $\mathbf{x}_{t+1} = \mathbf{f}(t, \mathbf{x}_t, \mathbf{u}_t)$ ,  $\mathbf{x}_0$  given,  $\mathbf{u}_t \in U$ , and terminal conditions on the form

$$\text{a) } x_{i,T} \text{ free, } \quad \text{b) } x_{i,T} \geq X_{i,T}, \quad \text{c) } x_{i,T} = X_{i,T} \tag{3.2.8}$$

The associated Hamiltonian may in case of so-called “normal” problems be defined as

$$H(t, \mathbf{x}, \mathbf{u}, \mathbf{p}) = \begin{cases} f_0(t, \mathbf{x}, \mathbf{u}) + \sum_{i=1}^n p_i f_i(t, \mathbf{x}, \mathbf{u}) & t < T \\ f_0(t, \mathbf{x}, \mathbf{u}) & t = T \end{cases} \tag{3.2.9}$$

where  $\mathbf{p} = (p_1, \dots, p_n)$  is the adjoint function.

Then we may formulate necessary and sufficient conditions for an optimal solution in the same way as we did in the one-dimensional case.

**Theorem 3.2.3.** Suppose that  $(\mathbf{x}_t^*, \mathbf{u}_t^*)$  is an optimal sequence for problem (3.2.7), (3.2.8) with Hamiltonian defined as in (3.2.9). Then there exists  $\mathbf{p}$  such that

$$\mathbf{u} = \mathbf{u}_t^* \text{ maximizes } \sum_{i=1}^m \frac{\partial H}{\partial u_i}(t, \mathbf{x}_t^*, \mathbf{u}_t^*, \mathbf{p}_t) u_i \quad (3.2.10)$$

Moreover

$$p_{i,t-1} = H'_{x_i}(t, \mathbf{x}_t^*, \mathbf{u}_t^*, \mathbf{p}_t), \quad t = 1, \dots, T-1 \quad (3.2.11a)$$

$$p_{i,T-1} = \frac{\partial f_0}{\partial x_i}(T, \mathbf{x}_T^*, \mathbf{u}_T^*) + p_{i,T} \quad (3.2.11b)$$

and

a)  $p_{i,T} = 0$  if the terminal condition is (3.2.9a).

b)  $p_{i,T} \geq 0$  ( $= 0$  if  $x_{i,T}^* > X_{i,T}$ ) (3.2.12)

if the condition is (3.2.9b).

c)  $p_{i,T}$  free if condition (3.2.9c) applies.

Finally, if  $H$  is concave in  $(\mathbf{x}, \mathbf{u})$  for each  $t$  then  $(\mathbf{x}_t^*, \mathbf{u}_t^*)$  solves problem (3.2.7), (3.2.8).  $\square$

As usual, we end with an example.

**Example 3.2.3.** Solve the problem

$$\max \sum_{t=0}^T (-u_t^2 - 2x_t) \text{ subject to } x_{t+1} = \frac{1}{2}y_t, \quad y_{t+1} = u_t + y_t$$

$x_0 = 2, y_0 = 1, u_t \in \mathbb{R}, x_T$  free,  $y_T$  free.

**Solution:** Denoting the adjoint functions by  $p$  and  $q$  respectively, the Hamiltonian becomes

$$H(t, x, y, u, p, q) = \begin{cases} -u^2 - 2x + \frac{1}{2}py + q(u + y) & t < T \\ -u^2 - 2x & t = T \end{cases}$$

which implies

$$\begin{aligned} H'_x &= -2 & H'_y &= \frac{1}{2}p + q & H'_u &= -2u + q & t < T \\ H'_x &= -2 & H'_y &= 0 & H'_u &= -2u & t = T \end{aligned}$$

Then, from (3.2.11a) it follows that  $p_{t-1} = -2$ ,  $q_{t-1} = (1/2)p_t + q_t$  and since  $x_T, y_T$  is free, (3.2.12a) implies  $p_T = q_T = 0$ . Thus (3.2.11b) reduces to  $p_{T-1} = -2$  and  $q_{T-1} = 0$ .

Consequently,  $p_t = -2$  for each  $t$  and if we insert this result into the difference equation for  $q$  we easily obtain the general solution  $q_t = C + t$ . Moreover, since  $q_{T-1} = 0$  it follows that  $0 = C + T - 1$  so  $C = 1 - T$  which means that  $q_t = t - T + 1$ .


Now, since the control region is open, it follows from (3.2.10) that  $H'_u^* = 0$ , thus  $-2u_t^* + q_t = 0$  if  $t < T$  and  $2u_T^* = 0$  whenever  $t = T$ . Hence at time  $t = T$ ,  $u_T^* = 0$  and in case of  $t < T$ ,  $u_t^* = (1/2)q_t = 1/2(t - T + 1)$ .

SIMPLY CLEVER

ŠKODA



We will turn your CV into  
an opportunity of a lifetime



Do you like cars? Would you like to be a part of a successful brand?  
We will appreciate and reward both your enthusiasm and talent.  
Send us your CV. You will be surprised where it can take you.

Send us your CV on  
[www.employerforlife.com](http://www.employerforlife.com)



Click on the ad to read more

Therefore, the problem is in many respects already solved. Indeed,  $y_t^*$  is now uniquely determined from the relation  $y_{t+1}^* = u_t^* + y_t^*$  (recall that  $y_0 = 1$ ) and  $x_t^*$  is subsequently found from  $x_{t+1}^* = (1/2)y_t^*$ . We leave the details to the reader.

Finally, observe that the Hesse determinant of  $H$  ( $t < T$ ) may be written as

$$\begin{vmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -2 \end{vmatrix}$$

so clearly  $(-1)^1 \Delta_1 \geq 0$ ,  $(-1)^2 \Delta_2 = 0$ ,  $(-1)^3 \Delta_3 = 0$  where  $\Delta_i$  is all possible principal minors of order  $i$  respectively. Consequently  $H$  is concave in  $(x, y, u)$ . (At time  $t = T$  the result is clear.) □

**Exercise 3.2.1.** Solve Exercises 3.1.2 and 3.1.3 by use of the maximum principle. □

### 3.3 Infinite horizon problems

In the previous two sections we considered discrete dynamic optimization problems where the planning period  $T$  was finite. Our goal here is to study problems where  $T \rightarrow \infty$ . Such problems are called infinite horizon problems. Note that the extension from the finite to the infinite case is by no means straightforward. Indeed, since the sum we want to maximize now consists of an infinite number of terms we may obviously face convergence problems which were absent in sections 3.1 and 3.2.

There are mainly two different solution methods available (along with some numerical alternatives) when we deal with infinite horizon problems. The first method which we will describe is based upon Theorem 3.1.1 (The fundamental equation of discrete dynamic programming).

Consider the problem

$$\underset{u}{\text{maximize}} \sum_{t=0}^{\infty} \beta^t f_0(x_t, u_t) \tag{3.3.1}$$

subject to  $x_{t+1} = f(x_t, u_t)$ ,  $\beta \in (0, 1)$ ,  $x_0$  given,  $u_t \in U$ . Clearly (3.3.1) is an autonomous system and it serves in many respects as a “standard” problem in the infinite horizon case. Especially economists study systems like (3.3.1). Indeed, they often assume that  $\beta = 1/(1 + r)$  is a discount factor where  $r$  is the interest rate. Under this assumption, (3.3.1) may be interpreted as maximizing the present value of a quantity like a profit or a utility function  $f_0(x, u)$  subject to  $x_{t+1} = f(x_t, u_t)$  over all times regardless of any terminal conditions.



Now, returning to (3.3.1), in order to ensure convergence of the series, we impose the restriction:

$$K_1 \leq f_0(x, u) \leq K_2 \quad (3.3.2a)$$

where  $K_1$  and  $K_2$  are constants, or

$$f_0(x_t, u_t) \leq c \theta^t \quad (3.3.2b)$$

where  $\theta \in (0, \beta^{-1})$  and  $0 < c < \infty$ .

Next (compare with section 3.1), define the (optimal) value function at time  $t = s$  as

$$J_s(x) = \max_u \sum_{t=s}^{\infty} \beta^t f_0(x_t, u_t) = \beta^s J^s(x) \quad (3.3.3)$$

where

$$J^s(x) = \max_u \sum_{t=s}^{\infty} \beta^{t-s} f_0(x_t, u_t) \quad (3.3.4)$$

Denoting  $J_0(x) = J(x)$  we now have the following result:

**Theorem 3.3.1 (Bellman's equation).** Consider problem (3.3.1) under the restriction(s) (3.3.2). Then the (optimal) value function  $J_0(x) = J(x)$  defined through (3.3.3) satisfies

$$J(x) = \max_u [f_0(x, u) + \beta J(f(x, u))] \quad (3.3.5)$$

□

**Proof.** Since the horizon is infinite,  $J^{s+1}(x) = J^s(x)$ . Hence,

$$J_{s+1}(x) = \beta^{s+1}(x) J^{s+1}(x) = \beta \beta^s J^s(x) = \beta J_s(x)$$

Now, using the same argument as we did in the last paragraph before Theorem 3.3.1 was established it now follows:

$$\begin{aligned} J(x) = J_0(x) &= \max_u \left[ f_0(x_0, u_0) + \max_{u_1, \dots} \sum_{t=1}^{\infty} \beta^t f_0(x_t, u_t) \right] \\ &= \max_u [f_0(x_0, u_0) + J_1(x)] = \max_u [f_0(x_0, u_0) + J_1(f(x_0, u_0))] \\ &= \max_u [f_0(x_0, u_0) + \beta J_0(f(x_0, u_0))] \end{aligned}$$

□

**Remark 3.3.1.** Note the fundamental difference between equations (3.1.7), (3.1.8) in Theorem 3.1.1 and equation (3.3.5) in Theorem 3.3.1. (3.1.7) relates the value function  $J$  at different times  $T, T - 1, \dots$  and as we have demonstrated, the (finite) optimization problem could then be solved recursively. Regarding (3.3.5), this is not the case. Bellman's equation is a functional equation and there are no general solution methods for such equations. Therefore, often the best one can do is to "guess" the appropriate form of  $J(x)$  for a given problem. □

**Remark 3.3.2.** In the proof of Theorem 3.3.1 it is implicitly assumed that the maximum exists at each time step. This is not necessarily true but (3.3.5) still holds if we use the supremum notation instead of the max notation. □

Let us now by way of examples show how Theorem 3.3.1 applies.

**Example 3.3.1.** Solve the problem

$$\max_u \sum_{t=0}^{\infty} \beta^t \sqrt{x_t u_t}$$

Cynthia | AXA Graduate

**AXA Global Graduate Program**

Find out more and apply

redefining / standards AXA

 Click on the ad to read more

subject to  $x_{t+1} = (1 - u_t)x_t$ ,  $\beta \in (0, 1)$ ,  $u \in (0, 1)$ ,  $x_0$  given. □

**Solution.** First, consider  $f_0(x_t, u_t) = \sqrt{x_t}\sqrt{u_t}$ . Clearly,  $0 < \sqrt{u_t} < 1$  and since  $x_{t+1} < x_t$  it follows that  $0 < f_0(x_t, u_t) < x_0$ . Hence, (3.3.2a) is satisfied.

Next, from Theorem 3.3.1:

$$J(x) = \max_u [\sqrt{xu} + \beta J((1 - u)x)]$$

Assume that  $J(x) = \alpha\sqrt{x}$ .  $\alpha > 0$ . Then

$$\alpha\sqrt{x} = \max_u [\sqrt{x}\sqrt{u} + \alpha\beta\sqrt{1 - u}\sqrt{x}]$$

Thus

$$\alpha = \max_u [\sqrt{u} + \alpha\beta\sqrt{1 - u}] \tag{3.3.6}$$

Defining  $g(u) = \sqrt{u} + \alpha\beta\sqrt{1 - u}$ , the maximum of [ ] occurs when  $g'(u) = 0$ , i.e. when

$$u = \frac{1}{1 + (\alpha\beta)^2} \tag{3.3.7}$$

and by inserting into (3.3.6) we eventually arrive at

$$\alpha = \sqrt{(1 - \beta^2)^{-1}} \tag{3.3.8}$$

Finally, by substituting (3.3.8) back into (3.3.7) we obtain  $u = 1 - \beta^2$  so consequently the solution is

$$J(x) = \alpha\sqrt{x} = \sqrt{\frac{x}{1 - \beta^2}} \tag{3.3.9}$$

with associated optimal control  $u^* = 1 - \beta^2$ .

For comparison reasons let us also compute the maximum value of the infinite series in another way. From the constraint it follows that

$$x_{t+1}^* = (1 - u_t^*)x_t^* = \beta^2 x_t^*$$

Thus  $x_t^* = \beta^{2t}x_0$ . Consequently, the series becomes

$$\sum_{t=0}^{\infty} \beta^{2t} \sqrt{1-\beta^2} \sqrt{x_0} = \sqrt{(1-\beta^2)x_0} \sum_{t=0}^{\infty} \beta^{2t} = \sqrt{\frac{x_0}{1-\beta^2}}$$

in accordance with (3.3.9) ( $x_0 = x$ ). □

**Example 3.3.2.** Assuming  $J(x) = -\alpha x^2$ ,  $\alpha > 0$ , solve the problem:

$$\max_u \sum_0^{\infty} \beta^t (-x_t^2 - u_t^2)$$

subject to  $x_{t+1} = x_t + u_t$ ,  $\beta \in (0, 1)$ ,  $u \in (-\infty, \infty)$ ,  $x_0 > 0$  given. □

**Solution.** From Theorem 3.3.1

$$J(x) = \max_u [-x^2 - u^2 + \beta J(x + u)]$$

Thus (due to the assumption)

$$-\alpha x^2 = \max_u [-x^2 - u^2 - \alpha\beta(x + u)^2] = \max_u [-x^2 - u^2 - \alpha\beta x^2 - 2\alpha\beta x u - \alpha\beta u^2]$$

The function  $g(u) = -u^2 - 2\alpha\beta x u - \alpha\beta u^2$  is clearly concave in  $u$ , hence [ ] attains its maximum where  $g'(u) = 0$  which gives

$$u = -\frac{\alpha\beta x}{1 + \alpha\beta} \tag{3.3.10}$$

Consequently,

$$-\alpha x^2 = -x^2 - \frac{\alpha^2 \beta^2}{(1 + \alpha\beta)^2} x^2 - \alpha\beta x^2 + \frac{2\alpha^2 \beta^2}{1 + \alpha\beta} x^2 - \frac{\alpha^3 \beta^3}{(1 + \alpha\beta)^2} x^2$$

so after cancelling by  $x^2$  and rearranging we eventually arrive at

$$(1 + \alpha\beta) [-\beta\alpha^2 + (2\beta - 1)\alpha + 1] = 0 \tag{3.3.11}$$

Now, since  $\alpha > 0$ , the only acceptable solution of (3.3.11) is

$$\alpha = \frac{2\beta - 1}{2\beta} + \frac{\sqrt{1 + 4\beta^2}}{2\beta} \tag{3.3.12}$$

Hence,

$$J(x) = -\frac{1}{2\beta} \left[ 2\beta - 1 + \sqrt{1 + 4\beta^2} \right] x^2 \quad (3.3.13)$$

It is still no yet clear that we have solved the problem. We must check if (3.3.2a) is satisfied. Clearly,  $f_0(x_t, y_t) = -x_t^2 - y_t^2 \leq 0$  so if the sum shall be maximized it is natural to assume that  $|x_{t+1}| \leq |x_t|$ . Hence  $|x_t| \leq x_0$ . In the same way  $|u_{t+1}| \leq |x_t| \leq x_0$ . Under this assumption (3.3.13) will solve the problem.  $\square$

**Exercise 3.3.1.** Solve the problem

$$\max_u \sum_0^{\infty} \beta^t (-x_t - u_t)$$

subject to  $x_{t+1} = \frac{1}{2}x_t + \frac{1}{2}u_t$ ,  $\beta \in (0, 1)$ ,  $u \geq 0$ ,  $x_0$  given.  $\square$

**Exercise 3.3.2.** Consider the problem

I joined MITAS because  
I wanted **real responsibility**

The Graduate Programme  
for Engineers and Geoscientists  
[www.discovermitas.com](http://www.discovermitas.com)



**Month 16**

I was a construction supervisor in the North Sea advising and helping foremen solve problems

Real work  
International opportunities  
Three work placements







$$\max_u \sum_0^{\infty} \beta^t (-u_t^2 x_t)$$

subject to  $x_{t+1} = (1 - u_t)x_t$ ,  $\beta \in \langle 0, 1 \rangle$ ,  $u \in \langle -\infty, \infty \rangle$ ,  $x_0$  given.

a) Suppose that  $J(x) = \alpha x$  and use Bellman's equation to show that

$$J(x) = \frac{4(1 - \beta)}{\beta^2} x$$

with associated optimal control

$$u^* = \frac{2(\beta - 1)}{\beta}$$

b) Try to evaluate the sum of the series in the same way as we did at the end of Example 3.3.1 and conclude whether the found  $J(x)$  solves the problem or not. □

**Exercise 3.3.3.** Find  $J(x)$  and  $u_t^*$  for the problem

$$\max_u \sum_0^{\infty} \beta^t (-e^{-2x_t})$$

subject to  $x_{t+1} = x_t - 2u_t$ ,  $\beta \in (0, 1)$ ,  $u \in [-1, 1]$ ,  $x_0$  given. □

—

Our next goal is to show how infinite horizon problems may be solved by use of the maximum principle.

Consider the problem

$$\maximize_u \sum_{t=0}^{\infty} f_0(t, x_t, u_t) dt \tag{3.3.14}$$

subject to  $x_{t+1} = f(t, x_t, u_t)$ ,  $x_0$  given together with one of the following terminal conditions:

$$\lim_{T \rightarrow \infty} x(T) = \bar{x} \tag{3.3.15a}$$

$$\underline{\lim}_{T \rightarrow \infty} x(T) \geq \bar{x} \tag{3.3.15b}$$

$$\underline{\lim}_{T \rightarrow \infty} x(T) \text{ free} \tag{3.3.15c}$$

**Remark 3.3.3.** Recall the definition of  $\underline{\lim}_{t \rightarrow \infty}$  :

$$\underline{\lim}_{t \rightarrow \infty} f(t) = \lim_{t \rightarrow \infty} \inf \{f(s) | s \in [t, \rightarrow)\}$$

which means that  $\underline{\lim}_{t \rightarrow \infty} f(t) \geq a$  implies that for each  $\varepsilon > 0$  there exists a  $t'$  such that  $t > t'$  implies that  $f(t) \geq a - \varepsilon$ . □

**Remark 3.3.4.** Note that both  $f_0$  and  $f$  may depend explicitly on  $t$  in problem (3.3.14), (3.3.15) which is in contrast to the case covered by Bellman's equation. Also note the more general terminal conditions (3.3.15a,b,c). □

Let the Hamiltonian  $H$  be defined just as in section 3.2. Then we have the following:

**Theorem 3.3.2 (Maximum principle, infinite horizon).** Suppose that  $(\{x_t^*\}, \{u_t^*\})$  is an optimal sequence for problem (3.3.14), (3.3.15). Then there exist numbers  $p_t$  such that for  $t = 0, 1, 2, \dots$

$$H'_u(t, x_t^*, u_t^*, p_t)(u_t - u_t^*) \leq 0 \tag{3.3.16}$$

$$p_{t-1} = H'_x(t, x_t^*, u_t^*, p_t) \tag{3.3.17}$$

□

**Theorem 3.3.3.** Assume that all conditions in Theorem 3.3.2 are satisfied and moreover, that  $H(t, x, u, p)$  is concave in  $(x, u)$  for every  $t$  and that

$$\underline{\lim}_{t \rightarrow \infty} p_t(x_t - x_t^*) \geq 0 \tag{3.3.18}$$

Then  $(\{x_t\}, \{u_t\})$  is optimal. □

**Example 3.3.3.**

$$\max_u \sum_{t=0}^{\infty} \beta^t \sqrt{x_t u_t}$$

subject to  $x_{t+1} = (1 - u_t)x_t$ ,  $x_0$  given,  $\lim_{t \rightarrow \infty} x_t = \bar{x}$  where  $0 < \bar{x} < x_0$ ,  $\beta \in (0, 1)$  and  $u \in (0, 1)$ . □



**Solution.** Let  $H(t, x_t, u_t, p_t) = \beta^t \sqrt{x_t u_t} + p_t(1 - u_t)x_t$ . Since  $u \in (0, 1)$  is an interior point, (3.3.16) simplifies to  $H'_u(t, x_t^*, u_t^*, p_t) = 0$ , thus

$$\frac{1}{2} \beta^t \sqrt{\frac{x_t^*}{u_t^*}} = p_t x_t^* \quad (3.3.19)$$

Further, from (3.3.17) it follows that

$$\frac{1}{2} \beta^t \sqrt{\frac{u_t^*}{x_t^*}} = p_{t-1} - p_t(1 - u_t^*) \quad (3.3.20)$$

and through division

$$\frac{x_t^*}{u_t^*} = \frac{p_t x_t^*}{p_{t-1} - p_t(1 - u_t^*)}$$

which again implies that  $p_{t-1} - p_t = 0$ . Hence,  $p_t = K$  and clearly  $K > 0$  (cf. (3.3.19)).

Further from (3.3.19)

$$u_t^* = \frac{1}{4K^2 x_t^*} \beta^{2t}$$

**ie** business school

**93%**  
OF MIM STUDENTS ARE  
WORKING IN THEIR SECTOR 3 MONTHS  
FOLLOWING GRADUATION

**MASTER IN MANAGEMENT**

- STUDY IN THE CENTER OF MADRID AND TAKE ADVANTAGE OF THE UNIQUE OPPORTUNITIES THAT THE CAPITAL OF SPAIN OFFERS
- PROPEL YOUR EDUCATION BY EARNING A DOUBLE DEGREE THAT BEST SUITS YOUR PROFESSIONAL GOALS
- STUDY A SEMESTER ABROAD AND BECOME A GLOBAL CITIZEN WITH THE BEYOND BORDERS EXPERIENCE

Length: 10 MONTHS  
Av. Experience: 1 YEAR  
Language: ENGLISH / SPANISH  
Format: FULL-TIME  
Intakes: SEPT / FEB

**5 SPECIALIZATIONS**  
PERSONALIZE YOUR PROGRAM

**#10 WORLDWIDE**  
MASTER IN MANAGEMENT  
FINANCIAL TIMES

**55 NATIONALITIES**  
IN CLASS

[www.ie.edu/master-management](http://www.ie.edu/master-management) | [mim.admissions@ie.edu](mailto:mim.admissions@ie.edu) | [f](#) [t](#) [y](#) Follow us on IE MIM Experience



Thus

$$x_{t+1}^* = \left(1 - \frac{1}{4K^2 x_t^*} \beta^{2t}\right) x_t^*$$

which gives

$$x_{t+1}^* - x_t^* = -\frac{1}{4K^2} \beta^{2t}$$

The general solution becomes

$$x_t^* = C - \frac{1}{4K^2(\beta^2 - 1)} \beta^{2t}$$

and moreover (since  $x_0$  is given)

$$x_t^* = x_0 - \frac{1}{4K^2(1 - \beta^2)} (1 - \beta^{2t})$$

Finally, from the terminal condition  $\lim_{T \rightarrow \infty} x_T = \bar{x}$  it follows that

$$x_0 - \frac{1}{4K^2(1 - \beta^2)} = \bar{x}$$

so

$$K^2 = \frac{1}{4(x_0 - \bar{x})(1 - \beta^2)}$$

Consequently,

$$x_t^* = \bar{x} + (x_0 - \bar{x})\beta^{2t} \quad u_t^* = \frac{(x_0 - \bar{x})(1 - \beta^2)\beta^{2t}}{\bar{x} + (x_0 - \bar{x})\beta^{2t}}$$

Note that if we substitute these solutions back into the original series we obtain

$$\sum_{t=0}^{\infty} \beta^t \sqrt{(x_0 - \bar{x})(1 - \beta^2)} \beta^t = \sqrt{x_0 - \bar{x}} \sqrt{1 - \beta^2} \sum_{t=0}^{\infty} \beta^{2t} = \sqrt{\frac{x_0 - \bar{x}}{1 - \beta^2}}$$

(This example should be compared with Example 3.3.1.)  $\square$

**Example 3.3.4.**

$$\max_u \sum_{t=0}^{\infty} \beta^t (-e^{2x_t})$$

subject to  $x_{t+1} = x_t + 2u_t$ ,  $x_0$  given,  $\lim_{T \rightarrow \infty} x_T$  free,  $u \in [0, 1]$ ,  $\beta \in (0, 1)$ . □

**Solution.** The Hamiltonian becomes  $H = -\beta^t e^{-2x_t} + p_t(x_t + 2u_t)$  and evidently  $H$  is concave in  $(x, u)$ . Moreover, (3.3.16), (3.3.17) may be expressed as

$$2p_t(u_t - u_t^*) \leq 0 \tag{3.3.21}$$

and

$$p_{t-1} = 2\beta^t e^{-2x_t} + p_t \tag{3.3.22}$$

Consequently (from (3.3.21)) we conclude that  $u_t^* = 1$  whenever  $p_t \geq 0$  and  $u_t^* = 0$  if  $p_t < 0$ .

First, suppose  $u_t^* = 1$ . Then  $x_{t+1}^* = x_t^* + 2$ . Thus  $x_t^* = C + 2t$  and the corresponding  $p_t$  may be obtained from (3.3.22) as

$$p_t = K + \frac{2\beta e^{-2(C+2)}}{1 - \beta e^{-4}} (\beta e^{-4})^t \tag{3.3.23}$$

and we observe that  $p_t$  is a decreasing sequence of points.

Next, assume  $u_t^* = 0$ . Then  $x_{t+1}^* = x_t^* \Rightarrow x_t^* = M$  and (3.3.22) implies that

$$p_t = W + \frac{2\beta e^{-2M}}{1 - \beta} \beta^t \tag{3.3.24}$$

and again we recognize that  $p_t$  is a decreasing sequence.

We are now left with three possibilities: (A)  $u_t^* = 1$  for every  $t$ , (B)  $u_t^* = 0$  for every  $t$ , or (C) there exists  $t = t^*$  such that  $u_t^*$  takes the value 1 (or 0) if  $t < t^*$  and the value 0 (or 1) if  $t \geq t^*$ .

Suppose (A). Then  $u_t^* = 1$ ,  $x_t^* = x_0 + 2t$  and (3.3.23) may be expressed as

$$p_t = K + \frac{2\beta e^{-2(x_0+2)}}{1 - \beta e^{-4}} (\beta e^{-4})^t$$

Further, since  $\lim_{T \rightarrow \infty} x_T$  is free, (3.3.18) implies that  $\lim_{T \rightarrow \infty} p_T = 0$ . Thus  $K = 0$  so clearly  $p_t > 0$ . Hence, possibility (A) satisfies both Theorem 3.3.2 and Theorem 3.3.3.

Next, consider (B). Then  $u_t^* = 0$ ,  $x_t^* = x_0$  and (3.3.24) becomes

$$p_t = W + \frac{2\beta e^{-2x_0}}{1 - \beta} \beta^t$$

and just as in the treatment of (A), (3.3.18) implies that  $W = 0$ , hence  $p_t > 0$  which contradicts (3.3.21).

Finally, assume (C), i.e. that there exists a  $t = t^*$  such that for  $t = 0, 1, \dots, t^* - 1$  we have  $u_t^* = 0$ ,  $x_t^* = x_0$  and for  $t = t^*, t^* + 1, \dots$  we have  $u_t^* = 1$ ,  $x_t^* = C + 2t$ . The relation  $x_{t^*}^* = x_{t^*-1}^* + u_{t^*-1}^*$  now implies  $C + 2t^* = x_0 + 0$ . Thus  $C = x_0 - 2t^*$ . But then (from (3.3.22))

$$p_{t^*-1} = 2\beta^{t^*} e^{-2x_{t^*}^*} + p_{t^*}^* > 0$$

"I studied English for 16 years but...  
...I finally learned to speak it in just six lessons"  
Jane, Chinese architect

ENGLISH OUT THERE

Click to hear me talking before and after my unique course download



(recall that  $p_t^* > 0$ ) which contradicts  $u_t^* = 0$ .

Consequently,  $x_t^* = x_0 + 2t$ ,  $u_t^* = 1$  and

$$p_t = \frac{2\beta e^{-2(x_0+2t)}}{1 - \beta e^{-4t}} (\beta e^{-4t})^t$$

solves the problem. The maximum value becomes

$$\sum_0^{\infty} \beta^t (-e^{-2(x_0+2t)}) = -e^{-2x_0} \sum_0^{\infty} (\beta e^{-4t})^t = -\frac{e^{-2x_0}}{1 - \beta e^{-4}}$$

□

### 3.4 Discrete stochastic optimization problems

In sections 3.1–3.3 we discussed various aspects of discrete deterministic optimization problems. The theme in this section is to include stochasticity in such problems, so, instead of assuming a relation of the form  $x_{t+1} = f(t, x_t, u_t)$  between the deterministic state variable  $x$  and the control  $u$  (cf. (3.1.1), see also Theorem 3.1.1), we shall from now on suppose that

$$X_{t+1} = f(t, X_t, u_t, V_{t+1}) \quad X_0 = x_0, \quad V_0 = v_0 \quad (3.4.1)$$

where  $x_0$  and  $v_0$  are given, and  $u_t \in U$ .

The use of capital letters  $X$  and  $V$  indicates that they are stochastic variables. Indeed,  $X$  will in general depend on the values of  $V$ .  $V$  is a random variable which may be interpreted as environmental noise or some other kind of disturbances. Regarding  $V$ , we may in some cases know the distribution of  $V$  explicitly, for example that  $V_{t+1}$ , is identically normal distributed with expected value  $E(V_{t+1}) = \mu$ . Alternatively, we may know the probability  $P(V_{t+1} = v)$ , for example  $P(V_{t+1} = 1) = p$ , and a third possibility is that we have a knowledge of the conditional probability  $P(V_{t+1} | V_t)$ . (Later, when we turn to examples, all cases mentioned above will be considered.) A final comment is that the control  $u$  may depend on both  $X$  and  $V$ , thus  $u_t = u_t(X_t, V_t)$  and from now on we shall refer to  $u_t$  as a Markov control. We further assume that we actually can observe the value of  $X_t$  before we choose  $u_t$ . (If we have to choose  $u_t$  before observing the value of  $X_t$ , that may lead to a different value of the optimal Markov control.)

Now, referring to section 3.1, in the deterministic case we studied optimization problems of the form

$$\max_u \sum_{t=0}^T f_0(t, x_t, u_t)$$

subject to  $x_{t+1} = f(t, x_t, u_t)$  where  $u_t \in U$ ,  $x_0$  given. In the stochastic approach which we consider here, it does not make sense to maximize  $f_0$  at each time  $t$ , so instead we have to maximize the expected value of  $f_0$  at each time. Consequently, we study the problem

$$\text{maximize}_{u_0, u_1, \dots, u_T} E \left( \sum_{t=0}^T f_0(t, X_t, u_t, V_t) \right) \quad (3.4.2)$$

subject to  $X_{t+1} = f(t, X_t, u_t, V_{t+1})$  where  $X_0 = x_0$ ,  $V_0 = v_0$  and  $u_t \in U$ .

Define

$$J_s(t, x_t, v_t) = \max_u E \left( \sum_{t=s}^T f_0(s, X_s, u_s(X_s, V_s) \mid x_t, v_t) \right) \quad (3.4.3)$$

Then, somewhat roughly, we have by the same argument that eventually lead to Theorem 3.1.1 the following:

**Theorem 3.4.1.** Let  $J_s(t, x_t, v_t)$  defined through (3.4.3) be the value function for problem (3.4.2).

Then

$$\begin{aligned} J(t-1, x_{t-1}, v_{t-1}) &= \max_{u_{t-1}} \{f_0(t-1, x_{t-1}, u_{t-1}) + E[J(t, X_t, V_t)]\} \\ &= \max_{u_{t-1}} \{f_0(t-1, x_{t-1}, u_{t-1}) \\ &\quad + E[J(t, f(t-1, x_{t-1}, u_{t-1}, V_t), V_t)]\} \end{aligned} \quad (3.4.4a)$$

and

$$J(T, x_T, v_T) = J(T, x_T) = \max_{u_T} f_0(T, x_T, u_T) \quad (3.4.4b)$$

□

**Remark 3.4.1.** Note that Theorem 3.4.1 works backwards in the same way as Theorem 3.1.1. First, we find the optimal Markov control  $u_T^*(x_T, v_T)$  and the associated value function  $J(T, x_T)$  from (3.4.4b). Then, through (3.4.4a) the Markov controls and corresponding optimal value functions at times  $T - 1, T - 2, \dots$  are found recursively.  $\square$

**Example 3.4.1.** Solve the problem

$$\max_u E \left( \sum_{t=0}^T (u_t + X_t) \right)$$

subject to  $X_{t+1} = X_t - 2u_t + V_{t+1}$ , where  $u_t \in [0, 1]$ ,  $x_0$  given, and  $V_{t+1} \geq 0$  is Rayleigh distributed with probability density  $h(v) = (v/\theta^2) \exp[-v^2/2\theta^2]$  and  $\theta > 0$ .

**Solution:** From (3.4.4b):  $J(T, x_T) = \max_u (x + u)$  so obviously we choose  $u = 1$ . Hence:  $J(T, x_T) = x_T + 1$  and  $u_T^* = 1$ .

Excellent Economics and Business programmes at:



university of  
 groningen




“The perfect start  
of a successful,  
international career.”

**CLICK HERE**  
to discover why both socially  
and academically the University  
of Groningen is one of the best  
places for a student to be

[www.rug.nl/feb/education](http://www.rug.nl/feb/education)





Now, using the fact that  $E(V_{t+1}) = \theta\sqrt{\pi/2} = K$  it follows from (3.4.4a):

$$\begin{aligned} J(T-1, x) &= \max_u \{x + u + E(X_t + 1)\} \\ &= \max_u \{x + u + x - 2u + K + 1\} = \max_u \{2x - u + K + 1\} \end{aligned}$$

so clearly, the optimal Markov control is 0 which implies

$$J(T-1, x_{T-1}) = 2x_{T-1} + 1 + K \quad \text{and} \quad u_{T-1}^* = 0$$

Proceeding in the same way, (3.4.4a) gives

$$\begin{aligned} J(T-2, x) &= \max_u \{u + x + E(2X_T + K + 1)\} \\ &= \max_u \{u + x + 2(x - 2u + K) + K + 1\} \\ &= \max_u \{3x - 3u + 3K + 1\} \end{aligned}$$

Again, the optimal Markov control is  $u = 0$ , so consequently:

$$J(T-2, x_{T-2}) = 3x_{T-2} + 3K + 1 \quad u_{T-2}^* = 0$$

From the findings above it is natural to suspect that in general:

$$J(T-k, x) = (k+1)x + \alpha_k K + 1 \quad \alpha_0 = 0$$

The formulae is certainly correct in case of  $k = 0$  and by induction

$$J(T-k, x) = (k+1)x + \alpha_k K + 1 \quad \alpha_0 = 0$$

Clearly, the optimal Markov control is  $u = 0$  so

$$J(T-(k+1), x) = (k+2)x + (\alpha_k + k + 1)K + 1 = (k+2)x + \alpha_{k+1}K + 1$$

which proves what we want.  $\alpha_k$  obeys the difference equation  $\alpha_{k+1} - \alpha_k = k + 1$ . The homogeneous solution is  $C$ , and by assuming a particular solution of the form  $(Ak + B)k$  together with the fact that  $\alpha_0 = 0$  it follows that  $\alpha_k = (k/2)(k + 1)$ . Thus

$$J(T, x_T) = x_T + 1 \quad u_T^* = 1$$

$$J(T-k, x_{T-k}) = (k+1)x_{T-k} + \frac{1}{2}(k^2 + k)K + 1 \quad u_{T-k}^* = 0, \quad k \geq 1$$

or alternatively:

$$J(T, x) = x + 1 \quad u_T^* = 1$$

$$J(t, x) = (T - t + 1)x + \frac{1}{2}(T - t)(T - t + 1)K + 1 \quad u_t^* = 0, \quad t < T$$

□

**Example 3.4.2.** Solve the problem:

$$\max_u E \left( \sum_{t=0}^{T-1} -u_t^2 - X_T^2 \right)$$

subject to  $X_{t+1} = (X_t + u_t)V_{t+1}$ ,  $V_{t+1} \in \{0, 1\}$ ,  $P(V_{t+1} = 1) = \frac{1}{2}$ ,  $P(V_{t+1} = 0) = \frac{1}{2}$ ,  $x_t > 0$ ,  $x_0$  given and  $u \in \mathbb{R}$ . (Note that an alternative way of expressing the probabilities above is to say that  $X_{t+1} = X_t + u_t$  with probability 1/2 and  $X_{t+1} = 0$  with probability 1/2.)

**Solution:**

$$J(T, x_T) = \max_u (-x_T^2) = -x_T^2 \quad u_T^* \text{ arbitrary}$$

$$\begin{aligned} J(T-1, x) &= \max_u \left\{ -u^2 + E(-X_T^2) \right\} = \max_u \left\{ -u^2 - (x+u)^2 \cdot \frac{1}{2} + O^2 \cdot \frac{1}{2} \right\} \\ &= \max_u \left\{ -u^2 - \frac{1}{2}(x+u)^2 \right\} \end{aligned}$$

Denoting  $g_1(u) = -u^2 - (1/2)(x+u)^2$ , the equation  $g_1'(u) = 0$  gives  $u = -(1/3)x$ .

(Note that  $g_1$  is concave.) Thus

$$J(T-1, x) = - \left( -\frac{1}{3}x \right)^2 - \frac{1}{2} \left( \frac{2}{3}x \right)^2 = -\frac{1}{3}x^2 \quad \text{and} \quad u_{T-1}^* = -\frac{1}{3}x$$

In the same way:

$$\begin{aligned} J(T-2, x) &= \max_u \left\{ -u^2 + E \left( -\frac{1}{3}X_{T-1}^2 \right) \right\} \\ &= \max_u \left\{ -u^2 - \frac{1}{3} \left[ (x+u)^2 \cdot \frac{1}{2} + O^2 \cdot \frac{1}{2} \right] \right\} = \max_u \left\{ -u^2 - \frac{1}{6}(x+u)^2 \right\} \end{aligned}$$

Letting  $g_2(u) = -u^2 - (1/6)(x + u)^2$ , we easily obtain the solution of  $g_2'(u) = 0$  as  $u = -(1/7)x$ , hence

$$J(T - 2, x) = - \left(-\frac{1}{7}x\right)^2 - \frac{1}{6} \left(x - \frac{1}{7}x\right)^2 = -\frac{1}{7}x^2 \quad u_{T-2}^* = -\frac{1}{7}x$$

Now, assume that  $J(T - k, x) = -\alpha_k x^2$  where  $\alpha_0 = 1$ . Then:

$$\begin{aligned} J(T - (k + 1), x) &= \max_u \{ -u^2 + E(-\alpha_k X_{T-k}^2) \} \\ &= \max_u \left\{ -u^2 - \alpha_k \left[ (x + u)^2 \cdot \frac{1}{2} + O^2 \cdot \frac{1}{2} \right] \right\} \\ &= \max_u \left\{ -u^2 - \frac{1}{2} \alpha_k (x + u)^2 \right\} \end{aligned}$$

## American online LIGS University

is currently enrolling in the  
Interactive Online **BBA, MBA, MSc,**  
**DBA and PhD** programs:

- ▶ enroll **by September 30th, 2014** and
- ▶ **save up to 16%** on the tuition!
- ▶ pay in 10 installments / 2 years
- ▶ Interactive **Online education**
- ▶ visit [www.ligsuniversity.com](http://www.ligsuniversity.com) to find out more!

**Note: LIGS University is not accredited by any nationally recognized accrediting agency listed by the US Secretary of Education. More info [here](#).**





The equation  $g'(u) = 0$  (where  $g(u)$  is the concave function inside the  $\{ \}$  bracket) has the solution  $u = -\alpha_k(2 + \alpha_k)^{-1}x$ . Thus,

$$\begin{aligned} J(T - (k + 1), x) &= - \left( -\frac{\alpha_k}{2 + \alpha_k}x \right)^2 - \frac{1}{2}\alpha_k \left( x - \frac{\alpha_k}{2 + \alpha_k}x \right)^2 \\ &= -\frac{\alpha_k}{2 + \alpha_k}x^2 = -\alpha_{k+1}x^2 \end{aligned}$$

which is in accordance with the assumption. Consequently,

$$J(T, x_T) = -x_T^2 \quad u_T^* \text{ arbitrary}$$

$$J(T - k, x_{T-k}) = -\alpha_k x_{T-k}^2 \quad u_{T-k}^* = -\frac{\alpha_k}{2 + \alpha_k}x \quad k \geq 1$$

where

$$\alpha_{k+1} = \frac{\alpha_k}{2 + \alpha_k}$$

□

In the previous examples we have considered the cases that  $V_{t+1}$  is from a known distribution (Example 3.4.1) and  $P(V_{t+1} = v)$  is known (Example 3.4.2). In the next example we present the solution of a problem found in Sydsæter et al. (2005), which incorporates conditional probabilities.

**Example 3.4.3.** Solve the problem

$$\max E \left( \sum_{t=0}^{T-1} -u_t^2 - X_T^2 \right)$$

subject to  $X_{t+1} = X_t V_{t+1} + u_t$ ,  $x_0 > 0$  given,  $u_t \in \mathbb{R}$ ,  $V_{t+1} \in \{0, 1\}$ ,

$$P(V_{t+1} = 1 \mid V_t = 1) = \frac{3}{4}, \quad P(V_{t+1} = 1 \mid V_t = 0) = \frac{1}{4}.$$

**Solution:** First, note that the conditional probabilities above also imply  $P(V_{t+1} = 0 \mid V_t = 1) = 1/4$  and  $P(V_{t+1} = 0 \mid V_t = 0) = 3/4$ . Clearly:

$$J(T, x_T) = \max_u (-x_T^2) = -x_T^2 \quad u_T^* \text{ arbitrary}$$

Regarding  $J(T-1, x_{T-1}, v_{T-1})$  there are two cases to consider, the case  $v_{T-1} = 1$  and the case  $v_{T-1} = 0$ . The former yields:

$$\begin{aligned} J(T-1, x, 1) &= \max_u \{-u^2 + E(-x_T^2)\} \\ &= \max_u \left\{ -u^2 - \frac{3}{4}(x \cdot 1 + u)^2 - \frac{1}{4}(x \cdot 0 + u)^2 \right\} \\ &= \max_u \left\{ -\frac{5}{4}u^2 - \frac{3}{4}(x + u)^2 \right\} \end{aligned}$$

Defining  $g_1(u) = -(5/4)u^2 - (3/4)(x + u)^2$ , the solution of  $g'_1(u) = 0$  is  $u = -(3/8)x$  which after some algebra gives

$$J(T-1, x_{T-1}, 1) = -\frac{15}{32}x_{T-1}^2 \quad u_{T-1}^* = -\frac{3}{8}x_{T-1}$$

In the same way, the latter yields

$$J(T-1, x_{T-1}, 0) = -\frac{7}{32}x_{T-1}^2 \quad u_{T-1}^* = -\frac{1}{8}x_{T-1}$$

Now, assume:

$$J(T-k, x, 1) = -\alpha_k x^2 \quad J(T-k, x, 0) = -\beta_k x^2 \tag{3.4.5}$$

Then, by induction:

$$\begin{aligned} J(T-(k+1), x, 1) &= \max_u \left\{ -u^2 - \alpha_k \left[ (x \cdot 1 + u)^2 \cdot \frac{3}{4} \right] - \beta_k \left[ (x \cdot 0 + u)^2 \cdot \frac{1}{4} \right] \right\} \\ &= \max_u \left\{ -u^2 - \frac{3}{4}\alpha_k(x + u)^2 - \frac{1}{4}\beta_k u^2 \right\} \end{aligned}$$

Letting  $g(u) = -u^2 - (3/4)\alpha_k(x + u)^2 - (1/4)\beta_k u^2$ , the equation  $g'(u) = 0$  implies  $u = -3\alpha_k(3\alpha_k + \beta_k + 4)^{-1}x$ . Substituting back into  $J(T-(k+1), x, 1)$  then gives after some algebra

$$J(T-(k+1), x, 1) = -\frac{3}{4} \frac{\alpha_k}{3\alpha_k + \beta_k + 4} (\beta_k + 4)x^2 = -\alpha_{k+1}x^2$$

and

$$u_{T-(k+1)}^*(1) = -\frac{3\alpha_k}{3\alpha_k + \beta_k + 4}x_{T-(k+1)}$$

By applying the same technique as above:

$$\begin{aligned} J(T - (k + 1), x, 0) &= \max_u \left\{ -u^2 - \frac{1}{4}\alpha_k(x \cdot 1 + u)^2 - \frac{3}{4}\beta_k(x \cdot 0 + u)^2 \right\} \\ &= \max_u \left\{ -u^2 - \frac{1}{4}\alpha_k(x + u)^2 - \frac{3}{4}\beta_k u^2 \right\} \end{aligned}$$

and we easily conclude that  $u = -\alpha_k(\alpha_k + 3\beta_k + 4)^{-1}x$  is the optimal Markov control.

Inserting back into  $J(T - (k + 1), x, 0)$  gives

$$J(T - (k + 1), x, 0) = -\frac{\alpha_k(3\beta_k + 4)}{4(\alpha_k + 3\beta_k + 4)}x^2 = -\beta_{k+1}x^2$$

and

$$u_{T-(k+1)}^*(0) = -\frac{\alpha_k}{\alpha_k + 3\beta_k + 4}x_{T-(k+1)}$$

**DON'T EAT YELLOW SNOW**

What will your advice be?

Some advice just states the obvious. But to give the kind of advice that's going to make a real difference to your clients you've got to listen critically, dig beneath the surface, challenge assumptions and be credible and confident enough to make suggestions right from day one. At Grant Thornton you've got to be ready to kick start a career right at the heart of business.

Sound like you? Here's our advice: visit [GrantThornton.ca/careers/students](http://GrantThornton.ca/careers/students)

Scan here to learn more about a career with Grant Thornton.

**Grant Thornton**  
An instinct for growth™

© Grant Thornton LLP. A Canadian Member of Grant Thornton International Ltd



Finally, since  $\alpha_0 = \beta_0 = 1$ , we may by iteration find  $\alpha_k, \beta_k$  for any  $k < T$  through the equations

$$\alpha_{k+1} = \frac{3\alpha_k(\beta_k + 4)}{4(3\alpha_k + \beta_k + 4)} \quad \beta_{k+1} = \frac{\alpha_k(3\beta_k + 4)}{4(\alpha_k + 3\beta_k + 4)}$$

so the solution is given by (3.4.5) and associated optimal Markov controls. □

**Exercise 3.4.1.** Solve the problem

$$\max_u E \left( \sum_{t=0}^T X_t \right)$$

subject to  $X_{t+1} = u_t X_t V_{t+1}$ , where  $u_t \in [0, 1]$ ,  $x_0$  given,  $P(V_{t+1} = 1) = 1/3$ ,  
 $P(V_{t+1} = 0) = 2/3$ . □

**Exercise 3.4.2.** Solve the problem

$$\max_u E \left( \sum_{t=0}^T \beta^t (-u_t^2 - X_t^2) \right)$$

subject to  $X_{t+1} = X_t + u_t + V_{t+1}$ ,  $\beta \in \langle 0, 1 \rangle$ ,  $u_t \in \mathbb{R}$ ,  $x_0$  given,  $V_{t+1}$  is normal distributed  
where  $E(V_{t+1}) = \mu = 0$  and  $\text{Var}(V_{t+1}) = \sigma^2 > 0$ .

Hint: referring to Remark 3.4.3,  $E(V_{t+1}^2) = v$ . □

**Exercise 3.4.3.** Solve the problem

$$\max_u E \left( \sum_{t=0}^T (X_t - u_t) \right)$$

subject to  $X_{t+1} = X_t + u_t + V_{t+1}$ ,  $u_t \in [0, 1]$ ,  $x_0$  given,  $V_{t+1} \geq 0$  is exponential distributed  
and  $E(V_{t+1}) = 1/\lambda$ ,  $\lambda > 0$  for all  $t$ . □

**Exercise 3.4.4.** Show that the solution of the problem

$$\max_u E \left( \sum_{t=0}^T X_t \right)$$



subject to  $X_{t+1} = u_t X_t V_{t+1}$ ,  $u_t \in [0, 1]$ ,  $V_{t+1} \in \{0, 1\}$ ,  $P(V_{t+1} = 1) \mid V_t = 1) = 2/3$ ,  
 $P(V_{t+1} = 1 \mid V_t = 0) = 1/3$  may be written as

$$J(t, x, 1) = \left( -2 \left( \frac{2}{3} \right)^{T-t} + 3 \right) x \quad J(t, x, 0) = \left( -\frac{1}{2} \left( \frac{1}{3} \right)^{T-t} + \frac{3}{2} \right) x$$

□

—

Next, let us briefly comment on the case  $T \rightarrow \infty$ , i.e. the infinite horizon case. As explained in section 3.3, the extension from  $T$  finite to  $T$  infinite is by no means straightforward (mainly due to convergence problems). Therefore, adopting the same strategy as in section 3.3 we now restrict the analysis to the autonomous problem

$$\max_u E \left( \sum_{t=0}^{\infty} \beta^t f_0(X_t, u_t(X_t, V_t)) \right) \quad (3.4.6)$$

subject to  $X_{t+1} = f(X_t, u_t(X_t, V_t), V_{t+1})$ ,  $x_0$  given.  $\beta \in (0, 1)$ ,  $u_t \in \mathbb{R}$  and where all probabilities  $P(V_{t+1} = v)$  are time independent. Moreover, cf. (3.3.2a), we also impose the boundedness condition  $K_1 \leq f_0(x, u) \leq K_2$ .

Now, define (se Remark 3.3.2)

$$J(s, x_s, v_s) = \sup E \left( \sum_{t=s}^{\infty} \beta^t f_0(X_t, u(X_t, V_t)) \right) \quad (3.4.7)$$

Then, (roughly) by using the same kind of arguments that lead to Theorem 3.3.1 we may formulate the stochastic version of the Bellman equation as:

**Theorem 3.4.2.** Consider problem (3.4.6) and let  $J(s, x_s, v_s)$  be defined through (3.4.7). Then

$$J(x, v) = \max_u \{f_0(x, u) + \beta E(J(X_1, V_1))\} \quad (3.4.8)$$

where  $J(x, v) = J(t = 0, x, v)$  and  $X_1 = f(X, u, V_1)$ . □

**Remark 3.4.2.** Just as in section 3.3, note the fundamental difference between (3.4.4a,b) and (3.4.8). The latter is a functional equation which may not be solved recursively. Therefore, often the best we can do is to “guess” the appropriate form of  $J(x, v)$  in (3.4.8). □

**Remark 3.4.3.** Before we turn to an example, let us briefly state a useful result. Suppose that  $V$  is a continuous stochastic variable with expected value

$$E(V) = \int_{-\infty}^{\infty} v f(v) dv = \mu$$

where  $f(v)$  is the probability density. Then:

$$\begin{aligned} \text{Var}(V) &= \int_{-\infty}^{\infty} (v - \mu)^2 f(v) dv \\ &= \int_{-\infty}^{\infty} v^2 f(v) dv - 2\mu \int_{-\infty}^{\infty} v f(v) dv + \mu^2 \int_{-\infty}^{\infty} f(v) dv \\ &= \int_{-\infty}^{\infty} v^2 f(v) dv - \mu^2 = E(V^2) - \mu^2 \end{aligned}$$

Thus,

$$E(V^2) = \text{Var}(V) + \mu^2 \tag{3.4.9}$$

□

**Example 3.4.4 (Stochastic extension of Example 3.3.2).** Find  $J(x)$  for the problem

.....Alcatel-Lucent 

[www.alcatel-lucent.com/careers](http://www.alcatel-lucent.com/careers)

What if you could build your future and create the future?

One generation's transformation is the next's status quo. In the near future, people may soon think it's strange that devices ever had to be "plugged in." To obtain that status, there needs to be "The Shift".



$$\max_u E \left( \sum_{t=0}^{\infty} \beta^t (-u_t^2 - X_t^2) \right)$$

subject to  $X_{t+1} = X_t + u_t + V_{t+1}$ , where  $\beta \in (0, 1)$ ,  $x_0$  given,  $u_t \in \mathbb{R}$  and  $V_{t+1}$  is normal distributed with expected value  $E(V_{t+1}) = \mu = 0$  and variance  $\text{Var}(V_{t+1}) = \sigma^2 = v$ .

**Solution:** Referring to the deterministic case (Example 3.3.2), we supposed a solution on the form  $J(x) = -\alpha x^2$ . Regarding our problem here, we shall assume that  $J(x)$  is on the form  $J(x) = -ax^2 + b$  since  $E(X_{t+1}^2)$  will contain terms where neither  $X$  nor  $u$  will occur. Thus, from (3.4.8):

$$\begin{aligned} -ax^2 + b &= \max_u \{ -u^2 - x^2 + \beta E [ -a(X + u + V_1)^2 + b ] \} \\ &= \max_u \{ -u^2 - x^2 - \beta a E [ (X + u)^2 - 2(X + u)V_1 + V_1^2 ] + \beta b \} \end{aligned}$$

Now, since  $E(V_1) = 0$ , it follows from (3.4.9) that  $E(V_1^2) = v$ . Hence,

$$-ax^2 + b = \max_u \{ -u^2 - x^2 - \beta a(x + u)^2 - \beta av + \beta b \}$$

Clearly,  $u = -\beta a(1 + \beta a)^{-1}x$  maximizes the expression within the bracket, so

$$-ax^2 + b = -\frac{1 + 2\beta a}{1 + \beta a} x^2 - \beta av + \beta b$$

Equating terms of equal powers yields

$$-a(1 + \beta a) = -(1 + 2\beta a) \tag{3.4.10a}$$

$$b = -\beta av + \beta b \tag{3.4.10b}$$

The solution of (3.7.10a) is easily found to be

$$a = \frac{-(1 - 2\beta) + \sqrt{1 + 4\beta^2}}{2\beta}$$

which implies

$$b = -\frac{\beta av}{1 - \beta}$$

Consequently

$$J(x) = - \left[ \frac{-(1 - 2\beta) + \sqrt{1 + 4\beta^2}}{2\beta} \right] x^2 - \frac{v}{1 - \beta} \left[ \frac{-(1 - 2\beta) + \sqrt{1 + 4\beta^2}}{2} \right]$$

with associated optimal Markov control  $u = -\beta a(1 + \beta a)^{-1}x$ . □

**Exercise 3.4.5.** Find  $J(x, v)$  for the problem:

$$\max_u E \left( \sum_{t=0}^{\infty} \beta^t (-e^{-2X_t}) \right)$$

subject to  $X_{t+1} = X_t - 2u_t + V_{t+1}$ .  $\beta \in (0, 1)$ ,  $x_0$  given,  $u_t \in [-1, 1]$ ,  $V_{t+1} \geq 0$  is

identically distributed with  $E(e^{-2V_{t+1}}) < \infty$ . □

Now, referring to Example 3.4.4 as well as Exercise 3.4.5, it is still not clear if the optimal value functions  $J(x)$  which we found really solve the given optimization problems. The problem is the boundedness condition. Neither of the  $f_0(x, u)$  functions from the example nor the exercise satisfy  $K_1 \leq f_0(x, u) \leq K_2$  (cf. Theorem 3.4.2). Still, there exists a few ways to show that  $J(x)$  can solve a given problem even if the boundedness condition fails (Bertsekas, 1976; Hernández-Lerma, 1996; Sydsæter et al., 2005). One way to proceed is to argue along the following line:

Suppose that  $f_0(x, u) \leq 0$  and  $\beta \in (0, 1)$  (which is the case both in Example 3.4.4 and Exercise 3.4.5). Moreover, assume that we have succeeded in solving the corresponding finite horizon problem (i.e.  $T$  finite), and that  $U$  is compact and  $f_0(x, u), f(x, u)$  are continuous functions of  $(x, u)$ . Denote the optimal value function in the finite case by  $J(0, x, v, T)$ . Then  $\lim_{T \rightarrow \infty} J(0, x, v, T)$  (if it exists!) is the optimal function which solves the infinite horizon problem. We shall now demonstrate (partly as an exercise) that  $J(x)$  found in Exercise 3.4.5 really solves the given optimization problem.

The optimal value function of the infinite horizon problem given in Exercise 3.4.5 is found to be

$$J(x) = -ae^{-2x} = - \frac{1}{1 - \beta K e^{-4}} e^{-2x}$$

where  $K = E(e^{-2V_{t+1}})$ .

Now, consider the corresponding finite horizon problem

$$\max_u E \left( \sum_{t=0}^T \beta^t (-e^{-2X_t}) \right)$$

We leave it as an exercise to the reader to show that the solution of this problem is:

$$J(T, x) = -\beta^T e^{-2x}$$

$u_T^*$  arbitrary and  $J(T - k, x) = -\beta^{T-k} \alpha_k e^{-2x}$ ,  $u_{T-k}^* = 1$ , where  $\alpha_{k+1} = 1 + \beta K e^{-4} \alpha_k$  and  $\alpha_T = 1$ , or alternatively

$$J(t, x) = -\beta^t \alpha_t e^{-2x}$$

where  $\alpha_{t-1} = 1 + \beta K e^{-4} \alpha_t$ . Clearly,  $J(0, x, T) = -\alpha_0 e^{-2x}$  (and  $\alpha_0 = \alpha_0(T)$ ). Our goal is to find  $\lim_{T \rightarrow \infty} J(0, x, T)$  which is the same as finding  $\lim_{T \rightarrow \infty} (-\alpha_0(T))$  which again is the same as finding  $\lim_{t \rightarrow \infty} (-\alpha_t)$  when  $T$  is fixed.

Note that  $(-\alpha_{t-1}) < (-\alpha_t)$  and when  $t \rightarrow \infty$ ,  $\alpha = 1 + \beta K e^{-4} \alpha$ , thus



## Join the best at the Maastricht University School of Business and Economics!

### Top master's programmes

- 33<sup>rd</sup> place Financial Times worldwide ranking: MSc International Business
- 1<sup>st</sup> place: MSc International Business
- 1<sup>st</sup> place: MSc Financial Economics
- 2<sup>nd</sup> place: MSc Management of Learning
- 2<sup>nd</sup> place: MSc Economics
- 2<sup>nd</sup> place: MSc Econometrics and Operations Research
- 2<sup>nd</sup> place: MSc Global Supply Chain Management and Change

Sources: Keuzegids Master ranking 2013; Elsevier 'Beste Studies' ranking 2012; Financial Times Global Masters in Management ranking 2012

Maastricht  
University is  
the best specialist  
university in the  
Netherlands  
(Elsevier)

Visit us and find out why we are the best!

Master's Open Day: 22 February 2014

www.mastersopenday.nl



$$\alpha = \frac{1}{1 - \beta K e^{-4}}$$

which is nothing but the quantity  $a$  in  $J(x)$  obtained in the infinite horizon problem. Consequently, the optimal value function found in Exercise 3.4.5 really solves the optimization problem.

# Appendix (Parameter Estimation)



The image shows the BI Norwegian Business School logo, which consists of a central blue square with the letters 'BI' in white. Surrounding this central square are numerous colorful, 3D-style rectangular bars of various colors (red, orange, yellow, green, blue, purple) radiating outwards. Each bar has a label for a business program: 'Business', 'Strategic Marketing Management', 'International Business', 'Leadership & Organisational Psychology', 'Shipping Management', and 'Financial Economics'. The labels are in different colors matching their respective bars.

**BI NORWEGIAN BUSINESS SCHOOL**

EFMD  
**EQUIS**  
ACCREDITED

## Empowering People. Improving Business.

BI Norwegian Business School is one of Europe's largest business schools welcoming more than 20,000 students. Our programmes provide a stimulating and multi-cultural learning environment with an international outlook ultimately providing students with professional skills to meet the increasing needs of businesses.

BI offers four different two-year, full-time Master of Science (MSc) programmes that are taught entirely in English and have been designed to provide professional skills to meet the increasing need of businesses. The MSc programmes provide a stimulating and multi-cultural learning environment to give you the best platform to launch into your career.

- MSc in Business
- MSc in Financial Economics
- MSc in Strategic Marketing Management
- MSc in Leadership and Organisational Psychology

[www.bi.edu/master](http://www.bi.edu/master)





Referring to both the linear and nonlinear population models presented in Part I and Part II, most of them share the common feature that they contain one or several parameters. Hence, if we want to apply such models on a concrete species (for example a fish stock) we have to use available data in order to estimate these parameters. In this appendix we shall briefly discuss how such estimations may be carried out.

Suppose that we know the size of a population  $x$  at times  $t = 0, 1, 2, \dots, n$ , i.e. that  $x_0, x_1, \dots, x_n$  is known, how do we for example estimate the growth rate  $r$  if the population obeys the difference equation

$$x_{t+1} = x_t e^{r(1-x_t)} \tag{A.1}$$

(the Ricker model)? The usual way to perform such an estimation is first to convert the deterministic model like (A.1) into a stochastic model. Now, following Dennis et al. (1995), ecologists draw a major distinction between different classes of factors which may influence the values of vital parameters and thereby impose stochastic variations in ecological models. Demographic factors such as intrinsic chance of variation of birth and death processes among population inhabitants are factors that occur at an individual level. Environmental factors, chance variations from extrinsic factors occur mainly at population (or age or stage class) level.

Moreover, it appears as a general ecological principle that stochastic fluctuations due to the latter type of factors seem to affect population persistence in a much more serious way than those of demographic type (Dennis et al., 1991).

Now, as is true for the analysis of almost all population models in Part I and Part II, we typically were interested in the population as a whole, not at individual levels. Thus, for our purposes we want to build stochasticity into models like (A.1) of the environmental type. Therefore, we consider the stochastic version of (A.1)

$$x_{t+1} = x_t e^{r(1-x_t)} e^{E_t} \tag{A.2}$$

where  $E_t$  is a normal distributed stochastic variable with expected value  $\mu = 0$  and variance  $\sigma^2$ .

(Recall that if  $Z$  is normal distributed with expected value  $\mu$  and variance  $\sigma^2$  the probability density is given by

$$f(z) = \frac{1}{\sqrt{2\pi}} \frac{1}{\sigma} \exp \left\{ -\frac{1}{2} \left( \frac{z - \mu}{\sigma} \right)^2 \right\} \tag{A.3}$$

and if  $Z_1, \dots, Z_n$  are all normal distributed stochastic variables with expected values and variances  $\mu_1, \dots, \mu_n$  and  $\sigma_1^2, \dots, \sigma_n^2$  respectively we may express the joint probability density function as

$$f(z_1, \dots, z_n) = \frac{1}{\sqrt{2\pi}^n} \frac{1}{\sqrt{|\Sigma|}} \exp \left\{ -\frac{(\mathbf{z} - \boldsymbol{\mu})^T \Sigma^{-1} (\mathbf{z} - \boldsymbol{\mu})}{2} \right\} \quad (\text{A.4})$$

where  $(\mathbf{z} - \boldsymbol{\mu}) = (z_1 - \mu_1, \dots, z_n - \mu_n)$  and the variance covariance matrix  $\Sigma$  is given by

$$\Sigma = \begin{pmatrix} \sigma_1^2 & Cov(Z_1, Z_2) & \cdots & \cdots & Cov(Z_1, Z_n) \\ Cov(Z_1, Z_2) & \sigma_2^2 & \cdots & \cdots & Cov(Z_2, Z_n) \\ \vdots & & \ddots & & \\ \vdots & & & \ddots & \\ Cov(Z_1, Z_n) & Cov(Z_n, Z_2) & & & \sigma_n^2 \end{pmatrix} \quad (\text{A.5})$$

Now, before we turn to (A.1), (A.2) let us first study the estimation problem in a more general context.

Consider

$$\begin{aligned} x_{1,t+1} &= f_1(x_{1,t}, \dots, x_{n,t}, \theta_1, \dots, \theta_q) e^{E_{1,t}} \\ x_{2,t+1} &= f_2(x_{1,t}, \dots, x_{n,t}, \theta_1, \dots, \theta_q) e^{E_{2,t}} \\ &\vdots \\ x_{n,t+1} &= f_n(x_{1,t}, \dots, x_{n,t}, \theta_1, \dots, \theta_q) e^{E_{n,t}} \end{aligned} \quad (\text{A.6})$$

Hence, there are  $n$  state variables  $\mathbf{x} = (x_1, \dots, x_n)^T$ ,  $q$  parameters  $\boldsymbol{\theta} = (\theta_1, \dots, \theta_q)$  and  $\mathbf{E}_t = (E_{1,t}, \dots, E_{n,t})^T$  is a stochastic “environmental noise” vector which is multivariate normal distributed with expected value  $\mathbf{0}$  and variance, covariance matrix  $\Sigma$ . (If there is one variable only, all covariance terms vanish so we are left with  $\mu = 0$  and variance  $v = \sigma^2$ .)

Now, defining

$$\mathbf{M}_{t+1} = (\ln x_{1,t+1}, \dots, \ln x_{n,t+1})^T \quad \mathbf{M}_t = (\ln x_{1,t}, \dots, \ln x_{n,t})^T$$

we may reformulate (A.6) on a logarithmic scale as

$$\mathbf{M}_{t+1} = \mathbf{h}(\mathbf{M}_t) + \mathbf{E}_t \quad (\text{A.7})$$

where  $\mathbf{h}(\mathbf{M}_t) = (\ln f_1(x_{1,t}, \dots, x_{n,t}, \theta_1, \dots, \theta_q), \dots, \ln f_n(x_{1,t}, \dots, x_{n,t}, \theta_1, \dots, \theta_q))^T$  and we may observe that the environmental noise is added to the original model on a logarithmic scale.

Next, assuming that  $\mathbf{y}_t, t = 0, \dots, k$  is  $k + 1$  consecutive time observations of  $\mathbf{x}_t$ , it follows that the conditional expected value  $E(\mathbf{M}_{t+1} | \mathbf{x}_t = \mathbf{y}_t)$  may be expressed as

$$E(\ln \mathbf{x}_{t+1} | \mathbf{x}_t = \mathbf{y}_t) = \ln \mathbf{f}(\mathbf{y}_t, \boldsymbol{\theta}) = \mathbf{h}(\mathbf{m}_t) = \mathbf{h}_t \quad (\text{A.8})$$

Hence, referring to Tong (1995), (A.8) expresses that the nonlinear deterministic skeleton  $\mathbf{x}_{t+1} = \mathbf{f}(\mathbf{x}_t, \boldsymbol{\theta})$  is preserved on a logarithmic scale.

The likelihood function for our problem now becomes

$$I(\boldsymbol{\theta}, \boldsymbol{\Sigma}) = \prod_{t=1}^k p(\mathbf{m}_t | \mathbf{m}_{t-1}) \quad (\text{A.9})$$

(where  $\mathbf{m}$  (as in (A.8)) contains the observation values  $\mathbf{y}$ ) and we may interpret  $I$  as a measure of the likelihood of the observations at each time as functions of the unknown parameters.

## Need help with your dissertation?

Get in-depth feedback & advice from experts in your topic area. Find out what you can do to improve the quality of your dissertation!

Get Help Now



Go to [www.helpmyassignment.co.uk](http://www.helpmyassignment.co.uk) for more info



Click on the ad to read more

Now, following Dennis et al. (1995), the probability  $p(\mathbf{m}_t | \mathbf{m}_{t-1})$  is the joint probability density for  $\mathbf{M}_t$  conditional of  $\mathbf{M}_{t-1} = \mathbf{m}_{t-1}$ . It is a multivariate normal probability density which according to (A.8) possesses the expected value  $E(\mathbf{M}_t) = \mathbf{h}(\mathbf{m}_{t-1})$  and variance, covariance matrix given by  $\Sigma$ . Therefore, by use of (A.8) and (A.4) we may express the joint probability distribution as

$$p(\mathbf{m}_t | \mathbf{m}_{t-1}) = \frac{1}{(2\pi)^{n/2} \sqrt{|\Sigma|}} \exp \left\{ \frac{(\mathbf{m}_t - \mathbf{h}_{t-1})^T \Sigma^{-1} (\mathbf{m}_t - \mathbf{h}_{t-1})}{-2} \right\} \quad (\text{A.10})$$

The maximum likelihood parameters are now obtained by computing zeros of derivatives of (A.9) with respect to  $\theta_1, \dots, \theta_q$  and  $\Sigma$ . Moreover, calculations are simplified if we first apply the logarithm. Thus, instead of computing the derivatives directly from (A.9) we compute the derivatives of

$$\begin{aligned} \ln I(\boldsymbol{\theta}, \Sigma) &= \sum_{t=1}^k \ln p(\mathbf{m}_t | \mathbf{m}_{t-1}) \\ &= -\frac{nk}{2} \ln 2\pi - \frac{k}{2} \ln |\Sigma| - \frac{1}{2} \sum_{t=1}^k (\mathbf{m}_t | \mathbf{h}_{t-1})^T \Sigma^{-1} (\mathbf{m}_t | \mathbf{h}_{t-1}) \end{aligned} \quad (\text{A.11})$$

Estimates obtained from (A.9), (A.11) are often referred to as maximum likelihood estimates. Evidently, the log-likelihood function (A.11) is complicated in case of several state variables  $x_1, \dots, x_n$ . Therefore, most estimations must be done by use of numerical algorithms. One such frequently used algorithm which has several desired statistical properties is the Nelder–Mead simplex algorithm which is described in Press et al. (1992). Here, we shall concentrate on cases where it is possible to estimate parameters without using numerical methods.

To this end, consider the stochastic difference equation with one state variable

$$x_{t+1} = f(x_t, \boldsymbol{\theta}) e^{E_t} \quad (\text{A.12})$$

which we may interpret as the stochastic version of almost all nonlinear maps considered in Part I. Now, since  $n = 1$ , the variance, covariance matrix  $\Sigma$  degenerates to only one term, namely the variance  $v$ . (We prefer  $v$  instead of  $\sigma^2$  for notation convenience.) If we in addition have  $k + 1$  observation points  $y_t$  of  $x_t$  at times  $0, 1, \dots, k$ , the log-likelihood function (A.11) may be cast in the form

$$\ln I(\theta_1, \dots, \theta_q, v) = -\frac{k}{2} \ln 2\pi - \frac{k}{2} \ln v - \frac{1}{2v} \sum_{t=1}^k u_t^2(\theta_1, \dots, \theta_q) \quad (\text{A.13})$$

where the log-residuals  $u_t = \ln y_t - \ln f(y_{t-1}, \theta_1, \dots, \theta_q)$ .

The maximum likelihood parameter estimators are then obtained from

$$\frac{\partial \ln I}{\partial \theta_i} = -\frac{1}{v} \sum_{t=1}^k u_t(\theta_1, \dots, \theta_q) \frac{\partial u_t}{\partial \theta_i}(\theta_1, \dots, \theta_q) = 0 \quad (\text{A.14a})$$

$$\frac{\partial \ln I}{\partial v} = -\frac{k}{2v} + \frac{1}{2v^2} \sum_{t=1}^k u_t^2(\theta_1, \dots, \theta_q) = 0 \quad (\text{A.14b})$$

or equivalently (by use of the definition of  $u_t$ ) from

$$\sum_{t=1}^k u_t(\theta_1, \dots, \theta_q) \frac{\frac{\partial f}{\partial \theta_i}(y_{t-1}, \theta_1, \dots, \theta_q)}{f(y_{t-1}, \theta_1, \dots, \theta_q)} = 0 \quad (\text{A.15a})$$

where  $i = 1, 2, \dots, q$  and

$$v = \frac{1}{k} \sum_{t=1}^k u_t^2(\theta_1, \dots, \theta_q) \quad (\text{A.15b})$$

**Example A.1.** Suppose that we have observations  $y_t$  of  $x_t$  at times  $t = 0, 1, \dots, k$  (i.e.  $k + 1$  observations  $y_t$ ) and estimate  $r$  in the nonlinear equation (A.1).

**Solution:** Consider the stochastic version of (A.1)

$$x_{t+1} = f(x_t, r) e^{E_t} = x_t e^{r(1-x_t)} e^{E_t}$$

(which is nothing but (A.2)). The log-residuals become

$$u_t = \ln y_t - \ln(y_{t-1} e^{r(1-y_{t-1})}) = \ln y_t - \ln y_{t-1} - r(1 - y_{t-1})$$

Thus, according to (A.15a)

$$\sum_{t=1}^k \left\{ \ln y_t - \ln y_{t-1} - r(1 - y_{t-1}) \right\} \frac{y_{t-1}(1 - y_{t-1}) e^{r(1-y_{t-1})}}{y_{t-1} e^{r(1-y_{t-1})}} = 0$$

or

$$\sum_{t=1}^k \left\{ \ln y_t - \ln y_{t-1} - r(1 - y_{t-1}) \right\} (1 - y_{t-1}) = 0$$

from which we obtain

$$r = \frac{\sum_{t=1}^k (1 - y_{t-1}) \ln \left( \frac{y_t}{y_{t-1}} \right)}{\sum (1 - y_{t-1})^2} \quad (\text{A.16})$$

The variance  $v$  may be estimated from (A.15b) as

$$v = \frac{1}{k} \sum_{t=1}^k u_t^2(\theta_1, \dots, \theta_q) = \frac{1}{k} \sum_{t=1}^k \left\{ \ln \left( \frac{y_t}{y_{t-1}} \right) - r(1 - y_{t-1}) \right\}^2$$

after we have first estimated  $r$  from (A.16). □

**Example A.2.** Suppose that we have observations  $y_t$  of  $x_t$  at consecutive times  $t = 0, \dots, k$  and estimate the parameters  $F$  and  $r$  in the equation

$$x_{t+1} = f(x_t, F, r) = Fx_t e^{-rx_t} \tag{A.17}$$

**Solution:** The stochastic version of (A.17) becomes

$$x_{t+1} = Fx_t e^{-rx_t} e^{E_t}$$

so the log-residuals may be expressed as

$$u_t = \ln y_t - \ln(Fy_{t-1}e^{-ry_{t-1}}) = \ln \left( \frac{y_t}{y_{t-1}} \right) - \ln F + ry_{t-1}$$

**Brain power**

By 2020, wind could provide one-tenth of our planet's electricity needs. Already today, SKF's innovative know-how is crucial to running a large proportion of the world's wind turbines.

Up to 25 % of the generating costs relate to maintenance. These can be reduced dramatically thanks to our systems for on-line condition monitoring and automatic lubrication. We help make it more economical to create cleaner, cheaper energy out of thin air.

By sharing our experience, expertise, and creativity, industries can boost performance beyond expectations. Therefore we need the best employees who can meet this challenge!

The Power of Knowledge Engineering

Plug into The Power of Knowledge Engineering.  
Visit us at [www.skf.com/knowledge](http://www.skf.com/knowledge)

**SKF**

Hence equation(s) (A.15a) may be cast in the form

$$\sum_{t=1}^k \left\{ \ln \left( \frac{y_t}{y_{t-1}} \right) - \ln F + ry_{t-1} \right\} \frac{y_{t-1} e^{-ry_{t-1}}}{F y_{t-1} e^{-ry_{t-1}}} = 0$$

$$\sum_{t=1}^k \left\{ \ln \left( \frac{y_t}{y_{t-1}} \right) - \ln F + ry_{t-1} \right\} \frac{(-F) y_{t-1}^2 e^{-ry_{t-1}}}{F y_{t-1} e^{-ry_{t-1}}} = 0$$

or equivalently

$$k \ln F - Ar = B \tag{A.18a}$$

$$A \ln F - Cr = D \tag{A.18b}$$

where

$$A = \sum_{t=1}^k y_{t-1}, \quad B = \sum_{t=1}^k \ln \left( \frac{y_t}{y_{t-1}} \right)$$

$$C = \sum_{t=1}^k y_{t-1}^2, \quad D = \sum_{t=1}^k y_{t-1} \ln \left( \frac{y_t}{y_{t-1}} \right)$$

Consequently, from (A.18)

$$\ln F = \frac{AD - BC}{A^2 - kC} \quad r = \frac{kD - AB}{A^2 - kC}$$

Finally, (A.15b) implies

$$v = \frac{1}{k} \sum_{i=1}^k \left\{ \ln \left( \frac{y_t}{y_{t-1}} \right) - \ln F + ry_{t-1} \right\}^2$$

$$= \frac{1}{k} \{ G - 2B \ln F + 2rD - k(\ln F)^2 - 2rA \ln F + r^2 C \}$$

where

$$G = \sum_{i=1}^k \ln^2 \left( \frac{y_t}{y_{t-1}} \right)$$

□



**Remark A.1.** Cushing (1998) considers a similar model as (A.17) where he generates data points at 60 consecutive times. There, he obtains estimates of parameters  $b$  and  $c$  (corresponding to  $F$  and  $r$  in (A.17)) which accurately recover the correct parameters used in the generation of data with seven significant digits. For further details, see Cushing (1998).  $\square$

**Exercise A.1.** Suppose that we have observations  $y_t$  of  $x_t$  at  $k + 1$  consecutive times  $t = 0, \dots, k$  and estimate  $\mu$  in equation (1.2.1) (the quadratic map).  $\square$

**Exercise A.2.** Use (A.15) and estimate parameters  $a$  and  $b$  in the Hassel family

$$x_{t+1} = \frac{ax_t}{(1+x_t)^b} \quad a > 1, \quad b > 1$$

by use of observation points  $y_t$  of  $x_t$  at times  $t = 0, \dots, k$ .  $\square$

—

In the previous examples (and exercises) the estimations have been carried out by use of the log-likelihood function (A.11). Another possibility is to apply conditional least squares and we close this appendix by giving a brief overview of the method. (We still denote state variables by  $\mathbf{x}$ , observations by  $\mathbf{y}$  and parameters by  $\boldsymbol{\theta}$ .)

Now, suppose that we have  $k + 1$  consecutive time observations  $\mathbf{y}_0, \dots, \mathbf{y}_k$ , the purpose of the method is to minimize log-residuals (recall that environmental noise is additive on a logarithmic scale, cf. (A.7)) so if we are dealing with a map  $\mathbf{x} \rightarrow \mathbf{f}(\mathbf{x}, \boldsymbol{\theta})$  we search for parameter estimates that minimize

$$D = \sum_{i=1}^k (\ln y_t - \ln \mathbf{f}(y_{t-1}, \boldsymbol{\theta}))^2 \tag{A.19}$$

Estimates found through (A.19) are often referred to as conditional least squares estimates because they are found (on a logarithmic scale) through a minimization of conditional sums of squares. We shall now by way of examples show how the method works.

**Example A.3.** Assuming  $k + 1$  time observations  $y_0, \dots, y_k$ , estimate parameter  $r$  in map (A.1).

**Solution:** In this case (A.19) becomes

$$D = \sum_{i=1}^k (\ln y_t - \ln(y_{t-1}e^{r(1-y_{t-1})}))^2 = \sum \left( \ln \left( \frac{y_t}{y_{t-1}} \right) - r(1 - y_{t-1}) \right)^2$$

Hence,

$$\frac{\partial D}{\partial r} = 0 \Leftrightarrow \sum 2 \left[ \ln \left( \frac{y_t}{y_{t-1}} \right) - r(1 - y_{t-1}) \right] (y_{t-1} - 1) = 0$$

which yields

$$r = \frac{\sum (1 - y_{t-1}) \ln \left( \frac{y_t}{y_{t-1}} \right)}{\sum (1 - y_{t-1})^2}$$

in accordance with the result we obtained by use of (A.15). (Also note that  $\partial^2 D / \partial r^2 = \sum (1 - y_{t-1})^2 > 0$ , hence the  $r$  estimate really corresponds to a minimum.)  $\square$

**Exercise A.3.** Given  $k + 1$  time observations  $y_0, \dots, y_k$  of  $x_t$ , estimate  $F$  and  $r$  in equation (A.17). (Compare with the results of Example A.2.)  $\square$

## TURN TO THE EXPERTS FOR SUBSCRIPTION CONSULTANCY

Subscribe is one of the leading companies in Europe when it comes to innovation and business development within subscription businesses.

We innovate new subscription business models or improve existing ones. We do business reviews of existing subscription businesses and we develop acquisition and retention strategies.

Learn more at [linkedin.com/company/subscribe](https://www.linkedin.com/company/subscribe) or contact  
Managing Director Morten Suhr Hansen at [mha@subscribe.dk](mailto:mha@subscribe.dk)

**SUBSCR**✓**BE** - to the future



Referring to the  $n$ -dimensional nonlinear population models considered in Part II, most of them have from an estimation point of view a desired property, namely that the various parameters in the model occur in one equation only. (See for example the three-dimensional model presented in Exercise 2.8.2. Here the fecundity  $F_2$  is in the first equation, parameter  $P_0$  is in the second equation only and  $P_1$  shows up in the third equation only.) In such cases the method of conditional least squares is particularly convenient to use because we may apply the method on each equation in the model separately. As an illustration, consider the following example:

**Example A.4.** Consider the nonlinear map or difference equation model

$$x_{1,t+1} = Fx_{2,t} \tag{A.20}$$

$$x_{2,t+1} = Pe^{-(x_{1,t}+x_{2,t})}x_{1,t}$$

Note that (A.20) is a special case of (2.8.2), ( $\alpha = 1$ ), which was extensively studied in Example 2.8.1. Since  $\alpha$  acts as a scaling factor only, (A.20) possesses the same dynamics as (2.8.2). In case of “small” values of  $F$  the dynamics is a stable nontrivial equilibrium. Nonstationary dynamics is introduced through a supercritical Hopf bifurcation and when  $F$  is increased beyond instability threshold, the various dynamical outcomes are displayed in Figures 16–20 (cf. Example 2.8.1).

Now, suppose a time series of  $k + 1$  observation points  $(y_{1,0}, y_{2,0}), \dots, (y_{1,k}, y_{2,k})$  of  $(x_{1,t}, x_{2,t})$ . Our goal is to use these points in order to estimate  $F$  and  $P$  by applying conditional least squares. To this end (cf. (A.19)), define

$$D_1 = \sum_{t=1}^k [\ln y_{1,t} - \ln(Fy_{2,t-1})]^2 = \sum \left[ \ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) - \ln F \right]^2$$

$$D_2 = \sum_{t=1}^k \left[ \ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) - \ln P + y_{1,t-1} + y_{2,t-1} \right]^2$$

The equations  $\partial D_1 / \partial F = 0$ ,  $\partial D_2 / \partial P = 0$  give respectively

$$\sum \ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) - k \ln F = 0$$

$$\sum \left( \ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) + y_{1,t-1} + y_{2,t-1} \right) - k \ln P = 0$$

Consequently, we may estimate  $F$  and  $P$  through

$$\ln F = \frac{1}{k} \sum_{t=1}^k \ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) \quad (\text{A.21a})$$

$$\ln P = \frac{1}{k} \sum_{t=1}^k \left( y_{1,t-1} + y_{2,t-1} + \ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) \right) \quad (\text{A.21b})$$

In order to investigate how good the estimates really are we have performed the following “experiment”. Let  $F = 27.0$  and  $P = 0.6$ . Then, from (A.20) we have generated a time series of 50 “observation points”  $(y_{1,t}, y_{2,t})$ . The points are located on a chaotic attractor as displayed in Figure 20. Next, “pretending” that  $F$  and  $P$  are unknown we have used the “observations” in (A.21) in order to estimate  $F$  and  $P$ . The result is,  $F = 27.00003065$  and  $P = 0.6000000143$  so the estimation appears to be excellent.  $\square$

Still considering the map (A.20) let us for comparison reasons also find the maximum likelihood estimates of  $F$  and  $P$ . Suppose that

$$\Sigma_2 = \begin{pmatrix} \sigma_1^2 & c \\ c & \sigma_2^2 \end{pmatrix}$$

Then, by use of (A.4), (A.5) we may express (A.11) as

$$\begin{aligned} \ln I(F, P, \Sigma_2) = & -k \ln 2\pi - \frac{k}{2} \ln |\sigma_1^2 \sigma_2^2 - c^2| \\ & - \frac{1}{2(1-\rho^2)} \sum_{t=1}^k \left\{ \left( \frac{\ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) - \ln F}{\sigma_1} \right)^2 + \left( \frac{\ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) + y_{t-1} - \ln P}{\sigma_2} \right)^2 \right. \\ & \left. - 2\rho \frac{\left[ \ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) - \ln F \right] \left[ \ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) + y_{t-1} - \ln P \right]}{\sigma_1 \sigma_2} \right\} \end{aligned}$$

where  $y_{t-1} = y_{1,t-1} + y_{2,t-1}$  and  $\rho = c/\sigma_1\sigma_2$ .

The equations  $\partial(\ln I)/\partial F = 0$  and  $\partial(\ln I)/\partial P = 0$  may be cast in the forms

$$\frac{k}{\sigma_1} \ln F - \frac{k\rho}{\sigma_2} \ln P = \frac{A}{\sigma_1} - \frac{\rho B}{\sigma_2} \quad (\text{A.22a})$$

$$- \frac{k\rho}{\sigma_1} \ln F + \frac{k}{\sigma_2} \ln P = \frac{B}{\sigma_2} - \frac{\rho A}{\sigma_1} \quad (\text{A.22b})$$

where

$$A = \sum \ln \left( \frac{y_{1,t}}{y_{2,t-1}} \right) \quad B = \sum \left[ \ln \left( \frac{y_{2,t}}{y_{1,t-1}} \right) + y_{t-1} \right]$$

The solution of (A.22a,b) is easily found to be

$$\ln F = \frac{1}{k} A \quad \text{and} \quad \ln P = \frac{1}{k} B$$

which is the same as we obtained by use of conditional least squares.

**Exercise A.4.** Given  $k + 1$  time observations  $(y_{1,0}, y_{2,0}), \dots, (y_{1,k}, y_{2,k})$  of  $(x_{1,t}, x_{2,t})$  and find the conditional least squares estimates of  $F$ ,  $P$  and  $\alpha$  in the age structured Ricker model

$$\begin{aligned} x_{1,t+1} &= Fx_{1,t}e^{-\alpha x_t} + Fx_{2,t}e^{-\alpha x_t} \\ x_{2,t+1} &= Px_{1,t} \end{aligned}$$

where  $x = x_1 + x_2$ . □

**Exercise A.5.** Given  $k + 1$  consecutive time observations, find the conditional least squares estimates of all parameters in the map

$$(x_1, x_2) \rightarrow (Fe^{-\alpha x} x_2, Pe^{-\beta x} x_1)$$

□

# References

Adler F. (1990) Coexistence of two types on a single resource in discrete time. *J. Math. Biol.* **28**, 695–713.

Barnsley M. (1988) *Fractals Everywhere*. Academic Press Inc.

Beddington J.R., Free C.A. and Lawton J.H. (1975) Dynamic complexity in predator–prey models framed in difference equations. *Nature*, **255**, 58–60.

Behncke H. (2000) Periodical cicadas. *J. Math. Biol.*, **40**, 413–431.

Bergé P., Pomeau Y. and Vidal C. (1984) *Order within Chaos*. John Wiley & Sons.

Bergh M.O. and Getz W.M. (1988) Stability of discrete age-structured and aggregated delay-difference population models. *J. Math. Biol.*, **26**, 551–581.

Bertsekas D.P. (1976) *Dynamic Programming and Stochastic Control*. Academic Press.

Bernardelli H. (1941) Population waves. *Journal of Burma Research Society*, **31**, 1–18.

What do you want to do?

No matter what you want out of your future career, an employer with a broad range of operations in a load of countries will always be the ticket. Working within the Volvo Group means more than 100,000 friends and colleagues in more than 185 countries all over the world. We offer graduates great career opportunities – check out the Career section at our web site [www.volvogroup.com](http://www.volvogroup.com). We look forward to getting to know you!

**VOLVO**  
AB Volvo (publ)  
[www.volvogroup.com](http://www.volvogroup.com)

VOLVO TRUCKS | RENAULT TRUCKS | MACK TRUCKS | VOLVO BUSES | VOLVO CONSTRUCTION EQUIPMENT | VOLVO PENTA | VOLVO AERO | VOLVO IT  
VOLVO FINANCIAL SERVICES | VOLVO 3P | VOLVO POWERTRAIN | VOLVO PARTS | VOLVO TECHNOLOGY | VOLVO LOGISTICS | BUSINESS AREA ASIA



Botsford L.W. (1986) Population dynamics of the Dungeness crab (*Cancer magister*). *Can. Spec. Publ. Fish. Aquat. Sci.*, **92**, 140–153.

Caswell H. (2001) *Matrix Population Models*. Sinauer Ass., Inc. Sunderland, Massachusetts.

Clark C.W. (1976) A delayed-recruitment model of population dynamics with an application to baleen whale populations. *J. Math. Biol.*, **3**, 381–391.

Collet P. and Eckmann J.P. (1980) Iterated maps on the interval as dynamical systems. *Progress on Physics*, Vol. 1, Birkhäuser – Boston, Boston.

Costantino R.F., Desharnais R.A., Cushing J.M., and Dennis B. (1997) Chaotic dynamics in an insect population. *Science*, **275**, 389–391.

Cushing J.M. (1998) An introduction to structured population dynamics. *SIAM*, Philadelphia.

Cushing J.M., Dennis B., Desharnais R.A. and Costantino R.F. (1996) An interdisciplinary approach to understanding nonlinear ecological dynamics. *Ecol. Model.*, **92**, 111–119.

Cushing J.M., Costantino R.F., Dennis B., Desharnais R. A. and Henson S. M. (1998) Nonlinear population dynamics: models, experiments and data. *J. Theor. Biol.*, **194**, 1–9.

Cvitanović P. (1996) *Universality of Chaos*. Institute of Physics Publishing.

Davydova N.V., Diekmann O. and van Gils S.A. (2003) Year class coexistence or competitive exclusion for strict biennials. *J. Math. Biol.*, **46**, 95–131.

Dennis B., Munholland P.L. and Scott J.M. (1991) Estimation of growth and extinction parameters for endangered species. *Ecological Monographs*, **61**, 115–143.

Dennis B., Desharnais R.A., Cushing J. M. and Costantino R.F. (1995) Nonlinear Demographic Dynamics: Mathematical Models, Statistical Methods, and Biological Experiments. *Ecological Monographs*, **65**, 261–281.

Dennis B., Desharnais R.A., Cushing J.M. and Costantino R.F. (1997) Transition in population dynamics: equilibria to periodic cycles to aperiodic cycles. *J. Anim. Ecol.*, **66**, 704–729.

Devaney R.L. (1989) *An Introduction to Chaotic Dynamical Systems*. Addison–Wesley Publishing Company, Inc.



- Diekmann O. and van Gils S. (2000) Difference Equations with Delay. *Japan J. Indust. Appl. Math.*, **17**, 73–84.
- Edelstein–Keshet L. (1988) *Mathematical Models in Biology*. Random House, New York.
- Feigenbaum M. J. (1978) Quantitative universality for a class of nonlinear transformations. *J. Stat. Phys.*, **19**, 25–52.
- Frauenthal J. (1986) Analysis of Age-Structure Models. *Biomathematics*, Vol. 17, *Mathematical Ecology*, eds. Hallan T.G. and Levin S.A. Springer Verlag, Berlin, Heidelberg.
- Govaerts W. and R. Khoshsiar Ghazani (2006) Numerical bifurcation analysis of a nonlinear stage structured cannibalism model. *J. difference eqn. and applications*, **12**, 1069–1085.
- Grimshaw R. (1990) *Nonlinear Ordinary Differential Equations*. Blackwell Scientific Publications.
- Guckenheimer J. (1977) On the bifurcation of maps of the interval. *Invent. Math.*, **39**, 165–178.
- Guckenheimer J. and Holmes P. (1983) *Nonlinear Oscillations, Dynamical Systems, and Bifurcation of Vector Fields*. Springer Verlag.
- Guckenheimer J., Oster G.F., and Ipaktchi A. (1977) The dynamics of density dependent population models. *J. Math. Biol.*, **4**, 101–147.
- Hartman P. (1964) *Ordinary Differential Equations*. Wiley, New York.
- Hassel M.P. (1978) *The Dynamics of Arthropod Predator–Prey Systems*. Princeton University Press.
- Hénon M. (1976) A two dimensional mapping with a stange attractor. *Comm. Math. Phys.*, **50**, 69–77.
- Hernández–Lerma O. and Lasserre J.B. (1996) *Discrete-Time Markov Control Processes*. Springer Verlag.
- Higgins K., Hastings A. and Botsford L.W. (1997) Density dependence and age structure: nonlinear dynamics and population behaviour. *Am. Nat.*, **149**, 247–269.
- Horn R.A. and Johnson C. R. (1985) *Matrix Analysis*. Cambrigde University Press.
- International Whaling Commission (1979) Report No. 29. International Whaling Commission, Cambridge.

Iooss G. (1979) Bifurcation of maps and applications. *North Holland Math. Studies*, **36**.

Iooss G. and Adelmeyer M. (1999) *Topics in Bifurcation Theory*. Adv. Series in Nonlinear Dynamics, 3, World Sci. 2nd ed.

Jost J. (2005) *Dynamical Systems*. Springer Verlag, Berlin, Heidelberg, New York.

Katok A. and Hasselblatt B. (1995) *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press.

King A. and Schaffer W. (1999) The rainbow bridge: Hamiltonian limits and resonance in Hamiltonian limits and resonance in predator–prey dynamics. *J. Math. Biol.*, **39**, 439–469.

Kon R. (2005) Nonexistence of synchronous orbits and class coexistence in matrix population models. *SIAM J. Appl. Math.*, **66**, 616–626.

Kon R., Saito Y. and Takeuchi T. (2004) Permanence of single-species stage-structured models. *J. Math. Biol.*, **48**, 515–528.

Kot M. (2001) *Elements of Mathematical Ecology*. Cambridge University Press.

 **gaieteye**<sup>®</sup>  
Challenge the way we run

**EXPERIENCE THE POWER OF  
FULL ENGAGEMENT...**

**RUN FASTER.  
RUN LONGER..  
RUN EASIER...**

**READ MORE & PRE-ORDER TODAY  
WWW.GAITEYE.COM**

- Kuznetsov Y.A. (2004) *Elements of Applied Bifurcation Theory*. 3rd ed., New York, Springer-Verlag.
- Kuznetsov Y.A. and Meijer H.G.E. (2005) Numerical normal forms for codim 2 bifurcations of maps with at most two critical eigenvalues. *SISC*, **26**, 1932–1954.
- Leslie P.H. (1945) On the use of matrices in certain population mathematics. *Biometrika*, **33**, 183–212.
- Leslie P.H. (1948) Some further notes on the use of matrices in population mathematics. *Biometrika*, **35**, 213–245.
- Levin S.A. and May R.M. (1976) A note on difference-delay equations. *Theor. pop. biol.*, **9**, 178–187.
- Levin S.A. and Goodyear P.H. (1980) Analysis of an age-structured fishery model. *J. Math. Biol.*, **9**, 245–274.
- Lewis E. G. (1942) On the generation and growth of a population. *Sankhya: The Indian Journal of Statistics*, **6**, 93–96.
- Li T.Y. and Yorke J.A. (1975) Period three implies chaos. *Am. Math. Monthly*, **82**, 985–992.
- Marsden J.E. and McCracken M. (1976) *The Hopf Bifurcation and its Applications*. Springer Verlag, New York, Heidelberg.
- May R.M. (1976) Simple mathematical models with very complicated dynamics. *Nature*, **261**, 459–467.
- Maynard Smith J. (1968) *Mathematical Ideas in Biology*. Cambridge University Press.
- Maynard Smith J. (1979) *Models in Ecology*. Cambridge University Press.
- Meyer C.D. (2000) *Matrix Analysis and Applied Linear Algebra*. SIAM, Philadelphia.
- Mills N.J. and Getz W.M. (1996) Modelling the biological control of insect pests: a review of host-parasitoid models. *Ecological Modelling*, **92**, 121–143.
- Mjølhus E., Wikan A., and Solberg, T. (2005) On synchronization in semelparous populations. *J. Math. Biol.*, **50**, 1–21.
- Moore G. (2008) From Hopf to Neimark Sacker bifurcation: a computational algorithm. *Int. J. Comp. Sci. Math.*, **2**, 132–180.

- Murdoch W.W. (1994) Population regulation in theory and practice. *Ecology*, **75**, 271–287.
- Murray J.D. (1989) *Mathematical Biology*. Springer, Berlin, Heidelberg.
- Nagashima H. and Baba Y. (1999) *Introduction to Chaos*. Institute of Physics Publishing, Bristol and Philadelphia.
- Neimark Y.I. and Landa P.S. (1992) *Stochastic and Chaotic Oscillations*. Dordrecht Kluwer Acc. Publ.
- Neubert M.G. and Kot M. (1992) The subcritical collapse of predator populations in discrete-time predator–prey models. *Math. Biosciences*, **110**, 45–66.
- Nicholson A.J. (1933) The balance of animal populations. *Journal of Animal Ecology* **2**, 132–178.
- Nicolson A.J. and Bailey V.A. (1935) The balance of animal populations. Part I. *Proceedings of Zoological Society of London*, **3**, 551–598.
- Saber E., Györi I. and Ladas G. (1998) Advances in Difference Equations. *Proceedings of the Second International Conference on Difference Equations*. CRC Press.
- Sacker R.J. (1964) On invariant surfaces and bifurcation of periodic solutions of ordinary differential equations. IMM–NYC, Courant Inst. Math. Sci., New York University.
- Sacker R.J. (1965) A new approach to the perturbation theory of invariant surfaces. *Comm. Pure and Appl. Math.*, **18**, 717–732.
- Seierstad A. and Sydsæter K. (1987) *Optimal Control Theory with Economic Applications*. North-Holland.
- Silva J.A.L. and Hallam T. (1992) Compensation and Stability in Nonlinear Matrix Models. *Math. Biosci.*, **110**, 67–101.
- Silva J.A. and Hallam T.G. (1993) Effects of delay, truncation and density dependence in reproduction schedules on stability of nonlinear Leslie matrix models. *J. Math. Biol.*, **31**, 367–395.
- Singer D. (1978) Stable orbits and bifurcation of maps of the interval. *SIAM, J. Appl. Math.*, **35**, 260–267.
- Smale S. (1963) Diffeomorphisms with many periodic points. *Differential and Combinatory Topology*, S. S. Cairns (ed.), 63–80, Princeton University Press, Princeton.
- Smale S. (1967) Differential Dynamical Systems. *Bull. Amer. Math. Soc.*, **73**, 747–817.

Spiegel M. (1974) *Complex Variables*. Schaum's Outline Series, McGraw-Hill.

Stuart A.M. and Humphries A.R. (1998) *Dynamical Systems and Numerical Analysis*. Cambridge University Press.

Sydsæter K. (2002) *Matematisk analyse*, Bind II, Universitetsforlaget (in Norwegian).

Sydsæter K., Hammond P., Seierstad A., and Strøm A. (2005) *Further Mathematics for Economic Analysis*. Prentice Hall.

Thieullen P.H., Tresser C. and Young L.S. (1994) Positive Lyapunov exponent for generic one-parameter families of unimodal maps. *Journal d'Analyse Mathématique*, **64**, 121–172.

Thunberg H. (2001) Periodicity versus Chaos in One-Dimensional Dynamics, *SIAM Rev.*, **43**, 3–30.

Tong H. (1995) *Non-linear Time Series*. Oxford Science Publications.

Tsuji M. (1993) Positive Lyapunov exponents in families of one dimensional dynamical systems. *Invent. math.*, **111**, 113–137.



This e-book  
is made with  
**SetaPDF**



**SETASIGN**

PDF components for PHP developers

[www.setasign.com](http://www.setasign.com)

- Tuljapurkar S., Boe C. and Wachter K.W. (1994) Nonlinear Feedback Dynamics in Fisheries. Analysis of the Deriso–Schnute Model. *Can. J. Fish. Aquat. Sci.*, **51**, 1462–1472.
- Vanderbauwhede A. (1987) Invariant manifolds in infinite dimensions. *Dynamics of Infinite Dimensional Systems*, ed. S.N. Chow and J.K. Hale, Springer Verlag, Berlin.
- Van Dooren T.J.M. and Metz J.A.J. (1998) Delayed maturation in temporally structured populations with non-equilibrium dynamics. *J. Evol. Biol.*, **11**, 41–62.
- Wan Y.H. (1978) Computations of the stability condition for the Hopf bifurcation of diffeomorphisms on  $\mathbb{R}^2$ . *SIAM, J. Appl. Math.*, **34**, 167–175.
- Wikan A. (1994) *Bifurcations, Nonlinear Dynamics, and Qualitative Behaviour in a Certain Class of Discrete Age-Structured Population Models*. University of Tromsø, Norway.
- Wikan A. (1997) Dynamic consequences of reproductive delay in Leslie matrix models with nonlinear survival probabilities. *Math. Biosci.*, **146**, 37–62.
- Wikan A. (1998) Four-periodicity in Leslie matrix models with density dependent survival probabilities. *Theor. Popul. Biol.*, **53**, 85–97.
- Wikan A. (2001) From chaos to chaos. An analysis of a discrete age-structured prey–predator model. *J. Math. Biol.*, **43**, 471–500.
- Wikan A. (2012a) Age or stage structure? A comparison of dynamic outcomes from discrete age- and stage-structured population models. *Bull. Math. Biol.*, **74(6)**, 1354–1378.
- Wikan A. (2012b) On nonlinear age- and stage-structured population models. *J. Math. & Stat.*, **8(2)**, 311–322.
- Wikan A. and Mjølhus E. (1995) Periodicity of 4 in age-structured population models with density dependence. *J. Theor. Biol.*, **173**, 109–119.
- Wikan A. and Mjølhus E. (1996) Overcompensatory recruitment and generation delay in discrete age-structured population models. *J. Math. Biol.*, **35**, 195–239.
- Wikan A. and Eide A. (2004) An analysis of a nonlinear stage-structured cannibalism model with application to the northeast Arctic cod stock. *Bull. Math. Biol.*, **66**, 1685–1704.
- Zhang Q. and Tian R. (2008) Calculation of Coefficients of Simplest Normal Forms of Neimark–Sacker and Generalized Neimark–Sacker Bifurcations. *Journal of Physics: Conference Series*, 96, 012152.