

Bretherick's
Handbook of
**Reactive
Chemical
Hazards**

Sixth Edition

Volume 1

AN INDEXED GUIDE TO PUBLISHED DATA

Bretherick's
Handbook of
**Reactive
Chemical
Hazards**

Sixth Edition — Volume 1

Edited by
P G Urben
Akzo-Nobel

Compiler
M J Pitt
Department of Chemical Engineering
University of Leeds

BUTTERWORTH
HEINEMANN

OXFORD AUCKLAND BOSTON JOHANNESBURG MELBOURNE NEW DELHI

Butterworth-Heinemann Ltd
Linacre House, Jordan Hill, Oxford OX2 8DP

 A member of the Reed Elsevier plc group

First published 1975
Second edition 1979
Reprinted 1979, 1981, 1984
Third edition 1985
Reprinted 1987, 1989
Fourth edition 1990
Fifth edition 1995
Sixth edition 1999
© Butterworth-Heinemann 1999

All rights reserved. No part of this publication may be reproduced in any material form (including photocopying or storing in any medium by electronic means and whether or not transiently or incidentally to some other use of this publication) without the written permission of the copyright holder except in accordance with the provisions of the Copyright, Designs and Patents Act 1988 or under the terms of a licence issued by the Copyright Licensing Agency Ltd, 90 Tottenham Court Road, London, England W1P 9HE. Applications for the copyright owners written permission to reproduce any part of this publication should be addressed to the Publishers.

British Library Cataloguing in Publication Data

A record for this title is available from the British Library

Library of Congress Cataloguing in Publication Data

A record for this title is available from the Library of Congress

ISBN 0 7506 3605 X

Typeset by Laser Words, Madras, India
Printed and bound in Great Britain by Bath Press

Foreword

It is now commonplace to say that the growth of knowledge of every sort is exponential and that keeping abreast with it becomes more difficult every year. As a result research can often become re-research, discovering again what is already described somewhere in the literature, perhaps in a little-known journal or little-read book.

In many fields the worst result is a waste of resources on repetitive work but when safety is concerned the lack of access to available knowledge can have tragic results. Those of us who have been working in the safety field for many years have seen the same accidents repeat themselves with distressing regularity. We welcome, therefore, every attempt to bring together scattered information on any aspect of safety and make it readily accessible.

In preparing the first four editions of this book Leslie Bretherick, almost unaided, produced a masterly summary of available information on reactive chemicals and their reactions. It was a remarkable achievement for one man, especially when we remember that the earlier editions were prepared in his spare time! Now that he has retired a team of editors, led by Peter Urben, has carried on the good work. They have increased the length of this edition by adding data on about 200 new compounds in Part 1, making the total nearly 5000, and 30 new groups in Part 2, making the total about 650. Leslie's high standard has been maintained.

It is a tribute to Leslie that very little of the new information in the 5th and this edition is old stuff that he missed; almost all comes from new publications. The entries on new groups range from acrylates to yeast passing through drums, fumes, mists and pnictides on the way. Familiar accidents continue to occur and new entries have been added on well-known hazards, such as nitric acid and azides.

In reviewing an earlier edition, I compared Leslie with those immortals, Beilstein and Perry, and said that he would become as well-known. Today I would go further and say that he has done more. He, and his successors, have not just made knowledge available but have saved lives and prevented injuries and damage and will continue to do so for many years.

Knowledge is effective only if it is used. Many coffee table books and cookery books are used to decorate the living room and kitchen rather than add to our knowledge. Some engineers are known to buy books to impress their visitors or

perhaps with good intentions that are never fulfilled. As an author I don't complain but I hope your copy of this book does not undergo this fate. Look up the entries on all the compounds and groups of compounds you use or are thinking of using (now, not after your accident). Also, look up the entries on substances you used in the past and think of what might have happened.

The editors have done their bit; the publishers have done their bit by making the data available in book form and on CD-ROM. It is now up to you. As a bonus you will find the data fascinating to browse through, you will come across many facts that you never knew before, or had forgotten, and you will be amused by the deadpan humour of a few entries such as those on air, environmentalism, safety literature, sunspots and superiors.

TREVOR KLETZ

April 1999

Preface to the Sixth Edition

Bretherick's Handbook remains broadly similar to the previous editions but older readers will notice some changes. There are, of course, some hundreds of additional entries and much supplementary information in existing entries. This is the second edition for which I have been responsible and readers will still regret the absence of Leslie Bretherick, who had to withdraw from compilation because of worsening sight but remains a support and stay. The bulk is still his work, which is an indication of his immense labours laying the foundations, when accidents were less often reported and databases harder to compile than they now are. The present editor and his assistants have a far easier task continuing the work.

The change in the cyclic structures, now drawn in the more contemporary notation for which Hampden Data Services are to be thanked continues. The book is also now available as an electronic database, with all the improved ease of searching for cross-references or related materials which that brings. In future, it is proposed to supplement this rather more frequently than the new editions of the book will appear.

The present edition includes the literature until the end of 1998. There are few organisational changes from Leslie Bretherick's model, although no new references to safety data sheets are given. There are ever proliferating series of these, it would be impossible to read all and invidious to distinguish some, while others appear to have been compiled by (mis)information (ne)scientists in the library rather than by chemists skilled and experienced in lab and plant.

Once again, we request all users to inform us of any hazards of which they are aware and of which we are not, as also of any errors they find (regretfully, I must admit that some will certainly have escaped detection). Thanks are given to all those who have contributed to this and previous editions.

I have been valiantly assisted in the compilation and evaluation of data by Dr Martin Pitt of the University of Sheffield. I am also indebted to the staff of Butterworth-Heinemann and Hampden Data Services for their ability to make my files manifest as book and database.

My employers, Courtaulds plc, (recently subsumed into Akzo-Nobel) have generously allowed me time to undertake this work which, while both yet existed,

benefited greatly from their library and from the Courtaulds Library at the University of Warwick. But, above all, thanks are due to Leslie Bretherick, not only for assistance and counsel but because, without him, the whole would be inconceivable. We again wish him a long and active (semi)retirement.

P. URBEN

March 1999

Preface to the First Edition

Although I had been aware during most of my career as a preparative chemist of a general lack of information relevant to the reactive hazards associated with the use of chemicals, the realisation that this book needed to be compiled came soon after my reading *Chemistry & Industry* for June 6th, 1964. This issue contained an account of an unexpected laboratory explosion involving chromium trioxide and acetic anhydride, a combination which I knew to be extremely hazardous from close personal experience 16 years previously.

This hazard had received wide publicity in the same journal in 1948, but during the intervening years had apparently lapsed into relative obscurity. It was then clear that currently existing arrangements for communicating 'well-known' reactive chemical hazards to practising chemists and students were largely inadequate. I resolved to try to meet this obvious need for a single source of information with a logically arranged compilation of available material. After a preliminary assessment of the overall problems involved, work began in late 1964.

By late 1971, so much information had been uncovered but remained to be processed that it was apparent that the compilation would never be finished on the spare-time basis then being used. Fortunately I then gained the support of my employers, the British Petroleum Company, Ltd., and have now been able to complete this compilation as a supporting research objective since January 1972.

The detailed form of presentation adopted has evolved steadily since 1964 to meet the dual needs for information on reactive chemical hazards in both specific and general terms, and the conflicting practical requirements of completeness and brevity. A comprehensive explanation of how this has been attempted, with suggestions on using this Handbook to best advantage, is given in the Introduction.

In an attempt to widen the scope of this work, unpublished information has been sought from many sources, both by published appeals and correspondence. In this latter area, the contribution made by a friend, the late Mr A. Kruk-Schuster, of Laboratory Chemicals Disposal Company, Ltd., Billericay, has been outstanding. During 1965–1968 his literature work and global letter campaign to 2000 University chemistry departments and industrial institutions yielded some 300 contributions.

The coverage attempted in this Handbook is wide, but is certainly incomplete because of the difficulties in retrieving relevant information from original literature when it does not appear in the indices of either primary or abstract journals. Details of such new material known to users of this Handbook and within the scope given in the Introduction will be welcomed for inclusion in supplementary or revised editions of this work.

L.B

October 1974

Contents

Volume 1

INTRODUCTION

Aims of the Handbook	xi
Scope and Source Coverage	xi
General Arrangement	xii
Specific Chemical Entries (Volume 1)	xiii
Grouping of Reactants	xiv
General Group Entries (Volume 2)	xv
Nomenclature	xv
Cross-reference System	xvii
Information Content of Individual Entries	xvii

REACTIVE CHEMICAL HAZARDS

Basics	xix
Kinetic Factors	xix
Adiabatic Systems	xxii
Reactivity vs. Composition and Structure	xxii
Reaction Mixtures	xxiii
Protective Measures	xxiv

SPECIFIC CHEMICALS

(Elements and compounds arranged in formula order)

APPENDIX 1 Source Title Abbreviations used in Handbook References	1927
APPENDIX 2 Tabulated Fire-related Data	1937
APPENDIX 3 Glossary of Abbreviations and Technical Terms	1947
APPENDIX 4 Index of Chemical Names and Serial Numbers used as Titles in Volume 1	1951
APPENDIX 5 Index of CAS Registry Numbers and Text Serial Numbers	2081

Introduction

**THIS SHOULD BE READ THROUGH CAREFULLY
TO GAIN FULL BENEFIT FROM WHAT FOLLOWS**

Aims of the Handbook

This compilation has been prepared and revised to give access to a wide and up-to-date selection of documented information to research students, practising chemists, safety officers and others concerned with the safe handling and use of reactive chemicals. This will allow ready assessment of the likely potential for reaction hazards which may be associated with an existing or proposed chemical compound or reaction system.

A secondary, longer-term purpose is to present the information in away which will, as far as possible, bring out the causes of, and interrelationships between, apparently disconnected facts and incidents. This is designed to encourage an increased awareness of potential chemical reactivity hazards in school, college and university teaching laboratories, and to help to dispel the relative ignorance of such matters which is still in evidence in this important area of safety training during the formative years of technical education.

Others involved in a more general way with the storage, handling, packing, transport and distribution of chemicals, or emergencies related thereto, are likely to find information of relevance to their activities.

Scope and source coverage

This Handbook includes all information which had become available to the editor by January 1999 on the reactivity hazards of individual elements or compounds, either alone or in combination. Appropriate source references are included to give access to more expansive information than that compressed into the necessarily abbreviated text entries.

A wide variety of possible sources of published information has been scanned to ensure maximum coverage. Primary sources have largely been restricted to journals known to favour or specialise in publication of safety matters, and the textbook series specialising in synthetic and preparative procedures.

Secondary sources have been a fairly wide variety of both specialised and general textbooks and encyclopaedic collections (notably those of Mellor, Sidgwick, Pascal and Bailar in the inorganic area, Houben-Weyl in the organic and

organometallic areas, and Kirk-Othmer in the industrial area). Section 50 of *Chemical Abstracts*, the CAS selection *Chemical Hazards, Health, & Safety*, the *Universities' Safety Association Safety News*, the CIA CISHC *Chemical Safety Summary*, (publication of which ceased in 1986 after 56 years), and the IChE *Loss Prevention Bulletin* have been rich sources, together with the more recent RSC *Laboratory Hazards Bulletin* and *Chemical Hazards in Industry*. Additionally, various safety manuals, compilations, summaries, data sheets and case histories have been used, and fuller details of all the sources used are set out in Appendix 1. References in the text to textbooks are characterised by absence of the author's initials after the surname.

More recently, some reports have been picked from the Internet, when two of the three following conditions obtained: the editor finds the report credible; it represents a hazard not already present in the handbook; or the source is authoritative. Information on toxic hazards has been specifically excluded because it is available elsewhere in many well-ordered and readily usable forms.

However, it should be remembered that many of the compounds included in this Handbook show high reactivity of one sort or another toward other materials, so may in general terms be expected to be reactive even in brief contact with animal organisms or tissue (including yours), with possible toxic effects, either acute or chronic. Also, no attempt has been made to include details of all flammable or combustible materials capable of burning explosively when mixed with air and ignited, nor of any incidents related to this most frequent cause of accidents, such information again being available elsewhere.

However, to focus attention on the potential hazards always associated with the use of flammable and especially highly flammable substances, some 560 gases and liquids with flash points below 25°C and/or autoignition temperature below 225°C have been included in the text, their names prefixed with a dagger. The numerical values of the fire hazard-related properties of flashpoint, autoignition temperature and explosive (flammability) limits in air where known are given in the tabular Appendix 2. Those elements or compounds which ignite on exposure to air are included in the text, but not in the Table.

General arrangement

The information presented on reactive hazards is of two main types, specific or general, and these types of information have been arranged differently in their respective separate volumes 1 and 2.

FOR CROSS REFERENCES IN CAPITALS, PAGE NUMBERS REFER TO VOLUME 2.

Specific information on instability of individual chemical compounds, and on hazardous interactions of elements and/or compounds, is contained in the main formula-based Volume 1 of the Handbook. For an example of an unstable compound,

see Ethyl perchlorate

For an example of a hazardous interaction between 2 compounds,

see Nitric acid: Acetone

or 2 separate examples involving the same compound,

see Nitric acid: Acetone, or: Ethanol
and one involving 3 compounds,

see Hydrogen peroxide: Nitric acid, Thiourea

General information relating to classes or groups of elements or compounds possessing similar structural or hazardous characteristics is contained in the smaller alphabetically based Volume 2.

See ACYL NITRATES

PYROPHORIC METALS

References in the text to these general classes or groups of materials is always in small capitals to differentiate them from references to specific chemicals, the names of which are given in normal roman typeface.

Some individual materials of variable composition (substances) and materials which cannot conveniently be formulated and placed in Volume 1 are also included in this general section.

See BLEACHING POWDER, CELLULOSE NITRATE

Both theoretical and practical hazard topics, some indirectly related to the main theme of this book, are also included.

See DISPOSAL, EXPLOSIBILITY

GAS CYLINDERS, OXYGEN ENRICHMENT

Several topics which bring together incidents involving a common physical cause or effect but different types of chemicals are now included in Volume 2.

See CATALYTIC IMPURITY INCIDENTS

GAS EVOLUTION INCIDENTS

Specific chemical entries (Volume 1)

A single unstable compound of known composition is placed in the main first volume and is located on the basis of its empirical molecular formula expressed in the Hill system used by *Chemical Abstracts* (C and H if present, then all other element symbols alphabetically). The use of this indexing basis permits a compound to be located if its structure can be drawn, irrespective of whether a valid name is known for it. A representation of the structure of each compound is given on the third bold title line while the name of the compound appears as the first bold title line. References to the information source are given, followed by a statement of the observed hazard, with any relevant explanation. Cross-reference to similar compounds, often in a group entry, completes the entry. *See* Trifluoroacetyl nitrite p. 244.

Where two or more elements or compounds are involved in a reactive hazard, and an intermediate or product of reaction is identifiable as being responsible for the hazard, both reacting substances are normally cross-referred to the identified product. The well-known reaction of ammonia and iodine to give explosive nitrogen triiodide-ammonia is an example of this type. The two entries

Ammonia: Halogens

Iodine: Ammonia

are referred back to the main entry under the identified material

Nitrogen triiodide-ammonia

No attempt has been made, however, to list all combinations of reactants which can lead to the formation of a particular main entry compound.

In a multi-reactant system where no identification of an unstable product was possible, one of the reactants had to be selected as primary reactant to prepare and index the main entry, with the other material(s) as secondary reactant(s). No strictly logical basis of choice for this is obvious.

However, it emerged during the compilation phase that most two component reaction hazard systems of this type involve a fairly obvious oxidant material as one of the reactants. Where this situation was recognised, the oxidant has normally been selected as primary (indexing) reactant, with the other as secondary reactant, following the colon.

See Potassium permanganate: Acetic acid, etc.

In the markedly fewer cases where an obvious reductant has been involved as one reactant, that was normally selected as primary reactant.

See Lithium tetrahydroaluminate: 3,5-Dibromocyclopentene

In the relatively few cases where neither (or none) of the reactants can be recognised as an oxidant or reductant, the choice was made which appeared to give the more informative main entry text.

See Chloroform: Acetone, etc.

Where some hazard has been noted during the preparation of a specific compound, but without it being possible to identify a specific cause, an entry for that compound states 'Preparative hazard', and back-refers to the reactants involved in the preparation.

See Sulfur dioxide

Occasionally, departures from these considerations have been made where such action appeared advantageous in bringing out a relationship between formally unrelated compounds or hazards. In all multi-component cases, however, the secondary reactants (except air and water) appear as formula entries back-referred to the main entry text, so that the latter is accessible from either primary or secondary reactants.

See Dimethyl sulfoxide: Acyl halides (main entry)

Acetyl chloride: Dimethyl sulfoxide (back reference)

Grouping of Reactants

There are advantages to be gained in grouping together elements or compounds showing similar structure or reactivity, because this tends to bring out the relationships between structure and activity more clearly than separate treatment. This course has been adopted widely for primary reactants (see next heading), and for secondary reactants where one primary reactant has been involved separately with a large number of secondary materials. Where possible, the latter have been collected together under a suitable general group title indicative of the composition or characteristics of those materials.

See Chlorine: Hydrocarbons

Hydrogen peroxide: Metals, Metal oxides, Metal salts

Hydrogen sulfide: Oxidants

This arrangement means, however, that some practice will be necessary on the user's part in deciding into what group an individual secondary reactant falls before the longer-term advantages of the groupings become apparent. The formal group titles listed in Volume 2, Appendix 3, and classified in Appendix 4, will be of use in this connection. However, it should be noted that sometimes informal group titles are used which do not appear in these Appendices.

General group entries (Volume 2)

In some cases literature references relating to well-defined groups of hazardous compounds or to hazard topics have been found, and these are given, with a condensed version of relevant information at the beginning of the topic or group entry, under a suitable bold title, the latter being arranged in alphabetical order in Volume 2.

Cross references to related group or sub-group entries are also included, with a group list of the names and serial (not page) numbers of the chemicals appearing in Volume 1 which lie within the structural or functional scope of the group entry title. Compounds which are closely similar to, but not in strict conformity with, the group definition are indicated by a prefixed asterisk.

The group entries thus serve as sub-indexes for each structurally based group of hazardous compounds. Conversely, each individual compound entry is back-referred to the group entry, and thence to all its strict structural analogues and related congeners included in Volume 1 of this Handbook. Note that these group lists of chemicals are now in alphabetical (not formula) order, and give the serial-number (not page number) for the chemical.

These features should be useful in attempts to estimate the stability or reactivity of a compound or reaction system which does not appear in this Handbook. The effects on stability or reactivity of changes in the molecular structure to which the destabilising or reactive group(s) is attached are in some cases discussed in the group entry. Otherwise such information may be gained from comparison of the information available from the individual compound entries listed collectively (now in alphabetical order, with serial number) in the group entry.

Care is, however, necessary in extrapolating from the described properties of compounds to others in which the user of this Handbook may be interested. Due allowance must be made for changes in elemental reactivity up or down the columns of the Periodic Table, and for the effects of variation in chain length, branching and point of group-attachment in organic systems. Purity of materials, possible catalytic effects (positive or negative) of impurities, and scale of operations may all have a direct bearing upon a particular reaction rate. These and other related matters are dealt with in more detail in the following Introductory Chapter.

Nomenclature

With the direct encouragement and assistance of the Publishers, an attempt has been made to use chemical names which conform to recent recommendations of IUPAC. While this has not been an essential part of the compilation, because

each title name has the corresponding structural and molecular formula adjacent, it seems none the less desirable to minimise possible confusion by adopting the unambiguous system of nomenclature presented in the IUPAC publications.

Where the IUPAC name for a compound is very different from a previously used recent trivial name, the latter is included as a synonym in parentheses (and in single quotes where no longer an acceptable name). Generally, retained trivial names have not been used as main entry titles, but they have been used occasionally in the entry texts. Rarely, on the grounds of brevity, names not conforming strictly to IUPAC principles but recommended for chemicals used in industry in BS 2474: 1983 have been used. The prefix mixo-, to represent the mixtures of isomers sometimes used as industrial materials, is a case in point.

Some of the rigidly systematic names selected by the Association for Science Education for their nomenclature list in 1985 from the IUPAC possibilities, and some of the systematic indexing names used by *Chemical Abstracts* since 1972, are given as synonyms in the Index of Chemical Names (Appendix 4). This should assist those coming into industry and research with a command of those nomenclature systems but who may be unfamiliar with the current variety of names used for chemicals. The inclusion where possible of the CAS Registry Number for each title compound should now simplify the clarification of any chemical name or synonym problems, by reference to the Registry Handbook or other CAS source.

In connection with the group titles adopted for the alphabetically ordered Volume 2, it has been necessary in some cases to devise groupnames (particularly in the inorganic field) to indicate in a very general way the chemical structures involved in various classes, groups or sub-groups of compounds.

For this purpose, all elements have been considered either as METALS or NON-METALS and of the latter, HALOGENS, HYDROGEN, NITROGEN, OXYGEN, and SULFUR were selected as specially important. Group names have then been coined from suitable combinations of these, such as the simple

METAL OXIDES, NON-METAL SULFIDES,
N-HALOGEN COMPOUNDS, NON-METAL HYBRIDES,
METAL NON-METALLIDES, COMPLEX HYBRIDES

or the more complex

METAL OXOHALOGENATES
AMMINECHROMIUM PEROXOCOMPLEXES
OXOSALTS OF NITROGENOUS BASES
METALOXONON-METALLATES

Organic group entries are fairly conventional, such as

HALOALKENES
NITROARL COMPOUNDS
DIAZONIUM SALTS

Where necessary, such group names are explained in the appropriate group entry, of which a full listing is given in Volume 2, Appendix 3, and a classified listing in Appendix 4.

Cross reference system

The cross-reference system adopted in this Handbook plays a large part in providing maximum access to, and use of, the rather heterogeneous collection of information herein. The significance of the five types of cross-reference which have been used is as follows.

See . . . refers to a directly related item.

See also . . . refers to an indirectly related item.

See other . . . refers to listed strict analogues of the compound etc.

See related . . . refers to listed related compounds (congeners) or groups not strictly analogous structurally.

See entry . . . points to a, or the relevant, reference in Volume 2.

Information content of individual entries

A conscious effort has been made throughout this compilation to exclude all fringe information not directly relevant to the involvement of chemical reactivity in the various incidents or observations, with just enough detail present to allow the reader to judge the relevance or otherwise of the quoted reference(s) to his or her particular reactivity problems or interests.

It must be stressed that this book can do no more than to serve as a guide to much more detailed information available via the quoted references. It cannot relieve the student, the chemist and their supervisors of their moral and now legal obligation to themselves and to their co-workers, to equip themselves with the fullest possible information from the technical literature resources which are widely available, *before* attempting any experimental work with materials known, or suspected, to be hazardous or potentially so. It could be impossible for you *after* the event.

THE ABSENCE OF A MATERIAL OR A COMBINATION OF MATERIALS FROM THIS HANDBOOK CANNOT BE TAKEN TO IMPLY THAT NO HAZARD EXISTS. LOOK THEN FOR ANALOGOUS MATERIALS USING THE GROUP ENTRY SYSTEM AND THE INDEXES THERETO.

One aspect which, although it is excluded from most entry texts, is nevertheless of vital importance, is that of the potential for damage, injury or death associated with the various materials and reaction systems dealt with in this Handbook.

Though some of the incidents have involved little or no damage (*see* CAN OF BEANS), others have involved personal injuries, often of unexpected severity (*see* SODIUM PRESS), and material damage is often immense. For example, the incident given under Perchloric acid: Cellulose derivatives, (reference 1) involved damage to 116 buildings and a loss approaching \$ 3M at 1947 values. The death-toll associated with reactive chemical hazards has ranged from 1 or 2 (*see* Tetrafluoroethylene: Iodine pentafluoride) to some 600 with 2000 injured in the incident at Oppau in 1921 (*see* Ammonium nitrate, reference 4), and now to several thousand

with more than 100,000 injured by methyl isocyanate fumes at Bhopal in 1984 (reference 7).

This sometimes vast potential for destruction again emphasises the need to gain the maximum of detailed knowledge *before* starting to use an unfamiliar chemical or reaction system.

Reactive Chemical Hazards

CROSS REFERENCES IN CAPITALS REFER TO PAGE NUMBERS IN VOLUME 2.

This introductory chapter seeks to present an overview of the complex subject of reactive chemical hazards, drawing attention to the underlying principles and to some practical aspects of minimising such hazards. It also serves in some measure to correlate some of the topic entries in the alphabetically arranged Volume 2 of the Handbook.

Basics

All chemical reactions implicitly involve energy changes (energy of activation + energy of reaction), for these are the driving force. The majority of reactions liberate energy as heat (occasionally as light or sound) and are termed exothermic. In a minority of reactions, energy is absorbed into the products, when both the reaction and its products are described as endothermic.

All reactive hazards involve the release of energy in quantities or at rates too high to be absorbed by the immediate environment of the reacting system, and material damage results. The source of the energy may be an exothermic multi-component reaction, or the exothermic decomposition of a single unstable (often endothermic) compound.

All measures to minimise the possibility of occurrence of reactive chemical hazards are therefore directed at controlling the extent and rate of release of energy in a reacting system. In an industrial context, such measures are central to modern chemical engineering practice. Some of the factors which contribute to the possibility of excessive energy release, and appropriate means for their control, are now outlined briefly, with references to examples in the text.

Kinetic Factors

The rate of an exothermic chemical reaction determines the rate of energy release. so factors which affect reaction kinetics are important in relation to possible reaction hazards. The effects of proportions and concentrations of reactants upon reaction rate are governed by the Law of Mass Action, and there are many examples where changes in proportion and/or concentration of reagents have transformed an

established uneventful procedure into a violent incident. For examples of the effect of increase in proportion,

see 2-Chloronitrobenzene: Ammonia

Sodium 4-nitrophenoxide

For the effect of increase in concentration upon reaction velocity,

see Dimethyl sulfate: Ammonia

Nitrobenzene: Alkali (reference 2)

The effects of catalysts (which effectively reduce the energy of activation), either intentional or unsuspected, is also relevant in this context. Increase in the concentration of a catalyst (normally used at 1–2%) may have a dramatic effect on reaction velocity.

See Trifluoromethanesulfonic acid: Acyl chlorides, etc.

2-Nitroanisole: Hydrogen

HYDROGEN CATALYSTS

The presence of an unsuspected contaminant or catalytic impurity may affect the velocity or change the course of reaction. For several examples,

See CATALYTIC IMPURITY INCIDENTS

In the same context, but in opposite sense, the presence of inhibitors (negative catalysts, increasing energy of activation) may seriously interfere with the smooth progress of a reaction. An inhibitor may initiate an induction period which can lead to problems in establishing and controlling a desired reaction. For further details and examples,

See INDUCTION PERIOD INCIDENTS

Undoubtedly the most important factor affecting reaction rates is that of temperature. It follows from the Arrhenius equation that the rate of reaction will increase exponentially with temperature. Practically, it is found that an increase of 10°C in reaction temperature often doubles or trebles the reaction velocity.

Because most reactions are exothermic, they will tend to accelerate as reaction proceeds unless the available cooling capacity is sufficient to prevent rise in temperature. Note that the exponential temperature effect accelerating the reaction will exceed the (usually) linear effect of falling reactant concentration in decelerating the reaction. When the exotherm is large and cooling capacity is inadequate, the resulting accelerating reaction may proceed to the point of loss of control (runaway), and decomposition, fire or explosion may ensue.

The great majority of incidents described in the text may be attributed to this primary cause of thermal runaway reactions. The scale of the damage produced is related directly to the size, and more particularly to the rate, of energy release.

See RUNAWAY REACTIONS

Reactions at high pressure may be exceptionally hazardous owing to the enhanced kinetic energy content of the system.

See HIGH-PRESSURE REACTION TECHNIQUES

Although detailed consideration of explosions is outside the scope of this Handbook, three levels of intensity of explosion (i.e. rates of fast energy release) can be discerned and roughly equated to the material damage potential.

Deflagration involves combustion of a material, usually in presence of air. In a normal liquid pool fire, combustion in an open situation will normally proceed

without explosion. Mixtures of gases or vapours with air within the explosive limits which are subsequently ignited will burn at normal flame velocity (a few m/s) to produce a 'soft' explosion, with minor material damage, often limited to scorching by the moving flame front. Injuries to personnel may well be more severe.

If the mixture (or a dust cloud) is confined, even if only by surface irregularities or local partial obstructions, significant pressure effects can occur. Fuel-air mixtures near to stoichiometric composition and closely confined will develop pressures of several bar within milliseconds, and material damage will be severe. Unconfined vapour explosions of large dimensions may involve higher flame velocities and significant pressure effects, as shown in the Flixborough disaster.

See DUST EXPLOSION INCIDENTS

PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

VAPOUR CLOUD EXPLOSIONS

Detonation is an extreme form of explosion where the propagation velocity becomes supersonic in gaseous, liquid or solid states. The temperatures and particularly pressures associated with detonation are higher by orders of magnitude than in deflagration. Energy release occurs in a few microseconds and the resulting shattering effects are characteristic of detonation. Deflagration may accelerate to detonation if the burning material and geometry of confinement are appropriate (endothermic compounds, long narrow vessels or pipelines).

See Acetylene (reference 9)

ENDOTHERMIC COMPOUNDS

EXPLOSIONS

UNIT PROCESS INCIDENTS

Factors of importance in preventing such thermal runaway reactions are mainly related to the control of reaction velocity and temperature within suitable limits. These may involve such considerations as adequate heating and particularly cooling capacity in both liquid and vapour phases of a reaction system; proportions of reactants and rates of addition (allowing for an induction period); use of solvents as diluents and to reduce viscosity of the reaction medium; adequate agitation and mixing in the reactor; control of reaction or distillation pressure; use of an inert atmosphere.

See AGITATION INCIDENTS

In some cases it is important not to overcool a reaction system, so that the energy of activation is maintained.

See Acetylene: Halogens (reference 1)

Adiabatic Systems

Because process heating is expensive, lagging is invariably applied to heated process vessels to minimise heat loss, particularly during long-term hot storage. Such adiabatic or near-adiabatic systems are potentially hazardous if materials of limited thermal stability, or which possess self-heating capability, are used in them. Insufficiently stabilised bulk-stored monomers come into the latter category.

See 1,2,4,5-Tetrachlorobenzene: Sodium hydroxide, Solvent

POLYMERISATION INCIDENTS

SELF-HEATING AND IGNITION INCIDENTS
THERMAL STABILITY OF REACTION MIXTURES
VIOLENT POLYMERISATION

Reactivity vs. Composition and Structure

The ability to predict reactivity and stability of chemical compounds from their composition and structure is as yet limited, so the ability accurately to foresee potential hazards during preparation, handling and processing of chemicals and their mixtures is also restricted. Although some considerable progress has been made in the use of computer programs to predict hazards, the best available approach for many practical purposes appears to be an initial appraisal based on analogy with, or extrapolation from, data for existing compounds and processes. This preliminary assessment should be supplemented with calorimetric instrumental examination, then bench-scale testing procedures for thermal stability applied to realistic reaction mixtures and processing conditions. A wide range of equipment and techniques is now available for this purpose.

See ACCELERATING RATE CALORIMETRY

ASSESSMENT OF REACTIVE CHEMICAL HAZARDS
COMPUTATION OF REACTIVE CHEMICAL HAZARDS
DIFFERENTIAL SCANNING CALORIMETRY
DIFFERENTIAL THERMAL ANALYSIS
MAXIMUM REACTION HEAT
REACTION SAFETY CALORIMETRY

It has long been recognised that instability in single compounds, or high reactivity in combinations of different materials, is often associated with particular groupings of atoms or other features of molecular structure, such as high proportions or local concentrations of oxygen or nitrogen. Full details of such features associated with explosive instability are collected under the heading EXPLOSIBILITY.

An approximate indication of likely instability in a compound may be gained from inspection of the empirical molecular formula to establish stoichiometry.

See HIGH-NITROGEN COMPOUNDS

OXYGEN BALANCE

Endothermic compounds, formed as the energy-rich products of endothermic reactions, are thermodynamically unstable and may be liable to energetic decomposition with low energy of activation.

See ENDOTHERMIC COMPOUNDS

Reaction Mixtures

So far as reactivity between different compounds is concerned, some subdivision can be made on the basis of the chemical types involved. Oxidants (electron sinks) are undoubtedly the most common chemical type to be involved in hazardous incidents, the other components functioning as fuels or other electron sources. Air (21% oxygen) is the most widely dispersed oxidant, and air-reactivity may lead to either short- or long-term hazards.

Where reactivity of a compound is very high, oxidation may proceed so fast in air that ignition occurs.

See PYROPHORIC MATERIALS

Slow reaction with air may lead to the longer-term hazard of peroxide formation.

See AUTOXIDATION

PEROXIDATION INCIDENTS

PEROXIDATION IN SOLVENTS

PEROXIDISABLE COMPOUNDS

Oxidants more concentrated than air are of greater hazard potential, and the extent of involvement of the common oxidants

Perchloric acid

Chlorine

Nitric acid

Hydrogen peroxide

Sulfuric acid

METAL CHLORATES

may be judged from the large number of incidents in the text involving each of them, as well as other OXIDANTS, p. 271.

At the practical level, experimental oxidation reactions should be conducted to maintain in the reacting system a minimum oxygen balance consistent with other processing requirements. This may involve adding the oxidant slowly with appropriate mixing and cooling to the other reaction materials to maintain the minimum effective concentration of oxidant for the particular reaction. It will be essential to determine by a suitable diagnostic procedure that the desired reaction has become established, to prevent build-up of unused oxidant and a possible approach to the oxygen balance point.

See OXYGEN BALANCE

Reducants (rich electron sources) in conjunction with reducible materials (electron acceptors) feature rather less frequently than oxidants in hazardous incidents.

See REDUCANTS

Interaction of potent oxidants and reducants is invariably highly energetic and of high hazard potential.

See Dibenzoyl peroxide: Lithium tetrahydroaluminate

Hydrazine: Oxidants

REDOX REACTIONS

ROCKET PROPELLANTS

Similar considerations apply to those compounds which contain both oxidising and reducing functions in the same molecular structure.

See REDOX COMPOUNDS

Water is, after air, one of the most common reagents likely to come into contact with reactive materials, and several classes of compounds will react violently, particularly with restricted amounts of water.

See WATER-REACTIVE COMPOUNDS

Most of the above has been written with deliberate processing conditions in mind, but it must be remembered that the same considerations will apply, and

perhaps to a greater degree, under the uncontrolled reaction conditions prevailing when accidental contact of reactive chemicals occurs in storage or transit.

Adequate planning is therefore necessary in storage arrangements to segregate oxidants from fuels and reductants, and fuels and combustible materials from compressed gases and water-reactive compounds. This will minimise the possibility of accidental contact and violent reaction arising from faulty containers or handling operations, and will prevent intractable problems in the event of fire in the storage areas.

See SAFE STORAGE OF CHEMICALS

Unexpected sources of ignition may lead to ignition of flammable materials during chemical processing or handling operations.

See FRICTIONAL IGNITION OF GASES

IGNITION SOURCES

SELF-HEATING AND IGNITION INCIDENTS

STATIC INITIATION INCIDENTS

Protective Measures

The need to provide protective measures will be directly related to the level of potential hazards which may be assessed from the procedures outlined above. Measures concerned with reaction control are frequently mentioned in the following text, but details of techniques and equipment for personal protection, though usually excluded from the scope of this work, are obviously of great importance.

Careful attention to such detail is necessary as a second line of defence against the effects of reactive hazards. The level of protection considered necessary may range from the essential and absolute minimum of effective eye protection, via the safety screen, fume cupboard or enclosed reactor, up to the ultimate of a remotely controlled and blast-resistant isolation cell (usually for high-pressure operations). In the absence of facilities appropriate to the assessed level of hazard, operations must be deferred until such facilities are available.

Volume 1

Specific Chemicals

(Elements and Compounds arranged in formula order)

EXPLANATORY NOTES

This volume gives detailed information on the hazardous properties of individual chemicals, either alone or in combination with other compounds. The items are arranged in order of the empirical formula (at right of second bold title line) which corresponds to the chemical name, or a synonym within parentheses, used as the first line bold title of each main entry; (nomenclature is now rather promiscuous since the systematisers have contrived to give many materials two or even three new names since the first edition was published, while the name used where chemicals are handled, as opposed to in lecture rooms, will be something else again). The 3 part number within square brackets at the left of the second title line is the CAS registry number, now being widely used to provide a reliable basis for establishing equivalence between differing chemical names and trade names for the same chemical compound (but note that one compound, within the terms of this work, may have numerous CAS numbers by virtue of isotopic composition, undefined stereo- and regio-chemistry or variant solvation levels). Lack of content within the square brackets indicates that a registry number has not yet been located, and (ion) after the number indicates that the main ion only has been located, rather than the specific title salt. Where possible, a linear or graphical representation of the structure of the title compound is given at the centre of the third title line, otherwise the reader is referred to one of the several lettered pages preceding on which the corresponding cyclic structural formula is set out.

A † prefixed to the chemical name indicates the existence of tabulated information on fire-related properties in Appendix 2. The † prefix is also appended to the entry (and any synonym) in the index in Appendix 4 of the chemicals appearing as title lines. Immediately under the title lines some references to sources of general safety-related data concerning use and handling precautions for the title chemical are given. The references to the series of *MCA Safety Data Sheets* are given in parentheses because the whole series was withdrawn in 1980, apparently on grounds other than obsolescence of the technical content. Since these data sheets are no longer available, alternative references are given where possible to the Data Sheets available from the National Safety Council (*NSC*), Chicago; the Fire Protection Association (*FPA*), London; to the appropriate page of 'Handling Chemicals Safely 1980' (*HCS 1980*), published in Holland; or to the Laboratory Hazard

Data Sheet series published by the Royal Society of Chemistry (*RSC*), now in Cambridge.

No new data sheets have been included since 1990, since distinction between the proliferation of sources would be invidious, and even supposedly authoritative bodies are putting out sheets which have evidently been compiled by (mis)information scientists at their computers, unchecked by anyone who has ever seen, smelt, or handled, the material in question. The first reference(s) and data given under the title lines refer to the hazards of the title material alone, or in the presence of air, unless stated otherwise. Where other (secondary) chemicals are involved with the title compound in a reactive incident, the name(s) follows in roman characters under the bold title entry. As in previous editions of this Handbook, where these secondary chemicals are described in group terms (e.g. Polynitroaryl compounds), reference to the alphabetical group entries now in Volume 2 may suggest other analogous possibilities of hazards. References to original or abstract literature then follow, and sufficient of the relevant information content is given to allow a general picture of the nature and degree of hazard to be seen.

Two features relevant to entries for pairs of reactive chemicals arise from the work of Prof. T. Yoshida in developing a method for the calculation of maximum reaction heats (MRH) possible for binary (or ternary) mixtures of chemicals, and the publication of his tabulated results. Where available for combinations existing in this text, these data are given opposite the name of the secondary chemical in the form MRH 2.9/22. This means that the calculated reaction heat is maximal at 2.9 kJ/g in a mixture containing 22% wt of the secondary reactant with 78% of the main (bold title) compound. The second feature is the inclusion of the secondary entry 'Other reactants' under which the extent of the information available in Yoshida's book for some 240 title compounds is given. More detail on the origin of these figures is given in Volume 2 under the entry MAXIMUM REACTION HEAT.

All temperatures in the text are expressed in degrees Celsius; pressures in bars, mbars or Pa; volumes in m³, litres or ml; and energy as joules, kJ or MJ. Where appropriate, attention is drawn to closely similar or related materials or events by *See* or *See also* cross-references. Finally, if a title compound is a member of one of the general classes or groups in Volume 2, it is related to those by a *See other* cross-reference. If the compound is not strictly classifiable, a *See related* cross-reference establishes a less direct link to the group compound index lists in Volume 2, such compounds being prefixed in the lists by an asterisk. In relatively few cases, literature references (or further references) for individual compounds are in the alphabetical entries in Volume 2, and a *See entry* cross-reference leads to that entry with the literature reference. An alphabetical index of the chemical names used as bold titles in Volume 1, together with synonyms, is given in Appendix 4.

Details of corrections of typographical or factual errors, or of further items for inclusion in the text, will be welcomed, and a page which can be photocopied for this purpose will be found at the back of the book.

0001. Silver

[7440-22-4]

Ag

Ag

Acetylenic compounds

MRH Acetylene 8.70/99+

See ACETYLENIC COMPOUNDS

Aziridine

See Aziridine: Silver

Bromine azide

See Bromine azide

3-Bromopropyne

See 3-Bromopropyne: Metals

Carboxylic acids

Koffolt, J. H., private comm., 1965

Silver is incompatible with oxalic or tartaric acids, since the silver salts decompose on heating. Silver oxalate explodes at 140°C, and silver tartrate loses carbon dioxide.

See other METAL OXALATES

Chlorine trifluoride

MRH 1.42/36

See Chlorine trifluoride: Metals

Copper, Ethylene glycol

See Ethylene glycol: Silvered copper wire

Electrolytes, Zinc

Britz, W. K. *et al.*, *Chem. Abs.*, 1975, **83**, 150293

Causes of spontaneous combustion and other hazards of silver–zinc batteries were investigated.

Ethanol, Nitric acid

Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 334

Action of silver on nitric acid in presence of ethanol may form the readily detonable silver fulminate.

See Nitric acid: Alcohols

See also SILVER-CONTAINING EXPLOSIVES

Ethylene oxide

MRH 3.72/99+

See Ethylene oxide: Reference 4

Ethyl hydroperoxide

See Ethyl hydroperoxide: Silver

Hydrogen peroxide

MRH 1.59/99+

See Hydrogen peroxide: Metals

Iodoform

Grignard, 1935, Vol. 3, 320

In contact with finely divided (reduced) silver, incandescence occurs.

Other reactants

Yoshida, 1980, 103

MRH values for 7 combinations, largely with catalytically susceptible materials, are given.

Ozonides

See OZONIDES

Peroxomonosulfuric acid

See Peroxomonosulfuric acid: Catalysts

Peroxyformic acid

MRH 5.69/100

See Peroxyformic acid: Metals

See other METALS

0002. Silver–aluminium alloy

[11144-29-9]

Ag–Al

Ag–Al

1. Popov, E. I. *et al.*, *Chem. Abs.*, 1977, **87**, 205143

2. Popov, E. I. *et al.*, *Chem. Abs.*, 1980, **94**, 35622

Combustion and explosion hazards of the powdered alloy used in batteries were studied. Increase in silver content leads to higher values of the ignition temperature and COI [1,2].

See other ALLOYS, SILVER COMPOUNDS

0003. Silvered copper

[37218-25-0]

Ag–Cu

Ag–Cu

Ethylene glycol

See Ethylene glycol: Silvered copper wire

See related ALLOYS

0004. Silver–thorium alloy

[[12785-36-3] (1:2)]

Ag–Th

Ag–Th

See entry PYROPHORIC ALLOYS

0005. Silver tetrafluoroborate
[14104-20-2]



Preparative hazard

1. Meerwein, H. *et al.*, *Arch. Pharm.*, 1958, **291**, 541–544
2. Lemal, D. M. *et al.*, *Tetrahedron Lett.*, 1961, 776–777
3. Olah, G. A. *et al.*, *J. Inorg. Nucl. Chem.*, 1960, **14**, 295–296

Experimental directions must be followed exactly to prevent violent spontaneous explosions during preparation of the salt from silver oxide and boron trifluoride etherate in nitromethane, according to the earlier method [1]. The later method [3] is generally safer than that in [2].

See other SILVER COMPOUNDS

0006. Silver tetrafluorobromate
[35967-89-6]



See entry METAL POLYHALOHALOGENATES *See other* SILVER COMPOUNDS

0007. Silver bromate
[7783-89-3]



Other reactants

Yoshida, 1980, 133

MRH values for 16 combinations with oxidisable materials are given.

Sulfur compounds

MRH Sulfur 2.0/12

1. Taradoire, F., *Bull. Soc. Chim. Fr.*, 1945, **12**, 94–95
2. Pascal, 1960, Vol. 13.1, 1004

The bromate is a powerful oxidant, and unstable mixtures with sulfur ignite at 73–75°C, and with disulfur dibromide on contact [1]. Hydrogen sulfide ignites on contact with the bromate [2].

See other METAL OXOHALOGENATES, SILVER COMPOUNDS

0008. Silver chloride
[7783-90-6]



Aluminium

See Aluminium: Silver chloride

Ammonia

1. Mellor, 1941, Vol. 3, 382
2. Kauffmann, G. B., *J. Chem. Educ.*, 1977, **54**, 132
3. Ranganathan, S. *et al.*, *J. Chem. Educ.*, 1976, **53**, 347

Exposure of ammoniacal silver chloride solutions to air or heat produces a black crystalline deposit of 'fulminating silver', mainly silver nitride, with silver diimide and silver amide also possibly present [1]. Attention is drawn [2] to the possible explosion hazard in a method of recovering silver from the chloride by passing an ammoniacal solution of the chloride through an ion exchange column to separate the $\text{Ag}(\text{NH}_3)^+$ ion, prior to elution as the nitrate [3]. It is essential to avoid letting the ammoniacal solution stand for several hours, either alone or on the column [2].

See Silver nitride

See other METAL HALIDES, SILVER COMPOUNDS

0009. Silver azide chloride

[67880-13-1]



Frierson, W. J. *et al.*, *J. Amer. Chem. Soc.*, 1943, **65**, 1698

It is shock sensitive when dry.

See other METAL AZIDE HALIDES, SILVER COMPOUNDS

0010. Silver chlorite

[7783-91-7]



Alone, or Iodoalkanes

Levi, G. R., *Gazz. Chim. Ital.* [2], 1923, **53**, 40

The salt is impact-sensitive, cannot be finely ground, and explodes at 105°C. Attempts to react silver chlorite with iodo-methane or -ethane caused explosions, immediate in the absence of solvents, or delayed in their presence.

Hydrochloric acid, or Sulfur

Mellor, 1941, Vol. 2, 284

It explodes in contact with hydrochloric acid or on rubbing with sulfur.

Non-metals

Pascal, 1960, Vol. 16, 264

Finely divided carbon, sulfur or red phosphorus are oxidised violently by silver chlorite.

See other CHLORITE SALTS, SILVER COMPOUNDS

0011. Silver chlorate

[7783-92-8]



Sorbe, 1968, 126

An explosive compound and powerful oxidant.

Ethylene glycol MRH 2.68/17
See Ethylene glycol: Oxidants

Other reactants
Yoshida, 1980, 69
MRH values for 17 combinations, largely with oxidisable materials, are given.
See other METAL CHLORATES, SILVER COMPOUNDS

0012. Silver perchlorate
[7783-93-9]



1. Anon., *Angew. Chem. (Nachr.)*, 1962, **10**, 2
It melts without decomposition although the enthalpy of conversion to silver chloride and oxygen appears to be about -0.5 kJ/g. An explosion while grinding the salt (which had not been in contact with organic materials) has been reported [1].
A powerful oxidant.

Acetic acid MRH 2.80/22
Mellor, 1956, Vol. 2, Suppl.1, 616
The salt solvated with acetic acid is impact sensitive.
See Aromatic compounds, below

Alkynes, Mercury
Comyns, A. E. *et al.*, *J. Amer. Chem. Soc.*, 1957, **79**, 4324
Concentrated solutions of the perchlorate in 2-pentyne or 3-hexyne (complexes are formed) explode on contact with mercury.
See METAL ACETYLIDES

Aromatic compounds MRH Aniline 3.47/11, toluene 3.51/9
1. Sidgwick, 1950, 1234
2. Brinkley, S. R., *J. Amer. Chem. Soc.*, 1940, **62**, 3524
3. Peone, J. *et al.*, *Inorg. Synth.*, 1974, **15**, 69
4. Stull, 1977, 22
Silver perchlorate forms solid complexes with aniline, pyridine, toluene, benzene and many other aromatic hydrocarbons [1]. A sample of the benzene complex exploded violently on crushing in a mortar. The ethanol complex also exploded similarly, and unspecified perchlorates dissolved in organic solvents were observed to explode [2]. Solutions of the perchlorate in benzene are said to be dangerously explosive [3], but this may be in error for the solid benzene complex. The energy released on decomposition of the benzene complex has been calculated as 3.4 kJ/g, some 75% of that for TNT [4].

Carbon tetrachloride, Hydrochloric acid
491M, 1975, 368

Silver perchlorate and carbon tetrachloride in presence of a little hydrochloric acid produce trichloromethyl perchlorate, which explodes at 40°C.

See Trichloromethyl perchlorate

1,2-Diaminoethane

491M, 1975, 368

Dropwise addition of the amine to the salt led to an explosion (possibly initiated by heat liberated by complex formation).

Diethyl ether

1. Heim, F., *Angew. Chem.*, 1957, **69**, 274

After crystallisation from ether, the material exploded violently on crushing in a mortar. It had been considered stable previously, since it melts without decomposition [1].

Dimethyl sulfoxide

Ahrland, S. *et al.*, *Acta. Chem. Scand. A*, 1974, **28**, 825

The crystalline complex solvated with 2DMSO explodes with extreme violence if rubbed or scratched.

See Dimethyl sulfoxide: Metal oxosalts

Ethanol

MRH 3.30/13

See Aromatic compounds, above

Other reactants

Yoshida, 1980, 81

MRH values for 20 combinations with oxidisable materials are given.

1,4-Oxathiane

Barnes, J. C. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 103–113

The perchlorate forms complexes with 2, 3 or 4 mols of oxathiane which explode on heating.

Tetrachlorosilane, or Tetrabromosilane, or Titanium tetrachloride, and Diethyl ether

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

Reaction gives explosive volatile organic perchlorates, probably ethyl perchlorate.

See ALKYL PERCHLORATES

Tetrasulfur tetraimide

See Tetrasulfurtetraimide–silver(I) perchlorate

See other METAL PERCHLORATES, OXIDANTS, SOLVATED OXOSALT INCIDENTS

0013. Silver fluoride

[7775-41-9]

AgF

AgF

Calcium hydride

See Calcium hydride: Silver halides

Non-metals

Mellor, 1941, Vol. 3, 389

Boron reacts explosively when ground with silver fluoride; silicon reacts violently.

Titanium

Mellor, 1941, Vol. 7, 20

Interaction at 320°C is incandescent.

See other METAL HALIDES, SILVER COMPOUNDS

0014. Silver difluoride

[7783-95-1]

AgF₂

AgF₂

Boron, Water

Tulis, A. J. *et al.*, *Proc. 7th Symp. Explos. Pyrotechnics*, 1971, **3**(4), 1–12

Mixtures of boron and silver difluoride function as detonators when contacted with water.

Dimethyl sulfoxide

See Iodine pentafluoride: Dimethyl sulfoxide

Hydrocarbons, or Water

Priest, H. F. *Inorg. Synth.*, 1950, **3**, 176

It reacts even more vigorously with most substances than does cobalt fluoride.

See other METAL HALIDES, SILVER COMPOUNDS

0015. Silver amide

[65235-79-2]

AgH₂N

AgNH₂

Brauer, 1965, Vol. 2, 1045

Extraordinarily explosive when dry.

See Nitrogen triiodide–silver amide

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0016. Silver N-nitrosulfuric diamidate

[]

AgH₂N₃O₄S

AgN(NO₂)SO₂NH₂

Sorbe, 1968, 120

The silver salt of the nitroamide is explosive.

See other N-METAL DERIVATIVES, N-NITRO COMPOUNDS, SILVER COMPOUNDS

0017. Silver phosphinate

[]

AgH₂O₂P

AgP(H)(O)OH

Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 335

Explosive, but less sensitive than the azide or fulminate.
See other METAL PHOSPHINATES, SILVER COMPOUNDS

0018. Diamminesilver permanganate

[]




Pascal, 1960, Vol. 16, 1062

It may explode on impact or shock.

See other AMMINEMETAL OXOSALTS, SILVER COMPOUNDS

0019. Dihydrazinesilver nitrate

[31247-72-0]



Gall, H. *et al.*, *Z. Anorg. Chem.*, 1932, **206**, 376

The salt explodes at -1.5°C .

See other AMMINEMETAL OXOSALTS, SILVER COMPOUNDS

0020. Silver iodate

[7783-97-3]



Metals

MRH Potassium 1.50/25, sodium 1.92/35

See Potassium: Oxidants

Sodium: Iodates

Other reactants

Yoshida, 1980, 194

MRH values for 16 combinations with oxidisable materials are given.

Tellurium

Pascal, 1960, Vol. 13.2, 1961

Interaction is violent.

See other METAL HALOGENATES, SILVER COMPOUNDS

0021. Silver permanganate

[7783-98-4]



Sulfuric acid

491M, 1975, 369

The moist salt exploded during drying over the concentrated acid in a vacuum desiccator, (presumably owing to formation of traces of manganese heptoxide from reaction with sulfuric acid vapour).

See other OXIDANTS, SILVER COMPOUNDS

0022. Silver nitrate
[7761-88-8]



HCS 1980, 822

Acetaldehyde

Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 336

Aqueous silver nitrate reacts with acetaldehyde to give explosive silver fulminate.

Acetylene and derivatives

Mellor, 1946, Vol. 5, 854

Silver nitrate (or other soluble salt) reacts with acetylene in presence of ammonia to form silver acetylide, a sensitive and powerful detonator when dry. In the absence of ammonia, or when calcium acetylide is added to silver nitrate solution, explosive double salts of silver acetylide and silver nitrate are produced. Mercury(I) acetylide precipitates silver acetylide from the aqueous nitrate.

See 1,3-Butadiyne, and Buten-3-yne, both below

See METAL ACETYLIDES

Acrylonitrile

See Acrylonitrile: Silver nitrate

Aluminium

Laing, M., *J. Chem. Educ.*, 1994, **71**, 270

It is warned that a mixture of aluminium powder and silver nitrate is potentially as dangerous as that with magnesium, both being capable of producing >8 kJ/g.

See Magnesium, Water; below

Ammonia

MRH 1.46/29

1. *MCA Case History No. 2116*

2. *CISHC Chem. Safety Summ.*, 1976, **47**, 31

3. MacWilliam, E. A. *et al.*, *Photogr. Sci. Eng.*, 1977, **21**, 221–224

A bottle containing Gomari tissue staining solution (ammoniacal silver nitrate) prepared 2 weeks previously exploded when disturbed. The solution must be prepared freshly each day, and discarded immediately after use with appropriate precautions [1]. A large quantity of ammoniacal silver nitrate solution exploded violently when disturbed by removing a glass rod [2]. However, it has now been shown that neither the solid precipitated during addition of ammonia to the nitrate, nor the redissolved complex, is sensitive to initiation by very severe shocks. This was so for fresh or aged solutions. The solids produced by total evaporation at 95°C or higher would explode only at above 100 kg cm shock force. A pH value above 12.9 is essential for separation of explosive precipitates, and this cannot be attained by addition of ammonia alone [3].

See SILVER-CONTAINING EXPLOSIVES (reference 2)

Silver(I) oxide: Ammonia, etc.

Ammonia, Ethanol

MCA Case History No. 1733

A silvering solution exploded when disturbed. This is a particularly dangerous mixture, because both silver nitride and silver fulminate could be formed.

See Ethanol, below

Ammonia, Sodium carbonate

Vasbinder, H., *Pharm. Weekblad*, 1952, **87**, 861–865

A mixture of the components in gum arabic solution (marking ink) exploded when warmed.

Ammonia, Sodium hydroxide

1. Milligan, T. W. *et al.*, *J. Org. Chem.*, 1962, **27**, 4663

2. *MCA Case History No. 1554*

3. Morse, J. R., *School Sci. Rev.*, 1955, **37**(131), 147

4. Baldwin, J., *School Sci. Rev.*, 1967, **48**(165), 586

5. MacWilliam, E. A. *et al.*, *Photogr. Sci. Eng.*, 1977, **21**, 221–224

6. Anon., *Univ. Safety Assoc. Safety News*, 1977, (8), 15–16

During preparation of an oxidising agent on a larger scale than described [1], addition of warm sodium hydroxide solution to warm ammoniacal silver nitrate with stirring caused immediate precipitation of black silver nitride which exploded [2]. Similar incidents had been reported previously [3], including one where explosion appeared to be initiated by addition of Devarda's alloy (Al–Cu–Zn) [4]. The explosive species separates at pH values above 12.9, only produced when alkali is added to ammoniacal silver solutions, or when silver oxide is dissolved with ammonia [5]. The Sommer & Market reagent mixture used to identify cellulose derivatives led to a severe explosion [6].

See Silver nitride, also Ammonia: Silver compounds

See also SILVERING SOLUTIONS, TOLLENS' REAGENT

Arsenic

Mellor, 1941, Vol. 3, 470

A finely divided mixture with excess nitrate ignited when shaken out onto paper.

1,3-Butadiyne

See 1,3-Butadiyne: Silver nitrate

Buten-3-yne

See Buten-3-yne: Silver nitrate

Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Chlorosulfuric acid

Mellor, 1941, Vol. 3, 470

Interaction is violent, nitrosulfuric acid being formed.

Disilver ketenide

See Disilver ketenide–silver nitrate

Ethanol

MRH 2.55/12

1. Tully, J. P., *Ind. Eng. Chem. (News Ed.)*, 1941, **19**, 250
2. Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 334
3. Garin, D. L. *et al.*, *J. Chem. Educ.*, 1970, **47**, 741
4. Perrin, D. D. *et al.*, *Chem. Brit.*, 1986, **22**, 1084; *Chem. Eng. News*, 1987, **65**(2), 2

Reclaimed silver nitrate crystals, damp with the alcohol used for washing, exploded violently when touched with a spatula, generating a strong smell of ethyl nitrate [1]. The explosion was attributed to formation of silver fulminate (which is produced on addition of ethanol to silver nitrate solutions). Ethyl nitrate may also have been involved. Alternatives to avoid ethanol washing of recovered silver nitrate are discussed [2], including use of 2-propanol [3]. Another case of explosion during filtration of silver nitrate purified by progressive dilution with ethanol of its aqueous solution has been reported. Initiation was by agitation of the slurry on a glass frit with a spatula [4].

See Silver fulminate

Magnesium, Water

MRH 6.94/30

1. Marsden, F., private comm., 1973
2. Lyness, D. J. *et al.*, *School Sci. Rev.*, 1953, **35**(125), 139

An intimate mixture of dry powdered magnesium and silver nitrate may ignite explosively on contact with a drop of water [1,2].

See other REDOX REACTIONS

Non-metals

MRH Carbon 2.46/10, phosphorus (y) 3.89/18, sulfur 1.67/16

Mellor, 1941, Vol. 3, 469–473

Under a hammer blow, a mixture with charcoal ignites, while mixtures with phosphorus or sulfur explode, the latter violently.

Other reactants

Yoshida, 1980, 194

MRH values for 19 combinations with oxidisable materials are given.

Phosphine

Mellor, 1941, Vol. 3, 471

Rapid passage of gas into a conc. nitrate solution caused an explosion, or ignition of a slower gas stream. The explosion may have been caused by rapid oxidation of the precipitated silver phosphide derivative by the co-produced nitric acid or dinitrogen tetraoxide.

Phosphonium iodide

See Phosphonium iodide: Oxidants

Polymers

491M, 1975, 366

To assess suitability of plastic storage containers for distribution of silver nitrate, behaviour under fire exposure conditions of various polymers in contact with the salt was examined. All polymers tested burned vigorously.

Silver acetylide

See Silver acetylide–silver nitrate

Thiophene

Southern, T., private communication, 1990

A black solid is produced from these two reagents under influence of ultrasound (but not otherwise) which explodes violently on warming. It is apparently not silver acetylide.

Titanium

Shanley, E. S., *Chem. Eng. News*, 1990, **68**(16), 2

A titanium-containing sludge from a nitric acid bath was separated, before completely dry it exploded, killing a workman. Investigation showed the dry sludge to be a powerful explosive sensitive to heat, friction and impact, composed of about 60:40 silver nitrate:titanium.

See Titanium: Nitric acid

See other METAL NITRATES, SILVER COMPOUNDS

0023. Silver azide

[13863-88-2]

AgN₃

AgN₃

1. Mellor, 1940, Vol. 8, 349; 1967, Vol. 8, Suppl. 2, 47
2. Gray, P. *et al.*, *Chem. & Ind.*, 1955, 1255
3. Kabanov, A. A. *et al.*, *Russ. Chem. Rev.*, 1975, **44**, 538
4. Ryabykh, S. M. *et al.*, *Chem. Abs.*, 1984, **100**, 194549

As a heavy metal azide, it is considerably endothermic ($\Delta H_f^\circ + 279.5$ kJ/mol, 1.86 kJ/g). While pure silver azide explodes at 340°C [1], the presence of impurities may cause explosion at 270°C. It is also impact-sensitive and explosions are usually violent [2]. Its use as a detonator has been proposed. Application of an electric field to crystals of the azide will detonate them, at down to –100°C [3], and it may be initiated by irradiation with electron pulses of nanosecond duration [4].
See other CATALYTIC IMPURITY INCIDENTS, IRRADIATION DECOMPOSITION INCIDENTS

Ammonia

Mellor, 1940, Vol. 8, 349

Solutions in aqueous ammonia become explosive around 100°C.

Chlorine azide

See Chlorine azide: Ammonia, etc.

Halogens

Mellor, 1940, Vol. 8, 336

Silver azide, itself a sensitive compound, is converted by ethereal iodine into the less stable and explosive compound, iodine azide. Similarly, contact with nitrogen-diluted bromine vapour gives bromine azide, often causing explosions.

See Silver azide chloride

Metal oxides, or Metal sulfides

Kurochin, E. S. *et al.*, *Chem. Abs.*, 1974, **83**, 201390

Pure silver azide explodes at 340°C, but presence of below 10% of copper(I) or (II) oxides or sulfides, copper(I) selenide or bismuth(III) sulfide reduces the detonation temperature to 235°C. Concentrations of 10% of copper(II) oxide, copper(I) selenide or sulfide further reduced it to 200, 190 and 170°C, respectively.

Photosensitising dyes

Aleksandrov, E. *et al.*, *Chem. Abs.*, 1974, **81**, 31755

In a study of dye-sensitised silver azide, it was found that many dyes caused explosions in the initial stages.

Sulfur dioxide

Mellor, 1940, Vol. 8, 349

Mixtures of the slightly soluble azide with liquid sulfur dioxide became explosive at elevated temperatures.

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES, SILVER COMPOUNDS

0024. Silver trisulfurpentanitride

[]

(Structure unknown)

AgN₅S₃

See 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazine: Ammonia

See other N-S COMPOUNDS, SILVER COMPOUNDS

0025. Silver(II) oxide

[1301-96-8]

AgO

AgO

Hydrogen sulfide

See Hydrogen sulfide: Metal oxides

See other METAL OXIDES, SILVER COMPOUNDS

0026. Silver sulfide

[21548-73-2]

AgS

AgS



Potassium chlorate

See Potassium chlorate: Metal sulfides

See other METAL SULFIDES, SILVER COMPOUNDS

0027. Silver hexahydrohexaborate(2-)

[]



Bailar, 1973, Vol. 1, 808


It is a detonable salt.

See other SILVER COMPOUNDS, COMPLEX HYDRIDES**0028. Silver perchlorylamide**

[25870-02-4]

*See entry* PERCHLORYLAMIDE SALTS *See other* SILVER COMPOUNDS**0029. Tetrasulfurtetraimide-silver perchlorate**

[64867-41-0]

Nabi, S. N., *J. Chem. Soc., Dalton Trans.*, 1977, 1156

The complex detonates violently at 120°C.

See related METAL PERCHLORATES, N-S COMPOUNDS*See other* SILVER COMPOUNDS**0030. Silver imide**

[]



Bailar, 1973, Vol. 3, 101

It explodes very violently when dry.

But see Silver(I) oxide: Ammonia (reference 2)*See other* N-METAL DERIVATIVES, SILVER COMPOUNDS

0031. Silver hyponitrite
[7784-04-5]

AgON:NOAg

Ag₂N₂O₂

See Hyponitrous acid

See other METAL OXONON-METALLATES, SILVER COMPOUNDS

0032. Silver(I) oxide
[20667-12-3]

Ag₂O

Ag₂O

Aluminium

See Copper(II) oxide: Metals

Ammonia

1. Vasbinder, H., *Pharm. Weekblad*, 1952, **87**, 861–865

2. MacWilliam, E. A., *Photogr. Sci. Eng.*, 1977, **21**, 221–224

The clear solution, obtained by centrifuging a solution of the oxide in aqueous ammonia which had been treated with silver nitrate until precipitation started, exploded on two occasions after 10–14 days' storage in closed bottles in the dark. This was ascribed to slow precipitation of amorphous silver imide, which is very explosive even when wet [1]. When silver oxide is dissolved in ammonia solution, an extremely explosive precipitate (probably Ag₃N₄) will separate. The explosive behaviour is completely inhibited by presence of colloids or ammonium salts (acetate, carbonate, citrate or oxalate). Substitution of methylamine for ammonia does not give explosive materials [2].

See Silver nitrate: Ammonia, etc., *also* Ammonia: Silver compounds

Ammonia or Hydrazine, Ethanol

Silver oxide and ammonia or hydrazine slowly form explosive silver nitride and, in presence of alcohol, silver fulminate may also be produced.

See entries SILVER-CONTAINING EXPLOSIVES, SILVERING SOLUTIONS

Boron trifluoride etherate, Nitromethane

See Silver tetrafluoroborate

Carbon monoxide

Mellor, 1941, Vol. 3, 377

Carbon monoxide is exothermically oxidised over silver oxide, and the temperature may attain 300°C.

Chlorine, Ethylene

See Ethylene: Chlorine

Dichloromethylsilane

See Dichloromethylsilane: Oxidants

Hydrogen sulfide

See Hydrogen sulfide: Metal oxides

Magnesium

Mellor, 1941, Vol. 3, 378

Oxidation of magnesium proceeds explosively when warmed with silver oxide in a sealed tube.

Metal sulfides

Mellor, 1941, Vol. 3, 376

Mixtures with gold(III) sulfide, antimony sulfide or mercury(II) sulfide ignite on grinding.

Nitroalkanes

See NITROALKANES: metal oxides

Non-metals

Mellor, 1941, Vol. 3, 376–377

Selenium, sulfur or phosphorus ignite on grinding with the oxide.

Potassium–sodium alloy

See Potassium–sodium alloy: Metal oxides

Seleninyl chloride

See Seleninyl chloride: Metal oxides

Selenium disulfide

Mellor, 1941, Vol. 3, 377

A mixture may ignite under impact.

See other METAL OXIDES, SILVER COMPOUNDS

0033. Silver peroxide

[25455-73-6]

AgOOAg

Ag₂O₂

Poly(isobutene)

Mellinger, T., *Arbeitsschutz*, 1972, 248

Mixtures of silver peroxide with 1% of polyisobutene exploded on 3 separate occasions. Use of a halogenated polymer was safe.

See other METAL PEROXIDES

0034. Silver osmate

[]

Ag₂OsO₄

Ag₂O₄Os

Sorbe, 1968, 126

Explodes on impact or heating.

See other HEAVY METAL DERIVATIVES, SILVER COMPOUNDS

0035. Silver sulfide
[21548-73-2]

Ag₂S

AgSAg

Potassium chlorate

See Potassium chlorate: Metal sulfides

See other METAL SULFIDES, SILVER COMPOUNDS

0036. Disilver pentatin undecaoxide

[]

Ag₂Sn₅O₁₁

(Structure unknown)

Mellor, 1941, Vol. 7, 418


The compound 'silver beta-stannate' is formed by long contact between solutions of silver and stannous nitrates, and loses water on heating, then decomposing explosively.

See other METAL OXIDES, SILVER COMPOUNDS

0037. Silver peroxochromate

[]

Ag₃CrO₈



Sulfuric acid

Riesefeld, E. H. *et al.*, *Ber.*, 1914, **47**, 548

In attempts to prepare 'perchromic acid', a mixture of silver (or barium) peroxochromate and 50% sulfuric acid prepared at -80°C reacted explosively on warming to about -30°C .

See other PEROXOACID SALTS, SILVER COMPOUNDS

0038. Silver nitride

[20737-02-4]

Ag₃N

Ag₃N

Hahn, H. *et al.*, *Z. Anorg. Chem.*, 1949, **258**, 77

Very sensitive to contact with hard objects, exploding when moist. An extremely sensitive explosive when dry, initiatable by friction, impact or heating. The impure product produced by allowing ammoniacal silver oxide solution to stand seems even more sensitive, often exploding spontaneously in suspension.



See Silver chloride: Ammonia

See also SILVER-CONTAINING EXPLOSIVES, SILVERING SOLUTIONS, TOLLENS' REAGENT

See other CATALYTIC IMPURITY INCIDENTS, N-METAL DERIVATIVES, SILVER COMPOUNDS

0039. Silver 2,4,6-tris(dioxoselena)perhydrotriazine-1,3,5-triide ('Silver triselenimidate')

[]



See Selenium difluoride dioxide: Ammonia

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0040. Trisilver tetranitride

[]



See Silver(I) oxide: Ammonia (reference 2)

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0041. Tetrasilver orthodiamidophosphate

[]



Sulfuric acid

Mellor, 1940, Vol. 8, 705

Ignites with sulfuric acid.

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0042. Tetrasilver diimidotriphosphate

[]



Alone, or Sulfuric acid

Mellor, 1940, Vol. 8, 705; 1971, Vol. 8, Suppl. 3, 787



The dry material explodes on heating, and ignites in contact with sulfuric acid.

The molecule contains one N–Ag bond.

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0043. Tetrasilver diimidodioxosulfate

[]




Nachbauer, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1973, **12**, 339

The dry salt explodes on friction or impact.
See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0044. Pentasilver orthodiamidophosphate

[]

$\text{Ag}_5\text{H}_2\text{N}_2\text{O}_3\text{P}$



Alone, or Sulfuric acid

Stokes, H. N., *Amer. Chem. J.*, 1895, **17**, 275

Explodes on heating, friction or contact with sulfuric acid.

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0045. Pentasilver diimidotriphosphate

[]

$\text{Ag}_5\text{H}_2\text{N}_2\text{O}_8\text{P}_3$



Alone, or Sulfuric acid

Mellor, 1940, Vol. 8, 705; 1971, Vol. 8, Suppl. 3, 787

The salt is explosive and may readily be initiated by friction, heat or contact with sulfuric acid. The molecule contains two N–Ag bonds.

0046. Pentasilver diamidophosphate

[]

$\text{Ag}_5\text{N}_2\text{O}_2\text{P}$



1. Bailar, 1973, Vol. 8, 455

2. Mellor, 1940, Vol. 8, 705


The salt contains 85% silver and four N–Ag bonds [1], and detonates readily on friction, heating or contact with sulfuric acid [2].

See other N-METAL DERIVATIVES, SILVER COMPOUNDS

0047. Heptasilver nitrate octaoxide

[12258-22-9]

$\text{Ag}_7\text{NO}_{11}$



Alone, or Sulfides, or Non-metals

Mellor, 1941, Vol. 3, 483–485

The crystalline product produced by electrolytic oxidation of silver nitrate (and possibly as formulated) detonates feebly at 110°C. Mixtures with phosphorus and sulfur explode on impact, hydrogen sulfide ignites on contact and antimony trisulfide ignites when ground with the salt.

See related METAL NITRATES See other SILVER COMPOUNDS

0048. Aluminium

[7429-90-5]

Al

Al

HCS 1980, 135 (powder)

1. *Haz. Chem. Data*, 1975, 44
2. *Dust Explosion Prevention: Aluminum Powder*, NFPA Standard Codes 65, 651, both 1987
3. Popov, E. I. *et al.*, *Chem. Abs.*, 1975, **82**, 61411
4. Ida, K. *et al.*, *Japan Kokai*, 1975, 75 13 233
5. Steffens, H. D. *et al.*, *Chem. Abs.*, 1979, **90**, 209354
6. Scherbakov, V. K. *et al.*, *Chem. Abs.*, 1979, **91**, 76322
7. Barton, J. A. *et al.*, *Chem. Brit.*, 1986, **22**, 647–650
8. May, D. C. *et al.*, *J. Haz. Mat.*, 1987, **17**(1), 61–88
9. See entry DUST EXPLOSION INCIDENTS (reference 22)

Finely divided aluminium powder or dust forms highly explosive dispersions in air [1], and all aspects of prevention of aluminium dust explosions are covered in 2 recent US National Fire Codes [2]. The effects on ignition properties of impurities introduced by recycled metal used to prepare dust were studied [3]. Pyrophoricity is eliminated by surface coating aluminium powder with polystyrene [4]. Explosion hazards involved in arc and flame spraying of the powder are analysed and discussed [5], and the effect of surface oxide layers on flammability was studied [6]. The causes of a severe explosion in 1983 in a plant producing fine aluminium powder are analysed, and improvements in safety practices discussed [7]. A number of fires and explosions involving aluminium dust arising from grinding, polishing and buffing operations are discussed, and precautions detailed [8]. Atomised and flake aluminium powders attain maximum explosion pressures of 5.7 and 8.6 bar, respectively, both with maximum rates of pressure rise exceeding 1.36 kbar/s [9].

Air, Hydrocarbons

Lamnevik, S., *Chem. Abs.*, 1982, **96**, 202034

Presence of dispersed aluminium powder in propane–air or butane–air mixtures enhances the detonative properties of stoichiometric mixtures.

Air, Water

1. Turetzky, M., private comm., 1987
2. *MARS Database*, 1998, short report 028

Fine aluminium dust and chippings from precision surface machining operations were air-transported to a cyclone collection system with outlet bag filters for the dust, and the cyclone contents were discharged into metal scrap bins. Sparks from a portable grinding machine ignited aluminium dust on a maintenance platform, and use of a carbon dioxide extinguisher on the fire transferred it into a metal scrap bin below. Heat from this larger fire activated the automatic water sprinkler system, and contact of the water (89% oxygen) with the burning metal liberated hydrogen, which, after mixing with air, exploded. This primary explosion created an aluminium dust cloud which exploded forcefully, creating larger dust clouds and eventually involving the contents of the other scrap bins and 2 of the dust filter bags, leading to at least 4 separate tertiary explosions in all. Structural damage to adjacent buildings and vehicles was extensive. The dry collection system was replaced by a wet one [1].

A different scenario involving these three occurred when lightning struck an aluminium foundry. It is supposed that this dispersed molten metal droplets in air, which then exploded with the estimated force of 200 kg TNT, causing damage which allowed remaining molten metal to fall into the wet casting pit, producing a second explosion of half the power of the first [2].

Aluminium halides, Carbon oxides

Guntz, A. *et al.*, *Compt. rend.*, 1897, **124**, 187–190

Aluminium powder burns when heated in carbon dioxide, and presence of aluminium chloride or aluminium iodide vapour in carbon monoxide or carbon dioxide accelerates the reaction to incandescence.

Ammonium nitrate

MRH 8.70/75

Mellor, 1946, Vol. 5, 219

Mixtures with the powdered metal are used as an explosive, sometimes with the addition of carbon or hydrocarbons, or other oxidants.

See Ammonium nitrate: Metals

Ammonium nitrate, Calcium nitrate, Formamide

See Calcium nitrate: Aluminium, etc.

Ammonium peroxodisulfate

See Ammonium peroxodisulfate: Aluminium

Antimony or Arsenic

Matignon, C., *Compt. rend.*, 1900, **130**, 1393

Powdered aluminium reacts violently on heating with antimony or arsenic.

Antimony trichloride

Matignon, C., *Compt. rend.*, 1900, **130**, 1393

The metal powder ignites in antimony trichloride vapour.

Arsenic trioxide, Sodium arsenate, Sodium hydroxide

MCA Case History No.1832

An aluminium ladder was used (instead of the usual wooden one) to gain access to a tank containing the alkaline arsenical mixture. Hydrogen produced by alkaline attack on the ladder generated arsine, which poisoned the three workers involved.

Barium peroxide

See Barium peroxide: Metals

Bismuth

Mellor, 1947, Vol. 9, 626

The finely divided mixture of metals produced by hydrogen reduction of coprecipitated bismuth and aluminium hydroxides is pyrophoric.

Butanol

Luberoff, B. J., private comm., 1964

Butanol, used as a solvent in an autoclave preparation at around 100°C, severely attacked the aluminium gasket, liberating hydrogen which caused a sharp rise in pressure. Other alcohols would behave similarly, forming the aluminium alkoxide.

See other CORROSION INCIDENTS

Calcium oxide, Chromium oxide, Sodium chlorate

MRH 10.7/67

Nolan, 1983, Case History 174

In a thermite process to produce chromium metal, the mechanically-mixed ingredients were ignited in a large crucible and the reaction proceeded smoothly. When the mixer broke down, manual mixing was used but gave poorer dispersion of the constituents. An explosion after ignition is attributed to a high local concentration of sodium chlorate and aluminium powder in the mixture.

See other THERMITE REACTIONS

Carbon, Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Carbon dioxide

See Carbon dioxide: Metals

Carbon dioxide, Sodium peroxide

See Sodium peroxide: Metals, etc.

Carbon disulfide

Matignon, C., *Compt. rend.*, 1900, **130**, 1391

Aluminium powder ignites in carbon disulfide vapour.

Carbon tetrachloride, Methanol

See Methanol: Carbon tetrachloride, Metals

Catalytic metals

1. *49IM*, 1975, 28

2. Kirk-Othmer (3rd ed.), 1982, Vol. 19, 495

At the m.p. of aluminium (600°C) an aluminium-sheathed palladium thermocouple formed an alloy with a flash and an exotherm to 2800°C [1]. The use of thin layers of palladium or platinum on aluminium foil or wire as igniters derives from the intense heat of alloy formation, which is sufficient to melt the intermetallic compounds [2].

Chloroformamidinium nitrate

See Chloroformamidinium nitrate (reference 2)

Copper, Sulfur

Donohue, P. C., US Pat. 3 932 291, 1973

During preparation of aluminium copper(I) disulfide from the elements in an air-free silica tube at 900–1000°C, initial heating must be slow to prevent explosion of the tube by internal pressure of unreacted sulfur vapour.

Copper(I) oxide

Haws, L. D. *et al.*, *Proc. 6th Int. Pyrotech. Semin.*, 1978, 209–222

Technical aspects of the use of aluminium–copper(I) oxide thermite devices are detailed.

See Copper(I) oxide: Aluminium

Copper(II) oxide

MRH 3.29/82

1. Anon., *Chem. Age.*, 1932, **27**, 23

2. Scamaton, W. B. *et al.*, *Chem. Abs.*, 1980, **92**, 8484

3. Peterson, W. S., *Light Met.* (Warrendale, PA), 1987, 699–701

A mixture of aluminium powder and hot copper oxide exploded violently during mixing with a steel shovel on an iron plate. The frictional mixing initiated the thermite-like mixture [1]. Such mixtures are now used in electro-explosive devices [2]. Two cases of violent explosions after adding scrap copper to molten aluminium are discussed. In both cases, when some undissolved copper with adhering aluminium and oxide dross was removed from the furnace, the explosions occurred outside the melting furnace [3].

See Metal oxides, etc., below

Diborane

See Diborane: Metals

Dichloromethane

Piotrowski, A. M. *et al.*, *J. Org. Chem.*, 1988, **53**, 2829–2835

Conditions are described for the safe reaction of aluminium powder with dichloromethane to give bis(dichloroaluminum)methane in high yield. Derivatives of this compound are effective methyleneating agents for ketones.

See Halocarbons, below

Diethyl ether

Murdock, T. O., Ph.D. Thesis, Univ. N. Dakota, 1977 (*Diss. Abs. B*, 1978, **39**, 1291)

The aluminium–solvent slurry produced by metal atom/solvent co-condensation at -196°C is so reactive that oxygen is abstracted from the solvent ether as the mixture is allowed to melt. Hydrocarbon solvents are more suitable (but halocarbon solvents would react explosively).

Disulfur dibromide

See Disulfur dibromide: Metals

Drain cleaner, Cola

1. Utterbuck, P., Internet, 1997

2. Editor's comments

A student mixed aluminium foil and drain cleaner in a soft drink bottle, which started emitting gas. Another student carried the bottle outside and was claimed to have been overcome by the toxic fumes [1]. Most drain cleaners are alkalis, so that aluminium will dissolve to produce hydrogen. The bleach that it is suggested may have been present will produce no toxic fumes in alkali, and one would be surprised to find arsenic or antimony compounds present. If the collapse was not

purely hysterical, the remaining, though remote, possibility would be phosphine. The soft drink the bottle had contained was one of the many perhaps best described as impure dilute phosphoric acid [2].

See Arsenic trioxide, etc., above

Explosives

1. Stettbacher, A., *Chem. Abs.*, 1944, **38**, 4445₄

2. Muraour, H., *Chem. Abs.*, 1944, **38**, 4445₇

The addition of substantial amounts (up to 32%) of aluminium powder to conventional explosives enhances the energy release by up to 100% [1], involving high temperature reduction of liberated carbon dioxide and water by the metal [2].

Formic acid

Matignon, C., *Compt. rend.*, 1900, **130**, 1392

The metal reduces the acid (itself a reductant) with incandescence

1. Anon., *Angew. Chem.*, 1950, **62**, 584

2. Anon., *Chem. Age*, 1950, **63**, 155

3. Anon., *Chem. Eng. News*, 1954, **32**, 258

4. *Pot. Incid. Rep.*, ASESB, 1968, **39**

5. Anon., *Chem. Eng. News*, 1955, **33**, 942

6. Eiseman, B. J., *J. Amer. Soc. Htg. Refr. Air Condg. Eng.*, 1963, **5**, 63

7. Eiseman, B. J., *Chem. Eng. News*, 1961, **39**(27), 44

8. Laccabue, J. R., *Fluorolube–Aluminium Detonation Point: Report 7E.1500*, San Diego, Gen. Dynamics, 1958

9. Atwell, V. J., *Chem. Eng. News*, 1954, **32**, 1824

10. ICI Mond Div., private comm., 1973

11. Anon., *Ind. Acc. Prev. Bull. RoSPA*, 1953, **21**, 60

12. Wendon, G. W., private comm., 1973

13. Coffee, R. D., *Loss Prev.*, 1971, **5**, 113

14. Schwab, R. F., *ibid.*, 113

15. Corley, R., *ibid.*, 114

16. Heinrich, H. J., *Arbeitsschutz*, 1966, 156–157

17. Lamoroux, A. *et al.*, *Mém. Poudres*, 1957, **39**, 435–445

18. Hamstead, A. C. *et al.*, *Corrosion*, 1958, **14**, 189t–190t

19. Hartmann, I., *Ind. Eng. Chem.*, 1948, **40**, 756

20. *MCA Case History No. 2160*

21. Arias, A., *Ind. Eng. Chem., Prod. Res. Dev.*, 1976, **15**, 150

22. Wendon, G. W., *Ind. Eng. Chem., Prod. Res. Dev.*, 1977, **16**, 112

23. Arias, A., *Ind. Eng. Chem., Prod. Res. Dev.*, 1977, **16**, 112

24. Archer, W. L., Paper 47, 1–11, presented at *Corrosion/78*, Houston, March 6–10, 1978

25. Anon. *Univ. Safety Assoc. Newsletter*, 1982, (16), 5

26. Stull, 1977, 22

27. Cardillo, P., *et al.*, *J. Loss Prevention*, 1992, **5**(2), 81

MRH Chloromethane 2.46/85, bromomethane 1.33/91, chloroform 3.84/82, carbon tetrachloride 4.15/91, carbon tetrafluoride 6.48/71, dichlorodifluoromethane

5.39/77, trichlorofluoromethane 4.76/79, trichloroethylene 4.10/83, trichlorofluoroethane 4.98/77, 1,2-dichloropropene 3.30/86, *o*-dichlorobenzene 2.67/89

Heating aluminium powder with carbon tetrachloride, chloromethane or carbon tetrachloride–chloroform mixtures in closed systems to 152°C may cause an explosion, particularly if traces of aluminium chloride are present [1]. A mixture of carbon tetrachloride and aluminium powder exploded during ball-milling [2], and it was later shown that heavy impact would detonate the mixture [3]. Mixtures with fluorotrichloroethane and with trichlorotrifluoroethane will flash or spark on heavy impact [4]. A virtually unvented aluminium tank containing a 4:1:2 mixture of *o*-dichlorobenzene, 1,2-dichloroethane and 1,2-dichloropropane exploded violently 7 days after filling. This was attributed to formation of aluminium chloride which catalysed further accelerating attack on the aluminium tank [5]. An analysis of the likely course of the Friedel-Crafts reaction and calculation of the likely heat release (29.1 kJ/mol) has been published [26].

In a dichlorodifluoromethane system, frictional wear exposed fresh metal surfaces on an aluminium compressor impellor, causing an exothermic reaction which melted much of the impellor. Later tests showed similar results, decreasing in order of intensity, with: tetrafluoromethane; chlorodifluoromethane; bromotrifluoromethane; dichlorodifluoromethane; 1,2-difluorotetrafluoroethane; 1,1,2-trichlorotrifluoroethane [6]. In similar tests, molten aluminium dropped into liquid dichlorodifluoromethane burned incandescently below the liquid [7]. Aluminium bearing surfaces under load react explosively with polytrifluoroethylene greases or oils. The inactive oxide film will be removed from the metal by friction, and hot spots will initiate reaction [8]. An attempt to scale up the methylation of 2-methylpropane with chloromethane in presence of aluminium chloride and aluminium went out of control and detonated, destroying the autoclave. The preparation had been done on a smaller scale on 20 previous occasions without incident [9].

Violent decomposition, with evolution of hydrogen chloride, may occur when 1,1,1-trichloroethane comes into contact with aluminium or its alloys with magnesium [10]. Aluminium-dusty overalls were cleaned by immersion in trichloroethylene. During subsequent drying, violent ignition occurred. This was attributed to presence of free hydrogen chloride in the solvent, which reacted to produce aluminium chloride [10]. This is known to catalyse polymerisation of trichloroethylene, producing more hydrogen chloride and heat. The reaction is self-accelerating and can develop a temperature of 1350°C [11]. Trichloroethylene cleaning baths must be kept neutral with sodium carbonate, and free of aluminium dust. Halocarbon solvents now available with added stabilisers (probably amines) show a reduced tendency to react with aluminium powder [12]

Aluminium powder undergoes an exothermic and uncontrollable reaction with dichloromethane above 95°C under appropriate pressure [13]. Several cases of violent reaction between aluminium and trichloroethylene or tetrachloroethylene in vapour degreasers have been noted [14]. Chloromethane in liquefied storage diffused 70 m along a nitrogen inerting line into the pressure regulator. Interaction with aluminium components of the regulator formed alkylaluminium compounds which ignited when the regulator was dismantled [15]. An explosion in an aluminium degreasing plant using tetrachloroethylene was attributed to

overheating of residues on the heating coils. Subsequent tests showed that simultaneous presence of water and aluminium chloride in an aluminium powder-tetrachloroethylene mixture lowered the initiation temperature to below 250°C. Presence of cutting oils reduced it below 150°C, and a temperature of 300°C was reached within 100 min [16]

Reaction of aluminium powder with hexachloroethane in alcohol is not initially violent, but may become so [17]. An aluminium transfer pipe failed after a few hours' service carrying refined 1,2-dichloropropane in warm weather. The corrosive attack was simulated and studied under laboratory conditions [18]. In dichlorodifluoromethane vapour, aluminium dust ignited at 580°C, and suspensions of the dust in the vapour gave strong explosions when sparked [19]. A fire occurred at a liquid outlet from a 40 m³ mild steel tanker of chloromethane. This was traced to the presence of trimethylaluminium produced by interaction of chloromethane and (unsuspected) aluminium baffle plates in the tanker [20]. A proposal to prepare pure aluminium chloride by ball-milling aluminium in carbon tetrachloride [21] was criticised [22] as potentially hazardous. Possible modifications (use of inert solvents, continuously fed mills, etc.) to improve the procedure were suggested [23]. The reaction mechanism of attack and inhibition in aluminium-1,1,1-trichloroethane systems has been investigated [24]. An attempt to clean a motor assembly containing an aluminium alloy gearwheel by soaking overnight in 1,1,1-trichloroethane led to gross degradation of the assembly [25].

A tabulation of theoretically hazardous halocarbon/aluminium combinations, and estimates of safe dilutions, calculated by the CHETAH program is reported [27].

See Bromomethane: Metals

See Dichloromethane, above

See COMPUTATION OF REACTIVE CHEMICAL HAZARDS, *See other* CORROSION INCIDENTS, METAL-HALOCARBON INCIDENTS

Halogens MRH Bromine 1.96/90, chlorine 5.23/80

1. Mellor, 1946, Vol. 2, 92, 135; Vol. 5, 209
2. Azmathulla, S. *et al.*, *J. Chem. Educ.*, 1955, **32**, 447; *School Sci. Rev.*, 1956, **38**(134), 107
3. Hammerton, C. M., *ibid*, 1957, **38**(136), 459

Aluminium powder ignites in chlorine without heating, and foil reacts vigorously with liquid bromine at 15°C, and incandesces on warming in the vapour [1]. The metal and iodine react violently in the presence of water, either as liquid, vapour or that present in hydrated salts [2]. Moistening a powdered mixture causes incandescence and will initiate a thermite mixture [3].

See Iodine: Aluminium, Diethyl ether

Hydrochloric acid or Hydrofluoric acid

Kirk-Othmer, 1963, Vol. 1, 952

The metal is attacked violently by the aqueous acids.

Hydrogen chloride

MRH 3.05/80

Batty, G. F., private comm., 1972

Erroneous use of aluminium instead of alumina pellets in a hydrogen chloride purification reactor caused a vigorous exothermic reaction which distorted the steel reactor shell.

Interhalogens

See Bromine pentafluoride: Acids, etc.

Chlorine fluoride: Aluminium

Iodine chloride: Metals

Iodine heptafluoride: Metals

Iodine pentafluoride: Metals

Iron, Water

Chen, W. Y. *et al.*, *Ind. Eng. Chem.*, 1955, **47**(7), 32A

A sludge of aluminium dust (containing iron and sand) removed from castings in water was found, during summer weather, to undergo sudden exotherms to 95°C with hydrogen evolution. Similar effects with aluminium-sprayed steel plates exposed to water were attributed to electrolytic action, as addition of iron filings to a slurry of aluminium in water caused self heating and hydrogen evolution to occur.

See Water, below

See other CORROSION INCIDENTS

Mercury(II) salts

Woelfel, W. C., *J. Chem. Educ.*, 1967, **44**, 484

Aluminium foil is unsuitable as a packing material in contact with mercury(II) salts in presence of moisture, when vigorous amalgamation ensues.

See Aluminium amalgam (reference 2)

See other CORROSION INCIDENTS

Metal nitrates, Potassium perchlorate, Water

Johansson, S. R. *et al.*, *Chem. Abs.*, 1973, **78**, 18435

A pyrotechnic mixture of aluminium powder with potassium perchlorate, barium nitrate, potassium nitrate and water exploded after 24 h storage under water. Tests revealed the exothermic interaction of finely divided aluminium with nitrate and water to produce ammonia and aluminium hydroxide. Under the conditions prevailing in the stored mixture, the reaction would be expected to accelerate, finally involving the perchlorate as oxidant and causing ignition of the mixture.

Metal nitrates, Sulfur, Water

Anon., *Chem. Eng. News*, 1954, **32**, 258

Aluminium powder, barium nitrate, potassium nitrate, sulfur and vegetable adhesives, mixed to a paste with water, exploded on 2 occasions. Laboratory investigation showed initial interaction of water and aluminium to produce hydrogen. It was supposed that nascent hydrogen reduced the nitrates present, increasing the alkalinity and thence the rate of attack on aluminium, the reaction becoming self-accelerating. Cause of ignition was unknown. Other examples of interaction of aluminium with water are known.

See Iron, Water, above; Water, below

See other CORROSION INCIDENTS

Metal oxides or Oxosalts or Sulfides

1. Mellor, 1946, Vol. 5, 217

2. Price, D. J. *et al.*, *Chem. Met. Eng.*, 1923, **29**, 878

MRH Chromium trioxide 6.15/79, iron(II) oxide 3.01/80, iron(II)(III) oxide 3.38/76, calcium sulfate 5.73/65, sodium carbonate 2.92/67, sodium nitrate 8.32/65, sodium sulfate 5.27/66

Many metal oxo-compounds (nitrates, oxides and particularly sulfates) and sulfides are reduced violently or explosively (i.e. undergo 'thermite' reaction) on heating an intimate mixture with aluminium powder to a suitably high temperature to initiate the reaction. Contact of massive aluminium with molten salts may give explosions [1]. Application of sodium carbonate to molten (red hot) aluminium caused an explosion [2].

See Sodium sulfate, below

See Iron(III) oxide: Aluminium, *See other* MOLTEN SALT BATHS, THERMITE REACTIONS

Methanol

Médard, L. *Mém. Poudres*, 1951, **33**, 490–503 (Engl. translation HSE 11270, 1986 available from HSE/LIS)

The explosive nature of mixtures of aluminum or magnesium with methanol or water is detailed.

See also Magnesium: Methanol

Niobium oxide, Sulfur

491M, 1975, 28

A mixture caused a serious fire.

See Metal oxides, etc., above

Nitro compounds, Water

Hajek, V. *et al.*, *Research*, 1951, **4**, 186–191

Dry mixtures of picric acid and aluminium powder are inert, but addition of water causes ignition after a delay dependent upon the quantity added. Other nitro compounds and nitrates are discussed in this context.

Non-metal halides

MRH Phosphorus pentachloride 2.76/83

1. Matignon, C., *Compt. rend.*, 1900, **130**, 1393

2. Lenher, V. *et al.*, *J. Amer. Chem. Soc.*, 1926, **48**, 1553

3. Berger, E., *Compt. rend.*, 1920, **170**, 29

Powdered aluminium ignites in the vapour of arsenic trichloride or sulfur dichloride, and incandescens in phosphorus trichloride vapour [1]. Above 80°C, aluminium reacts incandescently with diselenium dichloride [2]. The powder ignites in contact with phosphorus pentachloride [3].

Non-metals

Matignon, C., *Compt. rend.*, 1900, **130**, 1393–1394

Powdered aluminium reacts violently with phosphorus, sulfur or selenium, and a mixture of powdered metal with red phosphorus exploded when severely shocked.

Oleic acid

de Ment, J., *J. Chem. Educ.*, 1956, **36**, 308

Shortly after mixing the two, an explosion occurred, but this could not be repeated. The acid may have been peroxidised.

Other reactants

Yoshida, 1980, 29

MRH values calculated for 44 combinations, largely with oxidants, are given.

Oxidants

MRH values show % of oxidant

1. Kirshenbaum, 1956, 4, 13
2. Mellor, 1947, Vol. 2, 310
3. Annikov, V. E. *et al.*, *Chem. Abs.*, 1976, **85**, 145389
4. Nakamura, H. *et al.*, *Chem. Abs.*, 1985, **102**, 97779

Mixtures of aluminium powder with liquid chlorine, dinitrogen tetroxide or tetranitromethane are detonable explosives, but not as powerful as aluminium–liquid oxygen mixtures, some of which exceed TNT in effect by a factor of 3 to 4 [1]. Mixtures of the powdered metal and various bromates may explode on impact, heating or friction. Iodates and chlorates act similarly [2]. Detonation properties of gelled slurries of aluminium powder in aqueous nitrate or perchlorate salt solutions have been studied [3]. Reactions of aluminium powder with potassium chlorate or potassium perchlorate have been studied by thermal analysis [4].

For other combinations,

See Halogens, above; Oxygen, below

Ammonium peroxodisulfate: Aluminium, etc.

Nitryl fluoride: Metals

Potassium chlorate: Metals

MRH 9.20/70

Potassium perchlorate: Aluminium, or: Metal powders

MRH 9.96/67

Sodium chlorate: Aluminium, Rubber

MRH (no rubber) 10.71/67

Sodium nitrate: Aluminium

MRH 8.32/65

Sodium peroxide: Aluminium, etc., or: Metals

MRH 4.56/81

Zinc peroxide: Alone, or Metals

See other REDOX REACTIONS

Oxygen

Phillips, B. R. *et al.*, *Combust. Flame*, 1979, **35**(3), 249–258

Metal powders or fibres ignite in an oxygen-fed resonance tube.

Oxygen, Water

Aleshim, M. A. *et al.*, *Chem. Abs.*, 1979, **91**, 76312

Injection of oxygen is used to initiate the metal–water reaction in rocket propulsion systems.

Paint

See Zinc: Paint primer base

Palladium

See Catalytic metals, above

Polytetrafluoroethylene

1. Pittaluga, F. *et al.*, *Termotechnica*, 1981, **25**, 332–334
2. Schwartz, A., Internet, 1996

A mixture of aluminium powder and PTFE has been evaluated as an underwater missile propellant [1]. It is suggested that using PTFE tape to lubricate and seal aluminium screw threads is inadvisable [2].

See Halocarbons, above

2-Propanol

1. Wilds, A. L., *Org. React.*, 1944, **2**, 198
2. Muir, G. D., private comm., 1968
3. Anon., *Safety Digest Univ. Safety Assoc.* 1994, (50), 21

Dissolution of aluminium in 2-propanol to give the isopropoxide is rather exothermic, but often subject to an induction period similar to that in preparation of Grignard reagents [1]. Only small amounts of aluminium should be present until reaction begins [2]. A canister of Video Display Unit Cleaner exploded in a cupboard. This is thought to have been because of faulty interior lacquer which allowed the propanol containing cleaner to attack the aluminium can, generating heat and hydrogen [3].

See Magnesium: Methanol

See Butanol; Methanol; both above

Silicon steel

Partington, 1967, 418

Aluminium is added (0.1%) to molten steel to remove dissolved oxygen and nitrogen to prevent blowholes in castings. It reacts very violently with silicon steels.

Silver chloride

MRH 0.67/94

Anon., *Chem. Eng. News*, 1954, **32**, 258

An intimate mixture of the two powders may lead to reaction of explosive violence, unless excess aluminium is present (but note low MRH value).

Sodium acetylide

See Sodium acetylide: Metals

Sodium dithionite, Water

See Sodium dithionite: Aluminium, Water

Sodium diuranate

Gray, L. W., *Rept. DP-1485*, Richmond (Va), NTIS, 1978

During outgassing of scrap uranium–aluminium cermet reactor cores, powerful exotherms led to melting of 9 cores. It was found that the incident was initiated by reactions at 350°C between aluminium powder and sodium diuranate, which released enough heat to initiate subsequent exothermic reduction of ammonium uranyl hexafluoride, sodium nitrate, uranium oxide and vanadium trioxide by aluminium, leading to core melting.

Sodium hydroxide

1. *MCA Case History No. 1115*
2. *MCA Case History No. 1888*

In an incident involving corrosive attack on aluminium by sodium hydroxide solution, the vigorous evolution of hydrogen was noticed before a tank trailer (supposed to be mild steel) had perforated [1]. Corrosion caused failure of an aluminium coupling between a pressure gauge and a pump, causing personal contamination [2].

Sodium sulfate

MRH 5.27/66

Kohlmeyer, E. J., *Aluminium*, 1942, **24**, 361–362

The violent explosion experienced when an 8:3 molar mixture of metal powder and salt was heated to 800°C was attributed to thermal dissociation (at up to 3000°C) of the metal sulfide(s) formed as primary product(s).

See other THERMITE REACTIONS

Steel

1. *MCA Case Histories Nos. 2161, 2184* (same incident)

2. Bailey, J. C., *Met. Mater.*, 1978, (Oct.), 26–27

A joint between a mild steel valve screwed onto an aluminium pipe was leaking a resin–solvent mixture, and when the joint was tightened with a wrench, a flash fire occurred. This was attributed to generation of sparks by a thermite reaction between the rusted steel valve and the aluminium pipe when the joint was tightened [1]. Hazards involved in the use of aluminium-sprayed steel are reviewed [2].

Sulfur

MRH 4.81/64

Read, C. W. W., *School Sci. Rev.*, 1940, **21**(83), 977

The violent interaction of aluminium powder with sulfur on heating is considered to be too dangerous for a school experiment.

Uranium compounds

See Sodium diuranate, above

Water

1. *MCA Case History No. 462*

2. Shidlovskii, A. A., *Chem. Abs.*, 1947, **41**, 1105d

3. Bamberger, M. *et al.*, *Z. Angew. Chem.*, 1913, **26**, 353–355

4. Gibson, 1969, 2

5. Pittaluga, F. *et al.*, *Termotechnica*, 1977, **31**, 306–312

6. Sukhov, A. V., *Chem. Abs.*, 1978, **88**, 193939

7. Tompa, A. S. *et al.*, *J. Haz. Mat.*, 1978, **2**, 197–215

8. Meguro, T. *et al.*, *Chem. Abs.*, 1980, **90**, 8505

9. Lemmon, A. W., *Light Met.* (New York), 1980, 817–836

10. Boxley, G. *et al.*, *Light Met.* (New York), 859–868

11. Epstein, S. G. *et al.*, *Light Met.* (Warrendale Pa), 1987, 693–698; *ibid.*, 1993, 845

12. Epstein, S. G., *Light Met.* (Warrendale, Pa), 1995, 885

13. Ikata, K., *Chem. Abs.*, 1993, **118**, 127778

14. Breault, R. *et al.*, *Light Met.* (Warrendale, Pa), 1995, 903

15. Hughes, D. T. *et al.*, *IMechE Conf. Trans.*, 1995, (2), 73

16. Taylor, F. R. *et al.*, *Rept. ARAED-TR-87022*, 1–26, 1988;

Cans of aluminium paint contaminated with water contained a considerable pressure of hydrogen from interaction of finely divided metal and moisture [1]. Mixtures of

powdered aluminium and water can be caused to explode powerfully by initiation with a boosted detonator [2]. A propellant explosive composed of aluminium and water has been patented [13]. During granulation of aluminium by pouring the molten metal through a sieve into water, a violent explosion occurred. This was attributed to steam trapped in the cooling metal [3], but see references [9,10]. Moist finely divided aluminium may ignite in air [4]. The aluminium–steam reaction was evaluated for power generation purposes, a peak steam temperature of over 1000°C being attained [5]. The combustion of aluminium in steam has been studied in detail [6]. In a study of reaction of metal powder with water at 100–110°C in presence of various salts, pH values above 9.5 increased the rate of hydrogen evolution [7]. Hazards of storing aluminium powders and pastes in contact with water under alkaline or acid conditions were studied [8]. Investigations of the explosion mechanism when molten aluminium contacts water have been described [9,10,11], and plant incidents since 1940 are presented [11,12]. A study of hazards of transport of molten aluminium in possibly wet ‘torpedoes’ is available [14]. Investigation of aluminium alloyed with lithium or calcium revealed a more energetic explosion of the melt with water, but did not conclusively demonstrate easier initiation [15]. The use of organic coatings on aluminium or magnesium powder in pyrotechnic compositions prevents reaction with atmospheric moisture and problems resulting from hydrogen evolution [16].

See other CORROSION INCIDENTS, MOLTEN METAL EXPLOSIONS

Zinc

MCA Case History No. 1722

Ball-milling aluminium–zinc (not stated if alloy or mixture) with inadequate inerting arrangements led to fires during operation or discharge of the mill.

See other METALS, REDUCANTS

0049. Aluminium–cobalt alloy (Raney cobalt alloy)

[37271-59-3] 50:50; [12043-56-0] Al₅Co; [73730-53-7] Al₂Co Al–Co

Al–Co

The finely powdered Raney cobalt alloy is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

0050. Aluminium–copper–zinc alloy (Devarda’s alloy)

[8049-11-4] Al–Cu–Zn

Al–Cu–Zn

See DEVARDA’S ALLOY

Silver nitrate: Ammonia, etc.

See other ALLOYS

0051. Aluminium amalgam (Aluminium–mercury alloy)

[[12003-69-9] (1:1)] Al–Hg

Al–Hg

1. Neely, T. A. *et al.*, *Org. Synth.*, 1965, **45**, 109

2. Calder, A. *et al.*, *Org. Synth.*, 1975, **52**, 78

The amalgamated aluminium wool remaining from preparation of triphenylaluminium will rapidly oxidise and become hot on exposure to air. Careful disposal is necessary [1]. Amalgamated aluminium foil may be pyrophoric and should be kept moist and used immediately [2].

See other ALLOYS

0052. Aluminium–lithium alloy

[103760-93-6] (7:1 atom ratio, 3.7% Li)

Al–Li

Al–Li

Water

Jacoby, J. E. *et al.*, Can. Pat. CA 1 225 8166, 1987 (*Chem. Abs.*, 1988, **108**, 42436)

The molten alloy (2–3% Li) explodes violently if cooled by contact with water, but not if cooled by ethylene glycol containing less than 25% of water.

See other ALLOYS

0053. Aluminium–magnesium alloy

[12042-38-5] (1:3)

Al–Mg

Al–Mg

The finely powdered alloy is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

Barium nitrate

See Barium nitrate, Aluminium–magnesium alloy

Iron(III) oxide, Water

Maischak, K. D. *et al.*, *Neue Huette*, 1970, **15**, 662–665

Accidental contact of the molten alloy (26% Al) with a wet rusty iron surface caused violent explosions with brilliant light emission. Initial evolution of steam, causing fine dispersion of the alloy, then interaction of the fine metals with rust in a ‘thermite’ reaction, were postulated as likely stages. Direct interaction of the magnesium (74%) with steam may also have been involved.

See other THERMITE REACTIONS

Water

Long, G. C., *Spectrum* (Pretoria), 1980, **18**, 30

In a school demonstration, an alloy pencil sharpener body (ignited in a flame) continues to burn vigorously in steam.

See other ALLOYS

0054. Aluminium–magnesium–zinc alloy

[]

Al–Mg–Zn

Al–Mg–Zn

Rusted steel

Yoshino, H. *et al.*, *Chem. Abs.*, 1966, **64**, 14017j

Impact of an alloy containing 6% Al and 3% Zn with rusted steel caused incendive sparks which ignited LPG–air mixtures in 11 out of 20 attempts.

See IGNITION SOURCES, THERMITE REACTIONS

0055. Aluminium–nickel alloys (Raney nickel alloys)

[12635-29-9]

Al–Ni

Al–Ni

HCS 1980, 807

Water

Anon., *Angew. Chem. (Nachr.)*, 1968, **16**, 2

Heating moist Raney nickel alloy containing 20% aluminium in an autoclave under hydrogen caused the aluminium and water to interact explosively, generating 1 k bar pressure of hydrogen.

See also HYDROGENATION CATALYSTS See other ALLOYS

0056. Aluminium–titanium alloys

[12003-96-2] (1:1)

Al–Ti

Al–Ti

Oxidants

Mellor, 1941, Vol. 7, 20–21

Alloys ranging from Al₃Ti₂ to Al₄Ti have been described, which ignite or incandesce on heating in chlorine; or bromine or iodine vapour; (or hydrogen chloride); or oxygen.

See other ALLOYS

0057. Tetraiodoarsonium tetrachloroaluminate

[124687-11-2]

AlAsCl₄I₄

AsI₄⁺AlCl₄⁻

Klapötke, T. *et al.*, *Angew. Chem. (Int.)*, 1989, **101**(12), 1742

Decomposes spontaneously at –78°C. Low temperature photolysis led to violent explosions on warming.

See other IODINE COMPOUNDS

0058. Aluminium tetrahydroborate

[16962-07-5]

AlB₃H₁₂

Al[BH₄]₃

1. Schlessinger, H. I. *et al.*, *J. Amer. Chem. Soc.*, 1940, **62**, 3421

2. Badin, F. J. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 2950

The vapour is spontaneously flammable in air [1], and explodes in oxygen, but only in presence of traces of moisture [2].

Alkenes, Oxygen

Gaylord, 1956, 26

The tetrahydroborate reacts with alkenes and, in presence of oxygen, combustion is initiated even in absence of moisture. Butene explodes after an induction period, while butadiene explodes immediately.

See other INDUCTION PERIOD INCIDENTS

Dimethylaminoborane

Burg, A. B. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 953

Reaction with dimethylaminoborane or dimethylaminodiborane gives an oily mixture which ignites in air and reacts violently with water.

Water

Semenenko, K. N. *et al.*, *Russ. Chem. Rev.*, 1973, 4

Interaction at ambient temperature is explosive.

See other COMPLEX HYDRIDES

0059. Aluminium tetraazidoborate

[67849-01-8]

$\text{AlB}_3\text{N}_{36}$

$\text{Al}[\text{B}(\text{N}_3)_4]_3$

Mellor, 1967, Vol. 8, Suppl. 2, 2

A very shock sensitive explosive, containing nearly 90 wt% of nitrogen.

See other HIGH-NITROGEN COMPOUNDS, NON-METAL AZIDES

0060. Aluminium bromide

[7727-15-3]

AlBr_3

AlBr_3

Dichloromethane

Kramer, G. M. *et al.*, *Acc. Chem. Res.*, 1986, **19**(3), 78–84

Solutions of aluminium bromide in dichloromethane used as a catalyst in hydride-transfer equilibrium experiments should be kept cold, as a potentially dangerous exothermic halide exchange reaction occurs on warming.

Water

Nicholson, D. G. *et al.*, *Inorg. Synth.*, 1950, **3**, 35

The anhydrous bromide should be destroyed by melting and pouring slowly into running water. Hydrolysis is very violent and may destroy the container if water is added to it.

See other METAL HALIDES

0061. Aluminium dichloride hydride diethyl etherate
[13497-97-7] (solvent-free)

$\text{AlCl}_2\text{H}\cdot\text{C}_4\text{H}_{10}\text{O}$

$\text{Cl}_2\text{HAlOEt}_2$

Dibenzyl ether

Marconi, W. *et al.*, *Ann. Chim.*, 1965, **55**, 897

During attempted reductive cleavage of the ether with aluminium dichloride hydride etherate an explosion occurred. Peroxides may have been present in the susceptible ether.

See other PEROXIDATION INCIDENTS

See related COMPLEX HYDRIDES

0062. Aluminium chloride
[7446-70-0]

AlCl_3

AlCl_3

(MCA SD-62, 1956); NSC 435, 1978; HCS 1980, 130; RSC Lab. Hazards Data Sheet No 72, 1988

1. Popov, P. V., *Chem. Abs.*, 1947, **41**, 6723d
2. Kitching, A. F., *School Sci. Rev.*, 1930, **12**(45), 79
3. Bailar, 1973, Vol. 1, 1019

Long storage of the anhydrous salt in closed containers caused apparently spontaneous decomposition and occasional explosion on opening [1,2]. This seems likely to have arisen from slow diffusion of moisture (MW 18) in through the closure and pressurisation of the container by the liberated HCl which because of its higher MW (36.5) would diffuse out at a slower rate. The need is emphasised for care in experiments in which the chloride is heated in sealed tubes. High internal pressure may be generated, not only by its vapour pressure and pressure of desorbed hydrogen chloride, but also by the near doubling in volume which occurs when the chloride melts to the monomer [3].

Alkenes

Jenkins, P. A., private. comm., 1975

Mixtures of C_4 alkene isomers (largely isobutene) are polymerised commercially in contact with low levels of aluminium chloride (or other Lewis acid) catalysts. The highly exothermic runaway reactions occasionally experienced in practice are caused by events leading to the production of high local levels of catalyst. Rapid increases in temperature and pressure of 160°C and 18 bar, respectively, have been observed experimentally when alkenes are brought into contact with excess solid aluminium chloride. The runaway reaction appears to be more severe in the vapour phase, and a considerable amount of catalytic degradation contributes to the overall large exotherm.

See other POLYMERISATION INCIDENTS, RUNAWAY REACTIONS

Aluminium oxide, Carbon oxides

See Aluminium: Aluminium halides, Carbon oxides

Aluminium, Sodium peroxide

See Sodium peroxide: Aluminium, etc.

Benzene, Carbon tetrachloride

Nolan, 1983, Case history 105

Triphenylmethyl chloride was manufactured by a Friedel-Crafts reaction of benzene and carbon tetrachloride with excess aluminium chloride. Owing to an operating fault, all the carbon tetrachloride was added to the other reactants without agitation but with water cooling. When the agitator was started, two explosions followed. The first was from the sudden exotherm and gas evolution on mixing the reactants, and this damaged the heat exchangers and allowed ingress of cooling water, leading to a second explosion on contact with the aluminium chloride.

See GAS EVOLUTION INCIDENTS (reference 2); and Water, below

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

Benzoyl chloride, Naphthalene

Clar, E. *et al.*, *Tetrahedron*, 1974, **30**, 3296

During preparation of 1,5-dibenzoylnaphthalene, addition of aluminium chloride to a mixture of benzoyl chloride and naphthalene must be effected above the m.p. of the mixture to avoid a violent reaction.

Ethylenimine, Substituted anilines

See Ethylenimine, Aluminium chloride, etc.

Ethylene oxide

See Ethylene oxide: Contaminants

Nitrobenzene

1. Riethman, J. *et al.*, *Chem. Ing. Tech.*, 1974, **48**, 729

2. Eigenmann, K. *et al.*, *Proc. Int. Symp. Simulation*, 436, Zurich, Acta Press, 1975

3. Finck, P. *et al.*, *Proc. Symp. Safety Chem. Ind.*, 17–25, Mulhouse, 1978

4. See entry SELF-ACCELERATING REACTIONS

5. Nolan, 1983, Case history 104

Mixtures of nitrobenzene and aluminium chloride are thermally unstable and may lead to explosive decomposition. Subsequent to an incident involving rupture of a 4000 l vessel, the decomposition reaction was investigated and a 3-stage mechanism involving formation and subsequent polymerisation of 2- and 4-chloronitrosobenzene was proposed [1]. Further chemical and thermodynamic work [2,3] on the thermal degradation of nitrobenzene–aluminium chloride addition compounds formed in Friedel-Crafts reactions shows that it is characterised by a slow multi-step decomposition reaction above 90°C, which self-accelerates with high exothermicity producing azo- and azoxy-polymers. Simulation of the original incident suggested that the pressure in the reactor probably increased from 3 to 40 bar in 5 s [4]. Nitrobenzene was added to a 2.2 kl reactor in readiness for a Friedel-Crafts reaction, but it leaked away through a faulty bottom valve. Aluminium chloride and a reactant, hexamethyltetralin were added, but when the agitator stopped (solid complex formation), the absence of the nitrobenzene solvent was realised. More nitrobenzene was being sucked in to the reactor when the exotherm from the reaction with aluminium chloride pressurised

the reactor, boiled and ignited the limited amount of nitrobenzene, and shattered a glass fitting [5].

See other AGITATION INCIDENTS, POLYMERISATION INCIDENTS

Nitrobenzene, Phenol

Anon., *Chem. Eng. News*, 1953, **31**, 4915

Addition of aluminium chloride to a large volume of recovered nitrobenzene containing 5% of phenol caused a violent explosion. Experiment showed that mixtures containing all 3 components reacted violently at 120°C.

Nitromethane

See Aluminium chloride–Nitromethane

Other reactants

Yoshida, 1980, 52

MRH values calculated for 12 combinations with oxidants are given.

Oxygen difluoride

See Oxygen difluoride: Halogens, etc.

Phenyl azide

See Phenyl azide: Lewis acids

Perchlorylbenzene

See Perchlorylbenzene: Aluminium chloride

Sodium tetrahydroborate

See Sodium tetrahydroborate: Aluminium chloride, etc.

Water

1. Anon., *Ind. Eng. Chem. (News Ed.)*, 1934, **12**, 194

2. Anon., *Loss Prev. Bull.*, 1980, (035), 9

An unopened bottle of anhydrous aluminium chloride erupted when the rubber bung with which it was sealed was removed. The accumulation of pressure was attributed to absorption of moisture by the anhydrous chloride before packing (but see comment following text of masthead reference 2). The presence of an adsorbed layer of moisture in the bottle used for packing may have contributed. Reaction with water is violently exothermic [1]. The unsuspected presence of 100 kg of the chloride in a vessel led to bursting of a glass vent when the vessel was being flushed with water [2].

See other GLASS INCIDENTS, GAS EVOLUTION INCIDENTS

See other METAL HALIDES

0063. Aluminium chloride–nitromethane

[3495-54-3]

$\text{AlCl}_3 \cdot \text{CH}_3\text{NO}_2$

$\text{AlCl}_3 \cdot \text{CH}_3\text{NO}_2$

An alkene

Cowen, F. M. *et al.*, *Chem. Eng. News*, 1948, **26**, 2257

A gaseous alkene was passed into a cooled autoclave containing the complex, initially with agitation, and later without. Later, when the alkene was admitted to a pressure of

5.6 bar at 2°C, a slight exotherm occurred, followed by an explosion. The autoclave contents were completely carbonised. Mixtures of ethylene, aluminium chloride and nitromethane had exploded previously, but at 75°C.

See Ethylene: Aluminium chloride, Catalysts

Carbon monoxide, Phenol

Webb, H. F., *Chem. Eng. News*, 1977, **55**(12), 4

An attempt to formylate phenol by heating a mixture with nitromethane and aluminium chloride in an autoclave under carbon monoxide at 100 bar pressure at 110°C led to a high-energy explosion after 30 min.

See related NITROALKANES

0064. Trihydrazinealuminium perchlorate

[85962-45-4]

$\text{AlCl}_3\text{H}_{12}\text{N}_6\text{O}_{12}$

$[(\text{H}_4\text{N}_2)_3\text{Al}][\text{ClO}_4]_3$

Ind. Inst. Sci., Indian Pat. IN 150422, 1982

Useful as a propellant oxidant.

See other AMMINEMETAL OXOSALTS, REDOX COMPOUNDS

0065. Aluminium chlorate

[15477-33-5]

AlCl_3O_9

$\text{Al}(\text{ClO}_3)_3$

Sidgwick, 1950, 428

During evaporation, its aqueous solution evolves chlorine dioxide, and eventually explodes.

See other METAL CHLORATES

0066. Aluminium perchlorate

[14452-39-2]

$\text{AlCl}_3\text{O}_{12}$

$\text{Al}(\text{ClO}_4)_3$

Dimethyl sulfoxide

See Dimethyl sulfoxide: Metal oxosalts

Hydrazine

See Trihydrazinealuminium perchlorate

See other METAL PERCHLORATES

0067. Caesium hexahydroaluminate(3-)

[53436-80-9]

AlCs_3H_6

$\text{Cs}_3[\text{AlH}_6]$

See Potassium hexahydroaluminate(3-)

See other COMPLEX HYDRIDES

0068. Copper(I) tetrahydroaluminate
[62126-20-9]

AlCuH₄

Cu[AlH₄]

Aubry, J. *et al.*, *Compt. rend.*, 1954, **238**, 2535

The unstable hydride decomposed at -70°C , and ignited on contact with air.

See other COMPLEX HYDRIDES

0069. Lithium tetra-deuteroaluminate
[14128-54-2]

AlD₄Li

Li[AlD₄]

Leleu, Cahiers, 1977, (86), 99

It may ignite in moist air.

See related COMPLEX HYDRIDES

0070. Aluminium hydride
[7784-21-6]

AlH₃

AlH₃

1. Mirviss, S. B. *et al.*, *Ind. Eng. Chem.*, 1961, **54**(1), 54A

2. Gibson, 1968, 66

It is very unstable and has been known to decompose spontaneously at ambient temperature with explosive violence. Its complexes (particularly the diethyl etherate) are considerably more stable [1]. The hydride ignites in air with or without oxygen enrichment [2].

See Aluminium hydride–trimethylamine

Aluminium hydride–diethyl ether

Carbon dioxide, Methyl ethers

Barbaras, G. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 877

Presence of carbon dioxide in solutions of the hydride in dimethyl or bis(2-methoxyethyl) ether can cause a violent decomposition on warming the residue from evaporation. Presence of aluminium chloride tends to increase the vigour of decomposition to explosion. Lithium tetrahydroaluminate may behave similarly, but is generally more stable.

See Lithium tetrahydroaluminate (reference 8)

Carbon dioxide, or Sodium hydrogen carbonate

Thompson, B. T. *et al.*, *Polyhedron*, 1983, **2**, 619–621

At elevated temperatures, the hydride reduces carbon dioxide or sodium hydrogen-carbonate to methane and ethane. The latter are probably the explosive reaction products produced when carbon dioxide extinguishers are used on LAH fires.

See other GAS EVOLUTION INCIDENTS

Tetrazole derivatives

Fetter, N. R. *et al.*, US Pat. 3 396 170, 1968

The 1:1 complexes arising from interaction of the hydride (as a complex with ether or trimethylamine) and various tetrazole derivatives are explosive. Tetrazoles mentioned are 2-methyl-, 2-ethyl-, 5-ethyl-, 2-methyl-5-vinyl-, 5-amino-2-ethyl-, 1-alkyl-5-amino-, and 5-cyano-2-methyl-tetrazole.

See TETRAZOLES

See other METAL HYDRIDES

0071. Alane–*N,N*-dimethylethylamine ((*N,N*-Dimethylethanamine)trihydroaluminium)

[124330-23-0]

$\text{AlH}_3 \cdot \text{C}_4\text{H}_{11}\text{N}$

$\text{H}_3\text{Al} \cdot \text{Me}_2\text{NEt}$

Frigo, D. M., *et al.*, *Chem. Mater.*, 1994, **6**(2), 190

Not generally pyrophoric, unless dropped on vermiculite, the complex decomposes slowly in the liquid phase to generate considerable pressures of hydrogen. It appears much more stable in the vapour phase.

See related METAL HYDRIDES

0072. Aluminium hydride–trimethylamine

[17013-07-9]

$\text{AlH}_3 \cdot \text{C}_3\text{H}_9\text{N}$

$\text{H}_3\text{Al} \cdot \text{NMe}_3$

Water

Ruff, J. K., *Inorg. Synth.*, 1967, **9**, 34

It ignites in moist air and is explosively hydrolysed by water.

See related METAL HYDRIDES

0073. Aluminium hydride–diethyl ether

[26351-01-9]

$\text{AlH}_3 \cdot \text{C}_4\text{H}_{10}\text{O}$

$\text{H}_3\text{Al} \cdot \text{OEt}_2$

Water

Schmidt, D. L. *et al.*, *Inorg. Synth.*, 1973, **14**, 51

Interaction of the solid with water or moist air is violent and may be explosive.

See related METAL HYDRIDES

0074. Aluminium hydroxide

[21645-51-2]

AlH_3O_3

$\text{Al}(\text{OH})_3$

Chlorinated rubber

See CHLORINATED RUBBER: Metal oxides or hydroxides

0075. Lithium tetrahydroaluminate (Lithium aluminium hydride)
[16853-85-3]

AlH₄Li

Li[AlH₄]

HCS 1980, 593; RSC Lab. Hazard Data Sheet No. 5, 1982

1. Augustine, 1968, 12
2. Gaylord, 1956, 37
3. Sakaliene, A. *et al.*, *Chem. Abs.*, 1970, **73**, 19115
4. *MCA Case History No. 1832*
5. Walker, E. R. H., *Chem. Soc. Rev.*, 1976, **5**, 36
6. Brendel, G., *Chem. Eng. News*, 1979, **57**(36), 5
7. Anon., *Lab. Haz. Bull.*, 1985, (12), item 852
8. Green, M. L. H., private comm., 1986
9. Schatzschneider, U., Internet, 1996

Care is necessary in handling this powerful reductant, which may ignite if lumps are pulverised with a pestle and mortar, even in a dry box [1]. An actual explosion destroying the mortar, has been claimed as the result of attempting to grind down large pellets, though contributory factors seem probable [9]. A rubber mallet is recommended for breaking up lumps [2]. The explosive thermal decomposition of the aluminate at 150–170°C is due to its interaction with partially hydrolysed decomposition products [3]. A spilled mixture with ether ignited after the ether evaporated [4]. Sodium bis(2-methoxyethoxy)dihydroaluminate, which is of similar reducing capability to lithium tetrahydroaluminate, is safer in that it does not ignite in moist air or oxygen and is stable at 200°C [5]. A detailed comment on the latter states that the commercial (crystalline) lithium compound is not pyrophoric, even in contact with moist air of high humidity [6]. An attempt to decontaminate a polythene bag dusted with residual lithium tetrahydroaluminate by immersion in ethyl acetate caused a fire. Two alternative methods are proposed, one for light coatings of dust, which should be immersed completely in a large volume of water behind screens until reaction ceases. The other method for larger amounts involves suspension of the residue in an inert solvent (light petroleum—*flammable*), and dropwise treatment with ethyl acetate until reaction ceases. After standing, the treatment is repeated with ethanol, and then water [7]. Following the investigation of a laboratory explosion, precautions essential for the safe use of lithium aluminium hydride (LAH) have been defined [8]. The measures given below are to prevent overheating of the hydride and its dissociation to finely divided aluminium, which can then undergo thermite-like reactions with compounds or solvents containing combined oxygen or halogen.

a. All apparatus and reactants should be perfectly dry, and reactions should be run rigorously under nitrogen, with the reaction temperature below 60°C at all times.

b. Order of addition is important. Always first add the hydride to the solvent in the nitrogen-purged apparatus, before adding the other reactant last.

AlH₄Li

c. The hydride should never be allowed to form a crust above the level of the liquid or to settle to the bottom, so efficient but gentle stirring is absolutely essential.

d. To prevent local overheating of the reaction vessel, heating mantles should never be used: always use an oil bath as heat source.

e. After reduction has been effected, destroy excess LAH by slow and careful addition of dry ethyl acetate (preferably diluted with inert solvent), again under nitrogen and keeping the temperature below 60°C. All LAH reactions should be carried out behind suitable protective screens.

See Ethyl acetate, below

Alkyl benzoates

1. Tados, W. *et al.*, *J. Chem. Soc.*, 1954, 2353

2. Field, B. O. *et al.*, *J. Chem. Soc.*, 1955, 1111

Application of a method for reducing benzaldehydes to the corresponding alcohols with a fourfold excess of the aluminate [1] to alkyl benzoates proved to be difficult to control and frequently dangerous [2].

Bis(2-methoxyethyl) ether

1. Watson, A. R., *Chem. & Ind.*, 1964, 665

2. Adams, R. M., *Chem. Eng. News*, 1953, **31**, 2334

3. Barbaras, G. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 877

4. *MCA Case History No. 1494*

5. Author's comment, 1987

The peroxide-free ether, being dried by distillation at 162°C under inert atmosphere at ambient pressure, exploded violently when the heating bath temperature had been raised to 200°C towards the end of distillation. This was attributed to local overheating of an insulating crust of hydride in contact with oxygen-containing organic material [1]. Two previous explosions were attributed to peroxides [2] and the high solubility of carbon dioxide in such ethers [3]. Stirring during distillation would probably prevent crust formation. Alternatively, drying could be effected with a column of molecular sieve or activated alumina. During distillation of the solvent from the aluminate at 100°C at ambient pressure, the flask broke and the contents ignited explosively. The aluminate decomposes at 125–135°C [4]. It seems probable that at least some of these incidents are better explained as arising from formation of finely divided aluminium from overheating of the hydride, and a thermite-like reaction with the oxygen-containing solvent [5]. See initial reference 8 above for precautions to avoid this.

See Aluminium hydride: Carbon dioxide

Boron trifluoride diethyl etherate

1. Scott, R. B., *Chem. Eng. News*, 1967, **45**(28), 7; **45**(21), 51

2. Shapiro, I. *et al.*, *J. Amer. Chem. Soc.*, 1952, **74**, 90

Use of lumps of the solid aluminate, rather than its ethereal solution, and of peroxide-containing etherate [1], rather than the peroxide-free material specified [2], caused an explosion during the attempted preparation of diborane.

Dibenzoyl peroxide

MRH 4.39/85

See Dibenzoyl peroxide: Lithium tetrahydroaluminate

3,5-Dibromocyclopentene

Johnson, C. R. *et al.*, *Tetrahedron Lett.*, 1964, **45**, 3327

Preparation of the 4-bromo compound by partial debromination of crude 3,5-dibromocyclopentene by addition of its ethereal solution to the aluminate in ice-cold ether is hazardous. Explosions have occurred on 2 occasions about 1 h after addition of dibromide.

1,2-Dimethoxyethane

1. *MCA Case History No. 1182*

2. Hoffman, K. A. *et al.*, *Org. Synth.*, 1968, **48**, 62

The finely powdered aluminate was charged through a funnel into a nitrogen-purged flask. When the solvent was added through the same funnel, ignition occurred, possibly due to local absence of purge gas in the funnel caused by turbulence [1]. Distillation of the solvent from the solid must not be taken to dryness, to avoid explosive decomposition of the residual aluminate [2].

Dioxane

1. Anon., private comm., 1976

2. Author's comments, 1987

Dioxane was purified by distillation from the complex hydride in a glass still, and when the residue was cooling down, a severe explosion and fire occurred [1]. This may have been caused by ingress of air into the cooling dioxane vapour (flammability limits 2–22%), then subsequent oxidative heating of finely divided hydride deposited on the upper parts of the still on contact with air to above the rather low autoignition temperature of dioxane (180°C). Nitrogen purging will render the operation safe. However, the alternative explanation of a thermite-like reaction with dioxane, arising from overheating of the hydride (initial reference 8 above) may be equally probable. If this were the case, nitrogen purging alone would not necessarily render the operation safe [2]. See reference 8 above for preventive measures.

Ethyl acetate

1. Bessant, K. H. C., *Chem. & Ind.*, 1957, 432

2. Yardley, J. T., *Chem. & Ind.*, 1957, 433

Following a reductive dechlorination in ether, a violent explosion occurred when ethyl acetate was added to decompose excess aluminate [1]. Ignition was attributed to the strongly exothermic reaction occurring when undiluted (and reducible) ethyl acetate contacts the solid aluminate. Addition of a solution of ethyl acetate in inert solvent or of a moist unreactive solvent to destroy excess reagent is preferable [2].

See reference 1–7, above

Fluoroamides

1. Karo, W., *Chem. Eng. News*, 1955, **33**, 1368

2. Reid, T. S. *et al.*, *Chem. Eng. News*, 1951, **29**, 3042

AlH₄Li

The reduction of amides of fluorocarboxylic acids with the tetrahydroaluminate appears generally hazardous at all stages. During reduction of *N*-ethylheptafluorobutyramide in ether, violent and prolonged gas evolution caused a fire. Towards the end of reduction of trifluoroacetamide in ether, solid separated and stopped the stirrer. Attempts to restart the stirrer by hand caused a violent explosion [1]. During decomposition by water of the reaction complex formed by interaction with tetrafluorsuccinamide in ether, a violent explosion occurred. Reaction complexes similarly obtained from trifluoroacetic acid, heptafluorobutyramide and octafluoroadipamide also showed instability, decomposing when heated. General barricading of all reductions of fluorocompounds with lithium tetrahydroaluminate is recommended [2].

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

Hydrogen peroxide

See Hydrogen peroxide: Lithium tetrahydroaluminate

Nitrogen

1. Metts, L., *Chem. Eng. News*, 1981, **59**(31), 3
2. Grossman, M. I., *Chem. Eng. News*, 1981, **59**(41), 57
3. Deberitz, J., *Chem. Eng. News*, 1982, **60**(11), 3

The incident which originally described the glowing residue from combustion in air of a pellet of hydride as being able to burn under nitrogen [1] was later interpreted as 'ignition of the hydride in a nitrogen atmosphere' [2]. It is later stressed that the hydride itself does not react directly with nitrogen under any foreseeable conditions of normal use. After combustion, the residue of hot aluminium and lithium metals would be expected to react with nitrogen [3].

Nitromethane

See Nitromethane: Lithium tetrahydroaluminate

Pyridine

Augustine, 1968, 22–23

Addition of the aluminate (0.5 g) to pyridine (50 ml) must be effected very slowly with cooling. Addition of 1 g portions may cause a highly exothermic reaction.

Tetrahydrofuran

Moffett, R. B., *Chem. Eng. News*, 1954, **32**, 4328

The solvent had been dried over the aluminate and then stored over calcium hydride for 2 years 'to prevent peroxide formation'. Subsequent addition of more aluminate caused a strong exotherm and ignition of liberated hydrogen. Calcium hydride does not prevent peroxide formation in solvents.

See other PEROXIDATION INCIDENTS

1,2,3,4-Tetrahydro-1-naphthyl hydroperoxide

See 1,2,3,4-Tetrahydro-1-naphthyl hydroperoxide: Lithium tetrahydroaluminate

Water

1. Leleu, *Cahiers*, 1977, (86), 100
2. Anon., *Fire Prevention*, 1986, (191), 43

Interaction is very vigorous, and with limited water, incandescent [1]. An account of a serious fire caused by inadvertent addition of lithal to water [2]. In the editor's experience ignition of evolved hydrogen will occur with quantities as slight as 200 mg.

See other GAS EVOLUTION INCIDENTS

See other COMPLEX HYDRIDES

0076. Sodium tetrahydroaluminate (Sodium aluminium hydride)

[13770-96-2]



Tetrahydrofuran

Del Giudica, F. P. *et al.*, *Chem. Eng. News*, 1961, **39**(40), 57

During synthesis from its elements in tetrahydrofuran, a violent explosion occurred when absorption of hydrogen had stopped. This was attributed to deposition of solid above the liquid level, overheating and reaction with solvent to give butoxyaluminium hydrides. Vigorous stirring and avoiding overheating are essential.

Water

Gibson, 1969, 85

It may ignite and explode in contact with water.

See other COMPLEX HYDRIDES

0077. Potassium hexahydroaluminate(3-)

[17083-63-5]



Ashby, E. C., *Chem. Eng. News*, 1969, **47**(1), 9

A 20 g sample, prepared and stored in a dry box for several months, developed a thin crust of oxidation/hydrolysis products. When the crust was disturbed, a violent explosion occurred, later estimated as equivalent to 230 g TNT. A weaker explosion was observed with potassium tetrahydroaluminate. The effect was attributed to superoxidation of traces of metallic potassium, and subsequent interaction of the hexahydroaluminate and superoxide after frictional initiation. Precautions advised include use of freshly prepared material, minimal storage in a dry diluent under an inert atmosphere and destruction of solid residues. Potassium hydrides and caesium hexahydroaluminate may behave similarly, as caesium also superoxidises in air.

See other COMPLEX HYDRIDES

0078. Aluminium phosphinate

[24704-64-1]



Mellor, 1971, Vol. 8, Suppl. 3, 623

It decomposes at around 220°C liberating spontaneously flammable phosphine.
See other METAL PHOSPHINATES

0079. Aluminium iodide
[7784-23-8]



Alone, or Water

Bailar, 1973, Vol. 1, 1023

It reacts violently with water, and on heating produces flammable vapour which may explode if mixed with air and ignited.

Aluminium, Carbon oxides

See Aluminium: Aluminium halides, Carbon oxides

See other METAL HALIDES

0080. Aluminium–lanthanum–nickel alloy
[66459-02-7]



Mendelsohn, M. H. *et al.*, *Inorg. Synth.*, 1983, **22**, 99

The alloy powder (used in hydrogen-storage systems) may occasionally be pyrophoric after hydriding–dehydriding operations, igniting when placed on a combustible surface (e.g. weighing paper).

See other ALLOYS, PYROPHORIC ALLOYS

0081. Lithium tetraazidoaluminate
[67849-02-9]



Mellor, 1967, Vol. 8, Suppl. 2, 2

A shock-sensitive explosive.

See related METAL AZIDES *See other* HIGH-NITROGEN COMPOUNDS

0082. Aluminium azide
[39108-14-0]



Brauer, 1963, Vol. 1, 829

Containing 82.3% of nitrogen, it may be detonated by shock.

See other HIGH-NITROGEN COMPOUNDS, METAL AZIDES

0083. Aluminium phosphide
[20859-73-8]

AIP

AIP

HCS 1980, 133

Mineral acids

1. Wang, C. C. *et al.*, *J. Inorg. Nucl. Chem.*, 1963, **25**, 327

2. Mellor, 1971, Vol. 8, Suppl. 3, 306

Evolution of phosphine is slow in contact with water or alkali, but explosively violent in contact with dilute mineral acids [1]. However, reports of violent interaction with concentrated or dilute hydrochloric acid, and of explosive reaction with 1:1 aqua regia, have been questioned [2].

See other GAS EVOLUTION INCIDENTS

See other METAL NON-METALLIDES

0084. Aluminium copper(I) sulfide
[12003-23-5]

(S:AlSCu)₂

Al₂Cu₂S₄

Preparative hazard

See Aluminium: Copper, Sulfur

See other METAL SULFIDES

0085. Magnesium tetrahydroaluminate
[17300-62-8]

Mg[AlH₄]₂

Al₂H₈Mg

Gaylord, 1956, 25

It is similar to the lithium salt.

See other COMPLEX HYDRIDES

0086. Manganese(II) tetrahydroaluminate
[65776-39-8]

Mn[AlH₄]₂

Al₂H₈Mn

Aubry, J. *et al.*, *Compt. rend.*, 1954, **238**, 2535

The unstable hydride decomposes at -80°C and ignites in contact with air.

See other COMPLEX HYDRIDES

0087. Aluminium oxide ('Alumina')
[1344-28-1]

Al₂O₃

Al₂O₃

HCS 1980, 132

Chlorine trifluoride
See Chlorine trifluoride: Metals, etc.

Ethylene oxide
See Ethylene oxide: Contaminants

Halocarbons, Heavy metals
Burbidge, B. W., unpublished information, 1976
It is known that alumina is chlorinated exothermically at above 200°C by contact with halocarbon vapours, and hydrogen chloride, phosgene etc. are produced. It has now been found that a Co/Mo–alumina catalyst will generate a substantial exotherm in contact with vapour of carbon tetrachloride or 1,1,1-trichloroethane at ambient temperature in presence of air. In absence of air, the effect is less intense. Two successive phases appear to be involved: first, adsorption raises the temperature of the alumina; then reaction, presumably metal-catalysed, sets in with a further exotherm.

Oxygen difluoride
See Oxygen difluoride: Adsorbents

Sodium nitrate
See Sodium nitrate: Aluminium, etc.

Vinyl acetate
See Vinyl acetate: Desiccants
See other METAL OXIDES

0088. Dialuminium octavanadium tridecasilicide

[]

(Complex structure)



Hydrofluoric acid
Sidgwick, 1950, 833
The silicide reacts violently with the aqueous acid.
See other METAL NON-METALLIDES

0089. Cerium(III) tetrahydroaluminate

[65579-06-8]



Aubry, J. *et al.*, *Compt. rend.*, 1954, **238**, 2535
The unstable hydride decomposes at –80°C, and ignites in contact with air.
See other COMPLEX HYDRIDES

0090. Americium trichloride
[13464-46-5]

AmCl₃

AmCl₃

MCA Case History No. 1105

A multi-wall shipping container, holding 400 ml of a solution of americium chloride in a polythene bottle and sealed for over 3 months, exploded. The reason could have been a slow pressure build-up of radiolysis products. Venting and other precautions are recommended.

See Radon: Water

See other GAS EVOLUTION INCIDENTS, METAL HALIDES

0091. Argon
[7440-37-1]

Ar

Ar

Liquid nitrogen

1. Baker, K., *Chem. Brit.*, 1979, **15**, 65
2. Janiaut, H., *Actual Chim.*, 1986, (1-2), 59-61

The presence of argon in apparatus cooled by liquid nitrogen can lead to hazardous situations with the possibility of explosion. This is because argon solidifies at -189°C and has so low a vapour pressure at -196°C that the solid may survive exposure to high vacuum for a considerable period, and subsequently evaporate on warming with generation of high pressure. Use of solid CO_2 -solvent baths is recommended for argon [1]. A phenylhydrazone was vacuum-degassed and frozen in liquid nitrogen, probably during freeze-drying. Vacuum was broken by slow admission of argon (some of which would condense and solidify), and the flask, still under liquid nitrogen, was sealed. During subsequent warming to ambient temperature, the flask exploded violently from internal gas pressure [2].

See other GAS EVOLUTION INCIDENTS

Magnesium perchlorate

See Magnesium perchlorate: Argon

See other NON-METALS

0092. Arsenic
[7440-38-2]

As

As

NSC 499, 1979 (As and its inorganic compounds); *RSC Lab. Hazard Data Sheet No. 62*, 1987 (As and compounds)

Müller, W. J., *Z. Angew. Chem.*, 1914, **27**, 338

An explosive variety (or compound) of arsenic was produced as a surface layer on the exposed iron surfaces of a corroded lead-lined vessel which contained 35% sulfuric

acid with a high arsenic content. It exploded on friction or ignition, and contained no hydrogen, but variable small amount of iron and lead. It may have been analogous to explosive antimony.

Bromine azide

See Bromine azide

Halogens or Interhalogens

1. Mellor, 1946, Vol. 2, 92
2. Mellor, 1956, Vol. 2, Suppl. 1, 379

The finely powdered element inflames in gaseous chlorine or liquid chlorine at -33°C [1]. The latter is doubtful [2].

See Bromine trifluoride: Halogens, etc.

Bromine pentafluoride: Acids, etc.

Chlorine trifluoride: Metals, etc.

Iodine pentafluoride: Metals, etc.

Metals

Mellor, 1940, Vol. 4, 485–486; 1942, Vol. 15, 629; 1937, Vol. 16, 161

Palladium or zinc and arsenic react on heating with evolution of light and heat, and platinum with vivid incandescence.

See Aluminium: Antimony, etc.

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Oxidants

See Chromium trioxide: Arsenic

Dichlorine oxide: Oxidisable materials

Nitrosyl fluoride: Metals, etc.

Potassium dioxide: Metals

Potassium permanganate: Antimony, etc.

Silver nitrate: Arsenic

Sodium peroxide: Non-metals

Rubidium acetylide

See Rubidium acetylide: Non-metals

See other NON-METALS

0093. Difluoroperchloryl hexafluoroarsenate

[39003-82-2]

[F₂ClO₂] [AsF₆]

AsClF₈O₂

See entry DIFLUOROPERCHLORYL SALTS

0094. Arsenic trichloride
[7784-34-1]



HCS 1980, 165

Aluminium

See Aluminium: Non-metal halides

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

See other NON-METAL HALIDES

0095. Arsenic pentafluoride
[7784-36-3]



Benzene, Potassium methoxide

Kolditz, L. *et al.*, *Z. Anorg. Chem.*, 1965, **341**, 88–92

Interaction of the pentafluoride and methoxide proceeded smoothly in trichlorotrifluoroethane at 30–40°C, whereas in benzene as solvent repeated explosions occurred.

1,3-Butadiyne


See 1,3-Butadiyne: Arsenic pentafluoride

Krypton difluoride

See Krypton difluoride: Arsenic pentafluoride

See other NON-METAL HALIDES


0096. Fluorokrypton hexafluoroarsenate
[50859-36-2]



See Krypton difluoride: Arsenic pentafluoride

See XENON COMPOUNDS

0097. Monofluoroxonium hexafluoroarsenate
[]



Minkwitz, R. *et al.*, *Angew. Chem. Int.*, 1990, **29**(6), 689


The above salt is a powerful oxidant, reacting explosively with methanol even at –40°C. The antimonate is similar.

See other OXIDANTS

0098. Difluoroammonium hexafluoroarsenate

[56533-30-3]

AsF₈H₂N



Christe, K. O., *Inorg. Chem.*, 1975, **14**, 2821–2824

Solutions of this and the hexafluoroantimonate salt in hydrogen fluoride, kept for extended periods between –50 and +50°C, burst the Kel-F or Teflon FEP containers. This was attributed to excess pressure of hydrogen fluoride and nitrogen arising from decomposition of the salts. The variable rates of decomposition indicated catalysis by trace impurities. The salts also decompose exothermally after a short period at ambient temperature.

See other GAS EVOLUTION INCIDENTS, N-HALOGEN COMPOUNDS

0099. Trifluoroselenium hexafluoroarsenate

[59544-89-7]

AsF₉Se



Water

Bartlett, N. *et al.*, *J. Chem. Soc.*, 1956, 3423

Violent interaction.

See related NON-METAL HALIDES

†0100. Arsine

[7784-42-1]

AsH₃



HCS 1980, 167 (cylinders); *RSC Lab. Hazard Data Sheet No. 51*, 1986

1. Rüst, 1948, 301

2. Kayser, J. C., *Proc. 11th Int. Symp. Prev. Occup. Risks Chem. Ind.*, 289–315, Heidelberg, ISSA, 1987

Arsine is strongly endothermic, and can be detonated by suitably powerful initiation [1]. The potential hazards involved in setting up and operating a plant for manufacture of arsine for semi-conductor use have been studied [2].

Chlorine

See Chlorine: Non-metal hydrides

Nitric acid

See Nitric acid: Non-metal hydrides

See other ENDOTHERMIC COMPOUNDS, NON-METAL HYDRIDES

0101. Arsine–boron tribromide
[65313-32-8]



Oxidants

1. Stock, A., *Ber.*, 1901, **34**, 949
2. Mellor, 1939, Vol. 9, 57

Unlike arsine, the complex ignites on exposure to air or oxygen, even at below 0°C [1]. It is violently oxidised by nitric acid [2].

See related NON-METAL HYDRIDES, NON-METAL HALIDES

0102. Trisilylarsine
[15100-34-6]




Leleu, *Cahiers*, 1977, (88), 363

Ignites in air.

See related SILANES



0103. Triazidoarsine (Arsenous triazide)
[167771-41-7]



See entry Tetraazidoarsonium hexafluoroarsenate

See other NON-METAL AZIDES

0104. Tetraazidoarsonium hexafluoroarsenate
[171565-26-7]



Tornieporth-Oetting, I. C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(5), 511

Klapoetke, T. M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1995, (20), 3365

As might be expected, it is violently explosive, as is the precursor triazidoarsine; any other salts of this hitherto unknown arsonium ion may also prove so.

See other NON-METAL AZIDES

0105. Arsenic trioxide
[1327-53-3]



(*MCA SD-60*, 1956); *HCS 1980*, 166

Various reagents

See Chlorine trifluoride: Metals, etc.

Hydrogen fluoride: Oxides

Sodium nitrate: Arsenic trioxide, etc.

Zinc: Arsenic trioxide

See other NON-METAL OXIDES

0106. Arsenic pentaoxide
[1303-28-2]



Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

See other NON-METAL OXIDES

0107. Platinum diarsenide
[12044-52-9]



Preparative hazard

See Platinum: Arsenic

See other METAL NON-METALLIDES

0108. Arsenic trisulfide
[1303-33-9]




Oxidants

See Chloric acid: Metal sulfides

Potassium chlorate: Arsenic trisulfide

See other NON-METAL SULFIDES

0109. Tetraarsenic tetrasulfide
[12279-90-2]



Oxidants

See Chloric acid: Metal sulfides

Chlorine: Sulfides

Potassium nitrate: Metal sulfides

See other NON-METAL SULFIDES

0110. Gold
[7440-57-5]



Analytical hazard

See Gold(III) chloride, next below

Hydrogen peroxide
See Hydrogen peroxide: Metals
See other METALS

0111. Gold(III) chloride
[13453-07-1]



Ammonia and derivatives

1. Mellor, 1941, Vol. 3, 582–583
2. Sidgwick, 1950, 178
3. *491M*, 1975, 194

Action of ammonia or ammonium salts on gold chloride, oxide or other salts under a wide variety of conditions gives explosive or ‘fulminating’ gold [1]. Of uncertain composition but containing Au–N bonds, this is a heat-, friction- and impact-sensitive explosive when dry, similar to the related mercury and silver compounds [2]. In an attempt to precipitate finely divided gold from its solution in aqua regia (effectively gold chloride solution), ammonia solution was added instead of ammonium oxalate. The precipitated ‘gold’ subsequently exploded when heated in a furnace with other metals to prepare an alloy [3].

See FULMINATING GOLD

Potassium cyanate

Selig, W. S., *Microchem. J.*, 1990, **41**(3), 386

A precipitate, explosive when touched after drying, which appeared from a neutralised solution stood overnight with a large excess of potassium cyanate, was described as a fulminate. This is unlikely, though fulminating gold is probable, cyanate hydrolysing to release ammonia (see above). Existence of explosive cyanate complexes is also conceivable.

See other GOLD COMPOUNDS, METAL HALIDES

0112. Gold(III) hydroxide–ammonia

[]




1. Sorbe, 1958, 63
2. Ephraim, 1939, 463

Explosive gold, formed from the hydroxide and ammonia, is formulated as above [1]. Dry heating forms the equally explosive $\text{Au}_2\text{O}_3 \cdot 3\text{NH}_3$, then $3\text{Au}_2\text{O} \cdot 4\text{NH}_3$, while heating with water forms the more explosive $\text{Au}_2\text{O}_3 \cdot 2\text{NH}_3$.

See FULMINATING GOLD See other GOLD COMPOUNDS

0113. Sodium triazidoaurate(?)

[]




Rodgers, G. T., *J. Inorg. Nucl. Chem.*, 1958, **5**, 339–340

The material (of unknown structure, analysing as $\text{Au}_{1.5}\text{N}_9\text{Na}$ and possibly the impure title compound, explodes at 130°C .

See other GOLD COMPOUNDS, METAL AZIDES

0114. Bis(dihydroxygold)imide

[]



Mellor, 1940, Vol. 8, 259

An explosive compound.

See other GOLD COMPOUNDS, N-METAL DERIVATIVES

0115. Gold(III) oxide

[1303-58-8]



Ammonium salts

See Gold(III) chloride: Ammonia, etc.

See other ENDOTHERMIC COMPOUNDS, GOLD COMPOUNDS

0116. Gold(III) sulfide

[1303-61-3]





Silver oxide

See Silver(I) oxide: Metal sulfides

See other GOLD COMPOUNDS, METAL SULFIDES

0117. Gold(I) nitride–ammonia

[]





Raschig, F., *Ann.*, 1886, **235**, 349

An explosive compound, probably present in fulminating gold, produced from action of ammonia on gold(I) oxide.

See FULMINATING GOLD *See other* GOLD COMPOUNDS, N-METAL DERIVATIVES

0118. Gold(III) nitride trihydrate

[]



Mellor, 1940, Vol. 8, 101

Very explosive when dry.
See other GOLD COMPOUNDS, *N*-METAL DERIVATIVES

0119. Boron
[7440-42-8]

B

B

Bailar, 1973, Vol. 1, 692
Many of the previously described violent reactions of boron with a variety of reagents are ascribed to the use of impure or uncharacterised 'boron'. The general impression of the reactivity of pure boron, even when finely divided, is one of extreme inertness, except to highly oxidising agents at high temperatures.

Ammonia

Mellor, 1940, Vol. 8, 109
Boron incandescs when heated in dry ammonia, hydrogen being evolved.

Dichromates, Silicon

Howlett, S. *et al.*, *Thermochim. Acta*, 1974, **9**, 213–216
The mechanism of ignition and combustion of pyrotechnic mixtures of boron with potassium dichromate and/or sodium dichromate in presence or absence of silicon are discussed.

Rubidium acetylide

See Rubidium acetylide: Non-metals

Halogens or Interhalogens

1. Mellor, 1941, Vol. 2, 92
 2. Bailar, 1973, Vol. 1, 690–691
- Boron ignites in gaseous chlorine or fluorine at ambient temperature, attaining incandescence in fluorine [1]. Powdered boron reacts spontaneously with the halogens from fluorine to iodine at 20, 400, 600 and 700°C respectively [2].
See Bromine trifluoride: Halogens, etc.
Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Boron-containing materials
Iodine pentafluoride: Metals, etc.

Metals

Mellor, 1946, Vol. 3, 389; Vol. 5, 15
Explosive interaction when boron and lead fluoride or silver fluoride are ground together at ambient temperature.

Oxidants

See Halogens or Interhalogens, above
Calcium chromate: Boron
Dinitrogen oxide: Boron

Lead(II) oxide: Non-metals
Lead(IV) oxide: Non-metals
Nitric acid: Non-metals
Nitrogen oxide: Non-metals
Nitrosyl fluoride: Metals, etc.
Nitryl fluoride: Non-metals
Oxygen difluoride: Non-metals
Potassium nitrite: Boron
Potassium nitrate: Non-metals
Silver difluoride: Boron, etc.
Sodium peroxide: Non-metals

Water

1. Bailar, 1973, Vol. 1, 691
2. Shidlovskii, A. A., *Chem. Abs.*, 1963, **59**, 11178

Interaction of powdered boron and steam may become violent at red heat [1]. The highly exothermic reactions with water might become combustive or explosive processes at sufficiently high temperatures and pressures [2].

See other NON-METALS

0120. Boron bromide diiodide

[14355-21-6]



Water

Mellor, 1946, Vol. 5, 136

Interaction is violent, as for the tribromide or triiodide.

See other NON-METAL HALIDES

0121. Dibromoborylphosphine

[30641-57-7] (polymeric form)



491M, 1975, 146

Ignites in air.

See related NON-METAL HALIDES, NON-METAL HYDRIDES

0122. Boron tribromide

[10294-33-4]



Sodium

See Sodium: Non-metal halides (reference 7)

Tungsten trioxide

Levason, W. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1981, 2501–2507

Tungsten oxide tetrabromide was prepared by condensing a little of the bromide onto the oxide at -196°C , then allowing slow warming by immersion of the container in an ice bath. Omission of the ice bath or use of large amounts of bromide may lead to explosions.

Water

1. *BCISC Quart. Safety Summ.*, 1966, **37**, 22

2. Anon., *Lab. Pract.*, 1966, **15**, 797

Boron halides react violently with water, and particularly if there is a deficiency of water, a violent explosion may result. It is therefore highly dangerous to wash glass ampoules of boron tribromide with water under any circumstances. Following a serious accident, experiment showed that an ampoule of boron tribromide, when deliberately broken under water, caused a violent explosion, possibly a detonation. Only dry non-polar solvents should be used for cleaning or cooling purposes [1]. Small quantities of boron tribromide may be destroyed by cautious addition to a large volume of water, or water containing ice [2].

See other GLASS INCIDENTS

See other NON-METAL HALIDES

0123. Difluoroperchloryl tetrafluoroborate

[38682-34-7]

$[\text{F}_2\text{ClO}_2] [\text{BF}_4]$

BClF_6O_2

See entry DIFLUOROPERCHLORYL SALTS

0124. Dichloroborane

[10325-39-0]

Cl_2BH

BCl_2H

Bailar, 1973, Vol. 1, 742

Ignites in air.

See related NON-METAL HALIDES

0125. Dichlorodisilylaminoborane

[25573-61-9]

$\text{Cl}_2\text{BN}(\text{SiH}_3)_2$

$\text{BCl}_2\text{H}_6\text{NSi}_2$

491M, 1975, 149

Ignites in air.

See related NON-METAL HALIDES, NON-METAL HYDRIDES

0126. Boron azide dichloride

[]

N_3BCl_2

BCl_2N_3

1. Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 27

2. Paetzold, P. I., *Z. Anorg. Chem.*, 1963, **326**, 47

A hard crust of sublimed material exploded when crushed with a spatula [1]. Explosions on sublimation or during solvent removal were known previously [2].

See related NON-METAL AZIDES

0127. Boron trichloride

[10294-34-5]

BCl₃

BCl₃

HCS 1980, 215 (cylinder)

Aniline

Jones, R. G., *J. Amer. Chem. Soc.*, 1939, **61**, 1378

In absence of cooling or diluent, interaction is violent.

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Boron trichloride

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Methanol

Anon., *Safety Digest Univ. Safety Assoc.*, 1989 **34**, 16

On addition of boron trichloride, dropwise, to methanol an explosion and fire resulted. The academic reporter appears unaware that reaction of these was reported in 1834, and attributes the explosion to water in his methanol. Since the chloride is a gas, ability to add it dropwise is puzzling. Had a phonetic error meant PCl₃ was employed the fire would be intelligible, since phosphine would be formed as a pyrophoric byproduct.

Phosphine

Mellor, 1946, Vol. 5, 132

Interaction is energetic.

Triethylsilane

Matteson, D. S. *et al.*, *J. Org. Chem.*, 1990, **55**, 2274

A pressure build-up, septum expulsion and combustion was experienced on mixing these reagents at -78°C.

See other NON-METAL HALIDES

0128. Dicobalt boride

[12045-01-1]

BCo₂

Co₂B

Heinzmann, S. W. *et al.*, *J. Amer. Chem. Soc.*, 1982, **104**, 6801 (footnote 22)

The boride precipitated from sodium borohydride and cobalt(II) chloride in methanol becomes pyrophoric after vacuum drying. It can safely be stored solvent-moist.

See other METAL NON-METALLIDES, PYROPHORIC MATERIALS

0129. Boron trifluoride
[7637-07-2]



HCS 1980, 216 (cylinder)

Alkali metals, or Alkaline earth metals (not magnesium)

Merck Index, 1976, 175

Interaction hot causes incandescence.

Alkyl nitrates

See ALKYL NITRATES: Lewis acids

See other NON-METAL HALIDES

0130. Tetrafluoroboric acid
[16872-11-0]



Acetic anhydride

See Acetic anhydride: Tetrafluoroboric acid

See other INORGANIC ACIDS

0131. Nitronium tetrafluoroborate
[13826-86-3]



Tetrahydrothiophene-1,1-dioxide

See entry NITRATING AGENTS

0132. Dioxygenyl tetrafluoroborate
[12228-13-6]




Organic materials

Goetschel, C. T. *et al.*, *J. Amer. Chem. Soc.*, 1969, **91**, 4706

It is a very powerful oxidant, addition of a small particle to small samples of benzene or 2-propanol at ambient temperature causing ignition. A mixture prepared at -196°C with either methane or ethane exploded when the temperature was raised to -78°C .

See other OXIDANTS

0133. Tetrafluoroammonium tetrafluoroborate
[15640-93-4]



2-Propanol

Goetschel, C. T. *et al.*, *Inorg. Chem.*, 1972, **11**, 1700

When the fluorine used for synthesis contained traces of oxygen, the solid behaved as a powerful oxidant (causing 2-propanol to ignite on contact) and it also exploded on impact. Material prepared from oxygen-free fluorine did not show these properties, which were ascribed to the presence of traces of dioxygenyl tetrafluoroborate (above).
See other CATALYTIC IMPURITY INCIDENTS, *N*-HALOGEN COMPOUNDS

0134. Poly[borane(1)]

[13766-26-2]

(BH)_n

(BH)_n

Bailar, 1973, Vol. 1, 740

Ignites in air.

See other BORANES

0135. Borane

[13283-31-3]

BH₃

BH₃

The monomeric borane is extremely endothermic ($\Delta H_f^\circ +105.5$ kJ/mol, 7.62 kJ/g) and on formation apparently immediately dimerises to diborane (or higher boranes). It is usually stabilised as the monomer by the formation of various complexes with N, O, P or S donor molecules and many of these are available commercially.

See other BORANES, ENDOTHERMIC COMPOUNDS

0136. Borane–dimethylsulfide

[13292-87-0]

BH₃·C₂H₆S

(CH₃)₂S·BH₃

Clinton, F. L., *Chem. Eng. News*, 1992, **70**(35), 5

Bottles of this complex sometimes pressurise during ambient temperature storage. The cause, which it is hoped has now been eliminated, may be traces of thiols reacting to form hydrogen. Cold storage, careful opening and regular venting are desirable.

See other BORANES

0137. Borane–bis(2,2-dinitropropylhydrazine)

[]

BH₃·2C₃H₈N₄O₄

BH₃·2H₂NNHCH₂C(NO₂)₂Me

Gao, F. *et al.*, *Youji Huaxue*, 1984, (2), 123–124 (Ch.)

It decomposes rapidly at ambient temperature and ignites or explodes within 1–2 h.
See related BORANES, POLYNITROALKYL COMPOUNDS *See other* REDOX COMPOUNDS

0138. Borane–tetrahydrofuran

[14044-65-6]

BH₃.C₄H₈O**C₄H₈O.BH₃**

1. Bruce, M. I., *Chem. Eng. News*, 1974, **52**(41), 3
2. Hopps, H., *Chem. Eng. News*, 1974, **52**(41), 3
3. Gaines, D. F. *et al.*, *Inorg. Chem.*, 1963, **2**, 526
4. Kollonitsch, J., *Chem. Eng. News*, 1974, **52**(47), 3

A glass bottle containing a 1M solution of the complex in THF exploded after 2 weeks in undisturbed laboratory storage out of direct sunlight at 15°C [1]. The problem of pressure build-up during storage of such commercial solutions (which are stabilised with 5 mol% of sodium tetrahydroborate) at above 0°C had been noted previously, and was attributed to presence of moisture in the original containers [2]. However, by analogy with the known generation of hydrogen in tetrahydroborate–diborane–bis(2-methoxyethyl) ether systems [3], it is postulated that the tetrahydroborate content may in fact destabilise the borane–THF reagent, with generation of hydrogen pressure in the closed bottle [4]. Storage at 0°C and opening bottles behind a screen are recommended [2].

See other GLASS INCIDENTS, GAS EVOLUTION INCIDENTS

See related BORANES

0139. Borane–pyridine

[110-51-0]

BH₃.C₅H₅N**C₅H₅N.BH₃**

1. Brown, H. C. *et al.*, *J. Amer. Chem. Soc.*, 1956, **78**, 5385
2. Baldwin, R. A. *et al.*, *J. Org. Chem.*, 1961, **26**, 3550

Decomposition was rapid at 120°C/7.5 mbar [1], and sometimes violent on attempted distillation at reduced pressure [2].

See related BORANES

0140. Borane–phosphorus trifluoride

[14391-39-6]

BH₃.F₃P**H₃B.PF₃**

Mellor, 1971, Vol. 8, Suppl. 3, 442

The unstable gas ignites in air.

See related BORANES, NON-METAL HALIDES

0141. Borane–ammonia

[17596-45-1]


BH₃.H₃N**H₃B.NH₃**

Sorbe, 1968, 56

It may explode on rapid heating.

See related BORANES

0142. Borane–hydrazine
[14391-40-9]





Gunderloy, F. C., *Inorg. Synth.*, 1967, **9**, 13

It is shock-sensitive and highly flammable, like the bis(borane) adduct.


See related BORANES

0143. Bis(borane)–hydrazine
[13730-91-1]



See Borane–hydrazine *See related* BORANES

0144. Boric acid
[10043-35-3]



HCS 1980, 214

Acetic anhydride

See Acetic anhydride: Boric acid

Potassium

See Potassium: Oxidants

See other INORGANIC ACIDS

0145. Lithium tetrahydroborate (Lithium borohydride)
[16949-15-8]




Water

Gaylord, 1965, 22

Contact with limited amounts of water, either as liquid or that present as moisture in cellulose fibres, may cause ignition after a delay.

See other COMPLEX HYDRIDES

0146. Ammonium peroxoborate
[17097-12-0]



Menzel, H. *et al.*, *Österr. Chem. Z.*, 1925, **28**, 162

Explosive decomposition under vacuum.

See other PEROXOACID SALTS

0147. Sodium tetrahydroborate (Sodium borohydride)

[16940-66-2]

BH₄Na

Na[BH₄]

HCS 1980, 830; RSC Lab. Hazard Data Sheet No. 39, 1985

1. le Noble, W. J., *Chem. Eng. News*, 1983, **61**(19), 2
2. le Noble, W. J., private comm., 1983

A several year-old 100 ml glass bottle which had originally contained 25 g of the complex hydride exploded while being opened a few hours after the previous opening to remove a portion [1]. No plausible explanation can be deduced, and a dust explosion seems unlikely in view of the large initiation energy required [2].

See other GLASS INCIDENTS

Acetic acid, Dichloromethane, Methanol

Ward, D. E. *et al.*, *Tetrahedron Lett.*, 1988, **29**, 517–520

In a general method for the selective reduction of ketones in presence of conjugated enones, this is effected by the tetrahydroborate in 1:1 methanol–dichloromethane at 75°C. In favourable cases the reaction is carried out at 20°C in dichloromethane containing a little acetic acid. It should be noted that addition of acetic acid to sodium tetrahydroborate in methanol–dichloromethane leads to vigorous evolution of much hydrogen.

See other GAS EVOLUTION INCIDENTS

Acids

Bailar, 1973, Vol. 1, 768

Interaction of sodium and other tetrahydroborates with anhydrous acids (fluorophosphoric, phosphoric or sulfuric) to generate diborane is very exothermic, and may be dangerously violent with rapid mixing. Safer methods of making diborane are detailed.

Alkali

1. Anon., *Angew. Chem. (Nachr.)*, 1960, **8**, 238
2. Volpers, *Proc. 1st Int. Symp. Prev. Occup. Risks Chem. Ind.*, 188–193, Heidelberg, ISSA, 1970
3. Mikheeva, V. I. *et al.*, *Chem. Abs.*, 1969, **71**, 85064

A large volume of alkaline tetrahydroborate solution spontaneously heated and decomposed, liberating large volumes of hydrogen which burst the container. Decomposition is rapid when pH is below 10.5 [1]. A more detailed account of the investigation was published [2]. Dry mixtures with sodium hydroxide containing 15–40% of tetrahydroborate liberate hydrogen explosively at 230–270°C [3].

See other GAS EVOLUTION INCIDENTS

Aluminium chloride, Bis(2-methoxyethyl) ether

1. de Jongh, H. A. P., *Chem. Eng. News*, 1977, **55**(31), 31
2. Brown, H. C., *Chem. Eng. News*, 1977, **55**(35), 5

Addition of a 4% solution of sodium tetrahydroborate in diglyme containing 0.09% of water to a 27% solution of aluminium chloride in the same solvent led to a violent explosion, attributed to formation and ignition of hydrogen. The ignition source arose from contact of the hydroborate solution with the solid chloride, as demonstrated experimentally. Nitrogen purging is essential for all hydride reductions [1], and also for hydroboration, organoborane, Grignard and organometallic reactions generally [2]. Previous work had shown that clear solutions of the sodium tetrahydroborate–aluminium chloride reagent did not ignite in dry air, but the solid-containing reagent could lead to ignition [2].

Charcoal

Collin, P. A., Sidgwick, C., *Chemistry in Britain*, 1992, **28**(4), 324

Mixtures of charcoal and borohydride are liable to autoignition in air, the probability being higher if the charcoal has been exposed to a damp atmosphere.

See other PYROPHORIC MATERIALS

Diborane, Bis(2-methoxyethyl) ether

See Borane–tetrahydrofuran (reference 3)

Dimethylformamide

1. Yeowell, A. *et al.*, *Chem. Eng. News*, 1979, **57**(39), 4
2. Schwartz, J. *et al.*, *J. Org. Chem.*, 1993, **58**(18), 5005
3. Thiokol-Ventron announcement in *Cambrian News*, 1–2, May 1980, K & K Greeff Chemicals

Hot solutions of the tetrahydroborate (15.7% wt) in DMF will undergo a violent runaway thermal decomposition, the solid residue attaining a temperature of 310°C. The induction period depends on temperature, and is 45 h at 62, and 45 m at 90°C. In a plant-scale incident, an 83 kg batch led to a violent explosion, ascribed to spontaneous ignition of trimethylamine (AIT 190°C) produced by reduction of the solvent [1]. Further investigation confirmed these results, whether technical, reagent grade or redistilled dry solvent were used, the induction period being independent of added water, amine, metal salts or borate. However, addition of formic (or acetic) acid significantly reduced the induction period, and if the solvent containing 2.6% of formic acid is added to the hydride at ambient temperature, immediate and violent decomposition ensues, involving formation of sodium formyloxytrihydroborate. The latter is responsible for reduction of the amide solvent to trimethylamine. Traces of formic acid and dimethylamine are present in the commercial solvent from hydrolysis, and the latter is catalysed by water, acid or base. Hot solutions of the hydride in DMF above 2M concentration will soon begin to undergo hydrolysis/reduction reactions, both of which produce formic acid and/or di- or tri-methylamine, so the decomposition is autocatalytic and soon accelerates out of control, causing gross and violent reduction of the solvent. Further investigation is reported and the greater reduction powers of the intermediate exploited [2]. Dimethylacetamide does not react violently with sodium tetrahydroborate, even at 4.7M concentration, and should be considered as a substitute solvent, particularly at higher temperatures [3].

See other GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS

Glycerol

Epshtein, N. A. *et al.*, *Chem. Abs.*, 1987, **106**, 55145

Contact of a drop of glycerol with a flake of sodium tetrahydroborate leads to ignition, owing to thermal decomposition of the latter at above 200°. Other glycols and methanol also react exothermally, but do not ignite.

Palladium

See Palladium: Sodium tetrahydroborate

Ruthenium salts

Cusumano, J. A., *Nature*, 1974, **247**, 456

Use of borohydride solutions to reduce ruthenium salt solutions to the metal or an alloy gave solid products (possibly hydrides), which when dry, exploded violently in contact with water or when disturbed by a spatula. Hydrazine appears to be a safe reductant for ruthenium salt solutions.

Sulfuric acid

Pascal, 1961, Vol. 6, 337

Ignition may occur if the mixture is not cooled.

See other COMPLEX HYDRIDES

0148. Boron azide diiodide

[68533-38-0]



Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 510

Decomposes explosively in contact with water.

See related NON-METAL AZIDES, NON-METAL HALIDES

0149. Boron diiodophosphide

[12228-28-3]



Chlorine

See Chlorine: Phosphorus compounds

Metals

Mellor, 1947, Vol. 8, 845

It ignites in contact with mercury vapour or magnesium powder.

See related NON-METAL HALIDES

0150. Boron triiodide

[13517-10-7]



Though a moderately endothermic compound ($\Delta H_f^\circ +70.8$ kJ/mol), its high MW gives it the rather low specific energy content of 0.18 kJ/g.

Ammonia

Mellor, 1945, Vol. 5, 136
Strong exotherm on contact.

Phosphorus

Mellor, 1946, Vol. 5, 136
Warm red or white phosphorus reacts incandescently.

Water

1. Moissan, H., *Compt. rend.*, 1892, **115**, 204
2. Unpublished information
Violent reaction [1], particularly with limited amounts of water [2].
See other IODINE COMPOUNDS, NON-METAL HALIDES

0151. Lithium tetraazidoborate

[]



BLiN₁₂

Wiberg, E. *et al.*, *Z. Naturforsch.*, 1954, **9B**, 499
Highly explosive, sensitive to heat, impact and friction (e.g. of a spatula when removing solid from a flask).
See related NON-METAL AZIDES *See other* HIGH-NITROGEN COMPOUNDS

0152. Boron nitride

[10043-11-5]



BN

Peroxydisulfuryl difluoride
See Tetra(boron nitride) fluorosulfate

Sodium peroxide
See Sodium peroxide: Boron nitride
See related NON-METALS

0153. Boron triazide (Triazidoborane)

[21844-15-1]



BN₉

1. Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 27
2. Skillern, K. R. *et al.*, *Inorg. Chem.*, 1977, **16**, 3001
3. Miller, N. E. *et al.*, *Inorg. Chem.*, 1964, **3**, 1064
A sample of the vacuum distilled pyridine complex exploded in a heated capillary sampling tube [1]. Detonation of the trimethylamine complex [2], at or near 200°C [3] is also noted.

Diethyl ether, or Water

Wiberg, E. *et al.*, *Z. Naturforsch.*, 1954, **9B**, 498

The highly explosive material detonated in contact with ether vapour or water at ambient temperature, or with ether at -35°C (probably initiated by the heat of coordination to O).

See other NON-METAL AZIDES

0154. Sodium borate hydrogen peroxidate

[56892-92-5]

$\text{BNaO}_2 \cdot \text{H}_2\text{O}_2$

$\text{O}:\text{BONa} \cdot \text{H}_2\text{O}_2$

HCS 1980, 853


See entry CRYSTALLINE HYDROGEN PEROXIDATES

See Sodium peroxoborate (next below)

0155. Sodium peroxoborate

[7632-04-4]

BNaO_3



1. Anon., *Angew. Chem.*, 1963, **65**, 41

2. Castrantas, 1965, 5

The true peroxoborate has been reported to detonate on light friction [1]. The common 'tetrahydrate' is not a peroxoborate, but $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2 \cdot 3\text{H}_2\text{O}$, and while subject to catalytic decomposition by heavy metals and their salts, or easily oxidisable foreign matter, it is relatively stable under mild grinding with other substances [2].

See also CRYSTALLINE HYDROGEN PEROXIDATES

See other CATALYTIC IMPURITY INCIDENTS, PEROXOACID SALTS

0156. Boron phosphide

[20205-91-8]

BP

BP

Oxidants

See Nitric acid: Non-metals

Sodium nitrate: Boron phosphide

See related NON-METALS

0157. Beryllium tetrahydroborate

[17440-85-6]

B_2BeH_8

$\text{Be}[\text{BH}_4]_2$

Air, or Water

1. Mackay, 1966, 169

2. Semenenko, K. N. *et al.*, *Russ. Chem. Rev.*, 1973, 4

It ignites vigorously and often explodes in air [1], or on contact with cold water [2].
See other COMPLEX HYDRIDES

0158. Calcium tetrahydroborate

[17068-96-0]



B₂CaH₈

Köster, R. *et al.*, *Inorg. Synth.*, 1977, **17**, 18

Like other complex hydrides, it reacts vigorously with protic materials and ignition may occur.

Tetrahydrofuran


Köster, R. *et al.*, *Inorg. Synth.*, 1977, **17**, 18

Dissolution of the 90% pure hydride in dry THF is extremely exothermic.

See other COMPLEX HYDRIDES

0159. Bromodiborane

[23834-96-0]



B₂BrH₅


Drake, J. E., *Inorg. Synth.*, 1978, **18**, 146

May ignite violently in air.

See other HALOBORANES

0160. Chlorodiborane

[17927-57-0]



B₂ClH₅

Bailar, 1973, Vol. 1, 778

A gas above -11°C , it ignites in air.

See other HALOBORANES

0161. Diboron tetrachloride

[13701-67-2]



B₂Cl₄

Wartik, T. *et al.*, *Inorg. Synth.*, 1967, **10**, 125

Sudden exposure to air may cause explosion.

Dimethylmercury

Wartik, T. *et al.*, *Inorg. Chem.*, 1971, **10**, 650

The reaction, starting at -63°C under vacuum, exploded violently on 2 occasions after 23 uneventful runs.

See other HALOBORANES, NON-METAL HALIDES

0162. Diboron tetrafluoride
[13965-73-6]



Metal oxides

Holliday, A. K. *et al.*, *J. Chem. Soc.*, 1964, 2732

Mixtures with mercury(II) oxide and manganese dioxide prepared at -80°C ignited at 20° and reacted violently at 15°C , respectively. Copper(II) oxide reacted vigorously at 25°C without ignition.


Oxygen

Trefonas, L. *et al.*, *J. Chem. Phys.*, 1958, **28**, 54

The gas is extremely explosive in presence of oxygen.

See other HALOBORANES, NON-METAL HALIDES

0163. Potassium hypoborate
[63706-85-4]



Mellor, 1946, Vol. 5, 38

As a reductant stronger than the phosphinate, it may be expected to interact more vigorously with oxidants.

See Potassium phosphinate: Air, etc.

See other REDUCANTS

0164. Sodium hypoborate
[16903-32-5]




Mellor, 1946, Vol. 5, 38

As a reductant stronger than sodium phosphinate, it may be expected to interact more vigorously with oxidants.

See Sodium phosphinate: Oxidants


See other REDUCANTS

0165. Iododiborane
[20436-27-5]



Drake, J. E., *Inorg. Synth.*, 1978, **18**, 146
May ignite violently in air.
See other HALOBORANES

†0166. Diborane(6)
[19287-45-7]



(MCA SD-84, 1961); FPA H81, 1978; HCS 1980, 363 (cylinder)

Preparative hazard

1. Mellor, 1946, Vol. 5, 36
2. Schlessinger, H. I. *et al.*, *Chem. Rev.*, 1942, **31**, 8
3. *Hydrides of Boron and Silicon*, Stock, A. E., Ithaca, Cornell Univ. Press, 1933
4. Follet, M., *Chem. & Ind.*, 1986, 123–128
5. Hariguchi, S. *et al.*, *Chem. Abs.*, 1989, **110**, 62931

The endothermic gas ($\Delta H_f^\circ (g) +31.3 \text{ kJ/mol}$, 1.12 kJ/g) usually ignites in air unless dry and free of impurities [1], and ignition delays of 3–5 days, followed by violent explosions, have been experienced [2]. Explosion followed spillage of liquid diborane [3]. Problems in industrial-scale use of diborane are discussed, and the advantages of 2 new solid and non-flammable diborane complexes with 1,2-bis(*tert*-butylthio)ethane and 1,4-bis(benzylthio)butane, and the pyrophoric complex with dimethyl sulfide are outlined [4]. Flammability limits and explosion pressures of diborane in mixtures with air, nitrogen or helium, and with hydrogen and air, have been studied. Limits for diborane–air mixtures were 0.84 to 93.3 vol%, and presence of nitrogen or helium was not effective in suppressing explosion, the limiting oxygen value being 1.3 vol%. Explosion pressure in air was 11 bar max. at 10% diborane, with a rate of rise of 1200 bar/s. Presence of hydrogen increases the explosive range. Extraordinary precautions are required for industrial use of diborane [5].

See GAS HANDLING (reference 2)

See Sodium tetrahydroborate: Acids

See Benzene, or Moisture; Oxygen, both below

Ammonia

See Diammineboronium tetrahydroborate

Benzene, or Moisture

Simons, H. P. *et al.*, *Ind. Eng. Chem.*, 1958, **50**, 1665, 1659

Effects of presence of moisture or benzene vapour in air on the spontaneously explosive reaction have been studied.

Chlorine

See Chlorine: Non-metal hydrides

Dimethyl sulfoxide

See Dimethyl sulfoxide: Boron compounds

Halocarbons

Haz. Chem. Data, 1975, 114

Diborane reacts violently with halocarbon liquids used as vaporising fire-extinguishants.

Metals

Haz. Chem. Data, 1975, 114

Interaction with aluminium or lithium gives complex hydrides which may ignite in air.

Nitrogen trifluoride

See Nitrogen trifluoride: Diborane

Octanal oxime, Sodium hydroxide

Augustine, 1968, 78

Addition of sodium hydroxide solution during work-up of a reaction mixture of oxime and diborane in THF is very exothermic, a mild explosion being noted on one occasion.

Oxygen

Whatley, A. I. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 1997–1999

Mixtures at 105–165°C exploded spontaneously after an induction period dependent on temperature and composition.

See title reference 2

Tetravinyllead

Houben-Weyl, 1975, Vol. 13.3, 253


Interaction is explosively violent at ambient temperature.

See other BORANES, ENDOTHERMIC COMPOUNDS

0167. Diammineboronium tetrahydroborate

[23777-63-1]

B₂H₁₂N₂



1. Sidgwick, 1950, 354

2. Bailar, 1973, Vol. 1, 924

The diammine complex of diborane (formulated as above), though less reactive than diborane, ignites on heating in air [1,2].

See related BORANES *See other* COMPLEX HYDRIDES

0168. Magnesium boride
[12007-25-9]



Acids

Mellor, 1946, Vol. 5, 25

The crude product containing some silicide evolves, in contact with hydrochloric or sulfuric acids, a mixture of borane and silane which may ignite.

See other METAL NON-METALLIDES

0169. Diboron oxide
[12505-77-0]



1. Merck, 1976, 1365

2. Halliday, A. K. *et al.*, *Chem. Rev.*, 1962, **62**, 316

Probably polymeric at ambient temperature [1], at 400°C, traces of water react to cause a violent eruption and incandescence [2].

See other NON-METAL OXIDES

0170. Boron trioxide
[1303-86-2]



Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

See other NON-METAL OXIDES

0171. Thallium(I) peroxodiborate
[]



Bailar, 1973, Vol. 1, 1154

It liberates oxygen at 18°C and explodes on further warming.

See other PEROXOACID SALTS

0172. Boron trisulfide
[12007-33-9]



Chlorine

See Chlorine: Sulfides


Water

Partington, 1967, 415

Hydrolysis of the sulfide is violent.

See other NON-METAL SULFIDES

0173. 1,3,5-Trichloro-2,4,6-trifluoroborazine
[56943-26-1]





Water

Elter, G. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1975, **14**, 709

Hydrolysis of this *N*-chloro-*B*-fluoro compound is explosively violent.

See other N-HALOGEN COMPOUNDS, NON-METAL HALIDES

0174. B-1,3,5-Trichloroborazine
[26445-82-9]



Water

Niendenzu, K. *et al.*, *Inorg. Synth.*, 1967, **10**, 141

Hydrolysis of the *B*-chloro compound is violent.

See related NON-METAL HALIDES

0175. Triboron pentafluoride
[15538-67-7]



Air, or Water

Timms, P. L., *J. Amer. Chem. Soc.*, 1967, **89**, 1631

It reacts explosively with air or water.

Tetrafluoroethylene


Timms, P. L., *J. Amer. Chem. Soc.*, 1967, **89**, 1631

The pentafluoride catalyses polymerisation of tetrafluoroethylene smoothly below -100°C , but explosively above that temperature.

See other NON-METAL HALIDES

See other POLYMERISATION INCIDENTS

0176. Borazine
[6569-51-3]



Niedenzu, K. *et al.*, *Inorg. Synth.*, 1967, **10**, 144
Samples sealed into ampoules exploded when stored in daylight, but not in the dark.
See other IRRADIATION DECOMPOSITION INCIDENTS
See related NON-METAL HYDRIDES

0177. Sodium octahydrotriborate
[12007-46-4]



Solvents


Dewkett, W. J. *et al.*, *Inorg. Synth.*, 1974, **15**, 116
Air should not be drawn through solutions of the compound in ether, or through its solid complex with dioxane, because such materials have occasionally ignited in air.
See other COMPLEX HYDRIDES

0178. Uranium(III) tetrahydroborate
[]




Semenenko, K. N. *et al.*, *Russ. Chem. Rev.*, 1973, 4
It ignites in air and explodes on heating, unlike the U(VI) analogue.
See other COMPLEX HYDRIDES

0179. Tetraboron tetrachloride
[17156-85-3]




Urry, G. *et al.*, *Inorg. Chem.*, 1963, **2**, 398
Ignites in air.
See other HALOBORANES

0180. Tetra(boron nitride) fluorosulfate
[68436-99-7]



Bartlett, N. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1978, 201
Apparently thermodynamically unstable, because when the peroxodisulfonyl difluoride–boron nitride reaction mixture was heated to 40°C, detonations occurred.

0181. Tetraborane(10)
[18283-93-7]



Oxidants

1. Mellor, 1946, Vol. 5, 36
2. Bailar, 1973, Vol. 1, 790

It ignites in air or oxygen, and explodes with conc. nitric acid [1]. The pure compound is stated not to ignite in air [2].

See Nitric acid: Non-metal hydrides

See other BORANES

0182. Hafnium(IV) tetrahydroborate
[25869-93-6]




Gaylord, 1956, 58

Violent ignition on exposure to air.

See other COMPLEX HYDRIDES

0183. Uranium(IV) tetrahydroborate etherates
[65579-07-9]



Water

Rietz, R. R. *et al.*, *Inorg. Chem.*, 1978, **17**, 654, 658

In contact with water, the adduct with dimethyl ether ignites and that of diethyl ether often explodes, as does invariably the bis-THF adduct.

See other COMPLEX HYDRIDES

0184. Zirconium(IV) tetrahydroborate

[23840-95-1]




Gaylord, 1956, 58

Violent ignition on exposure to air.

See other COMPLEX HYDRIDES

0185. Sodium tetraborate

[1330-43-4]




Zirconium

See Zirconium: Oxygen-containing compounds

See related METAL OXONON-METALLATES

0186. 1-Bromopentaborane(9)

[23753-67-5]



Air, or Hexamethylenetetramine

Rommel, R. J. *et al.*, *Inorg. Synth.*, 1979, **19**, 248

It ignites in air and reacts explosively with hexamine above 90°C.


Dimethyl ether, Potassium hydride

See Potassium hydride: Bromopentaborane(9), Dimethyl ether

See other HALOBORANES

0187. 1,2-Dibromopentaborane(9)

[]




Rommel, R. J. *et al.*, *Inorg. Synth.*, 1979, **19**, 248

It may detonate above ambient temperature.
See other HALOBORANES

†0188. Pentaborane(9)
[19624-22-7]

B₅H₉



(MCA SD-84, 1961); NSC 508, 1979 (Boron hydrides)
The endothermic liquid ignites spontaneously in air if impure.

Ammonia

See Diammineboronium heptahydrotetraborate

Pentacarbonyliron, Pyrex glass

Shore, S. G. *et al.*, *Inorg. Chem.*, 1979, **18**, 670

In the preparation of 2-(tricarbonyliron)hexaborane(10) by co-pyrolysis of the reactants in a hot-cold Pyrex tube reactor, the latter was severely etched and weakened, sometimes splintering. At 230° a maximum cumulative service life of 4 months was observed, and at 260°C the reactor was replaced at the first signs of etching, usually after 6 runs.

See other GLASS INCIDENTS

Oxygen

Pentaborane, *Tech. Bull. LF202*, Alton (Il.), Olin Corp. Energy Div., 1960

Reaction of pentaborane with oxygen is often violently explosive.

Reactive solvents

1. Cloyd, 1965, 35

2. Miller, V. R. *et al.*, *Inorg. Synth.*, 1964, **15**, 118–122

Pentaborane is stable in inert hydrocarbon solvents but forms shock-sensitive solutions in most other solvents containing carbonyl, ether or ester functional groups and/or halogen substituents [1]. The later reference gives detailed directions for preparation and handling of this exceptionally reactive compound, including a list of 26 solvents and compounds rated as potentially dangerous in the presence of pentaborane. When large quantities are stored at low temperature in glass, a phase change involving expansion of the solid borane may rupture the container [2].

See other GLASS INCIDENTS

See Dimethyl sulfoxide: Boron compounds


Tris(difluoroamino)fluoromethane

See Tris(difluoroamino)fluoromethane: Pentaborane(9)

See other BORANES, ENDOTHERMIC COMPOUNDS

0189. Pentaborane(11)
[18433-84-6]


B₅H₁₁



Kit and Evered, 1960, 69
Ignites in air.
See other BORANES

0190. Diammineboronium heptahydrotetraborate (Pentaborane(9) diammoniate)
[28965-70-0]

B₅H₁₅N₂



Kodama, G., *J. Amer. Chem. Soc.*, 1970, **92**, 3482
The diammoniate of pentaborane(9) decomposes spectacularly on standing at ambient temperature.
See related BORANES

0191. Hexaborane(10)
[23777-80-2]

B₆H₁₀

(Complex Structure)

Rommel, R. J. *et al.*, *Inorg. Synth.*, 1979, **19**, 248
It ignites in air.
See other BORANES

0192. Hexaborane(12)
[28375-94-2]

B₆H₁₂

(Complex Structure)

Mellor, 1946, Vol. 5, 36
It is unstable and ignites in air.
See other BORANES

0193. Lanthanum hexaboride
[12008-21-8]

B₆La

LaB₆

See entry REFRACTORY POWDERS

0194. Caesium lithium tridecahydrononaborate
[[12430-27-2] (ion)] **B₉CsH₁₃Li**
(Complex structure)

Siedle, A. R. *et al.*, *Inorg. Chem.*, 1974, **13**, 2737
It ignites in air.
See other COMPLEX HYDRIDES

0195. Disodium tridecahydrononaborate(2-)
[119391-53-6] **B₉H₁₃Na₂**
(Complex structure)

See Sodium tetradecahydrononaborate
See other COMPLEX HYDRIDES

0196. Sodium tetradecahydrononaborate
[70865-40-6] **B₉H₁₄Na**
(Complex structure)

Getman, T. D. *et al.*, *Inorg. Chem.*, 1989, **28**(8), 1507
Explosions occurred during the preparation and handling of this compound and the disodium salt.
See other COMPLEX HYDRIDES

0197. 1,10-Bis(diazonio)decaboran(8)ate
[] **B₁₀H₈N₄**
(Complex structure)

The precursor is explosive.
See Ammonium decahydrodecaborate(2-): Nitrous acid
See related DIAZONIUM SALTS

0198. Decaborane(14)
[17702-41-9] **B₁₀H₁₄**
(Complex Structure)

Ether, or Halocarbons, or Oxygen

1. *MCA SD-84*, 1961
2. Hawthorne, M. F., *Inorg. Synth.*, 1967, **10**, 93–94
3. Shore, S. G., *Chem. Abs.*, 1986, **105**, 202122

It forms impact-sensitive mixtures with ethers (dioxane, etc.) and halocarbons (carbon tetrachloride) and ignites in oxygen at 100°C [1,2]. An improved and safer synthesis of decaborane from pentaborane is given [3].

See Dimethyl sulfoxide: Boron compounds
See other BORANES

0199. Ammonium decahydrodecaborate(2-)

[12008-61-6]

B₁₀H₁₈N₂

(NH₄)₂B₁₀H₁₀

Nitrous acid

Knoth, W. H., *J. Amer. Chem. Soc.*, 1964, **86**, 115

Interaction of the B₁₀H₁₀(2-) anion with excess nitrous acid gives an inner diazonium salt (of unknown structure, possibly containing a nitronium ion) which is highly explosive in the dry state. It is readily reduced wet to the non-explosive 1,10-bis(diazonio)decaboran(8)ate inner salt.

See related BORANES

0200. Barium

[7440-39-3]

Ba

Ba

HCS 1980, 171

Air

See Oxidising gases, below

Halocarbons

1. *Serious Accid. Ser.*, 1952, 23 and Suppl., Washington, USAEC
2. Anon., *Ind. Res.*, 1968, (9), 15
3. *Pot. Incid. Rep. 39*, ASES, 1968
4. Stull, 1977, 25

A violent reaction occurred when cleaning lump metal under carbon tetrachloride [1]. Finely divided barium, slurried with trichlorotrifluoroethane, exploded during transfer owing to frictional initiation [2]. Granular barium in contact with fluorotrichloromethane, carbon tetrachloride, 1,1,2-trichlorotrifluoroethane, tetrachloroethylene or trichloroethylene is susceptible to detonation [3]. Thermodynamic calculations indicated a heat of decomposition of 2.60 kJ/g of mixture and a likely adiabatic temperature approaching 3000°C, accompanied by a 30-fold increase in pressure [4].

See other FRICTIONAL INITIATION INCIDENTS, METAL-HALOCARBON INCIDENTS

Interhalogens

See Bromine pentafluoride: Acids, etc.

Iodine heptafluoride: Metals

Oxidising gases

Kirk-Othmer, 1964, Vol. 3, 78

The finely divided metal may ignite or explode in air or other oxidising gases.

Water

Sidgwick, 1950, 844

Interaction is more violent than with calcium or strontium, but less so than with sodium.

See other METALS, PYROPHORIC METALS

0201. Barium tetrafluorobromate
[35967-90-9]



See entry METAL POLYHALOHALOGENATES

0202. Barium bromate
[13967-90-3]



Hackspill, L. *et al.*, *Compt. rend.*, 1930, **191**, 663

Thermal decomposition with evolution of oxygen is almost explosive at 300°C.

Disulfur dibromide

See Disulfur dibromide: Oxidants

Metals

MRH Aluminium 6.44/22, magnesium 6.40/26

See METAL HALOGENATES: metals

Other reactants

Yoshida, 1980, 186

MRH values calculated for 28 combinations with oxidisable materials are given.

Sulfur

MRH 2.26/12

Taradoire, F., *Bull. Soc. Chim. Fr.*, 1945, **12**, 94, 447

Mixtures are unstable and may ignite 2–11 days after preparation if kept at ambient temperature, or immediately at 91–93°C. Presence of moisture (as water of crystallisation) accelerates ignition.

See other METAL HALOGENATES

0203. Barium perchlorylamide
[28815-10-3]



See entry PERCHLORYLAMIDE SALTS *See other* N-METAL DERIVATIVES

0204. Barium chlorite
[14674-74-9]



Solymosi, F. *et al.*, *Chem. Abs.*, 1968, **68**, 51465

When heated rapidly, barium chlorite decomposes explosively at 190°C, and the lead salt at 112°C.

Dimethyl sulfate

Pascal, 1960, Vol. 16.1, 264

The sulfate ignites in contact with the unheated chlorite, presumably owing to formation of very unstable methyl chlorite.

See other CHLORITE SALTS, OXIDANTS

0205. Barium chlorate

[13477-00-4]



HCS 1980, 175

Other reactants

Yoshida, 1980, 72

MRH values calculated for 19 combinations with oxidisable materials are given.

See other METAL CHLORATES

0206. Barium perchlorate

[13465-95-7]



HCS 1980, 180

Alcohols

Kirk-Othmer, 1964, Vol. 5, 75

Distillation of mixtures with C₁–C₃ alcohols gives the highly explosive alkyl perchlorates. Extreme shock-sensitivity is still shown by n-octyl perchlorate.

See ALKYL PERCHLORATES

Calcium 2,4-pentanedionate

Hamid, I. *et al.*, *Thermochim. Acta*, 1986, **101**, 189

If the ratio of the calcium chelate salt to the trihydrated barium salt in a mixture is above 2:1, the thermal decomposition during DTA/TG analysis may be explosive.

See other METAL PERCHLORATES

0207. Barium hydride

[13477-09-3]



Air, or Oxygen

1. Gibson, 1969, 74


2. Mellor, 1941, Vol. 3, 650

The finely divided hydride ignites in air [1], and coarser material when heated in oxygen [2].

Metal halogenates

See METAL HALOGENATES: Metals, etc. *See other* METAL HYDRIDES

0208. Barium hydroxide
[17194-00-2]




HCS 1980, 178

Chlorinated rubber

See CHLORINATED RUBBER: Metal oxides or hydroxides

See other INORGANIC BASES

0209. Barium amidosulfate
[13770-86-0]



Metal nitrates or nitrites

See entry METAL AMIDOSULFATES

0210. Barium phosphinate ('Barium hypophosphite')
[14871-79-5]



Potassium chlorate

See Potassium chlorate: Reducants

See other METAL PHOSPHINATES, REDUCANTS

0211. Barium iodate
[10567-69-8]



Other reactants

Yoshida, 1980, 383

MRH values calculated for 27 combinations with oxidisable materials are given.

See other METAL HALOGENATES

0212. Barium nitrate
[10022-31-8]



HCS 1980, 179

Aluminium, Potassium nitrate, Potassium perchlorate, Water

See Aluminium: Metal nitrates, etc.

Aluminium–magnesium alloy

MRH 4.93/22 or 5.31/27

Tomlinson, W. R. *et al.*, *J. Chem. Educ.*, 1950, **27**, 606

An intimate mixture of the finely divided components, once widely used as a photo-flash composition, is readily ignitable and extremely sensitive to friction or impact.

Other reactants

Yoshida, 1980, 202

MRH values calculated for 15 combinations with oxidisable materials are given.

See other METAL NITRATES

0213. Barium nitridoosmate

[25395-83-9]



Mellor, 1942, Vol. 15, 728

Explodes at 150°C, like the ammonium salt.

See other N-METAL DERIVATIVES

0214. Barium azide

[18810-58-7]



HCS 1980, 173

1. Fagan, C. P. *J. and Proc. R. Inst. Chem.*, 1947, 126
2. Ficherouille, H. *et al.*, *Mém. Poudres*, 1956, **33**, 7
3. Gyunter, P. L. *et al.*, *Chem. Abs.*, 1943, **37**, 1270₉
4. Verneker, V. R-P. *et al.*, *J. Phys. Chem.*, 1968, **72**, 778–783
5. Stull, 1977, 10

The material is impact-sensitive when dry and is supplied and stored damp with ethanol. It is used as a saturated solution and it is important to prevent total evaporation, or the slow growth of large crystals which may become dried and shock-sensitive. Lead drains must not be used, to avoid formation of the detonator, lead azide. Exposure to acid conditions may generate explosive hydrazoic acid [1]. It has been stated that barium azide is relatively insensitive to impact but highly sensitive to friction [2]. Strontium, and particularly calcium azides show much more marked explosive properties than barium azide. The explosive properties appear to be closely associated with the method of formation of the azide [3]. Factors which affect the sensitivity of the azide include surface area, solvent used and ageing. Presence of barium metal, sodium or iron ions as impurities increases the sensitivity [4]. Though not an endothermic compound ($\Delta H_f^\circ -22.17$ kJ/mol, 0.1 kJ/g), it may thermally decompose to barium nitride, rather than to the elements, when a considerable exotherm is produced (98.74 kJ/mol, 0.45 kJ/g of azide) [5].

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

0215. Barium oxide

[1304-28-5]



Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Barium oxide

Hydroxylamine

See Hydroxylamine: Oxidants

Sulfur trioxide

See Sulfur trioxide: Metal oxides

Triuranium octaoxide

See Triuranium octaoxide: Barium oxide

Water

Bailar, 1973, Vol. 1, 638

Interaction is so vigorous and exothermic as to be a potential ignition source and fire hazard.

See other METAL OXIDES

0216. Barium peroxide

[1304-29-6]

BaO₂

BaO₂

HSC 1980, 181

Acetic anhydride

Rüst, 1948, 337

The peroxide was substituted for (unavailable) potassium permanganate in a process for purifying the crude anhydride in an open vessel. After several operations, when only minor explosions occurred, a violent explosion and fire occurred. Acetyl peroxide would be produced.

Calcium–silicon alloy

Smolin, A. O. *et al.*, *Chem. Abs.*, 1976, **85**, 194906

Combustion of silico-calcium in a mixture with barium peroxide (title only translated).

Delay compositions

Stupp, J., *Chem. Abs.*, 1975, **82**, 113746

The spontaneous ignition of the peroxide in (unspecified) tracer-ignition delay compositions is described.

Hydrogen sulfide

See Hydrogen sulfide: Metal oxides

Hydroxylamine

See Hydroxylamine: Oxidants

Metals

Pascal, 1958, Vol. 4, 775

Powdered aluminium or magnesium ignite in intimate contact with the peroxide.

See also FLASH POWDER

Non-metal oxides

Pascal, 1958, Vol. 4, 773–774

The heated peroxide attains incandescence in a rapid stream of carbon dioxide or sulfur dioxide.

Organic materials, Water

Koffolt, J. H., private comm., 1966

Contact of barium peroxide and water will readily produce a temperature and a local oxygen concentration high enough to ignite many organic compounds.

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

Propane

Hoffmann, A. B. *et al.*, *J. Chem. Educ.*, 1974, **51**, 419 (footnote 7)

Heating barium peroxide under gaseous propane at ambient pressure caused a violent exothermic reaction which deformed the glass container.

See other GLASS INCIDENTS

Selenium

Johnson, L. B., *Ind. Eng. Chem.*, 1960, **52**, 241–244

Powdered mixtures ignite at 265°C.

Wood

1. Dupré, A., *J. Soc. Chem. Ind.*, 1897, **16**, 492

2. Anon., *Jahresber.*, 1987, 66

Friction of the peroxide between wooden surfaces ignited the latter [1]. A granulated and dried priming composition, largely consisting of barium peroxide, was being sieved by hand to break up agglomerated lumps. Towards the end of the operation, ignition occurred. This was caused by rubbing contact of the priming mixture with the sloping plywood frame of the sieve as the composition was pressed through [2].

See other FRICTIONAL INITIATION INCIDENTS

See other METAL PEROXIDES, OXIDANTS

0217. Barium sulfate

[7727-43-7]

BaSO₄

BaO₄S

Aluminium

See Aluminium: Metal oxides, or Oxosalts

Phosphorus

See Phosphorus: Metal sulfates

See other METAL OXONON-METALLATES

0218. Barium sulfide
[21109-95-5]

BaS

BaS

HCS 1980, 182

Dichlorine oxide

See Dichlorine oxide: Oxidisable materials

Oxidants

Mellor, 1941, Vol. 3, 745

Barium sulfide explodes weakly on heating with lead dioxide or potassium chlorate, and strongly with potassium nitrate. Calcium and strontium sulfides are similar.

Phosphorus(V) oxide

Pascal, 1958, Vol. 4, 832

Interaction is violent, attaining incandescence.

See other METAL SULFIDES

0219. Barium nitride
[12047-79-9]

Ba₃N₂

Ba₃N₂

Air, or Water

Sorbe, 1968, 34

It reacts violently with air or water.

See Barium azide (reference 5)

See other N-METAL DERIVATIVES

0220. Beryllium
[7440-41-7]

Be

Be

HCS 1980, 207; *RSC Lab. Hazards Data Sheet No. 67*, 1988 (Be and compounds)

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, Nitrogen

Halocarbons

Pot. Incid. Rep. 39, ASESB, 1968

Mixtures of powdered beryllium with carbon tetrachloride or trichloroethylene will flash on heavy impact.

See other METAL-HALOCARBON INCIDENTS

Halogens

Pascal, 1958, Vol. 4, 22

Warm beryllium incandesces in fluorine or chlorine.

Phosphorus
See Phosphorus: Metals
See other METALS

0221. Beryllium chloride
[7787-47-5]



Sulfur nitrides
See Disulfur dinitride: Metal chlorides

Tetrasulfur tetranitride: Metal chlorides
See other METAL HALIDES

0222. Beryllium perchlorate
[13597-95-0]



Laran, R. J., US Pat. 3 157 464, 1964
A powerful oxidant, insensitive to heat or shock and useful in propellant and igniter systems.
See other METAL PERCHLORATES

0223. Beryllium fluoride
[7787-49-7]



Magnesium
See Magnesium: Beryllium fluoride
See other METAL HALIDES

0224. Beryllium hydride
[7787-52-2]



Methanol, or Water

1. Barbaras, G. D., *J. Amer. Chem. Soc.*, 1951, **73**, 48
2. Brendel, G. J. *et al.*, *Inorg. Chem.*, 1978, **17**, 3589

Reaction of the ether-containing hydride with methanol or water is violent, even at -196°C [1]. A crystalline modification produced by pressure-compaction of the hydride reacts slowly with moist air or water [2].

See other METAL HYDRIDES

0225. Beryllium oxide
[1304-56-9]

BeO

BeO

HCS 1980, 208

Magnesium

See Magnesium: Metal oxides

See other METAL OXIDES

0226. Bismuth
[7440-69-9]

Bi

Bi

Aluminium

See Aluminium: Bismuth

Oxidants

See Ammonium nitrate: Metals

Bromine pentafluoride: Acids, etc.

Chloric acid: Metals, etc.

Iodine pentafluoride: Metals

Nitric acid: Metals

Nitrosyl fluoride: Metals

Perchloric acid: Bismuth

See other METALS

0227. Bismuth pentafluoride
[7787-62-4]

BiF₅

BiF₅

Water

von Wartenberg, H., *Z. Anorg. Chem.*, 1940, **224**, 344

It reacts vigorously with water, sometimes igniting.

See other METAL HALIDES

0228. Bismuthic acid (Bismuth oxide hydroxide)
[22750-47-6]

HOBiO₂ (? Bi₂O₅·H₂O)

BiHO₃

Hydrofluoric acid

Mellor, 1939, Vol. 9, 657

Interaction of the solid acid with 40% hydrofluoric acid is violent, ozonised oxygen being evolved.

See related INORGANIC ACIDS *See other* OXIDANTS

0229. Bismuth amide oxide

[]



Watt, G. W. *et al.*, *J. Amer. Chem. Soc.*, 1939, **61**, 1693

The solid, prepared in liquid ammonia, explodes when free of ammonia and exposed to air.

See other N-METAL DERIVATIVES

0230. Bismuth nitride

[12232-97-2]



Alone, or Water

1. Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1471

2. Franklin, E. C., *J. Amer. Chem. Soc.*, 1905, **27**, 847

Very unstable, exploded on shaking [1] or heating, or in contact with water or dilute acids [2].

See other N-METAL DERIVATIVES, METAL NON-METALLIDES

0231. Plutonium bismuthide

[12010-53-6]



Williamson, G. K., *Chem. & Ind.*, 1960, 1384

Extremely pyrophoric.

See other ALLOYS, PYROPHORIC MATERIALS

0232. Dibismuth dichromium nonaoxide ('Bismuth chromate')

[37235-82-8]



Hydrogen sulfide

Pascal, 1960, Vol. 13.1, 1025

The gas may ignite on contact with the 'chromate'.

See other METAL OXIDES, METAL OXOMETALLATES

0233. Bismuth trioxide

[1304-76-3]



Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Potassium

See Potassium: Metal oxides

Sodium

See Sodium: Metal oxides

See other METAL OXIDES

0234. Bismuth trisulfide

[1345-07-9]



Preparative hazard

Glatz, A. C. *et al.*, *J. Electrochem. Soc.*, 1963, **110**, 1231

Possible causes of explosions in direct synthesis are discussed.

See Sulfur: Metals

See other METAL SULFIDES

0235. Bromine perchlorate

[32707-10-1]



Schack, C. J. *et al.*, *Inorg. Chem.*, 1971, **10**, 1078

It is shock-sensitive.

Perfluorobutadiene

Schack, C. J. *et al.*, *Inorg. Chem.*, 1975, **14**, 151 (footnote 8)

During adduct formation, the perchlorate must be present in excess to prevent formation of a mono-adduct, which may well be explosive.

See Chlorine perchlorate: Chlorotrifluoroethylene

See other GLASS INCIDENTS *See other* HALOGEN OXIDES

0236. Caesium hexafluorobromate

[26222-92-4]



See entry METAL POLYHALOHALOGENATES

0237. Caesium bromoxenate

[]



Water

Jaselskis, B. *et al.*, *J. Amer. Chem. Soc.*, 1969, **91**, 1875

Aqueous solutions are extremely unstable and caution is required if isolation of the compound is contemplated.

See other XENON COMPOUNDS

0238. Bromine fluoride
[13863-59-7]

BrF

BrF

Sidgwick, 1950, 1149

Chemically it behaves like the other bromine fluorides, but is more reactive.

Hydrogen

Pascal, 1960, Vol. 16.1, 412

Hydrogen ignites in the fluoride at ambient temperature.

See other INTERHALOGENS

0239. Bromyl fluoride
[22585-64-4]

O₂BrF

BrFO₂

Water

Bailar, 1973, Vol. 2, 1388

Hydrolysis may proceed explosively.

See other HALOGEN OXIDES

0240. Perbromyl fluoride
[25251-03-0]

O₃BrF

BrFO₃

Fluoropolymers

Johnson, K. G. *et al.*, *Inorg. Chem.*, 1972, **11**, 800

It is considerably more reactive than perchloryl fluoride, and attacks glass and the usually inert polytetrafluoroethylene and polychlorotrifluoroethylene.

See other CORROSION INCIDENTS, GLASS INCIDENTS, HALOGEN OXIDES

0241. Bromine trifluoride
[7787-71-5]

BrF₃

BrF₃

1. Davis, R. A. *et al.*, *J. Org. Chem.*, 1962, **32**, 3478

2. Musgrave, W. K. R., *Advan. Fluorine Chem.*, 1964, **1**, 12

The hazards and precautions in use of this very reactive fluorinating agent are outlined [1]. Contact with rubber, plastics or other organic materials may be explosively violent and reaction with moisture is very vigorous [2].

Ammonium halides

Sharpe, A. G. *et al.*, *J. Chem. Soc.*, 1948, 2137

Explosive reaction.

Antimony(III) chloride oxide

Mellor, 1956, Vol. 2, Suppl. 1, 166

Interaction is violent, even more so than with antimony trioxide.

Carbon monoxide

Mellor, 1956, Vol. 2, Suppl. 1, 166

At temperatures rather above 30°C, explosions occurred.

Carbon tetrachloride

Dixon, K. R. *et al.*, *Inorg. Synth.*, 1970, **12**, 233

Excess bromine trifluoride may be destroyed conveniently in a hood by slow addition to a large volume of the solvent, interaction being vigorous but not dangerous.

See Solvents, below

Halogens, or Metals, or Non-metals, or Organic materials

1. Mellor, 1941, Vol. 2, 113; 1956, Vol. 2, Suppl. 1, 164–167

2. 'Chlorine Trifluoride Tech. Bull.', Morristown, Baker & Adamson, 1970

Incandescence is caused by contact with bromine, iodine, arsenic, antimony (even at –10°C); powdered molybdenum, niobium, tantalum, titanium, vanadium; boron, carbon, phosphorus or sulfur [1]. Carbon tetraiodide, chloromethane, benzene or ether ignite or explode on contact, as do organic materials generally. Silicon also ignites [2].

See Uranium, below

2-Pentanone

Stevens, T. E., *J. Org. Chem.*, 1961, **26**, 1629 (footnote 11)

During evaporation of solvent hydrogen fluoride, an exothermic reaction between residual ketone and bromine trifluoride set in and accelerated to explosion.

Potassium hexachloroplatinate

Dixon, K. R. *et al.*, *Inorg. Synth.*, 1970, **12**, 233–237

Interaction of the reagents in bromine as diluent to produce the trichlorotrifluoro- and then hexafluoro-platinates is so vigorous that increase in scale above 1 g of salt is not recommended.

Pyridine

Kirk-Othmer, 1966, Vol. 9, 592

The solid produced by action of bromine trifluoride on pyridine in carbon tetrachloride ignites when dry. 2-Fluoropyridine reacts similarly.

Silicone grease

Sharpe, A. G. *et al.*, *J. Chem. Soc.*, 1948, 2136

As it reacts explosively in bulk, the amount of silicone grease used on joints must be minimal.

Solvents

1. Sharpe, A. G. *et al.*, *J. Chem. Soc.*, 1948, 2135

2. Simons, J. H., *Inorg. Synth.*, 1950, **3**, 185

Bromine trifluoride explodes on contact with acetone or ether [1], and the frozen solid at -80°C reacts violently with toluene at that temperature [2].

See Halogens, etc., above

Tin(II) chloride

Mellor, 1956, Vol. 2, Suppl. 1, 165

Contact causes ignition.

Uranium, Uranium hexafluoride

Johnson, K. *et al.*, *6th Nucl. Eng. Sci. Conf.*, New York, 1960. Reprint Paper No. 23

Uranium may ignite or explode during dissolution in bromine trifluoride, particularly when high concentrations of the hexafluoride are present. Causative factors are identified.

See Halogens, etc. above

Water

1. Mellor, 1941, Vol. 2, 113

2. *491M*, 1975, 73


Interaction is violent, oxygen being evolved [1], and even at -50°C , reaction with 6N hydrochloric acid is explosive [2].

See other INTERHALOGENS

0242. Tetrafluoroammonium perbromate

[25483-10-7]

BrF_4NO_4



Christe, K. O. *et al.*, *Inorg. Chem.*, 1980, **19**, 1495

Solutions in hydrogen fluoride at -78°C exploded when isolation of the salt was attempted.

See other N-HALOGEN COMPOUNDS, OXOSALTS OF NITROGENOUS BASES

0243. Bromine pentafluoride

[7789-30-2]

BrF_5



Acetonitrile

1. Meinert, H. *et al.*, *Z. Chem.*, 1969, **9**, 190

2. Stein, L., *Chem. Eng. News*, 1984, **62**(28), 4

Although solutions of bromine pentafluoride in acetonitrile were reported stable at ambient temperature [1], it has been found that a 9% solution in the anhydrous solvent prepared at -196°C decomposed violently, bursting the container, about 1 h after attaining ambient temperature [2].

See Hydrogen-containing materials, below

Acids, or Halogens, or Metal halides, or Metals, or Non-metals, or Oxides

1. Mellor, 1956, Vol. 2, Suppl. 1, 172

2. Sidgwick, 1950, 1158

Contact with the following at ambient or slightly elevated temperatures is violent, ignition often occurring: strong nitric acid or sulfuric acids; chlorine (explodes on heating), iodine; ammonium chloride, potassium iodide; antimony, arsenic, boron powder, selenium, tellurium; aluminium powder, barium, bismuth, cobalt powder, chromium, iridium powder, iron powder, lithium powder, manganese, molybdenum, nickel powder, rhodium powder, tungsten, zinc; charcoal, red phosphorus, sulfur, arsenic pentoxide, boron trioxide, calcium oxide, carbon monoxide, chromium trioxide, iodine pentoxide, magnesium oxide, molybdenum trioxide, phosphorus pentoxide, sulfur dioxide or tungsten trioxide [1,2].

Hydrogen-containing materials

1. Mellor, 1956, Vol. 2, Suppl. 1, 172
2. Braker, 1980, 56

Contact with the following materials, containing combined hydrogen, is likely to cause fire or explosion: acetic acid, ammonia, benzene, ethanol, hydrogen, hydrogen sulfide, methane; cork, grease, paper, wax, etc. The carbon content further contributes to the observed reactivity [1]. Chloromethane reacts with explosive violence [2].

See Acetonitrile, above

Perchloryl perchlorate

See Perchloryl perchlorate: Bromine pentafluoride

Water

Sidgwick, 1950, 1158

Contact with water causes a violent reaction or explosion, oxygen being evolved.

See other INTERHALOGENS

0244. Potassium hexafluorobromate

[32312-22-4]



See entry METAL POLYHALOHALOGENATES

0245. Rubidium hexafluorobromate

[32312-23-5]



See entry METAL POLYHALOHALOGENATES

0246. Bromogermane

[13569-43-2]



Preparative hazard

See Bromine: Germane

See related METAL HYDRIDES

0247. Hydrogen bromide
[10035-10-6]

HBr

BrH

HCS 1980, 545 (cylinder gas), 538 (48% solution)

Preparative hazard

See Bromine: Phosphorus

Ammine-1,2-diaminoethanediperoxochromium(IV)

See Ammine-1,2-diaminoethanediperoxochromium(IV): Hydrogen bromide

Fluorine

See Fluorine: Hydrogen halides

Ozone

See Ozone: Hydrogen bromide

See other INORGANIC ACIDS, NON-METAL HALIDES, NON-METAL HYDRIDES

0248. Bromic acid
[7789-31-3]

HOBrO₂

BrHO₃

In contact with oxidisable materials, reactions are similar to those of the metal bromates.

See entry METAL HALOGENATES

See other INORGANIC ACIDS, OXOHALOGEN ACIDS

0249. Bromamine (Bromamide)
[14519-10-9]

BrNH₂

BrH₂N

Jander, J. *et al.*, *Z. Anorg. Chem.*, 1958, **296**, 117

The isolated material decomposes violently at -70°C , while an ethereal solution is stable for a few hours at that temperature.

See other N-HALOGEN COMPOUNDS

†0250. Bromosilane
[13465-73-1]

BrSiH₃

BrH₃Si

Ward, L. G. L., *Inorg. Synth.*, 1968, **11**, 161

Ignites in air (gas above 2°C).

See other HALOSILANES

0251. Ammonium bromide
[12124-97-9]



Bromine trifluoride

See Bromine trifluoride: Ammonium halides

See related METAL HALIDES

0252. Ammonium bromate
[13483-59-5]



1. Sorbe, 1968, 129

2. Shidlovskii, A. A. *et al.*, *Chem. Abs.*, 1968, **69**, 78870

It is a combustible and explosive salt which is very friction-sensitive [1], and may explode spontaneously [2].

Other reactants


Yoshida, 1980, 180

MRH values calculated for 16 combinations with oxidisable materials are given.

See other OXOSALTS OF NITROGENOUS BASES

0253. Poly(dimercuryimmonium bromate)

[]



Sorbe, 1967, 97

Highly explosive.

See entry POLY(DIMERCURYIMMONIUM) COMPOUNDS

0254. Iodine bromide
[7789-33-5]



Metals

1. Mellor, 1961, Vol. 2, Suppl. 2, 452; 1963, Suppl. 3, 1563

2. Pascal, 1963, Vol. 8.3, 308

A mixture with sodium explodes under a hammer blow, while potassium explodes strongly under the molten bromide [1]. Tin reacts violently with the bromide [2].

Phosphorus

Mellor, 1963, Vol. 2, Suppl. 3, 264

Phosphorus reacts violently with the molten bromide.

See other INTERHALOGENS, IODINE COMPOUNDS

0255. Potassium bromate
[7758-01-2]

BrKO₃

KOBrO₂

HCS 1980, 760

Anon., Personal communication, 1999

A factory using potassium bromate regularly in small portions found that it caked in storage and had to be broken up. A largely full, but year old, polythene lined fibre-board drum thereof was broken by stabbing the contents with a, possibly rusty, knife and the lumps needed for use removed by gloved hand. The 1.5 kg so removed was observed to be fizzing and was sprayed with water. Minutes later, the 18 kg remaining in the drum deflagrated brightly, described as “sunrise behind the reactor”, and ignited a sack of monomer nearby. Water extinguished the fire in a few minutes. Although bromates are thermodynamically unstable with respect to bromides, and there may have been slight surface contamination with organics, propagation of a surface reaction is hard to explain. The packaging would be a substantial fuel, but mixing with that is very poor. The exact cause remains mysterious. The business has decided to buy bromate in smaller lots and to subdivide those into individual 1.5 kg charges when received and before caking can occur. The operators have decided to no longer break up caked chemicals by jumping on the polythene liners containing them.

Aluminium, Dinitrotoluene

Yoshinaga, S. *et al.*, *Chem. Abs.*, 1980, **92**, 44080

The mixture reacts violently at 290°C, with enormous gas evolution and is used to fracture concrete. The mechanism was studied.

See other GAS EVOLUTION INCIDENTS

Azoformamide

Vidal, F. D. *et al.*, *Bakers Dig.*, 1979, **53**(3), 16–18

A mixture of the oxidant and blowing agent (2:1) is used as a dough improver. Potential problems of incompatibility during tableting operations were overcome by incorporating hydrated salts into the tableting formulation.

Ceric ammonium nitrate, Malonic acid, Water

Bartmess, J. *et al.*, *Chem. Eng. News*, 1998, **76**(24), 4

Subsequent to a fire in a teaching laboratory, it was discovered that a mixture of equal weights of the three dry solids, itself stable, reacted violently when wetted with up to two parts of water and was capable of igniting paper. All components (which exhibit an oscillating chemical reaction in solution) were necessary for this effect.

Disulfur dibromide

See Disulfur dibromide: Oxidants

Non-metals

MRH Sulfur 2.55/14

1. Taradoire, F., *Bull. Soc. Chim. Fr.*, 1945, **12**, 94, 466
2. Pascal, 1961, Vol. 6, 440

Mixtures with sulfur are unstable, and may ignite some hours after preparation, depending on the state of subdivision and atmospheric humidity [1]. Selenium reacts violently with aqueous solutions of the oxidant [2].

Other reactants

Yoshida, 1980, 181–182

MRH values calculated for 28 combinations with oxidisable materials are given.

See other METAL OXOHALOGENATES, OXIDANTS

0256. Bromine azide

[13973-87-0]

BrN₃

BrN₃

1. Hargittai, M. *et al.*, *Angew. Chem. (Int.)*, 1993, **32**, 759

2. Mellor, 1940, Vol. 8, 336

3. Tornieporth-Oetting, I. C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(5), 511

The solid, liquid and vapour are all very shock-sensitive [1]. The liquid explodes on contact with arsenic; sodium, silver foil; or phosphorus. Explosion is likely to be triggered by pressure fluctuations of around 10 Pa [3]. Concentrated solutions in organic solvents may explode on shaking [2].

See other ENDOTHERMIC COMPOUNDS, HALOGEN AZIDES

0257. Sodium bromate

[7789-38-0]

BrNaO₃

NaOBrO₂

HCS 1970, 831

Fluorine

See Fluorine: Sodium bromate

Grease

1. *MCA Case History No. 874*

2. Stull, 1977, 28–29

A bearing assembly from a sodium bromate crusher had been degreased at 120°C, and while still hot the sleeve was hammered to free it. The assembly exploded violently, probably because of the presence of a hot mixture of sodium bromate and a grease component (possibly a sulfurised derivative). It is known that mixtures of bromates and organic or sulfurous matter are heat- and friction-sensitive [1]. The energy of decomposition of the likely components has been calculated as 1.93 kJ/g, with an explosion temperature above 2000°C [2].

Other reactants

Yoshida, 1980, 184–185

MRH values calculated for 29 combinations with oxidisable materials are given.

See other METAL OXOHALOGENATES

0258. Bromine dioxide
[21255-83-4]




Brauer, 1963, Vol. 1, 306

Unstable unless stored at low temperatures, it may explode if heated rapidly.

See other HALOGEN OXIDES

0259. Bromine trioxide
[32062-14-4]



1. Lewis, B. *et al.*, *Z. Elektrochem.*, 1929, **35**, 648–652

2. Pflugmacher, A. *et al.*, *Z. Anorg. Chem.*, 1955, **279**, 313

The solid produced at -5°C by interaction of bromine and ozone is only stable at -80°C or in presence of ozone, and decomposition may be violently explosive in presence of trace impurities [1]. The structure may be the dimeric bromyl perbromate, analogous to Cl_2O_6 [2].

See other CATALYTIC IMPURITY INCIDENTS, HALOGEN OXIDES

0260. Thallium bromate
[14550-84-6]



Pascal, 1961, Vol. 6, 950

It decomposes explosively around 140°C .

See other HEAVY METAL DERIVATIVES, METAL OXOHALOGENATES

0261. Bromine
[7726-95-6]



(MCA SD-49, 1968); NSC 313, 1979; FPA H61, 1977; HCS 1980, 218
RSC Lab. Hazard Data Sheet No. 24, 1984

Berthelot, J. *et al.*, *J. Chem. Educ.*, 1986, **63**(11), 1011

The stable complex of bromine with tetrabutylammonium bromide is safer and more easily handled than bromine itself.

Acetone

MRH 0.46/16

Levene, P. A., *Org. Synth.*, 1943, Coll. Vol. 2, 89

During bromination of acetone to bromoacetone, presence of a large excess of bromine must be avoided to prevent sudden and violent reaction.

See Carbonyl compounds, below

Acetylene MRH 8.70/100
See Acetylene: Halogens

Acrylonitrile MRH 2.84/100
See Acrylonitrile: Halogens

Alcohols

1. Muir, G. D., *Chem. Brit.*, 1972, **8**, 16
2. Bush, E. L., private comm., 1968
3. Desty, D. H., private comm., 1986

Reaction with methanol may be vigorously exothermic. A mixture of bromine (9 ml) and methanol (15 ml) boiled in 2 m and in a previous incident such a mixture had erupted from a measuring cylinder [1]. The exotherm with industrial methylated spirits (ethanol containing 5% methanol) is much greater, and addition of 10 ml of bromine to 40 ml of IMS rapidly causes violent boiling [2]. A further case of ejection of a methanol solution of bromine from a measuring cylinder was described [3].

See other HALOGENATION INCIDENTS

Aluminium, Dichloromethane

Nolan, 1983, Case History 26

Bromochloromethane was being prepared in a 400 l reactor by addition of liquid bromine to dichloromethane in presence of aluminium powder (which would form some aluminium bromide to catalyse the halogen exchange reaction). The reaction was started and run for 1.5 h, stopped for 8 h, then restarted with addition of bromine at double the usual rate for 2.5 h, though the reaction did not appear to be proceeding. Soon afterwards a thermal runaway occurred, shattering the glass components of the reactor.

See other GLASS INCIDENTS, HALOGENATION INCIDENTS

Ammonia

Mellor, 1967, Vol. 8, Suppl. 2, 417

Interaction at normal or elevated temperatures, followed by cooling to -95°C , gives an explosive red oil.

See Nitrogen tribromide hexaammoniate

Boron

See Boron: Halogens

See Phosphorus: Halogens

3-Bromopropyne

See 3-Bromopropyne, (reference 3)

See Chlorine: 3-Chloropropyne

Carbonyl compounds

MCA SD-49, 1968

Br₂

Organic compounds containing active hydrogen atoms adjacent to a carbonyl group (aldehydes, ketones, carboxylic acids) may react violently in unmoderated contact with bromine.

See Acetone, above

Chlorotrifluoroethylene, Oxygen

Haszeldine, R. N. *et al.*, *J. Chem. Soc.*, 1959, 1085

Addition of bromine to the gas-phase mixture initiated an explosion, but see Oxygen: Halocarbons (reference 3).

Copper(I) hydride

See Copper(I) hydride: Halogens

Diethyl ether

1. Tucker, H., private comm., 1972

2. Anon., *Safety Digest Univ. Safety Assoc.*, 1989, **34**, 14

Shortly after adding bromine to ether the solution erupted violently (or exploded softly). Photocatalytic bromination may have been involved [1]. Spontaneous ignition occurred on addition of ether to impure bromine [2].

See Tetrahydrofuran, below; or Chlorine: Diethyl ether

Diethylzinc

Houben-Weyl, 1973, Vol. 13.2a, 757

Interaction without diluents may produce dangerous explosions. Even with diluents (ether), interaction of dialkylzincs with halogens is initially violent at 0 to -20°C.

Dimethylformamide

Tayim, H. A. *et al.*, *Chem. & Ind.*, 1973, 347

Interaction is extremely exothermic, and under confinement in an autoclave the internal temperature and pressure exceeded 100°C and 135 bar, causing failure of the bursting disc. The product of interaction is dimethylhydroxymethylenimmonium bromide, and the explosive decomposition may have involved formation of *N*-bromodimethylamine, carbon monoxide and hydrogen bromide.

See *N*-HALOGEN COMPOUNDS

Ethanol, Phosphorus

Read, C. W. W., *School Sci. Rev.*, 1940, **21**(83), 967

The vigorous interaction of ethanol, phosphorus and bromine to give bromoethane is considered too dangerous for a school experiment.

Fluorine

See Fluorine: Halogens

Germane

Swiniarski, M. F. *et al.*, *Inorg. Synth.*, 1974, **15**, 157-160

During the preparation of mono- or di-bromogermane, either the scale of operation or the rate of addition of bromine must be closely controlled to prevent explosive reaction occurring.

See Non-metal hydrides, below

Hydrogen

Mellor, 1956, Vol. 2, Suppl. 1, 707

Combination is explosive under appropriate temperature and pressure conditions.

Isobutyrophenone

MCA Guide, 1972, 307

Bromine was added dropwise at 20–31°C to a solution of the ketone in carbon tetrachloride. The completed reaction mixture was cooled in ice, but exploded after 15 m.

See other HALOGENATION INCIDENTS

Metal acetylides and carbides

Several of the mono- and di-alkali metal acetylides and copper acetylides ignite at ambient temperature or on slight warming, with either liquid or vapour. The alkaline earth, iron, uranium and zirconium carbides ignite in the vapour on heating.

See Calcium acetylide: Halogens

Caesium acetylide: Halogens

Dicopper(I) acetylide: Halogens

Iron carbide: Halogens

Lithium acetylide: Halogens

Rubidium acetylide: Halogens

Strontium acetylide: Halogens

Uranium dicarbide: Halogens

Zirconium dicarbide: Halogens

Metal azides

Mellor, 1940, Vol. 8, 336

Nitrogen-diluted bromine vapour passed over silver azide or sodium azide formed bromine azide, and often caused explosions.

Metals

MRH Aluminium 1.96/10

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549
2. Mellor, 1941, Vol. 2, 469; 1963, Vol. 2, Suppl. 2.2, 1563, 2174
3. *MCA SD-49*, 1968
4. Mellor, 1941, Vol. 7, 260
5. Mellor, 1939, Vol. 3, 379
6. Hartgen, C. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 70

Lithium is stable in contact with dry bromine, but heavy impact will initiate explosion, while sodium in contact with bromine needs only moderate impact for initiation [1]. Potassium ignites in bromine vapour and explodes violently in contact with liquid bromine, and rubidium ignites in bromine vapour [2]. Aluminium,

Br₂

mercury or titanium react violently with dry bromine [3]. Warm germanium ignites in bromine vapour [4], and antimony ignites in bromine vapour and reacts explosively with the liquid halogen [5]. During preparation of praseodymium bromide, accidental contact of liquid bromine with small particles of praseodymium led to a violent explosion [6].

See Gallium: Halogens

Nitromethane

Rochat, A. C., private comm., 1990.

A solution of bromine (116 g) in nitromethane (300 ml) was employed in an attempt to brominate a pigment (30 g). On heating the mix in an autoclave, runaway commenced at 70°C, soon shattering the vessel. Nitromethane being an explosive of low oxygen balance, the potential energy certainly, and the sensitivity probably, will be increased by bromine or bromination. Bromonitromethane salts and formaldehyde adducts thereof are intermediates in manufacture of some disinfectants; isolation of the first is considered most unsafe in the industry, the second are thought to need careful handling.

See Nitromethane, Chloronitromethane.

Non-metal hydrides

1. Stock, A. *et al.*, *Ber.*, 1917, **50**, 1739
2. Sujishi, S. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 4631
3. Geisler, T. C. *et al.*, *Inorg. Chem.*, 1972, **11**, 1710
4. Merck, 1976, 955

Interaction of silane and its homologues with bromine at ambient temperature is explosively violent [1] and temperatures of below -30°C are necessary to avoid ignition of the reactants [2]. Ignition of disilane at -95°C and of germane at -112°C emphasises the need for good mixing to dissipate the large exotherm [3]. Phosphine reacts violently with bromine at ambient temperature [4].

See Ethylphosphine: Halogens, or Phosphine: Halogens

Other reactants

Yoshida, 1980, 179

MRH values for 10 combinations with reactive materials are given.

Oxygen, Polymers

Groome, I. J., *Chem. Brit.*, 1983, **19**, 644-665

Bromination of polymers should be effected at temperatures below 120°C to avoid the possibility of explosion likely at higher temperatures if traces of oxygen were present in the highly unsaturated pyrolysis products.

Oxygen difluoride

See Oxygen difluoride: Halogens

Ozone

See Ozone: Bromine

Phosphorus

1. Bandar, L. S. *et al.*, *Zh. Prikl. Khim.*, 1966, **39**, 2304
2. 'Leaflet No. 2', Inst. of Chem., London, 1939

During preparation of hydrogen bromide by addition of bromine to a suspension of red phosphorus in water, the latter must be freshly prepared to avoid the possibility of explosion. This is due to formation of peroxides in the suspension on standing and subsequent thermal decomposition [1]. In the earlier description of such an explosion, action of bromine on boiling tetralin was preferred to generate hydrogen bromide [2], which is now available in cylinders.

Rubber

Pascal, 1960, Vol. 16.1, 371

Bromine reacts violently in contact with natural rubber, but more slowly with some synthetic rubbers.

Sodium hydroxide

MCA Case History No. 1636

A bucket containing 25% sodium hydroxide solution was used to catch and neutralise bromine dripping from a leak. Lack of stirring allowed a layer of unreacted bromine to form below the alkali. Many hours later, a violent eruption occurred when the layers were disturbed during disposal operations. Continuous stirring is essential to prevent stratification of slowly reacting mutually insoluble liquids, especially of such differing densities.

Tetracarbonylnickel

See Tetracarbonylnickel: Bromine

Tetrahydrofuran

Tinley, E. J., private comm., 1983

Rapid addition of bromine to the dried solvent to make a 10% solution caused a vigorous reaction with gas evolution. As this happened in a newly installed brightly illuminated fume cupboard lined with a reflective white finish, photocatalysed bromination of the solvent may have been involved, as has been observed in chlorine–ether systems.

See Diethyl ether, above; Chlorine: Diethyl ether

Tetraselenium tetranitride

See Tetraselenium tetranitride: Alone, or Halogens

Trialkyl boranes

Coates, 1967, Vol. 1, 199

The lower homologues tend to ignite in bromine or chlorine.

Trimethylamine

Bohme, H. *et al.*, *Chem. Ber.*, 1951, **84**, 170–181

The 1:1 adduct (presumably *N*-bromotrimethylammonium bromide) decomposes explosively when heated in a sealed tube.

Trioxxygen difluoride

See 'Trioxxygen difluoride': Various materials

Tungsten, Tungsten trioxide

Tillack, J., *Inorg. Synth.*, 1973, **14**, 116–120

During preparation of tungsten(IV) dibromide oxide, appropriate proportions of reactants are heated in an evacuated sealed glass ampoule to 400–500°C. Initially only one end should be heated to prevent excessive pressure bursting the ampoule.

See other GLASS INCIDENTS

See other HALOGENS, OXIDANTS

0262. Calcium bromide

[7789-41-5]



Potassium

See Potassium: Metal halides

See other METAL HALIDES

0263. Cobalt(II) bromide

[7789-43-7]




Sodium

See Sodium: Metal halides

See other METAL HALIDES

0264. Tetraamminecopper(II) bromate

[]



Rammelsberg, C. F., *Pogg. Ann.*, 1842, **55**, 63

A bright blue solid detonating at 140°C or if struck.

See other AMMINEMETAL OXOSALTS

0265. Copper(I) bromide

[7787-70-4]



tert-Butyl peroxybenzoate, Limonene

See tert-Butyl peroxybenzoate: Copper(I) bromide, etc.

See other METAL HALIDES, REDUCANTS

0266. Iron(II) bromide

[7789-46-0]



Potassium

See Potassium: Metal halides

Sodium

See Sodium: Metal halides

See other METAL HALIDES

0267. Dibromogermane

[13769-36-3]




Preparative hazard

See Bromine: Germane

See related METAL HALIDES, METAL HYDRIDES

0268. N,N-Bis(bromomercurio)hydrazine

[]



Hofmann, K. A. *et al.*, *Ann.*, 1899, **305**, 217

An explosive compound.

See other MERCURY COMPOUNDS, N-METAL DERIVATIVES

0269. Mercury(II) bromide

[7789-47-1]



HCS 1980, 614

Indium


Clark, R. J. *et al.*, *Inorg. Synth.*, 1963, **7**, 19–20

Interaction at 350°C is so vigorous that it is unsafe to increase the scale of this preparation of indium bromide.

See other MERCURY COMPOUNDS, METAL HALIDES

0270. Mercury(II) bromate

[[26522-91-8] (dihydrate)]



Janz, 1976, Table 2, 7

It deflagrates around 155°C.

See other MERCURY COMPOUNDS, METAL OXOHALOGENATES, OXIDANTS

0271. Mercury(I) bromate

[13465-33-3]



Hydrogen sulfide


Pascal, 1960, Vol. 13.1, 1004

Contact of the gas with the solid oxidant causes ignition.

See other MERCURY COMPOUNDS, METAL HALOGENATES

0272. N,N'-Dibromosulfurdiimide


[]

**Br₂N₂S**Seppelt, K. *et al.*, *Angew. Chem. (Int.)*, 1969, **8**, 771

This is more shock sensitive than the iodo-analogue, which explodes both on melting and impact. The chloro-compound is not known.

See other N-HALOGEN COMPOUNDS, N-S COMPOUNDS**0273. Titanium diazide dibromide**

[32006-07-8]

**Br₂N₆Ti**Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 507; *Chem. Ztg.*, 1982, **106**, 187-188

A highly explosive solid, (possibly polymeric).

See other METAL AZIDE HALIDES**0274. Sulfinyl bromide (Thionyl bromide)**

[507-16-4]

**Br₂OS**

1. Hodgson, P. K. G., private comm., 1981

2. Beattie, T. R., *Chem. Eng. News*, 1982, **60**(28), 5

Opening an unused but old commercial ampoule of the ice-cooled bromide led to eruption of the decomposed contents [1]. Similar occurrences on 2 occasions were reported later [2], though on both occasions hexabromonaphthalene was identified in the residue (suggesting contamination of the bromide by naphthalene as an additional source of internal pressure). Sulfinyl bromide is of limited stability, decomposing to sulfur, sulfur dioxide and bromine, and so should be stored under refrigeration and used as soon as possible.

See other GAS EVOLUTION INCIDENTS, NON-METAL HALIDES**0275. Seleninyl bromide**

[7789-51-7]

**Br₂OSe**

Metals

Mellor, 1947, Vol. 10, 912

Sodium and potassium react explosively (the latter more violently), and zinc dust ignites, all in contact with the liquid bromide.

See Sodium: Non-metal halides

Phosphorus

Mellor, 1947, Vol. 10, 912

Red phosphorus ignites, and white phosphorus explodes, in contact with the liquid bromide.

See other NON-METAL HALIDES

0276. Bromine bromate (Dibromine trioxide)

[]



Kuschel, R. *et al.*, *Angew. Chem. (Int.)*, 1993, **32**(11), 1632

Orange needles which decompose above -40°C and detonate if warmed rapidly to 0°C . The isomeric bromous anhydride would not be expected to be much more stable.

See other HALOGEN OXIDES

0277. Dibromine pentoxide

[]




Leopold, D. *et al.*, *Angew. Chem. (Int.)*, 1994, **33**(9), 975

The yellow powder decomposes above -40°C , sometimes by detonation.

See other HALOGEN OXIDES

0278. Lead bromate

[34018-28-5]



Sidgwick, 1950, 1227

An explosive salt.

See Lead acetate–lead bromate

See other HEAVY METAL DERIVATIVES, METAL HALOGENATES

0279. Zinc bromate

[14519-07-4]



See entry METAL HALOGENATES

0280. Sulfur dibromide

[14312-20-0]



Nitric acid

See Nitric acid: Sulfur halides

Sodium

See Sodium: Non-metal halides

See other NON-METAL HALIDES

0281. Silicon dibromide sulfide

[13520-74-6]

Br₂SSi

S:SiBr₂

Water

Bailar, 1973, Vol. 1, 1415

Hydrolysis of the sulfide is explosive.

See related NON-METAL HALIDES, NON-METAL SULFIDES

0282. Disulfur dibromide

[13172-31-1]

Br₂S₂

BrSSBr

HCS 1980, 873

Metals

1. Mellor, 1947, Vol. 10, 652

2. Pascal, 1960, Vol. 13.2, 1162

Thin sections of potassium or sodium usually ignite in the liquid bromide. Iron at about 650°C ignites and incandescs in the vapour [1]. Interaction with finely divided aluminium or antimony is violent [2].

Oxidants

Taradoire, F., *Bull. Soc. Chim. Fr.*, 1945, **12**, 95

Interaction with moist barium bromate is very violent, and mixtures with potassium bromate and water (3–4%) ignite at 20°C. In absence of water, ignition occurs at 125°C. Silver bromate also deflagrates.

See Silver bromate: Sulfur compounds

Nitric acid: Sulfur halides

Phosphorus

See Phosphorus: Non-metal halides

See other NON-METAL HALIDES

0283. Poly(dibromosilylene)

[14877-32-9]

(Br₂Si)_n

(SiBr₂)_n

Oxidants

Brauer, 1963, Vol. 1, 688

Ignites in air at 120°C and reacts explosively with oxidants such as nitric acid.

See related NON-METAL HALIDES

0284. Titanium dibromide
[13783-04-5]



Gibson, 1969, 60–61
It may ignite in moist air.
See other METAL HALIDES, PYROPHORIC MATERIALS

0285. Zirconium dibromide
[24621-17-8]



Air, or Water
Pascal, 1963, Vol. 9, 558
It ignites in air, reacts violently with water and incandescs in steam.
See other METAL HALIDES, PYROPHORIC MATERIALS

0286. Iron(III) bromide
[10031-26-2]



Potassium
See Potassium: Metal halides


Sodium
See Sodium: Metal halides
See other METAL HALIDES

†**0287. Tribromosilane**
[7789-57-3]



Schumb, W. C., *Inorg. Synth.*, 1939, **1**, 42
It usually ignites when poured in air (generating an extended liquid/air interface).
This ignition is remarkable in a compound containing over 89% of bromine.
See other HALOSILANES

0288. Molybdenum azide tribromide
[68825-98-9]



Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 510
It is highly explosive.
See other METAL AZIDE HALIDES

0289. Indium bromide
[13469-09-3]





Preparative hazard

See Mercury(II) bromide: Indium

See other METAL HALIDES

0290. Nitrogen tribromide hexaammoniate (Tribromamine hexaammoniate)

[]



Alone, or Non-metals

Mellor, 1967, Vol. 8, Suppl. 2, 417; 1940, Vol. 8, 605

The compound, formed by condensation of its vapour at -95°C explodes suddenly at -67°C . Prepared in another way, it is stable under water but explodes violently in contact with phosphorus or arsenic.

See other N-HALOGEN COMPOUNDS

0291. Tribromamine oxide (Nitrosyl tribromide)

[13444-89-8]



Sodium-antimony alloy

Mellor, 1940, Vol. 8, 621

The powdered alloy ignites when dropped into the vapour.

See other N-HALOGEN COMPOUNDS, N-OXIDES

0292. Vanadium tribromide oxide

[13520-90-6]



Water

Bailar, 1973, Vol. 3, 508

The bromide (and analogous chloride) is violently hygroscopic.

See related METAL HALIDES

0293. Phosphorus tribromide

[7789-60-8]



HCS 1980, 744

Calcium hydroxide, Sodium carbonate

Seager, J. F., *Chem. Brit.*, 1976, **12**, 105

During disposal of the tribromide by a recommended procedure involving slow addition to a mixture of soda ash and dry slaked lime, a violent reaction, accompanied by flame, occurred a few seconds after the first drop. Cautious addition of the bromide to a large volume of ice water is suggested for disposal.

Oxidants

See Chromyl chloride: Non-metal halides

Oxygen (Gas): Phosphorus tribromide

Ruthenium(VIII) oxide: Phosphorus tribromide

3-Phenylpropanol

Taylor, D. A. H., *Chem. Brit.*, 1974, **10**, 101–102

During dropwise addition of the bromide to the liquid alcohol, the mechanical stirrer stopped, presumably allowing a layer of the dense tribromide to accumulate below the alcohol. Later manual shaking caused an explosion, probably owing to the sudden release of gaseous hydrogen bromide on mixing.

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

Potassium

See Potassium: Non-metal halides

Sodium, Water

See Sodium: Non-metal halides

Sulfur acids

Dillon, K. B. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1979, 885–887

The tribromide is initially insoluble in 100% sulfuric acid, 25% oleum or fluoro-sulfuric acid, but violent exotherms occurred after contact for 11, 4 and 5 hours respectively. 65% Oleum reacts violently on contact, and chlorosulfuric acid vigorously after shaking.

See Phosphorus trichloride: Sulfur acids

Phosphorus triiodide: Sulfur acids

See other INDUCTION PERIOD INCIDENTS

1,1,1-Tris(hydroxymethyl)methane

1. Derfer, J. M. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 175

2. Farber, S. *et al.*, *Synth. Comm.*, 1974, **4**, 243

Interaction to form the corresponding tribromomethyl compound is extremely hazardous, even using previously specified precautions [1]. Several fires occurred in the effluent gases, and in reaction residues exposed to air [2], doubtless owing to phosphine or its derivatives.

Water

Mellor, 1940, Vol. 8, 1032

Interaction with warm water is very rapid and may be violent with limited quantities.

See other NON-METAL HALIDES

0294. Tungsten tetrabromide oxide
[13520-77-9] **Br₄OW**

O:WBr₄

Preparative hazard

See Boron tribromide: Tungsten trioxide

See related METAL HALIDES

0295. Selenium tetrabromide
[7789-65-3] **Br₄Se**

SeBr₄

Trimethylsilyl azide

See Trimethylsilyl azide: Selenium halides

See other NON-METAL HALIDES

0296. Tellurium tetrabromide
[10031-27-3] **Br₄Te**

TeBr₄

Ammonia

Sorbe, 1968, 154

Interaction gives a mixture of 'tritellurium' tetranitride and tellurium bromide nitride, which explodes on heating.

See Tetratellurium tetranitride

See other METAL HALIDES

0297. Tungsten azide pentabromide
[] **Br₅N₃W**

N₃WBr₅

Extremely explosive.

See entry METAL AZIDE HALIDES

0298. Carbon

[7440-44-0] (Amorphous)	C_n
[7782-42-5] (Graphite)	C_n
[99685-96-8;115383-22-7] (Fullerene)	C₆₀,C₇₀
[7782-40-3] (Diamond)	C_n
[126487-10-3] (Cyclooctadecanonayne)	C₁₈
C	

1. Anon., *Fire Prot. Assoc. J.*, 1964, 337

2. Cameron, A. *et al.*, *J. Appl. Chem.*, 1972, **22**, 1007

3. Zav'yalov, A. N. *et al.*, *Chem. Abs.*, 1976, **84**, 166521
4. Suzuki, E. *et al.*, *Chem. Abs.*, 1980, **93**, 78766
5. Beever, P. F., *Runaway Reactions*, 1981, Paper 4/X, 1–9
6. Van Liempt, J. H. M., *PT-Procestech.* (Neth.), 1988, **43**(9), 45–47
7. Anon., *Loss Prev. Bull.*, 1992, **105**, 15
8. Diederich, F. *et al.*, *Angewand. Chem. (Int.)*, 1994, **33**(9), 997
9. Editor's comments

Activated carbon exposed to air is a potential fire hazard because of its very high surface area and adsorptive capacity. Freshly prepared material may heat spontaneously in air, and presence of water accelerates this. Spontaneous heating and ignition may occur if contamination by drying oils or oxidising agents occurs [1]. The spontaneous heating effect has been related to the composition and method of preparation of activated carbon, and the relative hazards may readily be assessed [2]. Free radicals present in charcoal are responsible for auto-ignition effects, and charcoal may be stabilised for storage and transport without moistening by treatment with hot air at 50°C [3]. The causes of accidents involving activated carbon have been surveyed [4]. Fires originating in paper-bagged cargoes of active carbon in holds of ships travelling through tropical waters have been investigated by isothermal methods. Use of oxygen-impermeable plastic bags to limit oxidation and moisture uptake was proposed as a solution to the problem [5]. The potential hazards arising from use of activated carbon for various purposes are reviewed. Hazards usually arise from self-heating from adsorption of vapours or gases (especially oxygen), and may lead to autoignition, fire or explosions, including carbon dust explosions [6]. A fire in an activated charcoal odour abatement system, essentially caused by scale-up without considering the reduced heat loss thus occasioned, is reported [7].

Fullerenes may be expected to be at least as autoxidisable as charcoal (graphite) when suitably finely divided. Heats of formation (solid, w.r.t. graphite): C₆₀ 3.2 kJ/g; C₇₀ 3.0 kJ/g. They have hitherto proved surprisingly stable kinetically [8]. The cyclic C₁₈, when available in substantial quantity, will surely prove capable of spontaneous combustion and will probably be explosive [9].

Generation of pyrophoric carbon

See Barium acetylide

See also HIGH SURFACE-AREA SOLIDS, PETROLEUM COKE

Alkali metals

1. Bailar, 1973, Vol. 1, 443
2. Werner, H. *et al.*, *Fullerene Sci. Technol.*, 1993, **1**(2), 199

Graphite in contact with liquid potassium, rubidium or caesium at 300°C gives intercalation compounds (C₈M) which ignite in air and may react explosively with water [1]. 'Fullerene black'—probably a finely divided and distorted graphite—impregnated with potassium explodes spontaneously in air [2].

See Potassium: Carbon

Sodium: Non-metals

Chlorinated paraffins, Lead(IV) oxide, Manganese(IV) oxide

See Lead(IV) oxide: Carbon black, etc.

Dibenzoyl peroxide

See Dibenzoyl peroxide: Charcoal

1,4-Diazabicyclo[2.2.2]octane

Hardman, J. S. *et al.*, *Fuel*, 1980, **59**, 213–214

Activated carbon showed an auto-ignition temperature in flowing air of 452–518°C. Presence of 5% of the base ('triethylenediamine') adsorbed on the carbon reduced the AIT to 230–260°C. At high air flow rates an exotherm was seen at 230–260°, but ignition did not then occur until 500°C.

Iron(II) oxide, Oxygen (Liquid)

See Oxygen (Liquid): Carbon, Iron(II) oxide

Lithium, Lithium tetrachloroaluminate, Sulfinyl chloride

See Lithium: Carbon, etc.

Metal salts

MCA Case History No. 1094

Dry metal-impregnated charcoal catalyst was being added from a polythene bag to an aqueous solution under nitrogen. Static so generated ignited the charcoal dust and caused a flash fire. The risk was eliminated by adding a slurry of catalyst in water from a metal container.

See Cobalt(II) nitrate: Carbon

See other STATIC INITIATION INCIDENTS

Molybdenum(IV) oxide

See Molybdenum(IV) oxide: Graphite

2-Nitrobenzaldehyde

See 2-Nitrobenzaldehyde: Active carbon

Nitrogen oxide, Potassium hydrogen tartrate

See Nitrogen oxide: Carbon, Potassium hydrogen tartrate

Other reactants

Yoshida, 1980, 225

MRH values calculated for 25 combinations with oxidants are given.

Oxidants

MRH values show % of carbon

Carbon has frequently been involved in hazardous reactions, particularly finely divided or high-porosity forms exhibiting a high ratio of surface area to mass (up to 2000 m²/g). It then functions as an unusually active fuel which possesses adsorptive and catalytic properties to accelerate the rate of energy release involved in combustion reactions with virtually any oxidant. Less active forms of carbon will ignite or explode on suitably intimate contact with oxygen, oxides, peroxides, oxosalts, halogens, interhalogens and other oxidising species. Individual combinations are found under the entries listed below.

Ammonium perchlorate: Carbon	MRH 6.19/85
Bromine pentafluoride: Acids, etc.	
Bromine trifluoride: Halogens, etc.	
Chlorine trifluoride: Metals, etc.	
Cobalt(II) nitrate: Carbon	MRH 3.68/86
Dichlorine oxide: Carbon, or: Oxidisable materials	MRH 5.14/90
Fluorine: Non-metals	
Hydrogen peroxide: Carbon	MRH 6.19/15
Iodine heptafluoride: Carbon	
Iodine(V) oxide: Non-metals	
Nitrogen oxide: Non-metals	MRH 7.82/16
Nitrogen trifluoride: Charcoal	
Oxygen (Liquid): Charcoal	MRH (Gas) 8.95/27
Oxygen difluoride: Non-metals	
Ozone: Charcoal, Potassium iodide	
Peroxyformic acid: Non-metals	MRH 5.69/tr.
Peroxyfuroic acid: Alone, etc.	
Potassium chlorate: Charcoal, etc.	MRH 4.52/87
Potassium dioxide: Carbon	
Potassium nitrate: Non-metals	
Potassium permanganate: Non-metals	MRH 2.59/7
Silver nitrate: Non-metals	MRH 2.47/10
Sodium nitrate: Non-metals	
Sodium peroxide: Non-metals	MRH 2.30/7
'Trioxxygen difluoride': Various materials	
Zinc nitrate: Carbon	MRH 3.86/87

Potassium hydroxide

Hejduk, J., *Chem. Abs.*, 1989, **110**, 107229

Analytical decomposition of powdered diamond by fusion with potassium hydroxide may become explosive. This can be avoided by fusion with a potassium carbonate–sodium carbonate mixture, followed by addition of small portions of potassium nitrite or nitrate.

Sodium hydrogen carbonate

See Sodium hydrogen carbonate: Carbon, Water

Sodium tetrahydroborate

See Sodium tetrahydroborate

Turpentine

US Environmental Protection Agency, *Alert EPA 550-F-97-002e*, 1997

Fires have been caused when using activated carbon to de-odorise crude sulfate turpentine. This is another case of adsorption forming hot spots then exposed to air.

Unsaturated oils

1. Bahme, C. W., *NFPA Quart.*, 1952, **45**, 431
2. von Schwartz, 1918, 326

Unsaturated (drying) oils, like linseed oil, etc., will rapidly heat and ignite when distributed on active carbon, owing to the enormous increase in surface area of the oil exposed to air, and in the rate of oxidation, probably catalysed by metallic impurities [1]. A similar, but slower, effect occurs on fibrous materials such as cotton waste [2].


See other CATALYTIC IMPURITY INCIDENTS

See other NON-METALS

0299. Silver cyanide

[506-64-9]

CAgN



Phosphorus tricyanide

See Phosphorus tricyanide (reference 2)

Fluorine

See Fluorine: Metal salts

See other ENDOTHERMIC COMPOUNDS, METAL CYANIDES, SILVER COMPOUNDS

0300. Silver cyanate

[3315-16-0]

CAgNO



Sorbe, 1968, 125


It explodes on heating.

See related METAL CYANIDES (AND CYANO COMPLEXES) *See other* SILVER COMPOUNDS

0301. Silver fulminate

[5610-59-3]

CAgNO



1. Urbanski, 1967, Vol. 3, 157

2. Collins, P. H. *et al.*, *Propellants, Explos.*, 1978, **3**, 159–162

Silver fulminate is dimeric and rather endothermic ($\Delta H_f^\circ +361.5$ kJ/dimol, 1.21 kJ/g). It is readily formed from silver or its salts, nitric acid and ethanol, and is a much more sensitive and powerful detonator than mercuric fulminate [1]. The properties and applications have been reviewed [2].

Hydrogen sulfide

Boettger, A., *J. Pr. Chem.*, 1868, **103**, 309

Contact with hydrogen sulfide at ambient temperature initiates violent explosion of the fulminate.

See other ENDOTHERMIC COMPOUNDS, METAL FULMINATES, SILVER COMPOUNDS

- 0302. Silver trinitromethanide**
 [25987-94-4] **C_{Ag}N₃O₆**

$$\text{AgC(NO}_2)_3$$
 Witucki, E. F. *et al.*, *J. Org. Chem.*, 1972, **37**, 152
 The explosive silver salt may be replaced with advantage by the potassium salt in the preparation of 1,1,1-trinitroalkanes.
See other POLYNITROALKYL COMPOUNDS, SILVER COMPOUNDS
- 0303. Silver azidodithioformate**
 [74093-43-9] **C_{Ag}N₃S₂**

$$\text{AgSC(S)N}_3$$
 Sorbe, 1968, 126
 The tetrahydrated salt explodes on the slightest friction.
See Azidodithioformic acid
See related ACYL AZIDES *See other* SILVER COMPOUNDS
- 0304. Silver trichloromethanephosphonate**
 [] **C_{Ag}₂Cl₃O₃P**

$$(\text{AgO})_2\text{P(O)CCl}_3$$
 Yakubovich, A. Ya. *et al.*, *Chem. Abs.*, 1953, **47**, 2685i
 It explodes on heating.
See other SILVER COMPOUNDS
- 0305. Disilver cyanamide**
 [3884-87-0] **C_{Ag}₂N₂**

$$\text{Ag}_2\text{NC}\equiv\text{N}$$
 1. Chrétien, A. *et al.*, *Compt. rend.*, 1951, **232**, 1114
 2. Deb, S. K. *et al.*, *Trans. Faraday Soc.*, 1959, **55**, 106–113
 3. Cradock, S., *Inorg. Synth.*, 1974, **15**, 167
 During pyrolysis to silver (via silver dicyanamide), initial heating must be slow to avoid explosion [1]. High intensity illumination will also cause explosive decomposition of a confined sample [2]. Safety precautions for preparation and subsequent use of the explosive salt are detailed [3].
See other IRRADIATION DECOMPOSITION INCIDENTS
See other N-METAL DERIVATIVES, SILVER COMPOUNDS
- 0306. Disilver diazomethanide**
 [54086-40-7] **C_{Ag}₂N₂**

$$\text{Ag}_2\text{CN}_2$$
 Blues, E. T. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1974, 466–467

Both the disilver derivative and its precursory dipyridine complex are highly explosive and extremely shock-sensitive when dry.

See other DIAZO COMPOUNDS, SILVER COMPOUNDS

0307. Diazido(trifluoromethyl)arsine

[157951-76-3]

CAsF₃N₆



Ang, H. G. *et al.*, *Inorg. Chem.*, 1994, **33**(20), 4425

A vapour phase study of this azide showed it sometimes to explode when exposed to heat.

See other AZIDES

0308. Gold(I) cyanide

[506-65-0]

CAuN



Magnesium

See Magnesium: Metal cyanides

See other GOLD COMPOUNDS, METAL CYANIDES

0309. Carbon tetraboride

[12069-32-8]

CB₄



Chlorine trifluoride

See Chlorine trifluoride: Boron-containing materials

See related NON-METALS

0310. Bromotrichloromethane

[75-62-7]

CBrCl₃



Ethylene

See Ethylene: Bromotrichloromethane

See other HALOALKANES

0311. Bromotrifluoromethane

[75-63-8]

CBrF₃




Aluminium

See Aluminium: Halocarbons

See other HALOALKANES

0312. Bromine(I) trifluoromethanesulfonate
[70142-16-4]

CBrF₃O₃S



Readily oxidisable materials

Katsuhara, Y. *et al.*, *J. Org. Chem.*, 1980, **45**, 2442

Contact of this oxidant with readily oxidisable materials may lead to explosions.

See Chlorine(I) trifluoromethanesulfonate

See related HYPOHALITES

0313. Cyanogen bromide
[506-68-3]

CBrN



HCS 1980, 338

1. Grossman, M. I., *Chem. Eng. News*, 1980, **58**(35), 43

2. Harvey, W. R. *et al.*, *Chem. Eng. News*, 1981, **59**(5), 49

3. Crocker, H. P., *Chem. Eng. News*, 1981, **59**(5), 49

Cyanogen bromide is moderately endothermic ($\Delta H_f^\circ(g)$ +50 kJ/mol, 0.47 kJ/g) and shows evidence of instability. The plastic cap of a bottle stored in a laboratory for several years on a high shelf, occasionally at 31°C, shattered and drove fragments into the shelf above [1]. This instability was confirmed, and a procedure outlined to obviate the use of the bromide in autoanalysis by generating cyanogen chloride on demand from Chloramine-T and potassium cyanide [2]. A 50 wt% solution of the bromide in chloroform is a stable and convenient form for use [3].
See Cyanogen chloride

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0314. Tribromonitromethane
[464-10-8]

CBr₃NO₂



Sorbe, 1968, 40

It is used as an explosive.

See related NITROALKANES

0315. Carbon tetrabromide (Tetrabromomethane)
[558-13-4]

CBr₄



HCS 1980, 283

Hexacyclohexyldilead

See Hexacyclohexyldilead: Halocarbons

Lithium

See Lithium: Halocarbons

See other HALOALKANES

0316. Calcium cyanamide

[156-62-7]



CCaN₂

Water

Pieri, M., *Chem. Abs.*, 1952, **46**, 8335i

Absorption of water during handling or storage of technical calcium cyanamide may cause explosions, owing to liberation of acetylene from the calcium carbide content (up to 2%). Precautions are discussed.

See other CYANO COMPOUNDS, N-METAL DERIVATIVES

0317. Calcium carbonate

[471-34-1]



CCaO₃

HCS 1980, 265

Fluorine

See Fluorine: Metal salts

See other INORGANIC BASES

0318. (Chlorocarbonyl)imidodisulfur difluoride

[53654-97-0]



CClF₂NOS

Mews, R. *et al.*, *Inorg. Synth.*, 1986, **24**, 14–16

The title compound, particularly if impure, rapidly forms explosive decomposition products when stored at ambient temperature.

See other ACYL HALIDES, N-S COMPOUNDS

0319. Chloroperoxytrifluoromethane

[32755-26-3]



CClF₃O₂

Tetrafluoroethylene

Ratcliffe, C. T. *et al.*, *J. Amer. Chem. Soc.*, 1971, **93**, 3887–3888

The peroxy compound initiated explosive polymerisation of tetrafluoroethylene when a mixture prepared at –196°C warmed to –110°C.

See other POLYMERISATION INCIDENTS

See related HYPOHALITES See other ORGANIC PEROXIDES

0320. Chlorine(I) trifluoromethanesulfonate

[65597-24-2]

CCIF₃O₃S



Readily oxidisable materials

1. Katsuhara, Y. *et al.*, *J. Amer. Chem. Soc.*, 1979, **101**, 1040
2. Katsuhara, Y. *et al.*, *J. Org. Chem.*, 1980, **45**, 2442

Contact of the oxidant with readily oxidisable materials must be controlled and at low temperatures to prevent explosions.

See Bromine(I) trifluoromethanesulfonate

See related HYPOHALITES

0321. Trifluoromethyl perchlorate

[52003-45-9]

CCIF₃O₄



Preparative hazard

1. Schack, C. J. *et al.*, *Inorg. Chem.*, 1974, **13**, 2375
2. Schack, C. J. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1974, **10**, 449

Though apparently not explosively unstable [1], its synthesis [2] was occasionally accompanied by deflagrations.

See other ALKYL PERCHLORATES

0322. Trifluoromethanesulfonyl chloride

[421-17-0]

CCIF₃S



Chlorine fluorides

Sprenger, G. H. *et al.*, *J. Fluorine Chem.*, 1976, **7**, 335

In the preparation of trifluoromethylsulfur trifluoride, the chloride must be dissolved in a fully halogenated solvent to prevent explosion during treatment with chlorine fluoride or chlorine trifluoride.

See other HALOGENATION INCIDENTS

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

See other ACYL HALIDES

0323. Cyanogen chloride

[506-77-4]

CCIN



HCS 1980, 339

Preparative hazard

Good, R. J., private comm., 1979

Crude cyanogen chloride (endothermic, prepared from hydrogen cyanide and chlorine) may trimerise violently to cyanuric chloride, catalysed by traces of hydrogen chloride or ammonium chloride.

See other CATALYTIC IMPURITY INCIDENTS

See Ammonium chloride

See Cyanogen bromide

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0324. Chlorosulfonyl isocyanate

[1189-71-5]



CCINO₃S

Water

Graf, R., *Org. Synth.*, 1973, Coll. Vol. 5, 228


Interaction is violent.

See related CYANO COMPOUNDS

See other N-S COMPOUNDS, ACYL HALIDES

0325. Triazidocarbenium perchlorate (Triazidomethylum perchlorate)

[]



CCIN₉O₄

Petrie, M. A. *et al.*, *J. Amer. Chem. Soc.*, 1997, **119**, 8802

All salts of this cation were friction and shock sensitive, too sensitive for actual use as explosives. The most sensitive though, with estimated energy of 5.4 kJ/g, not the most powerful, was the perchlorate. Even though prepared on low milligram scale this invariably detonated spontaneously during work-up or soon after. The more powerful dinitramide (5.8 kJ/g) decomposed non-explosively on standing.

See other ORGANIC AZIDES

0326. Dichlorodifluoromethane (Freon 12)

[75-71-8]



CCl₂F₂

Aluminium

See Aluminium: Halocarbons

Magnesium

See Magnesium: Halocarbons

Water

See LIQUEFIED GASES: WATER *See other* HALOALKANES

0327. Phosphoryl dichloride isocyanate
[870-30-4]

CCl₂NO₂P



Preparative hazard

See N-Carbomethoxyiminophosphoryl chloride

See related CYANO COMPOUNDS, NON-METAL HALIDES

0328. Dichlorodinitromethane
[1587-41-3]

CCl₂N₂O₄



Hocking, M. B. *et al.*, *Chem. & Ind.*, 1976, 952

It exploded during attempted distillation at atmospheric pressure, but was distilled uneventfully at 31°C/13 mbar.

See other POLYNITROALKYL COMPOUNDS

0329. Carbonyl dichloride (Phosgene)
[75-44-5]

CCl₂O



(MCA SD-95, 1967); FPA H105, 1981; HCS 1980, 740

Anon., *Sichere Chemiarb.*, 1988, **40**(3), 34

Trichloromethyl chloroformate (diphosgene) is used as a safe substitute for highly toxic phosgene gas. The latter is generated in situ by addition of catalytic amounts of tertiary amines or amides, or active carbon. Diphosgene also disproportionates to 2 equivalents of phosgene on heating above 250°C.

N,N-Dimethylbenzeneamine, Water

Anon., *J. Loss Prev.*, 1994, **7**(3), 257

The aniline was being phosgenated in toluene as solvent, the reaction ran wild and ejected more than 3 tonnes of reactor contents. This is believed to have been due to water contamination, possibly as ice. An initial charge of only part of the phosgene failed to show the exotherm anticipated if water was present, however this may not have been enough, nor was the thermocouple immersed in the solvent.

See Diprotium monoxide

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Other reactants

Yoshida, 1980, 341

MRH values for 18 combinations with a range of materials are given.

Potassium

MRH 4.76/45

See Potassium: Non-metal halides

Sodium azide

See *tert*-Butyl azidoformate (reference 4)

See related ACYL HALIDES

0330. Trichlorofluoromethane (Freon 11)

[75-69-4]



CClF_3

HCS 1980, 921

Metals

See Aluminium: Halocarbons

Barium: Halocarbons

Lithium: Halocarbons

METAL-HALOCARBON INCIDENTS

See other HALOALKANES

0331. Trichloronitromethane ('Chloropicrin')

[76-06-2]



CCl_3NO_2

HCS 1980, 312

Anon., *Chem. Eng. News*, 1972, **50**(38), 13

Tests showed that above a critical volume, bulk containers of trichloronitromethane can be shocked into detonation. Containers below 700 kg content will now be the maximum size as against rail-tanks previously used.

Aniline

Jackson, K. E., *Chem. Rev.*, 1934, **14**, 269

Reaction at 145°C with excess aniline is violent.

3-Bromopropyne

BCISC Quart. Safety Summ., 1968, **39**, 12

An insecticidal mixture in a rail-tank exploded with great violence during pump-transfer operations, possibly owing to the pump running dry and overheating. Both components of the mixture are explosive and the mixture was also found to be shock- and heat-sensitive.

Sodium hydroxide

Scholtz, S., *Explosivstoffe*, 1963, **11**, 159, 181

During destruction of chemical warfare ammunition, pierced shells containing chloropicrin reacted violently with alcoholic sodium hydroxide.

Sodium methoxide

Ramsey, B. G. *et al.*, *J. Amer. Chem. Soc.*, 1966, **88**, 3059

During addition of the nitrocompound in methanol to sodium methoxide solution, the temperature must not be allowed to fall much below 50°C. If this happens, excess nitro compound will accumulate and cause a violent and dangerous exotherm.

See related NITROALKANES

0332. Carbon tetrachloride (Tetrachloromethane)

[56-23-5]

CCl₄

CCl₄

(MCA SD-3, 1963); HCS 1980, 284; RSC Lab. Hazard Data Sheet No. 59, 1987

Aluminium oxide, Heavy metals

See Aluminium oxide: Halocarbons, etc.

Aluminium chloride, Triethylaluminium

See Triethyldialuminium trichloride: Carbon tetrachloride

Boranes

See BORANES: carbon tetrachloride

Calcium disilicide

See Calcium disilicide: Carbon tetrachloride

Calcium hypochlorite

See Calcium hypochlorite: Carbon tetrachloride

Chlorine trifluoride

MRH 1.21/44

See Chlorine trifluoride: Carbon tetrachloride

Decaborane(14)

See Decaborane(14): Ether, etc.

1,11-Diamino-3,6,9-triazaundecane ('Tetraethylenepentamine')

1. Hudson, F. L., private comm., 1973
2. Collins, R. F., *Chem. & Ind.*, 1957, 704

A mixture erupted vigorously one hour after preparation [1]. Interaction (not vigorous) of amines and halocarbons at ambient temperature had been recorded previously [2]. The presence of 5 basic centres in the viscous amine would be expected to enhance exothermic effects.

Dibenzoyl peroxide, Ethylene

See Ethylene: Carbon tetrachloride

Dibenzoyl peroxide: Carbon tetrachloride, Ethylene

See also WAX FIRE

Dimethylformamide

MRH 1.30/28

1. 'DMF Brochure' Billingham, ICI, 1965
2. Cardillo, P. *et al.*, *Ann. Chim.* (Rome), 1984, 74, 129–133

There is a potentially dangerous reaction of carbon tetrachloride with dimethylformamide in presence of iron. The same occurs with 1,2,3,4,5,6-hexachlorocyclohexane, but not with dichloromethane or 1,2-dichloroethane under the same conditions [1]. A quantitative study of the reaction by DSC and ARC techniques shows that in a 1:1 wt. mixture with carbon tetrachloride in absence of iron, an exothermic reaction sets in below 100°C. Under adiabatic conditions, the heat release (207.6 J/g) would take a runaway reaction to over 240°C. In presence of 3% of iron powder, the same mixture shows 2 exotherms, one at 56°C (108 J/g) and the second at 94°C (275 J/g), a final adiabatic temperature exceeding 285°C being possible [2]. Dimethylacetamide behaves similarly but more so.

See N,N-Dimethylacetamide: Halogenated compounds

See other CATALYTIC IMPURITY INCIDENTS

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Halocarbons

Ethanol (or methanol), Potassium sulfide

See Potassium dithioformate

See Sodium: Methanol and Sodium methoxide, both below

Fluorine

See Fluorine: Halocarbons

Metals

MRH values show % of carbon tetrachloride

<i>See</i> Aluminium: Halocarbons	MRH 4.14/81
Barium: Halocarbons	MRH 4.18/36
Beryllium: Halocarbons	
Lithium: Halocarbons	MRH 8.24/85
Potassium: Halocarbons	MRH 5.19/50
Potassium–sodium alloy: Halocarbons	
Sodium: Halocarbons	MRH 6.10/63
Uranium: Carbon tetrachloride	
Zinc: Halocarbons	MRH 2.43/54

METAL–HALOCARBON INCIDENTS

Methanol

See Methanol: Carbon tetrachloride, Metals

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

Other reactants

Yoshida, 1980, 148–149

MRH values for 23 combinations with a range of materials are given.

2-Propen-1-ol

MRH 1.42/27

See 2-Propen-1-ol: Carbon tetrachloride

See other HALOALKANES

0333. Tetrachlorotrifluoromethylphosphorane

[1066-48-4]

CCl₄F₃P

Cl₄PCF₃

Tetramethyllead

Yap, N. T. *et al.*, *Inorg. Chem.*, 1979, **18**, 1304

Interaction under vacuum to form methyltrichlorotrifluoromethylphosphorane is hazardous, violent explosions having occurred twice after 30 min at ambient temperature.

See related ALKYLHALOPHOSPHINES

0334. Trichloromethyl perchlorate

[]

CCl₄O₄

Cl₃COCIO₃

Sidgwick, 1950, 1236

An extremely explosive liquid, only capable of preparation in minute amounts.

Oxygen balance is 100%

See Silver perchlorate: Carbon tetrachloride

See other ALKYL PERCHLORATES

0335. Triazidomethylum hexachloroantimonate

[19708-47-5]

CCl₆N₉Sb

[C(N₃)₃]⁺ [SbCl₆]⁻

Müller, U. *et al.*, *Angew. Chem.*, 1966, **78**, 825

The salt is sensitive to shock or rapid heating.

See related ORGANIC AZIDES

0336. Dideuterodiazomethane

[14621-84-2]

CD₂N₂

D₂CN₂

Gassman, P. G. *et al.*, *Org. Synth.*, 1973, **53**, 38–43

The explosive properties will be similar to those of diazomethane, for which precautions are extensively summarised.

See other DIAZO COMPOUNDS

0337. Poly(carbon monofluoride)

[25136-85-0]

(CF)_n

(Complex structure)

Preparative hazard

See Fluorine: Graphite

Hydrogen

Bailar, 1973, Vol. 1, 1269

Above 400°C in hydrogen, deflagration and flaming of the polymer occurs, the vigour depending on the fluorine content. Rapid heating to 500°C in an inert atmosphere causes explosive deflagration.

See related HALOALKANES

0338. Fluorocarbonylperoxonitrate

[]

CFNO₅

FC(O)OONO₂

Scheffler, D. *et al.*, *Inorg. Chem.*, 1997, **36**(3), 339

Preparation has been achieved; working with more than millimolar quantities is not advised. It is an explosive oxidant in contact with organics and can scarcely be safe in their absence.

See also Trifluoromethylperoxonitrate

See other PEROXYESTERS

0339. Azidocarbonyl fluoride (Fluorocarbonyl azide)

[23143-88-6]

CFN₃O

F.CO.N₃

Mack, H.-G. *et al.*, *J. Mol. Struct.*, 1994, **291**(2-3), 197

It is advised that the explosive gas be handled only in millimolar quantities and with due precautions.

See other ACYL AZIDES

0340. Fluorotrinitromethane

[1840-42-2]

CFN₃O₆

FC(NO₂)₃

Preparative hazard

See Fluorine: Trinitromethane


Nitrobenzene

Zotov, E. V. *et al.*, *Chem. Abs.*, 1980, **93**, 49732

Detonation characteristics of liquid explosive mixtures with nitrobenzene were studied.

See other POLYNITROALKYL COMPOUNDS

0341. Fluorodinitromethyl azide
[17003-82-6]



CFN₅O₄

Unstable at ambient temperature.

See entry FLUORODINITROMETHYL COMPOUNDS

See other ORGANIC AZIDES, POLYNITROALKYL COMPOUNDS

0342. Difluorodiazirine
[693-85-6]



CF₂N₂

Craig, N. C. *et al.*, *Spectrochim. Acta*, 1979, **35A**, 895

Explosive decomposition occurred during purification by GLC and when running the gas phase laser Raman spectrum.

See other IRRADIATION DECOMPOSITION INCIDENTS, DIAZIRINES

0343. Carbonyl difluoride
[353-50-4]



CF₂O

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

See related ACYL HALIDES

0344. Difluoro-*N*-fluoromethanimine
[338-66-9]



CF₃N

Ginsberg, V. A. *et al.*, *Zh. Obsch. Khim.*, 1967, **37**, 1413

The gas boils at -60°C and explodes in contact with flame.

See other *N*-HALOGEN COMPOUNDS

0345. Nitrosotrifluoromethane
[334-99-6]



CF₃NO

1. Spaziante, P. M., *Intern. Rev. Sci.: Inorg. Chem. Ser. 1*, 1972, **3**, 141

2. Banks, R. E. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1974, 2534–2535

Suggestions of untoward hazard inherent in the preparation of nitrosotrifluoromethane by pyrolysis of trifluoroacetyl nitrite [1] are discounted in the later reference, which gives full details of the equipment and procedure that had been used uneventfully during the previous decade [2].

See other NITROSO COMPOUNDS

0346. Trifluoromethyl peroxonitrate

[50311-48-3]



Hohorst, F. A. *et al.*, *Inorg. Chem.*, 1974, **13**, 715

A small sample exploded under a hammer blow.

See other PEROXYESTERS

0347. Trifluoromethyl azide (Azidotrifluoromethane)

[3802-95-7]



Schack, C. J., *J. Fluorine Chem.*, 1981, **18**, 584

Potentially hazardous, it must be handled with caution.

See other ORGANIC AZIDES

0348. Trifluoromethylsulfonyl azide

[3855-45-6]



Cavender, C. J. *et al.*, *J. Org. Chem.*, 1972, **37**, 3568

An explosion occurred when the azide separated during its preparation in the absence of solvent.

See other ACYL AZIDES

0349. Carbon tetrafluoride (Tetrafluoromethane)

[75-73-0]



Aluminium

See Aluminium: Halocarbons

Oxygen, Hydrogen

See HALOALKANES

See other HALOALKANES, FLUOROCARBONS

0350. Perfluoroformamidine

[14362-70-0]



CF_4N_2


Koshar, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 3859, 3862

It explodes on shock or phase change.

See other *N, N, N'*-TRIFLUOROAMIDINES

0351. Tetrafluorodiaziridine

[17224-09-8]



CF_4N_2

Firth, W. C., *J. Org. Chem.*, 1968, **33**, 3489, 3491

An explosive oxidant, readily initiated by shock or phase changes during condensation or evaporation, especially around -160°C . Handle only small quantities at moderately low temperatures.

See other *N*-HALOGEN COMPOUNDS

0352. Tetrafluorourea

[10256-92-5]



$\text{CF}_4\text{N}_2\text{O}$

Acetonitrile

Fraser, G. W. *et al.*, *Chem. Comm.*, 1966, 532

The solution of the difluoroamide in acetonitrile prepared at -40°C must not be kept at ambient temperature, since difluorodiazene is formed.

See Acetonitrile: *N*-Fluoro compounds

See other DIFLUOROAMINO COMPOUNDS

0353. Trifluoromethyl hypofluorite

[373-91-1]



CF_4O

Hydrocarbons

Allison, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1959, **81**, 1089–1091

In absence of nitrogen as diluent, interaction with acetylene, cyclopropane or ethylene is explosive on mixing, and solutions in benzene explode on sparking or UV irradiation.

See other IRRADIATION DECOMPOSITION INCIDENTS

Hydrogen-containing solvents

1. *Catalogue G-7*, 10, Gainesville (Fa.), Peninsular Chem Research, 1973
2. Barton, D. H. R. *et al.*, *Chem. Comm.*, 1968, 804
3. Robins, M. J. *et al.*, *J. Amer. Chem. Soc.*, 1976, **89**, 7389 (footnote 24)

Contact of the extremely reactive compound with hydrogen-containing solvents or conventional plastics tubing, even at -80°C , is undesirable [1]. Fully halogenated solvents are preferred, and some general precautions are described [2]. Violent explosions have occurred on introducing the gas directly into methanol [3].

Lithium

Porter, R. S. *et al.*, *J. Amer. Chem. Soc.*, 1957, **79**, 5625

Interaction set in at about 170°C with a sufficient exotherm to melt the glass container.

See other GLASS INCIDENTS

Polymers

Barton, D. H. R. *et al.*, *J. Org. Chem.*, 1972, **37**, 329

It is a powerful oxidant and only all-glass apparatus, free of polythene, PVC, rubber or similar elastomers should be used. Appreciable concentrations of the gas in oxidisable materials should be avoided.

Pyridine

Barton, D. H. R. *et al.*, *Chem. Comm.*, 1968, 804 (footnote)

Use of pyridine as an acid-acceptor in reactions involving trifluoromethyl hypofluorite is discouraged, as a highly explosive by-product is formed.

See other HYPOHALITES, OXIDANTS

0354. Trifluoromethanesulfinyl fluoride

[812-12-4]



CF_4OS

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

See other ACYL HALIDES

0355. Difluoromethylene dihypofluorite

[16282-67-0]



CF_4O_2

Haloalkenes

Hohorst, F. A. *et al.*, *Inorg. Chem.*, 1968, **7**, 624

Attempts to react the oxidant with *trans*-dichloroethylene or tetrafluoroethylene at ambient temperature in absence of diluent caused violent explosions. The oxidant should not be allowed to contact organic or easily oxidised material without adequate precautions.

See other HYPOHALITES

0356. Xenon(II) fluoride trifluoromethanesulfonate
[39274-39-0]

CF₄O₃SXe



Wechsberg, M. *et al.*, *Inorg. Chem.*, 1972, **11**, 3066

Unless a deficiency of xenon fluoride was used in the preparation at 0°C or below, the title product exploded violently on warming to ambient temperature.

See other XENON COMPOUNDS

0357. N,N-Difluorotrifluoromethylamine (Pentafluoromethanamine)
[335-01-3]

CF₅N




Schack, C. J., *J. Fluorine Chem.*, 1981, **18**, 584

Potentially hazardous, handle with caution.

See other DIFLUOROAMINO COMPOUNDS

0358. 3-Difluoroamino-1,2,3-trifluorodiaziridine
[17224-08-7]

CF₅N₃



Firth, W. C., *J. Org. Chem.*, 1968, **33**, 3489, 3491

An explosive oxidant, readily initiated by shock or phase change during condensation or evaporation. Only small samples should be used and at moderately low temperatures.

See other DIFLUOROAMINO COMPOUNDS, N-HALOGEN COMPOUNDS

0359. Pentafluoroguanidine
[10051-06-6]

CF₅N₃



Zollinger, J. L. *et al.*, *J. Org. Chem.*, 1973, **38**, 1070–1071

This, and several of its adducts with alcohols, are shatteringly explosive compounds, frequently exploding during phase changes at low temperatures, or on friction or impact.

Liquid fuels

Scurlock, A. C. *et al.*, US Pat. 3 326 732, 1967

This compound with multiple N–F bonding is useful as an oxidant in propellant technology.

See related *N, N, N'*-TRIFLUOROAMIDINES

See other DIFLUOROAMINO COMPOUNDS, *N*-HALOGEN COMPOUNDS

0360. Difluorotrifluoromethylphosphine oxide

[19162-94-8]



CF₅OP

Preparative hazard

See Dinitrogen tetraoxide: Difluorotrifluoromethylphosphine

See related ALKYLHALOPHOSPHINES

0361. Difluorotrifluoromethylphosphine

[1112-04-5]



CF₅P


Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Difluorotrifluoromethylphosphine

See other ALKYLHALOPHOSPHINES

0362. Bis(difluoroamino)difluoromethane

[4394-93-8]



CF₆N₂

Koshar, R. J. *et al.*, *J. Org. Chem.*, 1966, **31**, 4233

Explosions occurred during the handling of this material, especially during phase transitions. Use of protective equipment is recommended for preparation, handling and storage, even on the microscale.

See other DIFLUOROAMINO COMPOUNDS

0363. Trifluoromethylsulfur trifluoride

[374-10-7]



CF₆S


Preparative hazard

See Trifluoromethanesulfonyl chloride: Chlorine fluorides

See other ALKYLNON-METAL HALIDES

0364. Tris(difluoroamino)fluoromethane

[14362-68-6]



CF₇N₃

Koshar, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 3859, 3862

A shock-sensitive explosive, especially in the liquid state (b.p. 5.6°C).

Pentaborane(9)

Marcellis, A. W. *et al.*, US Pat. 4 376 665, 1983

Mixtures prepared at cryogenic temperatures, then allowed to warm, are air-sensitive and powerful explosives.

See other DIFLUOROAMINO COMPOUNDS

0365. Iron carbide

[12011-67-5]



CFe_3

Halogens

Mellor, 1946, Vol. 5, 898


Incandescens in chlorine below 100°C, and in bromine at that temperature.

See other METAL NON-METALLIDES

0366. Silver tetrazolide

[13086-63-0]

CHAgN_4



Thiele, J., *Ann.*, 1892, **270**, 59

It explodes on heating.

See other N-METAL DERIVATIVES, SILVER COMPOUNDS, TETRAZOLES

0367. Fluorohydrocyanokrypton(II) hexafluoroarsenate

[]



CHAsF_7KrN

Schrobilgen, G. J., *J. Chem. Soc., Chem. Comm.*, 1988, 863

A white solid precipitating from hydrogen cyanide and fluorokrypton in hydrogen fluoride is attributed this structure. It decomposes rapidly above -50°C , usually with violent detonation.

See other XENON COMPOUNDS

0368. Bromoform (Tribromomethane)

[75-25-2]



CHBr_3

Acetone, Potassium hydroxide

1. Willgerodt, C., *Ber.*, 1881, **14**, 2451

2. Weizmann, C. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 1189

Interaction in presence of powdered potassium hydroxide (or other bases) is violently exothermic [1], even in presence of diluting solvents [2].

See Chloroform: Acetone, etc.

Cyclic poly(ethylene oxides) ('Crown ethers'), Potassium hydroxide

Le Goaller, R. *et al.*, *Synth. Comm.*, 1982, **12**, 1163–1169

Crown ethers promote dihalocarbene formation from chloroform or bromoform and potassium hydroxide. However, in absence of diluent dichloromethane, dropwise addition of bromoform to the base in cyclohexane led to explosions.

Metals

See Potassium: Halocarbons

Lithium: Halocarbons

METAL–HALOCARBON INCIDENTS

See other HALOALKANES

0369. Chlorodifluoromethane (Freon 22)

[75-45-6]

CHClF₂

ClCHF₂

HCS 1980, 668 (cylinder)

Sand, J. R., *ASHRAE J.*, 1982, **24**(5), 38–40

At elevated pressures, mixtures of the 'refrigerant 22' gas with 50% of air are combustible (though ignition is difficult) and a 6–8-fold pressure increase may occur in closed systems if ignition occurs.

Aluminium


See Aluminium: Halocarbons

See other HALOALKANES

0370. Tetrazole-5-diazonium chloride

[27275-90-7]

CHCIN₆




Shevlin, P. B. *et al.*, *J. Amer. Chem. Soc.*, 1977, **99**, 2628

The crystalline diazonium salt will detonate at the touch of a spatula. An ethereal solution exploded violently after 1 h at -78°C , presumably owing to separation of the solid salt.

See 5-Aminotetrazole: Nitrous acid
See other DIAZONIUM SALTS, TETRAZOLES

0371. 1-Dichloroaminotetrazole
[68594-17-2]

CHCl₂N₅



Karrer, 1950, 804

1-Dichloroaminotetrazole and its 5-derivatives are extremely explosive, as expected in an *N,N*-dichloro derivative of a high-nitrogen nucleus.

See other *N*-HALOGEN COMPOUNDS, TETRAZOLES

0372. Chloroform (Trichloromethane)
[67-66-3]

CHCl₃

HCCl₃

(MCA SD-89, 1962); HCS 1980, 301; RSC Lab. Hazard Data Sheet No. 44, 1986

Acetone, Alkali

1. Willgerodt, C., *Ber.*, 1881, **14**, 258
2. King, H. K., *Chem. & Ind.*, 1970, 185
3. Hodgson, J. F., *Chem. & Ind.*, 1970, 380
4. Ekely, J. B. *et al.*, *J. Amer. Chem. Soc.*, 1924, **46**, 1253
5. Grew, E. L., *Chem. & Ind.*, 1970, 491
6. Grant, D. H., *Chem. & Ind.*, 1970, 919

Chloroform and acetone interact vigorously and exothermally in presence of solid potassium hydroxide or calcium hydroxide to form 1,1,1-trichloro-2-hydroxy-2-methylpropane [1], and a laboratory incident involving the bursting of a solvent residues bottle was attributed to this reaction. Addition of waste chloroform to a Winchester containing acetone, ether, and petroleum ether chromatography solvent residues led to a vigorous exothermic reaction, the effects of which would be greatly augmented by evolution of vapour of the much lower-boiling solvents in the narrow-necked bottle [2]. No reaction whatsoever occurs in the absence of base, even at 150°C under pressure [1], and the mechanism of the reaction was indicated [3]. Other haloforms and lower ketones react similarly in presence of base [4], but other halocarbons with a less activated hydrogen atom (e.g. dichloromethane) do not undergo the reaction, though there is an exotherm (physical effect) on mixing with acetone [5]. A minor eruption (or sudden boiling) of a chloroform-acetone mixture in new glassware may have been caused by surface alkali [6].

See Bromoform: Acetone, etc.

See other CATALYTIC IMPURITY INCIDENTS, GLASS INCIDENTS

Bis(dimethylamino)dimethylstannane

See Bis(dimethylamino)dimethylstannane: Chloroform

Dinitrogen tetroxide

See Dinitrogen tetroxide: Halocarbons

Ethanol (or methanol), Potassium sulfide

See Potassium dithioformate

See Sodium, Methanol; and Sodium methoxide, below

Fluorine

See Fluorine: Halocarbons

Metals

MRH Aluminium 3.85/18, magnesium 5.35/23, potassium 4.94/50, sodium 5.81/37

Davis, T. L. *et al.*, *J. Amer. Chem. Soc.*, 1938, **60**, 720–722

The mechanism of the explosive interaction on impact of chloroform with sodium or potassium has been studied.

See other METAL–HALOCARBON INCIDENTS

Nitromethane

See Nitromethane: Haloforms

Other reactants

Yoshida, 1980, 114

MRH values calculated for 14 combinations with a range of materials are given.

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

Sodium, Methanol

Unpublished information, 1948

During attempted preparation of trimethyl orthoformate, addition of sodium to an inadequately cooled chloroform–methanol mixture caused a violent explosion.

See Ethanol (or methanol), Potassium sulfide, above

See Sodium methoxide, below

Sodium hydroxide, Methanol

1. *MCA Case History No. 498*

2. *MCA Case History No. 1913*

A chloroform–methanol mixture was put into a drum contaminated with sodium hydroxide. A vigorous reaction set in and the drum burst. Chloroform normally reacts slowly with sodium hydroxide owing to the insolubility of the latter. The presence of methanol (or other solubiliser) increases the rate of reaction by increasing the degree of contact between chloroform and alkali [1]. Addition of chloroform

to a 4:1 mixture of methanol and 50 w/v% sodium hydroxide solution caused the drum to burst [2].

Sodium methoxide

1. *MCA Case History No. 693*

2. Kaufmann, W. E. *et al.*, *Org. Synth.*, 1944, Coll. Vol. 1, 258

For the preparation of methyl orthoformate, solid sodium methoxide, methanol and chloroform were mixed together. The mixture boiled violently and then exploded [1]. The analogous preparation of ethyl orthoformate [2] involves the slow addition of sodium or sodium ethoxide solution to a chloroform-ethanol mixture. The explosion was caused by the addition of the solid sodium methoxide as one portion.

See Sodium, etc., above

See Ethanol (or methanol), Potassium sulfide, above

Triisopropylphosphine

See Triisopropylphosphine: Chloroform

See other HALOALKANES

0373. Fluorodiiodomethane

[1493-01-2]



Preparative hazard

See Mercury(I) fluoride: Iodoform

See other HALOALKANES

0374. Fluorodinitromethane

[7182-87-8]



Potentially explosive.

See entry FLUORODINITROMETHYL COMPOUNDS

See other POLYNITROALKYL COMPOUNDS

0375. Trifluoromethanesulfonic acid ('Triflic acid')

[1493-13-6]



Acyl chlorides, Aromatic hydrocarbons

Effenberger, F. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1972, **11**, 300

Addition of catalytic amounts (1%) of the acid (stronger even than perchloric acid) to mixtures of acyl chlorides and aromatic hydrocarbons causes more or less violent evolution of hydrogen chloride, depending on the reactivity of the Friedel-Crafts components.

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

Perchlorate salts

Dixon, N. E. *et al.*, *Inorg. Synth.*, 1986, **24**, 245

Because trifluoromethanesulfonic acid is a stronger acid than perchloric acid, under no circumstances should perchlorate salts be used with the neat acid, because the hot anhydrous perchloric acid so formed represents an extreme explosion hazard, especially in contact with transition metal complexes (or with organic materials).

See Perchloric acid: Dehydrating agents

See other ORGANIC ACIDS

0376. Iodoform (Triiodomethane)

[75-47-8]

CHI₃

CHI₃

Acetone

See Chloroform: Acetone, etc.

Hexamethylenetetramine

Sorbe, 1968, 137

The 1:1 addition complex exploded at 178°C.

Mercury(I) fluoride

See Mercury(I) fluoride: Iodoform

Silver

See Silver: Iodoform

See other ENDOTHERMIC COMPOUNDS, HALOALKANES

0377. Potassium dinitromethanide

[32617-22-4]

KCH(NO₂)₂

CHKN₂O₄

Grakauskas, V. *et al.*, *J. Org. Chem.*, 1978, **43**, 3486

Alkali-metal salts of dinitromethane are sensitive to impact and should be handled remotely behind shields, in small quantity and with great care.

See other POLYNITROALKYL COMPOUNDS

0378. Potassium dithioformate

[30962-16-4]

KSCS.H

CHKS₂



Preparative hazard

1. Martin, K., *Chem. Brit.*, 1988, **24**, 427–428

2. Engler, K., *Z. Allgem. Anorg. Chem.*, 1972, **389**, 145

3. Yoshida, 1980, 387

Potassium dithioformate is prepared by interaction of chloroform with potassium sulfide in ethanol, via a carbene reaction, possibly involving the 3 main stages



The reaction has been found unpredictable in practice, 2 out of 3 attempts leading to eruption or explosion of the flask contents [1]. However, using a modified version of a published procedure [2], with methanol as solvent, the reaction can be performed without incident, provided that the working scale is restricted to one third of that published (and initially one twelfth until one gains experience of the reaction); that a flanged reaction flask is used; and that the rate of addition of chloroform to the methanolic potassium sulfide is carefully controlled to 4–5 drops/s. This is essential, for if too slow the reaction stops, and if too fast it becomes uncontrollable [1]. Though no figures for the more reactive chloroform are given, the MRH value for a mixture of potassium sulfide with 41 wt% of carbon tetrachloride is 2.05 kJ/g [3].

0379. Lithium diazomethanide

[67880-27-7]

CHLiN₂

LiCHN₂

Müller, E. *et al.*, *Chem. Ber.*, 1954, **87**, 1887

Alkali metal salts of diazomethane are very explosive when exposed to air in the dry state, and should be handled, preferably wet with solvent, under an inert atmosphere.

See other DIAZO COMPOUNDS

†0380. Hydrogen cyanide

[74-90-8]

CHN

HC≡N

(MCA SD-67, 1961); FPA H94, 1980; HCS 1980, 547 (neat), 542 (solutions)

1. MCA SD-67, 1961
2. Wöhler, L. *et al.*, *Chem. Ztg.*, 1926, **50**, 761, 781
3. Gause, E. H. *et al.*, *J. Chem. Eng. Data*, 1960, **5**, 351
4. Salomone, G., *Gazz. Chim. Ital. [I]*, 1912, **42**, 617–622
5. Anon., *Jahresber.*, 1979, 70–71
6. Bond, J., *Loss Prev. Bull.*, 1991, **101**, 3

Hydrogen cyanide is highly endothermic and of low MW (ΔH_f° (g) +130.5 kJ/mol, 4.83 kJ/g). A comprehensive guide to all aspects of industrial handling of anhydrous hydrogen cyanide and its aqueous solutions states that the anhydrous liquid is stable at or below room temperature if it is inhibited with acid (e.g. 0.1% sulphuric acid) [1]. Presence of alkali favours explosive polymerisation [2]. In absence of inhibitor, exothermic polymerisation occurs, and if the temperature attains 184°C, explosively rapid polymerisation occurs [3]. A 100 g sample of 95–96% material stored in a glass bottle shielded from sunlight exploded after 8 weeks [4]. The explosive polymerisation of a 33 kg cylinder was attributed to lack of sufficient phosphoric acid

as stabiliser [5]. A tank containing 4 or 5 tonnes of hydrogen cyanide exploded with the force of several kg of TNT, leading to an HCN fire. This was easily extinguished, the clear-up thereafter being impeded by the toxicity of the unburnt cyanide. The explosion was attributed to build up of polymer (the reaction is autocatalytic because of ammonia generation), insufficient stabiliser (oxalic acid), lower than usual purity (<93%) and possibly to mercury contamination. It was recommended that HCN should not be stored if overwet, that tanks should be regularly emptied to inspect for polymer, and that stabiliser should be added before running to storage [6].

See Mercury(II) cyanide: Hydrogen cyanide

See other GLASS INCIDENTS, POLYMERISATION INCIDENTS

Alcohols, Hydrogen chloride

See Hydrogen chloride: Alcohols, etc.

Hypochlorites

See CYANIDES

Other reactants

Yoshida, 1980, 136

MRH values for 15 combinations, mainly with oxidants, are given

See other CYANO COMPOUNDS ENDOTHERMIC COMPOUNDS, ORGANIC ACIDS

0381. Fulminic acid (Hydrogen cyanide *N*-oxide)

[506-85-4]

CHNO



1. Sorbe, 1968, 72

2. Wentrup, C. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 467

It is fairly stable as an ethereal solution, but the isolated acid is explosively unstable, and sensitive to heat, shock or friction [1]. In a new method of preparation of the acid or its salts, pyrolysis of 4-oximato-3-substituted-isoxazol-5(4*H*)-ones or their metal salts must be conducted with extreme care under high vacuum to prevent explosive decomposition [2].

See Silver 3-methylisoxazolin-4,5-dione-4-oximate

Sodium 3-methylisoxazolin-4,5-dione-4-oximate

Sodium 3-phenylisoxazolin-4,5-dione-4-oximate

METAL FULMINATES

See other *N*-OXIDES

0382. Monosodium cyanamide

[17292-62-5]

CHN₂Na



tert-Butyl hypochlorite

See Cyanonitrene

See other CYANO COMPOUNDS, *N*-METAL DERIVATIVES

0383. Sodium diazomethanide
[67880-28-8]




See Lithium diazomethanide
See other DIAZO COMPOUNDS

0384. Sodium dinitromethanide
[25854-41-5]



Grakauskas, V. *et al.*, *J. Org. Chem.*, 1978, **43**, 3486
Alkali-metal salts of dinitromethane are sensitive to impact and should be handled remotely behind shields in small quantity, and with great care.
See other POLYNITROALKYL COMPOUNDS

0385. Trinitromethane ('Nitroform')
[517-25-9]



1. Marans, N. S. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 5329
2. Stull, 1977, 20

Explosions occurred during distillation of this polynitro compound [1]. Though not an endothermic compound, it is of positive oxygen balance and the heat of decomposition (2.80 kJ/g) would give an adiabatic decomposition temperature exceeding 2200°C and a 40-fold increase in pressure [2].

See entry OXYGEN BALANCE

Divinyl ketone

Graff, M. *et al.*, *J. Org. Chem.*, 1968, **33**, 1247

One attempted reaction of trinitromethane with impure ketone caused an explosion at refrigerator temperature.

2-Propanol

MCA Case History No. 1010

Frozen mixtures of trinitromethane –2-propanol (9:1) exploded during thawing. The former (of positive oxygen balance) dissolves exothermally in the alcohol, the heat effect increasing directly with the concentration above 50% w/w. Traces of nitric acid may also have been present.

See other POLYNITROALKYL COMPOUNDS

0386. Azidodithioformic acid
[4472-06-4]



1. Mellor, 1947, Vol. 8, 338
2. Smith, G. B. L., *Inorg. Synth.*, 1939, **1**, 81

The isolated acid or its salts are shock- and heat-sensitive explosives [1]. Safe preparative procedures have been detailed. The heavy metal salts, though powerful detonators, are too sensitive for practical use [2].


See Carbon disulfide: Metal azides

See Bis(azidothiocarbonyl) disulfide

See other ACYL AZIDES, ORGANIC ACIDS

0387. 5-Nitrotetrazole
[55011-46-6]

CHN₅O₂




1. Jenkins, J. M., *Chem. Brit.*, 1970, **6**, 401
2. Bates, L. R. *et al.*, US Pat. 4 094 879, 1978
3. Koldobskii, G. I. *et al.*, *Russ. J. Org. Chem.*, 1997, **33**(12), 1771

An acidified solution of the sodium salt was allowed to evaporate during 3 days and spontaneously exploded 2 weeks later. The nature of the explosive species, possibly the *aci*-tetrazolic acid, was being sought [1]. The silver and mercury salts are explosive [2]. The chemical properties of the free nitrotetrazole have been studied. The sodium salt tetrahydrate loses water above 50°C, greatly increasing its friction and shock sensitivity [3].

See other C-NITRO COMPOUNDS, TETRAZOLES

0388. 5-Azidotetrazole
[35038-46-1]

CHN₇



Alone, or Acetic acid, or Alkali

1. Thiele, J. *et al.*, *Ann.*, 1895, **287**, 238
2. Lieber, E. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 1313

Though explosive, it (and its ammonium salt) are much less sensitive to impact or friction than its sodium or potassium salts [1]. A small sample of the latter exploded violently during vacuum filtration. The parent compound explodes spontaneously even in acetone (but not in ethanol or aqueous) solution if traces of acetic acid are

present [2]. The salts are readily formed from diaminoguanidine salts and alkali nitrites. The ammonium salt explodes on heating, and the silver salt is violently explosive even when wet [1]. The sodium salt is also readily formed from cyanogen azide.

See Sodium 5-azidotetrazolide

See other ORGANIC AZIDES, TETRAZOLES

0389. Triazidomethane

[187585-04-6]

CHN₉

CH(N₃)₃

Hassner, A., private comm., 1986

Hassner, A. *et al.*, *J. Org. Chem.*, 1990, **55**, 2304

The azide form of the quaternary ammonium ion exchange resin IR-400 (which exhibits low friction-sensitivity) reacts with bromoform, analogously to dichloro- and dibromo-methane, to form the highly explosive triazidomethane. Solutions of above 50% concentration explode in contact with a pipette or on injection into a GLC inlet port.

See other ORGANIC AZIDES

0390. Sodium hydrogen carbonate

[144-55-8]

CHNaO₃

NaOCO.OH

HCS 1980, 828

Carbon, Water

CISHC Chem. Safety Summ., 1978, **49**, 33

A mixture was being stirred and steam heated when power failure interrupted stirring, and heating was turned off for a hour before power was restored. When stirring was restarted, the hot contents of the pan erupted immediately. Carbon dioxide is evolved from warm aqueous solutions of the base, and absence of stirring and presence of the carbon adsorbent would lead to non-equilibrium retention of the gas, which would be released instantaneously on stirring.

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

2-Furaldehyde

See 2-Furaldehyde

See other INORGANIC BASES, METAL OXONON-METALLATES

0391. Silver nitroureide

[[74386-96-2] (ion)]


CH₂AgN₃O₃

AgN(NO₂)CO.NH₂

See Nitrourea

See other N-METAL DERIVATIVES, N-NITRO COMPOUNDS, SILVER COMPOUNDS

0392. Silver 5-aminotetrazolide
[[50577-64-5] (ion)]



Thiele, J., *Ann.*, 1892, **270**, 59

Similar to silver tetrazolide, it explodes on heating.

See other N-METAL DERIVATIVES, SILVER COMPOUNDS, TETRAZOLES

0393. Cyanoborane oligomer
[60633-76-3]



Gyori, B. *et al.*, *J. Organomet. Chem.*, 1984, **262**, C7


It explodes on mechanical shock, and decomposes violently on heating under nitrogen to 230°C. The scale of preparation may need to be limited.

Sodium chlorate

See Sodium chlorate: Cyanoborane oligomer

See related BORANES, CYANO COMPOUNDS

0394. Bis(difluoroboryl)methane
[55124-14-6]



Air, or Water

Maraschin, N. J. *et al.*, *Inorg. Chem.*, 1975, **14**, 1856

Highly reactive, it explodes on exposure to air or water.

See related ALKYLHALOBORANES

0395. Dibromomethane
[74-95-3]



Potassium

See Potassium: Halocarbons

See other HALOALKANES

0396. Chloronitromethane
[1794-84-9]



Preparative hazard

1. Seigle, L. W. *et al.*, *J. Org. Chem.*, 1940, **5**, 100
2. Libman, D. D., private comm., 1968

Chlorination of nitromethane following the published general method [1] gave a product which decomposed explosively during distillation at 95 mbar [2]. A b.p. of 122°C/1 bar is quoted in the literature.

See Sodium *aci*-nitromethanide: Carbon disulfide, Chlorine

See related NITROALKANES

†0397. Dichloromethane
[75-09-2]



(MCA SD-86, 1962); NSC 474, 1979; HCS 1980, 648

1. RSC Lab. Hazard Data Sheet No. 3, 1982
2. Downey, J. R., *Chem. Eng. News*, 1983, **61**(8), 2
3. Kelling, R. A., *Chem. Eng. News*, 1990, **68**(17), 2

Previously thought to be non-flammable except at elevated temperature or pressure or in oxygen-enriched air [1], it is in fact flammable in the range 12–19% in ambient air, given a sufficiently high level of ignition energy. Though it has no measurable flash point, it is calculated that flammable regions may exist above –9°C [2]. A surprisingly violent burst of a half full separating funnel occurred when dichloromethane was shaken with water, air also being present [3].

See FLASH POINTS (reference 19)

Air, Methanol

Coffee, R. D. *et al.*, *J. Chem. Eng. Data*, 1972, **17**, 89–93

Dichloromethane, previously considered to be non-flammable except in oxygen, becomes flammable in air at 102°C/1 bar, at 27°C/1.7 bar or at 27°C/1 bar in presence of less than 0.5 vol% of methanol (but see reference 2 above). Other data are also given.

Aluminium

See Aluminium: Dichloromethane

See Metals, below

Aluminium bromide

See Aluminium bromide: Dichloromethane

Azides

See Quaternary ammonium azides, below; Sodium azide, below

1,2-Diaminoethane

1. Nolan, 1983, Case history 145
2. Heskey, W. A., *Chem. Eng. News*, 1986, **64**(21), 2
3. Laird, T., *Chem. & Ind.*, 1986, 139

Dichloromethane was being distilled from its mixture with the amine at a bath temperature of 30°C when an exothermic reaction led to deflagration [1]. Heat of reaction, $\Delta H = -343$ kJ/mol (4.04 kJ/g) of dichloromethane [2]. Reaction of amines with dichloromethane at ambient temperature is common, exothermic in conc. solutions, and involves formation of an *N*-chloromethyl quaternary salt. Concentrating dichloromethane solutions of amines to low volumes should therefore be avoided [3].

Dimethyl sulfoxide, Perchloric acid

See Perchloric acid: Dichloromethane, Dimethyl sulfoxide

Dinitrogen pentaoxide

See NITRATING AGENTS

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Halocarbons

Metals

See Aluminium: Halocarbons

Lithium: Halocarbons

Sodium: Halocarbons

METAL-HALOCARBON INCIDENTS

Nitric acid

See Nitric acid: Dichloromethane

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

Quaternary ammonium azides

1. Hassner, A. *et al.*, *Angew. Chem. (Intern. Edn.)*, 1986, **25**, 479–480
2. Demer, F. R., *Lab. Safety Notes* (Univ. of Arizona), 1991(Spring), 2
3. Wood, W., private comm., 1986
4. Bretherick, L., *Chem. Eng. News*, 1986, **64**(51), 2

Quaternary ammonium azides will displace halogens in a synthesis of alkyl azides. Dichloromethane has been used as a solvent, although this can slowly form diazidomethane which may be concentrated by distillation during work-up, thereafter easily exploding [1]. An accident attributed to this cause is described, and acetonitrile recommended as a preferable solvent, supported polymeric azides, excess of which can be removed by filtration are also preferred in place of the tetrabutylammonium salt [2]. A similar explosion was previously recorded when the quaternary azide was generated in situ from sodium azide and a phase transfer catalyst in a part aqueous system [3,4].

Sodium azide

See Quaternary ammonium azides, above

See other HALOALKANES

0398. 1,1-Difluorourea
[1510-31-2]

CH₂F₂N₂O

F₂NCO.NH₂

Parker, C. O. *et al.*, *Inorg. Synth.*, 1970, **12**, 309

Concentrated aqueous solutions of difluorourea decompose above -20°C with evolution of tetrafluorohydrazine and difluorammine, both explosive gases.

See other DIFLUOROAMINO COMPOUNDS, GAS EVOLUTION INCIDENTS

0399. Trifluoromethylphosphine
[420-52-0]

CH₂F₃P

F₃CPH₂

491M, 1975, 428

Ignites in air.

See other ALKYLPHOSPHINES

0400. Diiodomethane
[75-11-6]

CH₂I₂

I₂CH₂

Alkenes, Diethylzinc

See Diethylzinc: Alkenes, Diiodomethane

Metals

See Copper–zinc alloys: Diiodomethane, etc.

Lithium: Halocarbons

Potassium: Halocarbons

METAL–HALOCARBON INCIDENTS

See other ENDOTHERMIC COMPOUNDS, HALOALKANES

0401. Methylenedilithium
[21473-42-1]

CH₂Li₂

H₂CLi₂

491M, 1975, 234

Ignites in air.

See other ALKYLMETALS

0402. Methylenemagnesium
[25382-52-9]



Ziegler, K. *et al.*, *Z. Anorg. Chem.*, 1955, **282**, 345
The polymeric form ignites in air.
See other ALKYL METALS

0403. Sodium *aci*-nitromethanide
[25854-38-0]



Meyer, V., *et al.*, *Ber.*, 1894, **27**, 1601; 3407
Sodium nitromethanoate is relatively stable when solvated. The dry salt is a sensitive and powerful explosive which may be detonated by warming to 100°C, by a strong blow or contact with traces of water. The potassium salt is even more sensitive. Lecture demonstrations of these properties are described.
See Nitromethane: Sodium hydride

Carbon disulfide, Chlorine

1. Stirling, C. J. M., *Chem. Brit.*, 1986, **22**, 524
2. Stirling, C. J. M., private comm., 1986

During the preparation of chloronitromethane by adding portions of dry sodium *aci*-nitromethanide to chlorine (40 mol of each) dissolved in carbon disulfide, a violent explosion occurred when the addition was half-complete. Similar reactions using bromine had been executed uneventfully many times previously [1]. No certain explanation has emerged, but the sodium salt is known to be explosively unstable, and mixtures of carbon disulfide vapour and air are of course extremely flammable and explosive. Contact of the dry salt with traces of chlorine above its carbon disulfide solution may have led to an exotherm and ignition of the vapour-air mixture in the flask [2].

Mercury(II) chloride, Acids

Nef, J. U., *Ann.*, 1894, **280**, 263, 305
Interaction gives mercury nitromethanide, which is converted by acids to mercury fulminate.
See Mercury(II) fulminate

Nitric oxide, Base

Traube, W., *Annalen*, 1898, **300**, 107
A slurry of sodium nitromethanoate in excess sodium ethoxide solution forms a (presumably even more) explosive salt on treatment with nitric oxide. It is claimed the parent acid is $\text{O}_2\text{NCH}_2\text{N}_2\text{O}_2\text{H}$.
See Nitrogen oxide

1,1,3,3-Tetramethyl-2,4-cyclobutanedione

See Sodium 1,3-dihydroxy-1,3-bis(*aci*-nitromethyl)-2,2,4,4-tetramethylcyclobutanediide

Water

1. Nef, J. U., *Ann.*, 1894, **280**, 273

2. Jung, M. E. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1987, 753 (footnote)

The *aci*-sodium salt, normally crystallising with one molecule of ethanol and stable, will explode if moistened with water. This is due to liberation of heat and conversion to sodium fulminate [1]. After several uneventful similar operations, during the destruction of excess sodium salt by pouring water onto it, a violent explosion occurred [2]. A safer procedure would be to add the salt in small portions to a bulk of stirred ice-water.

See also Sodium 1,3-dihydroxy-1,3-bis(*aci*-nitromethyl)-2,2,4,4-tetramethylcyclobutandiide

See other aci-NITRO SALTS

0404. Cyanamide

[420-04-2]

CH₂N₂



1. Anon., *Fire Prot. Assoc. J.*, 1966, 243

2. Anon., *Sichere Chemiearbeit*, 1976, **28**, 63

Cyanamide is endothermic ($\Delta H_f^\circ +58.8$ kJ/mol, 1.40 kJ/g), thermally unstable and needs storage under controlled conditions. Contact with moisture, acids or alkalis accelerates the rate of decomposition, and at temperatures above 40°C thermal decomposition is rapid and may become violent. A maximum storage temperature of 27°C is recommended [1]. Commercial cyanamide is stabilised with boric acid, phosphoric acid, sodium dihydrogen phosphate etc., but vacuum distillation produces a neutral (unstabilised) distillate, which immediately may decompose spontaneously. Small-scale storage tests showed that unstabilised cyanamide was 47% decomposed after 18 days at 20°C and 75% decomposed after 29 days at 30°C, whereas stabilised material showed only 1% decomposition under each of these conditions. Larger-scale tests with 1–2 kg unstabilised samples led to sudden and violent exothermic polymerisation after storage for 14 days at ambient temperature. If small samples of unstabilised cyanamide are required, they are best prepared by freezing out from aqueous solutions of the stabilised material. Such small samples should be used immediately or stored under refrigeration [2].

1,2-Phenylenediamine salts

Sawatari, K. *et al.*, Japan Kokai, 76 16 669, 1976 (*Chem. Abs.*, 1976, **85**, 63069)

During the preparation of 2-aminobenzimidazoles, reaction conditions are maintained below 90°C to prevent explosive polymerisation of cyanamide.

Water

Pinck, L. A. *et al.*, *Inorg. Synth.*, 1950, **3**, 41

Evaporation of aqueous solutions to dryness is hazardous, owing to the possibility of explosive polymerisation in conc. solution.

See other POLYMERISATION INCIDENTS

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0405. Diazirine
[157-22-2]




1. Graham, W. H., *J. Org. Chem.*, 1965, **30**, 2108
2. Schmitz, E. *et al.*, *Chem. Ber.*, 1962, **95**, 800

This cyclic isomer of diazomethane is also a gas (b.p., -14°C) which explodes on heating. Several homologues are also thermally unstable [1,2].

See other DIAZIRINES

0406. Diazomethane
[334-88-3]



RSC Lab. Hazard Data Sheet No. 4, 1982

1. Eistert, B., in *Newer Methods of Preparative Organic Chemistry*, 517–518, New York, Interscience, 1948
2. de Boer, H. J. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 250
3. Gutsche, C. D., *Org. React.*, 1954, **8**, 392–393
4. Zollinger, H., *Azo and Diazo Chemistry*, 22, London, Interscience, 1961
5. Fieser, L. *et al.*, *Reagents for Organic Synthesis*, Vol. 1, 191, New York, Wiley, 1967
6. Horák, V. *et al.*, *Chem. & Ind.*, 1961, 472
7. Ruehle, P. H. *et al.*, *Chem. & Ind.*, 1979, 255–256
8. Bowes, C. M., *Univ. Safety Assoc. Safety News*, 1981, **15**, 18–19
9. Hashimoto, N. *et al.*, *Chem. Pharm. Bull.*, 1981, **29**, 1475–1478; 1982, **30**, 119–124
10. Barnes, C. J. *et al.*, *J. Assoc. Off. Anal. Chem.*, 1982, **65**, 273–274
11. Black, H. T., *Aldrichimica Acta*, 1983, **16**(1), 3–10
12. Asyama, T. *et al.*, *Chem. Abs.*, 1984, **101**, 171306
13. Shiga, S., *Chem. Abs.*, 1987, **107**, 117404
14. Carlton, L. *et al.*, *S. African J. Chem.*, 1987, **40**, 203–204
15. Archibald, T. G. *et al.*, US Patent 5,817,778, 1998

Diazomethane is a highly endothermic small molecule ($\Delta H_f^{\circ}(\text{g}) +192.5$ kJ/mol, 4.58 kJ/g) which boils at -23°C and the undiluted liquid or concentrated solutions may explode if impurities or solids are present [1], including freshly crystallised products [2]. Gaseous diazomethane, even when diluted with nitrogen, may explode at elevated temperatures (100°C or above), or under high-intensity lighting, or if rough surfaces are present [1]. Ground glass apparatus or glass-sleeved stirrers are therefore undesirable when working with diazomethane. Explosive intermediates may also be formed during its use as a reagent, but cold dilute solutions have frequently been used uneventfully [1]. Further safety precautions have been detailed

[2,3,4]. Many precursors for diazomethane generation are available [5], including the stable water soluble intermediate *N*-nitroso-3-methylaminosulfolane [6]. Many of the explosions observed are attributed to uncontrolled or unsuitable conditions of contact between concentrated alkali and undiluted nitroso precursors [1].

A large scale generator and procedure for safe preparation and use of the reagent on 1.1 mol scale are detailed [7]. Insolubility of the substrate in ether led to omission of ether from a microgenerator chamber and use of dioxan to dissolve the substrate in the outer chamber. During addition of alkali to nitrosomethylguanidine (30 mg) in 0.125 ml water, the apparatus exploded. This was attributed to absence of ether vapour to dilute the diazomethane [8]. Trimethylsilyldiazomethane is presented as a safe and effective substitute for the hazardous parent compound [9], and two cheap and safe storage containers for small amounts of ethereal solutions of the latter are described [10]. The available data has recently been summarised with 116 references [11]. The preparation of trimethylsilyldiazomethane and its application as a safe substitute for diazomethane are reviewed [12]. Automated equipment for esterification of small samples of acidic materials for GLC analysis has been developed [13]. A convenient method for preparation of small amounts of pure diazomethane involves entrainment in a stream of helium and condensation as a solid, m.p. -145°C , on a liquid nitrogen-cooled cold finger. The solid may be kept almost indefinitely at LN temperature [14]. Procedures for small industrial scale (50 mole) preparation and use in solution have been patented [15].

See also GLASS INCIDENTS

Alkali metals

Eistert, B. in *Newer Methods of Preparative Organic Chemistry*, 518, New York, Interscience, 1948

Contact of diazomethane with alkali metals causes explosions.

See Lithium diazomethanide

Calcium sulfate

Gutsche, C. D., *Org. React.*, 1954, **8**, 392

Calcium sulfate is an unsuitable desiccant for drying tubes in diazomethane systems. Contact of diazomethane vapour and the sulfate causes an exotherm which may lead to detonation. Potassium hydroxide is a suitable desiccant.

Dimethylaminodimethylarsine, Trimethyltin chloride

Krommes, P. *et al.*, *J. Organomet. Chem.*, 1976, **110**, 195–200

Interaction in ether to produce diazomethyldimethylarsine is accompanied by violent foaming, and eye protection is essential.

See other DIAZO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0407. Isocyanoamide ('Isodiazomethane')

[4702-38-9]

CH_2N_2


$:\text{C}=\text{NNH}_2$

1. Müller, E. *et al.*, *Chem. Ber.*, 1954, **87**, 1887
2. Müller, E. *et al.*, *Ann.*, 1968, **713**, 87

This unstable liquid begins to decompose at 15°C and explodes exothermically at 35–40°C [1], but may be handled safely in ether solution [2].

See related CYANO COMPOUNDS

0408. N-Nitromethanimine
[67400-85-2]




Zhao, X. *et al.*, *J. Chem. Phys.*, 1988, **88**, 801

This nitrimine, the nominal monomer of the cyclic nitramine high explosives RDX and HMX, may be involved in their detonation and can be formed from them by pyrolysis.

See 1,3,5-Trinitrohexahydro-1,3,5-triazine

See other N-NITRO COMPOUNDS

0409. Nitrooximinomethane ('Methylnitrolic acid')
[625-49-0]




Sorbe, 1968, 147

An unstable and explosive crystalline solid, formally a nitro-oxime.

See related NITROALKANES, OXIMES

0410. Dinitromethane
[625-76-3]




1. Sorbe, 1968, 148

2. Bedford, C. D. *et al.*, *J. Org. Chem.*, 1979, **44**, 635

It explodes at 100°C [1], and attempted distillation of more than 1 g at 30–35°C/1.5 mbar led to a violent explosion [2].

See other POLYNITROALKYL COMPOUNDS

0411. Tetrazole
[288-94-8]



1. Benson, F. R., *Chem. Rev.*, 1947, **41**, 5

2. Stull, 1977, 22

3. *DOC* 5, 1982, 5032

4. Anon., *IST Sci. Tech.*, 1987, (4), 9

It explodes above its m.p., 155°C [1]. It is highly endothermic ($\Delta H_f^\circ +237.2$ kJ/mol, 3.39 kJ/g) with a heat of decomposition (3.27 kJ/g) which would give an adiabatic product temperature of some 1950°C and a 22-fold increase in pressure in a closed container [2]. Its solutions are also explosive when shocked [3]. An explosion during sublimation of tetrazole at ambient pressure was caused by overheating [4].

See other ENDOTHERMIC COMPOUNDS, TETRAZOLES


0412. Lead methylenebis(nitramide) (*N,N'*-Dinitromethanediamine, lead(II) salt)
[86202-43-9] **CH₂N₄O₄Pb**



See Methylenebis(nitramine)

See other HEAVY METAL DERIVATIVES, *N*-NITRO COMPOUNDS

0413. 5-Amino-1,2,3,4-thiazotriazole
[6630-99-5]




Lieber, E. *et al.*, *Inorg. Synth.*, 1960, **6**, 44

It decomposes with a slight explosion in a capillary tube at 136°C.

See other HIGH-NITROGEN COMPOUNDS, N-S COMPOUNDS

0414. Diazidomethane
[107585-03-5]



1. Hassner, A. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1986, **25**, 479–480


2. Hassner, A. *et al.*, *J. Org. Chem.*, 1990, **55**, 2304

The azide form of the quaternary ammonium ion exchange resin IR-400 (which exhibits low friction-sensitivity) reacts very slowly with dichloromethane or dibromomethane to produce the explosive diazidomethane [1]. Solutions of above 70% concentration will explode if a pipette is inserted or on injection into a GLC inlet port [2].

See other ORGANIC AZIDES

0415. 5-N-Nitroaminotetrazole
[18558-16-4]

CH₂N₆O₂



1. Lieber, E. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 2328
 2. O'Connor, T. E. *et al.*, *J. Soc. Chem. Ind.*, 1949, **68**, 309
- It and its monopotassium salt explode at 140°C and the diammonium salt explodes at 220°C after melting [1]. The disodium salt explodes at 207°C [2].

Amminemetals

Complexes with several ammine derivatives of metals are explosive.

See entry NITRAMINE–METAL COMPLEXES

See other N-NITRO COMPOUNDS, TETRAZOLES

†0416. Formaldehyde
[50-00-0]

CH₂O

O:CH₂

(*MCA SD-1*, 1960); *NSC 342*, 1982; *FPA H54*, 1977; *HCS 1980*, 506

van den Brink, M. J., *Chem. Mag.* (Rijswijk), 1982, 428

Pure formaldehyde, prepared by vacuum depolymerisation of paraldehyde, was collected as a solid at –189°C. When the flask was transferred to a Cardice–ethanol bath, the contents began to repolymerise exothermally and ignited.

See other POLYMERISATION INCIDENTS

See Glyoxal

Acrylonitrile

See Acrylonitrile, Formaldehyde

Hydrogen peroxide

MRH 6.44/69

See Hydrogen peroxide: Oxygenated compounds

Magnesium carbonate hydroxide

BCISC Quart. Safety Summ., 1965, **36**(143), 44

During neutralisation of the formic acid present in formaldehyde solution by shaking with the basic carbonate in a screw-capped bottle, the latter burst owing to pressure of liberated carbon dioxide. Periodical release of pressure should avoid this.

See other GAS EVOLUTION INCIDENTS, NEUTRALISATION INCIDENTS

Nitromethane

See Nitromethane: Formaldehyde

Other reactants

Yoshida, 1980, 346

MRH values for 14 combinations with oxidants are given, and are all high.

Peroxyformic acid

MRH 5.69/100

See Peroxyformic acid: Organic materials

Phenol

1. Taylor, H. D. *et al.*, *Major Loss Prevention in Process Industries*, (Symp. Ser. No. 34), 46, London, IChE, 1971
2. Anon., *Chem. Eng. News*, 1992, **70**(10), 7
3. Anon. *Chem. Eng. News*, 1997, **75**(38), 11
4. Starkie, A. *et al.*, *Chem. in Brit.*, 1996, **32**(2), 35
5. Gustin, J.-L. *et al.*, *J. Loss Prev.*, 1993, **6**(2), 103

At least 9 cases of catalysed plant-scale preparations of phenol–formaldehyde resin which ran away with sudden pressure development and failure of bursting disks or reactors are briefly mentioned [1]. No details of process conditions are given. A destructive incident of this type, which must have developed the force of some kg of TNT, is reported without real detail [2]. Another fatal and destructive explosion when preparing phenol formaldehyde resins is reported [3]. The editor calculates from analogues that this reaction could have above 1 kJ/g reagents, depending upon stoichiometry and concentration, and mostly associated with the second step, dehydrative condensation of the initial hydroxymethylphenol to dihydroxybiphenyls. A photograph of the aftermath of a runaway polymerisation is reproduced [4]. A very thorough study of hazards of the phenol/formaldehyde polymerisation reaction, leading to calculations of emergency vent size, is given [5].

See other POLYMERISATION INCIDENTS, RUNAWAY REACTIONS

Potassium permanganate

See Potassium permanganate: Formaldehyde

Sodium hydroxide

1. Ashby, E. C. *et al.*, *J. Amer. Chem. Soc.*, 1993, **115**, 1171
2. Chrisope, D. R. *et al.*, *Chem. Eng. News.*, 1995, **73**(2), 2
3. Kapoor, S. *et al.*, *J. Phys. Chem.*, 1995, **99**(18), 6857

In place of the well known Cannizzaro reaction (which is significantly exothermic) formaldehyde, at lower concentrations, produces hydrogen with alkalis, leading to possible pressurisation and ignition [1]. It has been demonstrated that this modified Cannizzaro reaction giving hydrogen can operate in real commercial situations (500 ppm formaldehyde, 1.5% sodium hydroxide). Caution and ventilated or inerted headspaces when storing alkaline formaldehyde containing products are advisable [2]. A detailed study of the kinetics and mechanism of hydrogen evolution has been published [3].

Water, Methanol, Air

Anon., *Jahresbericht*, 1994, 73

An explosion is recorded, consequent upon welding a ladder to the wall of a half-full tank, which had not been inerted with nitrogen and was vented to atmosphere, containing a 37% formaldehyde solution stabilised with 5.6% methanol at 72°C. The flash point of this mixture in air is about 71°C, autoignition temperature 420°C: the head space ignited, killing two workers, injuring nine and blowing off the top of the tank.

See other ALDEHYDES, REDUCANTS

0417. Paraformaldehyde

[9002-81-7]

(CH₂O)_n



The dry finely powdered linear polymer is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

See other ALDEHYDES, REDUCANTS

0418. Formic acid (Methanoic acid)

[64-18-6]

CH₂O₂



FPA H86, 1979; *HCS 1980*, 508

1. *BCISC Quart. Safety Summ.*, 1973, **44**, 18
2. Falconer, J. *et al.*, *Proc. 2nd Int. Conf. Solid. Surf.*, 1974, 525
3. Anon., *Jahresber.*, 1981, 74
4. Reynolds, R. J., *DNA Repair*, 1981, **1**(A), 11–21, NY, Dekker

The slow decomposition in storage of 98–100% formic acid with liberation of carbon monoxide led to rupture of the sealed glass containers. In absence of gas leakage, a full 2.5 l bottle would develop a pressure of over 7 bar during 1 year at 25°C [1]. Explosive decomposition of formic acid on a clean nickel (1.1.0) surface was studied, using deuterioformic acid [2]. A full 1 l bottle of 96% formic acid burst when the ambient temperature fell to –6°C overnight and the contents froze and expanded. Gas pressure from previous partial decomposition may also have contributed [3]. Hydrolysis of precipitated DNA with 97% acid at 180°C in sealed glass tubes is potentially hazardous because of rapid gas formation [4].

See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS

Aluminium

See Aluminium: Formic acid

Hydrogen peroxide

MRH 3.30/42

See Peroxyformic acid (references 5,6)

Hydrogen peroxide

See Hydrogen peroxide: Oxygenated compounds

4-Hydroxy-3-methoxybenzaldehyde, Thallium(III) nitrate

See Thallium(III) nitrate: Formic acid, etc.

Nitric acid

MRH 2.64/36

See Nitric acid: Formic acid

Nitric acid, Urea

See Nitric acid: Formic acid, Urea

Nitromethane

MRH 5.69/100

See Nitromethane: Acids

Other reactants

Yoshida, 1980, 100

MRH values calculated for 16 combinations, largely with oxidants, are given.

Palladium–carbon catalyst

Freifelder, 1971, 188

Addition of dry catalyst to 98% formic acid used as a hydrogenation solvent can be extremely hazardous, because hydrogen is released by decomposition of the acid. Addition of acid to the water-wetted catalyst is safer.

See other GAS EVOLUTION INCIDENTS, HYDROGENATION INCIDENTS

Phosphorus pentaoxide

Muir, G. D., private comm., 1968

Attempted dehydration of 95% acid to anhydrous formic acid caused rapid evolution of carbon monoxide.

See other GAS EVOLUTION INCIDENTS

Sodium hypochlorite


See Sodium hypochlorite: Formic acid

See other ORGANIC ACIDS, REDUCANTS

0419. Formaldehyde oxide polymer (Poly(dioxymethylene))

[107981-97-5]

$(\text{CH}_2\text{O}_2)_n$



Lapalme, R. *et al.*, *Can. J. Chem.*, 1979, **57**, 3272

The dry peroxide polymer (formed in ozonisation of vinyl acetate, and possibly a dioxirane polymer) is a shock-sensitive explosive, which needs care in handling.

See other POLYPEROXIDES

0420. Peroxyformic acid (Methaneperoxoic acid)

[107-32-4]

CH_2O_3



1. Greenspan, F. P., *J. Amer. Chem. Soc.*, 1946, **68**, 907

2. D'Ans, J. *et al.*, *Ber.*, 1915, **48**, 1136

3. Weingartshofer, A. *et al.*, *Chem. Eng. News*, 1952, **30**, 3041

4. Swern, 1970, Vol. 1, 337
5. Isard, A. *et al.*, *Chemical Tech.*, 1974, **4**, 380
6. Anon., *Chem. Eng. News*, 1950, **28**, 418
7. Stull, 1977, 19

Peroxyformic acid solutions are unstable and undergo a self-accelerating exothermic decomposition at ambient temperature [1]. An 80% solution exploded at 80–85°C [2]. A small sample of the pure vacuum-distilled material cooled to below –10°C exploded when the flask was moved [3]. Though the acid has occasionally been distilled, this is an extremely dangerous operation (it is formally the redox compound formyl hydroperoxide) [4]. During preparation of the acid by a patented procedure involving interaction of formic acid with hydrogen peroxide in the presence of meta-boric acid, an explosion occurred which was attributed to spontaneous separation of virtually pure peroxyformic acid [5]. Following an incident in which a 1:1 mixture of formic acid and 90% hydrogen peroxide had exploded violently when handled after use as an oxidant [6], it was calculated that the (redox) decomposition exotherm (1.83 kJ/g of acid) would attain an adiabatic reaction temperature of 1800°C with a 22-fold increase in pressure in a closed container [7].

Chlorine

See Chlorine: Chloromethane

Metals, or Metal oxides

D'Ans, J. *et al.*, *Ber.*, 1915, **48**, 1136

Violence of reaction depends on concentration of acid and scale and proportion of reactants. The following observations were made with additions to 2–3 drops of ca. 90% acid. Nickel powder, becomes violent; mercury, colloidal silver and thallium powder readily cause explosions; zinc powder causes a violent explosion immediately. Iron powder is ineffective alone, but a trace of manganese dioxide promotes deflagration. Barium peroxide, copper(I) oxide, impure chromium trioxide, iridium dioxide, lead dioxide, manganese dioxide and vanadium pentoxide all cause violent decomposition, sometimes accelerating to explosion. Lead(II) oxide, lead(II),(IV) oxide and sodium peroxide all cause an immediate violent explosion.

Non-metals

D'Ans, J. *et al.*, *Ber.*, 1915, **48**, 1136

Impure carbon and red phosphorus are oxidised violently, and silicon, promoted by traces of manganese dioxide, is oxidised with ignition.

Organic materials

1. D'Ans, J. *et al.*, *Ber.*, 1915, **48**, 1136
2. Anon., *Chem. Eng. News*, 1950, **28**, 418
3. Shanley, E. S., *Chem. Eng. News*, 1950, **28**, 3067
4. Öhlenschlager, A. *et al.*, *Chem. Eng. News*, 1987, **65**(15), 2

Formaldehyde, benzaldehyde and aniline react violently with 90% performic acid [1]. An unspecified organic compound was added to the acid (preformed from formic acid and 90% hydrogen peroxide), and soon after the original vigorous reaction had subsided, the mixture exploded violently [2]: (see references 6,7 above). Reaction

with alkenes is vigorously exothermic, and adequate cooling is necessary [3]. Reactions with performic acid can be more safely accomplished by slow addition of hydrogen peroxide to a solution of the compound in formic acid, so that the peroxy acid is used as it is formed. Adequate safety screens should be used with all peracid preparations [3]. After the hydroxylation of *trans*-2-pentanoic acid with performic acid according to a published general procedure, while formic acid and water were being distilled off under vacuum at 45°C, a violent explosion occurred. This was attributed to decomposition of unreacted performic acid [4].

Sodium nitrate


D'Ans, J. *et al.*, *Ber.*, 1915, **48**, 1139

The salt may lead to explosive decomposition of the peroxy acid.

See other PEROXYACIDS, REDOX COMPOUNDS

0421. Methylsilver

[75993-65-6]



Thiele, H., *Z. Elektrochem.*, 1943, **49**, 426

Prepared at -80°C , the addition compound with silver nitrate decomposes explosively on warming to -20°C .

See other ALKYL METALS

0422. Silver nitroguanidide

[]



See Nitroguanidine

See other SILVER COMPOUNDS

0423. Methylaluminium diiodide

[2938-46-7]



Nitromethane

See Nitromethane: Alkylmetal halides

See other ALKYLALUMINIUM HALIDES

0424. Dichloromethylarsine

[593-89-5]



Chlorine

See Chlorine: Dichloro(methyl)arsine

See other ALKYLNON-METAL HALIDES

0425. Methylborylene
[62785-41-5]

CH₃B

MeB:

Preparative hazard

See Dibromomethylborane: Sodium–potassium alloy (next below)

See related ALKYLBORANES

0426. Dibromomethylborane
[17933-16-2]

CH₃BBr₂

MeBBr₂

Nöth, H. *et al.*, *Inorg. Synth.*, 1983, **22**, 223

It may ignite in air if warm, or if the heat of oxidation or hydrolysis cannot be dissipated.

See other SELF-HEATING AND IGNITION INCIDENTS

Sodium–potassium alloy

Van der Kerk, S. M. *et al.*, *Polyhedron*, 1983, **2**, 1337

Use of potassium graphite or caesium graphite to generate methylborylene is uneventful, while use of sodium–potassium alloy (1:5 mol) caused an explosion in 2 out of 5 attempts.

See other ALKYLHALOBORANES

0427. Methaneboronic anhydride–pyridine complex
[79723-21-0]

CH₃BO.C₅H₅N

H₃CB=O.C₅H₅N

1. Mattesson, D. S. *et al.*, *J. Org. Chem.*, 1964, **29**, 3399

2. Mattesson, D. S. *et al.*, *Organometallics*, 1982, **1**, 27

The published procedure [1] for preparing the complex failed to note that the organic extracts will ignite spontaneously if exposed to air for a few seconds during workup. Inert blanketing with argon is essential throughout, and the preparation is hazardous on the scale described. Ignition probably arises from by-product trimethylborane [2].

See other AUTOIGNITION INCIDENTS

See related ACID ANHYDRIDES

0428. Methylbismuth oxide (Methyloxobismuthine)
[]

CH₃BiO

H₃CBi=O

Marquardt, A., *Ber.*, 1887, **20**, 1522

It ignites on warming in air.

See related ALKYLMETALS

†0429. Bromomethane (Methyl bromide)

[74-83-9]

CH₃Br

H₃CBr

(MCA SD-35, 1968); HCS 1980, 641

1. MCA SD-35, 1968
2. MCA Case History No. 746

Though bromomethane is used as a fire extinguishant, it does in fact form difficultly flammable mixtures between 10 and 15% (13.5–14.5% but also 8.6–20% have been noted), the limits in oxygen, or under pressure, being wider [1,2].

See entry FLASH POINTS (reference 19)

Aluminium

Lambert, P. G., *Chem. and Ind.*, 1990, (18), 562

Specially bottled methyl bromide, in an aluminium cylinder under nitrogen pressure, disgorged a black sludge when liquid was discharged. Inversion and venting, to blow free the line, gave a burst of flame from the vent. It is presumed that corrosion of the cylinder produced pyrophoric aluminium alkyls.

See Trimethylaluminium; Aluminium: halocarbons

See other CORROSION INCIDENTS, METAL–HALOCARBON INCIDENTS

Dimethyl sulfoxide

See Trimethylsulfoxonium bromide

Ethylene oxide

See Ethylene oxide: Air, Bromomethane

Metals

MCA Case History No. 746 and addendum

Metallic components of zinc, aluminium and magnesium (or their alloys) are unsuitable for service with bromomethane because of the formation of pyrophoric Grignard-type compounds. The Case History attributes a severe explosion to ignition of a bromomethane–air mixture by pyrophoric methylaluminium bromides produced by corrosion of an aluminium component.

See other CORROSION INCIDENTS, METAL–HALOCARBON INCIDENTS

See other HALOALKANES

0430. N,N-Dibromomethylamine

[10218-83-4]

CH₃Br₂N

Br₂NCH₃

Cooper, J. C. *et al.*, *Explosivstoffe*, 1969, 17(6), 129–130

Like the dichloro analogue, it appears to be more sensitive to impact or shock than N-chloromethylamine.

See other N-HALOGEN COMPOUNDS

0431. Methylcadmium azide
[7568-37-8]



Dehnicke, K. *et al.*, *J. Organomet. Chem.*, 1966, **6**, 298

Surprisingly it is thermally stable to 300°C, (cadmium azide is very heat sensitive) but is hygroscopic and very readily hydrolysed to explosive hydrogen azide.

See related METAL AZIDES

†**0432. Chloromethane (Methyl chloride)**
[74-87-3]



(*MCA SD-40*, 1970); *FPA H68*, 1979; *HCS 1980*, 644

Aluminium

See Aluminium: Halocarbons

Aluminium chloride, Ethylene

See Ethylene: Aluminium chloride

Interhalogens

See Bromine trifluoride: Halogens, etc.

Bromine pentafluoride: Hydrogen-containing materials

Metals

MCA SD-40, 1970

In presence of catalytic amounts of aluminium chloride, powdered aluminium and chloromethane interact to form pyrophoric trimethylaluminium. Chloromethane may react explosively with magnesium, or potassium, sodium or their alloys. Zinc probably reacts similarly to magnesium.

See Aluminium: Halocarbons

Sodium: Halocarbons

METAL-HALOCARBON INCIDENTS

See other CATALYTIC IMPURITY INCIDENTS

Other reactants

Yoshida, 1980, 366

MRH values calculated for 13 combinations with oxidants are given.

See other HALOALKANES

0433. Methylmercury perchlorate
[40661-97-0]



Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 214

On rubbing with a glass rod, a sample exploded violently. As the explosion could not be reproduced with a metal rod, initiation by static electricity was suspected.

See other STATIC INITIATION INCIDENTS

See related METAL PERCHLORATES

0434. Methyl hypochlorite

[593-78-2]

CH₃ClO

MeOCl

Sandmeyer, T., *Ber.*, 1886, **19**, 859

The liquid could be gently distilled (12°C) but the superheated vapour readily and violently explodes, as does the liquid on ignition.

See other HYPOHALITES

0435. Methanesulfinyl chloride

[676-85-7]

CH₃ClOS

MeS(O)Cl

Douglass, I. B. *et al.*, *J. Org. Chem.*, 1964, **29**, 951

A sealed ampoule burst after shelf storage for several months. Store only under refrigeration.

See other ACYL HALIDES

0436. Methyl perchlorate

[17043-56-0]

CH₃ClO₄

MeOClO₃

Sidgwick, 1950, 1236

The high explosive instability is due in part to the covalent character of the alkyl perchlorates, and also to the excess of oxygen in the molecule over that required to combust completely the other elements present (i.e. a positive oxygen balance).

See other ALKYL PERCHLORATES

0437. N,N-Dichloromethylamine

[7651-91-4]

CH₃Cl₂N

Cl₂NMe

Calcium hypochlorite, or Sodium sulfide, or Water

1. Bamberger, E. *et al.*, *Ber.*, 1895, **28**, 1683

2. Okon, K. *et al.*, *Chem. Abs.*, 1960, **54**, 17887

A mixture with water exploded violently on warming [1]. Contact with solid sodium sulfide or distillation over calcium hypochlorite also caused explosions [2].

See other N-HALOGEN COMPOUNDS

0438. Dichloromethylphosphine
[676-83-5]



Bis(2-hydroxyethyl)methylphosphine

Piccinni-Leopardi, C. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1986, 91

Cyclo-condensation of the 2 components gives 2,6-dimethyl-1,3-dioxo-2,6-diphosphacyclooctane, and its subsequent distillation (65–68°C/0.07 mbar) must be effected with bath temperatures below 120°C to avoid explosion.


See other ALKYLHALOPHOSPHINES

†**0439. Methyltrichlorosilane**
[75-79-6]



See other ALKYLHALOSILANES

0440. Methylcopper
[1184-53-8]



1. Coates, 1960, 348

2. Ikariya, T. *et al.*, *J. Organomet. Chem.*, 1974, **72**, 146

The dry solid is very impact-sensitive and may explode spontaneously on being allowed to dry out at room temperature [1]. Methylcopper decomposes explosively at ambient temperature, and violently in presence of a little air [2].


See other ALKYLMETALS

†**0441. Fluoromethane (Methyl fluoride)**
[593-53-3]



See other HALOALKANES

0442. Methyl hypofluorite (Fluoroxymethane)
[36336-08-0]



Appelman, E. *et al.*, *J. Amer. Chem. Soc.*, 1991, **113**(7), 2648

This compound, the first known alkyl hypofluorite, was found to be stable but reactive at –120°C, but samples sometimes exploded on warming to cardice temperatures (–80°C?).

0443. Xenon(II) fluoride methanesulfonate

[]



Wechsberg, M. *et al.*, *Inorg. Chem.*, 1972, **11**, 3066

The solid explodes on warming from 0°C to ambient temperature.

See other XENON COMPOUNDS

0444. Silyl trifluoromethanesulfonate

[2923-28-6]



Bassindale, A. R. *et al.*, *J. Organomet. Chem.*, 1984, **271**, C2

It cannot be isolated pure by distillation at ambient pressure as it disproportionates to spontaneously flammable silane.

See related SULFUR ESTERS

0445. Iodomethane (Methyl iodide)

[74-88-4]



HCS 1980, 562

Oxygen

Antonik, S., *Bull. Soc. Chim. Fr.*, 1982, **3-4**, 128-133

Second stage ignition during oxidation/combustion of iodomethane in oxygen at 300-500°C was particularly violent, occasionally causing fracture of the apparatus, and was attributed to formation and decomposition of a periodic species.

Silver chlorite

See Silver chlorite: Iodoalkanes

Sodium

1. Anon., *J. Chem. Educ.*, 1966, **43**, A236

2. Braidech, M. M., *J. Chem. Educ.*, 1967, **44**, A324

The first stage of a reaction involved the addition of sodium dispersed in toluene to a solution of adipic ester in toluene. The subsequent addition of iodomethane (b.p. 42°C) was too fast and vigorous boiling ejected some of the flask contents. Exposure of sodium particles to air caused ignition, and a violent toluene-air explosion followed [1]. When a reagent as volatile and reactive as iodomethane is added to a hot reaction mixture, controlled addition, and one or more wide-bore reflux condensers are essential. A similar incident involving benzene was also reported [2].

See Sodium: Halocarbons

Trialkylphosphines

Houben-Weyl, 1963, Vol. 12.1, 79

In the absence of a solvent, quaternation of trialkylphosphines with methyl iodide may proceed explosively.

See other HALOALKANES

0446. Methylmagnesium iodide (Iodomethylmagnesium)

[917-64-6]



Thiophosphoryl chloride

See Tetramethyldiphosphane disulfide

Vanadium trichloride

See Vanadium trichloride: Methylmagnesium iodide

See other ALKYL METAL HALIDES, GRIGNARD REAGENTS

0447. Methylzinc iodide (Iodomethylzinc)

[18815-73-1]



Nitromethane

See Nitromethane: Alkylmetal halides

See other ALKYL METAL HALIDES

0448. Methylpotassium

[17814-73-2]




Weiss, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1968, **7**, 133

The dry material is highly pyrophoric.

See other ALKYL METALS

0449. Potassium methanediazoate (Potassium methyldiazenoxide)

[19416-93-4]



Water

Hantzsch, A. *et al.*, *Ber.*, 1902, **35**, 901

Interaction with water is explosively violent.

See related ARENEDIAZOATES

See other N-O COMPOUNDS

0450. Potassium methoxide
[865-33-8]



It may ignite in moist air.
See entry METAL ALKOXIDES

Arsenic pentafluoride, Benzene
See Arsenic pentafluoride: Benzene, etc.

0451. Potassium methylselenide
[54196-34-8]



2-Nitroacetophenone
See 2-Nitroacetophenone: Potassium methylselenide
See related METAL ALKOXIDES

0452. Methyllithium
[917-54-4]



Sidgwick, 1950, 71
Ignites and burns brilliantly in air. The commercial solution in diethyl ether is also pyrophoric.
See ALKALI-METAL DERIVATIVES OF HYDROCARBONS
See other ALKYL METALS

0453. Formamide
[75-12-7]



HCS 1980, 507; *RSC Lab. Hazards Data Sheet No. 58*, 1987

Aluminium, Ammonium nitrate, Calcium nitrate
See Calcium nitrate: Aluminium, etc.

Di-*tert*-butyl hyponitrite
See Di-*tert*-butyl hyponitrite: Formamides

Iodine, Pyridine, Sulfur trioxide
Anon., *J. Chem. Educ.*, 1973, **50**, A293

Bottles containing a modified Karl Fischer reagent with formamide replacing methanol developed gas pressure during several months and burst. No reason was apparent, but slow formation of sulfuric acid, either by absorption of external water or by

dehydration of some of the formamide to hydrogen cyanide, and liberation of carbon monoxide from the formamide seems a likely sequence.

See other GAS EVOLUTION INCIDENTS

Organozinc compounds

Mistryukov, E. A. *et al.*, *Mendeleev Comm.*, 1993, (6) 242

Adding formamide to a reaction mixture of allylzinc bromide and a ketone as a proton source for a Barbier reaction in ether or tetrahydrofuran gave uncontrollable explosion-like reaction. The reaction proceeded smoothly when starting from zinc dust and allyl bromide, with both formamide and the ketone in situ.

0454. *N*-Hydroxydithiocarbamic acid

[66427-01-8]

HONHC(S)SH

CH₃NOS₂

Voigt, C. W. *et al.*, *Z. Anorg. Chem.*, 1977, **437**, 233–236

The free acid is unstable and may decompose explosively at sub-zero temperatures.

See other N–O COMPOUNDS, ORGANIC ACIDS

†0455. Methyl nitrite

[624-91-9]

MeON:O

CH₃NO₂

NSC 693, 1982; *HCS 1980*, 687

Rüst, 1948, 285

Explodes on heating, more powerfully than the ethyl homologue.

Other reactants

Yoshida, 1980, 14

MRH values calculated for 16 combinations, largely with oxidants, are given.

See other ALKYL NITRITES, N–O COMPOUNDS

0456. Nitromethane

[75-52-5]

MeNO₂

CH₃NO₂

1. *HCS 1980*, 687
2. McKitterick, D. S. *et al.*, *Ind. Eng. Chem. (Anal. Ed.)*, 1938, **10**, 630
3. Makovsky, A. *et al.*, *Chem. Rev.*, 1958, **58**, 627
4. Travis, J. R., *Los Alamos Rept. DC 6994*, Washington, USAEC, 1965
5. Sorbe, 1968, 148
6. Lieber, C. O., *Chem. Ing. Tech.*, 1978, **50**, 695–697
7. Perche, A. *et al.*, *J. Chem. Res.*, 1979, (S) 116; (M) 1555–1578
8. Lewis, D. J., *Haz. Cargo Bull.*, 1983, **4**(9), 36–38

9. Stull, 1977, 20
10. Anon., *Jahresber.*, 1984, 66–67
11. Coetzee, J. F. *et al.*, *Pure Appl. Chem.*, 1986, **58**, 1541–1545
12. Brewer, K. R., *J. Org. Chem.*, 1988, **53**, 3776–3779
13. Piermanini, G. J. *et al.*, *J. Phys. Chem.*, 1989, **93**(1), 457–462
14. Dick, J. J., *J. Phys. Chem.*, 1993, **97**(23), 6193
15. Ullmann, 1991, A 17, 405/6

Hazardous properties and handling procedures are summarised [1]. Conditions under which it may explode by detonation, heat or shock were determined. It was concluded that it is potentially very explosive and precautions are necessary to prevent its exposure to severe shock or high temperatures in use [2]. Later work, following two rail tank explosions, showed that shock caused by sudden application of gas pressure, or sudden forced flow through restrictions, could detonate the liquid. The stability and decomposition of nitromethane relevant to use as a rocket fuel are also reviewed [3]. The role of discontinuities in the initiation of shock-compressed nitromethane has been evaluated experimentally [4]. It explodes at about 230°C [5]. The role of pressure waves producing local resonance heating in bubbles and leading to detonation has been studied [6], as have the effects of nitrogen oxide, nitrogen dioxide and formaldehyde upon pyrolytic decomposition of nitromethane [7]. Bulk shipment in rail tanks is now permitted for solutions in appropriate diluents [8].

The heat of decomposition (238.4 kJ/mol, 3.92 kJ/g) has been calculated to give an adiabatic product temperature of 2150°C accompanied by a 24-fold pressure increase in a closed vessel [9]. During research into the Friedel-Crafts acylation reaction of aromatic compounds (components unspecified) in nitrobenzene as solvent, it was decided to use nitromethane in place of nitrobenzene because of the lower toxicity of the former. However, because of the lower boiling point of nitromethane (101°C, against 210°C for nitrobenzene), the reactions were run in an autoclave so that the same maximum reaction temperature of 155°C could be used, but at a maximum pressure of 10 bar. The reaction mixture was heated to 150°C and maintained there for 10 minutes, when a rapidly accelerating increase in temperature was noticed, and at 160°C the lid of the autoclave was blown off as decomposition accelerated to explosion [10]. Impurities present in the commercial solvent are listed, and a recommended purification procedure is described [11]. The thermal decomposition of nitromethane under supercritical conditions has been studied [12]. The effects of very high pressure and of temperature on the physical properties, chemical reactivity and thermal decomposition of nitromethane have been studied, and a mechanism for the bimolecular decomposition (to ammonium formate and water) identified [13]. Solid nitromethane apparently has different susceptibility to detonation according to the orientation of the crystal, a theoretical model is advanced [14]. Nitromethane actually finds employment as an explosive [15].

See Acetone below, or Haloforms below

Acetone

Varob'ev, A. A., *Chem. Abs.*, 1981, **94**, 49735

Mixtures can be detonated (title only translated).

Acids, or Bases

Makovky, A. *et al.*, *Chem. Rev.*, 1958, **58**, 631

Addition of bases or acids to nitromethane renders it susceptible to initiation by a detonator. These include aniline, diaminoethane, iminobispropylamine, morpholine, methylamine, ammonium hydroxide, potassium hydroxide, sodium carbonate, and formic, nitric, sulfuric or phosphoric acids.

See 1,2-Diaminoethane, *N*,2,4,6-Tetranitro-*N*-methylamine, below

Alkalis

Ullmann, 1991, A 17, 405/6

It will form *aci*-nitromethanoate salts, which are sensitive explosives when dry, and may form methazonates (salts of nitroacetaldehyde oxime) which are still more explosive.

See Sodium *aci*-nitromethanide, 2-Nitroacetaldehyde oxime

Alkylmetal halides

Traverse, G., US Pat. 2 775 863, 1957

Contact with R_mMX_n (R is methyl, ethyl; M is aluminium, zinc; X is bromide, iodide) causes ignition. Diethylaluminium bromide, dimethylaluminium bromide, ethylaluminium bromide iodide, methylzinc iodide and methylaluminium diiodide are claimed as specially effective.

Aluminium powder

MRH 9.33/23

Kato, Y. *et al.*, *Chem. Abs.*, 1979, **91**, 159867

Increase in concentration of aluminium powder in a mixture with poly(methyl methacrylate) increases the sensitivity to detonation.

Aluminium chloride

See Aluminium chloride–nitromethane

Aluminium chloride, Ethylene

See Ethylene: Aluminium chloride

Ammonium salts, Organic solvents

Runge, W. F. *et al.*, US Pat. 3 915 768, 1975

Presence of, for example, 5% of methylammonium acetate and 5% of methanol sensitises nitromethane to shock-initiation.

Bis(2-aminoethyl)amine

1. Runge, W. F. *et al.*, US Pat. 3 798 902, 1974

2. Walker, F. E., *Acta Astronaut.*, 1979, **6**, 807–813

Explosive solutions of nitromethane in dichloromethane, sensitised by addition of 10–12% of the amine, retained their sensitivity at –50°C [1]. Presence of 0–5% of the triamine considerably increases detonation sensitivity of nitromethane [2].

Bromine

See Bromine: Nitromethane

Boron trifluoride etherate, Silver oxide

See Silver tetrafluoroborate

Calcium hypochlorite

See Calcium hypochlorite: Nitromethane

Carbon Disulphide

Editor's comments

Salts of the reaction product, nitroethanedithioic acid, are intermediates in manufacture of pharmaceuticals. They are explosive and are to be avoided if not isolated. The free acid will surely be unstable, but not necessarily explosive.

1,2-Diaminoethane, *N*,2,4,6-Tetranitro-*N*-methylaniline

MCA Case History No. 1564

During preparations to initiate the explosion of nitromethane sensitised by addition of 20% of the diamine, accidental contact of the liquid mixture with the solid 'tetryl' detonator caused ignition of the latter.

See Acids, or Bases, above

Formaldehyde

Noland, W. E., *Org. Synth.*, 1961, **41**, 69

Interaction of nitromethane and formaldehyde in presence of alkali gives not only 2-nitroethanol, but also di- and tri-condensation products. After removal of the 2-nitroethanol by vacuum distillation, the residue must be cooled before admitting air into the system to prevent a flash explosion or violent fume-off.

Haloforms

Presles, H. N. *et al.*, *Acta Astronaut.*, 1976, **3**, 531–540

Mixtures with chloroform or bromoform are detonable.

Hydrazine, Methanol

Forshey, D. R. *et al.*, *Explosivstoffe*, 1969, **17**(6), 125–129

Addition of hydrazine strongly sensitises nitromethane and its mixtures with methanol to detonation.

Hydrocarbons

1. Watts, C. E., *Chem. Eng. News*, 1952, **30**, 2344

2. Makovsky, A. *et al.*, *Chem. Rev.*, 1958, **58**, 631

Nitromethane may act as a mild oxidant, and should not be heated with hydrocarbons or readily oxidisable materials under confinement [1]. Explosions may occur during cooling of such materials heated to high temperatures and pressures [2]. Mixtures of nitromethane and solvents which are to be heated above the b.p. of nitromethane should first be subjected to small-scale explosive tests [2].

Hydrocarbons, Oxygen

See Oxygen: Hydrocarbons, Promoters

CH₃NO₂

Lithium perchlorate

MRH 5.69/43

1. Titus, J. A., *Chem. Eng. News*, 1971, **49**(23), 6
2. 'Nitroparaffin Data Sheet TDS 1' New York, Comm. Solvents Corp., 1965
3. Egly, R. S., *Chem. Eng. News*, 1973, **51**(6), 30

Explosions which occurred at the auxiliary electrode during electro-oxidation reactions in nitromethane–lithium perchlorate electrolytes, may have been caused by lithium fulminate. This could have been produced by formation of the lithium salt of nitromethane and subsequent dehydration to the fulminate [1], analogous to the known formation of mercury(II) fulminate [2]. This explanation is not considered tenable, however [3].

Lithium tetrahydroaluminate

1. Nystrom, R. F. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 3739
2. Wollweber, H. *et al.*, private comm., 1988

Addition of nitromethane to ethereal lithium tetrahydroaluminate solution at ambient temperature gave an explosively violent reaction [1], and this was confirmed when addition of 0.5 ml of dry nitromethane to 10 ml of reductant solution led, after 30 s, to a violent explosion which pulverised the flask [2]. The violence of the explosion suggests that lithium *aci*-nitromethanide (or possibly lithium fulminate) may have been involved.

See Sodium hydride, below

See other *aci*-NITRO SALTS

Metal oxides

See NITROALKANES: metal oxides

Molecular sieve

1. Wollweber, H., private comm., 1979
2. Bretherick, L., *Chem. & Ind.*, 1979, 532

Nitromethane was dried and stored in a flask over 13X (large-pore) molecular sieve, and when a further portion of freshly activated sieve was added after several weeks, the contents erupted and ignited, breaking the flask [1]. This was attributed to slow formation of sodium *aci*-nitromethanide from the zeolitic sodium ions, which then decomposed in the exotherm arising from adsorption of nitromethane on the freshly activated sieve. Use of a small-pore (3A or 4A) sieve which would exclude nitromethane from the internal channels and greatly reduce the contact with sodium ions should avoid this problem. The possibility of other problems arising from slow release or exchange of ionic species under virtually anhydrous conditions during long term use of molecular sieves was noted [2].

See other MOLECULAR SIEVE INCIDENTS

Nickel

Benziger, J. B., *Appl. Surface Sci.*, 1984, **17**, 309–323

Decomposition of nitromethane on the (111) face of nickel has been studied.

Nitric acid

MRH 6.19/38

See Nitric acid: Nitromethane

Other reactants

Yoshida, 1980, 278

MRH values calculated for 17 combinations, largely with oxidants, are given.

Silver nitrate

Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 336

Aqueous silver nitrate may react to form silver fulminate.

Sodium hydride

1. Pearson, A. J. *et al.*, *J. Organomet. Chem.*, 1980, **202**, 178

2. Johnson, B. F. G. *et al.*, *J. Organomet. Chem.*, 1981, **204**, 221–228

Generation of sodium *aci*-nitromethanide by adding de-oiled sodium hydride powder to nitromethane in THF becomes violent at 40°C [1], and if the solvent is omitted, too fast addition of the hydride leads to a series of small explosions [2].

See Sodium *aci*-nitromethanide

Trimethylsilyl iodide

Voronkov, M. G. *et al.*, *Zh. Obshch. Khim.*, 1989, **59**(5), 1055. (*Chem. Abs.*, **112**, 77304)

Reaction of the silane with nitromethane is explosive, probably by intermediacy of fulminic acid (a dehydration product of nitromethane).

Uronium perchlorate

See Uronium perchlorate: Organic materials

See other NITROALKANES, C-NITRO COMPOUNDS

0457. Methyl nitrate

[598-58-3]

MeONO₂

CH₃NO₃

1. Kit and Evered, 1960, 268

2. Black, A. P. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 412

3. Goodman, H. *et al.*, *Combust. Flame*, 1972, **19**, 157

4. Griffiths, J. F. *et al.*, *Combust. Flame*, 1982, **45**, 54–66

5. Stull, 1977, 20

It has high shock- and thermal sensitivity, exploding at 65°C, and is too sensitive for use as a rocket mono-propellant [1]. Conditions during preparation of the ester from methanol and mixed nitric–sulfuric acids are fairly critical, and explosions may occur if it is suddenly heated or distilled in presence of acid [2]. Spontaneous ignition or explosion of the vapour at 250–316°C in presence of gaseous diluents [3], and the mechanism of exothermic decomposition and ignition [4] have been studied. The rather high decomposition exotherm (153.8 kJ/mol, 4.42 kJ/g) would raise the products to an adiabatic temperature approaching 2600°C, with a 31-fold pressure increase in a closed system [5].

Other reactants

Yoshida, 1980, 207

Of the MRH values calculated for 11 combinations, that with magnesium was most energetic at 8.70/38.

See other ALKYL NITRATES

†0458. Thallium(I) methanediazoate (Thallium *N*-nitrosomethylamide)

[113925-83-0]

CH₃N₂O₂l

TlON=NMe

1. Keefer, L. K. *et al.*, *J. Amer. Chem. Soc.*, 1988, **110**, 2800–2806, and footnote 20
2. Burns, M. E., private comm., 1985

It appears to be a stable covalent highly crystalline compound (unlike other metal methanediazoates). Alkanediazoates are easily converted to diazoalkanes, so should be regarded as capable of detonation. (Though named by the author as a methanediazoate, it is indexed and registered in *CA* as a nitrosomethylamide salt) [1]. A sample of the freshly synthesised compound was dissolved in dichlorodideuteromethane and sealed into an NMR tube. Four days later, when the tube was being opened for recovery of the sample, the tube exploded. This was attributed to diazomethane formation, possibly from reaction with traces of moisture sealed into the tube [2].

See other HEAVY METAL DERIVATIVES, N–O COMPOUNDS

0459. Methyl azide

[624-90-8]

CH₃N₃

MeN₃

1. *MCA Case History No. 887*
2. Boyer, J. H. *et al.*, *Chem. Rev.*, 1954, **54**, 323
3. Burns, M. E. *et al.*, *Chem. Eng. News*, 1984, **62**(2), 2

The product, prepared by interaction of sodium azide with dimethyl sulfate and sodium hydroxide, exploded during concurrent vacuum distillation. The explosion was attributed to formation and co-distillation with the product of hydrogen azide, owing to excursion of the pH to below 5 during the preparation. Free hydrogen azide itself is explosive, and it may also have reacted with mercury in a manometer to form the detonator mercuric azide [1]. Methyl azide is stable at ambient temperature, but may detonate on rapid heating [2]. A further explosion during the preparation as above led to incorporation of bromothymol blue in the reaction system to give a visible indication of pH in the generation flask [3].

Mercury

Currie, C. L. *et al.*, *Can. J. Chem.*, 1963, **41**, 1048

Presence of mercury in methyl azide markedly reduces the stability towards shock or electric discharge.

Methanol

Grundman, C. *et al.*, *Angew. Chem.*, 1950, **62**, 410

In spite of extensive cooling and precautions, a mixture of methyl azide, methanol and dimethyl malonate exploded violently while being sealed into a Carius tube. The vapour of the azide is very easily initiated by heat, even at low concentrations.

See other ORGANIC AZIDES

0460. Nitrourea
[556-89-8]




1. Urbanski, 1967, Vol. 3, 34
2. Medard, L., *Mém. Poudres*, 1951, **33**, 113–123; (English transl., HSE 11292, available from HSE/LIS, 1986)

A rather unstable explosive material, insensitive to heating or impact, which gives mercuric and silver salts which are rather sensitive to impact [1]; further detailed data are available [2].

See other N-NITRO COMPOUNDS

0461. 5-Aminotetrazole
[4418-61-5]



Nitrous acid

1. Elmore, D. T., *Chem. Brit.*, 1966, **2**, 414
2. Thiele, J., *Ann.*, 1892, **270**, 59
3. Gray, E. J. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1976, 1503
4. Cawkill, E. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1979, 727


Diazotised 5-aminotetrazole is unstable under the conditions recommended for its use as a biochemical reagent. While the pH of the diazotised material (the cation of which contains 87% nitrogen) at 0°C was being reduced to 5 by addition of potassium hydroxide, a violent explosion occurred [1]. This may have been caused by a local excess of alkali causing the formation of the internal salt, 5-diazoniotetrazolide, which will explode in concentrated solution at 0°C [2]. The diazonium chloride is also very unstable in concentrated solution at 0°C. Small-scale diazotisation (2 g of amine) and subsequent coupling at pH 3 with ethyl cyanoacetate to prepare ethyl 2-cyano-(1H-tetrazol-5-ylhydrazono)acetate proceeded uneventfully, but on double the scale a violent explosion occurred [3]. The importance of adequate dilution of the reaction media to prevent explosions during diazotisation is stressed [4].

See DIAZONIUM SALTS

See other HIGH-NITROGEN COMPOUNDS, TETRAZOLES

0462. 5-Hydrazino-1,2,3,4-thiaziazole
[99319-30-9]

CH₃N₅S



Martin, D., *Z. Chem.*, 1985, **25**(4), 136–137

The explosive title compound was not isolated after preparation from thiocarbazide and nitrous acid, but was condensed with various ketones to give the hydrazones.

See other HIGH-NITROGEN COMPOUNDS, N-S COMPOUNDS

See related TRIAZOLES

0463. Methylsodium
[18356-02-0]

CH₃Na

H₃CNa

Schlenk, W. *et al.*, *Ber.*, 1917, **50**, 262

Ignites immediately in air. The tendency to ignition decreases with ascent of the homologous series of alkylsodiums.

See other ALKYL METALS

0464. Sodium methoxide
[124-41-4]

CH₃NaO

NaOCH₃

1. *HCS 1980*, 850

2. *See entry* THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

It may ignite in moist air. Hazardous properties and handling procedures are summarised [1]. Energy of decomposition (in range 410–460 °C) measured as 0.77 kJ/g [2].

Chloroform

See Chloroform: Sodium methoxide

4-Chloronitrobenzene

See 4-Chloronitrobenzene: Sodium methoxide

Perfluorocyclopropene

See FLUORINATED CYCLOPROPENYL METHYL ETHERS

See other METAL ALKOXIDES

0465. Methanesulfonyl azide

[1516-70-7]

CH₃N₃O₂S**MeSO₂N₃**Brown, R. *et al.*, *Chem. Brit.*, 1994, **30**(6), 470

Gram quantities of the azide have twice detonated, once during distillation, the second time also possibly due to localised overheating. It is stated that there was hitherto only a general warning of possible explosivity.

See other ACYL AZIDES

†0466. Methane

[74-82-8]

CH₄**CH₄**

FPA H59, 1977; *HCS 1980*, 627 (cylinder)

Halogens, or Interhalogens

See Bromine pentafluoride: Hydrogen-containing materials

Chlorine: Hydrocarbons

Chlorine trifluoride: Methane

Fluorine: Hydrocarbons

Iodine heptafluoride: Carbon, etc.

Other reactants

Yoshida, 1980, 361

MRH values calculated for 12 combinations with oxidants are given.

Oxidants

See Dioxygen difluoride: Various materials

Dioxygenyl tetrafluoroborate: Organic materials

Oxygen (Liquid): Hydrocarbons, or: Liquefied gases

'Trioxygen difluoride': Various materials

See also LIQUEFIED NATURAL GAS

0467. Chloroformamidinium nitrate

[75524-40-2]

CH₄ClN₃O₃**ClC(:NH)N⁺H₃ NO₃⁻**

Alone, or Amines, or Metals

1. Sauermilch, W., *Explosivstoffe*, 1961, **9**, 71–74

2. Sorbe, 1968, 144

It is powerfully explosive, and also an oxidant which reacts violently with ammonia or amines, and causes explosive ignition of wet magnesium powder [1], or of powdered aluminium or iron [2].

See Chloroformamidinium chloride: Oxyacids, etc. (next below)

See other OXOSALTS OF NITROGENOUS BASES

0468. Chloroformamidinium chloride

[29671-92-9]

CH₄Cl₂N₂

Oxyacids and salts

Sauermilch, W., *Explosivstoffe*, 1961, **9**, 71–74, 256

The title compound ('cyanamide dihydrochloride') reacts with perchloric acid to form the perchlorate salt, and with ammonium nitrate, the nitrate salt, both being highly explosive, and the latter also an oxidant.

0469. Chloroformamidinium perchlorate (Methanimidamide monoperchlorate)

[40985-41-9]

CH₄Cl₂N₂O₄

See item above See other PERCHLORATE SALTS OF NITROGENOUS BASES

†0470. Dichloromethylsilane

[75-54-7]

CH₄Cl₂Si

Oxidants

1. Müller, R. *et al.*, *J. Prakt. Chem.*, 1966, **31**, 1–6

2. Sorbe, 1968, 128

The pure material is not ignited by impact, but it is in presence of potassium permanganate or lead(II),(IV) oxide [1], or by copper oxide or silver oxide, even under an inert gas [2].

See Trichlorosilane

See other ALKYLHALOSILANES

0471. Tetrakis(hydroxymercurio)methane

[]

CH₄Hg₄O₄Breitinger, D. K. *et al.*, *Z. Naturforsch. B*, 1984, **39B**, 123

Intensive and heated high vacuum drying of the title compound (Hofmann's base) over phosphorus(V) oxide may lead to an explosion. Unheated drying appears satisfactory and safer.

See other MERCURY COMPOUNDS

0472. Potassium methylamide

[[54448-39-4] (ion)]

CH₄KNMakhija, R. C. *et al.*, *Can. J. Chem.*, 1971, **49**, 807

Extremely hygroscopic and pyrophoric; may explode on contact with air.

See other N-METAL DERIVATIVES

0473. Formylhydrazine
[624-84-0]



See 4-Amino-4H-1,2,4-triazole
See other HIGH-NITROGEN COMPOUNDS

0474. Methyldiazene
[26981-93-1]



Oxygen

Ackermann, M. N. *et al.*, *Inorg. Chem.*, 1972, **11**, 3077

Interaction on warming a mixture from -196°C rapidly to ambient temperature is explosive.

See related AZO COMPOUNDS

0475. Urea (Carbamide)
[57-13-6]



NSC 691, 1980; *HCS 1980*, 949

Preparative hazard for ^{15}N -labelled compound.

See Oxygen (Liquid): Ammonia, etc.

Chromyl chloride

See Chromyl chloride: Urea

Dichloromaleic anhydride, Sodium chloride

See Dichloromaleic anhydride: Sodium chloride, etc.

Metal hypochlorites

491M, 1975, 213

Urea reacts with sodium hypochlorite or calcium hypochlorite to form explosive nitrogen trichloride.

See Chlorine: Nitrogen compounds

Nitrosyl perchlorate

See Nitrosyl perchlorate: Organic materials

Oxalic acid

See Oxalic acid: Urea

Phosphorus pentachloride


See Phosphorus pentachloride: Urea

Sodium nitrite

See Sodium nitrite: Urea

Titanium tetrachloride
See Titanium tetrachloride: Urea
See other ORGANIC BASES

0476. Urea hydrogen peroxidate
[124-43-6]




MCA Case History No. 719

The contents of a screw-capped brown glass bottle spontaneously erupted after 4 years' storage at ambient temperature. All peroxides should be kept in special storage and checked periodically.

See other CRYSTALLINE HYDROGEN PEROXIDATES, GLASS INCIDENTS

0477. Hydroxyurea
[127-07-1]



Cardillo, P. *et al.*, *Chem. Eng. News*, 1998, **76**(22), 6

An industrial reactor containing a concentrate of this in water was left over the weekend, at an initial temperature possibly around 50°C. On the Tuesday morning it exploded. Investigation of the thermal stability of this drug in pure form showed it to be thermally unstable from 85°C as the solid, 70°C in aqueous solution. The decomposition is probably autocatalytic, gas evolving and with an enthalpy of 2.2 kJ/g. Aged samples were still less stable, decomposing after a variable induction period (from an initial 50°C, 80 hours). Gas evolution combined with temperature elevation to boiling point explains the burst reactor.

See other N-O COMPOUNDS

0478. N-Nitromethylamine
[598-57-2]



Diels, O. *et al.*, *Berichte*, 1913, **46**, 2006.

A tiny quantity exploded with the utmost force when strongly heated in a capillary tube.

Amminemetals

Complexes with several ammine derivatives of metals are explosive.

See entry NITRAMINE-METAL COMPLEXES

Sulfuric acid

Urbanski, 1967, Vol. 3, 16

The nitroamine is decomposed explosively by conc. sulfuric acid.

See other N-NITRO COMPOUNDS

0479. Ammonium thiocyanate
[1762-95-4]



HCS 1980, 152

Other reactants

Yoshida, 1980, 227

MRH values calculated for 13 combinations, 12 with oxidants, are given.

Potassium chlorate

See Potassium chlorate: Metal thiocyanates

See related CYANO COMPOUNDS

0480. Thiourea
[62-56-6]



Acrylaldehyde

See Acrylaldehyde: Acids, etc.

Hydrogen peroxide, Nitric acid

See Hydrogen peroxide: Nitric acid, etc.

Potassium chlorate

See Potassium chlorate: Thiourea

See other ORGANIC BASES

0481. Nitrosoguanidine
[674-81-7]



Henry, R. A. *et al.*, *Ind. Eng. Chem.*, 1949, **41**, 846–849

The compound is normally stored and transported as a water-wet paste which slowly decomposes at elevated ambient temperatures, evolving nitrogen. Precautions are described to prevent pressure build-up in sealed containers.

See other NITROSO COMPOUNDS

0482. Nitroguanidine
[556-88-7]



1. Urbanski, 1967, Vol. 3, 31

2. McKay, A. F., *Chem. Rev.*, 1952, **51**, 301

Nitroguanidine is difficult to detonate, but its mercury and silver complex salts are much more impact-sensitive [1]. Many nitroguanidine derivatives have been considered as explosives [2].

See other N-NITRO COMPOUNDS

0483. Methylenebis(nitramine)

[14168-44-6]

CH₄N₄O₄

H₂C(NHNO₂)₂

Glowiak, B., *Chem. Abs.*, 1960, **54**, 21761e

It is a powerful and sensitive explosive which explodes at 217°C, and the lead salt at 195°C.

See other N-NITRO COMPOUNDS

†0484. Methanol

[67-56-1]

CH₄O

H₃COH

(MCA SD-22, 1970); NSC 407, 1979; FPA H42, 1975; HCS 1980, 628; RSC Lab. Hazard Data Sheet No. 25, 1984

Ferris, T. V., *Loss Prev.*, 1974, **8**, 15–19

Explosive behaviour on combustion of methanol–air mixtures at 1.8 bar and 120°C was studied, with or without addition of oxygen and water.

Acetyl bromide

See Acetyl bromide: Hydroxylic compounds

Alkylaluminium solutions

MCA Case History No. 1778

Accidental use of methanol in place of hexane to rinse out a hypodermic syringe used for a dilute alkylaluminium solution caused a violent reaction which blew the plunger out of the barrel.

See ALKYLALUMINIUM DERIVATIVES: alcohols

Beryllium hydride

See Beryllium hydride: Methanol

Boron trichloride

See Boron trichloride

Carbon tetrachloride, Metals

Kuppers, J. R., *J. Electrochem. Soc.*, 1978, **125**, 97–98

The rapid autocatalytic dissolution of aluminium, magnesium or zinc in 9:1 methanol–carbon tetrachloride mixtures is sufficiently vigorous to be rated as potentially hazardous. Dissolution of zinc powder is subject to an induction period of 2 h, which is eliminated by traces of copper(II) chloride, mercury(II) chloride or chromium(III) bromide.

See other INDUCTION PERIOD INCIDENTS

Chloroform, Sodium

See Chloroform: Sodium, Methanol

Chloroform, Sodium hydroxide
See Chloroform: Sodium hydroxide, Methanol

Cyanuric chloride
See 2,4,6-Trichloro-1,3,5-triazine: Methanol

Dichloromethane
See Dichloromethane: Air, etc.

Diethylzinc
See Diethylzinc: Methanol

Hydrogen, Raney nickel catalyst
Klais, O., *Hazards from Pressure, IChE Symp. Ser. No. 102*, 25–36, Oxford, Pergamon, 1987
During hydrogenation of an unspecified substrate in methanol solution under hydrogen at 100 bar with Raney nickel catalyst, a sudden temperature increase led to hydrogenolysis of methanol to methane, and the pressure increase led to an overpressure accident. Such incidents may be avoided by control of agitation, limiting the amount of catalyst, and checking thermal stability of starting materials and end products beforehand.
See other GAS EVOLUTION INCIDENTS

Metals
See Aluminium: Methanol
Magnesium: Methanol
Potassium (Slow oxidation)

Other reactants
Yoshida, 1980, 360
MRH values calculated for 17 combinations, largely with oxidants, are given.

Oxidants	MRH values show % of oxidant
<i>See</i> Barium perchlorate: Alcohols	
Bromine: Alcohols	
Chlorine: Methanol	
Chromium trioxide: Alcohols	
Hydrogen peroxide: Oxygenated compounds	MRH 5.98/76
Lead perchlorate: Methanol	
Nitric acid: Alcohols (reference 6)	MRH 5.31/70
Sodium hypochlorite: Methanol	MRH 2.47/90


Phosphorus(III) oxide
See Tetraphosphorus hexaoxide: Organic liquids

Potassium *tert*-butoxide
See Potassium *tert*-butoxide: Acids, etc.

Water
MCA Case History Nos. 1822, 2085

Static discharge ignited the contents of a polythene bottle being filled with a 40:60 mixture of methanol and water at 30°C, and a later similar incident in a plastics-lined metal tank involved a 30:70 mixture.

0485. Poly[oxy(methyl)silylene]
[9004-73-3]



Preparative hazard

Crivello, J. V. *et al.*, *Chem. Abs.*, 1990, **113**, 41011z; U.S. Pat. 4,895,967.

A procedure for cyclisation of the linear polymer to small oligomers without risk of explosion is described.

1-Allyloxy-2,3-epoxypropane

Anon., *Eur. Chem. News*, 1995, **63**(1656), 27

The two compounds were accidentally mixed in a reactor because of similar packaging. They reacted uncontrollably, rupturing the reactor, and the evolved hydrogen ignited, killing one worker and injuring four. It is probable a base catalyst was present depolymerisation/disproportionation to oxomethylsilane and methylsilane is also possible, with an increase in both vapour pressure and flammability.

See other SILANES

0486. Methyl hydroperoxide
[3031-73-0]



Alone, or Phosphorus(V) oxide, or Platinum

1. Rieche, A. *et al.*, *Ber.*, 1929, **62**, 2458, 2460

2. Hasegawa, K. *et al.*, *Proc. Int. Pyrotech. Semin.*, 1994, (19th), 684

The hydroperoxide is violently explosive and shock-sensitive, especially on warming; great care is necessary in handling. It explodes violently in contact with phosphorus pentoxide, and a 50% aqueous solution decomposed explosively on warming with spongy platinum. The barium salt is dangerously explosive when dry.

See other ALKYL HYDROPEROXIDES

0487. Hydroxymethyl hydroperoxide
[15932-89-5]



Rieche, A., *Ber.*, 1931, 64, 2328; 1935, 68, 1465

It explodes on heating, but is friction-insensitive. Higher homologues are not explosive.

See other 1-OXYPEROXY COMPOUNDS

0488. Methanesulfonic acid
[75-75-2]



Ethyl vinyl ether

See Ethyl vinyl ether: Methanesulfonic acid

Hydrogen fluoride

See Hydrogen fluoride: Methanesulfonic acid

See other ORGANIC ACIDS

†0489. Methanethiol
[74-93-1]



FPA H93, 1980; HCS 1980, 655

Mercury(II) oxide

See Mercury(II) oxide: Methanethiol

See other ALKANETHIOLS, ALKYLNON-METAL HYDRIDES

0490. Methanetellurol
[25284-83-7]




Hamada, K. *et al.*, *Synth. React. Inorg. Metal-Org. Chem.*, 1977, **7**, 364 (footnote 1)

It ignites in air and explodes with oxygen at ambient temperature.

See other ALKYLNON-METAL HYDRIDES

0491. Uronium perchlorate (Urea perchlorate)
[18727-07-6]



Aromatic nitro compounds

1. Shimio, K. *et al.*, *Chem. Abs.*, 1975, **83**, 134504

2. Fujiwara, S. *et al.*, *Japan Kokai*, 74 134 812, 1974

Conc. aqueous solutions of the urea salt will dissolve solid or liquid aromatic nitro compounds (e.g. picric acid [1], or nitrobenzene [2]) to give high velocity explosives.

See next item below

Organic materials

Kusakabe, M. *et al.*, *Chem. Abs.*, 1980, **92**, 79028

Liquid mixtures with detonable materials (picric acid, nitromethane) were extremely powerful explosives, and those with nitrobenzene or dimethylformamide less so.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

0492. Fluoromethylsilane

[10112-08-0]



CH_5FSi

Chlorine, Dichloromethane

See Chlorine: Fluoromethylsilane, Dichloromethane

See other SILANES: halogens

†0493. Methylamine (Methanamine)

[74-89-5]



CH_5N

(MCA SD-57, 1965); HCS 1980, 637

Nitromethane

See Nitromethane: Acids, etc.

Other reactants


Yoshida, 1980, 362

MRH values calculated for 13 combinations with oxidants are given.

See other ORGANIC BASES

0494. Uronium nitrate (Urea nitrate)

[124-47-0]



$\text{CH}_5\text{N}_3\text{O}_4$

1. Kirk-Othmer, 1970, Vol. 21, 38

2. Markalous, F. *et al.*, Czech Pat. 152 080, 1974

3. Lazarov, S. *et al.*, *Chem. Abs.*, 1973, **79**, 94169

4. Medard, L., *Mém. Poudres*, 1951, **33**, 113–123

(English transl. HSE 11292 available from HSE/LIS, 1986)

The nitrate decomposes explosively when heated [1]. Prepared in the presence of phosphates, the salt is much more stable, even when dry [2]. The manufacture and explosive properties of urea nitrate and its mixtures with other explosives are discussed in detail [3]. Explosive properties of the nitrate are detailed [4].

Heavy metals

Kotzerke, *Proc. 1st Int. Symp. Prev. Occ. Risks Chem. Ind.*, 201–207, Heidelberg, ISSA, 1970

During evaporation of an alkaline aqueous solution of the nitrate, decomposition led to gas evolution and a pressure explosion occurred. This was attributed to the use of recovered alkali containing high levels of lead and iron, which were found to catalyse the thermal decomposition of the nitrate. Precautions to prevent recurrence are detailed.

See other GAS EVOLUTION INCIDENTS, OXOSALTS OF NITROGENOUS BASES

0495. 1-Amino-3-nitroguanidine


[18264-75-0]

CH₅N₅O₂Lieber, E. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 2328

It explodes at the m.p., 190°C.

See other HIGH-NITROGEN COMPOUNDS, *N*-NITRO COMPOUNDS**0496. Hydrazinium trinitromethanide (Hydrazinium nitroformate)**


[14913-74-7]

CH₅N₅O₆1. Anan, T. *et al.*, *Chem. Abs.*, 1995, **123**, 1743512. Schoyer, H. F. R. *et al.*, *J. Propul. Power.*, 1995, **11**(4), 856

The compound was examined as a propellant explosive, it proved easily detonable and more sensitive than ammonium perchlorate. A formulation with hydroxyl terminated polybutadiene binder ignited spontaneously at room temperature [1]. Other workers have found it more tractable [2].

See other HYDRAZINIUM SALTS POLYNITROALKYL COMPOUNDS**0497. Methylbis(η^2 -peroxo)rhenium oxide hydrate**

[]

CH₅O₆ReHerrmann, W. A. *et al.*, *Angew. Chem. (Int.)*, 1993, **32**, 1157

This, the active species in methylrheniumtrioxide catalysed peroxidations by hydrogen peroxide, is fairly stable in solution but explosive when isolated.

See other ORGANOMETALLIC PEROXIDES**0498. Methylphosphine**

[5931-54-4]

CH₅P

Houben-Weyl, 1963, Vol. 12.1, 69

Primary lower-alkylphosphines readily ignite in air.

See other ALKYLPHOSPHINES**0499. Methylstibine**

[23362-09-6]

CH₅Sb


Sorbe, 1968, 28

Alkylstibines decompose explosively on heating or shock.

See other ALKYL METAL HYDRIDES

0500. Methylammonium chlorite


[15875-44-2]

Levi, G. R., *Gazz. Chim. Ital.* [1], 1922, **52**, 207

A conc. solution caused a slight explosion when poured onto a cold iron plate.

See other CHLORITE SALTS, OXOSALTS OF NITROGENOUS BASES**0501. Methylammonium perchlorate**

[16875-44-2]

1. Kasper, F., *Z. Chem.*, 1969, **9**, 342. Kempson, R. M., *Chem. Abs.*, 1980, **92**, 61259

The semi-crystalline mass exploded when stirred after standing overnight. The preparation was based on a published method used uneventfully for preparation of ammonium, dimethylammonium and piperidinium perchlorates [1]. Its use in explosives and propellants has been surveyed [2].

See other PERCHLORATE SALTS OF NITROGENOUS BASES**0502. Guanidinium perchlorate**

[10308-84-6]



1. Davis, 1943, 121

2. Schumacher, 1960, 213

3. Fujiwara, S. *et al.*, *Chem. Abs.*, 1980, **92**, 2004434. Udupa, M. R., *Propellants, Explos., Pyrotech.*, 1983, **8**, 109–111

Unusually sensitive to initiation and of high explosive power [1], it decomposes violently at 350°C [2]. The explosive properties have been determined [3], and thermal decomposition at 275–325° was studied in detail [4].

See DIFFERENTIAL THERMAL ANALYSIS (reference 1)

Iron(III) oxide

Isaev, R. N. *et al.*, *Chem. Abs.*, 1970, **73**, 132626

Addition of 10% of iron oxide reduces the thermal stability of the salt.

See other PERCHLORATE SALTS OF NITROGENOUS BASES**†0503. Methylhydrazine**

[60-34-4]



491M 1975, 259

When extensively exposed to air (as a thin film, or absorbed on fibrous or porous solids), it may ignite.

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

Oxidants


Kirk-Othmer, 1966, Vol. 11, 186

A powerful reducing agent and fuel, hypergolic with many oxidants such as dinitrogen tetroxide or hydrogen peroxide.

See ROCKET PROPELLANTS See other ORGANIC BASES, REDUCANTS

0504. Ammonium aci-nitromethanide

[]




Watts, C. E., *Chem. Eng. News*, 1952, **30**, 2344

The isolated salt is a friction-sensitive explosive.

See other aci-NITRO SALTS

0505. Methylammonium nitrite

[68897-47-2]




Hara, Y., *Chem. Abs.*, 1982, **97**, 55056

It is unstable at ambient temperature, and is formed on contact of the nitrate with sodium or potassium nitrites.

See other NITRITE SALTS OF NITROGENOUS BASES, OXOSALTS OF NITROGENOUS BASES

0506. Methylammonium nitrate (Methanaminium nitrate)

[1941-24-8]



1. Biasutti, 1981, 145

2. Miron, Y., *J. Haz. Mat.*, 1980, **3**, 301–321

3. Hara, Y., *Chem. Abs.*, 1982, **97**, 55056

Rail tanks of 86% aqueous solutions or slurries of the salt exploded, apparently during pump-transfer operations [1]. The course and mechanism or thermal decomposition has been investigated. Traces of rust or copper powder catalyse and accelerate the decomposition, so corrosion prevention is an important aspect of safety measures [2]. It is of higher thermal stability than the chlorate salt, or the nitrite, which decomposes at ambient temperature [3].

See other CATALYTIC IMPURITY INCIDENTS See other OXOSALTS OF NITROGENOUS BASES

0507. Aminoguanidine

[79-17-4]



Kurzer, F. *et al.*, *Chem. & Ind.*, 1962, 1585

All the oxoacid salts are potentially explosive, including the wet nitrate.

See Aminoguanidinium nitrate
See other HIGH-NITROGEN COMPOUNDS, ORGANIC BASES

0508. Carbonic dihydrazide
[497-18-7]

CH₆N₄O



Sorbe, 1968, 74
It explodes on heating.

Nitrous acid

Curtius, J. *et al.*, *Ber.*, 1894, **27**, 55
Interaction forms the highly explosive carbonic diazide.
See Carbonic diazide
See other HIGH-NITROGEN COMPOUNDS

0509. Guanidinium nitrate
[52470-25-4]

CH₆N₄O₃



1. Davis, T. L., *Org. Synth.*, 1944, Coll. Vol. 1, 302
2. Smith, G. B. L., and Schmidt, M. T., *Inorg. Synth.*, 1939, **1**, 96, 97
3. Schopf, C. *et al.*, *Angew. Chem.*, 1936, **49**, 23
4. Udupa, M. R., *Thermochim. Acta*, 1982, **53**, 383–385
5. Medard, L. *Mém. Poudres*, 1951, **33**, 113–123 (English transl. HSE 11292 available from HSE/LIS, 1986)
6. Indian Pat. IN 157 293, Projects and Developments (India) Ltd., 1986 (*Chem. Abs.*, 1986, **105**, 210788)

According to an O.S. amendment sheet, the procedure as described [1] is dangerous because the reaction mixture (dicyanodiamide and ammonium nitrate) is similar in composition to commercial blasting explosives. This probably also applies to similar earlier preparations [2]. An earlier procedure which involved heating ammonium thiocyanate, lead nitrate and ammonia demolished a 50 bar autoclave [3]. TGA and DTA studies show that air is not involved in the thermal decomposition [4]. Explosive properties of the nitrate are detailed [5]. An improved process involves catalytic conversion at 90–200°C of a molten mixture of urea and ammonium nitrate to give 92% conversion (on urea) of guanidinium nitrate, recovered by crystallisation. Hazards of alternative processes are listed [6].
See other OXOSALTS OF NITROGENOUS BASES

0510. Methylsilane
[992-94-9]

CH₆Si




Mercury, Oxygen
Stock, A. *et al.*, *Ber.*, 1919, **52**, 706

It does not ignite in air, but explodes if shaken with mercury in oxygen.
See other ALKYL SILANES

0511. Methylhydrazinium nitrate
[29674-96-2]

CH₇N₃O₃



Lawton, E. A. *et al.*, *J. Chem. Eng. Data*, 1984, **29**, 358
Impact sensitivity is comparable to that of ammonium perchlorate.
See other OXOSALTS OF NITROGENOUS BASES

0512. Aminoguanidinium nitrate
[10308-82-4]



CH₇N₅O₃



Koopman, H., *Chem. Weekbl.*, 1957, **53**, 97
An aqueous solution exploded violently during evaporation on a steam bath. Nitrate salts of many organic bases are unstable and should be avoided.
See Aminoguanidine
See other HIGH-NITROGEN COMPOUNDS, OXOSALTS OF NITROGENOUS BASES

0513. Methylidborane
[23777-55-1]


CH₈B₂



Bunting, *Inorg. Synth.*, 1979, **19**, 237
It ignites explosively in air.
See other ALKYL BORANES

0514. Diaminoguanidinium nitrate
[10308-83-5]


CH₈N₆O₃



Violent decomposition at 260°C.
See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1)
See other OXOSALTS OF NITROGENOUS BASES

0515. Triaminoguanidinium perchlorate


[4104-85-2]

CH₉ClN₆O₄

Violent decomposition at 317°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1)*See other* PERCHLORATE SALTS OF NITROGENOUS BASES**0516. O-Methyl-N,N-disilylhydroxylamine**


[]

CH₉NOSi₂Mitzel, N. W. *et al.*, *J. Amer. Chem. Soc.*, 1996, **118**(11), 2664

Reasonably stable at ambient temperature, the compound explodes if heated to 200°C

See other SILANES, N-O COMPOUNDS**0517. Triaminoguanidinium nitrate**

[4000-16-2]

CH₉N₇O₃

Violent decomposition at 230°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1)*See other* OXOSALTS OF NITROGENOUS BASES**0518. Caesium cyanotridecahydrodecaborate(2-)**

[71250-00-5]

CH₁₃B₁₀Cs₂N

Hydrochloric acid

Schultz, R. V. *et al.*, *Inorg. Chem.*, 1979, **18**, 2883

Addition of the salt to conc. hydrochloric acid is extremely exothermic.

See other CYANO COMPOUNDS**0519. Pentaamminethiocyanatocobalt(III) perchlorate**

[15663-42-0]

CH₁₅Cl₂CoN₆O₈S

Explodes at 325°C; medium impact-sensitivity.

See entry AMMINOMETAL OXOSALTS (reference 3)

0520. Pentaamminethiocyanatoruthenium(III) perchlorate
[38139-15-0] **CH₁₅Cl₂N₆O₈RuS**
 $[(\text{H}_3\text{N})_5\text{RuSC}\equiv\text{N}] [\text{ClO}_4]_2$

Armour, J. N., private comm., 1969
After washing with ether, 0.1 g of the complex exploded violently when touched with a spatula.
See other AMMINOMETAL OXOSALTS, REDOX COMPOUNDS

0521. Hafnium carbide
[12069-85-1] **CHf**
HfC

See entry REFRACTORY POWDERS

0522. Poly(diazomethylenemercury) ('Mercury diazocarbide')
[31724-50-2] **(CHgN₂)_n**
[HgCN₂]_n

Houben-Weyl, 1974, Vol. 13.2b, 19
Interaction of diazomethane with bis(trimethylsilylamino)mercury gives the polymeric explosive solid.
See other DIAZO COMPOUNDS, MERCURY COMPOUNDS

0523. Mercury(I) cyanamide
[72044-13-4] **CHg₂N₂**
Hg₂NC≡N

Deb, S. K. *et al.*, *Trans. Faraday Soc.*, 1959, **55**, 106–113
Relatively large particles explode on heating rapidly to 325°C, or under high-intensity illumination when confined.
See other IRRADIATION DECOMPOSITION INCIDENTS, MERCURY COMPOUNDS, *N*-METAL DERIVATIVES

0524. Iodine isocyanate
[3607-48-5] **CINO**
IN=C=O

Rosen, S. *et al.*, *Anal. Chem.*, 1966, **38**, 1394
On storage, solutions of iodine isocyanate gradually deposit a touch-sensitive, mildly explosive solid (possibly cyanogen peroxide).
See related *N*-HALOGEN COMPOUNDS *See other* IODINE COMPOUNDS

0525. Carbon tetraiodide (Tetraiodomethane)
[507-25-5]



CI_4

Bromine trifluoride

See Bromine trifluoride: Halogens, etc.

Lithium

See Lithium: Halocarbons

See other HALOALKANES

0526. Potassium cyanide
[151-50-8]



CKN

HCS 1980, 764

Mercury(II) nitrate

See Mercury(II) nitrate: Potassium cyanide

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Perchloryl fluoride

See Perchloryl fluoride: Calcium acetylide, etc.

Sodium nitrite

See Sodium nitrite: Metal cyanides

See entry METAL CYANIDES (AND CYANO COMPLEXES) (reference 1)

0527. Potassium cyanide–potassium nitrite
[]



CKN.KNO_2

Sorbe, 1968, 68

The double salt (a redox compound) is explosive.

See other REDOX COMPOUNDS

0528. Potassium cyanate
[590-28-3]



CKNO

Water

Prinsky, M. L. *et al.*, *J. Loss Prev. Process Ind.*, 1990, **3**, 345

A drum of 30% solution in water exploded an hour after filling at 50°C, despite having a vent. Calorimetry demonstrated an exothermic, autocatalytic hydrolysis to ammoniacal potassium bicarbonate. In theory, a pressure exceeding 30 bar is obtainable. Aqueous solutions are unstable even at room temperature. Similar hydrolysis may account for an explosive product with Gold(III) chloride.

See Gold(III) chloride

See related METAL CYANATES, METAL CYANIDES

0529. Potassium thiocyanate

[333-20-0]

CKNS



HCS 1980, 778

Calcium chlorite

See Calcium chlorite: Potassium thiocyanate

Other reactants

Yoshida, 1980, 228

MRH values calculated for 13 combinations with oxidants are given.

Perchloryl fluoride

See Perchloryl fluoride: Calcium acetylide, etc.

See related METAL CYANIDES

0530. Potassium trinitromethanide

[14268-23-6]

CKN₃O₆



1. Sandler, S. R. *et al.*, *Organic Functional Group Preparations*, 433, New York, Wiley, 1968

2. Shulgin, A. T., private comm., 1968

This intermediate, produced by action of alkali on tetranitromethane, must be kept damp and used as soon as possible with great care, as it may be explosive [1]. Material produced as a by-product in a nitration reaction using tetranitromethane was washed with acetone. It exploded very violently after several months' storage [2].

See other POLYNITROALKYL COMPOUNDS

0531. Potassium carbonate

[584-08-7]

CK₂O₃



Carbon

Druce, J. G. F., *School Sci. Rev.*, 1926, 7(28), 261

Potassium metal prepared by the old process of distilling an intimate mixture of the carbonate and carbon contained some 'carbonylpotassium' (actually potassium benzenehexoxide), and several explosions with old samples of potassium may have involved this compound (or, perhaps more likely, potassium superoxide).

See Potassium (Slow oxidation)

Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Magnesium

See Magnesium: Potassium carbonate

See other INORGANIC BASES, METAL OXONON-METALLATES

0532. Lithium thiocyanate

[556-65-0]

CLiNS



1,3-Dioxolane

Rao, B. L. M. *et al.*, *J. Electrochem. Soc.*, 1980, **127**, 2333–2335

Thermal decomposition of the thiocyanate in air at 650°C yields 0.01% of volatile cyanide, but presence of moisture or the solvent increases cyanide evolution to 0.5–1.5 wt%.

See related METAL CYANIDES

0533. Lithium carbonate

[554-13-2]

CLi₂O₃



HCS 1980, 594

Fluorine


See Fluorine: Metal salts

See other INORGANIC BASES, METAL OXONON-METALLATES

0534. Magnesium carbonate hydroxide

[39409-82-0]

CMgO₃· $\frac{1}{4}$ H₂MgO₂



Formaldehyde

See Formaldehyde: Magnesium carbonate hydroxide

See other INORGANIC BASES, METAL OXONON-METALLATES

0535. Carbonyl(pentasulfur pentanitrido)molybdenum

[]

CMoN₅OS₅



Wynne, K. J. *et al.*, *J. Inorg. Nucl. Chem.*, 1968, **30**, 2853

A shock- and heat-sensitive amorphous compound of uncertain structure. A sample exploded violently when scraped from a glass sinter, under impact from a hammer, on sudden heating to 260°C or when added to conc. sulfuric acid.

See related CARBONYLMETALS, N-S COMPOUNDS

0536. Sodium cyanide

[143-33-9]



CNNa

HCS 1980, 836, 837 (30% solution)

Benzyl chloride

See Benzyl chloride: Methanol, Sodium cyanide, etc.

Ethyl chloroacetate

See Ethyl chloroacetate: Sodium cyanide

Oxidants

See entry METAL CYANIDES (AND CYANO COMPLEXES) (reference 1)

0537. Sodium fulminate

[15736-98-8]



CNNaO

Smith, 1966, Vol. 2, 99

Even sodium fulminate detonates when touched lightly with a glass rod.

See other METAL FULMINATES

0538. Sodium thiocyanate

[540-72-7]



CNNaS

Other reactants

Yoshida, 1980, 229

MRH values calculated for 14 combinations with a variety of reagents are given.

Sodium nitrate

See Sodium nitrate: Sodium thiocyanate

Sodium nitrite

MRH 3.51/72

See Sodium nitrite: Sodium thiocyanate

Sulfuric acid

See Sulfuric acid: Sodium thiocyanate

See related METAL CYANIDES

0539. Thallium fulminate
[20991-79-1]

CNOTI



Boddington, T. *et al.*, *Trans. Faraday Soc.*, 1969, **65**, 509
Explosive, even more shock- and heat-sensitive than mercury(I) fulminate.
See other METAL FULMINATES

0540. Cyanonitrene
[1884-64-6]

CN₂



Hutchins, M. G. K. *et al.*, *Tetrahedron Lett.*, 1981, **22**, 4599–4602
Interaction of sodium hydrogen cyanamide and *tert*-butyl hypochlorite in methanol at -50°C to ambient temperature forms cyanonitrene. Isolation gave an orange dimeric residue which exploded on two occasions.
See related CYANO COMPOUNDS

0541. Nitrosyl cyanide
[4343-68-4]

CN₂O



Alone, or Nitrogen oxide

1. Kirby, G. W., *Chem. Soc. Rev.*, 1977, 9
2. Keary, C. M. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1978, 243

Direct preparation of the gas is potentially hazardous, and explosive decomposition of the impure gas in the condensed state (below -20°C) has occurred. A safe procedure involving isolation of the 1:1 adduct with 9,10-dimethylanthracene is preferred. The impure gas contains nitrogen oxide and it is known that nitrosyl cyanide will react with the latter to form an explosive compound [1]. The need to handle this compound of high explosion risk in small quantities, avoiding condensed states, is stressed [2].

See other CYANO COMPOUNDS, NITROSO COMPOUNDS

0542. Sulfinylcyanamide
[16438-87-3]

CN₂OS



Alone, or Ethanol, or Water

Scherer, O. J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1967, **6**, 701–702

The crude highly unsaturated material decomposes explosively if stored in a refrigerator or on attempted distillation under water-pump vacuum (moisture?). However, it is distillable under oil-pump vacuum and is then stable in refrigerated storage. It decomposes explosively with traces of water or ethanol, and polymerises in daylight at ambient temperature.

See other CYANO COMPOUNDS, N-S COMPOUNDS

0543. Thallium(I) azidodithiocarbonate (Thallium(I) dithiocarbonazidate)
[] **CN₃S₂Tl**



Sulfuric acid

Bailar, 1973, Vol. 1, 1155

The highly unstable explosive salt is initiated by contact with sulfuric acid.

See Azidodithioformic acid

See related ACYL AZIDES

0544. Cyanogen azide

[764-05-6]

CN₄



1. Marsh, F. D., *J. Amer. Chem. Soc.*, 1964, **86**, 4506

2. Marsh, F. D., *J. Org. Chem.*, 1972, **37**, 2966

3. Coppolino, A. P., *Chem. Eng. News*, 1974, **52**(25), 3

4. Mardell, L., *Chem. in Brit.*, 1995, **31**(8), 591

It detonates with great violence when subjected to mild mechanical, thermal or electrical shock [1]. Special precautions are necessary to prevent separation of the azide from solution and to decontaminate equipment and materials [2]. The azide is not safe in storage at -20°C as previously stated. A sample exploded immediately after removal from refrigerated (dark) storage [3]. A student was performing a preparation involving in situ generation of cyanogen azide in acetonitrile. Some unexpected white needles crystallised from the mixture, were filtered, dried on the filter (probably a glass sinter) and scraped off, this caused an explosion removing the student's thumb. It is hypothesised that water contamination, or excess sodium azide, had induced formation of diazidomethylenecyanamide or sodium 5-azidotetrazolide, which had crystallised [4].

See Diazidomethylenecyanamide

Sodium hydroxide

Marsh, F. D., *J. Org. Chem.*, 1972, **37**, 2966

Interaction with 10% alkali forms sodium 5-azidotetrazolide, which is violently explosive if isolated.


See 5-Azidotetrazole

See related HALOGEN AZIDES

0545. Dinitrodiazomethane

[25240-93-1]

CN₄O₄



Alone, or Sulfuric acid

Schallkopf, V. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1969, **8**, 612

It explodes on impact, rapid heating or contact with sulfuric acid.
See other DIAZO COMPOUNDS, POLYNITROALKYL COMPOUNDS

0546. Tetranitromethane
[509-14-8]

CN₄O₈



1. Liang, P., *Org. Synth.*, 1955, Coll. Vol. 3, 804
2. *MCA Guide*, 1972, 310 (reference 9)

During its preparation from fuming nitric acid and acetic anhydride, strict temperature control and rate of addition of anhydride are essential to prevent a runaway violent reaction [1]. An explosion occurred during preparation in a steel tank [2]. It should not be distilled, as explosive decomposition may occur [1].
See Nitric acid: Acetic anhydride

Aluminium

See Aluminium: Oxidants

Amines

Gol'binder, A. A., *Chem. Abs.*, 1963, **59**, 9730b

Mixtures of amines, for example aniline, with tetranitromethane (19.5%) ignite in 35–55 s, and will proceed to detonation if the depth of liquid is above a critical value.

Aromatic nitro compounds

1. Urbanski, 1964, Vol. 1, 592
2. Zotov, E. V. *et al.*, *Chem. Abs.*, 1982, **98**, 5965

Mixtures with nitrobenzene, 1- or 4-nitrotoluene, 1,3-dinitrobenzene or 1-nitronaphthalene were found to be high explosives of high sensitivity and detonation velocities [1]. Those with nitrobenzene are spark-detonable [2].

Dienes

Eberson, L. *et al.*, *Acta Chem. Scand.*, 1998, **52**(4), 450

Tetranitromethane adds across various non-conjugated dienes to give cyclic products. The products with norbornadiene and cycloocta-1,5-diene are subject to spontaneous violent decomposition.

Ferrocene

Jaworska-Augustyniak, A., *Transition Met. Chem.*, 1979, **4**, 207–208

Contact of tetranitromethane with ferrocene under various conditions leads to violent explosions, though not if the latter is dissolved in methanol or cyclohexane. Other ferrocene derivatives react violently, but not the dimethyl compound.

Hydrocarbons

1. Stettbacher, A., *Z. Ges. Schiess. Sprengstoffw.*, 1930, **25**, 439
2. Hager, K. F., *Ind. Eng. Chem.*, 1949, **41**, 2168

3. Tschinkel, J. G., *Ind. Eng. Chem.*, 1956, **48**, 732
4. Stettbacher, A., *Tech. Ind. Schweiz. Chem. Ztg.*, 1941, **24**, 265–271
5. Anon., *Chem. Ztg.*, 1920, **44**, 497
6. Zotov, E. V. *et al.*, *Chem. Abs.*, 1980, **93**, 49732, 49734

When mixed with hydrocarbons in approximately stoichiometric proportions, a sensitive highly explosive mixture is produced which needs careful handling [1,2]. The use of such mixtures as rocket propellants has also been investigated [3]. Explosion of only 10 g of a mixture with toluene caused 10 deaths and 20 severe injuries [4]. The mixture contained excess toluene in error [5]. Detonation characteristics of mixtures with benzene, toluene (and nitrobenzene) were studied [6].

Pyridine

Isaacs, N. S. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 2802

Mixtures occasionally have exploded several hours after preparation.

Sodium ethoxide

Macbeth, A. K., *Ber.*, 1913, **46**, 2537–2538

Addition of the last of several portions of the ethoxide solution caused the violent explosion of 30 g of tetranitromethane (possibly involving formation of sodium trinitromethanide).

See Potassium trinitromethanide

Toluene, Cotton



Winderlich, R., *J. Chem. Educ.*, 1950, **27**, 669

A demonstration mixture, to show combustion of cellulose by combined oxygen, exploded with great violence soon after ignition.

See Hydrocarbons, above

See other OXIDANTS, POLYNITROALKYL COMPOUNDS

0547. Sodium 5-nitrotetrazolide [67312-43-0]




1. Spear, R. J., *Aust. J. Chem.*, 1982, **35**, 10
2. Redman, L. D. *et al.*, *Chem. Abs.*, 1984, **100**, 194540

Extremely sensitive to mechanical initiation, it should be treated as a primary explosive [1]. A batch of the dihydrate exploded spontaneously [2].

See other C-NITRO COMPOUNDS, TETRAZOLES

0548. 5-Diazoniotetrazolide
[13101-58-1]

CN₆




Thiele, J., *Ann.*, 1892, **270**, 60

Conc. aqueous solutions of this internal diazonium salt explode at 0°C.

See other DIAZONIUM SALTS, TETRAZOLES

0549. Disodium tetrazole-5-diazoate
[]

CN₆Na₂O



Sulfuric acid

Thiele, J., *Ann.*, 1893, **273**, 148–149, 151

It decomposes non-explosively on heating, but ignites in contact with conc. sulfuric acid.

See other TETRAZOLES

0550. Carbonic diazide (Carbonyl azide)
[14435-92-8]

CN₆O



1. Chapman, L. E. *et al.*, *Chem. & Ind.*, 1966, 1266

2. Kesting, W., *Ber.*, 1924, **57**, 1321


3. Curtius, T. *et al.*, *Ber.*, 1894, **27**, 2684

It is a violently explosive solid, which should be used only in solution, and on a small scale [1]. It exploded violently under ice-water [2], or on exposure to bright light [3].

See other ACYL AZIDES, IRRADIATION DECOMPOSITION INCIDENTS

0551. Sodium 5-azidotetrazolide
[35038-45-0]

CN₇Na



Marsh, F. D., *J. Org. Chem.*, 1972, **37**, 2967, 2969

Readily formed from cyanogen bromide and sodium azide in aqueous solution at 0°C, it is extremely sensitive to friction, heat or pressure. A dry sample under vacuum at 1 mbar will detonate on rapid admission of air.

See 5-Azidotetrazole

See other FRICTIONAL INITIATION INCIDENTS *See other* HIGH-NITROGEN COMPOUNDS, *N*-METAL DERIVATIVES, TETRAZOLES

0552. Sodium carbonate
[497-19-8]

CNa₂O₃

NaOCO.ONa

HCS 1980, 832

Acidic reaction liquor

Anon., *Loss Prev. Bull.*, 1978, (023), 138

Use of sodium carbonate to neutralise the acid produced during methylation with dimethyl sulfate in a 2-phase system led to severe frothing, and the agitator was turned off to allow it to abate. When agitation was restarted, violent foaming from mixing of the separated acid and alkaline layers ejected 2 t of the reactor contents (remedy worse than ailment!) Use of magnesium oxide to neutralise acid without gas evolution was recommended.

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS, NEUTRALISATION INCIDENTS

Ammonia, Silver nitrate

See Silver nitrate: Ammonia, etc.

An aromatic amine, A chloronitro compound

MCA Case History No. 1964

An unspecified process had been operated for 20 years using synthetic sodium carbonate powder (soda-ash) to neutralise the hydrogen chloride as it was formed by interaction of the amine and chloro compound in a non-aqueous (and probably flammable) solvent in a steel reactor. Substitution of the powdered sodium carbonate by the crystalline sodium carbonate–sodium hydrogencarbonate double

salt ('trona, natural soda') caused a reduction in the rate of neutralisation, the reaction mixture became more acid, and attack on the steel vessel led to contamination by iron. These changed conditions initiated exothermic side reactions, which eventually ran out of control and caused failure of the reactor. Subsequent laboratory work confirmed this sequence and showed that presence of dissolved iron(III) was necessary to catalyse the side reactions.

See Sulfuric acid, below

See other CATALYTIC IMPURITY INCIDENTS, NEUTRALISATION INCIDENTS

2,4-Dinitrotoluene

See 2,4-Dinitrotoluene: Alkali

Fluorine

See Fluorine: Metal salts

Hydrogen peroxide

See CRYSTALLINE HYDROGEN PEROXIDATES

Lithium

See Lithium: Sodium carbonate

Phosphorus pentaoxide

See Tetraphosphorus decaoxide: Inorganic bases

Sodium sulfide, Water

See SMELT: Water

Sulfuric acid

MCA Case History No. 888

Lack of any mixing arrangements caused stratification of strong sulfuric acid and (probably) sodium carbonate solutions in the same tank. When gas evolution caused intermixing of the layers, a violent eruption of the tank contents occurred.

See An aromatic amine, etc., above

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

2,4,6-Trinitrotoluene

See 2,4,6-Trinitrotoluene: Added impurities

See other INORGANIC BASES, METAL OXONON-METALLATES

0553. Sodium carbonate hydrogen peroxidate ('Sodium percarbonate')

[15630-89-4]

CNa₂O₃·1.5H₂O₂

NaOCO.ONa.1.5H₂O₂

See entry CRYSTALLINE HYDROGEN PEROXIDATES

0554. Sodium monoperoxy carbonate

[4452-58-8]

NaOCO.OONa

CNa₂O₄

Acetic anhydride

See Sodium carbonate hydrogen peroxidate: Acetic anhydride

See other PEROXOACID SALTS

†0555. Carbon monoxide

[630-08-0]

CO

CO

NSC 415, 1976; FPA H57, 1977 (cylinder); HCS 1980, 282

Aluminium, Aluminium halides

See Aluminium: Aluminium halides, Carbon oxides

Copper(I) perchlorate

See Copper(I) perchlorate, Alkenes, etc.

Dinitrogen oxide

See Dinitrogen oxide: Carbon monoxide

Fluorine, Oxygen

See Bis(fluoroformyl) peroxide

Hydrogen, Oxygen

See Oxygen: Carbon monoxide, Hydrogen

Interhalogens

MRH values show % of interhalogen

See Bromine pentafluoride: Acids, etc.

MRH 2.47/71

Bromine trifluoride: Carbon monoxide

MRH 3.05/tr.

Iodine heptafluoride: Carbon, etc.

Metal oxides

See Caesium oxide: Halogens, etc.

Iron(III) oxide: Carbon monoxide

Silver(I) oxide: Carbon monoxide

Metals

See Potassium: Non-metal oxides

Sodium: Non-metal oxides

Other reactants

Yoshida, 1980, 42

MRH values calculated for 11 combinations with oxidants are given.

Oxidants

See Chlorine dioxide: Carbon monoxide

Oxygen (Liquid): Liquefied gases

Peroxodisulfuryl difluoride: Carbon monoxide

See other NON-METAL OXIDES

†0556. Carbonyl sulfide (Carbon oxide sulfide)
[463-58-1]

O:C:S

COS

See related NON-METAL OXIDES, NON-METAL SULFIDES

0557. Carbon dioxide
[124-38-9]

O:C:O

CO₂

NSC 682, 1980; HCS 1980, 279 (cylinder) 280 (solid)

Acrylaldehyde

See Acrylaldehyde: Acids

Aluminium, Aluminium halides

See Aluminium: Aluminium halides, Carbon oxides

Aziridine

See Aziridine: Acids

Barium peroxide

See Barium peroxide: Non-metal oxides

Caesium oxide

See Caesium oxide: Halogens, etc.

Flammable materials

1. *BCISC Quart. Safety Summ.*, 1973, **44**(174), 10
2. Leonard, J. T. *et al.*, *Inst. Phys. Conf. Ser.*, 1975, **27** (Static Electrification), 301–310
3. Anon., *Chem. Eng. News*, 1976, **54**(30), 18
4. Collocott, S. J. *et al.*, *IEE Proc. part A*, 1980, **127**, 119–120
5. Moffat, M., *Fire Prot.*, 1980, (7), 9
6. Kletz, T. A., *Loss Prev. Bull.*, 1984, (056), 9

Dangers attached to the use of carbon dioxide in fire prevention and extinguishing systems in confined volumes of air and flammable vapours are discussed. The main hazard arises from the production of very high electrostatic charges, due to the presence of small frozen particles of carbon dioxide generated during expanding discharge of the compressed gas. Highly incendive sparks (5–15 mJ at 10–20 kV) readily may be produced. Nitrogen is preferred as an inerting gas for enclosed

volumes of air and flammable gases or vapours [1]. The electrostatic hazard in discharge of carbon dioxide in tank fire-extinguishing systems is associated with the plastics distribution horns, and their removal eliminated the hazard [2]. Alternatively, a metal sleeve may be fitted inside the plastics horn [3]. The accumulation of charge during discharge of a portable extinguisher has also been studied and found potentially hazardous [4]. The latter hazard is minimised in a statement to clarify the overall situation, which emphasizes the need for avoiding the use of liquid carbon dioxide for inerting purposes [5]. Previous accidents involving carbon dioxide as the cause of explosions are reviewed [6].

Hydrazine, Stainless steel

See Hydrazine: Carbon dioxide, Stainless steel

Metal hydrides

See Aluminium hydride: Carbon dioxide

Lithium tetrahydroaluminate: Bis(2-methoxyethyl) ether

Metal acetylides

See Monolithium acetylide–ammonia: Gases

Monopotassium acetylide: Non-metal oxides

Rubidium acetylide: Non-metal oxides

Sodium acetylide: Non-metal oxides

Metals

1. Hartmann, I., *Ind. Eng. Chem.*, 1948, **40**, 756

2. Rhein, R. A., *Rept. No. CR-60125*, Washington, NASA, 1964

Dusts of magnesium, zirconium, titanium and some magnesium–aluminium alloys [1], and (when heated) of aluminium, chromium and manganese [2], when suspended in carbon dioxide atmospheres are ignitable and explosive, and several bulk metals will burn in the gas.

See Metals, Nitrogen below

Aluminium: Aluminium halides, Carbon oxides

Lithium: Non-metal oxides

Magnesium: Carbon dioxide, etc.

Potassium: Non-metal oxides

Potassium–sodium alloy: Carbon dioxide

Sodium: Non-metal oxides

Titanium: Carbon dioxide

Uranium: Carbon dioxide

Metals, Nitrogen

Rhein, R. A., *Rept. No. CR-60125*, Washington, NASA, 1964

Powdered beryllium, cerium, cerium alloys, thorium, titanium, uranium and zirconium ignite on heating in mixtures of carbon dioxide and nitrogen; ignition temperatures were determined. Aluminium, chromium, magnesium and manganese ignite only in the absence of nitrogen.

Metals, Sodium peroxide
See Sodium peroxide: Metals, Carbon dioxide
See other NON-METAL OXIDES

0558. Lead carbonate
[598-63-0]



CO₃Pb

HCS 198, 586

Fluorine
See Fluorine: Metal salts

0559. Carbon sulfide
[2944-05-0]



CS

Pearson, T. G., *School Sci. Rev.*, 1938, **20**(78), 189
The gaseous radical polymerises explosively.
See other NON-METAL SULFIDES

†0560. Carbon disulfide
[75-15-0]



CS₂

(*MCA SD-12*, 1967); *NSC 341*, 1977; *FPA H34*, 1975; *HCS 1980*, 281; *RSC Lab. Hazard Data Sheet No. 7*, 1982

1. Thiem, F., *Chem. Technik*, 1979, **31**, 313–314
2. Briggs, A. G. *et al.*, *Chem. Brit.*, 1982, **18**, 27
3. Bothe, H. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries* Vol. II (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 327, Amsterdam, Elsevier, 1995

Carbon disulfide vapour, alone or mixed with nitrogen, did not decompose explosively in the range 0.4–2.1 bar/88–142°C when initiated with high energy sparks or a hot wire at 700–900°C. The endothermic sulfide (ΔH_f° (l) +115.3 kJ/mol, 1.52 kJ/g) will, however, decompose explosively to its elements with mercury fulminate initiation [1]. A screening jacket filled with carbon disulfide was used to surround the reaction tube used in flash photolysis experiments. When the quartz lamp was discharged, some vapour of the disulfide which had leaked out ignited in the radiation flash and exploded. Care is also necessary with the vapour and other UV sources [2]. Ignition of vapour air mixes by laser has been studied [3].
See other IRRADIATION DECOMPOSITION INCIDENTS

Air, Rust

1. Mee, A. J., *School Sci. Rev.*, 1940, **22**(86), 95
2. Dickens, G. A., *School Sci. Rev.*, 1950, **31**(114), 264

3. Anon., *J. Roy. Inst. Chem.*, 1956, **80**, 664

4. Thompson, E., private comm., 1981

Disposal of 2 l of the solvent into a rusted iron sewer caused an explosion. Initiation of the solvent–air mixture by rust was suspected [1]. A hot gauze falling from a tripod into a laboratory sink containing some carbon disulfide initiated two explosions [2]. It is a very hazardous solvent because of its extreme volatility and flammability. The vapour or liquid has been known to ignite on contact with steam pipes, particularly if rusted [3]. When a winchester of the solvent fell off a high shelf and broke behind a rusted steel cupboard, ignition occurred [4].

Alkali metals

1. Steimecke, G. *et al.*, *Phos. Sulfur*, 1979, **7**, 49

2. Moradpour, A. *et al.*, *J. Chem. Educ.*, 1986, **62**, 1016

3. Staudinger, H., *Z. Angew. Chem.*, 1922, **35**, 659

Attempts to follow a published procedure for the preparation of 1,3-dithiole-2-thione-4,5-dithiolate salts [1], involving reductive coupling of carbon disulfide with alkali metals, have led to violent explosions with potassium metal, but not with sodium [2]. However, mixtures of carbon disulfide with potassium–sodium alloy, potassium, sodium, or lithium are capable of detonation by shock, though not by heating. The explosive power decreases in the order given above, and the first mixture is more shock-sensitive than mercury fulminate [3].

See Metals, below

Amines

1. Anon., Personal communication, 1991

2. Editor's comments.

Carbon disulphide was used as an extraction solvent when analysing epoxy resins. On one occasion, adding to a hardener produced a vigorous fume-off leaving a residue looking like sulfur [1]. Amines and complexes thereof are used as hardeners, and the reaction with, especially, polyamines to give dithiocarbamates is surprisingly exothermic [2].

Chlorine, Sodium *aci*-nitromethanide

See Sodium *aci*-nitromethanide: Carbon disulfide, etc.

Halogens

See Chlorine: Carbon disulfide

Fluorine: Sulfides

Metal azides

Mellor, 1947, Vol. 8, 338

Carbon disulfide and aqueous solutions of metal azides interact to produce metal azidodithioformates, most of which are explosive with varying degrees of power and sensitivity to shock or heat.

See Azidodithioformic acid, *also* Sodium azide: Carbon disulfide

Metals

MRH Zinc 2.43/63

See Alkali metals, above
Aluminium: Carbon disulfide
Sodium: Sulfides, etc.
Zinc: Carbon disulfide

Other reactants

Yoshida, 1980, 281
MRH values calculated for 16 combinations, largely with oxidants, are given.

Oxidants

See Halogens, above
Dinitrogen tetraoxide: Carbon disulfide
Nitrogen oxide: Carbon disulfide
Permanganic acid: Organic materials

Phenylcopper–triphenylphosphine complexes

Camus, A. *et al.*, *J. Organomet. Chem.*, 1980, **188**, 390
The bis- or tris-complexes of phenylcopper with triphenylphosphine react violently and exothermically with carbon disulfide, even at 0°C. Suitable control procedures are described.
See other ENDOTHERMIC COMPOUNDS, NON-METAL SULFIDES

0561. Titanium carbide

[12070-08-5]

TiC

CTi

MCA Case History No. 618

A violent, and apparently spontaneous, dust explosion occurred while the finely ground carbide was being removed from a ball-mill. Static initiation seems a likely possibility.

See entries DUST EXPLOSION INCIDENTS, REFRACTORY POWDERS

See other METAL NON-METALLIDES

0562. Uranium carbide

[12070-09-6]

CU

CU

Schmitt, C. R., *J. Fire Flamm.*, 1971, **2**, 163

The finely divided carbide is pyrophoric.

See other METAL NON-METALLIDES, PYROPHORIC MATERIALS

0563. Tungsten carbide
[12070-12-1]

WC

CW

Fluorine

See Fluorine: Metal acetylides and carbides

Nitrogen oxides

Mellor, 1946, Vol. 5, 890

At about 600°C, the carbide ignites and incandesces in dinitrogen mono- or tetra-oxides.

See other METAL NON-METALLIDES

0564. Ditungsten carbide
[12070-13-2]

W₂C

CW₂

Oxidants

Mellor, 1946, Vol. 5, 890

The carbide burns incandescently at red heat in contact with dinitrogen mono- or tetra-oxides.

See Fluorine: Metal acetylides and carbides

See other METAL NON-METALLIDES

0565. Zirconium carbide
[12070-14-3]

ZrC

CZr

See entry REFRACTORY POWDERS

0566. Silver chloroacetylide
[]

AgC≡CCl

C₂AgCl

Kirk-Othmer, 1964, Vol. 5, 204

Explosive, even when wet or under water.

See related HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

See other SILVER COMPOUNDS

0567. Silver cyanodinitromethanide
[12281-65-1]

AgC(CN)(NO₂)₂

C₂AgN₃O₄

Parker, C. O. *et al.*, *Tetrahedron*, 1962, **17**, 86

A sample in a m.p. capillary exploded at 196°C, shattering the apparatus.
See other POLY-NITROALKYL COMPOUNDS, SILVER COMPOUNDS

0568. Silver acetylide
[7659-31-6]




1. Miller, 1965, Vol. 1, 486
2. Bailar, 1973, Vol. 3, 102
3. Reynolds, R. J. *et al.*, *Analyst*, 1971, **96**, 319
4. Koehn, J., *Chem Abs.*, 1979, **91**, 7164

Silver acetylide is a more powerful detonator than the copper derivative, but both will initiate explosive acetylene-containing gas mixtures [1]. It decomposes violently when heated to 120–140°C [2]. Formation of a deposit of this explosive material was observed when silver-containing solutions were aspirated into an acetylene-fuelled atomic absorption spectrometer. Precautions to prevent formation are discussed [3]. The effect of ageing for 16 months on the explosive properties of silver and copper acetylides has been studied. Both retain their hazardous properties for many months, and the former is the more effective in initiating acetylene explosions [4].

See other METAL ACETYLIDES, SILVER COMPOUNDS

0569. Silver acetylide–silver nitrate
[15336-58-0]




1. Baker, W. E. *et al.*, *Chem. Eng. News*, 1965, **43**(49), 46
2. Hogan, V. D. *et al.*, *OTS Rept. AD 419 625*, Washington, US Dept. Comm., 1960
3. Benham, R. A., *Chem. Abs.*, 1981, **94**, 142109

The dry complex is exploded by high-intensity light pulses, or by heat or sparks. As a slurry in acetone, it is stable for a week if kept dark [1]. The complex explodes violently at 245°C/1 bar and at 195°C/1.3 mbar [2], or on exposure to light [3].

See related METAL ACETYLIDES

See other IRRADIATION DECOMPOSITION INCIDENTS, SILVER COMPOUNDS

0570. Disilver ketenide
[27278-01-4]





1. Blues, E. T. *et al.*, *Chem. Comm.*, 1970, 699
2. Bryce-Smith, D. *et al.*, *J. Chem. Soc., Perk. Trans. 2*, 1993, (9), 1631.

This and its pyridine complex explode violently if heated or struck [1] (but see below). More data on the stability of disilver ketenide and its complexes is contained in [2].

See other SILVER COMPOUNDS

0571. Disilver ketenide–silver nitrate

[]



1. Bryce-Smith, D., *Chem. & Ind.*, 1975, 154
2. Blues, E. T. *et al.*, *Chem. Comm.*, 1970, 701

The red complex is more dangerously explosive than the ketenide itself, now described as mildly explosive in the dry state when heated strongly or struck [1]. It was previously formulated as a 2:1 complex [2].

See other SILVER COMPOUNDS

0572. Silver oxalate

[533-51-7]



1. Sidgwick, 1950, 126
2. Anon., *BCISC Quart. Safety Summ.*, 1973, **44**, 19
3. Kabanov, A. A. *et al.*, *Russ. Chem. Rev.*, 1975, **44**, 538
4. MacDonald, J. Y. *et al.*, *J. Chem. Soc.*, 1925, **127**, 2625
5. Waltenberger-Razniewska, M. *et al.*, *Chem. Abs.*, 1982, **96**, 203117
6. Ullmann, 1993, Vol. A24, 161

Above 140°C its exothermic decomposition to metal and carbon dioxide readily becomes explosive [1]. A 1 kg batch which had been thoroughly dried at 50°C exploded violently when mechanical grinding in an end-runner mill was attempted [2]. Explosions have been experienced when drying the oxalate as low as 80°C [6]. It is a compound of zero oxygen balance. The explosion temperature of the pure oxalate is lowered appreciably (from 143 to 122°C) by application of an electric field [3]. The salt prepared from silver nitrate with excess of sodium oxalate is much less stable than that from excess nitrate [4]. Decomposition at 125°C in glycerol prevents explosion in the preparation of silver powder [5].

See ELECTRIC FIELDS

See other METAL OXALATES, SILVER COMPOUNDS

0573. Gold(I) acetylide

[70950-00-4]



1. Mellor, 1946, Vol. 5, 855
2. Matthews, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1900, **22**, 110


Produced by action of acetylene on sodium gold thiosulfate (or other gold salts), the acetylides are explosive and readily initiated by light impact, friction or rapid heating to 83°C [1]. This unstable detonator is noted for high brisance [2].

See other GOLD COMPOUNDS, METAL ACETYLIDES

0574. Digold(I) ketenide

[54086-41-8]

C_2Au_2O



Blues, E. T. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1974, 513–514

The ketenide is shock-sensitive when dry, and it or its complexes with tertiary heterocyclic bases (pyridine, methylpyridines, 2,6-dimethylpyridine or quinoline) explode when heated above 100°C.

See other GOLD COMPOUNDS

0575. Barium acetylide

[12070-27-8]

$(C_2Ba)_n$



Air, or Hydrogen, or Hydroxylic compounds

Masdupay, E. *et al.*, *Compt. rend.*, 1951, **232**, 1837–1839

It is much more reactive than the diacetylide, and ignites in contact with water or ethanol in air. It may incandesce on heating to 150°C under vacuum or hydrogen, the product from the latter treatment being very pyrophoric owing to the presence of pyrophoric carbon.

Halogens

Mellor, 1946, Vol. 5, 862

Barium acetylide incandesces with chlorine, bromine or iodine at 140, 130 or 122°C respectively.

Selenium

Mellor, 1946, Vol. 5, 862

A mixture incandesces when heated to 150°C.

See other METAL ACETYLIDES

0576. Barium thiocyanate

[2092-17-3]

$C_2BaN_2S_2$



Potassium chlorate

See Potassium chlorate: Metal thiocyanates


Sodium nitrite

See Sodium nitrite: Potassium thiocyanate

See related METAL CYANIDES

0577. Barium 5,5'-azotetrazolide
[87478-73-7]

C₂BaN₁₀



See entry METAL AZOTETRAZOLIDES *See other* TETRAZOLES

0578. Bromochloroacetylene
[25604-70-0]

C₂BrCl




Houben-Weyl, 1977, Vol. 5.2a, 604 (footnote 6)

Highly explosive and unstable.

See other HALOACETYLENE DERIVATIVES

†**0579. Bromotrifluoroethylene**
[598-73-2]

C₂BrF₃




Oxygen

See Oxygen (Gas): Halocarbons

See other HALOALKENES

0580. Lithium bromoacetylide
[]

C₂BrLi



See Sodium bromoacetylide (next below); Lithium chloroacetylide; Lithium trifluoropropynide; Sodium chloroacetylide

See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

0581. Sodium bromoacetylide
[]

C₂BrNa



Cradock, S. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1978, 760

The dry material (perhaps the ultimate 'carbene precursor') is an extremely shock-sensitive explosive.

See Lithium bromoacetylide; Lithium chloroacetylide; Lithium trifluoropropynide; Sodium chloroacetylide

See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

0582. Dibromoacetylene
[624-61-3]



Rodd, 1951, Vol. 1A, 284
It ignites in air and explodes on heating.
See other HALOACETYLENE DERIVATIVES

0583. Oxalyl dibromide
[15219-34-8]




Potassium
See Potassium: Oxalyl dihalides
See other ACYL HALIDES

0584. Hexabromoethane
[594-73-0]



Hexacyclohexyldilead
See Hexacyclohexyldilead: Halocarbons
See other HALOALKANES

0585. Calcium acetylide (Calcium carbide)
[75-20-7]



HCS 1980, 264

1. Anon., *Ind. Safety Bull.*, 1940, **8**, 41
2. Déribère, M., *Chem. Abs.*, 1936, **30**, 8169.8
3. Orlov, V. N., *Chem. Abs.*, 1975, **83**, 168138

Use of a steel chisel to open a drum of carbide caused an incendive spark which ignited traces of acetylene in the drum. The non-ferrous tools normally used for this purpose should be kept free from embedded ferrous particles [1]. If calcium carbide is warm when filled into drums, absorption of the nitrogen from the trapped air may enrich the oxygen content up to 28%. In this case, less than 3% of acetylene (liberated by moisture) is enough to form an explosive mixture, which may be initiated on opening the sealed drum. Other precautions are detailed [2]. Use of carbon dioxide to purge carbide drums, and of brass or bronze non-sparking tools to open them are advocated [3].

Calcium hypochlorite
See Calcium hypochlorite

Halogens

Mellor, 1946, Vol. 5, 862

The acetylide incandesces with chlorine, bromine or iodine at 245, 350 or 305°C, respectively. Strontium and barium acetylides are more reactive.

Hydrogen chloride

MRH 3.97/54

Mellor, 1946, Vol. 5, 862

Incandescence on warming; strontium and barium acetylides are similar.

Iron(III) chloride, Iron(III) oxide

Partington, 1967, 372

The carbide is an energetic reductant. A powdered mixture with iron oxide and chloride burns violently when ignited, producing molten iron.

See other THERMITE REACTIONS

Lead difluoride

Mellor, 1946, Vol. 5, 864

Incandescence on contact at ambient temperature.

Magnesium

Mellor, 1940, Vol. 4, 271

A mixture incandesces when heated in air.

Methanol

Unpublished observations, 1951

Interaction of calcium carbide with methanol to give calcium methoxide is very vigorous, but subject to an induction period of variable length. Once reaction starts, evolution of acetylene gas is very fast, and a wide-bore condenser and adequate ventilation are necessary.

See other GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS

Other reactants

Yoshida, 1980, 223

MRH values calculated for 8 combinations with various reagents are given.

Perchloryl fluoride

See Perchloryl fluoride: Calcium acetylide

Selenium

See Selenium: Metal acetylides

Silver nitrate

Luchs, J. K., *Photog. Sci. Eng.*, 1966, **10**, 334

Addition of calcium acetylide to silver nitrate solution precipitates silver acetylide, a highly sensitive explosive. Copper salt solutions would behave similarly.

See METAL ACETYLIDES

Sodium peroxide MRH 3.01/86
See Sodium peroxide: Calcium acetylide

Sulfur MRH 4.35/33
See Sulfur: Metal acetylides

Tin(II) chloride
Mellor, 1941, Vol. 7, 430
A mixture can be ignited with a match, and reduction to metallic tin proceeds with incandescence.

Water
1. Jones, G. W. *et al.*, *Explosions in Med. Press. Generators, Invest. Rept.* 3775, Washington, US Bur. Mines, 1944
2. Moll, H., *Chem. Weekbl.*, 1933, **30**, 108
At the end of a generation run, maximum temperature and high moisture content of acetylene may cause the finely-divided acetylide to overheat and initiate explosion of pressurised gas [1]. During analysis of technical carbide by addition of water, the explosive mixture formed in the unpurged reaction vessel exploded, ignited either by excessive local temperature or possibly by formation of crude phosphine from the phosphide present as impurity in the carbide [2].
See other METAL ACETYLIDES, REDUCANTS

0586. Bis(trifluoromethyl)cadmium **C₂CdF₆**
[3327-66-1] **F₃CCdCF₃**

Eujen, R. *et al.*, *J. Organomet. Chem.*, 1995, **503**(2), C51
The uncomplexed compound decomposes to difluorocarbene at -5° and explodes violently on warming to room temperature.
See METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

0587. Bis(trifluoromethyl)cadmium-1,2-dimethoxyethane adduct **C₂CdF₆.C₄H₁₀O₂**
[33327-66-1] **(F₃C)₂Cd.glyme**

Ontiveros, C. D. *et al.*, *Inorg. Synth.*, 1986, **24**, 57
It may ignite in air if not thoroughly dry and pure.
See related ALKYL METALS

0588. Cadmium cyanide **C₂CdN₂**
[542-83-6] **Cd(C≡N)₂**

It is moderately endothermic (ΔH_f° (s) +163.2 kJ/mol, 0.99 kJ/g)

Magnesium

See Magnesium: Metal cyanides

See other ENDOTHERMIC COMPOUNDS, METAL CYANIDES

0589. Cadmium fulminate

[42294-95-6]

$C_2CdN_2O_2$



It is a moderately endothermic compound (ΔH_f° (s) +163.2 kJ/mol, 0.80 kJ/g).

See entries ENDOTHERMIC COMPOUNDS, METAL FULMINATES

0590. Cerium carbide

[12012-32-7]

C_2Ce




See Lanthanum carbide

†0591. Chlorotrifluoroethylene

[79-38-9]

C_2ClF_3



Bromine, Oxygen

See Bromine: Chlorotrifluoroethylene, Oxygen

Chlorine perchlorate

See Chlorine perchlorate: Chlorotrifluoroethylene

1,1-Dichloroethylene

See 1,1-Dichloroethylene: Chlorotrifluoroethylene

Ethylene

See Ethylene: Chlorotrifluoroethylene

Oxygen


See Oxygen (Gas): Halocarbons (references 3,4)

See other HALOALKENES

0592. Poly(chlorotrifluoroethylene)

[9002-83-9]

$(C_2ClF_3)_n$




Aluminium

See Aluminium: Halocarbons

See related HALOALKANES

0593. 3-Chloro-3-trifluoromethyldiazirine
[58911-30-1]

$C_2ClF_3N_2$



Moss, R. A. *et al.*, *J. Amer. Chem. Soc.*, 1981, **103**, 6168–6169

The potentially explosive compound was used safely as a solution in pentane.

See other DIAZIRINES

0594. Trifluoroacetyl hypochlorite
[65597-25-3]

$C_2ClF_3O_2$

$F_3CCO.OCl$

1. Tari, I. *et al.*, *Inorg. Chem.*, 1979, **18**, 3205–3208

2. Schack, C. J. *et al.*, US Pat. Appl. 47,588, 1979

3. Gard, G. L. *et al.*, *Inorg. Chem.*, 1965, **4**, 594

4. Rozen, S. *et al.*, *J. Org. Chem.*, 1980, **45**, 677

Thermally unstable at 22°C and explosive in the gas phase at pressures above 27–62 mbar [1], there is also an explosion hazard during distillation and trapping at –78°C [2]. The crude (80%) material prepared by action of 3% fluorine in nitrogen on sodium trifluoroacetate [3] has, however, been used synthetically for 3 years without mishap [4].

See other ACYL HYPOHALITES

0595. 2-Chloro-1,1-bis(fluorooxy)trifluoroethane
[72985-56-9]

$C_2ClF_5O_2$

$ClCF_2CF(OF)_2$

Sekiya, A. *et al.*, *Inorg. Chem.*, 1980, **19**, 1330

Extremely explosive, detonation of a 1 mmol sample in a vacuum line led to 'impressive damage'.

See other BIS(FLUOROOXY)PERHALOALKANES

0596. N-Chlorobis(trifluoromethanesulfonyl)imide
[91742-17-5]

$C_2ClF_6NO_4S_2$

$ClN(O_2SCF_3)_2$

Foropoulos, J. *et al.*, *Inorg. Chem.*, 1984, **23**, 3721

During preparation of the title compound from the imide and chlorine fluoride, initially at –111°C with slow warming during 12 h to 0°C, the scale should be limited to 40 mmol or below. Forceful explosions occurred on several occasions when this scale was exceeded, but care is necessary on any scale of working.

See other N-HALOGEN COMPOUNDS

0597. Bis(trifluoromethyl)chlorophosphine
[650-52-2] **C₂ClF₆P**
 $(F_3C)_2PCl$

491M, 1975, 60
It ignites in air.
See other ALKYLHALOPHOSPHINES

0598. Chloroiodoacetylene (Chloroiodoethyne)
[25604-71-1] **C₂ClI**
 $ClC\equiv CI$

Houben-Weyl, 1977, Vol. 5.2a, 604 (footnote 6)
Highly unstable and explosive.
See other HALOACETYLENE DERIVATIVES

0599. Lithium chloroacetylide (Lithium chloroethynide)
[6180-21-8] **C₂ClLi**
 $LiC\equiv CCl$

1. Cadot, P., *BCISC Quart. Safety Summ.*, 1965, **36**(142), 27
2. Sinden, B. B. *et al.*, *J. Org. Chem.*, 1980, **45**, 2773
During its synthetic use in liquid ammonia, some of the salt separated as a crust owing to evaporation of solvent, and exploded violently. Such salts are stable in solution, but dangerous in the solid state. Evaporation of ammonia must be prevented or made good until unreacted salt has been decomposed by addition of ammonium chloride [1]. An improved preparation of the lithium derivative is available [2].
See Lithium bromoacetylide; Lithium trifluoropropynide; Sodium bromoacetylide; Sodium chloroacetylide

Methyl chloroformate
Acheson, R. M. *et al.*, *J. Chem. Res.; Synop.*, 1986, (10), 378–379
An attempt to prepare methyl chloropropiolate from chloroethynyllithium and methyl chloroformate led to an explosion.
See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

0600. Cyanoformyl chloride
[4474-17-3] **C₂ClNO**
 $N\equiv CCO.Cl$

Water
Appel, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1983, **22**, 785
Reaction with water is violent, forming triazinetricarboxylic acid.
See other ACYL HALIDES, CYANO COMPOUNDS

0601. Sodium chloroacetylide (Sodium chloroethynide)

[]

 C_2ClNa Viehe, H. G., *Chem. Ber.*, 1959, **92**, 1271

Though stable and usable in solution, the sodium salt, like the lithium and calcium salts, is dangerously explosive in the solid state.

See Lithium bromoacetylide; Lithium chloroacetylide; Sodium bromoacetylide; Lithium trifluoropropynide

See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

0602. Dichloroacetylene (Dichloroethyne)

[7572-29-4]

 C_2Cl_2 

1. Wotiz, J. H. *et al.*, *J. Org. Chem.*, 1961, **26**, 1626
2. Ott, E., *Ber.*, 1942, **75**, 1517
3. Kirk-Othmer, 1964, Vol. 5, 203–205
4. Siegel, J. *et al.*, *J. Org. Chem.*, 1970, **35**, 3199
5. Riemschneider, R. *et al.*, *Ann.*, 1961, **640**, 14
6. *MCA Case History No. 1989*
7. Kende, A. S. *et al.*, *Synthesis*, 1982, 455
8. Kende, A. S. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 2369, 2373
9. Denis, J. N. *et al.*, *J. Org. Chem.*, 1987, **52**, 3461–3462

Dichloroacetylene is rather endothermic (ΔH_f° (g) +149.4 kJ/mol, 1.57 kJ/g) and a heat-sensitive explosive gas which ignites in contact with air. However, its azeotrope with diethyl ether (55.4% dichloroacetylene) is not explosive and is stable to air [1,2]. It is formed on catalysed contact between acetylene and chlorine, or sodium hypochlorite at low temperature; or by the action of alkali upon polychloro-ethane and -ethylene derivatives, notably trichloroethylene [3]. A safe synthesis has been described [4]. Ignition of a 58 mol% solution in ether on exposure to air of high humidity, and violent explosion of a conc. solution in carbon tetrachloride shortly after exposure to air have been reported. Stirring the ethereal solution with tap water usually caused ignition and explosion [5]. Dichloroacetylene had been collected without incident on 6 previous occasions as a dilute solution. When the cooling system was modified by lowering the temperature in the water separator, liquid chloroacetylene separated there and exploded when a stopcock was turned [6]. A safer low temperature preparation of material free of trichloroethylene has been described [7,8]. A practical high-yielding synthesis of dichloroacetylene involves addition of a catalytic amount of methanol to a mixture of trichloroethylene and potassium hydride in THF [9].

See Tetrachloroethylene: Sodium hydroxide

Trichloroethylene: Alkali, or: Epoxides

See Chlorocyanoacetylene

See other HALOACETYLENE DERIVATIVES

0603. Chlorodifluoroacetyl hypochlorite


[68674-44-2]



Unstable above 22°C and explosive in gas phase at pressures above 27–63 mbar.
See entry ACYL HYPOHALITES (reference 1)

0604. 1,2-Dichlorotetrafluoroethane

[76-14-2]




Aluminium

See Aluminium: Halocarbons

See other HALOALKANES

0605. Oxalyl dichloride (Oxalyl chloride)

[79-37-8]



Alumina

Kropp. P. J. *et al.*, *J. Amer. Chem. Soc.*, 1993, **115**(8), 3071

Vigorous gas evolution (probably CO—Ed.) was produced when oxalyl chloride was adsorbed on alumina.

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides


Potassium

See Potassium: Oxalyl dihalides

See other ACYL HALIDES

0606. 1,1,2-Trichlorotrifluoroethane

[76-13-1]



Metals

See Aluminium: Halocarbons

Barium: Halocarbons

Lithium: Halocarbons

Samarium: 1,1,2-Trichlorotrifluoroethane

Titanium: Halocarbons

See other HALOALKANES

0607. Trichloroacetonitrile
[545-06-2]



Ammonium chloride, Sodium azide
See 5-Trichloromethyltetrazole
See other CYANO COMPOUNDS

0608. Sodium trichloroacetate
[650-51-1]



HCS 1980, 864


1. Doyle, W. H., *Loss Prev.*, 1969, **3**, 15

Energy of decomposition (in range 150–250 °C) was measured as 0.23 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 119°C by adiabatic Dewar tests, with an apparent energy of activation of 140 kJ/mol.

Bags of the salt ignited in storage (cause unknown) [1].

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

0609. Tetrachloroethylene
[127-18-4]



(MCA SD-24, 1971); HCS 1980, 887; RSC Lab. Hazard Data Sheet No. 6, 1982

Aluminium

See Aluminium: Halocarbons

Aluminium, Zinc oxide

Katz, S. *et al.*, *Chem. Abs.*, 1981, **94**, 33106

The mixture is burned militarily to produce dense smoke.

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Halocarbons

Metals

See Barium: Halocarbons

Lithium: Halocarbons

Sodium hydroxide

Mitchell, P. R., private comm., 1973

The presence of 0.5% of trichloroethylene as impurity in tetrachloroethylene during unheated drying over solid sodium hydroxide caused generation of dichloroacetylene. After subsequent fractional distillation, the volatile fore-run exploded.

See Dichloroacetylene; also Trichloroethylene: Alkali
See other HALOALKENES


Zinc

J. A. Young, private communication, 1997

Distillation of tetrachloroethylene (formerly a dry-cleaning solvent) in new galvanised steel equipment produces traces of dichloroacetylene. This is toxic and may cause ill-health in those exposed. Should the acetylene chance to be concentrated (as in the sub-entry above), it is also very explosive. The galvanised metal becomes passivated in a few days and the effect was not found with other steels.

See Dichloroacetylene

0610. 3-Chloro-3-trichloromethyldiazirine [35295-56-5]



Liu, M. T. H., *Chem. Eng. News*, 1974, **52**(36), 3

An ampoule of the compound exploded when scored with a file, indicating high shock-sensitivity.

See other DIAZIRINES

0611. Hexachloroethane [67-72-1]



HCS 1980, 530


Metals

See Aluminium: Halocarbons

Zinc: Halocarbons

See other HALOALKANES

0612. Chromyl isothiocyanate []



Forbes, G. S. *et al.*, *J. Amer. Chem. Soc.*, 1943, **65**, 2273

The salt, prepared in carbon tetrachloride solution, exploded feebly several times as the reaction proceeded. Oxidation of thiocyanate by chromium(VI) is postulated.

See related METAL CYANIDES

0613. Chromyl isocyanate

[]

Forbes, G. S. *et al.*, *J. Amer. Chem. Soc.*, 1943, **65**, 2273

The product is stable during unheated vacuum evaporation of its solutions in carbon tetrachloride. Evaporation with heat at ambient pressure led to a weak explosion of the salt.

See related METAL CYANIDES

0614. Caesium acetylide

[22750-56-7]



Mineral acids

Mellor, 1946, Vol. 5, 848

It ignites in contact with gaseous hydrogen chloride or its conc. aqueous solution, and explodes with nitric acid.

Iron(III) oxide

Mellor, 1946, Vol. 5, 849

Incandescence on warming.

Halogens

Mellor, 1946, Vol. 5, 848

It burns in all four halogens at or near ambient temperature.

Lead dioxide

See Lead(IV) oxide: Metal acetylides

Non-metals



Mellor, 1946, Vol. 5, 848

Mixtures with boron or silicon react vigorously on heating.

See other METAL ACETYLIDES

0615. Copper(II) acetylide

[12540-13-5]



1. Urbanski, 1967, Vol. 3, 228

2. Bond, J., *Hazards from Pressure, IChE Symp. Ser. No. 102*, 37–44, Oxford, Pergamon, 1987

3. Anon., *Jahresber.*, 1987, 64; 1988, 140 (same incident)

Copper(II) acetylide (black or brown) is much more sensitive to impact, friction or heat than copper(I) acetylide (red-brown) which is used in electric fuses or


detonators [1]. An explosion during maintenance work on a copper condenser in a methanol plant was attributed to copper acetylide deposits in the heat exchanger [2]. During internal inspection of an air-purged 15 year old 330 m³ liquefied gas cylinder, it was noticed that there was a 2–3 cm layer of rust-like deposit at the bottom. As the inspection proceeded and the layer was disturbed, small flames were seen, and then a detonation occurred. The deposit was found to contain 20% of copper but less than 1% of sulfur, so was thought to be a mixture of rust and copper acetylide, friction-sensitive when dry [3].

See Silver acetylide (reference 4)

See other METAL ACETYLIDES

0616. Copper(II) cyanide

[544-92-3]



Magnesium

See Magnesium: Metal cyanides

See other METAL CYANIDES (AND CYANO COMPLEXES)

0617. Copper(II) fulminate

[22620-90-2]



See entry METAL FULMINATES

0618. Copper(II) thiocyanate

[15192-76-4]



Tudela, D., *J. Chem. Educ.*, 1993, **70**(2), 174

On addition to boiling water, black cupric thiocyanate converts to the white cuprous salt with violent gas evolution. It is suggested that the gas is hydrogen cyanide, though carbon dioxide seems probable.

See related METAL CYANATES

0619. Dicopper(I) acetylide

[1117-94-8]



1. Mellor, 1946, Vol. 5, 851, 852

2. Rutledge, 1968, 84–85

3. Morita, S., *J. Soc. High Press. Gas Ind.*, 1955, **19**, 167–176

4. Bond, J., *Loss Prev. Bull.*, 1995, **121**, 17

Readily formed from copper or its compounds and acetylene, it detonates on impact or heating above 100°C. If warmed in air or oxygen, it explodes on subsequent contact with acetylene [1]. Explosivity of the precipitate increases with acidity of the salt solutions, while the stability increases in the presence of reducing agents (formaldehyde, hydrazine, or hydroxylamine). The form with a metallic lustre was the most explosive acetylide made. Catalysts with the acetylide supported on a porous solid are fairly stable [2]. The ignition temperature of the pure red acetylide is 260–270°C. On exposure to air or oxygen, it is converted to black copper(II) acetylide, which ignites and explodes at 100°C [3]. It may be formed inadvertently in copper chemical plant; an account of such an incident and a procedure for making such equipment safe (treatment with warm 5% hydrogen peroxide, followed by sodium diethyldithiocarbamate of the same concentration) is reported [4].

Halogens

Mellor, 1946, Vol. 5, 852

Ignition occurs on contact with chlorine, bromine vapour or finely divided iodine.

Silver nitrate

Mellor, 1946, Vol. 5, 853

Contact with silver nitrate solution transforms copper(I) acetylide into a sensitive and explosive mixture of silver acetylide and silver.

Talc

Chambionnat, A., *Chem. Abs.*, 1951, **45**, 7791e

The effect of adding talc as an inert desensitiser upon the sensitivity of the acetylide has been studied.

See other METAL ACETYLIDES

0620. Copper(I) cyanide

[544-92-3]




3-Methoxy-2-nitrobenzoyl chloride

See 3-Methoxy-2-nitrobenzoyl chloride

See other METAL CYANIDES

0621. Dicopper(I) ketenide

[[41084-90-6] (ion)]



Blues, E. T. et al., *J. Chem. Soc., Chem. Comm.*, 1973, 921

The dry compound is mildly explosive.

See other HEAVY METAL DERIVATIVES

- 0622. Copper(I) oxalate**
[53421-36-6] **C₂Cu₂O₄**
- CuOCO.CO.OCu**
- Sidgwick, 1950, 126
It explodes feebly on heating.
See other METAL OXALATES
- 0623. Difluoroethyne (Difluoroacetylene)**
[689-99-6] **C₂F₂**
- FC≡CF**
1. Feast, W. J., Personal communication, 1993
2. Bürger, H. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1991, 456
Like other substituted acetylenes, this is violently unstable. Attempts to prepare poly(difluoroacetylene) thence usually end in explosion [1]. Difluoroacetylene decomposes slowly even at liquid nitrogen temperatures [2].
See other HALOACETYLENE DERIVATIVES
- 0624. 2-(Difluoroamino)-2,2-dinitroacetonitrile**
[180570-35-8] **C₂F₂N₄O₄**
- NCC(NO₂)₂NF₂**
- See* 1,1-DINITRODIFLUORAMINES, *See other* N-HALOGEN COMPOUNDS, POLYNITRO-ALKYL COMPOUNDS
- 0625. Bis(fluoroformyl) peroxide**
[692-74-0] **C₂F₂O₄**
- FCO.OOCO.F**
1. Talbot, R. L., *J. Org. Chem.*, 1968, **33**, 2095
2. Czerepinski, R. *et al.*, *Inorg. Chem.*, 1968, **7**, 109
Several explosions occurred during the preparation, which involves charging carbon monoxide into a mixture of fluorine and oxygen [1]. It has been known to decompose or explode at elevated temperatures, and all samples should be maintained below 30°C and well shielded [2].
See other DIACYL PEROXIDES
- 0626. Trifluoroacetyl nitrite**
[667-29-8] **C₂F₃NO₃**
- F₃CCO.ON:O**
1. Taylor, C. W. *et al.*, *J. Org. Chem.*, 1962, **27**, 1064
2. Gibbs, R. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1972, 1340
3. Banks, R. E. *et al.*, *J. Chem. Soc. (C)*, 1956, 1350–1353
4. Banks, R. E. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1974, 2535
Though much more stable than acetyl nitrite even at 100°C, the vapour of trifluoroacetyl nitrite will explode at 160–200°C unless diluted with inert gas to below

about 50 vol% concentration. Higher perfluorohomologues are more stable [1]. A detailed examination of the explosion parameters has been made [2]. This and higher polyfluoroacyl nitrites tend to explode above 140°C at ambient pressure, and handling of large quantities should be avoided [3], especially during pyrolysis [4].
See other ACYL NITRITES

0627. Trifluoroacetyl azide
[23292-52-6]

$C_2F_3N_3O$




Sprenger, G. H. *et al.*, *Inorg. Chem.*, 1973, **12**, 2891

It explodes on exposure to mechanical or thermal shock. Care is necessary during preparation to eliminate hydrogen chloride from the precursory acid chloride, to prevent formation of hydrogen azide.

See other ACYL AZIDES

†0628. Tetrafluoroethylene
[116-14-3]

C_2F_4



1. Graham, D. P., *J. Org. Chem.*, 1966, **31**, 956
2. Muller, R. *et al.*, *Paste u. Kaut.*, 1967, **14**, 903
3. Kirk Othmer, 1994, Vol. 11, 624; Ullmann, 1988, **A11**, 396

A terpene inhibitor is usually added to the monomer to prevent spontaneous polymerisation, and in its absence, the monomer will spontaneously explode at pressures above 2.7 bar. The inhibited monomer will explode if ignited [1]. Explosion under thermal initiation is now held to be a disproportionation, that to tetrafluoromethane and carbon gives 3.2 kJ/g, the same energy as black powder [3]. Liquid tetrafluoroethylene, being collected in a liquid nitrogen-cooled trap open to air, formed a peroxidic polymer which exploded [2].

See other POLYMERISATION INCIDENTS

Air, Hexafluoropropene

Dixon, G. D., private comm., 1968

A mixture of the two monomers and air sealed in an ampoule formed a gummy peroxide during several weeks. The residue left after opening the ampoule exploded violently on warming.

See other POLYPEROXIDES

Chloroperoxytrifluoromethane

See Chloroperoxytrifluoromethane: Tetrafluoroethylene

Difluoromethylene dihypofluorite

See Difluoromethylene dihypofluorite: Haloalkenes

Dioxygen difluoride

See Dioxygen difluoride: Various materials

Iodine pentafluoride, Limonene

MCA Case History No. 1520

Accidental contamination of a tetrafluoroethylene gas supply system with iodine pentafluoride caused a violent explosion in the cylinders. Exothermic reaction of the limonene inhibitor with the contaminant present in the gas cylinders may have depleted the inhibitor and initiated explosive polymerisation.

See other POLYMERISATION INCIDENTS

Metal alkoxides

See Methyl trifluorovinyl ether

Oxygen

See Oxygen (Gas): Tetrafluoroethylene

Sulfur trioxide

See Sulfur trioxide: Tetrafluoroethylene


Triboron pentafluoride

See Triboron pentafluoride: Tetrafluoroethylene

See other HALOALKENES

0629. Poly(tetrafluoroethylene)

[9002-84-0]



Boron, or Magnesium, or Titanium

Shidlovskii, A. A. *et al.*, *Chem. Abs.*, 1978, **88**, 193944

The combustion of compressed mixtures of the polymer with the finely powdered elements under nitrogen or argon has been studied, reactivity decreasing in the order B, Mg, Ti.

Aluminium: Polytetrafluoroethylene

Fluorine

See Fluorine: Polymeric materials

Metal hydrides

Andreev, B. M. *et al.*, *Chem. Abs.*, 1987, **106**, 162978

In a study of thermal stability and hydrogen sorption characteristics of a series of sorbent tablets composed of hydride-forming metals dispersed in polymers under a 50% hydrogen in argon atmosphere, it was found that tablets of 80% palladium in PTFE, and 80% of 1:5 atom lanthanum–nickel alloy in PTFE could not be used above 247°C because of explosive decomposition of the PTFE.

Oxygen

See FLUOROCARBONS

See related HALOALKANES

0630. (Difluoroamino)difluoroacetonitrile

[5131-88-4]

C₂F₄N₂

Hydrazine

John, E. O. *et al.*, *Inorg. Chem.*, 1988, **27**, 3100–3104


The major product of reaction of the nitrile with anhydrous hydrazine in THF at -196° , followed by slow warming to 25°C for 2 h, is 2-difluoroamino-2,2-difluoro-1-iminoethylhydrazine, F₂NCF₂C(:NH)NHNH₂, which is stable only in solution. Evaporation of solvent led on each occasion to mild explosions, caused by decomposition of the product with considerable gas evolution, for which a mechanism is suggested.

See other GAS EVOLUTION INCIDENTS

See other CYANO COMPOUNDS, DIFLUOROAMINO COMPOUNDS

0631. 3-Fluoro-3-(trifluoromethyl)-3H-diazirine

[11713-32-0]


C₂F₄N₂Dailey, W. P., *Tetrahedron Lett.*, 1987, **28**, 5801–5804

As it is potentially explosive in neat liquid or solid forms, it is best handled in solution.

See other DIAZIRINES, N-HALOGEN COMPOUNDS

0632. Tetrafluorooxirane (Tetrafluoroethylene oxide)

[694-17-7]

C₂F₄O

Preparative hazard

Eleuterio, H. S., *J. Macromol. Sci., Chem.*, 1972, **6**, 1027

Formation from tetrafluoroethylene and oxygen may be explosive. It rearranges to trifluoroacetyl fluoride above its b.p., -63.5°C .

See Oxygen: Tetrafluoroethylene

See other 1,2-EPOXIDES

0633. Trifluoroacetyl hypofluorite

[359-46-6]

C₂F₄O₂

Alone, or Potassium iodide, Water

Cady, G. H. *et al.*, *J. Amer. Chem. Soc.*, 1953, **75**, 2501–2502

The gas explodes on sparking and often during preparation or distillation. Unless much diluted with nitrogen, it explodes on contact with aqueous potassium iodide.

See other ACYL HYPOHALITES

0634. Xenon(II) fluoride trifluoroacetate

[25710-89-8]



$\text{C}_2\text{F}_4\text{O}_2\text{Xe}$


Jha. N. K., *RIC Rev.*, 1971, **4**, 157

It explodes on thermal or mechanical shock.

See other XENON COMPOUNDS

0635. Tetrafluorooxathietane-2,2-dioxide (Tetrafluoroethane sultone)

[697-18-7]



$\text{C}_2\text{F}_4\text{O}_3\text{S}$

Preparative hazard

See Sulfur trioxide: Tetrafluoroethylene

0636. Pentafluoroethylithium (Perfluoroethylithium)

[91935-83-0]



$\text{C}_2\text{F}_5\text{Li}$

Roddick, D. M., *Chem. Eng. News*, 1997, **75**(40), 6

This reagent, prepared in situ from butyllithium and chloropentafluoroethane as an ether solution, has several times decomposed violently on addition of a few drops of a chlorophosphine. This is attributed to running the preparation too cool (< -90°C), so that it had not proceeded. However:

See other METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

0637. Perfluoro-N-cyanodiaminomethane

[16408-94-9]




$\text{C}_2\text{F}_5\text{N}_3$

See Fluorine: Sodium dicyanamide

See other CYANO COMPOUNDS, DIFLUOROAMINO COMPOUNDS

0638. Bis(trifluoromethyl) nitroxide

[2154-71-4]



$\text{C}_2\text{F}_6\text{NO}$

Benzoyl azide

Banks, R. E. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1981, 455–456

Interaction on warming from -196°C in absence of solvent led to a violent explosion.

Platinum hexafluoride


See Platinum hexafluoride: Bis(trifluoromethyl) nitroxide

See other N-O COMPOUNDS

0639. Bis(trifluoromethyl)phosphorus(III) azide

[1479-48-7]

$C_2F_6N_3P$



1. Allcock, H. R., *Chem. Eng. News*, 1968, **46**(18), 70

2. Tesi, G. *et al.*, *Proc. Chem. Soc.*, 1960, 219


Explosive, but stable if stored cold [1]. Previously it was found to be unpredictably unstable, violent explosions having occurred even at $-196^\circ C$ [2].

See other NON-METAL AZIDES

†**0640. N-(Trifluoromethylsulfinyl)trifluoromethylimidodisulfinyl azide**

[81341-46-0]

$C_2F_6N_4O_2S_2$



Bechtold, T. *et al.*, *J. Fluorine Chem.*, 1981-2, **19**, 385

An explosive liquid.

See other ACYL AZIDES, N-S COMPOUNDS

0641. 1,1-Bis(fluorooxy)tetrafluoroethane

[16329-92-3]

$C_2F_6O_2$



1. Thompson, P. G. *et al.*, *J. Amer. Chem. Soc.*, 1967, **89**, 2203

2. Sekiya, A. *et al.*, *Inorg. Chem.*, 1980, **19**, 1329-1330

Though reported to be stable up to $200^\circ C$ in metal [1], samples in glass at $22^\circ C$ exploded during vaporisation at ambient pressure, or when a sealed tube was broken. Higher homologues are less stable [2].

See other BIS(FLUOROOXY)PERHALOALKANES, GLASS INCIDENTS

0642. Bis(trifluoromethyl)trioxide

[1718-18-9]

$C_2F_6O_3$




Gobbato, K. I. *et al.*, *Angew. Chem. (Int.)*, 1995, **34** 20, 2244

This unstable gas (b.p. $-16^\circ C$) has never been reported as exploding, possibly because experimenters have always exercised the greatest caution. It is recommended that this continue. As well as being thermodynamically unstable, the immediate decomposition products being trifluoromethyl peroxide and oxygen, it should also be a strong oxidant and therefore a greater danger in presence of combustible organics.

See ORGANIC PEROXIDES

0643. Bis(trifluoromethyl) sulfide
[371-78-8]



$\text{C}_2\text{F}_6\text{S}$

Chlorine fluorides

Sprenger, G. H. *et al.*, *J. Fluorine Chem.*, 1976, **7**, 335

In the preparation of bis(trifluoromethyl)sulfur difluoride, explosions have occurred when chlorine mono- or tri-fluorides were treated with 2–3 mmol of the sulfide in absence of solvent. Because chlorine trifluoride is known to deprotonate solvents with formation of explosive carbene species, only fully halogenated solvents are suitable as diluents.

See other ALKYLNON-METALS

0644. Bis(trifluoromethyl) disulfide
[372-64-5]



$\text{C}_2\text{F}_6\text{S}_2$


Chlorine fluorides

Sprenger, G. H. *et al.*, *J. Fluorine Chem.*, 1976, **57**, 335

In the preparation of bis(trifluoromethyl)sulfur difluoride, the presence of fully halogenated solvents is essential to prevent explosion during treatment of the disulfide with chlorine mono- or tri-fluoride.

See related ALKYLNON-METALS

0645. Fluorobis(trifluoromethyl)phosphine oxide
[34005-83-9]




$\text{C}_2\text{F}_7\text{OP}$

Preparative hazard

See Fluorobis(trifluoromethyl)phosphine, next below

See related ALKYLHALOPHOSPHINES

0646. Fluorobis(trifluoromethyl)phosphine
[1426-40-0]



$\text{C}_2\text{F}_7\text{P}$

Mahler, W., *Inorg. Chem.*, 1979, **18**, 352

Admission of air to effect oxidation to the phosphine oxide must be slow to prevent ignition.

See other ALKYLHALOPHOSPHINES

0647. Perfluoro-1-aminomethylguanidine
[16408-93-8]



$\text{C}_2\text{F}_8\text{N}_4$

See Fluorine: Cyanoguanidine

See other *N,N,N'*-TRIFLUOROAMIDINES

0648. Bis(trifluoromethyl)sulfur difluoride
[30341-38-9] **C₂F₈S**
 $(F_3C)_2SF_2$

Preparative hazard

See Bis(trifluoromethyl) sulfide: Chlorine fluorides
Bis(trifluoromethyl) disulfide: Chlorine fluorides
See related ALKYLNON-METAL HALIDES

0649. C,N-Perfluoro-N-aminomethyltriaminomethane
[16408-92-7] **C₂F₁₀N₄**
 $F_2NCF_2NFCF(NF_2)_2$

See Fluorine: Cyanoguanidine
See other DIFLUOROAMINO COMPOUNDS

0650. Monosilver acetylide
[13092-75-6] **C₂HAg**
 $AgC\equiv CH$

Anon., *Chem. Trade J.*, 1966, **158**, 153

A poorly stoppered dropping bottle of silver nitrate solution absorbed sufficient acetylene from the atmosphere in an acetylene plant laboratory to block the dropping tube. A violent detonation occurred on moving the dropping tube.

See Silver nitrate: Acetylene
See other METAL ACETYLIDES

0651. Bromoacetylene
[593-61-3] **C₂HBr**
 $BrC\equiv CH$

1. Tanaka, R. *et al.*, *J. Org. Chem.*, 1971, **36**, 3856

2. Hucknall, D. J. *et al.*, *Chem. & Ind.*, 1972, 116

It is dangerous and may burn or explode in contact with air, even when solid at $-196^\circ C$ [1]. Procedures for safe generation, transfer and storage, all under nitrogen, are described [2].

See other HALOACETYLENE DERIVATIVES

0652. Chloroacetylene
[593-63-5] **C₂HCl**
 $ClC\equiv CH$

Tanaka, R. *et al.*, *J. Org. Chem.*, 1971, **36**, 3856

It is notably endothermic (ΔH_f° (g) +213 kJ/mol, 3.52 kJ/g). Chloroacetylene will burn and may explode in contact with air, and its greater volatility makes it more dangerous than bromoacetylene. Procedures for safe generation, handling and storage, all under nitrogen, are described.

See Trichloroethylene: Alkali

See other ENDOTHERMIC COMPOUNDS, HALOACETYLENE DERIVATIVES

0653. Difluoroacetyl hypochlorite

[71359-63-2]



See entry ACYL HYPOHALITES (reference 1)

0654. Chloratomercurio(formyl)methylenemercury(II)

[]




Whitmore, 1921, 154

An extremely sensitive crystalline solid, which explodes on shaking with its crystallisation liquor, or upon gentle mixing when dry with copper oxide (for analysis).

See other MERCURY COMPOUNDS

0655. 5-Chloro-1,2,3-thiadiazole

[4113-57-9]



1. Yoshida, T., private comm., 1982


2. Ouchi, H. *et al.*, *Kogyo Kayaku*, 1984, **45**, 73 (*Chem. Abs.*, 1985, **102**, 187475)

Two recent industrial-scale accidents which involved explosive decomposition of this intermediate caused great damage. The compound has considerable explosive potential, for though it lacks oxygen in its structure, it is of rather low oxygen balance [1]. A comprehensive examination using several hazard evaluation techniques revealed it as an extremely dangerous compound. DSC and the drop-hammer test on the solid do not indicate hazard potential, but ARC at 126°C under adiabatic conditions give a corrected time-to-explosion of 40 min.

The impact-sensitivity was determined as 2 kg cm [2].

See entry OXYGEN BALANCE See other N-S COMPOUNDS

†0656. Trichloroethylene
[79-01-6]



(MCA SD-14, 1956); NSC 389, 1979; HCS 1980, 919; RSC Lab. Hazard Data Sheet No. 13, 1983

1. Author's comments, 1988
2. Santon, R. C. *et al.*, *Proc. 6th Intl. Sympos. Loss Prev. Safety Prom. Proc. Ind.*, 1989, Oslo, Norweg. Soc C. Eng.

Though usually considered to be non-flammable in use, it has unusually wide flammability limits, and with an AIT of 410°C it can be ignited by a flame or other intense source. The most recent Data Sheet is devoted to all aspects of laboratory use [1]. The limits of flammability have been determined more accurately as 8.5 vol%/25°C to 14.5%/35°, widening to 6.0–47% at 150°C. Intermediate values are also given [2].

See FLASH POINTS (reference 19)

Alkali

1. Fabian, F., private comm., 1960
2. *ABCM Quart. Safety Summ.*, 1956, **27**, 17
3. Anon., *Jahresber.*, 1978, 70

An emulsion, formed during extraction of a strongly alkaline liquor with trichloroethylene, decomposed with evolution of the spontaneously flammable gas, dichloroacetylene [1]. This reaction could also occur if alkaline metal-stripping preparations were used in conjunction with trichloroethylene degreasing preparations, some of which also contain amines as inhibitors, which could also cause the same reaction [2]. Apparently accidental contact of the solvent with potassium hydroxide solution led to generation of flames in the charging port of a stirred reactor [3].

See Tetrachloroethylene: Sodium hydroxide

Aluminium

See Aluminium: Halocarbons

Epoxides

Dobinson, B. *et al.*, *Chem. & Ind.*, 1972, 214

1-Chloro-2,3-epoxypropane, the mono- and di-2,3-epoxypropyl ethers of 1,4-butanediol, and 2,2-bis[4(2',3'-epoxypropoxy)phenyl]propane can, in presence of catalytic quantities of halide ions, cause dehydrochlorination of trichloroethylene to dichloroacetylene, which causes minor explosions when the mixture is boiled under reflux. A mechanism is discussed.

See Alkali, above

Metals

See Aluminium: Halocarbons

Barium: Halocarbons

Beryllium: Halocarbons

Lithium: Halocarbons
Magnesium: Halocarbons
Titanium: Halocarbons

Oxidants

See Dinitrogen tetroxide: Halocarbons
Oxygen (Gas): Halocarbons
Oxygen (Liquid): Halocarbons
Perchloric acid: Trichloroethylene


Water

Brade, C., *Chem. Tech.* (Berlin), 1952, **4**, 506–507

A violent explosion occurred during routine recovery of wet trichloroethylene by distillation at ambient pressure. This was attributed to hydrolytic formation of hydrochloric acid and concentration of the latter, leading to corrosion and blocking of the vent line which caused pressure development and a consequential increase in distillation temperature of the vessel contents. The corrosion products may have catalysed the exothermic decomposition reaction which led to explosion. The vessel contained much carbonised residue.

See other CORROSION INCIDENTS See other HALOALKENES

0657. 5-Trichloromethyltetrazole [2925-21-5]



1. Beck, W. *et al.*, *Chem. Eng. News*, 1984, **62**(10), 39
2. Finnegan, W. G. *et al.*, *J. Amer. Chem. Soc.*, 1958, **80**, 3908
3. Howe, R. K. *et al.*, *Chem. Eng. News*, 1983, **61**(3), 4
4. Burke, L. A., *Chem. Eng. News*, 1983, **61**(17), 2

The later publication [1] reveals that the title compound is in fact a relatively stable compound. The previously attempted preparation of the then unknown compound from trichloroacetonitrile, sodium azide and ammonium chloride (0.14:0.42:0.2 mol) by an analogous established method [2], but at lower initial temperature because of the exothermic reaction, gave, after vacuum evaporation of solvent, an oily product. When sampled with a pipette, this evolved gas and then exploded violently. It was thought that an azidomethyltetrazole may have been formed by displacement of chloro-substituent(s) by the excess azide employed [3]. An alternative hypothesis which involved isomerisation of the title compound to the open chain azidoazomethine [4] was discounted, because no trace of this could be detected [1].

See other TETRAZOLES

0658. Trichloroacetic acid
[76-03-9]



Copper, Dimethyl sulfoxide
See Dimethyl sulfoxide: Copper, etc.

Methanol, Water

1. Mabbott, D. J., Personal communication, 1997
2. Ullmann, 1986, Vol. A6, 545.
3. Merck Index, 1989, 1515

A bottle labelled as 12% acid in aqueous methanol, and probably two years forgotten, exploded in storage, breaking adjacent bottles[1]. Trichloroacetic acid is known to be more unstable, with respect to carbon dioxide and chloroform, in aqueous solution than pure. The reaction usually requires either heat or base catalysis [2]. Storage of trichloroacetic acid at less than 30% concentration is not advised [3]. Hydrolysis of the trichlorogroup is also conceivable, which would yield intermediate oxalyl monochloride, which habitually breaks down to give carbon monoxide, dioxide and hydrogen chloride.

See other ORGANIC ACIDS

0659. Trichloroperoxyacetic acid
[7796-16-9]



Swern, D., *Chem. Rev.*, 1945, **45**, 10

Very unstable, forming mainly the toxic gaseous products phosgene, chlorine, carbon monoxide and hydrogen chloride.

See other GAS EVOLUTION INCIDENTS, PEROXYACIDS

0660. Pentachloroethane
[76-01-7]



Potassium

See Potassium: Halocarbons

See other HALOALKANES

0661. Monocaesium acetylide
[30180-52-0]



Mellor, 1946, Vol. 5, 849–850


It reacts similarly to the dicaesium compound.

See Caesium acetylide

See other METAL ACETYLIDES

0662. Fluoroacetylene
[2713-09-9]

C₂HF



Alone, or Bromine


Middleton, W. J. *et al.*, *J. Amer. Chem. Soc.*, 1959, **81**, 803–804

It is notably endothermic (ΔH_f° (g) +125 kJ/mol, 2.84 kJ/g), and liquid fluoroacetylene is treacherously explosive close to its b.p., -80°C . The gas does not ignite in air and is not explosive. Ignition occurred in contact with a solution of bromine in carbon tetrachloride. The mercury and silver salts were stable to impact, but the latter exploded on heating, whereas the former decomposed violently.

See other ENDOTHERMIC COMPOUNDS, HALOACETYLENE DERIVATIVES

†0663. Trifluoroethylene
[359-11-5]

C₂HF₃



Hulbert, J. D. *et al.*, *Chem. Eng. News* 1997, **75**(51), 6

Trifluoroethene at pressures above 14 bar, and possibly below, deflagrates with tenfold pressure rises when initiated by an exploding wire. The tests were on air-free and inhibited commercial material.

Ethylene, Tetrafluoroethylene

See other HALOALKENES

0664. 2,2,2-Trifluorodiazethane
[371-67-5]

C₂HF₃N₂



Fields, R. *et al.*, *J. Fluorine Chem.*, 1979, **13**, 154

The stabilising effect of the trifluoromethyl group is insufficient to prevent explosions, either of the compound itself, or of unstable reaction intermediates formed at low temperatures, especially in absence of solvent. Energy released on explosion is estimated at 50–100% TNT equivalent.

Ozone

See Ozone: Trifluoroethylene

See other DIAZO COMPOUNDS

0665. Trifluoroacetic acid
[76-05-1]

C₂HF₃O₂



HCS 1980, 935

Aromatic hydrocarbons, Hydrogen peroxide

See Hydrogen peroxide: Aromatic hydrocarbons, Trifluoroacetic acid

Lithium tetrahydroaluminate
See Lithium tetrahydroaluminate: Fluoroamides
See other ORGANIC ACIDS

0666. Peroxytrifluoroacetic acid
[359-48-8]




Sundberg, R. J. *et al.*, *J. Org. Chem.*, 1968, **33**, 4098
This extremely powerful oxidant must be handled and used with great care.

4-Iodo-3,5-dimethylisoxazole

Plepys, R. A. *et al.*, US Pat. 3 896 140, 1975
Interaction to produce 3,5-dimethyl-4-bis(trifluoroacetoxy)iodoisoxazole yields a detonable by-product, believed to be iodine pentaoxide contaminated with organic material.

See Iodine(V) oxide: Non-metals
See other OXIDANTS, PEROXYACIDS

0667. 3,3,5-Trifluoro-1,2,4-trioxolane (Trifluoroethylene ozonide)
[86013-87-8]




Explosive.
See Ozone: Trifluoroethylene
See other OZONIDES

0668. 1,1,2,2-Tetrafluoro-1-(fluoroxy)ethane (1,1,2,2-Tetrafluoroethyl hypofluorite)
[]




Randolph, B. B. *et al.*, *J. Fluorine Chem.*, 1993, **64**(1-2), 129
The hypofluorite exploded on attempted purification. A variety of related compounds were also unstable, if not explosive.
See other HYPOHALITES

0669. Poly(dimercuryimmonium acetylde)
[]



Sorbe, 1968, 97
Highly explosive.
See other POLY(DIMERCURYIMMONIUM) COMPOUNDS

0670. Iodoacetylene
[14545-08-5]



Sorbe, 1968, 65
It explodes above 85°C.
See other HALOACETYLENE DERIVATIVES


0671. Monopotassium acetylide
[1111-63-3]



Chlorine
Mellor, 1946, Vol. 5, 849
It ignites in chlorine.

Non-metal oxides
Mellor, 1946, Vol. 5, 849
Interaction with sulfur dioxide at ambient temperature, or with carbon dioxide on warming, causes incandescence.
See other METAL ACETYLIDES

0672. Dipotassium *aci*-nitroacetate
[19419-98-8]



Water
1. Lyttle, D. A., *Chem. Eng. News*, 1949, **27**, 1473
2. Whitmore, F. C., *Org. Synth.*, 1941, Coll. Vol. 1, 401
An inhomogeneous mixture of the dry salt with a little water exploded violently after 30 min [1]. This was probably owing to exothermic decarboxylation generating the *aci*-salt of nitromethane, which is explosively unstable. The decomposition of sodium nitroacetate proceeds exothermically above 80°C [2].
See other aci-NITRO SALTS

0673. Monolithium acetylide
[1111-64-4]



Mellor, 1946, Vol. 5, 849
It reacts similarly to the dilithium salt.
See Lithium acetylide
See other METAL ACETYLIDES

0674. Monolithium acetylide-ammonia

[]

 $C_2HLi.H_3N$ 

Gases, or Water

Mellor, 1946, Vol. 5, 849

The complex ignites in contact with carbon dioxide, sulfur dioxide, chlorine or water.

See related METAL ACETYLIDES**0675. Diazoacetonitrile**

[13138-21-1]

 C_2HN_3 1. Phillips, D. D. *et al.*, *J. Amer. Chem. Soc.*, 1956, **78**, 54522. *MCA Case History No. 2169*3. Roelants, F. *et al.*, *Tetrahedron*, 1978, **34**, 2231

4. Author's comments

Removal of a rubber stopper from a flask of a concentrated solution in methylene chloride initiated an explosion, probably through friction on solvent-free material between the flask neck and bung. Handling only in dilute solution is recommended [1]. An explosion also occurred during operation of the glass stopcock of a dropping funnel while 20 ml of the apparently undiluted nitrile was being added to a reaction mixture [2]. All attempts to isolate the nitrile from its solutions in chlorinated solvents caused explosions [3]. By inference from the thermodynamic properties of diazomethane and acetonitrile, diazoacetonitrile should be strongly endothermic, with heat of formation exceeding 200 kJ/mol, 3 kJ/g [4].


See other DIAZO COMPOUNDS, FRICTIONAL INITIATION INCIDENTS, GLASS INCIDENTS

0676. Dinitroacetonitrile

[921-22-2]

 $C_2HN_3O_4$ *See entry* POLYNITROALKYL COMPOUNDS**0677. 3-Diazo-3H-1,2,4-triazole**

[64781-78-8]

 C_2HN_5 

The precursory 1,2,4-triazole-3-diazonium nitrate is extremely unstable, even in ice-cold aqueous solution, and no attempt was made to isolate the title diazo


compound from its solution, prepared by cautious treatment of the cold diazonium nitrate solution with dilute aqueous potassium hydroxide with cooling.

See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

0678. 4-Diazo-1,2,3-triazole
[85807-68-7]

C₂HN₅



See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

0679. Diazoacetyl azide
[19932-64-0]

C₂HN₅O





Neunhöffer, G. *et al.*, *Ann.*, 1968, **713**, 97–98

The material may be distilled cautiously at 20–21°/0.27 mbar but is preferably used in solution as prepared. Either the solid (below 7°C) or the liquid explodes with great violence on impact or friction.

See other ACYL AZIDES, DIAZO COMPOUNDS

0680. 1,2,4-Triazole-3-diazonium nitrate
[59104-93-7]

C₂HN₆O₃




Magee, H. L. *et al.*, *J. Org. Chem.*, 1987, **52**, 5547

The diazonium nitrate is extremely unstable and may only be prepared as an aqueous solution of limited life at below 0°C.

See other DIAZONIUM SALTS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

0681. Sodium 5-(dinitromethyl)tetrazolide
[2783-96-2]


$C_2HN_6NaO_4$



Einberg, F., *J. Org. Chem.*, 1964, **29**, 2021
It explodes violently at the m.p., 160°C, and may be an *aci*-nitro salt.
See other POLYNITROALKYL COMPOUNDS, TETRAZOLES

0682. Sodium 5(5'-hydroxytetrazol-3'-ylazo)tetrazolide
[]

$C_2HN_{10}NaO$



Thiele, J. *et al.*, *Ann.*, 1893, **273**, 150
The salt explodes very violently on heating.
See other AZO COMPOUNDS, TETRAZOLES

0683. 1,1,1-Triazidodinitroethane
[56522-42-0]

$C_2HN_{11}O_4$



Frankel, M. B. *et al.*, UK Pat. Appl. 2 123 829, 1984
An energetic compound (with 63% N and of low oxygen balance).
See other POLYNITROALKYL COMPOUNDS, ORGANIC AZIDES

0684. Monosodium acetylide
[1066-26-8]

C_2HNa



1. Mellor, 1946, Vol. 5, 849
2. Greenlee, K. W., *Inorg. Synth.*, 1947, **2**, 81
3. Houben-Weyl, Vol. 13.1, 280

It reacts similarly to the disodium salt [1]. If heated to 150°C, it decomposes extensively, evolving gas which ignites in air owing to presence of pyrophoric carbon. The residual carbon is also highly reactive [2]. The dry powder (from solution in liquid ammonia) may ignite if exposed to air as an extended layer, e.g. on filter paper [3].

See Sodium acetylide

See other METAL ACETYLIDES

0685. Monorubidium acetylide

[20720-71-2]

C₂HRb



Mellor, 1946, Vol. 5, 849

It reacts similarly to the dirubidium salt.

See Rubidium acetylide

See other METAL ACETYLIDES

†0686. Acetylene (Ethyne)

[74-86-2]

C₂H₂



(MCA SD-7, 1957); NSC 494, 1982; FPA H6, 1972; HCS 1980, 110, 111

1. Mayes, H. A., *Chem. Engr.*, 1965, (185), 25
2. Nedwick, J. J., *Ind. Eng. Chem., Proc. Des. Dev.*, 1962, **1**, 137; Tedeschi, R. J. *et al.*, *ibid.*, 1968, **7**, 303
3. Miller, 1965, Vol. 1, 506
4. Kirk-Othmer, 1963, Vol. 1, 195–202
5. Rimarski, W., *Angew. Chem.*, 1929, **42**, 933
6. Foote, C. S., private comm., 1965
7. Sutherland, M. E. *et al.*, *Chem. Eng. Progr.*, 1973, **69**(4), 48–51
8. Williams, A. *et al.*, *Chem. Rev.*, 1970, **70**, 270–271
9. Menge, R. *et al.*, *Brandschutz, Explosionschutz*, 1987, **16**, 109
10. Yantovskii, S. A., *Chem. Abs.*, 1964, **61**, 13117f
11. Landesman, Ya. M. *et al.*, *Chem. Abs.*, 1974, **81**, 151352
12. Glikin, M. A. *et al.*, *Chem. Abs.*, 1975, **83**, 100289
13. Savitskaya, L. M. *et al.*, *Chem. Abs.*, 1978, **88**, 157743
14. Viallon, J., *Loss Prev. Safety Prom. Proc. Ind.*, 121–122, Frankfurt, Dechema, 1978
15. Lietze, D. *et al.*, *Chem. Ing. Tech.*, 1978, **50**, 808–809 (summary of 21 page paper)
16. Code of Practice for Acetylene Pipelines, Paris, Euro. Comm. Indust. Gases, 1978
17. Grosse-Wortmann, H. *et al.*, *Chem. Ing. Tech.*, 1981, **53**, 461–463
18. Lietze, D., *Chem. Abs.*, 1986, **105**, 48048

19. Ashmore, F. S. *et al.*, *Fire Prev.*, 1987, (204), 37–39; *Loss Prev. Bull.*, 1988, (080), 15–18
20. Marcks, G., *Proc. 11th Int. Symp. Prev. Occup. Risks Chem. Ind.*, 623–649, Heidelberg, ISSA, 1987
21. Dorfinger, S., *ibid.*, 653–663
22. Schildberg, H. P. *et al.*, *Chem. Ing. Tech.*, 1994, **66**(10), 1389

The extremely endothermic gas (ΔH_f° (g) +226.7 kJ/mol, 8.72 kJ/g) may decompose explosively in absence of air. Addition of up to 30% of miscible diluent did not sufficiently desensitise explosive liquid acetylene to permit of its transportation. The solid has similar properties [1]. However, safe techniques for the use of liquid acetylene in reaction systems at pressures up to 270 bar have been described [2]. Solid acetylene in admixture with liquid nitrogen at -181°C is sensitised by presence of grit (carborundum) and may readily explode on impact [3]. The hazards of preparing and using acetylene have been adequately described, including many reports of explosions in acetylene generators and distribution systems (formerly including domestic lighting installations) where faulty pressure control has allowed pressures to approach 1.4 bar, when explosion occurs in presence of moisture [3,4,5]. A glass flask which had contained acetylene was left open to atmosphere for a week, and then briefly purged with air. It still contained enough acetylene to form an explosive mixture [6]. The explosive decomposition of acetylene in absence of air readily escalates to detonation in tubular vessels. This type of explosive decomposition has been experienced in a 7 mile acetylene pipeline system [7].

In a review on combustion and oxidation of acetylene, the factors affecting spontaneous ignition at low temperatures are discussed [8]. The autoignition temperature may be as low as 85°C when mixed with 1–2% of nitric oxide (NO) [9]. The explosive decomposition of acetylene and conditions necessary for spontaneous ignition to occur have been reviewed [10,11,12]. Use of C₁–C₄ alcohols, C₃–C₄ ketones, C₁ and C₄ acids and C₃–C₄ diols or hydrocarbons to retard explosive decomposition was studied comparatively. The retardation mechanism involves thermal and chemical factors, related to heat content of diluent and to structure, respectively. Practical systems are detailed [13]. Acetylene at 170°C is somewhat stabilised by presence of acetic acid [14]. It is destabilised by potassium hydroxide [22]. The course of explosive decomposition of the gas in pipelines has been studied experimentally [15]. The EU Code of Practice divides pipeline systems into 3 ‘working ranges’, depending on internal pressure and diameter of the pipe, and specifies suitable design and construction criteria for each range [16]. The relationship between initiation energy and decomposition has been studied [17]. The explosibility of acetylene, alone or admixed with air, hydrogen or ethylene has been reviewed, including boundary limits for deflagration/detonation transformation [18]. Accidental local heating to 185°C or above of part of the wall (as little as 6 sq. cm may be enough) of a cylinder containing acetylene may lead to the development of an extremely dangerous situation. At this temperature, exothermic and self-sustaining decomposition, or polymerisation, of (endothermic) acetylene may set in, and if not stopped by rapid and effective cooling (large volume water spray), the cylinder may explode without warning. Flame flash-back into a cylinder from

a wrongly adjusted and/or damaged welding or cutting torch can cause the same effect. Methods of identifying such unstable cylinders and of emergency control are detailed [19]. The hazards associated with installations of banks of acetylene cylinders connected to a high-pressure manifold are discussed [20], and a detailed account of an accident involving explosion of two such overheated cylinders is given [21].

See Magnesium: Barium carbonate, Water

Bleaching powder

See Oxidants, below

Cobalt

MRH 8.70/tr.

Mellor, 1942, Vol. 14, 513

Finely divided (pyrophoric) cobalt decomposes acetylene on contact, becoming incandescent.

Copper

MRH 8.70/tr.

Anon., *ABCM Quart. Safety Summ.*, 1946, **17**, 24

Rubber-covered electric cable, used as a makeshift handle in the effluent pit of an acetylene plant, formed copper acetylide with residual acetylene and the former detonated when disturbed and initiated explosion of the latter. All heavy metals must be rigorously excluded from locations where acetylene may be present.

See METAL ACETYLIDES

Copper(I) acetylide

MRH 8.70/tr.

See Dicopper(I) acetylide

Dinitrogen oxide

See ATOMIC ABSORPTION SPECTROSCOPY

Ethylene, Hydrogen

See Hydrogen (Gas): Acetylene, Ethylene

Gaseous additives

1. Konovalov, E. N. *et al.*, *Chem. Abs.*, 1975, **82**, 111548

2. Detz, C. M., *Combust. Flame*, 1979, **34**, 187–193; US Pat. 4 161 495, 1979

Addition of 0.3–3.5 wt% of propane or butane reduces the explosion hazards of acetylene–air mixtures [1]. Out of 10 gaseous additives tested for effect on the threshold temperature for initiation of explosive decomposition of acetylene by a heated wire at 2–22 bar, nitrogen oxide, hydrogen bromide, hydrogen chloride, hydrogen iodide and vinyl bromide showed stabilising effects, and sulfur dioxide a mild destabilising effect. The patent specifies use of one or more of these radical scavenging agents at levels of 0.5–10% [2].

Halogens

MRH Chlorine or Iodine 8.70/tr.

1. Muir, G. D., private comm., 1968

2. von Schwartz, 1918, 142, 321

3. MCA SD-7, 1957
4. Humphreys, V., *Chem. & Ind.*, 1971, 681–682
5. Sokolova, E. I. *et al.*, *Chem. Abs.*, 1970, **72**, 71120
6. Davenport, A. P., *School Sci. Rev.*, 1973, **55**(191), 332
7. Long, G. C., *Spectrum* (Pretoria), 1980, **18**, 30
8. Nolan, 1983, Case History 22
9. Nolan, 1983, Case History 23

Tetrabromoethane is made by passing acetylene into bromine in carbon tetrachloride at reflux. The rate of reaction falls off rapidly below reflux temperature, and if the rate of addition of acetylene is insufficient to maintain the temperature, high concentrations of unreacted acetylene build up, with the possibility of a violent delayed reaction [1]. In absence of a diluent, reaction may be explosive [2]. Mixtures of acetylene and chlorine may explode upon initiation by sunlight or other UV source, or at high temperature, sometimes very violently [2]. Interaction with fluorine is very violent [3] and with iodine possibly explosive [2]. Dilution of equimolar mixtures of chlorine and acetylene with 55 mol% of nitrogen or 70% air prevented spontaneous explosion. At higher dilutions, sparking did not initiate explosion [4]. Explosive interaction of chlorine and acetylene in the dark is initiated by presence of oxygen at between 0.1 and about 40 vol%. The reaction is inhibited by inert gases or oxygen at higher concentrations [5]. Safe techniques for demonstrating explosive combination of acetylene and chlorine have been described [6,7], the latter involving dropping a small lump of calcium carbide into acidified sodium hypochlorite solution. Tetrachloroethane is manufactured by reacting excess chlorine with acetylene at 100°C in presence of ferric chloride. On one occasion the temperature fell to 60°C (low chlorine flow?) and there was an explosion. It was suggested that monochloroacetylene had formed and decomposed, initiating an acetylene/chlorine or gas/air explosion. A number of such explosions have occurred, particularly during start-up [8]. Another explosion was attributed to damp ferric chloride [9].

Heavy metals and salts

See METAL ACETYLIDES

Liquid nitrogen

Houseman, T. H. *et al.*, *J. Labelled Compd. Radiopharm.*, 1978, **14**, 164

Liquid nitrogen should not be used as a trap coolant with acetylene, owing to the explosive nature of liquid or solid acetylene (title reference 5 above).

Mercury(II) salts, Nitric acid, (Sulfuric acid)

See Nitric acid: Acetylene, Mercury(II) salts

Nitric acid, Perchloric acid

See Perchloric acid: Acetylene, Nitric acid

Other reactants

Yoshida, 1980, 15

MRH values calculated for 10 combinations with materials catalysing the decomposition of acetylene are given.

Oxidants

Since acetylene is endothermic (values above) and effectively a reducing agent, its reactions with oxidants are usually violent or explosive if uncontrolled.

See Dinitrogen oxide or Halogens, both above; Oxygen, below

Calcium hypochlorite: Acetylene

Nitric acid: Acetylene, Mercury(II) salts

Nitrogen oxide: Dienes, Oxygen

Ozone: Acetylene

Trifluoromethyl hypofluorite: Hydrocarbons

Oxides of nitrogen

See Nitrogen oxide: Dienes, Oxygen

Oxygen

1. Fowles, G. *et al.*, *School Sci. Rev.*, 1940, **22**(85), 6; 1962, **44**(152), 161; 1963, **44**(154), 706; 1964, **45**(156), 459
2. Kiyama, R. *et al.*, *Rev. Phys. Chem. Japan*, 1953, **23**, 43–48
3. Moye, A., *Chem. Ztg.*, 1922, **46**, 69
4. Ivanov, U., *Chem. Abs.*, 1976, **84**, 57968

The explosion of acetylene–oxygen mixtures in open vessels is a very dangerous experiment (stoichiometric mixtures detonate with great violence, completely shattering the container). Full precautions are essential for safety [1]. When a mixture of acetylene and oxygen (54:46) at 270°C/10.9 bar was compressed in 0.7 s to 56.1 bar, the resulting explosion attained a pressure of several kbar. In other tests rapid compression of acetylene or its mixtures with air caused no explosions [2]. Previously, passage of acetylene into liquid air to deliberately generate a paste of solid acetylene and liquid oxygen, ‘by far the most powerful of explosives’, had been proposed [3]. Acetylene had been collected for teaching purposes over water in a pneumatic trough. Later, oxygen was collected in the same way without changing the water, and the sample exploded violently when exposed to a glowing splint. Acetylene remaining dissolved in the water had apparently been displaced by the oxygen stream, the lower explosive limit for acetylene being only 2.5% in air, and less in oxygen [4].

Potassium

MRH 8.70/tr.

Berthelot, M., *Bull. Soc. Chim. Fr.* [2], 1866, **5**, 188

Molten potassium ignites in acetylene, then explodes.

Potassium hydroxide

Schildberg, H. P. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries*, (Mewis, Pasman and De Rademaker, eds.), Vol. 1, 401, Amsterdam, Elsevier, 1995

At temperatures above 100°C and pressures above 2 bar, potassium hydroxide powder can initiate self-propagating decomposition of acetylene. More detail on limits is given.

See other ALKYNES

0687. Poly(acetylene)

[25067-58-7]

$(C_2H_2)_n$

$(-CH=CH-)_n$

Feast, W. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1985, 202–203

The second stage of an improved synthesis of poly(acetylene), which involves disproportionation of a soluble polymer by heating a thin film at 75°C to give 1,2-bis(trifluoromethyl)benzene and poly(acetylene), must not be done in bulk because the reaction then becomes explosive. The earlier synthesis by direct polymerisation of acetylene was considerably more dangerous

See Poly([7,8-bis(trifluoromethyl)tetracyclo[4.2.0.0^{2,8}.0^{5,7}]octane-3,4-diyl]-1,2-ethenediyl)

Iodine

See Iodine: Poly(acetylene)

See related DIENES

Perchlorate-doping


See PERCHLORATE-DOPEDED CONDUCTING POLYMERS

See related DIENES

0688. Silver 1,2,3-triazolide

[]

$C_2H_2AgN_3$



See 1,2,3-Triazole

See other N-METAL DERIVATIVES, SILVER COMPOUNDS, TRIAZOLES

0689. Silver dinitroacetamide

[[26163-27-9] (ion)]

$C_2H_2AgN_3O_5$

$AgC(NO_2)_2CO.NH_2$


Parker, C. O. *et al.*, *Tetrahedron*, 1962, **17**, 108

The sodium salt explodes at 160°C and the potassium and silver salts at 159 and 130°C, respectively.

See other POLYNITROALKYL COMPOUNDS, SILVER COMPOUNDS

0690. Silver 1,3-di(5-tetrazolyl)triazenide

[]


Ag⁺

See 1,3-Di(5-tetrazolyl)triazene

See other SILVER COMPOUNDS, TETRAZOLES, TRIAZENES

0691. 2-Iodosovinyl chloride

[]



Alone, or Water


Thiele, J. *et al.*, *Ann.*, 1909, **369**, 131

It explodes at 63°C and contact with water disproportionates it to the more explosive iodyl compound (next below).

See other IODINE COMPOUNDS

0692. 2-Iodylvinyl chloride

[]

Thiele, J. *et al.*, *Ann.*, 1909, **369**, 131

Explodes violently on impact, friction or heating to 135°C.

See other IODINE COMPOUNDS

0693. Azidoacetyl chloride

[30426-58-5]

1. Wieland, T. *et al.*, *Chem. Ber.*, 1960, **93**, 12362. Prodan, K. A. *et al.*, *Chem. Eng. News*, 1981, **59**(14), 593. Hoppe, D. *et al.*, *Ann.*, 1980, 1519

Using a published procedure [1] for preparing azidoacetyl chloride, but at 1.7-fold higher concentration, azidoacetic acid and sulfinyl chloride were slowly heated up to reflux. After 40 min. decomposition started and accelerated to a violent explosion [2]. The cause seems likely to have been the higher initial concentration of azido acid, which would have produced a higher concentration (87%) of acid chloride in excess sulfinyl chloride, and at a significantly higher reflux temperature, than the original method (about 50%). A relatively safe way of preparing the explosively unstable material involves a low temperature partial distillation procedure, but full precautions are recommended [3].

See Azidoacetic acid
See other ORGANIC AZIDES



0694. Sodium chloroacetate
[3296-62-3]



Energy of decomposition (in range 130–220°C) was measured as 0.29 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 96°C by adiabatic Dewar tests, with an apparent energy of activation of 112 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

†**0695. 1,1-Dichloroethylene (Vinylidene chloride)**
[75-35-4]



HCS 1980, 378

1. Reinhardt, R. C., *Chem. Eng. News*, 1947, **25**, 2136
2. MCA Case History No. 1172
3. MCA Case History No. 1693
4. Harmon, 1974, iv, 2.21
5. Warren, H. S. *et al.*, *Chem. Abs.*, 1978, **89**, 219959

When stored at between -40° and $+25^\circ\text{C}$ in the absence of inhibitor and in presence of air, vinylidene chloride rapidly absorbs oxygen with formation of a violently explosive peroxide. The latter initiates polymerisation, producing an insoluble polymer which adsorbs the peroxide. Separation of this polymer in the dry state must be avoided, since if more than 15% of peroxide is present, the polymer may be detonable by slight shock or heat. Hindered phenols are suitable inhibitors to prevent peroxidation [1]. The earlier Case History describes an explosion during handling of a pipe used to transfer the chloride [2]. Two further cases of formation of the polymeric peroxide in bottles of vinylidene chloride in refrigerated storage, and subsequent explosions during handling or disposal, are recorded [3]. The monomer is normally handled at -10°C in absence of light or water, which tend to promote self-polymerisation [4]. Hazards from explosions when the chloride is stored or handled in presence of air are discussed [5].

See entries POLYMERISATION INCIDENTS, POLYPEROXIDES, VIOLENT POLYMERISATION

Chlorotrifluoroethylene

Raasch, M. S. *et al.*, *Org. Synth.*, 1962, **42**, 46

Condensation of the reactants at 180°C under pressure to give 1,1,2-trichloro-2,3,3-trifluorocyclobutane was effected smoothly several times in a 1 l autoclave. Scaling up to a 3 l preparation led to uncontrollable polymerisation which distorted the larger autoclave.



Ozone

'Vinylidene Chloride Monomer', 9, Midland (Mich.), Dow Chemical Co., 1968
The reaction products formed with ozone are particularly dangerous.
See other OZONIDES

Perchloryl fluoride



See Perchloryl fluoride: Hydrocarbons, etc.
See other HALOALKENES, PEROXIDISABLE COMPOUNDS

†0696. *cis*-1,2-Dichloroethylene
[156-60-5]



See other HALOALKENES

†0697. *trans*-1,2-Dichloroethylene
[156-59-2]



1. Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 37
2. Anon., *Sichere Chemiearb.*, 1964, **16**, 35

Under appropriate conditions, dichloroethylene, previously thought to be non-flammable, can cause a fire hazard [1]. Addition of a hot liquid to the cold solvent caused sudden emission of sufficient vapour to cause a flame to flash back 12 m from a fire. Although the bulk of the solvent did not ignite, various items of paper and wood in the room were ignited by the transient flame [2].

Alkalies

1. Anon., *Fire Accid. Prev.*, 1956, **42**, 28
2. Thron, H., *Chem. Ztg.*, 1924, **48**, 142

1,2-Dichloroethylene in contact with solid caustic alkalies or their conc. solutions will form chloroacetylene which ignites in air [1]. Distillation of ethanol containing 0.25% of the halocarbon with aqueous sodium hydroxide gave a product which ignited in air [2].

See Trichloroethylene: Alkali

1,1,2,2-Tetrachloroethane: Alkalies

Difluoromethylene dihypofluorite

See Difluoromethylene dihypofluorite: Haloalkenes


Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Halocarbons

See other HALOALKENES

0698. Trichloroacetaldehyde oxime
[1117-99-3]

$C_2H_2Cl_3NO$



Alkali

Rodd, 1965, Vol. 1C, 46

The oxime decomposes explosively with alkali, forming hydrogen cyanide, hydrogen chloride and carbon dioxide.

See other GAS EVOLUTION INCIDENTS, OXIMES

0699. 1,1,2,2-Tetrachloroethane
[79-34-5]

$C_2H_2Cl_4$



(MCA SD-34, 1949); HCS 1980, 886

Alkalis

MCA SD-34, 1949

It is not an inert solvent, and on heating with solid potassium hydroxide or other base, hydrogen chloride is eliminated and chloro- or dichloro-acetylene, which ignite in air, are formed.

Metals

See Potassium: Halocarbons


Sodium: Halocarbons

See METAL-HALOCARBON INCIDENTS

See other HALOALKANES

†**0700. 1,1-Difluoroethylene (Vinylidene fluoride)**
[75-38-7]

$C_2H_2F_2$



HCS 1980, 409

Hydrogen chloride

Jensen, J. H., *Chem. Eng. News*, 1981, **59**(14), 3, 59

As well as the normal addition reaction, an extremely exothermic decomposition reaction may occur, particularly at high vessel loadings. At loadings of 0.8 ml of 1:1 mixture per ml, the violent reaction, catalysed by iron(III) chloride, initiates at $-40^\circ C$ and will attain pressures above 0.7 kbar at the rate of 14 kbar/s. At 0.5 ml loading density, a maximum pressure of 68 bar, attained at 114 bar/s, was observed.

See entries CATALYTIC IMPURITY INCIDENTS, PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Ozone


See Ozone: 1,1-Difluoroethylene

See other HALOALKENES

0701. 2-(Difluoroamino)-2,2-dinitroacetamide

[180570-38-1]

$C_2H_2F_2N_4O_5$



See 1,1-DINITRODIFLUORAMINES, See other N-HALOGEN COMPOUNDS, POLY-NITRO-ALKYL COMPOUNDS

0702. Difluoroacetic acid

[381-73-7]

$C_2H_2F_2O_2$



Caesium fluoride, Fluorine


See Fluorine: Caesium fluoride, etc.

See other ORGANIC ACIDS

0703. 1,1-Difluoroethylene ozonide (3,3-Difluoro-1,2,4-trioxolane)

[69932-17-8]

$C_2H_2F_2O_3$



See Ozone: 1,1-Difluoroethylene

See other OZONIDES

0704. Trifluoroacetamide

[354-38-1]

$C_2H_2F_3NO$



Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Fluoroamides

0705. Sodium 2,2,2-trifluoroethoxide

[420-87-1]

$C_2H_2F_3NaO$




Schmutz, J. L. *et al.*, *Inorg. Chem.*, 1975, **14**, 2437

During preparation from sodium and excess alcohol in ether, attempts to remove last traces of the alcohol by warming under vacuum led to explosions.

See related METAL ALKOXIDES

0706. 1,1,2,2-Tetrafluoroethane
[811-97-2]




Schröder, V. *et al. Loss Prevention and Safety Promotion in the Process Industries* Vol II (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 339, Amsterdam, Elsevier, 1995

Although not flammable under normal conditions in air, it is at elevated pressure, or with oxygen enrichment. Explosive limits are reported and reviewed.

See also FLUOROCARBONS

0707. Mercury(II) aci-dinitromethanide
[]




Urbanski, 1967, Vol. 3, 158

This salt of *aci*-dinitromethane shows detonator properties.

See other MERCURY COMPOUNDS, *aci*-NITRO SALTS

0708. 1,2-Bis(hydroxomercurio)-1,1,2,2-bis(oxydimercurio)ethane ('Ethane hexamercarbide')
[67536-44-1]




Hofmann, K. A., *Ber.*, 1898, **31**, 1904

The compound, formulated as above, explodes very violently at 230°C.

See other MERCURY COMPOUNDS

0709. Potassium dinitroacetamide
[50650-92-5]



See Silver dinitroacetamide

See other POLYNITROALKYL COMPOUNDS

0710. Diazoacetaldehyde
[6832-13-9]



Arnold, Z., *Chem. Comm.*, 1967, 299

It may be distilled out continuously as formed at 40°C/13 mbar, but readily detonates with great violence if overheated.

See other ALDEHYDES, DIAZO COMPOUNDS

0711. 2-Nitroethanonitrile (Nitroacetonitrile)

[]



Koenig, L. *et al.*, Personal communication, 1997



Made by dehydration of nitroacetaldehyde oxime, this is an explosion hazard. It is improbable that the nitrile is not, itself, detonable.

See 2-Nitroacetaldehyde oxime

See other C-NITRO COMPOUNDS

0712. Sodium dinitroacetamide

[[26163-27-9] (ion)]




See Silver dinitroacetamide

See other POLYNITROALKYL COMPOUNDS

0713. Sodium 2,2,2-trinitroethanide

[]





See *N,N'*-Bis(trinitroethyl)urea

See other POLYNITROALKYL COMPOUNDS

0714. Azidoacetonitrile

[57707-64-9]





Freudenberg, K. *et al.*, *Ber.*, 1932, **65**, 1188

With over 51% nitrogen content, it is, as expected, explosive, sensitive to impact or heating to 250°C.

See other CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

0715. 1,2,4,5-Tetrazine

[290-96-0]




Sulfuric acid

Sorbe, 1968, 139

The solid base decomposes violently in contact with conc. acid.
See other HIGH-NITROGEN COMPOUNDS, ORGANIC BASES

0716. 3-Nitro-1,2,4-triazolone (Oxynitrotriazole)
[932-64-9]





Ullmann, 1987, A10, 162

Finds employment as an high explosive (oxygen balance -40%).

See EXPLOSIVES, OXYGEN BALANCE

See other HIGH-NITROGEN COMPOUNDS, NITRO COMPOUNDS, TRIAZOLES

0717. 3-Azido-1,2,4-triazole
[21041-85-0]




Denault, G. C. *et al.*, *J. Chem. Eng. Data*, 1968, **13**, 514

Samples exploded during analytical combustion.

See other ORGANIC AZIDES, TRIAZOLES

0718. 3-Amino-6-nitro-1,2,4,5-tetrazine-2,4-dioxide
[153757-95-0]




See 3,6-Diamino-1,2,4,5-tetrazine-1,4-dioxide

See other HIGH-NITROGEN COMPOUNDS, *See also* AMINE OXIDES

0719. 5-(Diazomethylazo)tetrazole

[]

$C_2H_2N_8$



Mayants, A. G. *et al.*, *Zh. Org. Khim.*, 1991, **27**(11), 2450; *Chem. Abs.*, 1992, **116**, 234937

The compound, produced as an hydrate from dilute acid hydrolyses of disodium azotetrazolide, exploded on heating for analytical purposes.

See other AZO COMPOUNDS, DIAZO COMPOUNDS, TETRAZOLES

0720. 1,2-Diazidocarbonylhydrazine

[67880-17-5]

$C_2H_2N_8O_2$



Kesting, W., *Ber.*, 1924, **57**, 1321


Similar in properties to lead or silver azide, it explodes on heating or impact.

See other ACYL AZIDES, HIGH-NITROGEN COMPOUNDS

0721. 5,5'-Tetrazolyl disulfide (5,5'-Dithiobis[1H]tetrazole)

[72848-18-1]

$C_2H_2N_8S_2$




Christensen, T. B. *et al.*, *Sulfur Lett.*, 1993, **16**(5-6), 205

In a paper devoted to tetrazole chemistry, the explosive nature of this compound excited comment.

See other TETRAZOLES; *See also* HIGH NITROGEN COMPOUNDS

0722. Ketene (Ethenone)

[463-51-4]



$\text{C}_2\text{H}_2\text{O}$

Hydrogen peroxide

See Hydrogen peroxide: Ketene

†0723. Ethanedial (Glyoxal)

[107-22-2]



$\text{C}_2\text{H}_2\text{O}_2$

HCS 1980, 522

Preparative hazard

See Nitric acid: 2,4,6-Trimethyltrioxane

Alone, or Water

1. Merck, 1983, 647

2. van den Brink, M. J., *Chemisch Magazine* (Rijswijk), 1982, (7-8), 428

A powerfully reducing and reactive solid (m.p., 15°C) of high vapour pressure (b.p., 50°C). Mixtures with air may explode, and contact with water causes violent polymerisation [1]. More usually encountered polymeric or as a hydrate. Like formaldehyde, pure glyoxal may polymerise exothermally and ignite in storage [2].

Nitric acid, Water

See Nitric acid: Glyoxal

See other ALDEHYDES, REDUCANTS

0724. Glyoxylic acid (Oxoacetic acid)

[298-12-4]



$\text{C}_2\text{H}_2\text{O}_3$

Preparative hazard

See Nitric acid: Glyoxal

See other ORGANIC ACIDS See related ALDEHYDES

0725. Oxalic acid (Ethanedioic acid)

[144-62-7]



$\text{C}_2\text{H}_2\text{O}_4$

NSC 406, 1978; HCS 1980, 706

CHETAH, 1990, 184

Surprisingly, even the dihydrate is apparently detonated by a 50 g tetryl booster, the anhydrous acid is thermally less stable and thus probably more sensitive. There is, however, no history of explosion.

Other reactants

Yoshida, 1980, 178

MRH values calculated for 5 combinations are given.

Silver

See Silver: Carboxylic acids


Sodium chlorite

See Sodium chlorite: Oxalic acid

Urea

von Bentzinger, R. *et al.*, *Praxis Naturwiss. Chem.*, 1987, **36**(8), 41–42

During the qualitative analysis of an ‘unknown’ mixture of oxalic acid, urea and a little charcoal, a sample was put in an ignition tube under a layer of granular copper oxide, topped by glass wool and a little anhydrous copper sulfate (to detect evolved water). The top end of the tube was drawn down obliquely into a capillary so that the pyrolysis gases could be passed into barium hydroxide solution (to detect carbon dioxide). It was not noticed that the end of the capillary was sealed, and when the sample was heated in a flame, an unexpectedly violent explosion followed. Later experiments showed that there was no explosion if oxalic acid alone, or urea alone, was heated with copper oxide in a sealed tube, but that a mixture of the two always led to a violent explosion. The same also happened in absence of copper oxide. Further work (DSC, and measurement of volumes of gas liberated) led to the conclusions that the reactions involved were:-



and that a gas pressure approaching 100 bar would be produced in the accidentally sealed ignition tube at 570°C.

See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS, REDOX REACTIONS

See other ORGANIC ACIDS

0726. Lead carbonate–lead hydroxide (Basic lead carbonate)

[598-63-0]



$\text{C}_2\text{H}_2\text{O}_8\text{Pb}_3$


Fluorine

See Fluorine: Metal salts

See other INORGANIC BASES

†0727. Bromoethylene (Vinyl bromide)

[593-60-2]



$\text{C}_2\text{H}_3\text{Br}$

1. *HCS* 1980, 956

2. *DOC* 5, 1982, 799

The highly flammable gas reacts violently with oxidants [1] and may polymerise in sunlight [2].

See other HALOALKENES

0728. Acetyl bromide
[506-96-7]

C₂H₃BrO



HCS 1980, 108

Hydroxylic compounds

Merck, 1976, 11

Interaction with water, methanol or ethanol is violent, hydrogen bromide being evolved.

See other ACYL HALIDES

0729. Acetyl hypobromite
[4254-22-2]

C₂H₃BrO₂



1. Reilly, J. J. *et al.*, *J. Org. Chem.*, 1974, **39**, 3992


2. Skell, P. S. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 4000, 4007

The instability of the hypobromite is noted, and protective shielding is recommended for work with more than 5 g quantities [1]. Though stable in fluorotrichloromethane solution at -23°C under nitrogen in the dark, the isolated solid exploded at -23°C at below 1.2 mbar [2].

See other ACYL HYPOHALITES

†0730. Chloroethylene (Vinyl chloride)
[75-01-4]

C₂H₃Cl



(*MCA SD-56*, 1972); *HCS 1980*, 957; *RSC Lab. Hazards Data Sheet No. 56*, 1987

1. *MCA Case History No. 625*

2. Harmon, 1974, 2.20, 4.74–4.77

3. *MCA Case History No. 1551*

4. Scali, C. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 1350–1358, Basle, SSCI, 1980

5. Anon., *Eur. Chem. News*, 1993, **60**(1589), 26

Discharge of a spray of vapour and liquid under pressure from a cylinder into a fume hood generated static and ignited the vapour. Discharge of the vapour phase only did not lead to ignition [1]. Vinyl chloride tends to self-polymerise explosively if peroxidation occurs, and several industrial explosions have been recorded [2]. Accidental exposure of the recovered monomer to atmospheric oxygen for a long period caused formation of an unstable polyperoxide which initiated an explosion. Suitable precautions are discussed, including the use of 20–30% aqueous sodium hydroxide solution to destroy the peroxide [3]. Problems of stability and control of batch reactors for the suspension polymerisation of vinyl chloride have been studied in relation to process kinetics and plant heat transfer coefficients [4]. A polymerisation reaction ran beyond control, generating back pressure which caused the monomer to leak through a valve, forming a vapour cloud which ignited and exploded [5].

See 1,1-Dichloroethylene

See VIOLENT POLYMERISATION

See *other* POLYMERISATION INCIDENTS, POLYPEROXIDES, STATIC INITIATION INCIDENTS, VAPOUR CLOUD EXPLOSIONS

Oxides of nitrogen

Povey, R., private comm., 1979

An explosion in a valve in a liquid monomer line was ascribed to traces of oxides of nitrogen remaining after the valve had been passivated by treatment with nitric acid.


Oxygen

Tartari, V., *Proc. 34th Ital. Chem. Congr. Safety Chem. Proc.*, 51–60, Milan, 1983
The stability of the monomer to heat in the presence (and absence) of oxygen has been studied by ARC. The intensity of the exotherm at 90–100°C depends on the partial pressure of oxygen.

See *other* HALOALKENES, PEROXIDISABLE COMPOUNDS

†0731. 1-Chloro-1,1-difluoroethane

[75-68-3]




Difficult to ignite vapour

See *other* HALOALKANES

See FLASH POINTS (reference 19)

0732. Ethenylmagnesium chloride (Vinylmagnesium chloride, Chloroethenylmagnesium)

[3536-96-7]





Rittmeyer, P. *et al.*, *Z. Naturforsch., B: Chem. Sci.*, 1993, **48**(9), 1223

An incident during concentration of a tetrahydrofuran solution on pilot plant led to a study of the concentration dependent exothermic decomposition of the reagent and generation of safety rules.

See *other* GRIGNARD REAGENTS

0733. 3-Chloro-3-methyldiazirine

[4222-21-3]



1. Liu, M. T. H., *Chem. Eng. News*, 1974, **53**(36), 10


2. Jones, W. E. *et al.*, *J. Photochem.*, 1976, **5**, 233–239

3. Frey, H. M. *et al.*, *J. Chem. Soc., Faraday Trans. 1*, 1977, **73**, 2011

Extremely shock-sensitive and violently explosive; initiation has been caused by prolonged freezing at -196°C , or by sawing a stopcock off a metal trap containing trace amounts [1]. It may be stored safely at -80°C [2]. Detonative explosion during trap-to-trap distillation of purified material is noted [3].

See other DIAZIRINES

0734. 3-Chloro-3-methoxydiazirine
[4222-27-9]



Smith, N. P. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1979, 213

Unpredictably explosive as the neat liquid, it may be handled safely in solution. All distillation traps must contain solvent.

See other DIAZIRINES

†0735. Acetyl chloride (Ethanoyl chloride)
[75-36-5]



FPA H65, 1978; *HCS 1980*, 109

Preparative hazard

See Phosphorus trichloride: Carboxylic acids

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides

Other reactants

Yoshida, 1980, 51

MRH values calculated for 13 combinations with oxidants are given.

Water

1. *Haz. Chem. Data*, 1975, 30

2. Anon., *CISHC Chem. Safety Summ.*, 1986, **57**(225), 3

3. Anon., personal communication 1997

Interaction is violent [1]. A polythene-lined drum which had previously contained acetyl chloride was inspected and appeared empty, but when it was washed out with water a violent eruption occurred. This was attributed to trapping of residual acetyl chloride between the split liner and the drum wall [2]. A very similar but more violent event, rupturing the drum and sending part of it 30 m. with sufficient force to puncture another drum which it hit, occurred in Scotland in 1997 [3].

See other ACYL HALIDES

- †0736. Methyl chloroformate (Methyl carbonochloridate)
[79-22-1] C₂H₃ClO₂
$$\text{H}_3\text{COCO.Cl}$$

HCS 1980, 645
See other ACYL HALIDES
0737. Peroxyacetyl perchlorate
[66955-43-9] C₂H₃ClO₆
$$\text{MeCO.OOClO}_3$$

Yakovleva, A. A. *et al.*, *Chem. Abs.*, 1979, **91**, 210847
The peroxyester of +9.1% oxygen balance exploded on detonation or friction.
See Peroxypropionyl perchlorate
See related ALKYL PERCHLORATES, PEROXYESTERS
0738. 1,1-Dichloro-1-fluoroethane
[1717-00-6] C₂H₃Cl₂F
$$\text{MeCCl}_2\text{F}$$

1. Anon., *Sichere Chemiearbeit*, 1993, **45**(8), 105
2. Bretherick, L., *Chem. Eng. News*, 1988, **66**(50), 2
An 'empty' drum which had contained this exploded when cut into by a grinder. It is presumably more flammable than might be anticipated [1]. Like trichloroethane the vapour is difficult to ignite and may not show a flash point, but can form explosive mixtures with air [2].
See other HALOALKANES
0739. N,N-Dichloroglycine
[58941-14-3] C₂H₃Cl₂NO₂
$$\text{Cl}_2\text{NCH}_2\text{CO.OH}$$

Vit, J. *et al.*, *Synth. Comm.*, 1976, **6**(1), 1–4
It explodes at 65°C, so solutions must be evaporated at below 40°C.
See other N-HALOGEN COMPOUNDS, ORGANIC ACIDS
- †0740. 1,1,1-Trichloroethane
[71-55-6] C₂H₃Cl₃
$$\text{Cl}_3\text{CCH}_3$$

(*MCA SD-90*, 1965); *HCS 1980*, 918; *RSC Lab. Hazard Data Sheet No. 1*, 1982
1. Wrightson, I. *et al.*, *Loss Prev. Bull.*, 1988, (083), 21–23
2. Santon, R. C. *et al.*, *Proc. 6th Int. Sympos. Loss Prev. Safety Prom. Process Ind.*, 1989, Oslo, Norweg. Soc. Chem. Eng.

Although no flash point is measurable by standard tests, the vapour can be ignited by a high-energy source, the limits being 8.0–10.5 vol% in air at 25°C, with an autoignition temperature of 500°C. A 1050 m³ storage tank containing 70 t of the inhibited solvent, normally blanketed with nitrogen to exclude moisture, had developed gas leaks in the roof, and the purge system had been switched to manual control. Previous withdrawal of 1150 t had drawn in air, which had not been effectively displaced by manual purging. Arc-welding to repair the roof leaks led to an explosion which separated the roof from the tank for 65% of the 36 m circumference. Only one other explosion, when the end of an ‘empty’ 200 l drum was cut with an oxy-acetylene torch, has been reported [1]. New work has extended downwards the temperature limits for ignition to 10°C for l_{el} of 8.0 vol%, and 22° for u_{el} of 14.2 vol%. These values diverge with increasing temperature to 6% and 15.5%, respectively, at 100°C [2].

See FLASH POINTS (reference 19)

Aluminium oxide, Heavy metals

See Aluminium oxide: Halocarbons, etc.

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Halocarbons

Inhibitors

See FLASH POINTS (reference 19)

Metals

See Aluminium: Halocarbons

Magnesium: Halocarbons

Potassium: Halocarbons

Potassium–sodium alloy: Halocarbons

METAL–HALOCARBON INCIDENTS

Molecular sieve

1. Salmon, J. *et al.*, *Chem. & Ind.*, 1995, 814
2. Wilson, W., *Chem. & Ind.*, 1995, 854
3. Bretherick, L., personal communication
4. Winterton, N. *et al.*, *Chem. & Ind.*, 1995, 902
5. Whitmore, M. W. *et al.*, *Chem. & Ind.*, 1995, 942

Some unstabilised trichloroethane had stood over molecular sieve (uncertain which grade) for a period of some years. The sieve was filtered out and the bottle, still containing some sieve, purged with nitrogen, after which it exploded when moved. The remaining sieve deflagrated on addition to a bucket of water [1]. It is suggested that the sieve catalysed dehydrochlorination to chloroacetylene, which is explosive [2]. This seems improbable since the acetylene is also volatile and not very polar, thus should not adhere to the dried sieve. An alternative explanation is that dehydrochlorination occurred, giving vinylidene chloride (1,1-dichloroethene) which polymerised with atmospheric oxygen as a comonomer, leading to a polyperoxide

which would adhere to the sieve [3,4]. Another sample of ancient trichloroethane, stored over sieves, showed no instability or peroxidation [5].

Oxygen

See Oxygen (Gas): Halocarbons

Oxygen (Liquid): Halocarbons

See other HALOALKANES

0741. 1,1,2-Trichloroethane

[79-00-5]



HCS 1980, 917

Potassium

See Potassium: Halocarbons

See other HALOALKANES

0742. Methyltrifluoromethyltrichlorophosphorane

[69517-30-2]



Preparative hazard

See Tetrachlorotrifluoromethylphosphorane: Tetramethyllead

0743. N-Carbomethoxyiminophosphoryl chloride

[25147-05-1]



Mellor, 1971, Vol. 8, Suppl. 3, 589

This intermediate (or its ethyl homologue), produced during the preparation of phosphoryl dichloride isocyanate from interaction of phosphorus pentachloride and methyl (or ethyl) carbamate, is unstable. Its decomposition to the required product may be violent or explosive unless moderated by presence of a halogenated solvent.

See related NON-METAL HALIDES

0744. 2,2,2-Trichloroethanol

[115-20-8]



Sodium hydroxide

MCA Case History No. 1574

Accidental contact of 50% sodium hydroxide solution with residual trichloroethanol in a pump caused an explosion. This was confirmed in laboratory experiments.

Chlorohydroxyacetylene, or the isomeric chloroketene, or chlorooxirene may have been formed by elimination of hydrogen chloride.

See related HALOALKANES

0745. 2,2,2-Trichloro-1,1-ethanediol (Chloral hydrate)
[302-17-0]




1. Marvel, C. S. *et al.*, *Org. Synth.*, 1941, Coll. Vol. 1, 377
2. Goodwin, B., *Chem. Brit.*, 1988, **24**, 336
3. Gandy, R. *et al.*, *Chem. Brit.*, 1988, **24**, 336

Following a published procedure for converting substituted anilines to isatins by reaction with chloral hydrate and hydroxylamine [1], it was noticed that at the end of the first stage (formation of an isonitrosoacetanilide), the odour of hydrogen cyanide was present, and this was confirmed by a Prussian blue test [2]. In related work, concentrations of 100–200 ppm of hydrogen cyanide were found [3]. A mechanism for its formation from chloral hydrate and hydroxylamine was proposed [2], and the need for appropriate precautions was stressed [2,3].

See other GAS EVOLUTION INCIDENTS

†0746. Trichlorovinylsilane
[74-94-5]



It may ignite in air.


Water

491M, 1975, 426

It reacts violently with water or moist air.

See related ALKYLHALOSILANES

†0747. Fluoroethylene (Vinyl fluoride)
[75-02-5]





HCS 1980, 958

Ozone

See Fluoroethylene ozonide

See other HALOALKENES

0748. 1-Fluoro-1,1-dinitroethane
[13214-58-9]



See entry FLUORODINITROMETHYL COMPOUNDS

0749. 2(?) -Fluoro-1,1-dinitroethane
[68795-10-8]




MCA Case History No. 784

During a prolonged fractional distillation of the material at 75°C/50 mbar, an exothermic decomposition began. As a remedial measure (!), air was admitted to the hot decomposing residue, causing a violent explosion. Admission of nitrogen and/or cooling the residue before admitting air might have avoided the incident.

See related FLUORODINITROMETHYL COMPOUNDS

0750. 2-Fluoro-2,2-dinitroethanol
[17003-75-7]



1. Cochoy, R. E. *et al.*, *J. Org. Chem.*, 1972, **37**, 3041
2. Cochoy, R. E. *et al.*, *J. Org. Chem.*, 1990, **55**(4) 1401

It is a potentially explosive vesicant, from which a series of esters, expected to be explosive, was prepared [1]. During concentration of a dichloromethane solution, there is a sensitivity maximum between 30% solution and the pure compound [2].

See other FLUORODINITROMETHYL COMPOUNDS, POLYNITROALKYL COMPOUNDS

0751. Acetyl hypofluorite
[78948-09-1]




1. Adam, M. J., *Chem. & Eng. News*, 1985, **63**(7), 2
2. Appelman, E. *et al.*, *J. Amer. Chem. Soc.*, 1985, **107**, 6516–6518

A minor explosion in the use of gaseous acetyl hypofluorite was caused by inadvertent contact of the gas inlet tube with a cooling bath, which condensed liquid hypofluorite which subsequently exploded on contact with organic material or air [1]. More detail on the isolation of this compound of limited stability, particularly in the liquid phase, is given [2].

See other ACYL HYPOHALITES, OXIDANTS

0752. Fluoroethylene ozonide (3-Fluoro-1,2,4-trioxolane)
[60553-18-6]



1. Mazur, W. *et al.*, *J. Org. Chem.*, 1979, **44**, 3183
2. Hillig, K. W. *et al.*, *J. Amer. Chem. Soc.*, 1982, **104**, 991

During ozonolysis of vinyl fluoride an explosive solid residue is produced, and the volatile ozonide, trapped at -63°C , may explode spontaneously or during removal by syringe [1]. During the cryogenic distillation of the ozonide (formulated as a trioxolane), several explosions occurred, and the explosive reaction residue was destroyed by digestion with 5% potassium iodide solution for 24 h [2].

See other OZONIDES

0753. 2-(Difluoroamino)-2,2-dinitroethane

[121832-04-0]



$\text{C}_2\text{H}_3\text{F}_2\text{N}_3\text{O}_4$

See 1,1-DINITRODIFLUORAMINES, *See other* *N*-HALOGEN COMPOUNDS, POLY-NITRO-ALKYL COMPOUNDS

†**0754. 1,1,1-Trifluoroethane**

[420-46-2]



$\text{C}_2\text{H}_3\text{F}_3$

See other HALOALKANES

0755. Pentafluorosulfur peroxyacetate

[60672-60-8]



$\text{C}_2\text{H}_3\text{F}_5\text{O}_3\text{S}$

Hopkinson, M. J. *et al.*, *J. Fluorine Chem.*, 1976, **7**, 505

The peroxyester and its homologues, like other fluoroperoxy compounds, are potentially explosive and may detonate on thermal or mechanical shock.

See related FLUORINATED PEROXIDES AND SALTS, PEROXYESTERS

0756. Potassium 1-nitroethane-1-oximate

[3454-11-3]



$\text{C}_2\text{H}_3\text{KN}_2\text{O}_3$


Rodd, 1965, Vol. 1B, 98

Isolated salts of 'nitrolic acids', produced by action of nitrous acid on 1-nitroalkanes, are explosive.

See related NITROALKANES, OXIMES

0757. Vinyl lithium

[917-57-7]



$\text{C}_2\text{H}_3\text{Li}$

Juenge, E. C. *et al.*, *J. Org. Chem.*, 1961, **26**, 564

When freshly prepared, it is violently pyrophoric but on storage it becomes less reactive and slow to ignite in air, possibly owing to polymerisation.

See related ALKYL METALS

†0758. Acetonitrile (Ethanenitrile)

[75-05-8]

C₂H₃N



NSC 683, 1981; *FPA H108*, 1981; *HCS 1980*, 104
RSC Lab. Hazard Data Sheet No. 31, 1984

A moderately endothermic compound (ΔH_f° (l) +53.1 kJ/mol, 1.29 kJ/g) which, though reactive, does not exhibit inherent instability under normal conditions.

Chlorine fluoride, Fluorine

See Fluorine: Acetonitrile, etc.

2-Cyano-2-propyl nitrate

See NITRATING AGENTS

Dinitrogen tetraoxide, Indium

See Dinitrogen tetraoxide: Acetonitrile, Indium

Diphenyl sulfoxide, Trichlorosilane

See Trichlorosilane: Acetonitrile, etc.

N-Fluoro compounds

Fraser, G. W. *et al.*, *Chem. Comm.*, 1966, 532

Nitrogen-fluorine compounds are potentially explosive in contact with acetonitrile.

See Tetrafluorourea: Acetonitrile

Iron(III) perchlorate

See Iron(III) perchlorate: Acetonitrile

Lanthanide perchlorate

Forsberg, J. H., *Chem. Eng. News*, 1984, **62**(6), 33

A solution of an unspecified lanthanide perchlorate in acetonitrile detonated while being heated under reflux.

See other METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS

Nitric acid

MRH 6.15/77

See Nitric acid: Acetonitrile

Other reactants

Yoshida, 1980, 19

MRH values calculated for 14 combinations with oxidants are given.

Perchloric acid

See Perchloric acid: Acetonitrile

Sulfuric acid, Sulfur trioxide

MRH Sulfuric acid 1.76/54

Lee, S. A., private comm., 1972

A mixture of acetonitrile and sulfuric acid on heating (or self-heating) to 53°C underwent an uncontrollable exotherm to 160°C in a few seconds. The presence of 28 mol% of sulfur trioxide reduces the initiation temperature to about 15°C. Polymerisation of the nitrile is suspected.

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0759. Methyl isocyanide

[593-75-9]

C₂H₃N

H₃CN:C

1. Lemoult, M. D., *Compt. rend.*, 1906, **143**, 902
2. Stein, A. R., *Chem. Eng. News*, 1968, **46**(45), 7–8
3. Collister, J. L., *Diss. Abs. Intl. B*, 1977, **37**, 4479
4. Pritchard, H. O., *Can. J. Chem.*, 1979, **57**, 2677–2679

It is very endothermic (ΔH_f° (g) +150.2 kJ/mol, 3.66 kJ/g). A sample exploded when heated in a sealed ampoule [1], and during redistillation at 59°C/1 bar, a drop of liquid fell back into the dry boiler flask and exploded violently [2]. The explosive decomposition has been studied in detail [3], and existing data on thermal explosion parameters have been re-examined and discrepancies eliminated [4].

See Ethyl isocyanide

See related CYANO COMPOUNDS See other ENDOTHERMIC COMPOUNDS

0760. Glycolonitrile (Hydroxyacetonitrile)

[107-16-4]

C₂H₃NO

HOCH₂C≡N

1. Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 2
2. Gaudry, R., *Org. Synth.*, 1955, Coll. Vol. 3, 436
3. Anon., *Chem. Eng. News*, 1966, **44**(49), 50
4. Parker, A. J. *et al.*, *Aust. J. Chem.*, 1978, **31**, 1187

A year-old bottled sample, containing syrupy phosphoric acid as stabiliser and which already showed signs of tar formation, exploded in storage. The pressure explosion appeared to be due to polymerisation, after occlusion of inhibitor in the tar, in a container in which the stopper had become cemented by polymer [1]. A similar pressure explosion occurred when dry redistilled nitrile, stabilised with ethanol [2], polymerised after 13 days [3]. The spontaneous and violent decomposition of the nitrile on standing for more than a week is usually preceded by formation of a red polymer [4].

See other POLYMERISATION INCIDENTS

Alkali

Sorbe, 1968, 146

Traces of alkali promote violent polymerisation.

See other CYANO COMPOUNDS

†0761. Methyl isocyanate
[624-83-9]

C₂H₃NO



HCS 1980, 570

1. Anon., *CISHC Chem. Safety Summ.*, 1979, **50**, 93
2. Author's comments
3. Anon., *Chem. & Ind.*, 1985, 202
4. Anon., *Loss Prev. Bull.*, 1985, (063), 1-8
5. Worthy, W., *Chem. Eng. News*, 1985, **63**(6), 27-33
6. D'Silva, T. D. J., *J. Org. Chem.*, 1986, **51**, 3781-3788
7. Bowander, B. *et al.*, *J. Haz. Mat.*, 1988, **19**, 237-269

Bottled samples had been supplied 3 years previously packed in vermiculite in sealed cans. The caps had disintegrated in storage allowing the volatile contents to escape into the external can. When this was pierced, a jet of isocyanate sprayed out uncontrollably. Improved caps are now used for this material [1].

There was a disastrous release of methyl isocyanate at Bhopal in December 1984 which caused over 2000 fatalities. Although making it in a continuous process, the plant stored several weeks supply of the isocyanate. An exothermic reaction in one of the buried 13 m³ methyl isocyanate storage vessels raised the internal temperature to above the boiling point (39°C). A cooling system existed but appears to have been turned off at the time. The continuing exotherm caused a further substantial increase in pressure, which was eventually relieved through the emergency alkali scrubber and flare stack system. Unfortunately the scrubbing system proved inadequate to deal with the rapid vapour release, and the flare was not alight, so that of the 35 t of isocyanate in the tank, some 23 t were released during the 2 h the safety valve was open.

The origin of the exothermic reaction was not clear at the time, but it was known that alkyl isocyanates would trimerise to trialkyl isocyanurates in the presence of various catalysts, would react with amines to form alkylureas, and were hydrolysed by water in presence of catalytic amounts of acids or bases, with evolution of carbon dioxide [2]. The key role of water has now been confirmed [3]. The Union Carbide report on the probable causes and course of the Bhopal disaster has been summarised [4,5]. Analysis of the residue in the distorted tank led to proposal of a mechanism, involving both water and chloroform as contaminants, to account for the detailed composition of the residue [6]. A detailed analysis of the many factors and failings which led to the ingress of water into the storage tank and to the subsequent disastrous large-scale vapour release has been presented [7]. Shortly after the event fatalities were estimated as 2,500, but such tolls do not shrink in retelling and figures up to 8,000 have since been given.

Bhopal exhibits several features of maladministration of chemical industry familiar to many who work in it on other continents. Construction of a fine chemical plant on a green field site soon causes local planners to regard the surrounding area as suitable for cheap high-density housing. As Trevor Kletz has reiterated in numerous books, add-on safety features, of which there were many, are a menace. If multiple, any one will be turned off as soon as it causes trouble or expense, and reliance be placed on the others which may be ill-designed and will, in any case, be later turned off in full confidence that the first is still working.

See related ORGANIC ISOCYANATES

0762. Acetyl nitrite
[5813-49-0]



Francesconi, L. *et al.*, *Gazz. Chim. Ital.* [I], 1895, **34**, 439

An unstable liquid, decomposed by light, of which the vapour is violently explosive on mild heating.

See other ACYL NITRITES

0763. Nitroacetaldehyde (Nitroethanal)
[44397-85-5]



Beilstein, Vol. 1, **III**, 2678

Decomposes on standing, explodes on contact with flame.

See other C-NITRO COMPOUNDS

0764. Poly(vinyl nitrate) (Poly(ethenyl nitrate))
[]



Durgapal, U. C. *et al.*, *Propellants, Explosives, Pyrotechnics*, 1995, **20**, 64

This compound, made by mild nitration of polyvinyl alcohol, is an explosive as shock sensitive as tetryl.

See other ALKYL NITRATES

0765. Acetyl nitrate
[591-09-3]



1. Pictet, A. *et al.*, *Ber.*, 1907, **40**, 1164

2. Bordwell, F. G. *et al.*, *J. Amer. Chem. Soc.*, 1960, **82**, 3588

3. Konig, W., *Angew. Chem.*, 1955, **667**, 517

4. Wibaut, J. P., *Chem. Weekbl.*, 1942, **39**, 534

Acetyl nitrate, readily formed above 0°C from acetic anhydride and conc. nitric acid, is thermally unstable and its solutions may decompose violently above 60°C (forming tetranitromethane, a powerful oxidant). The pure nitrate explodes violently on rapid heating to above 100°C [1]. Isolation before use as a nitrating agent at -10°C is not necessary, but the mixture must be preformed at 20–25° before cooling to -10°C to avoid violent reactions [2]. Spontaneous explosions of pure isolated material a few days old had been reported previously [3]. Spontaneous detonation of freshly distilled nitrate has also been recorded, as well as explosion on touching with a glass rod [4].

Ethyl 3,4-dihydroxybenzenesulfonate, Oleum

Dohmen, E. A. M. F. *et al.*, *Chem. Weekbl.*, 1942, **39**, 447–448

Attempted nitration of the sulfonate in 20% oleum led to a violent explosion, probably from decomposition of the nitrate.

Mercury(II) oxide

Chrétien, A. *et al.*, *Compt. rend.*, 1945, **220**, 823

Acetyl nitrate explodes when mixed with red mercuric oxide, or other 'active' (catalytically) oxides.

See other ACYL NITRATES

0766. Peroxyacetyl nitrate (Acetyl nitro peroxide)

[2278-22-0]

C₂H₃NO₅

H₃CCO.OONO₂

1. Stephens, E. R. *et al.*, *J. Air Poll. Control Ass.*, 1969, **19**, 261–264

2. Stephens, E. R., *Anal. Chem.*, 1964, **36**, 928

3. Louw, R. *et al.*, *J. Amer. Chem. Soc.*, 1975, **97**, 4396

Accidental production of the liquid nitrate during overcooled storage at 0°C of a rich mixture of the vapour with helium is thought to have caused a violent explosion [1]. Mixtures of up to 0.1 vol% at 7 bar stored at 10°C or above are quite safe. During dilution with nitrogen of vapour samples in evacuated bulbs, slow pressurisation is necessary to avoid explosion [2]. It is extremely explosive and may only be handled in high dilution with nitrogen. The propionyl analogue is similarly explosive, but higher homologues less so [3].

See related ACYL NITRATES, PEROXYESTERS

0767. Mercaptoacetonitrile (Acetonitrilethiol)

[54524-31-1]

C₂H₃NS

HSCH₂C≡N


Mathias, E. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1981, 569–570

Great caution is necessary when handling conc. solutions or the solvent-free nitrile. Phosphoric acid stabilises it to some extent, but sudden foaming polymerisation

may occur. Gaseous decomposition products may burst the container, even when kept in refrigerated storage.

See other CYANO COMPOUNDS, GAS EVOLUTION INCIDENTS

0768. 1,2,3-Triazole
[27070-49-1]





Baltzer, O. *et al.*, *Ann.*, 1891, **262**, 320, 322

The vapour readily explodes if superheated (above 200°C), and the silver derivative explodes on heating.

See other TRIAZOLES

0769. 1,2,4-Triazole
[288-88-0]




Energy of decomposition (in range 280–420°C) measured as 1.1 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other TRIAZOLES

0770. Vinyl azide
[7570-25-4]



Wiley, R. H. *et al.*, *J. Org. Chem.*, 1957, **22**, 995

A sample contained in a flask detonated when the ground joint was rotated. Literature statements that it is surprisingly stable are erroneous.

See other ORGANIC AZIDES

0771. Acetyl azide
[24156-53-4]




Smith, 1966, Vol. 2, 214

It is treacherously explosive.

See other ACYL AZIDES

0772. Azidoacetaldehyde
[67880-11-9]

$C_2H_3N_3O$




Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 1870

It decomposed with vigorous gas evolution below 80°C at 5 mbar. A preparative reaction mixture of chloroacetaldehyde hydrate and sodium azide had previously exploded mildly on heating in absence of added water.

See other 2-AZIDOCARBONYL COMPOUNDS

0773. 5-Methoxy-1,2,3,4-thiaziazole
[19155-52-3]

$C_2H_3N_3OS$



Jensen, K. A. *et al.*, *Acta. Chem. Scand.*, 1964, **18**, 825

It explodes at ambient temperature, and its higher homologues are unstable.

See related TRIAZOLES

0774. Azidoacetic acid
[18523-48-3]

$C_2H_3N_3O_2$



Iron, or Iron salts

Borowski, S. J., *Chem. Eng. News*, 1976, **54**(44), 5


The pure acid is insensitive to shock at up to 175°C, and not explosively unstable at up to 250°C. In contrast, the presence of iron or its salts leads to rapid exothermic decomposition of the acid at 25°C, and explosion at 90°C or lower under strong illumination with visible light.

See other CATALYTIC IMPURITY INCIDENTS, IRRADIATION DECOMPOSITION INCIDENTS

Azidoacetyl chloride *See other* 2-AZIDOCARBONYL COMPOUNDS, ORGANIC ACIDS

0775. 1,1,1-Trinitroethane
[595-86-8]

$C_2H_3N_3O_6$



Mukhametshin, F. M., *Russ. Pat.* 515 378, 1976

Safety aspects of the preparation from halotrinitromethanes or tetranitromethane by treatment with methyl iodide are improved by use of an aprotic solvent (DMF, DMSO or HMPA) in diethyl ether or carbon tetrachloride at 30–60°C.

See other POLYNITROALKYL COMPOUNDS

0776. 2,2,2-Trinitroethanol
[918-54-7]

$C_2H_3N_3O_7$




1. Marans, N. S. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 5329
2. Cochoy, R. E. *et al.*, *J. Org. Chem.*, 1972, **37**, 3041
3. Shiino, K., *Chem. Abs.*, 1978, **88**, 104580

It is a moderately shock-sensitive explosive which has exploded during distillation [1]. A series of its esters, expected to be explosive, was prepared [2], and explosive ether and ester derivatives are reviewed [3].

See other POLYNITROALKYL COMPOUNDS

0777. 4-Nitroamino-1,2,4-triazole
[52096-16-9]

$C_2H_3N_5O_2$




It explodes at the m.p., 72°C.

See entry N-AZOLIUM NITROIMIDATES *See other* TRIAZOLES

0778. 1,3-Di(5-tetrazolyl)triazene
[56929-36-3]

$C_2H_3N_{11}$



Hofman, K. A. *et al.*, *Ber.*, 1910, **43**, 1869–1870

The barium salt explodes weakly on heating, and the copper salt and silver salt strongly on heating or friction.

See other TETRAZOLES, TRIAZENES

0779. Sodium acetate
[127-09-3]

$C_2H_3NaO_2$



HCS 1980, 824

Diketene

See Diketene: Acids, etc.

Potassium nitrate

See Potassium nitrate: Sodium acetate

See other METAL OXONON-METALLATES

0780. Sodium peroxyacetate (Sodium ethaneperoxoate)

[64057-57-4]


 $C_2H_3NaO_3$ Humber, L. G., *J. Org. Chem.*, 1959, **24**, 1789

A sample of the dry salt exploded at room temperature.

See other PEROXOACID SALTS, METAL OXONON-METALLATES

†0781. Ethylene (Ethene)

[74-85-1]

 C_2H_4 

FPA H17, 1973; HCS 1980, 458 (cryogenic liquid), 477 (cylinder gas)

1. Waterman, H. I. *et al.*, *J. Inst. Petr. Tech.*, 1931, **17**, 505–510
2. Stull, 1977, 15–16
3. Conrad, D. *et al.*, *Chem. Ing. Tech.*, 1975, **47**, 265 (summary of full paper)
4. Luft, G. *et al.*, *Chem. Ing. Tech.*, 1978, **50**, 620–622
5. Lawrence, W. W. *et al.*, *Loss Prev.*, 1967, **1**, 10–12
6. McKay, F. F. *et al.*, *Hydrocarbon Proc.*, 1977, **56**(11), 487–494
7. Tanimoto, S., *Safety in Polyethylene Plants*, 1974, **2**, 14–21
8. Anon., *Oil Gas J.*, 1982, (1), 54
9. Watanabe, H. *et al.*, *Chem. Abs.*, 1983, **98**, 218185
10. Brown, D. G., *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 1, M1–M10, Rugby, IChE, 1983
11. Britton, L. G., *Process Safety Progr.*, 1994, **13**(3), 128
12. Britton, L. G. *et al.*, *Plant/Oper. Progr.*, 1986, **5**, 238–251
13. Gebauer, M. *et al.*, E. Ger. Pats DD 251 260–2, *Chem. Abs.*, 1988, **109**, 55464–6
14. Britton, L. G., *Process Safety Progr.*, 1996, **15**(3), 128
15. Black, D. E., *Int. Conf. Workshop Risk Anal. Process Saf.*, 1997, 461

Explosive decomposition in absence of air occurred at 350°C under a pressure of 170 bar [1]. An experimental investigation of the decomposition of ethylene (which is moderately endothermic, H_f° (g) +52.3 kJ/mol, 2.18 kJ/g) at 68 bar with high energy initiation (hot wire plus guncotton) gave results which were in close agreement with those calculated from thermodynamic data (exotherm of 4.33 kJ/g, flame temperature 1360°C, 6-fold pressure increase) [2]. The limiting pressures and temperatures for explosive decomposition of ethylene with electric initiation were determined in the ranges 100–250 bar and 120–250°C. Limiting conditions are much lower for high energy (exploding wire) initiation [3]. The study of spontaneous and initiated decomposition has been extended up to 5000 bar and 450°C, in the context of high pressure polymerisation of ethylene [4]. Previously, reaction parameters including effect of pipe size and presence of nitrogen as inert diluent upon the propagation of explosive decomposition had been studied using thermite initiation: 10 vol% of nitrogen markedly interferes with propagation [5].

Sudden pressurising with ethylene of part of an air-containing pipeline system from 1 to 88.5 bar led to adiabatic compressive heating and autoignition of the ethylene. The decomposition flame slowly propagated upstream into the main 30 cm pipeline and subsequent rupture of the latter occurred at an estimated wall temperature of 700–800°C. The AIT for ethylene in air at 1 bar is 492°C, but this falls with increase in pressure and could lie between 204 and 371°C at 68–102 bar, and the ethylene decomposition reaction may start at 315–371° within these pressure limits. Detonation was not, however, observed. Precautions against such incidents in ethylene installations are discussed [6]. In a published symposium mainly devoted to engineering aspects of large-scale polyethylene manufacture, this paper describes an extensive large-scale field experiment to determine the blast and destructive effects when various ethylene–air mixtures were detonated to simulate a compressor house explosion [7]. A fire and explosion in an ethylene separator was attributed to ingress of air which formed a catalytically active species which initiated exothermic polymerisation eventually leading to pipe rupture [8]. The decomposition of ethylene at 100–400 bar/20–149°C has been studied in detail [9]. An account of a decomposition incident at a high density polythene plant which was considered safe because it operated with ethylene diluted in a solvent is available [15].

In a review of low level ethylene decomposition since its discovery in 1842, the main factors are recognised as pressure–temperature–ignition relationships, heat balance in reaction zone and presence of impurities or contaminants. Case histories are discussed and safety practices and general recommendations are presented [10]. In a survey of incidents on high pressure ethylene plant, most appeared to involve overheating on molecular sieves in driers from polymerisation under conditions of reduced flow. Large pored sieves are not advisable [11]. In a review of the stability of ethylene at elevated pressures, the factors of thermal runaway, catalysis, influence of contaminants, and adsorptive heating, compressive ignition, and direct ignition are discussed, the effects of decomposition being described in terms of pressure and propagation [12]. A series of 3 patents describe start-up procedures to avoid explosive decomposition of ethylene during high-pressure polymerisation at 1600 bar using peroxide initiation. The procedures involve use of ethylene–hydrogen mixtures (6–30:1) with compression in several stages, interspersed with circulation through the reactor [13]. Another study of the slow deflagrative decomposition of ethene at around 300 bar to mostly methane, hydrogen and carbon is reported. Addition of hydrogen facilitates ignition, but not propagation, and increases the proportion of methane in the product. Hot spots, thence initiation, may develop over finely divided iron, starting from hydrogen reduction of rust from as low as 70°C [14].

Acetylene, Hydrogen

See Hydrogen (Gas): Acetylene, etc.

Air, Chlorine

Fujiwara, T. *et al.*, Jap. Pat., 74 32 841, 1974

During oxychlorination of ethylene to 1,2-dichloroethane, excess hydrogen chloride is used to maintain the reaction mixture outside the explosive limits.

Air, Polyethylene

Fonin, M. F. *et al.*, *Chem. Abs.*, 1984, **101**, 111718

The solubility of ethylene in freshly prepared polyethylene, and its diffusion out of the latter were studied in relation to the formation of explosive ethylene–air mixtures in storage. Explosive mixtures may be formed, because the solubility of ethylene in its polymer (e.g. 1130 ppm w/w at 30°C) considerably exceeds the concentration (30 ppm at 30°C) necessary to exceed the lower explosive limit above the gas-containing polymer in closed storage, and the diffusion coefficient is also 30% higher than for aged polymer samples.

Aluminium chloride, Catalysts

Waterman, H. I. *et al.*, *J. Inst. Pet.*, 1947, **33**, 254

Mixtures of ethylene and aluminium chloride, initially at 30–60 bar, rapidly heat and explode in presence of supported nickel catalysts, methyl chloride or nitromethane.

See Aluminium chloride: Alkenes

Aluminium chloride, Nitromethane

See Aluminium chloride–nitromethane: An alkene

Boron trioxide, Oxygen

Ts'olkovskii, T. I. *et al.*, *Chem. Abs.*, 1979, **90**, 8511

AITs for ethylene–oxygen mixtures at 1 bar in stainless steel were reduced by 30–40°C by coating the vessel walls with boron trioxide.

Bromotrichloromethane

MRH 1.21/90

Elsner, H. *et al.*, *Angew. Chem.*, 1962, **74**, 253

Following a literature method for preparation of 1-bromo-3,3,3-trichloropropane, the reagents were being heated at 120°C/51 bar. During the fourth preparation, a violent explosion occurred.

See Halocarbons, below

Carbon tetrachloride

MRH 1.55/85

Zakaznov, V. F. *et al.*, *Khim. Prom.*, 1968, **8**, 584

Mixtures of ethylene and carbon tetrachloride can be initiated to explode at temperatures between 25 and 105°C and pressures of 30–80 bar, causing a six-fold pressure increase. At 100°C and 61 bar, explosion initiated in the gas phase propagated into the liquid phase. Increase of halocarbon conc. in the gas phase decreased the limiting decomposition pressure.

See Halocarbons, below, *also* Dibenzoyl peroxide: Carbon tetrachloride, etc.

Chlorine

491M, 1975, 103

C₂H₄

Interaction is explosive when catalysed by sunlight or UV irradiation, or in presence of mercury(I) oxide, mercury(II) oxide or silver oxide.

See Chlorine: Hydrocarbons

Chlorotrifluoroethylene

Colombo, P. *et al.*, *J. Polymer Sci.*, 1963, **B1**(8), 435–436

Mixtures containing ratios of about 20:1 and 12:1 of ethylene:haloalkene undergoing polymerisation under gamma irradiation at 308 krad/h exploded violently after a total dose of 50 krad. Dose rate and haloalkene conc. were both involved in the initiation process.

See Halocarbons, below

Copper

Dunstan, A. E. *et al.*, *J. Soc. Chem. Ind.*, 1932, **51**, 1321

Polymerisation of ethylene in presence of metallic copper becomes violent above a pressure of 54 bar at about 400°C, much carbon being deposited.

Halocarbons

See Bromotrichloromethane, and Carbon tetrachloride, and Chlorotrifluoroethylene, all above; Tetrafluoroethylene, below

Lithium

See Lithium: Ethylene

Molecular sieve

See MOLECULAR SIEVE INCIDENTS: Ethylene

Other reactants

Yoshida, 1980, 46

The MRH values calculated and given for 5 combinations are noted above and below.

Oxides of nitrogen

See Nitrogen oxide: Dienes, Oxygen

Ozone

See Ozone: Combustible gases, or: Ethylene

Steel-braced tyres

Anon., *CISHC Chem. Safety Summ.*, 1974–5, **45–46**, 2–3

Two hours after a road tanker had crashed, causing the load of liquid ethylene to leak, one of the tyres of the tanker burst and ignited the spill, eventually causing the whole tanker to explode. The tyre failed because it froze and became embrittled, and it is known that such failure of steel-braced tyres gives off showers of sparks. This could therefore be a common ignition source in cryogenic transportation spillage incidents.

See other IGNITION SOURCES

Tetrafluoroethylene

Coffman, D. D. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 492

A violent explosion occurred when a mixture of tetrafluoroethylene and excess ethylene was heated at 160°C and 480 bar. Traces of oxygen must be rigorously excluded. Other alkenes reacted smoothly.

See Oxygen (Gas): Tetrafluoroethylene, *also* Halocarbons, above

Trifluoromethyl hypofluorite

See Trifluoromethyl hypofluorite: Hydrocarbons

Vinyl acetate

Albert J. *et al.*, *Chem. Eng. Process*, 1998, **37**(1), 55

At pressures above 25 Mpa and temperatures above 250°C, vinyl acetate admixture destabilises ethylene and increases the maximum explosion pressure from its decomposition.

See other ALKENES

0782. Poly(ethylene)

[9002-88-4]

(C₂H₄)_n

Complex structure

Experimental investigation of poly(ethylene) powder explosions.

See entry DUST EXPLOSION INCIDENTS (reference 18)

0783. 1-Bromoaziridine

[19816-89-8]

C₂H₄BrN



Graefe, A. F., *J. Amer. Chem. Soc.*, 1958, **80**, 3940

The compound is very unstable and always decomposes, sometimes explosively, during or shortly after distillation.

See other AZIRIDINES, *N*-HALOGEN COMPOUNDS

0784. *N*-Bromoacetamide

[79-15-2]

C₂H₄BrNO

H₃CCO.NHBr

‘Organic Positive Bromine Compounds’, Brochure, Boulder (Co.) Arapahoe Chemicals Inc., 1962


It tends to decompose rapidly at elevated temperatures in presence of moisture and light.

See other *N*-HALOGEN COMPOUNDS

0785. 1,2-Dibromoethane (Ethylene dibromide)

[106-93-4]

$C_2H_4Br_2$



HCS 1980, 365

Magnesium

See Magnesium: Halocarbons

See other HALOALKANES

0786. 1-Chloroaziridine

[25167-31-1]

C_2H_4ClN



1. Davies, C. S., *Chem. Eng. News*, 1964, **42**(8), 41

2. Graefe, A. F., *Chem. Eng. News*, 1958, **36**(43), 52

A sample of redistilled material exploded after keeping in an amber bottle at ambient temperature for 3 months [1]. A similar sample had exploded very violently when dropped [2].

See other AZIRIDINES, *N*-HALOGEN COMPOUNDS

0787. Chloroacetaldehyde oxime

[51451-05-9]

C_2H_4ClNO



1. Brintzinger, H. *et al.*, *Chem. Ber.*, 1952, **85**, 345

2. Anon., *Chem. Eng. News*, 1993, **71**(34), 6

Vacuum distillation of the product at 61°C/27 mbar must be interrupted when a solid separates from the residue, to avoid an explosion [1]. A serious explosion occurred in a cooling loop containing the oxime and hydrochloric acid, during shutdown to clear a blockage elsewhere in a pesticide plant [2].

See Bromoacetone oxime


See other OXIMES

See related ALDEHYDES

0788. Chloroacetamide

[79-07-2]

C_2H_4ClNO



Energy of decomposition (in range 170–300°C) measured as 0.67 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

0789. N-Chloroacetamide
[598-49-2]



Muir, G. D., private comm., 1968

It has exploded during desiccation of the solid or during concentration of its chloroform solution. It may be purified by pouring a solution in acetone into water and air-drying the product.

See other N-HALOGEN COMPOUNDS

†0790. 1,1-Dichloroethane
[75-34-3]



HCS 1980, 494

See other HALOALKANES

†0791. 1,2-Dichloroethane (Ethylene dichloride)
[107-06-2]



(MCA SD-18, 1971); NSC 350, 1977; FPA H77, 1979; HCS 1980, 482

Chlorine

See Chlorine: 1,2-Dichloroethane

3-Dimethylaminopropylamine

491M, 1975, 171

After the amine was charged into a tank containing some residual wet halocarbon, a violent explosion occurred later. Subsequent investigation showed it was an extremely hazardous combination (as dehydrochlorination of the solvent would give acetylene).

Dinitrogen tetroxide

See Dinitrogen tetroxide: Halocarbons

Metals

See Aluminium: Halocarbons

Potassium: Halocarbons

Nitric acid

See Nitric acid: 1,2-Dichloroethane

See other HALOALKANES

- 0792. Azo-*N*-chloroformamidine**
 [502-98-7] **C₂H₄Cl₂N₆**

$$\text{H}_2\text{NC}(:\text{NCl})\text{N}=\text{NC}(:\text{NCl})\text{NH}_2$$
- Braz, G. I. *et al.*, *Chem. Abs.*, 1946, **40**, 2267.9
 Decomposition is explosive at 155°C, and accelerated by contact with metals.
See other AZO COMPOUNDS, *N*-HALOGEN COMPOUNDS
- †0793. Bis(chloromethyl) ether**
 [542-88-1] **C₂H₄Cl₂O**

$$\text{ClCH}_2\text{OCH}_2\text{Cl}$$
- HCS 1980, 377; RSC Lab. Hazards Data Sheet No. 46, 1986*
- 0794. 1,1-Dichloroethyl hydroperoxide**
 [90584-32-0] **C₂H₄Cl₂O₂**

$$\text{MeCCl}_2\text{OOH}$$
- Gäb, S. *et al.*, *J. Org. Chem.*, 1984, **49**, 2714
 Samples of the oil, either neat or as conc. solutions, rapidly decomposed exothermally on warming to ambient temperature. Although no explosions were experienced, caution is advised.
See Ozone: *trans*-2,3-Dichloro-2-butene
See other ALKYL HYDROPEROXIDES
- 0795. Ethylene diperchlorate**
 [52936-25-1] **C₂H₄Cl₂O₈**

$$(-\text{CH}_2\text{OCIO}_3)_2$$
- Schumacher, 1960, 214
 A highly sensitive, violently explosive material, capable of detonation by addition of a few drops of water.
See other ALKYL PERCHLORATES
- 0796. Bis(chloromethyl)thallium chloride**
 [] **C₂H₄Cl₃Tl**

$$(\text{ClCH}_2)_2\text{TlCl}$$
- Yakubovich, A. Ya. *et al.*, *Dokl. Akad. Nauk SSSR*, 1950, **3**, 957
 An explosive solid of low stability.
See related ALKYL METAL HALIDES

0797. 1,2-Bis(dichlorophosphino)ethane
[28240-69-9]




Chatt, J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1985, 1131–1136

A modified method of preparation from ethylene, phosphorus trichloride and phosphorus in a stainless autoclave gives a highly pyrophoric reaction residue.

See related ALKYLHALOPHOSPHINES, PYROPHORIC MATERIALS

0798. Ethylenedicaesium
[65313-36-2]



Leleu, *Cahiers*, 1977, (88), 366

Reacts violently with water.

See related ALKYLMETALS

0799. Hydroxycopper(II) glyoximate
[63643-78-7]




Morpurgo, G. O. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1977, 746

The complex loses weight up to 140°C, then explodes.

See related OXIMES *See other* HEAVY METAL DERIVATIVES

0800. 2-Fluoro-2,2-dinitroethylamine
[18139-02-1]



Adolph, H. G. *et al.*, *J. Org. Chem.*, 1969, **34**, 47

Outstandingly explosive among fluorodinitromethyl compounds, samples stored neat at ambient temperatures regularly exploded within a few hours. Occasionally, concentrated solutions in dichloromethane have decomposed violently after long storage.

See other FLUORODINITROMETHYL COMPOUNDS

†0801. 1,1-Difluoroethane
[75-37-6]



HCS 1980, 408 (cylinder)

See other HALOALKANES

- 0802. 1,2-Bis(difluoroamino)ethanol**
 [13084-47-4] **C₂H₄F₄N₂O**

$$\text{F}_2\text{NCH}_2\text{CH}(\text{NF}_2)\text{OH}$$
- Reed, S. F., *J. Org. Chem.*, 1967, **32**, 2894
 It is slightly more impact-sensitive than glyceryl trinitrate.
See other DIFLUOROAMINO COMPOUNDS
- 0803. 1,2-Bis(difluoroamino)-N-nitroethylamine**
 [18273-30-8] **C₂H₄F₄N₄O₂**

$$\text{F}_2\text{NCH}_2\text{CH}(\text{NF}_2)\text{NHNO}_2$$
- Tyler, W. E., US Pat. 3 344 167, 1967
 The crude product tends to explode spontaneously on storage, though the triple-distilled material appears stable on prolonged storage. Generally, such nitroamines are unstable and explode at 75°C or above.
See other CATALYTIC IMPURITY INCIDENTS, DIFLUOROAMINO COMPOUNDS, N-NITRO COMPOUNDS
- 0804. Mercury(II) formohydroxamate**
 [] **C₂H₄HgN₂O₄**

$$\text{Hg}(\text{ONHCO.H})_2 \text{ or } \text{Hg}(\text{ON=CHOH})_2$$
- Urbanski, 1967, Vol. 3, 158
 It possesses detonator properties.
See other MERCURY COMPOUNDS, N-O COMPOUNDS
- 0805. Bis(iodomethyl)zinc**
 [14399-53-2] **C₂H₄I₂Zn**


$$\text{ICH}_2\text{ZnCH}_2\text{I}$$
- Dichloromethane
 Charette, A. *Chem. Eng. News*, 1995, **73**(6), 2
 An explosion was twice experienced when working with a suspension of this compound in dichloromethane at near ambient temperature.
See Copper-zinc alloys: Diiodomethane, Ether
See also MAGNETIC STIRRERS
- 0806. Potassium 1-nitroethoxide**
 [] **C₂H₄KNO₃**

$$\text{KOCH}(\text{NO}_2)\text{CH}_3$$
- Meyer, V. *et al.*, *Ber.*, 1872, **5**, 1032

The solid exploded on heating, and an aqueous solution sealed into a Carius tube exploded violently when the tube was cracked open with a red-hot glass rod. (The *aci*-salt may have been formed).

See other C-NITRO COMPOUNDS

0807. 2-Nitroethanol, sodium salt
[2406-51-1]



Beilstein, **H1**, 339
Deflagrates at 120°C.

See other C-NITRO COMPOUNDS

0808. 3-Methyldiazirine
[765-31-1]



Schmitz, E. *et al.*, *Chem. Ber.*, 1962, **95**, 795

The gas explodes on heating.

See other DIAZIRINES

0809. 2-Nitroacetaldehyde oxime ('Methazonic acid')
[5653-21-4]



1. Sorbe, 1968, 149
2. Cooke, F., *Chem. Eng. News*, 1981, **59**(34), 3
3. Hoglen, D. K. *et al.*, Personal communication, 1997
4. Editor's comments

The oxime (an isomer of 'ethylnitrolic acid') decomposes gradually at ambient temperature, but explosively above 110°C [1]. The product from treatment of nitromethane with strong sodium hydroxide solution at 50°C, followed by acidification and ether extraction, gave, after vacuum evaporation, a residue which exploded when air was admitted. On this occasion, a weekend elapsed between alkali and acid treatments, and it seems possible that formation of fulminic acid derivatives may have occurred [2]. A 75 g sample of methazonic acid, from a preparation which had transiently overheated to 70°C, during the addition of nitromethane to sodium hydroxide solution, was stored for a week at -15°C, then allowed to warm to room temperature. Twelve hours later it spontaneously detonated, destroying a fume hood. There is no evident cause for this explosion [3]. It seems sensible to

apply Occam's Razor, assume that methazonic acid is itself explosive, and cease to invoke impurities [4]. The salts of methazonic acid are also explosive.

See Nitromethane: Acids, or Bases; Alkalis

See other OXIMES

See related ALDEHYDES

0810. 1-Nitro-1-oximinoethane ('Ethylnitrolic acid')

[600-26-0]



Sorbe, 1968, 147

An explosive solid, formally a nitro-oxime.

See other OXIMES

0811. Ethylidene dinitrate

[55044-04-7]



Kacmarek, A. J. *et al.*, *J. Org. Chem.*, 1975, **40**, 1853


An analytical sample exploded in the hot zone of the combustion apparatus.

See Dinitrogen pentaoxide: Acetaldehyde

See other ALKYL NITRATES

0812. 4-Amino-4H-1,2,4-triazole

[584-13-4]



Preparative hazard

Tomann, J. *et al.*, *Chem. Prum.*, 1987, **37**, 489–492

The title compound (with 66.5% nitrogen content) is prepared by condensing formylhydrazine (2 mols, with elimination of 2H₂O) by heating to 170°C. During a pilot production run in a 500 l reactor, an explosion destroyed the vessel. The heat of decomposition of the compound was determined by thermal analysis as 1.5 kJ/g, with an energy of activation of 91 kJ/mol.

See other HIGH-NITROGEN COMPOUNDS, TRIAZOLES

0813. Cyanoguanidine ('Dicyanodiamide')

[461-58-5]



HCS 1980, 386


Oxidants

Baumann, J., *Chem. Ztg.*, 1920, **44**, 474

Mixtures of cyanoguanidine with ammonium nitrate, potassium chlorate, etc., were formerly proposed for use as powerful explosives.

See other CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

0814. 2-Methyltetrazole
[16681-78-0]





Aluminium hydride

See Aluminium hydride: Tetrazole derivatives

See other TETRAZOLES

0815. Azoformaldoxime
[74936-21-3]



Armand, J. *et al.*, *J. Chem. Res.; Synop.*, 1980, 304; *Microf.* 1980, 3853–3869

The dioxime exploded on melting, as did the cyclised isomeric triazole derivative, 1-hydroxy-2-hydroxylamino-1,3,4-triazole.

See other AZO COMPOUNDS, N–O COMPOUNDS, OXIMES

0816. Azoformamide (Azodicarboxamide)
[123-77-3]



Whitmore, M. W. *et al.*, *J. Loss Prev.*, 1993, **6**(3), 169

An 850 kg batch of a slightly doped form of azodicarbonamide exploded violently, with a TNT equivalence of 3.3 kg, 5 minutes after sampling at the end of drying. The probable initial temperature was 65°C, the lowest self accelerating decomposition temperature 90°C, and such decomposition is not explosive. Full explosibility tests, including detonability, had shown no hazard. Further study demonstrated that slightly contained azodicarboxamide, thermally initiated at the bottom of a column or conical vessel could explode even at the 5 kg scale. The above TNT equivalence corresponds to decomposition of 4% of the available charge. The cause of the presumptive hot spot is unknown.

Energy of decomposition (in range 160–230°C) measured as 1.36 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 120°C by adiabatic Dewar tests, with an apparent energy of activation of 131 kJ/mol.

See entries BLOWING AGENTS, PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


Potassium bromate

See Potassium bromate: Azoformamide

See other AZO COMPOUNDS

0817. 1-Hydroxy-2-hydroxylamino-1,3,4-triazole
[76002-04-5]

$C_2H_4N_4O_2$



Armand, J. *et al.*, *J. Chem. Res.; Synop.*, 1980, 304; *Microf.*, 1980, 3853–3869

The triazole exploded on melting, as did the isomeric azoformaldoxime, from which it is prepared by cyclisation.

See other HIGH-NITROGEN COMPOUNDS, N–O COMPOUNDS, TRIAZOLES

0818. 1,1-Diazidoethane
[67880-20-0]

$C_2H_4N_6$



Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 1070

The extreme instability and explosive behaviour of this diazide caused work on other *gem*-diazides to be abandoned.

See other ORGANIC AZIDES

0819. 1,2-Diazidoethane
[629-13-0]

$C_2H_4N_6$



Alone, or Sulfuric acid

Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 1070

Though less unstable than the 1,1-isomer (next above), it explodes on heating, and irreproducibly in contact with sulfuric acid.

See other ORGANIC AZIDES

0820. Azidocarbonylguanidine
[54567-24-7]

$C_2H_4N_6O$




Thiele, J. *et al.*, *Ann.*, 1898, **303**, 93

It explodes violently on rapid heating.


See related ACYL AZIDES

0821. 3,6-Diamino-1,2,4,5-tetrazine-1-oxide
[153757-93-8]




See 3,6-Diamino-1,2,4,5-tetrazine-1,4-dioxide
See other HIGH-NITROGEN COMPOUNDS, See also AMINE OXIDES

0822. 3,6-Diamino-1,2,4,5-tetrazine-1,4-dioxide
[153757-93-8]




Coburn, M. D. *et al.*, *J. Heterocycl. Chem.*, 1993, **30**(6), 1593
This, the mono-oxide and the 3-amino-6-nitro-2,4-dioxide are all explosives to be handled with care. All are thermally stable well above 100°C.
See other HIGH-NITROGEN COMPOUNDS, See also AMINE OXIDES

0823. Ammonium 3,5-dinitro-1,2,4-triazolide
[76556-13-3]



Selig, W., *Propellants Explos.*, 1981, **6**, 96-98
It is an explosive.
See other TRIAZOLES

0824. 1,3-Diazido-2-nitroazapropene (*N,N*-Bis(azidomethyl)nitric amide)
[67362-62-3] **C₂H₄N₈O₂**




Flanagan, J. E. *et al.*, US Pat. 4 085 123, 1978
It is useful as a high energy plasticiser for propellants.
See other N-NITRO COMPOUNDS, ORGANIC AZIDES

0825. Azo-*N*-nitroformamidine
[53144-64-2] **C₂H₄N₈O₄**




Wright, G. F., *Can. J. Chem.*, 1952, **30**, 64
Explosive decomposition at 165°C.
See other AZO COMPOUNDS, N-NITRO COMPOUNDS

0826. 5,5'-Hydrazotetrazole
[74999-19-2] **C₂H₄N₁₀**



1. Thiele, J. *et al.*, *Ann.*, 1898, **303**, 66
2. Reddy, G. O., *J. Haz. Mat.*, 1984, **9** 291–303
It explodes without melting when heated [1]. The title compound and its barium, lead(II), and mercury(II) salts were studied by DTA and DSC techniques. The lead salt is the least stable and most powerful primary explosive of the 4 compounds [2].
See other HIGH-NITROGEN COMPOUNDS, TETRAZOLES

0827. 1,6-Bis(5-tetrazolyl)hexaaza-1,5-diene
[68594-19-4] **C₂H₄N₁₄**



Hofman, K. A. *et al.*, *Ber.*, 1911, **44**, 2953
This very high-nitrogen compound (87.5%) explodes violently on pressing with a glass rod, or on heating to 90°C.
See other HIGH-NITROGEN COMPOUNDS, TETRAZOLES

†0828. Acetaldehyde (Ethanal)

[75-07-0]

C₂H₄O

H₃CCO.H

(MCA SD-43, 1952); FPA H29, 1974; HCS 1980, 97; RSC Lab. Hazards Data Sheet No. 38, 1985

1. White, A. G. *et al.*, *J. Soc. Chem. Ind.*, 1950, **69**, 206
2. Vervalin, 1973, 90–91
3. Garbuzynk, T. A., *Chem. Abs.*, 1978, **89**, 131835; 1989, **111**, 117216b
4. Nalbandyan, A. B. *et al.*, *Prog. Astronaut. Aeronaut.*, 1988, **113**, 58; *Chem. Abs.*, 1989, **111**, 22912; *Khim. Fiz.*, 1988, **7**(12), 1709; *Chem. Abs.*, 1989, **111**, 56791

(The MCA Data Sheet describes acetaldehyde as extremely or violently reactive with: acid anhydrides, alcohols, halogens, ketones, phenols, amines, ammonia, hydrogen cyanide or hydrogen sulfide). Mixtures of 30–60% of acetaldehyde vapour with air (or 60–80% with oxygen) may ignite on surfaces at 176 and 105°C, respectively, owing to formation and subsequent violent decomposition of peroxyacetic acid [1]. Acetaldehyde vapour leaking into a building equipped only with flameproof electrical equipment nevertheless ignited, possibly on contact with rusted steel, corroded aluminium or hot steam lines [2]. The minimum AIT (130°C) for acetaldehyde in air mixtures is at 55–57 vol% of aldehyde. The effect of presence of acetone or methane upon AIT was studied, they inhibit ignition during the oxidation to peroxyacetic acid; the AIT is surface sensitive and may be as low as 80°C in quartz [3]. The mechanism of this ignition has been studied [4], it is catalysed by peroxy-acids on the surface of the reactor.

See other CATALYTIC IMPURITY INCIDENTS

Acetic acid

MRH 1.13/tr.

MCA Case History No. 1764

A drum contaminated with acetic acid was filled with acetaldehyde. The ensuing exothermic polymerisation reaction caused a mild eruption lasting several hours.

Cobalt acetate, Oxygen

1. Phillips, B. *et al.* *J. Amer. Chem. Soc.*, 1957, **79**, 5982
 2. Bloomfield, G. F. *et al.*, *J. Soc. Chem. Ind.*, 1935, **54**, 129T
- Oxygenation of acetaldehyde in presence of cobalt acetate at –20°C caused precipitation of 1-hydroxyethyl peroxyacetate (acetaldehyde hemi-peracetate), which exploded violently on stirring [1]. Ozone or UV light also catalyses the autoxidation [2].

Desiccants, Hydrogen peroxide

See Hydrogen peroxide: Acetaldehyde, Desiccants

Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Acetaldehyde

Halocarbons

Jones, J. C. *et al.*, *Combust. Flame*, 1983, **52**, 211–213

C₂H₄O

Cool flame behaviour of acetaldehyde is apparently eliminated by *tert*-butyl bromide, and reduced by methyl iodide.

Hydrogen peroxide MRH 6.27/79
See Hydrogen peroxide: Oxygenated compounds

Mercury(II) oxosalts
Sorbe, 1968, 97
Some of the products of interaction of acetaldehyde and mercury(II) chlorate or mercury(II) perchlorate are highly explosive and extremely shock-sensitive.
See Chloratomercurio(formyl)methylenemercury(II)

Metals
Sorbe, 1968, 103
Impure material will polymerise readily in presence of trace metals (iron) or acids.
See Sulfuric acid, below
See other CATALYTIC IMPURITY INCIDENTS

Other reactants
Yoshida, 1980, 17
MRH values calculated for 18 combinations, largely with oxidants, are given.

Oxygen MRH 8.66/64
1. *MCA Case History No. 117*
2. Armstamyan, A. M. *et al.*, *Chem. Abs.*, 1984, **101**, 194603
Oxygen leaked into the free space in an acetaldehyde storage tank normally purged with nitrogen. Accelerating exothermic oxidation led to detonation [1]. The self-ignition temperature of acetaldehyde–oxygen mixtures depends on dimensions of the reactor and the partial pressure of peracetic acid accumulated on the walls. Spontaneous ignition temperatures of 71–73°C were observed [2].

Silver nitrate
See Silver nitrate: Acetaldehyde

Sulfuric acid MRH 1.34/43
Sorbe, 1968, 103
Acetaldehyde is polymerised violently by the conc. acid.
See Metals, above
See other POLYMERISATION INCIDENTS *See other* ALDEHYDES, PEROXIDISABLE COMPOUNDS

†0829. Ethylene oxide (Oxirane) [75-21-8]

C₂H₄O



(*MCA SD-38*, 1971); *FPA H32*, 1974; *HCS 1980*, 486; *RSC Lab. Hazard Data Sheet No. 2*, 1982

1. *Guidelines for Bulk Handling of Ethylene Oxide*, London, CIA, 1983
2. Hess, L. G. *et al.*, *Ind. Eng. Chem.*, 1950, **42**, 1251
3. Burgoyne, J. H. *et al.*, *Proc. 1st Symp. Chem. Proc. Hazards*, Symp. Ser. No. 7, 30–36, London, IChE, 1960; *Proc. 3rd Symp. Chem. Proc. Hazards*, Symp. Ser. No. 25, 1–7, London, IChE, 1967
4. *MCA SD-38*, 1971
5. *MCA Case History No. 1666*
6. Harmon, 1974, 2.9
- 6a Anon., *Loss Prev. Bull.*, 1991, **100**, 1
- 6b Mellin, B. E., *ibid.*, 13
- 6c Viera, G. A. *et al.*, *Chem. Eng. Progress*, 1993, **89**(8), 66
7. Heuser, S. G. B., *Chem. Brit.*, 1977, **13**, 317
8. Pogany, G. A., *Chem. & Ind.*, 1979, 16–21
9. Pesetsky, B. *et al.*, *Loss Prev.*, 1980, **13**, 123–131
10. Pesetsky, B. *et al.*, *Loss Prev.*, 1980, **13**, 132–141
11. Chen, L. D. *et al.*, *Combust. Flame*, 1981, **40**, 13–28
12. *See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION* (reference 2)
13. Loeffler U., *Prax. Sicherheitstechn.*, 1997, **4**, 31
14. Britton, L. G., *Plant/Oper. Progr.*, 1990, **9**(2), 75

The detailed code of practice [1] serves as a comprehensive guide to the industrial use of ethylene oxide to replace the US Safety Data Sheet withdrawn in 1980. Ethylene oxide vapour may readily be initiated into explosive decomposition in absence of air, and storage and handling requirements were detailed [2]. Explosive decomposition may be suppressed by many diluents, not necessarily inert [3]. Metal fittings containing copper, silver, mercury or magnesium should not be used in ethylene oxide service, since traces of acetylene in the oxide could produce metal acetylides capable of detonating the vapour [4]. Ethylene oxide exposed to heating and subsequently cooled (e.g. by exposure to fire conditions) may continue to polymerise exothermically, leading to container pressurisation and explosion. A mechanism consistent with several observed incidents is proposed [5]. Presence of hot spots in processing plants is identified as a particular hazard [6]. These can be generated by the polymerisation/autoxidation of leaks absorbed onto lagging material, with devastating results [6a, 6b, 13], or by catalysis of decomposition by γ -ferric oxide and γ -FeO(OH) [6c]. Appropriate column packing can inhibit propagation of ethylene oxide decomposition from hot spots in ethylene oxide stills [13]. Combustion properties, including ignition of leaks onto packing are reviewed thoroughly in [14]. The autoignition temperature may be as low as 140°C in presence of rust. Aerosols, detonated rather than ignited, are used in fuel/air weapons, giving a TNT equivalence of 2.4 kg/kg. Under conditions of ignition and deflagration TNT equivalence is more like zero.

Hazards attendant on use of ethylene oxide in steriliser chambers arise from difficulties in its subsequent removal by evacuation procedures, owing to its ready absorption or adsorption by the treated material. Even after 2 evacuation cycles the oxide may still be present. Safety is ensured by using the oxide diluted with up to 90% of Freon or carbon dioxide. If high concentrations of oxide are used, an inert gas purge between cycles is essential [7]. The main factors in safe handling

C₂H₄O

and use on laboratory or small pilot plant scales have been identified [8]. Liquid phase decomposition, previously thought not possible, was observed under conditions of temperature and pressure outside the normal working range with different submerged igniter wires, and a mechanism is discussed [9]. The use of methane as a diluent for ethylene oxide has been studied in detail [10]. The decomposition threshold temperature for ethylene oxide has been related to pressure, ignition source and geometry of the containment vessel, pressure having the greatest effect [11]. The energy of decomposition (in range 320–490°C) has been revised to 1.516 kJ/g [12].

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other POLYMERISATION INCIDENTS

See also VIOLENT POLYMERISATION; INSULATION

Air, Bromomethane

Baratov, A. N. *et al.*, p.24 of BLL translation (628.74, issued 1966) of Russian book on 'Fire Prevention and Firefighting Symposium'. The addition of bromomethane to ethylene oxide (used for germicidal sterilising) to reduce the risk of explosion is relatively ineffective, the inhibiting concentration being 31.2, as against 5.8 for hexane and 13.5 vol.% for hydrogen.

Air, Ethylene

See Oxygen, below

Alkanethiols, or An alcohol

1. Meigs, D. P., *Chem. Eng. News*, 1942, **20**, 1318
2. Personal experience (PGU)

Autoclave reactions involving ethylene oxide with alkanethiols or an (unspecified) alcohol went out of control and exploded violently. Similar previous reactions had been uneventful [1]. An arenethiol was being reacted with ethylene oxide under catalysis by a fraction of a percent of sodium hydroxide (solid) dissolved in the thiol to which the oxirane was slowly charged. After an initial exotherm a white solid precipitated, the exotherm died away and later resumed, with dissolution of the solid, the reaction then running out of control from the backlog of charged oxirane [2].

Ammonia

1. *MCA Case History No. 792*
2. Troyan, J. E. *et al.*, *Loss Prev.*, 1968, **2**, 125–130
3. Stull, 1977, 26–27

Accidental contamination by aqueous ammonia of an ethylene oxide feed tank containing 22 t caused violent polymerisation which ruptured the tank and led to a devastating vapour cloud explosion [1,2]. The close similarity to other base-catalysed incidents was stressed [3].

See Contaminants, and Trimethylamine, both below

See other POLYMERISATION INCIDENTS, RUNAWAY REACTIONS, VIOLENT POLYMERISATION

Contaminants

1. Gupta, A. K., *J. Soc. Chem. Ind.*, 1949, **68**, 179
2. Stull, 1977, 26–27
3. Freeder, B. G. *et al.*, *J. Loss Prev. Process Ind.*, 1988, **1**, 164–168

Accidental contamination of a 90 kg cylinder of ethylene oxide with a little sodium hydroxide solution led to explosive failure of the cylinder over 8 hours later [1]. Based on later studies of the kinetics and heat release of the polycondensation reaction, it was estimated that after 8 hours and 1 min, some 12.7% of the oxide had condensed with an increase in temperature from 20 to 100°C. At this point the heat release rate was calculated to be 2.1 MJ/min, and 100 s later the temperature and heat release rate would be 160° and 1.67 MJ/s respectively, with 28% condensation. Complete reaction would have been attained some 16 s later at a temperature of 700°C [2]. Precautions designed to prevent explosive polymerisation of ethylene oxide are discussed, including rigid exclusion of acids; covalent halides, such as aluminium chloride, iron(III) chloride, tin(IV) chloride; basic materials like alkali hydroxides, ammonia, amines, metallic potassium; and catalytically active solids such as aluminium oxide, iron oxide, or rust [1]. A comparative study of the runaway exothermic polymerisation of ethylene oxide and of propylene oxide by 10 wt% of solutions of sodium hydroxide of various concentrations has been done using ARC. Results below show onset temperatures/corrected adiabatic exotherm/maximum pressure attained; and heat of polymerisation for the least (0.125 M) and most (1 M) concentrated alkali solutions used as catalysts.

EO 55.0, 22.5°C/439, 415°/40.5, 44.6 bar; 1.06, 1.0 kJ/g.

PO 91, 53°/451, 452°/26.6, 20.9 bar; 1.09, 1.1 kJ/g

The more reactive ethylene oxide clearly shows much higher self-heating rates and hazard potential than its higher homologue, though more detailed work is needed to quantify the differences [3].

See Ammonia, above; Trimethylamine, below

See other CATALYTIC IMPURITY INCIDENTS, POLYCONDENSATION REACTION INCIDENTS, POLYMERISATION INCIDENTS

Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Strained ring heterocycles

Glycerol

Anon., *Loss Prev. Bull.*, 1979, (028), 92–93

Glycerol was to be ethoxylated at 115–125°C in a circulating reaction system with separate reactor, heat exchanger and catalyst units. The valve at the base of the reactor was still closed, but an inoperative flow indicator failed to indicate absence of circulation and a total of 3 tonnes of the oxide, plus glycerol, was charged to the reactor. Upon subsequent opening of the valve, the reaction mixture passed through the heater, now at 200°C, and a runaway reaction developed, the reactor burst and an explosion followed.

See other RUNAWAY REACTIONS

Iron(III) hexacyanoferrate(4–) ('Iron blue pigment')

See Iron(III) hexacyanoferrate(4–): Ethylene oxide

Lagging materials

Hilado, C. J., *J. Fire Flamm.*, 1974, **5**, 321–326

The self-heating temperature (effectively the AIT) of a 50:50 mixture of ethylene oxide and air is reduced from 456°C on passage through various thermal insulation (lagging) materials to 251–416°C, depending on the particular material (of which 13 were tested).

Magnesium perchlorate

See Magnesium perchlorate: Ethylene oxide

4-Methoxybenzenelithium

Taylor, D. A. H., personal communication, 1993

Oxirane was added to the organolithium (probably in ether solution) at a temperature perhaps –40°C. The mixture was then allowed to warm up, went out of control and distributed itself widely. Presumably too much ethylene oxide had accumulated because of sluggish reaction at the low temperature.

See Alkanethiols, above

3-Nitroaniline

Anon., *Angew. Chem. (Nachr.)*, 1958, **70**, 150

Interaction of the two compounds in an autoclave at 150–160°C is described as safe in Swiss Pat. 171 721. During careful repetition of the reaction with stepwise heating, an autoclave exploded at 130°C.

Nitrogen

Grosse-Wortmann, H., *Chem. Ing. Tech.*, 1970, **42**, 85–86

Concentration boundaries for explosive decomposition of the oxide and its mixtures with nitrogen at elevated temperatures and pressures are presented graphically.

Other reactants

Yoshida, 1980, 47

MRH values calculated for 13 combinations with oxidants are given.

Polyhydric alcohol, Propylene oxide

Vervalin, 1973, 82

A polyether-alcohol, prepared by co-condensation of ethylene oxide and propylene oxide with a polyhydric alcohol, was stored at above 100°C and exposed to air via a vent line. After 10–15 h, violent decomposition occurred, rupturing the vessel. It was subsequently found that exothermic oxidation of the product occurred above 100°C, and that at 300°C a rapid exothermic reaction set in, accompanied by vigorous gas evolution.

See other GAS EVOLUTION INCIDENTS

Oxygen

Griffiths, R. F. *et al.*, *Proc. 18th Int. Combust. Symp.*, 1981, 893–901

If rapid compression of dilute ethylene oxide vapours occurs, ignition and/or decomposition may occur. Thus 5% of oxide in argon will ignite if compressed

more than 11-fold, and presence of traces of oxygen enhances the effect. Even 1% or less of ethylene in the oxide in air may be hazardous if modest but rapid compression occurs.

Pyridine, Sulfur dioxide

Nolan, 1983, Case History 151

Ethylene sulfite is prepared from ethylene oxide, pyridine, and sulfur dioxide in excess to prevent polymerisation of ethylene oxide. Use of a deficiency of sulfur dioxide led to rupture of a reactor from that cause.

Sodium hydroxide

See Contaminants, above

Sucroglyceride

1. Slade, R. C., *Chem. Brit.*, 1977, **13**, 317
2. Parker, K. J., private comm., 1977

During ethoxylation of the glyceride by stirring and heating to 120°C with excess ethylene oxide in an autoclave fitted with a 300 bar bursting disc [1], exothermic reaction occurred, usually causing an increase in temperature to 190°C (even with ice cooling applied) when large excesses were used. In one such attempt, an operating error led to the initial ethylene oxide charge being distilled out of the autoclave via an open vent valve, and the dry residue being heated to a temperature considerably above the 100°C indicated by the (unwetted) pocketed thermocouple. After cooling, more oxide was added and the preparation was continued. At an indicated temperature of 126°C and a pressure reading above 10 bar, it was decided to cool the autoclave to moderate the accelerating reaction. At this point, the rapidly increasing pressure and temperature caused the autoclave lid bolts to stretch and relieve the explosive reaction [2]. It is surmised that during the 'false start' of the reaction, a layer of the sucroglyceride became distributed as a paste round the heated autoclave wall, and was then overheated, forming an insulating layer on the autoclave wall. After more ethylene oxide was added and heating restarted, the indicated liquid temperature (126°C) would be far below that of the heated autoclave wall behind the insulating layer of thermally degraded solid. Dissolution of the inner layers of solid in the hot ethylene oxide may have exposed a hot spot which initiated thermal decomposition. Additionally or alternatively, local concentrations of alkali arising from thermal decomposition of potassium soaps present as impurity in the sucroglyceride may have initiated exothermic polymerisation of the oxide.

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Trimethylamine

1. Anon., *BCISC Quart. Safety Summ.*, 1966, **37**, 44
2. Anon., *Chem. Trade J.*, 1956, **138**, 1376

Accidental contamination of a large ethylene oxide feed-cylinder by reaction liquor containing trimethylamine caused the cylinder to explode 18 h later. Contamination was possible because of a faulty pressure gauge and suck-back of froth from above the liquid level [1]. A similar incident had occurred previously [2].

See Ammonia, and Contaminants, both above

Water

Vanderwater, R. G., *Chem. Eng. Progress*, 1989, **85**(12), 16


A supposedly empty tankwagon, which may have still held 12 tonnes of oxirane, was filled with water as the first stage of a cleaning process and the valves then closed. Several hours later, an explosion shattered the tank, sending parts up to 600 metres distance. It appears that filling will not have given complete mixing but left an oxirane layer on top of the water, which would be heated, but not mixed, by reaction with water at the interface, possibly aided by catalysis at sites of corrosion. There should have been enough water to act as a heat sink had it mixed. Explosion appears to have taken place within the supernatant ethylene oxide layer.

See other CORROSION INCIDENTS

See other 1,2-EPOXIDES

0830. Cyclic poly(ethylene oxides) ('Crown ethers')

[[294-93-9] ($n = 4$) [33100-27-5] ($n = 5$) [17455-13-9] ($n = 6$)] $(C_2H_4O)_n$



Bromoform, Potassium hydroxide

See Bromoform: Cyclic poly(ethylene oxides), etc.

0831. Poly(vinyl alcohol)

[9002-89-5]


$[-CH_2CH(OH)-]_n$

$(C_2H_4O)_n$

Energy of decomposition (in range 125–430°C) measured as 0.59 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

†0832. Thiolacetic acid
[507-09-5]



See other ORGANIC ACIDS

0833. Acetic acid (Ethanoic acid)
[64-19-7]



(MCA SD-41, 1951); NSC 410, 1979; FPA H120, 1983; HCS 1980, 99

Acetaldehyde

See Acetaldehyde: Acetic acid

Acetic anhydride, Water

See Acetic anhydride: Acetic acid, Water

5-Azidotetrazole

See 5-Azidotetrazole: Alone, or Acetic acid

Other reactants

Yoshida, 1980, 121

MRH values calculated for 15 combinations are given.

Oxidants

MRH values show % of oxidant

See Bromine pentafluoride: Hydrogen-containing materials

Chromium trioxide: Acetic acid

MRH 2.17/12

Hydrogen peroxide: Acetic acid

MRH 5.10/69

Potassium permanganate: Acetic acid

MRH 2.34/84

Sodium peroxide: Acetic acid

MRH 2.80/84

Phosphorus trichloride

See Phosphorus trichloride: Carboxylic acids (references 1,2)

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids

See other ORGANIC ACIDS

†0834. Methyl formate (Methyl methanoate)
[107-31-3]



HCS 1980, 653

Methanol, Sodium methoxide

Pond, D. M., *Chem. Eng. News*, 1982, **60**(37), 43

The crude product of reaction of methanol and carbon monoxide at 100°C/70 bar in presence of 0.5% of sodium methoxide was discharged after cooling into a storage bottle, which burst 4 h later. This was attributed to extreme instability of the ester in presence of the base, leading to the reverse reaction with vigorous evolution of carbon monoxide. Immediate neutralisation of the reaction mixture would prevent the decomposition, which also occurs with ethyl formate and base.
See other GAS EVOLUTION INCIDENTS

0835. Poly(ethylidene peroxide)

[]




Rieche, A. *et al.*, *Angew. Chem.*, 1936, **49**, 101

The highly explosive polyperoxide is present in peroxidised diethyl ether and has been responsible for many accidents during distillation of the solvent.

See other POLYPEROXIDES

0836. Ethylene ozonide (1,2,4-Trioxolane)

[289-14-5]



1. Harries, C. *et al.*, *Ber.*, 1909, **42**, 3305
2. Briner, E. *et al.*, *Helv. Chim. Acta*, 1921, **12**, 154
3. Criegee, R., *Angew. Chem.*, 1958, **65**, 398–399

It explodes very violently on heating, friction or shock [1]. Stable at 0°C, but often decomposes explosively at ambient temperature [2]. Pouring the liquid from one vessel to another initiated a violent explosion [3].

See other OZONIDES

0837. Peroxyacetic acid (Ethaneperoxoic acid)

[79-21-0]



FPA H89, 1980; *HCS 1980*, 720 (40% solution in acetic acid)

1. Swern, D., *Chem. Rev.*, 1949, **45**, 7
2. Davies, 1961, 56
3. Phillips, B. *et al.*, *J. Org. Chem.*, 1959, **23**, 1823
4. Anon., *Angew. Chem. (Nachr.)*, 1957, **5**, 178
5. Smith, I. C. P., private comm., 1973
6. *MCA Case History No. 1804*
7. Slattery, G. H., US Pat. 4 137 256, 1979

It is insensitive to impact but explodes violently at 110°C [1]. The solid acid has exploded at -20°C [2]. Safe procedures (on basis of detonability experiments) for preparation of anhydrous peracetic acid solutions in chloroform or ester solvents have been detailed [3]. However, a case of explosion on impact has been recorded [4]. During vacuum distillation, turning a ground glass stopcock in contact with the liquid initiated a violent explosion, but the grease may have been involved as well as friction [5]. During a pilot-scale preparation of a solution of the acid by adding acetic anhydride slowly to 90% hydrogen peroxide in dichloromethane, an explosion of great violence occurred. Subsequent investigation revealed that during the early stages of the addition a 2-phase system existed, which was difficult to detect because of closely similar densities and refractive indices. The peroxide-rich phase became extremely shock-sensitive when between 10 and 30% of the anhydride had been added [6]. A safer process for the vapour phase oxidation of acetaldehyde to peroxyacetic acid has been claimed [7].

See Hydrogen peroxide: Acetic acid, or: Acetic anhydride, or: Vinyl acetate

See PEROXYACIDS (references 5,6) See other GLASS INCIDENTS

Acetic anhydride

1. *MCA Case History No. 1795*

2. Augustine 1969, Vol. 2, 217

During an attempt to prepare an anhydrous 25% solution of peroxyacetic acid in acetic acid by dehydrating a water-containing solution with acetic anhydride, a violent explosion occurred. Mistakes in the operational procedure allowed heated evaporation to begin before the anhydride had been hydrolysed. Acetyl peroxide could have been formed from the anhydride and peroxyacid, and the latter may have detonated and/or catalysed violent hydrolysis of the anhydride [1]. A technique for preparing the anhydrous acid in dichloromethane without acetyl peroxide formation has been described [2].

See Acetic anhydride: Water

5-*p*-Chlorophenyl-2,2-dimethyl-3-hexanone

Gillespie, J. S. *et al.*, *Tetrahedron*, 1975, **31**, 5

During the attempted preparation of 3-*p*-chlorophenylbutanoic acid by addition of the ketone (2 g mol) to peracetic acid (50%) in acetic acid at 65–70°C, a serious explosion occurred.

Ether solvents

Augustine, 1971, Vol. 2, 159

Ether solvents, such as THF, diethyl ether etc. are unsuitable solvents for peracetic acid oxidations, as interaction of the acid with the peroxidisable solvent is violent.

See other PEROXIDATION INCIDENTS

Metal chlorides

Wienhofer, E., *Chem. Ztg.*, 1980, **104**, 146

Addition of chloride ions (as solid calcium chloride, potassium chloride or sodium chloride) to aqueous solutions containing 40% of peroxyacetic acid and 1% of

acetic acid leads to a violently exothermic decomposition reaction. Chlorine is evolved, most of the liquid evaporates and the residue (often red coloured) deflagrates.

See other GAS EVOLUTION INCIDENTS

3-Methyl-3-buten-1-yl tetrahydropyranyl ether

See TETRAHYDROPYRANYL ETHER DERIVATIVES

1-Octene

1. Swern, D. et al., *J. Amer. Chem. Soc.*, 1946, **68**, 1506

2. van den Brink, M. J., private comm., 1983

Following a published procedure [1], octene was treated with a solution of peroxyacetic acid in acetic acid for 8 h to form the epoxide, but the reaction mixture was then allowed to stand uncooled overnight. Next morning, when a 3µl sample was injected into a heated GLC injection port, the syringe shattered. This was attributed to formation of diacetyl peroxide during the overnight standing, and its subsequent explosion in the heated port [2].

Paper

See other PACKAGING INCIDENTS

See other PEROXYACIDS

0838. Hydroperoxymethyl formate (Formyloxymethyl hydroperoxide)

[]




Thamm, J. et al., *Chem. Phys. Lett.*, 1996, **258**(1,2), 155

This compound was shown to be an intermediate in ozonolysis of some ethenes. In concentrated form it is very explosive; minimal handling and working with strong gloves behind a blast screen are recommended.

See other ALKYL HYDROPEROXIDES

0839. 2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide ('Carbyl sulfate')

[503-41-3]



N-Methyl-4-nitroaniline

1. Deucker, W. et al., *Chem. Ing. Tech.*, 1973, **45**, 1040–1041

2. Wooton, D. L. et al., *J. Org. Chem.*, 1974, **39**, 2112

The title cyclic ester (1 mol) and substituted aniline (>2 mol) condense when heated in nitrobenzene solution, initially at 45°C and eventually at 75°C under

vacuum in a sealed vessel, to give the ester-salt, *N*-methyl-4-nitroanilinium 2-(*N*-methyl-*N*-4-nitrophenylaminosulfonyl)ethyl sulfate. The reaction technique had been used uneventfully during 8 years, but the 3 m³ reactor was violently ruptured when the reaction ran away and developed an internal pressure exceeding 70 bar.

Subsequent DTA investigation showed that an exothermic reaction set in above 75°C after an induction period depending on the initial temperature and concentration of reactants, which attained nearly 300°C, well above the decomposition point of the cyclic ester component (170°C). The reaction conditions used could have permitted local over-concentration and overheating effects to occur, owing to slow dissolution of the clumped solid ester and aniline in the nitrobenzene solvent [1]. Crude 'carbonyl sulfate' contains excess sulfur trioxide [2].

See DIFFERENTIAL THERMAL ANALYSIS (reference 4)

See other INDUCTION PERIOD INCIDENTS, RUNAWAY REACTIONS

†0840. Thirane (Ethylene sulfide)

[420-12-2]

C₂H₄S



Acids

Sorbe 1968, 122

The sulfide may polymerise violently in presence of acids, especially when warm.

See related 1,2-EPOXIDES

0841. Ethylaluminium bromide iodide

[32673-60-2]

C₂H₅AlBrI

EtAl(Br)I

Nitromethane

See Nitromethane: Alkylmetal halides

See other ALKYLALUMINIUM HALIDES

0842. Ethylaluminium dibromide

[2386-62-1]

C₂H₅AlBr₂

EtAlBr₂

See entry ALKYLALUMINIUM HALIDES

0843. Ethylaluminium dichloride

[563-43-9]

C₂H₅AlCl₂

EtAlCl₂

HCS 1980, 465

See entry ALKYLALUMINIUM HALIDES

- 0844. Ethylaluminium diiodide**
 [2938-73-0] **C₂H₅AlI₂**
EtAlI₂
 491M, 1975, 153
 It ignites in air.
 See other ALKYLALUMINIUM HALIDES
- 0845. Dichloroethylborane**
 [1739-53-3] **C₂H₅BCl₂**
EtBCl₂
 491M, 1975, 149
 It ignites in air.
 See other ALKYLHALOBORANES
- †**0846. Bromoethane (Ethyl bromide)**
 [74-96-4] **C₂H₅Br**
EtBr
 FPA H98, 1981; HCS 1980, 472
 Preparative hazard
 See Bromine: Ethanol, etc.
 See other HALOALKANES
- 0847. Ethylmagnesium bromide**
 [925-90-6] **C₂H₅BrMg**
EtMgBr
 Water
 Nolan, 1983, Case History 108
 Ethylmagnesium bromide was prepared as usual on a large scale in one reactor, and then transferred by nitrogen pressurisation to another. The glass transfer line had not been completely dried; the ethane evolved on contact of the Grignard reagent with moisture overpressurised the line and it burst.
 See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS
 See other GRIGNARD REAGENTS
- †**0848. Chloroethane (Ethyl chloride)**
 [75-00-3] **C₂H₅Cl**
EtCl
 (MCA SD-50, 1953); FPA H114, 1982; HCS 1980, 474

Potassium

See Potassium: Halocarbons

See other HALOALKANES

0849. 2-Chloro-*N*-hydroxyacetamide (Chloroacetamide oxime)

[3272-96-6]

C₂H₅ClN₂O



Maytum, D. M., *Chem. Brit.*, 1978, **14**, 382

During vacuum evaporation at 60°C of ethyl acetate from a solution of the oxime, a violent explosion occurred. The oxime was found to be thermally unstable above 60°C, and at the m.p., 90°C, an exotherm of over 100°C occurred accompanied by rapid gas evolution.

See related OXIMES See other GAS EVOLUTION INCIDENTS

†0850. Chloromethyl methyl ether

[107-30-2]

C₂H₅ClO



HCS 1980, 302

See related HALOALKANES

0851. Ethyl hypochlorite

[624-85-1]

C₂H₅ClO



Alone, or Copper

Sandmeyer, T., *Ber.*, 1885, **18**, 1768

Though distillable slowly (at 36°C), ignition or rapid heating of the vapour causes explosion, as does contact of copper powder with the cold liquid.

See other HYPOHALITES

0852. Ethyl perchlorate

[22750-93-2]

C₂H₅ClO₄



Sidgwick, 1950, 1235

Reputedly the most explosive substance known, it is very sensitive to impact, friction and heat. It is readily formed from ethanol and perchloric acid.

See other ALKYL PERCHLORATES

0853. 2,2-Dichloroethylamine
[5960-88-3]



$\text{C}_2\text{H}_5\text{Cl}_2\text{N}$

Roedig, A. *et al.*, *Chem. Ber.*, 1966, **99**, 121

A violent explosion occurred during evaporation of an ethereal solution at 260 mbar from a bath at 80–90°C. No explosion occurred when the bath temperature was limited to 40–45°C, or during subsequent distillation at 60–64°C/76 mbar. (Aziridine derivatives may have been formed by dehydrohalogenation.)

†**0854. Trichloroethylsilane**
[115-21-9]



$\text{C}_2\text{H}_5\text{Cl}_3\text{Si}$

See other ALKYLHALOSILANES

†**0855. Fluoroethane (Ethyl fluoride)**
[353-36-6]



$\text{C}_2\text{H}_5\text{F}$

See other HALOALKANES

0856. Ethyl fluorosulfate
[371-69-7]



$\text{C}_2\text{H}_5\text{FO}_3\text{S}$

Kumar, R. C. *et al.*, *Inorg. Chem.*, 1984, **23**, 3113–3114

The compound, prepared by by co-condensation of diethyl sulfite (3.5 mmol) and chlorine fluoride (10 mmol) at –196°C, followed by slow warming to –78, then –20°, is unstable. Trap-to-trap distillation must be effected with great care, as violent explosions occurred (even on this small scale) when cryogenic cooling was removed from the traps. Scaling up is not recommended.

See other SULFUR ESTERS

0857. 2-Hydroxyethylmercury(II) nitrate
[51821-32-0]



$\text{C}_2\text{H}_5\text{HgNO}_4$

Whitmore, 1921, 110

It decomposes with a slight explosion on heating.

See other MERCURY COMPOUNDS *See other* ORGANOMETALLIC NITRATES

0858. Iodoethane (Ethyl iodide)

[75-03-6]

EtI

C₂H₅I

HCS 1980, 561

Preparative hazard

See Iodine: Ethanol, etc.

Silver chlorite

See Silver chlorite: Iodoalkanes

See other HALOALKANES

0859. Ethylmagnesium iodide

[10467-10-4]

EtMgI

C₂H₅IMg

Ethoxyacetylene

Jordan, C. F., *Chem. Eng. News*, 1966, **44**(8), 40

A stirred mixture of ethoxyacetylene with ethylmagnesium iodide in ether exploded when the agitator was turned off. The corresponding bromide had been used without incident in earlier smaller-scale attempts.

See other AGITATION INCIDENTS, ALKYL METAL HALIDES, GRIGNARD REAGENTS

0860. 2-Iodoethanol

[624-76-0]

IC₂H₄OH

C₂H₅IO

Trimethylstibine

See 2-Hydroxyethyltrimethylstibonium iodide

See related HALOALKANES

0861. Potassium ethoxide

[917-58-8]

KOEt

C₂H₅KO

1. *See entry* METAL ALKOXIDES

2. Anon. private comm., 1984

It may ignite in moist air [1]. The dry powder is very dusty, and suspensions in dry air have a low minimum ignition energy, 17.1 mJ. After an induction period depending on the temperature and humidity of the atmosphere, a sample confined in a Dewar flask decomposed rapidly, with an exotherm of 200°C. Heats of solution in ethanol and in water are 0.63 and 0.65 kJ/g, respectively. [2].

See other INDUCTION PERIOD INCIDENTS

0862. Ethyllithium
[811-49-4]

EtLi

C₂H₅Li

Sorbe, 1968, 82
It ignites in air.
See other ALKYL METALS

†0863. Aziridine (Ethylenimine)
[151-56-4]

C₂H₅N



HCS 1980, 169

Preparative hazard

Energy of decomposition (in range 130–390°C) measured as 2.02 kJ/g.
See 2-Chloroethylammonium chloride: Alkali
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Acids

'Ethylenimine', Brochure 125-521-65, Midland (Mich.), Dow Chemical Co., 1965
It is very reactive chemically and subject to aqueous auto-catalysed exothermic polymerisation, which may be violent if uncontrolled by dilution, slow addition or cooling. It is normally stored over solid caustic alkali, to minimise polymerisation catalysed by presence of carbon dioxide.
See other CATALYTIC IMPURITY INCIDENTS, POLYMERISATION INCIDENTS

Aluminium chloride, Substituted anilines

Lehman, D. *et al.*, *Chem. Abs.*, 1983, **98**, 125559
In the preparation of a series of substituted phenylethylenediamines, it is essential to add the reagents to an aromatic solvent at 30–80°C in the order aniline, then aluminium chloride, then ethylenimine to prevent uncontrollable exothermic reaction.

Chlorinating agents

Graefe, A. F. *et al.*, *J. Amer. Chem. Soc.*, 1958, **80**, 3939
It gives the explosive 1-chloroaziridine on treatment with, e.g. sodium hypochlorite solution.
See 1-Chloroaziridine

Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Strained ring heterocycles


Silver

'Ethylenimine' Brochure 125-521-65, Midland (Mich.), Dow Chemical Co., 1965
Explosive silver derivatives may be formed in contact with silver or its alloys,
including silver solder, which is therefore unsuitable in handling equipment.

See other AZIRIDINES

0864. Vinylamine (Etheneamine)

[593-67-9]



$\text{C}_2\text{H}_5\text{N}$


Isoprene

Seher, A., *Ann.*, 1952, **575**, 153–161

Attempts at interaction led to explosions.

†**0865. Acetaldehyde oxime (Hydroximinoethane)**

[107-29-9]



$\text{C}_2\text{H}_5\text{NO}$

HCS 1980, 98

See other OXIMES

0866. N-Methylformamide

[123-39-7]



$\text{C}_2\text{H}_5\text{NO}$

HCS 1980, 670

See other APROTIC SOLVENTS

†**0867. Ethyl nitrite**

[109-95-5]



$\text{C}_2\text{H}_5\text{NO}_2$

Haz. Chem. Data, 1975, 161

It may decompose explosively around 90°C.

See other ALKYL NITRITES

0868. Methyl carbamate

[598-55-0]



$\text{C}_2\text{H}_5\text{NO}_2$

Phosphorus pentachloride

See N-Carbomethoxyiminophosphoryl chloride

0869. Nitroethane

[79-24-3]

EtNO₂C₂H₅NO₂

HCS 1980, 682

1. Kirk-Othmer, 1996, Vol. 17, 216
2. *Beilsteins Handbuch der organischen Chemie*, H(1), 101, Springer, Berlin, 1918
Although it is certainly less explosive, being detonable only if both hot and heavily confined [1], many reference works recommend precautions similar to those for nitromethane. The sodium salt explodes on heating [2]
See Nitromethane

Metal oxides

See NITROALKANES: Metal oxides

Other reactants

Yoshida, 1980, 271

MRH values calculated for 13 combinations with oxidants are given.

See other NITROALKANES**†0870. Ethyl nitrate**

[625-58-1]

EtONO₂C₂H₅NO₃

Hydrocarbons, Oxygen

See Oxygen: Hydrocarbons, Promoters

Lewis acids

See ALKYL NITRATES (references 2,3)

Other reactants

Yoshida, 1980, 191

MRH values calculated for 7 combinations are given.

See other ALKYL NITRATES**0871. 2-Nitroethanol**

[625-48-9]

O₂NC₂H₄OHC₂H₅NO₃Anon., *ABCM Quart. Safety Summ.*, 1956, **27**, 24

An explosion occurred towards the end of vacuum distillation of a relatively small quantity of 2-nitroethanol. This was attributed to the presence of peroxides, but the presence of traces of alkali seems a possible alternative cause.

See other C-NITRO COMPOUNDS

0872. Ethyl azide
[871-31-8]



1. Boyer, J. H. *et al.*, *Chem. Rev.*, 1954, **54**, 32
2. Koch, E., *Angew. Chem. (Nachr.)*, 1970, **18**, 26
3. Burns, M. E. *et al.*, *Chem. Eng. News*, 1985, **63**(50), 2
4. Nielsen, C. J. *et al.*, *Spectrochim. Acta A*, 1988, **44A**, 409–422

Though stable at room temperature, it may detonate on rapid heating [1]. A sample stored at -55°C exploded after a few minutes' exposure at ambient temperature, possibly owing to development of internal pressure in the stoppered vessel. It will also explode if dropped from 1 m in a small flask on to a stone floor [2]. An explosion occurred during laboratory preparation of ethyl azide when bromomethyl blue indicator was added to the reaction mixture to show the pH. This was attributed either to the acidic nature of the indicator, or to exposure of the azide vapour to the ground glass joint. Precautions recommended are similar to those when using diazomethane, and include small scale working, no ground joints, protection from intense light, and dilution of the product with solvent [3]. A safe synthesis is described, and IR, NMR and Raman spectral data were recorded in vapour, liquid and solid states [4].

See other GLASS INCIDENTS, ORGANIC AZIDES

0873. 2-Azidoethanol
[1517-05-1]



1. Appleby, I. C., *Chem. & Ind.*, 1986, 337
2. Appleby, I. C., private comm., 1987

During the preparation of 2-azidoethanol from a stirred mixture of 2-bromoethanol (14.6 mol) and sodium azide (15.4 mol) heated on a steam bath, a violent explosion occurred after 100 min. The preparation had been carried out previously without mishap. The need for care in handling azides of low MW is stressed [1]. Later detailed studies showed that the most probable cause of the explosion was the extraordinarily high mechanical and thermal sensitivity of the compound, with initiation by vibration from the agitator [2].

See other ORGANIC AZIDES

0874. Imidodicarbonic diamide (Biuret)
[108-19-0]



Chlorine

See Chlorine: Nitrogen compounds

0875. N-Methyl-N-nitrosourea
[684-93-5]

$C_2H_5N_3O_2$




1. Arndt, F., *Org. Synth.*, 1943, Coll. Vol. 2, 462
2. Anon., *Angew. Chem. (Nachr.)*, 1957, **5**, 198
3. Anon., *Jahresber.*, 1978, 70–71
4. See entry SELF-ACCELERATING REACTIONS

This must be stored under refrigeration to avoid sudden decomposition after storage at ambient temperature or slightly above (30°C) [1]. More stable precursors for generation of diazomethane are now available. Material stored at 20°C exploded after 6 months [2]. Explosive decomposition and ignition of solvent ether occurred when 50 g of the compound in a nickel spoon was being added to 40% potassium hydroxide solution to generate diazomethane [3]. DSC examination of several N-nitroso compounds showed that most underwent self accelerating decomposition, some with extraordinary velocity. The time to maximum rate for the urea falls from 1200 min at 40° to near zero at 80°C, when the maximum heat flow is 8.7 J/s/g [4]. See Diazomethane

See other NITROSO COMPOUNDS

0876. 1-Methyl-3-nitro-1-nitrosoguanidine
[70-25-7]

$C_2H_5N_5O_3$



1. Eisendrath, J. N., *Chem. Eng. News*, 1953, **31**, 3016
2. Aldrich advertisement, *J. Org. Chem.*, 1974, **39**(7), cover iv

Formerly used as a diazomethane precursor, this material will detonate under high impact, and a sample exploded when melted in a sealed capillary tube [1]. Although the crude product from the aqueous nitrosation is pyrophoric, recrystallised material is stable (though powerfully mutagenic) [2].

See other N-NITRO COMPOUNDS, NITROSO COMPOUNDS

0877. Ethylsodium
[676-54-0]

C_2H_5Na



Houben-Weyl, 1970, Vol. 13.1, 384

The powder ignites in air.

See other ALKYL METALS

0878. Sodium ethoxide
[141-52-6]

C_2H_5NaO



May ignite in moist air.

See entry METAL ALKOXIDES

0879. Sodium dimethylsulfinate

[58430-57-2]



4-Chlorotrifluoromethylbenzene

See 4-Chlorotrifluoromethylbenzene: Sodium dimethylsulfinate*See* Dimethyl sulfoxide: Sodium hydride**0880. Sodium 2-hydroxyethoxide**

[7388-28-5]



Polychlorobenzenes

Milnes, M. H., *Nature*, 1971, **232**, 395–396

It was found that the monosodium salt of ethylene glycol decomposed at about 230°C, during an investigation of the 1968 Coalite explosion during the preparation of 2,3,5-trichlorophenol by hydrolysis of 1,2,4,5-tetrachlorobenzene with sodium hydroxide in ethylene glycol at 180°C. Decomposition in the presence of polychlorobenzenes was exothermic and proceeded rapidly and uncontrollably, raising the temperature to 410°C, and liberating large volumes of white vapours, one of which was identified as ethylene oxide. The decomposition proceeded identically in presence of the tetrachlorobenzene or the 3 isomeric trichlorobenzenes (and seems also likely to occur in the absence of these or the chlorophenols produced by hydrolysis). The latter view has now been confirmed.

See Sodium hydroxide: Glycols**†0881. Ethane**

[74-84-0]

*FPA H37*, 1975; *HCS 1980*, 456 (cylinder)

Chlorine

See Chlorine: Hydrocarbons

Dioxygenyl tetrafluoroborate

See Dioxygenyl tetrafluoroborate: Organic materials**0882. Dimethylaluminium bromide**

[3017-85-4]



Nitromethane

See Nitromethane: Alkylmetal halides*See other* ALKYLALUMINIUM HALIDES

- 0883. Dimethylaluminium chloride**
 [1184-58-3] **C₂H₆AlCl**
 Me_2AlCl
- Air, or Water
 Gaines, D. F. *et al.*, *Inorg. Chem.*, 1974, **15**, 203–204
 It ignites in air and reacts violently with water.
See other ALKYLALUMINIUM HALIDES
- 0884. Chlorodimethylarsine**
 [557-89-1] **C₂H₆AsCl**
 ClAsMe_2
- Leleu, *Cahiers*, 1977, (88), 362
 It ignites in air, even at 0°C.
See other ALKYLNON-METAL HALIDES
- 0885. Dimethylfluoroarsine**
 [420-23-5] **C₂H₆AsF**
 Me_2AsF
- Leleu, *Cahiers*, 1977, (88), 363
 It ignites in air.
See other ALKYLNON-METAL HALIDES
- 0886. Dimethyliodoarsine**
 [676-75-5] **C₂H₆AsI**
 Me_2AsI
- Millar, I. T. *et al.*, *Inorg. Synth.*, 1960, **6**, 117
 It ignites when heated in air.
See other ALKYLNON-METAL HALIDES
- 0887. Bromodimethylborane**
 [5158-50-9] **C₂H₆BBr**
 Me_2BBr
- Katz, H. E., *J. Org. Chem.*, 1985, 50, **50** 28
 It is pyrophoric in air.
See other ALKYLHALOBORANES, PYROPHORIC MATERIALS
- 0888. Azidodimethylborane**
 [2306-67-0] **C₂H₆BN₃**
 N_3BMe_2
- Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 27

A sample exploded on contact with a warm sampling capillary.
See other NON-METAL AZIDES

0889. Barium methyl hydroperoxide

[]



See Methyl hydroperoxide
See related ALKYL HYDROPEROXIDES

0890. Dimethylberyllium

[506-63-8]



Air, or Carbon dioxide, or Water

Coates, 1967, Vol. 1, 106

It ignites in moist air or in carbon dioxide, and reacts explosively with water.

See other ALKYL METALS

0891. Dimethylberyllium-1,2-dimethoxyethane

[]



Houben-Weyl, 1973, Vol. 13.2a, 38

It ignites in air.

See related ALKYL METALS

0892. Chloroethylbismuthine

[65313-33-9]



Leleu, *Cahiers*, 1977, (88), 364

It ignites in air.

See related ALKYL METAL HALIDES, ALKYL METAL HYDRIDES

0893. Dimethylbismuth chloride


[]



Sidgwick, 1950, 781

It ignites when warm in air.
See other ALKYL METAL HALIDES

0894. Dimethylcadmium
[506-82-1]




1. Egerton, A. *et al.*, *Proc. R. Soc.*, 1954, **A225**, 429
2. Davies, A. G., *Chem. & Ind.*, 1958, 1177
3. Sidgwick, 1950, 268

On exposure to air, dimethylcadmium peroxide is formed as a crust [1] which explodes on friction [2]. Ignition of dimethylcadmium may occur if a large area:volume ratio is involved, as when it is dripped onto filter paper [3]. It is a mildly endothermic compound (ΔH_f° (l) +67.8 kJ/mol, 0.47 kJ/g).

See other ALKYL METALS, ENDOTHERMIC COMPOUNDS

0895. N-Chlorodimethylamine
[1585-74-6]




Antimony chlorides

Weiss, W. *et al.*, *Z. Anorg. Chem.*, 1977, **433**, 207–210

The crystalline addition compounds formed with antimony trichloride and antimony pentachloride in dichloromethane at -78°C tend to explode at ambient temperature, or when shocked or heated.

See other N-HALOGEN COMPOUNDS

0896. 2-Chloroethylamine
[689-98-5]





It may polymerise explosively.

See 2-Chloroethylammonium chloride: Alkali *also* 2-Chloro-1,4-phenylenediamine

See other POLYMERISATION INCIDENTS


0897. Methyl iminioformate chloride
[15755-09-6]



Preparative hazard

See Hydrogen chloride: Alcohols, Hydrogen cyanide

0898. 2-Aza-1,3-dioxolanium perchlorate (Ethylenedioxyammonium perchlorate)
[] **C₂H₆ClNO₆**



MCA Case History No. 1622

A batch of this sensitive compound (low oxygen balance) exploded violently, probably during recrystallisation.

See other PERCHLORATE SALTS OF NITROGENOUS BASES, N-O COMPOUNDS

0899. Chlorodimethylphosphine
[811-62-1] **C₂H₆ClP**
ClPMe₂

1. Staendeke, H. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1973, **12**, 877

2. Parshall, G. W., *Inorg. Synth.*, 1974, **15**, 192–193

It ignites in contact with air [1], and handling procedures are detailed [2].

See other ALKYLHALOPHOSPHINES

0900. Dimethylantimony chloride
[18380-68-2] **C₂H₆ClSb**
Me₂SbCl

Sidgwick, 1950, 777

It ignites at 40°C in air.

See other ALKYLMETAL HALIDES

0901. Dimethyl N,N-dichlorophosphoramidate
[29727-86-4] **C₂H₆Cl₂NO₃P**
(MeO)₂P(O)NCl₂

Preparative hazard

See Chlorine: Dimethyl phosphoramidate

See other N-HALOGEN COMPOUNDS, PHOSPHORUS ESTERS

†0902. Dichlorodimethylsilane
[75-78-5] **C₂H₆Cl₂Si**
Cl₂SiMe₂

HCS 1980, 428

Water

Leleu, *Cahiers*, 1977, (88), 370

Violent reaction with water.


See other ALKYLHALOSILANES

†0903. Dichloroethylsilane
[1789-58-8]




See other ALKYLHALOSILANES

0904. (Dimethyl ether)oxodiperoxochromium(VI)
[]





Schwarz, R. *et al.*, *Ber.*, 1936, **69**, 575
The blue solid explodes powerfully at -30°C .
See related AMMINECHROMIUM PEROXOCOMPLEXES

0905. Copper(II) glycinate nitrate (Aquadiglycinatonitratocopper)
[94791-14-7]



Davies, H. O. *et al.*, *J. Chem. Soc. Chem. Comm.*, 1992, (3), 226
The mixed salt explodes on heating to 167°C , first losing water.
See related METAL NITRATES

†0906. 2,2,2,4,4,4-Hexafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine
[3880-04-4]



tert-Butyllithium

See *tert*-Butyllithium: 2,2,2,4,4,4-Hexafluoro-1,3,2,4-diazadiphosphetidine

0907. Dimethylmercury
[593-74-8]

C₂H₆Hg

Me₂Hg

It is mildly endothermic (ΔH_f° (l) +75.3 kJ/mol, 0.33 kJ/g).

Diboron tetrachloride

See Diboron tetrachloride: Dimethylmercury

See other ALKYL METALS, ENDOTHERMIC COMPOUNDS, MERCURY COMPOUNDS

0908. Dimethylmagnesium
[2999-74-8]

C₂H₆Mg

Me₂Mg

Air, or Water

Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 5049

Contact with moist air usually caused ignition of the dry powder, and water always ignited the solid or its ethereal solution.

See other ALKYL METALS, DIALKYL MAGNESIUMS

0909. Dimethylmanganese
[33212-68-9]

C₂H₆Mn

Me₂Mn

1. Bailar, 1973, Vol. 3, 851; Vol. 4, 792

2. *491M* 1975, 159

It is a readily explosive powder, (probably polymeric) [1], which also ignites in air [2].

See other ALKYL METALS

0910. Azomethane (Dimethyldiazene)
[503-28-6]

C₂H₆N₂

H₃CN=NCH₃

Allen, A. O. *et al.*, *J. Amer. Chem. Soc.*, 1935, **57**, 310–317

The slow thermal decomposition of gaseous azomethane becomes explosive above 341°C/250 mbar and 386°C/24 mbar.

See other AZO COMPOUNDS

0911. Disodium *N,N'*-dimethoxysulfonyldiamide
[]

C₂H₆N₂Na₂O₄S

O₂S[N(Na)OMe]₂

Goehring, 1957, 87

The dry solid readily explodes, as do many *N*-metal hydroxylamides.
See other N-METAL DERIVATIVES, N-O COMPOUNDS

0912. Acetohydrazide
[1068-57-1]




Energy of decomposition (in range 460–510°C) measured as 1.02 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

0913. Dimethyl hyponitrite (Dimethoxydiazene)
[29128-41-4]



Mendenhall, G. D. *et al.*, *J. Org. Chem.*, 1975, **40**, 1646
Samples of the ester exploded violently during low-temperature distillation, and on freezing by liquid nitrogen. The undiluted liquid is exceptionally unpredictable, but addition of mineral oil before vaporisation improved safety aspects.
See other DIALKYL HYPONITRITES

0914. Dimethyltin dinitrate
[40237-34-1]



Sorbe, 1968, 159
It decomposes explosively on heating.
See other ORGANOMETALLIC NITRATES

0915. *N*-Azidodimethylamine
[2156-66-3]



Bock, H. *et al.*, *Angew. Chem.*, 1962, **74**, 327
It is rather explosive.
See related ORGANIC AZIDES

0916. *N,N'*-Dinitro-1,2-diaminoethane
[505-71-5]



1. Urbanski, 1967, Vol. 3, 20
2. Stull, 1977, 20

This powerful but relatively insensitive explosive decomposes violently at 202°C, and gives lead and silver salts which are highly impact sensitive [1]. Though not endothermic ($\Delta H_f^\circ -103.3$ kJ/mol), as a bis-nitramine it has a rather high heat of decomposition (3.91 kJ/g) which it is calculated would attain an adiabatic decomposition temperature over 2250°C, with a 60-fold pressure increase in a closed system [2].


See other N-NITRO COMPOUNDS

Amminemetals

Complexes with several ammine derivatives of metals are explosive.

See entry NITRAMINE-METAL COMPLEXES

0917. S, S-Dimethylpentasulfur hexanitride [71901-54-7]



Sheldrick, W. S. *et al.*, *Inorg., Chem.*, 1980, **19**, 339

Powerfully explosive like the non-methylated compound, breaking the quartz combustion tubes during microanalysis.

See other N-S COMPOUNDS

0918. Diazidodimethylsilane [4774-73-6]



Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 26-27

A 3 year old sample exploded violently on removing the ground stopper.

See other FRICTIONAL INITIATION INCIDENTS, NON-METAL AZIDES

†0919. Dimethyl ether (Oxybismethane) [115-10-6]



FPA H113, 1983, HCS 1980, 429

†0920. Ethanol (Ethyl alcohol)

[64-17-5]



NSC 391, 1968; *FPA H2*, 1972; *HCS 1980*, 639; *RSC Lab. Hazard Data Sheet No. 50*, 1986

Acetic anhydride, Sodium hydrogensulfate

See Acetic anhydride: Ethanol, etc.

Acetyl bromide

See Acetyl bromide: Hydroxylic compounds

Ammonia, Silver nitrate

See Silver nitrate: Ammonia, Ethanol

Disulfuric acid, Nitric acid

See Nitric acid: Alcohols, Disulfuric acid

Dichloromethane, Sulfuric acid, (A nitrate or nitrite?)

See Sulfuric acid, Dichloromethane, etc.

Disulfuryl difluoride

See Disulfuryl difluoride: Ethanol

Magnesium perchlorate

See Magnesium perchlorate: Ethanol

Nitric acid, Silver

See Silver: Ethanol, Nitric acid

Other reactants

Yoshida, 1980, 43

MRH values calculated for 18 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Barium perchlorate: Alcohols

Bromine pentafluoride: Hydrogen-containing materials

Calcium hypochlorite: Hydroxy compounds

MRH 2.30/93

Chloryl perchlorate: Organic matter

Chromium trioxide: Alcohols

MRH 2.55/90

Chromyl chloride: Organic solvents

Dioxygen difluoride: Various materials

'Fluorine nitrate': Organic materials

Hydrogen peroxide: Alcohols

MRH 6.19/21

Iodine heptafluoride: Organic solvents

Nitric acid: Alcohols

MRH 5.56/76

Nitrosyl perchlorate: Organic materials
 Perchloric acid: Alcohols MRH 6.28/79
 Permanganic acid: Organic materials
 Peroxodisulfuric acid: Organic liquids
 Potassium dioxide: Ethanol
 Potassium perchlorate: Ethanol
 Potassium permanganate: Ethanol, etc. MRH 2.68/91
 Ruthenium(VIII) oxide: Organic materials
 Silver perchlorate: Aromatic compounds MRH 3.30/87
 Sodium peroxide: Hydroxy compounds MRH 2.43/91
 Uranium hexafluoride: Aromatic hydrocarbons, etc.
 Uranyl perchlorate: Ethanol
See N-HALOMIDES: Alcohols

Phosphorus(III) oxide
See Tetraphosphorus hexaoxide: Organic liquids

Platinum
See Platinum: Ethanol

Potassium
See Potassium: Slow oxidation

Potassium *tert*-butoxide
See Potassium tert-butoxide: Acids, etc.

Silver nitrate
See Silver nitrate: Ethanol

Silver oxide
See Silver(I) oxide: Ammonia, etc.

Sodium
See Sodium: Ethanol

Tetrachlorosilane
See Tetrachlorosilane: Ethanol

†0921. Dimethyl sulfoxide (Sulfinylbismethane) C₂H₆OS
 [69-68-5] Me₂S(:O)

HCS 1980, 435; RSC Lab. Hazard Data Sheet No. 11, 1983

1. Brogli, F. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 681–682, Basle, SSCI, 1980
2. Cardillo, P. *et al.*, *Chim. e Ind. (Milan)*, 1982, **44**, 231–234

3. Santosusso, T. M. *et al.*, *Tetrahedron Lett.*, 1974, 4255–4258

4. *See entry* THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
Two instances of used DMSO decomposing exothermally while being kept at 150°C prior to recovery by vacuum distillation were investigated. Traces of alkyl bromides lead to a delayed, vigorous and strongly exothermic reaction ($Q = 0.85$ kJ/g) at 180°C. Addition of zinc oxide as a stabiliser extends the induction period and markedly reduces the exothermicity [1]. ARC examination shows that exothermic decomposition sets in by a radical mechanism at 190°C, just above the b.p., 189°C. The proposed retardants, sodium carbonate and zinc oxide, do not affect the decomposition temperature, and a maximum decomposition pressure of 60 bar was attained (at up to 4 bar/min) at the low sample loading of 18 w/v% in the bomb [2]. The thermolytic degradation of the sulfoxide to give acidic products which catalyse further decomposition had been discussed previously [3]. T_{ait24} was determined as 213°C by adiabatic Dewar tests, with an apparent energy of activation of 243 kJ/mol. At elevated temperatures (200°C) DSC shows decomposition to be both faster and more energetic when chloroform or sodium hydroxide is present.

See other INDUCTION PERIOD INCIDENTS, SELF-ACCELERATING REACTIONS

Acid anhydrides

See Trifluoroacetic anhydride, below

Acids, or Bases

Hall, J., *Loss Prev. Bull.*, 1993, (114), 2

These may catalyse exothermic decomposition. A runaway of a mixture with 4-nitrobenzenesulfonic acid, possibly from as low as 60°C (though a steam valve may have been leaking) is reported by Hall.

See reference 2, next below

Acyl halides, or Non-metal halides

MRH Thionyl chloride 1.63/60

1. Buckley, A., *J. Chem. Educ.*, 1965, **42**, 674

2. Allan, G. G. *et al.*, *Chem. & Ind.*, 1967, 1706

3. Mancuso, A. J. *et al.*, *J. Org. Chem.*, 1979, **44**, 4148–4150

In absence of diluent or other effective control of reaction rate, the sulfoxide reacts violently or explosively with the following: acetyl chloride, benzenesulfonyl chloride, cyanuric chloride, phosphorus trichloride, phosphoryl chloride, tetrachlorosilane, sulfur dichloride, disulfur dichloride, sulfuryl chloride or thionyl chloride [1]. These violent reactions are explained in terms of exothermic polymerisation of formaldehyde produced under a variety of conditions by interaction of the sulfoxide with reactive halides, acidic or basic reagents [2]. Oxalyl chloride reacts explosively with DMSO at ambient temperature, but controllably in dichloromethane at –60°C [3].

See Carbonyl diisothiocyanate, and Dinitrogen tetraoxide, and Hexachlorocyclotriphosphazine, and Sodium hydride, all below

See Perchloric acid: Sulfoxides

Allyl trifluoromethanesulfonates

See ALLYL TRIFLUOROMETHANESULFONATES: Alone, or Aprotic solvents

Boron compounds

Shriver, 1969, 209

DMSO forms an explosive mixture with nonahydroxonaborate(2-) ion and with diborane. It is probable that other boron hydrides and hydroborates behave similarly.

Bromides

1. Hall, J., *Loss Prev. Bull.*, 1993, (114), 2

2. Aida, T., *et al.*, *Bull. Chem. Soc. Jap.*, 1976, **49**, 117

A review of runaway decomposition reactions of DMSO [1], several involving bromides, some alkaline and under vacuum, from temperatures possibly as low as 130°C. A bromide/bromine catalysed decomposition reaction is known [2].

4(4'-Bromobenzoyl)acetanilide

491M, 1975, 74

An explosion occurred after a solution of the anilide in DMSO had been held at 100°C for 30 min.

Bromomethane

See Trimethylsulfoxonium bromide

Carbonyl diisothiocyanate

Bunnenberg, R. *et al.*, *Chem. Ber.*, 1981, **114**, 2075–2086

Interaction of the solvent with the strong electrophile is explosive.

See Acyl halides, above

Copper, Trichloroacetic acid

1. Giessemann, B. W., *Chem. Eng. News*, 1981, **59**(28), 4

2. Dunne, T. G. *et al.*, *Chem. Eng. News*, 1981, **59**(50), 4

The acid was added to copper wool and rinsed down with DMSO. Within 20 s, the contents of the flask were ejected and the neck was distorted by intense heat [1]. Adding the copper wool to a solution of the acid in the solvent gave an exothermic but controlled reaction [2]. (Formation of a reactive carbene species by dehydrohalogenation seems a remote possibility.)

See Acyl halides, above

Dichloromethane, Perchloric acid

See Perchloric acid: Dichloromethane, Dimethyl sulfoxide

Dinitrogen tetroxide

Buckley, A., *J. Chem. Educ.*, 1965, **42**, 674

Interaction may be violent or explosive.

See Acyl halides

See Nitric acid: Dimethyl sulfoxide

Hexachlorocyclotriphosphazine

Pierce, T. *et al.*, *Lab. Haz. Bull.*, 1984, (4), item 215

C₂H₆OS

Addition of about 100 mg of hexachlorocyclotriphosphazine (also containing some phenoxyated products) to 0.5 ml of DMSO in an NMR tube led to an exothermic reaction which ejected the tube contents. General precautions are suggested.

See Acyl halides, above

Iodine pentafluoride

Lawless, E. M., *Chem. Eng. News*, 1969, **47**(13), 8

Interaction is explosive, after a delay, in either tetrahydrothiophene-1,1-dioxide (sulfolane) or trichlorofluoromethane as solvent, on 0.15 g mol scale, though not on one tenth this scale.

Magnesium perchlorate

MRH 5.52/78

See Magnesium perchlorate: Dimethyl sulfoxide

Metal alkoxides

MCA Case History No. 1718

Addition of potassium *tert*-butoxide or of sodium isopropoxide to the solvent led to ignition of the latter. This was attributed to presence of free metal in the alkoxides, but a more likely explanation seems to be that of direct interaction between the powerful bases and the sulfoxide.

See Acids, or Bases, above (reference 2)

Potassium *tert*-butoxide: Acids, etc.

Metal oxosalts

1. Martin, 1971, 435

2. Dehn, H., Brit. Pat. 1 129 777, 1968

3. Sandstrom, M. *et al.*, *Acta Chem. Scand.*, 1978, **A32**, 610

Mixtures of the sulfoxide with metal salts of oxoacids are powerful explosives. Examples are aluminium perchlorate, sodium perchlorate and iron(III) nitrate [1]. The water in hydrated oxosalts (aluminium perchlorate, iron(III) perchlorate, iron(III) nitrate) may be partially or totally replaced by dimethyl (or other) sulfoxide to give solvated salts useful as explosives [2]. Metal nitrates and perchlorates solvated with DMSO are generally powerfully explosive, and under certain conditions a violent reaction is easily triggered [3]. Several other explosions involving perchlorates and the sulfoxide have been reported.

See Chromium(III) perchlorate . 6 dimethyl sulfoxide

Magnesium perchlorate: Dimethyl sulfoxide

Mercury(II) perchlorate . 6(or 4)dimethyl sulfoxide

Silver perchlorate: Dimethyl sulfoxide

See Perchloric acid: Sulfoxides

Nitric acid

See Nitric acid: Dimethyl sulfoxide

Non-metal halides

See Acyl halides, etc., above

Other reactants

Yoshida, 1980, 171

MRH values calculated for 6 combinations are given.

Perchloric acid

MRH 6.19/77

See Perchloric acid: Sulfoxides

Periodic acid

See Periodic acid: Dimethyl sulfoxide

Phosphorus(III) oxide

See Tetraphosphorus hexaoxide: Organic liquids

Potassium

Houben-Weyl 1970, Vol. 13.1, 295

Interaction of potassium 'sand' and dimethyl sulfoxide is violent in the absence of a diluent, and leads to partial decomposition of the potassium dimethylsulfinat. THF is a suitable diluent.

See Metal alkoxides, above

Potassium permanganate

MRH 3.01/14

See Potassium permanganate: Dimethyl sulfoxide

Silver difluoride

Lawless, E. M., *Chem. Eng. News*, 1969, **47**(13), 8

Interaction is violent.

Sodium

Hall, J., *Loss Prev. Bull.*, 1993, (114), 2

A preparation of dimethyl sodium from sodium and DMSO, on 14 litre scale, overheated, then exploded causing two fatalities. It has not proved possible to duplicate the runaway.

Sodium hydride

1. French, F. A., *Chem. Eng. News*, 1966, **44**(15), 48

2. Olson, G. A., *Chem. Eng. News*, 1966, **44**(24), 7

3. Russell, G. A. *et al.*, *J. Org. Chem.*, 1966, **31**, 248

4. Batchelor, J. F., private comm., 1976

5. Anon., *Loss Prev. Bull.*, 1980, (030), 161

6. Allan, G. G. *et al.*, *Chem. & Ind.*, 1967, 1706

7. Itoh, M. *et al.*, *Lab. Haz. Bull.* 1985(5), item 304

Two violent pressure-explosions occurred during preparations of dimethylsulfinyl anion on 3–4 g mol scale by reaction of sodium hydride with excess solvent. In each case, the explosion occurred soon after separation of a solid. The first reaction involved addition of 4.5 g mol of hydride to 18.4 g mol of sulfoxide, heated to 70°C [1], and the second 3.27 and 19.5 g mol respectively, heated to 50°C [2]. A smaller scale reaction at the original lower hydride concentration [3], did not

explode, but methylation was incomplete. Explosions and fire occurred when the reaction mixture was overheated (above 70°C) [4]. Reaction of 1 g mol of hydride with 0.5 l of sulfoxide at 80°C led to an exotherm to 90°C with explosive decomposition [5]. These and similar incidents are explicable in terms of exothermic polymerisation of formaldehyde produced from sulfoxide by reaction with the hydride base [6]. The heat of reaction was calculated and determined experimentally. Thermal decomposition of the solution of hydride is not very violent, but begins at low temperatures, with gas evolution [7].

See other GAS EVOLUTION INCIDENTS

See Sodium dimethylsulfinate

Sulfur trioxide

MRH 2.51/59

See Sulfur trioxide: Dimethyl sulfoxide

Trifluoroacetic anhydride

Sharma, A. K. *et al.*, *Tetrahedron Lett.*, 1974, 1503–1506

Interaction of the sulfoxide with some acid anhydrides or halides may be explosive. The highly exothermic reaction with trifluoroacetic anhydride was adequately controlled in dichloromethane solution at below –40°C.

See Acyl halides, above

See other APROTIC SOLVENTS

0922. Poly(dimethylsiloxane)

[9016-00-6]

(C₂H₆OSi)_n

(-SiMe₂O-)_n

See SILICONE OIL

0923. Dimethyl peroxide

[690-02-8]

C₂H₆O₂

MeOOME

1. Rieche, A. *et al.*, *Ber.*, 1928, **61**, 951

2. Baker, G. *et al.*, *Chem. & Ind.*, 1964, 1988

Extremely explosive, heat- and shock-sensitive as liquid or vapour [1]. During determination of the impact-sensitivity of the confined material, rough handling of the container caused ignition. The material should only be handled in small quantity and with great care [2].

See other DIALKYL PEROXIDES

0924. Ethylene glycol (1,2-Ethanediol)

[107-21-1]

C₂H₆O₂

HOC₂H₄OH

FPA H47, 1976; *HCS 1980*, 483; *RSC Lab. Hazards Data Sheet No. 69*, 1988

Dimethyl terephthalate, Titanium butoxide

See Dimethyl terephthalate: Ethylene glycol, Titanium butoxide

Other reactants

Yoshida, 1980, 48

MRH values calculated for 15 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

1. Oouchi, H. *et al.*, *Chem. Abs.*, 1982, **97**, 8605

2. Inoue, Y. *et al.*, *Chem. Abs.*, 1987, **107**, 120353

In a study of hypergolic ignition of ethylene glycol by oxidants, chromium trioxide, potassium permanganate and sodium peroxide caused ignition on contact at ambient temperature, and ammonium dichromate, silver chlorate, sodium chlorite and uranyl nitrate at 100°C [1]. Results from an exothermicity test in which 1 g portions each of a glycol and an oxidising agent were mixed at 100°C in a Dewar flask were compared with those from DTA experiments [2].

See Chromium trioxide: Ethylene glycol

MRH 2.47/84

Perchloric acid: Glycols and their ethers

MRH 5.69/70

Potassium dichromate: Ethylene glycol

MRH 1.46/89

See Sodium hypochlorite: Ethanediol

Phosphorus pentasulfide

See Tetrathosphorus decasulfide: Alcohols

Silvered copper wire

1. Downs, W. R., *TN D-4327*, Washington, NASA Tech. Note, 1968

2. Stevens, H. D., *Chem. Abs.*, 1975, **83**, 134485

Contact of aqueous ethylene glycol solutions with d.c.-energised silvered copper wires causes ignition of the latter. Bare copper or nickel- or tin-plated wires were inert and silver-plated wire can be made so by adding benzotriazole as a metal deactivator to the coolant solution [1]. This problem of electrical connector fires in aircraft has been studied in detail to identify the significant factors [2].

Sodium hydroxide

See Sodium hydroxide: Glycols, and Sodium 2-hydroxyethoxide

Sodium hypochlorite

See Sodium hypochlorite: Ethanediol

0925. Ethyl hydroperoxide

[3031-74-1]

C₂H₆O₂

EtOOH

Baeyer, A. *et al.*, *Ber.*, 1901, **34**, 738

It explodes violently on superheating; the barium salt is heat- and impact-sensitive.

Hydriodic acid

Sidgwick, 1950, 873

The concentrated acid (a reducer) is oxidised explosively.

See other REDOX REACTIONS

Silver

Sidgwick, 1950, 873

Finely divided ('molecular') silver decomposes the hydroperoxide, sometimes explosively.

See other ALKYL HYDROPEROXIDES

0926. Hydroxymethyl methyl peroxide

[]




Rieche, A. *et al.*, *Ber.*, 1929, **62**, 2458

Violently explosive, impact-sensitive when heated.

See other 1-OXYPEROXY COMPOUNDS

0927. Dimethyl sulfite

[616-42-2]



Phosphorus (III) oxide

See Tetraphosphorus hexaoxide: Organic liquids

See other SULFUR ESTERS

0928. Bis(hydroxymethyl) peroxide

[17088-73-2]




Wieland, H., *et al.*, *Ber.*, 1930, **63**, 66

Highly explosive and very friction-sensitive. Higher homologues are more stable.

See other 1-OXYPEROXY COMPOUNDS

†0929. Dimethyl sulfate

[77-78-1]



(*MCA SD-19*, 1966); *HCS 1980*, 434

Ammonia

1. Lindlar, H., *Angew. Chem.*, 1963, **75**, 297

2. Claesson, P. *et al.*, *Ber.*, 1880, **13**, 1700

A violent reaction occurred which shattered the flask when litre quantities of dimethyl sulfate and conc. aqueous ammonia were accidentally mixed. Use dilute ammonia in small quantities to destroy dimethyl sulfate [1]. Similar incidents were noted previously with ammonia and other bases [2].

See Tertiary bases, below

See other AMINATION INCIDENTS

Barium chlorite

See Barium chlorite: Dimethyl sulfate

Other reactants

Yoshida, 1980, 395

MRH values calculated for 13 combinations with oxidants are given.

Tertiary bases

Sorbe, 1968, 123

In absence of diluent, quaternation of some tertiary organic bases may proceed explosively.

See Ammonia, above

Unnamed material


MCA Case History No. 1786

The product of methylating an unnamed material at 110°C was allowed to remain in the reactor of a pilot plant. and after 80 min the reactor exploded. This was ascribed to exothermic decomposition of the mixture above 100°C, and subsequent acceleration and boiling decomposition at 150°C.

See other SULFUR ESTERS

0930. Dimethyl selenate

[6918-51-0]



Sidgwick, 1950, 977

Dimethyl selenate, like its ethyl and propyl homologues, can be distilled under reduced pressure, but explodes at around 150°C under ambient pressure, though less violently than the lower alkyl nitrates.

See related SULFUR ESTERS

0931. Dimethanesulfonyl peroxide

[1001-62-3]



Haszeldine, R. N. *et al.*, *J. Chem. Soc.*, 1964, 4903

A sample heated in a sealed tube exploded violently at 70°C. Unconfined, the peroxide decomposed (sometimes explosively) immediately after melting at 79°C.

See other DIACYL PEROXIDES

†0932. Dimethyl sulfide (Thiobismethane)
[75-18-3] C₂H₆S
Me₂S

Dibenzoyl peroxide
See Dibenzoyl peroxide: Dimethyl sulfide

1,4-Dioxane, Nitric acid
See Nitric acid: Dimethyl sulfide, 1,4-Dioxane

Other reactants
Yoshida, 1980, 389
MRH values calculated for 14 combinations with oxidants are given.

Oxygen
See Oxygen (Gas): Dimethyl sulfide

Xenon difluoride
See Xenon difluoride: Dimethyl sulfide

†0933. Ethanethiol
[75-08-1] C₂H₆S
EtSH

HCS 1980, 497

Other reactants
Yoshida, 1980, 44
MRH values calculated for 14 combinations with oxidants are given.
See other ALKANETHIOLS

†0934. Dimethyl disulfide
[624-92-0] C₂H₆S₂
MeSSMe

0935. Dimethylzinc
[544-97-8] C₂H₆Zn
Me₂Zn

Air, or Oxygen
1. Egerton, A. *et al.*, *Proc. R. Soc.*, 1954, **A225**, 429
2. Frankland, E., *Phil. Trans. R. Soc.*, 1852, 417
It ignites in air (owing to peroxide formation) [1], and explodes in oxygen [2]. It is slightly endothermic (ΔH_f° (l) +25.1 kJ/mol, 0.26 kJ/g).

2,2-Dichloropropane
Houben-Weyl, 1973, Vol. 13.2a, 767
Uncontrolled reaction is explosive.
See DIALKYLZINCS: Alkyl chlorides

Ozone
See Ozone: Alkylmetals

Water
Leleu, *Cahiers*, 1977, (88), 371
Reaction is explosively violent.
See other ALKYLALUMINIUM ALKOXIDES AND HYDRIDES, ENDOTHERMIC COMPOUNDS

0936. Dimethylaluminium hydride
[865-37-2]



$\text{C}_2\text{H}_7\text{Al}$

Houben-Weyl, 1970, Vol. 13.4, 58
Slight contact with air or moisture causes ignition.
See other ALKYLALUMINIUM ALKOXIDES AND HYDRIDES


0937. Dimethylarsine
[593-57-7]



$\text{C}_2\text{H}_7\text{As}$

1. von Schwartz, 1918, 322
2. Sidgwick, 1950, 762
It inflames in air [1], even at 0°C [2].
See other ALKYLNON-METAL HYDRIDES

0938. 1,1-Dimethyldiazenium perchlorate
[[53534-20-6] (ion)]



$\text{C}_2\text{H}_7\text{ClN}_2\text{O}_4$

McBride, W. E. *et al.*, *J. Amer. Chem. Soc.*, 1957, **79**, 576 footnote 16
The salt is impact sensitive.
See other PERCHLORATE SALTS OF NITROGENOUS BASES

0939. 1-Methyl-3-nitroguanidinium perchlorate
[]



$\text{C}_2\text{H}_7\text{ClN}_4\text{O}_6$

McKay, A. F. *et al.*, *J. Amer. Chem. Soc.*, 1947, **69**, 3029
The salt is sensitive to impact, exploding violently.
See other N-NITRO COMPOUNDS, PERCHLORATE SALTS OF NITROGENOUS BASES

0940. 2-Chloroethylammonium chloride (2-Chloroethylamine hydrochloride)
[870-24-6] **C₂H₇Cl₂N**



Alkali

1. Wystrach, V. P. *et al.*, *J. Amer. Chem. Soc.*, 1955, **77**, 5915
2. Wystrach, V. P. *et al.*, *Chem. Eng. News*, 1956, **34**, 1274

The procedure described previously [1] for the preparation of ethylenimine is erroneous. 2-Chloroethylammonium chloride must be added with stirring as a 33% solution in water to strong sodium hydroxide solution. Addition of the solid salt to the alkali caused separation in bulk of 2-chloroethylamine, which polymerised explosively. Adequate dilution and stirring, and a temperature below 50°C are all essential [2].

See other POLYMERISATION INCIDENTS

†0941. Dimethylamine (N-Methylmethanamine)
[124-40-3]



(MCA SD-57, 1955); HCS 1980, 421 (40% solution)

Acrylaldehyde

See Acrylaldehyde: Acids, etc.

4-Chloroacetophenone

See 4-Chloroacetophenone: Dimethylamine

Fluorine

MRH 10.6/77

See Fluorine: Nitrogenous bases

Maleic anhydride

See Maleic anhydride: Bases, etc.

Other reactants

Yoshida, 1980, 169

MRH values calculated for 14 combinations with oxidants are given.

See other ORGANIC BASES

†0942. Ethylamine (Ethanamine)
[75-04-7]



FPA H67, 1978; HCS 1980, 467 (cylinder)

Cellulose nitrate

See CELLULOSE NITRATE: Amines

See other ORGANIC BASES

0943. 2-Hydroxyethylamine (Ethanolamine)

[141-43-5]

C_2H_7NO



HCS 1980, 457

Carbon dioxide, Pyrophoric deposits

1. Bond, J., *Hazards from Pressure*, IChE Symp. Ser. No. 102, 37–44, Oxford, Pergamon, 1987

2. Bond, J., *Loss Prev. Bull.*, 1995, **121**, 18

Ethanolamine has a large affinity for carbon dioxide, and is used to remove it from process gas streams. A storage tank used to hold the amine had a steel steam coil to warm the contents, and the coil developed a pyrophoric deposit, probably from contact with sulfur-containing impurities. When part of the coil was not covered by the amine, the surface deposit caused a fire which ignited some of the ethanolamine in the closed tank until the air was exhausted and the fire went out. The combined effect of cooling of the hot gases in the tank and of the rapid absorption of the carbon dioxide produced in the fire caused a significant pressure reduction and the tank collapsed [1]. It was also suggested that the pyrophoric material might have been tris(hydroxyethylamine)iron(0), a rather improbable species, or tris(aminoethoxy)iron(III) trihydrate which is perhaps more probable, though its postulated decomposition to pyrophoric iron is less so [2].

See PYROPHORIC IRON–SULFUR COMPOUNDS

Cellulose nitrate

See CELLULOSE NITRATE: Amines

N,N-Dimethyl-*N,N*-dinitrosoterephthalamide

See N,N'-Dimethyl-*N,N'*-dinitrosoterephthalamide: 2-Hydroxyethylamine

Other reactants

Yoshida, 1980, 370


MRH values calculated for 15 combinations are given.

See other ORGANIC BASES

0944. *O*-(2-Hydroxyethyl)hydroxylamine

[3279-95-6]

$C_2H_7NO_2$



Sulfuric acid

Campbell, H. F., *Chem. Eng. News*, 1975, **53**(49), 5

After 30 min at 120°C under vacuum, an equimolar mixture with the conc. acid exploded violently. Salts of unsubstituted hydroxylamine are thermally unstable.

See HYDROXYLAMINIUM SALTS *See other* N–O COMPOUNDS

- 0945. 1,3-Dimethyltriazene ('Diazoaminomethane')**
 [3585-32-8] **C₂H₇N₃**

$$\text{MeN}=\text{NNHMe}$$
 Dimroth, O., *Ber.*, 1906, **39**, 3910
 A drop explodes sharply in contact with flame.
See other TRIAZENES
- 0946. 1,2-Dimethylnitrosohydrazine**
 [101672-10-0] **C₂H₇N₃O**

$$\text{MeN}(\text{N}:\text{O})\text{NHMe}$$
 Smith, 1966, Vol. 2, 459
 The liquid deflagrates on heating.
See other NITROSO COMPOUNDS
- 0947. 1-Methyl-3-nitroguanidinium nitrate**
 [] **C₂H₇N₅O₅**

$$\text{MeN}^+\text{H}_2\text{C}(\text{:NH})\text{NHNO}_2\text{NO}_3^-$$
 McKay, A. F. *et al.*, *J. Amer. Chem. Soc.*, 1947, **69**, 3029
 The salt could be exploded by impact between steel surfaces.
See other N-NITRO COMPOUNDS, OXOSALTS OF NITROGENOUS BASES
- 0948. Dimethylphosphine**
 [676-59-5] **C₂H₇P**

$$\text{Me}_2\text{PH}$$
 1. Houben-Weyl, 1963, Vol. 12.1, 69
 2. Parshall, G. W., *Inorg. Synth.*, 1968, **11**, 158
 3. Trenkle, A. *et al.*, *Inorg. Synth.*, 1982, **21**, 180–181
 Secondary lower-alkylphosphines readily ignite in air [1]; preparative precautions are detailed [2,3].
See other ALKYLPHOSPHINES
- 0949. Ethylphosphine**
 [593-68-0] **C₂H₇P**

$$\text{EtPH}_2$$
 Houben-Weyl, 1963, Vol. 12.1, 69
 Primary lower-alkylphosphines readily ignite in air.
 Halogens, or Nitric acid
 von Schwartz, 1918, 324–325


It explodes on contact with chlorine, bromine or fuming nitric acid, and inflames with conc. acid.

See other ALKYLPHOSPHINES

0950. Dimethylammonium perchlorate

[14488-49-4]

$C_2H_8ClNO_4$



1. Gore, P. H., *Chem. Brit.*, 1976, **12**, 205
2. Menzer, M. *et al.*, *Z. Chem.*, 1977, **17**, 344
3. Gore, P. H. *et al.*, *Synth. Commun.*, 1980, **10**, 320

A violent explosion occurred during vacuum evaporation of an aqueous mixture of excess dimethylamine and perchloric acid [1], and a similar incident occurred when the moist solid from a like preparation was moved [2]. Use of this explosive salt as an intermediate for the preparation of 3-dimethylamino-2-propenylidenedimethylimmonium perchlorate should be avoided, the methylsulfate or triflate salts being preferred [3].

See other PERCHLORATE SALTS OF NITROGENOUS BASES

0951. 2-Hydroxyethylaminium perchlorate (Ethanolamine perchlorate)

[38092-76-1]

$C_2H_8ClNO_5$



Annikov, V. E. *et al.*, *Chem. Abs.*, 1983, **99**, 160860

Ethanolamine perchlorate was outstanding amongst other nitrogenous perchlorate salts for its capacity for detonation as an aqueous solution.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

0952. Dimethyl phosphoramidate

[2697-42-9]

$C_2H_8NO_3P$



Chlorine

See Chlorine: Dimethyl phosphoramidate

See other PHOSPHORUS ESTERS

0953. 1,2-Diaminoethane (Ethylenediamine)

[107-15-3]

$C_2H_8N_2$



Cellulose nitrate

See CELLULOSE NITRATE: Amines

Diisopropyl peroxydicarbonate

See Diisopropyl peroxydicarbonate: Amines, etc.

Nitromethane

See Nitromethane: Acids, etc., and: 1,2-Diaminoethane, etc.

Other reactants

Yoshida, 1980, 50

MRH values calculated for 13 combinations with oxidants are given.

Silver perchlorate

See Silver perchlorate: 1,2-Diaminoethane

See other ORGANIC BASES

†0954. 1,1-Dimethylhydrazine

[57-14-7]

$C_2H_8N_2$

Me_2NNH_2

HCS 1980, 431

Energy of decomposition (in range 280–380°C) measured as 1.15 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Dichloromethane

See Dichloromethane: 1,2-Diaminoethane

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

Oxidants

1. Kirk-Othmer, 1966, Vol. 11, 186

2. Wannagat, U. *et al.*, *Monatsh.*, 1969, **97**, 1157–1162

3. *49IM*, 1975, 158

A powerful reducing agent and fuel, hypergolic with many oxidants, such as dinitrogen tetroxide, hydrogen peroxide and nitric acid [1]. The ignition delay with fuming nitric acid was determined as 8 ms, explosion also occurring [2]. When spread (as a thin film, or absorbed on porous or fibrous material) to expose a large surface to air, ignition may occur [3].

See Hexanitroethane: 1,1-Dimethylhydrazine

See also ROCKET PROPELLANTS

See other ORGANIC BASES, REDUCANTS

†0955. 1,2-Dimethylhydrazine

[540-73-8]

$C_2H_8N_2$

$MeNHNHMe$

See other ORGANIC BASES, REDUCANTS

- 0956. 1-Hydroxyethylidene-1,1-diphosphonic acid**
 [2809-21-4] **C₂H₈O₇P₂**

$$\text{MeCOH[P(O)(OH)}_2\text{]}_2$$
 'Data Sheet ADPA 60A', Oldbury, Albright & Wilson, 1977
 The anhydrous acid decomposes, often violently, at above 200°C, to give phosphine, phosphoric acid and other products.
See other ORGANIC ACIDS
- 0957. 1,2-Diphosphinoethane**
 [5518-62-7] **C₂H₈P₂**

$$\text{H}_2\text{PC}_2\text{H}_4\text{PH}_2$$
 Taylor, R. C. *et al.*, *Inorg. Chem.*, 1973, **14**, 10–11
 It ignites in air.
See other ALKYLPHOSPHINES
- 0958. 2-Aminoethylammonium perchlorate**
 [25682-07-9] **C₂H₉ClN₂O₄**


$$\text{H}_2\text{NC}_2\text{H}_4\text{N}^+\text{H}_3\text{ClO}_4^-$$
 Hay, R. W. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1975, 1467
 Solutions of the perchlorate salts of 1,2-diaminoethane and other amines must be evaporated without heating to avoid the risk of violent explosions.
See other PERCHLORATE SALTS OF NITROGENOUS BASES
- 0959. Dimethyl hydrazidophosphate (Dimethylphosphorohydrazine)**
 [58816-51-8] **C₂H₉N₂O₃P**

$$(\text{MeO})_2\text{P(O)NHNH}_2$$
 Anon., *Sichere Chemiarbeit*, 1990, **42**, 56
 The compound decomposed explosively upon vacuum distillation at 150°C.
 Caution is advised with similar hydrazides.
See related PHOSPHORUS ESTERS
- †**0960. 1,1-Dimethyldiborane**
 [16924-32-6] **C₂H₁₀B₂**

$$\text{Me}_2\text{B:H}_2\text{:BH}_2$$
See other ALKYLBORANES
- †**0961. 1,2-Dimethylborane**
 [17156-88-6] **C₂H₁₀B₂**

$$\text{MeHB:H}_2\text{:BHMe}$$
See other ALKYLBORANES

0962. B-Chlorodimethylaminodiborane
[]



Burg, A. B. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 3454

It ignites in air.

See related BORANES, HALOBORANES

0963. 2,2-Dimethyltriazanium perchlorate
[20446-69-9]




Rubtsov, Yu. I. *et al.*, *Chem. Abs.*, 1981, **95**, 117860

It decomposes rapidly above 122°C.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

0964. 1,2-Ethylenebis(ammonium) perchlorate
[15718-71-5]




Lothrop, W. C. *et al.*, *Chem. Rev.*, 1949, **44**, 432

An explosive which appreciably exceeds the power and brisance of TNT.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

0965. Aqua-1,2-diaminoethanediperoxochromium(IV)
[17168-82-0]



1. Childers, R. F. *et al.*, *Inorg. Chem.*, 1968, **7**, 749


2. House, D. A. *et al.*, *Inorg. Chem.*, 1966, **5**, 840

The monohydrate is light-sensitive and explodes at 96–97°C if heated at 2°/min [1]. It effervesces vigorously on dissolution in perchloric acid [2].

See other AMMINECHROMIUM PEROXOCOMPLEXES

0966. 1,2-Diammonioethane nitrate (Ethylenediamine dinitrate)

[20829-66-7]



$\text{C}_2\text{H}_{10}\text{N}_4\text{O}_6$

Dobratz, B. M., *Chem. Abs.*, 1984, **100**, 88163


Ethylenediamine dinitrate was formerly used as a military explosive.

See other OXOSALTS OF NITROGENOUS BASES

0967. Dimethylaminodiborane

[23273-02-1]

$\text{C}_2\text{H}_{11}\text{B}_2\text{N}$



Keller, P. G., *Inorg. Synth.*, 1977, **17**, 34

It ignites in air, like the parent diborane.

See related BORANES

0968. Ammine-1,2-diaminoethanediperoxochromium(IV)

[17168-82-0]



$\text{C}_2\text{H}_{11}\text{CrN}_3\text{O}_4$

House, D. A. *et al.*, *Inorg. Chem.*, 1967, **6**, 1077

The monohydrate is potentially explosive at 25°C and decomposes or explodes at 115° during slow or moderate heating.

Hydrogen bromide

Hughes, R. G. *et al.*, *Inorg. Chem.*, 1968, **7**, 74

Interaction must be slow with cooling to prevent explosion.

See other AMMINECHROMIUM PEROXOCOMPLEXES

0969. Tetraamminedithiocyanatocobalt(III) perchlorate

[36294-69-6] (ion)



$\text{C}_2\text{H}_{12}\text{ClCoN}_6\text{O}_4\text{S}_2$

Tomlinson, W. R. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 375


Explodes at 335°C, medium impact-sensitivity.

See other AMMINOMETAL OXOSALTS

0970. Tetraamminebis(5-nitro-2H-tetrazolato)cobalt(1⁺) perchlorate

[117412-28-9]

C₂H₁₂ClCoN₁₄O₈



Fronabarger, J. W., *et al.*, *Proc. Int. Pyrotech. Semin.*, 1996, 22nd, 645


It is being developed as an initiating explosive.

See other AMMINEMETAL OXOSALTS, TETRAZOLES

0971. Guanidinium dichromate

[27698-99-3]

C₂H₁₂Cr₂N₆O₇



Ma, C., *J. Amer. Chem. Soc.*, 19512, **73**, 1333–1335

Heating causes orderly decomposition, but under confinement, a violent explosion.

See other DICROMATE SALTS OF NITROGENOUS BASES

0972. Ammonium 1,2-ethylenebis(nitramide)

[3315-89-7]

C₂H₁₂N₆O₄



Violent decomposition occurred at 191°C.


See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1)

See other N-NITRO COMPOUNDS

0973. Ethylpentaborane(9)

[28853-06-7]

C₂H₁₃B₅



Leleu, *Cahiers*, 1977, (88), 365

It ignites in air.

See other ALKYLBORANES

0974. 2(5-Cyanotetrazole)pentaamminecobalt(III) perchlorate
[70247-32-4] $C_2H_{16}Cl_3CoN_{10}O_{12}$




Loyola, V. M. *et al.*, *Chem. Abs.*, 1981, **95**, 222342

It is a practical explosive of zero oxygen balance, but much less sensitive to accidental explosion than conventional detonators.

See other AMMINEMETAL OXOSALTS, TETRAZOLES

0975. Mercury(II) acetylide

[37297-87-3]




Bailar, 1973, Vol. 3, 314–315

Possibly polymeric, it explodes when heated or shocked.

See other MERCURY COMPOUNDS, METAL ACETYLIDES

0976. Mercury(II) cyanide

[592-04-1]



Ullmann, A8, 176

It is a moderately endothermic compound but of high MW (ΔH_f° (s) +261.5 kJ/mol, 1.035 kJ/g). It may explode if detonated, although it is used as an inerting diluent for the more sensitive and more powerful oxycyanide, $Hg(CN)_2 \cdot HgO$.

Fluorine

See Fluorine: Metal salts

Hydrogen cyanide

Wöhler, L. *et al.*, *Chem. Ztg.*, 1926, **50**, 761

The cyanide is a friction- and impact-sensitive explosive, and may initiate detonation of liquid hydrogen cyanide. Other heavy metal cyanides are similar.

Magnesium

See Magnesium: Metal cyanides

Sodium nitrite

See Sodium nitrite: Metal cyanides

See other ENDOTHERMIC COMPOUNDS, MERCURY COMPOUNDS, METAL CYANIDES

0977. Mercury(II) cyanate

[3021-39-4]



Biasutti, 1981, 53

Two explosions during the crushing of the cyanate are recorded.

See related METAL CYANIDES (AND CYANO COMPLEXES) *See other* MERCURY COMPOUNDS

0978. Mercury(II) fulminate
[628-86-4]



NSC 309, 1978

Alone, or Sulfuric acid

1. Urbanski, 1967, Vol. 3, 135, 140
2. Carl, L. R., *J. Franklin Inst.*, 1945, **240**, 149

Mercury fulminate, readily formed by interaction of mercury(II) nitrate, nitric acid and ethanol, is endothermic (ΔH_f° (s) +267.7 kJ/mol, 0.94 kJ/g) and was a very widely used detonator. It may be initiated when dry by flame, heat, impact, friction or intense radiation. Contact with sulfuric acid causes explosion [1]. The effects of impurities on the preparation and decomposition of the salt have been described [2].

See other ENDOTHERMIC COMPOUNDS, MERCURY COMPOUNDS, METAL FULMINATES

0979. Mercury(II) thiocyanate
[592-85-8]




HCS 1980, 619

Unpublished information, 1950

It is a moderately endothermic and thermally unstable compound of high MW (ΔH_f° (s) +200.8 kJ/mol, 0.63 kJ/g). A large batch of the damp salt became overheated in a faulty steam drying oven, and decomposed vigorously, producing an enormous 'Pharaoh's serpent'.


See related METAL CYANIDES (AND CYANO COMPLEXES) *See other* ENDOTHERMIC COMPOUNDS, MERCURY COMPOUNDS

0980. Mercury 5,5'-azotetrazolide
[87552-19-0]



See entry METAL AZOTETRAZOLIDES *See other* MERCURY COMPOUNDS, TETRAZOLES

0981. Mercury(II) 5-nitrotetrazolide
[60345-95-1]



1. Gilligan, W. H. *et al.*, *Rept.AD-A036086*, Richmond (Va.), USNTIS, 1976
2. Farncomb, R. E. *et al.*, *Chem. Abs.*, 1977, **87**, 203672
3. Redman, L. D. *et al.* *Chem. Abs.*, 1984, **100**, 194540

It is a promising replacement for lead azide in detonators [1], and a remotely controlled procedure for the preparation of this oxygen-balanced compound is described [2]. Preparative methods have been assessed for safety features [3].

See other MERCURY COMPOUNDS, C-NITRO COMPOUNDS, TETRAZOLES

0982. Mercury(II) oxalate
[3444-13-1]




1. Anon., *ABCM Quart. Safety Summ.*, 1953, **24**, 30, 45
2. Muir, G. D., private comm., 1968

When dry, the oxygen-balanced salt explodes readily on percussion, grinding or heating to 105°C. This instability is attributed to presence of impurities (nitrate, oxide, or basic oxalate) in the product [1]. It is so thermally unstable that storage is inadvisable [2].

See other CATALYTIC IMPURITY INCIDENTS, MERCURY COMPOUNDS, METAL OXALATES

See entry OXYGEN BALANCE

0983. Dimercury dicyanide oxide (Mercury(II) oxycyanide)
[1335-31-5]



1. Merck Index, 1983, 840; *ibid.* 1989, 5781
2. Kast, H. *et al.*, *Chem. Abs.*, 1922, **16**, 4065
3. *MCA Guide*, 1972, 309

The compound is explosive when pure, and sensitive to impact or heat. It is stabilised for commerce by the presence of a major excess of mercury(II) cyanide [1]. Several explosive incidents have been described [2], most involving friction [3].

See related METAL CYANIDES (AND CYANO COMPLEXES) *See other* FRICTIONAL INITIATION INCIDENTS, MERCURY COMPOUNDS

0984. Thallium(I) iodacetylide

[]

C₂ITl

Fawcett, H. H., private comm., 1984

The acetylide, produced as an unexpected major product, is a shock- and friction-sensitive explosive. A few mg exploded when touched with a spatula, and the remaining 2 g sample detonated violently when destroyed by a controlled explosion.

See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

0985. Diiodoacetylene (Diiodoethyne)

[624-74-8]


C₂I₂1. Anon., *Sichere Chemiearb.*, 1955, **7**, 552. Vaughn, T. H. *et al.*, *J. Amer. Chem. Soc.*, 1932, **54**, 7893. Taylor, G. N., *Chem. Brit.*, 1981, **17**, 107

Pure recrystallised material exploded while being crushed manually in a mortar. The decomposition temperature is 125°C, and this may have been reached locally during crushing [1]. Explosion on impact, on heating to 84°C, and during attempted distillation at 98°C/5 mbar had been recorded previously [2]. During an attempt to repeat the published preparation [2], a relatively large amount of the very sensitive explosive, nitrogen triiodide (*ibid.*), was isolated [3].

See other HALOACETYLENE DERIVATIVES

0986. Tetraiodoethylene

[513-92-8]

C₂I₄

Iodine pentafluoride

See Iodine pentafluoride: Tetraiodoethylene

See other HALOALKENES, IODINE COMPOUNDS

0987. Potassium acetylide

[22754-96-7]

C₂K₂

Water


Bahme, 1972, 80

Contact with limited amounts of water may cause ignition and explosion of evolved acetylene.

See other METAL ACETYLIDES

0988. Potassium dinitrooxalatoplatinate(2-)
[15213-49-7]

C₂K₂N₂O₈Pt




Vèzes, M. *Compt. rend.*, 1897, **125**, 525

The salt decomposes violently at 240°C.

See related METAL OXALATES *See other* PLATINUM COMPOUNDS

0989. Potassium 1,1,2,2-tetranitroethanediide
[32607-31-1]

C₂K₂N₄O₈



1. Borgardt, F. G. *et al.*, *J. Org. Chem.*, 1966, **31**, 2806, 2810

2. Griffin, T. S. *et al.*, *J. Org. Chem.*, 1980, **45**, 2882

This anhydrous salt, and the mono- and di-hydrates of the analogous lithium and sodium salts, are all very impact-sensitive. The potassium salt, an intermediate in the preparation of hexanitroethane [1], was not allowed to become dry during isolation, but after precipitation was washed with methanol and dichloromethane and used wet with the latter for the succeeding nitration stage [2].

See other POLYNITROALKYL COMPOUNDS

0990. Potassium acetylene-1,2-dioxide (Potassium ethynediolate)
[2851-55-0]

C₂K₂O₂



Air, or Halocarbons, or Halogens, or Protic compounds

Taylor, C. K., *Chem. Abs.*, 1983, **99**, 32105

Produced by action of carbon monoxide on potassium in liquid ammonia at -50°C, the yellow powder burns explosively in contact with air, halocarbons, halogens, alcohols, water and any material with acidic hydrogen. Analogous metal derivatives are reviewed.

See Potassium benzenehexoxide

See other ACETYLENIC COMPOUNDS

0991. Lanthanum carbide
[12071-15-7]

C₂La



Borlas, R. A. *et al.*, *Chem. Abs.*, 1976, **84**, 182199

'On the flammability and explosiveness of lanthanum carbide and cerium carbide powders' (title only translated).

See other METAL NON-METALLIDES

0992. Lithium acetylide

[1070-75-3]



C_2Li_2

Halogens

Mellor, 1946, Vol. 5, 848

It burns brilliantly when cold in fluorine or chlorine, but must be warm before ignition occurs in bromine or iodine vapours.

Lead(II) oxide

See Lead(II) oxide: Metal acetylides

Non-metals

Mellor, 1946, Vol. 5, 848

It burns vigorously in phosphorus, selenium or sulfur vapours.

See other METAL ACETYLIDES

0993. Lithium ethynediolate

[88906-07-4]



$\text{C}_2\text{Li}_2\text{O}_2$

Weiss, E., *Angew. Chem. (Int.)*, 1993, **32**(11), 1518


Too shock sensitive to obtain crystallographic data.

See Lithium benzenhexoxide

See other ACETYLENIC COMPOUNDS

0994. Lithium 1,1,2,2-tetranitroethanediide

[]



$\text{C}_2\text{Li}_2\text{N}_4\text{O}_8$

See Potassium 1,1,2,2-tetranitroethanediide

See other POLYNITROALKYL COMPOUNDS

0995. Dicarbonylmolybdenum diazide

[68348-85-6]



$\text{C}_2\text{MoN}_6\text{O}_2$

Alone, or Water

Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 513

This and the tungsten analogue (both homopolymeric) are extremely sensitive, exploding on the slightest mechanical stress, or violently in contact with traces of water.

See related CARBONYLMETALS, METAL AZIDES

†0996. Dicyanogen (Ethanedinitrile)

[460-19-5]

C₂N₂



Oxidants

Dicyanogen is extremely endothermic (ΔH_f° (g) +307.9 kJ/mol, 5.91 kJ/g) and the potential energy of mixtures with powerful oxidants may be released explosively under appropriate circumstances.

See Dichlorine oxide: Dicyanogen

Fluorine: Halogens

Oxygen (Liquid): Liquefied gases

Ozone: Dicyanogen

ROCKET PROPELLANTS

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

0997. Nickel(II) cyanide

[557-19-7]

C₂N₂Ni



It is a somewhat endothermic compound (ΔH_f° (s) +113.4 kJ/mol, 1.02 kJ/g), though not notably thermally unstable.

Magnesium

See Magnesium: Metal cyanides

See other ENDOTHERMIC COMPOUNDS, METAL CYANIDES

0998. Dicyanogen *N,N'*-dioxide

[4331-98-0]

C₂N₂O₂



Grundmann, C. J., *Angew. Chem.*, 1963, **75**, 450; *Ann.*, 1965, **687**, 194

The solid decomposes at -45°C under vacuum, emitting a brilliant light before exploding.

See related CYANO COMPOUNDS, HALOGEN OXIDES

0999. Lead(II) cyanide

[592-09-2]

C₂N₂Pb



Magnesium

See Magnesium: Metal cyanides

See other METAL CYANIDES

1000. Lead(II) thiocyanate

[592-87-0]

 $C_2N_2PbS_2$ 

Urbanski, 1967, Vol. 3, 230

The explosive properties of lead thiocyanate have found limited use.

See related METAL CYANIDES (AND CYANO COMPLEXES) *See other* HEAVY METAL DERIVATIVES**1001. Thiocyanogen**

[505-14-6]

 $C_2N_2S_2$ Söderbäck, E., *Ann.*, 1919, **419**, 217Low-temperature storage is necessary, as it polymerises explosively above its m.p., $-2^\circ C$ ($-7^\circ C$ is also recorded).*See other* POLYMERISATION INCIDENTS*See related* CYANO COMPOUNDS**1002. Sulfur thiocyanate**

[57670-85-6]

 $C_2N_2S_3$ Lecher, H. *et al.*, *Ber.*, 1922, **55**, 1481

It decomposes explosively (but harmlessly) on storage at ambient temperature.

See related CYANO COMPOUNDS *See other* N-S COMPOUNDS**1003. Disulfur thiocyanate**

[]

 $C_2N_2S_4$ Lecher, H. *et al.*, *Ber.*, 1922, **55**, 1485It melts at $-3.5^\circ C$ and decomposes explosively (but harmlessly) after turning yellow.*See related* CYANO COMPOUNDS *See other* N-S COMPOUNDS**1004. Zinc cyanide**

[557-21-1]

 C_2N_2Zn It is modestly endothermic (ΔH_f° (s) $+77$ kJ/mol, 0.65 kJ/g) but not notably thermally unstable.

Magnesium

See Magnesium: Metal cyanides*See other* ENDOTHERMIC COMPOUNDS, METAL CYANIDES

1005. Dicyanodiazene (Azocarbonitrile)

[1557-57-9]

C₂N₄1. Marsh, F. D. *et al.*, *J. Amer. Chem. Soc.*, 1965, **87**, 18192. Ittel, S. D. *et al.*, *Inorg. Chem.*, 1975, **14**, 1183

The solid explodes when mechanically shocked or heated in a closed vessel [1]. Preparative methods are hazardous because of the need to heat the explosive precursor, cyanogen azide [1,2].

See Cyanonitrene See other AZO COMPOUNDS, CYANO COMPOUNDS

1006. Disodium dicyanodiazene

[]


C₂N₄Na₂Marsh, F. D. *et al.*, *J. Amer. Chem. Soc.*, 1965, **87**, 1820

The radical anion salt is an explosive powder.

See related AZO COMPOUNDS, CYANO COMPOUNDS

1007. Sodium 1,1,2,2-tetranitroethanediide

[4415-23-0]


C₂N₄Na₂O₈

See Potassium 1,1,2,2-tetranitroethanediide

See other POLYNITROALKYL COMPOUNDS

1008. 3,4-Dinitrofurazan-2-oxide

[153498-61-4]


C₂N₄O₆Godovikova, T. L., *Mendeleev Comm.*, 1993, (5) 209

This oxygen rich explosive is claimed to be the first pernitro heterocycle known.

See other FURAZAN N-OXIDES, C-NITRO COMPOUNDS

1009. Trinitroacetoneitrile

[630-72-8]

C₂N₄O₆1. Schischkow, A., *Ann. Chim. [3]*, 1857, **49**, 3102. Parker, C. O. *et al.*, *Tetrahedron*, 1962, **17**, 79, 84


It explodes if heated quickly to 220°C [1]. It is also a friction- and impact-sensitive explosive, which may be used conveniently in carbon tetrachloride solution to minimise handling problems [2].

See other CYANO COMPOUNDS, POLYNITROALKYL COMPOUNDS

1010. Tetranitroethylene

[13223-78-4]

C₂N₄O₈



Baum, K, *et al.*, *J. Org. Chem.*, 1985, **50**, 2736–2739

This exceptionally reactive dienophile (at least one order of magnitude greater than tetracyanoethylene) reacted normally with dilute solutions of unsaturated compounds in dichloromethane, but sometimes explosively in absence of solvent.

See other NITROALKENES

1011. Thiocarbonyl azide thiocyanate

[]

C₂N₄S₂



CHETAH, 1990, 182

Shock sensitive

Ammonia, or Hydrazine

Audrieth, L. F. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 2799–2805


The unstable (undoubtedly endothermic) compound reacts explosively with ammonia gas, and violently with conc. hydrazine solutions.

See related CYANO COMPOUNDS See other ORGANIC AZIDES

1012. Disodium 5-tetrazolazocarboxylate

[68594-24-1]

C₂N₆Na₂O₂



Thiele, J. *et al.*, *Ann.*, 1895, **287**, 238

It is explosive.

See other AZO COMPOUNDS, TETRAZOLES

1013. Dicarbonyltungsten diazide

[68379-32-8]

C₂N₆O₂W



Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 513

This and the molybdenum analogue (both homopolymeric) are extremely sensitive, exploding on the slightest mechanical stress, or violently in contact with traces of water.

See related CARBONYLMETALS, METAL AZIDES

1014. Hexanitroethane

[918-37-6]



$\text{C}_2\text{N}_6\text{O}_{12}$

1. Loewenschuss, A. *et al.*, *Spectrochim. Acta*, 1974, **30A**, 371–378
2. Sorbe, 1968, 149

Grinding the solid to record its IR spectrum was precluded on safety grounds [1]. It decomposes explosively above 140°C [2].

Boron

Finnerty, A. E. *et al.*, *Chem. Abs.*, 1979, **91**, 177486

There is an autocatalytic reaction at ambient temperature in this new explosive combination.

1,1-Dimethylhydrazine

Noble, P. *et al.*, *Am. Inst. Aeron. Astronaut. J.*, 1963, **1**, 395–397

Hexanitroethane is a powerful oxidant and hypergolic with dimethylhydrazine or other strong organic bases.

Organic compounds

Will, M., *Ber.*, 1914, **47**, 961–965

Though relatively insensitive to friction, impact or shock, it can be detonated. With hydrogen containing organic compounds, this oxygen-rich compound (+200% oxygen balanced) forms powerfully explosive mixtures. The addition compound with 2-nitroaniline ($\text{C}_8\text{H}_6\text{N}_8\text{O}_{14}$, –27% balance) is extremely explosive.

See other POLYNITROALKYL COMPOUNDS

1015. Palladium(II) azidodithioformate

[29149-89-1]



$\text{C}_2\text{N}_6\text{PdS}_4$

Fehlhammer, W. P. *et al.*, *Z. Naturforsch.*, 1983, **38B**, 547

In the first stage of the preparation of bis(triphenylphosphine)palladium(II) isothiocyanate, a large deficiency of aqueous palladium nitrate must be added with rapid stirring to an excess of sodium azidodithioformate solution to avoid the precipitation of explosive palladium(II) azidodithioformate.

See Azidodithioformic acid

See other ACYL AZIDES

1016. Bis(azidothiocarbonyl) disulfide

[]

C₂N₆S₄Smith, G. B. L., *Inorg Synth.*, 1939, **1**, 81


This compound, readily formed by iodine oxidation of azidodithioformic acid or its salts, is a powerful explosive. It is sensitive to mechanical impact or heating to 40°C, and slow decomposition during storage increases the sensitivity. Preparative precautions are detailed.

See Azidodithioformic acid

See other ACYL AZIDES

1017. Diazidomethylenecyanamide

[67880-22-2]


C₂N₈Marsh, F. D., *J. Org. Chem.*, 1972, **37**, 2967

This explosive solid may be produced during preparation of cyanogen azide.

See other CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1018. Disodium 5,5'-azotetrazolide


[41463-64-3]

C₂N₁₀Na₂

See entry METAL AZOTETRAZOLIDES See other TETRAZOLES

1019. Disodium 5,5'-azoxytetrazolide

[]


C₂N₁₀Na₂OThiele, J. *et al.*, *Ann.*, 1893, **273**, 151

Insensitive to friction or impact, but explodes violently on heating in a melting point tube or on analytical combustion, destroying the apparatus in both cases.

See related METAL AZOTETRAZOLIDES See other N-OXIDES, TETRAZOLES

1020. Lead 5,5'-azotetrazolide
[87489-52-9]

C₂N₁₀Pb



See entry METAL AZOTETRAZOLIDES See other TETRAZOLES

1021. Diazidomethyleneazine
[]

C₂N₁₄




Houben-Weyl, 1965, Vol. 10.3, 793

This very explosive bis-*gem*-diazide contains over 89% of nitrogen.

See other HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1022. Sodium acetylide (Sodium ethynide)
[2881-62-1]

C₂Na₂



1. Opolsky, S., *Bull. Acad. Cracow*, 1905, 548

2. Houben-Weyl, 1977, Vol. 5.2a, 361

A brown explosive form is produced if excess sodium is used in preparation of thiophene homologues — possibly because of sulfur compounds [1]. As normally produced, it is a dry stable solid, but material prepared from acetylene and sodium—oil dispersions ignites in air [2].

Halogens

Mellor, 1946, Vol. 5, 848

Sodium acetylide burns in chlorine and (though not stated) probably also in fluorine, and in contact with bromine and iodine on warming.

Metals

Mellor, 1946, Vol. 5, 848

Trituration in a mortar with finely divided aluminium, iron, lead or mercury may be violent, carbon being liberated.

Metal salts

Mellor, 1946, Vol. 5, 848

Rubbing in a mortar with some metal chlorides or iodides may cause incandescence or explosion. Sulfates are reduced, and nitrates would be expected to behave similarly.

Non-metal oxides

von Schwartz, 1918, 328

Sodium acetylide incandesces in carbon dioxide or sulfur dioxide.

Oxidants

Mellor, 1946, 5, 848

Ignites on warming in oxygen, and incandesces at 150°C in dinitrogen pentaoxide.

See Halogens, above

Phosphorus

See Phosphorus: Metal acetylides

Water

1. Mellor, 1946, Vol. 5, 848

2. Davidsohn, W. E., *Chem. Rev.*, 1967, **67**, 74

Excess water is necessary to prevent explosion [1]; the need for care in handling is stressed [2].

See other METAL ACETYLIDES

1023. Sodium ethynediolate

[2611-42-9]



See Sodium: Non-metal oxides

See other ACETYLENIC COMPOUNDS

1024. Sodium peroxydicarbonate

[3312-92-6]



Acetic anhydride

See Sodium carbonate hydrogen peroxide: Acetic anhydride

See other PEROXOACID SALTS

1025. Rubidium acetylide

[22754-97-8]



Acids

Mellor, 1946, Vol. 5, 848

With conc. hydrochloric acid ignition occurs, and contact with nitric acid causes explosion.

Halogens

Mellor, 1946, Vol. 5, 848

It burns in all four halogens.

Metal oxides

Mellor, 1946, Vol. 5, 848–850

Iron(III) oxide and chromium(III) oxide react exothermally, and lead oxide explosively. Copper oxide and manganese dioxide react at 350°C incandescently.

Non-metal oxides

Mellor, 1946, Vol. 5, 848

Warming in carbon dioxide, nitrogen oxide or sulfur dioxide causes ignition.

Non-metals

Mellor, 1946, Vol. 5, 848

It reacts vigorously with boron or silicon on warming, ignites with arsenic, and burns in sulfur or selenium vapours.

See other METAL ACETYLIDES

1026. Strontium acetylide

[12071-29-3]



$(\text{C}_2\text{Sr})_n$

Halogens

Mellor, 1946, Vol. 5, 862

Strontium acetylide incandesces with chlorine, bromine or iodine at 197, 174 and 182°C, respectively.

See other METAL ACETYLIDES

1027. Thorium dicarbide

[12674-40-7]



C_2Th

Non-metals, or Oxidants

Mellor, 1946, Vol. 5, 862

Contact with selenium or sulfur vapour causes the heated carbide to incandesce. Contact of the carbide with molten potassium chlorate, potassium nitrate or even potassium hydroxide causes incandescence.

See other METAL NON-METALLIDES

1028. Uranium dicarbide

[12071-33-9]



C_2U

1. Mellor, 1946, Vol. 5, 890

2. Sidgwick, 1950, 1071

Uranium carbide emits brilliant sparks on impact, ignites on grinding in a mortar [1], or on heating in air to 400°C [2].

Halogens

Mellor, 1946, Vol. 5, 891

Incandescens in warm fluorine, in chlorine at 300°C and weakly in bromine at 390°C.

Hydrogen chloride

See Hydrogen chloride: Metal acetylides or carbides

Nitrogen oxide

See Nitrogen oxide: Metal acetylides or carbides

Water

1. Sidgwick, 1950, 1071

2. Mellor, 1946, Vol. 5, 890–891

Interaction with hot water is violent [1], and the carbide ignites in steam at dull red heat [2].

See other METAL NON-METALLIDES

1029. Zirconium dicarbide

[12070-14-3]

C₂Zr

ZrC₂

Halogens

Mellor, 1946, Vol. 5, 885

It ignites in cold fluorine, and in chlorine, bromine and iodine at 250, 300 and 400°C, respectively.

See other METAL NON-METALLIDES

1030. Silver trifluoropropynide

[]

C₃AgF₃

AgC≡CCF₃

Henne, A. L. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 1042

Explosive decomposition on heating.

See other HALOACETYLENE DERIVATIVES, SILVER COMPOUNDS

1031. Aluminium carbide

[1299-86-1]

C₃Al₄

Al₄C₃


Oxidants

Mellor, 1946, Vol. 5, 872

Incandescence on warming with lead dioxide or potassium permanganate.


See other METAL NON-METALLIDES

1032. Potassium 1,3-dibromo-1,3,5-triazine-2,4-dione-6-oxide
[15114-46-2]



It may be expected to show similar properties to the chloro-analogue.
See Sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide
See other *N*-HALOIMIDES

1033. 1-Chlorodifluoromethyl-1-trifluoromethyldioxirane
[35356-48-3]




See Bis(trifluoromethyl)dioxirane
See other CYCLIC PEROXIDES

1034. Pentafluoropropionyl hypochlorite
[71359-61-0]



Tari, I. *et al.*, *Inorg. Chem.*, 1979, **18**, 3205–3208
Thermally unstable at 22°C, and explosive in the gas phase at pressures above 27–62 mbar.
See other ACYL HYPOHALITES

1035. Heptafluoroisopropyl hypochlorite
[22675-68-9]



Schack, C. J. *et al.*, *J. Amer. Chem. Soc.*, 1969, **91**, 2904

Material condensed at -95°C may suddenly decompose completely and vaporise.
See other HYPOHALITES

1036. Chlorocyanoacetylene (Chloropropyne nitrile)

[2003-31-8]

C_3ClN



Hashimoto, N. *et al.*, *J. Org. Chem.*, 1970, **35**, 675


Avoid contact with air at elevated temperatures because of its low (unstated) ignition temperature. Burns moderately in the open, but may explode in a nearly closed vessel. Presence of mono- and di-chloroacetylenes as impurities increases the flammability hazard, which may be reduced by addition of 1% of ethyl ether.

See other HALOACETYLENE DERIVATIVES

1037. Sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide

[2893-78-9]

$\text{C}_3\text{Cl}_2\text{N}_3\text{NaO}_3$



'Fi-Clor 60S', Brochure NH/FS/67.4, Loughborough, Fisons, 1967

This compound (sodium dichloroisocyanurate), used in chlorination of swimming pools, is a powerful oxidant and indiscriminate contact with combustible materials must be avoided. Ammonium salts and other nitrogenous materials are incompatible in formulated products. The dibromo analogue, used for the same purpose, will behave similarly.

Preparative hazard

See 1,3,5-Trichloro-1,3,5-triazinetrione: Cyanuric acid, etc.


Calcium hypochlorite

1. Jones, C., INTERNET: CatJones@FC. KERN.ORG
2. Kirk Othmer, 1998, Vol. 25, 592

Although calcium hypochlorite is used in conjunction with sodium dichloroisocyanurate (usually as the dihydrate) for swimming pool maintenance, they must not be mixed undiluted. Noxious fumes and violent explosion or fire can result [1]. Although the organic compound is, at first sight, at its highest oxidation level, it contains disguised ammonia, and nitrogen trichloride may appear on reaction with hypochlorites [2].

See other N-HALOIMIDES

1038. 2,4,6-Trichloro-1,3,5-triazine (Cyanuric chloride)
[108-77-0]



HCS 1980, 925

1. Anon., *Loss Prev. Bull.*, 1979, (025), 21–22
2. See entry SELF-ACCELERATING REACTIONS

Of the factors associated with the high reactivity of cyanuric chloride (high exothermicity, rapid hydrolysis in presence of water-containing solvents, acid catalysed reactions, liberation of up to 3 mol hydrogen chloride/mol of chloride, formation of methyl chloride gas with methanol, formation of carbon dioxide from bicarbonates), several were involved in many of the incidents recorded [1] (and given below). The acid catalysed self acceleration and high exothermicity are rated highest [2]. It is also a mildly endothermic compound (ΔH_f° (s) +91.6 kJ/mol, 0.49 kJ/g).

Acetone, Water

Anon., *Loss Prev. Bull.*, 1979, (025), 20

The chloride was to be purified by dissolution in dry acetone, but in error, acetone containing 40% of water was used. The acid-catalysed exothermic hydrolysis reaction of the chloride accelerated to runaway, and gas and vapour evolution ruptured the vessel, leading to fire and explosion.

See Methanol, also Water, both below

See other GAS EVOLUTION INCIDENTS, SELF-ACCELERATING REACTIONS

Allyl alcohol, Sodium hydroxide, Water

Anon., *Loss Prev. Bull.*, 1974, (001), 11; 1979, (029), 21

When aqueous sodium hydroxide was added to a mixture of the chloride and alcohol at 28°C instead of the normal 5°C, a rapidly accelerating reaction led to rupture of the bursting disc and a gasket, and subsequently to a flash-fire and explosion.

See Methanol, also Water, both below

2-Butanone, Sodium hydroxide, Water

Anon., *Loss Prev. Bull.*, 1979, (025), 21

The total product of a batch containing a chlorotriazine, water, sodium hydroxide and 2-butanone had not been discharged from the 11,000 l vessel when a further 1.5 t of cyanuric chloride was charged ready for the next batch. The ensuing rapid

exothermic hydrolysis led to eruption of the reactor contents, ignition and explosion of butanone vapour, and damage of 3 M dollars.

Dimethylformamide

Anon., *BCISC Quart. Safety Summ.*, 1960, **35**, 24

Cyanuric chloride reacts vigorously and exothermically with DMF after a deceptively long induction period. The 1:1 adduct initially formed decomposes above 60°C with evolution of carbon dioxide and formation of a dimeric unsaturated quaternary ammonium salt. Dimethylformamide is appreciably basic and is not a suitable solvent for acyl halides.

See other INDUCTION PERIOD INCIDENTS

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides

2-Ethoxyethanol

Anon., *Loss Prev. Bull.*, 1979, (025), 20

Alcoholysis of the chloride on the plant scale was effected at 40°C (with brine cooling) by adding portions to the alcohol alternately with finely crystalline disodium phosphate to neutralise the hydrogen chloride produced. On one occasion, use of coarsely crystalline sodium phosphate (of low surface area) reduced the rate of neutralisation, the mixture became acid, and a runaway exotherm to 170°C developed leading to eruption of vessel contents. On another occasion, accidental addition of sodium sulfate instead of phosphate led to a similar situation beginning to develop, but an automatic pH alarm allowed remedial measures to be instituted successfully.

See other NEUTRALISATION INCIDENTS

Methanol

Anon., *ABCM Quart. Safety Summ.*, 1960, **31**, 40

Cyanuric chloride dissolved in methanol reacted violently and uncontrollably with the solvent. This was attributed to the absence of an acid acceptor to prevent the initially acid catalysed (and later auto-catalysed) exothermic reaction of all 3 chlorine atoms simultaneously.

Methanol, Sodium hydrogen carbonate

Anon., *Loss Prev. Bull.*, 1979, (025), 19

A crust of residual cyanuric chloride left in a reactor from a previous batch reacted with the methanol (usually charged first) to form hydrogen chloride. When the base was added (usually before the chloride), vigorous evolution of carbon dioxide expelled some of the solvent. In a second incident, accidentally doubling the charge of cyanuric chloride but not the base, led to the development of free acid (which auto-catalyses the reaction with methanol), and a runaway reaction developed causing violent boiling of the solvent, methyl chloride evolution and damage to the plant.

See Methanol, above *See other* RUNAWAY REACTIONS

Sodium azide

See 2,4,6-Triazido-1,3,5-triazine (reference 2)

Water

MCA Case History No. 1869

A reaction mixture containing the chloride and water, held in abeyance before processing, developed a high internal pressure in the containing vessel. Hydrolysis (or alcoholysis) of the chloride becomes rapidly exothermic above 30°C.


See Allyl alcohol, also Methanol, both above

See related ACYL HALIDES See other ENDOTHERMIC COMPOUNDS

1039. 1,3,5-Trichloro-1,3,5-triazinetriene ('Trichloroisocyanuric acid')

[87-90-1]

$C_3Cl_3N_3O_3$



HCS 1980, 920

Combustible materials

'Fi-Clor 91' Brochure, Loughborough, Fisons, 1967

This compound, used in chlorination of swimming pools, is a powerful oxidant, and indiscriminate contact with combustible materials must be avoided. The tribromo analogue would be similar.

Cyanuric acid, Sodium hydroxide

Brennan, J. P., US Pat. 4 118 570, 1978

Formation of sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide ('sodium dichloroisocyanurate') from an aqueous slurry of the trione, cyanuric acid and sodium hydroxide is accompanied by evolution of nitrogen trichloride. Nitrogen purging will keep the concentration of the explosive gas below the lower explosive limit of 5–6%.

Water

Young, J. A., private comm., 1986

If mixed with a small amount of water, the conc. solution (with pH around 2) may explode, owing to evolution of nitrogen trichloride. It is believed that hydrolysis leads to formation of hypochlorous acid and dichloro-*s*-triazinetriene, and the protonated acid then attacks the C=N bonds in the triazine ring leading to formation of


chloramines and nitrogen trichloride. The dichloro compound is stable to acid in absence of hypochlorous acid.

See other N-HALOIMIDES

†1040. Trichloromethyl carbonate (Triphosgene)

[32315-10-9]

C₃Cl₆O₃



Hollingsworth, M. D., *Chem. Eng. News*, 1993, **70**(28), 2


A package of this, consisting of a wax-coated glass jar in vermiculite within a sealed can, was found to be pressurised when opened. The cap of the jar had cracked, and it was assumed that residual water on the vermiculite had reacted to form hydrogen chloride, phosgene and carbon dioxide. Similar mishaps had been experienced elsewhere. Less packing would have given rise to less danger; the supplier has, however, supplemented the four layers initially present with a fifth and a recommendation not to store long and then under refrigeration. (The editor would predict that there are possible contaminants which could rearrange this compound slowly to phosgene in absence of water).

See other PACKAGING INCIDENTS

1041. Tetrachloropropadiene

[18608-30-5]

C₃Cl₄



Tetracarbonylnickel


See Dicarbonyl- η -trichloropropenyldinickel chloride dimer

See other HALOALKENES

1042. Tetrachloroethylene carbonate

[22432-68-4]

C₃Cl₄O₃



Tributylamine


1. Fagenburg, D. R., *Chem. Eng. News*, 1982, **60**(33), 45

2. Melby, L. R., *Chem. Eng. News*, 1982, **60**(40), 2

Interaction of the two compounds led to the evolution of a toxic gas thought to be chlorine [1]. It is the far more poisonous phosgene, arising from the known base-catalysed disproportionation of the carbonate to oxalyl chloride and phosgene, which occurs even at ambient temperature [2]. (The editor knows that amides, too, catalyse this rearrangement and suspects that Lewis acids will also)

See other GAS EVOLUTION INCIDENTS

1043. 2,4,6-Tris(dichloroamino)-1,3,5-triazine (Hexachloromelamine)
[2428-04-8]



As a trifunctional dichloroamino compound, it is probably more reactive and less stable than the monochloroamino analogue.

Acetone, or Bases


491M, 1975, 197

Addition of acetone, ammonia, aniline or diphenylamine to the oxidant causes rapid exothermic reactions, with or without flame, and large amounts under confinement would explode. The trichloro analogue is similar, but less vigorous.

See 2,4,6-Tris(chloroamino)-1,3,5-triazine (next above)

See other N-HALOGEN COMPOUNDS

1044. Potassium tricyanodiperoxochromate(3-)
[65521-60-0]




Bailar, 1973, Vol. 4, 167

A highly explosive material, with internal redox features.

See other PEROXOACID SALTS, REDOX COMPOUNDS

1045. Trifluoromethyl iodine(III) isocyanate
[127510-65-0]




Naumann, D. *et al.*, *J. Fluor. Chem.*, 1989, **45**(3), 401

This compound explosively decomposes when warmed up to room temperature.

See other IODINE COMPOUNDS, ORGANIC ISOCYANATES

1046. Lithium trifluoropropynide
[14856-86-1]




See Lithium chloroacetylide

See other HALOACETYLENE DERIVATIVES, METAL ACETYLIDES

1047. 2-Trifluoroacetyl-1,3,4-dioxazalone
[87050-94-0]

C₃F₃NO₃




Middleton, W. J., *J. Org. Chem.*, 1983, **48**, 3845

A sample contained in a dropping funnel at 25°C exploded forcefully. Though not sensitive to mechanical shock, it could be detonated by a hot wire when unconfined, or if rapidly heated to 220°C when confined in a stainless steel capillary.

See 1,3,4-DIOXAZOLONES

1048. Dicyanoiodonium triflate
[]

C₃F₃IN₂O₃S



Steng, P. J. *et al.*, *Tetrahedron Lett.*, 1992, **33**(11), 1419

This iodinating reagent is storable under nitrogen at -20°C; it may explode after contact with air. It decomposes in minutes at ambient temperatures.

See other CYANO COMPOUNDS, IODINE COMPOUNDS

1049. Trifluoroacryloyl fluoride
[667-49-2]

C₃F₄O



Sodium azide

Middleton, W. J., *J. Org. Chem.*, 1973, **38**, 3294

The product of interaction was an unidentified highly explosive solid.

See other ACYL HALIDES

1050. O-Trifluoroacetyl-S-fluoroformyl thioperoxide
[27961-70-2]

C₃F₄O₃S




Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 378

It exploded spontaneously in a glass bomb closed with a PTFE-lined valve. No previous indications of instability had been noted during distillation, pyrolysis or irradiation.

See related DIACYL PEROXIDES

†1051. Hexafluoropropene
[116-15-4]



C_3F_6

Air, Tetrafluoroethylene

See Tetrafluoroethylene: Air, Hexafluoropropene

Grignard reagents

Dinowski, W., *J. Fluorine Chem.*, 1981, **18**, 25–30

Reaction of various substituted phenylmagnesium bromides with hexafluoropropene under pressure at ambient temperature had been effected on the 0.2–0.6 g mol scale without incident. An attempt to scale up to 0.8 g mol with phenylmagnesium bromide led to an explosion.

Oxygen, Oxygen difluoride

See Oxygen difluoride: Hexafluoropropene, etc.

See other HALOALKENES

1052. Hexafluoroisopropylideneaminolithium
[31340-36-0]




$\text{C}_3\text{F}_6\text{LiN}$

Non-metal halides

Swindell, R. F., *Inorg. Chem.*, 1972, **11**, 242

Interaction of the lithium derivative with a range of chloro- and fluoro-derivatives of arsenic, boron, phosphorus, silicon and sulfur during warming to 25°C tended to be violently exothermic in absence of solvent. Thionyl chloride reacted with explosion. See Hexafluoroisopropylideneamine: Butyllithium See other N-METAL DERIVATIVES

1053. Bis(trifluoromethyl)cyanophosphine
[431-97-0]



$\text{C}_3\text{F}_6\text{NP}$

491M, 1975, 60

It ignites in air.

See related ALKYLHALOPHOSPHINES, CYANO COMPOUNDS

1054. Pentafluoropropionyl fluoride
[422-61-7]



$\text{C}_3\text{F}_6\text{O}$


Fluorinated catalysts

Sorbe, 1968, 2

The acid fluoride may decompose explosively in contact with fluorinated catalysts.
See other ACYL HALIDES

1055. Bis(trifluoromethyl)dioxirane
[35357-46-1]

$C_3F_6O_2$



Ando, W. (Ed.), *Organic Peroxides*, 195, Wiley, New York, 1992.

Formed by fluorine oxidation of the dilithium salt of hexafluoroacetone hydrate, it is unstable and explosive. The chloropentafluoro homologue is similar. Proponents of their use as reagents claim that the dimethyl and methyltrifluoromethyl analogues are not explosive; this seems improbable, especially since the less stressed lower dioxetanes (homodioxiranes) are all dangerous.

See other CYCLIC PEROXIDES

1056. Pentafluoropropionyl hypofluorite

[]

$C_3F_6O_2$

$F_5C_2CO.OF$

Menefee, A. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 2020

Less stable than its lower homologue, the hypofluorite explodes on sparking, on distillation at ambient pressure (b.p., 2°C), though not at below 0.13 bar.

See Fluorine: Caesium heptafluoropropoxide

See other ACYL HYPOHALITES

1057. Trifluoroacetyl trifluoromethanesulfonate

[68602-57-3]

$C_3F_6O_4S$

$F_3CCO.OSO_2CF_3$

Forbus, T. R. *et al.*, *J. Org. Chem.*, 1979, **44**, 313

The mixed anhydride reacts almost explosively with water.

See other ACID ANHYDRIDES, SULFUR ESTERS

1058. 1-Fluoroiminohexafluoropropane

[78343-32-5]

C_3F_7N

$FN=CFC_2F_5$


Fluorine

See Fluorine: 1- or 2-Fluoroiminohexafluoropropane

See other N-HALOGEN COMPOUNDS

1059. 2-Fluoroiminohexafluoropropane

[2802-70-2]



C₃F₇N

Fluorine

See Fluorine: 1- or 2-Fluoroiminohexafluoropropane

See other N-HALOGEN COMPOUNDS

1060. Heptafluoropropyl hypofluorite

[2203-57-8]



C₃F₈O

1. Sorbe, 1968, 62

2. *MCA Case History No. 1045*

An explosive compound [1], and the preparation is hazardous, involving the fluorination of caesium perfluoropropoxide [2].

See Fluorine: Caesium heptafluoropropoxide

See other HYPOHALITES

1061. 1,1-Bis(fluorooxy)hexafluoropropane

[72985-54-7]



C₃F₈O₂

Explosive, less stable than its lower homologue.

See entry BIS(FLUOROOXY)PERHALOALKANES

1062. 2,2-Bis(fluorooxy)hexafluoropropane

[16329-93-4]




C₃F₈O₂

It is especially prone to explosive decomposition.

See entry BIS(FLUOROOXY)PERHALOALKANES

1063. Tris(trifluoromethyl)phosphine

[432-04-2]



C₃F₉P

Oxygen

Mahler, W., *Inorg. Chem.*, 1979, **18**, 352

Contact with oxygen at 25°C/0.5 bar causes ignition of the phosphine.
See related ALKYLPHOSPHINES

1064. Sodium tricarbonylnitrosylferrate
[25875-18-7]



Davies, S. G. *et al.*, *J. Organomet. Chem.*, 1990, **386**(2), 195

This compound is air sensitive and decomposes explosively. Use of the more stable tetrabutylammonium salt as a reagent for carbonylation of alkyl halides is recommended.

1065. Propioloyl chloride (Propynoyl chloride)
[50277-65-1]



Balfour, W. J. *et al.*, *J. Org. Chem.*, 1974, **39**, 726

This chloride, purified by distillation at 58–60°C/1 bar usually ignites spontaneously in air owing to presence of chloroacetylene (formed by decarbonylation), but vacuum distillation at cryogenic temperatures prevents formation of the impurity.

See other ACYL HALIDES, ACETYLENIC COMPOUNDS

1066. 3,3,3-Trifluoropropyne
[661-54-1]




Haszeldine, R. N., *J. Chem. Soc.*, 1951, 590

It tends to explode during analytical combustion, and the copper(I) and silver derivatives decomposed violently (with occasional explosion) on rapid heating.

See other HALOACETYLENE DERIVATIVES

†**1067. Hexafluoroisopropylideneamine (2-Iminohexafluoropropane)**
[1645-75-6]




Butyllithium

Swindell, R. F. *et al.*, *Inorg. Chem.*, 1972, **11**, 242

The exothermic reaction which set in on warming the reagents in hexane to 0°C sometimes exploded if concentrated solutions of butyllithium (above 2.5 M) were used, but not if diluted (to about 1.2 M) with pentane.

1068. Potassium hydrogen diazirine-3,3-dicarboxylate

[]




See Diazirine-3,3-dicarboxylic acid

See other DIAZIRINES

1069. 4-Azidocarbonyl-1,2,3-thiadiazole

[]




Pain, D. L. *et al. J. Chem. Soc.*, 1965, 5167

The azide is extremely explosive in the dry state, though the oxygen balance is -54%.

See OXYGEN BALANCE See other ACYL AZIDES

1070. Diazopropyne

[17761-23-8]




Wierlacher, S. *et al.*, *Chem. Phys. Lett.*, 1994, **222**(4), 319

This compound is highly explosive and has detonated violently when in use on laboratory scale.

See other ACETYLENIC COMPOUNDS, DIAZO COMPOUNDS

1071. Silver malonate

[57421-56-4]



Sorbe, 1968, 126

It explodes on heating.

See other SILVER COMPOUNDS

1072. 2,2-Dinitro-2-fluoroethoxycarbonyl chloride
[31841-79-9]

$C_3H_2ClFN_2O_6$



Potassium picrate

Becuwe, A. G. *et al.*, US Pat. 4 145 361, 1979

Interaction to give the (explosive) ester, 2,2-dinitro-2-fluoroethyl 2,4,6-trinitrophenyl carbonate must be effected at below 50°C to prevent explosion.

See other ACYL HALIDES, FLUORODINITROMETHYL COMPOUNDS

1073. 3-Chloro-1-iodopropyne
[109-71-7]

C_3H_2ClI




Houben-Weyl, 1977, Vol. 5.2a, 610

Distillation at 47°C/5 mbar should only be taken to a 33% residue; less may lead to violent decomposition, as may admission of air to the hot residue.

See other HALOACETYLENE DERIVATIVES

†**1074. 2-Chloroacrylonitrile**
[920-37-6]

C_3H_2ClN



See other CYANO COMPOUNDS *See related* HALOALKENES

1075. Cyanoacetyl chloride
[16130-58-8]

C_3H_2ClNO



491M, 1975, 140

After treating cyanoacetic acid with phosphorus pentachloride, volatile materials were distilled off and the crude chloride was left in a stoppered flask. After 24 h the flask exploded.

See other ACYL HALIDES, CYANO COMPOUNDS

1076. Sodium diformylnitromethanide hydrate
[34461-00-2]

$C_3H_2NNaO_4.H_2O$



Fanta, P. E., *Org. Synth.*, 1962, Coll. Vol. 4, 844

The monohydrate, possibly the *gem*-dihydroxy compound and an *aci*-nitro salt, is an impact-sensitive solid and must be carefully handled with precautions.


See related C-NITRO COMPOUNDS, *aci*-NITRO SALTS

1077. Diisocyanomethane

[]

 $C_3H_2N_2$ Buschmann, J. *et al.*, *Angew. Chem. (Int.)*, 1997, **36**(21), 2372A clear crystalline solid at $-55^\circ C$, this compound melts, with browning at $-15^\circ C$, and decomposes explosively at $-10^\circ C$, leaving a polymer.*See other* DIISOCYANIDE LIGANDS**†1078. Malononitrile (Propanedinitrile)**

[109-77-3]

 $C_3H_2N_2$ Energy of decomposition (in range $180-270^\circ C$) measured as 1.65 kJ/g.*See entry* THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


Alone, or Bases

1. Personal experience
2. 'Malononitrile' Brochure, p. 11, Basle, Lonza Ltd., 1974
3. Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 29

It may polymerise violently on heating at $130^\circ C$, or in contact with strong bases at lower temperatures [1]. The stability of the molten nitrile decreases with increasing temperature and decreasing purity, but no violent decomposition below $100^\circ C$ has been recorded [2]. However, a partially filled drum of malononitrile stored in an oven at $70-80^\circ C$ for 2 months exploded violently [3].

See other CYANO COMPOUNDS**†1079. Diisocyanatomethane (Methylene diisocyanate)**

[4747-90-4]

 $C_3H_2N_2O_2$ 


Dimethylformamide

'DMF Chemical Uses', Brochure, Wilmington, Du Pont, 1967

Contact with the basic solvent causes violent polymerisation of the isocyanate.

See other POLYMERISATION INCIDENTS *See related* CYANO COMPOUNDS**1080. Diazirine-3,3-dicarboxylic acid**

[76429-98-6]

 $C_3H_2N_2O_4$ Schustov, G. V. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1981, **20**, 200

While the dipotassium salt is stable for over 2 months at 20°C, the monopotassium salt exploded within a few hours at 0°C. Both salts exploded when triturated.

See other DIAZIRINES, ORGANIC ACIDS

1081. Diazomalonic acid

[59348-62-8]

$C_3H_2N_2O_4$

$N_2C(CO.OH)_2$

1. 491M, 1975, 144
2. DOC 5, 1982, 1588


During attempted distillation of the impure acid at 4 mbar, the (involatile) sample exploded [1]. The free acid is unknown, but the diethyl ester has exploded during low pressure distillation [2].

See other DIAZO COMPOUNDS, ORGANIC ACIDS

1082. 2-Diazo-2H-imidazole

[50846-98-5]

$C_3H_2N_4$



As an isolated solid, the compound is shock-sensitive and should be freshly prepared only as a solution for immediate use.

See entry DIAZAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

1083. Cyanohyrazonoacetyl azide

[115057-44-8]

$C_3H_2N_6O$

$N_3CO.C(:NNH_2)CN$

Kobolov, M. Yu. *et al.*, *Chem. Abs.*, 1988, **109**, 20908


It explodes on melting.

See other ACYL AZIDES, CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

1084. Bis(1,2,3,4-thiaziazol-5-ylthio)methane

[]

$C_3H_2N_6S_4$



Pilgram, K. *et al.*, *Angew. Chem.*, 1965, **77**, 348

This compound (which is 'sulfur balanced'), and its three longer chain homologues, explodes loudly with a flash on impact, or on heating to the m.p.

See OXYGEN BALANCE See other N-S COMPOUNDS

†1085. Propiolaldehyde (Propynal)
[624-67-9]

C₃H₂O



Bases

1. Sauer, J. C., *Org. Synth.*, 1963, Coll. Vol. 4, 814
2. Makula, D. *et al.*, *L'Actual. Chim.*, 1983, (6), 31-34

The acetylenic aldehyde undergoes vigorous polymerisation in presence of alkalis and, with pyridine, the reaction is almost explosive [1]. The ground glass stopper of a 1 l brown bottle containing 750 ml of the aldehyde in toluene (2:1) and stored for 8 years was found to be seized. During attempts to open it an explosion occurred, leaving 250 ml of a polymeric residue in the bottle. DSC experiments and modeling calculations led to the conclusion that polymerisation had been initiated at ambient temperature by flakes of (alkaline) soda glass and peroxidic material (which had seized the stopper) falling into the solution of aldehyde. It is recommended that propynal should not be stored in glass, and only as a dilute (10%) solution [2] and for strictly limited periods.

See Acrylaldehyde

See other ACETYLENIC COMPOUNDS, ALDEHYDES, GLASS INCIDENTS, POLYMERISATION INCIDENTS

1086. Propiolic acid (Propynoic acid)
[471-25-0]

C₃H₂O₂



Ammonia, Heavy metal salts


Baudrowski, E., *Ber.*, 1882, **15**, 2701

Interaction of the acid with ammoniacal solutions of copper(I) or silver salts gives precipitates which explode on warming or impact. The structures are not given, but may be aminometal acetylide salts.

See METAL ACETYLIDES See other ACETYLENIC COMPOUNDS, ORGANIC ACIDS

†1087. 1,3-Dioxol-4-en-2-one (Vinylene carbonate)
[872-36-6]

C₃H₂O₃



Archer, M., private comm., 1979

A 7 year old screw capped sample burst in storage. Peroxide formation seems a less likely cause than slow hydrolysis and carbon dioxide evolution, though both are possibilities.

See related GAS EVOLUTION INCIDENTS, PEROXIDISABLE COMPOUNDS

1088. Silver 3-hydroxypropynide

[]

C_3H_3AgO



Karrer, P., *Organic Chemistry*, 111, London, Elsevier, 4th Engl. Edn., 1950

The silver salt is explosive.

See other METAL ACETYLIDES, SILVER COMPOUNDS

1089. Aluminium formate

[7360-53-4]

$C_3H_3AlO_6$



Anon., *ABCM Quart. Safety Summ.*, 1939, **10**, 1

An aqueous solution of aluminium formate was being evaporated over a low flame. When the surface crust was disturbed, an explosion occurred. This seems likely to have been due to thermal decomposition of the solid, liberation of carbon monoxide and ignition of the latter admixed with air.

†1090. 3-Bromopropyne (Propargyl bromide)

[106-96-7]

C_3H_3Br



HCS 1980, 784, (785 in toluene)

1. Coffee, R. D. *et al.*, *Loss Prev.*, 1967, **1**, 6–9
2. Driedger, P. E. *et al.*, *Chem. Eng. News*, 1972, **50**(12), 51
3. Brown, J. P., 1991, personal communication
4. Forshey, *Fire Technol.*, 1969, **5**, 100–111

This liquid acetylenic endothermic compound (ΔH_f° estimated as 230–270 kJ/mol, ~ 2 kJ/g) may be decomposed by mild shock, and when heated under confinement, it decomposes with explosive violence and may detonate. Addition of 20–30 wt% of toluene makes the bromide insensitive in laboratory impact and confinement tests [1]. More recently, it was classed as extremely shock-sensitive [2]. It can be ignited by impact derived from the ‘liquid-hammer’ effect of accidental pressurisation of the aerated liquid, and will then undergo sustained (monopropellant) burning decomposition. Propargyl bromide, added dropwise to bromine, exploded as it neared the halogen [3]. The chloro analogue is similar, but less readily ignited [4].

See other ENDOTHERMIC COMPOUNDS

Metals

Dangerous Substances, 1972, Sect. 1, 27

There is a danger of explosion in contact with copper, high-copper alloys, mercury or silver (arising from metal acetylide formation).


See METAL ACETYLIDES

Trichloronitromethane

See Trichloronitromethane: 3-Bromopropyne

See other HALOACETYLENE DERIVATIVES

1091. 2,4,6-Tris(bromoamino)-1,3,5-triazine (Tribromomelamine)
[22755-34-6]



Vona, J. A. *et al.*, *Chem. Eng. News*, 1952, **30**, 1916

Bromination with this and similar *N*-halogen compounds may become violent or explosive after an induction period as long as 15 min. Small scale preliminary experiments, designed to avoid the initial presence of excess brominating agent, are recommended.

See other INDUCTION PERIOD INCIDENTS


Allyl alcohol

Vona, J. A. *et al.*, *Chem. Eng. News*, 1952, **30**, 1916

The components reacted violently 15 min after mixing at ambient temperature. This seems likely to have been a radical-initiated polymerisation of the alcohol (possibly peroxidised) in absence of diluent.

See other *N*-HALOGEN COMPOUNDS

†1092. 3-Chloropropyne (Propargyl chloride)
[624-65-7]



HCS 1980, 786 (787 in toluene)

Doyle, W. H., *Loss Prev.*, 1969, **3**, 15

Pumping the liquid against a closed valve caused the pump to explode, which detonated the contents of the reservoir tank-car.

Ammonia

Anon., *Sichere Chemiearb.*, 1956, **8**(6), 45

Interaction of 3-chloropropyne and liquid ammonia under pressure in a static steel bomb had been used several times to prepare the amine. On one occasion the usual

slow exothermic reaction did not occur, and the bomb was shaken mechanically. The increased reaction rate led to a rapid exothermic reaction, followed by an explosion. Other cases of instability in propyne derivatives are known.

See 3-Propynol, and 3-Propynethiol

See other AMINATION INCIDENTS

Chlorine

See Chlorine: 3-Chloropropyne

See other HALOACETYLENE DERIVATIVES

†1093. Acryloyl chloride (2-Propenoyl chloride)

[814-68-6]

C_3H_3ClO



1. Pyriadi, T. M., *Chem. Eng. News*, 1985, **63**(44), 4
2. Griffith, T. E., *Chem. Eng. News*, 1985, **63**(50), 2
3. Author's comment, 1986


A 500 ml bottle of the acid chloride stabilised with 0.05% of phenothiazine was shipped to a hot climate without refrigeration, and was stored on arrival in a fume cupboard for 2 days at temperatures approaching 50°C. The material polymerised, bursting the bottle and forming a solid foam [1]. The label recommended storage at 4°C. Another manufacturer proposed mislabeling or contamination by moisture to explain the incident [2], but the presence of polymeric foam appears to confirm that polymerisation at the extremely high ambient storage temperature was the primary cause, perhaps with some thermal decomposition and evolution of hydrogen chloride assisting in foam formation [3].

See other ACYL HALIDES, POLYMERISATION INCIDENTS

1094. 1,3-Dithiolium perchlorate

[3706-77-2]

$C_3H_3ClO_4S_2$




1. Ferraris, J. P. *et al.*, *Chem. Eng. News*, 1974, **52**(37), 3
2. Klingsberg, E., *J. Amer. Chem. Soc.*, 1964, **86**, 5292
3. Leaver, D. *et al.*, *J. Chem. Soc.*, 1962, 5109
4. Wudl, F. *et al.*, *J. Org. Chem.*, 1974, **39**, 3608–3609; *Inorg. Chem.*, 1979, **19**, 28–29
5. Melby, L. R. *et al.*, *J. Org. Chem.*, 1974, **39**, 2456

The salt (an intermediate in the preparation of 1,4,5,8-tetrahydro-1,4,5,8-tetrathiafulvalene) exploded violently during removal from a glass frit with a Teflon-clad spatula [1]. Previous references to the salt exploding at 250°C [2] or melting at

264°C [3] had been made. Use of a salt alternative to the perchlorate is urged [1]. Safer methods suitable for small-scale [4] and large-scale [5] preparations have been described.

See other NON-METAL PERCHLORATES

†1095. **2,4,6-Tris(chloroamino)-1,3,5-triazine (Trichloromelamine)**
[7673-09-8]



Acetone, or Bases

491M, 1975, 197

Addition of acetone, ammonia, aniline or diphenylamine to the oxidant causes, after a few seconds delay, a rapid reaction accompanied by smoke and flame.

See 2,4,6-Tris(bromoamino)-1,3,5-triazine

See other N-HALOGEN COMPOUNDS

1096. **Methyl trichloroacetate**
[598-99-2]



Trimethylamine

Anon., *Angew. Chem. (Nachr.)*, 1962, **10**, 197

A stirred uncooled mixture in an autoclave reacted violently, the pressure developed exceeding 400 bar. Polymerisation of a reactive species produced by dehydrochlorination of the ester or a decomposition product seems a possibility.

See Sodium hydride: Ethyl 2,2,3-trifluoropropionate

See other POLYMERISATION INCIDENTS

1097. **Trichloromethyl peroxyacetate**
[94089-34-6]




Turner, W. V. *et al.*, *J. Org. Chem.*, 1992, **57**(5), 1610

This compound has caused several explosions in the author's laboratory.

See other PEROXYESTERS



1098. **1-Propynylcopper(I)**
[30645-13-7]



Houben-Weyl, 1977, Vol. 5.2a, 570

Preparation and use of this very explosive compound are detailed.
See other METAL ACETYLIDES

†1099. **3,3,3-Trifluoropropene**
[677-21-4]



See other HALOALKENES

†1100. **1,1,1-Trifluoroacetone**
[421-50-1]



1101. **Methyl trifluorovinyl ether**
[3823-94-7]




1. Dixon, S., US Pat. 2 917 548, 1959
2. Anderson, A. W., *Chem. Eng. News*, 1976, **54**(16), 5
3. O'Neill, G. J., US Pat. 4 127 613, 1978

Methyl trifluorovinyl ether, b.p. 10.5–12.5°C, prepared from tetrafluoroethylene and sodium methoxide [1], has considerable explosive potential. On ignition, it decomposes more violently than acetylene and should be treated with extreme caution [2]. Other trifluorovinyl ethers are similarly available from higher alkoxides [1], and although not tested for instability, should be handled carefully. Presence of fluoro-haloalkanes boiling lower than the ether stabilises the latter against spark-initiated decomposition in both fluid phases [3].

See related HALOALKENES

1102. **Methyltrifluoromethyldioxirane**
[115464-59-0]



See Dimethyldioxirane, Bis(trifluoromethyl)dioxirane
See other CYCLIC PEROXIDES

1103. **Trifluoromethyl peroxyacetate**
[33017-08-2]



Bernstein, P. A. *et al.*, *J. Amer. Chem. Soc.*, 1971, **93**, 3885

A 1 g sample cooled to -196°C exploded violently when warmed in a bath at 22°C .
See other PEROXYESTERS

1104. 1,1-Bis(difluoroamino)-2,2-difluoro-2-nitroethyl methyl ether
[30957-47-2]

$\text{C}_3\text{H}_3\text{F}_6\text{N}_3\text{O}_3$



Ross, D. L. *et al.*, *J. Org. Chem.*, 1970, **35**, 3096–3098
A shock-sensitive explosive.
See other DIFLUOROAMINO COMPOUNDS

1105. 3-Iodopropyne
[659-86-9]


$\text{C}_3\text{H}_3\text{I}$



Whiting, M. C., *Chem. Eng. News*, 1972, **50**(23), 86–87
It explodes during distillation at 180°C .
See other HALOACETYLENE DERIVATIVES

1106. Potassium 1-tetrazolacetate
[51286-83-0]


$\text{C}_3\text{H}_3\text{KN}_4\text{O}_2$



Eizember, R. F. *et al.*, *J. Org. Chem.*, 1974, **39**, 1792–1793
During oven-drying, kg quantities of the salt exploded violently. Investigation showed that self-propagating and extremely rapid decomposition of a cold sample can be initiated by local heating to over 200°C by a flint spark, prolonged static spark or flame. The sodium salt could only be initiated by flame, and the free acid is much less sensitive.
See other TETRAZOLES

†**1107. Acrylonitrile (Propenenitrile)**
[107-13-1]

$\text{C}_3\text{H}_3\text{N}$



(*MCA SD-31*, 1964); *FPA H51*, 1976; *HCS 1980*, 116; *RSC Lab. Hazards Data Sheet No. 48*, 1986

1. *MCA SD-31*, 1964
2. Harmon, 1974, 2.3
3. Taylor, B. J., *Univ. Safety Assoc. Safety News*, 1972, (1), 5–6
4. Bond, J., *Loss Prev. Bull.*, 1985, (065), 26

C₃H₃N

The monomer is sensitive to light, and even when inhibited (with aqueous ammonia) it will polymerise exothermally at above 200°C [1]. It must never be stored uninhibited, or adjacent to acids or bases [2]. Polymerisation of the monomer in a sealed tube in an oil bath at 110°C led to a violent explosion. It was calculated that the critical condition for runaway thermal explosion was exceeded by a factor of 15 [3]. Runaway polymerisation in a distillation column led to an explosion and fire [4].

See other POLYMERISATION INCIDENTS

Acids MRH Nitric acid 6.32/78, sulfuric acid 2.26/48

1. 'Acrylonitrile', London, British Hydrocarbon Chemicals Ltd., 1965
2. Kaszuba, F. J., *J. Amer. Chem. Soc.*, 1945, **67**, 1227
3. Shirley, D. A., *Preparation of Organic Intermediates*, 3, New York, Wiley, 1951
4. Kaszuba, F. J., *Chem. Eng. News*, 1952, **30**, 824
5. Anon., *Loss Prev. Bull.*, 1978, (024), 167
6. Bond, J., *Loss Prev. Bull.*, 1985, (065), 26

Contact of strong acids (nitric or sulfuric) with acrylonitrile may lead to vigorous reactions. Even small amounts of acid are potentially dangerous, as these may neutralise the aqueous ammonia present as polymerisation inhibitor and leave the nitrile unstabilised [1]. Precautions necessary in the hydrolysis of acrylonitrile [2] are omitted in the later version of the procedure [3]. It is essential to use well-chilled materials (acrylonitrile, diluted sulfuric acid, hydroquinone, copper powder) to avoid eruption and carbonisation. A really wide-bore condenser is necessary to contain the vigorous boiling of unhydrolysed nitrile [4]. Leaking valves or operator error caused contamination of a 1000 l dosing vessel of the nitrile with conc. sulfuric acid, and an explosion ensued [5]. Contamination of a drum of acrylonitrile by nitric acid residues in the filling hose led to a slowly accelerating polymerisation reaction which burst the drum after 10 days [6].

See Bromine, etc., below

Bases MRH Potassium or sodium hydroxides 2.84/tr.

1. *MCA SD-31*, 1964
2. Castaneda Hernandez, H. R. *et al.*, *Chem. Abs.*, 1987, **107**, 78274

Acrylonitrile polymerises violently in contact with strong bases, whether stabilised or unstabilised [1]. Alkaline hydrolysis of acrylonitrile is exothermic and violent, especially when the temperature is above 60°C, the pressure is above atmospheric, and when heating at 60°C is prolonged above 10 mins. Polymerisation does not induce a violent reaction at 40–50°C at a concentration of 3% of sodium hydroxide in water [2].

Benzyltrimethylammonium hydroxide, Pyrrole

MCA Guide, 1972, 299

To catalyse the cyanoethylation of pyrrole, 3 drops of the basic catalyst solution were added to the reaction mixture of pyrrole (30%) in the nitrile. An exotherm developed and base-catalysed polymerisation of the nitrile accelerated to explosion.

See Bases, above, *also* Tetrahydrocarbazole, etc., below

See other POLYMERISATION INCIDENTS

Formaldehyde

Personal experience

The reaction between acrylonitrile and formaldehyde (as paraformaldehyde or trioxane), under strong acid catalysis (usually sulphuric) and most often in presence of catalytic quantities of acetic anhydride, to produce triacrylohexahydrotriazine, is inclined to violent exotherm after an induction period. The runaway can be uncontrollable on sub-molar scale. It may be due to acrylate polymerisation or to increasing reactivity of the formaldehyde equivalent due to progressive de-oligomerisation. Procedures claimed to prevent the risk have been described in the literature but do not seem reliable.

Halogens

MRH 2.84/tr. Bromine

1. *MCA Case History No. 1214*

2. Towell, G., 1989, Personal communication

Bromine was being added in portions to acrylonitrile with ice cooling, with intermediate warming to 20°C between portions. After half the bromine was added, the temperature increased to 70°C; then the flask exploded. This was attributed either to an accumulation of unreacted bromine (which would be obvious) or to violent polymerisation [1]. The latter seems more likely, catalysed by hydrogen bromide formed by substitutive bromination. Chlorine produces similar phenomena, even if the flask stays intact. The runaway is preceded by loss of yellow colouration and accompanied by formation of 3-chloroacrylonitrile and derivatives. It can be suppressed by presence of bases [2].

See Acids, above

Initiators

1. Zhulin, V. M. *et al.*, *Dokl. Akad. Nauk SSSR*, 1966, **170**, 1360–1363

2. Biesenberger, J. A. *et al.*, *Polymer Eng. Sci.*, 1976, **16**, 101–116

At pressures above 6000 bar, free radical polymerisation sometimes proceeded explosively [1]. The parameters were determined in a batch reactor for thermal runaway polymerisation of acrylonitrile initiated by azoisobutyronitrile, dibenzoyl peroxide or di-*tert*-butyl peroxide [2].

See other POLYMERISATION INCIDENTS

See VIOLENTPOLYMERISATION

Other reactants

Yoshida, 1980, 11

MRH values calculated for 17 combinations, mainly with oxidants, are given

Silver nitrate

Anon., *ABCM Quart. Safety Summ.*, 1962, **33**, 24

Acrylonitrile containing undissolved solid silver nitrate is liable, on long standing, to polymerise explosively and ignite. This was attributed to the slow deposition of a thermally insulating layer of polymer on the solid nitrate, which gradually gets hotter and catalyses rapid polymerisation. Photocatalysed decomposition of the salt with formation of traces of nitric acid may have been involved in the initial stages.

See other POLYMERISATION INCIDENTS

Tetrahydrocarbazole, Benzyltrimethylammonium hydroxide

Anon., *BCISC Quart. Safety Summ.*, 1968, **39**, 36

Cyanoethylation of 1,2,3,4-tetrahydrocarbazole initiated by the quaternary base had been effected smoothly on twice a published scale of working. During a further fourfold increase in scale, the initiator was added at 0°C, and shortly after cooling had been stopped and heating begun, the mixture exploded. A smaller proportion of initiator and very slow warming to effect reaction are recommended (to avoid rapid polymerisation of the nitrile by the base).

See Benzyltrimethylammonium hydroxide, etc., and Bases, both above

See other CYANO COMPOUNDS

See related HALOALKENES

1108. Poly(acrylonitrile)

[25014-41-9]

(C₃H₃N)_n



Methacrylate polymer, Nitric acid

See Nitric acid: Acrylonitrile–methacrylate copolymer

1109. Vinyl isocyanide (Isocynoethene)

[14668-82-7]

C₃H₃N



1. Matteson, D. S. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 3765

2. Bailey, R., *Chem. Abs.*, 1968, **69**, 113029

The molar heat of formation of this endothermic compound (+230–250 kJ, 4.5 kJ/g) is comparable with that of buten-3-yne (vinylacetylene). While no explosive decomposition of the isocyanide has been reported, the possibility should be borne in mind [1]. It is stable at –15°C, but isomerises to acrylonitrile and polymerises at ambient temperature [2].


See related CYANO COMPOUNDS

See other ENDOTHERMIC COMPOUNDS

†**1110. Isoxazole**

[288-14-2]

C₃H₃NO




Sealed samples decompose exothermally above 136°C.

See entry ISOXAZOLES

See other N–O COMPOUNDS

†1111. Oxazole
[288-42-6]


C₃H₃NO



See other N-O COMPOUNDS

1112. 2-Thioxo-4-thiazolidinone ('Rhodanine')
[141-84-4]

C₃H₃NOS₂



Merck, 1983, 1180
It may explode on rapid heating.
See other N-S COMPOUNDS

1113. Cyanoacetic acid
[372-09-8]

C₃H₃NO₂



Furfuryl alcohol
See Furfuryl alcohol: Acids
See other CYANO COMPOUNDS, ORGANIC ACIDS


1114. 3-Azidopropyne (Propargyl azide)
[14989-89-0]

C₃H₃N₃



Almlof, J. *et al.*, *J. Mol. Struct.*, 1987, **160**, 1
It is extremely explosive and must be handled with utmost care. Explosions during synthesis (100 mmol, 8.1 g) wrecked a ventilated hood. (The energy of decomposition of this endothermic acetylenic azide probably exceeds 250 kJ/mol, 3 kJ/g.)
See other ACETYLENIC COMPOUNDS, ENDOTHERMIC COMPOUNDS, ORGANIC AZIDES

1115. 1,3,5-Triazine
[290-87-9]



Nitric acid, Trifluoroacetic anhydride

See Nitric acid: Triazine, etc.

See other ORGANIC BASES

†**1116. 2-Carbamoyl-2-nitroacetonitrile ('Fulminuric acid')**
[475-08-1]




Sorbe, 1968, 74

It explodes on heating.

See other CYANO COMPOUNDS, C-NITRO COMPOUNDS

1117. 2-Amino-5-nitrothiazole
[121-66-4]




CHETAH, 1990, 183

Can be exploded with a detonator, not by mechanical shock.

Preparative hazard

See Nitric acid: 2-Aminothiazole, Sulfuric acid


†**1118. 2,4,6-Trihydroxy-1,3,5-triazine (Cyanuric acid)**
[108-80-5]



Chlorine


See Chlorine: Nitrogen compounds

1119. Sodium 1-tetrazolacetate
[51286-84-1]




See Potassium 1-tetrazolacetate
See other TETRAZOLES

1120. 5-Cyano-2-methyltetrazole
[91511-39-6]




Aluminium hydride
See Aluminium hydride: Tetrazole derivatives
See other TETRAZOLES

1121. 2,3-Diazidopropionitrile
[101944-90-5]



Owston, P. G. *et al.*, *J. Chem. Res.; Synop.*, 1985, (11), 352–353
It is unstable and potentially explosive: routine use of a fume cupboard and safety screens is recommended.
See other CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1122. 2-Hydroxy-4,6-bis(nitroamino)-1,3,5-triazine
[19899-80-0]



Atkinson, J. R., *J. Amer. Chem. Soc.*, 1951, **73**, 4443–4444
The explosive nitration product of ‘melamine’ was identified as the title compound of low oxygen balance, which is easily detonated on impact.
See other N-NITRO COMPOUNDS

1123. Sodium methoxyacetylide

[]



Houben-Weyl, 1970, Vol. 13.1, 649

It may ignite in air.


Brine

Jones, E. R. H. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 406

During addition of saturated brine at -20°C to the sodium derivative at -70°C , minor explosions occur. These may have been caused by particles of sodium igniting the liberated methoxyacetylene.

See other METAL ACETYLIDES**†1124. Propadiene (Allene)**

[463-49-0]

*HCS 1980*, 779 (cylinder)1. Bondor, A. M. *et al.*, *Khim. Prom.*, 1965, **41**, 9232. Forshey, D. R., *Fire Technol.*, 1969, **5**, 100–111

The diene is endothermic (ΔH_f° (g) +192.1 kJ/mol, 4.80 kJ/g) and the pure gas can decompose explosively under a pressure of 2 bar [1], but this is also quoted as the upper limiting pressure for flame propagation during sustained (monopropellant) burning at 25°C [2].

See Propyne (below)

Oxides of nitrogen

See Nitrogen oxide: Dienes, Oxygen*See other* DIENES, ENDOTHERMIC COMPOUNDS**†1125. Propyne**

[74-99-7]

*HCS 1980*, 633 (cylinder)1. *MCA Case History No. 632*2. Fitzgerald, F., *Nature*, 1960, **186**, 386–3873. Hurden, D., *J. Inst. Fuel.*, 1963, **36**, 50–54

4. Stull, 1977, 16

The liquid material in cylinders (which contains around 30% of propadiene) is not shock-sensitive, but a wall temperature of 95°C (even very localised) accompanied by pressures of about 3.5 bar, will cause a detonation to propagate from the hot spot [1]. Induced decomposition of the endothermic hydrocarbon leads to flame

propagation in absence of air above minimum pressures of 3.4 and 2.1 bar at 20 and 120°C, respectively [2]. Application as a monopropellant and possible hazards therefrom (including formation of explosive copper propynide) have been discussed [3]. Although the pure material is highly endothermic (ΔH_f° (g) +185.4 kJ/mol, 4.64 kJ/g), the commercial mixture with propadiene and propane (MAPP gas) is comparable with ethylene for handling requirements and potential hazard [4].

Silver nitrate

Grignard, 1935, Vol. 3, 175

Interaction in ammoniacal solution gives the silver derivative which ignites around 150°C.

See other ALKYNES, ENDOTHERMIC COMPOUNDS

1126. 3-Bromo-1,1,1-trichloropropane

[13749-37-6]



Preparative hazard

See Ethylene: Bromotrichloromethane

See other HALOALKANES

†1127. 1-Chloro-3,3,3-trifluoropropane

[460-35-5]



See other HALOALKANES

1128. 2-Chloro-*N,N,N'*-trifluoropropionamidine

[25238-02-2]



Ross, D. L. *et al.*, *J. Org. Chem.*, 1970, **35**, 3096–3097

A shock-sensitive explosive.

See other *N,N,N'*-TRIFLUOROAMIDINES

1129. *N*-Chloro-3-aminopropyne (*N*-Chloropropargylamine)

[103698-31-3]



It explodes at ambient temperature.

See entry *N*-HALOGEN COMPOUNDS (reference 4)

See other ACETYLENIC COMPOUNDS, *N*-HALOGEN COMPOUNDS

1130. 2-Chloro-1-cyanoethanol (3-Chlorolactonitrile)

[33965-80-9]



Scotti, F. *et al.*, *J. Org. Chem.*, 1964, **29**, 1800

Distillation (at 110°C/4 mbar) is hazardous, since slight overheating may cause explosive decomposition to 2-chloroacetaldehyde and hydrogen cyanide.

See other CYANO COMPOUNDS

1131. N-(2-Chloroethyl)-N-nitrosocarbamoyl azide
[60784-40-9]

C₃H₄ClN₅O₂



Eisenbrand, G., Ger. Offen. 2 659 862, 1978

It should not be isoated from solution, as it is potentially explosive.

See other ACYL AZIDES, NITROSO COMPOUNDS

†1132. **2,3-Dichloropropene**
[78-88-6]


C₃H₄Cl₂



See other HALOALKENES

1133. 2,2,3,3-Tetrafluoropropanol
[76-37-9]

C₃H₄F₄O



Potassium hydroxide, or Sodium


Bagnall, R. D., private comm., 1972

Attempted formation of sodium tetrafluoropropoxide by adding the alcohol to sodium (40 g) caused ignition and a fierce fire which melted the flask. This was attributed to alkoxide-induced elimination of hydrogen fluoride, and subsequent exothermic polymerisation. In an alternative preparation of the potassium alkoxide by adding the alcohol to solid potassium hydroxide, a vigorous exotherm occurred. This was not seen when the base was added slowly to the alcohol.

See other POLYMERISATION INCIDENTS

1134. 1-Hydroxyimidazole N-oxide
[35321-46-1]

C₃H₄N₂O₂



Flynn, A. P., *Chem. Brit.*, 1984, **20**, 30

On DSC examination it exhibits a large decomposition exotherm at 200–250°C, and is a sensitive detonating and deflagrating explosive.

See 1-Methoxyimidazole *N*-oxide

See other N–O COMPOUNDS, *N*-OXIDES

†1135. 3-Diazopropene (Vinyl diazomethane)

[2032-04-4]

C₃H₄N₂



Salomon, R. G. *et al.*, *J. Org. Chem.*, 1975, **40**, 758


Potentially explosive, it should be stored in solution at 0°C and shielded from light.

See other DIAZO COMPOUNDS

1136. 3-Aminoisoxazole

[1750-42-1]

C₃H₄N₂O




Sealed samples decompose exothermally above 115°C.

See entry ISOXAZOLES See other N–O COMPOUNDS

1137. 4-Amino-3-isoxazolidinone

[68-39-3]

C₃H₄N₂O₂



Sealed samples decompose exothermally above 78°C.

See entry ISOXAZOLES See other N–O COMPOUNDS

1138. Methyl diazoacetate

[6832-16-2]

C₃H₄N₂O₂




Searle, N. E., *Org. Synth.*, 1963, Coll. Vol. 4, 426

This ester must be handled with particular caution as it explodes with extreme violence on heating.


See other DIAZO COMPOUNDS

1139. 3-Nitro-2-isoxazoline
[1121-14-8]



CHETAH, 1990, 188
Shock sensitive
See other C-NITRO COMPOUNDS, N-O COMPOUNDS

1140. 2-Aminothiazole
[96-50-4]




MCA Case History No. 1587

Drying 2-aminothiazole in an oven without forced air circulation caused development of hot spots and eventual ignition. It has a low auto-ignition temperature and will ignite after 3.5 h at 100°C.

Nitric acid
See Nitric acid: 2-Aminothiazole

Nitric acid, Sulfuric acid
See Nitric acid: 2-Aminothiazole, Sulfuric acid

†1141. Imidazoline-2,4-dithione ('Dithiohydantoin')
[5789-17-3]




Pouwels, H., *Chem. Eng. News*, 1975, **53**(49), 5

A 70 g sample, sealed into a brown glass ampoule, exploded after storage at ambient temperature for 17 years. This was attributed to slow decomposition and gas generation (perhaps initiated by traces of alkali in the ampoule glass).

See Pyrimidine-2,4,5,6-(1*H*,3*H*)-tetraone
See other GLASS INCIDENTS

1142. 1,3-Dinitro-2-imidazolidinone
[2536-18-7]

C₃H₄N₄O₅



Violent decomposition occurred at 238°C.
See entry DIFFERENTIAL THERMAL ANALYSIS(DTA) (reference 1) See other *N*-NITRO COMPOUNDS

1143. 1,3-Diazidopropene
[22750-69-2]


C₃H₄N₆



Forster, M. O. *et al.*, *J. Chem. Soc.*, 1912, **101**, 489
A sample exploded while being weighed.
See other ORGANIC AZIDES

1144. Ammonium 2,4,5-trinitroimidazolid
[63839-60-1]

C₃H₄N₆O₆



Coburn, M. D. US Pat. 4 028 154, 1977
The ammonium salt is an explosive comparable to RDX but of higher thermal stability.
See related POLYNITROARYL COMPOUNDS

†**1145. Acrylaldehyde (Propenal)**
[107-02-8]

C₃H₄O



(MCA SD-85, 1961); NSC 436, 1978; NFPA H49, 1976; HCS 1980, 113; RSC Lab. Hazards Data Sheet No. 70, 1988

Energy of decomposition (in range 70–380°C) measured as 0.864 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 117°C by adiabatic Dewar tests, with an apparent energy of activation of 214 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Acids, or Bases

1. MCA SD-85, 1961
2. Hearsey, C. J., private comm., 1973
3. Personal experience (PGU)
4. Catalogue note, Hopkin & Williams, 1973
5. Bond, J., *Loss Prev. Bull.*, 1985, (065), 26

Acrylaldehyde (acrolein) is very reactive and will polymerise rapidly, accelerating to violence, in contact with strong acid or basic catalysts. Normally an induction period, shortened by increase in contamination, water content or initial temperature, precedes the onset of polymerisation. Uncatalysed polymerisation sets in at 200°C in the pure material [1]. Exposure to weakly acidic conditions (nitrous fumes, sulfur dioxide, carbon dioxide), some hydrolysable salts, or thiourea will also cause exothermic and violent polymerisation. A 2 year old sample stored in a refrigerator close to a bottle of dimethylamine exploded violently, presumably after absorbing enough volatile amine (which penetrates plastics closures) to initiate polymerisation [2]. Measurement of acrolein in a plastic measuring cylinder previously used for triethylamine (and allowed to evaporate dry) gave rise to violent polymerisation and a lacquered fume cupboard within less than a minute [3]. The stabilising effect of the added hydroquinone may cease after a comparatively short storage time. Such unstabilised material could polymerise explosively [4]. Violent polymerisation in a 250 kl storage tank led to an explosion and widespread damage [5].

See other ALDEHYDES, CATALYTIC IMPURITY INCIDENTS, INDUCTION PERIOD INCIDENTS, PEROXIDISABLE COMPOUNDS, POLYMERISATION INCIDENTS

Trienes

Anon., 1989, Personal communication

Calorimetric investigation of a Diels Alder reaction between propenal and a triene, which had caused problems on scale-up, showed, after the exotherm due to the Diels Alder reaction, and from a temperature a little above 200°C, a second, more exothermic, reaction with a very fast pressure rise which burst the ARC can employed. This is presumably aromatisation of the alkenylcyclohexenealdehyde first formed, with probable liberation of hydrogen and carbon monoxide.

See ACCELERATING RATE CALORIMETRY

†1146. Methoxyacetylene (Ethyne methyl ether)

[6443-91-0]

C_3H_4O



See Sodium methoxyacetylide: Brine

See other ACETYLENIC COMPOUNDS

†1147. 3-Propynol (Propargyl alcohol)
[107-19-7]

C₃H₄O



Alkalies

Anon., *Angew. Chem. (Nachr.)*, 1954, **2**, 209

If propynol and similar acetylenic compounds are dried with alkali before distillation, the residue may explode (probably owing to acetylenic salt formation). Sodium sulfate is recommended as a suitable desiccant.

Mercury(II) sulfate, Sulfuric acid, Water

1. Reppe, W. *et al.*, *Ann.*, 1955, **596**, 38
2. Nettleton, J., private comm., 1972

Following the published procedure [1], hydroxyacetone was being prepared on half the scale by treating propargyl alcohol as a 30% aqueous solution with mercury sulfate and sulfuric acid (6 g and 0.6 g/mol of alcohol, respectively). On stirring and warming the mixture to 70°C a violent exothermic eruption occurred. Quartering the scale of operations to 1 g mol and reducing the amount of acid to 0.37 g/mol gave a controllable reaction at 70°C. Adding the alcohol to the other reactants at 70°C is an alternative possibility to avoid the suspected protonation and polymerisation of the propargyl alcohol.

See other POLYMERISATION INCIDENTS

Phosphorus pentoxide

491M, 1975, 321

Addition of the (acidic) oxide to the alcohol caused ignition.

Sulfuric acid

1. Sauer, J. C., *Org. Synth.*, 1963, Coll. Vol. 4, 813
2. Crouse, D. M., *Chem. Eng. News*, 1979, **57**(41), 4
3. Hart, E. V., *Chem. Eng. News*, 1980, **58**(2), 4

During addition of 56% aqueous acid to the aqueous 33% alcohol, using a published procedure for conversion to propionaldehyde [1], a violent explosion was attributed to lack of adequate cooling [2]. This attribution is disputed, and contamination, possibly with a heavy metal ion, is postulated [3]. The published procedure is rated not hazardous.

See Mercury(II) sulfate, etc., above

See other ACETYLENIC COMPOUNDS

†1148. Acrylic acid (Propenoic acid)
[79-10-7]

C₃H₄O₂



FPA H95, 1980; *HCS 1980*, 115; *RSC Lab. Hazard Data Sheet No. 68*, 1988

1. *Haz Chem. Data*, 1975, 34
2. Anon., *RoSPA OS & H Bull.*, 1976, (10), 3

3. Anaorian, M. *et al.*, *Eur. Polym. J.*, 1981, **17**, 823–838
4. Bond, J., *Loss Prev. Bull.*, 1985, (065), 26
5. Kurland, J. J. *et al.*, *Plant/Oper. Progr.*, 1987, **6**, 203–207
6. Yassin, A. A. *et al.*, *Chem. & Ind.*, 1985, 275
7. Levy, L. B., *Plant/Oper. Progr.*, 1987, **6**, 188–189; *ibid.*, 1989, **8**(2), 105
8. Wamper, F. M., *Plant/Oper. Progr.*, 1988, **7**, 183–189

Acrylic acid is normally supplied as the inhibited monomer, but because of its relatively high freezing point (14°C) it often partly solidifies, and the solid phase (and the vapour) will then be free of inhibitor which remains in the liquid phase. Even the uninhibited acid may be stored safely below the m.p., but such material will polymerise exothermically at ambient temperature, and may accelerate to a violent or explosive state if confined. Narrow vents may become blocked by polymerisation of uninhibited vapour [1]. A 17 m³ tank trailer of glacial acrylic acid was being warmed by internal coils containing water from an unmonitored steam and water mixer to prevent the acid freezing in sub-zero temperatures. The extremely violent explosion which occurred later probably involved explosive polymerisation accelerated by both the latent heat liberated when the acid froze, and the uncontrolled deliberate heating [2]. The detailed mechanism of auto-accelerating polymerisation of the acid in hexane–methanol solution, which becomes explosive, has been studied [3]. Contamination of a 220 kl cargo of acrylic acid from an adjacent storage tank in a marine tanker led to an escalating polymerisation which could not be controlled, and the cargo could not be off-loaded. Eventually the tank had to be cut out of the ship [4]. It appears that the contaminant leaking into the acrylic acid was ethylenenorbornene, which reacted with the traces of oxygen necessary to activate the inhibitor in the acrylic acid, leading to onset of polymerisation [5]. Hexamethylenetetramine is an effective stabiliser for acrylic acid in both liquid and vapour phases, preventing polymerisation at concentrations of 50–100 ppm [6]. Dimerisation of acrylic acid is temperature and solvent dependent and cannot be controlled by inhibitors, but vinyl polymerisation can be inhibited by conventional radical-trapping compounds, such as hydroquinone or phenothiazine, both of which need access of some oxygen for activation [7]. The dimerisation process of moist acrylic acid has been further investigated, and an empirical equation permits prediction of the extent of dimerisation in storage at different temperatures [8].

See other POLYMERISATION INCIDENTS

Initiator, Water

Anon., *Sichere Chemiearb.*, 1978, **30**, 3

During the experimental large scale continuous polymerisation of acrylic acid in aqueous solution in presence of an initiator and a moderator, failure of one of the feed pumps led to an unusually high concentration of monomer in solution. This led to runaway polymerisation which burst a glass vent line and the escaping contents ignited and led to an explosion and fire.

See other GLASS INCIDENTS, POLYMERISATION INCIDENTS

Other reactants

Yoshida, 1980, 8

MRH values calculated for 13 combinations with oxidants are given.
See VIOLENT POLYMERISATION *See other* ORGANIC ACIDS

†1149. Vinyl formate (Ethenyl methanoate)
[692-45-5]



$\text{C}_3\text{H}_4\text{O}_2$

1150. Pyruvic acid (2-Oxopropanoic acid)
[127-17-3]



$\text{C}_3\text{H}_4\text{O}_3$

1. Anon., *Sichere Chemiearb.*, 1977, **29**, 87
2. Wolff, L., *Ann.*, 1901, **317**, 2

A bottle of analytical grade material exploded in laboratory storage at 25°C (undoubtedly from internal pressure of carbon dioxide). Pure material, protected from light and air, is only stable on a long term basis if kept refrigerated. Otherwise slow decomposition and decarboxylation occurs [1], possibly accelerated by enzymic catalysis from ingress of airborne yeasts. At ambient temperature, the acid dimerises and dehydrates to 2-oxo-4-carboxyvalerolactone [2].

See other GAS EVOLUTION INCIDENTS, ORGANIC ACIDS

1151. Malonic acid (Propanedioic acid)
[141-82-2]



$\text{C}_3\text{H}_4\text{O}_4$

Although undoubtedly capable of generating carbon dioxide by thermal decomposition, the editor has heard of no pressurisation problems arising from this.

Potassium bromate, Ceric ammonium nitrate, Water

See Potassium bromate: Ceric ammonium nitrate, Malonic acid, Water

1152. 3-Propynethiol
[27846-30-6]



$\text{C}_3\text{H}_4\text{S}$

1. Sato, K. *et al.*, *Chem. Abs.*, 1956, **53**, 5112b
2. Brandsma, 1971, 179

When distilled at ambient pressure, it polymerised explosively. It distils smoothly under reduced pressure at 33–35°C/127 mbar [1]. The polymer produced on exposure to air may explode on heating. Presence of a stabiliser is essential during handling or storage under nitrogen at –20°C [2].

See other ACETYLENIC COMPOUNDS, POLYMERISATION INCIDENTS

- †1153. **3-Bromo-1-propene (Allyl bromide)**
 [106-95-6] **C₃H₅Br**

$$\text{H}_2\text{C}=\text{CHCH}_2\text{Br}$$

HCS 1980, 124
See other ALLYL COMPOUNDS, HALOALKENES
- 1154. Propionyl hypobromite**
 [82198-80-9] **C₃H₅BrO₂**

$$\text{EtCO}\cdot\text{OBr}$$

 Skell, P. S. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 4000
 As an isolated solid it is unpredictably explosive.
See other ACYL HYPOHALITES
- 1155. Allylzinc bromide (Bromo-2-propenylzinc)**
 [18925-10-5] **C₃H₅BrZn**

$$\text{CH}_2=\text{CHCH}_2\text{ZnBr}$$

 Formamide, Ketones
See Formamide: Organozinc compounds
See also GRIGNARD REAGENTS
- †1156. **1-Chloro-1-propene**
 [590-21-6] **C₃H₅Cl**

$$\text{ClCH}=\text{CHMe}$$

See other HALOALKENES
- †1157. **2-Chloropropene**
 [557-98-2] **C₃H₅Cl**

$$\text{H}_2\text{C}=\text{CHClMe}$$

See other HALOALKENES
- †1158. **3-Chloropropene (Allyl chloride)**
 [107-05-1] **C₃H₅Cl**

$$\text{H}_2\text{C}=\text{CHCH}_2\text{Cl}$$

 (MCA SD-99, 1973); *HCS 1980*, 125; *RSC Lab. Hazards Data Sheet No. 28*, 1984
 Piccinini, N. *et al.*, *Plant/Operations Progr.*, 1982, **1**, 69–74
 An operability analysis to identify potential risks in an allyl chloride manufacturing plant has been published.

Aromatic hydrocarbons, Ethylaluminium chlorides

491M, 1975, 22

Friedel-Crafts alkylation of benzene or toluene by allyl chloride in presence of ethylaluminium chlorides is vigorous even at -70°C , and explosions have occurred.

See Lewis acids, etc., next below

Lewis acids, Metals

MCA SD-99, 1973

Contact with aluminium chloride, boron trifluoride, sulfuric acid etc., may cause violently exothermic polymerisation. Organometallic products of contact of the chloride with aluminium, magnesium, zinc (or galvanised metal) may produce similar results.

See other ALLYL COMPOUNDS, HALOALKENES

1159. Ethyl *N*-chloro-*N*-sodiocarbamate

[17510-52-0]

$\text{C}_3\text{H}_5\text{ClNaO}_2$

$\text{ClN}(\text{Na})\text{CO.OEt}$

1. Herranz, E. *et al.*, *J. Amer. Chem. Soc.*, 1978, **100**, 3598, footnote 8

2. Scully, F. E., *Tetrahedron Lett.*, 1990, **31**(37), 5261

It may decompose (non-explosively) exothermally with gas evolution, and the scale of preparation should be limited to 0.2 g mol [1]. Samples drying in desiccators several times shattered them. Use in solution is recommended [2].

See other *N*-HALOGEN COMPOUNDS

1160. 1-Chloro-2,3-propylene dinitrate

[2612-33-1]

$\text{C}_3\text{H}_5\text{ClN}_2\text{O}_6$

$\text{ClCH}_2\text{CH}(\text{ONO}_2)\text{CH}_2\text{ONO}_2$

Sorbe, 1968, 54

An explosive syrup.

See related ALKYL NITRATES

†1161. Chloroacetone (Chloro-2-propanone)

[78-95-5]

$\text{C}_3\text{H}_5\text{ClO}$

$\text{ClCH}_2\text{CO.Me}$

HCS 1980, 291


1. Allen, C. F. H. *et al.*, *Ind. Eng. Chem. (News Ed.)*, 1931, **9**, 184

2. Ewe, G. E., *Ind. Eng. Chem. (News Ed.)*, 1931, **9**, 229

Two separate incidents involved explosive polymerisation of chloroacetone stored in glass bottles under ambient conditions for extended periods were reported [1,2].

See Bromoacetone oxime

†1162. 1-Chloro-2,3-epoxypropane (Chloromethyloxirane, Epichlorohydrin)
[106-89-8] C₃H₅ClO



HCS 1980, 454

Energy of decomposition (in range 375–500°C) measured as 0.5 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Aniline

Hearfield, F., *Chem. Abs.*, 1980, **92**, 115649

In a review of thermal instability in chemical reactors, the explosive interaction of 'epichlorohydrin' with aniline is detailed.

Catalyst, Heterocyclic nitrogen compounds

Anon., *Sichere Chemiearb.*, 1985, **37**, 81

The epoxide and the nitrogen compound were mixed and stirred in a 250 l reactor fitted for reflux and protected by a rupture disk. After addition of the catalyst, subsequent heating to the required temperature seems to have been effected by direct application of steam to the jacket, rather than the usual hot water. The too-rapid heating led to a violently exothermic reaction which ruptured the disk and stretched the lid clamping bolts.

Contaminants

1. TV News item, ITN, 5th January, 1984
2. Sittig, 1981, 293

A road tanker was loaded with 1-chloro-2,3-epoxypropane and then driven 250 miles overnight to the delivery point. On arrival, the contents were found to have self heated (undoubtedly from polymerisation initiated by some unknown contaminant) to the boiling point (115°C at ambient pressure) and soon afterwards the relief valve lifted and discharged large volumes of vapour. Cooling with water sprays eventually restored thermal control over the remaining tanker contents [1]. The material is incompatible with strong acids, caustic alkalis, zinc, aluminium, aluminium chloride or iron(III) chloride, all of which catalyse exothermic polymerisation [2].

See other CATALYTIC IMPURITY INCIDENTS, POLYMERISATION INCIDENTS

Isopropylamine

Barton, N. *et al.*, *Chem. & Ind.*, 1971, 994

With slow mixing and adequate cooling, smooth condensation to 1-chloro-3-isopropylamino-2-propanol occurs. With rapid mixing and poor cooling, a variable

induction period with slow warming precedes a rapid, violent exotherm (to 350°C in 6 s). Other primary and secondary amines behave similarly. Moderating effects of water and the nature of the products are discussed.

See *N*-Substituted aniline, below

See other AMINATION INCIDENTS, INDUCTION PERIOD INCIDENTS

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

Sodium chlorite

Anon. *Chemical Engineer*, 1996, (620), 4; *ibid*, 1996, (621), 7

A tanker of aqueous 'sodium chlorite' was part charged to a tank of epichlorohydrin before it was reported it was mis-labelled. There was a very violent subsequent explosion, which blew down the wall of an adjacent factory. (The editor suspects it was, in fact, sodium hypochlorite, commonly known simply as hypochlorite, which sounds like epichlorohydrin. Sodium hypochlorite would be expected to be the more reactive of the two, generating an alkyl hypochlorite.)

See HYPOHALITES

N-Substituted aniline (unspecified)

Schierwater, F.-W., *Major Loss Prevention*, 1971, 47

Interaction is exothermic and the mixture was normally maintained at 60°C by stirring and cooling. Malfunction caused a temperature increase to 70°C and cooling capacity was insufficient to regain control. The temperature steadily increased to 120°C, when explosive decomposition occurred. This was attributed to thermal instability of the reaction system and inadequate pressure relief arrangements.

See Isopropylamine, above

Sulfuric acid

Leleu, *Cahiers*, 1974, (75), 276

Interaction is violent.

Trichloroethylene

See Trichloroethylene: Epoxides

See other 1,2-EPOXIDES

†1163. Propionyl chloride (Propanoyl chloride)

[79-03-8]

EtCO.Cl

C₃H₅ClO

HCS 1980, 794

Preparative hazard

See Phosphorus trichloride: Carboxylic acids

Diisopropyl ether

Koenst, W. M. B., *J. Haz. Mat.*, 1981, 4, 291–298

A facile exothermic reaction, catalysed by traces of zinc chloride and iron(III) chloride (4 and 40 ppm, respectively) to produce isopropyl propionate and 2-chloropropane (b.p., 35°C) led to pressure build up and bursting of a closed galvanised drum after 24 h. Similar reactions are thermodynamically possible with other acid chlorides and ethers (particularly if secondary or tertiary alkyl ethers), so such mixtures should only be prepared immediately prior to use.

See Sulfinyl chloride: Esters, Metals

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

See other ACYL HALIDES

†1164. Ethyl chloroformate (Ethyl carbonochloridate)

[541-41-3]



HCS 1980, 476

See other ACYL HALIDES

†1165. Methoxyacetyl chloride (2-Methoxyethanoyl chloride)

[38870-89-2]



Anon., *Jahresber.*, 1979, 75

A sample of redistilled material in a screw capped bottle exploded 15 weeks after capping. This was attributed to development of internal pressure (probably of hydrogen chloride arising from hydrolysis, and perhaps also of chloromethane from scission). (The editor has also known methoxyacetic mixed anhydrides to decarbonylate, releasing carbon monoxide)

See other ACYL HALIDES, GAS EVOLUTION INCIDENTS

1166. Methoxycarbonylmethyl perchlorate

[95407-67-3]



See entry ALKYL PERCHLORATES (reference 6)

1167. Peroxypropionyl perchlorate

[66955-44-0]



Yakarka, A. A. *et al.*, *Chem. Abs.*, 1979, **91**, 210847

The oxidant exploded on detonation or friction.

See related ACID ANHYDRIDES

See other NON-METAL PERCHLORATES, ORGANIC PEROXIDES

†1168. *N,N*-Dichloro- β -alanine (2-Dichloroaminopropanoic acid)
[58941-15-4] C₃H₅Cl₂NO₂
 $\text{Cl}_2\text{NCH}(\text{Me})\text{CO.OH}$

Vit, J. *et al.*, *Synth. Commun.*, 1976, **6**(1), 1–4
It is thermally unstable above 95°C.
See other N-HALOGEN COMPOUNDS


1169. Allyl phosphorodichloridite C₃H₅Cl₂OP
[1498-47-1]
 $\text{H}_2\text{C}=\text{CHCH}_2\text{OPCl}_2$

Kamler, M. *et al.*, *Can. J. Chem.*, 1985, **63**, 825–826
The title compound is formed by reaction of allyl alcohol with phosphorus trichloride and is reported to polymerise explosively if the material, when being purified by distillation, is taken down to less than half its bulk.
See other ALLYL COMPOUNDS, PHOSPHORUS ESTERS

1170. *N*-Carboethoxyiminophosphoryl chloride C₃H₅Cl₃NO₂P
[3356-63-6]
 $\text{EtOCO.N}=\text{PCl}_3$

See N-Carbomethoxyiminophosphoryl chloride
See related NON-METAL HALIDES

†1171. 1-Fluoro-2,3-epoxypropane (Fluoromethyloxirane) C₃H₅FO
[503-09-3]



See other 1,2-EPOXIDES

1172. *N,N,N'*-Trifluoropropionamidine C₃H₅F₃N₂
[21372-60-1]
 $\text{EtC}(\text{:NF})\text{NF}_2$

Ross, D. L. *et al.*, *J. Org. Chem.*, 1970, **35**, 3096–3097
A shock-sensitive explosive.
See other N, N, N''-TRIFLUOROAMIDINES

1173. Allylmercury(II) iodide
[2845-00-3]



$\text{C}_3\text{H}_5\text{HgI}$

Potassium cyanide

Whitmore, 1921, 122

Among the products of interaction, a minor one is an explosive liquid.

See other ALLYL COMPOUNDS, MERCURY COMPOUNDS

†**1174. 3-Iodopropene (Allyl iodide)**
[556-56-9]



$\text{C}_3\text{H}_5\text{I}$

See other ALLYL COMPOUNDS, HALOALKENES, IODINE COMPOUNDS

1175. Potassium 1,1-dinitropropanide
[30533-63-2]




$\text{C}_3\text{H}_5\text{KN}_2\text{O}_4$

Bisgrove, D. E. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 373

The potassium salt of 1,1-dinitropropane, isolated as a by-product during preparation of 3,4-dinitro-3-hexene, is a hazardous explosive.

See NITROALKANES: alkali metals See other POLYNITROALKYL COMPOUNDS

†**1176. Potassium O-ethyl dithiocarbonate ('Potassium ethyl xanthate')**
[140-89-6]



$\text{C}_3\text{H}_5\text{KOS}_2$

Diazonium salts

See DIAZONIUM SULFIDES AND DERIVATIVES

1177. Allyllithium
[3052-45-7]



$\text{C}_3\text{H}_5\text{Li}$

Leleu, *Cahiers*, 1977, (88), 367

As usually prepared, it is a pyrophoric solid.

See other ALKYL METALS, ALLYL COMPOUNDS

†**1178. Ethyl isocyanide (Isocynoethane)**
[624-79-3]



$\text{C}_3\text{H}_5\text{N}$

Lemoult, M. P., *Compt. rend.*, 1906, **143**, 903

This showed a strong tendency to explode while being sealed into glass ampoules.

See Methyl isocyanide

See related CYANO COMPOUNDS

†1179. Propionitrile (Propanonitrile)

[107-12-0]

C₃H₅N

EtCN

Preparative hazard

See Nitric acid: Aliphatic amines

N-Bromosuccinimide

See N-Bromosuccinimide: Propionitrile

See other CYANO COMPOUNDS

†1180. Acrylamide (Propenamide)

[79-06-1]

C₃H₅NO

H₂C=CHCO.NH₂

HCS 1980, 114

Bretherick, 1981, 165

It may polymerise with violence on melting at 86°C.

See other POLYMERISATION INCIDENTS

†1181. 2-Cyanoethanol (3-Hydroxypropanenitrile)

[109-78-4]

C₃H₅NO

N≡CC₂H₄OH

HCS 1980, 479

Acids, or Bases

HCS 1980, 479


Reacts violently with mineral acids, amines or inorganic bases, probably because of dehydration to acrylonitrile and subsequent catalysed polymerisation of the latter.

See other POLYMERISATION INCIDENTS See other CYANO COMPOUNDS

1182. 2,3-Epoxypropionaldehyde oxime (Oxiranecarboxaldehyde oxime)

[67722-96-7]

C₃H₅NO₂



Payne, G. B., *J. Amer. Chem. Soc.*, 1959, **81**, 4903

The residue from distillation at 48–49°C/1.3 mbar polymerised violently, and the distilled material polymerised explosively after 1–2 h at ambient temperature.

See 2,3-Epoxypropionaldehyde 2,4-dinitrophenylhydrazone
See other 1,2-EPOXIDES, OXIMES, POLYMERISATION INCIDENTS
See related ALDEHYDES

1183. 2-Nitropropene

[4749-28-4]

$C_3H_5NO_2$



Miyashita, M. *et al.*, *Org. Synth.*, 1981, **60**, 101–103

Vacuum distillation must be effected at below 80°C to avoid fume-offs, particularly if air be admitted to the warm residue. The lachrymatory material also polymerises in contact with alkalis.

See other NITROALKENES, POLYMERISATION INCIDENTS

†**1184. Nitroacetone (1-Nitro-2-propanone)**

[10230-68-9]

$C_3H_5NO_3$



1. Tegeler, J. J., *Chem. Eng. News*, 1987, **65**(1), 4

2. Author's comment, 1987

Nitroacetone (237 g) was generated in dichloromethane–ethyl acetate solution by treating its dicyclohexylamine *aci*-salt with sulfuric acid. Towards the end of concentration of the solution by rotary vacuum evaporation at 40–50°C, a small explosion occurred. Extreme caution and small scale handling of α -nitroketones are urged [1]. If exact neutralisation of the dicyclohexylamine salt had not been achieved, presence of excess sulfuric acid (or of some undecomposed *aci*-salt) may have reduced the stability of the nitroacetone [2].

See Nitromethane: Acids, or Bases;

See also *aci*-NITRO SALTS

See related NITROALKANES

1185. Propionyl nitrite

[28128-14-5]

$C_3H_5NO_3$




See entry ACYL NITRITES

†**1186. 2,3-Epoxypropyl nitrate (Oxiranemethanol nitrate)**

[6659-62-7]

$C_3H_5NO_4$



Urbanski, 1965, Vol. 2, 129

Shock-sensitive and explodes at 200°C.

See related ALKYL NITRATES, 1,2-EPOXIDES

1187. Peroxypropionyl nitrate
[5796-89-4]

$C_3H_5NO_5$



1. Stephens, E. R., *Anal. Chem.*, 1964, **36**, 928–929
2. Louw, R. *et al.*, *J. Amer. Chem. Soc.*, 1975, **97**, 4396

Like the lower homologues, it is extremely explosive [1], and may only be handled in high dilution with air or nitrogen. Higher homologues are less explosive [2].

See related ACID ANHYDRIDES, ACYL NITRATES, PEROXYESTERS

†**1188. 3-Azidopropene (Allyl azide)**
[821-13-6]

$C_3H_5N_3$




1. Anon., *Saf. Digest*, (Univ. Saf. Assoc.), 1985, (13), 2–3
2. Forster, M. O., *J. Chem. Soc.*, 1908, **93**, 1174

An explosion occurred in the final stages of evaporation of ether from allyl azide [1]. Peroxides in the ether may possibly have been involved, but the azide itself is of limited thermal stability, though apparently capable of distillation at 76°C/1 bar. The vapour is weakly explosive [2].

See other ALLYL COMPOUNDS, ORGANIC AZIDES

1189. 1-Methyl-1,2,3-triazole
[16681-65-5]

$C_3H_5N_3$



Anon., private comm., 1980

An explosion during the preparation of the triazole was attributed to local overheating or presence of impurities.

See other HIGH-NITROGEN COMPOUNDS, TRIAZOLES

†**1190. Azidoacetone (Azido-2-propanone)**
[4504-27-2]

$C_3H_5N_3O$



1. Spauschus, H. O. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 209
2. Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 72

A small sample exploded after storage in the dark for 6 months [1]. The freshly prepared material explodes when dropped on to a hotplate and burns brilliantly [2].


See other 2-AZIDOCARBONYL COMPOUNDS

1191. Glycidyl azide (Azidomethyloxirane)

[80044-09-3]

$C_3H_5N_3O$

[80044-10-6] polymer($-CH_2CHOHCH_2(N_3)-$)_n



Nazare, A. N. *et al.*, *J. Energ. Mater.*, 1992, **10**(1), 43


A review on the derived polymer as a propellant component, which use depends upon the energy of the azide. The monomer would be expected to have several kJ/g, including oxirane ring strain.

See 1,2-EPOXIDES, ORGANIC AZIDES

1192. 5-Amino-3-methylthio-1,2,4-oxadiazole

[55864-39-6]

$C_3H_5N_3OS$



Wittenbrook, L. S., *J. Heterocycl. Chem.*, 1975, **12**, 38

It undergoes rapid and moderately violent decomposition at the m.p., 97–99°C.

See other N–O COMPOUNDS

†1193. Ethyl azidoformate (Ethyl carbonazidate)

[817-87-8]

$C_3H_5N_3O_2$

$EtOCO.N_3$

Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 91


It is liable to explode if boiled at ambient pressure (at 114°C). The methyl ester (b.p. 103°C) can behave similarly.

See other ACYL AZIDES

1194. 2-Ammoniothiazole nitrate

[57530-25-3]


$C_3H_5N_3O_3S$



1. Rebsch, M., *Proc. Int. Symp. Prev. Occ. Risks Chem. Ind.*, 485–486, Heidelberg, ISSA., 1976

2. Soldarna Marlör, L., *Chem. Abs.*, 1977, **87**, 203834


During air-pressurised discharge of a hot 53% aqueous solution of the nitrate salt from a reaction vessel via a filter press, a violent explosion occurred. The nitrate salt begins to decompose below 100°C, and at the likely internal temperature of 142°C, decomposition would be expected to be very rapid, involving much gas/vapour generation according to the equation below.



Air-pressurisation would have prevented the water from boiling and absorbing the exotherm to moderate the thermal explosion. Decomposition of the solution may have been accelerated by the presence of impurities, or by the solid salt splashed onto the heated vessel wall [1]. There is an independent account of the incident [2].

See other GAS EVOLUTION INCIDENTS

1195. 3,3-Dinitroazetidine
[129660-17-9]



Hiskey, M. A. *et al.*, *J. Energ. Mater.*, 1993, **11**(3), 157.

Dinitroazetidine salts were studied as explosives: nitrate, dinitroimidate, 2,4-dinitroimidazolate, 4,4',5,5'-tetranitro-2,2'-biimidazolate, 5-nitro-1,2,4-triazolonate and 3,5-dinitrotriazolate.

See other C-NITRO COMPOUNDS, STRAINED-RING COMPOUNDS

†**1196. Glyceryl trinitrate (1,2,3-Propanetriyl nitrate)**
[55-63-0]



1. Urbanski, 1965, Vol. 2, 47, 53–54
2. Bond, J., *Loss Prev. Bull.*, 1983, (050), 18

The relatively high sensitivity of the pure liquid material to initiation by shock or heating (it explodes at 215–218°C, and decomposes energetically above 145°C) is enhanced by presence of impurities, especially nitric acid, arising from imperfect washing, hydrolysis or thermal aging (e.g. heating to 50°C). Intense UV radiation will explode a sample at 100°C. The sensitivity of the viscous liquid to shock is slightly reduced by freezing, but is greatly diminished by absorption onto porous solids, as in dynamite [1]. Many incidents, some dating back to 1870, are summarised, all involving violent explosions when cans or buckets of nitroglycerine were dropped. Some of these detonations initiated the detonation of bulk material nearby [2].

See other ALKYL NITRATES, IRRADIATION DECOMPOSITION INCIDENTS

†1197. Cyclopropane
[75-19-4]

C₃H₆



HCS 1980, 351 (cylinder)

Stull, 1977, 16


It is fairly endothermic and of low MW (ΔH_f° (g) +82.3 kJ/mol, 1.96 kJ/g), and a minor constituent of MAPP gas.

See Propyne

See other ENDOTHERMIC COMPOUNDS, STRAINED-RING COMPOUNDS

†1198. Propene (Propylene)
[115-07-1]

C₃H₆



(*MCA SD-59*, 1956); *FPA H75*, 1978; *HCS 1980*, 788 (cylinder)

1. Russell, F. R. *et al.*, *Chem. Eng. News*, 1952, **30**, 1239

2. Stull, 1977, 16

Several sets of values for flammability limits in air are quoted; 1.4–7.1, 2.4–11.1, 2.7–36.0 vol%.

Propene at 955 bar and 327°C was being subjected to further rapid compression. At 4.86 kbar explosive decomposition occurred, causing a pressure surge to 10 kbar or above. Decomposition to carbon, hydrogen and methane must have occurred to account for this pressure. Ethylene behaves similarly at much lower pressure, and cyclopentadiene, cyclohexadiene, acetylene and a few aromatic hydrocarbons have been decomposed explosively [1]. It is mildly endothermic (ΔH_f° (g) +20.4 kJ/mol, 0.49 kJ/g) and a minor constituent of MAPP gas [2].

Lithium nitrate, Sulfur dioxide

Pitkethly, R. C., private comm., 1973

A mixture under confinement in a glass pressure bottle at 20°C polymerised explosively, the polymerisation probably being initiated by access of light through the clear glass container. Such alkene–sulfur dioxide co-polymerisations will not occur above a ceiling temperature, different for each alkene.

See other GLASS INCIDENTS, POLYMERISATION INCIDENTS

Other reactants

Yoshida, 1980, 321

MRH values calculated for 13 combinations with oxidants are given.

Oxides of nitrogen

See Nitrogen oxide: Dienes, Oxygen

Trifluoromethyl hypofluorite
See Trifluoromethyl hypofluorite: Hydrocarbons

Water
See LIQUEFIED GASES: water
See other ALKENES

1199. Cyanodimethylarsine
[683-45-4]



C₃H₆AsN

491M, 1975, 140
It ignites in air.
See related ALKYLNON-METAL HALIDES

1200. Dimethylgold selenocyanate
[42494-76-1]



C₃H₆AuNSe

Stocco, F. *et al.*, *Inorg. Chem.*, 1971, **10**, 2640
Very shock-sensitive and explodes readily when precipitated from aqueous solution. Crystals obtained by slow evaporation of a carbon tetrachloride extract were less sensitive.
See other GOLD COMPOUNDS *See related* METAL CYANATES

1201. Bromoacetone oxime (1-Bromo-2-oximinopropane)
[62116-25-0]



C₃H₆BrNO

Forster, M. O *et al.*, *J. Chem. Soc.*, 1908, **93**, 84
It decomposes explosively during distillation.
See Chloroacetaldehyde oxime, and Chloroacetone
See other OXIMES

†1202. N-Chloroallylamine (N-Chloro-3-aminopropene)
[82865-33-6]



C₃H₆ClN

It explodes at ambient temperature.
See entry N-HALOGEN COMPOUNDS (reference 4)
See other ALLYL COMPOUNDS

†1203. 1,1-Dichloropropane
[78-99-9] Cl_2CHEt $\text{C}_3\text{H}_6\text{Cl}_2$

See other HALOALKANES

†1204. 1,2-Dichloropropane
[78-87-5] $\text{ClCH}_2\text{CHClCH}_3$ $\text{C}_3\text{H}_6\text{Cl}_2$

HCS 1980, 383

Aluminium

See Aluminium: Halocarbons (reference 18)

Aluminium, 1,2-Dichlorobenzene, 1,2-Dichloroethane

See Aluminium: Halocarbons (reference 5)

See other HALOALKANES

1205. 2,2-Dichloropropane
[594-20-7] $\text{H}_3\text{CCCl}_2\text{CH}_3$ $\text{C}_3\text{H}_6\text{Cl}_2$

Dimethylzinc

See Dimethylzinc: 2,2-Dichloropropane

See other HALOALKANES

1206. 1-Chloro-2-propyl perchlorate
[58426-27-0] $\text{ClCH}_2\text{CH}(\text{Me})\text{OCIO}_3$ $\text{C}_3\text{H}_6\text{Cl}_2\text{O}_4$

See entry ALKYL PERCHLORATES (reference 6)

1207. 3-Chloro-2-hydroxypropyl perchlorate
[101672-07-5] $\text{ClCH}_2\text{CH}(\text{OH})\text{CH}_2\text{OCIO}_3$ $\text{C}_3\text{H}_6\text{Cl}_2\text{O}_5$

Hofmann, K. A. *et al.*, *Ber.*, 1909, **42**, 4390

It explodes violently on shaking in a capillary.


See other ALKYL PERCHLORATES

†1208. Dichloromethylvinylsilane
[124-70-9] C₃H₆Cl₂Si



See related ALKYLHALOSILANES

1209. Dichloromethylenedimethylammonium chloride
[33843-02-3] C₃H₆Cl₃N



1. Bader, A. R., private comm., 1971
 2. Senning, A., *Chem. Rev.*, 1965, **65**, 388
- The compound does not explode on heating [1]; the earlier reference [2] is in error.

1210. Potassium *O*-propionhydroxamate
[71939-10-1] C₃H₆KNO₂



Anon., *Actual. Chim. (Fr.)*, 1983, **8**, 49–50
The solid explodes on attempted drying. DTA indicates that the rapid decomposition at 70–74°C to potassium hydroxide and ethyl isocyanate, accompanied by an exotherm of 1.37 kJ/g, is involved in the explosion.
See other N–O COMPOUNDS

1211. Dimethylthallium fulminate
[] C₃H₆NOTl



Beck, W. *et al.*, *J. Organomet. Chem.*, 1965, **3**, 55
Highly explosive, unlike the diphenyl analogue.
See related METAL FULMINATES


1212. 3-Aminopropiononitrile
[151-18-8] C₃H₆N₂



1. Shner, V. F., *Zhur. Prikl. Khim.*, 1966, **39**, 2386 (Engl. transl.)
 2. *MCA Guide*, 1972, 309
 3. Merck, 1976, 65
- Vacuum distilled material kept dark for 6 months was found to have largely polymerised to a yellow solid, which exploded 15 days later [1]. A bottle of uninhibited distilled material exploded after shelf storage for several months [2]. It may be stored for several months out of contact with air under refrigeration. In contact with ambient air, it polymerises slowly, but rapidly in contact with acidic materials [3].
See other CYANO COMPOUNDS, POLYMERISATION INCIDENTS

1213. 2-Ethyltetrazole
[82944-28-3]

C₃H₆N₄




Aluminium hydride

See Aluminium hydride: Tetrazole derivatives

See other TETRAZOLES

1214. 5-Ethyltetrazole
[50764-78-8]

C₃H₆N₄



Aluminium hydride

See Aluminium hydride: Tetrazole derivatives

See other TETRAZOLES

1215. Azidoacetone oxime
[101672-04-2]

C₃H₆N₄O



Forster, M. O. *et al.*, *J. Chem. Soc.*, 1908, **93**, 83

During distillation at 84°C/2.6 mbar, the large residue darkened and finally exploded violently.

See Bromoacetone oxime

See other 2-AZIDOCARBONYL COMPOUNDS

1216. 2,2-Diazidopropane
[85620-95-7]

C₃H₆N₆




Al-Khalil, S. I. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1986, 555–565

The liquid azide, b.p. 28–30°C/0.6 mbar, exploded during attempted analytical combustion.

See other ORGANIC AZIDES

1217. 1,3,5-Trinitrosohexahydro-1,3,5-triazine
[13980-04-6]




Sulfuric acid

Urbanski, 1967, Vol. 3, 122

It decomposes explosively in contact with conc. sulfuric acid.

See other NITROSO COMPOUNDS

1218. 5-Nitro-2-(nitroimino)hexahydro-1,3,5-triazine (1,4,5,6-Tetrahydro-*N*,5-dinitrotriazin-2-amine)
[130400-13-4]





Dagley, I. J., *et al.*, *J. Energ. Mater.*, 1995, **13**(1 & 2), 35

This nitroguanidine analogue is more sensitive than RDX by the drop-weight impact test.

See other *N*-NITRO COMPOUNDS

1219. 1,3,5-Trinitrohexahydro-1,3,5-triazine (Cyclotrimethylenetrinitramine)
[121-82-4]



Botcher, T. R. *et al.*, *J. Amer. Chem. Soc.*, 1992, **114**(2), 8303

Some studies relating to the explosion mechanism of this military explosive (RDX or Cyclonite) and its higher homologue HMX.

Calcium hydroxide, Water

Iida, M. *et al.*, *Chem. Abs.*, 1988, **109**, 112968

The self-heating and decomposition of the explosive and aqueous alkali was studied by DSC in a sealed capsule and in a larger scale furnace test. A rapid exothermic decomposition reaction can be initiated at 100°C or below, and may lead to spontaneous ignition and then deflagration or detonation.

See other N-NITRO COMPOUNDS

See also NITROAROMATIC-ALKALI HAZARDS

†1220. Acetone (2-Propanone)

[67-64-1]

C₃H₆O

H₃CCO.CH₃

(MCA SD-87, 1962); NSC 398, 1982; FPA H1, 1972; HCS 1980, 102;
RSC Lab. Hazard Data Sheet No. 21, 1984

Bromoform

See Bromoform: Acetone, etc.

Carbon, Air

Boiston, D. A., *Brit. Chem. Eng.*, 1968, **13**, 85

(In the absence of other ignition source), fires in plant to recover acetone from air with active carbon are due to the bulk surface effect of oxidative heating when air flow is too low to cool effectively.

Chloroform

See Chloroform: Acetone, etc.

2-Methyl-1,3-butadiene

See 2-Methyl-1,3-butadiene: Acetone

Nitric acid, Sulfuric acid

MRH Nitric 5.86/77, Sulfuric 1.05/53

Fawcett, H. H., *Ind. Eng. Chem.*, 1959, **51**(4), 89A

Acetone will be oxidised with explosive violence if brought into contact with the mixed (nitrating) acids, particularly under confinement.

See Oxidants, below

Other reactants

Yoshida, 1980, 20

MRH values calculated for 17 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Nitric acid, Sulfuric acid, above

Bromine: Acetone

Bromine trifluoride: Solvents
 Chromium trioxide: Acetone MRH 2.59/90
 Chromyl chloride: Organic solvents
 Dioxygen difluoride: Various materials
 Hydrogen peroxide: Acetone MRH 6.36/81
 Hydrogen peroxide: Ketones, or: Oxygenated compounds
 Nitric acid: Acetone MRH 5.86/77
 Nitrosyl chloride: Acetone, etc.
 Nitrosyl perchlorate: Organic materials
 Oxygen (Gas): Acetylene, Acetone
 Peroxomonosulfuric acid: Acetone

Peroxyacetic acid

See 5-Bromo-4-pyrimidinone

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

Sulfur dichloride

See Sulfur dichloride: Acetone

Thiotriithiazyl perchlorate

See Thiotriithiazyl perchlorate: Organic solvents

†1221. Methyl vinyl ether (Methoxyethene)

[107-25-5]

C₃H₆O



HCS 1980, 664 (cylinder)

Acids

1. Braker, 1980, 487

2. 'MVE' Brochure, Billingham, ICI, 1962

Methyl vinyl ether is rapidly hydrolysed by contact with dilute acids to form acetaldehyde, which is more reactive and has wider flammability limits than the ether [1]. Presence of base is essential during storage or distillation of the ether to prevent rapid acid-catalysed homopolymerisation, which is not prevented by antioxidants. Even mildly acidic solids (calcium chloride or some ceramics) will initiate exothermic polymerisation [2].

See other POLYMERISATION INCIDENTS

Halogens, or Hydrogen halides

Braker, 1980, 487

Addition reactions with bromine, chlorine, hydrogen bromide or hydrogen chloride are very vigorous and may be explosive if uncontrolled.

See other PEROXIDISABLE COMPOUNDS

†1222. Oxetane
[503-30-0]

C₃H₆O



Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Strained ring heterocycles

See other STRAINED-RING COMPOUNDS

†1223. 2-Propen-1-ol (Allyl alcohol)
[107-18-6]

C₃H₆O



HCS 1980, 122

Energy of decomposition (in range 360–500°C) measured as 0.69 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkali, 2,4,6-Trichloro-1,3,5-triazine MRH Sodium hydroxide 1.21/tr.
See 2,4,6-Trichloro-1,3,5-triazine: Allyl alcohol, etc.

Carbon tetrachloride MRH 1.42/73
491M, 1975, 21

Interaction gives an unstable mixture of halogenated C₄ epoxides which exploded during distillation.

Other reactants

Yoshida, 1980, 24

MRH values calculated for 15 combinations, mainly with oxidants, are given.

Sulfuric acid MRH 1.42/53

Senderens, J. B., *Compt. rend.*, 1925, **181**, 698–700

During preparation of diallyl ether by dehydration of the alcohol with sulfuric acid, a violent explosion may occur (possibly involving peroxidation and certainly polymerisation).

See other PEROXIDATION INCIDENTS, POLYMERISATION INCIDENTS

2,4,6-Tris(bromoamino)-1,3,5-triazine

See 2,4,6-Tris(bromoamino)-1,3,5-triazine: Allyl alcohol

See other ALLYL COMPOUNDS

†1224. Propionaldehyde (Propanal)
[123-38-6]

C₃H₆O



HCS 1980, 791

Methyl methacrylate
See Methyl methacrylate: Propionaldehyde
See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

†1225. Propylene oxide (Methyloxirane)
[75-56-9]

C₃H₆O



FPA H88, 1980; HCS 1980, 800

1. CHETAH, 1990, 183
 2. Kulik, M. K., *Science*, 1967, **155**, 400
 3. Smith, R. S., *Science*, 1967, **156**, 12
 4. Pogany, G. A., *Chem. & Ind.*, 1979, 16–21
 5. See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
- The liquid can be exploded by a detonator, though not by mechanical shock [1]. Use of propylene oxide as a biological sterilant is hazardous because of ready formation of explosive mixtures with air (2.8–37%). Commercially available mixtures with carbon dioxide, though non-explosive, may be asphyxiant and vesicant [1]. Such mixtures may be ineffective, but neat propylene oxide vapour may be used safely, provided that it is removed by evacuation using a water-jet pump [2]. The main factors involved in the use and safe handling on a laboratory scale have been discussed [3]. The energy of decomposition (in range 340–500°C) has been measured as 1.114 kJ/g [4].

Epoxy resin

Sheaffer, J., *CHAS Notes*, 1981, **1**(4), 5

Mixing of propylene oxide and epoxy resin in a waste bottle led to an explosion, probably owing to the polymerisation of the oxide catalysed by the amine accelerator in the resin.

See Sodium hydroxide, below

Ethylene oxide, Polyhydric alcohol

See Ethylene oxide: Polyhydric alcohol, etc.

Other reactants

Yoshida, 1980, 322

MRH values calculated for 13 combinations with oxidants are given.

Oxygen

1. Hassan, A. *et al.*, *Combust. Sci. Technol.*, 1983, **35**, 215–230
2. Peng, J., *Proc. Int. Pyrotech. Semin.*, 1991, (17th), Vol. 1, 353

The transition of deflagration to detonation in mixtures was studied with respect to mixing ratio, pressure and spark energy [1]. A study of TNT equivalences in propylene oxide fuel/air explosives is made [2].

Sodium hydroxide

1. *MCA Case History No. 31*

2. Freeder, B. G. *et al.*, *J. Loss Prev. Process Ind.*, 1988, **1**, 164–168

A drum of crude product containing unreacted propylene oxide and sodium hydroxide catalyst exploded and ignited, probably owing to base-catalysed exothermic polymerisation of the oxide [1]. A comparative ARC study of the runaway exothermic polymerisation of ethylene oxide and the less reactive propylene oxide in presence of sodium hydroxide solutions, as typical catalytically active impurities, has been done. The results suggest that the hazard potential for propylene oxide is rather less than that for the lower homologue, though more detailed work is needed to quantify the difference [2].

See Ethylene oxide: Contaminants

See other CATALYTIC IMPURITY INCIDENTS, POLYMERISATION INCIDENTS

See other 1,2-EPOXIDES

1226. Allyl hydroperoxide (2-Propenyl hydroperoxide)

[6069-42-7]

$C_3H_6O_2$



1. Dykstra, S. *et al.*, *J. Amer. Chem. Soc.*, 1957, **79**, 3474

2. Seyfarth, H. E. *et al.*, *Angew. Chem.*, 1965, **77**, 1078


When impure, the material is unstable towards heat or light and decomposes to give an explosive residue. The pure material is more stable to light, but detonates on heating or in contact with solid alkalis [1]. Preparation by action of oxygen on diallylzinc gives improved yields, but there is a risk of explosion. The peroxide is also impact-sensitive if sand is admixed [2].

See other ALKYL HYDROPEROXIDES, ALLYL COMPOUNDS

1227. Dimethyldioxirane

[74087-85-7]

$C_3H_6O_2$



1. Murray, R. W., private communication, 1992

This material, and homologues, are being recommended as oxidants, they are prepared in situ and very dilute. Thus far, the material has shown itself surprisingly stable during isolation and pyrolysis (decomposes slowly at ambient temperatures and quietly on heating) and even explosivity testing [1]. However several


known oligomers, presumably of lower (strain) energy, are dangerously sensitive explosives. Considerable caution during work-up of reactions is recommended.

See Bis(trifluoromethyl)dioxirane

See Hydrogen Peroxide: Acetone

See other CYCLIC PEROXIDES


†1228. 1,3-Dioxolane
[646-06-0]



Lithium perchlorate

See Lithium perchlorate: 1,3-Dioxolane

1229. 2,3-Epoxypropanol (Oxiranemethanol)
[556-52-5]



HCS 1980, 518

1. Anon., *Jahresber.*, 1982, 67–68; *Sichere Chemiearb.*, 1984, **36**, 80
2. Anon, *Loss Prevention Bull.*, 1989, **88**, 17
3. Cardillo, P. *et al.*, *J. Loss Prev. Process Ind.*, 1991, **4**(4), 242

Off-specification 'glycidol' from a production unit was collected in a 40 t storage tank and recovered occasionally by redistillation. As no stabiliser is known, the contents were water-cooled to prevent self-heating and polymerisation, and pump-circulated to ensure homogeneity. A 9 t quantity of the stored material was found to be quite warm, and soon after, the tank exploded, causing 3 fatalities. The contents of the tank had been subjected to storage tests at 60°C on 200 l scale, when decomposition but no runaway was seen, presumably the drums were not adiabatic enough. It is known that the epoxy compound will undergo explosive decomposition in presence of strong acids or bases, salts (aluminium chloride, iron(III) chloride, tin(IV) chloride), or metals (copper or zinc), but these were not likely to have been present. It was concluded that the water cooling had been turned off, and that the energy dissipated by the circulation pump would have raised the temperature of the tank contents to 50°C under the prevailing weather conditions, the latter conclusion being confirmed experimentally [1]. The energy of decomposition (in the range 130–450°C) has been measured by DSC as 1.365 kJ/g, and T_{ait24} was determined as the low figure of 65°C by adiabatic Dewar tests,

with an apparent energy of activation of 71 kJ/mol. A detailed study of the thermochemistry of glycidol, including CHETAH calculations and calorimetry (ARC and DSC), showed a two stage reaction; polymerisation from 60°C, followed by a decomposition with pressure generation from 280°C. Polymerisation was catalysed by NaOH and triethanolamine but not, apparently, by HCl or triethylamine [3].

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

See other 1,2-EPOXIDES

See also CALORIMETRY; COMPUTATION OF REACTIVE CHEMICAL HAZARDS

†1230. Ethyl formate (Ethyl methanoate)

[109-94-4]



$\text{C}_3\text{H}_6\text{O}_2$

HCS 1980, 488

Base

See Methyl formate: Methanol, Sodium methoxide

1231. Hydroxyacetone (Propan-1-ol-2-one)

[116-09-6]



$\text{C}_3\text{H}_6\text{O}_2$

Preparative hazard

See 3-Propynol: Mercury(II) sulfate, etc.

†1232. Methyl acetate (Methyl ethanoate)

[79-20-9]



$\text{C}_3\text{H}_6\text{O}_2$

FPA H124, 1983; *HCS 1980*, 426

Other reactants

Yoshida, 1980, 123

MRH values calculated for 13 combinations with oxidants are given.

†1233. Dimethyl carbonate

[616-38-6]



$\text{C}_3\text{H}_6\text{O}_3$

HCS 1980, 426

Other reactants

Yoshida, 1980, 224

MRH values calculated for 13 combinations with oxidants are given.

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

1234. Lactic acid (2-Hydroxypropanoic acid)

[598-82-3]



C₃H₆O₃

Hydrofluoric acid, Nitric acid


See Nitric acid: Hydrofluoric acid, Lactic acid

See other ORGANIC ACIDS

1235. 3-Methyl-1,2,4-trioxolane (Propene ozonide)

[38787-96-1]

C₃H₆O₃



Briner, E. *et al.*, *Helv. Chim. Acta*, 1929, **12**, 181

Stable at 0°C but often explosively decomposes at ambient temperature.

See other CYCLIC PEROXIDES, OZONIDES

1236. Peroxypropionic acid (Propaneperoxoic acid)

[4212-43-5]



C₃H₆O₃

1. Swern, D., *Chem. Rev.*, 1949, **45**, 9

2. Phillips, B. *et al.*, *J. Org. Chem.*, 1959, **23**, 1823

More stable than its lower homologues, it merely deflagrates on heating. Higher homologues appear to be still more stable [1]. Safe procedures (on the basis of detonability experiments) for preparation of anhydrous solutions of peroxypropionic acid in chloroform or ethyl propionate have been described [2].


Organic solvents

Manly, T. D., *Chem. Brit.*, 1982, **18**, 341

A violent explosion occurred during the distillation of solvents that had contained the peroxyacid. Although the solvents had been treated by addition of sodium sulfite to reduce the acid, dipropionyl peroxide (which is not reduced by sulfite, and which may have been produced by the sulfite treatment) remained in solution and subsequently decomposed on heating. If the solvent is added to the sulfite solution, the diacyl peroxide is not formed.

See DIACYL PEROXIDES *See other* PEROXYACIDS

1237. 1,3,5-Trioxane
[110-88-3]



HCS 1980, 942
MCA Case History No. 1129

Use of an axe to break up a large lump of trioxane caused ignition and a vigorous fire. Peroxides may have been involved.

Acrylonitrile
See Acrylonitrile: Formaldehyde

Hydrogen peroxide, Lead
See Hydrogen peroxide: Lead, Trioxane

Oxygen (Liquid)
See Oxygen (Liquid): 1,3,5-Trioxane
See PEROXIDISABLE COMPOUNDS

1238. 3-Thietanol-1,1-dioxide
[22524-35-2]



Preparative hazard
See Hydrogen peroxide: Acetic acid, 3-Thietanol

†**1239. 2-Propene-1-thiol (Allyl mercaptan)**
[870-23-5]



See other ALLYL COMPOUNDS

1240. Diazomethyldimethylarsine
[59871-26-0]



Preparative hazard
See Diazomethane: Dimethylaminodimethylarsine
See other DIAZO COMPOUNDS

†1241. 1-Bromopropane (Propyl bromide)
[106-94-5] C₃H₇Br
 BrCH_2Et

See other HALOALKANES

†1242. 2-Bromopropane (Isopropyl bromide)
[75-26-3] C₃H₇Br
 MeCHBrMe

See other HALOALKANES

†1243. 1-Chloropropane (Propyl chloride)
[540-54-5] C₃H₇Cl
 ClCH_2Et

See other HALOALKANES

†1244. 2-Chloropropane (Isopropyl chloride)
[75-29-6] C₃H₇Cl
 MeCHClMe

HCS 1980, 578

See other HALOALKANES

1245. *S*-Carboxymethylisothiuronium chloride (Carboxymethyl
carbamimioniothioate chloride)
[5425-78-5] C₃H₇ClN₂O₂S
 $\text{HOCO.CH}_2\text{SC}(\text{:N}^+\text{H}_2)\text{NH}_2\text{Cl}^-$

Chlorine

See Chlorine: Nitrogen compounds (reference 4)

See related ORGANIC ACIDS

†1246. Chloromethyl ethyl ether (Chloromethoxyethane)
[3188-13-4] C₃H₇ClO
 ClCH_2OEt

See related HALOALKANES

1247. Isopropyl hypochlorite
[53578-07-7] C₃H₇ClO
 Me_2CHOCl

1. Chattaway, F. D. *et al.*, *J. Chem. Soc.*, 1923, **123**, 3001

2. Mellor, 1956, Vol. 2, Suppl. 1, 550

Of extremely low stability; explosions occurred during its preparation if cooling was inadequate [1], or on exposure to light [2].

See other HYPOHALITES, IRRADIATION DECOMPOSITION INCIDENTS

1248. 3-Chloro-1,2-propanediol

[96-24-2]



$\text{C}_3\text{H}_7\text{ClO}_2$

Perchloric acid

See Perchloric acid: Glycols, etc. (reference 1)

1249. Propyl perchlorate

[22755-14-2]



$\text{C}_3\text{H}_7\text{ClO}_4$

See entry ALKYL PERCHLORATES

1250. 2-Propyl perchlorate

[52936-53-1]



$\text{C}_3\text{H}_7\text{ClO}_4$

See entry ALKYL PERCHLORATES (reference 6)

1251. Propylcopper(I)

[18365-12-3]



$\text{C}_3\text{H}_7\text{Cu}$

Houben-Weyl 1970, Vol. 13.1, 737

Small quantities only should be handled because of the danger of explosion.

See other ALKYL METALS

†**1252. 2-Iodopropane (Isopropyl iodide)**

[75-30-9]



$\text{C}_3\text{H}_7\text{I}$

See other HALOALKANES, IODINE COMPOUNDS

1253. Propyllithium

[2417-93-8]



$\text{C}_3\text{H}_7\text{Li}$

Leleu, *Cahiers*, 1977, (88), 368

It ignites in air.

See other ALKYL METALS

†1254. 3-Aminopropene (Allylamine)

[107-11-9]



$\text{C}_3\text{H}_7\text{N}$

HCS 1980, 123

See other ALLYL COMPOUNDS

†1255. Azetidine

[503-29-7]



$\text{C}_3\text{H}_7\text{N}$


Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Strained ring heterocycles

See other STRAINED-RING COMPOUNDS

†1256. Cyclopropylamine


[765-30-0]



$\text{C}_3\text{H}_7\text{N}$

†1257. 2-Methylaziridine (Propyleneimine)

[75-55-8]



$\text{C}_3\text{H}_7\text{N}$

Acids


Inlow, R. O. *et al.*, *J. Inorg. Nucl. Chem.*, 1975, **37**, 2353

Like the lower homologue ethyleneimine, it may polymerise explosively if exposed to acids or acidic fumes, so it must always be stored over solid alkali.

See other POLYMERISATION INCIDENTS

1258. Acetone oxime (2-Hydroxyiminopropane)

[127-06-0]



$\text{C}_3\text{H}_7\text{NO}$

Energy of decomposition (in range 200–420°C) measured as 1.5 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other OXIMES

1259. Dimethylformamide

[68-12-2]

C₃H₇NO

Me₂NCO.H

HCS 1980, 430; RSC Lab. Hazard Data Sheet No. 17, 1983

This powerful aprotic solvent is not inert and reacts vigorously or violently with a range of materials.

See *endo*-2,5-Dichloro-7-thiabicyclo[2.2.1]heptane: Dimethylformamide

Diisocyanatomethane: Dimethylformamide

Lithium azide: Alkyl nitrates, etc.

Tetraphosphorus decaoxide: Organic liquids

Sodium hydride: Dimethylformamide

Sodium tetrahydroborate: Dimethylformamide

Sodium: Dimethylformamide

Sulfinyl chloride: Dimethylformamide

Triethylaluminium: Dimethylformamide

2,4,6-Trichloro-1,3,5-triazine: Dimethylformamide

Allyl trifluoromethanesulfonates

See ALLYL TRIFLUOROMETHANESULFONATES: alone, or aprotic solvents

Halocarbons

See Carbon tetrachloride: Dimethylformamide

1,2,3,4,5,6-Hexachlorocyclohexane: Dimethylformamide

See *N,N*-Dimethylacetamide: Halogenated compounds, etc.

See other CATALYTIC IMPURITY INCIDENTS

Other reactants

Yoshida, 1980, 172

MRH values calculated for 13 combinations with oxidants are given.

Oxidants

Pal, B. C. *et al.*, *Chem. Eng. News*, 1981, **59**(17), 47

Explosions involving chromium trioxide or potassium permanganate and dimethylformamide are explained in terms of DMF being, like formic acid, a reducing agent.

Dimethylacetamide is recommended as a safer, less reducing substitute solvent for DMF. For other combinations listed

See Bromine: Dimethylformamide

Chlorine: Dimethylformamide

Chromium trioxide: Dimethylformamide

Magnesium nitrate: Dimethylformamide

Potassium permanganate: Dimethylformamide

Uronium perchlorate: Organic materials

See other APROTIC SOLVENTS

1260. S-Methylthioacetohydroximate

[13749-94-5]

C₃H₇NOS

Preparative hazard

Oren, Z. *et al.*, *Israeli Patent IL 89641*, 1995; *Chem. Abs.*, 1996, **124**, 260401g
It is claimed that a prior method of manufacture from acetonitrile, methanethiol and hydroxylamine hydrochloride gave uncontrolled reaction and explosion. A safer preparation starting from acetaldoxime is claimed.

1261. Ethyl carbamate

[51-79-6]

C₃H₇NO₂

Phosphorus pentachloride

See N-Carbomethoxyiminophosphoryl chloride

†**1262. Isopropyl nitrite**

[541-42-4]

C₃H₇NO₂

See other ALKYL NITRITES

1263. 1-Nitropropane

[108-03-2]

C₃H₇NO₂

Metal oxides

See NITROALKANES: metal oxides

Other reactants

Yoshida, 1980, 276

MRH values calculated for 14 combinations, mainly with oxidants, are given.

See other NITROALKANES

1264. 2-Nitropropane

[79-46-9]

C₃H₇NO₂

HCS 1980, 689; *RSC Lab. Hazard Data Sheet No. 10*, 1983

Carbon, Hopcalite

'Tech. Data Sheet No. 7', New York, Comml. Solvents Corp., 1968

The heat of adsorption of 2-nitropropane is very high, so carbon-containing respirators should not be used in high vapour concentrations. Also, if Hopcalite catalyst (co-precipitated copper(II) oxide and manganese (IV) oxide) is present in the respirator cartridge, ignition may occur.

Amines, Heavy metal oxides

'Tech Data Sheet No. 7', New York, Comml. Solvents Corp., 1968

Contact with amines and mercury oxide or silver oxide may lead to formation of unstable salts analogous to those of nitromethane.

See other NITROALKANES

†1265. Propyl nitrite

[543-67-9]

$C_3H_7NO_2$

PrON:O

Anon., private comm., 1985

Bottles of isopropyl nitrite decomposed during refrigerated storage, developing enough internal pressure to burst the containers and blow open the refrigerator door. Other alkyl nitrites lower than amyl nitrite may be expected to behave similarly.

See other ALKYL NITRITES

†1266. Isopropyl nitrate

[1712-64-7]

$C_3H_7NO_3$

Me₂CHONO₂

1. 'Tech. Bull. No. 2', London, ICI Ltd., 1964
2. Brochet, C., *Astronaut. Acta*, 1970, **15**, 419–425
3. Beeley, P. *et al.*, *Chem. Abs.*, 1977, **87**, 120000
4. Beeley, P. *et al.*, *Comb. Flame*, 1980, **39**, 255–268, 269–281

The self-sustaining exothermic decomposition/combustion characteristics of this fuel of negative oxygen balance (–68%) renders it suitable as a rocket monopropellant. Storage and handling precautions are detailed [1]. Conditions for detonation of the nitrate, an explosive of low sensitivity, have been investigated [2]. The pure vapour ignites spontaneously at very low temperatures and pressures, and other aspects were studied in detail [3], including spontaneous ignition on compression [4].

Lewis acids

See ALKYL NITRATES: lewis acids *See other* ALKYL NITRATES

†1267. Propyl nitrate

[627-13-4]

$C_3H_7NO_3$

PrONO₂

Other reactants

Yoshida, 1980, 204

MRH Values calculated for 11 combinations are given.

Solvents


Breslow, B. A., US Pat. 4 042 032, 1977

The shock-sensitive nitrate is desensitised by 1–2% of propane, butane, chloroform, dimethyl ether or diethyl ether.

See THERMAL EXPLOSIONS See other ALKYL NITRATES

1268. 5-Amino-2-ethyl-2H-tetrazole
[95112-14-4]

C₃H₇N₅



Aluminium hydride

See Aluminium hydride: Tetrazole derivatives

See other TETRAZOLES

1269. Propylsodium
[15790-54-2]

C₃H₇Na

PrNa

Houben-Weyl, 1970, Vol. 13.1, 384

The powder ignites in air.

See other ALKYL METALS

1270. Sodium isopropoxide
[683-60-3]

C₃H₇NaO

NaOCHMe₂

Dimethyl sulfoxide

See Dimethyl sulfoxide: Metal alkoxides

See other METAL ALKOXIDES

†**1271. Propane**
[74-98-6]

C₃H₈

H₃CCH₂CH₃

FPA H10, 1973 (cylinder); HCS 1980, 780 (cylinder)

Barium peroxide

See Barium peroxide: Propane

See also CRYOGENIC LIQUIDS (reference 10), LIQUEFIED PETROLEUM GASES

- 1272. Ethyliodomethylarsine**
 [65313-31-7] **C₃H₈AsI**

$$\text{EtAs(I)Me}$$
 Leleu, *Cahiers*, 1977, (88), 363
 It ignites in air.
See other ALKYLNON-METAL HALIDES
- 1273. Ethyl iminioformate chloride**
 [16694-46-5] **C₃H₈CINO**

$$\text{EtOC(:N}^+\text{H}_2\text{)H Cl}^-$$
 Preparative hazard
See Hydrogen chloride: Alcohols, Hydrogen cyanide
- 1274. Isopropyldiazene**
 [26981-95-3] **C₃H₈N₂**

$$\text{Me}_2\text{CHN=NH}$$
 Chlorine
 Abendroth, H. J., *Angew. Chem.*, 1959, **71**, 340
 The crude chlorination product of 'isoacetone hydrazone' (presumably an *N*-chloro compound) exploded violently during drying at 0°C.
See N-HALOGEN COMPOUNDS *See related* AZO COMPOUNDS
- 1275. 3-Hydrazinopropanenitrile (3-Cyanoethylhydrazine)**
 [353-07-1] **C₃H₇N₃**


$$\text{N}\equiv\text{CCH}_2\text{CH}_2\text{NHNH}_2$$
 Jasiewicz, M. L., *Chem. & Ind.*, 1996, (3), 70
 In the late stage of work up of a sample prepared from acrylonitrile and hydrazine hydrate and stripped of water by a procedure involving dichloromethane, the distillation flask pressurised and burst, shattering the front of the fume cupboard. Previously, slight pressurisation had once been observed. Involving both a nitrile and a hydrazino moiety, this molecule cannot be thermodynamically stable but it has not given previous problems.
See Dichloromethane: 1,2-Diaminoethane, 3-Aminopropionitrile
See CYANO COMPOUNDS
- 1276. *N,N'*-Dinitro-*N*-methyl-1,2-diaminoethane**
 [10308-90-4] **C₃H₈N₄O₄**

$$\text{MeN(NO}_2\text{)C}_2\text{H}_4\text{NHNO}_2$$
 Violent decomposition occurred at 210°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1) See other N-NITRO COMPOUNDS

1277. 2,2-Dinitropropylhydrazine

[]



$\text{C}_3\text{H}_8\text{N}_4\text{O}_4$

Borane

See Borane bis(2,2-dinitropropylhydrazine)

See other POLYNITROALKYL COMPOUNDS

†1278. Ethyl methyl ether (Methoxyethane)

[540-67-0]



$\text{C}_3\text{H}_8\text{O}$

HCS 1980, 498

†1279. Propanol (1-Hydroxypropane)

[71-23-8]



$\text{C}_3\text{H}_8\text{O}$

HCS 1980, 797

Other reactants

Yoshida, 1980, 315

MRH values calculated for 13 combinations with oxidants are given.

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

†1280. 2-Propanol (Isopropanol)

[67-63-0]



$\text{C}_3\text{H}_8\text{O}$

(MCA SD-98, 1972); FPA H45, 1976; HCS 1980, 783

1. Renfrew, M. M., *J. Chem. Educ.*, 1983, **60**(9), A229

2. Bonafede, J. D., *J. Chem. Educ.*, 1984, **61**, 632

3. Redemann, C. E., *J. Amer. Chem. Soc.*, 1942, **64**, 3049

4. Mirafzal, G. A. *et al.*, *J. Chem. Educ.*, 1988, **65**(9), A226–229

5. Bohanon, J. T., *Chem. Eng. News*, 1989, **67**(1), 4

Several explosions have occurred during laboratory distillation of isopropanol [1,2,4,5], some with a sample stored for 5 years in a part empty can [5]. No cause was apparent, but presence of traces of ketone(s) promoting peroxidation is a probability. Previously, the presence of 0.36 M peroxide had been reported

C₃H₈O

in a 99.5% pure sample of isopropanol stored for several months in a partially full clear glass bottle in strong daylight [3]. The reformation of peroxides in deperoxidised isopropanol 'within a few days' had been noted [2]. It appears that the tertiary H on the 2-position is susceptible to autoxidation, and that 2-propanol must be classed as peroxidisable. 2-Hydroperoxy-2-hydroxypropane has, in fact, been isolated from photocatalysed oxidation of isopropanol.

See PEROXIDISABLE COMPOUNDS (reference 7)

See 2-Butanone, below; *also* Oxygen (Gas): Alcohols

Aluminium

See Aluminium: 2-Propanol

Aluminium isopropoxide, Crotonaldehyde

Wagner-Jauregg, T., *Angew. Chem.*, 1939, **53**, 710

During distillation of 2-propanol recovered from the reduction of crotonaldehyde with isopropanol/aluminium isopropoxide, a violent explosion occurred. This was attributed to peroxidised diisopropyl ether (a possible by-product) or to peroxidised crotonaldehyde. An alternative or additional possibility is that the isopropanol may have contained traces of a higher secondary alcohol (e.g. 2-butanol) which would be oxidised during the Meerwein-Ponndorf reduction procedure to 2-butanone. The latter would then effectively sensitise the isopropanol or other peroxidisable species to peroxidation.

See 2-Butanone, next below; *also* Diisopropyl ether

2-Butanone

1. Bathie, H. M., *Chem. Brit.*, 1974, **10**, 143

2. Unpublished observations, 1974

3. Pitt, M. J., *Chem. Brit.*, 1974, **10**, 312

4. Schenk, G. O. *et al.*, *Angew. Chem.*, 1958, **70**, 504

Distillation to small volume of a small sample of a 4-year-old mixture of the alcohol with 0.5% of the ketone led to a violent explosion, and the presence of peroxides was subsequently confirmed [1]. Pure alcohols which can form stable radicals (secondary branched structures) may slowly peroxidise to a limited extent under normal storage conditions (isopropanol to 0.0015 M in brown bottle, subdued light during 6 months; to 0.0009 M in dark during 5 years) [2]. The presence of ketones markedly increases the possibility of peroxidation by sensitising photochemical oxidation of the alcohol. Acetone (produced during autoxidation of isopropanol) is not a good sensitiser, but the presence of even traces of 2-butanone in isopropanol would be expected to accelerate markedly peroxidation of the latter. Treatment of any mixture or old sample of a secondary alcohol with tin(II) chloride and then lime before distillation is recommended [3]. The product of photosensitised oxidation is 2-hydroperoxy-2-propanol [4].

See 2-Butanol

Hydrogen peroxide

Baratov, A. N. *et al.*, *Chem. Abs.*, 1978, **88**, 39537

Addition of a small amount of hydrogen peroxide may reduce sharply the AIT of the alcohol (455°C), probably to that of the hydroperoxide.

Other reactants

Yoshida, 198, 41

MRH values calculated for 13 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Barium perchlorate: Alcohols

Chromium trioxide: Alcohols

MRH 2.47/91

Dioxygenyl tetrafluoroborate: Organic materials

Hydrogen peroxide: Oxygenated compounds

MRH 6.32/84

Oxygen (Gas): Alcohols

Sodium dichromate: 2-Propanol, etc.


Trinitromethane: 2-Propanol

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

†1281. Dimethoxymethane

[109-87-5]



HCS 1980, 419

Oxygen

Molem, M. J. *et al.*, *Chem. Abs.*, 1975, **83**, 45495

The nature of the reaction and the products in methylal–oxygen mixtures during cool flame or explosive oxidation were studied.

See other PEROXIDISABLE COMPOUNDS

1282. Ethyl methyl peroxide

[70299-48-8]



Rieche, A., *Ber.*, 1929, **62**, 218

Shock-sensitive as a liquid or vapour, it explodes violently on superheating.

See other DIALKYL PEROXIDES

1283. Isopropyl hydroperoxide (2-Hydroperoxypropane)

[3031-75-2]



Medvedev, S. *et al.*, *Ber.*, 1932, **65**, 133

It explodes just above the b.p., 107–109°C.

See other ALKYL HYDROPEROXIDES

1284. 2-Methoxyethanol (Ethylene glycol monomethyl ether)
[109-86-4]

C₃H₈O₂



HCS 1980, 799

491M, 1975, 256

It forms explosive peroxides.

See other PEROXIDISABLE COMPOUNDS

1285. Propylene glycol (1,2-Propanediol)
[57-55-6]

C₃H₈O₂



HCS 1980, 782

Hydrofluoric acid, Nitric acid, Silver nitrate

See Nitric acid: Hydrofluoric acid, Propylene glycol, etc.

1286. Glycerol (1,2,3-Propanetriol)
[56-81-5]

C₃H₈O₃



HCS 1980, 516

Ethylene oxide

See Ethylene oxide: Glycerol

Other reactants

Yoshida 1980, 448

MRH values calculated for 16 combinations, mainly with oxidants, are given.

Oxidants

MRH values show % of oxidant

Unpublished comments, 1978

The violent or explosive reactions exhibited by glycerol in contact with many solid oxidants are due to its unique properties of having three centres of reactivity, of being a liquid which ensures good contact, and of high boiling point and viscosity which prevents dissipation of oxidative heat. The difunctional, less viscous liquid glycols show similar but less extreme behaviour.

See Calcium hypochlorite: Hydroxy compounds

MRH 2.13/88

Chlorine: Glycerol

Hydrogen peroxide: Organic compounds (reference 2)

MRH 5.56/71

Nitric acid: Glycerol

Nitric acid: Glycerol, Hydrochloric acid

Nitric acid: Glycerol, Hydrofluoric acid

†1289. Propanethiol
[107-03-9] C₃H₈S

PrSH

Calcium hypochlorite

See Calcium hypochlorite: Organic sulfur compounds

See other ALKANETHIOLS

†1290. 2-Propanethiol
[75-33-2] C₃H₈S

Me₂CHSH

See other ALKANETHIOLS

1291. Trimethylaluminium
[75-24-1] C₃H₉Al

Me₃Al

HCS 1980, 938

Kirk-Othmer 1963, Vol. 2, 40

Extremely pyrophoric, ignition delay 13 ms in air at 235°C/0.16 bar.

Dichlorodi- μ -chlorobis(pentamethylcyclopentadienyl)dirhodium

(Structure III, p. S-2)

See Dichlorodi- μ -chlorobis(pentamethylcyclopentadienyl)dirhodium: Air, Alkyl-
metals

Other reactants

Yoshida, 1980, 257

MRH values calculated for 15 combinations, largely with oxidants, are given.

Tetrafluorobenzene-1,4-diol

Spence, R., *Chem. Eng. News*, 1996, **74**(21), 4

The dried product of reaction of approximately equimolar quantities of the diol
and methyl aluminium exploded when crushed with a spatula.

See METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

See entry TRIALKYLALUMINIUMS

See other ALKYLMETALS

1292. Trimethyldialuminium trichloride (Aluminium chloride-trimethyl-
aluminium complex)
[12542-85-7] C₃H₉Al₂Cl₃

Me₃Al·AlCl₃

HCS 1980, 636

See entry ALKYLALUMINIUM HALIDES

1293. Ethylmethylarsine
[689-93-0] **C₃H₉As**

EtAsHMe

Leleu, *Cahiers*, 1977, (88), 362
It ignites in air.
See other ALKYLNON-METAL HYDRIDES

1294. Trimethylarsine
[593-88-4] **C₃H₉As**

Me₃As

Air, or Halogens
Sidgwick, 1950, 762, 769
It inflames in air, and interaction with halogens is violent.
See other ALKYLNON-METALS

1295. Trimethylborane
[593-90-8] **C₃H₉B**

Me₃B

Stock, A. *et al.*, *Ber.*, 1921, **54**, 535
The gas ignites in air.
See Chlorine: Trialkylboranes
See other ALKYLBORANES

†**1296. Trimethyl borate**
[121-43-7] **C₃H₉BO₃**

(MeO)₃B

1297. 2-(Tricarbonylferra)hexaborane(10)
[75952-58-8] **C₃H₉B₅FeO₃**

(Complex Structure)

Preparative hazard
See Pentaborane(9): Pentacarbonyliron, Pyrex glass

1298. Trimethylbismuthine
[593-91-9] **C₃H₉Bi**

Me₃Bi

1. Coates, 1967, Vol. 1, 536
2. Sorbe, 1968, 157

It ignites in air, but is unreactive with water [1], and explodes at 110°C [2].
See TRIALKYLBISMUTHS See other ALKYL METALS

1299. Trimethylsulfoxonium bromide
[25596-24-1]

C₃H₉BrOS



1. Scaros, M. G. *et al.*, *Chem. Brit.*, 1973, **9**, 523; private comm., 1974
2. Scaros, M. G. *et al.*, *Proc. 7th Conf. Catal. Org. Synth.* 1978, 1980, 301–309
3. Dryden, H. L. *et al.*, US Pat. 4 141 920, 1979

Preparation of the title compound by interaction of dimethyl sulfoxide and bromomethane, sealed into a resin-coated glass bottle and heated at 65°C, led to an explosion after 120 h. The isolated salt thermally decomposes above 180°C to produce formaldehyde and a residue of methanesulfonic acid. In solution in dimethyl sulfoxide, the salt begins to decompose after several hours at 74–80°C (after exposure to light), the exothermic reaction accelerating with vigorous evolution of vapour (including formaldehyde and dimethyl sulfide), the residue of methanesulfonic acid finally attaining 132°C. Some white solid, probably poly-formaldehyde, is also produced. The explosion seems likely to have been a pressure-burst of the container under excessive internal pressure from the decomposition products. A safe procedure has been developed in which maximum reaction temperatures and times of 65°C and 55 h are used. Once prepared, the salt should not be redissolved in dimethyl sulfoxide. In contrast, trimethylsulfoxonium iodide appears simply to dissolve on heating rather than decomposing like the bromide [1]. The decomposition mechanism involves isomerisation of some of the product to dimethylmethoxysulfonium bromide, which then degrades to formaldehyde, dimethyl sulfide and hydrogen bromide, and the latter catalyses further decomposition reactions [2]. Use of orthoformate or orthocarbonate esters as bromine and hydrogen bromide scavengers, and a cooling/pressure venting system avoids the possibility of exothermic decomposition of the reaction mixture [3].

See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS

1300. N-Bromotrimethylammonium bromide(?)

[]

C₃H₉Br₂N



See Bromine: Trimethylamine

See other N-HALOGEN COMPOUNDS

1301. Trimethylsilyl chlorochromate
[102488-24-4]

C₃H₉ClCrO₃Si



Aizpurna, J. M. *et al.*, *Tetrahedron Lett.*, 1983, **24**, 4367

The oxidant appears safe when prepared and used in dichloromethane solution, but attempted isolation by distillation led to a violent explosion.

See other OXIDANTS

1302. Trimethylsilyl perchlorate

[18204-79-0]

$C_3H_9ClO_4Si$

$Me_3SiOCIO_3$

See entry ORGANOSILYL PERCHLORATES *See other* NON-METAL PERCHLORATES

1303. Trimethylsulfonium chloride

[3086-29-1]

C_3H_9ClS

$Me_3S^+Cl^-$

Preparative hazard

Byrne, B. *et al.*, *Tetrahedron Lett.*, 1986, **27**, 1233–1236

During its preparation by heating dimethyl sulfide and methyl chloroformate in a glass pressure bottle at 80°C, it is essential to interrupt the reaction after 4 h, cool and release the internal pressure of carbon dioxide, before recapping the bottle and heating for a further 22 h. This avoids the possibility of excessive pressure build-up and failure of the cap seal.

See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS

†**1304. Chlorotrimethylsilane**

[75-77-4]

C_3H_9ClSi

$ClSiMe_3$

HCS 1980, 941

Preparative hazard

See Chlorine: Antimony trichloride, Tetramethylsilane

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Water

Leleu, *Cahiers*, 1977, (88), 370

It reacts violently with water.

See other ALKYLHALOSILANES

1305. Trimethylgallium

[1445-79-0]

C_3H_9Ga

Me_3Ga

Air, or Water

Sidgwick, 1950, 461

It ignites in air, even at -76°C , and reacts violently with water.
See other ALKYL METALS

†1306. Iodotrimethylsilane
[16029-98-4]



$\text{C}_3\text{H}_9\text{ISi}$

See other ALKYLHALOSILANES, IODINE COMPOUNDS

1307. Trimethylindium
[3385-78-2]



$\text{C}_3\text{H}_9\text{In}$

1. Bailar, 1973, Vol. 1, 1116

2. Houben-Weyl, 1970, Vol. 13.4, 351

Trimethylindium and other lower-alkyl derivatives ignite in air [1], including the 2:1 complex with diethyl ether [2].

See other ALKYL METALS

†1308. Isopropylamine (2-Propanamine)
[75-31-0]



$\text{C}_3\text{H}_9\text{N}$

FPA H72, 1978; *HCS 1980*, 577

1-Chloro-2,3-epoxypropane

See 1-Chloro-2,3-epoxypropane: Isopropylamine

Perchloryl fluoride

See Perchloryl fluoride: Nitrogenous bases

See other ORGANIC BASES

†1309. Propylamine (Propanamine)
[107-10-8]



$\text{C}_3\text{H}_9\text{N}$

HCS 1980, 796

Triethynylaluminium

See Triethynylaluminium: Diethyl ether, etc.

See other ORGANIC BASES

†1310. Trimethylamine (*N,N*-Dimethylmethanamine)
[75-50-3]



$\text{C}_3\text{H}_9\text{N}$

(*MCA SD-57*, 1955); *HCS 1980*, 939 (anhydrous), 940 (40% solution)

Bromine

See Bromine: Trimethylamine

Ethylene oxide

MRH 3.72/100

See Ethylene oxide: Trimethylamine

Other reactants

Yoshida, 1980, 256

MRH values calculated for 14 combinations, largely with oxidants, are given.

See other ORGANIC BASES

1311. 1-Amino-2-propanol

[78-96-6]

C₃H₉NO



Cellulose nitrate

See CELLULOSE NITRATE: amines

2,4-Hexadienal

See 2,4-Hexadienal: 1-Amino-2-propanol

See other ORGANIC BASES

†1312. 2-Methoxyethylamine

[109-85-3]

C₃H₉NO



Energy of decomposition (in range 140–200°C) measured as 0.23 kJ/g.


See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other AMINOMETHOXY COMPOUNDS

1313. Trimethylamine oxide

[1184-78-7]

C₃H₉NO



1. *Inorg. Synth.*, 1989, **25**, 107

2. Ringel, C., *Z. Chem.*, 1969, **9**, 188

3. Editor's comments

Energy of decomposition (in range 160–280°C) measured as 1.455 kJ/g.

The assessors experienced an explosion while drying the oxide in ethyl ether. Rather drastic precautions are recommended in handling it [1]. A preparation, allowed to stand for a week rather than the day specified, exploded during concentration [2]. Amine oxides from the standard preparation are inclined to retain hydrogen peroxide of 'hydration' unless it is destroyed during work-up. The peroxide (or diperoxidate) of dimethylamine oxide would be expected to be far more dangerous than the oxide itself [3].

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other N-OXIDES

†1314. Trimethylsilyl azide (Azidotrimethylsilane)
[4648-54-8]

$C_3H_9N_3Si$

Me_3SiN_3

1. Sundermeyer, W., *Chem. Ber.*, 1963, **96**, 1293
2. Thayer, J. S. *et al.*, *Inorg. Chem.*, 1964, **3**, 889
3. West, R. *et al.*, *Chem. Eng. News*, 1984, **62**(24), 4

A standard literature method [1,2] had been used frequently and uneventfully to prepare the azide from trimethylsilyl chloride and sodium azide in presence of aluminium chloride as catalyst. A recent duplication led to a violent detonation during distillation of the product, and this was attributed to carry-over of traces of aluminium azides into the distillation flask. Precautions are detailed [3].

Rhenium hexafluoride

1. Fawcett, J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 958
2. Fawcett, J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1987, 567–571

Interaction at $-60^\circ C$ in 1,1,2-trichloroethane to give rhenium nitride tetrafluoride is violent and an explosion is possible even with repeated cooling with liquid nitrogen. The final warming to $0^\circ C$ is also hazardous [1]. If the reactants are mixed at $-196^\circ C$, at any stage of warming, the reaction can become extremely vigorous and detonate, even with maximum cooling control [2].

Selenium halides

Kennett, F. A. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 851–857

In attempted preparation of poly(selenium nitride), the black solid formed by interaction with selenium tetrabromide in acetonitrile exploded violently within 1 min at $0^\circ C$. The solid produced from diselenium tetrachloride in acetonitrile exploded at around $100^\circ C$, and in dichloromethane the product exploded in contact with a nickel spatula.

Tungsten hexafluoride

See Azidopentafluorotungsten

See related NON-METAL AZIDES

1315. Trimethyl phosphite
[121-45-9]

$C_3H_9O_3P$

$(MeO)_3P$

Magnesium perchlorate

See Magnesium perchlorate: Trimethyl phosphite

Trimethylplatinum(IV) azide tetramer

See Dodecamethyltetraplatinum(IV)azide: Trimethyl phosphite

See other PHOSPHORUS ESTERS

- 1316. Trimethyl thiophosphate (Trimethyl phosphorothioate)**
[152-18-1] **C₃H₉O₃PS**
(MeO)₃P(:S)
- Chlorine
See Chlorine: Trimethyl thionophosphate
See other PHOSPHORUS ESTERS
- 1317. Titanium(III) methoxide**
[7245-18-3] **C₃H₉O₃Ti**
Ti(OMe)₃
- Bailar, 1973, Vol. 3, 384
A pyrophoric solid.
See other METAL ALKOXIDES
- 1318. Trimethyl phosphate**
[512-56-1] **C₃H₉O₄P**
(MeO)₃P(:O)
- Anon., *ABCM Quart. Safety Summ.*, 1953, **25**, 3
The residue from a large scale atmospheric pressure distillation of trimethyl phosphate exploded violently. This was attributed to rapid decomposition of the ester, catalysed by the acidic degradation products, with evolution of gaseous hydrocarbons. It is recommended that only small batches of alkyl phosphates should be vacuum distilled and in presence of magnesium oxide to neutralise any acid by-products, and to suppress the acid catalysed reaction.
See other CATALYTIC IMPURITY INCIDENTS, PHOSPHORUS ESTERS
- 1319. Trimethylphosphine**
[594-09-2] **C₃H₉P**
Me₃P
1. Personal experience
2. Labinger, J. A. *et al.*, *Inorg. Synth.*, 1978, **18**, 63
It may ignite in air [1]. It is readily regenerated by heating its air-stable complex with silver iodide [2], which became commercially available in 1983.
See other ALKYLPHOSPHINES
- 1320. Trimethylantimony (Trimethylstibine)**
[594-10-5] **C₃H₉Sb**
Me₃Sb
- Von Schwartz, 1918, 322
It ignites in air.
- Halogens
Sidgwick, 1950, 777
Interaction is violent.

2-Iodoethanol

See 2-Hydroxyethyltrimethylstibonium iodide

See other ALKYL METALS

1321. Trimethylthallium

[3003-15-4]

C_3H_9Tl

Me_3Tl

Sidgwick, 1950, 463

It ignites in air, and is liable to explode violently above 90°C.

Diethyl ether

Leleu, *Cahiers*, 1977, (88), 371

The complex with ether explodes at 0°C.

See other ALKYL METALS

1322. Trimethylammonium perchlorate

[15576-35-9]

$C_3H_{10}ClNO_4$

$Me_3N^+H ClO_4^-$

Jain, S. R. *et al.*, *Chem. Abs.*, 1982, **96**, 220110

The salt burns more readily as a propellant than does ammonium perchlorate.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

†1323. Trimethylhydroxylammonium perchlorate (Trimethylamine oxide perchlorate)

[22755-36-8]

$C_3H_{10}ClNO_5$

$Me_3N^+OH ClO_4^-$

Hofmann, K. A. *et al.*, *Ber.*, 1910, **43**, 2624


The salt, formed from trimethylamine oxide and perchloric acid, explodes on heating, or under a hammer blow.

See other PERCHLORATE SALTS OF NITROGENOUS BASES, REDOX COMPOUNDS

1324. Dihydrzinecadmium(II) malonate (Bis(hydrazine)[propanedioato(2-)] cadmium)

[159793-64-3]

$C_3H_{10}CdN_4O_4$



See HYDRAZINE METAL MALONATES AND SUCCINATES

†1325. 1,2-Diaminopropane (1,2-Propanediamine)
[78-90-0] C₃H₁₀N₂



HCS 1980, 798

See other ORGANIC BASES


†1326. 1,3-Diaminopropane (1,3-Propanediamine)
[109-76-2] C₃H₁₀N₂



HCS 1980, 361

See other ORGANIC BASES


1327. *S*-Ethylisothiuronium hydrogen sulfate (Ethyl carbamimonothioate
hydrogen sulfate)
[22722-03-8] C₃H₁₀N₂O₄S₂



Chlorine

See Chlorine: Nitrogen compounds (reference 4)

1328. Dihydrzinenickel(II) malonate (Bis(hydrazine)[propanedioato(2-)]nickel)
[159793-62-1] C₃H₁₀N₄NiO₄



See HYDRAZINE METAL MALONATES AND SUCCINATES

1329. Trimethylplatinum hydroxide
[14477-33-9] C₃H₁₀OPt



Höchstetter, M. N. *et al.*, *Inorg. Chem.*, 1969, **8**, 400

The compound (originally reported as tetramethylplatinum) and usually obtained in admixture with an alkoxy derivative, detonates on heating.

See other PLATINUM COMPOUNDS

1330. Trimethylsilyl hydroperoxide
[18230-75-6]



Buncel, E. *et al.*, *J. Chem. Soc.*, 1958, 1551

It decomposes rapidly above 35°C and may have been involved in an explosion during distillation of the corresponding peroxide, which is stable to 135°C.

See related ORGANOMETALLIC PEROXIDES

1331. Propylsilane
[13154-66-0]




Leleu, *Cahiers*, 1977, (88), 369

It ignites in air.

See other ALKYL SILANES

1332. Trimethylgermylphosphine
[20519-92-0]




Oxygen

Dahl, A. R. *et al.*, *Inorg. Chem.*, 1975, **14**, 1093

A solution in chloroform at -45°C reacts rapidly with oxygen, and at oxygen pressures above 25 mbar combustion occurs. Dimethyldiphosphinogermane behaves similarly.

See related ALKYL METALS, PHOSPHINES

1333. Di(hydroperoxy)trimethylantimony(V)
[]




Dodd, M. *et al.*, *Appl. Organomet. Chem.*, 1992, **6**(2), 207

This compound, prepared from trimethylstibine and hydrogen peroxide, exploded violently when heated under vacuum.

See other ORGANOMETALLIC PEROXIDES

1334. Trimethyldiborane
[21107-27-7]




Leleu, *Cahiers*, 1977, (88), 366

It ignites in air.

See other ALKYL BORANES

1335. 2,4,6-Trimethylborazine (B-Trimethylborazine)
[5314-85-2]





Nitryl chloride

See Nitryl chloride: B-Trimethylborazine

See related BORANES

1336. Aqua-1,2-diaminopropanediperoxochromium(IV) dihydrate
[17185-68-1]




House, D. A. *et al.*, *Inorg. Chem.*, 1966, **5**, 40

Several preparations of the dihydrate exploded spontaneously at 20–25°C, and the aqua complex explodes at 88–90°C during slow heating.

See other AMMINECHROMIUM PEROXOCOMPLEXES

1337. Beryllium tetrahydroborate–trimethylamine
[]



Air, or Water

Burg, A. B. *et al.*, *J. Amer. Chem. Soc.*, 1940, **62**, 3427

The complex ignites in contact with air or water.

See other COMPLEX HYDRIDES


1338. Cadmium hydrazinium tris(hydrazinecarboxylate) trihydrate
[]



Dhas, N. A. *et al.*, *Chem. Abs.*, 1994, **121**, 314276d

The hydrazinecarboxylate decomposed explosively on heating to 120°C.

1339. Triiodocyclopropenium iodide
[99796-78-8]



Weiss, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1986, **25**, 103


This first example of an ionic halocarbon is extremely explosive, not only when dry, but also upon dissolution in polar solvents (ethanol, acetonitrile). Only 1 mmol

of the salt should be prepared in a single experiment, and only samples moist with dichloromethane should be used or handled.

See HALOCARBONS

See other IODINE COMPOUNDS

1340. Dipotassium diazirine-3,3-dicarboxylate
[76429-97-5]



See Diazirine-3,3-dicarboxylic acid

See other DIAZIRINES

1341. Oxopropanedinitrile (Carbonyl dicyanide)
[1115-12-4]




Water

Martin, E. L., *Org. Synth.*, 1971, **51**, 70

The nitrile reacts explosively with water.

See related ACYL HALIDES See other CYANO COMPOUNDS

1342. Carbonyl diisothiocyanate
[6470-09-3]



Dimethyl sulfoxide

See Dimethyl sulfoxide: Carbonyl diisothiocyanate

See related CYANO COMPOUNDS

1343. Phosphorus tricyanide
[1116-01-4]



1. Absolom, R. *et al.*, *Chem. & Ind.*, 1967, 1593

2. Smith, T. D. *et al.*, *Chem. & Ind.*, 1967, 1969

3. Mellor, 1971, Vol. 8, Suppl. 3, 583

During vacuum sublimation at around 100°C, explosions occurred on 3 occasions. These were attributed to slight leakage of air into the sublimer. The use of joints remote from the heating bath, a high-melting grease, and a cooling liquid other than water (to avoid rapid evolution of hydrogen cyanide in the event of breakage) are recommended, as well as working in a fume cupboard [1]. However, the later publication suggests that the silver cyanide used in the preparation could be the cause of the explosions. On long keeping, it discolours and may then contain the nitride (fulminating silver). Silver cyanide precipitated from slightly acidic solution and stored in a dark bottle is suitable [2]. Phosphorus tricyanide ignites in air if touched with a warm glass rod, and reacts violently with water.


See related NON-METAL HALIDES AND THEIR OXIDES

See other CYANO COMPOUNDS

1344. Diazomalononitrile (Diazodicyanomethane)

[1618-08-2]

C₃N₄



Ciganek, E., *J. Org. Chem.*, 1965, **30**, 4200

Small (mg) quantities melt at 75°C, but larger amounts explode. It also has borderline sensitivity towards static electricity and must be handled with full precautions.

See other CYANO COMPOUNDS, DIAZO COMPOUNDS

1345. 5-Cyano-4-diazo-4H-1,2,3-triazole

[16968-06-2]

C₃N₆

The crystalline solid explodes at the m.p., 125–6°C.


See entry DIAZAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

1346. Cyanodiazoacetyl azide

[115057-40-4]

C₃N₆O



Kobolov, M. Yu. *et al.*, *Chem. Abs.*, 1988, **109**, 22908

It explodes on melting.

See other ACYL AZIDES, CYANO COMPOUNDS, DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

1347. Diazidomalononitrile (Diazidodicyanomethane)
[67880-21-1]

C_3N_8




MCA Case History No. 820

An ethereal solution of some 100 g of the crude nitrile was allowed to spontaneously evaporate and crystallise. The crystalline slurry so produced exploded violently without warning. Previously such material had been found not to be shock-sensitive to hammer blows, but dry recrystallised material was very shock-sensitive. Traces of free hydrogen azide could have been present, and a metal spatula had been used to stir the slurry, so metal azides could have been formed. See other CYANO COMPOUNDS, ORGANIC AZIDES

1348. 2,4,6-Triazido-1,3,5-triazine
[5637-83-2]


C_3N_{12}



1. Ott, *Ber.*, 1921, **54**, 183
 2. Simmonds, R. J. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1982, 1824
- This polyazide (82.3% N) explodes on impact, shock or rapid heating to 170–180°C [1]. When preparing 2,4-diazido-6-dimethylamino-1,3,5-triazine from the 2,4-dichloro compound and sodium azide, there is the possibility of forming the triazido derivative, which detonates violently when touched. Reactions involving organic chlorides and excess sodium azide are extremely dangerous [2].
- See other HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

†1349. Propadiene-1,3-dione ('Carbon suboxide')
[504-64-3]


C_3O_2



See related ACID ANHYDRIDES

1350. Propadienedithione (Carbon subsulfide)

[627-34-9]

C₃S₂

Mellor, 1940, Vol. 6, 88

A red oil decomposing to a black polymer, explosively if heated rapidly to 100–120°C. Probably highly endothermic.

See other ENDOTHERMIC COMPOUNDS, NON-METAL SULFIDES

1351. 1,4-Dibromo-1,3-butadiyne

[36333-41-2]

C₄Br₂Petterson, R. C. *et al.*, *Chem. Eng. News*, 1977, **55**(48), 36

An explosion temperature has been published, but a solvent-free sample standing unheated under nitrogen exploded after 1 h.

See other HALOACETYLENE DERIVATIVES

1352. Heptafluorobutyryl hypochlorite

[71359-62-1]

C₄ClF₇O₂Tari, I. *et al.*, *Inorg. Chem.*, 1979, **18**, 3205–3208

It is thermally unstable above 22°C and explosive in the gas phase at pressures above 27–62 mbar.

See entry ACYL HYPOHALITES (reference 1)

1353. 1,4-Dichloro-1,3-butadiyne

[51104-87-1]

C₄Cl₂

Sorbe, 1968, 50

It explodes above 70°C.

See other HALOACETYLENE DERIVATIVES

1354. Copper(I) chloroacetylide

[]


C₄Cl₂Cu₂

Kirk-Othmer, 1964, Vol. 5, 204

Explosive.

See other HALOACETYLENE DERIVATIVES, HEAVY METAL DERIVATIVES

1355. 4,5-Dichloro-3,3,4,5,6,6-hexafluoro-1,2-dioxane
[]




Haszeldine, R. N. *et al.*, *J. Chem. Soc.*, 1959, 1089

Formed in the controlled oxidation of chlorotrifluoroethylene, it explodes violently on heating.

See other CYCLIC PEROXIDES

1356. Mercury bis(chloroacetylide)
[64771-59-1]





Whitmore, 1921, 119

It explodes fairly violently above the m.p., 185°C.

See other HALOACETYLENE DERIVATIVES, MERCURY COMPOUNDS, METAL ACETYLIDES

1357. 3,4-Dichloro-2,5-dilithiothiophene
[29202-04-8]



Gilman, H., private comm., 1971

The dry solid is explosive, though relatively insensitive to shock.

See other ORGANOLITHIUM REAGENTS

1358. Tetracarbonylmolybdenum dichloride
[15712-13-7]




Newton, W. E. *et al.*, *Inorg. Synth.*, 1978, **18**, 54

If stored at ambient temperature, the complex decomposes to a dark pyrophoric powder.

See related CARBONYLMETALS

1359. Dichloromaleic anhydride
[1122-17-4]




Sodium chloride, Urea

Bott, D. C., private comm., 1980

To prepare dichloromaleimide, the 3 components are melted together with stirring. At 118°C the vigorous exothermic reaction sets in, and rapid ice cooling must then be applied to prevent explosion.

See other ACID ANHYDRIDES

1360. 1,1,4,4-Tetrachlorobutatriene
[19792-18-8]



Heinrich, B. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1968, **7**, 375

One of the by-products formed by heating this compound is an explosive polymer.

See other HALOALKENES *See related* DIENES

1361. Bis(trichloroacetyl) peroxide
[2629-78-9]



1. Swern, 1971, Vol. 2, 815

2. Miller, W. T., *Chem. Abs.*, 1952, **46**, 7889e

Pure material explodes on standing at ambient temperature [1], but the very shock-sensitive solid may be stored safely in trichlorofluoromethane solution at -20°C [2].

See other DIACYL PEROXIDES

1362. Tetracarbon monofluoride
[12774-81-1]



Bailar, 1973, Vol. 1, 1271

Though generally inert, rapid heating causes deflagration.

See related HALOALKANES, NON-METAL HALIDES

- 1363. 1,1,4,4-Tetrafluorobutatriene**
 [2252-95-1] C₄F₄

$$\text{F}_2\text{C}=\text{C}=\text{C}=\text{CF}_2$$
 Martin, E. L. *et al.*, *J. Amer. Chem. Soc.*, 1959, **81**, 5256
 In the liquid state, it explodes above -5°C .
See other HALOALKENES *See related* DIENES
- 1364. Perfluorobutadiene**
 [87-68-3] C₄F₆

$$\text{F}_2\text{C}=\text{CFCF}=\text{CF}_2$$
 Feast, W. J., personal communication, 1993.
 Attempts to use this material as a monomer are not recommended as reaction is uncontrollable and it is liable to explode.
 Bromine perchlorate
See Bromine perchlorate: Perfluorobutadiene
See other DIENES, HALOALKENES
- 1365. Bis(trifluoroacetoxy)xenon**
 [21961-22-8] C₄F₆O₄Xe

$$\text{Xe}(\text{OOC}\text{CF}_3)_2$$
 Kirk Othmer, 1980, Vol. 12, 293
 Detonates easily and should be handled with extreme care.
See other XENON COMPOUNDS
- 1366. Trifluoroacetic anhydride**
 [407-25-0] C₄F₆O₃

$$\text{F}_3\text{CCO}\cdot\text{OCO}\cdot\text{CF}_3$$
 Dimethyl sulfoxide
See Dimethyl sulfoxide: Trifluoroacetic anhydride
 Nitric acid, 1,3,5-Triacetylhexahydro-1,3,5-triazine
See Nitric acid: 1,3,5-Triacetylhexahydrotriazine
 Nitric acid, 1,3,5-Triazine
See Nitric acid: 1,3,5-Triazine, etc.
See other ACID ANHYDRIDES
- 1367. Bis(trifluoroacetyl) peroxide**
 [383-73-3] C₄F₆O₄

$$\text{F}_3\text{CCO}\cdot\text{OOCO}\cdot\text{CF}_3$$
 Swern, 1971, Vol. 2, 815


Pure material explodes on standing at ambient temperature.
See other DIACYL PEROXIDES

1368. Heptafluorobutyryl nitrite
[663-25-2]




Banks, R. E. *et al.*, *J. Chem. Soc. (C)*, 1966, 1351
This may be explosive like its lower homologues, and suitable precautions are necessary during heating or distillation.
See other ACYL NITRITES

1369. Heptafluorobutyryl hypofluorite
[]



Cady, G. H. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 2020
The vapour slowly decomposes at ambient temperature, but will decompose explosively on spark initiation.
See other ACYL HYPOHALITES


1370. Perfluoro-*tert*-nitrosobutane (Tris(trifluoromethyl)nitrosomethane)
[354-93-8]



Nitrogen oxides

Sterlin, S. R. *et al.*, Russ. Pat. 482 432, 1975
The danger of explosions during oxidation of the nitroso compound to perfluoro-*tert*-butanol with nitrogen oxides and subsequent hydrolysis, was reduced by working in a flow system at 160–210°C with 8–10% of nitrogen oxides in air, and using conc. sulfuric acid for hydrolysis.
See other NITROSO COMPOUNDS

1371. Decafluorobutyramidine
[41409-50-1]



Ross, D. L. *et al.*, *J. Org. Chem.*, 1970, **35**, 3096–3097
A shock-sensitive explosive
See other *N, N, N''*-TRIFLUOROAMIDINES

1372. Decafluoro-2,5-diazahehexane 2,5-dioxyl
[36525-64-1] **C₄F₁₀N₂O₂**
F₃CN(O)C₂F₄N(O)CF₃

Preparative hazard

Banks, R. E. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1974, 2535–2536

During an increased-scale preparation of the dioxyl by permanganate oxidation of the hydrolysate of a nitrosotrifluoromethane–tetrafluoroethylene–phosphorus trichloride adduct, an impurity in the dioxyl, trapped out at -96°C (? -196°C) and <2.5 mbar, caused a violent explosion to occur when the trap content was allowed to warm up. A procedure to eliminate the hazard is detailed.

See related N-OXIDES

1373. Perfluoro-tert-butyl peroxyhypofluorite
[66793-67-7] **C₄F₁₀O₂**
(F₃C)₃COOF

Yu, S.-L. *et al.*, *Inorg. Chem.*, 1978, **17**, 2485

It explodes when warmed to 22°C in a sealed tube with a liquid phase present, but is apparently stable in the gas phase alone.

See related HYPOHALITES See other ORGANIC PEROXIDES

1374. 1,1,4,4-Tetrakis(fluoroxy)hexafluorobutane
[22410-18-0] **C₄F₁₀O₄**
(FO)₂CFC₂F₄CF(OF)₂

Sekiya, A. *et al.*, *Inorg. Chem.*, 1980, **19**, 1329

It explodes energetically at -20°C .

See other BIS(FLUOROOXY)PERHALOALKANES

1375. Di[bis(trifluoromethyl)phosphido]mercury
[] **C₄F₁₂HgP₂**
[(F₃C)₂P]₂Hg

Grobe, J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1972, **11**, 1098

It ignites in air.

See related ALKYLPHOSPHINES See other MERCURY COMPOUNDS

1376. Sodium tetracarbonylferrate(2-)
[14878-31-0] **C₄FeNa₂O₄**
Na₂[Fe(CO)₄]

1. Collman, J. P., *Accts. Chem. Res.*, 1975, **8**, 343

2. Finke, R. G. *et al.*, *Org. Synth.*, 1979, **59**, 102–112


It is extremely oxygen-sensitive and ignites in air [1]. The salt solvated with 1.5 mols of dioxane is similarly reactive [2].

See related CARBONYLMETALS

1377. Germanium isocyanate (Tetraisocyanatogermane)

[4756-66-5]

$C_4GeN_4O_4$



Water

Johnson, D. H., *Chem. Rev.*, 1951, **48**, 287


The rate of exothermic hydrolysis becomes dangerously fast above 80°C.

See related CYANO COMPOUNDS, METAL CYANATES

1378. Dichloromaleimide (3,4-Dichloro-2,5-pyrrolidinedione)

[1193-54-0]

$C_4HCl_2NO_2$



Preparative hazard

See Dichloromaleic anhydride: Sodium chloride, Urea

1379. Poly(1-pentafluorothio-1,2-butadiyne)

[84864-30-2]

$(C_4HF_5S)_n$

See Structure

Kovacina, T. A. *et al.*, *Ind. Eng. Chem., Prod. Res. Dev.*, 1983, **22**, 170–172


The insoluble polymer formed in the liquid phase was pressure- and shock-sensitive and should not be heated above 25°C. Cutting with a razor or breaking the polymer causes detonation and ignition.

See related HALOALKENES, DIENES

1380. Perfluoro-*tert*-butanol

[2378-02-1]

C_4HF_9O



Preparative hazard

See Perfluoro-*tert*-nitrosobutane: Nitrogen oxides

1381. 1-Iodo-1,3-butadiyne
[6088-91-1]

C₄HI



1. Schlubach, H. H. *et al.*, *Ann.*, 1951, **573**, 118, 120
2. Kloster-Jensen, E., *Tetrahedron*, 1966, **22**, 969

Crude material exploded violently at 35°C during attempted vacuum distillation, and temperatures below 30°C are essential for safe handling [1]. A sample of pure material exploded on scratching under illumination [2].

See other HALOACETYLENE DERIVATIVES, IRRADIATION DECOMPOSITION INCIDENTS

1382. Potassium hydrogen acetylenedicarboxylate
[928-84-1]

C₄HKO₄




Energy of decomposition (in range 170–360°C) measured as 0.775 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 202°C by adiabatic Dewar tests, with an apparent energy of activation of 155 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ACETYLENIC COMPOUNDS *See related* ORGANIC ACIDS

1383. Cyanoform (Tricyanomethane)
[454-50-2]

C₄HN₃




1. Stull, 1977, 19
2. Rodd, 1976, **IE**, 333

As expected for a tricyano compound, it is highly endothermic (ΔH_f° (s) +348.1 kJ/mol, 3.78 kJ/g), so its stability seems dubious [1]. However, because of isomerisation to the acidic dicyano-enimine (NC)₂C=C=NH, it is considerably stabilised and the silver and sodium salts surprisingly are described as stable [2].

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

1384. 2-Azido-3,5-dinitrofuran
[70664-49-2]

C₄HN₅O₅



Sitzmann, M. E., *J. Heterocycl. Chem.*, 1979, **16**, 477–480

Very sensitive to heat and impact.

See other ORGANIC AZIDES, POLYNITROARYL COMPOUNDS

†1385. 1,3-Butadiyne
[460-12-8]

C_4H_2



1. Armitage, J. B. *et al.*, *J. Chem. Soc.*, 1951, 44
2. Mushii, R. Y. *et al.*, *Khim. Prom.*, 1963, **39**, 109
3. Moshkovich, F. B. *et al.*, *Khim. Prom.*, 1965, **41**, 137–139
4. Anon., *Chem. Age*, 1951, **64**(1667), 955–958
5. Schilling, H. *et al.*, Ger. Pat. 860 212, 1956

Potentially very explosive, it may be handled and transferred by low temperature distillation. It should be stored at -25°C to prevent decomposition and formation of explosive polymers [1]. The critical pressure for explosion is 0.04 bar, but presence of 15–40% of diluents (acetylene, ammonia, carbon dioxide or nitrogen) will raise the critical pressure to 0.92 bar [2]. Further data on attenuation by inert diluents of the explosive decomposition of the diyne are available [3]. During investigation of the cause of a violent explosion in a plant for separation of higher acetylenes, the most important finding was to keep the concentration of 1,3-butadiyne below 12% in its mixtures. Methanol is a practical diluent [4]. The use of butane (at 70 mol%) or other diluents to prevent explosion of 1,3-butadiyne when heated under pressure has been claimed [5]. It polymerises rapidly above 0°C .

Arsenic pentafluoride

Russo, P. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 53–54

Polymerisation of the diyne by arsenic pentafluoride at low temperature may be explosive if too rapid mixing in the gas phase, or too rapid warming of the solids from -196°C , is permitted for 200 mg quantities.

See other POLYMERISATION INCIDENTS

Silver nitrate

Grignard, 1935, Vol. 3, 181

Reaction with ammoniacal silver nitrate gives a very explosive friction-sensitive silver salt.

See METAL ACETYLIDES

See other ALKYNES, PEROXIDISABLE COMPOUNDS

1386. Poly(butadiyne)
[61565-16-0]

$(C_4H_2)_n$



Preparative hazard

See 1,3-Butadiyne: Arsenic pentafluoride

See related ALKYNES

1387. 1,2-Dibromo-1,2-diisocyanatoethane polymers

[51877-12-4]

$(\text{C}_4\text{H}_2\text{Br}_2\text{N}_2\text{O}_2)_2$ or 3



2-Phenyl-2-propyl hydroperoxide


See 2-Phenyl-2-propyl hydroperoxide: 1,2-Dibromo-1,2-diisocyanatoethane polymers

See related CYANO COMPOUNDS, ORGANIC ISOCYANATES

1388. 4-Nitrothiophene-2-sulfonyl chloride

[40358-04-1]

$\text{C}_4\text{H}_2\text{ClNO}_4\text{S}_2$



Libman, D. D., private comm., 1968

After distilling the chloride up to 147°C/6 mbar, the residue decomposed vigorously.

See other NITROACYL HALIDES

1389. 1,1,2,3-Tetrachloro-1,3-butadiene

[1637-31-6]

$\text{C}_4\text{H}_2\text{Cl}_4$



It autoxidises to an unstable peroxide.


See 3,3,4,5-Tetrachloro-3,6-dihydro-1,2-dioxin

See other DIENES, HALOALKENES, PEROXIDISABLE COMPOUNDS

1390. 3,3,4,5-Tetrachloro-3,6-dihydro-1,2-dioxin

[]

$\text{C}_4\text{H}_2\text{Cl}_4\text{O}_2$



Akopyan, A. N. *et al.*, *Chem. Abs.*, 1975, **82**, 111537

This autoxidation product of 1,1,2,3-tetrachloro-1,3-butadiene exploded during attempted vacuum distillation.

See other CYCLIC PEROXIDES

- 1391. Perfluorosuccinic acid (Tetrafluorobutanedioic acid)**
 [377-38-8] **C₄H₂F₄O₂**
 (HOCO.CF₂-)₂
- Caesium fluoride, Fluorine
See Fluorine: Caesium fluoride, etc.
See other ORGANIC ACIDS
- 1392. Heptafluorobutyramide**
 [662-50-0] **C₄H₂F₇NO**
 F₇C₃CO.NH₂
- Lithium tetrahydroaluminate
See Lithium tetrahydroaluminate: Fluoroamides
- 1393. Iron(II) maleate**
 [7705-12-6] **C₄H₂FeO₄**
 Fe(-OCO.CH:)₂
- Schwab, R. F. *et al.*, *Chem. Eng. Prog.*, 1970, **66**(9), 53
 The finely divided maleate, a by-product of phthalic anhydride manufacture, is subject to rapid aerial oxidation above 150°C, and has been involved in plant fires.
See other HEAVY METAL DERIVATIVES, PYROPHORIC MATERIALS
- 1394. Potassium diethynylpalladate(2-)**
 [] **C₄H₂K₂Pd**
 K₂[Pd(C≡CH)₂]
- Air, or Water
 Immediately pyrophoric in air, explosive decomposition with aqueous reagents; the sodium salt is similar.
See entry COMPLEX ACETYLIDES
- 1395. Potassium diethynylplatinate(2-)**
 [] **C₄H₂K₂Pt**
 (Complex Structure)
- Air, or Water
 Pyrophoric in air, explosive decomposition with water; the sodium salt is similar.
See entry COMPLEX ACETYLIDES
- 1396. Manganese(II) bis(acetylide)**
 [] **C₄H₂Mn**
 Mn(C≡CH)₂
- Bailar, 1973, Vol. 3, 855

A highly explosive compound.
See other METAL ACETYLIDES

1397. Fumarodinitrile

[766-42-1]

$C_4H_2N_2$




Energy of decomposition (in range 340–380°C) measured as 0.454 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other CYANO COMPOUNDS

†**1398. Pyrimidine-2,4,5,6-(1H,3H)-tetrone ('Alloxan')**

[50-71-5]

$C_4H_2N_2O_4$



1. Wheeler, A. S., and Bogert, M. T., *J. Amer. Chem. Soc.*, 1910, **32**, 809
2. Gartner, R. A., *J. Amer. Chem. Soc.*, 1911, **33**, 85
3. Franklin, E. C., *J. Amer. Chem. Soc.*, 1901, **23**, 1362
4. Sorbe, 1968, 140

The tetrone slowly decomposes during storage at ambient temperature with generation of carbon dioxide. Two separate pairs of incidents involving bursting of bottles [1], and of pressure generation [2,3], were reported. It also explodes at temperatures above 170°C [4].


See Imidazoline-2,4-dithione

See other GAS EVOLUTION INCIDENTS

1399. 3-Diazoniopyrazolide-4-carboxamide

[102613-59-2]

$C_4H_2N_4O$




Cheng, C. C. *et al.*, *J. Med. Chem.*, 1986, **29**, 1546

Though the preparative method is suitable for large scale synthesis, the product must be handled with care. On rapid heating, the internal diazonium salt decomposes violently at 155–160°C accompanied by a sharp sound. The dried powder should not be scratched with a metal spatula or ground finely.

See other DIAZONIUM SALTS, HIGH-NITROGEN COMPOUNDS

1400. 2,6-Diazidopyrazine
[74273-75-9]


C₄H₂N₈



Shaw, J. T. *et al.*, *J. Heterocycl. Chem.*, 1980, **17**, 14
It is heat- and impact-sensitive, exploding at 200°C, or under a hammer-blow.
See other HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1401. 3,3'-Azo-(1-nitro-1,2,4-triazole)
[104364-15-0]


C₄H₂N₁₀O₄



Lee, K. Y., *Energy Res. Abstr.*, 1985, **10**(17), 34672
A candidate for high-energy propellant applications.
See other AZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, *N*-NITRO COMPOUNDS, TRIAZOLES

1402. Poly(furan-2,5-diyl)
[51325-04-3]

(C₄H₂O)_n



Perchlorate ions
See entry PERCHLORATE-DOPED CONDUCTING POLYMERS

1403. Acetylenedicarboxaldehyde (1,4-Dioxobut-2-yne)
[21251-20-7]

$C_4H_2O_2$



Gorgues, A. *et al.*, *Tetrahedron*, 1990, **46**(8), 2817


The solid explodes, flamelessly, even under nitrogen, at its melting point, $-11^\circ C$.

The vapour was handled at reduced pressure up to $60^\circ C$.

See other ACETYLENIC COMPOUNDS, ALDEHYDES

1404. Maleic anhydride (2,5-Dihydrofuran-2,5-dione)
[108-31-6]

$C_4H_2O_3$



(MCA SD-88, 1962); HCS 1980, 605; RSC Lab. Hazards Data Sheet No. 53, 1983

Vaughn, C. B. *et al.*, *J. Loss Prevention*, 1993, **6**(1), 61

Energy of decomposition (in range $260-370^\circ C$) measured as 0.92 kJ/g.

A fire within a maleic acid plant left pipework glowing red hot. Investigation of possible sources of ignition showed that the anhydride, spread on insulation materials, undergoes runaway autoxidation/decomposition from $190^\circ C$.

See also INSULATION, THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Bases, or Cations

1. Vogler, C. E. *et al.*, *J. Chem. Eng. Data*, 1963, **8**, 620

2. Davie, W. R., *Chem. Eng. News.*, 1964, **42**(8), 41

3. MCA Case History No. 622

4. MCA Case History No. 2032

Maleic anhydride decomposes exothermically, evolving carbon dioxide, in the presence of alkali- or alkaline earth-metal or ammonium ions, dimethylamine, triethylamine, pyridine or quinoline, at temperatures above $150^\circ C$ [1]. Sodium ions and pyridine are particularly effective, even at concentrations below 0.1%, and decomposition is rapid [2]. An industrial incident involved gas-rupture of a large insulated tank of semi-solid maleic anhydride which had been contaminated with sodium hydroxide. Use of additives to reduce the sensitivity of the anhydride has been described [3]. Accidental transfer of an aqueous solution of sodium 2-benzothiazolethiolate into a bulk storage tank of the anhydride led to eventual explosive destruction of the tank [4].

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

1-Methylsilacyclopenta-2,4-diene

See 1-Methylsilacyclopenta-2,4-diene: Dienophiles

See other ACID ANHYDRIDES

1405. Acetylenedicarboxylic acid

[142-45-0]

C₄H₂O₄



Energy of decomposition (in range 140–270°C) measured as 1.355 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Copper salts

Zamudio, W. *et al.*, *Bol. Soc. Chil. Quim.*, 1994, **39**(4), 339

Aqueous solutions of copper acetylenedicarboxylates decompose thermally to precipitate explosive copper acetylides. Presumably other elements readily forming explosive acetylides might do likewise.

See Copper(II) acetylide


See also METAL ACETYLIDES

See other ACETYLENIC COMPOUNDS, ORGANIC ACIDS

1406. Maleic anhydride ozonide

[101672-17-7]

C₄H₂O₆



Briner, E. *et al.*, *Helv. Chim. Acta*, 1937, **20**, 1211


It explodes on warming to –40°C.

See other OZONIDES

1407. Poly(thiophene)

[25233-34-5]

(C₄H₂S)_n



Perchlorate ions

See entry PERCHLORATE-DOPED CONDUCTING POLYMERS

1408. Silver buten-3-ynide
[15383-68-3]




Alone, or Ammonia, or Nitric acid

1. Willstätter, R. *et al.*, *Ber.*, 1913, **46**, 535–538
2. Grignard, 1935, Vol. 3, 187

The silver salt, which deflagrates on heating, explodes if moistened with fuming nitric acid [1], or with ammonia solution [2].

See other METAL ACETYLIDES, SILVER COMPOUNDS

1409. Silver 3-methylisoxazolin-4,5-dione-4-oximate
[70247-51-7]




Wentrup, C. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 467

It explodes on rapid heating. On slow heating under vacuum, it gives dangerously explosive silver fulminate.

See Fulminic acid (reference 2), also Sodium 3-methylisoxazolin-4,5-dione-4-oximate

See related OXIMES See other N–O COMPOUNDS, SILVER COMPOUNDS

1410. 5-Bromo-4-pyrimidinone
[19808-30-1]



Preparative hazard


Dietsche, T. J. *et al.*, *Chem. Eng. News*, 1995, **73**(28), 4

A literature procedure whereby bromopyrimidine is oxidised by excess peroxyacetic acid in acetone, with sulfuric acid catalysis, was being scaled up. The crude product from the fourth batch at two molar scale was filtered out and allowed to dry to dry in the sintered glass funnel over the weekend. An explosion occurred when it was scraped out to complete purification on the Monday. This was considered due to acetone peroxides, which had probably concentrated locally by wicking or sublimation.

See CYCLIC PEROXIDES, SINTERED GLASS

1411. 1-Chloro-3,3-difluoro-2-methoxycyclopropene
[59034-34-3]


C₄H₃ClF₂O



See entry FLUORINATED CYCLOPROPENYL METHYL ETHERS

1412. Acetoxymercurio(perchloratomercurio)ethenone
[73399-68-5]

C₄H₃ClHg₂O₇



Blues, E. T. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 1043

It is dangerously explosive.

See related METAL PERCHLORATES *See other* MERCURY COMPOUNDS

1413. Acetoxymercurio(perchloratodimercurio)ethenone
[73399-71-0]

C₄H₃ClHg₄O₇




Blues, E. T. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 1043

It is dangerously explosive.

See related METAL PERCHLORATES *See other* MERCURY COMPOUNDS

1414. 1,3,3-Trifluoro-2-methoxycyclopropene
[59034-32-1]


C₄H₃F₃O



See entry FLUORINATED CYCLOPROPENYL METHYL ETHERS

1415. Potassium 3-methylfuran-4-carboxylate 2-oxide
[37895-51-5]

C₄H₃KN₂O₄



Gasco, A. *et al.*, *Tetrahedron Lett.*, 1974, 630 (footnote 2)
The dried salt explodes violently on heating, impact or friction.
See other FURAZAN N-OXIDES

1416. 3-Cyanopropyne (1-Butyne-4-nitrile)
[2235-08-7]

C_4H_3N




Reddy, G. S. *et al.*, *J. Amer. Chem. Soc.*, 1961, **83**, 4729
After preparation from interaction of 3-bromopropyne with copper(I) cyanide and filtration from copper salts, an explosion occurred during distillation of the evaporated filtrate at 45–60°C/66 mbar. This was attributed to explosion of some dissolved copper acetylide(s). After refiltration the product was again distilled at 45–48°C/53 mbar without incident, and it appeared to be stable, unlike true haloalkynes. However it is undoubtedly an endothermic compound with its two triple bonds.

See related HALOACETYLENE DERIVATIVES

See other ACETYLENIC COMPOUNDS, CYANO COMPOUNDS

1417. Poly(pyrrole)
[30604-81-0]

$(C_4H_3N)_n$




Perchlorate ions

See entry PERCHLORATE-DOPED CONDUCTING POLYMERS

1418. Maleimide
[541-59-3]

C_4H_3NO




Energy of decomposition (in range 140–300°C) measured as 0.49 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

1419. Sodium 3-methylisoxazolin-4,5-dione-4-oximate
[70247-50-6]

$C_4H_3N_2NaO_3$



Wentrup, C. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 467


It explodes on rapid heating, like the analogous silver salt and 3-phenyl derivative.

See related OXIMES *See other* N-O COMPOUNDS

1420. 5-(Prop-2-ynyloxy)-1,2,3,4-thiatriazole

[]

C₄H₃N₃OS



Banert, K. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(7), 866


Decomposes in solution at 0°C but can be handled, with caution even sublimed, as an explosive crystalline solid, m.p. 66°C.

See other ACETYLENIC COMPOUNDS, N-S COMPOUNDS

1421. 2-Amino-3,5-dinitrothiophene

[2045-70-7]

C₄H₃N₃O₄S




Fast flame propagation occurs on heating the powder moderately.

See entry HIGH RATE DECOMPOSITION

1422. 3-Diazoniopyrazolide-4-carboxamide

[102613-59-2]

C₄H₃N₅O



Cheng, C. C. *et al.*, *J. Med. Chem.*, 1986, **29**, 1546

Though the preparative method is suitable for large scale synthesis, the product must be handled with care. On rapid heating, the internal diazonium salt decomposes violently at 155–160°C accompanied by a sharp sound. The dried powder should not be scratched with a metal spatula or ground finely.

See other DIAZONIUM SALTS, HIGH-NITROGEN COMPOUNDS

†1423. Buten-3-yne (Vinylacetylene)
[689-97-4]

C₄H₄



1. Hennion, G. F. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 684
2. Rutledge, 1968, 28
3. Strizhevskii, I. I. *et al.*, *Chem. Abs.*, 1973, **78**, 98670
4. Vervalin, 1973, 38–43

It must be stored out of contact with air to avoid the formation of explosive (peroxidic) compounds [1]. The sodium salt is a safe source of butenyne for synthetic use [2]. Explosive properties of the liquid and gaseous hydrocarbon alone, or diluted with decahydronaphthalene, were determined [3]. Mechanical failure of a compressor circulating the gaseous hydrocarbon through a reaction system led to local overheating and explosive decomposition, which propagated throughout the large plant [4]. As anticipated for a multiply unsaturated hydrocarbon, vinylacetylene is strongly endothermic (ΔH_f° (g) +143.5 kJ/mol, 2.76 kJ/g).

See other ENDOTHERMIC COMPOUNDS

1,3-Butadiene

1. Jarvis, H. C., *Chem. Eng. Progr.*, 1971, **67**(6), 41–44
2. Freeman, R. H. *et al.*, *Chem. Eng. Progr.*, 1971, **67**(6), 45–51
3. Keister, R. G. *et al.*, *Loss Prev.*, 1971, **5**, 67–75
4. Carver, F. W. S. *et al.*, *Proc. 5th Symp. Chem. Proc. Haz.*, No. 39a, 99–116, London, IChE, 1975

Unusual conditions in the main fractionation column separating product butadiene from by-product butenyne (vinylacetylene, thought to be safe at below 50 mol% concentration) caused the concentration of the latter to approach 60% in part of the column as fractionation proceeded. Explosive decomposition, possibly initiated by an overheated unstable organic material derived from sodium nitrite, destroyed the column and adjacent plant [1,2]. Subsequent investigation showed that all mixtures of butenyne and butadiene can reproducibly be caused to react exothermically and then decompose explosively at appropriately high heating rates under pressure. Butadiene alone will behave similarly at higher heating rates and pressures [3]. Another study of the detailed mechanism and type of decomposition of C₄ hydrocarbons containing butenyne showed that both were dependent on energy of initiation and time of application, as well as on the composition and phases present [4].

See Sodium nitrite: 1,3-Butadiene

Oxygen

1. Dolgopolskii, I. M. *et al.*, *Chem., Abs.*, 1958, **52**, 19904g
2. Gosa, V. *et al.*, *Rev. Chim.* (Bucharest), 1981, **32**, 1103–1105
3. Ionescu, N. I. *et al.*, *Rev. Chim.* (Bucharest), 1982, **33**, 1021–1026

The rate of absorption of oxygen by liquid butenyne increased with time, and eventually a yellow liquid phase separated. After evaporation of excess hydrocarbon, the yellow peroxidic liquid was explosive. Presence of 5% of chloroprene

increased the rate of absorption 5–6-fold, and of 2% of water decreased the rate by 50%, but residues were explosive in each case [1]. Explosive combustion in admixture with oxygen has been studied [2], and the effects of presence of nitrogen upon explosion parameters were determined [3].

Silver nitrate

Grignard, 1935, Vol. 3, 187

An explosive silver salt is formed.


See Silver buten-3-ynide

See other ALKYNES, PEROXIDISABLE COMPOUNDS

1424. Barium 1,3-di(5-tetrazolyl)triazenide

[]

$C_4H_4BaN_{22}$




See 1,3-Di(5-tetrazolyl)triazene

See other HIGH-NITROGEN COMPOUNDS, N-METAL DERIVATIVES, TETRAZOLES, TRIAZENES

1425. N-Bromosuccinimide (1-Bromo-2,5-pyrrolidinedione)

[128-08-5]

$C_4H_4BrNO_2$



Aniline, or Diallyl sulfide, or Hydrazine hydrate

See N-HALOIMIDES: alcohols, etc.

Dibenzoyl peroxide, 4-Toluic acid

1. Tcheou, F. K. *et al.*, *J. Chinese Chem. Soc.*, 1950, **17**, 150

2. Mould, R. W., private comm., 1975

Thirtyfold increase in scale of a published method [1] for radical-initiated side-chain bromination of the acid in carbon tetrachloride led to violent reflux and eruption of the flask contents through the condenser [2].

See other HALOGENATION INCIDENTS

Propionitrile

MCA Guide, 1972, 309

After refluxing for 24 hours at 105°C, a mixture exploded, possibly due to dehydrohalogenation of the bromonitrile to acrylonitrile, and polymerisation of the latter.

See other *N*-HALOGEN COMPOUNDS, *N*-HALOIMIDES

1426. 2-Chloro-2-propenyl trifluoromethanesulfonate
[62861-56-7]


C₄H₄ClF₃O₃S



See entry ALLYL TRIFLUOROMETHANESULFONATES See other SULFUR ESTERS

1427. *N*-Chlorosuccinimide (1-Chloro-2,5-pyrrolidinedione)
[128-09-6]

C₄H₄ClNO₂



Energy of decomposition (in range 160–380°C) measured as 1.61 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alcohols, or Benzylamine

See *N*-HALOIMIDES: alcohols, etc.

Dust

Boscott, R. J., private comm., 1968

Smouldering in a stored drum of the chloroimide was attributed to dust contamination.

See other *N*-HALOGEN COMPOUNDS, *N*-HALOIMIDES

1428. 1,4-Dichloro-2-butyne
[821-10-3]

C₄H₄Cl₂



1. Ford, M. *et al.*, *Chem. Eng. News*, 1972, **50**(3), 67

2. Herberz, T., *Chem. Ber.*, 1952, **85**, 475–482


3. *CHETAH*, 1990, 182 and 188

Preparation of the acetylenic dichloride by conversion of the diol in pyridine with neat thionyl chloride is difficult to control, and hazardous on a large scale. Use of dichloromethane as diluent and operation at –30°C renders the preparation reproducible and safer [1]. During distillation at up to 110°C/7–8 mbar, slight overheating of the residue to 120°C caused explosive decomposition [2]. Sensitive to detonator, sometimes to mechanical shock [3].

See other HALOACETYLENE DERIVATIVES

1429. Copper(II) 1,3-di(5-tetrazolyl)triazenide
[32061-49-7]

$C_4H_4CuN_{22}$



See 1,3-Di(5-tetrazolyl)triazene

See other HIGH-NITROGEN COMPOUNDS, N-METAL DERIVATIVES, TETRAZOLES, TRIAZENES

1430. Tetrafluorosuccinamide
[377-37-7]

$C_4H_4F_4N_2O_2$



Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Fluoroamides

1431. Poly[bis(2,2,2-trifluoroethoxy)phosphazene]
[28212-50-2]

$(C_4H_4F_6NO_2P)_n$



Allcock, H. R. *et al.*, *Macromolecules*, 1974, **7**, 284–290

Sealed tubes containing the linear P=N polymer exploded after heating at 300°C for 24–30 h (or after shorter times at higher temperatures), owing to pressure build-up from formation of the more volatile cyclic tri- and tetra-mers.

1432. 2-Cyano-1,2,3-tris(difluoroamino)propane
[16176-02-6]

$C_4H_4F_6N_4$



Reed, S. F., *J. Org. Chem.*, 1968, **33**, 1864

The 95% pure material is shock-sensitive.

See other CYANO COMPOUNDS, DIFLUOROAMINO COMPOUNDS

1433. Succinodinitrile (1,4-Butanedinitrile)
[110-61-2]

$C_4H_4N_2$



1. Kniess, H., private comm., 1983

2. Berthold, W. *et al.*, *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 3, C1–C4, Rugby, IChE, 1983


After vacuum distillation of several t of the nitrile at 150°C, lack of a receiver necessitated hot storage of the product at 80°C for 46 h, and the drainage line blocked with solid. Though the bulk of the material was still liquid, heating to 195°C did not clear the line, and shortly afterwards decomposition occurred. It was subsequently found that the distillation residue had also decomposed at 175°C, 2 h after distillation ended [1]. DTA showed that the nitrile is unusual in that no heat is evolved during the induction period of around 33 h/200°C, or 1.5 h/280°C, and the rapid onset and narrow peak width suggested a self-accelerating decomposition process, with $Q = 700$ J/g. No stabilising additives were found, but addition of 1% of hydrogen cyanide reduced the induction period for the nitrile to 20% of its value, while 1% of potassium cyanide reduced it to 2%. This supports the proposed mechanism of cyanide ion-catalysed decomposition to hydrogen cyanide and acrylonitrile, followed by their simultaneous polymerisation, again by self-accelerating processes. Adiabatic storage tests on 150 g samples gave the same induction period and a decomposition peak of 400°C/40 bar [2].

See entry SELF-ACCELERATING REACTIONS

See other CYANO COMPOUNDS, INDUCTION PERIOD INCIDENTS

1434. α -Diazo- γ -thiobutyrolactone
[79472-87-0]

C₄H₄N₂OS




See α -Bromo- γ -thiobutyrolactone

See other DIAZO COMPOUNDS

1435. Maleic hydrazide (1,2-Dihydropyridazine-3,6-dione)
[123-33-1]


C₄H₄N₂O₂



Energy of decomposition (in range 280–300°C) measured as 0.59 kJ/g by DSC, and T_{ait24} was determined as 261°C by adiabatic Dewar tests, with an apparent energy of activation of 257 kJ/mol.

1436. 2-Amino-4,6-dihydroxy-5-nitropyrimidine
[80466-56-4]

C₄H₄N₄O₄



Madeley, J. P., personal communication 1989

A 200 g sample, prepared by addition of substrate to mixed acid, followed by quenching into water, filtration and washing, decomposed vigorously at a late stage of drying in a vacuum oven, which was pressurised and the seal forced open. This behaviour might have been due to inadequate washing and residual sulfuric acid.

See other C-NITRO COMPOUNDS

1437. 2,3-Diazido-1,3-butadiene
[91686-86-1]

C₄H₄N₆



Mielsen, C. J. *et al.*, *J. Mol. Struct.*, 1986, **147**, 217–229

Although it is extremely explosive, like other polyunsaturated azides, it was possible to run gaseous electron diffraction and IR spectra at cryogenic temperatures. Sample decomposition occurred during laser Raman spectral determination.

See other DIENES, ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1438. Succinoyl diazide
[40428-75-9]

C₄H₄N₆O₂



1. Maclaren, J. A., *Chem. & Ind.*, 1971, 3952

2. France, A. D. G. *et al.*, *Chem. & Ind.*, 1962, 2065

This intermediate for the preparation of ethylene diisocyanate exploded violently during isolation [1]. Previously the almost dry solid had exploded violently on stirring with a spatula [2].

See other ACYL AZIDES, FRICTIONAL INITIATION INCIDENTS

†**1439. Furan**
[110-00-9]


C₄H₄O



See other APROTIC SOLVENTS, PEROXIDISABLE COMPOUNDS

1440. 1,2-Cyclobutanedione
[33689-28-0]

C₄H₄O₂




Dennis, J. M. *et al.*, *Org. Synth.*, 1981, **60**, 18–19

The dione must be stored cold and handled out of direct light to prevent polymerisation.

†**1441. Diketene (4-Methylene-2-oxetanone)**
[674-82-8]

C₄H₄O₂



HCS 1980, 416

1. Vervalin, 1973, 86
2. Zdenek, F. *et al.*, Czech Pat. 156 584, 1975
3. *See entry* PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Diketene residues in a tank trailer awaiting incineration decomposed violently on standing, blew off the dome cover and ignited [1]. The risk of autoignition during the exothermic dimerisation of diketene to dehydroacetic acid is eliminated by operating in a non-flammable solvent [2]. It produces a moderate pressure rise on exothermic decomposition at 125°C [3].

Acids, or Bases, or Sodium acetate

1. *Haz. Chem. Data*, 1975, 136
2. Laboratory Chemical Disposal Co. Ltd., confid. information, 1968

Presence of mineral or Lewis acids, or bases including amines, will catalyse violent polymerisation of this very reactive dimer, accompanied by gas evolution [1]. Sodium acetate is sufficiently basic to cause violent polymerisation at 0.1% concentration when added to diketene at 60°C [2].

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS, POLYMERISATION INCIDENTS

†**1442. Methyl propiolate (Methyl propynoate)**
[922-67-8]

C₄H₄O₂




Octakis(trifluorophosphine)dirhodium

See Octakis(trifluorophosphine)dirhodium: Acetylenic esters

See other ACETYLENIC COMPOUNDS

1443. Succinic anhydride
[108-30-5]

C₄H₄O₃



Sodium hydroxide

1. Hoekstra, M. C., *Synth. Comm.*, 1984, **14**, 1379–1380
2. Kirk Othmer, 1997, (4th edn.), Vol. 22, 1076

Succinic anhydride is dimerised to 1,6-dioxaspiro [4.4] nonane-2,7-dione by heating with sodium hydroxide. Modification of an existing procedure by adding further sodium hydroxide after the initial reaction led to a severe exothermic reaction after heating for some 30 h which fused the glass flask to the heating mantle, probably at a temperature approaching 550°C. The reason for this was not known [1]. At elevated temperatures and under influence of alkali, succinic acid condenses decarboxylatively beyond the dimeric spiroacetal, sometimes explosively. Contamination of the anhydride with base is to be avoided [2].

See other GLASS INCIDENTS

See other ACID ANHYDRIDES

1444. 2,3,7-Trioxabicyclo[2,2,1]hept-5-ene
[6824-18-6]

C₄H₄O₃


Lorenca, P. *et al.*, *J. Phys. Chem.*, 1989, **93**(6), 2276

This, the ozonide of cyclobutadiene (and also a likely peroxide of furan), may be stored in liquid nitrogen. It can explode with considerable force on warming.

See other CYCLIC PEROXIDES, OZONIDES

1445. 3,6-Dioxo-1,2-dioxane (Succinyl peroxide)
[]

C₄H₄O₄



CHETAH, 1990, 183


This cyclic peroxide, readily formed from succinate derivatives and peroxides, is claimed to be more stable than its molecular formula might indicate. It is, however, shock sensitive.

See other DIACYL PEROXIDES, CYCLIC PEROXIDES

1446. Fumaric acid (*trans*-2-Butene-1,4-dioic acid)

[110-17-8]

C₄H₄O₄



Energy of decomposition (in range 290–340°C) measured as 0.925 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ORGANIC ACIDS

1447. Dihydroxymaleic acid (Dihydroxybutenedioic acid)

[526-84-1]

C₄H₄O₆



Axelrod, B., *Chem. Eng. News*, 1955, **33**, 3024


A glass ampoule of the acid exploded during storage at ambient temperature. It is unstable and slowly loses carbon dioxide, even at 4°C.

See other GAS EVOLUTION INCIDENTS, GLASS INCIDENTS, ORGANIC ACIDS

†1448. Thiophene

[110-02-1]

C₄H₄S



HCS 1980, 899

Nitric acid

See Nitric acid: Thiophene

N-Nitrosoacetanilide

See *N*-Nitrosoacetanilide: Thiophene

1449. Ethynyl vinyl selenide

[101672-11-1]

C₄H₄Se




Brandsma, L. *et al.*, *Rec. Trav. Chim.*, 1962, **81**, 539

Distillable at 30–40°C/80 mbar, it decomposes explosively on heating at ambient pressure.

See other ACETYLENIC COMPOUNDS

1450. α -Bromo- γ -thiobutyrolactone (3-Bromo-2-oxotetrahydrothiophene)
[20972-64-9] **C₄H₅BrOS**



Miller, G. A. *et al.*, *J. Org. Chem.*, 1981, **46**, 4751–4753
In the preparation *via* diazotisation and bromination of homocysteine thiolactone, presence of unreacted diazo intermediate led to an explosion during distillation of the product.
See DIAZO COMPOUNDS

†1451. 2-Chloro-1,3-butadiene (Chloroprene)
[126-99-8] **C₄H₅Cl**




1. Bailey, H. C., in *Oxidation of Organic Compounds-I*, ACS No. 75 (Gould, R. F., Ed.), 138–149, Washington, ACS, 1968
2. Bailey, H. C., private comm., 1974

Chloroprene monomer will autoxidise very rapidly with air, and even at 0°C it produces an unstable peroxide (a mixed 1,2- and 1,4-addition copolymer with oxygen), which effectively will catalyse exothermic polymerisation of the monomer. The kinetics of autoxidation have been studied [1]. It forms ‘popcorn polymer’ at a greater rate than does butadiene [2].

See other DIENES, HALOALKENES, PEROXIDISABLE COMPOUNDS, POLYMERISATION INCIDENTS

Preparative hazard
See 2-Chloro-1-nitro-4-nitroso-2-butene

1452. 2-Chloro-1-nitro-4-nitroso-2-butene
[58675-02-8] **C₄H₅ClN₂O₃**




Simmons, H. E. *et al.*, *Chem. Eng. News*, 1995, **73**(32), 4
A worker was injured by an explosion when cutting into a “decontaminated” steel pipe removed from a storage tank in a chloroprene plant. The pipe was found to be coated by a deposit of this compound, existing as a dimer. Investigation showed it to be unstable from 80°C, deflagrating with an energy of 2.5 kJ/g (about that of black powder). It is thought it may be capable of true detonation. It was formed by reaction of chloroprene with dinitrogen trioxide, generated during distillation of the chloroprene by decomposition of an ill-defined nitrosated additive used for vapour phase stabilisation against polymerisation reactions. Although originally present

at negligible concentration the nitroso-dimer, being very insoluble in chloroprene, accumulated in the pipe. Warning is given that reaction of alkenes with nitrogen oxides, to form similar products, is a general reaction.

See Nitrogen dioxide: Alkenes

1453. 4-Chloro-1-methylimidazolium nitrate
[[4897-21-6] (base)]

C₄H₅ClN₃O₃



Personal experience

Towards the end of evaporation of a dilute aqueous solution, the residue decomposed violently. Nitrate salts of many organic bases are thermally unstable and should be avoided.

See other OXOSALTS OF NITROGENOUS BASES

1454. 1-Chloro-1-buten-3-one
[7119-27-9]

C₄H₅ClO



Pohland, A. E. *et al.*, *Chem. Rev.*, 1966, **66**, 164

If prepared from vinyl chloride, this unstable ketone will decompose almost explosively within one day, apparently owing to the presence of some *cis*-isomer.

See related HALOALKENES

1455. 4-Chloro-2-butyne-1-ol
[13280-07-4]

C₄H₅ClO



1. Taylor, S. K. *et al.*, *J. Org. Chem.*, 1983, **48**, 595

2. Harmon, D. P. G. *Chem. Brit.* 1998, (7), 19

It exploded during distillation [1]. Another explosion during distillation of a crude sample is reported. It is claimed that there were no previous warnings of the hazard [2].

See other HALOACETYLENE DERIVATIVES

1456. Ethyl oxalyl chloride (Ethyl chloroglyoxylate)
[4755-77-5]

C₄H₅ClO₃



Urban, P. G., Personal experience

A drum of clear and bright material which had been vacuum distilled pressurised on standing overnight. Investigation revealed a crack in the heat exchanger which had been leaking water into the oxalyl chloride, forming partial anhydrides. These are unstable and decompose to give carbon monoxide and dioxide, other oxalyl halides will behave similarly. On another occasion, a vacuum distillation went suddenly to positive pressure on heating to about 70°C. This was catalysis by traces of iron or aluminium chlorides, which induce break up to, probably, carbon monoxide, carbon dioxide and ethyl chloride.

See Oxalyl dichloride

See other ACYL HALIDES, CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

1457. 2-Methyl-2-propenoyl chloride (Methacryloyl chloride)

[920-46-7]

C₄H₅ClO



Frazier, J., *Chem. Eng. News*, 1996, **74**(23), 4

A five year old bottle, originally stabilised with phenothiazine, was found to have polymerised explosively in storage. The resultant black goo was not acidic, suggesting something more complicated than an acrylate polymerisation. It is recommended that this monomer be stored under refrigeration.

See VIOLENT POLYMERISATION

1458. 4-Fluoro-4,4-dinitrobutene

[19273-49-5]

C₄H₅FN₂O₄




See entries FLUORODINITROMETHYL COMPOUNDS, NITROALKENES

1459. Bis(2-fluoro-2,2-dinitroethyl)amine

[18139-03-2]

C₄H₅F₂N₅O₈



1. Gilligan, W. H., *J. Org. Chem.*, 1972, **37**, 3947

2. Gilligan, W. H., *J. Chem. Eng. Data*, 1982, **27**, 97–99

The amine and several derived amides are explosives, and need appropriate care in handling [1]. Several other derivatives, most of them explosive, are prepared and described [2].

See other FLUORODINITROMETHYL COMPOUNDS, ORGANIC BASES

†1460. Ethyl trifluoroacetate

[383-63-1]

C₄H₅F₃O₂



Diethyl succinate, Sodium hydride

See Sodium hydride: Diethyl succinate, etc.

1461. Prop-2-enyl trifluoromethanesulfonate

[41029-45-2]

$C_4H_5F_3O_3S$



It explodes at 20°C.

See entry ALLYL TRIFLUOROMETHANESULFONATES See other SULFUR ESTERS

1462. Potassium hydrogen tartrate

[868-14-4]

$C_4H_5KO_6$



Carbon, Nitrogen oxide

See Nitrogen oxide: Carbon, etc.

See related ORGANIC ACIDS

1463. Cyanocyclopropane (Cyclopropanecarbonitrile)

[5500-21-0]

C_4H_5N



Preparative hazard


See 4-Chlorobutyronitrile: Sodium hydroxide

See other CYANO COMPOUNDS, STRAINED-RING COMPOUNDS

†**1464. 1-Cyanopropene (2-Butenitrile)**

[627-26-9]

C_4H_5N



See other CYANO COMPOUNDS

†**1465. 3-Cyanopropene (3-Butenitrile)**

[109-75-1]

C_4H_5N



See other ALLYL COMPOUNDS, CYANO COMPOUNDS

1466. Pyrrole

[109-97-7]

C_4H_5N



2-Nitrobenzaldehyde

See 2-Nitrobenzaldehyde: Pyrrole

1467. 1-Cyano-2-propen-1-ol (2-Hydroxy-3-butenitrile)
[5809-59-6]

C₄H₅NO




Unpublished information

It shows a strong tendency to exothermic polymerisation of explosive violence in presence of light and air above 25–30°C. Presence of an inhibitor and inert atmosphere are essential for stable storage.

See other CYANO COMPOUNDS

1468. 5-Methylisoxazole
[576-44-6]

C₄H₅NO




Sealed samples decompose exothermally above 171°C.

See entry ISOXAZOLES See other N–O COMPOUNDS

1469. N-Hydroxysuccinimide
[6066-82-6]

C₄H₅NO₃




Energy of decomposition (in range 180–370°C) measured as 1.51 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other N–O COMPOUNDS

1470. Methyl isocyanoacetate
[39687-95-1]

C₄H₅NO₂



Heavy metals

Naef, H. *et al.*, private comm., 1983

Crude material prepared in glass on 3 g mol scale was distilled uneventfully at 40°C/0.067 mbar from a bath at 70–80°C. A 30 mol batch prepared in a glass-lined vessel with a stainless steel thermo-probe (and later found to contain 15 ppm of iron) decomposed very violently during distillation at 75°C/13 mbar from a bath at 130°C. Thermal analysis showed that the stability of the methyl (and ethyl) ester

was very sensitive to traces of heavy metals (iron, copper, chromium, etc.) and was greatly reduced. Addition of traces of hydrated iron(II) sulfate led to explosive decomposition at 25°C.

See related CYANO COMPOUNDS *See other* CATALYTIC IMPURITY INCIDENTS

1471. Allyl isothiocyanate (3-Isothiocyanatopropene)

[57-06-7]

C₄H₅NS



Anon., *Ind. Eng. Chem. (News Ed.)*, 1941, **19**, 1408

A routine preparation by interaction of allyl chloride and sodium thiocyanate in an autoclave at 5.5 bar exploded violently at the end of the reaction. Peroxides were not present or involved and no other cause could be found, but extensive decomposition occurred when allyl isothiocyanate was heated to 250°C in glass ampoules. Exothermic polymerisation seems a likely possibility.


See related CYANO COMPOUNDS, ORGANIC ISOCYANATES

See other ALLYL COMPOUNDS

1472. 2-Azido-1,3-butadiene

[01686-88-3]

C₄H₅N₃



Schel, S. A. *et al.*, *J. Mol. Struct.*, 1986, **147**(3-4), 203-215


Although it is highly explosive, like other polyunsaturated azides, it was possible to record spectral data under the following conditions: gaseous electron diffraction; IR spectra of matrix-isolated species in argon at 15°K; of amorphous and crystalline solids at 90°K and Raman spectra of the liquid at 240°K.

See other DIENES, ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1473. Azido-2-butyne

[105643-77-4]

C₄H₅N₃



Nielsen, C. J. *et al.*, *J. Mol. Struct.*, 1987, **16**(1-2), 41-56

Precautions appropriate to a potentially explosive compound were observed during synthesis and handling. Electron diffraction studies on the vapour, and IR, NMR and Raman spectral studies on the vapour, liquid and solid (-183°C) phases were effected without incident.

See other ACETYLENIC COMPOUNDS, ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1474. Iminobisacetonitrile

[628-87-5]

C₄H₅N₃




Anon., *Loss. Prev. Bull.*, 1993, (109), 7

Iminobisacetonitrile crystallised in a holding tank of a production plant. Cleaning was by filling with water and heating to dissolve the nitrile. It was heated to a supposed 75°C. An hour later the tank exploded causing severe structural damage. The pyrometer used to measure the temperature was found inaccurate. The nitrile decomposes exothermically from perhaps as low as 70°C in alkaline water (135°C dry) to a variety of products including HCN.

See other CYANO COMPOUNDS

1475. 2-Methyl-4-nitroimidazole
[696-23-1]



Zmojdzin, A. *et al.*, Brit. Pat. 1 418 538, 1974

2-Methylimidazole is difficult to nitrate, and use of conventional reagents under forcing conditions (excess sulfuric/nitric acids, high temperature) involves a high risk of cleavage and violent runaway oxidation. A new and safe process involves the use of nitration liquor from a previous batch (and optional use of excess nitric acid), both of which moderate the nitration reaction, in conjunction with balancing amounts of sulfuric acid, which tends to accelerate the nitration of 2-methylimidazole. Accurate control of the highly exothermic reaction is readily effected.

See other NITRATION INCIDENTS


Nitric acid, Sulfuric acid

Zmojdzin, A. *et al.*, Fr. Demande 2 220 523, 1974

The danger of explosion during further nitration with nitrating acid is eliminated by addition of excess nitric acid as the reaction proceeds.

See NITRATION INCIDENTS *See related* NITROARYL COMPOUNDS

1476. 5-Aminoisoxazole-3-carbonamide
[3445-52-1]




Sealed samples decompose exothermally above 155°C.

See entry ISOXAZOLES *See other* N-O COMPOUNDS

1477. 1-Hydroxyimidazole-2-carboxaldoxime 3-oxide
[35967-34-3]

C₄H₅N₃O₃




Hayes, K. J., *J. Heterocycl. Chem.*, 1974, **11**, 615

It is an extremely pyrophoric, containing three types of N–O bonds.

See other N–O COMPOUNDS, OXIMES

1478. Sodium ethoxyacetylide
[73506-39-5]

C₄H₅NaO



1. Jones, E. R. H. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 405


2. Brandsma, 1971, 120

It is extremely pyrophoric, apparently even at -70°C [1], and may explode after prolonged contact with air [2].

See other METAL ACETYLIDES, PEROXIDISABLE COMPOUNDS

†**1479. 1,2-Butadiene**
[590-19-2]

C₄H₆



Highly endothermic, ΔH_f° (g) +165.5 kJ/mol, 3.06 kJ/g.

Cyclopentadiene (or its dimer)

Donike, W., Ger. Offen. DE 3 617 461, 1987 (*Chem. Abs.*, 1988, **109**, 210597)

In the Diels-Alder condensation of the 2 neat endothermic dienes to give 5-ethylidene- and 5-methyl-6-methylene-bicyclo[2.2.1]hept-2-ene, there is a serious risk of explosive decomposition arising from local overheating of the reactor walls. This hazard is eliminated by the presence of various hydrocarbons and their mixtures as diluents.

See other DIENES, ENDOTHERMIC COMPOUNDS

†**1480. 1,3-Butadiene**
[106-99-0]

C₄H₆



(MCA SD-55, 1954); FPA H63, 1977; HCS 1980, 223 (cylinders); RSC Lab. Hazards Data Sheet No. 45, 1986

1. Scott, D. A., *Chem. Eng. News*, 1940, **18**, 404
2. Hendry, D. G. *et al.*, *Ind. Eng. Chem., Prod. Res. Dev.*, 1968, **7**, 136, 1145
3. Mayo, F. R. *et al.*, *Prog. Rept. No. 40*, Sept. 1971, Stamford Res. Inst. Project PRC-6217
4. Keister, R. G. *et al.*, *Loss Prev.*, 1971, **5**, 69
5. Penkina, O. M. *et al.*, *Chem. Abs.*, 1975, **83**, 29457
6. Bailey, H. C., private comm., 1974
7. Miller, G. H. *et al.*, *J. Polymer Sci. C*, 1964, 1109–1115
8. Vervalin, 1964, 335–338, 358–359
9. Harmon, 1974, 2.5–2.6
10. Alexander, D. S., *Ind. Eng. Chem.*, 1959, **51**, 733–738

Solid butadiene at below -113°C will absorb enough oxygen at reduced pressure to make it explode violently when allowed to melt. The peroxides formed on long contact with air are explosives sensitive to heat or shock, but may also initiate polymerisation [1]. The hazards associated with peroxidation of butadiene are closely related to the fact that the polyperoxide is insoluble in butadiene and progressively separates. If local concentrations build up, self-heating from the (initially slow) spontaneous decomposition begins, and when a large enough mass of peroxide accumulates, explosion occurs. Critical mass at 27°C is a 9 cm sphere; the size decreases rapidly with increasing temperature [2]. Although isoprene and styrene also readily peroxidise, the peroxides are soluble in the monomers, and the degree of hazard is correspondingly less [3].

Butadiene is rather endothermic ($\Delta H_f^{\circ}(\text{g}) +111.9 \text{ kJ/mol}$, 2.07 kJ/g) and will decompose explosively if heated under pressure at $30\text{--}40^{\circ}\text{C/min}$ to exceed critical temperatures of $200\text{--}324^{\circ}\text{C}$ and critical pressures of $1.0\text{--}1.2 \text{ kbar}$ simultaneously [4]. A more recent study of the explosion properties of butadiene at elevated temperatures and pressures has been published (but title only translated) [5]. Phenolic antioxidants (e.g. *tert*-butylcatechol at 0.02 wt.%) are effective in stabilising butadiene against autoxidation in clean storage at moderate ambient temperatures. The presence of rust and/or water in steel storage cylinders rapidly consumes the antioxidant [6]. In prolonged storage, butadiene will (even when very pure and in sealed glass containers) produce ‘popcorn’ polymer [7]. Growth of this involves continued diffusion of monomer into an existing polymeric matrix which continues to increase in bulk. The expansion eventually may rupture the container [6]. Case histories of two industrial explosions involving peroxide formation with air have been detailed [8], and the literature relating to violent polymerisation of butadiene has been reviewed [9]. A detailed account of the experimental investigation following a large-scale explosion in 1951 of a partially full 159 m^3 butadiene storage vessel is given. The involvement of peroxide formation in the explosion was established, as was the effectiveness of the presence of aqueous sodium hydroxide in destroying peroxide and preventing explosion [10].

See Poly(1,3-butadiene peroxide): Butadiene

See other SELF-HEATING AND IGNITION INCIDENTS, PEROXIDATION INCIDENTS

See entries POLYMERISATION INCIDENTS, VIOLENT POLYMERISATION

See other GLASS INCIDENTS

Aluminium tetrahydroborate

See Aluminium tetrahydroborate: Alkenes, etc.

Boron trifluoride etherate, Phenol

MCA Case History No. 790

The hydrocarbon-phenol reaction, catalysed by the etherate, was being run in petroleum ether solution in a sealed pressure bottle. The bottle burst, possibly owing to exothermic polymerisation of the diene.

Buten-3-yne

See Buten-3-yne: 1,3-Butadiene

Cobalt

See reference 7 above

Crotonaldehyde

See Crotonaldehyde: Butadiene

Ethanol, Iodine, Mercury oxide

See 2-Ethoxy-1-iodo-3-butene

Other reactants

Yoshida, 1980, 296

MRH values calculated for 13 combinations with oxidants are given.

Oxides of nitrogen, Oxygen

MRH Nitrogen oxide 8.74/85

Vervalin, 1973, 63-65

An explosion and fire occurred in the pipework of a vessel in which dilute butadiene was stored under an 'inert' atmosphere, generated by the combustion of fuel gas in a limited air supply. The 'inert' gas, which contained up to 1.8% of oxygen and traces of oxides of nitrogen, reacted in the vapour phase over an extended period to produce concentrations of gummy material containing up to 64% of butadiene peroxide and 4.2% of a butadiene-nitrogen oxide complex. The deposits eventually decomposed explosively.

See Nitrogen oxide: Dienes, Oxygen

Oxygen

MCA Case History No. 303

A leaking valve allowed butadiene to accumulate in a pipeline exposed to an inerting gas containing up to 2% of oxygen. Peroxide formed and initiated 'popcorn' polymerisation which burst the pipeline.


See reference 6 above

Sodium nitrite

See Sodium nitrite: 1,3-Butadiene

See other DIENES, ENDOTHERMIC COMPOUNDS, PEROXIDISABLE COMPOUNDS

†1481. 1-Butyne
[107-00-6]



C_4H_6

Highly endothermic (ΔH_f° (g) +166.1 kJ/mol, 3.07 kJ/g).
See other ALKYNES, ENDOTHERMIC COMPOUNDS

†1482. 2-Butyne
[503-17-3]



C_4H_6

HCS 1980, 253

Highly endothermic (ΔH_f° (g) +148.0 kJ/mol, 2.74 kJ/g).
See other ALKYNES, ENDOTHERMIC COMPOUNDS


†1483. Cyclobutene
[822-35-3]



C_4H_6

See other ALKENES

1484. *cis*-Poly(butadiene)
[9003-17-2]




$(\text{C}_4\text{H}_6)_n$

Sedov, V. V. *et al.*, *Chem. Abs.*, 1976, **85**, 109747

Stereoregular *cis*-poly(butadiene) compositions may explode when heated at 337–427°C/1–0.01 μbar, presumably owing to exothermic cyclisation.

See related DIENES

1485. Dimethylaminobis(trifluoromethyl)borane (*N,N*-Dimethyl-1,1-bis(trifluoromethyl)boranamine)
[105224-90-6]



$\text{C}_4\text{H}_6\text{BF}_6\text{N}$

Primary alkynes

Bürger, H. *et al.*, *Main Group Met. Chem.*, 1995, **18**(5), 235

The adducts of primary alkynes with the borane may explode when heated to 90°C.
See other ACETYLENIC COMPOUNDS

1486. Barium acetate
[543-80-6]



Copper(II) oxide, Yttrium oxide


1. Fleischer, N., *Chem. Eng. News*, 1988, **66**(15), 2
2. Author's comments
3. Castrillon, J., *Chem. Eng. News*, 1988, **66**(25), 2

A pelleted mixture containing barium acetate, copper(II) oxide and yttrium oxide, 3 g in all in a quartz tube, was heated in a furnace, and a small explosion occurred during the early stages, 'from formation of pyrolysis products'. It was suggested that the tube should in future be purged, preferably with an oxygen-containing gas [1]. This suggestion is hazardous, however, as the pyrolysis product is acetone. Only inert gas should be used for purging [2]. Pyrolysis of barium acetate gives high yields of acetone, and is catalysed by the metal oxides present in the mixture [3].

See Cadmium propionate

See other HEAVY METAL DERIVATIVES

1487. 4-Chlorobutyronitrile (4-Chlorobutanonitrile)
[628-20-6]




Sodium hydroxide

Brogli, F. *et al.*, *Runaway Reactions*, 1981, Paper 3/M, 5–6, 10

Unstable plant-scale operation in the catalysed cyclisation by sodium hydroxide to cyclopropanecarbonitrile was investigated using a bench scale calorimeter. Crust formation on the reactor wall, which caused the erratic operation, was eliminated by using liquid alkali instead of solid.

See other CYANO COMPOUNDS

1488. N-Chloro-5-methyl-2-oxazolidinone
[25480-76-6]




Wallis, W. E., US Pat. 3 850 920, 1974

It exploded at 160°C.

See other N-HALOGEN COMPOUNDS

1489. N-Chloro-3-morpholinone
[33744-03-5]


$C_4H_6ClNO_2$



Wallis, W. E., US Pat 3 850 920, 1974
It exploded at 115°C.
See other N-HALOGEN COMPOUNDS

1490. N-Chloro-4,5-dimethyltriazole
[72040-09-6]

$C_4H_6ClN_3$



Gallagher, T. C. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 420
The solid (or a concentrated solution) decomposes vigorously on standing at ambient temperature and needs careful handling.
See other N-HALOGEN COMPOUNDS, TRIAZOLES

1491. 1,1-Dichloroethyl peroxyacetate
[59183-18-5]


$C_4H_6Cl_2O_3$

$MeCO.OOCCl_2Me$

Griesbaum, K. *et al.*, *J. Amer. Chem. Soc.*, 1976, **98**, 2880
A friction- and heat-sensitive viscous liquid, it exploded violently during attempted injection into a gas chromatograph.
See other PEROXYESTERS

1492. 3,6-Dichloro-3,6-dimethyltetraoxane
[59183-17-4]

$C_4H_6Cl_2O_4$



Griesbaum, K. *et al.*, *J. Amer. Chem. Soc.*, 1976, **98**, 2880

Extremely sensitive to shock, heat or minor friction, it has exploded very violently when touched with a glass rod or spatula.

See other CYCLIC PEROXIDES

1493. Chromium(II) acetate
[628-52-4]

C₄H₆CrO₄

Cr(OAc)₂

1. Ocone, L. R. *et al.*, *Inorg. Synth.*, 1966, **8**, 129–131

2. Young, G. C., *J. Chem. Educ.*, 1988, **65**, 918–919

The anhydrous salt is pyrophoric in air, or chars slowly in lower oxygen concentrations [1]. A safe and convenient synthesis of the dimeric monohydrate is described [2].

See Bis(2,4-pentanedionato)chromium

See other PYROPHORIC MATERIALS

1494. Chromyl acetate
[4112-22-5]

C₄H₆CrO₆

O₂Cr(OAc)₂


Preparative hazard

See Chromium trioxide: Acetic anhydride

See other OXIDANTS

1495. 3,3-Dinitroazetidinium trifluoromethanesulfonate
[]

C₄H₆F₃N₃O₇S



Hiskey, M. A. *et al.*, *J. Heterocycl. Chem.*, 1992, **29**(7), 1855

The *gem*-dinitro compound is shock sensitive, deflagrating under a heavy hammer blow.

See related POLYNITROALKYL COMPOUNDS

1496. 1,2-Bis(difluoroamino)ethyl vinyl ether
[13084-45-2]

C₄H₆F₄N₂O

F₂NCH₂CH(NF₂)OCH=CH₂

Reed, S. F., *J. Org. Chem.*, 1967, **32**, 2894

It is slightly less impact-sensitive than glyceryl nitrate.

See other DIFLUOROAMINO COMPOUNDS

1497. Di-1,2-bis(difluoroaminoethyl) ether

[13084-46-3]



C₄H₆F₈N₄O


Reed, S. F., *J. Org. Chem.*, 1967, **32**, 2894

It is slightly more impact-sensitive than glyceryl nitrate.

See other DIFLUOROAMINO COMPOUNDS

1498. Divinylmagnesium

[6928-74-1]



C₄H₆Mg


Houben-Weyl, 1973, Vol. 13.2a, 203

The solid may ignite in air.

See related ALKYL METALS

1499. 1-Methylimidazole

[616-47-7]




C₄H₆N₂

Osmium(VIII) oxide

See Osmium(VIII) oxide: 1-Methylimidazole

1500. 2-Amino-4-methyloxazole

[35629-70-0]



C₄H₆N₂O


Hydrogen peroxide

See Hydrogen peroxide: 2-Amino-4-methyloxazole

See related ISOXAZOLES

1501. 3-Amino-5-methylisoxazole

[1072-67-9]



C₄H₆N₂O

Cardillo, P., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 46-48; *Chem. Eng. News*, 1988, **66**(1), 4


The molten compound (100 kg) at 80°C was put into a 200 l drum for storage, but the closed drum exploded after 2 h. Analysis showed the presence of 5-amino-3-methylisoxazole as the main impurity. Thermal stability of the title compound, of the isomeric impurity, and of a mixture of the former with 5% of the latter was investigated by DSC and ARC. Very high self-heating rates were seen, and while the pure 3-amino compound decomposed from 140–160°C, the impure mixture and the prepared mixture of isomers started to decompose at 62°C, the decomposition exotherms being above 1.9 kJ/g in all cases, accompanied by gas evolution. The rapid rate of pressure rise and the final pressure (above 170 bar) which were observed explain the violent bursting of the drum of exothermically decomposing material. The thermal stability of the isomeric 5-amino-3-methylisoxazole was lower than that of the title compound (though the decomposition exotherm was lower at 1.42 kJ/g), and the presence of a few % of the 5-amino compound led to violent exothermic decomposition of the molten compound at 80°C.

See entry ISOXAZOLES See other N–O COMPOUNDS

1502. 5-Amino-3-methylisoxazole

[14678-02-5]

C₄H₆N₂O



See 3-Amino-5-methylisoxazole (next above)

See entry ISOXAZOLES See other N–O COMPOUNDS

1503. Ethyl diazoacetate

[623-73-4]

C₄H₆N₂O₂

N₂CHCO.OEt

1. Searle, N. E., *Org. Synth.*, 1962, Coll. Vol. 4, 426

2. Soos, R. *et al.*, *Chem. Abs.*, 1984, **101**, 230032

It is explosive, and distillation, even under reduced pressure as described, may be dangerous [1]. A Hungarian patent describes a safe procedure for *in-situ* generation of the ester, azeotropic dehydration and subsequent metal-catalysed reaction with 1,3-dienes to give alkyl cyclopropanecarboxylates [2].

Tris(dimethylamino)antimony


Krommes, P. *et al.*, *J. Organomet. Chem.*, 1975, **97**, 63, 65

Interaction at ambient temperature, either initially or on warming from –20°C or below, becomes explosively violent.

See other DIAZO COMPOUNDS

1504. 3-Ethyl-4-hydroxy-1,2,5-oxadiazole (4-Ethyl-3-furazanone)
[34529-29-8]

$C_4H_6N_2O_2$



Sodium hydroxide


Barker, M. D., *Chem. & Ind.*, 1971, 1234

The sodium salt obtained by vacuum evaporation at 50°C of an aqueous alcoholic mixture of the above ingredients exploded violently when disturbed.

See related FURAZAN *N*-OXIDES See other *N*-O COMPOUNDS

1505. 1-Methoxyimidazole *N*-oxide
[90052-22-5]

$C_4H_6N_2O_2$



Flynn, A. P., *Chem. Brit.*, 1984, **20**, 30

Attempted removal of water from a 3 g sample of the hydrate by vacuum distillation at 150–180°C/0.027 mbar caused a violent explosion as distillation started. DSC examination showed a large decomposition exotherm at 140–180°C.

See 1-Methoxy-3,4,5-trimethylpyrazole *N*-oxide, also 1-Hydroxyimidazole *N*-oxide

See other *N*-OXIDES

1506. 2-Cyano-2-propyl nitrate
[40561-27-1]

$C_4H_6N_2O_3$

$Me_2C(CN)ONO_2$

Freeman, J. P., *Org. Synth.*, 1963, **43**, 85


It is a moderately explosive material of moderate impact-sensitivity.

See NITRATING AGENTS

See other ALKYL NITRATES, CYANO COMPOUNDS, *N*-O COMPOUNDS

†1507. 1,2:3,4-Diepoxybutane
[298-18-0]

C₄H₆O₂



See other 1,2-EPOXIDES

1508. 1,1-Dinitro-3-butene
[10229-09-1]

C₄H₆N₂O₄




Witucki, E. F. *et al.*, *J. Org. Chem.*, 1972, **37**, 152

A fume-off during distillation of this compound (b.p., 63–65°C/2 mbar) illustrates the inherent instability of this type of compound.

See other NITROALKENES, POLYNITROALKYL COMPOUNDS

1509. 2,3-Dinitro-2-butene
[28103-68-6]

C₄H₆N₂O₄




Bisgrove, D. E. *et al.*, *Org. Synth.*, 1962, Coll. Vol. 4, 374

Only one explosion has been recorded during vacuum distillation at 135°C/14 mbar.

See other NITROALKENES, POLYNITROALKYL COMPOUNDS

1510. Diacetatoplatinum(II) nitrate
[]

C₄H₆N₂O₁₀Pt




Preparative hazard

See Nitric acid: Acetic acid, etc.

See other PLATINUM COMPOUNDS

1511. 2-Methyl-5-vinyltetrazole
[15284-39-6]

C₄H₆N₄



Aluminium hydride

See Aluminium hydride: Tetrazole derivatives

See other TETRAZOLES

1512. *N,N'*-Dimethyl-*N,N'*-dinitrosooxamide

[7601-87-8]

C₄H₆N₄O₄



Preussmann, R., *Angew. Chem.*, 1963, **75**, 642

Removal of the solvent carbon tetrachloride (in which nitrosation had been effected) at ambient, rather than reduced, pressure caused a violent explosion at the end of distillation. Lowest possible temperatures should be maintained in the preparation. Other precursors seem more suitable as sources of diazomethane.

See Diazomethane

See other NITROSO COMPOUNDS

1513. *N,N'*-Dimethyl-*N,N'*-dinitrooxamide

[14760-99-7]

C₄H₆N₄O₆



Allenby, O. C. W. *et al.*, *Can. J. Res.*, 1947, **25B**, 295–300


It is an explosive.

See other *N*-NITRO COMPOUNDS

1514. Octahydro-2,5-bis(nitroimino)imidazo[4,5-*d*]imidazole (1.3a,4,6a-Tetrahydro-*N,N'*-dinitroimidazo[4,5-*d*]imidazole-2,5-diamine)

[139394-51-7]

C₄H₆N₈O₄



Dagley, I. J. *et al.*, *J. Energ. Mater.*, 1995, **13**(1 & 2), 35

This nitroguanidine analogue is more sensitive than RDX by the drop-weight impact test.

See other *N*-NITRO COMPOUNDS

†1515. 1-Buten-3-one (Methyl vinyl ketone)

[78-94-4]

C₄H₆O



HCS 1980, 665

Haz. Chem. Data, 1975, 207

The uninhibited monomer polymerises on exposure to heat or sunlight. The inhibited monomer may also polymerise if heated sufficiently (by exposure to fire) and lead to rupture of the containing vessel.

See other POLYMERISATION INCIDENTS

Other reactants

Yoshida, 1980, 368

MRH values calculated for 13 combinations with oxidants are given.

†1516. Crotonaldehyde (2-Butenal)
[4170-30-3]

C₄H₆O



HCS 1980, 330

Butadiene

1. Greenlee, K. W., *Chem. Eng. News*, 1948, **26**, 1985
2. Hanson, E. S., *Chem. Eng. News*, 1948, **26**, 2551
3. Watson, K. M., *Ind. Eng. Chem.*, 1943, **35**, 398

An autoclave without a bursting disk and containing the two poorly mixed reactants was wrecked by a violent explosion which occurred on heating the autoclave to 180°C [1]. This was attributed to not allowing sufficient free space for liquid expansion to occur [2]. The need to calculate separate reactant volumes under reaction conditions for all autoclave preparations is stressed [3].

See 1,3-Butadiene

Ethyl acetoacetate

Soriano, D. S. *et al.*, *J. Chem. Educ.*, 1988, **65**, 637

A procedure using a phase-transfer catalyst is employed to prevent the rapid polymerisation of crotonaldehyde during the Robinson annulation reaction.

See related POLYMERISATION INCIDENTS

Nitric acid

See Nitric acid: Crotonaldehyde

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

†1517. 2,3-Dihydrofuran
[1191-91-7]


C₄H₆O



See other PEROXIDISABLE COMPOUNDS

1518. Dimethylketene (2-Methyl-1-propene-1-one)
[598-26-5]

C₄H₆O



1. Smith, G. W. *et al.*, *Org. Synth.*, 1962, Coll. Vol. 4, 348
2. Staudinger, H., *Ber.*, 1925, **58**, 1079–1080

Dimethylketene rapidly forms an extremely explosive peroxide when exposed to air at ambient temperatures. Drops of solution allowed to evaporate may explode. Inert atmosphere should be maintained above the monomer [1]. The peroxide is

polymeric and very sensitive, exploding on friction at -80°C . Higher homologues are very unstable and unisolable [2].

See other FRICTIONAL INITIATION INCIDENTS


Oxygen

See Poly(peroxyisobutyrolactone)

See other POLYPEROXIDES

†1519. Divinyl ether (1,1'-Oxybisethene)

[109-93-3]



$\text{C}_4\text{H}_6\text{O}$

Anon., *Chemist & Druggist*, 1947, **157**, 258

The presence of *N*-phenyl-1-naphthylamine as inhibitor considerably reduces the development of peroxide in the ether.


Nitric acid

See Nitric acid: Divinyl ether

See other PEROXIDISABLE COMPOUNDS

†1520. 3,4-Epoxybutene (Vinyloxirane)

[930-22-3]



$\text{C}_4\text{H}_6\text{O}$

See other 1,2-EPOXIDES

†1521. Ethoxyacetylene (Ethoxyethyne)

[927-80-0]



$\text{C}_4\text{H}_6\text{O}$

Jacobs, T. L. *et al.*, *J. Amer. Chem. Soc.*, 1942, **64**, 223

Small samples rapidly heated in sealed tubes to around 100°C exploded.

Ethylmagnesium iodide

See Ethylmagnesium iodide: Ethoxyacetylene

See other ACETYLENIC COMPOUNDS

†1522. Methacrylaldehyde (2-Methylpropenal)

[78-85-3]



$\text{C}_4\text{H}_6\text{O}$

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

1523. 3-Methoxypropyne (Methyl propargyl ether)
[627-41-8]

C₄H₆O



Brandsma, 1971, 13

It explodes on distillation at its atmospheric b.p., 61°C.

See other ACETYLENIC COMPOUNDS

†**1524. Allyl formate (3-Propenyl methanoate)**
[1838-59-1]

C₄H₆O₂



See other ALLYL COMPOUNDS

†**1525. Butane-2,3-dione**
[431-03-8]

C₄H₆O₂



HCS 1980, 228

1526. 2-Butyne-1,4-diol
[110-65-6]

C₄H₆O₂



Alkalies, or Halide salts, or Mercury salts and acids


Kirk-Othmer, 1960, Suppl. Vol. 2, 45

The pure diol may be distilled unchanged, but traces of alkali or alkaline earth hydroxides or halides may cause explosive decomposition during distillation. In presence of strong acids, mercury salts may cause violent decomposition of the diol.

See other ACETYLENIC COMPOUNDS, CATALYTIC IMPURITY INCIDENTS

1527. Butyrolactone (2(3H)-Dihydrofuranone)
[96-48-0]

C₄H₆O₂



Butanol, 2,4-Dichlorophenol, Sodium hydroxide

Anon., *CISHC Chem. Safety Summ.*, 1977, **48**, 3

In an altered process to prepare 2,4-dichlorophenoxybutyric acid, the lactone was added to the other components, and soon after, the reaction temperature reached 165°C, higher than the usual 160°C. Application of cooling failed to check the

thermal runaway, and soon after reaching 180°C the vessel began to fail and an explosion and fire occurred.

See other RUNAWAY REACTIONS

1528. Dimethyl azoformate (Dimethyl diazenedicarboxylate)
[2446-84-6]

C₄H₆N₂O₄



1. US Pat. 3 347 845, 1967
2. Kauer, J. C., *Org. Synth.*, 1962, Coll. Vol. 4, 412
3. Young, J. A., private communication, 1993
4. Diels, O. *et al.*, *Berichte*, 1913, **46**, 2006

It is shock-sensitive [1]. An undergraduate student was preparing a 200 g batch following the Organic Syntheses procedure [2]. The first stage quality is believed to have been poor. The final product exploded during vacuum distillation, costing her the sight of an eye [3]. Since the first man to describe it compared it with guncotton, when heated under slight confinement [4], it might be considered unsuitable for undergraduates. The editor suspects that hydrazoic acid reported as an hydrolysis product may indicate that procedure [2] can produce methyl azidoformate as a byproduct. Diels [4] also demonstrated formation of another very unstable compound from its further reaction with nitric acid, which is the oxidant used in the second stage of preparation [2].

See Ethyl azidoformate

See Diethyl azoformate

See other AZO COMPOUNDS

†**1529. 1,2:3,4-Diepoxybutane**
[298-18-0]

C₄H₆O₂



See other 1,2-EPOXIDES

1530. Methacrylic acid (2-Methylpropenoic acid)
[79-41-4]

C₄H₆O₂



1. Anon., *CISHC Chem. Safety Summ.*, 1979, **50**, 34
2. Bond, J., *Loss Prev. Bull.*, 1991, **101**, 1
3. Anderson, S. E. *et al.*, *Plant/Oper. Progress*, 1992, **11**(3), 151
4. Nicolson, A., *Plant/Oper. Progress.*, 1991, **10**(3), 171

A drum of the uninhibited acid (m.p., 16°C) which had been stored outside under winter conditions was transferred into a warm room to liquefy the acid. Later, exothermic polymerisation led to bulging of the drum and leakage of the acid

vapour [1]. A more serious accident involved a railtank of acid-washed crude technical methacrylic acid (liable to throw down a dilute H₂SO₄ layer) which exploded, throwing debris 300 metres, some 20 hrs after a pressure relief valve was seen to lift, despite being sprayed with water meanwhile [2,3]. It was shipped in unlined steel without corrosion tests because that had previously been used for purified inhibited product and it contained negligible stabiliser. A study of the complex interactions of oxygen (inhibitory at low but not at high concentrations) and stabilisers is reported [3]. Phenothiazine seems a better inhibitor than the usual phenols.

See Acrylic acid

See other CORROSION INCIDENTS

See other ORGANIC ACIDS, CORROSION INCIDENTS, POLYMERISATION INCIDENTS

†1531. Methyl acrylate (Methyl propenoate)

[96-33-3]

C₄H₆O₂



(MCA SD-79, 1960); HCS 1980, 634

1. MCA Case History No. 2033
2. Harmon, 1974, 2.11
3. *See entry* SELF-ACCELERATING REACTIONS

A 4 l glass bottle of the ester (sealed and partly polymerised old stock) exploded several hours after being brought from storage into a laboratory. An inhibitor had originally been present, but could have been consumed during prolonged storage, and peroxides may have initiated exothermic polymerisation of the remaining monomer [1]. The monomer is normally stored and handled inhibited and at below 10°C, but not under inert atmosphere, because traces of oxygen are essential to the inhibition process [2]. DSC investigation showed that the temperature-dependent induction period (425 min/125°C, 25/165, 6/185, in which the stabiliser becomes consumed) is followed by autocatalytic and rapidly accelerating polymerisation, with $Q = 0.89$ kJ/g [3].

See VIOLENT POLYMERISATION

See other GLASS INCIDENTS, INDUCTION PERIOD INCIDENTS, POLYMERISATION INCIDENTS

Other reactants

Yoshida, 1980, 10

MRH values calculated for 13 combinations with oxidants are given.

†1532. Vinyl acetate (Ethenyl ethanoate)

[108-05-4]

C₄H₆O₂



(MCA SD-75, 1970); FPA H36, 1975; HCS 1980, 955

1. Harmon, 1974, 2.19
2. Levy, L. B., *Process Safety Progress*, 1993, **12** (1), 47
3. Gustin, J. L. *et al.*, *Chem. Abs.*, 1998, **129**, 264630g

The monomer is volatile and tends to self-polymerise, and is therefore stored and handled cool and inhibited, with storage limited to below 6 months. Several industrial explosions have been recorded [1]. Unlike acrylic monomers, oxygen is not involved in stabilisation and is detrimental at higher temperatures [2]. The polymerisation has been modelled and causes of accidents proposed [3].

Air, Water

MCA SD-75, 1970

Vinyl acetate is normally inhibited with hydroquinone to prevent polymerisation. A combination of too low a level of inhibitor and warm, moist storage conditions may lead to spontaneous polymerisation. This process involves autoxidation of acetaldehyde (a normal impurity produced by hydrolysis of the monomer) to a peroxide which initiates exothermic polymerisation as it decomposes. In bulk, this may accelerate to a dangerous extent. Other peroxides or radical sources will initiate the exothermic polymerisation.

Desiccants

MCA SD-75, 1970

Vinyl acetate vapour may react vigorously in contact with silica gel or alumina.

Dibenzoyl peroxide, Ethyl acetate

Vervalin, 1973, 81

Polymerisation of the ester with dibenzoyl peroxide in ethyl acetate accelerated out of control and led to discharge of a large volume of vapour which ignited and exploded.

See other POLYMERISATION INCIDENTS

Ethylene

See Ethylene: Vinyl acetate

Hydrogen peroxide

MRH 6.19/78

See Hydrogen peroxide: Vinyl acetate

Other reactants

Yoshida, 1980, 122

MRH values calculated for 15 combinations with oxidants are given.

Oxygen

Barnes, C. E. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 210

The unstabilised polymer exposed to oxygen at 50°C generated an ester-oxygen interpolymeric peroxide which, when isolated, exploded vigorously on gentle heating.

See other POLYPEROXIDES

Ozone

See Vinyl acetate ozonide

Toluene

MCA Case History No. 2087

The initial exotherm of a solution polymerisation of the ester in boiling toluene in a 10 m³ reactor was too great for the cooling and vent systems, and the reaction began to accelerate out of control. Failure of a gasket released a quantity of the flammable reaction mixture which became ignited and destroyed the containing building.

See other POLYMERISATION INCIDENTS, PEROXIDISABLE COMPOUNDS

1533. Poly(1,3-butadiene peroxide)

[28655-95-0]

(C₄H₆O₂)_n

Complex structure

Butadiene

1. Alexander, D. S., *Ind. Eng. Chem.*, 1959, **51**, 733
2. Hendry, D. G. *et al.*, *Ind. Eng. Chem., Prod. Res. Dev.*, 1968, **7**, 136, 145
3. *ASTM D1022-76*
4. Bailey, H. C., private comm., 1974
5. *CISHC Chem. Safety Summ.*, 1980, **51**, 39
6. *Sichere Chemiearb.*, 1980, **32**, 6–8
7. *MCA Case History No. 2270*

A violent explosion in a partially full butadiene storage sphere was traced to butadiene peroxide. The latter had been formed by contact with air over a long period, and had eventually initiated the exothermic polymerisation of the sphere contents. A tank monitoring and purging system was introduced to prevent a recurrence [1]. The polyperoxide (formed even at 0°C) is a mixed 1,2- and 1,4-addition copolymer of butadiene with oxygen [2], effectively a dialkyl peroxide. Its concentration in butadiene is very seriously underestimated by a standard method applicable to hydroperoxides [3], which indicates only 5% of the true value. Alternative methods of greater reliability are likely to form the basis of a revised standard method currently under consideration [4]. A bolt falling into an 'empty' rail tanker during maintenance work led to a serious explosion. Other rail tankers used to convey unstabilised butadiene were later found to contain several kg of the gelatinous polyperoxide [5]. Peroxide formation in uninhibited monomer is very rapid, air-saturated monomer giving in 24 h 124 ppm of peroxide/18°C and 460 ppm/50°C. The peroxide ($n = 7-9$) has very low solubility in the monomer, and separates as a lower layer which is very shock-sensitive and powerfully explosive [6]. Solidified peroxide plugging a tube condenser decomposed explosively during drilling operations to clear the tubes [7].

See 1,3-Butadiene

See other POLYPEROXIDES

1534. Acetic anhydride

[108-24-7]

C₄H₆O₃

MeCO.OCO.Me

(*MCA SD-15*, 1962); *HCS 1980*, 101

The principal reaction hazard attached to use of acetic anhydride is the possibility of rapid and exothermic acid-catalysed hydrolysis unless the conditions prevailing (temperature, agitation, order of mixing, proportion of water) are such as to promote smooth and progressive hydrolysis with adequate heat removal. The examples below illustrate these factors.

Acetic acid, Water

MCA Case History No. 1865

Erroneous addition of aqueous acetic acid into a tank of the anhydride caused violent exothermic hydrolysis of the latter.

See Water, below

Ammonium nitrate, Hexamethylenetetraminium acetate, Nitric acid

See Nitric acid: Acetic anhydride, etc.

Barium peroxide

See Barium peroxide: Acetic anhydride

Boric acid

1. Lerner, L. M., *Chem. Eng. News*, 1973, **51**(34), 42

2. *Experiments in Organic Chemistry*, 281, Fieser, L. F., Boston, Heath, 1955

Attempted preparation of acetyl borate by slowly heating a stirred mixture of the anhydride and solid acid led to an eruptive explosion at 60°C [1]. The republished procedure being used [2] omitted the reference to a violent reaction mentioned in the German original. Modifying the procedure by adding portions of boric acid to the hot stirred anhydride should give a smoother reaction.

Bromohydroxybiphenyl, Pyridine, Water

Nolan, 1983, Case History 36

Bromohydroxybiphenyl was *O*-acetylated with excess acetic anhydride in pyridine. The excess anhydride was to be hydrolysed by addition of water, but this was done without proper control using a hosepipe. The hydrolysis reaction ran away, causing boiling and evaporation of the reactor contents.

See Acetic acid, above; Water, below

N-*tert*-Butylphthalamic acid, Tetrafluoroboric acid

Boyd, G. V. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1978, 1346

Interaction to give *N*-*tert*-butylphthalisoimidium tetrafluoroborate was very violent, possibly because of exothermic hydrolysis of the anhydride by the 40% aqueous tetrafluoroboric acid.

See Tetrafluoroboric acid, below

Chromic acid

1. Dawkins, A. E., *Chem. & Ind.*, 1956, 196

2. Baker, W., *Chem. & Ind.*, 1956, 280

C₄H₆O₃

Addition of acetic anhydride to a solution of chromium trioxide in water caused violent boiling [1], due to the acid-catalysed exothermic hydrolysis of the anhydride [2].

Chromium trioxide MRH 2.38/84
See Chromium trioxide: Acetic anhydride

1,3-Diphenyltriazene
See 1,3-Diphenyltriazene: Acetic anhydride

Ethanol, Sodium hydrogen sulfate
Staudinger, H., *Angew. Chem.*, 1922, **35**, 657
Accidental presence of the acid salt vigorously catalysed a large scale preparation of ethyl acetate, causing violent boiling and emission of vapour which became ignited and exploded.
See other CATALYTIC IMPURITY INCIDENTS

Glycerol, Phosphoryl chloride
Bellis, M. P., *Hexagon Alpha Chi Sigma* (Indianapolis), 1949, **40**(10), 40
Violent acylation occurs in catalytic presence of phosphoryl chloride, because the high viscosity of the mixture in absence of solvent prevents mixing and dissipation of the high heat of reaction.

Hydrochloric acid, Water
1. Vogel, 1957, 572–573
2. Batchelor, J. F., private comm., 1976
Crude dimethylaniline was being freed of impurities by treatment with acetic anhydride according to a published procedure [1]. However, three times the recommended proportion of anhydride was used, and the reaction mixture was ice cooled before addition of diluted hydrochloric acid to hydrolyse the excess anhydride. Hydrolysis then proceeded with explosive violence.

Hydrogen peroxide MRH 5.52/71
See Hydrogen peroxide: Acetic anhydride

Hypochlorous acid
See Hypochlorous acid: Acetic anhydride

Metal nitrates
1. Davey, W. *et al.*, *Chem. & Ind.*, 1948, 814
2. Collman, J. P. *et al.*, *Inorg. Synth.*, 1963, **7**, 205–207
3. James, B. D., *J. Chem. Educ.*, 1974, **51**(9), 568
Use of mixtures of metal nitrates with acetic anhydride as a nitrating agent may be hazardous, depending on the proportions of reactants and on the cation; copper nitrate or sodium nitrate usually cause violent reactions [1]. An improved procedure for the use of the anhydride–copper(II) nitration mixture [2] has been further modified [3] to improve safety aspects.

Nitric acid

MRH 4.93/66

See Nitric acid: Acetic anhydride

Other reactants

Yoshida, 1980, 356

MRH values calculated for 16 combinations with oxidants are given.

Perchloric acid, Water

1. Anon., *ABCM Quart. Safety Summ.*, 1954, **25**, 3

2. Turner, H. S. *et al.*, *Chem. & Ind.*, 1965, 1933

3. McG. Tegart, W. J., *The Electrolytic and Chemical Polishing of Metals in Research and Industry*, London, Pergamon, 2nd Ed., 1959

Anhydrous solutions of perchloric acid in acetic acid are prepared by using acetic anhydride to remove the diluting water from 72% perchloric acid. It is essential that the anhydride is added slowly to the aqueous perchloric-acetic acid mixture under conditions where it will react readily with the water, i.e. at about 10°C. Use of anhydride cooled in a freezing mixture caused delayed and violent boiling to occur. Full directions for the preparation are given [1]. A violent explosion occurred during the preparation of an electropolishing solution by addition of perchloric acid solution to a mixture of water and acetic anhydride. The cause of the explosion, the vigorously exothermic acid-catalysed hydrolysis of acetic anhydride, is avoided if water is added last to the mixture produced by adding perchloric acid solution to acetic anhydride [2]. The published directions [3] are erroneous and insufficiently detailed for safe working.

See Perchloric acid: Acetic anhydride, or: Dehydrating agents

Peroxyacetic acid

See Peroxyacetic acid: Acetic anhydride

Polyphosphoric acid, Water

Montfort, B. *et al.*, *Bull. Soc. Chim. Fr.*, 1987, 848–854

2-Aryloxy-1,2-diarylethanones can be cyclodehydrated to diarylbenzofurans by heating with sodium acetate and acetic anhydride in polyphosphoric acid. Quenching the hot reaction mixture with water leads to initially violent acid-catalysed hydrolysis of the excess anhydride.

Potassium permanganate

MRH 2.55/86

See Potassium permanganate: Acetic acid, etc.

Sodium percarbonate

See Sodium carbonate hydrogen peroxidate: Acetic anhydride

Tetrafluoroboric acid

1. Lichtenberg, D. W. *et al.*, *J. Organomet. Chem.*, 1975, **94**, 319

2. Wudl, F. *et al.*, *Inorg. Synth.*, 1979, **19**, 29

Dehydration of the aqueous 48% acid by addition to the anhydride is rather exothermic, and caution is advised [1]. Operation at 0°C led to an explosion [2].

See Chromic acid, also Hydrochloric acid, also Perchloric acid, all above

4-Toluenesulfonic acid, Water

1. Jones, B. J. *et al.*, *Clin. Chem.*, 1955, **1**, 305
2. Pearson, S. *et al.*, *Anal. Chem.*, 1953, **25**, 813–814

Caution is advised [1] to prevent explosions when using an analytical method involving sequential addition of acetic acid, aqueous 4-toluenesulfonic acid and acetic anhydride to serum [2]. It is difficult to see why this should happen, unless the anhydride were all added before the sulfonic acid solution.

See Water, below

Water

1. Leigh, W. R. D. *et al.*, *Chem. & Ind.*, 1962, 778
2. Benson, G., *Chem. Eng. News*, 1947, **25**, 3458

Accidental slow addition of water to a mixture of the anhydride and acetic acid (85:15) led to a violent, large scale explosion. This was simulated closely in the laboratory, again in the absence of mineral-acid catalyst [1]. If unmoderated, the rate of acid-catalysed hydrolysis of (water insoluble) acetic anhydride can accelerate to explosive boiling [2]. Essentially the same accident, fortunately with no injuries or fatalities this time, was repeated in 1990.

See other ACID ANHYDRIDES

1535. Peroxycrotonic acid (Peroxy-2-butenic acid)

[5813-77-4]

C₄H₆O₃



Vasilina, T. U. *et al.*, *Chem. Abs.*, 1974, **81**, 151446

The decomposition of the acid has been mentioned in a safety context, but the details were not translated.

See other PEROXYACIDS

1536. Poly(peroxyisobutyrolactone) (Poly(dimethylketene peroxide))

[67772-28-5]

(C₄H₆O₃)_n



Turro, N. J. *et al.*, *J. Amer. Chem. Soc.*, 1978, **100**, 5580

Autoxidation of dimethylketene with oxygen in ether at -20°C gives the poly(peroxylactone) which as a dry solid is liable to undergo unpredictable and violent detonation.

See Dimethylketene

See related PEROXYESTERS

See other POLYPEROXIDES

1537. Diacetyl peroxide

[110-22-2]

C₄H₆O₄



1. Shanley, E. S., *Chem. Eng. News*, 1949, **27**, 175
2. Kuhn, L. P., *Chem. Eng. News*, 1948, **26**, 3197

3. Kharasch, M. S. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 1269

4. Moore, C. G., *J. Chem. Soc.*, 1951, 236

Acetyl peroxide may readily be prepared and used in ethereal solution. It is essential to prevent separation of the crystalline peroxide even in traces, since, when dry, it is shock-sensitive and a high explosion risk [1]. Crystalline material, separated and dried deliberately, detonated violently [2]. The commercial material, supplied as a 30% solution in dimethyl phthalate, is free of the tendency to crystallise and is relatively safe. It is, however, a powerful oxidant [1]. Precautions necessary for the preparation and thermolysis of the peroxide have been detailed [3,4].

See Fluorine: Sodium acetate


Other reactants

Yoshida, 1980, 134

MRH values calculated for 15 combinations with oxidisable elements and compounds are given.

See other DIACYL PEROXIDES

1538. *cis*-1,4-Dioxenedioxetane (2,5,7,8-Tetraoxa[4.2.0]bicyclooctane) [59261-17-5]




Wilson, T. *et al.*, *J. Amer. Chem. Soc.*, 1976, **98**, 1087

A small sample of the solid peroxide exploded on warming to ambient temperature. Storage at -20°C in solution appears safe.

See other CYCLIC PEROXIDES

1539. Poly(vinyl acetate peroxide)

[]




See Vinyl acetate: Oxygen

See other POLYPEROXIDES

1540. Lead acetate–lead bromate

[]



1. Berger, A., *Arbeits-Schutz.*, 1934, **2**, 20

2. Leymann, –, *Chem. Fabrik*, 1929, 360–361

The compound (formulated as the double salt, (fuel + oxidant), rather than the mixed salt) may be formed during the preparation of lead bromate from lead acetate and

potassium bromate in acetic acid, and is explosive and very sensitive to friction. Although lead bromate is stable up to 180°C, it is an explosive salt [1]. Further details of the incident are available [2].

See related METAL OXOHALOGENATES

1541. Palladium(II) acetate
[375-31-3]




Phenylacetylene

See Phenylacetylene: Palladium(II) acetate

See related PLATINUM COMPOUNDS

1542. Monoperoxsuccinic acid
[3504-13-0]





1. Lombard, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, **12**, 2800

2. Castrantas, 1965, 16

It explodes in contact with flame [1] and is weakly shock-sensitive [2].

See other PEROXYACIDS

1543. Vinyl acetate ozonide (3-Acetoxy-1,2,4-trioxolane)
[101672-23-5]




Kirk-Othmer, 1970, Vol. 21, 320

The ozonide formed by vinyl acetate is explosive when dry.

See other OZONIDES

1544. Dimethyl peroxydicarbonate
[15411-45-7]



Explodes on heating to 55–60°C, or readily under a hammer blow, and more powerfully than dibenzoyl peroxide.

See entry PEROXYCARBONATE ESTERS

1545. Tartaric acid (2,3-Dihydroxybutanedioic acid)
[87-69-4] **C₄H₆O₆**
(-CHOHCO.OH)₂

HCS 1980, 883

Silver

See Silver: Carboxylic acids

See other ORGANIC ACIDS

1546. 2-Butyne-1-thiol
[101672-05-3] **C₄H₆S**
EtC≡CSH

Brandsma, 1971, 180

Presence of a stabiliser is essential during handling or storage at -20°C under nitrogen. Exposure to air leads to formation of a polymer which may explode on heating.

See other ACETYLENIC COMPOUNDS

1547. Divinylzinc
[1119-22-8] **C₄H₆Zn**
(H₂C=CH)₂Zn

491M, 1975, 164

It ignites in air.

See related ALKYL METALS

†**1548. 1-Bromo-2-butene**
[4784-77-4] **C₄H₇Br**
BrCH₂CH=CHMe

See other ALLYL COMPOUNDS, HALOALKENES

†**1549. 4-Bromo-1-butene**
[5162-44-7] **C₄H₇Br**
H₂C=CHCH₂CH₂Br

Chloromethylphenylsilane, Chloroplatinic acid

See Chloromethylphenylsilane: 4-Bromobutene, etc.

See other HALOALKENES


1550. 1,1,1-Tris(bromomethyl)methane (3-(Bromomethyl)-1,3-dibromopropane)
[62127-48-4] **C₄H₇Br₃**



Preparative hazard

See Phosphorus tribromide: 1,1,1-Tris(hydroxymethyl)methane

†**1551. 2-Chloro-2-butene**
[4461-41-0] **C₄H₇Cl**



See other HALOALKENES

†**1552. 3-Chloro-1-butene**
[563-52-0] **C₄H₇Cl**




See other HALOALKENES

†**1553. 3-Chloro-2-methyl-1-propene**
[563-47-3] **C₄H₇Cl**



See other HALOALKENES

1554. N-Chloro-4-methyl-2-imidazolinone
[55341-15-6] **C₄H₇ClN₂O**



Walles, W. E., US Pat. 3 850 920, 1974

It exploded after several hours at ambient temperature.


See other N-HALOGEN COMPOUNDS

†**1555. Butyryl chloride (Butanoyl chloride)**
[141-75-3] **C₄H₇ClO**



See other ACYL HALIDES

1556. 1-Chloro-2-butanone
[616-27-3]



$\text{C}_4\text{H}_7\text{ClO}$

Tilford., C. H. private comm., 1965
An amber bottle of stabilised material spontaneously exploded.
See Chloroacetone

†**1557. 2-Chloroethyl vinyl ether (2-Chloroethoxyethene)**
[110-75-8]



$\text{C}_4\text{H}_7\text{ClO}$

See other PEROXIDISABLE COMPOUNDS

†**1558. Isobutyryl chloride (2-Methylpropanoyl chloride)**
[79-30-1]



$\text{C}_4\text{H}_7\text{ClO}$

See other ACYL HALIDES

1559. Ethyl chloroacetate
[105-39-5]



$\text{C}_4\text{H}_7\text{ClO}_2$

HCS 1980, 475

Sodium cyanide

Brown, E. A. B., private comm., 1982
A well-established procedure (1 g mol, 20 runs) for preparation of ethyl cyanoacetate by heating the reactants together suddenly erupted out of control.

†**1560. Isopropyl chloroformate**
[108-23-6]



$\text{C}_4\text{H}_7\text{ClO}_2$

1. *491M*, 1975, 223

2. Young, J. A., *Chem. Health Saf.*, 1998, **5** (5), 4

A sample exploded in refrigerated storage. It is known that if iron salts are present during the preparation from 2-propanol and phosgene, catalysed thermal decomposition may occur [1]. A 2 litre glass bottle, presumably less than a year old and stored unrefrigerated at eye-level, exploded during stocktaking, splashing the eyes and face of a researcher [2]. Iron is usually present in glass, especially brown glass, while most eye protection is designed primarily to stop fast particles and shields poorly against slow liquids.

See other ACYL HALIDES, CATALYTIC IMPURITY INCIDENTS

1561. Isopropylisocyanide dichloride

[29119-58-2]

C₄H₇Cl₂N

Iron(III) chloride, Metal oxides, or Water

Fuks, R. *et al.*, *Tetrahedron*, 1973, **29**, 297–298

Interaction of the 3:2 complex with iron(III) chloride and calcium oxide, mercury oxide or silver oxide was usually too violent for preparative purposes, but zinc oxide was satisfactory. Reaction with water was violent.

See related CYANO COMPOUNDS**1562. 1-Fluoro-1,1-dinitrobutane**

[19273-47-3]

C₄H₇FN₂O₄*See entry* FLUORODINITROMETHYL COMPOUNDS†**1563. Butyronitrile (Butanonitrile)**

[109-74-0]

C₄H₇N*See other* CYANO COMPOUNDS**1564. (Dimethylamino)acetylene (N,N-Dimethylethynamine)**

[24869-88-3]

C₄H₇N

Water

Brandsma, 1971, 139

The amine reacts extremely vigorously with neutral water.

See other ACETYLENIC COMPOUNDS, ORGANIC BASES†**1565. Isobutyronitrile (2-Methylpropanonitrile)**

[78-82-0]

C₄H₇N*See other* CYANO COMPOUNDS**1566. 2-Cyano-2-propanol**

[75-86-5]

C₄H₇NO*HCS 1980*, 103


Sulfuric acid

See Sulfuric acid: 2-Cyano-2-propanol

See other CYANO COMPOUNDS

1567. 3,6-Dihydro-1,2,2H-oxazine
[3686-43-9]

C₄H₇NO



Nitric acid

See Nitric Acid: 3,6-Dihydro-1,2,2H-oxazine

See other N-O COMPOUNDS

1568. 2-Methylacryaldehyde oxime
[28051-68-5]

C₄H₇NO




Mowry, D. T. *et al.*, *J. Amer. Chem. Soc.*, 1947, **69**, 1831

The viscous distillation residue may decompose explosively if overheated. A polymeric peroxide may have been involved.

See other OXIMES, POLYPEROXIDES

†**1569. 2-Methyl-2-oxazoline**
[1120-64-5]


C₄H₇NO



See other N-O COMPOUNDS

1570. 2,3-Butanedione monoxime
[57-71-6]

C₄H₇NO₂



Muir, G. D., private comm., 1968

Distillation from a mantle heated flask at 6.5 mbar caused explosions on several occasions. Distillation from a steam heated flask at below 1.3 mbar appeared safe.

See other OXIMES

1571. N-Hydroxymethylacrylamide (N-Hydroxymethyl-2-propenamide)
[924-42-5] **C₄H₇NO₂**



491M, 1975, 213

Excessive heat, smoke, flames and crackling noises were emitted from stored fibre drums of the monomer, some unopened. Polymerisation may have been initiated by minor contaminants (perhaps as vapours), and/or by excessively warm storage conditions.

See other POLYMERISATION INCIDENTS

1572. 4-Nitro-1-butene
[32349-29-4] **C₄H₇NO₂**



Witucki, E. F. *et al.*, *J. Org. Chem.*, 1972, **37**, 152

A fume-off during distillation of one sample of this compound (b.p., 55°C/25 mbar) illustrates the inherent instability of this type of compound.

See other NITROALKENES

1573. Butyryl nitrate
[101672-06-4] **C₄H₇NO₄**



Francis, F. E., *Ber.*, 1906, **39**, 3798

It detonates on heating.

See other ACYL NITRATES

1574. Butyryl peroxonitrate
[5796-89-4] **C₄H₇NO₅**




Stephens, E. R., *Anal. Chem.*, 1964, **36**, 928

It is highly explosive, like its lower homologues.

See related PEROXYESTERS

1575. 4-Hydroxy-3,5-dimethyl-1,2,4-triazole
[35869-74-0] **C₄H₇N₃O**




Armand, J. *et al.*, *J. Chem. Res. (M)*, 1980, 3868

It explodes on melting at 122°C.
See other N-O COMPOUNDS, TRIAZOLES


1576. 1,3-Bis(5-amino-1,3,4-triazol-2-yl)triazene
[3751-44-8]

C₄H₇N₁₁



Hauser, M., *J. Org. Chem.*, 1964, **29**, 3449
It explodes at the m.p., 187°C.
See other HIGH-NITROGEN COMPOUNDS, TRIAZOLES

†**1577. 1-Butene**
[106-98-9]




C₄H₈

HCS 1980, 229 (cylinder)

Aluminium tetrahydroborate
See Aluminium tetrahydroborate: Alkenes, etc.
See other ALKENES


†**1578. cis-2-Butene**
[624-64-6]



C₄H₈

See other ALKENES

†**1579. trans-2-Butene**
[590-18-1]



C₄H₈

See other ALKENES

†**1580. Cyclobutane**
[287-23-0]

C₄H₈



†1581. Methylcyclopropane
[594-11-6]


C₄H₈



See other STRAINED-RING COMPOUNDS

†1582. 2-Methylpropene (Isobutene)
[115-11-7]

C₄H₈



FPA H108, 1981; HCS 1980, 566

See other ALKENES

1583. Poly(isobutene)
[9003-27-4]

(C₄H₈)_n




Silver peroxide

See Silver peroxide: Poly(isobutene)

1584. N-Chloropyrrolidine
[19733-68-7]

C₄H₈ClN



It explodes at ambient temperature.

See entry N-HALOGEN COMPOUNDS (reference 4)

1585. N,N-Dimethyl-2-chloroacetamide
[2675-89-0]

C₄H₈ClNO




Energy of decomposition (in range 180–440°C) measured as 0.59 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

1586. 4-Morpholinesulfenyl chloride
[2958-89-6]

C₄H₈ClNOS



MCA Case History No. 1806

The product from chlorination of 4,4'-dithiodimorpholine exploded violently after vacuum stripping of the solvent carbon tetrachloride. As it was a published process that previously had been operated uneventfully, no cause was apparent. It seems remotely possible that an unstable *N*-chloro derivative could have been produced by N–S bond cleavage if chlorination conditions had differed from those previously employed.

See other N–S COMPOUNDS

†**1587. mixo-Dichlorobutane**
[26761-81-9]


C₄H₈Cl₂



See other HALOALKANES

1588. 1,1-Bis(dimethyl sulfoxide)telluracyclopentane diperchlorate
[100242-83-9]

C₄H₈Cl₂O₈·2C₂H₆OS



Srivastava, T. *et al.*, *Polyhedron*, 1985, **4**, 1223–1229

The complex explodes at its m.p., 116°C.

See other ORGANOMETALLIC PERCHLORATES

1589. Bis(2-chloroethyl)sulfide (1,1'-Thiobis(2-chloroethane))
[505-60-2]

C₄H₈Cl₂S



Bleaching powder

See BLEACHING POWDER

1590. N-Chloro-bis(2-chloroethyl)amine
[63915-60-6]

C₄H₈Cl₃N



Butters, M. *et al.*, UK Pat. Appl. GB 2 165 244, 1986

The *N*-chloro derivative is potentially explosive when neat, often exploding during distillation at 60°C/2.6 mbar.

See other *N*-HALOGEN COMPOUNDS

1591. Bis(1-chloroethylthallium chloride) oxide

[]



Yakubovich, A. Ya. *et al.*, *Doklad. Akad. Nauk SSSR*, 1950, **73**, 957–958

An explosive solid of low stability.

See related ALKYL METALS, METAL HALIDES, METAL OXIDES

1592. 2-Methyl-1-nitratodimercurio-2-nitratomercuriopropane

[]



Whitmore, 1921, 116

The compound, prepared from isobutene and mercury(II) nitrate, explodes on impact at 80°C.

See other ORGANOMETALLIC NITRATES

1593. 1,1'-Biaziridinyl

[4388-03-8]



Oxygen


Graefe, A. F., *J. Amer. Chem. Soc.*, 1958, **80**, 3941

It exploded violently during analytical combustion in oxygen.

See other AZIRIDINES

1594. 3-Propyldiazirine

[70348-66-2]



Schmitz, E. *et al.*, *Ber.*, 1962, **95**, 800

It exploded on attempted distillation from calcium chloride at about 75°C.

See other DIAZIRINES

1595. Butane-2,3-dione dioxime
[95-45-4]

$C_4H_8N_2O_2$



Energy of decomposition (in range 220–360°C) measured as 1.98 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other OXIMES

**1596. 1-Methylamino-1-methylthio-2-nitroethene (*N*-Methyl-1-(methylthio)-
2-nitroethenamine)**
[61832-41-5]

$C_4H_8N_2O_2S$



Preparative hazard

1. Koshy, A. *et al.*, *J. Hazard. Mater.*, 1995, 42(3), 215
Goulding, J. P. in *Safety and Runaway Reactions*, Smeder, B & Mitchison, N. (Eds.),
EUR 17723 EN, European Commission, 1997, p31

This pharmaceutical intermediate, itself unstable, is prepared via intermediates which are actually explosive. A calorimetric study of risks is reported [1]. The byproducts are also hazardous, a study into an explosion and fire consequent upon isopropanol (solvent) recovery from a manufacturing process is reported. This was done by distillation, leaving a concentrated residue, more than one distillation being performed before the residue was discharged for incineration. A fortnight's accumulated residue, as about a 25% solution or suspension in isopropanol was left in the reactor for a week. This was probably below minimum stir volume and below the level of the temperature probe. During this week, there was a power cut which appears to have stopped the flow of coolant to the reactor; the coolant coils may in any event not have dipped far into the residue. Two days later, shortly preceded by some indications of modest warming, the explosion occurred, rupturing storage tanks, igniting their contents and thus causing considerable secondary damage. Calorimetry suggested that the deposited solids became unsafe at 40°C, the stirred bulk from about 65°C. It was concluded that both the design and operation of the recovery process had been inadequate. (Elsewhere in this book, an account of another explosion and destructive sequence of secondary fires, when treating wastes from a slightly different manufacturing process, is to be found. In this other case neither product nor byproducts were directly responsible — Ed.)

See Nitromethane; Carbon disulfide

See other C-NITRO COMPOUNDS

1597. Ethyl *N*-methyl-*N*-nitrosocarbamate
[615-53-2]

$C_4H_8N_2O_3$




1. Hartman, W. W., *Org. Synth.*, 1943, Coll. Vol. 2, 465
2. Druckrey, H. *et al.*, *Nature*, 1962, **195**, 1111

The material is unstable and explodes if distilled at ambient pressure [1], or may become explosive if stored above 15°C [2]. Its use for preparation of diazomethane has been superseded by more stable intermediates.

See Diazomethane

See other NITROSO COMPOUNDS

1598. *trans*-4-Hydroperoxy-5-hydroxy-4-methylimidazolin-2-one
[85576-52-9]



Preparative hazard

See Hydrogen peroxide: 2-Amino-4-methyloxazole

See related ALKYL HYDROPEROXIDES

1599. 2,2'-Oxybis(ethyl nitrate)
[693-21-0]



1. Kit and Evered, 1960, 268
2. Toso, R. *et al.*, *Chem. Abs.*, 1988, **108**, 24140

A powerful explosive, sensitive to vibration and mechanical shock, too heat-sensitive for a rocket propellant [1]. In the industrial preparation of this explosive by the continuous nitration of diethylene glycol by nitric-sulfuric acid mixtures, the main factor limiting the overall safety of the process is the instability of the spent acid mixture, and handling of the latter is the most important specific problem. Close control of the acid composition and of the glycol:acid ratio reduces the nitric acid and water contents of the organic phase in the reaction mixture, and so increases the stability of the organic phase. Continuous addition of oleum to the spent acid coming from the separator increases its stability and permits safe handling and storage [2].

See other NITRATION INCIDENTS, ALKYL NITRATES

1600. *N*-Allylthiourea
[101-57-9]



Energy of decomposition (in range 170–270°C) measured as 0.549 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ALLYL COMPOUNDS

1601. Azo-*N*-methylformamide


[18880-14-3]

C₄H₈N₄O₂Smith, R. F. *et al.*, *J. Org. Chem.*, 1975, **40**, 1855

Thermolysis is vigorously exothermic, the temperature increasing from 176 to 259°C during decomposition.

See other AZO COMPOUNDS, BLOWING AGENTS**1602. Bis(2-nitratoethyl)nitric amide**

[4185-47-1]

C₄H₈N₄O₈

Chromium compounds

Aleksandrov, V. V. *et al.*, *Combust. Flame*, 1979, **35**, 1–15

The burning rate of the solid explosive is increased by various chromium compounds.

See related ALKYL NITRATES, *N*-NITRO COMPOUNDS**1603. 2,2-Diazidobutane**

[90329-44-5]

C₄H₈N₆Kirchmeyer, S. *et al.*, *Synthesis*, 1983, 501Potentially explosive (a *gem*-diazide).*See other* ORGANIC AZIDES**1604. 1,1'-Oxybis-2-azidoethane**


[24345-74-2]

C₄H₈N₆O1. Rankine A., *Safety Digest Univ. Safety Assn.*, 1992, **43**, 222. Peacock, R. D., *Chemistry in Britain*, 1992, **28** (4), 327

An explosion was experienced during isolation of supposed diamine produced by reduction of the diazide with hydrogen sulfide. This was attributed to incomplete reduction [1]. An explosion, apparently during reduction, of a closely related compound by this method has earlier been reported [2].

See 1,2-Bis(2-azidoethoxy)ethane*See other* ORGANIC AZIDES**1605. 1,3,5,7-Tetranitroperhydro-1,3,5,7-tetrazocine**

[2691-41-0]

C₄H₈N₈O₈

The military explosive HMX. Violent decomposition occurred at 279°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1)

See 1,3,5-Trinitrohexahydro-1,3,5-triazine

See other N-NITRO COMPOUNDS

†1606. 2-Butanone (Ethyl methyl ketone)

[78-93-3]

C₄H₈O

MeCO.Et

(MCA SD-83, 1961); FPA H16, 1973; HCS 1980, 650; RSC Lab. Hazard Data Sheet No. 18, 1983

Chloroform, Alkali

See Chloroform: Acetone, etc.

Hydrogen peroxide

MRH ,

MRH 6.36/83

Nitric acid

MRH 5.77/79

See Hydrogen peroxide: Ketones, etc.

Other reactants

Yoshida, 1980, 365

MRH values calculated for 13 combinations with oxidants are given.

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

2-Propanol

See 2-Propanol: 2-Butanone

†1607. Butyraldehyde (Butanal)

[123-72-8]

C₄H₈O

PrCO.H


(MCA SD-78, 1960); FPA H70, 1978; HCS 1980, 254

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

†1608. Cyclopropyl methyl ether (Methoxycyclopropane)

[540-47-6]

C₄H₈O



See other PEROXIDISABLE COMPOUNDS, STRAINED-RING COMPOUNDS

†1609. 1,2-Epoxybutane (Ethyloxirane)
[106-88-7]

C₄H₈O



HCS 1980, 455

Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Oxygen heterocycles

See other 1,2-EPOXIDES, STRAINED-RING COMPOUNDS

†1610. Ethyl vinyl ether (Ethoxyethene)
[109-92-2]

C₄H₈O



Methanesulfonic acid

Eaton, P. E. *et al. J. Org. Chem.*, 1972, **37**, 1947

Methanesulfonic acid is too powerful a catalyst for *O*-alkylation with the vinyl ether, causing explosive polymerisation of the latter on multimolar scale. Dichloroacetic acid is a satisfactory catalyst on the 3 g mol scale.

Other reactants

Yoshida, 1980, 288

MRH values calculated for 13 combinations with oxidants are given.

See other PEROXIDISABLE COMPOUNDS

†1611. Isobutyraldehyde (2-Methylpropanal)
[78-84-2]

C₄H₈O




(*MCA SD-78*, 1960)

See other PEROXIDISABLE COMPOUNDS

†1612. Tetrahydrofuran
[109-99-9]

C₄H₈O



FPA H60, 1977; *HCS 1980*, 891; *RSC Lab. Hazard Data Sheet No. 12*, 1983

C₄H₈O

1. *Recommended Handling Procedures*, THF Brochure FC3-664, Wilmington, Du Pont, 1964
2. Anon., *Org. Synth.*, 1966, **46**, 105
3. Schurz, J. *et al.*, *Angew. Chem.*, 1956, **68**, 182
4. Davies, A. G., *J. R. Inst. Chem.*, 1956, **80**, 386
5. Halonbrenner, R. *et al.*, *Chem. Eng. News*, 1978, **56**(6), 3
6. Coates, J. S., *Chem. Eng. News*, 1978, **56**(6), 3
7. Wissink, H. G., *Chem. Eng. News*, 1997, **75**(9), 6
8. Schwartz, A. M., *Chem. Eng. News*, 1978, **56**(24), 88
9. Hanson, E. S., *Chem. Eng. News*, 1978, **56**(24), 88
10. Nagano, H., *Chem. Abs.*, 1980, **92**, 163833

Like many other ethers and cyclic ethers, in absence of inhibitors tetrahydrofuran is subject to autoxidation on exposure to air, when initially the 2-hydroperoxide forms. This tends to decompose smoothly when heated, but if allowed to accumulate for a considerable period, it becomes transformed into other peroxidic species which will decompose violently [1]. Commercial material is supplied stabilised with a phenolic antioxidant which is effective under normal closed storage conditions in preventing the formation and accumulation of peroxide. Procedures for testing for the presence of peroxides and also for their removal are detailed [1,2,4]. The last of these references recommends the use of copper(I) chloride for removal of trace amounts of peroxide. If more than trace amounts are present, the peroxidised solvent (small quantities only) should be discarded by dilution and flushing away with water [2]. An attempt to remove peroxides by shaking with solid ferrous sulfate before distillation did not prevent explosion of the distillation residue [3]. Alkali treatment to destroy peroxides [1] appears not to be safe [2]. (See Caustic alkalies, below). Distillation or alkali treatment of stabilised THF removes the involatile antioxidant and the solvent must be restabilised or stored under nitrogen to prevent peroxide formation during storage, which should not exceed a few days' duration in the absence of stabiliser [2]. The use of lithium tetrahydroaluminate is only recommended for drying THF which is peroxide-free and is not grossly wet [2]. A violent explosion during reflux of the solvent with calcium hydride [5] was attributed to cleavage of the cyclic ether by overheated excess hydride [6]. Other metal hydrides (and some other strong bases) can also cause decomposition to ethene and enolate salts of acetaldehyde [7], pressurising containers by gas emission. The use of sodium benzophenone ketyl [8], or of an activated alumina column [9] to remove moisture and peroxides are detailed. Peroxides in THF may be destroyed by passage through activated carbon at 20–66°C with contact time >2 min [10].

See 2-Tetrahydrofuryl hydroperoxide

See also ETHERS (references 9,10)

2-Aminophenol, Potassium dioxide

See Potassium dioxide: 2-Aminophenol, Tetrahydrofuran

Borane

See Borane–tetrahydrofuran

Bromine

See Bromine: Tetrahydrofuran

Calcium hydride

See (references [5,6])

Caustic alkalis

1. *NSC Newsletter, Chem. Sect.*, 1964(10); 1967(3)

2. Anon., *Org. Synth.*, 1966, **46**, 105

It is not safe to store quantities of THF which have been freed of the phenolic inhibitor (e.g. by alkali treatment) since dangerous quantities of peroxides may build up in prolonged storage. Peroxidised materials should not be dried with sodium hydroxide or potassium hydroxide, as explosions may occur [1,2].

Diisobutylaluminium hydride

See Diisobutylaluminium hydride: Tetrahydrofuran

Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Tetrahydrofuran

Metal halides

See Hafnium tetrachloride: Tetrahydrofuran

Titanium tetrachloride: Tetrahydrofuran

Zirconium tetrachloride: Tetrahydrofuran

Other reactants

Yoshida, 1980, 238

MRH values calculated for 12 combinations with oxidants are given.

Oxygen, strong base

Butora, G. *et al.*, *J. Amer. Chem. Soc.*, 1997, **119**(33), 7694

As part of a synthesis, a rather dilute THF solution of an heterocycle, converted into a dianion with *sec*-butyllithium, was oxidised by bubbling gaseous oxygen through at -78 C . In the light of above entries, formation of peroxides is not surprising; however, the reaction tended to inflame or explode while still warming to ambient temperature for treatment with deperoxidising reagents. The technique is not recommended and an electrochemical alternative, far superior both in yield and safety, was devised.

Sodium tetrahydroaluminate

See Sodium tetrahydroaluminate: Tetrahydrofuran


Sulfinyl chloride

See Sulfinyl chloride: Tetrahydrofuran

See *other* APROTIC SOLVENTS, PEROXIDISABLE COMPOUNDS

1613. 1,4-Oxathiane
[15980-15-1]

C₄H₈OS



Metal perchlorates

See Silver perchlorate: 1,4-Oxathiane

Copper(I) perchlorate: 1,4-Oxathiane

1614. Butyric acid (Butanoic acid)
[107-92-6]

C₄H₈O₂

PrCO.OH

HCS 1980, 255


Chromium trioxide

See Chromium trioxide: Butyric acid

See other ORGANIC ACIDS

1615. 3,3-Dimethyl-1,2-dioxetane
[32315-88-1]

C₄H₈O₂




Adam, W. *et al.*, *J. Amer. Chem. Soc.*, 1994, **116**(17), 7581

This molecule sometimes detonated spontaneously even below 0°C. The trimethyl and tetramethyl homologues should not be handled above 0°C and the precursor bromohydroperoxides are also hazardous.

See other DIOXETANES, STRAINED-RING COMPOUNDS; CYCLIC PEROXIDES


†**1616. 1,3-Dioxane**
[505-22-6]

C₄H₈O₂



See other PEROXIDISABLE COMPOUNDS

†1617. 1,4-Dioxane
[123-91-1]



HCS 1980, 442; RSC Lab. Hazard Data Sheet No. 29, 1984

1. Dasler, W. *et al.*, *Ind. Eng. Chem. (Anal. Ed.)*, 1946, **18**, 52

Like other monofunctional ethers but more so because of the four susceptible hydrogen atoms, dioxane exposed to air is susceptible to autoxidation with formation of peroxides which may be hazardous if distillation (causing concentration) is attempted. Because it is water-miscible, treatment by shaking with aqueous reductants (iron(II) sulfate, sodium sulfide, etc.) is impracticable. Peroxides may be removed, however, under anhydrous conditions by passing dioxane (or any other ether) down a column of activated alumina. The peroxides (and any water) are removed by adsorption onto the alumina, which must then be washed with methanol or water to remove them before the column material is discarded [1]. The heat of decomposition of dioxane has been determined (130–200°C) as 0.165 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Decaborane(14)

See Decaborane(14): Ether, etc.

Nickel

Mozingo, R., *Org. Synth.*, 1955, Coll. Vol. 3, 182

Dioxane reacts with Raney nickel catalyst almost explosively above 210°C.

Nitric acid, Perchloric acid

MRH Nitric acid 5.81/74

See Perchloric acid: Dioxane, Nitric acid

Other reactants

Yoshida, 1980, 150

MRH values calculated for 14 combinations with oxidants are given.

Sulfur trioxide

MRH 2.26/38

See Sulfur trioxide: Diethyl ether, etc.

Triethynylaluminium

See Triethynylaluminium: Dioxane

See other APROTIC SOLVENTS, PEROXIDISABLE COMPOUNDS

†1618. Ethyl acetate (Ethyl ethanoate)
[141-78-6]



(MCA SD-51, 1953); FPA H8, 1973; HCS 1980, 462; RSC Lab. Hazards Data Sheet No. 41, 1985

1. Anon., *Loss. Prev. Bull.*, 1977, (013), 16

2. Bond, J., *Loss Prev. Bull.*, 1994, (119), 9

During rotary evaporation of solvent (from an ethyl acetate extract of a fermentation culture) at 55°C/50 mbar, the flask exploded. (It seems likely that some unsuspected peroxidised material may have been present in the extract, or perhaps the flask was scratched or cracked, and imploded [1].) Several fires and explosions suggest that ethyl acetate may be a worse generator of static electricity than its measured conductivity would indicate [2].

See Hydrogen peroxide: Ethyl acetate

See other STATIC INITIATION INCIDENTS

Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Ethyl acetate

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

†1619. Isopropyl formate
[625-55-8]



$\text{C}_4\text{H}_8\text{O}_2$

†1620. Methyl propionate
[554-12-1]



$\text{C}_4\text{H}_8\text{O}_2$

HCS 1980, 659

†1621. Propyl formate
[110-74-7]



$\text{C}_4\text{H}_8\text{O}_2$

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

1622. Tetrahydrothiophene-1,1-dioxide (Sulfolane)
[126-33-0]

$\text{C}_4\text{H}_8\text{O}_2\text{S}$




Nitronium tetrafluoroborate

See NITRATING AGENTS *See other* APROTIC SOLVENTS

1623. *trans*-2-Butene ozonide (3,5-Dimethyl-1,2,4-trioxolane)
[16187-15-8]


C₄H₈O₃



See *trans*-2-Hexene ozonide
See other OZONIDES

1624. 2-Tetrahydrofuryl hydroperoxide
[4676-82-8]

C₄H₈O₃



1. Rein, H., *Angew. Chem.*, 1950, **62**, 120
2. Criegee, R., *Angew. Chem.*, 1950, **62**, 120


This occurs as the first product of the ready autoxidation of tetrahydrofuran, and is relatively stable [1]. It readily changes, however, to a highly explosive polyalkylidene peroxide, which is responsible for the numerous explosions observed on distillation of peroxidised tetrahydrofuran [2].

See Tetrahydrofuran

See other ALKYL HYDROPEROXIDES

1625. 3,6-Dimethyl-1,2,4,5-tetraoxane
[cis [102502-32-9]; trans [118171-56-5]]

C₄H₈O₄



Rieche, A. *et al.*, *Ber.*, 1939, **72**, 1933

This dimeric 'ethylidene peroxide' is an extremely shock-sensitive solid which explodes violently at the slightest touch. Extreme caution in handling is required.

See other CYCLIC PEROXIDES

1626. 1-Hydroxyethyl peroxyacetate
[7416-48-0]

C₄H₈O₄




An explosive low-melting solid, readily formed during autoxidation of acetaldehyde.

See Acetaldehyde: Cobalt acetate, etc.
See other PEROXYESTERS

†1627. Tetrahydrothiophene
[110-01-0]

C₄H₈S



Hydrogen peroxide
See Hydrogen peroxide: Tetrahydrothiophene

1628. Acetyldimethylarsine
[21380-82-5]

C₄H₉AsO

AcAsMe₂

DOC 5, 1982, 38

It ignites in air.
See related ALKYLNON-METALS

1629. Butyldichloroborane
[14090-22-3]

C₄H₉BCl₂

BuBCl₂

Air, or Water
Niedenzu, K. *et al.*, *Inorg. Synth.*, 1967, **10**, 126
It ignites on prolonged exposure to air; hydrolysis may be explosive.
See other ALKYLHALOBORANES

1630. But-2-en-2-ylboronic acid
[[125261-72-5] E -]
[[125261-73-6] Z -]

C₄H₉BO₂

CH₃CH=C(CH₃).B(OH)₂

Trombini, C. *et al.* *J. Chem. Soc. Perk. 1*, 1989, 1025
The compound is pyrophoric when dry.
See other PYROPHORIC COMPOUNDS

†1631. 1-Bromobutane (*n*-Butyl bromide)
[109-65-9]

C₄H₉Br

BrCH₂Pr

Bromobenzene, Sodium
See Sodium: Halocarbons (reference 7)
See other HALOALKANES

- †1632. 2-Bromobutane (*sec*-Butyl bromide)
[78-76-2] C₄H₉Br
MeCHBrEt
See other HALOALKANES
- †1633. 1-Bromo-2-methylpropane (Isobutyl bromide)
[78-77-3] C₄H₉Br
BrCH₂CHMe₂
See other HALOALKANES
- †1634. 2-Bromo-2-methylpropane (*tert*-Butyl bromide)
[507-19-7] C₄H₉Br
BrCMe₃
See other HALOALKANES
- †1635. 2-Bromoethyl ethyl ether
[592-55-2] C₄H₉BrO
BrC₂H₄OEt
1636. 1,1-Dimethylethyldibromamine (*tert*-Butyldibromamine)
[51655-36-8] C₄H₉Br₂N
Me₃CNBr₂
Kille, G. *et al.*, *Ind. Chem. Libr.*, 1991, 3(1), 75; *Chem. Abs.*, 1992, 116620(11), 105572
Differential scanning calorimetry (DSC) showed this to be prone to highly exothermic decomposition (100 J/g) at ambient temperatures. Solutions are a little more stable.
See other N-HALOGEN COMPOUNDS
See also DIFFERENTIAL SCANNING CALORIMETRY
- †1637. 1-Chlorobutane (*n*-Butyl chloride)
[109-69-3] C₄H₉Cl
ClCH₂Pr
HCS 1980, 297
See other HALOALKANES
- †1638. 2-Chlorobutane (*sec*-Butyl chloride)
[78-86-4] C₄H₉Cl
MeCHClEt
See other HALOALKANES

†1639. 1-Chloro-2-methylpropane (Isobutyl chloride)
[513-36-0] C₄H₉Cl
ClCH₂CHMe₂

See other HALOALKANES

†1640. 2-Chloro-2-methylpropane (*tert*-Butyl chloride)
[507-20-0] C₄H₉Cl
ClCMe₃

HCS 1980, 304

See other HALOALKANES

1641. Butylmagnesium chloride
[693-04-9] C₄H₉ClMg
BuMgCl

Water

Nolan, 1983, Case history 109

Butylmagnesium chloride was prepared by adding a mixture of ether, butyl chloride and butyl bromide to magnesium, then cyclohexane was added prior to adding stannic chloride to form tetrabutyltin. However, the cyclohexane contained water and this reacted with the Grignard reagent to liberate butane which ruptured the bursting disk and ignited.

See other GAS EVOLUTION INCIDENTS, GRIGNARD REAGENTS

1642. *tert*-Butyl hypochlorite
[507-40-4] C₄H₉ClO
Me₃COCl

1. Lewis, J. C., *Chem. Eng. News*, 1962, **40**(43), 62
2. Teeter, H. M. *et al.*, *Org. Synth.*, 1962, Coll. Vol. 4, 125
3. Mintz, M. J. *et al.*, *Org. Synth.*, 1973, Coll. Vol. 5, 185

This material, normally supplied or stored in sealed ampoules and used as a paper chromatography spray reagent, is photo-sensitive. Exposure to UV light causes exothermic decomposition to acetone and chloromethane. Ampoules have burst because of pressure build-up after exposure to fluorescent or direct day light. Store cool and dark, and open ampoules with personal protection. The material also reacts violently with rubber [1,2]. It should not be heated to above its boiling point [3]. There is also a preparative hazard.

See Chlorine: tert-Butanol

Sodium hydrogen cyanamide

See Cyanonitrene

See other HYPOHALITES, IRRADIATION DECOMPOSITION INCIDENTS

- 1643. 2(2-Hydroxyethoxy)ethyl perchlorate**
 [] **C₄H₉ClO₆**

$$\text{HOC}_2\text{H}_4\text{OC}_2\text{H}_4\text{OCIO}_3$$
- Hofmann, K. A. *et al.*, *Ber.*, 1909, **42**, 4390
 It explodes violently on heating in a capillary.
See other ALKYL PERCHLORATES
- 1644. tert-Butyl peroxophosphoryl dichloride**
 [] **C₄H₉Cl₂O₃P**

$$\text{Me}_3\text{COOP(O)Cl}_2$$
- The product decomposed violently after isolation.
See entry tert-BUTYL PEROXOPHOSPHATE DERIVATIVES
- 1645. N-Fluoro-N-nitrobutylamine**
 [14233-86-4] **C₄H₉FN₂O₂**

$$\text{BuN(F)NO}_2$$
- Grakauskas, V. *et al.*, *J. Org. Chem.*, 1972, **37**, 334
 A sample exploded on vacuum distillation at 60°C, though not at 40°C/33 mbar.
See other N-HALOGEN COMPOUNDS, N-NITRO COMPOUNDS
- 1646. tert-Butyldifluorophosphine**
 [29149-32-4] **C₄H₉F₂P**

$$\text{Me}_3\text{CPF}_2$$
- Stelzer, O. *et al.*, *Inorg. Synth.*, 1978, **18**, 174
 It ignites in air.
See other ALKYLHALOPHOSPHINES
- †**1647. 2-Iodobutane (sec-Butyl iodide)**
 [513-48-4] **C₄H₉I**

$$\text{MeCHI Et}$$
- See other* HALOALKANES
- †**1648. 1-Iodo-2-methylpropane (Isobutyl iodide)**
 [513-38-2] **C₄H₉I**

$$\text{ICH}_2\text{CHMe}_2$$
- See other* HALOALKANES

†1649. 2-Iodo-2-methylpropane (*tert*-Butyl iodide)
[558-17-8]

Me₃CI

C₄H₉I

See other HALOALKANES

1650. Potassium *tert*-butoxide
[865-47-4]

KOCMe₃

C₄H₉KO

Fieser, 1967, Vol. 1, 911

This extremely powerful base may ignite if exposed to air (or oxygen) at elevated temperatures.

See other PYROPHORIC MATERIALS

Acids, or Reactive solvents

Manwaring, R. *et al.*, *Chem. & Ind.*, 1973, 172

Contact of 1.5 g portions of the solid butoxide with drops of the liquid (l) or with the vapours (v) of the reagents below caused ignition after the indicated period (min).

Acetic acid, v, 3; sulfuric acid, l, 0.5

Methanol, l, 2; ethanol, v, 7; propanol, l, 1; isopropanol, l, 1

Ethyl acetate, v, 2; butyl acetate, v, 2; propyl formate, v, 4; dimethyl carbonate, l, 1; diethyl sulfate, l, 1

Acetone, v, 4, l, 2; 2-butanone, v, 1, l, 0.5; 4-methyl-2-butanone, v, 3

Dichloromethane, l, 2; chloroform, v, 2, l, 0; carbon tetrachloride, l, 1; 1-chloro-2,3-epoxypropane, l, 1

The potentially dangerous reactivity with water, acids or halocarbons was already known, but that arising from contact with alcohols, esters or ketones was unexpected. Under normal reaction conditions, little significant danger should exist where excess of solvent will dissipate the heat, but accidental spillage of the solid butoxide could be hazardous.

Dimethyl sulfoxide

See Dimethyl sulfoxide: Metal alkoxides

See other METAL ALKOXIDES

1651. Butyllithium
[109-72-8]

BuLi

C₄H₉Li

Air, or Carbon dioxide, or Water

1. (*MCA SD-91*, 1966)

2. Dezmelyk, E. W. *et al.*, *Ind. Eng. Chem.*, 1961, **53**(6), 56A

This reactive liquid is now normally supplied commercially as a solution (up to 25 wt.%) in pentane, hexane or heptane, because it reacts with ether and ethereal solutions must be stored under refrigeration. Reaction with atmospheric oxygen or

water vapour is highly exothermic and solutions of 20% concentration will ignite on exposure to air at ambient temperature and about 70% relative humidity. Solutions of above 25% concentration will ignite at any humidity. Contact with liquid water will cause immediate ignition of solutions in these highly flammable solvents [1]. Though contact with carbon dioxide causes an exothermic reaction, it is less than with atmospheric oxygen, so carbon dioxide extinguishers are effective on butyllithium fires. Full handling, operating and disposal procedures are detailed [2].

Styrene

See Styrene: Butyllithium

Water

See Air, etc., above

See other ALKYL METALS

1652. *tert*-Butyllithium

[594-19-4]

C₄H₉Li

Me₃CLi

Turbitt, T. D. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1974, 183

The 2 M solution in heptane ignites in air.

Copper(I) iodide, Tetrahydrofuran

Kurzen, H. *et al.*, *J. Org. Chem.*, 1985, **50**, 3222

t-Butyllithium in presence of copper(I) iodide in THF is used for alkylation. If an excess of the lithium derivative is present, introduction of oxygen or air can cause a violent explosion, even at -78°C.

2,2,2,4,4-Hexafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine

Harris, R. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1976, 23

Interaction in hexane to produce the 2,4-di-*tert*-butyl derivative often starts only after an induction period and may then proceed very violently. Careful temperature control is imperative.


See other INDUCTION PERIOD INCIDENTS See other ALKYL METALS

†1653. Pyrrolidine

[123-75-1]

C₄H₉N

Benzaldehyde, Propionic acid



See Benzaldehyde: Propionic acid, etc.

1654. 2-Butanone oxime (2-Oximinobutane)

[96-29-7]

C₄H₉NO

MeC(:NOH)Et

HCS 1980, 652

Tyler, L. J., *Chem. Eng. News*, 1974, **52**(35), 3

Investigation following two violent explosions involving the oxime or its derivatives showed that it may be distilled at 152°C at ambient pressure only if highly purified. Presence of impurities, especially acidic impurities (e.g. the oxime hydrochloride) drastically lowers the temperature at which degradation occurs.

See 2-Butanone oxime hydrochloride

See other CATALYTIC IMPURITY INCIDENTS

Sulfuric acid

MCA Guide, 1972, 300

The oxime was hydrolysed with aqueous sulfuric acid and the 2-butanone liberated was distilled out at 15 mbar from a bath at 110–115°C. Soon after release of vacuum and bath removal, the residue (crude hydroxylaminium sulfate) decomposed violently.

See HYDROXYLAMINIUM SALTS See other OXIMES

1655. Butyraldehyde oxime (1-Hydroxyiminobutane)

[151-00-8]

C₄H₉NO

PrCH=NOH

1. Anon., *Sichere Chemiearb.*, 1966, **18**(3), 20

2. Kröger, *Proc. 1st Int. Symp. Prev. Occ. Risks Chem. Ind.*, 180–187, Heidelberg, ISSA, 1970

A large batch exploded violently (without flame) during vacuum distillation at 90–100°C/20–25 mbar. Since the distilled product contained up to 12% butyronitrile, it was assumed that the oxime had undergone the Beckman rearrangement to butyramide and then dehydrated to the nitrile. The release of water into a system at 120°C would generate excessive steam pressure which the process vessel could not withstand. The rearrangement may have been catalysed by metallic impurities [1]. This hypothesis was confirmed in a detailed study, which identified lead oxide and rust as active catalysts for the rearrangement and dehydration reactions [2].

See Ethyl 2-formylpropionate oxime: Hydrogen chloride

Sulfuric acid: Cyclopentanone oxime

See other CATALYTIC IMPURITY INCIDENTS, OXIMES

1656. N,N-Dimethylacetamide

[127-19-5]

C₄H₉NO

Me₂NAc

HCS 1980, 420

Halogenated compounds

1. 'DMAC Brochure A-79931', Wilmington, Du Pont, 1969
2. Cardillo, P. *et al.*, *3rd Euro. Symp. Therm. Anal. Calorim.*, Interlaken, 1984 (private comm.,)

The tertiary amide acts as a dehydrohalogenating agent, and reaction with some highly halogenated compounds (carbon tetrachloride, hexachlorocyclohexane) is very exothermic and may become violent, particularly if iron is present [1]. A detailed investigation by ARC shows that 1:1 w/w mixtures with carbon tetrachloride are most reactive and that 2 exotherms are evident. One starting at 91° and ending at 97°C is relatively weak (40 J/g) while the second, starting at 97° and ending at 172° has its maximum rate at 147°C, with an exotherm of 508 J/g, which under adiabatic runaway conditions would give a final temperature exceeding 450°C. A considerable release of gas generates a maximum pressure of 12.8 bar at 172 °C. Presence of 1% of iron powder initiates the first exotherm at 71°C and increases it to 60 J/g, the second exotherm being marginally reduced by presence of iron. The exothermic effect is much greater than in the similar reactions of DMF [2].

See Carbon tetrachloride: Dimethylformamide

See 3,4-Dichloronitrobenzene

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

Other reactants

Yoshida, 1980, 165


MRH values calculated for 13 combinations with oxidants are given.

See other APROTIC SOLVENTS

1657. Morpholine (Tetrahydro-1,4-oxazine)

[110-91-8]

C₄H₉NO



HCS 1980, 671; *RSC Lqab. Hazards Data Sheet No. 73*, 1988

Cellulose nitrate

See CELLULOSE NITRATE: amines

Nitromethane

See Nitromethane: Acids, etc.

See other ORGANIC BASES

†1658. Butyl nitrite

[544-16-1]

BuON:O

C₄H₉NO₂

See other ALKYL NITRITES

- †1659. *tert*-Butyl nitrite
[540-80-7] C₄H₉NO₂
Me₃CON:O
- Lopez, F., *Chem. Eng. News*, 1992, **70**(51), 2
DSC showed that exothermic decomposition (−1200 J/g) begins at about 110°C. A falling hammer test indicated shock induced decomposition. Considerable caution is advised when employing this useful reagent.
See DIFFERENTIAL SCANNING CALORIMETRY
See other ALKYL NITRITES
1660. *tert*-Nitrobutane (2-Methyl-2-nitropropane)
[594-70-7] C₄H₉NO₂
O₂NCMe₃
- Preparative hazard
Mee, A. J., *School Sci. Rev.*, 1940, **22**(85), 96
A sample exploded during distillation.
See Potassium permanganate: Acetone, *tert*-Butylamine
See other NITROALKANES
1661. Butyl nitrate
[928-45-0] C₄H₉NO₃
BuONO₂
- Lewis acids
See ALKYL NITRATES: lewis acids
- Other reactants
Yoshida, 1980, 203
MRH values calculated for 10 combinations with various reagents are given.
See other ALKYL NITRATES
1662. Ethyl 2-nitroethyl ether (2-Nitroethoxyethane)
[31890-52-5] C₄H₉NO₃
EtOC₂H₄NO₂
- Cann, P. F. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1974, 817–819
Addition of diphenyl ether as an inert diluent to the crude ether before distillation at 0.13 mbar is essential to prevent violent explosion of the residue after it has cooled.
See related NITROALKANES
1663. *tert*-Butyl peroxyhydrate (1,1-Dimethylethyl peroxyhydrate)
[] C₄H₉N₄
Me₃COONO₂
- Like other low molecular weight peroxyesters, or alkyl nitrates, this is a material to avoid isolating even when you can make it.

Preparative hazard


See *tert*-Butylhydroperoxide: Toluene, Dinitrogen pentaoxide

See other PEROXYESTERS

1664. Tris(hydroxymethyl)nitromethane

(2-Hydroxymethyl-2-nitropropane-1,3-diol)

[126-11-4]



C₄H₉NO₅

Hydrogen, Nickel

Ostis, K. *et al.*, *Chem. Abs.*, 1975, **82**, 36578

Catalytic hydrogenation of the title compound to the amine with hydrogen at 5–10 bar is described as hazardous, and an electrochemical reduction process is recommended.

See other CATALYTIC NITRO REDUCTION PROCESSES, HYDROGENATION INCIDENTS

See related NITROALKANES

†1665. Cyanotrimethylsilane

[7677-24-9]



C₄H₉NSi

See related ALKYLHALOSILANES See other CYANO COMPOUNDS

1666. 1-Azidobutane (Butyl azide)

[7332-00-5]



C₄H₉N₃

See Butyl toluenesulfonate

See AZIDES

1667. Butylsodium

[3525-44-8]



C₄H₉Na

Preparative hazard

See Sodium: 1-Chlorobutane See other ALKYLMETALS

†1668. Butane

[106-97-8]



C₄H₁₀

FPA H15, 1973 (cylinder); *HCS 1980*, 224 (cylinder)

See LIQUEFIED PETROLEUM GASES

†1669. Isobutane (2-Methylpropane)
[75-28-5]



FPA H119, 1983 (cylinder); *HCS 1980*, 565

1670. Diethylaluminium bromide
[760-19-0]



Nitromethane

See Nitromethane: Alkylmetal halides

See entry ALKYLALUMINIUM HALIDES

1671. Diethylaluminium chloride
[96-10-6]



HCS 1980, 390

Chlorine azide

Houben-Weyl 1970, Vol. 13.4, 76

Interaction produces distillable ethylaluminium azide chloride, but the residue is explosive.

See other ALKYLALUMINIUM HALIDES

1672. Diethylgold bromide
[26645-10-3]



Sorbe, 1968, 63

It explodes at 70°C.

See other ALKYLMETAL HALIDES, GOLD COMPOUNDS

1673. Chlorodiethylborane
[5314-83-0]




Koster, R. *et al.*, *Inorg. Synth.*, 1974, **15**, 149–152

It ignites in air; preparative hazard also described.

See other ALKYLHALOBORANES

1674. Boron trifluoride diethyl etherate

[109-63-7]

C₄H₁₀BF₃O*HCS 1980*, 217

Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Boron trifluoride diethyl etherate*See related* NON-METAL HALIDES**1675. Diethylberyllium**

[542-63-2]

C₄H₁₀Be

Air, or Water

Coates, 1967, Vol. 1, 106

Ignites in air, even when containing ether, and reacts explosively with water.

See other ALKYL METALS**1676. Diethylbismuth chloride**

[65313-34-0]

C₄H₁₀BiClGilman, H. *et al.*, *J. Org. Chem.*, 1939, **4**, 167

It ignites in air.

See other ALKYL METAL HALIDES**1677. Diethylcadmium**

[592-02-9]

C₄H₁₀Cd1. Krause, E., *Ber.*, 1918, **50**, 1813

2. Houben-Weyl, 1973, Vol. 13.2a, 903 (footnote 3), 908

The vapour decomposes explosively at 180°C. Exposure to ambient air produces white fumes which turn brown and then explode violently [1]. Moderate quantities decompose explosively when heated rapidly to 130, not 180°C as stated previously.

This, apart from allylcadmium derivatives, is the only pyrophoric dialkylcadmium [2] (but *see* Dimethylcadmium). It is mildly endothermic (ΔH_f° (l) +61.5 kJ/mol, 0.36 kJ/g).*See other* ALKYL METALS, ENDOTHERMIC COMPOUNDS**1678. 2-Butanone oxime hydrochloride (2-Hydroxyiminiobutane chloride)**

[4154-69-2]

C₄H₁₀CINOTyler, L. J., *Chem. Eng. News*, 1974, **52**(35), 3

The salt undergoes violent degradation at 50–70°C, and its presence in trace quantities may promote degradation of the oxime.

See related OXIMES

1679. Diethylaminosulfinyl chloride

[34876-30-7]



$\text{C}_4\text{H}_{10}\text{ClNOS}$


Preparative hazard

See Sulfinyl chloride: Bis(dimethylamino) sulfoxide

See other N-S COMPOUNDS

1680. Morpholinium perchlorate

[35175-75-8]



$\text{C}_4\text{H}_{10}\text{ClNO}_5$


Udupa, M. R., *Thermochim. Acta*, 1980, **38**, 241–143

It decomposes exothermally at 230°C.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

1681. Diethyl phosphorochloridate (Diethyl chlorophosphate)

[814-49-3]



$\text{C}_4\text{H}_{10}\text{ClO}_3\text{P}$

Silbert, L. A., *J. Org. Chem.*, 1971, **36**, 2162

Presence of hydrogen chloride as impurity causes an uncontrollable exothermic reaction during preparation of diethyl phosphate from the title compound.

See related PHOSPHORUS ESTERS See other CATALYTIC IMPURITY INCIDENTS

1682. Diethylthallium perchlorate

[[22392-07-0] (ion)]



$\text{C}_4\text{H}_{10}\text{ClO}_4\text{Tl}$

Cook, J. R. *et al.*, *J. Inorg. Nucl. Chem.*, 1964, **26**, 1249

It explodes at the m.p., 250°C.

See related METAL PERCHLORATES

†**1683. Dichlorodiethylsilane**

[1719-53-5]



$\text{C}_4\text{H}_{10}\text{Cl}_2\text{Si}$

See other ALKYLHALOSILANES

1684. Diethylaminosulfur trifluoride
[38078-09-0]

$C_4H_{10}F_3NS$

Et_2NSF_3

Alone, or Water

1. Cochran, J., *Chem. Eng. News*, 1979, **57**(12), 4, 74
2. Middleton, W. J., *Chem. Eng. News*, 1979, **57**(21), 43
3. Middleton, W. J. *et al.*, *J. Org. Chem.*, 1980, **45**, 2884 (footnote 3)
4. Middleton, W. J. *et al.*, *Org. Synth.*, 1977, **57**, 50
5. Breckenridge, R. J. *et al.*, *Chem. Brit.*, 1982, **18**, 619
6. Suckling, C. J., private comm., 1982
7. Pivawer, P. M., private comm., 1983

Explosive decomposition occurred when stripping solvent or attempting vacuum distillation. Exothermic decomposition begins at about 50°C and is violent above 90°C. Contact with water causes explosive decomposition [1]. Later comments suggest that the compound may be distilled safely at pressures below 13 mbar and handled safely in solution, or in bulk below 50°C [2,3]. While repeating an earlier procedure [4], two explosions occurred. One was when admitting air to the residue from vacuum distillation, and the second immediately prior to redistillation of the cold product from vacuum stripping of solvent [5]. This was attributed to formation of some impurities related to a dialkylaminofluorosulfur imine [6]. During preliminary distillation to remove sulfur tetrafluoride, etc., the line from the cold trap to the alkaline scrubber must end in an inverted funnel to avoid blockage with solid [7].

See *S*-Diethylamino(methylimino)sulfur(IV) fluoride

See other N-S COMPOUNDS See related NON-METAL HALIDES

1685. Potassium diethylamide
[2245-68-3]

$C_4H_{10}KN$

$KNEt_2$

Pez, G. P. *et al.*, *Pure Appl. Chem.*, 1986, **57**, 1925

The solid is highly pyrophoric and will react violently even with limited exposure to air

See other N-METAL DERIVATIVES, PYROPHORIC MATERIALS

1686. Lithium diethylamide
[816-43-3]

$C_4H_{10}LiN$

$LiNEt_2$

1. Fischer, E. O. *et al.*, *Inorg. Synth.*, 1979, **19**, 165
2. Pez, G. P. *et al.*, *Pure Appl. Chem.*, 1986, **57**, 1924

Extremely pyrophoric in air [1], and will react violently upon even brief contact [2].

See other N-METAL DERIVATIVES, PYROPHORIC MATERIALS

1687. Diethylmagnesium
[557-18-6]

C₄H₁₀Mg

Et₂Mg

Air, or Water

Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 5048

Contact with moist air usually caused ignition of the dry powder, and water always ignited the solid or its ethereal solution.

Carbon dioxide

Merck, 1983, 453

It will glow and ignite in carbon dioxide.

See other ALKYL METALS, DIALKYL MAGNESIUMS

1688. Sodium diethylamide
[34992-80-8]

C₄H₁₀NNa

NaNEt₂


Pez, G. P. *et al.*, *Pure Appl. Chem.*, 1986, **57**, 1925

The solid is highly pyrophoric and will react violently even on limited exposure to air.

See other N-METAL DERIVATIVES, PYROPHORIC MATERIALS

1689. Piperazine (Hexahydropyrazine)
[110-85-0]

C₄H₁₀N₂



HCS 1980, 756

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

See other ORGANIC BASES

1690. Diethyl hyponitrite (Diethoxydiazene)
[4549-46-6]

C₄H₁₀N₂O₂

EtON=NOEt

See entry DIALKYL HYPONITRITES

1691. N-Nitrosoethyl-2-hydroxyethylamine
[13147-25-6] **C₄H₁₀N₂O₂**




Preussman, R., *Chem. Ber.*, 1962, **95**, 1572

It decomposed explosively during attempted distillation at 18 mbar from a bath at 170°C, but was distilled with slight decomposition at 103–105°/0.4 mbar.

See other NITROSO COMPOUNDS

1692. Diethyllead dinitrate
[17498-10-1] **C₄H₁₀N₂O₆Pb**



Potts, D. *et al.*, *Can. J. Chem.*, 1969, **47**, 1621

The salt is unstable above 0°C and explodes on heating.

See related ALKYL METALS, HEAVY METAL DERIVATIVES

1693. N-Butylamidodisulfuryl azide
[13449-22-4] **C₄H₁₀N₄O₂S**



Shozda, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 2876

It exploded during analytical combustion.

See other ACYL AZIDES

1694. 1-Butanol
[78-83-1] **C₄H₁₀O**



FPA H64, 1977; *HCS 1980*, 236

Aluminium MRH 1.17/11

See Aluminium: Butanol

Chromium trioxide MRH 2.55/92

See Chromium trioxide: Alcohols

Other reactants

Yoshida, 1980, 297

MRH values calculated for 15 combinations, largely with oxidants, are given.

†**1695. 2-Butanol**
[78-92-2] **C₄H₁₀O**



HCS 1980, 237

1. Posdev, V. S. *et al.*, *Chem. Abs.*, 1977, **86**, 194347
2. Peterson, D., *Chem. Eng. News*, 1981, **59**(19), 3
3. Sharpless, T. W., *J. Chem. Educ.*, 1984, **61**, 476
4. Doyle, R. R., *Chem. Eng. News*, 1986, **64**(7), 4; *J. Chem. Educ.*, 1986, **63**(20), 186

A powerful explosion which occurred during distillation of a 10-year-old sample of the alcohol was attributed to presence of peroxy compounds formed by autoxidation, possibly involving 2-butanone as an effective photochemical sensitiser [1]. After a later explosion, it was found that the sample being distilled contained 12% of peroxide [2]. A further incident involved a 12-year old sample which exploded at the end of distillation, and which also contained a high level of peroxide. Several other stock alcohols were found to contain much lower levels of peroxide than the 2-butanol, and recommendations on clean-up or disposal, depending on the level of peroxide, are made [3]. A further report of an explosion at the end of laboratory distillation confirms the potential for peroxide formation on prolonged storage of 2-butanol [4].

See 2-Propanol: 2-Butanone

Chromium trioxide

See Chromium trioxide: Alcohols

See other PEROXIDISABLE COMPOUNDS

†1696. *tert*-Butanol

[75-65-0]

Me₃COH

C₄H₁₀O

HCS 1980, 238

Potassium–sodium alloy

See Potassium–sodium alloy: *tert*-Butanol

†1697. Diethyl ether (Ethoxyethane)

[60-29-7]

EtOEt

C₄H₁₀O

(*MCA SD-29*, 1965); *NSC* 396, 1968; *FPA H40*, 1975; *HCS* 1980, 487; *RSC Lab. Hazards Data Sheet No. 48*, 1965

1. Criegee, R. *et al.*, *Angew. Chem.*, 1936, **49**, 101
2. Criegee, R. *et al.*, *Angew. Chem.*, 1958, **70**, 261
3. Mallinckrodt, E. *et al.*, *Chem. Eng. News*, 1955, **33**, 3194
4. Ray, A., *J. Appl. Chem.*, 1955, 188
5. Anon., *Chemist & Druggist*, 1947, **157**, 258
6. Feinstein, R. N., *J. Org. Chem.*, 1959, **24**, 1172
7. Davies, A. G., *J. R. Inst. Chem.*, 1956, **80**, 386
8. Anon., *Lab. Accid. in Higher Educ.*, Item 27, HSE, Barking, 1987

The hydroperoxide initially formed by autoxidation of ether is not particularly explosive, but on standing and evaporation, polymeric 1-oxyperoxides are formed which are dangerously explosive, even below 100°C [1]. Numerous laboratory explosions have been caused by evaporation of peroxidised ether [2]. Formation of peroxide in stored ether may be prevented by presence of sodium diethyldithiocarbamate (0.05 ppm), [4], which probably deactivates traces of metals which catalyse peroxidation [2]; of pyrogallol (1 ppm) [4]; or by larger proportions (5 to 20 ppm) of other inhibitors [5]. Once present in ether, peroxides may be detected by the iodine–starch test, and removed by percolation through anion exchanger resin [6] or activated alumina [7], which leaves the ether dry, or by shaking with aqueous ferrous sulfate or sodium sulfite solutions. Many other methods have been described [7]. A small quantity of ether stored in an unmodified domestic refrigerator in a biochemistry laboratory led to an explosion, with ignition by the thermostat contacts [8].

See 1-OXYPEROXY COMPOUNDS

See other PEROXIDISABLE COMPOUNDS, POLYPEROXIDES

Boron triazide

See Boron triazide: Diethyl ether, etc.

Halogens, or Interhalogens

See Bromine trifluoride: Halogens, etc., or: Solvents

Bromine pentafluoride: Hydrogen-containing materials

Bromine: Diethyl ether

Chlorine: Diethyl ether

Iodine heptafluoride: Organic solvents

Other reactants

Yoshida, 1980, 140

MRH values calculated for 13 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Halogens, or Interhalogens, above

Chromyl chloride: Organic solvents

‘Fluorine nitrate’: Organic materials

Hydrogen peroxide: Diethyl ether

MRH 6.53/84

Iodine(VII) oxide: Diethyl ether

Lithium perchlorate: Diethyl ether

Nitric acid: Diethyl ether

MRH 5.94/79

Nitrosyl perchlorate: Organic materials

Nitryl perchlorate: Organic solvents

Ozone: Diethyl ether

Perchloric acid: Diethyl ether

Permanganic acid: Organic materials

Peroxodisulfuric acid: Organic liquids

Silver perchlorate: Diethyl ether

Sodium peroxide: Organic liquids

MRH 2.55/92

Liquid Air: Diethyl ether

Peat soils

Walkley, A., *Austral. Chem. Inst. J.*, 1939, **6**, 310

Explosions occurred during the extraction of fats and waxes from the soils with ether, as well as when heating the extract at 100°C. Although the latter is scarcely surprising (the ether contained 230 ppm of peroxides), the former observation is unusual.

Sulfur, or Sulfur compounds

See Sulfonyl chloride: Diethyl ether

Sulfur: Diethyl ether

Thiothiazyl perchlorate: Organic solvents

Uranyl nitrate

See Uranyl nitrate: Diethyl ether

Wood pulp extracts

Urso, D. F., *Chem. Eng. News*, 1957, **35**(34), 91, 115

Ethereal extracts of pulp exploded during or after concentration by evaporation. Although the ether used for the extraction previously had been freed from peroxides by treatment with cerium(III) hydroxide, the ethereal extracts had been stored for 3 weeks before concentration was effected. (During this time the ether and/or extracted terpenes would be expected to again form peroxides, but no attempt seems to have been made to test for, or to remove them before distillation was begun).

See other PEROXIDISABLE COMPOUNDS

1698. *tert*-Butyl hydroperoxide (2-Hydroperoxy-2-methylpropane)

[75-91-2]

C₄H₁₀O₂

Me₃COOH

1. Milas, N. A. *et al.*, *J. Amer. Chem. Soc.*, 1946, **68**, 205
2. Castrantas, 1965, 15
3. Sharpless, K. B. *et al.*, *Aldrichimica Acta*, 1979, **12**, 63–73
4. Verhoeff, J., *Runaway Reactions*, 1981, Paper 3/T, 1–12
5. Hill, J. H. *et al.*, *J. Org. Chem.*, 1983, **48**, 3607–3608

Though relatively stable, explosions have been caused by distillation to dryness [1], or attempted distillation at ambient pressure [2]. In a comprehensive review of the use of the hydroperoxide as a selective metal-catalysed oxygenator for alkenes and alkynes, attention is drawn to several potential hazards in this application. One specific hazard to be avoided stems from the fact that Lucidol TBHP-70 contains 19% of di-*tert*-butyl peroxide which will survive the catalysed reaction and may lead to problems in the work-up and distillation [3]. A thorough investigation of the stability and explosive properties of the 70% solution in water has been carried out [4]. The anhydrous peroxide as a solution in toluene may now readily be prepared azeotropically, and the solutions are stable in storage at ambient temperature. This solution is now a preferred method for using the anhydrous hydroperoxide [5].

See 1,2-Dichloroethane, below,

See also THERMAL EXPLOSIONS

Preparative hazard

See Hydrogen peroxide: *tert*-Butanol, Sulfuric acid

Acids

1. Sharpless, K. B. *et al.*, *Aldrichimica Acta*, 1979, **12**, 71
2. Andreozzi, R. *et al.*, *J. Haz. Mat.*, 1988, **17**, 305–313

Not even traces of strong acids should be added to high strength solutions of the hydroperoxide [1]. The thermal stability and mechanism of cleavage of mixtures with 4-toluenesulfonic acid have been studied under adiabatic conditions, and there is potential for development of a thermal runaway [2].

1,2-Dichloroethane

Hill, J. H. *et al.*, *J. Org. Chem.*, 1983, **48**, 3607

Solutions of the hydroperoxide in halogenated solvents, and especially dichloroethane are much less stable than in toluene (reference 5 above). On a large scale, the azeotropic drying of solutions of the hydroperoxide in dichloroethane may present a thermal hazard.

Molecular sieves

Sharpless, K. B. *et al.*, *Aldrichimica Acta*, 1979, **12**, 71

Use of the 4A sieve to dry the hydroperoxide (6% water) had been used routinely without incident, but inadvertent use of the larger-pored 13X sieve gave a high exotherm in the packed bed which soon led to autoignition of the hydroperoxide. This was attributed to the heat of adsorption occurring within the larger sieve pores (10 Angstroms in 13X), but caution is advised when using sieve to dry any high strength (90%) solutions of the hydroperoxide, and an azeotropic method is preferred (see reference 5 above).

See Nitromethane: Molecular sieve

See other MOLECULAR SIEVE INCIDENTS

Toluene, Dinitrogen pentoxide

1. Owens, K. A. *et al.*, *J. Loss Prevention*, 1995, **8**(5), 296
2. Editor's comments

Toluene was used, in place of tetrachloromethane, as solvent when preparing *t*-Butyl peroxyxynitrate. An explosion ensued, attributed to trinitrotoluene formation and detonation [1]. Since TNT is not easy to form and difficult to explode, this seems improbable. The intended product will certainly be a more dangerous explosive than TNT, as can be the expected byproduct, an unreacted solution of anhydrous nitric acid in toluene. Most explosions in nitration systems are due to free radical oxidation of an organic substrate or solvent by the nitrating agent; this is quite probable here. Though nitrogen pentoxide is itself more an ionic nitration reagent than a radical oxidant, all peroxy species are radical generators par excellence, and commonly used to initiate free radical reaction sequences [2].

Transition metal salts

Sharpless, K. B. *et al.*, *Aldrichimica Acta*, 1979, **12**, 71

Transition metal salts known to be good autoxidation catalysts (of cobalt, iron or manganese) should never be added to high-strength solutions of the hydroperoxide to avoid the near-certainty of vigorous evolution of a large volume of oxygen.

See other ALKYL HYDROPEROXIDES

1699. Diethyl peroxide

[628-37-5]

EtOOEt

C₄H₁₀O₂

1. Baeyer, A. *et al.*, *Ber.*, 1900, **33**, 3387
2. Castrantas, 1965, 15
3. Baker, G. *et al.*, *Chem. & Ind.*, 1964, 1988
4. Gray, P. *et al.*, *Proc. R. Soc.*, 1971, **A325**, 175

While it is acknowledged as rather explosive, the stated lack of shock-sensitivity at ambient temperature [1] is countered by its alternative description as shock-sensitive [2] and detonable [3]. The vapour explodes above a certain critical pressure, at temperatures above 190°C [4].

See other DIALKYL PEROXIDES

†1700. 1,1-Dimethoxyethane

[534-15-6]

(MeO)₂CHCH₃

C₄H₁₀O₂

HCS 1980, 417

See other PEROXIDISABLE COMPOUNDS

†1701. 1,2-Dimethoxyethane

[110-71-4]

MeOC₂H₄OMe

C₄H₁₀O₂

HCS 1980, 418

Phifer, L. H., *CHAS Notes*, 1982, **1**(3), 2

A seven-year-old peroxidised sample exploded during distillation.

Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: 1,2-Dimethoxyethane

See other PEROXIDISABLE COMPOUNDS

1702. 2-Ethoxyethanol

[110-80-5]

EtOC₂H₄OH

C₄H₁₀O₂

HCS 1980, 459

Hydrogen peroxide, Poly(acrylamide) gel, etc.
See Hydrogen peroxide: 2-Ethoxyethanol, etc.
See other PEROXIDISABLE COMPOUNDS

†1703. Methyl propyl ether (1-Methoxypropane)
[557-17-5]



HCS 1980, 660

1704. Zinc ethoxide
[3851-22-7]



Nitric acid

Solomartin, V. S. *et al.*, *Chem. Abs.*, 1977, **87**, 210678

In the analysis of diethylzinc, a 1 ml sample is cooled to -196°C and treated with 2 ml of ethanol to give the ethoxide. During subsequent conversion to zinc nitrate (prior to pyrolysis to the oxide) by treatment with 3 ml of 30% nitric acid, cooling must be continued to avoid an explosion hazard.

See other METAL ALKOXIDES

1705. Bis(2-hydroxyethyl) ether (Diethylene glycol)
[111-46-6]



HCS 1980, 397

Other reactants

Yoshida, 1980, 144

MRH values calculated for 14 combinations, largely with oxidants, are given.

Sodium hydroxide

See Sodium hydroxide: Glycols

1706. 1-Hydroxy-3-butyl hydroperoxide (3-Hydroperoxy-1-butanol)
[]



Rieche, A., *Ber.*, 1930, **63**, 2462

It explodes on heating.

1707. 2-Methoxyprop-2-yl hydroperoxide (2-Methoxy-1-methylethyl hydroperoxide)

[10027-74-4]

C₄H₁₀O₃



Dussault, P., *Chem. Eng. News*, 1993, **71**(32), 2

This reagent, made by methanolic ozonolysis of tetramethylethylene, may deflagrate if isolated.

See Hydrogen peroxide: Acetone

See related 1-OXYPEROXY COMPOUNDS

†**1708. Trimethyl orthoformate (Trimethoxymethane)**

[149-73-5]

C₄H₁₀O₃




Preparative hazard

See Chloroform: Sodium, Methanol, or: Sodium methoxide

1709. Diethyl sulfite

[623-81-4]

C₄H₁₀O₃S



Chlorine fluoride


See Ethyl fluorosulfate

See other SULFUR ESTERS

1710. Diethyl sulfate

[64-67-5]

C₄H₁₀O₄S



HCS 1980, 406

3,8-Dinitro-6-phenylphenanthridine, Water

Hodgson, J. F., *Chem. & Ind.*, 1968, 1399

Accidental ingress of water to the heated mixture liberated sulfuric acid or its half ester which caused a violent reaction with generation of a large volume of solid black foam.

See 4-Nitroaniline-2-sulfonic acid, also Sulfuric acid: Nitroaryl bases, etc.

Iron, Water

Siebeneicher, K., *Angew. Chem.*, 1934, **47**, 105

Moisture in a sealed iron drum of the ester caused hydrolysis to sulfuric acid, leading to corrosion of the metal and development of a high internal pressure of hydrogen, which ruptured the drum.

See other CORROSION INCIDENTS, GAS EVOLUTION INCIDENTS

Potassium *tert*-butoxide
See Potassium *tert*-butoxide: Acids, etc.
See other SULFUR ESTERS

1711. Zinc ethylsulfinate
[24308-92-7]



Preparative hazard
See Diethylzinc: Sulfur dioxide
See related SULFUR ESTERS

†**1712. Butanethiol**
[109-79-5]



Nitric acid
See Nitric acid: Alkanethiols
See other ALKANETHIOLS

†**1713. 2-Butanethiol**
[513-53-1]



See other ALKANETHIOLS

†**1714. Diethyl sulfide (3-Thiapentane)**
[352-93-2]



HCS 1980, 407

†**1715. 2-Methylpropanethiol**
[513-44-0]



Calcium hypochlorite
See Calcium hypochlorite: Organic sulfur compounds
See other ALKANETHIOLS

†**1716. 2-Methyl-2-propanethiol**
[75-66-1]



See other ALKANETHIOLS

1717. Diethyl telluride
[627-54-3]



Ellern, 1968, 24–25
It ignites in air.
See other ALKYL METALS

1718. Diethylzinc
[557-20-0]



Aluminium Alkyls and other Organometallics, 5, New York, Ethyl Corp., 1967

It is immediately pyrophoric in air.

Alkenes, Diiodomethane

Houben-Weyl, 1973, Vol. 13.2a, 845

During preparation of cyclopropane derivatives, it is important to add the diiodomethane slowly to a solution of diethylzinc in the alkene. Addition of diethylzinc to an alkene–diiodomethane mixture may be explosively violent.

Halogens

See Bromine: Diethylzinc

Chlorine: Diethylzinc

Methanol

Houben-Weyl, 1973, Vol. 13.2a, 855

Interaction is explosively violent and ignition ensues.

Nitro compounds

Leleu, *Cahiers*, 1977, (88), 371

Interaction is usually violent.

Non-metal halides

Leleu, *Cahiers*, 1977, (88), 372

Interaction with arsenic trichloride or phosphorus trichloride is violent, forming pyrophoric triethylarsine or triethylphosphine.

Ozone

See Ozone: Alkylmetals

Sulfur dioxide

Houben-Weyl, 1973, 13.2a, 709

During preparation of zinc ethylsulfinate, addition of diethylzinc to liquid sulfur dioxide at –15°C leads to an explosively violent reaction. Condensation of sulfur dioxide into cold diethylzinc leads to a controllable reaction on warming.

Water

Ito, Y. *et al.*, *Org. Synth.*, 1979, **59**, 113, 116

It reacts violently with water, but a procedure for destroying excess reagent with ice-water is detailed.

See other ALKYL METALS, DIALKYLZINC

1719. Diethylaluminium hydride

[871-27-2]



$\text{C}_4\text{H}_{11}\text{Al}$

HCS 1980, 391

See entry ALKYLALUMINIUM ALKOXIDES AND HYDRIDES

1720. Diethylarsine

[692-42-2]



$\text{C}_4\text{H}_{11}\text{As}$

Sidgwick, 1950, 762

It ignites in air, even at 0°C.

See other ALKYLNON-METAL HYDRIDES

1721. Diethylgallium hydride (Diethylhydrogallium)

[93481-56-2]



$\text{C}_4\text{H}_{11}\text{Ga}$

Air, or Water

Eisch, J. J., *J. Amer. Chem. Soc.*, 1962, **84**, 3835

It ignites in air, and reacts violently with water.

See other ALKYL METAL HYDRIDES

1722. Trimethylsilylmethyl lithium

[1872-00-0]



$\text{C}_4\text{H}_{11}\text{LiSi}$

Tessier-Youngs, C. *et al.*, *Inorg. Synth.*, 1986, **24**, 92

It ignites spontaneously in air and reacts violently with water, like the aluminium analogue.

See related ALKYL METALS, ALKYL SILANES

†**1723. Butylamine (1-Aminobutane)**

[109-73-9]



$\text{C}_4\text{H}_{11}\text{N}$

FPA H100, 1982; *HCS* 1980, 239

Other reactants

Yoshida, 1980, 305

MRH values calculated for 13 combinations with oxidants are given.

Perchloryl fluoride

See Perchloryl fluoride: Nitrogenous bases

See other ORGANIC BASES

†1724. 2-Butylamine (2-Aminobutane)

[13952-84-6]



C₄H₁₁N

See other ORGANIC BASES

†1725. *tert*-Butylamine (2-Amino-2-methylpropane)

[75-64-9]



C₄H₁₁N

HCS 1980, 240


2,2-Dibromo-1,3-dimethylcyclopropanoic acid

See 2,2-Dibromo-1,3-dimethylcyclopropanoic acid: *tert*-Butylamine

See other ORGANIC BASES

†1726. Diethylamine

[109-89-7]



C₄H₁₁N

(MCA SD-97, 1971); FPA H85, 1979; HCS 1980, 392

Cellulose nitrate

See CELLULOSE NITRATE: amines

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

Other reactants

Yoshida, 1980, 139

MRH values calculated for 13 combinations with oxidants are given.

Sulfuric acid

See Sulfuric acid: Diethylamine

See other ORGANIC BASES

†1727. Ethyldimethylamine

[598-56-1]



C₄H₁₁N

See other ORGANIC BASES

†1728. Isobutylamine (2-Methypropylamine)
[78-81-9] C₄H₁₁N
 $\text{Me}_2\text{CHCH}_2\text{NH}_2$

See other ORGANIC BASES

1729. N-2-Hydroxyethyl dimethylamine (2-Dimethylaminoethanol)
[108-01-0] C₄H₁₁NO
 $\text{HOC}_2\text{H}_4\text{NMe}_2$

HCS 1980, 422

Cellulose nitrate

See CELLULOSE NITRATE: amines

Other reactants

Yoshida, 1980, 167

MRH values calculated for 14 combinations with oxidants are given.

See other ORGANIC BASES

1730. 3-Methoxypropylamine
[5332-73-0] C₄H₁₁NO
 $\text{MeOC}_3\text{H}_6\text{NH}_2$

Energy of decomposition (in range 150–550°C) measured as 0.57 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other AMINOMETHOXY COMPOUNDS

1731. Tris(hydroxymethyl)methylamine
[77-86-1] C₄H₁₁NO₃
 $(\text{HOCH}_2)_3\text{CNH}_2$

Preparative hazard

See Tris(hydroxymethyl)nitromethane: Hydrogen, etc.

See other ORGANIC BASES

1732. Dimethyl ethanephosphonite
[15715-42-1] C₄H₁₁O₂P
 $(\text{MeO})_2\text{PEt}$

Arbuzov, B. A. *et al.*, *Chem. Abs.*, 1953, **47**, 3226a

It ignites in air when absorbed onto filter paper.

See other PHOSPHORUS ESTERS, PYROPHORIC MATERIALS

- 1733. Diethyl phosphite**
[762-04-9] **C₄H₁₁O₃P**
(EtO)₂P(:O)H
- 4-Nitrophenol
 Phifer, L. H., private comm., 1982
 Interaction in absence of solvent in a stirred flask heated by a regulated mantle led to a runaway reaction and explosion.
See other PHOSPHORUS ESTERS
- 1734. Diethylphosphine**
[627-49-6] **C₄H₁₁P**
Et₂PH
- Houben-Weyl 1963, Vol. 12.1, 69
 Secondary lower-alkylphosphines readily ignite in air.
See other ALKYLPHOSPHINES
- 1735. Ethyldimethylphosphine**
[1605-51-2] **C₄H₁₁P**
EtPMe₂
- Smith, J. F., private comm., 1970
 May ignite in air.
See other ALKYLPHOSPHINES
- 1736. Tetramethyldiarsane (Tetramethyldiarsine)**
[471-35-2] **C₄H₁₂As₂**
Me₂AsAsMe₂
- Sidgwick, 1950, 770
 Inflames in air.
See other ALKYLNON-METALS
- 1737. Bis(dimethylarsinyl) oxide**
[503-80-0] **C₄H₁₂As₂O**
(Me₂As)₂O
- von Schwartz, 1918, 322
 Ignites in air.
See related ALKYLNON-METALS

1738. Bis(dimethylarsinyl) sulfideCAS591-10-6C₄H₁₂As₂S

von Schwartz, 1918, 322

Ignites in air.

See related ALKYLNON-METALS**1739. Tetramethyldigold diazide**


[22653-19-6]

Beck, W. *et al.*, *Inorg. Chim. Acta*, 1968, **2**, 468

The dimeric azide is extremely sensitive and may explode under water if touched.

See other GOLD COMPOUNDS, METAL AZIDES**1740. Lithium tetramethylborate**

[2169-38-2]




491M, 1975, 238


It may ignite in moist air.

See related ALKYLNON-METALS**1741. Dihydrzinecadmium(II) succinate (Bis(hydrazine)[butanedioato(2-)] cadmium)**

[159793-67-6]

*See* HYDRAZINE METAL MALONATES AND SUCCINATES**1742. Tetramethylammonium chlorite**


[67922-18-3]

Levi, *Gazz. Chim. Ital.* [1], 1922, **52**, 207

The dry solid explodes on impact.

See other CHLORITE SALTS, OXOSALTS OF NITROGENOUS BASES**1743. 1,2-Dichlorotetramethyldisilane**

[39437-99-5]

Rich, J. D., *Chem. Eng. News*, 1990, **68**(13), 2


Explosions were experienced in the still-head when distilling this compound. The cause is not very clear and may have to do with impurities.

See Chloropentamethyldisilane,

See other ALKYLHALOSILANES

1744. Lithium tetramethylchromate(II)

[]




Kurras, E. *et al.*, *J. Organomet. Chem.*, 1965, **4**, 114–118

Isolated as a dioxane complex, it ignites in air.

See other ALKYLMETALS

1745. Tetramethylammonium monoperchromate

[]



Mellor, 1943, Vol. 11, 358

It explodes on moderate heating or in contact with sulfuric acid. Now thought probably to have the structure shown (and corresponding empirical formula) it has in the intervening period been considered to be a dimer, a pentaperoxydichromate.

See Tetramethylammonium pentaperoxydichromate

See other OXOSALTS OF NITROGENOUS BASES, PEROXOACID SALTS, PEROXOCHROMIUM COMPOUNDS, QUATERNARY OXIDANTS

1746. 1,2-Diamino-2-methylpropaneoxodiperoxychromium(VI)

[]





House, D. A. *et al.*, *Inorg. Chem.*, 1967, **6**, 1078 (footnote 6)

This blue precipitate is formed intermediately during preparation of 1,2-diamino-2-methylpropaneaquadiperoxychromium(IV) monohydrate, and its analogues are dangerously explosive and should not be isolated without precautions.

See other AMMINECHROMIUM PEROXOCOMPLEXES

1747. Tetramethylammonium pentafluoroxenoxide

[161535-67-6]




Christe, K. O. *et al.*, *Inorg. Chem.*, 1995, **34**(7), 1868

Not surprisingly, this salt is highly explosive.

See other XENON COMPOUNDS, See also QUATERNARY OXIDANTS

1748. Tetramethyldigallane
[65313-37-3]





Leleu, *Cahiers*, 1977, (88), 366

It ignites in air.

See other ALKYL METALS

1749. Tetramethylammonium periodate (*N,N,N*-Trimethylmethaniminium periodate)
[55999-69-4]



1. Wagner, R. I. *et al.*, *Inorg. Chem.*, 1997, **36**(12), 2564



2. Malpass, J. R. *et al.*, *Chem. Brit.*, 1998, **34**(12), 18

3. Cullis, P. M., Personal communication, 1998

Explodes violently on heating above 240°C [1]. A 14g sample exploded while being transferred to a cold, empty, unagitated flask via a polythene powder funnel, injuring the worker. The isolated salt was essentially neutral, so excess periodic acid cannot be involved. Subsequent attempts to explode the material deliberately have failed, but it is felt that static electricity may be the cause [2], [3].

See other QUATERNARY OXIDANTS

1750. Tetramethylammonium diazidoiodate(I)
[68574-13-0]




Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 512

Highly explosive crystals.

See related NON-METAL AZIDES

1751. Tetramethylammonium superoxide
[3946-86-9]



1. McElroy, A. O. *et al.*, *Inorg. Chem.*, 1964, **3**, 1798

2. Foote, C. S., *Photochem. Photobiol.*, 1978, **28**, 718–719

During the preparation by solid phase interaction of tetramethylammonium hydroxide and potassium superoxide [1] by tumbling for several days in a rotary evaporator flask, a violent explosion occurred [2]. This may have been caused by ingress of grease or other organic material leading to contact with potassium superoxide, a powerful oxidant.

See Potassium dioxide: Organic materials

See other QUATERNARY OXIDANTS

- 1752. *N*-(Diethylphosphinoyl)hydroxylamine**
 [] **C₄H₁₂NO₂P**
 $\text{Et}_2\text{P}(\text{O})\text{NHOH}$
 Harger, M. J. P. *et al.*, *Tetrahedron Lett.*, 1991, **32**(36), 4769
 Described as a hazardously unstable oil.
See other N–O COMPOUNDS
- 1753. Tetramethylammonium ozonate (*N,N,N*-Trimethylmethanaminium ozonide)**
 [78657-29-1] **C₄H₁₂NO₃**
 $\text{Me}_4\text{N}^+\text{O}_3^-$
 Preparative hazard
See Ozone: Tetramethylammonium hydroxide
See other QUATERNARY OXIDANTS
- 1754. Tetrakis(hydroxymethyl)phosphonium nitrate**
 [24748-25-2] **C₄H₁₂NO₇P**
 $(\text{HOCH}_2)_4\text{P}^+\text{NO}_3^-$
 Calamari, T. A. *et al.*, *Chem. Eng. News*, 1979, **57**(12), 74
 A violent explosion occurred during prolonged azeotropic drying at 105–110°C of a 75 wt% benzene solution of the salt. Traces of a nitrate ester may have been formed from a slight excess of nitrate ion.
See related OXOSALTS OF NITROGENOUS BASES, REDOX COMPOUNDS
- †**1755. 2-Dimethylaminoethylamine**
 [108-00-9] **C₄H₁₂N₂**
 $\text{Me}_2\text{NC}_2\text{H}_4\text{NH}_2$
See other ORGANIC BASES
- 1756. *N*-2-Hydroxyethyl-1,2-diaminoethane (2-(2-Aminoethyl)aminoethanol)**
 [111-41-1] **C₄H₁₂N₂O**
 $\text{HOC}_2\text{H}_4\text{NHC}_2\text{H}_4\text{NH}_2$
HCS 1980, 555
 Cellulose nitrate
See CELLULOSE NITRATE: amines *See other* ORGANIC BASES
- 1757. Bis(dimethylamino) sulfoxide**
 [3768-60-3] **C₄H₁₂N₂OS**
 $\text{Me}_2\text{NS}(\text{:O})\text{NMe}_2$
 Sulfinyl chloride
See Sulfinyl chloride: Bis(dimethylamino) sulfoxide
See other N–S COMPOUNDS

1758. Seleninyl bis(dimethylamide)

[2424-09-1]


C₄H₁₂N₂OSe

Bailar, 1973, Vol. 2, 975

Vacuum distillation at temperatures around 50–60°C leads to explosive decomposition.

See related N-METAL DERIVATIVES, N-S COMPOUNDS**1759. Tetramethyl-2-tetrazene**

[6130-87-6]


C₄H₁₂N₄

1. Houben-Weyl, 1967, Vol. 10.2, 830

2. Stull, 1977, 22

It explodes above its b.p., 130°C/1 bar [1], and is strongly endothermic (ΔH_f° (l) +226.4 kJ/mol, 1.98 kJ/g) [2].*See other* ENDOTHERMIC COMPOUNDS, TETRAZENES**1760. Dihydrzinenickel(II) succinate (Bis(hydrazine)[butanedioato(2-)]nickel)**

[45101-14-2]

C₄H₁₂N₄NiO₄*See* HYDRAZINE METAL MALONATES AND SUCCINATES**1761. Bis(dimethylstibinyl) oxide**

[]


C₄H₁₂OSb₂

Sidgwick, 1950, 777

It ignites in air.

See related ALKYL METALS**1762. 1,1-Diethoxy-1,3-disiladioxetane**

[]

C₄H₁₂O₄Si₂

491M, 1975, 150

580

It ignites in air.
See related NON-METAL HYDRIDES

1763. Diethylhydroxytin hydroperoxide


[]



Bailar, 1973, Vol. 2, 68
It is explosive.
See other ORGANOMETALLIC PEROXIDES

†**1764. Tetramethoxysilane (Tetramethyl silicate)**

[681-84-5]



HCS 1980, 662

Metal hexafluorides

Jacob, E., *Angew. Chem. (Intern. Ed.)*, 1982, **21**, 143, (footnote 6)

During the preparation of the hexamethoxides of rhenium, molybdenum and tungsten by co-condensation with excess tetramethoxysilane on a cold surface, simultaneous co-condensation is necessary to avoid the danger of explosion present when sequential condensation of the reactants is employed. In the latter case, the high concentrations of hexafluoride at the interface leads to violent reaction with the silane.

See related SILANES

1765. Tetramethyldiphosphane (Tetramethyldiphosphine)

[3676-91-3]



1. Kardosky, G. *et al.*, *Inorg. Synth.*, 1973, **14**, 15
2. Goldsberry, R. *et al.*, *Inorg. Synth.*, 1972, **13**, 28


It ignites in air [1], and unreacted tetramethyldiphosphane residues may also ignite [2].

See Bis(tetramethyldiphosphane disulfide)cadmium perchlorate

See other ALKYLPHOSPHINES

1766. Tetramethyldiphosphane disulfide

[3676-97-9]



Preparative hazard

1. Parshall, G. W., *Org. Synth.*, 1975, Coll. Vol. 5, 1016
2. Butter, A. S. *et al.*, *Inorg. Synth.*, 1974, **15**, 186
3. Davies, S. G., *Chem. Brit.*, 1984, **20**, 403
4. Bercauw, J. E., *Chem. Eng. News*, 1984, **62**(18), 4
5. Burt, R. J. *et al.*, *J. Organomet. Chem.*, 1979, **182**, 203

6. Bercaw, J. E. *et al.*, *Inorg. Synth.*, 1985, **25**, 199

7. Parshall, G. W., *Org. Synth.*, 1985, **63**, 226

The published preparation of the disulfide from thiophosphoryl chloride (4.28 mole) and methylmagnesium iodide (4.0 mole) [1,2] is hazardous. On the stated scale and at the recommended rate of addition of the 47% solution of thiophosphoryl chloride in ether, it is difficult to maintain the temperature at 0–5°C. Even at below one tenth of the recommended addition rate, after half of the solution had been added during 9 hours, the temperature rose uncontrollably from –5 to +60 °C in 10 s and a violent explosion followed, pulverising the apparatus [3]. A closely similar accident was simultaneously reported [4]. An alternative method is recommended [5], the product of which, though air-sensitive, does not ignite in air as previously stated. In the light of recent experience, the procedure in [1] is preferred to that in [2] because more dilute solutions are used in the former method [6]. To avoid explosions in the preparation of the title compound from methylmagnesium bromide and thiophosphoryl chloride in ether, the chloride should be redistilled and reaction vessel cooling should be with ice–salt rather than acetone–carbon dioxide, to assist in maintaining the reaction temperature in the range 0–5°C. If the temperature drops below –5°C, stop the addition and cautiously warm to 0–5°C [7].

See related ALKYLPHOSPHINES, NON-METAL SULFIDES

1767. Tetramethyllead (Tetramethylplumbane)

[75-74-1]

C₄H₁₂Pb

Me₄Pb

HCS 1980, 895

Sidgwick, 1950, 463

It is liable to explode violently above 90°C.

Tetrachlorotrifluoromethylphosphorane

See Tetrachlorotrifluoromethylphosphorane: Tetramethyllead

See other ALKYLMETALS

1768. Tetramethylplatinum

[22295-11-0]

C₄H₁₂Pt

Me₄Pt

Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1953, **75**, 2065

It explodes weakly on heating.

See Trimethylplatinum hydroxide

See other ALKYLMETALS, PLATINUM COMPOUNDS

1769. Tetramethyldistibane (Tetramethyldistibine)

[41422-43-9]

C₄H₁₂Sb₂



Me₂SbSbMe₂

Sidgwick, 1950, 779

- It ignites in air.
See other ALKYL METALS
- †1770. **Tetramethylsilane**
 [75-76-3] **C₄H₁₂Si**
Me₄Si
- Chlorine
See Chlorine: Antimony trichloride, etc.
See other ALKYL SILANES
- †1771. **Tetramethyltin (Tetramethylstannane)**
 [594-27-4] **C₄H₁₂Sn**
Me₄Sn
- Dinitrogen tetraoxide
See Dinitrogen tetraoxide: Tetramethyltin
See other ALKYL METALS
1772. **Tetramethyltellurium(IV)**
 [123311-08-0] **C₄H₁₂Te**
Me₄Te
- Gedridge, R. W. *et al.*, *Organometallics*, 1989, **8**(12), 2817
 Extremely pyrophoric, sometimes explosively so.
See other ALKYL METALS
1773. **Bis(2-aminoethyl)aminesilver nitrate**
 [] **C₄H₁₃AgN₄O₃**
[HN(C₂H₄NH₂)₂Ag] NO₃
- Tennhouse, G. J. *et al.*, *J. Inorg. Nucl. Chem.*, 1966, **28**, 682–684
 The complex explodes at 200°C.
See other AMMINEMETAL NITRATES, SILVER COMPOUNDS
1774. **1,1,3,3-Tetramethylsiloxalane**
 [] **C₄H₁₃AlOSi**
(Me)₂SiHOAl(Me)₂
- This compound, which exists as the dimer, explodes on contact with water.
See ALKYL ALUMINIUM DERIVATIVES, *See related* ALKYL SILANES
1775. **Bis(2-aminoethyl)aminecobalt(III) azide**
 [26493-63-0] **C₄H₁₃CoN₁₂**
[HN(C₂H₄NH₂)₂Co] [N₃]₃
- Druding, L. F. *et al.*, *Acta Cryst.*, 1974, **B30**, 2386


The shock-sensitivity and dangerously explosive nature is stressed.
See AMMINECOBALT(III) AZIDES See related METAL AZIDES

1776. Bis(2-aminoethyl)aminediperoxochromium(IV)
[59419-71-5]



1. House, D. A. *et al.*, *Inorg. Chem.*, 1966, **5**, 840
 2. Ghosh, S. K. *et al.*, *J. Organomet. Chem.*, 1988, **340**(1), 59
- The monohydrate explodes at 109–110°C during slow heating [1]. The solid complex exploded when manipulated on a glass sinter with a metal spatula [1]. Preparation on small scale and use of plastic equipment is recommended [2].
See other AMMINECHROMIUM PEROXOCOMPLEXES

1777. Bis(2-aminoethyl)amine (Diethylenetriamine)
[111-40-0]



HCS 1980, 399

Cellulose nitrate

See CELLULOSE NITRATE: amines

Nitromethane

See Nitromethane: Bis(2-aminoethyl)amine


Other reactants

Yoshida, 1980, 147

MRH values calculated for 13 combinations with oxidants are given.

See other ORGANIC BASES

1778. Tetramethyldialuminium dihydride
[33196-65-5]




Wiberg, W. E. *et al.*, *Angew. Chem.*, 1939, **52**, 372

It ignites and burns explosively in air.



See other ALKYLALUMINIUM ALKOXIDES AND HYDRIDES

1779. Tetramethyldiborane
[21482-59-7]




Leleu, *Cahiers*, 1977, (88), 365
It ignites in air.
See other ALKYLBORANES

1780. Diacetatotetraaquocobalt (Cobalt(II) acetate tetrahydrate)
[6147-53-1]




Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455
Energy of decomposition (270–340°C) was determined by DSC as 2.42 kJ/g, peaking at 328°C. The exotherm is largely from combustion of the evolved carbon monoxide and acetone in air.
See other HEAVY METAL DERIVATIVES, REDUCANTS

1781. 1,2-Diamino-2-methylpropaneaquadiperoxochromium(IV)
[17168-83-1]



House, D. A. *et al.*, *Inorg. Chem.*, 1967, **6**, 1078
The monohydrate exploded at 83–84°C during slow heating, and is potentially explosive at 20–25°C. There is also a preparative hazard.
See 1,2-Diamino-2-methylpropaneoxodiperoxochromium(VI)
See other AMMINECHROMIUM PEROXOCOMPLEXES

1782. Tetramethylammonium amide
[13422-81-6]



Ammonia
Musker, W. K., *J. Org. Chem.*, 1967, **32**, 3189


During the preparation, the liquid ammonia used as solvent must be removed completely at -45°C . The compound decomposes explosively at ambient temperature in presence of ammonia.

See Sodium: Ammonia, Aromatic hydrocarbons

See related N-METAL DERIVATIVES

1783. Tetramethyldisiloxane

[3277-26-7]



Allylphenols, Platinum

See SILANES: olefins, platinum


Oxygen, Sodium hydroxide, Water

See Oxygen (Gas): Tetramethyldisiloxane

See related ALKYL SILANES

1784. Bis(1,2-diaminoethane)dinitrocobalt(III) perchlorate

[[14781-32-9] (c-) [14781-33-0] (t-)]



Seel, F. *et al.*, *Z. Anorg. Chem.*, 1974, **408**, 281

The *cis*- and *trans*-isomers are both dangerously explosive compounds.

See other AMMINOMETAL OXOSALTS

1785. Tetrakis(thiourea)manganese(II) perchlorate

[50831-29-3]





Karnaikhov, A. S. *et al.*, *Chem. Abs.*, 1973, **79**, 140195

The complex decomposes explosively at 257°C .

See related AMMINOMETAL OXOSALTS

1786. Bis(1,2-diaminoethane)dichlorocobalt(III) chlorate

[26388-78-3]




It explodes at 320°C .

See entry AMMINOMETAL OXOSALTS (reference 2)

1787. Bis(1,2-diaminoethane)dichlorocobalt(III) perchlorate

[14932-06-0]



It explodes at 300°C ; low impact-sensitivity.

See entry AMMINOMETAL OXOSALTS (reference 2)

1788. cis-Bis(1,2-diaminoethane)dinitrocobalt(III) iodate

[]



Lobanov, N. I., *Zh. Neorg. Khim.*, 1959, **4**, 151

It dissociates explosively on heating.

See other AMMINEMETAL OXOSALTS

1789. Diazido-bis(1,2-diaminoethane)ruthenium(II) hexafluorophosphate

[30459-28-0]





Kane-Maguire, L. A. P. *et al.*, *Inorg. Synth.*, 1970, **12**, 24

Small explosions were observed on scratching the crystals with a metal spatula.

See related METAL AZIDES

1790. 2,4,6,8-Tetramethylcyclotetrasiloxane

[2370-88-9]



See 2,4,6,8-Tetraethenyl-2,4,6,8-tetramethylcyclotetrasiloxane

See other SILANES

1791. Bis(1,2-diaminoethane)hydroxooxorhenium(V) perchlorate

[19267-68-6]




Murmann, R. K. *et al.*, *Inorg. Synth.*, 1966, **8**, 174–175

It explodes violently when dried at above ambient temperature and is also shock-sensitive. Several explosions occurred during analytical combustion.

See other AMMINEMETAL OXOSALTS


1792. Pentaamminepyrazineruthenium(II) perchlorate

[41481-90-7]




Creutz, C. A., private comm., 1969


After ether washing, 30 mg of the salt exploded violently when disturbed.
See other AMMINEMETAL OXOSALTS




Seel, F. *et al.*, *Z. Anorg. Chem.*, 1974, **408**, 281
During evaporation of a solution of the complex salt, a very violent explosion occurred.
See other AMMINEMETAL OXOSALTS




Evans, I. P. *et al.*, *J. Amer. Chem. Soc.*, 1977, **98**, 8042
It is explosive.
See other AMMINEMETAL OXOSALTS



Air, or Water
Burg, A. B. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 957
The impure oily product ignites in air and reacts violently with water.
See other COMPLEX HYDRIDES *See related* BORANES



See entry AZOCARBABORANES



Ammonia
Pieters, 1957, 30
Contact may be explosive.
See Potassium hexacyanoferrate(III): Ammonia
See other MERCURY COMPOUNDS *See related* METAL CYANIDES

1798. μ -1,2-Bis(cyanomercurio)ethanediylidenedimercury(II)
[] **C₄Hg₄N₂**
NCHgC(:Hg)C(:Hg)HgCN

Whitmore, 1921, 128
It explodes slightly on heating.
See related METAL CYANIDES *See other* MERCURY COMPOUNDS

1799. 1,4-Diiodo-1,3-butadiyne
[53214-97-4] **C₄I₂**
IC \equiv CC \equiv CI

Sorbe, 1968, 66
It explodes at about 100°C.
See other HALOACETYLENE DERIVATIVES

1800. Potassium tetracyanotitanate(IV)
[75038-71-0] **C₄K₄N₄Ti**
K₄[Ti(CN)₄]

Water
Nicholls, D. *et al.*, *J. Chem. Soc., Chem Comm.*, 1974, 635–636
Interaction is violent.
See other METAL CYANIDES

†**1801. Dicyanoacetylene (2-Butynedinitrile)**
[1071-98-3] **C₄N₂**
N \equiv CC \equiv CC \equiv N

1. Kirshenbaum, A. D. *et al.*, *J. Amer. Chem. Soc.*, 1956, **78**, 2020
2. Robinson, Rodd, 1965, Vol. ID, 338, 366
3. Ciganek, E. *et al.*, *J. Org. Chem.*, 1968, **33**, 542
As anticipated for a conjugated multiple triply bound structure, it is extremely endothermic (ΔH_f° (g) +626.4 kJ/mol, 8.22 kJ/g), and may decompose explosively to carbon powder and nitrogen. When burnt in oxygen a flame temperature exceeding 4700°C is attained [1], and it ignites in air at 130°C [2]. It is potentially explosive in the pure state or in conc. solutions, but is fairly stable in dilute solution [3]. It rapidly polymerises with water, weak bases, alcohols, etc. [2].
See related HALOACETYLENE DERIVATIVES
See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS


1802. Sodium tetracyanatopalladate(II)
[] **C₄N₄Na₂O₄Pd**
Na₂[Pd(OCN)₄]

Bailar, 1973, Vol. 3, 1288

It explodes on heating or impact.
See related METAL CYANATES

1803. Dicyanofurazan
[55644-07-0]

C₄N₄O



Homewood, R. H. *et al.*, US Pat. 3 832 249, 1974

Although highly endothermic (ΔH_f° (s) +456.3 kJ/mol, 3.80 kJ/g), it is a stable and relatively insensitive but powerful explosive.

Nitrogenous bases


Denson, D. B. *et al.*, US Pat. 3 740 947, 1973

Contact of dicyanofurazan, or its *N*-oxide (dicyanofuroxan), with hydrazine, mono- or di-methylhydrazine, piperidine, piperazine, diethylamine or their mixtures is instantaneously explosive.

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS, N-O COMPOUNDS

1804. Dicyanofurazan *N*-oxide (Dicyanofuroxan)
[55644-07-0]

C₄N₄O₂



It is highly endothermic (ΔH_f° (s) +565.5 kJ/mol, 3.43 kJ/g).

Nitrogenous bases

Denson, D. B. *et al.*, US Pat. 3 740 947, 1973

Contact of dicyanofurazan *N*-oxide or the parent furazan with hydrazine, mono- or di-methylhydrazine, piperidine, piperazine, diethylamine or their mixtures is instantaneously explosive.

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS, FURAZAN *N*-OXIDES

†1805. Tetracarbonylnickel (Nickel tetracarbonyl)
[13463-39-3]

C₄NiO₄

(CO)₄Ni

HCS 1980, 885

Bromine

Blanchard, A. A. *et al.*, *J. Amer. Chem. Soc.*, 1926, **48**, 872

The two compounds interact explosively in the liquid state, but smoothly as vapour.

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Tetracarbonylnickel

Mercury, Oxygen

Mellor, 1946, Vol. 5, 955

A mixture of the dry carbonyl and oxygen will explode on vigorous shaking with mercury (presumably catalysed by mercury or its oxide).

Oxygen, Butane

1. Egerton, A. *et al.*, *Proc. R. Soc.*, 1954, **A225**, 427

2. Badin, E. J. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 2055

The carbonyl on exposure to atmospheric oxygen produces a deposit which becomes peroxidised and may ignite. Mixtures with air or oxygen at low partial and total pressures explode after a variable induction period [1]. Addition of the carbonyl to a butane–oxygen mixture at 20–40°C caused explosive reaction in some cases [2].

See other INDUCTION PERIOD INCIDENTS


Tetrachloropropadiene

See Dicarbonyl- η -trichloropropenyldinickel chloride dimer

See other CARBONYLMETALS

1806. Hexafluoroglutaryl dihypochlorite

[71359-64-3]




Tari, I. *et al.*, *Inorg. Chem.*, 1979, **18**, 3205–3208

Explosive above -10°C .

See other ACYL HYPOHALITES

1807. Tetrachlorodiazocyclopentadiene

[21572-61-2]



Abel, E. W. *et al.*, *Inorg. Chim. Acta*, 1980, **44**(3), L161–163


Conversion to octachloronaphthalene by heating at 150°C in absence of solvent was explosive.

See other DIAZO COMPOUNDS

1808. Hexachlorocyclopentadiene

[77-47-4]

C₅Cl₆



Sodium


See Sodium: Halocarbons

See other HALOALKENES

1809. Potassium azidopentacyanocobaltate(3-)

[14705-99-8]

C₅CoK₃N₈



Huan, J. *et al.*, *Thermochim. Acta*, 1988, **130**, 77–85


Thermal decomposition under hydrogen of a series of pentacyanocobaltate complexes (CN-, NO₂-, NO- or N₃-ligands) revealed that the latter complex is the most exothermic by far. Presence of iron powder suppresses hydrogen cyanide formation.

See related METAL AZIDES, METAL CYANIDES

1810. Potassium pentacyanodiperoxochromate(5-)

[]

C₅CrK₅N₅O₄





Bailar, 1973, Vol. 4, 167

A highly explosive material, with internal redox features.

See related AMMINECHROMIUM PEROXOCOMPLEXES See other REDOX COMPOUNDS

1811. Caesium pentacarbonylvanadate(3-)
[78937-12-9]




Air, or Hydroxy compounds

Ellis, J. F. *et al.*, *J. Amer. Chem. Soc.*, 1981, **103**, 6101

Pyrophoric in air, deflagrates under nitrogen on scratching with a metal spatula, and reacts explosively with alcohols or water.

See related CARBONYLMETALS

1812. 2-Heptafluoropropyl-1,3,4-dioxazolone
[87050-95-1]




Middleton, W. J., *J. Org. Chem.*, 1983, **48**, 3845

It could not be detonated by hot wire initiation, but a sample exploded during distillation at ambient pressure at 102°C.

See 1,3,4-DIOXAZOLONES See other N-O COMPOUNDS

1813. Sodium pentacyanonitrosylferrate(2-)
[14402-89-2]



Sodium nitrite

See Sodium nitrite: Metal cyanides

See related METAL CYANIDES

†1814. Pentacarbonyliron (Iron pentacarbonyl)

[13463-40-6]

(OC)₅Fe

C₅FeO₅

Merck, 1983, 736

Pyrophoric in air.

Acetic acid, Water

Braye, E. H. *et al.*, *Inorg. Synth.*, 1966, **8**, 179

A brown pyrophoric powder is produced if the carbonyl is dissolved in acetic acid containing above 5% of water.

Nitrogen oxide

See Nitrogen oxide: Pentacarbonyliron

Pentaborane(9), Pyrex glass

See Pentaborane(9): Pentacarbonyliron, etc.

Transition metal halides, Zinc

Lawrenson, M. J., private comm., 1970


The preparation of carbonylmetals by treating a transition metal halide either with carbon monoxide and zinc, or with iron pentacarbonyl is well-known and smooth. However, a violent eruptive reaction occurs if a methanolic solution of a cobalt halide, a rhodium halide or a ruthenium halide is treated with both zinc and iron pentacarbonyl.

See other CARBONYLMETALS

1815. 1,3-Pentadiyn-1-ylsilver

[]

C₅H₃Ag



Alone, or Sulfuric acid


Schlubach, H. H. *et al.*, *Ann.*, 1950, **568**, 155

Very sensitive to impact or friction, and explodes when moistened with sulfuric acid.

See other METAL ACETYLIDES, SILVER COMPOUNDS

1816. 2-Chloro-3-pyridinediazonium tetrafluoroborate
[70682-07-4]

$C_5H_3BClF_4N_3$




Verde, C., Span. Pat. ES 548 695, 1986

The diazonium salt decomposes violently above 25°C or when dry.

See other DIAZONIUM TETRAHALOBORATES

1817. 2-Furylchlorodiazirine
[]

$C_5H_3ClN_2$




Khasanova, T. *et al.*, *J. Amer. Chem. Soc.*, 1998, **120**(1), 233

This previously unknown compound is, not surprisingly, detonable.

See other DIAZIRINES

1818. Furoyl chloride
[1300-32-9]

$C_5H_3ClO_2$



Preparative hazard

Wrigley, T. C., private comm., 1979

A sample of freshly distilled material exploded violently during overnight storage. Self-acylation or polymerisation catalysed by hydrogen chloride may have been involved.


See Phosphorus trichloride: Carboxylic acids

See also 2-HALOMETHYL-FURANS OR -THIOPHENES

See other ACYL HALIDES, POLYCONDENSATION REACTION INCIDENTS

1819. 1,3-Pentadiyn-1-ylcopper
[115609-87-5]


C₅H₃Cu



Schlubach, H. H. *et al.*, *Ann.*, 1950, **568**, 155
Explodes on impact or friction.
See other METAL ACETYLIDES

1820. Lithium 3-(1,1,2,2-tetrafluoroethoxy)propynide
[82906-06-7]


C₅H₃F₄LiO



Von Werner, K., *J. Fluorine Chem.*, 1982, **20**, 215–216
The explosion during work-up of a substantial amount of the lithiated fluorocompound emphasises the need for great care and full precautions in handling such materials.
See related HALOACETYLENE DERIVATIVES *See other* METAL ACETYLIDES

1821. 2-Furoyl azide
[20762-98-5]


C₅H₃N₃O₂



Dunlop, 1953, 544
The azide explodes violently on heating in absence of a solvent or diluent.
See other ACYL AZIDES

1822. 2-Hydroxy-3,5-dinitropyridine
[2980-33-8]

$C_5H_3N_3O_5$




Glowiak, B., *Chem. Abs.*, 1963, **58**, 498h

Various heavy metal salts show explosive properties, and the lead salt might be useful as an initiating explosive.

See other POLYNITROARYL COMPOUNDS

1823. 2-Formylamino-3,5-dinitrothiophene
[]

$C_5H_3N_3O_5S$




Fast flame propagation on heating the powder moderately.

See entry HIGH RATE DECOMPOSITION See other POLYNITROARYL COMPOUNDS

1824. Methyl 3,3-diazido-2-cyanoacrylate
[82140-87-2]

$C_5H_3N_7O_2$




Saalfank, R. W. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1987, **26**, 1161 (footnote 14)

The crystalline solid decomposes explosively around 70°C, so working scale was limited to 0.5 g.

See other CYANO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, ORGANIC AZIDES

1825. 1,3-Pentadiyne
[4911-55-1]

C₅H₄




Brandsma, 1971, 7, 36

It explodes on distillation at ambient pressure.

See other ACETYLENIC COMPOUNDS

1826. 3-Pyridinediazonium tetrafluoroborate
[586-92-5]

C₅H₄BF₄N₃



1. Johnson, E. P. *et al.*, *Chem. Eng. News*, 1967, **45**(44), 44


2. Roe, A. *et al.*, *J. Amer. Chem. Soc.*, 1947, **69**, 2443

A sample, air-dried on aluminium foil, exploded spontaneously and another sample exploded on heating to 47°C [1]. The earlier reference describes the instability of the salt above 15°C if freed from solvent, and both the 2- and 4-isomeric salts were found to be very unstable and incapable of isolation [2].

See other DIAZONIUM TETRAHALOBORATES

1827. 3-Bromopyridine
[626-55-1]

C₅H₄BrN




Acetic acid, Hydrogen peroxide

See Hydrogen peroxide: Acetic acid, *N*-Heterocycles

1828. 3-Bromopyridine *N*-oxide
[2402-97-3]

C₅H₄BrNO




Preparative hazard

See Hydrogen peroxide: Acetic acid, *N*-Heterocycles

1829. 2-Chloropyridine *N*-oxide
[2402-95-1]

C₅H₄ClNO




1. Kotoyori, T., private comm., 1983
2. *MARS Database*, 1998, short report 025

A 500 kg batch had been prepared in glacial acetic acid using 35% aqueous hydrogen peroxide with 2% sulfuric acid as catalyst. When the reaction was complete, solvents were removed by vacuum distillation, steam was turned off and the reaction vessel jacket filled with water. After standing overnight, violent decomposition of the contents occurred, causing the safety valve to lift, even although the vessel was vented via its condenser/receiver system. An external fire ensued, and carbonised residue was found inside the reactor. It was surmised that as the product began to solidify inwards from the outer cooled reactor wall, the inner liquid material effectively went into adiabatic storage. The compound is of limited thermal stability and begins to decompose exothermally at 90–100°C, and salt formation from liberated hydrogen chloride will tend to accelerate the process. The hot weather (and possibly a leak of steam into the jacket) may have exaggerated the adiabatic effect. The necessity to assess thermal stabilities of reactants and products, to minimise heating times and to maximise cooling rates, and to monitor and supervise processes is stressed [1]. Another report of a process explosion during manufacture of this, using maleic anhydride as catalyst and without a solvent, was attributed to adding a shot of 70% hydrogen peroxide slightly faster than normal. Since the reaction was almost complete and no product remained in the reactor afterwards, it is possible that the amine oxide was also implicated [2]

See other N-OXIDES

1830. 2-Chloro-5-chloromethylthiophene
[23784-96-5]


$C_5H_4Cl_2S$



Rosenthal, N. A., *J. Amer. Chem. Soc.*, 1951, **73**, 590
Storage at ambient temperature may lead to explosively violent decomposition.
See other 2-HALOMETHYL-FURANS OR -THIOPHENES

1831. 2-Fluoropyridine
[372-48-5]


C_5H_4FN



Bromine trifluoride
See Bromine trifluoride: Pyridine

1832. Diazocyclopentadiene
[1192-27-4]

$C_5H_4N_2$



1. Ramirez, F., *J. Org. Chem.*, 1958, **23**, 2036–2037
2. Wedd, A. G., *Chem. & Ind.*, 1970, 109
3. Aarons, L. J. *et al.*, *J. Chem. Soc., Faraday Trans. 2*, 1974, **70**, 1108
4. DeMore, W. B. *et al.*, *J. Amer. Chem. Soc.*, 1959, **81**, 5875
5. Weil, T., *J. Org. Chem.*, 1963, **28**, 2472
6. Regitz, M. *et al.*, *Tetrahedron*, 1967, **23**, 2706
7. Gillespie, R. J. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1979, 2626
8. Reimer, K. J. *et al.*, *Inorg. Synth.*, 1980, **20**, 189

A violent explosion occurred after distillation at 47–49°C/60–65 mbar [1], and also during distillation at 48–53°C/66 mbar; use of solutions of undistilled material was recommended [2]. The compound exploded on one occasion in the solid state after condensation at –196°C, probably owing to fortuitous tribomechanical shock [3]. A similar explosion on chilling a sample in liquid nitrogen had been noted previously [4]. Of two simplified methods of isolating the material without distillation [4,5],

the former appears to give higher yields. A further explosion during distillation has been reported [7], but solutions in pentane are considered convenient and safe [8].


See other DIAZO COMPOUNDS

1833. 4-Nitropyridine N-oxide

[1124-33-0]

$C_5H_4N_2O_3$

$O_2NC_5H_4N:O$



Diethyl 1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate

Kurbatova, A. S. *et al.*, *Chem. Abs.*, 1978, **88**, 120946


Dehydrogenation of the 'Hantzsch' ester in a melt with the oxide at 130–140°C proceeded explosively.

See other N-OXIDES

1834. 1,2-Dihydropyrido[2,1-e]tetrazole

[]

$C_5H_4N_4$



Fargher, R. G. *et al.*, *J. Chem. Soc.*, 1915, **107**, 695


Explodes on touching with a hot rod.

See other TETRAZOLES

1835. 1,2-Bis(azidocarbonyl)cyclopropane

[[68979-48-6; 114752-52-2] (cis, trans, resp.)]

$C_5H_4N_6O_2$



1. Landgrebe, J. A., *Chem. Eng. News*, 1981, **59**(17), 47

2. Majchrzak, M. W. *et al.*, *Synth. Comm.*, 1981, **11**, 493–495

3. Johnson, R. E., *Chem. Eng. News*, 1988, **66**(13), 2


Trituration of the crude *cis*-azide with hexane at ambient temperature caused detonation to occur. Isolation of low molecular weight carbonyl azides should be avoided, or extreme precautions taken. A similar incident with the analogous cyclobutane diazide was reported [1]. During use as an intermediate in preparation of the 1,2-diamine, the explosive diazide was never isolated or handled free of solvent [2]. The *trans* isomer (0.25 mole) was prepared by the latter technique, but as a 45 w/v% solution in toluene, rather than as the recommended 10% solution. When a magnetic spin-bar was introduced, the solution detonated. Initiation may have involved friction from adventitious presence of iron filings adhering to the magnetic bar [3].

See other ACYL AZIDES, FRICTIONAL INITIATION INCIDENTS

1836. 2-Furaldehyde (Furfural)

[98-01-1]

C₅H₄O₂



HCS 1980, 511

Other reactants

Yoshida, 1980, 313

MRH values calculated for 13 combinations with oxidants are given.

Sodium hydrogen carbonate

Unpublished information, 1977

Several cases of spontaneous ignition after exposure to air of fine coke particles removed from filter strainers on a petroleum refinery furfural extraction unit have been noted. This has been associated with the use of sodium hydrogen carbonate (bicarbonate) injected into the plant for pH control, which produced a pH of 10.5 locally. This would tend to resinify the aldehyde, but there is also the possibility of a Cannizzaro reaction causing conversion of the aldehyde to furfuryl alcohol and furoic acid. The latter, together with other acidic products of autoxidation of the aldehyde, would tend to resinify the furfuryl alcohol. Pyrolysis GLC showed the presence of a significant proportion of furfuryl alcohol-derived resins in the coke. The latter is now discarded into drums of water, immediately after discharge from the strainers, to prevent further incidents.

See Furfuryl alcohol: Acids

See other PYROPHORIC MATERIALS


Sodium hypochlorite

See Sodium hypochlorite: Furfuraldehyde

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

1837. Peroxyfuroic acid
[5797-06-8]

$C_5H_4O_4$



Alone, or Metal salts, or Organic materials

Milas, N. A. *et al.*, *J. Amer. Chem. Soc.*, 1934, **56**, 1221

Thermal decomposition of the acid becomes violently explosive at 40°C. Intimate contact with finely divided metal salts (mainly halides), ergosterol, pyrogallol or animal charcoal led to explosive decomposition (often violent) in 18 out of 28 experiments.

See other PEROXYACIDS

1838. Silver cyclopropylacetylide
[]

C_5H_5Ag




Slobodin, Ya. M. *et al.*, *Chem. Abs.*, 1952, **46**, 10112e

It explodes on heating.

See other METAL ACETYLIDES, SILVER COMPOUNDS

1839. Cyclopentadienylgold(I)
[21254-73-9]

C_5H_5Au




Huttel, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1967, **6**, 862

It is sensitive to friction and heat, often deflagrating on gentle warming.

See other ALKYLMETALS, GOLD COMPOUNDS

1840. 2-Bromomethylfuran
[4437-18-7]

C_5H_5BrO




Dunlop, 1953, 231

It is very unstable, and the liberated hydrogen bromide accelerates further decomposition to explosive violence.

See other 2-HALOMETHYL-FURANS OR -THIOPHENES

1841. 2-Chloromethylfuran
[615-88-9]

C_5H_5ClO




Anon., *ABCM Quart. Safety Summ.*, 1962, **33**, 2

A small sample of freshly prepared and distilled material, when stored over a weekend, exploded violently owing to polymerisation or decomposition (probably arising from the facile hydrolysis with traces of moisture and liberation of hydrogen chloride). Material should be prepared and used immediately, but if brief storage is inevitable, refrigeration is essential.

See other 2-HALOMETHYL-FURANS OR -THIOPHENES

1842. 2-Chloromethylthiophene
[617-88-9]

C_5H_5ClS



1. Bergeim, F. H., *Chem. Eng. News*, 1952, **30**, 2546

2. Meyer, F. C., *Chem. Eng. News*, 1952, **30**, 3352

3. Wiberg, K. B., *Org. Synth.*, 1955, Coll. Vol. 3, 197

The material is unstable and gradually decomposes, even when kept cold and dark, with liberation of hydrogen chloride which accelerates the decomposition [1]. If kept in closed containers, the pressure increase may cause an explosion [2]. Amines stabilise the material, which may then be kept cold in a vented container for several months [3].

See other GAS EVOLUTION INCIDENTS, 2-HALOMETHYL-FURANS OR -THIOPHENES

1843. Poly(cyclopentadienyltitanium dichloride)
[35398-20-0]

$(C_5H_5Cl_2Ti)_n$

(Complex structure)


Lucas, C. R. *et al.*, *Inorg. Synth.*, 1976, **16**, 238

It is pyrophoric in air.

See other ORGANOMETALLICS, PYROPHORIC MATERIALS

1844. Oxodiperoxopyridinechromium *N*-oxide
[38293-27-5]

$C_5H_5CrNO_6$



1. Westland, A. D. *et al.*, *Inorg. Chem.*, 1980, **19**, 2257
2. Daire, E. *et al.*, *Nouv. J. Chim.*, 1984, **8**, 271–274


This chromium complex, and the molybdenum and tungsten analogues with two *N*-oxide ligands, were all explosive [1]. The chromium complex has exploded during desiccation [2].

See Triphenylphosphine oxide-oxodiperoxochromium(VI)

See other AMMINECHROMIUM PEROXOCOMPLEXES

1845. 1-Iodo-3-penten-1-yne
[]

C_5H_5I




Vaughn, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1934, **56**, 1208

After distillation at 72°C/62 mbar, the residue always exploded if heating were continued.

See other HALOACETYLENE DERIVATIVES

1846. Potassium cyclopentadienide
[30994-24-2]

C_5H_5K




Angus, P. L. *et al.*, *Inorg. Synth.*, 1977, **17**, 177

The dry powder is very air-sensitive and pyrophoric.

See other ALKYLMETALS, PYROPHORIC MATERIALS

1847. Potassium 2,5-dinitrocyclopentanonide
[26717-79-3]




Wieland, H. *et al.*, *Ann.*, 1928, **461**, 304

The salt (possibly an *aci*-nitro salt) explodes at 154–158°C and is less stable than the di-potassium salt (241–245°C).

See other C-NITRO COMPOUNDS

†**1848. Pyridine**
[110-86-1]



NSC 310, 1978; *FPA H58*, 1977; *HCS 1980*, 802; *RSC Lab. Hazard Data Sheet No. 8*, 1983

Formamide, Iodine, Sulfur trioxide
See Formamide: Iodine, etc.

Maleic anhydride
See Maleic anhydride: Bases, etc.


Other reactants
Yoshida, 1980, 292
MRH values calculated for 15 combinations with oxidants are given.

Oxidants	MRH values show % of oxidant
<i>See</i> Bromine trifluoride: Pyridine	MRH 3.05/75
Chromium trioxide: Pyridine	MRH 3.05/75
Dinitrogen tetroxide: Heterocyclic bases	MRH 7.07/78
Fluorine: Nitrogenous bases	
Trifluoromethyl hypofluorite: Pyridine	

See other ORGANIC BASES

1849. Pyridine *N*-oxide
[694-59-7]

C₅H₅NO



Energy of decomposition (in range 250–380°C) measured as 1.076 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 205°C by adiabatic Dewar tests, with an apparent energy of activation of 124 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Hexamethyldisilane, Tetrabutylammonium fluoride

Vorbrueggen, H. *et al.*, *Tetrahedron Lett.*, 1983, **24**, 5337

Slow addition of hexamethyldisilane in THF to pyridine *N*-oxide and tetrabutylammonium fluoride in THF effected smooth reduction to pyridine, while addition of undiluted fluoride led to an explosion on two occasions.

See other N-OXIDES

1850. Methylnitrothiophene

[]

(Unstated isomer)

C₅H₅NO₂S


Preparative hazard

See Nitric acid: Methylthiophene

See other C-NITRO COMPOUNDS

1851. 2-(*N*-Nitroamino)pyridine *N*-oxide
[85060-25-9]

C₅H₅N₃O₃




Talik, T. *et al.*, *Chem. Abs.*, 1983, **98**, 160554

Powerfully explosive.


See other N-NITRO COMPOUNDS, N-OXIDES

1852. 4-(*N*-Nitroamino)pyridine *N*-oxide (*N*-Nitro-4-pyridinamine 1-oxide)
[85060-27-1] C₅H₅N₃O₃




Talik, T. *et al.*, *Chem. Abs.*, 1983, **98**, 160554
Powerfully explosive.
See other N-NITRO COMPOUNDS, N-OXIDES

1853. Ethyl 4-diazo-1,2,3-triazole-5-carboxylate
[85807-69-8] C₅H₅N₅O₂



See entry DIAZAZOLES
See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

1854. 1,1,1,3,5,5,5-Heptanitropentane
[20919-99-7] C₅H₅N₇O₁₄



Klager, K. *et al.*, *Propellants, Explos., Pyrotech.*, 1983, **8**, 25–28
This explosive compound has a +12% oxygen balance so can function as an oxidant.
See other OXIDANTS, POLYNITROALKYL COMPOUNDS

1855. Cyclopentadienylsodium
[4984-82-1]

C_5H_5Na



Na^+

King, R. B. *et al.*, *Inorg. Synth.*, 1963, **7**, 101

The solid obtained by evaporation of the air-sensitive solution is pyrophoric in air.

See ALKALI-METAL DERIVATIVES OF HYDROCARBONS

Ammonium hexanitrocerate

See Sodium nitrate: Tetracyclopentadienylcerium

Lead(II) nitrate

See Lead(II) nitrate: Cyclopentadienylsodium

See other ALKYL METALS, ORGANOMETALLICS

1856. Bicyclo[2.1.0]pent-2-ene
[5164-35-2]

C_5H_6



Andrist, A. H. *et al.*, *Org. Synth.*, 1976, **55**, 18

A purified undiluted sample was reported to explode.

See other STRAINED-RING COMPOUNDS

†**1857. Cyclopentadiene**
[542-92-7]

C_5H_6



HCS 1980, 348

1. Kirk-Othmer, 1965, Vol. 6, 690

2. Raistrick, B. *et al.*, *J. Chem. Soc.*, 1939, 1761–1773

3. Starkie, A. *et al.*, *Chem. in Brit.*, 1996, **32**(2), 35

Dimerisation is highly exothermic, the rate increasing rapidly with temperature, and may cause rupture of a closed uncooled container. The monomer may largely be prevented from dimerising by storage at $-80^\circ C$ or below [1]. The polymerisation of the undiluted diene may become explosive within the range $0-40^\circ C$ and at pressures above 2300 bar. The effect of diluents was also studied [2]. A polymerisation was

conducted with insufficient solvent, leading to a runaway and an explosion which killed three workers and destroyed plant up to 300m away [3].

See Propene

See other POLYMERISATION INCIDENTS

Nitric acid

See Nitric acid: Hydrocarbons (reference 9)

Oxides of nitrogen

See Nitrogen oxide: Dienes, Oxygen

Dinitrogen tetroxide: Hydrocarbons (reference 10)

Oxygen

Hock, H. *et al.*, *Chem. Ber.*, 1951, **84**, 349

Exposure of the diene to oxygen gives peroxidic products containing some monomeric, but largely polymeric, peroxides, which explode strongly on contact with a flame.

See other POLYPEROXIDES

Oxygen, Ozone

See Ozone: Dienes, Oxygen

Potassium hydroxide

Wilson, P. J. *et al.*, *Chem. Rev.*, 1944, **34**, 8

Contact with the ethanolic base causes vigorously exothermic resin formation.

See other POLYMERISATION INCIDENTS

Sulfuric acid

See Sulfuric acid: Cyclopentadiene

See other DIENES

†1858. 2-Methyl-1-buten-3-yne

[78-80-8]



C_5H_6

See other ALKYNES

1859. Cyclopentadiene-silver perchlorate

[]




$\text{C}_5\text{H}_6\text{AgClO}_4$

Ulbricht, T. L. V., *Chem. & Ind.*, 1961, 1570

The complex explodes on heating.

See related METAL PERCHLORATES See other SILVER COMPOUNDS

1860. 3,5-Dibromocyclopentene
[1890-04-6]




Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: 3,5-Dibromocyclopentene

See other HALOALKENES


1861. 4,4-Dibromo-3,5-dimethylpyrazole
[84691-20-3]



Adam, W. A. *et al.*, *J. Org. Chem.*, 1994, **59**(23), 7067

This compound decomposed in a few days at -18°C . On warming to ambient it decomposed rapidly, exploding on one occasion.

1862. Pyridinium chlorochromate
[26299-14-9]




1. Glaros, G., *J. Chem. Educ.*, 1978, **55**, 410
2. Singh, J. H. *et al.*, *Chem. & Ind.*, 1986, 751

It is a stable and safe oxidant which can replace oxodiperoxodipyridine-chromium(VI) in oxidation of alcohols to aldehydes [1]. The quinolinium analogue has also found similar application [2].

See Oxodiperoxodipyridinechromium(VI)

See other OXIDANTS, QUATERNARY OXIDANTS

1863. Pyridinium perchlorate
[15598-34-2]




1. Kuhn, R. *et al.*, *Chem. Ztg.*, 1950, **74**, 139
2. Anon., *Sichere Chemiearb.*, 1963, **15**(3), 19
3. Arndt, F. *et al.*, *Chem. Ztg.*, 1950, **74**, 140
4. Schumacher, 1960, 213
5. Zacherl, M. K., *Mikrochemie*, 1948, **33**, 387–388
6. Anon., Private communication, 1991

It can be detonated on impact, but is normally considered a stable intermediate (m.p. 288°C), suitable for purification of pyridine [1]. Occasionally explosions have occurred when the salt was disturbed [2], which have been variously attributed to presence of ethyl perchlorate, ammonium perchlorate, or chlorates. A sample was being dried in a glass vacuum desiccator. Over-enthusiastic release of vacuum distributed some of the salt onto the lid. The grinding action as the lid was slid off initiated explosion of this, shattering the lid [6]. A safer preparative modification is described [3]. It explodes on heating to above 335°C, or at a lower temperature if ammonium perchlorate is present [4]. A violent explosion which occurred during the final distillation according to the preferred method [3] was recorded [5].

See other PERCHLORATE SALTS OF NITROGENOUS BASES


1864. 5-Chloro-1,3-dimethyl-4-nitro-1H-pyrazole
[13551-73-0]



See NITRATION INCIDENTS (reference 8) See other C-NITRO COMPOUNDS

1865. 1,3-Dichloro-5,5-dimethyl-2,4-imidazolidindione
[118-52-5]

$C_5H_6Cl_2N_2O_2$



Xylene


1. Anon., *Chem. Trade J.*, 1951, **129**, 136
2. *See entry* SELF-ACCELERATING REACTIONS

An attempt to chlorinate xylene with the 'dichlorohydantoin' caused a violent explosion [1]. The haloimide undergoes immediate self accelerating decomposition in presence of solvents. Safe conditions (including lower temperatures and progressive addition of reagent to match its consumption) can be developed for its use [2].

See other N-HALOIMIDES

1866. 3,3-Difluoro-1,2-dimethoxycyclopropene
[59034-33-2]


$C_5H_6F_2O_2$



See entry FLUORINATED CYCLOPROPENYL METHYL ETHERS

1867. 4,4-Bis(difluoroamino)-3-fluoroimino-1-pentene
[33364-51-1]

$C_5H_6F_5N_3$




Parker, C. O. *et al.*, *J. Org. Chem.*, 1972, **37**, 922

Samples of this and related poly-difluoroamino compounds exploded during analytical combustion.

See other DIFLUOROAMINO COMPOUNDS, N-HALOGEN COMPOUNDS

1868. 4-Iodo-3,5-dimethylisoxazole
[10557-85-4]

C₅H₆INO




Peroxytrifluoroacetic acid

See Peroxytrifluoroacetic acid: 4-Iodo-3,5-dimethylisoxazole

See other N-O COMPOUNDS, IODINE COMPOUNDS

1869. 1,3-Diisocyanopropane
[72399-85-0]


C₅H₆N₂



See entry DIISOCYANIDE LIGANDS

1870. Glutarodinitrile
[544-13-8]

C₅H₆N₂




Energy of decomposition (in range 200–340°C) measured as 0.098 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other CYANO COMPOUNDS

1871. 4,5-Cyclopentanofurazan-N-oxide (Trimethylenefuroxan)
[54573-23-8]

C₅H₆N₂O₂




Barnes, J. F. *et al.*, *J. Chem. Res. (S)*, 1979, 314–315

It decomposes explosively on attempted distillation at 150°C at ambient pressure.

See other FURAZAN N-OXIDES

1872. Furan-2-amidoxime
[50892-99-4]




Pearse, G. A., *Chem. Brit.*, 1984, **20**, 30

Attempted vacuum distillation of the oxime at a pot temperature of 100°C led to an explosion as distillation began. When heated at ambient pressure to above 65°C, a rapid exotherm occurs with gas evolution.

See other GAS EVOLUTION INCIDENTS, OXIMES

1873. Pyridinium nitrate
[543-53-3]




Goldstein, N. *et al.*, *J. Chem. Phys.*, 1979, **70**, 5073

Like the perchlorate salt, pyridinium nitrate explodes on heating, but not with shock or friction.

See Pyridinium perchlorate

See other OXOSALTS OF NITROGENOUS BASES

1874. Glutaryl diazide
[64624-44-8]




Curtiss, T. *et al.*, *J. Prakt. Chem.*, 1900, **62**, 196

A very small sample exploded sharply on heating.

See other ACYL AZIDES

1875. *N,N'*-Bis(2,2,2-trinitroethyl)urea
[918-99-0]

C₅H₆N₈O₁₃



Sodium hydroxide

Biasutti, G. S., private comm., 1981


A pilot plant for manufacture of the title compound (an explosive of zero oxygen balance) was decommissioned and treated with concentrated alkali, but not completely drained. After 4 years, an explosion occurred, attributed to the hydrolytic formation of trinitroethanol or its sodium salt(s).

See 2,2,2-Trinitroethanol

See other POLY-NITROALKYL COMPOUNDS

1876. 2,4-Diazido-6-dimethylamino-1,3,5-triazine
[83297-43-2]

C₅H₆N₁₀




Preparative hazard

See 2,4,6-Triazido-1,3,5-triazine

See other ORGANIC AZIDES

1877. Divinyl ketone (1,4-Pentadien-3-one)
[890-28-4]

C₅H₆O




Trinitromethane

See Trinitromethane: Divinyl ketone


†1878. 2-Methylfuran
[534-22-5]

C₅H₆O



1879. 2-Penten-4-yn-3-ol
[101672-20-2]

C₅H₆O




Brandsma, 1971, 73

The residue from distillation at 20 mbar exploded vigorously at above 90°C.

See other ACETYLENIC COMPOUNDS

1880. Furfuryl alcohol (2-Furanemethanol)
[98-00-0]

C₅H₆O₂



Acids

1. Tobie, W. C., *Chem. Eng. News*, 1940, **18**, 72
2. *MCA Case History No. 858*
3. Dunlop, 1953, 214, 221, 783
4. Golosova, L. V., *Chem. Abs.*, 1975, **83**, 194098

A mixture of the alcohol with formic acid rapidly self-heated, then reacted violently [1]. A stirred mixture with cyanoacetic acid exploded violently after application of heat [2]. Contact with acids causes self-condensation of the alcohol, which may be explosively violent under unsuitable physical conditions. The general mechanism has been discussed [3]. The explosion hazards associated with the use of acidic catalysts to polymerise furfuryl alcohol may be avoided by using as catalyst the condensation product of 1,3-phenylenediamine and 1-chloro-2,3-epoxypropane [4]. See Nitric acid: Alcohols (reference 6)

See other POLYMERISATION INCIDENTS

Hydrogen peroxide


See Hydrogen peroxide: Alcohols

Sulfur tetrafluoride, Triethylamine

See Sulfur tetrafluoride: 2-(Hydroxymethyl)furan, Triethylamine

1881. Methyl 2-butynoate
[23326-27-4]


C₅H₆O₂



Cochran, J., *Chem. Eng. News* 1994, **72**(46), 4
Attempts to manufacture above the gram scale by several different routes have often resulted in violent decomposition during vacuum distillation. Distillation of more than a few grams demands caution.
See other ACETYLENIC COMPOUNDS

†**1882. 2-Methylthiophene**
[554-14-3]


C₅H₆S



Nitric acid
See Nitric acid: Methylthiophene

1883. 2-Propynyl vinyl sulfide
[21916-66-5]


C₅H₆S



Brandsma, 1971, 7, 182
It decomposes explosively above 85°C.
See other ACETYLENIC COMPOUNDS

1884. 4-Bromocyclopentene
[1781-66-4]


C₅H₇Br



Preparative hazard
See Lithium tetrahydroaluminate: 3,5-Dibromocyclopentene
See other HALOALKENES

1885. 3-Chlorocyclopentene
[96-40-2]

C₅H₇Cl




MCA Guide, 1972, 305

A 35 g portion decomposed (or polymerised) explosively after storage for 1 day.

See other HALOALKENES

1886. Ethyl 2,2,3-trifluoropropionate
[28781-86-4]

C₅H₇F₃O₂




Sodium hydride

See Sodium hydride: Ethyl 2,2,3-trifluoropropionate


†**1887. 1-Methylpyrrole**
[96-54-8]

C₅H₇N



1888. 3,5-Dimethylisoxazole
[300-87-8]

C₅H₇NO




Sealed samples decompose exothermally above 210°C.

See entry ISOXAZOLES *See other* N-O COMPOUNDS

1889. Ethyl cyanoacetate
[105-56-6]

C₅H₇NO₂




Preparative hazard

See Ethyl chloroacetate: Sodium cyanide

See other CYANO COMPOUNDS

1890. Ethyl 2-azido-2-propenoate
[81852-50-8]

C₅H₇N₃O₂



Kakimoto, M. *et al.*, *Chem. Lett.*, 1982, 525–526

Attempted distillation caused a severe explosion; such materials should be purified without heating.

See other 2-AZIDOCARBONYL COMPOUNDS

†**1891. Cyclopentene**
[142-29-0]

C₅H₈




HCS 1980, 350

See other ALKENES

†**1892. 3-Methyl-1,2-butadiene**
[598-25-4]

C₅H₈



It is rather endothermic (ΔH_f° (g) +129.7 kJ/mol, 1.90 kJ/g).

See other DIENES, ENDOTHERMIC COMPOUNDS

†1893. 2-Methyl-1,3-butadiene (Isoprene)

[78-79-5]

C₅H₈



FPA H112, 1982; *HCS 1980*, 573

Kirk-Othmer, 1967, Vol. 12, 79

It is somewhat endothermic, ΔH_f° (g) +75.7 kJ/mol, 1.11 kJ/g. In absence of inhibitors, isoprene absorbs atmospheric oxygen to form peroxides which do not separate from solution. Although the solution is not detonable, the gummy peroxidic polymer obtained by evaporation can be detonated by impact under standard conditions.

See 1,3-Butadiene (reference 2)

Acetone

Lokhmacheva, I. K. *et al.*, *Chem. Abs.*, 1975, **82**, 63856

Prevention of peroxidation of isoprene-acetone mixtures, and other hazards involved in the industrial preparation of synthetic citral, are discussed.

Other reactants

Yoshida, 1980, 40

MRH values calculated for 14 combinations with oxidants are given.

Oxygen, Ozone

See Ozone: Dienes, Oxygen

Ozone

Loveland, J. W., *Chem. Eng. News*, 1956, **34**(3), 292

Isoprene (1 g) dissolved in heptane was ozonised at -78°C . Soon after cooling was stopped, a violent explosion, followed by a lighter one, occurred. This was attributed to high concentrations of peroxides and ozonides building up at the rather low temperature employed. Operation at a higher temperature would permit the ozonides and peroxides to decompose, so avoiding high concentrations in the reaction mixture.

See other OZONIDES

Vinylamine

See Vinylamine: Isoprene

See other DIENES, ENDOTHERMIC COMPOUNDS

†1894. 3-Methyl-1-butyne

[598-23-2]


C₅H₈



It is considerably endothermic, ΔH_f° (g) +136.4 kJ/mol, 2.0 kJ/g.

See other ALKYNES, ENDOTHERMIC COMPOUNDS

†1895. 1,2-Pentadiene
[591-95-7]



C_5H_8

It is considerably endothermic, $\Delta H_f^\circ (\text{g}) +145.6 \text{ kJ/mol}$, 2.14 kJ/g.
See other DIENES, ENDOTHERMIC COMPOUNDS

†1896. 1,3-Pentadiene
[504-60-9]



C_5H_8

It is somewhat endothermic, $\Delta H_f^\circ (\text{g}) +77.8 \text{ kJ/mol}$, 1.14 kJ/g.
See other DIENES, ENDOTHERMIC COMPOUNDS

†1897. 1,4-Pentadiene
[591-93-5]



C_5H_8

It is somewhat endothermic, $\Delta H_f^\circ (\text{g}) +105.4 \text{ kJ/mol}$, 1.55 kJ/g.
See other DIENES, ENDOTHERMIC COMPOUNDS

†1898. 1-Pentyne
[627-19-0]



C_5H_8

It is somewhat endothermic, $\Delta H_f^\circ (\text{g}) +144.3 \text{ kJ/mol}$, 2.12 kJ/g.
See other ALKYNES, ENDOTHERMIC COMPOUNDS

†1899. 2-Pentyne
[627-21-4]



C_5H_8

It is somewhat endothermic, $\Delta H_f^\circ (\text{g}) +128.9 \text{ kJ/mol}$, 1.89 kJ/g.


Mercury, Silver perchlorate

See Silver perchlorate: Alkynes, etc.

See other ALKYNES, ENDOTHERMIC COMPOUNDS

1900. Tetrakis(*N,N*-dichloroaminomethyl)methane
[]

$C_5H_8Cl_8N_4$




Lévy, R. S. *et al.*, *Mém. Poudres*, 1958, **40**, 109

Shock- and heat-sensitive, it is a brisant more powerful than mercury fulminate.

See other N-HALOGEN COMPOUNDS

1901. 2,2-Bis[(nitrooxy)methyl]propane-1,3-diol dinitrate
[78-11-5]

$C_5H_8N_4O_{12}$




This heart drug (−14.3% oxygen balance) enjoys larger scale use as the explosive PETN (PentaErythritolTetraNitrate).

See other ALKYL NITRATES

1902. 3,3-Bis(azidomethyl)oxetane
[17607-20-4]

$C_5H_8N_6O$




Frankel, M. B. *et al.*, *J. Chem. Eng. Data*, 1981, **26**, 219

Explosive, of considerable sensitivity.

See other ORGANIC AZIDES

1903. Ethyl 2,3-diazidopropionate
[85590-62-1]

C₅H₈N₆O₂




Owston, P. G. *et al.*, *J. Chem. Res., Synop.*, 1985, 352–353

It is unstable and potentially explosive: routine use of a fume cupboard and safety screens is recommended.

See other 2-AZIDOCARBONYL COMPOUNDS, ORGANIC AZIDES

†1904. Allyl vinyl ether (3-Ethenoxypropene)
[3917-15-5]


C₅H₈O



See other ALLYL COMPOUNDS

1905. Cyclopentanone
[120-92-3]

C₅H₈O




Hydrogen peroxide, Nitric acid

See Hydrogen peroxide: Ketones, etc.

†1906. Cyclopentene oxide (6-Oxabicyclo[3.1.0]hexane)
[285-67-6]


C₅H₈O



See other 1,2-EPOXIDES


†1907. Cyclopropyl methyl ketone
[765-43-5]

C₅H₈O



†1908. 2,3-Dihydropyran
[110-87-2]


C₅H₈O



HCS 1980, 410

1909. 1-Ethoxy-2-propyne
[628-33-1]

C₅H₈O



Brandsma, 1971, 172

Distillation of a 1 kg quantity at 80°C/1 bar led to a violent explosion. As the compound had not been stored under nitrogen during the 3 weeks since preparation, peroxide formation was suspected.

See other ACETYLENIC COMPOUNDS, PEROXIDISABLE COMPOUNDS

†1910. 2-Methyl-3-butyn-2-ol
[115-19-5]

C₅H₈O



Sulfur tetrafluoride

See Sulfur tetrafluoride: 2-Methyl-3-butyn-2-ol

See other ACETYLENIC COMPOUNDS

†1911. Methyl isopropenyl ketone (2-Methylbuten-3-one)
[814-78-8]

C₅H₈O



†1912. Allyl acetate
[591-87-7]

C₅H₈O₂



Energy of decomposition (in range 170–470°C) measured as 0.45 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other ALLYL COMPOUNDS

†1913. Ethyl acrylate (Ethyl propenoate)
[140-88-5]

C₅H₈O₂



FPA H73, 1978; *HCS 1980*, 464

MCA Case History No. 1759

Inhibited monomer was transferred from a steel drum into a 4 l clear glass bottle exposed to sunlight in a laboratory in which the ambient temperature was temporarily higher than usual. Exothermic polymerisation set in and caused the bottle to burst. Precautions recommended included increase in inhibitor concentration tenfold (to 200 ppm) for laboratory-stored samples, and use of metal or brown glass containers.
See other POLYMERISATION INCIDENTS

†1914. Methyl crotonate (Methyl 2-butenate)
[623-43-8]

C₅H₈O₂



†1915. Methyl methacrylate (Methyl 2-methylpropenoate)
[80-62-6]

C₅H₈O₂



FPA H62, 1977; *HCS 1980*, 656

1. Harmon, 1974, 2.13
2. Barnes, C. E. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 210
3. Bond, J., *Loss Prev. Bull.*, 1985, (065), 26
4. Huschke, D., *Chem. Processing* (Chicago), 1988, **51**(1), 136–137

The monomer tends to self polymerise and this may become explosive. It must be stored inhibited [1]. Exposure of the pure (unstabilised) monomer to air at ambient temperature for 2 months generated an ester–oxygen interpolymers, which exploded on evaporation of the surplus monomer at 60°C (but not at 40°C) [2]. Polymerisation, probably initiated by rust, in a drum of the monomer led to development of overpressure which sheared off the base and propelled the drum into the roof of the building [3]. Some oxygen must be present in the nitrogen used to inert storage tanks, to activate the stabiliser and prevent gelling of the monomer. Use of an oxygen-selective permeable membrane to produce enriched nitrogen from compressed air for this purpose is described [4].

See Initiators, below;

See also VIOLENT POLYMERISATION

See other POLYPEROXIDES

Dibenzoyl peroxide

See Dibenzoyl peroxide: Methyl methacrylate

Initiators

1. Schulz, G. V. *et al.*, *Z. Elektrochem.*, 1941, **47**, 749–761

2. Biesenberg, J. S. *et al.*, *J. Polym. Eng. Sci.*, 1976, **16**, 101–116

Polymerisation of methyl methacrylate initiated by oxygen or peroxides proceeds with a steady increase in velocity during a variable induction period, at the end of which a violent 90°C exotherm occurs. This was attributed to an increase in chain branching, and not to a decrease in heat transfer arising from the increasing viscosity [1]. The parameters were determined in a batch reactor for thermal runaway polymerisation of methyl methacrylate, initiated by azoisobutyronitrile, dibenzoyl peroxide or di-*tert*-butyl peroxide [2].

Other reactants

Yoshida, 1980, 359

MRH values calculated for 13 combinations with oxidants are given.

Propionaldehyde

CISHC Chem. Safety Summ., 1983, **54**(216), 498

Following an incident in which a drum containing bulked drainings (from other drums awaiting reconditioning) fumed and later exploded after sealing, it was found that methyl methacrylate and propionaldehyde can, under certain conditions of mixing, lead to a rapid exothermic reaction. Precautions are discussed.

See other POLYMERISATION INCIDENTS

†1916. Isopropenyl acetate


[108-72-5]



$\text{C}_5\text{H}_8\text{O}_2$

†1917. Methyl cyclopropanecarboxylate

[2868-37-3]



$\text{C}_5\text{H}_8\text{O}_2$

See other STRAINED-RING COMPOUNDS

†1918. Vinyl propionate

[105-38-4]




$\text{C}_5\text{H}_8\text{O}_2$

1919. Poly(methyl methacrylate peroxide)

[]

$(C_5H_8O_4)_n$




See Methyl methacrylate (reference 2) also

See POLYPEROXIDES

1920. Isoprene diozonide (3-(3-Methyl-1,2,4-trioxolan-3-yl)-1,2,4-trioxolane)

[[118112-41-7] [118112-40-6] (diastereoisomers)]

$C_5H_8O_6$




See 2-Methyl-1,3-butadiene: Ozone

See other OZONIDES

1921. 1-Methylsilacyclopenta-2,4-diene (1-Methylsilole)

[73132-51-5]

C_5H_8Si




Dienophiles

Barton, T. J., *J. Organomet. Chem.*, 1979, **179**, C19

Thermal cracking at 150°C of the dimer (mixed isomers) of the silole in presence of reactive dienophiles (maleic anhydride, tetracyanoethylene or dimethyl acetylenedicarboxylate) inevitably produced violent explosions arising from exothermic Diels-Alder reactions.

1922. 2,2,2-Tris(bromomethyl)ethanol
[1522-92-5]

C₅H₉Br₃O




Ethyl acetoacetate, Zinc

See Ethyl acetoacetate: 2,2,2-Tris(bromomethyl)ethanol, Zinc

†1923. Chlorocyclopentane
[930-28-9]


C₅H₉Cl



See other HALOALKANES

1924. N-(Chlorocarbonyloxy)trimethylurea
[52716-12-8]

C₅H₉ClN₂O₃



Groebner, P. *et al.*, *Euro. J. Med. Chem., Chim. Ther.*, 1974, **9**, 32–34

A small sample decomposed explosively during vacuum distillation.

See other ACYL HALIDES, N–O COMPOUNDS

†1925. Pivaloyl chloride (Trimethylacetyl chloride)
[3282-30-2]


C₅H₉ClO

Me₃CCO.Cl

See other ACYL HALIDES

1926. *tert*-Butyl chloroperoxyformate
[56139-33-4]

C₅H₉ClO₃




1. Bartlett, P. D. *et al.*, *J. Amer. Chem. Soc.*, 1963, **85**, 1858
2. Hanson, P., *Chem. Brit.*, 1975, **11**, 418

The peroxyester is stable in storage at -25°C , and samples did not explode on friction, impact or heating. However, a 10 g sample stored at ambient temperature heated spontaneously, exploded and ignited [1]. A sample stored at -30°C exploded during manipulation at $0-5^{\circ}\text{C}$ [2].

See other ACYL HALIDES, PEROXYESTERS

1927. 5-Trichloromethyl-1-trimethylsilyltetrazole
[72385-44-5]

C₅H₉Cl₃N₄Si




Lazukina, L. A. *et al.*, *J. Org. Chem. (USSR)*, 1980, **15**, 2009

It explodes at $80-90^{\circ}\text{C}$.

See other TETRAZOLES

1928. 2(2-Iodoethyl)-1,3-dioxolane
[83665-55-8]

C₅H₉IO₂



Stowell, J. C. *et al.*, *J. Org. Chem.*, 1983, **48**, 5382

Chromatographic purification on alumina is recommended because the compound decomposes with some violence on distillation at $55^{\circ}\text{C}/1.6$ mbar.

See other IODINE COMPOUNDS

†1929. Pivalonitrile (Trimethylacetoneitrile)
[630-18-2]


C₅H₉N

Me₃CCN

See other CYANO COMPOUNDS

†1930. 1,2,3,6-Tetrahydropyridine
[694-05-3]


C₅H₉N



See other ORGANIC BASES

†1931. Butyl isocyanate
[111-36-4]

C₅H₉NO




Energy of decomposition (in range 160–450°C) measured as 0.55 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ORGANIC ISOCYANATES

1932. Cyclopentanone oxime
[1192-28-5]

C₅H₉NO



Sulfuric acid

1. Brookes, F. R., private comm., 1968


2. Peacock, C. J., private comm., 1978

Heating the oxime with 85% sulfuric acid to prepare 2-piperidone caused eruption of the stirred flask contents. Benzenesulfonyl chloride in alkali is a less vigorous reagent [1]. A similar reaction using 70% acid and methanol solvent proceeded uneventfully until vacuum distillation to remove volatiles had been completed at 90°C (bath)/27 mbar when the dark residue exploded [2].

See other OXIMES

1933. 2-Ethylacryldehyde oxime (2-Methylenebutanal oxime)
[99705-27-8]

C₅H₉NO




Marvel, C. S. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 5408

An explosion during distillation was recorded, possibly attributable to peroxide formation (or to Beckmann rearrangement).

See other OXIMES, PEROXIDISABLE COMPOUNDS

1934. 2-Piperidone
[675-20-7]

C_5H_9NO




Preparative hazard

See Cyclopentanone oxime: Sulfuric acid

1935. Pivaloyl azide (Trimethylacetyl azide)
[4981-48-0]

$C_5H_9N_3O$




Bühler, A. *et al.*, *Helv. Chim. Acta*, 1943, **26**, 2123

The azide exploded very violently on warming, and on one occasion, on standing.

See other ACYL AZIDES

1936. *tert*-Butyl azidoformate
[1070-19-5]

$C_5H_9N_3O_2$



1. Carpino, L. A. *et al.*, *Org. Synth.*, 1964, **44**, 15
2. Sakai, K. *et al.*, *J. Org. Chem.*, 1971, **36**, 2387
3. Insalaco, M. A. *et al.*, *Org. Synth.*, 1970, **50**, 12
4. Yajima, H. *et al.*, *Chem. Pharm. Bull.*, 1970, **18**, 850–851


5. Fenlon, W. J., *Chem. Eng. News*, 1976, **54**(22), 3
6. Koppel, H. C., *Chem. Eng. News*, 1976, **54**(39), 5
7. Aldrich advertisement, *J. Chem. Soc., Chem. Comm.*, 1976, (23), i
8. Anon., *Org. Synth.*, 1977, **57**, 122
9. Aldrich advertisement, *J. Org. Chem.*, 1979, **44**(3), cover iv
10. Feyen, P., *Angew. Chem. (Intern. Ed.)*, 1977, **16**, 115

Explosion during distillation at 74°C/92 mbar had been recorded on only one out of several hundred occasions [1], but an alternative procedure avoiding distillation has been described [2]. Although distillation at 56–71°C/53 mbar has been accomplished on many occasions, the structure of the compound suggests that it should be considered as potentially explosive [3]. It is important in the preparation to eliminate all traces of phosgene before adding sodium azide to avoid formation of the dangerously explosive carbonyl azide [4]. The undiluted compound is a shock-sensitive explosive, which is thermally unstable between 100 and 135°C and autoignites at 143°C [5]. A safer reagent for *tert*-butoxycarbonylation of amino groups was then made available [6,7], and other safer alternatives are also available [8], including di-*tert*-butyl dicarbonate [9]. An explosion during evaporation of an ethereal solution of the azide at 50°C has also been reported [10].

See other ACYL AZIDES

1937. 1,1,1-Tris(azidomethyl)ethane
[31044-86-7]

C₅H₉N₉



Hydrogen, Palladium catalyst.

Zompa, L. J. *et al.*, *Org. Prep. Proced. Int.*, 1974, **6**, 103–106


There is an explosion hazard during the palladium-catalysed hydrogenation of the tris-azide in ethanol at 2 bar to the tris-amine.

See other CATALYTIC NITRO REDUCTION PROCESSES, HYDROGENATION INCIDENTS, ORGANIC AZIDES

1938. Dimethyl-1-propynylthallium

[]

C₅H₉Tl



Nast, R. *et al.*, *J. Organomet. Chem.*, 1966, **6**, 461

It explodes on heating and is sensitive to stirring or impact.

See other ALKYL METALS, METAL ACETYLIDES

†1939. Cyclopentane

[287-92-3]

C₅H₁₀



HCS 1980, 349

Other reactants

Yoshida, 1980, 154

MRH values calculated for 13 combinations with oxidants are given.

†1940. 1,1-Dimethylcyclopropane

[1630-94-0]

C₅H₁₀



†1941. Ethylcyclopropane

[1191-96-4]


C₅H₁₀



†1942. 2-Methyl-1-butene

[563-46-2]


C₅H₁₀



†1943. 2-Methyl-2-butene

[513-35-9]

C₅H₁₀




†1944. 3-Methyl-1-butene
[563-45-1]




C_5H_{10}

†1945. Methylcyclobutane
[598-61-8]



C_5H_{10}


†1946. 1-Pentene
[109-76-1]



C_5H_{10}


HCS 1980, 718

†1947. 2-Pentene
[646-04-8]



C_5H_{10}

1948. *N*-Chloropiperidine
[2156-71-0]




$\text{C}_5\text{H}_{10}\text{ClN}$

Claxton, G. P. *et al.*, *Org. Synth.*, 1977, **56**, 120

To avoid rapid spontaneous decomposition of the compound, solvent ether must only be partially distilled out of the extracted product, and from a water bath maintained at below 60°C.

See other N-HALOGEN COMPOUNDS

1949. *N*-Perchlorylpiperidine
[768-34-3]



$\text{C}_5\text{H}_{10}\text{ClNO}_3$

Alone, or Piperidine


Gardner, D. L. *et al.*, *J. Org. Chem.*, 1964, **29**, 3738–3739

It is a dangerously sensitive oil which has exploded on storage, heating or contact with piperidine. Absorption in alumina was necessary to desensitise it to allow non-explosive analytical combustion.

See other PERCHLORYL COMPOUNDS

1950. Dichloromethylenediethylammonium chloride
[33842-02-3]

C₅H₁₀Cl₃N



1. Bader, A. R., private comm., 1971

2. Viehe, H. G. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1971, **10**, 573

The compound does not explode on heating [1], the published statement [2] is in error.

1951. N-(2-Cyanoethyl)ethylhydroxylamine
[]

C₅H₁₀N₂O

EtN(OH)CH₂CH₂CN

Krueger H., *Chem. Abs.*, 1998, **128**, 92406z

Although this compound would undoubtedly be unstable, it here appears to be a misprint for the isomeric ethanolamine entered directly below.

See 3-(2-Hydroxyethylamino)propionitrile

1952. 3-(2-Hydroxyethylamino)propionitrile (N-(2-Cyanoethyl)ethanolamine)
[33759-44-3]


C₅H₁₀N₂O

HOCH₂CH₂NHCH₂CH₂CN

Krueger H., *Prax. Sicherheitstech.*, 1997, **4**, 83


A storage tank containing this compound self-heated, with evolution of ammonia, then burst with generation of polymer. The reaction appears to be autocatalytic generation of 2-vinyloxazoline, which then polymerises. Thermal analysis showed it unsafe for storage at 50°C, although storable at room temperature. Other homologues became increasingly stable as the distance between hydroxyl and amino groups increases.

See other CYANO COMPOUNDS

1953. 2,4-Dinitropentane isomers[109745-04-2] [109745-05-3] (**R*,S*** and **R*,R*** (\pm), resp.)**C₅H₁₀N₂O₄**Wade, P. A. *et al.*, *J. Amer. Chem. Soc.*, 1987, **109**, 5052–5055

A grossly impure sample of the (\pm) mixed isomers decomposed vigorously when distilled at 130°C. During steam distillation of the **R*,S*** isomer, overheating of the still pot caused vigorous decomposition: destruction of the pot residues with nitric acid was also vigorous for both.

See other POLYNITROALKYL COMPOUNDS

1954. 3,7-Dinitroso-1,3,5,7-tetraazabicyclo[3.3.1]nonane
[101-25-7]**C₅H₁₀N₆O₂**1. *MCA Case History No. 841*

2. See entry HIGH RATE DECOMPOSITION

3. Hancyk, B. *et al.*, *Chem. Abs.*, 1988, **109**, 212215

A cardboard drum of the blowing agent ‘dinitrosopentamethylenetetramine’ (as a 40% dispersion in fine silica) ignited when roughly handled in storage [1]. It exhibits fast flame propagation on moderate heating, either neat or mixed with 20% chalk [2]. Mixtures with mineral dust (1:4) eliminated neither fire nor explosion hazards, and mixtures with liquid wax (1:4) eliminated explosion hazard but increased the flammability hazard. The neat blowing agent is highly sensitive to explosion if impacted or dropped. Thermal analysis showed that presence of desensitising agents had no effect on thermal stability and course of decomposition of the title compound, which at above 180°C undergoes self-accelerating decomposition [3]. The recently calculated value of 78°C for the critical ignition temperature is in close agreement with the previous value of 75°C.

See entry CRITICAL IGNITION TEMPERATURE

Other reactants

Yoshida, 1980, 158

MRH values calculated for 11 mixtures with other materials (usually present in catalytic proportions) are given.

See other BLOWING AGENTS, NITROSO COMPOUNDS

†1955. Allyl ethyl ether (3-Ethoxypropene)
[557-31-3] C₅H₁₀O



HCS 1980, 127

Anon., *ABCM Quart. Safety Summ.*, 1963, **34**, 7

A commercial sample was distilled without being tested for peroxides and exploded towards the end of distillation. It was later shown to contain peroxides.

See other PEROXIDISABLE COMPOUNDS

†1956. Ethyl propenyl ether (1-Ethoxypropene)
[928-55-2] C₅H₁₀O



See other PEROXIDISABLE COMPOUNDS

†1957. Isopropyl vinyl ether
[926-65-8] C₅H₁₀O



See other PEROXIDISABLE COMPOUNDS

†1958. Isovaleraldehyde
[590-86-3] C₅H₁₀O



See other ALDEHYDES, PEROXIDISABLE COMPOUNDS


†1959. 3-Methyl-2-butanone
[563-80-4] C₅H₁₀O



†1960. 2-Methyl-3-buten-2-ol
[115-18-4] C₅H₁₀O



†1961. 2-Methyltetrahydrofuran
[96-47-9] C₅H₁₀O



See other PEROXIDISABLE COMPOUNDS

†1962. 2-Pentanone
[107-87-9]

MeCO.Pr

C₅H₁₀O

FPA H110, 1982; *HCS 1980*, 661

Bromine trifluoride
See Bromine trifluoride: 2-Pentanone

Other reactants
Yoshida 1980, 369
MRH values calculated for 13 combinations with oxidants are given.

†1963. 3-Pentanone
[96-22-0]

Et₂C:O

C₅H₁₀O

HCS 1980, 401

Hydrogen peroxide
MRH 6.44/84

MRH ,

Nitric acid
See Hydrogen peroxide: Ketones, etc.

MRH 5.82/79

Other reactants
Yoshida, 1980, 338
MRH values calculated for 13 combinations with oxidants are given.


†1964. 4-Penten-1-ol
[821-09-0]

H₂C=CH[CH₂]₃OH

C₅H₁₀O

†1965. Tetrahydropyran
[142-68-7]

C₅H₁₀O



See other PEROXIDISABLE COMPOUNDS

†1966. Valeraldehyde (Pentanal)
[110-62-3] **BuCO.H** **C₅H₁₀O**

See ALDEHYDES


†1967. Butyl formate
[592-84-7] **BuOCO.H** **C₅H₁₀O₂**

HCS 1980, 247

†1968. 3,3-Dimethoxypropene
[6044-68-4] **H₂C=CHCH(OMe)₂** **C₅H₁₀O₂**

See other ALLYL COMPOUNDS, PEROXIDISABLE COMPOUNDS

†1969. 2,2-Dimethyl-1,3-dioxolane
[2916-31-6] **C₅H₁₀O₂**



†1970. Ethyl propionate (Ethyl propanoate)
[105-37-3] **EtOCO.Et** **C₅H₁₀O₂**

HCS 1980, 499

†1971. Isobutyl formate
[542-55-2] **Me₂CHCH₂OCO.H** **C₅H₁₀O₂**

†1972. Isopropyl acetate
[108-21-4] **Me₂CHOCO.Me** **C₅H₁₀O₂**

†1973. 2-Methoxyethyl vinyl ether
[1663-35-0]



$\text{C}_5\text{H}_{10}\text{O}_2$


See other PEROXIDISABLE COMPOUNDS

†1974. Methyl butyrate
[623-42-7]



$\text{C}_5\text{H}_{10}\text{O}_2$

†1975. 4-Methyl-1,3-dioxane
[1120-97-4]



$\text{C}_5\text{H}_{10}\text{O}_2$

†1976. Methyl isobutyrate
[547-63-7]



$\text{C}_5\text{H}_{10}\text{O}_2$


†1977. Propyl acetate
[109-60-4]



$\text{C}_5\text{H}_{10}\text{O}_2$

HCS 1980, 795

1978. Tetrahydrofurfuryl alcohol
[97-99-4]



$\text{C}_5\text{H}_{10}\text{O}_2$

HCS 1980, 892

3-Nitro-*N*-bromophthalimide
See N-HALOIMIDES: alcohols

Other reactants


Yoshida, 1980, 239

MRH values calculated for 13 combinations with oxidants are given.

1979. Trimethyl-1,2-dioxetane

[22688-10-6]

C₅H₁₀O₂




See 3,3-Dimethyl-1,2-dioxetane

See other DIOXETANES, STRAINED-RING COMPOUNDS; CYCLIC PEROXIDES

1980. Tellurane-1,1-dioxide

[]

C₅H₁₀O₂Te



Alone, or Acids

Morgan, G. T. *et al.*, *J. Chem. Soc.*, 1928, 321

It exploded on rapid heating, and decomposed violently in contact with nitric or sulfuric acids.

See related ALKYL METALS

†1981. Diethyl carbonate

[105-58-8]

C₅H₁₀O₃

(EtO)₂C:O


HCS 1980, 396

Norton, C., *Chemical Engineer*, 1995, (595), 6

A drum containing distillation wastes of ethyl carbonate, ethyl bromoacetate and another bromoester (not intelligibly named) burst spontaneously, releasing a cloud which caused considerable lachrymation and respiratory irritation in nearby housing. Carbon dioxide was doubtless generated, but whether by water contamination or a previously unknown catalytic reaction of the known contents is unclear to the editor.

1982. *trans*-2-Pentene ozonide (3-Ethyl-5-methyl-1,2,4-trioxolane)
[16187-03-4]


C₅H₁₀O₃



See trans-2-Hexene ozonide
See other OZONIDES

1983. Allyldimethylarsine
[691-35-0]

C₅H₁₁As



Ellern, 1968, 24–25
It ignites in air if exposed on filter paper.
See related ALKYLNON-METALS *See other* ALLYL COMPOUNDS

†**1984. 1-Bromo-3-methylbutane**
[107-82-4]

C₅H₁₁Br



See other HALOALKANES

†**1985. 2-Bromopentane**
[107-81-3]

C₅H₁₁Br



See other HALOALKANES

†**1986. 1-Chloro-3-methylbutane**
[107-84-6]

C₅H₁₁Cl



See other HALOALKANES

†**1987. 2-Chloro-2-methylbutane**
[594-36-5]

C₅H₁₁Cl



See other HALOALKANES

†1988. 1-Chloropentane
[543-59-9]




$\text{C}_5\text{H}_{11}\text{Cl}$

See other HALOALKANES

1989. 2-Chlorovinyltrimethyllead
[]

$\text{C}_5\text{H}_{11}\text{ClPb}$



Steingross, W. *et al.*, *J. Organomet. Chem.*, 1966, **6**, 109
It may explode violently during vacuum distillation at 102°C/62.5 mbar.
See related ALKYL METALS

†1990. 2-Iodopentane
[637-97-8]




$\text{C}_5\text{H}_{11}\text{I}$

See other HALOALKANES

†1991. Cyclopentylamine
[1003-03-8]


$\text{C}_5\text{H}_{11}\text{N}$



See other ORGANIC BASES

†1992. *N*-Methylpyrrolidine
[120-94-5]


$\text{C}_5\text{H}_{11}\text{N}$



See other ORGANIC BASES

†1993. Piperidine
[110-89-4]

C₅H₁₁N



HCS 1980, 757

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

N-Nitrosoacetanilide

See N-Nitrosoacetanilide: Piperidine

Other reactants

Yoshida, 1980, 291

MRH values calculated for 13 combinations with oxidants are given.


N-Perchlorylpiperidine

See N-Perchlorylpiperidine: Alone, etc.

See other ORGANIC BASES

1994. *N*-*tert*-Butylformamide
[2425-74-3]

C₅H₁₁NO




Di-*tert*-butyl hyponitrite

See trans-Di-*tert*-butyl hyponitrite: Formamides

†1995. 4-Methylmorpholine
[109-02-4]

C₅H₁₁NO



HCS 1980, 657



See other ORGANIC BASES

†1996. Isopentyl nitrite
[110-46-3]



See other ALKYL NITRITES

1997. *N*-Methylmorpholine oxide (4-Methylmorpholine-4-oxide)
[7529-22-8]
[70187-32-5], [85489-61-8], [80913-66-2], [80913-65-1] (hydrates);
[95650-61-6] (cellulose complex)




Cellulose

Peguy, A. *et al.*, *J. Appl. Poly. Science*, 1990, **40**(3/4), 429

Solutions of cellulose in wet methylmorpholine oxide can undergo exothermic reaction to the point of explosion if confined at elevated temperatures from about 120°C or if otherwise heated to 180°C. The reaction is catalysed by some metals, notably copper.

See other CATALYTIC IMPURITY INCIDENTS, *N*-OXIDES

†1998. Pentyl nitrite
[463-04-7]



Sorbe, 1968, 146

It explodes on heating to above 250°C.

See other ALKYL NITRITES

†1999. 2,2-Dimethylpropane (Neopentane)
[463-82-1]



Preparative hazard

See Aluminium: Halocarbons (reference 9)

†2000. 2-Methylbutane (Isopentane)
[78-78-4]



HCS 1980, 572

646

†2001. Pentane
[109-66-0]


BuMe

C₅H₁₂

FPA H55, 1977; HCS 1980, 716

2002. *N*-Bromotetramethylguanidine
[6926-40-5]

C₅H₁₂BrN₃




Papa, A. J., *J. Org. Chem.*, 1966, **31**, 1426

The material is unstable even at 0°C, and explodes if heated above 50°C at ambient pressure.

See other *N*-HALOGEN COMPOUNDS

2003. 1-Chloro-3-dimethylaminopropane
[109-54-6]

C₅H₁₂ClN




Lithium, Sodium

See Lithium: 1-Chloro-3-dimethylaminopropane, etc.

2004. *N*-Chlorotetramethylguanidine
[6926-39-2]

C₅H₁₂ClN₃





Papa, A. J., *J. Org. Chem.*, 1961, **31**, 1426

The material is unstable even at 0°C, and explodes if heated above 50°C at ambient pressure.

See other *N*-HALOGEN COMPOUNDS

2005. Tetramethylammonium azidocyaniodate(I)
[68574-17-4]




Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 512

This, and the cyanato and selenocyanato (pseudohalogen) analogues are explosive in the solid state, but may be handled with comparative safety. They are also sensitive to laser light.

See related METAL AZIDES


2006. Tetramethylammonium azidocyanatoiodate(I)
[68574-15-2]



See Tetramethylammonium azidocyaniodate(I) (next above)

See related METAL AZIDES, METAL CYANATES

2007. Tetramethylammonium azidoselenocyanatoiodate(I)
[]




See Tetramethylammonium azidocyaniodate(I)

See related METAL AZIDES, METAL CYANATES

2008. Dimethylthallium N-methylacetohydroxamate


[]

C₅H₁₂NO₂TlSchwering, H. U. *et al.*, *J. Organomet. Chem.*, 1975, **99**, 22

It exploded below 160°C, unlike the other dialkylmetal derivatives, which showed high thermal stability.

See other N-O COMPOUNDS**2009. O, O-Dimethyl S-methylcarbamoylmethyl phosphorodithioate**

[60-51-5]

C₅H₁₂NO₃PS₂1. Matos, E. *et al.*, *Chem. Abs.*, 1983, **98**, 949552. Ludovisi, G. *et al.* in *Safety and Runaway Reactions*, Smeder, B. & Mitchison, N. (Eds.), EUR 17723 EN, European Commission, 1997, p11

Accidental exothermic decomposition of the insecticide Dimethoate in a 6000 l vessel led to gross contamination; (no further details of circumstances available from author) [1]. A second incident of bulk decomposition in cyclohexanone solution (concentration not given) is slightly better reported. This was a formulation solution; the details of the processing are not clear from the report but appear to have involved azeotroping the dimethoate dry, then discharging the warm solution (no temperatures given) to the insulated, uncooled, storage vessel which burst. It is claimed that out of specification material had been processed at above previous temperatures. In any event, there was a runaway exotherm, loss of containment, fire, the attentions of regulatory authorities and closure of the factory. The manufacturer's safety study went beyond the then legal requirements, but, as so often, concentrated on toxicity while neglecting the reactivity which could cast that toxicity abroad. A laboratory sample of the liquid residue of the accident itself later burst its container [2].

See other PHOSPHORUS ESTERS**†2010. Butyl methyl ether**

[628-28-4]

C₅H₁₂O**BuOMe**

649

†2011. *tert*-Butyl methyl ether
[1634-04-4]

C₅H₁₂O

Me₃COME

1. Hage, T., *Chem. Abs.*, 1987, **107**, 21833
2. Pearson, H., *Chem. Abs.*, 1988, **108**, 115254

tert-Butyl methyl ether, now manufactured in bulk as a gasoline component, is a less hazardous extracting solvent than diethyl ether, as it scarcely forms peroxides, is less volatile (b.p. 55°C) and has a narrower flammable range (2.5–15%) in air [1]. However, fires involving it are difficult to extinguish with foam, as no film-forming effect is shown [2].

Sulfuric acid

Anon., *Eur. Chem. News*, 1995, **64**(1678), 5

An explosion and fire, costing the life of one worker, occurred when sulphuric acid entered a distillation during scale-up (in glass) of a carotenoid synthesis. This caused a sudden rise in temperature and pressure, bursting the distillation column and an adjacent vessel containing methyl *t*-butyl ether. Reading between the lines: this was the solvent being distilled off; acid catalysis will cause it to revert to methanol and butene. Large scale glass apparatus may burst with overpressures as low as 0.2 bar.

†2012. Ethyl isopropyl ether
[625-54-7]

C₅H₁₂O

EtOCHMe₂

See other PEROXIDISABLE COMPOUNDS


†2013. Ethyl propyl ether
[628-32-0]

C₅H₁₂O

EtOPr

2014. Isopentanol (3-Methylbutanol)
[123-51-3]

C₅H₁₂O



FPA H123, 1983; HCS 1980, 642

Hydrogen trisulfide

See Hydrogen trisulfide: Pentanol

†2015. *tert*-Pentanol (1,1-Dimethylpropanol)
[75-85-4] C₅H₁₂O
 $\text{Me}_2\text{C}(\text{OH})\text{Et}$

HCS 1980, 395

†2016. Diethoxymethane
[462-95-3] C₅H₁₂O₂
 $(\text{EtO})_2\text{CH}_2$

†2017. 1,1-Dimethoxypropane
[4744-10-9] C₅H₁₂O₂
 $(\text{MeO})_2\text{CHEt}$

†2018. 2,2-Dimethoxypropane
[77-76-9] C₅H₁₂O₂
 $(\text{MeO})_2\text{CMe}_2$

Becerra, R. *et al.*, *An. Quim., Ser. A*, 1983, **79**, 163–167

The relationship between the cool flames observed at 210°C and subsequent explosions were studied.

See entry COOL FLAMES

Metal perchlorates

1. Dickinson, R. C. *et al.*, *Chem. Eng. News*, 1970, **48**(28), 6

2. Cramer, R., *Chem. Eng. News*, 1970, **48**(45), 7


3. Mikulski, S. M., *Chem. Eng. News*, 1971, **49**(4), 8

During dehydration of manganese(II) perchlorate [1] or nickel(II) perchlorate [2] with dimethoxypropane, heating above 65°C caused violent explosions, probably involving oxidation by the anion [1] (possibly of the methanol liberated by hydrolysis). Triethyl orthoformate is recommended as a safer dehydrating agent [2] (but methanol would still be liberated).

See Dimethyl sulfoxide: Metal oxosalts

See also METAL PERCHLORATES (reference 2)

2019. 2-(2-Methoxyethoxy)ethanol C₅H₁₂O₃
[111-77-3]




Calcium hypochlorite

See Calcium hypochlorite: Hydroxy compounds

2020. Pentanesulfonic acid
[35452-30-3]

C₅H₁₂O₃S




Preparative hazard

See Nitric acid: Alkanethiols (reference 3)

See other ORGANIC ACIDS

2021. Pentaerythritol
[115-77-5]

C₅H₁₂O₄



Thiophosphoryl chloride

See Thiophosphoryl chloride: Pentaerythritol

†**2022. Tetramethyl orthocarbonate (Tetramethoxymethane)**
[1850-14-2]

C₅H₁₂O₄



†**2023. 2-Methylbutane-2-thiol**
[1679-09-0]

C₅H₁₂S



See other ALKANETHIOLS

†**2024. 3-Methylbutanethiol**
[541-31-1]

C₅H₁₂S



See other ALKANETHIOLS

†2025. Pentanethiol
[110-66-7]





Nitric acid

See Nitric acid: Alkanethiols (reference 3)

See other ALKANETHIOLS

2026. *N,N,N',N'*-Tetramethylformamidinium perchlorate
[2506-80-1]





Menzer, M. *et al.*, *Z. Chem.*, 1977, **17**, 344

The undeuterated salt was refluxed with deuterium oxide to effect deuterium exchange, then the excess was evaporated off under vacuum. Towards the end of the evaporation the moist residue exploded violently. The presence of D is not likely to affect instability of the structure.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

2027. *S*-Diethylamino(methylimino)sulfur(IV) fluoride
[]



von Halasz, A. *et al.*, *Chem. Ber.*, 1971, **104**, 1250, 1253

Care is necessary in the preparation, because if coloured impurities are present, the compound may polymerise violently and very exothermically at ambient temperature.

See other N-S COMPOUNDS

†2028. 1,1-Dimethylpropylamine
[594-39-8]



See other ORGANIC BASES

†2029. 1,2-Dimethylpropylamine
[598-74-3] C₅H₁₃N



See other ORGANIC BASES

†2030. 2,2-Dimethylpropylamine (Neopentylamine)
[5813-64-9] C₅H₁₃N



See other ORGANIC BASES

†2031. *N,N*-Dimethylpropylamine
[926-63-6] C₅H₁₃N



See other ORGANIC BASES

†2032. Isopentylamine
[107-85-7] C₅H₁₃N




See other ORGANIC BASES

†2033. *N*-Methylbutylamine
[110-68-9] C₅H₁₃N




See other ORGANIC BASES

†2034. Pentylamine
[110-58-7] C₅H₁₃N



See other ORGANIC BASES

2035. Bis(2-hydroxyethyl)methylphosphine
[53490-67-8] C₅H₁₃O₂P




Dichloromethylphosphine

See Dichloromethylphosphine: Bis(2-hydroxyethyl)methylphosphine

See related ALKYLPHOSPHINES

2036. Diethylmethylphosphine
[1605-58-9]


C₅H₁₃P



Personal experience
May ignite in air with long exposure.
See other ALKYLPHOSPHINES

2037. 2-Hydroxyethyltrimethylstibonium iodide
[81924-93-8]

C₅H₁₄IOSb



Meyer, E. M. *et al.*, *Biochem. Pharm.*, 1981, **30**, 3003–3005
An attempt to prepare this antimony analogue of choline by heating trimethylstibine with iodoethanol in bis(2-methoxyethyl) ether in a sealed tube at 150°C led to an explosion.


†**2038. 2-Dimethylamino-*N*-methylethylamine**
[142-25-6]

C₅H₁₄N₂



2039. 3-Dimethylaminopropylamine
[109-55-7]

C₅H₁₄N₂



HCS 1980, 423


Cellulose nitrate
See CELLULOSE NITRATE: amines

1,2-Dichloroethane
See 1,2-Dichloroethane: 3-Dimethylaminopropylamine

Other reactants
Yoshida, 1980, 168
MRH values calculated for 14 combinations with oxidants are given.

2040. Pentamethylbismuth
[148739-67-7]

C₅H₁₅Bi




Seppelt, K. *et al.*, *Angew. Chem. (Int.)*, 1994, **33**(9), 976

This compound, formed at -90°C is unstable in solution and the solid explodes on rapid warming to room temperature.

See other ALKYL METALS

2041. Lithium 2,2-dimethyltrimethylsilylhydrazide
[13529-75-4]

C₅H₁₅LiN₂Si




Oxidants

See entry SILYLHYDRAZINES

2042. Lithium pentamethyltitanate-bis(2,2'-bipyridine)
[50662-24-3]

C₅H₁₅LiTi.2C₁₀H₈N₂



Houben-Weyl, 1975, Vol. 13.3, 288

Friction-sensitive, it decomposes explosively.

See related ALKYL METALS

†2043. Dimethylaminotrimethylsilane
[2083-91-2]



$\text{C}_5\text{H}_{15}\text{NSi}$


Xenon difluoride

See Xenon difluoride: Silicon–nitrogen compounds

See related ALKYL SILANES

2044. 1,1,1-Tris(aminomethyl)ethane
[15995-42-3]

$\text{C}_5\text{H}_{15}\text{N}_3$




Preparative hazard

See 1,1,1-Tris(azidomethyl)ethane: Hydrogen, etc.

2045. Dimethyltrimethylsilylphosphine
[26464-99-3]

$\text{C}_5\text{H}_{15}\text{PSi}$



Air, or Water


Goldsberry, R. *et al.*, *Inorg. Chem.*, 1972, **13**, 29–32

It ignites in air and is hydrolysed to dimethylphosphine, also spontaneously flammable.

See related ALKYLPHOSPHINES, ALKYL SILANES

2046. Pentamethyltantalum
[53378-72-6]

$\text{C}_5\text{H}_{15}\text{Ta}$




Mertis, K. *et al.*, *J. Organomet. Chem.*, 1975, **97**, C65

It should be handled with extreme caution, even in absence of air, because dangerous explosions have occurred on warming frozen samples or during transfer operations.

See other ALKYL METALS

2047. 1,2-Dimethyl-2-trimethylsilylhydrazine
[13271-93-7]

C₅H₁₆N₂Si




Oxidants

See entry SILYLHYDRAZINES

2048. Pentaamminepyridineruthenium(II) perchlorate
[19482-31-6]

C₅H₂₀Cl₂N₆O₈Ru




Creutz, C. A., private comm., 1969

The dry salt exploded on touching.

See other AMMINOMETAL OXOSALTS

2049. Potassium pentacarbonylvandate(3-)
[78937-14-1]

C₅K₃O₅V



Alone, or Poly(chlorotrifluoroethylene)

Ellis, J. E. *et al.*, *J. Amer. Chem. Soc.*, 1981, **103**, 6101


It is a pyrophoric and treacherously shock-sensitive explosive, especially as the dry black form. It will deflagrate under nitrogen or argon when scratched with a spatula, and ignites in contact with poly(chlorotrifluoroethylene), (Fluorolube). The caesium and rubidium salts are not shock-sensitive but will deflagrate on friction and are

pyrophoric. An insoluble by-product from the preparation will explode if treated with limited amounts of water or alcohol.

See related CARBONYLMETALS

2050. 2-Diazonio-4,5-dicyanoimidazolid (Diazodicyanoimidazole)
[40953-35-3]

C_5N_6



1. Sheppard, W. A. *et al.*, *J. Amer. Chem. Soc.*, 1973, **95**, 2696


2. Sheppard, W. A. *et al.*, *J. Org. Chem.*, 1979, **44**, 1718

It is highly shock-sensitive when dry and explodes above 150°C [1]. It must not be scraped or rubbed, or used in above 2 g portions. The crystalline 1:1 complex with 18-crown-6 ether is significantly less hazardous [2].

See other CYANO COMPOUNDS, DIAZONIUM SALTS, HIGH-NITROGEN COMPOUNDS

2051. Sodium pentacarbonylrhenate
[33634-75-2]

C_5NaO_5Re




Bailar, 1973, Vol. 3, 955

This and other alkali-metal salts are pyrophoric.

See related CARBONYLMETALS

2052. Silver 1,3,5-hexatriynide
[]

C_6Ag_2




Hunsman, W., *Chem. Ber.*, 1950, **83**, 216

The silver salt may be handled moist, but when dry it explodes violently on touching with a glass rod.

See other METAL ACETYLIDES, SILVER COMPOUNDS

2053. Pentafluorophenylaluminium dibromide
[4457-90-3]

$C_6AlBr_2F_5$



Air, or Water


1. Chambers, R. D. *et al.*, *J. Chem. Soc. (C)*, 1967, 2185
2. Chambers, R. D. *et al.*, *Tetrahedron Lett.*, 1965, 2389

It ignites in air [1], and explodes violently on rapid heating to 195°C or during uncontrolled hydrolysis [2].

See other HALO-ARYLMETALS

2054. 2,5-Diazido-3,6-dichlorobenzoquinone
[26157-96-0]

$C_6Cl_2N_6O_2$




1. Gilligan, W. H., *Tetrahedron Lett.*, 1978, 1676
2. Winkelman, E., *Tetrahedron*, 1969, **25**, 2427

While it does not explode on heating, it is moderately impact-sensitive [1], contradicting an earlier report [2] of absence of explosive properties.

See other 2-AZIDOCARBONYL COMPOUNDS

2055. Hexachlorobenzene (Perchlorobenzene)
[118-74-1]


C_6Cl_6



Calcium hydride



See Calcium hydride: Hexachlorobenzene

2056. Hexacarbonylchromium
[13007-92-6]



Weast, 1972, B-83
It explodes at 210°C.
See other CARBONYLMETALS



2057. 1,3,5-Trifluorotrinitrobenzene
[1423-11-6]



Nucleophiles

Koppes, W. M. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1981, 1817
The title compound readily undergoes substitution reactions with a variety of N, O, C and X nucleophiles. Many of the products of such reactions, especially with hydrazine and its derivatives, are explosive.
See other POLYNITROARYL COMPOUNDS

2058. Pentafluoroiodosylbenzene
[14353-90-3]



1. Collmann, J. P., *Chem. Eng. News*, 1985, **63**(11), 2
2. Groves, J. T. *et al.*, *Chirality*, 1998, **10**, 106


A 12 g sample decomposed explosively on heating under vacuum at 60°C, though it has been described as stable up to 210°C when it decomposes violently. Avoid heating it to remove volatile impurities [1]. In later safety warnings it has been described as able to detonate spontaneously [2].

See other IODINE COMPOUNDS

2059. Pentafluorophenyllithium

[1076-44-4]

C₆F₅Li



Krafft, T. E. USA Pat. 5,679,289, 1996

A procedure is described for safe preparation and use of this compound in bulk. It depends upon keeping inventories low.

Deuterium oxide

Kinsella, E. *et al.*, *Chem. & Ind.*, 1971, 1017

During addition of an ethereal solution of deuterium oxide (containing some peroxide) to a suspension of the organolithium reagent in pentane, a violent explosion occurred. This may have been initiated by the peroxide present, but probably involved elimination of lithium fluoride.


See other HALO-ARYLMETALS

See other FLUORINATED ORGANOLITHIUM COMPOUND

†2060. Hexafluorobenzene

[392-56-3]

C₆F₆



Metals

See HALOARENEMETAL π -COMPLEXES

Silane


Koga, Y. *et al.*, *J. Phys. Chem.*, 1987, **91**(2), 298–305

Reaction of hexafluorobenzene and silane under single pulse irradiation of a MW CO₂ laser was explosive.

See other IRRADIATION DECOMPOSITION INCIDENTS

2061. Trifluoromethyl 3-fluorocarbonylhexafluoroperoxybutyrate
[32750-98-4]


C₆F₁₀O₄



Bernstein, P. A. *et al.*, *J. Amer. Chem. Soc.*, 1971, **93**, 3885
A small sample exploded at 70°C.
See other PEROXYESTERS

2062. Perfluorohexyl iodide
[355-43-1]

C₆F₁₃I




Sodium

See Sodium: Halocarbons (reference 6)

2063. Potassium hexacyanoferrate(III) ('Potassium ferricyanide')
[13746-66-2]

C₆FeK₃N₆



HCS 1980, 765

Ammonia

1. Pieters, 1957, 30
2. Sidgwick, 1950, 1359

Contact may be explosive [1], possibly owing to rapid oxidation of ammonia by alkaline 'ferricyanide' [2].

See Potassium tetracyanomercuroate(II)

See other AMINATION INCIDENTS

Chromium trioxide

See Chromium trioxide: Potassium hexacyanoferrate(III)

Hydrochloric acid

Ephraim, 1939, 303

Treatment of the complex salt with acid liberates the corresponding complex ferri-cyanic acid, an oxidant which is rather endothermic (ΔH_f° (aq) +640.5 kJ/mol, 2.96 kJ/g), and which forms solid complexes with ether, etc.

Sodium nitrite


See Sodium nitrite: Metal cyanides

See other METAL CYANIDES (AND CYANO COMPLEXES)

2064. Potassium hexacyanoferrate(II) ('Potassium ferrocyanide')

[13943-58-3]

C₆FeK₄N₆



HCS 1980, 766

Copper(II) nitrate

See Copper(II) nitrate: Potassium hexacyanoferrate(II)

Hydrochloric acid

Ephraim, 1939, 303

Treatment of the complex salt with acid produces the corresponding complex ferro-cyanic acid, which is rather endothermic (ΔH_f° (aq) +534.7 kJ/mol, 2.48 kJ/g), and which forms solid complexes with ether, etc.

See other ENDOTHERMIC COMPOUNDS

Sodium nitrite


See Sodium nitrite: Metal cyanides

See other METAL CYANIDES (AND CYANO COMPLEXES)

2065. Iron(III) oxalate

[2944-66-3]

C₆Fe₂O₁₂




Weinland, R. *et al.*, *Z. Anorg. Chem.*, 1929, **178**, 219

The salt, probably complex, decomposes at 100°C.
See other METAL OXALATES

2066. 4-Chloro-2,5-dinitrobenzenediazonium 6-oxide


[]



See Nitric acid: 4-Chloro-2-nitroaniline
See other ARENEDIAZONIUM OXIDES

2067. 3,4-Difluoro-2-nitrobenzenediazonium 6-oxide


[]



Finger, G. C. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 148
Diazotisation of 3,4,6-trifluoro-2-nitroaniline under differing conditions gave either this isomer or the 3,6-difluoro 4-oxide (next below), both of which exploded on heating, ignition or impact.
See other ARENEDIAZONIUM OXIDES

2068. 3,6-Difluoro-2-nitrobenzenediazonium 4-oxide


[]



Finger, G. C. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 148
Diazotisation of 3,4,6-trifluoro-2-nitroaniline under differing conditions gave either this isomer or the 3,4-difluoro 6-oxide (next above), both of which exploded on heating, ignition or impact.
See other ARENEDIAZONIUM OXIDES

2069. 1,2,3,-4,5-Pentafluorobicyclo[2.2.0]hexa-2,5-diene
[21892-31-9]

C₆HF₅




1. Rotajezak, E., *Roczn. Chem.*, 1970, **44**, 447–449
2. Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455

Both isomers were stored as frozen solids, for as liquids they are more explosive than the hexafluoro ‘Dewar benzene’ analogue [1]. Energy of decomposition (320–540°C) was determined by DSC as 4.60 kJ/g.

See other HALOALKENES, STRAINED-RING COMPOUNDS

2070. 1,2,3,-5,6-Pentafluorobicyclo[2.2.0]hexa-2,5-diene
[28663-71-0]

C₆HF₅




1. Rotajezak, E., *Roczn. Chem.*, 1970, **44**, 447–449
2. Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455

Both isomers were stored as frozen solids, for as liquids they are more explosive than the hexafluoro ‘Dewar benzene’ analogue [1]. Energy of decomposition (320–540°C) was determined by DSC as 4.60 kJ/g.

See other HALOALKENES, STRAINED-RING COMPOUNDS

2071. Lead 2,4,6-trinitroresorcinoxide (‘Lead styphnate’)
[15245-44-0]

C₆HN₃O₈Pb



1. *MCA Case History No. 957*
2. Okazaki, K. *et al.*, *Chem. Abs.*, 1977, **86**, 57638
3. Biasutti, 1981, 63, 73, 75, 86, 108

Three beakers containing lead styphnate were being heated in a laboratory oven to dry the explosive salt. When one of the beakers was moved, all 3 detonated. Other heavy metal salts of polynitrophenols are dangerously explosive when dry [1]. The desensitising effect of presence of water upon the friction sensitivity of this priming explosive was studied. There is no effect up to 2% content and little at 5%. Even

at 20% water content it is still as sensitive as dry pentaerythritol tetranitrate [2]. In one incident in 1927, 20 kg of the dry salt exploded while being poured into paper bags. Initiation by static sparks in a very dry atmosphere was supposed. In another incident, 1 kg exploded during drying in thin layers in an oven held below 65°C. In a third incident, sifting of the dry material also led to the explosion of adjacent wet material, though in a fourth incident, the latter did not also explode. The other incidents involved explosions during drying or handling operations [3].


See LEAD SALTS OF NITRO COMPOUNDS

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, POLYNITROARYL COMPOUNDS

2072. 2,3,5-Trinitrobenzenediazonium-4-oxide

[]

C₆HN₅O₇



Meldola, R. *et al.*, *J. Chem. Soc.*, 1909, **95**, 1383


It is extremely explosive.

See other ARENEDIAZONIUM OXIDES

2073. 3,4-Bis(1,2,3,4-thiatriazol-5-ylthio)maleimide

[1656-16-2]

C₆HN₇O₂S₄



Pilgram, K. *et al.*, *Angew. Chem.*, 1965, **77**, 348


This compound of low oxygen balance explodes loudly on impact or at the m.p.

See other HIGH-NITROGEN COMPOUNDS, N-S COMPOUNDS

2074. 1,3,5-Hexatriyne

[3161-99-7]

C₆H₂



1. Hunsman, W., *Chem. Ber.*, 1950, **83**, 213–217


2. Armitage, J. B. *et al.*, *J. Chem. Soc.*, 1952, 2013

3. Bjornov, E. *et al.*, *Spectrochim. Acta*, 1974, **30A**, 1255–1256

It polymerises slowly at -20°C , but rapidly at ambient temperature in air to give a friction-sensitive explosive solid, probably a peroxide [1]. Although stable under vacuum at -5°C for a limited period, exposure at 0°C to air led to violent explosions soon afterwards [2]. Explosive hazards have again been stressed [3].

See other ALKYNES

2075. Silver 2-azido-4,6-dinitrophenoxide
[82177-80-8]




Spear, R. J. *et al.*, *Aust. J. Chem.*, 1982, **35**, 5

Of a series of 17 compounds examined as stab-initiation sensitisers, the title compound had the lowest ignition temperature of 122°C .

See other METAL NITROPHENOXIDES, ORGANIC AZIDES, POLYNITROARYL COMPOUNDS, SILVER COMPOUNDS

2076. 2,6-Dibromobenzoquinone-4-chloroimide
[537-45-1]



1. Hartmann, W. W. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 177

2. Horber, D. F. *et al.*, *Chem. & Ind.*, 1967, 1551



3. Taranto, B. J., *Chem. Eng. News*, 1967, **45**(24), 54

The chloroimide decomposed violently on a drying tray at 60°C [1], and a bottle accidentally heated to 50°C exploded [2]. Decomposition of the heated solid occurs after a temperature-dependent induction period. Unheated material eventually exploded after storage at ambient temperature [3].

See 2,6-Dichlorobenzoquinone-4-chloroimide

See other N-HALOGEN COMPOUNDS, INDUCTION PERIOD INCIDENTS

2077. 1,5-Dichloro-2,4-dinitrobenzene
[3698-83-7]



Dimethyl sulfoxide, Potassium fluoride

Koch-Light Laboratories Ltd., private comm., 1976

A mixture of the (highly activated) dichloro compound with potassium fluoride in the solvent was heated to reflux to effect replacement of chlorine by fluorine. The reaction accelerated out of control and exploded, leaving much carbonised residue. Analogous reactions had been effected uneventfully on many previous occasions.


See Dimethyl sulfoxide: Acyl halides, etc.

See other HALOARYL COMPOUNDS, POLYNITROARYL COMPOUNDS

2078. 2,6-Dichlorobenzoquinone-4-chloroimide

[101-38-2]

C₆H₂Cl₃NO



Taranto, B. J., *Chem. Eng. News*, 1967, **45**(52), 54

The heated solid material decomposed violently after a temperature-dependent induction period. Unheated material eventually exploded after storage at ambient temperature.


See 2,6-Dibromobenzoquinone-4-chloroimide

See other N-HALOGEN COMPOUNDS, INDUCTION PERIOD INCIDENTS

2079. 1,2,4,5-Tetrachlorobenzene

[95-94-3]

C₆H₂Cl₄



Sodium hydroxide, Solvent

1. Anon., *Chem. Eng. News*, 1983, **61**(23), 44
2. *MCA Case History No. 620*
3. Hofmann, H. T., *Arch. exptl. Pathol. Pharmacol.*, 1957, **232**, 228–230
4. Cardillo, P. *et al.*, *J. Haz. Mat.*, 1984, **9**, 221–234
5. Sambeth, J. A., *Chimia*, 1982, **36**, 128–132
6. Anon., *ABCM Quart. Safety Summ.*, 1952, **23**, 5
7. May, G., *Brit. J. Ind. Med.*, 1973, **30**, 276–277
8. Milnes, M. H., *Nature*, 1971, **232**, 396
9. Calnan, A. B., private comm., 1972
10. Theofanous, T. G., Statement at special meeting of Society of Chemical Industry's Industrial Health & Safety Group, London, 23rd March, 1981; *Chem. & Ind.*, 1981, 197; *Nature*, 1981, **291**, 640

11. Theofanous, T. G., *Chem. Engrg. Sci.*, 1983, **38**, 1615
12. Salomon, C. M., *Chimia*, 1982, **36**, 133–139
13. Künzi, H., *Chimia*, 1982, **36**, 162–168
14. Cardillo, P. *et al.*, *Chim. e Ind. (Milan)*, 1983, **65**, 611–615
15. Theofanous, T. G., *Chem. Engrg. Sci.*, 1983, **38**, 1631
16. Cardillo, P. *et al.*, *Chim. e Ind. (Milan)*, 1982, **64**, 781–784
17. Marshall, V. C., *Loss Prevention Bull.*, 1992, **104**, 15

Eight serious accidents occurred during the commercial preparation of 2,4,5-trichlorophenol by alkaline partial hydrolysis of 1,2,4,5-tetrachlorobenzene during the period 1949 to 1976 [1], involving two distinct processes, each with their particular potential hazards. The earlier process used methanolic alkali under autogenous pressure to effect the hydrolysis, and on 2 occasions around 1949 the reaction at 125°C went out of control, one attaining 400°C [2]. One of the principal hazards of this pressurised process is associated with the fact that no reflux is possible, so the reaction exotherm must be removed by applied cooling in a batch reactor, (or by controlled flow rate in a continuous reactor) to prevent undue temperature rise. A second potential hazard resides in the fact that if the sodium hydroxide normally used as base (and in excess) is not completely dissolved in the methanol before the temperature of onset of hydrolysis (165–180°C) is reached, the hydrolysis exotherm (108.4 kJ/mol of tetrachlorobenzene) is increased by the solution exotherm (perhaps 30–90 kJ/mol) of undissolved alkali, and a runaway reaction, accompanied by a rapid and uncontrollable pressure increase may develop. In a further incident in 1953, the explosion was associated with the post-reaction stage during distillation of methanol from the reaction mixture [3]. Under these conditions a virtually solid crust of the sodium phenoxide is formed and may become overheated [4]. Four explosions occurred during the commercial use of the methanol process [5].

The later process used ethylene glycol as solvent to effect hydrolysis with sodium hydroxide and operated essentially at or near atmospheric pressure. Hydrolysis was effected at between 140 and 180°C during several hours, and the excess glycol (that portion which did not condense to relatively involatile diethylene glycol) might or might not be recovered by vacuum distillation, depending on the process design. When another solvent was present in the reaction system (dichlorobenzene to dissolve sublimed tetrachlorobenzene from the condenser in the Coalite system, or xylene to assist in water and glycol removal in the Seveso system), some inherent temperature control by reflux during the hydrolysis reaction was possible. Violent decomposition could, however, occur during the subsequent solvent distillation phase in the absence of effective temperature control. A laboratory residue from vacuum stripping using electric heating (without knowledge of the liquid temperature) exploded when the vapour temperature had reached 160°C [6]. In the Coalite plant incident in 1968, where the hydrolysis was run at about 180°C in a reaction vessel heated by circulating oil at 300°C, failure of the manually regulated oil heating system led to an uncontrollable temperature increase during 50 m to above 250°C, when a violent explosion occurred [7]. Subsequent work showed that an accelerating exothermic decomposition sets in at 230°C, capable of attaining 410°C. This involves sodium hydroxide and ethylene glycol (or diethylene glycol) and produces hydrogen, thus pressurisation, and sodium carbonate [8,16].

The Seveso accident in 1976 also involved the glycol-based process, but differed fundamentally from the 1968 incident. While the latter apparently involved a thermal runaway initiated during the hydrolysis reaction by application of excessive heat by the faulty hot oil system [7], the process design adopted by Icmesa at Seveso featured heating the reaction vessel by steam at 12 bar (192°C if saturated) to ensure a minimum 40°C safety margin below the known decomposition temperature of 230°C [5]. At Seveso the exothermic hydrolysis reaction had been completed, but not the ensuing xylene–glycol solvent distillation, before the weekend shutdown, when the liquid temperature was 158°C. The bursting disk set at 3.8 bar failed some 7.5 h after processing operations, agitation and heating had been terminated [5]. The accident was followed by an unprecedented amount of investigational work; giving a highly probable explanation if several factors and new information are taken into account.

It was found that the steam supply to the reactor was often superheated (just prior to shutdown to 330°C) [10]. Although this degree of superheat would not grossly increase the temperature of the inner reactor wall in contact with the liquid (or the bulk liquid temperature) [11], it seems probable that any reaction material splashed onto and dried out at the top of the coil-heated wall would have become heated to a much higher temperature. Further detailed work on the thermal stability of the mixture showed that a previously unsuspected very slow exothermic decomposition existed, beginning at 180°C and proceeding at an appreciable rate only above 200°C, so that the exotherm was insufficient to heat the contents of the reactor from the last recorded temperature of 158°C to the decomposition temperature of 230°C in 7.5 h [12,13,14]. It was concluded that an alternative (effectively an external) source of heat was necessary to account for the observed effect, and the residual superheat from the steam at 330°C seems to have been that source.

Any superheating effect would be localised near the steam inlet at the top of the vessel, and there would not have been enough excess heat to raise the temperature of the whole reactor contents by more than a few °C. A new physicochemical process involving conduction, convection and radiation was proposed, and verified experimentally, which concentrated the available excess heat into the top surface layer of a few cm (a few % of the total depth) of the reactor contents, so that the critical temperature of 230°C would be attained in that top layer [11]. The exothermic decomposition reaction, (perhaps catalysed by any thermal degradation products falling into the top layer from the overheated top of the reactor) would then have propagated rapidly downwards into the reactor contents (some 6 t), leading to runaway reaction, pressure build-up and bursting disk failure with release of the reactor contents to atmosphere [11,15].

A publication summarises all the then available technical evidence related to the Seveso accident, and recommends operational criteria to ensure safety in commercial processes to produce trichlorophenol [4]. All the plant scale incidents were characterised [1] by the subsequent occurrence of chloracne arising from the extremely toxic and dermatitic compound 2,3,7,8-tetrachlorodibenzodioxin (structure IX, p. S-3), formed during the thermal runaway reaction and dispersed in the ensuing explosion. It is also extremely resistant to normal chemical decontamination procedures, and after the 1968 explosion, further cases occurred after transient contact with plant



which had been decontaminated and allowed to weather for 3 years, and which appeared free of 'dioxin' [7,9]. The consequences at Seveso: 447 cases of chemical burns (NaOH); and 179 of chloracne, only 34 with both [17].

See Sodium hydroxide: Glycols

See other GAS EVOLUTION INCIDENTS, RUNAWAY REACTIONS

See other HALOARYL COMPOUNDS

2080. 1,5-Difluoro-2,4-dinitrobenzene
[327-92-4]





Preparative hazard

See 1,5-Dichloro-2,4-dinitrobenzene: Dimethyl sulfoxide, etc.



See other HALOARYL COMPOUNDS, POLYNITROARYL COMPOUNDS

†2081. 1,2,4,5-Tetrafluorobenzene
[327-54-8]




See other HALOARYL COMPOUNDS

2082. Tetrafluorobenzene-1,4-diol
[771-63-1]



See Trimethylaluminium: Tetrafluorobenzene-1,4-diol

2083. Sodium 3-hydroxymercurio-2,6-dinitro-4-*aci*-nitro-2,5-cyclohexadienonide
 [] **C₆H₂HgN₃NaO₅**





Hantzsch, A. *et al.*, *Ber.*, 1906, **39**, 1111

This salt of the mono-*aci p*-quinonoid form of 3-hydroxymercurio-2,4,6-trinitrophenol explodes on rapid heating.

See entry *aci*-NITROQUINONOID COMPOUNDS

2084. 3,4,5-Triiodobenzediazonium nitrate
 [68596-99-6]




Kalb, L. *et al.*, *Ber.*, 1926, **59**, 1867

Unstable on warming, explodes on heating in a flame.

See other DIAZONIUM SALTS

2085. Potassium picrate (Potassium 2,4,6-trinitrophenoxide)
 [573-83-1]





2,2-Dinitro-2-fluoroethoxycarbonyl chloride

See 2,2-Dinitro-2-fluoroethoxycarbonyl chloride: Potassium picrate

See other PICRATES


2086. Sodium picrate (Sodium 2,4,6-trinitrophenoxide)
[73771-13-8]




1. Coningham, H., *H. M. Insp. Expl. Rept.* 236, London, 1919
 2. Nakagawa, K. *et al.*, *Bull. Chem. Soc. Japan*, 1987, **60**, 2038, footnote
- A large scale explosion involving initiation of wet sodium picrate by impact was investigated [1]. A small sample heated to above 250°C exploded with sufficient violence to destroy the DTA thermocouple [2].
See other METAL NITROPHENOXIDES, PICRATES

2087. 3,5-Dinitrobenzenediazonium 2-oxide (5,7-Dinitrobenzoa-1,2,3-diazole)

[4682-03-5]



[87-31-0] (oxadiazole form)




It appears uncertain which structure this explosive has, quite possibly both, since it can be made either by diazotisation or nucleophilic cyclisation.

See N,2,3,5-Tetranitroaniline

See other ARENEDIAZONIUM OXIDES

2088. 4,6-Dinitrobenzenediazonium 2-oxide

[]




Urbanski, 1967, Vol. 3, 204

This priming explosive, as sensitive as mercury fulminate, is much more powerful than metal-containing initiators.

See other ARENEDIAZONIUM OXIDES

2089. 4,6-Dinitrobenzofurazan N-oxide
[5128-28-9]



Norris, W. P. *et al.*, *Aust. J. Chem.*, 1983, **36**, 306

A powerful high explosive, of comparable sensitivity to dry picric acid.

Cysteine

See 4-(2'-Ammonio-2'-carboxyethylthio)-5,7-dinitro-4,5-dihydro-benzofurazanide N-oxide

Furan, or N-Methylindole, or N-Methylpyrrole

Halle, J. C. *et al.*, *Tetrahedron Lett.*, 1983, **24**, 2258

The mono- or bis-furoxan adducts are explosive, and work on them was discontinued.

Ketones

Terrier, F. *et al.*, *J. Org. Chem.*, 1981, **46**, 3538

The adducts formed with acetone, cyclopentanone, cyclopentanedione, 2,4-pentanedione or 3-methyl-2,4-pentanedione (See Structure str01, p. S-1, R=CH₃CO.CH₂, etc.) form insoluble potassium salts (M=K), but these are highly explosive and shock-sensitive and should not be isolated, but handled only in solution.

Nucleophilic reagents


Norris, W. P. *et al.*, *Aust. J. Chem.*, 1983, **36**, 297–309

Reaction of the title compound with a series of nucleophilic oxygen reagents (potassium hydrogencarbonate in water or methanol, potassium hydroxide in methanol), or nitrogen reagents (ammonia, hydroxylamine or hydrazine hydrate) gives a series of mono- or bi-nuclear 7-substituted Meisenheimer complexes (structure II, p. S-4, R=OH, OMe, NH₂, NHOH or NHNH₂). All of these are explosives sensitive to initiation by impact, friction or electrostatic discharge, which need due care in handling.

See other FURAZAN N-OXIDES, POLYINITROARYL COMPOUNDS

2090. 3,5-Dinitro-4-hydroxybenzenediazonium 2-oxide
[29906-35-2]

$C_6H_2N_4O_6$




Dynamit Nobel, Belg. Pat. 874 187, 1979

The barium, lead and potassium salts (on the 4-position) are sensitive detonators.

See other ARENEDIAZONIUM OXIDES, POLYNITROARYL COMPOUNDS

2091. 2,3,4,6-Tetranitrophenol
[641-16-7]

$C_6H_2N_4O_9$




Sorbe, 1968, 152

Exceptionally explosive.

See other POLYNITROARYL COMPOUNDS

2092. Picryl azide (2,4,6-Trinitrophenyl azide)
[1600-31-3]

$C_6H_2N_6O_6$




Schrader, E., *Ber.*, 1917, **50**, 778

Explodes weakly under impact, but not on heating.

See other ORGANIC AZIDES, POLYNITROARYL COMPOUNDS

2093. Pentanitroaniline
[21958-87-5]





Koppes, W. M. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1981, 1816

It is a very sensitive explosive.

See other POLYNITROARYL COMPOUNDS

2094. 2,4-Hexadiyne-1,6-dioic acid
[]





Sorbe, 1968, 109

Explodes on heating.

See other ACETYLENIC COMPOUNDS, ORGANIC ACIDS

2095. Triethynylaluminium
[61204-16-8]



Diethyl ether, or Dioxane, or Trimethylamine

1. Chini, P. *et al.*, *Chim. e Ind. (Milan)*, 1962, **44**, 1220

2. Houben-Weyl, 1970, Vol. 13.4, 159


The residue from sublimation of the complex with dioxane is explosive, and the complex should not be dried by heating. The trimethylamine complex may also

explode on sublimation [1]. Triethynylaluminium or its complex with diethyl ether may decompose explosively on heating. Sublimation is not therefore advised as a purification method [2].

See other METAL ACETYLIDES

2096. Triethynylarsine
[687-78-5]

C_6H_3As




Voskuil, W. *et al.*, *Rec. Trav. Chim.*, 1964, **83**, 1301

It explodes on strong friction.

See related METAL ACETYLIDES

2097. 6-Bromo-2,4-dinitrobenzenediazonium hydrogen sulfate
[65036-47-7]

$C_6H_3BrN_4O_8S$




See entry DIAZONIUM SULFATES

See other POLYNITROARYL COMPOUNDS

2098. 1-Chloro-2,4-dinitrobenzene
[97-00-7]

$C_6H_3ClN_2O_4$



HCS 1980, 299

Sorbe, 1968, 56

$T_{\text{ait}24}$ was determined as 250°C by adiabatic Dewar tests, with an apparent energy of activation of 185 kJ/mol.

It has been used as an explosive.

See 1-Fluoro-2,4-dinitrobenzene

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Ammonia

1. Uhlmann, P. W., *Chem. Ztg.*, 1914, **38**, 389–390

2. Wells, F. B. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 221–222

During the preparation of 2,4-dinitroaniline by a well-established procedure involving heating the reactants in a direct-fired autoclave (170°C and 40 bar were typical conditions), a sudden increase in temperature and pressure preceded a violent explosion [1]. An alternative process avoiding the use of a sealed vessel is now used [2].

See 2-Chloronitrobenzene: Ammonia

See POLYNITROARYL COMPOUNDS: bases, or salts

See other AMINATION INCIDENTS

Hydrazine hydrate

491M, 1975, 201

In absence of diluent, the reaction is sufficiently exothermic and violent to shatter the flask.

Other reactants

Yoshida, 1980, 157


MRH values calculated for 7 combinations with materials catalysing its decomposition are given.

See other HALOARYL COMPOUNDS, POLYNITROARYL COMPOUNDS

2099. 2,4-Dinitrobenzenesulfenyl chloride

[528-76-7]

$\text{C}_6\text{H}_3\text{ClN}_2\text{O}_4\text{S}$




Kharasch, N. *et al.*, *Org. Synth.*, 1964, **44**, 48

During removal of solvent under vacuum, the residual chloride must not be overheated, as it may explode.

See related NITROACYL HALIDES See other POLYNITROARYL COMPOUNDS

2100. 2,4-Dinitrobenzenesulfonyl chloride
[1656-44-6]

C₆H₃ClN₂O₆S




Preparative hazard

See Bis(2,4-dinitrophenyl) disulfide: Chlorine

See other NITROACYL HALIDES, POLYNITROARYL COMPOUNDS

2101. 2,6-Dinitro-4-perchlorylphenol
[]

C₆H₃ClN₂O₈




Gardner, D. M. *et al.*, *J. Org. Chem.*, 1963, **28**, 2652

This analogue of picric acid is dangerously explosive and very shock-sensitive.

See other PERCHLORYL COMPOUNDS, POLYNITROARYL COMPOUNDS

2102. 6-Chloro-2,4-dinitrobenzenediazonium hydrogen sulfate
[68597-05-7]

C₆H₃ClN₄O₈S



See entry DIAZONIUM SULFATES

See other POLYNITROARYL COMPOUNDS

2103. Dichloronitrobenzene

[?]




Aniline

Anon., *Chem. Hazards Ind.*, 1992, (Feb.), 171

A reaction with an unspecified isomer of dichloronitrobenzene ran wild on overheating.

See other RUNAWAY REACTIONS *See other* NITROARYL COMPOUNDS**2104. 2,4,-Dichloronitrobenzene**

[611-06-3]




Acetic acid, Potassium fluoride, Dimethylacetamide

Mooney, D. G., *Hazards XI*, 38132, Symp. Ser. 124, Rugby (UK), IChE, 1991van Reijndam, J. W. *et al.*, *J. Loss Prev. Proc. Ind.*, 1992, **5**(4), 211

A Halex process, exchanging the chlorine for fluorine in solvent dimethylacetamide, on 15,000l scale with above a tonne of substrate, showed a runaway exotherm from 160°C, followed by rupture of the vessel and formation of a fireball by the contents. Fragments and damage extended up to 500m distance. Examination suggested that a complex chain of events, starting with large scale contamination of a previous batch by water during work up, had produced recycled dimethylacetamide contaminated with acetic acid. Acetate then acted as nucleophile, subsequently eliminating ketene to produce a phenol, itself another nucleophile producing diphenyl ethers. The limited exotherm produced by these reactions was enough to take the mix to 250°C, where a more exothermic reaction, probably involving the oxidant nitro group, sets in. Presumably similar reactions could take place with other halonitrobenzenes.

See 5-Fluoro-2-nitrophenol*See other* RUNAWAY REACTIONS *See other* HALOARYL COMPOUNDS, NITROARYL COMPOUNDS**2105. 3,4-Dichloronitrobenzene**

[99-54-7]




Catalyst, Hydrogen

See 3,4-Dichlorophenylhydroxylamine*See other* HALOARYL COMPOUNDS, NITROARYL COMPOUNDS

2106. 2,6-Dichloro-4-nitrobenzenediazonium hydrogen sulfate
[68597-06-8]


C₆H₃Cl₂N₃O₆S



See entry DIAZONIUM SULFATES

2107. 2,4,5-Trichlorophenol
[95-95-4]

C₆H₃Cl₃O




Preparative hazard

See 1,2,4,5-Tetrachlorobenzene: Sodium hydroxide, Solvent

See other HALOARYL COMPOUNDS

2108. 1-Fluoro-2,4-dinitrobenzene
[70-34-8]

C₆H₃FN₂O₄



HCS 1980, 505

1. Halpern, B. D., *Chem. Eng. News*, 1951, **29**, 2666
2. Muir, G. D., private comm., 1969
3. Reinheimer, J. D. *et al.*, *J. Amer. Chem. Soc.*, 1961, **83**, 837

The residue left from conversion of the chloro to the fluoro compound exploded during distillation at 1.3 mbar [1]. Reheating the residue from distillation caused violent decomposition [2]. Traces of an *aci*-nitroquinonoid species (structure str02, p. S1-S7) may have been formed in the chlorodinitrobenzene–potassium fluoride reaction system [3].

See related aci-NITROQUINONOID COMPOUNDS



Ether peroxides

Shafer, P. R., private comm., 1967

When air was admitted after vacuum evaporation of a (peroxidic) ether solution of the dinitro compound (5 g), a violent explosion occurred. Exploding ether peroxide may have initiated the dinitro compound.

See other HALOARYL COMPOUNDS, POLYNITROARYL COMPOUNDS

2109. 2,4-Difluoronitrobenzene [446-35-5]





Sodium hydroxide, Dioxane, Water

See 5-Fluoro-2-nitrophenol


See other NITROARYL COMPOUNDS

†2110. 1,2,4-Trifluorobenzene [367-23-7]



See other HALOARYL COMPOUNDS

2111. Sodium 2-hydroxymercurio-6-nitro-4-*aci*-nitro-2,5-cyclohexadienonide
 [] **C₆H₃HgN₂NaO₆**




Hantzsch, A. *et al.*, *Ber.*, 1906, **39**, 1113

This salt of the mono-*aci-p*-quinonoid form of 2-hydroxymercurio-4,6-dinitrophenol explodes violently on strong heating.

See entry *aci*-NITROQUINONOID COMPOUNDS


See other MERCURY COMPOUNDS, POLYNITROARYL COMPOUNDS

2112. Potassium 1,3,5-trinitrobenzene
 [] **C₆H₃KN₃O₆**



See entry NITROAROMATIC-ALKALI HAZARDS; potassium radical salts

2113. Potassium 6-*aci*-nitro-2,4-dinitro-2,4-cyclohexadieniminide
 [[12244-59-6] (ion)] **C₆H₃KN₄O₆**




Green, A. G. *et al.*, *J. Chem. Soc.*, 1913, **103**, 508–513

This salt of the *aci-o*-iminoquinonoid form of 2,4,6-trinitroaniline explodes violently at 110°C.

See entry *aci*-NITROQUINONOID COMPOUNDS

See other POLYNITROARYL COMPOUNDS

2114. Potassium 4-hydroxy-5,7-dinitro-4,5-dihydrobenzofurazanide 3-oxide
[[57891-85-7] (ion)] $C_6H_3KN_4O_7$




1. Boulton, A. J. *et al.*, *J. Chem. Soc.*, 1965, 5414
2. Spear, R. J., *Aust. J. Chem.*, 1982, **35**, 4
3. Spear, R. J. *et al.*, *Prop. Explos., Pyrotech.*, 1983, **8**, 95–98

The complex formed between 4,6-dinitrobenzofurazan 3-oxide and potassium hydroxide has been evaluated as an explosive [1]. The structure is now believed to be that of a Meisenheimer complex [2]. The analogous sodium, silver and barium salts are also explosive, the latter being less sensitive to initiation [3].

See other FURAZAN *N*-OXIDES, POLYNITROARYL COMPOUNDS

2115. Sodium 2,4-dinitrophenoxide
[38892-09-0] $C_6H_3N_2NaO_5$




1. Hickson, B., *Chem. Age*, 1926, **14**, 522
2. Uhlmann, P. W., *Chem. Ztg.*, 1914, **38**, 388–390

A violent explosion occurred during centrifugal dehydration of an aqueous paste of the sodium salt [1], which was known to be explosive [2]. The instability may be associated with the presence of some of the *aci-o*- or *p*-quinonoid salt.

See other METAL NITROPHENOXIDES, NITROAROMATIC-ALKALI HAZARDS, POLYNITROARYL COMPOUNDS

2116. 1,3,5-Trinitrobenzene
[99-35-4] $C_6H_3N_3O_6$



Alkylmetallates, or Arylmetallates

Artamkina, G. M. *et al.*, *J. Organomet. Chem.*, 1979, **182**, 185

The THF-containing complexes formed between trinitrobenzene and the lithium or potassium salts of trimethyl-, triethyl- or triphenyl-germanate, -silanate or -stannate decompose explosively on heating, though trinitrobenzene-potassium trimethylstannate decomposes explosively at ambient temperature.

Methanol, Potassium hydroxide


See Potassium 4-methoxy-1-*aci*-nitro-3,5-dinitro-2,5-cyclohexadienide

See other POLYNITROARYL COMPOUNDS

2117. Trinitrosophloroglucinol (2,4,6-Trinitrosobenzene-1,3,5-triol)

[]

$C_6H_3N_3O_6$



The isomeric cyclohexanehexone 1,3,5-trioxime is identified as [14378-99-5].

Heavy metal compounds

Heavy metal salts (possibly of the oxime form) may be hazardous.


See Lead(II) trinitrosobenzene-1,3,5-trioxide

See other NITROSO COMPOUNDS See related OXIMES

2118. Picric acid (2,4,6-Trinitrophenol)

[88-89-1]

$C_6H_3N_3O_7$



NSC 351, 1979; FPA H128, 1983; HCS 1980, 755; RSC Lab. Hazards Data Sheet No. 34, 1985

Alone, or Heavy metals, or Bases

1. Cooper-Key, A., *Home Office Rept. 211*, 9, HMSO, 1914
2. Kirk-Othmer, 1965, Vol. 8, 617
3. Urbanski, 1964, Vol. 1, 518
4. Garey, H. E., *Chem. Eng. News*, 1979, **57**(41), 51
5. Ventrone, T. A., *CHAS Notes*, 1982, **1**(3), 1-2; (4), 4

Picric acid, in common with several other polynitrophenols, is an explosive material in its own right and is usually stored as a water-wet paste. Several dust explosions of dry material have been reported [1]. It forms salts with many metals, some of which (lead, mercury, copper or zinc) are rather sensitive to heat, friction or impact. The salts with ammonia and amines, and the molecular complexes with aromatic hydrocarbons, etc. are, in general, not so sensitive [2]. Contact of picric acid with concrete floors may form the friction-sensitive calcium salt [3]. Contact of molten picric acid with metallic zinc or lead forms the metal picrates which can detonate the acid. Picrates of lead, iron, zinc, nickel, copper, etc. should be considered dangerously sensitive. Dry picric acid has little effect on these metals at ambient temperature. Picric acid of sufficient purity is of the same order of stability as TNT, and is not considered unduly hazardous in regard to sensitivity [4]. Details of handling and disposal procedures have been collected and summarised [5].

See Sodium picrate, *also* Ammonium picrate

Aluminium, Water

See Aluminium: Nitro compounds, etc.


Uronium perchlorate

See Uronium perchlorate: Organic materials

See other ORGANIC ACIDS, POLYNITROARYL COMPOUNDS

2119. Trinitroresorcinol (2,4,6-Trinitrobenzene-1,3-diol)
[82-71-3]

C₆H₃N₃O₈




Sorbe, 1968, 152

The compound ('styphnic acid', +6.7% oxygen balance) and its salts are very explosive.

See other ORGANIC ACIDS, POLYNITROARYL COMPOUNDS

2120. Trinitrophenol (2,4,6-Trinitrobenzene-1,3,5-triol)
[4328-17-0]

C₆H₃N₃O₉




1. Sorbe, 1968, 153

2. DeFusco, A. A., *Org. Prep. Proced. Int.*, 1982, **14**, 393–395

It explodes on heating [1], and an improved preparation is described [2]. The compound has +20% oxygen balance, so can function as an overall oxidant. It would also be expected to form explosive metal salts.

See other POLYNITROARYL COMPOUNDS

2121. N,2,3,5-Tetranitroaniline
[57284-58-7]



1. Mudge, P. R. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1975, 569

2. Meldola, R. *et al.*, *J. Chem. Soc.*, 1905, **87**, 1204, 1205

Thermal rearrangement to the internal salt, 3,5-dinitrobenzenediazonium 2-oxide, proceeds explosively in absence of solvent [1]. The oxide itself explodes violently at 190°C [2].

See other N-NITRO COMPOUNDS, POLYNITROARYL COMPOUNDS

2122. 2,3,4,6-Tetranitroaniline
[3698-54-2]




Preparative hazard

See 3-Nitroaniline: Nitric acid, Sulfuric acid

See other POLYNITROARYL COMPOUNDS

2123. N,2,4,6-Tetranitroaniline
[4591-46-2]




Olsen, R. E. *et al.*, *ACS* 54, 1966, 50

Impure tetryl must be recrystallised with care as it may deflagrate at only 50°C.

See other N-NITRO COMPOUNDS, POLYNITROARYL COMPOUNDS

2124. Triethynylphosphine
[687-80-9]

C₆H₃P




Voskuil, W. *et al.*, *Rec. Trav. Chim.*, 1964, **83**, 1301

It explodes on strong friction, and on standing it decomposes and may then explode spontaneously.

See related ALKYLPHOSPHINES *See other* ACETYLENIC COMPOUNDS

2125. Triethynylantimony
[687-81-0]

C₆H₃Sb




Voskuil, W. *et al.*, *Rec. Trav. Chim.*, 1964, **83**, 1301

It explodes on strong friction.

See other METAL ACETYLIDES

2126. Silver 4-nitrophenoxide
[86255-25-6]

C₆H₄AgNO₃




Spiegel, L. *et al.*, *Ber.*, 1906, **39**, 2639

A sample decomposed explosively after intensive drying at 110°C. The *aci*-nitro-*p*-quinonoid salt may have been formed.

See related aci-NITROQUINONOID COMPOUNDS

See other METAL NITROPHENOXIDES, SILVER COMPOUNDS

2127. Silver benzo-1,2,3-triazole-1-oxide
[93693-72-2]




Deorha, D. S. *et al.*, *J. Indian Chem. Soc.*, 1964, **41**, 793

It is unsuitable as a gravimetric precipitate, as it explodes on heating rather than smoothly giving weighable silver.


See other N-O COMPOUNDS, SILVER COMPOUNDS, TRIAZOLES

2128. 2-Nitrobenzenediazonium tetrachloroborate
[]



See entry DIAZONIUM TETRAHALOBORATES


2129. 3-Bromophenyllithium
[6592-86-1]



See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2130. 4-Bromophenyllithium
[22480-64-4]

C₆H₄BrLi




Anon., *Angew. Chem. (Nachr.)*, 1962, **10**, 65

It explodes if traces of oxygen are present in the inert atmosphere needed for the preparation.

See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2131. 4-Bromobenzenediazonium salts
[17333-82-2] (ion)

C₆H₄BrN₂Z




Hydrogen sulfide

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 2) *See other* DIAZONIUM SALTS

2132. 2-Bromo-4,6-dinitroaniline
[1817-73-8]

C₆H₄BrN₃O₄




This and the 4-bromo-2,6-dinitro analogue show local decomposition on moderate heating.

See entry HIGH RATE DECOMPOSITION

2133. 4-Bromo-2,6-dinitroaniline
[62554-90-9]

$C_6H_4BrN_3O_4$




This and the 2-bromo-4,6-dinitro analogue show local decomposition on moderate heating.

See entry HIGH RATE DECOMPOSITION

†**2134. 2-Chlorofluorobenzene**
[348-51-6]


C_6H_4ClF



See other HALOARYL COMPOUNDS

†**2135. 3-Chlorofluorobenzene**
[625-98-9]


C_6H_4ClF



See other HALOARYL COMPOUNDS


†**2136. 4-Chlorofluorobenzene**
[352-33-0]

C_6H_4ClF




See other HALOARYL COMPOUNDS

2137. 4-Chlorobenzenediazonium triiodide
[68596-93-0]




See entry DIAZONIUM TRIIODIDES

2138. 3-Chlorophenyllithium
[25077-87-6]



See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2139. 4-Chlorophenyllithium
[14774-78-8]




Anon., *Angew. Chem. (Nachr.)*, 1962, **10**, 65

It explodes if traces of oxygen are present in the inert atmosphere needed for the preparation.

See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2140. 1,4-Benzoquinone-4-chloroimine
[637-61-6]

C₆H₄ClNO




Sorbe, 1968, 112

It explodes on heating, like the bis-chloroimine.

See other N-HALOGEN COMPOUNDS

2141. 2-Chloronitrobenzene
[88-73-3]

C₆H₄ClNO₂



Energy of decomposition (in range 350–450°C) measured as 1.83 kJ/g by DSC, and T_{ait24} was determined as 264°C by adiabatic Dewar tests, with an apparent energy of activation of 173 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkalis, Alcohols

See 4-Nitrochlorobenzene, next below

Ammonia

1. MCA Case History No. 1624
2. Vincent, G. C., *Loss. Prev.*, 1971, **5**, 46–52
3. Jacobson, O. F., *Proc. 3rd Int. Sympos. Loss Prev. Safety Prom. Proc. Ind.*, 417–421, SSCI, Basle, 1982
4. Gordon, M. D. *et al.*, *Plant/Oper. Progr.*, 1982, **1**, 27–33

During the large-scale preparation of 2-nitroaniline at 160–180°C/30–40 bar in a jacketed autoclave, several concurrent processing abnormalities (excess chloro compound, too little ammonia solution, failure to apply cooling or to vent the autoclave and non-failure of a rupture-disk) led to a runaway reaction and pressure-explosion of the vessel [1]. A full analysis of the incident is published [2]. A retrospective hazard and operability study revealed that 6 simultaneous fault conditions had been involved [3], and the thermal stability of the amination process for manufacturing 2-nitroaniline was studied using mathematical modelling techniques [4]. The results of modelling the process abnormalities closely followed the actual events [1,2].

See other AMINATION INCIDENTS

Methanol, Sodium hydroxide

Klais, O. *et al.*, *Prax. Sicherheitstech.*, 1997, **4**, 41

Klais, O. *et al.* in *Safety and Runaway Reactions*, Smeder, B & Mitchison, N. (Eds.), EUR 17723 EN, European Commission, 1997

A very large scale reaction to produce 2-nitroanisole from chloronitrobenzene and methanolic sodium hydroxide ran out of control and painted the town orange. This was attributed to reduction of the nitro group at temperatures above 100°C, a far more exothermic reaction than intended. This temperature was reached because the methanolic alkali was charged without agitation, and, the reaction not starting, the batch was heated to 90°C. The agitation was only then switched on.

See also 4-Nitrochlorobenzene, next below


See other AGITATION INCIDENTS

See other HALOARYL COMPOUNDS, NITROARYL COMPOUNDS

2142. 4-Chloronitrobenzene

[100-00-5]

$C_6H_4ClNO_2$



HCS 1980, 308

1. Doyle, W. H., *Loss. Prev.*, 1969, **3**, 16

A steam-heated still used to top the crude material exploded

Energy of decomposition (in range 300–450°C) measured as 2.05 kJ/g by DSC, and T_{ait24} was determined as 275°C by adiabatic Dewar tests, with an apparent energy of activation of 172 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkalis, Alcohols

Anderson, H. L. *et al.*, *Thermochim. Acta*, 1996, 271, 195

Anderson, H. L. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries*, Vol II, (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 387, Amsterdam, Elsevier, 1995

A calorimetric study of reaction with sodium or potassium hydroxides in ethanol or 2-propanol is given. At starting temperatures below 70°C the product is the appropriate nitrophenyl ether; above that temperature, reduction of the nitro groups may come into play, to give much more energy and a variety of other products. This reaction is inhibited by oxygen. There is potential for runaway if such reactions are operated industrially with poor temperature control. The editor suspects that the stimulus for this study was an accident which sprayed the German environment with 2-nitroanisole.

See Sodium methoxide; below

Potassium hydroxide

Mixtures with potassium hydroxide (1:1.5 mol) deflagrate readily at a rate of 1.3 cm/min.

See entries PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 3), NITROAROMATIC-ALKALI HAZARDS

See other DEFLAGRATION INCIDENTS

Sodium methoxide

Anon., *ABCM Quart. Safety Summ.*, 1944, **15**, 15

Addition of the chloro compound to a solution of sodium methoxide in methanol caused an unusually exothermic reaction to occur. The lid of the 450 l vessel was blown off, and a fire and explosion followed. No cause for the unusual vigour of the reaction was found.


See Alkalis, Alcohols; above

See other HALOARYL COMPOUNDS, NITROARYL COMPOUNDS

2143. 3-Nitroperchlorylbenzene

[20731-44-6]

$C_6H_4ClNO_5$



McCoy, G., *Chem. Eng. News*, 1960, **38**(4), 62


The nitration product of perchloryl benzene is explosive, comparable in shock-sensitivity with lead azide, with a very high propagation rate.

See other NITROARYL COMPOUNDS, PERCHLORYL COMPOUNDS

2144. 2-Chloro-5-nitrobenzenesulfonic acid

[96-73-1]

$C_6H_4ClNO_5S$



Preparative hazard

Exothermic decomposition of the acid at 150°C ($Q = 0.708$ kJ/g) is time-delayed, may be faster in a closed than an open system, and is dangerous.

See Sulfuric acid: 4-Chloronitrobenzene


See entry SELF-ACCELERATING REACTIONS

See other HALOARYL COMPOUNDS, NITROARYL COMPOUNDS, ORGANIC ACIDS

2145. 4-Chloro-3-nitrobenzenesulfonic acid

[71799-32-1]

$C_6H_4ClNO_5S$



Négyesi, G., *Process Safety Progress*, 1996, **15**(1), 42


This compound shows signs of impact and friction sensitivity.

See other NITROARYL COMPOUNDS

2146. 4-Hydroxy-3-nitrobenzenesulfonyl chloride

[]

$C_6H_4ClNO_5S$



Vervalin, C. H., *Hydrocarbon Proc.*, 1976, **55**(9), 321–322


The chloride, produced by interaction of 2-nitrophenol and chlorosulfuric acid at 4°C, decomposed violently during discharge operations from the 2000 l vessel, leaving a glowing residue. It was subsequently found that accelerating exothermic decomposition sets in at 24–27°C.

See related SULFONATION INCIDENTS See other NITROACYL HALIDES

2147. Dicarbonylpyrazinerhodium(I) perchlorate

[]

$C_6H_4ClN_2O_6Rh$



Uson, R. *et al.*, *J. Organomet. Chem.*, 1981, **220**, 109


This (notably) and other polynuclear complexes exploded violently on heating during microanalysis.

See related AMMINEMETAL OXOSALTS

2148. 2-Chlorobenzenediazonium salts

[17333-83-4] (ion)

$C_6H_4ClN_2Z$




Potassium 2-chlorothiophenoxide

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 9) *See other* DIAZONIUM SALTS

2149. 3-Chlorobenzenediazonium salts

[17333-84-5] (ion)

$C_6H_4ClN_2Z$




Potassium thiophenoxide, or Sodium polysulfide

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 9) *See other* DIAZONIUM SALTS

2150. 1-Chlorobenzotriazole

[21050-95-3]

$C_6H_4ClN_3$




Hopps, H. B., *Chem. Eng. News*, 1971, **49**(30), 3

Spontaneous ignition occurred during packing operations.

See other N-HALOGEN COMPOUNDS


2151. 3-Nitrobenzenediazonium chloride
[2028-76-4]



Potassium *O*-ethyldithiocarbamate

See DIAZONIUM SULFIDES AND DERIVATIVES *See other* DIAZONIUM SALTS

2152. 4-Chlorobenzenesulfonyl azide
[4547-68-6]




N-Methyltetrahydro- β -carboline

Bailey, A. S. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1980, 1513

In absence of solvent chloroform, interaction at ambient temperature is explosive.

See other ACYL AZIDES

2153. 2-Chloro-4,6-dinitroaniline
[3531-19-9]



The solid amine can be caused to detonate by heating, or by a powerful initiating charge, and fast local decomposition occurs on heating moderately.

See entry HIGH RATE DECOMPOSITION

Nitrosylsulfuric acid

Bersier, P. *et al.*, *Chem. Ing. Tech.*, 1971, **43**, 1311–1315


Diazotisation of the potentially explosive 2-chloro-4,6-dinitroaniline in the third batch of a new process (at a higher than usual concentration in 40% nitrosylsulfuric acid) led to a violent explosion soon after the temperature had been increased to 50°C. Subsequent DSC work showed that the temperature at which thermal decomposition of the diazonium sulfate solution or suspension sets in is inversely proportional to the concentration of amine, falling from 160°C at 0.3 mmol/g to 80°C at 2 mmol/g. Thermal stability of 17 other diazonium derivatives was similarly investigated.

See other DIAZONIUM SULFATES See other POLYNITROARYL COMPOUNDS

2154. 4-Chloro-2,6-dinitroaniline

[5388-62-5]

$C_6H_4ClN_3O_4$



CHETAH, 1990, 183

Can be caused to detonate by 50 g tetryl booster.

Preparative hazard

See Nitric acid: 4-Chloro-2-nitroaniline

Nitrosylsulfuric acid

Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 62


During large-scale diazotisation of the amine, severe local overheating is thought to have caused the explosion observed. The effect could not be reproduced in the laboratory.

See DIAZOTISATION See other POLYNITROARYL COMPOUNDS

2155. 3-Nitrobenzenediazonium perchlorate

22751-24-2

$C_6H_4ClN_3O_6$




Schumacher, 1960, 205

Explosive, very sensitive to heat or shock.
See other DIAZONIUM PERCHLORATES

2156. 1,2-Dichlorobenzene
[95-50-1]

$C_6H_4Cl_2$



Aluminium, Halocarbons

See Aluminium: Halocarbons (reference 5)

Other reactants


Yoshida, 1980, 156

MRH values calculated for 13 combinations with oxidants are given.

See other HALOARYL COMPOUNDS

2157. 1,6-Dichloro-2,4-hexadiyne
[16260-59-6]

$C_6H_4Cl_2$




1. Driedger, P. E. *et al.*, *Chem. Eng. News*, 1972, **50**(12), 51
2. Whiting, M. C., *Chem. Eng. News*, 1972, **50**(23), 86
3. Ford, M *et al.*, *Chem. Eng. News*, 1972, **50**(30), 67
4. Walton, D. R. M., private comm., 1987
5. Anon., *Inst. Univ. Saf. Officers Bull.*, 1987, (8), 9

It is extremely shock-sensitive, a 4.0 kg cm shock causing detonation in 50% of test runs (cf. 3.5 kg cm for propargyl bromide; 2.0 kg cm for glyceryl nitrate). The intermediate bis-chlorosulfite involved in the preparation needs low temperatures to avoid vigorous decomposition. The corresponding diiodo derivative was expected to be similarly hazardous [1], and this has been confirmed [2]. Improvements in preparative techniques (use of dichloromethane solvent at $-30^{\circ}C$) to avoid violent reaction have also been described [3]. An attempt to distill the compound (b.p. $55-58^{\circ}C/0.6$ mbar, equivalent to about $230^{\circ}C/1$ bar) at atmospheric pressure from a heating mantle led to a violent explosion [4]. The compound involved was erroneously given as 1,6-dichloro-2,4-hexadiene [5].

See other HALOACETYLENE DERIVATIVES

2158. 4-Nitrophenylphosphorodichloridate
[777-52-6]




Frommer, M. A. *et al.*, 1990, US Pat. 4910190; *Chem. Abs.*, 1991, **114**, 122705b

This compound may decompose violently during distillation.

See other NITROARYL COMPOUNDS

See related NON-METAL HALIDES (AND THEIR OXIDES)

2159. Benzoquinone 1,4-bis(chloroimine) (1,4-Bis(chlorimido)-2,5-cyclohexadiene)
[637-70-7]





Sorbe, 1968, 112

It explodes on heating, like the mono-chloroimine.

See other N-HALOGEN COMPOUNDS

2160. Benzene-1,4-bis(diazonium perchlorate)
[43008-25-9]




Hofman, K. A. *et al.*, *Ber.*, 1910 **43**, 2624

Described by Hofman as the most powerfully explosive compound of which he knew. Since it is of better oxygen balance than the aromatic high explosives, and the diazonium unit is also high energy, this is quite probable.

See DIAZONIUM PERCHLORATES

2161. Dichlorophenol mixed isomers
[25167-81-1]





HCS 1980, 381 (2,4-Dichlorophenol)

Laboratory Chemical Disposal Co. Ltd., confidential information, 1968

During vacuum fractionation of the mixed dichlorophenols produced by partial hydrolysis of trichlorobenzene, rapid admission of air to the receiver caused the column contents to be forced down into the boiler at 210°C, and a violent explosion ensued.

See other HALOARYL COMPOUNDS

2162. 2,4-Hexadiynylene chlorosulfite
[]




Driedger, P. E. *et al.*, *Chem. Eng. News*, 1972, **50**(12), 51

This is probably an intermediate in the preparation of the 1,6-dichloro compound from 2,4-hexadiyne-1,6-diol and thionyl chloride in DMF. The reaction mixture must be kept at a low temperature to avoid vigorous decomposition and charring. This may be due to interaction with the solvent.

See 2,4-Hexadiynylene chloroformate also Dimethylformamide

See other ACETYLENIC COMPOUNDS, ACYL HALIDES

2163. 2,3,4-Trichloroaniline
[634-67-3]




Energy of decomposition (in range 220–400°C) measured as 0.21 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other HALOANILINES

2164. Cobalt(II) chelate of 1,3-bis(*N*-nitrosohydroxylamino)benzene
[27662-11-9]


C₆H₄CoN₄O₄



Battei, R. S. *et al.*, *J. Inorg. Nucl. Chem.*, 1970, **32**, 1525
The cobalt(II) chelate of 'dicupferron' explodes on heating at 205°C.
See Cobalt(II) chelate of bi(1-hydroxy-3,5-diphenylpyrazol-4-yl *N*-oxide)
See other N-O COMPOUNDS

2165. Dicopper(I) 1,5-hexadiynide
[86425-12-9]


C₆H₄Cu₂



Chow, Y. L. *et al.*, *Can. J. Chem.*, 1983, **61**, 795–800
It exploded when dried in a desiccator.
See other METAL ACETYLIDES


2166. 4-Fluorophenyllithium
[1493-23-8]

C₆H₄FLi



See entry ORGANOLITHIUM REAGENTS See other HALO-ARYLMETALS

2167. 5-Fluoro-2-nitrophenol
[446-36-6]




Preparative hazard

Robinson, N., *Chem. Brit.*, 1987, **23**, 837

Following a patented procedure for the conversion of 2,4-dinitrochlorobenzene to 5-chloro-2-nitrophenol, 2,4-difluoronitrobenzene was treated with sodium hydroxide in hot aqueous dioxane containing a phase transfer catalyst. On the small scale, the reaction and isolation of 5-fluoro-2-nitrophenol, including vacuum distillation, were uneventful. On the 20 l scale, vacuum distillation of combined batches of the crude product led to onset of decomposition at 150°C, which could not be controlled, and the residue erupted with explosive violence and a small fire ensued. Thermal examination of fresh small-scale crude material has shown that it is capable of highly exothermic decomposition, with onset of the exotherm at 150°C (ARC). It was then realised that difficulty in controlling the reaction temperature had been experienced on the 20 l scale. It is recommended that this procedure and purification should not be attempted on so large a scale.


See other NITROARYL COMPOUNDS

†2168. 1,3-Difluorobenzene
[372-18-9]





See other HALOARYL COMPOUNDS

†2169. 1,4-Difluorobenzene
[540-36-3]



See other HALOARYL COMPOUNDS

2170. Octafluoroadipamide
[355-66-8]




Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Fluoroamides

2171. Sodium 2-hydroxymercurio-4-*aci*-nitro-2,5-cyclohexadienonide

[]




Hantzsch, A. *et al.*, *Ber.*, 1906, **39**, 1115

This salt of the *aci-p*-quinonoid form of 2-hydroxymercurio-4-nitrophenol explodes on heating.

See other MERCURY COMPOUNDS, *aci*-NITROQUINONOID COMPOUNDS

2172. 1,2-Diiodobenzene
[615-42-9]




Rüst, 1948, 302

A small sample in a sealed glass capillary exploded violently at 181°C, breaking the surrounding oil bath.

See other HALOARYL COMPOUNDS, IODINE COMPOUNDS

2173. 1,6-Diiodo-2,4-hexadiyne
[44750-17-6]


$C_6H_4I_2$



See 1,6-Dichloro-2,4-hexadiyne
See other HALOACETYLENE DERIVATIVES

2174. Potassium 4-nitrophenoxide
[1124-31-8]


$C_6H_4KNO_3$



1. L. Bretherick, Personal experience
 2. Thorpe's Dictionary of Applied Chemistry; 4th Edn., London, 1940, **IV**, 475
- A sample of this potassium salt exploded after long storage in a non-evacuated desiccator [1]. This may have been caused by formation of some tautomeric *aci*-quinonoid salt by prolonged desiccation. This compound had earlier been described as exploding on percussion [2].
See 4-Nitrophenol: Potassium hydroxide
See other METAL NITROPHENOXIDES, NITROARYL COMPOUNDS, *aci*-NITROQUINONOID COMPOUNDS

2175. Potassium 4-nitrobenzeneazosulfonate
[]

$C_6H_4KN_3O_5S$




491M, 1975, 345

During examination of a 10 g batch of the crystalline material on a filter paper using a hand lens, it exploded. This was attributed to sudden rearrangement of the

metastable syn- to the anti-form (perhaps triggered by light energy concentrated by the lens).

See other AZO COMPOUNDS, N-S COMPOUNDS

2176. Potassium 4-hydroxyamino-5,7-dinitro-4,5-dihydrobenzofurazanide 3-oxide
[86341-95-9] **C₆H₄KN₅O₇**




Norris, W. P. *et al.*, *Chem. Abs.*, 1983, **99**, 40662

Sensitivity to mechanical, electrostatic and thermal shock is typical of a primary explosive.

See other FURAZAN N-OXIDES

2177. Lithium 4-nitrothiophenoxide
[78350-94-4] **C₆H₄LiNO₂S**



1. Julia, M., *Chem. & Ind.*, 1981, 130; *Chem. Eng. News*, 1981, **59**(14), 3


2. Bretherick, L., *Chem. & Ind.*, 1981, 202; *Chem. Eng. News*, 1981, **59**(31), 3

A vacuum dried sample of the salt exploded violently when exposed to air [1]. It was postulated that the *aci*-quinonoid species may have been the unstable species [2].

See other NITROAROMATIC-ALKALI HAZARDS, *aci*-NITROQUINONOID COMPOUNDS



See related METAL NITROPHENOXIDES

2178. 1,3-Dilithiobenzene
[2592-85-0] **C₆H₄Li₂**




See entry ORGANOLITHIUM REAGENTS See other ARYLMETALS

2179. 1,4-Dilithiobenzene
[7279-42-0]




See entry ORGANOLITHIUM REAGENTS See other ARYLMETALS

2180. N,N,4-Trilithioaniline
[]




Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 2934
Highly explosive in contact with air.
See other N-METAL DERIVATIVES, ORGANOMETALLICS

2181. Sodium 4-nitrosophenoxide
[823-87-0]



Sorbe, 1968, 86
The dry solid tends to ignite.
See other NITROSO COMPOUNDS


2182. Sodium 2-nitrothiophenoxide
[22755-25-5]



Davies, H. J., *Chem. & Ind.*, 1966, 257

The material exploded when the temperature of evaporation of the slurry was increased by adding a higher-boiling solvent. This may have been caused by presence of some of the tautomeric *aci*-nitro-*o*-thioquinone. Such salts are thermally unstable. See other NITROAROMATIC-ALKALI HAZARDS, *aci*-NITROQUINONOID COMPOUNDS. See related METAL NITROPHENOXIDES.

2183. Sodium 4-nitrophenoxide
[824-78-2]




1. Semiganowski, N., *Z. Anal. Chem.*, 1927, **72**, 29

2. Baltzer, private comm., 1976

An ammoniacal solution of 4-nitroaniline containing ammonium chloride was being treated with 50% sodium hydroxide solution to displace ammonia. In error, double the amount required to give the usual 10% of free alkali was added, and during the subsequent degassing operation (heating to 130°C under pressure, followed by depressuring to vent ammonia), complete conversion to sodium 4-nitrophenoxide occurred unwittingly [1]. This solid product, separated by centrifuging, was then heated to dry it, when it decomposed violently and was ejected through the vessel opening like a rocket [2]. Some of the tautomeric *aci*-nitroquinonoid salt may have been produced during the drying operation.

See other METAL NITROPHENOXIDES, NITROAROMATIC-ALKALI HAZARDS, *aci*-NITROQUINONOID COMPOUNDS

2184. Sodium 3-nitrobenzenesulfonate
[127-68-4]




1. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987
2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

It ignites at 335°C and the deflagration rate of the solid is 4.5 cm/min [1]. T_{ait24} was determined as 305°C by adiabatic Dewar tests, with an apparent energy of activation of 441 kJ/mol [2].

See 3-Nitrobenzenesulfonic acid

See other DEFLAGRATION INCIDENTS, NITROARYL COMPOUNDS

2185. Disodium 4-nitrophenylphosphate
[4264-83-9]




Preparative hazard

See Nitric acid: Disodium phenyl orthophosphate

See other NITROARYL COMPOUNDS

2186. Thallium(I) 2-nitrophenoxide
[32707-02-1]



Langhans, A., *Z. Ges. Schiess.-Sprengstoff W.*, 1936, **31**, 359, 361

Both solids explode on heating.

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, NITROARYL COMPOUNDS

2187. Thallium(I) 4-nitrophenoxide
[32707-03-2]

$C_6H_4NO_3TI$

$TlOC_6H_4NO_2$


Langhans, A., *Z. Ges. Schiess.-Sprengstoff W.*, 1936, **31**, 359, 361

Both solids explode on heating.

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, NITROARYL COMPOUNDS

2188. Sodium 1,4-bis(*aci*-nitro)-2,5-cyclohexadienide
[]

$C_6H_4N_2Na_2O_4$




Meisenheimer, J. *et al.*, *Ber.*, 1906, **39**, 2529

The sodium and potassium salts of the bis-*aci*-quinonoid form of 1,4-dinitrobenzene deflagrate on heating.

See other aci-NITROQUINONOID COMPOUNDS

2189. Benzenediazonium-4-oxide
[6925-01-5]

$C_6H_4N_2O$




Hantzsch, A. *et al.*, *Ber.*, 1896, **29**, 1530

It decomposes violently at 75°C.

See other ARENEDIAZONIUM OXIDES

2190. Benzofurazan-N-oxide
[480-96-6]


$C_6H_4N_2O_2$



CHETAH, 1990, 188
Shock sensitive (probably not very).
See other FURAZAN N-OXIDES

2191. Benzo-1,2,3-thiadiazole 1,1-dioxide
[37150-27-9]


C₆H₄N₂O₂S



1. Wittig, G. *et al.*, *Chem. Ber.*, 1962, **95**, 2718
 2. Wittig, G. *et al.*, *Org. Synth.*, 1967, **47**, 6
- The solid explodes at 60°C, on impact or friction, or sometimes spontaneously. Solutions decompose below 10°C to give benzyne [1]. Full preparative and handling details are given [2].
See other N-S COMPOUNDS

2192. Benzenediazonium-2-sulfonate
[612-31-7]


C₆H₄N₂O₃S



Franklin, E. C., *Amer. Chem. J.*, 1899, **20**, 459
The internal diazonium salt explodes on contact with flame or on percussion.
See related DIAZONIUM CARBOXYLATES See other DIAZONIUM SALTS

2193. Benzenediazonium-4-sulfonate
[305-80-6]

C₆H₄N₂O₃S




1. Wichelhaus, H., *Ber.*, 1901, **34**, 11
2. Anon., *Sichere Chemiearb.*, 1959, **11**, 59
3. Anon., *Sichere Chemiearb.*, 1978, **30**, 77

This internal salt of diazotised sulfanilic acid (Pauly's reagent) exploded violently on touching when thoroughly dry [1]. Use of a metal spatula to remove a portion of a refrigerated sample of the solid salt caused a violent explosion [2], and a similar incident was reported 19 years later [3]. All solid diazo compounds must be stored only in small quantity under refrigeration in loosely plugged containers. Handle gently with non-metallic spatulae using personal protection [2].

See related DIAZONIUM CARBOXYLATES *See other* DIAZONIUM SALTS

2194. 1,2-Dinitrobenzene
[528-29-0]




Nitric acid

See Nitric acid: Nitroaromatics

See other POLYNITROARYL COMPOUNDS

2195. 1,3-Dinitrobenzene
[99-65-0]



Anon., *Sichere Chemiearbeit*, 1993, **45**(7), 82

Heat of decomposition was determined as 4.6 kJ/g (c.f. MRH calculated value of 5.02). $T_{\text{ait}24}$ was determined as 284°C by adiabatic Dewar tests, with an apparent energy of activation of 284 kJ/mol.

A vessel containing dinitrobenzene for drying, after washing out styphnic and picric acid byproducts in alkali, retained a heel of some hundreds of litres, with a residue of rust, after emptying. This was cleaned out with low pressure steam at 130°C every two years. During cleaning, the tank burst and burnt out. It was supposed that residual nitrophenol salts had accumulated on the rust, which was shown to thermally sensitise them, and deflagrated, initiating the dinitrobenzene.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (references 1,2)

Nitric acid

See Nitric acid: Nitroaromatics

Other reactants

Yoshida, 1980, 160

MRH values calculated for 7 combinations with materials catalysing its decomposition are given.

Tetranitromethane


See Tetranitromethane: Aromatic nitro compounds

See other POLYNITROARYL COMPOUNDS

2196. 1,4-Dinitrobenzene

[100-25-4]

$C_6H_4N_2O_4$



Nitric acid


See Nitric acid: Nitroaromatics

See other POLYNITROARYL COMPOUNDS

2197. 2,4-Dinitrophenol

[51-28-5]

$C_6H_4N_2O_5$



NSC 690, 1981; HCS 1980, 439

T_{ait24} was determined as 189°C by adiabatic Dewar tests, with an apparent energy of activation of 144 kJ/mol. Dinitrophenol is now classified as an explosive in the UK, and is normally available from laboratory suppliers wetted with 15% water, as is picric acid.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Bases

1. Uhlmann, P. W., *Chem. Ztg.*, 1914, **38**, 389–390

2. *Thorpe's Dictionary of Applied Chemistry*; London, 1940, **IV**, 477

Dinitrophenol forms explosive (*aci*-quinonoid ?) salts with alkalis or ammonia, and should not be heated with them in closed vessels [1]. Dinitrophenol forms explosive salts with most metals [2].

See Sodium 2,4-dinitrophenoxide


See other NITROAROMATIC–ALKALI HAZARDS, *aci*-NITROQUINONOID COMPOUNDS

See other POLYNITROARYL COMPOUNDS

2198. 4,6-Dinitro-1,3-benzenediol (4,6-Dinitroresorcinol)

[616-74-0]

$C_6H_4N_2O_6$



Preparative hazard

Schmitt, R. J. *et al.*, *J. Org. Chem.*, 1988, **53**, 5568–5569


The reaction conditions necessary to obtain a good yield of the title compound (a difficult isomer), and to avoid hazards during the nitration of resorcinol, are critical and strict adherence to those specified is essential. The necessary 80% white fuming nitric acid must be completely free from oxides of nitrogen and nitrous acid, and procedures for this are detailed. Then the temperature during addition of the diacetate must be kept between -10 and 0°C by regulating the rate of addition. The alternative use of 80% sulfuric acid as solvent for the 80% nitric acid (5 equiv.) is preferred as more reliable, but both methods have led to violent exothermic decomposition, accompanied by fume-off, after an induction period. In any event, the explosive 2,4,6-trinitroresorcinol ('styphnic acid') is produced as a by-product.

See other INDUCTION PERIOD INCIDENTS, NITRATION INCIDENTS, POLYNITROARYL COMPOUNDS

2199. Benzo-1,2,3-thiadiazole

[273-77-8]

$C_6H_4N_2S$



Jordis, U. *et al.*, *Phos. Sulfur*, 1984, **19**, 279


Purification by extraction and then distillation led to violent decomposition. Steam distillation avoided this problem.

See other N-S COMPOUNDS

2200. Sodium 6-*aci*-nitro-4-nitro-2,4-cyclohexadieniminide

[]

C₆H₄N₃NaO₄



Green, A. G. *et al.*, *J. Chem. Soc.*, 1913, **103**, 508–513


The sodium and potassium salts of the mono-*aci*-*o*-quinonoid form of 2,4-dinitroaniline readily deflagrate on heating.

See other *aci*-NITROQUINONOID COMPOUNDS, POLYNITROARYL COMPOUNDS

2201. 2-Nitrobenzenediazonium salts

[25910-37-6] (ion)

C₆H₄N₃O₂Z




Sodium disulfide, or Sodium polysulfide

See DIAZONIUM SULFIDES AND DERIVATIVES

2202. 4-Nitrobenzenediazonium salts

[14368-49-1] (ion)


C₆H₄N₃O₂Z



Hydrogen sulfide, or Sodium sulfide, or Potassium *O, O*-diphenyl phosphorodithioate

See DIAZONIUM SULFIDES AND DERIVATIVES

2203. Sodium 3,5-bis(*aci*-nitro)cyclohexene-4,6-diiminide
[]




Alone, or Water

Meisenheimer, J. *et al.*, *Ber.*, 1906, **39**, 2538

The disodium salt of the bis-*aci-o*-quinonoid form of 2,4-dinitro-1,3-phenylenediamine deflagrates violently in contact with heat or moisture.


See other aci-NITROQUINONOID COMPOUNDS

2204. 4-Nitrobenzenesulfinyl azide
[53852-44-1]



See entry SULFINYL AZIDES See other ACYL AZIDES

2205. 4-Nitrobenzenediazonium nitrate
[42238-29-9]




Bamberger, E., *Ber.*, 1895, **28**, 239

The isolated dry salt explodes on heating, but not on friction.

See other DIAZONIUM SALTS

2206. 2,4-Dinitrobenzenediazonium hydrogen sulfate
[64445-49-4]


C₆H₄N₄O₈S



See entry DIAZONIUM SULFATES See other POLYNITROARYL COMPOUNDS

2207. 1,3-Diazidobenzene
[13556-50-8]

C₆H₄N₆



Acids


Forster, M. O. *et al.*, *J. Chem. Soc.*, 1907, **91**, 1953

The diazide ignites and explodes mildly with concentrated acid.

See other ORGANIC AZIDES

2208. 1,4-Diazidobenzene
[2294-47-5]

C₆H₄N₆




Griess, P., *Ber.*, 1888, **21**, 1561

The diazide explodes very violently on heating.

See other ORGANIC AZIDES

2209. 4-Nitrobenzenediazonium azide
[68560-64-3]


$C_6H_4N_6O_2$



Hantzsch, A., *Ber.*, 1903, **36**, 2058
The dry azide salt exploded violently with a brilliant flash.
See other DIAZONIUM SALTS, NON-METAL AZIDES

2210. Benzene-1,3-bis(sulfonyl azide)
[4547-69-7]


$C_6H_4N_6O_4S_2$



Balabanov, G. P. *et al.*, *Chem. Abs.*, 1969, **70**, 59427
The title compound was the most explosive of a series of 7 substituted benzenesulfonyl azides.
See other ACYL AZIDES

2211. 1,4-Di(1,2,3,4-thiaziazol-5-yloxy)but-2-yne
[]

$C_6H_4N_6O_2S_2$




Banert, K. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(7), 866

An explosive crystalline solid.
See other ACETYLENIC COMPOUNDS, N-S COMPOUNDS

2212. 2-Ethynylfuran
[18649-64-4]


C₆H₄O



Alone, or Nitric acid
Dunlop, 1953, 77
It explodes on heating or in contact with conc. nitric acid.
See other ACETYLENIC COMPOUNDS

2213. 1,5-Hexadiyn-3-one
[66737-76-7]


C₆H₄O



Dowd, P. *et al.*, *Synth. Commun.*, 1978, **8**, 20
The reactive material should be vacuum distilled at ambient temperature to prevent the near-explosive decomposition which occurs if it is heated.
See other ACETYLENIC COMPOUNDS

2214. 1,4-Benzoquinone (Cyclohexadiene-1,4-dione)
[106-51-4]

C₆H₄O₂



Anon., *Loss Prev. Bull.*, 1977, (013), 4
Drums containing moist quinone self-heated, generating smoke and internal pressure. Investigation showed that the moist material showed an exotherm at 40–50°C and decomposed at 60–70°C, possibly owing to presence of impurities. Drums exposed to direct sunlight soon attain a surface temperature of 50–60°C.
See other SELF-HEATING AND IGNITION INCIDENTS


Other reactants

Yoshida, 1980, 102

MRH values calculated for 13 combinations with oxidants are given.

2215. 2,2'-Bi-1,3-dithiole (1,4,5,8-Tetrahydro-1,4,5,8-tetrathiafulvalene)
[31366-25-3]

$C_6H_4S_2$




Preparative hazard

See 1,3-Dithiolium perchlorate

2216. 6-Fulvenoselone
[72443-10-8]

C_6H_4Se




Schulz, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1980, **19**, 69

It polymerises explosively at $-196^\circ C$.

See other POLYMERISATION INCIDENTS

2217. Phenylsilver
[5274-48-6]

C_6H_5Ag




1. Houben-Weyl, 1970, Vol. 13.1, 774

2. Krause, E. *et al.*, *Ber.*, 1923, **56**, 2064

The dry solid explodes on warming to room temperature, or on light friction or stirring [1]. It is more stable wet with ether [2].


See other ARYLMETALS, SILVER COMPOUNDS

2218. Silver phenoxide
[61514-68-9]




Macomber, R. S. *et al.*, *Synth. React. Inorg. Met.-Org. Chem.*, 1977, **7**, 111–122
The dried solid product from interaction of lithium phenoxide and silver perchlorate in benzene (probably largely silver phenoxide) exploded on gentle heating. Other silver alkoxide derivatives were unstable.
See Silver perchlorate
See other SILVER COMPOUNDS

2219. Silver benzeneselenonate
[39254-49-4]



See Benzeneselenonic acid
See other SILVER COMPOUNDS

2220. 4-Hydroxy-3,5-dinitrobenzenearsonic acid
[6269-50-7]



Phillips, M. A., *Chem. & Ind.*, 1947, 61


A heated, unstirred water-wet sludge of this analogue of picric acid exploded violently, apparently owing to local overheating in the containing flask.

See other ORGANIC ACIDS, POLYNITROARYL COMPOUNDS

2221. Phenylgold

[]

C₆H₅Au



De Graaf, P. W. J. *et al.*, *J. Organomet. Chem.*, 1977, **141**, 353


The impure solid at -70°C exploded when touched with a spatula.

See other ARYLMETALS, GOLD COMPOUNDS

2222. Dichlorophenylborane

[873-51-8]

C₆H₅BCl₂



Gerwanch, U. W. *et al.*, *Inorg. Synth.*, 1983, **22**, 207


It is pyrophoric when hot, and extremely sensitive to hydrolysis.

See related ALKYLHALOBORANES

2223. Benzenediazonium tetrafluoroborate

[369-57-3]

C₆H₅BF₄N₂




Caesium fluoride, Difluoroamine

See Difluoroamine: Benzenediazonium tetrafluoroborate, etc.

See other DIAZONIUM TETRAHALOBORATES

2224. Bromobenzene
[106-86-1]

C₆H₅Br




HCS 1980, 667

Bromobutane, Sodium
See Sodium: Halocarbons (reference 7)

Other reactants
Yoshida, 1980, 323
MRH values calculated for 13 combinations with oxidants are given.
See other HALOARYL COMPOUNDS

2225. Phenylmagnesium bromide
[100-58-3]

C₆H₅BrMg




Chlorine
See Chlorine: Phenylmagnesium bromide

Hexafluoropropene
See Hexafluoropropene: Grignard reagents


Water
Wosnick, J., INTERNET, 1996
An overconcentrated ether solution of the reagent was poured down a sink. Reaction was violent.
See other GRIGNARD REAGENTS

2226. 2,4-Dibromoaniline
[615-57-6]





Energy of decomposition (in range 150–310°C) measured as 0.24 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other HALOANILINES

2227. Benzenediazonium tribromide
[19521-84-7]



Sorbe, 1968, 40
An explosive compound.
See other DIAZONIUM SALTS

2228. Chlorobenzene
[108-90-7]




Other reactants
Yoshida, 1980, 113
MRH values calculated for 13 combinations, largely with oxidants, are given.

Phosphorus trichloride, Sodium
See Sodium: Chlorobenzene, etc.

Sodium
See Sodium: Halocarbons (reference 8)
See other HALOARYL COMPOUNDS

2229. Sodium *N*-chlorobenzenesulfonamide ('Chloramine B')
[127-52-6]


$C_6H_5ClNNaO_2S$



Piotrowski, T., *Chem. Abs.*, 1984, **101**, 60040
In a study of explosive potential of 20 pharmaceutical products, 'Chloramine B' was found to present the greatest fire hazard, decomposing explosively at 185°C.
See other *N*-HALOGEN COMPOUNDS, *N*-METAL DERIVATIVES

2230. Benzenediazonium chloride
[100-34-5]

$C_6H_5ClN_2$



1. Hantzsch, A. *et al.*, *Ber.*, 1901, **34**, 3338
 2. Grever, T., *Chem. Ing. Tech.*, 1975, **47**, 233–234
 3. *See* entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION
- The dry salt was more or less explosive, depending on the method of preparation [1]. DTA examination of a 15% solution of the diazonium salt in hydrochloric acid showed an exothermic decomposition reaction with the exotherm peak at 65°C. Adiabatic decomposition of the solution at 20°C took 80 min to attain the maximum

temperature of 80°C [2]. The heat of decomposition Q was determined as 1.5 kJ/g [3].
See Benzenediazonium tetrachlorozincate


Potassium *O*-methyldithiocarbonate

See Thiophenol

See also DIAZONIUM SULFIDES AND DERIVATIVES


See other DIAZONIUM SALTS

2231. *N*-Chloro-4-nitroaniline
[59483-61-3]



1. Goldschmidt, S. *et al.*, *Ber.*, 1922, **55**, 2450
 2. Pyetlewski, L. L., *Rept. AD A028841*, 24, Richmond (Va.), NTIS, 1976
- Though stable at -70°C , it soon decomposes explosively at ambient temperature [1], and violently on heating slightly [2].
See other *N*-HALOGEN COMPOUNDS, NITROARYL COMPOUNDS

2232. Benzenediazonium perchlorate
[114328-95-9]



Alone, or Sulfuric acid


1. Vorlander, D., *Ber.*, 1906, **39**, 2713–2715
2. Hofmann, K. A. *et al.*, *Ber.*, 1906, **39**, 3146–3148

The salt explodes on friction, impact or contact with sulfuric acid when still wet with liquor [1], and is frightfully explosive when dry [2].

See other DIAZONIUM PERCHLORATES

2233. Phenylphosphonic azide chloride

[]



Utvary, K., *Inorg. Nucl. Chem. Lett.*, 1965, **1**, 77–78



A violent explosion occurred during distillation at 80–88°C/0.46 mbar of mixtures containing the azide chloride and the diazide

See Phenylphosphonic diazide

See other ACYL AZIDES

2234. Benzenesulfinyl chloride

[4972-29-6]




Anon., *Chem. Eng. News*, 1957, **35**, 57

A glass bottle, undisturbed for several months, exploded, probably from internal gas pressure arising from photolytic or hydrolytic decomposition.

See other ACYL HALIDES, GAS EVOLUTION INCIDENTS

2235. Benzenesulfonyl chloride

[98-09-9]



HCS 1980, 188

Anon., *Chem. Eng. News*, 1957, **35**(47), 57

A glass container exploded after shelf storage for several months, (probably from internal gas pressure arising from slow hydrolysis by atmospheric or adsorbed moisture).

See other GAS EVOLUTION INCIDENTS

Dimethyl sulfoxide


See Dimethyl sulfoxide: Acyl halides

See other ACYL HALIDES

2236. Perchlorylbenzene

[5390-07-8]

$C_6H_5ClO_3$



Aluminium chloride

Bruce, W. F., *Chem. Eng. News*, 1960, **38**(4), 63


A mixture is quiescent for some time, then suddenly explodes. Many perchloryl aromatics appear to be inherently shock-sensitive.

See other PERCHLORYL COMPOUNDS

2237. 2,3-Dichloroaniline

[608-27-5]

$C_6H_5Cl_2N$




T_{ait24} was determined as 212°C by adiabatic Dewar tests, with an apparent energy of activation of 119 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HALOANILINES

2238. 2,4-Dichloroaniline
[554-00-7]

C₆H₅Cl₂N




T_{ait24} was determined as 211°C by adiabatic Dewar tests, with an apparent energy of activation of 129 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HALOANILINES

2239. 2,5-Dichloroaniline
[98-82-9]

C₆H₅Cl₂N




T_{ait24} was determined as 216°C by adiabatic Dewar tests, with an apparent energy of activation of 126 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HALOANILINES

2240. 2,6-Dichloroaniline
[608-31-1]

C₆H₅Cl₂N




T_{ait24} was determined as 217°C by adiabatic Dewar tests, with an apparent energy of activation of 227 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other HALOANILINES

2241. 3,4-Dichloroaniline
[95-76-1]

$C_6H_5Cl_2N$



HCS 1980, 374

T_{ait24} was determined as 187°C by adiabatic Dewar tests, with an apparent energy of activation of 146 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Preparative hazard

Kotoyori, T., private comm., 1983


During pilot-scale vacuum distillation from a mild steel still of the crude hydrogenation product of 3,4-dichloronitrobenzene, normal conditions had been established as 160–170°C/43–63 mbar with steam heating at 194°C. While the seventh 8 t batch was being distilled, distillation conditions were 187°C/135 mbar, and later 194°C/165 mbar, at which point heating was stopped but the temperature and pressure kept rising. The vessel (without a pressure relief) was isolated for 2.5 h while the vacuum line was examined for blockage. Before vacuum could be reapplied, the vessel ruptured at the lid flange with eruption of white smoke and black tar. Several contributory factors for the decomposition were established later. 3,4-Dichloroaniline undergoes dehydrochlorination-polymerisation reactions, forming hydrochloride salts, at temperatures above 180°C in presence of ferric chloride, and the salts dissociate above 260°C with violent liberation of hydrogen chloride gas. Leaks of water into the still from a condenser, and of air from a hardened and cracked gasket, tended to increase the rate of formation of ferric chloride by hydrolysis of the aniline and oxidation of iron dissolved from the vessel, as well as causing increases in distillation temperature and pressure. The level of dissolved iron in the residue after the accident was around 1% (80 kg Fe), owing to the fact that distillation residues were not removed after each batch was completed. Recommended precautionary measures included fitting a pressure relief, distilling under alkaline conditions, better gasket maintenance to prevent air leaks, and removal of the residue after each batch.

See 3,4-Dichlorophenylhydroxylamine

See other HALOANILINES

2242. *N,N*-Dichloroaniline
[70278-00-1]

$C_6H_5Cl_2N$




Goldschmidt, S., *Ber.*, 1913, **46**, 2732

The oil prepared at $-20^{\circ}C$ explodes while warming to ambient temperature.

See other *N*-HALOGEN COMPOUNDS

2243. 3,4-Dichlorophenylhydroxylamine
[33175-34-7]

$C_6H_5Cl_2NO$



1. Tong, W. R. *et al.*, *Loss Prev.*, 1977, **11**, 71–75

2. MacNab, J. I., *Runaway Reactions*, 1981, 3/S, 1–12


A violent explosion during the plant-scale catalytic hydrogenation of 3,4-dichloronitrobenzene to the aniline (heat of hydrogenation is 523 kJ/mol, 2.72 kJ/g) was traced to thermal decomposition of one of the reaction intermediates, 3,4-dichlorophenylhydroxylamine. The latter is normally present to the extent of about 5%, but contamination of the nitro compound by unseparated wash-water containing 2% of nitrite ion had slowed the reaction and increased the proportion of hydroxylamine to 35%. Investigation of the thermal stability of the reaction mixture as hydrogenation proceeded showed that the minimum autodecomposition temperature of $260^{\circ}C$ coincided with the maximum concentration of the hydroxylamine at 60% of the theoretical hydrogen uptake. The pure hydroxylamine begins to decompose at $80^{\circ}C$. Shortly before the explosion occurred, the heat input had been increased to offset the slower reaction rate and had triggered the decomposition. The sequence was confirmed experimentally, and maximum rates of increase of $6.7^{\circ}C/s$ and 36 bar/s were recorded for the decomposition [1]. The initial temperature for exothermic decomposition of this reduction intermediate, alone or diluted with the nitro or amino compounds, was $80^{\circ}C$ or below, with exotherms up to $60^{\circ}C$ [2].

See 2-Chloro-5-methylphenylhydroxylamine, also 3,4-Dichloroaniline

See other *N*-O COMPOUNDS, CATALYTIC NITRO REDUCTION PROCESSES

2244. *N,N*-Dichlorobenzenesulfonamide
[473-34-7]


C₆H₅Cl₂NO₂S



Energy of decomposition (in range 160–260°C) measured as 0.92 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other *N*-HALOGEN COMPOUNDS

2245. Phenylphosphonyl dichloride
[824-72-6]


C₆H₅Cl₂OP



Koch-Light Ltd., private comm., 1976
A bottle of the dichloride exploded violently while on a bench. No cause was established, but diffusive ingress of moisture, or an intramolecular Friedel-Crafts reaction, both of which would form free hydrogen chloride, seem possible contributory factors.
See related ACYL HALIDES, NON-METAL HALIDES

2246. Phenylvanadium(V) dichloride oxide
[28597-01-5]


C₆H₅Cl₂OV



Houben-Weyl, 1975, Vol. 13.3, 365
Thermal decomposition may be explosive.
See related ARYLMETALS, METAL HALIDES

2247. Phenyliodine(III) chromate ('[(Chromyldioxy)iodo]benzene')

[]




Sidgwick, 1950, 1250

It explodes at 66°C.

See other IODINE COMPOUNDS

†2248. Fluorobenzene


[462-06-6]



See other HALOARYL COMPOUNDS

2249. Iodobenzene

[591-50-4]




Sorbe, 1968, 66

It explodes on heating above 200°C.

See other HALOARYL COMPOUNDS, IODINE COMPOUNDS

2250. Benzenediazonium iodide
[57965-32-9]


$C_6H_5IN_2$



See entries DIAZONIUM SALTS, DIAZONIUM TRIIODIDES

2251. Phenyliodine(III) nitrate
[58776-08-2]


$C_6H_5IN_2O_6$



Sorbe, 1968, 66
It explodes above 100°C.
See other IODINE COMPOUNDS


2252. Iodosylbenzene
[536-80-1]

C_6H_5IO



1. Saltzman, A. *et al.*, *Org. Synth.*, 1963, **43**, 60
2. Banks, D. F., *Chem. Rev.*, 1966, **66**, 255
It explodes at 210°C [1], and homologues generally explode on melting [2].
See other IODINE COMPOUNDS


2253. Iodylbenzene
[696-33-3]



1. Sharefkin, J. G. *et al.*, *Org. Synth.*, 1963, **43**, 65
2. Banks, D. F., *Chem. Rev.*, 1966, **66**, 259


It explodes at 230°C [1]. Extreme care should be used in heating, compressing or grinding iodyl compounds, as heat or impact may cause detonation of homologues [2].
See other IODINE COMPOUNDS

2254. Benzenediazonium triiodide
[68596-95-2]




See entry DIAZONIUM TRIIODIDES

2255. Potassium nitrobenzene
[]



See entry NITROAROMATIC-ALKALI HAZARDS; potassium radical salts

2256. Potassium 3,5-dinitro-2(1-tetrazenyl)phenoxide
[70324-35-5]




Kenney, J. F., US Pat. 2 728 760, 1952

It is explosive.

See other HIGH-NITROGEN COMPOUNDS, POLYNITROARYL COMPOUNDS

2257. Potassium benzenesulfonylperoxosulfate

[]





Davies, 1961, 65

It explodes on friction or warming.

See other DIACYL PEROXIDES

2258. Potassium citrate tri(hydrogen peroxidate)

[31266-90-1]





Anon., *Jahresber.*, 1978, 75

Treatment of the citrate monohydrate with a 20% excess of 70% hydrogen peroxide gave a crystalline product which, after vacuum drying at 50°C corresponded approximately to the tri(hydrogen peroxidate). It decomposed violently when disturbed with a porcelain spatula.

See other CRYSTALLINE HYDROGEN PEROXIDATES

2259. Phenyllithium

[591-51-5]



Preparative hazard

See Lithium: Bromobenzene

Titanium tetraethoxide


Gilman, H. *et al.*, *J. Org. Chem.*, 1945, **10**, 507

The product of interaction at 0°C (of unknown composition) ignited in air and reacted violently with water.

See other ARYLMETALS

2260. 1,4-Benzoquinone monoimine (2,5-Cyclohexadienone-4-imine)
[3009-34-5]

C₆H₅NO




Willstätter, R. *et al.*, *Ber.*, 1906, **37**, 4607

The solid decomposes with near-explosive violence.

See related DIENES

2261. Nitrosobenzene
[586-86-9]

C₆H₅NO




Energy of decomposition (in range 190–380°C) measured as 0.823 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROSO COMPOUNDS

2262. Nitrobenzene
[98-95-3]

C₆H₅NO₂



(MCA SD-21, 1967); HCS 1980, 681; 32RSC Lab Hazards Data Sheet No. 77, 1988

CHETAH, 1990, 182

It is marginally detonable; requiring a 50 g tetryl booster.

The heat of decomposition of nitrobenzene was determined (360–490°C) as 1.76 kJ/g by DSC, and T_{ait24} was as 280°C by adiabatic Dewar tests, with an apparent energy of activation of 216 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkali

1. Hatton, J. P., private comm., 1976
2. Anon., *Sichere Chemiearb.*, 1975, **27**, 89; Bilow, W., private comm., 1976
3. Wohl, A., *Ber.*, 1899, **32**, 3846; 1901, **34**, 2444
4. Anon., *ABCM Quart. Safety Summ.*, 1953, **24**, 42
5. Bretherick, L., *Chem. & Ind.*, 1976, 576
6. *CHSO Rept. Y4648-181*, Washington (DC), OSHA, 1981

Heating a mixture of nitrobenzene, flake sodium hydroxide and a little water in an autoclave led to an explosion [1]. During the technical-scale preparation of a warm solution of nitrobenzene in methanolic potassium hydroxide (flake 90% material), accidental omission of most of the methanol led to an accelerating exothermic reaction which eventually ruptured the 6 m³ vessel. Laboratory investigation showed that no exothermic reaction occurred between potassium hydroxide and nitrobenzene, either alone, or with the full amount (3.4 vol.) of methanol, but that it did if only a little methanol were present. The residue was largely a mixture of azo- and azoxy-benzene [2]. The violent conversion of nitrobenzene to (mainly) 2-nitrophenol by heating with finely powdered and anhydrous potassium hydroxide had been described earlier [3]. Accidental substitution of nitrobenzene for aniline as diluent during large-scale fusion of benzanthrone with potassium hydroxide caused a violent explosion [4]. Related incidents involving aromatic nitro compounds and alkali are also discussed [5]. Drying wet nitrobenzene in presence of flake sodium hydroxide (probably by distilling out the nitrobenzene–water azeotrope) led to separation of finely divided solid base which remained in contact with the hot dry nitrobenzene under essentially adiabatic conditions. A violent explosion subsequently blew out a valve and part of the 10 cm steel pipeline. These conditions, when simulated in an ARC test run, also led to thermal runaway and explosion [6].

See other NITROAROMATIC–ALKALI HAZARDS

Aluminium chloride

See Aluminium chloride: Nitrobenzene, or :Nitrobenzene, Phenol

Aniline, Glycerol, Sulfuric acid

See Quinoline

Other reactants

Yoshida, 1980, 277

MRH values calculated for 2 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Dinitrogen tetraoxide: Nitrobenzene

Fluorodinitromethane: Nitrobenzene

Nitric acid: Nitroaromatics, or: Nitrobenzene
Peroxydisulfuric acid: Organic liquids
Sodium chlorate: Nitrobenzene
Tetranitromethane: Aromatic nitro compounds
Uronium perchlorate: Organic materials

MRH 4.94/12
MRH 7.36/76

Phosphorus pentachloride

See Phosphorus pentachloride: Nitrobenzene

Potassium

See Potassium: Nitrogen-containing explosives

Sulfuric acid

1. MCA Case History No. 678
2. Silverstein, J. L. *et al.*, *Loss Prev.*, 1981, **14**, 78

Nitrobenzene was washed with dilute (5%) sulfuric acid to remove amines, and became contaminated with some tarry emulsion that had formed. After distillation, the hot tarry acidic residue attacked the iron vessel with hydrogen evolution, and an explosion eventually occurred. It was later found that addition of the nitrobenzene to the diluted acid did not give emulsions, while the reversed addition did. A final wash with sodium carbonate solution was added to the process [1]. During hazard evaluation of a continuous adiabatic process for manufacture of nitrobenzene, it was found that the latter with 85% sulfuric acid gave a violent exotherm above 200°C, and with 69% acid a mild exotherm at 150–170°C [2].

Sulfuric acid, Sulfur trioxide

See 3-Nitrobenzenesulfonic acid

Tin(IV) chloride

Unpublished observations, 1948

Mixtures of nitrobenzene and tin(IV) chloride undergo exothermic decomposition with gas evolution above 160°C.

See other GAS EVOLUTION INCIDENTS

See other NITROARYL COMPOUNDS

2263. 2-Nitrosophenol (1,2-Benzoquinone monoxime)

[13168-78-0] ([637-62-7])

C₆H₅NO₂



Energy of decomposition (in range 80–320°C) measured as 1.214 kJ/g by DSC, and T_{ait24} was determined as the very low figure of 53°C by adiabatic Dewar tests, with an apparent energy of activation of 462 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other DEFLAGRATION INCIDENTS

Alone, or Acids

Baeyer, A. *et al.*, *Ber.*, 1902, **35**, 3037

It explodes on heating or in contact with concentrated acids (the latter possibly involving Beckmann rearrangement and/or polymerisation of the oxime form, which is effectively a 1,3-diene).

See other NITROSO COMPOUNDS, OXIMES

See related DIENES

2264. 4-Nitrosophenol (1,4-Benzoquinone monoxime)

[637-62-7] ([104-91-6])

$C_6H_5NO_2$



Alone, or Acids, or Alkalies

1. Milne, W. D., *Ind. Eng. Chem.*, 1919, **11**, 489
2. Kuznetsov, V., *Chem. Abs.*, 1940, **34**, 3498i
3. Anon., *Sichere Chemiearb.*, 1980, 3532, 6
4. Klais, O. *et al.*, *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 3, C24–C34, Rugby, IChE, 1983
5. Grever, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987


Stored barrels heated spontaneously and caused a fire. Contamination of the bulk material by acid or alkali may cause ignition [1]. The tendency to spontaneous ignition was found to be associated with the presence of nitrates in the sodium nitrite used in preparation. A diagnostic test for this tendency is to add sulfuric acid, when impure material will effervesce or ignite. The sodium salt is more suitable for storage purposes [2] (*but see* Sodium 4-nitrosophenoxide). Compressed material has a greater tendency to self-ignition than the loose solid, and a 20–30 kg quantity of centrifuged material ignited in a drum located near a steam heater [3]. Material for laboratory use is now supplied wet with 30–40% water. The heat of decomposition was determined as 1.2 kJ/g, and the pressure rise occurring during exothermic decomposition is low [4]. However, the deflagration rate for the solid of 8 cm/min is relatively high [5].

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

See other NITROSO COMPOUNDS

See related OXIMES

2265. 2-Nitrophenol
[88-75-5]



Energy of decomposition (in range 260–365°C) measured as 2.15 kJ/g by DSC, and T_{ait24} was determined as 182°C by adiabatic Dewar tests, with an apparent energy of activation of 135 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Chlorosulfuric acid

See 4-Hydroxy-3-nitrobenzenesulfonyl chloride


Potassium hydroxide

491M, 1975, 342

The molten phenol reacts violently with commercial 85% potassium hydroxide pellets (possibly involving formation of the *aci-o*-nitroquinonoid salt).

See NITROAROMATIC-ALKALI HAZARDS See other NITROARYL COMPOUNDS

2266. 3-Nitrophenol
[554-81-7]




T_{ait24} was determined as 213°C by adiabatic Dewar tests, with an apparent energy of activation of 148 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROARYL COMPOUNDS

2267. 4-Nitrophenol
[100-02-7]



HCS 1980, 688

1. Grewer, T. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 657–664, Basle, SSCI, 1980
2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn, 1986

There is a high rate of pressure increase during exothermic decomposition [1]. Energy of decomposition (in range 240–440°C) measured as 1.58 kJ/g by DSC, and T_{ait24} was determined as 185°C by adiabatic Dewar tests, with an apparent energy of activation of 117 kJ/mol [2].

Diethyl phosphite

See Diethyl phosphite: 4-Nitrophenol

Potassium hydroxide

Solid mixtures of 4-nitrophenol with potassium hydroxide (1:1.5 mol) readily deflagrate, at the rapid rate of 30 cm/min.

See entries METAL NITROPHENOXIDES, PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 3)

See Potassium 4-nitrophenoxide


See other DEFLAGRATION INCIDENTS

See other NITROARYL COMPOUNDS

2268. 3-Nitrobenzenesulfonic acid

[984-47-5]

$C_6H_5NO_5S$



Sulfuric acid, Sulfur trioxide


1. *MCA Case History No. 1482*
2. *MCA Case History No. 944*
3. See entry SELF-ACCELERATING REACTIONS
4. Nolan, 1983, Case history 102
5. Bergroth, K., *Loss Prev. Bull.*, 1993, (109), 1

The acid is prepared by sulfonation of nitrobenzene with oleum, and the reaction product consists essentially of a hot solution of the acid in sulfuric acid. A completed 270 l batch exploded violently after hot storage at ~150°C for several hours. An exotherm develops at 145°C, and the acid is known to decompose at ~200°C [1]. A similar incident arose from water leaking from a cooling coil into the fuming sulfuric acid reaction medium, which caused an exotherm to over 150°C and subsequent violent decomposition [2]. Detailed examination of the thermal decomposition of the acid shows that it is much slower for the isolated acid than for the reaction mass, and that the concentration of sulfur trioxide in the oleum used for sulfonation bears

directly upon the decomposition rate. Care must be exercised when interpreting short-term DSC test results, because the self-acceleration features are not immediately evident. A Q value of 1 kJ/g of reaction mass after sulfonation with 25% oleum was found experimentally [3]. A 2000 l batch of the acid prepared with 20% excess of 65% oleum had been cooled to 70°C prior to being blown over into a tank for treatment with sodium carbonate. Application of air pressure at 2.75 bar to the reactor failed to transfer the product, so the transfer lines were isolated and steamed at 2 bar to clear them. On reconnecting the lines, the condensed water reacted with the excess oleum, leading to exothermic decomposition of the acid, rupture of the vessel and a fire [4]. An explosion in a plant resulted because of crystallisation from the oleum sulfonation mixture in a control valve, which failed. Manual operation led to steam flowing back from the quench to a nitration-product holding tank, which overheated and exploded [5].

See other NITROARYL COMPOUNDS, ORGANIC ACIDS, SULFONATION INCIDENTS

2269. Benzotriazole
[95-14-7]





1. Anon., *Chem. Eng. News*, 1956, **34**, 2450
2. *CHETAH*, 1990, 182
3. Stull, 1977, 22–23

A 1 t batch decomposed exothermally, then detonated at 220°C, during distillation at 160°C/2.5 mbar. No cause was found, and similar batches had previously distilled satisfactorily. The multiple N–N bonding would tend to cause instability in the molecule, particularly in presence of heavy metals, but these were absent in this case [1]. It is shock sensitive (probably not very) [2]. Benzotriazole is an endothermic compound (ΔH_f° (s) +249.8 kJ/mol, 2.1 kJ/g) and this energy on release would attain an adiabatic decomposition temperature approaching 1100°C, with an 18 bar pressure increase in the closed system [3].

See other ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES


2270. 1,2,4-Triazolo[4,3-a]pyridine silver nitrate
[]



Fargher, R. G. *et al.*, *J. Chem. Soc.*, 1915, **107**, 696

The base [274-80-6] forms an addition compound with silver nitrate which decomposes at 228°C in a m.p. capillary, but explodes when heated in larger amount.
See other AMMINOMETAL NITRATES, SILVER COMPOUNDS

2271. Phenyl azide (Azidobenzene)
[622-37-7]




1. Lindsay, R. O. *et al.*, *Org. Synth.*, 1955, Coll. Vol. 3, 710
 2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
 3. Lee, S. W. *et al.*, *Inorg. Chem.*, 1988, **27**, 1213–1219
- Though distillable at considerably reduced pressure, phenyl azide explodes when heated at ambient pressure, and occasionally at lower pressures [1]. The energy of decomposition has been determined (130–260°C) as 1.36 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 87°C by adiabatic Dewar tests, with an apparent energy of activation of 119 kJ/mol [2]. It is best stored as a 3M solution in toluene. Exposure to heavy metals should be avoided [3].

Lewis acids

Boyer, J. H. *et al.*, *Chem. Rev.*, 1954, **54**, 29

The ready decomposition of most aryl azides with sulfuric acid and Lewis acids may be vigorous or violent, depending on structure and conditions. In absence of a diluent (carbon disulfide), phenyl azide and aluminium chloride exploded violently.
See other ORGANIC AZIDES

2272. 1-Hydroxybenzotriazole
[2592-95-2]



Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 52–53

A sample of the compound (which can isomerise to two *N*-oxide forms) was inadvertently heated, probably to well over 160°C, during vacuum drying and exploded violently. Several salts have been reported as powerfully explosive.


Copper(II) nitrate

See Copper(II) nitrate: Hydroxybenzotriazole, Ethanol

See other HIGH-NITROGEN COMPOUNDS, N-O COMPOUNDS, TRIAZOLES

2273. Benzenesulfinyl azide
[21230-20-6]


C₆H₅N₃OS



See entry SULFINYL AZIDES See other ACYL AZIDES

2274. Benzenesulfonyl azide
[938-10-3]

C₆H₅N₃O₂S




Dermer, O. C. *et al.*, *J. Amer. Chem. Soc.*, 1955, **77**, 71

Whereas the pure azide decomposes rapidly but smoothly at 105°C, the crude material explodes violently on heating.

See other ACYL AZIDES

2275. Benzenediazonium nitrate
[619-96-7]

C₆H₅N₃O₃




1. Urbanski, 1967, Vol. 3, 201
2. Stull, 1977, 19

Although not a practical explosive, the isolated salt is highly sensitive to friction and impact, and explodes at 90°C [1]. It is, like most diazonium salts, endothermic, with ΔH_f° (s) +200.8 kJ/mol, 1.2 kJ/g [2].

See other DIAZONIUM SALTS, ENDOTHERMIC COMPOUNDS

2276. Dinitroanilines (: all isomers except 3,5-)

2,3- [602-03-9]; 2,4- [97-02-9]; 2,5- [619-18-1]; 2,6- [606-22-4]; $C_6H_5N_3O_4$
3,4- [610-41-3]



Chlorine, Hydrochloric acid


See Hydrochloric acid: Chlorine, Dinitroanilines

See other POLYNITROARYL COMPOUNDS

2277. 2,4-Dinitroaniline

[97-02-9]

$C_6H_5N_3O_4$



HCS 1980, 437

Preparative hazard

1. Sorbe, 1968, 151
2. CHETAH, 1990, 183
3. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

It is a compound with high fire and explosion hazards [1], showing fast flame propagation on heating the powder moderately. Not sensitive to mechanical shock,

it can be exploded by a detonator [2]. T_{ait24} was determined as 217°C by adiabatic Dewar tests, with an apparent energy of activation of 129 kJ/mol [3].

See 1-Chloro-2,4-dinitrobenzene: Ammonia

See entry HIGH RATE DECOMPOSITION


Charcoal

Leleu, *Cahiers*, 1980, (99), 278

A mixture with powdered charcoal smoked above 280°C and ignited at 350°C.

See other POLYNITROARYL COMPOUNDS

2278. 2-Amino-4,6-dinitrophenol [96-91-3]




Sorbe, 1968, 152

This partially reduced derivative of picric acid, 'picramic acid', explodes very powerfully when dry.

See other POLYNITROARYL COMPOUNDS

2279. O-(2,4-Dinitrophenyl)hydroxylamine [17508-17-7]



Potassium hydride

Radhakrishna, A. S. *et al.*, *J. Org. Chem.*, 1979, **44**, 4838


Interaction in THF gives 2,4-dinitrophenol and ammonia, and on one occasion a violent detonation occurred; potassium dinitrophenoxide may have been involved.

See Sodium 2,4-dinitrophenoxide

See other N-O COMPOUNDS, POLYNITROARYL COMPOUNDS

2280. 2-Acetylamino-3,5-dinitrothiophene
[51249-08-2]


C₆H₅N₃O₅S



Fast flame propagation on heating the powder moderately.
See entry HIGH RATE DECOMPOSITION *See other* POLYNITROARYL COMPOUNDS

2281. Benzotriazolium 1-nitroimidate
[]


C₆H₅N₅O₂



A dangerous impact-sensitive solid which explodes at the m.p., 74°C.
See entry N-AZOLIUM NITROIMIDATES

2282. Benzotriazolium 2-nitroimidate
[52096-20-5]


C₆H₅N₅O₂



A dangerous solid which explodes at the m.p., 84°C.
See entry N-AZOLIUM NITROIMIDATES

2283. 1-(Nitroamino)benzotriazole
[52096-18-1]

C₆H₅N₅O₂



This is the structure assigned by *Chem. Abs.* to the claimed benzotriazolium nitroimidate below. The editor suspects they are right and that the nitroamine

structure should also be considered for the 2-isomer. The *aci*-nitro structures are also possible.


See Benzotriazolium 1-nitroimidate

See also N-NITRO COMPOUNDS

2284. Phenylphosphonic diazide

[]

$C_6H_5N_6OP$



Baldwin, R. A., *J. Org. Chem.*, 1965, **30**, 3866


Although a small sample had been distilled at 72–74°C/0.13 mbar without incident, it decomposed vigorously on exposure to flame, and the liquid was impact-sensitive, exploding vigorously. The material can be prepared and handled safely in pyridine solution.

See other ACYL AZIDES

2285. Phenylthiophosphonic diazide

[]

$C_6H_5N_6PS$



Baldwin, R. A., *J. Org. Chem.*, 1965, **30**, 3866


During an attempt to distil crude material at 80°C/0.13 mbar, a violent explosion occurred. The material can be prepared and handled safely in pyridine solution.

See other ACYL AZIDES

2286. Phenylthallium diazide

[]

C₆H₅N₆Tl




Pascal, 1961, Vol. 6, 1019
It decomposes violently at 200°C.
See related METAL AZIDES

2287. Phenylsodium

[1623-99-0]

C₆H₅Na



Preparative hazard

Sorbe, 1968, 86

Suspensions containing phenylsodium may ignite and burn violently in moist air.


See Sodium: Halocarbons (reference 8)

See other ARYLMETALS

†2288. Benzene

[71-43-2]

C₆H₆



(MCA SD-2, 1960); NSC 308, 1979; FPA H18, 1973; HCS 1980, 187; RSC Lab.
Hazard Data Sheet No. 20, 1984

Arsenic pentafluoride, Potassium methoxide

See Arsenic pentafluoride: Benzene, etc.

Diborane

See Diborane: Benzene, etc.

Hydrogen, Raney nickel

Anon., *Loss Prev. Bull.*, 1979, (028), 91–92

Hydrogenation of benzene to cyclohexane was effected in a fixed bed reactor at 210–230°C, but a fall in conversion was apparent. Increasing the bed temperature by 10°C and the hydrogen flow led to a large increase in reaction rate which the interbed cooling coils could not handle, and an exotherm to 280°C developed, with a hot spot of around 600°C which bulged the reactor wall.

See other HYDROGENATION INCIDENTS

Interhalogens

MRH value shows % of interhalogen

See Bromine trifluoride: Halogens, etc.

Bromine pentafluoride: Hydrogen-containing materials MRH 3.85/73

Iodine pentafluoride: Benzene

Iodine heptafluoride: Organic solvents

Other reactants

Yoshida, 1980, 334

MRH values calculated for 17 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Dioxygen difluoride: Various materials

Dioxygenyl tetrafluoroborate: Organic materials

Nitric acid: Hydrocarbons MRH 8.15/82

Nitryl perchlorate: Organic solvents

Oxygen (Liquid): Hydrocarbons MRH 9.83/75

Ozone: Aromatic compounds, or: Benzene, etc. MRH Benzene 12.05/75

Permanganic acid: Organic materials

Peroxodisulfuric acid: Organic liquids

Peroxomonosulfuric acid: Aromatic compounds

Silver perchlorate MRH 3.51/91

Sodium peroxide: Organic liquids, etc. MRH 2.76/93

Ozone: Aromatic compounds, or: Benzene, etc. MRH Benzene 12.05/75


Uranium hexafluoride

See Uranium hexafluoride: Aromatic hydrocarbons

2289. Benzvalene

[659-85-8]

C₆H₆




Katz, T. J. *et al.*, *J. Amer. Chem. Soc.*, 1971, **93**, 3782

When isolated, this highly strained valence tautomer of benzene exploded violently when scratched. It may be handled safely in solution in ether.

See other STRAINED-RING COMPOUNDS

2290. 1,3-Hexadien-5-yne
[10420-90-3]

C₆H₆



Brandsma, 1971, 132

Distillation of this rather unstable material at normal pressure involves risk of an explosion.

See other ALKYNES, DIENES

†**2291. 1,5-Hexadien-3-yne (Divinylacetylene)**
[821-08-9]

C₆H₆



1. Nieuwland, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1931, **53**, 4202

2. Cupery, M. E. *et al.*, *J. Amer. Chem. Soc.*, 1934, **56**, 1167–1169


3. Handy, C. T. *et al.*, *J. Org. Chem.*, 1962, **27**, 41

The extreme hazards involved in handling this highly reactive material are stressed. Freshly distilled material rapidly polymerises at ambient temperature to produce a gel and then a hard resin. These products can neither be distilled nor manipulated without explosions ranging from rapid decomposition to violent detonation. The hydrocarbon should be stored in the mixture with catalyst used to prepare it, and distilled out as required [1]. The dangerously explosive gel is a peroxidic species not formed in absence of air, when some 1,2-di(3-buten-1-ynyl)cyclobutane is produced by polymerisation [2]. The dienyne reacts readily with atmospheric oxygen, forming an explosively unstable polymeric peroxide. Equipment used with it should be rinsed with a dilute solution of a polymerisation inhibitor to prevent formation of unstable residual films. Adequate shielding of operations is essential [3].

See other ALKYNES, DIENES, POLYPEROXIDES

2292. 1,5-Hexadiyne
[628-16-0]

C₆H₆




Grignard, 1935, Vol. 3, 181

At the end of distillation at 86°C/1 bar, the last drop ignited. It explodes if heated to 100–120°C, and the copper(I) salt also ignites.

See other ALKYNES

2293. 2,4-Hexadiyne
[2809-69-0]

C₆H₆




1. Bohlman, F., *Chem. Ber.*, 1951, **84**, 785–794
2. Armitage, J. B., *J. Chem. Soc.*, 1951, 45

Filtration through alumina, which prevents explosion on subsequent distillation at 55°C/13 μbar [1], or at ambient pressure [2], presumably removes peroxidic impurities.

See other ALKYNES

2294. Prismane
[650-42-0]

C₆H₆



Katz, T. J. *et al.*, *J. Amer. Chem. Soc.*, 1973, **95**, 2739

This tetracyclic isomer of benzene is an explosive liquid.

See other STRAINED-RING COMPOUNDS

2295. Iodosylbenzene tetrafluoroborate
[120312-56-3]

C₆H₆BF₄ 4IO



van Look G., *Chem. Eng. News*, 1989 **67**(30), 2


A compound with tetrafluoroboric acid, previously claimed to be a useful safe oxidant, proved not to be on preparation at 0.5 molar scale. It exploded spontaneously while drying in vacuo at room temperature. *Chem. Abs.* (**112** 98095p) identifies it with this title compound, which was claimed to be too unstable to explode by the original users. It seems more likely that the explosive was the "anhydride", oxybisphenyliodonium tetrafluoroborate. Several related compounds have also proved to be explosive.

See Oxybisphenyliodonium bistetrafluoroborate, also Iodosylbenzene

See other IODINE COMPOUNDS

2296. 4-Bromoaniline
[106-40-1]

C₆H₆BrN



HCS 1980, 220

Energy of decomposition (in range 220–270°C) measured as 0.428 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


Nitrous acid

See 4-Bromobenzenediazonium salts

See other HALOANILINES

2297. 2-Bromo-4-methylpyridine N-oxide
[17117-12-3]

C₆H₆BrNO




Preparative hazard

See 3-Chloroperoxybenzoic acid: 2-Bromo-4-methylpyridine

See other N-OXIDES

2298. 2,3-Dibromo-5,6-epoxy-7,8-dioxabicyclo[2.2.2]octane
[56411-66-6]

C₆H₆Br₂O₃




Sasaoka, M. *et al.*, *J. Org. Chem.*, 1979, **44**, 373 (footnote 10)

It explodes on heating.

See other CYCLIC PEROXIDES, 1,2-EPOXIDES

2299. (Hydroxy)(oxo)(phenyl)- λ^3 -iodanium perchlorate (Iodylbenzene perchlorate)
[101672-14-4]




Masson, I., *J. Chem. Soc.*, 1935, 1674

A small sample exploded very violently while still damp.

See other IODINE COMPOUNDS, NON-METAL PERCHLORATES

2300. 2-Chloroaniline
[95-51-2]



HCS 1980, 295

Heat of decomposition was determined as 0.41 kJ/g.


See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

Nitrous acid

See 2-Chlorobenzenediazonium salts

See other HALOANILINES

2301. 3-Chloroaniline
[108-42-9]



Heat of decomposition was determined as zero.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

Nitrous acid


See 3-Chlorobenzenediazonium salts

See other HALOANILINES

2302. 4-Chloroaniline

[106-47-8]

C₆H₆ClN



HCS 1980, 294

Energy of exothermic decomposition in range 210–400°C was measured as 0.63 kJ/g by DSC, and T_{ait24} was determined as 200°C by adiabatic Dewar tests, with an apparent energy of activation of 124 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


Nitrous acid

See DIAZONIUM SULFIDES AND DERIVATIVES (references 8,9) *See other* HALOANILINES

2303. 4-Chloro-2-aminophenol

[95-85-2]

C₆H₆ClNO



T_{ait24} was determined as 110°C by adiabatic Dewar tests, with an apparent energy of activation of 140 kJ/mol.


See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HALOANILINES

2304. 4-Aminobenzediazonium perchlorate

[]

C₆H₆ClN₃O₄



Hofmann, K. A. *et al.*, *Ber.*, 1910, **43**, 2624


Extremely explosive.

See other DIAZONIUM PERCHLORATES

2305. 2,5-Bis(chloromethyl)thiophene

[28569-48-4]

C₆H₆Cl₂S



Griffing, J. M. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 3417


It polymerises at ambient temperature and must therefore be stored cold.

See other 2-HALOMETHYL-FURANS OR -THIOPHENES

2306. 1,2,3,4,5,6-Hexachlorocyclohexane

[608-73-1]

C₆H₆Cl₆



Dimethylformamide

'DMF Brochure', Billingham, ICI, 1965

There is a potentially dangerous reaction of hexachlorocyclohexane with DMF in presence of iron. The same occurs with carbon tetrachloride, but not with dichloromethane or 1,2-dichloroethane under the same conditions.


See *N,N*-Dimethylacetamide: Halogenated compounds

See related HALOALKANES

See other CATALYTIC IMPURITY INCIDENTS

2307. *N*-Ethylheptafluorobutyramide
[70473-76-6]


$C_6H_6F_7NO$



Lithium tetrahydroaluminate
See Lithium tetrahydroaluminate: Fluoroamides

2308. Potassium bis(propynyl)palladate
[]


$C_6H_6K_2Pd$



Air, or Water
Immediately pyrophoric in air, and explosive decomposition with aqueous reagents.
The sodium salt is similar.
See entry COMPLEX ACETYLIDES

2309. Potassium bis(propynyl)platinate
[]


$C_6H_6K_2Pt$



Air, or Water
Pyrophoric in air, explosive decomposition with water, and the sodium salt is similar.
See entry COMPLEX ACETYLIDES

2310. 1,4-Benzoquinone diimine (1,4-Diimido-2,5-cyclohexadiene)
[4377-73-5]

$C_6H_6N_2$



Acids


Willstätter, R. *et al.*, *Ber.*, 1906, **37**, 4607

The unstable solid decomposes explosively in contact with conc. hydrochloric acid or sulfuric acid.

See related DIENES

2311. 1,4-Dicyano-2-butene
[1119-69-3]

$C_6H_6N_2$




MCA Case History No. 1747

Overheating in a vacuum evaporator initiated accelerating polymerisation-decomposition of (endothermic) dicyanobutene, and the rapid gas evolution eventually caused pressure-failure of the process equipment.

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS, GAS EVOLUTION INCIDENTS, POLYMERISATION INCIDENTS

2312. Bis(acrylonitrile)nickel(0)
[12266-58-9]

$C_6H_6N_2Ni$




Schrauzer, G. N., *J. Amer. Chem. Soc.*, 1959, **81**, 5310

Pyrophoric in air.

See other CYANO COMPOUNDS, PYROPHORIC MATERIALS

2313. 2-Nitroaniline
[88-74-4]

C₆H₆N₂O₂



Energy of decomposition (in range 280–380°C) measured as 1.81 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 223°C by adiabatic Dewar tests, with an apparent energy of activation of 171 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Preparative hazard

Duck, M. W. *et al.*, *Plant/Oper. Progr.*, 1982, **1**, 19–26

Exothermic decomposition of 2-nitroaniline in chemical processes was studied by DSC and ARC techniques. The stability in reaction mixtures was markedly less than for the pure, isolated compound.

See 2-Chloronitrobenzene: Ammonia

Hexanitroethane

See Hexanitroethane: Organic compounds

Magnesium, Nitric acid

See Nitric acid: Magnesium, 2-Nitroaniline

Nitrous acid

See 2-Nitrobenzenediazonium salts


Sulfuric acid

See Sulfuric acid: Nitroaryl bases

See other NITROARYL COMPOUNDS

2314. 3-Nitroaniline
[99-09-2]

C₆H₆N₂O₂



Hakl, J. *et al.*, *Runaway Reactions*, 1981, Paper 3/L, 4, 10

Energy of decomposition (in range 280–380°C) measured as 1.882 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 213°C by adiabatic Dewar tests, with an apparent energy of activation of 149 kJ/mol.

Initial exothermic decomposition occurs at 247°C.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Ethylene oxide

See Ethylene oxide: 3-Nitroaniline

Nitric acid, Sulfuric acid

Vanderah, D. J., *J. Energ. Mater.*, 1990, **58**(5), 378

Nitration to 2,3,4,6-tetranitroaniline proved unsafe in mixed acid at 30–50°C. Efficient and safe nitration at 20–30°C was obtained by dissolving the substrate in sulfuric acid, then adding mixed acid (made with oleum) thereto.


See other NITRATION INCIDENTS

See other NITROARYL COMPOUNDS

2315. 4-Nitroaniline

[100-01-6]

$\text{C}_6\text{H}_6\text{N}_2\text{O}_2$



HCS 1980, 680

Energy of decomposition (in range 280–380°C) measured as 1.882 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 217°C by adiabatic Dewar tests, with an apparent energy of activation of 160 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Nitrous acid

See 4-Nitrobenzenediazonium salts

Sodium hydroxide

See Sodium 4-nitrophenoxide


Sulfuric acid

See Sulfuric acid: Nitroaryl bases

See other NITROARYL COMPOUNDS

2316. 2-Amino-5-nitrophenol
[121-88-0]

$C_6H_6N_2O_3$



Nitrous acid


Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 53

An explosion in a poorly maintained and infrequently cleaned rubber lined vessel used for diazotisation of 2-amino-5-nitrophenol was attributed to decomposition of traces of a diazonium derivative (possibly 4-nitrobenzenediazonium-2-oxide) which had accumulated and dried out on the underside of the lid.

See ARENEDIAZONIUM OXIDES See other NITROARYL COMPOUNDS

2317. 3-Methyl-4-nitropyridine N-oxide
[1074-98-2]

$C_6H_6N_2O_3$



Preparative hazard


1. Ross, W. C. J., *J. Chem. Soc. C*, 1966, 1816
2. Kuilen, A. v. d. *et al.*, *Chem. Eng. News*, 1989, **67**(5), 2
3. Baumgarten, H. E., *J. Amer. Chem. Soc.*, 1957, **79**, 3145

An explosion occurred during attempted preparation of the title compound following a published procedure. This involved conversion of 3-methylpyridine to the *N*-oxide by treatment with hydrogen peroxide in glacial acetic acid, vacuum evaporation of the volatiles, and then addition of sulfuric and nitric acids, followed by slight warming to effect nitration [1]. Soon after warming the final mixture, a vigorous reaction set in, followed by an explosion. This was attributed to the presence of residual conc. peracetic acid in the *N*-oxide, and its subsequent detonation by the nitration exotherm [2]. An alternative procedure for preparing *N*-oxides precludes formation of peracetic acid [3].

See other *N*-OXIDES, NITRATION INCIDENTS

2318. Benzenediazonium hydrogen sulfate
[36211-73-1]

C₆H₆N₂O₄S




Sorbe, 1968, 122

The salt explodes at about 100°C, but not on impact.

See other DIAZONIUM SULFATES

2319. 4-Nitroaniline-2-sulfonic acid
[96-75-3]

C₆H₆N₂O₅S




See Sulfuric acid: Nitroaryl bases

See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2320. η⁶-Benzeneruthenium(II) dinitrate
[]

C₆H₆N₂O₆Ru




Bennett, M. A. *et al.*, *Organometallics*, 1992, **11**(9), 3069

This compound is a shock sensitive explosive.

See other ORGANOMETALLIC NITRATES


2321. 2,4-Dinitrophenylhydrazine
[119-26-6]



This spot-test reagent is now supplied water-wet since it has been found to be detonable and legally classifiable as an explosive. The editor has not heard of any problems in use.

See other POLYNITROARYL COMPOUNDS

2322. Ammonium picrate
[131-74-8]



Unpublished data.


A small sample, isolated incidentally during decomposition of a picrate with ammonia, exploded during analytical combustion. The presence of traces of metallic picrates (arising from metal contact) increases heat sensitivity. The salt may also explode on impact.

See other PICRATES, POLYNITROARYL COMPOUNDS

2323. *cis*-1,2-Bis(azidocarbonyl)cyclobutane

[]

C₆H₆N₆O₂



Landgrebe, J. A., *Chem. Eng. News*, 1981, **59**(17), 47


Explosion during preparation is reported for this and its cyclopropane analogue.

See other ACYL AZIDES

2324. 1,3,5-Triaminotrinitrobenzene

[3058-38-0]

C₆H₆N₆O₆



Hydroxylaminium perchlorate


See Hydroxylaminium perchlorate: 1,3,5-Triaminotrinitrobenzene

See other POLYNITROARYL COMPOUNDS

2325. Hexanitrohexaazaisowurtzitane (Octahydro-1,3,4,7,8,10-hexanitro-5,2,6-(iminomethenimino-1*H*-imidazo[4,5-*b*]pyrazine))

[135285-90-4]

C₆H₆N₁₂O₁₂



Borman, S., *Chem. Eng. News*, 1994, **72**(3), 18


This nitramine is claimed to be the most energetic molecular explosive known to man (which the editor doubts).

See other *N*-NITRO COMPOUNDS

2326. Di(1-propynyl) ether

[]

C₆H₆O



Brandsma, L. *et al.*, *Rec. Trav. Chim.*, 1962, **81**, 510


It is unstable, exploding at ambient temperature.

See other ACETYLENIC COMPOUNDS

2327. Di(2-propynyl) ether (Dipropargyl ether)

[6921-27-3]

C₆H₆O



Guide for Safety, 1972, 302


Distillation of the ether in a 230 l still led to an explosion, attributable to peroxidation.

See other ACETYLENIC COMPOUNDS, PEROXIDISABLE COMPOUNDS

2328. 4,5-Hexadien-2-yn-1-ol

[2749-79-3]

C₆H₆O



Brandsma, 1971, 8, 54


Dilution with white mineral oil before distillation is recommended to prevent explosion of the concentrated distillation residue.

See other ACETYLENIC COMPOUNDS, DIENES, PEROXIDISABLE COMPOUNDS

2329. 2,5-Hexadiyn-1-ol

[28255-99-4]

C₆H₆O



Houben-Weyl, 1977, Vol. 5.2a, 480


The residue after vacuum distillation (at 59–62°C/0.7 mbar) may explode on strong heating or on admission of air, unless mineral oil diluent is present.

See other ACETYLENIC COMPOUNDS, PEROXIDISABLE COMPOUNDS

2330. Phenol (Hydroxybenzene)

[108-95-2]

C₆H₆O



(MCA SD-4, 1964); NSC 405, 1978; FPA H82, 1979; HCS 1980, 725; RSC Lab. Hazards Data Sheet No. 42, 1985

Aluminium chloride, Nitrobenzene

See Aluminium chloride: Nitrobenzene, etc.

Aluminium chloride–nitromethane complex, Carbon monoxide

See Aluminium chloride–nitromethane: Carbon monoxide, etc.

Formaldehyde

See Formaldehyde: Phenol

Other reactants

Yoshida, 1980, 295

MRH values calculated for 15 combinations, largely with oxidants, are given.

Peroxodisulfuric acid

See Peroxodisulfuric acid: Organic liquids

Peroxomonosulfuric acid

See Peroxomonosulfuric acid: Aromatic compounds

Sodium nitrate, Trifluoroacetic acid

See Sodium nitrate: Phenol, etc.


Sodium nitrite

See Sodium nitrite: Phenol

2331. 1,2-Benzenediol (Pyrocatechol)

[120-80-9]

C₆H₆O₂



HCS 1980, 285


Nitric acid
See Nitric acid: Pyrocatechol

MRH 5.73/74

Other reactants
Yoshida, 1980, 89
MRH values calculated for 13 combinations with oxidants are given.

2332. 1,3-Benzenediol (Resorcinol)
[108-46-3]


C₆H₆O₂



Nitric acid
See Nitric acid: Resorcinol

2333. 1,4-Benzenediol (Hydroquinone)
[123-31-9]

C₆H₆O₂



HCS 1980, 554


Other reactants
Yoshida, 1980, 287
MRH values calculated for 13 combinations, largely with oxidants, are given.

Oxygen
See Oxygen (Gas): 1,4-Benzenediol

Sodium hydroxide
491M, 1975, 385
Accidental mixing of the hot crude hydroquinone with conc. sodium hydroxide solution led to extensive exothermic decomposition.
See other REDUCANTS


MRH 1.09/tr.

2334. 1,4-Benzenediol–oxygen complex (Hydroquinone–oxygen clathrate)
[20471-57-2] $C_6H_6O_2 \cdot O_2$



Preparative hazard
See Oxygen: 1,4-Benzenediol

2335. Benzeneseleninic acid
[6996-92-5] $C_6H_6O_2Se$



Preparative hazard
See Hydrogen peroxide: Diphenyl diselenide


Hydrazine derivatives

Back, T. G. *et al.*, *J. Org. Chem.*, 1981, **46**, 1568

The acid or its anhydride react very vigorously with hydrazine derivatives in absence of solvent. A solid mixture of the acid with benzohydrazide reacted violently after an induction period of several minutes.

See other INDUCTION PERIOD INCIDENTS See other ORGANIC ACIDS

2336. endo-2,3-Epoxy-7,8-dioxabicyclo[2.2.2]oct-5-ene (Oxepin-3,6-endoperoxide)
[39597-90-5] $C_6H_6O_3$



Foster, C. H. *et al.*, *J. Org. Chem.*, 1975, **40**, 3744


Evaporation of (peroxide-contacted) solvent ether from 9.8 g of crude peroxide led to an explosion. Subsequent preparations were effected on a smaller scale.

See other CYCLIC PEROXIDES, 1,2-EPOXIDES

2337. Benzeneperoxyseleninic acid

[62865-97-6]

$C_6H_6O_3Se$



Syper, L. *et al.*, *Tetrahedron*, 1987, **43**, 207–213


It explodes at the m.p., 52°C.

See other PEROXYACIDS

2338. Benzeneselenonic acid

[39254-48-3]

$C_6H_6O_3Se$



Stöcker, M. *et al.*, *Ber.*, 1906, **39**, 2917


It explodes feebly at 180°C but the silver salt is more explosive.

See other ORGANIC ACIDS

2339. 2,3:5,6-Diepoxy-7,8-dioxabicyclo[2.2.2]octane

[56411-67-7]

$C_6H_6O_4$




Sasaoka, M. *et al.*, *J. Org. Chem.*, 1979, **44**, 373 (footnote 10)

It explodes when heated.

See other CYCLIC PEROXIDES, 1,2-EPOXIDES

2340. Dimethyl acetylenedicarboxylate (Dimethyl 2-butynedioate)
[762-42-5]

C₆H₆O₄



Energy of decomposition (in range 150–430°C) measured as 1.224 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 86°C by adiabatic Dewar tests, with an apparent energy of activation of 91 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

1-Methylsilacyclopenta-2,4-diene

See 1-Methylsilacyclopenta-2,4-diene: Dienophiles

Octakis(trifluorophosphine)dirhodium


See Octakis(trifluorophosphine)dirhodium: Acetylenic esters

See other ACETYLENIC COMPOUNDS

2341. Benzeneperoxysulfonic acid

[]

C₆H₆O₄S



Preparative hazard


See Hydrogen peroxide: Benzenesulfonic anhydride

See other PEROXYACIDS

2342. *E*-Propene-1,2,3-tricarboxylic acid (Aconitic acid)

[4023-65-8]

C₆H₆O₆



Hydrogen peroxide

See Hydrogen peroxide: Aconitic acid

See other ORGANIC ACIDS

2343. Benzene triozonide (Ozobenzene)

[71369-69-2]

C₆H₆O₉

(Uncertain structure)

1. Harries, C. *et al.*, *Ber.*, 1904, **37**, 3431
2. *DOC* 5, 1982, Vol. 4, 4471

The product of ozonising benzene exhaustively is extremely explosive at the slightest touch [1]. The structure is not well defined as previously thought, but is probably polymeric [2].


See Ozone: Aromatic compounds

See other OZONIDES

2344. Benzenethiol (Thiophenol)

[108-98-5]

C₆H₆S



HCS 1980, 189

Preparative hazard

1. Leuckart, R., *J. Prakt. Chem.*, 1890, **41**, 179
2. Graesser, R., private comm., 1968


During preparation of thiophenol by addition of a cold solution of potassium *O*-methylthiocarbonate to a cold solution of benzenediazonium chloride, a violent explosion accompanied by an orange flash occurred [1]. This was attributed to the formation and decomposition of bis(benzenediazo) disulfide. A preparation in which the diazonium solution was added to the 'xanthate' solution proceeded smoothly [2].

See also DIAZONIUM SULFIDES AND DERIVATIVES

2345. 4-Bromo-1,2-diaminobenzene

[1575-37-7]

C₆H₇BrN₂




T_{ait24} was determined as the rather low value of 70°C by adiabatic Dewar tests, with an apparent energy of activation of 133 kJ/mol.

See related HALOANILINES

2346. 2-Bromomethyl-5-methylfuran
[57846-03-4]


C₆H₇BrO



Dunlop, 1953, 261
It distills with violent decomposition at 70°C/33 mbar.
See other 2-HALOMETHYL-FURANS OR -THIOPHENES

2347. 4-Chloro-1,2-benzenediamine
[95-83-0]


C₆H₇ClN₂



Energy of decomposition (in range 160–260°C) measured as 0.78 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 100°C by adiabatic Dewar tests, with an apparent energy of activation of 140 kJ/mol.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See related HALOANILINES

2348. 4-Chloro-1,3-benzenediamine
[5131-60-2]

C₆H₇ClN₂




Energy of decomposition (in range 200–250°C) measured as 1.13 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 1006°C by adiabatic Dewar tests, with an apparent energy of activation of 121 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See related HALOANILINES

2349. 2-Chloro-1,4-benzenediamine
[615-66-7]

C₆H₇ClN₂



Frankenberg, P. E., *Chem. Eng. News*, 1982, **60**(3), 97


Violently explosive decomposition at the start of distillation at 165°C/33 mbar of a 2 kg sample was attributed to autocatalytic decomposition which can occur at 125°C in small samples. Increased lability of the 2-chloro substituent when hydrochloride formation has occurred is postulated as a possible cause. Caution on heating any halogenated amine is advised.

See 2-Chloroethylamine

See other HALOANILINES

2350. 4-Nitroanilinium perchlorate
[15873-50-4]

C₆H₇ClN₂O₆




Jain, S. R. *et al.*, *Combust. Flame*, 1979, **35**, 289–295

Of a series of substituted anilinium perchlorate salts, that of 4-nitroaniline showed the lowest thermal stability (DTA peak temperature 201°C) and highest shock-sensitivity.

See other NITROARYL COMPOUNDS, PERCHLORATE SALTS OF NITROGENOUS BASES

2351. 2,4-Dinitrophenylhydrazinium perchlorate
[104503-71-1]

C₆H₇ClN₄O₈



Anon., *Angew. Chem. (Nachr.)*, 1967, **15**, 78


Although solutions of the perchlorate are stable as prepared, explosive decomposition may occur during concentration by heated evaporation.

See also HYDRAZINIUM SALTS

See other POLY-NITROARYL COMPOUNDS, PERCHLORATE SALTS OF NITROGENOUS BASES

2352. 2-Chloromethyl-5-methylfuran
[52157-57-0]

C₆H₇ClO




Dunlop, 1953, 261

It is even more unstable than its lower homologue, 2-chloromethylfuran.

See other 2-HALOMETHYL-FURANS OR -THIOPHENES

2353. 3-Methoxycarbonylpropen-2-yl trifluoromethanesulfonate
[62861-57-8]


C₆H₇F₃O₅S



See entry ALLYL TRIFLUOROMETHANESULFONATES

2354. Aniline (Benzeneamine)
[62-53-3]

C₆H₇N



(MCA SD-17, 1963); NSC 409, 1979; FPA H80, 1979; HCS 1980, 156; RSC Lab. Hazard Data Sheet No. 23, 1984

Anilinium chloride

Anon., *Chem. Met. Eng.*, 1922, **27**, 1044

Heating a mixture of the components in an autoclave at 240–260°C/7.6 bar to produce diphenylamine led to a violent explosion, following a sudden increase in pressure to 17 bar. The equipment and process had been used previously and uneventfully.

Benzenediazonium-2-carboxylate

See Benzenediazonium-2-carboxylate

Benzenediazonium-2-carboxylate: Aniline, etc.

Boron trichloride

See Boron trichloride: Aniline

1-Chloro-2,3-epoxypropane

See 1-Chloro-2,3-epoxypropane: Aniline

Dibenzoyl peroxide

See Dibenzoyl peroxide: Aniline

Nitromethane

See Nitromethane: Acids, etc.

MRH 5.69/99+

Nitrous acid

See DIAZONIUM SALTS

Other reactants

Yoshida, 1980, 22

MRH values calculated for 20 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Diisopropyl peroxydicarbonate: Amines, etc.

'Fluorine nitrate': Organic materials

Fluorine: Nitrogenous bases

Hydrogen peroxide: Organic compounds (reference 2) MRH 6.44/83
Nitric acid: Aromatic amines MRH 5.98/81
Nitrosyl perchlorate: Organic materials
Ozone: Aromatic compounds MRH 11.5/72
Perchloric acid: Aniline, etc. MRH 6.65/82
Perchloryl fluoride: Nitrogenous bases
Peroxydisulfuric acid: Organic liquids
Peroxomonosulfuric acid: Aromatics
Peroxyformic acid: Organic materials
Sodium peroxide: Organic liquids, etc. MRH 2.76/93
N-HALOIMIDES: Alcohols, Amines

Tetranitromethane

See Tetranitromethane: Amines

Trichloronitromethane


MRH 2.38/64

See Trichloronitromethane: Aniline

2355. 2-Methylpyridine (2-Picoline)

[109-06-8]

C₆H₇N



HCS 1980, 754

Hydrogen peroxide, Iron(II) sulfate, Sulfuric acid


See Hydrogen peroxide: Iron(II) sulfate, etc.

See other ORGANIC BASES

2356. *N*-Phenylhydroxylamine (*N*-Hydroxybenzeneamine)

[100-65-2]

C₆H₇NO




Preparative hazard

See Zinc: Nitrobenzene

See Phenylhydroxylaminium chloride *See other* N-O COMPOUNDS, ORGANIC BASES

2357. 4-Oximino-4,5,6,7-tetrahydrobenzofurazan N -oxide
[]

C₆H₇N₃O₃




Preparative hazard

See Sulfinyl chloride: 1,2,3-Cyclohexanetrione trioxime, etc.

See other FURAZAN N-OXIDES, OXIMES

2358. 5,6-Dinitro-2-dimethylaminopyrimidinone
[]

C₆H₇N₅O₅




Tennant, G. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 61

It decomposed violently on melting at 190°C.

See related POLYNITROARYL COMPOUNDS

2359. Ethyl 2-cyano-2-(1-H-tetrazol-5-ylhydrazono)acetate
[61033-15-6]

C₆H₇N₇O₂




Preparative hazard

See 5-Aminotetrazole (reference 3)

See other CYANO COMPOUNDS, TETRAZOLES

2360. Phenylphosphine
[638-21-1]

C₆H₇P




Haszeldine, R. N. *et al.*, *J. Fluorine Chem.*, 1977, **10**, 38

It is pyrophoric in air.

See related ALKYLPHOSPHINES

†**2361. 1,3-Cyclohexadiene**
[592-57-4]

C₆H₈



1. Bodendorf, K., *Arch. Pharm.*, 1933, **271**, 11

2. Hock, H. *et al.*, *Chem. Ber.*, 1951, **84**, 349


Cyclohexadiene autoxidises slowly in air, but the residual (largely polymeric) peroxide explodes very violently on ignition [1]. The monomeric peroxide has also been isolated [2].

See Propene

See other DIENES, POLYPEROXIDES

†**2362. 1,4-Cyclohexadiene**
[628-41-1]

C₆H₈




See Propene

See other DIENES

2363. Methylcyclopentadiene
[26519-91-5]


C₆H₈



Bis(pentafluorophenyl)ytterbium
See Bis(pentafluorophenyl)ytterbium: Methylcyclopentadiene
See other DIENES

2364. 2,2-Dibromo-1,3-dimethylcyclopropanoic acid
[72957-64-3]


C₆H₈Br₂O₂



tert-Butylamine
Baird, M. S. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1979, 2324
Interaction is highly exothermic.
See other ORGANIC ACIDS

2365. Anilinium chloride
[142-04-1]


C₆H₈ClN



HCS 1980, 157

Aniline
See Aniline: Anilinium chloride

2366. Phenylhydroxylaminium chloride
[22755-09-5]




MCA Guide, 1972, 299

A brown glass bottle containing 700 g of the salt exploded forcefully after shelf storage for 2 weeks.

See HYDROXYLAMINIUM SALTS See other N-O COMPOUNDS

2367. Anilinium perchlorate
[14796-11-3]




Metal oxides

Rao, M. V. *et al.*, *Combust. Flame*, 1980, **39**, 63–68

The effect of metal oxides in sensitising the thermal decomposition and explosion of the salt is in the order: manganese dioxide > copper oxide > nickel oxide.

See other AMINIUM PERCHLORATES, PERCHLORATE SALTS OF NITROGENOUS BASES

2368. 2-Azatricyclo[2.2.1.0^{2,6}]hept-7-yl perchlorate
[98566-27-9]




Durrant, M. L. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1985, 1879

A crystalline sample exploded without warning in the freezer compartment of a refrigerator.

See related ALKYL PERCHLORATES, PERCHLORATE SALTS OF NITROGENOUS BASES, STRAINED-RING COMPOUNDS

2369. endo-2,5-Dichloro-7-thiabicyclo[2.2.1]heptane
[6522-40-3]

C₆H₈Cl₂S



Dimethylformamide, Sodium tetrahydroborate

1. Corey, E. J. *et al.*, *J. Org. Chem.*, 1969, **34**, 1234 (footnote 5)
2. Paquette, L. A., private comm., 1975
3. Brown, H. C. *et al.*, *J. Org. Chem.*, 1962, **27**, 1928–1929

To effect reduction to the parent heterocycle, a solution of the dichloro compound in DMF was being added to a hot solution of sodium tetrahydroborate in the same solvent, when a violent explosion occurred [1,2]. This may have arisen either from interaction of the dichloro compound with the solvent, or from the known instability of hot solutions of the tetrahydroborate in DMF. Use of aqueous diglyme as an alternative solvent [3] would only be applicable to this and other hydrolytically stable halides [2].

See Dimethylformamide: Halocarbons


Sodium tetrahydroborate: Dimethylformamide

See HALOCARBONS

2370. N,N,N',N'-Tetrachloroadipamide

[]

C₆H₈Cl₄N₂O₂




Water

See Dichloramine See other N-HALOGEN COMPOUNDS

2371. 1,4-Diaminobenzene (p-Phenylenediamine)

[106-50-3]

C₆H₈N₂




The finely powdered base is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)
See other ORGANIC BASES

2372. 1,4-Diisocyanobutane
[929-25-9]


C₆H₈N₂



See entry DIISOCYANIDE LIGANDS See related CYANO COMPOUNDS

2373. Phenylhydrazine
[100-63-0]

C₆H₈N₂



HCS 1980, 735

et al., *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

Energy of decomposition (in range 280–375°C) measured as 0.662 kJ/g by DSC, and T_{ait24} was determined as 227°C by adiabatic Dewar tests, with an apparent energy of activation of 165 kJ/mol.

Heat of decomposition was determined as 0.65 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

Lead(IV) oxide

MRH 0.46/95

See Lead(IV) oxide: Nitrogen compounds

Other reactants

Yoshida, 1980, 294

MRH values calculated for 13 combinations with oxidants are given.

Perchloryl fluoride

See Perchloryl fluoride: Nitrogenous bases


α -Phenylazo hydroperoxide

See α -Phenylazo hydroperoxide: Phenylhydrazine

See other ORGANIC BASES, REDUCANTS

2374. 2,3-Diazabicyclo[2.2.2]octa-2,5-diene N-oxide
[37436-17-2]

C₆H₈N₂O




Preparative hazard

See Hydrogen peroxide: 4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2,6}]undeca-8-ene-3,5-dione, etc.

See other N-OXIDES

2375. 2-Diazocyclohexanone
[3242-56-6]

C₆H₈N₂O




Regitz, M. *et al.*, *Org. Synth.*, 1971, **51**, 86

It may explode on being heated, and a relatively large residue should be left during vacuum distillation to prevent overheating.

See other DIAZO COMPOUNDS

2376. Butane-1,4-diisocyanate
[4538-37-8]

C₆H₈N₂O₂




Zlobin, V. A. *et al.*, *Chem. Abs.*, 1987, **107**, 39144

Conditions for the safe preparation of the diisocyanate from adipoyl chloride and sodium azide in acetonitrile-toluene mixtures were established.

See other ORGANIC ISOCYANATES

2377. 2-Buten-1-yl diazoacetate
[14746-03-3]

C₆H₈N₂O₂




Blankley, C. J. *et al.*, *Org. Synth.*, 1969, **49**, 25

Precautions are necessary during vacuum distillation of this potentially explosive material.

See other DIAZO COMPOUNDS

2378. 2,5-Dimethylpyrazine 1,4-dioxide
[6890-38-6]

C₆H₈N₂O₂




Preparative hazard

See Hydrogen peroxide: Acetic acid, *N*-Heterocycles

See other *N*-OXIDES

2379. Anilinium nitrate
[542-15-4]

C₆H₈N₂O₃




Nitric acid

See Nitric acid: Anilinium nitrate

See other OXOSALTS OF NITROGENOUS BASES

2380. Isosorbide dinitrate (1,4:3,6-Dianhydro-D-glucitol dinitrate)
[87-33-2]

$C_6H_8N_2O_8$




Reddy, G. O. *et al.*, *J. Hazard. Mater.*, 1992, **32**(1), 87
This heart drug is detonable when dry but non-explosive with 30% of water.
See other ALKYL NITRATES

2381.

Dodecahydro-4,8-dinitro-2,6-bis(nitroimino)diimidazo[4,5-*b*:4',5'-*e*]pyrazine
[134927-04-1] [160693-88-2] (stereoisomer)


$C_6H_8N_{12}O_8$



Dagley, I. J. *et al.*, *J. Energ. Mater.*, 1995, **13**(1 & 2), 35
This nitroguanidine analogue is more sensitive than RDX to friction, but equivalent by the drop-weight impact test.
See other N-NITRO COMPOUNDS

†2382. 2,5-Dimethylfuran
[625-86-5]

C_6H_8O



2383. 2,4-Hexadienal
[142-83-6]

C_6H_8O




1-Amino-2-propanol
Steele, A. B. *et al.*, *Chem. Engrg.*, 1959, **66**(8), 166

Following a plant incident in which the crude aldehyde had polymerised during distillation, it was found that the presence of 2% of isopropanolamine (a probable impurity) led to an explosive polymerisation reaction. It is conceivable that aldehydic peroxides may also have been involved.

See other PEROXIDISABLE COMPOUNDS, POLYMERISATION INCIDENTS

2384. 3-Methyl-2-penten-4-yn-1-ol
[105-29-3]

C₆H₈O



1. Lorentz, F., *Loss Prev.*, 1967, **1**, 1-5
2. Fesenko, G. V. *et al.*, *Chem. Abs.*, 1975, **83**, 47605

Temperature control during pressure hydrogenation of *cis*- or *trans*-isomers is essential, since at 155°C violent decomposition to carbon, hydrogen and carbon monoxide with development of over 1 kbar pressure will occur. The material should not be heated above 100°C, particularly if acid or base is present, to avoid exothermic polymerisation [1]. The *cis*-isomer is readily cyclised to 2,3-dimethylfuran, which promotes fire and explosion hazards. These were measured for the *cis*- and *trans*-isomers, and for *trans*-3-methyl-1-penten-4-yn-3-ol [2].

See other GAS EVOLUTION INCIDENTS, HYDROGENATION INCIDENTS, POLYMERISATION INCIDENTS

Sodium hydroxide


1. *MCA Case History No. 363*
2. Silver, L., *Chem. Eng. Progr.*, 1967, **63**(8), 43

Presence of traces of sodium hydroxide probably caused formation of the acetylenic sodium salt, which exploded during high-vacuum distillation in a metal still [1]. A laboratory investigation which duplicated the explosion, without revealing the precise cause, was also reported [2].

See other ACETYLENIC COMPOUNDS

2385a. 1,2-Cyclohexanedione
[765-87-7]

C₆H₈O₂



Preparative hazard

See Nitric acid: 4-Methylcyclohexanone

2385b. 2,4-Hexadienoic acid (Sorbic acid)

[110-44-1]



$\text{C}_6\text{H}_8\text{O}_2$

The dry finely powdered acid is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

See other ORGANIC ACIDS

2386. Poly(1,3-cyclohexadiene peroxide)

[]

(Complex structure)

$(\text{C}_6\text{H}_8\text{O}_2)_n$


See 1,3-Cyclohexadiene

See other POLYPEROXIDES

2387. 2,5-Dimethyl-2,5-dihydrothiophene-2,5-endoperoxide

[67711-62-0]

$\text{C}_6\text{H}_8\text{O}_2\text{S}$



Golnick, K. *et al.*, *Tetrahedron Lett.*, 1984, **25**, 4291–4294

The endoperoxide of 2,5-dimethylthiophene (effectively a thiazonide) when neat decomposes violently at ambient temperature.


See other CYCLIC PEROXIDES

See related OZONIDES

2388. 1,4-Dimethyl-2,3,7-trioxabicyclo[2.2.1]hept-5-ene (2,5-Dimethyl-2,5-dihydrofuran-2,5-endoperoxide)

[45722-89-2]

$\text{C}_6\text{H}_8\text{O}_3$




Gollnick, K. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1983, **22**, 726

The solvent-free peroxide always explodes if amounts above 100 mg are handled.

See other CYCLIC PEROXIDES

2389. Citric acid (2-Hydroxypropane-1,2,3-tricarboxylic acid)
[77-92-9]

C₆H₈O₇




Metal nitrates

See METAL NITRATES: citric acid *See other* ORGANIC ACIDS

2390. Triacetyl borate
[4887-24-5]

C₆H₉BO₆




Preparative hazard

See Acetic anhydride: Boric acid

2391. Trivinylbismuth
[65313-35-1]

C₆H₉Bi




Coates, 1967, Vol. 1, 538

It ignites in air.

See related ALKYLMETALS, TRIALKYLBISMUTHS

2392. 2-Bromocyclohexanone
[822-85-5]


C₆H₉BrO



Energy of decomposition (in range 80–130°C) measured as 0.29 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

2393. 1-Iodohepta-2,4-diene
[]


C₆H₉I



Hayashi, T., *et al.*, *J. Chem. Soc., Chem. Comm.*, 1990, (19), 1362
This compound undergoes explosive decomposition when freed of solvent. Homologues are also unstable.
See other DIENES, HALOALKENES, IODINE COMPOUNDS

2394. Iodine triacetate
[6540-76-7]


C₆H₉IO₆



1. Schutzenberger, P., *Compt. rend.*, 1861, **52**, 135
2. Oldham, F. W. H. *et al.*, *J. Chem. Soc.*, 1941, 368
The acetate explodes at 140°C [1], and higher homologues decompose at about 120°C, the violence decreasing with ascent of the homologous series [2].
See other IODINE COMPOUNDS

2395. 3,3-Dimethyl-1-nitro-1-butyne (tert-Butylnitroacetylene)
[22691-91-4]

C₆H₉NO₂



Amines


Jäger, V. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1969, **8**, 273

Reaction with primary, secondary or tertiary amines proceeds explosively with ignition in absence of a solvent.

See other ACETYLENIC COMPOUNDS See related NITROALKENES

2396. 1,4,3,6-Dianhydroglucitol 2-nitrate (Isosorbide mononitrate)
[16016-20-0]

C₆H₉NO₆



Négyesi, G., *Process Safety Progress*, 1996, **15**(1), 42


Lead block tests show this to be a feeble explosive, it is not impact sensitive but shows some decomposition on friction.

See Isosorbide dinitrate

See other ALKYL NITRATES

2397. Bis(2-cyanoethyl)amine
[111-94-4]

C₆H₉N₃




Anon., *BCISC Quart. Safety Summ.*, 1967, **38**, 42

Some 18-month old bottles of 'iminodipropionitrile' exploded, probably owing to slow hydrolysis and release of ammonia.

See other CYANO COMPOUNDS, GAS EVOLUTION INCIDENTS

2398. 3,6,9-Triazatetracyclo[6.1.0.0^{2,4}.0^{5,7}]nonane (cis-Benzene triimine)
[52851-26-0]

C₆H₉N₃




Schwesinger, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1973, **12**, 989

It explodes if rapidly heated to 200°C.

See other STRAINED-RING COMPOUNDS

2399. Ammonium *N*-nitrosophenylaminoxide ('Cupferron')
[135-20-6]




Thorium salts

Pittwell, L. R., *J. R. Inst. Chem.*, 1956, **80**, 173

Solutions of this reagent are destabilised by the presence of thorium ions. If a working temperature of 10–15°C is much exceeded, the risk of decomposition, not slowed by cooling and accelerating to explosion, exists. Titanium and zirconium salts also cause slight destabilisation, but decomposition temperatures are then 35 and 40°C, respectively.

See other NITROSO COMPOUNDS, N–O COMPOUNDS

2400. 1,2,3-Cyclohexanetrione trioxime
[3570-93-2]




Sulfinyl chloride

See Sulfinyl chloride: 1,2,3-Cyclohexanetrione trioxime, etc.

See other OXIMES

2401. 1,3,5-Cyclohexanetrione trioxime
[622-22-7]




Baeyer, A., *Ber.*, 1886, **19**, 160

The oxime of phloroglucinol explodes rather violently at 155°C.
See other OXIMES

2402. Lead(IV) acetate azide
[]


C₆H₉N₃O₆Pb



See Lead(IV) azide
See related METAL AZIDES

2403. 3(2,3-Epoxypropoxy)2,2-dinitropropyl azide
[76828-34-7]


C₆H₉N₅O₆



Frankel, M. B. *et al.*, *J. Chem. Eng. Data*, 1981, **26**, 219
The monomeric azide and several precursors are explosives of moderate to considerable sensitivity to initiation by impact, shock, friction or heat.
See other 1,2-EPOXIDES, ORGANIC AZIDES, POLYNITROALKYL COMPOUNDS

2404. Ethyl sodioacetoacetate
[19232-39-4]


C₆H₉NaO₃



2-Iodo-3,5-dinitrophenyl
See 2-Iodo-3,5-dinitrophenyl: Ethyl sodioacetoacetate

2405. Trivinylantimony
[5613-68-3]


C₆H₉Sb



Leleu, *Cahiers*, 1977, (88), 362
It ignites in air.
See related ALKYL METALS

†**2406. Cyclohexene**
[110-83-8]

C₆H₁₀




HCS 1980, 346

See other ALKENES

2407. 2,3-Dimethyl-1,3-butadiene
[513-81-5]

C₆H₁₀




Bodendorf, K., *Arch. Pharm.*, 1933, **271**, 33
A few mg of the autoxidation residue, a largely polymeric peroxide, exploded violently on ignition.
See other POLYPEROXIDES

Oxygen, Ozone
See Ozone: Dienes, Oxygen

Thiazyl fluoride
See Thiazyl fluoride: Alkylbutadienes
See other DIENES

2408. 3,3-Dimethyl-1-butyne
[917-92-0]

C₆H₁₀



Preparative hazard

See 2,2-Dichloro-3,3-dimethylbutane: Sodium hydroxide

See other ALKYNES

†**2409. 1,3-Hexadiene**
[592-48-3]

C₆H₁₀



See other DIENES

†**2410. 1,4-Hexadiene**
[592-45-0]


C₆H₁₀



See other DIENES

†**2411. 1,5-Hexadiene**
[592-42-7]

C₆H₁₀



See other DIENES

†**2412. cis-2-trans-4-Hexadiene**
[5194-50-3]

C₆H₁₀



See other DIENES

†**2413. trans-2-trans-4-Hexadiene**
[5194-51-4]


C₆H₁₀



See other DIENES

†**2414. 1-Hexyne**
[693-02-7]


C₆H₁₀



See other ALKYNES

†2415. 3-Hexyne
[928-49-4]

C_6H_{10}



Mercury, Silver perchlorate
See Silver perchlorate: Alkynes, etc.
See other ALKYNES

†2416. 2-Methyl-1,3-pentadiene
[1118-58-7]

C_6H_{10}



See other DIENES

†2417. 4-Methyl-1,3-pentadiene
[926-56-7]


C_6H_{10}



See other DIENES

2418. Cadmium propionate
[16986-83-7]

$C_6H_{10}CdO_4$




Anon., *Chem. Age*, 1957, **77**, 794

The salt exploded during drying at 60–100°C in an electric oven, presumably because of overheating which caused pyrolysis to 3-pentanone and ignition of its vapour.

See other HEAVY METAL DERIVATIVES

2419. 1-Chloro-1-nitrosocyclohexane
[695-64-7]

$C_6H_{10}ClNO$




Labaziewicz, H. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1979, 2928

There is a risk of explosion if the pressure rises during its vacuum distillation.
See other NITROSO COMPOUNDS

2420. *trans*-2-Chlorocyclohexyl perchlorate
[81971-84-8]


C₆H₁₀Cl₂O₄



See entry ALKYL PERCHLORATES (reference 6)

2421. Bis(2-fluoro-2,2-dinitroethoxy)dimethylsilane
[73526-98-4]

C₆H₁₀F₂N₄O₁₀Si




Neale, R. S., *Ind. Eng., Chem., Prod. Res. Dev.*, 1980, **19**, 634

This substituted silane (intended as an explosive plasticiser), though relatively stable to heat and impact of a hammer, is exceedingly sensitive to compressive shock. Great care must therefore be exercised in syringing samples of the viscous liquid.

See other ALKYL SILANES, FLUORODINITROMETHYL COMPOUNDS

2422. 3,3-Pentamethylenediazirine
[930-82-5]

C₆H₁₀N₂




1. Schmitz, E. *et al.*, *Org. Synth.*, 1965, **45**, 85
2. Schmitz, E. *et al.*, *Chem. Ber.*, 1961, **94**, 2168

Small-scale preparation is recommended [1], in view of a previous explosion through superheating of the liquid during distillation at 109°C [2].

See other DIAZIRINES

2423. *tert*-Butyl diazoacetate
[35059-50-8]

$C_6H_{10}N_2O_2$




Regitz, M. et al., *Org. Synth.*, 1968, **48**, 36

Vacuum distillation of the compound is potentially hazardous.

See other DIAZO COMPOUNDS

2424. *N*-Nitroso-6-hexanelactam
[35784-01-1]

$C_6H_{10}N_2O_2$




Preparative hazard

See 6-Hexanelactam: Acetic acid, etc.

See other NITROSO COMPOUNDS

2425. Diethyl azoformate (1,2-Bisethoxycarbonyldiazene)
[1972-28-7]

$C_6H_{10}N_2O_4$



1. US Pat. 3 347 845, 1967
2. Kauer, J. C., *Org. Synth.*, 1963, Coll. Vol. 4, 412
3. Houben Weyl, 1967, 10/2, 809

Shock-sensitive [1], it explodes on heating in a (sealed?) capillary, like its lower homologue. The crude product prepared by chlorine oxidation of the corresponding


hydrazide may detonate violently during distillation. This is thought due to either excess chlorine or inadequate washing leaving chloramine byproducts [3]. However, the editor believes the alternative preparation of azodicarboxylates by fuming nitric acid oxidation of the hydrazide [2] can give azide and nitramine impurities; crude methyl ester from this route has also exploded during distillation.

See Dimethyl azoformate

See other AZO COMPOUNDS

2426. *N,N'*-Diacetyl-*N,N'*-dinitro-1,2-diaminoethane
[922-89-4]

C₆H₁₀N₄O₆




Violent decomposition occurred at 142°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (reference 1) See other *N*-NITRO COMPOUNDS

2427. *trans*-1,4,5,8-Tetranitro-1,4,5,8-tetraazadecahydronaphthalene
[83673-81-8]

C₆H₁₀N₈O₈



Willer, R. L., *Propellants, Explos. Pyrotech.*, 1983, **8**, 65

A new explosive compound.

See other *N*-NITRO COMPOUNDS

2428. Butoxyacetylene
[3329-56-4]

C₆H₁₀O

BuOC≡CH


Jacobs, T. L. *et al.*, *J. Amer. Chem. Soc.*, 1942, **64**, 223

Small samples rapidly heated in sealed tubes to around 100°C exploded.

See other ACETYLENIC COMPOUNDS

2429. Cyclohexanone
[108-94-1]

C₆H₁₀O



HCS 1980, 344; RSC Lab. Hazards Data Sheet No. 80, 1989


Hydrogen peroxide, Nitric acid
See Hydrogen peroxide: Ketones, etc.

Nitric acid
See Nitric acid: 4-Methylcyclohexanone

Other reactants
Yoshida, 1980, 152
MRH values calculated for 14 combinations with oxidants are given.

2430. Diethylketene (2-Ethyl-1-butene-1-one)
[24264-08-2]

C₆H₁₀O



Sorbe, 1968, 118
Like the lower homologue, it readily forms explosive peroxides with air at ambient temperature.
See other PEROXIDISABLE COMPOUNDS

†**2431. Diallyl ether (Di-2-propenyl ether)**
[557-40-4]

C₆H₁₀O



HCS 1980, 126

Preparative hazard
MCA Case History No. 412
The ether was left exposed to air and sunlight for 2 weeks before distillation, and became peroxidised. During distillation to small bulk, a violent explosion occurred.

See 2-Propen-1-ol: Sulfuric acid

See other ALLYL COMPOUNDS, PEROXIDATION INCIDENTS, PEROXIDISABLE COMPOUNDS

2432. 2-Hexenal
[6728-26-3]




$\text{C}_6\text{H}_{10}\text{O}$

Nitric acid

See Nitric acid: 2-Hexenal

See other PEROXIDISABLE COMPOUNDS

†**2433. 2-3-Methyl-2-methylenebutanal (Isopropylacrylaldehyde)**
[4417-80-5]




$\text{C}_6\text{H}_{10}\text{O}$

See other PEROXIDISABLE COMPOUNDS

2434. 1-Allyloxy-2,3-epoxypropane (Allyl glycidyl ether, 2-Propenyloxy-methyloxirane)
[106-92-3]

$\text{C}_6\text{H}_{10}\text{O}_2$



Peroxide test strips

Anon., *Jahresber.*, 1987, 65

Peroxide test strips may not be sufficiently reliable as the sole means of testing for presence of peroxides in a wide range of susceptible liquids.

A sample of the readily peroxidisable allyl ether, tested with a peroxide test strip gave a negative test result, so was charged into a distillation apparatus, but there was a detonation during distillation. It was recommended that an additional test (potassium iodide–acetic acid) should be used unless it is certain that the test strips will give a positive result with a particular peroxide-containing liquid.

See PEROXIDES IN SOLVENTS, PEROXIDISABLE COMPOUNDS


Poly[oxy(methyl)silylene]

See Poly[oxy(methyl)silylene]: 1-Allyloxy-2,3-epoxypropane

See other ALLYL COMPOUNDS, 1,2-EPOXIDES

2435. 2-Cyclohexenyl hydroperoxide
[4845-05-0]

C₆H₁₀O₂



Leleu, *Cahiers*, 1973, (71), 226

Accidental contact of a mixture containing the peroxide with the hot top of the freshly sealed glass container led to an explosion.

See other ALKYL HYDROPEROXIDES


†2436. Ethyl crotonate
[623-70-1]

C₆H₁₀O₂



†2437. Ethyl cyclopropanecarboxylate
[4606-07-9]

C₆H₁₀O₂



See other STRAINED-RING COMPOUNDS

†2438. Ethyl methacrylate
[97-63-2]

C₆H₁₀O₂




†2439. Vinyl butyrate
[123-20-6]

C₆H₁₀O₂



2440. Ethyl acetoacetate (Ethyl 3-oxobutanoate)
[141-97-9]

C₆H₁₀O₃



HCS 1980, 463

2,2,2-Tris(bromomethyl)ethanol, Zinc

1. Reeder, J. A., U S Pat. 3 578 619, 1971
2. *491M*, 1975, 167

The patent describes the formation of complex metal chelates by treatment of the ketoester simultaneously with an alcohol and a metal to effect trans-esterification and chelate formation by distilling out the by-product ethanol [1]. This process was being applied to produce the zinc chelate of 2-tris(bromomethyl)ethyl acetoacetate, and when 80% of the ethanol had been distilled out (and the internal temperature had increased considerably), a violent decomposition occurred [2]. This presumably involved interaction of a bromine substituent with excess zinc to form a Grignard-type reagent, and subsequent exothermic reaction of this with one or more of the bromo or ester functions present.

See related GRIGNARD REAGENTS

2441. Adipic acid (Hexane-1,6-dioic acid)

[124-04-9]

$C_6H_{10}O_4$



Kaiser, M. A., *Plant/Oper. Progr.*, 1991, **10**(2), 100


After two minor dust explosions in an industrial adipic acid dryer, evidence was obtained that adipic acid forms an iron complex capable of both decarboxylation/dehydration of adipic acid to cyclopentanone and of catalysing air oxidation, giving exotherms from as low as 135°C.

See other CATALYTIC IMPURITY INCIDENTS, SELF-HEATING AND IGNITION INCIDENTS, ORGANIC ACIDS

2442. Dipropionyl peroxide

[3248-28-0]

$C_6H_{10}O_4$



Swern, 1971, Vol. 2, 815

Pure material explodes on standing at ambient temperature.

Other reactants


Yoshida, 1980, 317

MRH values calculated for 13 combinations, largely with materials catalysing its decomposition, are given.

See other DIACYL PEROXIDES

2443. Diallyl sulfate
[27063-40-7]

C₆H₁₀O₄S



von Braun, J. *et al.*, *Ber.*, 1917, **50**, 293

The explosive decomposition during distillation may well have been caused by polymerisation initiated by acidic decomposition products.

See other ALLYL COMPOUNDS, POLYMERISATION INCIDENTS, SULFUR ESTERS

2444. Diethyl dicarbonate (Ethyl pyrocarbonate)
[1609-47-8]

C₆H₁₀O₅

EtOCO.OCO.OEt


Anon., *Univ. Safety Assoc. Safety Digest*, 1991, **42**, 7

Unopened bottles of this, all originating from the same batch and well within the manufacturer's expiry date, exploded while warming to room temperature. This was attributed to water contamination leading to hydrolysis and carbon dioxide formation, without supporting evidence—a number of other contaminants, both nucleophiles and acids, might catalyse carbon dioxide formation (Editor).

See other GAS EVOLUTION INCIDENTS

2445. 2-Hydroxy-2-methylglutaric acid
[503-49-1]

C₆H₁₀O₅




Preparative hazard

See 4-Hydroxy-4-methyl-1,6-heptadiene: Ozone

See other ORGANIC ACIDS

2446. Diethyl peroxydicarbonate
[14666-78-5]

C₆H₁₀O₆




An oil of extreme instability when impure, and sensitive to heat or impact, exploding more powerfully than dibenzoyl peroxide.

See entry PEROXYCARBONATE ESTERS

2447. Diallyl sulfide
[592-88-1]

C₆H₁₀S




N-Bromosuccinimide

See *N*-HALOIMIDES: alcohols, etc. See other ALLYL COMPOUNDS

2448. *N,N,N'*-Trifluorohexanamidine
[31330-22-0]

C₆H₁₁F₃N₂




Ross, D. L. *et al.*, *J. Org. Chem.*, 1970, **35**, 3096–3097

A shock-sensitive explosive.

See other *N,N,N'*-TRIFLUOROAMIDINES

2449. 2-Ethoxy-1-iodo-3-butene
[13957-21-6]

C₆H₁₁IO



Trent, J. *et al.*, *Chem. Eng. News*, 1966, **44**(33), 7

During a 1 g mol-scale preparation by a published method from butadiene, ethanol, iodine and mercury oxide, a violent explosion occurred while ethanol was being distilled off at 35°C under slight vacuum. The cause of the explosion could not be established, and several smaller-scale preparations had been uneventful.

See related HALOALKENES See other IODINE COMPOUNDS

†2450. Diallylamine
[124-02-7]

C₆H₁₁N




HCS 1980, 356

See other ALLYL COMPOUNDS, ORGANIC BASES

2451. Caprolactam (6-Hexanelactam)
[105-60-2]

C₆H₁₁NO



HCS 1980, 258


Acetic acid, Dinitrogen trioxide

Huisgen, R. *et al.*, *Ann.*, 1952, **575**, 174–197

During preparation of the *N*-nitroso derivative from the lactam in acetic acid solution, the treatment with dinitrogen trioxide must be very effectively cooled to prevent explosive decomposition.

2452. Cyclohexanone oxime
[100-64-1]

C₆H₁₁NO



HCS 1980, 345

Ullmann, 1986, **A5** 34

It can decompose violently, for unknown reasons, during vacuum distillation (probably exothermic Beckmann rearrangement and subsequent polymerisation — Editor)

Oleum (fuming sulfuric acid)


Ale, B. J. M., *Proc. 3rd Intl. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 4, 1541–1550, Basle, SSCI, 1980

During startup of the Beckmann rearrangement stage in the caprolactam process, a mixture of caprolactam and oleum is preheated to operating temperature, then oxime feed and cooling are begun. On one occasion, the exothermic rearrangement ($Q = 230.8$ kJ/mol, 2.04 kJ/g of oxime) failed to start, and the temperature decreased. On application of steam to the cooler to start the reaction, the reactor vessel burst. At temperatures above 150–160°C, oleum oxidises the oxime with evolution of carbon dioxide and sulfur trioxide. An investigation of the operating stability of the reactor system and transient behaviour during startup allowed safe starting procedures to be clearly defined.

See other GAS EVOLUTION INCIDENTS, OXIMES

2453. 2-Isopropylacrylaldehyde oxime

[]


C₆H₁₁NOMarvel, C. S. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 5408–5409

Hydroquinone must be added to the oxime before distillation, to prevent formation and subsequent violent decomposition of a peroxide (there being 2 susceptible hydrogen atoms in the molecule).

See other OXIMES, PEROXIDISABLE COMPOUNDS

2454. Ethyl 2-formylpropionate oxime

[]

C₆H₁₁NO₃

Hydrogen chloride (?)


Loftus, F., private comm., 1972

The formyl ester and hydroxylamine were allowed to react in ethanol, which was then removed by vacuum evaporation at 40°C. The dry residue decomposed violently 5 min later. Analysis of the residue suggested that the oxime had undergone an exothermic Beckmann rearrangement, possibly catalysed by traces of hydrogen chloride in the reaction residue.

See Butyraldehyde oxime

See other OXIMES


See other CATALYTIC IMPURITY INCIDENTS

2455. 5-Methyl-1(1-methylethyl)-1,2,3-azadiphosphole (3(1-Methylethyl)-4-methyldiphosphazole)**126330-30-1****C₆H₁₁NP₂**Gueth, W., *New J. Chem.* 1989, **13**(4–5), 309

Extremely pyrophoric
See other PYROPHORIC MATERIALS, *See related* ALKYLPHOSPHINES

2456. Diallyl phosphite (Di-2-propenyl phosphonite)
[23679-20-1]


C₆H₁₁O₃P



Houben-Weyl, 1964, Vol. 12.2, 22
The ester is liable to explode during distillation unless thoroughly dry allyl alcohol is used for the preparation, and if more than 60% of the product is distilled over. Acid-catalysed polymerisation is probably involved.
See other ALLYL COMPOUNDS, PHOSPHORUS ESTERS, POLYMERISATION INCIDENTS

†2457. Cyclohexane
[110-82-7]

C₆H₁₂




(MCA SD-68, 1957); FPA H25, 1974; HCS 1980, 342; RSC Lab. Hazards Data Sheet No. 54, 1986

Dinitrogen tetroxide
See Dinitrogen tetroxide: Hydrocarbons

Other reactants
Yoshida, 1980, 153
MRH values calculated for 14 combinations with oxidants are given.


†2458. Ethylcyclobutane
[4806-61-5]

C₆H₁₂




†2459. 1-Hexene
[592-41-6]

C₆H₁₂



See other ALKENES


†2460. 2-Hexene
[592-43-8]



C_6H_{12}


See other ALKENES

†2461. Methylcyclopentane
[96-37-7]



C_6H_{12}

†2462. 2-Methyl-1-pentene
[763-29-1]



C_6H_{12}

See other ALKENES


†2463. 4-Methyl-1-pentene
[691-37-2]



C_6H_{12}

See other ALKENES

†2464. *cis*-4-Methyl-2-pentene
[691-38-3]



C_6H_{12}

See other ALKENES

†2465. *trans*-4-Methyl-2-pentene
[4461-48-7]



C_6H_{12}

See other ALKENES

2466. Diethylaluminium 2,2,2-trifluoroethoxide
[116778-64-4]



$\text{C}_6\text{H}_{12}\text{AlF}_3\text{O}$


Malpass, D. B., *Chem. Eng. News*, 1990, **68**(14), 2

This compound was found, on investigation of an explosion in a syringe during transfer, to have a decomposition energy equivalent to that of commercial explosives. Slow decomposition, even at room temperature, becomes explosive above 100°C. Shock sensitive. Stable as hydrocarbon solution below 25% concentration. Recommended that this and related compounds be handled only in such solution.

See FLUORINATED ORGANOLITHIUM COMPOUND

See related ALKYLALUMINIUM ALKOXIDES AND HYDRIDES

2467. Bis(dimethylarsinyldiazomethyl)mercury
[63382-64-9]





Glozbach, E. *et al.*, *J. Organomet. Chem.*, 1977, **132**, 359

It is explosive.

See related ALKYLNON-METALS *See other* DIAZO COMPOUNDS, MERCURY COMPOUNDS

2468. N-Ethyl-N-propylcarbamoyl chloride
[98456-61-2]




Water

Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 78

The chloride (60 l) and 2 volumes of water were mixed ready for subsequent addition of alkali to effect controlled hydrolysis. Before alkali was added, the internal temperature rose from 15 to 25° during 30 min, and then to 35°C in 5 min, when gas was suddenly evolved. This was attributed to the effect of liberated hydrochloric acid causing autocatalytic acceleration of the hydrolysis and then rapid release of carbon dioxide arising from decarboxylation of the carbamic acid. Hydrolysis by addition of the chloride to excess alkali would prevent the gas evolution.

See other ACYL HALIDES, GAS EVOLUTION INCIDENTS

2469. 2,2-Dichloro-3,3-dimethylbutane
[594-84-3]



Sodium hydroxide


1. Bartlett, P. D. *et al.*, *J. Amer. Chem. Soc.*, 1942, **64**, 543

2. Kocienski, P. J., *J. Org. Chem.*, 1974, **39**, 3285–3286

A previous method [1] of preparing 3,3-dimethylbutyne by dehydrochlorination of the title compound in a sodium hydroxide melt is difficult to control and hazardous on the large scale. Use of potassium *tert*-butoxide as base in DMSO is a high-yielding, safe and convenient alternative method of preparation of the alkyne [2].
See Dimethyl sulfoxide: Metal alkoxides


See other HALOALKANES

2470. 1,6-Hexanediyl perchlorate
[95407-64-0]



See entry ALKYL PERCHLORATES (reference 6)

2471. Hexamethylenetetramine tetraiodide
[12001-65-9]




Merck, 1976, 779

The hexamine–iodine complex deflagrates at 138°C.

See other IODINE COMPOUNDS

2472. (Oxidiperoxy(1,3-dimethyl-2,4,5,6-tetrahydro-2-1H)-pyrimidinone) molybdenum
[128568-96-7]




Paquette, L. A. *et al.*, *Chem. Eng. News*, 1992, **70**(37), 2

A well dried sample of this complex exploded violently when pushed through a funnel with a steel spatula. The pyridine adduct, which was being prepared, is claimed to be a useful, safe, oxidant, replacing the hexamethylphosphoramide complex of MoO_5 (*ibid.*).

See other METAL PEROXIDES

2473. 1,4-Diazabicyclo[2.2.2]octane
[280-57-9]



An exceptionally strong base.

Carbon

See Carbon: 1,4-Diazabicyclo[2.2.2]octane



Cellulose nitrate

See CELLULOSE NITRATE: amines

Hydrogen peroxide

See next below *See other* ORGANIC BASES

2474. 1,4-Diazabicyclo[2.2.2]octane hydrogen peroxidate
[38910-25-8]




Davies, A. G. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1981, 1512

A sample of the 1:1 complex exploded while being dried overnight in a desiccator.
See other CRYSTALLINE HYDROGEN PEROXIDATES

2475. 1,3-Dimethylhexahydropyrimidone (Dimethylpropyleneurea)
[7226-23-5]

C₆H₁₂N₂O




Chromium trioxide

See Chromium trioxide: 1,3-Dimethylhexahydropyrimidone

2476. 1,6-Diaza-3,4,8,9,12,13-hexaoxabicyclo[4.4.4]tetradecane ('Hexamethylene-triperoxydiamine')
[283-66-9]

C₆H₁₂N₂O₆



Alone, or Bromine, or Sulfuric acid


Leulier, *J. Pharm. Chim.*, 1917, **15**, 222–229

The compound, precipitated by interaction of hexamethylene tetramine and acidic 30% hydrogen peroxide, is a heat- and shock-sensitive powerful explosive when dry, much more shock sensitive than mercury fulminate. It explodes in contact with bromine or sulfuric acid.

See other CYCLIC PEROXIDES

2477. Hexamethylenetetramine (1,3,5,7-Tetraazatricyclo[3.3.1.1^{3,7}]decane)
[100-97-0]

C₆H₁₂N₄



Finely powdered dry hexamine is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

Acetic acid, Acetic anhydride, Ammonium nitrate, Nitric acid

See Nitric acid: Acetic anhydride, Hexamethylenetetramine acetate

1-Bromopentaborane(9)

See 1-Bromopentaborane(9): Air, or Hexamethylenetetramine

Hydrogen peroxide

See 1,6-Diaza-3,4,8,9,12,13-hexaoxabicyclo[4.4.4]tetradecane

Iodine

See Hexamethylenetetramine tetraiodide


Iodoform

See Iodoform: Hexamethylenetetramine

See other ORGANIC BASES

2478. 4a,8a,9a,10a-Tetraaza-2,3,6,7-tetraoxaperhydroanthracene
[262-38-4]

$C_6H_{12}N_4O_4$




Katritzky, A. R. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1979, 1136

The bis-peroxy tricyclic heterocycle explodes at 120°C.

See other CYCLIC PEROXIDES

2479. 1,2-Bis(2-azidoethoxy)ethane
[59559-04-7]

$C_6H_{12}N_6O_2$



Ethanol, Hydrogen sulfide


Anon., *Lab. Accid. Higher Educ.* Item 32, HSE, Barking, 1987

During the reduction of the azide in ethanol with hydrogen sulfide (no regulation other than main cylinder valve) under reflux (with insufficient flow of condenser water), the mixture apparently overheated, then exploded violently.

See other ORGANIC AZIDES

2480. 1,4-Diazabicyclo[2.2.2]octane 1,4-bis(nitroimidate)
[51470-74-7]


$C_6H_{12}N_6O_4$



See *N*-AZOLIUM NITROIMIDATES

2481. Bis(2-azidoethoxymethyl)nitramine
[88487-87-6]


$C_6H_{12}N_8O_4$



See entry ENERGETIC COMPOUNDS See other *N*-NITRO COMPOUNDS, ORGANIC AZIDES


2482. Tris(2-azidoethyl)amine
[84928-99-4]

$C_6H_{12}N_{10}$



See entry ENERGETIC COMPOUNDS See other ORGANIC AZIDES

2483. 1,1,3,3,5,5-Tris-spiro(*N,N'*-dinitroethylenediamino)cyclotriphosphazene
(1,4,8,11,14,17-Hexanitro-1,4,6,8,11,12,14,17,18-nonaaza-5λ⁵,7λ⁵,13λ⁵-
triphosphatatripiro[4.1.4.1.4.1]octadeca-5,7(12),13(18)-triene)
[155793-91-2] **C₆H₁₂N₁₅O₁₂P₃**



Paritosh, D. R. *et al.*, *Chem. Abs.*, 1994, **121**, 35869j; U.S. Pat. Appl. 42,229
 Paritosh, D. R. *et al.*, *Phosphorus, Sulfur, Silicon Relat. Elem.*, 1994, **90**(1-4), 175
 This and several other compounds with fewer nitro groups and partial replacement
 of the spiro rings by halogen are patented as explosives.
See other N-NITRO COMPOUNDS

†**2484. Butyl vinyl ether**
[111-34-2]




C₆H₁₂O

Glikin, M. A. *et al.*, *Chem. Abs.*, 1978, **89**, 30115
 'Explosion hazard of industrial synthesis of butyl vinyl ether' (title only translated).
See other PEROXIDISABLE COMPOUNDS

2485. Cyclohexanol
[108-93-0]

C₆H₁₂O



FPA H91, 1980; *HCS 1980*, 343

Other reactants

Yoshida, 1980, 151

MRH values calculated for 12 combinations with oxidants are given.

Oxidants
See Chromium trioxide: Alcohols
 Nitric acid: Alcohols

MRH values show % of oxidant
 MRH 2.55/92
 MRH 5.86/81

†2486. 3,3-Dimethyl-2-butanone
 [75-97-8] C₆H₁₂O
 MeCO.CMe₃

†2487. 2-Ethylbutanal
 [97-96-1] C₆H₁₂O
 Et₂CHCO.H

See other PEROXIDISABLE COMPOUNDS

†2488. 2-Hexanone
 [591-78-6] C₆H₁₂O
 MeCO.Bu

HCS 1980, 643

†2489. 3-Hexanone
 [589-38-8] C₆H₁₂O
 EtCO.Pr

†2490. Isobutyl vinyl ether
 [109-53-5] C₆H₁₂O
 Me₂CHCH₂OCH=CH₂

†2491. 2-Methylpentanal
 [123-15-9] C₆H₁₂O
 Me₂CHC₂H₄CO.H

†2492. 3-Methylpentanal
 [15877-57-3] C₆H₁₂O
 EtCHMeCH₂CO.H

†2493. 2-Methyl-3-pentanone
 [565-69-5] C₆H₁₂O
 Me₂CHCO.Et

†2494. 3-Methyl-2-pentanone
 [565-61-7] C₆H₁₂O
 MeCOCHMeEt

†2495. 4-Methyl-2-pentanone
[108-10-1]



FPA H97, 1981; *HCS 1980*, 576

1. Smith, G. H., *Chem. & Ind.*, 1972, 291
2. Bretherick, L., *Chem. & Ind.*, 1972, 363

4-Methyl-2-pentanone had not been considered prone to autoxidation, but an explosion during prolonged and repeated aerobic hot evaporation of the solvent [1] was attributed to formation and explosion of a peroxide [2].

Other reactants

Yoshida, 1980, 363

MRH values calculated for 13 combinations with oxidants are given.

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids, etc.

See other PEROXIDISABLE COMPOUNDS

†2496. Butyl acetate
[123-86-4]



FPA H116, 1982; *HCS 1980*, 233

Potassium *tert*-butoxide


See Potassium *tert*-butoxide: Acids, etc.


†2497. 2-Butyl acetate
[105-46-4]



HCS 1980, 234

†2498. 2,6-Dimethyl-1,4-dioxane
[10138-17-7]





†2499. Ethyl isobutyrate [97-62-1]	EtOCO.CHMe ₂	C ₆ H ₁₂ O ₂
†2500. 2-Ethyl-2-methyl-1,3-dioxolane [126-39-6]		C ₆ H ₁₂ O ₂
†2501. 4-Hydroxy-4-methyl-2-pentanone [123-42-2]	MeCO.CH ₂ CH(OH)Me ₂	C ₆ H ₁₂ O ₂
<i>HCS 1980, 355</i>		
†2502. Isobutyl acetate [110-19-0]	Me ₂ CHCH ₂ OCO.Me	C ₆ H ₁₂ O ₂
<i>HCS 1980, 567</i>		
†2503. Isopentyl formate [110-45-2]	Me ₂ CHC ₂ H ₄ OCO.H	C ₆ H ₁₂ O ₂
<i>HCS 1980, 564</i>		
†2504. Isopropyl propionate [637-78-5]	Me ₂ CHOCO.Et	C ₆ H ₁₂ O ₂
†2505. Methyl isovalerate [556-24-1]	MeOCO.CH ₂ CHMe ₂	C ₆ H ₁₂ O ₂
†2506. Methyl pivalate [598-98-1]	MeOCO.CMe ₃	C ₆ H ₁₂ O ₂

†2507. Methyl valerate
[624-24-8]



2508. Tetramethyl-1,2-dioxetane
[35856-82-7]



Kopecky, K. R. *et al.*, *Can. J. Chem.*, 1975, **53**, 1107

Several explosions occurred in vacuum-sealed samples kept at ambient temperature. Storage under air at -20°C , or as solutions up to 2 Mol appear safe.

See 3,3-Dimethyl-1,2-dioxetane

See DIOXETANES See *other* CYCLIC PEROXIDES

2509. *tert*-Butyl peroxyacetate
[107-71-1]





1. Martin, J. H., *Ind. Eng. Chem.*, 1960, **52**(4), 49A
2. Castrantas, 1965, 16
3. *CHETAH*, 1990, 188

The peroxyester explodes with great violence when rapidly heated to a critical temperature. Previous standard explosivity tests had not shown this behaviour. The presence of benzene (or preferably a less toxic solvent) as diluent prevents the explosive decomposition, but if the solvent evaporates, the residue is dangerous [1]. The pure ester is also shock-sensitive and detonable, but the commercial 75% solutions are not [2]. However, a 75% benzene solution has been exploded with a detonator, though not by mechanical shock [3].

See *other* PEROXYESTERS

2510. Cyclobutylmethyl methanesulfonate
[22524-45-4]




Explodes during distillation at $81-89^{\circ}\text{C}/0.8$ mbar.

See *entry* SULFONIC ACID ESTERS See *other* SULFUR ESTERS

2511. 2-Ethoxyethyl acetate
[111-15-9]

C₆H₁₂O₃



HCS 1980, 461

1. Anon., private comm., 1975
2. Author's comments

Mild explosions have occurred at the end of technical-scale distillations of 'ethyl glycol acetate' in a copper batch still about 20 min after the kettle heater was shut off. It is possible that air was drawn into the distillation column as it cooled, creating a flammable mixture, but no mechanism for ignition could be established. Oxidation studies did not indicate the likely formation of high peroxide concentrations in the liquid phase of this ether-ester [1]. It seems remotely possible that small amounts of dioxane (formed by dehydration of traces of ethylene glycol) or of acetaldehyde (arising from traces of ethanol or from cracking of the ester, catalysed perhaps by copper), and both having very low autoignition temperatures, may have been involved in the incidents [2].

†**2512. Isobutyl peroxyacetate**
[55153-36-1]


C₆H₁₂O₃



See other PEROXYESTERS

2513. *trans*-2-Hexene ozonide (3-Methyl-5-propyl-1,2,4-trioxolane)
[]

C₆H₁₂O₃



Loan, L. D. *et al.*, *J. Amer. Chem. Soc.*, 1965, **87**, 41

Attempts to get C and H analyses by combustion of the ozonides of 2-butene, 2-pentene and 2-hexene caused violent explosions, though oxygen analyses were uneventful.

See other OZONIDES

2514. Peroxyhexanoic acid
[5106-46-7]

C₆H₁₂O₃




Swern, D., *Chem. Rev.*, 1949, **45**, 10

Fairly stable at ambient temperature, it explodes and ignites on rapid heating.
See other PEROXYACIDS

†2515. 2,4,6-Trimethyltrioxane
[123-63-7]


$C_6H_{12}O_3$



Nitric acid
See Nitric acid: 2,4,6-Trimethyltrioxane

2516. Tetramethoxyethylene
[1069-12-1]


$C_6H_{12}O_4$



Preparative hazard
See 1,2,3,4-Tetrachloro-7,7-dimethoxy-5-phenylbicyclo[2.2.1]hepta-2,5-diene


2517. 3,3,6,6-Tetramethyl-1,2,4,5-tetraoxane
[1073-91-2]

$C_6H_{12}O_4$



Baeyer, A. *et al.*, *Ber.*, 1900, **33**, 858
Action of hydrogen peroxide or permonosulfuric acid on acetone produces this dimeric acetone peroxide, which explodes violently on impact, friction or rapid heating.
See other CYCLIC PEROXIDES

2518. Glucose
[50-99-7]



Energy of decomposition (in range 224–330°C) measured as 0.406 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkali

See SUGARS



Potassium nitrate, Sodium peroxide

See Sodium peroxide: Glucose, Potassium nitrate

See other REDUCANTS

2519. Bis(dimethylthallium)acetylide

[]





Nast, R. *et al.*, *J. Organomet. Chem.*, 1966, **6**, 461

Extremely explosive, heat- and friction-sensitive.

See related ALKYLMETALS, METAL ACETYLIDES


2520. Hexyl perchlorate
[52936-24-0]



See entry ALKYL PERCHLORATES (reference 6)

†2521. Cyclohexylamine
[108-91-8]

$C_6H_{13}N$




HCS 1980, 347

Nitric acid

See Nitric acid: Cyclohexylamine See other ORGANIC BASES

†2522. 1-Methylpiperidine
[626-67-5]


$C_6H_{13}N$



See other ORGANIC BASES

†2523. 2-Methylpiperidine
[109-05-7]

$C_6H_{13}N$



See other ORGANIC BASES


†2524. 3-Methylpiperidine
[626-56-2]

$C_6H_{13}N$

See other ORGANIC BASES

†2525. 4-Methylpiperidine
[626-58-4]


$C_6H_{13}N$



See other ORGANIC BASES

†2526. Perhydroazepine (Hexamethyleneimine)
[111-49-9]


C₆H₁₃N



See other ORGANIC BASES

2527. *N*-Methylpiperidine *N*-oxide
[17206-00-7]

C₆H₁₃NO




Cope, A. C. *et al.*, *J. Amer. Chem. Soc.*, 1960, **82**, 4656

This amine oxide is thermally surprisingly stable, decomposing explosively at 215°C.

See other *N*-OXIDES

2528. *N*-Butyl-*N*-2-azidoethylnitramine
[84928-98-3]

C₆H₁₃N₅O₂



See entry ENERGETIC COMPOUNDS

See other *N*-NITRO COMPOUNDS, ORGANIC AZIDES

†2529. 2,2-Dimethylbutane
[75-83-2]

C₆H₁₄



†2530. 2,3-Dimethylbutane
[79-29-8]

C₆H₁₄



†2531. Hexane
[110-54-3]

C₆H₁₄



FPA H83, 1979; HCS 1980, 533 (covers isomers); RSC Lab. Hazards Data Sheet No. 63, 1987

Dinitrogen tetroxide

See Dinitrogen tetroxide: Hydrocarbons

Other reactants

Yoshida, 1980, 326

MRH values calculated for 13 combinations with oxidants are given.

†2532. Isohexane (2-Methylpentane)

[107-83-5]



C_6H_{14}

†2533. 3-Methylpentane

[96-14-0]




C_6H_{14}

2534. Chlorodipropylborane

[22086-53-9]

$\text{C}_6\text{H}_{14}\text{BCl}$



Leleu, *Cahiers*, 1977, (88), 365

It ignites in air

See other ALKYLHALOBORANES

2535. Diisopropylberyllium

[15721-33-2]

$\text{C}_6\text{H}_{14}\text{Be}$



Water

Coates, G. E. *et al.*, *J. Chem. Soc.*, 1954, 22


Uncontrolled reaction with water is explosive.

See other ALKYLMETALS

2536. 1,4-Bis(isothiuronio)-2-butene dichloride

[67405-47-4]

$\text{C}_6\text{H}_{14}\text{Cl}_2\text{N}_4\text{S}_2$



Harrp, D. N. *et al.*, *Org. Prep. Proceed. Int.*, 1978, **10**, 13

Crystallisation from 95% ethanol is strongly exothermic.

2537. Dipropylmercury
[628-85-3] **C₆H₁₄Hg**




Iodine

Whitmore, 1921, 100

Interaction is violent.

See other ALKYL METALS, MERCURY COMPOUNDS

2538. Diisopropyl hyponitrite (Bis(2-propyloxy)diazene)
[82522-47-2]; E-[86886-16-0] **C₆H₁₄N₂O₂**



Ogle, C. A. *et al.*, *J. Org. Chem.*, 1983, **48**, 3729

It detonated violently when deliberately struck.

See other DIALKYL HYPONITRITES

2539. Dipropyl hyponitrite (Bis(propyloxy)diazene)
[] **C₆H₁₄N₂O₂**



See entry DIALKYL HYPONITRITES


†**2540. Butyl ethyl ether**
[628-81-9] **C₆H₁₄O**



HCS 1980, 473

See other PEROXIDISABLE COMPOUNDS

†**2541. *tert*-Butyl ethyl ether**
[637-92-3] **C₆H₁₄O**



†**2542. Diisopropyl ether**
[108-20-3] **C₆H₁₄O**



FPA H101, 1981; *HCS 1980*, 579

1. Anon., *Chem. Eng. News*, 1942, **20**, 1458

2. *MCA Case Histories Nos. 603, 1607*

3. Douglass, I. B., *J. Chem. Educ.*, 1963, **40**, 469
4. Rieche, A. *et al.*, *Ber.*, 1942, **75**, 1016
5. Hamstead, A. C. *et al.*, *J. Chem. Eng. Data.*, 1960, **5**, 583; *Ind. Eng. Chem.*, 1961, **53**(2), 63A
6. Walton, G. C., *CHAS Notes*, 1988, **6**, 3

There is a long history of violent explosions involving peroxidised diisopropyl ether, with initiation by disturbing a drum [1], unscrewing a bottle cap [2], or accidental impact [3]. This ether, with two susceptible hydrogen atoms adjacent to the oxygen link, is extremely readily peroxidised after only a few hours' exposure to air, initially to a dihydroperoxide which disproportionates to dimeric and trimeric acetone peroxides. These separate from solution and are highly explosive [4]. Presence of a crystalline solid in a sample of the ether should be a cause for great concern, no attempt being made to open the container. Professional assistance should be sought for the safe disposal of substantially peroxidised samples. It has been reported that the ether may be inhibited completely against peroxide formation for a considerable time by addition of *N*-benzyl-4-aminophenol at 16 ppm, or by addition of diethylenetriamine, triethylenetetramine or tetraethylenepentamine at 50 ppm [5]. Further information on the extreme hazards of peroxidised diisopropyl ether is presented, with directions for safe disposal in a remote location by controlled explosion. The dry crystalline peroxide is photo-sensitive and will detonate on exposure to sunlight. Four additional literature references are included [6]. See 3,3,6,6-Tetramethyl-1,2,4,5-tetraoxane, and 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxonane

See also ETHERS

Propionyl chloride

See Propionyl chloride: Diisopropyl ether

See other PEROXIDATION INCIDENTS, PEROXIDISABLE COMPOUNDS

†2543. Dipropyl ether

[111-43-3]

PrOPr

C₆H₁₄O

See other PEROXIDISABLE COMPOUNDS

†2544. 2-Methyl-2-pentanol

[590-36-3]

Me₂C(OH)Pr

C₆H₁₄O

†2545. 1,1-Diethoxyethane (Diethyl acetal)

[105-57-7]

(EtO)₂CHMe

C₆H₁₄O₂

Mee, A. J., *School Sci. Rev.*, 1940, **22**(86), 95

A peroxidised sample exploded violently during distillation.
See other PEROXIDISABLE COMPOUNDS

†2546. 1,2-Diethoxyethane
[629-14-1]



HCS 1980, 400

See other PEROXIDISABLE COMPOUNDS

2547. Dipropyl peroxide
[29914-92-9]





Swern, 1972, Vol. 3, 21

Unexpected explosions have occurred with dipropyl peroxides.

See other DIALKYL PEROXIDES

2548. 2,6-Dimethyl-1,3-dioxo-2,6-diphosphacyclooctane
[79251-56-2; 79251-57-3] *cis*-, *trans*-, resp.




Piccini-Leopardi, C. *et al.*, *J. Chem. Soc., Perkin Trans 2*, 1986, 85–92

The isomers could not be separated by distillation at 65–68°C/0.07 mbar, and bath temperatures above 120°C caused explosions.

See related ALKYLPHOSPHINES, PHOSPHORUS ESTERS

2549. Bis-(2-methoxyethyl) ether
[111-96-6]



'Diglyme Properties and Uses', Brochure, London, ICI, 1967

Details of properties, applications and safe handling are given. Light, heat and oxygen (air) promote formation of potentially explosive peroxides. These may be removed by stirring with a suspension of iron oxide in aqueous alkali.

Metal hydrides

See Aluminium hydride: Carbon dioxide, etc.


Lithium tetrahydroaluminate: Bis(2-methoxyethyl) ether

See other PEROXIDISABLE COMPOUNDS

2550. 1-Pentyl methanesulfonate

[7958-20-3]

C₆H₁₄O₃S



It decomposes vigorously at 185°C.

See entry SULFONIC ACID ESTERS *See other* SULFUR ESTERS

2551. Dipropylzinc

[628-91-1]

C₆H₁₄Zn

PrZnPr

Ellern, 1968, 24

It ignites in air if exposed on an extended surface.

See other DIALKYLZINCS

2552. Dipropylaluminium hydride

[2036-15-9]

C₆H₁₅Al

Pr₂AlH

HCS 1980, 444

See entry ALKYLALUMINIUM ALKOXIDES AND HYDRIDES

†2553. Triethylaluminium

[97-93-8]

C₆H₁₅Al

Et₃Al

HCS 1980, 929

It ignites in air.

See TRIALKYLALUMINIUMS

Alcohols

MRH Methanol, 1.45/46

Mirviss, S. B. *et al.*, *Ind. Eng. Chem.*, 1961, **53**(1), 54A

Interaction of methanol, ethanol and 2-propanol with the undiluted trialkylaluminium is explosive, while *tert*-butanol reacts vigorously.

See ALKYLALUMINIUM DERIVATIVES

Carbon tetrachloride

MRH 1.51/85

Reineckel, H., *Angew. Chem. (Intern. Ed.)*, 1964, **3**, 65

A mixture prepared at 0°C with a 3:1 molar excess of halocarbon exploded violently soon after removal of the ice bath. Formation of a 1:1 chlorine-bridged adduct was assumed.

See ALKYLALUMINIUM DERIVATIVES: halocarbons

Dimethylformamide

MRH 1.38/66

'DMF Brochure', Billingham, ICI, 1965

A mixture of the amide solvent and triethylaluminium explodes when heated.

Other reactants

Yoshida, 1980, 248

MRH values calculated for 14 combinations, largely with oxidants, are given.

Tris(pentafluorophenyl)boron

Pohlmann, J. L. W. *et al.*, *Z. Naturforsch.*, 1965, **20b** (1), 5

A mixture detonated on warming to 70°C.


See METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

See *other* ALKYLMETALS, TRIALKYLALUMINIUMS

2554. Diethylethoxyaluminium

[1586-92-1]

C₆H₁₅AlO



Dangerous Loads, 1972


The pure material, or solutions of above 20% concentration, ignite in air.

See *other* ALKYLALUMINIUM ALKOXIDES AND HYDRIDES

2555. Triethoxydialuminium tribromide

[65232-69-1]

C₆H₁₅Al₂Br₃O₃




Air, or Ethanol, or Water

Anon., *Chem. Eng. Progr.*, 1966, **62**(9), 128

It ignites in air and explodes with ethanol and water.

See related ALKYLALUMINIUM ALKOXIDES AND HYDRIDES, METAL HALIDES

2556. Triethyldialuminium trichloride
[12075-68-2]



HCS 1980, 466


Carbon tetrachloride

Reineckel, H., *Angew. Chem. (Intern. Ed.)*, 1964, **3**, 65

A mixture exploded when warmed to room temperature.

See other ALKYLALUMINIUM HALIDES See related METAL HALIDES

2557. Triethylarsine
[617-75-4]




Sidgwick, 1950, 762

It inflames in air.

See other ALKYLNON-METALS

2558. Triethylphosphinegold nitrate
[14243-51-7]





Coates, G. E. *et al.*, *Aust. J. Chem.*, 1966, **19**, 541

After thorough desiccation, the crystalline solid exploded spontaneously.

See other GOLD COMPOUNDS See related ORGANOMETALLIC NITRATES

2559. Triethylborane
[97-94-9]



HCS 1980, 931

Meerwein, H., *J. Prakt. Chem.*, 1937, **147**, 240
It ignites on exposure to air.

Halogens, or Oxygen

Coates, 1960, 84

It ignites in contact with chlorine or bromine, and explodes in oxygen.


Triethylaluminium

Hurd, D. T., *J. Amer. Chem. Soc.*, 1948, **70**, 2053


Mixtures with triethylaluminium have been used as hypergolic igniters in rocket propulsion systems.

See other ALKYLBORANES

†**2560. Triethyl borate**
[150-46-9]



2561. Triethylbismuth
[617-77-6]




Coates, 1967, Vol. 1, 537

It ignites in air, and explodes at about 150°C, before distillation begins.

See other TRIALKYLBISMUTHS

2562. Triethylsilyl perchlorate
[18244-91-2]



1. Wannagat, U. *et al.*, *Z. Anorg. Chem.*, 1950, **302**, 185–198


2. Barton, T. J. *et al.*, *J. Org. Chem.*, 1978, **43**, 3651

Several trialkyl- or triaryl-silyl perchlorates explode on heating [1]. A syringe used to inject a sample of the title perchlorate ester into a reaction flask exploded soon after addition was complete [2].

See other ORGANOSILYL PERCHLORATES

2563. Triethylgallium
[1115-99-7]

C₆H₁₅Ga



Air, or Water


Dennis, L. M. *et al.*, *J. Amer. Chem. Soc.*, 1932, **54**, 182

The compound ignites in air. Breaking a bulb containing 0.2 g under cold water shattered the container. The monoetherate behaved similarly under 6 M nitric acid

See other ALKYL METALS

2564. Triethylindium
[923-34-2]

C₆H₁₅In



Leleu, *Cahiers*, 1977, (88), 367

It ignites in air.

See other ALKYL METALS

†**2565. Butylethylamine**
[13360-63-9]

C₆H₁₅N

BuNH₂

See other ORGANIC BASES

†**2566. Diisopropylamine**
[108-18-9]

C₆H₁₅N

(Me₂CH)₂NH

See other ORGANIC BASES

†**2567. 1,3-Dimethylbutylamine**
[108-09-8]

C₆H₁₅N

Me₂CHCH₂CHMeNH₂

See other ORGANIC BASES

†2568. Dipropylamine
[142-84-7]



See other ORGANIC BASES

†2569. Triethylamine
[121-44-8]



FPA H92, 1980; HCS 1980, 930; RSC Lab. Hazard Data Sheet No. 27, 1984



2-Bromo-2,5,5-trimethylcyclopentanone, Potassium hydroxide
See 2-Bromo-2,5,5-trimethylcyclopentanone: Potassium hydroxide, etc.

Dinitrogen tetraoxide
See Dinitrogen tetraoxide: Triethylamine

Maleic anhydride
See Maleic anhydride: Bases, etc.


Other reactants
Yoshida, 1980, 247
MRH Values calculated for 13 combinations with oxidants are given.
See other ORGANIC BASES

2570. Triethylamine hydrogen peroxidate
[]



See entry CRYSTALLINE HYDROGEN PEROXIDATES

2571. 2-Diethylammonioethyl nitrate nitrate
[]




1. Barbière, J., *Bull. Soc. Chim. Fr.*, 1944, **11**, 470-480
2. Fakstorp, J. *et al.*, *Acta Chem. Scand.*, 1951, **5**, 968-969

Repetition of the original preparation [1] involving interaction of diethylaminoethanol with fuming nitric acid, followed by vacuum distillation of excess acid, invariably caused explosions during this operation. A modified procedure without evaporation is described [2].

See related ALKYL NITRATES, OXOSALTS OF NITROGENOUS BASES

2572. Diethyl ethanephosphonite
[2651-85-6]

C₆H₁₅O₂P




Arbuzov, B. A. *et al.*, *Chem. Abs.*, 1953, **47**, 3226a

Absorbed on to filter paper it ignites in air.

See other PHOSPHORUS ESTERS *See related* ALKYLPHOSPHINES

2573. Triethylphosphine
[554-70-1]

C₆H₁₅P



Oxygen

Engler, C. *et al.*, *Ber.*, 1901, **34**, 365

Action of oxygen at low temperature on the phosphine produces an explosive product.


Palladium(II) perchlorate

See Perchloratotris(triethylphosphine)palladium(II) perchlorate

See other ALKYLPHOSPHINES

2574. Triethylantimony
[617-85-6]


C₆H₁₅Sb



von Schwartz, 1918, 322

It ignites in air.
See other ALKYL METALS

2575. Sodium dihydrobis(2-methoxyethoxy)aluminat
[22722-98-1]




Preparative hazard

1. Casensky, B. *et al.*, *Inorg. Synth.*, 1978, **18**, 150
2. Strauf, O. *et al.*, *J. Organomet. Chem. Library* 15, 1985

The preparation is effected by heating sodium, aluminium powder and 2-methoxyethanol in toluene at 150°C under 70 bar pressure of hydrogen in an autoclave. The reaction is exothermic, and hydrogen is first evolved by the dissolving metals, initially increasing the pressure, before absorption of hydrogen into the complex hydride occurs to reduce the pressure. Care in controlling temperature and the volume of reagents charged to the autoclave is necessary to avoid hazards. The solid complex hydride is not pyrophoric with air or water, but concentrated solutions (above 70%) may ignite when exposed to air on dry textiles [1]. It is, however a novel and versatile reducing agent which can with advantage (including that of increased safety in use) replace other complex hydrides [2].


See other COMPLEX HYDRIDES, GAS EVOLUTION INCIDENTS

2576. Bis(methyl 1-methylhydrazinocarbodithioate *N*²,*S*'(perchlorato-*O*, *O*') copper(1+) perchlorate
[67870-97-7]



See Bis-*O*, *N*[(*N*'-pent-2-en-2-oxo-4-ylidene)-*N,S*-dimethyldithiocarbamate] copper(II) perchlorate

2577. Ammonium hexacyanoferrate(II)
[14481-29-9]



Hydrochloric acid

Ephraim, 1939, 303

Contact with acids liberates the solid complex ferrocyanic acid which is endothermic (ΔH_f° (aq) +534.7 kJ/mol, 2.48 kJ/g), and forms complexes with diethyl ether etc.


Metal nitrates

See Cobalt(II) nitrate: Ammonium hexacyanoferrate

Copper(II) nitrate: Ammonium hexacyanoferrate

See related ENDOTHERMIC COMPOUNDS, METAL CYANIDES (AND CYANO COMPLEXES)

2578. Lithium triethylsilylamide
[13270-95-6]



Oxidants

It is hypergolic with fluorine or fuming nitric acid, and explodes with ozone.


See SILYLHYDRAZINES See other N-METAL DERIVATIVES

†**2579. 1,2-Bis(dimethylamino)ethane**
[110-18-9]



See other ORGANIC BASES

2580. Triethylammonium nitrate
[27096-31-7]




Dinitrogen tetroxide

See Dinitrogen tetroxide: Triethylammonium nitrate

See other OXOSALTS OF NITROGENOUS BASES

2581. Diazido(*N,N,N,N*-tetramethylethanediamine)palladium
[158668-93-0]

C₆H₁₆N₈Pd



Kim, Y. *et al.*, *Bull. Korean Chem. Soc.*, 1994, **15**(8), 690

Exploded during melting point determination.

See AZIDES

†**2582. Diethoxydimethylsilane**
[78-62-6]


C₆H₁₆O₂Si

(EtO)₂SiMe₂

See related ALKYL SILANES

2583. Triethyltin hydroperoxide
[]

C₆H₁₆O₂Sn



Hydrogen peroxide

Aleksandrov, Y. A. *et al.*, *Zh. Obsch. Khim.*, 1965, **35**, 115

While the hydroperoxide is stable at ambient temperature, the addition compound which it readily forms with excess hydrogen peroxide is not, decomposing violently.

See other ORGANOMETALLIC PEROXIDES

2584. Triethoxysilane
[998-30-1]

C₆H₁₆O₃Si

(EtO)₃SiH

1. Xin, S. *et al.*, *Can. J. Chem.*, 1990, **68**, 471

2. Buchwald, S. L., *Chem. Eng. News*, 1993, **71**(13), 2


This compound can disproportionate, generating silane (SiH₄) which is liable to be pyrophoric. Presumably other alkoxy silanes can do likewise [1]. Its use as a reducing agent for the preparation of alcohols from esters is considered safer in air than under

an inert atmosphere, which latter permits accumulation of silane to hazardous levels and later fire or explosion. [2]

See related SILANES

2585a. Ethylenebis(dimethylphosphine)
[23936-60-9]

C₆H₁₆P₂



1. Burt, R. J., *J. Organomet. Chem.*, 1979, **182**, 205
 2. Butter, S. A. *et al.*, *Inorg. Synth.*, 1974, **15**, 190
- Though air sensitive [1], it is not pyrophoric as reported previously [2].
See other ALKYLPHOSPHINES

2585b. Triethylsilane
[617-86-7]

C₆H₁₆Si

Et₃SiH


Boron trichloride

See Boron trichloride: Triethylsilane

See other ALKYL SILANES

2586. 1-Ethyl-1,1,3,3-tetramethyltetrazenium tetrafluoroborate
[65651-91-4]

C₆H₁₇BF₄N₄



Stödt, E. *et al.*, *Angew. Chem. (Intern., Ed.)*, 1978, **17**, 203

An explosive salt.

See other PERCHLORATE SALTS OF NITROGENOUS BASES, TETRAZENES

2587. 3,3'-Iminobispropylamine (Norspermidine)
[56-18-8]

C₆H₁₇N₃


HN(CH₂CH₂CH₂NH₂)₂

Editor's comments, 1993

Various secondary sources of safety data are now listing this as an explosive. I can find no primary source for this classification, which seems very improbable. Simple minded use of many computational hazard prediction procedures would show thermodynamically that this compound, like most lower amines, could hypothetically convert to alkane, ammonia and nitrogen with sufficient energy (about 3 kJ/g) to count as an explosion hazard. This reaction is not known to happen. (Simple minded thermodynamicists would rate this book, or computer, and its reader as a severe hazard in an air environment.) Like other bases, iminobispropylamine certainly sensitises many nitro-explosives to detonation. It is used experimentally to study the effect, which may have found technical exploitation and, garbled, could have led to description of the amine as itself an explosive.

See C-NITRO COMPOUNDS, COMPUTATION OF REACTIVE CHEMICAL HAZARDS

2588. Triethyldiborane
[62133-36-2]




Leleu, *Cahiers*, 1977, (88), 365

It ignites in air.

See other ALKYLBORANES

2589. 1,2-Bis(ethylammonio)ethane perchlorate
[53213-78-8]




Sunderlin, K. G. R., *Chem. Eng. News*, 1974, **52**(31), 3

It exploded mildly under an impact of 11.5 kgm.

See other PERCHLORATE SALTS OF NITROGENOUS BASES


2590. Lithium hexamethylchromate(3-)
[14931-97-6] (tris-dioxane complex)



Kurras, E. *et al.*, *J. Organomet. Chem.*, 1965, **4**, 114–118

Isolated as a dioxane complex, it ignites in air.
See related ALKYL METALS


2591. Bis(trimethylsilyl) chromate
[1746-09-4]



1. Anon., *ABCM Quart. Safety Summ.*, 1960, **31**, 16
2. Schmidt, M. *et al.*, *Angew. Chem.*, 1958, **70**, 704

Small quantities can be distilled at about 75°C/1.3 mbar, but larger amounts are liable to explode violently owing to local overheating [1]. An attempt to prepare an analogous poly(dimethylsilyl) chromate by heating a polydimethylsiloxane with chromium trioxide at 140°C exploded violently after 20min at this temperature [2].
See related ALKYL SILANES, METAL OXIDES



2592. Hexamethylerbium–hexamethylethylenediaminelithium complex
[66862-11-1]



Schumann, H. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1981, **20**, 120–121

It ignites in air, as do the analogous derivatives of lutetium and other lanthanoids
See related ALKYL METALS

2593. Bis(trimethylsilyl)mercury
[4656-04-6]




Eisch, J. J. *et al.*, *J. Organomet. Chem.*, 1979, **171**, 149

Momentary exposure of traces of the product on a stopper to air usually causes ignition.

See related ALKYL SILANES *See other* MERCURY COMPOUNDS

2594. Lithium bis(trimethylsilyl)amide
[4039-32-1]




Amonoo-Neizer, E. H. *et al.*, *Inorg. Synth.*, 1966, **8**, 21

It is unstable in air and ignites when compressed.

See other N-METAL DERIVATIVES


2595. Dilithium 1,1-bis(trimethylsilyl)hydrazide
[15114-92-8]



It ignites in air.

See entry SILYLHYDRAZINES *See other* N-METAL DERIVATIVES, METAL HYDRAZIDES

2596. Molybdenum hexamethoxide
[135840-44-7]




Preparative hazard

See Tetramethoxysilane: Metal hexafluorides

See other METAL ALKOXIDES

2597. Bis(dimethylamino)dimethylstannane
[993-74-8]

C₆H₁₈N₂Sn



Chloroform


Randall, E. W. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1965, **1**, 105

Gentle heating of a 1:1 chloroform solution of the stannane led to a mild explosion. Similar explosions had been experienced at the conclusion of fractional distillation of other bis(dialkylamino)stannanes where the still pot temperature had risen to about 200°C.

See other N-METAL DERIVATIVES See related ALKYL METALS

2598. Tris(dimethylamino)antimony
[7289-92-1]

C₆H₁₈N₃Sb



Ethyl diazoacetate

See Ethyl diazoacetate: Tris(dimethylamino)antimony

See other N-METAL DERIVATIVES

2599. N,N'-Bis(2-aminoethyl)1,2-diaminoethane (Triethylenetetramine)
[112-24-3]

C₆H₁₈N₄



Cellulose nitrate

See CELLULOSE NITRATE: amines


Other reactants

Yoshida, 1980, 251

MRH values calculated for 13 combinations with oxidants are given.

See other ORGANIC BASES

2600. 2,4,6,8,9,10-Hexamethylhexaaza-1,3,5,7-tetraphosphaadamantane
 [10369-17-2] **C₆H₁₈N₆P₄**



Oxidants

Verdier, F., *Rech. Aerosp.*, 1970, (137), 181–189

Among other solid P–N compounds examined, the title compound ignited immediately on contact with nitric acid, hydrogen peroxide or dinitrogen trioxide.


See ROCKET PROPELLANTS

†**2601. Bis(trimethylsilyl) oxide**
 [107-46-0] **C₆H₁₈OSi₂**



See related ALKYL SILANES

2602. Bis(trimethylsilyl) peroxomonosulfate
 [23115-33-5] **C₆H₁₈O₅SSi₂**




Blaschette, A. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1969, **8**, 450

It is stable at –30°C, but on warming to ambient temperature it decomposes violently evolving sulfur trioxide.

See other GAS EVOLUTION INCIDENTS, PEROXYESTERS, SULFUR ESTERS

2603. Rhenium hexamethoxide
 [] **C₆H₁₈O₆Re**




Preparative hazard

See Tetramethoxysilane: Metal hexafluorides

See other METAL ALKOXIDES

2604. Tungsten hexamethoxide
[35869-33-1]




Preparative hazard

See Tetramethoxysilane: Metal hexafluorides

See other METAL ALKOXIDES

2605. (Dimethylsilylmethyl)trimethyllead
[]




Schmidbaur, H., *Chem. Ber.*, 1964, **97**, 270

It decomposes slowly above 100°C, and with explosive violence in presence of oxygen.

See related ALKYL METALS, ALKYL SILANES

2606. Hexamethyldiplatinum
[4711-74-4]




Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1953, **75**, 2065

It explodes sharply in a shower of sparks on heating.

See other ALKYL METALS, PLATINUM COMPOUNDS

2607. Hexamethylrhenium
[56090-02-9]

C₆H₁₈Re




1. Mertis, K. *et al.*, *J. Organomet. Chem.*, 1975, **97**, C65
2. Mertis, K. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1976, 1489
3. Edwards, P. G. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 2474

It should be handled with extreme caution, even in absence of air, because dangerous explosions have occurred on warming frozen samples, or during transfer operations [1]. It is unstable above -20°C , and on one occasion admission of nitrogen to the solid under vacuum caused a violent explosion [2]. It should be prepared only as required and in small amounts, and in use, oxygen and moisture must be rigorously excluded [3].

See other ALKYLMETALS

2608. Hexamethyldisilane
[1450-14-2]

C₆H₁₈Si₂




Pyridine *N*-oxide, Tetrabutylammonium fluoride

See Pyridine *N*-oxide: Hexamethyldisilane, Tetrabutylammonium fluoride

See other ALKYL SILANES

2609. Hexamethyltungsten
[36133-73-0]

C₆H₁₈W




Galyer, L. *et al.*, *J. Organomet. Chem.*, 1975, **85**, C37

Several unexplained and violent explosions during preparation and handling indicate that the compound must be handled with great care and as potentially explosive, particularly during vacuum sublimation.



See other ALKYL METALS

†2610. Hexamethyldisilazane
[999-97-3]



See related ALKYL SILANES

2611. Bis(trimethylsilyl) phosphonite
[30148-50-6]




Hata, T. *et al.*, *Chem. Lett.*, 1977, 1432

It ignites in air.

See other PHOSPHORUS ESTERS, PYROPHORIC MATERIALS

2612. *N,N*-Bis(trimethylsilyl)aminoborane
[73452-31-0]





Wisian-Nielsen, P., *J. Inorg. Nucl. Chem.*, 1979, **41**, 1546

It ignites in direct contact with air.

See related ALKYL SILANES, BORANES

2613. Bis(1,2-diaminopropane)-*cis*-dichlorochromium(III) perchlorate
[59598-02-6]



Perchloric acid

Anon., *Chem. Eng. News*, 1963, **41**(27), 47

A mixture of the complex and 20 volumes of 70% acid was being stirred at 22°C when it exploded violently. Traces of ether or ethanol may have been present. Extreme care must be exercised when concentrated perchloric acid is contacted with organic materials with agitation or without cooling.

See other AMMINOMETAL OXOSALTS

2614. (Benzenesulfinato-S)pentaamminecobalt(III) trichloro(perchlorato)-stannate(II)

[64825-48-5]

$C_6H_{20}Cl_4CoN_5O_6SSn$



Elder, R. C. *et al.*, *Inorg. Chem.*, 1978, **17**, 427

It explodes at temperatures around 150°C. Thermal decomposition appears to involve an internal redox reaction in the anion leading to formation of trichloro(chlorato)oxostannate(IV).


See other AMMINEMETAL OXOSALTS, REDOX COMPOUNDS

See related METAL PERCHLORATES, REDOX REACTIONS

2615. cis-Bis(trimethylsilylamino)tellurium tetrafluoride

[86045-52-5]

$C_6H_{20}F_4N_2Si_2Te$



Hartl, H. *et al.*, *Inorg. Chem.*, 1983, **22**, 2183–2184

The colourless solid explodes mildly at 100°C, but a 50 g sample on keeping for 6 months in a stoppered cylinder was converted to a yellow extremely explosive solid, probably a hydrolysis product.


See other AMMINEMETAL HALIDES

See related ALKYL SILANES, METAL HALIDES

2616. 1,2-Bis(trimethylsilyl)hydrazine

[692-56-8]

$C_6H_{20}N_2Si_2$




Oxidants

It is hypergolic with fluorine or fuming sulfuric acid.

See entry SILYLHYDRAZINES

2617. Hexamethyltrisiloxane
[1189-93-1]





See entry SILANES; olefins, platinum
See related ALKYL SILANES

2618. Isonicotinamidopentaammineruthenium(II) perchlorate
[31279-70-6]





Stanbury, D. M. *et al.*, *Inorg Chem.*, 1980, **19**, 519
It is explosive.
See other AMMINOMETAL OXOSALTS

2619. Tris(1,2-diaminoethane)chromium(III) perchlorate
[15246-55-6]



Olander, D. E., *Chem. Eng. News*, 1984, **62**(10), 2, 39
It is a treacherously sensitive, very brisant primary explosive (while its cobalt analogue is a well behaved secondary explosive).
See other AMMINOMETAL OXOSALTS

2620. Hexaureagallium(III) perchlorate
[31332-72-6]



Lloyd, D. J. *et al.*, *J. Chem. Soc.*, 1943, 76
It decomposes violently when heated above its m.p., 179°C.
See related AMMINOMETAL OXOSALTS


2621. Tris(1,2-diaminoethane)ruthenium(III) perchlorate
[67187-50-2]



Lavallee, U. *et al.*, *Inorg. Chem.*, 1978, **17**, 2218
An attempt to prepare the perchlorate salt led to an explosion.
See other AMMINOMETAL OXOSALTS

2622. Tris(1,2-diaminoethane)cobalt(III) nitrate
[6865-68-5]

$C_6H_{24}CoN_9O_9$



Tomlinson, W. R. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 375
No explosion on heating, but medium impact-sensitivity.
See other AMMINEMETAL NITRATES

2623. Hexaureachromium(III) nitrate
[22471-42-7]

$C_6H_{24}CrN_{15}O_{15}$



Tomlinson, W. R. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 375
Explodes at 265°C, medium impact-sensitivity.
See related AMMINEMETAL NITRATES

2624. 1,1'-Azo-[2-methyl-1,2-dicarbadeborane(14)]
[73469-45-1]

$C_6H_{26}B_{20}N_2$

Complex Structure

Preparative hazard


Totari, T. *et al.*, *J. Chem. Soc., Chem. Commun.*, 1979, 1051

The reaction mixture from permanganate oxidation of the aminocarbaborane in liquid ammonia must be diluted with toluene at $-60^\circ C$ to avoid detonation when the ammonia is subsequently evaporated.

See AZOCARBABORANES *See related* BORANES

2625. Potassium cyclohexanehexone 1,3,5-trioximate
[]

$C_6K_3N_3O_6$



Alone, or Acids


Benedikt, R., *Ber.*, 1878, **11**, 1375

The salt explodes on heating above $130^\circ C$ or if moistened with sulfuric or nitric acids. The lead salt also explodes violently on heating.

See related OXIMES

See other N-O COMPOUNDS

2626. Potassium benzenehexoxide
[3264-86-6]




Air, or Oxygen, or Water

1. Sorbe, 1968, 68, 69
2. Mellor, 1963, Vol. 2, Suppl. 2.2, 1567–1568; 1946, Vol. 5, 951
3. Weiss, E., *Angew. Chem. (Int.)*, 1993, **32**(11), 1518

The compound 'potassium carbonyl' is now thought to be the hexameric potassium salt of hexahydroxybenzene, and reacts with moist air to give a very explosive product [1]. It reacts violently with oxygen, and explodes on heating in air or in contact with water [2]. There are probably two compounds described, the original preparation from carbon monoxide and liquid potassium may well give the aromatic structure. X-ray crystallography shows the product from potassium and carbon monoxide in liquid ammonia to be dipotassium ethynediolate [3].

See Potassium: Non-metal oxides (reference 4)

2627. Lithium benzenehexoxide
[101672-16-6]




Water

1. Mellor, 1961, Vol. 2, Suppl. 2.1, 88
2. Weiss, E., *Angew. Chem. (Int.)*, 1993, **32**(11), 1518

Like the potassium compound, it explodes with water [1]. It is probable that this compound is actually dilithium ethynediolate [2].

2628. Hexacarbonylmolybdenum
[13939-06-5]

C_6MoO_6



Diethyl ether


Owen, B. B. *et al.*, *Inorg. Synth.*, 1950, **3**, 158

Solutions of hexacarbonylmolybdenum in ether have exploded after extended storage.

See other CARBONYLMETALS

2629. Tetracyanoethylene
[670-54-2]

C_6N_4




1-Methylsilacyclopenta-2,4-diene

See 1-Methylsilacyclopenta-2,4-diene: Dienophiles

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS

2630. 3(3-Cyano-1,2,4-oxadiazol-5-yl)-4-cyanofurazan 2-(or 5-) oxide
[56092-91-2]

$C_6N_6O_3$



Hydrazine derivatives

Lupton, E. C. *et al.*, *J. Chem. Eng. Data*, 1975, **20**, 136


The compound is highly endothermic (ΔH_f° (s) +565.5 kJ/mol, 2.77 kJ/g), and interaction with hydrazines is explosive.

See Dicyanofurazan: Nitrogenous bases

See other CYANO COMPOUNDS, ENDOTHERMIC COMPOUNDS, FURAZAN N-OXIDES

2631. Benzotri(furazan *N*-oxide) (Hexanitrosobenzene)
[3470-17-5]

$C_6N_6O_6$




1. Selig, W., *Z. Anal. Chem.*, 1978, **289**, 44–46
2. McGuire, R. R., *Chem. Abs.*, 1978, **89**, 131857
3. *CHETAH*, 1990, 182

It is explosive [1], and the limited thermal stability is reduced by traces of triazido-trinitrobenzene as impurity [2]. It is shock sensitive [3].

See other FURAZAN *N*-OXIDES

2632. Hexanitrobenzene
[13232-74-1]

$C_6N_6O_{12}$




Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1979, **44**, 1181

The compound (of zero oxygen balance) is a powerful explosive, as is pentanitrobenzene. As a nitrocarbon, it has similar properties to tetranitromethane and hexanitroethane.

See other POLYNITROARYL COMPOUNDS

2633. Tetraazido-1,4-benzoquinone
[22826-61-5]

$C_6N_{12}O_2$



1. Friess, K. *et al.*, *Ber.*, 1923, **56**, 1304


2. Gilligan, W. H. *et al.*, *Tetrahedron Lett.*, 1978, 1675–1676

3. Winkelmann, E., *Tetrahedron*, 1969, **25**, 2427

Extremely explosive, sensitive to heat, friction and impact. A single crystal heated on a spatula exploded and bent it badly [1]. It explodes at 120°C, and the impact-sensitivity was too high to be measured, extreme friction sensitivity causing explosions when samples were being arranged on sandpaper. Quantities above 100–200 mg are too dangerous to prepare and handle under ordinary laboratory conditions [2], the 24 g scale previously described [3] being dangerously excessive. *See other 2-AZIDOCARBONYL COMPOUNDS, HIGH-NITROGEN COMPOUNDS*

2634. Sodium benzenhexoxide
[101672-22-4]

C₆Na₆O₆



See Sodium: Non-metal oxides (reference 4)

2635. Hexacarbonylvandium
[14024-00-1]

C₆O₆V

(CO)₆V

1. Pruett, R. L. *et al.*, *Chem. & Ind.*, 1960, 119

2. Bailar, 1973, Vol. 3, 529

The pyrophoric compound previously regarded as 'dodecacarbonyl divandium' [1] is now known to be the monomeric hexacarbonylvandium [2].

See other CARBONYLMETALS

2636. Hexacarbonyltungsten
[14040-11-0]

C₆O₆W

(CO)₆W

Preparative hazard


Hurd, D. T. *et al.*, *Inorg. Chem.*, 1957, **5**, 136

It is dangerous to attempt the preparation from tungsten hexachloride, aluminium powder and carbon monoxide in an autoclave of greater than 0.3 l capacity.

See other CARBONYLMETALS

2637. 2-Chloro-1,3-dinitro-5-trifluoromethylbenzene
[393-75-9]

C₇H₂ClF₃N₂O₄




Nazarov, V. P. *et al.*, *Chem. Abs.*, 1980, **93**, 2387

Mathematical modelling was used to develop the recommendation to use a cascade of 3 reactors for the highly exothermic dinitration of 4-chloro-trifluoromethylbenzene.

See other NITRATION INCIDENTS, POLYNITROARYL COMPOUNDS

2638. 5-Nitro-2-picryltetrazole
[82177-75-1]

C₇H₂N₈O₈




Spear, R. J. *et al.*, *Chem. Abs.*, 1983, **98**, 91969

Powerfully explosive but of low impact-sensitivity.

See other POLYNITROARYL COMPOUNDS, TETRAZOLES

2639. 4-Iodobenzenediazonium-2-carboxylate
[]

C₇H₃IN₂O₂




Gompper, R. *et al.*, *Chem. Ber.*, 1968, **101**, 2348

It is a highly explosive solid.

See other DIAZONIUM CARBOXYLATES

2640. 2,4,6-Trinitrobenzoic acid
[129-66-8]

$C_7H_3N_3O_8$



Preparative hazard

See Sodium dichromate: Sulfuric acid, Trinitrotoluene

Heavy metals


Kranz, A. *et al.*, *Chem. Age* (London), 1925, **13**, 392

All heavy metal salts prepared were explosive on heating or impact.

See other POLYNITROARYL COMPOUNDS

2641. Hepta-1,3,5-triyne
[66486-68-8]

C_7H_4




Cook, C. L. *et al.*, *J. Chem. Soc.*, 1952, 2890

It explodes very readily (in absence of air) above 0°C, and the distillation residues also exploded on admission of air.

See other ALKYNES

2642. Silver 3,5-dinitroanthranilate
[58302-42-4]

$C_7H_4AgN_3O_6$




Gupta, S. P. *et al.*, *Def. Sci. J.*, 1975, **25**, 101–106

It explodes at 394°, and the copper salt at 371°C, each after a 10 s delay. They are not impact-sensitive.

See other POLY-NITROARYL COMPOUNDS, SILVER COMPOUNDS

2643. 2-, 3- or 4-Trifluoromethylphenylmagnesium bromide
[395-47-1, 402-26-6, 402-51-7, resp.]




1. Appleby, J. C., *Chem. & Ind.*, 1971, 120
2. 'Benzotrifluorides Catalog 6/15' West Chester (Pa.), Marshallton Res. Labs., 1971
3. Taylor, R. *et al.*, private comm., 1973
4. Moore, *et al.*, *Chem. Eng. News*, 1997, **75** (11), 6

During a 2 kg preparation under controlled conditions, the 3-isomer exploded violently [1]. The same had occurred previously in a 1.25 kg preparation 30 min after all magnesium had dissolved. A small preparation of the 2-isomer had also exploded. It was found that 3-preparations in ether decomposed violently at 75° and in benzene-THF at 90°C. Molar preparations or less have been done several times uneventfully when temperatures below 40°C were maintained, but caution is urged [2]. The 3- and 4-reagents prepared in ether exploded when the temperature was raised by adding benzene and distilling the ether out [3]. Twentyfive years after the establishment of a hazard, a sample of the 4-isomer, which had been all but isolated by evaporation of ether, exploded; the experimenters sought to blame hypothetical residues of highly active magnesium [4].

See other GRIGNARD REAGENTS, HALO-ARYLMETALS, FLUORINATED ORGANO-LITHIUM COMPOUNDS

2644. 4-Bromobenzoyl azide
[14917-59-0]




Curtiss, T. *et al.*, *J. Prakt. Chem.*, 1898, **58**, 201

It explodes violently above its m.p., 46°C.
See other ACYL AZIDES

2645. 3-Bromo-3(4-nitrophenyl)-3H-diazirine
[115127-49-6]

C₇H₄BrN₃O₂



Tetrabutylammonium fluoride


Terpinski, J. *et al.*, *Magn. Reson. Chem.*, 1987, **25**, 924

Direct introduction of the bromodiazirine into molten tetrabutylammonium fluoride (at 63°C) to effect halogen exchange leads to an explosion. Use of acetonitrile at 0°C as solvent gives a low yield of the required fluorodiazirine.

See other DIAZIRINES, N-HALOGEN COMPOUNDS, NITROARYL COMPOUNDS

2646. 4-Chlorotrifluoromethylbenzene
[98-56-6]

C₇H₄ClF₃



Sodium dimethylsulfinate

Meschino, J. A. *et al.*, *J. Org. Chem.*, 1971, **36**, 3637, footnote 9


Interaction with dimethylsulfinate anion (from the sulfoxide and sodium hydride) at -5°C is very exothermic, and addition of the chloro compound must be slow to avoid violent eruption.

See other HALOARYL COMPOUNDS

2647. 2-Chloro-5-trifluoromethylbenzenediazonium hydrogen sulfate

[29362-18-3] (ion)

C₇H₄ClF₃N₂O₄S




See entry DIAZONIUM SULFATES

2648. 4-Chlorophenyl isocyanate

[104-12-1]

C₇H₄ClNO



Preparative hazard


See 4-Chlorobenzoyl azide

See other ORGANIC ISOCYANATES

2649. 2-Chloro-5-nitrobenzaldehyde

[6361-21-3]

C₇H₄ClNO₃




Initial decomposition temperature by ARC was 156°C.

See entry NITROBENZALDEHYDES *See other* NITROARYL COMPOUNDS

2650. 2-Chloro-6-nitrobenzaldehyde
[6361-22-4]


$C_7H_4ClNO_3$



Initial decomposition temperature by ARC was 146°C.
See entry NITROBENZALDEHYDES
See other ALDEHYDES, NITROARYL COMPOUNDS

2651. 4-Chloro-3-nitrobenzaldehyde
[16588-34-4]


$C_7H_4ClNO_3$



Initial decomposition temperature by ARC was 116°C.
See entry NITROBENZALDEHYDES
See other ALDEHYDES, NITROARYL COMPOUNDS

2652. 5-Chloro-2-nitrobenzaldehyde
[6628-86-0]


$C_7H_4ClNO_3$



Initial decomposition temperature by DSC was 240°C.
See entry NITROBENZALDEHYDES
See other ALDEHYDES, NITROARYL COMPOUNDS

2653. 2-Nitrobenzoyl chloride
[610-14-0]

$C_7H_4ClNO_3$




1. Cook, N. C. *et al.*, *Chem. Eng. News*, 1945, **23**, 2394
2. Bonner, W. D. *et al.*, *J. Amer. Chem. Soc.*, 1946, **68**, 344; *Chem. & Ind.*, 1948, 89
3. *MCA Case History No. 1915*
4. Lockemann, G. *et al.*, *Ber.*, 1947, **80**, 488
5. *CHETAH*, 1990, 189

The hot material remaining after vacuum stripping of solvent up to 130°C decomposed with evolution of gas and then exploded violently 50 min after heating had ceased. Further attempts to distil the acid chloride even in small amounts at below 1.3 mbar caused exothermic decomposition at 110°C. It was, however, possible to flash-distil the chloride in special equipment [1]. Two later similar publications recommend use in solution of the unisolated material [2]. Smaller scale distillation of the chloride at 94–95°C/0.03 mbar had been uneventful, but a 1.2 mol scale preparation exploded during distillation at 128°C/1.2 mbar [3], even in presence of phosphorus pentachloride, previously recommended to reduce the danger of explosion during distillation [4]. Many previous explosions had been reported, and the need for adequate purity of intermediates was stressed. It is shock sensitive [5].

See other GAS EVOLUTION INCIDENTS, NITROACYL HALIDES

2654. 4-Chlorobenzoyl azide
[14848-01-2]

$C_7H_4ClN_3O$




1. Cobern, D. *et al.*, *Chem. & Ind.*, 1965, 1625; 1966, 375
2. Tyabji, M. M., *Chem. & Ind.*, 1965, 2070

A violent explosion occurred during vacuum distillation of 4-chlorophenyl isocyanate, prepared by Curtius reaction from the azide. It was found by IR spectroscopy that this isocyanate (as well as others prepared analogously) contained some unchanged azide, to which the explosion was attributed. The use of IR spectroscopy to check for absence of azides in isocyanates is recommended before distillation [1]. Subsequently, the explosion was attributed to free hydrogen azide, produced by hydrolysis of the unchanged acyl azide [2].

See other ACYL AZIDES

2655. 2-Trifluoromethylphenyllithium
[49571-31-5]

C₇H₄F₃Li




(Preparative hazard for 2-)

See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2656. 3-Trifluoromethylphenyllithium
[368-49-0]


C₇H₄F₃Li



See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2657. 4-Trifluoromethylphenyllithium
[2786-01-8]


C₇H₄F₃Li



See entry ORGANOLITHIUM REAGENTS *See other* HALO-ARYLMETALS

2658. Sodium 2-benzothiazolylthiolate
[26249-01-4]

C₇H₄NNaS₂




Maleic anhydride

See Maleic anhydride: Bases, etc.

2659. Benzenediazonium-2-carboxylate
[17333-86-7]

$C_7H_4N_2O_2$



1. Yaroslavsky, S., *Chem. & Ind.*, 1965, 765
2. Sullivan, J. M., *Chem. Eng. News*, 1971, **49**(16), 5
3. Embree, H. D., *Chem. Eng. News*, 1971, **49**(30), 3
4. Mich, T. K. *et al.*, *J. Chem. Educ.*, 1968, **45**, 272
5. Matuszak, C. A., *Chem. Eng. News*, 1971, **49**(24), 39
6. Logullo, F. M. *et al.*, *Org. Synth.*, 1973, Coll. Vol. 5, 54–59
7. Stiles, R. M. *et al.*, 1963, **85**, 1795, footnote 30a
8. Rabideau, P. W., *Org. Prep. Proced. Int.*, 1986, **18**, 113–116

The isolated internal salt is explosive, and should only be handled in small amounts [1]. This has been amply confirmed [2,3]. The salt may be precipitated during the generation of benzyne by diazotisation of anthranilic acid at ambient temperature; some heating is essential for complete decomposition [4]. During decomposition of the isolated salt, too-rapid addition to hot solvent caused a violent explosion [5]. The published procedure [6] must be closely followed for safe working. It was claimed that, contrary to earlier beliefs, diazonium carboxylate hydrohalide salts are also shock-sensitive explosives [7], but this is now firmly refuted. The hydrochloride salt cannot be detonated by impact, but if deliberately ignited by flame contact, it will flash. Thus, when a 30 g quantity was ignited, it burned away very rapidly without explosion. The hydrochloride appears to have other practical advantages, in that the yield of dibenzobarrelene, formed from benzyne and naphthalene is much higher when the hydrochloride is used as benzyne precursor, rather than the internal salt [8].

Aniline, or Isocyanides

Huisgen, R. *et al.*, *Chem. Ber.*, 1965, **98**, 4104

It reacts explosively with aniline, and violently with aryl isocyanides.

See other AMINATION INCIDENTS

1-Pyrrolidinylcyclohexene


Kuehne, M. E., *J. Amer. Chem. Soc.*, 1962, **84**, 841

Decomposition of benzenediazonium-2-carboxylate in the enamine at 40°C caused a violent explosion.

See other DIAZONIUM CARBOXYLATES

2660. Benzenediazonium-3-carboxylate
[17333-87-8]

$C_7H_4N_2O_2$




Potassium *O, O*-diphenyl dithiophosphate

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 10)

See other DIAZONIUM CARBOXYLATES

2661. Benzenediazonium-4-carboxylate
[17333-88-9]

$C_7H_4N_2O_2$




Potassium *O, O*-diphenyl dithiophosphate

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 10)

See other DIAZONIUM CARBOXYLATES

2662. 2-Nitrobenzonitrile
[612-24-8]

$C_7H_4N_2O_2$



1. Miller, C. S., *Org. Synth.*, 1955, Coll. Vol. 3, 646


2. Partridge, M. W., private comm., 1968

When the published method [1] for preparing the 4-isomer is used to prepare 2-nitrobenzonitrile, a moderate explosion often occurs towards the end of the reaction period. (This may be owing to formation of nitrogen trichloride as a by-product.) Using an alternative procedure, involving heating 2-chloronitrobenzene with copper(I) cyanide in pyridine for 7 h at 160°C, explosions occurred towards the end of the heating period in about 20% of the preparations [2].

See 4-Chloro-3-nitrotoluene: Copper(I) cyanide, Pyridine
See other CYANO COMPOUNDS, NITROARYL COMPOUNDS

2663. 4-Hydroxybenzenediazonium-3-carboxylate
[68596-89-4]


C₇H₄N₂O₃



1. Auden, W., *Chem. News*, 1899, **80**, 302
 2. Puxeddu, E., *Gazz. Chim. Ital.*, 1929, **59**, 14
- It explodes at either 155° [1] or 162°C [2], probably depending on the heating rate.
See other DIAZONIUM CARBOXYLATES

2664. 3-Nitrobenzoyl nitrate
[101672-19-9]


C₇H₄N₂O₆



Francis, F. E., *J. Chem. Soc.*, 1906, **89**, 1
The nitrate explodes if heated rapidly, like benzoyl nitrate.
See other ACYL NITRATES, NITROARYL COMPOUNDS

2665. 3,5-Dinitrosalicylic acid (2-Hydroxy-3,5-dinitrobenzoic acid)
[609-99-4]

C₇H₄N₂O₇




The editor would be surprised if it proved impossible to detonate this analytical reagent, the carboxylation product of 2,4-dinitrophenol, especially in contact with bases. There is no record of problems in use.

See other POLYNITROARYL COMPOUNDS
See 2,4-Dinitrophenol; 2,4-Dinitrophenylhydrazine

2666. 3,5-Dinitro-2-methylbenzenediazonium-4-oxide
[70343-16-7]


C₇H₄N₄O₅



Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1979, **44**, 2503–2504
A very shock-sensitive explosive.
See other ARENEDIAZONIUM OXIDES, POLYNITROARYL COMPOUNDS

2667. 3,5-Dinitro-6-methylbenzenediazonium-2-oxide
[70343-16-7]


C₇H₄N₄O₅



Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1979, **44**, 2503–2504
A very shock-sensitive explosive which melts at 152°C, then decomposes explosively.
See other ARENEDIAZONIUM OXIDES, POLYNITROARYL COMPOUNDS

2668. 2-Cyano-4-nitrobenzenediazonium hydrogen sulfate
[68597-10-4]

C₇H₄N₄O₆S



Sulfuric acid
Grewer, T., *Chem. Ing. Tech.*, 1975, **47**, 233–234

DTA examination of a 35% solution of the diazonium salt in sulfuric acid showed 3 exotherms, corresponding to hydrolysis of the nitrile group (peak at 95°), decomposition of the diazonium salt (peak at 160°) and loss of the nitro group (large peak at 240°C). Adiabatic decomposition of the solution from 50°C also showed 3 steps, with induction periods of around 30, 340 and 380 min, respectively.

See THERMAL STABILITY OF REACTION MIXTURES AND SYSTEMS

See other CYANO COMPOUNDS, DIAZONIUM SULFATES, INDUCTION PERIOD INCIDENTS, NITROARYL COMPOUNDS

2669. Poly(2,4-hexadiyne-1,6-ylene carbonate) (Poly(oxycarbonyl-2,4-hexadiyne-1,6-diyl))

[63354-77-8]

$(C_7H_4O_3)_n$



Kuehling, S. *et al.*, *Macromolecules*, 1993, **23**(19), 4192


This polymer, prepared by copper catalysed air oxidation of dipropynyl carbonate, exploded readily on warming or manipulation. The corresponding adipate and sebacate esters are non-explosive. The authors wonder if peroxide formation is involved; such explanation seems otiose.

See other ACETYLENIC COMPOUNDS

2670. Silver 3-cyano-1-phenyltriazen-3-ide

[70324-20-8]

$C_7H_5AgN_4$



Bretschneider, H., *Monats.*, 1950, **81**, 981


It is explosive.

See other CYANO COMPOUNDS, SILVER COMPOUNDS, TRIAZENES

2671. 6-Chloro-2-nitrobenzyl bromide

[56433-01-3]

$C_7H_5BrClNO_2$




See entry NITROBENZYL COMPOUNDS (reference 2)

See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2672. 2,6-Dinitrobenzyl bromide
[3013-38-5]

$C_7H_5BrN_2O_4$



Potassium phthalimide


Reich, S. *et al.*, *Bull. Soc. Chim. Fr.* [4], 1917, **21**, 119

Reaction temperature in absence of solvent must be below 130–135°C to avoid explosive decomposition.

See entry NITROBENZYL COMPOUNDS See other POLYNITROARYL COMPOUNDS

2673. Phenylchlorodiazirine
[4460-46-2]

$C_7H_5ClN_2$



1. Wheeler, J. J., *Chem. Eng. News*, 1970, **48**(30), 10
2. Manzara, A. P., *Chem. Eng. News*, 1974, **52**(42), 5
3. Padwa, A. *et al.*, *Org. Synth.*, 1981, **60**, 53–57

The neat material is about three times as shock-sensitive as glyceryl nitrate, and should not be handled undiluted [1]. It exploded during vacuum distillation at 3.3 mbar from a bath at 140°C. Impact- and spark-sensitivities were determined, and autoignition occurred after 30 s at 107°C [2]. An explosion occurred during distillation at a pot temperature of 80°C. Full handling precautions are detailed. Dilution of neat material with solvent eliminates shock-sensitivity [3].

See other AUTOIGNITION INCIDENTS


Phenylacetylene

See 3-Chloro-1,3-diphenylcyclopropene

See other DIAZIRINES

2674. N-Chloro-5-phenyltetrazole
[65037-43-6]


C₇H₅ClN₄



Gallacher, T. C. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 420
Thermal decomposition is vigorous, and was explosive on one occasion.
See other N-HALOGEN COMPOUNDS, TETRAZOLES

2675. Benzoyl chloride
[98-88-4]

C₇H₅ClO



Aluminium chloride, Naphthalene
See Aluminium chloride: Benzoyl chloride, etc.


Dimethyl sulfoxide
See Dimethyl sulfoxide: Acyl halides

MRH 1.30/tr.

Other reactants
Yoshida, 1980, 58
MRH values calculated for 13 combinations with oxidants are given.
See other ACYL HALIDES

2676. 3-Chloroperoxybenzoic acid
[937-14-4]

C₇H₅ClO₃




2-Bromo-4-methylpyridine
1. Brand, W. W., *Chem. Eng. News*, 1978, **56**(24), 88
2. Shanley, E. S., *Chem. Eng. News*, 1978, **56**(29), 43

Vacuum evaporation of the product of unheated conversion of the pyridine to its *N*-oxide with 5% excess of the peroxy acid in chloroform gave a residue which decomposed violently [1]. This was attributed to the relative stability of the peroxy acid in the cold pure state, which when concentrated and finally heated with other materials underwent accelerating decomposition [2].

See other PEROXYACIDS

2677. 1,3-Benzodithiolium perchlorate
[32283-21-9]




1. Pelter, A., *Chem. Abs.*, 1981, **95**, 97643 (ref. there to *Tetrahedron Lett.* original is incorrect)
2. Degari, I. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1976, 1886
3. Anon., *Univ. Safety Assoc. Safety News*, 1978, (10), 11–12

Highly friction sensitive, with considerable explosion hazard [1]. Repeating a published method ([2], where explosion at the m.p., 180°C is noted), the product was dried in a desiccator. When the lid (ground glass flange) was slid off, the friction initiated explosion of a little of the compound deposited on the flange, and a severe explosion resulted [3].

See other GLASS INCIDENTS, NON-METAL PERCHLORATES

2678. 2-Chloro-4-nitrobenzyl chloride
[50274-95-8]




See entry NITROBENZYL COMPOUNDS (reference 2)

See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2679. 4-Chloro-2-nitrobenzyl chloride
[938-71-6]


$C_7H_5Cl_2NO_2$



See entry NITROBENZYL COMPOUNDS (reference 2)
See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2680. α -(Difluoroamino)- α,α -dinitrotoluene ((Difluoroaminodinitromethyl) benzene, *N,N*-Difluoro- α,α -dinitrobenzenemethanamine)
[157066-32-5]


$C_7H_5F_2N_3O_4$



See 1,1-DINITRODIFLUORAMINES, See other *N*-HALOGEN COMPOUNDS, POLYNITRO-ALKYL COMPOUNDS

2681. 2-Iodylbenzoic acid (1-Hydroxy-1,2-benziodoxol-3-one-1-oxide)
[64297-64-9] (Acid)
[131-62-4] (Iodoxolone)

$C_7H_5IO_4$




Plumb, J. B., *Chem. Eng. News*, 1990, **68**(29), 3

The pure dry acid explodes at 233°C, and violently if confined, possibly <200°C. It is also impact-sensitive, exploding under a hammer blow, or under impact of a 534 g steel ball falling from a height of 1 m. Several of its salts (ammonium, potassium, sodium, silver, barium, calcium and magnesium) are also explosive. The acid is precursor to the 'Dess-Martin periodinane' mild oxidant, and is produced when the latter is treated with water.

Calcium 2-iodylbenzoate
See 1,1,1-Triacetoxyl-1,2-benziodoxol-3-one
See other IODINE COMPOUNDS, ORGANIC ACIDS

2682. Potassium phenyldinitromethanide
[2918-51-8]


C₇H₅KN₂O₄



Rohman, A. *et al.*, *J. Org. Chem.*, 1976, **41**, 124
It is unwise to use more than 0.02 mol of potassium salts of dinitroalkanes as they are explosive.
See related *aci*-NITRO SALTS, POLYNITROALKYL COMPOUNDS

2683. Potassium 2,4,6-trinitrotoluene
[]


C₇H₅KN₃O₆



See entry NITROAROMATIC-ALKALI HAZARDS; potassium radical salts

2684. Potassium *O*-*O*-benzoylmonoperoxosulfate
[]

C₇H₅KO₅S



Alone, or Sulfuric acid
Willstätter, R. *et al.*, *Ber.*, 1909, **42**, 1839


The anhydrous salt explodes on grinding, and the monohydrate on heating to 70–80°C, or in contact with sulfuric acid.

See other DIACYL PEROXIDES, PEROXOACID SALTS

2685. Phenyl isocyanate

[103-71-9]

C₇H₅NO



Energy of decomposition (in range 240–460°C) measured as 0.41 kJ/g.


See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ORGANIC ISOCYANATES

2686. 2-Nitrobenzaldehyde

[552-89-6]

C₇H₅NO₃



1. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn, 1986
2. Grewer, T. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 657–664, Basle, SSCI, 1980
3. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987
4. Anon., *Loss Prev. Bull.*, 1989, (087), 27–29

Energy of decomposition (in range 220–370°C) measured as 2.124 kJ/g by DSC, and T_{ait24} was determined as 123°C by adiabatic Dewar tests, with an apparent energy of activation of 116 kJ/mol [1]. Initial decomposition temperature by ARC was 176°C. A very high rate of pressure increase was observed in exothermic decomposition, and further work on homogeneous decomposition under confinement has been reported [2,3]. Exothermic decomposition during vacuum distillation of an atypical 600 kg batch of 2-nitrobenzaldehyde containing other isomers, and which accelerated to explosion, has been reported and discussed [4].

See entry NITROBENZALDEHYDES

Active carbon

Mixtures with active carbon (1.5:1 mol) deflagrate at 1.25 cm/min.

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 3)

Pyrrole

Sorrell, T. N., *Inorg. Synth.*, 1980, **20**, 162


In the preparation of 5,10,15,20-tetrakis(2-nitrophenyl)porphyrin by dropwise addition of the aldehyde in acetic acid, the exothermic reaction may become so violent that solvent sprays out of the condenser. However, a low yield may ensue if the reaction is insufficiently vigorous, so a careful compromise is required.

See other ALDEHYDES, NITROARYL COMPOUNDS

2687. 3-Nitrobenzaldehyde

[99-61-6]

$C_7H_5NO_3$



1. Lange, J. *et al.*, *Chem. & Ind.*, 1967, 1424
2. Klais, O. *et al.*, *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 3, C24–C34, Rugby, IChE, 1983
3. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

Delays in working up the crude product caused violent explosions during attempted vacuum distillation. An alternative method of crystallisation is described [1]. There is a very high rate of pressure increase in exothermic decomposition [2]. Energy of decomposition (in range 180–420°C) measured as 2.19 kJ/g by DSC, and T_{ait24} was determined as 147°C by adiabatic Dewar tests, with an apparent energy of activation of 168 kJ/mol [3]. The initial decomposition temperature by ARC was 166°C.


See entry NITROBENZALDEHYDES

See other ALDEHYDES, NITROARYL COMPOUNDS

2688. 4-Nitrobenzaldehyde

[555-16-5]

$C_7H_5NO_3$



1. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

Energy of decomposition (in range 220–450°C) measured as 1.602 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 171°C by adiabatic Dewar tests, with an apparent energy of activation of 202 kJ/mol [1], The initial decomposition temperature by ARC was 226°C. A very high rate of pressure increase was observed in exothermic decomposition.

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION


See entry NITROBENZALDEHYDES

See other ALDEHYDES, NITROARYL COMPOUNDS

2689. Benzoyl nitrate

[6786-32-9]

$C_7H_5NO_4$



1. Francis, F. E., *J. Chem. Soc.*, 1906, **89**, 1
2. Ferrario, E., *Gazz. Chim. Ital.* [1], 1901, **40**, 99

An unstable liquid which is capable of distillation under reduced pressure, but which explodes violently on rapid heating at ambient pressure [1]. It may also explode on exposure to light [2].

See other IRRADIATION DECOMPOSITION INCIDENTS

Water

Francis, F. E., *J. Chem. Soc.*, 1906, **89**, 1


Reactivity of benzoyl nitrate towards moisture is so great that attempted filtration through an undried filter paper causes explosive decomposition (possibly involving cellulose nitrate?).

See other ACYL NITRATES

2690. 3-Hydroxy-4-nitrobenzaldehyde

[704-13-2]

$C_7H_5NO_4$




Initial decomposition temperature by DSC was 200°C.

See entry NITROBENZALDEHYDES See other ALDEHYDES, NITROARYL COMPOUNDS

2691. 4-Hydroxy-3-nitrobenzaldehyde
[3011-34-5]

C₇H₅NO₄




Initial decomposition temperature by DSC was 200°C.

See entry NITROBENZALDEHYDES *See other* ALDEHYDES, NITROARYL COMPOUNDS

2692. 5-Hydroxy-2-nitrobenzaldehyde
[42454-06-8]

C₇H₅NO₄




Initial decomposition temperature by DSC was 175°C.

See entry NITROBENZALDEHYDES *See other* NITROARYL COMPOUNDS

2693. 2-Nitrobenzoic acid
[552-16-9]

C₇H₅NO₄




Energy of decomposition (in range 230–450°C) measured as 1.72 kJ/g by DSC, and T_{ait24} was determined as 189°C by adiabatic Dewar tests, with an apparent energy of activation of 156 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2694. 3-Nitrobenzoic acid
[121-92-6]

C₇H₅NO₄




Energy of decomposition (in range 320–410°C) measured as 1.734 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 245°C by adiabatic Dewar tests, with an apparent energy of activation of 151 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2695. 4-Nitrobenzoic acid
[62-23-7]

C₇H₅NO₄



Energy of decomposition (in range 320–360°C) measured as 2.234 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 247°C by adiabatic Dewar tests, with an apparent energy of activation of 131 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Potassium hydroxide


Mixtures of the acid with potassium hydroxide (1:2 mol) readily deflagrated, at a rate of 5.1 cm/min.

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 3)

See other DEFLAGRATION INCIDENTS, NITROARYL COMPOUNDS, ORGANIC ACIDS

2696. 2,5-Pyridinedicarboxylic acid
[100-26-5]

C₇H₅NO₄




Preparative hazard

See Nitric acid: 5-Ethyl-2-methylpyridine

See other ORGANIC ACIDS

2697. 4-Azidobenzaldehyde
[24173-36-2]

C₇H₅N₃O



Lahti, P. M., *Chem. Eng. News*, 1998, **76** (28), 8

Very shock sensitive; explodes in drop-weight test at 11 kg-cm.


Energy of decomposition (in range 110–220°C) measured as 1.67 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (REFERENCE 2)

See other ALDEHYDES, ORGANIC AZIDES

2698. Benzoyl azide
[582-61-6]

C₇H₅N₃O




Bergel, F., *Angew. Chem.*, 1927, **40**, 974

The sensitivity towards heat of this explosive compound is increased by previous compression, confinement and presence of impurities. Crude material exploded violently between 120 and 165°C.

See other ACYL AZIDES, CATALYTIC IMPURITY INCIDENTS

2699. Carboxybenzenesulfonyl azide
[56743-33-0]

C₇H₅N₃O₄S




Anon., *Res. Discl.*, 1975, **134**, 44

The azide (structure not stated) is not impact-sensitive, but decomposes explosively at 120°C. Blending with 75% of polymer as diluent eliminated the explosive decomposition.

See other ACYL AZIDES

2700. 2-Nitrophenylsulfonyldiazomethane
[49558-46-5]

C₇H₅N₃O₄S




Wagenaar, A. *et al.*, *J. Org. Chem.*, 1974, **39**, 411–413

Severe explosions occurred when the cold crystalline solid was allowed to warm to ambient temperature. Normal manipulation of dilute solutions appears feasible, but the solid must be handled at lowest possible temperatures with full safety precautions. The 4-isomer appears much more stable.

See other DIAZO COMPOUNDS, NITROARYL COMPOUNDS

2701. 2,4,6-Trinitrotoluene
[118-96-7]

$C_7H_5N_3O_6$



NSC 314, 1981

Heat of decomposition was determined as 5.1 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (REFERENCE 1)

Added impurities

Häuptli, H., private comm., 1972

During investigation of effect of 1% of added impurities on the thermal explosion temperature of TNT (297°C), it was found that fresh red lead, sodium carbonate and potassium hydroxide reduced the explosion temperatures to 192, 218, and 192°C, respectively.

See Potassium: Nitrogen-containing explosives

See other CATALYTIC IMPURITY INCIDENTS

Bases

MRH Sodium hydroxide 5.40/tr.

Capellos, C. *et al.*, *Chem. Abs.*, 1982, **96**, 88003

The formation of *aci*-quinonoid transient intermediates by electronic excitation has been identified spectroscopically. Proton abstractors (bases) such as sodium hydroxide, potassium iodide or tetramethylammonium octahydrotriborate induce deflagration in molten TNT.

See NITROAROMATIC-ALKALI HAZARDS

Metals, Nitric acid

Kovache, A. *et al.*, *Mé m. Poudres*, 1952, **34**, 369–378

Trinitrotoluene in contact with nitric acid and lead or iron produces explosive substances which may readily be ignited by shock, friction or contact with nitric or sulfuric acids. Such materials have been involved in industrial explosions.

See HEAVY METAL DERIVATIVES

Other reactants

Yoshida, 1980, 254

MRH values calculated for 3 combinations with catalytically active materials are given.

Potassium hydroxide

Copisarow, M., *Chem. News*, 1915, **112**, 283–284

TNT and potassium hydroxide in methanol will interact even at -65°C to give explosive *aci*-nitro salts (presumably *o*-quinonoid, or possibly Meisenheimer complexes). The explosion temperature is lowered to 160°C by the presence of a little potassium hydroxide.

See entry aci-NITROQUINONOID COMPOUNDS

Sodium carbonate, Water

Iida, M. *et al.*, *Chem. Abs.*, 1997, **127**, 222627q

TNT reacts vigorously with sodium carbonate, in presence of water at 100°C and above. The solid residue resulting ignites on contact with strong acid.

See 2,4,5-Trinitrotoluene; next below

See NITROAROMATIC-ALKALI HAZARDS

Sodium dichromate, Sulfuric acid


See Sodium dichromate: Sulfuric acid, Trinitrotoluene

See other POLYNITROARYL COMPOUNDS

2702. 2,4,5-Trinitrotoluene

[610-25-3]

$\text{C}_7\text{H}_5\text{N}_3\text{O}_6$



Sodium carbonate

Biasutti, 1981, 78


Sodium carbonate was added to a heated tank of washed TNT to neutralise excess acidity. It is known that the assym. isomer of TNT reacts with sodium carbonate to form unstable compounds, and decomposition led to a fire and detonation of 5 tonnes of explosive.

See other NITROAROMATIC-ALKALI HAZARDS, POLYNITROARYL COMPOUNDS

2703. 3-Methyl-2,4,6-trinitrophenol

[602-99-3]

$\text{C}_7\text{H}_5\text{N}_3\text{O}_7$




Sorbe, 1968, 153

It explodes above 150°C.
See other POLYNITROARYL COMPOUNDS

2704. Benzimidazolium 1-nitroimidate
[52096-22-7]


C₇H₅N₄O₂



It explodes at the m.p., 169°C.
See entry N-AZOLIUM NITROIMIDATES

2705. 1-Nitro-3-(2,4-dinitrophenyl)urea
[22751-18-4]


C₇H₅N₅O₇



McVeigh, J. L. *et al.*, *J. Chem. Soc.*, 1945, 621
It is an explosive of lower power, but of greater impact- or friction-sensitivity than picric acid, so should not be stored in bottles with ground-in stoppers.
See other N-NITRO COMPOUNDS, POLYNITROARYL COMPOUNDS

2706. N,2,4,6-Tetranitro-N-methylaniline (Tetryl)
[479-45-8]

C₇H₅N₅O₈



Hydrazine
See Hydrazine: Oxidants

Trioxxygen difluoride


See 'Trioxxygen difluoride': Various materials

See other N-NITRO COMPOUNDS, POLYNITROARYL COMPOUNDS

2707. 1-Heptene-4,6-diyne

[]

C₇H₆



1. Armitage, J. B. *et al.*, *J. Chem. Soc.*, 1952, 1994

2. *CHETAH*, 1990, 182


It decomposes explosively with great ease [1] but was shown to be exploded by a detonator, not by mechanical shock [2].

See other ALKYNES, ALLYL COMPOUNDS

2708. 2-Azidomethylbenzenediazonium tetrafluoroborate

[59327-98-9]

C₇H₆BF₄N₅



Trichloroacetonitrile

Kreher, R. *et al.*, *Tetrahedron Lett.*, 1976, 4260

Thermal decomposition of the salt in nitriles produces 2-substituted quinazolines.


In trichloroacetonitrile it proceeded explosively.

See other DIAZONIUM TETRAHALOBORATES, ORGANIC AZIDES

2709. 5-Fluoro-2-methylphenylmagnesium bromide (Bromo(5-fluoro-2-methylphenyl)magnesium)

[]

C₇H₆BrFMg



Preparative Hazard

Jasiewicz, M. L. *Chem. & Ind.*, 1995, 246


A considerable explosion resulted from preparation of this compound (which presumably exists with several tightly complexed solvent molecules) in limited tetrahydrofuran under reflux. Investigation showed that there is an exotherm at 130°C, giving 440 J/g, followed by a more violent one at 300°C giving 690 J/g. Using a large excess of solvent no problems are observed.

See other GRIGNARD REAGENTS

2710. 2-Nitrobenzyl bromide

[3958-60-9]

C₇H₆BrNO₂



1. Biasutti, 1981, 114
2. Cardillo, P. *et al.*, *Rivista Combust.*, 1982, **36**, 304
3. Cardillo, P. *et al.*, *Proced. 34th Ital. Chem. Congr.*, 'Safety of Chemical Processes', 71–82, Milan, 1983


During the first large scale preparation of the bromide from 2-nitrotoluene, the reactor exploded violently, though hundreds of runs had been done uneventfully in a pilot scale plant [1]. Following a thermal explosion during drying of the compound at a slightly elevated temperature, its thermal stability has been determined by TGA, DSC and ARC techniques. The latter shows that exothermic decomposition begins at 100°C and proceeds in two stages with abundant gas evolution. The former techniques indicate onset of decomposition at 130° and 112°C, respectively [2]. Further work on the 3- and 4-isomers showed them to be marginally more stable than the 2-nitro compound.

See 2-Nitrobenzyl chloride

See other GAS EVOLUTION INCIDENTS, NITROBENZYL COMPOUNDS, NITROARYL COMPOUNDS

2711. 2-Chloro-4-nitrotoluene
[121-86-8]

C₇H₆ClNO₂



Other reactants

Yoshida, 1980, 112

MRH values calculated for 13 combinations with oxidants are given.

Sodium hydroxide

MRH 3.39/tr.

1. Anon., *ABCM Quart. Safety Summ.*, 1962, **33**, 20
2. *MCA Case History No. 907*


The residue from vacuum distillation of crude material (contaminated with sodium hydroxide) exploded after showing signs of decomposition. Experiment showed that 2-chloro-4-nitrotoluene decomposes violently when heated at 170–200°C in presence of alkali. Thorough water washing is therefore essential before distillation is attempted [1]. A similar incident was reported with mixed chloronitrotoluene isomers [2].

See other NITROAROMATIC-ALKALI HAZARDS

See other NITROARYL COMPOUNDS

2712. 4-Chloro-3-nitrotoluene
[890-60-1]

C₇H₆ClNO₂



Copper(I) cyanide, Pyridine

1. Hub, L., *Proc. Chem. Process. Haz. Symp. VI* (Manchester, 1977), No. 49, 39–46, Rugby, IChE, 1977
2. Hub, L., private comm., 1977

Industrial preparation of 4-cyano-3-nitrotoluene by heating the reaction components at around 170°C for 6 h led to an explosion in 1976. Subsequent investigation by DSC showed that the cyano compound in presence of the starting materials exhibited an exotherm at 180°C. After 6 h reaction, this threshold temperature fell to 170°C. Isothermal use of a safety calorimeter showed that a large exotherm occurred during the first hour of reaction and that, in absence of strong cooling,

the reaction accelerated and the vessel contents were ejected by the vigour of the decomposition [2].


See -nitrobenzonitrile, REACTION SAFETY CALORIMETRY

See other NITROARYL COMPOUNDS

2713. 2-Nitrobenzyl chloride

[612-23-7]

$C_7H_6ClNO_2$



Cardillo, P. *et al.*, *Proc. 34th Ital. Chem. Congr., 'Safety of Chem. Processes'*, 71–82, Milan, 1983

Following an explosion during drying of 2-nitrobenzyl bromide, the thermal stability of all six isomeric nitrobenzyl halides was investigated by TGA, DSC and ARC techniques. All the halides decomposed exothermally with abundant gas evolution, and the 2-nitro chloride is less thermally stable than the 3- or 4-isomers.

See 2-Nitrobenzyl bromide

See other BENZYL COMPOUNDS, NITROBENZYL COMPOUNDS, NITROARYL COMPOUNDS

2714. 4-Nitrobenzyl chloride

[100-14-1]

$C_7H_6ClNO_2$

$O_2NC_6H_4CH_2Cl$

T_{ait24} was determined as 189°C by adiabatic Dewar tests, with an apparent energy of activation of 124 kJ/mol.


See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROARYL COMPOUNDS, NITROBENZYL COMPOUNDS

2715. 2-Chloro-5-nitrobenzyl alcohol

[80866-80-4]

$C_7H_6ClNO_3$




See entry NITROBENZYL COMPOUNDS (reference 3)

See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2716. 4-Chloro-2-nitrobenzyl alcohol
[22996-18-5]


C₇H₆ClNO₃



See entry NITROBENZYL COMPOUNDS (reference 3)
See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2717. 4-Chloro-3-nitrobenzyl alcohol
[55912-20-4]


C₇H₆ClNO₃



See entry NITROBENZYL COMPOUNDS (reference 3)
See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS


2718. 5-Chloro-2-nitrobenzyl alcohol
[73033-58-6]

C₇H₆ClNO₃




See entry NITROBENZYL COMPOUNDS (reference 3)
See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2719. 4-Chloro-2-methylbenzenediazonium salts
[27165-08-8] (ion)



Sodium hydrogen sulfide, or Sodium disulfide, or Sodium polysulfide
See entry DIAZONIUM SULFIDES AND DERIVATIVES (references 1–4)
See other DIAZONIUM SALTS

2720. Sodium 4-chloro-2-methylphenoxide
[52106-86-2]




Anon., private comm., 1984

A dry solution of the sodium salt in n-butanol was usually prepared by azeotropic drying. Use of excessively wet recovered butanol led to complete removal of the butanol with the water and heating of the dry salt at 200°C, when rapid decomposition occurred, leaving a glowing carbonised residue.

See 4-Chloro-2-methylphenol: Sodium hydroxide


2721. Potassium 2,6-dinitrotoluene
[]



The o- and m-isomers are similar.

See entry NITROAROMATIC-ALKALI HAZARDS; potassium radical salts

2722. Potassium 4-methoxy-1-*aci*-nitro-3,5-dinitro-2,5-cyclohexadienonide
[1270-21-9] **C₇H₆KN₃O₇**




1. Meisenheimer, J., *Ann.*, 1902, **323**, 221
2. Millor, H. D. *et al.*, *Sidgwick's Organic Chemistry of Nitrogen*, 396, Oxford, Clarendon Press, 3rd edn, 1966

The product of interaction of trinitrobenzene and conc. aqueous potassium hydroxide in methanol is explosive, and analyses as the hemihydrate of a hemiacetal of the *aci-p*-quinonoid form of picric acid [1], and/or the mesomeric *o*-forms [2].

See other aci-NITROQUINONOID COMPOUNDS, POLYNITROARYL COMPOUNDS

2723. Thallium *aci*-phenylnitromethanide
[53847-48-6] **C₇H₆NO₂Tl**




McKillop, A. *et al.*, *Tetrahedron*, 1974, **30**, 1369

No attempt should be made to isolate or dry this compound, as it is treacherously explosive.

See other HEAVY METAL DERIVATIVES, aci-NITRO SALTS

2724. Benzimidazole (Benzodiazole)
[51-17-2] **C₇H₆N₂**



CHETAH, 1990, 188


Shock sensitive (which seems unlikely but not impossible).

See Benzotriazole

2725. Phenyl diazomethane

[766-91-6]

$C_7H_6N_2$



Schneider, M. *et al.*, *Tetrahedron*, 1976, **36**, 621


The need for care in distillation of phenyl diazomethane is stressed. Previously it has been used as prepared in solution.

See other DIAZO COMPOUNDS

2726. 2,4-Dinitrotoluene

[121-14-2]

$C_7H_6N_2O_4$



(MCA SD-93, 1966); NSC 658, 1976

1. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
2. MCA SD-93, 1966
3. Biasutti, 1981, 143
4. Bond, J., *Loss Prev. Bull.*, 1983, 050, 18
5. Anon., *Loss Prevention Bull.*, 1989, **88**, 13

T_{ait24} was determined as 181°C by adiabatic Dewar tests, with an apparent energy of activation of 153 kJ/mol [1]. The commercial material, containing some 20% of the 2,6-isomer, decomposes at 250°, but at 280°C decomposition becomes self-sustaining. Prolonged heating below these temperatures may also cause some decomposition, and the presence of impurities may decrease the decomposition temperatures [2]. Although not considered to be explosive, several cases of detonation in manufacture or storage have been reported [3], including an explosion when a jug of dinitrotoluene was dropped into a machine [4]. An isolated pipe filled with crude dinitrotoluene at a temperature between 125 and 135°C ran away to

detonation over 3.5 hours. Having been separated from spent acid, but not washed, it contained traces of both sulfuric and nitric acids. Investigation showed that this combination destabilised the material more than either separately. A sample (2 phase) of approximate composition dinitrotoluene 45%, nitric acid 5%, sulfuric acid 40%, water 10% showed an exothermic reaction on ARC as low as 66°C [5].

Alkali

Bateman, T. L. *et al.*, *Loss. Prev.*, 1974, **8**, 117–122

Dinitrotoluene held at 210°C (rather than the 125° intended) for 10 days in a 50 mm steam-heated transfer pipeline exploded. Subsequent tests showed decomposition at 210°C (producing a significant pressure rise) in 1 day, and presence of sodium carbonate (but not rust) reduced the induction period. A maximum handling temperature of 150°C was recommended, (when the induction period was 32 days, or 14 days for alkali-contaminated material).

See other CATALYTIC IMPURITY INCIDENTS, INDUCTION PERIOD INCIDENTS, NITROAROMATIC–ALKALI HAZARDS

Nitric acid

MRH 4.56/tr.

See Nitric acid: Nitroaromatics

Other reactants

Yoshida, 1980, 159

MRH values calculated for 10 combinations, largely with catalytically active materials, are given.

Sodium oxide

Anon., *Rept. of Explos. Div.*, Ottawa, Dept. of Mines, 1929

Admixture of the two solids caused a rapid reaction and fire.


See other NITROAROMATIC–ALKALI HAZARDS

See other POLYNITROARYL COMPOUNDS

2727. 5-Phenyltetrazole

[3999-10-8]

C₇H₆N₄



Wedekind, E., *Ber.*, 1898, **31**, 948


It explodes on attempted distillation.

See other TETRAZOLES

2728. *N*-Azidocarbonylazepine

[]

C₇H₆N₄O



Chapman, L. E. *et al.*, *Chem. & Ind.*, 1966, 1266


It explodes on attempted distillation.

See other ACYL AZIDES

2729. *N*-Phenyl-1,2,3,4-thiaziazolamine (Phenylaminothia-2,3,4-triazole)

[13078-30-3]

C₇H₆N₄S



CHETAH, 1990, 182


Shock sensitive.

See other HIGH NITROGEN COMPOUNDS, N-S COMPOUNDS

2730. Phenyl diazidomethane

[17122-97-3]

C₇H₆N₆




Barash, L. *et al.*, *J. Amer. Chem. Soc.*, 1967, **89**, 3391

A sample of the *gem*-diazide detonated on contact with a hot glass surface during ampoule sealing operations.

See other ORGANIC AZIDES

†2731. Benzaldehyde
[100-52-7]

C₇H₆O



HCS 1980, 191

Other reactants

Yoshida, 1980, 333

MRH values calculated for 14 combinations with oxidants are given.

Peroxyformic acid

MRH 5.69/99+

See Peroxyformic acid: Organic materials

Propionic acid, Pyrrolidine

Kaufman, J. A., private comm., 1987


An explosion occurred when benzaldehyde, pyrrolidine (pyrrole ?) and propionic acid were heated to form porphins.

See 2-Nitrobenzaldehyde: Pyrrole

See other ALDEHYDES, REDUCANTS

2732. Benzoic acid
[65-85-0]

C₇H₆O₂



HCS 1980, 191

The finely powdered dry acid is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

Other reactants

Yoshida, 1980, 32


MRH values calculated for 12 combinations with oxidants are given.

Oxygen

Oxygen (Gas): Benzoic acid See other ORGANIC ACIDS

2733. Peroxybenzoic acid
[93-59-4]

C₇H₆O₃




1. Baeyer, A. *et al.*, *Ber.*, 1900, **33**, 1577
2. Silbert, L. S. *et al.*, *Org. Synth.*, 1963, **43**, 95

It explodes weakly on heating [1]. A chloroform solution of peroxybenzoic acid exploded during evaporation of solvent. Evaporation should be avoided or conducted with full precautions [2].

See other PEROXYACIDS

2734. Silver *N*-perchlorylbenzylamide
[89521-43-7]

C₇H₇AgClNO₃



Hennrichs, W. *et al.*, *Z. Anorg. Chem.*, 1983, **50** 6, 203–207


It explodes on impact or on heating above 105°C.

See entry PERCHLORYLAMIDE SALTS (reference 2)

See other *N*-METAL DERIVATIVES, SILVER COMPOUNDS

2735. Benzyl bromide
[100-39-0]

C₇H₇Br



HCS 1980, 200

Molecular sieve

Harris, T. D., *Chem. Eng. News*, 1979, **57**(12), 74

Redistilled bromide was stored over 4A molecular sieve previously activated at 300°C. After 8 days the closed bottle burst from internal pressure of hydrogen bromide. This was attributed to the sieve catalysing a Friedel-Craft type intermolecular condensation-polymerisation with liberation of hydrogen bromide.

See Benzyl chloride: Catalytic impurities


Benzyl alcohol: Hydrogen bromide, Iron

See other GAS EVOLUTION INCIDENTS, MOLECULAR SIEVE INCIDENTS, POLYCONDENSATION REACTION INCIDENTS

See other BENZYL COMPOUNDS

2736. 2-Toluenediazonium bromide
[54514-12-4]

C₇H₇BrN₂



Copper powder


Vogel, 1956, 606–607

During the decomposition of the diazonium solution by warming with copper bronze to give 2-bromotoluene, strong cooling must be applied immediately to prevent the nitrogen evolution becoming violent and ejecting the flask contents.

See other DIAZONIUM SALTS, GAS EVOLUTION INCIDENTS

2737. Methyl 4-bromobenzenediazoate
[67880-26-6]

C₇H₇BrN₂O




Bamberger, E., *Ber.*, 1895, **28**, 233

It explodes on heating.

See other ARENEDIAZOATES, N–O COMPOUNDS

2738. Benzyl chloride
[100-44-7]

C₇H₇Cl



HCS 1980, 202

Oehme, F., *Chem. Tech.* (Berlin), 1952, **4**, 404

During distillation of the technical chloride, hydrogen chloride was evolved, air was bubbled through to remove it, and there was an explosion. This was attributed to a polycondensation (intermolecular Friedel-Craft) reaction, possibly catalysed by oxidation products.

Energy of decomposition (in range 290–370°C) measured as 0.14 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Catalytic impurities

Naef, H., private comm., 1980

The chloride is usually (but not always) stabilised in storage by addition of aqueous alkali or anhydrous amines as acid acceptors. A 270 kg batch which was not stabilised polymerised violently when charged into a reactor. Contact of the chloride (slightly hydrolysed and acidic) with rust led to formation of ferric chloride which catalysed an intermolecular Friedel-Craft reaction to form polybenzyls with evolution of further hydrogen chloride. Contact of unstabilised benzyl chloride with aluminium, iron or rust should be avoided to obviate the risk of polycondensation.

See Benzyl bromide: Molecular sieve

Benzyl alcohol: Hydrogen bromide, Iron

1,2-Bis(chloromethyl)benzene: Catalytic impurities

See other GAS EVOLUTION INCIDENTS, POLYCONDENSATION REACTION INCIDENTS

Other reactants

Yoshida, 1980, 331

MRH values for 13 combinations with oxidants are given, many values being high.

Sodium acetate, Pyridine

Bast, -, *Proc. 1st Int. Symp. Prev. Occ. Risks Chem. Ind.*, 194–199, Heidelberg, ISSA, 1970

Benzyl acetate was prepared by addition of benzyl chloride (containing 0.6% pyridine as stabiliser) to preformed sodium acetate at 70°, followed by heating at 115°, then finally up to 135°C to complete the reaction. On one occasion, gas began to be evolved at the end of the dehydration phase, and the reaction accelerated to a violent explosion, rupturing the 25 mm thick cast iron vessel. This was attributed to presence of insufficient pyridine to maintain basicity, dissolution of iron by the

acidic mixture, and catalysis by ferric chloride of a Freidel-Craft type polycondensation reaction to polybenzyls, with evolution of hydrogen chloride, which at 130°C would produce an overpressure approaching 100 bar. Previously the chloride had been supplied in steel drums containing 10% sodium carbonate or 3% sodium hydroxide solutions as stabilisers, but the packing was changed to polythene-lined kegs and the stabiliser omitted. However, higher pyridine concentrations effectively stabilise the chloride in steel drums.

See Catalytic impurities, above

Sodium cyanide

Nolan, 1983, Case history 192


Benzyl cyanide was prepared from the chloride and sodium cyanide in aqueous methanol. Inadequate cooling of the reactor led to a fire.

See other BENZYL COMPOUNDS, GAS EVOLUTION INCIDENTS, POLYCONDENSATION REACTION INCIDENTS

2739. Sodium *N*-chloro-4-toluenesulfonamide

[127-65-1]

$C_7H_7ClNaO_2S$



1. Klundt, I. L., *Chem. Eng. News*, 1977, **55** (49), 56

2. Kirk Othmer, (4th Edn), 1993, Vol IV, 921

The anhydrous salt ('Chloramine-T') explodes at 175°C, but a small quantity exploded after storage in a bottle at ambient temperature [1]. The trihydrate may explode as low as 130°C, the anhydrous compound is less stable still [2].

Calcium carbonate, Isonitriles

Sorbe, 1968, 137

Mixtures explode when warmed on the steam-bath in presence of calcium carbonate.

Isonitriles, Anthranilates, Tetraethylammonium chloride


Bossio, R. *et al.*, *J. Heterocycl. Chem.*, 1995, **32** (4), 1115

A one pot synthesis of heterocycles involving the above reagents and catalyst sometimes showed an induction period followed by violent reaction.

See other N-HALOGEN COMPOUNDS

2740. 2-Toluenediazonium perchlorate
[69597-04-6]


C₇H₇ClN₂O₄



Hofmann, K. A. *et al.*, *Ber.*, 1906, **39**, 3146
The solid salt is explosive even when wet.
See other DIAZONIUM PERCHLORATES

2741. 4-Chloro-2-methylphenol
[1570-64-5]

C₇H₇ClO



Sodium hydroxide

Anon., *ABCM Quart. Safety Summ.*, 1957, **28**, 39


A large quantity (700 kg) of the chlorophenol, left in contact with conc. sodium hydroxide solution for 3 days, decomposed, attaining red heat and evolving fumes which ignited explosively. Although this could not be reproduced under laboratory conditions, it is believed that exothermic hydrolysis to the hydroquinone (possibly with subsequent aerobic oxidation to the quinone) occurred, the high viscosity of the liquid preventing dissipation of heat.

See Sodium 4-chloro-2-methylphenoxide

See other HALOARYL COMPOUNDS

2742. Tropylium perchlorate
[25230-72-2]

C₇H₇ClO₄



1. Ferrini, P. G. *et al.*, *Angew. Chem.*, 1962, **74**, 488


2. Barltrop, J., *Chem. Brit.*, 1980, **16**, 452

Pressing the salt through a funnel with a glass rod caused a violent explosion [1], and grinding 19 g of the dry salt in a mortar led to the same result.

See other NON-METAL PERCHLORATES

2743. 2-Tolylcopper
[20854-03-9]

C₇H₇Cu




Camus, A. *et al.*, *J. Organomet. Chem.*, 1968, **14**, 442–443

The solid 2-isomer usually exploded strongly on exposure to oxygen at 0°C, or weakly on heating above 100°C under vacuum.

See other ARYLMETALS

2744. 3-Tolylcopper
[20854-05-1]

C₇H₇Cu




Camus, A. *et al.*, *J. Organomet. Chem.*, 1968, **14**, 442–443

The solid 3-isomers usually exploded strongly on exposure to oxygen at 0°C, or weakly on heating above 100°C under vacuum.

See other ARYLMETALS

2745. 4-Tolylcopper
[5588-74-9]

C₇H₇Cu




Camus, A. *et al.*, *J. Organomet. Chem.*, 1968, **14**, 442–443

The solid 4-isomers usually exploded strongly on exposure to oxygen at 0°C, or weakly on heating above 100°C under vacuum.

See other ARYLMETALS

2746. Benzyl fluoride (α -Fluorotoluene)
[25496-08-6]

C₇H₇F




Szucs, S. S., *Chem. Eng. News.*, 1990 **68**(34), 2

A bottle exploded on storage, coating the surroundings with polymer. It is considered that glass has catalytic properties for the Friedel Crafts polymerisation reaction.

See other BENZYL COMPOUNDS, POLYCONDENSATION REACTION INCIDENTS

†2747. **2-Fluorotoluene**
[952-52-3]


C₇H₇F



See other HALOARYL COMPOUNDS

†2748. **3-Fluorotoluene**
[352-70-5]


C₇H₇F



See other HALOARYL COMPOUNDS

†2749. **4-Fluorotoluene**
[352-32-9]


C₇H₇F



See other HALOARYL COMPOUNDS

2750. 4-Iodotoluene
[624-31-7]

C₇H₇I




Sorbe, 1968, 66

It explodes above 200°C.
See other HALOARYL COMPOUNDS, IODINE COMPOUNDS

2751. 2-Toluenediazonium iodide
[54514-15-7]


C₇H₇IN₂



See 3-Toluenediazonium salts
See also DIAZONIUM TRIIODIDES
See other DIAZONIUM SALTS

2752. 3-Toluenediazonium iodide
[54514-15-7]


C₇H₇IN₂



See 3-Toluenediazonium salts
See also DIAZONIUM TRIIODIDES
See other DIAZONIUM SALTS

2753. 4-Iodosyltoluene
[69180-59-2]


C₇H₇IO



Sorbe, 1968, 66
It explodes at above 175°C
See other IODINE COMPOUNDS

2754. 4-Iodyltoluene
[16825-72-2]


C₇H₇IO₂



Sorbe, 1968, 66
It explodes at above 200°C.
See other IODINE COMPOUNDS

2755. 4-Iodylanisole
[16825-74-4]


C₇H₇IO₃



Leymann, -, *Chem. Fabrik.*, 1928, 705
A 1.5 kg quantity of pure material, prepared as a disinfectant, exploded with great violence when a door was opened, possibly during drying operations. It explodes at 225°C.
See other IODINE COMPOUNDS

2756. 4-Toluenediazonium triiodide
[68596-94-1]


C₇H₇I₃N₂



See entry DIAZONIUM TRIIODIDES

2757. 2-Methoxybenzenediazonium triiodide
[68596-92-9]


C₇H₇I₃NO



See entry DIAZONIUM TRIIODIDES

2758. 4-Methoxybenzenediazonium triiodide
[68596-91-8]

C₇H₇I₃NO



See entry DIAZONIUM TRIIODIDES

2759. Vinylpyridine
[1337-81-1]



C₇H₇N

Aminium nitrites


Karakuleva, G. I. *et al.*, Russ. Pat. 396 349, 1973

Nitrite salts inhibit the sometimes explosive spontaneous polymerisation of 'N-vinylpyridine derivatives'.

See other ORGANIC BASES, POLYMERISATION INCIDENTS

2760. Benzaldehyde oxime
[932-90-1]

C₇H₇NO




Energy of decomposition (in range 160–350°C) measured as 1.37 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 108°C by adiabatic Dewar tests, with an apparent energy of activation of 96 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other OXIMES

2761. 1-Nitrocyclohepta-1,3,5-triene
[156545-33-4]

C₇H₇NO₂




Burnett, I. J. *et al.*, *J. Chem. Soc., Chem Comm.*, 1994, (10) 1187.

This compound can decompose explosively if stored at room temperature.

See other C-NITRO COMPOUNDS

2762. mixo-Nitrotoluene
[1321-12-6]

C₇H₇NO₂



1. Staudinger, H., *Z. Angew. Chem.*, 1922, **35**, 657–659
2. Harris, G. F. P. *et al.*, *Runaway Reactions*, 1981, Paper 4/W, 1–11
3. Anon., *Euro. Chem. News*, 1992, 3558(1548), 27; *ibid.*, 1993, **59**(1562), 22; *Chem. Brit.*, 1993, **29**(1), 15

4. Anon., *Chem. & Ind.*, 1994, 485

5. The Fire at Hickson and Welch Ltd., HSE Books, Sudbury UK, 1994


The combined residues (300–400 kg) from several vacuum distillations at up to 180°C/40 mbar of a mixture of the 2- and 4-isomers exploded several hours after distillation had been completed. It appeared probable that admission of air to cool the hot residue initiated autoxidative heating, possibly involving nitrogen oxide and catalysis by iron, which eventually led to explosion of the residue, which probably contained polynitro compounds [1]. Explosions have occurred during fractional distillation to separate mixed nitrotoluene isomers when excessive heating was employed, or when the materials were held at more moderate temperatures but for long periods. The factors involved were identified by a series of systematic thermal stability tests on various mixtures, some including dinitrotoluenes. It was established that decomposition would not occur on the large scale if still temperatures were limited to 190°C [2]. An explosion occurred, killing five workers, when raking out (with metal rakes) the 40 cm accumulated residues from thirty years nitrotoluene distillations. It appears to have been initiated by runaway decomposition from hot-spots developed when a steam coil, possibly at 165°C, was used to soften the still tars, which were not in contact with the thermocouple, by heating to a specified 80°C. There is a suggestion that nitroresol byproducts were involved [3,4]. A full report on this event has been published [5].

See other NITROARYL COMPOUNDS

2763. 2-Nitrotoluene

[88-72-2]

C₇H₇NO₂



HCS 1980, 690

T_{ait24} was determined as 232°C by adiabatic Dewar tests, with an apparent energy of activation of 162 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (REFERENCE 2)

Alkali

1. Anon., *Loss Prev. Bull.*, 1977, (013), 2

2. Hauptli, H., private comm., 1976

1,2-Bis(2-nitrophenyl)ethane (2,2'-dinitrobibenzyl) is prepared industrially by action of alkali on 2-nitrotoluene, and the highly exothermic reaction is controlled by the rate of addition of the nitro compound to the other cooled reaction components, to keep the temperature at 5–10°C. During one batch, failure of the agitator caused increase in temperature and at 15° the temperature monitor stopped addition of nitrotoluene and increased cooling, the temperature rose to 20° and the

building was evacuated. After the exotherm had subsided, the agitator was used very intermittently to assist cooling and mixing. At 12°C full agitation was restored and processing continued to completion [1]. Similar caution is necessary during the preparation of other analogous nitrobenzyls [2].

See other AGITATION INCIDENTS

Sodium hydroxide

Anon., *ABCM Quart. Safety Summ.*, 1945, **16**, 2


Crude 2-nitrotoluene, containing some hydrochloric and acetic acids, was charged into a vacuum still with flake sodium hydroxide to effect neutralisation prior to distillation. An explosion occurred later. Similar treatments had been used uneventfully previously, using the weaker bases lime or sodium carbonate for neutralisation. It seems likely that the explosion involved formation and violent decomposition of the sodium salt of an *aci*-nitro species, possibly of quinonoid type.

See *ACI-NITROQUINONOID COMPOUNDS, NITROAROMATIC-ALKALI HAZARDS*

See other NITROARYL COMPOUNDS

2764. 4-Nitrotoluene [99-99-0]

C₇H₇NO₂



HCS 1980, 691

1. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
2. Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455
3. Anon., *ABCM Quart. Safety Summ.*, 1938, **9**, 65; 1939, **10**, 2

Energy of decomposition (in range 300–420°C) was measured as 1.74 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 232°C by adiabatic Dewar tests, with an apparent energy of activation of 162 kJ/mol [1]. Energy of decomposition of pure 4-nitrotoluene in the range 240–460°C was determined by micro DSC as 3.52 kJ/g, peaking at 409°C [2]. The residue from large-scale vacuum distillation of 4-nitrotoluene was left to cool after turning off heat and vacuum, and after 8 h a violent explosion occurred. This was variously attributed to: presence of an excessive proportion of dinitrotoluenes in the residue, which would decompose on prolonged heating; presence of traces of alkali in the crude material, which would form unstable *aci*-nitro salts during distillation; ingress of air to the hot residue on shutting off the vacuum supply, and subsequent accelerating oxidation of the

residue. All these factors may have contributed, and several such incidents have occurred [3].

Sodium

Schmidt, J., *Ber.*, 1899, **32**, 2970

One of the products as treating 4-nitrotoluene in ether with sodium is a dark brown sodium derivative which ignites in air.

See other NITROAROMATIC-ALKALI HAZARDS

Sulfuric acid

1. Hunt, J. K., *Chem. Eng. News*, 1949, **27**, 2504

2. *Chem. Abs.*, 1949, **43**, 8681a

3. McKeand, G., private comm., 1974

4. McKeand, G., *Chem. & Ind.*, 1974, 425

Solutions of 4-nitrotoluene in 93% sulfuric acid decompose very violently if heated to 160°C. This happened on plant-scale when automatic temperature control failed [1], but the temperature was erroneously abstracted as 135°C [2]. The explosion temperature of 160°C for the mixture (presumably containing a high proportion of 4-nitrotoluene-2-sulfonic acid) is 22°C lower than that observed for onset of decomposition when 4-nitrotoluene and 93% sulfuric acid are heated at a rate of 100°C/h [3]. Mixtures of 4-nitrotoluene with 98% acid or 20% oleum begin to decompose at 180 and 190°C, respectively [3,4]. Thereafter, decomposition accelerates (190–224° in 14 min, 224–270 in 1.5 min) until eruption occurs with evolution of much gas [4].

See also 3-NITROBENZENESULFONIC ACID: SULFURIC ACID

Sulfuric acid, Sulfur trioxide

Vervalin, H. C., *Hydrocarbon Process.*, 1976, **55**(9), 323

During sulfonation of 4-nitrotoluene at 32°C with 24% oleum in a 2000 l vessel, a runaway decomposition reaction set in and ejected the contents as a carbonaceous mass. The thermal decomposition temperature was subsequently estimated as 52°C (but see above).

See other SULFONATION INCIDENTS

Tetranitromethane


See Tetranitromethane: Aromatic nitro compounds

See other NITROARYL COMPOUNDS

2765. Benzyl nitrate

[15285-42-4]

C₇N₇NO₃



1. Nef, J. U., *Ann.*, 1899, **309**, 172
2. Verbal comment on presentation by Verhoeff, J., *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 3, C5-13, Rugby, IChE, 1983


It decomposed explosively at 180–200°C [1], but there is a pressure band between ambient and 64 bar in which thermal explosion does not occur [2].

Lewis acids

See entry ALKYL NITRATES *See other* BENZYL COMPOUNDS

2766. 3-Methyl-4-nitrophenol
[2581-34-2]

C₇H₇NO₃



Dartnell, R. C. *et al.*, *Loss Prev.*, 1971, **5**, 53–56; *MCA Case History No. 1649*
A batch of 8 t of material accumulated in storage at 154°C during 72 h decomposed explosively. Stability tests showed that thermal instability developed when 3-methyl-4-nitrophenol is stored molten at temperatures above 140°C. Decomposition set in after 14 h at 185° or 45 h at 165°, with peak temperatures of 593 and 521°C, respectively. In a closed vessel, a peak pressure of 750 bar was attained, with a maximum rate of increase of 40 kbar/s. Thermal degradation involves an initially slow exothermic free radical polymerisation process, followed by a rapid and violently exothermic decomposition at take-off.


See THERMAL STABILITY OF REACTION MIXTURES AND SYSTEMS

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

See other NITROARYL COMPOUNDS

2767. 4-Methyl-2-nitrophenol
[119-33-5]

C₇H₇NO₃



Sodium hydroxide, Sodium carbonate, Methanol


MCA Case History No. 701

Failure to agitate a large-scale mixture of the reagents caused an eruption owing to action of the exotherm when mixing did occur.

See entry NITROAROMATIC-ALKALI HAZARDS *See other* AGITATION INCIDENTS

2768. 2-Nitroanisole
[91-23-6]

C₇H₇NO₃



Hydrogen

1. Carswell, T. S., *J. Amer. Chem. Soc.*, 1931, **53**, 2417

2. Adkins, H., *J. Amer. Chem. Soc.*, 1931, **53**, 2808

Catalytic hydrogenation on 400 g scale at 34 bar under excessively vigorous conditions (250°C, 12% catalyst, no solvent) caused the thin autoclave without bursting disk to rupture [1]. Under more appropriate conditions the hydrogenation is safe [2].

See other CATALYTIC NITRO REDUCTION PROCESSES

Sodium hydroxide, Zinc


Anon., *Angew. Chem. (Nachr.)*, 1955, **3**, 186

In the preparation of 2,2'-dimethoxyazoxybenzene, solvent ethanol was distilled out of the mixture of 2-nitroanisole, zinc and sodium hydroxide before reaction was complete. The exothermic reaction continued unmoderated, and finally exploded.

See other NITROAROMATIC-ALKALI HAZARDS *See other* NITROARYL COMPOUNDS

2769. 2-Nitrobenzyl alcohol
[612-25-9]

C₇H₇NO₃




See entry NITROBENZYL COMPOUNDS (reference 3)

See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2770. 3-Nitrobenzyl alcohol
[619-25-0]

C₇H₇NO₃




See entry NITROBENZYL COMPOUNDS (reference 3)

See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2771. 4-Nitrobenzyl alcohol
[619-37-8]


C₇H₇NO₃



See entry NITROBENZYL COMPOUNDS (reference 3)
See other BENZYL COMPOUNDS, NITROARYL COMPOUNDS

2772. 2-Methyl-5-nitrobenzenesulfonic acid
[121-03-9]

C₇H₇NO₅S




Preparative hazard

See 4-Nitrotoluene: Sulfuric acid, or: Sulfuric acid, Sulfur trioxide
See other NITROARYL COMPOUNDS, ORGANIC ACIDS


2773. 4-Methyl-3-nitrobenzenesulfonic acid
[97-06-3]

C₇H₇NO₅S



Hakl, J., *Runaway Reactions*, 1981, paper 3/L, 5
Initial exothermic decomposition sets in at 170°C.
See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2774. 3-Toluenediazonium salts
[14604-29-6] (ion)



Ammonium sulfide, or Hydrogen sulfide

See entry DIAZONIUM SULFIDES AND DERIVATIVES

Potassium *O*-ethyl dithiocarbonate

1. Tarbell, D. S. *et al.*, *Org. Synth.*, 1955, Coll. Vol. 3, 810

2. Parham, W. E. *et al.*, *Org. Synth.*, 1967, **47**, 107

During interaction of the diazonium sulfide and the *O*-ethyl dithiocarbonate ('xanthate') solutions, care must be taken to ensure that the intermediate diazonium dithiocarbonate decomposes to 2-thiocresol as fast as it is formed [1]. This can be assured by presence of a trace of nickel in the solution to effect immediate catalytic decomposition [2]. When the 2 solutions were mixed cold and then heated to effect decomposition, a violent explosion occurred.

See other DIAZONIUM SULFIDES AND DERIVATIVES

Potassium iodide


Trumbull, E. R., *J. Chem. Educ.*, 1971, **48**, 640

Addition of potassium iodide solution to diazotised 3-toluidine was accompanied on 3 occasions by an eruption of the beaker contents. This was not observed with the 2- and 4-isomers or other aniline derivatives.

See DIAZONIUM TRIIODIDES

See other DIAZONIUM SALTS

2775. 2-Toluenediazonium salts
[17333-74-3] (ion)




Ammonium sulfide, or Hydrogen sulfide

See entry DIAZONIUM SULFIDES AND DERIVATIVES

See other DIAZONIUM SALTS


2776. 4-Toluenediazonium salts
[14597-45-6] (ion)



Ammonium sulfide, or Hydrogen sulfide

See entry DIAZONIUM SULFIDES AND DERIVATIVES *See other* DIAZONIUM SALTS

2777. 7-Azidobicyclo[2.2.1]hepta-2,5-diene
[104172-76-1]




Hoffmann, R. W. *et al.*, *Chem. Ber.*, 1986, **119**, 3309

It is explosive, and when distilled at 41°C/1.3 mbar, the bath temperature must not exceed 100°C.

See other DIENES, ORGANIC AZIDES

2778. 5-Nitrosalicylhydrazide
[946-32-7]




Khadikar, P. V. *et al.*, *Thermochim Acta*, 1986, (99), 109–118

Several heavy metal complexes of the hydrazide ($\text{ML}_2 \cdot 2\text{H}_2\text{O}$) exploded around 250°C during thermal analysis.

See related HEAVY METAL DERIVATIVES *See other* NITROARYL COMPOUNDS

2779. Benzyl azide
[622-79-7]

C₇N₇N₃




Sorbe, 1968, 138
A particularly heat-sensitive explosive oil.

Bis(trifluoromethyl) nitroxide
See Bis(trifluoromethyl) nitroxide: Benzoyl azide
See other BENZYL COMPOUNDS, ORGANIC AZIDES

2780. 4-Toluenesulfinyl azide
[40560-76-7]


C₇H₇N₃OS



See entry SULFINYL AZIDES *See other* ACYL AZIDES

2781. 4-Toluenesulfonyl azide
[941-55-9]

C₇H₇N₃O₂S



1. Rewicki, D. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1972, **11**, 44

2. Roush, W. R. *et al.*, *Tetrahedron. Lett.*, 1974, 1391–1392


3. Spencer, H., *Chem. Brit.*, 1981, **17**, 106

The impure compound, present as a majority in the distillation residue from preparation of 1-diazoindene, will explode if the bath temperature exceeds 120°C [1]. A polymer-bound sulfonyl azide reagent has been described as safer in use than the title compound [2]. Distillation of the azide at 130–132°C/~0.5 mbar from an oil bath at 145°C led to darkening, then violent explosion of the 30–40% residue [3].

See other ACYL AZIDES

2782. Methyl 2-nitrobenzenediazoate
[62375-91-1]

C₇H₇N₃O₃




Bamberger, E., *Ber.*, 1895, **28**, 237

It explodes violently on heating, or on disturbing after 24 h confinement in a sealed tube at ambient temperature. The 4-nitro analogue is more stable.

See other ARENEDIAZOATES

2783. 3-Amino-2,5-dinitrotoluene
[65321-68-8]

C₇H₇N₃O₄




Preparative hazard

See 2,5-Dinitro-3-methylbenzoic acid: Oleum, Sodium azide

See other POLYNITROARYL COMPOUNDS

2784. 2-Methanesulfonyl-4-nitrobenzenediazonium hydrogen sulfate
[68597-08-0]


C₇H₇N₃O₈S₂



See entry DIAZONIUM SULFATES

2785. 4-Methylaminobenzene-1,3-bis(sulfonyl azide)
[87425-02-3]

C₇H₇N₇O₄S₂




Lenox, R. S., US Pat. 4 393 006, 1983

It detonates on melting.

See other ACYL AZIDES

2786. Tris(2,2,2-trinitroethyl) orthoformate
[14548-59-5]


C₇H₇N₉O₂₁



See entry TRINITROETHYL ORTHOESTERS
See other POLYNITROALKYL COMPOUNDS

2787. Benzylsodium
[1121-53-5]

C₇H₇Na




Leleu, *Cahiers*, 1977, (88), 370

It ignites in air.

See related ALKYLMETALS *See other* BENZYL COMPOUNDS, ORGANOMETALLICS

2788. Sodium 4-methylphenoxide (Sodium 4-cresolate)
[22113-51-5] (ion)

C₇H₇NaO




May & Baker Ltd, private comm., 1968

Sodium 4-methylphenoxide solution was dehydrated azeotropically with chlorobenzene, and the filtered solid was dried in an oven, where it soon ignited and glowed locally. This continued for 30 min after it was removed from the oven. A substituted potassium phenoxide, prepared differently, also ignited on heating. Finely divided and moist alkali phenoxides may be prone to vigorous oxidation (or perhaps reaction with carbon dioxide) when heated in air.

†2789. **Bicyclo[2.2.1]hepta-2,5-diene**
[121-46-0]

C₇H₈




HCS 1980, 209

See related DIENES

†2790. **1,3,5-Cycloheptatriene**
[544-25-2]

C₇H₈




HCS 1980, 341

See related DIENES

†2791. **Toluene (Methylbenzene)**
[108-88-3]

C₇H₈



(*MCA SD-63*, 1956); *NSC 204*, 1968; *FPA H11*, 1973; *HCS 1980*, 906;
RSC Lab. Hazard Data Sheet No. 15, 1983

Other reactants

Yoshida, 1980, 261

MRH Values calculated for 15 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Bromine trifluoride: Solvents

MRH 2.97/80

1,3-Dichloro-5,5-dimethyl-2,4-imidazolidindione: Xylene

Dinitrogen tetroxide: Hydrocarbons

MRH 7.15/81

Nitric acid: Hydrocarbons

MRH 6.07/82

Tetranitromethane: Hydrocarbons

Uranium hexafluoride: Aromatic hydrocarbons


Sulfur dichloride

See Sulfur dichloride: Toluene

2792. Ethyl 4-bromo-1,1,3-trioxoisothiazoleacetate

[126623-64-1]

C₇H₈BrNO₅S



Ethyl diazoacetate

Burri, K. F., *Helv. Chim. Acta*, 1989, **72**(6), 1416

It is suggested that both adducts of these compounds are explosive.

See Ethyl 6-ethoxycarbonyl-3,4-dihydro-1,1,3-trioxo-2-pyrazolo [3,4-d]isothiazole-2-acetate


See Ethyl diazoacetate

See other N-S COMPOUNDS

2793. 2-Chloro-5-methylaniline

[95-81-8]

C₇H₈ClN




Preparative hazard

See 2-Chloro-5-methylphenylhydroxylamine

See other HALOANILINES

2794. 4-Chloro-2-methylaniline
[95-69-2]

C₇H₈ClN



See 4-Chloro-2-methylbenzenediazonium salts

Copper(II) chloride


Kotoyori, T. *et al.*, *J. Haz. Mat.*, 1976, 252–262

At the end of vacuum distillation of 4-chloro-2-methylaniline, the 1300 l still burst owing to internal gas pressure from an unexpected decomposition reaction. Subsequent investigation showed that copper(I) oxide present in the still charge would react with hydrogen chloride and air leaking in to produce copper(II) chloride. The latter is an effective catalyst for decomposition of the haloaniline at 239°C to produce hydrogen chloride. The physical conditions of the installation were compatible with this mechanism, and enough chloroaniline was present to exceed the bursting pressure of the vessel by a factor of 3. It is concluded that halogenated aromatic amines should be distilled under alkaline conditions in absence of air.

See CATALYTIC IMPURITY INCIDENTS See *other* HALOANILINES

2795. 3-Chloro-4-methoxyaniline
[5345-54-0]

C₇H₈ClNO




Heat of decomposition was determined as 0.82 kJ/g by DSC, and T_{ait24} was determined as 158°C by adiabatic Dewar tests, with an apparent energy of activation of 135 kJ/mol.

See *entry* THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See *other* AMINOMETHOXY COMPOUNDS, HALOANILINES

2796. 2-Chloro-5-methylphenylhydroxylamine
[65039-20-5]

C₇H₈ClNO




Rondevstedt, C. S., *Chem. Eng. News*, 1977, **55**(27), 38; *Synthesis*, 1977, 851–852
In an attempt to reduce 2-chloro-5-methylnitrobenzene to the aniline by treatment with excess hydrazine and Pd/C catalyst, the hydroxylamine was unexpectedly produced as major product. During isolation of the product, after removal of solvent it was heated to 120°C under vacuum and exploded fairly violently. Many arylhydroxylamines decompose violently when heated above 90–100°C, especially in presence of acids. GLC is not suitable as an analytical diagnostic for arylhydroxylamines because of this thermal decomposition.

See 3,4-Dichlorophenylhydroxylamine

See other CATALYTIC NITRO REDUCTION PROCESSES, N–O COMPOUNDS

2797. N-Chlorotoluene-4-sulfonamide
[144-86-5]

C₇H₈ClNO₂S




Energy of exothermic decomposition in range 110–330°C was measured as 1.36 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

2798. Benzohydrazide
[613-94-5]

C₇H₈N₂O




Benzeneseleninic acid

See Benzeneseleninic acid: Hydrazine derivatives

2799. Methyl benzenediazoate (1-Phenyl-2-methoxydiazene)
[66217-76-3]

C₇H₈N₂O




Bamberger, E., *Ber.*, 1895, **28**, 228

It explodes on heating or after storage for 1 h in a sealed tube at ambient temperature.

See other ARENEDIAZOATES

2800. N-Methyl-4-nitroaniline
[100-15-2]

C₇H₈N₂O₂




2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide ('Carbyl sulfate')

See 2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide: *N*-Methyl-4-nitroaniline

See other NITROARYL COMPOUNDS

2801. 2-Methyl-5-nitroaniline
[99-55-8]

C₇H₈N₂O₂




Hakl, J., *Runaway Reactions*, 1981, Paper 3/L, 5

Initial exothermic decomposition sets in at 150°C.

See other NITROARYL COMPOUNDS

2802. 2-Methoxy-5-nitroaniline
[99-59-2]

C₇H₈N₂O₃




Preparative hazard

See 2-Methoxyanilinium nitrate: Sulfuric acid

See other AMINOMETHOXY COMPOUNDS, NITROARYL COMPOUNDS

2803. Ammonium 3-methyl-2,4,6-trinitrophenoxide
[58696-86-9]

C₇H₈N₄O₇




Biasutti, 1981, 63

An old store of the explosive material exploded violently with no apparent cause. A similar incident had been recorded previously. Some isomeric *aci*-nitroquinonoid salt may have been formed.

See *aci*-NITROQUINONOID COMPOUNDS See other POLYNITROARYL COMPOUNDS

2804. 3-Phenyl-1-(5-tetrazolyl)-1-tetrazene
[]

C₇H₈N₈




Thiele, J., *Ann.*, 1892, **270**, 60

It explodes on warming, as do C-substituted homologues.
See other HIGH-NITROGEN COMPOUNDS, TETRAZENES, TETRAZOLES

2805. Methylenebis(3-nitramino-4-methylfuzazan)

[]

C₇H₈N₈O₆




Willer, R. L. *et al.*, *J. Heterocycl. Chem.*, 1992, **29**(7), 1835

An explosive of moderate sensitivity, thermally decomposing (non-explosively) from 120°C. The 4-nitro analogue was also explosive and decomposed from 90°C.
See other N-NITRO COMPOUNDS, HIGH NITROGEN COMPOUNDS

2806. Benzyl alcohol

[100-51-6]

C₇H₈O



HCS 1980, 198

Other reactants

Yoshida, 1980, 329

MRH values calculated for 13 combinations with oxidants are given.

Sulfuric acid

MRH 1.13/48

See Sulfuric acid: Benzyl alcohol

Hydrogen bromide, Iron

Iselin, E. *et al.*, *Chem. Ing. Tech.*, 1972, **44**, 462


Benzyl alcohol contaminated with 1.4% of hydrogen bromide and 1.1% of dissolved iron(II) polymerises exothermally above 100°C. Bases inhibit the polymerisation reaction. In a laboratory test, alcohol containing 1% of HBr and 0.04% of Fe polymerised at about 150° with an exotherm to 240°C. Formation and iron-catalysed poly-condensation of benzyl bromide seems to have been implicated.

See Benzyl bromide: Molecular sieve, or: Catalytic impurities

See other BENZYL COMPOUNDS, POLYCONDENSATION REACTION INCIDENTS

2807. 2-Acetyl-3-methyl-4,5-dihydrothiophen-4-one
[]

C₇H₈O₂S




MCA Guide, 1972, 307

An explosion occurred during vacuum distillation. The highly conjugated compound could have been susceptible to peroxidation as well as capable of condensative polymerisation.

See other PEROXIDATION INCIDENTS, POLYCONDENSATION REACTION INCIDENTS

2808. 4-Toluenesulfonic acid
[104-15-4]

C₇H₈O₃S



HCS 1980, 908


Acetic anhydride, Water

See Acetic anhydride: 4-Toluenesulfonic acid, etc.

See other ORGANIC ACIDS

2809. 3-Thiocresol
[108-40-7]

C₇H₈S




Preparative hazard

See 3-Toluenediazonium salts: Potassium *O*-ethyl dithiocarbonate

2810. Chloromethylphenylsilane
[1631-82-9]

C₇H₉ClSi



4-Bromobutene, Chloroplatinic acid


Brook, A. G. *et al.*, *J. Organomet. Chem.*, 1975, **87**, 265

Interaction of chloromethylphenylsilane and 4-bromobutene, catalysed by chloroplatinic acid at 100°C, is extremely exothermic (to 165°C) and is accompanied by vigorous frothing.

See other ALKYLHALOSILANES

2811. 2-Deuterobicyclo[2.2.1]hept-2-ene
[694-94-0]

C₇H₉D



Preparative hazard


Shackelford, S. A., *Tetrahedron Lett.*, 1977, 4268, note 12

The metallation reaction involved in the preparation proved hazardous and unpredictable.

See related ALKENES

2812. Benzylamine
[100-46-9]

C₇H₉N




N-Chlorosuccinimide

See N-HALOIMIDES: alcohols, etc. *See other* BENZYL COMPOUNDS, ORGANIC BASES

2813. 2-Toluidine
[95-53-4]

C₇H₉N



HCS 1980, 910

See 2- or 4-Toluenediazonium salts

Nitric acid

MRH 6.11/81

See Nitric acid: Aromatic amines


Other reactants

Yoshida, 1980, 260

MRH values calculated for 13 combinations with oxidants are given.

2814. 3-Toluidine
[108-44-1]


C₇H₉N



See 3-Toluenediazonium salts

2815. 4-Toluidine
[106-49-0]


C₇H₉N



See 2- or 4-Toluenediazonium salts

2816. 2-Methoxyaniline (*o*-Anisidine)
[90-04-0]


C₇H₉NO



Energy of decomposition (in range 190–480°C) measured as 0.387 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS

2817. 3-Methoxyaniline (*m*-Anisidine)
[536-90-3]


C₇H₉NO



Energy of decomposition (in range 410–500°C) measured as 0.518 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS

2818. 4-Methoxyaniline (*p*-Anisidine)
[100-94-9]


C₇H₉NO



Energy of decomposition (in range 340–490°C) measured as 0.429 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS

2819. 2,6-Dimethylpyridine N-oxide
[1073-23-0]

C₇H₉NO




Phosphoryl chloride

See Phosphoryl chloride: 2,6-Dimethylpyridine N-oxide

See other N-OXIDES

2820. 4(1-Hydroxyethyl)pyridine N-oxide
[34277-48-0]

C₇H₉NO₂




Taylor, R., *J. Chem. Soc., Perkin Trans. 2*, 1975, 279

The compound exploded during vacuum distillation, possibly owing to the presence of traces of peracetic acid.

See other N-OXIDES

2821. O-4-Toluenesulfonylhydroxylamine
[52913-14-1]

C₇H₉NO₃S



Carpino, L. A., *J. Amer. Chem. Soc.*, 1960, **82**, 3134


It deflagrated spontaneously on attempted isolation and drying at ambient temperature.

See O-Mesitylenesulfonylhydroxylamine

See other N-O COMPOUNDS

2822. 4-Chloro-2,6-diamino-*N*-methylaniline
[48114-68-7]

C₇H₁₀ClN₃




T_{ait24} determined as 148°C in adiabatic Dewar tests with apparent energy of activation of 123 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HALOANILINES

2823. *N*-(Methylphenylphosphinoyl)hydroxylamine
[94193-96-3]

C₇H₁₀NO₂P




Harger, M. J. P., *J. Chem. Soc., Perkin Trans. 1*, 1987, 685

A sample decomposed vigorously after several hours at warm ambient temperature. The ethyl and isopropyl homologues appeared stable, but all (including the derived *O*-sulfonic acids) were treated as potentially unstable and were only heated briefly during crystallisation, and were stored at -20°C.

See related HYDROXYLAMINIUM SALTS See other N-O COMPOUNDS

2824. Dicyclopropyldiazomethane
[16102-24-2]

C₇H₁₀N₂




Ensslin, H. M *et al.*, *Angew. Chem. (Intern. Ed.)*, 1967, **6**, 702-703

Storable in ether at -50°C , but violent decomposition of undiluted material sets in at -15°C .

See other DIAZO COMPOUNDS, STRAINED-RING COMPOUNDS

2825. 1,5-Diisocyanopentane
[97850-58-3]


$\text{C}_7\text{H}_{10}\text{N}_2$



See entry DIISOCYANIDE LIGANDS See related CYANO COMPOUNDS

2826. 4-Methoxy-1,3-phenylenediamine
[615-05-4]

$\text{C}_7\text{H}_{10}\text{N}_2\text{O}$




Energy of decomposition (in range $260-420^{\circ}\text{C}$) measured as 0.42 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 207°C by adiabatic Dewar tests, with an apparent energy of activation of 114 kJ/mol .

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other AMINOMETHOXY COMPOUNDS

2827. 4-Toluenesulfonylhydrazide
[877-66-7]

$\text{C}_7\text{H}_{10}\text{N}_2\text{O}_2\text{S}$




1. Hansell, D. P. *et al.*, *Chem. & Ind.*, 1975, 464

While drying in a tray dryer at 60°C, a batch decomposed fairly exothermically but without fire. As a blowing agent, it is designed to have limited thermal stability [1]. One of the recently calculated values of 87° and 78°C for induction periods of 7 and 60 days, respectively, for critical ignition temperature coincides with the previous value of 87°/7 days. Autocatalytic combustion is exhibited.

See other BLOWING AGENTS

2828. Diethyl diazomalonate
[5256-74-6]

C₇H₁₀N₂O₄




DOC 5, 1982, Vol. 2, 1588

It may explode during distillation at low pressure.

See other DIAZO COMPOUNDS

2829. 2-Methoxyanilinium nitrate
[]

C₇H₁₀N₂O₄



1. Vervalin, C. H., *Hydrocarb. Process.*, 1976, **55**(9), 321
2. *See entry* HIGH RATE DECOMPOSITION

Use of a screw feeder to charge a reactor with the 2-anisidine salt led to ignition of the latter. It was known that the salt would decompose exothermically above 140°C, but later investigation showed that lower-quality material could develop an exotherm above 46°C under certain conditions [1]. Fast flame propagation occurs on moderately heating anisidine nitrate powder [2].

Sulfuric acid

Vervalin, C. H., *Hydrocarb. Process.*, 1976, **55**(9), 321

In a process for preparation of 2-methoxy-5-nitroaniline the anisidine salt was added to stirred sulfuric acid. An accidental deficiency of the latter prevented


proper mixing and dissipation of the heat of solution, and local decomposition spread through the entire content of the 2000 l vessel, attaining red heat.

See other OXOSALTS OF NITROGENOUS BASES, SULFONATION INCIDENTS

See related AMINOMETHOXY COMPOUNDS

2830. Methyl 3-methoxycarbonylazocrotonate
[63160-33-8]

C₇H₁₀N₂O₄




Sommer, S., *Tetrahedron Lett.*, 1977, 117

Attempted high-vacuum fractional distillation of an isomeric mixture (10% *cis*-) led to explosive decomposition. The 'diene' polymerises if stored at 20°C.

See other AZO COMPOUNDS

2831. 5-*tert*-Butyl-3-diazo-3*H*-pyrazole
[62072-11-1]

C₇H₁₀N₄




The light-sensitive and piezosonic crystalline solid explodes at 97°C.

See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

2832. Benzylsilane
[766-06-3]

C₇H₁₀Si



491M, 1975, 58


It ignites in air.

See related ALKYL SILANES See other BENZYL COMPOUNDS

2833. Cyclopentadienyldimethylborane

[]

C₇H₁₁B



Herberich, G. E. *et al.*, *Organometallics*, 1996, **15**(1), 58


Spontaneously inflames in air.

See other ALKYL BORANES

2834. Lithium 1-heptynide

[42017-07-2]

C₇H₁₁Li



Ammonia, Iodine


See Iodine: Ammonia, Lithium 1-heptynide

See other METAL ACETYLIDES

2835. 3-Methyl-4-nitro-1-buten-3-yl acetate

[61447-07-2]

C₇H₁₁NO₄



Wehrli, P. A. *et al.*, *J. Org. Chem.*, 1977, **42**, 2940


This and the isomeric nitroester (below), both produced in the nitroacetoxylation of isoprene, are of limited thermal stability and it is recommended that neither be

heated above 100°C, either neat or in solution. Vacuum distillation of the nitro esters should be limited to 1 g portions, as decomposition 'fume-offs' have been observed.

See related NITROALKENES

2836. 3-Methyl-4-nitro-2-buten-1-yl acetate
[1447-06-1]

C₇H₁₁NO₄




See 3-Methyl-4-nitro-1-buten-3-yl acetate

See related NITROALKENES

†2837. Cycloheptene
[628-92-2]

C₇H₁₂



See related ALKENES

†2838. 1-Heptyne
[628-71-7]


C₇H₁₂



See other ALKYNES

†2839. 4-Methylcyclohexene
[591-47-9]


C₇H₁₂



See related ALKENES

2840. 1-Methoxy-3,4,5-trimethylpyrazole *N*-oxide
[39753-42-9]

C₇H₁₂N₂O₂



Boyle, F. T. *et al.*, *J. Chem. Soc., Perkin Trans. I*, 1973, 167
It is thermally unstable and exploded during attempted distillation.
See 1-Methoxyimidazole *N*-oxide
See other N-O COMPOUNDS, *N*-OXIDES

2841. 2-Heptyn-1-ol
[1002-36-4]


C₇H₁₂O



Brandsma, 1971, 69
Dilution of the alcohol with white mineral oil before vacuum distillation is recommended to avoid the possibility of explosion of the undiluted distillation residue.
See other ACETYLENIC COMPOUNDS

2842. 3-Methylcyclohexanone
[591-24-2]


C₇H₁₂O



Hydrogen peroxide, Nitric acid
See Hydrogen peroxide: Ketones, etc.

2843. 4-Methylcyclohexanone
[589-92-4]

C₇H₁₂O




Nitric acid

See Nitric acid: 4-Methylcyclohexanone

2844. 4-Ethoxy-2-methyl-3-butyn-2-ol
[2041-76-1]

C₇H₁₂O₂



Brandsma, 1971, 78


Traces of acid adhering to glassware are sufficient to induce explosive decomposition of the alcohol during distillation, and must be neutralised by pre-treatment with ammonia gas. Low pressures and temperatures are essential during distillation. Explosions during distillation using water pump vacuum and bath temperatures above 115°C were frequent.

See ETHOXYETHYNYL ALCOHOLS

See other ACETYLENIC COMPOUNDS, CATALYTIC IMPURITY INCIDENTS

2845. 1,2-Dimethylcyclopentene ozonide (1,5-Dimethyl-6,7,8-trioxabicyclo[3.2.1]octane)
[19987-14-5]

C₇H₁₂O₃



Criegee, R. *et al.*, *Ber.*, 1953, **86**, 3

Though the ozonide appeared stable to small-scale high-vacuum distillation, the residue exploded violently at 130°C.

See other OZONIDES

2846. N-Cyano-2-bromoethylbutylamine


[]

C₇H₁₃BrN₂Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 23


A small sample heated to 160°C decomposed exothermically, reaching 250°C. It had previously been distilled at 95°/0.65 mbar without decomposition.

*See N-Cyano-2-bromoethylcyclohexylamine**See other* CYANO COMPOUNDS**2847. 3-Trimethylsilylprop-2-enyl trifluoromethanesulfonate**

[62905-84-4]

C₇H₁₃F₃O₃SSi*See entry* ALLYL TRIFLUOROMETHANESULFONATES *See other* ALLYL COMPOUNDS, SULFUR ESTERS**2848. N-Dimethylethyl-3,3-dinitroazetidine (N-tert-Butyl-3,3-dinitroazetidine)**


[125735-38-8]

C₇H₁₃N₃O₄Archibald, T. G. *et al.*, *J. Org. Chem.*, 1990, **55**, 2920

The above compound, prepared as an intermediate to 1,3,3-trinitroazetidine for study as an explosive, decomposed explosively when heated above 120°C for distillation. Precautions when handling nitroazetidines are counselled.


See related POLYNITROALKYL COMPOUNDS *See other* STRAINED RING COMPOUNDS**†2849. Cycloheptane**

[291-64-5]

C₇H₁₄

†2850. Ethylcyclopentane
[1640-89-7]

C₇H₁₄



†2851. 1-Heptene
[592-76-7]


C₇H₁₄



See other ALKENES

†2852. 2-Heptene
[592-77-8]


C₇H₁₄



See other ALKENES

†2853. 3-Heptene
[592-78-9]


C₇H₁₄



See other ALKENES

†2854. Methylcyclohexane
[108-87-2]

C₇H₁₄



HCS 1980, 646

†2855. 2,3,3-Trimethylbutene
[594-56-9]

C₇H₁₄




See other ALKENES

2856. Dimethylmethylenexosulfanenickel(0) diethylene complex

[]

C₇H₁₄NiOS



Pörschke, K. R., *Chem. Ber.*, 1987, **120**, 425–427

The complex decomposes violently at 0°C, evolving ethane, cyclopropane and methane.

See other GAS EVOLUTION INCIDENTS, HEAVY METAL DERIVATIVES

†2857. 2,4-Dimethyl-3-pentanone

[565-80-0]

C₇H₁₄O



†2858. 3,3-Diethoxypropene

[3054-95-3]

C₇H₁₄O₂



See other PEROXIDISABLE COMPOUNDS

†2859. Ethyl isovalerate

[108-64-5]

C₇H₁₄O₂



†2860. Isobutyl propionate

[540-42-1]

C₇H₁₄O₂



†2861. Isopentyl acetate

[123-92-2]

C₇H₁₄O₂



NSC 208, 1977; *FPA H117*, 1982; *HSC 1980*, 563

†2862. Isopropyl butyrate

[638-11-9]

C₇H₁₄O₂



†2863. Isopropyl isobutyrate
[617-50-5]



$\text{C}_7\text{H}_{14}\text{O}_2$

†2864. Pentyl acetate
[628-63-7]



$\text{C}_7\text{H}_{14}\text{O}_2$

HCS 1980, 153

†2865. 2-Pentyl acetate
[53496-15-4]




$\text{C}_7\text{H}_{14}\text{O}_2$

HCS 1980, 154

2866. 4(2,3-Epoxypropoxy)butanol
[4711-95-9]

$\text{C}_7\text{H}_{14}\text{O}_3$




Trichloroethylene

See Trichloroethylene: Epoxides

See other 1,2-EPOXIDES

2867. Cyclopentylmethyl methanesulfonate
[73017-74-2]




$\text{C}_7\text{H}_{14}\text{O}_3\text{S}$

It decomposes vigorously at 128°C.

See entry SULFONIC ACID ESTERS *See other* SULFUR ESTERS

†2868. 2,6-Dimethylpiperidine
[504-03-0]


$\text{C}_7\text{H}_{15}\text{N}$



See other ORGANIC BASES

†2869. 1-Ethylpiperidine
[766-09-6]


$C_7H_{15}N$



See other ORGANIC BASES

†2870. 2-Ethylpiperidine
[1484-80-6]

$C_7H_{15}N$



See other ORGANIC BASES

†2871. 2,3-Dimethylpentane
[565-59-3]

C_7H_{16}



†2872. 2,4-Dimethylpentane
[108-08-7]

C_7H_{16}



†2873. Heptane
[142-82-5]

C_7H_{16}



HCS 1980, 524 (includes isomers)

†2874. 2-Methylhexane
[591-76-4]

C_7H_{16}



†2875. 3-Methylhexane
[589-34-4]

C_7H_{16}




†2876. 2,2,3-Trimethylbutane
[464-06-2]



C_7H_{16}

2877. 3-Diethylaminopropylamine
[14642-66-1]

$\text{C}_7\text{H}_{18}\text{N}_2$



Cellulose nitrate

See CELLULOSE NITRATE: amines See other ORGANIC BASES

2878. *N*-*tert*-Butyl-*N*-trimethylsilylaminoborane
[73452-32-1]



$\text{C}_7\text{H}_{20}\text{BNSi}$


Wisian-Nielsen, P. *et al.*, *J. Inorg. Nucl. Chem.*, 1979, **41**, 1546

It ignites in direct contact with air.

See related BORANES, ALKYL SILANES

2879. Tetrakis(chloroethynyl)silane
[]

$\text{C}_8\text{Cl}_4\text{Si}$



Viehe, H. G., *Chem. Ber.*, 1959, **92**, 3075


Though thermally stable, it exploded violently on grinding (with potassium bromide) or on impact.

See other FRICTIONAL INITIATION INCIDENTS, HALOACETYLENE DERIVATIVES

2880. Potassium octacyanodicobaltate(8-)

[23705-25-1]

$C_8Co_2K_8N_8$



Bailar, 1973, Vol. 4, 175

A very unstable, pyrophoric material.

See other METAL CYANIDES (AND CYANO COMPLEXES), PYROPHORIC MATERIALS

2881. Octacarbonyldicobalt

[10210-68-1]

$C_8Co_2O_8$



Wender, I. *et al.*, *Inorg. Synth.*, 1957, **5**, 191

When the carbonyl is prepared (by rapid cooling) as a fine powder, it decomposes in air to give pyrophoric dodecacarbonyltetracobalt.

See other CARBONYLMETALS

2882. Caesium graphite

[12079-66-2]

C_8Cs




See Carbon: Alkali metals

See other METAL NON-METALLIDES

2883. 3-Bromo-2,7-dinitrobenzo[b]thiophene-5-diazonium-4-oxide

[]

$C_8HBrN_4O_5S$



Brown, I. *et al.*, *Chem. & Ind.*, 1962, 982

It is an explosive compound.


See Nitric acid: Benzo[b]thiophene derivatives

See other ARENEDIAZONIUM OXIDES

2884. N-Bromo-3-nitrophthalimide

[]

$C_8H_3BrN_2O_4$




Tetrahydrofurfuryl alcohol

See N-HALOIMIDES: alcohols See other N-HALOGEN COMPOUNDS, NITROARYL COMPOUNDS

2885. 1,3-Bis(trifluoromethyl)-5-nitrobenzene

[328-75-6]

$C_8H_3F_6NO_2$



Preparative hazard


See Nitric acid: 1,3-Bis(trifluoromethyl)benzene

See other NITROARYL COMPOUNDS

2886. 4-Nitro-1-picryl-1,2,3-triazole

[31123-28-1]

$C_8H_3N_7O_8$



Storm, C. B. *et al.*, *J. Phys. Chem.*, 1989, **93**, 1000–1007


Though the 1- and 2-picryl isomers are similarly explosive, the former has a much higher impact-sensitivity than the latter. This is attributed to the former

having a shorter N2 to N3 interatomic distance, longer adjacent bonds, and a lower activation energy for elimination of molecular nitrogen from the 2,3 positions.

See 5-nitro-2-picryltetrazole, 1-Picryl-1,2,3-triazole

See other POLY-NITROARYL COMPOUNDS, TRIAZOLES

2887. Silver isophthalate
[57664-97-8]




Fields, E. K., *J. Org. Chem.*, 1976, **41**, 918–919

Decomposition of the salt at 375°C under nitrogen, with subsequent cooling under hydrogen, gives a black carbon-like polymer containing metallic silver which ignites at 25°C on exposure to air.

See related PYROPHORIC MATERIALS See other SILVER COMPOUNDS

2888. Isophthaloyl chloride
[99-63-8]



Preparative hazard

See 1,3-Bis(trichloromethyl)benzene: Oxidants


Methanol

Morrell, S. H., private comm., 1968

Violent reaction occurred between isophthaloyl chloride and methanol when they were accidentally added in succession to the same waste solvent bottle.

See other ACYL HALIDES

2889. Terephthaloyl chloride (1,4-Benzenedicarbonyl chloride)
[100-20-9]




Water

Anon., *Loss Prev. Bull.*, 1986, (071), 9

Hydrolysis of 7 t of the acid chloride by water in a herbicide processing unit ran out of control, gas evolution eventually rupturing the reactor violently.

See other ACYL HALIDES, GAS EVOLUTION INCIDENTS

2890. 2,4-Hexadiynylene chloroformate
[]





Driedger, P. E. *et al.*, *Chem. Eng. News*, 1972, **50**(12), 51

One sample exploded with extreme violence at 15°C/0.2 mbar, while another smaller sample had been distilled uneventfully at 114–115°C/0.2 mbar.

See other ACYL HALIDES, ACETYLENIC COMPOUNDS

2891. 1,3-Bis(trichloromethyl)benzene
[881-99-2]



Oxidants


Rondevstedt, C. S., *J. Org. Chem.*, 1976, **41**, 3574–3577

Heating the bis(trichloromethyl)benzene with potassium nitrate, selenium dioxide or sodium chlorate to effect conversion to the bis(acyl chloride) led to eruptions at higher temperatures, and was too dangerous to pursue.

See related HALOALKANES, HALOARYL COMPOUNDS

2892. N-Trifluoroacetoxy-2,4-dinitroaniline
[127526-90-3]

C₈H₄F₃N₃O₆




Bryant, I. R. *et al.*, *Austr. J. Chem.*, 1989, **42**(12), 2275

The crude compound was isolated, began to evolve trifluoroacetic acid and, after 15 minutes, exploded.

See other N–O COMPOUNDS

2893. 1-Picryl-1,2,3-triazole
[18922-71-9]

C₈H₄N₆O₆



Storm, C. B. *et al.*, *J. Phys. Chem.*, 1989, **93**, 1000–1007


Though the 1- and 2-picryl isomers are similarly explosive, the former has a much higher impact-sensitivity than the latter. This is attributed to the former having a shorter N2 to N3 interatomic distance, longer adjacent bonds, and a lower activation energy for elimination of molecular nitrogen from the 2,3 positions.

See 4-Nitro-1-picryl-1,2,3-triazole

See other POLYNITROARYL COMPOUNDS, TRIAZOLES

2894. 1,3-Bis(trifluoromethyl)benzene
[402-31-3]

$C_8H_4F_6$




Nitric acid, Sulfuric acid

See Nitric acid: 1,3-Bis(trifluoromethyl)benzene, etc.

See related HALOALKANES, HALOARYL COMPOUNDS

2895. Tetraethynylgermanium
[4531-35-5]

C_8H_4Ge



1. Chokiewicz, W. *et al.*, *Compt. rend.*, 1960, **250**, 866


2. Shikkiev, I. A. *et al.*, *Doklad. Akad. Nauk SSSR*, 1961, **139**, 1138 (Engl. transl., 830)

It explodes on rapid heating [1], or friction [2].

See other METAL ACETYLIDES

2896. Potassium tetraethynynickelate(2-)
[65664-23-5]

$C_8H_4K_2Ni$



Air, or Water


Vigorously pyrophoric in air, it may explode on contact with drops of water.

See entry COMPLEX ACETYLIDES

2897. Potassium tetraethynynickelate(4-)

[]

$C_8H_4K_4Ni$




The diammoniate (and that of the sodium salt) explodes on exposure to impact, friction or flame.

See entry COMPLEX ACETYLIDES

2898. Phthaloyl diazide

[50906-29-1]

$C_8H_4N_6O_2$




Lindemann, H. *et al.*, *Ann.*, 1928, **464**, 237

The *symm*-diazide is extremely explosive, while the *asymm*-isomer (*gem*-diazidoisobenzofuranone) is less so, melting at 56°C, but exploding on rapid heating.

See other ACYL AZIDES

2899. Phthalic anhydride
[85-44-9]



HCS 1980, 752

Preparative hazard

1. Kratochvil, V., *Chem. Abs.*, 1974, **80**, 3215
2. English transl. by D. S. Rosenberg, 10.2.69 of German transl. by S. Vedrilla (Chemiebau) of 'Low Temperature Pyrophoric Compounds in Process for Preparing Phthalic Anhydride', a Japanese paper in *Anzen Kogyo Kogaku*, 1968
3. Schwab, R. F. *et al.*, *Chem. Eng. Progr.*, 1970, **66**(9), 49-53

Investigation of an explosion in a phthalic anhydride plant showed that naphthoquinones (by-products from naphthalene oxidation) reacted with iron(III) salts of phthalic, maleic or other acids (corrosion products) to form labile materials. The latter were found to undergo exothermic oxidation and self-ignition at around 200°C. Process conditions to minimise hazards are discussed, with 27 references [1]. Fires and explosions in the condenser sections of phthalic anhydride plants were traced to the presence of deposits of thermally unstable basic iron salts of maleic or phthalic acids and iron sulfides derived from sulfur in the naphthalene feedstock. A small-scale laboratory test procedure was developed to measure the exotherm point and maximum temperature rise in various mixtures of materials found to be present in the condensing section. A basic iron(III) maleate hydrate showed an exotherm point of 165°C and temperature rise of 369°C, changed by pre-treatment with hydrogen sulfide to 164 and 392°C, respectively. The corresponding figures for basic iron(III) phthalate were 236 and 477°C, both unchanged by pre-sulfiding treatment. The part played by acid-corrosion of the mild steel condensers in formation of the unstable pyrophoric compounds was also investigated. It was concluded that the origins of the many mishaps which have occurred soon after plant start-up were explicable in terms of these experimental findings [2]. Other hazards in phthalic anhydride production units have also been discussed [3].

See PYROPHORIC IRON-SULFUR COMPOUNDS See other CORROSION INCIDENTS

Other reactants

Yoshida, 1980, 357

MRH Values calculated for 11 combinations with oxidants are given.

Oxidants

See Copper(II) oxide: Phthalic acid, etc.

Nitric acid: Phthalic anhydride

Sodium nitrite: Phthalic acid, etc.

See other ACID ANHYDRIDES

MRH values show % of oxidant


MRH 2.09/tr.

MRH 5.52/71

MRH 2.09/tr.

2900. Phthaloyl peroxide (2,3-Benzodioxin-1,4-dione)
[4733-52-2]

$C_8H_4O_4$




Jones, M. *et al.*, *J. Org. Chem.*, 1971, **36**, 1536

Detonable by impact or by melting at 123°C. It may be a linear polymer, rather than the cyclic structure implied by the alternative name.

See other DIACYL PEROXIDES

2901. Tetraethynyltin
[16413-88-0]

C_8H_4Sn




Jenkner, H., Ger. Pat. 115 736, 1961

It explodes on rapid heating.

See other METAL ACETYLIDES

2902. 2,4-Dinitrophenylacetyl chloride
[109799-62-4]

$C_8H_5ClN_2O_5$



Bonner, T. G., *J. R. Inst. Chem.*, 1957, **81**, 596

Explosive decomposition occurred during attempted vacuum distillation, attributed either to the presence of some trinitro compound in the unpurified dinitrophenylacetic acid used, or to the known instability of *o*-nitro acid chlorides. A previous publication (ibid., 407) erroneously described the decomposition of 2,4-dinitrobenzoyl chloride.

See other NITROACYL HALIDES

2903. Trifluoroacetylminoiodobenzene

[94286-09-6]



$\text{C}_8\text{H}_5\text{F}_3\text{INO}$

Mansuy, D. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1984, 1163


The solid undergoes explosive decomposition at 100°C.

See other N-HALOGEN COMPOUNDS, IODINE COMPOUNDS

2904. 3-Benzocyclobutenylpotassium

[]

$\text{C}_8\text{H}_5\text{K}$



Perchloryl fluoride


See Perchloryl fluoride: Benzocyclobutene, etc.

See related ARYLMETALS

2905. 3-Nitrophenylacetylene

[3034-94-4]

$\text{C}_8\text{H}_5\text{NO}_6$




Sorbe, 1968, 152

The compound (isomer not stated) and its metal derivatives are explosive.

See other ALKYNES, NITROARYL COMPOUNDS

2906. 3-Nitrophthalic acid
[603-11-2]

C₈H₅NO₆




Preparative hazard

See Nitric acid: Phthalic anhydride, etc.

See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2907. Nitroterephthalic acid
[603-11-2]

C₈H₅NO₆




Preparative hazard

See Nitric acid: Sulfuric acid, Terephthalic acid

See other NITROARYL COMPOUNDS, ORGANIC ACIDS

2908. 3-Sodio-5-(5'-nitro-2'-furfurylideneamino)imidazolidin-2,4-dione
[54-87-5]

C₈H₅N₄NaO₅




Boros, L. et al., Ger. Offen., 2 328 927, 1974

The crystalline *N*-sodium salt is explosive.

See other *N*-METAL DERIVATIVES

2909. 3-Diazo-5-phenyl-3H-1,2,4-triazole
[80670-36-6]

$C_8H_5N_5$




It is a light-sensitive and highly shock-sensitive solid, which can be weighed or stored in the dark at ambient temperature as the dry solid, but it decomposes in solution in a few hours.

See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

2910. 4-Diazo-5-phenyl-1,2,3-triazole
[64781-77-7]

$C_8H_5N_5$




The crystalline solid explodes at the m.p., 133 – 5°C

See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, TRIAZOLES

2911. Sodium phenylacetylide
[1004-22-4]

C_8H_5Na



Nef, J. U., *Ann.*, 1899, **308**, 264


The ether-moist powder ignites in air.

See ALKALI-METAL DERIVATIVES OF HYDROCARBONS See other METAL ACETYLIDES

2912. Phenylacetylene

[536-74-3]

C₈H₆



Palladium(II) acetate

Trumbo, D. L. *et al.*, *J. Polym. Sci., A, Polymer Chem.*, 1987, **25**, 1027–1034

Among palladium(II) salts used to polymerise phenylacetylene, the acetate led to rapid and very exothermic polymerisation, sometimes leading to explosion.

See other POLYMERISATION INCIDENTS

Perchloric acid


See Perchloric acid: Phenylacetylene

See other ALKYNES

2913. 3-Methyl-2-nitrobenzoyl chloride

[50424-93-6]

C₈H₆ClNO₃



1. Dahlbom, R., *Acta Chem. Scand.*, 1960, **14**, 2049


2. Anon., *Loss Prev. Bull.*, 1977, (013), 16

Attempted distillation caused an explosion [1], and a 200 g sample deflagrated at 120–130°C under high vacuum [2].

See other NITROACYL HALIDES

2914. 2-Nitrophenylacetyl chloride
[22751-23-1]

$C_8H_6ClNO_3$




Hayao, S., *Chem. Eng. News*, 1964, **42**(13), 39

Distillation of solvent chloroform from the preparation caused the residue to explode violently on 2 occasions. Previous publications had recommended use of solutions of the chloride, in preference to isolated material.

See other NITROACYL HALIDES

2915. 3-Methoxy-2-nitrobenzoyl chloride
[15865-57-3]

$C_8H_6ClNO_4$



1. Goodwin, B., *Chem. Brit.*, 1987, **23**, 1180–1181

2. Bretherick, L., *Chem. Brit.*, 1988, **24**, 125


During an attempt to prepare 3-methoxy-2-nitrobenzoyl cyanide by heating the title chloride with copper(I) cyanide to over 200°C, the reaction mixture had decomposed vigorously at around 60°C. This was attributed to thermal instability of the intended product [1], but available evidence suggests that instability of the title nitroacyl halide is a much more likely cause. However, there is also the possibility of a reaction involving demethylation of the methoxynitrobenzoyl chloride by traces of hydrogen chloride, and of a subsequent polycondensation reaction of the hydroxynitroacyl chloride so formed and of its subsequent exothermic decomposition, analogous to that of the 4-methoxy-3-nitro isomer. The presence of the copper salt would be expected to catalyse either of the possibilities of exothermic reaction [2].

See 4-Methoxy-3-nitrobenzoyl chloride

See entry NITROACYL HALIDES

2916. 4-Methoxy-3-nitrobenzoyl chloride
[10397-28-1]

$C_8H_6ClNO_4$



Preparative hazard


Grewer, T. *et al.*, *Chem. Ing. Tech.*, 1977, **49**, 562–563

Preparation of the acid chloride by an established procedure involved heating the acid with sulfinyl chloride, finally at 100°C for 4 h, when evolution of hydrogen chloride and sulfur dioxide was complete (and the vessel appears then to have been isolated from the gas absorption system). An apparently normally completed batch led to a double explosion with rupture of the process vessel. Subsequent thermoanalytical investigation revealed that the nitroacyl chloride is demethylated by hydrogen chloride produced in the primary reaction, methyl chloride being formed as well as the hydroxyacyl chloride. The latter condenses to form a poly(nitrophenylcarboxylic ester) and hydrogen chloride, which can lead to further demethylation. The sequence of exothermic reactions leads to accelerating decomposition, which becomes violent at about 350–380°C, with generation of pressure up to 40 bar by the volatiles produced. The double explosion was attributed to ignition of chloromethane, followed by that of the decomposing poly(nitro ester). Preventive measures are discussed.

See other GAS EVOLUTION INCIDENTS, NITROACYL HALIDES, POLYCONDENSATION REACTION INCIDENTS

2917. 4-Cyano-3-nitrotoluene
[26830-95-5]

$C_8H_6N_2O_2$




Preparative hazard

See 4-Chloro-3-nitrotoluene: Copper(I) cyanide, Pyridine

See other CYANO COMPOUNDS, NITROARYL COMPOUNDS

2918. 1,4-Dinitropentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane
[87830-30-6]


C₈H₆N₂O₄



See entry CUBANES

2919. 2,5-Dinitro-3-methylbenzoic acid
[70343-15-6]

C₈H₆N₂O₆



Oleum, Sodium azide


Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1979, **44**, 2504

Addition of sodium azide to a solution of the acid in 20% oleum at 5–10°C to produce 3-amino-2,5-dinitrotoluene must be slow (0.1 g portions during 1 h) to avoid explosion.

See other ORGANIC ACIDS, POLY-NITROARYL COMPOUNDS

2920. Phenoxyacetylene
[4279-76-9]

C₈H₆O




Jacobs, T. L. *et al.*, *J. Amer. Chem. Soc.*, 1942, **64**, 223

Small samples heated rapidly in sealed tubes to around 100°C exploded.

See other ACETYLENIC COMPOUNDS

2921. *cis*-3,4-Diethynylcyclobut-3-ene-1,2-diol
[125358-28-3]

C₈H₆O₂



Diederich, F., *et al.*, *J. Amer. Chem. Soc.*, 1990, **112**, 1616


The title compound explodes on heating to 55°C. The ethylene ketal of the corresponding diketone explodes at 118°C.

See 11,12-Diethynyl-1,4,7,10-tetraoxadispiro[4.0.4.2]dodec-11-ene

See *other* ACETYLENIC COMPOUNDS

†2922. Phenylglyoxal
[1074-12-0]


C₈H₆O₂



2923. Phthalic acid
[88-99-3]

C₈H₆O₄

1,2-C₆H₄(CO.OH)₂



HCS 1980, 753

Other reactants

Yoshida, 1980, 298

MRH values calculated for 12 combinations with oxidants are given.

Sodium nitrite


MRH 3.64/79

See Sodium nitrite: Phthalic acid

See *other* ORGANIC ACIDS

2924. Terephthalic acid
[100-21-0]

$C_8H_6O_4$




Preparative hazard

See Nitric acid: Hydrocarbons (reference 6)

See other ORGANIC ACIDS

2925. Diperoxyterephthalic acid
[1711-42-8]

$C_8H_6O_6$




Baeyer, A. *et al.*, *Ber.*, 1901, **34**, 762

It explodes when heated or struck.

See other PEROXYACIDS

2926. 4-Bromomethylbenzoic acid
[6232-88-8]

$C_8H_7BrO_2$




Preparative hazard

See *N*-Bromosuccinimide: Dibenzoyl peroxide, etc.

See other BENZYL COMPOUNDS, ORGANIC ACIDS

2927. Sodium 4-chloroacetophenone oximate
[1956-39-4] (oxime)

C₈H₇ClNNaO




van Dijk, J. *et al.*, *J. Med. Chem.*, 1977, **20**, 1201, 1205

Admission of air to the evacuated container of the dry oximate led to explosion on 2 occasions: nitrogen purging is advised.

See related OXIMES *See other* N-O COMPOUNDS

2928. Chloro-(4-methoxyphenyl)diazirine
[4222-26-8]

C₈H₇ClN₂O




Griller, D. *et al.*, *J. Amer. Chem. Soc.*, 1982, **104**, 5549

A cold sample decomposed violently on rapid warming to ambient temperature.

See other DIAZIRINES

2929. 4-Chloroacetophenone
[99-91-2]

C₈H₇ClO



Dimethylamine

Lundstedt, T. *et al.*, *Acta Chem. Scand., Ser. B*, 1984, **B38**, 717-719


During optimisation of a preparative procedure, heating mixtures of 4-chloroacetophenone and dimethylamine (4.22:1) at 234°C led to two explosions. Use of a safety screen is advised.

See other AMINATION INCIDENTS

2930. 4-Methoxybenzoyl chloride (Anisoyl chloride)

[100-07-2]

$C_8H_7ClO_2$



1. Carroll, D. W., *Chem. Eng. News*, 1960, **38**(34), 40

2. Ager, J. H. *et al.*, *Chem. Eng. News*, 1960, **38**(43), 5


Two incidents involving explosion of bottles of chloride stored at ambient temperature are described [1,2]. Safe preparation with storage at 5°C is detailed [2].

See other ACYL HALIDES

2931. Benzyl chloroformate

[501-53-1]

$C_8H_7ClO_2$



Iron salts

Anon., *Loss Prev. Bull.*, 1975, (003), 2

The ester is made by adding benzyl alcohol slowly to a preformed solution of phosgene in toluene at 12–16°C, toluene solvent finally being distilled off under vacuum. When discoloured phosgene was used (probably containing iron salts from corrosion of the cylinder), a violent explosion occurred during the distillation phase, presumably involving iron-catalysed decomposition of the chloroformate ester.


See other CATALYTIC IMPURITY INCIDENTS, CORROSION INCIDENTS

Water

See ARYL CHLOROFORMATES: water *See related* ACYL HALIDES

2932. Tricarbonyl(1-methylpyrrole)chromium(0) (Tricarbonyl[(1,2,3,4,5- η)-1-methyl-1H-pyrrole]chromium)
[33506-43-3]


C₈H₇CrNO₃



Goti, A. *et al.*, *J. Organomet. Chem.*, 1994, **470**(1-2), C4
This half-sandwich compound is extremely pyrophoric.
See other ORGANOMETALLICS

2933. Copper 1,3,5-octatrien-7-ynide
[]


C₈H₇Cu



Georgieff, K. K. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 5495
It deflagrates on heating in air.
See other METAL ACETYLIDES

2934. 1-Fluoro-1,1-dinitro-2-phenylethane
[22692-30-4]


C₈H₇FN₂O₄



See entry FLUORODINITROMETHYL COMPOUNDS (reference 4)

2935. Phenylacetonitrile
[140-29-4]

C₈H₇N



HCS 1980, 203


Sodium hypochlorite

See Sodium hypochlorite: Phenylacetonitrile

See other BENZYL COMPOUNDS, CYANO COMPOUNDS

2936. 2-Nitroacetophenone
[614-21-1]

C₈H₇NO₃



Potassium methylselenide

Leitem, L. *et al.*, *Compt. rend. C*, 1974, **278**, 276


Interaction in dimethylformamide was explosive.

See other NITROARYL COMPOUNDS

2937. 3-Methoxy-2-nitrobenzaldehyde
[53055-05-3]

C₈H₇NO₄

MeOC₆H₃(NO₂)CO.H



Initial decomposition temperature by DSC was 245°C.


See entry NITROBENZALDEHYDES

See other ALDEHYDES, NITROARYL COMPOUNDS

2938. 4-Nitrophenylacetic acid
[104-03-0]

C₈H₇NO₄

O₂NC₆H₄CH₂CO.OH



Acetic anhydride, Pyridine

1. Batchelor, J. F., 1992, Personal communication

2. Smith, J., *J. Amer. Chem. Soc.*, 1953, **75**, 1134


The distillation residue from 0.8 molar preparation of 2-acetoxy-1(4-nitrophenyl) prop-1-ene from reaction of the above followed by distillation, according to [2],

erupted on being left to cool, spraying a fume cupboard with tar. The preparation had previously been uneventful at smaller scale [1]. The reaction is not high yielding, indicating much residue and a pot temperature perhaps in excess of 200°C to distil the product.

See other NITROARYL COMPOUNDS

2939. Methyl 2-azidobenzoate
[16714-23-1]

C₈H₇N₃O₂




491M, 1975, 257

It exploded during distillation.

See other ORGANIC AZIDES

2940. 2-Methyl-5-nitrobenzimidazole
[1792-40-1]

C₈H₇N₃O₂




Safe preparation

See Nitric acid: 2-Methylbenzimidazole, etc.

See other NITROARYL COMPOUNDS

2941. 3,5-Dinitro-2-toluamide
[148-01-6]

C₈H₇N₃O₅



1. *The Explosion at the Dow Chemical Factory, King's Lynn 27th June 1976*, London, Health & Safety Executive, HMSO, 1977

2. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

A 1.3 t batch of off-spec. material had been redried in a rotating double cone vacuum drier heated by steam to around 120–130°C for 23 h. It was left in the

drier after heating had been discontinued (though cooling was not applied) and air had been allowed in, to await discharge the next working day. After 2 h, a violent explosion occurred causing one fatality and much destruction. Later investigation of the stability of the compound, notably by ARC techniques, showed that the excessively long drying time had probably caused the onset of thermal decomposition while heat was still being applied, and that further storage under virtually adiabatic conditions would have led to the steady acceleration of the exothermic decomposition to the point of explosion. At this point, it was estimated that a temperature in excess of 800°C had been attained in the drier at detonation, when the shock wave would generate a pressure approaching 800 bar. DSC techniques had shown exotherms at 273° for unheated material, and of 248–259 for material which had been preheated using conditions thought to be present in the drier [1]. The finely powdered amide is also a significant dust explosion hazard, maximum explosion pressures of 10.4 bar and a maximum rate of rise above 680 bar/s having been determined [2].

See other POLYNITROARYL COMPOUNDS

See DRYING, THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

2942. Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane
[277-10-1]


C₈H₈



See entry CUBANES

†2943. 1,3,5,7-Cyclooctatetraene
[629-20-9]

C₈H₈



The tetraene is endothermic by some 2.5 kJ/g; which is about the energy of explosion of old-fashioned black powder.

Dimethyl acetylenedicarboxylate

1. Bianchi, G., *Chem. in Brit.*, 1996, **32**(8), 22

In an attempt to obtain a 1:1 Diels Alder adduct, small quantities were sealed into a steel bomb and incubated in an oven at 140°C. After a time there was a violent explosion, converting the oven to an irregular perforated sphere [1]. The editor

suspects that the bursting bomb released flammable vapours that ignited in the cubic oven, distorting the shapnel punctured body.

See Lithium perchlorate: Diethyl ether

Oxygen


See Oxygen: Cyclooctatetraene

See related DIENES

2944. 1,3,7-Octatrien-5-yne

[16607-77-5]

C₈H₈



1. Nieuwland, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1931, **53**, 4201

2. Dolgopolski, I. M. *et al.*, *Chem. Abs.*, 1948, **42**, 4517e

3. Georgieff, K. K. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 5495


This tetramer of acetylene decomposes violently on distillation at 156°C, but not at reduced pressure [1]. It polymerises on standing to a solid detonable by shock [2]. Explosions occurred during attempted analytical combustion [3].

See related ALKYNES, DIENES

2945. Styrene (Ethenylbenzene)

[100-42-5]

C₈H₈



(MCA SD-37, 1971); NSC 627, 1982; FPA H44, 1976; HCS 1980, 866

RSC Lab. Hazard Data Sheet No. 22, 1984

1. Bond, J., *Loss Prev. Bull.*, 1985, (065), 25

2. MCA SD-37, 1971

3. Harmon, 1974, 2.17

4. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

The autocatalytic exothermic polymerisation reaction exhibited by styrene was involved in a plant-scale incident where accidental heating caused violent ejection of liquid and vapour from a storage tank [1]. Polymerisation becomes self-sustaining above 95°C [2]. The monomer has been involved in several plant-scale explosions, and must be stored at below 32°C, and for less than 3 months [3]. Styrene is a moderately endothermic compound (ΔH_f° (l) +147.7 kJ/mol,

1.42 kJ/g), and the energy of decomposition (110–300°C) has been measured as 0.67 kJ/g [4].

See other ENDOTHERMIC COMPOUNDS

Air, Polymerising styrene

Cullis, C. F. *et al.*, *Combust. Flame*, 1978, **31**, 1–5

Introduction of polymerising liquid styrene into a heated styrene vapour–air mixture can, despite the temporary cooling effect, cause ignition to occur under conditions where it would not normally do so. This effect had been noted previously in plant incidents.

See other POLYMERISATION INCIDENTS

Alkali-metal–graphite compounds

Williams, N. E., private comm., 1968

Interlaminar compounds of sodium or potassium in graphite will ionically polymerise styrene (and other monomers) smoothly. The occasional explosions experienced were probably due to rapid collapse of the layer structure and release of very finely divided metal.

Butyllithium

Roper, A. N. *et al.*, *Br. Polym. J.*, 1975, **7**, 195–203

Thermal explosion which occurred during fast anionic polymerisation of styrene, catalysed by butyllithium, was prevented by addition of low MW polystyrene before the catalyst.

See other POLYMERISATION INCIDENTS

Chlorine, Iron(III) chloride

See Chlorine: Iron(III) chloride, Monomers

Dibenzoyl peroxide

Sebastian, D. H. *et al.*, *Polym. Eng. Sci.*, 1976, **16**, 117–123

The conditions were determined for runaway/non-runaway polymerisation of styrene in an oil-heated batch reactor at 3 bar, using dibenzoyl peroxide as initiator at 3 concentrations. Results are presented diagrammatically.

See other POLYMERISATION INCIDENTS

Initiators

Biesenberger, J. A. *et al.*, *Polym. Eng. Sci.*, 1976, **16**, 101–116

The parameters were determined in a batch reactor for thermal runaway polymerisation of styrene, initiated by azoisobutyronitrile, dibenzoyl peroxide or di-*tert*-butyl peroxide.

See other POLYMERISATION INCIDENTS

Other reactants

Yoshida, 1980, 218

MRH values calculated for 14 combinations with oxidants are given.

Oxygen

Barnes, C. E. *et al.*, *J. Amer. Chem. Soc.*, 1950, **72**, 210

MRH 9.75/73

Exposure of unstabilised styrene to oxygen at 40–60°C generated a styrene–oxygen interpolymeric peroxide which, when isolated, exploded violently on gentle heating.

See other POLYPEROXIDES

Trichlorosilane


See Trichlorosilane: Styrene

See related ALKENES

2946. 1,2-Bis(chloromethyl)benzene

[612-12-4]

$C_8H_8Cl_2$



Catalytic impurities

Naef, H., private comm., 1980

The molten material, after holding for 4 h at 78°C in a stainless steel vessel, underwent a thermal runaway reaction and 500 kg erupted through the vent line. It was later found that addition of 0.1% of rust to the hot material led to an accelerating self-condensation Friedel-Craft reaction, catalysed by iron(III) chloride, which led to formation of poly-benzyls accompanied by evolution of hydrogen chloride.

See Benzyl bromide: Molecular sieve


Benzyl chloride: Catalytic impurities

See other BENZYL COMPOUNDS, GAS EVOLUTION INCIDENTS, POLYCONDENSATION REACTION INCIDENTS

2947. Dipotassium μ -cyclooctatetraene

[59391-85-4]

$C_8H_8K_2$



Air, or Oxygen

1. Starks, D. F. *et al.*, *Inorg. Chem.*, 1974, **13**, 1307


2. Evans, W. J. *et al.*, *J. Org. Chem.*, 1981, **46**, 3927

The dry solid is violently pyrophoric, exploding on contact with oxygen [1], or air [2]. The solid produced by passing oxygen into a THF solution of the potassium complex must be kept wet with solvent, as it is shock-sensitive when dry [2].

See other ORGANOMETALLICS

2948. Dilithium μ -cyclooctatetraene
[37609-69-1]


$C_8H_8Li_2$



Spencer, J. L., *Inorg. Synth.*, 1979, **19**, 214
The solid (but not an ether solution) is pyrophoric in air.
See other ORGANOMETALLICS

2949. N-Nitrosoacetanilide
[938-81-8]

$C_8H_8N_2O_2$



Piperidine


Huisgen, R., *Ann.*, 1951, **573**, 163–181
The dry anilide explodes on contact with a drop of piperidine.

Thiophene

Bamberger, E., *Ber.*, 1887, **30**, 367
A mixture prepared at 0°C exploded when removed from the cooling bath.
See other NITROSO COMPOUNDS

2950. 4-Nitroacetanilide
[104-04-1]


$C_8H_8N_2O_3$



Sulfuric acid

See Sulfuric acid: Nitroaryl bases, etc.
See other NITROARYL COMPOUNDS

2951. Sodium 4,4-dimethoxy-1-*aci*-nitro-3,5-dinitro-2,5-cyclohexadienide
[12275-58-0] **C₈H₈N₃NaO₈**




Jackson, C. L. *et al.*, *Amer. Chem. J.*, 1898, **20**, 449

The compound is produced from interaction of trinitroanisole and sodium methoxide and explodes with great violence on heating in a free flame. It should probably be formulated as shown, a dimethyl acetal of the *aci-p*-quinonoid form of picric acid.

See other aci-NITROQUINONOID COMPOUNDS, POLYNITROARYL COMPOUNDS

2952. 5-Amino-3-phenyl-1,2,4-triazole
[4922-98-9] **C₈H₈N₄**



Nitrous acid

Polya, J. B., *Chem. & Ind.*, 1965, 812


The solution, obtained by conventional diazotisation of the amine, contained some solid which was removed by filtration on a glass frit. The solid, thought to be precipitated diazonium salt (possibly an internal salt), exploded violently when disturbed with a metal spatula.

See 5-Aminotetrazole

See other DIAZONIUM SALTS, DIAZOZOLES, TRIAZOLES

2953. 1,1-Diazidophenylethane
[87272-16-0]


C₈H₈N₆



Kinlager, S. *et al.*, *Synthesis*, 1983, 501
The *gem*-diazide is potentially explosive.
See other ORGANIC AZIDES

2954. 1,2-Diazido-1-phenylethane
[22710-73-2]


C₈H₈N₆



Preparative hazard
See Manganese(III) azide: Styrene
See other ORGANIC AZIDES

2955. 7,8-Dioxabicyclo[4.2.2]-2,4,7-decatriene
[83931-97-8]


C₈H₈O₂



Preparative hazard
See Oxygen (Gas): Cyclooctatetraene
See other CYCLIC PEROXIDES

2956. 4-Methoxybenzaldehyde (Anisaldehyde)
[123-11-5]

C₈H₈O₂



Preparative hazard

See Oxygen (Gas): 4-Methoxytoluene

See ALDEHYDES

2957. Poly(styrene peroxide)
[33873-53-9]

(C₈H₈O₂)_n




See Styrene: Oxygen

See other POLYPEROXIDES

2958. 4-Hydroxy-3-methoxybenzaldehyde (Vanillin)
[121-33-5]

C₈H₈O₃




Formic acid, Thallium(III) nitrate

See Thallium(III) nitrate: Formic acid, etc.

See other ALDEHYDES

2959. 4-Methylbenzyl chloride
[104-82-5]

C₈H₉Cl



Iron


Hakl, J., *Runaway Reactions*, 1981, Paper 3/L, 4

Initial exothermic decomposition sets in at 55°C in presence of 0.02% of iron.

See other BENZYL COMPOUNDS, CATALYTIC IMPURITY INCIDENTS

2960. Dimethyl 2-chloro-4-nitrophenyl thionophosphate
[2643-84-5]

C₈H₉ClNO₅PS




McPherson, J. B. *et al.*, *J. Agric. Food Chem.*, 1956, **4**, 42

Thermal decomposition of a 1.5 g sample at about 270°C leads to effervescence after 20 s, with later deflagration and combustion, but a 5 g sample ignites immediately during decomposition. The induction period increases with decreasing temperature, to 384 s at 174°C. The 3-chloro analogue behaves similarly.

See other INDUCTION PERIOD INCIDENTS, PHOSPHORUS ESTERS

2961. Dimethyl 3-chloro-4-nitrophenyl thionophosphate
[500-28-7]


C₈H₉ClNO₅PS



See Dimethyl 2-chloro-4-nitrophenyl thionophosphate (next above)
See other INDUCTION PERIOD INCIDENTS, PHOSPHORUS ESTERS

2962. 4-Methoxybenzyl chloride (Anisyl chloride)
[824-94-2]


C₈H₉ClO



1. Bryant, R. J., personal communication 1990
 2. Harger, M. J. P. *et al.*, *J. Chem. Soc., Perk. I*, 1997, (4), 527
- This compound is usually made and used in situ. A 5 g sample was allowed to sit in a screw top vial for a couple of days, one morning it was found to have blown a hole in the lid and extruded a 20 ml plume of polymeric foam [1]. An attempt to dry anisyl chloride over molecular sieves pressurised the container, presumably with hydrogen chloride evolved during polymerisation [2].
- See other* BENZYL COMPOUNDS, POLYCONDENSATION REACTION INCIDENTS

2963. 2,4-Dimethylbenzenediazonium triiodide
[]

C₈H₉I₃N₂



See entry DIAZONIUM TRIIODIDES

2964. Potassium 1-phenylethanediazoate

[]

C₈H₉KN₂O



White, E. H. *et al.*, *J. Amer. Chem. Soc.*, 1992, **114**(221), 8023


The *syn*-isomer decomposed exothermically to styrene when handled in a nitrogen atmosphere.

See related DIAZONIUM SALTS *See other* N-O COMPOUNDS

2965. 5(1,1-Dinitroethyl)-2-methylpyridine

[]

C₈H₉N₃O₄




See Nitric acid: 5-Ethyl-2-methylpyridine

See other POLYNITROALKYL COMPOUNDS

2966. 1,3,5-Cyclooctatriene

[1871-52-9]

C₈H₁₀




See Diethyl acetylenedicarboxylate: 1,3,5-Cyclooctatriene

See related DIENES

2967. 6,6-Dimethylfulvene

[2175-91-9]

C₈H₁₀



Air, or Ether

Engler, C. *et al.*, *Ber.*, 1901, **34**, 2935


Dimethylfulvene readily peroxidised to give an insoluble (polymeric?) peroxide which exploded violently on heating to 130°C, and also caused ether used to triturate it to ignite.

See 1,4-Epidioxy-1,4-dihydro-6,6-dimethylfulvene

See other DIENES, POLYPEROXIDES

†2968. Ethylbenzene
[100-41-4]


C_8H_{10}



FPA H99, 1981; HCS 1980, 470; RSC Lab. Hazards Safety Data Sheet No. 74, 1988

†2969. *m*ixo-Xylene
[1330-20-7]


C_8H_{10}



NSC 204, 1968; FPA H76, 1978; RSC Lab. Hazard Data Sheet No. 16, 1983

†2970. *o*-Xylene
[95-47-6]

C_8H_{10}



Gorecki, J. *et al.*, *Arch. Combust.*, 1988, **8**, 123–127

The aerobic oxidation of *o*-xylene to phthalic acid or anhydride at elevated temperatures is industrially important, and the flammability limits and explosion parameters at 350°C under a range of pressures have been redetermined.

See also OXIDATION PROCESSES


Other reactants

Yoshida, 1980, 101

MRH values calculated for 13 combinations with oxidants are given.

†2971. *m*-Xylene
[108-38-3]

C_8H_{10}




HCS 1980, 962

Sodium

See Sodium: Xylene

†2972. *p*-Xylene
[106-42-3]

C_8H_{10}



HCS 1980, 963

Acetic acid, Air

Shraer, B. I., *Khim. Prom.*, 1970, **46**(10), 747–750

In liquid phase aerobic oxidation of *p*-xylene in acetic acid to terephthalic acid, it is important to eliminate the inherent hazards of this fuel–air mixture. Effects of temperature, pressure and presence of steam on the explosive limits of the mixture have been studied.

See OXIDATION PROCESSES

1,3-Dichloro-5,5-dimethyl-2,4-imidazolidindione


See 1,3-Dichloro-5,5-dimethyl-2,4-imidazolidindione: Xylene

Nitric acid

See Nitric acid: Hydrocarbons

2973. 4-Bromodimethylaniline
[586-77-6]

C₈H₁₀BrN




Anon., *Chem. Eng. News*, 1961, **39**(13), 37

During vacuum distillation, self-heating could not be controlled, and proceeded to explosion. Internal dehydrohalogenation to a benzyne seems a possibility.

See other HALOANILINES

2974. 2-Bromo-3,5-dimethoxyaniline
[70277-99-5]

C₈H₁₀BrNO₂




491M, 1975, 74

During laboratory-scale fractional distillation of a mixture of the aniline and its mono- and di-brominated derivatives, an explosion occurred, attributed to the lability of the bromine substituent.

See other AMINOMETHOXY COMPOUNDS, HALOANILINES

2975. 2-Chloro-N-(2-hydroxyethyl)aniline
[94-87-1]

C₈H₁₀ClNO



CISHC Chem. Safety Summ., 1977, **48**, 25; Shaw, A. W., private comm., 1978


During the vacuum fractional distillation of bulked residues (7.2 t containing 30–40% of the bis(hydroxyethyl) derivative, and up to 900 ppm of iron) at 210–225°C/445–55 mbar in a mild steel still, a runaway decomposition set in and accelerated to explosion. Laboratory work on the material charged showed that exothermic decomposition on the large scale would be expected to set in around 210–230°C, and that the induction time at 215°C of 12–19 h fell to 6–9 h in presence of mild steel. Quantitative work in sealed tubes showed a maximum rate of pressure rise of 45 bar/s, to a maximum developed pressure of 200 bar. The thermally induced decomposition produced primary amine, hydrogen chloride, ethylene, methane, carbon monoxide and carbon dioxide.

See related HALOANILINES

See other GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS

2976. 2(3-Chlorophenoxy)ethylamine
[6488-00-2]

C₈H₁₀ClNO




Energy of exothermic decomposition in range 270–360°C was measured as 0.27 kJ/g

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

2977. N-(4-Chloro-2-nitrophenyl)-1,2-diaminoethane
[59320-16-0]

C₈H₁₀ClN₃O₂




See N-(2-Nitrophenyl)-1,2-diaminoethane

See other HALOANILINES, NITROARYL COMPOUNDS

2978. Bis(ethoxycarbonyldiazomethyl)mercury
[20539-85-9]

$C_8H_{10}HgN_4O_4$




Buchner, E., *Ber.*, 1895, **28**, 217

It decomposes with foaming on melting at 104°C, and will explode under a hammer-blow.

See other DIAZO COMPOUNDS, MERCURY COMPOUNDS

2979. Dimethyl 4-nitrophenyl thionophosphate
[298-00-0]

$C_8H_{10}NO_5PS$




McPherson, J. B. *et al.*, *J. Agric. Food Chem.*, 1956, **4**, 42–49

Overheating during removal of solvent by distillation from a pilot batch of 'methyl parathion' led to explosive decomposition, and the course of the 2-stage decomposition was studied. A 1.5 g sample immersed at 270°C decomposes after an induction period of 54 s and the residue later deflagrates, but a 5 g sample deflagrates during the initial decomposition.

See other INDUCTION PERIOD INCIDENTS, PHOSPHORUS ESTERS

2980. *N,N*-Dimethyl-4-nitrosoaniline
[138-89-6]

$C_8H_{10}N_2O$



Energy of exothermic decomposition (in range 120–310°C) measured as 1.92 kJ/g by DSC, and $T_{\text{ait}24}$ was determined as 87°C by adiabatic Dewar tests, with an apparent energy of activation of 123 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Acetic anhydride, Acetic acid

Bain, P. J. S. *et al.*, *Chem. Brit.*, 1971, **7**, 81


An exothermic reaction, sufficiently violent to expel the flask contents, occurs after an induction period of 15–30 s when acetic anhydride is added to a solution of the nitroso compound in acetic acid. 4,4'-Azobis(*N,N*-dimethylaniline), isolated from the reaction tar, may have been formed in a redox reaction, possibly involving an oxime derived from the nitroso compound.

See other INDUCTION PERIOD INCIDENTS, REDOX REACTIONS

See other NITROSO COMPOUNDS

2981. 2-Ethylpyridine-4-carbothioamide
[536-33-4]


$\text{C}_8\text{H}_{10}\text{N}_2\text{S}$



See FRICTIONAL INITIATION INCIDENTS (reference 3)

2982. 1-Acetyl-4-(4'-sulfophenyl)-3-tetrazene (4-(4-Acetyl-1-tetrazenyl)benzene-sulfonic acid)
[70324-27-5]

$\text{C}_8\text{H}_{10}\text{N}_4\text{O}_4\text{S}$




Houben-Weyl, 1965, Vol. 10.3, 740

It explodes on heating.

See other HIGH-NITROGEN COMPOUNDS, ORGANIC ACIDS, TETRAZENES

2983. Benzeneethanol (Phenethyl alcohol)
[60-12-8]

$C_8H_{10}O$




Nitric acid

See Nitric acid: Benzeneethanol, Sulfuric acid

2984. 1,4-Epidioxy-1,4-dihydro-6,6-dimethylfulvene (7-(1-Methylethylidene)-2,3-dioxabicyclo[2.2.1]hept-5-ene)
[51027-90-8]

$C_8H_{10}O_2$



Harada, N. *et al.*, *Chem. Lett. (Japan)*, 1973, 1173–1176


When free of solvent it decomposes explosively above $-10^{\circ}C$.

See 6,6-Dimethylfulvene: Air, or Ether

See other CYCLIC PEROXIDES

2985. 1-Hydroperoxphenylethane
[3071-32-7]

$C_8H_{10}O_2$




Druliner, J. D., *Chem. Eng. News*, 1981, **59**(14), 3

A 9 g sample of the freshly prepared hydroperoxide decomposed after 20 min at ambient temperature, bursting the 20 ml glass container. A 30% solution of the hydroperoxide in ethylbenzene is stable.

See other ALKYL HYDROPEROXIDES

2986. Dicrotonoyl peroxide
[93506-63-9]

C₈H₁₀O₄




Guillet, J. E. *et al.*, Ger. Pat. 1 131 407, 1962

The diacyl peroxide and its lower homologues, though of relatively high thermal stability, are very shock-sensitive. Replacement of the hydrogen atoms adjacent to the carbonyl groups with alkyl groups renders the materials non-shock-sensitive.
See other DIACYL PEROXIDES

2987. Diethyl acetylenedicarboxylate
[762-21-0]

C₈H₁₀O₄




1,3,5-Cyclooctatriene

Foote, C. S., private comm., 1965

A mixture of the reactants being heated at 60°C to effect a Diels-Alder addition exploded. Onset of this vigorously exothermic reaction was probably delayed by an induction period, and presence of a solvent and/or cooling would have moderated it.
See other INDUCTION PERIOD INCIDENTS, ACETYLENIC COMPOUNDS

2988. Ethyl 3,4-dihydroxybenzenesulfonate
[]

C₈H₁₀O₅S




Acetyl nitrate

See Acetyl nitrate: Ethyl 3,4-dihydroxybenzenesulfonate

See other SULFUR ESTERS

2989. Diallyl peroxydicarbonate
[34037-79-1]


C₈H₁₀O₆



The distilled oil exploded on storage at ambient temperature.
See entry PEROXYCARBONATE ESTERS *See other* ALLYL COMPOUNDS

2990. Bis(3-carboxypropionyl) peroxide
[123-23-9]


C₈H₁₀O₈



Lombard, R. *et al.*, *Bull. Soc. Chim. Fr.*, 1963, **12**, 2800
It explodes on contact with flame. The commercial dry 95% material ('succinic acid peroxide') is highly hazard-rated.
See other COMMERCIAL ORGANIC PEROXIDES, DIACYL PEROXIDES, ORGANIC ACIDS

2991. Dimethylphenylarsine
[696-26-4]


C₈H₁₁As



Hydrogen peroxide
See Hydrogen peroxide: Dimethylphenylarsine
See related ALKYLNON-METALS

2992. 2,4,6-Trimethylpyrilium perchlorate
[940-93-2]

C₈H₁₁ClO₅




Hafner, K. *et al.*, *Org. Synth.*, 1964, **44**, 102

The crystalline solid is an impact- and friction-sensitive explosive and must be handled with precautions. These include use of solvent-moist material and storage in a corked rather than glass-stoppered vessel.

See other GLASS INCIDENTS, NON-METAL PERCHLORATES

2993. N,N-Dimethylaniline
[121-69-7]

C₈H₁₁N



HCS 1980, 425

Dibenzoyl peroxide

MRH 2.26/99+

See Dibenzoyl peroxide: N,N-Dimethylaniline

Diisopropyl peroxydicarbonate

See Diisopropyl peroxydicarbonate: Amines, etc.

Other reactants


Yoshida, 1980, 166

MRH values calculated for 14 combinations with oxidants are given.

See other ORGANIC BASES

2994. N-Ethylaniline
[103-69-5]

C₈H₁₁N




HCS 1980, 469

Nitric acid

See Nitric acid: Aromatic amines

2995. 5-Ethyl-2-methylpyridine
[104-90-5]

C₈H₁₁N




Nitric acid

See Nitric acid: 5-Ethyl-2-methylpyridine

See other ORGANIC BASES

2996. 3,3-Dimethyl-1-phenyltriazeno
[7227-91-0]

C₈H₁₁N₃



1. Baeyer, O. *et al.*, *Ber.*, 1875, **8**, 149


2. Heusler, F., *Ann.*, 1890, **260**, 249

It decomposes explosively on attempted distillation at ambient pressure [1], but may be distilled uneventfully at reduced pressure [2].

See other TRIAZENES

2997. 3-Methoxybenzylamine
[5071-96-5]

C₈H₁₁NO




Energy of decomposition (in range 280–430°C) measured as 0.22 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS, BENZYL COMPOUNDS

2998. 2,4-Dimethoxyaniline
[2735-04-8]


$C_8H_{11}NO_2$



Energy of decomposition (in range 160–440°C) measured as 0.55 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS

2999. 3,5-Dimethoxyaniline
[10272-07-8]


$C_8H_{11}NO_2$



Energy of decomposition (in range 320–450°C) measured as 0.83 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AMINOMETHOXY COMPOUNDS

3000. 1-(4-Methoxyphenyl)-3-methyltriazenes
[53477-43-3]



$C_8H_{11}N_3O$



Rondevstedt, C. S. *et al.*, *J. Org. Chem.*, 1957, **22**, 203
The liquid product exploded on attempted vacuum distillation at below 1 mbar.
See other TRIAZENES See related AMINOMETHOXY COMPOUNDS

3001. N-(2-Nitrophenyl)-1,2-diaminoethane
[51138-16-0]

C₈H₁₁N₃O₂



Doleschall, G. *et al.*, *Tetrahedron*, 1976, **32**, 59


This, and the corresponding 5-chloro and 5-methoxy derivatives, tend to explode during vacuum distillation, and minimal distillation pressures and temperatures are recommended. This instability, (which was not noted for the homologues in which the hydrogen atom of the secondary amino group was replaced by alkyl) may be connected with the possibility of isomerisation to the *aci*-nitro iminoquinone internal salt species.

See NITROAROMATIC-ALKALI HAZARDS, *aci*-NITROQUINONOID COMPOUNDS

See *other* NITROARYL COMPOUNDS

3002. Dimethylphenylphosphine oxide
[10311-08-7]

C₈H₁₁OP




Preparative hazard

See Hydrogen peroxide: Dimethylphenylphosphine

See *related* ALKYLPHOSPHINES

3003. Dimethylphenylphosphine
[672-66-2]

C₈H₁₁P




Hydrogen peroxide

See Hydrogen peroxide: Dimethylphenylphosphine

See *other* ALKYLPHOSPHINES

†3004. 4-Vinylcyclohexene
[100-40-3]


C₈H₁₂



See other PEROXIDISABLE COMPOUNDS

3005. 1-(4-Chlorophenyl)biguanidinium hydrogen dichromate
[15842-89-4]


C₈H₁₂ClCr₂N₅O₇



See entry DICHROMATE SALTS OF NITROGENOUS BASES See other OXOSALTS OF NITROGENOUS BASES

3006. 1,6-Diisocyanohexane
[929-57-7]


C₈H₁₂N₂



See entry DIISOCYANIDE LIGANDS

3007. Tetramethylsuccinodinitrile
[3333-52-6]

C₈H₁₂N₂




Preparative hazard

Azoisobutyronitrile: Heptane See other CYANO COMPOUNDS

3008. 1,6-Diisocyanatohexane
[822-06-0]

$C_8H_{12}N_2O_2$



Alcohols


491M, 1975, 19

Base-catalysed reactions of isocyanates, such as the title compound, with alcohols may be explosively violent in absence of diluting solvents.

See other ORGANIC ISOCYANATES

3009. tert-Butyl 2-diazoacetoacetate
[13298-76-5]

$C_8H_{12}N_2O_3$




Regitz, M. *et al.*, *Org. Synth.*, 1968, **48**, 38

During low-temperature crystallisation, scratching to induce seeding of the solution must be discontinued as soon as the sensitive solid separates.

See other DIAZO COMPOUNDS

3010. 6-Aminopenicillanic acid S-oxide
[4888-97-5]

$C_8H_{12}N_2O_4S$



Preparative hazard

1. Micetich, R. G., *Synthesis*, 1976, **4**, 264
2. Noponen, A., *Chem. Eng. News*, 1977, **55**(8), 5
3. Micetich, R. G., *Chem. Brit.*, 1977, **13**, 163


According to a published procedure, the sulfur-containing acid is oxidised to the sulfoxide with hydrogen peroxide and precipitated as the 4-toluenesulfonate salt in the presence of acetone [1]. During subsequent purification, trimeric acetone peroxide was precipitated and exploded violently after filtration [2]. The use of acetone in this and similar preparations involving hydrogen peroxide is not now recommended, and is highly dangerous [3].

See Hydrogen peroxide: Acetone, etc., also 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxonane

3011. Azoisobutyronitrile (AZDN)

[78-67-1]

C₈H₁₂N₄



1. Lamy, P., *J. Calorim. Anal. Therm.*, 1980, **11**, 2-10-1–2-10-11
2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
3. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987
4. Kilattab, M. A. *et al.*, *Bull. Fac. Sci. Alexandria Univ.*, 1995, **35**(2), 229; *Chem. Abs.*, 1996, **124**, 184211c
5. *CHETAH*, 1990, 187
6. Gladwell, P., *Loss Prev. Bull.*, 1998, 139, 3

AZDN is a much used polymerisation initiator. A mathematical model of thermal dissociation leading to explosion was verified experimentally with solutions of the nitrile in dibutyl phthalate and silicone oil [1]. It shows a very high rate of pressure increase during exothermic decomposition. Energy of decomposition (78–130°C) measured by DSC as 1.3 kJ/g [2], and further work on thermal decomposition under confinement has been done [3]. The recently calculated value of 56°C for the critical ignition temperature is in agreement with the previous value of 54°C and a more recent calorimetric study of runaway hazards [4]. It is shock sensitive [5]. A review of numerous fires and explosions caused by, or involving, AZDN has been published. Several appear to have started below the 40°C temperature now required by transport regulations. Refrigeration is desirable and confinement must be avoided. Solutions are even less stable than the solid and are not normally transported or stored [6].

See PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

See entry CRITICAL IGNITION TEMPERATURE

See BLOWING AGENTS

Acetone

Carlisle, P. J., *Chem. Eng. News*, 1949, **27**, 150

During recrystallisation of technical material from acetone, explosive decomposition occurred. Non-explosive decomposition occurred when the nitrile was heated alone, or in presence of methanol.

Ammonium peroxodisulfate

See Ammonium peroxodisulfate: Azoisobutyronitrile



See other AZO COMPOUNDS, BLOWING AGENTS, CYANO COMPOUNDS

Heptane

Scheffold, R. *et al.*, *Helv. Chim. Acta*, 1975, **58**, 60

During preparation of tetramethylsuccinodinitrile by thermal decomposition of 100 g of the azonitrile by slow warming in unstirred heptane, an explosion occurred. Successive addition of small portions of the nitrile to heptane at 90–92°C is a safer, preferred method.

3012. Diazido[(1,2,5,6- η)-1,5-Cyclooctadiene]platinum [144673-73-4]





Kim, Y. *et al.*, *Bull. Korean Chem. Soc.*, 1994, **15**(8), 690

Exploded during melting point determination.

See other AZIDES

3013. 1,3,4,6-Tetrakis(2-methyltetrazol-5-yl)-hexaaza-1,5-diene [83195-98-6]




Butler, R. N. *et al.*, *J. Chem. Res., Synop.*, 1982, 183

With 74% N content, it is explosive, sensitive to shock, friction or rapid heating to 121–125°C.

See other HIGH-NITROGEN COMPOUNDS, TETRAZOLES

3014. 3-Ethoxymethylene-2,4-pentanedione
[33884-41-2]

C₈H₁₂O₃




Riley, D. P. *et al.*, *Inorg. Synth.*, 1978, **18**, 37

The residue from distillation of the title compound (prepared from ethyl orthoformate, pentanedione and acetic anhydride) at 165–170°C/13 mbar, is pyrophoric when hot.

3015. O-O-tert-Butyl hydrogen monoperoxymaleate
[1931-62-0]

C₈H₁₂O₅




Castrantas, 1965, 17

Slightly shock-sensitive, the commercial dry 95% material is highly hazard-rated.

See other COMMERCIAL ORGANIC PEROXIDES, PEROXYESTERS

3016. Tetravinyllead
[866-87-5]

C₈H₁₂Pb



Holliday, A. K. *et al.*, *Chem. & Ind.*, 1968, 1699

In the preparation from lead(II) chloride and vinylmagnesium bromide in THF-hexane, violent explosions occurred during isolation of the product by distillation of solvent. This could be avoided by a procedure involving steam distillation of the tetravinyllead, no significant loss of yield by hydrolysis being noted. It is likely to be considerably more endothermic than tetraethyllead.

Diborane

See Diborane: Tetravinyllead

Phosphorus trichloride

Houben-Weyl 1975, Vol. 13.3, 244


In absence of solvent, interaction may be explosive

See related ALKYL METALS, ENDOTHERMIC COMPOUNDS

3017. 3-Buten-1-ynyl-diethylaluminum

[]

$C_8H_{13}Al$



Petrov, A. A. *et al.*, *Zh. Obsch. Khim.*, 1962, **32**, 1349


It ignites in air.

See other METAL ACETYLIDES, TRIALKYLALUMINIUMS

3018. 2-Bromo-2,5,5-trimethylcyclopentanone

[69167-99-3]

$C_8H_{13}BrO$



Potassium hydroxide, Triethylene glycol


Adam, W., *J. Amer. Chem. Soc.*, 1986, **108**, 4559

In the preparation of 2,5,5-trimethylcyclopentan-3-one by dehydrobromination of the title bromoketone, the scale should be limited to below 10 g of the latter, to avoid a vigorous exotherm and loss of control on addition of the bromoketone to potassium hydroxide in triethylene glycol at 60–70°C.

See related HALOALKANES

3019. 1-Phenylbiguanidinium hydrogen dichromate
[15760-45-9]

C₈H₁₃Cr₂N₅O₇




See entry DICHROMATE SALTS OF NITROGENOUS BASES

See other OXOSALTS OF NITROGENOUS BASES

3020. 1-Diethylamino-1-buten-3-yne
[1809-53-6]

C₈H₁₃N



Brandsma, 1971, 163

Dilution of the amine with white mineral oil is recommended before vacuum distillation, to avoid explosive decomposition of the residue.

See other ACETYLENIC COMPOUNDS

†**3021. 1,7-Octadiene**
[3710-30-3]

C₈H₁₄



See other DIENES

†**3022. 1-Octyne**
[629-05-0]

C₈H₁₄



See other ALKYNES


†**3023. 2-Octyne**
[2809-67-8]

C₈H₁₄



See other ALKYNES

†3024. 3-Octyne
[15232-76-5]



C_8H_{14}

See other ALKYNES


†3025. 4-Octyne
[1942-45-6]



C_8H_{14}

See other ALKYNES

†3026. Vinylcyclohexane
[695-12-5]




C_8H_{14}

See related ALKENES

3027. Di-2-butenylcadmium
[7544-40-3]

$\text{C}_8\text{H}_{14}\text{Cd}$




Houben-Weyl, 1973, Vol. 13.2a, 881

When rapidly warmed from -5°C to ambient temperature, it decomposes explosively.

See related ALKYL METALS, ALLYL COMPOUNDS

3028. μ -Cyclopentadienyl(methyl)-bis-(*N*-methyl-*N*-nitrosohydroxylamino)titanium
[58659-15-7]

$\text{C}_8\text{H}_{14}\text{N}_4\text{O}_4\text{Ti}$



Clark, R. J. H. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1974, 122


The complex was stable under nitrogen or vacuum, but exploded quite readily and decomposed in air.

See other NITROSO COMPOUNDS, N-O COMPOUNDS, ORGANOMETALLICS

3029. 4-Hydroxy-4-methyl-1,6-heptadiene

[25021-40-5]

C₈H₁₄O



Ozone

1. Rabinowitz, J. L., *Biochem. Prep.*, 1958, **6**, 25

2. Miller, F. W., *Chem. Eng. News*, 1973, **51**(6), 29


Following a published procedure [1], but using 75 g of diene instead of the 12.6 g specified, the ozonisation product exploded under desiccation. Eight previous preparations on the specified scale had been uneventful.

See OZONIDES

3030. Poly(vinyl butyral)

[63148-65-2]

(C₈H₁₄O₂)_n




Korshova, I. T. *et al.*, *Chem. Abs.*, 1983, **99**, 195868

Ignition and autoignition temperatures, lower explosion limit and oxygen index were determined for the powdered butyral in relation to dispersibility and viscosity.

3031. Diethyl succinate

[123-25-1]

C₈H₁₄O₄



HCS 1980, 405

Ethyl trifluoroacetate, Sodium hydride

See Sodium hydride: Diethyl succinate, etc.

3032. Diisobutyryl peroxide

[3437-84-1]


C₈H₁₄O₄

1. Kharasch, S. *et al.*, *J. Amer. Chem. Soc.*, 1941, **63**, 526
2. *MCA Case History No. 579*

A sample of the peroxide in ether, prepared according to a published procedure [1], was being evaporated to dryness with a stream of air when it exploded violently. Handling the peroxide as a dilute solution at low temperature is recommended [2].
See other DIACYL PEROXIDES

3033. Acetyl cyclohexanesulfonyl peroxide

[3179-56-4]

C₈H₁₄O₅S

1. 'Code of Practice for Storage of Organic Peroxides', GC14, Widnes, Interox Chemicals Ltd, 1970
2. *CHETAH*, 1990, 182

While the commercial material damped with 30% water is not shock- or friction-sensitive, if it dries out it may become a high-hazard material. As a typical low-melting diacyl peroxide, it may be expected to decompose vigorously or explosively on slight heating or on mechanical initiation [1]. It is shock sensitive (presumably when dry) [2].

See other COMMERCIAL ORGANIC PEROXIDES, DIACYL PEROXIDES

3034. Diisopropyl peroxydicarbonate

[105-64-6]

C₈H₁₄O₆

1. Strong, W. A., *Ind. Eng. Chem.*, 1964, **56**(12), 33
 2. 'Bulletin, T. S. 350', Pittsburgh, Pittsburgh Plate Glass Co., 1963
- When warmed slightly above its m.p., 10°C, the ester undergoes slow but self-accelerating decomposition, which may become dangerously violent under confinement. Bulk solutions of the peroxyester (45%) in benzene-cyclohexane stored at 5°C developed sufficient heat to decompose explosively after 1 day, and 50–90% solutions were found to be impact-sensitive [1]. The solid is normally stored and transported at below –18°C in loose-topped trays [2].

Amines, or Potassium iodide

Strain, F., *J. Amer. Chem. Soc.*, 1950, **72**, 1254

At 20–30°C, decomposition occurred in 10–30 min. Addition of 1% of aniline, 1,2-diaminoethane or potassium iodide caused instant decomposition, and of dimethylaniline, instant explosion.

See other COMMERCIAL ORGANIC PEROXIDES, DIACYL PEROXIDES, PEROXYCARBONATE ESTERS

3035. Dipropyl peroxydicarbonate

[16066-38-9]

C₈H₁₄O₆

(PrOCO.O-)₂

Barter, J. A., US Pat. 3 775 341, 1973


Methylcyclohexane is a suitable solvent to reduce the hazardous properties of the peroxyester.

See other DIACYL PEROXIDES, PEROXYCARBONATE ESTERS

3036. 4-Hydroxy-4-methyl-1,6-heptadiene diozonide

[]

C₈H₁₄O₇




See 4-Hydroxy-4-methyl-1,6-heptadiene: Ozone

See other OZONIDES

3037. Bis(2-methoxyethyl) peroxydicarbonate

[22575-95-7]

C₈H₁₄O₈




It exploded at 34°C.

See entry PEROXYCARBONATE ESTERS

3038. μ -Cyclopentadienyltrimethyltitanium
[38386-55-9]


C₈H₁₄Ti



Bailar, 1973, Vol. 3, 395
Pyrophoric at ambient temperature.
See other ALKYLMETALS, ORGANOMETALLICS

†3039. Dimethylcyclohexanes (5 isomers)
[27195-67-1]

C₈H₁₆



†3040. 1-Octene
[111-66-0]

C₈H₁₆



See other ALKENES

†3041. 2-Octene
[111-67-1]

C₈H₁₆



See other ALKENES

†3042. 2,3,4-Trimethyl-1-pentene
[565-76-4]

C₈H₁₆



See other ALKENES

†3043. 2,4,4-Trimethyl-1-pentene
[107-39-1]



C_8H_{16}

HCS 1980, 414

See other ALKENES


†3044. 2,3,4-Trimethyl-2-pentene
[565-77-5]



C_8H_{16}

See other ALKENES


†3045. 2,4,4-Trimethyl-2-pentene
[107-40-4]



C_8H_{16}

See other ALKENES

†3046. 3,4,4-Trimethyl-2-pentene
[598-96-9]




C_8H_{16}

See other ALKENES

3047. 4,4'-Dithiodimorpholine
[103-34-4]

$\text{C}_8\text{H}_{16}\text{N}_2\text{O}_2\text{S}_2$




Chlorine

See 4-Morpholinesulfonyl chloride

3048. 3-Azoniabicyclo[3.2.2]nonane nitrate
[10308-93-7]

C₈H₁₆N₂O₃




Violent decomposition at 258°C.

See entry DIFFERENTIAL THERMAL ANALYSIS

See other OXOSALTS OF NITROGENOUS BASES

3049. 1,12-Diazido-3,10-dioxa-5,8-dinitrazadodecane
[88487-88-7]

C₈H₁₆N₁₀O₆



See entry ENERGETIC COMPOUNDS

See other N-NITRO COMPOUNDS, ORGANIC AZIDES

†**3050. 2-Ethylhexanal**
[123-05-7]

C₈H₁₆O

BuCHEtCO.H

HCS 1980, 489

1. Steele, A. B. *et al.*, *Chem. Engrg.*, 1969, **66**, 160

2. Urben, P. G., private comm., 1989

The aldehyde ignited in air [1]. This is typical of mid-range aldehydes if sorbed on paper or cloth which increases surface exposure, ignition occurring within 2 hours [2].

See α -Pentylcinnamaldehyde

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

3051. O-O-tert-Butyl isopropyl monoperoxycarbonate
[2372-21-6]




$\text{C}_8\text{H}_{16}\text{O}_4$

See entry THERMAL EXPLOSIONS *See other* PEROXYCARBONATE ESTERS

3052. 3,6-Diethyl-3,6-dimethyl-1,2,4,5-tetraoxane
[33817-92-4]

$\text{C}_8\text{H}_{16}\text{O}_4$




1. Castrantas, 1965, 18
2. Swern, 1970, Vol. 1, 37

By analogy, this dimeric 2-butanone peroxide and the corresponding trimer probably contribute largely to the high shock-sensitivity of the commercial 'MEK peroxide' mixture, which contains these and other peroxides [1,2].

See other CYCLIC PEROXIDES, KETONE PEROXIDES

3053. Octylsodium
[2875-36-7]

$\text{C}_8\text{H}_{17}\text{Na}$



Houben-Weyl, 1970, Vol. 13.1, 384

The powder ignites in air.

See other ALKYL METALS

†**3054. 2,3-Dimethylhexane**
[584-94-1]



C_8H_{18}

†**3055. 2,4-Dimethylhexane**
[589-43-5]



C_8H_{18}

†3056. 3-Ethyl-2-methylpentane [609-26-7]	$\text{Me}_2\text{CHCHEt}_2$	C_8H_{18}
†3057. 2-Methylheptane [529-27-8]	$\text{Me}_2\text{CHCH}_2\text{Bu}$	C_8H_{18}
†3058. 3-Methylheptane [589-81-1]	EtCHMeBu	C_8H_{18}
†3059. Octane [111-65-9]	$\text{Me}[\text{CH}_2]_6\text{Me}$	C_8H_{18}
<i>HCS 1980, 696</i>		
†3060. 2,2,3-Trimethylpentane [564-02-3]	$\text{Me}_3\text{CCHMeEt}$	C_8H_{18}
†3061. 2,2,4-Trimethylpentane [540-84-1]	$\text{Me}_3\text{CCH}_2\text{CHMe}_2$	C_8H_{18}
<i>HCS 1980, 696</i>		
†3062. 2,3,3-Trimethylpentane [560-21-4]	$\text{Me}_2\text{CHCMe}_2\text{Et}$	C_8H_{18}
†3063. 2,3,4-Trimethylpentane [565-75-3]	$\text{Me}_2\text{CHCHMeCHMe}_2$	C_8H_{18}
3064. Diisobutylaluminium chloride [1779-25-5]	$(\text{Me}_2\text{CHCH}_2)_2\text{AlCl}$	$\text{C}_8\text{H}_{18}\text{AlCl}$

HCS 1980, 411

See entry ALKYLALUMINIUM HALIDES

3065. Chlorodibutylborane
[1730-69-4]



491M, 1975, 146

It ignites in air.

See other ALKYLHALOBORANES

3066. Di-*tert*-butyl chromate
[1189-85-1]



1. Anon., *ABCM Quart. Safety Summ.*, 1953, **24**, 2
2. Koenst, W. M. B. *et al.*, *Synth. Comm.*, 1980, **10**, 905–909

Large-scale preparation by addition of *tert*-butanol to chromium trioxide in a full unstirred flask with poor cooling detonated owing to local overheating. Effective cooling and stirring are essential [1]. It may safely be prepared by addition of a 40% aqueous solution of chromium trioxide to the alcohol [2].

Valencene

1. Hunter, G. L. K. *et al.*, *J. Food Sci.*, 1965, **30**, 876
2. Wilson, C. W. *et al.*, *J. Agric. Food Chem.*, 1978, **26**, 1430–1432

Oxidation of the sesquiterpene valencene to the 4-en-3-one proceeds explosively when at sub-ambient temperatures [1]. A safer alternative oxidation procedure is detailed [2].

See other OXIDANTS

3067. Di-*tert*-butylfluorophosphine
[29146-24-5]



Stelzer, O. *et al.*, *Inorg. Synth.*, 1978, **18**, 176

It may ignite in air.

See other ALKYLHALOPHOSPHINES

3068. Dibutylmagnesium
[1191-47-5]



491M, 1975, 147

It ignites in air.
See other ALKYL METALS


3069. Dibutyl hyponitrite
[86886-17-1]



$\text{C}_8\text{H}_{18}\text{N}_2\text{O}_2$

See entry DIALKYL HYPONITRITES

3070. *trans*-Di-*tert*-butyl hyponitrite
[14976-54-6]



$\text{C}_8\text{H}_{18}\text{N}_2\text{O}_2$

Mendenhall, G. D. *et al.*, *J. Amer. Chem. Soc.*, 1982, **104**, 5113
It is unstable and shock-sensitive, and needs careful handling.

Formamides

Protasiewicz, J. *et al.*, *J. Org. Chem.*, 1985, **50**, 3222

An equimolar mixture with *N-tert*-butylformamide underwent a sudden exotherm after placing in a bath at 60°C and the tube contents were ejected vigorously as a small cloud. A 1:2 mixture of the ester with formamide emitted a loud pop on heating in a bath at 70°C.

See other DIALKYL HYPONITRITES

†**3071. Dibutyl ether**
[142-96-1]



$\text{C}_8\text{H}_{18}\text{O}$

HCS 1980, 369

Dasler, W. *et al.*, *Ind. Eng. Chem. (Anal. Ed.)*, 1946, **18**, 52

Peroxides formed in storage are removed effectively by percolation through alumina.

See ETHERS, PEROXIDES IN SOLVENTS *See other* PEROXIDISABLE COMPOUNDS

3072. 1-(1,1-Dimethylethoxy)-2-methylpropane (*tert*-Butyl isobutyl ether)
[33021-02-2]



$\text{C}_8\text{H}_{18}\text{O}$


Bowen, R. D. *et al.*, *J. Chem. Soc. Perk. 2*, 1990, 147

Despite testing negative for peroxides, a sample exploded during distillation. Precautions are recommended.

See other ETHERS, PEROXIDISABLE COMPOUNDS

3073. 3,5-Dimethyl-3-hexanol
[4209-91-0]

C₈H₁₈O



Hydrogen peroxide, Sulfuric acid
See Hydrogen peroxide: 3,5-Dimethyl-3-hexanol, etc.

†**3074. Di-tert-butyl peroxide**
[110-05-4]

C₈H₁₈O₂



HCS 1980, 370

Preparative hazard

1. Griffiths, J. F. *et al.*, *Combust. Flame*, 1984, **56**, 135–148
2. Stull, 1977, 21
3. Sime, R. J., *Chem. Eng. News*, 1988, **66**(24), 4

Thermal decomposition is exothermic and self-ignition may result, especially if oxygen is present [1]. Though the heat of exothermic decomposition (1.32 kJ/g) is not exceptionally high, the weakness of the peroxide link tends to ready decomposition, and the energy release, coupled with the high volume of gaseous products (7.6 mol) at the adiabatic maximum of 550°C would give a 21-fold pressure increase in a closed vessel [2]. A routine student experiment on the gas-phase decomposition of the peroxide (0.2 ml) by injection into a 250 ml flask immersed in an oil bath at 155°C. The procedure was identical to a published method except that a mercury manometer was used to measure the pressure in place of a Bourdon gauge, and the procedure had been used on 30 occasions without incident. During a run at 165°C a violent explosion occurred [3].

See Hydrogen peroxide: *tert*-Butanol, etc.


See entry THERMAL EXPLOSIONS

See other DIALKYL PEROXIDES, SELF-HEATING AND IGNITION INCIDENTS

3075. 3,5-Dimethyl-3-hexyl hydroperoxide

[]

C₈H₁₈O₂



Preparative hazard

See Hydrogen peroxide: 3,5-Dimethyl-3-hexanol, etc.

See other ALKYL HYDROPEROXIDES

†3076. Bis(2-ethoxyethyl) ether
[112-36-7]



$\text{C}_8\text{H}_{18}\text{O}_3$

HCS 1980, 398

See other PEROXIDISABLE COMPOUNDS

†3077. 2-(2-Butoxyethoxy)ethanol (Diethyleneglycol monobutyl ether)
[112-34-5]




$\text{C}_8\text{H}_{18}\text{O}_3$

HCS 1980, 231

3078. Bis(2-hydroperoxy-2-butyl) peroxide (Dioxybis(1-methylpropyl hydroperoxide))
[126-76-1]

$\text{C}_8\text{H}_{18}\text{O}_6$



Leleu, *Cahiers*, 1973, (71), 238


The triperoxide, the main constituent of 'MEK peroxide', is explosive in the pure state, but insensitive to shock as the commercial 50% solution in dimethyl phthalate. The solution will explode at about 85°C, and slowly liberates oxygen at ambient temperature.

See other COMMERCIAL ORGANIC PEROXIDES, KETONE PEROXIDES

See related ALKYL HYDROPEROXIDES

3079. Diammonium Aquabis(peroxotartartrate)(2-)
[]

$\text{C}_8\text{H}_{18}\text{N}_2\text{O}_{19}\text{V}_2$



Schwendt, P, *et al.*, *Polyhedron*, 1998, **17**(13-14), 2161

This compound, which exists as a pentahydrate and for which a considerably weirder structure than shown has been inferred from X-ray crystallography on a homologue, may decompose explosively at room temperature. Handling more than 20 mg is discountenanced.

See related PEROXOMOLYBDATES AND TUNGSTATES, PEROXOCHROMIUM COMPOUNDS

3080. Dibutylzinc
[1119-90-0]



Leleu, *Cahiers*, 1977, (88), 371

It fumes in air, and may ignite under warm conditions.

See other DIALKYLZINCS, ALKYL METALS

3081. Diisobutylzinc
[1854-19-9]



Gibson, 1969, 180

It ignites in air.

See other DIALKYLZINCS, ALKYL METALS

3082. Diisobutylaluminium hydride
[1191-15-7]



1. Mirviss, S. B. *et al.*, *Ind. Eng. Chem.*, 1961, **53**(1), 54A

2. 'Specialty Reducing Agents', Brochure TA-2002/1, New York, Texas Alkyls, 1971

The higher thermal stability of dialkylaluminium hydrides over the corresponding trialkylaluminiums is particularly marked in this case with 2 branched alkyl groups [1]. Used industrially as a powerful reductant, it is supplied as a solution in hydrocarbon solvents. The undiluted material ignites in air unless diluted to below 25% concentration [2].

Tetrahydrofuran

Wissink, H. G., *Chem. Eng. News*, 1997, **75**(9), 6

The reagent can react with this solvent to give gaseous byproducts, capable of bursting glass containers. Initial complex formation is exothermic, so good cooling is needed to prevent decomposition when solutions are prepared, they should not be stored.

See other ALKYLALUMINIUM ALKOXIDES AND HYDRIDES, REDUCANTS

3083. Dibutylamine
[111-92-2]



$\text{C}_8\text{H}_{19}\text{N}$

HCS 1980, 367

Cellulose nitrate
See CELLULOSE NITRATE: amines

MRH 4.81/99+

Other reactants
Yoshida, 1980, 163
MRH values calculated for 14 combinations with oxidants are given.
See other ORGANIC BASES

†**3084. Di-2-butylamine**
[626-23-3]



$\text{C}_8\text{H}_{19}\text{N}$

See other ORGANIC BASES


†**3085. Diisobutylamine**
[110-96-3]



$\text{C}_8\text{H}_{19}\text{N}$

See other ORGANIC BASES

3086. Dibutyl hydrogen phosphite
[1809-19-4]




$\text{C}_8\text{H}_{19}\text{O}_3\text{P}$

Morrell, S. H., private comm., 1968
The phosphite was being distilled under reduced pressure. At the end of distillation the air-bleed was opened more fully, when spontaneous combustion occurred inside the flask, probably of phosphine formed by thermal decomposition.
See other PHOSPHORUS ESTERS

3087. Tetraethyldiarsane
[612-08-8]


$\text{C}_8\text{H}_{20}\text{As}_2$



Sidgwick, 1950, 770



It ignites in air.
See other ALKYLNON-METALS

3088. Tetraethylammonium perchlorate
[2567-83-1]




Udupa, M. R., *Propellants, Explos. Pyrotech.*, 1982, **7**, 155–157
It explodes at 298°C after an energetic phase transformation (10.4 kJ/mol) at 98°C.
See other QUATERNARY OXIDANTS, PERCHLORATE SALTS OF NITROGENOUS BASES

3089. 2,2'-Azobis(2-amidiniopropane) chloride
[2997-92-4]



Sodium peroxodisulfate
See 2,2'-Azobis(2-amidiniopropane) peroxodisulfate
See other AZO COMPOUNDS, POLYMERISATION INCIDENTS


3090. Tetraethylammonium periodate
[5492-69-3]



Preparative hazard
See Periodic acid: Tetraethylammonium hydroxide
See also TETRAMETHYLAMMONIUM PERIODATE
See other QUATERNARY OXIDANTS, OXOSALTS OF NITROGENOUS BASES

3091. 2,2'-Azobis(2-amidiniopropane) peroxodisulfate

[]


C₈H₂₀N₆O₈S₂Rausch, D. A., *Chem. Eng. News*, 1988, **66**(5), 2

When aqueous solutions of the polymerisation initiators 2,2'-azobis(2-amidiniopropane) chloride and sodium peroxodisulfate are mixed, the title compound separates as a water insoluble shock-sensitive salt. The shock-sensitivity increases as the moisture level decreases, and is comparable with that of lead azide. Stringent measures should be used to prevent contact of the solutions outside the polymerisation environment. (The instability derives from the high nitrogen (21.4%) and oxygen (31.6%) contents, and substantial oxygen balance, as well as the structural factors present in the salt.)

See other AZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS, OXOSALTS OF NITROGENOUS BASES

3092. 1,3,6,8-Tetraazatricyclo[6.2.1.1^{3,6}]dodecane tetranitrate

[10308-95-9]


C₈H₂₀N₈O₁₂

Violent decomposition occurred at 260°C.

See entry DIFFERENTIAL THERMAL ANALYSIS (DTA)*See other* OXOSALTS OF NITROGENOUS BASES

3093. Bis(trimethylphosphine)nickel(0)–acetylene complex
[98316-02-0]


C₈H₂₀NiP₂



Pörschke, K. R. *et al.*, *Z. Naturforsch. B*, 1985, **40B**, 199–209
The complex with acetylene (1 mol) decomposed explosively at 0°C.
See related COMPLEX ACETYLIDES, ALKYLPHOSPHINES

3094. Ethoxytriethylidiphosphinyl oxide
[]

C₈H₂₀O₂P₂




491M, 1975, 167

It ignites in air.
See related ALKYLPHOSPHINES

3095. Tetraethyllead
[78-00-2]

C₈H₂₀Pb



HCS 1980, 890

Doyle, F. P., *Loss Prev.*, 1969, **3**, 16
It is mildly endothermic (ΔH_f° (l) +217.5 kJ/mol, 0.56 kJ/g).

Failure to cover the residue with water after emptying a tank of the compound caused explosive decomposition after several days.

Other reactants



Yoshida, 1980, 234

MRH values calculated for 12 combinations with oxidants are given.

See other ALKYLMETALS, ENDOTHERMIC COMPOUNDS

3096. Tetrakis(ethylthio)uranium

[]




Bailar, 1973, Vol. 5, 416

It ignites in air.

See related METAL ALKOXIDES

3097. Tetraethyltin

[597-64-8]





Sorbe, 1968, 160

It tends to ignite in air.

See other ALKYL METALS

3098. 'Tetraethyldiborane'

[12081-54-8]




Köster, R. *et al.*, *Inorg. Synth.*, 1974, **15**, 141–146

The material (actually an equilibrium mixture of diborane and highly ethylated homologues) ignites in air. The propyl analogue behaves similarly.

See other ALKYL BORANES

3099. Bis(trimethylsilylmethyl)magnesium
[51329-17-0]




Oxygen, or Water

Andersen, R. A. *et al.*, *Inorg. Synth.*, 1979, **19**, 264

It is sensitive to oxygen and reacts violently with water.

See related ALKYL METALS, ALKYL SILANES

3100. 3,6,9-Triaza-11-aminoundecanol ('Hydroxyethyltriethylenetetramine')
[38361-85-2]




HCS 1980, 889

Cellulose nitrate

See CELLULOSE NITRATE: amines See other ORGANIC BASES

3101. 1,11-Diamino-3,6,9-triazaundecane ('Tetraethylenepentamine')
[112-57-2]



Carbon tetrachloride

See Carbon tetrachloride: 1,11-Diamino-3,6,9-triazaundecane

Cellulose nitrate

See CELLULOSE NITRATE: amines

See other ORGANIC BASES

3102. Bis(tetramethyldiphosphane disulfide)cadmium perchlorate
[71231-59-9]




McQuillan, G. P. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1979, 899

This and the nickel analogue are shock-sensitive and violently explosive. Other metal complexes (except cobalt) decompose slowly liberating the dimethyldiphosphane which tends to ignite when old sample bottles are opened.

See related AMMINOMETAL OXOSALTS

3103. Tetramethylammonium pentaperoxodichromate

[]





Mellor, 1943, Vol. 11, 358

The structure given is a 1960's vintage re-interpretation of the salt of CrO_5^- given by Mellor.

See Tetramethylammonium monoperoxochromate

3104. Tetrakis(dimethylamino)titanium

[]



Hydrazine

See Hydrazine: Titanium compounds

See other N-METAL DERIVATIVES

3105. 1,2-Diaminoethanebis(trimethylgold)

[]



Alone, or Nitric acid



Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 550

The solid complex is very sensitive to light and explodes violently on heating in an open crucible. A drop of conc. nitric acid added to the dry compound causes explosion.

See related ALKYL METALS See other GOLD COMPOUNDS

3106. Bis(diethylenetriamine)cobalt(III) perchlorate

[28872-17-5]



It explodes at 325°C; high impact-sensitivity.

See entry AMMINEMETAL OXOSALTS

3107. Potassium graphite

[12081-88-8]

(Complex Structure)



Water

Bailar, 1973, Vol. 1, 1276


It may explode with water.

See Carbon: Alkali metals

See other METAL NON-METALLIDES

3108. Oligo(octacarbonyl) (Cyclo-oligo-butadiynyl-2,3-dioxocyclobutenylidene)

[123002-86-8] $n = 3$; [123002-87-9] $n = 4$; [] $n = 5$



Diederich, F. *et al.*, *J. Amer. Chem. Soc.*, 1991, **113**(2), 495

These compounds exploded violently on heating above 85°C in an attempt to obtain melting points.

See 11,12-Diethynyl-1,4,7,10-tetraoxadispiro[4.0.4.2]dodec-11-ene

See other ACETYLENIC COMPOUNDS

3109. Lithium octacarbonylnickelate

[81181-41-5] (ion)



Bailar, 1973, Vol. 3, 1117

A pyrophoric salt.

See related CARBONYLMETALS

3110. Rubidium graphite

[12193-29-2]

(Complex Structure)



See Carbon: Alkali metals

See other METAL NON-METALLIDES

3111. Nonacarbonyldiiron
[20982-74-5]


C₉Fe₂O₉



Anon., *ABCM Quart. Safety Summ.*, 1943, **14**, 18
Commercial iron carbonyl (fl.p.35°C) has an autoignition temperature in contact with brass of 93°C, lower than that of carbon disulfide.
See other CARBONYLMETALS

3112. 2-Nonen-4,6,8-triyn-1-ol
[]


C₉H₄O



Bohlman, F. *et al.*, *Chem. Ber.*, 1963, **96**, 2586
Extremely unstable, explodes after a few minutes at ambient temperature and ignites at 110°C.
See other ACETYLENIC COMPOUNDS


3113. 2-Nitrophenylpropionic acid
[530-85-8]

C₉H₅NO₄



Sorbe, 1968, 152
It explodes above 150°C.
See other ACETYLENIC COMPOUNDS, NITROARYL COMPOUNDS, ORGANIC ACIDS

3114. Sodium 3-phenylisoxazolin-4,5-dione-4-oximate
[70247-49-3]



Wentrup, C. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 467



It explodes on rapid heating, like the 5-methyl analogue and the silver salt.

See Fulminic acid (reference 2)

See other N-O COMPOUNDS

See related OXIMES

3115. 3,5-Dimethyl-4-[I,I-bis(trifluoroacetoxy)iodo]isoxazole
[57508-69-7]




Preparative hazard

See Peroxytrifluoroacetic acid: 4-Iodo-3,5-dimethylisoxazole

See [I,I-Bis(trifluoroacetoxy)iodo]benzene

See other IODINE COMPOUNDS

3116. 1-Diazoindene
[35847-40-6]



Quinn, S. *et al.*, *Inorg. Chim. Acta*, 1981, **50**, 141–146


The preparation of this is hazardous, and it is recommended that diazoindene always be kept in solution.

See 4-Toluenesulfonyl azide

See other DIAZO COMPOUNDS

3117. 2,4-Diisocyanatotoluene
[584-84-9]

C₉H₆N₂O₂



(MCA SD-73, 1971); NSC 489, 1977; FPA H5, 1972; HCS 1980, 907

Acyl chlorides, or Bases

MCA SD-73, 1971

The diisocyanate may undergo exothermic polymerisation in contact with bases or more than traces of acyl chlorides, sometimes used as stabilisers.

Other reactants

Yoshida, 1980, 259

MRH values calculated for 13 combinations with oxidants are given.

Water


Kirk-Othmer, 1981, Vol. 13, 812

Polythene containers holding the diisocyanate may harden and burst in prolonged storage, because of slow absorption of water vapour through the wall leading to urea deposition from hydrolysis and generation of pressure of liberated carbon dioxide.

See other ORGANIC ISOCYANATES

3118. 3-Diazo-5-phenyl-3H-pyrazole
[62072-18-8]

C₉H₆N₄




The light-sensitive crystalline solid is shock-sensitive and explodes at 115°C.

See entry DIAZOAZOLES

See other DIAZO COMPOUNDS, HIGH-NITROGEN COMPOUNDS

3119. 5-(4-Diazoniobenzenesulfonamido)thiazole tetrafluoroborate
[41334-40-1] (ion)

C₉H₇BF₄N₄O₂S




Zuber, F. *et al.*, *Helv. Chim. Acta*, 1950, **33**, 1269–1271

It explodes at 135–137°C.

See entry DIAZONIUM TETRAHALOBORATES

3120. 3-Iodo-1-phenylpropyne
[73513-15-2]

C₉H₇I




Whiting, M. C., *Chem. Eng. News*, 1972, **50**(23), 86

It detonated on distillation at around 180°C.
See other HALOACETYLENE DERIVATIVES

3121. Quinoline
[91-22-5]

C₉H₇N



HCS 1980, 805

1. Blumann, A., *Proc. R. Aust. Chem. Inst.*, 1964, **31**, 286
2. *MCA Case History No. 1008*

The traditional unpredictably violent nature of the Skraup reaction (preparation of quinoline and derivatives by treating anilines with glycerol, sulfuric acid and an oxidant, usually nitrobenzene) is attributed to lack of stirring and adequate temperature control in many published descriptions [1]. A reaction on 450 l scale, in which sulfuric acid was added to a stirred mixture of aniline, glycerol, nitrobenzene, ferrous sulfate and water, went out of control soon after the addition. A 150 mm rupture disk blew out first, followed by the manhole cover of the vessel. The violent reaction was attributed to doubling the scale of the reaction, an unusually high ambient temperature (reaction contents at 32°C) and the accidental addition of excess acid. Experiment showed that a critical temperature of 120°C was attained immediately on addition of excess acid under these conditions [2].

Dinitrogen tetraoxide

See Dinitrogen tetraoxide: Heterocyclic bases

Hydrogen peroxide

See Hydrogen peroxide: Organic compounds (reference 2)

Linseed oil, Thionyl chloride

See Sulfinyl chloride: Linseed oil, etc.


Maleic anhydride

See Maleic anhydride: Bases, etc.

See other ORGANIC BASES

3122. 3-Phenyl-5-isoxazolone
[7713-79-3]

C₉H₇NO₂



Sealed samples decompose exothermally above 105°C.
See entry ISOXAZOLES
See other N–O COMPOUNDS

3123. Dicarbonyl- π -cycloheptatrienyltungsten azide
[59589-13-8]


C₉H₇N₃O₂W



Hoch, G. *et al.*, *Z. Naturforsch. B*, 1976, **31B**, 295
It was insensitive to shock, but decomposed explosively at 130°C.
See related CARBONYLMETALS, METAL AZIDES, ORGANOMETALLICS

3124. 3-Methoxy-2-nitrobenzoyldiazomethane
[24115-83-1]

C₉H₇N₃O₄




Alone, or Sulfuric acid

Musajo, L. *et al.*, *Gazz. Chim. Ital.*, 1950, **80**, 171–176
The diazoketone explodes at 138–140°C, or on treatment with conc. sulfuric acid.
See other DIAZO COMPOUNDS, NITROARYL COMPOUNDS

3125. Tetracyclo[4.3.0^{3,5}.0^{4,6}]nona-1,7-diene (2,4-Dihydro-1H-1,2,6a-methanopentalene)
[85861-34-3]

C₉H₈




Burger, U. *et al.*, *Helv. Chim. Acta*, 1983, **66**, 60–67

The highly strained hydrocarbon, produced from dilithiopentalene and chlorocarbene, explodes violently at temperatures as low as -40°C when concentrated. Other isomers are probably similar.

See other DIENES, STRAINED-RING COMPOUNDS

3126. N-Chlorocinnamaldimin (3-Chlorimino-1-phenylpropene)
[]

C₉H₈ClN



Hauser, C. R. *et al.*, *J. Amer. Chem. Soc.*, 1935, **57**, 570


It decomposes vigorously after storage at ambient temperature for 30 m.

See other N-HALOGEN COMPOUNDS

See related DIENES

3127. Spiro(homocubane-9,9'-diazirine) (Spiro[3H-diazirine-3,9'-pentacyclo[4,3,0,0^{2,5},3,8,0^{4,7}]nonane])
[129215-84-5]

C₉H₈N₂




Platz, M. S. *et al.*, *J. Amer. Chem. Soc.*, 1991, **113**, 4991

A white, dangerously explosive solid, apparently storable at room temperature but decomposing before melting.

See other DIAZIRINES, STRAINED-RING COMPOUNDS

3128. 3,5-Dimethylbenzenediazonium-2-carboxylate
[68596-88-3]

C₉H₈N₂O₂



Wilcox, C. F. *et al.*, *J. Amer. Chem. Soc.*, 1984, **106**, 7196

It is more sensitive to thermal decomposition than other diazonium carboxylates, and will decompose readily when inadequate stirring, large scale or excessively rapid diazotisation produces too large a temperature rise. There is significant potential for violent eruption of suspensions, or detonation of the filtered solid, and all recommended precautions should be used.


See 2-Carboxy-3,6-dimethylbenzenediazonium chloride

See other DIAZONIUM CARBOXYLATES

3129. 4,6-Dimethylbenzenediazonium-2-carboxylate

[]

C₉H₈N₂O₂



Gompper, R. *et al.*, *Chem. Ber.*, 1968, **101**, 2348


It is a highly explosive solid.

See other DIAZONIUM CARBOXYLATES

3130. 2,3-Epoxypropionaldehyde 2,4-dinitrophenylhydrazone

[]

C₉H₈N₄O₅



Payne, G. B., *J. Amer. Chem. Soc.*, 1959, **81**, 4903

Although it melts at 96–98°C in a capillary, 0.5 g in a test tube decomposed explosively at 100°C.


See 2,3-Epoxypropionaldehyde oxime

See other 1,2-EPOXIDES, POLYNITROARYL COMPOUNDS See related HYDRAZONES

3131. 3-(2-Nitrophenyl)-2-hydroxyiminopropanoic acid (*o*-Nitrophenylpyruvic acid oxime)

[27878-36-0]

C₉H₈N₂O₅



CHETAH, 1990, 187


Shock sensitive

See other NITROARYL COMPOUNDS, ORGANIC ACIDS, OXIMES

3132. Tetrakis(2,2,2-trinitroethyl)orthocarbonate

[14548-58-4]


C₉H₈N₁₂O₂₈



See entry TRINITROETHYL ORTHOESTERS

3133. Benzyloxyacetylene
[40089-12-1]


C₉H₈O



Olsman, H. *et al.*, *Rec. Trav. Chim.*, 1964, **83**, 305
If heated above 60°C during vacuum distillation, explosive rearrangement occurs.
See other ACETYLENIC COMPOUNDS, BENZYL COMPOUNDS

3134. Cinnamaldehyde (3-Phenylpropenal)
[104-55-2]

C₉H₈O



HCS 1980, 316

Sodium hydroxide


Morrell, S. H., private comm., 1968

Rags soaked in sodium hydroxide and in the aldehyde overheated and ignited owing to aerobic oxidation when they came into contact in a waste bin.

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS, SELF-HEATING AND IGNITION INCIDENTS

3135. 2-Methyl-3,5,7-octatriyn-2-ol
[]

C₉H₈O




Cook, C. J. *et al.*, *J. Chem. Soc.*, 1952, 2885, 2890

The crude material invariably deflagrated at ambient temperature, and once during drying at 0°C/0.013 mbar.

See other ACETYLENIC COMPOUNDS

3136. 4-Hydroxy-*trans*-cinnamic acid
[7400-08-0]

C₉H₈O₃



Tanaka, Y. *et al.*, *Proc. 4th Int. Conf. High Press.* (1974), 704, 1975


During spontaneous solid-state polymerisation at 325°C/8–10 kbar, the acid exploded violently.

See HIGH PRESSURE REACTION TECHNIQUES

See other ORGANIC ACIDS, POLYMERISATION INCIDENTS

3137. *O*-Acetylsalicylic acid
[50-78-2]

C₉H₈O₄




For experimental investigation of aspirin powder explosions,

See entry DUST EXPLOSION INCIDENTS (reference 18)

See other ORGANIC ACIDS

3138. 2-Carboxy-3,6-dimethylbenzenediazonium chloride
[36794-93-1]

C₈H₉ClN₂O₂



Hart, H. *et al.*, *J. Org. Chem.*, 1972, **37**, 4272


The compound (the hydrochloride of the internal carboxylate salt) appears to be stable for considerable periods at ambient temperature, but explodes on melting at 88°C.

See related DIAZONIUM CARBOXYLATES

See other ORGANIC ACIDS

3139. 2-Phenylethyl isocyanate
[14649-03-7]

C₉H₉NO




Energy of decomposition (in range 220–350°C) measured as 0.50 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ORGANIC ISOCYANATES

3140. *miso*-Nitroindane
[34701-14-9, 7436-07-9] (4- and 5-isomers, resp.)

C₉H₉NO₂



1. Lindner, J. *et al.*, *Ber.*, 1927, **60**, 435


2. Gribble, G. W., *Chem. Eng. News*, 1973, **51**(6), 30, 39

The crude mixture of 4- and 5-nitroindanes produced by mixed acid nitration of indane following a literature method [1] is hazardous to purify by distillation. The warm residue from distillation of 15 mmol at 80°C/1.3 mbar exploded on admission of air, and a 1.3 mol batch exploded as distillation began under the same conditions. Removal of higher-boiling poly-nitrated material before distillation is recommended.

See other NITROARYL COMPOUNDS

3141. 3-Nitropropiophenone
[17408-16-1]

C₉H₉NO₃




Preparative hazard

See Nitric acid: Propiophenone, etc.

See other NITROARYL COMPOUNDS

3142. 4-Nitrophenylpropan-2-one
[5332-96-7]

C₉H₉NO₃




Preparative Hazard

4-Nitrophenylacetic acid. *See other* NITROARYL COMPOUNDS

3143. 2-Isocyanoethyl benzenesulfonate
[57678-14-5]

C₉H₉NO₃S



Matteson, D. S. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 3761


Heating the product under vacuum to remove pyridine solvent caused a moderately forceful explosion.

See other ORGANIC ISOCYANATES, SULFUR ESTERS

3144. 3-Phenylpropionyl azide

[]

C₉H₉N₃O



Curtiss, T. *et al.*, *J. Prakt. Chem.*, 1901, **64**, 297


A sample exploded on a hot water bath.

See other ACYL AZIDES

3145. 1,3,5-Tris(nitromethyl)benzene

[]

C₉H₉N₃O₆




See Nitric acid: Hydrocarbons (reference 8)

See related POLYNITROALKYL COMPOUNDS, POLYNITROARYL COMPOUNDS

3146. 4-(2-Ammonio-2-carboxyethylthio)-5,7-dinitro-4,5-dihydrobenzofurazanide N-oxide (4,6-Dinitrobenzofurazan N-oxide.cysteine complex)

[79263-47-1]

C₉H₉N₅O₈S




Strauss, M. J. *et al.*, *Tetrahedron Lett.*, 1981, **22**, 1945–1948

When thoroughly dry, the cysteine complex of 5,7-dinitrobenzofurazan *N*-oxide exploded on heating.

See other FURAZAN N-OXIDES

3147. 2-Chloro-1-nitroso-2-phenylpropane
[6866-10-0]

C₉H₁₀ClNO




MCA Case History No. 747

A sample of the air-dried material (which is probably the dimer) decomposed vigorously on keeping in a closed bottle at ambient temperature overnight.

See other NITROSO COMPOUNDS

3148. 1,3-Dimethyl-2,1-benzisoxazolium perchlorate
[63609-41-6]

C₉H₁₀ClNO₅




Haley, N. F., *J. Org. Chem.*, 1978, **43**, 1236

The salt, m.p. 152°, explodes at 154°C.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

3149. Methoxy-1,3,5,7-cyclooctatetraene
[7176-89-8]

C₉H₁₀O



Oxygen

Adam, W. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 2837–2840


The evaporated residue from sensitised photochemical oxidation of the polyene ignited spontaneously, on several occasions explosions occurring.

See related DIENES

See other PEROXIDISABLE COMPOUNDS, SELF-HEATING AND IGNITION INCIDENTS

3150. 3-Phenoxy-1,2-epoxypropane (Glycidyl phenyl ether)
[122-60-1]

C₉H₁₀O




Energy of decomposition (in range 360–450°C) measured as 0.626 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

3151. Propiophenone
[93-55-0]

C₉H₁₀O




Nitric acid, Sulfuric acid

See Nitric acid: Propiophenone, etc.

3152. 3,5-Dimethylbenzoic acid
[499-06-9]

C₉H₁₀O₂



Preparative hazard


See Nitric acid: Hydrocarbons (reference 8)

See other ORGANIC ACIDS

3153. 1-(*cis*-Methoxyvinyl)-1,4-endoperoxy-2,5-cyclohexadiene (*Z*-1-(2-Methoxyethenyl)-2,3-dioxabicyclo[2.2.2]octa-5,7-diene)

[]

C₉H₁₀O₃




Matsumoto, M. *et al.*, *Tetrahedron. Lett.*, 1979, 1607–1610

The product of sensitised photooxidation of *cis*-2-methoxystyrene is explosive.

See other CYCLIC PEROXIDES

3154. 3- or 4-Methoxy-5,6-benzo-6*H*-1,2-dioxin
[69681-97-6, 69681-98-7, resp.]

C₉H₁₀O₃




Lerdal, D. *et al.*, *Tetrahedron. Lett.*, 1978, 3227–3228

Both isomers decomposed violently on warming in absence of solvent.

See other CYCLIC PEROXIDES

3155. Allyl benzenesulfonate
[7575-57-7]

C₉H₁₀O₃S



Dye, W. T. *et al.*, *Chem. Eng. News*, 1950, **28**, 3452


The residue from vacuum distillation at 92–135°C/2.6 mbar darkened, thickened, then exploded after removal of the heat source.

For precautions, see Triallyl phosphate

See other ALLYL COMPOUNDS, SULFUR ESTERS

3156. 2-(4-Bromophenylazo)-2-propyl hydroperoxide
[72447-41-7]


C₉H₁₁BrN₂O₂



See entry α -PHENYLAZO HYDROPEROXIDES (reference 4)
See related ALKYL HYDROPEROXIDES, AZO COMPOUNDS

3157. 4-Nitroisopropylbenzene
[1817-47-6]


C₉H₁₁NO₂



High rate of pressure increase during exothermal decomposition.
See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION
See other NITROARYL COMPOUNDS

3158. Nitromesitylene
[603-71-4]


C₉H₁₁NO₂



Preparative hazard
See Nitric acid: Hydrocarbons (references 7,8)
See other NITROARYL COMPOUNDS

3159. 2-Azido-2-phenylpropane
[32366-26-0]


C₉H₁₁N₃



Energy of exothermic decomposition in range 180–270°C was measured as 1.2 kJ/g
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other ORGANIC AZIDES

3160. 4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2,6}]undeca-8-ene-3,5-dione (5,8-Dihydro-2-methyl-5,8-ethano-1H-[1,2,4]triazolo[1,2a]pyridazine-1,3(2H)dione)
[78790-57-5]

C₉H₁₁N₃O₂




Hydrogen peroxide, Potassium hydroxide

See Hydrogen peroxide: 4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2,6}]undeca-8-ene-3,5-dione

3161. 5-(4-Dimethylaminobenzeneazo)tetrazole
[53004-03-8]

C₉H₁₁N₇




Thiele, J., *Ann.*, 1892, **270**, 61

It explodes at 155°C.
See other AZO COMPOUNDS, TETRAZOLES

3162. Mesitylene (1,3,5-Trimethylbenzene)
[108-67-3]

C₉H₁₂




HCS 1980, 621

Nitric acid
See Nitric acid: Hydrocarbons (reference 8)

3163. 2-Propylpyridine-4-carbothioamide (2-Propyl(thioisonicotinamide))
[14222-60-7]

C₉H₁₂N₂S



See FRICTIONAL INITIATION INCIDENTS (REFERENCE 3)

3164. 2-Methoxyethylbenzene
[3558-60-9]


C₉H₁₂O

C₆H₅CH₂CH₂OMe

Nitric acid
See Nitric acid: 2-Methoxyethylbenzene, Sulfuric acid

3165. 3-Phenylpropanol
[122-97-4]


C₉H₁₂O



Phosphorus tribromide
See Phosphorus tribromide: 3-Phenylpropanol

3166. 2-Phenyl-2-propyl hydroperoxide (Cumyl hydroperoxide)
[80-15-9]

C₉H₁₂O₂



1. Leroux, A., *Mém. Poudres*, 1955, **37**, 49
2. Simon, A. H. *et al.*, *Chem. Ber.*, 1957, **90**, 1024
3. Hulanicki, A., *Chem. Anal. (Warsaw)*, 1973, **18**, 723–726
4. *MCA Case History No. 906*
5. Leleu, *Cahiers*, 1973, (71), 226
6. Fleming, J. B. *et al.*, *Hydrocarbon Process.*, 1976, **55**(1), 185–196
7. Redoshkin, B. A. *et al.*, *Chem. Abs.*, 1964, **60**, 14359h
8. Antonovskii, V. L. *et al.*, *Chem. Abs.*, 1964, **60**, 14360c
9. Kletz, T. A., *Loss Prev.*, 1979, **12**, 98
10. Anon., *Loss Prev. Bull.*, 1986, (071), 9
11. Schwab, R. F., *Loss. Prev. Bull.*, 1988, (083), 20–32
12. Anon., *Loss Prev. Bull.*, 1992, **103**, 31
13. Francisco, M. A., *Chem. Eng. News*, 1993, **71**(22), 4
14. Shanley, E. S., *Chem. Eng. News*, 1993, **71**(35), 61

The explosibility of this unusually stable hydroperoxide has been investigated [1]. It is difficult, but not impossible, to induce explosive decomposition [2]. A colorimetric method of analysing mixtures of cumene and its hydroperoxide which involves alkaline decomposition of the latter is safer than the use of acidic conditions, or of reductants, which may cause explosive reactions [3]. During vacuum concentration of the hydroperoxide by evaporation of cumene, a 6.5 m³ quantity of the 36% material decomposed explosively after storage at 109°C for 5 h. Catalytic decomposition under near-adiabatic conditions may have been involved [4]. At the end of concentration from hexane, after purification, 100 ml of peroxide exploded violently for no apparent reason; isolation of no more than 5 g is recommended [13]. Contact with copper, copper or lead alloys, mineral acids or reductants may lead to violent decomposition [5]. Cobalt also catalyses the decomposition reaction which is autocatalytic. The acid cleavage reaction used industrially is highly exothermic, and safety aspects of commercial phenol/acetone processes are discussed fully [6]. Much detailed kinetic work has also been published [7,8]. Five explosions due to overheating and subsequent decomposition of the concentrated solutions in cumene have been noted. Another explosion arose from reduced flow of acid gas used in the cleavage reaction because the cylinder was outdoors on a cold night. The reduced flow led to spurious control signals which eventually caused the explosion [9]. During a shut-down, a 100 t quantity of hydroperoxide solution in intermediate storage was being kept hot

with steam but became overheated, causing cumene to vent. The vapour ignited explosively and ruptured the storage tank [10]. A detailed analysis of the incident is available [11]. A drum of the hydroperoxide exploded, this was attributed to possible contamination with sodium hydroxymethylsulfonate. This might form catalytic acids if oxidised, but the drum had been sampled with a, possibly corroded, stainless steel dip pipe [12].

CHETAH calculations for energy/hazard of an hypothetical decomposition reaction to carbon, water and methane are reported [14].

See THERMAL EXPLOSIONS, COMPUTATION OF REACTIVE CHEMICAL HAZARDS

See other CATALYTIC IMPURITY INCIDENTS, CORROSION INCIDENTS

Charcoal

Leleu, *Cahiers*, 1980, (99), 278

Contact with charcoal powder at ambient temperature gives a strong exotherm (probably owing to traces of heavy metals present).

1,2-Dibromo-1,2-diisocyanatoethane polymers

Lapshin, N. M. *et al.*, *Chem. Abs.*, 1974, **81**, 64013

A mixture of the dimeric and trimeric dibromo compounds with the hydroperoxide in benzene may react vigorously or explode if the solution is heated to concentrate it.

Iodides

Duswalt, A. A., *Chem. Eng. News*, 1990, **68**(6), 2

Following a laboratory incident, cumene hydroperoxide was shown to be explosively decomposed by catalytic sodium iodide.


Other reactants

Yoshida, 1980, 105

MRH values calculated for 15 combinations with oxidisable materials are given.

See other ALKYL HYDROPEROXIDES, COMMERCIAL ORGANIC PEROXIDES

3167. 2,3,4,6-Tetramethylpyrilium perchlorate [35941-38-9]




Balaban, A. T. *et al.*, *Org. Prep. Proced. Int.*, 1982, **14**, 31–38

It must be handled cautiously.

See other NON-METAL PERCHLORATES

3168. Benzyltrimethylamine
[28262-13-7]

C₉H₁₃N




Cellulose nitrate

See CELLULOSE NITRATE: amines See *other* BENZYL COMPOUNDS, ORGANIC BASES

3169. 2,4,6-Trimethylaniline
[88-05-1]

C₉H₁₃N




Nitrosyl perchlorate

See Nitrosyl perchlorate: Organic materials

3170. O-Mesitylenesulfonylhydroxylamine
[36016-40-7]

C₉H₁₃NO₃S



1. Tamura, Y. *et al.*, *Tetrahedron Lett.*, 1972, **40**, 4133
2. Ning, R. Y., *Chem. Eng. News*, 1973, **51**(51), 36-37
3. Carpino, L. A., *J. Amer. Chem. Soc.*, 1960, **82**, 3134
4. Ning, R. Y., private comm., 1974

5. Johnson, C. R. *et al.*, *J. Org. Chem.*, **39**, 2459, footnote 15
6. Scopes, D. I. C. *et al.*, *J. Org. Chem.*, 1977, **42**, 376
7. Dillard, R. D. *et al.*, *J. Med. Chem.*, 1980, **23**, 718
8. Fluka advertisement, *J. Org. Chem.*, 1980, **45**(9), cover ii


The title compound was prepared following a published procedure [1] and a dried sample decomposed soon after putting it into an amber bottle for storage, the screw cap being shattered [2]. Although the instability of the compound had been mentioned [3], no suggestion of violent decomposition had previously been made. It seems likely that traces of surface alkali in the soda-glass bottle had catalysed both the formation of the highly active imidogen radical (HN:) from the base-labile compound, and its subsequent exothermic decomposition. Storage in dichloromethane solution appears safe [4]. The small crystals produced by a modified method appear to be safe in storage at 0°C, or in use [5]. Attempted vacuum drying of the compound at ambient temperature led to a mild explosion. Subsequently, a solution of the wet solid in dimethoxyethane was dried over molecular sieve and used in further work [6]. The solid may explode violently when heated to 60°C. Large amounts should be used either in solution or as the crystalline solid. The neat molten liquid form should be avoided, as it evolves heat on crystallisation. It is best to avoid storage and generate the reagent as needed [7]. A stable precursor, *tert*-butyl *N*-mesitylenesulfonyloxycarbamate has become available [8].

See *O*-4-Toluenesulfonylhydroxylamine

See other CATALYTIC IMPURITY INCIDENTS, GLASS INCIDENTS, N-O COMPOUNDS

3171. 3-Ethyl-1(4-methylphenyl)triazene
[50707-40-9]

C₉H₁₃N₃




Anon., *Loss Prev. Bull.*, 1980, (030), 160

A 5 g sample was distilled under vacuum uneventfully from a bulb tube with air-bath heating, but a 50 g portion exploded violently when distilled at 0.13 mbar from an oil-bath at 120°C.

See other TRIAZENES

3172. 2-(2-Aminoethylamino)-5-methoxynitrobenzene
[13556-31-5]


C₉H₁₃N₃O₃



See *N*-(2-Nitrophenyl)-1,2-diaminoethane
See other NITROARYL COMPOUNDS

3173. 3,3,6,6-Tetrakis(bromomethyl)-9,9-dimethyl-1,2,4,5,7,8-hexoxonane
[16007-16-2]

C₉H₁₄Br₄O₆




Schulz, M. *et al.*, *Chem. Ber.*, 1967, **100**, 2245

It explodes on impact or friction, as do the tetrachloro- and 9-ethyl-9-methyl analogues.

See other CYCLIC PEROXIDES

3174. 1-(4-Methyl-1,3-diselenonylidene)piperidinium perchlorate
[53808-72-3]

C₉H₁₄ClNO₄Se₂




See entry 1-(1,3-DISELENONYLIDENE)PIPERIDINIUM PERCHLORATES

See other PERCHLORATE SALTS OF NITROGENOUS BASES

3175. 6,6-Dimethylbicyclo[3.1.1]heptan-2-one (Nopinone)
[24903-95-5]; S-[7782-63-9]

C₉H₁₄O



Preparative hazard


1. Gordon, P. M., *Chem. Eng. News*, 1990, **68**(30), 2
2. Ferreira, J. T. B., *ibid.*, (50), 2
3. Weissman, S., *ibid.*, 1991, **69**(4), 2

Several reports have been received of explosion during work-up of the products obtained from ozonolysis of pinene, despite prior treatment with reductants. A safe alternative procedure is given in [3].

See OZONIDES

3176. 2,6-Dimethyl-2,5-heptadien-4-one diozonide
[]

C₉H₁₄O₇




Harries, G. *et al.*, *Ann.*, 1910, **374**, 338

'Phorone' diozonide ignites on warming to ambient temperature.

See other OZONIDES

3177. Tri-2-propenylborane (Triallylborane)
[688-61-9]

C₉H₁₅B




Borane

Alder, R. W. *et al.*, *J. Chem. Soc., Perk. I*, 1996, (7), 657

The 1:1 adduct is highly pyrophoric, so too is its pyrolysis product. Both are polymeric, diborabicycloundecane is not.

See other ALKYLBORANES

3178. N-Cyano-2-bromoethylcyclohexylamine
[53182-16-4]






Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 23

The reaction product from *N*-cyclohexylaziridine and cyanogen bromide (believed to be the title compound) exploded violently on attempted distillation at 160°C/0.5 mbar.

See *N*-Cyano-2-bromoethylbutylamine

See other CYANO COMPOUNDS

3179. Tris(ethylthio)cyclopropenium perchlorate
[50744-08-6]




Sunderlin, K. G. R., *Chem. Eng. News*, 1974, **52**(31), 3

Some 50 g of the compound had been prepared by a method used for analogous compounds, and the solvent-free oil had been left to crystallise. Some hours later it exploded with considerable violence. Preparation of this and related compounds by other workers had been uneventful.

See other NON-METAL PERCHLORATES

3180. Triallylchromium
[12082-46-1]



O'Brien, S. *et al.*, *Inorg. Chem.*, 1972, **13**, 79


Triallylchromium and its thermal decomposition products are pyrophoric in air.

See related ALKYL METALS

See other ALLYL COMPOUNDS

3181. Nitrilotris(oxiranemethane) (Triglycidylamine)
[61014-23-1]

C₉H₁₅NO₃




McKelvey, J. *et al.*, *J. Org. Chem.*, 1960, **25**, 1424

The crude product polymerises violently during vacuum distillation. The pure is storable cool

See other 1,2-EPOXIDES

3182. 4-Tolylbiguanidium hydrogen dichromate
[15760-46-0]


C₉H₁₅Cr₂N₅O₇



See entry DICHROMATE SALTS OF NITROGENOUS BASES

3183. 1,3,5-Triacetylhexahydro-1,3,5-triazine
[26028-46-6]

C₉H₁₅N₃O₃




Nitric acid, Trifluoroacetic anhydride

See Nitric acid: 1,3,5-Triacetylhexahydro-1,3,5-triazine, etc.

3184. Triallyl phosphate
[1624-19-4]

C₉H₁₅O₄P



1. Dye, W. T. *et al.*, *Chem. Eng. News*, 1950, **28**, 3452

2. Steinberg, G. M., *Chem. Eng. News*, 1950, **28**, 3755

Alkali-washed material, stabilised with 0.25% of pyrogallol, was distilled at 103°C/4 mbar until slight decomposition began. The heating mantle was then removed and the still-pot temperature had fallen below its maximum value of 135°C when the residue exploded violently [1]. The presence of solid alkali [2] or 5% of phenolic inhibitor is recommended, together with low-temperature high-vacuum distillation, to avoid formation of acidic decomposition products, which catalyse rapid exothermic polymerisation.

See other ALLYL COMPOUNDS, CATALYTIC IMPURITY INCIDENTS, PHOSPHORUS ESTERS

3185. 1,4,7-Triazacyclononanetricarbonylmolybdenum hydride perchlorate

[]

[C₆H₁₅N₃MoH(CO)₃] ClO₄

C₉H₁₆ClMoN₃O₇

See 1,4,7-Triazacyclononanetricarbonyltungsten hydride perchlorate (next below)

See related AMMINOMETAL OXOSALTS, CARBONYLMETALS

3186. 1,4,7-Triazacyclononanetricarbonyltungsten hydride perchlorate

[88253-43-4]

[C₆H₁₅N₃WH(CO)₃] ClO₄

C₉H₁₆ClN₃O₇W


Chaudhuri, P. *et al.*, *Inorg. Chem.*, 1984, **23**, 429

The dry tungsten and molybdenum complexes are extremely explosive when dry; handle no more than 10 mg portions. The corresponding Mo complexes with Br or I replacing H are also explosive.

See related AMMINOMETAL OXOSALTS, CARBONYLMETALS

3187. Azelaic acid
[123-99-9]

C₉H₁₆O₄



Hydrogen peroxide, Sulfuric acid
See Hydrogen peroxide: Azelaic acid, Sulfuric acid
See other ORGANIC ACIDS

3188. Diisopropyl malonate
[13196-64-7]


C₉H₁₆O₄



Sulfinyl chloride
See Sulfinyl chloride: Diisopropyl malonate

3189. Diperoxyazelaic acid
[1941-79-3]


C₉H₁₆O₆



Preparative hazard
See Hydrogen peroxide: Azelaic acid, Sulfuric acid
See other PEROXYACIDS

†**3190. 2,6-Dimethyl-3-heptene**
[2738-18-3]


C₉H₁₈



See other ALKENES


†**3191. 1,3,5-Trimethylcyclohexane**
[1795-27-3]

C₉H₁₈



1051

3192. 1,4-Bis(2-chloroethyl)-1,4-bis(azonia)bicyclo[2.2.1]heptane periodate
 [77628-04-7] **C₉H₁₈Cl₂I₂N₂O₈**



Pettit, G. R. *et al.*, *Can. J. Chem.*, 1981, **59**, 218, 220
 It exploded violently at about 150°C, or when ignited openly or on gentle impact.
See related PERCHLORATE SALTS OF NITROGENOUS BASES


3193. Dibutylthallium isocyanate
 [74637-34-6] **C₉H₁₈NOTl**
Bu₂TlN:C:O

Srinastava, T. N. *et al.*, *Indian J. Chem. Sect. A*, 1980, **19A**, 480
 It explodes at 35°C.
See related ALKYL METALS, METAL CYANATES *See other* N-METAL DERIVATIVES

3194. Di-tert-butyl diperoxycarbonate
 [3236-56-4] **C₉H₁₈O₅**
(Me₃COO)₂CO

A partially decomposed sample exploded violently at 135°C.
See entry PEROXYCARBONATE ESTERS *See other* PEROXYESTERS

3195. 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxonane
 [17088-37-8] **C₉H₁₈O₆**



1. Dilthey, W. *et al.*, *J. Prakt. Chem.*, 1940, **154**, 219
2. Rohrlich, M. *et al.*, *Z. Ges. Schiess u. Sprengstoffw.*, 1943, **38**, 97-99

This trimeric acetone peroxide is powerfully explosive [1], and will perforate a steel plate when heated on it [2].

See 6-Aminopenicillanic acid *S*-oxide

Hydrogen peroxide: Ketones, Nitric acid


Ozone: Citronellic acid

See other CYCLIC PEROXIDES

3196. 3,6,9-Triethyl-1,2,4,5,7,8-hexoxonane

[]

C₉H₁₈O₆



Rieche, A. *et al.*, *Ber.*, 1939, **72**, 1938


This trimeric 'propylidene peroxide', formed from propanal and hydrogen peroxide, is an extremely explosive and friction-sensitive oil.

See other CYCLIC PEROXIDES

3197. 2,2,4,4,6,6-Hexamethyltrithiane

[828-26-2]

C₉H₁₈S₃




Nitric acid

See Nitric acid: 2,2,4,4,6,6-Hexamethyltrithiane

3198. Pivaloyloxydiethylborane

[32970-52-8]

C₉H₁₉O₂



Air, or Water

Köster, R. *et al.*, *Inorg. Synth.*, 1983, **22**, 196, 189

It reacts violently with air or water.
See related ACID ANHYDRIDES, ALKYLBORANES

†3199. 3,3-Diethylpentane [1067-20-5]	Et ₄ C	C ₉ H ₂₀
†3200. 2,5-Dimethylheptane [2216-30-0]	Me ₂ CHC ₂ H ₄ CHMeEt	C ₉ H ₂₀
†3201. 3,5-Dimethylheptane [926-82-9]	EtMeCHCH ₂ CHMeEt	C ₉ H ₂₀
†3202. 4,4-Dimethylheptane [1069-19-5]	Me ₂ CPr ₂	C ₉ H ₂₀
†3203. 3-Ethyl-2,3-dimethylpentane [16747-33-4]	Me ₂ CHC(Me)Et ₂	C ₉ H ₂₀
†3204. 3-Ethyl-4-methylhexane [3074-77-9]	Et ₂ CHCHMeEt	C ₉ H ₂₀
†3205. 4-Ethyl-2-methylhexane [3074-75-7]	Me ₂ CHCH ₂ CHEt ₂	C ₉ H ₂₀
†3206. 2-Methyloctane [3221-61-2]	Me ₂ CH[CH ₂] ₅ Me	C ₉ H ₂₀
†3207. 3-Methyloctane [2216-33-3]	Me ₂ CH[CH ₂] ₅ Me	C ₉ H ₂₀

†3208. 4-Methyloctane
[2216-34-4 resp.] C₉H₂₀



†3209. Nonane
[111-84-2] C₉H₂₀



†3210. 2,2,5-Trimethylhexane
[3522-94-9] C₉H₂₀




†3211. 2,2,3,3-Tetramethylpentane
[7154-79-2] C₉H₂₀



†3212. 2,2,3,4-Tetramethylpentane
[1186-53-4] C₉H₂₀



3213. 1-Methoxy-2-methyl-1-(trimethylsilyloxy)-1-butene ((1-Methoxy-2-methyl-1butenyl)oxy]trimethylsilane)
[84393-12-4] C₉H₂₀O₂Si



Bromoanisole, Sodamide


See Sodamide: Aryl halide, 1-Alkoxy-1-(trimethylsilyloxy)alkenes

3214. Triisopropylaluminium
[2397-67-3] C₉H₂₁Al




See entry TRIALKYLALUMINIUMS

3215. Tripropylaluminium
[102-67-0]



See entry TRIALKYLALUMINIUMS

3216. Aluminium isopropoxide
[555-31-2]



Crotonaldehyde, 2-Propanol


See 2-Propanol: Aluminium isopropoxide, etc.

Hydrogen peroxide

See Hydrogen peroxide: Aluminium isopropoxide

See other METAL ALKOXIDES

3217. Tripropylborane
[1116-61-6]



Gibson, 1969, 147

It ignites in air.



See other ALKYLBORANES

3218. Tripropylsilyl perchlorate
[]



See entry ORGANOSILYL PERCHLORATES

3219. Tripropylindium
[3015-98-3]




Leleu, *Cahiers*, 1977, (88), 367

It ignites in air.

See other ALKYLMETALS

3220. Triisopropylphosphine
[6476-36-4]

C₉H₂₁P



Chloroform, or Oxidants


Catalogue entry, Deutsche Advance Produktion, 1968

Particularly this phosphine reacts, when undiluted, rather vigorously with most peroxides, ozonides, *N*-oxides, and also chloroform. It may be safely destroyed by pouring into a solution of bromine in carbon tetrachloride.

See other ALKYLPHOSPHINES

3221. Tripropylantimony
[5613-69-4]

C₉H₂₁Sb




Gibson, 1969, 171

The solid ignites or chars when exposed to air on filter paper.

See other ALKYLMETALS

3222. Tris(2-propylthio)silane
[17891-55-3]

C₉H₂₂S₃Si




Lambert, J. A., *J. Amer. Chem. Soc.*, 1988, **110**, 2210

The silane, prepared by reduction of tris(2-propylthio)silylenium perchlorate with diisobutylaluminium hydride at -78°C , is subsequently isolated by high-vacuum distillation from the mixture. Heating must be very mild to prevent explosion.

See related ALKYL SILANES

3223. *N,N*-Bis(diethylboryl)methylamine
[19162-21-1]

C₉H₂₃B₂N




491M, 1975, 258

It ignites in air.

See related ALKYLBORANES

3224. Tris(dimethylfluorosilylmethyl)borane
[62497-91-0]

C₉H₂₄BF₃Si₃




Hopper, S. P. *et al.*, *Synth. React. Inorg. Met.-Org. Chem.*, 1976, **6**, 378

The crude or pure material will ignite in air, especially if exposed on filter paper or a similar extended surface.

See related ALKYLBORANES, ALKYLHALOSILANES

3225. *N,N,N',N'*-Tetramethylethane-1,2-diamine, trimethylpalladium(IV) bromide (Bromotrimethyl-*N,N,N',N'*-tetramethylethanediamine-*N,N'*-palladium)
[123147-95-5]


C₉H₂₅BrN₂Pd



De Graaf, W. *et al.*, *Organometallics*, 1990, **9**(5), 1479


Decomposes explosively when heated to 50°C.
See other AMMINEMETAL HALIDES, *See related* PLATINUM COMPOUNDS

3226. *N,N,N,N*-Tetramethylethane-1,2-diamine, trimethylpalladium(IV) iodide
[123147-84-2] **C₉H₂₅IN₂Pd**



De Graaf, W. *et al.*, *Organometallics*, 1989, **8**(12), 2907
Decomposes explosively when brought to room temperature.
See other AMMINEMETAL HALIDES
See related PLATINUM COMPOUNDS

3227. Tris(trimethylsilyl)aluminium
[65343-66-0] **C₉H₂₇AlSi₃**




Rösch, L., *Angew. Chem. (Intern. Ed.)*, 1977, **16**, 480
The crystalline solid ignites in air.
See related ALKYL SILANES

3228. Potassium dinitrotris(trimethylphosphine)cobaltate(1-)
[63181-09-9] **C₉H₂₇CoKN₂P₃**
K[N₂Co(PMe₃)₃]

Hammer, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1977, **16**, 485
The crystalline solid ignites immediately on exposure to air.
See related ALKYLPHOSPHINES, *N*-METAL DERIVATIVES

3229. Tris(trimethylsilyl)phosphine
[5573-38-3]

$C_9H_{27}PSi_3$




Dahl, O., *Acta Chem. Scand. Ser. B*, 1976, **30**, 799

It is pyrophoric.

See related ALKYL SILANES

3230. Tris(trimethylsilyl)hydrazine
[13272-02-1]

$C_9H_{28}N_2Si_3$



Oxidants

It is hypergolic with fluorine or fuming nitric acid, and explodes with ozone.

See entry SILYLHYDRAZINES

3231. Tris(trimethylsilyl)silane (1,1,1,3,3,3-Hexamethyl-2-(trimethylsilyl)trisilane)
[1873-77-4]

$C_9H_{28}Si_4$

$(Me_3Si)_3SiH$

Preparative hazard


Dickhaut, J. *et al.*, *Org. Synth.*, 1992, **70**, 164

The solid residue of the preparation from lithium powder, trimethylsilyl chloride, tetrachlorosilane and methyllithium may be highly pyrophoric. Caution in quenching into dilute acid is advised.

See other ALKYL SILANES

3232. *N,N,N'*-Tris(trimethylsilyl)diaminophosphine
[63104-54-1]


$C_9H_{29}N_2PSi_3$



Niecke, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1977, **16**, 487
The finely divided solid ignites in air.
See related ALKYL SILANES, PHOSPHINES

3233. Dicarbonyl- η -trichloropropenyldinickel chloride dimer
[64297-36-5]


$C_{10}Cl_8Ni_4O_4$



Posey, R. G. *et al.* *J. Amer. Chem. Soc.*, 1977, **99**, 4865 (footnote 13)
Nickel carbonyl and tetrachloropropadiene react in a wide variety of solvents to give the crystalline title compound which is extremely shock-sensitive, and explodes if touched when dry.
See related CARBONYLMETALS

3234. 2-Azidoperfluoronaphthalene
[74415-60-4]

$C_{10}F_7N_3$




Banks, R. E. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1980, 151

Thermolyses at 300°C/7 mbar to give 2-cyanoperfluorindene were terminated by explosions.

See other ORGANIC AZIDES

3235. Octatetrayne-1,8-dicarboxylic acid
[40575-26-6]

$C_{10}H_2O_4$




Sorbe, 1968, 151

An explosive solid, extremely heat-sensitive.

See other ACETYLENIC COMPOUNDS, ORGANIC ACIDS

3236. Copper(I) benzene-1,4-bis(ethynide)
[40575-26-6]

$(C_{10}H_4Cu_2)_n$




Royer, E. C. *et al.*, *J. Inorg. Nucl. Chem.*, 1981, **43**, 708

Explosive, and pyrophoric at 200°C.

See other METAL ACETYLIDES

3237. 1,3,6,8-Tetranitronaphthalene
[28995-89-3]


C₁₀H₄N₄O₈



Sorbe, 1968, 151
An explosive solid, most sensitive to heating.
See other POLYNITROARYL COMPOUNDS

3238. 2,2'-Azo-3,5-dinitropyridine
[55106-91-7]


C₁₀H₄N₈O₈



Explosive.
See entry POLYNITROAZOPYRIDINES
See other AZO COMPOUNDS, POLYNITROARYL COMPOUNDS

3239. [I,I-Bis(trifluoroacetoxy)iodo]benzene
[2712-78-9]


C₁₀H₅F₆IO₄



Increase in pH converts the title compound into an explosive 'dimer'.
See μ -OxoI,I-bis(trifluoroacetato-O)-I,I-diphenyldiiodine(III)
See 3,5-Dimethyl-4[I,I-bis(trifluoroacetoxy)iodo]isoxazole
See other I,I-(DIBENZOYLPEROXY)ARYLIODINESIODINE COMPOUNDS

3240. 2-Hydroxy-6-nitro-1-naphthalenediazonium-4-sulfonate
[5366-84-7]

C₁₀H₅N₃O₆S



HCS 1980, 362


Fast smouldering propagation occurs on heating the powder moderately.

See entry HIGH RATE DECOMPOSITION

See related DIAZONIUM CARBOXYLATES

3241. 1,4-Diethynylbenzene
[935-14-8]

C₁₀H₆




Preparative hazard

See 1,4-Bis(1,2-dibromoethyl)benzene: Potassium hydroxide

See other ALKYNES

3242. 1,5-Dinitronaphthalene
[605-71-0]

C₁₀H₆N₂O₄



Sulfur, Sulfuric acid

Hub, L., *Proc. Chem. Process Haz. Symp. VI*, (Manchester, 1977), No. 49, 43,
Rugby, IChE, 1977


For industrial conversion to 5-aminonaphthoquinone derivatives, dinitronaphthalene had been mixed cold with sulfuric acid and sulfur (to form sulfur dioxide), then heated to 120°C on over 100 occasions without incident. When dinitronaphthalene from a different supplier was used, the mixture exploded violently. Investigation in the safety calorimeter showed that an exothermic reaction begins at only 30°C, and that the onset and intensity of the exotherm (up to 400°C) markedly depends on quality of the dinitronaphthalene.

See REACTION SAFETY CALORIMETRY *See other* POLYNITROARYL COMPOUNDS

3243. 6-Quinolinecarbonyl azide

[78141-00-1]

C₁₀H₆N₄O



Houben-Weyl, 1952, Vol. 8, 682


A sample heated above its m.p. (88°C) exploded violently.

See other ACYL AZIDES

3244. 2,4-Diethynylphenol

[30427-53-3]

C₁₀H₆O




Kotlyarevskii, I. L. *et al.*, *Chem. Abs.*, 1971, **74**, 53198

Oxidation of the methyl or propyl ethers gave insoluble polymeric solids which exploded on heating.

See other ACETYLENIC COMPOUNDS, POLYPEROXIDES

3245. Di-2-furoyl peroxide
[25639-45-6]


C₁₀H₆O₆



Castrantas, 1965, 17
It explodes violently on friction and heating.
See other DIACYL PEROXIDES

3246. 1-Naphthalenediazonium perchlorate
[68597-02-4]


C₁₀H₇ClN₂O₄



Hofmann, K. A. *et al.*, *Ber.*, 1906, **39**, 3146
The salt explodes under light friction or pressure when dry.
See other DIAZONIUM PERCHLORATES

3247. 2-Naphthalenediazonium perchlorate
[69587-03-5]



C₁₀H₇ClN₂O₄



Hofmann, K. A. *et al.*, *Ber.*, 1906, **39**, 3146
The salt explodes under light friction or pressure when dry.
See other DIAZONIUM PERCHLORATES

3248. 2-Naphthalenediazonium trichloromercurate
[68448-47-5]

C₁₀H₇Cl₃HgN₂




Nesmeyanov, A. M., *Org. Synth.*, 1943, Coll. Vol. 2, 433

The isolated double salt precipitated by mercury(II) chloride explodes violently if heated during drying.

See other DIAZONIUM SALTS, MERCURY COMPOUNDS

3249. 1-Nitronaphthalene
[86-57-7]

C₁₀H₇NO₂



Nitric acid, Sulfuric acid

See Nitric acid: 1-Nitronaphthalene, Sulfuric acid


Tetranitromethane

See Tetranitromethane: Aromatic nitro compounds

See other NITROARYL COMPOUNDS

3250. 1-Nitroso-2-naphthol
[131-91-9]

C₁₀H₇NO₂



1. Starobinskii, V. A. *et al.*, *Chem. Abs.*, 1980, **92**, 7828

2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

3. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987

4. *CHETAH*, 1990, 187


1-Nitroso-2-naphthol, an intermediate in the preparation of ‘Pigment green’, undergoes an exothermic rearrangement at 124°C to an unstable material which appears to ignite spontaneously. Pigment green (an iron complex of a bisulfite derivative) decomposes at 230°C with a large exotherm, and the instances of ignition observed during drying of the pigment may be connected with the presence of some free nitrosonaphthol in it [1]. Energy of decomposition (in range 110–180°C) measured by DSC as 1.409 kJ/g by DSC, and T_{ait24} was determined as the low figure of 78°C by adiabatic Dewar tests, with an apparent energy of activation of 209 kJ/mol [2]. The solid deflagrates at 0.74 cm/min, and will attain a maximum of 68 bar pressure in a closed system [3]. It is shock sensitive [4].

See other NITROSO COMPOUNDS, SELF-HEATING AND IGNITION INCIDENTS

3251. 1-Naphthalenediazonium salts

[15511-25-8] (ions)

$C_{10}H_7N_2Z$




Ammonium sulfide, or Hydrogen sulfide, or Sodium sulfide

See entry DIAZONIUM SULFIDES AND DERIVATIVES

3252. 2-Naphthalenediazonium salts

[36097-38-8] (ions)

$C_{10}H_7N_2Z$




Ammonium sulfide, or Hydrogen sulfide, or Sodium sulfide

See entry DIAZONIUM SULFIDES AND DERIVATIVES

3253. Naphthylsodium
[25398-08-7]

C₁₀H₇Na



Chlorinated biphenyl

MCA Case History No. 565


A chlorinated biphenyl heat-transfer liquid was added to a burning batch of naphthylsodium to help to extinguish it. An exothermic reaction, followed by an explosion occurred. Sodium is known to react violently with many halogenated materials.

See Sodium: Halocarbons

See other ARYLMETALS

3254. Naphthalene
[91-20-3]

C₁₀H₈



(MCA SD-58, 1956); NSC 370, 1978; FPA H69, 1978; HCS 1980, 672

Zavetskii, A. D., Chem. Abs., 1983, 98, 128923

Fires inside wood-packed benzene scrubbers in coke oven gas plants were attributed to saturation of the wood with naphthalene, and vapour-phase oxidation of the latter to phthalic anhydride, which participates in exothermic free radical chain reactions.

Aluminium chloride, Benzoyl chloride

See Aluminium chloride: Benzoyl chloride, etc.

Dinitrogen pentaoxide


See Dinitrogen pentaoxide: Naphthalene

Other reactants

Yoshida, 1980, 266

MRH values calculated for 13 combinations with oxidants are given.

3255. 2,2'-Bipyridyldichloropalladium(IV) perchlorate
[14871-92-2] (ion)




Gray, L. R. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1983, 133–141

Though no explosions occurred during preparation (by addition of 72% perchloric acid to solutions of the complex in conc. nitric/hydrochloric mixtures), this and related amino and phosphino analogues are potentially unstable.

See other AMMINOMETAL OXOSALTS

3256. μ ,1,1'-Ferrocenediydilithium (Dilithioferrocene)
[33272-09-2]
[174872-99-2] tetramethylethanediamine complex




Anhydrides

Rulkens, R. *et al.*, *J. Amer. Chem. Soc.*, 1997, **119**(45), 10976

A synthetic procedure was devised which involved mixing this organometallic, as its TMEDA complex, with either solid benzenesulphonothioic acid anhydrosulphide (PhSO_2)₂S, or the equivalent selenide, at liquid nitrogen temperatures, and adding tetrahydrofuran. The first few drops of THF, which presumably allowed the dog to see the rabbit, produced vigorous interaction, with possible pressure generation. Reacting anhydrides with alkyl metals, in absence of diluents, may surely be expected to be violent.

See other ALKYL METALS


3257. 1,3-Bis(isocyanomethyl)benzene
[]



See entry DIISOCYANIDE LIGANDS *See related* CYANO COMPOUNDS

3258. 2,2'-Bipyridyl 1-oxide
[33421-43-1]

C₁₀H₈N₂O




Preparative hazard

See Hydrogen peroxide: Acetic acid, *N*-Heterocycles

See other *N*-OXIDES

3259. 2-Furaldehyde azine
[5428-37-5]

C₁₀H₈N₂O₂



Nitric acid


See Nitric acid: Aromatic amines (reference 5), and : Hydrazines

See related HYDRAZONES

3260. 2,2'-Oxybis(iminomethylfuran) mono-*N*-oxide (Dehydrofurfural oxime)

[]

C₁₀H₈N₂O₄




Ponzio, G. *et al.*, *Gazz. Chim. Ital.*, 1906, **36**, 338–344

It explodes at 130°C.

See other *N*-OXIDES, *N*-O COMPOUNDS

3261. Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane-1,2-dicarboxylic acid (Cubane-1,2-dicarboxylic acid)
[129103-51-1]

C₁₀H₈O₄



Eaton, P. E. *et al.*, *J. Org. Chem.*, 1990, **55**(22), 5746


Attempts to dehydrate this to the anhydride were unsuccessful; attempting to distil off monomeric products under vacuum at 80°C after a dehydration attempt with ethoxyacetylene led to a violent explosion. This is said to be atypical of cubanes, but:

See other STRAINED-RING COMPOUNDS

See entry CUBANES

3262. Poly(ethylene terephthalate)
[25038-59-9]

(C₁₀H₈O₄)_n




Preparative hazard

See Dimethyl terephthalate: Ethylene glycol, Titanium butoxide

3263. 2,2'-Bi-5,6-dihydro-1,3-dithiolo[4,5-b][1,4]dithiinylidene
[66946-48-3]

C₁₀H₈S₈



Preparative hazard


Larsen, J. *et al.*, *Org. Synth.*, 1995, **72**, 265

A procedure avoiding the sometimes explosive reaction of carbon disulfide with alkali metals is detailed.

See Carbon disulfide: Alkali metals

3264. 1-Naphthylamine
[134-32-7]

C₁₀H₉N




Nitrous acid

See 1-Naphthalenediazonium salts and 2-Naphthalenediazonium salts

3265. 2-Naphthylamine
[91-59-8]

C₁₀H₉N




Nitrous acid

See 1-Naphthalenediazonium salts and 2-Naphthalenediazonium salts

3266. 4-Acetoxy-3-methoxy-2-nitrobenzaldehyde
[2698-69-3]

C₁₀H₉NO₆




Preparative hazard

See Nitric acid: 4-Acetoxy-3-methoxybenzaldehyde

See *other* ALDEHYDES, NITROARYL COMPOUNDS

3267. Dipyridinesilver(I) perchlorate

[2152-13-7]



Acids


Kauffman, G. B. *et al.*, *Inorg. Synth.*, 1960, **6**, 7, 8

Contact with acids, especially hot, must be avoided to prevent the possibility of violent explosion.

See other AMMINOMETAL OXOSALTS, SILVER COMPOUNDS

3268. 1,4-Bis(1,2-dibromoethyl)benzene

[25393-98-0]



Potassium hydroxide


Kotlyarevskii, I. L. *et al.*, *Chem. Abs.*, 1978, **89**, 23858

Tetra-dehydrobromination of the compound in benzene with ethanolic potassium hydroxide for the commercial production of 1,4-diethynylbenzene is highly exothermic and needs careful control to avoid hazard.

See related HALOALKANES

3269. Ferrocenium perchlorate

[12312-33-3]




See 1,3-Di[bis(cyclopentadienyl)iron]-2-propen-1-one: Perchloric acid, etc.

See other ORGANOMETALLIC PERCHLORATES

3270. 1-Chloro-4-(2-nitrophenyl)-2-butene
[7318-34-5]


C₁₀H₁₀ClNO₂



Acheson, R. M. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1978, 1119
It may explode during distillation at 112–114°C/60 mbar.
See other ALLYL COMPOUNDS, NITROARYL COMPOUNDS

3271. Dicyclopentadienylperoxyniobium chloride
[75109-11-4]


C₁₀H₁₀ClNbO₂



Sala-pala, J. *et al.*, *J. Mol. Catal.*, 1980, **7**, 142–143
The compound and its methyl homologue exploded occasionally when scraped from the walls of a tube, and invariably when exposed to a Raman laser beam, even at –80°C.
See other IRRADIATION DECOMPOSITION INCIDENTS, ORGANOMETALLIC PEROXIDES

3272. 4-(2,4-Dichlorophenoxy)butyric acid
[94-82-6]


C₁₀H₁₀Cl₂O₃



Preparative hazard
See γ -Butyrolactone: 2,4-Dichlorophenol, etc.

3273. Dicyclopentadienylchromium (Chromocene)
[1271-24-5]

C₁₀H₁₀Cr



Alcohols

Votinsky, J. *et al.*, *Coll. Czech. Chem. Comm.*, 1979, **44**, 82

Reaction with C₁–C₄ alcohols at 20°C produces pyrophoric dialkoxochromiums and cyclopentadiene.

See METAL ALKOXIDES See other ORGANOMETALLICS

3274. Oxodiperoxodipyridinechromium(VI)

[]

[(C₅H₅N)₂CrO(O₂)₂]

C₁₀H₁₀CrN₂O₅

1. Caldwell, S. H. *et al.*, *Inorg. Chem.*, 1969, **8**, 151
2. Adams, D. M. *et al.*, *J. Chem. Educ.*, 1966, **43**, 94
3. Collins, J. C. *et al.*, *Org. Synth.*, 1972, **52**, 5–8
4. Wiede, O. F., *Ber.*, 1897, **30**, 2186
5. Glaros, G., *J. Chem. Educ.*, 1978, **55**, 410


This complex, formerly called 'pyridine perchromate' and now finding application as a powerful and selective oxidant, is violently explosive when dry [1]. Use while moist on the day of preparation and destroy any surplus with dilute alkali [2]. Preparation and use of the reagent have been detailed further [3]. The analogous complexes with aniline, piperidine and quinoline may be similarly hazardous [4]. The damage caused by a 1 g sample of the pyridine complex exploding during desiccation on a warm day was extensive. Desiccation of the aniline complex had to be at ice temperature to avoid violent explosion [4]. Pyridinium chlorochromate is commercially available as a safer alternative oxidant of alcohols to aldehydes [5].

See Chromium trioxide: Pyridine

Dipyridinium dichromate

See other AMMINECHROMIUM PEROXOCOMPLEXES

3275. Bis(cyclopentadienyldinitrosylchromium)
[66649-16-9]




Flitcroft, N. *et al.*, *Chem. & Ind.*, 1969, 201

A small sample exploded violently upon laser irradiation for Raman spectroscopy.

See other IRRADIATION DECOMPOSITION INCIDENTS, ORGANOMETALLICS

3276. Ferrocene
[102-54-5]



Mercury(II) nitrate


See Mercury(II) nitrate: Ferrocene

Tetranitromethane

See Tetranitromethane: Ferrocene

See other ORGANOMETALLICS

3277. Bis(cyclopentadienyl)magnesium
[1284-72-6]




Barber, W. A., *Inorg. Synth.*, 1960, **6**, 15

It may ignite on exposure to air.
See other ORGANOMETALLICS

3278. Bis(cyclopentadienyl)manganese
[1271-27-8]

C₁₀H₁₀Mn




491M, 1975, 248

It ignites in air.
See other ORGANOMETALLICS

3279. Oxodiperoxodi(pyridine N-oxide)molybdenum
[73680-23-6]

C₁₀H₁₀MoN₂O₇




Westland, A. D. *et al.*, *Inorg. Chem.*, 1980, **19**, 2257

The analogous complex of tungsten, and that of chromium with a single *N*-oxide ligand, were also explosive.

See related AMMINECHROMIUM PEROXOCOMPLEXES

3280. Dipyridinesodium
[101697-88-5]

C₁₀H₁₀N₂Na



Sidgwick, 1950, 89


The addition product of sodium and pyridine (or its 2-methyl derivative) ignites in air.

See related N-METAL DERIVATIVES

**3281. 4a,5,7a,8-Tetrahydro-4,8-methano-4H -indeno[5,6-c]-1,2,5-oxadiazole
1-oxide (Dicyclopentadienefurazan N-oxide)**

[]

$C_{10}H_{10}N_2O_2$



Crosby, J. *et al.*, Br. Pat. 1 474 693, 1977


The solid isomeric mixture of the 1-oxides explodes at 80–85°C, but may be handled safely in solution.

See other FURAZAN N-OXIDES

**3282. 4a,5,7a,8-Tetrahydro-4,8-methano-4H -indeno[5,6-c]-1,2,5-oxadiazole
3-oxide (Dicyclopentadienefurazan N-oxide)**

[]

$C_{10}H_{10}N_2O_2$



Crosby, J. *et al.*, Br. Pat. 1 474 693, 1977


The solid isomeric mixture of the 3-oxide explodes at 80–85°C, but may be handled safely in solution.

See other FURAZAN N-OXIDES

3283. Oxodiperoxodi(pyridine N-oxide)tungsten

[]

$C_{10}H_{10}N_2O_7W$




Westland, A. D. *et al.*, *Inorg. Chem.*, 1980, **19**, 2257

The analogous complex of molybdenum, and that of chromium with a single *N*-oxide ligand, were also explosive.

See related AMMINECHROMIUM PEROXOCOMPLEXES

3284. *N,N'*-Dimethyl-*N,N'*-dinitrosoterephthalamide
[133-55-1]

C₁₀H₁₀N₄O₄



2-Hydroxyethylamine

MCA Case History No. 2308


Addition of the dry nitrosamide to a flask apparently containing residual ethanolamine led to evolution and ignition of diazomethane and the dry nitrosamide powder.

See Diazomethane

See other NITROSO COMPOUNDS

3285. Bis(cyclopentadienyl)tungsten diazide oxide
[53504-80-6]

C₁₀H₁₀N₆O_W




Anand, S. P., *J. Inorg. Nucl. Chem.*, 1974, **36**, 926

It is highly explosive.

See related METAL AZIDES

3286. Bis(cyclopentadienyl)vanadium diazide
[85836-55-1]


C₁₀H₁₀N₆V



Blandy, C. *et al.*, *Tetrahedron. Lett.*, 1983, **24**, 4189–4192
It is explosive in the solid state.
See related METAL AZIDES

3287. Nickelocene
[1271-28-9]


C₁₀H₁₀Ni



See 1-(1-Methyl-1-phenylethyl)-4-(2-propynyloxy)benzene
See other ORGANOMETALLICS

3288. 2-Indanecarboxaldehyde
[37414-44-1]

C₁₀H₁₀O




1. Black, T. H., *Chem. Eng. News*, 1988, **66**(32), 2
2. Black, T. H., private comm., 1988

A violent explosion occurred at the start of bulb-tube distillation of a 2 g sample of the crude aldehyde (b.p. 97°/4 mbar) at 75°C/0.13 mbar [1]. It seems likely that the crude product could have formed peroxide during the 2 day's storage (in a closed but not hermetically sealed vessel) between preparation and distillation. The structure is such that the proton on C-2, which effectively has 2 benzyl substituents and an adjacent carbonyl function, seems rather susceptible to autoxidation [2].

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

3289. 1-Oxo-1,2,3,4-tetrahydronaphthalene
[529-34-0]

C₁₀H₁₀O




Preparative hazards

See Chromium trioxide: Acetic anhydride
Hydrogen peroxide: Acetone, etc.

3290. 4-Acetoxy-3-methoxybenzaldehyde
[881-68-5]

C₁₀H₁₀O₄




Nitric acid

See Nitric acid: 4-Acetoxy-3-methoxybenzaldehyde
See other ALDEHYDES

3291. 1,3-Diacetoxybenzene (Resorcinol diacetate)
[108-58-7]

C₁₀H₁₀O₄




Nitric acid

See 4,6-Dinitro-1,3-benzenediol

3292. Dimethyl terephthalate
[120-61-6]

C₁₀H₁₀O₄



The finely powdered ester is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)


Ethylene glycol, Titanium butoxide

1. Rix, G. C., private comm., 1977
2. Goodman, I. in *Encyclopaedia of Polymer Science and Technology*, Mark, H. F. (Ed.), Vol. 11, 112–113, New York, Interscience, 1969

The polymer had been produced for 12 years by transesterification under nitrogen of the dimethyl ester with ethylene glycol at 250°C in presence of titanium butoxide catalyst. After increasing the heating capacity of the vessel from a half coil to a full coil, 3 incidents of ignition of vapour after opening the vessel were noted [1]. This is attributed to formation and ignition of mixtures of acetaldehyde or dioxane with ingressing air on the hot vessel surfaces. Acetaldehyde is produced by thermal degradation of the polymer [2], and has rather wide flammability limits (4–57%) and is readily ignited on hot or corroded surfaces. Dioxane, readily formed by heating ethylene glycol with acids, metal oxides or other catalysts, also has wide limits (2–22%) and a low autoignition temperature (180°C). It seems likely that if either of these were present, perhaps with residual traces of methanol vapour, migration towards an open vent would produce turbulence and admixture with air.
See related SELF-HEATING AND IGNITION INCIDENTS

3293. Bis(cyclopentadienyl)titanium selenate
[87612-48-4]

C₁₀H₁₀O₄SeTi





Malek, J. *et al.*, *Z. Chem.*, 1983, **23**, 189

It is explosive, sensitive to heat or friction.
See other ORGANOMETALLICS *See related* ORGANOMETALLIC NITRATES

3294. Bis(cyclopentadienyl)lead
[1294-74-2]


C₁₀H₁₀Pb



Preparative hazard
See Lead(II) nitrate: Cyclopentadienylsodium
See other ORGANOMETALLICS

3295. Bis(cyclopentadienyl)titanium
[1271-29-9]

C₁₀H₁₀Ti




Bailar, 1973, Vol. 3, 389
Pyrophoric crystals (but see Polar solvents, below).

Polar solvents
Watt, G. W. *et al.*, *J. Amer. Chem. Soc.*, 1966, **88**, 1139–1140
Though not pyrophoric, it reacts very vigorously with oxygen-free water and other polar solvents.
See other ORGANOMETALLICS

3296. Bis(cyclopentadienyl)zirconium
[12116-83-5]


C₁₀H₁₀Zr



Watt, G. W. *et al.*, *J. Amer. Chem. Soc.*, 1966, **88**, 5926
A pyrophoric solid.
See other ORGANOMETALLICS

3297. 3-(4-Chlorophenyl)butanoic acid
[1012-17-5]

C₁₀H₁₁ClO₂




Preparative hazard

See Peroxyacetic acid: 5-(4'-Chlorophenyl)-2,2-dimethyl-3-hexanone
See other ORGANIC ACIDS

3298. Dimethyl(phenylethynyl)thallium
[10158-43-7]


C₁₀H₁₁Tl



Nast, R. *et al.*, *J. Organomet. Chem.*, 1966, **6**, 461
May explode on stirring, heating or impact.
See other ALKYL METALS, METAL ACETYLIDES

†3299. Dicyclopentadiene
[77-73-6]

C₁₀H₁₂




HCS 1980, 387

Ahmed, M., *et al.*, *Plant/Oper. Progr.*, 1991, **10**(3), 143
The compound is capable of exothermic pressure generating decompositions, from about 170°C.
See other DIENES, GAS EVOLUTION INCIDENTS

3300. Tetrahydronaphthalene
[119-64-2]

C₁₀H₁₂



HCS 1980, 893; *RSC Lab. Hazards Safety Data Sheet No. 78*, 1989



1. Grignard, 1949, Vol. 17.1, 673
2. Duckworth, D. F., private comm., 1977
3. McDonald, J. S., *Chem. Abs.*, 1960, **54**, 15933

Readily peroxidises in air, and explosions have occurred during large-scale distillations [1], particularly towards the end when sufficiently concentrated [2]. Use of tetralin as secondary coolant in a molten sodium heat transfer system led to an explosion due to peroxide formation [3].

See 1,2,3,4-Tetrahydro-1-naphthyl hydroperoxide

See other PEROXIDISABLE COMPOUNDS

3301. 4-Chloro-2,5-diethoxynitrobenzene
[91-43-0]





CHETAH, 1990, 183

Although not sensitive to mechanical shock, this compound can be exploded by a detonator.

See other HALOARYL COMPOUNDS, NITROARYL COMPOUNDS

3302. Ethylphenylthallium(III) acetate perchlorate
[34521-47-6]





Preparative hazard

See Perchloric acid: Ethylbenzene, Thallium triacetate

See related METAL PERCHLORATES


3303. 6,6'-Dihydrazino-2,2'-bipyridylnickel(II) perchlorate
[53346-21-7]



Lewis, J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1977, 738

It detonates violently on heating, but is stable to shocks.
See other AMMINEMETAL OXOSALTS

3304. Dipyridinium dichromate
[20039-37-6]





Preparative hazard

1. Coates, W. M. *et al.*, *Chem. & Ind.*, 1969, 1594
2. Fieser, M. *et al.*, *Reagents for Organic Synth.*, 1981, Vol. 9, 399
3. Corey, E. J. *et al.*, *Tetrahedron Lett.*, 1979, 399
4. Salmon, J., *Chem. & Ind.*, 1982, 616; *Chem. Brit.*, 1982, **18**, 703
5. Gill, P. *et al.*, *Thermochim. Acta*, 1984, **80**, 193–196

It is known that preparation of the oxidant salt under anhydrous conditions is explosion-prone [1], but during repetition of a supposedly safe preparative method [2], recommended for large-scale use [3], ignition and a violent explosion occurred. Use of more water to dissolve completely the chromium trioxide, and a reaction temperature below 25°C, are recommended [4]. During a study by TGA of the thermal degradation of the salt, too-rapid heating of the samples led to explosions with fire [5].

See other DICHROMATE SALTS OF NITROGENOUS BASES, OXIDANTS

3305. Sodium ethylenediaminetetraacetate
[67401-50-7]




Sodium hydroxide, Sodium hypochlorite

See Sodium hypochlorite: Sodium ethylenediaminetetraacetate, etc.

3306. 4,6-Dinitro-2-*sec*-butylphenol
[88-85-7]

C₁₀H₁₂N₂O₅




Nolan, 1983, Case history 99

A batch became overheated during drying (leaking steam valve) and decomposed violently.

See other POLYNITROARYL COMPOUNDS

3307. *O, O*-Dimethyl-*S*-[(4-oxo-1,2,3-benzotriazin-3[4*H*]-yl)methyl]phosphorodithioate (Azinphos-methyl)
[86-50-0]

C₁₀H₁₂N₃O₃PS₂




Kirschner, E., *Chem. Eng. News*, 1997, **75**(20), 12

A smouldering bag in a pesticide warehouse, believed to be of this, led to an explosion, killing three firemen, and fire which took six days to extinguish (possibly because of caution concerning anti-cholinesterase toxicity). This is a moderately high energy compound by virtue of the triazene function. Other pesticides present were of less energetic structure.

See other TRIAZENES

3308. 2-Dimethylamino-3-methylbenzoyl azide
[161616-67-7]

C₁₀H₁₂N₄O




Waldron, N. M. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1995, 81

With only 20% azide nitrogen by weight, this azide might be thought safe for cautious isolation. A 1 g sample isolated by evaporation of solvent at -10°C , exploded violently and spontaneously on warming to 0°C .

See other ACYL AZIDES

3309. 3,4-Dimethyl-4-(3,4-dimethyl-5-isoxazolylazo)isoxazolin-5-one
[4100-38-3]

$\text{C}_{10}\text{H}_{12}\text{N}_4\text{O}_3$




Boulton, A. J. *et al.*, *J. Chem. Soc.*, 1965, 5415

It invariably decomposed explosively if heated rapidly to 100°C , but was stable to impact or friction.

See other AZO COMPOUNDS, ISOXAZOLES, N-O COMPOUNDS

3310. 1,3,5,7-Tetranitroadamantane
[75476-36-7]

$\text{C}_{10}\text{H}_{12}\text{N}_4\text{O}_8$




Gilbert, E. E. *et al.*, US Pat. 4 329 522, 1982

It shows explosive properties

See related POLYNITROALKYL COMPOUNDS

3311. Isobutyrophenone
[611-70-1]

$\text{C}_{10}\text{H}_{12}\text{O}$




Bromine

See Bromine: Isobutyrophenone

3312. 1,2,3,4-Tetrahydro-1-naphthyl hydroperoxide
[771-29-9]

C₁₀H₁₂O₂



Hock, H. *et al.*, *Ber.*, 1933, **66**, 61
It explodes on superheating the liquid.

Lithium tetrahydroaluminate


Sutton, D. A., *Chem. & Ind.*, 1951, 272

Interaction in ether is vigorously exothermic.

See other REDOX REACTIONS See related ALKYL HYDROPEROXIDES

3313. 3-Methoxy-3-methyl-5,6-benzo-6H-1,2-dioxin
[69681-99-8, *cis*-; 69682-00-4, *trans*-]


C₁₀H₁₂O₃



Lerdal, D. *et al.*, *Tetrahedron Lett.*, 1978, 3228
Both isomers decomposed violently on heating in absence of solvent.
See other CYCLIC PEROXIDES

3314. 3-Methoxy-4-methyl-5,6-benzo-6H-1,2-dioxin
[69682-01-5]


C₁₀H₁₂O₃



Lerdal, D. *et al.*, *Tetrahedron Lett.*, 1978, 3228
It decomposed violently on heating in absence of solvent.
See other CYCLIC PEROXIDES

3315. Allyl 4-toluenesulfonate
[4873-09-0]

C₁₀H₁₂O₃S




Rüst, 1948, 302

Explosion and charring of the ester during high-vacuum distillation from an oil-bath at 110°C was ascribed to exothermic polymerisation, (probably catalysed by acidic decomposition products).

See other ALLYL COMPOUNDS, POLYMERISATION INCIDENTS, SULFUR ESTERS

3316. 2-Buten-1-yl benzenesulfonate
[20443-67-8]

C₁₀H₁₂O₃S




Sorbe, 1968, 122

After evaporation of solvent, the ester may explode in contact with air (possibly owing to exothermic polymerisation catalysed by peroxide formation).

See other POLYMERISATION INCIDENTS, SULFUR ESTERS

3317. 1-Phenylboralane
[1075-17-8]

C₁₀H₁₃B




Leleu, *Cahiers*, 1977, (88), 365

It ignites in air.

See related ALKYLBORANES

3018. 2-(4-Chlorophenyl)-1,1-dimethylethyl hydroperoxide
[71356-36-0]

$C_{10}H_{13}ClO_2$



Preparative hazard

1. Kochi, J. B. *et al.*, *J. Amer. Chem. Soc.*, 1964, **86**, 5268

2. Parris, M. *et al.*, *Can. J. Chem.*, 1979, **57**, 1233–1234

Preparation by the method used for the 2-phenyl analogue [1] is hazardous [2].


(No further details)

See Hydrogen peroxide: 2-Phenyl-1,1-dimethylethanol, etc.

See other ALKYL HYDROPEROXIDES

3319. 4-tert-Butyliodylbenzene
[64297-75-2]

$C_{10}H_{13}IO_2$




Ranganathan, S. *et al.*, *Tetrahedron Lett.*, 1985, **26**, 4955–4956

Although a stable crystalline compound (m.p. 217–221°C), its oxidation potential approximates to that of ozone, and it should only be used in a suitably inert solvent (chlorobenzene or nitrobenzene).

See other IODINE COMPOUNDS, OXIDANTS

3320. Diethyl trifluoroacetosuccinate
[94633-25-7]

C₁₀H₁₃F₃O₅




Preparative hazard

See Sodium hydride: Diethyl succinate, etc.

3321. 4-Dimethylaminoacetophenone
[18925-69-4]

C₁₀H₁₃NO




Preparative hazard

See 4-Chloroacetophenone: Dimethylamine

3322. 2-Formylamino-1-phenyl-1,3-propanediol
[51317-78-3]

C₁₀H₁₃NO₃




Nitric acid

See Nitric acid: 2-Formylamino-1-phenyl-1,3-propanediol

3323. Butylbenzene
[104-51-8]


C₁₀H₁₄



Preparative hazard
See Sodium: Halocarbons (reference 7)

3324. Bis(cyclopentadienyl)niobium tetrahydroborate
[37298-41-2]

C₁₀H₁₄BNb



Lucas, C. R., *Inorg. Synth.*, 1976, **16**, 109–110
It is pyrophoric in air, particularly after sublimation.
See other COMPLEX HYDRIDES, ORGANOMETALLICS

3325. Calcium 2,4-pentanedionate
[19372-47-2]


C₁₀H₁₄CaO₄

Ca(OCMe=CHCO.Me)₂ bicyclic chelate

Barium perchlorate
See Barium perchlorate: Calcium 2,4-pentanedionate

3326. Bis(2,4-pentanedionato)chromium
[14024-50-1]

C₁₀H₁₄CrO₄




Ocone, L. R. *et al.*, *Inorg. Synth.*, 1966, **8**, 131

It ignites in air.
See Chromium(II) acetate
See related METAL ALKOXIDES, ORGANOMETALLICS

3327. Diaquabis(cyclopentadienyl)titanium dichromate
[87612-49-5]


C₁₀H₁₄Cr₂O₉Ti



Malek, J. *et al.*, *Z. Chem.*, 1983, **23**, 189
Explosive, sensitive to heat or friction.
See other ORGANOMETALLICS

3328. Diethyl 4-nitrophenyl thionophosphate
[25737-28-4]

C₁₀H₁₄NO₅PS



1. Bright, N. H. F. *et al.*, *J. Agr. Food Chem.*, 1950, **1**, 344
2. Gweek, B. F. *et al.*, *Ind. Eng. Chem.*, 1959, **51**, 104–112

Vacuum distillation of 'parathion' at above 100°C is hazardous, frequently leading to violent decomposition [1]. Following a plant explosion, the process design was modified and featured a high degree of temperature sensing and control to avoid a recurrence [2].


Endrin

During the blending of the two insecticides (endrin is a halogenated polycyclic epoxide) into a petroleum solvent, an unexpected exothermic reaction occurred which vaporised some solvent and led to a vapour–air explosion. Faulty agitation was suspected.

See other AGITATION INCIDENTS, NITROARYL COMPOUNDS, PHOSPHORUS ESTERS

3329. Diethyl 4-nitrophenyl phosphate
[311-45-5]

C₁₀H₁₄NO₆P




Preparative hazard

See Diethyl phosphite: 4-Nitrophenol

See other NITROARYL COMPOUNDS, PHOSPHORUS ESTERS

3330. 1,4-Bis(isocyanomethyl)cyclohexane
[]


C₁₀H₁₄N₂



See entry DIISOCYANIDE LIGANDS

3331. 4-Azido-*N,N*-dimethylaniline
[24573-95-3]

C₁₀H₁₄N₄




491M, 1975, 51

It decomposed violently on attempted distillation.
See other ORGANIC AZIDES

3332. 2-Phenyl-1,1-dimethylethyl hydroperoxide
[1944-83-8]

C₁₀H₁₄O₂




Preparative hazard

See Hydrogen peroxide: 2-Phenyl-1,1-dimethylethanol, etc.
See other ALKYL HYDROPEROXIDES

3333. 2-Chlorobenzylidenemalononitrile ('CS gas')
[2698-41-1]


C₁₀H₁₅ClN₂



The finely powdered nitrile is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)
See other CYANO COMPOUNDS

3334. N-Methoxy-N-methylbenzylidenimmonium methylsulfate
[]

C₁₀H₁₅NO₅S




Volpiana, -, *Proc. 1st Int. Symp. Prev. Risks Chem. Ind.*, 176-179, Heidelberg, ISSA, 1970

Quaternation of *O*-methylbenzaldoxime was effected by addition to dimethyl sulfate at 110°C with cooling to maintain that temperature for 2 hours more. Cessation of temperature control allowed a slow exotherm to proceed unnoticed, and at 140–150°C a violent decomposition set in.

See related OXIMES *See other* N–O COMPOUNDS

3335. 1-Phenyl-3-*tert*-butyltriazene
[63246-75-3]

C₁₀H₁₅N₃




Rondevstedt, C. S. *et al.*, *J. Org. Chem.*, 1957, **22**, 203

It exploded on attempted distillation at below 1 mbar.

See other TRIAZENES

3336. Δ 3-Carene (3,7,7-Trimethylbicyclo[4.1.0]hept-3-ene)
[13466-78-9]

C₁₀H₁₆




Nitric acid

See Nitric acid: Hydrocarbons (reference 15)

See related ALKENES

3337. Limonene (4-Isopropenyl-1-methylcyclohexene)
[5989-27-5]

C₁₀H₁₆



HCS 1980, 591

Iodine pentafluoride, Tetrafluoroethylene

See Tetrafluoroethylene: Iodine pentafluoride, etc.


Sulfur

See Sulfur: Limonene

See other DIENES, PEROXIDISABLE COMPOUNDS

3338. 1,5-*p*-Menthadiene (5-Isopropyl-2-methyl-1,3-cyclohexadiene)
[99-83-2]

C₁₀H₁₆




Bodendorf, K., *Arch. Pharm.*, 1933, **271**, 28

The terpene readily peroxidises with air, and the (polymeric) peroxidic residue exploded violently on attempted distillation at 100°C/0.4 mbar.

See other DIENES, PEROXIDISABLE COMPOUNDS, POLYPEROXIDES

3339. 2-Pinene (2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene)
[80-56-8]

C₁₀H₁₆




Nitrosyl perchlorate

See Nitrosyl perchlorate: Organic materials

See related ALKENES

3340. 1-(4,5-Dimethyl-1,3-diselenonylidene)piperidinium perchlorate
[53808-74-5]


C₁₀H₁₆ClNO₄Se₂



See entry 1-(1,3-DISELENONYLIDENE)PIPERIDINIUM PERCHLORATES


See other PERCHLORATE SALTS OF NITROGENOUS BASES

3341. Sebacoyl chloride
[111-19-3]





Hüning, S. *et al.*, *Org. Synth.*, 1963, **43**, 37
During vacuum distillation of the chloride at 173°C/20 mbar, the residue frequently decomposes spontaneously, producing a voluminous black foam.
See other ACYL HALIDES

3342. Benzyltrimethylammonium permanganate
[76710-75-3]



It explodes at 80–90°C.
See entry QUATERNARY OXIDANTS

3343. 1,8-Diisocyanooctane
[]




See entry DIISOCYANIDE LIGANDS

3344. Sodium 1,3-dihydroxy-1,3-bis(*aci*-nitromethyl)-2,2,4,4-tetramethylcyclobutane-1,3-diolide

[]

C₁₀H₁₆N₂Na₂O₆



Water

Dauben, H. J., *Org. Synth.*, 1963, Coll. Vol. 4, 223

The dry powdered condensation product of sodium *aci*-nitromethanide (2 mol) with dimethylketene dimer exploded violently when added to crushed ice.


See Sodium *aci*-nitromethanide: Water

See other *aci*-NITRO SALTS

3345. 2,2'-Azoisovaleronitrile

[70843-45-7]

C₁₀H₁₆N₄




See entry BLOWING AGENTS See other AZO COMPOUNDS, CYANO COMPOUNDS

3346. 3,7-Dimethyl-2,6-octadienal (Citral)

[5392-40-5]

C₁₀H₁₆O




Preparative hazard

See 2-Methyl-1,3-butadiene: Acetone

See related DIENES

3347. 1,4-Epidioxy-2-*p*-menthene (Ascaridole)
[512-85-6]

C₁₀H₁₆O₂




1. Castrantas, 1965, 15
2. Henry, T. A. *et al.*, *J. Chem. Soc.*, 1921, **119**, 1722
3. Walton, F. P., private comm., 1980

Explosive decomposition occurs on heating from 130 to 150°C [1]. When heated above 160°C, explosive rearrangement occurs with an exotherm to 300°C. An explosion occurred during fractional distillation at 108–110°C/20 mbar. It may be distilled without explosion by a flash evaporation technique [2,3].

See other CYCLIC PEROXIDES

**3348. 3-Peroxycamphoric acid (1,2,2-Trimethyl-1-carboxycyclopentane-3-
peroxycarboxylic acid)**
[31921-03-6]

C₁₀H₁₆O₅




1. Milas, N. A. *et al.*, *J. Amer. Chem. Soc.*, 1933, **55**, 350
2. Pirkle, W. H. *et al.*, *J. Org. Chem.*, 1977, **42**, 2080–2082

Explosive decomposition occurs at 80–100°C [1], but the isomeric 1-peroxy acid may also have been present in the sample [2].

See other ORGANIC ACIDS, PEROXYACIDS

3349. 7,7,10,10-Tetramethyl-1,2,5,6-tetroxecane-3,4-dione (2,5-Dimethyl-2,5-hexylenebis diperoxyoxalate)
[35551-00-7]

C₁₀H₁₆O₆



Preparative hazard


Adam, W. *et al.*, *J. Amer. Chem. Soc.*, 1980, **102**, 4803

The preparation is hazardous because a highly explosive peroxidic polymer is formed as the major product. After extraction of the title product with pentane, the residue must be disposed of forthwith to avoid the possibility of detonation of the polymer as it dries, as happened on one occasion.

See other CYCLIC PEROXIDES, PEROXYESTERS

3350. 1-Pyrrolidinylcyclohexene
[1125-99-1]

C₁₀H₁₇N




Benzenediazonium-2-carboxylate

See Benzenediazonium-2-carboxylate: 1-Pyrrolidinylcyclohexene

See related ALKENES

3351. cis-Cyclododecene
[2198-20-1]

C₁₀H₁₈




Preparative hazard

See Hydrazine: Metal salts (reference 3)

See other ALKENES

3352. Citronellic acid (3,7-Dimethyl-6-hexenoic acid)
[502-47-6]

C₁₀H₁₈O₂




Ozone

See Ozone: Citronellic acid

See other ORGANIC ACIDS

3353. 1,4-Bis(2,3-epoxypropoxy)butane
[2425-79-8]

C₁₀H₁₈O₄



Trichloroethylene

See Trichloroethylene: Epoxides

See other 1,2-EPOXIDES

3354. Di-2-methylbutyryl peroxide
[1607-30-3]

C₁₀H₁₈O₄




Swern, 1971, Vol. 2, 815

Pure material explodes on standing at ambient temperature.

See other DIACYL PEROXIDES

3355. 1,2-Bis(acetoxyethoxy)ethane (Triethylene glycol diacetate)
[111-21-7]

C₁₀H₁₈O₆



Acetic acid

Kolodner, H. J. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 345-355, Basle, SSCI, 1980

When fractional distillation at 150°C/27 mbar of a mixture of the ester with acetic acid (36% mol) was interrupted by admission of air, smoking and gas evolution occurred. This was attributed to cool flame auto-ignition and detailed investigation confirmed this. Nitrogen should always be used to break vacuum, and further precautions are recommended.

See 2-Ethoxyethyl acetate

See COOL FLAMES

3356. tert-Butyl diperoxyoxalate

[14666-77-4]

C₁₀H₁₈O₆



1. Castrantas, 1965, 17

2. Nutter, D. E. *et al.*, *Chem. Abs.*, 1997, **126**, 263799v


It exploded on removal from a freezing mixture [1]. Now attracts warnings as 'highly explosive' [2].

See other PEROXYESTERS

3357. 3-Buten-1-ynyltriethyllead

[30478-96-7]

C₁₀H₁₈Pb



Zavagorodnii, S. V. *et al.*, *Dokl. Akad. Nauk SSSR*, 1962, **143**, 855 (Engl. transl. 268)


It explodes on rapid heating.

See other ALKYLMETALS, METAL ACETYLIDES

3358. Manganese(II) N,N-diethyldithiocarbamate

[15685-17-3]

C₁₀H₂₀MnN₂S₄




Bailar, 1973, Vol. 3, 872

A pyrophoric solid.

See other PYROPHORIC MATERIALS

3359. 2,2'-Di-tert-butyl-3,3'-bioxaziridine
[54222-33-2]

C₁₀H₂₀N₂O₂



Putnam, S. J. *et al.*, *Chem. Eng. News*, 1958, **36**(23), 46

A sample prepared by an established method and stored overnight at 2°C exploded when disturbed with a metal spatula.

See 2,2''-Ethylenebis(3-phenyloxaziridine)

See other N-O COMPOUNDS, STRAINED-RING COMPOUNDS

†3360. 2-Ethylhexyl vinyl ether
[103-44-6]

BuCHEtCH₂OCH=CH₂

C₁₀H₂₀O

†3361. Isopentyl isovalerate
[659-70-1]

Me₂CHC₂H₄OCO.CH₂CHMe₂

C₁₀H₂₀O₂

3362. '10-Carbon oxime'
[]

(Unknown structure)

C₁₀H₂₁NO

Knies, H., private comm., 1983

A 10-carbon oxime at 60°C was stored in an insulated container, and within 24 h the temperature had risen to 125°C, possibly owing to an exothermic Beckmann rearrangement to an amide. Application of cooling prevented a thermal runaway.

See other OXIMES

3363. Tripropyllead fulminate
[43135-83-7]

Pr₃PbC≡N → O


C₁₀H₂₁NOPb

Houben-Weyl, 1975, Vol. 13.3, 101

An extremely explosive solid.

See related ALKYL METALS, METAL FULMINATES

3364. Decyl nitrite
[1653-57-2]



Hakl, J., *Runaway Reactions*, 1981, Paper 3/L, 4
Initial exothermic decomposition occurs at 115°C.
See other ALKYL NITRITES


†3365. Decane
[124-18-5]



†3366. 2-Methylnonane
[34464-38-5]




3367. Oxodiperoxodipiperidinechromium(VI)
[]



See Oxodiperoxodipyridinechromium(VI)
See other AMMINECHROMIUM PEROXOCOMPLEXES

3368. Diisopentylmercury
[24423-68-5]




Iodine

Whitmore, 1921, 103

Interaction is violent, accompanied by hissing.


See other ALKYLMETALS, MERCURY COMPOUNDS

- †3369. Dipentyl ether
[693-65-2] C₁₀H₂₂O
- BuCH₂OCH₂Bu**
- HCS 1980, 719*
3370. Formyl(triisopropyl)silane
[] C₁₀H₂₂OSi
- (Me₂CH)₃SiCO.H**
- Soderquist, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1992, **114**(25), 10078
Pyrophoric.
See other ALKYL SILANES
3371. Diisopentylzinc
[21261-07-4] C₁₀H₂₂Zn
- (Me₂CHC₂H₄)₂Zn**
- Gibson, 1969, 181
It ignites in air.
See other ALKYL METALS, DIALKYL ZINCS
3372. 4-Ethoxybutyldiethylaluminium
[65235-78-1] C₁₀H₂₃Al
- 
- Bahr, G. *et al.*, *Chem. Ber.*, 1955, **88**, 256
It ignites in air.
See related ALKYL ALUMINIUM ALKOXIDES AND HYDRIDES
3373. Ethoxydiisobutylaluminium
[15769-72-9] C₁₀H₂₃AlO
- EtOAl(CH₂CHMe₂)₂**
- May ignite in air.
See entry ALKYL ALUMINIUM ALKOXIDES AND HYDRIDES
3374. Di(*O*-*O*-*tert*-butyl) ethyl diperoxophosphate
[] C₁₀H₂₃O₆P
- (Me₃COO)₂P(O)OEt**
- Rieche, A. *et al.*, *Chem. Ber.*, 1962, **95**, 385

The liquid deflagrated soon after isolation.

See other *tert*-BUTYL PEROXOPHOSPHATE DERIVATIVES, PHOSPHORUS ESTERS

3375. *N*-Trimethylsilyl-*N*-trimethylsilyloxyacetamide
[87318-59-0]





Oster, T. A. *et al.*, *J. Org. Chem.*, 1983, **48**, 4309

It may decompose spontaneously if too high a distillation temperature is attained.

See other N-O COMPOUNDS

See related ALKYL SILANES

3376. 1,4,8,11-Tetraazacyclotetradecanenickel(II) perchlorate
[15220-72-1]





Barefield, E. K., *Inorg. Chem.*, 1972, **11**, 2274

It exploded violently during analytical combustion.

See TETRAAZAMACROCYCLANEMETAL PERCHLORATES

See other AMMINOMETAL OXOSALTS

3377. 2,4-Di-*tert*-butyl-2,2,4,4-tetrafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine
[58656-11-4]



Preparative hazard

See Butyllithium: 2,2,2,4,4,4-Hexafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine

3378. 1,2-Bis(*tert*-butylphosphino)ethane

[25196-24-9]

C₁₀H₂₄P₂



Benn, R. *et al.*, *Z. Naturforsch. B*, 1987, **41B**, 680–691

The title compound and the bis(diisopropylphosphino) homologue were pyrophoric in air.

See related ALKYLPHOSPHINES

3379. 1,2-Bis(diethylphosphino)ethane

[6411-21-8]

C₁₀H₂₄P₂



Mays, M. J. *et al.*, *Inorg. Synth.*, 1974, **15**, 2

It ignites in air.

See other ALKYLPHOSPHINES

3380. Potassium [8-(4-nitrophenylthio)undecahydrodicarbaundecaborato]undecahydrodicarbaundecaboratocobaltate(1–) (Potassium [(7,8,9,10,11- η)-1,2,3,4,5,6,7,8,9,11-decahydro-10-(4-nitrobenzenethiolato-*S*)-7,8-dicarbaundecaborato(2–)][(7,8,9,10,11- η)undecahydro-7,8-dicarbadeborato(2–)cobaltate(1–)])

[155147-43-6]

C₁₀H₂₅B₁₈CoKNO₂S




See Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato)cobaltate(1–)

See other BORANES

3381. Tris(trimethylsilyl)methylcaesium

[165610-13-9]

C₁₀H₂₇CsSi₃




Eaborn, C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(6), 687

This compound was isolated with 3.5 benzenes of crystallisation as moisture sensitive, pyrophoric, yellow plates.



See other ORGANOMETALLICS

3382. Tetramethylbis(trimethylsilyloxy)digold
[15529-45-0]



Schmidbaur, H. *et al.*, *Inorg. Chem.*, 1966, **5**, 2069
Sublimed crystals decomposed explosively at 120°C.
See related ALKYL METALS *See other* GOLD COMPOUNDS


3383. *trans*-Diammine(1,4,8,11-tetraazacyclotetradecane)chromium(III) perchlorate
[94484-09-0]



Kane-Maguire, N. A. P. *et al.*, *Inorg. Chem.*, 1985, **24**, 597–605
It exploded on heating. Hexafluorophosphate is suggested as a more stable anion for this type of salt.
See other AMMINOMETAL OXOSALTS

3384. Pentakis(dimethylamino)tantalum

[]




Preparative hazard

Chestnut, R. W. *et al.*, *Chem. Eng. News*, 1990, **68**(31), 2
Preparation by reaction of tantalum pentachloride with pure lithium dimethylamide in pentane is unsafe. Initial non-reaction is followed by explosion during manipulation of the slurry. Presence of ether or dimethylamine gives smooth reaction. Stepwise displacement of chlorine, starting with the free amine, is also recommended.

See other N-METAL DERIVATIVES

3385. 4,4'-Bipyridyl-bis(pentaammineruthenium(III) perchlorate)

[69042-75-7]

C₁₀H₃₈Cl₆N₁₂O₂₄Ru₂Sutton, J. E. *et al.*, *Inorg. Chem.*, 1979, **18**, 1018

Use of the perchlorate salt gives a cleaner product than the tosylate, but is more hazardous.

See other AMMINOMETAL OXOSALTS**3386. Decacarbonyldirhenium**

[14285-68-8]


C₁₀O₁₀Re₂**(OC)₅ReRe(CO)₅**

Bailar, 1973, Vol. 3, 953

It tends to ignite in air above 140°C.

See other CARBONYLMETALS**3387. 3,5-Dinitro-2-(picrylazo)pyridine**


[55106-93-9]

C₁₁H₄N₈O₁₀

Explosive.

See entry POLYNITROAZOPYRIDINES**3388. 1-Chloro-2,3-di(2-thienyl)cyclopropenium perchlorate**

[]

C₁₁H₆Cl₂O₄S₂


Komitsu, K. *et al.*, *Tetrahedron Lett.*, 1978, 806

When isolated and dried, a violent explosion occurred. The tetrachloroaluminate salt is safer.

See other NON-METAL PERCHLORATES, STRAINED-RING COMPOUNDS

3389. 1-(3,5-Dinitro-2-pyridyl)-2-picrylhydrazine
[55106-92-8]

C₁₁H₆N₈O₁₀




Explosive.

See entry POLYNITROAZOPYRIDINES

3390. 1-Naphthyl isocyanate
[1984-04-9]

C₁₁H₇NO




Energy of decomposition (in range 250–380°C) measured as 0.31 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other ORGANIC ISOCYANATES

3391. (1-Naphthyl)-1,2,3,4-thiaziazole
[95322-62-6]


C₁₁H₇N₃S



CHETAH, 1990, 182
Shock sensitive.
See other N-S COMPOUNDS

3392. 1-(4-Diaziophenyl)-1,2-dihydropyridine-2-iminosulfinate
[61224-59-7]

C₁₁H₈N₄O₂S




Hoffmann, H. *et al.*, US Pat. 3 985 724, 1977
Diazotisation of 2-*p*-aminophenylsulfonamidopyridine with subsequent adjustment of pH to 3–6 gives the internal diazonium salt, which decomposes violently at 230°C.

See other DIAZONIUM SALTS, N-S COMPOUNDS
See related DIAZONIUM CARBOXYLATES

3393. 2,4-Diethynyl-5-methylphenol
[32229-04-2]


C₁₁H₈O



Myasnikova, R. N. *et al.*, *Izv. Akad. Nauk SSSR*, 1970, **11**, 2637
Oxidation of the phenol or its methyl or propyl ethers in pyridine gave insoluble polymers. Those of the ethers exploded on heating.
See other ACETYLENIC COMPOUNDS, POLYPEROXIDES

3394. 1-(2-Naphthyl)-3-(5-tetrazoly)triazene
[68499-65-0]


C₁₁H₉N₇



Thiele, J., *Ann.*, 1892, **270**, 54; 1893, **273**, 144
It explodes at 184°C.
See other TETRAZOLES, TRIAZENES

3395. (Nitrobenzene)(cyclopentadienyl)iron(II) hexafluorophosphate
[77486-94-3]


C₁₁H₁₀F₆FeNO₂P



Gassman, P. G. *et al.*, *Organometallics*, 1994, **13**(7), 2890
Heating this complex above 200°C may lead to explosion.
See other ORGANOMETALLICS

3396. η-Benzenecyclopentadienyliron(II) perchlorate
[]

C₁₁H₁₁ClFeO₄




Denning, R. G. *et al.*, *J. Organomet. Chem.*, 1966, **5**, 292

The dry material is shock-sensitive and detonated on touching with a spatula.
See other ORGANOMETALLIC PERCHLORATES

3397. Ethyl α -azido-*N*-cyanophenylacetimidate
[51688-25-6]

C₁₁H₁₁N₅O




Petersen, H. J., *J. Med. Chem.*, 1974, **17**, 104

The crude imidoester should be used as a concentrated solution in ether. A small solvent-free sample exploded violently.

See related 2-AZIDOCARBONYL COMPOUNDS

3398. 3,3-Dimethyl-1-(3-quinolyl)triazene
[70324-23-1]

C₁₁H₁₂N₄




Rondevstedt, C. S. *et al.*, *J. Org. Chem.*, 1957, **22**, 201

The crude material decomposes violently if allowed to dry, and purified material explodes at 131°C or during analytical combustion.

See other TRIAZENES

3399. 3-Butyn-1-yl 4-toluenesulfonate
[23418-85-1]

C₁₁H₁₂O₃S




Eglinton, G. *et al.*, *J. Chem. Soc.*, 1950, 3653

The material could be distilled in small amounts at below 0.01 mbar, but exploded on attempted distillation at 0.65 mbar.

See other ACETYLENIC COMPOUNDS, SULFUR ESTERS

3400. *tert*-Butyl 4-nitroperoxybenzoate
[16166-61-3]

C₁₁H₁₃NO₅




Criegee, R. *et al.*, *Ann.*, 1948, **560**, 135

This and the 4-nitrobenzoates of homologous *tert*-alkyl hydroperoxides explode in contact with flame.

See other NITROARYL COMPOUNDS, PEROXYESTERS

†3401. *tert*-Butyl peroxybenzoate
[614-45-9]

C₁₁H₁₄O₃



1. Criegee, R., *Angew. Chem.*, 1953, **65**, 398–399
2. Schnur, R. C., *Chem. Eng. News*, 1981, **59**(19), 3
3. Verhoeff, J., *Experimental Study of the Thermal Explosion of Liquids*, Doctoral thesis, University of Delft, 1983
4. *CHETAH*, 1990, 182

Shortly after interruption of vacuum distillation from an oil-bath at 115°C to change a thermometer, the ester exploded violently and this was attributed to overheating [1]. A commercial sample of undetermined age exploded violently just after vacuum distillation had begun [2]. In an examination of thermal explosion behaviour, the title compound was used as a model compound in autoclave

experiments at low, high or constant pressure. Three stages in the overall process were identified: thermal runaway, initiation, and then explosion, and these are studied and discussed in detail. In the high pressure experiments, maximum rates of pressure rise approaching 1000 kbar/s were observed [3]. It is shock sensitive [4].
See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Copper(I) bromide, Limonene


Wilson, C. W. *et al.*, *J. Agric. Food Chem.*, 1975, **23**, 636

Addition of all the perester in one portion to limonene and catalytic amounts of copper(I) bromide before oxidation had begun, as shown by development of a blue-green colour, led to an explosion.

See other PEROXYESTERS

3402. 1-Chloro-3-phenylpent-1-en-4-yn-3-ol
[33543-69-0]

C₁₁H₉ClO




Acheson, R. M. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1987, 2322. 2326

In spite of very careful temperature control, explosive decomposition always occurred towards the end of vacuum distillation at 133–136°C/1 mbar.

See other ACETYLENIC COMPOUNDS, HALOACETYLENE DERIVATIVES

3403. Ethyl 6-ethoxycarbonyl-3,4-dihydro-1,1,3-trioxo-2-pyrazolo[3,4-d]isothiazole-2-acetate
[126623-77-6]


C₁₁H₁₃N₃O₇S



Burri, K. F., *Helv. Chim. Acta*, 1989, **72**(6), 1416
It is suggested that this compound, and two precursors, can explode.
See Ethyl 4-bromo-1,1,3-trioxoisothiazoleacetate
See other N-S COMPOUNDS

3404. Cyclobutyl 4-methylbenzenesulfonate
[10437-85-1]


C₁₁H₁₄O₃S



It decomposes vigorously at 118°C.
See entry SULFONIC ACID ESTERS *See other* SULFUR ESTERS

3405. Methyl spiro[5-bromobicyclo[2.2.1]heptane 2,2' 1,3-dioxolane]-7-carboxylate
[71155-12-9]


C₁₁H₁₅BrO₄



Perchloryl fluoride
See Perchloryl fluoride: Methyl 2-bromo-5,5-(ethylenedioxy)[2.2.1]bicycloheptane-7-carboxylate

3406. 2-tert-Butyl-3-phenyloxaziridine
[7731-34-2]

C₁₁H₁₅NO




Emmons, W. D. *et al.*, *Org. Synth.*, 1969, **49**, 13

Vacuum distillation of this active oxygen compound is potentially hazardous and precautions are necessary.

See other N-O COMPOUNDS, STRAINED-RING COMPOUNDS

3407. N-Phenylazopiperidine
[16978-76-0]

C₁₁H₁₅N₃



Hydrofluoric acid


Wallach, O. *et al.*, *Ann.*, 1886, **235**, 258; 1888, **243**, 219

Interaction to give fluorobenzene is violent and is not suitable for above 10 g quantities.

See other TRIAZENES

3408. 2-(1,3-Diselena-4,5,6,7-tetrahydroindanylidene)piperidinium perchlorate
[76371-73-8]


C₁₁H₁₆ClNO₄Se₂



See entry 1-(1,3-DISELENONYLIDENE)PIPERIDINIUM PERCHLORATES
See other PERCHLORATE SALTS OF NITROGENOUS BASES

3409. Butyl toluenesulfonate (Butyl 4-methylbenzenesulfonate)
[778-28-9]

C₁₁H₁₆O₃S



Boyer, J. H. *et al.*, *J. Org. Chem.* 1958, **23**, 1051

Eight grams of the crude ester, which had been prepared from tosyl azide and sodium butoxide, then stored for four weeks at 0°C, was to be distilled at 0.2 mm Hg. On the bath reaching 70°C, there was a violent explosion. This was attributed to a toluenesulphonic acid autocatalysed elimination. However, presence of butyl azide as an impurity seems probable.

See other SULFONIC ACID ESTERS, See also AZIDES

**3410. Oxodiperoxy(pyridine)(1,3-dimethyl-2,4,5,6-tetrahydro-2-1H)-
(pyrimidinone)molybdenum**
[128575-71-3]

C₁₁H₁₇MoN₃O₆

Complex structure


Preparative hazard

See Oxodiperoxy(1,3-dimethyl-2,4,5,6-tetrahydro-2-1H)-(pyrimidinone)
molybdenum

See related AMMINECHROMIUMPEROXO COMPLEXES

3411. Di-tert-butyl diazomalonate
[35207-75-1]

C₁₁H₁₈N₂O₄




Ledon, H. J., *Org. Synth.*, 1979, **59**, 70

Although no explosions have been encountered during the distillation of 10 g portions under argon at 54–58°C/0.003 mbar, it should be treated as potentially explosive. Contact with rough or metallic surfaces should be avoided.

See other DIAZO COMPOUNDS

3412. 3-Iodo-4-methoxy-4,7,7-trimethylbicyclo[4.1.0]heptane
[65082-64-6]

C₁₁H₁₉IO




See vic-iodo-alkoxy compounds

See other IODINE COMPOUNDS

3413. 3-Methyl-3-buten-1-ynyltriethyllead
[]

C₁₁H₂₀Pb




Zavgorodnii, S. V. *et al.*, *Doklad. Akad. Nauk SSSR*, 1962, **143**, 855 (Engl. transl. 268)

It explodes on rapid heating.

See other ALKYL METALS, METAL ACETYLIDES

3414. Diethyl-3-diethylaminopropylaluminium
[73566-71-5]

C₁₁H₂₆AlN




491M, 19758, 151

It ignites in air.

See related TRIALKYLALUMINIUMS

3415. *N,N*-Bis(3-aminopropyl)-1,4-diazacycloheptanenickel(II) perchlorate
[70212-12-3] **C₁₁H₂₆Cl₂N₄NiO₈**



Billo, E. J., *Inorg. Nucl. Chem. Lett.*, 1978, **14**, 502
 An explosive solid.
 See other AMMINOMETAL OXOSALTS

3416. 3-Dibutylaminopropylamine
[30734-83-9] **C₁₁H₂₆N₂**
Bu₂N[CH₂]₃NH₂

Cellulose nitrate
 See CELLULOSE NITRATE: amines See other ORGANIC BASES

3417. 1-Trimethylsilyloxy-1-trimethylsilylphosphylidene-2,2-dimethylpropane
(4-(Dimethylethyl)-2,2,6,6-tetramethyl-3-oxa-5-phospha-2,6-disilahept-4-ene)
[78114-26-8] **C₁₁H₂₇OPSi₂**
Me₃SiP=C(CMe₃)OSiMe₃

Becker, G. *et al.*, *Inorg. Synth.*, 1990, **27**, 249
 Extremely oxygen sensitive and pyrophoric.
 See other ALKYLPHOSPHINES


3418. Bis(pentafluorophenyl)aluminium bromide
[4457-91-4] **C₁₂AlBrF₁₀**
(F₅C₆)₂AlBr

Air, or Ether, or Water
 1. Chambers, R. D. *et al.*, *J. Chem. Soc. (C)*, 1967, 2185
 2. Chambers, R. D. *et al.*, *Tetrahedron Lett.*, 1965, 2389
 3. Cohen, C. S. *et al.*, *Advan. Fluorine Chem.*, 1970, **6**, 156
 It ignites in air, explodes during uncontrolled hydrolysis [1], and chars during controlled hydrolysis [2]. When isolated as the etherate, attempts to remove solvent ether caused violent decomposition [3].
 See other HALO-ARYLMETALS

3419. Bis(hexafluorobenzene)cobalt(0)

[]

$C_{12}CoF_{12}$



It decomposes explosively at 10°C

See entry HALOARENEMETAL π -COMPLEXES

3420. Dodecacarbonyltetracobalt

[19212-11-4]

$C_{12}Co_4O_{12}$

$(OC)_{12}Co_4$

Blake, E. J., private comm., 1974


Cobalt catalysts discharged from 'oxo'-process reactors are frequently pyrophoric, owing to the presence of the carbonylcobalt.

See other CARBONYLMETALS, PYROPHORIC CATALYSTS

3421. Bis(hexafluorobenzene)chromium(0)

[59420-01-8]

$C_{12}CrF_{12}$



It decomposes explosively at 40°C.

See entry HALOARENEMETAL π -COMPLEXES

3422. Graphite hexafluorogermanate
[76321-00-0]

C₁₂F₆Ge

C₁₂GeF₆

McCarron, E. M. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1980, 890–891


The solid product of intercalating graphite with germanium tetrafluoride–fluorine mixtures at 20°C is thought to have an oxidising and fluorinating capability close to that of elemental fluorine itself.

See Fluorine

See other OXIDANTS

3423. Bis(pentafluorophenyl)ytterbium
[66080-22-6]

C₁₂F₁₀Yb



Methylcyclopentadiene


Deacon, G. B. *et al.*, *Aust. J. Chem.*, 1985, **38**, 1763

The THF-solvated product of the ligand exchange reaction between bis(pentafluorophenyl)ytterbium and methylcyclopentadiene was explosive. The expected product, bis(methylcyclopentadiene)ytterbium, obtained by another exchange route, is not explosive.

See other ORGANOMETALLICS

3424. Bis(hexafluorobenzene)iron(0)
[37279-26-8]

C₁₂F₁₂Fe




It decomposes explosively at –40°C.

See entry HALOARENEMETAL π -COMPLEXES

3425. Bis(hexafluorobenzene)nickel(0)
[74167-05-8]

C₁₂F₁₂Ni



Alone, or Air, or Hydrogen, or Carbon monoxide
or Allyl bromide, or Trifluoromethyl iodide


Klabunde, K. J. *et al.*, *J. Fluorine Chem.*, 1974, **4**, 114–115

The complex decomposes explosively on slight provocation (flakes falling into the reactor, static charge or uneven warming) and at 70°C with careful heating. At ambient temperature, interaction with air or the reagents above is explosive, but can be controlled by preliminary cooling to -196°C, followed by controlled slow warming.

See entry HALOARENEMETAL π -COMPLEXES

3426. Bis(hexafluorobenzene)titanium(0)
[]

C₁₂F₁₂Ti




It decomposes explosively at -50°C.

See entry HALOARENEMETAL π -COMPLEXES

3427. Bis(hexafluorobenzene)vanadium(0)

[]

C₁₂F₁₂V



It decomposes explosively at 100°C.

See entry HALOARENEMETAL π -COMPLEXES

3428. Dodecacarbonyltriiron

[17658-52-8]

C₁₂Fe₃O₁₂



King, R. B. *et al.*, *Inorg. Synth.*, 1963, **7**, 195


On prolonged storage, pyrophoric decomposition products are formed.

See other CARBONYLMETALS

3429. Silver hexanitrodiphenylamide

[32817-38-2] (ion)

C₁₂H₄AgN₇O₁₂




Taylor, C. A. *et al.*, *Army Ordnance*, 1926, **7**, 68–69

It had been evaluated as a detonator.

See other *N*-METAL DERIVATIVES, POLY-NITROARYL COMPOUNDS, SILVER COMPOUNDS

3430. Calcium picrate
[16824-78-5]


C₁₂H₄CaN₆O₁₄



See Picric acid (reference 3)
See other PICRATES, POLYNITROARYL COMPOUNDS

3431. Bis(2,4,5-trichlorobenzediazo) oxide
[]

C₁₂H₄Cl₆N₄O



Alone, or Benzene


Kaufmann, T. *et al.*, *Ann.*, 1960, **634**, 77

The dry solid explodes under a hammer-blow, or on moistening with benzene.

See other BIS(ARENEDIAZO) OXIDES, HALOARYL COMPOUNDS


3432. Copper(II) picrate
[33529-09-8]

C₁₂H₄CuN₆O₁₄



See Picric acid (reference 2)
See other PICRATES, POLYNITROARYL COMPOUNDS


3433. Mercury(II) picrate
[19528-48-4]



See Picric acid (reference 2)

See other MERCURY COMPOUNDS, PICRATES, POLYNITROARYL COMPOUNDS

3434. 1,3-Dicyano-2-diazo-2,6-azulene quinone
(2-Diazo-2,6-dihydro-6-oxo-1,3-azulenedicarbonitrile)
[53535-47-0]





Nosoe, T. *et al.*, *J. Org. Chem.*, 1995, **60**(18), 5919

The compound sometimes decomposes explosively at 155°C.

See other ARENEDIAZOMIUM OXIDES

3435. Nickel picrate
[63085-06-3]




Bernardi, A., *Gazz. Chim. Ital.*, 1930, **60**, 169–171

It deflagrates or explodes very violently on heating.

See other HEAVY METAL DERIVATIVES, PICRATES, POLYNITROARYL COMPOUNDS

3436. Lead(II) picrate
[6477-64-1]





Belcher, R., *J. R. Inst. Chem.*, 1960, **84**, 377

During the traditional qualitative inorganic analytical procedure, samples containing the lead and salicylate radicals can lead to the formation and possible detonation of lead picrate. This arises during evaporation of the filtrate with nitric acid, after precipitation of the copper–tin group metals with hydrogen sulfide. Salicylic acid is converted under these conditions to picric acid, which in presence of lead gives explosive lead picrate. An alternative (MAQA) scheme is described which avoids this possibility.

See Picric acid (reference 2)

See other HEAVY METAL DERIVATIVES, PICRATES, POLYNITROARYL COMPOUNDS


3437. Zinc picrate
[16824-81-0]



See Picric acid (reference 2)

See other PICRATES, POLYNITROARYL COMPOUNDS

3438. 4,6-Dinitro-1-picrylbenzotriazole
[50892-86-9]




Coburn, M. D., *J. Heterocycl. Chem.*, 1973, **10**, 743–746

The isomeric dinitro derivatives, and some of the mononitro derivatives, are impact-sensitive explosives.

See other POLYNITROARYL COMPOUNDS, TRIAZOLES

3439. 5,7-Dinitro-1-picrylbenzotriazole
[50892-90-5]

$C_{12}H_4N_8O_{10}$




Coburn, M. D., *J. Heterocycl. Chem.*, 1973, **10**, 743–746

The isomeric dinitro derivatives, and some of the mononitro derivatives, are impact-sensitive explosives.

See other POLYNITROARYL COMPOUNDS, TRIAZOLES

**3440. 5,12-Dioxo-4,4,11,11-tetrahydroxy-14H-[1,2,5]oxadiazolo[3,4-e]
[1,2,5]oxadiazolo[3',4':4,5]benzotriazolo[2,1-a]benzotriazolo-b-ium
1,8-dioxide**
[]

$C_{12}H_4N_8O_{10}$




Subramanian, G. *et al.*, *J. Org. Chem.*, 1996, **61**(5), 1898

This, and the tetraketone resulting from dehydration of the gem-diol groups, both explode under a hammer blow.

See other FURAZAN N-OXIDES

3441. 4,6-Dinitro-2-(2,4,6-trinitrophenyl)benzotriazole *N*-oxide
[87604-86-2]

C₁₂H₄N₈O₁₁




Renfrow, R. A. *et al.*, *Aust. J. Chem.*, 1983, **36**, 1847

It is explosive and a super-electrophile, like 4,6-dinitrobenofurazan *N*-oxide.

See other N-OXIDES, POLYNITROARYL COMPOUNDS, TRIAZOLES

3442. 1,3,6,8-Tetranitrocarbazole
[4543-33-3]

C₁₂H₅N₅O₈




Sorbe, 1968, 153

It may readily explode when dry, so it should be stored wet with water (10% is normally added to commercial material).

See other POLYNITROARYL COMPOUNDS

3443. 1,3,5-Triethynylbenzene
[17814-74-3]

C₁₂H₆



Shvartsberg, M. S. *et al.*, *Izv. Akad. Nauk SSSR, Ser. Khim.*, 1963, **110**, 1836


It polymerised explosively on rapid heating and compression.

See other ACETYLENIC COMPOUNDS, POLYMERISATION INCIDENTS

3444. 2,6-Diperchloryl-4,4-diphenoquinone

[]

C₁₂H₆Cl₂O₈



Gardner, D. M. *et al.*, *J. Org. Chem.*, 1963, **28**, 2650


A shock-sensitive explosive.

See other PERCHLORYL COMPOUNDS

3445. Bis(3,4-dichlorobenzenesulfonyl) peroxide

[]

C₁₂H₆Cl₄O₆S₂




Dannley, R. *et al.*, *J. Org. Chem.*, 1966, **31**, 154

The peroxide was too unstable to dry thoroughly; such samples often exploded spontaneously.

See other DIACYL PEROXIDES

3446. Potassium hexaethynylcobaltate(4-)

[]



Alone, or Ammonia


Nast, R. *et al.*, *Z. Anorg. Chem.*, 1955, **282**, 210

It is moderately stable at below $-30^{\circ}C$, is very shock- and friction-sensitive, and explodes violently on contact with water. At ambient temperature, it rapidly forms explosive decomposition products. Its addition compound with ammonia behaves similarly, exploding on contact with air.

See other COMPLEX ACETYLIDES

3447. Iron(III) maleate

[14451-00-4]



Iron(III) hydroxide

Thermally unstable basic iron(III) maleates produced from mixtures of the above may be pyrophoric.


See Phthalic anhydride (reference 2)

Sulfur compounds

See Phthalic anhydride (reference 2)


See other HEAVY METAL DERIVATIVES

3448. Potassium hexaethynylmanganate(3-)
[]



Bailar, 1973, Vol. 3, 855
A highly explosive solid.
See other COMPLEX ACETYLIDES

3449. Bis(2,4-dinitrophenyl) disulfide
[2217-55-2]



Chlorine

MCA Guide, 1972, 317

The disulfide was chlorinated in wet tetrachloroethylene to give dinitrobenzenesulfonyl chloride. When the solvent was being removed by evaporation, the residue exploded. Some of the less stable sulfenyl chloride may have been present.


See 2,4-Dinitrobenzenesulfonyl chloride

See other NON-METAL SULFIDES, POLYNITROARYL COMPOUNDS

3450. 2-Iodo-3,5-dinitrophenyl

[]

C₁₂H₇IN₂O₄



Ethyl sodioacetoacetate

Zaheer, S. H. *et al.*, *J. Indian Chem. Soc.*, 1955, **32**, 491


Interaction of 2-halo-3,5-dinitrophenyls with the sodio ester should be limited to 5–6 g of the title compound to avoid explosions experienced with larger amounts.

See other HALOARYL COMPOUNDS, POLYNITROARYL COMPOUNDS

3451. Potassium 6-*aci*-nitro-2,4-dinitro-1-phenylimino-2,4-cyclohexadienide

[[12245-12-4] (ion)]

C₁₂H₇KN₄O₆



Sudborough, J. J. *et al.*, *J. Chem. Soc.*, 1906, **89T**, 586


The salt of the mono-*aci*-quinonoid form of 2,4,6-trinitrodiphenylamine is explosive, like the *N*-1- and -2-naphthyl analogues.

See other aci-NITROQUINONOID COMPOUNDS

3452. 1,4-Dihydrodicyclopropa[*b*, *g*]naphthalene

[52781-75-6]

C₁₂H₈




Ippen, J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1974, **13**, 736

It undergoes explosive decomposition at 132–133°C, and is a shock-sensitive solid which requires handling with caution.

See other STRAINED-RING COMPOUNDS

3453. Bis(bromobenzenesulfonyl) peroxide
[29342-58-3]



1. Bolte, J. *et al.*, *Tetrahedron Lett.*, 1965, 1529

2. Dannley, R. L. *et al.*, *J. Org. Chem.*, 1966, **31**, 154

The peroxide was too unstable to dry thoroughly; such samples often exploded spontaneously.

See other DIACYL PEROXIDES

3454. 2,2'-Bipyridine *N,N'*-dioxide-dicarbonylrhodium(I) perchlorate
[84578-88-1]




Uson, R. *et al.*, *J. Organomet. Chem.*, 1982, **240**, 433 footnote f

It exploded violently during analytical combustion.

See related AMMINOMETAL OXOSALTS

3455. Thianthrenium perchlorate
[21299-20-7]



Shine, H. J. *et al.*, *Chem. & Ind.*, 1969, 782; *J. Org. Chem.*, 1971, **36**, 2925


A 1–2 g portion of the freshly prepared suction-dried material exploded violently during transfer from a sintered filter. Initiation may have been caused by friction of transfer or rubbing with a glass rod. Preparation of 50–100 mg quantities only is recommended.

See other FRICTIONAL INITIATION INCIDENTS, NON-METAL PERCHLORATES

3456. Bis(4-chlorobenzenediazo) oxide

[]

C₁₂H₈Cl₂N₄O



Alone, or Benzene

Bamberger, E., *Ber.*, 1896, **29**, 464


More stable than unsubstituted analogues, it may be desiccated at 0°C, but is then extremely sensitive and violently explosive. Contact with benzene (even at 0°C) is violent and the reaction may become explosive.

See other BIS(ARENEDIAZO) OXIDES, HALOARYL COMPOUNDS

**3457. 4,4'-Biphenylenebis(diazonium) perchlorate
(1,1'-Biphenyl-4,4'-bisdiazonium perchlorate)**

[]

C₁₂H₈Cl₂N₄O₈




Vorlander, D., *Ber.*, 1906, **39**, 2713–2715

The perchlorate derived from tetrazotised benzidine is unstable and explosive.

See other DIAZONIUM PERCHLORATES

3458. Bis(4-chlorobenzenesulfonyl) peroxide
[1886-69-7]

$C_{12}H_8Cl_2O_6S_2$




Dannley, R. L. *et al.*, *J. Org. Chem.*, 1966, **31**, 154

The peroxide was too unstable to dry thoroughly; such samples often exploded spontaneously.

See other DIACYL PEROXIDES

3459. Endrin (1,2,3,4,10,10-Hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-endo,endo-1,4:5,8-dimethanonaphthalene)
[72-20-8]

$C_{12}H_8Cl_6O$




Diethyl 4-nitrophenyl thionophosphate

See Diethyl 4-nitrophenyl thionophosphate: Endrin

See other 1,2-EPOXIDES

3460. Cobalt(II) picramate
[33393-70-3]



Agrawal, J. P. *et al.*, *Propellants, Explos.*, 1981, **6**, 112–116


The picramate is less stable than the picrate.

See related METAL NITROPHENOXIDES, PICRATES

See other HEAVY METAL DERIVATIVES, POLYNITROARYL COMPOUNDS

3461. Bis(1,*n*-difluorobenzene)chromium(0) isomers

[53504-66-8] [53504-62-4] [39422-94-1] (1,2-, 1,3-, 1,4- resp.)





All isomers unstable and explosive.

See entry HALOARENEMETAL π -COMPLEXES

3462. 3,6-Bis(trifluoromethyl)pentacyclo [6.2.0.0^{2,4}.0^{3,6}.0^{5,7}]dec-9-ene

[24516-71-0], [96570-67-1] homopolymer




Editor's comments

The monomer is the highest energy stage of the Durham synthesis of polyacetylene; since the next stage, metathesized polymer, is known to be potentially explosive, the monomer is not likely to be absolutely safe.

See Poly(acetylene)

3463. Poly([7,8-bis(trifluoromethyl)tetracyclo [4.2.0.0^{2.8}.0^{5.7}]octane-3,4-diy]-1,2-ethenediy]
[102868-70-2]

(C₁₂H₈F₆)_n



Feast, W. J., personal communication, 1993


This, the second stage of the Durham polyacetylene synthesis (and presumably identical to the homopolymer [96570-67-1]) is capable of explosion with one sixth the energy of TNT.

See Poly(acetylene)

See other STRAINED-RING COMPOUNDS

3464. Nickel 2-nitrophenoxide
hydrates: [54217-99-1], [54217-98-0]

C₁₂H₈N₂NiO₆




Bernardi, A., *Gazz. Chim. Ital.*, 1930, **60**, 166

It deflagrates violently or explodes on heating.

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, NITROARYL COMPOUNDS

3465. Bis(2-nitrophenyl) disulfide
[115-00-6]




Anon., *Jahresber.*, 1977, 72–74

This is prepared from 2-nitrochlorobenzene and sodium disulfide in ethanol, and the reaction residue after recovery of the ethanol by steam distillation is known to be of limited thermal stability. On one occasion, a container (atypically with closed vents) of bulked residues totalling over 2 t exploded violently while awaiting collection for disposal. It seems remotely possible that an *aci*-nitro thioquinonimine species derived from 2-nitrobenzenethiol may have been involved.

See Sodium 2-nitrothiophenoxide

See other NON-METAL SULFIDES, POLYNITROARYL COMPOUNDS

3466. 2-trans-1-Azido-1,2-dihydroacenaphthyl nitrate
[68206-59-5]



Trahanovsky, W. S. *et al.*, *J. Amer. Chem. Soc.*, 1971, **93**, 5257


Although several other 1-azido-2-nitrato-alkanes appeared thermally stable, the acenaphthene derivative exploded violently on heating (probably during analytical combustion).

See other ALKYL NITRATES, ORGANIC AZIDES

3467. Bis(4-nitrobenzenediazo) sulfide

[]

C₁₂H₈N₆O₄S



Tomlinson, W. R., *Chem. Eng. News*, 1951, **29**, 5473

The dry material is extremely sensitive and can be exploded by very light friction. It is too sensitive to handle other than as a solution, or as a dilute slurry in excess solvent, and then only on 1 g scale.

See other BIS(ARENEDIAZO) SULFIDES

3468. 4,4'-Biphenylene-bis-sulfonylazide

[4712-19-0]

C₁₂H₈N₆O₄S₂



CHETAH, 1990, 182


Shock sensitive.

See other ACYL AZIDES

3469. Lead 2-amino-4,6-dinitrophenoxide

[11802-21-2]

C₁₂H₈N₆O₁₀Pb



Sobela, B. *et al.*, Czech Pat. CS 245 822, 1987


The lead salt is used as a minor component in compositions for low-power electric detonators.

See other HEAVY METAL DERIVATIVES, POLYNITROARYL COMPOUNDS

See related METAL NITROPHENOXIDES

3470. (2,2-Dichloro-1-fluorovinyl)ferrocene
[58281-24-6]

C₁₂H₉Cl₂FFe




Okohura, K., *J. Org. Chem.*, 1976, **41**, 1493, footnote 14

Two attempts to purify the crude material by vacuum distillation led to sudden exothermic decomposition of the still contents. Distillation in steam was satisfactory.

See related HALOALKENES *See other* ORGANOMETALLICS

3471. Potassium tricarbonyltris(propynyl)molybdate(3-)
[]

C₁₂H₉K₃MoO₃




Houben-Weyl, 1975, Vol. 13.3, 470

It is pyrophoric.

See related CARBONYLMETALS, COMPLEX ACETYLIDES

3472. 4-Nitrodiphenyl ether
[2216-12-8]

C₁₂H₉NO₃




T_{ait24} was determined as 310°C by adiabatic Dewar tests, with an apparent energy of activation of 314 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROARYL COMPOUNDS

3473. Chloromercuriodiphenylgold
[65381-02-4]


C₁₂H₁₀AuClHg



De Graaf, P. W. J. *et al.*, *J. Organomet. Chem.*, 1977, **141**, 346
When dry it explodes violently, even at -20°C .
See other GOLD COMPOUNDS, MERCURY COMPOUNDS

3474. 3,5-Dibromo-7-bromomethylene-7,7a-dihydro-1,1-dimethyl-1H-azirino[1,2-a]indole
[34489-60-6]

C₁₂H₁₀Br₃N



Brown, J. P., *et al.*, *J. Chem. Soc. (C)*, 1971, 3631
A melting point sample exploded at 150°C . Larger quantities showed signs of autocatalytic decomposition from -30°C . This presumably involves release of strain in the aziridine ring.
See other STRAINED-RING COMPOUNDS


3475. Oxybisphenyliodonium bistetrafluoroborate
[119701-48-3]

C₁₂H₁₀B₂F₈I₂O

(PhI⁺)₂O 2BF₄⁻

1. van Look G., *Chem. Eng. News*, 1989 **67**(30), 2
 2. Zhdankin, V. V. *et al.*, *J. Org. Chem.*, 1989, **54**, 2607; 2609
- An explosion [1] of "iodosylbenzene tetrafluoroborate" attributed by *Chem. Abs.* to another structure is more likely to be of this, in view of the original synthetic papers [2].
See Iodosylbenzene tetrafluoroborate
See other IODINE COMPOUNDS

3476. Benzenediazonium tetrachlorozincate
[15727-43-2]





Muir, G. D., *Chem. & Ind.*, 1956, 58

A batch of the 'double salt' exploded, either spontaneously or from slight vibration, after thorough drying under vacuum at ambient temperature overnight. Although dry diazonium salts are known to be light-, heat- and shock-sensitive when dry, the double salts with zinc chloride were thought to be more stable. Presence of traces of solvent reduces the risk of frictional heating and deterioration.

See other DIAZONIUM SALTS

3477. Bis(fluorobenzene)chromium(0)
[42087-90-1]




Unstable, potentially explosive

See entry HALOARENEMETAL π -COMPLEXES

3478. Bis(fluorobenzene)vanadium(0)
[53966-09-9]


C₁₂H₁₀F₂V



Unstable, potentially explosive.
See entry HALOARENEMETAL π -COMPLEXES

3479. Ferrocene-1,1'-dicarboxylic acid
[1293-87-4]


C₁₂H₁₀FeO₄



Phosphoryl chloride
See Phosphoryl chloride: Ferrocene-1,1'-dicarboxylic acid
See related ORGANIC ACIDS *See other* ORGANOMETALLICS

3480. Diphenylmercury
[587-85-9]

C₁₂H₁₀Hg




Chlorine monoxide, or Sulfur trioxide

1. Dreher, E. *et al.*, *Ann.*, 1870, **154**, 127
2. Otto, R., *J. Prakt. Chem. [2]*, 1870, **1**, 183

Chlorine monoxide reacts violently [1], and sulfur trioxide very violently [2], with diphenylmercury.

See other ARYLMETALS, MERCURY COMPOUNDS

3481. Potassium *O,O*-diphenyl dithiophosphate
[3514-82-7]




Arenediazonium salts

See DIAZONIUM SULFIDES AND DERIVATIVES (reference 10)

See related PHOSPHORUS ESTERS

3482. Diphenylmagnesium
[555-55-4]




Air, or Carbon dioxide, or Water

1. Sidgwick, 1950, 234
2. Leleu, *Cahiers*, 1977, (88), 368

It ignites in moist (but not dry) air, reacts violently with water, reaching incandescence [1], and ignites in carbon dioxide [2].

See other ARYLMETALS, DIALKYL MAGNESIUMS

3483. Azobenzene (Diphenyldiazene)
[103-33-3]



HCS 1980, 170


Heat of decomposition is determined as 0.80 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

See other AZO COMPOUNDS

3484. cis-Azobenzene
[17082-12-1]

C₁₂H₁₀N₂




Stull, 1977, 19

It is highly endothermic (ΔH_f° (s) +362.3 kJ/mol, 1.99 kJ/g). The *trans*-isomer [1080-16-6] will be less so.

See other ENDOTHERMIC COMPOUNDS

3485. Azoxybenzene
[495-48-7]

C₁₂H₁₀N₂O



1. Stull, 1977, 18


2. Grever, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

Azoxybenzene is moderately endothermic (ΔH_f° (s) +152.3 kJ/mol, 0.77 kJ/g). (The isolated stereoisomeric components [21650-65-7] '*cis*-', [20972-43-4] '*trans*-' may differ significantly from these values) [1]. Energy of decomposition in range 245–420°C measured by DSC as 1.303 kJ/g by DSC, and T_{ait24} was determined as 191°C by adiabatic Dewar tests, with an apparent energy of activation of 194 kJ/mol [2].

See other ENDOTHERMIC COMPOUNDS

3486. N-Nitrosodiphenylamine
[86-30-6]

C₁₂H₁₀N₂O




Energy of decomposition (in range 300–500°C) measured as 0.65 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other NITROSO COMPOUNDS

3487. 4-Aminophenylazobenzene
[60-09-3]


C₁₂H₁₁N₃



Energy of decomposition (in range 145–365°C) measured as 0.522 kJ/g.
See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)
See other AZO COMPOUNDS

3488. Diphenylphosphorus(III) azide
[4230-63-1]


C₁₂H₁₀N₃P



Allcock, H. R. *et al.*, *Macromolecules*, 1975, **8**, 380
The azide monomer, prepared in solution from the chlorophosphine and sodium azide, should not be isolated because it is potentially explosive.
See other NON-METAL AZIDES

3489. Diphenyl azidophosphate
[26386-83-9]

C₁₂H₁₀N₃O₃P




Nebuloni, M. *et al.*, *J. Calorim., Anal. Therm. Thermodyn.*, 1986, **17**, 398–
Various methods were used to assess the thermal stability and potential hazards in
handling the title compound.

See related NON-METAL AZIDES, PHOSPHORUS ESTERS

3490. Bis(benzeneazo) oxide

[]

C₁₂H₁₀N₄O



Bamberger, E., *Ber.*, 1896, **29**, 460


Extremely unstable, it explodes on attempted isolation from the liquor, or on
allowing the latter to warm to -18°C .

See other BIS(ARENEDIAZO) OXIDES

3491. Bis(benzenediazo) sulfide

[22755-07-3]

C₁₂H₁₀N₄S



Tomlinson, W. R., *Chem. Eng. News*, 1951, **29**, 5473


The wet solid can be exploded by impact or heating, and explodes while handling
in air at ambient temperature. The material is too sensitive to handle other than as
a solution or dilute slurry in excess solvent, and then only on 1 g scale.

See other BIS(ARENEDIAZO) SULFIDES

3492. Diphenyl ether

[101-84-8]

C₁₂H₁₀O




HCS 1980, 731

Chlorosulfuric acid

See Chlorosulfuric acid: Diphenyl ether

3493. Diphenyl sulfoxide
[945-51-7]


C₁₂H₁₀OS



Acetonitrile, Trichlorosilane
See Trichlorosilane: Acetonitrile, etc.

3494. Diphenylselenone
[10504-99-1]


C₁₂H₁₀O₂Se



Krafft, F. *et al.*, *Ber.*, 1896, **29**, 424
It explodes feebly on heating in a test tube.
See related NON-METAL OXIDES

3495. Benzeneseleninic anhydride
[17697-12-0]

C₁₂H₁₀O₃Se₂




Preparative hazard
See Hydrogen peroxide: Diphenyl diselenide

Benzhydrazide
See Benzeneseleninic acid: Hydrazine derivatives
See other ACID ANHYDRIDES

3496. 11,12-Diethynyl-1,4,7,10-tetraoxadispiro[4.0.4.2]dodec-11-ene
[123002-93-7]


C₁₂H₁₀O₄



See *cis*-3,4-Diethynylcyclobut-3-ene-1,2-diol
See Oligo(octacarbondioxide)
See *other* ACETYLENIC COMPOUNDS

3497. Bis(4-hydroxyphenyl) sulfone
[80-09-1]

C₁₂H₁₀O₄S




When prepared by thermal dehydration of 4-hydroxybenzenesulfonic acid, the reaction mixture begins to decompose exothermally around 240°C. Decomposition is delayed but still occurs at lower temperatures (160°C), and the presence of iron reduces the time to maximum rate of decomposition. Above 800 ppm of iron, the time to maximum rate is less than the dehydration reaction time, leading to severe control problems. Improved processing conditions were developed.

See *entry* SELF-ACCELERATING REACTIONS See *other* INDUCTION PERIOD INCIDENTS

3498. Benzenesulfonic anhydride
[512-35-6]

C₁₂H₁₀O₅S₂




Hydrogen peroxide

See Hydrogen peroxide: Benzenesulfonic anhydride

See *other* ACID ANHYDRIDES

3499. Dibzenesufonyl peroxide
[29342-61-8]

C₁₂H₁₀O₆S₂



Alone, or Nitric acid, or Water


1. Davies, 1961, 65
2. Crovatt, L. W. *et al.*, *J. Org. Chem.*, 1959, **24**, 2032
3. Weinland, R. F. *et al.*, *Ber.*, 1903, **36**, 2702

It explodes at 53–54°C [1], and decomposes somewhat violently after storage overnight at ambient temperature, but may be stored unchanged for several weeks at –20°C [2]. It is also shock-sensitive, and explodes with fuming nitric acid, or addition to boiling water [3].

See other DIACYL PEROXIDES

3500. Diphenyldistibene (Stibobenzene)
[5702-61-4]

C₁₂H₁₀Sb₂



Air, or Nitric acid


Schmidt, H., *Ann.*, 1920, **421**, 235

This antimony analogue of azobenzene ignites in air, and is oxidised explosively by nitric acid.

See related ARYLMETALS, AZO COMPOUNDS

3501. Diphenyl diselenide
[1666-13-3]

C₁₂H₁₀Se₂




Hydrogen peroxide

See Hydrogen peroxide: Diphenyl diselenide

3502. Diphenyltin
[6381-06-2]

C₁₂H₁₀Sn




Nitric acid

Nitric acid: Diphenyltin *See other* ARYLMETALS

3503. Lithium diphenylhydridotungstate(2-)

[]

C₁₂H₁₁Li₂W



Sanny, B., *Z. Anorg. Chem.*, 1964, **329**, 221


It is pyrophoric in air.

See related ARYLMETALS, METAL HYDRIDES

3504. Diphenylamine

[122-39-4]

C₁₂H₁₁N




Preparative hazard

See Aniline: Anilinium chloride

3505. Sodium 2-allyloxy-6-nitrophenylpyruvate oxime

[80841-08-3]

C₁₂H₁₁N₂NaO₆



Hydrochloric acid


Lading, P. *et al.*, *Dan. Kemi*, 1981, **62**, 242–244

Conversion of the sodium salt to the free acid by hydrochloric acid treatment led to an explosion. This was attributed to the thermal instability of the acid, and its Claisen rearrangement to 3-allyl-2-hydroxynitrotoluene, identified in the residue.

See other ALLYL COMPOUNDS, NITROARYL COMPOUNDS, OXIMES

3506. 1,3-Diphenyltriazeno
[136-35-6]

C₁₂H₁₁N₃



1. Müller, E. *et al.*, *Chem. Ber.*, 1962, **95**, 1257
2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
3. Grewer, T. *et al.*, *Proc. 4th Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 3, A1–A11, Rugby, IChE, 1983
4. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74
5. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987

It decomposes explosively at the m.p., 98°C [1], and shows a high rate of pressure increase on exothermic decomposition [2]. The heat of decomposition was determined as 1.34 kJ/g by DSC, and T_{ait24} was determined as 88°C by adiabatic Dewar tests, with an apparent energy of activation of 223 kJ/mol [3]. When finely divided, it also shows significant dust explosion hazards, with a maximum explosion pressure of 7.75 bar, and a maximum rate of pressure rise above 680 bar/s [4]. Further work on homogeneous decomposition under confinement has been reported [5].

Acetic anhydride


Heusler, F., *Ber.*, 1891, **24**, 4160

A mixture exploded with extraordinary violence on warming.

See other TRIAZENES

3507. 1,5-Diphenylpentaazadiene
[]

C₁₂H₁₁N₅




Griess, P., *Ann.*, 1866, **137**, 81

The dry solid explodes violently on warming, impact or friction, and *C*-homologues behave similarly.

See other HIGH-NITROGEN COMPOUNDS *See related* TRIAZENES

3508. Diphenylphosphine
[829-85-6]

C₁₂H₁₁P




Ireland, R. F. *et al.*, *Org. Synth.*, 1977, **56**, 47

If the phosphine is spilled onto a paper towel (or other extended surface), it may ignite in air.

See related ALKYLPHOSPHINES

3509. 1,2-Di(3-buten-1-ynyl)cyclobutane
cis [80605-27-2], trans [80605-36-3]

C₁₂H₁₂




Cupery, M. E. *et al.*, *J. Amer. Chem. Soc.*, 1934, **56**, 1167

During isolation of the title product from polymerised 1,5-hexadiene-3-yne, high vacuum distillation must be carried only to a limited extent to prevent sudden explosive decomposition of the more highly polymerised residue.

See other ACETYLENIC COMPOUNDS, POLYMERISATION INCIDENTS

3510. Tetraacrylonitrilecopper(I) perchlorate
[]

C₁₂H₁₂ClCuN₄O₄




Ondrejovic, G., *Chem. Zvesti*, 1964, **18**, 281

Like the copper(II) analogue, it decomposes explosively on heating

See related AMMINOMETAL OXOSALTS *See other* CYANO COMPOUNDS

3511. Bis(η -benzene)chromium(0)
[1271-54-1]

C₁₂H₁₂Cr



Oxygen


Anon., *Chem. Eng. News*, 1964, **42**(38), 55

The orange-red complex formed with oxygen in benzene decomposes vigorously on friction or heating in air.

See related ARYLMETALS See other ORGANOMETALLICS

3512. Bis(η -benzene)iron(0)
[68868-87-1]

C₁₂H₁₂Fe




Timms, P. L., *Chem. Eng. News*, 1969, **47**(18), 43

Prepared in the vapour phase at low temperature, the solid explodes at -40°C .

See related ARYLMETALS See other ORGANOMETALLICS

3513. Potassium tetrakis(propynyl)nickelate(4-)
[]

C₁₂H₁₂K₄Ni



Ammonia

The diammoniate ignites on friction, impact or flame contact.

See entry COMPLEX ACETYLIDES

3514. Bis(η -benzene)molybdenum(0)
[12129-68-9]

C₁₂H₁₂Mo




Silverthorn, W. E., *Inorg. Synth.*, 1977, **17**, 56

The dry crude material is highly pyrophoric in air.

See related ARYLMETALS See other ORGANOMETALLICS

3515. Dimethyl 4-acetamino-5-nitrophthalate
[52412-56-3]

C₁₂H₁₂N₂O₇




Preparative Hazard

See Nitric acid: Dimethyl 4-acetamidophthalate

See other NITROARYL COMPOUNDS

3516. Benzidine (4,4'-Diaminobiphenyl) (4,4'-Diaminobiphenyl)
[92-87-5]

$C_{12}H_{12}N_2$




Nitric acid

See Nitric acid: Benzidine

3517. 1,2-Diphenylhydrazine
[530-50-7]

$C_{12}H_{12}N_2$



Energy of decomposition (in range 135–405°C) measured as 0.442 kJ/g by DSC, and T_{ait24} was determined as 131°C by adiabatic Dewar tests, with an apparent energy of activation of 139 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


Perchloryl fluoride

See Perchloryl fluoride: Nitrogenous bases

See other REDUCANTS

3518. Amminebarium bis(nitrophenylide)
[91410-67-2]

$C_{12}H_{13}BaN_3O_4$




Stevenson, G. R. *et al.*, *J. Org. Chem.*, 1984, **49**, 3444

A 5 day old sample of the crystalline radical complex exploded violently on agitation.

See related N-METAL DERIVATIVES, NITROARYL COMPOUNDS

3519. 1,1-Diphenylhydrazinium chloride
[530-47-2]

C₁₂H₁₃ClN₂




Energy of decomposition (in range 140–200°C) measured as 0.59 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

See other HYDRAZINIUM SALTS

3520. Dimethyl 4-acetamidophthalate
[51832-56-5]

C₁₂H₁₃NO₅



Nitric acid


Anon., *Jahresber.*, 1987, 64

Nitration of the ester with 90% nitric acid was effected according to a literature method. Examination of the reaction mixture by TLC showed that both expected mononitro derivatives had been formed, and it was allowed to stand at ambient temperature overnight. Ice and dichloromethane were then added, and the separated solvent layer was washed with aqueous sodium hydrogen carbonate, dried, and then freed of solvent by evaporation. The viscous oily residue decomposed and ignited in the flask. The preparation was repeated, but worked up immediately and under nitrogen while still cold, and furnished samples of both of the pure isomeric nitro derivatives. Thermal analysis of these showed both to be thermally stable. It was concluded that the decomposition was due to an impurity formed during the 16 h interval between nitration and work-up.

See other NITRATION INCIDENTS

3521. *N*-tert-Butylphthalisoimidium tetrafluoroborate
(3-*N*-tert-Butyliminioisobenzofuranone tetrafluoroborate)
[69338-55-2]

$C_{12}H_{14}BF_4NO_2$




Preparative hazard

See Acetic anhydride: *N*-tert-Butylphthalamic acid, etc.

3522. Bis(methylcyclopentadienyl)peroxoniobium chloride
[77625-37-7]

$C_{12}H_{14}ClNbO_2$



Sala-Pala, J. *et al.*, *J. Mol. Catal.*, 1980, **7**, 142

Occasionally the dry product exploded violently when scraped off the glass container. It invariably exploded when irradiated with a Raman laser beam, even at $-80^{\circ}C$.

See related ORGANOMETALLIC PEROXIDES See other IRRADIATION DECOMPOSITION INCIDENTS

3523. Dianilineoxodiperoxochromium(VI)

[]

$[(PhNH_2)_2CrO(O_2)_2]$

$C_{12}H_{14}CrN_2O_5$


See Oxodiperoxodipyridinechromium(VI)

See other AMMINECHROMIUM PEROXOCOMPLEXES

3524. Bis(2-methylpyridine)sodium

[]

C₁₂H₁₄N₂Na




Sidgwick, 1950, 89

The addition product of sodium and 2-methylpyridine (or pyridine) ignites in air.

3525. 4,4'-Oxybis(benzenesulfonylhydrazide)

[80-51-3]

C₁₂H₁₄N₄O₅S₂



The recently calculated value of 78°C for critical ignition temperature is appreciably lower than the previous value of 90°C.


See entry CRITICAL IGNITION TEMPERATURE (reference 2)

See entry BLOWING AGENTS *See other* N-S COMPOUNDS

3526. Bis(2-hydroxyethyl) terephthalate

[959-26-2]

C₁₂H₁₄O₆




Ethanol, Perchloric acid

See Perchloric acid: Bis(2-hydroxyethyl) terephthalate, etc.

3527. 6-Isopropylideneethiacyclodeca-3,8-diyne
[]


C₁₂H₁₄S



Gleiter, R. *et al.*, *Synthesis*, 1993, (6), 558
The compound exploded during Kugelrohr distillation at about 1 millibar, temperature not stated but probably about 100°C.
See other ACETYLENIC COMPOUNDS

3528. 1,3,4,7-Tetramethylisoindole
[20944-65-4]

C₁₂H₁₅N




Cobalt

Dolphin, D. *et al.*, *Inorg. Synth.*, 1980, **20**, 157
The scale of the procedure described for preparing [octamethyltetrabenzoporphinato (2-)]cobalt(II) by heating the reagents in an evacuated Carius tube at 390°C must not be increased or the tube will explode. Personal protection is also necessary when opening the sealed tube.
See other ORGANIC BASES

3529. Acetoacet-4-phenetidide
[122-82-7]


C₁₂H₁₅NO₃



The fine powder is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)

3530. 2,4,6-Triallyloxy-1,3,5-triazine ('Triallylcyanurate')
[1025-15-6]

$C_{12}H_{15}N_3O_3$




Preparative hazard

See 2,4,6-Trichloro-1,3,5-triazine: Allyl alcohol

See other ALLYL COMPOUNDS

3531. 1,3,5-Tris(1-oxoprop-2-enyl)-hexahydro-1,3,5-triazine
(1,3,5-Triacrylylhexahydro-*sym*-triazine)
[959-52-4]

$C_{12}H_{15}N_3O_3$




Preparative Hazard

See Acrylonitrile: Formaldehyde


3532. 5-*tert*-Butyl-2,4,6-trinitro-1,3-xylene
[81-15-2]

$C_{12}H_{15}N_3O_6$



See Nitric acid: *tert*-Butyl-*m*-xylene, Sulfuric acid
See other POLYNITROARYL COMPOUNDS

3533. 1,3,5-Tris(2,3-epoxypropyl)triazine-2,4,6-trione ('Triglycidylisocyanurate')
[2589-01-7] **C₁₂H₁₅N₃O₆**




Anon., *Loss Prev. Bull.*, 1977, (013), 3

The epoxy resin component is made by a 2-stage process involving reaction of 1-chloro-2,3-epoxypropane (epichlorhydrin) with isocyanuric acid to give the 1,3,5-tris(2-hydroxy-3-chloropropyl) derivative, which is then treated with sodium hydroxide to eliminate hydrogen chloride to form the title compound. One batch contained more than the normal amount of hydrolysable chlorine, and when excess epichlorhydrin was distilled off, the residual material decomposed with explosive violence. It was later established that the abnormal chlorine content was associated with reduced thermal stability, and criteria for hydrolysable chlorine, epoxy content and pH have been set to prevent distillation of off-spec. material.

See other 1,2-EPOXIDES

3534. Pentamethylcyclopentadienyl-bis(thiolatothiazolyl)rhodium
[71178-86-4] **C₁₂H₁₅N₆RhS₄**




Rigby, W. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1979, 378

The material, which is probably polymeric, may detonate.

See other N-S COMPOUNDS, ORGANOMETALLICS

3535. 2,4,6-Tri(2-acetylhydrazino)-1,3,5-trinitrobenzene
[78925-61-8]

C₁₂H₁₅N₉O₉



Hydrochloric acid


Lawrence, G. W. *et al.*, *Tetrahedron Lett.*, 1980, **21**, 1618

Hydrolysis of the acetylhydrazide by 12 M hydrochloric acid at ambient temperature gave a solid product which was surprisingly sensitive to a mild hammer-blow. Formation of a diazo structure was suspected.

See other DIAZONIUM SALTS *See other* POLYNITROARYL COMPOUNDS

3536. Chloro(η^8 1,3,5,7-cyclooctatetraene)(tetrahydrofuran)lutetium
[96504-50-6]

C₁₂H₁₆ClLuO






Wayda, A. L., *Inorg. Synth.*, 1990, **27**, 150

The dry powder may ignite spontaneously in air, partially oxidised samples can explode.

See other ORGANOMETALLICS

3537. Dianilinium dichromate
[101672-09-7]

C₁₂H₁₆Cr₂N₂O₇




Gibson, G. M., *Chem. & Ind.*, 1966, 553

It is unstable on storage.

See other DICHROMATE SALTS OF NITROGENOUS BASES

3538. Tetrakis(pyrazole)manganese(II) sulfate

[]




C₁₂H₁₆MnN₈O₄S

See entry AMMINOMETAL OXOSALTS (reference 9)

3539. 1-Phenylazocyclohexyl hydroperoxide

[950-32-3]

C₁₂H₁₆N₂O₂



See entry α -PHENYLAZO HYDROPEROXIDES (reference 4)

3540. Tris(3-methylpyrazole)zinc sulfate

[55060-80-5]




C₁₂H₁₈N₆O₄SZn

See entry AMMINOMETAL OXOSALTS (reference 9)

3541. 1,2-Bis(2-hydroperoxy-2-propyl)benzene
[29014-32-2]

C₁₂H₁₈O₄




Velenskii, M. S. *et al.*, *Chem. Abs.*, 1974, **81**, 108066

This difunctional analogue of cumyl hydroperoxide appears to be no more hazardous than the latter. Though impact-sensitive, the decomposition was mild and incomplete.

See other ALKYL HYDROPEROXIDES

3542. 1,4-Bis(2-hydroperoxy-2-propyl)benzene
[3159-98-6]

C₁₂H₁₈O₄




Velenskii, M. S. *et al.*, *Chem. Abs.*, 1974, **81**, 108066

This difunctional analogue of cumyl hydroperoxide appears to be no more hazardous than the latter. Though impact-sensitive, the decomposition was mild and incomplete.

See other ALKYL HYDROPEROXIDES

3543. 1,10-Di(methanesulfonyloxy)deca-4,6-diyne
[42404-59-1]

C₁₂H₁₈O₆S₂



CHETAH, 1990, 189


Shock sensitive. The parent diol is not but shows thermal risks on calorimetry.

See other ACETYLENIC COMPOUNDS, SULFUR ESTERS

3544. 3-Acetoxy-4-iodo-3,7,7-trimethylbicyclo[4.1.0]heptane

[]

$C_{12}H_{19}IO_2$




See vic-iodo-alkoxy compounds

See other IODINE COMPOUNDS

3545. Tetraallyl-2-tetrazene (Tetra-2-propenyl-2-tetrazene)

[52999-19-6]

$C_{12}H_{20}N_4$



Houben-Weyl, 1967, Vol. 10.2, 831


It explodes with great violence when heated above its b.p., 113°C/1 bar.

See other ALLYL COMPOUNDS, TETRAZENES

**3546. 3,6-Di(spirocyclohexane)-1,2,4,5-tetraoxane
(7,8,15,16-Tetraoxadispiro[5.2.5.2]hexadecane)**

[183-84-6]

$C_{12}H_{20}O_4$



Dilthey, W. *et al.*, *J. Prakt. Chem.*, 1940, **154**, 219


This dimeric cyclohexanone peroxide explodes on impact.

See other CYCLIC PEROXIDES

3547. 1-Acetoxy-6-oxo-cyclodecyl hydroperoxide

[]

C₁₂H₂₀O₅



Criegee, R. *et al.*, *Ann.*, 1949, **564**, 15


The crystalline compound prepared at -70°C explodes mildly after a few minutes at ambient temperature.

See related ALKYL HYDROPEROXIDES, 1-OXYPEROXY COMPOUNDS

3548. Tetraallyluranium

[28711-64-0]

C₁₂H₂₀U



Bailar, 1973, Vol. 5, 405


Only stable below -20°C , it ignites in air.

See related ALKYL METALS *See other* ALLYL COMPOUNDS

3549. 3-Buten-1-ynyl-diisobutylaluminum

[18864-05-6]

C₁₂H₂₁Al




Petrov, A. A. *et al.*, *Zh. Obsch. Khim.*, 1962, **32**, 1349

It ignites in air.

See other METAL ACETYLIDES, TRIALKYLALUMINIUMS

3550. Perhydro-9b-boraphenalene
[16664-33-8]

C₁₂H₂₁B




Negishi, E.-I. *et al.*, *Org. Synth.*, 1983, **61**, 103–111

The title product and several precursors are highly pyrophoric, and all leaks must be eliminated from vacuum handling equipment.

See related ALKYLBORANES

3551. 6-Azidohexyl 6-azidohexanoate
[84487-84-3]

C₁₂H₂₂N₆O₂



See entry ENERGETIC COMPOUNDS

See other ORGANIC AZIDES

3552. 1-Butoxyethyl 3-trimethylplumbylpropiolate
[21981-95-3]

C₁₂H₂₂O₃Pb




Houben-Weyl, 1975, Vol. 13.3, 80

It is explosive.

See related ALKYLMETALS, METAL ACETYLIDES

3553. Bis(1-hydroxycyclohexyl) peroxide (1,1'-Dioxybiscyclohexanol)
[2407-94-5]

C₁₂H₂₂O₄



1. Stoll, M. *et al.*, *Helv. Chim. Acta*, 1930, **13**, 142
2. *CHETAH*, 1990, 183

Normally stable, it explodes on attempted vacuum distillation [1] and is shock sensitive [2].

See other 1-OXYPEROXY COMPOUNDS

3554. Dihexanoyl peroxide
[2400-59-1]

C₁₂H₂₂O₄




Castrantas, 1965, 17

It explodes at 85°C.

See other DIACYL PEROXIDES

3555. 1-Hydroperoxy-1'-hydroxydicyclohexyl peroxide
(1-[(1-Hydroperoxycyclohexyl)dioxy]cyclohexanol)
[78-18-2]

C₁₂H₂₂O₅




1. Davies, 1961, 74
2. Cardinale, G. *et al.*, *Tetrahedron*, 1985, **45**, 2901

This appears to be a main constituent of commercial 'cyclohexanone peroxide' together with the bis(hydroperoxy) peroxide (below), and is also known to be hazardous [1]. Prepared from cyclohexanone and 30% hydrogen peroxide in presence of hydrochloric acid catalyst, it must be kept moist after isolation as it may explode on drying out [2].

See entry COMMERCIAL ORGANIC PEROXIDES *See other* 1-OXYPEROXY COMPOUNDS

3556. Bis(1-hydroperoxycyclohexyl) peroxide
(Dioxydicyclohexylidenebishydroperoxide)
[2699-12-9]

C₁₂H₂₂O₆




Criegee, R. *et al.*, *Ann.*, 1949, **565**, 17–18

One of the components in 'cyclohexanone peroxide', it explodes violently at elevated temperatures (during vacuum distillation or on exposure to flame).

See entry COMMERCIAL ORGANIC PEROXIDES *See other* 1-OXYPEROXY COMPOUNDS

3557. Lactose
[63-42-3]

C₁₂H₂₂O₁₁



Alkalis

See SUGARS



Oxidants

Treumann, H., *Chem. Abs.*, 1981, **94**, 177551

Explosion hazards of mixtures of lactose monohydrate with potassium chlorate, potassium nitrate or potassium perchlorate are assessed.

See Potassium chlorate: Sugars

3558. Sucrose
[57-50-1]



1. Geysen, W. J. *et al.*, *ASTM Spec. Tech. Publ.* 958, 234–242, Philadelphia, ASTM, 1987

2. Beck, H. *Gefahrstoffe-Reinhalt. Luft*, 1996, **56**(10), 369

The investigation of a sugar dust explosion in a Belgian factory is discussed and preventive and protective measures are described in detail [1]. Sugar dust explosions and safety measures against them are reviewed [2].

Energy of decomposition (in range 280–350°C) measured as 0.48 kJ/g.

See entry THERMOCHEMISTRY AND EYOTHERMIC DECOMPOSITION (reference 2)

See other DUST EXPLOSION INCIDENTS


Nitric Acid

See Nitric Acid: Sucrose

Sulfuric Acid

See Sulfuric Acid: Sucrose


3559. *trans*-Bromoazido(meso-5,12-dimethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) perchlorate
[74190-87-7]



Lawrence, G. A. *et al.*, *Aust. J. Chem.*, 1980, **33**, 274
 Perchlorate salts of the azide complexes, and also of the dichloro analogue, may present an explosion hazard.
See related [14] DIENE-N₄COMPLEXES

3560. 2,4,6,8-Tetraethenyl-2,4,6,8-tetramethylcyclotetrasiloxane.
 [2554-06-5]


C₁₂H₂₄O₄Si₄



Rubinsztajn, S. *et al.*, *Chem. Eng. News.*, 1994, **72**(17), 4
 Hydrosilylation with 2,4,6,8-tetramethylcyclotetrasiloxane has sometimes given uncontrollable exotherms or explosions in the laboratory. Use of safety screens and the lowest possible (<20 ppm) level of platinum catalyst are recommended.
See VINYL-SILOXANES

3561. 1,4,7,10,13,16-Hexaoxacyclooctadecane ('18-Crown-6')
 [17455-13-9]

C₁₂H₂₄O₆



Preparative hazard

1. Cram, D. J. *et al.*, *J. Org. Chem.*, 1974, **39**, 2445
2. Stott, P. E., *Chem. Eng. News*, 1976, **54**(37), 5
3. Gouw, T. H., *Chem. Eng. News*, 1976, **54**(44), 5
4. Stott, P. E., *Chem. Eng. News*, 1976, **54**(51), 5
5. de Jong, F. *et al.*, *J. R. Neth. Chem. Soc.*, 1981, **100**, 449–452


There is a potential explosion hazard during larger scale operation of the published (small-batch) procedure [1]. This arose during thermal decomposition of the crown

ether–potassium chloride complex under reduced pressure, when the crude ether distilled out. The larger batch size involved more extensive heating to complete the decomposition of the complex, and a considerable amount of 1,4-dioxane was produced by cracking and blocked the vacuum pump trap. Admission of air to the overheated residue led to a violent explosion, attributed to autoignition of the dioxane–air mixture. Dioxane has a relatively low AIT (180°C) and rather wide explosive limits. Practical precautions are detailed [2]. Subsequently the observed hazards were attributed to poor distillation procedures [3], but further more detailed information on the experimental procedure was published to refute this view [4]. A safe method of purifying the product without distillation involves formation of complexes with alkaline earth metal alkanedisulfonates [5].

See other APROTIC SOLVENTS

3562. 3,6,9-Triethyl-3,6,9-trimethyl-1,2,4,5,7,8-hexoxonane
[24748-23-0]

C₁₂H₂₄O₆



See 3,6-Diethyl-3,6-dimethyl-1,2,4,5-tetraoxane

See other CYCLIC PEROXIDES

†3563. Dihexyl ether
[112-58-3]

(C₆H₁₃)₂O

C₁₂H₂₆O

3564. Di-2-butoxyethyl ether
[112-73-2]

(BuOC₂H₄)₂O

C₁₂H₂₆O₃

Nitric acid

See Nitric acid: Di-2-butoxyethyl ether

3565. 2,2-Di(*tert*-butylperoxy)butane
[2167-23-9]

(Me₃COO)₂CMeEt

C₁₂H₂₆O₄

Dickey, F. H. *et al.*, *Ind. Eng. Chem.*, 1949, **41**, 1673

The pure material explodes on heating to about 130°C, on sparking or on impact.
See related DIALKYL PEROXIDES

3566. Bis(2-hydroperoxy-4-methyl-2-pentyl) peroxide
[53151-88-5]

C₁₂H₂₆O₆



Leleu, *Cahiers*, 1973, (71), 238


The triperoxide, main component in 'MIBK peroxide', is explosive in the pure state, but insensitive to shock as the commercial 60% solution in dimethyl phthalate. The solution will explode at about 75°C, and slowly liberates oxygen at ambient temperatures.

See KETONE PEROXIDES

See related ALKYL HYDROPEROXIDES, DIALKYL PEROXIDES

3567. Dodecanethiol
[1322-36-7]

C₁₂H₂₆S



HCS 1980, 449

Nitric acid

See Nitric acid: Alkanethiols

See other ALKANETHIOLS

†3568. Triisobutylaluminium
[100-99-2]

C₁₂H₂₇Al



HCS 1980, 937

1. 'Specialty Reducing Agents', Brochure TA-2002/1, New York, Texas Alkyls, 1971
2. Fischer, T. S. *et al.*, *Int. Annu. Conf. ICT*, 1996, **27th** (Energetic Materials), 107


Used industrially as a powerful reductant, it is supplied as a solution in hydrocarbon solvents. The undiluted material is of relatively low thermal stability (decomposing above 50°C) and ignites in air unless diluted to below 25% concentration [1]. A thermal explosion of the compound has been studied from a theoretical standpoint [2].

See other TRIALKYLALUMINIUMS

- 3569. mixo-Tributylborane**
 [122-56-5, 1116-39-8] (n-, iso- resp.) **C₁₂H₂₇B**
Bu₃B, (Me₂CHCH₂)₃B
1. Hurd, D. T., *J. Amer. Chem. Soc.*, 1948, **70**, 2053
 2. Brown, C. A., *Inorg. Synth.*, 1977, **17**, 27
 A mixture of the n- and iso-compounds ignited on exposure to air [1]. In bulk, the liquid does not ignite in air, but may do so when absorbed onto a porous surface like paper, or if spread as a thin film [2]. *See other* ALKYLBORANES
- 3570. Tri-2-butylborane**
 [1113-78-6] **C₁₂H₂₇B**
(MeEtCH)₃B
- Brown, C. A., *Inorg. Synth.*, 1977, **17**, 27
 It ignites in air.
See other ALKYLBORANES
- 3571. Tributylbismuth**
 [3692-81-7] **C₁₂H₂₇Bi**
Bu₃Bi
- Air, or Oxygen
 Gilman, H. *et al.*, *J. Amer. Chem. Soc.*, 1939, **61**, 1170
 It ignites in air and explodes violently in oxygen.
See other TRIALKYLBISMUTHS
- 3572. Tributylgallium**
 [15677-44-8] **C₁₂H₂₇Ga**
Bu₃Ga
- Kovar, R. A. *et al.*, *Inorg. Chem.*, 1980, **19**, 3265
 Tributylgallium and dibutylgallium chloride are probably pyrophoric in air.
See other ALKYLMETALS
- 3573. Tributylindium**
 [15676-66-1] **C₁₂H₂₇In**
Bu₃In
- Houben-Weyl, 1970, Vol. 13.4, 347
 It ignites in air.
See other ALKYLMETALS

3574. Tributyl phosphate
[126-73-8]

C₁₂H₂₇O₄P



Nitric acid, Nitrates
See NUCLEAR WASTES

†**3575. Tributylphosphine**
[998-40-3]


C₁₂H₂₇P

Bu₃P

Steele, A. B. *et al.*, *Chem. Engrg.*, 1969, **66**(4), 160
It ignites in air.
See other ALKYLPHOSPHINES

3576. Tris(2,2-dimethylethyl)triphosphirane
[61695-12-3]

C₁₂H₂₇P₃



Baudler, M. *et al.*, *Inorg. Synth.*, 1989, **25**, 1
The compound is reported as pyrophoric when finely divided.
See related ALKYLPHOSPHINES

3577. Tetraisopropylchromium
[38711-69-2]


C₁₂H₂₈Cr

(Me₂CH)₄Cr

Müller, J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1975, **14**, 761
The finely crystalline form is pyrophoric.
See other ALKYLMETALS

3578. 1,10-Phenanthroline-5,6-dione
[27318-90-7]

C₁₂H₈N₂O₂



Preparative hazard

Black, K. J. *et al.*, *Inorg. Chem.*, 1993, **32**(24), 5591–6

Preparation by oxidising tris(phenanthroline)cobalt(III) tetrafluoroborate with nitric acid in sulfuric acid with potassium bromide catalysis is potentially explosive.

See Nitric acid

3579. Titanium tetraisopropoxide
[546-68-9]

C₁₂H₂₈O₄Ti



HCS 1980, 894

Hydrazine

See Hydrazine: Titanium compounds

See other METAL ALKOXIDES

3580. ‘Tetrapropyldiborane’
[22784-01-6]


C₁₂H₃₀B₂

(Complex mixture)

‘Tetraethyldiborane’ See other ALKYLBORANES

3581. Bis(triethyltin) peroxide
[4403-63-8]


C₁₂H₃₀O₂Sn₂



Sorbe, 1968, 160

Readily decomposed, it explodes at 50°C.
See other ORGANOMETALLIC PEROXIDES

3582. Tetra(3-aminopropanethiolato)trimercury perchlorate
[95247-08-1]





Barrera, H. *et al.*, *Chem. Abs.*, 1985, **102**, 159407

It decomposed violently but non-explosively.

See other MERCURY COMPOUNDS



See related AMMINOMETAL OXOSALTS

3583. 1,2-Bis(triethylsilyl)hydrazine
[13271-97-1]



See entry SILYLHYDRAZINES See related ALKYL SILANES

3584. Tris(trimethylsilylmethyl)aluminium
[41924-27-0]




Tessier-Youngs, C. *et al.*, *Inorg. Synth.*, 1986, **24**, 92, 94

It burns spontaneously in air and reacts violently with water, as does the congener, bis(trimethylsilylmethyl)aluminium bromide.

See related TRIALKYLALUMINIUMS, ALKYL METALS, ALKYL SILANES

3585. Tris(trimethylsilylmethyl)indium
[69833-15-4]

$C_{12}H_{33}InSi_3$




Kopasz, J. P *et al.*, *Inorg. Synth.*, 1986, **24**, 89

It burns spontaneously in air.

See related TRIALKYLALUMINIUMS, ALKYLMETALS, ALKYLSILANES

3586. Triethylsilyl-1,2-bis(trimethylsilyl)hydrazine
[13272-03-2]


$C_{12}H_{34}N_2Si_3$



See entry SILYLHYDRAZINES See related ALKYLSILANES

3587. Dodecamethyltetraplatinum(IV) perchlorate (Trimethylplatinum(IV) perchlorate tetramer)
[23411-89-4] (monomer)

$C_{12}H_{36}Cl_4O_{16}Pt_4$



Preparative hazard


Neruda, B. *et al.*, *J. Organomet. Chem.*, 1977, **131**, 318

The salt is a heat and shock-sensitive explosive. Attempts to prepare it from trimethylplatinum hydroxide and perchloric acid led to violent explosions at higher acid concentrations.

See related ALKYLMETALS, METAL PERCHLORATES See other PLATINUM COMPOUNDS

3588. Tetrakis(trimethylsilyl)diaminodiphosphene
[84521-55-1]

$C_{12}H_{36}N_2P_2$



Niecke, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1983, **22**, 155

It is pyrophoric in air.

See related ALKYL SILANES, PHOSPHINES

3589. Tris(2,3-diaminobutane)nickel(II) nitrate
[66599-67-5]

$C_{12}H_{36}N_8NiO_6$



Billo, E. J. *et al.*, *Inorg. Chim. Acta*, 1978, **26**, L10


The complex is potentially explosive.

See other AMMINOMETAL OXOSALTS

3590. Dodecamethyltetraplatinum(IV) azide (Trimethylplatinum(IV) azide tetramer)

[52732-14-6]

$C_{12}H_{36}N_{12}Pt_4$



von Dahlen, K. H. *et al.*, *J. Organomet. Chem.*, 1974, **65**, 267

The azide is not shock-sensitive, but detonates violently on rapid heating or exposure to flame.

Trimethyl phosphite

Neruda, B. *et al.*, *J. Organomet. Chem.*, 1976, **111**, 241–248

In the exothermic reaction with trimethyl phosphite to give *cis*-dimethylbis(trimethyl phosphito)platinum, the azide must be added to the phosphite in small portions with stirring. Addition of the phosphite to the solid azide led to a violent explosion, probably involving the transitory by-product methyl azide.


See related METAL AZIDES See other PLATINUM COMPOUNDS

3591. μ -Peroxobis[ammine(2,2',2''-triaminotriethylamine)cobalt(III)](4+) perchlorate




1. Yang, C.-H. *et al.*, *Inorg. Chem.*, 1973, **12**, 666
 2. Mori, M. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 619
- The salt exploded at 220°C [1], and other oxosalts of this type (permanganates, possibly nitrates) are also explosive [2].
See related AMMINECHROMIUM PEROXOCOMPLEXES, AMMINOMETAL OXOSALTS

3592. Di[tris(1,2-diaminoethane)cobalt(III)] triperoxodisulfate




- Beacom, S. E., *Nature*, 1959, **183**, 38
 It explodes upon ignition, or after UV irradiation and then heating to 120°C.
See other AMMINOMETAL OXOSALTS

3593. Di[tris(1,2-diaminoethane)chromium(III)] triperoxodisulfate



- Beacom, S. E., *Nature*, 1959, **183**, 38
 It explodes upon ignition, or after UV irradiation and then heating to 115°C.
See other AMMINOMETAL OXOSALTS

3594. Lead(II) trinitrosobenzene-1,3,5-trioxide




- Freund, H. E., *Angew. Chem.*, 1961, **73**, 433
 An air dried sample of the lead salt of trinitrosophoroglucinol exploded when disturbed, possibly owing to aerobic oxidation to the trinitro compound.
See other HEAVY METAL DERIVATIVES, NITROSO COMPOUNDS

3595. 4,5,11,12-Tetraoxo-14H-[1,2,5]oxadiazolo[3,4-*e*][1,2,5]oxadiazolo[3',4':4,5]benzotriazolo[2,1-*a*]benzotriazolo-*b*-ium 1,8-dioxide

[]


C₁₂N₈O₈



See 5,12-Dioxo-4,4,11,11-tetrahydroxy-14H-[1,2,5]oxadiazolo[3,4-*e*][1,2,5]oxadiazolo[3',4':4,5]benzotriazolo[2,1-*a*]benzotriazolo-*b*-ium 1,8-dioxide
See other FURAZAN N-OXIDES

3596. *N,N'*-dichlorobis(2,4,6-trichlorophenyl)urea
[2899-02-7]

C₁₃H₄Cl₈N₂O



Ammonia

Pytlewski, L. L., *Rep. AD-A028841*, 13, Richmond (Va.), USNTIS, 1976
Contact of gaseous ammonia with the *N*-chlorourea, either alone, or mixed with zinc oxide, leads to ignition. The same could happen in contact with conc. aqueous ammonia, solid ammonium carbonate or organic amines.

Dimethyl sulfoxide

1. Pytlewski, L. L., *Rep. AD-A028841*, 13, Richmond (Va.), USNTIS, 1976
2. Owens, C. *et al.*, *Phosphorus, Sulfur*, 1976, **2**, 177–180

Violent ignition occurs on mixing [1]. Interaction is explosive, and the products have been identified and a homolytic mechanism proposed for the reaction [2].

1-(4-Nitrophenylazo)-2-naphthol, Zinc oxide

Pytlewski, L. L., *Rep. AD-A028841*, 13, Richmond (Va.), USNTIS, 1976


Spontaneous combustion in storage (occasionally at high ambient temperatures) of clothing impregnation kits containing the three title compounds was investigated. The *N*-chlorourea when heated evolves chlorine to give the isocyanate and a nitrene. Chlorine and the azo-dye react violently and serve as an initiation source of heat. Zinc oxide is converted to the chloride, which catalyses violently exothermic polymerisation of the isocyanate, the main contribution to the total very high exotherm (some 2 MJ/mol), which leads to vigorous smouldering decomposition of the whole mass, temperatures in excess of 315°C being attained. In absence of the dye, heating a mixture of the urea and zinc oxide at 5°/min leads to ignition at 130°C.

See other *N*-HALOGEN COMPOUNDS

3597. 5-Benzoylbenzenediazonium-2-oxide

[]

C₁₃H₈N₂O₂



Berry, W. L., *J. Org. Chem.*, 1961, **26**, 27


Red crystals, exploding on melting (115°C). It was the highest molecular weight of several homologues to do this. Smaller compounds were often too unstable to isolate, larger are not reported as exploding. Also exploding were: 5-aceto, red, 96°; 5-acetamino, orange, 120°C; 5-phenylmethyl, yellow, temperature unstated.

See other ARENEDIAZONIUM OXIDES

3598. Bis(1-benzo[d]triazolyl) carbonate

[88544-01-8]

C₁₃H₈N₆O₃



Keller, O. *et al.*, *Chimia*, 1985, **39**(2-3), 63

It is explosive in the dry state, but if prepared at below 60°C, it may be handled wet with 30% of 1,1,2-trichloroethane.


See other HIGH-NITROGEN COMPOUNDS, N-O COMPOUNDS, TRIAZOLES

3599. 1-Bromo-1,2-cyclotridecadien-4,8,10-triyn

[17530-57-3]

Leznoff, C. C. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 731

C₁₃H₉Br




It explodes at 85°C and slowly decomposes in the dark at 0°C.

See other HALOACETYLENE DERIVATIVES

3600. 2-Benzylideneamino-4,6-dinitrophenol

[53088-06-5]

C₁₃H₉N₃O₅



Metal salts


See entry 2-ARYLIDENEAMINO-4,6-DINITROPHENOL SALTS

See other POLYNITROARYL COMPOUNDS

3601. 2-(4-Nitrophenoxyazo)benzoic acid

[]

C₁₃H₉N₃O₅




Griess, P., *Ber.*, 1884, **17**, 338

It explodes on heating.

See other ARENEDIAZOATES, N-O COMPOUNDS, ORGANIC ACIDS

3602. 1-(2-Nitrophenyl)-5-phenyltetrazole
[55761-75-8]

C₁₃H₉N₅O₂




Houghton, P. G. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 771

Thermolysis of the undiluted tetrazole is violent, but controlled if dispersed in sand or in solution.

See other NITROARYL COMPOUNDS, TETRAZOLES

3603. Diazidodiphenylmethane
[17421-82-8]

C₁₃H₁₀N₆




Kirchmeyer, S. *et al.*, *Synthesis*, 1983, 301

Potentially explosive (a *gem*-diazide).

See other ORGANIC AZIDES

3604. Sodium diphenylketyl
[3463-17-0]

C₁₃H₁₀NaO




Nitrogen oxide

See Nitrogen oxide: Sodium diphenylketyl

3605. Silver 1-benzeneazothiocarbonyl-2-phenylhydrazide
[12154-56-2]

$C_{13}H_{11}AgN_4S$




Sorbe, 1968, 126

The silver derivative of dithizone decomposes explosively at higher temperatures.

See other AZO COMPOUNDS, SILVER COMPOUNDS

3606. α -(4-Bromophenylazo)benzyl hydroperoxide
[72437-42-4]

$C_{13}H_{11}BrN_2O_2$




See entry α -PHENYLAZO HYDROPEROXIDES (reference 4)

See related ALKYL HYDROPEROXIDES, AZO COMPOUNDS

3607. α -Phenylazo-4-bromobenzyl hydroperoxide
[83844-92-2]

$C_{13}H_{11}BrN_2O_2$

$BrC_6H_4CH(OOH)N=NPh$




See entry α -PHENYLAZO HYDROPEROXIDES (reference 3)

See related ALKYL HYDROPEROXIDES, AZO COMPOUNDS

3608. α -Phenylazo-4-fluorobenzyl hydroperoxide
[83844-91-1]


$C_{13}H_{11}FN_2O_2$



See entry α -PHENYLAZO HYDROPEROXIDES (reference 3)
See related ALKYL HYDROPEROXIDES, AZO COMPOUNDS

3609. α -Phenylazobenzyl hydroperoxide
[2829-31-4]

$C_{13}H_{12}N_2O_2$



Alone, or Acids

1. Swern, 1971, Vol. 2, 19
2. Busch, M. *et al.*, *Ber.*, 1914, **47**, 3277

The phenylhydrazones of benzaldehyde and its homologues, (or of acetone) are readily autoxidised in solution and rearrange to give the azo-hydroperoxides, isolable as solids which may explode after a short time on standing, though not on friction or impact [1]. Contact with flame or with conc. sulfuric or nitric acids also initiates explosion [2].

Phenylhydrazine

Bergman, M. *et al.*, *Ber.*, 1923, **56**, 681

The hydroperoxide reacts violently after warming for a few min. with phenylhydrazine.


See other REDOX REACTIONS

See entry α PHENYLAZO HYDROPEROXIDES

See related ALKYL HYDROPEROXIDES, AZO COMPOUNDS

3610. 1,1,1-Triacetoxy-1,2-benziodoxol-3-one
[87413-09-0]

C₁₃H₁₃IO₈



Plumb, J. B., *Chem. Eng. News*, 1990, **68**(29), 3


The title compound is produced by treatment of 2-iodylbenzoic acid with acetic anhydride in acetic acid, and has found wide application as a mild oxidant ('Dess-Martin periodinane') for 1y and 2y alcohols. Although it appears not to be sensitive to impact, unlike the precursor acid, both explode violently when heated under confinement. The oxidant, on treatment with water is hydrolysed back to the explosive 2-iodylbenzoic acid. Forethought and caution are advised before using these explosive materials on any scale of working.

See 2-Iodylbenzoic acid

See other IODINE COMPOUNDS, OXIDANTS

3611. 3-Ethoxycarbonyl-4,4,5,5-tetracyano-3-trimethylplumbyl-4,5-dihydro-3H - pyrazole
[52762-90-0]

C₁₃H₁₄N₆O₂Pb




Houben-Weyl, 1975, VI. 13.3, 221

It tends to explode on heating.

See related ALKYLMETALS, CYANO COMPOUNDS

3612. 2-(Dimethylaminomethyl)fluoroferrocene
[54747-08-9]

C₁₃H₁₆FFeN




Preparative hazard

See Perchloryl fluoride: 2-Lithio(dimethylaminomethyl)ferrocene

See other ORGANOMETALLICS

3613. Dimethylaminomethylferrocene
[1271-86-9]

C₁₃H₁₇FeN




Nitric acid, Water

See Nitric acid: Dimethylaminomethylferrocene, etc.

See other ORGANOMETALLICS

3614. 2-Methyl-2-[4-(2-methylpropyl)phenyl]oximinoethane
(*p*-Isobutylphenyl- α -methylacetaldehyde oxime)
[58609-72-6]

C₁₃H₁₉NO




P. Cardillo, P., 1992, Personal Communication

On the first full-scale run of a modified process for the nickel catalysed isomerisation of the oxime to the corresponding amide, on 1200 kg scale in toluene solution under reflux in place of the previous water, the reaction overheated (to $>180^{\circ}\text{C}$), pressurised, and then escaped confinement. Investigation showed the explosion to be purely a runaway reaction caused by a slower start than previously. It was recommended that the oxime be charged in portions, and the solvent changed to the higher-boiling xylene.

See other OXIMES

See related ALDEHYDES


3615. Diethyl 1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate
[1149-23-1]



4-Nitropyridine *N*-oxide

See 4-Nitropyridine *N*-oxide: Diethyl 1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate

3616. Iron(II) chelate of bis-*N,N*-(2-pentan-4-ylidene)-1,3-diamino-2-hydroxypropane
[22281-49-8]




Berenbaum, M. B., US Pat. 3 388 141, 1968

The solid chelate is pyrophoric in air, burning to iron(III) oxide.

See other PYROPHORIC MATERIALS

3617. Benzyltriethylammonium permanganate
[68844-25-7]




1. Schmidt, H.-J. *et al. Angew. Chem. (Intern. Ed.)*, 1979, **18**, 69
2. Jäger, H. *et al., Angew. Chem. (Intern. Ed.)*, 1979, **18**, 786–787
3. Schmidt, H.-J. *et al., Angew. Chem. (Intern. Ed.)*, 1979, **18**, 787
4. Leddy, B. P. *et al., Tetrahedron Lett.*, 1980, **21**, 2261–2262
5. *See entry SELF-ACCELERATING REACTIONS*
6. Graefe, J. *et al., Angew. Chem. (Intern. Ed.)*, 1983, **23**, 625

The quaternary oxosalt, previously reported as insensitive to hammer-blows or heating to 100°C for 5 min. [1], exploded violently during vacuum drying at 80°C/1.3 mbar. Detailed work has shown that it explodes at 120°C when heated at 4°/min, or at above 50° after various induction periods, e.g. 14 h/60°, 90 min/80°, 25, 28 min/90°, 7 min/100°C [2]. It is also sensitive to heavy blows [2,3]. It explodes in a m.p. capillary at 80–90°C, but may be stabilised by absorption on alumina [4]. An induction period of 80 h/60°C has also been reported [5]. In a further laboratory accident, material dried for 36 h at only 20°C under high vacuum ignited during transfer from a flask to another container, generating flames 1 m long. Since it undergoes violent decomposition in the absence of heating, extreme care and precautions are obviously necessary. It appears that drying conditions may be critical [5].

See other INDUCTION PERIOD INCIDENTS, QUATERNARY OXIDANTS

3618. Dibutyl-3-methyl-3-buten-1-ynylborane
[]




Davidsohn, W. E., *Chem. Rev.*, 1967, **67**, 75

It ignites in air.

See other ACETYLENIC COMPOUNDS, ALKYLBORANES

3619. Tridecanal

[]

C₁₃H₂₆O

1. Steele, A. B. *et al.*, *Chem. Engrg.*, 1959, **66**(4), 160

2. Urben, P. G., private comm., 1989

Reported to ignite in air [1]. This is typical of medium-range aldehydes, particularly if exposure is increased by sorption on paper or cloth, ignition often occurring within 2 hours [2].

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

3620. Tris(trimethylsilyl)aluminium etherate ([1,1'-Oxybis[ethane]]tris(trimethylsilyl)aluminium)

[75441-10-0]

C₁₃H₃₇AlOSi₃**(Me₃Si)₃Al.OEt₂**

Avery, M. A. *et al.*, *J. Amer. Chem. Soc.*, 1992, **114**(3), 974


This compound was pyrophoric even as a relatively dilute pentane solution (Editor's Note: This was a crude preparation probably containing many other organometallics).

See Tris(trimethylsilyl)aluminium

See other ALKYL METALS, ALKYL SILANES

3621. 1,8-Dihydroxy-2,4,5,7-tetranitroanthraquinone

[]

C₁₄H₄N₄O₁₂

Sorbe, 1968, 153


It is explosive.

See other POLYNITROARYL COMPOUNDS

3622. 2,2'-[1,4-Phenylenebis(azidomethylidene)]bis(propanedinitrile)

[]

C₁₄H₄N₁₀



Moore, J. A., *et al.*, *Macromolecules*, 1993, **26**(5), 916


It exploded when filtered on a sinter, destroying the apparatus.

See related BENZYL COMPOUNDS, CYANO COMPOUNDS, ORGANIC AZIDES

3623. Bis(2,4-dichlorobenzoyl) peroxide

[133-14-2]

C₁₄H₆Cl₄O₄



'Lucidol Data Sheet', Buffalo, Wallace & Tiernan, 1963

Whereas the pure compound is extremely shock-sensitive and decomposes rapidly at 80°C, the commercial 50% dispersion in plasticiser is not shock-sensitive.

Charcoal


Leleu, *Cahiers*, 1980, (99), 279

A mixture with a solution of the peroxide in a dialkyl phthalate plasticiser decomposes exothermically but moderately, (probably catalysed by trace heavy metals in the carbon).

See other COMMERCIAL ORGANIC PEROXIDES, DIACYL PEROXIDES

3624. 1-Nitroanthraquinone
[82-34-8]

C₁₄H₇NO₄




MARS Database, 1998, short report 007

A runaway decomposition in a melting vessel associated with distillation of the crude product burst the vessel and led to a major fire. The short report does not include detailed circumstances. The decomposition was attributed to the catalytic effects of impurities. However, it seems that nitric acid was also present.

See Nitric acid: most sub-entries

3625. Calcium 2-iodylbenzoate
[59643-77-5]

C₁₄H₈CaI₂O₈



1. Unpublished information, 1948

2. Merck, 1976, 55


Formulated granules accidentally dried to below normal moisture content exploded violently [1]. The ammonium salt is similarly unstable to heating [2].

See 2-Iodylbenzoic acid

See other IODINE COMPOUNDS

3626. Dicarbonyl(phenanthroline N-oxide)rhodium(I) perchlorate
[84578-90-5]


C₁₄H₈ClN₂O₇Rh



Uson, R. *et al.*, *J. Organomet. Chem.*, 1982, **240**, 433, footnote f
It exploded violently during microanalytical combustion.
See related AMMINEMETAL OXOSALTS

3627. Copper(II) 3,5-dinitroanthranilate
[58302-41-3]


C₁₄H₈CuN₆O₁₂



See Silver 3,5-dinitroanthranilate
See other POLYNITROARYL COMPOUNDS

3628. Bis(2-azidobenzoyl) peroxide
[20442-99-3]

C₁₄H₈N₆O₄



1. Leffler, J. E., *Chem. Eng. News*, 1963, **41**(48), 45


2. Hoffman, J., *Chem. Eng. News*, 1963, **41**(52), 5

A small sample of crystalline material on a sintered glass funnel detonated with extreme violence when touched with a metal spatula [1]. Static electrical initiation may have been involved [2].

See related DIACYL PEROXIDES, ORGANIC AZIDES, STATIC INITIATION INCIDENTS

3629. Bis(1-benzo[d]triazolyl) oxalate
[89028-37-5]

C₁₄H₈N₆O₄




ReftitleKeller, O. *et al.*, *Chimia*, 1985, **39**(2-3), 63

It is explosive in the dry state, but if prepared at below 50°C, it may be handled wet with 40% of 1,1,2-trichloroethane.

See other HIGH-NITROGEN COMPOUNDS, N-O COMPOUNDS, TRIAZOLES

3630. 9,10-Epidioxyanthracene
[220-42-8]

C₁₄H₈O₂




Dufraisse, C. *et al.*, *Compt. rend.*, 1935, **201**, 428

It decomposes explosively at 120°C.

See other CYCLIC PEROXIDES

3631. 2,2-Biphenyldicarbonyl peroxide (Dibenzo-d,f-1,2-dioxocin-3,8-dione)
[6109-04-2]

C₁₄H₈O₄




Ramirez, F., *J. Amer. Chem. Soc.*, 1964, **86**, 4394

It explodes violently on heating to 70°C or on impact, but can be preserved at low temperature.

See other CYCLIC PEROXIDES, DIACYL PEROXIDES

3632. Anthracene
[120-12-7]

C₁₄H₁₀



HCS 1980, 159

Fluorine

See Fluorine: Hydrocarbons


Other reactants

Yoshida, 1980, 35

MRH values calculated for 13 combinations with oxidants are given.

3633. 2,3:5,6-Dibenzobicyclo[3.3.0]hexane (9,10 Dewar anthracene)
[79403-75-1]

C₁₄H₁₀



Pritschins, W. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 1153

It reverts violently to anthracene at 73°C in a tube sealed under argon.

See other STRAINED-RING COMPOUNDS

3634. Diphenylethyne (Diphenylacetylene)
[501-65-5]

C₁₄H₁₀



Aluminium chloride, Nitrobenzene

1. Eisch, J. J. *et al.*, *Chem. Eng. News*, 1998, **76**(6), 2

2. Eisch, J. J., Personal communication, 1998.

On adding one drop of nitrobenzene to an equimolar, ten millimolar, mixture of the other two solids a violent reaction produced gas and carbonaceous material. This was initially attributed to the oxidative powers of the nitrobenzene [1]. However, diphenylacetylene is a high energy molecule, $\Delta H_f^\circ +315$ kJ/mole. At least 98% of the potential chemical energy present will have been the diphenylacetylene. It is probable that the nitrobenzene merely provided a liquid phase in which the aluminium chloride could interact with the acetylene, catalysing reaction beyond the intended azulene dimerisation product[2].

See other ACETYLENIC COMPOUNDS

3635. Phenyl, phenylethynyliodonium perchlorate


[126208-50-2]

C₁₄H₁₀ClO₄Stang, P. J., *Chem. Eng. News*, **68**(34), 2

This salt is explosive, which was attributed to the iodonium group.

See other ACETYLENIC COMPOUNDS, IODINE COMPOUNDS, NON-METAL PERCHLORATES**3636. Bis(cyclopentadienyl)hexafluoro-2-butynechromium**


[63618-84-8]

C₁₄H₁₀CrF₆Chisholm, M. H. *et al.*, *Synth. React. Inorg. Metal-Org. Chem.*, 1977, **7**, 283

This adduct of chromocene with hexafluoro-2-butyne decomposes after a few mins. at ambient temperature, either under nitrogen or vacuum. The process is autocatalytic and violently exothermic, several explosions having occurred.

See other ACETYLENIC COMPOUNDS, ORGANOMETALLICS**3637. Mercury(II) peroxybenzoate**

[18918-17-7]

C₁₄H₁₀HgO₆


Castrantas, 1965, 19

It explodes if heated above its normal decomposition temperature of 100–110°C.

See other MERCURY COMPOUNDS, PEROXOACID SALTS

3638. 1-Benzoyl-1-phenyldiazomethane
[3469-17-8]


C₁₄H₁₀N₂O



Nenitzescu, C. D. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 497
The material may explode if heated to above 40°C.
See other DIAZO COMPOUNDS

3639. Dibenzoyl peroxide
[94-36-0]

C₁₄H₁₀O₄



(*MCA SD-81*, 1960); *FPA H56*, 1977; *HCS 1980*, 196 (both latter relate to commercial material dispersed in water or plasticiser)

1. *MCA SD-81*, 1960
2. Lappin, G. R., *Chem. Eng. News*, 1948, **26**, 3518
3. Taub, D., *Chem. Eng. News*, 1949, **27**, 46
4. Nametkin, S. S., *Chem. Abs.*, 1931, **25**, 4127₈
5. Nozaki, K. *et al.*, *J. Amer. Chem. Soc.*, 1946, **68**, 1692
6. Fine, D. J. *et al.*, *Combust. Flame*, 1967, **11**, 71–78
7. McCloskey, C. M. *et al.*, *Chem. Abs.*, 1967, **66**, 12613c
8. Uetake, K. *et al.*, *Chem. Abs.*, 1974, **81**, 5175
9. Sekida, O., *Jap. Pat.* 40 220, 1974

10. Anon., *Sichere Chemiearb.*, 1976, **28**, 49

11. Anon., *Jahresber.*, 1980, 66

The dry material is readily ignited, burns very rapidly and is moderately sensitive to heat, shock, friction or contact with combustible materials. When heated above its m.p. (103–105°C), instantaneous and explosive decomposition occurs without flame, but the decomposition products are flammable. If under confinement (or in large bulk), decomposition may be violently explosive [1]. An explosion which occurred when a screw-capped bottle of the peroxide was opened was attributed to friction initiating a mixture of peroxide and organic dust in the cap threads [2]. Waxed paper tubs are recommended to store this and other sensitive solids [3]. Crystallisation of the peroxide from hot chloroform solution involves a high risk of explosion. Precipitation from cold chloroform solution by methanol is safer [4]. Water- or plasticiser-containing pastes of dibenzoyl peroxide are much safer for industrial use.

The explosive decomposition of the solid has been studied in detail [6]. The effect of moisture upon ignitibility and explosive behaviour under confinement was studied. A moisture content of 3% allowed slow burning only, and at 5% ignition did not occur [7]. Thermal instability was studied using a pressure vessel test, ignition delay time, TGA and DSC, and decomposition products were identified [8]. The presence of acyl chlorides renders dibenzoyl peroxide impact-sensitive [9]. There is a further report of a violent explosion during purification of the peroxide by Soxhlet extraction with hot chloroform [10]. Residual traces of the peroxide in a polythene feed pipe exploded when it was cut with a handsaw [11]. The heat of decomposition has been determined as 1.39 kJ/g. The recently calculated value of 69°C for critical ignition temperature coincides with that previously recorded.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

Aniline

Bailey, P. S. *et al.*, *J. Chem. Educ.*, 1975, **52**, 525

Addition of a drop of aniline to 1 g of the peroxide leads to mildly explosive decomposition after a short delay.

N-Bromosuccinimide, 4-Toluic acid

See N-Bromosuccinimide: Dibenzoyl peroxide, etc.

Carbon tetrachloride, Ethylene

Bolt, R. O. *et al.*, *Chem. Eng. News*, 1947, **25**, 1866

Interaction of ethylene and carbon tetrachloride at elevated temperatures and pressures, initiated with benzoyl peroxide as radical source, caused violent explosions on several occasions. Recommended precautions include use of minimum pressure and quantity of initiator, maximum agitation, and presence of water as an inert moderator of high specific heat.

See Ethylene: Carbon tetrachloride

Charcoal

Leleu, *Cahiers*, 1980, (99), 279

At 50°C, a mixture reacts violently, evolving white fumes. (Catalytic decomposition by traces of heavy metals in the charcoal seems likely to be involved.)

N,N-Dimethylaniline

Horner, L. *et al.*, *Chem. Ber.*, 1953, **86**, 1071–1072

The solid peroxide exploded on contact with a drop of dimethylaniline.

Dimethyl sulfide

Pryor, W. A. *et al.*, *J. Org. Chem.*, 1972, **37**, 2885

The rapid decomposition of benzoyl peroxide by dimethyl sulfide is explosive in absence of solvent.

Lithium tetrahydroaluminate

MRH 4.39/15

Sutton, D. A., *Chem. & Ind.*, 1951, 272

One or two attempts to reduce the diacyl peroxide in ether led to a moderately violent explosion.

See other REDOX REACTIONS

Methyl methacrylate

MCA Case History No. 996

Local overheating and ignition occurred when solid benzoyl peroxide was put into a beaker which had been rinsed out with methyl methacrylate. Contact between the peroxide, a powerful oxidant and radical source, and oxidisable or polymerisable materials should only be under controlled conditions.

Vinyl acetate


See Vinyl acetate: Dibenzoyl peroxide, etc.

See other DIACYL PEROXIDES

3640. Bis-3-(2-furyl)acryloyl peroxide

[22978-89-8]

C₁₄H₁₀O₆



Milas, N. A. *et al.*, *J. Amer. Chem. Soc.*, 1934, **56**, 1219


It explodes violently on heating.

See other DIACYL PEROXIDES

3641. N-(3-Methylphenyl)-2-nitrobenzimidyl chloride

[]

C₁₄H₁₁ClN₂O₂




Preparative hazard

See Phosphorus pentachloride: 3'-Methyl-2-nitrobenzanilide

3642. 1,1-Diphenylethylene

[530-48-3]

C₁₄H₁₂



Oxygen

Staudinger, H., *Ber.*, 1925, **58**, 1075


Exposure of the alkene to oxygen at ambient temperature and pressure produces an alkene-oxygen interpolymeric peroxide which explodes lightly on heating. An attempt to react the alkene with oxygen at 100 bar and 40–50°C caused a violent explosion in the autoclave.

See other POLYPEROXIDES See related ALKENES

3643. trans-1,2-Diphenylethylene (Stilbene)

[103-30-0]

C₁₄H₁₂




It is mildly endothermic (ΔH_f° (s) +135.4 kJ/mol, 0.75 kJ/g), and the *cis*-isomer [645-49-8] will be rather more so.

See related ALKENES See other ENDOTHERMIC COMPOUNDS

3644. 5-Chlorotoluene-2-diazonium tetrachlorozincate
[89452-69-0]

C₁₄H₁₂Cl₆N₄Zn



Anon., *ABCM Quart Safety Summ.*, 1953, **24**, 42


A batch containing only half the normal water content (60%) exploded violently during ball-milling. Tests later showed the dry material to be shock-sensitive.

See Benzenediazonium tetrachlorozincate

See other DIAZONIUM SALTS

3645. 3'-Methyl-2-nitrobenzanilide
[50623-64-8]

C₁₄H₁₂N₂O₃




Phosphorus pentachloride

See Phosphorus pentachloride: 3'-Methyl-2-nitrobenzanilide

See other NITROARYL COMPOUNDS

3646. 1,2-Bis(2-nitrophenyl)ethane (2,2'-Dinitrobibenzyl)
[16968-19-7]

C₁₄H₁₂N₂O₄




Preparative hazard

See 2-Nitrotoluene: Alkali

See other POLYNITROARYL COMPOUNDS

3647. 2,2-Diphenyl-1,3,4-thiadiazoline
[79999-60-3]

C₁₄H₁₂N₂S




Kolwisch, I. *et al.*, *J. Amer. Chem. Soc.*, 1981, **103**, 7032

The crystalline solid isolated at -78°C suddenly decomposed with nitrogen evolution at -20°C .

See related AZO COMPOUNDS, N-S COMPOUNDS

3648. α -(4-Bromophenylazo)phenylethyl α -hydroperoxide
[91364-94-2]

C₁₄H₁₃BrN₂O₂




See entry α -PHENYLAZO HYDROPEROXIDES (reference 4)

See related AZO COMPOUNDS, ALKYL HYDROPEROXIDES

3649. Barium *N*-perchlorylbenzylamide
[89521-42-6]

C₁₄H₁₄BaCl₂N₂O₆



It explodes on impact.


See entry PERCHLORYLAMIDE SALTS (reference 2)

See other BENZYL COMPOUNDS, N-METAL DERIVATIVES

3650. Dibenzyl phosphorochloridate ('Dibenzyl chlorophosphate')

[538-37-4]

C₁₄H₁₄ClO₃P



Atherton, F. R. *et al.*, *J. Chem. Soc.*, 1945, 382


It is too unstable to be distilled, and the precursory phosphite also tends to decompose on distillation.

Dibenzyl phosphite See other BENZYL COMPOUNDS, PHOSPHORUS ESTERS

3651. Mercury(II) N-perchlorylbzylamide

[89521-44-8]

C₁₄H₁₄Cl₂HgN₂O₆



It explodes on impact, or on heating above 120°C.


See entry PERCHLORYLAMIDE SALTS (reference 2)

See other BENZYL COMPOUNDS, MERCURY COMPOUNDS, N-METAL DERIVATIVES

3652. Bis(1-methylbenzotriazole)cobalt(II) nitrate

[123668-66-6]


C₁₄H₁₄CoN₈O₆



Zafropoulos, T. F. *et al.*, *Monatsh.*, 1989, **120**(4), 357
Explodes at 260 C.
See other AMMINOMETAL NITRATES, TRIAZOLES

3653. 2-Azoxyanisole
[13620-57-0]

C₁₄H₁₄N₂O₃




Preparative hazard

See 2-Nitroanisole: Sodium hydroxide, etc.
See related AZO COMPOUNDS

3654. Bis(toluenediazo) oxide (1,1-Oxybis(4-methylphenyldiazene))
[90238-04-3]

C₁₄H₁₄N₄O



Alone, or Toluene


Bamberger, E., *Ber.*, 1896, **39**, 452, 458

Extremely unstable, it explodes under its reaction liquor at above -4°C. Very shock- and friction-sensitive, a small sample exploded when dried on a porous tile and set off the moist material some distance away. Contact with toluene, even at -5°C, causes an explosive reaction with flame.

See other BIS(ARENEDIAZO) OXIDES

3655. Dibenzyl ether
[103-50-4]

C₁₄H₁₄O




HCS 1980, 204

Aluminium dichloride hydride
See Aluminium dichloride hydride diethyl etherate

Other reactants
Yoshida, 1980, 330
MRH values calculated for 13 combinations with oxidants are given.
See other BENZYL COMPOUNDS, PEROXIDISABLE COMPOUNDS

3656. Di-4-toluenesulfonyl peroxide
[1886-68-6]


C₁₄H₁₄O₆S₂



1. Bolte, J. *et al.*, *Tetrahedron Lett.*, 1965, **6**, 1929
 2. Dannley, R. L. *et al.*, *J. Org. Chem.*, 1966, **31**, 154
- The peroxide was too unstable to dry thoroughly [1]; such samples often exploded spontaneously [2].
See other DIACYL PEROXIDES

3657. 1-Benzyl-3-(4-tolyl)triazene
[17683-09-9]


C₁₄H₁₅N₃



White, E. H. *et al.*, *Org. Synth.*, 1973, Coll. Vol. 5, 799
Vacuum sublimation at 90–100°C led to a violent explosion.
See other BENZYL COMPOUNDS, TRIAZENES

3658. Dibenzyl phosphite
[17176-77-1]


C₁₄H₁₅O₃P



Atherton, F. R. *et al.*, *J. Chem. Soc.*, 1945, 382; 1948, 1106
It decomposes at 160°C, but prolonged heating at 120°C may have the same effect. Not more than 50 g should be distilled at one time, using high-vacuum conditions (b.p 100–120°C/0.001 mbar) unless a preliminary treatment to remove acidic impurities has been used.
See other BENZYL COMPOUNDS, PHOSPHORUS ESTERS

3659. 1-(4-Phenyl-1,3-diselenonylidene)piperidinium perchlorate
[53808-76-7]


C₁₄H₁₆ClNO₄Se₂



See entry 1-(1,3-DISELENONYLIDENE)PIPERIDINIUM PERCHLORATES

3660. 3,3'-Dimethoxy-4,4'-diaminobiphenyl (*o*-Dianisidine)
[119-90-4]


C₁₄H₁₆N₂O₂



The finely powdered carcinogen is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)
See other ORGANIC BASES

3661. 1-Ferrocenyl-2-methylpropyl azide
[105017-95-6]


C₁₄H₁₇FeN₃



Cushman, M. *et al.*, *J. Org. Chem.*, 1987, **52**, 1517–1521
Heating the title azide at 100°C caused explosive decomposition.
See related ORGANIC AZIDES, ORGANOMETALLICS

3662. 2,10-Dimethyl-1,2,3,4,5,10-hexahydroazepino[3-4,b]indole
[75142-81-3]

C₁₄H₁₈N₂




4-Chlorobenzenesulfonyl azide

See 4-Chlorobenzenesulfonyl azide: 2,10-Dimethyl-1,2,3,4,

3663. α -Pentylcinnamaldehyde (3-Phenyl-2-pentylpropenal)
[122-40-7]

C₁₄H₁₈O




Anon., *Chem. Trade. J.*, 1937, **100**, 362

This is very prone to spontaneous oxidative heating. A mixture with absorbent cotton attained a temperature of 230°C 4 min after exposure to air.

See other ALDEHYDES, PEROXIDISABLE COMPOUNDS

3664. 5-*p*-Chlorophenyl-2,2-dimethyl-3-hexanone
[55058-76-9]

C₁₄H₁₉ClO




Peroxyacetic acid

See Peroxyacetic acid: 5-*p*-Chlorophenyl-2,2-dimethyl-3-hexanone

3665. Bis(2-dimethylaminopyridine N-oxide)copper(II) perchlorate

[67517-99-7]

$C_{14}H_{20}Cl_2CuN_4O_{10}$



West, D. X., *Inorg. Nucl. Chem. Lett.*, 1978, **14**, 155–159


It decomposes explosively at 270°C without prior weight loss.

See related AMMINEMETAL OXOSALTS

3666. 2,6-Di-*tert*-butyl-4-nitrophenol

[728-40-5]

$C_{14}H_{21}NO_3$



1. *ASESB Expl. Incid. Rept.*, 1961, 24

2. Barnes, T. J. *et al.*, *J. Chem. Soc.*, 1961, 953


A sample of the compound exploded violently after short heating to 100°C. Although this was attributed to presence of polynitro derivatives [1], the thermal decomposition of this type of nitro compound is known [2].

See other NITROARYL COMPOUNDS

3667. Dicyclohexylcarbonyl peroxide

[4904-55-6]

$C_{14}H_{22}O_4$



Castrantas, 1965, 17
Large quantities may explode without apparent reason.
See other DIACYL PEROXIDES

3668. 2,2'-Azobis(2,4-dimethylvaleronitrile)
[4419-11-8]




$\text{C}_{14}\text{H}_{24}\text{N}_4$

Yoshida, T., private comm., 1984
The low-melting solid of limited thermal stability (max. safe storage temperature 20°C, self-heating decomposition detected above 30°C) will explode on heavy impact (150 kg cm), but is of low sensitivity to ignition. However, 3 cases of spontaneous decomposition in storage or transportation have been noted.
See entry BLOWING AGENTS *See other* AZO COMPOUNDS, CYANO COMPOUNDS

3669. 1-Acetoxy-1-hydroperoxy-6-cyclodecanone
[]

$\text{C}_{14}\text{H}_{24}\text{O}_5$



Criegee, R. *et al.*, *Ann.*, 1949, **564**, 9
It explodes on removal from a freezing mixture.
See other 1-OXYPEROXY COMPOUNDS

3670. Bis(dipropylborino)acetylene
[]



$\text{C}_{14}\text{H}_{28}\text{B}_2$

Hartmann, H. *et al.*, *Z. Anorg. Allgem. Chem.*, 1959, **299**, 179
Both n- and iso-propyl derivatives ignite in air.
See other ACETYLENIC COMPOUNDS, ALKYLBORANES


†**3671. Tetradecane**
[629-59-4]



$\text{C}_{14}\text{H}_{30}$

3672. Acetylenebis(triethyllead)
[5120-07-0]


C₁₄H₃₀Pb₂



Beerman, C. *et al.*, *Z. Anorg. Chem.*, 1954, **276**, 20
It is very sensitive to heat, oxygen or light, and should not be dried.
See related ALKYLMETALS, METAL ACETYLIDES

3673. 1,2-Bis-(di-2-propylphosphino)ethane
[87532-69-2]


C₁₄H₃₂P₂



Benn, R. *et al.*, *Z. Naturforsch. B*, 1986, **41 B**, 680–691
The title ligand, and the bis(*tert*-butylphosphino) lower homologue are pyrophoric in air.
See related ALKYLPHOSPHINES

3674. Acetylenebis(triethyltin)
[994-99-0]

C₁₄H₃₀Sn₂




Stannic chloride
Beerman, C. *et al.*, *Z. Anorg. Chem.*, 1954, **276**, 20
The product of interaction is highly explosive.
See related ALKYLMETALS, METAL ACETYLIDES

3675. *N,N'*-Di-*tert*-butyl-*N,N'*-bis(trimethylsilyl)diaminophosphene

[85923-32-6]

$C_{14}H_{36}N_2P_2Si_2$



Niecke, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1983, **22**, 486

Highly pyrophoric, like the tetrakis(trimethylsilyl) analogue.

See related ALKYL SILANES, PHOSPHINES

3676. Heptakis(dimethylamino)trialuminium triboron pentahydride

[28016-59-3]

$C_{14}H_{47}Al_3B_3N_7$

Complex structure

Hall, R. E., *et al.*, *Inorg. Chem.*, 1969, **8**, 270

The crystalline solid ignites in air.


See other COMPLEX HYDRIDES

See related BORANES

3677. 5,6-Diphenyl-1,2,4-triazine-3-diazonium tetrafluoroborate 2-oxide

[98571-91-6]

$C_{15}H_{10}BF_4N_5O$




Jovanovic, V., *Heterocycles*, 1985, **23**, 1969–1981

The isolated diazonium salt exploded when exposed to air, and also on impact.

See other DIAZONIUM TETRAHALOBORATES, HIGH-NITROGEN COMPOUNDS, *N*-OXIDES

3678. Pyrocatecholato(2-)-(quinolin-8-olato-N,O)-trioxygenido(2-)phosphorus
[82434-14-8] **C₁₅H₁₀NO₆P**



Koenig, M. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 422


The hexavalent phosphorus ozonide, stable at -20°C , exploded in contact with air or on warming to ambient temperature.

See other OZONIDES

See related PHOSPHORUS ESTERS

3679. 3-Chloro-1,3-diphenylcyclopropene
[20421-00-5]

C₁₅H₁₁Cl



Preparative hazard


Padwa, A. *et al.*, *J. Org. Chem.*, 1969, **34**, 2732

If the crude title product (from interaction of 3-chloro-3-phenyldiaziridine and diphenylacetylene) is not purified immediately after preparation, violent decomposition occurs.

See other STRAINED-RING COMPOUNDS

3680. Diphenylcyclopropenylium perchlorate
[37647-36-2]

C₁₅H₁₁ClO₄




Hughes, R. P. *et al.*, *Organometallics*, 1985, **4**, 1761–1766
It should not be air dried, and as an explosive salt it must be handled with care.
See other NON-METAL PERCHLORATES

3681. 4(4'-Bromobenzoyl)acetanilide

[]

C₁₅H₁₂BrNO₂




Dimethyl sulfoxide

See Dimethyl sulfoxide: 4(4'-Bromobenzoyl)acetanilide

3682. 1,2,3,4-Tetrachloro-7,7-dimethoxy-5-phenylbicyclo[2.2.1]hepta-2,5-diene

[15584-65-3]

C₁₅H₁₂Cl₄O₂




Hoffmann, R. W. *et al.*, *Tetrahedron*, 1965, **21**, 900

Pyrolysis of the material at 130°C under nitrogen at low pressure to give tetra-methoxyethylene may be explosive if more than 25 g is used.

3683. Tris(cyclopentadienyl)cerium
[1298-53-9]

C₁₅H₁₅Ce




Preparative hazard

See Sodium nitrate: Tris(cyclopentadienyl)cerium

See other ORGANOMETALLICS

3684. Tris(cyclopentadienyl)plutonium
[12216-68-9]

C₁₅H₁₅Pu




Bailar, 1973, Vol. 5, 407

It ignites in air.

See other ORGANOMETALLICS

3685. Tris(cyclopentadienyl)uranium
[54007-00-0]

C₁₅H₁₅U




Bailar, 1973, Vol. 5, 407

It ignites in air.

See other ORGANOMETALLICS

3686. 2,2-Bis(4-hydroxyphenyl)propane
[80-05-7]

C₁₅H₁₆O₂




The finely powdered resin component (Bisphenol-A) is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

3687. *trans*-Aquadioxo(terpyridine)ruthenium(2+) diperchlorate

[]

C₁₅H₁₇Cl₂N₅O₁₀Ru




Meyer, T. J. *et al.*, *Inorg. Chem.*, 1992, **31**(8), 1375

The complex sometimes explodes when scraped against a glass frit with a metal spatula.

See other AMMINEMETAL OXOSALTS

3688. Tricyclopentadienyluranium tetrahydroaluminate
[107633-87-4]

$C_{15}H_{19}AlU$




Ossola, S. *et al.*, *J. Organomet. Chem.*, 1986, **310**(1), C1

It ignites in air, and reacts with benzene, dimethoxyethane, THF and toluene.

See related COMPLEX HYDRIDES, HEAVY METAL DERIVATIVES, ORGANOMETALLICS, PYROPHORIC MATERIALS

3689. *cis*-Dicarbonyl(cyclopentadienyl)cyclooctenemanganese
[49716-47-4]

$C_{15}H_{19}MnO_2$



Butler, I. S. *et al.*, *J. Inorg. Nucl. Chem.*, 1978, **40**, 1937

Overheating during high-vacuum sublimation may produce pyrophoric manganese which explodes on opening the sublimer to atmosphere.

See PYROPHORIC METALS *See other* ORGANOMETALLICS

3690. Tris(2,4-pentanedionato)molybdenum(III)
[14284-90-3]

$C_{15}H_{21}MoO_6$


Larson, M. L. *et al.*, *Inorg. Synth.*, 1966, **8**, 153

It rapidly oxidises in air, sometimes igniting.

See related ORGANOMETALLICS

3691. 2,6-Di-*tert*-butyl-4-cresol
[120-37-0]


C₁₅H₂₄O



The finely powdered anti-oxidant is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)

3692. *tert*-Butyl 1-adamantaneperoxyoxycarboxylate
[21245-43-2]


C₁₅H₂₄O₃



Razuvajev, G. A. *et al.*, *Tetrahedron*, 1969, **25**, 4925
It explodes on heating to 90–100°C.
See other PEROXYESTERS

3693. Tri(spirocyclopentane)1,1,4,4,7,7-hexoxonane
[4884-18-8]


C₁₅H₂₄O₆



Bjorklund, G. H. *et al.*, *Trans. R. Soc. Can. (Sect. III)*, 1950, **44**, 25
A violent explosive, very sensitive to shock, friction and rapid heating.
See Hydrogen peroxide: Ketones, etc.
See other CYCLIC PEROXIDES

3694. 2,4,6-Tris(dimethylaminomethyl)phenol
[90-72-2]

C₁₅H₂₇N₃O




Cellulose nitrate

See CELLULOSE NITRATE: amines *See other* ORGANIC BASES

3695. Hexaethyltrialuminium trithiocyanate
[17548-36-6]


C₁₅H₃₀Al₃N₃S₃



Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1967, **6**, 947
On heating at 210° under vacuum it disproportionates explosively, but smoothly at 180°C.
See related ALKYLALUMINIUM HALIDES, METAL CYANATES

3696. 1,2-Bis(trimethylsilyl)trimethylsilylhydrazine
[13272-04-3]


C₁₅H₄₀N₂Si₃



See entry SILYLHYDRAZINES See related ALKYL SILANES

3697. 1-(4-Chloro-2-nitrobenzeneazo)-2-hydroxynaphthalene
[6410-13-5]

C₁₆H₁₀ClN₃O₃




Lead chromate

See Lead chromate: Azo-dyestuffs

See other AZO COMPOUNDS

3698. μ -Oxo-*I,I*-bis(trifluoroacetato-*O*)-*I,I*-diphenyldiodine(III)
[91879-79-7]

C₁₆H₁₀F₆I₂O₅



Boutin, R. H. *et al.*, *J. Org. Chem.*, 1984, **49**, 4278, 4284


It detonated at 220°C during determination of the m.p.

See [*I,I*-Bis(trifluoroacetoxy)iodo]benzene

See other IODINE COMPOUNDS

3699. Potassium bis(phenylethynyl)palladate(2-)
[66986-75-2]

C₁₆H₁₀K₂Pd




Air, or Water

Immediately pyrophoric in air, and explosive decomposition occurs with aqueous reagents; the sodium salt is similar.

See entry COMPLEX ACETYLIDES (reference 2)

3700. Potassium bis(phenylethynyl)platinate(2-)
[]

C₁₆H₁₀K₂Pt




Air, or Water

Immediately pyrophoric in air, and explosive decomposition occurs with water; the sodium salt is similar.

See entry COMPLEX ACETYLIDES (reference 2)

3701. mixo-Dimethoxydinitroanthraquinone
[6407-56-3] (1,5,4,8-)
[53939-55-2] (1,8:4,5-)

C₁₆H₁₀N₂O₈



Sulfuric acid

Hildreth, J. D., *Chem. & Ind.*, 1970, 1592

During hydrolysis of crude dimethoxy compound by heating in sulfuric acid, a runaway exothermic decomposition occurred causing vessel failure. Experiment showed a threshold decomposition temperature of 150–155°C, and an oxidising effect of the nitro groups, yielding CO and CO₂ above 162°C.

Oleum

White, D. L. *et al.*, Ger. Offen. 2 451 569, 1975


In an alternative method of preparing *miso*-dihydroxyanthraquinonedisulfonic acid by heating the title compound with oleum to effect simultaneous hydrolysis, denitration and sulfonation of the nucleus, there is the possibility of formation of methyl nitrate from the scission fragments.

See other POLYNITROARYL COMPOUNDS

3702. 1-(2,4-Dinitrobenzeneazo)-2-hydroxynaphthalene (Pigment orange 5)

[3468-63-1]

C₁₆H₁₀N₄O₅



The solid deflagrates at 1.8 cm/min.

See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 3)

See other DEFLAGRATION INCIDENTS

Lead chromate


See Lead chromate: Azo-dyestuffs

See other AZO COMPOUNDS, POLYNITROARYL COMPOUNDS

3703. Bis(cyclopentadienyl)pentafluorophenylzirconium hydroxide

[]

$C_{16}H_{11}F_5OZr$



Chaudhari, M. A. *et al.*, *J. Chem. Soc. A*, 1966, 840


Decomposition on heating in air above 260°C is sometimes explosive.

See related HALO-ARYLMETALS *See other* ORGANOMETALLICS

3704. Copper 1,3,5-octatrien-7-yneide

[]

$C_{16}H_{14}Cu_2$



Georgieff, K. K. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 5495


It deflagrates on heating in air.

See other METAL ACETYLIDES

3705. Bis(cyclopentadienyl)phenylvanadium

[12212-56-5]

$C_{16}H_{15}V$




Gibson, 1969, 178

It ignites in air.


See related ARYLMETALS *See other* ORGANOMETALLICS

3706. Bis(cyclooctatetraenyl)cerium (Bis(η^8 -1,3,5,7-cyclooctatetraene)cerium)
[37205-27-9] **C₁₆H₁₆Ce**



Edelmann, F. T. *et al.*, *Angew. Chem. (Int.)*, 1994, **33**(15/16), 1618
This sandwich compound, nominally the Ce(IV) derivative of cyclooctatetraene(2-) is highly pyrophoric.
See other ORGANOMETALLICS

3707. 2,2'-(1,2-Ethylenebis)3-phenyloxaziridine
[54222-34-3] **C₁₆H₁₆N₂O₂**




MCA Case History No. 2175

The washed crude product from oxidation of 1,2-dibenzylideneaminoethane in methylene chloride solution exploded violently during vacuum evaporation at a relatively low temperature.
See 2,2'-Di-*tert*-butyl-3,3'-bioxaziridine
See other N-O COMPOUNDS, STRAINED-RING COMPOUNDS

3708. Bis(cyclooctatetraene)uranium(0)
[11079-26-8]

C₁₆H₁₆U




Streitwieser, A. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 7364

It inflames in air.

See other ORGANOMETALLICS

3709. 2-(4-Bromophenyl)-2-propyl 1-(1,1-dimethyl-2-pentyn-4-enyl) peroxide
[88519-63-5]


C₁₆H₁₉BrO₂



See 2-(4-Chlorophenyl)-2-propyl 1-(1,1-dimethyl-2-pentyn-4-enyl) peroxide
(below)

See other ACETYLENIC PEROXIDES, DIALKYL PEROXIDES

3710. 1,4,7-Trithia[7]ferrocenophene–acetonitrilecopper(I) perchlorate
[110725-79-6] **C₁₆H₁₉ClCuFe₂NO₄S₃**




Sato, M. *et al.*, *Bull. Chem. Soc. Japan*, 1986, **59**, 3611–3615

It exploded on heating.

See other AMMINOMETAL OXOSALTS, NON-METAL PERCHLORATES

3711. 2-(4-Chlorophenyl)-2-propyl 1-(1,1-dimethyl-2-pentyn-4-enyl) peroxide
[] **C₁₆H₁₉ClO₂**




Vilenskaya, M. R. *et al.*, *Zh. Org. Khim.*, 1981, **17**, 959–961

The acetylenic peroxide and its bromo analogue explode at 120–130°C.

See other ACETYLENIC PEROXIDES, DIALKYL PEROXIDES

3712. O-O-tert-Butyl diphenyl monoperoxophosphate
[20914-03-0] **C₁₆H₁₉O₅P**



Rieche, A. *et al.*, *Chem. Ber.*, 1962, **95**, 385


The material deflagrated as a solid, or decomposed exothermically in its reaction mixture soon after preparation.

See other *tert*-BUTYL PEROXOPHOSPHATE DERIVATIVES, PHOSPHORUS ESTERS

3713. *N*-Methyl-*p*-nitroanilinium 2(*N*-methyl-*N*-*p*-nitrophenylaminosulfonyl) ethylsulfate

[]

C₁₆H₂₀N₄O₁₀S₂



There is a preparative hazard for this reaction product of *N*-methyl-*p*-nitroaniline and 'carbonyl sulfate'.


See 2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide: *N*-Methyl-4-nitroaniline, etc.

See related NITROARYL COMPOUNDS, SULFUR ESTERS

3714. Dibutyl phthalate

[84-74-2]

C₁₆H₂₂O₄




Chlorine

See Chlorine: Dibutyl phthalate

3715. Di-*tert*-butyl diperoxyphthalate
[2155-71-7]


C₁₆H₂₂O₆



Castrantas, 1965, 17
It is shock-sensitive.
See other PEROXYESTERS

3716. Tetrakis(3-methylpyrazole)cadmium sulfate
[55060-82-7]


C₁₆H₂₄CdN₈O₄S



See entry AMMINEMETAL OXOSALTS (reference 9)

3717. Tetrakis(3-methylpyrazole)manganese(II) sulfate
[55060-79-2]


C₁₆H₂₄MnN₈O₄S



See entry AMMINEMETAL OXOSALTS (reference 9)

3718. Bis-*O*, *N*[(*N*'-pent-2-en-2-oxo-4-ylidene)-*N*,*S*-dimethyldithiocarbazate] copper(II) perchlorate

[]



Preparative hazard

Akbar Ali, M. *et al.*, *J. Inorg. Nucl. Chem.*, 1978, **40**, 452
Attempts to prepare this and the corresponding *S*-benzyl complex gave highly explosive products. It appears that the ligand hydrolysed and that the bis(*N*,*S*-dialkyldithiocarbazate)copper perchlorates were obtained and it was these which exploded.

See related AMMINEMETAL OXOSALTS

3719. Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato) cobaltate(1-)

[155147-44-7]

$C_{16}H_{28}B_{18}CoKN_2O_4S_2$



Knyazev, S. P. *et al.*, *Chem. Abs.*, 1994, **120** 298911d

Nitro compounds of this series are reported as exploding on grinding or on rapid heating to $>200^\circ C$. The equivalent ferrate and a mono(2-nitrophenyl) substituted bisboratocobaltate were also prepared.

See other BORANES

3720. Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato) ferrate(1-)

[155147-41-4]

$C_{16}H_{28}B_{18}FeKN_2O_4S_2$



See Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato) cobaltate(1-)

See other BORANES

3721. Bis (2-phenyl-1,2-dicarbododecaborane(12)-1-yl)diazene

[73399-75-4]

$C_{16}H_{30}B_{20}N_2$


Complex structure

See entry AZOCARBABORANES

3722. Bromo-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclo-tetra decadieneiron(II) perchlorate


[31122-43-7]

$C_{16}H_{32}BrClFeN_4O_4$





See entry [14] DIENE-N₄ COMPLEXES

3723. Iodo-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II) perchlorate
[31122-35-7]





See entry [14] DIENE-N₄ COMPLEXES

3724. Chloro-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II) perchlorate
[31122-42-6]




See entry [14] DIENE-N₄ COMPLEXES

3725. Dichloro-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(III) perchlorate
[36691-97-1]



See entry [14] DIENE-N₄ COMPLEXES


†3726. Hexadecane
[544-76-3]



$\text{C}_{16}\text{H}_{34}$

3727. Hexadecanethiol
[2917-26-2]

$\text{C}_{16}\text{H}_{34}\text{S}$



Nitric acid

See Nitric acid: Alkanethiols

3728. Tetrabutylammonium fluoride
[429-41-4]




$\text{C}_{16}\text{H}_{36}\text{FN}$

Hexamethyldisilazane, Pyridine *N*-oxide

See Pyridine *N*-oxide: Hexamethyldisilazane, etc.

3729. Tetrabutylammonium permanganate
[35638-41-6]



$\text{C}_{16}\text{H}_{36}\text{MnNO}_4$

1. Sala, T. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1978, 254

2. Morris, J. A. *et al.*, *Chem. & Ind.*, 1978, 446

Portions of a 20 g sample of the oxidant salt, prepared exactly as the published description [1], had been used uneventfully during a few days. Subsequently, as a 2 g portion was tipped out onto glazed paper, it ignited and burned violently, leading also to ignition of the bottled material [2].

See other QUATERNARY OXIDANTS

3730. Titanium butoxide
[5593-70-4]



$\text{C}_{16}\text{H}_{36}\text{O}_6\text{Ti}$

HCS 1980, 884


Ethyl terephthalate, Ethylene glycol

See Dimethyl terephthalate: Ethylene glycol, Titanium butoxide

See other METAL ALKOXIDES

3731. Tetrakis(butylthio)uranium

[]




Bailar, 1973, Vol. 5, 416

It ignites in air.

See related METAL ALKOXIDES, METAL SULFIDES**3732. Tetrabutylammonium hydrogen monoperoxysulfate**


[104548-30-3]

Trost, B. M. *et al.*, *J. Org. Chem.*, 1987, **53**, 532–537

Although no instability was noted, and a small sample was insensitive to a hammer blow, it should be treated as potentially unstable as an organic oxidant.

See other QUATERNARY OXIDANTS**3733. Tetrakis(diethylphosphino)silane**

[]




Air, or Water

Fritz, G. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1963, **2**, 262

It ignites in air and is extremely sensitive to water.

See related ALKYLPHOSPHINES, ALKYL SILANES**3734. 2,6-Bis(picrylazo)-3,5-dinitropyridine**

[55106-96-2]




Explosive.

See entry POLYNITROAZOPYRIDINES

3735. 2,6-Bis(2-picrylhydrazino)-3,5-dinitropyridine
[55106-95-1]

C₁₇H₉N₁₃O₁₆




Explosive.

See entry POLYNITROAZOPYRIDINES

3736. Benzanthrone
[82-05-3]

C₁₇H₁₀O




Nitrobenzene, Potassium hydroxide

See Nitrobenzene: Alkali (reference 4)

3737. S-7-Methylnonylthiuronium picrate
[]

C₁₇H₂₇N₅O₇S




Leese, C. L. *et al. J. Chem. Soc.*, 1950, 2739

The salt explodes if heated rapidly.

See other PICRATES

3738. Tris(pentafluorophenyl)boron
[1109-15-5]

C₁₈BF₁₅



See Triethylaluminium: Tris(pentafluorophenyl)boron
See also METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

3739. Iron(III) hexacyanoferrate(4-)
[14038-43-8]

C₁₈Fe₇N₁₈



Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455

The energy of decomposition of Prussian blue in air in the range 210–360°C was determined by DSC as 2.42 kJ/g, peaking at 271°C. When nitrogen was used as inerting gas in the sealed micro-crucible, decomposition was delayed to 250–350°C with a broad peak.

Blown castor oil, Turkey red oil (sulfonated castor oil)

Gärtner, K., *Farben-Ztg.*, 1938, **43**, 1118

A colour mixture containing the 3 components ignited spontaneously. Oxidation products in the air-blown oil may have reacted exothermally with the complex cyanide, a reductant.

Ethylene oxide

1. Wagle, N. G., *Chem. Brit.*, 1977, **13**, 317

2. Heuser, S. G. B., *Chem. Brit.*, 1977, **13**, 436

Interaction of 'iron blue' pigment with ethylene oxide vapour at ambient temperature is highly exothermic and gives a product which ignites in contact with air [1]. The hazards of using undiluted ethylene oxide in sterilisers, and the need to use inert purging between vacuum degassing stages is stressed [2].


Lead chromate

See Lead chromate: Iron(III) hexacyanoferrate(4-)

See other METAL CYANIDES (AND CYANO COMPLEXES)

**3740. 1,7,13-Cyclooctadecatriene-3,5,9,11,15,17-hexayne
(Hexahydro[18]annulene)
[16668-69-2]**

C₁₈H₆




Diederich, F. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(9), 1101

Explodes on heating to 85°C.

See other ALKYNES

3741. Lanthanum picrate




Tucholskii, T., *Rept. AD 633414*, Springfield (Va.), USNTIS, 1966

In a study of the thermal stability of the picrates of group III metals, that of lanthanum occasionally exploded prematurely on heating.

See other PICRATES

3742. Manganese picrate hydroxide
[41570-79-0]

C₁₈H₇Mn₂N₉O₂₂




Bernardi, A., *Gazz. Chim. Ital.*, 1930, **60**, 169–171

The basic manganese picrate double salt, and its 2,4-dinitro- and 2-nitro-analogues, all deflagrated violently on heating.

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, PICRATES, POLYNI-TROARYL COMPOUNDS

3743. 1,6-Bis(4-chlorophenyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane
[76695-68-6]

C₁₈H₈Cl₂N₄O₄




See 1,6-Diphenyl-2,5-bis(diazo)-1,3,4,6-tetraoxohexane (next below)

See other DIAZO COMPOUNDS

3744. 1,6-Diphenyl-2,5-bis(diazo)-1,3,4,6-tetraoxohexane
[76695-64-2]

$C_{18}H_{10}N_4O_4$



Rubin, M. B. *et al.*, *J. Chem. Soc., Perkin Trans. I*, 1980, 2671


The title compound and its 4,4'-chloro-, methyl- and methoxy-derivatives all decomposed violently at 155–165°C.

See other DIAZO COMPOUNDS

3745. Nickel 2,4-dinitrophenoxide hydroxide

[]

$C_{18}H_{10}N_6Ni_2O_{16}$




Bernardi, A., *Gazz. Chim. Ital.*, 1930, **60**, 166

It deflagrates very violently on heating.

See other HEAVY METAL DERIVATIVES, METAL NITROPHENOXIDES, POLYNITROARYL COMPOUNDS

3746. 4,4'-Diphenyl-2,2'-bi(1,3-dithiol)-2'-yl-2-ylum perchlorate
[53213-77-7]

C₁₈H₁₂ClO₄S₄




Sunderlin, K. G. R., *Chem. Eng. News*, 1974, **52**(31), 3

A few mg exploded violently during determination of the m.p., shattering the apparatus.

See other NON-METAL PERCHLORATES

3747. 9-Phenyl-9-iodafluorene
[32714-73-5]

C₁₈H₁₃I



Banks, D. F., *Chem. Rev.*, 1966, **66**, 248


It explodes at 105°C.

See other IODINE COMPOUNDS

3748. Oxodiperoxodiquinolinechromium(VI)

[]

C₁₈H₁₄CrN₂O₅




See Oxodiperoxodipyridinechromium(VI)

See other AMMINECHROMIUM PEROXOCOMPLEXES

3749. Triphenylaluminium
[841-76-9]

C₁₈H₁₅Al



Water


Neely, T. A. *et al.*, *Org. Synth.*, 1965, **45**, 107

Triphenyl aluminium and its etherate evolved heat and sparks on contact with water.

See other ARYLMETALS

3750. Triphenylsilyl perchlorate
[101652-99-7]

C₁₈H₁₅ClO₄Si



See entry ORGANOSILYL PERCHLORATES

3751. Triphenylchromium tetrahydrofuranate
[16462-53-6]

C₁₈H₁₅Cr.3C₄H₈O

Ph₃Cr.3C₄H₈O

Diethyl ether

Bailar, 1973, Vol. 4, 974

On warming, or on treatment with ether, the solvated complex give a black pyrophoric material of unknown constitution.

See other ARYLMETALS

3752. Triphenylphosphine oxide-oxodiperoxochromium(VI)
[93228-65-0]

C₁₈H₁₅CrO₆P


[Ph₃P(O)Cr(O₂)₂O]

Daire, E. *et al.*, *Nouv. J. Chim.*, 1984, **8**, 271–274

It effectively hydroxylates hydrocarbons but is free of the explosion risk of the analogous pyridine *N*-oxide complex.

See related AMMINECHROMIUM PEROXOCOMPLEXES

3753. Triphenyllead nitrate
[21483-09-8] (ion)




Sulfuric acid

Gilman, H. *et al.*, *J. Org. Chem.*, 1951, **16**, 466

It ignites in contact with conc. acid

See related ARYLMETALS

3754. 1,3,5-Triphenyl-1,4-pentaazadiene
[30616-12-7]




Sorbe, 1968, 141

It explodes at 80°C and is shock-sensitive.

See other HIGH-NITROGEN COMPOUNDS

3755. Triphenylphosphine oxide hydrogen peroxidate
[3319-45-7]



Bradley, D. C. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1974, 4


The 2:1 complex may explode.

See CRYSTALLINE HYDROGEN PEROXIDATES

See related REDOX COMPOUNDS

3756. Triphenylphosphine
[603-35-0]

C₁₈H₁₅P




Preparative hazard

See Sodium: Chlorobenzene, Phosphorus trichloride

See related ALKYLPHOSPHINES

3757. 1,3-Bis(phenyltriazeno)benzene
[70324-26-4]

C₁₈H₁₆N₆




Kleinfeller, H., *J. Prakt. Chem.* [2], 1928, **119**, 61

It explodes if heated rapidly.

See other TRIAZENES

3758. Triphenyltin hydroperoxide
[4150-34-9]

C₁₈H₁₆O₂Sn



Dannley, R. L. *et al.*, *J. Org. Chem.*, 1965, **30**, 3845


It explodes reproducibly at 75°C.

See other ORGANOMETALLIC PEROXIDES

3759. Sodium hexakis(propynyl)ferrate(4-)

[]

C₁₈H₁₈FeNa₄



Bailar, 1973, Vol. 3, 1025


An explosive complex salt.

See other COMPLEX ACETYLIDES

3760. 1-(1-Methyl-1-phenylethyl)-4-(2-propynyloxy)benzene

[]

C₁₈H₁₈O



Douglas, W. E. *et al.*, *J. Organomet. Chem.*, **444**(1-2), C62

Explosively polymerises when heated >110°C in presence of nickelocene catalyst.


See VIOLENT POLYMERISATION

See other ACETYLENIC COMPOUNDS, POLYMERISATION INCIDENTS

3761. 1,5-Cyclooctadiene-bis(4-chloropyridine N-oxide)rhodium(I) perchlorate

[80005-73-8]

C₁₈H₂₀Cl₃N₂O₆Rh




Uson, R. *et al.*, *J. Organomet. Chem.*, 1981, **217**, 252

The complex, one of a series of substituted pyridine derivatives, exploded violently during microanalytical combustion.

See related AMMINOMETAL OXOSALTS

3762. 1,2-Dihydroperoxy-1,2-bis(benzeneazo)cyclohexane
[71404-13-2]

C₁₈H₂₀N₄O₄




See Oxygen: Cyclohexane-1,2-dione bis(phenylhydrazone)

See other α -PHENYLAZO HYDROPEROXIDES

3763. O-O-tert-Butyl di(4-tolyl) monoperoxophosphate
[]

C₁₈H₂₃O₅P




Rieche, A. *et al.*, *Chem. Ber.*, 1962, **95**, 385

The material deflagrated as a solid, or decomposed exothermally in its reaction mixture soon after preparation.

See other tert-BUTYL PEROXOPHOSPHATE DERIVATIVES, PHOSPHORUS ESTERS

3764. 2-(1-Methylheptyl)-4,6-dinitrophenyl crotonate
[27661-44-5]

C₁₈H₂₄N₂O₆



Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 2

Decomposition and a pressure explosion occurred while a 10 kl tanker with steam-heating coils was being unloaded.


See related PEROXIDISABLE COMPOUNDS

See other POLYNITROARYL COMPOUNDS

3765. 2,2,4-Trimethyldecahydroquinolinium picrate

[]

C₁₈H₂₃N₄O₇





2-(2-Butoxyethoxy)ethanol

Franklin, N. C., private comm., 1967

Evaporation of a solution of the picrate in the diether (b.p. 230°C) caused a violent explosion. The solvent had probably peroxidised during open storage and the residual mixture of peroxide and picrate had exploded as evaporation proceeded. Use of peroxide-free solvent and lower (vacuum) evaporation temperature appeared to be safe.



See other PICRATES

**3766. Bis(benzyl 1-methylhydrazinocarbodithioate
*N*²,*S*¹)(perchlorato-*O*,*O*¹)copper(1+) perchlorate
[67870-99-9]**



See Bis-*O,N*[(*N*'-pent-2-en-2-oxy-4-ylidene)-*N,S*-dimethyldithiocarbazate]
copper(II) perchlorate

**3767. Dicumenechromium(0)
[12001-89-7]**





Aldrich catalogue, 1986

It is pyrophoric.




See other ARYLMETALS, ORGANOMETALLICS

3768. 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetra-
cyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-
hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²cobalt(II) perchlorate
[72644-04-3]





See entry CLATHROCHELATED METAL PERCHLORATES

3769. 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetra-
cyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-
hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²iron(II) perchlorate
[75516-37-9]




See entry CLATHROCHELATED METAL PERCHLORATES

3770. 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetra-
cyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-
hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²nickel(II) perchlorate
[63128-08-5]



See entry CLATHROCHELATED METAL PERCHLORATES

3771. 9,12,15-Octadecatrienoic acid (Linolenic acid)
[463-40-1]





Cobalt naphthenate

Kotoyori, T., *Chem. Abs.*, 1980, **93**, 221950

Model experiments on the mixture (simulating paint tailings) exposed to air showed that storage at low temperature (7°C) led to a larger build-up of peroxide than occurred at 35°C. This would account for spontaneous ignition of paint tailings (residues) stored at low temperature and subsequently heated to 60°C.

See AUTOXIDATION See other ORGANIC ACIDS

3772. Oleoyl chloride (cis-9-Octadecenoyl chloride)
[112-77-6]



'Acid tars'

Anon., *Jahresbericht*, 1994, 74

The residue of a sample of 'olein chloride' (crude oleoyl chloride) was tipped into a waste container containing 'acid tars' (70–80% waste sulfuric acid). There was a vigorous reaction, with gas evolution, which sprayed the analytical technician involved with the liquid contents of the container, causing burns.

Nitric acid


See Nitric acid: Oleoyl chloride

See other ACYL HALIDES

3773. 1,4-Octadecanolactone

[502-26-1]

C₁₈H₃₄O₂




Preparative hazard

See Perchloric acid: Oleic acid

3774. *cis*-9-Octadecenoic acid (Oleic acid)

[112-80-1]

C₁₈H₃₄O₂



HCS 1980, 701

Aluminium

See Aluminium: Oleic acid


Perchloric acid

See Perchloric acid: Oleic acid

See other ORGANIC ACIDS

3775. Bis(dibutylborino)acetylene

[]

C₁₈H₃₆B₂Hartmann, H. *et al.*, *Z. Anorg. Chem.*, 1959, **299**, 174

Both the n- and iso-derivatives ignite in air.

See other ACETYLENIC COMPOUNDS, ALKYLBORANES**3776. 1,2-Bis(di-tert-butylphosphino)ethane**

[107783-62-0]

C₁₈H₄₀P₂Benn, R. *et al.*, *Z. Naturforsch. B*, 1986, **41B**, 680–691

The ligand, like its lower homologues, is pyrophoric in air.

See related ALKYLPHOSPHINES**3777. 1,2-Bis(tripropylsilyl)hydrazine**

[13271-98-2]

C₁₈H₄₄N₂Si₂*See entry* SILYLHYDRAZINES *See related* ALKYL SILANES**3778. Perchloratotris(triethylphosphine)palladium(II) perchlorate**


[94288-47-8]

C₁₈H₄₅Cl₂O₈P₃PdBruce, D. W. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1984, 2252

It explodes on heating.

See related AMMINOMETAL OXOSALTS, METAL PERCHLORATES, PLATINUM COMPOUNDS**3779. Hexamethylenetetrammonium tetraperoxochromate(V)?**

[]


C₁₈H₄₈Cr₄N₁₂O₃₂House, D. A. *et al.*, *Inorg. Chem.*, 1966, **6**, 1078, footnote 8

Material (believed to be the title compound) recrystallised from water, and washed with methanol to dry by suction, ignited and then exploded on the filter funnel.

See related AMMINECHROMIUM PEROXOCOMPLEXES, AMMINOMETAL OXOSALTS

3780. 3,8-Dinitro-6-phenylphenanthridine
[82921-86-6]

C₁₉H₁₁N₃O₄




Diethyl sulfate

See Diethyl sulfate: 3,8-Dinitro-6-phenylphenanthridine

See other POLYNITROARYL COMPOUNDS

3781. Triphenylmethylpotassium
[1528-27-4]

C₁₉H₁₅K





Houben-Weyl, 1970, Vol. 13.1, 269

The dry powder ignites in air.

See other ORGANOMETALLICS

3782. Triphenylmethyl nitrate
[111422-41-4]

C₁₉H₁₅NO₃




Lewis acids

See entry ALKYL NITRATES

3783. Triphenylmethyl azide
[14309-25-2]

C₁₉H₁₅N₃




Robillard, J. J., Ger. Offen. 2 345 787, 1974

The unstable explosive azide may be stabilised by adsorption for reprographic purposes.

See other ORGANIC AZIDES

3784. Methyltriphenylphosphonium permanganate
[73335-41-8]

C₁₉H₁₈MnO₄P



1. Reischl, W. *et al.*, *Tetrahedron*, 1979, **35**, 1109–1110


2. Leddy, B. P. *et al.*, *Tetrahedron Lett.*, 1980, **21**, 2261–2262

It decomposes explosively above 70° [1], or at 80–90°C [2].

See other QUATERNARY OXIDANTS

3785. Hexadecyltrimethylammonium permanganate (Cetyltrimethylammonium permanganate)
[73257-07-5]

C₁₉H₄₂MnNO₄




Dash, S. *et al.*, *Chem. Abs.*, 1995, **123**, 227457m

Stable when kept in the dark at room temperature, it decomposes violently at 115°C.

See other QUATERNARY OXIDANTS

3786. Dibenzocyclododeca-1,7-dienetetrayne
(5,6,7,8,13,14,15,16-Octadehydrodibenzo[*a,g*]cyclododecene)
[7203-21-6]


$C_{20}H_8$



Guo, L. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1994, (3), 243
An unstable synthetic byproduct, attributed the above structure, explodes when scratched in air, leaving carbon as the residue. It blackens rapidly even in vacuum.
See other ACETYLENIC COMPOUNDS

3787. 2',4',5',7'-Tetrabromo-3',6'-dihydroxyspiro[isobenzofuran-1(3*H*),9'-(9*H*)]xanthen-3-one (Eosin)
[15876-39-8], [122530-01-2]


$C_{20}H_8Br_4O_5$



Lead salts, Red lead
See Dilead(II)lead(IV) oxide: Eosin

3788. 1,8-Diphenyloctatetrayne
[4572-12-7]

$C_{20}H_{10}$



1259


Nakagawa, M., *Chem. Abs.*, 1951, **45**, 7082a

Though stable in the dark at ambient temperature for a year, when exposed to light on a metal plate it decomposed explosively.

See other ALKYNES, IRRADIATION DECOMPOSITION INCIDENTS

3789. *I,I*-Bis(4-nitrobenzoylperoxy)-4-chlorophenyl iodine
[56391-49-2]


C₂₀H₁₂ClI₂N₂O₁₀



See entry *I,I*-DI(BENZOYLPEROXY)ARYLIODINES

3790. Perylenium perchlorate
[12576-63-5]

C₂₀H₁₂Cl₂O₈



Rosseinsky, D. R. *et al.*, *J. Chem. Soc., Perkin Trans. 2*, 1985, 135–138


The title compound, prepared by electrolysis of perylene and tetrabutylammonium perchlorate in nitrobenzene, exploded on contact with nickel. Co-produced compounds should also be handled with caution and in small amounts, especially in contact with metals.

See other NON-METAL PERCHLORATES

3791. Bis(1,8-naphthyridine)tetracarbonyldirhodium(I) perchlorate

[90635-49-7]

$C_{20}H_{12}Cl_2N_4O_{12}Rh_2$



Camellini, A. T. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1984, 130


Attempted microanalytical combustion of old samples of the salt or its 2-methyl homologue led to violent explosions.

See related CARBONYLMETALS, AMMINEMETAL OXOSALTS

3792. 11,12-Diethynyl-9,10-dihydro-9,10-ethenoanthracene

[126487-13-6]

$C_{20}H_{12}$



Diederich, F. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(9), 1101


The crystalline solid has been known to explode spontaneously.

See other ALKYNES

3793. 1,1-Bis(3-chlorobenzoylperoxy)-4-chlorophenyliodine

[]


$C_{20}H_{12}Cl_3IO_6$



See entry 1,1-DI(BENZOYLPEROXY)ARYLIODINES

3794. *I,I*-Bis(3-chlorobenzoylperoxy)phenyliodine
[30242-75-2]


$C_{20}H_{13}Cl_2IO_6$



See entry 1,1-DI(BENZOYLPEROXY)ARYLIODINES

3795. *I,I*-Bis(4-nitrobenzoylperoxy)phenyliodine
[30030-30-9]


$C_{20}H_{13}IN_2O_{10}$



See entry 1,1-DI(BENZOYLPEROXY)ARYLIODINES

3796. 1,6-Di(4'-tolyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane
[76695-67-5]


$C_{20}H_{14}N_4O_4$



See 1,6-Diphenyl-2,5-bis(diazo)-1,3,4,6-tetraoxohexane

3797. 1,6-Di(4'-methoxyphenyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane
[77695-66-4]


$C_{20}H_{14}N_4O_6$



See 1,6-Diphenyl-2,5-bis(diazo)-1,3,4,6-tetraoxohexane

3798. 2,5-Diphenyl-3,4-benzofuran-2,5-endoperoxide
(1,4-Epoxy-1,4-dihydro-1,4-diphenyl-2,3-benzodioxin)
[41337-62-6]

$C_{20}H_{14}O_3$



Alone, or Carbon disulfide

1. Dufraisse, C. *et al.*, *Compt. rend.*, 1946, **223**, 735


2. Rio, G. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 72

Formally an ozonide, this photo-peroxide explodes at 18°C [1], and its dimerisation, catalysed by carbon disulfide, was exothermic and sometimes explosive at ambient temperature [2].

See other CYCLIC PEROXIDES, OZONIDES

3799. *cis*-Dichlorobis(2,2'-bipyridyl)cobalt(III) chloride
[23380-38-3 or 14522-39-5]

$C_{20}H_{16}Cl_3CoN_4$




Preparative hazard

See Chlorine: Cobalt(II) chloride, Methanol

3800. 1,1-Bis(4-nitrobenzoylperoxy)cyclohexane

[]

$C_{20}H_{18}N_2O_{10}$



Criegee, R. *et al.*, *Ann.*, 1948, **560**, 135

It explodes at 120°C.

See other PEROXYESTERS

3801. Tetrapyrindinecobalt(II) chloride

[13985-87-0]

$C_{20}H_{20}Cl_2CoN_4$

$[(C_5H_5N)_4Co]Cl_2$


Chlorine, Methanol

See Chlorine: Methanol, Tetrapyrindinecobalt(II) chloride

3802. *trans*-Dichlorotetrapyrindinecobalt(III) chloride

[144077-02-2 or 27883-34-7]

$C_{20}H_{20}Cl_3CoN_4$




Preparative hazard

See Chlorine: Methanol, Tetrapyrindinecobalt(II) chloride

3803. Oxybis[bis(cyclopentadienyl)titanium]
[51269-39-7]


$C_{20}H_{20}OTi_2$



Bottomley, F. *et al.*, *J. Amer. Chem. Soc.*, 1980, **102**, 5241
The dry solid is explosively oxidised in contact with air.
See other ORGANOMETALLICS

3804. 1,1-Bis(benzoylperoxy)cyclohexane
[13213-29-1]


$C_{20}H_{20}O_6$



Criegee, R. *et al.*, *Ann.*, 1948, **560**, 135
It explodes sharply in a flame.
See other PEROXYESTERS

3805. Sodium abietate
[14351-66-7]


$C_{20}H_{29}NaO_2$



See entry METAL ABIETATES

3806. Dichlorodi- μ -chlorobis(pentamethylcyclopentadienyl)dirhodium

[12354-85-7]

C₂₀H₃₀Cl₄Rh₂


Air, Alkylmetals

Isobe, K. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1983, 1445

In preparation of di- μ -methylenebis(methyl-pentamethylcyclopentadienyl)dirhodium complexes by aerobic oxidation of a solution of the halocomplex and methyl-lithium or trimethylaluminium in ether-benzene, the reaction mixture occasionally ignited and burned violently. Full precautions and a working scale below 1 mmol are recommended.

See related ALKYL METAL HALIDES**3807. Di-3-camphoroyl peroxide**


[]

C₂₀H₃₀O₈Milas, N. A. *et al.*, *J. Amer. Chem. Soc.*, 1933, **55**, 350-351

Explosive decomposition occurred at the m.p., 142°C, or on exposure to flame.

See other DIACYL PEROXIDES

3808. *trans*-Dichlorobis[1,2-phenylenebis(dimethylarsine)]palladium(IV) diperchlorate
[60489-76-1]





Gray, L. R. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1983, 33–41

Though no explosions occurred during the preparation by addition of 75% perchloric acid to solutions of the complex in conc. nitric/hydrochloric acid mixtures, this and the amino- and phosphino-analogues are potentially unstable.

See related AMMINOMETAL OXOSALTS

3809. Sodium 5,8,11,14-eicosatetraenoate (Sodium arachidonate)
[6610-25-9]





Henderson, J. D. *et al.*, *Chem. Eng. News*, 1983, **61**(33), 2

An opened 100 mg ampoule of the salt was stored dark in a desiccator at -20°C between occasional samplings during a few months. A few minutes after return to storage, the ampoule contents decomposed violently. It was supposed that peroxide formation (and perhaps polymerisation of the polyene) was involved in the incident.

See other PEROXIDISABLE COMPOUNDS

3810. 1,1,6,6-Tetrakis(acetylperoxy)cyclododecane
[]



Criegee, R. *et al.*, *Ann.*, 1948, **560**, 135

Weak friction causes strong explosion.
See other PEROXYESTERS

3811. Tributyl(phenylethynyl)lead
[21249-40-0]




$\text{C}_{20}\text{H}_{32}\text{Pb}$

Houben-Weyl., 1975, Vol. 13.3, 80
Violent decomposition occurred during attempted distillation.
See related ALKYLMETALS, METAL ACETYLIDES

**3812. B-Iododiisopinocampheylborane (Iodobis(2,6,6-trimethylbicyclo
[3.1.1]hept-3-yl)borane)**
[116005-09-5]


$\text{C}_{20}\text{H}_{34}\text{BI}$



Brown, H. C. *et al.*, *Heteroat. Chem.*, 1995, **6**(2), 117
Pyrophoric, very hygroscopic, fumes in air.
See other ALKYLHALOBORANES

**3813. Diacetonitrile-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-
4,11-cyclotetradecadieneiron(II) perchlorate**
[50327-56-5]


$\text{C}_{20}\text{H}_{38}\text{Cl}_2\text{FeN}_6\text{O}_8$



See entry [14] DIENE-N₄ COMPLEXES

3814. Tetracyanooctaethyltetragold

[]


C₂₀H₄₀Au₄N₄Burawoy, A. *et al.*, *J. Chem. Soc.*, 1935, 1026

This tetramer of cyanodiethylgold is friction-sensitive and also decomposes explosively above 80°C. The propyl homologue is not friction-sensitive, but also decomposes explosively on heating in bulk.

See other CYANO COMPOUNDS, GOLD COMPOUNDS

3815. Lanthanum 2-nitrobenzoate

[114515-72-9]


C₂₁H₁₂LaN₃O₁₂Ferenc, W. *et al.*, *Monatsh. Chem.*, 1987, **118**, 1087–1100

Preparation of the 2-nitrobenzoate salts of yttrium and the lanthanide metals (except praseodymium) as mono- or di-hydrates was studied. All melted and decomposed explosively above 250°C.

See other HEAVY METAL DERIVATIVES, NITROARYL COMPOUNDS

3816. Scandium 3-nitrobenzoate

[111016-97-8]

C₂₁H₁₂N₃O₁₂Sc

Brzyska, W. *et al.*, *J. Therm. Anal.*, 1987, **32**, 671–678


After loss of the hemi-hydrate water, the anhydrous complex decomposes explosively on further heating.

See other HEAVY METAL DERIVATIVES, NITROARYL COMPOUNDS

3817. Yttrium 4-nitrobenzoate trihydrate

[115627-20-8]

C₂₁H₁₂N₃O₁₂Y.3H₂O



Ferenc, W. *et al.*, *Monatsh. Chem.*, 1988, **119**, 407–424


The yttrium, lanthanum and other lanthanide salts exploded after dehydration during heating to above 300°C.

See other HEAVY METAL DERIVATIVES, NITROARYL COMPOUNDS

3818. 1,3,5-Tris(4-azidosulfonylphenyl)-1,3,5-triazinetrione

[31328-33-3]

C₂₁H₁₂N₁₂O₉S₃



Ulrich, H. *et al.*, *J. Org. Chem.*, 1975, **40**, 804


This trimer of 4-azidosulfonylphenylisocyanate melts at 200°C with violent decomposition.

See other ACYL AZIDES

3819. 1,1-Bis(4-nitrobenzoylperoxy)-2-methoxyphenyliodine

[]

C₂₁H₁₅IN₂O₁₁




See entry 1,1-DI(BENZOYLPEROXY)ARYLIODINES

3820. N,N-Diphenyl-3-phenylpropenyldenimmonium perchlorate

[77664-36-9]

C₂₁H₁₈ClNO₄



Qureshi, M. *et al.*, *J. Indian Chem. Soc.*, 1980, **57**, 915–917


The immonium salt explodes on heating.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

3821. 1-(2-Azido-2-phenylvinyl)phenyliodonium 4-toluenesulfonate

[118235-47-5]

C₂₁H₁₈IN₃O₃S




Kitamura, T. *et al.*, *Tetrahedron Lett.*, 1988, **29**, 1887–1890

It decomposes explosively on melting.

See other IODINE COMPOUNDS, ORGANIC AZIDES

3822. Tribenzylarsine
[5888-61-9]

C₂₁H₂₁As




Dondorov, J. *et al.*, *Ber.*, 1935, **68**, 1255

The pure material oxidises slowly at first in air at ambient temperature, but reaction becomes violent through autocatalysis.

See other ALKYLNON-METALS, BENZYL COMPOUNDS

3823. Tri-4-methylphenylsilyl perchlorate
[]


C₂₁H₂₁ClO₄Si



See entry ORGANOSILYL PERCHLORATES

3824. Tri(4-tolyl)ammonium perchlorate
[34729-49-2]

C₂₁H₂₂ClNO₄




Weitz, E. *et al.*, *Ber.*, 1926, **59**, 2307

It explodes violently when heated above the m.p., 123°C.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

3825. 2,2-Bis[4(2',3'-epoxypropoxy)phenyl]propane
[1675-54-3]

C₂₁H₂₄O₄




Trichloroethylene

See Trichloroethylene: Epoxides

See other 1,2-EPOXIDES

**3826. (η^8 1,3,5,7-Cyclooctatetraene)[2-(dimethylaminomethyl)phenyl-
C,N](tetrahydrofuran)lutetium**
[84582-81-0]

C₂₁H₂₈LuNO




Wayda, A. L., *Inorg. Synth.*, 1990, **27**, 150

The dry powder may ignite spontaneously in air, partially oxidised samples can explode.

See related ORGANOMETALLICS

3827. N-Hexadecylpyridinium permanganate
[76710-77-5]

C₂₁H₃₈MnNO₄




It explodes at 80–90°C.

See entry QUATERNARY OXIDANTS

3828. Acetonitrileimidazole-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II) perchlorate
[33691-99-3]


C₂₁H₃₉Cl₂FeN₇O₈



See entry [14] DIENE-N₄ COMPLEXES

3829. [1,1'-Oxybis[ethane]]tris(pentafluorophenyl)aluminium
(Tris(pentafluorophenyl)aluminium etherate)
[14524-44-8]


C₂₂H₁₀AlF₁₅O



Pohlmann, J. L. W. *et al.*, *Z. Naturforsch.*, 1965, **20b**(1), 5
This compound sometimes explodes on heating.
See other METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

3830. Bis(cyclopentadienyl)bis(pentafluorophenyl)zirconium
[12097-97-1]


C₂₂H₁₀F₁₀Zr



Chaudhari, M. A. *et al.*, *J. Chem. Soc. (A)*, 1966, 838
It explodes in air (but not nitrogen) above the m.p., 219°C.
See other HALO-ARYLMETALS, ORGANOMETALLICS

3831. Di-1-naphthoyl peroxide
[29903-04-6]


C₂₂H₁₄O₄



Castrantas, 1965, 17
It explodes on friction.
See other DIACYL PEROXIDES

3832. Tetraspiro[2.2.2.2.2.2.4]docosa-4,9,14,19,21pentyne
[159148-53-5]

$C_{22}H_{16}$




Scott, L. T. *et al.*, *J. Amer. Chem. Soc.*, 1994, **116**(22), 10275

This compound melted at 155°C with decomposition, one sample exploded in an oil bath at 192°C.

See also ALKYNES, See also ROTANES

3833. Bis(2-methyl-1,8-naphthyridine)tetracarbonyldirhodium(I) perchlorate
[90635-53-3]


$C_{22}H_{16}Cl_2N_4O_{12}Rh_2$



See Bis(1,8-naphthyridine)tetracarbonyldirhodium(I) perchlorate


See related CARBONYLMETALS, AMMINOMETAL OXOSALTS

3834. Bis(benzeneiron)–fulvalenediyl complex
[78398-48-8]





Desbois, M. H. *et al.*, *J. Amer. Chem. Soc.*, 1985, **107**, 5280–5282
Unstable above $-10^{\circ}C$, it burns explosively on contact with air.
See other ORGANOMETALLICS

3835. Bis(dicarbonylcyclopentadienyliron)–bis(tetrahydrofuran)magnesium
[55800-08-3]



McVicker, G. B., *Inorg. Synth.*, 1976, **16**, 56–58
This, like other transition metal carbonyl derivatives of magnesium, is pyrophoric.
See other CARBONYLMETALS, ORGANOMETALLICS

**3836. 4-[2-(4-Hydrazino-1-phthalazinyl)hydrazino]-4-methyl-2-pentanone
(4-hydrazino-1-phthalazinyl)hydrazonickel(II) tetra perchlorate**
[9018-45-5]




Rosen, W., *Inorg. Chem.*, 1971, **10**, 1833

The green polymeric complex isolated directly from the reaction mixture explodes fairly violently at elevated temperatures. The tetrahydrate produced by recrystallisation from aqueous methanol is also moderately explosive when heated rapidly.

See other AMMINOMETAL OXOSALTS

See related TETRAAZAMACROCYCLANEMETAL PERCHLORATES

3837. Bis(trimethylphosphine)di(3,5-dibromo-2,6-dimethoxyphenyl)nickel
[83459-96-5]




Wada, S. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 1446

The complex decomposed explosively at 194°C.

See related ALKYLPHOSPHINES, HALO-ARYLMETALS

3838. 4,4'-(Butadiyne-1,4-diyl)bis(2,2,6,6-tetramethyl-4-hydroxypiperidine-1-oxyl)
[14306-88-8] **C₂₂H₃₄N₂O₄**




Miller, J. S. *et al.*, *Chem. Mater.*, 1990, **2**(1), 60

Both crystalline forms decompose explosively to lower molecular weight products at 140°C under nitrogen.

See related ALKYNES

3839. 2,4,6-Triphenylpyrilium perchlorate
[1484-88-4]

C₂₃H₁₇ClO₅




Pelter, A., *Chem. Abs.*, 1981, **95**, 97643 (reference to *Tetrahedron Lett.* original is incorrect)

Though not as sensitive as 1,3-benzodithiolium perchlorate, this salt is also explosive.

See other NON-METAL PERCHLORATES

3840. 1,3-Di[bis(cyclopentadienyl)iron]-2-propen-1-one
[12171-97-0]

C₂₃H₂₀Fe₂O




Perchloric acid, Acetic anhydride, Cyclopentanone, Ether, Methanol
Anon., *Chem. Eng. News*, 1966, **44**(49), 50

Condensation of the iron complex with cyclopentanone in perchloric acid–acetic anhydride–ether medium had been attempted. The non-crystalline residue, after methanol washing and drying in air for several weeks, exploded on being disturbed. This was attributed to possible presence of a derivative of ferrocenium perchlorate, a powerful explosive and detonator. However, methyl or ethyl perchlorates alternatively may have been involved.

See Perchloric acid: Acetic anhydride, etc., or: Diethyl ether

See other ORGANOMETALLICS

3841. 3,3,5-Triphenyl-4,4-dimethyl-5-hydroperoxy-4,5-dihydro(3H)pyrazole
(4,5-Dihydro-4,4-dimethyl-3,5,5-triphenyl-3H-pyrazol-3-yl hydroperoxide)
[133610-01-2] **C₂₃H₂₂N₂O₂**




Baumstark, A. L., *et al.*, *J. Heterocyclic Chem.*, 1991, **28**(1), 113

This, the most stable of 5 homologues, the others having one or two aryl substituents replaced by methyl, exploded at 90°C in a melting point determination. It appears that azohydroperoxy compounds are even more explosive than the sum of their parts might indicate.

See related α -PHENYLAZO HYDROPEROXIDES, ORGANIC PEROXIDES


3842. Tetrakis(pentafluorophenyl)titanium
[37759-21-0] **C₂₄F₂₀Ti**



Houben-Weyl, 1975, Vol. 13.3, 300
It explodes at 120–130°C.
See other HALO-ARYLMETALS

3843. 1,7,13,19-Cyclotetracosatetraene-3,5,9,11,15,17,21,23-octayne
(Octahydro[24]annulene)
[30047-26-8]


C₂₄H₈



Diederich, F. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(9), 1101
Explodes on heating.
See other ALKYNES

3844. Iron(III) phthalate
[52118-12-4]


C₂₄H₁₂Fe₂O₁₂



Sulfur compounds
See Phthalic anhydride (reference 2)

3845. 1,5-Dibenzoylnaphthalene
[83-80-7]

C₂₄H₁₆O₂




Preparative hazard

See Aluminium chloride: Benzoyl chloride, etc.

3846. Aquafluorobis(1,10-phenanthroline)chromium(III) perchlorate
[77812-44-3]

C₂₄H₁₈Cl₂CrFN₄O₁₀




Delover, M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1981, 985

Its explosive properties prevented analysis.

See *other* AMMINEMETAL OXOSALTS

3847. 1-(2'-Diazoniophenyl)2-methyl-4,6-diphenylpyridinium diperchlorate
[55358-25-3]

C₂₄H₁₉Cl₂N₃O₈




Dorofenko, G. N. *et al.*, *Chem. Abs.*, 1975, **82**, 139908

The title compound, its 3'- and 4'-isomers, and their 2,4,6-triphenyl analogues, exploded on heating.

See other DIAZONIUM PERCHLORATES

3848. Tetraphenylarsonium permanganate
[4312-28-1]

C₂₄H₂₀AsMnO₄




It explodes at 120–130°C.

See entry QUATERNARY OXIDANTS

3849. 1,3,6,8-Tetraphenyloctaazatriene
[]

C₂₄H₂₀N₈




Wohl, A. *et al.*, *Ber.*, 1900, **33**, 2741

This compound (and several C-homologues) is unstable and explodes sharply on heating, impact or friction. In a sealed tube, the explosion is violent.

See other HIGH-NITROGEN COMPOUNDS

3850. Tetraphenyllead
[595-89-1]

C₂₄H₂₀Pb



Potassium amide

See Potassium amide: Tetraphenyllead

Sulfur



Houben-Weyl, 1975, Vol. 13.3, 236

Interaction may be explosive.

See other ARYLMETALS

3851. Bis(benzene)chromium dichromate

[]





Anon., *Chem. Eng. News*, 1964, **42**(38), 55

This catalyst exists as explosive orange-red crystals.

See other ORGANOMETALLICS See related METAL OXOMETALLATES

3852. 3,3,6,6-Tetraphenylhexahydro-3,6-disilatetrazine


[13272-05-4]



See entry SILYLHYDRAZINES

3853. Bis(*O*-salicylidinaminopropylaziridine)iron(III) perchlorate

[79151-63-6]



Federer, W. D. *et al.*, *Inorg. Chem.*, 1984, **23**, 3863


Heat- and mildly shock-sensitive, a sample of the complex exploded after hand-grinding in a mortar for 30 min.

See other AZIRIDINES, AMMINOMETAL OXOSALTS

3854. 2-Tetrahydrofuranylidene(dimethylphenylphosphine-trimethylphosphine)-2,4,6-trimethylphenylnickel perchlorate

[]

$C_{24}H_{37}ClNiO_5P_2$



Wada, M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 794

It explodes on heating, like a tetrahydropyranilydene analogue.


See other ORGANOMETALLIC PERCHLORATES

See related ALKYLPHOSPHINES

3855. Di- μ -methylenebis(methylpentamethylcyclopentadienyl)dirhodium

[80410-45-3]

$C_{24}H_{40}Rh_2$




Preparative hazard

See Dichlorodi- μ -chlorobis(pentamethylcyclopentadienyl)dirhodium: Air, Alkylmetals

See related ALKYLMETALS

3856. Tris(bis-2-methoxyethyl ether)potassium hexacarbonylniobate(1-)
[57304-94-6] **C₂₄H₄₂KNbO₁₅**




Solvents

Attempts to recover the complex from solutions in solvents other than diglyme led to the formation of pyrophoric solids.

See related CARBONYLMETALS

3857. Didodecanoyl peroxide (Dilauroyl peroxide)
[105-74-8] **C₂₄H₄₆O₄**



1. *Haz. Chem. Data*, 1975, 137
2. *CHETAH*, 1990, 183

Though regarded as one of the more stable peroxides, it becomes shock-sensitive on heating, and self-accelerating decomposition sets in at 49°C [1]. One of the recently calculated values of 46 and 42°C for induction periods of 7 and 60 days, respectively, for critical ignition temperatures is closely similar to that (45°/7 days) previously recorded. Autocatalytic combustion of the polymerisation initiator is exhibited. Although not ordinarily shock sensitive, it responds to a detonator [2].

Charcoal


Leleu, *Cahiers*, 1980, (99), 279

Mixtures react exothermally at 90°C, sometimes igniting. (Catalysis by trace metals in the carbon may well be involved).

See other DIACYL PEROXIDES

3858. Tetracyclo[20.2.0.0^{6,9}.0^{14,17}]tetracos-1(22),6(9),14(17)-triene-2,4,10,12,18,20-hexayne-7,8,15,16,23,24-hexone
[123002-86-8]


C₂₄O₆



See Oligo(octacarbondioxide)
See other ACETYLENIC COMPOUNDS

3859. 4,4-Diferrocenylpentanoyl chloride
[56386-22-2]


C₂₅H₇ClFe₂O



Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1976, **41**, 657
The acid chloride is thermally unstable much above 25°C. At 80–100°C it rapidly evolves hydrogen chloride and forms a black solid.
See other ACYL HALIDES, ORGANOMETALLICS

3860. 1,5-(or 1,8-)Bis(dinitrophenoxy)-4,8-(or 4,5-)dinitroanthraquinone


[]

C₂₆H₁₀N₆O₁₆

Fast flame propagation occurs on heating the powders moderately.
See entry HIGH RATE DECOMPOSITION

3861. Copper bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide)

[12213-13-7]


C₂₆H₂₂CuN₈S₂

Sorbe, 1968, 80

The copper derivative of dithizone explodes at 150°C.



See other N-METAL DERIVATIVES, METAL HYDRAZIDES**3862. Lead bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide)**

[21519-20-0]

C₂₆H₂₂N₈PbS₂


Sorbe, 1968, 36
 The lead derivative of dithizone explodes at 215°C.
See other N-METAL DERIVATIVES, METAL HYDRAZIDES

3863. Zinc bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide)
 [21790-54-5]




Sorbe, 1968, 159
 The zinc derivative of dithizone explodes at 215°C.
See other N-METAL DERIVATIVES, METAL HYDRAZIDES

3864. 1,2-Bis(diphenylphosphino)ethanepalladium(II) perchlorate
 [69720-86-1]



Davies, J. A. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 2246–2249
 Extremely sensitive to thermal and mechanical shock, occasionally detonating spontaneously without external stimulus, it should only be prepared and used in solution. The solid is an extreme hazard, and the acetone-solvated complex is also explosive.
See related AMMINEMETAL OXOSALTS

3865. Didodecyl peroxydicarbonate
 [24356-04-5]



D'Angelo, A. J., US Pat. 3 821 273, 1974


It decomposes violently after 90 min at 40°C, while the hexadecyl homologue is stable for a week at 40°C.

See other PEROXYCARBONATE ESTERS

3866. Methylaluminiumbis(pentamethylcyclopentadienyltrimethyliridium)

[]

$C_{27}H_{51}AlIr_2$



Isobe, K. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1984, 932

When the complex in pentane at ambient temperature was deliberately exposed to air to produce pentamethylcyclopentadienyltetramethyliridium, it occasionally ignited. Using proper precautions it is not dangerous, but increase in scale above 0.4 mmol is not advised.

See related ALKYL METALS See other ORGANOMETALLICS

3867. Tetrakis(pyridine)bis(tetracarbonylcobalt)magnesium

[51006-26-9]

$C_{28}H_{20}Co_2MgN_4O_8$



McVicker, G. B., *Inorg. Synth.*, 1976, **16**, 58–59


This, like other transition metal carbonyl derivatives of magnesium, is pyrophoric.

See related CARBONYL METALS

3868. Bis[1,5-bis(4-methylphenyl)-1,3-pentaazadienato-N3,N5]-(T-4) cobalt

[113634-30-3]

$C_{28}H_{28}CoN_{10}$




Schmid, R. *et al.*, *Z. Naturforsch. B*, 1987, **42B**, 911–916

The red complex is explosive.

See other HEAVY METAL DERIVATIVES, HIGH-NITROGEN COMPOUNDS, *N*-METAL DERIVATIVES, PENTAAZADIENES

3869. 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadiene-1,10-phenanthrolineiron(II) perchlorate
[36732-77-1]


$C_{28}H_{40}Cl_2FeN_6O_8$



See entry [14] DIENE-N₄ COMPLEXES

3870. 1(2'-, 3'-, or 4'-Diazoniophenyl)-2,4,6-triphenylpyridinium diperchlorate
[55358-23-1, 55358-29-7, 55358-33-3, resp.]

$C_{29}H_{21}Cl_2N_3O_8$



Dorofenko, G. N. *et al.*, *Chem. Abs.*, 1975, **82**, 139908


The three diazonium perchlorate isomers, and their 2-methyl-4,6-diphenyl analogues, all exploded on heating.

See other DIAZONIUM PERCHLORATES

3871. Sodium tris(*O,O'*-1-oximatonaphthalene-1,2-dione)ferrate (Pigment green B)

[16143-80-9]

$C_{30}H_{18}FeN_3NaO_6$



Piotrowski, T., *Chem. Abs.*, 1983, **99**, 159938


The pigment (containing below 1% of inorganic salts) may ignite or explode during milling or drying. The causes of ignition and impact sensitivity were determined and safe handling procedures are proposed.

See related OXIMES See other N-O COMPOUNDS

3872. Cobalt(II) chelate of bi(1-hydroxy-3,5-diphenylpyrazol-4-yl *N*-oxide)

[75027-71-3]

$C_{30}H_{20}CoN_4O_4$



Battei, R. S. *et al.*, *J. Inorg. Nucl. Chem.*, 1980, **42**, 494




The title cobalt chelate exploded at 175°C during DTA studies, 30° lower than the Co chelate of 'dicupferron'.

See Cobalt(II) chelate of 1,3-bis(*N*-nitrosohydroxylamino)benzene

See other N-O COMPOUNDS, *N*-OXIDES

3873. Tris(2,2'-bipyridine)silver(II) perchlorate

[]







Morgan, G. T. *et al.*, *J. Chem. Soc.*, 1930, 2594

It explodes on heating.

See other AMMINEMETAL OXOSALTS

3874. Tris(2,2'-bipyridine)chromium(II) perchlorate

[15388-46-2]




Holah, D. G. *et al.*, *Inorg. Synth.*, 1967, **10**, 34

It explodes violently on slow heating to 250°C and can be initiated by static sparks, but not apparently by impact.

See other AMMINEMETAL OXOSALTS, STATIC INITIATION INCIDENTS

3875. Tris(2,2'-bipyridine)chromium(0)
[14751-89-4]

C₃₀H₂₄CrN₆




Herzog, S. *et al.*, *Z. Naturforsch. B*, 1957, **12B**, 809

It ignites in air.

See other PYROPHORIC MATERIALS

3876. Benzenediazonium tetraphenylborate
[2200-13-7]

C₃₀H₂₅BN₂




Sakakura, T. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1994, (3), 283

On two occasions this salt exploded during drying. The dry material is dangerous to handle. In view of the high molecular weight, the diazonium function, alone, represents a low energy per unit mass for explosivity

See other DIAZONIUM SALTS

**3877. 2-Tetrahydropyranylidene-bis(dimethylphenylphosphine)-
3,4,6-trimethylphenyl-nickel perchlorate**
[82647-51-6]

C₃₀H₄₁ClNiO₅P₂




Wada, M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 794

It explodes on heating, like a tetrahydrofurylidene analogue.

See other ORGANOMETALLIC PERCHLORATES



See related ALKYLPHOSPHINES

3878. 2,12,18,28-Tetramethyl-3,7,11,19,23,27,33,34-octaazatricyclo
 [27.3.1.1(13,17)]tetratriaconta-1(33),2,11,13,15,17(34)18,27,29,31-
 decaenetetraaquadiron(II) tetraperchlorate
 [94643-57-9]



Cabral, M. F. *et al.*, *Inorg. Chim. Acta*, 1984, **90**, 170
 The dinuclear complex exploded readily on slight friction.
 See other POLYAZACAGED METAL PERCHLORATES

3879. Bis(μ_3 -methylidyne) triangulo tris(pentamethylcyclopentadienylrhodium)
 [83350-11-2]




Vasquez de Miguel, A. *et al.*, *Organometallics*, 1982, **1**, 1606

Preparation of hexamethyldialuminium in benzene, necessary for reaction with $(C_5H_5Rh)_2Cl_4$ to give the title compound, is hazardous and should only be attempted by experienced workers.

See other ORGANOMETALLICS

3880. 2,7,12,18,23,28-Hexamethyl-3,7,11,19,23,27,33,34-octaazatricyclo [27.3.1.1(13,17)]tetratriaconta-1(33),2,11,13,15,17(34),18,27,29,31-decaenediaquadichlorodiiron(II) diperchlorate
[94643-57-9]





Cabral, M. F. *et al.*, *Inorg. Chim. Acta*, 1984, **90**, 170

The dinuclear complex exploded readily on slight friction.


See other POLYAZACAGED METAL PERCHLORATES

3881. Hydroxobis[5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II)] triperchlorate
[36913-76-5]




See other [14] DIENE-N₄ COMPLEXES

3882. Oxybis[aqua-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II)] tetraperchlorate
[36841-23-3]



See other [14] DIENE-N₄ COMPLEXES

3883. Tetrakis(μ₃-2-amino-2-methylpropanolato)tetrakis(μ₂-2-amino-2-methylpropanolato)hexacopper(II) perchlorate
[100113-85-7]




Complex structure

Muhonen, H. *et al.*, *Inorg. Chem.*, 1986, **25**, 800

The complex, containing a bicapped cubane Cu₆O₈ cluster, may explode if heated when dry.


See other AMMINOMETAL OXOSALTS

3884. Octacarbondioxide tetramer
[123002-87-9]



Oligo(octacarbondioxide) See other ACETYLENIC COMPOUNDS

3885. Triferrocenylcyclopropenium perchlorate
[56581-58-9]





Agranat, I. *et al.*, *Tetrahedron*, 1979, **3** 5, 738

The salt is thermally stable below 171°C, but may explode on friction at ambient temperature.

See other FRICTIONAL INITIATION INCIDENTS, NON-METAL PERCHLORATES

3886. Tris(4-methoxy-2,2''bipyridine)ruthenium(II) perchlorate
[]



Thomas, R. N., *Univ. Safety Assoc. Safety News*, 1981, **15**, 16–17


Failure of the product (0.5 g) to crystallise out from the aqueous DMF reaction liquor led to vacuum evaporation of the solution at 60–70°C. During evaporation the mixture exploded violently, shattering the fume cupboard sash of toughened glass. The product may well be thermally unstable, but reaction of DMF with excess warm perchloric acid, possibly in near-absence of water, may also have been involved.

See other AMMINOMETAL OXOSALTS

3887. Pentaspiro[2.4.2.4.2.4.2.4.2.4]pentatriaconta-4,6,11,13,18,20,25,27,32,34-
decayne

[164077-35-4]

C₃₅H₂₀



de Meijere, A. *et al.*, *Chem. Eur. J.* 1995, **1**(2), 124


Like others of the series, it is explosive. A sample destroyed the apparatus during melting point determination.

See other ROTANES

3888. 1,2:5,6:11,12:15,16-Tetrabenzocycloocta-1,5,11,15-tetraene-3,7,9,13,17,19-hexayne (1,2:5,6:11,12:15,16-Tetrabenzo-3,7,9,13,17,19-hexahydro[20]annulene)

[]

$C_{36}H_{16}$






Matzger, A. J. *et al.*, *J. Amer. Chem. Soc.*, 1997, **119**, 2052

Explodes at 245°C to give various forms of carbon, hydrogen and methane and 1 kJ/g of heat.

See other ACETYLENIC COMPOUNDS

3889. Tris(1,10-phenanthroline)ruthenium(II) perchlorate [27778-27-4]

$C_{36}H_{24}Cl_2N_6O_8Ru$








Sulfuric acid

Gillard, R. D. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1974, 1235

Dissolution of the salt in the conc. acid must be very slow with ice-cooling to prevent an explosive reaction.

See other AMMINEMETAL OXOSALTS

3890. Tris(1,10-phenanthroline)cobalt(III) perchlorate
[14516-66-6]




1. Hunt, H. R., *J. Chem. Educ.*, 1977, **54**, 710
2. Kildall, L. K., *J. Chem. Educ.*, 1978, **55**, 476
3. Hunt, H. R., *J. Chem. Educ.*, 1978, **55**, 476

The use of the perchlorate anion to precipitate the cobalt complex to determine the yield [1] is deprecated on the grounds of potential hazard [2], though it was not found possible to cause the salt to detonate by pounding, but it will burn if ignited.

A spectroscopic assay method is suggested as an alternative to precipitation [3].

See other AMMINEMETAL OXOSALTS

3891. Hexaphenylhexaarsane
[20738-31-2]



Oxygen



1. Maschmann, E., *Ber.*, 1926, **59**, 1143
2. Blicke, F. F. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 2946–2950

The explosively violent reaction of trimeric 'arsenobenzene' with oxygen [1] was later ascribed to the presence of various catalytic impurities, because the pure compound is stable towards oxygen [2].

See related ALKYLNON-METALS

3892. Lithium hexaphenyltungstate(2-)

[]





Sarry, B. *et al.*, *Z. Anorg. Chem.*, 1964, **329**, 218

The trietherate is pyrophoric in air.

See related ARYLMETALS

**3893. Di[*N,N'*-Ethylenebis(2-oxidoacetophenoneiminato)copper(II)]
oxovanadium(IV) diperchlorate**


[93895-68-2]



Bencini, A. *et al.*, *Inorg. Chem.*, 1985, **24**, 696

It is thermally unstable and may easily explode.
See related AMMINEMETAL OXOSALTS

3894. Hexacyclohexyldilead
[6713-82-2]




Halocarbons

Houben-Weyl, 1975, Vol. 13.3, 168

In absence of other solvent and in presence of air, interaction with carbon tetrabromide is explosive, and with hexabromoethane, more so.

See related ALKYL METALS

3895. Lead oleate
[1120-46-3]





Mineral oil

Williams, C. G., *Mech. Eng.*, 1932, **54**, 128–129

Lead oleate greases tend to cause violent explosions when used on hot-running bearings. The cause is not immediately apparent, but peroxidation may well have been involved.

See other HEAVY METAL DERIVATIVES

3896. Calcium stearate
[1592-23-0]





1. Schmutzler, G. *et al.*, *Plaste Kaut.*, 1967, **14**, 827–829
2. Gossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

Dust explosions and spontaneous ignition hazards for calcium stearate and related plastics additives are detailed and discussed [1]. A maximum pressure increase of 6.6 bar and a maximum rate of rise above 680 bar/s have been recorded [2].


See other DUST EXPLOSION INCIDENTS, AUTOIGNITION INCIDENTS

3897. Zinc stearate
[557-05-1]




The finely powdered soap is a significant dust explosion hazard.
See entry DUST EXPLOSION INCIDENTS (reference 22)

3898. Carbonyl-bis(triphenylphosphine)iridium–silver diperchlorate
[]




Kuyper, J. *et al.*, *J. Organomet. Chem.*, 1976, **107**, 130
Highly explosive, detonated by heat or shock.
See related AMMINEMETAL OXOSALTS

3899. 1,2-Bis(triphenylphosphonio)ethane permanganate
[76710-76-4]




It explodes at 80–90°C.
See entry QUATERNARY OXIDANTS

3900. Calcium abietate
[13463-98-4]



See entry METAL ABIETATES



3901. Manganese abietate
[54675-76-2]



See entry METAL ABIETATES

3902. Lead abietate
[5434-72-0]



C₄₀H₅₈O₄Pb



See entry METAL ABIETATES

3903. Zinc abietate
[6798-76-1]

C₄₀H₅₈O₄Zn




See entry METAL ABIETATES

3904. (Octacarbondioxide) pentamer

[]


C₄₀O₁₀



See Oligo(octacarbondioxide)
See other ACETYLENIC COMPOUNDS

**3905. Hexaspiro[2.4.2.4.2.4.2.4.2.4]dotetraconta-
4,6,11,13,18,20,25,27,32,34,39,41-dodecayne**
[155797-52-7]

C₄₂H₂₄



1. de Meijere, A. *et al.*, *Angew. Chem. (Int.)*, 1994, **33**(8), 869


2. de Meijere, A. *et al.*, *Chem. Eur. J.* 1995, **1**(2), 124

The white solid explodes on friction or impact (photo on the cover of this *Angewandte*) and decomposes at 130°C. The instability is associated with the cyclopropane rings as it is claimed to be absent from the dodecamethyl homologue [1]. Calculation indicates 6 kJ/g available energy [2].

See other STRAINED-RING COMPOUNDS, ACETYLENIC COMPOUNDS, ROTANES

3906. Hexakis(pyridine)iron(II) tridecacarbonyltetraferrate(2-)

[23129-50-2]



C₄₃H₃₀Fe₅N₆O₁₃

Brauer, 1965, Vol. 2, 1758


Extremely pyrophoric in air.

See related AMMINOMETAL OXOSALTS, CARBONYLMETALS

3907. 5,10,15,20-Tetrakis(2-nitrophenyl)porphine

[22834-73-8]

C₄₄H₂₆N₈O₈









Preparative hazard

See 2-Nitrobenzaldehyde: Pyrrole







See other NITROARYL COMPOUNDS

3908. Tetrakis(4-*N*-methylpyridinio)porphinecobalt(III)(5+) perchlorate
[53149-77-2] **C₄₄H₃₆Cl₅CoN₈O₂₀**



Pasternack, R. F. *et al.*, *J. Inorg. Nucl. Chem.*, 1974, **36**, 600
 A sample obtained by precipitation exploded.
 See other AMMINEMETAL OXOSALTS

3909. Tetrakis(4-*N*-methylpyridinio)porphineiron(III)(5+) perchlorate
[64365-00-0] **C₄₄H₃₆Cl₅FeN₈O₂₀**




Harris, F. L. *et al.*, *Inorg. Chem.*, 1978, **17**, 71

A small sample exploded when scraped with a metal spatula on a glass frit.

See other AMMINEMETAL OXOSALTS

3910. 1,4,8,11,15,18,22,25-Octamethyl-29H,31H-tetrabenzo[b.g.l.q]porphinato(2-)cobalt(II)
[27662-34-6]

$C_{44}H_{36}N_4Co$




Preparative hazard

See 1,3,4,7-Tetramethylisoindole: Cobalt

3911. Oxybis(N,N-dimethylacetamidetriphenylstibonium) perchlorate

[37035-60-2]

$C_{44}H_{50}Cl_2N_2O_{11}Sb_2$




Goel, R. G. *et al.*, *Inorg. Chem.*, 1972, **11**, 2143

It exploded on several occasions during handling and attempted analysis.

See related AMMINEMETAL OXOSALTS

3912. Heptaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]nonatetraconta-4,6,11,13,18,20,25,27,32,34,39,41,46,48-tetradecayne
[168848-73-5]


C₄₉H₂₈



See other ROTANES

3913. Bis[dicarbonyl(cyclopentadienyl)tributylphosphinemolybdenum]-tetrakis(tetrahydrofuran)magnesium
[82148-84-3]

C₅₄H₉₆MgMo₂O₈P₂




McVicker, G. B., *Inorg. Synth.*, 1976, **16**, 59–61

This and other transition metal carbonyl derivatives of magnesium is pyrophoric.

See related CARBONYLMETALS See other ORGANOMETALLICS

3914. Aluminium stearate
[637-12-7]

C₅₄H₁₀₅AlO₆




The finely powdered soap is a significant dust explosion hazard.

See entry DUST EXPLOSION INCIDENTS (reference 22)

3915. Octaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]hexapentaconta-4,6,11,13,18,20,25,27,32,34,39,41,46,48,53,55-hexadecayne
 [164077-36-5]

C₅₆H₃₂




See other ROTANES

3916. Buckminsterfullerene
 [99685-96-8]
See Carbon

C₆₀

3917. Aluminium abietate
 [32454-63-0]

C₆₀H₈₇AlO₆



See entry METAL ABIETATES

3918. Nonaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]trihexaconta-4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62-octadecayne
 [155797-58-3]

C₆₃H₃₆



See other ROTANES


3919. Fullerene
 []
See Carbon

C₇₀

1313

3920. Decaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]heptaconta-4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62,67,69-eicosayne [168848-75-7]


C₇₀H₄₀



See other ROTANES

3921. Dodecaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]tetraoctaconta-4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62,67,69,74,76,81,83-tetracosayne [168848-76-8]

C₈₄H₄₈



See other ROTANES

3922. Calcium [7440-70-2]

Ca

Ca

FPA H121, 1983; *HCS 1980*, 262

'Product Information Sheet No. 212', Sandwich, Pfizer Chemicals, 1969
Calcium is pyrophoric when finely divided.

See other PYROPHORIC METALS

Alkalies

Merck, 1983, 227

Interaction with alkali-metal hydroxides or carbonates may be explosive.

Ammonia

1. Partington, 1967, 369

2. Gibson, 1969, 369

At ambient temperature, ammonia gas reacts exothermally with calcium, but if warmed the latter becomes incandescent [1]. The metal dissolves unchanged in liquid ammonia, but if the latter evaporates, the finely divided metal is pyrophoric [2].

See other PYROPHORIC METALS

Asbestos cement

Scott, P. J., *School Sci. Rev.*, 1967, **49**(167), 252

Drops of molten calcium falling on to hard asbestos cement sheeting caused a violent explosion which perforated the sheet. Interaction with sorbed water seems likely to have occurred.

See Water, below

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, Nitrogen

Chlorine fluorides

See Chlorine trifluoride: Metals

Chlorine pentafluoride: Metals

1,2-Diaminoethane

Benkeser, R. A. *et al.*, *Tetrahedron Lett.*, 1984, **25**, 2089–2092

The use of calcium in 1,2-diaminoethane as a safer substitute for sodium or lithium in liquid ammonia for the improved Birch reduction of aromatic hydrocarbons is described in detail.

Dinitrogen tetroxide

See Dinitrogen tetroxide: Metals

Halogens

MRH Chlorine 7.15/64

Mellor, 1941, Vol. 3, 638

Massive calcium ignites in fluorine at ambient temperature, and finely divided (but not massive) calcium ignites in chlorine.

Lead chloride

Mellor, 1941, Vol. 3, 369

Interaction is explosive on warming.

Mercury

Pascal, 1958, Vol. 4, 290

Amalgam formation at 390°C is violent.

Other reactants

Yoshida, 1980, 98

MRH values calculated for 16 combinations, largely with oxidants, are given.

Phosphorus(V) oxide

See Tetraphosphorus decaoxide: Metals

Silicon

Mellor, 1940, Vol. 6, 176–177

Interaction is violently incandescent above 1050°C after a short delay.

Sodium, Mixed oxides

Anon., *BCISC Quart. Safety Summ.*, 1966, **37**(145), 6

An operator working above the charging hole of a sludge reactor was severely burned when a quantity of burning sludge containing calcium and sodium metals and their oxides was ejected. This very reactive mixture is believed to have been ignited by drops of perspiration falling from the operator.

See Water, below

Sulfur

MRH 6.69/44

Mellor, 1941, Vol. 3, 639

A mixture reacts explosively when ignited.

Sulfur, Vanadium(V) oxide
See Vanadium(V) oxide: Calcium, etc.

Water

'Product Information Sheet No. 212', Sandwich, Pfizer Chemicals, 1969
Calcium or its alloys react violently with water (or dilute acids) and the heat of reaction may ignite evolved hydrogen under appropriate contact conditions.

See ASBESTOS CEMENT, ABOVE *See other* METALS

3923. Calcium chloride

[10043-52-4]

CaCl₂

CaCl₂

HCS 1980, 247

Methyl vinyl ether

See Methyl vinyl ether: Acids

Water

1. *MCA Case History No. 69*
2. *MCA Case History No. 2300*

The exotherm produced by adding solid calcium chloride to hot water caused violent boiling [1], and the same happened 25 years later [2].

Zinc

Anon., *ABCM Quart. Safety Summ.*, 1932, **3**, 35

Prolonged action of calcium chloride solution upon the zinc coating of a galvanised iron vessel caused slow evolution of hydrogen, which became ignited and exploded.

See other CORROSION INCIDENTS, GAS EVOLUTION INCIDENTS

See other METAL HALIDES

3924. Calcium hypochlorite

[7778-54-3]

CaCl₂O₂

Ca(OCl)₂

FPA H46, 1976; HCS 1980, 269

1. Sidgwick, 1950, 1217
2. Clancey, V. J., *J. Haz. Mat.*, 1975, **1**, 83–94
3. Cave, R. F., *Chem. in Austr.*, 1978, **45**, 313
4. Uehara, Y. *et al.*, *Chem. Abs.*, 1979, **91**, 159834
5. Uehara, Y. *et al.*, *Chem. Abs.*, 1979, **91**, 180693
6. Uehara, Y. *et al.*, *Chem. Abs.*, 1980, **92**, 220207
7. Clancey, V. J., *Hazards of Pressure*, IChE Symp. Ser. No. 102, 11–23, Oxford, Pergamon, 1987
8. Wejtowicz, J. A., *ibid.*, 323–351

This powerful oxidant is technically of great importance for bleaching and sterilisation applications, and contact with reductants or combustible materials must be

CaCl₂O₂

under controlled conditions. It is present in diluted state in bleaching powder, which is a less powerful oxidant with lower available chlorine content [1]. There have been several instances of mild explosions and/or intense fires on ships carrying cargoes of commercial hypochlorite, usually packed in lacquered steel drums and of Japanese origin. Investigational work was still in progress, but preliminary indications suggested that presence of magnesium oxide in the lime used to prepare the hypochlorite may have led to the presence of magnesium hypochlorite which is known to be of very limited stability. Unlike many other oxidants, the hypochlorite constitutes a hazard in the absence of other combustible materials, because after initiation, local rapid thermal decomposition will spread through the contained mass of hypochlorite as a vigorous fire which *evolves*, rather than consumes, oxygen [2]. Thermal decomposition with evolution of oxygen and chlorine sets in around 175°C and will propagate rapidly through the whole bulk.

Several instances of containers or drums igniting or erupting on reopening after previous use have been reported, and may well have involved formation by slow hydrolysis of a relatively high concentration of dichlorine monoxide in the containers. This may have been caused to decompose from sudden exposure to light, by friction on opening the drum, or by a static spark. In one incident, a dull (gas) explosion was heard immediately before the solid erupted. Dry storage is essential [3]. Extrapolation from small-scale determinations of critical thermal ignition temperatures to commercial dimensions (35.4 or 38.2 cm diameters) gave values of 76.3 and 75°, as compared with experimental values of 77 and 75°C, respectively [4]. Fire and explosion hazards of high strength (70%) calcium hypochlorite were studied, with measurements of thermal properties, ignition temperatures, shock-sensitivity, flame propagation rates and explosive power for mixtures with organic materials [5]. A review of the results, and their comparison with predictions from computer simulation has been published [6]. Further possible causes of the instability of commercial 70% calcium hypochlorite are discussed [7], but it is held that the material, if stored and used correctly, is a relatively stable product in the absence of impurities [8]. The recently calculated value of 75°C for critical ignition temperature agrees with the previously recorded value of 76°C.

See BLEACHING POWDER

See related FRICTIONAL INITIATION INCIDENTS, IRRADIATION DECOMPOSITION INCIDENTS, STATIC INITIATION INCIDENTS

Acetylene

Rüst, 1948, 338

Contact of acetylene with bleaching powder, etc., may lead (as with chlorine itself) to formation of explosive chloroacetylenes. Application of hot water to free a partially frozen acetylene purification system which contained bleaching powder caused a violent explosion to occur.

See related HALOACETYLENE DERIVATIVES

Acetic acid, Potassium cyanide

Pitt, M. J., private comm., 1984

A jug containing calcium hypochlorite (probably as moist solid) was used as a disposal receptacle for cyanide wastes from student preparations of benzoin. When a little acetic acid residue was inadvertently added, an explosion occurred, attributed to a cyanide–chlorine redox reaction.

See other REDOX REACTIONS

Algaecide

Cave, R. F., *Chem. in Austr.*, 1978, **45**, 314

When algaecide (possibly a conc. solution of a quaternary ammonium salt) was spilt on a drum of swimming pool sterilant, it ignited and flared up immediately.

Ammonium chloride

MRH 1.05/27

1. Morris, D. L., *Science Teach.*, 1968, **35**(6), 4

2. Anderson, M. B., *Science Teach.*, 1968, **35**(9), 4

3. Rai, H., *Chem. News*, 1918, **117**, 253

When calcium hypochlorite was inadvertently used in place of calcium hydroxide to prepare ammonia gas, an explosion occurred [1], attributed to formation of nitrogen trichloride [2]; this was early claimed to be one of the most efficient methods of preparing the trichloride [3].

See Nitrogenous bases, below

Calcium carbide

Saville, B. W., 1988, private communication

A mixture explodes violently on addition of a drop of water.

Carbon

Mellor, 1941, Vol. 2, 262

A confined intimate mixture of hypochlorite and finely divided charcoal exploded on heating.

Carbon tetrachloride

491M, 1975, 84

Use of a carbon tetrachloride extinguisher on a fire in an open container of hypochlorite caused a severe explosion.

See Oxygen (Gas): Halocarbons

Contaminants

1. *MCA Case History No. 666*

2. Cave, R. F., *Chem. in Austr.*, 1971, **45**, 314

The contents of a drum erupted and ignited during intermittent use. This was attributed to contamination of the soldered metal scoop (normally kept in the drum) by oil, grease or water, or all three, and subsequent exothermic reaction with the oxidant [1]. Use of a bucket, possibly contaminated with traces of engine oil, to transport the oxidant led to eruption and ignition of the contents [2].

See Iron oxide, below; *and* Organic matter, below

N,N-Dichloromethylamine

See N,N-Dichloromethylamine: Calcium hypochlorite, etc.

Hydrocarbons

Young, J. A., *CHAS Notes*, 1992, **X**(5), 3

'Gasoline' inflames some minutes after being poured onto the hypochlorite. A fire consequent upon transporting the two in adjacent containers in a car is reported. The editor imagines that non-hydrocarbon fuel additives may also be involved.

See Organic matter, etc., below

Hydroxy compounds

MRH values below reference

1. Fawcett, H. H., *Ind. Eng. Chem.*, 1959, **51**(4), 90A

2. Kirkbride, K. P. *et al.*, *J. Forensic Sci.*, 1991, **36**(3), 902

MRH values: Glycerol 13/12, phenol 2.26/7, ethanol 2.30/7, methanol 2.34/10

Contact of the solid hypochlorite with glycerol, diethylene glycol monomethyl ether or phenol causes ignition within a few min, accompanied by irritant smoke, especially with phenol (formation of chlorophenols). Ethanol may cause an explosion, as may methanol, undoubtedly owing to formation of the alkyl hypochlorites [1]. Reaction with polyethylene glycol hydraulic fluid may produce ignition, with a fireball. A mechanism for creation of combustible gases is proposed [2].

Iron oxide (rust)

Gill, A. H., *Ind. Eng. Chem.*, 1924, **16**, 577

Presence of rust in metal containers has caused many accidental explosions. Metal oxides catalyse the oxygen-evolving decomposition of the oxidant.

Nitrogenous bases

MRH Ethylamine 2.92/15

Kirk-Othmer, 1963, Vol. 2, 105

Primary aliphatic or aromatic amines react with calcium (or sodium) hypochlorite to form *N*-mono- or di-chloroamines which are explosively unstable, but less so than nitrogen trichloride.

Nitromethane

MRH 5.69/99+

Fawcett, H. H., *Trans. Nat. Safety Congr. Chem. Fertilizer Ind.*, Vol. 5, 32, Chicago, NSC, 1963

They interact, after a delay, with extreme vigour.

Oil, Polyethylene film, Water, Wood

1. Armstrong, B. S., *Loss Prev. Bull.*, 1988, (084), 25–31

2. Sampson, R. G., private comm., 1989

Impact damage occurred to one of 48 10 kg polypropylene tubs of calcium hypochlorite, each closed with a polythene lid, all supported on a wooden pallet and stretch-wrapped round the base with polyethylene film. The spilled material was swept up and discarded, and the pallet was moved to another part of the warehouse, and some 30 mins. later flames were seen at the base of the pallet [1]. Ignition was attributed to contact between residual solid hypochlorite on the pallet and lubricant drips on the concrete floor from fork lift traffic. Application of a dry powder extinguisher had little effect on the fire, but use of a 40 l water extinguisher led to rapid escalation of the fire as contact between hypochlorite

and wood was increased. Flash-over occurred and the fire eventually involved a considerable fraction of the 1400 t of assorted warehouse contents [2].

See Organic matter, etc., below

See other IGNITION SOURCES

Other reactants

Yoshida, 1980, 133

MRH values calculated for 11 combinations with oxidisable materials are given.

Organic matter

1. 'Halane' Information Sheet, Wyandotte Chem. Co., Michigan, 1958

2. Weicherz, J., *Chem. Ztg.*, 1928, **52**, 729

3. Cave, R. F., *Chem. in Austr.*, 1978, **45**, 313

Mixtures of the solid hypochlorite with 1% of admixed organic contaminants are sensitive to heat in varying degree. Wood caused ignition at 176°, while oil caused violent explosion at 135°C [1]. When solid hypochlorite was transferred to a bucket which had been cleaned out with an alcoholic cleaning composition, it ignited violently. This could not be repeated, but it was found that many organic liquids and fibrous materials caused ignition if warmed to 100°C with the oxidant [2]. Heating a little solid oxidant in a lightly greased crucible led to violent ignition around 180°C [3].

See reference 5 above; Contaminants, above; and Turpentine, below

Organic matter, Water

1. Tatara, S. *et al.*, Ger. Offen., 2 450 816, 1975

2. Lawrence, S., Internet, 1996

Calcium hypochlorite containing over 60% active chlorine normally ignites in contact with lubricating oils, but addition of 16–22% of water will prevent this. Examples show the effect of 18% (but not 15%) of added water in preventing ignition when glycerol was dripped onto a hypochlorite (79% active chlorine) containing 2% of oil [1]. Ignition with brake fluid is also reported, in this case reaction is accelerated by cola soft drinks — which are dilute acids rather than just water [2].

Organic sulfur compounds

1. Stephenson, F. G., *Chem. Eng. News*, 1973, **51**(26), 14

2. Laboratory Waste Disposal Manual, 142, Washington, MCA, 1969–1972 edns (withdrawn in 1980)

3. *MCA Guide for Safety in the Chemical Laboratory*, 464, New York, Van Nostrand-Reinhold, 1972

4. *491M*, 1975, 85

5. Wacker, J. V. *et al.*, *Rec. Trav. Chim.*, 1986, **105**(3), 99–107

Contact of the solid oxidant with organic thiols or sulfides may cause a violent reaction and flash fire [1]. This procedure was recommended formerly for treating spills of sulfur compounds [2,3], but is now withdrawn as potentially hazardous. Use of an aqueous solution of up to 15% concentration, or of 5% sodium hypochlorite solution is recommended [1]. Addition of 10 g of oxidant to 5 ml portions of

1-propanethiol or isobutanethiol led to explosions [4]. Application of factorial design techniques to experimental planning gave specific conditions for the safe oxidation of organic sulfides to sulfoxides using calcium hypochlorite or sodium chlorate [5].

See Sulfur, below

Sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide

See Sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide: Calcium hypochlorite

Sodium hydrogen sulfate, Starch, Sodium carbonate

Anon., *Ind. Eng. Chem. (News Ed.)*, 1937, **15**, 282

Shortly after a mixture of the four ingredients had been compressed into tablets, incandescence and an explosion occurred. This may have been caused by interaction of the oxidant and starch, accelerated by the acid salt, but may also have involved dichlorine monoxide liberated by the same salt.

Sulfur

Katz, S. A. *et al.*, *Chem. Eng. News*, 1965, **46**(29), 6

Admixture of damp sulfur and 'swimming pool chlorine' caused a violently exothermic reaction, and ejection of molten sulfur.

Turpentine

491M, 1975, 85

An explosion occurred soon after the oxidant was put into a can which had contained turpentine but was thought to be empty.

See other METAL HYPOCHLORITES, OXIDANTS

3925. Calcium chlorite

[14674-72-7]



Potassium thiocyanate

Pascal, 1960, Vol. 16, 264

Mixtures may ignite spontaneously.

See other CHLORITE SALTS, METAL OXOHALOGENATES

3926. Calcium chromate

[13816-48-3]



Boron

1. Rogers, J. W. *et al.*, *Energy Res. Abs.*, 1981, **6**, 20813

2. Rogers, J. W., *Proc. 8th Int. Pyrotech. Seminar*, 1982, 556–573

Pyrotechnic properties of blends with 20% boron were studied [1], and the performance of several other blends has been assessed [2].

See other METAL OXOMETALLATES, OXIDANTS

3927. Calcium hydride
[7789-78-8]

CaH₂

CaH₂

HCS 1980, 268

Halogens

Mellor, 1941, Vol. 16, 651

Heating the hydride strongly with chlorine, bromine or iodine leads to incandescence.

Hexachlorobenzene

Loiselle, S. *et al.*, *Environ. Sci. Technol.* 1997, **31**(1), 261; *J. Solid State Chem.*, 1997, **129**(2), 263

It was discovered that milling hexachlorobenzene with calcium hydride, to effect dechlorination, could give rise to explosive reaction, the conditions under which this happens were studied.

Manganese dioxide

See Manganese(IV) oxide: Calcium hydride

Metal oxohalogenates

MRH values: Ammonium bromate 4.52/82, ammonium chlorate 6.36/76, ammonium perchlorate 6.53/70, barium bromate 3.97/76, barium chlorate 4.85/72, calcium bromate 5.02/70, calcium chlorate 6.53/62, potassium bromate 4.56/73, potassium chlorate 5.81/56, potassium perchlorate 6.07/63, silver chlorate 4.60/73, sodium bromate 4.85/70, sodium chlorate 6.90/63, sodium perchlorate 6.69/60

1. Mellor, 1946, Vol. 3, 651

2. Yoshida, 1980, 214

Mixtures of the hydride with various bromates, chlorates and perchlorates explode on grinding [1]. MRH values calculated for 14 such combinations are given [2].

Other reactants

See next above

Silver halides

Mellor, 1941, Vol. 3, 389, 651

A mixture of silver fluoride and the hydride incandescens on grinding, and the iodide reacts vigorously on heating.


Tetrahydrofuran

See Tetrahydrofuran: Lithium tetrahydroaluminate

See other METAL HYDRIDES

3928. Calcium hydroxide

[1305-62-0]



Polychlorinated phenols, Potassium nitrate

Kozloski, R. P., *Chem. Eng. News*, 1982, **60**(11), 3

When chlorinated phenols are heated for analytical purposes with calcium hydroxide–potassium nitrate mixtures, chlorinated benzodioxins analogous to the extremely toxic tetrachlorodibenzodioxin may be formed.

Sugars

See SUGARS

1,3,5-Trinitrohexahydro-1,3,5-triazine

See 1,3,5-Trinitrohexahydro-1,3,5-triazine: Calcium hydroxide, Water*See other* INORGANIC BASES**3929. Calcium hydroxide *O*-hydroxylamide**

[]



Sorbe, 1968, 43

It explodes on heating (and presumably would be formed if distillation of hydroxylamine from calcium oxide were attempted).

See Hydroxylamine: Metals*See other* N–O COMPOUNDS**3930. Calcium bis(*O*-hydroxylamide)**

[]

*See* Hydroxylamine: Metals*See other* N–O COMPOUNDS**3931. Calcium phosphinate (Calcium hypophosphite)**

[7789-79-9]



Other reactants

Yoshida, 1980, 344

MRH values calculated for 8 combinations with various reagents are given.

Potassium chlorate

MRH 2.88/49

See Potassium chlorate: Reducants*See other* METAL PHOSPHINATES, REDUCANTS

3932. Calcium hydrazide

[]




Hydrazine and its Derivatives, Schmidt, F. W., New York, Wiley, 1984, 75

Liabile to explode on preparation from the metal, or amide, and hydrazine. Other authors, however, report a 1:1 hydrazide [28330-64-5] as being fairly stable.

See other METAL HYDRAZIDES

3933. Hexaamminecalcium

[12133-31-2]



491M, 1975, 83

It ignites in air.

See related N-METAL DERIVATIVES

3934. Calcium permanganate

[10118-76-0]



Acetic acid

MRH 2.93/18

MRH ,

Acetic anhydride

See Potassium permanganate: Acetic acid, etc.

MRH 3.14/15

Cellulose

Kirk-Othmer, (3rd ed) 1982, Vol. 19, 494

Calcium permanganate is a more active oxidant than the potassium salt, and ignites paper or cotton on contact.

Hydrogen peroxide

See Hydrogen peroxide: Metals, etc.

MRH 1.59/99+

Other reactants

Yoshida, 1980, 92

MRH values calculated for 18 combinations, largely with oxidisable materials, are given.

See other OXIDANTS

3935. Calcium nitrate
[10124-37-5]



HCS 1980, 270

Aluminium, Ammonium nitrate, Formamide, Water MRH Aluminium 9.33/36

1. Wilson, J. F. *et al.*, S. Afr. Pat. 74 03 305, 1974
2. Slykehouse, T. E., Can. Pat. 1 011 561, 1977

A mixture containing 51% of calcium nitrate and 12% ammonium nitrate with 27% formamide and 10% water is detonable at -20°C [1]. Addition of aluminium powder improves performance as a blasting explosive [2].

Ammonium nitrate, Hydrocarbon oil

Haid, A. *et al.*, *Jahresber. Chem. Tech. Reichsanst.*, 1930, **8**, 108–115

The explosibility of the double salt, calcium ammonium nitrate, is enhanced by presence of oil.

See Ammonium nitrate: Organic fuels

Ammonium nitrate, Water-soluble fuels

Clark, W. F. *et al.*, US Pat. 3 839 107, 1970

Up to, or over 40% of the ammonium nitrate content of explosive mixtures with water-soluble organic fuels may be replaced with advantage by calcium nitrate.

Organic materials

Sorbe, 1968, 423

Mixtures of the nitrate with organic materials may be explosive.

Other reactants

Yoshida, 1980, 193

MRH values calculated for 16 combinations with oxidisable materials are given.

See other METAL NITRATES, OXIDANTS

3936. Calcium azide
[19465-88-8]



1. Mellor, 1940, Vol. 8, 349
2. Stull, 1977, 10

Calcium, strontium and barium azides are not shock-sensitive, but explode on heating at about 150, 170 and 225 (or 152) $^\circ\text{C}$, respectively. In sealed tubes, the explosion temperatures are higher [1]. Although calcium azide is rather mildly endothermic (ΔH_f° (s) +46 kJ/mol, 0.37 kJ/g), it can decompose much more exothermally to the nitride (189.9 kJ/mol, 1.53 kJ/g) than to the elements [2].

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

3937. Calcium oxide
[1305-78-8]

CaO

CaO

NSC 241, 1982; *HCS 1980*, 272

Ethanol

Keusler, V., *Apparatebau*, 1928, **40**, 88–89

The lime–alcohol residue from preparation of anhydrous alcohol ignited on discharge from the still and caused a vapour explosion. The finely divided and reactive lime may have heated on exposure to atmospheric moisture and caused ignition.

See Water, below

Hydrogen fluoride

See Hydrogen fluoride: Oxides

Interhalogens

See Bromine pentafluoride: Acids, etc.

MRH 3.23/56

Chlorine trifluoride

MRH 4.06/52

Other reactants

Yoshida, 1980, 126

MRH values calculated for 6 combinations with various reagents are given.

Phosphorus(V) oxide

See Tetraphosphorus decaoxide: Inorganic bases

Water

1. Mellor, 1941, Vol. 3, 673
2. Anon., *Fire*, 1935, **28**, 30
3. Amos, T., *Zentr. Zuckerind.*, 1923, **32**, 103
4. Anon., *BCISC Quart. Safety Summ.*, 1967, **38**, 15
5. Anon., *BCISC Quart. Safety Summ.*, 1971, **42**(168), 4
6. Anon., *Engrg. Digest*, 1908, **4**, 661

Crystalline calcium oxide reacts imperceptibly slowly with water, but the powdered material reacts with explosive violence after a few minutes delay [1]. Quicklime, when mixed with $\frac{1}{3}$ its weight of water, will reach 150–300°C (depending on quantity) and may ignite combustible material. Occasionally 800–900°C has been attained [2]. Moisture present in wooden storage bins caused ignition of the latter [3]. A water jet was used unsuccessfully to clear a pump hose blocked with quicklime. On standing, the exothermic reaction generated enough steam to clear the blocked hose explosively [4]. Two glass bottles of calcium oxide powder burst while on a laboratory shelf, owing to the considerable increase in bulk which occurs on hydration. Storage in plastic bottles is recommended, and granular oxide should cause fewer problems in this respect than the powder [5]. The powerful expansion effect was formerly used in coal-mining operations [6].

See other GLASS INCIDENTS
See other INORGANIC BASES, METAL OXIDES

3938. Calcium peroxide
[1305-79-9]

CaO₂

CaO₂

Chlorinated paraffin

Anon., *Chem. Brit.*, 1997, **33**(6), 7

A fire is reported subsequent to industrial mixing of calcium peroxide and a chlorinated paraffin. Even completely chlorinated hydrocarbons can be oxidised, to phosgene.

See Oxidisable materials; below

Other reactants

Yoshida, 1980, 84

MRH values calculated for 14 combinations with oxidisable materials are given.

Oxidisable materials

MRH Magnesium 6.02/25

1. Castrantas, 1965, 4
2. Sorbe, 1968, 43

Grinding the peroxide with oxidisable materials may cause fire [1], and the octahydrated solid explodes at high temperature [2].

Polysulfide polymers

491M, 1975, 87

The dry peroxide, added to cause cross-linking in liquid polysulfide polymers with pendant thiol groups, caused sparking or ignition, depending on the scale of the process.

See other METAL PEROXIDES, OXIDANTS

3939. Calcium sulfate
[7778-18-9]

CaSO₄

CaO₄S

Aluminium

See Aluminium: Metal oxides, or Oxosalts

Diazomethane

See Diazomethane: Calcium sulfate

Phosphorus

See Phosphorus: Metal sulfates

See other METAL OXONON-METALLATES

- 3940. Calcium peroxodisulfate**
 [13235-16-0] **CaO₈S₂**
 $\text{Ca}(\text{OSO}_2\text{O})_2$
- Castrantas, 1965, 6
 It is shock-sensitive and explodes violently.
See other OXIDANTS, PEROXOACID SALTS
- 3941. Calcium sulfide**
 [20548-54-3] **CaS**
CaS
- HCS 1980, 274*
- Oxidants
 Mellor, 1941, Vol. 3, 745
 Alkaline-earth sulfides react vigorously with chromyl chloride, lead dioxide, potassium chlorate (explodes lightly) and potassium nitrate (explodes violently).
See other METAL SULFIDES
- 3942. Calcium polysulfide**
 [1332-67-8] (tetrasulfide) **CaS_x**
 [1332-69-9] (pentasulfide)
CaS_x
- Edwards, P. W., *Chem. Met. Eng.*, 1922, **27**, 986
 The powdered sulfide, admixed with small amounts of calcium thiosulfate and sulfur, has been involved in several fires and explosions, some involving initiation by static discharges.
See other METAL SULFIDES, STATIC INITIATION INCIDENTS
- 3943. Calcium silicide**
 [12013-55-7] **CaSi**
CaSi
- Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74
 - Yan, J. *et al.*, *Chem. Abs.*, 1987, **107**, 182628
 - Deng, X. *et al.*, *Arch. Combust.*, 1987, **7**(1–2), 19–31
- The finely powdered silicide is a significant dust explosion hazard [1]. The lower explosion limit for a calcium–silicon dust cloud of mean particle size 9.7 μm was measured as 79 g/m³, in good agreement with a calculated value [2]. Other dust cloud parameters are presented and related to predictions [3].
- Acids
 Mellor, 1940, Vol. 6, 177
 Interaction is vigorous, and the evolved silanes ignite in air.

Boron

See Boron: Calcium–silicon alloy

See other METAL NON-METALLIDES, PYROPHORIC MATERIALS

3944. Calcium disilicide

[12013-56-8]

CaSi₂

CaSi₂

Carbon tetrachloride

Zirconium Fire and Explosion Incidents, TID-5365, Washington, USAEC, 1956

Calcium disilicide exploded when milled in the solvent.

Iron(III) oxide

See Iron(III) oxide: Calcium disilicide

Metal fluorides

Berger, E., *Compt. rend.*, 1920, **170**, 29

Calcium disilicide ignites in close contact with alkali metal fluorides, forming silicon tetrafluoride.

Other reactants

Yoshida, 1980, 99

MRH values calculated for 14 combinations with oxidants are given.

Potassium nitrate

See Potassium nitrate: Calcium silicide

See other METAL NON-METALLIDES

3945. Calcium triperoxochromate

[]

Ca₃Cr₂O₁₂

Ca₃[OO)₃Cr]₂

Raynolds, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 1851

A buff solid, prepared by means appropriate to a tetraperoxochromate(3–), it analysed as the dodecahydrate of this structure. It explodes at 100°C.

See other OXIDANTS, PEROXOACID SALTS

3946. Calcium nitride

[12013-82-0]

Ca₃N₂

Ca₃N₂

von Schwartz, 1918, 322

Spontaneously flammable in air (probably when finely divided and in moist air).

Halogens

Mellor, 1940, Vol. 8, 103

Reaction with incandescence in chlorine gas or bromine vapour.
See other N-METAL DERIVATIVES, NITRIDES

3947. Calcium phosphide
[1305-99-3]

Ca₃P₂

Ca₃P₂

Dichlorine oxide

See Dichlorine oxide: Oxidisable materials

Oxygen

Van Wazer, 1958, Vol. 1, 145

Calcium and other alkaline-earth phosphides incandesce in oxygen at about 300°C.

Water

Mellor, 1940, Vol. 8, 841

Calcium and other phosphides on contact with water liberate phosphine, which usually ignites in air, owing to the diphosphane content.

See other METAL NON-METALLIDES

3948. Columbium
[7440-03-1]

Cb

Cb

See Niobium

3949. Cadmium
[7440-43-9]

Cd

Cd

NSC 312, 1978; HCS 1980, 257; RSC Lab. Hazards Safety Data Sheet No. 55, 1987
(element and compounds)

The finely divided metal is pyrophoric.

See CADMIUM HYDRIDE *See other* PYROPHORIC METALS

Hydrazoic acid

See Cadmium azide (reference 2)

Oxidants

See Nitryl fluoride: Metals

Ammonium nitrate: Metals

Selenium, or Tellurium

Mellor, 1940, Vol. 4, 480

Reaction on warming powdered cadmium with selenium or tellurium is exothermic, but less vigorous than that of zinc.

See other METALS

3950. Trihydrazinecadmium chlorate

[]




Mellor, 1956, Vol. 2, Suppl. 2.1, 592

It explodes on impact.

See other AMMINEMETAL OXOSALTS

3951. Basic trihydrazinecadmium perchlorate

[]



Mellor, 1967, Vol. 8, Suppl. 2, 88

The monohydrated basic salt is extremely explosive.

See related AMMINEMETAL OXOSALTS

3952. Cadmium chlorate

[10137-74-3]



Sulfides

Mellor, 1956, Vol. 2, Suppl. 1, 584

Interaction with copper(II) sulfide is explosive, and with antimony(II) sulfide, arsenic(III) sulfide, tin(II) sulfide and tin(IV) sulfide, incandescent.

See METAL CHLORATES: phosphorus, etc.

also Chloric acid: Metal sulfides

See other METAL CHLORATES, METAL OXOHALOGENATES

3953. Cadmium hydride

[72172-64-6]



Barbaras, G. D. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 4585

The hydride, prepared at -78°C , suddenly decomposes during slow warming at 2°C , leaving a residue of pyrophoric cadmium.

See other METAL HYDRIDES

3954. Cadmium amide
[22750-53-4]




Alone, or Water

Mellor, 1940, Vol. 8, 261

When heated rapidly, the amide may explode. Reaction with water is violent.

See other N-METAL DERIVATIVES

3955. Trihydrazinecadmium(II) nitrate
[32434-37-5]



See HYDRAZINE METAL NITRATES

3956. Tetraamminecadmium permanganate
[34410-88-3]




Mellor, 1942, Vol. 12, 335

It explodes on impact.

See other AMMINOMETAL OXOSALTS

3957. Cadmium azide
[14215-29-3]



1. Turney, T. A., *Chem. & Ind.*, 1965, 1295

2. Mellor, 1967, Vol. 8, Suppl. 2.2, 25, 50

3. Schechkov, G. T., *Chem. Abs.*, 1983, **98**, 146010

A solution, prepared by mixing saturated solutions of cadmium sulfate and sodium azide in a 10 ml glass tube, exploded violently several hours after preparation [1].

The dry solid is extremely hazardous, exploding on heating or light friction. A violent explosion occurred with cadmium rods in contact with aqueous hydrogen azide [2].

A DTA study showed a lesser thermal stability than lead azide [3]. It is strongly endothermic (ΔH_f° (s) +451 kJ/mol, 2.32 kJ/g).

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

3958. Cadmium oxide
[1306-19-0]



HCS 1980, 260

Magnesium

See Magnesium: Metal oxides *See other* METAL OXIDES

3959. Cadmium selenide
[1306-24-7]

CdSe

CdSe

Preparative hazard

See Selenium: Metals

See related METAL SULFIDES

3960. Cadmium nitride
[12380-95-9]

Cd₃N₂

Cd₃N₂

Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1469

The shock of the violent explosion caused by heating a sample of the nitride caused an unheated adjacent sample to explode. It is much less endothermic (ΔH_f° (s) +161.5 kJ/mol, 0.44 kJ/g) than the azide.

Acids, or Bases

Mellor, 1964, Vol. 8, Suppl. 2.1, 161

It reacts explosively with dilute acids or bases.

Water

Mellor, 1940, Vol. 8, 261

It explodes on contact.

See other ENDOTHERMIC COMPOUNDS, *N*-METAL DERIVATIVES, NITRIDES

3961. Cerium
[7440-45-1]

Ce

Ce

Alone, or Metals

Mellor, 1945, Vol. 5, 602–603

Cerium or its alloys readily give incandive sparks (pyrophoric particles) on frictional contact, this effect of iron alloys being widely used in various forms of 'flint' lighters. The massive metal ignites and burns brightly at 160°C, and cerium wire burns in a Bunsen flame more brightly than magnesium. Of its alloys with aluminium, antimony, arsenic, bismuth, cadmium, calcium, copper, magnesium, mercury, sodium and zinc, those containing major proportions of cerium are often extremely pyrophoric. The mercury amalgams ignite spontaneously in air without the necessity for frictional generation of small particles. The interaction of cerium with zinc is explosively violent, and with antimony or bismuth, very exothermic.

See other PYROPHORIC ALLOYS

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, Nitrogen

Halogens

Mellor, 1946, Vol. 5, 603

Cerium filings ignite in chlorine or in bromine vapour at about 215°C.

Phosphorus

See Phosphorus: Metals

Silicon

Mellor, 1946, Vol. 5, 605

Interaction at 1400°C to form cerium silicide is violently exothermic, often destroying the containing vessel.

Water

Uno, T. *et al.*, *Anzen Kagaku*, 1988, **27**(3), 157–161 (*Chem. Abs.*, 1989, **110**, 12857)

The mixed alloy of cerium with lanthanum and other rare-earth metals ('Mischmetall') may ignite spontaneously in contact with aqueous solutions, owing to its oxidation by water and ignition of evolved hydrogen.

See other GAS EVOLUTION INCIDENTS, REDOX REACTIONS

See other METALS, PYROPHORIC METALS

3962. Cerium dihydride

[13569-50-1]



Libowitz, G. G. *et al.*, *Inorg. Synth.*, 1973, **14**, 189–192

The polycrystalline hydride is frequently pyrophoric, particularly at higher hydrogen contents (up to H_{2.85}), while monocrystalline material appears to be less reactive to air.

See other METAL HYDRIDES

3963. Cerium trihydride

[13864-02-3]



Muthmann, W. *et al.*, *Ann.*, 1902, **325**, 261

The hydride is stable in dry air, but may ignite in moist air.

See other METAL HYDRIDES

3964. Ammonium hexanitrocerate (Ceric ammonium nitrate, Ammonium cerium(IV) nitrate)

[16774-21-3]



Cyclopentadienylsodium

See Sodium nitrate: Tris(cyclopentadienyl)sodium

Potassium bromate, Malonic acid, Water

See Potassium bromate: Ceric ammonium nitrate, Malonic acid, Water.

See related METAL NITRATES *See other* OXIDANTS

3965. Cerium nitride

[25764-08-3]

CeN

CeN

Water

Mellor, 1940, Vol. 8, 121

Contact with water vapour slowly causes incandescence, while a limited amount of water or dilute acid causes rapid incandescence with ignition of evolved ammonia and hydrogen.

See other N-METAL DERIVATIVES, NITRIDES

3966. Cerium azide

[]

Ce(N₃)₃

CeN₉

Mellor, 1940, Vol. 8, 354

The precipitate from cerium nitrate and sodium azide is explosive.

See other METAL AZIDES

3967. Cerium trisulfide

[12014-93-6]

Ce₂S₃

Ce₂S₃

Mellor, 1946, Vol. 5, 649

It is pyrophoric in ambient air when finely divided.

See other METAL SULFIDES, PYROPHORIC MATERIALS

3968. Chromyl azide chloride

[14259-67-7]

CrO₂(N₃)Cl

ClCrN₃O₂

An explosive solid.

See entry METAL AZIDE HALIDES

3969. Caesium tetrafluorochlorate(1-)

[15321-67-7]

Cs[ClF₄]

ClCsF₄

Highly reactive oxidant.

See entry METAL POLYHALOHALOGENATES

3970. Caesium chloroxenate
[26283-13-6]

ClCsO_3Xe

$\text{Cs}[\text{ClXeO}_3]$

Jaselskis, B. *et al.*, *J. Amer. Chem. Soc.*, 1967, **89**, 2770
It explodes at 205°C under vacuum.
See other XENON COMPOUNDS

3971. Chlorine fluoride
[7790-89-8]

ClF

ClF

1. Sidgwick, 1950, 1149
2. 'Product Information Sheet ClF', Tulsa, Ozark-Mahoning Co., 1970
This powerful oxidant reacts with other materials similarly to chlorine trifluoride or fluorine [1], but more readily than the latter [2].
See Chlorine trifluoride, or Fluorine

Acetonitrile, Fluorine
See Fluorine: Acetonitrile, etc.

Aluminium
Mellor, 1956, Vol. 2, Suppl. 1, 63
Aluminium burns more readily in chlorine fluoride than in fluorine.

Bis(trifluoromethyl) disulfide
See Bis(trifluoromethyl) disulfide: Chlorine fluorides

Bis (trifluoromethyl) sulfide
See Bis(trifluoromethyl) sulfide: Chlorine fluorides

tert-Butanol
Young, D. F. *et al.*, *J. Amer. Chem. Soc.*, 1970, **92**, 2314
A mixture prepared at -196°C exploded at -100°C during slow warming.

N-Chlorosulfinylamine
Kuta, G. S. *et al.*, *Intern. J. Sulfur Chem.*, 1973, **8**, 335-340
Interaction of the reagents in equimolar proportions produced a highly explosive and strongly oxidising material (possibly an N-F compound).

Diethyl sulfite
See Ethyl fluorosulfate

Fluorocarbon polymers
'Product Information Sheet ClF', Tulsa, Ozark-Mahoning Co., 1970
Chlorine fluoride can probably ignite Teflon and Kel-F at high temperatures, or under friction or flow conditions.

Phosphorus trifluoride

1. Fox, W. B. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1971, **7**, 861

2. Neilson, R. H. *et al.*, *Inorg. Chem.*, 1975, **14**, 2019

Interaction of the fluorides to produce chlorotetrafluorophosphorane [1] is uncontrollably violent even at -196°C [2]. An improved method of making the phosphorane from phosphorus pentafluoride and boron trichloride is detailed [2].

Tellurium

Mellor, 1943, Vol. 11, 26

Interaction is incandescent.

Trifluoromethanesulfenyl chloride

See Trifluoromethanesulfenyl chloride: Chlorine fluorides

Trifluorosulfur nitride

Waterfeld, A. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1981, **20**, 1017

In the preparation of chloriminosulfur tetrafluoride or its dimer by slowly warming a mixture at -196° to -78°C and then ambient temperature, too rapid warming leads to violent explosions.

Trifluorosulfur nitride

See *N,N*-Dichloropentafluorosulfanylamine

Water


Pascal, 1960, Vol. 16.1, 189

Interaction with liquid water or that bound in crystalline hydrates is violent.

See other INTERHALOGENS

3972. Fluoronium perchlorate

[]



Water

Hantzsch, A., *Ber.*, 1930, **63**, 97

This hydrogen fluoride–perchloric acid complex reacts explosively with water.

See other NON-METAL PERCHLORATES

3973. Chloryl hypofluorite

[101672-08-6]



1. Hoffman, C. J., *Chem. Rev.*, 1964, **64**, 97

2. Christe, K. O. *et al.*, *Inorg. Chem.*, 1987, **56**, 920

Not then completely purified or characterised, its explosive nature was in contrast to the stability of the isomeric perchloryl fluoride [1]. It is also shock-sensitive [2].

See other HALOGEN OXIDES, HYPOHALITES

3974. Perchloryl fluoride
[7616-94-6]

CIFO₃

O₃ClF

1. 'Booklet DC-1819', Philadelphia, Pennsalt Chem. Corp., 1957
2. Anon., *Chem. Eng. News*, 1960, **38**, 62
3. Barton, D. H. R., *Pure Appl. Chem.*, 1970, **21**, 185
4. Sharts, C. M. *et al.*, *Org. React.*, 1974, **21**, 232–234

Procedures relevant to safe handling and use are discussed. Perchloryl fluoride is stable to heat, shock and moisture, but is a powerful oxidiser comparable with liquid oxygen. It forms flammable and/or explosive mixtures with combustible gases and vapours [1,2]. It only reacts with strongly nucleophilic centres, and the by-product, chloric acid is dangerously explosive in admixture with organic compounds [3]. Safety aspects of practical use of perchloryl fluoride have been reviewed [4].

Benzocyclobutene, Butyllithium, Potassium *tert*-butoxide

Adcock, W. *et al.*, *J. Organomet. Chem.*, 1975, **91**, C20

An attempt to convert benzocyclobutenylpotassium (prepared from the reagents above) to the fluoro derivative at -70°C with excess perchloryl fluoride led to a violent explosion when the reaction mixture was removed from the cooling bath after stirring for an hour.

Calcium acetylide, or Potassium cyanide, or Potassium thiocyanate, or Sodium iodide
Kirk-Othmer, 1966, Vol. 9, 602

Unreactive at 25° , these solids react explosively in the gas at $100\text{--}300^{\circ}\text{C}$.

Charcoal

Inman, C. E. *et al.*, *Friedel-Craft and Related Reactions*, (Olah, G. A., Ed.), Vol. 3, 1508, New York, Interscience, 1964

Adsorption of perchloryl fluoride on charcoal can, like liquid oxygen, produce a powerful 'Sprenkel' explosive.

Ethyl 4-fluorobenzoylacetate

Fuqua, S. A. *et al.*, *J. Org. Chem.*, 1964, **29**, 395, footnote 3

Fluorination of the ester with perchloryl fluoride by an established method led to a violent explosion.

Finely divided solids

McCoy, G., *Chem. Eng. News*, 1960, **38**(4), 62

Oxidisable organic materials of high surface to volume ratio (carbon powder, foamed elastomers, lampblack, sawdust) react very violently, even at -78°C , with perchloryl fluoride, which should be handled with the same precautions as liquid oxygen.

Hydrocarbons, or Hydrogen sulfide, or Nitrogen oxide, or Sulfur dichloride, or Vinylidene chloride

Braker, 1980, 578

At ambient temperature, perchloryl fluoride is unreactive with the compounds above, but reaction is explosive at 100–300°C, or if the mixtures are ignited.

3 α -Hydroxy-5 β -androstane-11,17-dione 17-hydrazone

Nomine, G. *et al.*, *J. Chem. Educ.*, 1969, **46**, 329

A reaction mixture in aqueous methanol exploded violently at –65°C. (Hydrazine–perchloryl fluoride redox reaction?) A previous reaction at 20°C had been uneventful, and the low-temperature explosion could not be reproduced.

Laboratory materials

Schlosser, M. *et al.*, *Chem. Ber.*, 1969, **102**, 1944

Contact of perchloryl fluoride with laboratory greases or rubber tubing, etc., has led to several explosions.

Lithiated compounds

Houben-Weyl, 1970, Vol. 13.1, 223

There is a danger of explosion during replacement of a lithium substituent by fluorine using perchloryl fluoride.

See Benzocyclobutene, etc., above

2-Lithio(dimethylaminomethyl)ferrocene

Peet, J. H. J. *et al.*, *J. Organomet. Chem.*, 1974, **82**, C57

Preparation of dimethylaminomethyl-2-fluoriferrocene, by interaction of the lithio compound in THF at –70°C with perchloryl fluoride diluted with helium, unexpectedly exploded violently.

Methyl 2-bromo-5,5-(ethylenedioxy)[2.2.1]bicycloheptane-7-carboxylate

Wang, C.-L. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1976, 468

In an attempt to improve isomer distribution from the fluorination reaction, one run was cooled to below –40°C, but on quenching with water, a violent explosion occurred.

Nitrogenous bases

1. Scott, F. L. *et al.*, *Chem. & Ind.*, 1960, 258

2. Gardner, D. M. *et al.*, *J. Org. Chem.*, 1967, **32**, 1115

3. Parish, E. J. *et al.*, *J. Org. Chem.*, 1980, **45**, 4036–4037

Interaction, in presence of diluent below 0°C, with isopropylamine or isobutylamine caused separation of explosive liquids, and with aniline, phenylhydrazine and 1,2-diphenylhydrazine, explosive solids [1]. In absence of diluents, contact with most aliphatic or non-aromatic heterocyclic amines often leads to uncontrolled oxidation and/or explosions [2]. During oxidation of two steroidal dienes in dry pyridine at –35 to –40°C, on one occasion each of the reactions was accompanied by violent explosions [3].

See 3 α -Hydroxy-5 β -androstane-11,17-dione 17-hydrazone, above

Podophyllotoxin enolate tetrahydropyranyl ether

Glinski, M. B. *et al.*, *J. Org. Chem.*, 1987, **53**, 2749–2753

Attempted preparation of 2-fluoropodophyllotoxin by treatment of the tetrahydropyranyl ether with perchloryl fluoride led to a violent explosion.

Sodium methoxide, Methanol

Papesch, V., *Chem. Eng. News*, 1959, **36**, 60

Addition of solid methoxide to a reaction vessel containing methanol vapour and gaseous perchloryl fluoride caused ignition and explosion. This could be avoided by adding all the methoxide first, or by nitrogen purging before addition of methoxide.

See other HALOGEN OXIDES, PERCHLORYL COMPOUNDS

3975. Chlorine fluorosulfate

[13997-90-5]



ClFO_3S

Bailar, 1973, Vol. 2, 1469

Hydrolysis is violent, producing oxygen.

See related ACYL HALIDES, HYPOHALITES

3976. Fluorine perchlorate (Perchloryl hypofluorite)

[10049-03-3]



ClFO_4

Alone, or Laboratory materials, or Potassium iodide

1. Rohrback, G. H. *et al.*, *J. Amer. Chem. Soc.*, 1949, **69**, 677

2. Christe, K. O. *et al.*, *Inorg. Chem.*, 1982, **21**, 2938

The pure liquid explodes on freezing at -167°C , and the gas is readily initiated by sparks, flame or contact with grease, dust or rubber tubing. Contact of the gas with aqueous potassium iodide also caused an explosion [1]. The pure material is now readily accessible via the relatively stable intermediate, tetrafluoroammonium perchlorate [2].

Difluorimide–potassium fluoride adduct

Christe, K. O. *et al.*, *Inorg. Chem.*, 1987, 56, 925

A mixture prepared at -196°C exploded when warmed towards -78°C .

Bis(trifluoromethanesulfonyl)imide

See N-Chlorobis(trifluoromethanesulfonyl)imide

Hydrogen

Hoffman, C. J., *Chem. Rev.*, 1964, **64**, 97

Ignition of the perchlorate occurs in excess hydrogen gas.

See other HALOGEN OXIDES, HYPOHALITES

- 3977. Xenon(II) fluoride perchlorate**
 [25582-86-9] **CIFO₄Xe**
 XeFCIO_4
- Bartlett, N. *et al.*, *Chem. Comm.*, 1969, 703
 Thermodynamically unstable, it explodes readily and sometimes with violence.
See other NON-METAL PERCHLORATES, XENON COMPOUNDS
- 3978. Nitrogen chloride difluoride**
 [13637-87-1] **CIF₂N**
 CINF_2
- Petry, R. C., *J. Amer. Chem. Soc.*, 1960, **82**, 2401
 Caution in handling is recommended for this *N*-halogen compound.
See other *N*-HALOGEN COMPOUNDS
- 3979. Phosphorus chloride difluoride**
 [14335-40-1] **CIF₂P**
 CIPF_2
- Hexafluoroisopropylideneaminolithium
See Hexafluoroisopropylideneaminolithium: Non-metal halides
See other NON-METAL HALIDES
- 3980. Thiophosphoryl chloride difluoride**
 [2524-02-9] **CIF₂PS**
 CIP(S)F_2
- Mellor, 1971, Vol. 8, Suppl. 3, 536
 Mixtures with air explode spontaneously at certain concentrations.
See other AUTOIGNITION INCIDENTS *See related* NON-METAL HALIDES (AND THEIR OXIDES)
- 3981. Chlorine trifluoride**
 [7790-91-2] **CIF₃**
 CIF_3
1. Anon., *J. Chem. Educ.*, 1967, **44**, A1057–1062
 2. O'Connor, D. J. *et al.*, *Chem. & Ind.*, 1957, 1155
 3. Farrar, R. L. *et al.*, *Some Considerations in Handling Fluorine and Chlorine Fluorides*, Oak Ridge Rept. K/ET-252, 1979, 41pp
- Handling procedures for this highly reactive oxidant gas have been detailed [1].
 Surplus gas is best burned with town- or natural gas, followed by absorption in alkali [2]. More recent and comprehensive information is available [3].

Acids

MRH Sulfuric acid 1.17/34

Mellor, 1956, Vol. 2, Suppl. 1, 157

Strong nitric and sulfuric acids reacted violently, (possibly owing to water content).

Ammonium fluoride, or Ammonium hydrogen fluoride

Gardner, D. M. *et al.*, *Inorg. Chem.*, 1963, **2**, 413

The reaction gases (containing chlorodifluoramine) must be handled at below -5°C to avoid explosion. Ammonium hydrogen fluoride behaves similarly.

Bis(trifluoromethyl) sulfide, or disulfide

See Bis(trifluoromethyl) sulfide: Chlorine fluorides

Bis(trifluoromethyl) disulfide: Chlorine fluorides

Boron-containing materials

Bryant, J. T. *et al.*, *J. Spacecr. Rockets*, 1971, **8**, 192–193

Finely divided boron, tetraboron carbide, and boron–aluminium mixtures will ignite on exposure to the gas.

See Metals, etc., below

Carbon tetrachloride

MRH 1.21/56

Mellor, Vol. 2, Suppl. 1, 156

Chlorine trifluoride will dissolve in carbon tetrachloride at low temperatures without reaction. Such solutions are dangerous, being capable of detonation. If it is used as a solvent for fluorination with the trifluoride, it is therefore important to prevent build-up of high concentrations of the latter.

Chromium trioxide

Mellor, 1943, Vol. 11, 181

Interaction of the two oxidants is incandescent.

Deuterium, or Hydrogen

MRH Hydrogen 6.82/3.0

Haberland, H. *et al.*, *Chem. Phys.*, 1975, **10**, 36

Studies of the interaction of chlorine trifluoride with deuterium or hydrogen atoms in a scattering chamber were accompanied by frequent flashes or explosions, within 3 h if the reactor had been vented, or after 8 h if it had not.

Fluorinated solvents

'Chlorine Trifluoride', Tech. Bull. TA 8522-3, Morristown, N. J., Baker & Adamson Div. of Allied Chemicals Corp., 1968

Bulk surfaces of polytetrafluoroethylene or polychlorotrifluoroethylene are resistant to the liquid or vapour under static conditions, but breakdown and ignition may occur under flow conditions.

See Polychlorotrifluoroethylene, below

Fuels

See ROCKET PROPELLANTS

Halocarbons

Brower, K. R., *J. Fluorine Chem.*, 1986, **31**, 333–349

Combinations of liquid chlorine trifluoride with several halocarbons except perfluorohexane exploded immediately when suddenly mixed at all temperatures between 25° and –70°C. Poly(chlorotrifluoroethylene), below

See Carbon tetrachloride, Fluorinated solvents, both above

Hydrocarbons

Brower, K. R., *J. Fluorine Chem.*, 1986, **31**, 333–349

Combinations of liquid chlorine trifluoride and several hydrocarbons exploded immediately when suddenly mixed at all temperatures between 25° and –70°C.

See Organic materials, below

Hydrogen-containing materials

MRH Ammonia 5.52/16

Hydrogen sulfide

MRH 4.68/13

Mellor, 1956, Vol. 2, Suppl. 1, 157

Explosive reactions occur with ammonia, coal-gas, hydrogen or hydrogen sulfide.

Ice

See Water, below

Iodine

Mellor, 1956, Vol. 2, Suppl. 1, 157

Ignition on contact.

Metals, or Metal oxides, or Metal salts, or Non-metals, or Non-metal oxides

MRH Iron 5.10/68, potassium 2.90/63, Calcium oxide 4.05/48, magnesium oxide 3.89/39, manganese dioxide 1.80/48, Calcium oxide 4.05/48, magnesium oxide 3.89/39, manganese dioxide 1.80/48, Sulfur 4.10/15, silver nitrate 0.5/73

1. Mellor, 1956, Vol. 2, Suppl. 1, 155–157

2. Sidgwick, 1950, 1156

3. Rhein, R. A., *J. Spacecr. Rockets*, 1969, **6**, 1328–1329

4. Chlorine Trifluoride' Tech. Bull. TA 8532-3, Morristown, N. J., Baker & Adamson Div. Allied Chem. Corp., 1968

Chlorine trifluoride is a hypergolic oxidiser with recognised fuels, and contact with the materials following at ambient or slightly elevated temperatures is violent, ignition often occurring. The state of subdivision may affect the results.

Antimony, arsenic, selenium, tellurium, iridium, iron, molybdenum, osmium, potassium, rhodium, tungsten; (and when primed with charcoal,) aluminium, copper, lead, magnesium, silver, tin, zinc. Interaction of lithium or calcium with chlorine tri- or penta-fluorides is hypergolic and particularly energetic.

Aluminium oxide, arsenic trioxide, bismuth trioxide, calcium oxide, chromic oxide, lanthanum oxide, lead dioxide, magnesium oxide, manganese dioxide, molybdenum trioxide, phosphorus pentoxide, stannic oxide, sulfur dioxide (explodes), tantalum pentoxide, tungsten trioxide, vanadium pentoxide.

Red phosphorus, sulfur; but with carbon, the observed ignition has been attributed to presence of impurities; mercury iodide, potassium iodide, silver nitrate, potassium carbonate.

Methane

Baddiel, C. B. *et al.*, *Proc. 8th Combust. Symp.*, 1960, 1089–1095

The explosive interaction of chlorine trifluoride with methane and its homologues has been studied in detail.

Nitroaryl compounds

Mellor, 1956, Vol. 2, Suppl. 1, 156

Several nitro compounds are soluble in chlorine trifluoride, but the solutions are extremely shock-sensitive. These include trinitrotoluene, hexanitrobiphenyl, hexanitrodiphenyl-amine, -sulfide or -ether. Highly chlorinated compounds behave similarly.

Organic materials MRH Acetic acid 3.05/19, benzene 5.35/36, ether 5.56/28

Mellor, 1956, Vol. 2, Suppl. 1, 155

Violence of the reaction, sometimes explosive, with e.g., acetic acid, benzene, ether, is associated with both their carbon and hydrogen contents. If nitrogen is also present, explosive fluoroamino compounds may be involved. Fibrous materials—cotton, paper, wood—invariably ignite.

Other reactants

Yoshida, 1980, 129

MRH values calculated for 18 combinations with oxidisable materials are given.

Polychlorotrifluoroethylene

Anon., *Chem. Eng. News*, 1965, **43**(20), 41

An explosion occurred when chlorine trifluoride was being bubbled through the fluorocarbon oil at -4°C . Moisture (snow) may have fallen into the mixture, reacted exothermally with the trifluoride and initiated the mixture.

See Water, below

Refractory materials

1. Cloyd, 1965, 58

2. Mellor, 1956, Vol. 2, Suppl. 1, 157

Fibrous or finely divided refractory materials, asbestos, glass wool, sand, or tungsten carbide, may ignite with the liquid and continue to burn in the gas [1]. The presence of adsorbed or lattice water seems necessary for attack on the siliceous materials to occur [2].

See other GLASS INCIDENTS

Ruthenium

Burns, R. C. *et al.*, *J. Inorg. Nucl. Chem.*, 1980, **42**, 1614

Formation of the adduct chlorine trifluoride–ruthenium pentafluoride by reaction with ruthenium metal at ambient temperature is extremely violent.

Selenium tetrafluoride

See Selenium tetrafluoride: Chlorine trifluoride

Trifluoromethanesulfonyl chloride

See Trifluoromethanesulfonyl chloride: Chlorine fluorides

Water

1. Sidgwick, 1950, 1156
2. Mellor, 1956, Vol. 2, Suppl. 1, 156, 158
3. Ruff, O. *et al.*, *Z. Anorg. Chem.*, 1930, **190**, 270
4. Farrar, R. L. *et al.*, USDOE Rept. K/ET-252, 1979, 41

Interaction is violent and may be explosive, even with ice, oxygen being evolved [1]. Part of the water dropped into a flask of the gas was expelled by the violent reaction ensuing [2]. An analytical procedure, involving absorption of chlorine trifluoride into 10% sodium hydroxide solution from the open capillary neck of a quartz ampoule to avoid explosion, was described [3]. Inadvertent collection of chlorine trifluoride and ice in a cryogenic trap led to a small but violent explosion when the trap began to warm up overnight [4].

See other INTERHALOGENS

3982. Chlorine trifluoride oxide

[30708-80-6]

ClF₃O

ClF₃O

Bougon, R. *et al.*, Fr. Pat. 2 110 555, 1972

It is a powerful oxidant, potentially useful in rocketry.

See other HALOGEN OXIDES

3983. Chlorine dioxygen trifluoride

[12133-60-7]

Cl(O₂)F₃

ClF₃O₂

Organic materials

1. Streng, A. G., *Chem. Rev.*, 1963, **63**, 607
2. Christe, K. O. *et al.*, *Inorg. Chem.*, 1973, **12**, 1357

It is a very powerful oxidant, but of low stability, which reacts explosively with organic materials [1]; such combinations should be avoided [2].

3984. Potassium tetrafluorochlorate(1-)

[19195-69-8]

K[ClF₄]

ClF₄K

See entry METAL POLYHALOHALOGENATES

3985. Nitrosyl tetrafluorochlorate
[13815-21-9] **ClF₄NO**



Organic compounds

Sorbe, 1968, 133

The powerful oxidant reacts explosively with many organic compounds.

See other NITROSO COMPOUNDS, OXIDANTS

3986. Tetrafluoroammonium perchlorate
[13706-14-4] **ClF₄NO₄**



Christe, K. O. *et al.*, *Inorg. Chem.*, 1980, **19**, 1495

It decomposes rapidly at 25°C, giving the very shock-sensitive chloryl hypofluorite.

See other N-HALOGEN COMPOUNDS, PERCHLORATE SALTS OF NITROGENOUS BASES

3987. Chlorotetrafluorophosphorane
[13637-88-2] **ClF₄P**



Preparative hazard

See Chlorine fluoride: Phosphorus trifluoride

See other NON-METAL HALIDES

3988. Rubidium tetrafluorochlorate(1-)
[15321-10-5] **ClF₄Rb**



See entry METAL POLYHALOHALOGENATES

3989. Chlorine pentafluoride
[13637-63-3] **ClF₅**



Metals

See Chlorine trifluoride: Metals

Nitric acid

Christe, K. O., *Inorg. Chem.*, 1972, **11**, 1220

Interaction of anhydrous nitric acid with chlorine pentafluoride vapour at -40°C, or with the liquid at above -100°C, is very vigorous.

Water

1. Pilipovich, D. *et al.*, *Inorg. Chem.*, 1967, **6**, 1918

2. Christe, K. O., *Inorg. Chem.*, 1972, **11**, 1220

Interaction of liquid chlorine pentafluoride with ice at -100°C [1], or of the vapour with water vapour above 0°C [2], is extremely vigorous.

See other INTERHALOGENS

3990. Pentafluorosulfur peroxyhypochlorite

[58249-49-3]



$\text{ClF}_5\text{O}_2\text{S}$

Haloalkenes


Hopkinson, M. J. *et al.*, *J. Org. Chem.*, 1976, **41**, 1408

The peroxyhypochlorite is especially reactive, and the fluoroperoxy compounds produced by its interaction with haloalkenes can detonate when subjected to thermal or mechanical shock. However, no explosions were experienced during this work.

See related HYPOHALITES, FLUORINATED PEROXIDES AND SALTS

3991. Difluoroperchloryl hexafluoroplatinate

[36609-92-4]



$\text{ClF}_8\text{O}_2\text{Pt}$

See entry DIFLUOROPERCHLORYL SALTS

3992. Chlorogermane

[13637-65-5]



ClGeH_3

Ammonia

Johnson, O. H., *Chem. Rev.*, 1951, **48**, 274

Both mono- and di-chlorogermanes react with ammonia to give involatile products (presumably with N-Ge bonds) which explode on heating.

See related METAL HALIDES, METAL HYDRIDES

See other AMINATION INCIDENTS

3993. Hydrogen chloride (Hydrochloric acid)

[7647-01-0]



ClH

FPA H41, 1975 (gas); *HCS 1980*, 546 (cylinder); *HCS 1980*, 541 (solution)

Preparative hazard

See Sulfuric acid, below

Alcohols, Hydrogen cyanide

1. Pinner, A., *Die Imidoäther*, Berlin, Openheim, 1892
2. Erickson, J. G., *J. Org. Chem.*, 1955, **20**, 1573

Preparation of alkyliminioformate chlorides (imidoester hydrochlorides) by passing hydrogen chloride rapidly into alcoholic hydrogen cyanide proceeds explosively (probably owing to a rapid exotherm), even with strong cooling [1]. Alternative procedures involving very slow addition of hydrogen chloride into a well-stirred mixture kept cooled to ambient temperature, or rapid addition of cold alcoholic hydrogen cyanide to cold alcoholic hydrogen chloride, are free of this hazard [2].

Aluminium

MRH 3.05/20

See Aluminium: Hydrogen chloride

Aluminium–titanium alloys: Oxidants

Caesium telluroacylates

Kato, S. *et al.*, *J. Amer. Chem. Soc.*, 1995, **116**(12), 1262

While generating tellurobenzoic acids by acidifying the corresponding cesium salts with hydrogen chloride in tubes immersed in liquid nitrogen, it was found that explosions sometimes occurred while later allowing the reaction tube to warm. This is due to overcooling; HCl freezes, and therefore becomes unreactive, well above liquid nitrogen temperature, permitting sudden and exothermic reaction during the warming period.

Chlorine, Dinitroanilines

1. Sihlbohm, L., *Acta Chem. Scand.*, 1953, **7**, 1197–1206
2. Harris, G. P. F. *et al.*, *Chem. & Ind.*, 1983, 183

The previously reported cleavage of dinitroanilines to chloronitrodiazonium salts by hydrochloric acid [1] is apparently catalysed by chlorine or other oxidants. The reaction takes place vigorously with copious gas evolution, but at low temperatures (40–25°C or lower) there can be a very long induction period before onset of the vigorous reaction. This could be hazardous on attempting to scale-up laboratory processes. The reactivity of the isomeric dinitroanilines varies, 2,3- being most reactive and 3,5- being unreactive. The cleavage reaction is specific for conc. hydrochloric acid, and has not been observed with hydrobromic acid or 30% sulfuric acid [2].

See other GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS

1,1-Difluoroethylene

See 1,1-Difluoroethylene: Hydrogen chloride

Fluorine

MRH 3.18/35

See Fluorine: Hydrogen halides

Hexalithium disilicide

See Hexalithium disilicide: Acids

Metal acetylides or carbides

Calcium carbide 3.97/46

See Caesium acetylide: Mineral acids

Rubidium acetylide: Acids

Uranium dicarbide: Hydrogen chloride

Metals

Anon., *Loss Prev. Bull.*, 1994, (116), 22

Hydrochloric acid is capable of evolving hydrogen on reaction with many common metals. A magnetic flow meter is reported as exploding after the acid reached the electric amplifier compartment.

Other reactants

Yoshida, 1980, 55

MRH values calculated for 11 combinations, largely with oxidants, are given.

Potassium permanganate

MRH 0.75/37

See Potassium permanganate: Hydrochloric acid

Silicon dioxide

See Silicon dioxide: Hydrochloric acid

Sodium

See Sodium: Acids

Sulfuric acid.

1. *MCA Case History No. 1785*

2. Libman, D. D., *Chem. & Ind.*, 1948, 728

3. Smith, G. B. L. *et al.*, *Inorg. Synth.*, 1950, **3**, 132

Accidental addition of 6500 l of conc. hydrochloric acid to a bulk sulfuric acid storage tank released sufficient hydrogen chloride gas by dehydration to cause the tank to burst violently [1]. Complete dehydration of hydrochloric acid solution releases some 250 volumes of gas. A laboratory apparatus for effecting this safely has been described [2], which avoids the possibility of layer formation in unstirred flask generators [3].

See other GAS EVOLUTION INCIDENTS

Tetraselenium tetranitride

See Tetraselenium tetranitride: Alone, or Halogens and derivatives

See other INORGANIC ACIDS, NON-METAL HALIDES, NON-METAL HYDRIDES

3994. Monopotassium perchlorylamide

[16971-92-9]

KNHClO₃

ClHKNO₃

Sorbe, 1968, 67

Like the dipotassium salt, it will explode on impact or exposure to flame.
See entry PERCHLORYLAMIDE SALTS *See other* N-METAL DERIVATIVES

3995. Hypochlorous acid
[7790-92-3]

ClHO

HOCl

Acetic anhydride

Rüst, 1948, 341

A mixture of the anhydride and hypochlorous acid exploded violently while being poured. Some acetyl hypochlorite and/or dichlorine monoxide may have been formed.

Alcohols

Mellor, 1956, Vol. 2, Suppl. 1, 560

Contact of these, or of chlorine and alcohols, readily forms unstable alkyl hypochlorites.

See HYPOHALITES

Ammonia

Mellor, 1940, Vol. 8, 217

The violent explosion occurring on contact with ammonia gas is due to formation of nitrogen trichloride and its probable initiation by the heat of solution of ammonia.

See Chloramine (reference 3)

Arsenic

Mellor, 1941, Vol. 2, 254

Ignition occurs on contact.

See other HYPOHALITES, OXOHALOGEN ACIDS

3996. Chloric acid
[7790-93-4]

ClHO₃

HOClO₂

Muir, G. D., private comm., 1968

Aqueous chloric acid solutions decompose explosively if evaporative concentration is carried too far.

Cellulose

Mellor, 1946, Vol. 2, 310

Filter paper ignites after soaking in chloric acid.

Metal sulfides

Mellor, 1956, Vol. 2, Suppl. 1, 584

Copper sulfide explodes with concentrated chloric acid solution (and also with cadmium, magnesium or zinc chlorates). Antimony trisulfide or arsenic trisulfide, or tin(II) and (IV) sulfides react incandescently with concentrated solutions.

Metals, or Organic materials

Majer, V., *Chemie* (Prague), 1948, **3**, 90–91

The recorded explosions of chloric acid have been attributed to the formation of explosive compounds with antimony, bismuth and iron (including hydrogen also in the latter case). Organic materials (and ammonia) are violently oxidised.

Oxidisable materials

In contact with oxidisable materials, reactions are similar to those of the metal chlorates.

See entry METAL CHLORATES *See other* OXOHALOGEN ACIDS

3997. Chlorosulfuric acid

[7790-94-5]

ClHO₃S

ClSO₂OH

FPA H122, 1983; *HCS 1980*, 314

Diphenyl ether

Ehama, T. *et al.*, *Jap. Pats.* 74 45 034, 74 45 035, 1974

Presence of various nitrogen-containing compounds, or of fatty acids or their derivatives, controls the vigorous interaction of the ether and chlorosulfuric acid at above 40°C, producing higher yields of 4,4'-oxybisbenzenesulfonyl chloride.

Hydrocarbons

MRH Hexane, 0.96/24; Toluene, 1.29/37

1. Anon., *CIHSC Chem. Safety Summ.*, 1986, **57**(226), 65

2. Baker, J. D., *Chem. Eng. News*, 1989, **67**(50), 2

Use of heptane to wash sludge from a chlorosulfuric acid weigh tank led to violent rupture of the 2.5 kl vessel. Laboratory work showed that the two compounds react vigorously with gas evolution (HCl) sufficient to overpressure the tank which was fitted with a 40 mm relief line [1]. Another(?) tank containing chlorosulfonic acid ruptured after the lines to it were washed out with heptane, subsequent investigation showed that chlorosulfonic acid gives reaction, with large gas evolution, when stirred with commercial heptane fractions (and with hexane, but not cyclohexane, in the editor's experience.) Subsequent correspondence sought to attribute the observation to water contamination, which seems unlikely [2].

See other GAS EVOLUTION INCIDENTS, SULFONATION INCIDENTS

Other reactants

Yoshida, 1980, 115–116

MRH values calculated for 21 combinations, mainly with oxidisable materials, are given.

Phosphorus

Heumann, K. *et al.*, *Ber.*, 1882, **15**, 417

The acid is a strong oxidant, and above 25–30°C, interaction is vigorous and accelerates to explosion.

Silver nitrate

See Silver nitrate: Chlorosulfuric acid

Sulfuric acid

Anon., *Loss Prev. Bull.*, 1977, (013), 2–3

Failure to start an agitator before charging 500 kg of 98% sulfuric acid ($d = 1.84$) into 560 kg of chlorosulfuric acid ($d = 1.44$) led to formation of a lower layer of sulfuric acid. Subsequent starting of the agitator led to eruption of part of the contents owing to rapid formation of hydrogen chloride gas from interaction of the chlorosulfuric acid with the water content (2%) in the sulfuric acid.

See other GAS EVOLUTION INCIDENTS

See other ACYL HALIDES, INORGANIC ACIDS, OXIDANTS

3998. Perchloric acid

[7601-90-3]

HOClO_3

ClHO_4

(MCA SD-11, 1965); NSC 311, 1982; FPA H53, 1977; HCS 1980, 721

1. MCA SD-11, 1965
2. Lazerte, D., *Chem. Eng. News*, 1971, **49**(3), 33
3. Muse, L. A., *J. Chem. Educ.*, 1972, **49**, A463
4. Graf, F. A., *Chem. Eng. Progr.*, 1966, **62**(10), 109
5. Friedmann, W., *Chem. Eng. News*, 1947, **25**, 3458
6. Anon., *ABCM Quart. Safety Summ.*, 1936, **7**, 51
7. Annikov, V. E. *et al.*, *Chem. Abs.*, 1983, **99**, 160860
8. Anon., *Indian Chem. Manuf.*, 1976, **14**(5), 9–13
9. Schilt, A. A., *Perchloric Acid and Perchlorates*, Columbus (Oh.), G. F. Smith Chemical Co., 1979
10. Furr, A. K. (Ed.), *Handbook of Laboratory Safety*. 3rd Edn., 207, CRC Press, Boca Raton (Florida), 1990.

Most of the numerous and frequent hazards experienced with perchloric acid have been associated with either its exceptional oxidising power or the inherent instability of its covalent compounds, some of which form readily. Although the 70–72% acid of commerce behaves when cold as a very strong, but non-oxidising acid, it becomes an extreme oxidant and powerful dehydrator at elevated temperatures (160°C) or when anhydrous [1].

Where an equally strong but non-oxidising acid can be used, trifluoromethanesulfonic acid is recommended [2]. Safe handling procedures have been detailed [3] and an account of safe handling of perchloric acid in large-scale preparation of hydrazinium diperchlorate, with recommendations for materials of construction, has been published [4]. The lack of oxidising power of the unheated 72% aqueous acid was confirmed during analytical treatment of oil sludges on over 100 occasions [5]. A report of two explosions during evaporation of aqueous perchloric acid in a method for estimating potassium infers that the aqueous acid may be unstable. It is more likely that oxidisable material (probably ethanol) was present during the

ClHO₄

evaporation [6]. The aqueous acid can be caused to detonate with a booster charge [7]. Safety procedures for using perchloric acid for various purposes have been reviewed [8], and the section on safety aspects in a text [9] is rated by a reviewer as particularly pertinent. Precautions for using perchloric acid and perchlorates are given, and many incidents described in [10].

Acetic anhydride

See Dehydrating agents, below (references 2–4)

Acetic anhydride, Acetic acid, Organic materials

MRH Acetic acid 4.98/35

1. Burton, H. *et al.*, *Analyst*, 1955, **80**, 4
2. Schumacher, 1960, 187, 193
3. Kuney, J. H., *Chem. Eng. News*, 1947, **25**, 1659
4. *Tech. Survey No. 2, Fire Hazards and Safeguards for Metalworking Industries*, US Board of Fire Underwriters, 1954

Mixtures of 70% (dihydrated) perchloric acid with enough acetic anhydride will produce a solution of anhydrous perchloric acid in acetic acid/anhydride, which is of high catastrophic potential [1]. Sensitivity to heat and shock depend on the composition of the mixture, and the vapour evolved on heating is flammable [2]. Such solutions have been used for electropolishing operations, and during modifications to an electropolishing process by the unqualified supervisor, a cellulose acetate rack was introduced into a large volume of an uncooled mixture of perchloric acid and acetic anhydride. Dissolution of the rack introduced organic materials (largely cellulose acetate, capable of transesterification to cellulose perchlorate) into the virtually anhydrous acid and caused the whole to explode disastrously, causing enormous damage. This cause was confirmed experimentally [3,4].

See Alcohols; Cellulose and derivatives; Vegetable matter, all below

See Acetic anhydride: Perchloric acid

Acetic anhydride, Carbon tetrachloride, 2-Methylcyclohexanone

Gall, M. *et al.*, *Org. Synth.*, 1972, **52**, 40

During acetylation of the enolised ketone, the 70% perchloric acid must be added last to the reaction mixture to provide maximum dilution and cooling effect.

Acetic anhydride, Organic materials, Transition metals

Hikita, T. *et al.*, *J. Chem. Soc. Japan, Ind. Chem. Sect.*, 1951, **54**, 253–255

The stability ranges of mixtures of the acid, anhydride and organic materials (ethanol, gelatine) used in electropolishing were studied. Presence of transition metals (chromium, iron, nickel) increases the possibility of explosion. (This is why such mixtures must not be stored after use for etching metals.)

Acetonitrile

Andrussow, L., *Chim. Ind. (Paris)*, 1961, **86**, 542–545

The latent hazards in storing and handling the explosive mixtures of perchloric acid with acetonitrile or dimethyl ether are discussed.

Acetylene, Nitrous oxide

1. Hobson, J. D., *Chem. Brit.*, 1985, **21**, 632
2. Anon., *Lab. Haz. Bull.*, 1987, (9), item 735
3. Andrew, B. E., *J. Anal. At. Spectrom.*, 1988, **3**, 401–405

A number of explosions have been experienced when using sample solutions containing perchloric acid in atomic absorption spectrometers using acetylene–nitrous oxide flames. Further information is being sought from other AAS users [1]. A warning against using perchloric acid digests in AA instruments using such flames has also been given by an American source [2]. The role of perchlorate accumulation and the effects of instrumental and experimental variables upon such explosions has been studied and preventive measures proposed [3].

See ATOMIC ABSORPTION SPECTROSCOPY

Alcohols

MRH Ethanol 6.27/21

1. Sweasey, D., *Lab. Practice*, 1968, **17**, 915
2. Moureu, H. *et al.*, *Arch. Mal. Prof. Med.*, 1951, **12**, 157–159
3. Michael, A. T. *et al.*, *Amer. Chem. J.*, 1900, **23**, 444
4. Bailar, 1973, Vol. 2, 1451

During digestion of a lipid extract (1 mg) with 72% acid (1.5 ml), a violent explosion occurred. This was attributed to residual traces of the extraction solvent (methanol–chloroform) having formed methyl perchlorate [1]. In the analytical determination of potassium as perchlorate, heating the solid containing traces of ethanol and perchloric acid caused a violent explosion [2]. Contact of drops of anhydrous perchloric acid and ethanol caused immediate violent explosion [3]. Partial esterification of polyfunctional alcohols (ethylene glycol, glycerol, pentaerythritol) with the anhydrous acid gives extremely explosive liquids which may explode on pouring from one vessel to another [4].

See Glycols and their ethers; and Methanol, Triglycerides; both below

See ALKYL PERCHLORATES

Aniline, Formaldehyde

MRH Aniline 6.65/18

Aniline, 86, New York, Allied Chem. Corp., 1964

Aniline reacts with perchloric acid and then formaldehyde to give an explosively combustible condensed resin.

Antimony(III) compounds

Burton, H. *et al.*, *Analyst*, 1955, **80**, 4

Treatment of trivalent compounds of antimony with perchloric acid can be very hazardous.

See Bismuth, below

Azo-pigment, Orthoperiodic acid

1. Anon., *ABCM Quart. Safety Summ.*, 1961, **32**, 125
2. Smith, G. F. *et al.*, *Talanta*, 1960, **4**, 185

During the later stages of the wet oxidation of an azo-pigment with mixed perchloric and orthoperiodic acids, a violent reaction, accompanied by flashes of

ClHO₄

light, set in and terminated in an explosion [1]. The general method upon which this oxidation was based is described as hazard-free [2].

Bis(1,2-diaminopropane)-*cis*-dichlorochromium(III) perchlorate

See Bis(1,2-diaminopropane)-*cis*-dichlorochromium(III) perchlorate: Perchloric acid

1,3-Di[bis(cyclopentadienyl)iron]-2-propene-1-one

See 1,3-Di[bis(cyclopentadienyl)iron]-2-propene-1-one: Perchloric acid

Bis(2-hydroxyethyl) terephthalate, Ethanol, Ethylene glycol

491M, 1975, 303

When a solution of the ester in ethanol containing 5% of perchloric acid was evaporated to dryness, there was a violent explosion (probably owing to formation of ethyl perchlorate, perhaps with some ester perchlorate or ethylene glycol perchlorate also being formed). A similar mixture containing ethylene glycol flashed brightly after refluxing for 18 h.

See Alcohols, above; Glycols and their ethers, below

Bismuth

Nicholson, D. G. *et al.*, *J. Amer. Chem. Soc.*, 1935, **57**, 817

Attempts to dissolve bismuth and its alloys in hot perchloric acid carry a very high risk of explosion. At 110°C a dark brown coating is formed, and if left in contact with the acid (hot or cold), explosion occurs sooner or later. The same is true of antimony and its trivalent compounds.

Calcium compounds

Akerman, L. A. *et al.*, *Combustion, Explos., & Shockwaves*, 1987, **23**, 178

Calcium chloride and oxide catalyse the second, violent, stage of ammonium perchlorate decomposition and increase the shock sensitivity of mixtures with sugar.

Carbon

MRH 6.19/17

1. Mellor, 1941, Vol. 2, 380

2. Pascal, 1960, Vol. 16, 300–301

Contact of a drop of the anhydrous acid with wood charcoal causes a very violent explosion [1], and carbon black also reacts violently [2].

See Graphitic carbon, Nitric acid, below

Cellulose and derivatives

MRH Cellulose 5.81/33

1. Schumacher, 1960, 187, 195

2. Harris, E. M., *Chem. Eng.*, 1949, **56**, 116

3. Sutcliffe, *J. Textile Ind.*, 1950, **41**, 196T

Contact of the hot concentrated acid or the cold anhydrous acid with cellulose (as paper, wood fibre or sawdust, etc.) is very dangerous and may cause a violent explosion. Many fires have been caused by long-term contact of diluted acid with wood (especially in fume cupboards) with subsequent evaporation and ignition

[1,2]. Contact of cellulose acetate with 1200 l of uncooled anhydrous acid in acetic anhydride led to a particularly disastrous explosion [1], and interaction of benzyl cellulose with boiling 72% acid was also explosive [3]. Perchlorate esters of cellulose may have been involved in all these incidents.

See Hydrofluoric acid, Structural materials, below

Charcoal, Chromium trioxide

Randall, W. R., private comm., 1977

A wet-ashing technique used for dissolution of graphite in perchloric acid involved boiling a mixture of 70% perchloric acid and 1% of chromium trioxide as an aqueous solution. This method was later applied to 6–14 mesh charcoal, and after boiling for 30 min the reaction rate increased (foaming) and accelerated to explosion. The charcoal contained traces of extractable tar.

Clay

Atkinson, G. F., *Chem. Eng. News*, 1987, **65**(39), 2

Digestion of clay samples with perchloric acid led to 3 explosions, each of which destroyed a ceramic hotplate used as heat source. There seems a good case for using metal hotplates for reactions involving risk of explosion.

See Hydrogen peroxide: Acetic acid, Jute

Combustible materials

Elliot, M. A. *et al.*, *Rept. Invest. No. 4169*, Washington, US Bur. Mines, 1948

Tests of sensitivity to initiation by heat, impact, shock or ignition sources were made on mixtures of a variety of absorbent materials containing a stoichiometric amount of 40–70% perchloric acid. Wood meal with 70% acid ignited at 155°C and a mixture of coal and 60% acid which did not ignite below 200°C ignited at 90°C when metallic iron was added. Many of the mixtures were more sensitive and dangerous than common explosives.

Copper dichromium tetraoxide

Solymosi, F. *et al.*, *Proc. 14th Combust. Symp.*, 1309–1316, 1973

The mixed oxide (copper chromite) was the most effective of several catalysts for the vapour-phase decomposition of perchloric acid, decomposition occurring above 120°C.

Dehydrating agents

1. Mellor, 1941, Vol. 2, 373, 380; 1956, Vol. 2, Suppl. 2.1, 598, 603
2. Kuney, J. H., *Chem. Eng. News*, 1947, **25**, 1659
3. Burton, H. *et al.*, *Analyst*, 1955, **80**, 4
4. Schumacher, 1960, 71, 187, 193
5. Wirth, C. M. P., *Lab. Practice*, 1966, **15**, 675
6. Plesch, P. H. *et al.*, *Chem. & Ind.*, 1971, 1043
7. Musso, H. *et al.*, *Angew. Chem.*, 1970, **82**, 46

Although commercial 70–72% perchloric acid (approximating to the dihydrate) itself is stable, incapable of detonation (except by a booster charge) and readily

ClHO₄

stored, it may be fairly readily dehydrated to the anhydrous acid by contact with drying agents. This is not safe when stored at ambient temperature, since it slowly decomposes, even in the dark, with accumulation of chlorine dioxide in the solution, which darkens and finally explodes after some 30 days [1,2,5]. The 72% acid (or perchlorate salts) may be converted to the anhydrous acid by heating with sulfuric acid, phosphorus pentoxide or phosphoric acid, or by distillation under reduced pressure [1,4,5]. In contact with cold acetic anhydride, mixtures of the anhydrous acid with excess anhydride and acetic acid are formed, which are particularly dangerous, being sensitive to mechanical shock, heating, or the introduction of organic contaminants [2,3,4]. A solution of the monohydrated acid in chloroform exploded in contact with phosphorus pentoxide [1]. A safer method of preparing anhydrous solutions of perchloric acid in dichloromethane, which largely avoids the risk of explosion, has been described [6]. Further precautions are detailed in an account of an explosion during a similar preparation [7]. Solutions of anhydrous acid of less than 5% concentration in acetic acid or anhydride are relatively stable.

Deoxyribonucleic acid

Cochrane, A. R. G. *et al.*, *School Sci. Rev.*, 1977, **58**, 706–708

Hazards of using perchloric acid to hydrolyse DNA are stressed. Perchloric acid can cause ignition of any organic material, even a considerable time after contact. Other acids to effect hydrolysis are suggested.

Dichloromethane, Dimethyl sulfoxide

MRH Dimethyl sulfoxide 6.19/23

Anon., *Univ. Safety Assoc. Safety News*, 1978, (9), 10

When a syringe used for DMSO and rinsed with dichloromethane was being filled with perchloric acid, a violent explosion resulted.

See Sulfoxides, below

Diethyl ether

MRH 6.61/18

Michael, A. T. *et al.*, *Amer. Chem. J.*, 1900, **23**, 444

The explosions sometimes observed on contact of the anhydrous acid with ether are probably owing to formation of ethyl perchlorate by scission of the ether, (or possibly to formation of diethyloxonium perchlorate).

Dimethyl ether

See Acetonitrile, above

Dioxane, Nitric acid

MCA Guide, 1972, 312–313

Boron trifluoride in aqueous dioxane was being evaporated with nitric acid treatment (3 portions) for analysis. Final addition of perchloric acid with continued heating led to an explosion while unattended. Ring scission of dioxane to diethylene glycol and formation of diethylene glycol nitrate and/or perchlorate may have been involved.

See Nitric acid, Organic matter, below

Ethylbenzene, Thallium triacetate

Uemura, S. *et al.*, *Bull. Chem. Soc., Japan.*, 1971, **44**, 2571

Application of a published method of thallation to ethylbenzene caused a violent explosion. A reaction mixture of thallium triacetate, acetic acid, perchloric acid and ethylbenzene was stirred at 65°C for 5 h, then filtered from thallos salts. Vacuum evaporation of the filtrate at 60°C gave a pasty residue which exploded. This preparation of ethylphenylthallic acetate perchlorate monohydrate had been done twice previously and uneventfully, as had been analogous preparations involving thallation of benzene, toluene, three isomeric xylenes and anisole in a total of 150 runs, where excessive evaporation had been avoided.

Faecal material, Nitric acid

Anon., *Univ. Safety Assoc. Safety News*, 1979, (12), 24

In the determination of metal ions in animal faeces, digestion with 12% perchloric and 56% nitric acids was in progress when an explosion occurred. This was attributed to one of the samples going to dryness on a sand tray heater.

See Nitric acid, Organic matter, below

Fluorine

Rohrbock, G. H. *et al.*, *J. Amer. Chem. Soc.*, 1947, **69**, 677–678

Contact of fluorine and 72% acid at ambient temperature produces a high yield of the explosive gas, fluorine perchlorate.

Fume cupboards

Furr, A. K. (Ed.), *Handbook of Laboratory Safety*. 3rd Edn., 207, CRC Press, Boca Raton (Florida), 1990.

Fume cupboards or their vent lines which have frequently been used with perchloric acid have often spontaneously deflagrated or exploded because of spillage or absorbed vapour.

See Hydrofluoric acid, Structural materials, below; Cellulose and derivatives, above

Glycerol, Lead oxide

MCA Case History No. 799

During maintenance work on casings of fans used to extract perchloric acid fumes, seven violent explosions occurred when flanges sealed with lead oxide–glycerol cement were disturbed. The explosions, attributed to formation of explosive compounds by interaction of the cement with perchloric acid, may have involved perchlorate esters and/or lead salts. Use of an alternative inorganic silicate–hexafluorosilicate cement is recommended.

Glycols and their ethers

MRH Ethylene glycol 5.69/30

1. Schumacher, 1960, 195, 214

2. Comas, S. M. *et al.*, *Metallography*, 1974, **7**, 45–47

Glycols and their ethers undergo violent decomposition in contact with ~70% perchloric acid. This seems likely to involve formation of the glycol perchlorate esters (after scission of ethers) which are explosive, those of ethylene glycol

and 3-chloro-1,2-propanediol being more powerful than glyceryl nitrate, and the former so sensitive that it explodes on addition of water [1]. Investigation of the hazards associated with use of 2-butoxyethanol for alloy electropolishing showed that mixtures with 50–95% of acid at 20°C, or 40–90% at 75°C, were explosive and initiatable by sparks. Sparking caused mixtures with 40–50% of acid to become explosive, but 30% solutions appeared safe under static conditions of temperature and concentration.

See Alcohols, above

Graphitic carbon, Nitric acid

Buzzelli, G. *et al.*, *Talanta*, 1977, **24**, 383–385

Dissolution procedures are described for gram samples of graphite or pyrolytic carbon, milligram samples of irradiated fuel particles, and for more readily oxidised forms of carbon, such as charcoal. The first two methods involve heating the samples with mixtures of 70% perchloric and 90% nitric acids (10:1), and must only be used for graphite or pyrolytic carbon. Other forms of carbon must not be oxidised in this way (to avoid explosions), but by a preliminary treatment with nitric acid alone and in portions.

See Carbon, above

Hydrofluoric acid, Structural materials

1. Anon., *Sichere Chemiearb.*, 1983, **35**, 85–86

2. Anon., *Sichere Chemiearb.*, 1984, **36**, 59

A violent explosion in a fume hood, in which inorganic siliceous materials had been digested with perchloric and hydrofluoric acids on a hotplate-heated sandbath during several years, originated from the tiled working surface. This consisted of ceramic tiles set in acid-proof organic resin-bond cement with mineral filler and supported on a 4 cm chipboard base. It appeared that prolonged use of the acids had led to development of hair cracks in the tiled surface, slow penetration of the acids into the underlying chipboard base and progressive formation of cellulose perchlorate and other perchlorate esters from the resin binder beneath the tiled surface. The explosion was probably initiated by impact or friction, perhaps from moving the hotplate on the tiled surface. The significance of such factors in preventing such accidents is discussed [1]. A new piece of equipment should prevent the occurrence of such accidents caused by long-term use of perchloric acid. This is designed to accommodate 8 separate digestion vessels (PTFE for up to 250°C, other material in development for higher temperatures) heated in an aluminium block. Each vessel is connected by a PTFE line into a gas manifold which leads to an alkali-containing absorption vessel to completely neutralise the gases or vapours evolved during acid digestion [2].

See Cellulose and derivatives, above

Hydrogen

1. Dietz, W., *Angew. Chem.*, 1939, **52**, 616

2. Schumacher, 1960, 189

Occasional explosions experienced during use of hot perchloric acid to dissolve steel samples for analysis [1] were attributed to formation of hydrogen–perchloric

acid vapour mixtures and their ignition by steel particles at temperatures as low as 215°C [2].

Hydrogen halides

See Fluoronium perchlorate, *also* Chloronium perchlorate

Iodides

Michael, A. T. *et al.*, *Amer. Chem. J.*, 1900, **23**, 444

The anhydrous acid ignites in contact with sodium iodide or hydriodic acid.

Iron(II) sulfate

Tod, H., private comm., 1968

During preparation of iron(II) perchlorate, a mixture of iron sulfate and perchloric acid was being strongly heated when a most violent explosion occurred. Heating should be gentle to avoid initiating this redox system.

See *other* REDOX REACTIONS

Ketones

MRH Acetone 6.48/21

Schumacher, 1960, 195

Ketones may undergo violent decomposition in contact with 70% acid.

Methanol, Triglycerides

1. Mavrikos, P. J. *et al.*, *J. Amer. Oil Chem. Soc.*, 1973, **50**(5), 174

2. Wharton, H. W., *J. Amer. Oil Chem. Soc.*, 1974, **51**(2), 35–36

Attention is drawn to the hazards involved in the use of perchloric acid in a published method [1] for transesterification of triglycerides with methanol. Alternative acid catalysts and safety precautions are suggested [2].

2-Methylpropene, Metal oxides

Lesnikovich, L. I. *et al.*, *Chem. Abs.*, 1975, **83**, 149760; 1976, 85, 7929

Mixtures of the alkene and perchloric acid vapour (5:1 molar) in nitrogen ignite spontaneously at 250°C. Some metal oxides of low specific surface reduced the ignition temperature below 178°C.

Nitric acid, Organic matter

1. Anon., *Ind. Eng. Chem. (News Ed.)*, 1937, **15**, 214

2. Lambie, D. A., *Chem. & Ind.*, 1962, 1421

3. Mercer, E. R., private comm., 1967

4. Muse, L. A., *Chem. Eng. News*, 1973, **51**(6), 29–30

5. Cooke, G. W., private comm., 1967

6. Kuney, J. H., *Chem. Eng. News*, 1947, **25**, 1659

7. Balks, R. *et al.*, *Chem. Abs.*, 1939, **33**,2438.9

8. Rooney, R. C., *Analyst*, 1975, **100**, 471–475

9. Martinie, G. D. *et al.*, *Anal. Chem.*, 1976, **48**, 70–74

10. *MCA Case History No. 2145*

11. Anon., *Univ. Safety Assoc. Safety News*, 1978, (9), 7

12. Satzger, R. D. *et al.*, *J. Assoc. Off. Anal. Chem.*, 1983, **66**, 985–988

13. Frank, A., *Chem. Abs.*, 1983, **99**, 197442

14. Anon., *Lab. Accid Higher Educ.*, Item 15, HSE, Barking, 1987

The mixed acids have been used to digest organic materials prior to analysis, but several explosions have been reported, including those with vegetable oil [1], milk [2], and calcium oxalate precipitates from plants [3]. To avoid trace metal contamination by homogenising rat carcasses in a blender, the whole carcasses were dissolved in nitric acid. After separation of fat and addition of perchloric acid (125 ml), evaporation to near-dryness caused a violent explosion [4]. Finely ground plant material in contact with perchloric/nitric acid mixture on a heated sand bath became hot before all the plant material was saturated with acid and it exploded. Subsequent digests left overnight in contact with cold acids proceeded smoothly [5]. Cellulose nitrate and/or perchlorate may have been involved. Treatment with nitric acid before adding perchloric acid was a previously used and well-tried safe procedure [6]. Application to animal tissues of a method previously found satisfactory for plant material caused a violent explosion [7].

A method which involves gradually increasing the liquid temperature by controlled distillation during the digestion of organic matter with the hot mixed acids had been described as safe provided that strict control of the distillation process were observed [4]. During oxidation of large, high-fat samples of animal matter by the usual technique, the existence of a layer of separated fat on the perchloric acid mixture created a highly hazardous situation. A modified procedure is free of explosion risk [8]. The effectiveness of mixed perchloric and nitric acids in wet oxidation of a wide range of organic materials has been studied. Violent oxidation occasionally occurred, but addition of sulfuric acid prevented any explosions or ignition during digestions [9]. The case history describes an explosion during wet-ashing of a phosphorus-containing polymer. One of the flasks may not have been topped up (i.e. diluted) with nitric acid during the digestion phase [10]. Digestion of vegetable matter with a 4:1 mixture of nitric/perchloric acids led to an explosion [11]. A safer alternative to the use of mixed acids for digestion of bone-meal for lead determination is pressurised dissolution in hydrochloric acid, then stripping voltammetry [12]. Equipment and procedures to minimise the risk of explosion during wet digestion of fat-containing materials are detailed [13]. An untried procedure to digest fresh cows milk with the mixed acids at 180°C led to a violent explosion [14].

Nitric acid, Pyridine, Sulfuric acid

Randall, W. L., *Safety in Handling Hazardous Chemicals*, 8–10, UCID-16610, Lawrence Livermore Lab., Univ. Calif., 1974

Some rare-earth fluoride samples had been wet-ashed incompletely with the three mixed acids and some gave low results. These samples, now containing pyridine, were reprocessed by addition of more acids and slow evaporation on a hot-plate. One of the samples frothed up and then exploded violently. Pyridinium perchlorate seems likely to have been involved.

Nitrogenous epoxides

Harrison, G. E., private comm., 1966

Traces of perchloric acid used as hydration catalyst for ring opening of nitrogenous epoxides caused precipitation of an organic perchlorate salt which was highly explosive. The concentration of acid was less than 1% by vol.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

Oleic acid

1. Swern, D. *et al.*, US Pat. 3 054 804; *Chem. Eng. News*, 1963, **41**(12), 39
2. Anon., *Chem. Eng. News*, 1963, **41**(27), 47

The improved preparation of 1,4-octadecanolactone [1] involves heating oleic acid (or other C₁₈ acids) with 70% perchloric acid to 115°C. This is considered to be a potentially dangerous method [2].

Organic materials, Sodium hydrogen carbonate

1. Zderic, J. A. *et al.*, *J. Amer. Chem. Soc.*, 1960, **82**, 446
2. Anon., *Univ. Safety Assoc. Safety News*, 1978, (9), 8–9

Following a published procedure [1], perchloric acid was used as catalyst in preparing a diol ketal, and was neutralised with sodium hydrogencarbonate before workup. Some sodium perchlorate remained dissolved after filtering the reaction mixture, and during subsequent vacuum distillation (with the bath temperature increasing to 200°C) an explosion occurred.

Organophosphorus compounds

Anderson, R. L. *et al.*, *Clin. Chim. Acta*, 1982, **121**, 111–116

The standard method for assaying organophosphorus compounds can be modified to use sulfuric acid to digest the samples and hydrogen peroxide as oxidant in place of perchloric acid.

Other reactants

Yoshida, 1980, 78

MRH values calculated for 19 combinations with oxidisable materials are given.

Phenylacetylene

1. Martheard, J.-P. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 3484
2. Martheard, J.-P. *et al.*, *J. Chem. Res., Synop.*, 1983, 224–225

A mixture of phenylacetylene and the pure (?anhydrous) acid prepared at –180°C exploded when allowed to warm to –78°C, possibly owing to formation of unstable 1-phenylethenyl perchlorate [1]. Reaction of various phenylacetylenes in acetic acid at 40°C is controllable [2].

Phosphine

See Phosphonium perchlorate

Pyridine

See Pyridinium perchlorate

Sodium phosphinate

Smith, G. F., *Analyst*, 1955, **80**, 16

MRH 3.68/63

ClHO₄

Although no interaction occurs in the cold, these powerful oxidising and reducing agents violently explode on heating.

See other REDOX REACTIONS

Sulfinyl chloride

Bailar, 1973, Vol. 2, 1442

The anhydrous acid ignites the chloride.

Sulfoxides

MRH Dimethyl sulfoxide 6.19/23

1. Therésa, J. de B., *Anales Soc. Espan. Fis. y Quim.*, 1949, **45B**, 235

2. Graf, F. A., *Chem. Eng. Progr.*, 1966, **62**(10), 109

3. Uemura, S. *et al.*, *Bull. Chem. Soc. Japan*, 1971, **44**, 2571

4. *471M*, 1975, 302

5. Anon., *Jahresber.*, 1990, 64

6. Eigenmann, K. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1975, **14**, 647–648

Lower members of the series of salts formed between organic sulfoxides and perchloric acid are unstable and explosive when dry. That from dibenzyl sulfoxide explodes at 125°C [1]. Dimethyl sulfoxide explodes on contact with 70% perchloric acid solution [2]; one drop of acid added to 10 ml of sulfoxide at 20°C caused a violent explosion [3], and dibutyl sulfoxide behaves similarly [4]. A fatal explosion resulted from mistakenly connecting a DMSO reservoir to an autopipette previously used with perchloric acid [5]. (The editor has met a procedure for methylthiolation of aromatics where DMSO was added to excess 70% perchloric acid; he did not feel justified in trying to scale it up.) Explosions reported seem usually to result from addition to excess sulfoxide. Aryl sulfoxides condense uneventfully with phenols in 70% perchloric acid, but application of these conditions to the alkyl sulfoxide (without addition of the essential phosphoryl chloride) led to a violent explosion [4]. Subsequent investigation showed that mixtures of phenol and perchloric acid are thermally unstable (ester formation?) and may decompose violently, the temperature range depending on composition. DSC measurements showed that sulfoxides alone may decompose violently at elevated temperatures; e.g. dimethyl sulfoxide, 270–355, cyclohexyl methyl sulfoxide, 181–255, or methyl phenyl sulfoxide, 233–286°C, respectively [6].

See Dimethyl sulfoxide: Metal oxosalts

Sulfuric acid, Organic materials

Young, E. G. *et al.*, *Science* (New York), 1946, **104**, 353

Precautions are necessary to prevent explosions when using the mixed acids to oxidise organic materials for subsequent analysis. The sulfuric acid probably tends to dehydrate the 70% perchloric acid to produce the hazardous anhydrous acid.

See Nitric acid, etc., above

Sulfur trioxide

Pascal, 1960, Vol. 16, 301

Interaction of the anhydrous acid and sulfur trioxide is violent and highly exothermic, even in presence of chloroform as diluent, and explosions are frequent.

See Dehydrating agents, above

Trichloroethylene

Prieto, M. A. *et al.*, *Research on the Stabilisation and Characterisation of Highly Concentrated Perchloric Acid*, 36, Whittier, Calif., American Potash and Chem. Corp., 1962

The solvent reacts violently with the anhydrous acid.

Trimethylplatinum hydroxide

See Dodecamethyltetraplatinum(IV) perchlorate

Vegetable matter

Tabatabai, M. A. *et al.*, *Agron. J.*, 1982, **74**, 404–406

As a safer alternative to digestion of vegetable matter with perchloric acid, alkaline oxidation of sulfur compounds to sulfate by sodium hypobromite, and reduction of sulfate to hydrogen sulfide by hydriodic acid/formaldehyde/phosphinic acid is recommended.

See Sulfuric acid, Organic materials, above

Zinc phosphide

Muir, G. D., private comm., 1968

Use of perchloric acid to assist solution of a sample for analysis caused a violent reaction.

See other REDOX REACTIONS

See other INORGANIC ACIDS, OXIDANTS, OXOHALOGEN ACIDS

3999. Mercury(II) amide chloride

[10124-48-8]



Halogens

Schwarzenbach, V., *Ber.*, 1875, **8**, 1231–1234

Schwarz min after addition of ethanol to a mixture of the amide chloride ('fusible white precipitate') and iodine, an explosion occurs. Addition of the compound to chlorine gas or bromine vapour leads to a delayed violent or explosive reaction. Amminmetal salts behave similarly, and formation of *N*-halogen compounds is involved in all cases.

See other *N*-METAL DERIVATIVES, MERCURY COMPOUNDS

4000. Chloramine (Chloramide)

[10599-90-3]



1. Marckwald, W. *et al.*, *Ber.*, 1923, **56**, 1323

2. Coleman, G. H. *et al.*, *Inorg. Synth.*, 1939, **1**, 59

3. Walek, W. *et al.*, *Tetrahedron*, 1976, **32**, 627

The solvent-free material, isolated at -70°C , disproportionates violently (sometimes explosively) at -50°C to ammonium chloride and nitrogen trichloride [1]. Ethereal solutions of chloramine are readily handled [2]. In the preparation of chloramine by reaction of sodium hypochlorite with ammonia, care is necessary to avoid excess chlorine in the preparation of the hypochlorite from sodium hydroxide, because nitrogen trichloride may be formed in the subsequent reaction with ammonia [3].

See other N-HALOGEN COMPOUNDS

4001. Ammonium chloride

[12125-02-9]

ClH_4N

NH_4Cl

HCS 1980, 142

1. Alfenaar, M. *et al.*, Neth. Appl. 73 07 035, 1974
2. Knothe, M., *Schriftenr. GDMB*, 1998, (82), 101

Ammonium chloride is generally inoffensive. However, subjected to oxidation, especially electrochemically, it has the potential to form nitrogen trichloride. Cyanogen halides may be prepared by electrolysis of hydrogen cyanide or its salts mixed with halide salts. If ammonium chloride is used as the halide salt, precautions to prevent formation of explosive nitrogen trichloride are necessary [1]. Nitrogen trichloride is an under-recognised hazard of electrolytic operations in hydrometallurgy, where it may appear at $\text{pH} < 9$ from electrolysis of ammonium or amine salts in chloride electrolytes [2].

See ELECTROLYSIS

Interhalogens

See Bromine trifluoride: Ammonium halides

Bromine pentafluoride: Acids, etc.

Other reactants

Yoshida, 1980, 53

MRH values calculated for 10 combinations with various reagents are given.

Potassium chlorate

MRH 1.46/60

See Potassium chlorate: Ammonium chloride

See related METAL HALIDES

4002. Hydroxylaminium chloride

[5470-11-1]

ClH_4NO

$\text{HON}^+\text{H}_3 \text{Cl}^-$

FPA H66, 1978

Manganese dioxide

Chaterjee, B. P. *et al.*, *Talanta*, 1977, **24**, 180–181


In a method for titrimetric determination of manganese in pyrolusite ore, addition of the powdered ore to a cold 20% solution of the salt causes vigorous decomposition to occur.

See entry HYDROXYLAMINIUM SALTS

4003. Ammonium chlorate

[10192-29-7]

ClH_4NO_3



1. Brauer, 1963, Vol. 1, 314
2. Urbanski, 1965, Vol. 2, 476

It occasionally explodes spontaneously, and invariably above 100°C [1]. It will explode after 11 h at 40°C, and after 45 min at 70°C. Ammonium and chlorate salts should not be mixed together [2].

Water

Mellor, 1941, Vol. 2, 339; 1956, Vol. 2, Suppl. 1, 591


A cold saturated solution may decompose explosively after a few days if much excess solid is present. Hot aqueous solutions have exploded during evaporation in steam heated vessels.

See other OXOSALTS OF NITROGENOUS BASES

4004. Ammonium perchlorate

[7790-98-9]

ClH_4NO_4



1. MCA Case History No. 1002
2. Barret, P., *Cah. Therm.*, 1974, **4**, 13–22
3. Seltzer, R. J., *Chem. Eng. News*, 1988, **66**(32), 7

Materials for a batch of ammonium perchlorate castable propellant were charged into a mechanical mixer. A metal spatula was left in accidentally, and the contents ignited when the mixer was started, owing to local friction caused by the spatula. A tool-listing safety procedure was instituted [1]. The literature on the kinetics of thermal decomposition has been reviewed critically [2].

For an account of the destruction of a plant producing ammonium perchlorate, which was not generally considered to be explosive, see [3]. Below 400°C it behaves as an oxidising agent but decomposition is not violent. The event suggests that tonnage quantities can detonate in some circumstances. The fire was started in spilt waste by sparks from welding and spread to storage areas on asphalt (fuel!); two aluminium bins, each containing five tonnes of perchlorate then detonated.

Aluminium

MRH 9.99/27

Loftus, H. J. *et al.*, *Rept. AD-769283/3GA*, 1–98, Springfield (Va.), USNTIS, 1973

The powdered solid materials have been evaluated as a practical propellant pair.

Carbon

MRH 4.52/9

Galwey, A. K. *et al.*, *Trans. Faraday Soc.*, 1960, **56**, 581

Below 240°C intimate mixtures with sugar charcoal undergo exothermic decomposition, while mild explosions occur above 240°C.

Catalysts

1. Solymosi, F., *Acta Phys. Chem.*, 1974, **20**, 83–103

2. Shadman-Yazdi, F., *Proc. 1973 Iran. Congr. Chem. Eng.*, 1974, **1**, 353–356

3. Glazkova, A. P., *Chem. Abs.*, 1976, **85**, 162846

4. Rastogi, R. P. *et al.*, *J. Appl. Chem. Biotechnol.*, 1978, **28**, 889–894

In a review of the course and mechanism of the catalytic decomposition of ammonium perchlorate, the considerable effects of metal oxides in reducing the explosion temperature of the salt are described [1]. Solymosi's previous work had shown reductions from 440° to about 270° by dichromium trioxide, to 260° by 10 mol% of cadmium oxide and to 200°C by 0.2% of zinc oxide. The effect of various concentrations of 'copper chromite', copper oxide, iron oxide and potassium permanganate on the catalysed combustion of the propellant salt was studied [2]. Similar studies on the effects of compounds of 11 metals and potassium dichromate in particular, have been reported [3]. Presence of calcium carbonate or calcium oxide has a stabilising effect on the salt, either alone or in admixture with polystyrene [4].

Copper

MRH 2.17/57

Anon., *Chem. Engrg.*, 1955, **62**(12), 335

Crystalline ammonium perchlorate ignited in contact with hot copper pipes.

Ethylene dinitrate

MCA Case History No. 1768

Samples of mixtures of ammonium perchlorate and the highly explosive liquid nitrate kept at 60°C ignited after 7 days. Many adverse criticisms of the general planning and execution of the experiments were made.

Impurities

1. Jacobs, P. W. M. *et al.*, *Chem. Rev.*, 1969, **69**, 590

2. Jain, S. R. *et al.*, *Combust. Flame*, 1981, **40**, 113–120

The medium impact-sensitivity of this solid propellant component is greatly increased by co-crystallisation of certain impurities, notably nitryl perchlorate, potassium periodate and potassium permanganate [1]. The presence of certain minimum amounts of mono-, di-, tri- or tetra-methylammonium perchlorates in the salt leads to a single step decomposition, at around 290°C for the mono-derivative [2].

See other CATALYTIC IMPURITY INCIDENTS

Metal perchlorates

1. Solymosi, F., *Combust. Flame*, 1966, **10**, 398–399

2. Solymosi, F., *Z. Phys. Chem.*, 1969, **67**, 76–85

3. Patel, K. C. *et al.*, *Combust. Flame*, 1976, **27**, 295–298
4. Al Fakir, M. S., *Progr. Astronaut. Aeronaut.*, 1981, **76**, 5512–564

Admixture of lithium perchlorate [1] or zinc perchlorate [2] leads to decomposition with explosion at 290° or ignition at 240°C, respectively. The role of ammine derivatives of lithium and magnesium perchlorates in catalysing the thermal decomposition of ammonium perchlorate has been studied [3], and lithium perchlorate has a strong catalytic effect on the burning rate [4].

Metals, or Organic materials,
or Sulfur

MRH Magnesium 12.97/49
MRH Sulfur 3.59/27

1. *Haz. Chem. Data*, 1975, 51
2. Klager, K. *et al.*, *Chem. Abs.*, 1982, **97**, 94903

The powdered oxidant functions as an explosive when mixed with finely divided metals, organic materials or sulfur, which increase the shock-sensitivity up to that of picric acid [1]. The hazardous properties of such mixtures increase as the particle size of the oxidant salt decreases [2].

Nitrophenol–formaldehyde polymer

Girdhar, H. L. *et al.*, *Indian J. Technol.*, 1981, **19**, 531–532

The combination (or one with potassium perchlorate) is an explosive propellant.

Other reactants

Yoshida, 1980, 79


MRH values calculated for 19 combinations with oxidisable materials are given.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

4005. Hydroxylaminium perchlorate

[15588-62-2]

ClH_4NO_5



1. Rafeev, V. A. *et al.*, *Chem. Abs.*, 1981, **95**, 83166
2. Bokii, V. A. *et al.*, *Chem. Abs.*, 1981, **94**, 49734

The decomposition and combustion of the redox salt [1], and its mechanism [2] were studied.

1,3,5-Triaminotrinitrobenzene

Quong, R., *Chem. Abs.*, 1975, **82**, 113753

A saturated aqueous solution of the perchlorate, and the solid fuel, are individually non-explosive, but form a viable explosive composition on admixture.

See other PERCHLORATE SALTS OF NITROGENOUS BASES, REDOX COMPOUNDS

4006. Phosphonium perchlorate

[101672-21-3]

$\text{ClH}_4\text{O}_4\text{P}$



1. Fichter, F. *et al.*, *Helv. Chim. Acta*, 1934, **17**, 222
2. Mellor, 1971, Vol. 8, Suppl. 3, 274

The crystalline salt obtained by action of phosphine on 68% perchloric acid at -20°C is dangerously explosive, sensitive to moist air, increase in temperature, or friction [1] and cannot be dried [2].

See related PERCHLORATE SALTS OF NITROGENOUS BASES

See other REDOX COMPOUNDS

4007. Hydrazinium chloride

[2644-70-4]




ClH_5N_2

See entry THERMAL EXPLOSIONS *See other* HIGH-NITROGEN COMPOUNDS

4008. Hydrazinium chlorite

[66326-45-2]



$\text{ClH}_5\text{N}_2\text{O}_4$

Levi, G. R., *Gazz. Chim. Ital.* [2], 1923, **53**, 105–108

It is spontaneously flammable when dry.

See other CHLORITE SALTS, REDOX COMPOUNDS

4009. Ammonium perchlorylamide

[20394-65-4]




$\text{ClH}_5\text{N}_2\text{O}_3$

See entry PERCHLORYLAMIDE SALTS

4010. Hydrazinium chlorate

[66326-46-3]



$\text{ClH}_5\text{N}_2\text{O}_3$


Salvadori, R., *Gazz. Chim. Ital.* [2], 1907, **37**, 32–40

It explodes violently at its m.p., 80°C .

See other OXOSALTS OF NITROGENOUS BASES, REDOX COMPOUNDS

4011. Hydrazinium perchlorate

[13762-80-6]



$\text{ClH}_5\text{N}_2\text{O}_4$

1. Levy, J. B. *et al.*, *ACS* 54, 1966, 55

2. Grelecki, C. J. *et al.*, *ACS* 54, 1966, 73

3. Salvadori, R., *Gazz. Chim. Ital.* [2], 1907, **37**, 32–40

The deflagration [1] and thermal decomposition [2] of the salt, a component of solid rocket propellants, have been studied. It also explodes on impact [3].

Metal salts, or Metal oxides

1. Shidlovskii, A. A. *et al.*, *Zh. Priklad. Chim.*, 1962, **35**, 756

2. Probhakaran, C. *et al.*, *Def. Sci. J.*, 1982, **31**, 285–291

Presence of 5% of copper(II) chloride caused explosion to occur at 170°C [1]. Of the series of additives copper chromite, copper chloride, nickel oxide, iron oxide, magnesium oxide, the earlier members have the greatest effect in increasing the sensitivity of the perchlorate to heat, impact and friction.

See other PERCHLORATE SALTS OF NITROGENOUS BASES, REDOX COMPOUNDS

4012. Poly(dimercuryimmonium perchlorate)

[]

(ClHg₂NO₄)_n

(Hg=N⁺=Hg ClO₄⁻)_n

Sorbe, 1968, 97

Highly explosive.

See other POLY(DIMERCURYIMMONIUM) COMPOUNDS, PERCHLORATE SALTS OF NITROGENOUS BASES

4013. Iodine chloride

[7790-99-0]

ClI

ICI

Metals

Mellor, 1940, Vol. 2, 119; 1956, Vol. 2, Suppl. 1, 452; 1963, Vol. 2, Suppl. 2.2, 1563

Mixtures containing sodium explode only on impact, while potassium explodes on contact with the chloride. Aluminium foil ignites after prolonged contact (probably after the surface layer of oxide has been dissolved).

Phosphorus trichloride

Mellor, 1956, Vol. 2, Suppl. 1, 502

Reaction is intensely exothermic.

See other INTERHALOGENS, IODINE COMPOUNDS

4014. Indium(I) perchlorate

[62763-56-8]

ClInO₄

InClO₄

Ashraf, M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1977, 1723

The solvent-free material detonated when crushed with a glass rod. The nitrate (also a redox compound) probably would behave similarly.

See other METAL PERCHLORATES, REDOX COMPOUNDS

4015. Potassium chloride
[7447-40-7]

KCl

ClK

Potassium permanganate, Sulfuric acid

See Potassium permanganate: Potassium chloride, etc.

See other METAL HALIDES

4016. Potassium chlorite
[14314-27-3]

KClO₂

ClKO₂

Other reactants

Yoshida, 1980, 3

MRH values calculated for 18 combinations with oxidisable materials are given.

Sulfur

MRH 3.22/13

Leleu, *Cahiers*, 1974, (74), 137

Interaction is violent.

See other CHLORITE SALTS

4017. Potassium chlorate
[3811-04-9]

KClO₃

ClKO₃

HCS 1980, 762

1. Rüst, 1948, 294

2. Biasutti, 1981, 151

3. Anon., *Fire Prevention*, 1986, (186), 48

Although most explosive incidents have involved mixtures of the chlorate with combustible materials, the exothermic decomposition of the chlorate to chloride and oxygen can accelerate to explosion if a sufficient quantity and powerful enough heating are involved. A case history of a fire-heated explosion of a store of 80 t of chlorate is given. The more stable sodium chlorate will also explode under similar conditions [1]. The enthalpy of conversion of potassium chlorate to either chloride or perchlorate is slight, about $\frac{1}{4}$ kJ/g, less than is the case for the sodium salt. During sieving of the dry chlorate in a store attached to a fuse igniter plant, a violent explosion of 75 kg of the salt occurred. It seems likely that some oxidisable material was involved in the incident, though the possibility of dropping a drum onto the concrete floor is noted as a potential cause of ignition [2]. An account of an explosion of 16 tonnes during a fire in a warehouse, demolishing 100 metres of the wall is given. The chlorate does not seem to have been isolated from other goods [3].

Agricultural materials

See Sodium chlorate: Agricultural materials

Aluminium, MRH 9.20/30
Antimony trisulfide MRH 2.84/39

Crozier, T. H. *et al.*, *52nd Ann. Rept. HM Insp. Explos.*, London, Home Office, 1928

A pyrotechnic mixture containing the powdered ingredients was found dangerously sensitive to frictional initiation and highly explosive.

See Metals; *and* Metal sulfides; both below

Ammonia, or Ammonium sulfate MRH Ammonia 4.23/21

Mellor, 1941, Vol. 2, 702; 1940, Vol. 8, 217

High concentrations of ammonia in air react so vigorously with potassium chlorate as to be dangerous. Mixtures with ammonium sulfate when heated decompose with incandescence.

Ammonium chloride

1. Potjewijd, T., *Pharm. Weekbl.*, 1935, **72**, 68–69

2. Ellern, 1968, 155

Addition of ammonium chloride to a drum of weed-killer was suspected as the cause of a violent explosion (involving formation of ammonium chlorate) [1].

Mixtures, used for smoke compositions, are hazardous [2].

Aqua regia, Ruthenium

Sidgwick, 1950, 1459

Ruthenium is insoluble in aqua regia, but addition of potassium chlorate causes explosive oxidation.

Arsenic trisulfide

Ganguly, A., *J. Indian Acad. Forensic Sci.*, 1973, **12**, 29–30

Dry powdered mixtures containing over 30% oxidant exploded under a hammer-blow.

See Metal sulfides, below

Carbon MRH 4.52/13

Read, C. W. W., *School Sci. Rev.*, 1941, **22**(87), 341

Accidental substitution of powdered carbon for manganese dioxide in 'oxygen mixture' caused a violent explosion when the mixture was heated.

Cellulose

1. Watanabe, M. *et al.*, *Chem. Abs.*, 1988, **108**, 207227

2. Ishida, H. *et al.*, *Chem. Abs.*, 1988, **109**, 56797

Mixtures of stoichiometric proportions (zero oxygen balance) are a high deflagration hazard and show remarkable pressure increase effects on ignition [1], as well as lowest ignition temperatures by ARC [2].

See PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION (reference 4)

Charcoal, Potassium nitrate, Sulfur

Biasutti, 1981, 4

ClK₃

A batch of black powder, modified by addition of potassium chlorate, was being mixed mechanically with added water. A friction-sensitive crust appears to have formed, leading to initiation of the explosion which followed.

See Non-metals, below

Charcoal, Sulfur

Biasutti, 1981, 15, 42, 46

A demonstration by Berthollet in 1788 of replacement of potassium nitrate in gunpowder by the chlorate led to a violent explosion during the crushing operation which caused two fatalities. Later incidents involving factories for chlorate-containing explosives led to widespread destruction.

Cyanides

See METAL CYANIDES (AND CYANO COMPLEXES) (reference 1)

Cyanoguanidine

See Cyanoguanidine: Oxidants

Dinickel trioxide

Mellor, 1942, Vol. 15, 395

Interaction at 300°C is violently exothermic, red heat being attained.

Fabric

Anon., *Accidents*, 1968, 24

Fabric gloves (wrongly used in place of impervious plastics gloves) became impregnated during handling operations and were subsequently ignited from cigarette ash.

See Cellulose, above

Fluorine

Pascal, 1960, Vol. 16, 316

Interaction at low pressure leads to formation of the explosive gas, fluorine perchlorate.

See Fluorine: Sodium bromate

Gallic acid, Gum tragacanth

Anon., *Lab. Accid. Higher Educ.*, item 7, Barking, HSE, 1987

A 'pyrotechnic device' was being prepared by loading 16 g of a mixture of 73% potassium chlorate, 24% gallic acid and 3% gum tragacanth into a 9 mm copper tube 280 mm long, when it exploded violently and caused shrapnel injuries. Such mixtures are very friction-sensitive.

Hydrocarbons

MRH Hexane 4.77/10, Toluene 4.81/12

1. Bjorkman, P. O., *Chem. Abs.*, 1934, **28**, 6311a

2. von Feilitzen, G., *Chem. Abs.*, 1934, **28**, 6311b

Mixtures of powdered chlorate and hydrocarbons explode as violently as nitro compound explosives [1]. Porous masses of chlorate impregnated with hydrocarbons are friction-sensitive explosives [2].

Hydrogen iodide

Mellor, 1941, Vol. 2, 310

Molten potassium chlorate ignites hydrogen iodide gas.

Manganese dioxide

1. Mellor MIC, 1961, 333

2. Partington, 1967, 651

3. Renfrew, M. M., *J. Chem. Educ.*, 1983, **60**, A229

When oxygen is generated in the laboratory by heating potassium chlorate with manganese dioxide as decomposition catalyst, the latter must be free of organic matter or an explosion will occur [1]. The mixture (17% wt of manganese dioxide, 'oxygen mixture') must be cautiously heated at 250°C with a small flame to prevent the exothermic decomposition reaction becoming violent [2]. Mixing the components dry with a partially plastic-coated micro-spatula caused a fire [3].

See other CATALYTIC IMPURITY INCIDENTS

Manganese dioxide, Potassium hydroxide

Molinari, E. *et al.*, *Inorg. Chem.*, 1964, **3**, 898

The oxidation of manganese dioxide to manganate by solid alkali-chlorate mixtures becomes explosive above 80–90°C at pressures above 19 kbar.

Metal, Wood

Davis, G. C., *J. Ind. Eng. Chem.*, 1909, **1**, 118

Explosions were caused by transportation of metal castings in wooden kegs previously used to store potassium chlorate, impact or friction of the metal causing initiation of the chlorate-impregnated wood.

Metal phosphides

Mellor, 1940, Vol. 8, 839, 844

Tricopper diphosphide and trimercury tetraphosphide form impact-sensitive mixtures with potassium chlorate. By analogy, the phosphides of aluminium, magnesium, silver and zinc, etc., would be expected to form similar mixtures with metal halogenates.

Metal phosphinates ('hypophosphites')

Mellor, 1940, Vol. 8, 881, 883

Dry mixtures of barium phosphinate and potassium chlorate burn rapidly with a feeble report if unconfined, but even under the slight confinement of enclosing in paper, a sharp explosion occurs. The mixture is readily initiated by sparks, impact or friction. A mixture of calcium phosphinate, potassium chlorate and quartz exploded during mixing. Mixtures of various phosphinates and chlorates have been proposed as explosives, but they are very sensitive to initiation by sparks, friction or shock. Admixture of powdered magnesium causes a brilliant flash on initiating the mixture.

See Sodium phosphinate: Oxidants

See other FRICTIONAL INITIATION INCIDENTS, STATIC INITIATION INCIDENTS

Metals MRH Aluminium 9.20/30, magnesium 9.50/37, iron 3.64/47

1. Mellor, 1941, Vol. 2, 310
2. Anon., *Chem. Eng. News*, 1936, **14**, 451
3. Mellor, 1940, Vol. 4, 480
4. Mellor, 1943, Vol. 11, 163
5. Mellor, 1941, Vol. 7, 20, 116, 260
6. Jackson, H., *Spectrum*, 1969, **7**, 82
7. Ganguly, A., *J. Indian Acad. Forensic Sci.*, 1973, **12**, 29–30

Mixtures of finely divided aluminium, copper, magnesium [1,2] and zinc [3] with potassium chlorate (or other metal halogenates) are explosives and may be initiated by heat, impact or light friction. Chromium incandesces in the molten salt [4] and germanium explodes on heating with potassium chlorate. Titanium explodes on heating, while zirconium gives mild explosions on heating, and ignites when the mixture is impacted [5]. Contact of molten chlorate with steel wool causes violent combustion of the latter [6]. Qualitative experiments showed that hammer impact would explode mixtures with aluminium powder containing over 10% of chlorate [7].

See also FLASH POWDER

See other METAL OXOHALOGENATES, METAL CHLORATES

Metal sulfides MRH Antimony trisulfide 2.34/39, silver sulfide 1.38/62

1. Mellor, 1941, Vol. 2, 310
2. Mellor, 1939, Vol. 9, 523
3. Mellor, 1941, Vol. 3, 447
4. Rüst, 1948, 335

Many metal sulfides when mixed intimately with metal halogenates form heat-, impact- or friction-sensitive explosive mixtures [1]. That with antimony trisulfide can be initiated by a spark [2] and with silver sulfide a violent reaction occurs on heating [3]. For the preparation of 'oxygen mixture', antimony trisulfide was used in error instead of manganese dioxide, and during grinding, the mixture of sulfide and chlorate exploded very violently [4].

See also METAL HALOGENATES

Metal thiocyanates

1. von Schwartz, 1918, 299–300
2. Anon., *Chem. Age.*, 1936, **35**, 42

Mixtures of thiocyanates with chlorates (or nitrates) are friction- and heat-sensitive, and explode on rubbing, heating to 400°C, or initiation by spark or flame [1]. A violent explosion occurred when a little chlorate was ground in a mortar contaminated with ammonium thiocyanate. A similar larger-scale explosion involving traces of barium thiocyanate is also described [2].

Nitric acid, Organic materials

Asthana, S. S. *et al.*, *Chem. & Ind.*, 1976, 953–954

In a new method of degrading organic matter prior to analysis, small portions (0.3 g only) of chlorate are added to a hot suspension of the organic matter in

conc. nitric acid. Use of larger portions of chlorate may lead to explosive oxidation of the organic matter.

Non-metals MRH Carbon 4.52/13, phosphorus 5.98/23, sulfur 3.18/18

1. Mellor, 1941, Vol. 2, 310
2. Mellor, 1940, Vol. 8, 785–786
3. Mellor, 1946, Vol. 5, 15
4. Ganguly, A., *J. Indian Acad. Forensic Sci.*, 1973, **12**, 29–30

Potassium chlorate (or other metal halogenate) intimately mixed with arsenic, carbon, phosphorus, sulfur or other readily oxidised materials gives friction-, impact- and heat-sensitive mixtures which may explode violently [1]. When potassium chlorate is moistened with a solution of phosphorus in carbon disulfide, it eventually explodes as the solvent evaporates and oxidation proceeds [2]. Boron burns in molten chlorate with dazzling brilliance [3]. Mixtures of the chlorate and finely powdered sulfur containing over 20% of the latter will explode under a hammer-blow [4].

See Sulfur, below

See also METAL CHLORATES, METAL HALOGENATES

Other reactants

Yoshida, 1980, 65–66

MRH values calculated for 24 combinations with oxidisable materials are given.

Phosphorus (red)

1. Haarmann, D. J., *American Fireworks News*, 1985, #51; *ibid.*, 1986, #54,
2. *Explosives and Their Power*, Berthelot, M., (translated and condensed from the French), London, 1892

This combination has, in the past, been the base of various impact sensitive pyrotechnics, described by Haarmann [1] and more recently on the Internet, including reference to a report [2] of a mere 60–70 kg, dispersed in children's toy caps, demolishing a building in an accidental explosion. Red phosphorus is a material of variable composition and reactivity, so unreliability is to be anticipated. There have been passivating components in most of these mixtures, it is suggested that the pure dry powders will often react on contact. For the more reactive white allotrope:

See Non-metals, above

Reducants

Mellor, 1941, Vol. 3, 651; 1940, Vol. 8, 881, 883

Mixtures with calcium hydride or strontium hydride may explode readily, and interaction of the molten chlorate is, of course, violent. A mixture of syrupy sodium phosphinate ('hypophosphite') and the powdered chlorate on heating eventually explodes as powerfully as glyceryl nitrate. Calcium phosphinate mixed with the chlorate and quartz detonates (the latter producing friction to initiate the mixture). Dried mixtures of barium phosphinate and the chlorate are very sensitive and highly explosive under the lightest confinement (screwed up in paper).

Sugars

1. Zaehring, A. J., *Rocketscience*, 1949, **3**, 64–68
2. Ganguly, A., *J. Indian Acad. Forensic Sci.*, 1973, **12**, 29–30
3. Scanes, F. S., *Combust. Flame*, 1974, **23**, 363–371

A stoichiometric mixture with sucrose ignites at 159°C, and has been evaluated as a rocket propellant [1]. Dry powdered mixtures with glucose containing above 50% of chlorate explode under a hammer-blow [2]. Pyrotechnic mixtures with lactose begin to react exothermally at about 200°C, when the lactose melts, and carbon is formed. This is then oxidised by the chlorate at about 340°C. The mechanism was studied by DTA [3].

Sugar, Sulfuric acid

Hanson, R. M., *J. Chem. Educ.*, 1976, **53**, 578

Addition of a drop of sulfuric acid to a mound of the chlorate–sugar mixture leads to ignition.

Sulfur

MRH 3.18/18

1. Tanner, H. G., *J. Chem. Educ.*, 1959, **36**, 59
2. Storey, P. D., *Proc. 13th Int. Pyrotech. Semin.*, 1988, 765–784
3. Chapman, D. *et al.*, *J. Pyrotech.*, 1998 (7), 51

The chemistry involved in this explosively unstable system is reviewed [1]. The mechanism of the trigger reactions that initiate the exothermic decomposition of chlorate–sulfur mixtures has been studied. Mixtures containing 1–30% of sulfur can decompose well below the m.p. of sulfur, and addition of sulfur dioxide, the suspected chemical trigger, causes immediate onset of the reaction [2]. Autoignition of stoichiometric mixtures can be as low as 115°C, with frictional sensitivity at 5N, the lowest load the test apparatus permitted. Both were dependent upon the history of the sulphur used [3].

See Non-metals, above

Sulfur, Metal derivatives

McKown, G. L. *et al.*, *Chem. Abs.*, 1977, **87**, 103955

The sensitivity of chlorate–sulfur mixtures to initiation is increased by cobalt and its oxide, greatly so by copper nitride, copper sulfate, and extremely so by copper chlorate. Implications for manufacturing operations are discussed.

Sulfur dioxide

MRH 1.80/56

Mellor, 1947, Vol. 10, 217; 1941, Vol. 2, 311

Contact at temperatures above 60°C causes flashing of the evolved chlorine dioxide. Solutions of sulfur dioxide in ethanol or ether cause an explosion on contact at ambient temperature.

Sulfuric acid

1. Mellor, 1947, Vol. 10, 435
2. Bailar, 1973, Vol. 2, 1362

Addition of potassium chlorate in portions to sulfuric acid maintained at below 60 or above 200°C causes brisk effervescence. At intermediate temperatures,

explosions are caused by the chlorine dioxide produced, and these reach maximum intensity at 120–130°C [1]. Uncontrolled contact of any chlorate with sulfuric acid may be explosive. The need for great caution with this system was stressed by Davy in 1815 [2].

See METAL CHLORATES: acids

Sodium amide

MRH 2.63/24

Mellor, 1940, Vol. 8, 258

A mixture explodes.

Tannic acid

Rüst, 1948, 336

A mixture exploded during grinding in a mortar.

See METAL CHLORATES

Thiourea

1. Soothill, D., *Safety Management*, 1992, **8**(6), 11.

2. Wharton, R. K. *et al.*, *Propellants, Explos., Pyrotech.*, 1993, **18**(2), 77

An explosion is mentioned consequent upon grinding potassium chlorate in equipment previously used for thiourea. It is claimed the mixture explodes spontaneously [1]. This is an exaggeration, it is very sensitive to friction, shock and heat (ignition <155°C) [2].

Thorium dicarbide

See Thorium dicarbide: Non-metals, etc.

See other METAL CHLORATES, OXIDANTS

4018. Potassium perchlorate

[7778-74-7]

KClO₄

KClO₄

Aluminium, Aluminium fluoride

1. Hahma, A., *Propellant, Explosives, Pyrotechnics*, 1996, 21, 100

2. Freeman, E. S. *et al.*, *Combust. Flame*, 1966, **80**, 16

Whether mixtures of the perchlorate with aluminium truly detonate is under debate. However, appropriate compositions certainly explode violently even when uncontained, so the deflagration/detonation dispute is essentially academic [1]. Presence of aluminium fluoride increases the ease of ignition of aluminium–perchlorate mixtures, owing to complex fluoride formation [2].

Aluminium, Barium nitrate, Potassium nitrate, Water

See Aluminium: Metal nitrates, etc.

Aluminium powder, Titanium dioxide

Anon., *Fire Prot. Assoc. J.*, 1957, (36), 9

ClKO₄

A mixture of the 3 compounds exploded violently during mixing. Previously the mixture had been accidentally ignited by a spark. Aluminium powder is incompatible with oxidants, and its mixture with titanium dioxide is a thermite-like combination.

See Aluminium: Metal oxides

See also THERMITE REACTIONS

Barium chromate, Tungsten and/or Titanium

1. Carrazza, J. A. *et al.*, *Chem. Abs.*, 1975, **82**, 113751

2. Nagaishi, T., *Chem. Abs.*, 1978, **88**, 173088

Mixtures containing 65% of tungsten with organic binders were developed as priming charges for surface flares [1]. Ignition of the dual metal-oxidant mixtures by IR irradiative heating was studied [2].

See other IRRADIATION DECOMPOSITION INCIDENTS

Boron, Magnesium, Silicone rubber

Paulin, C. J. *et al.*, US Pat, 4 101 352, 1978

Ignition of a compressed mixture, applied as a 6 mm layer on electronic components generates enough heat effectively to destroy them.

Combustible materials

MRH Ethylene glycol 4.48/27

Grodzinski, J., *J. Appl. Chem.*, 1958, **8**, 523-528

Explosion temperatures were determined for a wide range of combustible liquid and solid organic materials sealed into glass tubes with potassium perchlorate. The lowest temperatures were shown by mixtures with ethylene glycol (240°), cotton linters (245°) and furfural (270°C).

Ethanol

MRH 4.85/18

Burton, H. *et al.*, *Analyst*, 1955, **80**, 16

Many explosions have been experienced during the gravimetric determination of either perchlorates or potassium as potassium perchlorate by a standard method involving an ethanol extraction. During subsequent heating, formation and explosion of ethyl perchlorate is very probable.

Ferrocenium diamminetetrakis(thiocyanato-*N*)chromate(1-)

Guslev, V. G. *et al.*, *Chem. Abs.*, 1974, **86**, 57615

Presence of 25% of the organometallic salt (ferrocenium reineckate) considerably increases the rate of thermal decomposition of the perchlorate, involving hydrogen cyanide arising from the thiocyanato groups.

Lactose

See Lactose: Oxidants

Metal powders

MRH Aluminium 9.96/33, iron 3.76/51, magnesium 10.25/41, sodium 4.27/45

1. Schumacher, 1960, 210

2. Riffault, M. L., *Chem. Abs.*, 1975, **82**, 127130

3. Ywenkeng, H., *Chem. Abs.*, 1981, **94**, 194474
4. Rittenhouse, C. *et al.*, *Proc. 6th Int. Pyrotech. Semin.*, 1978, 519–536
5. Wild, R. *J. Haz. Mat.*, 1982, **7**, 75–79
6. Munger, A. C. *et al.*, *Chem. Abs.*, 1973, **98**, 91971
7. Parks, R. L., *Chem. Abs.*, 1981, **95**, 100052
8. Sarawadekar, R. G. *et al.*, *Proc. 8th Int. Protech. Semin.*, 1982, 574–587

The mixture of aluminium and/or magnesium powders with potassium perchlorate (a photoflash composition) is very readily ignited and three industrial explosions have occurred. Mixtures of nickel and titanium powders with the oxidant and infusorial earth are very friction-sensitive, causing severe explosions, and easily ignited by very small (static) sparks [1]. A mixture containing 70% of molybdenum powder ignites at 330°C [2], and the rate of combustion of a mixture containing 48% is accelerated by 1% of various additives [3]. The pyrotechnic mixture with aluminium is insensitive to spark discharges but ignited consistently with a hot wire source [4], and blast waves produced on ignition of mixtures with 70% oxidant have been investigated [5]. The mixture with iron powder is sufficiently exothermic after ignition for use in metal welding, and an iron oxide–aluminium–nickel thermite mixture was also developed for the same purpose [6]. Data on detonation and deflagration of the pyrotechnic mixture with titanium were determined in open and closed vessels, and subsequently used to design a containment room for blending the powdered components [7]. The pyrotechnic/explosive perchlorate–tantalum system has been studied [8].

See Aluminium, etc., Boron, etc. and Barium chromate, etc., all above

See other FRICTIONAL INITIATION INCIDENTS, STATIC INITIATION INCIDENTS

See also THERMITE REACTIONS

Other reactants

Yoshida, 1980, 80

MRH values calculated for 18 combinations with oxidisable materials are given.

Potassium hexacyanocobaltate(3–)

Massis, T. M. *et al.*, *Chem. Abs.*, 1977, **87**, 8142

Mixtures serve as gasless pyrotechnic compositions.

Reducers

See Metal powders, above; Titanium hydride, below

Sulfur

MRH 3.39/20

Schumacher, 1960, 211–212

Mixtures of sulfur and potassium perchlorate, used in pyrotechnic devices, can be exploded by moderate impact. All other inorganic perchlorates form such impact-sensitive mixtures.

Titanium hydride

1. Massis, T. M. *et al.*, *Rept. SAND-75-5889*, Richmond (Va.), USNTIS, 1976

2. Collins, L. W., *J. Haz. Mat.*, 1982, **5**, 325–333

3. Parks, R. L., *J. Haz. Mat.*, 1982, **5**, 359–371

The stability of the pyrotechnic mixture has been studied, including the effect of hydrogen content of the hydride [1]. Presence of moisture and impurities adversely affects stability [2], and remote handling facilities for the mixture have been developed [3].

See other METAL PERCHLORATES, OXIDANTS

4019. Potassium perchlorylamide
[15320-25-9]



Sorbe, 1968, 67

Like the monopotassium salt, it will explode on impact or exposure to flame.

See entry PERCHLORYLAMIDE SALTS *See other* N-METAL DERIVATIVES

4020. Lithium chlorite
[27505-49-3]



Janz, 1976, Table 2, p. 10

It disproportionates violently on heating.

Chlorine, 2-Chloroalkyl aryl sulfides

See Chlorine: 2-Chloroalkyl aryl sulfides, etc.

See other CHLORITE SALTS, OXIDANTS

4021. Lithium perchlorate
[7791-03-9]



Diethyl ether

1. Urben, P. G., *Chemtech*, 1991, (May), 259

2. Silva, R. A., *Chem. Eng. News*, 1992 **70**(51), 2

3. Handy, S. T. *et al.*, *Synlett.*, 1995, 565

A warning was given that the 5 molar solution in ether used as a solvent for Diels-Alder reactions would lead to explosions [1]. Such a reaction of dimethyl acetylenedicarboxylate and cyclooctatetraene in this solvent exploded very violently on heating. The cyclooctatetraene was blamed, with no supporting evidence [2]. It would appear desirable to find the detonability limits of the mixture with ether before any attempt is made to scale up. A safe alternative to lithium perchlorate/ether as a solvent for Diels-Alder reactions is proposed [3].

See Dimethyl acetylenedicarboxylate, 1,3,3,7-Cyclooctatetraene

See Magnesium perchlorate

1,3-Dioxolane

Newman, G. H. *et al.*, *J. Electrochem. Soc.*, 1980, **127**, 2025–2027

Lithium perchlorate–dioxolane electrolyte systems are unsafe for secondary battery applications, as an explosion occurred during overnight cyclic testing of a Li/TiS₂ system. The effect was duplicated under all over-discharge or cell-reversal conditions.

Hydrazine

Rosolovskii, V. Ya., *Chem., Abs.*, 1969, **70**, 53524

Interaction gives only the dihydrazine complex, which explodes on grinding, but loses hydrazine uneventfully on heating at atmospheric or reduced pressure. The sodium salt is similar.

Nitromethane

See Nitromethane: Lithium perchlorate

Olefins, Electrophiles.

Zefirov, N. S., XIII Int. Symp. Org. Chem. Sulfur, 1988 (lecture; this information not in the later published proceedings).

Electrophilic additions to olefins, especially of XCl species, can give substantial quantities of perchlorate esters as by-products if performed in the presence of lithium perchlorate. These esters are extremely explosive, and may concentrate during distillation of crude products.

See other METAL PERCHLORATES, OXIDANTS

4022. Manganese chloride trioxide

[15605-27-3]

ClMnO₃

ClMnO₃

Briggs, S. T., *J. Inorg. Nucl. Chem.*, 1968, **30**, 2867–2868

Explosively unstable if isolated as a liquid at ambient temperature, it may be handled safely in carbon tetrachloride solution.

See Manganese dichloride dioxide, Manganese trichloride oxide

See related METAL OXIDES

4023. Nitrosyl chloride

[2696-92-6]

ClNO

O:NCl

It is a moderately endothermic compound (ΔH_f° (g) +52.6 kJ/mol, 0.80 kJ/g).

Acetone, Platinum

Kaufmann, G. B., *Chem. Eng. News*, 1957, **35**(43), 602

A cold sealed tube containing nitrosyl chloride, platinum wire and traces of acetone exploded violently on being allowed to warm up.

Hydrogen, Oxygen

See Nitrogen oxide: Hydrogen, etc.

See other ENDOTHERMIC COMPOUNDS, N-HALOGEN COMPOUNDS, NITROSO COMPOUNDS, OXIDANTS

4024. N-Chlorosulfinylimide
[25081-01-0]

CINOS

O:S:NCI

Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 318

The ampouled solid exploded violently on melting. Distillation at ambient pressure and impact tests had not previously indicated instability.

Chlorine fluoride

See Chlorine fluoride: N-Chlorosulfinylamine

See other N-HALOGEN COMPOUNDS, N-S COMPOUNDS

4025. Nitryl chloride
[13444-90-1]

CINO₂

O₂NCI

It is feebly endothermic (ΔH_f° (g) +12.1 kJ/mol, 0.15 kJ/g) but a powerful oxidant gas.

Inorganic materials

Batey, H. H. *et al.*, *J. Amer. Chem. Soc.*, 1952, **74**, 3408

Interaction of the chloride with ammonia or sulfur dioxide is very violent, even at -75°C, and is vigorous with tin(II) bromide or tin(II) iodide.

Organic matter

Kaplan, R. *et al.*, *Inorg. Synth.*, 1954, **4**, 54

It attacks organic matter rapidly, sometimes explosively.

B-Trimethylborazine

Hirata, T., *Rept, AD-7293939*, Richmond (Va.), USNTIS, 1971

Interaction is violent in absence of a diluent.

See other ENDOTHERMIC COMPOUNDS, N-HALOGEN COMPOUNDS, OXIDANTS

4026. Nitryl hypochlorite ('Chlorine nitrate')
[14545-72-3]

CINO₃

O₂NOCl

Metals, or Metal chlorides

Bailar, 1973, Vol. 2, 379

Interaction with most metals and metal chlorides is explosive at ambient temperature, but controllable at between -40 and -70°C.

Organic materials

Schmeisser, M., *Inorg. Synth.*, 1967, **9**, 129

Not inherently explosive, but it reacts explosively with alcohols, ethers and most organic materials.

See Nitryl hypofluorite

See other HYPOHALITES, OXIDANTS

4027. Nitrosyl perchlorate

[15605-28-4]

ClNO₅

O:NOCIO₃

Gerding, H. *et al.*, *Chem. Weekbl.*, 1956, **52**, 282–283

Although stable at ambient temperature, it begins to decompose below 100°C, and at 115–120°C the decomposition becomes a low-order explosion.

Pentaammineazidocobalt(III) perchlorate, Phenyl isocyanate

Burmeister, J. L. *et al.*, *Chem. Eng. News*, 1968, **46**(8), 39

During an attempt to introduce phenyl isocyanate into the Co coordination sphere, a mixture of the 3 components exploded violently when stirring was stopped.

Organic materials

Hofmann, K. A. *et al.*, *Ber.*, 1909, **42**, 2031

As the anhydride of nitrous and perchloric acids, it is a very powerful oxidant. Pinene explodes sharply; acetone and ethanol ignite, then explode; ether evolves gas, then explodes after a few s delay. Small amounts of primary aromatic amines—*aniline*, *toluidines*, *xylidines*, *mesidine*—ignite on contact, while larger amounts exploded dangerously, probably owing to rapid formation of diazonium perchlorates. Urea ignites on stirring with the perchlorate, (probably for a similar reason).

See other NON-METAL PERCHLORATES, OXIDANTS

4028. Nitronium perchlorate

[12051-08-0]

ClNO₆

O₂N⁺ ClO₄⁻

Albright, Hanson, 1976, 2

The explosively unstable behaviour of stored nitronium perchlorate is attributed to the formation of small equilibrium concentrations of the isomeric covalent nitryl perchlorate ester (below).

1,2-Epoxides

Golding, P. *et al.*, *Tetrahedron Lett.*, 1988, **29**, 2733

Reaction with epoxides gives the dangerously unstable and explosive mixed nitrate–perchlorate diesters, such as 1,2-ethanediyl nitrate perchlorate from ethylene oxide.

See related NITRATION INCIDENTS
See other NON-METAL PERCHLORATES

4029. Nitryl perchlorate
[17495-81-7]



CINO₆

Ammonium perchlorate
See Ammonium perchlorate: Impurities

Organic solvents

1. Spinks, J. W. T., *Chem. Eng. News*, 1960, **38**(15), 5
2. Gordon, W. E. *et al.*, *Can. J. Res.*, 1940, **18B**, 358

Interaction with benzene gave a slight explosion and flash [1], while sharp explosions with ignition were observed with acetone and ether[2].

See other NON-METAL PERCHLORATES, OXIDANTS

4030. Chlorine azide
[13973-88-1]



CIN₃

Alone, or Ammonia, or Phosphorus, or Silver azide, or Sodium

1. Frierson, W. J. *et al.*, *J. Amer. Chem. Soc.*, 1943, **65**, 1696, 1698
2. Rice, W. J. *et al.*, *J. Chem. Educ.*, 1971, **48**, 659
3. Combourieu, J. *et al.*, *Chem. Abs.*, 1977, **87**, 123313
4. Tornieporth-Oetting, I. C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(5), 511

The undiluted material is extremely unstable, usually exploding violently without cause at any temperature, even as solid at -100°C [1]. Explosion is likely to be triggered by pressure fluctuations of around 10 Pa [4]. It gives an explosive yellow liquid with liquid ammonia; when condensed on to yellow phosphorus at -78°C an extremely violent explosion soon occurs. Addition of phosphorus to a solution of the azide in carbon tetrachloride at 0°C causes a series of mild explosions if the mixture is stirred, or a violent explosion without stirring. Contact of the liquid or gaseous azide with silver azide at -78°C gave a blue colour, soon followed by explosion, and sodium reacted similarly under the same conditions [1]. When chlorine azide (25 mol %) is used as a thermally activated explosive initiator in a chemical gas laser tube, the partial pressure of azide should never exceed 16 mbar [2]. The explosive decomposition has been studied in detail [3].

See entry HALOGEN AZIDES

4031. Sulfuryl azide chloride
[13449-15-5]



CIN₃O₂S



Preparative hazard
Shozda, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 2876

During the preparation of this explosive liquid by interaction of sulfuryl chloride fluoride and sodium azide, traces of chlorine must be eliminated from the former to avoid detonation. The product is nearly as shock-sensitive as glyceryl nitrate and may explode on rapid heating. Solutions (25 wt%) in solvents may be handled safely. The corresponding fluoride is believed to behave similarly.

See other ACYL AZIDES, ACYL HALIDES, NON-METAL AZIDES

4032. Thiotrithiazyl perchlorate

[]



Organic solvents


Goehring, 1957, 74

The precipitated perchlorate salt exploded on washing with acetone or ether.

See other NON-METAL PERCHLORATES, N-S COMPOUNDS

4033. Thiotrithiazyl chloride

[12015-30-4]



Alone, or Ammonia

1. Mellor, 1940, Vol. 8, 631-632

2. Bailar, 1973, Vol. 2, 903


The dry chloride, which explodes on heating in air, will rapidly absorb ammonia gas and then explode [1]. The structure of the cation is now known to be a 7-membered ring with only 2 adjacent sulfur atoms. Thiotrithiazyl salts other than the chloride are also explosive [2].

See Thiotrithiazyl nitrate, Thiotrithiazyl perchlorate

See other N-S COMPOUNDS, NON-METAL HALIDES

4034. Cyclopentaazathienium chloride

[88433-73-2]




Banister, A. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1983, 1017

It detonates on percussion and decomposes rapidly above 90°C to give a residue, largely thiotrithiazyl chloride, which explodes above 170°C.

See other N-S COMPOUNDS

4035. Triazidochlorosilane
[67880-25-5]



CIN₉Si

It detonates on percussion and decomposes rapidly above 90°C to give a residue, largely thiotriiazyl chloride, which explodes above 170°C.

See Silicon tetraazide

See other NON-METAL AZIDES

See other N-S COMPOUNDS

4036. Sodium chloride
[7647-14-5]



CINa

HCS 1980, 834

Dichloromaleic anhydride, Urea

See Dichloromaleic anhydride: Sodium chloride, etc.

Lithium

See Lithium: Sodium carbonate, etc.

Nitrogen compounds

Rosinski, M. *et al.*, *Chem. Abs.*, 1980, **93**, 15512

Electrolysis of sodium chloride in presence of nitrogenous compounds to produce chlorine may lead to formation of explosive nitrogen trichloride. Precautions are detailed.

Water

Aleshin, G. Ya., *Chem. Abs.*, 1982, **96**, 220112

The mechanism of explosion of molten salt at 1100°C in accidental contact with water was studied.

See MOLTEN SALTS See other METAL HALIDES

4037. Sodium hypochlorite
[7681-52-9]



CINaO

FPA H52, 1976; HCS 1980, 213, 847, 848 (5, 10, 15% solns, resp.)

1. Brauer, 1963, Vol. 1, 311

2. Sorbe, 1968, 85

The anhydrous solid obtained by desiccation of the pentahydrate will decompose violently on heating or friction [1,2].

Amines

Kirk-Othmer, 1963, Vol. 2, 105

MRH Aniline 3.10/26

Primary aliphatic or aromatic amines react with sodium hypochlorite (or calcium hypochlorite) to form *N*-mono- or di-chloroamines which are explosively unstable, but less so than nitrogen trichloride.

Ammonium salts

1. Anon., *Loss Prev. Bull.*, 1978, (022), 116
2. Anon., *Occ. Safety Health*, 1976, **6**(4), 2
3. Anon., *Sichere Chemiearbeit*, 1996, **48**(10), 114

Contact in the drains of an effluent containing the hypochlorite with one containing ammonium salts and acid led to formation of nitrogen trichloride which decomposed explosively. The effect was reproduced under laboratory conditions [1]. Cleaning a brewery tank with an acidified ammonium sulfate cleaning preparation, then sodium hypochlorite solution without intermediate rinsing, led to nitrogen trichloride formation and a violent explosion [2]. The reservoir of a scrubber system which had previously been employed to scrub ammonia containing off-gas with dilute acid was charged with sodium hypochlorite solution for another process. There was a violent explosion, destroying the reservoir and scrubber assembly. Investigation showed that the reservoir had not been fully cleaned from the previous operation [3].

See Amines, above; Phenylacetonitrile, below

Aziridine

Graefe, A. F. *et al.*, *J. Amer. Chem. Soc.*, 1958, **80**, 3839

Interaction of ethyleneimine with sodium (or other) hypochlorite gives the explosive *N*-chloro compound.

See 1-Chloroaziridine

Carbonised residue

Dokter, T., *J. Haz. Mat.*, 1985, **10**, 85

A saucepan of vegetable stew had been heated too long and had formed a thick carbonised adherent cake. In an attempt to clean the pan, 1 l of domestic bleach was added and the pan was left to heat on an electric hotplate. Again it was left too long and after all the water had evaporated, the residue exploded violently. This was attributed to formation of sodium chlorate during evaporation, and ignition of the overheated chlorate-impregnated carbonised mass.

See Sodium chlorate: Organic matter

Ethanediol

Bickerton, J., *Chem. Brit.*, 1991, **27**(6), 504

Mixtures of aqueous sodium hypochlorite (presumably the 15% available chlorine commercial product) and ethylene glycol were observed to erupt violently after an induction period of 4 to 8 minutes. Caution is advised in view of the use of glycol as a cooling fluid in industrial reactors.

Formic acid

Khristoskova, S., *Chem. Abs.*, 1983, **98**, 95125

Removal of formic acid from industrial waste streams with sodium hypochlorite solution becomes explosive at 55°C.

See other REDOX REACTIONS

Furfuraldehyde

Birchwood, P. J., *Chem. Brit.*, 1985, **21**, 29

In a variation of the usual reaction conditions for oxidising furfuraldehyde to 2-furoic acid with hypochlorite, the aldehyde was added dropwise to a 10% excess of commercial sodium hypochlorite solution at 20–25°C, but without the inclusion of additional sodium hydroxide. When aldehyde addition was almost complete, a violent explosion occurred. Subsequent investigation showed that the pH of the reaction mixture fell progressively with addition of aldehyde, and at pH 8.5 the reaction mixture erupted violently, the temperature increased by 70°C and the pH fell to 2. Similar results were seen with benzaldehyde, but not with thiophene-2-aldehyde.

'Imidate'

Rayner, P., *Safety Digest Univ. Safety Assoc.*, 1991, **40**, 14

A methyl but-3-enylimidate ester hydrochloride, was charged to a tenfold excess of stirred 14% hypochlorite solution cooled in ice. After 50 min the flask was removed to replenish the ice. Shortly after returning the flask to the icebath a violent explosion shattered both flask and icebath. This was attributed to thermal runaway (although available energy is scarcely sufficient to boil the water in the flask). It seems more likely that trichloroamine was generated by the excess hypochlorite, settled when removed from the magnetic stirrer, and detonated from friction when this restarted. The reaction is said to have been performed many times previously without incident.

See Nitrogen trichloride

See other AGITATION INCIDENTS

Methanol

MRH 2.47/10

ICI Mond Div., private comm., 1968

Several explosions involving methanol and sodium hypochlorite were attributed to formation of methyl hypochlorite, especially in presence of acids or other esterification catalyst.

Other reactants

Yoshida, 1980, 132

MRH values calculated for 17 combinations with oxidisable materials are given.

Phenylacetonitrile

Libman, D. D., private comm., 1968

Use of sodium hypochlorite solution to destroy acidified benzyl cyanide residues caused a violent explosion, thought to have been due to formation of nitrogen trichloride.

Photographic developer

Anon., *Environment, Safety & Health Bull.*, 1993, **93**(2), 1

As part of an analytical procedure prior to silver recovery, 10 ml commercial hypochlorite solution (15% available chlorine?) was added to highly alkaline waste developer solution. The redox reaction was violent enough to spray the worker, causing alkali burns.

Sodium ethylenediaminetetracetate, Sodium hydroxide

Schierwater, F.-W., *Sichere Chemiearb.*, 1987, **39**, 35

Operating instructions specified that 50% sodium hydroxide solution, sodium hypochlorite solution and sodium EDTA solution were to be added separately by pumping into an off-gas scrubber unit, when the exotherms would be dissipated slowly in the solution tank. An attempt to simplify the operation by premixing the 3 solutions in a drum before pumping the mixture, led to vigorous foaming decomposition. Mixing the conc. alkali with the bleach caused the oxidant to become heated by the heat of dilution of the former, and oxidation of the EDTA component then proceeded exothermically with decomposition (and evolution of carbon dioxide).

See other GAS EVOLUTION INCIDENTS

Water

Hazard Note HN(76)189, Dept. of Health and Social Security, London, 1976

Two 2.5 l bottles of strong sodium hypochlorite solution (10–14% available chlorine) burst in storage, owing to failure of the cap designed to vent oxygen slowly evolved during storage. This normal tendency may have been accelerated by the unusually hot summer. Vent caps should be checked with full personal protection, and material should be stored at 15–18°C and out of direct sunlight, which accelerates decomposition.

See other METAL HYPOCHLORITES

4038. Sodium chlorite

[7758-19-2]

ClNaO₂

NaClO₂

FPA H109, 1972; *HCS 1980*, 835

Alone

1. Brauer, 1963, Vol. 1, 312
2. Leleu, *Cahiers*, 1974, (74), 137
3. Mellor, 1956, Vol. 2 (Suppl. II, Pt. 1), 573
4. Ullmann, 1986, A6, 500
5. Simoyi, R. H. *et al.*, *Chem. Eng. News*, 1993, **71**(12), 4

The anhydrous salt explodes on impact [1], and decomposes violently at 200°C [2]. The trihydrate is also percussion sensitive [3] though other sources suggest that in clean, grease- and oil-free equipment the anhydrous salt is shockproof [4]. A bottle of the purified anhydrous salt exploded then 'burnt' on opening, (transition to

sodium chloride and oxygen has an enthalpy of 1.2 kJ/g), presumably initiated by friction in the screw cap (most screw caps are combustible). Various contaminants, including iron and ammonium salts, may sensitise sodium chlorite [5].

Acids MRH Several inorganic acids, all 1.17/tr.

Bailar, 1973, Vol. 2, 1413

Under normal conditions, solutions of sodium (and other) chlorites when acidified do not evolve chlorine dioxide in dangerous amounts. However, explosive concentrations may result if acid is dropped onto solid chlorites.

See Oxalic acid, below

Carbon MRH 3.60/9

Leleu, *Cahiers*, 1980, (99), 278

A mixture with powdered charcoal ignited above 60°C.

Ethylene glycol MRH 3.51/17

See Ethylene glycol: Oxidants

Oils MRH Octane 4.60/22, toluene 4.35/33

Karzhenyak, I. G. *et al.*, *Chem. Abs.*, 1981, **95**, 67122

In presence of oils, the chlorite is friction- and shock-sensitive.

Organic matter MRH Cellulose 3.76/24

The Diox Process, Newark, N. J., Wallace and Tiernan, 1949

Intimate mixtures of the solid chlorite with finely divided or fibrous organic matter may be explosive and very sensitive to heat, impact or friction.

Other reactants

Yoshida, 1980, 4–7

MRH Values calculated for no fewer than 79 combinations, largely with oxidisable materials, are given.

Oxalic acid

1. *MCA Case History No. 839*

2. Stull, 1977, 20

A bleach solution was being prepared by mixing solid sodium chlorite, oxalic acid, and water, in that order. As soon as water was added, chlorine dioxide was evolved and later exploded. The lower explosive limit of the latter is 10%, and the mixture is photo- and heat-sensitive [1]. It was calculated that the heat of reaction (1.88 kJ/g of dry mixture) would heat the expected products to an adiabatic temperature approaching 1500°C with an 18-fold increase in pressure in a closed vessel [2].

Phosphorus MRH (Yellow) 6.23/21

Mellor, 1971, Vol. 8, Suppl. 2.2, 645

Red phosphorus and the chlorite react very exothermally in aqueous suspension above 50°C, and there may be a sudden and near-explosive stage in the redox reaction after an induction period.

See *other* INDUCTION PERIOD INCIDENTS, REDOX REACTIONS

Sodium dithionite

Anon., *Chem. Trade J.*, 1953, **132**, 564

Use of a scoop contaminated with sodium dithionite for sodium chlorite caused ignition of the latter. Materials containing sulfur (dithionite, natural rubber gloves) cause decomposition of sodium chlorite and contact should be avoided.

Sulfur-containing materials MRH Sulfur 3.64/15, diethyl sulfide 4.27/28

1. Leleu, *Cahiers*, 1974, (74), 138

2. Weber, J. U. *et al.*, *Rec. Trav. Chim.*, 1986, **105**(3), 99–102

Sodium chlorite reacts very violently with organic compounds of divalent sulfur, or with free sulfur (which may ignite), even in presence of water. Contact of the chlorite with rubber vulcanised with sulfur or a divalent sulfur compound should therefore be avoided [1]. Application of factorial design techniques to experimental planning gave specific conditions for the safe oxidation of organic sulfides to sulfoxides using sodium chlorite or calcium hypochlorite [2].

See Sodium dithionite, above

Zinc

MRH 3.60/58

Karzhenyak, I. G. *et al.*, *Chem. Abs.*, 1981, **95**, 64606

Mixtures containing less than 0.1% of zinc dust or other assorted technical materials are not explosive or combustible on impact or friction.

See other CHLORITE SALTS, METAL OXOHALOGENATES, OXIDANTS

4039. Sodium chlorate

[775-09-9]

ClNaO₃

NaClO₃

(MCA SD-52, 1952); FPA H7, 1972; HCS 1980, 833

1. *The Fire and Explosion at Braehead Container Depot, Renfrew, 1977*, London, HMSO, 1979

2. *The Fire and Explosion at River Road, Barking, Essex, 1980*, London, HSE, 1980

3. Anon., *New Scientist*, 1982, 595 (1325), 892

Following a violent explosion during a fire involving 67 t of drummed sodium chlorate, experiments confirmed that pure sodium chlorate would decompose explosively under intense fire conditions. An appended report by HM Inspector of Explosives detailed six incidents since 1899 involving explosions during fires in stores holding potassium or sodium chlorate [1]. A later incident involving explosive thermal decomposition of 2.5 t of the salt led to a drastic downward revision of the minimum quantity thought to pose this hazard [2]. A third incident in 1982 again involved explosion of drums of sodium chlorate in a warehouse fire [3]. It is difficult to understand this apparent reluctance to gain benefit from the recorded experience of over 80 years.

See Potassium chlorate

Preparative hazard

Kirk-Othmer, 1994, 4th Edn., **9**, 153

There is a risk of generating hydrogen/chlorine/oxygen mixtures during electrolytic preparation from brine. An explosive limits diagram for this ternary system is given.

Agricultural materials

Reimer, B. *et al.*, *Chem. Technik*, 1974, **26**, 447–447 (condensed paper)

The potential for explosive combustion of mixture of sodium chlorate-based herbicides with other combustible agricultural materials was determined. Initiation temperatures and maximum combustion temperatures were measured for mixtures of sodium (or potassium) chlorate with peat, powdered sulfur, sawdust, urotropine (hexamethylenetetramine), thiuram and other formulated materials. With many combinations, maximum temperature increases of 500–1000°C at rates of 400–1200°/s were recorded for 2 g samples.

Alkenes, Potassium osmate

Lloyd, W. D. *et al.*, *Synthesis*, 1972, 610

The hydroxylation of alkenes to diols with potassium osmate–oxidant mixtures has been described, with either hydrogen peroxide or sodium chlorate as the oxidant. The sodium chlorate method is not applicable where the diol is to be distilled from the mixture, because of the danger of explosive oxidation of the product diol by the chlorate.

Aluminium, Rubber

Olson, C. M., *J. Electrochem. Soc.*, 1949, **116**, 34C

The rubber belt of a bucket elevator, fitted with aluminium buckets and used for transporting solid chlorate, jammed during use. Friction from the rotating drive pulley heated and powdered the jammed belt. A violent explosion consumed all the rubber belt and most of the 90 aluminium buckets. Bronze and steel equipment is now installed.

Ammonium salts, or Metals, or Non-metals, or Sulfides

MRH values below references

1. *MCA SD-42*, 1952

2. *MCA Case History No. 2019*

MRH Aluminium 10.71/33, iron 4.35/50, magnesium 10.88/40, manganese 5.06/50, sodium 5.56/55, phosphorus 7.32/25, sulfur 4.27/20

Mixtures of the chlorate with ammonium salts, powdered metals, phosphorus, silicon, sulfur or sulfides are readily ignited and potentially explosive [1]. Residues of ammonium thiosulfate in a bulk road tanker contaminated the consignment of dry sodium chlorate subsequently loaded, and exothermic reaction occurred with gas evolution during several hours. Laboratory tests showed that such a mixture could be made to decompose explosively. A reaction mechanism is suggested.

See other GAS EVOLUTION INCIDENTS

Arsenic trioxide

Ellern, 1968, 51

Ignition may occur on contact.

1,3-Bis(trichloromethyl)benzene

See 1,3-Bis(trichloromethyl)benzene: Oxidants

Cyanides

See METAL CYANIDES (reference 1)

Cyanoborane oligomer

Gyori, B. *et al.*, *J. Organomet. Chem.*, 1984, **262**(1), C7

A mixture exploded violently on mild mechanical agitation.

Diols

See Alkenes, etc., above

Grease

Olson, C. M., *J. Electrochem. Soc.*, 1969, **116**, 34C

The greased bearing of a small grinder exposed to chlorate dust exploded violently during cleaning. Fluorocarbon-based greases and armoured bearings are recommended for chlorate service, with full operator protection during cleaning operations.

See Organic matter, below; *also* Sodium bromate: Grease

Leather

MCA Case History No. 1979

Shoes became contaminated with a chlorate weed-killer solution which dried out in wear. A welding spark later fell into the shoe and the front was blown off.

Nitrobenzene

MRH 4.94/88

Hodgson, J. F., private comm., 1973

The combination is powerfully explosive and has been widely used in recent terrorist activities.

Organic matter

MRH Carbon 4.73/12

1. *MCA SD-42*, 1952

2. Cook, W. H., *Can. J. Res.*, 1933, **8**, 509

3. Biasutti, 1981, 30

4. Anon., *Chem. Haz. in Ind.*, 1987, (7), item 1688

5. Anon., *Chem. Haz. in Ind.*, 1987, (10), item 2666

Mixtures of sodium (or other) chlorate with fibrous or absorbent organic materials (charcoal, flour, shellac, sawdust, sugar) are hazardous. If the chlorate concentration is high, the mixtures may be ignited or caused to explode by static sparks, friction or shock [1]. Even at 10–15% concentration, low relative humidity may allow easy ignition and rapid combustion to occur [2]. A fire originating thus led to destruction of part of an explosives factory [3]. Sparks from a grinder ignited newly-cleaned overalls, which had previously been contaminated with sodium chlorate [4]. It was later confirmed that the overalls had been dry-cleaned in perchloroethylene with added water, rather than being laundered in hot water.

The latter cleaning method would have removed completely the sodium chlorate, which is soluble in water but not in halogenated solvents [5].

See Paper, etc; Wood, both below *See also* OXIDANTS AS HERBICIDES

See other FRICTIONAL INITIATION INCIDENTS, STATIC INITIATION INCIDENTS

Osmium

Rogers, D. B. *et al.*, *Inorg. Synth.*, 1972, **13**, 141

During the preparation of osmium dioxide, the ampoule containing the reaction mixture must be cooled during sealing operations to prevent violent reaction occurring. Subsequent heating, first to 300 and then 600°C must also be effected slowly.

Other reactants

Yoshida, 1980, 70–71

MRH values calculated for 33 combinations with oxidisable materials are given.

Paint, Polythene

Lamb, J. A., *Chem. Brit.*, 1979, **15**, 125–126

A violent explosion occurred when the lid was levered off an old paint tin used to contain a polythene bag of sodium chlorate. Subsequent tests showed that low energy input (0.2 J) would cause explosion of unconfined chlorate on a rusty steel surface. The presence of contaminants (sawdust, copper acetate, paint flakes or shredded polythene) increases the sensitivity, and a mixture with paint flakes and polythene caused 100 mg portions to explode under 0.2 J impact in 24 out of 100 attempts.

Paper, Static electricity

1. Ewing, O. R., *Chem. Eng. News*, 1952, **30**, 3210

2. Grimmett, R. E. R., *New Zealand J. Agric.*, 1938, **57**, 224–225

3. Biasutti, 1981, 92

Paper impregnated with sodium chlorate and dried can be ignited by static sparks, but not by friction or impact. Paper bags or card cartons are unsuitable packing materials [1]. A previous incident involving paper sacks which had formerly contained mixed sodium and calcium chlorates had been noted [2]. The tendency of sodium chlorate to deliquesce or effloresce, depending on atmospheric temperature and humidity may lead, if the solid is packed in pervious containers, to the appearance of very fine crystals on the outside of a paper bag or cardboard box, etc. These fine crystals of high specific surface area are very sensitive to friction and initiation of ignition [3].

See other FRICTIONAL INITIATION INCIDENTS, STATIC INITIATION INCIDENTS

Phosphorus

MRH (Yellow) 7.32/25

Anon., *Angew. Chem. (Nachr.)*, 1957, **5**, 78

A mixture of red phosphorus and sodium chlorate exploded violently.

Sodium phosphinate

MRH 2.89/55

See Sodium phosphinate: Oxidants

Sulfuric acid

MRH 1.30/tr.

1. Anon., *ABCM Quart. Safety Summ.*, 1944, **15**, 3
2. *MCA Case History No. 282*

Erroneous addition of conc. sulfuric acid to sodium chlorate instead of sodium chloride caused an explosion owing to formation of chlorine dioxide [1]. Accidental contact of 93% acid on clothing previously splashed with sodium chlorate caused immediate ignition [2].

Titanium

See Titanium: Oxidants

Titanium: Sodium chlorate, etc.

Triethylene glycol, Wood

Anon., *CISHC Chem. Safety Summ.*, 1980, **51**, 112

A wooden pallet which ignited and burned fiercely was found to be contaminated with both the glycol and sodium chlorate to a level of 27%.

Wood

1. Anon., *ABCM Quart. Safety Summ.*, 1947, **18**, 25
2. Anon., *BCISC Quart. Safety Summ.*, 1967, **38**, 42
3. Kletz, T. A., *J. Loss Prev.Proc.Ind.*, 1989, **2**, 117

Various fires and explosions caused by use of wooden containers with chlorates, and precautions necessary during handling and storage, are discussed [1,2]. A wooden pallet burst into flames as it was dragged across ground contaminated with sodium chlorate [3].

See other FRICTIONAL INITIATION INCIDENTS

See Organic matter, above

See other METAL CHLORATES, OXIDANTS

4040. Sodium perchlorate

[7601-89-0]

ClNaO₄

NaClO₄

Hydrazine

Rosolovskii, V. Ya., *Chem. Abs.*, 1969, **70**, 53524

Interaction gives only a 1:1 complex which explodes on grinding, but dissociates uneventfully on heating at atmospheric or reduced pressure. The lithium salt is similar.

Other reactants

Yoshida, 1980, 82

MRH values calculated for 18 combinations with oxidisable materials are given.

Water

Moureu, H. *et al.*, *Chem. Abs.*, 1951, **45**, 5929h

During concentration of an aqueous solution, an explosion occurred. Involvement of an unsuspected impurity seems probable.

ClO₂

Water-soluble fuels MRH values below reference

Annikov, V. E. *et al.*, *Chem. Abs.*, 1983, **99**, 73190

MRH Acetone 4.89/18, ethanol 4.73/18, ethylene glycol 4.35/26

The detonation and combustion limits of mixtures of sodium perchlorate, water and ethylene glycol, glycerol, 1,3-butylene glycol, 2,3-butylene glycol, formamide, dimethylformamide, ethanolamine, diaminoethane, acetone, urea and galactose have been studied.

See other METAL PERCHLORATES, OXIDANTS

4041. Antimony(III) chloride oxide

[7791-08-4]

ClOSb

O:SbCl

Bromine trifluoride

See Bromine trifluoride: Antimony(III) chloride oxide

See related METAL HALIDES, METAL OXIDES

4042. Chlorine dioxide

[10049-04-4]

ClO₂

ClO₂

HCS 1980, 289

1. Sidgwick, 1950, 1203
2. Mellor, 1941, Vol. 2, 288
3. Stedman, R. F., *Chem. Eng. News*, 1951, **29**, 5030
4. Cameron, A. E., *Chem. Eng. News*, 1951, **29**, 3196
5. Fawcett, H. H., *Chem. Eng. News*, 1951, **29**, 4459
6. McHale, E. T. *et al.*, *J. Phys. Chem.*, 1968, **72**, 1849
7. Jansen, M., *et al.*, *Angew. Chemie (Int.)*, 1991, **30**(11), 1510
8. *Hazards of Chlorine Dioxide*, New York, Natl. Board of Fire Underwriters, 1950; *Chem. Eng. News*, 1950, **28**, 611
9. Derby, R. I. *et al.*, *Inorg. Synth.*, 1954, **4**, 152
10. Bielz, S., *Ger. Offen.* 2 917 132, 1980
11. Cowley, G., *Loss Prev. Bull.*, 1993, (113), 1
12. Simpson, G. D. *et al.*, *Ind. Water Treat.*, 1995, **27**(5), 48

Chlorine dioxide is considerably endothermic (ΔH_f° (g) +103.3 kJ/mol, 1.53 kJ/g) and of limited stability. It is a powerful oxidant and explodes violently on the slightest provocation as gas or liquid [1]. It is initiated by contact with several materials (below), on heating rapidly to 100°C or on sparking [2], or by impact as solid at -100°C [3]. A small sample exploded during vacuum distillation at below -50°C [4], and it was stated that decomposition by sparking begins to become hazardous at concentrations of 7-8% in air [3], and that at 10% concentration in air (0.1 bar partial pressure) explosion may occur from any source of initiation energy, such as sunlight, heat or electrostatic discharge [5]. A kinetic study of the decomposition shows that it is explosive above 45°C even in absence of light, and subject to long induction periods due to formation of intermediate dichlorine

trioxide. UV irradiation greatly sensitises the dioxide to explosion [6]. The solid (A dimer) can be relatively safely handled below -40°C and the gas at pressures below 50 mbar [7]. A guide on fire and explosion hazards in industrial use of chlorine dioxide is available [8], and preparative precautions have been detailed [9]. An improved and safer method for continuous production of chlorine dioxide is claimed [10]. A thorough review has been written, detailing numerous incidents, of hazards attending industrial preparation and use of chlorine dioxide (now much used as a low chlorine bleach). Liquid ClO_2 can separate from aqueous solutions >60 g/l, it is exceedingly shock sensitive. A partial pressure of 130 mbar in air is thought entirely safe, as are aqueous solutions (but not necessarily the head space above them) [11]. The safe use of chlorine dioxide and sodium chlorite has been reviewed [12].

See other ENDOTHERMIC COMPOUNDS, IRRADIATION DECOMPOSITION INCIDENTS, STATIC INITIATION INCIDENTS

Carbon monoxide MRH 5.35/45

Mellor, 1941, Vol. 2, 288
Explosion on mixing.

Fluoramines

Lawless, 1968, 171
Interaction with difluoramine or trifluoramine in the gas phase is explosive.

Hydrocarbons

MRH Hexane, 6.61/12, Toluene 6.65/13

Leleu, *Cahiers*, 1979, (95), 297
Mixtures with butadiene, ethane, ethylene, methane or propane always explode spontaneously.

Hydrogen

Mellor, 1941, Vol. 2, 288
Near-stoichiometric mixtures detonate on sparking, or on contact with platinum sponge.

Mercury

Mellor, 1941, Vol. 2, 288
The gas explodes on shaking with mercury.

Non-metals

Mellor 1941, Vol. 2, 289; 1956, Vol. 2, Suppl. 1, 532
Phosphorus, sulfur, sugar or combustible materials ignite on contact and may cause explosion.

Other reactants

Yoshida, 1980, 268
MRH values calculated for 19 combinations, largely with oxidisable materials, are given.

Phosphorus pentachloride MRH 1.55/71
See Phosphorus pentachloride: Chlorine dioxide

Potassium hydroxide MRH 1.92/37
Mellor, 1941, Vol. 2, 289
The liquid or gaseous oxide will explode in contact with solid potassium hydroxide or its conc. solution.

Sodium nitrite, Acid
Briggs, T. S., *J. Chem. Educ.*, 1991, **68**(11), 938
An experiment is described which purports to show the detonation of a mixture of chlorine dioxide (described as a stable gas!) and nitric oxide formed from the above reagents. It seems likely it demonstrates detonation of chlorine dioxide, itself, initiated by either the nitrite or nitric oxide.
See other HALOGEN OXIDES, OXIDANTS

4043. Thallium(I) chlorite **ClO₂Tl**
[40898-91-7] **TlClO₂**

Bailar, 1973, Vol. 1, 1167
It is detonable by shock.
See other CHLORITE SALTS, HEAVY METAL DERIVATIVES

4044. Chlorine trioxide **ClO₃**
[13932-10-0] **ClO₃**

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 498
The dimeric form is formulated as chloryl perchlorate.
See Chloryl perchlorate
See other HALOGEN OXIDES

4045. Rhenium chloride trioxide (Perrhenyl chloride) **ClO₃Re**
[42246-25-3] **ClReO₃**

Preparative hazard
See Oxygen: Trirhenium nonachloride
See related METAL HALIDES, METAL OXIDES

4046. Antimony(III) oxide perchlorate

[]

ClO₅Sb

O:SbClO₄

Mellor, 1956, Vol. 2, Suppl. 1, 613

Decomposes with decrepitation above 60°C.

See related METAL PERCHLORATES

4047. Chlorine

[7782-50-3]

Cl₂

Cl₂

(MCA SD-80, 1970); NSC 207, 1982; FPA H39, 1975; HCS 1980, 288 (cyl);
RSC Lab. Hazard Data Sheet No. 30, 1984

1. Dokter, T., *J. Haz. Mat.*, 1985, **10**, 73–87

2. Anon., *Sichere Chemiearbeit*, 1993, **45**(1), 9

The fire and explosion hazards of chlorine-containing systems have been reviewed [1]. A rail tanker of chlorine was being emptied to a chlorine gas distribution network at 7 bar, when air supply to a pneumatic valve failed, closing the system down. After repair, valve operation caused ignition of the pipework in a chlorine–iron fire, resulting in a leak of chlorine and almost complete destruction of a valve. Ignition is believed to have occurred because the electrolytic chlorine in the system contained 0.7% of hydrogen. Condensation of gaseous chlorine trapped in the pipework (at 0°C) during the shutdown produced a gas phase enriched with hydrogen to within explosive limits and easily ignited, in turn igniting the steelwork [2].

See Hydrogen, below; Steel, below

See CHLORINE-CONTAINING SYSTEMS

See other IGNITION SOURCES

See also REPAIR AND MAINTENANCE

Preparative hazard (electrolysis)

1. Anon., *Fire Prevention*, 1986, (189), 43.

2. Kucinski, R. E. *et al.*, *Mod. Chlor-Alkali Technol.*, 1995, 6, 89

A fire and explosions caused by hydrogen from a chloralkali plant seeping through a defective non-return valve, then spontaneously igniting is reported [1]. A consideration of the risks of producing hydrogen chlorine mixtures, related to the design of plant in question, claims the ion-exchange membrane plant to be safest [2].

See Sodium chloride: Nitrogen compounds

See Ammonia, below

Acetylene

MRH 8.70/99+

See Acetylene: Halogens

Air, Ethylene

See Ethylene: Air, Chlorine

Alcohols

See *tert*-Butanol, below

See also HYPOHALITES

Aluminium

MRH 5.23/20

Ann. Rep. Chief Insp. Factories, 1953 (Cmd. 9330), 171, HMSO

Corrosive failure of a vaporiser used in manufacture of aluminium chloride caused liquid chlorine to contact molten aluminium. A series of explosions occurred.

See Aluminium: Halogens, or: Oxidants

See Metals, below

See other CORROSION INCIDENTS

Amidosulfuric acid

MCA Guide, 1972, 306

Chlorination of aqueous sulfamic acid led to an explosion from formation of nitrogen trichloride.

See Nitrogen compounds, below (reference 6)

Ammonia

Khattak, M. A., *Engineering Horizons* (Pakistan), 1991, (Aug.), 33

Chlorine reacts with ammonia and compounds to form the treacherously explosive nitrogen trichloride. A change was made to ammonia as refrigerant in the production of liquid chlorine; some months later minor explosions when transferring the chlorine culminated in several fatalities when the delivery pipe, in which the less volatile trichloride had presumably concentrated by partial evaporation of chlorine, shattered. Further explosions were experienced while decontaminating the plant.

See Preparative hazard, above

See Nitrogen trichloride

Antimony trichloride, Tetramethylsilane

Bush, R. P., Br. Pat. 1 388 991, 1975

Antimony trichloride-catalysed chlorination of the silane to chlorotrimethylsilane in absence of diluent was explosive at 100°C, but controllable at below 30°C.

See other HALOGENATION INCIDENTS

N-Arylsulfonamides

Johnson, C. R. *et al.*, *J. Org. Chem.*, 1983, **48**, 1

Oxidation by chlorine to the arylimidosulfonyl chlorides is sometimes violent, and *tert*-butyl hypochlorite is a milder reagent.

Aziridine

MRH 36.98/99+ (?3.6)

See Aziridine: Chlorinating agents

Benzene

MRH 2.09/27

1. *491M*, 1975, 102

2. Yakushina, E. P. *et al.*, *Chem. Abs.*, 1981, **94**, 3389

A mixture in the vapour phase exploded when exposed to light [1]. Benzene–chlorine mixtures, though combustible, represent a low explosion hazard [2].

See Hydrocarbons, below

Bis(2,4-dinitrophenyl) disulfide

See Bis(2,4-dinitrophenyl) disulfide: Chlorine

Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

tert-Butanol

1. Bradshaw, C. P. C. *et al.*, *Proc. Chem. Soc.*, 1963, 213

2. Mintz, M. J. *et al.*, *Org. Synth.*, 1969, **49**, 9

Rate of admission of chlorine into the alcohol during preparation of *tert*-butyl hypochlorite must be regulated to keep the temperature below 20°C to prevent explosion [1]. A safer and simpler preparation uses hypochlorite solution in place of chlorine [2].

See *tert*-Butyl hypochlorite

Butyl rubber, Naphtha

Komarovskii, N. A. *et al.*, *Chem. Abs.*, 1979, **91**, 158791

Chlorination of butyl rubber in naphtha with chlorine–nitrogen mixtures may lead to explosion if N contents below 77% or Cl contents above 16% are used.

Caesium oxide

See Caesium oxide: Halogens

Carbon disulfide

MRH 1.17/99+

MCA Case History No. 971

When liquid chlorine was added to carbon disulfide in an iron cylinder, an explosion occurred, owing to the iron-catalysed chlorination of carbon disulfide to carbon tetrachloride. The operation had been done previously in glassware without incident.

See *other* CATALYTIC IMPURITY INCIDENTS

Carbon disulfide, Sodium *aci*-nitromethanide

See Sodium *aci*-nitromethanide: Carbon disulfide, etc.

Chlorinated pyridine, Iron powder

1. Anon., *SciQuest*, 1979, **52**(7), 26–27

2. De Haven, E. S., *Plant/Oper. Progr.*, 1990, **9**(2), 131

An explosion occurred during the preparation of iron(III) chloride from iron powder and chlorine gas in a chlorinated pyridine solvent. This was attributed to formation of iron(II) chloride, its interaction with the solvent to give iron(III) chloride, then reduction of the latter by iron to iron(II) chloride. The exotherm and increasing evolution of hydrogen chloride caused the reactor to fail [1].

Cl₂

Another account of what seems to be the same incident attributes the runaway to Friedel Crafts type polymerisation of trichloromethylpyridines largely present in the reactor [2].

See other GAS EVOLUTION INCIDENTS

2-Chloroalkyl aryl sulfides, Lithium perchlorate

Zefirov, N. S. *et al.*, *Chem. Abs.*, 1986, **104**, 168050

Chlorine scission of 2-chloroheptyl 4-nitrophenyl sulfide or of the 2,4-dinitro analogue in presence of lithium perchlorate formed the explosive 2-chloroheptyl perchlorate in 5–7% yield.

See other ALKYL PERCHLORATES, HALOGENATION INCIDENTS

Chloromethane

Dokter, T., *J. Haz. Mat.*, 1985, **10**, 73–87

The autoignition temperature of chloromethane in presence of chlorine is 215°C, some 400° lower than in presence of air.

3-Chloropropyne

Anon., *Loss Prev. Bull.*, 1974, (001), 10

A vigorous explosion during chlorination of 3-chloropropyne in benzene at 0°C over 4 h was attributed to presence of excess chlorine arising from the slow rate of reaction at low temperature.

See Acetylene: Halogens, *also* Hydrocarbons, below (reference 10)

See other HALOGENATION INCIDENTS

Cobalt(II) chloride, Methanol

1. Vlcek, A. A., *Inorg. Chem.*, 1967, **6**, 1425–1427

2. Gillard, R. D. *et al.*, *Chem. & Ind.*, 1973, 77

3. Gillard, R. D., private comm., 1973

During the preparation of *cis*-dichlorobis(2,2'-bipyridyl)cobalt(III) chloride by repeating a published procedure [1], passage of chlorine into an ice-cold solution of cobalt chloride, bipyridyl and lithium chloride in methanol soon caused an explosion followed by ignition of the methanol inside the reaction vessel. The fire could not be extinguished with carbon dioxide, but went out when the flow of chlorine stopped [2]. This is consistent with the formation and self ignition of methyl hypochlorite in the system at a slightly elevated temperature. An alternative route to the cobalt complex not involving chlorine is available[3].

See Methanol, *also* Methanol, Tetrapyridinecobalt(II)chloride, both below

See other HALOGENATION INCIDENTS

Dibutyl phthalate

Statesir, W. A., *Chem. Eng. Progr.*, 1973, **69**(4), 54

A mixture of the ester and liquid chlorine confined in a stainless steel bomb reacted explosively at 118°C.

1,2-Dichloroethane

Steblen, A. V., *Chem. Abs.*, 1981, **95**, 67123

Although some mixtures of the two components will burn, even that with 34% of haloalkane leads only to a 2-fold pressure increase. Such mixtures are comparatively safe.

Dichloro(methyl)arsine

Dillon, K. B. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1976, 1479

In the attempted preparation of tetrachloromethylarsinane (MeAsCl₄) by interaction of chlorine with the arsine while slowly warming from -196°C, several sealed ampoules exploded at well below 0°C, probably owing to liberation of chloromethane.

See other GAS EVOLUTION INCIDENTS

Diethyl ether

MRH 12.97/21

1. Harrison, G. E., private comm., 1966
2. Unpublished observation, 1949

Chlorine caused ignition of ether on contact [1]. Exposure of an ethereal solution of chlorine to daylight on removal from a fume cupboard caused a mild photocatalysed explosion [2].

See Bromine: Diethyl ether, or: Tetrahydrofuran

See other IRRADIATION DECOMPOSITION INCIDENTS

Diethylzinc

MRH 2.22/64

Weast, 1979, C-721

Ignition occurs on contact.

Dimethylformamide

Woltornist, A., *Chem. Eng. News*, 1983, **61**(6), 4

After a thermal runaway reaction during chlorination in DMF solution, investigation revealed that saturated solutions of chlorine in DMF are hazardous, and will self-heat and erupt under either adiabatic or non-adiabatic conditions. Principal products are tetramethylformamidium chloride and carbon dioxide, with dimethylammonium chloride and carbon monoxide in small amounts. A detailed account of the mechanism is to be published.

Dimethyl phosphoramidate

1. Block, H. D. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1971, **10**, 491
2. Zwierak, A. *et al.*, *Tetrahedron*, 1970, **26**, 3521
3. Walker, B. J. *et al.*, *Chem. Brit.*, 1979, **15**, 66
4. Walker, B. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1980, 462

In a 1.5 g mol preparation of dimethyl *N,N*-dichlorophosphoramidate by chlorination of the ester, a violent explosion occurred during the period of stirring after the reaction. This did not occur on 0.5 g mol scale, and the longer reaction time may have led to liberation of explosive nitrogen trichloride. Suggested precautions include a working scale below 0.2 g mol and short reaction time. The diethyl homologue appears more stable and may be prepared on a kg scale[1]. Another incident involving explosion while following the same preparative procedure [2] was reported some years later [3,4].

Dioxan

1. Leclerc-Battin, F. *et al.*, *J. Loss Prevention*, 1991, **4**(3), 170
2. Elaissi, A. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries Vol II* (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 349, Amsterdam, Elsevier, 1995

A study of autoignition of dioxan vapour and chlorine, in the gas phase at rather below 1 atm. Ignition was once observed at 25°C in daylight. In the dark it occurred from 100°C. The maximum pressure rise observed was 26 MPa per sec [1]. Detonation properties of gaseous mixtures of chlorine and dioxane, sometimes diluted with argon, are studied in detail [2]. The combination is more sensitive than most fuel/air mixtures.

Dioxygen difluoride

See Dioxygen difluoride: Various materials

Disilyl oxide

Bailar, 1973, Vol. 1, 1377
Interaction is explosive.

4,4'-Dithiodimorpholine

See 4-Morpholinesulfonyl chloride

Fluorine

See Fluorine: Halogens

Fluoromethylsilane, Dichloromethane

Bürger, H. *et al.*, *Organometallics*, 1993, **12**(12), 4930
Chlorination of the silane in dichloromethane proved uncontrollable even at -100°C. Explosions with decomposition and generation of carbon occurred.

Glycerol

Statesir, W. A., *Chem. Eng. Progr.*, 1973, **69**(4), 54
A mixture of liquid chlorine and glycerol confined in a stainless steel bomb exploded at 70–80°C.

Hexachlorodisilane

Martin, G., *J. Chem. Soc.*, 1914, **105**, 2859
Hexachlorodisilane vapour ignited in chlorine above 300°C; violent explosions sometimes occurred.

Halocarbons

1. Dokter, T., *J. Haz. Mat.*, 1985, **10**, 75–87
2. *See entry* CHLORINE-CONTAINING SYSTEMS

The auto-ignition temperatures of various halogenated hydrocarbons in presence of chlorine are considerably below the corresponding values in air. Examples are (in °C) chloromethane 215 (618 in air); dichloromethane 262 (556); 1,2-dichloropropane 180 (555°C). Flammability limits are usually wider in chlorine

than in air. Gaseous mixtures of chloromethane and chlorine are detonable in the range 12.5–55% [1]. Liquid mixtures of dichloromethane and chlorine may well also be detonable [2].

Hydrocarbons, MRH Benzene 2.09/27, acetylene 8.70/99+
(Inorganics)

1. Mellor, 1956, Vol. 2, Suppl. 1, 380
2. Eisenlohr, D. H., US Pat. 2 989 571, 1961
3. von Schwartz, 1918, 142, 321
4. von Schwartz, 1918, 324
5. Mamadaliev, Y. G. *et al.*, *Chem. Abs.*, 1937, **31**, 8502⁵
6. Brooks, B. T., *Ind. Eng. Chem.*, 1924, **17**, 752
7. Johnson, J. H., *Hydrocarbon Proc. Petr. Ref.*, 1963, **42**(2), 174
8. Statesir, W. A., *Chem. Eng. Progr.*, 1973, **69**(4), 52–54
9. de Oliveria, D. B., *Hydrocarbon Proc.*, 1973, **53**(3), 112–126
10. Kilby, J. L., *Chem. Eng. Progr.*, 1968, **64**(6), 52
11. Lawrence, W. W. *et al.*, *Ind. Eng. Chem., Proc. Des. Develop.*, 1970, **9**, 47–49
12. Kokochashvili, V. I. *et al.*, *Chem. Abs.*, 1976, **84**, 20030
13. Rozlovskii, A. I. *et al.*, *Chem. Abs.*, 1976, **84**, 49418
14. Jenkins, P. A., unpublished information, 1974

Interaction of chlorine with methane is explosive at ambient temperature over yellow mercury oxide [1], and mixtures containing above 20 vol% of chlorine are explosive [2]. Mixtures of acetylene and chlorine may explode on initiation by sunlight, other UV source, or high temperatures, sometimes very violently [3]. Mixtures with ethylene explode on initiation by sunlight, etc., or over mercury, mercury oxide or silver oxide at ambient temperature, or over lead oxide at 100°C [1,4]. Interaction with ethane over activated carbon at 350°C has caused explosions, but added carbon dioxide reduces the risk [5]. Accidental introduction of gasoline into a cylinder of liquid chlorine caused a slow exothermic reaction which accelerated to detonation. This effect was verified [6]. Injection of liquid chlorine into a naphtha–sodium hydroxide mixture (to generate hypochlorite in situ) caused a violent explosion. Several other incidents involving violent reactions of saturated hydrocarbons with chlorine were noted [7].

In a review of incidents involving explosive reactivity of liquid chlorine with various organic auxiliary materials, two involved hydrocarbons. A polypropylene filter element fabricated with zinc oxide filler reacted explosively, rupturing the steel case previously tested to over 300 bar. Zinc chloride derived from the oxide may have initiated the runaway reaction. Hydrocarbon-based diaphragm pump oils or metal-drawing waxes were violently or explosively reactive [8]. A violent explosion in a wax chlorination plant may have involved unplanned contact of liquid chlorine with wax or chlorinated wax residues in a steel trap. Corrosion products in the trap may have catalysed the runaway reaction, but hydrogen (also liberated by corrosion in the trap) may also have been involved [9].

During maintenance work, simultaneous release of chlorine and acetylene from two plants into a common vent line leading to a flare caused an explosion in the line [10]. The violent interaction of liquid chlorine injected into ethane at 80°C/10 bar

Cl₂

becomes very violent if ethylene is also present [11]. The relationship between critical pressure and composition for self-ignition of chlorine–propane mixtures at 300°C was studied, and the tendency is minimal for 60:40 mixtures. Combustion is explosive under some conditions [12]. Precautions to prevent explosions during chlorination of solid paraffin hydrocarbons are detailed [13]. In the continuous chlorination of polyisobutene at below 100°C in absence of air, changes in conditions (increase in chlorine flow, decrease in polymer feed) leading to over-chlorination caused an exotherm to 130°C and ignition [14].

See Benzene, above; Methane, Oxygen, etc., or Synthetic rubber, both below

See also TURPENTINE: halogens

Hydrocarbons, Lewis acids

Howard, W. B., *Loss Prev.*, 1973, **7**, 78

During chlorination of hydrocarbons with Lewis acid catalysis, the catalyst must be premixed with the hydrocarbon before admission of chlorine. Addition of catalyst to the chlorine–hydrocarbon mixture is very hazardous, causing instantaneous release of large volumes of hydrogen chloride.

See Iron(III) chloride, Monomers, below

See other GAS EVOLUTION INCIDENTS, HALOGENATION INCIDENTS

Hydrochloric acid, Dinitroanilines

See Hydrogen chloride: Chlorine, Dinitroanilines

Hydrogen

MRH 2.55/3

1. Mellor, 1956, Vol. 2, Suppl. 1, 373–375
2. Weissweiler, A., *Z. Elektrochem.*, 1936, **42**, 499
3. Eichelberger, W. C. *et al.*, *Chem. Eng. Progr.*, 1961, **57**(8), 94
4. Wood, J. L., *Loss Prev.*, 1969, **3**, 45–47
5. Stephens, T. J. R. *et al.*, *Paper 62B*, 65th AIChE Meeting, New York, 1972
6. de Oliveria, D. B., *Hydrocarbon Proc.*, 1973, **52**(3), 112–126
7. Antonov, V. N. *et al.*, *Chem. Abs.*, 1974, **81**, 172184
8. Frolov, U. E., *Chem. Abs.*, 1977, **87**, 119999
9. Anon., *Fire Prev.*, 1986, (189), 43

Combination of the elements may be explosive over a wide range of physical conditions, with initiation by sparks, radiant energy or catalysis, e.g. by yellow mercuric oxide at ambient temperature [1]. There is a narrow range of concentrations in which the mixture is supersensitive to initiation [2]. Explosion–detonation phenomena in chlorine production cells have been investigated [3,4]. The explosive limits of the mixture vary with the container shape and method of initiation, but are usually within the range 5–89% of hydrogen by volume [1]. After an explosion in a chlorine distillate receiver where hydrogen had been produced by corrosion, no initiation source could be identified or reasonably postulated following a thorough investigation [5]. Several other hydrogen–chlorine explosions without identifiable ignition sources are also mentioned. Hydrogen may have been involved in a severe wax chlorinator explosion [6]. Pressure increase during explosive combustion of mixtures within the critical concentration region of 5–15 vol% of hydrogen was studied, and found less than anticipated [7]. Critical self-ignition temperatures are

397–284°C for mixtures containing 8–18% of hydrogen [8]. A faulty non-return valve allowed hydrogen to leak into a fibreglass-reinforced pipeline containing chlorine, and spontaneous ignition and explosion occurred [9].

Hydrogen, Nitrogen trichloride

See Nitrogen trichloride: Chlorine, Hydrogen

Hydrogen, Other gases

Munke, K., *Chem. Technik*, 1974, **26**, 292–295

Available data on explosibility of chlorine and hydrogen in admixture with air, hydrogen chloride, oxygen or inert gases is discussed and presented as triangular or rectangular diagrams.

See Sodium chlorate: Preparative hazard

Hydrogen(?), Sulfuric acid

Tabata, Y. *et al.*, *Chem. Abs.*, 1984, **100**, 90601

Chlorine gas, produced by electrolysis of brine, was dried by passage through a mist of sulfuric acid in a drying tower, where static charge was generated. An explosion in the tower, (presumably involving hydrogen as fuel) was attributed to initiation by static spark.

See Hydrogen: Oxygen, Sulphuric acid

See other STATIC INITIATION INCIDENTS

Iron(III) chloride, Monomers

1. *MCA Case History No. 2115*

2. *MCA Case History No. 2147*

During chlorination of styrene in carbon tetrachloride at 50°C, a violent reaction occurred when some 10% of the chlorine gas had been fed in. Laboratory examination showed that the eruption was caused by a rapid decomposition reaction catalysed by ferric chloride [1]. Various aromatic monomers decomposed in this way when treated with gaseous chlorine or hydrogen chloride (either neat, or in a solvent) in the presence of steel or iron(III) chloride. Exotherms of 90°C (in 50% solvent) to 200°C (no solvent) were observed, and much gas and polymeric residue was forcibly ejected.

See Hydrocarbons, Lewis acids, above

Mercury(II) oxide

See Mercury(II) oxide: Chlorine

Metal acetylides and carbides

The mono- and di-alkali metal acetylides, copper acetylides, iron, uranium and zirconium carbides all ignite in chlorine, the former often at ambient temperature.

See Caesium acetylide: Halogens

Dicopper(I) acetylide: Halogens

Iron carbide: Halogens

Lithium acetylide: Halogens
 Monocaesium acetylide
 Monorubidium acetylide
 Strontium acetylide: Halogens
 Uranium dicarbide: Halogens
 Zirconium dicarbide: Halogens
See Monolithium acetylide–ammonia: Gases

Metal hydrides MRH Calcium hydride 5.35/37, Potassium hydride 4.98/52
 Mellor, 1941, Vol. 2, 483; Vol. 3, 73
 Potassium, sodium and copper hydrides all ignite in chlorine at ambient temperatures.

Metal phosphides MRH Magnesium phosphide 4.04/39
See Copper(II) phosphide: Oxidants

Metals MRH values below references
 1. Mellor, 1941, Vol. 2, 92, 95; 1956, Vol. 2, Suppl. 1, 380, 469; 1941, Vol. 3, 638; 1940, Vol. 4, 267, 480; 1941, Vol. 7, 208, 260, 436; 1941, Vol. 9, 379, 626, 849; 1942, Vol. 12, 312; Vol. 15, 146
 2. Hanson, B. H., *Process Eng.*, 1975, (2), 77
 3. Davies, D. J., *Process Eng.*, 1975, (2), 77
 4. Anon., *Loss Prev. Bull.*, 1975, (006), 1
 5. Anon., *Loss Prev. Bull.*, 1991, (097), 9

MRH Aluminium 5.23/20, bismuth 1.21/66, calcium 7.15/36, copper 1.55/47, germanium 2.51/34, magnesium 6.78/25, manganese 3.85/44, nickel 2.34/45, potassium 5.81/53, sodium 6.98/40, tin 2.47/45, vanadium 3.72/42, zinc 3.05/48

Tin ignites in liquid chlorine at —34°C, aluminium powder in the gas at —20°C, while vanadium powder explodes on contact at 0°C with the pressurised liquid. A solution in heptane ignites with powdered copper well below 0°C. Aluminium, brass foil, calcium powder, copper foil, iron wire, manganese powder, and potassium all ignite in the dry gas at ambient temperature, as do powdered antimony, bismuth and germanium sprinkled into the gas, while magnesium, sodium and zinc ignite in the moist gas. Thorium, tin and uranium ignite and incandesce on warming (uranium to 150°C), powdered nickel burns at 600°C, while mercury at 200—300°C ignites in a stream of chlorine. Aluminium—titanium alloys also ignite on heating in chlorine, and niobium ignites on gentle warming [1]. Unlike most other metals, titanium is not suitable for components in contact with *dry* gaseous or liquid chlorine, as ignition may occur [2]. Minimum water content to prevent attack may be from 0.015 to 1.5%, depending on conditions [3]. A statement on ignition of titanium in wet chlorine is incorrect [4]. Titanium washers were fitted to a liquid chlorine line due to mis-labelling. Within an hour of exposure to chlorine they burnt through in a 'puff of smoke' followed by a chlorine leak.

See Aluminium, above; Steel, below; *also* Beryllium: Halogens

Methane, Oxygen

Mal'tseva, A. S. *et al.*, *Chem. Abs.*, 1982, **96**, 90922

Ignition in an oxychlorination reactor was attributed to the presence of free space above the packed reaction zone. Other hazards are discussed.

Methanol

Trans. US Natl. Safety Congr., 1937, 273

Passage of chlorine through cold recovered methanol (but not fresh methanol) led to a mild explosion and ignition, formation of methyl hypochlorite apparently being catalysed by an impurity.

See Cobalt(II) chloride, Methanol, above; Methanol, Tetrapyrindinecobalt(II) chloride, below

See Bromine: Alcohols

See other CATALYTIC IMPURITY INCIDENTS

Methanol, Tetrapyrindinecobalt(II) chloride

Glerup, J. *et al.*, *Acta Chem. Scand., Ser. A*, 1978, **A32**, 673

In preparation of dichlorotetrapyrindinecobalt(III) chloride by oxidation of the Co(II) salt in methanol solution with gaseous chlorine, severe explosions (formation of methyl hypochlorite) have occurred. Use of liquid chlorine at -40° See Cobalt(II) chloride, Methanol, above

Nitrogen compounds

MRH values below references

1. Mellor, 1941, Vol. 2, 95; 1940, Vol. 8, 99, 288, 313, 607
2. Bowman, W. R. *et al.*, *Chem. & Ind.*, 1963, 979
3. Bainbridge, E. G., *Chem. & Ind.*, 1963, 1350
4. Folkers, K. H. *et al.*, *J. Amer. Chem. Soc.*, 1941, **63**, 3530
5. King, J. F. *et al.*, *Synthesis*, 1980, 285
6. Schierwater, F.-W., *Major Loss Prevention*, 1971, 49
7. Sorbe, 1968, 120
8. Kirk-Othmer, 1964, Vol. 5, 2
9. Dokter, T., *J. Haz. Mat.*, 1985, **10**, 207–224
10. Nothe, M. *et al.*, *Inorg. Chem.*, 1996, **35**(15), 4529

MRH Ammonia 1.88/14, hydrazine 2.43/19, hydroxylamine 3.22/49, calcium nitride 5.90/41

Ammonia—chlorine mixtures are explosive if warmed or if chlorine is in excess, owing to formation of nitrogen trichloride. Hydrazine, hydroxylamine and calcium nitride ignite in chlorine, and nitrogen triiodide may explode on contact with chlorine [1]. During chlorination of impure biuret in water at 20°C , a violent explosion occurred [2]. This was attributed to conversion of the cyanuric acid impurity (3%) to nitrogen trichloride and spontaneous explosion of the latter [3]. During interaction of chlorine and alkylthiuronium salts to give alkanesulfonyl chlorides, the dangerously explosive nitrogen trichloride may be produced if excess chlorine or slow chlorination is used. General precautions are discussed [4]. More recently it has been recommended that prompt working up of the reaction mixture

Cl₂

will avoid formation of nitrogen trichloride [5]. Aziridine readily gives the explosive *N*-chloro compound. During chlorination of 2,4,6-triketo-hexahydro-1,3,5-triazine (cyanuric acid), presence of the diaminoketo and aminodiketo analogues as impurities, or of an unusually low pH value, may lead to formation of nitrogen trichloride. UV irradiation may be used to destroy this in a continuous circulation reactor and prevent build-up of dangerous concentrations [6]. Chlorination of amidosulfuric acid [7] or acidic ammonium chloride solutions [8] gives the powerfully explosive oil, nitrogen trichloride. The formation and separation of nitrogen trichloride have been studied in detail [9], [10].

See N-HALOGEN COMPOUNDS

Non-metal hydrides MRH Diborane 3.59/12, phosphine 2.13/11
Mellor, 1939, Vol. 9, 55, 396; 1939, Vol. 8, 65; 1940, Vol. 6, 219; 1941, Vol. 5, 37

Arsine, phosphine and silane all ignite in contact with chlorine at ambient temperature, while diborane and stibine react explosively, the latter also with chlorine water.

See Ethylphosphine: Halogens

Non-metals

Mellor, 1941, Vol. 2, 292; 1956, Vol. 2, Suppl. 1, 380; 1943, Vol. 11, 26

Liquid chlorine at -34°C explodes with white phosphorus, and a solution in heptane at 0°C ignites red phosphorus. Boron, active carbon, silicon and phosphorus all ignite in contact with gaseous chlorine at ambient temperature. Arsenic incandesces on contact with liquid chlorine at -34°C, and the powder ignites when sprinkled into the gas at ambient temperature. Tellurium must be warmed slightly before incandescence occurs.

Other reactants

1. Yoshida, 1980, 61–62
2. Gustin, J. J. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries*, Vol I, (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 157, Amsterdam, Elsevier, 1995

MRH values calculated for 31 combinations with oxidisable materials are given [1]. Flammability limits and auto-ignition temperatures for a number of mixtures with organic fuels are measured and reviewed. All experimental work reported is in the vapour phase, detonation is common. Precautions for chlorination reactions are considered [2].

Oxygen difluoride

See Oxygen difluoride: Halogens

Phenylmagnesium bromide

1. Datta, R. L. *et al.*, *J. Amer. Chem. Soc.*, 1919, **41**, 287
2. Zakharkin, L. I. *et al.*, *J. Organomet. Chem.*, 1970, **21**, 271

After treatment of the Grignard reagent with chlorine, the solid which separated exploded when shaken [1]. This solid was not seen later, using either diethyl ether or THF as solvent [2]. (The explosive solid could have been magnesium hypochlorite if moisture were present in the chlorine.)

See Magnesium hypochlorite

Phosphorus compounds MRH values below references

1. Mellor, 1940, Vol. 8, 812, 842, 844–845, 897

2. von Schwartz, 1918, 324

MRH Phosphine 2.13/11, phosphorus trioxide 0.46/43, magnesium phosphide 4.02/39

Borondiodophosphide, phosphine, phosphorus trioxide and trimercury tetraphosphide all ignite in contact with chlorine at ambient temperature. Trimagnesium diphosphide and trimanganese diphosphide ignite in warm chlorine [1], while ethylphosphine explodes with chlorine [2]. Unheated boron phosphide incandescens in chlorine.

See Trimethyl thionophosphate, below

Polychlorobiphenyl

Statesir, W. A., *Chem. Eng. Progr.*, 1973, **69**(4), 53–54

A mixture of a polychlorobiphenyl process oil and liquid chlorine confined in a stainless steel bomb reacted exothermally between 25 and 81°C.

Silicones

Statesir, W. A., *Chem. Eng. Progr.*, 1973, **69**(4), 53–54

Silicone process oils mixed with liquid chlorine confined in a stainless steel bomb reacted explosively on heating; polydimethylsiloxane at 88–118°C, and polymethyltrifluoropropylsiloxane at 68–114°C. Previously, leakage of a silicone pump oil into a liquid chlorine feed system had caused rupture of a stainless steel ball valve under a pressure surge of about 2 kbar.

Sodium hydroxide

MCA Case History No. 1880

Attempted disposal of a small amount of liquid chlorine by pouring it into 20% sodium hydroxide solution caused a violent reaction leading to personal contamination.

Steel

1. *MCA Case History No. 608*

2. Stephens, T. J. R. *et al.*, *Paper 62B*, 65th AIChE Meeting, New York, 1972

3. Vervalin, 1973, 85

4. Schwarz, E., *Proc. 3rd Int. Symp. Prev. Occ. Risks Chem. Ind.*, 429–434, Heidelberg, ISSA, 1976

5. Fishwick, T., *Loss Prev. Bull.*, 1998, 139, 12

6. Anon., *Loss. Prev. Bull.*, 1983, (052), 7

Cl₂

Chlorine leaking into a steam-heated mild steel pipe caused ignition of the latter at about 250°C [1]. Sheet steel in contact with chlorine usually ignites at 200–250°C, but the presence of soot, rust, carbon or other catalysts may reduce the ignition temperature to 100°C. Dry steel wool ignites in chlorine at only 50°C [2]. Use of a carbon steel inlet pipe in a paraffin chlorination system led to rupture of the pipe and a fire [3]. Presence of an organic contaminant in a steel liquid chlorine vaporiser operating at 50°C led to local exothermic heating to 140–150°C and an iron–chlorine fire ensued. In another incident, admission of chlorine gas into a recently-welded (warm) steel pipe again led to a fire [4]. Another account of the same, or a very similar, vaporiser fire is found in [5]. When steel tanks containing chlorine are exposed to fire conditions, the tank wall above the liquid level may soon attain a temperature at which combustion of steel in chlorine will occur, perforating the tank. Two incidents involving derailed chlorine tank cars are described [6].

See other CORROSION INCIDENTS, IGNITION SOURCES

Sulfides

Mellor, 1940, Vol. 4, 952; 1946, Vol. 6, 144; 1939, Vol. 9, 270

Arsenic disulfide, boron trisulfide and mercuric sulfide all ignite in chlorine at ambient temperature, the first only in a rapid stream.

Synthetic rubber

Murray, R. L., *Chem. Eng. News*, 1948, **26**, 3369

During interaction of synthetic rubber and liquid chlorine, a violent explosion occurred. It is known that natural and synthetic rubbers will burn in liquid chlorine.

See Hydrocarbons, above

Tetraselenium tetranitride

See Tetraselenium tetranitride

Trialkyl boranes

Coates, 1967, Vol. 1, 199

The lower homologues tend to ignite in chlorine or bromine.

Trimethyl thionophosphate

MCA Case History No. 371

At an early stage in the preparation of methyl parathion, it is supposed that the phosphorus ester was being chlorinated to give dimethyl thionophosphorochloridate. Thermocouple failure indicated a low reaction temperature and the process controller boosted the chlorine feed rate, but when this fault situation was realised, the chlorine flow and agitator were stopped. However, an exothermic runaway reaction developed, eventually leading to a violent explosion.

See other RUNAWAY REACTIONS

Tungsten dioxide

Mellor, 1943, Vol. 11, 851

Incandescence on warming.

Water

1. Kosharov, P., *Chem. Abs.*, 1940, **34**, 3917q

2. Bray, A. W., private comm., 1982

The statement that 'Mixtures of chlorine and water at certain concentrations are capable of explosion by spark ignition' [1] should read 'Mixtures of chlorine and hydrogen' [2].

See other HALOGENS, OXIDANTS

4048. Cobalt(II) chloride

[7646-79-9]

CoCl₂

Cl₂Co

HCS 1980, 320

Metals

See Potassium: Metal halides

Sodium: Metal halides

See other METAL HALIDES

4049. Dihydrzinecobalt(II) chlorate

[]

[(H₄N₂)₂Co] [ClO₃]₂

Cl₂CoH₈N₄O₆

Salvadori, R., *Gazz. Chim. Ital.* [2], 1910, **40**, 9

It explodes powerfully on the slightest impact or friction, or on heating to 90°C.

See other AMMINEMETAL OXOSALTS

4050. Pentaamminephosphinatocobalt(III) perchlorate

[]

[(H₃N)₅CoH₂PO₂] [ClO₄]₂

Cl₂CoH₁₇N₅O₁₀P

Anon., *BCISC Quart. Safety Summ.*, 1965, **36**, 58

When a platinum wire (which may have been hot) was dipped for a flame test into a sintered funnel containing the air-dried complex, detonation occurred. This may have been due to heat and/or friction on a compound containing both strongly oxidising and reducing radicals. Avoid dipping (catalytically active) platinum wire into bulk samples of materials of unknown potential.

See other AMMINEMETAL OXOSALTS, REDOX COMPOUNDS

4051. Cobalt(II) perchlorate hydrates

[13478-33-6, 33827-55-3, 73954-59-3] (6, 4, 2H₂O, resp.) Cl₂CoO₈·2H₂O
Co(ClO₄)₂·2H₂O

1. Salvadori, R., *Gazz. Chim. Ital.* [2], 1910, **40**, 15–16
2. Robinson, P. J., private comm., 1982
3. Robinson, P. J., *Nature*, 1981, **293**, 696
4. Cook, R. E. *et al.*, *J. Chem. Res. (S)*, 1982, 267 (*M*, 2772–2783)
5. Cook, R. E. *et al.*, *Chem. Brit.*, 1982, **18**, 859

In view of the ready commercial availability and apparent stability of the hexahydrate, it is probable that the earlier report of explosion on impact, and deflagration on rapid heating [1] referred to the material produced by partial dehydration at 100°C, rather than the hexahydrate [2]. The caked crystalline hydrated salt, prepared from aqueous perchloric acid and excess cobalt carbonate with subsequent heated evaporation, exploded violently when placed in a mortar and tapped gently to break up the crystalline mass, when a nearby dish of the salt also exploded [3]. Subsequent investigation revealed the probable cause as heating the solid stable hexahydrate to a temperature (~150°C) at which partial loss of water produced a lower and endothermic hydrate (possibly a trihydrate) capable of explosive decomposition. This hazard may also exist for other hydrated metal perchlorates, and general caution is urged [4,5].

See other METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS

4052. Chromium(II) chloride

[10049-05-5] Cl₂Cr
CrCl₂

Water

1. *MCA Case History No. 1660*
2. Bretherick, L., *Chem. Brit.*, 1976, **12**, 204

An unopened bottle of chromous chloride solution exploded after prolonged storage [1]. This was most likely caused by internal pressure of hydrogen developed by slow reduction of the solvent water by the powerfully reducing Cr(II) ion [2].

See Chromium(II) sulfate

See other GAS EVOLUTION INCIDENTS, METAL HALIDES, REDUCANTS

4053. Pentaamminephosphinatochromium(III) perchlorate

[65901-39-5] Cl₂CrH₁₇N₅O₁₀P
[(H₃N)₅CrH₂PO₂] [ClO₄]₂

Coronas, J. M. *et al.*, *Inorg. Chim. Acta*, 1977, **25**, 110

A sample detonated violently on heating.

See other AMMINOMETAL OXOSALTS, REDOX COMPOUNDS

4054. Chromyl chloride
[14977-61-8]



1. Sidgwick, 1950, 1004
2. Pitwell, L. R., private comm., 1964

Though a powerful and often violent oxidant of inorganic and organic materials in absence of a diluent, it has found use as solutions in preparative organic chemistry for the controlled oxidation of alkyl aromatic derivatives [1]. In such a reaction, failure of a stirrer during addition of the chloride caused a build-up of unreacted material, followed by a violent explosion [2].

See other AGITATION INCIDENTS

Ammonia

Mellor, 1943, Vol. 11, 394

Contact with ammonia causes incandescence.

Isopropylcyclopropane

Wang, K. *et al.*, *J. Org. Chem.*, 1997, **62**(13), 4248

A violent reaction was experienced on adding this hydrocarbon to the chloride at room temperature. They can be safely mixed at -78°C .

Non-metal halides

1. Mellor, 1943, Vol. 11, 394–395

2. Pascal, 1959, Vol. 14, 153

A jet of chromyl chloride vapour ignites in the vapour of disulfur dichloride, and addition of drops of chromyl chloride to cooled phosphorus trichloride causes incandescence and sometimes explosion [1]. Phosphorus tribromide may also ignite with the chloride [2].

Non-metal hydrides

Pascal, 1959, Vol. 14, 153

Hydrogen sulfide or phosphine may ignite in contact with the chloride.

Non-metals

Mellor, 1943, Vol. 11, 394–395

Moist phosphorus explodes in contact with chromyl chloride, while flowers of sulfur ignites.

Organic solvents

Mellor, 1943, Vol. 11, 396

Acetone, ethanol or ether ignite on contact with the chloride, and turpentine behaves similarly.

Sodium azide

Mellor, 1967, Vol. 8, Suppl. 2.2, 36

Interaction of chromyl chloride and sodium azide to form chromyl azide is explosive in absence of a diluent.

Urea

Pascal, 1959, Vol. 14, 153

Urea ignites in contact with chromyl chloride.

Water

Partington, 1966, 751


Hydrolysis of chromyl chloride is violent.

See related METAL HALIDES

See other OXIDANTS

4055. Chromyl perchlorate

[62597-99-3]



Alone, or Organic solvents

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

A powerful oxidant which may explode violently above 80°C, and which causes organic solvents to ignite on contact. It decomposes on exposure to light, usually slowly, sometimes explosively.

See related METAL PERCHLORATES

See other OXIDANTS

4056. Copper(I) chloride

[7758-89-6]



HCS 1980, 337

Preparative hazard

See Copper: Complexing agents, Water

Lithium nitride

See Lithium nitride: Copper(I) chloride

See other METAL HALIDES

4057. Copper(II) perchlorate

[10294-46-9]



Polyfunctional amines

Fabbrizzi, L. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1984, 1496

Complexes formed with the extended chain polyfunctional amines $\text{H}_2\text{N}[\text{CH}_2]_n\text{NH}[\text{CH}_2]_m\text{NH}[\text{CH}_2]_n\text{NH}_2$ (n or $m = 2$ or 3) may be explosive and must be handled with care.

See related AMMINOMETAL OXOSALTS, [14] DIENE- N_4 COMPLEXES

N-(2-Pyridyl)acetylacetamides

Illiopoulos, P. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1986, 437–443

The dihydrated binuclear complexes between copper(II) perchlorate and *N*-(2-pyridyl)-acetoacetamide and -benzoylacetamide, $[\text{Cu}_2\text{L}_2(\text{ClO}_4)_2]\cdot 2\text{H}_2\text{O}$, exploded on heating to 260°C .

See related AMMINOMETAL OXOSALTS, [14] DIENE- N_4 COMPLEXES

See other METAL PERCHLORATES

4058. Copper(I) perchlorate

[15061-57-1]

$\text{Cl}_2\text{Cu}_2\text{O}_8$

$\text{O}_4\text{ClCuCuClO}_4$

Alkenes, or Carbon monoxide

1. Ogura, T., *Inorg. Chem.*, 1976, **15**, 2301–2303

2. Thompson, J. S. *et al.*, *Inorg. Chem.*, 1984, **23**, 2814

If preparation of the perchlorate (by stirring copper(II) perchlorate in water with copper powder) is effected under an atmosphere of the ligands ethylene, allene, 1,3-butadiene or carbon monoxide, two types of explosive complexes, $\text{L}_2\cdot\text{Cu}_2(\text{ClO}_4)_2\cdot n\text{H}_2\text{O}$, (L represents one double bond), are formed. There is a crystalline series with $n = 4$, e.g. $2\text{CO}\cdot\text{Cu}_2(\text{ClO}_4)_2\cdot 4\text{H}_2\text{O}$, and an amorphous series with $n = 3$, e.g. $\text{C}_3\text{H}_4\cdot\text{Cu}_2(\text{ClO}_4)_2\cdot 3\text{H}_2\text{O}$. All the complexes are explosive, but unpredictably so. Sometimes a complex would not explode under a hammer-blow, but sometimes it would explode on stirring with a Teflon-coated stirrer bar. On one occasion a 10 mg sample stored in glass exploded, propelling glass fragments 10 m [1]. When the ethylene complex is prepared in methanol solution, it must not be allowed to dry out or it will explode [2].

1,4-Oxathiane

Barnes, J. C. *et al.*, *J. Chem. Soc. Pak.*, 1982, **4**, 103–113

Of the two complexes which the perchlorate forms with oxathiane (1:3, 1:4), the latter explodes on heating.

See other METAL PERCHLORATES, REDOX COMPOUNDS

4059. Dichlorofluoramine

[17417-38-8]

Cl_2FN

Cl_2NF

1. Sukornick, B. *et al.*, *Inorg. Chem.*, 1963, **2**, 875

2. Batty, W. E., *Chem. & Ind.*, 1969, 1232

In the liquid phase, it is an extremely friction- and shock-sensitive explosive [1], like the trichloro analogue [2].

See other *N*-HALOGEN COMPOUNDS

4060. *N,N*-Dichloropentafluorosulfanylamine
[22650-46-0] **Cl₂F₅NS**



Kirk Othmer, 4th Edn., 1993, Vol IV, 917
Shock sensitive and unstable from 80°C.

Preparative hazard

Waterfeld, A. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 839

Reaction of chlorine fluoride with trifluorosulfur nitride is rather hazardous and may lead to violent explosions. A safer alternative preparation is to use chlorine and mercuric fluoride in place of chlorine fluoride.

See other N-HALOGEN COMPOUNDS, N-S COMPOUNDS

4061. Iron(II) chloride
[7758-94-3] **Cl₂Fe**



Ozonides

See OZONIDES: metals, etc.

See other METAL HALIDES, REDUCANTS

4062. Iron(II) perchlorate
[13933-23-8] **Cl₂FeO₈**



Preparative hazard

See Perchloric acid: Iron(II) sulfate

See other METAL PERCHLORATES, REDOX COMPOUNDS

4063. Dichloramine (Chlorimide)
[3400-09-7] **Cl₂HN**



Eckert, P. *et al.*, *Chem. Abs.*, 1951, **45**, 7527g

Dissolving *N,N,N',N'*-tetrachloroadipamide in boiling water gives a highly explosive oil, probably dichloramine.

See other N-HALOGEN COMPOUNDS

4064. *N,N'*-Bis(chloromercurio)hydrazine
[] **Cl₂H₂Hg₂N₂**





Mellor, 1940, Vol. 4, 874, 881; Vol. 8, 318

It explodes when heated or struck, and the bromo and other analogues are similar.
See Mercury(II) chloride: Hydrazine salts, Base
See other MERCURY COMPOUNDS See related METAL HYDRAZIDES

4065. Chloronium perchlorate

[]



Hantzsch, A., *Ber.*, 1930, **63**, 1789

This hydrogen chloride–perchloric acid complex spontaneously dissociates with explosive violence.

See other NON-METAL PERCHLORATES

†4066. Dichlorosilane

[4109-96-0]



1. Sharp, K. G. *et al.*, *J. Electrochem. Soc.*, 1982, **129**



2. Britton, L. G., *Plant/Oper. Progr.*, 1990, **9**(1), 16

It is unusual in being flammable with so high a chlorine content (70%), and mixtures with air may detonate if confined when ignited. One case of spontaneous ignition under ambient conditions was observed [1]. It has an unusually low autoignition temperature ($43 \pm 3^\circ\text{C}$). A survey of hazards and combustion is found in [2]. There is a risk of combustion with fluorochloromethanes. Autoignition was observed on contact with traces of a mixture of alkali nitrates/nitrite.

See other HALOSILANES

4067. Bis(hydroxylamine)zinc chloride

[33774-96-8]




Walker, J. E. *et al.*, *Inorg. Synth.*, 1967, **9**, 2

It explodes at 170°C .

See related AMMINOMETAL HALIDES

4068. Hydrazinium diperchlorate

[13812-39-0]



Grelecki, C. J. *et al.*, *ACS* **54**, 1966, 73

The thermal decomposition of this solid rocket propellant component has been studied.

Metal compounds

Probhakaran, C. *et al.*, *Def. Sci. J.*, 1982, **31**, 285–291

Of the series of additives copper chromate, copper chloride, nickel oxide, iron(III) oxide or magnesium oxide, the earlier members have the greatest effect in increasing the sensitivity of the explosive salt towards heat, impact or friction.

See other PERCHLORATE SALTS OF NITROGENOUS BASES

4069. Bis(hydrazine)nickel perchlorate

[]




Maissen, B. *et al.*, *Helv. Chim. Acta*, 1951, **34**, 2084–2085

The salt, known to be explosive when heated dry (but not under a hammer-blow), exploded violently when stirred as a dilute aqueous suspension.

See other AMMINEMETAL OXOSALTS

4070. Bis(hydrazine)tin(II) chloride

[55374-98-6]




Mellor, 1941, Vol. 7, 430

It explodes on heating.

See related AMMINEMETAL HALIDES, METAL HYDRAZIDES

4071. Tetraamminebis(dinitrogen)osmium(II) perchlorate

[]




Creutz, C. A., private comm., 1969

The dry salt may explode on touching.

See next entry below *See other* AMMINEMETAL OXOSALTS

4072. Pentaamminedinitrogenosmium(II) perchlorate

[20611-53-4]



Bohr, J. D. *et al.*, *Inorg. Chem.*, 1980, **19**, 2417

It can be detonated.

See entry next above *See other* AMMINEMETAL OXOSALTS

4073. Pentaammineazidoruthenium(III) chloride
[28223-30-5] (ion) **Cl₂H₁₅N₈Ru**

Hazardous intermediate

See Pentaamminechlororuthenium chloride: Sodium azide

See related METAL AZIDES

4074. Hexaamminenickel chlorate
[] **Cl₂H₁₈N₆NiO₆**
[(H₃N)₆Ni] [ClO₃]₂

Mellor, 1956, Vol. 2, Suppl. 1, 592

It explodes on impact.

See other AMMINOMETAL OXOSALTS

4075. Hexaamminenickel perchlorate
[14322-50-0] **Cl₂H₁₈N₆NiO₈**
[(H₃N)₆Ni] [ClO₄]₂

Mellor, 1956, Vol. 2, Suppl. 1, 592

It explodes on impact, but is less sensitive than the chlorate.

See other AMMINOMETAL OXOSALTS

4076. Mercury(II) chloride
[7847-94-7] **Cl₂Hg**
HgCl₂

HCS 1980, 615

Hydrazine salts, Base

Hofmann, K. A. *et al.*, *Annalen*, 1899, **305**, 191

A yellowish precipitate was obtained from the chloride and hydrazinium chlorides when basified with sodium acetate. It had N—Hg bonds and was perhaps ClHgNHNHHgCl. It exploded on heating or shock. A similar compound resulted from the bromides.

Sodium *aci*-nitromethanide

See Sodium *aci*-nitromethanide: Mercury(II) chloride

Sodium azide

See Sodium azide: Heavy metals

See other MERCURY COMPOUNDS, METAL HALIDES

4077. Mercury(II) chlorite
[73513-17-4]



Levi, G. R., *Gazz. Chim. Ital.* [2], 1915, **45**, 161

It is extremely unstable when dry, exploding spontaneously.

See other CHLORITE SALTS, MERCURY COMPOUNDS

4078. Mercury(II) perchlorate
[7616-83-3]




Samuels, G. J. *et al.*, *Inorg. Chem.*, 1980, **19**, 233

A filtered solution, prepared by heating mercuric oxide with a slight excess of perchloric acid, after standing for several months precipitated a little unidentified white solid. This (but not the supernatant liquid) was very shock sensitive, and detonated as it was being rinsed out with water. (Traces of a volatile amine may have been absorbed into the acid liquor to give an aminomercury perchlorate, expected to be explosive).

See other AMMINOMETAL OXOSALTS, MERCURY COMPOUNDS, METAL PERCHLORATES

4079. Mercury(II) perchlorate . 6 (or 4)dimethyl sulfoxide
[62909-81-3] (.6DMSO)



1. Rayner, P. N. G., *Chem. Brit.*, 1977, **13**, 396; private comm., 1977
2. Cotton, F. A. *et al.*, *J. Amer. Chem. Soc.*, 1960, **82**, 2886
3. Brown, A. J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1978, 1778
4. Sandstrom, M. *et al.*, *Acta Chem. Scand.*, 1978, **A32**, 610

The hexasolvated compound was being prepared from a mixture of the hydrated salt, methanol and dimethyl sulfoxide but inexplicably mercury(I) oxide came out of solution. This was redissolved by addition of perchloric acid and the solution was refrigerated for 60 h. After addition of molecular sieve to remove the water present, the solution was filtered and the sieve treatment repeated four more times. During the final filtration, a violent explosion occurred in the funnel. The most likely cause was formation of methyl perchlorate promoted by the dehydrating action of the sieve, but dimethyl sulfoxide also forms an unstable salt with perchloric acid, so this also may have been involved [1]. During preparation of the hexasolvate by a published method [2], a violent explosion occurred [3]. The solid tetrasolvated product from treatment of a solution of the perchlorate in methanol-perchloric acid with DMSO at ambient temperature exploded with extreme violence after prolonged desiccation (possibly owing to formation of methyl perchlorate). Materials prepared at -78°C appeared to be free of this hazard, though they will explode under mechanical impact, etc. [4].

See Dimethyl sulfoxide: Metal oxosalts

See other METAL PERCHLORATES

See other MOLECULAR SIEVE INCIDENTS, SOLVATED OXOSALT INCIDENTS

4080. Mercury(I) chlorite
[101672-18-8]



Levi, G. R., *Gazz. Chim. Ital.* [2], 1915, **45**, 161

It is extremely unstable when dry, exploding spontaneously.

See other CHLORITE SALTS, MERCURY COMPOUNDS, REDOX COMPOUNDS

4081. Magnesium chloride
[7786-30-3]



Air, Mild steel, Water

1. Anon., *CISHC Chem. Safety Summ.*, 1984, **55**, 66

2. Haudt, P. R., *Mater. Perform.*, 1986, **25**(9), 49–55

3. Melcher, R. G. *et al.*, *Am. Ind. Hyg. Assoc. J.*, 1987, **48**, 608–612

A large steel evaporator used for magnesium chloride solution was shut down for maintenance. During maintenance operations a fatality occurred from atmospheric oxygen deficiency inside the evaporator. It was found later that the oxygen content in the evaporator fell from the normal 21% to about 1% in under 24 h, and this was confirmed in laboratory tests. This was attributed to very rapid rusting of the steel under warm humid conditions in the presence of traces of magnesium chloride [1]. Further work shows that other salts (calcium bromide, calcium chloride, magnesium sulfate, potassium chloride) behave similarly, and that presence of scale is a contributory factor [2]. Magnetite scale (Fe_3O_4) on mild steel increases the depletion rate by a factor of 10, while the rust formed during the corrosion has little effect [3].

See other CORROSION INCIDENTS

Jute, Sodium nitrate

See Sodium nitrate: Jute, Magnesium chloride

See other METAL HALIDES

4082. Magnesium hypochlorite
[10233-03-1]




Clancey, V. J., *J. Haz. Mat.*, 1975, **1**, 83–94

The compound is very unstable, and its presence may be one of the causes of the observed explosive and apparently spontaneous decomposition of calcium hypochlorite, if produced from magnesia-containing lime (derived from dolomite).

See Chlorine: Phenylmagnesium bromide
See other METAL HYPOCHLORITES

4083. Magnesium chlorate
[10326-21-3]



Sulfides

Mellor, 1956, Vol. 2, Suppl. 1, 584

Interaction with copper(I) sulfide is explosive, and with antimony(III) sulfide, arsenic(III) sulfide, tin(II) sulfide or tin(IV) sulfide, incandescent.

See other METAL CHLORATES

4084. Magnesium perchlorate
[10034-81-8]



Marusch, H., *Chem. Tech.* (Berlin), 1956, **8**, 482–485

The drying agent may contain traces of perchloric acid remaining from manufacturing operations, and owing to the great desiccating power of the salt, the acid will be anhydrous. Many of the explosions experienced with magnesium perchlorate have their origins in contact of the anhydrous acid with oxidisable materials, or materials able to form unstable perchlorate esters or salts.

Alkenes

Heertjes, P. M. *et al.*, *Chem. Weekbl.*, 1941, **38**, 85

The anhydrous salt which had been used for drying unsaturated hydrocarbons exploded on heating to 220°C for reactivation. The need to avoid contact with acidic materials is stressed. (Traces of an alkyl perchlorate may conceivably have been formed from free perchloric acid).

See Organic materials, below

Ammonia

491M, 1975, 244

Intensive drying of ammonia gas by passing it over the desiccant in a steel drying tube led to an exotherm, followed by a violent explosion. (An ammine derivative may have been formed).

See related AMMINOMETAL OXOSALTS

Argon

Dam, J. W., *Chem. Weekbl.*, 1958, **54**, 277

An explosion which occurred when argon was being dried obviously involved some (unstated) impurity in the system.

Arylhydrazine, Ether

Belcher, R., private comm., 1968

Anhydrous magnesium perchlorate was used to dry thoroughly an ethereal solution of an arylhydrazine. During evaporation of the filtered solution it exploded completely and violently. Magnesium perchlorate is rather soluble in ether and may contain traces of free perchloric acid (probably in the anhydrous form, as the magnesium salt is a powerful dehydrator). It is entirely unsuitable for drying organic solvents. (The hydrazine may have complexed with the magnesium to form a redox salt).

See related AMMINOMETAL OXOSALTS

Cellulose, Dinitrogen tetroxide, Oxygen

Anon., *ABCM Quart. Safety Summ.*, 1961, **32**, 6

Magnesium perchlorate contained in a glass tube between wads of cotton wool was used to dry a mixture of oxygen and dinitrogen tetroxide. After several days the drying tube exploded violently. It seems probable that the acidic fumes and cotton produced cellulose nitrate, aided by the dehydrating action of the perchlorate.

Dimethyl sulfoxide

1. Tobe, M. L. *et al.*, *J. Chem. Soc.*, 1964, 2991

2. Anon., *Chem. Eng. News*, 1965, **43**(47), 62

3. Dessy, R. E. *et al.*, *J. Amer. Chem. Soc.*, 1964, **86**, 28

In the preparation of anhydrous DMSO by a literature method [1], an explosion occurred during distillation from anhydrous magnesium perchlorate [2]. This may have been due to the presence of some free methanesulfonic acid as an impurity in the solvent, which could liberate traces of perchloric acid. It is known that sulfoxides react explosively with 70% perchloric acid, (but several metal perchlorates also form unstable solvates with DMSO) The alternative procedure for drying DMSO with calcium hydride [3] seems preferable, as this would also remove any acidic impurities.

See Dimethyl sulfoxide: Metal oxosalts

Ethanol

Raymond, K. R., private comm., 1983

The desiccant in a drying tube, accidentally exposed to ethanol vapour, was left for several months. The explosion which occurred when the desiccant was scraped out was certainly due to formation of ethyl perchlorate.

See ALKYL PERCHLORATES

Ethylene oxide

491M, 1975, 314

Desiccation of gaseous ethylene oxide led to an explosion, (possibly involving formation of 2-hydroxyethyl perchlorate).

Fluorobutane, Water

Anon., *Ind. Eng. Chem. (News Ed.)*, 1939, **17**, 70

A violent explosion followed the use of magnesium perchlorate to dry wet fluoro-butane. The latter was presumed to have hydrolysed to give hydrogen fluoride which had liberated perchloric acid, explosively unstable when anhydrous. (This explanation seems unlikely in view of the large disparity between dissociation constants of the two acids). Magnesium perchlorate is unsuitable for drying acidic or flammable materials; calcium sulfate would be suitable.

Organic materials

1. Hodson, R. J., *Chem. & Ind.*, 1965, 1873
2. *MCA Case History No. 243*

The use of the perchlorate as desiccant in a drybag where contamination with organic compounds is possible is considered dangerous [1]. Magnesium perchlorate ('Anhydrone') was inadvertently used instead of calcium sulfate (anhydrite) to dry an unstated reaction product before vacuum distillation. The error was realised and all solid was filtered off. Towards the end of the distillation, decomposition and an explosion occurred, possibly owing to the presence of dissolved magnesium perchlorate, or more probably to perchloric acid present as impurity in the salt [2].

Phosphorus

See Phosphorus: Magnesium perchlorate

Trimethyl phosphite

Jercinovic, L. M. *et al.*, *J. Chem. Educ.*, 1968, **45**, 751

Contact between the salt and ester caused violent explosions on several occasions. (Methyl perchlorate may have been formed).

See related ALKYL PERCHLORATES

See other METAL PERCHLORATES

4085. Manganese(II) chloride

[7773-01-5]



Zinc

See Zinc: Manganese dichloride

See other METAL HALIDES

4086. Manganese dichloride dioxide

[51819-69-3]



Briggs, T. S., *J. Inorg. Nucl. Chem.*, 1968, **30**, 2867–2868

Explosively unstable if isolated as liquid at ambient temperature, it may be handled safely in carbon tetrachloride solution.

See Manganese chloride trioxide, Manganese trichloride oxide

See related METAL HALIDES, METAL OXIDES

4087. Manganese(II) chlorate
[104813-96-9]



Bailar, 1973, Vol. 3, 835

The hexahydrated salt decomposes explosively above 6°C, producing chlorine dioxide.

See other METAL CHLORATES, REDOX COMPOUNDS

4088. Manganese(II) perchlorate
[13770-16-6]



Sidgwick, 1950, 1285

It explodes at 195°C.



2,2-Dimethoxypropane

See 2,2-Dimethoxypropane: Metal perchlorates

See other METAL PERCHLORATES, REDOX COMPOUNDS

4089. Chloro-1,2,4-triselenadiazolium chloride (3-Chloro-1,3,4-triselenadiazolium chloride)

[]



1. Chivers, T. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(11), 1518



2. Chivers, T. *et al.*, *Inorg. Chem.*, 1993, **32**(20), 4391

The black explosive powder previously described as Se_4N_2 is now assigned this structure. It is very sensitive when dry and handling it damp with an hydrocarbon is recommended. There is brown dimer, also explosive [2].

See related NITRIDES, NON-METAL HALIDES, N-S COMPOUNDS

4090. Vanadyl azide dichloride

[]




An explosive solid.

See entry METAL AZIDE HALIDES

4091. Di(triselenadiazolium) dichloride

[]

 $\text{Cl}_2\text{N}_4\text{Se}_6$ Chivers, T. *et al.*, *Inorg. Chem.*, 1993, **32**(20), 4391

A brown explosive powder

See Chloro-1,2,4-triselenadiazolium chloride**4092. Diazidodichlorosilane**

[67880-19-7]

 $\text{Cl}_2\text{N}_6\text{Si}$ $(\text{N}_3)_2\text{SiCl}_2$ *See* Silicon tetraazide *See other* NON-METAL AZIDES**4093. Nickel chlorite**

[72248-72-7]

 Cl_2NiO_4 $\text{Ni}(\text{ClO}_2)_2$

1. Mellor, 1956, Vol. 2, Suppl. 1, 574

2. Levi, G. R., *Gazz. Chim. Ital.* [2], 1923, **53**, 245

The dihydrate explodes at 100°C [1], or on percussion, even when wet [2].

See other CHLORITE SALTS**4094. Nickel perchlorate**

[13637-71-3]

 Cl_2NiO_8 $\text{Ni}(\text{ClO}_4)_2$

2,2-Dimethoxypropane

See 2,2-Dimethoxypropane: Metal perchlorates*See other* METAL PERCHLORATES**4095. Dichlorine oxide**

[7791-21-1]

 Cl_2O Cl_2O

1. Sidgwick, 1950, 1202

2. Jacobs, P. W. M. *et al.*, *Chem. Rev.*, 1969, **69**, 5593. Pilipovich, D. *et al.*, *Inorg. Chem.*, 1972, **11**, 21904. Gray, P. *et al.*, *Combust. Flame*, 1972, **18**, 361–3715. Cady, G. H., *Inorg. Synth.*, 1957, **5**, 156

1429

It is somewhat endothermic (ΔH_f° (g) +87.5 kJ/mol, 1.0 kJ/g), the liquid may explode on pouring or sparking at 2°C, and the gas readily explodes on rapid heating or sparking [1,2], on adiabatic compression in a U-tube, or often towards the end of slow thermal decomposition. Kinetic data are summarised [3]. The spontaneously explosive decomposition of the gas was studied at 42–86°C, and induction periods up to several hours were noted [4]. Preparative precautions have been detailed [5].

See other ENDOTHERMIC COMPOUNDS, INDUCTION PERIOD INCIDENTS

Alcohols

Gallais, 1957, 677

Alcohols are oxidised explosively.

Carbon

Mellor, 1946, Vol. 5, 824

Addition of charcoal to the gas causes an immediate explosion, probably initiated by the heat of adsorption of the gas on the solid.

Carbon disulfide

Mellor, 1940, Vol. 6, 110

The vapours explode on contact.

Dicyanogen

Brotherton, T. K. *et al.*, *Chem. Rev.*, 1959, **59**, 843

Contact causes ignition or explosion.

Diphenylmercury

See Diphenylmercury: Chlorine monoxide, etc.

Ethers

Gallais, 1957, 677

Ethers are oxidised explosively.

Hydrocarbons

Ip, J. K. K. *et al.*, *Combust. Flame*, 1972, **19**, 117–129

The spontaneously explosive interaction of dichlorine oxide with methane, ethane, propane, ethylene or butadiene was investigated at 50–150°C. Self-heating occurs with ethylene, ethane and propane mixtures.

See other SELF-HEATING AND IGNITION INCIDENTS

Nitrogen oxide

Mellor, 1940, Vol. 8, 433

Interaction is explosive.

Oxidisable materials

1. Mellor, 1941, Vol. 2, 241–242; 1946, Vol. 5, 824

2. Cady, G. H., *Inorg. Synth.*, 1957, **5**, 156

The heat sensitivity (above) may explain the explosions which occur on contact of many readily oxidisable materials with this powerful oxidant. Such materials include ammonia, potassium; arsenic, antimony; sulfur, charcoal (adsorptive

heating may also contribute); calcium phosphide, phosphine, phosphorus; hydrogen sulfide, antimony sulfide, barium sulfide, mercury sulfide and tin sulfide [1]. Various organic materials (paper, cork, rubber, turpentine, etc.) behave similarly [2]. Mixtures with hydrogen detonate on ignition [1].

Potassium

See Potassium: Non-metal oxides

See other HALOGEN OXIDES, OXIDANTS

4096. Sulfinyl chloride (Thionyl chloride)

[7719-09-7]

Cl₂OS

O:SCl₂

HCS 1980, 898; RSC Lab. Hazard Data Sheet No. 26, 1984

Ammonia

MRH 0.84/16

Foote, C. S., private comm., 1965

Addition of a solution of 4-nitrobenzoyl chloride (1 g) in a large excess (10 ml) of sulfinyl chloride to ice-cold conc. ammonia solution caused a violent explosion. This may certainly be attributed to the instantaneous hydrolysis of the excess sulfinyl chloride by the aqueous ammonia with production of several l of un-neutralised acid gases in a test tube.

See Water, below

Bis(dimethylamino) sulfoxide

Armitage, D. A. *et al.*, *J. Inorg. Nucl. Chem.*, 1974, **36**, 993

Interaction of the chloride with the sulfoxide or its higher homologues to form dialkylaminosulfinyl chlorides causes extensive decomposition, possibly explosive above 80°C.

Chloryl perchlorate

See Chloryl perchlorate: Thionyl chloride

1,2,3-Cyclohexanetrione trioxime, Sulfur dioxide

1. Tokura, N. *et al.*, *Bull. Chem. Soc. Japan*, 1962, **35**, 723

2. Lewis, J. J., *J. Heterocycl. Chem.*, 1975, 12, 601

A previous method of making 4-oximino-4,5,6,7-tetrahydrobenzofurazan by cyclising the oxime with sulfinyl chloride in liquid sulfur dioxide sometimes led to explosive reactions [1]. A new procedure involving aqueous calcium carbonate is quite safe [2].

Diisopropyl malonate

McKeown, R. H., private comm., 1985

During reflux of a mixture to produce malonoyl chloride, vivid sparks were seen in the flask, and the reaction was closed down without mishap. No explanation is apparent, but the diisopropyl ester structure appears likely to be susceptible to autoxidation on storage, and peroxides may possibly have been involved in the phenomenon.

See related PEROXIDATION INCIDENTS

Dimethylformamide

1. Spitulnik, M. J., *Chem. Eng. News*, 1977, **55**(31), 31
2. Joshi, M. S., *Chem. Eng. News*, 1986, **64**(14), 2
3. Zemlicka, J., *Chem. Eng. News*, 1986, **64**(25), 4

Some 200 kg of a mixture of sulfinyl chloride and DMF decomposed vigorously after storage for several h at ambient temperature. This was attributed to the presence of 90 ppm each of iron and zinc in the chloride used for the preparation. Mixtures of the pure components remained unchanged for 48 h, but addition of 200 ppm of iron powder led to exothermic decomposition after stirring for 22 h [1]. During vacuum removal of sulfur dioxide from the reaction mixture in preparation of *N,N*-dimethylchloromethyliminium chloride, a sudden exotherm and pressure increase occurred in a 400 l reactor. Study of the reaction in an adiabatic calorimeter showed that a runaway reaction developed within 8 h at 60°C, or within 20 min at 85°C at ambient pressure. Reaction slurry from which part of the sulfur dioxide had been removed decomposed even more rapidly [2]. The use of phosgene in the reaction in place of sulfinyl chloride gives a more controlled reaction which does not need heating [3].

See other GAS EVOLUTION INCIDENTS

See Dimethylformamide: Halocarbons, etc.

See other CATALYTIC IMPURITY INCIDENTS

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides

Esters, Metals

1. Spagnolo, C. J. *et al.*, *Chem. Eng. News*, 1992, **70**(22), 2
2. Wang, S. S. Y. *et al.*, *Process Safety Progr.*, 1994, **13**(3), 153

It was found that a solution of thionyl chloride in ethyl acetate burst galvanised drums on storage. Further investigation demonstrated an exothermic gas-evolving reaction between the ester and the chloride, initiated by iron or zinc, to give chloroethane, acetyl chloride and sulfur dioxide. A similar reaction is sometimes used synthetically on lactones with Lewis acid catalysis, especially zinc chloride. This is therefore probably a general ester reaction[1]. A more comprehensive, but scarcely more comprehending, study of the ethyl acetate/thionyl chloride reaction demonstrates involvement of zinc. Elemental sulphur is apparently among the products [2].

See Diisopropyl malonate (above)

See Propionoyl chloride: Diisopropyl ether

See Sulfonyl chloride: Organic materials, metals

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

Heterocyclic amines

See HETEROCYCLIC *N*-SULFINYLAMINES

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Linseed oil, Quinoline MRH Quinoline 1.63/99+

1. Fieser, L. F. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 570

2. Laws, G. F., private comm., 1966

It is important to add the quinoline and linseed oil, used in purifying the chloride [1], to the chloride. If reversed addition is used, a vigorous decomposition may occur [2].

Lithium

See Lithium: Sulfinyl chloride

Other reactants

Yoshida, 1980, 57

MRH values calculated for 14 combinations with various reagents are given.

Sodium

MRH 5.27/33

See Sodium: Non-metal halides

Tetrahydrofuran

Jackson, R. W. *et al.*, *Chem. Eng. News*, 1988, **66**(27), 2

Unstirred mixtures of sulfinyl chloride and tetrahydrofuran may react extremely exothermally and with gas evolution when heated to 60°C, the products of ring-scission being bis(4-chlorobutyl) ether, 1,4-dichlorobutane and sulfur dioxide. Some ratios of the two reactants enhance the tendency, and relatively confined mixtures may explode. Sufficient heat to initiate the reaction may arise from self-mixing of the layered liquids, the chloride being twice as dense as THF.

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

Toluene, Ethanol, Water

Anon., *Univ. Safety Assoc., Safety Newsletter*, 1982–1984

A solution of the chloride (120 ml) in toluene (750 ml) was treated (apparently without effective stirring) with excess sodium bicarbonate solution to destroy it. When reaction had ceased, the organic layer was poured into a waste solvent drum. Vigorous evolution of sulfur dioxide and hydrogen chloride then ensued from reaction with ethanol (toluene-soluble) in the waste drum. For destruction of solutions of sulfinyl chloride in water-insoluble solvents, extremely good agitation is necessary to ensure proper contact with a basic reagent. Ammonia is more soluble in toluene than is water, so ammonia solution should be used after bicarbonate treatment to ensure complete destruction.


Water

MCA Case History No. 1808

Passage of thionyl chloride through a flexible metal transfer hose which was contaminated with water or sodium hydroxide solution caused the hose to burst. Interaction with water violently decomposes the chloride to hydrogen chloride (2 mol) and sulfur dioxide (1 mol), the total expansion ratio from liquid to gas being 993:1 at 20°C, so very high pressures may be generated.

See other ACYL HALIDES, NON-METAL HALIDES

4097. Seleninyl chloride
[7791-23-3]



Antimony

Mellor, 1947, Vol. 10, 906

Powdered antimony ignites on contact with the chloride.

Metal oxides

Mellor, 1947, Vol. 10, 909

In contact with silver oxide, light is evolved and sufficient heat to decompose some of it. Similar effects were observed with lead(II) oxide, lead(IV) oxide and lead (II)(IV) oxide.

Potassium

Mellor, 1947, Vol. 10, 908

Potassium explodes violently in contact with the liquid.

Phosphorus



Mellor, 1947, Vol. 10, 906

Red phosphorus evolves light and heat in contact with the chloride, while white phosphorus explodes.

See other NON-METAL HALIDES

4098. Lead(II) hypochlorite

[]



Hydrogen sulfide

Pascal, 1960, Vol. 13.1, 1004

Ignition occurs on contact.

See other METAL HYPOCHLORITES

4099. Sulfonyl chloride (Sulfuryl chloride)

[7791-25-5]



Alkalies

Brauer, 1963, Vol. 1, 385

Reaction with alkalies may be explosively violent.

Diethyl ether

Dunstan, I. *et al.*, *Chem. & Ind.*, 1966, 73

A solution of sulfuryl chloride in ether vigorously decomposed, evolving hydrogen chloride. This was shown to be accelerated by the presence of peroxides. Peroxide-free ether should be used, and with care.

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides

Dinitrogen pentaoxide

See Dinitrogen pentaoxide: Sulfur dichloride, etc.

Lead dioxide

See Lead(IV) oxide: Non-metal halides

Organic materials, Metals

Anon., personal communication, 1996

Waste ethyl acetate containing sulfuryl chloride and thionyl chloride was stored in a galvanised steel drum. This later burst, although sulfuryl chloride is supplied in such drums they are unsafe with contaminants. The material can chlorinate many active organics, including aromatics, ketones and probably esters. This will release sulphur dioxide and hydrogen chloride, a fairly soluble gas, which will react with zinc and iron to produce hydrogen, an insoluble gas, and zinc or iron chlorides, both catalysts for many reactions including ester cleavage.

See Sulfinyl chloride: Esters, Metals

Phosphorus

See Phosphorus: Non-metal halides

See other ACYL HALIDES, NON-METAL HALIDES

4100. Dichlorine trioxide

[17496-59-2]

Cl₂O₃

Cl₂O₃

1. McHale, E. T. *et al.*, *J. Amer. Chem. Soc.*, 1967, **89**, 2796

2. McHale, E. T. *et al.*, *J. Phys. Chem.*, 1968, **72**, 1849–1856

It is endothermic (ΔH_f° (g) +76.2 kJ/mol, 0.64 kJ/g), very unstable, and the vapour explodes at about 2 mbar and well below 0°C [1]. It has been identified as the intermediate involved in the delayed explosion of chlorine dioxide [2].

See other ENDOTHERMIC COMPOUNDS, HALOGEN OXIDES

4101. Chlorine perchlorate

[27218-16-2]

Cl₂O₄

ClOClO₃

1. Schack, C. J. *et al.*, *Inorg. Chem.*, 1970, **9**, 1387

2. Schack, C. J. *et al.*, *Inorg. Chem.*, 1971, **10**, 1078

A shock-sensitive explosive compound [1]. Bromine perchlorate is also unstable [2].

Chlorotrifluoroethylene

Schack, C. J. *et al.*, *Inorg. Chem.*, 1973, **12**, 897

Interaction of the reactants (pre-mixed at –196°C) during warming to –78°, then ambient temperature, exploded. Progressive addition of the alkene to the perchlorate at –78° was uneventful.

Perfluoroalkyl iodides

Schack, C. J. *et al.*, *Inorg. Chem.*, 1975, **14**, 145–151

Reaction mixtures of chlorine perchlorate with perfluoromethyl iodide, 1,2-diiodoperfluoroethane or 1,3-diiodoperfluoropropane occasionally deflagrated at or below ambient temperature.

See other HALOGEN OXIDES

4102. Lead(II) chlorite

[13453-57-1]

$\text{Cl}_2\text{O}_4\text{Pb}$



Alone, or Antimony sulfide, or Sulfur

Mellor, 1941, Vol. 2, 283

It explodes on heating above 100°C or on rubbing with antimony sulfide or fine sulfur.

Non-metals

Pascal, 1960, Vol. 16, 264

Carbon, red phosphorus, or sulfur are oxidised violently by the chlorite.

See other CHLORITE SALTS, METAL OXOHALOGENATES

4103. Disulfuryl dichloride

[7791-27-7]

$\text{Cl}_2\text{O}_5\text{S}_2$



Phosphorus

See Phosphorus: Non-metal halides

Water

Sveda, M., *Inorg. Synth.*, 1950, **3**, 127


The pure acid chloride reacts slowly with water, but if more than a few % of chlorosulfuric acid (a usual impurity) is present, the reaction is rapid and could become violent with large quantities.

See other ACYL HALIDES, NON-METAL HALIDES (AND THEIR OXIDES) *See related* ACID ANHYDRIDES

4104. Chloryl perchlorate

[12442-63-6]

Cl_2O_6



The value of the heat of formation quoted for the monomeric form, chlorine trioxide, indicates considerable endothermicity; (ΔH_f° (g) +154.8 kJ/mol, 1.85 kJ/g).

Organic matter, or Water

1. Mellor, 1956, Vol. 2, Suppl. 1, 539

2. Wechsberg, M. *et al.*, *Inorg. Chem.*, 1972, **11**, 3066

3. Sorbe, 1968, 45

4. Pascal, J. L. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1985, 297

Though the least explosive of the chlorine oxides, being insensitive to heat or shock, it is a very powerful oxidant and needs careful handling. It violently or explosively oxidises ethanol, stopcock grease, wood and organic matter generally, and explodes on contact with water [1]. It explodes on heating [2], and explosions in contact with organic matter below -70°C have been reported [3]. A later reference indicates that it is shock-sensitive, however [4], and stresses the need for care in its use as a most effective perchlorylating agent.

Tetrachlorosilane, or Tetrabromosilane

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

Reaction gives explosive solids, apparently perchloratosiloxanes.

See Tetra-perchloratosilicon

Thionyl chloride


Schmeisser, M. *Angew. Chem.*, 1955, **67**, 499

It is liable to explode on contact with thionyl chloride in absence of inert solvent.

See other ENDOTHERMIC COMPOUNDS, HALOGEN OXIDES, OXIDANTS

4105. Lead(II) chlorate

[10294-47-0]



Bailar, 1973, Vol. 2, 130

Thermal decomposition may be explosive.

See other METAL CHLORATES

4106. Zinc chlorate

[10361-95-2]



Sorbe, 1968, 158

The tetrahydrated salt decomposes explosively at 60°C .

Other reactants

Yoshida, 1980, 63–64

MRH values calculated for 30 combinations with oxidisable materials are given.

Sulfides

MRH values below reference

Mellor, 1956, Vol. 2, Suppl. 1, 584

MRH Copper(II) sulfide 2.84/38, antimony(III) sulfide 2.89/41, tin(II) sulfide 3.05/43, tin(IV) sulfide 3.1/37


Interaction with copper(II) sulfide is explosive, and incandescent with antimony(III) sulfide, arsenic(III) sulfide, tin(II) sulfide and tin(IV) sulfide.

See other METAL CHLORATES

4107. Perchloryl perchlorate (Dichlorine heptaoxide)

[10294-48-1]

Cl_2O_7



1. Mellor, 1956, Vol. 2, Suppl. 1, 542
2. Schmeisser, M., *Angew. Chem.*, 1955, **67**, 498

It is rather endothermic (ΔH_f° (g) +265.3 kJ/mol, 1.45 kJ/g) and explodes violently on impact or rapid heating, but is a less powerful oxidant than other chlorine oxides [1]. It is now formulated as perchloryl perchlorate [2].

Bromine pentafluoride

Lawless, 1968, 173

Mixtures are shock-sensitive explosives.

Copper phthalocyanine

Osadchaya, L. I. *et al.*, *Chem. Abs.*, 1997, **126**, 309255v

A shock-sensitive explosive complex is formed, which could not be characterised.

Iodine

Kirk-Othmer, 1964, Vol. 5, 5

Explosion on contact.

See other ENDOTHERMIC COMPOUNDS, HALOGEN OXIDES

4108. Lead perchlorate

[13637-76-8]

$\text{Cl}_2\text{O}_8\text{Pb}$



Methanol

Willard, H. H. *et al.*, *J. Amer. Chem. Soc.*, 1930, **52**, 2396

A saturated solution of anhydrous lead perchlorate in dry methanol exploded violently when disturbed. Methyl perchlorate may have been involved.

See ALKYL PERCHLORATES *See other* METAL PERCHLORATES

4109. Tin(II) perchlorate

[25253-54-7]

$\text{Cl}_2\text{O}_8\text{Sn}$



Bailar, 1973, Vol. 2, 76

The trihydrated salt decomposes explosively at 250°C.

Aromatic Schiff bases

Dwivedi, B. K. *et al.*, *Indian J. Chem.*, 1987, **26A**, 618–620

Twelve tris-complexes [$\text{Sn}(\text{ClO}_4)_2 \cdot 3\text{L}$] between tin(II) perchlorate and Schiff bases derived from salicylaldehyde, anisaldehyde, or 2-hydroxy-1-naphthaldehyde and aromatic amines were investigated for thermal instability.

See related AMMINEMETAL OXOSALTS
See other METAL PERCHLORATES, REDOX COMPOUNDS

4110. Xenon(II) perchlorate (Bis(perchloryloxy)xenon)
[25523-79-9] **Cl₂O₈Xe**
Xe(ClO₄)₂

1. Wechsberg, M. *et al.*, *Inorg. Chem.*, 1972, **11**, 3066
2. Kirk Othmer, 1980, **Vol. 12**, 293
During preparation from perchloric acid and xenon difluoride at -50°C , violent explosions occurred if the reaction mixture was allowed to warm up rapidly [1]. Detonates easily and should be handled with extreme care [2].
See other XENON COMPOUNDS, NON-METAL PERCHLORATES

4111. Uranyl perchlorate
[13093-00-0] **Cl₂O₁₀U**
O₂U(ClO₄)₂

Ethanol
Erametsa, O., *Suomen Kemist*, 1942, **15B**, 1
Attempted recrystallisation of the salt from ethanol caused an explosion (probably involving ethyl perchlorate).
See ALKYL PERCHLORATES *See other* METAL PERCHLORATES

4112. Lead chloride
[7758-95-4] **Cl₂Pb**
PbCl₂

Calcium
See Calcium: Lead chloride *See other* METAL HALIDES

4113. Sulfur dichloride
[10545-99-0] **Cl₂S**
ClSCl

(MCA SD-77, 1960); HCS 1980, 875

Acetone
Fawcett, F. S. *et al.*, *Inorg. Synth.*, 1963, **7**, 121
Acetone is an effective solvent for cleaning traces of sulfur chlorides from reaction vessels, but care is necessary, as the reaction is vigorous if more than a trace is present.

Dimethyl sulfoxide
See Dimethyl sulfoxide: Acyl halides, etc.

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Metals

See Potassium: Non-metal halides, and Sodium: Non-metal halides

Oxidants

See Dinitrogen pentoxide: Sulfur dichloride

Nitric acid: Sulfur halides

Perchloryl fluoride: Hydrocarbons, etc.

Toluene

1. Caporossi, J. C., *Chem. Eng. News*, 1988, **66**(32), 2

2. Author's comments, 1988

3. Urben, P. G., private comm., 1989

Toluene was added to a closed mild steel storage tank containing sulfur dichloride, and the vessel was subsequently ruptured by overpressurisation. It was later found that the exothermic reaction between toluene and sulfur dichloride was catalysed by iron or iron(III) chloride [1]. Several reactions may have been involved in this chlorination system, many capable of iron catalysis arising from the inevitable formation of traces of iron(III) chloride from contact of sulfur dichloride with mild steel. Initially at low temperatures, exothermic formation of 2- and 4-chlorotoluene with evolution of hydrogen chloride and formation of more iron(III) chloride would occur. As the temperature and concentration of catalyst rose, side-chain chlorination would set in, leading to formation of chloromethyl-, dichloromethyl- and trichloromethyl-benzenes and much more heat and hydrogen chloride. The direct catalysed reaction of toluene with sulfur dichloride to form toluenesulfonyl chloride could also occur [2], with the subsequent possibility of oxidation to the disulfide [3]. It is also known that chloromethylbenzene (benzyl chloride) will undergo a Friedel-Crafts self-condensation reaction, catalysed by iron(III) chloride, to form polybenzyls. Both of these reactions would again liberate hydrogen chloride. With so many possibilities for evolution of hydrogen chloride gas, it is perhaps not surprising that the mild steel vessel ruptured [2].

See Benzyl chloride: Catalytic impurities

Water

MCA SD-77, 1960

Exothermic reaction with water or steam.

See other NON-METAL HALIDES

4114. Disulfur dichloride

[10025-67-9]

CISSCI

Cl₂S₂

(*MCA SD-77*, 1960); *HCS 1980*, 874

Aluminium

See Aluminium: Non-metal halides

Antimony, or Antimony sulfide, or Arsenic sulfide

Mellor, 1947, Vol. 10, 641

Interaction at ambient temperature is surprisingly energetic.

Chromyl chloride

See Chromyl chloride: Non-metal halides

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides, etc.

Mercury oxide

Mellor, 1947, Vol. 10, 643

Interaction is rapid and very exothermic.

Phosphorus(III) oxide

See Tetraphosphorus decaoxide: Disulfur dichloride

Potassium

Mellor, 1947, Vol. 10, 642

A mixture of potassium and the liquid chloride is shock-sensitive and explodes violently on heating.

Sodium peroxide

See Sodium peroxide: Non-metal halides

Tin

Pascal, 1963, Vol. 8.3, 309

Interaction is violent.

Unsaturated materials

1. Mellor, 1947, Vol. 10, 64–642

2. Huestis, B. L., *Safety Eng.*, 1927, **54**, 95

Alkenes, terpenes and unsaturated glycerides react exothermically, some vigorously [1]. Ignition may occur with some organic materials [2].

Water

MCA SD-77, 1960

As with sulfinyl chloride, the exothermic reaction with limited amounts of water may be dangerously violent under confinement because of rapid gas evolution.

See other GAS EVOLUTION INCIDENTS, NON-METAL HALIDES

4115. Diselenium dichloride
[10025-68-0]

ClSeSeCl

Cl₂Se₂

Alkali metals or oxides

See Potassium: Non-metal halides

Potassium dioxide: Diselenium dichloride

Sodium: Non-metal halides

Sodium peroxide: Non-metal halides

Aluminium

See Aluminium: Non-metal halides

Trimethylsilyl azide

See Trimethylsilyl azide: Selenium halides

See other NON-METAL HALIDES

4116. Tin(II) chloride
[7772-99-8]

SnCl₂

Cl₂Sn

HCS 1980, 900

Bromine trifluoride

See Bromine trifluoride: Tin(II) chloride

Calcium acetylide

MRH 1.51/25

See Calcium acetylide: Tin(II) chloride

Hydrazine

Mellor, 1941, Vol. 7, 430

Stannous chloride reacts with hydrazine in ethanol to give a precipitate, believed the dihydrazino adduct, which explodes on heating.

Hydrogen peroxide

MRH 1.21/15

See Hydrogen peroxide: Tin(II) chloride

Metal nitrates

MRH Ammonium nitrate 1.21/30

See METAL NITRATES: esters, etc.

Other reactants

Yoshida, 1980, 56

MRH values calculated for 16 combinations, largely with oxidants, are given.

See other METAL HALIDES, REDUCANTS

4117. Titanium(II) chloride
[10049-06-6]



1. Mellor, 1941. Vol. 7, 75
2. Brauer, 1965, Vol. 2, 1187
3. *NSC Data Sheet 485*, 1966

It readily ignites in air, particularly if moist [1,2]. The dichloride on heating under inert atmospheres disproportionates into the tetrachloride and pyrophoric titanium [3].

Water

Gallais, 1957, 431

Interaction at ambient temperature is violent, and water is reduced with evolution of hydrogen.

See other METAL HALIDES, REDUCANTS

4118. Vanadium dichloride
[10580-52-6]



Platinum, Water

Ephraim, 1954, 261

Vanadium(II) chloride dissolved in water is slowly oxidised by the solvent with evolution of hydrogen. Contact with platinum foil accelerates the reaction to violence.

See other METAL HALIDES, REDUCANTS

4119. Tungsten dichloride
[13470-12-7]



Preparative hazard

Mussell, R. D. *et al.*, *Inorg. Chem.*, 1990, **29**(19), 3711

Explosions sometimes resulted on opening sealed tubes in which complex mixed chloride salts of this compound had been prepared. There is no obvious source of pressure in the reaction mixture of tungsten hexachloride, sodium and aluminium chlorides and aluminium metal.

See other METAL HALIDES, REDUCANTS

4120. Zinc chloride
[7646-85-7]



HCS 1980, 965

Zinc

See Zinc: Zinc chloride
See other METAL HALIDES

4121. Zirconium(II) chloride
[13762-26-0]



Sidgwick, 1950, 652
When warm it ignites in air.
See other METAL HALIDES, REDUCANTS

4122. Cobalt(III) chloride
[10241-04-4]




Pentacarbonyliron, Zinc
See Pentacarbonyliron: Transition metal halides, etc.
See other METAL HALIDES

4123. Pentaamminechlorocobalt(III) perchlorate
[15156-18-0]



Explodes at 320°C; high impact sensitivity.
See entry AMMINOMETAL OXOSALTS

4124. Pentaammineaquacobalt(III) chlorate
[13820-81-0]



Salvadori, R., *Gazz. Chim. Ital.* [2], 1910, **40**, 9
It is of zero oxygen balance and explodes on impact, or heating to 130°C.
See entry OXYGEN BALANCE See other AMMINOMETAL OXOSALTS

4125. Hexaamminecobalt(III) chlorate
[26156-56-9]



Friederich, W. *et al.*, *Z. ges. Schiess-Sprengstoffw.*, 1926, **21**, 49
The complex salt is explosive.
See other AMMINOMETAL OXOSALTS

4126. Hexaamminecobalt(III) perchlorate

[13820-83-2]

$\text{Cl}_3\text{CoH}_{18}\text{N}_6\text{O}_{12}$



It explodes at 360°C, is highly impact-sensitive and has +14.2% oxygen balance.

See entries AMMINEMETAL OXOSALTS, OXYGEN BALANCE

4127. Chromium(III) chloride

[10025-73-7]

Cl_3Co



Lithium, Nitrogen

See Lithium: Metal chlorides, etc.

See other METAL HALIDES

4128. Bis(hydrazine)diperchloratochromium(III) perchlorate

[74311-45-7]

$\text{Cl}_3\text{CrH}_8\text{N}_4\text{O}_{12}$



Patil, K. C. *et al.*, *Chem. & Ind.*, 1979, 902

It ignites at 240°C in air and decomposes explosively, but surprisingly is not sensitive to friction or impact, unlike its nickel analogue. It shows +118% oxygen balance.

See entry OXYGEN BALANCE See other AMMINEMETAL OXOSALTS, REDOX COMPOUNDS

4129. Hexaamminechromium(III) perchlorate

[15203-80-2]

$\text{Cl}_3\text{CrH}_{18}\text{N}_6\text{O}_{12}$



Ethanol, Ether

1. Pennington, D. E., *Chem. Eng. News*, 1982, **60**(33), 55

2. Cartwright, R. V., *Chem. Eng. News*, 1983, **61**(6), 4

A second crop of material (+14.2% oxygen balance), precipitated by addition of excess perchloric acid to the liquor, was collected on a sinter, washed and dried with ethanol, then ether. Stirring the filter cake led to a violent explosion [1]: formation of ethyl perchlorate seems to have been the most likely cause. The advantages of using plastic sintered funnels in reducing friction and the extent of fragmentation if an explosion occurs were stressed later [2].


See ALKYL PERCHLORATES See entry OXYGEN BALANCE

See other AMMINEMETAL OXOSALTS, FRICTIONAL INITIATION INCIDENTS

4130. Chromium(III) perchlorate . 6dimethyl sulfoxide

[59675-70-6]

$\text{Cl}_3\text{Cr} \cdot 6\text{C}_2\text{H}_6\text{OS}$



Langford, C. H. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1977, 139

The hexasolvated complex may be hazardous with respect to detonation. Other metal perchlorates and nitrates solvated with DMSO are explosive.

See Dimethyl sulfoxide: Metal oxosalts

See other METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS

4131. Dysprosium perchlorate

[14017-53-9]

$\text{Cl}_3\text{DyO}_{12}$

$\text{Dy}(\text{ClO}_4)_3$

Birnbaum, E. R. *et al.*, *Inorg. Chem.*, 1973, **12**, 379

The anhydrous salt was stable to 300°C or to impact of a steel hammer, but a mild explosion occurred when a grease-containing sample was disturbed with a metal spatula.

See other FRICTIONAL INITIATION INCIDENTS, METAL PERCHLORATES

4132. Erbium perchlorate

[14017-55-1]

$\text{Cl}_3\text{ErO}_{12}$

$\text{Er}(\text{ClO}_4)_3$

Acetonitrile

Wolsey, W. C., *J. Chem. Educ.*, 1973, **50**(6), A336–337

The shock-sensitive glassy residue (containing traces of acetonitrile), left after heating the tetrasolvated salt to above 150°C on a vacuum line, exploded violently when scraped with a spatula. (Oxygen balance of salt +700%).

See other FRICTIONAL INITIATION INCIDENTS, METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS

4133. Iron(III) chloride

[7705-08-0]

Cl_3Fe

FeCl_3

HCS 1980, 501

Chlorine, Monomers

See Chlorine: Iron(III) chloride, Monomers

Ethylene oxide

See Ethylene oxide: Contaminants

Metals

See Potassium: Metal halides

Sodium: Metal halides

See other METAL HALIDES

4134. Iron(III) perchlorate
[13537-24-1]



Acetonitrile

Bancroft, G. M. *et al.*, *Can. J. Chem.*, 1974, **52**, 783

The violent reaction which occurred on dissolution of the anhydrous salt in acetonitrile did not occur with the hydrated salt.

Dimethyl sulfoxide

See Dimethyl sulfoxide: Metal oxosalts

See other METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS

4135. Gallium perchlorate
[17835-81-3]



1. Foster, L. S., *J. Amer. Chem. Soc.*, 1933, **61**, 3123

2. Foster, L. S., *Inorg. Synth.*, 1946, **2**, 28

During preparation by dissolving the metal in 72% perchloric acid, the hexahydrate separates as a crystalline solid. After filtration, the damp crystals must not contact any organic material (filter paper, horn spatula, or recrystallisation solvent) as the adherent perchloric acid is above 72% concentration owing to the hexahydrate formation [1,2].

See Perchloric acid: Dehydrating agents

See other METAL PERCHLORATES

See related SOLVATED OXOSALT INCIDENTS

†**4136. Trichlorosilane**
[10025-78-2]



HCS 1980, 924

1. Müller, R. *et al.*, *J. Prakt. Chem.*, 1966, **31**, 1–6

2. Britton, L. G., *Plant/Oper. Progr.*, **9**(1), 16

The pure material is not impact-ignitable in absence of electrostatic charges, but technical material (possibly containing dichlorosilane) is [1]. Hazards are reviewed and an Ait of 182°C established [2].

See other STATIC INITIATION INCIDENTS

Acetonitrile, Diphenyl sulfoxide

Benkeser, R. A., *Chem. Eng. News*, 1978, **56**(32), 107

Use of a single portion of trichlorosilane added to acetonitrile at 10°C to reduce diphenyl sulfoxide led to a violent explosion. The reaction previously had been effected uneventfully in a wide range of other solvents. The explosion was

attributed to use of acetonitrile as solvent and/or the addition of the trichlorosilane as a single portion.

Other reactants

Yoshida, 1980, 252

MRH values calculated for 13 combinations with oxidants are given.

Styrene

Anon., *Chem. Eng. News*, 1994, **72**(26), 6

A drum of these two, apparently freshly mixed as feed for a reaction, pressurised and exploded. The subsequent fire destroyed the plant and injured 31 people. Styrene is polymerisable and the two probably react by Friedel-Crafts type chemistry, if suitably catalysed. It is not reported whether the drum was lined, plain steel, or galvanised.

See other HALOSILANES

4137. Pentaamminechlororuthenium chloride

[18532-87-1]

Cl₃H₁₅N₅Ru



Sodium azide

Allen, A. D. *et al.*, *Inorg. Synth.*, 1970, **12**, 5–6

During treatment with sodium azide of an intermediate (believed to be pentaammineaquaruthenium(III) derived from the title compound) to produce pentaaminedinitrogenruthenium(II) solutions, a dangerously explosive red solid may be produced. The solid, pentaammineazidoruthenium(III) will, however, decompose on standing to give the desired dinitrogen species.


See Hydrazine: Ruthenium(III) chloride

See other AMMINOMETAL HALIDES See related METAL HALIDES

4138. Octaammine- μ -hydroxy[μ -(peroxy-*O* : *O'*)]dirhodium(3+) perchlorate

[113728-41-9]

Cl₃H₂₅N₈O₁₅Rh₂



Springborg, J. *et al.*, *Acta Chem. Scand. Sec. A*, 1987, **A41**, 485
The complex may detonate when subjected to mechanical shock.
See other AMMINEMETAL OXOSALTS

4139. Iodine trichloride
[865-44-1]



Phosphorus

Pascal, 1960, Vol. 16.1, 578

Phosphorus ignites on contact with iodine trichloride.

See other INTERHALOGENS

4140. Iodine(III) perchlorate
[38005-31-1]



Christe, K. O. *et al.*, *Inorg. Chem.*, 1972, **11**, 1683

The solid was stable at -45°C but exploded on laser irradiation at low temperature.

See other IODINE COMPOUNDS, IRRADIATION DECOMPOSITION INCIDENTS, NON-METAL PERCHLORATES

4141. Manganese trichloride oxide
[23097-77-0]




Briggs, T. S., *J. Inorg. Nucl. Chem.*, 1968, **30**, 2867–2868

Explosively unstable if isolated as a liquid at ambient temperature, it may be handled safely in carbon tetrachloride solution.

See Manganese chloride trioxide, Manganese dichloride dioxide

See related METAL HALIDES, METAL OXIDES

4142. 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazinemolybdenum
[]



Water

Wynne, K. J. *et al.*, *J. Inorg. Nucl. Chem.*, 1968, **30**, 2851

A 2 g quantity of the product of interaction of hexacarbonylmolybdenum and thiazyl chloride, possibly constituted as shown, reacted explosively with water.

See other N-HALOGEN COMPOUNDS, N-S COMPOUNDS

4143. Nitrogen trichloride (Trichloramine)

[10025-85-1]

Cl₃N

NCl₃

HCS 1980, 686

1. Mellor, 1940, Vol. 8, 598–604; 1967, Vol. 8, Suppl. 2.2, 411
2. Sidgwick, 1950, 705
3. Anon., *ABCM Quart. Safety Summ.*, 1946, **17**, 17
4. Brauer, 1963, Vol. 1, 479
5. Schlessinger, G. G., *Chem. Eng. News*, 1966, **44**(33), 46
6. Kovacic, P. *et al.*, *Org. Synth.*, 1968, **48**, 46
7. Dokter, T., private comm., 1985
8. Dokter, T., *J. Haz. Mat.*, 1985, **10**, 205–222

Contact above 0°C of excess chlorine or a chlorinating agent with aqueous ammonia, ammonium salts or a compound containing a hydrolysable amino-derivative, or electrolysis of ammonium chloride solution produces the highly endothermic (ΔH_f° (g) +230.1 kJ/mol, 1.91 kJ/g) and explosive nitrogen trichloride as a water-insoluble yellow oil [1,2,3]. It is usually prepared [4] in solution in a solvent, and such solutions in chloroform are reported as stable up to 18% concentration. However, an 18% solution in dibutyl ether exploded on cooling in a refrigerator, and a 12% solution prepared without cooling had decomposed vigorously [5]. In absence of other materials, explosion of the trichloride may be initiated in a wide variety of ways. The solid frozen under vacuum in liquid nitrogen explodes on thawing, and the liquid explodes on heating to 60 or 95°C. Exposure to impact, light or ultrasonic irradiation will cause (or sensitise) detonation [1]. The preparation and synthetic use of solutions of nitrogen trichloride in dichloromethane is described in detail [6]. Trichloramine is surprisingly stable, can be formed at 80°C and above, and has been made in 200 kg lots for possible use as a chlorinating agent [7]. The formation and separation of the trichloride have been described in detail [8].

See Chlorine: Nitrogen compounds, or: Dimethyl phosphoramidate

Phosphorus pentachloride: Urea

Sodium chloride: Nitrogen compounds

See other IRRADIATION DECOMPOSITION INCIDENTS

Chlorine, Hydrogen

Ashmore, P. G., *Nature*, 1953, **172**, 449–450

Equimolar mixtures of chlorine and hydrogen containing 0.1–0.2% of the trichloride will explode in absence of light if the pressure is below a limiting value dependent on temperature (30 mbar at 20°, 132 mbar at 57°C).

Initiators

1. Mellor, 1940, Vol. 8, 601–604; 1967, Vol. 8, Suppl. 2.2, 412
2. Pascal, 1956, Vol. 10, 264

A wide variety of solids, liquids and gases will initiate the violent and often explosive decomposition of nitrogen trichloride. These include conc. ammonia, arsenic, dinitrogen tetroxide (above -40°C , with more than 25% solutions of trichloride in chloroform), hydrogen sulfide, hydrogen trisulfide, nitrogen oxide, organic matter (including grease from the fingers), ozone, phosphine, phosphorus (solid, or in carbon disulfide solution), potassium cyanide (solid, or aqueous solution), potassium hydroxide solution or selenium [1]. All four hydrohalide acids will also initiate explosion of the trichloride [2].

See other ENDOTHERMIC COMPOUNDS, *N*-HALOGEN COMPOUNDS

4144. Nitrosylruthenium trichloride [18902-42-6]




Sidgwick, 1950, 1486

It decomposes violently at 440°C .

See other NITROSO COMPOUNDS, *N*-METAL DERIVATIVES

4145. 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazine (Thiazyl chloride) [18428-81-4]



Bailar, 1973, Vol. 2, 908

It explodes violently on sudden heating.

Ammonia, Silver nitrate

Goehring, 1957, 67

Thiazyl chloride, treated with aqueous ammonia and then silver nitrate, gives a compound AgN_5S_3 (of unknown structure) which is shock-sensitive and explodes violently.

Hexacarbonylmolybdenum


See Carbonyl(pentasulfur pentanitrido)molybdenum

1,3,5-Trichlorotrithiahexahydro-1,3,5-triazinemolybdenum

See other *N*-HALOGEN COMPOUNDS, *N*-S COMPOUNDS

4146. Tin azide trichloride

[]



An explosive solid.

See entry METAL AZIDE HALIDES**4147. Titanium azide trichloride**



[]



An explosive solid.

See entry METAL AZIDE HALIDES**4148. Neodymium perchlorate . 2acetonitrile**

[13498-06-1]



1. Forsberg, J. H. *et al.*, *Inorg. Chem.*, 1971, **10**, 2656
2. Eigenbrot, C. W. *et al.*, *Inorg. Chem.*, 1982, **21**, 2867
3. Raymond, K. N., *Chem. Eng. News*, 1983, **61**(49), 4
4. Bretherick, L., *Chem. Eng. News*, 1983, **61**(50), 2

The perchlorate, prepared from neodymium oxide and perchloric acid [1] was purified and isolated as the salt tetrasolvated with acetonitrile [2], which has not been found to be shock-sensitive. On vacuum drying at 80°C/24h, it is converted to the disolvated salt which is very shock-sensitive and a 1 g sample exploded violently in a hand-held flask [3]. Other examples of partially desolvated perchlorate salts becoming friction- and shock-sensitive are known [4]. (The disolvate is very close to zero oxygen balance).

See other METAL PERCHLORATES, SOLVATED OXOSALT INCIDENTS**4149. Phosphoryl chloride**

[10025-87-3]



(MCA SD-26, 1968); HCS 1980, 751

Carbon disulfide

MRH Carbon disulfide 1.17/99+

Disposal of a benzene solution of phosphoryl chloride into a waste drum containing carbon disulfide (and other solvents) caused an instantaneous reaction, with evolution of (probably) hydrogen chloride. Presence of a hydroxylic compound seems likely.

Dimethylaniline

Anon., private comm., 1984

Cl₃OP

In a chlorination reaction using phosphoryl chloride with dimethylaniline as acid acceptor, the reagents were added all at once. After an induction period, most of the mixture was ejected from the flask. On a smaller scale, the reaction had been uneventful.

Dimethylformamide, 2,5-Dimethylpyrrole

MCA Case History No. 1460

Poor stirring during formylation of 2,5-dimethylpyrrole with the preformed complex of dimethylformamide with phosphoryl chloride caused eruption of the flask contents. Reaction of the complex with a local excess of the pyrrole may have been involved.

2,6-Dimethylpyridine *N*-oxide

1. Kato, T. *et al.*, *J. Pharm. Soc. Japan.*, 1951, **71**, 217
2. Anon., *Chem. Eng. News*, 1965, **43**(47), 40
3. Evans, R. F. *et al.*, *J. Org. Chem.*, 1962, **27**, 1333

Interaction of the reagents in absence of diluent, according to a published procedure [1], caused an explosion [2]. The use of a chlorinated solvent as diluent prevented explosion, confirming an earlier report [3].

See Pyridine *N*-oxide, below

Dimethyl sulfoxide

MRH 1.80/99+

See Dimethyl sulfoxide: Acyl halides

Ferrocene-1,1'-dicarboxylic acid

Marvel, C. S., *Chem. Eng. News*, 1962, **40**(3), 55

An explosion occurred immediately after pouring and capping of the chloride recovered from preparation of ferrocene-1,1'-dicarbonyl chloride. The storage bottle contained phosphoryl chloride recovered from similar preparations and which had been stored for some 3 months. No explanation was apparent.

Other reactants

Yoshida, 1980, 76

MRH values calculated for 10 combinations with various reagents are given.

Pyridine *N*-oxide

MCA Guide 1972, 321

Dropwise addition of the chloride caused a steady exotherm to 60–65°C when a runaway reaction accelerated to explosion.

See 2,6-Dimethylpyridine *N*-oxide, above

Sodium

MRH 4.27/31

See Sodium: Non-metal halides (reference 7)

Water

1. *MCA SD-26*, 1968
2. Unpublished observations

3. MCA Case History No. 1274
4. MCA Case History No. 2286
5. Anon., *Loss Prev. Bull.*, 1972, (013), 22–23
6. Anon., *Loss Prev. Bull.*, 1980, (038), 27
7. Anon., *Sichere Chemiearb.*, 1985, **37**(11), 130
8. Davies, R., *Loss Prev Bull.*, 1995, 121, 13

The hazards arising from interaction of phosphoryl chloride and water derive from there being often a considerable delay in onset of the exothermic hydrolysis reaction, which may proceed with enough vigour to generate steam and liberate hydrogen chloride gas. Conditions tending to favour delayed or violent reaction include limited quantities of water and/or ice for hydrolysis, lack of stirring, cold or frozen phosphoryl chloride, and reaction in closed or virtually closed containers [1]. A layer of the dense and cold liquid may survive for several minutes under water before violent, almost instantaneous hydrolysis occurs, particularly when disturbed [2]. The Case History describes a violent explosion which occurred when water was added to a drum containing some phosphoryl chloride which had been stored below its freezing point, 2°C [3]. Bursting of a 300 kg drum during filling was attributed to delayed reaction of the chloride with water contamination in the drum [4]. Methods for the safe decontamination of non-returnable drums which have contained the chloride are detailed, and involve thorough draining of the drum and internal inspection before careful application of large amounts of water [5]. When 10 l of the chloride was sucked into a measuring vessel, a violent explosion occurred which ruptured it. It was assumed that water previously used to clean the vessel had not fully drained out [6]. In outdoor storage, rainwater found its way into the space between the outer container and the polythene liner containing solid methylchlorouracil. When this was tipped into phosphoryl chloride in a reactor, some 4–5 l of water also ran in and caused an explosive evolution of hydrogen chloride gas which led to a fatality [7]. An account of an explosion when a transfer hose largely full of phosphoryl chloride was flushed with water is given [8]

See other GAS EVOLUTION INCIDENTS, PLANT CLEANING INCIDENTS

Zinc

MRH 1.34/39

Mellor, 1940, Vol. 8, 1025

Zinc dust ignites in contact with a little phosphoryl chloride, and subsequent addition of water liberates phosphine which ignites.

See other NON-METAL HALIDES (AND THEIR OXIDES)

4150. Antimony trichloride oxide (Antimonyl chloride)
[14459-54-2]

Cl₃OSb

O:SbCl₃

Bromine trifluoride

See Bromine trifluoride: Antimony trichloride oxide

See related METAL HALIDES, METAL OXIDES

4151. Vanadium trichloride oxide (Vanadyl chloride)

[7727-18-6]

Cl_3OV

$\text{O}:\text{VCl}_3$

HCS 1980, 953

Bailar, 1973, Vol. 3, 508

The chloride (and analogous bromide) is violently hygroscopic.

Rubidium

See Rubidium: Vanadium trichloride oxide

Sodium

Quam, G. N. *et al.*, in *Safety in the Chemical Laboratory*, (Ed. Steere, N. V.), Vol. 1, 1967, 82 (expanded articles reprinted from *J. Chem. Educ.*, 1964–67)

In a procedure to purify the chloride oxide with sodium, too-rapid addition of the metal led to a violent explosion.

See Sodium: Metal halides (reference 2)

See related METAL HALIDES, METAL OXIDES

4152. Vanadyl perchlorate

[63672-69-3]

$\text{Cl}_3\text{O}_{13}\text{V}$

$\text{O}:\text{V}(\text{ClO}_4)_3$

Alone, or Organic solvents

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

A powerful oxidant which may explode above 80°C, and which ignites organic solvents on contact. It decomposes, sometimes explosively, on exposure to light.

See other METAL PERCHLORATES, OXIDANTS

4153. Phosphorus trichloride

[7719-12-2]

Cl_3P

PCl_3

(*MCA SD-27, 1972*); *HCS 1980, 749*

Carboxylic acids

1. Coghill, R. D., *J. Amer. Chem. Soc.*, 1938, **60**, 488

2. Peacocke, T. A., *School Sci. Rev.*, 1962, **44**(152), 217

3. Scrimgeour, C. M., *Chem. Brit.*, 1975, **11**, 267

4. Taylor, D. A. H. *et al.*, *Chem. Brit.*, 1976, **12**, 105

5. Bretherick, L., *Chem. Brit.*, 1976, **12**, 26

Use of a free flame instead of a heating bath to distil acetyl chloride produced from acetic acid and phosphorus trichloride caused the residual phosphonic ('phosphorous') acid to decompose violently to give spontaneously flammable phosphine

[1]. Two later explosions in the same preparative system after reflux but before distillation from a water-bath may have been due to ingress of air into the cooling flask and ignition of traces of phosphine [2]. During preparation of furoyl chloride, the excess trichloride was distilled off at atmospheric pressure, then vacuum was applied prior to intended distillation at 100°C/100 mbar, and an explosion occurred shortly after [3]. A similar incident occurred during conversion of propionic acid to the chloride [4]. These incidents were attributed to thermal decomposition of the by-product phosphonic acid to give a phosphine–diphosphane mixture and leakage of air into the evacuated system to produce a spontaneously explosive mixture. Sulfinyl chloride is a safer chlorinating agent for carboxylic acids [5].

Chlorobenzene, Sodium

See Sodium: Chlorobenzene, etc.

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides, etc.

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

Hydroxylamine

See Hydroxylamine: Phosphorus chlorides

Metals

MRH Potassium 2.80/46, sodium 3.10/33

Mellor, 1940, Vol. 8, 1006; 1941, Vol. 2, 470

Potassium ignites in phosphorus trichloride, while molten sodium explodes on contact.

See Aluminium: Non-metal halides

Nitroheterocycle

Hallam, S., 1993, personal communication

A multi-kg mixture of the trichloride and a nitroheterocycle obtained after deoxygenation of the corresponding N-oxide was being worked up by distillation under vacuum. It burst the glass vessel. Investigation showed that this was due to a higher temperature (85-90°C) than previously employed, permitting the phosphorus trichloride to react vigorously with the product.

Other reactants

Yoshida, 1980, 59

MRH values calculated for 8 combinations with various reagents are given.

Oxidants

MRH values show % of oxidant

See Chromium pentafluoride: Phosphorus trichloride

Chromyl chloride: Non-metal halides

Fluorine: Phosphorus halides

Iodine chloride: Phosphorus trichloride

Lead(IV) oxide: Non-metal halides

MRH 0.67/69

Cl₃P

Nitric acid: Phosphorus halides	MRH 0.63/31
Selenium dioxide: Phosphorus trichloride	
Sodium peroxide: Non-metal halides	MRH 2.55/46

Oxygen

Vorob'ev, N. I. *et al.*, *Izv. Akad. Nauk SSSR, Neorg. Mater.*, 1974, **10**, 2039–2042
Interaction to form phosphoryl chloride at 100–700°C was investigated, and the limiting concentrations to prevent explosions were determined.

Sodium Carbonate

Holmes, W. S., 1989, personal communication

This is often recommended for clearing up spillage. A dangerously vigorous reaction can result (probably depending upon the hydration level of the carbonate?). Large quantities of dry sand followed by addition to water are preferred.

See Water, below

Sulfur acids

Dillon, K. B. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1979, 883–885

The trichloride is initially insoluble in 100% sulfuric acid, fluorosulfuric acid or 25% oleum, but dissolves very exothermally after delays of 8, 1 and 0.5 h, respectively. Reaction with 65% oleum is immediately violent.

See Phosphorus tribromide: Sulfur acids *and* Phosphorus triiodide: Sulfur acids

See other INDUCTION PERIOD INCIDENTS

Tetravinyllead

See Tetravinyllead: Phosphorus trichloride

Water

1. *MCA SD-27*, 1972
2. Albright and Wilson, (Manufacturer's safety sheet)
3. Coghill, R. D., *J. Amer. Chem. Soc.*, 1938, **60**, 488
4. *MCA Case History No. 520*
5. Anon., *Loss. Prev. Bull.*, 1977, (013), 21–23

Reaction with water is exothermic and immediately violent (unlike phosphoryl chloride), and is accompanied by evolution of some diphosphane which ignites [1]. There can sometimes be an induction period, especially if cold. Yellow solids produced by hydrolysis may ignite spontaneously if dried [2]. Evaporation of the trichloride from an open beaker on a steam-bath led to ignition, which did not occur if a hotplate were used as a dry heat source [3]. Interaction of the trichloride and water in a virtually sealed container caused the latter to burst under the pressure of hydrogen chloride generated [4]: (1 vol. trichloride gives 830 vol. of hydrogen chloride at 20°C). Methods for the safe decontamination of non-returnable drums which have contained the chloride are detailed, and involve careful application of large amounts of water to the well-drained drum [5].

See other GAS EVOLUTION INCIDENTS *See other* NON-METAL HALIDES (AND THEIR OXIDES)

4154. Thiophosphoryl chloride
[3982-91-0]

Cl₃PS

S:PCl₃

Möller, T. *et al.*, *Inorg. Synth.*, 1954, **4**, 72

During the preparation from phosphorus trichloride and sulfur, the quantity and quality of the aluminium chloride catalyst is critical to prevent the exothermic reaction going out of control.

Methylmagnesium iodide

See Tetramethyldiphosphane disulfide

Pentaerythritol

MCA Case History No. 1315

On two occasions violent explosions occurred after heating of equimolar proportions of the reagents (for 4 h at 160°C according to a literature method) had been discontinued. (This suggests spontaneous ignition of traces of phosphine derivatives as air was drawn into the cooling reaction vessel).

See Phosphorus trichloride: Carboxylic acids (reference 5)

See related NON-METAL HALIDES

4155. Rhodium(III) chloride
[10049-07-7]

Cl₃Rh

RhCl₃

Pentacarbonyliron, Zinc

See Pentacarbonyliron: Transition metal halides

See other METAL HALIDES

4156. Ruthenium(III) chloride
[10049-08-8]

Cl₃Ru

RuCl₃

Hydrazine

See Hydrazine: Ruthenium(III) chloride

Pentacarbonyliron, Zinc

See Pentacarbonyliron: Transition metal halides

See other METAL HALIDES

4157. Antimony trichloride
[10025-91-9]

Cl₃Sb

SbCl₃

(*MCA SD-66*, 1957); *HCS 1980*, 164

Aluminium

See Aluminium: Antimony trichloride

N-Chlorodimethylamine

See N-chlorodimethylamine: Antimony chlorides

See other METAL HALIDES

4158. Titanium trichloride

[7705-07-9]



Cl_3Ti

HCS 1980, 905

Air, or Water

1. Lerner, R. W., *Ind. Eng. Chem.*, 1961, **53**(12). 56A

2. Ingraham, T. K. *et al.*, *Inorg. Synth.*, 1960, **5**, 56

It reacts vigorously with air and/or water (vapour or liquid) and adequate handling precautions are necessary [1]. The finely divided powder is pyrophoric in air [2].

See other METAL HALIDES, PYROPHORIC MATERIALS, REDUCANTS

4159. Vanadium trichloride

[7718-98-1]



Cl_3V

Methylmagnesium iodide

Cotton, F. A., *Chem. Rev.*, 1955, **55**, 560

Reaction of vanadium trichloride and similar halides with Grignard reagents is almost explosively violent under a variety of conditions.

See other METAL HALIDES, REDUCANTS

4160. Zirconium trichloride

[10241-03-9]



Cl_3Zr

Water

Pascal, 1963, Vol. 9, 540

Interaction is very violent, hydrogen being evolved.

See other METAL HALIDES, REDUCANTS

4161. Caesium tetraperchloratiodate

[53078-10-7]



$\text{Cl}_4\text{CsIO}_{16}$

Christe, K. O. *et al.*, *Inorg. Chem.*, 1972, **11**, 683

Though stable at ambient temperature, samples exploded under laser irradiation at low temperatures.

See other IODINE COMPOUNDS, IRRADIATION DECOMPOSITION INCIDENTS, NON-METAL PERCHLORATES

4162. Germanium tetrachloride

[10038-98-9]

Cl₄Ge

GeCl₄

Water

Mellor, 1941, Vol. 7, 270

Interaction is very exothermic, accompanied by crackling if the chloride is dropped into water.

See other METAL HALIDES

4163. Hafnium tetrachloride

[13499-05-3]

Cl₄Hf

HfCl₄

Tetrahydrofuran

Manzer, L. E., *Inorg. Synth.*, 1982, **21**, 137

Addition of the anhydrous chloride directly to THF causes a violent exothermic reaction. Add THF dropwise to a suspension of the chloride in dichloromethane.

See other METAL HALIDES

4164. Molybdenum diazide tetrachloride

[14259-66-6]

Cl₄MoN₆

(N₃)₂MoCl₄

Highly explosive.

See entry METAL AZIDE HALIDES

4165. Chloriminovanadium trichloride

[14989-38-9]

Cl₄NV

ClN:VCl₃

1. Strähle, J. *et al.*, *Angew. Chem.*, 1966, **78**, 450

2. Strähle, J. *et al.*, *Z. Anorg. Chem.*, 1965, **338**, 287

A new method of preparation from vanadium nitride and chlorine [1] is free of the explosion hazards of chlorine azide and vanadium azide tetrachloride present in an earlier method [2].

See other AMMINOMETAL HALIDES, N-HALOGEN COMPOUNDS

See related METAL HALIDES

4166. Vanadium azide tetrachloride


[]



Explosive solid.

See entry METAL AZIDE HALIDES**4167. Rhenium tetrachloride oxide**

[13814-76-1]





Ammonia

Sidgwick, 1950, 1302

Interaction with gaseous or liquid ammonia is violent.

See related METAL HALIDES, METAL OXIDES*See other* AMINATION INCIDENTS**4168. Diphosphoryl chloride**

[13498-14-1]




Water

Mellor, 1971, Vol. 8, Suppl. 3, 505

Hydrolysis is as vigorous as that of phosphorus pentoxide.

See other NON-METAL HALIDES**4169. Tetraperchloratosilicon**

[]




1. Sorbe, 1968, 127

2. Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

A highly explosive liquid [1]. Early attempts failed to isolate it but prepared numerous other explosive compounds. Reaction of dichlorine hexoxide with silicon tetrachloride or tetrabromide gave an explosive solid, apparently a perchlorato oligosiloxane. Silver perchlorate and silicon tetrahalides in ether gave explosive volatile organics, perhaps ethyl perchlorate. Replacing ether by acetonitrile as solvent, a solid (di)acetonitrile adduct of the tetraperchlorate precipitated, described as exceptionally explosive even in the smallest quantities [2].

See other NON-METAL PERCHLORATES

4170. Titanium tetraperchlorate
[60580-20-3]



Kirk Othmer, 4th. Edn., 1996, Vol. 18, 161
The compound sublimes from 70°C and explodes on heating to 130°C at atmospheric pressure.

Diethyl ether, or Formamide, or Dimethylformamide

Laran, R. J., US Pat. 3 157 464, 1964

Described as insensitive to heat or shock, this powerful oxidant explodes on contact with diethyl ether, and ignites with formamide or DMF.

See other METAL PERCHLORATES, OXIDANTS

4171. Tetrachlorodiphosphane
[13497-91-1]



Besson, A. *et al.*, *Compt. rend.*, 1910, **150**, 102

It oxidises rapidly in air, sometimes igniting.

See other HALOPHOSPHINES, NON-METAL HALIDES

4172. Lead tetrachloride
[13463-30-4]



1. Bailar, 1973, Vol. 2, 136, 1066

2. Kaufmann, G. B. *et al.*, *Inorg. Synth.*, 1983, **22**, 150

It may disproportionate explosively to lead(II) chloride and chlorine above 100°C [1]. Preparative precautions are detailed [2].

Potassium

See Potassium: Metal halides

Sulfuric acid

Friedrich, H., *Ber.*, 1893, **26**, 1434

It explodes on warming with diluted sulfuric acid or on attempted distillation from the concentrated acid in a stream of chlorine.

See other METAL HALIDES

4173. Tetrachlorosilane (Silicon tetrachloride)
[10026-04-7]



HCS 1980, 821

1462

Dimethyl sulfoxide

See Dimethyl sulfoxide: Acyl halides, etc.

Ethanol, Water

Skinner, S. J., *Loss. Prev.*, 1981, **14**, 178–180

In the preparation of ethyl polysilicate by mixing tetrachlorosilane and industrial methylated spirit containing some water, failure of the agitator is thought to have led to layering of the alcohol over the dense chloride. Evolution of hydrogen chloride led to mixing of the layers, and a greatly increased rate of reaction and self-accelerating gas evolution which burst the reactor.

See other AGITATION INCIDENTS, GAS EVOLUTION INCIDENTS

Sodium

See Sodium: Non-metal halides (reference 8)

See other HALOSILANES, NON-METAL HALIDES

4174. Tin(IV) chloride

[7646-78-8]

SnCl₄

Cl₄Sn

HCS 1980, 902

Alkyl nitrates

See ALKYL NITRATES: lewis acids

Ethylene oxide

See Ethylene oxide: Contaminants

Nitrobenzene

See Nitrobenzene: Tin(IV) chloride

Turpentine

Mellor, 1941, Vol. 7, 446

Interaction is strongly exothermic and may lead to ignition.

See other METAL HALIDES

4175. Tellurium tetrachloride

[10026-07-0]

TeCl₄

Cl₄Te

Ammonia

Mellor, 1943, Vol. 11, 58

Interaction with liquid ammonia at -15°C forms tellurium nitride (?), which explodes at 200°C .

See other NON-METAL HALIDES

4176. Titanium tetrachloride
[7550-45-0]



HCS 1980, 904

Lithium nitride

See Lithium nitride: Transition metal halides

Sulfur nitrides

See Disulfur dinitride: Metal chlorides

Tetrasulfur tetranitride: Metal chlorides

Tetrahydrofuran

Manzer, L. E., *Inorg. Synth.*, 1982, **21**, 135

Addition of the anhydrous chloride directly to THF causes a violent exothermic reaction. Add THF dropwise to a suspension of the chloride in dichloromethane.

Urea

Ionova, E. A. *et al.*, *Chem. Abs.*, 1966, **64**, 9219b

The liquid hexaurea complex formed during 6 weeks at 80°C decomposed violently at above 90°C. *N*-Halogen compounds may have been formed.

See other METAL HALIDES

4177. Vanadium tetrachloride
[7632-51-1]



Lithium nitride

See Lithium nitride: Transition metal halides

4178. Zirconium tetrachloride
[10026-11-6]



Ethanol, or Water

Rosenheim, A. *et al.*, *Ber.*, 1907, **40**, 811

Interaction with either is violent.

Lithium, Nitrogen

See Lithium: Metal chlorides, etc.

Tetrahydrofuran

Manzer, L. E., *Inorg. Synth.*, 1982, **21**, 136

Addition of the anhydrous chloride directly to THF causes a violent exothermic reaction. Add THF dropwise to a suspension of the chloride in dichloromethane.

See other METAL HALIDES

- 4179. Diamminedichloroaminotrichloroplatinum(IV)**
[18815-16-2] **Cl₅H₆N₃Pt**
[(H₃N)₂Pt(NCl₂)Cl₃]
- Kukushkin, Yu. N., *Chem. Abs.*, 1958, **52**, 13509i
 It tended to decompose violently.
See other N-HALOGEN COMPOUNDS, PLATINUM COMPOUNDS
-
- 4180. Molybdenum pentachloride**
[10241-05-1] **Cl₅Mo**
MoCl₅
- Sodium, or Sodium sulfide
 1. Berry, D. H., *Chem. Eng. News*, 1989, **67**(47), 2
 2. Kaner, R. B., *Nature*, 1991, **349**, 510
 Placing the chloride in contact with finely divided sodium caused an explosion after a short induction period [1]. Reaction with sodium sulfide gives molybdenum disulfide in a vigorous deflagration; autoignition will occur on mixing if the sodium sulfide is finely divided. Using sintered sulfide, which is recommended, initiation may take place on local heating to less than 60°C [2].
See Sodium: Metal halides
See other METAL HALIDES
-
- 4181. Uranium azide pentachloride**
[55042-15-4] **Cl₅N₃U**
N₃UCl₅
- Explosive solid.
See entry METAL AZIDE HALIDES
-
- 4182. Tungsten azide pentachloride**
[88495-99-2] **Cl₅N₃W**
N₃WCl₅
- Extremely explosive.
See entry METAL AZIDE HALIDES
-
- 4183. Phosphorus pentachloride**
[10026-13-8] **Cl₅P**
PCl₅
- HCS 1980, 746*
- Aluminium MRH 2.76/17
See Aluminium: Non-metal halides
- 1465

Carbamates

See *N*-Carbomethoxyiminophosphoryl chloride

Chlorine dioxide, Chlorine

MRH Chlorine dioxide 1.55/29

Mellor, 1941, Vol. 2, 281; 1940, Vol. 8, 1013

Contact between phosphorus pentachloride and a mixture of chlorine and chlorine dioxide (previously considered to be dichlorine trioxide) usually causes explosion, possibly owing to formation of the more sensitive chlorine monoxide.

Fluorine

See Fluorine: Phosphorus halides

Hydroxylamine

See Hydroxylamine: Phosphorus chlorides

Magnesium oxide

Mellor, 1940, Vol. 8, 1016

A heated mixture incandesces brilliantly.

3'-Methyl-2-nitrobenzanilide

Partridge, M. W., private comm., 1968

The residue from interaction of the chloride and anilide in benzene after removal of solvent and phosphoryl chloride under vacuum exploded violently on admission of air.

Nitrobenzene

MRH 4.52/99+

Unpublished observations

A solution of phosphorus pentachloride in nitrobenzene is stable at 110° but begins to decompose with accelerating violence above 120°C, with evolution of nitrous fumes.

Other reactants

Yoshida, 1980, 60

MRH values calculated for 8 combinations with various reagents are given.

Phosphorus(III) oxide

Mellor, 1940, Vol. 8, 898

Interaction is rather violent at ambient temperature.

Sodium

MRH 4.89/35

See Sodium: Non-metal halides

Urea

Anon., *Angew. Chem. (Nachr.)*, 1960, **8**, 33

A dry mixture exploded after heating, probably owing to formation of nitrogen trichloride. Other chlorinating agents will react similarly with nitrogenous materials.

See Chlorine: Nitrogen compounds

Water

Mellor, 1940 Vol. 8, 1012

Interaction with water in limited quantities is violent, and the hydrolysis products may themselves react violently with more water.

See Phosphoryl chloride: Water

See other NON-METAL HALIDES

4184. Antimony pentachloride

[7647-18-9]

Cl₅Sb

SbCl₅

HCS 1980, 160

N-Chlorodimethylamine

See N-Chlorodimethylamine: Antimony chlorides

Oxygen difluoride

See Oxygen difluoride: Halogens, etc.

Phosphonium iodide

See Phosphonium iodide: Oxidants

See other METAL HALIDES, OXIDANTS

4185. Tantalum pentachloride

[7721-01-9]

Cl₅Ta

TaCl₅

Lithium dimethylamide

See Pentakis(dimethylamino)tantalum(V)

See other METAL HALIDES

4186. Ammonium hexachloroplatinate

[1332-76-9]

Cl₆H₈N₂Pt

[NH₄]₂ [PtCl₆]

Potassium hydroxide, Combustible materials

Mellor, 1942, Vol. 16, 336

Boiling ammonium hexachloroplatinate with alkali gives a product (possibly potassium hexahydroxyplatinate) which after drying will explode violently on heating alone or with combustible materials.

See other PLATINUM COMPOUNDS

4187. Potassium hexachloroplatinate
[1307-80-8] **Cl₆K₂Pt**



Bromine trifluoride

See Bromine trifluoride: Potassium hexachloroplatinate

See other PLATINUM COMPOUNDS

4188. Bis(trichlorophosphoranylidene)sulfamide
[14259-65-5] **Cl₆N₂O₂P₂S**



Cellulose, or Ethanol, or Water

Vandi, A. *et al.*, *Inorg. Synth.*, 1966, **8**, 119

It is extremely reactive with water or alcohol, and causes filter paper to ignite.

See related NON-METAL HALIDES (AND THEIR OXIDES) *See other* N-S COMPOUNDS


4189. Hexachlorocyclotriphosphazine
[940-71-6] **Cl₆N₃P₃**

Dimethyl sulfoxide

See Dimethyl sulfoxide: Hexachlorocyclotriphosphazine

See related NON-METAL HALIDES

4190. Bis(triperchloratosilicon) oxide
[] **Cl₆O₂₅Si₂**



Schmeisser, M., *Angew. Chem.*, 1955, **67**, 499

This solid decomposition product of tetraperchloratosilicon was so explosive, even in small amounts, that work was discontinued.

See related NON-METAL PERCHLORATES

4191. Hexachlorodisilane
[13465-77-5] **Cl₆Si₂**



See entry HALOSILANES

Chlorine

See Chlorine: Hexachlorodisilane

4192. Uranium hexachloride
[13763-23-0]



Water

Bailar, 1973, Vol. 5, 189

Interaction is violent.

See other METAL HALIDES

4193. Tungsten hexachloride
[13283-01-7]



Sodium sulfide

Kaner, R. B., *Nature*, 1991, **349**, 510

Reacts with sodium sulfide even more exothermically than molybdenum pentachloride.

See Molybdenum pentachloride.

See other METAL HALIDES

4194. Octachlorotrisilane
[13596-23-1]



See entry HALOSILANES

4195. Tetrazirconium tetraoxide hydrogen nonaperchlorate
[]



Mellor, 1946, Vol. 2, 403

The salt, 'zirconyl perchlorate', explodes if heated rapidly. Later work suggests alternative formulations for the salt.

See other METAL PERCHLORATES

4196. Trirhenium nonachloride
[14973-59-2]



Oxygen

See Oxygen: Trirhenium nonachloride

See other METAL HALIDES

4197. Decachlorotetrasilane
[13763-19-4]



See entry HALOSILANES

4198. Dodecachloropentasilane
[13596-24-2]



See entry HALOSILANES

4199. Cobalt
[7440-48-4]



1. Bailar, 1973, Vol. 3, 1056

2. Chadwell, A. J. *et al.*, *J. Phys. Chem.*, 1956, **60**, 1340

Finely divided cobalt is pyrophoric in air [1]. Raney cobalt catalyst appears to be less hazardous than Raney nickel [2].

Acetylene

MRH 8.70/99+

See Acetylene: Cobalt

Hydrazinium nitrate

See Hydrazinium nitrate: Alome, or Metals, etc.

Other reactants

Yoshida, 1980, 120

MRH values calculated for 11 combinations, largely with oxidants, are given.

Oxidants

MRH value shows % of oxidant

See Ammonium nitrate: Metals

MRH 2.55/58

Bromine pentafluoride: Acids, etc.

Nitryl fluoride: Metals

1,3,4,7-Tetramethylisoindole

See 1,3,4,7-Tetramethylisoindole: Cobalt

See other METALS, PYROPHORIC METALS

4200. Cobalt trifluoride
[10026-18-3]



Hydrocarbons, or Water

Priest, H. F., *Inorg. Synth.*, 1950, **3**, 176

1470

It reacts violently with hydrocarbons or water, and finds use as a powerful fluorinating agent.

Silicon

Mellor, 1956, Vol. 2, Suppl. 1, 64

A gently warmed mixture reacts exothermally, attaining red-heat.

See other METAL HALIDES

4201. Cobalt(III) amide

[]



Schmitz-Dumont, O. *et al.*, *Z. Anorg. Chem.*, 1941, **284**, 175

Heating converts it to pyrophoric cobalt(III) nitride.

See other N-METAL DERIVATIVES

4202. Diamminenitratocobalt(II) nitrate

[33362-98-0]



McPherson, G. L. *et al.*, *Inorg. Chem.*, 1971, **10**, 1574

A sample of the molten salt exploded at 200°C.

See other AMMINEMETAL OXOSALTS, REDOX COMPOUNDS

4203. Cobalt tris(dihydrogenphosphide)

[]



Zehr, J., *Staub*, 1962, **22**, 494–508

It ignites in air, especially when finely divided.

See related NON-METAL HYDRIDES, PHOSPHINES

4204. Triamminetrinitrocobalt(III)

[13600-88-9]



Explodes at 305°C, medium impact-sensitivity.

See entry AMMINEMETAL OXOSALTS (reference 2)

4205. Trihydrazinecobalt(II) nitrate

[82434-33-1]



Franzen, H. *et al.*, *Z. Anorg. Chem.*, 1908, **60**, 247, 274

It is explosive.
See other AMMINEMETAL NITRATES

4206. Ammonium hexanitrocobaltate(3-)
[14652-46-1] **CoH₁₂N₉O₁₂**
[NH₄]₃[Co(NO₂)₆]

Explodes at 230°C, medium impact-sensitivity.
See entry AMMINEMETAL OXOSALTS (reference 2)

4207. Pentaamminenitratocobalt(III) nitrate
[14404-36-5] **CoH₁₅N₈O₉**
[(H₃N)₅CoNO₃] [NO₃]₂

Explodes at 310°C, medium impact-sensitivity.
See entry AMMINEMETAL OXOSALTS (reference 2)

4208. Hexaamminecobalt(III) iodate
[14589-65-2] **CoH₁₈I₃N₆O₉**
[(H₃N)₆Co] [IO₃]₃

Explodes at 355°C, low impact-sensitivity.
See entry AMMINEMETAL OXOSALTS (reference 2)

4209. Hexaamminecobalt(III) permanganate
[22388-72-3] **CoH₁₈Mn₃N₆O₁₂**
[(H₃N)₆Co] [MnO₄]₃


1. Mellor, 1942, Vol. 12, 336
2. Joyner, T. B., *Can. J. Chem.*, 1969, **47**, 2730
It explodes on heating [1], and is of treacherously high impact-sensitivity [2].
See other AMMINEMETAL OXOSALTS

4210. Hexaamminecobalt(III) nitrate
[10534-86-8] **CoH₁₈N₉O₉**
[(H₃N)₆Co] [NO₃]₃

Explodes at 295°C, medium impact-sensitivity.
See entry AMMINEMETAL OXOSALTS (reference 2)

4211. Hexahydroxylaminocobalt(III) nitrate


[18501-44-5] (ion)

CoH₁₈N₉O₁₅Werner, A. *et al.*, *Ber.*, 1905, **38**, 897

It usually explodes during preparation or handling.

See other AMMINEMETAL OXOSALTS**4212. Potassium triazidocobaltate(1-)**


[52324-65-9]

CoKN₉1. Fritzer, H. P. *et al.*, *Angew. Chem. (Intern Ed.)*, 1971, **10**, 8292. Fritzer, H. P. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1974, **10**, 247-252

The complex azide is highly explosive and must be handled with extreme care. The analogous potassium and caesium derivatives of zinc azide and nickel azide deflagrate strongly in a flame and some are shock-sensitive [1]. The potassium salt alone out of 8 azido-complexes exploded during X-irradiation in an ESCA study [2].

See other IRRADIATION DECOMPOSITION INCIDENTS*See* AZIDE COMPLEXES OF COBALT(III) *See related* METAL AZIDES**4213. Potassium hexanitrocobaltate(3-)**

[13782-01-9]

CoK₃N₆O₁₂1. Broughton, D. B. *et al.*, *Anal. Chem.*, 1947, **19**, 722. Horowitz, O., *Anal. Chem.*, 1948, **20**, 893. Tomlinson, W. R. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 375

Evaporation by heating a filtrate from precipitation of 'potassium cobaltinitrite' caused it to turn purple and explode violently [1]. This was attributed to interaction of nitrite, nitrate, acetic acid and residual cobalt with formation of fulminic or methylnitric acids or their cobalt salts, all of which are explosive [2]. Mixtures containing nitrates, nitrites and organic materials are potentially dangerous, especially in presence of acidic materials and heavy metals. A later publication confirms the suggestion of formation of nitro- or nitrito-cobaltate(III) [3].

See related METAL NITRITES

- 4214. Cobalt(III) nitride**
[12139-70-7] **CoN**
- CoN**
- Schmitz-Dumont, O., *Angew. Chem.*, 1955, **67**, 231
 A pyrophoric powder.
See Cobalt(III) amide
See other NITRIDES, PYROPHORIC MATERIALS
- 4215. Cobalt(II) nitrate**
[10026-22-9] **CoN₂O₆**
- Co(NO₃)₂**
- HCS* 1980, 321
- Ammonium hexacyanoferrate(II)
 Wolski, W. *et al.*, *Acta Chim.* (Budapest), 1972, **72**, 25–32
 Interaction is explosive at 220°C.
See Copper(II) nitrate: Ammonium hexacyanoferrate(II), or: Potassium hexacyanoferrate(II)
- Carbon MRH 3.68/14
 Crowther, J. R., private comm., 1970
 Charcoal impregnated with the nitrate exploded lightly during sieving. Possibly a dust or ‘black powder’ explosion.
- Other reactants
 Yoshida, 1980, 195
 MRH values calculated for 16 combinations with oxidisable materials are given.
See other METAL NITRATES, OXIDANTS
- 4216. Cobalt(II) azide**
[14215-31-7] **CoN₆**
- Co(N₃)₂**
- Mellor, 1940, Vol. 8, 355
 It explodes at 200°C.
See also AMMINEMETAL AZIDES
See other METAL AZIDES
- 4217. Cobalt(II) oxide**
[1307-96-6] **CoO**
- CoO**
- Hydrogen peroxide
See Hydrogen peroxide: Metals, etc.
See other METAL OXIDES

4218. Cobalt(II) sulfide
[1317-42-6]

CoS

CoS

Brauer, 1965, Vol. 2, 1523
Material dried at 300°C is pyrophoric.
See other METAL SULFIDES, PYROPHORIC MATERIALS

4219. Hexaamminecobalt(III) hexanitrocobaltate(3-)
[15742-33-3]

$[(\text{H}_3\text{N})_6\text{Co}][\text{Co}(\text{NO}_2)_6]$

$\text{Co}_2\text{H}_{18}\text{N}_{12}\text{O}_{12}$

1. Tomlinson, W. R. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 376
2. Shidlovskii, A. A. *et al.*, *Chem. Abs.*, 1977, **87**, 70416
An unstable compound of low impact-sensitivity [1]. In a comparative study of a series of cobalt complexes ranging from triamminecobalt(III) nitrite to ammonium hexanitrocobaltate(3-), the title compound burned the fastest [2].
See other AMMINEMETAL OXOSALTS

4220. trans Tetraamminediazidocobalt(III)
trans-diamminetetraazidocobaltate(1-)
[54689-17-7]

$[(\text{H}_3\text{N})_4\text{Co}(\text{N}_3)_2][(\text{H}_3\text{N})_2\text{Co}(\text{N}_3)_4]$

$\text{Co}_2\text{H}_{18}\text{N}_{24}$

Druding, L. F. *et al.*, *Inorg. Chem.*, 1975, **14**, 1365
During its preparation, solutions of the salt, which is a dangerous detonator, must not be evaporated to dryness. Surprisingly, mixtures with potassium bromide could be compressed (for IR examination) to 815 bar without decomposition. It contains 71% of nitrogen.
See AMMINEMETAL AZIDES, AZIDE COMPLEXES OF COBALT(III)
See related METAL AZIDES *See other* HIGH-NITROGEN COMPOUNDS

4221. Cobalt(III) oxide
[1308-04-9]

Co_2O_3

Co_2O_3

(Commercial samples contain some cobalt(II) oxide)

Hydrogen peroxide
See Hydrogen peroxide: Metals, etc.

Nitroalkanes
See NITROALKANES: metal oxides *See other* METAL OXIDES

4222. Chromium
[7440-47-3]

Cr

Cr

RSC Lab. Hazards Safety Data Sheet No. 65, 1987 (metal and compounds)

1. Sidgwick, 1950, 1013
2. Uel'skii, A. A. *et al.*, *Chem. Abs.*, 1978, **89**, 116214
Evaporation of mercury from mercury amalgam leaves pyrophoric chromium [1]. Increasing the temperature at which hexacarbonylchromium is thermally decomposed increases the surface area and pyrophoricity of the chromium powder produced [2].

Carbon dioxide

See Carbon dioxide: Metals

Oxidants

See Ammonium nitrate: Metals

Bromine pentafluoride: Acids, etc.

Nitrogen oxide: Metals

Sulfur dioxide: Metals

See other METALS, PYROPHORIC METALS

4223. Copper chromate oxide
[1308-09-4]

$\text{CuCrO}_4 \cdot 2\text{CuO}$

$\text{CrCuO}_4 \cdot 2\text{CuO}$

Hydrogen sulfide

Pascal, 1960, Vol. 13.1, 1025

The gas may ignite on contact with the basic chromate.

See other METAL OXOMETALLATES

4224. Ammonium fluorochromate
[58501-09-0]

NH_4FCrO_3

CrFH_4NO_3

Ribas Bernat, J. G., *Ion* (Madrid), 1975, **35**, 573–576

It decomposes violently at 220°C.

See related OXOSALTS OF NITROGENOUS BASES

4225. Chromyl fluorosulfate
[33497-88-0]

$\text{O}_2\text{Cr}(\text{OSO}_2\text{F})_2$

$\text{CrF}_2\text{O}_8\text{S}_2$

Water

Rochat, W. V. *et al.*, *Inorg. Chem.*, 1969, **8**, 158

1476

Interaction is violent.
See related ACYL HALIDES, OXIDANTS

4226. Chromium pentafluoride
[14884-42-5] **CrF₅**
CrF₅

Bailar, 1973, Vol. 2, 1086
It reacts violently in halogen-exchange and oxidation-reduction reactions.

Phosphorus trichloride
O'Donnell, T. A. *et al.*, *Inorg. Chem.*, 1966, **5**, 1435, 1437
Interaction is violent after slight warming.
See other METAL HALIDES, OXIDANTS

4227. Potassium hydroxyoxodiperoxochromate(1-)
[40330-52-7] (ion) **CrHKO₆**
K[HO(O)Cr(O₂)₂]

See entry PEROXOCHROMIUM COMPOUNDS

4228. Thallium hydroxyoxodiperoxochromate(1-)
[40330-52-7] (ion) **CrHO₆Tl**
Tl[HO(O)Cr(O₂)₂]

See entry PEROXOCHROMIUM COMPOUNDS

4229. Chromic acid
[7738-94-5] **CrH₂O₄**
H₂CrO₄

HCS 1980, 385 (dichromate/sulfuric acid ± added water)

1. Bryson, W. R., *Chem. Brit.*, 1975, **11**, 377
2. Pitt, M. J., *Chem. Brit.*, 1975, **11**, 456
3. Downing, S., *Chem. Brit.*, 1975, **11**, 456
4. Baker, P. B., *Chem. Brit.*, 1975, **11**, 456
5. Kipling, B., *Chem. Brit.*, 1976, **12**, 169
6. Kaufman, J. A., private comm., 1987

A closed bottle of unused potassium dichromate-sulfuric acid mixture exploded after several months in storage [1]. Previous similar incidents were summarised, and the possibility of the bottle having burst from internal pressure of carbon dioxide arising from trace contamination by carbon compounds was advanced [1,2]. Two further reports of incidents within 1 or 2 days of preparation of

the mixture were reported [3,4], the latter involving exothermic precipitation of chromium trioxide. Presence of traces of chloride in the dichromate leads to formation of chromyl chloride, which may be unstable [2,5]. The use of other glass-cleaning agents, and non-storage of chromic acid mixtures is again recommended [2]. Used chromic acid cleaning solution was returned to the bottle and capped; it exploded 2–3 mins. later [6].

See other GAS EVOLUTION INCIDENTS

Acetone

MCA Case History No. 1583

During glass cleaning operations, acetone splashed into a beaker used previously to contain potassium dichromate–sulfuric acid mixture and the solvent ignited. Alcohols behave similarly.

Alcohols, Silica gel

Lou, J.-D. *et al.*, *Chem. & Ind.*, 1987, 531–532

Chromic acid adsorbed on silica gel ($\text{CrO}_3 : \text{SiO}_2 = 3 : 10$) is a mild and safe reagent to oxidise saturated or unsaturated alcohols in carbon tetrachloride to the corresponding aldehydes.

Oxidisable material

MCA Case History No. 1919

A waste plating solution (containing 22% sulfuric acid and 40% w/v of chromium) was being sucked into an acid-disposal tanker. When 500 l had been transferred, a mild explosion in the tanker blew back through the transfer pump and hose. The oxidisable component in the tanker was not identified.

See other INORGANIC ACIDS, OXIDANTS

4230. Ammonium hydroxyoxodiperoxochromate(1–)


[40330-52-7] (ion)



See entry PEROXOCHROMIUM COMPOUNDS

4231. Triamminediperoxochromium(IV)

[17168-85-3]



1. Hughes, R. G. *et al.*, *Inorg. Chem.*, 1968, **7**, 882

2. Kauffman, G. B., *Inorg. Synth.*, 1966, **8**, 133

It must be handled with care because it may explode or become incandescent on sudden heating or shock. Heating at 20°/min caused a violent explosion at around 120°C [1]. Preparative and handling precautions have been detailed [2].

See other AMMINECHROMIUM PEROXOCOMPLEXES

4232. Ammonium tetraperochromate(3-)
[67165-30-4] **CrH₁₂N₃O₈**
 $[\text{NH}_4]_3[\text{Cr}(\text{O}_2)_4]$

Alone, or Sulfuric acid
Mellor, 1943, Vol. 11, 356
Explodes at 50°C, on impact, or in contact with sulfuric acid.
See other PEROXOACID SALTS

4233. Pentaamminenitrochromium(III) nitrate
[31255-94-4] **CrH₁₅N₈O₈**
 $[(\text{H}_3\text{N})_5\text{CrNO}_2][\text{NO}_3]_2$

Mellor, 1943, Vol. 11, 477
It explodes on heating, and may be impact-sensitive.
See other AMMINEMETAL OXOSALTS

4234. Hexaamminechromium(III) nitrate
[15263-28-7] **CrH₁₈N₉O₉**
 $[(\text{H}_3\text{N})_6\text{Cr}][\text{NO}_3]_3$

Moderately impact-sensitive, explodes at 263°C.
See entry AMMINEMETAL OXOSALTS (reference 2)

4235. Potassium tetraperochromate(3-)
[12331-76-9] **CrK₃O₈**
 $\text{K}_3[\text{Cr}(\text{O}_2)_4]$

Alone, or Sulfuric acid
Mellor, 1943, Vol. 11, 356
Not sensitive to impact, but explodes at 178°C or in contact with sulfuric acid.
The impure salt is less stable, and explosive.
See other PEROXOACID SALTS

4236. Lithium chromate
[14307-35-8] **CrLi₂O₄**
 Li_2CrO_4

Zirconium
de Boer, H. J. *et al.*, *Z. Anorg. Chem.*, 1930, **191**, 113
During reduction of the chromate to lithium at 450–600°C, a considerable excess
of zirconium must be used to avoid explosions.
See other METAL OXOMETALLATES

4237. Chromium nitride
[24094-93-7]

CrN

CrN

Potassium nitrate

See Potassium nitrate: Chromium nitride

See other NITRIDES

4238. Chromyl nitrate
[16017-38-2]

$\text{O}_2\text{Cr}(\text{NO}_3)_2$

CrN_2O_8

Organic materials

1. Schmeisser, M., *Angew. Chem.*, 1955, **67**, 495

2. Harris, A. D. *et al.*, *Inorg. Synth.*, 1967, **9**, 87

Many hydrocarbons and organic solvents ignite on contact with this powerful oxidant and nitrating agent [1], which reacts like fuming nitric acid in contact with paper, rubber or wood [2].

See related METAL NITRATES

4239. Chromyl azide
[]

$\text{O}_2\text{Cr}(\text{N}_3)_2$

CrN_6O_2

Preparative hazard

See Chromyl chloride: Sodium azide

See related METAL AZIDES

4240. Sodium tetraperoxo chromate(3-)
[12206-14-3]

$\text{Na}_3[\text{Cr}(\text{O}_2)_4]$

CrNa_3O_8

Mellor, 1943, Vol. 11, 356

It explodes at 115°C.

See other PEROXOACID SALTS

4241. Chromium(II) oxide
[12018-00-7]

CrO

CrO

1. Férée, J., *Bull. Soc. Chim, Fr.*, 1901, **25**, 620

2. Ellern, 1961, 33

The black powder ignites if ground or heated in air [1]. The oxide obtained by oxidation of chromium amalgam is pyrophoric [2].

See other METAL OXIDES, PYROPHORIC MATERIALS, REDUCANTS

CrO₃

4242. Chromium trioxide
[1333-82-0]

CrO₃

CrO₃

(MCA SD-52, 1944); FPA H48, 1976; HCS 1980, 315

1. Baker, W., *Chem. & Ind.*, 1956, 280
2. Goertz, A., *Arbeitschutz*, 1935, 323

Presence of nitric acid or nitrates in chromium trioxide may cause oxidation reactions to accelerate out of control, possibly owing to formation of chromyl nitrate. Samples of the oxides should be tested by melting before use, and those evolving oxides of nitrogen should be discarded [1]. A closed container of the pure oxide exploded violently when laid down on its side. This was attributed to unsuspected contamination of the container [2].

See Chromyl nitrate

Acetic acid

MRH 2.22/18

1. Anon., *BCISC Quart. Safety Summ.*, 1966, **37**, 30
2. Anon., *Sichere Chemiearb.*, 1987, **39**, 70

An explosion occurred during initial heating of a large volume of glacial acetic acid being treated with chromium trioxide. This was attributed to violent interaction of solid chromium trioxide and liquid acetic acid on a hot, exposed steam coil, and subsequent initiation of an explosive mixture of acetic acid vapour and air. The risk has been obviated by using a solution of dichromate in sulfuric acid as oxidant, in place of the trioxide. The sulfuric acid is essential, as the solid dichromate, moist with acetic acid obtained by evaporating an acetic acid solution to near-dryness, will explode [1]. Use of a mixture of chromium trioxide and acetic acid as an oxidant shattered a glass reaction vessel [2].

See other GLASS INCIDENTS

See Butyric acid, below

Acetic anhydride

MRH 2.38/16

1. Dawber, J. G., *Chem. & Ind.*, 1964, 973
2. Bretherick, L., *Chem. & Ind.*, 1964, 1196
3. Baker, W., *Chem. & Ind.*, 1956, 280
4. Eck, C. R. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1974, 865
5. *MCA Guide*, 1972, 297

A literature method for preparation of chromyl acetate by interaction of chromium trioxide and acetic anhydride was modified by omission of cooling and agitation. The warm mixture exploded violently when moved [1]. A later publication emphasised the need for cooling, and summarised several such previous occurrences [2]. An earlier reference attributes the cause of chromium trioxide–acetic anhydride oxidation mixtures going out of control to presence of nitric acid or nitrates in the chromium trioxide, and a simple test to check this point is given [3]. Mixtures used as a reagent for the remote oxidation of carboxylic esters are potentially explosive, and must be made up and used at below 25°C under controlled conditions [4]. An attempt to purify the anhydride by warming with 2% w/v of trioxide led to an explosion at 30°C [5].

Acetic anhydride, 3-Methylphenol

Thorne, J. G., private comm., 1978

Addition of the oxide in portions to the reaction mixture (for preparation of 3-hydroxybenzaldehyde or its acetate) at 75–80°C proceeded smoothly, but the final portion caused a large exotherm leading to eruption of the flask contents.

Acetic anhydride, Tetrahydronaphthalene

Peak, D. A., *Chem. & Ind.*, 1949, 14

Use of an anhydride solution of the trioxide to prepare tetralone caused a vigorous fire. This was attributed to use of the more hygroscopic granular trioxide, which is less preferable than the flake type.

Acetone

MRH 2.59/10

Delhez, R., *Chem. & Ind.*, 1956, 931

The use of chromium trioxide to purify acetone is hazardous, ignition on contact occurring at ambient temperature. Methanol behaves similarly when used to reduce the trioxide in preparing hexaaquachromium(III) sulfate.

Acetylene

Grignard, 1935, Vol. 3, 167

Acetylene is oxidised violently.

Alcohols

MRH Ethanol 2.55/10

1. Newth, F. H. *et al.*, *Chem. & Ind.*, 1964, 1482
2. Neumann, H., *Chem. Eng. News*, 1970, **48**(24), 4
3. *491M*, 1975, 126
4. Kishore, K., *Def. Sci. J.*, 1978, **28**, 13–14

When methanol was used to rinse a pestle and mortar which had been used to grind coarse chromium trioxide, immediate ignition occurred due to vigorous oxidation of the solvent. The same occurred with ethanol, 2-propanol, butanol and cyclohexanol. Water is a suitable cleaning agent for the trioxide [1]. For oxidation of *sec*-alcohols in DMF, the oxide must be finely divided, as lumps cause violent local reaction on addition to the solution [2]. Use of methanol to reduce the Cr(VI) oxide to a Cr(III) derivative led to an explosion and fire [3]. The ignitability of the butanols decreases from *n*-through *sec*- to *tert*-butanol [4].

See Dimethylformamide, below

Alkali metals

MRH Potassium 1.46/23, sodium 3.39/21

Mellor, 1943, Vol. 11, 237

Sodium or potassium reacts with incandescence.

Ammonia

MRH Gas 2.51/15, solution 1.63/26

1. Mellor, 1943, Vol. 11, 233
2. Pascal, 1959, Vol. 14, 215

Gaseous ammonia leads to incandescence [1], and the aqueous solution is oxidised very exothermally [2].

Arsenic

Mellor, 1943, Vol. 11, 234

Interaction is incandescent.

Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

Butyric acid

Wilson, R. D., *Chem. & Ind.*, 1957, 758

A mixture of chromium trioxide and butyric acid became incandescent on heating to 100°C.

See Acetic acid, above

Chlorine trifluoride

See Chlorine trifluoride: Chromium trioxide

Chromium(II) sulfide

Mellor, 1943, Vol. 11, 430

Interaction causes ignition.

Dimethylformamide

MRH 2.64/11

1. Neumann, H. *Chem. Eng. News*, 1970, **48**(28), 4

2. Heathcock, C. H., *Chem. Eng. News*, 1981, **59**(8), 9

During oxidation of a *sec*-alcohol to ketone in cold DMF solution, addition of solid trioxide caused ignition. Addition of lumps of trioxide was later found to cause local ignition on addition to ice-cooled DMF under nitrogen [1]. Addition of 2 g of chromium trioxide to 18 ml of solvent to form a 10 wt% solution caused immediate ignition and ejection of the flask contents [2].

See Dimethylformamide: Oxidants

1,3-Dimethylhexahydropyrimidone

Mukhopadhyay, T. *et al.*, *Helv. Chim. Acta*, 1982, **65**, 386 (footnote 9)

Contact of the trioxide with the aprotic amide solvent ('dimethylpropyleneurea') is always explosive, with fire, whereas this only occurs with hexamethylphosphoramide (below) if the oxide is previously crushed. However the former is much less toxic than the latter solvent.

Ethylene glycol

See Ethylene glycol: Oxidants

Glycerol

MRH 2.38/17

Pieters, 1957, 30

Interaction is violent; the mixture may ignite owing to oxidation of the trihydric alcohol, which is viscous and unable to dissipate the exotherm.

See Glycerol: Oxidants

Hexamethylphosphoramide

Cardillo, G. *et al.*, *Synthesis*, 1976, **6**, 394–396

Stirring chromium trioxide (added in small portions) with the unheated solvent leads to the formation of a complex useful for oxidising alcohols to carbonyl derivatives. The trioxide must not be crushed before being added to the solvent, because violent decomposition may then occur.

See 1,3-Dimethylhexahydropyrimidone, above

Hydrazine MRH 3.01/19

See Hydrazine: Oxidants (reference 2)

Hydrogen sulfide MRH 2.05/12

Mellor, 1943, Vol. 11, 232

Contact with the heated oxide causes incandescence.

Organic materials, MRH values below references show % of organic
or Solvents

1. Leleu, *Cahiers*, 1976, (83), 286–287

2. Fawcett, H. H., *Ind. Eng. Chem.*, 1959, **51**(4), 90A

3. Mikhailov, V., *Chem. Abs.*, 1960, **54**, 23331f

4. Mellor, 1943, Vol. 11, 235

MRH Aniline 2.59/8, diethyl ether 2.59/8, hexane 2.46/6, phenol 2.51/9, toluene 2.55/7

Combustible materials may ignite or explode on contact with the oxide. A few drops of oil which fell into a container of the oxide led to an explosion which produced fatal burns [1]. If a few drops of an organic solvent (acetone, 2-butanone, ethanol) contact solid chromium trioxide, a few seconds' delay ensues while some of the oxidant attains the critical temperature of 330°C. Then combustion occurs with enough vigour to raise a fire-ball several feet, and spattering also occurs [2]. Possible ignition hazards were studied for a range of 60 organic liquids and solids in contact with the solid oxide. Hot liquids were added to the oxide; solids were covered with a layer of the oxide. The most dangerous materials were methanol, ethanol, butanol, isobutanol, acetaldehyde, propionaldehyde, butyraldehyde, benzaldehyde, acetic acid, pelargonic acid, ethyl acetate, isopropyl acetate, pentyl acetate, diethyl ether, methyldioxane, dimethyldioxane, acetone and benzylethylaniline. Other materials evolved heat, especially in presence of water. Segregation in storage or transport is essential [3]. Benzene ignites in contact with the powdered oxide [4].

See 1,3-Dimethylhexahydropyrimidone, or Dimethylformamide, or Glycerol, or Hexamethylphosphoramide, all above

Other reactants

Yoshida, 1980, 354–355

MRH values calculated for 28 combinations, largely with oxidisable materials, are given.

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

- Phosphorus MRH (Yellow) 3.68/16
Moissan, H., *Ann. Chim. Phys. [6]*, 1885, **5**, 435
Phosphorus and the molten trioxide react explosively.
- Potassium hexacyanoferrate(III) MRH 2.59/25
Anon., *BCISC Quart. Safety Summ.*, 1965, **36**(144), 55
Mixtures of the ferrate ('ferricyanide') and chromium trioxide explode and inflame when heated above 196°C. Friction alone is sufficient to ignite violently the mixture when ground with silver sand.
See other FRICTIONAL INITIATION INCIDENTS
- Pyridine MRH 2.63/92
1. Dauben, W. G. *et al.*, *J. Org. Chem.*, 1969, **34**, 3587
2. Poos, G. I. *et al.*, *J. Amer. Chem. Soc.*, 1953, **75**, 427
3. Collins, J. C. *et al.*, *Org. Synth.*, 1972, **52**, 7
4. Ratcliffe, R. *et al.*, *J. Org. Chem.*, 1970, **35**, 4001
5. Stensio, K. E., *Acta Chem. Scand.*, 1971, **25**, 1125
6. *MCA Case History No. 1284*
During preparation of the trioxide-pyridine complex (a powerful oxidant), lack of really efficient stirring led to violent flash fires as the oxide was added to the pyridine at -15 to -18°C. Reversed addition of pyridine to the oxide is extremely dangerous [1], ignition usually occurring [2]. A later preparation specifies temperature limits of 10-20°C to avoid an excess of unreacted trioxide [3]. A safe method of preparing the complex in solution has been described [4], and preparation and use of solutions of the isolated complex in dichloromethane [3] or acetic acid [5] have been detailed. The Case History gives further information on preparation of the complex. Solution of the oxide is not smooth; it first swells, then suddenly dissolves in pyridine with an exotherm. This hazard may be eliminated without loss of yield by dissolving the trioxide in an equal volume of water before adding it to 10 volumes of pyridine. Pulverising chromium trioxide before use is not recommended, as this increases its rate of reaction with organic compounds to a hazardous level [6].
See Oxodiperoxypyridinechromium *N*-oxide
- Selenium
Mellor, 1943, Vol. 11, 233
Interaction is violent.
- Sodium MRH 3.39/21
See Sodium: Metal oxides
See other REDOX REACTIONS
- Sodium amide
Mellor, 1943, Vol. 11, 234
Grinding a mixture leads to violent reaction.
- Sulfur MRH 1.76/14
Mellor, 1943, Vol. 11, 232

A mixture ignites on warming.
See other METAL OXIDES, OXIDANTS
See related ACID ANHYDRIDES

4243. Lead chromate
[7758-97-6]

CrO₄Pb

PbCrO₄

HCS 1980, 587

Aluminium, Dinitronaphthalene

Nagaishi, T. *et al.*, *Chem. Abs.*, 1977, **86**, 59602

The considerable energy released by the mixture derives from chromate-catalysed exothermic decomposition of the nitro compound, coupled with a thermite-type reaction of the aluminium and chromate. It is useful for cracking concrete.

Azo-dyestuffs

Anon., *Loss Prev. Bull.*, 1978, (022), 117

Under certain conditions, dry mixtures of lead chromate pigments with the azo-dyes 1-(2',4'-dinitrobenzeneazo)-2-hydroxynaphthalene (dinitroaniline orange) or 1-(4'-chloro-2'-nitrobenzeneazo)-2-hydroxynaphthalene (chlorinated para-red) may lead to violent explosions during mixing/blending operations.

See N,N''-Dichlorobis(2,4,6-trichlorophenyl)urea: 1-p-Nitrobenzeneazo-2-naphthol

Iron(III) hexacyanoferrate(4-)

1. Anon., *Chem. Processing*, 1967, **30**, 118

2. Twitchett, H. J., *Chem. Brit.*, 1977, **13**, 437

During grinding operations, the intimate mixture of pigments was ignited by a spark and burned fiercely [1]. Spontaneous ignition of Brunswick Green pigment (which also contains lead sulfate) soon after grinding was not uncommon, and similar incidents had led to the loss of ships with cargoes of Prussian Blue or Brunswick Green in wooden casks [2].

See Azo-dyestuffs, above

See other SELF-HEATING AND IGNITION INCIDENTS

Sulfur

Jackson, H., *Spectrum*, 1969, **7**(2), 82

The mixture is pyrophoric.

Tantalum

Sarawadekar, R. G. *et al.*, *Thermochim. Acta*, 1983, **70**, 133

The mixture is a pyrotechnic composition.

See other METAL OXOMETALLATES, OXIDANTS

4244. Chromium(II) sulfate
[13825-86-0]

CrO₄S

CrSO₄

Water

1. van Bemmelen, J. M., *Rec. Trav. Chim.*, 1887, **6**, 202–204
2. Bailar, 1973, Vol. 3, 657–658

Crystals of the heptahydrate, damp with surplus water, were sealed into a glass tube and stored in darkness, and after a year the tube exploded. This was attributed to the pressure of hydrogen liberated by reduction of water by the chromium(II) salt [1]. More recent information [2] confirms this hypothesis.

See Chromium(II) chloride *See other* METAL OXONON-METALLATES, REDUCANTS

4245. Chromium(II) sulfide
[12018-06-3]

CrS

CrS

Chromium trioxide

See Chromium trioxide: Chromium(II) sulfide

Fluorine

See Fluorine: Sulfides

See other METAL SULFIDES

4246. Ammonium dichromate
[7789-09-5]

Cr₂H₈N₂O₇

NH₄[OCrO₂OCrO₂O]NH₄

(MCA SD-45, 1952); HCS 1980, 143

1. Mellor, 1943, Vol. 11, 324
2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986
3. Anon., *Chem. Eng. News*, 1986, **64**(4), 6
4. Bretherick, L., *Chem. Eng. News*, 1986, **64**(22), 4

Thermal decomposition of the salt (which is of zero oxygen balance) is initiated by locally heating to 190°C, and flame and sparks spread rapidly through the mass, and if confined, it may become explosive [1]. Under close confinement, the deflagrating salt shows the extraordinarily high rate of pressure increase of 68 kbar/s, attaining a final pressure of 510 bar in about 10 ms, and further study of its deflagration and homogeneous decomposition under confinement have been reported [2]. Energy of decomposition (in range 230–260°C) was measured as 0.76 kJ/g by DSC, and T_{ait24} was determined as 96°C by adiabatic Dewar tests, with an apparent energy of activation of 112 kJ/mol. Following a report of an explosion during heated vacuum drying of a 1 t quantity of the salt [3], attention was drawn [4] to the extreme potential hazard involved in heating the salt under confinement.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION

See other DEFLAGRATION INCIDENTS

Ethylene glycol
See Ethylene glycol: Oxidants
See other DICHROMATE SALTS OF NITROGENOUS BASES

4247. Ammonium pentaperoxodichromate(2-)
[] **Cr₂H₈N₂O₁₂**
NH₄[O(O₂)₂CrOOCr(O₂)₂O]NH₄

Brauer, 1965, Vol. 2, 1392
It explodes at 50°C.
See other PEROXOACID SALTS

4248. Potassium dichromate
[7778-50-9] **Cr₂K₂O₇**
K₂[OCrO₂OCrO₂O]

FPA H43, 1976

Boron, Silicon
See Boron: Dichromates, etc.

Ethylene glycol MRH 1.46/11
Yoshida, T., private comm., 1982
Mixing of equal weights at ambient temperature is uneventful, but at 100° an exotherm of 170°C occurs.

Hydrazine MRH 1.59/99+
Mellor, 1943, Vol. 11, 234
Explosive interaction.
See Hydrazine: Oxidants (reference 2)

Hydroxylamine MRH 3.85/99+
See Hydroxylamine: Oxidants

Iron MRH 0.42/34
Laye, P. G. *et al.*, *Thermochim. Acta*, 1980, **39**, 3567–359
Pyrotechnic mixtures (1:1 wt) attained a maximum temperature of about 1090°C on ignition.
See Tungsten, below

Sulfuric acid
See Chromic acid

Tungsten
Boddington, T. *et al.*, *Combust. Flame*, 1975, **24**, 137–138
Combustion of a pyrotechnic mixture of the two materials (studied by DTA and temperature profile analysis) attains a temperature of about 1700°C in 0.1–0.2 s.
See other METAL OXOMETALLATES, OXIDANTS

4249. Potassium pentaperoxodichromate

[]




Mellor, 1943, Vol. 11, 357; Sidgwick, 1950, 1007

The powdered salt explodes above 0°C.

See other PEROXOACID SALTS

4250. Sodium dichromate

[10588-01-9]



HCS 1980, 838

Acetic acid, 2-Methyl-2-pentenal

Nolan, 1983, Case history 73

2-Methyl-2-pentenal was oxidised to the acid in a process involving addition over a period into a mixture of acetic acid and sodium dichromate at 50°C. Addition of the aldehyde, effected by sucking it into the reactor by application of vacuum, was entrusted to an inexperienced operator, but was apparently too fast and the exotherm led to a runaway and eruption of the reactor contents.

Acetic anhydride

MRH 1.84/13

Marszalek, G., private comm., 1973

Addition of the dehydrated salt to acetic anhydride caused an exothermic reaction which accelerated to explosion. Presence of acetic acid (including that produced by hydrolysis of the anhydride by the hydrate water) has a delaying effect on the onset of violent reaction, which occurs where the proportion of anhydride to acid (after hydrolysis) exceeds 0.37:1, with an initial temperature above 35°C. Mixtures of dichromate (30 g) with anhydride-acid mixtures (70 g, to give ratios of 2:1, 1:1, 0.37:1) originally at 40°C accelerated out of control after 18, 43 and 120 min, to 160, 155 and 115°C, respectively.

See other INDUCTION PERIOD INCIDENTS

Boron, Silicon

See Boron: Dichromates, etc.

Ethanol, Sulfuric acid

1. Annable, E. H., *School Sci. Rev.*, 1951, **32**(117), 249

2. Anon., *Education in Science*, 1990, (140), 30

During preparation of acetic acid by acid dichromate oxidation of ethanol according to a published procedure, minor explosions occurred on two occasions after refluxing had been discontinued. This possibly may have involved formation of acetaldehyde (which has an AIT of 140°C) and ingress of air into the reaction vessel as it cooled [1]. Runaway reactions during small scale oxidation of ethanol have apparently been experienced by many teachers, poor initial mixing or starting too cool may be the cause [2].

See 2-Propanol, Sulfuric acid, below

Hydrazine

Mellor, 1943, Vol. 11, 234

Interaction is explosive.

See Hydrazine: Oxidants

Hydroxylamine

See Hydroxylamine: Oxidants

Organic residues, Sulfuric acid

MRH Acetone 1.97/8

1. HCS 1980, 385

2. Bradshaw, J. R., *Process Biochem.*, 1970, **5**(11), 19

3. Anon., *CISHC Chem. Safety Summ.*, 1979, **50**, 95

The well-known 'chromic acid mixture' of dichromate and sulfuric acid [1] for cleaning glassware is by design a powerful oxidant, and contact with large amounts of tarry or other organic residues in process vessels should be avoided as it may lead to a violent reaction. Further, if solvents are first used to clean glassware roughly before acid treatment, traces of readily oxidisable solvents must be removed before adding the oxidant mixture. In many cases treatment with a properly formulated detergent will ensure adequate cleanliness and avoid possible hazard [2]. Addition of 1 l of oxidant mixture to a vessel containing residues of acetone or similar solvent led to a violent exothermic reaction [3].

See Chromic acid: Acetone, also Chromium trioxide: Acetic acid

Other reactants

Yoshida, 1980, 177

MRH values calculated for 16 combinations with oxidisable materials are given.

2-Propanol, Sulfuric acid

Cochrane, A., private comms., 1982, 1983

An established school preparation of 2-propanone (acetone) involves the small-scale (and rather exothermic) oxidation of the alcohol with dichromate(VI). It was observed in several laboratories that when the acidified dichromate solution was added to the alcohol in small portions (1–2 cc) rather than dropwise as specified, small sparks or incandescent particles were produced which sometimes survived long enough to escape from the neck of the flask. This also happened if the alcohol and/or the oxidant solution were diluted with extra water, with old or new samples of alcohol, and if air were displaced from the flask by carbon dioxide. It is therefore important not to exceed the specified dropwise rate of addition of oxidant solution. It is very unusual for glowing particles to be produced from a homogeneous liquid reaction system.

See Ethanol, Sulfuric acid, above

Sulfuric acid, Trinitrotoluene

Clarke, H. T. *et al.*, *Org. Synth.*, 1941, Coll. Vol. 1, 543–544

During oxidation of TNT in sulfuric acid to trinitrobenzoic acid, stirring of the viscous reaction mixture must be very effective to prevent added portions of solid dichromate causing local ignition.

See other AGITATION INCIDENTS, SELF-HEATING AND IGNITION INCIDENTS
See other METAL OXOMETALLATES, OXIDANTS

4251. Chromium(III) oxide
[1308-38-9]



Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Copper oxide

See COPPER CHROMITE CATALYST

Lithium

See Lithium: Metal oxides

Nitroalkanes

See NITROALKANES: metal oxides

Rubidium acetylide

See Rubidium acetylide: Metal oxides

See other METAL OXIDES

4252. Ammonium trichromate(2-)
[32390-97-1]



Mellor, 1943, Vol. 11, 349

It explodes at 190°C.

See other OXOSALTS OF NITROGENOUS BASES

4253. Ammonium tetrachromate(2-)
[54153-83-2]



Mellor, 1943, Vol. 11, 352

It decomposes suddenly at 175°C.

See other OXOSALTS OF NITROGENOUS BASES

4254. Caesium
[7440-46-2]



1. Mellor, 1941, Vol. 2, 468; 1963, Vol. 2, Suppl. 2.2, 2291

2. Pepmiller, P. L. *et al.*, *Rev. Sci. Instr.*, 1985, **56**, 1832–1833

It ignites immediately in air or oxygen [1]. A simple device to open glass vials of caesium and dispense the contents safely gives a good margin of safety to the user [2].

Acids

Pascal, 1957, Vol. 3, 94

Most acids react violently, even when anhydrous.

Halogens

Gibson, 1969, 8

Interaction at ambient temperature is violent with all halogens.

Non-metals

Pascal, 1957, Vol. 3, 94

Interaction with sulfur or phosphorus attains incandescence.

Water

1. Mellor, 1941, Vol. 2, 468; 1963, Vol. 2, Suppl. 2.2, 2328

2. Markowitz, M. M., *J. Chem. Educ.*, 1963, **40**, 633–636

Caesium reacts very violently with cold water and the evolved hydrogen ignites [1]. The reactivity of caesium and other alkali metals with water has been discussed in detail [2].

See other METALS, PYROPHORIC METALS

4255. Caesium fluoride

[13400-13-0]

CsF

CsF

Benzenediazonium tetrafluoroborate, Difluoramine

See Difluoramine: Benzenediazonium tetrafluoroborate, etc.

See other METAL HALIDES

4256. Caesium fluoroxysulfate

[70806-67-6]

CsOSO₂OF

CsFO₄S

Appelman, E. H., *Inorg. Synth.*, 1986, **24**, 22–23

Caesium fluoroxysulfate is thermodynamically unstable. Mild detonations have occurred occasionally during handling of 100 mg portions of salt, usually when deliberate or violent crushing or scraping occurred. It is likely that detonations were brought about by surface impurities on containers or spatulae. Amounts of 10–20 g have been used routinely over several years without detonations, though great care, shields and gloves were always used. It is also a powerful oxidant and reacts violently with organic solvents, including dimethylformamide, dimethyl sulfoxide, ethylene dichloride and pyridine.

See Rubidium fluoroxysulfate

See other ACYL HYPOHALITES, OXIDANTS

4257. Caesium pentafluorotelluramide

[42081-47-0]

CsF₅HNTe

CsNHTeF₅

Seppelt, K., *Inorg. Chem.*, 1973, **12**, 2838

Heating must be avoided during the preparation or subsequent drying, as occasional explosions occurred. It exploded immediately upon laser irradiation for Raman spectroscopy.

See other IRRADIATION DECOMPOSITION INCIDENTS, *N*-METAL DERIVATIVES

4258. Caesium hydride

[13772-47-9]

CsH

CsH

Oxygen

Gibson, 1969, 76

The unheated hydride ignites in oxygen.

See other METAL HYDRIDES

4259. Caesium hydrogen xenate

[[73378-56-0], Sesquihydrate]

CsHO₄Xe

CsOXeO₂OH

Alone, or Alcohols

Jaselskis, B. *et al.*, *J. Amer. Chem. Soc.*, 1966, **88**, 2150; 1969, **91**, 1874

It is unstable to friction, thermal or mechanical shock, and may explode on contact with alcohols.

See other XENON COMPOUNDS

4260. Caesium amide

[22205-57-8]

CsH₂N

CsNH₂

Pez, G. P. *et al.*, *Pure Appl. Chem.*, 1985, **57**, 1917–1926

Caesium amide is extremely pyrophoric and must be handled in an atmosphere virtually free from oxygen and moisture (each below 5 ppm).

Water

Mellor, 1940, Vol. 8, 256

Interaction is incandescent in presence of air.

See other INORGANIC BASES, *N*-METAL DERIVATIVES

4261. Caesium nitrate
[7789-18-6]



Xenon hexafluoride

See Xenon hexafluoride: Caesium nitrate

Xenon tetrafluoride

See Xenon tetrafluoride: Caesium nitrate

See other METAL NITRATES

4262. Caesium azide
[22750-57-8]



Sulfur dioxide

Kennet, F. A. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 853

The azide ignites in contact with sulfur dioxide at ambient temperature.

See other METAL AZIDES

4263. Caesium trioxide (Caesium ozonide)
[12053-67-7]



Water

Whaley, T. P. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 79

Reaction of caesium or potassium ozonides with water or aqueous acids is violent, producing oxygen and flashes of light.

See other METAL OXIDES

4264. Caesium oxide
[20281-00-9]



Ethanol

Pascal, 1957, Vol. 3, 104

Contact of a little alcohol with the oxide may ignite the solvent.

Halogens, or Non-metal oxides

Mellor, 1941, Vol. 2, 487

Above 150–200°C, incandescence occurs with fluorine, chlorine or iodine. In presence of moisture, contact at ambient temperature with carbon monoxide or carbon dioxide causes ignition, while dry sulfur dioxide causes incandescence on heating.

Water

Pascal, 1957, Vol. 3, 104

Incandescence on contact.
See other METAL OXIDES

4265. Caesium selenide
[31052-46-7]

CsSeCs

Cs₂Se

Sidgwick, 1950, 92
When warm it ignites in air.
See related METAL SULFIDES

4266. Caesium nitride
[12134-29-1]

Cs₃N

Cs₃N

Alone, or Chlorine, or Non-metals
Mellor, 1940, Vol. 8, 99
It burns in air, and is readily attacked by chlorine, phosphorus or sulfur.
See other NITRIDES

4267. Copper
[7440-50-8]

Cu

Cu

Acetylenic compounds
See ACETYLENIC COMPOUNDS: metals

Aluminium, Sulfur
See Aluminium: Copper, Sulfur

3-Bromopropyne
See 3-Bromopropyne: Metals

Complexing agents, Water

Shah, I. U., *Chem. Eng. News*, 1991, **69**(30), 2

A bottle of 'cuprous chloride' solution prepared by standing cupric chloride in strong hydrochloric acid over excess copper burst on standing. In the presence of some complexing agents, copper can react with aqueous media to form hydrogen. Slow pressurisation by this means explains the above explosion (Editor's comments). The metal is also known to dissolve in cyanides and some amine solutions.

Dimethyl sulfoxide, Trichloroacetic acid

See Dimethyl sulfoxide: Copper, Trichloroacetic acid

Ethylene oxide

See Ethylene oxide (reference 3)

Lead azide

See Lead(II) azide: Copper

Other reactants

Yoshida, 1980, 243

MRH values calculated for 11 combinations with oxidants are given.

Oxidants

MRH Sodium chlorate 2.0/36

Mellor, 1941, Vol. 2, 310

Mixtures of finely divided copper with chlorates or iodates explode on friction, shock or heating.

See Ammonium nitrate: Metals

Chlorine: Metals

Chlorine trifluoride: Metals, etc.

Fluorine: Metals

Hydrazinium nitrate: Alone, or Metals

Hydrogen sulfide: Metals

Potassium dioxide: Metals

Sulfuric acid: Copper

Water

Zyszkowski, W., *Int. J. Heat Mass Transf.*, 1976, **19**, 849–868

The vapour explosion which occurs when liquid copper is dumped into water has been studied.

See MOLTEN METAL EXPLOSIONS *See other* METALS

4268. Copper–zinc alloys

[12019-27-1]

Cu–Zn

Cu–Zn

Alkyl halides

See DIALKYLZINCS

Diiodomethane, Ether

1. Foote, C. S., private comm., 1965

2. Repic, O. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 2729–2732

Lack of cooling during preparation of the Simmons-Smith organozinc reagent caused the reaction to erupt. The possibly pyrophoric nature of organozinc compounds and the presence of ether presents a severe fire hazard [1]. An alternative, safer method of activating the zinc for the reaction involves use of ultrasonic irradiation rather than the copper–zinc couple [2].

See other ALLOYS

4269. Copper iron(II) sulfide
[12015-76-8]



Ammonium nitrate

See Ammonium nitrate: Copper iron(II) sulfide

Water

Gribin, A. A., *Chem. Abs.*, 1943, **37**, 1272₈

A large dump of copper pyrites ore ignited after heavy rain. The thick layer (6–7 m) and absence of ventilation were contributory factors to the accelerating aerobic oxidation which finally led to ignition.

See other METAL SULFIDES, SELF-HEATING AND IGNITION INCIDENTS

4270. Copper(II) azide hydroxide
[22887-15-6]



Cirulis, A., *Z. Anorg. Chem.*, 1943, **251**, 332–334

It exploded at 203°C or on impact. The dimeric double salt is similarly sensitive.

See related METAL AZIDES

4271. Lithium dihydrocuprate
[53201-99-3]



Ashby, E. C. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1974, 157–158

The solid is highly pyrophoric, but stable as a slurry in ether at ambient temperature.

See other COMPLEX HYDRIDES

4272. Copper(II) phosphinate
[34461-68-2]



Mellor, 1940, Vol. 8, 883; 1971, Vol. 8, Suppl. 3, 623

The solid suddenly explodes at about 90°C, and forms impact-sensitive priming mixtures.

See other METAL PHOSPHINATES, REDOX COMPOUNDS

4273. Lithium pentahydrocuprate(4–)
[64010-65-7]




Ashby, E. C. *et al.*, *J. Org. Chem.*, 1978, **43**, 187

Of a series of lithium hydrocuprates $\text{Li}_n\text{CuH}_{n+1}$ ($n = 1-5$), only the title compound (when solvated with THF) appeared a more powerful reductant than lithium tetrahydroaluminate. Safety precautions similar to those adopted for the complex aluminium hydride seem appropriate.

See other COMPLEX HYDRIDES

4274. Tetraamminecopper(II) sulfate

[14283-05-7]



$\text{CuH}_{12}\text{N}_4\text{O}_4\text{S}$

Iodine

See Iodine: Tetraamminecopper(II) sulfate

See other AMMINEMETAL OXOSALTS

4275. Tetraamminecopper(II) nitrite

[39729-81-2]



$\text{CuH}_{12}\text{N}_6\text{O}_4$

Mellor, 1940, Vol. 8, 480

The salt is nearly as shock-sensitive as picric acid. When pure it does not explode on heating, but traces of nitrate cause explosive decomposition.

See other AMMINEMETAL OXOSALTS

4276. Tetraamminecopper(II) nitrate

[31058-64-7]



$\text{CuH}_{12}\text{N}_6\text{O}_6$

Explodes at 330°C, high impact-sensitivity.

See entry AMMINEMETAL OXOSALTS (reference 2)

4277. Tetraamminecopper(II) azide

[70992-03-9]



$\text{CuH}_{12}\text{N}_{10}$

Mellor, 1940, Vol. 8, 348; 1967, Vol. 8, Suppl.2, 26

Explosive on heating or impact.

See related METAL AZIDES

4278. Lithium hexaazidocuprate(4-)

[]



$\text{CuLi}_4\text{N}_{18}$

Urbanski, 1967, Vol. 3, 185

It is, like copper(II) azide, an exceptionally powerful initiating detonator, containing 61.5% of nitrogen.

See related METAL AZIDES *See other* HIGH-NITROGEN COMPOUNDS

4279. Copper(II) nitrate

[3251-23-8]

CuN₂O₆

[10031-43-3]; [13478-38-1] hydrates

Cu(NO₃)₂

HCS 1980, 336

1. Priestley J., *Experiments and Observations on Different Kinds of Air*, 1790, 3, 203, Birmingham

Although the anhydrous nitrate is relatively stable and can be distilled under partial pressure, as is remarked in most recent text books, these do not tell that the hydrate actually obtainable by purchase or by normal preparations is among the least stable and most oxidising of main valence nitrates. It was early observed that paper contaminated with it ignited easily [1].

Acetic anhydride

MRH 3.56/26

See Acetic anhydride: Metal nitrates

Ammonia, Potassium amide

Sorbe, 1968, 80

Interaction gives an explosive precipitate containing the Cu(I) derivatives Cu₃N, Cu₃N.*n*NH₃, Cu₂NH, CuNK₂.NH₃.

Ammonium hexacyanoferrate(II)

1. Wolski, J. *et al.*, *Z. Chem.*, 1973, **13**, 95–97

2. Wolski, J. *et al.*, *J. Therm. Anal.*, 1973, **5**, 67

At 220°C interaction is explosive in wet mixtures [1], or in dry mixtures if the nitrate is in excess [2].

See other REDOX REACTIONS

Ammonium nitrate

Mellor, 1923, Vol. III, 284

Evaporation of mixed solutions gives violent decomposition during concentration.

Hydrazine, Ethanol

Hofmann, K. A. *et al.*, *Annalen*, 1899, **305**, 222

A blue crystalline complex, assigned a monohydrazino structure, was obtained from the nitrate and hydrazine hydrate in ethanol. It exploded on heating or, sometimes, after thorough drying in a desiccator, on shaking.

See related AMMINOMETAL NITRATES

1-Hydroxybenzotriazole, Ethanol

Anon., *Safety Digest Univ. Safety Assoc.*, 1992, **44**, 24

A mixture of the above was evaporated to dryness, on scraping the residue with a spatula a 'dust explosion', presumably of only a small part of the charge, resulted, covering the experimenter with powder.

See 1-Hydroxybenzotriazole

See other REDOX REACTIONS

See also AMMINOMETAL NITRATES

Organometallic materials

Buffinger, D. R., *et al.*, *J. Coord. Chem.*, 1988, **19**(1–3), 197

A double salt of the nitrate and the copper(II) salt of an organometallic ligand was found to explode at around 100°C. In view of the oxidising powers of the nitrate, complexes of this nature are not likely to be stable.

See ORGANOMETALLIC NITRATES

Other reactants

Yoshida, 1980, 198

MRH values calculated for 19 combinations with oxidisable materials are given.

Potassium hexacyanoferrate(II)

Wolski, J. *et al.*, *Explosivstoffe*, 1969, **17**(5), 103–110

Interaction is explosive at 220°C.

Tin

MRH 2.68/62

Ellern, 1968, 46

Tin foil in contact with a solution of the nitrate may ignite or give sparks.

See other METAL NITRATES

4280. Copper(II) azide

[14215-30-6]

CuN₆

Cu(N₃)₂

1. Mellor, 1940, Vol. 8, 348; 1967, Vol. 8, Suppl. 2.2, 42–50

2. Urbanski, 1967, Vol. 3, 185

3. Jenkins, C. L. *et al.*, *J. Org. Chem.*, 1971, **36**, 3095–3102

4. Sood, R. K. *et al.*, *Chem. Abs.*, 1997, **126**, 48910

The azide is very explosive, even when moist. Loosening the solid from filter paper caused frictional initiation. Explosion initiated by impact is very violent, and spontaneous explosion has also been recorded [1]. It is also an exceptionally powerful initiator [2]. Detonation of the azide when dry has been confirmed [3]. Good crystals are considerably more sensitive to shock, friction and electric discharge than is powder [4].

See Sodium azide: Heavy metals

See other METAL AZIDES

4281. Copper(II) oxide
[1317-38-0]

CuO

CuO

Aluminium

MRH 3.89/18

1. Krah, W., *Chem Abs.*, 1978, **89**, 89652
2. Haws, L. D. *et al.*, *Chem. Abs.*, 1979, **90**, 189294

In demonstrating the use of powdered aluminium to reduce the oxide in a thermite-type reaction, the mixture must be heated behind a safety screen because of the small explosion produced [1]. Consolidation of the thermite mixture into a high-density composite gives chemical heat sources which are safe to handle [2].

See Metals, below

See other THERMITE REACTIONS

Anilinium perchlorate

See Anilinium perchlorate: Metal oxides

Barium acetate, Yttrium oxide

See Barium acetate: Copper(II) oxide, etc.

Boron

Mellor, 1946, Vol. 5, 17

The exothermic reaction on heating a mixture melted the glass container.

See other GLASS INCIDENTS

Dichloromethylsilane

See Dichloromethylsilane: Oxidants

Rubidium acetylide

See Rubidium acetylide: Metal oxides

Hydrogen

Read, C. W. W., *School Sci. Rev.*, 1941, **22**(87), 340

Reduction of the heated oxide in a combustion tube by passage of hydrogen caused a violent explosion. (The hydrogen may have been contaminated with air.)

Hydrogen sulfide

MRH 0.59/30

See Hydrogen sulfide: Metal oxides

Hydrogen trisulfide: Metal oxides

Metals

MRH values below show % of metal

1. Browne, T. E., *School Sci. Rev.*, 1967, **48**(166), 921
2. Vella, A. J., *J. Chem. Educ.* 1994, **71**(4), 328

An attempted thermite reaction with aluminium powder and copper(II) oxide in place of iron(III) oxide caused a violent explosion. An anonymous comment suggests that a greater reaction rate and exothermic effect were involved, and adds that attempted use of silver oxide would be even more violent [1]. An explosion

resulted from heating a mixture of copper(II) oxide and magnesium in a school laboratory. Avoidance of magnesium thermites in teaching is counselled [2].

See Aluminium, above

Aluminium: Metal oxides MRH 3.89/18

Magnesium: Metal oxides MRH 4.30/23

Potassium: Metal oxides

Sodium: Metal oxides MRH 2.05/37

Other reactants

Yoshida, 1980, 127

MRH values calculated for 7 combinations, largely with oxidisable materials, are given.

Phospham

See Phospham: Oxidants

Phthalic anhydride

'Leaflet No. 5', Inst. of Chem., London, 1940

A mixture of the anhydride and anhydrous oxide exploded violently on heating.

Reducants

MRH Hydrazine 1.17/17

Mellor, 1941, Vol. 3, 137

Interaction with hydroxylamine or hydrazine is vigorous.

See other METAL OXIDES

4282. Copper(II) sulfate

[18939-61-2]



HCS 1980, 327

Hydroxylamine

See Hydroxylamine: Copper(II) sulfate

See other METAL OXONON-METALLATES

4283. Copper monophosphide

[12517-41-8]



See Copper(II) phosphide *See other* METAL NON-METALLIDES

4284. Copper diphosphide

[12019-11-3]



See Copper(II) phosphide *See other* METAL NON-METALLIDES

4285. Copper(II) sulfide
[1317-40-4]

CuS

CuS

Chlorates

Mellor, 1956, Vol. 2, Suppl. 1, 584

Copper(II) sulfide explodes in contact with magnesium chlorate, zinc chlorate or cadmium chlorate, or with a conc. solution of chloric acid.

See METAL HALOGENATES

Other reactants

Yoshida, 1980, 391

MRH values calculated for 12 combinations with various reagents are given.

See other METAL SULFIDES

4286. Copper(I) hydride
[13517-00-5]

HCuCuH

Cu₂H₂

Gibson, 1969, 77

The dry hydride ignites in air.

Halogens

Mellor, 1941, Vol. 3, 73

Ignition occurs on contact with fluorine, bromine or iodine.

See other METAL HYDRIDES

4287. Copper(I) azide
[14336-80-2]

N₃CuCuN₃

Cu₂N₆

1. Mellor, 1940, Vol. 8, 348; 1967, Vol. 8, Suppl. 2.2, 42–50

2. Urbanski, 1967, Vol. 3, 185

It is highly endothermic (ΔH_f° (s) 253.1 kJ/mol, 2.40 kJ/g). One of the more explosive metal azides, it decomposes at 205°C [1], and is very highly impact-sensitive [2].

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

4288. Copper(I) oxide
[1317-39-1]

Cu₂O

Cu₂O

Aluminium

Moddeman, W. E. *et al.*, *Chem. Abs.*, 1981, **84**, 159186

1503

The role of surface chemistry in reactivity of this thermite combination has been studied.

See other THERMITE REACTIONS

Lithium nitride

See Lithium nitride: Copper(I) chloride, etc.

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

See other METAL OXIDES

4289. Copper(I) nitride

[1308-80-1]

Cu₃N

Cu₃N

It is mildly endothermic (ΔH_f° (s) +74.5 kJ/mol, 0.36 kJ/g).

Mellor, 1940, Vol. 8, 100

It may explode on heating in air.

Nitric acid

See Nitric acid: Copper(I) nitride

See other N-METAL DERIVATIVES

4290. Copper(II) phosphide

[12134-35-9]

Cu₃P₂

Cu₃P₂

Oxidants

Mellor, 1940, Vol. 8, 839

The powdered phosphide burns vigorously in chlorine. Mixtures with potassium chlorate explode on impact, and with potassium nitrate, on heating. The monophosphide and diphosphide behave similarly.

See Potassium chlorate: Metal phosphides

See other METAL NON-METALLIDES

†4291. Deuterium oxide

[7789-20-0]

D₂O

D₂O

1. Anderson, C., *Nature*, 1992, **355**(6356), 102

2. Baum, R., *Chem. Eng. News*, 1992, **70**(26), 8

3. Zhang, X. *et al.*, *Front. Sci. Ser.*, 1993, (4[Frontiers of Cold Fusion]), 381

Electrolysis of heavy water over palladium electrodes is claimed to give inexplicable energy release—'Cold Fusion'. It can certainly kill [1], even though the mechanism be ignition of an explosive mixture of deuterium and oxygen resulting

from past electrolysis after a platinum recombination catalyst for the off-gases failed and a safety valve blocked. The cell then exploded when the experimenter picked it up; ignition presumably at a catalytic surface exposed by moving liquid [2]. Others hold out for less probable causes [3].

Pentafluorophenyllithium

See Pentafluorophenyllithium: Deuterium oxide

See Diprotium monoxide

See other NON-METAL OXIDES

4292. Europium

[7740-53-1]

Eu

Eu

Bailar, 1973, Vol. 4, 69

Europium is the most reactive lanthanide metal, and may ignite on exposure to air if finely divided.

See LANTHANIDE METALS: oxidants

See other METALS, PYROPHORIC METALS

4293. Europium(II) sulfide

[12020-65-4]

EuS

EuS

Sidgwick, 1950, 454

Pyrophoric in air.

See other METAL SULFIDES, PYROPHORIC MATERIALS

4294. Hydrogen fluoride

[7664-39-3]

HF

HF

(MCA SD-25, 1970); Soln. NSC 459, 1978; HCS 1980, 543; Gas HCS 1980, 548

1. Keen, M. J. *et al.*, *Chem. & Ind.*, 1957, 805

2. Braker, 1980, 391

3. Anon., *Lab. Hazards Bull.*, 1991, **11**(09), 556

4. Watson, E. F., *Chem. Eng. News*, 1997, **75**(17), 28, 6

Handling precautions for the gas or anhydrous liquid are detailed [1], and a polythene condenser for disposal of hydrogen fluoride is described [2]. A steel cylinder of the nominally anhydrous material exploded after 42 years. This is attributed to ppm levels of water, which catalyses reaction with the steel to produce hydrogen, slowly increasing the pressure. It is recommended that cylinders should be vented at yearly intervals, and those more than two years old returned to the manufacturer [3]. More warnings against prolonged storage in steel are given [4].

Bismuthic acid

See Bismuthic acid: Hydrofluoric acid

Cyanogen fluoride

See Cyanogen fluoride: Hydrogen fluoride

Glycerol, Nitric acid

See Nitric acid: Glycerol, etc.

Mercury(II) oxide, Organic materials

Ormston, J., *School Sci. Rev.*, 1944, **26**(98), 32

During the fluorination of organic materials by passing hydrogen fluoride into a vigorously stirred suspension of the oxide (to form transiently mercury difluoride, a powerful fluorinator), it is essential to use adequate and effective cooling below 0°C to prevent loss of control of the reaction system.

Metal alloys

Young, J. A., *Chem. Health and Safety*, 1995, **2**(2), 6

It is reported that an aluminium cleaner containing low concentrations of hydrofluoric acid can generate stibine from antimony containing bearing-metal alloys, to the permanent detriment of the health of nearby workers. Presumably arsine could appear from arsenic containing alloys; both are gases and extremely toxic.

Methanesulfonic acid

Gramstad, T. *et al.*, *J. Chem., Soc.*, 1956, 173–180

Electrolysis of a mixture produced oxygen difluoride which exploded.

See Oxygen difluoride

Nitric acid, Lactic acid

See Nitric acid: Hydrofluoric acid, Lactic acid

Nitric acid, Propylene glycol

See Nitric acid: Hydrofluoric acid, Propylene glycol

Other reactants

Yoshida, 1980, 311

MRH values calculated for 7 combinations with various reagents are given.

Oxides

MRH Calcium oxide 2.84/59

Mellor, 1956, Vol. 2, Suppl. 1, 122; 1939, Vol. 9, 101

Arsenic trioxide and calcium oxide incandesce in contact with liquid hydrogen fluoride.

N-Phenylazopiperidine

See *N*-Phenylazopiperidine: Hydrofluoric acid

Phosphorus(V) oxide

See Tetraphosphorus decaoxide: Hydrogen fluoride

Potassium permanganate
See Potassium permanganate: Hydrofluoric acid

Potassium tetrafluorosilicate(2-)
Mellor, 1956, Vol. 2, Suppl. 1, 121
Contact with liquid hydrogen fluoride causes violent evolution of silicon tetrafluoride. (The same is probably true of metal silicides and other silicon compounds generally.)
See Dialuminium octavanadium tridecasilicide

Sodium MRH 6.94/53
See Sodium: Acids

Sulfuric acid
See Sulfuric acid: Hydrofluoric acid
See other INORGANIC ACIDS, NON-METAL HALIDES, NON-METAL HYDRIDES

4295. Fluoroselenic acid
[14986-53-9]



Cellulose
Bartels, H. *et al.*, *Helv. Chim. Acta*, 1962, **45**, 179
A strong oxidant which reacts violently with filter paper or similar organic matter, igniting it if dry.
See other INORGANIC ACIDS, OXIDANTS

4296. Fluoramine (Fluoramide)
[15861-05-9]



1. Hoffmann, C. J., *et al.*, *Chem. Rev.*, 1962, **62**, 7
2. Minkwitz, R. *et al.*, *Z. Naturforsch. B, Chem. Sci.*, 1988, **43B**, 1478–1480
The impure material is very explosive [1]. The pure material, prepared by a thermolytic method, is stable as a solid at -103°C , but on melting it decomposes to ammonium hydrogen difluoride and nitrogen [2].
See other N-HALOGEN COMPOUNDS

4297. Fluorophosphoric acid
[13537-32-1]



Sodium tetrahydroborate
See Sodium tetrahydroborate: Acids
See other INORGANIC ACIDS

4298. Fluorosilane
[13537-33-2] **FH₃Si**



Azidogermane
See Azidogermane: Fluorosilane
See other HALOSILANES

4299. Ammonium fluoride
[16099-75-5] **FH₄N**



Chlorine trifluoride
See Chlorine trifluoride: Ammonium fluoride
See related METAL HALIDES

4300. Potassium fluoride hydrogen peroxidate
[32175-44-3] **FK.H₂O₂**



See entry CRYSTALLINE HYDROGEN PEROXIDATES

4301. Manganese fluoride trioxide
[22143-20-0] **FMnO₃**



Alone, or Organic compounds, or Water
Engelbrecht, A. *et al.*, *J. Amer. Chem. Soc.*, 1954, **76**, 2042–2044
It decomposes, usually explosively, above 0°C or in contact with moisture. It is a powerful oxidant and reacts violently with organic compounds.
See related METAL HALIDES, METAL OXIDES *See other* OXIDANTS

4302. Nitrosyl fluoride
[7789-25-5] **FNO**



Haloalkene (unspecified)
MCA Case History No. 928
Interaction of a mixture in a pressure vessel at –78°C caused it to rupture when moved from the cooling bath.

Metals, or Non-metals
1. Schmutzler, R., *Angew. Chem. (Intern. Ed.)*, 1968, **7**, 442
2. Pascal, 1956, Vol. 10, 346

Antimony, bismuth, arsenic, boron, red phosphorus, silicon [1] and tin [2] all react with incandescence.

Oxygen difluoride

Ruff, O. *et al.*, *Z. Anorg. Chem.*, 1932, **208**, 293

Explosion occurs on mixing, even at low temperatures.

Sodium

Mellor, 1940, Vol. 8, 612

Reaction is incandescent.

See other N-HALOGEN COMPOUNDS, NON-METAL HALIDES, OXIDANTS

4303. Nitryl fluoride

[10022-50-1]

FNO₂

O₂NF

Metals

Aynsley, E. E. *et al.*, *J. Chem. Soc.*, 1954, 1122

When nitryl fluoride is passed at ambient temperature over molybdenum, potassium, sodium, thorium, uranium or zirconium, glowing or white incandescence occurs. Mild warming is needed to initiate similar reactions of aluminium, cadmium, cobalt, iron, nickel, titanium, tungsten, vanadium or zinc, and 200–300°C for lithium or manganese.

Non-metals

Aynsley, E. E. *et al.*, *J. Chem. Soc.*, 1954, 1122

Boron and red phosphorus glow in the fluoride at ambient temperature, while hydrogen explodes at 200–300°C. Carbon and sulfur are also attacked.

Sodium azide

See Sodium azide: Nitryl fluoride

See other N-HALOGEN COMPOUNDS, NON-METAL HALIDES, OXIDANTS

4304. Nitryl hypofluorite ('Fluorine nitrate')

[7789-26-6]

FNO₃

O₂NOF

1. Schmutzler, R., *Angew. Chem. (Intern. Ed.)*, 1968, **7**, 454

2. Engelbrecht, A., *Monatsh.*, 1964, **95**, 633

3. Lustig, M. *et al.*, *Adv. Fluorine Chem.*, 1973, **7**, 189

4. Christe, K. O. *et al.*, *Inorg. Chem.*, 1987, **56**, 920

It is mildly endothermic (ΔH_f° (g) +10.4 kJ/mol, 0.12 kJ/g) but a powerful oxidant.

It is a toxic colourless gas which is dangerously explosive in the gaseous, liquid and solid states [1]. It is produced during electrolysis of nitrogenous compounds in hydrogen fluoride [2]. Later work (perhaps with purer material?) did not show the explosive instability [3]. The shock-sensitivity is confirmed [4].

Gases

Hoffmann, C. J., *Chem. Rev.*, 1964, **64**, 94

Immediate ignition in the gas phase occurs with ammonia, dinitrogen oxide or hydrogen sulfide.

Organic materials

Brauer, 1963, 189

The very powerful liquid oxidant explodes when vigorously shaken, or immediately on contact with alcohol, ether, aniline or grease. It is also sensitive in the vapour or solid state (but see reference 3 above).

See Nitryl hypochlorite

See other ENDOTHERMIC COMPOUNDS, OXIDANTS

4305. Sulfur oxide-(N-fluorosulfonyl)imide (Sulfinylsulfamoyl fluoride)

[16829-30-4]

FNO₃S₂



Water

Roesky, H. W., *Angew. Chem. (Intern. Ed.)*, 1967, **6**, 711

It reacts explosively with water at ambient temperature, but smoothly at -20°C .

See other ACYL HALIDES, N-S COMPOUNDS

4306. Thiazyl fluoride

[18820-63-8]

FNS



Alkylbutadienes

Bludssus, W. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 35–36

Interaction with, for example, 2,3-dimethylbutadiene is explosive.

See other NON-METAL HALIDES, N-S COMPOUNDS

4307. Fluorine azide

[14986-60-8]

FN₃



1. Bauer, S. H., *J. Amer. Chem. Soc.*, 1947, **69**, 3104

2. Gholivand, K. *et al.*, *Inorg. Chem.*, 1987, **26**, 2137–2140

This unstable material usually explodes on vaporisation (at -82°C) [1]. It is extremely explosive in the liquid and solid states. A safe method has been developed for preparing the pure gas on 20 mg scale. It may be stored safely at 10–20 mbar/ -80°C for several months. Cooling the gas to -196°C , or evaporation of the liquid at a fast rate may lead to very violent explosions. Fluorine azide is now described as triazadienyl fluoride [2], as shown above.

See other HALOGEN AZIDES

4308. Fluorothiophosphoryl diazide

[38005-27-5]

FN₆PSO'Neill, S. R. *et al.*, *Inorg. Chem.*, 1972, **11**, 1630

The explosion of the glassy material at -183°C was attributed to crystallisation of the glass.

See other ACYL AZIDES**4309. Rubidium fluoroxy sulfate**

[[73347-64-5] (ion)]

FO₄RbSAppelman, E. H. *et al.*, *J. Amer. Chem. Soc.*, 1979, **101**, 388

This and the caesium analogue, both powerful oxidants, detonate mildly at 100°C, evolving oxygen.

See other ACYL HYPOHALITES, OXIDANTS**4310. Fluorine**

[7782-41-4]

F₂

FPA H79, 1979 (cylinder); HCS 1980, 503 (cylinder)

1. Kirk-Othmer, 1966, Vol. 9, 506
2. Gall, J. F. *et al.*, *Ind. Eng. Chem.*, 1947, **39**, 262
3. Landau, R. *et al.*, *Ind. Eng. Chem.*, 1947, **39**, 262
4. Turnbull, S. G. *et al.*, *Ind. Eng. Chem.*, 1947, **39**, 286
5. Long, G., *Apparatus for Disposal of Fluorine on a Laboratory Scale*, Harwell, AERE, 1956
6. Gordon, J. *et al.*, *Ind. Eng. Chem.*, 1960, **52**(5), 63A
7. Farrar, R. L. *et al.*, *Some Considerations in the Handling of Fluorine and the Chlorine Fluorides*, Union Carbide Oak Ridge Rept. K/ET-252 for US DoE, 1979
8. Ring, R. J. *et al.*, *Review of Fluorine Cells and Production*, Lucas Heights, Aus. At. En. Comm., 1973
9. Bailar, 1973, Vol. 2, 1015–1019
10. DesMarteau, D. D. *et al.*, *J. Amer. Chem. Soc.*, 1987, **109**, 7194–7196

Fluorine is the most electronegative and reactive element known, reacting, often violently, with most of the other elements and their compounds (note the large MRH values quoted below). Handling hazards and disposal of fluorine on a laboratory scale are adequately described [1,2,3,4,5][6], and a more general review is also available [7]. Safety practices associated with the use of laboratory- and industrial-scale fluorine cells and facilities have been reviewed [8]. Equipment and procedures for the laboratory use of fluorine and volatile fluorides have been

detailed [9]. A new series of *N*-fluorosulfonimides shows promise as milder and safer reagents than elemental fluorine for aromatic fluorination [10].

Acetonitrile, Chlorine fluoride

Sekiya, A. *et al.*, *Inorg. Chem.*, 1981, **20**, 1–2

When fluorine was condensed onto acetonitrile and chlorine fluoride frozen at -196°C , a small explosion occurred in the reactor.

Acetylene

MRH 11.92/41

See Acetylene: Halogens

Alkanes, Oxygen

Von Elbe, G., US Pat. 3 957 883, 1976

Interaction of propane, butane or 2-methylpropane with fluorine and oxygen produces peroxides. Appropriate reaction conditions are necessary to prevent explosions.

Ammonia

MRH 10.25/23

1. Mellor, 1940, Vol. 8, 216

2. Müller, W., *Umsch. Wiss. Tech.*, 1974, **74**(15), 485–486

Ammonia ignites in contact with fluorine [1], and the anhydrous liquids have been used as a propellant pair [2]. Aqueous ammonia ignites or explodes on contact [1].

See Water, below

Boron nitride

Moissan, 1900, 232

Unheated interaction leads to incandescence.

Caesium fluoride, Fluorocarboxylic acids

Sekiya, A. *et al.*, *Inorg. Chem.*, 1980, **19**, 1329

Low temperature fluorination of fluorocarboxylic acids to give explosive 1,1-bis(fluoroxy)perfluoroalkanes occasionally led to explosive reactions. Thus difluoroacetic acid led to explosion at -195° , and perfluorosuccinic acid at -111° or -20°C .

See BIS(FLUOROOXY)PERHALOALKANES

Caesium heptafluoropropoxide

1. Gumprecht, W. H., *Chem. Eng. News*, 1965, **43**(9), 36

2. *MCA Case History No. 1045*

Fluorination of caesium heptafluoropropoxide at -40°C with nitrogen-diluted fluorine exploded violently after 10 h. This may have been caused by ingress of moisture, formation of some pentafluoropropionyl derivative and conversion of this to pentafluoropropionyl hypofluorite, known to be explosive if suitably initiated. Other possible explosive intermediates are peroxides or peresters.

See Pentafluoropropionyl hypofluorite

Ceramic materials

Farrar, R. L. *et al.*, *US DoE Rept. K/ET-252*, 1979, 9

Even finely divided ceramic materials may be ignited in fluorine.

Covalent halides

1. Mellor, 1940, Vol. 2, 12; 1956, Vol. 2, Suppl. 1, 64; 1940, Vol. 8, 995, 1003, 1013

2. Leleu, *Cahiers*, 1973, (73), 509; 1974, (74), 427

Chromyl chloride at high concentration ignites in fluorine, while phosphorus pentachloride, phosphorus trichloride and phosphorus trifluoride ignite on contact [1]. Boron trichloride ignites in cold fluorine, and silicon tetrachloride on warming [2].

Cyanogen (Dicyanogen)

Klapötke, T. M. *et al.*, *Angew. Chemie (Int.)*, 1991, **30**(11), 1485

An explosion was experienced on allowing a mix of these reagents, with arsenic pentafluoride, which had been uv irradiated at -196°C , to warm to -85°C .

See Halogens, below

Cyanoguanidine

Rausch, D. A. *et al.*, *J. Org. Chem.*, 1968, **33**, 2522

The products, perfluoro-1-aminomethylguanidine and perfluoro-*N*-aminomethyltri-aminomethane, and by-products of the reaction of fluorine with cyanoguanidine are extremely explosive in gas, liquid and solid states.

See Sodium dicyanamide, below

1- or 2-Fluoriminoperfluoropropane

Sekiya, A. *et al.*, *J. Fluorine Chem.*, 1981, **17**, 463–468

Interaction is explosive.

Graphite

1. Lagow, R. J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1974, 1270

2. Ruff, O. *et al.*, *Z. Anorg. Chem.*, 1934, **217**, 1

3. Simons, J. H. *et al.*, *J. Amer. Chem. Soc.*, 1939, **61**, 2962–2966

4. Farrar, R. L. *et al.*, *US DoE Rpt. K/ET-252*, 1979, 33–34

5. Kita, Y. *et al.*, *Chem. Abs.*, 1983, **99**, 107531

During interaction at ambient temperature in a bomb to produce poly(carbon monofluoride), admission of fluorine beyond a pressure of 13.6 bar must be extremely slow and carefully controlled to avoid a violently exothermic explosion [1]. Previously it had been shown that explosive interaction of carbon and fluorine was due to the formation and decomposition of the graphite intercalation compound, poly(carbon monofluoride) [2]. Presence of mercury compounds prevents explosion during interaction of charcoal and fluorine [3]. Reaction of surplus fluorine with graphite or carbon pellets was formerly used as a disposal method, but is no longer recommended. Violent reactions observed when an exhausted trap was opened usually involved external impact on the metal trap, prodding the trap contents to empty the trap, or possibly ingress of moist air

[4]. Removal of higher fluorocarbons (above C₄) from the circulating gas stream prevents explosive decomposition of the graphite fluoride [5].

Halocarbons MRH Carbon tetrachloride 2.34/67, chloroform 3.77/56

1. Mellor, 1956, Vol. 2, Suppl. 1, 198
2. Schmidt, 1967, 82
3. Fletcher, E. A. *et al.*, *Combust. Inst. Paper WSS/CI-67-23*, 1967
4. Moissan, 1900, 241
5. Farrar, R. L. *et al.*, *US DoE Rept. K/ET-252*, 1979, 36–40

The violent or explosive reactions which carbon tetrachloride, chloroform, etc., exhibit on direct local contact with gaseous fluorine [1], can be moderated by suitable dilution, catalysis and diffused contact [2]. Combustion of perfluorocyclobutane–fluorine mixtures was detonative between 9.04 and 57.9 vol% of the halocarbon [3]. Iodoform reacts very violently with fluorine owing to its high iodine content [4]. Explosive properties of mixtures with 1,2-dichlorotetrafluoroethane have been studied [5].

See Poly(tetrafluoroethylene), etc., below

Halogens, or Dicyanogen MRH values below references

1. Mellor, 1940, Vol. 2, 12
2. Sidgwick, 1950, 1148
3. Tari, I. *et al.*, *Inorg. Chem.*, 1979, **18**, 3205

MRH Bromine 1.38/89, chlorine 1.67/38, iodine 3.89/43

While bromine, iodine and dicyanogen all ignite in fluorine at ambient temperature [1], a mixture of chlorine and fluorine (containing essential moisture) needs sparking before ignition occurs, though an explosion immediately follows [2]. Heating fluorine and chlorine in a Monel pressure vessel gives contained explosions at around 100°C [3].

See Cyanogen, above

Hexalithium disilicide

Mellor, 1940, Vol. 6, 169

It incandescens on warming in fluorine.

Hydrocarbons MRH Anthracene 7.57/49, acetylene 11.92/41

1. Mellor, 1940, Vol. 2, 11; 1956, Vol. 2, Suppl.1, 198
2. Sidgwick, 1950, 1117
3. Schmidt, 1967, 82
4. Moissan, 1900, 240–241

Violent explosions occur when fluorine directly contacts liquid hydrocarbons, even at –210 with anthracene or turpentine, or solid methane at –190°C with liquid fluorine. Many lubricants ignite in fluorine [1,2]. Contact and reaction under carefully controlled conditions with catalysis can now be effected smoothly [3]. Gaseous hydrocarbons (town gas, methane) ignite in contact with fluorine, and mixtures with unsaturated hydrocarbons (ethylene, acetylene) may explode on exposure to sunlight. Each bubble of fluorine passed through benzene causes ignition, but a rapid stream may lead to explosion [4].

Hydrofluoric acid

See Hydrogen halides, below

Hydrogen

MRH 13.39/5

1. Mellor, 1940, Vol. 2, 11; 1956, Vol. 2, Suppl.1, 55
2. Sidgwick, 1950, 1102
3. Kirshenbaum, 1956, 46
4. Gmelin, 1980, Fluorine, Suppl. Vol. 2
5. Truby, F. K., *J. Appl. Phys.*, 1978, **49**, 3481–3484

The violently explosive reactions which sometimes occur when the 2 elements come into contact under conditions ranging from solid fluorine and liquid hydrogen at -252°C to the mixed gases at ambient temperature [1] are caused by the catalytic effects of impurities or the physical nature of the walls of the containing vessel [2]. Even in absence of such impurities, spontaneous explosions still occur in the range of 75–90 mol% fluorine in the gas phase [3]. Recent data, including the inhibiting effect of small amounts of oxygen, has been reviewed [4], but presence of relatively large amounts does not necessarily inhibit explosions [5].

Hydrogen, Oxygen

Getzinger, R. W., *Rept. LA-5659*, Los Alamos Sci. Lab., 1974

The conditions under which mixtures of the gases at 1 bar and ambient temperature will react non-explosively have been studied.

Hydrogen fluoride, Seleninyl fluoride

Seppelt, K., *Angew. Chem. (Intern. Ed.)*, 1972, **11**, 630

Preparation of pentafluoroorthoselenic acid from the above reagents in an autoclave above ambient temperature caused occasional explosions. A safer alternative preparation is described.

Hydrogen halides

MRH HBr 2.30/8, HCl 3.18/65, HI 1.97/87

Mellor, 1940 Vol. 2, 12

Hydrogen bromide, hydrogen chloride and hydrogen iodide ignite in contact with fluorine, and the conc. aqueous solutions, including that of hydrogen fluoride, also produce flame.

Hydrogen sulfide

See Sulfides, below

Ice

1. English, W. D. *et al.*, *Cryog. Technol.*, 1965, **1**, 260
2. *UK Scientific Mission Report 68/79*, Washington, UKSM, 1968

Mixtures of liquid fluorine and ice are highly impact-sensitive, with a power comparable to that of TNT. Contact of moist air or water with liquid fluorine can thus be very hazardous [1,2].

See Water, below

Metal acetylides and carbides

Mellor, 1946, Vol. 5, 849, 885, 890–891

Monocaesium acetylide and caesium acetylide, lithium acetylide and rubidium acetylide, tungsten carbide and ditungsten carbide, and zirconium dicarbide all ignite in cold fluorine, while uranium dicarbide ignites in warm fluorine.

Metal borides

Bailar, 1973, Vol. 1, 729

Interaction frequently attains incandescence.

Metal cyanocomplexes

Moissan, 1900, 228

Potassium hexacyanoferrate(II), lead hexacyanoferrate(III) and potassium hexacyanoferrate(III) become incandescent in fluorine, and the liberated dicyanogen also ignites.

Metal hydrides

Mellor, 1940, Vol. 2, 12, 483; 1956, Vol. 2, Suppl. 1, 56; 1941, Vol. 3, 73

Copper hydride, potassium hydride and sodium hydride all ignite on contact with fluorine at ambient temperature.

Metal iodides

Moissan, 1900, 227

Fluorine decomposes calcium iodide, lead iodide, mercury iodide and potassium iodide at ambient temperature, and the liberated iodine ignites, evolving much heat.

Metal oxides

MRH Calcium oxide 6.15/60

Mellor, 1940, Vol. 2, 13, 469; 1941, Vol. 3, 663; 1942, Vol. 13, 715; 1936, Vol. 15, 380, 399

Oxides of the alkali and alkaline earth metals and nickel(II) oxide incandesce in cold fluorine, and iron(II) oxide when warmed. Nickel(IV) oxide also burns in fluorine.

Metals

MRH Caesium 3.43/28, calcium 15.56/51, copper 3.01/76, lead 3.39/76, lithium 23.51/27, magnesium 17.78/39, manganese 8.91/49, molybdenum 7.57/46, potassium 9.58/68, rubidium 5.19/82, sodium 13.43/55, uranium 6.02/72, zinc 2.45/63

1. Mellor, 1940, Vol. 2, 13, 469; 1941, Vol. 3, 638; 1940, Vol. 4, 267, 476; 1946, Vol. 5, 421; 1939, Vol. 9, 379, 891; 1943, Vol. 11, 513, 730; 1942, Vol. 12, 344; 1942, Vol. 15, 675

2. Schmidt, 1967, 78–80

3. Kirk-Othmer, 1966, Vol. 9, 507–508

The vigour of reaction is greatly influenced by the state of sub-division of the metals involved. Massive calcium, moist magnesium, manganese powder, molybdenum powder, potassium, sodium, rubidium and antimony all ignite in cold fluorine gas. Warm tantalum powder or cold thallium ignites on contact with fluorine. Fine copper wire (as wool) ignites at 121°C, and osmium and tin begin to burn at 100°C, while iron powder (100-mesh, but not 20-mesh) ignites in liquid

fluorine. Titanium will ignite if impacted under the liquid at -188°C and has ignited in presence of catalysts in the gas at -80°C , but in all cases the fluoride film prevents further propagation. Tungsten and uranium powders ignite in the gas without heating, while zinc ignites at about 100°C . Molybdenum, tungsten and Monel wires ignited in atmospheric fluorine at 205, 283 and 396°C (averaged values), respectively. Generally, strongly electropositive metals, or those forming volatile fluorides are attacked the most vigorously.

Metal salts MRH Potassium dichromate 2.97/66, potassium permanganate 3.18/67, sodium chlorate 2.34/85, sodium nitrate 1.34/56

1. Mellor, 1940, Vol. 2, 13; 1956, Vol. 2, Suppl. 1, 63
2. Schmidt, 1967, 83–84
3. Moissan, 1900, 228–239
4. Pascal, 1960, Vol. 16, 57–60
5. Fichter, F. *et al.*, *Helv. Chim. Acta*, 1930, **13**, 99–102

Calcium carbonate, lead carbonate, basic lead carbonate and sodium carbonate all ignite and burn fiercely in contact with fluorine. Chlorides and cyanides are vigorously attacked by cold fluorine, including lead fluoride and thallium(I) chloride, both of which become molten. Mercury(II) cyanide ignites in fluorine when warmed gently, and silver cyanide reacts explosively when cold [1]. Sodium metasilicate ignites in fluorine [2]. Unheated calcium phosphate [3], sodium thio-sulfate and sodium diphosphate [4] all incandesce in contact with fluorine, and barium thiocyanate or mercury thiocyanate ignites [3]. On warming, chromium(III) chloride [4], calcium arsenate or copper borate incandesce, and sodium arsenate ignites [3]. Introduction of fluorine into solutions of silver fluoride, silver nitrate, silver perchlorate or silver sulfate causes violent exothermic reactions to occur, with liberation of ozone-rich oxygen [5].

Metal silicides

Mellor, 1940, Vol. 6, 169, 178

Calcium disilicide readily ignites, and lithium hexasilicide becomes incandescent, when warmed in fluorine.

Miscellaneous materials

1. Mellor, 1940, Vol. 2, 13
2. Schmidt, 1967, 84, 110
3. Lafferty, R. H. *et al.*, *Chem. Eng. News*, 1948, **26**, 3336

Town gas ignites in contact with gaseous fluorine, as does a mixture of lead oxide and glycerol (formerly used as a jointing compound) [1]. Spillage tests involving action of liquid fluorine alone or as a 30% solution in liquid oxygen caused asphalt and crushed limestone to ignite, and coke and charcoal to burn, the latter brilliantly, while JP4 liquid hydrocarbon fuel produced violent explosions and a large fireball. Humus-rich soil also burned with a bright flame [2]. Immersion of various glove materials in liquid fluorine was examined. Cotton exploded violently and Neoprene slightly with ignition, while leather charred but did not ignite [3].

Nitric acid

MRH 0.50/53

1. Cady, G. H., *J. Amer. Chem. Soc.*, 1934, **56**, 2635
2. Schmutzler, R., *Angew. Chem. (Intern. Ed.)*, 1968, **8**, 453

Interaction of fluorine with either the concentrated or very dilute acid caused explosions, while use of 4 N acid did not [1]. Later and safer methods of preparing nitryl hypofluorite are summarised [2].

Nitrogenous bases

MRH Dimethylamine 10.63/23, pyridine 8.28/45

Hoffman, C. J., *Chem. Rev.*, 1962, **62**, 12

Aniline, dimethylamine and pyridine incandesce on contact with fluorine.

See Ammonia, and Cyanoguanidine, both above

Non-metal oxides

MRH Carbon monoxide 6.44/42, dinitrogen tetraoxide 1.00/45,
nitrogen oxide 2.93/99+, sulfur dioxide 5.10/36

1. Mellor, 1940, Vol. 2, 11; 1939, Vol. 9, 101
2. Hoffman, C. J. *et al.*, *Chem. Rev.*, 1962, **62**, 10
3. Pascal, 1960, Vol. 16, 53, 62
4. Moissan, 1900, 138

Arsenic trioxide reacts violently and nitrogen oxide ignites in excess fluorine. Bubbles of sulfur dioxide explode separately on contacting fluorine, while addition of the latter to sulfur dioxide causes an explosion at a certain concentration [1]. Reaction of fluorine with dinitrogen tetraoxide usually causes ignition [2]. Interaction with carbon monoxide may be explosive. Anhydrous silica incandesces in the gas, and interaction with liquid fluorine at -80°C is explosive [3,4]. Boron trioxide also incandesces in the gas [3].

See Bis(fluoroformyl) peroxide

Non-metals

MRH Arsenic 7.19/57, silicon 14.73/27, sulfur 8.28/22

1. Mellor, 1940, Vol. 2, 11, 12; 1956, Vol. 2, Suppl. 1, 60; 1946, Vol. 5, 785, 822; 1940, Vol. 6, 161; 1939, Vol. 9, 34; 1943, Vol. 11, 26
2. Schmidt, 1967, 52, 107
3. Pascal, 1960, Vol. 16, 58
4. Moissan, 1900, 125–128

Boron, phosphorus (yellow or red), selenium, tellurium and sulfur all ignite in contact with fluorine at ambient temperature, silicon attaining a temperature above 1400°C [1]. The reactivity shown by various forms of carbon (charcoal, lampblack, soot) all of which ignite and burn vigorously in fluorine [1] has been reported to be due to presence of various impurities, moisture and hydrocarbons [1,2]. Carefully purified carbon (massive graphite) is inert to fluorine at ambient or slightly elevated temperatures for a short period but may then react explosively [2]. Phosphorus [3] and sulfur incandesce in liquid fluorine, and sulfur ignites even at -188°C [4].

See Graphite, above

Other reactants

Yoshida, 1980, 307–310

MRH values calculated for 58 combinations with a very wide variety of other reagents are given. Many of the values are extremely high.

Oxygenated organic compounds MRH Acetaldehyde 8.33/19, butanol 9.12/33,
dimethylformamide 8.28/43, methanol 8.37/22

1. Pascal, 1960, Vol. 16, 65
2. Moissan, 1900, 242–245
3. Mellor, 1940, Vol. 2, 13

Methanol, ethanol and 3-methylbutanol [1], acetaldehyde, trichloroacetaldehyde [2] and acetone [3] all ignite in contact with gaseous fluorine. Lactic acid, benzoic acid and salicylic acid ignite, while gallic acid becomes incandescent. Ethyl acetate and methyl borate ignite in fluorine [2].

Perchloric acid

See Perchloric acid: Fluorine

Phosphorus halides

See Covalent halides, above

Polymeric materials, Oxygen

Schmidt, 1967, 87

Various polymeric materials were tested statically with both gaseous and liquefied mixtures of fluorine and oxygen containing from 50 to 100% of the former. The materials which burned or reacted violently were: phenol–formaldehyde resins (Bakelite); polyacrylonitrile–butadiene (Buna N); polyamides (Nylon); polychloroprene (Neoprene); polyethylene; polytrifluoropropylmethylsiloxane (LS63); polyvinyl chloride–vinyl acetate (Tygan); polyvinylidene fluoride–hexafluoropropylene (Viton); polyurethane foam. Under dynamic conditions of flow and pressure, the more resistant materials which burned were: chlorinated polyethylenes, polymethyl methacrylate (Perspex); polytetrafluoroethylene (Teflon).

Polytetrafluoroethylene

Appelman, E. H., *Inorg. Chem.*, 1969, **8**, 223

Teflon tubing, when used to conduct fluorine into a reaction mixture, sometimes ignites. Combustion stops when the flow of fluorine is shut off.

Polytetrafluoroethylene, Trichloroethylene

Lawler, A. E., *Advan. Cryog. Eng.*, 1967, **12**, 780–783

Tests showed that Teflon gaskets containing more than 0.35 wt% of sorbed trichloroethylene were potentially hazardous in contact with liquid fluorine.

See Halocarbons, above

Potassium chlorate

See Potassium chlorate: Fluorine

Potassium hydroxide

1. Fichter, F. *et al.*, *Helv. Chim. Acta*, 1927, **10**, 551
2. Kacmarek, A. J. *et al.*, *Inorg. Chem.*, 1962, **1**, 659–661
3. Bailar, 1973, Vol. 2, 789–791

Interaction at -20°C produces potassium trioxide, a spontaneously explosive solid [1]. Later references suggest that these compounds are not spontaneously explosive [2,3].

Purge gases

Farrar, R. L. *et al.*, *US DoE Rept. K/ET-252*, 1979, 36

Purge gases used with liquid fluorine (-188°C or below) must be scrupulously dry and of low hydrocarbon content (<5 ppm), to prevent formation of ice crystals or solid hydrocarbons.

See Hydrocarbons, and Ice, both above

Sodium acetate

MRH 6.44/35

Mellor, 1956, Vol. 2, Suppl. 1, 562

Application of fluorine to aqueous sodium acetate solution causes an explosion, involving formation of diacetyl peroxide.

See Diacetyl peroxide

Sodium bromate

MRH 1.30/89

Appelman, E. H., *Inorg. Chem.*, 1969, **8**, 223

The oxidation of alkaline bromate by fluorine to perbromate is not smooth and small explosions may occur in the vapour above the solution. The reaction should not be run unattended.

Sodium dicyanamide

Rausch, D. A. *et al.*, *J. Org. Chem.*, 1968, **33**, 2522

The product, perfluoro-*N*-cyanodiaminomethane, and many of the by-products from interaction of fluorine and sodium dicyanamide, are extremely explosive in gaseous, liquid and solid states.

See Cyanoguanidine, above

Stainless steel

Stewart, J. W., *Proc. 7th Int. Conf. Low Temp. Phys.* (Toronto), 1960, 671

During the study of phase transitions of solidified gases at high pressures, solid fluorine reacted explosively with apparatus made from stainless steel.

Sulfides MRH Antimony(III) sulfide 6.40/44, carbon disulfide 8.24/25, hydrogen sulfide 9.29/18

1. Mellor, 1940, Vol. 2, 11, 13; 1940, Vol. 6, 110; 1938, Vol. 9, 522; 1947, Vol. 10, 133; 1943, Vol. 11, 430

2. Moissan, 1900, 231–232

Antimony trisulfide, carbon disulfide vapour, chromium(II) sulfide and hydrogen sulfide all ignite in contact with fluorine at ambient temperature, the solids becoming incandescent [1]. Iron(II) sulfide reacts violently on mild warming, and barium sulfide, potassium sulfide or zinc sulfide all incandesce in the gas, as does molybdenum(III) sulfide at 200°C [2].

Trinitromethane

Smith, W. L. *et al.*, *Chem. Abs.*, 1976, **84**, 46818

During preparation of fluorotrinitromethane, an instrumental method can be used to avoid occurrence of dangerous over-fluorination of nitroform.

Water

1. Mellor, 1940, Vol. 2, 11
2. Schmidt, 1967, 53, 119

Treatment of liquid air (containing condensed atmospheric moisture) with fluorine give a potentially explosive precipitate, thought to be fluorine hydrate [1]. Contact of liquid fluorine with a bulk of water causes violent explosions. Ice tends to react explosively with fluorine gas after an indeterminate induction period [2].

See other INDUCTION PERIOD INCIDENTS

Xenon, Catalysts

Levec, J. *et al.*, *J. Inorg. Nucl. Chem.*, 1974, **36**, 997–1001

Interaction may be explosive in the presence of finely divided nickel fluoride or silver difluoride, or nickel(III) oxide or silver(I) oxide, or if initiated by local heating. The mechanism is discussed.

See other HALOGENS, OXIDANTS

4311. Difluoramine (Fluorimide)

[10405-27-3]

F₂HN

F₂NH

1. Stevens, T. E., *J. Org. Chem.*, 1968, **33**, 2664, 2671
2. Petry, R. C. *et al.*, *J. Org. Chem.*, 1967, **32**, 4034
3. Rosenfeld, D. D. *et al.*, *J. Org. Chem.*, 1968, **33**, 2521
4. Martin, K. J., *J. Amer. Chem. Soc.*, 1965, **87**, 395
5. *MCA Case History No. 768*
6. Parker, C. O. *et al.*, *Inorg. Synth.*, 1970, **12**, 310–311
7. Christe, K. O., *Inorg. Chem.*, 1975, **14**, 2821
8. Christe, K. O. *et al.*, *Inorg. Chem.*, 1987, **26**, 920, 924
9. Chapman, R. D. *et al.*, *J. Org. Chem.*, 1998, **63**(5), 1566

It is a dangerous explosive and must be handled with skill and care and appropriate precautions [1,2]. Explosions have occurred when it was condensed at -196°C [3] or allowed to melt [4], and a glass bulb containing the gas exploded violently when accidentally dropped [5]. Although difluoramine may be condensed safely at -78 or -130° [6], or at -142° [7], liquid nitrogen should not be used to give a trapping temperature of -196°C , as explosions are very likely to occur [6]. The solid adduct with caesium fluoride prepared at -142°C always explodes if allowed to warm towards 0° , and the adduct with rubidium fluoride sometimes exploded [8]. A safer procedure for handling difluoroamine is reported [9].

Benzenediazonium tetrafluoroborate, Caesium fluoride

Baum, K., *J. Org. Chem.*, 1968, **33**, 4333

Use of caesium fluoride as base to effect condensation caused an explosion in absence of solvent. Pyridine or potassium fluoride, and use of dichloromethane gave satisfactory results.

See other N-HALOGEN COMPOUNDS

4312. Mercury(I) fluoride

[13967-25-4]

F₂Hg₂

FHgHgF

Iodoform

Andrews, L. *et al.*, *J. Fluorine Chem.*, 1979, **13**, 273–278

Attempted fluorination of iodoform with a 10 year-old sample of mercury(I) fluoride led to an explosion immediately on heating.

See other MERCURY COMPOUNDS, METAL HALIDES

4313. Krypton difluoride

[13773-81-4]

F₂Kr

KrF₂

ΔH_f° (g) 60 kJ/mol.

Ullmann, 1991, **A17**, 497

Krypton difluoride is a fluorinating agent some 50 kJ/mol more powerful than fluorine itself. It forms adducts, salts of FKr⁺, with high valency fluorides such as AsF₅ and SbF₅, these react explosively with organic compounds.

Arsenic pentafluoride

Christe, K. O. *et al.*, *Inorg. Chem.*, 1983, **22**, 3056

Interaction can lead to spontaneous exothermic decomposition of the fluoride accompanied by a bright flash and gas evolution. Safety precautions are required for this reaction system.

See other NON-METAL HALIDES (AND THEIR OXIDES)

See related XENON COMPOUNDS

4314. Difluorodiazene

[10578-16-2]

F₂N₂

FN=NF

Bensoam, J. *et al.*, *Tetrahedron Lett.*, 1977, 2797–2800

An explosion may occur when the diazene vapour is condensed, either to liquid or solid phases.

Both geometrical isomers are mildly endothermic (ΔH_f° (g) *cis*- +68.35 kJ/mol, 1.04 kJ/g; *trans*- +80.9 kJ/mol, 1.22 kJ/g).

Hydrogen

Kuhn, L. P. *et al.*, *Inorg. Chem.*, 1970, **9**, 60

Explosive interaction occurs above 90°C.
See other ENDOTHERMIC COMPOUNDS, *N*-HALOGEN COMPOUNDS

4315. Phosphorus azide difluoride
[37388-50-4]

F₂N₃P

N₃PF₂

1. O'Neill, S. R. *et al.*, *Inorg. Chem.*, 1972, **11**, 1630
2. Lines, E. L. *et al.*, *Inorg. Chem.*, 1972, **11**, 2270

It is photolytically and thermally unstable and has exploded at 25°C. It is also explosively sensitive to sudden changes in pressure, as occur on expansion into a vacuum or in surging during boiling [1]. It also ignites in air [2].

See related HALOPHOSPHINES, NON-METAL AZIDES, NON-METAL HALIDES

4316. Phosphorus azide difluoride–borane
[38115-19-4]

F₂N₃P.BH₃

N₃PF₂.BH₃

Lines, E. L. *et al.*, *Inorg. Chem.*, 1972, **11**, 2270

The liquid complex exploded violently during transfer operations.

See related BORANES, NON-METAL AZIDES, NON-METAL HALIDES

4317. Oxygen difluoride
[7783-41-7]

F₂O

OF₂

It is mildly endothermic (ΔH_f° (g) +23.0 kJ/mol, 0.43 kJ/g).

Adsorbents

1. Metz, F. I., *Chem. Eng. News*, 1965, **43**(7), 41
2. Streng, A. G., *Chem. Eng. News*, 1965, **43**(12), 5
3. Streng, A. G., *Chem. Rev.*, 1963, **63**, 611

Mixtures of silica gel and the liquid difluoride sealed in tubes at 334 mbar exploded above -196°C, presence of moisture rendering the mixture shock-sensitive at this temperature [1]. Reaction of oxygen difluoride with silica, alumina, molecular sieve or similar surface-active solids is exothermic, and under appropriate conditions may be explosive [2]. A quartz fibre can be ignited in the difluoride [3].

See other MOLECULAR SIEVE INCIDENTS

Combustible gases

Streng, A. G., *Chem. Rev.*, 1963, **63**, 612

Mixtures with carbon monoxide, hydrogen and methane are stable at ambient temperatures, but explode violently on sparking. Hydrogen sulfide explodes with oxygen difluoride at ambient temperature and, though interaction is smooth at

−78°C under reduced pressure, the white solid produced exploded violently when cooling was stopped.

Diborane

Rhein, R. A., *Combust. Inst. Paper WSCI-67-10*, 73–80, 1967

Although it reacts slowly at ambient temperature, a mixture of the components which is stable at −195°C could explode during warming to ambient conditions.

Diboron tetrafluoride

Holliday, A. K. *et al.*, *J. Chem. Soc.*, 1964, 2732

Ignition occurred on mixing at, or on warming mixtures to, −80°C.

Halogens, or Metal halides

Streng, A. G., *Chem. Rev.*, 1963, **63**, 611

Mixtures with chlorine, bromine or iodine explode on warming. A mixture with chlorine passed through a copper tube at 300°C exploded with variable intensity. Aluminium chloride explodes in the difluoride, and antimony pentachloride lightly at 150°C.

Hexafluoropropene, Oxygen

Afonso, M. Dos Santos *et al.*, *Chem. Abs.*, 1987, **107**, 175302

The thermal homogeneous chain reaction to give mainly octafluoropropane and hexafluoropropylene oxide becomes explosive above a minimum oxygen pressure of 26 mbar.

See other CATALYTIC IMPURITY INCIDENTS

Metals

Streng, A. G., *Chem. Rev.*, 1963, **63**, 611

Finely divided platinum group metals react on gentle warming, and coarser materials at higher temperatures; aluminium, barium, cadmium, magnesium, strontium, zinc and zirconium evolving light. Lithium, potassium and sodium incandesce brilliantly at 400°C, while tungsten explodes.

Nitrogen oxide

Streng, A. G., *Chem. Rev.*, 1963, **63**, 612

Gaseous mixtures may explode on sparking. The mixed gases slowly react to give a mixture (NO, NO₂) which, if liquefied by cooling, will explode on warming.

Nitrosyl fluoride

Hoffman, C. J. *et al.*, *Chem. Rev.*, 1962, **62**, 9

Solid mixture explodes on melting, and the gaseous components ignite on mixing.

Non-metals

1. Sidgwick, 1950, 1136

2. Streng, A. G., *Chem. Rev.*, 1963, **63**, 611

Pressure of the gas must be limited during concentration by contact with cooled charcoal to avoid violent explosions [1]. Red phosphorus ignites when gently

warmed, and powdered boron and silicon generate sparks on heating in the difluoride [2].

Phosphorus(V) oxide

Streng, A. G., *Chem. Rev.*, 1963, **63**, 611
Ignition occurs spontaneously on contact.

Sulfur tetrafluoride

Oberhammer, H. *et al.*, *Inorg. Chem.*, 1978, **17**, 1435
A mixture of the 2 fluorides, used at low temperature to prepare bis(pentafluoro-sulfur) oxide, is described as possibly explosive.

Water

Streng, A. G., *Chem. Rev.*, 1963, **63**, 610
Presence of water or water vapour in oxygen difluoride is dangerous, the mixture (even when diluted with oxygen) exploding violently on spark ignition, especially at 100°C (i.e. with steam).
See Adsorbents, above
See other ENDOTHERMIC COMPOUNDS, HALOGEN OXIDES, OXIDANTS, OXYGEN FLUORIDES

4318. Sulfinyl fluoride

[7783-42-8]



F₂OS

Sodium

See Sodium: Non-metal halides (reference 8)
See other NON-METAL HALIDES

4319. Xenon difluoride oxide

[13780-64-8]



F₂OXe

Alone, or Mercury, or Fluorides

1. Jacob, E. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1976, **15**, 158–159
2. Gillespie, R. J. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1977, 595–597
Although stable at below –40°C in absence of moisture, it will explode if warmed rapidly (>20°C/h). Explosive decomposition of the solid difluoride oxide at –196°C occurs on contact with mercury, or antimony pentafluoride or arsenic pentafluoride [1]. The fluoride explodes at about 0°C, and also in contact with arsenic pentafluoride in absence of hydrogen fluoride at –78°C [2].
See other NON-METAL HALIDES, XENON COMPOUNDS

4320. Dioxygen difluoride
[7783-44-0]

F₂O₂

O₂F₂

It is mildly endothermic (ΔH_f° (g) +19.8 kJ/mol, 0.28 kJ/g) but a powerful oxidant.

Sulfur trioxide

Solomon, I. J. *et al.*, *J. Chem. Eng. Data*, 1968, **13**, 530

Interaction of the endothermic fluoride with the trioxide is very vigorous, and explosive in absence of solvent.

Various materials

Streng, A. G., *Chem. Rev.*, 1963, **63**, 615

Though not shock-sensitive, it is of limited thermal stability, decomposing below its b.p., -57°C , and explosively in contact with fluorided platinum at -113°C . It is a very powerful oxidant and reacts vigorously or violently with many materials at cryogenic temperatures. It explodes with methane at -194° , with ice at -140° , with solid ethanol at below -130° and with acetone–solid carbon dioxide at -78°C . Ignition or explosion may occur with chlorine, phosphorus trifluoride, sulfur tetrafluoride or tetrafluoroethylene, in the range -130 to -190°C . Even a 2% solution in hydrogen fluoride ignites solid benzene at -78°C .

See other ENDOTHERMIC COMPOUNDS, HALOGEN OXIDES, OXIDANTS, OXYGEN FLUORIDES

4321. Selenium difluoride dioxide
[14984-81-7]

F₂O₂Se

F₂SeO₂

Ammonia

1. Bailar, 1973, Vol. 2, 966

2. Engelbrecht, A. *et al.*, *Monatsh.*, 1962, **92**, 555, 581

Interaction is violent [1], and many of the products and derivatives are both shock- and heat-sensitive explosives [2]. These include the ammonium, potassium, silver and thallium salts of the ‘triselenimidate’ ion, systematically 2,4,6-tris(dioxoselena)perhydrotriazine-1,3,5-triide.

See other NON-METAL HALIDES

4322. Xenon difluoride dioxide
[13875-06-4]

F₂O₂Xe

F₂XeO₂

Kirk Othmer, 1980, Vol. 12, 292

It is explosively unstable.

Preparative hazard

See Xenon hexafluoride: Water (reference 2)

Xenon tetrafluoride oxide: Caesium nitrate

See other XENON COMPOUNDS

4323. 'Trioxxygen difluoride'

[16829-28-0]

F₂O₃

F₂O₃

Bailar, 1973, Vol. 2, 761–763

This material is now considered to be an equimolar mixture of dioxygen difluoride and tetraoxygen difluoride, rather than the title species.

Various materials

Streng, A. G., *Chem. Rev.*, 1963, **63**, 619

Though thermally rather unstable, decomposing above its m.p., -190°C , it appears not to be inherently explosive. It is, however, an extremely potent oxidiser and contact with oxidisable materials causes ignition or explosions, even at -183°C . At this temperature, single drops added to solid hydrazine or liquid methane cause violent explosions, while solid ammonia, bromine, charcoal, iodine, red phosphorus and sulfur react with ignition and/or mild explosion. It is also extremely effective at initiating ignition of combustible materials in liquid oxygen, even at 0.1% concentration, whereas mixtures of ozone and fluorine in liquid oxygen are ineffective. This effect has been examined for use in hypergolic rocket propellant systems. Tetryl detonates spontaneously on contact with the difluoride.

See other HALOGEN OXIDES, OXIDANTS, OXYGEN FLUORIDES

4324. Fluorine fluorosulfate (Fluorosulfuryl hypofluorite)

[13536-85-1]

F₂O₃S

FSO₂OF

1. Cady, G. H., *Chem. Eng. News*, 1966, **44**(8), 40

2. Cady, G. H. *et al.*, *Inorg. Synth.*, 1968, **11**, 155

3. Cady, G. H., *Intra-Sci. Chem. Rep.*, 1971, **5**, 1

4. Lustig, M. *et al.*, *Adv. Fluorine Chem.*, 1973, **7**, 175

The crude fluorosulfate, produced as by-product in preparation of peroxodisulfuryl difluoride, was distilled into a cooled steel cylinder and, on warming to ambient temperature, the cylinder exploded. It decomposes at 200°C , but not explosively [1]. Preparative and handling procedures are detailed [2]. Further warnings on the need to handle with care have been given [3,4].



See other ACYL HYPOHALITES, OXIDANTS See related ACYL HALIDES

4325. Difluorotrioxoxenon
[15192-14-0]



Kirk Othmer, 1980, **Vol. 12**, 292
Explosively unstable.
See other XENON COMPOUNDS

4326. Disulfuryl difluoride
[13036-75-4]



Ethanol

Hayek, E., *Monatsh.*, 1951, **82**, 942
Violent reaction on mixing at ambient temperature.
See other ACYL HALIDES *See related* ACID ANHYDRIDES

4327. Hexaoxygen difluoride
[12191-80-9]



Bailar, 1973, Vol. 2, 764
Flashlight illumination or rapid warming of the solid at -213° to -183°C may lead to explosion.
See other HALOGEN OXIDES, IRRADIATION DECOMPOSITION INCIDENTS, OXYGEN FLUORIDES

4328. Peroxodisulfuryl difluoride
[13709-32-5]



Shreeve, J. M. *et al.*, *Inorg. Synth.*, 1963, **7**, 124
A powerful oxidant which ignites organic materials on contact. Preparative and handling procedures are detailed.

Boron nitride

See Tetra(boron nitride) fluorosulfate

Carbon monoxide

Gatti, R. *et al.*, *Z. Phys. Chem.*, 1965, **47**, 323–336
Interaction proceeds explosively above 20°C .

Dichloromethane

Kirchmeier, R. L. *et al.*, *Inorg. Chem.*, 1973, **12**, 2889

Equimolar amounts explode while warming together to ambient temperature after initial contact at -183°C . Dilution of the dichloromethane with trichlorofluoromethane prevented explosion at -20°C .

See Fluorine fluorosulfate

N-Fluoroiminosulfur tetrafluoride

See *N*-Fluoroiminosulfur tetrafluoride: Alone, etc.

See other ACYL HALIDES, DIACYL PEROXIDES, OXIDANTS

4329. Lead(II) fluoride

[7783-46-2]



F_2Pb

Fluorine

See Fluorine: Metal salts See other METAL HALIDES

4330. Poly(difluorosilylene)

[30582-57-1]



$(\text{F}_2\text{Si})_n$

1. Bailar, 1973, Vol. 1, 1352

2. Perry, D. L. *et al.*, *J. Chem. Educ.*, 1976, **53**, 696–699

Produced by condensation at low temperature, the rubbery polymer ignites in air [1], and preparation, handling and reactions have been detailed [2].

See related HALOSILANES, NON-METAL HALIDES (AND THEIR OXIDES) SEE OTHER PYROPHORIC MATERIALS

4331. Tin(II) fluoride

[7783-47-3]



F_2Sn

Magnesium nitrate

See Magnesium nitrate: Tin(II) fluoride

See other METAL HALIDES, REDUCANTS

4332. Xenon difluoride

[13709-36-9]



F_2Xe

Shackelford, S. A., *J. Org. Chem.*, 1979, **44**, 3490–3492

Though a powerful oxidant, the difluoride is not explosively unstable. Safe procedures for the use of xenon difluoride in fluorination reactions are detailed. Residual traces of the fluoride are rapidly destroyed by dichloromethane at ambient temperatures.

Alkylaluminiums

Bulgakov, R. G., *et al.*, *Chem. Abs.*, 1990, **112**, 7548

Reaction with diethoxyethylaluminium is explosive in the absence of solvent.

See ALKYLALUMINIUM ALKOXIDES AND HYDROXIDES, ALKYL METALS

Combustible materials

Klimov, B. D. *et al.*, *Chem. Abs.*, 1970, **72**, 85784

Xenon difluoride (or the tetrafluoride, or their mixtures) could not be caused to detonate by impact. Xenon difluoride and xenon tetrafluoride both may cause explosion in contact with acetone, aluminium, pentacarbonyliron, styrene, polyethylene, lubricants, paper, sawdust, wool or other combustible materials. Their vigorous reactions with ethanol, potassium iodate or potassium permanganate are not explosive, however.

Dimethyl sulfide

Forster, A. M., *J. Chem. Soc., Dalton Trans.*, 1984, 2827

Interaction in absence of a solvent is explosive at ambient temperature.

Silicon–nitrogen compounds

Gibson, J. A. *et al.*, *Can. J. Chem.*, 1975, **53**, 3050

Interaction of xenon difluoride and dimethylaminotrimethylsilane in presence or absence of solvent became explosive at sub-zero temperatures.

See Xenon tetrafluoride

See other NON-METAL HALIDES, XENON COMPOUNDS

†4333. Trifluorosilane

[13465-71-9]



See other HALOSILANES

4334. Iodine dioxide trifluoride

[25402-50-0]



Organic materials

Engelbrecht, A. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1969, **8**, 769

It ignites in contact with flammable organic materials.

See other HALOGEN OXIDES, IODINE COMPOUNDS, OXIDANTS

4335. Manganese trifluoride

[7783-53-1]



Glass

Mellor, 1942, Vol. 12, 344

When heated in contact it attacks glass violently, silicon tetrafluoride being evolved. It is a powerful fluorinating agent.

See other GLASS INCIDENTS, METAL HALIDES

4336. Nitrogen trifluoride

[7783-54-2]

F₃N

NF₃

Bromotrifluoromethane, Ethylene

Wyatt, J. R., *Chem. Abs.*, 1982, **97**, 75008

The presence of the halocarbon extinguishant significantly reduces the severity of the explosive oxidation of ethylene.

Charcoal

Massonne, J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1966, **5**, 317

Adsorption of nitrogen trifluoride on to activated granular charcoal at -100°C caused an explosion, attributed to the heat of adsorption not being dissipated on the porous solid and causing decomposition to nitrogen and carbon tetrafluoride. No reaction occurs at +100°C in a flow system, but incandescence occurs at 150°C.

Chlorine dioxide

Lawless, 1968, 171

Interaction in the gas phase is explosive.

Diborane

Lawless, 1968, 34-35

No interaction occurred at ambient temperature and at pressures up to 8 bar, but violent explosions occurred at low temperatures in the liquid phase, even in absence of the impurity oxygen difluoride.

Hydrogen-containing materials.

1. Ruff., O., *Z. Angew. Chem.*, 1929, **42**, 807

2. Hoffman, C. J. *et al.*, *Chem. Rev.*, 1962, **62**, 4

Sparking of mixtures with ammonia or hydrogen causes violent explosions, and with steam, feeble ones [1]. Mixtures with ethylene, methane and hydrogen sulfide (also carbon monoxide) explode on sparking [2].

Metals

Richter, R. F. *et al.*, *Chem. Health & Safety*, 1995, **2**(2), 18

It is claimed that in storage nitrogen fluoride can slowly react with metals, including stainless steel, to generate tetrafluorohydrazine.

See Tetrafluorohydrazine, and next below

Tetrafluorohydrazine

MCA Case History No. 683

A crude mixture of the 2 compounds, kept for 3 days in a stainless steel cylinder, exploded violently during valve manipulation.

See other N-HALOGEN COMPOUNDS

4337. Trifluoroamine oxide

[13847-65-9]

F₃NO



Gupta, O. D. *et al.*, *Inorg. Chem.*, 1990, **29**(3), 573

A strong oxidant, mixtures with both organic and inorganic compounds are potentially explosive. It is recommended that the synthesis be not scaled up and be performed with due safety precautions.

See AMINE OXIDES

See other N-HALOGEN COMPOUNDS, N-O COMPOUNDS, OXIDANTS

4338. Trifluorosulfur nitride

[15930-75-3]

F₃NS



Chlorine fluoride

See Chlorine fluoride: Trifluorosulfur nitride

See other NON-METAL HALIDES, NITRIDES, N-S COMPOUNDS

4339. Phosphorus trifluoride

[7783-55-3]

F₃P



Borane

See Borane-phosphorus trifluoride

Dioxygen difluoride

See Dioxygen difluoride: Various materials

Fluorine

See Fluorine: Covalent halides

Hexafluoroisopropylideneaminolithium

See Hexafluoroisopropylideneaminolithium: Non-metal halides

See other HALOPHOSPHINES, NON-METAL HALIDES

4340. Thiophosphoryl fluoride

[2404-52-6]

F₃PS



Air, or Sodium

Mellor, 1940, Vol. 8, 1072–1073

In contact with air, the fluoride ignites or explodes, depending on contact conditions. Heated sodium ignites in the gas.

See related NON-METAL HALIDES

4341. Palladium trifluoride
[13842-82-5]



F₃Pd

Hydrogen

Sidgwick, 1950, 1574

Contact with hydrogen causes the unheated fluoride to be reduced incandescently.

See other METAL HALIDES

4342. Bis-*N* (imidosulfurdifluoridato)mercury
[23303-78-8]



F₄HgN₂S₂

Preparative hazard

Mews, R. *et al.*, *Inorg. Synth.*, 1986, 24, 14–16

By-products formed during the preparation of the title compound from (fluorocarbonyl)iminosulfur difluoride, mercury(II) fluoride and bromine often react violently with water: cleaning of equipment should initially be effected with carbon tetrachloride.

See other MERCURY COMPOUNDS, N–S COMPOUNDS

4343. Manganese tetrafluoride
[15195-58-1]



F₄Mn

Petroleum oil

Sorbe, 1968, 84

Interaction leads to fire.

See other METAL HALIDES, OXIDANTS

4344. Rhenium nitride tetrafluoride
[84159-29-5]



F₄NRe

Preparative hazard

See Trimethylsilyl azide: Rhenium hexafluoride

See other NITRIDES *See related* METAL HALIDES

†4345. Tetrafluorohydrazine
[10036-47-2]

F₄N₂

F₂NNF₂

1. Logothetis, A. L., *J. Org. Chem.*, 1966, **31**, 3686, 3689
2. Modica, A. P. *et al.*, *Rept. No. 357-275*, Princetown Univ., 1963
3. Martin, K. J., *J. Amer. Chem. Soc.*, 1965, **87**, 394
4. Richter, M. F. *et al.*, *Chem. Health & Safety*, 1995, **2**(2), 18

General precautions for use of the explosive gas tetrafluorohydrazine and derived reaction products include: reactions on as small a scale as possible and behind a barricade; adequate shielding during work-up of products because explosions may occur; storage of tetrafluorohydrazine at -80°C under 1–2 bar pressure in previously fluorinated Monel or stainless steel cylinders with Monel valves; distillation of volatile difluoroamino products in presence of an inert halocarbon oil to prevent explosions in dry distilling vessels [1]. Light-initiated explosion of the gas has been reported [2], and it explodes on contact with air and combustible vapours, so careful inerting is essential [3]. Disposal of some aged tetrafluorohydrazine cylinders is described, by puncturing the cylinder with explosives in a pit full of lime[4].

Alkenyl nitrates

Reed, S. F. *et al.*, *J. Org. Chem.*, 1972, **37**, 3329

The products of interaction of tetrafluorohydrazine and alkenyl nitrates, bis(difluoroamino)alkyl nitrates, are heat- and impact-sensitive explosives.

See DIFLUOROAMINO COMPOUNDS

Hydrocarbons

Petry, R. C. *et al.*, *J. Org. Chem.*, 1967, **32**, 4034

Mixtures are potentially highly explosive, approaching the energy of hydrogen–oxygen systems.

Hydrocarbons, Oxygen

See Oxygen: Hydrocarbons, Promoters

Hydrogen

Kuhn, L. P. *et al.*, *Inorg. Chem.*, 1970, **9**, 602

Explosive interaction is rather unpredictable, the initiation temperature required ($20-80^{\circ}\text{C}$) depending on the condition of the vessel wall.

Nitrogen trifluoride

See Nitrogen trifluoride: Tetrafluorohydrazine

Organic materials

Reed, S. F., *J. Org. Chem.*, 1968, **33**, 2634

Mixtures with organic materials in presence of air constitute explosion hazards. Appropriate precautions are essential.

Ozone

Sessa, P. A. *et al.*, *Inorg. Chem.*, 1971, **10**, 2067

When tetrafluorohydrazine was pyrolysed at 310°C to generate NF₂ radicals and the mixture contacted liquid ozone at -196°C, a violent explosion occurred.

See other DIFLUOROAMINO COMPOUNDS, N-HALOGEN COMPOUNDS

4346. Xenon tetrafluoride oxide

[13774-85-1]

F₄OXe

OXeF₄

Preparative hazard

See Xenon hexafluoride: Silicon dioxide

Xenon hexafluoride: Caesium nitrate

Caesium nitrate

Christe, K. O., *Inorg. Chem.*, 1988, **27**, 3763

In the preparation of xenon difluoride dioxide from caesium nitrate and xenon tetrafluoride oxide, the latter must always be used in excess to prevent formation of explosive xenon trioxide.

See Xenon trioxide, below

Graphite, Potassium iodide, Water

Selig, H. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1975, **11**, 75-77

The graphite-xenon tetrafluoride oxide intercalation compound exploded in contact with potassium iodide solution.

Polyacetylene

Selig, H. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1981, 1288

During doping of polyacetylene films, contact with the liquid tetrafluoride led to ignition of the film.

Xenon trioxide

Christe, K. O., *Inorg. Chem.*, 1988, **27**, 3763

Liquid mixtures of the 2 xenon compounds, cooled in liquid nitrogen, tend to flash, then explode after a few seconds.

See other OXIDANTS, XENON COMPOUNDS

4347. Palladium tetrafluoride

[13709-55-2]

F₄Pd

PdF₄

Water

Bailar, 1973, Vol. 3, 1278

Interaction is violent.

See other METAL HALIDES

1535

4348. Platinum tetrafluoride
[13455-15-7]



F₄Pt

Water

Sidgwick, 1950, 1614
Interaction is violent.
See other METAL HALIDES

4349. Rhodium tetrafluoride
[60617-65-4]



F₄Rh

Water

Bailar, 1973, Vol. 3, 1235
Interaction is violent.
See other METAL HALIDES

4350. Sulfur tetrafluoride
[7783-60-0]



F₄S

2-(Hydroxymethyl)furan, Triethylamine

Jenzen, A. F. *et al.*, *J. Fluorine Chem.*, 1988, **38**, 205–208

Reaction of sulfur tetrafluoride with 2-hydroxymethylfuran in presence of triethylamine at -50°C is explosive in absence of a solvent.

See other HALOGENATION INCIDENTS

2-Methyl-3-butyn-2-ol

Boswell, G. A. *et al.*, *Org. React.*, 1974, **21**, 8

Interaction at -78°C is explosively vigorous.

Dioxygen difluoride

See Dioxygen difluoride: Various materials

See other NON-METAL HALIDES

4351. Selenium tetrafluoride
[13465-66-2]



F₄Se

Chlorine trifluoride

Olah, G. A. *et al.*, *J. Amer. Chem. Soc.*, 1974, **96**, 927

The tetrafluoride is prepared by interaction of chlorine trifluoride and selenium in selenium tetrafluoride as solvent. The crude tetrafluoride must be substantially free from excess chlorine trifluoride to avoid danger during subsequent distillation at $106^{\circ}\text{C}/1$ bar.

Water

Aynsley, E. E. *et al.*, *J. Chem. Soc.*, 1952, 1231

Interaction is violent.

See other NON-METAL HALIDES

4352. Silicon tetrafluoride

[7783-61-1]



F₄Si

Lithium nitride

See Lithium nitride: Silicon tetrafluoride

Sodium

See Sodium: Non-metal halides (reference 8)

See other NON-METAL HALIDES

4353. Xenon tetrafluoride

[13709-61-0]



F₄Xe

1. Shieh, J. C. *et al.*, *J. Org. Chem.*, 1970, **35**, 4022

2. Malm, J. G. *et al.*, *Inorg. Synth.*, 1966, **8**, 254

3. Chernick, C. L., *J. Chem. Educ.*, 1966, **43**, 619

4. Holloway, J. H., *Talanta*, 1967, **14**, 871

5. Falconer, E. E. *et al.*, *J. Inorg. Nucl. Chem.*, 1967, **29**, 1380

Moisture converts it to highly shock-sensitive xenon oxides [1]. Precautions necessary for various aspects of its use and application are detailed [2,1,1,5].

See Xenon trioxide

Flammable materials

Klimov, B. D. *et al.*, *Chem. Abs.*, 1970, **72**, 85784

Xenon tetrafluoride (or the difluoride, or their mixtures) could not be caused to detonate by impact. Xenon difluoride and xenon tetrafluoride both may cause explosion in contact with acetone, aluminium, pentacarbonyliron, styrene, polyethylene, lubricants, paper, sawdust, wool or other combustible materials. Their vigorous reactions with ethanol, potassium iodate or potassium permanganate are not explosive, however.

See other NON-METAL HALIDES, OXIDANTS, XENON COMPOUNDS

4354. Pentafluoroorthoselenic acid

[38989-47-8]



F₅HOSe

Preparative hazard

See Fluorine: Hydrogen fluoride, Seleninyl fluoride

See other INORGANIC ACIDS *See related* NON-METAL HALIDES

4355. Iodine pentafluoride
[7783-66-6]

IF₅

F₅I

Benzene

Ruff, O. *et al.*, *Z. Anorg. Chem.*, 1931, **201**, 245
Interaction becomes violent above 50°C.

Calcium carbide, or Potassium hydride

Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 425
Both incandesce on contact, the carbide when warmed.

Diethylaminotrimethylsilane

Oates, G. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1974, 1383
A mixture exploded at around -80°C. Reactions with other silanes were very exothermic.

Dimethyl sulfoxide

Lawless, E. M., *Chem. Eng. News*, 1969, **47**(13), 8, 109
Unmoderated reaction with the sulfoxide is violent, and in the presence of diluents the reaction may be delayed and become explosively violent. Although small-scale reactions were uneventful, reactions involving about 0.15 g mol of the pentafluoride and sulfoxide in presence of trichlorotrifluoromethane or tetrahydrothiophene-1,1-dioxide as diluents caused delayed and violent explosions. Silver difluoride and other fluorinating agents also react violently with the sulfoxide.
See Dimethyl sulfoxide: Acyl halides, etc.
See other INDUCTION PERIOD INCIDENTS

Limonene, Tetrafluoroethylene

See Tetrafluoroethylene: Iodine pentafluoride, etc.

Metals, or Non-metals

1. Sidgwick, 1950, 1159
2. Mellor, 1940, Vol. 2, 114
3. Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 424-425
Contact with boron, silicon, red phosphorus, sulfur, or arsenic, antimony or bismuth usually causes incandescence [1]. Solid potassium or molten sodium explode with the pentafluoride, and aluminium foil ignites on prolonged contact [2]. Molybdenum and tungsten incandesce when warmed [3].

Organic materials, or Potassium hydroxide

Pascal, 1960, Vol. 16.1, 582
The pentafluoride chars and usually ignites organic materials, and interaction with potassium hydroxide is violently exothermic.


Tetraiodoethylene

Mellor, 1956, Vol. 2, Suppl. 1, 176
Rapid mixing leads to explosion.

Water

Ruff, O. *et al.*, *Z. Anorg. Chem.*, 1931, **201**, 245
Reaction with water or water-containing materials is violent.
See other INTERHALOGENS, IODINE COMPOUNDS

4356. Tetrafluoroiodosyl hypofluorite
[72151-31-6]



F_5IO_2

Christe, K. O. *et al.*, *Inorg. Chem.*, 1981, **20**, 2105
Two explosions were encountered in reactions involving the hypofluorite.
See other HALOGEN OXIDES, HYPOHALITES, IODINE COMPOUNDS

4357. N-Fluoroiminosulfur tetrafluoride
[74542-20-4]



F_5NS

Alone, or Peroxydisulfuryl difluoride

O'Brien, B. A. *et al.*, *Inorg. Chem.*, 1984, **23**, 2189
Samples of the *N*-fluoro compound, either alone or in admixture with the difluoride, exploded forcefully while warming from -196°C to $+22^\circ\text{C}$. Sample size should not exceed 3 mmol.
See other *N*-HALOGEN COMPOUNDS, *N-S* COMPOUNDS


4358. Azidopentafluorotungsten
[75900-58-2]



$\text{F}_5\text{N}_3\text{W}$

Fawcett, J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 2294–2296
Trimethylsilyl azide reacts with excess tungsten hexafluoride in solvent to give the title compound, which explodes at 63°C or if dried in dynamic vacuum. A deficiency of the hexafluoride gave a mixture (presumably containing poly-azides) which exploded at ambient temperature.
See other METAL AZIDE HALIDES

4359. Tripotassium hexafluoroferrate(3-)
[13815-30-0]



F_6FeK_3

Wright, S. W., *J. Chem. Educ.*, 1994, **71**(3), 251
The anhydrous salt will undergo a 'thermite' reaction with aluminium to produce molten iron.
See other THERMITE REACTIONS

4360. Hydrogen hexafluorophosphate (Hexafluorophosphoric acid)
[16940-81-1] (anhydrous)

F₆HP

HPF₆

Borosilicate glass

1. Pearse, G. A., *Chem. Brit.*, 1989, **25**, 30

2. Woolf, A. A., *Chem. Brit.*, 1989, **25**, 361

Copper(II) hexafluorophosphate was being prepared by adding 60–65% aqueous acid to aqueous copper carbonate, followed by evaporation of the reaction mixture at 80–90°C by heating on a ceramic hotplate. During evaporation, the Pyrex vessel was dissolved at the liquid level, and the leaking solution also dissolved the ceramic hotplate. This was attributed to presence of hydrofluoric acid, arising from hydrolysis of the fluorophosphoric acid. Teflon-lined vessels are recommended for such operations [1]. Aqueous solutions contain but little of the acid, in hydrolytic equilibrium with mono- and di-fluorophosphoric acids, orthophosphoric acid and hydrofluoric acid, and such a mixture is clearly incompatible with glass containers, and warnings have been published. Anhydrous hexafluorophosphate salts may be prepared in glass by metathesis in anhydrous solvents [2].

See other GLASS INCIDENTS *See other* INORGANIC ACIDS

4361. Azidoiodoiodonium hexafluoroantimonate

[]

F₆I₂N₃Sb

N₃I⁺I SbF₆⁻

Schleyer, P. von R. *et al.*, *Angew. Chem. (Int.)*, 1992, **31**(10), 1338

This compound, which is more sensitive than iodonium azide, can explode spontaneously below –20°C. It is the first known I–N containing cationic species.

See other N-HALOGEN COMPOUNDS, IODINE COMPOUNDS, NON-METAL AZIDES

4362. Iridium hexafluoride

[7783-75-7]

F₆Ir

IrF₆

Silicon

See Silicon: Metal hexafluorides

See other METAL HALIDES, OXIDANTS

4363. Potassium hexafluoromanganate(IV)

[16962-31-5]

F₆K₂Mn

K₂[MnF₆]

Preparative hazard

See Potassium permanganate: Hydrofluoric acid

See related METAL HALIDES

4364. Potassium hexafluorosilicate(2-)
[16871-90-2] **F₆K₂Si**
K₂[SiF₆]

Hydrogen fluoride
See Hydrogen fluoride: Potassium tetrafluorosilicate(2-)
See related NON-METAL HALIDES

4365. Molybdenum hexafluoride
[7783-77-9] **F₆Mo**
MoF₆

Tetramethoxysilane
See Tetramethoxysilane: Metal hexafluorides
See other METAL HALIDES

4366. Neptunium hexafluoride
[14521-05-2] **F₆Np**
NpF₆

Water
Bailar, 1973, Vol. 5, 168
Interaction with water at ambient temperature is violent.
See other METAL HALIDES, OXIDANTS

4367. Pentafluorosulfur hypofluorite
[15179-32-5] **F₆OS**
F₅SOF

Ruff, J. K., *Inorg. Synth.*, 1968, **11**, 137
It is considered to be potentially explosive.
See other HYPOHALITES

4368. Pentafluoroselenium hypofluorite
[27218-12-8] **F₆OSe**
F₅SeOF

1. Mitra, G. *et al.*, *J. Amer. Chem. Soc.*, 1959, **81**, 2646
2. Smith, J. E. *et al.*, *Inorg. Chem.*, 1970, **9**, 1442
A cooled sample exploded when allowed to dry rapidly [1], but this may have been owing to impurities, as it did not happen in the later work [2].
See other HYPOHALITES

4369. Pentafluorotellurium hypofluorite
[83314-21-0] **F₆OTe**



Christe, K. O. *et al.*, *Inorg. Chem.*, 1987, **26**, 920
A shock-sensitive explosive, which needs full handling precautions.
See other HYPOHALITES

4370. Osmium hexafluoride
[13768-38-2] **F₆Os**



Organic materials
Sorbe, 1968, 91
It causes ignition of paraffin oil and other organic materials.

Silicon
See Silicon: Metal hexafluorides
See other METAL HALIDES, OXIDANTS

4371. Platinum hexafluoride
[13693-05-5] **F₆Pt**



Bis(trifluoromethyl) nitroxide
Christe, K. O. *et al.*, *J. Fluorine Chem.*, 1974, **4**, 425
Interaction of the nitroxide radical and this powerful oxidant was very violent during warming from -196°C .
See other METAL HALIDES, OXIDANTS

4372. Plutonium hexafluoride
[13693-06-6] **F₆Pu**



Water
Bailar, 1973, Vol. 5, 168
Interaction with water at ambient temperature is violent.
See other METAL HALIDES, OXIDANTS

4373. Rhenium hexafluoride
[10049-17-9] **F₆Re**



Silicon
See Silicon: Metal hexafluorides

Tetramethoxysilane
See Tetramethoxysilane: Metal hexafluorides

Trimethylsilyl azide
See Trimethylsilyl azide: Rhenium hexafluoride
See other METAL HALIDES, OXIDANTS

4374. Sulfur hexafluoride
[2551-62-4]

SF₆

F₆S

Disilane
See Disilane: Non-metal halides, etc.
See other NON-METAL HALIDES

4375. Uranium hexafluoride
[7783-81-5]

UF₆

F₆U

Aromatic hydrocarbons, or Hydroxy compounds
Sidgwick, 1950, 1072
Interaction with benzene, toluene, or xylene is very vigorous, with separation of carbon, and violent with ethanol or water.
See other METAL HALIDES, OXIDANTS

4376. Tungsten hexafluoride
[7783-82-6]

WF₆

F₆W

Aitchison, K. A. *et al.*, *Chem. Abs.*, 1987, **106**, 147494
Safety procedures for use of tungsten hexafluoride in CVD processing operations are emphasized.

Tetramethoxysilane
See Tetramethoxysilane: Metal hexafluorides

Trimethylsilyl azide
See Azidopentafluorotungsten
See other METAL HALIDES, OXIDANTS

4377. Xenon hexafluoride
[13693-09-9]

F₆Xe

XeF₆

Caesium nitrate

Christe, K. O., *Inorg. Chem.*, 1988, **27**, 3763

In the preparation of xenon tetrafluoride oxide from the hexafluoride and caesium nitrate, the former must always be used in excess to prevent formation of explosive xenon trioxide.

See Xenon tetrafluoride oxide: Xenon trioxide

Fluoride donors, Water

Bailar, 1973, Vol. 1, 317

Adducts of the hexafluoride with sodium fluoride, potassium fluoride, rubidium fluoride, caesium fluoride or nitrosyl fluoride react violently with water.

Hydrogen

Malm, J. G. *et al.*, *J. Amer. Chem. Soc.*, 1963, **85**, 110

Interaction is violent.

Silicon dioxide

1. McKee, D. E. *et al.*, *Inorg. Chem.*, 1973, **12**, 1722

2. Aleinikov, N. N. *et al.*, *J. Chromatog.*, 1974, **89**, 367

Interaction of the yellow hexafluoride with silica to give xenon tetrafluoride oxide must be interrupted before completion (disappearance of colour) to avoid the possibility of formation and detonation of xenon trioxide [1]. An attempt to collect the hexafluoride in fused silica traps at -20°C after separation by preparative gas chromatography failed because of reaction with the silica and subsequent explosion of the oxygen compounds of xenon so produced [2].

See Xenon trioxide

Water

1. Gillespie, R. J. *et al.*, *Inorg. Chem.*, 1974, **13**, 2370–2374

2. Schumacher, G. A. *et al.*, *Inorg. Chem.*, 1984, **23**, 2923

Although uncontrolled reaction of xenon hexafluoride and moisture produces explosive xenon trioxide, controlled action by progressive addition of limited amounts of water vapour with agitation to a frozen solution of the hexafluoride in anhydrous hydrogen fluoride at -196°C to give xenon oxide tetrafluoride or xenon dioxide difluoride is safe [1]. Controlled hydrolysis in solution in hydrogen fluoride is, however, described as hazardous [2].

See *other* NON-METAL HALIDES, OXIDANTS, XENON COMPOUNDS

4378. Iodine heptafluoride
[16921-96-3]

F₇I

IF₇

Carbon, or Combustible gases

Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 428

Activated carbon ignites immediately in the gas, mixtures with methane ignite, and those with carbon monoxide ignite on warming, while those with hydrogen explode on heating or sparking.

Metals

Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 427

Interaction with barium, potassium and sodium is immediate, accompanied by evolution of light and heat. Aluminium, magnesium, and tin are passivated on contact, but on heating react similarly to the former metals.

Organic solvents

Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 428

Benzene, light petroleum, ethanol and ether ignite in contact with the gas, while the exotherm with acetic acid, acetone or ethyl acetate causes rapid boiling. General organic materials (cellulose, grease, oils) ignite if excess heptafluoride is present. *See other* INTERHALOGENS

4379. Potassium heptafluorotantalate(V)

[16924-00-8]



F₇K₂Ta

Water


Ephraim, 1939, 353

Hydrolysis is less violent than that of tantalum pentafluoride.

See related METAL HALIDES

4380. Difluoroammonium hexafluoroantimonate

[56533-31-4]



F₈H₂NSb

See Difluoroammonium hexafluoroarsenate

See related N-HALOGEN COMPOUNDS

4381. Bis(S,S-difluoro-N-sulfinido)sulfur tetrafluoride

[52795-23-0]



F₈N₂S₃

Water

Hofer, R. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1973, **12**, 1000

Either the liquid tetrafluoride or its viscous polymer decomposes explosively in contact with water.

See other NON-METAL HALIDES, N-S COMPOUNDS

4382. Xenon(II) pentafluoroorthoselenate
[38344-58-0]



Oxidisable materials

Seppelt, K., *Angew. Chem. (Intern. Ed.)*, 1972, **11**, 724

Interaction is explosive.

See other OXIDANTS, XENON COMPOUNDS

4383. Xenon(II) pentafluoroorthotellurate
[25005-56-5]




Organic solvents

Sladky, F., *Angew. Chem. (Intern. Ed.)*, 1969, **8**, 523; *Inorg. Synth.*, 1986, **24**, 36

Explosive or very vigorous reactions occur on contact with acetone, benzene or ethanol.

See other OXIDANTS, XENON COMPOUNDS

4384. Tetrafluoroammonium hexafluoromanganate
[74449-37-9]




Water

Christe, K. O. *et al.*, *Inorg. Chem.*, 1980, **19**, 3254

Reaction of the powerful oxidant with water is extremely violent and needs proper precautions. In absence of fuels, it is not shock-sensitive.

See other N-HALOGEN COMPOUNDS, OXIDANTS

4385. Tetrafluoroammonium hexafluoronickelate
[63105-40-8]



Christe, K. O., US Pat. 4 108 965, 1978

This high energy oxidant is useful in propellants.

Water


Christe, K. O., *Inorg. Chem.*, 1977, **16**, 2238

The hydrolysis of this powerful oxidant is very violent, and may be explosive if attempted appreciably above -180°C .

See other N-HALOGEN COMPOUNDS, OXIDANTS

4386. Tetrafluoroammonium octafluoroxenate

[82963-15-3]

F₁₆N₂XeChriste, K. O. *et al.*, US Pat. 4 447 407, 1984

It produces the highest theoretical detonation pressures in explosive formulations (maximal yields of fluorine and nitrogen trifluoride).

See other N-HALOGEN COMPOUNDS, OXIDANTS, XENON COMPOUNDS**4387. Octakis(trifluorophosphine)dirhodium**

[14876-96-1]

F₂₄P₈Rh₂

Acetylenic esters

Bennett, M. A. *et al.*, *Inorg. Chem.*, 1976, **15**, 107–108

Formation of complexes with excess methyl propiolate or dimethyl acetylenedicarboxylate must not be allowed to proceed at above +20°C, or violently explosive polymerisation of the acetylene esters will occur.

See other POLYMERISATION INCIDENTS**4388. Iron**

[7439-89-6]

Fe**Fe**

1. Bailar, 1973, Vol. 3, 985

2. Gusein, M. A. *et al.*, *Chem. Abs.*, 1974, **81**, 1247743. Jakusch, H. *et al.*, *Chem. Abs.*, 1983, **98**, 1304714. Brinza, V. N. *et al.*, *Chem. Abs.*, 1983, **99**, 78725. Bandyopadhyay, A. *et al.*, *Steel India*, 1987, **10**, 58–636. Ganguly, A. *et al.*, *Tool Alloy Steels*, 1988, **22**(1), 19–26

The known pyrophoric [1] and explosive properties of ultrafine iron powder were examined in detail [2]. Pyrophoric iron particles may be stabilised by heat treatment with oxygen-containing gases in 2 stages at 25–45°, then 50–70°C [3]. The ignition temperature of sponge iron (183–203°C) is independent of heating rate and coincides with the dissociation temperature of iron hydroxides formed on the surface [4]. An incident involving spontaneous combustion of hydrogen-reduced sponge iron (Direct Reduced Iron, DRI) is analysed. The surface area and porosity of coal-reduced DRI prepared at over 1000°C is lower, and the product, a more dense, sintered DRI has greater oxidation resistance. Fire control and salvage techniques are discussed [5]. Other aspects of storing, handling and shipping of sponge iron are reviewed [6].

See entry HIGH SURFACE-AREA SOLIDS *See other* PYROPHORIC METALS

Air, Oil

Glaser, A., *Arbeitschutz*, 1941, 134–135

Oxidative heating of oily iron dust in a collecting vessel caused vaporisation of oil and subsequent ignition, causing an explosion.

See other SELF-HEATING AND IGNITION INCIDENTS

Air, Water

1. Brimelow, H. C., private comm., 1972

2. Unpublished observations, 1949

2-Nitrophenylpyruvic acid was reduced to oxindole using iron pin-dust–ferrous sulfate in water. The iron oxide–iron residues, after filtering and washing with chloroform, rapidly heated in contact with air and shattered the Buchner funnel [1]. Previously, rapid heating effects had been observed on sucking air through the iron oxide residues from hot filtration of aqueous liquor from reduction of a nitro compound with reduced iron powder [2].

See Water, below

Acetaldehyde

See Acetaldehyde: Metals

Chloric acid

See Chloric acid: Metals, etc.

Chloroformamidinium nitrate

See Chloroformamidinium nitrate: Alone, etc.

Halogens or Interhalogens

MRH values show % of halogen

See Bromine pentafluoride: Acids, etc.

MRH 4.23/54

Chlorine: Metals

MRH 2.68/56

Chlorine trifluoride: Metals

Fluorine: Metals

MRH 7.49/40

Other reactants

Yoshida, 1980, 233

MRH values calculated for 19 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Halogens or Interhalogens above

Ammonium nitrate: Metals

MRH 3.35/68

Ammonium peroxodisulfate: Iron

MRH 0.92/78

Dinitrogen tetraoxide: Metals

MRH 4.43/38

Hydrogen peroxide: Metals

MRH 4.60/48

Nitryl fluoride: Metals

Peroxyformic acid

MRH 5.69/99+

Potassium dichromate: Iron

MRH 0.42/66

Potassium perchlorate: Metal powders

Sodium peroxide: Metals

MRH 1.55/68

Polystyrene

Unpublished observation, 1971

Iron flake powder and polystyrene beads had been blended in a high-speed mixer. The mixture ignited and burned rapidly when discharged into a polythene bag. Rapid oxidation of the finely divided metal and/or static discharge may have initiated the fire. No ignition occurred when the iron powder was surface coated with stearic acid.

See other STATIC INITIATION INCIDENTS

Sodium acetylide

See Sodium acetylide: Metals

Water

1. Asakura, S. *et al.*, *Chem. Abs.*, 1983, **99**, 110033
2. Anon, *Safety Digest Univ. Safety Assoc.*, 1991, **41**, 10
3. Clancey, V. J., *Proc. 41st Ironmaking Conf.*, 1982, 244–250

In a study of the spontaneous exothermic reaction of iron powder or turnings with water in absence of chlorides, rust was found to catalyse the reaction [1]. On drilling holes into steel tubes filled with possibly wet ferrous scrap and sealed some years, they were found to be pressurised. The third tube produced an explosion and jet of flame causing some injury to the driller. This was probably attributable to hydrogen generation [2]. Hazards associated with shipping of direct-reduced iron and arising from reactivity with condensation etc. in ship's holds, with liberation and ignition of hydrogen, are discussed [3].

See Air, Water, above

See other CORROSION INCIDENTS, GAS EVOLUTION INCIDENTS *See other* METALS

4389. Ferromanganese (Iron–manganese alloy)

[73202-12-7]

Fe–Mn

Fe–Mn

See entry FERROALLOY POWDERS *See other* ALLOYS

4390. Ferrosilicon (Iron–silicon alloy)

[50645-52-8]

Fe–Si

Fe–Si

1. Delyan, V. I. *et al.*, *Chem. Abs.*, 1983, **99**, 25024
2. Babaitsev, I. V. *et al.*, *Chem. Abs.*, 1987, **107**, 241804

The flammability and explosive hazard of ferrosilicon powder is increased substantially during grinding in a vibratory mill [1]. Explosion hazards from air–hydrogen, —acetylene, or —propane mixtures formed during preparation of ferrosilicon containing alkaline earth additives are attributed to contact of barium or magnesium carbide or silicide additive with atmospheric moisture [2].

See FERROALLOY POWDERS

Sodium hydroxide, Water

Kirk-Othmer (3rd edn.), 1982, Vol. 19, 494

Intimate mixtures of ferrosilicon with solid sodium hydroxide incandesce when moistened.

Water

1. Anon., *Chem. Trade J.*, 1956, **139**, 1180
2. Horn, Q.C. *et al.*, *Chem. Abs.* 1998, **128**, 37646c; 37653c

Ferrosilicon containing 30–75% of silicon is hazardous, particularly when finely divided, and must be kept in a moisture-tight drum. In contact with water, the impurities present (arsenide, carbide, phosphide) evolve extremely poisonous arsine, combustible acetylene and spontaneously flammable phosphine [1 & 2].

See related METAL NON-METALLIDES

4391. Ferrotitanium (Iron–titanium alloy)

[87490-22-0]

Fe–Ti

Fe–Ti

See entry FERROALLOY POWDERS *See other* ALLOYS

4392. Iron(II) hydroxide

[18624-44-7]

Fe(OH)₂

FeH₂O₂

Gibson, 1969, 121

Prepared under nitrogen, it is pyrophoric in air, producing sparks.

See other PYROPHORIC MATERIALS, REDUCANTS

4393. Ammonium iron(III) sulfate

[7783-83-7]

NH₄Fe(SO₄)₂·12H₂O

FeH₄NO₈S₂·12H₂O

Sulfuric acid

See Sulfuric acid: Ammonium iron(III) sulfate

See related METAL OXONON-METALLATES

4394. Iron(III) phosphinate

[7783-84-8]

Fe[OP(H)OH]₃

FeH₆O₆P₃

Mellor, 1981, Vol. 8, Suppl. 3, 626

It is used in impact-sensitive priming compositions.

See other METAL PHOSPHINATES, REDOX COMPOUNDS

4395. Iron(II) iodide
[7783-86-0]




Alkali metals

See Potassium: Metal halides

Sodium: Metal halides

See other METAL HALIDES

4396. Potassium peroxoferrate(2-)
[]



Alone, or Non-metals, or Sulfuric acid

Goralevich, D. K., *J. Russ. Phys. Chem. Soc.*, 1926, **58**, 1115

It explodes on heating or impact, or in contact with charcoal, phosphorus, sulfur or sulfuric acid.

See other PEROXOACID SALTS

4397. Iron(III) nitrate
[10421-42-4]



Dimethyl sulfoxide

See Dimethyl sulfoxide: Metal oxosalts

See other METAL NITRATES

4398. Iron(II) oxide
[1345-25-1]



1. Mellor, 1941, Vol. 13, 715

2. Bailar, 1973, Vol. 3, 1008–1009

The oxide (prepared at 300°C) burns in air above 200°C, while the finely divided oxide prepared by reduction may be pyrophoric at ambient temperature [1]. That prepared by thermal decomposition under vacuum of iron(II) oxalate is also pyrophoric [2].

Oxidants

See Nitric acid: Iron(II) oxide

Hydrogen peroxide: Metals, etc.

Sulfur dioxide

Mellor, 1941, Vol. 13, 715

The oxide incandescens when heated in sulfur dioxide.
See other METAL OXIDES, REDUCANTS

4399. Iron(II) sulfate
[7720-78-7]



HCS 1980, 502

Arsenic trioxide, Sodium nitrate
See Sodium nitrate: Arsenic trioxide, Iron(II) sulfate

Methyl isocyanoacetate
See Methyl isocyanoacetate: Heavy metals
See other METAL OXONON-METALLATES, REDUCANTS

4400. Iron(II) sulfide
[1317-37-9]



1. Mellor, 1942, Vol. 14, 157
2. Anon., *Chem. Age.*, 1939, **40**, 267
3. Davie, F. M. *et al.*, *J. Loss Prev.*, 1993, **6**(3), 139 & 145
4. Walker, R. *et al.*, *Ind. Eng. Chem. Res.*, 1996, **35**(5), 1736

The moist sulfide readily oxidises in air exothermally, and may reach incandescence. Grinding in a mortar hastens this [1]. The impure sulfide formed when steel processing equipment is used with materials containing hydrogen sulfide or volatile sulfur compounds is pyrophoric, and has caused many fires and explosions when such equipment is opened without effective purging. Various methods of purging are discussed [2]. Formation of pyrophoric FeS in bitumen tanks is considered as a cause of spontaneous ignition and explosion in the head space [3]. A detailed study of formation of possibly pyrophoric sulphides from rust in crude oil tankers has been made [4].

Lithium
See Lithium: Metal oxides, etc.
See other IGNITION SOURCES, METAL SULFIDES

4401. Iron disulfide (Iron pyrites)
[1309-36-0]



1. Anon., *Angew. Chem. (Nachr.)*, 1954, **2**, 219
2. Bowes, P. C., *Ind. Chemist*, 1954, **30**, 12–14

3. Ruiss, I. G. *et al.*, *J. Chem. Ind. (Moscow)*, 1935, **12**, 692–696

4. Guedes de Carvalho, R. A. *et al.*, *Chem. Abs.*, 1980, **93**, 223677

Finely powdered pyrites, especially in presence of moisture, will rapidly heat spontaneously and ignite, particularly in contact with combustible materials [1]. Inert gas blanketing will prevent this [2]. Precautions to reduce the self-ignition hazards of powdered pyrites, and the explosion hazards of pyrites–air mixtures in the furnaces of sulfuric acid plants have been detailed and discussed [3]. Further studies on minimum moisture content of Portuguese pyrites for safe transportation and storage are reported [4].

See other SELF-HEATING AND IGNITION INCIDENTS

Carbon

Ruiss, I. G. *et al.*, *J. Chem. Ind. (Moscow)*, 1935, **12**, 696

The presence of carbon in pyrites lowers the ignition temperature to 228–42°C and increases the explosivity of dust suspensions in air.

See other METAL SULFIDES

4402. Di- μ -iodotetranitrosyldiiron

[15002-08-1]

$\text{Fe}_2\text{I}_2\text{N}_4\text{O}_4$



Kegzdins, P. *et al.* US Pat. 5,631,284, 1997; *Chem. Abs.*, 1997, **127**, 50793x

This compound may decompose explosively during sublimation, if overheated.

4403. Iron(III) oxide

[1309-37-1]

Fe_2O_3



Aluminium

1. Mellor, 1946, Vol. 5, 217

2. Bodur, G. M., *J. Chem. Educ.*, 1985, **62**, 1107

An intimately powdered mixture, usually ignited by magnesium ribbon as a high-temperature fuse, reacts with an intense exotherm to produce molten iron and was used formerly (before the advent of gas or arc welding) in the commercial ‘thermite’ welding process. Incendive particles have been produced by this reaction on impact between aluminium and rusty iron. (The term ‘thermite reaction’ has now been extended to include many combinations of reducing metals and metal oxides) [1]. Some accidents in demonstrating the thermite reaction are described [2].

See Calcium disilicide, below

See Aluminium: Metal oxides, etc. *also* Magnesium: Metal oxides

See also LIGHT ALLOYS

See other THERMITE REACTIONS

Aluminium, Propene

Batty, G. F., private comm., 1972

Use of a rusty iron tool on an aluminium compressor piston caused incendive sparks which ignited residual propene–air mixture in the cylinder.

See Aluminium, above

Aluminium–magnesium alloy, Water

See Aluminium–magnesium alloy: Iron(III) oxide, etc.

Aluminium–magnesium–zinc alloys

See Aluminium–magnesium–zinc alloy: Rusted steel

Calcium disilicide

Berger, E., *Compt. rend.*, 1920, 170, 29

The mixture ('silicon thermite') attains a very high temperature when heated, producing molten iron like the usual thermite mixture.

See Aluminium, above

Carbon monoxide

Othen, C. W., *School, Sci. Rev.*, 1964, **45**(156), 459

The reason for a previously reported explosion during reduction of iron oxide with carbon monoxide is given as the formation of pentacarbonyliron at temperatures between 0 and 150°C. Suitable heating arrangements and precautions will eliminate this hazard.

See Pentacarbonyliron

Ethylene oxide

See Ethylene oxide: Contaminants

Guanidinium perchlorate

See Guanidinium perchlorate: Iron(III) oxide

Hydrogen peroxide

See Hydrogen peroxide: Metals, etc.

Magnesium

See Magnesium: Metal oxides

Metal acetylides

See Calcium acetylide: Iron(III) chloride, etc.

Caesium acetylide: Iron(III) oxide

Rubidium acetylide: Metal oxides

See other METAL OXIDES

4404. Iron(III) sulfide

[12063-27-3]



Moore, F. M., *Proc. Gas Cond. Conf.*, 1976, **26**, 1

Hydrogen sulfide is removed from natural gas by passage over iron sponge, when flammable iron sulfide is produced. Handling precautions during regeneration of the reactor beds are detailed.

See other METAL SULFIDES, PYROPHORIC IRON-SULFUR COMPOUNDS

4405. Iron(II,III) oxide ('Magnetite')

[1317-61-9]

Fe₃O₄

FeO.Fe₂O₃

Aluminium, Calcium silicide, Sodium nitrate.

Schierwater, F-W., *Sichere Chemiearb.*, 1976, **28**, 30-31

During the preparation of a foundry mixture of the finely divided oxide with aluminium powder and small amounts of sodium nitrate and calcium silicide with calcium fluoride as flux, a violent explosion occurred in the conical mixer. It had been established previously that the mixture could not be ignited by impact or friction and that if ignited by a very high-energy source (magnesium ribbon), it burned rather slowly to a glowing liquid. The possibility of ignition of the solid mixture by silane produced from water acting on the silicide content was discounted, but an aluminium dust explosion may have been involved with this ignition source.

Aluminium, Sulfur

Crozier, T. H., *HM Insp. Expl. Spec. Rept. 237*, London, HMSO, 1929

A 20 t quantity of an incendiary bomb mixture of the finely powdered oxide, aluminium and sulfur became accidentally ignited and burned with almost explosive violence. It is similar to thermite mixture.

See other THERMITE REACTIONS

Hydrogen trisulfide

See Hydrogen trisulfide: Metal oxides

See other METAL OXIDES

4406. Gallium

[7440-55-3]

Ga

Ga

Aluminium alloys

Marshall, C., private comm., 1981

Serious problems may arise if gallium or its liquid alloys contact aluminium alloy structural components in aircraft, when rapid 'amalgamation' and weakening occurs.

See other CORROSION INCIDENTS

Halogens

1. Walker, H. L., *School Sci. Rev.*, 1956, **37**(132), 196

2. Bailar, 1973, Vol. 1, 1084

The metal reacts with cold chlorine strongly exothermically, and the compact metal with bromine even at -33°C , reaction being violent at ambient temperature [1]. Interaction of gallium with liquid bromine at 0°C proceeds with a flash, resembling the action of alkali metals with water [2].

Hydrogen peroxide

See Hydrogen peroxide: Gallium, etc.

See other METALS

4407. Lithium tetrahydrogallate

[17836-90-7]

GaH₄Li

Li[H₄Ga]

Gaylord, 1956, 26

Though of lower stability than the analogous aluminate, its reactivity is generally similar to that of the latter.

See other COMPLEX HYDRIDES

4408. Sodium tetrahydrogallate

[32106-51-7]

GaH₄Na

Na[H₄Ga]

Water

McKay, 1966, 169

It is explosively hydrolysed by water.

See other COMPLEX HYDRIDES

4409. Gallium azide (Triazidogallane)

[]

GaN₉

Ga(N₃)₃

Fischer, R. A. *et al.*, *Chem. Eur. J.*, 1996, **2**(11), 1353

The solid detonates violently when heated rapidly to above 280°C .

See other METAL AZIDES

4410. Digallane

[12140-58-8]

Ga₂H₆

H₃GaGaH₃

Leleu, *Cahiers*, 1976, (85), 585

According to some authors it ignites in air.

See other METAL HYDRIDES

4411. Gallium(I) oxide
[12024-20-3]

Ga₂O

Ga₂O

Bromine

Bailar, 1973, Vol. 1, 1091

It is a strong reductant, reacting violently with bromine.

See other METAL OXIDES, REDUCANTS

4412. Germanium
[7440-56-4]

Ge

Ge

Halogens

Mellor, 1941, Vol. 7, 260

The powdered metal ignites in chlorine, and lumps will ignite on heating in chlorine or bromine.

Oxidants

Mellor, 1941, Vol. 7, 260–261

The powdered metal reacts violently with nitric acid, and mixtures with potassium chlorate or nitrate explode on heating. Heated germanium burns with incandescence in oxygen.

See Potassium hydroxide: Germanium

Sodium peroxide: Metals (reference 2)

See other METALS

4413. Poly(germanium monohydride)
[13572-99-1]

(GeH)_n

(GeH)_n

Jolly, W. L. *et al.*, *Inorg. Synth.*, 1963, **7**, 39

The solid polymeric hydride sometimes decomposes explosively into its elements on exposure to air.

See other METAL HYDRIDES

4414. Germanium imide
[26257-00-1]

Ge:NH

GeHN

Air, or Oxygen

Johnson, O. H., *Chem. Rev.*, 1952, **51**, 449

On exposure to air it reacts violently, and in oxygen incandescence occurs.

See other N-METAL DERIVATIVES

4415. Poly(germanium dihydride)

[32028-94-7]



Bailar, 1973, Vol. 2, 13

Impact may cause explosive decomposition to the elements, with ignition of the liberated hydrogen.

See other METAL HYDRIDES

4416. Azidogermane

[21138-22-7]



Fluorosilane

Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 272

An attempt to prepare azidosilane by interaction of azidogermane and fluorosilane exploded.

See related METAL AZIDES, METAL HYDRIDES

†**4417. Germane**

[7782-65-2]



ΔH_f° 90 kJ/mol

1. Brauer, 1963, Vol. 1, 715

2. *Chem. Abs.*, 1994, **120**, 80876

Germane and its higher homologues decompose in air, often igniting [1]. The Japanese have studied the explosive decomposition of germane, the minimum pressure for this was a little more than 0.1 bar [2].

Bromine

See Bromine: Germane

See other METAL HYDRIDES

4418. Sodium germanide

[12265-93-9]



Air, or Water

Johnson, O. H., *Chem. Rev.*, 1952, **51**, 452

The binary alloy is pyrophoric and may ignite in contact with water, as do other alkali metal germanides.

See other ALLOYS, PYROPHORIC MATERIALS

4419. Germanium(II) sulfide
[12025-32-0]

GeS

GeS

Potassium nitrate

See Potassium nitrate: Metal sulfides

See other METAL SULFIDES

†**4420. Digermane**
[13819-89-8]

H₃GeGeH₃

Ge₂H₆

1. Mellor, 1941, Vol. 7, 264
2. Brauer, 1963, Vol. 1, 715
3. McKay, K. M., *Inorg. Synth.*, 1974, **15**, 170

It may ignite in air [1], particularly if air is admitted suddenly into the gas at reduced pressure [2]. Although digermane and its homologues do not usually ignite on exposure to air, their autoignition temperatures appear to be about 50°C and combustion is rapid or explosive [3].

See other METAL HYDRIDES

4421. Trigermane
[14691-44-2]

H₃GeGeH₂GeH₃

Ge₃H₈

Brauer, 1963, Vol. 1, 715
Air-sensitive, it may ignite.
See other METAL HYDRIDES

4422. Poly(dimercuryimmonium hydroxide) ('Millon's base anhydride')
[12529-66-7]

(Hg:N⁺:Hg OH⁻)_n

(HHg₂NO)_n

Sidgwick, 1950, 318

The anhydride of Millon's base explodes if touched or heated to 130°C.

See Mercury: Ammonia

See other FULMINATING METALS, MERCURY COMPOUNDS, POLY(DIMERCURYIMMONIUM) COMPOUNDS, N-METAL DERIVATIVES

4423. Hydriodic acid
[10034-85-2]

HI

HI

Hydrogen iodide is mildly endothermic (ΔH_f° (g) +25.9 kJ/mol, 0.20 kJ/g).

1559

Muir, G. D., private comm., 1968

During preparation of hydriodic acid by distillation of phosphorus and wet iodine, the condenser became blocked with by-product phosphonium iodide, and an explosion, possibly also involving phosphine, occurred. There is also a purification hazard.

See Phosphorus: Hydriodic acid

Metals

See Magnesium: Hydrogen halides

Potassium: Hydrogen halides

Oxidants

Leleu, *Cahiers*, 1974, (75), 271

Hydrogen iodide ignites in contact with fluorine, dinitrogen trioxide, dinitrogen tetraoxide and fuming nitric acid.

See Ethyl hydroperoxide: Hydriodic acid

Perchloric acid: Iodides

Potassium chlorate: Hydrogen iodide

Phosphorus

See Phosphorus: Hydriodic acid

See other INORGANIC ACIDS, NON-METAL HALIDES, NON-METAL HYDRIDES

4424. Iodic acid

[7782-68-5]



Non-metals

1. Mellor, 1946, Vol. 5, 15

2. Partington, 1967, 813

Interaction with boron below 40°C is vigorous, attaining incandescence [1]. Charcoal, phosphorus and sulfur deflagrate on heating [2].

See other INORGANIC ACIDS, OXIDANTS, OXOHALOGEN ACIDS, IODINE COMPOUNDS

4425. Periodic acid

[13444-71-8]



Dimethyl sulfoxide

Rowe, J. J. M. *et al.*, *J. Amer. Chem. Soc.*, 1968, **90**, 1924

Although 1.5 M solutions of periodic acid in dimethyl sulfoxide explode after a few min, 0.15 M solutions appear stable.

See Dimethyl sulfoxide: Metal oxosalts, or: Perchloric acid

Tetraethylammonium hydroxide

1. Cookson, R. C. *et al.*, *Chem. Brit.*, 1979, **15**, 329
2. Kirby, G. W. *et al.*, *J. Chem. Soc., Perkin Trans.*, 1981, 3252

During the preparation of the quaternary oxidant tetraethylammonium iodate from the aqueous reagents, the residue after vacuum evaporation of most of the water exploded, breaking the flask. This was attributed to possible presence of excess periodic acid in the reaction mixture [1]. Further details and precautions to avoid heating the salt, normally stable in storage, are given [2].

See QUATERNARY OXIDANTS

See other INORGANIC ACIDS, OXIDANTS, OXOHALOGEN ACIDS, IODINE COMPOUNDS

4426. Diiodamine (Iodimide)

[15587-44-7]

HI₂N

I₂NH

Mellor, 1940, Vol. 8, 607

Explosive, formed on prolonged contact of nitrogen triiodide with water.

See Nitrogen triiodide–ammonia *See other* N-HALOGEN COMPOUNDS

4427. Potassium hydride

[7693-26-7]

HK

KH

1. Sorbe, 1968, 67

2. Brown, C. A., *J. Org. Chem.*, 1974, **39**, 3913–3918

It ignites on exposure to air [1], and the hydride dispersed in oil is much more highly reactive than sodium hydride dispersions, and rather more careful handling is necessary for safe working. Such precautions are detailed. Contact with water of even traces of the dispersion in flammable solvents will lead to ignition [2].

See Potassium hexahydroaluminate

Bromopentaborane(9), Dimethyl ether

Anon., *Lab. Accid. Higher Educ.*, Item 11, HSE, Barking, 1987

During the reaction of bromopentaborane and potassium hydride in dimethyl ether at -78°C , the reaction became uncontrollable, shattering the glassware and igniting. Cause may have been contamination or effect of scale-up.

O-2,4-Dinitrophenylhydroxylamine

See O-2,4-Dinitrophenylhydroxylamine: Potassium hydride

Fluoroalkene (unspecified)

MCA Case History No. 2134

After a few minutes' reflux at 12°C , a mixture of the hydride (0.01 mol) and a fluoroalkene (0.02 mol) exploded violently. This was attributed to possible presence of metallic potassium in the hydride, causing polymerisation or formation of a fluoroacetylene.

Oxidants

See Fluorine: Metal hydrides

Oxygen (Gas): Metal hydrides

See other METAL HYDRIDES

4428. Potassium hydroxide

[1310-58-3]

HKO

KOH

HCS 1980, 768

Acids

MCA Case History No. 920

Incautious addition of acetic acid to a vessel contaminated with potassium hydroxide caused eruption of the acid.

See other NEUTRALISATION INCIDENTS

Ammonium hexachloroplatinate

See Ammonium hexachloroplatinate: Potassium hydroxide

1,4-Bis(1,2-dibromoethyl)benzene

See 1,4-Bis(1,2-dibromoethyl)benzene: Potassium hydroxide

Bromoform, Cyclic polyethylene oxides

See Bromoform: Cyclic polyethylene oxides, etc.

2-Bromo-2,5,5-trimethylcyclopentanone, Triethylamine

See 2-Bromo-2,5,5-trimethylcyclopentanone: Potassium hydroxide, etc.

Chlorine dioxide

MRH 1.88/55

See Chlorine dioxide: Potassium hydroxide

Cyclopentadiene

See Cyclopentadiene: Potassium hydroxide

Diamond

See Carbon: Potassium hydroxide

Germanium

Partington, 1967, 181

Germanium is oxidised by the fused hydroxide with incandescence.

Glass

Lloyd, A., *Chem. Brit.*, 1987, **23**, 208

During the distillation of hexane from potassium hydroxide pellets, it was found that the 2 l flask had become perforated by alkaline attack, and that hexane was leaking through two pinholes, generating much flammable vapour. The high rate

HKO

of corrosion of glass is probably associated with the fact that fresh pellets of potassium hydroxide already contain some 14% of water, and absorption of further water from solvents during drying over potash pellets leads to formation of drops of very concentrated aqueous potassium hydroxide solution.

See other CORROSION INCIDENTS, GLASS INCIDENTS

Glycols

There are no reports of mishap, but:

See Sodium hydroxide: Glycols

Hyponitrous acid

See Hyponitrous acid: Alone, or Potassium hydroxide

Maleic anhydride

MRH 1.67/99+

See Maleic anhydride: Bases, etc.

Nitroalkanes

MRH Nitromethane 5.69/99+, nitroethane 4.52/99+

See Nitromethane: Acids, etc.

See NITROALKANES: inorganic bases

Nitroaryl compounds

Solid mixtures may deflagrate readily.

See 4-Chloronitrobenzene: Potassium hydroxide

Nitrobenzene: Alkali

4-Nitrobenzoic acid: Potassium hydroxide

2-Nitrophenol: Potassium hydroxide

4-Nitrophenol: Potassium hydroxide

See DEFLAGRATION INCIDENTS

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Other reactants

Yoshida, 1980, 212

MRH values calculated for 10 combinations, largely with catalytically decomposed materials, are given.

Potassium peroxodisulfate

See Potassium peroxodisulfate: Potassium hydroxide

Sugars

See SUGARS

2,2,3,3-Tetrafluoropropanol

See 2,2,3,3-Tetrafluoropropanol: Potassium hydroxide

Tetrahydrofuran

See Tetrahydrofuran: Caustic alkalies

Thorium dicarbide

See Thorium dicarbide: Non-metals, etc.

2,4,6-Trinitrotoluene

See 2,4,6-Trinitrotoluene: Added impurities

Water

Anon., *CISHC Chem. Safety Summ.*, 1976, **46**, 8–9

A mixture of flake potassium hydroxide and sodium hydroxide was added to a reaction mixture without the agitator running. When this was started the batch erupted, owing to the sudden solution exotherm. Although this is a physical hazard rather than a chemical hazard, similar incidents have occurred frequently.


See Sodium hydroxide: Water.

See other AGITATION INCIDENTS

See other INORGANIC BASES

4429. Potassium hydrogen xenate

[[73378-54-8], sesquihydrate]




HKO₄Xe

See other XENON COMPOUNDS

4430. Potassium hydrogen peroxomonosulfate

[10058-23-8]



HKO₅S

Alone, or Organic matter

Castrantas, 1965, 5

It melts with decomposition at 100°C, and forms explosive mixtures with as little as 1% of organic matter.

See other PEROXOACID SALTS

4431. Dipotassium phosphinate

[13492-26-7]



HK₂O₂P

Water

Mellor, 1971, Vol. 8, Suppl. 3, 623

The salt ignites in contact with a little water.

See other METAL PHOSPHINATES, REDUCANTS

4432. Lithium hydride
[60380-67-8]

LiH

HLi

1. Bailar, 1973, Vol. 1, 344
2. Brinza, V. A. *et al.*, *Chem. Abs.*, 1979, **91**, 96308

The powdered material burns readily on exposure to air [1], and the rate of combustion of air suspensions has been studied with respect to concentrations of hydride and moisture in the air [2].

Dinitrogen oxide

See Dinitrogen oxide: Lithium hydride

Oxygen

See Oxygen (Gas): Metal hydrides

Oxygen (Liquid): Lithium hydride

See other METAL HYDRIDES

4433. Manganese diazide hydroxide

[]

(N₃)₂MnOH

HMnN₆O

Mellor, 1940, Vol. 8, 354

It explodes at 203°C.

See related METAL AZIDES

4434. Permanganic acid

[13465-41-3]

HOMnO₃

HMnO₄

Organic materials

1. Frigerio, N. A., *J. Amer. Chem. Soc.*, 1969, **91**, 6201
2. von Schwartz, 1918, 327

The crystalline acid and its dihydrate are very unstable, often exploding at about 3° and 18°C, respectively, but they may be stored virtually unchanged at -75°C. The anhydrous acid ignited explosively every organic compound with which it came into contact except mono-, di- or tri-chloromethanes [1]. The solution of permanganic acid (or its explosive anhydride, dimanganese heptoxide) produced by interaction of permanganates and sulfuric acid, will explode on contact with benzene, carbon disulfide, diethyl ether, ethanol, flammable gases, petroleum or other organic substances [2].

See Dioxonium hexamanganato(VII)manganate

See other INORGANIC ACIDS, OXIDANTS

4435. Nitrous acid
[7782-77-6]

HNO₂

HON:O

2-Amino-5-nitrophenol

See 2-Amino-5-nitrophenol: Nitrous acid

Ammonium decahydrodecaborate(2-)

See Ammonium decahydrodecaborate(2-): Nitrous acid

Anilines

Diazonium compounds or solutions produced by action of nitrous acid on the anilines below have been involved in various incidents.

See Aniline, 4-Bromoaniline, 2-Chloroaniline, 3-Chloroaniline, 2-Nitroaniline, 3-Nitroaniline, 4-Nitroaniline.

A semicarbazone, Silver nitrate

Mitchell, J. J., *Chem. Eng. News*, 1956, **34**, 4704

Use of nitrous acid to liberate a free keto-acid from its semicarbazone caused formation of hydrogen azide which was co-extracted into ether with the product. Addition of silver nitrate to precipitate the silver salt of the acid also precipitated silver azide, which later exploded on scraping from a sintered disc. The possibility of formation of free hydrogen azide from interaction of nitrous acid and hydrazine or hydroxylamine derivatives is stressed.

Phosphine

See Phosphine (reference 3)

Phosphorus trichloride

Mellor, 1940, Vol. 8, 1004

The trichloride explodes with nitrous (or nitric) acid.

See *other* INORGANIC ACIDS, OXIDANTS

4436. Nitric acid
[7697-37-2]

HNO₃

HONO₂

(*MCA SD-5*, 1961); *FPA H23*, 1974; *HCS 1980*, 678 (<70%), 679 (>70%); *RSC Lab. Hazards Data Sheet No. 32*, 1985

1. *MCA SD-5*, 1961
2. Bretherick, L., *CISHC Chem. Safety Summ.*, 1978, **49**(197), 3
3. Merck, 11th edn., 6497
4. Kirk Othmer, 3rd edn., Vol. 6, 866
5. Editor's comments, 1999

The oxidising power and hazard potential of nitric acid increases progressively with increase in strength from the conc. acid (70 wt%) through fuming acids (above

HNO₃

85 wt%) to the anhydrous 100% acid. Nitric acid is the common chemical most frequently involved in reactive incidents (note the extent of this compound entry), and this is a reflection of its exceptional ability to function as an effective oxidant even when fairly dilute (unlike sulfuric acid) or at ambient temperature (unlike perchloric acid). Its other notable ability to oxidise most organic compounds to gaseous carbon dioxide, coupled with its own reduction to gaseous 'nitrous fumes' has been involved in many of the incidents in which closed, or nearly closed, reaction vessels or storage containers have failed from internal gas pressure [2]. Screw-capped winchesters of fuming acid may therefore develop internal pressure if the inert liner to the plastic cap fails or is missing.

There is a potentially dangerous confusion over the meaning of fuming nitric acid. Once, and sometimes still [3], this meant the red fuming acid which consists of nitrogen dioxide dissolved in nitric acid. This is a potent oxidising agent [1] and best avoided for nitrations, where oxidation is an unwanted and dangerous side-reaction. Increasingly, however, the white fuming acid of American Chemical Society analytical specification [4] is meant. This is a good nitrating agent but a poor oxidant, showing induction periods of hours in length followed by auto-catalytic runaway. When following previous literature it is wise to consider which was meant. Nitric acid generally is becoming whiter and less oxidant, which can be partially rectified by adding sodium nitrite.

Anhydrous nitric acid is miscible even with hydrocarbons, and the increased availability of high quality white grades increases the danger of inadvertently forming such mixtures. If anywhere near stoichiometric composition, an homogeneous mixture of nitric acid and virtually any organic is a sensitive high explosive. So much was obvious to learned men of the last century, and many such mixtures were patented as explosives, by Sprengel and others, from 1871 onwards. Recent correspondence makes it apparent that this is neither known nor obvious to many Professors of Chemistry in the late 1990s: though learning has departed, danger has not [5]!

See Polyalkenes, below

See other GAS EVOLUTION INCIDENTS

Preparative hazard

Crawley, F. K., *Loss Prev. Bull.*, 1992, (104), 4

Corrosion permitted leak of nitric acid into the ammonia feed passing through the cooling tubes of a nitric acid plant. An ammonium nitrate deposit built up and exploded when the tubes were cut out with a welding torch for replacement.

Acetic acid

1. Raikova, V. M., *Chem. Abs.*, 1983, **98**, 5958

2. Vidal, P. *et al.*, *C. R. Acad. Sci. Ser. II*, 1991, **313**(12), 1383; *J. Energ. Mater.*, 1993, **11**(21), 135.

Critical detonation diameters for mixtures have been determined.

Acetic acid, Sodium hexahydroxyplatinate(IV)

1. Davidson, J. M. *et al.*, *Chem. & Ind.*, 1966, 306

2. Malerbi, B. W., *Chem. & Ind.*, 1970, 796

During preparation of diacetatoplatinum(II) by alternative procedures, the hexahydroxyplatinate in mixed nitric–acetic acids was evaporated to a syrup and several explosions were experienced [1], possibly owing to formation of acetyl nitrate. On one occasion a brown solid was isolated and dried, but subsequently exploded with great violence when touched with a glass rod. The material was thought to be a mixture of platinum (IV) acetate–nitrate species [2].

Acetic anhydride

MRH 4.94/34

1. Brown, T. A. *et al.*, *Chem. Brit.*, 1967, **3**, 504
2. Dubar, J. *et al.*, *Compt. rend. C*, 1968, **266**, 1114
3. Dingle, L. E. *et al.*, *Chem. Brit.*, 1968, **4**, 136
4. Ullmann, A10, 157
5. *MCA Case History No. 103*
6. Venters, K. *et al.*, *Chem. Abs.*, 1982, **97**, 45372
7. Raikova, V. M., *Chem. Abs.*, 1983, **98**, 5958

Mixtures containing 50–85 wt% of fuming nitric acid are detonable and very sensitive to initiation by friction or shock (possibly owing to formation of acetyl nitrate and/or tetranitromethane). For preparation of mixtures outside these limits, the order of mixing is important (below 50%, acid into anhydride; above 85%, vice versa; and all below 10°C) [1]. Similar information is also presented diagrammatically [2]. Mixtures containing less than 50% of nitric acid are also dangerous in that addition of small amounts of water (or water-containing mineral acids) readily initiates an uncontrollable exothermic fume-off, which will evaporate most of the liquid present. Equimolar mixtures of 38% nitric acid with acetic anhydride can be detonated at room temperature after aging for a few h [3]. Others put the lower limit of detonability at 30% [4]. Accidental contact of the 2 materials caused a violent explosion [5]. Thermodynamic calculations show maximum values for heats of decomposition for acetyl nitrate or its equivalent mixtures of the components [6]. Critical detonation diameters for mixtures have been determined [7].

See 1,3,5-Triacetylhexahydrotriazine, etc., below

Acetic anhydride, Hexamethylenetetramine acetate

Leach, J. T., *J. Haz. Mat.*, 1981, **4**, 271–281

The military explosives RDX and HMX are manufactured from the 3 components using the Bachman process. Some of the possible mixtures may lead to fires in open vessels and explosions under confinement, and the exothermic and other effects (some calculated by the CHETAH program) for a wide range of mixtures are presented as ternary diagrams. It was also found that acetic anhydride layered onto solutions of ammonium nitrate in nitric acid exploded, owing to formation of acetyl nitrate.

Acetone

MRH 5.86/23

1. Kennedy, R., private comm., 1975
2. Anon., *Safety Digest Univ. Safety Assoc.*, 1989 **34**, 14

A winchester of fuming nitric acid with a plastics cap burst, probably owing to internal pressure build-up and uneven wall thickness. The explosion fractured

HNO₃

an adjacent bottle of acetone which ignited on contact with the oxidant [1]. Segregation of oxidants and fuels in storage is essential to prevent such incidents, which have occurred previously elsewhere. Another explosion resulted, when attempting to clear a jammed glass stopper by successive application of acetone and nitric acid [2].

See Polyalkenes, below

See also SAFE STORAGE OF CHEMICALS

Acetone, Acetic acid

1. Frant, M. S., *Chem. Eng. News*, 1960, **38**(43), 56

2. Secunda, W. J., *Chem. Eng. News*, 1960, **38**(46), 5

A mixture of equal parts of nitric acid, acetone and 75% acetic acid, used to etch nickel, will explode 1.5–6 h after mixing if kept in a closed bottle. The presence of the diluted acetic acid would probably slow the known violent oxidation of acetone by nitric acid [1] (but would not prevent slow oxidation with gas evolution). Alternatively, the formation of tetranitromethane and subsequent oxidation of acetone was postulated [2], but this is perhaps unlikely, because presence of acetic anhydride is normally necessary for formation of tetranitromethane.

Acetone, Sulfuric acid

Fawcett, H. H., *Ind. Eng. Chem.*, 1959, **51**, 89A

Acetone is oxidised violently by mixed nitric–sulfuric acids, and if the mixture is confined in a narrow-mouthed vessel, it may be ejected or explode.

Acetonitrile

MRH 6.15/23

Andrussow, L., *Chim. Ind. (Paris)*, 1961, **86**, 542

Mixtures of fuming nitric acid and acetonitrile are high explosives.

4-Acetoxy-3-methoxybenzaldehyde

Legrand, M. *et al.*, *J. Chem. Educ.*, 1978, **55**, 769

The dangerously exothermic nitration of the aldehyde with conc. nitric acid is adequately controlled on a several hundred g scale by an automated assembly of laboratory apparatus.

Acetylene, Mercury(II) salts, Sulfuric acid

Orton, K. J. P. *et al.*, *J. Chem. Soc.*, 1920, **117**, 29

Contact of acetylene with the conc. acid in presence of mercury salts forms trinitromethane, explosive above its m.p., 15°C. Subsequent addition of sulfuric acid produces tetranitromethane, a powerful oxidant of limited stability, in high yield.

See Trinitromethane, also Tetranitromethane

Acrylonitrile

See Acrylonitrile: Acids

Acrylonitrile–methacrylate copolymer

Caprio, V. *et al.*, *J. Haz. Mat.*, 1983, **7**, 383–391

The Asahi process for wet-spinning the copolymer involves water dilution at below 0°C of a solution of the copolymer in aqueous 68 wt% nitric acid (the azeotropic composition). The potential for slow self heating and decomposition have been investigated experimentally with variations in several parameters in a Sikarex safety calorimeter. At 20% polymer content, the slow self heating starts even at ambient temperature, and later involves evolution of 30 mol of uncondensable gas per kg of copolymer with concomitant boiling of the nitric acid. A 2 step mechanism has been proposed.

See other GAS EVOLUTION INCIDENTS

Alcohols MRH Ethanol 5.56/24, furfuryl alcohol 5.86/28

1. Fawcett, H. H., *Chem. Eng. News*, 1949, **27**, 1396
2. Fromm, F. *et al.*, *Chem. Eng. News*, 1949, **27**, 1956
3. Unpublished observations, 1956
4. *MCA Case History No. 1152*
5. Potter, C. R., *Chem. & Ind.*, 1971, 501
6. Spengler, G. *et al.*, *Brennst. Chem.*, 1965, **46**, 117
7. Long, L. A., private comm., 1972
8. Kurbangalina, R. K., *Zh. Prikl. Khim.*, 1959, **32**, 1467–1470
9. Albright, Hanson, 1976, 341–343
10. Anon., *CIHSC Chem. Safety Summ.*, 1980, **51**, 4
11. Anon., *Loss Prev. Bull.*, 1980, (035), 7–9
12. Anon., *Univ. Safety Assoc. Safety News*, 1982, (16), 4
13. Mabbott, D. J., 1996, personal communication

A 15% solution of nitric acid in ethanol was used to etch a bismuth crystal. After removing the metal, the mixture decomposed vigorously [1]. Mixtures of nitric acid and alcohols ('Nital') are quite unstable when the concentration of acid is above 10%, and mixtures containing over 5% should not be stored [2]. The use of a little alcohol and excess nitric acid to clean sintered glassware (by 'nitric acid fizzing') is not recommended. At best it is a completely unpredictable approximation to a nitric acid–alcohol rocket propulsion system. At worst, if heavy metals are present, fulminates capable of detonating the mixture may be formed [3]. Chromic acid mixture is less hazardous for such cleaning operations. The Case History describes a violent explosion caused by addition of conc. acid to a tank car contaminated with a little ethanol [4]. During oxidation of cyclohexanol to the 1,2-dione by an established process, a violent explosion occurred. Two intermediates are possible suspects [5]. Furfuryl alcohol is hypergolic with high-strength nitric acid [6] and methanol has been used as a propellant fuel. It also readily forms the explosive ester, methyl nitrate [7]. Mixtures with ethylene glycol are easily detonated by heat, impact or friction [8]. The injector process for safe continuous nitration of glycerol and other alcohols is reviewed [9]. Use of the wrong proportions of denatured ethanol and nitric acid (5:95 instead of 95:5) led to an explosion [10]. During attempts to clean a reactor, unclear instructions led to 120 l of conc. nitric acid, rather than the aqueous 2.6% solution intended, being charged via a pump which contained some 5 l of 2-propanol from previous use. Rapid oxidation of the alcohol, (probably accompanied by formation of explosive isopropyl nitrate)

HNO₃

and gas evolution burst the reactor [11]. A stock solution of conc. nitric acid in methanol for cleaning glassware was stored in a screw-capped bottle which later burst from internal gas pressure. Such mixtures are dangerously unstable and should not be used or stored [12]. A 33% by volume solution of nitric acid in ethanol was mixed for metallurgical purposes in a large open bottle, after two hours the still warm mixture was capped. Perhaps 7 hours after mixing, there was an exceedingly violent explosion. It was decided not to use this mixture in future but to substitute one containing perchloric acid, (which may be no less hazardous) [13].

See Methanol, Sulfuric acid, below

See other GAS EVOLUTION INCIDENTS, NITRATION INCIDENTS, PLANT CLEANING INCIDENTS

Alcohols, Disulfuric acid

Rastogi, R. P. *et al.*, *Amer. Inst. Aero. Astronaut. J.*, 1966, **4**, 1083–1085

Ignition delays were determined for contact of various aliphatic alcohols with mixtures of disulfuric acid (*d*, 1.9) and red fuming nitric acid (*d*, 1.5). With ethanol, a minimum of 30 wt% of disulfuric acid was required for ignition.

Alcohols, Potassium permanganate

See Potassium permanganate: Alcohols, Nitric acid

Aliphatic amines

1. Schalla, R. L. *et al.*, *Amer. Rocket Soc. J.*, 1959, **29**, 33–39

2. Schneebeil, P. *et al.*, *Rech. Aeron.*, 1962, **87**, 33–46

In contact with white fuming nitric acid, ignition delays for tributylamine, tripropylamine and triethylamine were less than those of the corresponding dialkylamines, and under good mixing conditions, butylamine did not ignite [1]. Triethylamine ignites on contact with the conc. acid [2].

See Aromatic amines (reference 2); Cyclohexylamine; and Dinitrogen tetraoxide: Nitrogenous fuels, all below

Alkanethiols

1. Filby, W. G., *Lab. Practice*, 1974, **23**, 355

2. McCullough, F. *et al.*, *Proc. 5th Comb. Symp.*, 181, New York, Reinhold, 1955

3. Illingworth, B., private comm., 1980

4. Quam, G. N., *Safety Practice for Chemical Laboratories*, 52–53, Villanova (Pa.), Villanova Press, 1963

5. Reychler, A., *Bull. Soc. Chim. Belg.*, 1913, **110**, 217, 300

Oxidation of alkanethiols to alkanesulfonic acids with excess conc. acid as usually described is potentially hazardous, the exotherm often causing ignition of the thiol. A modified method involving oxidation under nitrogen and at temperatures 1–2°C above the m.p. of the thiol is safer and gives purer products [1]. Technical butanethiol (containing 28% of propane- and 7% of pentane-thiols) is hypergolic with 96% acid [2]. Oxidation of several thiols to the sulfonic acids by addition to stirred conc. acid had been effected normally, but when 2 new

batches of pentanethiol were used, flame was observed in the vapour phase a few s after addition. No unusual impurities were detected. The oxidation can be effected safely with nitric acid–water (2:3) with rapid stirring at below 35°C [3]. Oxidation of dodecanethiol [4] and hexadecanethiol [5] with fuming nitric acid proceeded explosively.

2-Alkoxy-1,3-dithia-2-phospholane ('O-Alkyl ethylene dithiophosphate')

Arbuzov, A. E. *et al.*, *Chem. Abs.*, 1953, **47**, 4833e,f

The methoxy and ethoxy derivatives ignite with the conc. acid.

See Phosphorus compounds, below

Alkylamines

1. Vouillamox, L., *Chimia*, 1995, **49**(11), 439

During propionitrile manufacture by oxidative amination of propanol, nitric acid was used for neutralisation. This eventually led to concentration of nitrates, which exploded, at the bottom of the final distillation column [1]. Many amines form highly crystalline nitrates, which are usually explosive once isolated.

See AMINIUM NITRATES

2-Aminothiazole

Biasutti, G. S. *et al.*, *Loss. Prev.*, 1974, **8**, 123–125

In an attempt to produce the nitrate salt, 2-aminothiazole was added to the required amount of nitric acid and the mixture stirred and heated: the mixture exploded without warning.

See other NITRATION INCIDENTS

2-Aminothiazole, Sulfuric acid

Silver, L., *Chem. Eng. Progr.*, 1967, **63**(8), 43

Nitration of 2-aminothiazole with nitric–sulfuric acids was normally effected by mixing the reactants at low temperature, heating to 90°C during 30 min and then applying cooling. When the cooling was omitted, a violent explosion occurred. Experiment showed that this was due to a slow exothermic reaction accelerating out of control under the adiabatic conditions. *N*-Nitroamines were not involved.

See other NITRATION INCIDENTS, SELF-ACCELERATING REACTIONS

Ammonia

Mellor, 1940, Vol. 8, 219

A jet of ammonia will ignite in nitric acid vapour.

Ammonium nitrate

Biasutti, 1981, 135

Decomposition of a 70% nitric acid–ammonium nitrate slurry explosive led to overflow, contact with wood and a fire. This spread to detonators, which initiated detonation of the slurry.

Anilinium nitrate

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

HNO₃

Although aniline may be hypergolic with nitric acid (below), anilinium nitrate dissolves unchanged in 98% acid and can be stored for long periods, though the solution has explosive properties.

See Cyclohexylamine, below

Anilinium nitrate, Inorganic additives

Munjal, M. N. *et al.*, *Indian J. Chem.*, 1967, **5**, 320–322

The effect of inorganic additives upon ignition delay in anilinium nitrate–red fuming nitric acid systems was examined. The insoluble compounds copper(I) chloride, potassium permanganate, sodium pentacyanonitrosylferrate and vanadium(V) oxide were moderately effective promoters, while the soluble ammonium or sodium metavanadates were very effective, producing vigorous ignition.

Aromatic amines

MRH Aniline 5.98/19

1. Kit and Evered, 1960, 239, 242
2. *Aniline*, 85, Allied Chem. Corp., New York, 1964
3. Miller, R. O., *Tech. Note No. 3884*, 1–32, Washington, Nat. Advisory Comm. Aeronaut., 1956
4. Spengler, G. *et al.*, *Brennst. Chem.*, 1965, **46**, 1172
5. Bernard, M. L. *et al.*, *Compt. rend. C*, 1966, **263**, 24–26
6. Schalla, R. L. *et al.*, *Amer. Rocket Soc. J.*, 1959, **29**, 38

Many aromatic amines (aniline, *N*-ethylaniline, *o*4-toluidine, xylidine, etc., and their mixtures) are hypergolic with red fuming nitric acid [1]. When the amines are dissolved in triethylamine, ignition occurs at –60°C and below [2]. Addition of a mixture of aniline, dimethylaniline, xylidine and pentacarbonyliron renders hydrocarbons hypergolic with conc. nitric acid [3]. Although aniline is not hypergolic with 96% acid, presence of sulfuric acid (5% or above) renders it so. Presence of dinitrogen tetroxide further reduces ignition delay [4]. The mechanism of ignition of *p*-phenylenediamine (or furaldehyde azine) on contact with red fuming nitric acid is quantified [5]. Traces of sulfuric acid may be essential for ignition to occur [6].

See Benzidine, below

Aromatic amines, Metal compounds

Rastogi, R. P. *et al.*, *Indian J. Chem.*, 1964, **2**, 301–307

The effects of various metal oxides and salts which promote ignition of amine–red fuming nitric acid systems were examined. Among soluble catalysts, copper(I) oxide, ammonium metavanadate, sodium metavanadate, iron(III) chloride (and potassium hexacyanoferrate(II) with *o*-toluidine) are most effective. Of the insoluble materials, copper(II) oxide, iron(III) oxide, vanadium(V) oxide, potassium chromate, potassium dichromate, potassium hexacyanoferrate(III) and sodium pentacyanonitrosylferrate(II) were effective.

See Anilinium nitrate, etc., also Aromatic amines (reference 3), both above

Aromatic compound

Anon., *Safety Digest Univ. Safety Assoc.*, 1989, **34**, 13

An explosion resulted on nitrating an unrevealed aromatic, possibly a phenethyl alcohol derivative, with fuming nitric at -40°C , adding the acid to substrate. Probably too cold and too basic for initial reaction, leading to runaway (oxidation?) when almost all the nitric acid had been charged. Nitration only proceeds when the acidity is sufficient to protonate much of the nitric acid present, in this context any oxygen in the substrate is a base. The editor's experience is that, for monoalkyl-benzenes, nitration is very slow, even with substantial sulfuric acid also present as catalyst and dehydrating agent, at -20°C and below. Red fuming nitric acid, which contains dissolved oxides, encourages oxidation (much more exothermic, autocatalytic and gas evolving) in place of nitration.

See Benzeneethanol, below

See other NITRATION INCIDENTS

Arsine–boron tribromide

See Arsine–boron tribromide: Oxidants

Benzeneethanol, Sulfuric acid

Rashevskaya, S. K. *et al.*, *Zh. Obshch. Khim.*, 1963, **33**(12), 3998; *Chem. Abs.*, 1964, 9178f

Indifferent yields of mononitro-products resulted from nitration in 1:1 nitric and sulfuric acid at 0°C . At 18°C there was an explosion. (The editor has dinitrated this compound with no trouble, but he used far more sulfuric acid and added the nitric acid to the other components).

See 2-Methoxyethylbenzene, Sulfuric acid; below

See other NITRATION INCIDENTS

Benzidine

Krishna, P. M. M. *et al.*, *Chem. Abs.*, 1981, **94**, 124107

Benzidine is hypergolic with red fuming nitric acid.

See Aromatic amines, above

Benzonitrile, Sulfuric acid

Anon., *Loss. Prev. Bull.*, 1978, (024), 168

Nitration of benzonitrile by addition to mixed nitrating acid had been effected uneventfully in 250 batches. Plant 'improvements' led to the same dosing vessel being used alternately for the nitrile and mixed acids. Some nitration occurred in the uncooled and unstirred dosing vessel containing a little residual nitrile and an explosion occurred during the second batch in the modified plant.

Benzo[*b*]thiophene derivatives

Brown, I. *et al.*, *Chem. & Ind.*, 1962, 982

During nitration of several amino derivatives, diazotisation and oxidation occurred to produce internal diazonium phenoxide compounds. 5-Acetylamino-3-bromobenzo[*b*]thiophene unexpectedly underwent hydrolysis, diazotisation and oxidation to the explosive compound below.

See 3-Bromo-2,7-dinitro-5-benzo[*b*]thiophenediazonium-4-oxide

See 4-Chloro-2-nitroaniline, below

See other NITRATION INCIDENTS, PLANT CLEANING INCIDENTS

N-Benzyl-*N*-ethylaniline, Sulfuric acid

Anon., *Loss. Prev. Bull.*, 1978, (024), 169; 1980, (038), 27

Addition of 2–3 l of the amine to a dosing vessel not completely free of traces of mixed acids led to a vigorous decomposition which burst a vent line.

See other NITRATION INCIDENTS, PLANT CLEANING INCIDENTS

1,4-Bis(methoxymethyl)2,3,5,6-tetramethylbenzene

Anon., *Jahresber.*, 1972, 84

Oxidation of the durene derivative to benzenehexacarboxylic acid (mellitic acid) in an autoclave is normally effected in stages, initially by heating at 80–104°C with the vent open to allow escape of the evolved gases. Subsequent heating to a higher temperature with the vent closed completes the reaction. On one occasion omission of the first vented heating phase led to explosive rupture of the autoclave at 80°C.

See other GAS EVOLUTION INCIDENTS

See Hydrocarbons, below

1,3-Bis(trifluoromethyl)benzene, Sulfuric acid

Grewer, T., *Proc. 1st Int. Symp. Prev. Occ. Risks in Chem. Ind.*, 153–160, Heidelberg, ISSA, 1970

The fluoro compound is resistant to nitration and an operating temperature of 90°C is necessary to ensure formation of the 5-nitro derivative. Under these conditions, the atmosphere (containing the fluoro compound, its nitro derivative and nitric acid vapours) in the nitration vessel is explosive and above the flash point. An unknown ignition source led to an explosion and rupture of the 3 cu. m vessel, and a maximum explosion pressure of 50 bar was confirmed experimentally. Such explosive atmospheres are not found in low temperature nitration reactions.

See Chlorobenzene, below

See other NITRATION INCIDENTS

Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

tert-Butyl-*m*-xylene, Sulfuric acid

Kotoyori, T., private comm., 1983

Normally, the hydrocarbon (360 kg) was tri-nitrated by slow addition during 15 h to premixed 97% nitric–98% sulfuric acids (wt ratio 1.04:1, 1470 kg) agitated at 35–40°C, maintaining the temperature (checked at 30 min intervals) by controlling the jacket cooling and rate of addition. Because of a confusing and inadequate control panel layout, stirring was accidentally stopped after 3.3 h and addition was continued for a further 2 h before the absence of agitation was noticed, when the temperature was 34°C, and addition was stopped. At this stage, a layer of the liquid hydrocarbon (45 kg, *d* = 0.8) would be on top of the solid trinitro compound (150 kg, m.p. 112°C), itself floating on the mixed acid (*d* about 1.75).

As the operator realised it would cause an immediate violent exotherm if full agitation were restored, he attempted to start mild agitation by flicking the agitator on for 1 s only. After some 20 s, yellow-brown fumes began to escape from the vent pipe, indicating the onset of exothermic decomposition, so he went to open the dump valve but was prevented by the thick fumes. During the next couple of minutes the velocity of the fumes (now white) and the associated noise increased to a very high level and the vessel finally exploded, the lid being blown off and the contents becoming ignited.

See other AGITATION INCIDENTS, NITRATION INCIDENTS

Cadmium phosphide

Juza, R. *et al.*, *Z. Anorg. Chem.*, 1956, **283**, 230

Reaction with conc. acid is explosive.

Cellulose

MCA SD-5, 1961

Cellulose may be converted to the highly flammable nitrate ester on contact with the vapour of nitric acid, as well as by the liquid acid.

Chlorobenzene

Anon., *Jahresber.*, 1974, 86

In a plant for the continuous nitration of chlorobenzene, maloperation during start-up caused the addition of substantial amounts of reactants into the reactor before effective agitation and mixing had been established. The normal reaction temperature of 60°C was rapidly exceeded by at least 60° and an explosion occurred. Subsequent investigation showed that at 80°C an explosive atmosphere was formed above the reaction mixture, and that the adiabatic vapour-phase nitration would attain a temperature of 700°C and ignite the explosive atmosphere in the reactor.

See 1,3-Bis(trifluoromethyl)benzene, above

See other AGITATION INCIDENTS, NITRATION INCIDENTS, SELF-HEATING AND IGNITION INCIDENTS

4-Chloro-2-nitroaniline

1. Elderfield, R. C. *et al.*, *J. Org. Chem.*, 1946, **11**, 820

2. *MCA Case History No. 1489*

The literature procedure for preparation of 4-chloro-2,6-dinitroaniline [1], involving direct nitration in 65% nitric acid, was modified by increasing the reaction temperature to 60°C 1 h after holding at 30–35°C as originally specified. This procedure was satisfactory on the bench scale, and was then scaled up into a 900 l reactor. After the temperature had reached 30°C, heating was discontinued, but the temperature continued to rise to 100–110°C. Decomposition set in with copious evolution of nitrous fumes and production of a very shock-sensitive explosive solid. This was identified as 4-chloro-2,5-dinitrobenzenediazonium-6-oxide, produced by hydrolysis of a nitro group in the expected product by the diluted nitric acid at high temperature, diazotisation of the free amino group by the nitrous acid produced in the hydrolysis (or by the nitrous fumes), and introduction of a further nitro

HNO₃

group under the prevailing reaction conditions. It is recommended that primary aromatic amines should be protected by acetylation before nitration, to prevent the possibility of accidental diazotisation.

See Benzo[*b*]thiophene derivatives, above

See also ARENEDIAZONIUM OXIDES

See other NITRATION INCIDENTS

Coal

Green, J. B. *et al.*, *Anal. Chem.*, 1979, **51**, 1126

The mixtures of bituminous coal and conc. nitric acid used to prepare nitrohumic acids are potentially explosive, and appropriate care is necessary.

See Organic materials, below

Contaminants

NSC Safety Newsletter, Chem. Sect., July, 1977

A 50% nitric acid solution was used to clean glass containers and then stored in a screw-capped bottle. Some time later the bottle burst, owing to gas generation by the dissolved contaminant(s).

See other GAS EVOLUTION INCIDENTS

Copper

1. Anon., *Loss Prev. Bull.*, 1987, (074), 23–27

2. Anon., *Loss Prev. Bull.*, 1988, (080), 31–34

A 'canned' electric pump was being used to pump 89% nitric acid during a plant malfunction when a violent explosion occurred. This was attributed to penetration of nitric acid into the sealed copper windings of the motor and subsequent generation of nitric oxide which pressurised the stator enclosure to some 60–280 bar, causing it to burst. Preventive precautions are detailed [1]. The pump manufacturers, however, subsequently challenged this account and explanation and revealed that the pump, which had been returned for service and repair on previous occasions, had shown signs of reverse rotation and presence of nitric acid in the windings. Such pressure build-up was most unlikely, since the stator enclosure had both an external vent and a separate pressure relief valve. The remains of the exploded pump also showed signs of reverse rotation and the impellor appeared to have come loose on the shaft prior to the explosion. The latter occurred when the motor was energised via replacement fuses 50% above recommended capacity. The precise cause of the explosion is not clear, but may well have been spark-initiated [2].

See IGNITION SOURCES

See other GAS EVOLUTION INCIDENTS

Copper(I) nitride

Mellor, 1940, Vol. 8, 100

Interaction with conc. acid is very violent.

Cotton, Rubber, Sulfuric acid, Water

Biasutti, 1981, 76

A rubber glove and a box of cotton waste fell into a tank of spent nitrating acid from the preparation of TNT. The total contents of the tank ignited, leading to a series of explosions which caused severe damage and some 900 casualties, including 82 killed.

See other NITRATION INCIDENTS

Crotonaldehyde

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

Crotonaldehyde is hypergolic with conc. nitric acid, ignition delay being 1 ms.

See ROCKET PROPELLANTS

Cycloalkanones

1. Dye, W. T., *Chem. Eng. News*, 1959, **37**, 48

2. Field, K. W. *et al.*, *J. Chem. Educ.*, 1985, **62**, 637

Oxidation of 4-methylcyclohexanone by addition of nitric acid at about 75°C caused a detonation to occur. These conditions had been used previously to oxidise the corresponding alcohol, but although the ketone is apparently an intermediate in oxidation of the alcohol, the former requires a much higher temperature to start and maintain the reaction. An OTS report, PB73591, mentions a similar violent reaction with cyclohexanone [1]. Presence of nitrous acid is essential for the smooth oxidation of cycloalkanones with nitric acid to α , ω -hexanedioic acids. Because high-purity nitric acid (free of nitrous acid) is now commonly available, addition of a little sodium or potassium nitrite to the acid is necessary before its use to oxidise cycloalkanones [2].

See Acetone, etc., above

Cyclohexanol, Cyclohexanone

1. Wilson, B. J. *et al.*, *Loss. Prev.*, 1979, **12**, 27–29

2. Vidal, P. *et al.*, *Thermochim. Acta*, 1993, **225**(2), 223

The process for manufacturing adipic acid by nitric acid oxidation of KA (the ketone–alcohol mixture produced by air oxidation of cyclohexane in the liquid phase) at 68°C is potentially hazardous, in that reactor or cooling failure could lead to a large and vigorous evolution of hot toxic nitrous fumes. The overall process has been studied and modified to reduce these potential hazards [1]. Solutions of organic diacids in nitric acid, such as result in adipic acid manufacture, were studied for exotherm, gas evolution, deflagration and detonation. Critical diameters were plotted against composition. Recommendations for plant design result [2].

Cyclohexylamine

MRH 6.07/18

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

Although cyclohexylamine has been used as a fuel with nitric acid in rocket motors, cyclohexylammonium nitrate dissolves unchanged in fuming nitric acid to give a solution stable for long periods.

See Anilinium nitrate, above

See also ROCKET PROPELLANTS

1,3-Diacetoxybenzene

See 4,6-Dinitro-1,3-benzenediol

See other INDUCTION PERIOD INCIDENTS, NITRATION INCIDENTS

1,2-Diaminoethanebis(trimethylgold)

See 1,2-Diaminoethanebis(trimethylgold)

Di-2-butoxyethyl ether

Hanson, C. *et al.*, *Proc. Int. Solvent Extr. Conf.* (Liege), 1980, 2, paper 80–70

The solvent ('Butex'), an extractant in nuclear reprocessing may decompose violently in contact with nitric acid. Hydrolysis to butanol, followed by violent oxidation, catalysed by nitrous acid, is involved.

See Alcohols, above

1,2-Dichloroethane

Kurbangalina, R. K., *Zh. Prikl. Khim.*, 1959, **32**, 1467

Mixtures are easily detonated by heat, impact or friction.

Dichloroethylene

Raikova, V. M., *Chem. Abs.*, 1983, **98**, 5958

Critical detonation diameters for mixtures have been determined.

Dichloromethane

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

Dichloromethane dissolves endothermically in conc. nitric acid to give a detonable solution.

Dichromates, Organic fuels

Munjal, N. L. *et al.*, *Amer. Inst. Aero. Astronaut. J.*, 1972, **10**, 1722–1723

The effects of dichromates in promoting ignition of non-hypergolic mixtures of red fuming nitric acid with cyclohexanol, 2-cresol or 3-cresol, and furfural were studied. Ammonium dichromate was most effective in all cases, and the only effective catalyst for cyclohexanol. Potassium chromate and potassium dichromate were also examined.

Diethylaminoethanol

See 2-Diethylammonioethyl nitrate nitrate

Diethyl ether

MRH 5.94/21

1. Foote, C. S., private comm., 1965

2. Holt, C., *School. Sci. Rev.*, 1976, **58**(202), 159–160

3. Fischer, E., *Ber.*, 1902, **35**, 3794

4. Anon., *Sichere Chemiearb.*, 1981, **33**, 94–95

5. Anon., *Univ. Saf. Assoc., Saf. Digest*, 1988, (25), 6–8

Addition of ether to a nitration mixture (2-bromotoluene and conc. nitric acid) diluted with an equal volume of water in a separating funnel led to a low order explosion. This was attributed to oxidation of the ether (possibly containing alcohol) by the

acid. Addition of more water before adding ether was recommended [1]. Attention is drawn to the delayed vigorous decomposition reaction which occurs some time after the initially homogeneous mixture of conc. nitric acid and ether has separated into 2 phases (possibly involving formation of ethyl nitrate?) [2]. In the preparation of 3-chlorolactic acid by oxidation of epichlorhydrin with nitric acid [3], the product was extracted into ether. Vacuum evaporation of solvent at 40°C led to a violent explosion, wrongly attributed to formation of peroxides. Experiment showed that extraction of 200 ml of a 30% aqueous solution of nitric acid with 8 × 200 ml portions of ether extracted 88% of the acid into the ether [4]. Upon concentration, this would produce very concentrated warm acid in contact with organic material, which would be oxidised violently or explosively (and could form ethyl nitrate). It is dangerous to extract nitric acid solutions with ether or other organic solvents without first removing the excess acid, either by neutralisation or ion exchange.

Nitric acid was being used to clean laboratory glassware to a very high state of surface cleanliness by soaking in a bucket of conc. acid for a week. During this time, one of the empty winchesters was inadvertently used as a solvent residues bottle and some 250 ml of diethyl ether was put in. The acid was to be recovered for re-use and it was put back into the original bottles. When the bottle containing ether was refilled and screw capped, after a few seconds it exploded violently [5].
See Glassware, below

Diethyl ether, Sulfuric acid

1. Kirk-Othmer, 1965, Vol. 8, 479
2. Van Alphen, J., *Rec. Trav. Chim.*, 1930, 492–500

Interaction of ether with anhydrous nitric acid to produce ethyl nitrate may proceed explosively [1], and in presence of conc. sulfuric acid, ether and conc. nitric acid explode violently [2].

See Dimethyl ether, below

3,6-Dihydro-1,2,2*H*-oxazine

Wichterle, O. *et al.*, *Coll. Czech. Chem. Comm.*, 1950, **15**, 309–321

The product of interaction (possibly an *N*-nitro and/or ring-opened nitrate ester derivative) is explosive.

Dimethyl 4-acetamidophthalate

See Dimethyl 4-acetamidophthalate: Nitric acid

Dimethylaminomethylferrocene, Water

Koch-Light Ltd., private comm., 1976

In an assay procedure the amine was heated with diluted nitric acid, and near-explosive decomposition occurred.

Dimethyl ether

Andrussow, L., *Chim. Ind. (Paris)*, 1961, **86**, 542–545

The latent hazards in storing and handling the explosive mixtures with the conc. acid are discussed (methyl nitrate may be formed).

See Diethyl ether, above

1,1-Dimethylhydrazine

See Hydrazine and derivatives, below

1,1-Dimethylhydrazine, Organic compounds

Spengler, G. *et al.*, *Brennst. Chem.*, 1965, **46**, 117

Contact of nitric acid (or dinitrogen tetroxide) with dimethylhydrazine is hypergolic and well described in rocket technology. While hydrocarbons and several other classes of organic compounds are not hypergolic with these oxidisers, addition of a proportion of dimethylhydrazine to a wide range of hydrocarbons, alcohols, amines, esters and heterocyclic compounds renders them hypergolic in contact with nitric acid or dinitrogen tetroxide.

Dimethyl sulfide, 1,4-Dioxane

Rudakov, E. S. *et al.*, *Chem. Abs.*, 1972, **76**, 13515

The mechanism which leads to delayed explosion in the system with nitric acid, even when cooled in liquid nitrogen, was investigated.

Dimethyl sulfoxide, Water

Fiola, J. *et al.*, *Czech. Pat.* 170 169, 1977

During the oxidation of the sulfoxide to the sulfone, the concentration of water in the reaction mixture must be maintained above 14% to prevent the risk of detonation.

1,3-Dinitrobenzene

MRH 5.02/99+

Biasutti, 1981, 92

After a detonation during manufacture of an explosive solution of the nitro compound in conc. nitric acid, manufacture was discontinued as being too dangerous.

Dinitrogen tetroxide, Nitrogenous fuels

1. Durgapal, U. C. *et al.*, *Amer. Inst. Aero. Astronaut. J.*, 1974, **12**, 1611

2. Munjal, N. L. *et al.*, *J. Spacecr. Rockets*, 1974, **11**, 4286–430

3. Durgapal, U. C. *et al.*, *J. Spacecr. Rockets*, 1974, **11**, 447–448

The effect on decrease in hypergolic ignition delay of increasing concentrations of dinitrogen tetroxide in red fuming nitric acid was studied with triethylamine, dimethylhydrazine or *miso*-xylydine as fuels [1]. The effect of various catalysts on ignition delay after contact of red fuming nitric acid with various arylamine–formaldehyde condensation products was also studied [2,3].

Dioxane, Perchloric acid

See Perchloric acid: Dioxane, Nitric acid

Diphenyldistibene

See Diphenyldistibene: Air, etc.

Diphenylmercury

Whitmore, 1921, 43, 168

Interaction in carbon disulfide, even at -15°C , is violent.

Diphenyltin

Krause, E. *et al.*, *Ber.*, 1920, **53**, 177

Ignition occurs on contact with fuming nitric acid.

Disodium phenyl orthophosphate

Muir, G. D., private comm., 1968

Concentration by evaporation of the nitration product of the phosphate ester caused a violent explosion, possibly involving picric acid derivatives from over-nitration.

Divinyl ether

MRH 6.44/22

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

Divinyl ether is hypergolic with conc. nitric acid, ignition delay being 1 ms.

See ROCKET PROPELLANTS

Ethanesulfonamide

Suter, C. M., *Organic Chemistry of Sulfur*, 111, London, Wiley, 1944

Reaction with nitric acid at a moderate temperature is explosive, dinitrogen oxide being evolved.

5-Ethyl-2-methylpyridine

1. Frank, R. L., *Chem. Eng. News*, 1952, **30**, 3348

2. Rubinstein, H. *et al.*, *J. Chem. Eng. Data*, 1967, **30**, 3348

3. Elam, E. V. *et al.*, *Chem. Eng. News*, 1973, **51**(34), 42

Following a patented procedure, the two reactants were being heated together at 145°C/14.5 bar to produce 2,5-pyridinedicarboxylic acid. The temperature rose to 160°C/43.5 bar, and the autoclave was vented and cooled, but 90 s later a violent explosion occurred, although both rupture disks (rated 105 and 411 bar) had relieved. General precautions are discussed [1]. Up to 20% of 5-(1,1-dinitroethyl)-2-methylpyridine, probably an explosive compound, is produced in this reaction. However, in presence of added water, no instability was seen in a series of reactions at temperatures up to 160°C and pressures up to 102 bar [2]. A further incident in an electrically heated titanium-lined autoclave involved heating the pyridine with excess 70% acid to 140°C during 1 h. When heating was discontinued, the temperature continued to rise in spite of air cooling, and an explosion occurred at 172°C which bulged the autoclave [3]. It is remotely possible that reaction of the titanium liner with the pressurised hot acid may have contributed to the incident. *See* Metals (reference 1), below *also* Titanium: Nitric acid

Fat, Sulfuric acid

1. Tyler, L. J., *Chem. Eng. News*, 1973, **51**(31), 32

2. Engan, W. L., *Chem. Eng. News*, 1973, **51**(51), 37

A wet-ashing procedure for analysis of fatty animal tissue was modified by using Teflon-lined bombs rated for use at 340 bar instead of open crucibles. Bombs cooled to well below 0°C were charged with fuming nitric and fuming sulfuric acids (1 ml of each) and adipose tissue (0.5 g), removed from the cooling bath and sealed. After 10 min delay, the bombs exploded, probably owing to development of

HNO₃

high internal temperature and pressure (calculated as 4000 K and 1000 bar max.) from complete oxidation of all the organic material [1]. Formation of glyceryl nitrate from the lipid content may have contributed to the violence of the explosion. The presence of 2–5% of water in the mixed acids is recommended to reduce the nitrating potential when in contact with organic fuels [2].

See Organic materials

See Organic materials

See Sulfuric acid

Formaldehyde

1. Pirie, C. J. *et al.*, *Chem. Brit.*, 1979, **15**, 11–12
2. Bretherick, L., *CISHC Chem. Safety Summ.*, 1979, **50**(197), 3
3. Vergnano, L. P., *CISHC Chem. Safety Summ.*, 1979, **50**(197), 2
4. Thurger, E. T., *CISHC Chem. Safety Summ.*, 1979, **50**(197), 3
5. Walker, J. F., *Formaldehyde, ACS Monograph 159*, 254–255, New York, Reinhold, 3rd ed. 1964
6. Winsor, L., *Chem. Brit.*, 1979, **15**, 281–282
7. Anon., *Loss. Prev. Bull.*, 1978, (020), 42
8. Haas, P. A., *Chem. Eng. News*, 1987, **65**(16), 2

A closed bottle of decalcifying fluid (a 10% nitric acid–formalin mixture) exploded a few h after the contents had been used and returned to the bottle for storage [1]. The incident was attributed to the interaction of an oxidant with a reductant with evolution of gas, promoted by dissolved impurities arising from its use, leading to pressurisation of the bottle to its bursting point [2,3,4]. This reaction is discussed in detail [5]. 5–10% Solutions of nitric acid or formic acid are preferred decalcifying fluids [6]. After failure of a pump which continuously fed nitric acid into circulating formaldehyde solution (used as a plant cleaning mixture), 30 l of 60% nitric acid was batch-charged into a filter vessel in the fluid circuit. When circulation was restarted to mix in the acid and continue the cleaning operation, a violent redox reaction blew off the lid of the filter vessel [7]. Process development wastes containing urea and hexamethylenetetramine were acidified with nitric acid to dissolve hydrated metal oxides. Under these conditions, hexamine hydrolyses to formaldehyde and ammonium nitrate, and it was calculated that the composition of the acidified waste would be 2M nitric acid, 2M formaldehyde, 1M ammonium nitrate and 0.5M metal nitrate. Such mixtures stored for 2 months in a loosely capped polythene bottle decomposed with vigorous gas evolution which ruptured the container [8].

See other GAS EVOLUTION INCIDENTS, PLANT CLEANING INCIDENTS, REDOX REACTIONS

Formic acid

1. Halze, K. *et al.*, *Proc. 3rd. Int. Symp. Loss. Prev. Safety Prom. Proc. Ind.*, 1413–1424, Basle, SSCI, 1980

The use of excess formic acid to destroy excess nitric acid (5M) in nuclear fuel reprocessing waste solutions at 100°C is potentially hazardous because of an induction period, high exothermicity and the evolution of large amounts of gas, mainly

carbon dioxide, dinitrogen oxide and nitrogen oxide, with some nitrogen and dinitrogen tetraoxide. The system has been studied thermokinetically, and the effects of various salts (which decrease the reaction rate) and sulfuric acid (which increases the rate) were determined [1].

See other GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS, REDOX REACTIONS

Formic acid, Oxalic acid

Cardillo, P. *et al.*, *Chim. Ind.* (Milan) 1989 **71**(4) 61

A study of, and details of how to control, the stop–start nature of the oxidation of formic acid. Probably relevant to other nitric acid oxidations, especially preparation of glyoxylic acid from glyoxal.

Formic acid, Urea

Mejdell, G. T., *Proc. 3rd. Int. Symp. Loss. Prev. Safety Prom. Proc. Ind.*, 1325–1338, Basle, SSCI, 1980

In the production of formic acid, a slurry of calcium formate in 50% aqueous formic acid containing urea is acidified with strong nitric acid to convert the calcium salt to free acid, and interaction of formic acid (reducant) with nitric acid (oxidant) is inhibited by the urea. When only 10% of the required amount of urea had been added (unwittingly, because of a blocked hopper), addition of the nitric acid caused a thermal runaway (redox) reaction to occur which burst the (vented) vessel. A small-scale repeat indicated that a pressure of 150–200 bar may have been attained. A mathematical model was developed which closely matched experimental data.

See other REDOX REACTIONS, RUNAWAY REACTIONS

2-Formylamino-1-phenyl-1,3-propanediol

1. Biasutti, G. S. *et al.*, *Loss Prev.*, 1974, **8**, 123

2. Biasutti, 1981, 131

Large scale nitration to the *p*-nitro derivative in a stirred and pump-circulated reactor had been effected uneventfully for 15 years when a violent explosion occurred. Subsequent investigation showed that with sufficient priming energy, the reaction mixture could be caused to detonate with an energy release of 3.77 kJ/g [1]. It was supposed that the priming energy originated from a bolt from the agitation gear being sucked into the reactor circulating pump [2].

See other AGITATION INCIDENTS, NITRATION INCIDENTS

Fluorine

MRH 0.50/47

See Fluorine: Nitric acid

Furfural

Giller, S. A. *et al.*, *Brit. Pat.*, 1 466 043, 1977

A safe method of nitrating furfuraldehyde or its *O,O*-diacetate to 5-nitrofurfural diacetate is described which minimises the formation of acetyl nitrate.

Furfurylidene ketones

Panda, S. P. *et al.*, *J. Armam. Stud.*, 1974, **10**(2), 44–45; 1976, **12**, 138–139

HNO₃

In 10 out of 12 cases, furfurylidene derivatives of ketones were hypergolic with red fuming nitric acid. Difurfurylidene hydrazine behaves similarly, and a mechanism involving polymerisation of the compounds is proposed.

Glassware

Anon., *Univ. Saf. Assoc., Saf. Digest*, 1988, (25), 6–8

Nitric acid has long been used to ensure absolute surface cleanliness on laboratory glassware where this is essential for technical reasons. Chromic acid cannot be used for this purpose because a surface film of adsorbed chromium ions remains after cleaning. Following an accident in which some diethyl ether was inadvertently put into the bottle used to store the conc. nitric acid after use and which led to an explosion, a Code of Practice covering the use of acid cleaning procedures for glassware was developed. This proposed that formulated detergents should be used for routine cleaning, and that use of acid processes would only be permitted if it could be practically demonstrated that they were essential for the proposed work. The Code covers the preparation of written procedures dealing with all aspects of the use, storage and disposal of acids, and personal protection requirements.

See Diethyl ether, above;

See CLEANING BATHS FOR GLASSWARE

See other GLASS INCIDENTS

Glycerol

Sanders, R. E., *Chem. Eng. Progress*, 1992, 88(5), 78

Glycerol was inadvertently used as the fluid in a gas bubbler on a nitric acid plant. It nitrated and detonated two days later.

Glycerol, Hydrofluoric acid

1. Buck, R. H., *J. Electrochem. Soc.*, 1966, **113**, 1352–1353

2. Wright, J. L., private comm., 1984

3. Anon., *Safety Digest Univ. Safety Assoc.*, 1992, **44**, 25

A chemical polishing solution consisting of nitric acid and hydrofluoric acid (1 Vol. each) and glycerol (2 vols.) generated enough pressure during storage for 4 h to rupture the closed plastics container. This was caused by gas evolution from oxidation of glycerol by the strongly oxidising mixture [1]. A mixture of nitric acid (80 ml), hydrofluoric acid (80 ml) and glycerol (240 ml) was used immediately for etching metal, again the next day, and then stored in a stoppered flask. After some 2–3 days, the stopper was ejected and 300 ml was sprayed around the fume cupboard containing the flask [2]. The metals dissolved during use further destabilise the mixture, which should not be stored under any circumstances. A freshly made etching mixture with hydrochloric acid, which one would expect to catalyse the oxidant power of nitric acid, burst its bottle in 30 minutes [3].

See Alcohols, above; Hydrofluoric acid, Lactic acid, below

See other GAS EVOLUTION INCIDENTS

Glycerol, Sulfuric acid

Dunn, B. W., *Bur. Expl. Accid. Bull.*, 35, Pennsylvania, Amer. Rail Assoc., 1916

Charging mixed nitrating acid into an insufficiently cleaned glycerol drum led to a violent explosion. Formation and detonation of glyceryl nitrate may have added to the oxidation energy release.

Glyoxal

Urban, P. G., personal experience

The oxidation of aqueous ethanedial to glyoxylic acid is inclined to induction periods and then runaway, cessation and renewed runaway, as the nitric acid is progressively added. Probably this is the same problem as with formic acid, oxalic acid (above).

Hexalithium disilicide

See Hexalithium disilicide: Acids

2,2,4,4,6,6-Hexamethyltrithiane

Baumann, E. *et al.*, *Ber.*, 1889, **22**, 2596

Oxidation of 'tri(thioacetone)' by conc. nitric acid is explosively violent.

See Thioaldehydes, etc., below

2-Hexenal

Lobanov, V. I., *Chem. Abs.*, 1966, **64**, 14013c

Hexenal is determined by photolorimetry after oxidation with nitric acid. The yellow-orange oxidation product explodes on heating.

Hydrazine and derivatives

1. Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

2. Jain, S. R., *Combust. Flame*, 1977, **28**, 542

3. Jain, S. R. *et al.*, *J. Spacecr. Rockets*, 1979, **16**, 69–73

Hydrazine is hypergolic with conc. nitric acid [1]. Of a series of hydrazones and azines derived from aldehydes and ketones, only those which decomposed when heated alone were hypergolic with the acid when heated at 12.5°C/min [2]. Solid hydrazones formed from various aldehydes with dimethylhydrazine or phenylhydrazine are hypergolic with the acid [3].

See ROCKET PROPELLANTS

Hydrocarbons

1. Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

2. Wilson, P. J. *et al.*, *Chem. Rev.*, 1944, **34**, 8

3. Trent, C. H. *et al.*, *J. Amer. Rocket Soc.*, 1951, **21**, 129–131

4. Ingolgaonkar, M. B. *et al.*, *J. Armam. Stud.*, 1979, **15**(2), 33–42

5. Sykes, W. G. *et al.*, *Chem. Eng. Progr.*, 1963, **59**(1), 70–71

6. Mason, C. M. *et al.*, *J. Chem. Eng. Data*, 1965, **10**, 173

7. Urbanski, 1961, Vol. 1, 140

8. Derbyshire, D. H., *Proc. 1st Symp. Chem. Process Haz. Ref. Plant Des.*, 37–41, Symp. Ser. No. 7, London, IChE, 1960

9. Powell, G. *et al.*, *Org. Synth.*, 1943, Coll. Vol. 2, 450

10. Wilms, H. *et al.*, *Angew. Chem.*, 1962, **74**, 465

11. Chaigneau, M. *Compt. rend.*, 1951, **223**, 657–659
 12. Anon., *Chemisches Ind.*, 1914, **37**, 337–342
 13. Biasutti, 1981, 48
 14. Bryant, T. J. *et al.*, *Rept. AD-B-000447*, Richmond (Va.), USNTIS, 1974
 15. Panda, S. P. *et al.*, *Chem. Abs.*, 1981, **94**, 194475
 16. Dutta, P. K. *et al.*, *J. Armam. Stud.*, 1976, **12**, 108–117
 17. Mellor, 1964, Vol. 8, Suppl. 2, 341
- MRH Benzene 8.15/18, cyclopentadiene 6.44/18, divinylbenzene 6.23/19, hexane 6.07/18, toluene 6.07/18, *o*-xylene 6.07/17

Dienes and acetylene derivatives are hypergolic in contact with conc. nitric acid, ignition delay being 1 ms [1]. Cyclopentadiene reacts explosively with fuming nitric acid [2], igniting under nitrogen [3], and dicyclopentadiene is highly hypergolic with red or white fuming nitric acid, also in presence of sulfuric acid [4]. Burning fuel oil and other petroleum products detonate immediately on contact with conc. nitric acid [5]. Very high sensitivity is shown by mixtures with benzene close to the stoichiometric proportions of around 84% acid [6]. Lack of proper control in nitration of toluene with mixed acids may lead to runaway or explosive reaction. A contributory factor is the oxidative formation, and subsequent nitration and decomposition, of nitrocresols [7]. Oxidation of *p*-xylene with nitric acid under pressure in manufacture of terephthalic acid carries explosion hazards in the autoclaves and condensing systems [8]. During nitration of mesitylene in acetic acid–acetic anhydride solution, fuming nitric acid must be added slowly to the cooled mixture to prevent the temperature exceeding 20°C, when an explosive reaction may occur [9]. During oxidation of mesitylene with nitric acid in an autoclave at 115°C to give 3,5-dimethylbenzoic acid, a violent explosion occurred. This was attributed to local overheating, formation of 1,3,5-tris(nitromethyl)benzene and violent decomposition of the latter. Smaller scale preparations with better temperature control were uneventful [10].

The explosive hazards involved in the preparation of mellitic acid by fuming nitric acid oxidation of hexamethylbenzene make the procedure only suitable for small scale operation [11]. Large-scale addition of too-cold nitrating acid to benzene without agitation later caused an uncontrollably violent reaction to occur when stirring was started. The vapour–air mixture produced was ignited by interaction of benzene and nitric acid at 100–170°C and caused an extremely violent explosion [12]. Failure to start the agitator or to apply water cooling in nitration of benzene led to a pressure burst of the nitrating vessel [13]. In the impact-initiated combustion of hydrocarbons in acid at high pressure, the effects of impact, hydrocarbon structure and stoichiometry upon ignition were studied [14]. The pure terpene 3-carene is more hypergolic with red fuming nitric acid than the mixture with other terpenes in turpentine [15]. The use of turpentine and cashew nutshell oil as bases for hypergolic rocket fuels with red fuming nitric acid has been evaluated [16]. The nitric acid–gasoline propellant system is non-hypergolic, but can be made so by addition of 15–25% of unsaturated hydrocarbons to the gasoline [17]. *See other* AGITATION INCIDENTS, NITRATION INCIDENTS

Hydrocarbons, 1,1-Dimethylhydrazine

Ingolgaonkar, M. B. *et al.*, *J. Armam. Stud.*, 1979, **15**(2), 33–42

Presence of the hydrazine confers hypergolicity upon mixtures of several non-hypergolic hydrocarbons with red fuming nitric acid.

Hydrofluoric acid, Lactic acid

Bubar, S. F. *et al.*, *J. Chem. Educ.*, 1966, **43**, A956

Mixtures of the 3 acids, used as metal polishing solutions, are unstable and should not be stored. Lactic acid and nitric acid react autocatalytically after a quiescent period, attaining a temperature of about 90°C with vigorous gas evolution after about 12 h. Prepare freshly, discard after use and handle carefully.

See Glycerol, etc., above

See other INDUCTION PERIOD INCIDENTS

Hydrofluoric acid, Propylene glycol, Silver nitrate

Leleu, *Cahiers*, 1979, (94), 125

A chemical polishing mixture was put into a closed glass bottle which burst 30 min later, and formation of silver fulminate was suggested. However, in absence of the silver salt such mixtures evolve gas and should not be stored in any event, especially after use for metal polishing, when the dissolved metal(s) tend to further destabilise the mixture.

See Glycerol, etc., above

Hydrogen iodide, or Hydrogen selenide, or Hydrogen sulfide

See Non-metal hydrides, below

Hydrogen peroxide, Ketones

See Hydrogen peroxide, Ketones, etc.

Hydrogen peroxide, Mercury(II) oxide

See Hydrogen peroxide: Mercury(II) oxide, etc.

Hydrogen peroxide, Soils

Krishnamurty, K. V. *et al.*, *At. Abs. Newslett.*, 1976, **15**, 68–70

When preparing soil and sediment samples for atomic absorption spectral analysis for trace metals, pre-oxidation with nitric acid before addition of hydrogen peroxide eliminates the danger of explosion.

Ion exchange resins

1. Barghusen, J. *et al.*, *Reactor Fuel Process.*, 1964, **7**, 297

2. McBride, J. A. *et al.*, *3rd Geneva Conf. Peaceful Uses At. En. A/CONF. 28 28/P/278*, 1965

3. Kalkworf, D. R., *Rept. BNWL-2931*, Richland (Wa.), Battelle, 1977

4. Colman, C., *Chem. Eng. (New York)*, 1980, **87**, 271–274

Several cases of interaction between anion exchange resins and nitric acid causing rapid release of energy or explosion have occurred [1]. The cause has been attributed to oxidative degradation of the organic resin matrix and/or nitration of

HNO₃

the latter. Suggested precautions include control of temperature, acid concentration and contact time. Presence of heavy ions (Pu) or oxidising agents (dichromates) tends to accelerate the decomposition [2]. The present state of knowledge was reviewed and relevant process parameters identified [3]. Several case histories are given, and safety aspects of regeneration of large-volume beds of strong-base anion exchange resins or cationic exchangers with nitric acid are discussed, but weak-base resins should not be treated with nitric acid. Storage of nitric acid-containing resins may lead to ignition [4].

See other SELF-HEATING AND IGNITION INCIDENTS

Iron(II) oxide

Mellor, 1941, Vol. 13, 716

The finely divided (pyrophoric) oxide incandesces with nitric acid.

Lead-containing rubber

Johnson, T. C. *et al.*, *Rep. RFP-1354*, 1–8, Washington (DC), USAEC, 1969

Radiation resistant lead-containing dry-box gloves may ignite in nitric acid environments.

Magnesium, 2-Nitroaniline

Gune, S. G. *et al.*, *J. Armam. Stud.*, 1983, **18–19**, 39–41

Though neither of the components is individually hypergolic with red fuming nitric acid, mixtures with magnesium containing 20% and 30% of 2-nitroaniline ignite with 67 and 23 ms delay, respectively.

Magnesium silicide

Mayes, R. B. *et al.*, *School Sci. Rev.*, 1975, **56**(197), 819–820

When the residue from combustion of magnesium in air was removed from the porcelain crucible, a grey stain remained. Addition of cold conc. nitric acid to remove the stain led to a violent reaction. This was found to be caused by the presence of magnesium silicide in the stain.

Magnesium phosphide

Mellor, 1940, Vol. 8, 842

Oxidation of the phosphide proceeds with incandescence.

Manufacture hazard

Lawrence, G. M. , *Plant/Oper. Progr.*, 1989, **8**(1), 33

An explosion in the vent of an ammonia combustion plant was attributed to deposition of ammonium nitrite/nitrate crystals. It is considered that the very unstable nitrite acts as a sensitiser to the nitrate, and that explosion is triggered by contact with acid.

See Ammonium nitrate, Ammonium nitrite

Metal acetylides

Mellor, 1946, Vol. 5, 8482

Caesium acetylide and rubidium acetylide explode in contact with nitric acid, and the sodium and potassium analogues probably react violently.

Metal cyanides

491M, 1975, 268

Explosive reactions.

Metal hexacyanoferrates

Sidgwick, 1950, 1344

The action of 30% nitric acid on hexacyanoferrates(II) or (III) to produce pentacyanonitrosylferrate(II) ('nitroprusside') is violent.

Metals

1. Mellor, 1940, Vol. 2, 470; 1940, Vol. 4, 270, 483; 1941, Vol. 7, 260; 1940, Vol. 9, 627; 1942, Vol. 12, 32, 188

2. Pascal, 1956, Vol. 10, 504

3. Condike, G. F., *Chemistry*, 1974, **47**(4), 27–28

4. McGarman, A. R., *Chemistry*, 1974, **47**(11), 27–28

MRH Calcium 9.33/61, lithium 15.27/40, magnesium 11.96/48, manganese 4.88/59, nickel 2.64/70, sodium 5.48/64, zinc 3.60/72

Bismuth powder glows red-hot in contact with fuming nitric acid, while the molten metal (271°C) explodes in contact with conc. acid. Powdered germanium reacts violently with the latter, and lithium ignites. Manganese powder incandesces and explodes feebly with nitric acid, and sodium ignites with nitric acid of density above 1.056. Titanium alloys form an explosive deposit with fuming nitric acid. Although uranium powder reacts vigorously with red fuming nitric acid, under some conditions explosive deposits may be formed. Addition of conc. acid to molten zinc (419°C) causes it to incandesce, and magnesium burns brilliantly in nitric acid vapour [1]. Antimony may be attacked violently by fuming nitric acid [2]. The experimental quantities of tin and nitric acid specified for a laboratory demonstration [3] are likely to lead to formation of the explosive nitrate oxide [4].
See Tetrahydroxotritin(2+) nitrate, Silver nitrate: Titanium

See Copper, above

Metal salicylates

Belcher, R., private comm., 1968

Metal salicylates are occasionally incorporated into mixtures of 'unknowns' for qualitative inorganic analysis. During the conventional group separation, organic radicals are removed by evaporation with nitric acid. When salicylates are present, this can lead to formation of trinitrophenol through nitration and decarboxylation. This may react with any heavy metal ions present to form unstable or explosive picrates, if the evaporation is taken to dryness. The MAQA alternative scheme of analysis obviates this danger.

See Lead(II) picrate

Metal thiocyanate

MCA Case History No. 853

MRH Potassium thiocyanate 4.31/42

HNO₃

When the (unspecified) thiocyanate solution was pumped through an 80 mm pipeline containing nitric acid, a violent explosion occurred. This was confirmed experimentally and attributed to the redox reaction between the reducing solution and the oxidant acid.

See other REDOX REACTIONS

Methanol, Sulfuric acid

Anon., *Chem. & Ind.*, 1986, 725

A nitration reactor was charged with methanol which was wrongly labeled as xylene. Addition of nitrating acid mixture led to a runaway exothermic reaction which eventually ruptured the reactor as well as the bursting disk, allowing 2700 l of mixed acid to spray out.

See Alcohols, above

See related NITRATION INCIDENTS

2-Methoxyethylbenzene, Sulfuric acid

Urban, P. G., personal experience

This substrate, dissolved in 3 molar equivalents of sulfuric acid, was batch nitrated by slow addition of 1.05 moles of concentrated nitric acid at 5–20°C followed by a stir-out and quenching into water. No problems were seen in the laboratory but on pilot plant ($\frac{1}{2}$ kilomole) there was twice a sharp exotherm 2 $\frac{1}{2}$ hours after completion of addition, blowing the bursting disk with gas evolution, leaving 4-nitrobenzoic acid as the (minor) isolable product. With good thermal insulation and poor agitation it was possible to reproduce this in the lab. No evidence of dinitration could be found and mononitration appeared almost instantaneous. Reduction of the nitric acid to 0.99 equivalents prevented recurrence, even on a much larger scale with day long delays, while not changing the near quantitative yield.

The end point corresponded to generation of sulfuric acid monohydrate; the experimenter subsequently adopted the rule of thumb that oxidative side reactions do not occur while the availability of water is less than this, and that dinitration side-products are a consequence of charging substrate, traditionally but irrationally, to nitration mixture.

See related NITRATION INCIDENTS

2-Methylbenzimidazole, Sulfuric acid

Zmojdzin, A. *et al.*, Ger. Offen. 2 310 414, 1974

A safe method of preparing 5-nitro-2-methylbenzimidazole involves preliminary addition of the heterocycle to nitric acid (*d* 1.40) with subsequent addition of sulfuric acid, keeping the temperature below 110°C.

See related NITRATION INCIDENTS

4-Methylcyclohexanone

Dye, W. T., *Chem. Eng. News*, 1959, **37**, 48

Oxidation of 4-methylcyclohexanone by addition of nitric acid at about 75°C caused a detonation to occur. These conditions had been used previously to oxidise the corresponding alcohol, but although the ketone is apparently an intermediate in oxidation of the alcohol, the former requires a much higher temperature to start

and maintain the reaction. An OTS report, PB73591, mentions a similar violent reaction with cyclohexanone.

See Acetone, etc., above

Methylthiophene

Rüst, 1948, 318

During the nitration of methylthiophene, direct liquid contact caused ignition to occur, so air saturated with the organic vapour was passed into the cooled conc. acid. After a while, the whole apparatus exploded violently, probably owing to ignition of the air–fuel mixture being passed in.

See other NITRATION INCIDENTS *SEE* THIOPHENE, BELOW

Molybdenum nitride

See Molybdenum nitride: Nitric acid

Naphthalene-1-sulfonic acid, Sulfuric acid

Anon. *Loss Prev. Bull.*, 1994, (116), 21

Production of dye intermediates by nitration of the sulfonic acid was conducted under automatic control, nitric acid being added automatically so long as the temperature was below 30°C. There was no other control on nitric acid addition. The agitation and probably also the thermocouple reading being inadequate, excess nitric acid appears to have charged, layered, then reacted, blowing apart the vessel.

See other AGITATION INCIDENTS

Nitroaromatics

1. Urbanski, 1967, Vol. 3, 290

2. Raikova, V. M., *Chem. Abs.*, 1983, **98**, 5958

MRH Nitrobenzene 6.32/29, nitroxylyene 6.23/25, dinitrobenzene

5.02/99+, 2,4-dinitrotoluene 4.56/99+

A series of mixtures of nitric acid with one or more of mono- and di-nitrobenzenes, di- and tri-nitrotoluenes have been shown to possess high-explosive properties [1]. Critical detonation diameters for mixtures of nitric acid with dinitrotoluene and TNT have been determined [2].

Nitrobenzene, Sulfuric acid

Fritz, E. J., *Loss Prev.*, 1969, **3**, 41–44

Failure of the agitator during addition of mixed nitrating acid allowed unreacted reagents to build up in the reaction system which, owing to absence of the usual reaction exotherm, became under-cooled. Application of heat and agitation caused a runaway reaction, terminating in explosion, to occur. Laboratory-scale repetition of this sequence showed an exotherm to 200°C in 0.1 s, a rate of almost 2000°/s. Operating improvements are detailed.

See other AGITATION INCIDENTS, NITRATION INCIDENTS, RUNAWAY REACTIONS

Nitrobenzene, Water

1. Anon., *J. R. Inst. Chem.*, 1960, **84**, 451

2. Van Dolah, R. W., *Loss. Prev.*, 1969, **3**, 32

HNO₃

A plant explosion involved a mixture of nitrobenzene, nitric acid and a substantial quantity of water. Detonation occurred with a speed and power comparable to TNT. This was unexpected in view of the presence of water in the mixture [1]. The later reference deals with a detailed practical and theoretical study of this system and determination of the detonability limits and shock-sensitivity. The limits of detonability coincided with the limits of miscibility over a wide portion of the ternary composition diagram. In absence of water, very high sensitivity (similar to that of glyceryl nitrate) occurred between 50 and 80% nitric acid, the stoichiometric proportion being 73% [2].

See Nitroaromatics, above

Nitromethane

MRH 6.19/62

Olah, G. A. *et al.*, *Org. Synth.*, 1957, **47**, 60

Mixtures are extremely explosive.

1-Nitronaphthalene, Sulfuric acid

Biasutti, 1981, 83

Towards the end of nitration of mononitronaphthalene to trinitronaphthalene, a drain valve became blocked with tarry solid. Raising the temperature to 60°C to melt the obstruction led to separation of more solid, failure of the agitator, then a runaway reaction and detonation.

See other AGITATION INCIDENTS, NITRATION INCIDENTS, RUNAWAY REACTIONS

Non-metal hydrides

1. Mellor, 1946, Vol. 5, 36; 1940, Vol. 6, 814; 1939, Vol. 9, 56, 397

2. Hofmann, A. W., *Ber.*, 1870, **3**, 658–660

3. Pascal, 1956, Vol. 10, 505

Arsine, phosphine and tetraborane(10) are all oxidised explosively by fuming nitric acid, while stibine behaves similarly with the conc. acid [1]. Hydrogen iodide, hydrogen selenide, hydrogen sulfide and phosphine all ignite when the fuming acid is dripped into the gas [2]. Hydrogen telluride ignites with cold conc. acid, sometimes exploding [3].

See Ammonia, above; and Phosphine derivatives, below

Non-metals

1. Mellor, 1946, Vol. 5, 16; 1940, Vol. 8, 787, 845

2. Bailar, 1973, Vol. 1, 1337

3. Pascal, 1956, Vol. 10, 504

4. *MCA Case History No. 1969*

Boron (finely divided forms) reacts violently with conc. acid and may attain incandescence. The vapour of phosphorus, heated in nitric acid in presence of air, may ignite. Boron phosphide ignites with the conc. acid [1]. Silicon crystallised from its eutectic with aluminium reacts violently with conc. acid [2], arsenic may react violently with the fuming acid, and finely divided carbon similarly with conc. acid [3]. Use of conc. acid to clean a stainless steel hose contaminated with phosphorus led to an explosion [4].

Oleoyl chloride

Nolan, 1983, Case history 62

3-Nitro-4-cresol was to be manufactured by adding 4-toluidine to nitric acid. One carboy of nitric acid was charged into the reactor, but the next one charged actually contained oleoyl chloride in error. The violent reaction ensuing ruptured the reactor and there was a fire.

See related NITRATION INCIDENTS

Organic diacids

See Cyclohexanol, Cyclohexanone, above

Organic materials

1. Bowen, H. J. M., private comm., 1968
2. Turnbull, A. J., *Chem. Brit.*, 1979, **15**, 377
3. Williams, E. V., *J. Food Technol.*, 1978, **13**, 167
4. Everson, R. J., *Chem. Eng. News*, 1980, **58**(11), 5
5. Coleman, C. B., *Chem. Eng. News*, 1980, **58**(11), 5
6. Katz, S. A., *Chem. Eng. News*, 1980, **58**(17), 51
7. Borggaard, O. K. *et al.*, *J. Assoc. Offic. Anal. Chem.*, 1982, **65**, 762–763
8. Anon., *Inst. Univ. Safety Officers Bull.*, 1987, (8), 9
9. Ward, N. I., *Internat. Analyst*, 1987, **1**(2), 7

When 16 M (70%) nitric acid was poured down a sink without diluting water, interaction with (unspecified) organic material in the trap caused a delayed explosion [1]. Digestion of biological materials with nitric acid in Teflon-lined bombs has led to several violent explosions. A suggested improvement was to leave the sample in contact with nitric acid overnight before sealing and heating the bomb [2]. An alternative procedure of digesting samples with nitric acid at 20°C for 24 h in polystyrene tubes with hand-tightened polythene screw caps [3] also led to explosions when generous samples of dry-powdered food products were digested [2]. Digestion of 0.8 g of dry dog-food with 5 cc of nitric acid at 130°C in Teflon lined stainless bombs led to explosions [4], or of 0.2 g of sewage sludge with 2.5 cc of acid [5]. The amount of organic material is normally restricted to 0.1 g in the bomb manufacturer's instructions [6]. Note that 0.1 g of carbon on oxidation with nitric acid gives a minimum volume of 540 cc of gas measured at 1 bar/120°C, so a static pressure of many bar would be attained in a 10–20 cc bomb. The hazards of digesting organic samples with nitric acid in closed systems for elemental analysis have been stressed. Digestion in an open flask may be a safer alternative in many cases [7]. In preparation for the analysis of powdered milk for trace metals, 1 g portions were heated at 80°C with 5 ml of conc. nitric acid in 25 ml Teflon lined stainless digestion vessels made to a design maximum of 500 bar, with maximum working pressure limited to 125 bar. With all 3 samples explosions occurred, rupturing the digestion vessels and causing local damage. In one account of the incident, the explosions were attributed to formation of glyceryl trinitrate [8], but this is extremely unlikely, as its formation normally requires presence of conc. sulfuric acid as dehydrating agent. In a second account the correct explanation of a runaway exothermic oxidation reaction of an excessive sample

HNO₃

by undiluted nitric acid accompanied by gas formation was deduced. The strong recommendation to limit sample size to 0.1 g maximum [9] coincides with that made 7 years previously [6].

See other GAS EVOLUTION INCIDENTS

Organic materials, Perchloric acid

See Perchloric acid: Nitric acid, etc.

Organic materials, Potassium chlorate

See Potassium chlorate: Nitric acid, etc.

Organic materials, Sulfuric acid

Anon., *ABCM Quart. Safety Summ.*, 1934, **5**, 17

Use of the mixed conc. acids to dissolve an organic residue caused a violent explosion. Nitric acid is a very powerful and rapid oxidant and may form unstable fulminic acid or polynitro compounds under these conditions.

Other reactants

Yoshida, 1980, 188–189

MRH values calculated for 28 combinations, largely with oxidisable materials, are given.

Phenylacetylene, 1,1-Dimethylhydrazine

Spengler, G. *et al.*, *Sci. Tech. Aerospace Rept.*, **2**(17), 2392, Washington, NASA, 1964

Phenylacetylene does not itself ignite on contact with nitric acid, but addition of 1,1-dimethylhydrazine renders it hypergolic.

See Hydrocarbons, above

Phosphine derivatives

MRH Ethylphosphine 6.15/23

1. von Schwartz, 1918, 325

2. Mellor, 1947, Vol. 8, 827, 1041

3. Graham, T., *Trans. Roy. Soc. Edinburgh*, 1835, **13**, 88

Phosphine ignites in conc. nitric acid and addition of warm fuming nitric acid to phosphine causes explosion [1]. Phosphonium iodide ignites with nitric acid, and ethylphosphine explodes with fuming acid [2]. Tris(iodomercuri)phosphine is violently decomposed by nitric acid or aqua regia [3].

Phosphorus compounds

Mellor, 1947, Vol. 8, 1061; 1971, Vol. 8, Suppl. 3, 348, 335, 373

Tetraphosphorus tetraoxide trisulfide or neodymium phosphide are violently oxidised, nickel tetraphosphide ignites with the fuming acid, and tetraphosphorus diiodo triselenide reacts explosively with nitric acid.

See 2-Alkoxy-1,3-dithia-2-phospholane, above

Phosphorus halides

Mellor, 1947, Vol. 8, 827, 1004, 1038

Tetraphosphorus iodide ignites in contact with cold conc. nitric acid, and phosphorus trichloride explodes with nitric (or nitrous) acid.

Phthalic anhydride, Sulfuric acid

1. Tyman, J. H. P. *et al.*, *Chem. & Ind.*, 1972, 664
2. Bentley, R. K., *Chem. & Ind.*, 1972, 767
3. Bretherick, L., *Chem. & Ind.*, 1972, 790

Attempts to follow a published method for nitrating phthalic anhydride in sulfuric acid at 80–100°C with fuming nitric acid caused an eruptive decomposition to occur after 2 h delay [1]. The hazard can be eliminated by use of a smaller excess of nitrating acid at 55–65°C [2]. Acyl nitrates were suggested as a possible cause of the delayed eruption [3].

See Sulfuric acid, Terephthalic acid, below

Polyalkenes

1. Marsh, J. R., *Chem. & Ind.*, 1968, 1718
2. Dabeka, R. W. *et al.*, *Anal. Chem.*, 1976, **48**, 2048
3. Kaufman, J. A., private comm., 1987
4. Weker, R. A. *et al.*, *Chem. Eng. News*, 1995, **73**(25), 4

Fuming nitric acid had seeped past the protective polytetrafluoroethylene liner inside the polyethylene or polypropylene screw cap and attacked the latter, causing pressure build-up in the glass bottle [1]. Polypropylene bottles are unsuitable for long term storage of nitric acid, because slow embrittlement and cracking occur [2]. A plastic bottle (polythene?) used to store conc. nitric acid collapsed while being carried [3]. Digestion of airborne particulate metals caught on a PTFE filter by nitric acid in a microwave oven led to explosion. The filter membrane was mounted in a poly(methylpentene) support, which was the cause of the violent reaction. Use of either an all PTFE filter, or cutting away the support ring, is advisable [4].

Poly(dibromosilylene)

See Poly(dibromosilylene): Oxidants

Poly(ethylene oxide) derivatives

1. Corby, M. P., *Chem. Brit.*, 1975, **11**, 334; private comm., 1975
2. Corby, M. P., *Chem. Brit.*, 1975, **11**, 456

A mixture of nitric and phosphoric acids (50, 17%, respectively) with a primary alcohol ethoxylate surfactant (0.1%) and water exploded after storage for 7 months in a glass bottle [1]. Progressive hydrolysis under these conditions would be expected to lead to production of the readily oxidised ethylene glycol, and gaseous decomposition products leading to pressure build-up. A general warning against mixing surfactants and oxidising acids is given [2]. The formation of ethylene glycol dinitrate is perhaps unlikely in view of the 33% water content in the mixture.

Polymer (unspecified)

Anon., *Safety Digest Univ. Safety Assoc.*, 1989, **34**, 14

HNO₃

Throat cut, almost fatally, by shards of a test tube from which a polymeric residue was being cleared. Explosion apparently almost immediate; were there also miscible solvent residues?

Poly(silylene)

See Poly(silylene): Oxidants

Polyurethane foam

Anon., *Loss Prev. Bull.*, 1980, (035), 10–11

Mechanical cleaning of a multitubular stainless steel condenser (blocked by a rigid polyurethane foam) by rodding the tubes was laborious, so chemical cleaning with conc. nitric acid was attempted. When the initial vigorous reaction ('fireworks') subsided, owing to crust formation, the rod was again inserted, but a sudden explosion occurred which ruptured the condenser.

See other PLANT CLEANING INCIDENTS

Propiophenone, Sulfuric acid

1. Baker, J. W., *J. Chem. Soc.*, 1932, 1155

2. Dowell, R. I., *Chem. Brit.*, 1981, **17**, 56–57

3. Keneford, J. R., *et al.*, *J. Chem. Soc.*, 1948, 356

During the nitration of propiophenone following a published procedure [1] by adding the ketone to the mixed acids at –5 to 0°C, an uncontrollable exotherm developed and finally accelerated to explosion [2]. The nitration was subsequently effected by an alternative procedure operated at –10 to –5°C [3].

See other NITRATION INCIDENTS

Pyrocatechol

Andrussow, L., *Chim. Ind.* (Paris), 1961, **86**, 542

The phenol is hypergolic with conc. nitric acid, with a 1 ms ignition delay.

Pumps

Anon. *Loss Prev. Bull.*, 1994, 116, 21

Two incidents of pump explosions in chemical plants are reported, both occasioned by nitric acid reaching parts of a pump it should not and there corroding metal or contacting electrics. Proper pump selection, installation and maintenance would have avoided these incidents.

Reducants

A variety of reducants ignite or explode with nitric acid.

See Hydrazine and derivatives, above

Hydrogen sulfide: Oxidants

Non-metal hydrides, above

Potassium phosphinate: Air, or Nitric acid

Sulfur dioxide, below

Resorcinol

Biasutti, 1981, 104

During the preparation of dinitroresorcinol, too low a concentration of nitric acid (82%) led to formation of tarry material, some of which remained in the nitrator discharge line. In the following batch, the tar decomposed in contact with higher strength acid, leading to an explosion.

See other NITRATION INCIDENTS, PLANT CLEANING INCIDENTS

Rubber

1. Read, T. R., *School Sci. Rev.*, 1976, **57**(200), 592
2. Mishra, B. *et al.*, *J. Hazard. Mater.*, 1997, 56, 107

In a demonstration of the corrosive effect of fuming nitric acid on rubber tubing, soon after the initial vigorous reaction had subsided, a small explosion followed by ignition occurred. (The sulfur compounds used to compound the rubber were probably involved) [1]. Lead neoprene rubber gloves, (the lead present as red lead, Pb_3O_4) used in the nuclear industry, may form explosive compounds in contact with nitric acid. It is hypothesised this could be lead fulminate formation. Since mixtures of lead nitrate with organics, or with sulfur from vulcanisers, would be explosive, this seems unduly complicated [2]

See other CORROSION INCIDENTS

Salicylic acid

See Lead(II) picrate

Silicone oil

Phillips, C. V. *et al.*, *Lab. Pract.*, 1981, **30**, 598

A mineral was being leached by heating with 7 M nitric acid in a PTFE-lined bomb heated by immersion in a silicone oil bath, and at 195°C a violent explosion occurred. This was attributed to prior leakage of oil into the pressure vessel, which had been immersed in the oil at 120°C, then allowed to cool, before being heated to the higher temperature. Appropriate precautions are recommended.

Silver buten-3-ynide

See Silver buten-3-ynide: Alone, or Ammonia, etc.

Steel gas cylinder

Anon., *CISHC Chem. Safety Summ.*, 1974–5, **45–46**, 3–4

Conc. nitric acid leaking from a faulty road tanker became partially diluted with water and was prevented from running away along the roadside gully by a full oxygen cylinder in the horizontal position. After several h, the cylinder was sufficiently weakened by corrosion to split open under the internal gas pressure.

See other CORROSION INCIDENTS

Sucrose

Poncini, L., *School Sci. Rev.*, 1981, **62**(200), 515–516

Use of aqueous solutions of sucrose rather than the solid in class demonstrations of preparation of oxalic acid is described as safer.

Sulfur dioxide

Coleman, G. H. *et al.*, *Inorg. Synth.*, 1939, **1**, 55

HNO₃

Presence of dinitrogen tetraoxide appears to be essential to catalyse smooth formation of nitrosylsulfuric acid from sulfur dioxide and nitric acid. In its absence, reaction may be delayed and then proceed explosively.

See other CATALYTIC IMPURITY INCIDENTS

Sulfur halides

Mellor, 1947, Vol. 10, 646, 651–652

Interaction with sulfur dichloride, sulfur dibromide or disulfur dibromide is violent, the hydrogen halide being liberated.

See other GAS EVOLUTION INCIDENTS

Sulfuric acid

1. Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 3

2. *FPA H50*, 1976; *HCS 1980*, 666

The gland of a centrifugal pump being used to pump nitrating acid (nitric:sulfuric, 1:3) exploded after 10 min use. This was attributed to nitration of the gland packing, followed by frictional detonation. Inert shaft sealing material is advocated [1]. General handling precautions for nitrating acid are detailed [2].

See 2,2-Oxybis(ethyl nitrate)

Sulfuric acid, Terephthalic acid

Withers, C. V., *Chem. & Ind.*, 1972, 821; private comm., 1972

During nitration of the acid with fuming nitric acid in oleum, a delayed exotherm increased the temperature after 2 h from 100° to 160°C, causing eruption of the contents. At 120° the delay was 1 h and at 130°C, 30 min.

See Phthalic anhydride, etc., above

See other INDUCTION PERIOD INCIDENTS

Thioaldehydes, or Thioketones

Campaigne, E., *Chem. Rev.*, 1946, **39**, 57

Nitric acid generally reacts too violently with thials or thiones for the reactions to be of preparative interest.

Thiols

See Alkanethiols, above

Thiophene

1. Meyer, V., *Ber.*, 1883, **16**, 1472

2. Babasinian, V. S., *Org. Synth.*, 1944, Coll. Vol. 2, 467

Interaction of thiophene with fuming nitric acid is very violent if uncontrolled, extensive oxidation occurring [1]. Use of a diluent and close control of temperature is necessary for preparation of nitrothiophene [2].

See Methylthiophene, above

1,3,5-Triacetylhexahydro-1,3,5-triazine, Trifluoroacetic anhydride

1. Albright and Hanson, 1976, 327

2. Bedford, C. D., *Chem. Eng. News*, 1980, **58**(35), 33, 43

3. Gilbert, E. E., *Chem. Eng. News*, 1980, **58**(40), 5
4. Bretherick, L., *Chem. Eng. News*, 1981, **59**(14), 59

During the attempted nitrolysis of the triacetyl compound to the 1-acetyl-3,5-dinitro compound with 99% nitric acid in trifluoroacetic anhydride at 30°C, following a general procedure [1], a violent explosion occurred on the 1 g scale [2]. This was ascribed to the formation of acetyl nitrate, expected to be formed under the reaction conditions [3,4]. Caution with nitrolysis of any acetyl compound is urged [3].

See Acetic anhydride, above; Triazine, etc., below

Triazine, Trifluoroacetic anhydride

491M, 1975, 272

N-Nitration of triazine with 99% nitric acid in trifluoroacetic anhydride at 36°C proceeded explosively.

Tributyl phosphate

See NUCLEAR WASTES

Triethylgallium etherate

See Triethylgallium

2,4,6-Trimethyltrioxane

Muir, G. D., private comm., 1968

Oxidation of the trioxane ('paraldehyde') to glyoxal by action of nitric acid is subject to an induction period, and the reaction may become violent if addition of the trioxane is too fast. Presence of nitrous acid eliminates the induction period.

See other INDUCTION PERIOD INCIDENTS

Turpentine

Hermoni, A., *Appl. Chem.*, 1958, **8**, 670–672

Turpentine and fuming nitric acid do not ignite on contact in absence of added catalysts (fuming sulfuric acid, iron(III) chloride, ammonium metavanadate or copper(II) chloride).

See Hydrocarbons, above

Uranium disulfide

Sidgwick, 1950, 1081

Interaction is violent.

Wood

1. Personal experience, 1974
2. Anon., *Chem. Eng. News*, 1975, **53**(11), 14
3. *MCA Case History No. 1797*
4. Anon., *Chem. & Ind.*, 1994, (6), 212

A cracked winchester of conc. acid leaked into sawdust packing and caused a fire [1]. A similar incident was involved in a freight-plane crash [2]. Fuming acid, leaking from a cracked bottle, ignited a wooden truck [3]. A retrospective survey on nitric acid packed in sawdust reveals an incident before 1894 [4].

See other PACKAGING INCIDENTS

Zinc ethoxide

See Zinc ethoxide: Nitric acid

See other INORGANIC ACIDS, OXIDANTS

4437. Peroxonitric acid

[26604-66-0]

HNO₄

O₂NOOH

1. Schwarz, R., *Z. Anorg. Chem.*, 1948, **256**, 3

2. Kenley, R. A. *et al.*, *J. Amer. Chem. Soc.*, 1981, **103**, 2205

The pure peroxyacid, prepared at -80°C , decomposes explosively at -30°C . Solutions in acetic acid or water of below the limiting concentration (corresponding to a stoichiometric mixture of 70% aqueous nitric acid and 100% hydrogen peroxide) are stable, while those above the limit decompose autocatalytically, eventually exploding [1]. Explosion of the vapour when passed into a mass spectrometer inlet at 427°C (but not at 327°) was noted [2].

See other PEROXOACIDS

4438. Nitrosylsulfuric acid

[7782-78-7]

HNO₅S

O:NOSO₂OH

Preparative hazard

See Nitric acid: Sulfur dioxide

6-Chloro-2,4-dinitroaniline

See 2-Chloro-4,6-dinitroaniline: Nitrosylsulfuric acid

Dinitroaniline

Anon., *BCISC Quart. Safety Summ.*, 1970, **41**, 28

During plant-scale diazotisation of a dinitroaniline hydrochloride, local increase in temperature, owing to high concentration of the reaction mixture, caused a violent explosion.

See Nitric acid: 4-Chloro-2-nitroaniline

See other INORGANIC ACIDS, NITROSO COMPOUNDS, OXIDANTS

4439. Lead(II) imide

[12397-26-1]

HNPb

Pb:NH

Alone, or Acids, or Water

Mellor, 1940, Vol. 8, 265

It explodes on heating, or in contact with water or dilute acids.

See other N-METAL DERIVATIVES

4440. Phospham
[33849-97-1]

HN₂P

(N:PNH)_n

Hydrogen sulfide

Mellor, 1940, Vol. 8, 270

The solid produced by interaction of phospham and hydrogen sulfide at red heat is probably a trimeric triphosphatriazine such as phospham. The solid ignites in slightly warm air or in dinitrogen tetraoxide, and is violently oxidised by nitric acid.

Oxidants

Mellor, 1940, Vol. 8, 269–270

Interaction with copper(II) oxide or mercury(II) oxide proceeds incandescently. Mixtures with a chlorate or nitrate explode on heating. Phospham ignites in dinitrogen tetraoxide.

See related NON-METAL HYDRIDES

4441. Hydrogen azide (Hydrazoic acid)
[7782-79-8]

HN₃

HN₃

1. Smith, 1966, Vol. 2, 214
2. Bowden, F. P. *et al.*, *Endeavour*, 1962, **21**, 121
3. Audreith, L. F. *et al.*, *Inorg. Synth.*, 1939, **1**, 77
4. Kemp, M. D., *J. Chem. Educ.*, 1960, **37**, 142
5. Birkofer, L. *et al.*, *Org. Synth.*, 1970, **50**, 109
6. Shapiro, E. L., *Chem. Eng. News*, 1974, **52**(2), 5
7. Chandross, E. A. and Gunderson, H., *Chem. Eng. News*, 1974, **52**(24), 5
8. Sood, R. K. *et al.*, *J. Therm. Anal.*, 1981, **20**, 491–493

Hydrogen azide is quite safe in dilute solution, but is violently explosive and of variable sensitivity in the concentrated (17–50%) or pure states. Wherever possible, a low-boiling solvent (ether, pentane) should be added to its solutions to prevent inadvertent concentration by evaporation and recondensation. If this is not possible, no unwetted part of apparatus containing its solutions should be kept at a temperature appreciably below the boiling point (35°C) of the pure acid. The pure acid has often been isolated by distillation, but appears to undergo rapid sensitisation on standing, so that after an hour, faint vibrations or speech are enough to initiate detonation [1]. The solid acid (–80°C) is also very unstable [2]. Preparative procedures have been detailed [3]. It is readily formed on contact of hydrazine to its salts with nitrous acid or its salts. A safe procedure for the preparation of virtually anhydrous hydrogen azide has been described [4]. Trimethylsilyl azide serves as a safe and stable substitute for hydrogen azide in many cases [5]. A laboratory explosion [6] seems likely to have been caused by the use of a huge (90-fold) excess of azide in too-concentrated solution and at too low an ambient temperature, leading to condensation of highly concentrated hydrogen azide. This

penetrated into a ground-glass joint, and explosion was initiated on removing the stopper [7]. It is claimed that it may be distilled without fear of explosion if the trapping of the liquid azide in ground joints is prevented by thorough pregreasing of all joints [8].

As the lowest MW azide, hydrogen azide is extremely endothermic (ΔH_f° (g) +294.1 kJ/mol, 6.83 kJ/g).

See Methyl azide, also 4-Chlorophenylisocyanate, also Nitrous acid: A semicarbazone, etc.

See other GLASS INCIDENTS

Heavy metals

1. Napier, D. H., private comm., 1972
2. Cowley, B. R. *et al.*, *Chem. & Ind.*, 1973, 444
3. Mellor, 1967, Vol. 8, Suppl. 2, 4

Great care is necessary to prevent formation of explosive heavy metal azides from unsuspected contact of hydrogen azide with heavy metals. Interaction of hydrazine and nitrite salts in a copper drainage system caused formation and explosion of copper azide deposits [1]. Use of a brass water-pump and vacuum gauge during removal of excess hydrogen azide under vacuum formed deposits which exploded when the pump and gauge were handled later [2]. Raney nickel catalyses the vigorous decomposition of solutions of hydrogen azide [3].

See Cadmium azide (reference 2), Sodium azide: Heavy metals

See other ENDOTHERMIC COMPOUNDS, INORGANIC ACIDS, NON-METAL AZIDES

4442. Dinitramine (Dinitramide)

[115045-20-4]

HN₃ O₄

HN(NO₂)₂

1. Borman, S., *Chem. Eng. News*, 1994, **72**(3), 18
2. Dawe, J. R. *et al.*, *Proc. Int. Pyrotech. Semin.*, 1998, **24th**, 789

Salts of the acidic dinitramine find use as oxidants in propellant formulations. The dinitramine itself is explosively unstable, though more stable than nitramide [1]. Although described as relatively stable the sodium salt has an autoignition temperature of 123°C, the potassium 140°C [2].

See Nitric amide

See other N-NITRO COMPOUNDS

4443. Pentazole

[289-19-0]

HN₅

Sorbe, 1968, 140

Pentazole (98.7% nitrogen) and its compounds are explosive.

See other HIGH-NITROGEN COMPOUNDS

See related TETRAZOLES

4444. Sodium hydride
[7646-69-7]

HNa

NaH

1. Plesek, J. *et al.*, *Sodium hydride* (Engl. transl., Jones, G.) 5, 8, London, Iliffe, 1968
2. Bigou, F., personal communication, 1993
3. Gibb, T. R. P., US Pat. 2 702 281, 1955
4. Urben, P. G., personal experience

Commercial sodium hydride may contain traces of sodium which render it spontaneously flammable in moist air, or air enriched with carbon dioxide. The very finely divided dry powder ignites in dry air. Dispersions of the hydride in mineral oil are safe to handle. All normal extinguishers are unsuitable for solid sodium hydride fires: powdered sand, ashes or sodium chloride are suitable [1]. Commercial oil-coated sodium hydride was washed with dichloromethane to remove the oil. During later charging to a reactor containing more dichloromethane and an involatile substrate the hydride ignited. Ignition was probably by traces of moisture, but combustion seems to have continued in an atmosphere of hydrogen and dichloromethane vapour. It seems inadvisable to mix sodium hydride with most chlorinated solvents. Although a protective coating of non-hydridic material may prevent immediate reaction, it is thermodynamically capable of reacting energetically with halocarbons [2].

Many acylation reactions of esters using sodium hydride as base appear autocatalytic, with considerable potential for runaway, since the active base in solution is an alkoxide and the alcohol is a product of reaction [4]. A safe form of sodium hydride (as a solid solution in a halide) for large-scale industrial use has been claimed [3].

See Sodium: Halocarbons

Acetylene

Mellor, 1940, Vol. 2, 483

Dry acetylene does not react with sodium hydride below 42°C, but in presence of moisture, reaction is vigorous even at -60°C.

Diethyl succinate, Ethyl trifluoroacetate

Harris, A. R. *et al.*, *Chem. Brit.*, 1983, **19**, 645

Acylation of diethyl succinate by ethyl trifluoroacetate in presence of sodium hydride and in absence of a solvent is hazardous, fire or explosion occurring on 2 occasions some 10–20 min after adding a little of the succinate to the hydride–trifluoroacetate premixture at 60°C. Presence of a solvent appears to eliminate the hazard.

See Ethyl trifluoropropionate, below

Dimethylformamide

1. Buckley, J. *et al.*, *Chem. Eng. News*, 1982, **60**(28), 5
2. De Wall, G. *Chem. Eng. News*, 1982, **60**(37), 5, 43

HNa

A mixture of the hydride and solvent was heated to, and held at 50°C. An exothermic reaction, which increased the temperature to 75°C, could not be controlled by external cooling, and the pilot-scale reactor contents erupted. It was later found that an exotherm began to develop at 26° (or at 40–50° with carefully dried solvent), and the subsequent reaction accelerated rapidly. Avoidance of holding mixtures hot is recommended, particularly when scaling up reactions. Similar behaviour was seen with dimethylacetamide [1]. A further similar plant-scale incident was reported, with onset of the exotherm at 40°C, followed by self-heating to 100°C in 10 min, even with maximum cooling applied [2].

See other SELF-ACCELERATING REACTIONS

Dimethyl sulfoxide

See Dimethyl sulfoxide: Sodium hydride

Ethyl 2,2,3-trifluoropropionate

Bagnall, R. D., private comm., 1972

The ester decomposes violently in presence of sodium hydride, probably owing to hydride-induced elimination of hydrogen fluoride and subsequent exothermic polymerisation.

See Methyl trichloroacetate: Trimethylamine

Glycerol

Unpublished observations, 1956

Exothermic interaction of granular hydride with undiluted (viscous) glycerol with inadequate stirring caused charring to occur. Dilution with THF to reduce viscosity and improve mixing prevented local overheating during formation of monosodium glyceroxide.

Halogens

MRH Chlorine 3.10/30

Mellor, 1940, Vol. 2, 483

Interaction with chlorine or fluorine is incandescent at ambient temperature, and with iodine at 100°C.

Other reactants

Yoshida, 1980, 215

MRH values calculated for 8 combinations with various materials are given.

Oxygen

MRH 2.59/31

See Oxygen (Gas): Metal hydrides

Sulfur

MRH 1.25/16

See Sulfur: Sodium hydride

Sulfur dioxide

MRH 1.21/43

Moissan, H., *Compt. rend.*, 1902, **135**, 647

Sulfur dioxide reacts explosively in contact with sodium hydride unless diluted with hydrogen.

Water

1. Braidech, M. M., *J. Chem. Educ.*, 1967, **44**, 321

2. *MCA Case History No. 1587*

Addition of sodium hydride to a damp reactor which had not been purged with inert gas caused evolution of hydrogen and a violent explosion. Solid dispersions of the hydride in mineral oil are more easily and safely handled [1]. When an unprotected polythene bag containing the hydride was moved, some of the powder leaked from a hole, contacted moisture and immediately ignited. Such materials should be kept in tightly closed containers in an isolated, dry location [2].

See other GAS EVOLUTION INCIDENTS

See other METAL HYDRIDES, REDUCANTS

4445. Sodium hydroxide

[1310-73-2]

HNaO

NaOH

(*MCA SD-9*, 1968); *NSC 373*, 1982; *HCS 1980*, 845 (solid), 846 (33% solution);
RSC Lab. Hazards Safety Data Sheet No. 36, 1985

Aluminium, Arsenical materials

See Aluminium: Arsenic trioxide, etc.

Ammonia, Silver nitrate

See Silver nitrate: Ammonia, Sodium hydroxide

1,4-Benzenediol

See 1,4-Benzenediol: Sodium hydroxide

N,N'-Bis(2,2,2-trinitroethyl)urea

See N,N'-Bis(2,2,2-trinitroethyl)urea: Sodium hydroxide

Bromine

See Bromine: Sodium hydroxide

4-Chlorobutyronitrile

See 4-Chlorobutyronitrile: Sodium hydroxide

Chloroform, Methanol

MRH Chloroform 2.05/43

See Chloroform: Sodium hydroxide, etc.

4-Chloro-2-methylphenol

See 4-Chloro-2-methylphenol: Sodium hydroxide

Cinnamaldehyde

See Cinnamaldehyde: Sodium hydroxide

Cyanogen azide

See Cyanogen azide: Sodium hydroxide

Diborane

See Diborane: Octanal oxime, etc.

2,2-Dichloro-3,3-dimethylbutane

See 2,2-Dichloro-3,3-dimethylbutane: Sodium hydroxide

Glycols

Cardillo, P. *et al.*, *Chim. e Ind.* (Milan), 1982, **64**, 781–784

Various mixtures of the base with ethylene glycol or diethylene glycol when heated in DSC capsules show exothermic decomposition around 230°C with rapid evolution of hydrogen. The exotherms increase with the base:glycol ratio, and the principal hazard arises from the rapid increase of pressure in closed systems caused by hydrogen evolution.

See Sodium 2-hydroxyethoxide

See entries GAS EVOLUTION INCIDENTS, PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Maleic anhydride

MRH 1.67/99+

See Maleic anhydride: Bases, etc.

4-Methyl-2-nitrophenol

See 4-Methyl-2-nitrophenol: Sodium hydroxide

3-Methyl-2-penten-4-yn-1-ol

See 3-Methyl-2-penten-4-yn-1-ol: Sodium hydroxide

Nitrobenzene

See Nitrobenzene: Alkali

Other reactants

Yoshida, 1980, 213

MRH values calculated for 12 combinations with various materials are given.

Sodium tetrahydroborate

See Sodium tetrahydroborate: Alkali

Succinic anhydride

See Succinic anhydride: Sodium hydroxide

Sugars

See SUGARS

1,2,4,5-Tetrachlorobenzene

See 1,2,4,5-tetrachlorobenzene: Sodium hydroxide, Solvent

12,2,2-Trichloroethanol

See 2,2,2-Trichloroethanol: Sodium hydroxide

Trichloroethylene

MRH 2.26/45

See Trichloroethylene: Alkali

Trichloronitromethane

MRH 1.80/46

See Trichloronitromethane: Sodium hydroxide

Water

1. *MCA SD-9*, 1968
2. *Haz. Chem. Data*, 1975, 267
3. *MCA Case History No. 2166*
4. Anon., *Loss Prev. Bull.*, 1987, (078), 26

The heat of solution is very high, and with limited amounts of water, violent boiling or even ignition of adjacent combustibles may occur [1,2]. The case history describes the disposal of open ended drums of waste caustic soda by putting into an empty hopper and sluicing out the solid alkali with a hot water jet. When the outlet became blocked, the hopper partly filled with hot water, and addition of a further drum caused a violent eruption to occur [3]. A tank holding 73% caustic soda solution at 121°C was found to be leaking and some of the contents were pumped out. A residue of 50 m³ was to be diluted with water before removal and 100 m³ of hot water was introduced via a hose into the top of the tank, where it formed a top layer. A compressed air lance was then introduced to mix the contents. After 30 min. a violent eruption occurred as the layers suddenly mixed to give a very large exotherm, and the steam so produced buckled the top of the tank [4].

See Potassium hydroxide: Water

Zinc

See Zinc: Sodium hydroxide

Zirconium

See Zirconium: Oxygen-containing compounds

See other INORGANIC BASES

4446. Sodium hydrogen sulfate

[7681-38-1]

NaOSO₂OH

HNaO₄S

Acetic anhydride, Ethanol

See Acetic anhydride: Ethanol etc.

Calcium hypochlorite

See Calcium hypochlorite: Sodium hydrogen sulfate

See other METAL OXONON-METALLATES *See related* INORGANIC ACIDS

4447. Sodium hydrogen xenate
[73378-53-7], sesquihydrate **HNaO₄Xe**
NaOXe(O)₂OH.1.5H₂O

See other XENON COMPOUNDS

4448. Rubidium hydrogen xenate
[73378-55-9], sesquihydrate **HO₄RbXe**
RbOXe(O)₂OH.1.5H₂O

See other XENON COMPOUNDS

4449. 'Solid Phosphorus hydride'
[] **HP₂**
(Indeterminate structure)

1. Mellor, 1940, Vol. 8, 851

2. Sidgwick, 1950, 730

This material (possibly phosphine adsorbed on phosphorus, and produced by decomposition of diphosphane in light) ignites in air, on impact, or on sudden heating to 100°C [1,2].

See other NON-METAL HYDRIDES

4450. Rubidium hydride
[13446-75-8] **HRb**
RbH

Acetylene

Mellor, 1940, Vol. 2, 483

In presence of moisture, interaction of the hydride and acetylene is vigorous at -60°C, while in dry acetylene, reaction only occurs above 42°C.

Oxygen

See Oxygen (Gas): Metal hydrides

Water

Mellor, 1963, Vol. 2, Suppl. 2.2, 2187

Interaction with water is too violent to permit of safe use of the hydride as a drying agent. When dispersed as a solid solution in a metal halide, it can be used as a drying or reducing agent.

See other METAL HYDRIDES

4451. Silicon monohydride (Silyldiyne)

[13774-94-2]

(HSi)_n**(SiH)_n**

Alkali

Stock, G. *et al.*, *Angew. Chem.*, 1956, **68**, 213

The polymeric hydride is relatively stable to water, but reacts violently with alkali, evolving hydrogen.

See other NON-METAL HYDRIDES**4452. 'Unsaturated' Silicon hydride**

[]

(H_{1.5}Si)_n**(SiH_{1.5})_n**

Bailar, 1973, Vol. 1, 1350

The polymeric material (composition varies from SiH_{1.42} to SiH_{1.56}) burns with a shower of sparks if heated in air.*See other* NON-METAL HYDRIDES**4453. Hydrogen (Gas)**

[1333-74-0]

H₂**H₂***FPA H20*, 1974; *HCS 1980*, 544 (cylinder)

1. Pignot, A., *Chaleur Ind.*, 1939, **20**, 251–259
2. Anon., *Loss Prev. Bull.*, 1977, (015), 2
3. Fenning, R. W. *et al.*, *Engineering*, 1930, **130**, 252
4. Messelt, W., *Engineering*, 1922, **113**, 502
5. Neer, M. E., *Diss. Abstr. Int. B*, 1972, **33**, 686
6. Carhart, H. W., *et al.*, *AD Rep. No. 781862/IGA*, 1–39, Richmond (Va.), USNTIS, 1974
7. Neer, M. E., *Amer. Inst. Aero. Astron. Paper 74-1159*, 1974, 1–7
8. *NASA Tech. Memo. TMX-71565*, Richmond (Va.), USNTIS, 1971
9. Andrew, H. G., *School Sci. Rev.*, 1979, **61**(215), 292–294
10. Boeck, H., *Chem. Sich.*, 1982, **29**(4), 158–160
11. McCue, J. C., *Altern. Energ. Sources*, 1980, 1983, **3**, 419–427
12. Hydrogen Safety Manual, Balthasar, W. *et al.*, Luxembourg, Comm. Euro. Commun., 1983
13. Angus, H. C., *Chem. & Ind.*, 1984, 68–72
14. Lee, J. H. S. *et al.*, *Adv. Heat Transfer*, 1997, **25**, 59

Several of the combustion-related properties of hydrogen in air, such as its wide flammability limits (4–75 vol%), wide detonation range (20–65 vol%), very low spark ignition energy (0.02 mJ), high heat of combustion (121 kJ/g) and high flame temperature (2050°C) combine to emphasise the high fire-related hazards of the

lightest element. Sudden release of hydrogen into the atmosphere from storage at above 79 bar may cause spontaneous ignition, owing to the inverse Joule-Thompson effect [1]. This, however, does not always occur in practice, largely because of the low value of the effect (1.8°C/110 bar drop at 20°C; 3.9°/100 bar at 100°), so that ignition is unlikely unless the gas temperature is close to the auto-ignition temperature of 530–590°C [2] (or unless catalytic impurities are involved). Release of hydrogen at 47.5 bar into a vented 17.5 l chromium-plated sphere had caused explosive ignition [3]. Earlier, it had been found that a release of hydrogen for filling balloons would ignite under certain circumstances when the aperture was rusted, a brush discharge then being visible [4]. Spontaneous ignition of flowing hydrogen–air mixtures has been studied [5]. All aspects of hazards involved in the production, storage, handling and use of hydrogen as a Navy fuel are discussed [6]. In a study of the causes of auto-ignition of fast flowing hydrogen–air mixtures, at velocities above 750 m/s ignition occurred at temperatures 400°C below those established for static mixtures [7]. All recorded incidents and accidents involving hydrogen ignition have been reviewed and analysed [8]. Safety aspects of the generation and use of hydrogen in school demonstrations are detailed [9,10]. A compact portable ultrasafe storage unit for hydrogen for laboratory use has been developed, based on a lanthanum–nickel alloy hydride storage capsule, to eliminate hazards associated with use of high pressure storage of hydrogen [11]. A safety manual devoted to hydrogen has been published [12], and safety matters are included in a review of current technological aspects of storage, distribution and compression of hydrogen [13]. A detailed review of combustion and explosion properties of hydrogen, and of associated parameters, with a particular eye to nuclear reactor safety, has been published more recently [14].

Acetylene, Ethylene

Anon., *BCISC. Quart. Safety Summ.*, 1974, **45**, 2–3

In a plant producing 200 kt/a of ethylene from cracked naphtha, acetylene in the product was hydrogenated to ethylene in a catalytic unit operated under conditions mild enough not to hydrogenate ethylene. During a temporary shut-down and probably owing to operating error, the internal temperature in the catalytic unit rose to about 400°C, though the external wall temperature was recorded as excessive at 120°C. Attempts to reduce the temperature by passing in additional ethylene were unsuccessful, as the conditions were now severe enough to hydrogenate ethylene. This exothermic reaction increased the temperature, finally to 950°C, and the extensive cracking to methane, carbon and hydrogen now occurring was accompanied by further pressure increase. Plant failure was followed by an explosion and fire which took 4 days to extinguish, and damage totalled 6 M sterling.

See other HYDROGENATION INCIDENTS

Air

1. Berman, M., *Nucl. Sci. Eng.*, 1986, **93**(4), 321–347
2. Edeskuty, F. J. *et al.*, *Adv. Hydrogen Energy*, 1986, **5** (Hydrogen Energy Prog. 6, Vol. 1), 147–158

3. Cummings, J. C. *et al.*, *Sandia Nat. Lab. Report, SAND-86-0173*, 1987, DE87011907 (USNTIS. Richmond, Va)
4. Mottin, M. C., *Proc. 11th Int. Sympos. Prev. Occup. Risks Chem. Ind.*, 383–385, Heidelberg, ISSA, 1987

Recent large-scale experimental work indicating that transition of hydrogen–air mixtures from deflagration to detonation is considerably affected by geometrical factors and scale, and that transition to detonation may be rather more likely in large-scale incidents than previously supposed [1]. Safety aspects of large-scale combustion of hydrogen–air mixtures, especially in confined volumes, have been investigated [2]. Factors involved in the flame acceleration observed in combustion of hydrogen–air mixtures were investigated experimentally in closed and vented vessels, and the results cast doubt on the validity of existing explosion-venting guidelines [3]. Based on several accidents and on large-scale experimental investigations of unconfined explosions of hydrogen–air mixtures, it was concluded that 0.1% of the energy potentially available was effectively released [4].

Air, Catalysts

Mellor, 1942, Vol. 1, 325, 1937, Vol. 16, 146

Catalytically active platinum and similar metals containing adsorbed oxygen or hydrogen will heat and cause ignition in contact with hydrogen or air, respectively. Nitrogen purging before exposure to atmosphere will eliminate the possibility.

Air, Various vapours

Schumacher, H. J., *Angew. Chem.*, 1951, **63**, 560–561

The effects of the presence of 44 gaseous or volatile materials upon the upper explosion limits of hydrogen–air mixtures have been tabulated.

Benzene, Raney nickel catalyst

See Benzene: Hydrogen, Raney nickel

Calcium carbonate, Magnesium

See Magnesium: Calcium carbonate, Hydrogen

Carbon monoxide, Oxygen

See Oxygen: Carbon monoxide, Hydrogen

Catalyst

Bretherick, L., *CISHC Chem. Safety Summ.*, 1980, **51**, 111

Ignition of hydrogen leaking from a hydrogenation autoclave stirrer gland was attributed to traces of hydrogenation catalyst dust outside the reactor.

Catalyst, 3,4-Dichloronitrobenzene

See 3,4-Dichloronitrobenzene: Catalyst, Hydrogen

Catalysts, Vegetable oils

Smirnov, V. M., *Chem. Abs.*, 1938, **32**, 4368₆

H₂

Flash fires and explosions which frequently occurred on discharge of the hot products of catalytic hydrogenation of vegetable oils were attributed to formation of phosphine from the phosphatides present to a considerable extent in, e.g., rape-seed and linseed oils.

See other HYDROGENATION INCIDENTS

Ethylene, Nickel catalyst

Anon., *Loss Prev. Bull.*, 1977, (015), 4–5

Hydrogen from a naphtha cracker normally contains 10% of methane, practically no ethylene and a little carbon monoxide, and the latter is removed by passage over a heated nickel catalyst when it is hydrogenated to methane. Failure of the ethylene refrigeration compressor led to the presence of 8.5% of ethylene in the hydrogen. Presence of catalyst dust in the inlet tube caused rapid hydrogenation of ethylene to occur in this confined volume and the temperature rapidly increased. A temperature monitor set at 500°C also failed, and when the inlet tube reached 800°C it burst under the prevailing pressure and the hydrogen released immediately ignited. Preventive measures are detailed.

See other HYDROGENATION INCIDENTS

Halogens, or Interhalogens

MRH values show % of halogen

See Bromine: Hydrogen 0.45/99

Bromine fluoride: Hydrogen

Bromine trifluoride: Halogens, etc.

Chlorine: Hydrogen 2.55/97

Chlorine trifluoride: Hydrogen-containing materials 6.82/97

Fluorine: Hydrogen 13.39/95

Iodine heptafluoride: Carbon, etc.

Hydrogen peroxide, Palladium catalyst

See Hydrogen peroxide: Hydrogen, Palladium catalyst

Liquid nitrogen

1. Mel'nik, B. D. *et al.*, *Chem Abs.*, 1963, **59**, 7309g

2. Bohlken, S. F., *Chem. Eng. Progr.*, 1961, **57**(4), 49–52

During the purification of washed hydrogen from cracking of natural gas, cooling with liquid nitrogen can lead to trapping of explosive products from interaction of alkenes in the gas with oxides of nitrogen arising from biologically derived ammonium nitrate or nitrite in the scrubbing water. Various measures to prevent this are discussed. Similar effects may be observed when alkenes are oxidised in presence of nitrogen [1]. Analysis of a similar incident involving hydrogen derived from fuel oil implicated resins derived from interaction of nitrogen oxide (and possibly dinitrogen oxide) in a low-temperature heat exchanger section operating at –130 to –145°C [2].

See Nitrogen oxide: Dienes, Oxygen

Metals

Mellor, 1942, Vol. 1, 327

Lithium, calcium, barium and strontium react readily, sometimes igniting, in hydrogen above 300°C, while sodium and potassium react more slowly to form the hydrides.

Nickel, Oxygen

See Nickel: Hydrogen, Oxygen

2-Nitroanisole

See 2-Nitroanisole: Hydrogen

Oxidants

See Chlorine dioxide: Hydrogen

Copper(II) oxide: Hydrogen

Dichlorine oxide: Oxidisable materials

Difluorodiazene: Hydrogen

Dinitrogen oxide: Hydrogen, etc.

Dinitrogen tetroxide: Hydrogen, etc.

Fluorine: Hydrogen, or: Hydrogen, Oxygen

Fluorine perchlorate: Hydrogen

Iodine heptafluoride: Carbon, etc.

Nitrogen oxide: Hydrogen, etc.

Nitryl fluoride: Non-metals

Oxygen (Gas): Hydrogen

Palladium(II) oxide: Hydrogen

Xenon hexafluoride: Hydrogen

Oxygen, Sulfuric acid

Tabata, Y. *et al.*, *J. Haz. Mat.*, 1987, **17**, 47–59

Three PVC chlorine drying towers in a mercury amalgam cell chlorine plant suddenly exploded violently some time after the AC auxiliary power supply to the plant had failed, while the DC cell and brine feed supply had remained on. The failure of the pumped mercury circulation (and its flow alarm) through the electrolysis cells caused hydrogen (instead of sodium) to be liberated at the cathodes and oxygen at the anodes, so the chlorine gas output fed into the PVC drying towers trickle-fed with conc. sulfuric acid contained hydrogen and oxygen also. Ignition of the explosive mixture in the towers was caused by static sparks generated by the falling sulfuric acid drops: this was confirmed by a measurement of a static negative potential of 5kV inside and near the base of the towers. Other possible hazardous malfunctions in amalgam chlorine cells and in non-conductive drying towers are described, and precautions to avoid such hazards discussed.

See other STATIC INITIATION INCIDENTS

Palladium, 2-Propanol

491M, 1975, 205

A stream of hydrogen containing entrained 2-propanol vapour and catalyst particles ignited in contact with air.

See other HYDROGENATION INCIDENTS

See HYDROGENATION CATALYSTS

Palladium trifluoride

See Palladium trifluoride: Hydrogen

Platinum catalyst

Dimitrov, I. *et al.*, *Chem. Abs.*, 1981, **94**, 17876

Some characteristics of ignition of hydrogen on platinum catalyst (title only translated).

Poly(carbon monofluoride)

See Poly(carbon monofluoride): Hydrogen

1,1,1-Tris(azidomethyl)ethane, Palladium catalyst

See 1,1,1-Tris(azidomethyl)ethane: Hydrogen, etc.

Tris(hydroxymethyl)nitromethane

See other NON-METALS, REDUCANTS

See Tris(hydroxymethyl)nitromethane: Hydrogen etc.

4454. Hydrogen (Liquid)

1. Kit and Evered, 1960, 123
2. *Rept. UCRL-3072*, Univ. of California, Berkeley, 1955
3. Weintraub, A. A. *et al.*, *Health Phys.*, 1962, **8**, 11
4. Edutsky, F. J., *Progr. Refrig. Sci. Technol., Proc. 12th Int. Congr. Refrig.*, 1969, **1**, 283–300
5. Lodini, M. A. K. *et al.*, *Int. J. Hydrogen Energy*, 1989, **14**, 35–43
6. *NFPA 50B*, Quincy (Ma), Natl. Fire Prot. Assoc., 1989

The main precaution necessary for use of liquid hydrogen is to prevent air leaking into the system, where it will be condensed and solidified. Fracture of a crystal of solid air or oxygen could produce a spark to initiate explosion [1]. Procedures for the safe handling of liquid hydrogen in the laboratory [2] and in liquid hydrogen bubble chambers [3] have been detailed. Safety problems in large-scale handling and transport of liquid hydrogen have been discussed and safety codes described [4]. Sudden rupture of a 34 kl vacuum jacketed tank of liquid hydrogen appears to have been caused by accidental admission of air to the jacket space, causing excessive boil-off of gas beyond the capacity of the pressure relief [5]. A new US National Fire Code covers all aspects of construction, siting, piping, components and safety devices in consumer systems for liquid hydrogen [6].

Oxygen

1. Cai, T.-J., *Chem. Abs.*, 1981, **94**, 158861
 2. Gamezo, V. N., *et al.*, *Fizika Goreniya i Vzryva*, 1992, **28**(1), 106
- Procedures to remove solid oxygen from liquefied hydrogen before storage are described [1]. Solid oxygen–liquid hydrogen mixtures are very detonable, with low critical diameters and anomalously high detonation velocities [2].

Ozone

See Ozone: Hydrogen (Liquid)

See other CRYOGENIC LIQUIDS

4455. Poly(dimercurymmonium iodide hydrate)

[]

$(\text{H}_2\text{Hg}_2\text{INO})_n$

$(\text{Hg}:\text{N}^+:\text{HgI}^-\text{H}_2\text{O})_n$

Sorbe, 1968, 97

It explodes on heating.

See other IODINE COMPOUNDS, POLY(DIMERCURYIMMONIUM) COMPOUNDS

4456. Potassium amide

[17242-52-3]

H_2KN

KNH_2

1. Brandsma, 1971, 20–21

2. Sorbe, 1968, 68

3. Sanders, D. R., *Chem. Eng. News*, 1986, **64**(21), 2

It has similar properties to the much more widely used and investigated sodium amide, but may be expected on general grounds to be more violently reactive than the former. The frequent fires or explosions observed during work-up of reaction mixtures involving the amide were attributed to presence of unreacted (oxide-coated) particles of potassium in the amide solution in liquid ammonia. A safe filtration technique for removal of the particles is described [1]. It also ignites on heating or friction in air [2]. After preparation from liquid ammonia and potassium, the dry product after evaporation exploded while being chiselled out of the evaporator flask [3].

Ammonia, Copper(II) nitrate

See Copper(II) nitrate: Ammonia, Potassium amide

Potassium nitrite

See Potassium nitrite: Potassium amide

Tetraphenyllead

Houben-Weyl, 1975, Vol. 13.3, 241

One of the by-products of interaction is a highly explosive lead(IV) compound.

Water

Mellor, 1940, Vol. 8, 255

Interaction is violent and ignition may occur, even in contact with humid air. Old samples may explode with considerable delay after contact with water.

See Sodium amide: Water

See other N-METAL DERIVATIVES


4457. Potassium amidosulfate
[13823-50-2]



Metal nitrates or nitrites

See entry METAL AMIDOSULFATES

4458. Potassium hydroxylamine-*O*-sulfonate
[49559-20-8]



1. Anon., private comm., 1985
2. Urben, P. G., private comm., 1989

The salt was isolated (possibly as a hydrate) by unheated vacuum rotary evaporation. The stopper was later ejected from the flask by an exothermic decomposition. It was thought that the salt had been hydrolysed by the water of hydration to give potassium hydrogen sulfate and anhydrous hydroxylamine which is unstable at ambient temperature [1]. It seems more likely that the warm salt would undergo a bimolecular disproportionation to give potassium hydrogen sulfate and the *N*-amino derivative of the title compound, which would decompose to give more potassium hydrogen sulfate and the very reactive and unstable diazene. Hydroxylamine is now thought to be stable when pure [2].

See Diazene

See other N-O COMPOUNDS

4459. Potassium phosphinate ('Potassium hypophosphite')
[7782-87-8]



Air, or Nitric acid

Mellor, 1940, Vol. 8, 882

The salt burns (owing to evolution of phosphine) when heated in air, and explodes when evaporated with nitric acid.

See other REDOX REACTIONS

See other METAL PHOSPHINATES, REDUCANTS

4460. Potassium dihydrogenphosphide
[13659-67-1]



Mellor, 1971, Vol. 8, Suppl. 3, 283

The solid ignites in air.

See related PHOSPHINES

4461. Lanthanum dihydride

[13823-36-4]

H₂La

Kirk-Othmer, 1966, Vol. 11, 207

The very reactive hydride ignites in air.

See other METAL HYDRIDES

4462. Lithium amide

[7782-89-0]

H₂LiN

LiNH₂

Water

Bergstrom, F. W. *et al.*, *Chem. Rev.*, 1933, **12**, 61

It reacts readily with water with a potentially dangerous exotherm.

See other N-METAL DERIVATIVES

4463. Magnesium hydride

[7693-27-8]

H₂Mg

MgH₂

Air, or Water

1. Bailar, 1973, Vol. 1, 34–35

2. Ashby, E. C., *Inorg. Synth.*, 1977, **17**, 4

3. Anon., *Chem. Brit.*, 1985, **21**, 711

4. Vigeholm, B., *Chem. Abs.*, 1987, **106**, 165264

The finely divided hydride produced by pyrolysis is pyrophoric in air, while synthesis from the elements produces a substantially air-stable product [1]. That prepared by reduction of butylmagnesium bromide with lithium tetrahydroaluminate is pyrophoric and reacts violently with water and other protic compounds [2]. The hydride produced from magnesium anthracene has a very large specific surface area and is pyrophoric [3]. In the context of use of the hydride for energy storage purposes, ignition and combustion behaviour of 100–400 g portions were studied, as well as the reaction with water [4].

See other PYROPHORIC MATERIALS

Oxygen (Gas)

See Oxygen (Gas): Metal hydrides

See other METAL HYDRIDES

4464. Magnesium–nickel hydride

[67016-28-8]

H₂Mg.H₂Ni

MgH₂.NiH₂

Hariguchi, S. *et al.*, *Chem. Abs.*, 1981, **94**, 159191

Of several mixed hydrides, the magnesium–nickel hydrides were the most hazardous in terms of dust explosions.

See other DUST EXPLOSION INCIDENTS, METAL HYDRIDES

4465. Sodium amide
[7782-92-5]

NaNH₂

H₂NNa

FPA H102, 1981; HCS 1980, 826

1. Bergstrom, F. W. *et al.*, *Chem. Rev.*, 1933, **12**, 61
2. Brauer, 1963, Vol. 1, 467
3. Krüger, G. R. *et al.*, *Inorg. Synth.*, 1966, **8**, 15
4. Shreve, R. N. *et al.*, *Ind. Eng. Chem.*, 1940, **32**, 173
5. Sandor, S., *Munkavédelem*, 1960, **6**(4-6), 20
6. Rüst, 1948, 283
7. Anon., *Univ. Safety Assoc. Safety News*, 1977, (8), 16
8. Ullmann, 1993, **A24**, 272

It frequently ignites or explodes on heating or grinding in air, particularly if previously exposed to air or moisture to produce degradation products (possibly peroxidic) [1,2,3]. Only one explosion not involving exposure to air has been recorded, during pulverisation [4]. The following oxidation products, all explosively unstable, have been identified: sodium hyponitrite, sodium trioxodinitrate, sodium tetraoxodinitrate, sodium pentaoxodinitrate and sodium hexaoxodinitrate [5]. Several cases of explosive incidents during packing or use of air-exposed material are described [6]. When a half-used 500 g bottle of sodamide was opened, sparks, fumes and intense heat were produced, leading to collapse of the bottle with subsequent explosion [7]. A test for safety of use consists of burning a gram on a spoon, if there are crepitations it is unsafe to use and should be cautiously destroyed [8].

Aryl halide, 1-Alkoxy-1-(trimethylsilyloxy)alkenes

Ferguson, J. R. *et al.*, *Chem. Brit.*, 1997, **33**(6), 21

A mixture of sodamide, bromoanisole, and 1-methoxy-2-methyl-1-(trimethylsilyloxy)-1-butene were reacted at room temperature at a 50 millimolar scale. After some two hours, with slight emission of ammonia, the reaction suddenly became exothermic, with violent gas emission and on one occasion a fire. This was a modest scale-up of a literature procedure for synthesis of 2-alkylbenzoic acids, via a benzyne intermediate. It is advised that this reaction be employed only on smaller scale, with safety precautions. The reaction must pass through a benzocyclobutane intermediate, this, or another, high energy species might accumulate and then decompose.

Halocarbons

MRH Carbon tetrachloride 3.31/50

Sorbe, 1968, 85

Interaction is explosively violent.

Other reactants

Yoshida, 1980, 264

MRH values calculated for 14 combinations with various materials are given.

Oxidants MRH values show % of oxidant
See Chromium trioxide: Sodium amide
 Dinitrogen tetraoxide: Sodium amide MRH 4.31/48
 Potassium chlorate: Sodium amide MRH 2.64/76
 Sodium nitrite: Sodium amide

Water

1. Mellor, 1940, Vol. 2, 255
2. Personal experience

Fresh material behaves like sodium with water, hissing, forming a diminishing floating globule, and often finally exploding [1]. Old, degraded (yellow) samples may be immersed in water for appreciable periods with little action, and then explode very violently. Disposal by controlled burning is safer [2].

See other N-METAL DERIVATIVES

4466. Sodium *O*-hydroxylamide

[22755-22-2]



See Hydroxylamine: Metals

4467. Sodium amidosulfate

[13845-18-6]



Metal nitrates or nitrites

See entry METAL AMIDOSULFATES

4468. Lead(II) nitrate phosphinate

[]



Bailar, 1973, Vol. 2, 131

It is powerfully explosive.

See other HEAVY METAL DERIVATIVES, REDOX COMPOUNDS

4469. Diazene

[3618-05-1]




Its extreme instability is matched by its extremely high endothermicity (ΔH_f° (g) +212.1 kJ/mol, 7.07 kJ/g)

See Potassium hydroxylamine-*O*-sulfonate (reference 2)

See other ENDOTHERMIC COMPOUNDS

4470. Hyponitrous acid
[14448-38-5]



Alone, or Potassium hydroxide


1. Sidgwick, 1950, 693
2. Mellor, 1940, Vol. 8, 407

An extraordinarily explosive solid, of which the sodium salt also explodes on heating to 260°C [1]. An attempt to prepare the acid by treating its silver salt with hydrogen sulfide caused explosive decomposition. Contact with solid potassium hydroxide caused ignition [2].

See Lead hyponitrite

See other N-O COMPOUNDS, REDUCANTS

4471. Nitric amide (Nitramide)
[7782-94-7]



Canis, C., *Rev. Chim. Minerale*, 1964, **1**, 521

Nitramide is quite unstable and various reactions in which it is formed are violent. Attempts to prepare it by interaction of various nitrates and sulfamates showed that the reactions became explosive at specific temperatures.

Alkalies

Thiele, J. *et al.*, *Ber.*, 1894, **27**, 1909

A drop of conc. alkali solution added to solid nitramide causes a flame and explosive decomposition.

Sulfuric acid

Urbanski, 1967, Vol. 3, 16

Nitramide decomposes explosively on contact with conc. sulfuric acid.

See other N-NITRO COMPOUNDS

4472. Sulfamoyl azide (Amidosulfuryl azide)
[13449-16-6]



Shozda, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 2876

It is a low-melting solid, as shock-sensitive as glyceryl nitrate.

See other ACYL AZIDES

4473. Sodium phosphinate ('Sodium hypophosphite')
[7681-53-0]

H₂NaO₂P

NaOP(O)H₂

Mellor, 1940, Vol. 8, 881

Evaporation of aqueous solutions by heating may cause an explosion, phosphine being evolved.

Other reactants

Yoshida, 1980, 345

MRH values calculated for 6 combinations, largely with oxidants, are given.

Oxidants

MRH Sodium chlorate 2.88/49, sodium nitrate 1.00/44

1. Costa, R. L., *Chem. Eng. News*, 1947, **25**, 3177

2. Mellor, 1940, Vol. 8, 881

3. Mellor, 1971, Vol. 8, Suppl. 3, 624

Evaporation of a moist mixture of sodium phosphinate with a trace of sodium chlorate by slow heating caused a violent explosion. It was concluded that, once started, the decomposition of the phosphinate proceeds spontaneously [1]. Similar reactions have been reported with nitrates instead of chlorates. Such mixtures had previously been proposed as explosives [2]. Interaction of iodine with the anhydrous salts is violently exothermic, causing ignition [3].

See Perchloric acid: Sodium phosphinate

See other REDOX REACTIONS

See other METAL PHOSPHINATES, REDUCANTS

4474. Sodium dihydrogen phosphide
[24167-76-8]

H₂NaP

NaPH₂

Albers, H. *et al.*, *Ber.*, 1943, **76**, 23

It ignites in air.

See related PHOSPHINES

4475. Diprotium monoxide (Water)
[7732-18-5]

H₂O

HOH

1. Carson, P. J. *et al.*, *Loss Prev. Bull.*, 1992, **102**, 7

2. Anon. *Safety Digest Univ. Safety Assoc.*, 1994, (50), 22 & 25

3. Editor's comments, 1999

An extremely reactive liquid, solid or vapour with a dangerously high thermal capacity in both liquid and vapour state. When heated, the commonest cause of mid-19th century industrial explosions; it also bursts containers on cooling to low ambient temperatures. Still a frequent cause of vapour explosions today.

H₂O

Reacts with many metals to give hydrogen, sometimes violently. With non-metals pyrophoric hydrides may result. Frequently initiates explosive reactions between other substances. Violent reactions with many non-metal and some metal halides and oxyhalides, also with many organometallic compounds. Many metal nonmetal-ides produce toxic, flammable or pyrophoric gases on contact with diprotium monoxide.

The oxide interacts exothermically with strong acids and bases. When heated to decomposition, diprotium monoxide evolves the dangerous materials hydrogen, oxygen and hydrogen peroxide (all have individual entries in this Handbook). For a few incidents involving diprotium monoxide reactions see [1]. An opposed pair of explosions from careless handling of diprotium monoxide are reported. A student cooled some glass bottles of impure material in liquid nitrogen, they exploded surprisingly violently, causing lacerations. A technician removed a heated carboy of the pure material from an autoclave, a few seconds later it exploded (having presumably been superheated) causing severe burns (these were lessened because she was not wearing safety type footwear, which might have filled with the hot liquid, but open shoes able to drain) [2]. The editor has known injury result from safety footwear when diprotium monoxide at only 70°C ran into it. He has himself emerged almost unscathed in sandals when it was poured on his ankle at 100°C; a similar accident to a colleague wearing shoes immobilised the victim for a week. Reading those Internet sites which cover chemical safety, anecdotally, it is apparent that diprotium monoxide is almost level with nitric acid as a cause of memorable mishap, usually by contact with unsuspected metal or metal hydride traces, sometimes as a catalyst to solid mixtures of an oxidant and a reductant, and occasionally through explosive boiling when heated [3]. In terms of human deaths caused, the most deadly material in this Handbook, also the initiator of the single chemical accident with the highest death toll. Too many potential cross-references to list

See Deuterium oxide

See Methyl isocyanate

Aluminium, Sodium dithionite

Kirschner, E., *Chem. Eng. News*, 1995, **73**(18), 6

During attempts to clear a blocked benzaldehyde pipe, water was allowed to flow into a vessel containing 400 kg of aluminium powder and 2,500 kg of 'sodium hydrosulfite'. The water reacted slowly with the hydrosulfite, evolving heat and hydrogen sulfide. Little attempt was made to deal with the reaction for 10 h. About 2 h later there was an explosion and fire, killing four operators who were attempting to nitrogen blanket the reactor. This was attributed to the aluminium having attained a temperature to react (in which case nitrogen would be useless, since air is unlikely to have been involved).

See Aluminium, Sodium dithionite

Incinerator dust

Takatsuki, H., *Chem. Abs.*, 1995, **122**, 168826k

Evolution of hydrogen, attributed to presence of aluminium particles, formed an explosive atmosphere over wetted incinerator residues.

Metals

U.S. Dept of Energy, 1994, HDBK-1081-94, (*Spontaneous Heating and Pyrophoricity*), 51

Several incidents in which moisture has contributed to fires or explosions, in some of which water was definitely the sole oxidant, in zirconium, magnesium, uranium and thorium scraps or powders are detailed, largely sourced from an earlier paper. However, a plutonium fire was extinguished with water.

See also WATER-REACTIVE COMPOUNDS

4476. Oxosilane

[22755-01-7]

H₂OSi

O:SiH₂

Kautsky, K., *Z. Anorg. Chem.*, 1921, **117**, 209

It ignites in air.

See related SILANES

4477. Hydrogen peroxide

[7722-84-1]

H₂O₂

HOOH

(MCA SD-53, 1969); FPA H3, 1972; HCS 1980, 549 (50%), 550 (30%), 551 (10%); RSC Lab. Hazard Safety Data Sheet No. 57, 1987

1. Shanley, E. S. *et al.*, *Ind. Eng. Chem.*, 1947, **39**, 1536
2. Naistat, S. S. *et al.*, *Chem. Eng. Progr.*, 1961, **57**(8), 76
3. Kirk-Othmer, 1966, Vol. 11, 407; MCA SD-53, 1969
4. Anon., *Fire Prot. Ass. J.*, 1954, 215
5. Smith, I. C. P., private comm., 1973
6. MCA Case History No. 1121
7. McClure, J. D., *J. Org. Chem.*, 1962, **27**, 627
8. Campbell, G. A. *et al.*, *Proc. 4th Symp. Chem Process Haz. Ref. Plant Des.*, 37–43, London, IChE, 1972
9. Clark, M. C. *et al.*, *Chem. & Ind.*, 1974, 113
10. Cookson, P. G. *et al.*, *J. Organomet. Chem.*, 1975, **99**, C31–32
11. *Manufacture of Organic Chemicals using Hydrogen Peroxide and Related Compounds*, AO 1.2, Widnes, Interlox Chemicals Ltd., 1979
12. Berthold, W. *et al.*, *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Chem. Ind.*, 1431–1434, Basle, SSCI, 1980
13. Clemens, D., *Chem. Eng. News*, 1986, **64**(50), 2
14. Manning, D., *Chem. Eng. News*, 1987, **65**(9), 4
15. Merrifield, 1988, 23 pp.
16. 'Hydrogen Peroxide in Organic Chemistry', Schirmann, J. P., Delavarenne, S. T., Paris, Informations Chimie, 1987 (English edn.)
17. Ullmann, 1989, **A13**, 462
18. Mackenzie, J., *Chem. Eng.*, 1990, **97**(6), 84

19. Tojo, G. INTERNET, 1997

20. Davies, A. G. *et al.*, *J. Chem. Soc., Perkin Trans, 2*, 1981, 1512

The hazards attendant upon use of conc. hydrogen peroxide solutions have been reviewed [1,2,3]. Salient points include:

Release of enough energy during catalytic decomposition of 65% peroxide to evaporate all water present and formed, and subsequent liability of ignition of combustible materials.

Most cellulosic materials contain enough trace-metal catalysts to cause spontaneous ignition with 90% peroxide.

Contamination of conc. peroxide causes possibility of explosion. Readily oxidisable materials, or alkaline substances containing heavy metals, may react violently.

Soluble fuels (acetone, ethanol, glycerol) will detonate on admixture with peroxide of over 30% concentration, the violence increasing with concentration.

Handling systems must exclude fittings of iron, brass, copper, Monel, and screwed joints caulked with red lead preparations.

Concentrated peroxide may decompose violently in contact with iron, copper chromium and most other metals and their salts, and dust (which frequently contains rust). Absolute cleanliness, suitable equipment (PVC, butyl or Neoprene rubber, Teflon) and personal protection are essential for safe handling [4].

During concentration under vacuum of aqueous [5] or of aqueous–alcoholic [6] solutions of hydrogen peroxide, violent explosions occurred when the concentration was sufficiently high (probably over 90%) [3]. It is possible to accidentally concentrate 30% hydrogen peroxide to explosion on a laboratory rotary evaporator [19]. A suspension of 90% hydrogen peroxide in dichloromethane is shock-sensitive [7]. Detonation of hydrogen peroxide vapour has been studied experimentally [8]. Explosion of a screw capped winchester of 35% peroxide solution after 2 years owing to internal pressure of liberated oxygen emphasises the need to date-label materials of limited stability, and to vent the container automatically by fitting a Bunsen valve or similar device [9]. It has been suggested that hazards in use and handling of concentrated hydrogen peroxide can be avoided by using the solid 2:1 complex of hydrogen peroxide with 1,4-diazabicyclo[2.2.2]octane. This is hygroscopic but supposedly stable for at least several months in storage, although it has been reported to decompose above 60°C [10]. The 1:1 complex, which is of considerably lower energy, has proved explosive, however [20].

A survey, with many references, of 14 classes of preparative reactions involving hydrogen peroxide or its derivatives emphasises safety aspects of the various procedures [11]. Following the decomposition of 100 l of 50% aqueous hydrogen peroxide which damaged the 630 l stainless vessel rated at 6 bar, the effect of added contaminants and variations in temperature and pH on the adiabatic decomposition was studied in a 1 l pressure vessel, where a final temperature of 310°C and a pressure around 200 bar were attained. Rust had little effect, but addition of a little ammonia (pH increased from 1.8 to 6.0) caused the induction period to fall dramatically, effectively from infinity to a few h at 40°C and a few min at 80°C. Addition of sodium hydroxide to pH 7.5 reduced the induction period at 24°C from infinity to about 4 min [12].

Several 0.5–1 l bottles of an unspecified plastic containing 30% hydrogen peroxide which had been stored for over a year became brittle and developed cracks during normal handling operations. One bottle stored for more than 4 years broke into small fragments when squeezed by hand [13]. However, these bottles were undoubtedly of the wrong material, because storage of up to 54% hydrogen peroxide solutions in suitable plastic bottles with no signs of container deterioration is a well-established commercial practice [14]. A recent report and review update and extend general information on storage and handling of the powerful oxidant. The second pays considerable attention to headspace explosions caused by decomposition in the presence of volatile fuels. It is calculated that during adiabatic decomposition of 90% peroxide solution, a temperature of 740°C could be attained, accompanied by the release of 5233 volumes of oxygen and steam [15]. Detonation limit diagrams for hydrogen peroxide vapours against liquid phase composition, temperature and pressure are given [17]. Each chapter of a recent monograph is devoted to a specific functional group, and the final chapter is concerned with safety aspects of the use of hydrogen peroxide as oxidant in mixtures with organic compounds [16].

See other INDUCTION PERIOD INCIDENTS

Acetal, Acetic acid

Ashley, J. N. *et al.*, *Chem. & Ind.*, 1957, 702

An organic sulfur compound containing an acetal function had been oxidised to the sulfone with 30% hydrogen peroxide in acetic acid. After the liquor had been concentrated by vacuum distillation at 50–60°C, the residue exploded during handling. This was attributed to formation of the peroxide of the acetal (formally a *gem*-diether) or of the aldehyde formed by hydrolysis, but formation and explosion of peracetic acid seems a more likely explanation.

Acetaldehyde, Desiccants

Karnojitzky, V. J., *Chim. Ind.* (Paris), 1962, **88**, 235

Interaction gives the extremely explosive poly(ethylidene) peroxide, also formed on warming peroxidised diethyl ether.

Acetic acid

1. Grundmann, C. *et al.*, *Ber.*, 1939, **69**, 1755

2. *491M*, 1975, 207

During preparation of peracetic acid, the temperature should not be too low to prevent reaction as the reagents are mixed, because reaction may begin later with explosive violence [1]. Heating well-diluted peroxide and acid will exothermically produce the acid which may explode at 110°C [2].

Acetic acid, *N*-Heterocycles

1. Wommack, J. B., *Chem. Eng. News*, 1977, **55**(50), 5

2. Dholakia, S. *et al.*, *Chem. & Ind.*, 1977, 963

3. Daisley, R. W. *et al.*, *Org. Prep. Proc. Int.*, 1983, **15**, 281

During isolation of the di-*N*-oxide of 2,5-dimethylpyrazine [1] and of the mono-*N*-oxide of 2,2-bipyridyl [1], prepared by action of hydrogen peroxide in acetic

acid on the heterocycles, violent explosions occurred on evaporation of the excess peroxyacetic acid. In the preparation of 3-bromopyridine *N*-oxide, the necessity to take adequate precautions against the possibility of explosion while vacuum distilling off excess reagents from a water bath maintained at 50°C is stressed [3]. For a method not involving this considerable explosion risk.

See N-OXIDES

See 3-Methyl-4-nitropyridine *N*-oxide

Acetic acid, Jute

De Forest, P. *et al.*, *Chem. Eng. News*, 1987, **65**(31), 2; *J. Chem. Educ.*, 1988, **65**(8), 728–729

Jute fibre (0.1 g) was being prepared for a standard forensic examination by treatment in glacial acetic acid (20 ml) with hydrogen peroxide solution (20 ml, but in error 30% peroxide was used in place of the specified 6% '20-vol') strength. After heating on a water bath for 4 h, there was a violent explosion which shattered the ceramic hotplate below the bath with formation of 'ceramic shrapnel'. The available hydrogen peroxide would lead to formation of 14.7 g of peracetic acid as a 34 wt% solution in aqueous 33% acetic acid. This would be expected to become more concentrated by evaporation during prolonged heating, and concentrated solutions are known to be thermally unstable.

See Perchloric acid: Clay

Acetic acid, 3-Thietanol

1. Dittmer, D. C. *et al.*, *J. Org. Chem.*, 1961, **26**, 1324
2. Dittmer, D. C. *et al.*, *J. Org. Chem.*, 1971, **36**, 1324
3. Chang, P. *et al.*, *J. Org. Chem.*, 1969, **34**, 2791
4. Lamm, B. *et al.*, *Acta Chem. Scand. Ser. B*, 1974, **28**, 701–703

When following the original route [1] to 3-hydroxythietane-1,1-dioxide, it is essential to dilute and evaporate the hydrogen peroxide–acetic acid reaction mixture slowly from a large dish, to prevent explosions arising from concentration of peroxyacetic acid [2]. Alternative routes to avoid this hazard are available [3,4].

See Peroxyacetic acid

Acetic anhydride

1. Prett, K., *Textilveredlung*, 1966, **1**, 288–290
2. Swern, D., *Chem. Rev.*, 1945, **45**, 5

During preparation of peracetic acid solutions for textile bleaching operations, the reaction mixture must be kept acid. Under alkaline conditions, highly explosive diacetyl peroxide separates from solution [1]. An excess of the anhydride has the same effect [2].

Acetone, Other reagents

MRH Acetone 6.36/19

1. Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 3
2. *MCA Case History No. 233*
3. *MCA Case History No. 223*
4. Stirling, C. J. M., *Chem. Brit.*, 1969, **5**, 36

5. Seidl, H. *Angew. Chem. (Intern. Ed.)*, 1964, **3**, 640
6. Treibs, W., *Angew. Chem. (Intern. Ed.)*, 1963, **3**, 802
7. Brewer, A. D., *Chem. Brit.*, 1975, **11**, 335
8. Schurz, J. *et al.*, *Angew. Chem.*, 1956, **68**, 182
9. Ghilardi, C. A. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1986, 12075–2081
10. Anon., *Sichere Chemiearb.*, 1987, **39**, 70
11. Ullmann, 1989, **A13**, 462

Acetone and hydrogen peroxide readily form explosive dimeric and trimeric cyclic peroxides, particularly during evaporation of the mixture. Many explosions have occurred during work-up of peroxide reactions run in acetone as solvent, including partial hydrolysis of a nitrile [1] and oxidation of 2,2-thiodiethanol [2] and of an unspecified material [3]. The reaction mixture from oxidation of a sulfide with hydrogen peroxide in acetone exploded violently during vacuum evaporation at 90°C. On another occasion, oxidation of a sulfide in acetone in presence of molybdate catalyst proceeded with explosive violence. A general warning against using acetone as a solvent for peroxide oxidations is given [4]. During the isolation of 1-tetralone, produced by oxidation of tetralin with hydrogen peroxide in acetone, a violent explosion occurred which was attributed to acetone peroxide [5]. The originator of the method later gave detailed instructions for a safe procedure, which feature the exclusion of mineral acids, even in traces [6].

During oxidation of an unspecified sulfur heterocycle in acetone with excess 35% hydrogen peroxide, a white solid separated during 3 days standing in a cool place. The solid (20 g) appeared to have been acetone peroxide, because it exploded with great violence during drying in a vacuum oven. The previous warning [4] on incompatibility of acetone and hydrogen peroxide was repeated [7]. A cyclic diketone was oxidised with alkaline peroxide in acetone solution and the product isolated by a solvent extraction procedure. During vacuum evaporation of the ethyl acetate extract, gassing was noted and the concentrate was treated twice with platinum catalyst to decompose excess hydrogen peroxide. After filtration and further vacuum evaporation, further gassing was noted and heating was discontinued, but the residue exploded [8]. This seems likely to have happened because the cyclic acetone peroxides, as dialkyl peroxides (and unlike hydroperoxides) would largely survive the catalytic decomposition treatment.

During the oxidation of the ligand tris(diphenylphosphinylethyl)amine with hydrogen peroxide in acetone to give tris(diphenylphosphonylethyl)amine, overheating must be avoided to prevent damaging explosions (of acetone peroxide). A Teflon beaker is recommended for the reaction [9]. A thiol was being oxidised in acetone to the disulfide with hydrogen peroxide. No product separated on standing, so the solution was concentrated to one fifth of its bulk in a rotary vacuum evaporator, and a white solid separated. During subsequent operations, the solid (trimeric acetone peroxide) exploded with great violence [10].

Detonation limit diagrams for hydrogen peroxide, water, acetone systems are given [11].

See 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxonane

See also KETONE PEROXIDES

Aconitic acid

Anon., private communication, 1993

An attempt was made to convert aconitic acid (2 g) to isocitric acid using 25 ml. of 30% aqueous hydrogen peroxide, time and temperature unspecified. A crystalline product precipitated which exploded on grinding in a mortar.

Alcohols

MRH Ethanol 6.19/19

1. MCA SD-53, 1969

2. Spengler, G. *et al.*, *Brennst. Chem.*, 1965, **46**, 117

3. Heemskerck, A. H. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries*, Vol I, (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 411, Amsterdam, Elsevier, 1995

Homogeneous mixtures of concentrated peroxide with alcohols or other peroxide-miscible organic liquids are capable of detonation by shock or heat [1]. Furfuryl alcohol ignites in contact with 85% peroxide within 1 s [2]. Detonability limits of mixtures with 2-propanol have been measured. Approximately stoichiometric combinations of 50% hydrogen peroxide and the alcohol could be made to detonate, as could a wider range of mixtures with higher test peroxide [3].

See Oxygenated compounds, etc., below

Alcohols, Sulfuric acid.

1. Hedaya, E. *et al.*, *Chem. Eng. News*, 1967, **45**(43), 73

2. Merrifield, 1988, 5–6, 16

During conversion of alcohols to hydroperoxides, the order of mixing of reagents is important. Addition of conc. acid to mixtures of an alcohol and conc. peroxide almost inevitably leads to explosion, particularly if the mixture is inhomogeneous and the alcohol is a solid [1]. In presence of concentrated acid, hydrogen peroxide forms peroxymonosulfuric acid (Caro's acid), especially if the rate of conversion of the alcohol is low. The peroxyacid is a very powerful and unstable oxidant, and its reactions are often violent. Thus in the hydrogen peroxide–sulfuric acid–alcohol ternary system, an autodetonation region may exist. This is shown in a triangular diagram for 2-propanol and *tert*-butanol [2].

See tert-Butanol, etc.; 2-Phenyl-1,1-dimethylethanol; 3,5-Dimethyl-3-hexanol, all below

Aluminium isopropoxide, Heavy metal salts

Ward, D. S., unpublished information, 1974

During preparation of an alumina catalyst, the isopropoxide was stirred with 6% hydrogen peroxide to generate a slurry of pseudo-boehmite alumina, to which was added a solution of heavy metal salts. The heating and foaming of the mixture was excessive and foam overflowed into a safety tray, then ignited. The incident was attributed to either spontaneous or static-induced ignition of the foam consisting of oxygen-rich bubbles in the isopropanol-containing liquid medium. The heavy metal salts would catalyse decomposition of the hydrogen peroxide and may also have been involved in the ignition process.

See Metals, etc., below

2-Amino-4-methyloxazole

Rapi, G. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1982, 1339–1340

Oxidative rearrangement of the oxazole to 4-hydroperoxy-5-hydroxy-4-methylimidazolidin-2-one in presence of iron(II) catalysts at ambient temperature may become explosive if not controlled effectively.

Ammonia

Matlow, S. L., *Chem. Eng. News*, 1990, **68**(30), 2

A sealed container of mixed 30% aqueous solution and aqueous ammonia exploded on storage, driving shards into a wall 18 m away.

See Oxygen(Gas): Ammonia and Oxygen(Liquid): Ammonia

Aromatic hydrocarbons, Trifluoroacetic acid

MRH Toluene 6.62/15

Demo, N. C. *et al.*, *Tetrahedron Lett.*, 1977, 1703–1704

A solution of 30% aqueous hydrogen peroxide in trifluoroacetic acid is useful for destructive oxidation of the aromatic ring in preference to the side chains as is usual with most oxidants. During work-up operations, the excess peroxide must be catalytically decomposed with manganese dioxide before removal of solvent to prevent explosions.

Azelaic acid, Sulfuric acid

Berkowitz, S., US Pat. 4 147 720, 1979

Conversion of the acid to diperoxyazelaic acid in hydrogen peroxide/sulfuric acid medium at 45–50°C was uncontrollably exothermic and led to explosion. Use of a peroxide/phosphoric acid medium gives a safe and effective conversion of C₆–C₁₆ diacids to the diperoxy acids, though up to 5% of sulfuric acid is needed as catalyst for C₁₂–C₁₆ acids.

See Carboxylic acids, below

Benzene

Merrifield, 1988, 13

A triangular diagram shows the range of explosive mixtures in the hydrogen peroxide–benzene–water ternary system, and also includes data for various oxygenated water-soluble organic compounds.

See Oxygenated compounds, below

Benzenesulfonic anhydride

Fichter, F., *Helv. Chim. Acta*, 1924, **7**, 1072

Attempts to prepare benzeneperoxy-sulfonic acid by interaction of the anhydride and 90–95% peroxide led to explosively violent decomposition within a few s. When diluted with acetic acid, the reaction mixture soon began to decompose, leading to violent boiling. It was concluded that the peroxyacid is too unstable for more than transitory existence.

tert-Butanol, Sulfuric acid

Schenach, T. A., *Chem. Eng. News*, 1973, **51**(6), 39

Preparation of di-*tert*-butyl peroxide by addition of *tert*-butanol to 50% hydrogen peroxide–78% sulfuric acid (1:2 by wt) is a dangerously deceptive procedure. On the small scale, and with adequate cooling capacity it may be possible to prevent the initial stage (exothermic formation of *tert*-butyl hydroperoxide) getting out of control and initiating violent or explosive decomposition of the peroxide–peroxomonosulfuric acid mixture. This hazard diminishes as the reaction proceeds with consumption of hydrogen peroxide and dilution by the water of reaction. On the plant scale several severe explosions occurred, preceded only by a gradual temperature increase, during attempted process development work.
See Alcohols, above

Carbon

MRH 6.19/15

1. Mellor, 1939, Vol. 1, 936–938
2. Schumb, 1955, 402, 478

The violent decomposition observed on adding charcoal to conc. hydrogen peroxide is mainly owing to catalysis by metallic impurities present and the active surface of the charcoal, rather than to direct oxidation of the carbon [1]. Charcoal mixed with a trace of manganese dioxide ignites immediately on contact with conc. peroxide [2].

Carboxylic acids

MRH Formic acid 3.30/58

Kuchta, J. M. *et al.*, *Rep. Invest. No. 5877*, Washington, US. Bur. Mines, 1961
Admixture produces peroxyacids, some of which are unstable and explosive. Aqueous peroxide solutions containing formic acid, acetic acid or tartaric acid above certain concentrations can be caused to detonate by a severe explosive shock.

See Azelaic acid, etc., above; Oxygenated compounds, etc., below

See PEROXYACIDS

Catalyst (unspecified)

1. Anon., *Sichere Chemiearb.*, 1979, **31**, 63
2. Merrifield, 1988, 16

Aqueous cyanide residues were being detoxified by treatment with a catalyst and then 50% peroxide solution to oxidise cyanide to cyanate at pH 10 ± 0.5 at above 40°C. Apparently all the alkaline catalyst solution had not been completely drained from the 100 l dosing flask before the peroxide solution (28 kg) was charged. The latter solution began to decompose and this accelerated to explosion, destroying the flask [1]. In a continuous bleaching plant, a 200 l stainless dosing vessel, unrelieved except for a small overflow line, was connected directly to a supply of 35 wt% peroxide solution and to the bleaching process vessel. Back-contamination of the peroxide solution in the dosing vessel led to exothermic decomposition, and the evolved gases caused the vessel to pressure-burst [2].

See other GAS EVOLUTION INCIDENTS

Coal

Wood, C. R., *Fuel*, 1974, **53**, 220

Coal (72 mesh) slurried with hydrogen peroxide solution (30 wt%) reacts steadily on warming to 85°C, but if much pyrites is present, the oxidation may accelerate to a dangerous extent.

See other CATALYTIC IMPURITY INCIDENTS

Copper(II) chloride

Merrifield, 1988, 7

In a printed circuit board etching line using copper(II) chloride solution, 45 wt% hydrogen peroxide solution was used to recover the copper salts. The peroxide header tank became contaminated with trace amounts of the etching solution, and catalytic decomposition of the peroxide led to a pressure burst of the tank.

See other CATALYTIC IMPURITY INCIDENTS, GAS EVOLUTION INCIDENTS

Cotton waste

Merrifield, 1988, 6

Spillage of a weak (5%) aqueous peroxide solution onto cotton waste led, after some time, to ignition. This probably involved concentration of the peroxide by evaporation/absorption of the water, the heat insulating effect of the fibrous mass preventing heat loss as oxidation proceeded, and possibly the presence of trace metals in the waste promoting catalytic decomposition.

See Wood, below

Diethyl ether

Bruhl, J. W., *Ber.*, 1895, **28**, 2856–2857

Evaporation of an ethereal solution of hydrogen peroxide gave a residue of which a drop on a platinum spatula exploded weakly on exposure to flame. When the sample (1–2 g) was stirred with a glass rod (not fire polished), an extremely violent detonation occurred.

See other GLASS INCIDENTS

See ETHERS

3,5-Dimethyl-3-hexanol, Sulfuric acid

MCA Guide, 1972, 315

The alcohol was treated with 90% hydrogen peroxide and a trace of sulfuric acid at 0°C. While warming to ambient temperature overnight it exploded violently. Some 3,5-dihydroperoxide may have formed from autoxidation at the 5-position.
See Alcohols, etc.; *also* *tert*-Butanol, etc.; both above, *and* 2-Methyl-1-phenyl-2-propanol, etc., below

Dimethylphenylarsine

Anon., *Lab. Accid. Higher Educ.*, item 8, Barking, HSE, 1987

During the preparation of the As-oxide in glassware by oxidation with hydrogen peroxide (conditions unknown), an explosion occurred, possibly due to presence of impurities.

See Dimethylphenylphosphine, next below

Dimethylphenylphosphine

Dennister, M. L. *et al.*, *Inorg. Synth.*, 1977, **17**, 185

Oxidation of the phosphine to the oxide by adding its solution in ether to stirred 15% aqueous peroxide may become very violent if the rate of addition is too fast.
See Dimethylphenylarsine, next above

Diphenyl diselenide

Bloodworth, A. J., *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1983, 471–473

Attempted thermal dehydration of benzeneseleninic acid, formed by oxidation of diphenyl diselenide with hydrogen peroxide, gave a solid which exploded at 53–55°C. The solid may have been the complex of the acid with hydrogen peroxide.

2-Ethoxyethanol, Polyacrylamide gel, Toluene

1. Darnall, D. W., *Chem. Eng. News*, 1978, **56**(47), 47

2. Shanley, E. S., *Chem. Eng. News*, 1979, **57**(21), 43

Polyacrylamide gels were dissolved in 30% peroxide solution and added to a scintillation mixture in 1:1 2-ethoxyethanol–toluene. After counting, the mixtures were bulked and evaporated intermittently with heat during a 4 week period, and the accumulated peroxidised residues eventually exploded violently [1]. A subsequent comment indicated that peroxidised materials may not necessarily have been formed, because solutions of organic materials in aqueous peroxide are themselves potentially explosive [2].

See Oxygenated compounds, below

Ethyl acetate

Schierwater, F.-W., *Jahresber.*, 1981, 77

Organic material was extracted from a sample with methanol, and after evaporation the small residue was refluxed with aqueous peroxide. Extraction with ethyl acetate, separation and evaporation gave a residue which was scarcely visible but which exploded violently when the flask was moved. Several peroxidic species, including methyl hydroperoxide, peroxyacetic acid or its ethyl ester, may have been involved.

See Ethyl acetate

Fatty acid

Merrifield, 1988, 7

In a continuous operation to bleach a fat-derived acid, the latter was stirred and treated at 80°C with a slow stream of 50 wt% hydrogen peroxide. Un-noticed failure of the agitator led to peroxide build-up, layer formation and eventual formation of an explosive mixture. When this passed into a centrifugal pump, it detonated.

See other AGITATION INCIDENTS

Formic acid, Metaboric acid

See Peroxyformic acid

Gadolinium hydroxide

Bogdanov, G. A. *et al.*, *Chem. Abs.*, 1976, **85**, 52253

Interaction gives a hydrated basic peroxide which decomposes explosively at 80–90°C.

Gallium, Hydrochloric acid

Titze, H., *Mikrochem. Acta*, 1977, 480

During dissolution for analysis of 1–9 g samples of gallium in conc. hydrochloric acid containing small portions of 30% hydrogen peroxide solution, cooling may be necessary to prevent development of explosively violent reactions at higher temperatures.

Hexamethylenetetramine

See 1,6-Diaza-3,4,8,9,12,13-hexaoxabicyclo[4.4.4]tetradecane

Hydroboration product of 2-methyl-2-propenyl tetrahydropyranyl ether

See TETRAHYDROPYRANYL ETHER DERIVATIVES

Hydrogen, Palladium catalyst

Anon., *Jahresber.*, 1982, 65–66

In the cyclic anthraquinone process for preparing hydrogen peroxide, a hydrocarbon solution of a 2-alkylanthraquinone is catalytically reduced with hydrogen to the 9,10-diol, which is then air-oxidised to the quinone with formation of hydrogen peroxide. The latter is removed from the process solution by water washing and phase separation, and the process stream, now virtually free of peroxide is recycled to the reactor. Failure of the compressed air supply prevented the separation of the aqueous hydrogen peroxide from the hydrocarbon phase, and both layers were recycled into the process. Contact of the peroxide solution with the catalyst released oxygen, which mixed with hydrogen and was ignited by the catalyst. The initial explosion released some of the flammable process solution which ignited, and eventually some 70 m³ of the latter was involved in a major fire, with a further explosion in the hydrogenation plant.

See other HYDROGENATION INCIDENTS

Hydrogen selenide

Mellor, 1939, Vol. 1, 941

Interaction is very fast.

Immiscible organic liquids

Merrifield, 1988, 5

Emulsions formed from immiscible organic liquids in aqueous peroxide mixtures may behave in the same way as miscible organic liquids, but if the emulsion breaks and separation of the organic phase occurs, passage into an explosive region of the peroxide–water–organic liquid ternary system may occur, and this is potentially very dangerous.

Iron(III) chloride, Hydrocarbons

Merrifield, 1988, 7–8

A 6000 m³ aqueous refinery effluent containing sulfides and traces of hydrocarbons was to be treated to remove sulfides before discharge. Aqueous ferric chloride was added, followed by 1800 l of 50 wt% peroxide added over 40 min, and after a further 30 min an explosion occurred which blew off the lid of the treatment vessel. This was attributed to ignition of the explosive mixture of hydrocarbon vapours and oxygen (from iron-catalysed decomposition of peroxide) formed above the liquid surface.

See other GAS EVOLUTION INCIDENTS

Iron(II) sulfate, 2-Methylpyridine, Sulfuric acid

Mond. Div., ICI, private. comm., 1969

Addition of 30% peroxide and sulfuric acid to 2-methylpyridine and iron(II) sulfate caused a sudden exotherm, followed by a vapour phase explosion and ignition. Lack of stirring is thought to have caused local overheating, vaporisation of the base and its ignition in the possibly oxygen-enriched atmosphere.

See other AGITATION INCIDENTS

Iron(II) sulfate, Nitric acid, Sodium carboxymethylcellulose

1. Sansoni, B. *et al.*, *Z. Anal. Chem.*, 1968, **243**, 209

2. Cela Torrijos, R. *et al.*, *Chem. & Eng. News*, 1978, **56**(49), 51

A published analytical procedure [1] for decomposing the sodium salt with ferrous-sulfate/peroxide in nitric acid at pH 2 led to deflagration or explosion during the evaporation stage when applied to 10 g samples, but not with 5 g samples. Presence of sulfuric acid avoids the problem [2].

Ketene

Swern, D., *Chem. Rev.*, 1949, 45, 7

Interaction with excess ketene rapidly forms explosive diacetyl peroxide.

Ketones, Nitric acid

Bjorklund, G. H. *et al.*, *Trans. R. Soc. Can., (Sect. III)*, 1950, **44**, 25

Unless the temperature and concentrations of reagents were carefully controlled, mixtures of hydrogen peroxide, nitric acid and acetone overheated and exploded violently. Under controlled conditions, the explosive dimeric or trimeric acetone peroxides were produced. 2-Butanone and 3-pentanone gave shock- and heat-sensitive oily peroxides. Cyclopentanone reacts vigorously, giving a solid which soon produces a series of explosions if left in contact with the undiluted reaction liquor. The isolated trimeric peroxide is very sensitive to shock, slight friction or rapid heating, and explodes very violently. Cyclohexanone and 3-methylcyclohexanone gave oily, rather explosive peroxides.

See CYCLIC PEROXIDES, KETONE PEROXIDES

Lead, Trioxane

Bamberger, M. *et al.*, *Z. Ges. Schiess-u. Sprengstoffw.*, 1927, **22**, 125–128

Mixtures of trioxane with 60% hydrogen peroxide solution are detonable by heat or shock, or spontaneously after contact with metallic lead. The latter may be owing to the heat of oxidation of lead.

See Metals *also* Oxygenated compounds, etc., both below

Lithium tetrahydroaluminate

Osmon, R. V., *Chem. Eng. Progr. Symp. Ser.*, 1966, **62**(61), 92–102

This oxidiser–fuel combination showed promise in rocketry.

See other REDOX REACTIONS

Mercury(II) oxide, Nitric acid

MRH Mercury(II) oxide 1.59/tr.

Mellor, 1940, Vol. 4, 781

Although red mercuric oxide usually vigorously decomposes hydrogen peroxide, the presence of traces of nitric acid inhibits decomposition and promotes formation of red mercury(II) peroxide. This explodes on impact or friction, even when wet, if the mercury oxide was finely divided.

Metals, or Metal oxides, or Metal salts

1. Mellor, 1939, Vol. 1, 936–944

2. Schumb, 1955, 480

3. Kit and Evered, 1960, 136

4. *Hydrogen Peroxide Data Manual*, Laporte Chemicals Ltd., Luton, 1960

5. Hardin, M. C. *et al.*, *Chem. Abs.*, 1963, **59**, 2583h

6. Winnacker, 1970, Vol. 2, 552

MRH Aluminium 12.76/35, iron 4.60/52, lithium 13.64/29, magnesium 12.68/41, silver, 1.59/tr., sodium 5.81/57; manganese dioxide, mercury(I) oxide, mercury(II) oxide, calcium permanganate, all 1.59/tr.

The noble metals are all very active catalysts, particularly when finely divided or dispersed on high-area surfaces, for the decomposition of hydrogen peroxide, silver being used for this purpose in peroxide-powered rocket motors. Gold and the platinum group metals behave similarly [2,3,4]. Addition of platinum black to conc. peroxide solution may cause an explosion, and powdered magnesium and iron, promoted by traces of manganese dioxide, ignite on contact [1]. Potassium and sodium are oxidised violently by conc. peroxide solutions [1], and interaction of lithium and hydrogen peroxide is hypergolic and controllable under rocket motor conditions [5]. The effectiveness of metals for catalytic decomposition of peroxide solutions decreases in the order: osmium, palladium, platinum, iridium, gold, silver, manganese, cobalt, copper, lead [6]. Oxides of cobalt, iron (especially rust), lead (also the hydroxide), manganese, mercury and nickel are also very active and the parent metals and their alloys must be rigorously excluded from peroxide handling systems [1,4]. Soluble derivatives of many other metals, particularly under alkaline conditions, will also catalyse the exothermic decomposition, even at low concentrations [4]. Calcium permanganate has been used, either as a solid or in conc. solution, to ignite peroxide rocket motors [3]. Comprehensive data on all aspects of handling and use of conc. peroxide are available [4].

See Carbon, above

Methanol, *tert*-Amine, Platinum catalyst

Laughlin, R. G., *Chem. Eng. News*, 1978, **56**(19), 38

To solubilise a dialkylmethylamine during oxidation to the *N*-oxide with hydrogen peroxide, methanol was used as diluent in place of water. After oxide formation

was complete, platinum black was added to decompose excess peroxide, and an explosion occurred. This was attributed to ignition of the methanol vapour–oxygen mixture by glowing catalyst particles.

See Oxygenated compounds, below

Methanol, Copper sulfate, Cyanide residues

Merrifield, 1988, 8

Aqueous cyanide effluent containing a little methanol in a 2 m³ open tank was being treated to destroy cyanide by oxidation to cyanate with hydrogen peroxide in the presence of copper sulfate as catalyst. The tank was located in a booth with doors. Addition of copper sulfate (1 g/l) was followed by the peroxide solution (27 l of 35 wt%), and after the addition was complete an explosion blew off the doors of the booth. This was attributed to formation of a methanol vapour–oxygen mixture above the liquid surface, followed by spontaneous ignition. It seems remotely possible that unstable methyl hydroperoxide may have been involved in the ignition process.

See Waste treatment, below.

See other GAS EVOLUTION INCIDENTS

Methanol, Phosphoric acid

1. Anon., *J. Electrochem. Soc.*, 1965, **112**, 251C

2. King, C. V., *J. Electrochem. Soc.*, 1965, **112**, 1057; 1966, **113**, 519

In mixtures of the 3 components intended for use as chemical polishing or etching solutions [1], conc. peroxide must not be used [2].

See Oxygenated compounds, below

4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2,6}]undeca-8-ene-3,5-dione, Potassium hydroxide

1. Olsen, H. *et al.*, *J. Amer. Chem. Soc.*, 1977, **99**, 1533

2. Engel, P. S. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 7107

Treatment of the dione dissolved in ethylene glycol–30% hydrogen peroxide with potassium hydroxide leads to a violently exothermic hydrolysis—*N*-oxidation reaction [1]. Simultaneous addition of peroxide and alkali solutions from separate funnels gives a controllable and higher yielding reaction [2].

Nitric acid, Soils

See Nitric acid: Hydrogen peroxide, Soils

Nitric acid, Thiourea

Bjorklund, G. H. *et al.*, *Trans. R. Soc. Can., (Sect. III)*, 1950, **44**, 28

The solid peroxide produced by action of hydrogen peroxide and nitric acid on thiourea (and possibly a hydrogen peroxidate of thiourea dioxide) decomposed violently on drying in air, with evolution of sulfur dioxide and free sulfur.

See CRYSTALLINE HYDROGEN PEROXIDATES

Nitriles

Wakakura, M. *et al.*, *Netsu Sokutsei*, 1994, **21**(2), 68; *Chem. Abs.*, 1994, **121**, 57411f

A calorimetric study of the alkaline hydrogen peroxide conversion of organic nitriles to amides was undertaken. The reaction showed potential for runaway if carelessly scaled up.

Nitrogenous bases

MRH values below references

1. Stone, F. S. *et al.*, *J. Chem. Phys.*, 1952, **20**, 1339
2. Urbanski, 1967, Vol. 3, 306
3. *Haz. Chem. Data*, 1975, 296

MRH Ammonia, 5.86/25, aniline 6.44/17, dimethylhydrazine 6.69/19

Ammonia dissolved in 99.6% peroxide gave an unstable solution which exploded violently [1]. In the absence of catalysts, conc. peroxide does not react immediately with hydrazine hydrate. This induction period has caused a number of explosions and accidents owing to sudden reaction of accumulated materials [2]. 1,1-Dimethylhydrazine is hypergolic with high-test peroxide [3].

See other INDUCTION PERIOD INCIDENTS

Organic compounds

MRH Acetic acid 5.10/31, ethanol 6.19/19

1. Hutton, E., *Chem. Brit.*, 1969, **5**, 287
2. *The Manufacture of Organic Chemicals using Hydrogen Peroxide*, AO 1.2, 36–37, Widnes, Interox Chemicals Ltd., 1979
3. Nolan, 1983, Case history 156
4. Merrifield, 1988, 8
5. Mackenzie, J., *Plant/Oper. Progr.*, 1991, **10**(3), 143
6. Klais, O., *Thermochim. Acta*, 1993, **225**(2), 213

Although under certain circumstances mixtures of hydrogen peroxide and organic compounds are capable of developing more explosive power than an equivalent wt of TNT, in many cases interaction can be effected safely and well under control by applying well-established procedures, to which several references are given [1]. A triangular diagram specifically for the system hydrogen peroxide–glycerol–water, but also generally applicable to other organic compounds, such as acetic acid, ethanol, aniline or quinoline, shows the range of detonable compositions, and gives guidance and examples on avoiding the hazardous regions in reaction systems [2]. A polyester was made from propylene glycol, adipic acid and aluminium stearate catalyst by distilling out the water of reaction at 230°C under nitrogen. A portion of hydrogen peroxide, added to bleach the product, failed to do this, so a second portion was added. The explosion which occurred later was attributed to accumulation of highly concentrated excess peroxide and traces of organic material (either volatile or entrained) in the reflux dividing head [3]. Only a few ppm of impurities are necessary to cause autodetonation of very concentrated hydrogen peroxide [4]. A survey of explosion hazards likely to be encountered from reagent use of hydrogen peroxide in organic chemistry is given [5]. Charging too concentrated peroxide to organic materials in surfactant manufacture caused a pressure increase which burst a pipe, starting a catalogue of mishap which destroyed the plant. Laboratory investigations showed that 10% peroxide by weight with ‘organic’ material could cause pressure surges in excess of the plant rating. The editor suspects this is base catalysed peroxide decomposition, the organic material being tertiary amines for amine oxide surfactants [6].

Organic materials, Sulfuric acid

Analytical Methods Committee, *Analyst*, 1976, **101**, 62–66

Advantages and potential hazards in the use of mixtures of 50% hydrogen peroxide solution and conc. sulfuric acid to destroy various types of organic materials prior to analysis are discussed in detail. The method is appreciably safer than those using perchloric and/or nitric acids, but the use of an adequate proportion of sulfuric acid with a minimum of peroxide is necessary to avoid the risk of explosive decomposition. The method is not suitable for use in pressure-digestion vessels (PTFE lined steel bombs), in which an explosion occurred at 80°C.

Other reactants

Yoshida, 1980, 85–86

MRH values calculated for a wide range of 25 oxidisable or catalytically active materials are given.

Oxygenated compounds, Water

1. Monger, J. M. *et al.*, *J. Chem. Eng. Data*, 1961, **6**(1), 23

2. Shanley, E. S., *Chem. Eng. News*, 1979, **57**(21), 43

MRH Acetaldehyde 6.27/21, acetic acid 5.27/31, acetone 6.26/19, ethanol 6.19/19, formaldehyde 6.44/31, formic acid 3.30/58, methanol 5.98/34, 2-propanol 6.32/16, propionaldehyde 6.44/19

The explosion limits have been determined for liquid systems containing hydrogen peroxide, water and acetaldehyde, acetic acid, acetone, ethanol, formaldehyde, formic acid, methanol, 2-propanol or propionaldehyde, under various types of initiation [1]. In general, explosive behaviour is noted where the ratio of hydrogen peroxide to water is >1, and if the overall fuel–peroxide composition is stoichiometric, the explosive power and sensitivity may be equivalent to those of glyceryl nitrate [2].

See Alcohols, above

2-Phenyl-1,1-dimethylethanol, Sulfuric acid

1. Winstein, S. *et al.*, *J. Amer. Chem. Soc.*, 1967, **89**, 1661

2. Hedaya, E. *et al.*, *Chem. Eng. News*, 1967, **45**(43), 73

3. Hiatt, R. R. *et al.*, *J. Org. Chem.*, 1963, **28**, 1893

Directions given [1] for the preparation of 2-phenyl-1,1-dimethylethyl hydroperoxide by adding sulfuric acid to a mixture of the alcohol and 90% hydrogen peroxide are wrong [2] and will lead to explosion [3]. The acidified peroxide (30–50% solution is strong enough) is preferably added to the alcohol, with suitable cooling and precautions [2].

See Alcohols, above, *also* Oxygenated compounds, etc., below

See 2-(4-Chlorophenyl)-1,1-dimethylethyl hydroperoxide

α -Phenylselenoketones

Reich, H. J. *et al.*, *J. Amer. Chem. Soc.*, 1975, **97**, 5442

During conversion of the selenoketone to the nor-phenylselenoenones with hydrogen peroxide, care is necessary to control the strongly exothermic reaction.

Under no circumstances should oxidation of amounts above 5 mmol be effected by adding the full amount of peroxide before oxidation has started.

Phosphorus MRH (Yellow) 7.78/27

Anon., *J. R. Inst. Chem.*, 1957, **81**, 473

If yellow or red phosphorus is incompletely immersed while undergoing oxidation in hydrogen peroxide solutions, heating at the air–solution interface can ignite the phosphorus and lead to a violent reaction. Such behaviour has been observed with peroxide solutions above 30% (110 vol) concentration.

Phosphorus(V) oxide MRH 1.59/tr.

Toennies, G., *J. Amer. Chem. Soc.*, 1937, **59**, 555

The extremely violent interaction of phosphorus(V) oxide and conc. hydrogen peroxide to give peroxomonophosphoric acid may be moderated by using acetonitrile as a diluent.

Poly(acetoxyacrylic acid lactone), Poly(2-hydroxyacrylic acid)

Anon., *Jahresber.*, 1978, 75

The complex peroxide formed by stirring the polymeric compounds with 30% peroxide solution and sodium hydroxide at 5°C for an hour, according to a patented procedure, was precipitated with methanol. After intensive vacuum drying at 1.3 mbar without heating, the product suddenly exploded.

See other POLYPEROXIDES

Polymer residues

Merrifield, 1988, 7

A jacketed polymerisation vessel had become coated internally by a build up of polymer residues, and the vessel was being cleaned by treatment with aqueous hydrogen peroxide. To 5000 l of water in the vessel was added 150 kg of 27 wt% peroxide solution, and the vessel was heated by application of 10 bar steam (180°C) to the jacket. After a few minutes an explosion occurred, attributed to spontaneous ignition of a mixture of oxygen from the decomposing peroxide and monomer vapours produced by depolymerisation of the residue on the heated walls of the vessel.

See other GAS EVOLUTION INCIDENTS

2-Propanol

Merrifield, 1988, 7

A large volume (11.25 m³) of mixed fatty acids was to be bleached by treatment with successive portions of 50 wt% hydrogen peroxide. 2-Propanol (450 l) was added to the acids (to improve the mutual solubility of the reactants). The first 20 l portion of peroxide (at 51°C) was added, followed after 1 min by a second portion. Shortly afterwards an explosion occurred, which was attributed to spontaneous ignition of a 2-propanol vapour–oxygen mixture formed above the surface of the liquid. Oxygen is almost invariably evolved from hydrogen peroxide reactions, and volatile flammable solvents are therefore incompatible components in peroxide systems.

See Alcohols, above

See other GAS EVOLUTION INCIDENTS

Sulfides

Editor's personal experience

Addition of 30% hydrogen peroxide to moderately strong sodium sulfide solutions produced superheating below the surface, followed by cavitation explosions sufficient to break laboratory glass beakers.

Sulfuric acid

1. Analytical Methods Committee, *Analyst*, 1967, **92**, 404
2. Dobbs, D. A. *et al.*, *Chem. Eng. News*, 1990, **68**(17), 2
3. Erickson, T. V., *Chem. Eng. News*, 1990, **68**(33), 2
4. Anon., *Environment, Safety & Health Bull.*, 1993, **93**(2), 1
5. Stuart, R., Internet, 1991

Evaporation of mixtures of excess 50% hydrogen peroxide solutions with sulfuric acid (10:1) leads to loud but non-shattering explosions of the peroxomonosulfuric acid formed [1]. A freshly-made mixture of equal volumes of sulfuric acid and 30% hydrogen peroxide, "Piranha Solution" used for cleaning glass frits, exploded violently on storage. Explosions have also been experienced when cleaning frits, when contamination with organic solvent was suspected. The mixture will, in any case, slowly evolve oxygen to pressurise sealed containers [2,3]. Piranha solution also burst its container after addition of nitric acid [4]. A similar mixture, also containing small quantities of lithium sulfate and selenious acid and intended as a digesting agent for soil samples, burst its container after twelve days undisturbed in a refrigerator. Such solutions are best not stored and, if stored, should not be sealed [5].

See Organic materials, Sulfuric acid, above

Tetrahydrothiophene

Koppel, H. C., *Chem. Eng. News*, 1974, **52**(39), 3

Preparative reactions involving oxidation of tetrahydrothiophene to the sulfoxide by slow addition of 37% peroxide solutions exploded violently on 3 occasions. No explanation is apparent, and similar reactions had been run uneventfully over a period of 10 years.

Tin(II) chloride

MRH 1.21/85

Vickery, R. C. *et al.*, *Chem. & Ind.*, 1949, 657

Interaction is strongly exothermic, even in solution. Addition of peroxide solutions of above 3 wt% strength causes a violent reaction.

Unsaturated compounds

Swern, 1971, Vol. 2, 432

In commercial-scale epoxidation of unsaturated organic compounds, 50–70% hydrogen peroxide is added during 2–8 h to the stirred mixture of unsaturated compound, aliphatic acid and acid catalyst at 50–80°C, the rate of addition being

dependent on heat-transfer capacity of the reaction system. When 70% peroxide is being used, care must be taken to avoid the dangerous detonable region by ensuring that an exotherm has occurred before more than 25% of the hydrogen peroxide is added.

See Carboxylic acids, *also* Organic compounds, *also* Oxygenated compounds, all above

Vinyl acetate

Anon., *ABCM Quart. Safety Summ.*, 1948, **19**, 18

Vinyl acetate had been hydroxylated by treatment with excess hydrogen peroxide in presence of osmium tetroxide catalyst. An explosion occurred while excess vinyl acetate and solvent were being removed by vacuum distillation. This was attributed to the presence of peracetic acid, formed by interaction of excess hydrogen peroxide with acetic acid produced by hydrolysis of the vinyl acetate.

Waste treatment

1. Anon., 1989, private communication.
2. Anon., *Eur. Chem. News*, 1992, **58**(1537), 34
3. Wehrum, W. L., *Process Safety Progr.*, 1993, **12**(4), 199
4. Pratt, T. H., *Process Safety Progr.*, 1993, **12**(4), 203
5. Anon., *Loss Prev. Bull.*, 1991, (102), 17

In the interests of greater safety and environmental consciousness, a chemical factory which had previously used sodium hypochlorite to treat alkaline aqueous wastes containing methanethiol, other volatile sulfur compounds, methanol and possibly dimethyl ether, changed to 30% hydrogen peroxide. Shortly after, an explosion in the headspace shattered the vessel, killing a worker and breaking a pipe containing flammable solvent, leading to a fire which destroyed the factory. It appears that decomposition of peroxide in alkali had produced an oxygen enriched headspace, allowing the organic vapours therein to enter explosive limits despite a nitrogen purge. The actual source of ignition is unknown [1]. Another fatal explosion when treating effluent, NO containing off gas, with peroxide is reported, presumably by creation and ignition of an oxygen rich atmosphere containing organic vapours [2]. A third waste treatment incident, possibly fuelled by acetone is reported [3]. An ignition mechanism, by means of electrostatic effects from bursting oxygen bubbles, is postulated [4]. A related accident which killed 17 men was due to an explosion in a waste tank from a propylene oxide/styrene plant. This contained hydrocarbons over aqueous waste and evolved oxygen from decomposition of 'light peroxide' by-product decomposition. The headspace was nitrogen flushed. During maintenance the nitrogen flow was cut off for 34 hours, but oxygen readings remained low. Minutes after nitrogen purging was restarted the tank exploded. It is thought the oxygen meter may have been ill positioned [5].

See Methanol, Copper sulfate, Cyanide residues, above

Wood

1. *MCA Case History No. 1626*
2. *MCA Case History No. 1648*

Leakage from drums of 35% hydrogen peroxide onto a wooden pallet caused ignition of the latter when it was moved. Combustion, though limited in area, was fierce and took some time to extinguish [1]. Leakage of 50% peroxide onto supporting pallets under polythene sheeting led to spontaneous ignition and a fierce fire. Contact of 50% peroxide with wood does not usually lead to spontaneous ignition, but hot weather, dry wood (possibly catalytically contaminated) and the thermal insulation of the cover may have contributed to ignition [2].

See other PACKAGING INCIDENTS

Yeast

Bodenbaugh, J. H. *et al.*, *J. Chem. Educ.*, 1988, **66**(5), 461–463


Hydrogen peroxide solution may be catalytically decomposed to give oxygen gas safely using dry yeast powder. (The active component is enzymatic.)

See other GAS EVOLUTION INCIDENTS

See other OXIDANTS, PEROXIDES

4478. Zinc hydroxide

[20427-58-1]



Chlorinated rubber

See CHLORINATED RUBBER: metal oxides or hydroxides

4479. Sulfuric acid

[7664-93-9]



(*MCA SD-20*, 1963); *NSC 325*, 1982; *NSC 210*, 1967 (oleum); *FPA H31*, 1974; *HCS 1980*, 878, 879 (solutions), 702–704 (oleum); *RSC Lab. Hazards Safety Data Sheet No. 33*, 1985

The MCA Data Sheet covers safe handling procedures for all grades of sulfuric acid from fuming (oleum) through conc. to diluted acid. Coverage of the other series is as shown.

For an unusual incident involving static charge generation by sulfuric acid drops
See Hydrogen: Oxygen, Sulfuric acid

Acetaldehyde

MRH 1.34/57

See Acetaldehyde: Sulfuric acid

Acetone, Nitric acid

See Nitric acid: Acetone, etc.

Acetonitrile, Sulfur trioxide

See Acetonitrile: Sulfuric acid

Acrylonitrile MRH 2.26/52
See Acrylonitrile: Acids

Alkyl nitrates MRH Methyl nitrate 6.53/99+
See ALKYL NITRATES: LEWIS ACIDS

2-Aminoethanol
Nolan, 1983, Case history 101
The two components are immiscible and must be stirred to effect conversion at 110°C to 2-aminoethyl hydrogen sulfate. A shift change led to the base being added to the acid without agitation. When the agitator was started later, reaction of the two cold and viscous layers of acid and base proceeded explosively.
See other AGITATION INCIDENTS

Ammonium iron(III) sulfate dodecahydrate
Clark, R. E. D., private comm., 1973
A few dense crystals heated with sulfuric acid exploded, owing to the exotherm in contact with water liberated as the crystals disintegrated.

Aniline, Glycerol, Nitrobenzene
See Quinoline

Benzyl alcohol MRH 1.13/52
Grignard, 1960, Vol. 5, 1005
A mixture of the alcohol with 58% sulfuric acid decomposes explosively at about 180°C.

Bromine pentafluoride
See Bromine pentafluoride: Acids, etc.

tert-Butyl-*m*-xylene, Nitric acid
See Nitric acid: *tert*-Butyl-*m*-xylene, etc.

1-Chloro-2,3-epoxypropane
See 1-Chloro-2,3-epoxypropane: Sulfuric acid

4-Chloronitrobenzene, Sulfur trioxide
1. Grewer, T., *Chem. Ing. Tech.*, 1975, **47**, 233
2. Vervalin, C. H., *Hydrocarbon Process.*, 1976, **55**(9), 323
3. *See entry* SELF-ACCELERATING REACTIONS
4. Nolan, 1983, Case history 100
DTA shows that the reaction mixture from sulfonation of the nitro compound in 20% oleum, containing 35 wt% of 2-chloro-5-nitrobenzenesulfonic acid, exhibits 2 exothermic stages at 100 and 220°C, respectively, the latter being violently rapid. The adiabatic reaction mixture, initially at 89° attained 285°C with boiling after 17 h. At 180° the induction period was about 20 min [1]. Sulfonation of 4-chloronitrobenzene with 65% oleum at 46°C led to a runaway decomposition

H₂O₄S

reaction in a 2000 l vessel. The original process using 20% oleum was less sensitive to heating rate and temperature. Knowledge that the reaction could be dangerous above 50°C had not been applied [2]. More thermoanalytical work has established safer operating conditions with a greater margin for remedial action [3]. In a further incident, the sulfonation (an established process) had been effected with 65% oleum at below 80°C, and the reaction mixture had been heated to, and held at 115°C, and then cooled to 80°C. Half of the batch was quenched in water with cooling to crystallise the product, but the remaining half in the reactor overheated and exploded. This was attributed to backflow of water from the quench tank into the reactor, and the exotherm from reaction of water with oleum led to exothermic decomposition of the sulfonic acid at around 160°C [4].

See THERMAL STABILITY OF REACTION MIXTURES AND SYSTEMS

See other INDUCTION PERIOD INCIDENTS, SULFONATION INCIDENTS

Copper

Campbell, D. A., *School Sci. Rev.*, 1939, **20**(80), 631

The generation of sulfur dioxide by reduction of sulfuric acid with copper is considered too dangerous for a school experiment.

2-Cyano-4-nitrobenzenediazonium hydrogen sulfate

See 2-Cyano-4-nitrobenzenediazonium hydrogen sulfate: Sulfuric acid

2-Cyano-2-propanol

1. *Occupancy Fire Record, FR 57-5*, 5, Boston, NFPA, 1957

2. Kirk-Othmer, 1967, Vol. 13, 333

Addition of sulfuric acid to the cyano-alcohol caused a vigorous reaction which pressure-ruptured the vessel [1]. This seems likely to have been caused by insufficient cooling to prevent dehydration of the alcohol to methacrylonitrile and lack of inhibitors to prevent exothermic polymerisation of the nitrile [2].

Cyclopentadiene

MRH 1.75/40

Wilson, P. J. *et al.*, *Chem. Rev.*, 1944, **34**, 8

It reacts violently with charring, or explodes in contact with conc. sulfuric acid.

See Nitric acid: Hydrocarbons

Cyclopentanone oxime

See Cyclopentanone oxime: Sulfuric acid

1,3-Diazidobenzene

See 1,3-Diazidobenzene: Acids

Dichloromethane, Ethanol, Nitrate or nitrite

Dickie, R. J., private comm., 1981

The residues from a cortisol assay procedure (5 cc dichloromethane, 2.5 cc of a fluorescent reagent in 15:85 ethanol-sulfuric acid) were added to a 500 cc bottle and screw capped. After a 90 s delay, the bottle burst violently and brown fumes were seen. It was surmised that a nitrate or nitrite contaminant in the bottle had

liberated oxides of nitrogen or nitric acid which had oxidised the organics exothermically.

See Nitric acid: Alcohols, *also* Dinitrogen tetraoxide: Halocarbons

Diethylamine

Anon., *CISHC Chem. Safety Summ.*, 1984, **55**(218), 34–35

Diethylamine fumes from a reactor were usually absorbed in a glass scrubber through which sulfuric acid was circulated, but an unresolved fault in the level sensor caused the acid circulation pump to operate intermittently. While the pump was not running, amine fumes condensed in the dip pipe, forming a solid crust (of the sulfate) which allowed a quantity of condensed amine to accumulate out of contact with the acid. When the pump was restarted, the neutralisation exotherm was sufficient to shatter the scrubber and distort the mesh guard around it.

See *other* NEUTRALISATION INCIDENTS

Dimethoxydinitroanthraquinone

See *mixo*-Dimethoxydinitroanthraquinone: Sulfuric acid

4-Dimethylaminobenzaldehyde

MCA Case History No. 2101

During the preparation of a solution of the aldehyde in dilute sulfuric acid, the latter should be prepared before addition of the aldehyde. An attempt to prepare the solution by adding a slurry of the basic aldehyde in a little water to conc. sulfuric acid caused the stoppered flask to burst from the large exotherm generated by this procedure.

2,5-Dinitro-3-methylbenzoic acid, Sodium azide

See 2,5-Dinitro-3-methylbenzoic acid: Oleum, Sodium azide

1,5-Dinitronaphthalene, Sulfur

See 1,5-Dinitronaphthalene: Sulfur, etc.

Ethoxylated nonylphenol

Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 73

Breakage of glassware allowed 92% sulfuric acid into a heating bath containing hot Synperonic NX (nearly anhydrous ethoxylated nonylphenol, a liquid surfactant). Hydrolysis of the latter lowered the flashpoint and the bath ignited (possibly from formation of dioxane, flash point 12°C).

Hexalithium disilicide

See Hexalithium disilicide: Acids

Hydrofluoric acid

Bentzinger, von R. *et al.*, *Praxis Naturwiss. Chem.*, 1987, 36, 37

A student was attempting to prepare anhydrous hydrogen fluoride by dehydrating aqueous 60% hydrogen fluoride solution with conc. sulfuric acid. Addition of 200 ml of sulfuric acid to 500 ml of hydrofluoric acid in a 1 l copper flask led to

H₂O₄S

a rumbling noise, then a fountain from the flask neck of hot mixed acids which severely corroded the window glass and the floor tiles.

See Water, below

See other CORROSION INCIDENTS, GLASS INCIDENTS

Hydrogen peroxide MRH 1.59/99+

See Hydrogen peroxide: Sulfuric acid

Mercury nitride

See Mercury nitride: Alone, or Sulfuric acid

Metal acetylides or carbides MRH Calcium carbide 4.35/56

1. Mellor, 1946, Vol. 5, 849

2. *MCA SD-50*, 1963

Monocaesium and monorubidium acetylides ignite with conc. sulfuric acid [1].

Other carbides are hazardous in contact [2].

Metal chlorates MRH Sodium chlorate 1.30/99+

See METAL CHLORATES: ACIDS, ALSO METAL HALOGENATES: METALS, ETC.

Metal perchlorates

See METAL PERCHLORATES: SULFURIC ACID

4-Methylpyridine

Anon., *Loss. Prev. Bull.*, 1979, (029), 130–131

The base was added to 60 l of acid to convert it to the salt, but malfunction of the swivelling blade glass stirrer caused no agitation and the liquid base remained layered above the conc. acid. Switching off the stirrer caused it to become effective and the whole of the contents erupted from the 100 l flask as an acid mist.

See other AGITATION INCIDENTS, NEUTRALISATION INCIDENTS

Nitramide

See Nitric amide: Sulfuric acid

Nitric acid, Organic materials

See Nitric acid: Organic materials, Sulfuric acid

Nitric acid, Toluene

See Nitric acid: Hydrocarbons (reference 7)

Nitroaryl bases and derivatives MRH Nitroaniline 3.30/99+

1. Hodgson, J. F., *Chem. & Ind.*, 1968, 1399; private comm., 1973

2. Poshkus, A. C. *et al.*, *J. Appl. Polym. Sci.*, 1970, **14**, 2049–2052

A series of 2- and 4-nitroaniline derivatives and analogues when heated with conc. sulfuric acid to above 200°C undergo, after an induction period, a vigorous reaction. This is accompanied by gas evolution which produces up to a 150-fold increase in volume of a solid foam, and is rapid enough to be potentially hazardous if confined.

2-Nitroaniline reacts almost explosively [1], and 4-nitroaniline, 4-nitroacetanilide, aminonitrobiphenyls, aminonitronaphthalenes and their various derivatives [2], as well as some nitro-N-heterocycles [1,2], also react vigorously. 4-Nitroanilinium sulfate and 4-nitroaniline-2-sulfonic acid and its salts also generate foams when heated without sulfuric acid. The mechanism is not clear, but involves generation of a polymeric matrix foamed by sulfur dioxide and water eliminated during the reaction [1].

See Diethyl sulfate: 2,7-Dinitro-9-phenylphenanthridine

See other INDUCTION PERIOD INCIDENTS, GAS EVOLUTION INCIDENTS

Nitrobenzene

See Nitrobenzene: Sulfuric acid

3-Nitrobenzenesulfonic acid

See 3-Nitrobenzenesulfonic acid: Sulfuric acid

Nitro-explosives

Gamezo, V. N. *et al.*, *J. Phys. IV*, 1995, **5** (C4), 395

The effect of sulphuric acid and oleum on the explosive properties of nitromethane, dinitrotoluene, trinitrobenzene and trinitrotoluene was studied. They are considerably sensitised.

See OTHER NITRO-DERIVATIVES, ABOVE AND BELOW

Nitromethane

See Nitromethane: Acids

N-Nitromethylamine

See N-Nitromethylamine: Sulfuric acid

4-Nitrotoluene

See 4-Nitrotoluene: Sulfuric acid

Other reactants

Yoshida, 1980, 393

MRH values calculated for 15 combinations with various materials are given.

Permanganates

MRH Potassium permanganate 0.50/76

Interaction produces the powerful oxidant, permanganic acid.

See Permanganic acid: Organic materials

Potassium permanganate: Sulfuric acid

Phosphorus

MRH (Red) 2.64/28

1. Mellor, 1940, Vol. 8, 786

2. Zschalich, A. *et al.*, *Chem. Abs.*, 1993, 11532b

3. Urben, P. G., personal experience

White phosphorus ignites in contact with boiling sulfuric acid or its vapour [1].

An explosion hazard exists at industrial scale under milder conditions [2]. Even

H₂O₄S

red phosphorus reacts vigorously with sulfuric acid in the presence of iodine or iodides, to produce hydrogen sulfide, from about 80°C [3].

Phosphorus(III) oxide

See Tetrphosphorus hexaoxide: Sulfuric acid

Poly(silylene)

See Poly(silylene): Oxidants

Potassium

See Potassium: Sulfuric acid

Potassium *tert*-butoxide

See Potassium *tert*-butoxide: Acids

3-Propynol

See 3-Propynol: Sulfuric acid

Silver permanganate

See Silver permanganate: Sulfuric acid

Silver peroxochromate

Sodium

See Sodium: Acids

MRH 3.89/31

Sodium carbonate

See Sodium carbonate: Sulfuric acid

Sodium tetrahydroborate

See Sodium tetrahydroborate: Acids

Sodium thiocyanate

Meyer, 1977, 39

Violent exotherm on contact, carbonyl sulfide being evolved.

Steel

MARS Database, 1998, short report 021

Steel is normally considered a safe material for contact with (reasonably pure) sulphuric acid. However this report is of a fatal explosion when working on a sulphuric acid tank with cutting torches, attributed to hydrogen in the headspace arising from corrosion of the metal ascribed to poor maintenance.

Sucrose

Meeks, E. G., *School Sci. Rev.*, 1979, **61**(215), 281–283

In the demonstration of the foaming carbonisation of sucrose by sulfuric acid, it has been found that about half of the gas produced is carbon monoxide and not

carbon dioxide as previously supposed. (Other sugars and carbohydrates may be expected to behave similarly.)

Tetramethylbenzenes

1. Birch, S. F. *et al.*, *J. Amer. Chem. Soc.*, 1949, **71**, 1364

2. Smith, L. I., *Org. Reactions*, 1942, Vol. 1, 382

Sulfonation of the mixed isomers of 1,2,3,5- and 1,2,4,5-tetramethylbenzenes was too violent [1], for shaking in a closed glass vessel as originally described [2].

See other SULFONATION INCIDENTS

1,2,4,5-Tetrazine

See 1,2,4,5-Tetrazine: Sulfuric acid

Thallium(I) azidodithiocarbonate

See Thallium(I) azidodithiocarbonate: Sulfuric acid

1,3,5-Trinitrosohexahydro-1,3,5-triazine

See 1,3,5-Trinitrosohexahydro-1,3,5-triazine: Sulfuric acid

Water

Mellor, 1947, Vol. 120, 405–408

Dilution of conc. sulfuric acid by water is vigorously exothermic and must be effected by adding acid to water to avoid local boiling. Mixtures of sulfuric acid and excess snow form powerful freezing mixtures. Fuming sulfuric acid (oleum, containing sulfur trioxide) reacts violently with water.

See 3-Nitrobenzenesulfonic acid: Sulfuric acid

Zinc iodide

Pascal, 1962, Vol. 5, 168

Interaction with the conc. acid is violent.

See other INORGANIC ACIDS, OXIDANTS

4480. Poly(dihydroxydioxodisilane)

[]

$(\text{H}_2\text{O}_4\text{Si}_2)_n$

$(\text{Si}(\text{O})(\text{OH})\text{Si}(\text{O})(\text{OH}))_n$


1. Sorbe, 1968, 127

2. Mellor, 1940, Vol. 6, 230

The dry compound explodes on impact or heating [1]. This polymeric product of hydrolysis of hexahalo- or hexaethoxy-disilane decomposes with more or less violence if heated in air (when it ignites) or in a test tube (when it explodes) [2].

See related NON-METAL OXIDES

4481. Peroxomonosulfuric acid
[7722-86-3]



1. Edwards, J. O., *Chem. Eng. News*, 1955, **33**, 3336
2. Brauer, 1963, Vol. 1, 389

A small sample had been prepared from chlorosulfuric acid and 90% hydrogen peroxide; the required acid phase was separated and stored at 0°C overnight. After warming slightly, it exploded [1]. The handling of large amounts is dangerous owing to the possibility of local overheating (e.g., from contact with moisture) and explosive decomposition [2].

Acetone

1. *MCA Case History No. 662*
2. Bayer, A., *Ber.*, 1900, **33**, 858

Accidental addition of a little acetone to the residue from wet-ashing a polymer with mixed nitric–sulfuric acids and hydrogen peroxide caused a violent explosion [1]. The peroxyacid would be produced under these conditions, and is known to react with acetone to produce the explosive acetone peroxide [2], but direct oxidation of acetone may have been responsible.

Alcohols

1. Toennies, G., *J. Amer. Chem. Soc.*, 1937, **59**, 552
2. Merrifield, 1988, 6, 16

Contact of the acid with *sec*- or *tert*-alcohols, even with cooling, may lead to violent explosions [1]. The autodetonation region which exists for the hydrogen peroxide–sulfuric acid –2-propanol or –*tert*-butanol system is shown on a triangular diagram [2].

Aromatic compounds

- Sidgwick, 1950, 939
Mixtures with aniline, benzene, phenol etc., explode.

Catalysts

- Mellor, 1946, Vol. 5, 483–484
The 92% acid is decomposed explosively on contact with massive or finely divided platinum, manganese dioxide or silver. Neutralised solutions of the acid also froth violently on treatment with silver nitrate, lead dioxide or manganese dioxide.

Fibres

- Ahrle, H., *Z. Angew. Chem.*, 1909, **23**, 1713
Wool and cellulose are rapidly carbonised by the 92% acid, while cotton ignites after a short delay.
See other INORGANIC ACIDS, OXIDANTS, PEROXOACIDS

4482. Peroxodisulfuric acid
[13445-49-3]

H₂O₈S₂

HOSO₂OOSO₂OH

Aromatic amines, Sulfuric acid

Nielsen, A. T. *et al.*, *J. Org. Chem.*, 1980, **45**, 2341–2347

The acid, prepared from 90–98% hydrogen peroxide and oleum or 100% sulfuric acid, is one of the most powerful known oxidants and its use for oxidising aromatic amines to nitro compounds has been studied. Some mono- di- and tri-amines are destroyed exothermically with violent fume-off. Precautions for use are detailed.

Organic liquids

1. D'Ans, J. *et al.*, *Ber.*, 1910, **43**, 1880; *Z. Anorg. Chem.*, 1911, **73**, 1911

2. Sidgwick, 1950, 938

A very powerful oxidant; uncontrolled contact with aniline, benzene, ethanol, ether, nitrobenzene or phenol may cause explosion [1]. Alkanes are slowly carbonised [2].

See other OXIDANTS, PEROXOACIDS

†4483. Hydrogen sulfide
[7783-06-4]

H₂S

H₂S

(MCA SD-36, 1968); NSC 284, 1977; FPA H78, 1979; HCS 1980, 553 (cylinder); RSC Lab. Hazards Safety Data Sheet No. 35, 1985

1,2-Bis(2-azidoethoxy)ethane, Ethanol

See 1,2-Bis(2-azidoethoxy)ethane: Ethanol, etc.

4-Bromobenzenediazonium chloride

See DIAZONIUM SULFIDES AND DERIVATIVES

Copper, Oxygen

Merz, V. *et al.*, *Ber.*, 1880, **13**, 722

Addition of powdered copper to a 1:2 mixture of hydrogen sulfide and oxygen causes the metal to become incandescent and ignite the explosive mixture.

See Metals, below

Metal oxides MRH Copper oxide 1.05/70, chromium trioxide 2.05/88, lead dioxide 0.54/78, manganese dioxide 0.84/94, sodium peroxide 3.01/70 Mellor, 1947, Vol. 10. 129, 141

Hydrogen sulfide is rapidly oxidised, and may ignite in contact with a range of metal oxides, including barium peroxide, chromium trioxide, copper oxide, lead dioxide, manganese dioxide, nickel oxide, silver(I) oxide, silver(II) oxide, sodium peroxide, and thallium(III) oxide. In the presence of air, contact with mixtures of calcium oxide or barium oxide with mercury oxide or nickel oxide may cause vivid incandescence or explosion.

See Rust, below

Metals

Mellor, 1947, Vol. 10, 140; 1943, Vol. 11, 731

A mixture with air passed over copper powder may attain red heat. Finely divided tungsten glows red hot in a stream of hydrogen sulfide.

See Copper, Oxygen, above *See also* Sodium: Sulfides

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Other reactants

Yoshida, 1980, 390

MRH values calculated for 11 combinations with oxidants are given.

Oxidants

MRH values show % of oxidant

See Metal oxides, above; Oxygen, and Rust, both below

Bromine pentafluoride: Hydrogen-containing materials MRH 3.01/70

Chlorine trifluoride: Hydrogen-containing materials MRH 2.05/88

Chromium trioxide: Hydrogen sulfide

Copper chromate oxide: Hydrogen sulfide

Dibismuth dichromium nonaoxide: Hydrogen sulfide

Dichlorine oxide: Oxidisable materials MRH 3.22/91

Fluorine: Hydrogen sulfide MRH 9.29/72

Heptasilver nitrate octaoxide: Alone, or Sulfides

Lead(II) hypochlorite: Hydrogen sulfide

Lead(IV) oxide: Hydrogen sulfide MRH 0.54/78

Mercury(I) bromate: Hydrogen sulfide

Nitric acid: Non-metal hydrides MRH 4.43/74

Oxygen difluoride: Combustible gases

Perchloryl fluoride: Hydrocarbons, etc.

Silver bromate: Sulfur compounds

Sodium peroxide: Hydrogen sulfide MRH 3.01/70

Oxygen

Gray, P. *et al.*, *J. Chem. Soc., Faraday Trans. 1*, 1974, **70** 2338–2350

During an investigation of the spontaneously explosive oxidation of near-stoichiometric gas mixtures in the range 280–360°C, extensive self-heating was observed before ignition occurred. The second and third ignition limits were also investigated.

See Oxidants, above

Rust

Mee, A. J., *School Sci. Rev.*, 1940, **22**(85), 95

Hydrogen sulfide may ignite if passed through rusty iron pipes.

See Metal oxides, above

Silver fulminate

See Silver fulminate: Hydrogen sulfide

Soda-lime

1. Mellor, 1947, Vol. 10, 140
2. Bretherick, L., *Chem. & Ind.*, 1971, 1042

Interaction is exothermic, and if air is present, incandescence may occur with freshly prepared granular material. Admixture with oxygen causes a violent explosion [1]. Soda-lime, used to absorb hydrogen sulfide, will subsequently react with atmospheric oxygen and especially carbon dioxide (from the solid coolant) with a sufficient exotherm in contact with moist paper wipes (in a laboratory waste bin) to cause ignition [2]. Spent material should be saturated with water before separate disposal. Mixture analogous to soda-lime, such as barium hydroxide with potassium or sodium hydroxides, also behave similarly [1].

See other SELF-HEATING AND IGNITION INCIDENTS

See other NON-METAL HYDRIDES, NON-METAL SULFIDES

†4484. **Hydrogen disulfide**

[13465-07-1]

H₂S₂

HSSH

Alkalies

Bloch, I. *et al.*, *Ber.*, 1908, **41**, 1977

Rather more flammable than hydrogen trisulfide, it is decomposed violently by alkalies.

See other NON-METAL HYDRIDES, NON-METAL SULFIDES

4485. Hydrogen trisulfide

[13845-23-3]

H₂S₃

HSSSH

Benzenediazonium chloride

King, W. B. *et al.*, *J. Amer. Chem. Soc.*, 1932, **54**, 3073

Addition of the dry diazonium salt to the crude liquid sulfide causes explosively violent interaction. Slow addition of the sulfide to the cooled salt, or dilution with a solvent, moderates the reaction.

See other DIAZONIUM SULFIDES AND DERIVATIVES

Metal oxides

Mellor, 1947, Vol. 10, 159

Contact with copper oxide, lead(II) oxide, lead(IV) oxide, mercury(II) oxide, tin(IV) oxide or iron(II,III) oxide causes violent decomposition and ignition. Dry powdered silver oxide causes an explosion.

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Pentanol

Mellor, 1947, Vol. 10, 158

Interaction is explosively violent.

Potassium permanganate
See Potassium permanganate: Hydrogen trisulfide
See other NON-METAL HYDRIDES, NON-METAL SULFIDES

†4486. Hydrogen selenide
[7783-07-5]



It is moderately exothermic (ΔH_f° (g) +85.8 kJ/mol, 1.06 kJ/g).

Oxidants

See Hydrogen peroxide: Hydrogen selenide
Nitric acid: Non-metal hydrides
See other ENDOTHERMIC COMPOUNDS, NON-METAL HYDRIDES
See related NON-METAL SULFIDES

4487. Poly(silylene)
[32078-95-8]



Brauer, 1963, Vol. 1, 682
The dry solid ignites in air.

Oxidants

Bailar, 1973, Vol. 1, 1352
It ignites in contact with conc. nitric acid and explodes with sulfuric acid.
See related SILANES

†4488. Hydrogen telluride
[7783-09-7]



It is moderately endothermic (ΔH_f° (g) +154.4 kJ/mol, 1.19 kJ/g).

Nitric acid

See Nitric acid: Non-metal hydrides
See other ENDOTHERMIC COMPOUNDS, METAL HYDRIDES *See related* NON-METAL HYDRIDES

4489. Thorium dihydride
[16689-88-6]



1. Hartmann, I. *et al.*, *Rept. Invest. No. 4835*, Washington, US Bur. Mines, 1951
2. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

Extremely pyrophoric when powdered [1], and is also a significant dust explosion hazard [2].

See other METAL HYDRIDES, PYROPHORIC MATERIALS

4490. Titanium dihydride

[7704-98-5]

H₂Ti

TiH₂

1. Alekseev, A. G. *et al.*, *Chem. Abs.*, 1974, **81**, 123688
2. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

Pyrophoricity and detonation behaviour of titanium hydride powders of various particle sizes were studied in comparison with those of titanium metal powders [1]. Maximum dust explosion pressures of 8.2 bar, with a maximum rate of rise of 816 bar/s have been recorded [2].

See Zirconium (reference 6)

See other METAL HYDRIDES, PYROPHORIC MATERIALS

4491. Titanium–zirconium hydride

[57425-12-4]

H₂Ti.H₂Zr

TiH₂.ZrH₂

See Zirconium (reference 6)

See other METAL HYDRIDES

4492. Zinc hydride

[14018-82-7]

H₂Zn

ZnH₂

Barbaras, G. D. *et al.*, *J. Amer. Chem. Soc.*, 1951, **73**, 4587

Fresh samples reacted slowly with air, but aged and partly decomposed samples (containing finely divided zinc) may ignite in air.

Acids, Water

Ashby, E. C. *et al.*, *Inorg. Synth.*, 1977, **17**, 9

Hydrolysis with water occurs slowly, but is violent with aqueous acids.

See other METAL HYDRIDES

4493. ‘Zirconium hydride’

[7704-99-6]

H₂Zr

ZrH₂

1. Mellor, 1941, Vol. 7, 114
2. Sidgwick, 1950, 633

3. Baumgart, H., *Chem. Abs.*, 1985, **10** 3, 73412
 4. Kuroda, E. *et al.*, *Chem. Abs.*, 1997, **127**, 150727x

The product of sorbing hydrogen on to hot zirconium powder burns with incandescence when heated in air [1,2]. Ignition and combustion properties of pyrophoric zirconium and zirconium hydride powder ($<10 \mu$), and safety measures are discussed [3]. Ignition energies for zirconium hydrides, ZrH_x ; $x = 0 - 2$, in air have been studied in Japan [4].

See Zirconium (reference 6)

See other METAL HYDRIDES

4494. Dipotassium triimidotellurite

[]

H₃K₂N₃Te



1. Schmitz-Du Mont, O. *et al.*, *Angewand. Chem. (Int.)*, 1967, **6**, 1071

2. Chivers, T. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(22), 2549

Colourless crystals, exploding on contact with water unless benzene damp. Ammonolysis converts it to the still more explosive tritellurium tetranitride [1]. Highly explosive [2].

See also N-S COMPOUNDS

4495. Lanthanum hydride

[13864-01-2]

H₃La



1. Mackay, 1966, 66

It ignites in air.

See other METAL HYDRIDES

4496. Lithium hydrazide

[37067-42-8]

H₃LiN₂



Hydrazine and its Derivatives, Schmidt, F. W., New York, Wiley, 1984, 75

Merely smouldering on contact with air, it is claimed to be a safe metal hydrazide.

See other METAL HYDRAZIDES

†**4497. Ammonia**

[7664-41-7]

H₃N



(MCA SD-8, 1960); NSC 251, 1979; FPA H13, 1973; HCS 1980, 139 (cylinder), 140 (25% solution); RSC Lab. Hazards Safety Data Sheet No. 37, 1985

1657

1. MCA SD-8, 1960
2. Anon., *ABCM Quart. Safety Summ.*, 1950, **45**, 14
3. Anon., *ABCM Quart. Safety Summ.*, 1974, **45**, 14
4. Unpublished observations, 1968–1974
5. Henderson, D., *Ammonia Plant Safety*, 1975, **17**, 132–134
6. Krivulin, V. N. *et al.*, *Chem. Abs.*, 1976, **84**, 124132
7. Harris, G. F. P. *et al.*, *Proc. 6th Symp. Chem. Proc. Haz. Ref. Plant Des.*, 29–38, Symp. Ser. 49, Rugby, IChE, 1977
8. McRae, M. H., *Plant/Oper. Progr.*, 1987, **6**(1), 17–19
9. Klem, T. J., *Ammonia Plant Saf.*, 1987, (26), 145–149
10. Andersson, B. O., *Hazards XI*, 15, Symp. Ser. **124**, Rugby(UK), IChE, 1991.
11. Andersson, B. O., *Ammonia Plant Safety*, 1991, **31**, 5
12. Carlsson, T. O. *et al.*, *Sci. Tech Froid*, 1996, (3), 277

Although there is a high lower explosive limit in air and ignition is not easy, there is a long history of violent gas–air explosions in refrigeration practice, in which ammonia previously was used widely [1]. Mixtures of ammonia and air lying within the explosive limits can occur above aqueous solutions of certain strengths. Welding operations on a vessel containing aqueous ammonia caused a violent explosion [2]. Several incidents involving sudden ‘boiling’ of conc. ammonia solution (d , 0.880, 35 wt%) have occurred when screw-capped winchesters are opened [3,4]. These are attributable to supersaturation of the solution with gas, caused by increase in temperature subsequent to preparation and bottling. The effect is particularly marked with winchesters filled in winter and opened in summer. Usually boiling-off begins immediately the cap is loosened, and if this is done carefully using gloves and eye-protection in a fume cupboard or under local extract ventilation, no great problems arise [4]. However a bottle which had been gently and successfully opened in this way without boiling-off subsequently erupted violently when the surface of the liquid was disturbed by a pipette [3]. This hazard may be avoided by using the slightly less concentrated solution (d , 0.990, 25 wt%) where this is technically acceptable.

There is an account of an incident involving combustion/explosion in the free space of a weak ammonia liquor tank where detailed examination revealed no evidence for an ignition source [5]. The existence of ammonia–air mixtures able to burn only in sufficiently large enclosed volumes was established [6]. Previous data on flammability has been summarised and extended by determination of flammability characteristics of ammonia admixed with dry air and oxygen-deficient air at temperatures from ambient up to 400°C. Similar work with added water vapour at 80°C shows that aqueous solutions of below 5% ammonia content do not produce flammable vapours at any temperature, and that above 49°C no flammable vapours are produced by ammonia solution of any concentration. Some data on ignition energy and explosibility are also given [7]. Two recent US incidents serve as a reminder of the considerable explosive potential of ammonia–air mixtures in confined volumes. An ice-cream factory with ammonia refrigeration plant which had been installed over 50 years previously had not been upgraded to meet modern engineering practice and electrical fire codes. There were no check valves or oil separators in the compressor discharge lines, and the plant was lit by unprotected

H₃N

tungsten filament lamps. During the night an ammonia leak developed in the basement, and a few minutes later there was a violent explosion which largely destroyed the building. The major devastation prevented identification of the cause of the leak or of the ignition source, and it was postulated that oil mist may have been involved in initiating the explosion [8]. During an attempt to isolate a leak in the anhydrous ammonia refrigeration system in a cold store warehouse, an electric fork lift truck was to be used to replace a faulty valve. The truck crashed into an internal wall and the large explosion which followed no doubt involved spark ignition of the flammable atmosphere [9]. Both incidents revealed a widespread belief that ammonia is non-flammable, possibly arising from the US classification of anhydrous ammonia in cylinders as 'a non-flammable gas'. In Europe, it is realistically classified as flammable, as also are highly concentrated aqueous solutions. With returning interest in it as a domestic refrigerant, studies of ignition energies and flame propagation in mixtures with air have been performed [12]

A report is given of explosion of a tank containing 7000 t of liquid ammonia, in Lithuania, consequent upon thermal disequilibrium and 'bumping' of the liquid [10,11]. The liberated ammonia caught fire in the open air, the first time this is thought to have been observed.

See Nitric acid: Manufacture hazard

Air, Hydrocarbons

Kalkert, N. *et al.*, *Chem. Ing. Tech.*, 1979, **51**, 895

Explosion limits have been estimated for mixtures containing C₁–C₃ hydrocarbons.

Air, Rust

Verduijn, W. D., *Process Safety Progress*, 1996, **15**(2), 89; Verduijn, W. D. *Ammonia Plant Safety*, 1996

An account is given of a sequence of mishaps starting with a burst pipe in a nitric acid plant, apparently due to pre-ignition by rust catalysis of an ammonia/air mixture at 220°C and 11 bar.

Boron halides

Sidgwick, 1950, 380

The boron halides react violently with ammonia.

Calcium

See Calcium: Ammonia

1-Chloro-2,4-dinitrobenzene

See 1-Chloro-2,4-dinitrobenzene: Ammonia

Chlorine azide

See Chlorine azide: Alone, or Ammonia

Chloroformamidinium nitrate

See Chloroformamidinium nitrate: Alone, or Amines, etc.

2-, or 4-Chloronitrobenzene

See 2-Chloronitrobenzene: Ammonia or 4-Chloronitrobenzene: Ammonia

1,2-Dichloroethane

491M, 1975, 36

Liquid ammonia and the solvent may explode when mixed. (It is possible this was a liquefied gas (physical) explosion, rather than an exothermic chemical reaction.)

See LIQUEFIED GASES

Ethanol, Silver nitrate

See Silver nitrate: Ammonia, Ethanol

Ethylene oxide

See Ethylene oxide: Ammonia

Germanium derivatives

Paschenko, I. S. *et al.*, *Chem. Abs.*, 1976, **84**, 8568

'Causes of the Explosiveness of Gas-Liquid Mixtures during Neutralisation of Germanium-containing Solutions by Ammonia' (title only translated).

See Germanium imide

Gold(III) chloride

See Gold(III) chloride: Ammonia, etc.

Halogens, or Interhalogens

Ammonia either reacts violently, or produces explosive products, with all 4 halogens and some of the interhalogens.

See Bromine: Ammonia

Bromine pentafluoride: Hydrogen-containing materials

Chlorine: Nitrogen compounds

Chlorine trifluoride: Hydrogen-containing materials

Fluorine: Hydrides

Iodine: Ammonia

Heavy metals

Ammonia is capable of reacting with some heavy metal compounds (silver, gold, mercury) to produce materials, some of uncertain constitution, which may explode violently when dry.

See Gold(III) chloride: Ammonia

Mercury: Ammonia

Potassium triamidothallate ammoniate

Silver azide: Ammonia

Silver chloride: Ammonia

Silver nitrate: Ammonia

Silver(I) oxide: Ammonia

See N-METAL DERIVATIVES

Iodine, Potassium

Staley, S. W. *et al.*, *Tetrahedron*, 1975, **31**, 1133

During the reductive cleavage of cyclopolyenes with potassium in liquid ammonia, the intermediate anionic species are quenched with iodine–pentane mixtures. The possibility of formation of explosive nitrogen triiodide and the need for precautions are stressed.

Magnesium perchlorate

See Magnesium perchlorate: Ammonia

Mercury

1. Marsch, H. D., *Ammonia Plant Safety*, 1991, **31**, 8

2. Wilhelm, S. M., *Ammonia Plant Safety*, 1991, **31**, 20

An explosion rupturing an ammonia separator (still) in an ammonia production unit, probably because mercury vapour from geological sources entered with hydrogen syngas originating from natural gas and reacted to give explosive nitride deposits. The separator remains cracked when scraped [1]. For a more academic study of the effects of mercury on ammonia plants, including embrittlement and corrosion, as well as explosive deposits [2].

See other CORROSION INCIDENTS

See Mercury: Ammonia

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Other reactants

Yoshida, 1980, 36

MRH values calculated for 16 combinations, largely with oxidants, are given.

Oxidants

MRH values show % of oxidant

Maslonka, F., *Chem. Abs.*, 1978, **88**, 193935

The flammability and explosion susceptibility of ammonia is reviewed, with thermochemical data, explosion limits for various oxidants, and explosion parameters, both for the gas and its conc. aqueous solutions.

See Ammonium peroxodisulfate: Ammonia, etc.

Chromium trioxide: Ammonia

Chromyl chloride: Ammonia

Dichlorine oxide: Oxidisable materials

Dinitrogen tetraoxide: Ammonia

Hydrogen peroxide: Nitrogenous bases

MRH 5.86/75

Nitric acid: Ammonia

MRH 5.10/69

Nitrogen oxide: Ammonia

Nitryl chloride: Inorganic materials

Oxygen (Liquid): Ammonia, etc.

Potassium chlorate: Ammonia

MRH 4.31/78

‘Trioxxygen difluoride’: Various materials

Oxygen, Platinum
See Oxygen (Gas): Ammonia, Platinum

Ozone
See Ozone: Ammoniacal vapours

Pentaborane(9)
See Pentaborane(9): Ammonia

Silver compounds
See Silver azide: Ammonia
Silver chloride: Ammonia
Silver nitrate: Ammonia
Silver(I) oxide: Ammonia, etc.

Stibine
See Stibine: Ammonia

Tellurium halides
See Tellurium tetrabromide: Ammonia
Tellurium tetrachloride: Ammonia

Tetramethylammonium amide
See Tetramethylammonium amide: Ammonia

Thiocarbonyl azide thiocyanate
See Thiocarbonyl azide thiocyanate: Ammonia, etc.

Sulfinyl chloride
See Sulfinyl chloride: Ammonia

Thiotriethiazyl chloride
See Thiotriethiazyl chloride: Ammonia
See other INORGANIC BASES, NON-METAL HYDRIDES

4498. Hydroxylamine
[7803-49-8]



HCS 1980, 557

1. Ashford, J. S., private comm., 1967
2. Brauer, 1963, Vol. 1, 502
3. Vervalin, C. H., *Hydrocarbon Proc. Petr. Ref.*, 1963, **42**(2), 174
4. Rüst, 1948, 302
5. Bailar, 1973, Vol. 2, 272

H₃NO

The base was being prepared by distilling a mixture of hydroxylamine hydrochloride and sodium hydroxide in methanol under reduced pressure, and a violent explosion occurred towards the end of distillation [1], probably owing to an increase in pressure above 53 mbar. It explodes when heated under atmospheric pressure [2]. Traces of hydroxylamine remaining after reaction with acetonitrile to form acetamide oxime caused an explosion during evaporation of solvent. Traces can be removed by treatment with diacetyl monoxime and ammoniacal nickel sulfate, forming nickel dimethylglyoxime [3]. An account of an extremely violent explosion towards the end of vacuum distillation had been published previously [4]. Anhydrous hydroxylamine is usually stored at 10°C to prevent internal oxidation–reduction reactions which occur at ambient temperature [5].

See other REDOX REACTIONS

Aldehyde (unspecified), Pyridine

MCA Guide, 1972, 300

During the preparation of an unspecified aldoxime, the ethanolic reaction mixture was extracted with ether and the extract was concentrated by vacuum distillation from a bath at 70–80°C. Towards the end of distillation violent decomposition occurred (probably of traces of unreacted hydroxylamine in the extract).

See above

Barium oxide

MRH 3.86/tr.

See Oxidants, below

Carbonyl compound (unspecified), Pyridine

Anon., *Chem. Processing* (Chicago), 1963, **26**(24), 30

During preparation of an unspecified oxime, the carbonyl compound, pyridine, hydroxylamine hydrochloride and sodium acetate were heated in a stainless steel autoclave. At 90°C a sudden reaction caused a pressure surge to 340 bar, when the bursting disk failed. The reaction had been run previously and uneventfully on one-tenth scale in a glass lined autoclave.

See Metals, below

Copper(II) sulfate

Mellor, 1940, Vol. 8, 290

Hydroxylamine ignites with the anhydrous salt (possibly owing to heat of coordination).

Hydrogenation catalyst, Acid, Fuel

Riesthuis, P. *et al.*, *J. Loss Prev. Process Ind.*, 1997, **10**(10), 67

In the presence of precious metal hydrogenation catalyst, hydroxylamine salts may disproportionate and form dinitrogen monoxide. Such a mixture is present in a process whereby the hydroxyamine is formed by hydrogenation of nitrate. An explosion in the degassing line, after a period of abnormal operation, was attributed to nitrous oxide build-up. Fuel, in the form of hydrogen and methane diluent, was already present.

See Dinitrogen oxide

Metals

MRH Calcium, sodium, zinc all 3.85/tr.

1. Mellor, 1940, Vol. 8, 292
2. Sorbe, 1968, 158

Sodium ignites in contact with hydroxylamine alone, but reacts smoothly in ether solution to give *N*-sodium hydroxylamide which may be pyrophoric in air. Calcium reacts to give the bis(hydroxylamide) which explodes at 180°C. Finely divided zinc either ignites or explodes when warmed in contact with hydroxylamine [1]. Zinc and hydroxylamine form the bis(hydroxylamide) solvated with hydroxylamine (3 mol), an explosive compound [2].

See also OXIMES

Other reactants

Yoshida, 1980, 286

MRH values calculated for 16 combinations with various materials are given.

Oxidants

MRH values below references show % of oxidant

1. Brauer, 1963, Vol. 1, 502
2. Mellor, 1930, Vol. 3, 670; 1940, Vol. 8, 287–294

MRH Barium peroxide 3.85/tr., calcium hypochlorite 3.77/52, lead dioxide 1.92/64, potassium dichromate 3.85/tr., potassium permanganate 2.80/49, sodium hypochlorite 3.97/53

Hydroxylamine is a powerful reductant, particularly when anhydrous, and if exposed to air on a fibrous extended surface (filter paper) it rapidly heats by aerobic oxidation. It explodes in contact with air above 70°C [1]. Barium peroxide will ignite aqueous hydroxylamine, while the solid ignites in dry contact with barium oxide, barium peroxide, lead dioxide and potassium permanganate, but with chlorates, bromates and perchlorates only when moistened with sulfuric acid. Contact of the anhydrous base with potassium dichromate or sodium dichromate is violently explosive, but less so with ammonium dichromate or chromium trioxide. Ignition occurs in gaseous chlorine, and vigorous oxidation occurs with hypochlorites.

See other REDOX REACTIONS

Phosphorus chlorides

MRH Phosphorus(III) & (V) chlorides both 3.85/tr.

Mellor, 1940, Vol. 8, 290

The base ignites in contact with phosphorus trichloride and phosphorus pentachloride.

See also HYDROXYLAMINIUM SALTS

See other N–O COMPOUNDS, REDUCANTS

4499. Amidosulfuric acid (Sulfamic acid)

[5329-14-6]

H₃NO₃S

H₂NSO₂OH

HCS 1980, 869

Chlorine

See Chlorine: Nitrogen compounds (reference 7)

Metal nitrates or nitrites

See METAL AMIDOSULFATES

See other INORGANIC ACIDS, N-S COMPOUNDS

4500. Sodium hydrazide

[13598-47-5]

$\text{H}_3\text{N}_2\text{Na}$

NaNHNH_2

Air, or Ethanol, or Water

1. Mellor, 1940, Vol. 8, 317

2. Kauffmann, T., *Angew. Chem. (Intern. Ed.)*, 1964, **3**, 342

Contact with traces of air, ethanol or water causes a violent explosion [1], as does heating to 100°C [2].

See other METAL HYDRAZIDES

4501. Azidosilane

[13847-60-4]

$\text{H}_3\text{N}_3\text{Si}$

N_3SiH_3

Preparative hazard

See Azidogermane: Fluorosilane

See other NON-METAL AZIDES See related SILANES

4502. Hydrogen hexaazidothallate(III)

[63338-74-9]

$\text{H}_3\text{N}_{18}\text{Tl}$

$\text{H}_3[\text{Tl}(\text{N}_3)_6]$

Krommes, P. *et al.*, *J. Organomet. Chem.*, 1977, **131**, 418

Not more than 200–500 mg portions of this extremely heat- and impact-sensitive explosive should be prepared.

See related METAL AZIDES See other N-METAL DERIVATIVES

4503. Phosphinic acid ('Hypophosphorous acid')

[6303-21-5]

$\text{H}_3\text{O}_2\text{P}$

$\text{HOP}(\text{O})\text{H}_2$

1. Mellor, 1940, Vol. 8, Suppl. 3, 614

2. Albright and Wilson (Manufacturer's safety sheet)

3. *MARS Database*, 1998, short report 181

The explosion hazard associated with the usual laboratory preparation from white phosphorus and alkali may be avoided by an alternative method involving oxidation of phosphine with an aqueous iodine solution [1]. The commercial 50% solution reacts violently with oxidants. On heating, it decomposes rapidly above 100°C evolving phosphine, which is liable to explode with air. It is recommended it

be kept below 50°C [2]. During vacuum concentration of a dilute solution on small industrial scale, a crust formed, overheated and decomposed, pressurising the system with phosphine which escaped and produced two explosions on mixing with air, causing modest damage to the factory installations [3].

Mercury(II) nitrate

See Mercury(II) nitrate: Phosphinic acid

Mercury(II) oxide

MRH 0.71/86

Mellor, 1940, Vol. 4, 778

The redox reaction is explosive.

See other REDOX REACTIONS

Other reactants

Yoshida, 1980, 343

MRH values calculated for 6 combinations with various other materials are given.

See other INORGANIC ACIDS, REDUCANTS

4504. Phosphonic acid (Phosphorous acid)

[13598-36-2]

H₃O₃P

P(OH)₃ or HP(O)(OH)₂

1. Partington, 1946, 614

2. Albright and Wilson, (Manufacturer's safety sheet)

It decomposes at around 200°C with evolution of phosphine which may ignite [1].

Yellow deposits may form in aqueous solution which are spontaneously flammable on drying [2].

See Phosphorus trichloride: Carboxylic acids

See other INORGANIC ACIDS, REDUCANTS

4505. Orthophosphoric acid

[7664-38-2]

H₃O₄P

P(O)(OH)₃

(MCA SD-7, 1958); NSC 674, 1978; HCS 1980, 742, 743, MCA SD-70, 1958

Although it is not an oxidant, it is an active acid which, because of its low equivalent wt. and availability in high concentration and density (90% soln. is 50 N), can develop a large exotherm on neutralisation. Handling precautions are detailed.

Chlorides, Stainless steel

Piekarz, J., *Chem. in Can.*, 1961, **13**(4), 40–41

Presence of traces of chloride ion in technical 75% phosphoric acid in a closed storage tank caused corrosion and liberation of hydrogen which later exploded at a sparking contact.

See other CORROSION INCIDENTS, GAS EVOLUTION INCIDENTS

Nitromethane

See Nitromethane: Acids, etc.

Sodium tetrahydroborate

See Sodium tetrahydroborate: Acids

See other INORGANIC ACIDS

4506. Peroxomonophosphoric acid

[13598-52-5]

H₃O₅P



Preparative hazard

See Hydrogen peroxide: Phosphorus(V) oxide

Coal, Potassium permanganate

See Potassium permanganate: Coal, etc.

Organic material

Castrantas, 1965, 5


The 80% solution causes ignition when dropped onto organic material.

See other INORGANIC ACIDS, PEROXOACIDS

4507. Hydroxyaqua(oxo)diperoxorheniumVII

[]

H₃O₇Re



Herrmann, W. A. *et al.*, *Chem. Eur. J.*, 1996, **2**(2), 168

A highly explosive red/orange solid, obtained from rhenium trioxide and hydrogen peroxide reacting in hexamethylphosphoramide as solvent, was tentatively assigned this structure.

See other INORGANIC PEROXIDES

†4508. Phosphine

[7803-51-2]

H₃P



HCS 1980, 745 (cylinder); RSC Lab. Hazards Safety Data Sheet No. 49⁰, 1986

1. Rüst, 1948, 301
2. Sidgwick, 1950, 729
3. Mellor, 1940, Vol. 8, 811, 814
4. McKay, H. A. C., *Chem. & Ind.*, 1964, 1978
5. Anon., *Lab. Haz. Bull.*, 1982, (3), item 120

6. Green, A. R. *et al.*, *Dev. Agric. Eng.*, 1984, **5** (Controlled Atmos. Fumig. Grain Storage), 433–449

7. Ethenakis, V. M. *et al.*, *Sol. Cells*, 1987, **22**, 305–317

Liquefied phosphine (a mildly endothermic compound, ΔH_f° (g) +22.8 kJ/mol, 0.67 kJ/g) can be detonated by powerful initiation [1]. Pure phosphine does not spontaneously ignite in air below 150°C unless it is thoroughly dried, when it ignites in cold air. The presence of traces (0.2%) of diphosphane in phosphine as normally prepared causes it to ignite spontaneously in air, even at below –15°C [2]. Traces of oxidants promote pyrophoricity [3]. Phosphine, generated by action of water on calcium phosphide, was dried by passage through towers packed with the latter. Soon after refilling the generator (but not the towers) and starting purging with argon, a violent explosion occurred. This was attributed to the air, displaced from the generator by argon, reacting explosively with dry phosphine present in the drying towers, possibly catalysed by the orange-yellow polyphosphine formed on the surface of calcium phosphide. Fresh calcium phosphide in both generator and drying towers, with separate purging, is recommended [4]. During vacuum transfer of the supposedly pure gas, air leaked into the cold-trap system and caused an explosion [5]. Lower flammability limit of pure phosphine in moist air (0.39 vol% water vapour) at 1037 mbar was 2.1% at 10° and 1.85% at 50°C. Presence of ammonia may tend to reduce the limit value [6]. Potential hazards involved in use of phosphine in manufacture of photovoltaic cells are reviewed [7].

See Oxygen, below, *also* Phosphonium iodide: Potassium hydroxide

Boron trichloride

See Boron trichloride: Phosphine

Dichlorine oxide

MRH 5.36/84

See Dichlorine oxide: Oxidisable materials

Halogens

MRH Chlorine 2.13/89

Mellor, 1940, Vol. 8, 812

Ignition occurs on contact with chlorine or bromine or their aqueous solutions.

Metal nitrates

MRH Silver nitrate 3.77/87

Mellor, 1941, Vol. 3, 471; 1940, Vol. 4, 993

Passage of phosphine into silver nitrate solution causes ignition or explosion, depending on the gas rate. Mercury(II) nitrate solution gives a complex phosphide, explosive when dry.

Nitric acid

MRH 7.78/74

See Nitric acid: Non-metal hydrides

Nitrogen trichloride

MRH 3.64/91

Other reactants

Yoshida, 1980, 342

MRH values calculated for 10 combinations with oxidants are given.

Oxidants MRH values below reference
Mellor, 1940, Vol. 8, 811, 814
MRH Dinitrogen monoxide 6.99/84, dinitrogen trioxide 9.25/75, nitrogen oxide 9.71/78, nitrous acid 7.11/79
Pure phosphine is rendered pyrophoric by traces of dinitrogen trioxide, nitrous acid, or similar oxidant.
See Dichlorine oxide; Halogens; Metal nitrates; Nitric acid, all above, Oxygen, below

Oxygen
1. Fischer, E. O. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1968, **7**, 136
2. Mellor, 1971, Vol. 8, Suppl. 3, 281–282
Even small amounts of oxygen present in phosphine give an explosive mixture, in which autoignition occurs at low pressures [1]. The effects of other materials upon the explosive interaction have been studied [2].
See other ENDOTHERMIC COMPOUNDS, NON-METAL HYDRIDES

4509. Plutonium(III) hydride
[15457-77-9]



H₃Pu

Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64
The hydride (and the metal) when finely divided are spontaneously flammable, and burning causes a specially dangerous contamination problem, in view of the radioactive and toxic hazards.

Water
Bailar, 1973, Vol. 5, 150
Rapid addition of water often causes ignition of the hydride.
See other METAL HYDRIDES

†4510. Stibine
[7803-52-3]



H₃Sb

Mellor, 1939, Vol. 9, 394
During evaporation of liquid stibine at -17°C , a relatively weak and isothermal explosive decomposition may occur. Gaseous stibine at ambient temperature may propagate an explosion from a hot spot on the retaining vessel wall, and it autocatalytically decomposes, sometimes explosively, at 200°C .
It is moderately endothermic (ΔH_f° (g) +145.2 kJ/mol, 1.16 kJ/g).

Ammonia
Mellor, 1939, Vol. 9, 397
A heated mixture explodes

Oxidants

See Chlorine: Non-metal hydrides

Nitric acid: Non-metal hydrides

Ozone: Stibine

See related NON-METAL HYDRIDES See other ENDOTHERMIC COMPOUNDS, METAL HYDRIDES

4511. Uranium(III) hydride

[13598-56-6]

H₃U

UH₃

Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64

The dry powdered hydride readily ignites in air.

Halocarbons

Bailar, 1973, Vol. 5, 150

Contact of the hydride with halogen-containing solvents may lead to explosive interaction.

Water

Bailar, 1973, Vol. 5, 150

Rapid addition of water often causes ignition of the hydride.

See other METAL HYDRIDES

4512. Ammonium iodide

[12027-06-4]

H₄IN

NH₄I

Bromine trifluoride

See Bromine trifluoride: Ammonium halides

See related METAL HALIDES

4513. Ammonium iodate

[13446-09-8]

H₄INO₃

NH₄IO₃

Anon., *ABCM Quart. Safety Summ.*, 1955, **26**, 24

Violent decomposition occurred on touching with a scoop (which presumably was contaminated). A similar batch of material contained less than 100 ppm of periodate.

Other reactants

Yoshida, 1980, 375

MRH values calculated for 15 combinations with oxidisable materials are given.

See related METAL OXOHALOGENATES See other IODINE COMPOUNDS

4514. Ammonium periodate
[13446-09-8]

H₄INO₄

NH₄IO₄

1. Anon., *Chem. Eng. News*, 1951, **29**, 1770
2. Mellor, 1940, Vol. 2, 408

It exploded while being transferred by scooping [1]. The sensitivity towards heat, but not abrasive impact, was known previously [2].

See related METAL OXOHALOGENATES

See other IODINE COMPOUNDS

4515. Phosphonium iodide
[12125-09-6]

H₄IP

PH₄I

Oxidants

Mellor, 1947, Vol. 8, 827

Bromates, chlorates or iodates ignite in contact with phosphonium iodide at ambient temperature if dry, or in presence of acid to generate bromic acid, etc. Ignition also occurs with nitric acid, and reaction with dry silver nitrate is very exothermic. Interaction with antimony pentachloride at ambient temperature proceeds explosively.

See other REDOX REACTIONS

Potassium hydroxide

Anon., *Univ. Safety Assoc., Safety Newsletter*, 1983–1984

Phosphine, generated by action of the alkali on phosphonium iodide, was shown to be pure by mass spectrometry. During a second preparation, an air leak developed during cold-trap transfer and an explosion occurred. Presence of diphosphane was suspected.

See Phosphine (reference 2)

Water

See Phosphorus: Hydriodic acid

See other REDUCANTS

4516. Potassium tetrahydrozincate
[34324-17-9]

H₄K₂Zn

K₂[H₄Zn]

Ashby, E. C. *et al.*, *Inorg. Chem.*, 1971, **10**, 2486

Extremely reactive, pyrophoric in air.

See other COMPLEX HYDRIDES

4517. Magnesium phosphinate

[7783-17-7]



Mellor, 1940, Vol. 8, 885

On heating it evolves phosphine which ignites in air.

See other METAL PHOSPHINATES, REDUCANTS**4518. Ammonium permanganate**

[13446-10-1]



1. Mellor, 1942, Vol. 12, 302

2. Pavlychenko, M. M. *et al.*, *Chem. Abs.*, 1962, **57**, 2897e

The dry material is friction-sensitive and explodes at 60°C in air [1]. Under vacuum, decomposition becomes explosive above 100°C, and sparks will also initiate decomposition.

See other OXOSALTS OF NITROGENOUS BASES (reference 1), OXIDANTS**4519. Manganese(II) phosphinate**

[10043-84-2]



Mellor, 1940, Vol. 8, 889

It explodes above 200°C.

See other HEAVY METAL DERIVATIVES, METAL PHOSPHINATES, REDUCANTS**†4520. Hydrazine**

[302-01-2]

*HCS 1980*, 536 (anhydrous), 537 (hydrate); *RSC Lab. Hazard Data Sheet No. 14*, 19831. Troyan, J. E., *Ind. Eng. Chem.*, 1953, **45**, 26082. Day, A. C. *et al.*, *Org. Synth.*, 1970, **50**, 423. *491M*, 1975, 1994. Mitchell, J. W., *Anal. Chem.*, 1980, **52**, 774–7765. Bellerby, J. M. *et al.*, *Chem. Abs.*, 1982, **96**, 2225356. Bunker, R. L. *et al.*, *Prog. Astronaut. Aeronaut.*, 1991, **133**, 3247. *Hydrazine and its Derivatives*, Schmidt, F. W., New York, Wiley, 1984It is rather endothermic (ΔH_f° (l) +50.4 kJ/mol, 1.57 kJ/g) and readily decomposed.

Hydrazine is an endothermic compound and the vapour is exceptionally hazardous in that once it is ignited, it will continue to burn by exothermic decomposition in complete absence of air or other oxidant [1]. Distillation of anhydrous hydrazine (prepared by dehydrating hydrazine hydrate with solid sodium

H₄N₂

hydroxide) must be carried out under nitrogen to avoid the possibility of an explosion if air is present [2]. If spread with an extended surface (as a thin film, or absorbed on porous or fibrous material), anhydrous hydrazine may autoxidise with sufficient heat generation and retention to cause ignition. Presence of traces of metal catalysts (as in some cellulosic materials) will greatly accelerate this effect [3]. Procedures for safe handling and purification of aqueous hydrazine solutions are detailed [4], and autoxidation processes of hydrazine and its methylated derivatives have been reviewed [5]. The mechanism by which hydrazine can be detonated during pumping operations has been studied [6]. A comprehensive monograph includes sections on decomposition and other hazard-related aspects of the preparation and applications of hydrazine [7].

See Barium oxide; Cotton waste, Heavy metals; Metal catalysts, all below

Barium oxide, or Calcium oxide

1. Gmelin, 1935, Syst. 4, 318
2. *ABCM Quart. Safety Summ.*, 1950, **21**, 18

The residue from dehydrating hydrazine with barium oxide slowly decomposes exothermically in daylight and finally explodes [1]. Dehydration of 95% hydrazine by boiling with calcium oxide under nitrogen caused violent explosions on 2 occasions. It was concluded that use of calcium or barium oxides for this purpose is potentially dangerous, and that boiling under reduced pressure may be advisable to lower the liquid temperature. General precautions in handling hydrazine are discussed [2].

See Sodium hydroxide, below

Benzeneseleninic acid or anhydride

See Benzeneseleninic acid: Hydrazine derivatives

Calcium

Hydrazine and its Derivatives, Schmidt, F. W., New York, Wiley, 1984, 75

Use of excess calcium metal, or its solution in ammonia (presumable calcamide) to dry hydrazine leads to formation of calcium hydrazide and explosion. Calcium hydride and carbide are, apparently, safe drying agents

See METAL HYDRAZIDES

Carbon dioxide, Stainless steel

Bellerby, J. M., *J. Haz. Mat.*, 1983, **7**, 187–197

The rate of decomposition of hydrazine in stainless steel vessels (which is accompanied by corrosion) is directly proportional to carbon dioxide concentration over 20 ppm and below 250 ppm. The species responsible for the catalytic decomposition is not one of the expected corrosion products, iron(II) carbazate or its nickel or chromium(II) analogues.

See other CATALYTIC IMPURITY INCIDENTS, CORROSION INCIDENTS

1-Chloro-2,4-dinitrobenzene

See 1-Chloro-2,4-dinitrobenzene: Hydrazine hydrate

2-Chloro-5-methylnitrobenzene

See 2-Chloro-5-methylphenylhydroxylamine

Cotton waste, Heavy metals

Orme, R. J., private comm., 1980

In a factory manufacturing organo-iron and organo-manganese catalysts, use of cotton waste to mop up a spill of dilute hydrazine solution led to a spontaneous fire later. The fire was attributed to onset of rapid metal-catalysed decomposition of the hydrazine after sufficient water had evaporated from the waste, the traces of heavy metals originating from dust contamination, etc.

See Metal catalysts, below

Dicyanofurazan

See Dicyanofurazan: Nitrogenous bases

(Difluoroamino)difluoroacetonitrile

See (Difluoroamino)difluoroacetonitrile: Hydrazine

Dinitrogen oxide

See Dinitrogen oxide: Hydrazine

Iodine pentoxide

Speeds, S. J. *et al.*, US Pat. Appl. 844 082, 1977

The oxide is used to effect rapid decomposition of hydrazine in a high-temperature gas generator which produces exit pressures (of nitrogen, hydrogen and ammonia) of 550 bar.

Metal catalysts

1. Mellor, 1940, Vol. 8, 317; 1967, Vol. 8, Suppl. 2.2, 83
2. Audrieth, L. F. *et al.*, *Ind. Eng. Chem.*, 1951, **43**, 1774
3. Troyan, J. E., *Ind. Eng. Chem.*, 1953, **45**, 2608
4. Blumenthal, J. L. *et al.*, US Pat. 3 846 339, 1974
5. Rao, K. R. *et al.*, *Trans. Powder Metall. Assoc. India*, 1975, **2**, 64–66
6. Armstrong, W. E. *et al.*, US Pat. 4 124 538, 1978
7. Bellerly, J. M., *J. Haz. Mat.*, 1986, **13**(1), 57–60

In contact with metallic catalysts (platinum black, Raney nickel) hydrazine is catalytically decomposed, yielding ammonia, hydrogen and nitrogen. With conc. hydrazine the reaction may be violent, and the ammonia and hydrogen evolved could be ignited by particles of dry catalyst [1]. Measures to prevent autoxidation of hydrazine catalysed by traces of copper are discussed, including displacement of air from transportation containers with nitrogen [2]. Catalytic decomposition is effected by iron oxide, molybdenum-stabilised stainless steel, molybdenum and its oxides, and finely divided solids. Handling procedures are discussed [3]. A rhenium–alumina catalyst causes immediate ignition of undiluted hydrazine [4]. The effect upon ignition delay of thermal treatment and particle size of powdered transition metal catalysts has been studied [5]. Iridium or iridium–ruthenium on alumina also effect catalytic decomposition of hydrazine [6]. Soluble chromium and/or manganese species do not increase the rate of decomposition of anhydrous hydrazine, but presence of traces of acid as well leads to significantly increased rates [7].

Metal salts

1. Mellor, 1941, Vol. 7, 430; 1940, Vol. 8, 318; 1967, Vol. 8, Suppl. 2.2, 882
2. Traynham, J. G. *et al.*, *J. Org. Chem.*, 1967, **32**, 3285
3. McCoy, P. O., *Chem. Eng. News*, 1970, **48**(48), 9

Several of its complexes with metallic salts are unstable, including those of basic cadmium perchlorate (highly explosive), cadmium nitrite (explosive), copper chlorate (explodes on drying without heat), manganese nitrate (ignites at 150°C), mercury(I) chloride, mercury(II) chloride, mercury(I) nitrate, mercury(II) nitrate (all are explosive) and tin(II) chloride (explodes on heating): other examples are given [1]. There is a published method for reducing *cis-trans*-cyclododecadiene to the *cis*-monoene with diazene ('diimide') generated in situ from hydrazine, copper(II) salts and air [2]. During a modified reaction sequence using crude diene, much sludge containing polymer, copper(II) acetate and hydrazine was produced. When filtered off in air, the sludge heated and eventually glowed. Use of purified diene and filtration under nitrogen avoids the problem [3].

See Trihydrazinealuminium perchlorate

See also AMMINOMETAL OXOSALTS

Methanol, Nitromethane

See Nitromethane: Hydrazine, Methanol

Other reactants

Yoshida, 1980, 283

MRH values calculated for 16 combinations with oxidants are given.

Oxidants

MRH values below references show % of oxidant

1. *Haz. Chem Data*, 1975, 166
2. Mellor, 1941, Vol. 3, 137; 1940, Vol. 8, 313–319; 1943, Vol. 11, 234
3. *ASESB Op. Incid. Rept. No. 105*
4. Troyan, J. E., *Ind. Eng. Chem.*, 1953, **45**, 4608
5. Wannagat, U. *et al.*, *Monatsh. Chem.*, 1969, **97**, 1157–1162
6. Ephraim, 1939, 651

MRH Ammonium nitrate 3.97/84, chromium trioxide 3.01/81, hydrogen peroxide 6.36/68, nitric acid 5.84/61, potassium dichromate 1.59/tr., sodium chlorate 6.07/69, sodium nitrate 3.18/68

Hydrazine is a powerful, endothermic reducing agent, and its interaction with oxidants may be expected to be violent if unmoderated, as in rocket propulsion systems (often hypergolic). Mixtures of hydrazine vapour with air are explosive over a very wide range (4.7—100%) and ignition temperature can be very low (24°C on a rusty iron surface). When mixed with, or spilt on, highly porous materials (asbestos, cloth, earth, wood), autoxidation in air may proceed fast enough to cause ignition. Contact with hydrogen peroxide or nitric acid may cause ignition with conc. reagents and with dinitrogen tetroxide is hypergolic [1]. Contact with chromate salts or chromium trioxide causes explosive decomposition. Copper oxide or lead oxide cause a vigorous decomposition, while dropping hydrazine hydrate on to mercury oxide can cause an explosion. This may be owing to known oxidation of hydrazine to hydrogen azide by two electron transfer, and

formation of explosive mercury azide [2]. Contact of hydrazine with the explosive *N*, 2,4,6-tetranitroaniline caused ignition [3]. Contact of iron oxide, chlorates or peroxides may lead to violent reactions [4]. The ignition delay with fuming nitric acid was 8 ms, explosion also occurring [5]. Contact with potassium permanganate leads to explosion [6].

See Lithium perchlorate: Hydrazine

Potassium peroxodisulfate: Hydrazine, etc.

Silver(I) oxide: Ammonia, etc.

Sodium perchlorate: Hydrazine

MRH 5.86/65

'Trioxigen difluoride': Various materials

See *N*-HALOIMIDES

See ROCKET PROPELLANTS

See also Silylhydrazines: Oxidants

Potassium

Gmelin, 1935, Syst. 4, 318

Explosive interaction.

See Sodium hydrazide

Rust

MCA Case History No. 1893

Use of rusty tweezers, rather than the glass rod specified to handle specimens being immersed in hot 64% hydrazine, caused the hydrazine vapour to ignite.

Ruthenium(III) chloride

Allen, A. D. *et al.*, *Inorg. Synth.*, 1970, **12**, 3

During the preparation of pentaamminedinitrogenruthenium(II) solutions, the initial gas producing reaction is so vigorous that increase in scale above that described (9 mmol) is not advised.

See Metal catalysts, above

See Pentaamminechlororuthenium chloride: Sodium azide

Silver compounds

MRH Silver nitrate 3.10/78

Anon., *CISHC Chem. Safety Summ.*, 1978, **49**(194), 29

For analysis, a silver-containing solution was made alkaline with 25% sodium hydroxide solution and filtered, then the precipitate was washed with ammonium hydroxide to redissolve the silver. Hydrazine (as sulfate) was added to precipitate the silver, and when the mixture was heated, an explosion occurred. This could have been caused by precipitation of explosive silver nitride, rapid catalytic decomposition by silver compounds of the hydrazine salt, and/or ignition of the hydrogen evolved.

Sodium

See Sodium: Hydrazine

Sodium hydroxide

1. Anon., *Univ. Safety Assoc., Safety News*, 1978, (10), 12

2. Day, A. C. *et al.*, *Org. Synth.*, 1970, **50**, 42

While drying hydrazine over sodium hydroxide pellets, explosive decomposition occurred, spraying molten sodium hydroxide around. This was attributed to possible leakage of air into the evacuated flask.

Thiocarbonyl azide thiocyanate

See Thiocarbonyl azide thiocyanate: Ammonia, etc.

Titanium compounds


Bains, M. S., *Can. J. Chem.*, 1966, **44**, 534–538

Interaction of anhydrous hydrazine and titanium isopropoxide is explosive at 130°C in absence of solvent. Evaporation of solvent ether from the reaction product of tetrakis(dimethylamino)titanium and anhydrous hydrazine caused an explosion, attributed to formation and ignition of dimethylamine. *N*-Metal derivatives may also have been formed.

See other NON-METAL HYDRIDES, REDUCANTS

4521. Ammonium nitrite

[13446-48-5]




See Nitric Acid: Manufacture Hazard

See entry NITRITE SALTS OF NITROGENOUS BASES

4522. Ammonium nitrate

[6484-52-2]



FPA H33, 1975; HCS 1980, 147

1. Federoff, 1960, 35
2. Urbanski, 1965, Vol. 2, 460
3. Popper, H., *Chem. Eng.*, 1962, **70**, 91
4. Sykes, W. G. *et al.*, *Chem. Eng. Progr.*, 1963, **59**(1), 66
5. Yoshida, T., private comm., 1983
6. Urben, P. G., *Loss Prev. Bull.*, 1992, **108**, 31
7. Mellor, 1964, Vol. 8, Suppl. 1, 543
8. *MCA Guide*, 1972, 308
9. Croysdale, L. G. *et al.*, *Chem. Eng. Progr.*, 1965, **61**(1), 76
10. *MCA Case History No. 873*
11. Bailar, 1973, Vol. 2, 317
12. Vakhrushev, Yu. A. *et al.*, *Chem. Abs.*, 1975, **82**, 113745
13. Pascal, 1956, Vol. 10, 216
14. Rozman, B. Yu., *Chem. Abs.*, 1960, **54**, 12587c
15. Lee, P. R., *J. Appl. Chem.*, 1969, **19**, 345–351
16. Freeman, R., *Chem. Eng. Progr.*, 1975, **71**(11), 71–74
17. David, A. *et al.*, *Hydrocarbon Process.*, 1978, **57**(11), 149–175
18. Shah, K. D., *Fert. Sci. Tech. Ser.*, 1985, **4**, 261–273

19. Kirschner, E. *Chem. Eng. News*, 1994, **72**(51), 4; *ibid.*, **73**(37), 7
20. Anon., *Chem. Eng. News*, 1996, **74**(5), 11
21. Wiswesser, W. J., private comm., 1988
22. Anon., *Fire Prevention*, 1990, (231), 40
23. Li, R. *et al.*, *Arch. Combust.*, 1988, **8**(1), 65–70
24. Buczkowski, D. *et al.*, *Chem Abs.*, 1996, **125**, 176043e

The decomposition, fire and explosion hazards of this salt of positive oxygen balance have been adequately reviewed [1]. In the absence of impurities it is difficult, but not impossible, to cause ammonium nitrate to detonate. Use of explosives to break up the caked double salt, ammonium nitrate–ammonium sulfate (2:1), caused a 4.5 Mkg dump to detonate [4], even though some 45% of ammonium sulfate was present, effectively as a diluent, in the salt. Although the explosion tests available in 1921 did not give positive results for the nitrate–sulfate salt, it has recently been shown, using the BAM 50/60 steel tube test, that it will explode [5]. The ammonium sulphate is unlikely to be totally inert, having a negative enthalpy of decomposition to steam, nitrogen and sulfur, which last is capable of oxidation to sulfur dioxide, bringing the double salt into oxygen balance [6]. The effect of various impurities and additives on the thermal stability of ammonium nitrate has been widely studied [1,2,7]. Impure ammonium nitrate now constitutes more than 95% of all civil explosives used in the USA. A few incidents involving explosive decomposition of aqueous solutions of the salt during evaporation [8], or transfer operations (some in presence of oil) have been recorded [9]. Flame cutting a mild steel pipe blocked by the solid (impure) salt caused it to explode [10]. During preparation of dinitrogen oxide by the exothermic pyrolysis of the nitrate at 170°C, the temperature must not be allowed to exceed 250°C, as explosion may occur [11].

The thermal stability of ammonium nitrate solutions is decreased by presence of sodium 1,1'-methylenebis(6-naphthalenesulfonate) at 0.15%, but zinc oxide or zinc sulfate–apatite mixtures act as stabilisers [12]. Presence of ammonium chloride (0.1%) causes thermal decomposition to become violent around 175°C, and some chlorine is evolved [13]. Magnesium nitrate is claimed to significantly desensitise ammonium nitrate [24]. The parameters involved in the self-ignition of the nitrate during storage or transport are described, and a mechanism for instantaneous decomposition is proposed [14]. The factors affecting potentially explosive decomposition in bulk storage of the salt are discussed, and steady state thermal explosion theory is applied to the prediction of critical masses in relation to temperature in bulk storage. Application of the results to current storage practice is also discussed [15]. A detailed account of the investigation of a fire and explosion (basically unexplained) involving some 5 t of nitrate of a 14 kt warehouse stock has been published [16]. A 2-stage production process is designed to minimise danger of decomposition of the salt [17]. Accidents involving ammonium nitrate and the factors causing them are reviewed, and procedures for manufacturing, handling, storage and transportation are discussed [18]. A large explosion, probably of some tons TNT equivalent with several fatalities, during concentration of ammonium nitrate solution for fertiliser manufacture is discussed. There had been 750 tons of the nitrate present, the initiation mechanism is disputed though acidity,

H₄N₂O₃

impurities, heat and lack of supervision were involved [19,20]. Road trailers laden with 14 t and 7 t of nitrate were involved in a fire and 3 subsequent detonations [21]. Another account makes it clear that it was, for the most part, ammonium nitrate/fuel oil (ANFO) explosive rather than the pure nitrate which exploded [22]. During pneumatic pipeline conveyance of the powdered salt, electrostatic generation is unlikely to be a problem under smooth flow conditions. However, the possibility of fast charge accumulation if flow discontinuities arise must not be overlooked [23].

Acetic acid

MRH 3.30/16

von Schwartz, 1918, 322

Conc. mixtures ignite on warming.

Acetic anhydride, Hexamethylenetetramine acetate, Nitric acid

See Nitric acid: Acetic anhydride, Hexamethylenetetramine acetate

Aluminium, Calcium nitrate, Formamide

See Calcium nitrate: Aluminium, etc.

Ammonia

Kohczkowski, A. *et al.*, *J. Chem. Technol. Biotechnol.*, 1981, **31**, 327–332

Depending on the conditions, presence of free ammonia in ammonium nitrate may either stabilise, or tend to destabilise, the salt.

Ammonium chloride, (Barium nitrate), Water, Zinc

1. Bailey, P. S. *et al.*, *J. Chem. Educ.*, 1975, **52**, 525

2. Jackson, H., *Spectrum*, 1969, **7**(2), 82

3. Bodner, G. M., *J. Chem. Educ.*, 1985, **62**(12), 1106

Addition of water to an intimate mixture of zinc powder and the salts causes spontaneous ignition [1], and a mixture of ammonium nitrate and ammonium chloride (9:1) sprinkled with zinc dust ignites vigorously when moistened [2]. Premature spontaneous ignition of the mixture was attributed to absorption of moisture by the hygroscopic salt mixture [3].

See Metals, below

See other REDOX REACTIONS

Ammonium phosphate, Potassium sulfate

Gryzlov, L. D. *et al.*, *Chem. Abs.*, 1986, **104**, 185477

During manufacture of 26/16/10 N/P/K fertiliser by ammoniation of nitric–phosphoric acid mixtures, followed by concentration, addition of potassium sulfate, then granulation at up to 250°C, the possibility of a thermal explosion exists. Kinetic studies showed that 240°C appears to be a safe granulation temperature, but that changes in the composition and pH of the mixture may decrease this critical temperature.

Ammonium sulfate, Potassium

Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

Ammonium nitrate containing the sulfate readily explodes on contact with potassium or its alloy with sodium.

Alkali metals

MRH Sodium 4.15/36

Mellor, 1964, Vol. 8, Suppl. 1, 546

Sodium progressively reduces the nitrate, eventually forming a yellow explosive solid, probably sodium hyponitrite.

Calcium superphosphate, Lignin, Phosphoric acid, Urea

Dobrova, Ts. *et al.*, *Chem. Abs.*, 1988, **109**, 37213

The products of thermal decomposition of organic–mineral fertiliser mixtures were studied by IR spectroscopy, and implications for fire and explosion hazards considered.

See Urea, below

Charcoal, Metal oxides

MRH Carbon 3.64/7

Herbst, H., *Chem. Ztg.*, 1935, **59**, 744–745

The pelleted explosive ('ammonpulver', containing 10% charcoal) normally ignites at 160–165°C, but presence of rust, or copper oxide or zinc oxide lowers the temperature to 80–120°C.

See Non-metals, below

Chloride salts

1. Pascal, 1956, Vol. 10, 216

2. Pany, V. *et al.*, *Chem. Abs.*, 1976, **85**, 56018

The nitrate containing 0.1% of ammonium chloride decomposes vigorously below 175°C [1]. Presence of 0.1% of calcium chloride or iron(III) chloride in the nitrate lowers its initiation temperature sufficiently to give violent or explosive decomposition. Thermal analysis plots for aluminium chloride, calcium chloride and iron(III) chloride are given [2].

Copper iron(II) sulfide

Kuznetsov, G. V. *et al.*, *Chem. Abs.*, 1975, **82**, 75133

During preparations for blasting the sulfide mineral copper pyrites, ammonium nitrate-based blasting cartridges exploded prematurely in the blast holes. This was attributed to exothermic interaction of acid ground-water with the sulfide–oxidant combination.

See Metal sulfides, below

Cyanoguanidine

See Cyanoguanidine: Oxidants

Fertiliser materials

Davies, R. O. E. *et al.*, *Ind. Eng. Chem.*, 1945, **37**, 59–63

Mixtures of ammonium nitrate, superphosphate and organic materials stored in bulk may ignite if the internal temperature exceeds 90°C. This is owing to the free acid in the superphosphate, and may be prevented by neutralisation with ammonia.

Hydrocarbon oils

MRH Hexane 3.80/5, Toluene 3.80/7

Goffart, P., *Chem. Abs.*, 1975, **82**, 61420

Detonability of several ammonium nitrate-based fertilisers, with or without the addition of fuel oil, was studied.

Metals

1. Mellor, 1964, Vol. 8, Suppl. 1, 543–546
2. Soda, N. *et al.*, *Chem. Abs.*, 1969, **70**, 49151
3. Peterson, W. S. *Light Met.* (Warrendale, Pa) 1995, 899

MRH Aluminium 8.49/25, lead 1.38/56, magnesium 8.79/31, sodium 4.15/36, zinc 3.22/tr.

Many of the following powdered metals reacted violently or explosively with fused ammonium nitrate below 200°C: aluminium, antimony, bismuth, cadmium, chromium, cobalt, copper, iron, lead, magnesium, manganese, nickel, tin, zinc; also brass and stainless steel. Mixtures with aluminium powder are used as the commercial explosive Ammonal. Sodium reacts to form the yellow explosive compound sodium hyponitrite, and presence of potassium sensitises the nitrate to shock [1]. Shock-sensitivity of mixtures of ammonium nitrate and powdered metals decreases in the order titanium, tin, aluminium, magnesium, zinc, lead, iron, antimony, copper [2]. Contact between molten aluminium and the salt is violently explosive, apparently there is a considerable risk of this happening in scrap remelting [3].

See Ammonium sulfate, etc., above

Aluminium: Ammonium nitrate

Potassium: Nitrogen-containing explosives

Metal salts

1. Glazkova, A. P., *Chem. Abs.*, 1970, **73**, 132646
2. Rosser, W. A. *et al.*, *Trans. Faraday Soc.*, 1964, **60**, 1618–1625

Catalytic effects on the thermal decomposition and burning under nitrogen of the nitrate were determined for ammonium dichromate, potassium dichromate, potassium chromate, barium chloride, sodium chloride and potassium nitrate. Chromium(VI) salts are most effective in decomposition, and the halides salts during burning of the nitrate [1]. The effect of chromium compounds soluble in the molten nitrate, all of which promote decomposition of the latter, was studied (especially using ammonium dichromate) in kinetic experiments [2].

Metal sulfides

Proc. 13th Symp. Explos. Pyrotech., 1986, **VII**, 1–7; (*Chem. Abs.*, 1989 **110**, 98227s)

In presence of sulfide ores, specifically pyrites, explosives containing ammonium nitrate may undergo runaway reaction, leading to detonation at temperatures below 40°C if pH is less than 2. The reaction is acid catalysed.

Nitric acid

Rubtsov, Yu. I. *et al.*, *Chem. Abs.*, 1987, **106**, 158775

In the context of safety of the process of neutralisation of nitric acid with ammonia, the effects of temperature (160–230°C), pressure (2.3–9.8 bar), and concentrations of ammonium nitrate (86–94%) and of nitric acid (0–4%) upon decomposition rate were studied.

See Nitric acid: Manufacture Hazard

Non-metals MRH Carbon 3.64/7, phosphorus (y) 4.60/14
Mellor, 1940, Vol. 2, 841–842; 1946, Vol. 5, 830

The powdered salt in admixture with charcoal explodes at 170°C, or the solid salt on contact with glowing charcoal. Phosphorus ignites on the fused salt.

See Sulfur, below, *also* Phosphorus: Nitrates

Organic fuels MRH values below references

1. Sykes, W. G. *et al.*, *Chem. Eng. Progr.*, 1963, **59**(1), 66

2. Urbanski, 1965, Vol. 2, 461

3. Bernoff, R. A. *et al.*, US Pat. 3 232 940, 1969

MRH Acetone 3.77/8, aniline 3.77/8, carbon disulfide 3.26/11, ethanol 3.68/9, ethylene glycol 3.56/14

In a review of ammonium nitrate explosion hazards, several incidents involving mixtures of this positive oxygen balance material with various organic materials were analysed [1]. In general, fire incidents involving ammonium nitrate admixed with 1% of wax, oil, or stearates (as anti-caking additives) tended more towards explosion than the pure salt, although the degree of confinement is also important [1,2]. The ease of detonation of mixtures is much greater when 2–4% of oil is present, and such mixtures are used as commercial explosives [2]. The preparation of ammonium nitrate–fuel systems at a molecular level is covered by the patent, which describes several alkylammonium nitrates as high-energy materials [3].

Other reactants

Yoshida, 1980, 190

MRH values calculated for 19 combinations with oxidisable materials are given.

Potassium nitrite

Mellor, 1940, Vol. 2, 842

Contact of the solid nitrate with fused nitrite causes incandescence.

See NITRITE SALTS OF NITROGENOUS BASES

Potassium permanganate

See Potassium permanganate: Ammonium nitrate

Sawdust

Biasutti, 1981, 151

A violent explosion in an ammonium nitrate store with sawdust-covered floors (to absorb spillage and prevent sparks) was attributed to local decomposition of the moist nitrate-containing sawdust, leading to temperature rise and spontaneous ignition. The observation of red-brown fumes just before the explosion supports this hypothesis.

Sugar

Gonzalez, M. M., *Chem. Abs.*, 1981, **65**, 83230

Mixtures of the salt with sugar syrup are effective explosives.

Sulfide ores

Liao, M. *et al.*, *Proc. 13th Symp. Explos. Pyrotech.*, 1988, VII.1–VII.7

The main factors leading to spontaneous explosions of blasting explosives in sulfide ore mines are identified as presence of ammonium nitrate, presence of pyrites and their weathering products, as well as the pH and temperature. At pH below 2, the critical temperature for spontaneous explosion may be lower than 40°.

Sulfur

1. Mason, C. M. *et al.*, *J. Agric. Food Chem.*, 1967, **15**, 954
2. Prugh, R. W., *Chem. Eng. Progr.*, 1967, **63**(11), 53–55

The fire risks of nitrate–sulfur mixtures have been discussed [1], and explosion risks assessed by DTA [2].

Trinitroanisole

Biasutti, 1981, 54

A mixture of ammonium nitrate and 2,4,6-trinitroanisole, prepared as an explosive by mixing the hot components, ignited spontaneously and later exploded violently.

Urea

1. Croysdale, L. C. *et al.*, *Chem. Eng. Progr.*, 1965, **61**(1), 72
2. Anderson, J. F., *Safety Air Ammon. Plants*, 1967, **9**, 70–71
3. Leach, J. W., *Rept. AD-A075338*, Richmond (Va.), USNTIS, 1979
4. Perbal, G., *Proc. 4th Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, Vol. 2, M1–M19, Rugby, IChE, 1983
5. Rubstov, Yu. I. *et al.*, *Chem. Abs.*, 1988, **108**, 137111

Conc. solutions of ammonium nitrate and urea exploded during large-scale mixing operations. Although the cause was not established, the hazards of these operations were discussed in relation to the circumstances [1]. A further explosion under similar circumstances was reported [2]. Some of the liquid streams involved in preparation of nitroguanidine, which are somewhat similar to those in the present process, are detonable if less than 25% of water is present in the solutions. Thermal characteristics and detonation data were developed for the nitroguanidine process and that to produce guanidinium nitrate [3]. The circumstances of the 3 explosions [1,2] have been reviewed and detailed experimental work has confirmed that the solutions (54–46% ammonium nitrate, 35–39% urea and 15–20% water) cannot be detonated. However, dehydration of the solution gives a solid residue capable of deflagration at 240°C, and the residue from prolonged heating at 158°C is capable of detonation. It is concluded that the explosions were most likely caused by leakage and dehydration in the lagging of the steam-traced pipelines at 158°C, slow development of the unstable product and of a hot spot, with eventual initiation of detonation. Any traces of urea nitrate would not have contributed significantly to the events [4]. There is no danger of violent reaction as long as the components remain in solution. Processing temperatures should not exceed 120°C [5].

Water

Kolaczkowski, A. *et al.*, *Chem. Abs.*, 1982, **97**, 78074

Hot aqueous solutions of the nitrate of above 50% concentration may decompose explosively under adiabatic conditions and under conditions of confinement (small, or no vents). A 9 m³ tank of 85% solution also containing 0.9% of nitric acid

and 0.2% of chloride ion (which catalyses decomposition) at 110°C exploded powerfully.

See related METAL NITRATES

See other OXOSALTS OF NITROGENOUS BASES

4523. Ammonium nitridoosmate

[22493-01-2]

$\text{H}_4\text{N}_2\text{O}_3\text{Os}$



Mellor, 1942, Vol. 15, 727


It decomposes explosively at 150°C, like the barium salt.

See related NITRIDES

4524. Hydroxylaminium nitrate

[13465-08-2]

$\text{H}_4\text{N}_2\text{O}_4$



1. Mellor, 1940, Vol. 8, 303

2. Kirk-Othmer, 1981, **16**, 179

3. Asaki, A., *Chem. Abs.*, 1983, **98**, 48238

4. Van Disk, C. A., *Chem. Abs.*, 1984, **100**, 70566

The redox salt which decomposes at 100°C, evolving nitrogen and oxygen [1] is overall an oxidant but has been used as a reductant in actinide processing [2]. It has been used recently as a missile monopropellant [3], and also as an oxidant, with its isopropyl derivative as the fuel [4].

See other HYDROXYLAMINIUM SALTS

Amine nitrates

Klein, N., *Progr. Astronaut. Aeronaut.*, 1988, **109** (Gun Propell. Technol.), 473–497


The use of aqueous mixtures of hydroxylammonium nitrate with various butylammonium nitrates as gun propellants is reviewed.

See other OXOSALTS OF NITROGENOUS BASES, REDOX COMPOUNDS

4525. Tetrahydroxotritin(2+) nitrate ('Basic stannous nitrate')

[73593-33-6] (ion)

$\text{H}_4\text{N}_2\text{O}_{10}\text{Sn}_3$



1. Anon., *J. Soc. Chem. Ind.*, 1922, **41**, 423–434R

2. Donaldson, J. D. *et al.*, *J. Chem. Soc.*, 1961, 1996

An explosion in flour-bleaching operations was attributed to violent decomposition of the basic nitrate [1], which is an impact-, friction- and heat-sensitive explosive [2]. The instability is associated with the presence of reductant and oxidant functions in the same molecule. The previous formulation as tin(II) nitrate oxide is revised to that above.

See other REDOX COMPOUNDS

See related METAL NITRATES

4526. Ammonium azide
[12164-94-2]




1. Mellor, 1940, Vol. 8, 344
2. Sorbe, 1968, 129
3. Obenland, C. O. *et al.*, *Inorg. Synth.*, 1966, **8**, 53

It explodes on rapid heating [1] and contains ~93% of nitrogen. It is also friction- and impact-sensitive [2], and preparative precautions have been detailed [3].

See other HIGH-NITROGEN COMPOUNDS, NON-METAL AZIDES

4527. Tetraimide (Tetraazetidine)
[58674-00-3]




Rice, F. O. *et al.*, *J. Amer. Chem. Soc.*, 1957, **79**, 1880–1881

This blue solid (a cyclic dimer of diazene, 93.3% of nitrogen), obtained by freezing out at -195°C the pyrolysis products of hydrogen azide, is extremely explosive above this temperature. An explosion at -125°C destroyed the apparatus.

See other HIGH-NITROGEN COMPOUNDS

4528. Ammonium dinitramide
[140456-78-6]



See Dinitramine

See other N-NITRO COMPOUNDS

4529. 'Sodium perpyrophosphate'
[18178-05-7]



See Sodium pyrophosphate hydrogen peroxidate

4530. Oxodisilane
[22755-00-6]



Kautsky, K., *Z. Anorg. Chem.*, 1921, **117**, 209

It ignites in air.

See related SILANES

- 4531. Lead(II) phosphinate**
 [10294-58-3] **H₄O₄P₂Pb**

$$\text{Pb(OP(O)H}_2)_2$$
 Mellor, 1971, Vol. 8, Suppl. 3, 623
 Impact-sensitive, used as priming explosive.
See other HEAVY METAL DERIVATIVES, METAL PHOSPHINATES, REDUCANTS
- 4532. Ruthenium(IV) hydroxide**
 [12181-34-9] **H₄O₄Ru**

$$\text{Ru(OH)}_4$$
 Mellor, 1942, Vol. 15, 516
 It deflagrates incandescently above 300°C.
- 4533. Xenon(IV) hydroxide**
 [] **H₄O₄Xe**

$$\text{Xe(OH)}_4$$
 Sorbe, 1968, 158
 Very unstable, it explodes violently on heating.
See other NON-METAL OXIDES, XENON COMPOUNDS
- 4534. Zirconium hydroxide**
 [14475-63-9] **H₄O₄Zr**

$$\text{Zr(OH)}_4$$
 Zirconium
See Zirconium: Oxygen-containing compounds
- 4535. Lead(II) phosphite**
 [15521-60-5] **H₄O₆P₂Pb**

$$\text{Pb(OPH(O)OH)}_2$$
 491M, 1975, 228
 A fibre drum of the salt ignited spontaneously. This was attributed to slow decomposition with formation of phosphine which accumulated and eventually ignited inside the drum.
See other HEAVY METAL DERIVATIVES, REDUCANTS
- 4536. Tetrahydroxydioxotrisilane**
 [] **H₄O₆Si₃**

$$\text{HOSi(O)Si(OH)}_2\text{Si(O)OH}$$
 1. Mellor, 1940, Vol. 6, 230
 2. Müller, R. *et al.*, *J. Prakt. Chem.*, 1966, **31**, 1–6
- 1686

The hydrolysis product of octachlorotrisilane (and probably polymeric), it decomposes violently or explosively when heated in air, or sometimes spontaneously [1], and ignites when subjected to friction [2].


See Poly(dihydroxydioxodisilane)

See related NON-METAL OXIDES

4537. Oxobis[aqua(oxo)diperoxorheniumVIII]

[174753-06-1]

H₄O₁₃Re₂



Herrmann, W. A. *et al.*, *Chem. Eur. J.*, 1996, **2**(2), 168

An explosive red/orange solid, obtained from dirhenium heptoxide and hydrogen peroxide reacting in ether as solvent.

See other INORGANIC PEROXIDES

†4538. Diphosphane

[13445-50-6]

H₄P₂

H₂PPH₂

Mellor, 1940, Vol. 8, 829

It ignites in air, and will cause other flammable gases to ignite when present at 0.2 vol% concentration.

See other PHOSPHINES

†4539. Silane

[7803-62-5]

H₄Si

SiH₄

1. Mellor, 1940, Vol. 6, 220–221

2. Braker, 1980, 632

3. Gale, H. J., *Lab. Haz. Bull.*, 1984, (3), item 140

4. Britton, L. G., *Plant/Oper. Progr.*, 1990, **9**(1), 16

5. Katou, Y., *Chem. Abs.* 1990, **113**, 84061h

6. Kondo, S. *et al.*, *Combust. Flame*, 1994, **97**(3–4), 296.

7. Horiguchi, S., *Chem. Abs.* 1997, **127**, 143180k

It is mildly endothermic (ΔH_f° (g) +32.5 kJ/mol, 1.01 kJ/g).

Pure material is said not to ignite in air unless the temperature be increased or the pressure reduced. Presence of other hydrides as impurities causes ignition always to occur on contact [1]. However, 99.95% pure material, even at concentrations down to 1% in hydrogen and/or nitrogen, ignites in contact with air unless emerging at

very high gas velocity, when mixtures with up to 10% silane content may not ignite [2]. Silane from a cylinder at 24 bar contained inside a ventilated cabinet inadvertently flowed back to the low pressure side of a nitrogen cylinder regulator outside the cabinet. When the regulator failed, the leaking silane ignited and burned outside the cabinet, preventing access to the main silane valve, and the leak reignited after extinguishing the fire. The system has been redesigned to eliminate the problems [3]. An exhaustive review of combustion and associated hazards is found in [4]. This also reveals potentially dangerous deflagration with the chlorofluoromethanes used in some fire extinguishers. A detailed study of silane flammability and blast effects in mixtures with nitrogen discharged to air [5]. The lower flammable limit of silane in air declines with decreasing oxygen content [6]. A Japanese review of explosion hazards with silane has been published. Spontaneous combustion in (somewhat oxygen depleted) air has been observed down to -162 C. In addition to the materials listed separately below, trifluorammine, interhalogens and a range of fluorocarbons and halofluorocarbons form explosive mixtures, as do sulfur tetra- and hexa-fluorides. Most of these are described as pyrophoric with silane, which may be a mistranslation [7].

See other IGNITION SOURCES

Air, *cis*-2-Butene

Urtiew, P. A. *et al.*, *Rept. UCRL-52007*, Richmond (Va.), USNTIS, 1976

Presence of the alkene delays or prevents spontaneous ignition of silane–air mixtures.

Halogens, or Covalent chlorides

MRH values below references

1. Mellor, 1940, Vol. 6, 220–221

2. Braker, 1980, 632

MRH Bromine 1.21/91, carbon tetrachloride 2.89/83, chlorine 3.85/82

Silane burns in contact with bromine, chlorine or covalent chlorides (carbonyl chloride, antimony pentachloride, tin(IV) chloride, etc.) [1]. Extreme caution is necessary when handling silane in systems with halogenated compounds, as a trace of free halogen may cause violent explosions [2].

Nitrogen oxides

Horiguchi, S. *et al.*, *Chem. Abs.*, 1990, **112** 121711

Mixtures of silane and nitrogen oxides have been studied, that with nitrous oxide detonates very easily.

Other reactants

Yoshida, 1980, 371

MRH values calculated for 4 combinations with the materials indicated are given.

Oxygen

MRH 14.31/66

1. Bailar, 1973, Vol. 1, 1366

2. Ring, M. A. *et al.*, *AIP Conf. Proc.*, 1988, **166** (Photovolt. Safety), 175–182

Very pure silane does not immediately explode with oxygen, but the decomposition products may ignite after a delay [1]. Mixtures of silane with 30% oxygen are metastable and potentially explosive under all pressure conditions studied, and

become explosive at above 80°C. Addition of a few % of disilane renders the mixtures much more explosive, and mechanisms are proposed [2].

See other SILANES

4540. Thorium hydride

[15457-87-1]

H₄Th

ThH₄

1. Mellor, 1941, Vol. 7, 207
2. Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64
3. Hartmann, I. *et al.*, *Rept. Invest. No. 4835*, Washington, US Bur. Mines, 1951
Thorium hydride explodes on heating in air [1], and the powdered hydride readily ignites on handling in air [2]. Layers of thorium or uranium hydrides ignited spontaneously after exposure to ambient air for a few min [3].

See other METAL HYDRIDES

4541. Uranium(IV) hydride

[51680-55-8]

H₄U

UH₄

1. Hartmann, I. *et al.*, *Rept. Invest. No. 4835*, Washington, US Bur. Mines, 1951
2. Starks, D. F. *et al.*, *J. Amer. Chem. Soc.*, 1973, **95**, 3423
Layers of uranium or thorium hydrides ignited spontaneously after exposure to ambient air for a few min [1]. Thermal decomposition yields pyrophoric uranium [2].

Oxygen

See Oxygen (Gas): Metal hydrides

See other METAL HYDRIDES

4542. Orthoperiodic acid

[10450-60-9]

H₅IO₆

O:I(OH)₅

Azo-pigment, Perchloric acid

See Perchloric acid: Azo-pigment, etc.

See other INORGANIC ACIDS, OXOHALOGEN ACIDS

4543. Iododisilane

[14380-76-8]

H₅ISi₂

ISiH₂SiH₃

Bailar, 1973, Vol. 1, 1371

It ignites in air.

See other HALOSILANES, IODINE COMPOUNDS

4544. Ammonium hydroxide

[1336-21-6]

 H_5NO

(MCA SD-13, 1947); NSC 701, 1983; HCS 1980, 140

Nitromethane

See Nitromethane: Acids, etc.

Other reactants

Yoshida, 1980, 211

MRH values calculated for 11 combinations with oxidants are given.

See other INORGANIC BASES See related NON-METAL HYDRIDES

4545. Ammonium hydrogen sulfite

[10192-30-0]

 $\text{H}_5\text{NO}_3\text{S}$

Self accelerating decomposition ($Q = 0.43$ kJ/g) occurs at 130°C in closed systems, such as are used for the Bucherer reaction.

See entry SELF-ACCELERATING REACTIONS

4546. Hydrazinium salts

[]

 $\text{H}_5\text{N}_2^+ \text{Z}^-$


Metal nitrites

See Hydrogen azide

See also NITRITE SALTS OF NITROGENOUS BASES

4547. Diamidophosphorous acid

[22750-67-0]

 $\text{H}_5\text{N}_2\text{OP}$

Water


Mellor, 1940, Vol. 8, 704

Interaction is violent with incandescence.

See other INORGANIC ACIDS


4548. Hydrazinium nitrite

[]

 $\text{H}_5\text{N}_3\text{O}_2$

See entries HYDRAZINIUM SALTS, NITRITE SALTS OF NITROGENOUS BASES

4549. Hydrazinium nitrate
[13464-97-6]



Alone, or Metals, or Metal compounds

Mellor, 1940, Vol. 8, 327; 1967, Vol. 8, Suppl. 2.2, 84, 96

It is an explosive of positive oxygen balance, less stable than ammonium nitrate, and has been studied in detail. Stable on slow heating to 300°C, it decomposes explosively on rapid heating or under confinement. Presence of zinc, copper, most other metals and their acetylides, nitrides, oxides or sulfides cause flaming decomposition above the m.p. (70°C). Commercial cobalt (cubes) causes an explosion also.

See entry HYDRAZINIUM SALTS

2-Hydroxyethylamine

Fujihara, S. *et al.*, *Chem. Abs.*, 1976, **84**, 7212

Mixtures of the salt (80%) and amine (15%) with water are useful as an impact-insensitive but powerful liquid explosive.

Potassium dichromate

Shidlovskii, A. A. *et al.*, *Chem. Abs.*, 1960, **54**, 22132g

Thermal decomposition becomes explosive above 270°C, or above 100°C in presence of 5% of potassium dichromate.

See other OXOSALTS OF NITROGENOUS BASES, REDOX COMPOUNDS


4550. Hydrazinium azide
[38551-29-0]



It contains some 94% of nitrogen.

See other HIGH-NITROGEN COMPOUNDS, HYDRAZINIUM SALTS, NON-METAL AZIDES

4551. Potassium triamidothallate ammoniate
[]



Alone, or Acids, or Water

Mellor, 1940, Vol. 8, 262

Like thallium(I) amide from which it is derived by treatment with potassium amide in liquid ammonia, the ammoniated salt ($x = 2$ or less) explodes violently on heating, friction, or contact with dilute acids or water.

See other N-METAL DERIVATIVES

4552. Lanthanum pentanickel hexahydride


[54847-21-1]

H₆LaNi₅Imamoto, T. *et al.*, *J. Org. Chem.*, 1987, **52**, 5695–5699

The alloy hydride has been investigated as a useful hydrogenation catalyst for a wide variety of substrates under mild conditions. It is however pyrophoric in air, and an experimental procedure has been developed to avoid this hazard. A related hydride, LaNi_{4.5}Al_{0.5}H₅ has similar properties.

See Lanthanum–nickel alloy*See* LANTHANIDE–TRANSITION METAL ALLOY HYDRIDES**4553. Dioxonium hexamanganato(VII)manganate**

[51184-04-4]

H₆Mn₁₄O₅₂·11H₂O[54065-28-0] (new unsolv. structure H₂Mn₇O₂₄)Krebs, B. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1974, **13**, 603

It has been shown that 'solid permanganic acid dihydrate', the powerful and unstable oxidant, is an undecahydrate as formulated above.

See Permanganic acid*See other* INORGANIC ACIDS, OXIDANTS**4554. Ammonium phosphinate**

[7803-65-8]


H₆NO₂P

Mellor, 1940, Vol. 8, 880

The salt evolves spontaneously flammable phosphine around 240°C.

See other REDUCANTS *See related* METAL PHOSPHINATES**4555. Hydroxylaminium phosphinate**

[]

H₆NO₃P

Mellor, 1940, Vol. 8, 880

The salt detonates above its m.p., 92°C.

See other HYDROXYLAMINIUM SALTS, REDUCANTS *See related* METAL PHOSPHINATES

4556. Ammonium amidosulfate (Ammonium sulfamate)

[7773-06-0]

H₆N₂O₃S



1. Hunt, J. K., *Chem. Eng. News*, 1952, **30**, 707
2. Rogers, M. G., private comm., 1973

A 60% solution of ammonium sulfamate (pH above 4.5) will not undergo rapid hydrolysis below 200°C. Addition of acid (to pH 2) causes a runaway exothermic hydrolysis to set in at 130°C. Superheating and vigorous boiling can occur under appropriate physical conditions [1]. The use of urea-formaldehyde resins as temporary binders in the firing of refractories and ceramics at high temperatures can lead to the formation of substantial deposits of ammonium sulfamate in the cooler parts of kilns, should a fuel oil containing appreciable amounts of sulfur be used for firing. Ammonia from decomposition of the resin combines with sulfur trioxide to form ammonium sulfamate which accumulates as either a solidified deposit in flues or as a white deposit on walls, causing corrosion and handling problems [2]. See other CORROSION INCIDENTS, N-S COMPOUNDS, OXOSALTS OF NITROGENOUS BASES

4557. Ammonium amidoselenate

[13767-10-0]

H₆N₂O₃Se



Dostal, K. *et al.*, *Z. Anorg. Chem.*, 1958, **296**, 29–35


It explodes if rapidly heated to 120°C.

See related N-S COMPOUNDS See other OXOSALTS OF NITROGENOUS BASES

4558. Hydrazinium hydrogen selenate

[]

H₆N₂O₄Se



Meyer, J. *et al.*, *Ber.*, 1928, **61**, 1839

The salt explodes in contact with a hot glass rod.

See other HYDRAZINIUM SALTS, OXOSALTS OF NITROGENOUS BASES

4559. Diamminepalladium(II) nitrite

[28068-05-5]

H₆N₄O₄Pd



Fountain, N. O., private comm., 1983

The dry salt is not readily ignitable, but is marginally more sensitive to impact than dinitrobenzene and can be ignited by impacted friction. The deflagration rate, once confined ignition had begun, was just greater than that of typical deflagrating explosives.

See other AMMINOMETAL OXOSALTS See related PLATINUM COMPOUNDS

4560. cis-Diammineplatinum(II) nitrite

[14286-02-3]

H₆N₄O₄Pt



1. Brauer, 1965, Vol. 2, 1560
2. Holifield, P. J., private comm., 1974
3. Fountain, N. O., private comm., 1983


It decomposes explosively at 200°C [1]. Dry material stored in clear bottles in sunlight for several weeks became sensitive and exploded violently on slight mechanical shock. The material is now supplied commercially moistened with water [2]. The dry salt may be readily ignited, and deflagration under confinement is similar to that of gunpowder. Presence of 10% of water reduced ignitability but did not slow the deflagration rate thereafter. Sensitivity to impact or friction was greater than that of nitrobenzene [3].

See other AMMINEMETAL OXOSALTS, PLATINUM COMPOUNDS

4561. Hydrazinium dinitrate

[13464-98-7]

H₆N₄O₆



Sorbe, 1968, 130

Like the mono-salt, it explodes on heating or impact.

See other HYDRAZINIUM SALTS, OXOSALTS OF NITROGENOUS BASES

4562. Diamminepalladium(II) nitrate

[34090-17-0]

H₆N₄O₆Pd



White, J. H., private comm., 1965

There is a danger of explosion if the nitrate is dried.

See other AMMINEMETAL OXOSALTS *See related* PLATINUM COMPOUNDS

4563. Zinc dihydrazide

[25546-98-9]

H₆N₄Zn



Mellor, 1940, Vol. 8, 315

It explodes at 70°C.

See other METAL HYDRAZIDES

4564. Sodium hexahydroxyplatinate(IV)

[12325-31-4]



Acetic acid, Nitric acid

See Nitric acid: Acetic acid, Sodium hexahydroxyplatinate(IV)*See other* PLATINUM COMPOUNDS**4565. Disilyl oxide**



[13597-73-4]



Chlorine

See Chlorine: Disilyl oxide*See related* SILANES**4566. 2,4,6-Trisilatrioxane (Cyclotrisiloxane)**

[291-50-9]




Air, or Chlorine

Mellor, 1940, Vol. 6, 234

The solid polymer (approximating to the trimer) ignites in air or chlorine.

See related SILANES**4567. Tetraphosphoric acid**

[13813-62-2]



Water

Mellor, 1971, Vol. 8, Suppl. 3, 736

Dilution of polyphosphoric acids with water in absence of cooling may lead to a large exotherm. Thus, tetraphosphoric acid diluted from 84 to 54% phosphorus pentoxide content rapidly attains a temperature of 120–140°C.

See other INORGANIC ACIDS**4568. Disilyl sulfide**

[16544-95-9]



Sorbe, 1968, 127

It ignites in moist air.

See related NON-METAL SULFIDES, SILANES

†4569. Disilane
[1590-87-0]



Bromine

See Bromine: Non-metal hydrides

Non-metal halides, or Oxygen

Mellor, 1940, Vol. 6, 220–224

It explodes on contact with carbon tetrachloride or sulfur hexafluoride, and contact with chloroform causes incandescence. Disilane ignites spontaneously in air, even when pure, and ingress of air or oxygen into a volume of disilane causes explosion.

See other SILANES

4570. Dihydrazinemanganese(II) nitrate

[39957-12-5]



Bailar, 1973, Vol. 3, 827

It ignites at 150°C, but is not shock-sensitive.

See other AMMINOMETAL NITRATES

4571. Amminepentahydroxyplatinum

[]



Jacobsen, J., *Compt. rend.*, 1909, **149**, 575

It explodes fairly violently above 250°C, as does the pyridine analogue.

See other N-METAL DERIVATIVES, PLATINUM COMPOUNDS

4572. Diamminedihydroxyosmium

[]




Ephraim, 1939, 463

It decomposes vigorously on heating.

See other N-METAL DERIVATIVES *See related* PLATINUM COMPOUNDS

4573. Ammonium thiosulfate

[7783-18-8]




Sodium chlorate

See Sodium chlorate: Ammonium salts, etc.

See related N-S COMPOUNDS

4574. Ammonium sulfate
[7783-20-2]



Other reactants

Yoshida, 1980, 394

MRH values calculated for 14 combinations with various materials are given.

Ammonium nitrate

See Ammonium Nitrate

Potassium chlorate

See Potassium chlorate: Ammonia, etc.

Sodium hypochlorite

See Sodium hypochlorite: Ammonium salts

See related METAL OXONON-METALLATES

4575. Hydroxylaminium sulfate
[10039-54-0]



FPA H66, 1978; *HCS 1980*, 559

1. Grewer, T. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 1–9, Oxford, Pergamon, 1987

2. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn, 1986

The solid deflagrates at 5.6 cm/min, and will attain a maximum of 250 bar pressure in a closed system [1]. Energy of decomposition (in range 180–280°C) measured as 1.6 kJ/g [2].


See other DEFLAGRATION INCIDENTS

Aniline, Chloral hydrate

See 2,2,2-Trichloro-1,1-ethanediol

See entry HYDROXYLAMINIUM SALTS

4576. Ammonium peroxodisulfate (Ammonium persulphate)
[7727-54-0]



HCS 1980, 149

Energy of decomposition (in range 140–210°C) measured as 0.323 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Aluminium, Water MRH Aluminium 5.90/24
Pieters, 1957, 30
A mixture including the powdered metal may explode.

Ammonia, Silver salts
Mellor, 1947, Vol. 10, 466
In conc. solutions the silver catalysed oxidation of ammonia to nitrogen may be very violent.

Azoisobutyronitrile

1. Anon., *Chemical Engineer*, 1993, **536**, 4

2. *The Fire at Allied Colloids Ltd.*, HSE Books, Sudbury UK, 1994

The azonitrile (AZDN) and ammonium persulphate were stored together as polymerisation initiators. The AZDN was warmed by a supposedly blanked-off steam pipe and partially decomposed, rupturing its kegs and falling onto the persulphate in sacks below. Some degree of mixing was effected by a falling keg lid piercing a sack and carrying the AZDN with it. Subsequent ignition of the explosive mix (exactly how is uncertain) caused explosion and a major fire. Though they serve the same purpose, oxidants should not be stored with reductants.

See Azoisobutyronitrile

See RADICAL INITIATORS

Iron MRH 0.92/22
Mellor, 1947, Vol. 10, 470
Iron exposed to the action of a slightly acid conc. solution of ammonium peroxodisulfate dissolves violently.

Other reactants

Yoshida, 1980, 328

MRH values calculated for 11 combinations, largely with oxidisable materials, are given.

Sodium peroxide MRH 2.72/40
See Sodium peroxide: Ammonium peroxodisulfate

Sodium sulphide

Zhang, G-S. *et al.*, *Loss Prevention and Safety Promotion in the Process Industries*, Vol I, (Mewis, J. J., Pasman, H. J. & De Rademaker, E. E. Eds.), 277, Amsterdam, Elsevier, 1995

An account of a serious warehouse explosion (15 dead, 141 injured). The two principal detonations were mostly due to ammonium nitrate, of which some hundred tonnes had been present, but the initiating fire was first observed in ammonium persulfate. This had been promiscuously stored alongside potassium permanganate, matches, potassium nitrate and sodium sulphide (or possibly sulphite), inter alia. None of these would improve the safety of ammonium persulfate. It was shown that the persulphate gives an immediate exothermic reaction with the sulphide. This was ascribed as the ultimate initiation. It was concluded that oxidants and

reducants should not be mixed in storage [many other oxidants, such as permanganates, should not be trusted near a peroxy compound—Ed.] and that storage areas should be better separated one from another.

See also POTASSIUM PERMANGANATE: AMMONIUM NITRATE

Sulfuric acid

See CLEANING BATHS FOR GLASSWARE

Zinc, Ammonia

Rept. HM Insp. Explos., 33, London, HMSO, 1950

The salt exploded during drying, but no cause was determined. However, the reputed explosive character of tetraamminezinc peroxodisulfate, possibly formed from interaction of the ammonium salt, galvanised iron and ammonia, was mentioned as a possible cause.

See Tetraamminezinc peroxodisulfate

See other OXIDANTS, PEROXOACID SALTS

4577. Ammonium sulfide

[12135-76-1] : polysulfide is [12259-52-6]
(NH₄)₂S

H₈N₂S

Zinc

Anon., *Sichere Chemiarb.*, 1960, **12**(4), 29

A closed zinc container filled with a conc. solution of ammonium sulfide exploded, owing to liberation of hydrogen sulfide and hydrogen, accompanying the formation of zinc ammonium sulfide.

See related METAL SULFIDES

4578. Ammonium tetranitroplatinate(II)

[22289-82-3] (ion)
[NH₄]₂[Pt(NO₂)₄]

H₈N₆O₈Pt

Nilson, L. F., *J. Prakt. Chem.* [2], 1877, **16**, 249

It decomposes explosively on heating.

See other OXOSALTS OF NITROGENOUS BASES, PLATINUM COMPOUNDS

†4579. Trisilane

[7783-26-8]
H₃SiSiH₂SiH₃

H₈Si₃

Air, or Carbon tetrachloride, or Oxygen

Mellor, 1940, Vol. 6, 224

It ignites and explodes in air or oxygen, and reacts vigorously with carbon tetrachloride.

See other SILANES

4580. Trisilylamine

[13862-16-3]

 H_9NSi_3

Air, or Ammonia, or Hydrogen

Mellor, 1940, Vol. 8, 262

The liquid ignites in air, and reacts vigorously with ammonia, hydrogen or moisture.

See related SILANES**4581. Sodium triammine**

[84937-00-8]

 $\text{H}_9\text{N}_3\text{Na}$ Winter, R. *et al.*, *Ber. Bunsenges. Phys. Chem.*, 1982, **86**, 1093–1097

The new crystalline solid, metastable in liquid nitrogen, dissociates at ambient temperature and caused several sample tubes to explode from internal pressure of ammonia.

See other GAS EVOLUTION INCIDENTS*See related* N-METAL DERIVATIVES**4582. Triamminenitratoplatinum(II) nitrate**

[17524-18-4]

 $\text{H}_9\text{N}_5\text{O}_6\text{Pt}$

Mellor, 1942, Vol. 16, 409

It decomposes violently on heating.

See other AMMINOMETAL OXOSALTS, PLATINUM COMPOUNDS**4583. Trisilylphosphine**

[15110-33-5]

 H_9PSi_3 Amberger, E. *et al.*, *Angew. Chem.*, 1962, **74**, 32–33

The liquid ignites in air.

See related PHOSPHINES, SILANES**4584. Tetrasilane**

[7783-29-1]

 H_{10}Si_4


Air, or Carbon tetrachloride, or Oxygen

Mellor, 1940, Vol. 6, 224

It ignites and explodes in air or oxygen, and reacts vigorously with carbon tetrachloride.

See other SILANES

4585. Tetrasilylhydrazine
[25573-59-5]




Bailar, 1973, Vol. 1, 1377

It explodes in air.

See SILYLHYDRAZINES See related SILANES

4586. Tetraamminezinc peroxodisulfate
[39733-13-6]



Barbieri, G. A. *et al.*, *Z. Anorg. Chem.*, 1911, **71**, 347

It explodes on heating or impact.

See other AMMINOMETAL OXOSALTS

4587. Tetraamminenickel(II) nitrate
[15651-35-1]



Explosive.

See other AMMINOMETAL NITRATES

4588. Tetraamminepalladium(II) nitrate
[13601-08-6]



White, J. H., private comm., 1965

Evaporation of a solution of the salt used for plating gave a moist residue which ignited and burned violently. Reclamation of palladium from such solutions by direct reduction is recommended. It was subsequently shown to be explosive

See other AMMINOMETAL NITRATES See related PLATINUM COMPOUNDS

4589. Tetraammineplatinum(II) nitrate (Tetraamminedinitratoplatinum)
[20634-12-2] tetrahedral



Similar to the palladium complex (above)


See other AMMINOMETAL NITRATES, PLATINUM COMPOUNDS

4590. Tetraammineplatinum(II) nitrate (Tetraamminedinitratoplatinum)
 [133163-34-5] octahedral **H₁₂N₆O₆Pt**

$$\text{O}_2\text{NOPt}(\text{NH}_3)_4\text{ONO}_2$$

Similar to the palladium complex (above)
See other AMMINEMETAL NITRATES, PLATINUM COMPOUNDS

4591. Ammonium 2,4,6-tris(dioxoselena)hexahydrotriazine-1,3,5-triide
 ('Ammonium triselenimidate') **H₁₂N₆O₆Se₃**
 []



Explosive.
See Selenium difluoride dioxide: Ammonia
See related N-S COMPOUNDS

4592. Trihydrazinenickel(II) nitrate
 [69101-54-8] **H₁₂N₈NiO₆**

$$[(\text{H}_4\text{N}_2)_3\text{Ni}][\text{NO}_3]_2$$

Ellern, H. *et al.*, *J. Chem. Educ.*, 1955, **32**, 24
 A dry sample exploded violently and a moist sample spontaneously deflagrated.
See other AMMINEMETAL NITRATES

4593. Amminedecahydroxydiplatinum
 [] **H₁₃NO₁₀Pt**

$$[\text{H}_3\text{NPt}_2(\text{OH})_{10}]$$


Jacobsen, J., *Compt. rend.*, 1909, **149**, 574–577
 The compound, of unknown structure, explodes violently on heating.
See other PLATINUM COMPOUNDS

4594. Tetraamminehydroxynitratoplatinum(IV) nitrate
 [] **H₁₃N₇O₁₀Pt**

$$[(\text{H}_3\text{N})_4\text{Pt}(\text{OH})\text{NO}_3][\text{NO}_3]_2$$

Mellor, 1942, Vol. 16, 411
 It explodes violently on heating.
See other AMMINEMETAL OXOSALTS, PLATINUM COMPOUNDS

4595. Tetraamminelithium dihydrogenphosphide
[44023-01-6] **H₁₄LiN₄P**



Water

Legoux, C., *Compt. rend.*, 1938, **207**, 634

It reacts vigorously with water, evolving phosphine and ammonia which may ignite.

See related METAL NON-METALLIDES

4596. Pentaamminedinitrogenruthenium(II) salts
[19504-40-6] (ion) **H₁₅N₇Ru²⁺**



Preparative hazard

See Pentaamminechlororuthenium chloride: Sodium azide

Hydrazine: Ruthenium(III) chloride

See other N-METAL DERIVATIVES

4597. Octaammine- μ -hydroxy[μ -superoxo-*O,O'*]dirhodium(4+) nitrate
[113753-12-1] **H₂₅N₁₂O₁₅Rh₂**



Springborg, J. *et al.*, *Acta Chem Scand. Sec. A*, 1987, **A41**, 485

The complex may detonate when subjected to mechanical shock.

See other AMMINEMETAL OXOSALTS

4598. Undecaamminetetraruthenium dodecaoxide
[] **H₃₃N₁₁O₁₂Ru₄**



Preparative hazard

See Ruthenium(VIII) oxide: Ammonia

See related METAL OXIDES

4599. Hafnium
[7440-58-6] **Hf**

Hf

Alone, or Non-metals, or Oxidants

(MCA SD-52, 1966)

Although the massive metal is relatively inert, when powdered it becomes very reactive. The dry powder may react explosively at elevated temperatures with nitrogen, phosphorus, oxygen, sulfur and other non-metals. The halogens react similarly, and in contact with hot conc. nitric acid and other oxidants it may explode (often after a delay with nitric acid). The powder is pyrophoric and readily ignitable by friction, heat or static sparks, and if dry burns fiercely. Presence of

water (5–10%) slightly reduces the ease of ignition, but combustion of the damp powder proceeds explosively (the oxygen content of water, 89%, being much higher than that of air). A minimum of 25% water is necessary to reduce handling hazards to a minimum. Full handling precautions are detailed.

See other METALS, PYROPHORIC METALS

4600. Mercury
[7439-97-6]

Hg

Hg

NSC 203, 1976; HCS 1980, 620; RSC Lab. Hazards Data Sheet No. 47, 1986

Acetylenic compounds

See 3-Bromopropyne: Metals 2.18/99+

2-Butyne-1,4-diol: Alkalies, etc.

Sodium acetylide: Metals

See ACETYLENIC COMPOUNDS: metals

Alkynes, Silver perchlorate

See Silver perchlorate: Alkynes, etc.

Ammonia

1. Sampey, J. J., *Chem. Eng. News.*, 1947, **25**, 2138

2. Braidech, M. M., *J. Chem. Educ.*, 1967, **44**, A324

3. Thodos, G., *Amer. Inst. Chem. Engrs. J.*, 1964, **10**, 274

4. Henderson, L. M., *Ind. Eng. Chem. (News Ed.)*, 1932, **10**, 73

5. Brunt, C. Van, *Science*, N. Y., 1927, **65**, 63–64

A mercury manometer used with ammonia became blocked by deposition of a grey-brown solid, which exploded during attempts to remove it mechanically or on heating. The solid appeared to be a dehydration product of Millon's base and was freely soluble in sodium thiosulfate solution. This method of cleaning is probably safer than others, but the use of mercury manometers with ammonia should be avoided as intrinsically unsafe [1,2]. Although pure dry ammonia and mercury do not react even under pressure at 340 kbar and 200°C, the presence of traces of water leads to the formation of an explosive compound, which may explode during depressurisation of the system [3]. Explosions in mercury–ammonia systems had been reported previously [4,5].

See Poly(dimercurymmonium hydroxide)

See also FULMINATING METALS

See other CORROSION INCIDENTS

Boron diiodophosphide

See Boron diiodophosphide: Metals

Ethylene oxide

MRH 3.72/99+

See Ethylene oxide(reference 4)

Metals

Bretherick, L., *Lab. Pract.*, 1973, **22**, 533

The high mobility and tendency to dispersion exhibited by mercury, and the ease with which it forms alloys (amalgams) with many laboratory and electrical contact metals, can cause severe corrosion problems in laboratories. A filter-cyclone trap is described to contain completely mercury ejected accidentally by overpressuring of mercury manometers and similar items.

See Aluminium: Mercury

Calcium: Mercury

Lithium: Mercury

Potassium: Mercury

Rubidium: Mercury

Sodium: Mercury

See other CORROSION INCIDENTS

Methyl azide

See Methyl azide: Mercury

Methylsilane, Oxygen

See Methylsilane: Mercury, etc.

Other reactants

Yoshida, 1980, 210

MRH values calculated for 6 combinations, largely with materials showing catalytic decomposition, are given.

Oxidants

MRH 5.69/99+

See Bromine: Metals

Chlorine: Metals (reference 1)

Chlorine dioxide: Mercury

Nitric acid: Alcohols (reference 2)

Peroxyformic acid: Metals, etc.

Tetracarbonylnickel, Oxygen

MRH Tetracarbonylnickel 0.88/99+

See Tetracarbonylnickel: Mercury, etc.

See other METALS

4601. Zinc amalgam

[72780-86-0]

Hg-Zn

Hg-Zn

Brimelow, H. C., private comm., 1972

Amalgamated zinc residues isolated from Clemmensen reduction of an alkyl aryl ketone in glacial acetic acid were pyrophoric, and had to be immediately dumped into water after filtration to prevent ignition.

See other PYROPHORIC ALLOYS

4602. Mercury(II) iodide
[7774-29-0]



Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

See other MERCURY COMPOUNDS, METAL HALIDES

4603. Mercury(II) nitrate
[10045-94-0]



HCS 1980, 616

Acetylene

Mellor, 1940, Vol. 4, 993

Contact of acetylene with the nitrate solution gives mercury acetylide, an explosive sensitive to heat, friction or contact with sulfuric acid.

Ethanol

Mellor, 1940, Vol. 4, 993

Addition of mercury(II) nitrate solution to ethanol gives mercury fulminate.

Ferrocene

Sallott, G. P. *et al.*, *Proc. Int. Pyrotech. Semin.*, 1984, 589–602

Compositions prepared from mercury(II) nitrate and ferrocene or its derivatives show promise as explosive priming mixtures, but such mixtures are fairly sensitive to electrostatic initiation and should be handled in the wet state.

Isobutene

See 2-Methyl-1-nitratodimercurio-2-nitratomercuriopropane

Petroleum hydrocarbons

1. Mixer, R. Y., *Chem. Eng. News*, 1948, **26**, 2434

2. Ball, J., *Chem. Eng. News*, 1948, **26**, 3300

Gas oil was stirred vigorously with the finely divided solid nitrate to complete the removal of sulfur compounds. After the mixture had congealed, preventing further stirring, a violent reaction set in which reached incandescence, accompanied by vigorous evolution of nitrous fumes. This was attributed to the self-catalysed nitrating action of mercury nitrate in a semi-solid environment unable to lose heat effectively. A similar occurrence was observed when crushed nitrate was just covered with cracked naphtha [1]. The second publication reveals that this type of hazard was known in cracked distillates containing a high proportion of unsaturates and aromatics, when allowed to stand in prolonged contact with mercury(II) nitrate. The hazard may be avoided by using several small portions of the salt sequentially, and working with 100 g portions of hydrocarbons [2].

Phosphine

Mellor, 1940, Vol. 9, 993

Phosphine reacts with the aqueous salt solution to give a complex nitrate-phosphide, which when dry explodes on heating or impact.

Phosphinic acid

Mellor, 1940, Vol. 4, 993

Phosphinic ('hypophosphorous') acid violently reduces the salt to the metal.

Potassium cyanide

Rüst, 1948, 337

Mixtures exploded when heated, but only if contained in narrow ignition tubes. Formation of nitrite, a more powerful oxidant than nitrate, may have been involved.

See other MERCURY COMPOUNDS, METAL NITRATES, OXIDANTS

4604. Mercury(II) azide

[14215-33-9]



HgN₆

1. Mellor, 1940, Vol. 8, 351; 1967, Vol. 8. Suppl. 2, 43, 50

2. Barton, A. F. M. *et al.*, *Chem. Rev.*, 1973, **73**, 138

3. Heathcock, C. H., *Angew. Chem. (Intern. Ed.)*, 1969, **8** 134

The very high friction-sensitivity, particularly of large crystals, and brisance on explosion are to be expected from the thermodynamic properties of the salt. Its great sensitivity, even under water, renders it unsuitable as a practical detonator [1]. Spontaneous explosions during intercrystalline transformations have been observed, or on crystallisation from hot water [2]. A safe method of preparing solutions in aqueous THF for synthetic purposes is available [3].

It is strongly endothermic (ΔH_f° (s) +556.5 kJ/mol, 1.96 kJ/g).

See other MERCURY COMPOUNDS, METAL AZIDES

4605. Mercury(II) oxide

[21908-53-2]



HgO

HCS 1980, 617

Under appropriate conditions, it can function as a powerful oxidant and/or catalyst, owing to the tendency to dissociate to metal and oxygen.

Acetyl nitrate

See Acetyl nitrate: Mercury(II) oxide

Butadiene, Ethanol, Iodine

See 2-Ethoxy-1-iodo-3-butene

Chlorine, Hydrocarbons

See Methane: Halogens, etc.

Ethylene: Chlorine

Diboron tetrafluoride

See Diboron tetrafluoride: Metal oxides

Disulfur dichloride

See Disulfur dichloride: Mercury oxide

Hydrogen peroxide

See Hydrogen peroxide: Mercury(II) oxide, or: Metals, etc.

Hydrogen trisulfide

See Hydrogen trisulfide: Metal oxides

Chlorine

1. Tabata, Y. *et al.*, *J. Haz. Mat.*, 1987, **17**, 55

2. Shilov, E. A., *Compt. Rend. Acad. Sci. USSR*, 1945, **46**, 64–65

Mercury(II) oxide reacts with chlorine to form dichlorine oxide which can explode at temperatures close to ambient [1]. This confirms a previous report [2].

Metals

1. Staudinger, H., *Z. Elektrochem.*, 1925, **312**, 549

2. Mellor, 1940, Vol. 4, 272

Mixtures of the red or yellow oxides with sodium–potassium alloy explode violently on impact, the yellow (more finely particulate) oxide giving the more sensitive mixture [1]. Mixtures with magnesium or potassium may explode on heating [2].

Methanethiol

Klason, P., *Ber.*, 1887, **20**, 3410

Interaction is rather violent in absence of diluent.

Non-metals

Mellor, 1940, Vol. 4, 777–778

Mixtures with phosphorus explode on impact or on boiling with water. A mixture with sulfur explodes on heating.

Phospham

See Phospham: Oxidants

Reducants

Mellor, 1940, Vol. 4, 778; Vol. 8, 318

Hydrazine hydrate and phosphinic acid both explosively reduce the oxide when dropped on to it.

See *other* MERCURY COMPOUNDS, METAL OXIDES, OXIDANTS

4606. Mercury peroxide
[12298-67-8]



Preparative hazard

See Hydrogen peroxide: Mercury(II) oxide, etc.

See other MERCURY COMPOUNDS, METAL PEROXIDES

4607. Mercury(II) sulfide
[1344-48-5]



Oxidants

Mellor, 1941, Vol. 2, 242; 1941, Vol. 3, 377; 1940, Vol. 4, 952

The sulfide causes dichlorine oxide to explode, and it incandescens in chlorine.

Grinding with silver oxide ignites the mixture.

See other MERCURY COMPOUNDS, METAL SULFIDES

4608. Poly(dimercuryimmonium permanganate)
[]




Sorbe, 1968, 97

It is highly explosive.

See other POLY(DIMERCURYIMMONIUM) COMPOUNDS

4609. Mercury(I) nitrate
[10415-75-5]



Carbon

Mellor, 1940, Vol. 4, 987



Contact with red-hot carbon causes a mild explosion.

Phosphorus

See Phosphorus: Nitrates

See other MERCURY COMPOUNDS, METAL NITRATES

4610. Mercury(I) thionitrosylate
[]





Goehring, M. *et al.*, *Z. Anorg. Chem.*, 1956, **285**, 702

It explodes on heating in a flame. The structure may be the indicated linear polymer of tetrasulfur tetraimidate rings linked by pairs of dimercury bonds.

See other MERCURY COMPOUNDS, N-S COMPOUNDS

4611. Poly(dimercuryimmonium azide)

[]




Sorbe, 1968, 97

It is highly explosive.

See other POLY(DIMERCURYIMMONIUM) COMPOUNDS *See related* METAL AZIDES

4612. Mercury(I) azide

[38232-63-2]



Mellor, 1940, Vol. 8, 351; 1967, Vol. 8, Suppl. 2, 25, 50

It is less sensitive and a less powerful explosive than silver azide or lead azide. It explodes on heating in air to above 270°C, or after an induction period at 140°C in the dark under vacuum.

See other INDUCTION PERIOD INCIDENTS, MERCURY COMPOUNDS, METAL AZIDES

4613. 'Mercury(I) oxide'

[15829-53-5]



Sidgwick, 1950, 292

The material is known to be an intimate mixture of mercury(II) oxide and finely dispersed metallic mercury.

Alkali metals

Mellor, 1940, Vol. 4, 771

Interaction with molten potassium or sodium is accompanied by a brilliant light and a light explosion.

Chlorine, Ethylene

See Ethylene: Chlorine

Hydrogen peroxide

MRH 1.59/99+

Antropov, V., *J. Prakt. Chem.*, 1908, **77**, 316

Contact cause explosive decomposition of the peroxide.

See Hydrogen peroxide: Metals, etc.

Non-metals

Mellor, 1940, Vol. 4, 771

Mixtures with phosphorus explode on impact, and that with sulfur ignites on frictional initiation.

See other MERCURY COMPOUNDS, METAL OXIDES

4614. Tris(iodomercurio)phosphine

[]



Nitric acid

See Nitric acid: Phosphine derivatives

See related MERCURY COMPOUNDS, PHOSPHINES

4615. Mercury nitride

[12136-15-1]



Alone, or Sulfuric acid.

1. Mellor, 1940, Vol. 8, 108

2. Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1469

3. Wilhelm, S. M., *Plant/Oper. Progr.*, 1991, **10**(4), 192

It explodes on friction, impact, heating or in contact with sulfuric acid [1]. A sample at below -40°C exploded when disturbed [2]. It may form from the metal and ammonia in some circumstances [3]

See other FULMINATING METALS, MERCURY COMPOUNDS, N-METAL DERIVATIVES

4616. Trimercury tetraphosphide

[12397-29-4]



Oxidants

Mellor, 1940, Vol. 8, 844



It ignites when warmed in air, or cold on contact with chlorine. A mixture with potassium chlorate explodes on impact.

See Potassium chlorate: Metal phosphides

See other MERCURY COMPOUNDS, METAL NON-METALLIDES

4617. Mercury(I) hypophosphate

[]



Mellor, 1971, Vol. 8, Suppl. 3, 651
Explosive decomposition occurs on heating.
See other MERCURY COMPOUNDS

4618. Potassium iodide

[7861-11-0]

KI

IK

HCS 1980, 770

Charcoal, Ozone

See Ozone: Charcoal, Potassium iodide

Diazonium salts

See DIAZONIUM TRIIODIDES

Diisopropyl peroxydicarbonate

See Diisopropyl peroxydicarbonate (reference 2)

Oxidants

See Bromine pentafluoride: Acids, etc.

Chlorine trifluoride: Metals, etc.

Fluorine perchlorate: Alone, etc.

Trifluoroacetyl hypofluorite

See other METAL HALIDES

4619. Potassium iodate

[7758-05-6]

KOIO₂

IKO₃

HCS 1980, 769

Charcoal, Ozone

See Ozone:Charcoal, Potassium iodide

Metals and oxidisable derivatives

See METAL HALOGENATES: METALS AND OXIDISABLE DERIVATIVES

Phosphorus: Metal halogenates

Other reactants

Yoshida, 1980, 376

MRH values calculated for 27 combinations with oxidisable materials are given.

Sodium disulfite, Water

Association for Science Education, Internet, 1997

A student, who should have mixed solutions of these two salts, mixed the solids and added water, which provoked a violent reaction. It is suggested that pupils should not be allowed access to solid, undiluted, oxidants and reductants at the same time.

See other IODINE COMPOUNDS, METAL HALOGENATES, OXIDANTS

4620. Potassium periodate
[7790-21-8]

KOIO₃

IKO₄

HCS 1980, 773

Ammonium perchlorate

See Ammonium perchlorate: Impurities

See other IODINE COMPOUNDS, METAL OXOHALOGENATES, OXIDANTS

4621. Iodine azide
[14696-82-3]

IN₃

IN₃

1. Hantzsch, A., *Ber.*, 1900, **33**, 525
2. Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1976, **15**, 553
3. Dehnicke, K., *Angew. Chem. (Intern. Ed.)*, 1979, **18**, 507–514
4. Klapötke, T. M. *et al.*, *Angew. Chem. (Int.)*, 1993, **32**(2), 275

The isolated solid is a very shock- and friction-sensitive explosive [1], but the preparation and safe handling of dilute solutions in solvents other than ether have been described [2]. The need to use appropriate techniques and precautions when using iodine azide as a reagent is stressed [3]. The purer the more explosive; explosive properties are characterised (lead-block test, etc.) in a footnote to [4].

Sulfur-containing alkenes

Hassner, A. *et al.*, *J. Org. Chem.*, 1968, **33**, 2690

Interaction is accompanied by violent decomposition of the azide.

See other HALOGEN AZIDES

4622. Iodinated poly(sulfur nitride)

[]

(Indeterminate structure)

(IN_{4.2}S₆)_n

Banister, A. J. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 937

The polymeric product is potentially explosive.

See other IODINE COMPOUNDS, N–S COMPOUNDS

4623. Sodium iodide
[7681-82-5]

NaI

INa

HCS 1980, 849

1713

Oxidants

See Perchloryl fluoride: Calcium acetylide, etc.

Perchloric acid: Iodides

See other METAL HALIDES

4624. Sodium iodate

[7681-55-2]

INaO₃

NaOIO₂

Metals

MRH Magnesium 5.98/27

Webster, H. A. *et al.*, *Rept. AD-782510/2GA*, Springfield (Va.), USNTIS, 1974

The use of mixtures with magnesium in pyrotechnic flares is discussed.

See Potassium: Oxidants

MRH 1.42/31

Sodium: Iodates

MRH 3.10/42

Other reactants

Yoshida, 1980, 381–382

MRH values calculated for 27 combinations with oxidisable materials are given.

See other IODINE COMPOUNDS, METAL HALOGENATES, OXIDANTS

4625. Iodine

[7553-56-2]

I₂

I₂

(*MCA SD-43*, 1952); *NSC 457*, 1968; *HCS 1980*, 560

Acetaldehyde

MCA SD-43, 1952

Interaction may be violent.

Acetylene

MRH 8.70/99+

See Acetylene: Halogens

Aluminium, Diethyl ether

Garcia, F. J. A., *Inorg. Chem.*, 1980, **20**, 83

In the preparation of aluminium iodide etherate from the elements in ether, the aluminium must be as turnings, rather than fine powder, to keep the reaction under control.

See Aluminium: Halogens

Ammonia

Mellor, 1940, Vol. 8, 605; 1967, Vol. 8, Suppl. 2.2, 416

Ammonia solutions react with iodine (or potassium iodide) to produce highly explosive addition compounds of nitrogen triiodide and ammonia.

See Nitrogen triiodide–ammonia

Ammonia, Lithium 1-heptynide

Houben-Weyl, 1977, Vol. 5.2a, 605

Reaction of iodine with the lithium heptynide in liquid ammonia to give (explosive) 1-iodoheptyne may lead to formation of nitrogen triiodide as a black precipitate. Low temperatures minimise the formation, and it may be destroyed by adding sodium ethoxide solution.

Ammonia, Potassium

See Ammonia: Iodine, Potassium

Butadiene, Ethanol, Mercuric oxide

See 2-Ethoxy-1-iodo-3-butene

Caesium oxide

See Caesium oxide: Halogens

Dipropylmercury

See Dipropylmercury: Iodine

Ethanol, Phosphorus

1. Read, C. W. W., *School Sci. Rev.*, 1940, **21**(83), 967

2. Llowarch, D., *School Sci. Rev.*, 1956, **37**(133), 434

Interaction of ethanol, phosphorus and iodine to form iodoethane was considered too dangerous for a school experiment [1]. A safer modification is now available [2].

See Phosphorus: Halogens

Formamide, Pyridine, Sulfur trioxide

See Formamide: Iodine, etc.

Halogens, or Interhalogens

See Bromine trifluoride: Halogens

Bromine pentafluoride: Acids, etc.

Chlorine trifluoride: Iodine

Fluorine: Halogens, etc.

Metal acetylides or carbides

Several react very exothermally with iodine.

See Barium acetylide: Halogens

Calcium acetylide: Halogens

MRH 1.46/20

Caesium acetylide: Halogens

Dicopper(I) acetylide: Halogens

Lithium acetylide: Halogens

Rubidium acetylide: Halogens

Strontium acetylide: Halogens

MRH 1.34/31

Zirconium dicarbide: Halogens

Metals

MRH Lithium 2.01/5, potassium 1.97/24

1. Mellor, 1941, Vol. 2, 469; 1963, Vol. 2, Suppl. 2.2, 1563; 1939, Vol. 9, 379

2. Tarnqvist, E. G. M. *et al.*, *Inorg. Chem.*, 1979, **18**, 1793

Iodine and antimony powder react so violently as to cause ignition or explosion of the bulk of the mixture. A mixture of potassium and iodine explodes weakly on impact, while potassium ignites in contact with molten iodine [1]. Interaction of molten iodine with titanium above 113°C under vacuum to form titanium tetraiodide is highly exothermic and sparks are produced. The preparative technique described permits the progressive reaction of 0.5 g portions of the titanium powder charged (7.2 g) to minimise hazard [2].

See Aluminium: Oxidants

Aluminium–titanium alloys: Oxidants

Hafnium: Alone, etc.

Sodium: Halogens (reference 5)

Metals, Water

Jackson, H., *Spectrum*, 1969, **7**, 82

Flash-ignition occurs when mixtures of iodine with powdered aluminium, magnesium or zinc are moistened with a drop of water.

Non-metals

See Boron: Halogens

Phosphorus: Halogens

Other reactants

Yoshida, 1980, 372

MRH values calculated for 8 combinations, largely with oxidisable materials, are given.

Oxygen difluoride

See Oxygen difluoride: Halogens, etc.

Poly(acetylene)

Pekker, S. *et al.*, *Chem. Abs.*, 1983, **99**, 213009

In TGA studies on the decomposition of iodine-doped polyacetylene, at high heating rates (30°C/min), decomposition becomes explosive at the m.p. of iodine, 113°C. This was attributed to exothermic reaction of liquid iodine with polyacetylene.

Silver azide

MRH 2.05/99+

See Silver azide: Halogens

Sodium phosphinate

See Sodium phosphinate: Oxidants

Tetraamminecopper(II) sulfate, Ethanol

Schwarzenbach, V., *Ber.*, 1875, **8**, 1233

Addition of ethanol to a mixture of iodine and the salt soon led to explosions of variable intensity, involving formation of *N*-iodo derivatives.

See Mercury(II) amide chloride: Halogens


Trioxxygen difluoride

See 'Trioxxygen difluoride': Various materials

See other HALOGENS, OXIDANTS

4626. *N,N'*-Diiodosulfurdiimide

[]



I₂N₂S

Seppelt, K. *et al.*, *Angew. Chem. (Int.)*, 1969, **8**, 771

This explodes both on melting and impact. The bromo-analogue is more sensitive, the chloro-compound is not known.

See other *N*-HALOGEN COMPOUNDS, N-S COMPOUNDS, IODINE COMPOUNDS

4627. Iodine(V) oxide

[12029-98-0]



I₂O₅

Aluminium

Ivanov, V. G. *et al.*, *Chem. Abs.*, 1981, **95**, 9423

Combustion of compressed mixtures of metal powder and the oxide is very vigorous, a layer of molten aluminium being produced initially, followed by a jet of flame.

Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

Hydrazine

See Hydrazine: Iodine pentoxide

Non-metals

Mellor, 1941, Vol. 2, 295

Iodine pentoxide reacts explosively with warm carbon, sulfur, rosin, sugar or powdered, easily combustible elements

See other HALOGEN OXIDES, IODINE COMPOUNDS, OXIDANTS

4628. Iodine(VII) oxide

[12055-24-2]



I₂O₇

Diethyl ether

Mishra, H. C. *et al. J. Chem. Soc.*, 1962, 1195–1196

1717

Washing of the (incompletely characterised) solid with ether occasionally led to explosive decomposition.

See other HALOGEN OXIDES, IODINE COMPOUNDS, OXIDANTS

4629. Phosphorus diiodide triselenide
[30911-12-7]



Nitric acid

See Nitric acid: Phosphorus compounds

See related IODINE COMPOUNDS, NON-METAL HALIDES, NON-METAL SULFIDES

4630. Titanium diiodide
[13783-07-8]



Preparative hazard

Gibson, 1969, 63

It may ignite in moist air.

See Titanium: Halogens (reference 3)

See other METAL HALIDES, PYROPHORIC MATERIALS

4631. Tungsten diiodide
[13470-17-2]



Preparative hazard

Mussell, R. D. *et al.*, *Inorg. Chem.*, 1990, **29**(19), 3711

Explosions sometimes resulted on opening sealed tubes in which complex mixed halide salts of this compound had been prepared. There is no obvious source of pressure in the reaction mixture of dipotassium tetrachlorotungstate, lithium iodide and potassium iodide.

See also Tungsten dichloride *See other* METAL HALIDES

4632. Zinc iodide
[10139-47-6]



Sulfuric acid

See Sulfuric acid: Zinc iodide

See other METAL HALIDES

4633. Nitrogen triiodide (Nitrogen iodide)

[13444-85-4]

I₃NKlapötke, T. M. *et al.*, *Angew. Chem. (Int.)*, 1990, **29**(6), 677

The uncomplexed material has been prepared and is very explosive, decomposing spontaneously at 0°C or below. Like its bromine and chlorine analogues, it is highly endothermic.

See next two below See other N-HALOGEN COMPOUNDS, ENDOTHERMIC COMPOUNDS, IODINE COMPOUNDS

4634. Nitrogen triiodide–silver amide

[]

I₃N·AgH₂N


Mellor, 1967, Vol. 8. Suppl. 2, 418

The dry complex may explode.

See other N-HALOGEN COMPOUNDS, IODINE COMPOUNDS, N-METAL DERIVATIVES, SILVER COMPOUNDS

4635. Nitrogen triiodide–ammonia

[34641-74-2], [14014-86-9] (1:1)

I₃N·H₃N

Alone, or Halogens, or Oxidants, or Concentrated acids

1. Mellor, 1940, Vol. 8, 607; 1967, Vol. 8, Suppl. 2, 418
2. Garner, W. E. *et al.*, *Nature*, 1935, **135**, 832
3. Taylor, G. N., *Chem. Brit.*, 1981, **17**, 107
4. Houben-Weyl, 1977, Vol. 5.2a, 605
5. Bodner, G. M., *J. Chem. Educ.*, 1985, **62**(12), 1107
6. Bentzinger, von R. *et al.*, *Praxis Naturwiss. Chem.*, 1987, **36**, 37

Readily formed in systems containing ammonia and iodine or some of its derivatives, the addition compound of nitrogen triiodide and ammonia when dry is an extremely sensitive, unstable detonator capable of initiation by minimal amounts of any form of energy (light, heat, sound, nuclear radiation, mechanical vibration), even at sub-zero temperatures, and occasionally even with moisture present. It may be handled cautiously when wet, but heavy friction will still initiate it. It explodes in boiling water and is decomposed by cold water to explosive diiodamine. It explodes, possibly owing to heat-initiation on contact with virtually any conc. acid, with chlorine or bromine, ozone or hydrogen peroxide solution [1]. Crystals desiccated in vacuum spontaneously explode when dry [2]. It has been formed in relatively large amounts during preparation of iodoacetylene derivatives in liquid ammonia [3,4], and may be destroyed by addition of sodium ethoxide [4]. Accidents involving demonstration of the explosive properties have been summarised [5]. A student prepared some to spread about on the floor as a joke, but it exploded

prematurely while still in contact with the solution, causing eye injuries to the joker [6].

See other N-HALOGEN COMPOUNDS, IODINE COMPOUNDS

4636. Phosphorus triiodide

[13455-01-1]



I₃P

Hydroxylic compounds, or Oxygen

Leleu, *Cahiers*, 1974, (75), 273

Interaction with water or glycerol is violent, and the iodide ignites in oxygen.

Sulfur acids

Dillon, K. B. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1979, 887–888

Contact with 25 or 50% oleum led to violently exothermic reactions, and vigorous reaction with chlorosulfuric acid.

See Phosphorus tribromide: Sulfur acids

Phosphorus trichloride: Sulfur acids

See other IODINE COMPOUNDS, HALOPHOSPHINES, NON-METAL HALIDES

4637. Tetraiododiphosphane

[13455-00-0]



I₄P₂

Preparative hazard

See Phosphorus: Halogens (references 3,4)

See other HALOPHOSPHINES, IODINE COMPOUNDS, NON-METAL HALIDES

4638. Titanium tetraiodide

[7720-83-4]



I₄Ti

Preparative hazard

Denes, G., *J. Solid State Chem.*, 1988, **77**(1), 54–59

Use of a 'bent copper tube' reactor to prepare the iodide is hazardous.

See Titanium: Halogens

See other METAL HALIDES

4639. Zirconium tetraiodide

[13986-26-0]



I₄Zr

Ethanol

Pascal, 1963, Vol. 9, 565

Interaction is very violent.
See other METAL HALIDES

4640. Indium
[7440-74-6]

In

In

Acetonitrile, Dinitrogen tetraoxide
See Dinitrogen tetraoxide: Acetonitrile, etc.

Mercury(II) bromide
See Mercury(II) bromide: Indium

Sulfur
See Sulfur: Metals
See other METALS

4641. Indium(II) oxide
[12136-26-4]

InO

InO

Ellern, 1968, 33
The lower oxide, prepared by hydrogenation, incandesces on exposure to air.
See other METAL OXIDES, PYROPHORIC MATERIALS

4642. Indium phosphide
[22398-80-7]

InP

InP

Preparative hazard
See Phosphorus: Metals
See METAL Pnictides

4643. Iridium
[7439-88-5]

Ir

Ir

The finely divided catalytic metal may be pyrophoric.
See Zinc: Catalytic metals *See entry* HYDROGENATION CATALYSTS

Interhalogens
See Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Metals
See other METALS, PYROPHORIC METALS

4644. Iridium(IV) oxide
[12030-49-8]

IrO₂

IrO₂

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

See other METAL OXIDES

4645. Potassium
[7440-09-7]

K

K

FPA H38, 1975; HCS 1980, 759

1. Gilbert, H. N., *Chem. Eng. News*, 1948, **26**, 2604
2. Johnson, W. S. *et al.*, *Org. Synth.*, 1963, Coll. Vol. 4, 134–135
3. *MCA Case History No. 1891*
4. Taylor, D. A. H., *Chem. Brit.*, 1974, **10**, 101
5. Diaper, D. G. M., *Chem. Brit.*, 1974, **10**, 312
- 5a. Anon., *Fire Precaut.*, 1988, (213), 46
6. Houben-Weyl, 1970, Vol. 13.1, 264
7. Burfield, D. R. *et al.*, *Chem. & Ind.*, 1979, 89
8. Davis, A. C., *Chem. Brit.*, 1979, **15**, 179
9. Levy, J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1981, **20**, 1033
10. Mellor, 1941, Vol. 2, 468; 1963, Vol. 2, Suppl. 2.2, 1559
11. Mellor, 1941, Vol. 2, 493
12. Short, J. F., *Chem. & Ind.*, 1964, 2132
13. Brazier, A. D., *Chem. & Ind.*, 1965, 220; Balfour, A. E., *ibid.*, 353; Bil, M. S., *ibid.*, 812; Cole, R. J., *ibid.*, 944
14. March, R. G., *Chem. Brit.*, 1979, **15**, 65
15. Brock, T. H. *et al.*, *Nachr. Chem. Tech. Lab.*, 1998, **46**(1), 16
16. Brandsma, 1971, 10, 21

In a review of the comparative properties of sodium and potassium, the latter is rated as invariably the more hazardous [1]. Laboratory procedures for safe handling of potassium have been detailed [2]. A safe method for disposal of potassium residues in bulk storage and processing vessels has been developed. Basically the method involves reaction of the metallic residues with dry (condensate-free) steam under closely controlled conditions. It may be used where small-scale techniques (dissolution in higher alcohols) are inapplicable [3]. Hazards associated with storage of potassium in aluminium containers are discussed. The severe corrosion leading to perforation of such commercially supplied containers [4] was attributed to the deliquescent and subsequently corrosive nature of potassium carbonate formed from atmospheric carbon dioxide inside the container. Sodium may be safely stored in aluminium cans because the derived sodium carbonate is not deliquescent and causes no corrosion [5]. A serious laboratory fire originating in a store cupboard containing potassium in an aluminium container may have arisen

K

through corrosive failure of the latter [5]. There is a considerable risk of fire if powdered potassium ('potassium sand') dispersed in benzene is exposed to air [6]. A practical survey of methods of safe handling and disposal of potassium covers the separation of metal from adhering oxide by melting under xylene containing 1% of isopropanol, disposal of small metal residues by treatment with 1:1 ethyl acetate–hydrocarbon diluent mixtures, and disposal of 30 g blocks by dropping into a 2 m deep hole in the ground containing 0.5 m of water [7]. Several fires (but not explosions) occurred when potassium was 'blotted' free of solvent (used to remove traces of oil) with previously used filter paper. The fires were attributed to rapid formation of potassium hydroxide solution from traces of potassium on the filter paper under humid conditions, and ignition of the dry metal on contact with the wet alkali [8]. Potassium dispersed on silica is relatively air stable [9], unlike the dispersions on sodium carbonate or potassium carbonate which are pyrophoric.

It is convenient to consider interaction of potassium and air under two headings.

Rapid oxidation

Reaction with air or oxygen in complete absence of moisture does not occur, even on heating to the boiling point. However, in contact with normal (moist) air, oxidation may become so fast that melting or ignition occurs, particularly if pressure is applied locally to cause melting and exposure of a fresh surface, as when potassium is pressed through a die to form potassium wire [10].

Slow oxidation

Metallic potassium on prolonged exposure to air forms a coating of yellowish potassium superoxide (KO_2) under which is a layer of potassium oxide in contact with the metal [11]. The previous statements that normal contact of potassium with the superoxide causes ignition to occur [11], and that if the layer of superoxide is impacted into the underlying metal by dry cutting operations or a hammer blow an explosion occurs [1,12] are now known to be incorrect. The explosions are caused by interaction of residual traces of mineral oil or other organic contaminant, rather than potassium metal, with the surface layer of superoxide, initiated by the blade pressure or impact. Cases of flashes of light and fires [13] or explosion [14] when potassium (some were old samples) was cut under oil have been reported. A review with references on explosion hazards of potassium, gives particular attention to oxidised crusts [15].

Fresh potassium should be stored under dry xylene in airtight containers to prevent oxidation [11]. Old stocks, where the coating is orange or yellow, should not be cut, but destroyed by burning on an open coke fire, or by addition of *tert*-butanol to small portions under xylene in a hood. It is dangerous to use methanol or ethanol (either wet or dry) as a replacement for *tert*-butanol. Recommended procedures include cutting and handling the metal with forceps under dry xylene, and disposal of scraps in xylene by addition of *tert*-butanol [1,12,13], even the latter being capable of violent reaction [16]. Alternatively, ethyl acetate–hydrocarbon mixtures may be used for small disposals [7].

See Potassium dioxide (reference 3)

See Potassium–sodium alloy

Acetylene

See Acetylene: Potassium

See other CORROSION INCIDENTS

Alcohols

Pratt, E. F. *et al.*, *J. Amer. Chem. Soc.*, 1954, **75**, 54

Interaction with a range of alcohols (n-propanol to n-octanol, benzyl alcohol, cyclohexanol) to form the alkoxides usually led to explosions unless air in the containing vessel was displaced by nitrogen before addition of potassium in small portions with stirring.

See Slow oxidation, above

Aluminium, Air

Anon., *Fire Prevention*, 1988, (213), 46

After a fire which started in a disused fume cupboard employed as a storage space for various inorganic chemicals, including potassium, it is suggested that the potassium oxides which inevitably coat aged potassium may have corroded the aluminium can, releasing the oil in which the metal is stored and exposing it to accelerated oxidation. This argument is thermodynamically sound, but kinetically questionable as it involves a solid/solid interaction of potassium and aluminium oxides.

Carbon

1. Mellor, 1963, Vol. 2, Suppl. 2.2, 1566

2. Mellor, D. P., *Chem. & Ind.*, 1965, 723

3. *Alkali metals*, 1957, 169

Reaction of various forms of carbon (soot, graphite or activated charcoal) is exothermic and vigorous at elevated temperatures, and if the carbon is finely divided and air is present, ignition may occur leading to explosions, possibly owing to the potassium superoxide which would be formed [1]. Explosions caused by attempts to extinguish potassium fires with graphite powder have been so attributed [2]. Potassium cannot be produced by electrolysis of potassium chloride because of interaction of the metal with graphite electrodes and formation of explosive 'carbonylpotassium' (potassium benzenehexoxide) [3].

Carbon disulfide

See Carbon disulfide: Alkali metals

Dimethyl sulfoxide

See Dimethyl sulfoxide: Potassium

Ethylene oxide

MRH 3.72/99+

See Ethylene oxide: Contaminants

Halocarbons

MRH values below references

1. Staudinger, H., *Z. Angew. Chem.*, 1922, **35**, 657

2. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

3. Lenze, F. *et al. Chem. Ztg.*, 1932, **56**, 921

4. Rampino, L. D., *Chem. Eng. News*, 1958, **36**(32), 62

MRH Bromoform 3.14/68, carbon tetrachloride 5.19/50, chloroform 4.93/50, pentachloroethane 4.98/51, tetrachloroethane 4.77/52, tetrachloroethylene 5.23/51
Although apparently stable on contact, mixtures of potassium (or its alloys) with a wide range of halocarbons are shock-sensitive and may explode with great violence on light impact. Chloroethane, dichloroethane, trichloroethane, pentachloroethane, bromoform, dibromomethane and diiodomethane are among those investigated. Sensitivity increases generally with the degree of substitution, and potassium–sodium alloy gives extremely sensitive mixtures. The mixture with carbon tetrachloride is 150–200 times as shock-sensitive as mercury fulminate, and a mixture of potassium with bromoform was exploded by a door slamming nearby [1,2]. Mixtures with tetrachloroethane and pentachloroethane will often explode after a short delay during which a voluminous solid separates out [3]. When heated together, potassium and tetrachloroethylene exploded at 97–99°C, except when the metal had been very recently freed of its usual oxide film. Sodium did not explode under the same conditions [4].

See Oxalyl dihalides, below

Halogens or Interhalogens MRH Bromine 3.22/68, chlorine 5.77/48

1. Mellor, 1941, Vol. 2, 114, 469; 1963, Vol. 2, Suppl. 2.2, 1563

2. Pascal, 1960, Vol. 16, 578

Potassium ignites in fluorine and in dry chlorine (unlike sodium). In bromine vapour it incandescs, and explodes violently in liquid bromine. Mixtures with iodine incandescs on heating, and explode weakly on impact. Potassium reacts explosively with molten iodine bromide and iodine, and a mixture with the former is shock-sensitive and explodes strongly. Molten potassium reacts explosively with iodine pentafluoride [1]. Contact with iodine trichloride causes ignition [2].

Hydrazine

See Hydrazine: Potassium

Hydrogen halides MRH Hydrogen chloride 3.72/48

1. Cueilieron, J., *Bull. Soc. Chim. Fr.*, 1945, **12**, 88

2. Pascal, 1963, Vol. 2.2, 31

Impact causes a mixture of potassium and anhydrous hydrogen chloride to explode very violently [1]. Molten potassium ignites in contact with hydrogen chloride, hydrogen bromide or hydrogen iodide [2].

Magnesium halides, Potassium iodide

Rieke, R. D. *et al.*, *Org. Synth.*, 1979, **59**, 85–94

Interaction of potassium with magnesium bromide, magnesium chloride or magnesium iodide in refluxing THF produces very finely divided and highly reactive magnesium which will ignite if exposed long to air.

See PYROPHORIC METALS

Maleic anhydride

See Maleic anhydride: Bases, etc.

Mercury

Mellor, 1941, Vol. 2, 469

Interaction to form amalgams is vigorously exothermic and may become violent if too much potassium is added at once.

Metal halides

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549
2. Cueilleron, J., *Bull. Soc. Chim. Fr.*, 1945, **12**, 88
3. Mellor, 1963, Vol. 2, Suppl. 2.2, 1571
4. Rieke, R. D., *Accts. Chem. Res.*, 1977, **10**, 301–305

Cobalt(II) chloride 2.59/62, iron(II) bromide 1.80/74, iron(III) bromide 1.88/72, iron(III) chloride 3.22/58, iron(II) iodide 1.38/80, aluminium chloride 2.43/53, copper(I) chloride 2.18/72, copper(II) chloride 3.10/63, silver fluoride 2.18/76

Mixtures of potassium with metal halides are sensitive to mechanical shock, and the ensuing explosions have been graded. Very violent explosions occurred with calcium bromide, iron(III) bromide, iron(III) chloride, iron(II) bromide, iron(II) iodide or cobalt(II) chloride. Strong explosions occurred with silver fluoride, mercury(II) bromide, mercury(II) chloride, mercury(II) fluoride, mercury(II) iodide, copper(I) bromide, copper(I) chloride, copper(I) iodide, copper(II) bromide, copper(II) chloride, and ammonium tetrachlorocuprate; zinc and cadmium chlorides, bromides and iodides; aluminium fluoride, chloride, or bromide, thallium(I) bromide; tin(II) or (IV) chloride, tin(IV) iodide (with sulfur), arsenic trichloride and triiodide, antimony and bismuth tribromides, trichlorides and triiodides; vanadium(V) chloride, chromium(IV) chloride, manganese(II) and iron(II) chlorides, and nickel bromide, chloride or iodide. Weak explosions occurred with a wide variety of other halides [1,2,3]. These reactions may be moderated by the use of ether or hydrocarbon solvents. The reactions are still highly exothermic but controllable, and give finely divided and highly reactive metals, some pyrophoric in air, such as magnesium and aluminium [4].

See Magnesium halides, etc., above

Metal oxides

1. Mellor, 1963, Vol. 2, Suppl. 2.2, 1571
2. Mellor, 1941, Vol. 3, 138
3. Mellor, 1940, Vol. 4, 770, 779
4. Mellor, 1941, Vol. 7, 401
5. Mellor, 1943, Vol. 11, 237, 542
6. Mellor, 1941, Vol. 9, 649

Lead peroxide reacts explosively [1] and copper(II) oxide incandescently [2] with warm potassium. Mercury(II) and (I) oxides react with molten potassium with incandescence and explosion, respectively [3]. Tin(IV) oxide is reduced incandescently on warming [4] and molybdenum(III) oxide on heating [5]. Warm bismuth trioxide is reduced with incandescence [6].

See 'Mercury(I) oxide': Alkali metals

Nitric acid

Pascal, 1963, Vol. 2.2, 31

Interaction with conc. nitric acid is explosive.

Nitrogen-containing explosives

Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

Nitro or nitrate explosives, normally shock-insensitive, are rendered extremely sensitive by addition of traces of potassium or potassium–sodium alloy. Ammonium nitrate, and nitrate–sulfate mixtures, picric acid, and even nitrobenzene respond in this way.

Non-metal halides

1. Mellor, 1963, Vol. 2, Suppl. 2.2, 1564–1568

2. Mellor, 1940, Vol. 8, 1006

3. Mellor, 1947, Vol. 10, 642–646, 908, 912

Diselenium dichloride and seleninyl chloride both explode on addition of potassium [1,3], while the metal ignites in contact with phosphorus trichloride vapour or liquid [2]. Mixtures of potassium with sulfur dichloride or sulfur dibromide, phosphorus tribromide or phosphorus trichloride, and with phosgene are shock-sensitive, usually exploding violently on impact. Potassium also explodes violently on heating with disulfur dichloride, and with sulfur dichloride or seleninyl bromide without heating [3].

Non-metal oxides

1. Gilbert, H. N., *Chem. Eng. News*, 1948, **26**, 2604

2. Mellor, 1941, Vol. 2, 241

3. Mellor, 1940, Vol. 8, 436, 544, 554, 945

4. Pascal, 1963, Vol. 2.2, 31

MRH Dinitrogen oxide 3.72/54, nitrogen oxide 4.60/61

Mixtures of potassium and solid carbon dioxide are shock-sensitive and explode violently on impact, and carbon monoxide readily reacts to form explosive ‘carbonylpotassium’ (potassium benzenehexoxide) [1]. Dichlorine oxide explodes on contact with potassium [2]. Potassium ignites in dinitrogen tetraoxide or dinitrogen pentaoxide at ambient temperature and incandescs when warmed with nitrogen oxide or phosphorus(V) oxide [3]. At -50°C , potassium and carbon monoxide react to give dicarbonylpotassium, which explodes in contact with air or water, or at 100°C . At 150°C , the product is a trimer of this, potassium benzenehexoxide. The just-molten metal ignites in sulfur dioxide [4].

Organic samples for qualitative analysis

See LASSAIGNE TEST (reference 2)

Other reactants

Yoshida, 1980, 96–97

MRH values calculated for 49 combinations with a wide variety of other materials are given.

Oxalyl dihalides

Staudinger, H., *Z. Angew. Chem.*, 1922, **35**, 657; *Ber.*, 1913, **46**, 1426

In absence of mechanical disturbance, potassium or potassium–sodium alloy appears to be stable in contact with oxalyl dibromide or oxalyl dichloride, but the mixtures are very shock-sensitive and explode very violently.

See Halocarbons, above

Oxidants

Mellor, 1963, Vol. 2, Suppl. 2.2, 1571

The potential for violence of interaction between the powerful reducing agent potassium and oxidant classes has been well described. Other miscellaneous oxygen-containing substances which react violently or explosively include sodium iodate, silver iodate, lead sulfate and boric acid.

See Metal oxides, *also* Non-metal oxides, both above

Selenium, etc., *also* Sulfur, *and* Sulfuric acid, all below

Ammonium nitrate: Ammonium sulfate, etc.

Chlorine trifluoride: Metals

Chromium trioxide: Alkali metals

Dichlorine oxide: Oxidisable materials

Nitryl fluoride: Metals

Selenium, or Tellurium

Mellor, 1947, Vol. 10, 767; 1943, Vol. 11, 40

Interaction of selenium and potassium is exothermic and ignition occurs, or, with excess potassium, a slight explosion. Tellurium and potassium become incandescent when warmed in a hydrogen atmosphere to prevent aerobic oxidation.

Sulfur

Pascal, 1963, Vol. 2.2, 30

Interaction is violent on warming.

Sulfuric acid

MRH 3.57/56

Kirk-Othmer, 1961, Vol. 16, 362

Interaction is explosive.

Water

1. Mellor, 1941, Vol. 2, 469; 1963, Vol. 2, Suppl. 2.2, 1560

2. Angus, L. H., *School Sci. Rev.*, 1950, **31**(115), 402

3. Markowitz, M. M., *J. Chem. Educ.*, 1963, **40**, 633–636

4. Ashman, A., *School Sci. Rev.*, 1983, **65**(231), 387

Interaction is violently exothermic, and the heat evolved with water at 20°C is enough to ignite evolved hydrogen. Larger pieces of potassium invariably explode in water and scatter burning potassium particles over a wide area. Aqueous alcohols should not be used for waste metal disposal [1]. Small pieces of potassium will also explode with a restricted amount of water [2]. The reactivity of potassium and other alkali metals with water has been discussed in detail. The vigour of

reaction is scarcely reduced by contact with ice–water slurries [3]. A convenient demonstration of ignition of potassium by placing a small piece on an ice cube is due to Faraday [4].

See ALKALI METALS See other METALS, REDUCANTS

4646. Potassium–sodium alloy
[12532-69-3]

K–Na

K–Na

1. *Health and Safety Information*, 251, Washington, USAEC, 1967
2. Ellis, J. E. *et al.*, *Inorg. Synth.*, 1976, **16**, 70
3. Mellor, 1961, Vol. 2, Suppl. 2.1, 562
4. Anon., *Fire Prot. Assoc. J.*, 1965, **68**, 1402

Appropriate precautions in handling the alloy, used as a thermally stable and radiation-resistant liquid coolant for reactor cores, are described. It is generally more hazardous than either of the component metals, because alloys in the range 50–80 wt% of potassium are liquid at ambient temperature and can therefore come into more intimate contact with reagents than the solid metals. Most of the entries under Potassium above, and some of those under Sodium may be expected to apply to their alloys, with allowance for the composition [1]. For the destruction of residual small amounts of the alloy, treatment with a 1:1 mixture of isopropanol and heptane over a safety tray is recommended [2]. The alloy usually ignites on exposure to air, with which it reacts in any case to form potassium dioxide (or superoxide), a very powerful oxidant [3]. Fires may be extinguished with dried sodium carbonate, calcium carbonate, sand or resin-coated sodium chloride. Graphite is not suitable as violent reaction with the superoxide is possible. Carbon dioxide or halocarbon extinguishers must not be used, to avoid explosions [4].

See Carbon dioxide, *also* Halocarbons, both below

See Potassium dioxide: Carbon

tert-Butanol

Renfrew, M. M., *J. Chem. Educ.*, 1983, **60**, A229

Contact of the alloy with *tert*-butanol caused ignition.

See reference 2, main entry above

Carbon dioxide

Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

Mixtures of the alloy and solid carbon dioxide are powerful explosives, some 40 times more sensitive to shock than mercury fulminate.

Carbon disulfide

See Carbon disulfide: Alkali metals

Fluoropolymers

1. *491M*, 1975, 394

2. Kimura, T. *et al.*, *J. Chem. Educ.*, 1973, **50**, A85

Poly-tetrafluoroethylene or -hexafluoropropylene sealing tapes burned vigorously in contact with the alloy in a helium atmosphere [1]. A teflon-coated magnetic stirrer bar used to stir the alloy under propane atmosphere ignited when the speed was increased and generated enough heat to melt the glass. Triboelectric initiation was postulated [2].

See other GLASS INCIDENTS

Halocarbons

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 550
2. *Inform. Exch. Bull.*, Lawrence Radiation Lab., 1961, **1**, 22
3. Schmidt, E. *et al.*, *Chem. Abs.*, 1975, **83**, 163034
4. *491M*, 1975, 394

The liquid alloy gives mixtures with halocarbons even more shock-sensitive than those with potassium. Highly chlorinated methane derivatives are more reactive than those of ethane, often exploding spontaneously after a delay [1]. Contact of 1,1,1-trichloroethane with a trapped alloy residue in a valve caused an explosion [2]. It is to be expected that chlorofluorocarbons will also form hazardous mixtures in view of their reactivity with barium. Precautionary measures for demonstrating the explosion of the alloy with chloroform are detailed [3]. Addition of 2 drops of 1,1,2-trichlorotrifluoroethane to the alloy caused a violent explosion [4].

See Fluoropolymers, above

See other METAL-HALOCARBON INCIDENTS

Metal halides

1. Rieke, R. D., *Accts. Chem. Res.*, 1977, **10**, 302
2. Ellern, 1968, 43

Use of the alloy to reduce metal halides in solvents to the finely divided and highly reactive metals is not recommended for cases where the halide is highly soluble in the solvent (e.g. zinc chloride or iron(III) chloride in THF). Explosive reaction may ensue [1]. The alloys explode violently in contact with silver halides.

See Potassium: Metal halides

Metal oxides

Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 551

Mixtures of the alloy with silver oxide or mercury oxide are shock-sensitive powerful explosives. The red form of mercury(I) oxide gives mixtures 40 times, and the yellow form 140 times as sensitive as mercury fulminate.

Nitrogen-containing explosives

See Potassium: Nitrogen-containing explosives

Water

Anon., *Loss Prev. Bull.*, 1977, (018), 18

The liquid alloy was used in an unlabelled Dreschel bottle to dry inert gas. A violent explosion occurred when the bottle was being cleaned, owing to contact of the alloy with water. Other desiccants are much safer (and would avoid the possibility of mistaking the alloy for mercury).

See other ALLOYS, PYROPHORIC MATERIALS

KMnO₄

4647. Potassium permanganate
[7722-64-7]

KMnO₄

KMnO₄

FPA H28, 1974; *HCS 1980*, 774

Acetic acid

MRH 2.34/84

or Acetic anhydride

MRH 2.55/86

von Schwartz, 1918, 34

Cooling is necessary to prevent possible explosion from contact of potassium permanganate (or the sodium or calcium salts) with acetic acid or acetic anhydride.

See Oxygenated organic compounds, below

Acetone, *tert*-Butylamine

MRH Acetone 2.76/8

1. Kornblum, N. *et al.*, *Org. Synth.*, 1973, Coll. Vol. 5, 847

2. Kornblum, N. *et al.*, *Org. Synth.*, 1963, **43**, 89

3. Haynes, R. K. *et al.*, *Lab. Equip. Digest*, 1974, **12**(6), 98; private comm., 1974

In the reprinted description [1] of a general method of oxidising *tert*-alkylamines to the corresponding nitroalkanes, a superscript reference indicating that *tert*-butylamine should be oxidised in water alone, rather than in acetone containing 20% of water, is omitted, although it was present in the original description [2]. This appears to be important, because running the reaction in 20% aqueous acetone led to a violent reaction with eruption of the flask contents. This was attributed to caking of the solid permanganate owing to inadequate agitation, and onset of an exothermic reaction between oxidant and solvent [3].

Alcohols, Nitric acid

Numjal, N. L. *et al.*, *Amer. Inst. Aero. Astronaut. J.*, 1972, **10**, 1345

Methanol, ethanol, isopropanol, pentanol or isopentanol do not ignite immediately upon mixing with red fuming nitric acid, but addition of potassium permanganate (20%) to the acid before mixing causes immediate ignition.

See Nitric acid: Alcohols

Aluminium carbide

Mellor, 1946, Vol. 5, 872

Incandescence on warming.

Ammonia, Sulfuric acid

Mellor, 1941, Vol. 1, 907

Ammonia is oxidised with incandescence in contact with the permanganic acid formed in the mixture.

Ammonium nitrate

Urbanski, 1965, Vol. 2, 491

A mixture of 0.5% of potassium permanganate with an ammonium nitrate explosive caused an explosion 7 h later. This was owing to formation and exothermic decomposition of ammonium permanganate, leading to ignition.

Ammonium perchlorate

See Ammonium perchlorate: Impurities

Antimony, or Arsenic

MRH 0.25/86, 0.25/91, resp.

Mellor, 1942, Vol. 12, 322

Antimony ignites on grinding in a mortar with the solid oxidant, while arsenic explodes.

Coal, Peroxomonosulfuric acid

Rawat, N. S., *Chem. & Ind.*, 1976, 743

In a new method for determination of sulfur in coal, the samples are oxidised with an aqueous mixture of permanganate and the peroxyacid. During the digestion, a reflux condenser is essential to prevent loss of water, which could lead to explosively violent oxidation.

Dichloromethylsilane

See Dichloromethylsilane: Oxidants

Dimethylformamide

Finlay, J. B., *Chem. Eng. News*, 1980, **58**(38), 65

Addition of potassium permanganate to dimethylformamide to give a 20% (approx. saturated) solution led to an explosion after 5 min. Subsequent tests on 1 g of oxidant with 5 g of solvent showed a rapid exotherm after 3–4 min, accompanied by popping noises from undissolved oxidant.

See Oxygenated organic compounds, below; also Dimethylformamide: Oxidants

Dimethyl sulfoxide

MRH 3.01/86

Kirk-Othmer (3rd ed.), 1982, Vol. 19, 494

The solvent ignites in contact with the solid oxidant.

See Oxygenated organic compounds, below

Ethanol, Sulfuric acid

Koch, K. R., *J. Chem. Educ.*, 1982, **58**, 973–974

In a demonstration of the powerful oxidant effect of manganese heptoxide on ethanol layered on top of sulfuric acid, it is essential to observe all the precautions given to prevent violent exothermic reactions.

Formaldehyde

1. Piefel, W. *et al.*, *Chem. Abs.*, 1977, **86**, 60508

2. Robinson, P. J., *Chem. & Ind.*, 1978, 723

3. Wainmann, H. E., *Chem. & Ind.*, 1978, 744

4. Craven, J. R., *Chem. Brit.*, 1979, **15**, 66

5. Young, J. A., *CHAS Notes*, 1988, **6**(3), 6

Formaldehyde gas for disinfection purposes may be released from the aqueous solution ('formalin') by treatment with potassium permanganate (the heat for evaporation arising from the redox reaction), but the quantities must be limited to avoid

KMnO₄

the risk of fire or explosion [1]. There is an account of an incident involving an extremely exothermic reaction overnight after addition of 0.1 l of formalin to 50 g of oxidant in a plastic beaker, which was melted [2]. Electrically heated evaporation of formalin is a safer fumigation technique [3]. The wisdom of using plastic containers for any unsupervised exothermic reaction is questioned [4]. An attempt to disinfect a large building and its hundreds of incubators all at once by running formalin from a 180 l drum onto several kg of potassium permanganate in a small waste container led, predictably, to a large fire which destroyed the building and its contents [5].

Glycerol

1. 'Leaflet No. 5', London, Inst. of Chem., April, 1940
2. Anon., *BCISC Quart. Safety Summ.*, 1974, **45**, 11–122
3. Bozzelli, J. W. *et al.*, *J. Chem. Educ.*, 1979, **56**, 675–676
4. Kauffman, G. B. *et al.*, *J. Chem. Educ.*, 1981, **58**, 802

Contact of glycerol with solid potassium permanganate caused a vigorous fire [1]. During the preparation of a solution for the decontamination of tetramethyllead spills, addition of the solid oxidant to a bucket contaminated with glycerol caused ignition to occur after a few s [2]. The trifunctionality and high viscosity of glycerol give high heat release and poor dissipation of reaction heat, respectively. The combination is a reliable method of igniting thermite mixture demonstrations [3], which should be conducted out of doors [4].

See Oxygenated organic compounds, below

Hydrochloric acid

MRH 0.75/63

1. Curry, J. C., *School Sci. Rev.*, 1965, **46**(160), 770
2. Adair, A., *Chem. & Ind.*, 1965, 1723
3. Jenkins, E. W., private comm., 1976
4. Ephraim, 1939, 162

During preparation of chlorine by addition of the conc. acid to solid permanganate, a sharp explosion occurred on one occasion [1]. Sulfuric acid was not used in error, nor was tube blockage involved [2]. A similar incident was reported later [3]. It appears remotely possible that permanganate may be able to oxidise chlorine to chlorine oxide [4], which as a dilute mixture with chlorine would be mildly explosive in the gas phase.

Hydrofluoric acid

Black, A. M. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1974, 977

During preparation of potassium hexafluoromanganate(IV), addition of the solid oxidant to exceptionally conc. hydrofluoric acid (60–90%, rather than 40% previously used) caused a violent exotherm with light emission.

Hydrogen peroxide

Anon., *J. Pharm. Chim.*, 1927, **6**, 410

Contact of hydrogen peroxide from a broken bottle with pervious packages of permanganate caused a violent reaction and fire.

See Hydrogen peroxide: Metals, etc. (reference 3)

Hydrogen trisulfide

Mellor, 1947, Vol. 10, 159

Contact with solid permanganate ignites the liquid sulfide.

Hydroxylamine

MRH 3.85/99+

See other REDOX REACTIONS

Non-metals

MRH values below reference

Mellor, 1942, Vol. 12, 319–323

MRH Carbon 2.59/7, phosphorus (yellow) 6.88/62, (red) 2.34/16, sulfur 2.80/12

A mixture of carbon and potassium permanganate is not friction-sensitive, but burns vigorously on heating. Mixtures with phosphorus or sulfur react explosively on grinding or heating, respectively.

Organic nitro compounds

Blinov, I., *J. Chem. Ind. USSR*, 1937, **14**, 1151–1153

Mixtures ignite easily on heating, shock, or contact with sulfuric acid.

Other reactants

Yoshida, 1980, 90

MRH values calculated for 26 combinations with oxidisable materials are given.

Oxygenated organic compounds

1. Rathsbury, H. *et al.*, *Chem. Ztg.*, 1941, **65**, 426–427

2. Gallais, 1957, 697

3. Partington, 1967, 830–831

Contact of several liquid organic compounds, or of aqueous solutions of solids, with the powdered oxidant leads to ignition. Presence of hydroxy and /or keto groups causes an extraordinary increase in sensitivity. Such compounds include ethylene glycol, propane-1,2-diol, erythritol, mannitol, triethanolamine, 3-chloropropane-1,2-diol; acetaldehyde, isobutyraldehyde, benzaldehyde; acetylacetone; esters of ethylene glycol, lactic acid, acetic acid, oxalic acid. The necessary presence of hydroxy or keto groups may be connected with solubility factors [1]. Mixtures of the solid oxidant with solid reducing sugars may react violently or explosively [2]. Powdered oxalic acid and the oxidant ignite soon after mixing [3]. *See* Ethylene glycol: Oxidants

Polypropylene

MCA Case History No. 1842

While using a screw conveyor to move the solid oxidant, ignition of a polypropylene tube in the feed system occurred. This could not be reproduced, even when likely contaminants were present.

Potassium chloride, Sulfuric acid

491M, 1975, 347

An attempt to prepare permanganyl chloride by cautiously adding conc. sulfuric acid to an intimate mixture of the salts at 0°C in clean glass apparatus caused a violent explosion.

Reducants

Interaction may be violent or explosive.

See Formaldehyde, above

Hydrazine: Oxidants

Hydroxylamine: Oxidants

Slag wool

Hallam, B. J. *et al.*, *School Sci. Rev.*, 1975, **56**(197), 820

Generation of oxygen by heating a layer of potassium permanganate retained in a test tube by a plug of 'Rocksil' slag wool led to minor explosions. The wool liberated an organic distillate on heating, and the explosions were attributed to combustion of the distillate in the liberated oxygen. Roasting the wool before using it with oxidants is recommended.

Sulfuric acid, Water

1. Archer, J. R., *Chem. Eng. News*, 1948, **26**, 205
2. Anon., *ABCM Quart. Safety Summ.*, 1946, **17**, 2
3. Stevens, G. C., private comm., 1971
4. Haight, G. P. *et al.*, *Chem. Eng. News*, 1980, **58**(13), 3
5. Mann, L. T., *Chem. Eng. News*, 1980, **58**(18), 2
6. Gulick, W. M., *Chem. Eng. News*, 1980, **58**(28), 5
7. Olley, R. H., *J. Polym. Sci. Polym. Phys. Edn.*, 1979, **176**, 627
8. Olley, R. H., *Chem. Brit.*, 1988, **24**, 31
9. Anon., *Polymer*, 1982, **23**, 1707

Addition of conc. sulfuric acid to the slightly damp permanganate caused an explosion. This was attributed to the formation of permanganic acid, dehydration to dimanganese heptoxide and explosion of the latter, caused by heat liberated from interaction of sulfuric acid and moisture [1]. A similar incident was reported previously when a solution of potassium permanganate in sulfuric acid, prepared as a cleaning agent, exploded violently [2]. There is, however a reputedly safe procedure for the final ultra-cleaning of glassware in cases where the adsorbed chromium film left by chromic acid cleaning would be intolerable. This involves treatment of the glassware with conc. sulfuric acid (10 ml) to which one or two small crystals (not more) of permanganate have been added. If the solution changes to a brown colour, it must immediately be discarded into excess water [3]. A lecture demonstration of the 'permanganate volcano' involving preparation of manganese heptoxide exploded with great violence. It was concluded that the demonstration was too dangerous to use [4]. Similar comments followed [5,6]. When preparing a 7% solution of potassium permanganate in conc. sulfuric acid as an etchant for polyalkenes, it is essential to use only small quantities and to stir the acid rapidly while adding the oxidant and until it is completely dissolved [7]. An attempt to do this with inadequate instructions (3 l of acid; no agitation) led to a violent explosion which demolished a fume cupboard [8]. A safer, more dilute etchant solution has been described [9].

Titanium

Mellor, 1941, Vol. 7, 20

A mixture of powdered metal and oxidant explodes on heating.



Trifluoroacetic anhydride

1. Perrin, D. D. *et al.*, *Purification of Laboratory Chemicals*, 448, Sydney, Pergamon, 2nd ed., 1980

2. Mook, R. A., *Chem. Eng. News*, 1987, **65**(13), 2

3. Author's comments

A bottle of the anhydride had been purified uneventfully by distillation from potassium permanganate according to a published procedure [1]. When a second batch was attempted, a brown colour was observed in the distillation flask and the distillate was green, so distillation was stopped. Several hours later the distillation flask exploded spontaneously and with great violence. This was attributed to the formation of dimanganese heptaoxide (which caused the brown colouration). Trifluoroacetic acid may also explode if the same purification procedure is followed [2]. It seems likely that the first bottle of anhydride had a low content of trifluoroacetic acid, so that little of the dimanganese heptaoxide would be produced by liberation of permanganic acid and its subsequent dehydration by the anhydride. If the second sample of anhydride had an appreciably higher acid content, a substantial amount of the explosive heptaoxide could have been produced during the several hours standing by the progressive reaction sequence below. [3]



3,4,4'-Trimethyldiphenyl sulfone

Adduci, J. M. *et al.*, *Macromol. Synth.*, 1982, **8**, 53–56

During oxidation of the sulfone to the tricarboxylic acid, addition of oxidant must be in small portions to avoid an exothermic vigorous reaction.

Wood

MRH Cellulose 2.68/15

1. Anon., *Chem. Ztg.*, 1927, **51**, 221

2. Dunn, B. W., *Amer. Rail Assoc., Bur Explos., Rept. No. 10*, 1917

Contact between the solid oxidant and wood, in presence of either moisture [1] or mechanical friction [2] may cause a fire.

See other METAL OXOMETALLATES, OXIDANTS

4648. Potassium thiazate

[73400-02-9]

KNOS

KN:S:O

Alone, or Water

Armitage, D. A. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1979, 1078–1079

It explodes upon ignition dry, and ignites in contact with water.

See other N-METAL DERIVATIVES, N-S COMPOUNDS

KNO₃

4649. Potassium nitrite
[7758-09-0]

KNO₂

KON:O

HCS 1980, 772

Ammonium salts

Mellor, 1941, Vol. 2, 702

Addition of ammonium sulfate to the fused nitrite causes effervescence and ignition.

See NITRITE SALTS OF NITROGENOUS BASES

Boron

Mellor, 1946, Vol. 5, 16

Addition of boron to the fused nitrite causes violent decomposition.

Potassium amide

Bergstrom, F. W., *Chem. Rev.*, 1933, **12**, 64

Heating a mixture of the solids under vacuum causes a vigorous explosion.

Potassium cyanide

See Sodium nitrite: Metal cyanides

See other METAL NITRITES, METAL OXONON-METALLATES, OXIDANTS

4650. Potassium nitrate
[7757-79-1]

KNO₃

KONO₂

FPA H74, 1978; HCS 1980, 771

Aluminium, Barium nitrate, Potassium perchlorate, Water

See Aluminium: Metal nitrates, etc.

1,3-Bis(trichloromethyl)benzene

See 1,3-Bis(trichloromethyl)benzene: Oxidants

Boron, 'Laminac', Trichloroethylene

MCA Case History No. 745

A 14 kg batch of the mixture, mainly of boron and potassium nitrate, with a minority of the synthetic resin adhesive and solvent, exploded 5 min after blending had started. Several possibilities of a frictional initiation source were considered.

Calcium silicide

1. Berger, F., *Compt. rend.*, 1920, **170**, 1492

2. Smolin, A. O., *Chem. Abs.*, 1977, **86**, 173758

A mixture of potassium nitrate (or sodium nitrate) and calcium silicide (60:40) is a readily ignited primer which burns at a very high temperature. It is capable of

initiating many high-temperature reactions [1]. The topic has been discussed more recently but no details were translated [2].

Calcium hydroxide, Polychlorinated phenols

See Calcium hydroxide: Polychlorinated phenols, etc.

Cellulose

Ishida, H. *et al.*, *Chem. Abs.*, 1988, **109**, 56797

Thermal reaction hazards of potassium nitrate–cellulose mixtures were evaluated by ARC. Stoichiometric mixtures (zero oxygen balance) showed the lowest ignition temperatures.

Chromium nitride

Partington, 1967, 744

The nitride deflagrates with the molten nitrate.

Honey, Sulfur

Winfield, J., *The Gunpowder Mills of Fernilee*, 20 (Private publ. by Author: 21 New Rd, Whaley Bridge, SK23 7JG, UK), 1996

An early Chinese manuscript warns that seething this mixture endangers the experimenter's beard. It is a probable precursor of gunpowder.

Lactose

See Lactose: Oxidants

Metals

Mellor, 1941, Vol. 7, 20, 116, 261; 1939, Vol. 9, 382

MRH Aluminium 7.15/35, iron 1.55/53, magnesium 7.57/42, sodium 3.10/58

Mixtures of potassium nitrate and powdered titanium, antimony or germanium explode on heating, and with zirconium at the fusion temperature of the mixture.

Metal sulfides

1. Mellor, 1939, Vol. 9, 270, 524

2. Mellor, 1941, Vol. 3, 745; Vol. 7, 91, 274

3. Mellor, 1943, Vol. 11, 647

4. Pascal, 1963, Vol. 8.3, 404

MRH Antimony trisulfide 2.30/37, titanium disulfide 3.42/26

Mixtures of potassium nitrate with antimony trisulfide [1], barium sulfide, calcium sulfide, germanium monosulfide or titanium disulfide all explode on heating [2]. The mixture with arsenic disulfide is detonable, and addition of sulfur gives a pyrotechnic composition [2]. Mixtures with molybdenum disulfide are also detonable [3]. Interaction with sulfides in molten mixtures is violent [4].

Non-metals

1. *MCA Case History No. 1334*

2. Mellor, 1946, Vol. 5, 16

3. Brede, U., *Chem. Abs.*, 1981, **94**, 86603

KNO₃

4. Leleu, *Cahiers*, 1980, (99), 278
5. Mellor, 1941, Vol. 2, 820, 825; 1963, Vol. 2, Suppl. 2.2, 1939
6. Mellor, 1940, Vol. 8, 788
7. Mellor, 1939, Vol. 9, 35

MRH Carbon 3.26/13, Phosphorus (yellow) 3.14/27, sulfur 2.97/21

A pyrotechnic blend of a finely divided mixture with boron ignited and exploded when the aluminium container was dropped [1]; (the aluminium container also may have been involved). Boron is not attacked at below 400°C, but is at fusion temperature or at lower temperatures if decomposition products (nitrites) are present [2]. The mixture has also been evaluated as a propellant [3]. Contact of powdered carbon with the nitrate at 290°C causes vigorous combustion [4] and a mixture explodes on heating. Gunpowder is the oldest known explosive and contains potassium nitrate, charcoal and sulfur, the latter to reduce ignition temperature and to increase the speed of combustion [5]. Mixtures of white phosphorus and potassium nitrate explode on percussion, and a mixture with red phosphorus reacts vigorously on heating [6]. Mixtures of potassium nitrate with arsenic explode vigorously on ignition [7].

Organic materials

1. Smith, A. J., *Quart. Natl. Fire Prot. Assoc.*, 1930, **24**, 39–44
2. Unpublished information, 1979

MRH Aniline 3.51/13, acetone 3.47/15, ethanol 3.31/16, toluene 3.56/12

Potassium nitrate in cloth sacks stowed next to baled peat moss became involved in a ship fire and caused rapid flame spread and explosions [1]. Heat transfer salt from a new supplier was added to a pilot plant reactor salt bath. Some 12 h after start of heating to melt the bath contents a muffled explosion occurred, attributed to presence of organic impurities in the new salt [2].

See Cellulose, above.

See Sodium nitrate: Jute, etc.

Other reactants

Yoshida, 1980, 192

MRH values calculated for 19 combinations with oxidisable materials are given.

Phosphides

Mellor, 1940, Vol. 8, 839, 845

Boron phosphide ignites in molten nitrates; mixtures of the nitrate with copper(II) phosphide explode on heating, and that with copper monophosphide explodes on impact.

Reducants

Mellor, 1941, Vol. 2, 820

Mixtures of potassium nitrate with sodium phosphinate and sodium thiosulfate are explosive, the former being rather powerful.

See other REDOX REACTIONS

Sodium acetate MRH 2.55/34
Pieters, 1957, 30
Mixtures may be explosive.

Thorium dicarbide
See Thorium dicarbide: Non-metals, etc.
See other METAL NITRATES, OXIDANTS

4651. Potassium nitridoosmate **KNO₃Os**
[21774-03-8] **K[NOsO₃]**

Clifford, A. F. *et al.*, *Inorg. Synth.*, 1960, **6**, 205
Heating above 180°C at ambient pressure causes it to explode.
See related NITRIDES

4652. Potassium azide **KN₃**
[20762-60-1] **KN₃**

Mellor, 1940, Vol. 8, 347
It is insensitive to shock; on heating progressively it melts, then decomposes evolving nitrogen, and the residue ignites with a feeble explosion.

Carbon disulfide
See Carbon disulfide: Metal azides

Manganese dioxide
See Manganese(IV)oxide: Potassium azide

Sulfur dioxide
Mellor, 1940, Vol. 8, 347
Potassium azide explodes at 120°C when heated in liquid sulfur dioxide.
See other METAL AZIDES

4653. Potassium azidosulfate **KN₃O₃S**
[] **KOSO₂N₃**



Mellor, 1940, Vol. 8, 314
It explodes on heating.
See other ACYL AZIDES

4654. Potassium dinitramide
[140456-79-7]



See Dinitramine
See other N-NITRO COMPOUNDS

4655. Potassium azidodisulfate
[67880-14-2]



Alone, or Water

Mellor, 1967, Vol. 8, Suppl. 2.2, 36

On keeping or slow heating, the salt produces explosive disulfuryl azide, and the salt reacts explosively with water.

See Sulfuryl diazide See other ACYL AZIDES

4656. Potassium dioxide (Potassium superoxide)
[12030-88-5]



1. *Alkali Metals*, 1957, 174
2. Gilbert, H. N., *Chem. Eng. News*, 1948, **26**, 2604
3. Madaus, private comm., 1976
4. Kirk-Othmer, 1968, Vol. 16, 365–366
5. Commander, J. C., *ERDA Rep. ANCR 1217; TID-4500*, Springfield (Va.), USNTIS, 1975
6. Sloan, S. A., *Proc. Int. Conf. Liquid Met. Technol.*, Champion (Pa.), Int. At. En. Auth., 1976

The earlier references, which state that this powerful oxidant is stable when pure, but explosive when formed as a layer on metallic potassium [1,2], are not wholly correct [3], because the superoxide is manufactured uneventfully by spraying the molten metal into air to effect oxidation [4]. Previous incidents appear to have involved the explosive oxidation of unsuspected traces of mineral oil or solvents [3]. However, mixtures of the superoxide with liquid or solid potassium–sodium alloys will ignite spontaneously after an induction period of 18 min, but combustion while violent is not explosive [3]. The additional presence of water (which reduces the induction period) or hydrocarbon contaminant did produce explosion hazards under various circumstances [5]. Contact of liquid potassium with the superoxide gives no obvious reaction below 117°C and a controlled reaction between 117 and 177°C, but an explosive reaction occurs above 177°C. Heating at 100°C/min from 77° caused explosion at 208°C [6].

See other INDUCTION PERIOD INCIDENTS

See Potassium:- *Slow oxidation*

See Dipotassium μ cyclooctatetraene

2-Aminophenol, Tetrahydrofuran

1. Crank, G. *et al.*, *Aust. J. Chem.*, 1984, **37**, 845–855

2. Crank, G. *et al.*, *Chem. Eng. News*, 1988, **68**(35), 2

2-Aminophenol was being oxidised in THF solution at 65°C using a larger than normal proportion of potassium dioxide. When stirring was stopped after 6 h, a violent explosion occurred. This was attributed to formation of tetrahydrofuranyl hydroperoxide by the excess dioxide. THF was described as an unsafe solvent for superoxide reactions [1]. A later attempt at the same reaction in toluene also led to explosion, now blamed on the substrate [2].

See Hydrocarbons, below.

Carbon

Mellor, D. P., *Chem. & Ind.*, 1965, 723

Residues from extinguishing small sodium–potassium alloy fires with graphite were accumulated in an airtight drum. Later, burning alloy fell into the drum and caused a violent explosion. This was attributed to formation of potassium superoxide during storage, and explosive reaction of the latter with graphite, initiated by the burning alloy. Graphite is not a suitable extinguishant for fires involving potassium or its alloys.

Diselenium dichloride

Mellor, 1947, Vol. 10, 897

Interaction is very violent.

Ethanol

Health & Safety Inf., 251, Washington, USAEC, 1967

Disposal of a piece of potassium–sodium alloy under argon in a glove box by addition of ethanol caused violent gas evolution which burnt a glove and produced a flame. A piece of highly oxidised potassium exploded when dropped into ethanol. Both incidents were attributed to violent interaction of potassium superoxide and ethanol.

Hydrocarbons

Health & Safety Inf., 251, Washington, USAEC, 1967

Residues of potassium–sodium alloy in metal containers were covered with oil prior to later disposal. When a lid was removed later, a violent explosion occurred. This was attributed to frictional initiation of the mixture of potassium superoxide (formed on long standing of the alloy) and oil.

Metals

Mellor, 1941, Vol. 2, 493

Oxidation of arsenic, antimony, copper, potassium, tin or zinc proceeds with incandescence.

See reference 6 above

Organic compounds

Frimer, A. A. *et al.*, *Photochem. Photobiol.*, 1978, **28**(4–5. Singlet oxygen), 718

Potassium superoxide must not be added to neat oxidation substrates, or ignition may occur, and weighing it out on filter paper is also hazardous.

Potassium–sodium alloy

Sloan, S. A. *et al.*, *Microscope*, 1977, **25**, 237–243

A runaway self heating reaction occurs 3.8 s after mixing at 201°C or above, and at that temperature an exotherm to 304°C at a rate above 2000°/s was observed, but no pressure effects were seen.

See related METAL PEROXIDES *See other* METAL OXIDES, OXIDANTS

4657. Potassium trioxide (Potassium ozonide)

[12030-89-6]



KO₃

See Caesium trioxide, *also* Fluorine: Potassium hydroxide

See related METAL PEROXIDES *See other* METAL OXIDES

4658. Potassium silicide

[16789-24-5]



KSi


Bailar, 1973, Vol. 1, 1357

It ignites in air.

See other METAL NON-METALLIDES

4659. Potassium diperoxomolybdate

[58412-05-8]



K₂MoO₆


Mellor, 1943, Vol. 11, 607

It explodes on grinding.

See other PEROXOACID SALTS

4660. Potassium tetraperoxomolybdate

[56094-15-6]




K₂MoO₈

Jahr, K. F., *FIAT Rev. Ger. Sci: Inorg. Chem., Pt. III*, 1948. 170

It is explosive.

See other PEROXOACID SALTS

4661. Potassium nitrosodisulfate
[14293-70-0]





1. Zimmer, H. *et al.*, *Chem. Rev.*, 1971, **71**, 230, 243
2. Wehrli, P. A. *et al.*, *Org. Synth.*, 1972, **52**, 86

The observed instability of the solid radical on storage, ranging from slow decomposition to violent explosion (probably depending on the degree of confinement), depends largely upon the degree of contamination by nitrite ion. Storage conditions to enhance stability are detailed [1]. Synthetic use of solutions of the salt has been preferred, and the storage of any isolated salt as an alkaline slurry is recommended [2].

See related NITROSO COMPOUNDS

4662. Potassium N-nitrosohydroxylamine-N-sulfonate ('Potassium dinitrososulfite')
[26241-10-1]



Weitz, E. *et al.*, *Ber.*, 1933, **66**, 1718

The salt (formerly named a dinitrososulfite) explodes on heating.

See other N-O COMPOUNDS, N-S COMPOUNDS

4663. Potassium azodisulfonate
[]



Alone, or Water

Konrad, E. *et al.*, *Ber.*, 1926, **59**, 135

There is a possibility of explosion during vacuum desiccation of the salt, which evolves nitrogen violently in contact with water.

See related AZO COMPOUNDS *See other* N-S COMPOUNDS

4664. Potassium sulfurdiimidate
[[79796-14-8] (ion)]



Alone, or Solvents, or Water

1. Goehring, 1957, 39, 6


2. Hornemann, K. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1982, **21**, 633

It ignites in air, and reacts violently and may explode with traces of water [1]. It is stable up to 180°C, but reacts explosively and ignites with water, methanol or chloromethane, methylene chloride or carbon tetrachloride [2].

See other N-METAL DERIVATIVES, N-S COMPOUNDS

4665. Potassium hexaazidoplatinate(IV)

[41909-62-0] (ion)



Mellor, 1940, Vol. 8, 355

Evaporation of a solution of hexachloroplatinic acid with a deficiency of potassium azide, or with an equivalence of ammonium azide gives explosive residues. Evaporation of a solution of the acid with an equivalence (8 mol) of potassium azide leads to explosion of the conc. solution of the title compound.

See other PLATINUM COMPOUNDS *See related* METAL AZIDES

4666. Potassium peroxide

[17014-71-0]



Other reactants

Yoshida, 1980, 83

MRH values calculated for 15 combinations with oxidisable materials are given.

Water


Pascal, 1963, Vol. 2.2, 59

Interaction is violent or explosive.

See other METAL PEROXIDES, OXIDANTS

4667. Potassium diperoxoorthovanadate

[]



1. Mellor, 1939, Vol. 9, 795


2. Emeléus, 1960, 425

It explodes on heating [1,2].

See other PEROXOACID SALTS

4668. Potassium peroxodisulfate

[7727-21-1]



FPA H106, 1981; *HCS 1980*, 775

Energy of decomposition (in range 190–840°C) measured as 0.29 kJ/g.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)

Alkali, Hydrazine salts

Mellor, 1947, Vol. 10, 466

Addition of alkali to the mixed salts liberates hydrazine which is vigorously oxidised to nitrogen gas.

Potassium hydroxide

Anon., *BCISC Quart. Safety Summ.*, 1965, **36**, 41

Surface contamination of 2 kg of the dry salt with as little as 2 flakes of moist potassium hydroxide caused a vigorous self-sustaining fire, which was extinguished with water, but not by carbon dioxide or dry powder extinguishers.

Water

Castrantas, 1965

The salt rapidly liberates oxygen above 100°C when dry, but at only 50°C when wet.

See other PEROXOACID SALTS, OXIDANTS

4669. Potassium tetraperoxotungstate

[37346-96-6]



$\text{K}_2\text{O}_8\text{W}$

Mellor, 1943, Vol. 11, 836

It explodes on friction or rapid heating to 80°C.

See other PEROXOACID SALTS

4670. Potassium sulfide

[1312-73-8]



K_2S

HCS 1980, 387

Merck Index, 1983, p1104

The anhydrous sulfide is unstable, and may explode on percussion or rapid heating.

Chloroform: Ethanol (or Methanol)

See Potassium dithioformate

Nitrogen oxide

See Nitrogen oxide: Potassium sulfide

Other reactants

Yoshida, 1980, 83

MRH values calculated for 13 combinations with various reagents are given.

See other METAL SULFIDES

4671. Potassium nitride
[29285-24-3]



Mellor, 1940, Vol. 8, 99
It usually ignites in air.

Phosphorus



See Phosphorus: Potassium nitride

Sulfur

See Sulfur: Potassium nitride

See other NITRIDES

4672. Potassium 2,4,6-tris(dioxoselena)hexahydrotriazine-1,3,5-triide ('Potassium triselenimidate')
[51593-99-8]



Explosive.

See Selenium difluoride dioxide: Ammonia

See other N-METAL DERIVATIVES

4673. Potassium antimonide
[16823-94-2]



1. Mellor, 1939, Vol. 9, 403

2. Rüst, 1948, 342

It usually ignites when broken in air [1] and explodes on exposure to moisture (in breath) [2].

See other ALLOYS, PYROPHORIC MATERIALS

4674. Potassium hexaoxoxenonate-xenon trioxide
[12273-50-6]



1. Grosse, A. V., *Chem. Eng. News*, 1964, **42**(29), 42

2. Spittler, T. M. *et al.*, *J. Amer. Chem. Soc.*, 1966, **88**, 2942

3. Appelman, E. H. *et al.*, *J. Amer. Chem. Soc.*, 1964, **86**, 2141

The complex salt is very sensitive to mechanical shock and explodes violently [1,2], even when wet [3].

See other XENON COMPOUNDS

See related NON-METAL OXIDES

- 4675. Lithium tripotassium tetrasilicide**
 [102210-64-0] **K₃LiSi₄**

$$\text{LiK}_3\text{Si}_4$$
 von Schering, H. G. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1986, **25**, 566
 It ignites in air.
See other METAL NON-METALLIDES
- 4676. Lithium heptapotassium di(tetrasilicide)**
 [102210-65-1] **K₇LiSi₈**

$$\text{LiK}_7(\text{Si}_4)_2$$
 von Schering, H. G. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1986, **25**, 566
 It ignites in air.
See other METAL NON-METALLIDES
- 4677. Lanthanum**
 [7439-91-0] **La**

$$\text{La}$$
 Nitric acid
 Mellor, 1946, Vol. 5, 603
 Oxidation is violent.
 Phosphorus
See Phosphorus: Metals
See other METALS
- 4678. Lanthanum–nickel alloy**
 [12196-72-4] **LaNi₅**

$$\text{LaNi}_5$$
 Hydrogen
See Lanthanum pentanickel hexahydride
 Hydrogen, Poly(tetrafluoroethylene)
See LANTHANIDE–TRANSITION METAL ALLOY HYDRIDES
See Poly(tetrafluoroethylene): Metal hydrides
See other ALLOYS, HYDROGENATION CATALYSTS
- 4679. Lanthanum oxide**
 [1312-81-8] **La₂O₃**

$$\text{La}_2\text{O}_3$$
 Chlorine trifluoride
See Chlorine trifluoride: Metals, etc.
- 1748

Water

Sidgwick, 1950, 441

Interaction is vigorously exothermic, accompanied by hissing, as for calcium oxide.

See Calcium oxide: Water

See other METAL OXIDES

4680. Lithium

[7439-93-2]

Li

Li

NSC 566, 1978; HCS 1980, 592

1. Mellor, 1941, Vol. 2, 468; 1961, Vol. 2, Suppl. 2.1, 71
2. *Haz. Chem. Data*, 1975, 187
3. Bullock, A., *Chem. Brit.*, 1975, **11**, 115
4. Bullock, A., *School Sci. Rev.*, 1975, **57**(119), 311–314; private comm., 1975
5. Lloyd, W. H., and Emley, E. F., *Chem. Brit.*, 1975, **11**, 334
6. Ireland, R. E. *et al.*, *Org. Synth.*, 1977, **56**, 47
7. Fritsch, *Proc. 1st Int. Symp. Prev. Occup. Risks Chem. Ind.*, 208–211, Heidelberg, ISSA, 1970

MRH Oxygen 19.95/54

Finely divided metal may ignite in air at ambient temperature and massive metal above the m.p., 180°C, especially if oxide or nitride is present. Since lithium will burn in oxygen, nitrogen or carbon dioxide, and when alight it will remove the combined oxygen in sand, sodium carbonate, etc., it is difficult to extinguish once alight [1]. (*Note the very high MRH value above.*) Molten lithium is extremely reactive and will attack concrete and refractory materials. Use of normal fire extinguishers (containing water, foam, carbon dioxide, halocarbons, dry powders) will either accelerate combustion or cause explosion. Powdered graphite, lithium chloride, potassium chloride or zirconium silicate are suitable extinguishants [2]. A well-tried and usually uneventful demonstration of atmospheric oxidation of molten lithium led to an explosion [3]. A high degree of correlation of incidence of explosions with high atmospheric humidity was demonstrated, with the intensity of explosion apparently directly related to the purity of the sample of metal [4]. Other possible factors were also identified [5]. While cleaning lithium wire by washing with hexane, the wire must be dried carefully with a paper towel. Too-vigorous rubbing will cause a fire [6].

Lithium blocks containing traces of nitride but which had been supplied under argon in sealed tins were cut into 1 cm strips and stored under air in closed tins overnight until used. Lithium containing some nitride reacts slowly with nitrogen at ambient temperature to form more nitride, which autocatalyses the reaction which progressively accelerates and becomes exothermic. The strips of lithium reacted with the nitrogen of the air in the closed tins, causing a partial vacuum and an oxygen-enriched atmosphere. When the tins collapsed, the impact and/or compression of the oxygen-enriched atmosphere caused ignition and fierce burning of the lithium, which was very difficult to extinguish inside the crushed tins [7]. See Metal chlorides, also Water, both below

Acetonitrile, Sulfur dioxide

Ebner, W. B. *et al.*, *Proc. 8th Power Sources Symp.*, 119–124, 1982

An ARC study of the thermal and pressure behaviour of actual electric batteries under various atypical conditions showed the major contributions to the exothermic behaviour as the reactions between lithium and acetonitrile, lithium and sulfur and the decomposition of lithium dithionite. The first reaction can generate enough heat to trigger other exothermic reactions. The hazards associated with the various parameters are quantified.

See Sulfur dioxide, below

Bromine pentafluoride

See Bromine pentafluoride: Acids, etc.

Bromobenzene

Koch-Light Labs., Ltd., private comm., 1976

In a modified preparation of phenyllithium, bromobenzene was added to finely powdered lithium (rather than coarse particles) in ether. The reaction appeared to be proceeding normally, but after about 30 min it became very vigorous and accelerated to explosion. It was thought that the powdered metal may have been partially coated with oxide or nitride which abraded during stirring, exposing a lot of fresh metal surface on the powdered metal.

See Sodium: Halocarbons (reference 8) *See also* Halocarbons, below

Carbon, Lithium tetrachloroaluminate, Sulfinyl chloride

Kilroy, W. P. *et al.*, *J. Electrochem. Soc.*, 1981, **128**, 934–935

In electric battery systems, lithium is inert to the electrolyte components in absence of carbon, but in presence of over 10% of carbon (pre-mixed by grinding with the metal), contact with the electrolyte mixture leads to ignition or explosion.

See Sulfinyl chloride, below

Carbon, Sulfinyl chloride

James, S. D. *et al.*, *J. Electrochem. Soc.*, 1983, **130**, 2037–2040

Pregrinding lithium with carbon leads to ignition on contact with sulfinyl chloride in electric battery systems. The effect of moisture and purity of the carbon on reactivity was studied.

Carbon disulfide

See Carbon disulfide: Alkali metals

Chlorine tri- or penta-fluorides

See Chlorine trifluoride: Metals, etc.

1-Chloro-3-dimethylaminopropane

1. Eisenbach, A. *et al.*, *Euro. Polym. J.*, 1975, **11**, 699

2. Service, D., *Chem. Brit.*, 1987, **23**, 27

3. Bell, J. A. *et al.*, *J. Org. Chem.*, 1959, **24**, 2036

4. Kamienskis, C. W., *Chem. Brit.*, 1987, **23**, 531–532

Li

The preparation of 1-lithio-3-dimethylaminopropane according to a published procedure [1] involved reaction of the chloro compound with a mineral oil dispersion of lithium (30%) in hexane at 0°C. It had previously been found that lithium with a sodium content of some 0.3% reacted slowly, even at temperatures above 15°C. When a new batch of lithium (later found to contain 1.9% of sodium) was used, a vigorous reaction which set in at between 0° and -35°C led to ignition of the reaction mixture [2]. Although it was known that the presence of sodium increased the reactivity of lithium towards organohalides up to a sodium content of 2% [3], it had not been appreciated that such wide variations in reactivity were likely, or that a hazardous situation could develop [1,2]. However, a more likely cause proposed for the runaway exothermic reaction was the fact that all the reagents were mixed at 0°C, rather than the more usual course of adding the halide slowly to the lithium dispersion in a hydrocarbon solvent lower-boiling than hexane, so as to maintain gentle reflux at 35–50°C. Some general precautions for this type of reaction are given [4].

See Halocarbons, below

1,2-Diaminoethane, Tetralin

Reggel, A. *et al.*, *J. Org. Chem.*, 1957, **22**, 892

Reduction of tetralin to octalin with lithium and ethylenediamine proceeds slowly, but if heated to 85°C it becomes violent, with rapid evolution of hydrogen.

See other GAS EVOLUTION INCIDENTS

Diazomethane

See Diazomethane: Alkali metals

Diborane

MRH 1.13/99+

See Diborane: Metals

Ethylene

Pascal, 1966, Vol. 2.1, 38

Passage of the gas over heated lithium causes the latter to incandesce, producing a mixture of lithium hydride and lithium acetylide.

See Metal chlorides, etc., below

Halocarbons

MRH Bromoform 3.77/92, carbon tetrabromide 3.89/92,
carbon tetrachloride 8.24/85, carbon tetraiodide 1.42/95,
chloroform 7.78/85, fluorotrichloromethane 9.29/83,
tetrachloroethylene 8.16/86

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549
2. *Pot. Incid. Rep.*, 1968, **39**, ASES, Washington
3. Pittwell, L. R., *J. R. Inst. Chem.*, 1959, **80**, 552
4. BDH Catalogue Safety Note, 969DD/14.0/0773, 1973
5. *491M*, 1975, 230

Mixtures of lithium shavings and several halocarbon derivatives are impact-sensitive and will explode, sometimes violently [1,2]. Such materials include:

bromoform, carbon tetrabromide, carbon tetrachloride, carbon tetraiodide, chloroform, dichloromethane, diiodomethane, fluorotrichloromethane, tetrachloroethylene, trichloroethylene and 1,1,2-trichlorotrifluoroethane. In an operational incident, shearing samples off a lithium billet immersed in carbon tetrachloride caused an explosion and continuing combustion of the immersed metal [3]. Lithium which had been washed in carbon tetrachloride to remove traces of oil exploded when cut with a knife. Hexane is recommended as a suitable washing solvent [4]. A few drops of carbon tetrachloride on burning lithium was without effect, but a 25 cc portion caused a violent explosion [5].

See 1-Chloro-3-dimethylaminopropane, above

Poly(1,1-difluoroethylene-hexafluoropropylene), below

See entry HALOCARBONS: metals

Halogens

MRH Chlorine 9.62/83

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

2. Mellor, 1961, Vol. 2, Suppl. 2.1, 82

Mixtures of lithium and bromine are unreactive unless subject to heavy impact, when explosion occurs [1]. Lithium and iodine react above 200°C with a large exotherm [2].

Hydrogen

See Hydrogen: Metals

Mercury

1. Smith, G. McP. *et al.*, *J. Amer. Chem. Soc.*, 1909, **31**, 799

2. Alexander, J. *et al.*, *J. Chem. Educ.*, 1970, **47**, 277

Interaction to form lithium amalgam is violently exothermic and may be explosive if large pieces of lithium are used [1]. An improved technique, using *p*-cymene as inerting diluent, is described in the later reference [2].

Metal chlorides, Nitrogen

Anon., *BCISC Quart. Safety Summ.*, 1969, **40**, 16

Accidental contamination of lithium strip with anhydrous chromium trichloride or zirconium tetrachloride caused it to ignite and burn vigorously in the nitrogen atmosphere of a glove box.

Metal oxides and chalcogenides

MRH Chromium trioxide 7.20/83,
sodium peroxide 13.64/71

1. Mellor, 1967, Vol. 2, Suppl. 2.1, 81–82

2. *Alkali Metals*, 1957, 11

Lithium is used to reduce metallic oxides in metallurgical operations, and the reactions, after initiation at moderate temperatures, are violently exothermic and rapid. Chromium(III) oxide reacts at 185°C, reaching 965°; similarly molybdenum trioxide (180 to 1400°), niobium pentoxide (320 to 490°), titanium dioxide (200–400 to 1400°), tungsten trioxide (200 to 1030°), vanadium pentoxide (394 to 768°); also iron(II) sulfide (260 to 945°), and manganese telluride (230 to 600°C)

Li

[1]. Residual mixtures from lithium production cells containing lithium and rust sometimes ignite when left as thin layers exposed to air [2].

Metals

1. Bailar, 1973, Vol. 1, 343–344

2. Popov, E. I. *et al.*, *Chem. Abs.*, 1989, **110**, 100000

Formation of various intermetallic compounds of lithium by melting with aluminium, bismuth, calcium, lead, mercury, silicon, strontium, thallium or tin may be very vigorous and dangerous to effect [1]. Ignition and combustion hazards of alloys of lithium with aluminium or magnesium have been studied, the latter being more reactive than the former. Use of nitrogen as a protective medium for alloy powders is ineffective, mixed nitrides being formed [2].

See Mercury, above; Platinum, below

Nitric acid

MRH 15.27/60

Accident & Fire Prev. Info., March, 1974, Washington, USAEC

Lithium ignited on contact with nitric acid and the reaction became violent, ejecting burning lithium.

Nitryl fluoride

See Nitryl fluoride: Metals

Non-metal oxides

Mellor, 1961, Vol. 2, Suppl. 2.1, 74, 84, 88

Although carbon dioxide reacts slowly with lithium at ambient temperature, the molten metal will burn vigorously in the gas, which cannot be used as an extinguisher on lithium fires. Carbon monoxide reacts in liquid ammonia to give the carbonyl which reacts explosively with water or air. Lithium rapidly attacks silica or glass at 250°C.

See Sulfur dioxide, below

See other GLASS INCIDENTS

Other reactants

Yoshida, 1980, 385–386

MRH values calculated for 24 combinations with various reducible materials are given.

Platinum

See Platinum: Lithium

Poly(1,1-difluoroethylene–hexafluoropropylene) (Viton)

Markowitz, M. M. *et al.*, *Chem. Eng. News*, 1961, **39**(32), 4

Mixtures of finely divided metal and shredded polymer ignited in air on contact with water, or on heating to 369°C, or at 354° under argon.

See Halocarbons, above

Sodium carbonate
or Sodium chloride

MRH Sodium carbonate 4.27/73

Mellor, 1961, Vol. 2, Suppl. 2.1, 25

Sodium carbonate and sodium chloride are unsuitable to use as extinguishers for lithium fires, since burning lithium will liberate the more reactive sodium in contact with them.

Sulfur

Mellor, 1961, Vol. 2, Suppl. 2.1, 75

Interaction when either is molten is very violent and, even in presence of inert diluent, the reaction begins explosively. Reaction of sulfur with lithium dissolved in liquid ammonia at -33°C is also very vigorous.

Sulfinyl chloride

1. Abraham, K. M. *et al.*, *Proc. 28th Power Sources Symp.*, 1978, 255–257

2. Subbarao, S. *et al.*, *Proc. Electrochem. Soc.*, 1988, **88-6**, 187–200

3. Ducatman, A. M. *et al.*, *J. Occup. Med.*, 1988, **30**, 309–311

4. Butakov, A. A. *et al.*, *J. Power Sources*, 1992, **39**(3), 375

An experimental investigation of explosion hazards in lithium–sulfinyl chloride cells on forced discharge showed cathode limited cells are safe, but anode limited cells may explode without warning signs [1]. Extended reversal at -40°C caused explosion on warming to ambient temperature, owing to thermal runaway caused by accelerated corrosion of lithium [2]. The violent explosion of a large prismatic cell of a battery is described [3]. Another study of explosion mechanisms in lithium/thionyl chloride batteries is reported [4]

See Carbon, etc., above

Sulfur dioxide

1. Dey, A. N. *et al.*, *Chem. Abs.*, 1980, **93**, 10716, 10717

2. Crofts, C. C., *Chem. Abs.*, 1983, **99**, 56425

3. Subbarao, S. *et al.*, *J. Power Sources*, 1987, **21**(3–4), 221–237

4. Donaldson, G. J., *et al.*, *Power Sources*, 1991, **13**, 363

Two reports cover safety studies on lithium–sulfur dioxide batteries [1]. The cause of violent venting of discharged lithium–sulfur dioxide cells was ascribed to corrosion in a glass to metal seal and formation of lithium–aluminium alloys and other cathode reaction product(s) which are both shock-sensitive [2] The pyrophoric charged anodes of lithium–sulfur dioxide batteries are covered with smooth crystalline platelets, but partially discharged anodes are covered with a rough, non-adherent layer of lithium dithionite. The explosions which may occur during charging are attributed to thermal runaway reactions of lithium and sulfur dioxide to form lithium dithionite, $\text{LiOSO}\cdot\text{SO}\cdot\text{OLi}$ [3]. A study of battery explosions consequent upon anode breakage in lithium–sulfur dioxide cells is reported [4].

See Acetonitrile, etc., also Non-metal oxides, both above

Trifluoromethyl hypofluorite

See Trifluoromethyl hypofluorite: Lithium

Water

1. Mellor, 1961, Vol. 2, Suppl. 2.1, 72
2. Bailar, 1973, Vol. 1, 337
3. Markowitz, M. M., *J. Chem. Educ.*, 1963, **40**, 633–636

Reaction with cold water is of moderate vigour, but violent with hot water, and the liberated hydrogen may ignite [1]. The powdered metal reacts explosively with water [2]. The reactivity of lithium and other alkali metals with various forms of water has been discussed in detail. Prolonged contact with steam forms a thermally insulating layer which promotes overheating of the metal and may lead to a subsequent explosion as the insulating layer breaks up [3].

See ALKALI METALS

See other METALS

4681. Lithium–magnesium alloy

[12384-02-0]

Li–Mg

Li–Mg

Danyshova, T. A. *et al.*, *Chem. Abs.*, 1982, **97**, 167510

Fire hazard (title only translated).

See other ALLOYS

4682. Lithium–tin alloys

[12359-06-7]

Li–Sn

Li–Sn

Alkali metals, 1957, 13

Lithium–tin alloys are pyrophoric.

See other ALLOYS, PYROPHORIC MATERIALS

4683. Lithium sodium nitroxylate

[]

Li[O:NONa]

LiNNaO₂

Mellor, 1961, Vol. 2, Suppl. 2.1, 78

It decomposes violently at 130°C.

See other N–O COMPOUNDS

4684. Lithium nitrate

[7790-69-4]

LiONO₂

LiNO₃

Ethanol

See VAPOUR EXPLOSIONS (reference 3)

Other reactants

Yoshida, 1980, 208

MRH values calculated for 15 combinations with oxidisable materials are given.

Propene, Sulfur dioxide

See Propene: Lithium nitrate, etc.

See other METAL NITRATES

4685. Lithium azide

[19597-69-4]



Mellor, 1940, Vol. 8, 345

Insensitive to shock, the moist or dry salt decomposes explosively at 115–298°C, depending on the rate of heating.

Alkyl nitrates, Dimethylformamide

Polansky, O. E., *Angew. Chem. (Intern. Ed.)*, 1971, **10**, 412

Although these reaction mixtures are stable at 25°C during preparation of *tert*-alkyl azides, above 200°C the mixtures are shock-sensitive and highly explosive.

See other METAL AZIDES

4686. Lithium oxide

[12057-24-8]



Metal halide hydrates


Rowley, A. T. *et al.*, *Inorg. Chem. Acta*, 1993, **211**(1), 77

Preparation of metal oxides by fusing metal halides with lithium oxide in a sealed tube leads to explosions if halide hydrates are employed, particularly lanthanide trihalide hydrates. The preparation succeeds with anhydrous halides. This will be purely a question of vapour pressure above an exothermic reaction; the question is whether the vapour is water, or metal halide, and the reaction oxide formation, or hydration of lithium oxide. Like other alkali metal oxides, hydration is extremely energetic.

See Diprotium monoxide

4687. Lithium dithionite

[59744-77-3]



See Lithium: Sulfur dioxide (reference 3)

See other METAL OXONON-METALLATES, REDUCANTS

4688. Lithium nitride
[26134-62-3]

Li₃N

Li₃N

Air, or Water

1. 'Data Sheet TD 121', Exton, Foote Mineral Co., 1965
2. Schönher, E. *et al.*, *Inorg. Synth.*, 1983, **22**, 48–55

The finely divided powder may ignite if sprayed into moist air [1]. Procedures for preparing polycrystalline and single crystal materials are detailed, with precautions to prevent ignition of material deposited on the walls of the reaction chamber when it is opened to air, or cleaned with water [2].

Copper(I) chloride

Mellor, 1940, Vol. 8, 100

Heated interaction to produce metallic copper is violent.

Silicon tetrafluoride

Porritt, L. J., *Chem. Brit.*, 1979, **15**, 282–283

Interaction is violently exothermic, causing sealed glass tubes at 350°C to explode, and a flow reaction system partially to melt.

See other GLASS INCIDENTS

Transition metal halides

Hector, A. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1993, (13), 1095; *J. Chem. Soc., Dalton Trans.*, 1993, (16), 2435

Titanium and vanadium nitrides may be prepared by a metathesis reaction of their tetrachlorides with the nitride, initiated by heat or friction. The reaction is potentially explosive. Other transition metal halides may cause ampules to explode after thermal initiation when anhydrous and were invariably found to do so when the hydrates were used.

See METAL Pnictides, METATHESIS

See other NITRIDES

4689. Hexalithium disilicide
[12136-61-7]

Li₆Si₂

Li₃SiSiLi₃

Acids

Mellor, 1940, Vol. 6, 170

The silicide incandesces in conc. hydrochloric acid, and with dilute acid evolves silanes which ignite. It explodes with nitric acid and incandesces when floated on sulfuric acid.

Halogens

Mellor, 1940, Vol. 6, 169

When warmed, it ignites in fluorine, but higher temperatures are necessary with chlorine, bromine and iodine.

Non-metals

Mellor, 1940, Vol. 6, 169

Interaction with phosphorus, selenium or tellurium causes incandescence. Sulfur also reacts vigorously.

Water

Mellor, 1940, Vol. 6, 170

Reaction with water is very violent, and the silanes evolved ignite.

See other METAL NON-METALLIDES

4690. Magnesium

[7439-95-4]

Mg

Mg

NSC 426, 1981; *FPA H21*, 1974 (not powder); *HCS 1980*, 598

Preparative hazard

1. *Dust Explosion Prevention: Magnesium Powder, NFPA Standard Codes 480, 651*, both 1987

2. Popov, E. I., *Chem. Abs.*, 1975, **82**, 61415

3. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

All aspects of prevention of magnesium (and aluminium) dust explosions in storage, handling or processing operations are covered in two recent US National Fire Codes [1]. Effects of various parameters on ignition of magnesium powders were studied [2]. Maximum explosion pressures of 7.9 bar, with maximum rate of rise of 884 bar/s have been recorded [3].

See Potassium: Magnesium chloride, etc., *also* Metal halides, below

Acetylenic compounds

See ACETYLENIC COMPOUNDS: metals

Aluminium, Rusted steel, Zinc

See Aluminium–magnesium–zinc alloy: Rusted steel

Ammonium salts, Chlorate salts

MRH Ammonium chlorate 9.07/74

Kirk-Othmer (3rd ed.), 1982, Vol. 19, 494

The mixture ignites when wet owing to formation of unstable ammonium chlorate, and combustion of the metal is very intense.

Atmospheric gases, Water

Darras, R. *et al.*, *Chem. Abs.*, 1961, **55**, 13235

The temperature at which massive magnesium and its alloys will ignite in air depends on the heating programme and the presence or absence of moisture.

Barium carbonate, Water

MCA Case History No. 1849

Mg

Fusion of the metal and salt formed barium acetylide, to which water was added without effective cooling. The vigorous evolution of acetylene blew off the reactor lid and the hot acetylene ignited in air.

See other GAS EVOLUTION INCIDENTS

Beryllium fluoride

Walker, H. L., *School Sci. Rev.*, 1954, **35**(127), 348

Interaction is violently exothermic.

Boron diiodophosphide

See Boron diiodophosphide: Metals

Calcium carbonate, Hydrogen

Mellor, 1940, Vol. 4, 271

Heating an intimate mixture of the powdered metal and carbonate in a stream of hydrogen leads to a violent explosion.

Carbon dioxide

MRH 8.66/49

1. Partington, 1967, 475

2. Driscoll, J. A., *J. Chem. Educ.*, 1978, **55**, 450–451

3. Darras, R. *et al.*, *Chem. Abs.*, 1961, **55**, 13235

A mixture of solid carbon dioxide and magnesium burns very rapidly and brilliantly when ignited [1]. Details for a safe demonstration of the violent combustion have been published [2]. The temperature at which massive magnesium and its alloys will ignite carbon dioxide at 1 bar or 15 bar pressure depends on the heating programme and the presence or absence of moisture. Steady progressive heating caused ignition in carbon dioxide at 900°C (780° if moist), or at 800°C after 4 h (650°/40 min if moist), combustion being violent [3].

Carbon tetrachloride

MRH 5.65/76

See Methanol: Carbon tetrachloride, Metals

Chloroformamidinium nitrate

See Chloroformamidinium nitrate: Metals

Copper compounds

Gorbunov, V. V. *et al.*, *Combustion, Expls, & Shockwaves*, 1986, **22**, 726

Not merely cupric oxide, but the fluoride, chloride and sulfide all form mixtures with heats of combustion above 2 kJ/g. The equivalent lead compounds are less active.

Ethylene oxide

MRH 6.82/79

See Ethylene oxide(reference 3)

Fluorocarbon polymers

Kirk-Othmer (3rd ed.), 1982, Vol. 19, 495

Compressed mixtures are used as special igniters.

See Halocarbons, below

Glass powders

See Metal oxides, below (reference 3)

Halocarbons MRH Carbon tetrachloride 5.65/76, trichloroethylene 5.48/78

1. Clogston, C. C., *Bull. Res. Underwriters Lab.*, 1945, **34**, 5
2. *Pot. Incid. Rept.*, 39, ASESAB, Washington, 1968
3. Mond. Div. ICI, private comm., 1968
4. Hartmann, I., *Ind. Eng. Chem.*, 1948, **40**, 756
5. *Haz. Chem. Data.*, 1975, 115

The powdered metal reacts vigorously and may explode on contact with chloromethane, chloroform, or carbon tetrachloride, or mixtures of these [1]. Mixtures of powdered metal with carbon tetrachloride or trichloroethylene will flash on heavy impact [2]. Violent decomposition with evolution of hydrogen chloride can occur when 1,1,1-trichloroethane comes into contact with magnesium or its alloys with aluminium [3]. Magnesium dust ignited at 400°C in dichlorodifluoromethane (used as an extinguishant for hydrocarbon fires) and suspensions of the dust exploded violently on sparking in the vapour [4]. Interaction with 1,2-dibromoethane may become violent, and gives air-sensitive Grignard compounds [5].

See Fluorocarbon polymers, above; Polytetrafluoroethylene, below

See Bromomethane: Metals

See other CORROSION INCIDENTS

Halogens, or Interhalogens MRH Chlorine 6.78/75, fluorine 17.78/61

Mellor, 1940, Vol. 4, 267

It ignites if moist and burns violently in fluorine, and ignites in moist or warm chlorine. It burns not very readily in bromine vapour and may ignite if finely divided on heating in iodine vapour.

See Chlorine trifluoride: Metals

Iodine heptafluoride: Metals

Iodine: Metals, Water

Hydrogen iodide

Mellor, 1941, Vol. 2, 206

Contact causes momentary ignition.

Lead compounds

See Copper compounds, above

Magnesium sulfate

See Magnesium sulfate: Magnesium

Metal cyanides

Mellor, 1940, Vol. 4, 271

Magnesium reacts with incandescence on heating with cadmium cyanide, cobalt cyanide, copper cyanide, lead cyanide, nickel cyanide and zinc cyanide. With gold

Mg

cyanide or mercury cyanide, the cyanogen released by thermal decomposition of these salts reacts explosively with magnesium.

Metal oxides MRH Chromium trioxide 6.44/73, lead dioxide 3.22/83

1. Mellor, 1941, Vol. 3, 138, 378; 1940, Vol. 4, 272; 1941, Vol. 7, 401
2. Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64
3. Lawrence, K. D. *et al.*, *Chem. Abs.*, 1979, **91**, 41576

Magnesium will reduce violently metal oxides on heating, similarly to aluminium powder in the thermite reaction. Beryllium oxide, cadmium oxide, copper oxide, mercury oxide, molybdenum oxide, tin oxide and zinc oxide are all reduced explosively on heating. Silver oxide reacts with explosive violence when heated with magnesium powder in a sealed tube [1]. Interaction of molten magnesium and iron oxide scale is violent [2]. Use of glass powders as suppressants for burning magnesium showed that some lithium oxide–magnesium oxide frits reacted violently [3].
See other GLASS INCIDENTS, THERMITE REACTIONS

Metal oxosalts MRH Ammonium nitrate 8.79/69, potassium nitrate 7.57/58, silver nitrate 6.95/70, barium sulfate 4.27/71, copper(II) sulfate 7.03/58, sodium sulfate 5.86/60, potassium carbonate, 3.10/66.

1. Mellor, 1940, Vol. 4, 272; Vol. 8, Suppl. 2.1, 545
2. Pieters, 1957, 30
3. Shimizu, T., *Chem. Abs.*, 1991, **114**, 9175

Interaction with fused ammonium nitrate or with metal nitrates, phosphates or sulfates may be explosively violent [1]. Lithium and sodium carbonates may also react vigorously [2]. The mixture with magnesium sulfate has been described as a noisy but low power bursting charge for pyrotechny [3].

See Silver nitrate: Magnesium

Potassium perchlorate: Metal powders

See other THERMITE REACTIONS

Methanol MRH 5.02/73

1. Vogel, 1957, 169
2. Personal experience, 1952
3. Shidlovskii, A. A., *Chem Abs.*, 1947, **41**, 1105d

The reaction of magnesium and methanol to form magnesium methoxide and used to prepare dry methanol [1] is very vigorous, but often subject to a lengthy induction period. Sufficient methanol must be present to absorb the violent exotherm which sometimes occurs [2]. Mixtures of powdered magnesium (or aluminium) and methanol are capable of detonation and are more powerful than military explosives [3].

See other INDUCTION PERIOD INCIDENTS

Methanol, 3-Methyl-2-butenoylanilide

Brettle, R. *et al.*, *Tetrahedron Lett.*, 1980, **21**, 2915

Reduction of the double bond in the anilide by magnesium in methanol is vigorously exothermic after an induction period, and efficient cooling is then necessary.

See Methanol, above.

See other INDUCTION PERIOD INCIDENTS

Molybdenum disulfide

See Molybdenum disulfide, Magnesium

Nitric acid, 2-Nitroaniline

See Nitric acid: Magnesium, 2-Nitroaniline

Nitrogen

See Nitrogen (Liquid): Magnesium

Other reactants

Yoshida, 1980, 347–348

MRH values calculated for 31 combinations, largely with oxidants, are given.

Oxidants

MRH values show % of oxidant

See Carbon dioxide; Halogens etc.; Metal oxides; Metal oxosalts, all above;
and Sulfur, etc., below

Ammonium nitrate: Metals	MRH 8.79/69
Barium peroxide: Metals	
Dinitrogen tetraoxide: Metals	MRH 12.97/50
Hydrogen peroxide: Metals	MRH 12.68/59
Lead(IV)oxide: Metals	MRH 3.22/83
Nitric acid: Metals	MRH 11.97/52
Oxygen (Liquid): Metals	
Potassium chlorate: Metals	MRH 9.50/63
Potassium perchlorate: Powdered metals	
Sodium iodate: Metals	
Sodium nitrate: Magnesium	MRH 8.58/58

Poly(tetrafluoroethylene)

1. Anon., *Indust. Res.*, 1969(9), 15
2. Diercks, B. V., *J. Haz. Mat.*, 1986, **12**(1), 3
3. Potter, L. J., *Proc. 13th Int. Pyrotech. Semin.*, 1988, 639–659

Finely divided magnesium and Teflon are described as a hazardous combination of materials [1]. The compressed pellets of powdered metal and polymer used as infra-red decoy flares have been found sensitive to ignition by electrostatic sparks during manufacture [2]. The ignition sensitivity of igniter mixtures containing magnesium, Teflon and Viton was studied, following two fires [3].

See Fluoropolymers, also Halocarbons, both above

Potassium carbonate

MRH 3.10/66

Winckler, C., *Ber.*, 1890, **23**, 442

The mixture of magnesium and potassium carbonate recommended by Castellana as a safe substitute for molten sodium in the Lassaigne test can itself be hazardous,

Mg

as an equimolar mixture gives an explosive substance (possibly 'carbonylpotassium', potassium benzenehexoxide) on heating.

Rusty steel

Billinge, K., *Fire Prev. Sci. Technol.*, 1981, (24), 13–19

Impact of a magnesium anode on rusty steel led to ignition and explosion in a tanker.

See other THERMITE REACTIONS

Silicon dioxide

1. Barker, W. D., *School Sci. Rev.*, 1938, **20**(77), 150

2. Marle, E. R., *Nature*, 1909, **82**, 428

Heating a mixture of powdered magnesium and silica (later found not to be absolutely dry) caused a violent explosion rather than the vigorous reaction anticipated [1]. A warning had been published previously [2].

See also FLASH POWDER

Sulfur, or Tellurium

MRH Sulfur 6.15/57

1. Brauer, 1963, Vol. 1. 913

2. Hutton, K., *School Sci. Rev.*, 1950, **31**(114), 265

3. Bodner, G. M., *J. Chem. Educ.*, 1985, **62**(12), 1107

Interaction of magnesium with sulfur or tellurium at elevated temperatures may be violent [1] or even explosive [2], even with tiny amounts [3].

See Sulfur: Metals

Water

1. Shidlovskii, A. A., *Chem. Abs.*, 1947, **41**, 1105d

2. Taylor, F. R. *et al.*, *Rept. ARAED-T-87022*, 1–26, 1988; O/No. AD A187158, NTIS, Richmond (Va.)

Mixtures of magnesium (or aluminium) powder with water can be caused to explode powerfully by initiation with a boosted detonator [1]. The use of organic coatings on magnesium or aluminium powder in pyrotechnic compositions prevents reaction with atmospheric moisture and problems resulting from hydrogen evolution [2].

See Methanol, above; *also* Aluminium–magnesium alloy: Water

Water, Uranium

1. Anon., *Loss Prev. Bull.*, 1992, **103**, 19

2. Pitt, M. J., personal communication, 1993

A drum of graphite with magnesium and uranium, stored underwater with a 6mm diameter relief hole, burst and scattered its contents. This was attributed to insufficient venting of hydrogen evolved by reaction of magnesium and water [1]. If the uranium was present as metal, it seems likely to have been an even more potent source of hydrogen [2].

See other CORROSION INCIDENTS

See other METALS, REDUCANTS

4691. Magnesium permanganate

[10377-62-5]



Organic compounds

See Zinc permanganate: Organic compounds

See other METAL OXOMETALLATES, OXIDANTS

4692. Magnesium nitrite

[15070-34-5]



Ephraim, 1939, 686–687

Less stable, more hygroscopic and readily hydrolysed than alkali metal nitrites, it may function as a more powerful oxidant.

See other METAL NITRITES, OXIDANTS

4693. Magnesium nitrate

[10377-60-3]



HCS 1980, 601

Dimethylformamide

MRH 5.06/22

'DMF Brochure', Billingham, ICI Ltd., 1965

Magnesium nitrate has been reported to undergo spontaneous decomposition in DMF, (possibly as a result of hydrolysis of the hexahydrate above its m.p., 90°C to liberate nitric acid). Although this effect has not been observed with other nitrates, reaction mixtures with hydrolysable nitrates should be treated with care.

See Sodium nitrate: Jute, Magnesium chloride

Other reactants

Yoshida, 1980, 205

MRH values calculated for 16 combinations with oxidisable materials are given.

Tin(II) fluoride

Denes, G., *Chem. Eng. News*, 1988, **66**(14), 2

Addition of an aqueous solution of the fluoride to one of the nitrate unexpectedly gave no precipitate of magnesium fluoride, and the solution was allowed to evaporate, first at ambient temperature, and then in a warm place, giving a wet white solid. While being manipulated with a spatula, this exploded with some violence, brown fumes were evolved and the solid was transformed into a dry tan-coloured powder. It was thought that oxidation of tin(II) to (IV) had occurred, the nitrate being reduced to nitrogen oxide.

See Tetrahydroxotritin(2+) nitrate

See other REDOX REACTIONS
See other METAL NITRATES, OXIDANTS

4694. Magnesium azide
[39108-12-8]



Mellor, 1940, Vol. 8, 350
It explodes on heating.
See other METAL AZIDES

4695. Magnesium oxide
[1309-48-4]



HCS 1980, 602

Interhalogens

See Bromine pentafluoride: Acids, etc., Chlorine trifluoride: Metals, etc.

Phosphorus pentachloride

See Phosphorus pentachloride: Magnesium oxide

See other INORGANIC BASES, METAL OXIDES

4696. Magnesium sulfate
[18939-43-0]



HCS 1980, 603

Ethoxyethynyl alcohols

See ETHOXYETHYNYL ALCOHOLS *See other* METAL OXONON-METALLATES

Magnesium

Shimizu, T., *Chem. Abs.*, 1991, **114**, 9175

The mixture with magnesium powder has been described as a pyrotechnic explosive.

See Magnesium: Metal oxosalts

4697. Magnesium silicide
[22831-39-6]



Acids, Water

Sorbe, 1968, 84

1765

Contact with moisture under acidic conditions generates silanes which ignite in air.
See other METAL NON-METALLIDES

4698. Magnesium nitride
[12057-71-5]



See entry REFRACTORY POWDERS
See other NITRIDES

4699. Magnesium phosphide
[12057-74-8]



Oxidants

Mellor, 1940, Vol. 8, 842

It ignites on heating in chlorine, or in bromine or iodine vapours at higher temperatures. Reaction with nitric acid causes incandescence.

Water

Mellor, 1940, Vol. 8, 842

Phosphine is evolved and may ignite.

See other METAL NON-METALLIDES

4700. Manganese
[7429-96-5]



NSC 306, 1982

Aluminium

Occup. Hazards, 1966, **28**(11), 44

The finely divided metal is pyrophoric, and a mixture of manganese and aluminium dusts accidentally released from a filter bag exploded violently.

Carbon dioxide

See Carbon dioxide: Metals

Other reactants

Yoshida, 1980, 352

MRH values calculated for 16 combinations with oxidants are given.

Oxidants MRH Chlorine 3.85/56, hydrogen peroxide 5.36/48, nitric acid 4.85/41
Mellor, 1942, Vol. 12. 186–188

Powdered manganese ignites and becomes incandescent in fluorine or on warming in chlorine. Contact with conc. hydrogen peroxide causes violent decomposition

and/or ignition, and with nitric acid incandescence and a feeble explosion were observed.

See Ammonium nitrate: Metals MRH 3.80/69

Bromine pentafluoride: Acids, etc. MRH 4.52/65

Nitryl fluoride: Metals

Phosphorus

See Phosphorus: Metals

Sulfur dioxide

Mellor, 1942, Vol. 12, 187

Pyrophoric manganese burns brilliantly on warming in sulfur dioxide.

See other METALS, PYROPHORIC METALS

4701. Manganese(II) nitrate

[10377-66-9]



Other reactants

Yoshida, 1980, 206

MRH values calculated for 16 combinations with oxidisable materials are given.

Urea

Novikov, A. V. *et al.*, *Chem. Abs.*, 1975, **82**, 48151


The anhydrous complex with urea finally decomposes at 240°C with a light explosion.

See also AMMINOMETAL OXOSALTS

See other METAL NITRATES

4702. Manganese(III) azide

[[] (not isolated)]



Alkenes

1. Fristad, W. F. *et al.*, *J. Org. Chem.*, 1985, **50**, 3647–3649

2. *Chem. Abs.*, **109**, 82979g

The preparation and immediate use of manganese(III) azide species generated slowly in situ by refluxing manganese(III) acetate and sodium azide in acetic acid in presence of alkene reaction substrates to prepare 1,2-diazoalkanes avoids the need to isolate manganese(III) azide which has a high probability of explosive instability [1]. The (II) azide is known [20260-90-6], isolable, detonable and has been patented as a power source for lasers [2].

Styrene

See Sodium azide: Manganese(III) salts, Styrene

See related METAL AZIDES

4703. Sodium permanganate
[10101-50-5]



Acetic acid, or Acetic anhydride
See Potassium permanganate: Acetic acid

Methanol

US Government Source, Internet (www.dne.bnl.gov/etd/csc), 1996
An exothermic reaction from these two is reported, causing boiling and spraying of the mixture. It is not clear if anything else (acid?) was involved.

Other reactants

Yoshida, 1980, 93
MRH values calculated for 20 combinations with oxidisable materials are given.
See other METAL OXOMETALLATES, OXIDANTS

4704. Manganese(II) oxide
[1344-43-0]



Hydrogen peroxide
See Hydrogen peroxide: Metals, etc.
See other METAL OXIDES

4705. Manganese(IV) oxide (Manganese dioxide)
[1313-13-9]



HCS 1980, 609

Aluminium

Sidgwick, 1950, 1265
Interaction on heating is very violent (a thermite reaction).
See Aluminium: Metal oxides
See other THERMITE REACTIONS

Anilinium perchlorate

See Anilinium perchlorate: Metal oxides

Calcium hydride

Pascal, 1958, Vol. 4, 304
Interaction becomes incandescent on warming.
See other REDOX REACTIONS

Carbon black, Chlorinated paraffin, Lead(IV) oxide, Manganese(IV) oxide
See Lead(IV) oxide: Carbon black, etc.

Diboron tetrafluoride

See Diboron tetrafluoride: Metal oxides

Rubidium acetylide

See Rubidium acetylide: Metal oxides

Hydrogen sulfide

See Hydrogen sulfide: Metal oxides

Hydroxylaminium chloride

See Hydroxylaminium chloride: Manganese dioxide

Lithium

Hamporzumian, K. *et al.*, *J. Power Sources*, 1982, **8**, 35–40

Manganese dioxide–lithium batteries are generally safer than sulfur dioxide–lithium cells.

See Lithium: Sulfur dioxide

Oxidants MRH Chlorine trifluoride 1.80/52, hydrogen peroxide 1.59/99+

1. Mellor, 1942, Vol. 12, 254

2. Mellor, 1941, Vol. 1, 936

3. Ahrle, H., *Z. Angew. Chem.*, 1909, **22**, 713

4. von Schwartz, 1918, 323

5. Blanco Prieto, J., *Chem. Abs.*, 1976, **84**, 46791

Action of chlorine trifluoride causes incandescence [1]. Manganese dioxide catalytically decomposes powerful oxidising agents, often violently. Dropped into conc. hydrogen peroxide, the powdered oxide may cause explosion [2]. Either the massive or the powdered oxide explosively decomposes 92% peroxomonosulfuric acid [3], and mixtures with chlorates ('oxygen mixture', heated to generate the gas) may react with explosive violence [4]. Cuban pyrolusite can be used in place of potassium dichromate to promote thermal decomposition of potassium chlorate in match-head formulations [5].

See Peroxyformic acid: Metals, etc.

Potassium azide

Mellor, 1940, Vol. 8, 347

On gentle warming, interaction is violent.

See other METAL OXIDES, OXIDANTS

4706. Manganese(II) sulfide

[18820-29-6]

MnS

MnS

Mellor, 1942, Vol. 12, 394

The vacuum-dried red sulfide becomes red-hot on exposure to air.

See other METAL SULFIDES, PYROPHORIC MATERIALS

1769

4707. Manganese(IV) sulfide
[12125-23-4]



Sorbe, 1968, 84
It explodes at 580°C.
See other METAL SULFIDES

4708. Manganese(II) telluride
[12032-88-1]



Lithium
See Lithium: Metal oxides, etc.
See other METAL NON-METALLIDES


4709. Manganese(VII) oxide
[12057-92-0]



1. Mellor, 1942, Vol. 12, 293
2. Bailar, 1973, Vol. 3, 805
An unstable powerful oxidant, it explodes between 40 and 70°C, or on friction or impact, sensitivity being as great as that of mercury fulminate [1]. Detonation occurs at 95°C, and under vacuum explosive decomposition occurs above 10°C [2].
See Potassium permanganate: Sulfuric acid

Organic material
1. Patterson, A. M., *Chem. Eng. News*, 1948, **26**, 711
2. Mellor, 1940, Vol. 12, 293
A sample of the heptoxide exploded in contact with the grease on a stopcock [1], and explosion or ignition has been noted with various solvents, oils, fats and fibres [2].
See other METAL OXIDES, OXIDANTS

4710. Zinc permanganate
[23414-72-4]



Cellulose
Sörgel, U. C., *Pharm. Prax., Beil. Pharmaz.*, 1960, (2), 30
An explosion involved contact of the permanganate with cellulose.

Organic compounds
Wolfe, S. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 7755–7757

Solid zinc permanganate and magnesium permanganate react violently with many classes of organic compounds. The reactions may be controlled successfully by supporting the oxidants on silica. The zinc salt is more effective than the magnesium salt, but both are much more reactive than potassium permanganate.

See other METAL OXOMETALLATES, OXIDANTS

4711. Manganese phosphide

[12263-33-1]



Chlorine

See Chlorine: Phosphorus compounds

See other METAL NON-METALLIDES

4712. Molybdenum

[7439-98-7]



Oxidants

MRH value shows % of oxidant

See Bromine pentafluoride: Acids, etc.

Bromine trifluoride: Halogens, etc.

Chlorine trifluoride: Metals

Fluorine: Metals

MRH 7.57/54

Iodine pentafluoride: Metals

Lead(IV) oxide: Metals

Nitryl fluoride: Metals

Potassium perchlorate: Metal powders

Sodium peroxide: Metals

See other METALS

4713. Sodium molybdate

[10102-40-6]



Aniline, Nitrobenzene

Nolan, 1983, Case history 144

In the preparation of a dyestuff from aniline, nitrobenzene (as oxidant), hydrochloric acid and sodium hydroxide, ferric chloride is often used as catalyst, but sodium molybdate was substituted as a more effective catalyst. The materials were charged into a 4.5 m³ reactor and heating was started after addition of nitrobenzene, but the temperature controller was mis-set, and overheating at a high rate ensued. The exotherm was much higher than normal because of the more effective catalyst, and partial failure of the cooling water led to an uncontrollable exotherm.

See other METAL OXOMETALLATES, RUNAWAY REACTIONS

4714. Molybdenum nitride

[37245-81-1]

Mo_nN

[12033-19-1] MoN; [12033-31-7] Mo₂N; [97953-19-0] Mo₄N

Complex structures

Nitric acid

Chorley, R. W., *Chem. Brit.*, 1993, **29**(10), 867

High surface area samples may be pyrophoric, but can be passivated by exposure to oxygen. A passivated sample (50 mg) caused a small explosion when concentrated nitric acid was added to it.

See other NITRIDES

See also MIXING

4715. Sodium tetraperoxomolybdate

[42489-15-6] (ion)

MoNa₂O₈

Na₂[Mo(O₂)₄]

Castrantas, 1965, 5

It decomposes explosively under vacuum.

See other PEROXOACID SALTS

4716. Molybdenum(IV) oxide

[18868-43-4]

MoO₂

MoO₂

Gibson, 1969, 102

It becomes incandescent in contact with air.

See other METAL OXIDES, PYROPHORIC MATERIALS

4717. Molybdenum(VI) oxide

[1313-27-5]

MoO₃

MoO₃

Graphite

1. Young, J. A., private comm., 1989; *CHAS Notes*, 1992, **X**(6), 2

2. Partington, 1967, 753

During preparation of an alloy steel, molybdenum trioxide (372 kg), and then graphite (340 kg, 1.56 equiv., in 4 portions), were charged into an electric furnace containing a charge of melted scrap steel at 1500–1600°C, using a charging ram and rotatable box for the additions through the furnace door. Immediately after the last graphite addition, as the ram and box were being withdrawn, a large fireball emerged through the furnace door and fatally burned the ram operator. This was caused by the very rapid exothermic redox reaction between the oxide and graphite to give molybdenum and a huge volume (some 800 m³) of very

hot carbon monoxide) which emerged at high velocity from the 27 m³ furnace and ignited, giving a flame-jet some 15 m long and 2–3 m in diameter [1]. It is possible that some of the graphite particles were ejected in the gas jet and augmented the combustion intensity. Reduction of the trioxide to molybdenum by strongly heating with carbon has long been a commercial metallurgical process [2].
See other REDOX REACTIONS

Interhalogens

See Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Metals. etc.

Metals

Mellor, 1943, Vol. 11, 542
Reduction of the oxide by heated sodium or potassium proceeds with incandescence, and explosion occurs in contact with molten magnesium.
See Lithium: Metal oxides
See other METAL OXIDES

4718. Oxodiperoxomolybdenum–hexamethylphosphoramide

[53474-95-6 + 680-31-9] MoO₅·C₆H₁₈N₃OP
 $\text{MoO}(\text{O}_2)_2 \cdot \text{O} \cdot \text{P}(\text{NMe}_2)_3$

Fieser, 1974, Vol. 4, 203
The monohydrated complex exploded after storage at ambient temperature for a month. Refrigerated storage is recommended.
See related METAL PEROXIDES, AMMINECHROMIUM PEROXOCOMPLEXES

4719. Molybdenum(IV) sulfide

[1317-33-5] MoS₂
 MoS_2

Magnesium

Danielyan, N. G. *et al.*, *Mod. Phys. Lett. B.*, 1991, **5**(19), 1301
Molybdenum disulphide is capable of forming a thermite type mixture with magnesium, which may explode on grinding.
See also THERMITE REACTIONS

Potassium nitrate

See Potassium nitrate: Metal sulfides
See other METAL SULFIDES

4720. Sodium nitrite

[7632-00-0] NNaO₂
 $\text{NaON} \cdot \text{O}$

FPA H133, 1984; *HCS 1980*, 852

Aminoguanidine salts

Urbanski, 1967, Vol. 3, 207

Interaction, without addition of acid, produces tetrazolylguanidine ('tetrazene'), a primary explosive of equal sensitivity to mercury(II) azide, but more readily initiated.

Ammonium salts

MRH Ammonium chloride 1.80/44, nitrate 1.46/99+

1. von Schwartz, 1918, 299

2. Mellor, 1967, Vol. 8, Suppl. 2.1, 388

3. *RoSPA Occ. Safety and Health Suppl.*, 1972, **2**(10), 32

Heating a mixture of an ammonium salt with a nitrite salt causes a violent explosion on melting [1], owing to formation and decomposition of ammonium nitrite. Salts of other nitrogenous bases behave similarly. Mixtures of ammonium chloride and sodium nitrite are used as commercial explosives [2]. Accidental contact of traces of ammonium nitrate with sodium nitrite residues caused wooden decking on a truck to ignite [3].

See wood, below; also NITRITE SALTS OF NITROGENOUS BASES

1,3-Butadiene

1. Beer, R. N., private comm., 1972

2. Keister, R. G. *et al.*, *Loss. Prev.*, 1971, **5**, 69

Sodium nitrite solution is used to inhibit 'popcorn' polymerisation of butadiene in processing plants. If conc. nitrite solutions (5%) are used, a black sludge is produced, which when dry, will ignite and burn when heated to 150°C, even in absence of air [1]. The sludge produced from use of nitrite solutions to scavenge oxygen in butadiene distillation systems contained 80% of organic polymer and a nitrate:nitrite ratio of 2:1. Use of dilute nitrite solution (0.5%) or pH above 8 prevents sludge formation [2].

See Buten-3-yne: 1,3-Butadiene, and Nitrogen oxide: Dienes

Hydroxylamine hydrochloride

Scott, E. S., *J. Chem. Ed.*, 1992, **69**(12), 1028

If hydroxylamine hydrochloride and sodium nitrite, of normal laboratory particle size, are swiftly shaken together and left to stand, a vigorous gas evolving reaction will start after a short induction period. This is a rare example of spontaneous reaction of solids

See Ammonium salts, above

Maize husks, Sodium nitrate

See Sodium nitrate: Maize husks, etc.

Metal amidosulfates

Heubel, J. *et al.*, *Compt. rend.*, 1963, **257**, 684–686

Interaction of nitrites when heated with metal amidosulfates (sulfamates) may become explosively violent owing to liberation of nitrogen and steam. Mixtures with ammonium sulfamate form ammonium nitrite which decomposes violently around 80°C.

See Sodium nitrate: Metal amidosulfates

Metal cyanides

1. von Schwartz, 1918, 299
2. Mellor, 1940, Vol. 8, 478
3. Greenwood, P. H. S., *J. Proc. R. Inst. Chem.*, 1947, 137
4. Elson, C. H. R., *J. Proc. R. Inst. Chem.*, 1947, 19
5. Eiter, K. *et al.*, Austrian Pat. 176 784, 1953

Mixtures of sodium nitrite and various cyanides [1] explode on heating, including potassium cyanide [2], potassium hexacyanoferrate(III), sodium pentacyanonitrosylferrate(II) [3], potassium hexacyanoferrate(II) [4], or mercury(II) cyanide [5]. Such mixtures have been proposed as explosives, initiatable by heat or a detonator [5].

Other reactants

Yoshida, 1980, 13

MRH values calculated for 19 combinations with oxidisable materials are given.

Paper, Sulfur

Anon., *CIHSC Chem. Safety Summ.*, 1984, **55**(218), 36

Empty paper sacks, some of which had contained sulfur or inorganic salts but mostly sodium nitrite, were dumped (with general rubbish) into a skip located inside by windows in direct sunlight. The fire which resulted was probably caused by exothermic interaction of the sodium nitrite with sulfur and damp paper, with further self heating aided by the generally insulating nature of the mound of paper sacks in the draught-free skip warmed by sunlight.

See Wood, below

See other SELF-HEATING AND IGNITION INCIDENTS

Phenol

Hatton, J. P., private comm., 1976

A mixture exploded violently on heating in a test tube.

Phthalic acid, or Phthalic anhydride

Hawes, B. V. W., *J. R. Inst. Chem.*, 1955, **79**, 668

Mixtures of sodium nitrite and phthalic acid or phthalic anhydride explode violently on heating. A nitrite ester may have been produced.

See ACYL NITRITES

Potassium thiocyanate

MRH (sodium salt) 3.57/28

1. Mellor, 1940, Vol. 8, 478
2. Baum, R., *Chem. Eng. News*, 1982, **60**(41), 29
3. Fawcett, H. H., *Chem. Eng. News*, 1982, **60**(46), 4
4. Shriver, D. F., also Coleman, L. F., *Chem. Eng. News*, 1982, **60**(47), 2
5. Pinsky, B., *Chem. Eng. News*, 1982, **60**(47), 2
6. Viola, A., *Chem. Eng. News*, 1983, **61**(3), 76

A mixture of the nitrite with sodium thiocyanate explodes on heating [1]. Preparation of a molten salt bath from 0.45 kg of potassium thiocyanate (reducing agent) and 1.35 kg of sodium nitrite (oxidant) led to a violent explosion on melting, which

caused severe structural damage to the laboratory [2]. It was claimed that this could not have been foreseen from available information [2], but this was vigorously refuted [3,4]. Use of sand baths, rather than salt baths for laboratory heating purposes is to be preferred [5]. The relative lack of descriptive chemistry in modern curricula is cited as the major contributory factor to the general ignorance which led to the explosion [6].

See METAL THIOCYANATES: oxidants, MOLTEN SALT BATHS

See other REDOX REACTIONS

Reducants

See Potassium thiocyanate, above

Sodium disulfite; Sodium thiosulfate, both below

See Sodium nitroxylate

Sodium amide

Bergstrom, F. W. *et al.*, *Chem. Rev.*, 1933, **12**, 64

Addition of solid sodium nitrite to the molten amide caused immediate gas evolution, followed by a violent explosion.

Sodium disulfite

MRH 1.84/68

MCA Case History No. 183

Large-scale addition of solid sodium disulfite to an unstirred and too-concentrated solution of sodium nitrite caused a vigorous exothermic reaction.

See other REDOX REACTIONS

Sodium nitrate, Sodium sulfide

Janz, 1976, Table 3, 22

Accidental mixing of the 3 molten salts caused a violent explosion.

See other REDOX REACTIONS

Sodium thiocyanate

See Potassium thiocyanate, above

See other REDOX REACTIONS

Sodium thiosulfate

Mellor, 1940, Vol. 8, 478; 1947, Vol. 10, 501

There is no interaction between solutions, but evaporation of the mixture gave a residue which explodes on heating. The mixed solids behave similarly.

See other REDOX REACTIONS

Urea

MRH 2.80/30

Bucci, F., *Ann. Chim. (Rome)*, 1951, **41**, 587

Fusion of urea (2 mol) with sodium nitrite (or potassium nitrite, 1 mol of either) to give high yields of the cyanate must be carried out exactly as described to avoid the risk of explosion.

Wood

Anon., *ABCM Quart. Safety Summ.*, 1944, **15**, 30

Wooden staging, which had become impregnated over a number of years with sodium nitrite, became accidentally ignited and burned as fiercely as if impregnated with potassium chlorate. Although the effect of impregnating cellulosic material with sodium nitrate is well known, that due to sodium nitrite was unexpected.

See Ammonium salts (reference 3), *also* Paper, Sulfur, both above

See other METAL NITRITES, METAL OXONON-METALLATES, OXIDANTS

4721. Sodium nitrate

[7631-99-4]



NNaO_3

FPA H26, 1983; *HCS 1980*, 851

Acetic anhydride

See Acetic anhydride: Metal nitrates

Aluminium

MRH Aluminium 4.43/27

1. Anon., *Fire*, 1935, **28**, 30
2. Farnell, P. L. *et al.*, *Chem. Abs.*, 1972, **76**, 129564
3. Gerling, H. W., *Metall* (Berlin), 1982, **36**, 698–699

Mixtures of the nitrate with powdered aluminium or its oxide (the latter seems unlikely) were reported to be explosive [1], and the performance characteristics of flares containing compressed mixtures of the metal and nitrate have been evaluated [2]. A violent explosion in a copper smelting works was caused mainly by reaction of aluminium with sodium nitrate [3].

Aluminium, Water

Jackson, B. *et al.*, *Chem. Abs.*, 1976, **84**, 46798

During investigation of pyrotechnic flare formulations, it was found that mixtures of the metal powder and oxidant underwent a low-temperature exothermic reaction at 70–135°C in presence of moisture.

Antimony

Mellor, 1939, Vol. 9, 382

Powdered antimony explodes when heated with an alkali metal nitrate.

Arsenic trioxide, Iron(II) sulfate

Dunn., B. W., *Bur. Expl. Rept. 13*, Pennsylvania, Amer. Rail Assoc., 1920

A veterinary preparation containing the oxidant and reductant materials (possibly with some additional combustibles) ignited spontaneously.

See other REDOX REACTIONS

Barium thiocyanate

Pieters, 1957, 30

Mixtures may explode

See Sodium thiocyanate, below

Bitumen

Xu, Y.-C. *et al.*, *Chem. Abs.*, 1980, **93**, 209761

The induction periods for the reaction of sodium nitrate–bitumen mixtures (43:57 wt) heated at 195, 234 or 260°C are 44, 2 and 0.5 h, respectively. Further study of sodium nitrate/bitumenised waste systems held at these temperatures showed an initial weak exotherm around 260°C and a larger exotherm (0.96–1.21 kJ/g) accompanied by 50% wt loss around 430°C.

Boron phosphide

Mellor, 1940, Vol. 8, 845

Deflagration occurs in contact with molten alkali metal nitrates.

Calcium–silicon alloy

Smolin, A. O., *Chem. Abs.*, 1974, **81**, 138219

Combustion of the alloy in admixture with sodium nitrate is mentioned in an explosives context, but no details are translated.

Fibrous material

MRH Cellulose 3.85/30

1. Mellor, 1961, Vol. 2, Suppl. 2.1, 1244

2. Anon., *ABCM Quart. Safety Summ.*, 1944, **15**, 30

Fibrous organic material (jute storage bags) is oxidised in contact with sodium nitrate above 160°C and will ignite below 220°C [1]. Wood and similar cellulosic materials are rendered highly combustible by nitrate impregnation [2].

Jute, Magnesium chloride

van Hoogstraten, C. W., *Chem. Abs.*, 1947, **41**, 3692b

Solid crude sodium nitrate packed in jute bags sometimes ignited the latter in storage. Normally ignition did not occur below 240°C, but in cases where magnesium chloride (up to 16%) was present, ignition occurred at 130°C. This was attributed to formation of magnesium nitrate hexahydrate, which hydrolyses above its m.p. (90°C) liberating nitric acid. The latter was thought to have caused ignition of the jute bags under unusual conditions of temperature and friction.

See Magnesium nitrate: Dimethylformamide

Magnesium

MRH 8.58/42

Bond, B. D. *et al.*, *Combust. Flame*, 1966, **10**, 349–354

A study of the kinetics in attack of magnesium by molten sodium nitrate indicates that decomposition of the nitrate releases oxygen atoms which oxidise the metal so exothermally that ignition ensues.

See Magnesium: Metal oxosalts

Maize husks, Sodium nitrite

Anon., *CISHC Chem. Safety Summ.*, 1985, **52**(222), 66

A rotary drum had been used previously to dry metal components which had been heat-treated in nitrate–nitrite molten salt baths, washed, then tumble dried with ground maize husks to absorb adhering water. When the drum was taken out of

NNaO₃

service, it was not cleaned out. After some 10 months it was recommissioned, but while being heated up to operating temperature, an explosion occurred which ejected flame jets for several m. This was attributed to presence of considerable contamination of the maize husks by metal nitrate–nitrite residues, and ignition on heating of such material was confirmed experimentally.

See MOLTEN SALT BATHS

Metal amidosulfates

Heubel, J. *et al.*, *Compt. rend.*, 1962, **255**, 708–709

Interaction of nitrates when heated with amidosulfates (sulfamates) may become explosively violent owing to liberation of dinitrogen oxide and steam.

See Sodium nitrite: Metal amidosulfates

Metal cyanides

MRH Sodium cyanide 4.31/37

See MOLTEN SALT BATHS

Non-metals

MRH Phosphorus (r) 4.43/27, sulfur 3.39/24

1. Mellor, 1941, Vol. 2, 820

2. Leleu, *Cahiers*, 1980, (99), 279

Contact of powdered charcoal with the molten nitrate, or of the solid nitrate with glowing charcoal, causes vigorous combustion of the carbon. Mixtures with charcoal and sulfur have been used as black powder [1]. Charcoal powder–nitrate mixtures burn briskly at 200°C [2].

Other reactants

Yoshida, 1980, 199

MRH values calculated for 19 combinations with oxidisable materials are given.

Peroxyformic acid

See Peroxyformic acid: Sodium nitrate

Phenol, Trifluoroacetic acid

Spitzer, U. A. *et al.*, *J. Org. Chem.*, 1974, **39**, 3936

When the salt–acid nitration mixture was applied to phenol, a potentially hazardous rapid exothermic reaction occurred producing tar.

See other NITRATION INCIDENTS

Sodium

MRH 3.85/57

Mellor, 1961, Vol. 2, Suppl. 2.1, 518

Interaction of sodium nitrate and sodium alone, or dissolved in liquid ammonia, eventually gives a yellow explosive compound.

See Sodium nitroxylate

Sodium nitrite, Sodium sulfide

See Sodium nitrite: Sodium nitrate, Sodium sulfide

Sodium phosphinate

MRH 1.00/56

Mellor, 1941, Vol. 2, 820

A mixture exploded violently on warming.
See other REDOX REACTIONS

Sodium thiocyanate

Anon., *Sichere Chemiearbeit*, 1994, **46**, 80
Sodium nitrate and sodium thiocyanate were being blended in a ploughshare mixer to produce a plating formulation. The blend deflagrated, burning four workers and blowing apart the mixer. Subsequent laboratory investigation showed such mixtures to be friction sensitive. The firm has ceased to prepare such mixtures. Caution is advised when mixing oxidants and reductants.

Sodium thiosulfate

Mellor, 1941, Vol. 2, 820
A mixture is explosive when heated.
See other REDOX REACTIONS

Tris(cyclopentadienyl)cerium

1. Jakob, K., *Z. Anorg. Allg. Chem.*, 1988, **556**, 170
2. Grodett, P. S., *Chem. Eng. News*, 1988, **66**(31), 2
Reaction of ammonium hexanitrocerate and cyclopentadienylsodium under inert conditions gives tris(cyclopentadienyl)cerium and sodium nitrate, removed by filtration before evaporation of solvent [1]. When the filtration step was omitted, and the evaporated solid mixture was heated to 75°C, a violent explosion occurred. This may have involved complexes of the type $\text{Ce}(\text{NO}_3)\text{Cp}_2 \cdot \text{NaNO}_3$ [2], but a direct redox reaction between the reactive CeCp_3 and the oxidant is also possible.
See other REDOX REACTIONS

Wood

See Fibrous material, above
See other METAL NITRATES, OXIDANTS

4722. Sodium nitroxylate [13968-14-4]



Air, or Carbon dioxide, or Water

Mellor, 1963, Vol. 2, Suppl. 2.2, 1566
The solid is very reactive towards air, moisture or carbon dioxide, and tends to explode readily, also decomposing violently on heating. It is produced from sodium nitrate or sodium nitrite by electrolytic reduction, or action of sodium.
See other N-O COMPOUNDS

4723. Sodium nitride
[12136-83-3]



NNa_3

It decomposes explosively on gentle warming.
See other NITRIDES

4724. Nitrogen oxide ('Nitric oxide')
[10102-43-9]



NO

1. *UK Sci. Mission Rept. 68/79*, Washington, UKSM, 1968
2. Miller, R. O., *Ind. Eng. Chem., Proc. Res. Dev.*, 1968, **7** 590–593
3. Ribovich, J. *et al.*, *J. Haz. Mat.*, 1977, **1**, 275–287
4. Ramsay, J. B. *et al.*, *Proc. 6th Int. Symp. Detonation*, 1976, 723–728

Liquid nitrogen oxide and other cryogenic oxidisers (ozone, fluorine in presence of water) are very sensitive to detonation in absence of fuel, and can be initiated as readily as glyceryl nitrate [1,2]. Detonation of the endothermic liquid oxide close to its b.p. (-152°C) generated a 100 kbar pulse and fragmented the test equipment. It is the simplest molecule that is capable of detonation in all 3 phases [3]. The liquid oxide is sensitive and may explode during distillation [4]. It is highly endothermic (ΔH_f° (g) $+90.4$ kJ/mol, 3.01 kJ/g) and an active oxidant.

Acetylene, Perchloric acid

See Perchloric acid: Acetylene, Nitric oxide

Alkenes, Oxygen

See Nitrogen dioxide: Alkenes

Ammonia

Checkel, M. D. *et al.*, *J. Loss Prevention*, 1995, **8**(4), 215

Flammability limits of mixtures of nitric oxide and ammonia were studied. The limits are 15–70% ammonia; tenfold pressure rise and explosion are possible. Introduction of oxygen into flammable compositions will produce autoignition, via nitrogen dioxide formation, contamination with air might.

Carbon, Potassium hydrogen tartrate

Pascal, P. *et al.*, *Mém. Poudres*, 1953, **35**, 335–347

The ignition temperature of 400°C for carbon black in nitrogen oxide was reduced by the presence of the tartrate.

Carbon disulfide

Winderlich, R., *J. Chem. Educ.*, 1950, **27**, 669

A demonstration of combustion of carbon disulfide in nitrogen oxide (both endothermic compounds) exploded violently.

Dichlorine oxide

See Dichlorine oxide: Nitrogen oxide

Dienes, Oxygen

1. Haseba, S. *et al.*, *Chem. Eng. Progr.*, 1966, **62**(4), 92
2. Schuftan, P. M., *Chem. Eng. Progr.*, 1966, **62**(7), 8

Violent explosions which occurred at -100 to -180°C in ammonia synthesis gas units were traced to the formation of explosive addition products of dienes and oxides of nitrogen, produced from interaction of nitrogen oxide and oxygen. Laboratory experiments showed that the addition products from 1,3-butadiene or cyclopentadiene formed rapidly at about -150°C , and ignited or exploded on warming to -35 to -15°C . The unconjugated propadiene, and alkenes or acetylene reacted slowly and the products did not ignite until $+30$ to $+50^{\circ}\text{C}$ [1]. This type of derivative ('pseudo-nitrosite') was formerly used (Wallach) to characterise terpene hydrocarbons. Further comments were made later [2].

See Sodium nitrite: 1,3-Butadiene

Fluorine

See Fluorine: Non-metal oxides

Hydrogen, Nitrous oxide

Glowinski, J. *et al.*, *Przem. Chem.*, 1996, **75**(2), 63

Explosive limits for this combination have been studied from 1 to 20 bar at 50°C .

Hydrogen, Oxygen

Chanmugam, J. *et al.*, *Nature*, 1952, **170**, 1067

Pre-addition of nitrogen oxide (or nitrosyl chloride as its precursor) to stoichiometric hydrogen-oxygen mixtures at 240 mbar/ 360°C will cause immediate ignition under a variety of circumstances.

Metal acetylides or carbides

Mellor, 1946, Vol. 5, 848, 891

Rubidium acetylide ignites on heating, and uranium dicarbide incandesces in the gas at 370°C .

Metals

Mellor, 1940, Vol. 8, 436; 1943, Vol. 11, 162; 1942, Vol. 12, 32

Pyrophoric chromium attains incandescence in the oxide, while calcium, potassium and uranium need heating before ignition occurs, when combustion is brilliant in the 53% oxygen content.

Methanol

Partel, G., *Riv. Ing.*, 1965, **15**, 969-976

This oxidant-fuel system was evaluated as a rocket propellant combination.

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Non-metals

Mellor, 1940, Vol. 8, 109, 433, 435

Amorphous (not crystalline) boron reacts with brilliant flashes at ambient temperature, and charcoal or phosphorus continue to burn more brilliantly than in air (which has a much lower oxygen content).

Ozone

See Ozone: Nitrogen oxide

Pentacarbonyliron

Manchot, W. *et al.*, *Ann.*, 1929, **470**, 275

Rapid heating to above 50°C in an autoclave caused an explosive reaction.

Perchloryl fluoride

See Perchloryl fluoride: Hydrocarbons, etc.

Phosphine, Oxygen

Mellor, 1940, Vol. 8, 435

Addition of oxygen to a mixture of phosphine and nitrogen oxide causes explosion.

See Phosphine: Oxygen

Potassium sulfide

Mellor, 1940, Vol. 8, 434

The pyrophoric sulfide ignites in the oxide.

Sodium diphenylketyl

1. Mendenhall, D., *J. Amer. Chem. Soc.*, 1974, **96**, 5000

2. Mendenhall, D., *Chem. Eng. News*, 1982, **60**(20), 2

A procedure for preparation of sodium hyponitrite involving reduction of nitrogen oxide by the ketyl, followed by extraction into water [1], had been operated routinely on the small scale. A 41-fold scaled up run exploded and ignited after 200 ml of water had been added as part of the work-up. This was attributed to the presence of an unusually large proportion of coagulated unreacted sodium in the ketyl. Small-scale operation with precautions is urged [2].

Vinyl chloride

See Vinyl chloride: Oxides of nitrogen

See other ENDOTHERMIC COMPOUNDS, NON-METAL OXIDES, OXIDANTS

4725. Nitrogen dioxide (Dinitrogen tetraoxide)

[10102-44-0]

NO₂

NO₂

At ambient temperatures the title gases exist as an equilibrium mixture; for details

See Dinitrogen tetraoxide

1783

Alkenes

Rozlovskii, A. I., *Chem. Abs.*, 1975, **83**, 118122

The mechanisms of explosions in solidified gas mixtures at low temperatures containing unsaturated hydrocarbons and oxides of nitrogen is discussed. Fast radical addition of nitrogen dioxide to double bonds is involved, and with dienes it is a fast reaction of very low energy of activation. Possibilities of preventing explosions are discussed.

See Nitrogen oxide: Dienes, Oxygen

See other NON-METAL OXIDES, OXIDANTS

4726. Plutonium nitride

[12033-54-4]

PuN

NPu

Oxygen, Water

Bailar, 1973, Vol. 5, 343

The ignition temperature of the nitride in oxygen is lowered from 300° to below 100°C by the presence of moisture.

See other NITRIDES

4727. Rubidium nitride

[12136-85-5]

Rb₃N

NRb₃

Mellor, 1940, Vol. 8, 99

The alkali nitrides burn in air.

See other N-METAL DERIVATIVES

4728. Poly(sulfur nitride)

[56422-03-8]

(Complex structure)

(NS)_n

Kennet, F. A. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1982, 851–857

Hazards involved in various routes to the polymer are detailed.

See Caesium azide: Sulfur dioxide, *also* Poly(selenium nitride)

See other NITRIDES, N-S COMPOUNDS

4729. Antimony(III) nitride

[12333-57-2]

SbN

NSb

Alone, or Water

1. Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1471

2. Franklin, E. C., *J. Amer. Chem. Soc.*, 1905, **27**, 850

Explosive decomposition occurs on warming under vacuum [1] and impure material explodes mildly on heating in air, or on contact with water or dilute acids.

See other NITRIDES

4730. Poly(selenium nitride)

[12033-59-9]

(Complex structure)

(NSe)_n

Preparative hazard

See Trimethylsilyl azide: Selenium halides

See also Poly(sulfur nitride) *See other* NITRIDES

See related N-S COMPOUNDS

4731. Thallium(I) nitride

[12033-67-9]

TlN

NTl

Alone, or Water

Mellor, 1940, Vol. 8, 262

The nitride explodes violently on exposure to shock, heat, water or dilute acids.

See other NITRIDES

4732. Uranium(III) nitride

[25658-43-9]

UN

NU

Bailar, 1973, Vol. 5, 340

Finely powdered material is pyrophoric on exposure to air at ambient temperature.

See other NITRIDES, PYROPHORIC MATERIALS

4733. Zirconium nitride

[25658-42-8]

ZrN

NZr

See entry REFRACTORY POWDERS *See other* NITRIDES

4734. Nitrogen (Gas)

[7727-37-9]

N₂

N₂

Kletz, T. A., *Proc. 3rd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 1519–1527, Basle, SSCI, 1980

The benefits of large scale use of nitrogen as an inerting gas to prevent fire or explosion have been at the expense of many fatal asphyxiations arising from its

accidental use in place of air or from its unexpected presence in confined spaces. At an oxygen concentration below 10%, collapse is almost instantaneous. Various accidents are described with appropriate precautions, and it was concluded that nitrogen may have killed more people than any other single substance in industry.

Lithium

See Lithium(reference 1), *also* Lithium: Metal chlorides

Lithium tetrahydroaluminate

See Lithium tetrahydroaluminate: Nitrogen

Ozone

See Ozone: Nitrogen

Titanium

See Titanium: Nitrogen

4735. Nitrogen (Liquid)

1. Anon., *BCISC Quart. Safety Summ.*, 1964, **35**, 25
2. Anon., *BCISC Quart. Safety Summ.*, 1974, **45**, 11
3. Lindsay, W. N., *Chem. Eng. News*, 1987, **65**(24), 2

Although liquid nitrogen is inherently safer than liquid oxygen or liquid air as a coolant, its ability to condense liquid oxygen out of the atmosphere can create hazards. A distillation tube containing a little solvent was cooled in liquid nitrogen while being sealed off in a blowpipe flame. A few minutes later the tube exploded, probably owing to high internal pressure caused by evaporation of liquid oxygen which had condensed into the tube during sealing. It is possible the atmosphere close to the blowpipe was oxygen-enriched. Open vessels which contain organic materials, or which are to be hermetically sealed should not be cooled in liquid nitrogen but in a coolant at a higher temperature. Liquid nitrogen should normally be used only for cooling evacuated or closed vessels where extreme cooling is necessary, and it should be removed from around the vacuum trap or other vessel before opening them to atmosphere [1]. The later reference describes 2 incidents of forceful explosion of 75 l cryogenic containers of liquid nitrogen. The cause of the first appeared to be blockage of the 19 mm neck by an ice-plug formed in a very humid environment. The cause of the second was probably 'roll-over' of a surface layer of superheated liquid, and fracture of the vessel by the ensuing violent boil-off of gas. Superheating of 10–15°C is known to occur in high-purity liquid nitrogen, but much higher levels of superheating may be possible. Various remedies (stirring, generation of a warm spot or vertical submerged conductive gauzes) are discussed [2]. With the current heightened interest in new materials which become superconducting at liquid nitrogen temperature (–195.8°C), the attention of workers (not necessarily chemists) in this area is drawn to the fact that containers of liquid nitrogen exposed to atmosphere inevitably will condense some oxygen (b.p. –183°C) into the nitrogen coolant. Such mixtures may behave

as an oxidant and react violently with traces of oil, grease or other organic material. Vacuum flasks of liquid nitrogen should not be left about unused or uncovered [3].

Argon

See Argon: Liquid nitrogen

Fatty materials

Anon., *CISHC Chem Safety Summ.*, 1979, **50**, 91

Use of liquid nitrogen in cryogenic grinding of fatty materials led to an explosion. Condensation of liquid oxygen onto the fatty material, with initiation by the grinding friction seems a likely causative sequence.

Hydrogen

See Hydrogen (Gas): Liquid nitrogen

Magnesium

Ephraim, 1939, 625

A mixture of magnesium powder and liquid nitrogen reacts very violently when lit with a fuse, forming magnesium nitride.

Oxygen, Radiation

1. Chen, C. W. *et al.*, *Cryogenics*, 1969, **9**, 131–132

2. Takehisa, M. *et al.*, *Chem. Abs.*, 1977, **87**, 30593

3. Perdue, P. T., *Health Phys.*, 1972, **23**, 116–117

4. Watanabe, H. *et al.*, *Chem. Abs.*, 1980, **92**, 30294

Liquid nitrogen subject to nuclear radiation (high neutron and gamma fluxes) must be kept free of oxygen to prevent explosions occurring in reactor cryostats. The explosive species generated is not nitrogen oxide or nitrogen dioxide, because solutions of these will not cause explosion on contact with a drop of acrylonitrile, whereas irradiated oxygen-containing nitrogen does so. Ozone was thought to be responsible [1], and this was confirmed by a detailed experimental investigation [2], though trace organic impurities may also have been involved [1]. The subject has been reviewed, and oxygen must also be eliminated from liquid helium, hydrogen or noble gases before irradiation [3]. Similar effects and conclusions were associated with an explosion which occurred when liquid nitrogen was used to cool a target during electron-beam irradiation [4].

See other CRYOGENIC LIQUIDS, IRRADIATION DECOMPOSITION INCIDENTS

4736. Sodium hyponitrite

[13517-64-7]

$\text{N}_2\text{Na}_2\text{O}_2$

NaON:NONa

Preparative hazard

See Nitrogen oxide: Sodium diphenylketyl

See Hyponitrous acid, *also* Sodium amide (reference 4)

- 4737. Sodium trioxodinitrate**
 [13826-64-7] **N₂Na₂O₃**

$$\text{NaON(O):NONa}$$

See Sodium amide(reference 4)
See other METAL NITRATES
- 4738. Sodium tetraoxodinitrate**
 [] **N₂Na₂O₄**

$$\text{NaON(O):N(O)ONa}$$

See Sodium amide (reference 4)
See other METAL NITRATES
- 4739. Sodium pentaoxodinitrate**
 [59795-18-5] **N₂Na₂O₅**

$$\text{NaON(O}_2\text{):N(O)ONa}$$

See Sodium amide(reference 4)
See other METAL NITRATES
- 4740. Sodium hexaoxodinitrate**
 [] **N₂Na₂O₆**

$$\text{NaON(O}_2\text{):N(O}_2\text{)ONa}$$

See Sodium amide(reference 4)
See other METAL NITRATES
- 4741. Dinitrosylnickel**
 [] **N₂NiO₂**

$$\text{ONNiNO}$$

 Mond, R. L. *et al.*, *J. Chem. Soc.*, 1922, **121**, 38
 A blue, impure and probably polymeric solid, produced from reaction of nitric oxide and nickel carbonyl, decomposed with incandescence at 90°C. The structure is very doubtful but a dinitrosyl was tentatively postulated. A trinitrosyl, [115380-62-6], has been listed recently.
See other HEAVY METAL DERIVATIVES, NITROSO COMPOUNDS
- 4742. Nitritonitrosylnickel**
 [] **N₂NiO₃**

$$\text{O:NNiON:O}$$

 Water
 Bailar, 1973, Vol. 3, 1119
 1788


It is probably polymeric and ignites on contact with water.

See other NITROSO COMPOUNDS

See related METAL NITRITES

4743. Nickel(II) nitrate

[13138-45-9]



N_2NiO_6

Oxidant. The tetrammine and tetrahydrazine complexes are explosive.

See AMMINOMETAL NITRATES, METAL NITRATES: organic matter

1-Methylbenzotriazole

Diamantopoulou, E. *et al.*, *Polyhedron*, 1994, **13**(10), 1593

A study of the 1-methylbenzotriazole complexes, of various stoichiometry and hydration, showed several to be explosive.

See other METAL NITRATES, OXIDANTS

4744. Dinitrogen oxide ('Nitrous oxide')

[10204-97-2]



N_2O

FPA H111, 1982 (cylinder); *HCS 1980*, 692 (cylinder)

1. Rüst, 1948, 278

2. Conrad, D. *et al.*, *Chem. Abs.*, 1983, **99**, 55875

3. Editorial feature, *Reading Times* (PA), 14th May, 1987

4. Bretherick, L., *Chem. Eng. News*, 1987, **65**(31), 3

5. Hamsen, B. *et al.*, *Chem. Eng. News*, 1991, **69**(29), 2

During transfer of the liquefied gas from a stock steel cylinder into smaller cylinders, the effective expansion caused cooling of the stock cylinder and a fall in pressure to occur. Application of a flame to the stock cylinder of the endothermic oxide led to decomposition and explosive rupture of the cylinder [1]. The explosive decomposition of the gas at 0.05–8 bar at ambient temperature after initiation by electric discharge can be prevented by addition of 30 vol% of air, nitrogen or oxygen [2]. Oxyacetylene welding repair work on or near to a 6 t tank of liquefied oxide led to a violent explosion [3], effectively a large scale rerun 85 years after the original incident described above [4]. When using nitrous oxide, containing organic solutes, as a supercritical fluid for hplc purposes, a 1 ml sample exploded, causing considerable shrapnel damage. It is recommended that it be not used as a supercritical fluid [5].

It is highly endothermic (ΔH_f° (g) +90.4 kJ/mol, 2.01 kJ/g) and an active oxidant.

See Fuels, below

See other ENDOTHERMIC COMPOUNDS

See also INDIGESTION

Preparative hazard

Bailar, 1973, Vol. 2, 317

Acetylene

See ATOMIC ABSORPTION SPECTROSCOPY

Boron

Mellor, 1940, Vol. 8, 109

Amorphous boron (not crystalline) ignites on heating in the dry oxide.

Carbon monoxide

Wilton, C. *et al.*, *Chem. Abs.*, 1975, **82**, 113724

Possible blast hazards associated with use of the propellant combination liquid dinitrogen oxide–liquid carbon monoxide have been evaluated.

Combustible gases

Sorbe, 1968, 131

The oxide (with an oxygen content 1.7 times that of air) forms explosive mixtures with ammonia, carbon monoxide, hydrogen, hydrogen sulfide and phosphine.

Fuels

1. L. Bretherick's comments

2. Young, J. A., *CHASNotes*, 1993, **XI**(6), 2

Although the oxygen in dinitrogen oxide is chemically (but endothermically) bound to nitrogen, it is present in a much higher concentration (36.4%) than the 21% present in air in admixture with nitrogen, so combustion or oxidation in the oxide will be much faster than in air. Advantage of this has been taken by feeding the oxide together with air into carburettor intakes to boost internal combustion engine performance, but strict control is necessary to prevent premature ignition. Nitrogen oxide, though of even higher oxygen content (53.3%), is too reactive and corrosive for this application [1].

A man habitually enjoyed the euphoric effects of inhaling whiffs of nitrous oxide in seclusion, and kept a cylinder of the gas in his sedan for that purpose. He decided to spray the faded car seats with an aerosol can of vinyl dressing (propane/butane propellant) with the windows closed. Then he had a whiff of gas from the briefly opened cylinder, and settled back to enjoy the euphoria and a cigarette. He was lucky to survive the resulting explosion of the fuel/oxidant mixture in a closed vessel [2].

See OXYGEN ENRICHMENT

Hydrazine

Mellor, 1967, Vol. 8, Suppl. 2.2, 214

Ignition occurs on contact.

See other REDOX REACTIONS

Hydrogen, Nitric oxide

See Nitrogen oxide: Hydrogen, etc.

Hydrogen, Oxygen

Bailar, 1973, Vol. 2, 321–322

The ignition temperature of mixtures of hydrogen with dinitrogen oxide is lower than that of hydrogen admixed with air or oxygen. The oxide also sensitises mixtures of hydrogen and oxygen, so that addition of oxygen to a hydrogen–oxide mixture will cause instantaneous ignition or explosion. Explosive limits are extremely wide.

Lithium hydride

Mellor, 1967, Vol. 8, Suppl. 2.2, 214

The hydride ignites in the gas.

Phosphine

Thénard, J., *Compt. rend.*, 1844, **18**, 652

A mixture with excess oxide can be exploded by sparking.

Plastic tubes

See Oxygen (Gas): Plastic tubes

Silane

See Silane: Nitrogen oxides

Tin(II) oxide

See Tin(II) oxide: Non-metal oxides

Tungsten carbides

See Tungsten carbide: Nitrogen oxides

Ditungsten carbide: Oxidants

See other NON-METAL OXIDES, OXIDANTS

4745. Lead hyponitrite

[19423-89-3]

N₂O₂Pb

(-PbON:NO-) _n

Partington, J. R. *et al.*, *J. Chem. Soc.*, 1932, 135, 2589

The lead salt decomposes with explosive violence at 150–160°C and should not be dried by heating. The salt prepared from lead acetate explodes more violently than that from the nitrate.

See other HEAVY METAL DERIVATIVES, N–O COMPOUNDS

4746. Dinitrogen trioxide

[10544-73-7]

N₂O₃

O:NON:O

Acetic acid, 6-Hexanelactam

See 6-Hexanelactam: Acetic acid, etc.

Chloroprene

See 2-Chloro-1-nitro-4-nitroso-2-butene

Phosphine

See Phosphine (reference 3)

Phosphorus

See Phosphorus: Non-metal oxides

See other NON-METAL OXIDES, OXIDANTS

4747. Dinitrogen tetraoxide (Nitrogen dioxide)

[10544-72-6]

N₂O₄

N₂O₄

HCS 1980, 675 (cylinder)

The equilibrium mixture of nitrogen dioxide and dinitrogen tetraoxide is completely associated at -9°C to the latter form which is marginally endothermic (ΔH_f° (g) $+9.7$ kJ/mol, 0.10 kJ/g). Above 140°C it is completely dissociated to nitrogen dioxide, which is moderately endothermic (ΔH_f° (g) $+33.8$ kJ/mol, 0.74 kJ/g).

Acetonitrile, Indium

MRH Acetonitrile 7.87/25

Addison, C. C. *et al.*, *Chem. & Ind.*, 1958, 1004

Shaking a slow-reacting mixture caused detonation, attributed to indium-catalysed oxidation of acetonitrile.

Alcohols

Daniels, F., *Chem. Eng. News*, 1955, **33**, 2372

A violent explosion occurred during the ready interaction to produce alkyl nitrates.

Ammonia

MRH 6.61/33

Mellor, 1940, Vol. 8, 541

Liquid ammonia reacts explosively with the solid tetraoxide at -80°C , while aqueous ammonia reacts vigorously with the gas at ambient temperature.

Barium oxide

Mellor, 1940, Vol. 8, 545

In contact with the gas at 200°C the oxide suddenly reacts, reaches red heat and melts.

Boron trichloride

Mellor, 1946, Vol. 5, 132

Interaction is energetic.

Carbon disulfide

1. Mellor, 1940, Vol. 8, 543

2. Sorbe, 1968, 132

Liquid mixtures proposed for use as explosives are stable up to 200°C [1], but can be detonated by mercury fulminate, and the vapours by sparking [2].

Carbonylmetals

Cloyd, 1965, 74
Combination is hypergolic.

Cellulose, Magnesium perchlorate

See Magnesium perchlorate: Cellulose, etc.

Cycloalkenes, Oxygen

Lachowicz, D. R. *et al.*, US Pat. 3 621 050, 1971
Contact of cycloalkenes with a mixture of dinitrogen tetroxide and excess oxygen at temperatures of 0°C or below produces nitroperoxonitrates of the general formula—CHNO₂—CH(OONO₂)— which appear to be unstable at temperatures above 0°C, owing to the presence of the peroxonitrate group.
See Hydrocarbons, below

Difluorotrifluoromethylphosphine

Mahler, W., *Inorg. Chem.*, 1979, **18**, 352
A reaction, to produce the phosphine oxide on 12 mmol scale, ignited.

Dimethyl sulfoxide

MRH 6.99/36

See Dimethyl sulfoxide: Dinitrogen tetroxide

Formaldehyde

1. Pollard, F. H. *et al.*, *Trans. Faraday Soc.*, 1949, **45**, 767–770
2. Rastogi, R. P. *et al.*, *Chem. Abs.*, 1975, **83**, 12936
The slow (redox) reaction becomes explosive around 180°C [1], or even lower [2].
See other REDOX REACTIONS

Halocarbons

MRH Chloroform 2.38/67, 1,2-dichloroethane 5.06/42,
1,1-dichloroethylene 5.06/46, trichloroethylene 3.97/56

1. Turley, R. E., *Chem. Eng. News*, 1964, **42**(47), 53
2. Benson, S. W., *Chem. Eng. News*, 1964, **42**(51), 4
3. Shanley, E. S., *Chem. Eng. News*, 1964, **42**(52), 5
4. Kuchta, J. M. *et al.*, *J. Chem. Eng. Data*, 1968, **13**, 421–428
Mixtures of the tetroxide with dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, trichloroethylene and tetrachloroethylene are explosive when subjected to shock of 25 g TNT equivalent or less [1]. Mixtures with trichloroethylene react violently on heating to 150°C [2]. Partially fluorinated chloroalkanes were more stable to shock. Theoretical aspects are discussed in the later reference [2,3]. The effect of pressure on flammability limits has been studied [4].
See Vinyl chloride: Oxides of nitrogen
See Uranium: Nitric acid

Heterocyclic bases MRH Pyridine 7.82/22, quinoline 7.87/22

Mellor, 1940, Vol. 8, 543

Pyridine and quinoline are attacked violently by the liquid oxide.

Hydrazine derivatives

1. Cloyd, 1965, 74

2. Miyajima, H. *et al.*, *Combust. Sci. Technol.*, 1973, **8**, 199–200

Combinations with hydrazine, methylhydrazine, 1,1-dimethylhydrazine or mixtures thereof are hypergolic and used in rocketry [1]. The hypergolic gas-phase ignition of hydrazine at 70–160°C/53–120 mbar has been studied [2].

See ROCKET PROPELLANTS

Hydrocarbons

MRH values below references

1. Mellor, 1967, Vol. 8, Suppl. 2.2, 264

2. Fierz, H. E., *J. Soc. Chem. Ind.*, 1922, **41**, 114R

3. Raschig, F., *Z. Angew. Chem.*, 1922, **35**, 117–119

4. Berl, E. *Z. Angew. Chem.*, 1923, **36**, 87–91

5. Schaarschmidt, A., *Z. Angew. Chem.*, 1923, **36**, 533–536

6. Berl, E., *Z. Angew. Chem.*, 1924, **37**, 164–165

7. Schaarschmidt, A., *Z. Angew. Chem.*, 1925, **38**, 537–541

8. *MCA Case History No. 128*

9. Folecki, J. *et al.*, *Chem. & Ind.*, 1967, 1424

10. Cloyd, 1965, 74

11. Urbanski, 1967, Vol. 3, 289

12. Biasutti, 1981, 50

13. Biasutti, 1981, 53–54

MRH Benzene 7.99/19, hexane 7.91/17, isoprene 8.28/18, methylcyclohexane 7.87/17

A mixture of the tetroxide and toluene exploded, possibly initiated by unsaturated impurities [1]. During attempted separation by low temperature distillation of an accidental mixture of light petroleum and the oxide, a large bulk of material awaiting distillation became heated by unusual climatic conditions to 50°C and exploded violently [2]. Subsequently, discussion of possible alternative causes involving either unsaturated or aromatic compounds was published [3] [4] [5] [6] [7]. Erroneous addition of liquid in place of gaseous nitrogen tetroxide to hot cyclohexane caused an explosion [8]. During kinetic studies, one sample of a 1:1 molar solution of tetroxide in hexane exploded during (normally slow) decomposition at 28°C [9]. Cyclopentadiene is hypergolic with the oxide [10]. These incidents are understandable because of their similarity to rocket propellant systems and liquid mixtures previously used as bomb fillings [11]. The liquid oxide leaking from a ruptured 6 t storage tank ran into a gutter containing toluene and a violent explosion ensued [12]. An alternative account describes the hydrocarbon as benzene [13].

See Cycloalkenes, above; Unsaturated hydrocarbons, below

Hydrogen, Oxygen

Lewis, B., *Chem. Rev.*, 1932, **10**, 60

N₂O₄

The presence of small amounts of the oxide in non-explosive mixtures of hydrogen and oxygen renders them explosive.

Isopropyl nitrite, Propyl nitrite

Safety in the Chemical Laboratory, Vol. 1, 121, Steere, N. V. (Ed.), Easton (Pa.) *J. Ch. Ed.*, 1967

A pressurised mixture of the cold components exploded very violently during a combustion test run. The mixture was known to be autoexplosive at ambient temperature, and both of the organic components are capable of violent decomposition in absence of added oxidant.

Laboratory grease

Arapava, L. D. *et al.*, *Chem. Abs.*, 1985, **102**, 169310

Contact of the lubricating grease Litol-24 with the oxidant at below 80°C led to explosion on subsequent impact. This involved nitration products of the antioxidant present, 4-hydroxydiphenylamine. Above 80°C decomposition superceded nitration, and no explosion occurred.

See other NITRATION INCIDENTS

Metal acetylides or carbides

MRH values show % of oxidant

Mellor, 1946, Vol. 5, 849

Caesium acetylide ignites at 100°C in the gas.

See Tungsten carbide: Nitrogen oxides

MRH 4.02/63

Ditungsten carbide: Oxidants

MRH 3.85/67

Metals

MRH Magnesium 12.97/50, potassium 3.72/46

1. Mellor, 1940, Vol. 8, 544–545; 1942, Vol. 13, 342

2. Pascal, 1956, Vol. 10, 382; 1958, Vol. 4, 291

Reduced iron, potassium and pyrophoric manganese all ignite in the gas at ambient temperature. Magnesium filings burn vigorously when heated in the gas [1]. Slightly warm sodium ignites in contact with the gas, and interaction with calcium is explosive [2].

See Aluminium: Oxidants

Nitroaniline

Anon., *CISHC Chem. Safety Summ.*, 1978, **49**, 3–4

Process errors led to discharge of copious amounts of nitrous fumes into the glass reinforced plastic ventilation duct above a diazotisation vessel. On two occasions fires were caused in the duct by vigorous reaction of the dinitrogen tetroxide with nitroaniline dusts in the duct. Laboratory tests confirmed this to be the cause of the fires, and precautions are detailed.

Nitroaromatics

1. Urbanski, 1967, Vol. 3, 288

2. Kristoff, F. T. *et al.*, *J. Haz. Mat.*, 1983, **7**, 199–210

Mixtures with nitrobenzene were formerly used as liquid high explosives, with addition of carbon disulfide to lower the freezing point, but high sensitivity to mechanical stimulus was disadvantageous [1]. During the recovery of acids from nitration of toluene, mixtures of the oxide with nitrotoluene or dinitrotoluene may

be isolated under certain process conditions. While such mixtures are not unduly sensitive to impact, friction or thermal initiation, when oxygen-balanced they are extremely sensitive to induced shock and are capable of explosive propagation at film thicknesses below 0.5 mm. It is suspected that many explosions in TNT acid recovery operations, previously attributed to tetranitromethane, may have been caused by such mixtures [2].

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Organic compounds

Riebsomer, J. L., *Chem. Rev.*, 1945, **36**, 158

In a review of the interaction of the oxidant with organic compounds, attention is drawn to the possibility of formation of unstable or explosive products.

Other reactants

Yoshida, 1980, 269

MRH values calculated for 18 combinations with oxidisable materials are given.

Ozone

See Ozone: Nitrogen oxide

Phospham

See Phospham: Oxidants

Phosphorus

MRH 9.12/35

See Phosphorus: Non-metal oxides

Sodium amide

Beck, G., *Z. Anorg. Chem.*, 1937, **233**, 158

Interaction with the oxide in carbon tetrachloride is vigorous, producing sparks.

Tetracarbonylnickel

Bailar, 1973, Vol. 3, 1130

Interaction of the liquids is rather violent.

See Carbonylmetals, above

Tetramethyltin

Bailar, 1973, Vol. 2, 355

Interaction is explosively violent even at -80°C , and dilution with inert solvents is required for moderation.

2-Toluidinium nitrate

Rastogi, R. P. *et al.*, *Indian J. Chem., Sect. A*, 1980, **19** A, 317–321

Reaction in this hybrid rocket propellant system is enhanced by presence of ammonium vanadate.

Triethylamine

Davenport, D. A. *et al.*, *J. Amer. Chem. Soc.*, 1953, **75**, 4175

The complex, containing excess oxide over amine, exploded at below 0°C when free of solvent.

Triethylammonium nitrate

Addison, C. C. *et al.*, *Chem. & Ind.*, 1953, 1315

The two component form an addition complex with diethyl ether, which exploded violently after partial desiccation: an ether-free complex is also unstable.

See Triethylamine, above

Unsaturated hydrocarbons

MRH Isoprene 8.28/18

1. Sergeev, G. P. *et al.*, *Chem. Abs.*, 1966, **65**, 3659g

2. Biasutti, 1981, 123

Dinitrogen tetroxide reacts explosively between -32° and -90°C with propene, 1-butene, isobutene, 1,3-butadiene, cyclopentadiene and 1-hexene, but 6 other unsaturates failed to react [1]. Reaction of propene with the oxide at 2 bar/ 30°C to give lactic acid nitrate was proceeding in a pump-fed tubular reactor pilot plant. A violent explosion after several hours of steady operation was later ascribed to an overheated pump gland which recently had been tightened. A similar pump with a tight gland created a hot-spot at 200°C [2].

See Nitrogen dioxide: Alkenes

Vinyl chloride

See Vinyl chloride: Oxides of nitrogen

Xenon tetrafluoride oxide

Christe, K. O., *Inorg. Chem.*, 1988, **27**, 3764

In the reaction of the pentaoxide with xenon tetrafluoride oxide to give xenon difluoride dioxide and nitryl fluoride, the xenon tetrafluoride oxide must be used in excess to avoid formation of xenon trioxide, which forms a sensitive explosive mixture with xenon difluoride dioxide.

See Xenon tetrafluoride oxide: Caesium nitrate

See other ENDOTHERMIC COMPOUNDS, NON-METAL OXIDES, OXIDANTS

4748. Dinitrogen pentaoxide (Nitryl nitrate)

[10102-03-1]

N_2O_5

NO_2+NO_3-

Partington, 1967, 569

If suddenly heated, the solid (m.p. 30°C) explodes. The structure is now considered to be nitryl nitrate.

Acetaldehyde

Kacmarek, A. J. *et al.*, *J. Org. Chem.*, 1975, **40**, 1853

Direct combination to produce ethylidene dinitrate at -196°C is violently explosive, but uneventful when the acetaldehyde is diluted with nitrogen.

Aromatic compounds

Editor's comments, 1999

Increased availability of this compound means it is finding use as a clean and mild nitrating agent. A stoichiometric byproduct in many procedures is anhydrous nitric acid, which is often blithely ignored. Anhydrous nitric acid is a very hazardous

material in its own right, being surprisingly unreactive—except as an oxidant during detonations! In particular, work-up by distillation is virtually certain to form detonable compositions in the condenser at some stage. The editor is not yet aware of any accidents during such distillation, but counsels forethought.

See Nitric acid

tert-Butylhydroperoxide, Toluene

See tert-Butylhydroperoxide: Toluene, Dinitrogen pentaoxide

Dichloromethane

See NITRATING AGENTS

Ethylene

Nitro Compounds, Feuer H. & Nielsen A. T. (Eds), New York, VCH, 1990, 315

The crude product is too explosive to be worked up. Analogy with other olefins indicates that the final product will be ethylene glycol dinitrate, a known explosive though safe enough to have largely replaced nitroglycerine. That will be preceded by 1,2-dinitroethane and nitratonitroethane, more sensitive if less powerful.

Metals

Mellor, 1940, Vol. 8, 554

Potassium and sodium burn brilliantly in the gas, while mercury and arsenic are vigorously oxidised.

Naphthalene

Mellor, 1940, Vol. 8, 554

Naphthalene explodes and other organic materials react vigorously with the powerful oxidant.

Sodium acetylide

See Sodium acetylide: Oxidants

Strained ring heterocycles

Golding, P. *et al.*, *Tetrahedron Lett.*, 1988, **29**, 2731–2734; 2735–2736

Strained ring oxygen heterocycles such as epoxides and oxetanes react with the oxide in dichloromethane at sub-ambient temperatures to form high yields of the powerfully explosive glycol dinitrate esters. Epoxides give 1,2-dinitrates (such as those from ethylene, propene, butene and 4 other oxides), while oxetanes (6 examples) give 1,3-dinitrates. Strained ring nitrogen heterocycles give explosive *N*-nitroaminoalkyl nitrates, and several derived from aziridines and 3 derived from azetidines are exemplified.

See other STRAINED-RING COMPOUNDS

See NITRATING AGENTS

Sulfur dichloride, or Sulfuryl chloride

Schmeisser, M., *Angew. Chem.*, 1955, **67**, 495, 499

Interaction is explosively violent.

See other NON-METAL OXIDES, OXIDANTS

See related ACID ANHYDRIDES

4749. Lead(II) nitrate
[10099-74-8]



HCS 1980, 589

Carbon

MRH 1.76/7

Mellor, 1941, Vol. 7, 863

Contact with red hot carbon causes an explosion with showers of sparks.

Calcium–silicon alloy

Smolin, A. O., *Chem. Abs.*, 1975, **85**, 7591

‘Combustion of Calcium–Silicon in a Mixture with Lead Nitrate’ (title only translated)

Cyclopentadienylsodium

Houben-Weyl, 1975, Vol. 13.3, 200

If lead nitrate is used rather than the chloride or acetate as the source of divalent lead, the crude dicyclopentadienyllead may explode violently during purification by high-vacuum sublimation at 100–130°C.

Other reactants

Yoshida, 1980, 200

MRH values for 17 combinations with oxidisable materials are given.


Potassium acetate

Mee, A. J., *School Sci. Rev.*, 1940, **28**(86), 95

A heated mixture of the oxidant and organic salt exploded violently.

See other METAL NITRATES, OXIDANTS

4750. Tin(II) nitrate
[22755-27-7]



Bailar, 1973, Vol. 2, 74

During attempted isolation of the nitrate by evaporation of its aqueous solutions, a number of explosions have occurred.

See other METAL NITRATES, REDOX COMPOUNDS

4751. Zinc nitrate
[7779-88-6]



HCS 1980, 967

Carbon

Mellor, 1940, Vol. 4, 655

When the nitrate is sprinkled on to hot carbon, an explosion occurs.

See other METAL NITRATES, OXIDANTS

4752. Uranyl nitrate
[10102-06-4]



Cellulose

Clinton, T. G., *J. R. Inst. Chem.*, 1958, **82**, 633

The analytical use of cellulose fibre to absorb uranyl nitrate solution prior to ignition has led to explosions during ignition, owing to formation of cellulose nitrate. An alternative method is described.

Diethyl ether

1. Ivanov, W. N., *Chem. Ztg.*, 1912, **36**, 297

2. Andrews, L. W., *J. Amer. Chem. Soc.*, 1912, **34**, 1686–1687

3. Müller, A., *Chem. Ztg.*, 1916, **40**, 38; 1917, **41**, 439

The mild detonations reported when the crystalline salt was disturbed [1] were thought to have been caused by presence of solvent ether in the crystals (6 mols of water may be replaced by 2 of ether) [2]. This was later confirmed [3], and the formation of ethyl nitrate or diethyloxonium nitrate may have been involved, as the anhydrous salt functions as a powerful nitrator. Solutions of the nitrate in ether should not be exposed to sunlight to avoid the possibility of explosions.

See Organic solvents, below

See other IRRADIATION DECOMPOSITION INCIDENTS

Ethylene glycol

See Ethylene glycol: Oxidants

Organic solvents

Burriss, W. H., *Nucl. Sci. Abs.*, 1976, **33**, 19790

During routine operations to reduce the hexahydrate to uranium trioxide, an excessive amount of organic solution entered the denitrator pots and ignited.

See Diethyl ether, above

See also NITRATING AGENTS

See other METAL NITRATES

4753. Lead nitride
[58572-21-7]



Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1470

Very unstable, it decomposes explosively during vacuum degassing.

See other NITRIDES

4754. Disulfur dinitride
[25474-92-4]



1. Brauer, 1963, Vol. 12, 410
2. Williams, J. D., *Polyhedron*, 1987, **6**(5), 939–941

Explosive, initiated by shock, friction, pressure or temperatures over 30°C [1]. It may also detonate spontaneously at ambient temperature [2].


Metal chlorides

Roesky, H. W. *et al.*, *Inorg. Chem.*, 1984, **23**, 75

The 1:1 complexes with beryllium chloride or titanium tetrachloride may explode violently.

See other NITRIDES, N–S COMPOUNDS

4755. Tetrasulfur dinitride
[32607-15-1]




1. Brauer, 1963, Vol. 1, 408
2. Heal, H. G. *et al.*, *J. Inorg. Nucl. Chem.*, 1975, **37**, 286

It decomposes explosively to its elements at 100°C. Previously prepared from the explosive tetranitride [1], it may now safely be prepared from the stable intermediate, heptasulfur imide [2].

See other NITRIDES, N–S COMPOUNDS

4756. Tetraselenium dinitride
[132724-39-1]



Dehnicke, K. *et al.*, *Angew. Chem. (Int.)*, 1991, **30**(5), 577

Made from diselenium dichloride and trimethylsilyl azide, the crystalline nitride explodes under a hammer blow, or on heating to 80°C. Can apparently be handled safely at room temperature.

See related N–S COMPOUNDS *See other* NITRIDES

4757. Poly(disilicon nitride)
[12438-96-9]



(Complex Structure with Si-Si bonds)

Alone, or Water

Gmelin, 1959, Vol. 15.B, 610

It ignites in air, explodes on heating and contact with drops of water generates sparks.

See other NITRIDES

4758. Sodium azide
[26628-22-8]

N₃Na

NaN₃

HCS 1980, 827

1. Mellor, 1940, Vol. 8, 345
2. Lambert, B., *School Sci. Rev.*, 1927, **8**(31), 218
3. Anon., *Jahresber.*, 1981, 77–79
4. Anon., *Safety*, 1982, (3), 2
5. Grewer, T. *et al.*, *Exothermic Decomposition*, Technical Report 01VD 159/0329 for Federal German Ministry for Res. Technol., Bonn. 1986

Insensitive to impact, it decomposes, sometimes explosively, above its m.p. [1], particularly if heated rapidly [2]. Although used in aqueous solutions as a preservative in pharmaceutical preparations, application of freeze-drying techniques to such solutions has led to problems arising from volatilisation of traces of hydrazoic acid from non-neutral solutions, condensation in metal lines, traps or filters, and formation of heavy metal azides in contact with lead, copper or zinc components in the drying plant [3,4].

Energy of exothermic decomposition in range 230–260°C was measured as 0.76 kJ/g by DSC, and T_{ait24} was determined as 253°C by adiabatic Dewar tests, with an apparent energy of activation of 145 kJ/mol [5].

See Heavy metals, below

Acids

See Hydrogen azide

Ammonium chloride, Trichloroacetonitrile

See 5-Trichloromethyltetrazole

Barium carbonate

Henneburg, G. O. *et al.*, *Can. J. Res.*, 1950, **28B**, 345

Interaction to form cyanide ion requires careful control of temperature at 630°C to prevent explosions.

Bromine

See Bromine: Metal azides

Carbon disulfide

Stohlmeier, M. *et al.*, *Chem. Ind. Dig.*, 1993, **6**(2), 124

A review on all the more or less explosive products which may be prepared from the above, with advice on safe handling and preparation.

See also Carbon disulfide: Metal azides

Carbonyl dichloride (Phosgene)

See tert-Butyl azidoformate (reference 4)

Chloroform

See Dichloromethane: Quaternary ammonium azides, etc.

Cyanuric chloride

See 2,4,6-Triazido-1,3,5-triazine (reference 2)

Chromyl chloride

See Chromyl chloride: Sodium azide

Dichloromethane, Dimethyl sulfoxide

1. Peet, N. P. *et al.*, *Chem. Eng. News*, 1993, **71**(16), 4

2. Hruby, V. J. *et al.*, *Chem. Eng. News*, 1993, **71**(41), 2

An explosion was experienced during work up of an epoxide opening reaction involving acidified sodium azide in a dichloromethane/dimethyl sulfoxide solvent. The author ascribes this to diazidomethane formation from dichloromethane [1]. A second report of an analogous accident, also attributed to diazidomethane, almost certainly involved hydrogen azide for the cold traps of a vacuum pump on a rotary evaporator were involved; this implies an explosive more volatile than dichloromethane. It is recommended that halogenated solvents be not used for azide reactions [2].

See Dichloromethane: Quaternary ammonium azides, etc.

2,5-Dinitro-3-methylbenzoic acid, Oleum

See 2,5-Dinitro-3-methylbenzoic acid: Oleum, Sodium azide

Heavy metals

1. Wear, J. O., *J. Chem. Educ.*, 1975, **52**, A23–25

2. Becher, H. H., *Naturwiss.*, 1970, **57**, 671

3. Anon., *Chem. Eng. News*, 1976, **54**(36), 6

4. Pobiner, H., *Chem. Eng. News*, 1982, **60**(125), 4

5. Anon., *Safety Digest Univ. Safety Assoc.*, 1992, **44**, 22

6. Kurtz, L., Internet, 1995

7. Mellor, 1967, Vol. 8, Suppl. 2, 4

The effluent from automatic blood analysers in which 0.01–0.1% sodium azide solutions are used, may lead over several months to formation of explosive heavy metal azides in brass, copper or lead plumbing lines, especially if acids are also present. Several incidents during drain-maintenance are described, and preventative flushing and decontamination procedures are discussed [1]. Brass plates, exposed to sodium azide solution during several months in soil-percolation tests and then dried, caused explosions owing to formation of copper and/or zinc azides [2]. During repairs to a metal thermostat bath in which sodium azide had been used as a preservative, a violent explosion occurred [3]. Use of sodium azide in automatic sulfur titrators led to explosions arising from formation of copper azide in a copper/brass valve assembly [4]. Mixing sodium azide solutions with heavy metal salts will produce the metal azides with greater facility; an explosion with mercuric chloride is reported [5]. A similar incident is reported later, apparently

the mixture has been recommended as a biocide for aqueous solutions. In this case an explosion occurred while stirring water containing perhaps 5% of each [6]. Raney nickel catalyses the vigorous decomposition of solutions of hydrogen azide or azide salts [7].

Manganese(III) salts, Styrene

Fristad, W. E. *et al.*, *J. Org. Chem.*, 1985, **50**, 3468

After applying an apparently safe procedure for diazidation of alkenes to styrene, an explosion occurred as the clear solution was cooling.

Nitryl fluoride

Klapötke, T. M. *et al.*, *Chem. Ber.*, 1994, **127**(11), 2181

Reaction with sodium azide even at -78°C leads to explosions attributed to fluorine azide formation.

See Fluorine azide

Sulfuric acid

Ross, F. F., *Water & Waste Treatment*, 1964, **9**, 528; private comm., 1966

One of the reagents required for the determination of dissolved oxygen in polluted water is a solution of sodium azide in 50% sulfuric acid. It is important that the diluted acid should be quite cold before adding the azide, since hydrogen azide boils at 36°C and is explosive in the condensed liquid state.

Trifluoroacetyl fluoride

See Trifluoroacetyl fluoride: Sodium azide

Water

Anon., *Angew. Chem.*, 1952, **64**, 169

Addition of water to sodium azide which had been heated strongly caused a violent reaction. This was attributed to formation of metallic sodium or sodium nitride.

See other METAL AZIDES

4759. Sodium azidosulfate

[67880-15-3]

$\text{N}_3\text{NaO}_3\text{S}$



Shozda, R. J. *et al.*, *J. Org. Chem.*, 1967, **32**, 2876


It is a weak explosive with variable sensitivity to mechanical shock and heating.

See other ACYL AZIDES

4760. Sodium trisulfurtrinitride

[65107-36-0] (ion)

N_3NaS_3



Bojes, J. *et al.*, *Inorg. Chem.*, 1978, **17**, 319

Like tetrasulfur tetranitride, salts of trisulfur trinitride and of pentasulfur tetranitride (particularly alkali-metal salts) are heat- and friction-sensitive explosives. Preparation on a scale limited to 1 g, and care in use of spatulae or in preparation of samples for IR examination is recommended.

See Sodium tetrasulfur pentanitride


See related NITRIDES

See other N-S COMPOUNDS

4761. Thallium 2,4,6-tris(dioxoselena)hexahydrotriazine-1,3-5-triide ('Thallium triselenimidate')

[]

$\text{N}_3\text{O}_6\text{Se}_3\text{Tl}_3$



Explosive.

See Selenium difluoride dioxide: Ammonia

See other N-METAL DERIVATIVES

4762. Thallium(III) nitrate

[13746-98-0]

$\text{Tl}(\text{NO}_3)_3$

$\text{N}_3\text{O}_9\text{Tl}$

Formic acid, 4-Hydroxy-3-methoxybenzaldehyde

491M, 1975, 416

Addition of a little of the aldehyde (vanillin) to a strong solution of the trihydrated nitrate in 90% formic acid led to a violent (redox) reaction.

See other METAL NITRATES, REDOX REACTIONS

4763. Vanadium trinitrate oxide

[16017-37-1]

$\text{VO}(\text{NO}_3)_3$

$\text{N}_3\text{O}_{10}\text{V}$

Organic materials

1. Schmeisser, M., *Angew. Chem.*, 1955, **67**, 495

2. Harris, A. D. *et al.*, *Inorg. Synth.*, 1967, **9**, 87



Many hydrocarbons and organic solvents ignite on contact with this powerful oxidant and nitrating agent [1], which reacts like fuming nitric acid with paper, rubber or wood [2].

See related METAL NITRATES

See other OXIDANTS

4764. Tris(thionitrosyl)thallium

[]



Bailar, 1973, Vol. 1, 1162

The compound and its adduct with ammonia explode very easily with heat or shock.

See other N-METAL DERIVATIVES, N-S COMPOUNDS**4765. Thallium(I) azide**

[13847-66-0]



Mellor, 1940, Vol. 8, 352

A relatively stable azide, it can be exploded on fairly heavy impact, or by heating at 350–400°C.

See other METAL AZIDES**4766. Nitrosyl azide**



[62316-46-5]

1. Lucien, H. W., *J. Amer. Chem. Soc.*, 1958, **80**, 44582. Klapötke, T. M., *et al.*, *Angew. Chem. (Int.)*, 1993, **32**(11), 16103. Tornieporth-Oetting, I. C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(5), 511

Explosions were experienced on several occasions during preparation of nitrosyl azide by various methods [1]. A yellow solid at –85°C, it decomposes even at –50°C [2]. It has an enthalpy of detonation of 6.3 kJ/g, the same as nitroglycerine, and a cavity volume greater than 600ml in the lead block test (TNT = 300; Nitroglycerine = 520 ml) [3].

See other HIGH-NITROGEN COMPOUNDS, NON-METAL AZIDES**4767. Thiotrithiazyl nitrate**

[79796-40-0] (ion)



Goehring, 1957, 74

It explodes on friction or impact.

See other N-S COMPOUNDS

4768. Plutonium(IV) nitrate
[13823-27-2]

N₄O₁₂Pu



MCA Case History No. 1498

Polythene bottles are not suitable for long-term storage of plutonium nitrate solutions as radiation-induced stress cracks appeared in the bases of several 10 l bottles during 6 months' storage. Short-term storage and improved venting are recommended.


See Americium trichloride

See other IRRADIATION DECOMPOSITION INCIDENTS, METAL NITRATES

4769. Diseleniumdisulfur tetranitride

[]

N₄S₂Se₂



Maaninen, A. *et al.*, *Phosphorus, Sulfur, Silicon Relat. Elem.*, 1997, (124–125), 457

Like the homologues below, this compound is explosive on heating or mechanical stress.


See other SULFUR, SELENIUM AND TELLURIUM TETRANITRIDES IN IMMEDIATELY SUBSEQUENT ENTRIES

See other N–S COMPOUNDS, NITRIDES

4770. Tetrasulfur tetranitride

[28950-34-7]

N₄S₄



1. Mellor, 1940, Vol. 8, 625
2. Brauer, 1963, Vol. 1, 406
3. Villema-Blanco, M. *et al.*, *Inorg. Synth.*, 1967, **9**, 101
4. Banister, A. J., *Inorg. Synth.*, 1977, **17**, 197–198
5. Witt, M. W. R. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 1668
6. Mark, H. *et al.*, *Chem. Eng. News*, 1984, **62**(5), 4
7. Buntain, G. A., *Chem. Eng. News*, 1988, **66**(18), 2

The endothermic nitride is susceptible to explosive decomposition on friction, shock or heating above 100°C [1]. Explosion is violent if initiated by a detonator [2]. Sensitivity toward heat and shock increases with purity. Preparative precautions have been detailed [3], and further improvements in safety procedures and handling described [4]. An improved plasma pyrolysis procedure to produce poly(sulfur nitride) films has been described [5]. Light crushing of a small sample of impure material (m.p. below 160°C, supposedly of relatively low sensitivity) prior to purification by sublimation led to a violent explosion [6] and a restatement of the need [4] for adequate precautions. Explosive sensitivity tests have shown it to be more sensitive to impact and friction than is lead azide, used in detonators. Spark-sensitivity is, however, relatively low [7].

Metal chlorides

Roesky, H. W. *et al.*, *Inorg. Chem.*, 1984, **23**, 75–76

The 1:1 complex with titanium chloride or the 1:2 complex with beryllium chloride may explode violently. Careful pyrolysis of these gives mixtures of disulfur dinitride and its 1:1 complexes with the metal chlorides, which may also explode, so screening is essential.

Oxidants

Pascal, 1956, Vol. 10, 645–646


Contact with fluorine leads to ignition, and interaction with metal chlorates or oxides is violent.

See other ENDOTHERMIC COMPOUNDS, NITRIDES, N-S COMPOUNDS

4771. Tetraselenium tetranitride

[12033-88-6]

N_4Se_4



Alone, or Halogens and derivatives

1. Mellor, 1947, Vol. 10, 789

2. Kelly, P. F. *et al.*, *Polyhedron*, 1990, **9**(13), 1567

The dry material explodes on slight compression, or on heating at 130–230°C. Contact with bromine, chlorine or a little fuming hydrochloric acid also causes explosion [1]. Explosions when manipulated with a metal spatula are reported [2].

See related N-S COMPOUNDS

See other NITRIDES

4772. Tritellurium tetranitride

[]

N_4Te_3

Structure unknown

Schmitz-Du Mont, O. *et al.*, *Angew. Chem. (Int.)*, 1967, **6**, 1071

1808

This yellow compound, prepared by ammonolysis of dipotassium triimidotellurite, was too explosive to be able to weigh it out for analysis.


See Tetratellurium tetranitride

See other NITRIDES

4773. Tetratellurium tetranitride

[12164-01-1]

N_4Te_4



1. Fischer, F. *et al.*, *Ber.*, 1910, **43**, 1472

2. Garcia-Fernandez, H., *Bull. Soc. Chim. Fr. [4]*, 1973, 1210–1215

The compound, previously formulated as tritellurium tetranitride is now shown to have the title structure (though probably polymeric, with inter-ring bonding). Two forms were originally described, one black which explodes on impact, and one yellow which explodes at 200°C [1]. It is explosive when dry, but may be stored safely under carbon tetrachloride [2].

See related N–S COMPOUNDS

See other NITRIDES

4774. Trithorium tetranitride

[12033-90-8]

N_4Th_3

Th_3N_4

Air, or Oxygen

Mellor, 1940, Vol. 8, 122


It burns incandescently on heating in air, and very vividly in oxygen.

See other NITRIDES

4775. Sodium tetrasulfur pentanitride

[60241-33-0]

N_5NaS_4



Scherer, O. J. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1975, **14**, 485

It explodes at about 180°C, or under pressure, or by friction of a spatula on a sintered glass filter.

See other NITRIDES, N-METAL DERIVATIVES, N–S COMPOUNDS

4776. Triphosphorus pentanitride
[12316-91-3]



Metals

Mellor, 1971, Vol. 8, Suppl. 3, 370–371

Interaction with molten magnesium at 651°C is explosive, and more so with calcium at 200°C.

Oxidants

Partington, 1946, 618

It ignites on heating in chlorine or oxygen.

See also ‘Tetraphosphorus hexanitride’ *See other* NITRIDES

4777. Nickel azide
[59865-91-7]



Mellor, 1940, Vol. 8, 355

It explodes at 200°C.

See other METAL AZIDES

4778. Sulfinyl azide
[]



Pascal, 1961, Vol. 10, 634

An explosive liquid.

See other ACYL AZIDES, NON-METAL AZIDES

4779. Sulfuryl diazide
[72250-07-8]




1. Curtius, T. *et al.*, *Ber.*, 1922, **55**, 1571

2. Nojima, M., *J. Chem. Soc., Perkin Trans. 1*, 1979, 1813

It explodes violently when heated and often spontaneously at ambient temperature [1]. A safe method of preparation in solution has been described [2].

See other ACYL AZIDES, NON-METAL AZIDES

4780. Disulfuryl diazide
[73506-23-7]



Alone, or Alkali

Lehmann, H. A. *et al.*, *Z. Anorg. Chem.*, 1957, **293**, 314

The azide decomposes explosively below 80°C and should only be stored in 1 g quantities. In contact with dilute alkali at 0°C an explosive deposit is formed (? sodium azidosulfate?).

See other ACYL AZIDES, NON-METAL AZIDES

4781. ‘Tetraphosphorus hexanitride’

[]

‘N₆P₄’

Unknown structure

Mellor, 1971, 8, Suppl. 3, 371

Supposedly impure triphosphorus pentanitride, it ignites in air.

See related NITRIDES

4782. Lead(II) azide

[13424-46-9]

N₆Pb

Pb(N₃)₂

1. Mellor, 1940, Vol. 8, 353; 1967, Vol. 8, Suppl. 2, 21, 50

2. Taylor, G. W. C. *et al.*, *J. Crystal Growth*, 1968, **3**, 391

3. Barton, A. F. M. *et al.*, *Chem. Rev.*, 1973, **73**, 138

4. *MCA Case History No. 2053*

As a heavy metal azide, it is rather endothermic (ΔH_f° (s) +436.4 kJ/mol, 1.50 kJ/g).

As a widely used detonator, its properties have been studied in great detail. Although quantitatively inferior to mercury fulminate as a detonator, it has proved to be more reliable in service. The pure compound occurs in 2 crystalline forms, one of which appears to be much more sensitive to initiation [1]. Factors which suppressed spontaneous explosions of lead azide during crystallisation were vigorous agitation and use of hydrophilic colloids [2]. These aspects have been reviewed [3]. A vacuum-desiccated sample of 5 g of the azide exploded violently when touched with a metal spatula [4] (possibly owing to static charge, generated by rapid evaporation of the solvent from the glass-insulated system, being earthed by the spatula).

See Sodium azide: Heavy metals (reference 1)

Calcium stearate

MCA Case History No. 949

An explosion occurred during blending and screening operations on a mixture of lead azide and 0.5% of calcium stearate. If free stearic acid were present as impurity in the calcium salt, free hydrogen azide may have been involved.

Copper, or Zinc

Federoff, 1960, A532, 551

Lead azide, on prolonged contact with copper, zinc or their alloys, forms traces of the extremely sensitive copper or zinc azides which may initiate detonation of the whole mass of azide.

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

4783. Palladium(II) azide
[13718-25-7]




N₆Pd

Ukhin, L. Yu. *et al.*, *J. Organomet. Chem.*, 1981, **210**, 270
The explosive precipitate needs careful handling.
See other METAL AZIDES

4784. Pentasulfur hexanitride
[68993-02-2]

N₆S₅



Sheldrick, W. S. *et al.*, *Inorg. Chem.*, 1980, **19**, 539
Powerfully explosive, like its dimethyl homologue. A m.p. apparatus was wrecked, and a large glass funnel containing 50 mg was pulverised.
See other NITRIDES, N-S COMPOUNDS


4785. Strontium azide
[19465-89-5]



N₆Sr

It is somewhat endothermic (ΔH_f° (s) +204.6 kJ/mol, 1.19 kJ/g).
See Barium azide, *also* Calcium azide
See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

4786. Zinc azide
[14215-28-2]



N₆Zn

Bailar, 1973, Vol. 3, 217
Zinc azide is said to explode easily.
See Sodium azide: Heavy metals *See other* METAL AZIDES

4787. Poly(diazidophosphazene)
[]



(N₇P)_n

Dillon, K. B. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 1037
This polymer of phosphonitrilic diazide is a viscous explosive oil.
See 1,1,3,3,5,5-Hexaazido-2,4,6-triaza-1,3,5-triphosphorine
See other NON-METAL AZIDES

4788. Phosphorus triazide oxide

[4635-46-5]

N₉OPBuder, W. *et al.*, *Z. Anorg. Chem.*, 1975, **415**, 263

The liquid may explode violently on warming from 0°C to ambient temperature, but can be used safely in solution.

See related ENDOTHERMIC COMPOUNDS, NON-METAL AZIDES

4789. Phosphorus triazide

[56280-76-3]

N₉P1. Lines, E. L. *et al.*, *Inorg. Chem.*, 1972, **11**, 22702. Buder, W. *et al.*, *Z. Anorg. Chem.*, 1975, **415**, 263

It is a highly explosive liquid [1], which may explode on warming from 0°C to ambient temperature, but can be used safely in solution [2].

See other NON-METAL AZIDES

4790. Lead(IV) azide

[73513-16-3]

N₁₂Pb

1. Mellor, 1967, Vol. 8, Suppl. 2, 22

2. Möller, H., *Z. Anorg. Chem.*, 1949, **260**, 103. Zbirol, E., *Synthesis*, 1972, (6), 285–302

The crystalline product appears less stable than the diazide, spontaneously decomposing, sometimes explosively [1]. It was rated as too unstable for use as a practical detonator or explosive [2]. Lead(IV) acetate azide (probably the triacetate azide) is also rather unstable, evolving nitrogen above 0°C with precipitation of lead(II) azide [3]. Lead(IV) azide will be considerably more endothermic than the lead(II) salt.

See other ENDOTHERMIC COMPOUNDS, METAL AZIDES

4791. Silicon tetraazide

[27890-58-0]

N₁₂Si

1. Mellor, 1967, Vol. 8, Suppl. 2, 20

2. Wiberg, E. *et al.*, *Z. Naturforsch. B*, 1954, **9B**, 5003. Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 27

Spontaneous explosions have been observed [1] with this dangerously explosive material, especially when pure. A sample at 0°C exploded during removal of traces of benzene under high vacuum [2]. A residue containing the tetraazide, silicon chloride triazide and probably silicon dichloride diazide, exploded on standing for 2 or 3 days, possibly owing to hydrazoic acid produced by hydrolysis.

See other ENDOTHERMIC COMPOUNDS, NON-METAL AZIDES

4792. Thallium(I) tetraazidothallate

[]

N₁₂Tl₂

Bailar, 1973, Vol. 1, 1152

A heat-, friction- and shock-sensitive explosive.

See related ENDOTHERMIC COMPOUNDS, HEAVY METAL DERIVATIVES, METAL AZIDES**4793. Pentaazidophosphorane**

[56295-87-5]

N₁₅P*See* Sodium hexaazidophosphate (next below)**4794. Sodium hexaazidophosphate**


[60955-41-1]

N₁₈NaP1. Volgnandt, P. *et al.*, *Z. Anorg. Chem.*, 1976, **425**, 189–1922. Buder, W. *et al.*, *Z. Anorg. Chem.*, 1975, **415**, 263

The product of interaction of sodium azide and phosphorus trichloride occasionally exploded on warming from 0°C to ambient temperature, but was examined safely in solution. The structure of the explosive product is determined as the title compound [1], rather than pentaazidophosphorane as originally reported [2]. It contains some 82% of nitrogen.

See related NON-METAL AZIDES*See other* ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS**4795. 1,1,3,3,5,5-Hexaazido-2,4,6-triaza-1,3,5-triphosphorine**

[22295-99-4]

N₂₁P₃1. Grundmann, C., *Z. Naturforsch. B*, 1955, **10B**, 1162. Tornieporth-Oetting, I. C. *et al.*, *Angewand. Chem. (Int.)*, 1995, **34**(5), 511

Na

This viscous liquid (containing 73.5% N) is violently explosive when subjected to shock, friction [1] or electric discharge [2].

See Poly(diazidophosphazene)

See other ENDOTHERMIC COMPOUNDS, HIGH-NITROGEN COMPOUNDS, NON-METAL AZIDES

4796. Sodium

[7440-23-5]

Na

Na

(MCA SD-47, 1952); NSC 231, 1978; FPA H4, 1972; HCS 1980, 823

1. Hawkes, A. S. *et al.*, *J. Chem. Educ.*, 1953, **30**, 467
2. *Alkali Metals*, 1957
3. Houben-Weyl, 1979, Vol. 13.1, 287–289
4. Muir, G. D., private comm., 1968
5. Reuillon, M. *et al.*, *Proc. 2nd Int. Symp. Loss Prev. Safety Prom. Proc. Ind.*, 481–486, Frankfurt, Dechema, 1978
6. Anon., *Fire Prevention*, 1995, (May), 39

In a discussion of safe methods for laboratory use of sodium, disposal of small quantities (up to 5–10 g) by immersion in isopropanol, which may contain up to 2% of water to increase the rate of reaction, is recommended. Quantities up to 50 g should be burned in a heavy metal dish, using a gas flame [1], (with suitable arrangements for dispersion or absorption of the alkaline particulate smoke so produced). Handling techniques and safety precautions for large-scale operations have also been detailed for this reactive metal [2]. Techniques for the preparation and precautions in use of finely dispersed sodium ('sodium sand') are given. It is to some extent pyrophoric [3]. Dispersions of sodium in volatile solvents may become pyrophoric if the solvent evaporates round the neck of a container. Serum cap closures are safer than screw caps [4]. Improved methods for extinction of liquid sodium (and other metal) fires are described which involve application of low-melting inorganic powder mixtures which melt and form a protective crust. Mixtures of sodium carbonate or graphite, with lithium carbonate, sodium chloride or sodium fluoride are very effective for sodium [5]. A fire involving a leak of bulk liquid sodium, which ignited, is reported. Autoignition in air can be as low as 100°C [6].

Acids

MRH values below reference

Mellor, 1941, Vol. 2, 469–470

MRH Hydrogen chloride 5.36/61, hydrogen fluoride 6.95/47, nitric acid 5.48/64, sulfuric acid 3.89/69

Anhydrous hydrogen chloride, hydrogen fluoride or sulfuric acid react slowly with sodium, while the aqueous solutions react explosively. Nitric acid of density above 1.056 causes ignition of sodium.

Ammonia, Aromatic hydrocarbons

Stevenson, J. R. *et al.*, *J. Org. Chem.*, 1984, **49**, 3444

When naphthalene is reduced in liquid ammonia by metallic potassium, evaporation of excess ammonia gives the solvated solid potassium naphthalide ($K_2 \cdot$

$\text{NH}_3 \cdot \text{Naphth}_2$). If sodium is used in place of potassium, the product detonates as crystallisation starts. This is attributed to energetic expulsion of ammonia held endothermically in the growing crystal lattice. The same also occurs with anthracene and sodium, and nitrobenzene and barium. Caution in preparing and using these compounds is urged.

Ammonia, 1,2-Dimethoxyethane, Pyridine

Nolan, 1983, Case history 215

Pyridine is converted to a slurry of a dihydrobipyridyl by treatment with sodium in liquid ammonia at -33°C , then warming to 10°C to evaporate excess ammonia. The slurry was then transferred to a decomposition vessel (for subsequent aerobic oxidation to liberate bipyridyl), the transfer pipe being flushed through with cold dimethoxyethane. When warm solvent was used for this purpose on one occasion, oxidation proceeded so fast that ignition occurred.

Aqua Regia

See AQUA REGIA: SODIUM

Calcium, Mixed oxides

See Calcium: Sodium, etc.

Chlorobenzene, Phosphorus trichloride

Venczel, A., Rom. Pat. RO 72 608, 1980

In the preparation of triphenylphosphine from chlorobenzene, phosphorus trichloride and sodium dispersed in toluene or xylene, the possibility of explosion is avoided by adding about 1 mol% of a lower alcohol, based on sodium usage.

1-Chlorobutane, Light petroleum

Shackelford, S. A., *J. Org. Chem.*, 1979, **44**, 3492, footnote 30

In formation of butylsodium, the temperature of interaction of the chloro compound with the dispersion of sodium in light petroleum must not be too low. At -23°C , reaction is smooth, but at -78° excess of chlorobutane accumulated, and when reaction did start, it accelerated to violent explosion.

See Halocarbons (reference 7), below

Chloroform, Methanol

See Chloroform: Sodium, Methanol

Diazomethane

See Diazomethane: Alkali metals

Diethyl ether

Hey, P., private comm., 1965

While sodium wire was being pressed into ether, the die-hole blocked. Increasing the pressure on the ram to free it caused ignition of the ejected sodium and explosion of the flask of ether. Pressing the sodium into less flammable xylene or toluene and subsequent replacement of solvent with ether was recommended.

Dimethylformamide

'DMF Brochure', Billingham, ICI, 1965

A vigorous reaction occurs on heating DMF with sodium metal.

Ethanol

Brändstrom, A., *Acta Chem. Scand.*, 1951, **4**, 1608–1610

Air must be excluded during exothermic interaction of ethanol with sodium finely dispersed in hydrocarbons to avoid the possibility of hydrogen–air mixture explosions.

See Potassium: Alcohols

Fluorinated alkanols

1. Anon., personal communication, 1996

2. Editor's comments

By mistake, sodium was added to excess liquid perfluoroalkylmethanol in the absence of the diluent which had previously moderated the reaction. Vigorous initial reaction culminated in an explosion which destroyed the flask and attached glassware and cracked the fume cupboard window, this was attributed to ignition of evolved hydrogen [1]. Since air would rapidly be swept from the flask by the gas flow, it seems probable that a hydrogen explosion sufficient to damage the hood would be external to the reaction flask. The desired fluoroalkoxide is undoubtedly thermodynamically unstable with respect to sodium fluoride formation and a number of other metal derivatives of fluorinated organics have proved explosively metastable [2].

See METAL DERIVATIVES OF ORGANOFLUORINE COMPOUNDS

Fluorinated compounds

Herring, D. E., private comm., 1964

Fusion of fluorinated compounds with sodium for qualitative analysis requires a high temperature for reaction because of their unreactivity. When reaction does occur, there is often an explosion and precautions must be taken.

Halocarbons

MRH values below references

1. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

2. Lenze, F. *et al.*, *Chem. Ztg.*, 1932, **56**, 921

3. Ward, E. R., *Proc. Chem. Soc.*, 1963, 15

4. *MCA SD-34*, 1949

5. Rayner, P. N. G., private comm., 1968

6. Anon., *Angew. Chem. (Nachr.)*, 1970, **18**, 397

7. Read, R. R. *et al.*, *Org. Synth.*, 1955, Coll. Vol. 3, 158

8. Houben-Weyl, 1970, Vol. 13.1, 394–395

9. Firat, Y., *Makromol. Chem.*, 1975, Suppl. 1, 207

10. Grattan, D. W. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1977, 1734–1744

MRH Carbon tetrachloride 6.10/63, chloroform 5.82/63, 1,2-dichloroethylene 5.52/68, trichloroethylene 5.94/66

Although apparently stable standing in contact, mixtures of sodium with a range of halogenated alkane solvents are metastable and capable of initiation to explosion by shock or impact. Carbon tetrachloride [1,2,3], chloroform, dichloromethane

and chloromethane [1,2], tetrachloroethane [1,2,4] have been investigated, among others. Generally the sensitivity to initiation and the force of the explosion increase with the degree of halogen substitution; the two former less than in the corresponding systems with potassium or potassium–sodium alloy. Any aliphatic halo-carbon (except fully fluorinated alkanes) may be expected to behave in this way. Small portions of sodium and hexachlorocyclopentadiene were mixed in preparation for a sodium fusion test. On shaking a few min later, the tube exploded [5]. On the small scale, no reaction occurred on boiling perfluorohexyl iodide in contact with metallic sodium. With 140 g of iodide and 7 g of sodium an explosion occurred after 30 min [6]. The temperature range for smooth interaction of bromobenzene, 1-bromobutane and sodium in ether to give butylbenzene is critical. Below 15°C reaction is delayed but later becomes vigorous, and above 30°C the reaction becomes violent [7]. Sodium wire and chlorobenzene react exothermally in benzene under nitrogen to give phenylsodium, and the reaction must be controlled by cooling. Use of finely divided sodium will lead to an uncontrollable, explosive reaction [8]. The hazards of contacting dichloromethane with a sodium film during an ultra-drying procedure are stressed. Thorough pre-drying and operations under vacuum are essential precautions [9]. Purification of chloromethane, chloroethane, or dichloromethane by condensation onto a fresh sodium mirror is extremely dangerous unless the halocarbons are very dry and oxygen-free [10].

See Chlorobenzene, etc.

1-Chlorobutane

Chloroform, etc.

Chloroform: Metals

Iodomethane: Sodium

Lithium: Bromobenzene

See entry METAL–HALOCARBON INCIDENTS

Halogen azides

See HALOGEN AZIDES: METALS

Halogens, or Interhalogens

MRH Chlorine 6.90/61, Fluorine 13.43/45

1. Mellor, 1961, Vol. 2, Suppl. 2.1, 450–452
2. Mellor, 1941, Vol. 2, 114, 469
3. Mellor, 1941, Vol. 2, 92, 469
4. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549
5. Booth, H. S. *et al.*, *Chem. Rev.*, 1947, **41**, 427
6. Pascal, 1966, Vol. 2.1, 221

Sodium ignites in fluorine gas but is inert in the liquefied gas [1]. Cold sodium ignites in moist chlorine [2] but may be distilled unchanged in the dry gas [1]. Sodium and liquid bromine appear to be unreactive on prolonged contact [3], but mixtures may be detonated violently by mechanical shock [4]. Finely divided sodium luminesces in bromine vapour [1]. Iodine bromide or iodine chloride react slowly with sodium, but mixtures will explode under a hammer-blow [1]. Interaction of iodine pentafluoride with solid sodium is initially vigorous, but soon slows with film-formation, while that with molten sodium is explosively violent

[2]. Sodium reacts immediately and incandescently with iodine heptafluoride [5]. Mixtures of solid sodium and iodine explode lightly when initiated by shock [6].

Hydrazine

Mellor, 1940, Vol. 8, 316

Anhydrous hydrazine and sodium react in ether to form sodium hydrazide, which explodes in contact with air. Hydrazine hydrate and sodium react very exothermally, generating hydrogen and ammonia.

Hydroxylamine

MRH 3.85/99+

See Hydroxylamine: Metals

Iodates

Cueilleron, J., *Bull. Soc. Chim. Fr.*, 1945, **12**, 88–89

Mixtures of sodium with silver iodate or sodium iodate explode when initiated by shock.

See Oxidants, below

Mercury

Brauer, 1965, Vol. 2, 1802

Interaction of sodium and mercury to form sodium amalgam is violently exothermic, and moderation of the reaction with an inert liquid, or by adding mercury slowly to the sodium is necessary. Even so, temperatures of 400°C may be attained.

Metal halides

1. *Alkali Metals*, 1957, 129

2. Mellor, 1961, Vol. 2, Suppl. 2.1, 494–496

3. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549

4. Cueilleron, J., *Bull. Soc. Chim. Fr.*, 1945, **12**, 88

MRH Ammonium chloride 2.64/66, iron(III) chloride, 3.60/70, manganese(II) chloride 1.97/73, tin(II) chloride 1.38/84, vanadium(III) chloride 2.38/83, vanadyl chloride 2.30/82

Sodium dispersions will reduce many metal halides exothermically. Iron(III) chloride is reduced fairly smoothly at ambient temperature or below in presence of 1,2-dimethoxyethane. Nickel chloride, cobalt chloride, lead chloride or calcium chloride require higher temperatures to initiate the reaction, the exotherm with cobalt chloride increasing the temperature from 325 to 375°C and causing evaporation of most of the dispersing oil [1]. The finely powdered metals produced by reduction of halides of cadmium, chromium, cobalt, copper, iron, manganese, molybdenum, nickel, silver, tin or zinc with sodium dispersed in ether or toluene, are all pyrophoric [2]. Interaction of sodium with vanadyl chloride at 180°C is violent [2]. Mixtures of sodium with metal halides are sensitive to mechanical shock [3]; the ensuing explosions have been graded for intensity. Very violent explosions occurred with iron(III) chloride, iron(III) bromide, iron(II) bromide, iron(II) iodide, cobalt(II) chloride or cobalt(II) bromide. Strong explosions occurred with the various halides of aluminium, antimony, arsenic, bismuth, copper(II), mercury, silver, and tin (including a mixture of tin(IV) iodide and

sulfur), also vanadium(V) chloride and ammonium tetrachlorocuprate. Ammonium, copper(I), cadmium and nickel halides generally gave weak explosions, and most of the alkali- and alkaline earth-metals were insensitive [4].

See Molybdenum pentachloride: Sodium, or Sodium sulfide

See Tungsten hexachloride: Sodium sulfide

Metal oxides

1. Mellor, 1939, Vol. 9, 649
 2. Mellor, 1943, Vol. 11, 237, 542
 3. Mellor, 1941, Vol. 3, 138
 4. Mellor, 1941, Vol. 7, 658, 401
 5. Mellor, 1940, Vol. 4, 770
 6. Brewer, L. *et al.*, *Rept. UCRL 1864*, 7, Washington, USAEC, 1952
 7. Mellor, 1961, Vol. 2, Suppl. 2.1, 474
- MRH Chromium trioxide 3.39/79, copper(II) oxide 2.05/63, sodium peroxide 2.55/63, vanadium(V) oxide 1.30/84

Bismuth(III) oxide[1], chromium trioxide [2], and copper(II) oxide [3], are reduced with incandescence on heating with sodium. Finely divided sodium ignites on admixture with fine lead oxide without heating, while the coarse material reacts vigorously with molten sodium [4]. The latter reduces mercury(I) oxide [5] or molybdenum trioxide [2] with incandescence, the former producing a light explosion also. Sodium reduces sodium peroxide vigorously at 500°C [6] and tin(IV) oxide incandescently on gentle heating [4]. Finely dispersed sodium reduces metal oxides on heating at temperatures between 100 and 300°C, producing, e.g., pyrophoric iron, nickel and zinc [4].

Nitrogen-containing explosives

Leleu, *Cahiers*, 1975, (79), 266

Mixtures of sodium (or its alloy with potassium) and nitromethane, trichloronitromethane, nitrobenzene, dinitrobenzene, dinitronaphthalene, ethyl nitrite, ethyl nitrate or glyceryl nitrate are shock-sensitive, the sensitivity increasing with the number of nitro groups.

Non-metal halides

MRH values below references

1. Mellor, 1940, Vol. 8, 1033
2. Mellor, 1941, Vol. 2, 470
3. Mellor, 1940, Vol. 8, 1016
4. Mellor, 1961, Vol. 2, Suppl. 2.1, 455, 460
5. Mellor, 1940, Vol. 8, 1073
6. Cueilieron, J., *Bull. Soc. Chem. Fr.*, 1945, **12**, 88
7. Mellor, 1947, Vol. 10, 912
8. Kirk-Othmer, 1969, Vol. 18, 438
9. Leleu, *Cahiers*, 1975, (79), 270

MRH Phosphorus trichloride 3.10/67, phosphorus pentachloride 4.89/65, phosphoryl chloride 4.27/69, sulfinyl chloride 5.27/67

Sodium floats virtually unchanged on phosphorus tribromide, but added drops of water caused a violent explosion [1]. Molten sodium explodes with phosphorus trichloride [2] and may ignite or explode with phosphorus pentachloride

Na

[3]. Diselenium dichloride reacts vigorously with sodium on heating, emitting light and heat [4]. Sodium ignites in sulfinyl chloride vapour at 300°C [4], or in a stream of thiophosphoryl fluoride [5]. The shock-sensitive mixtures of sodium with phosphorus pentachloride, phosphorus tribromide or sulfur dichloride gave violent explosions on impact, while those with boron tribromide or sulfur dibromide gave strong explosions [6]. Sodium and phosphoryl chloride interact explosively on heating [8], and mixtures of sodium with sulfinyl fluoride or silicon tetrachloride or silicon tetrafluoride are shock-sensitive explosives [9].

Non-metal oxides

MRH values below references

1. Mellor, 1940, Vol. 6, 70; 1961, Vol. 2, Suppl. 2.1, 468
2. Gilbert, H. N., *Chem. Eng. News*, 1948, **26**, 2604
3. Mellor, 1940, Vol. 8, 554, 945; 1961, Vol. 2, Suppl. 2.1, 458
4. Leleu, *Cahiers*, 1975, (79), 269
5. Weiss, E., *Angew. Chem. (Int.)*, 1993, **32**(11), 1518

MRH Carbon monoxide 6.15/65, chlorine dioxide 5.69/75, sulfur dioxide 4.56/74, sulfur trioxide 5.06/70

Sodium and carbon dioxide are normally unreactive till red-heat has been attained [1], but mixtures of the 2 solids are impact-sensitive and explode violently [2]. Carbon dioxide is unsuitable as an extinguishant for the burning metal alone, as the intensity of combustion is increased by replacing air with carbon dioxide (72.7% oxygen content). However it has been used successfully to extinguish solvent fires where sodium was also present, since it often fails to ignite because of the blanketing effect of solvent vapour. Conversely addition of kerosene to burning sodium enables the whole to be extinguished with carbon dioxide [1]. Finely divided silica (sand) will often react with burning sodium, so is not entirely suitable as an extinguishant [1,2]. (Anhydrous sodium carbonate and sodium chloride are suitably unreactive for this purpose.) Solid sodium is inert to dry liquid or gaseous sulfur dioxide, but molten sodium reacts violently [2]. The moist gas reacts almost as vigorously as water [1]. Phosphorus pentoxide becomes incandescent on warming with sodium, which also ignites in dinitrogen pentoxide [3]. The product of interaction of carbon monoxide with sodium (not sodium carbonyl as previously thought [2], but the hexamer), sodium benzenehexoxide, is shock sensitive, explodes at 90°C and contact with water causes explosion or ignition to occur [4]. Later work [5] makes it likely that this compound is, in fact, the dimer disodium ethynediolate.

Non-metals

MRH Sulfur 4.81/41

1. Mellor, 1961, Vol. 2, Suppl. 2.1, 466–467, 454–455
2. Mellor, 1941, Vol. 2, 469
3. Mellor, 1947, Vol. 10, 766

Explosions have occasionally occurred when carbon powder is in contact with evaporating sodium and air [1]. The violent interaction of ground or heated mixtures of sodium and sulfur may be moderated by the presence of sodium chloride or boiling toluene [1,2]. Selenium reacts incandescently with sodium when heated [3], and molten tellurium reacts vigorously when poured on to solid sodium [1].

Organic samples for qualitative analysis

See LASSAIGNE TEST

Other reactants

Yoshida, 1980, 262–263

MRH values calculated for 19 combinations with various materials are given.

Oxidants

MRH values show % of oxidant

See Halogens or Interhalogens; Metal halides;

Metal oxides; Non-metal oxides, all above

Ammonium nitrate: Metals

MRH 4.15/64

Nitric acid: Metals

MRH 5.84/32

Nitrosyl fluoride: Sodium

Nitryl fluoride: Metals

Sodium nitrate: Sodium

MRH 3.85/43

Sodium nitrite: Reducants

MRH 3.42/50

Sodium peroxide: Metals

MRH 2.55/63

Oxygenated compounds

Leleu, *Cahiers*, 1975, (79), 267

Mixtures of inorganic oxygenated compounds (halide oxides or oxide sulfides) or oxygen-rich organic compounds (alkyl oxalates) with sodium (or its alloy with potassium) are shock-sensitive explosives.

See Non-metal oxides, above

Sulfides

Leleu, *Cahiers*, 1975, (79), 268, 270

Passage of moist hydrogen sulfide over unheated sodium causes melting and then usually ignition of the metal. Mixtures of sodium (or its alloy with potassium) and carbon disulfide are shock-sensitive explosives.

Sulfur

Hames, D. *et al.*, *Proc. Inst. Electr. Eng.*, 1979, **126**, 1157–1161

Safety aspects of large arrays of sodium–sulfur batteries for rail traction are considered among technical aspects.

2,2,3,3-Tetrafluoropropanol

See 2,2,3,3-Tetrafluoropropanol: Potassium hydroxide, etc.

Tetrahydrofuran, Water

Anon., *Lab. Accid. Higher Educ.*, item 5, Barking, HSE, 1987

THF was being dried by refluxing over metallic sodium, and after the reflux period, heat was turned off and a stopper was put in the top of the condenser. The presence of the stopper was forgotten, and when heating was restarted, the condenser was ejected from the flask and broke. Escaping water came into contact with the sodium, evolving hydrogen which ignited and then set fire to the escaping THF vapour in a fume cupboard, and an explosion ensued.

See Water, below

Toluene

See Xylene, below

Water

1. Mellor, 1941, Vol. 2, 469
2. Mellor, 1961, Vol. 2, Suppl. 2.1, 362
3. *MCA Case History No. 456*
4. Markowitz, M. M., *J. Chem. Educ.*, 1963, **40**, 633–636
5. *MCA Case History No. 1653*
6. *MCA Case History No. 2082*
7. Peacocke, T. A. H., *Chem. Brit.*, 1979, **15**, 233

Small pieces of sodium in contact with water react vigorously but do not usually ignite the evolved hydrogen unless the water is above 40°C, or if rapid dissipation of heat is prevented by immobilising the sodium by use of a viscous aqueous solution (starch paste, etc.) or wet filter paper [1]. In contact with ice, sodium explodes violently [2]. Small, hot particles of sodium (remaining from dissolution of larger pieces) may finally explode as do large lumps of the metal in water [1]. Sodium residues from a Wurtz reaction were treated with alcohol to destroy sodium, but later accidental contact with water caused a fire, showing that the alcohol treatment was incomplete. This may have been owing to a crust of alcohol-insoluble halide over the residual sodium [3]. Reactivity of sodium and other alkali-metals with water or steam has been discussed in detail [4]. Rolling a drum containing a 12 cm layer of sodium sludge mixed with soda ash led to a mild pressure explosion. Moisture present in the drum contacted the sodium residue when rolled [5]. Preparing to dispose of a few l of old sodium dispersion by adding a solvent and then burning it, an operator opened the container under cover but in an atmosphere rendered humid by very recent rain. The dispersion immediately ignited and exploded [6]. An explosion of a small piece of sodium retained under water (out of contact with air) was attributed to formation of organic peroxides from arylsodium derivatives produced in storage of sodium under naphtha [7].

Xylene

Anon., *Sichere Chemiarb.*, 1989, **41**(5), 58

After storage for 16 years in a tin, a sealed bottle originally holding sodium dispersed in xylene was found to contain a yellow/white solid layer in place of the expected supernatant xylene. Scraping the solid out caused a violent explosion. The force of the explosion leads to a suspicion of peroxide formation, but there is no obvious explanation. Reactive materials like alkali-metal dispersions in volatile solvents should not be stored indefinitely, but clearly labeled after receipt or preparation to show the disposal date. Disposal of such dispersions by burning is recommended. Sodium dispersed in toluene might behave in the same way.

See ALKALI METALS

See other METALS

4797. Sodium–antimony alloy

[[12300-03-7] (1:1 alloy)]

Na–Sb

Na–Sb

See Nitrosyl tribromide: Sodium–antimony alloy

See other ALLOYS

- 4798. Sodium–zinc alloy**
 [[39422-73-6] (1:6 alloy)] **Na–Zn**
Na–Zn
 Houben-Weyl, 1973, Vol. 13.2a, 571
 Preparation by addition of sodium to molten zinc (1:4) in absence of air is rather violent.
See other ALLOYS
- 4799. Sodium dioxide (Sodium superoxide)**
 [12034-12-7] **NaO₂**
NaO₂
 Sodium peroxide
 Gausing, W. *et al.*, *Angew. Chem. (Intern. Ed.)*, 1978, **17**, 372
 A by-product mixture of sodium dioxide and sodium peroxide separated by filtration was sensitive to warmth and shock when dry.
See related METAL PEROXIDES *See other* METAL OXIDES, OXIDANTS
- 4800. Sodium trioxide**
 [12058-54-7] **NaO₃**
NaO₃
 Petrocelli, A. W. *et al.*, *J. Chem. Educ.*, 1962, **39**, 557
 This is the least stable of the alkali-metal ozonides.
See related METAL PEROXIDES *See other* METAL OXIDES, OXIDANTS
- 4801. Sodium silicide**
 [12164-12-4] **NaSi**
NaSi
 1. Mellor, 1961, Vol. 2, Suppl. 2.1, 564
 2. Bailar, 1973, Vol. 1, 446
 Sodium silicide ignites in air [1], and like its potassium, rubidium and caesium analogues, ignites explosively on contact with water or dilute acids [2].
See other METAL NON-METALLIDES
- 4802. Sodium oxide**
 [1313-59-3] **Na₂O**
Na₂O
 2,4-Dinitrotoluene
See 2,4-Dinitrotoluene: Sodium oxide
 Phosphorus(V) oxide
See Tetraphosphorus decaoxide: Inorganic bases
 Water
 Interaction is likely to be violently exothermic.
See other INORGANIC BASES, METAL OXIDES

4803. Sodium peroxide
[1313-60-6]

Na₂O₂

NaOONa

FPA H104, 1981; *HCS 1980*, 854

1. Allen, C. F. H. *et al.*, *J. Chem. Educ.*, 1942, **19**, 72
2. Rüst, 1948, 297

Hazards attendant on the use of this powerful oxidant may in many cases be eliminated by substitution with 'sodium perborate' (actually sodium borate hydrogen peroxidate) [1]. One of several wooden boxes of the peroxide (not clearly labelled as such) exploded with great violence during handling operations. It seems likely that contamination with a combustible material, or possibly with moisture, had occurred [2].

See Fibrous materials, *also* Wood, both below

Acetic acid MRH 2.80/16

von Schwartz, 1918, 321

Admixture causes explosion, owing either to direct oxidation of acetic acid by the highly concentrated hydrogen peroxide produced, or perhaps to formation of concentrated peroxyacetic acid.

Aluminium MRH 4.56/19

Aluminium chloride MRH 2.30/53

Anon., *Chem. Eng. News*, 1954, **32**, 258

A mixture of the three slowly reacted, creating a pressure of 122 bar (? of chlorine and possibly hydrogen chloride) in 41 days, and the residue reacted spontaneously on exposure to air.

See Metals, etc., below

Ammonium peroxodisulfate MRH 2.72/60

Mellor, 1947, Vol. 10, 464

A mixture exploded on being subjected to friction in a mortar, heating above 75°C, or exposure to carbon dioxide or drops of water.

Boron nitride

Mellor, 1940, Vol. 8, 111

Addition of powdered nitride to the molten peroxide caused incandescence.

Calcium acetylide MRH 3.01/14

Mellor, 1941, Vol. 2, 490

A mixture is explosive.

Carbon dioxide MRH 1.84/36

Cotton wool MRH Cellulose 1.97/99+

1. Bulavin, Yu. I., *Chem. Abs.*, 1978, **89**, 75003
2. Anon., *Chemist-Analyst*, 1978, **67**(3), 7

Cotton wool sprinkled with the peroxide (used in guard tubes) will ignite a few s after exposure to moist carbon dioxide. Exothermic interaction of sodium peroxide and carbon dioxide heats the cotton which then ignites in the liberated oxygen [1]. Cotton contaminated with peroxide should not be stored under any circumstances as it may ignite with atmospheric carbon dioxide [2].

Fibrous materials, Water MRH Cellulose 1.97/99+

1. von Schwartz, 1918, 328
2. Betteridge, D., private comm., 1973

Contact of the solid peroxide with moist cloth, paper or wood often causes ignition [1], and addition of water to intermixed cotton wool and peroxide causes violent ignition [2].

Glucose, Potassium nitrate MRH 2.93/15
491M, 1975, 391

A micro-bomb calorimeter exploded when the wrong proportions of sample and oxidants were used. Instead of 4 g of peroxide and 0.2 g of nitrate for 0.2 g of the sugar sample, 0.35 g of peroxide and 2.6 g of dextrose were used. The deficiency of peroxide to absorb the decomposition gases and excess of organic matter led to a rapid rise in temperature and pressure, which burst the bomb calorimeter.

Hexamethylenetetramine MRH 2.47/10
Ellern, 1968, 46

A mixture ignites when moistened with water.

Hydrogen sulfide MRH 3.01/30

1. Barrs, C. E., *J. R. Inst. Chem.*, 1955, **79**, 43
2. Mellor, 1947, Vol. 10, 129

Solid sodium peroxide causes immediate ignition in contact with gaseous hydrogen sulfide [1]. Barium peroxides and other peroxides behave similarly [2].

Hydroxy compounds MRH values below reference
Castrantas, 1965, 4

MRH Ethanol 2.43/9, ethylene glycol 2.68/14, glycerol 2.93/17, allose 2.93/17, acetic acid 2.80/16

The exothermic oxidation of ethanol, ethylene glycol, glycerol, sugar or acetic acid may lead to fire or explosion.

See Water, below, *also* Ethylene glycol: Oxidants

Magnesium

Bentzinger, von R. *et al.*, *Praxis Naturwiss. Chem.*, 1987, **36**, 38

A priming mixture of magnesium powder and barium peroxide to ignite thermite mixture was to be prepared, but as no barium peroxide was available, sodium peroxide was used instead. Some time after preparation, the mixture ignited spontaneously, because sodium peroxide, unlike barium peroxide, is very hygroscopic and forms hot conc. hydrogen peroxide from contact with atmospheric moisture.

See Carbon dioxide, etc., above

Metal halides

Wiley, J. B. *et al.*, *Mater. Res. Bull.*, 1993, **28**(9), 893

Metathesis reaction to prepare crystalline transition and B metal oxides from halides and sodium peroxide is initiated by a hot wire and can be highly exothermic and explosive in some instances.

See METATHESIS

Metals

MRH Aluminium 4.56/19, iron 1.55/32, magnesium 4.945/24,
sodium 2.55/37, tin 1.38/44

Bunzel, E. G. *et al.*, *Z. Anorg. Chem.*, 1947, **254**, 20

At 240°C mixtures of finely divided metals (aluminium, iron, tungsten) with the peroxide ignite under high friction, and molybdenum powder reacts explosively.

Metals, Carbon dioxide, Water

Mellor, 1941, Vol. 2, 490; 1961, Vol. 2, Suppl. 2.1, 633; 1946, Vol. 5, 217

Intimate mixtures of sodium peroxide with aluminium, magnesium or tin powders ignite on exposure to moist air and become incandescent on heating in air or on moistening with water. Exposure of such mixtures to carbon dioxide causes an explosion. (Interaction of the peroxide and carbon dioxide is highly exothermic.) Sodium is oxidised vigorously at 500°C.

Non-metal halides

1. Mellor, 1947, Vol. 10, 897; 1961, Vol. 2, Suppl. 2.1, 634

2. Comanducci, E., *Chem. Ztg.*, 1911, **15**, 706

Violent interactions occur with diselenium dichloride or disulfur dichloride, the latter emitting light and heat [1]. The very exothermic reaction with phosphorus trichloride may accelerate to explosion [2].

Non-metals

MRH Carbon 3.05/7, phosphorus 2.30/14, sulfur 3.35/29

1. von Schwartz, 1918, 328

2. Mellor, 1941, Vol. 2, 490

3. Anon., *ABCM Quart. Safety Summ.*, 1948, **19**, 13

Intimate mixtures with carbon or phosphorus may ignite or explode [1]. Other readily oxidisable materials (probably antimony, arsenic, boron, sulfur, selenium) also form explosive mixtures [2]. Use of finely powdered carbon, rather than the granular carbon specified for a reagent, mixed with sodium peroxide caused an explosion [3].

Organic liquids, Water

MRH Aniline 2.76/7, benzene 2.76/7, diethyl
ether 2.55/8, glycerol 2.72/15

von Schwartz, 1918, 328

Simultaneous contact of sodium peroxide with water and aniline, or benzene, diethyl ether or glycerol, causes ignition: (equivalent to contact with conc. hydrogen peroxide).

Organic materials

Rüst, 1948, 337

A 'medicinal' mixture of the peroxide (30%) with liquid paraffin (18%), dried soap (42%) and almond oil (10%) ignited explosively during preparation.

Organic material, Water

Bentzinger, von R. *et al.*, *Praxis Naturwiss. Chem.*, 1987, 36, 40

In a combustion demonstration, 50 g of sodium peroxide and half its volume of sago flour are cautiously mixed with a feather. Addition of a few drops of hot water immediately ignites the mixture, which burns vigorously. (The scale of the experiment, using 0.64 mol of sodium peroxide, seems excessive.)

Other reactants

Yoshida, 1980, 87–88

MRH values calculated for 30 combinations with oxidisable materials are given.

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

Silver chloride

MRH 0.21/78

Charcoal

MRH 3.05/7

Mellor, 1941, Vol. 3, 401

An intimate mixture ignites after a short delay (but the same is probably true if silver chloride is omitted).

See Non-metals, above

Sodium dioxide

See Sodium dioxide: Sodium peroxide

Water

1. *Haz. Chem. Data*, 1975, 269

2. Friend, J. N. *et al.*, *Nature*, 1934, **134**, 778

3. Cheeseman, G. H. *et al.*, *Nature*, 1934, **134**, 971

Reaction with water is vigorous, and with large amounts of peroxide it may be explosive. Contact of the peroxide with combustibles and traces of water may cause ignition [1]. Violent explosions on two occasions during attempted preparation of oxygen were attributed to traces of sodium in the peroxide. The former would liberate hydrogen and ignite the detonable mixture [2,3].

Wood


MRH Cellulose 1.97/99+

Dupré, A., *J. Soc. Chem. Ind.*, 1897, **16**, 492

Friction of the peroxide between wooden surfaces caused ignition of the latter.

See other METAL PEROXIDES, OXIDANTS

4804. Sodium thiosulfate
[7772-98-7]



HCS 1980, 863

Widmann, G. *et al.*, *Thermochim. Acta*, 1988, **134**, 451–455

Energy of decomposition (320–520°C) was determined by DSC as 2.42 kJ/g, peaking at 427°C.

Metal nitrates

MRH values show % of nitrate

See Potassium nitrate: Reducants

MRH 2.22/50

Sodium nitrate: Sodium thiosulfate

MRH 2.38/46

Other reactants

Yoshida, 1980, 230

MRH values calculated for 9 combinations with various materials are given.

Sodium nitrite

MRH 2.26/47

Stevens, H. P., *J. Proc. R. Inst. Chem.*, 1946, 285

A mixture of these oxidising and reducing salts will explode violently after most of the water of crystallisation has been driven off by heating.

See other REDOX REACTIONS

See other METAL OXONON-METALLATES, REDUCANTS

4805. Sodium metasilicate
[6834-92-0]



Fluorine

See Fluorine: Metal salts

See other METAL OXONON-METALLATES

4806. Sodium sulfate
[7681-38-1]



Aluminium

See Aluminium: Sodium sulfate

See other METAL OXONON-METALLATES

4807. Sodium dithionite ('Sodium hydrosulfite')

[7775-14-6]

$\text{Na}_2\text{O}_4\text{S}_2$

NaOS(O)S(O)ONa

1. *MCA Case History No. 882*
2. Tyler, B. J. *et al.*, *Hazards from Pressure*, IChE Symp. Ser. No. 102, 45–59, Oxford, Pergamon, 1987
3. Henderson, D. K. *et al.*, *J. Haz. Mat.*, 1988, **19**, 155–159

A batch decomposed violently during drying in a graining bowl. No explanation was offered but contamination with water and/or an oxidant seems likely. Thermal decomposition of pure material occurs violently at 190°C [1]. Spontaneous ignition of thin layers of sodium dithionite dust was studied experimentally. Ignition temperatures were measured for 5–40 mm layers of mixtures with less than 75% of inert dust. The overall importance of the effect of layer depth on heat transfer parameters at the cold surface was established, and the results were compared with theoretical predictions [2]. Using increasing temperature regimes to determine ignition temperature of dithionite dust layers confirmed the previous value of 190°C, but when decreasing regimes were used the much higher figure of 400°C was found. This was explained on the basis of a 2 stage decomposition characteristic [3].

Aluminium, Water

Kirschner, E., *Chem. Eng. News*, 1995, **73**(18), 6

As part of the process for manufacture of a gold-precipitant, 0.5 tonnes of aluminium powder and 3.5 tonnes of sodium dithionite were in a blender to which benzaldehyde was to be added (other sources also claim presence of potassium carbonate). Either from a leaking seal, or in trying to clear a benzaldehyde feed line, some water was admitted, starting a slow reaction emitting sulphurous fumes. After about eleven hours an attempt was made to blanket with nitrogen (if air had initially been involved, it seems probable the blender had long since purged itself with sulphur dioxide). There was an explosion, followed by fire, killing five workers. In view of the known properties of 'hydros', admission of water might warm localised regions to a temperature where self-sustaining decomposition would ensue in the absence of air. Although a reducing agent in wide use, where aluminium is concerned sodium dithionite and its decomposition products are most definitely oxidants (as is potassium carbonate). Aluminium oxidation was probably responsible for the main blast.

See Aluminium: Metal oxides, or Oxosalts, or Sulfides

See also Water, below

Sodium chlorite

See Sodium chlorite: Sodium dithionite

Water

1. Anon., *Chem. Trade J.*, 1939, **104**, 355
2. *MCA Case History No. 2292*
3. Ullmann, 1994, Vol A25, 483

Addition of 10% of water to the solid anhydrous material caused a vigorous exotherm and spontaneous ignition. Bulk material may decompose at 135°C [1]. A loose drum lid allowed ingress of water and smouldering started. As the drum was tipped over for disposal, ignition occurred [2]. Air is usually involved in the exotherm leading to fire after water contamination. The dihydrate may be pyrophoric when finely divided. In the absence of air, exothermic decomposition can occur from 90°C after prolonged heating [3].

See other METAL OXONON-METALLATES, REDUCANTS

4808. Sodium disulfite ('Sodium metabisulfite')

[7681-57-4]



Potassium iodate

See Potassium iodate: Sodium disulfite, Water


Sodium nitrite

See Sodium nitrite: Sodium disulfite

See other METAL OXONON-METALLATES, REDUCANTS

4809. Sodium peroxodisulfate

[7775-27-1]



$T_{\text{ait}24}$ was determined as 148°C by adiabatic Dewar tests, with an apparent energy of activation of 75 kJ/mol.

See entry THERMOCHEMISTRY AND EXOTHERMIC DECOMPOSITION (reference 2)


2,2'-Azobis(2-amidiniopropane) chloride

See 2,2'-Azobis(amidiniopropane) peroxodisulfate

See other PEROXOACID SALTS, OXIDANTS

4810. Sodium tetraperoxotungstate

[12501-01-8]



Mellor, 1943, Vol. 11, 835

It explodes feebly on warming.

See other PEROXOACID SALTS

4811. Sodium sulfide

[1313-82-2, 27610-45-2 (anhydr., 9H₂O, resp.)]

Na₂S

Na₂S

FPA H115, 1982; *HCS 1980*, 860

1. Anon., *ABCM. Quart. Safety Summ.*, 1942, **13**, 5

2. Dunn, B. W., *Bur. Explos. Accid. Bull., Amer. Rail Assoc.*, 1917, **25**, 36

Fused sodium sulfide in small lumps is liable to spontaneous heating from aerobic oxidation, temperatures of up to 120°C being observed after exposure to moisture and air. Packing in hermetically closed containers is essential [1]. Previously, similar material packed in wooden barrels had ignited in transit [2].

See also SULFUR BLACK

Ammonium persulfate

See Ammonium peroxydisulfate: Sodium sulfide

Carbon

MRH 0.041/tr.

Creevey, J., *Chem. Age*, 1941, **44**, 257

Mixtures of sodium sulfide and finely divided carbon exhibit an exotherm on exposure to air. As indicated by the low MRH value, this is probably not a direct interaction, but arises from co-promotion of aerobic oxidation of the individual components.

Diazonium salts

See DIAZONIUM SULFIDES


N,N-Dichloromethylamine

See N,N-Dichloromethylamine: Calcium hypochlorite, etc.

Glass

L.B., Personal experience, 1964

An old soda glass bottle of the hygroscopic nonhydrate had become severely etched and corroded internally by the contents, which had leaked through the perforated wall of the bottle. It is probable that the old material had become oxidised by air to a strongly alkaline mixture (below), corrosive to the soft glass.



See other CORROSION INCIDENTS, GLASS INCIDENTS

Hydrogen peroxide

See Hydrogen peroxide: Sulfides

Metal halides

See Tungsten hexachloride: Sodium sulfide

Other reactants

Yoshida, 1980, 392

MRH values calculated for 18 combinations, largely with oxidants, are given.

Sodium carbonate, Water
See SMELT: WATER
See other METAL SULFIDES

4812. Sodium disulfide
[22868-13-9]



Diazonium salts
See DIAZONIUM SULFIDES *See other* METAL SULFIDES

4813. Sodium polysulfide
[1344-08-7]



Diazonium salts
See DIAZONIUM SULFIDES AND DERIVATIVES *See other* METAL SULFIDES

4814. Sodium orthophosphate
[7601-54-9]




Sugars
See SUGARS

4815. Sodium phosphide
[12058-85-4]



Water
Mellor, 1940, Vol. 8, 834
Sodium phosphide (or potassium phosphide) is decomposed by moist air or water, evolving phosphine, which often ignites.
See other METAL NON-METALLIDES

4816. Sodium pyrophosphate hydrogen peroxidate
[15039-07-3]



See entry CRYSTALLINE HYDROGEN PEROXIDATES

4817. Niobium
[7440-03-1]

Nb

Nb

Halogens, or Interhalogens

Mellor, 1939, Vol. 9, 849; 1956, Vol. 2, Suppl. 1, 165

Niobium ignites in cold fluorine, and in chlorine at 205°C, and incandescs in contact with bromine trifluoride.

See other METALS

4818. Niobium(V) oxide
[1313-96-8]

O₂NbONbO₂

Nb₂O₅

Lithium

See Lithium: Metal oxides

See other METAL OXIDES

4819. Neodymium
[7440-00-8]

Nd

Nd

Phosphorus

See Phosphorus: Metals See other METALS

4820. Nickel
[7440-02-0]

Ni

Ni

RSC Lab. Hazards Data Sheet No. 60, 1987 (Ni and compounds)

1. Sasse, W. H. F., *Org. Synth.*, 1966, **46**, 5
2. Whaley, T. P., *Inorg. Synth.*, 1957, **5**, 197
3. Brown, C. A. *et al.*, *J. Org. Chem.*, 1973, **38**, 2226
4. Savel'eva, O. N., *Chem. Abs.*, 1982, **96**, 165547
5. Anon., *Jahresber.*, 1987, 65

Raney nickel catalysts must not be degassed by heating under vacuum, as large amounts of heat and hydrogen may be evolved suddenly and dangerous explosions may be caused [1]. Nickel powder prepared by several alternative methods may be pyrophoric if particles are fine enough [2]. A non-pyrophoric highly active colloidal hydrogenation catalyst (P-2 nickel) is produced by reduction of nickel acetate in ethanol by sodium tetrahydroborate [3]. Catalytic Raney nickel electrodes are rendered non-pyrophoric by aqueous metal nitrate treatment [4]. After storage for some time, a drum containing Raney nickel catalyst under water was opened for inspection. As the lid was being removed, there was an explosion which blew it up

into the air. This is attributed to desorption of hydrogen into the air-space to form an explosive mixture, and ignition of the latter by traces of dry (and pyrophoric) catalyst in the top of the drum which sparked in contact with the ingress of air as the lid was opened. It was suggested that such stored drums should be shaken to wet any catalyst at the top of the drum before opening. Displacing any air with carbon dioxide before sealing would seem also to be advisable [5].

Aluminium

See Aluminium–nickel alloys

Aluminium chloride, Ethylene

See Ethylene: Aluminium chloride

1,4-Dioxane

MRH 1.46/99+

See 1,4-Dioxane: Nickel

Hydrogen

1. Adkins, H. *et al.*, *J. Amer. Chem. Soc.*, 1948, **70**, 695
2. Adkins, H. *et al.*, *Org. Synth.*, 1955, Coll. Vol. 3, 176
3. Chadwell, A. J. *et al.*, *J. Phys. Chem.*, 1956, **60**, 1340

During hydrogenation of an unspecified substrate (possibly 4-nitrotoluene) at high pressure with the highly active W6 type of Raney nickel catalyst at 150°C, a sudden exotherm caused the initial pressure to double rapidly to 680 bar [1]. This does not happen at 100°C or below. Care is necessary with selection of reaction conditions for highly active catalysts [2]. Hydrogen-laden Raney nickel catalyst when heated strongly under vacuum undergoes explosive release of the sorbed hydrogen [3].

Hydrogen, Oxygen

Lee, W. B., *Ind. Eng. Chem., Prod. Res. Dev.*, 1967, **6**(1), 59–64

Raney nickel powder entrained in cryogenic hydrogen gas causes ignition on contact with liquid or cryogenic gaseous oxygen.

See other CATALYTIC IMPURITY INCIDENTS

Magnesium silicate

Blake, E. J., private comm., 1974

The pyrophoricity of nickel-on-sepiolite catalysts after use in petroleum processing operations may be caused by the presence of finely divided nickel and/or carbon.

See other PYROPHORIC CATALYSTS

Methanol

1. *MCA Case History No. 1225*
2. Anon., *Jahresber.*, 1983, 77–78
3. Anon., *Sichere Chemiearbeit*, 1990, **42**, 45

Ignition occurred when methanol was poured through the open manhole of a 230 l reactor containing Raney nickel catalyst. Although the reactor had been purged thoroughly with nitrogen before opening, it was later shown that air was entrained

during pouring operations, and this would cause nickel particles round the manhole opening to glow. A closed charging system was recommended [1]. A slurry of Raney nickel catalyst in methanol was to be added to a nitrogen-filled reactor containing a methanol solution. A valve connecting the reactor to a closed circuit acid-scrubber was opened to equalise the pressure, but the scrubber was at a rather lower pressure and reduced the pressure in the total system below ambient. When the charging port was opened, air was drawn in carrying residual catalyst particles into the vessel and ignition occurred. Remedial measures are outlined [2]. An identical ignition injuring two workers is reported subsequently [3].

Non-metals

Mellor, 1942, Vol. 15, 148, 151

On heating, mixtures of powdered nickel with sulfur or selenium react incandescently.

See Sulfur compounds, below

Organic solvents

Hotta, K. *et al.*, *Chem. Abs.*, 1969, **70**, 81308

Raney nickel catalyst evaporated with small amounts of methanol, ethanol, isopropanol, pentanol, acetone, benzene, cyclohexane or *p*-dioxane and then heated towards 200°C eventually explodes

See 1,4-Dioxane: Nickel

Other reactants

Yoshida, 1980, 270

MRH values calculated for 14 combinations, largely with oxidants, are given.

Oxidants

values show % of oxidant

See Ammonium nitrate: Metals

MRH 2.59/58

Bromine pentafluoride: acids, etc.

MRH 3.64/55

Chlorine: Metals

MRH 2.34/55

Nitryl fluoride: Metals

Peroxyformic acid: Metals

MRH 5.69/99+

Potassium perchlorate: Metal powders

MRH 2.64/37

Sulfur compounds

Kornfeld, E. C., *J. Org. Chem.*, 1951, **16**, 137

Raney nickel catalyst, containing appreciable amounts of the sulfide (after use to desulfurise thioamides) is rather pyrophoric.

See other HYDROGENATION CATALYSTS, METALS, PYROPHORIC METALS

4821. Nickel(II) oxide [1313-99-1]	NiO	NiO
Anilinium perchlorate <i>See</i> Anilinium perchlorate: Metal oxides		
Fluorine <i>See</i> Fluorine: Metal oxides		
Hydrogen peroxide <i>See</i> Hydrogen peroxide: Metals, etc.		
Hydrogen sulfide <i>See</i> Hydrogen sulfide: Metal oxides <i>See other</i> METAL OXIDES		
4822. Nickel(IV) oxide [12035-36-8]	NiO₂	NiO₂
Fluorine <i>See</i> Fluorine: Metal oxides <i>See other</i> METAL OXIDES		
4823. Nickel(III) oxide [1314-06-3]	Ni₂O₃	Ni₂O₃
Hydrogen peroxide <i>See</i> Hydrogen peroxide: Metals, etc.		
Nitroalkanes <i>See</i> NITROALKANES: metal oxides <i>See other</i> METAL OXIDES		
4824. Lead(II) oxide [1317-36-8]	PbO	OPb
Aluminium carbide Mellor, 1946, Vol. 5, 872 The carbide is oxidised with incandescence on warming with lead oxide. <i>See</i> Metal acetylides, below		
		1837

Chlorinated rubber

See CHLORINATED RUBBER: metal oxides

Chlorine, Ethylene

See Chlorine: Hydrocarbons (references 1,4)

Dichloromethylsilane

See Dichloromethylsilane: Oxidants

Fluorine, Glycerol

See Fluorine: Miscellaneous materials (reference 1)

Fluoroelastomers

Simpson, M. B. H., *Kautsch. Gummi, Kunstst.*, 1980, **33**, 83–85

Dispersions of lead oxide in fluoroelastomers underwent severe exothermic decomposition when heated above the normal mixing temperature of below 200°C, forming elemental lead.

Glycerol, Perchloric acid

See Perchloric acid: Glycerol, etc.

Hydrogen trisulfide

See Hydrogen trisulfide: Metal oxides

Linseed oil

Stolyarov, A. A., *Chem. Abs.*, 1935, **29**, 3860₃

Spontaneous ignition occurring during the early stages of mixing and grinding the components was traced to individual grades of lead oxide and the presence of undispersed lumps in the mixture.

Metal acetylides

Mellor, 1945, Vol. 5, 849

Interaction at 200°C with rubidium acetylide is explosive, and with lithium acetylide, incandescent.

Metals

Mellor, 1946, Vol. 5, 217; 1941, Vol. 7, 116, 656–658

Mixtures of the oxide with aluminium powder give a violent or explosive 'thermite' reaction on heating. Finely divided sodium ignites on admixture with the oxide, and a mixture of the latter with zirconium explodes on heating. Titanium is also oxidised violently on warming.

Non-metals

1. Mellor, 1941, Vol. 7, 657

2. Goodfield, J. A. C. *et al.*, *Fuel*, 1982, **61**, 843–847

A mixture with boron incandesces on heating, and with silicon the reaction is vigorous. If aluminium is present the mixture explodes on heating (but the same

is true if silicon is absent) [1]. The pyrotechnic reaction of boron and lead oxide mixtures has been studied by DSC [2].

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

Seleninyl chloride

See Seleninyl chloride: Metal oxides

See other METAL OXIDES, OXIDANTS

4825. Palladium(II) oxide

[1314-08-5]

PdO

OPd

Hydrogen

Sidgwick, 1950, 1558

It is a strong oxidant and glows in contact with hydrogen at ambient temperature.

See other METAL OXIDES, OXIDANTS

4826. Thorium oxide sulfide

[12218-77-8]

O:Th:S

OSTh

Mellor, 1941, Vol. 7, 240

It ignites in contact with air.

See related METAL OXIDES, METAL SULFIDES *See other* PYROPHORIC MATERIALS

4827. Zirconium oxide sulfide

[12164-95-3]

O:Zr:S

OSZr

Sorbe, 1968, 160

It ignites in air.

See related METAL OXIDES, METAL SULFIDES *See other* PYROPHORIC MATERIALS

4828. Silicon oxide

[10097-28-6]

SiO

OSi

Zintl, E. et al., *Z. Anorg. Chem.*, 1940, **245**, 1

The freshly prepared material ignites in air.

See other NON-METAL OXIDES, PYROPHORIC MATERIALS

4829. Tin(II) oxide
[21651-19-4]

SnO

OSn

Bailar, 1973, Vol. 2, 64

On heating at 300°C in air, oxidation proceeds incandescently.

Non-metal oxides

Mellor, 1941, Vol. 7, 388

The oxide ignites in nitrous oxide at 400°C, and incandesces when heated in sulfur dioxide.

See other METAL OXIDES

4830. Zinc oxide
[1314-13-2]

ZnO

OZn

NSC 267, 1978; *HCS 1980*, 968

Aluminium, Hexachloroethane

See Hexachloroethane: Metals

Chlorinated rubber

See CHLORINATED RUBBER: metal oxides

Linseed oil

Anon., *Chem. Trade J.*, 1933, **92**, 278

Slow addition of zinc white (a voluminous oxide containing much air) to cover the surface of linseed oil varnish caused generation of heat and ignition. Lithopone, a denser (air-free) grade of oxide did not cause heating.

Magnesium

See Magnesium: Metal oxides

See other METAL OXIDES

4831. Oxygen (Gas)
[7782-44-7]

O₂

O₂

NSC 472, 1977; *FPA H12*, 1973; *HCS 1980*, 708

1. Anon., *Sichere Chemiearbeit*, 1994, **46(8)**, 94

Even a slight increase in the oxygen content of air above its usual 21 vol.% will greatly increase the rate of oxidation or combustion of many substances, including the human anatomy. An autoclave for hot oxygen treatment (900°C, 406 bar) was located in a bunker. To purge it with argon for cooling it was necessary to enter

the bunker and open a valve. This discharged hot oxygen into a pipe and ignited a plastic seal, thence the metal of the pipe, then combustible contents of the bunker burnt explosively in the escaped oxygen. The operator was killed immediately [1]. Low concentrations of oxygen in other gases may be concentrated as liquid when working at low temperatures.

See Oxygen (Liquid)

See OXYGEN ENRICHMENT

Acetaldehyde

See Acetaldehyde (references 2,4)

Acetone

Anon., private comm., 1988

To 'ensure proper ignition' during oxygen-flask analysis, a technician put a single drop of acetone onto the filter paper enclosing the sample, before igniting the tail and insertion into the oxygen-filled flask. An explosion followed which shattered the flask, fortunately enclosed in a mesh screen which retained the fragments. It was calculated that 0.07 ml of acetone would have formed an explosive vapour-air mixture in the flask above the lower explosive limit, so considerably less acetone would have given an explosive mixture in the oxygen atmosphere.

See Barium acetate: Copper(II) oxide, etc.

Acetone, Acetylene

Anon., *CISHC Chem. Safety Summ.*, 1976, **47**, 33–34

A combination of faulty equipment and careless working led to an extremely violent explosion during oxy-acetylene cutting work. The oxygen cylinder was nearly empty and the regulator had a cracked diaphragm. The acetylene cylinder was lying on its side and was feeding a mixture of liquid acetone and acetylene gas to the burner head. When the oxygen ran out, the excess pressure from the acetylene line forced the acetone-acetylene mixture back up the oxygen line and into the cylinder via the cracked diaphragm. The explosion destroyed the whole plant.

See Acetylene: Oxygen

Acetylene

Skobelar, V. F., *Chem. Abs.*, 1981, **94**, 120243

A safe method for demonstrating explosive combustion of acetylene-oxygen mixtures in bubbles is described.

sec-Alcohols

1. Davies, 1961, 80

2. Redemann, E. G., *J. Amer. Chem. Soc.*, 1942, **64**, 3049

Secondary alcohols are readily autoxidised in contact with oxygen or air, forming ketones and hydrogen peroxide [1]. A partly full bottle of 2-propanol exposed to sunlight for a long period became 0.36 M in peroxide and potentially explosive [2].

See 2-Propanol: 2-Butanone

See 2-Butanol

See Hydrogen peroxide: Acetone, or: Alcohols

Alkali metals

1. Mellor, 1941, Vol. 2, 469

2. Sidgwick, 1950, 65

Reactivity towards air or oxygen increases from lithium to caesium, and the intensity depends on state of subdivision and on presence or absence of moisture. Lithium normally ignites in air above its m.p., while potassium may ignite after exposure to atmosphere, unless it is unusually dry. Rubidium and caesium ignite immediately on exposure [1]. It is reported that sodium and potassium may be distilled unchanged under perfectly dried oxygen [2].

Alkaline earth metals

Mellor, 1941, Vol. 3, 637–638

Finely divided calcium may ignite in air, and the massive metal ignites on heating in air, and burns vigorously at 300°C in oxygen. Strontium and barium behave similarly.

Aluminium–titanium alloys

See Aluminium–titanium alloys: Oxidants

Ammonia

Rüst, 1948, 332–333

Accidental connection of an oxygen cylinder to top-up an ammonia-containing refrigeration system led to explosive destruction of the compressor.

Ammonia, Platinum

1. Nealy, D., and Williams, E. J. *et al.*, *School Sci. Rev.*, 1935, **16**(63), 410

2. Long, G. C., *Spectrum* (Pretoria), 1980, **18**, 30

In school demonstrations of the oxidation of ammonia to nitric acid over platinum catalysts, substitution of oxygen for air causes fairly vigorous explosions to occur [1]. Practical details are given [2].

1,4-Benzenediol, 1-Propanol

1. Burgiel, J. C. *et al.*, *J. Chem. Phys.*, 1965, **43**, 4291

2. Anon., *Univ. Safety Assoc. Safety News*, 1978, (9), 11–12; Peacock, C. J., private comm., 1978

The solid inclusion (clathrate) complex of oxygen with hydroquinone had been prepared twice previously by a published method which involved saturating a solution of hydroquinone in propanol in an autoclave at 70°C with oxygen at 20–150 bar, followed by slow cooling under oxygen pressure (and most probably without stirring to allow large crystals of the inclusion complex to form) [1]. In a third attempt, (to produce smaller crystals free of the ‘quinhydrone complex’ impurity) a solution was prepared and saturated with oxygen at 80 bar, then heated to 90°C (when the partial pressure of oxygen would increase to around 100 bar),

then allowed to cool with agitation. After 20 min the bursting disk failed at 450 bar and an explosion damaged the autoclave and premises [2]. Without agitation, the rate of oxidation of the solvent and hydroquinone would be controlled by diffusion of oxygen through the (small) gas liquid interface. With agitation, the rate of oxidation would be expected to increase greatly, and lead to runaway oxidation with very fast increase in temperature and pressure and explosive combustion when the autoignition temperature (371°C for propanol in air at 1 bar, less in oxygen) were attained.

See Hydrocarbons (reference 3), below

Benzoic acid

1. Corbett, R. A., private comm., 1976
2. Wilson, L. Y. et al., *J. Chem. Educ.*, 1985, **62**(10), 902

Benzoic acid is burned in oxygen as a primary thermochemical standard to calibrate oxygen bomb calorimeters used in the IP12/ASTM D240 standard tests for determination of calorific value of liquid hydrocarbon fuels. If the benzoic acid is powdered (rather than pelleted as IP12 recommends), very rapid combustion occurs and the flame front may burn through the non-metallic (Teflon) seals on valve seats and the bomb may be destroyed [1]. Ignition of a pelleted sample in a bomb with a faulty closure valve led to sudden venting of combustion gases which blew off the insulating cover and thermometer. Safety precautions are listed for bomb calorimetry work [2].

Biological material, Ether

Napier, D. H., private comm., 1972

Biological material in a polythene bag filled with oxygen and being prepared for analytical combustion exploded. Diethyl ether used to anaesthetise the experimental animal from which the sample was derived may have still been present, and ignition from static charge on the plastics bag may have been involved.

See related STATIC INITIATION INCIDENTS

Boron tribromide

See Arsine–boron tribromide

Bromine, Chlorotrifluoroethylene

See Bromine: Chlorotrifluoroethylene, Oxygen

Calcium phosphide

See Calcium phosphide: Oxygen

Carbon disulfide

Dudkin, V. A., *Chem. Abs.*, 1982, **97**, 203762

The lower limit for spontaneous ignition of mixtures with oxygen has been studied.

Carbon disulfide, Mercury, Anthracene

Anon., *ABCM Quart. Safety Summ.*, 1953, **25**, 2

Shortly after mercury was accidentally introduced into a system containing a solution of anthracene in carbon disulfide under an atmosphere of oxygen, an explosion occurred. Presence of mercury may have catalysed rapid oxidation of carbon disulfide.

See other CATALYTIC IMPURITY INCIDENTS

Carbon monoxide, Hydrogen

Schierwater, F.-W., *Sichere Chemiearb.*, 1985, **37**(9), 101–102

Synthesis gas (CO + H₂) at 40 bar containing a low level of hydrogen sulfide was to be freed of the latter impurity by adding the theoretical quantity of oxygen and passing the mixture over a catalyst. Introduction of oxygen (from a supply at 60 bar) via a simple T-piece (instead of through the recommended small bore coaxial injection nozzle to ensure thorough mixing with the gas stream) caused development of an intense inverse flame in the locally very high oxygen concentration which burned through the reactor side wall opposite the oxygen inlet and ejected a metre-long flame-jet.

Copper, Hydrogen sulfide

See Hydrogen sulfide: Copper, Oxygen

Cyclohexane-1,2-dione bis(phenylhydrazone)

1. Schulze, M. *et al.*, *Z. Chem.*, 1979, **19**, 210–211

2. Adam, W. A. *et al.*, *Tetrahedron*, 1985, **41**, 2045–2056

Autoxidation of the hydrazone (O₂/C₆H₆/UV) gives the explosive isomeric 1,2-dihydroperoxy-1,2-bis(benzeneazo)cyclohexane [1], and the same is true for COT derivatives [2].

See 1,2-Dihydroperoxy-1,2-bis(benzeneazo)cyclohexane

See α -PHENYLAZO HYDROPEROXIDES

Cyclooctatetraene

Adam, W. A. *et al.*, *Tetrahedron Lett.*, 1982, **23**, 3156

In a procedure to oxygenate photochemically COT to the 1,4-endoperoxide, highly explosive undefined polymeric peroxides are formed as by-products.

See other POLYPEROXIDES

Diborane

See Diborane: Oxygen

Diboron tetrafluoride

See Diboron tetrafluoride: Oxygen

Dimethoxymethane

See Dimethoxymethane: Oxygen

Dimethylketene

See Poly(peroxyisobutyrolactone)

Dimethyl sulfide

Harkness, A. C. *et al.*, *Atmos. Environ.*, 1967, **1**, 491–497

Interaction is explosive at 210°C or above, and the mechanism has been studied.

Ethers

Davies, 1961, 79

Many ethers, either of open chain (diethyl or diisopropyl ether) or cyclic type (tetrahydrofuran, dioxane), are readily autoxidised on exposure to air or oxygen in presence of light. The hydroperoxides formed are less volatile than the parent ether and may be concentrated to a dangerous extent if distillation of peroxidised material is attempted.

See AUTOXIDATION, PEROXIDES IN SOLVENTS

Fibrous materials

1. Anon., *Loss Prev. Bull.*, 1978, (020), 49

2. Anon., *Sichere Chemiearbeit*, 1991, **43**(10), 112

3. Anon., *Loss Prev. Bull.*, 1978, (023), 141–143

Most fibrous fabrics will sorb oxygen when exposed to concentrations greater than the normal 21% in air, and this is retained for a long time after excess oxygen is no longer present, greatly increasing the possibility of ignition. Hose leakage led to oxygen enrichment inside a vessel in which welding was taking place. A welder lit a cigarette but did not appreciate the significance of its rapid combustion and the unusually long lighter flame. Sparks from welding later ignited clothing, leading to fatal burns [1]. Accidental connection of a pneumatic grinder to an oxygen line instead of a compressed air line led to ignition of the apprentices' clothing when the grinder was started. Another grinding incident occurred in the open air near the vent of a liquid oxygen tank, not only clothing but the worker's hair and beard ignited [2]. Accidental connection of a breathing set to an oxygen cylinder rather than to a medical air cylinder caused ignition of the facepiece [3].

See OXYGEN ENRICHMENT

Fluorine, Hydrogen

See Fluorine: Hydrogen, Oxygen

Fluorocarbons (including perfluorocarbons)

See FLUOROCARBONS: OXYGEN

Fuel gases

NFPA 51, Quincy (Ma), Natl. Fire Prot. Assocn., 1984

The US fire code covers installation and use of gaseous oxygen–fuel gas systems for welding and cutting, for the thermodynamically unstable fuels acetylene, MAPP (methylacetylene–allene–propene–propane mixtures), and the stable hydrocarbons propane or butane.

Fuels

1. Ivanov, B. A. *et al.*, *Chem. Abs.*, 1975, **82**, 75130

2. Rasbash, D. J., *Fire Saf. J.*, 1986, **11**, 113–125
3. Plewinsky, B. *et al.*, *Prax. Sicherheitstech.*, 1997, **4**, 95

Possible reasons for explosions of 40 l steel cylinders containing oxygen at or below 150 bar are discussed. Presence of a substantial (0.1–0.2 mm) layer of mineral oil, or more likely, ingress of a combustible gas from a higher-pressure source, are possibilities [1]. The major factors affecting the occurrence and progress of explosions in mixtures of air with flammable gases, vapours, mists and dusts are discussed. These include flammability limits, burning velocity in laminar and turbulent conditions, possible onset of detonation, pressure piling, etc. [2]. A review, with references, of detonation in systems with gaseous oxidants and condensed fuels, with particular relevance to the design of oxidation reactors, has been published [3].

See Hydrocarbons, and Oil films, both below

Glycerol

Smith, B. S., *Chem. Abs.*, 1978, **89**, 47649

Glycerol is oxidised at an appreciable rate by high-pressure oxygen in the presence of copper alloys to give glyceric acid, oxalic acid and formic acid, which lead to severe corrosion of the alloy. Malfunction of aircraft oxygen gauges operating at 125 bar was attributed to blockage of the brass Bourdon tubes by copper salts of the acids produced by long-term oxidation of residual traces of glycerol–water solutions used for gauge testing.

See other CORROSION INCIDENTS

Halocarbons

1. Fawcett, H. H., *Chem. Eng. News*, 1957, **35**(43), 60
2. Weber, U., *Chim. Ind.* (Paris), 1963, **90**, 178
3. Bashford, L. A. *et al.*, *J. Chem. Soc.*, 1938, 1064
4. Haszeldine, R. N. *et al.*, *J. Chem. Soc.*, 1959, 1085–1086

1,1,1-Trichloroethane exploded after heating under oxygen at 54 bar and 100°C for 3 h [1]. Trichloroethylene, remaining in a pipe after cleaning operations, exploded under 27 bar pressure of oxygen at ambient temperature. It was later found possible to explode stoichiometric mixtures [2]. Chlorotrifluoroethylene and bromotrifluoroethylene each react explosively with oxygen at ambient temperature [3], but controlled oxidation of the former produces an explosive cyclic peroxide, 4,5-dichloro-3,3,4,5,6,6-hexafluoro-1,2-dioxane [4].

See Tetrafluoroethylene

See Oxygen (Liquid): Halocarbons

Iodomethane: Oxygen

Hydrocarbons

1. Davies, 1961, 11
2. Staudinger, H., *Z. Elektrochem.*, 1925, **31**, 549
3. Unpublished information, 1968
4. Burgess, G. K., *Ind. Eng. Chem.*, 1930, **22**, 473
5. Francis, C. K., *Ind. Eng. Chem.*, 1930, **22**, 896

6. Hunn, E. B., *Ind. Eng. Chem.*, 1930, **22**, 804
7. Dunn., B. W., *Bur. Expl. Accid. Bull., Amer. Rail Assoc.*, 1924, **32**(1), 1–2
8. Anon., *CISHC Chem. Safety Summ.*, 1977, **48**, 8
9. Anon., *Loss Prev. Bull.*, 1978, (020), 49
10. Davies, R., *Loss Prev.*, 1979, **12**, 103–110
11. Kletz, T. A., *Loss Prev.*, 1979, **12**, 96–102
12. Herrmann, G. *et al.*, Ger. Offen. DE 3 514 924, 1986

Interaction of hydrocarbons with gaseous oxygen may be slow or fast, depending on conditions and substrate, but peroxidic products are always involved, and the rate of formation is highest at a C–H link adjacent to an aromatic ring or a double bond [1]. Predictable conditions for explosion are those relevant to the internal combustion engine (Otto or Diesel types). Unpredicted conditions include the following cases. During studies on autoxidation of 1,1-diphenylethylene with oxygen at high pressure and low temperature an explosion occurred [2]. Partial oxidation of a gasoline fraction in an autoclave under oxygen, initially at 22 bar and 100°C, ran wild and exploded; several smaller reactions had proceeded uneventfully [3]. Previously oxidation reactions of gasoline at 20 bar/105°C had exploded [4], and a severe explosion occurred while admitting oxygen, virtually at 1 bar pressure, to an autoclave containing a glass bottle of gasoline [5]. Precautions were detailed later [6].

An attempt to decarbonise a gasoline motor cycle engine by inserting a lighted match through the spark plug hole into the cylinder, and then blowing in oxygen, caused a violent explosion to occur [7]. Comment seems superfluous. Lack of a duplicate oxygen monitor on a plant in which oxygen and hydrocarbon were mixed led to a violent explosion during the hour-long daily monitor-checking procedure [8]. Use of oxygen instead of compressed air to start a diesel engine, or to clear a blocked petrol pipe, led to violent explosions [9]. In an overview of hazards of industrial air-oxidation processes, they are all identified as inherently hazardous, and the majority extremely so. Particular hazards are detailed, and major incidents involving liquid phase oxidation of cyclohexane or cumene, and vapour phase oxidation of ethylene to the oxide are listed, together with the causes [10]. Further examples of the causes of fires in the last 3 processes, and in the liquid phase oxidation of *p*-xylene to terephthalic acid are given [11]. A technique for liquid-phase oxidation of hydrocarbons in a vertical column containing a bottom water layer and no gas-phase above the hydrocarbon layer is claimed to be free of the danger of detonation [12].

See Fuel gases, Fuels, *also* Hydrogen, all above
and Oil films, below

Buten-3-yne: Oxygen

See Tetracarbonylnickel: Oxygen

See also BITUMEN

Hydrocarbons, Promoters

Barisov, A. A. *et al.*, *Chem. Abs.*, 1984, **101**, 93812

Among other compounds, methyl nitrate, nitromethane, ethyl nitrate and tetrafluorohydrazine function as promoters of spontaneous ignition of mixtures of methane or propane with oxygen and argon.

Hydrogen

1. Bailey, D. J. C., *Chem. & Ind.*, 1954, 492
2. Cook, M. A. *et al.*, *Chem. Eng. News*, 1956, **34**, 4436
3. Weissveiler, A., *Z. Elektrochem.*, 1936, **42**, 499
4. Carty, P., *School Sci. Rev.*, 1978, **59**(208), 517–518
5. Loubeyre, P. *et al.*, *Nature*, 1995, 378, 44
6. Editor's comments

When oxygen was passed through a drying tower containing activated alumina previously used to dry hydrogen, explosions occurred. Nitrogen purging between changing gases would prevent this [1]. During preparation of stoichiometric mixtures in a steel mixing tank (at 17–82 bar, with or without 15% argon), several spontaneous explosions occurred during valve manipulation at 30 min after mixing, but not 10 min after mixing. A possible catalytic effect of the surface of the steel tank was eliminated by coating it thinly with silicone grease [2]. There is a narrow range of concentrations in which the mixture is supersensitive to initiation [3]. A simple and safe method of demonstrating the explosive combustion of stoichiometric hydrogen–oxygen mixtures with spark ignition in a polythene bottle is described [4]. It is claimed that, at ultra-high pressure, hydrogen and oxygen mixtures become non-explosive [5]. Since an extremely small test apparatus is involved and no consideration given to critical diameter or other surface effects, this is open to doubt [6].

Hydrogen sulfide

See Hydrogen sulfide: Oxygen

Lithiated dialkylnitrosamines

Mochizuki, M., *Tetrahedron Lett.*, 1980, **21**, 1766

Oxygenation of lithiated dialkylnitrosamines in THF at -78°C is fast and gives good yields of the *N*-alkyl-*N*-(1-hydroperoxyalkyl)nitrosamines. If oxygenation is too prolonged, poor yields and explosive by-products result.

Mercury, Tetracarbonylnickel

See Tetracarbonylnickel: Mercury, etc.

Metal hydrides

1. Mellor, 1941, Vol. 2, 483
2. Mackay, 1966, 30, 67

Sodium hydride ignites in oxygen at 230°C , and finely divided uranium hydride ignites on contact. Lithium hydride, sodium hydride and potassium hydride react slowly in dry air, while rubidium and caesium hydrides ignite. Reaction is accelerated in moist air, and even finely divided lithium hydride ignites then [1]. Finely divided magnesium hydride, prepared by pyrolysis, ignites immediately in air [2].

See also COMPLEX HYDRIDES

Metals

1. Heinicke, G., *Technik*, 1969, **24**, 313–319
2. Hirano, T. *et al.*, *J. Loss Prev.*, 1993, **6**(3), 151

General data on fire hazards present in industrial oxygen-producing plants are given with details of experimental study of factors involved in combustion of various metals and alloys used in such plant. Safety in selection of materials is discussed [1]. A study of combustion of structural metals (Fe, Al) in oxygen is given [2].

See Oxygen (Liquid): Metals

Methoxy-1,3,5,7-cyclooctatetraene

See Methoxy-1,3,5,7-cyclooctatetraene: Oxygen

4-Methoxytoluene

Imamura, J. *et al.*, Brit. Pat. 1 546 397, 1979

Liquid phase oxidation of 4-methoxytoluene to anisaldehyde at 60 bar/115°C in acetic acid containing heavy metal salts is violent, and must be controlled by the rate of addition of oxygen gas.

Non-metal hydrides

1. Mellor, 1946, Vol. 5, 36
2. Sidgwick, 1950, 344, 553
3. Mellor, 1940, Vol. 6, 220–225

The reported ignition of diborane and tetraborane(10) in contact with air or oxygen is due to the presence of traces of silicon hydrides [1]. The lower members of the latter class ignite or explode in air or oxygen, especially at reduced pressure [2]. Phosphine is also sensitive [3].

See Aluminium tetrahydroborate: Alkenes, Oxygen

Decaborane(14)

Pentaborane(11)

Phosphine: Oxygen

Oil films

Werley, B. M., *ASTM Spec. Tech. Publ.*, 1983, (812), 108–125

The hazards arising from presence of oil films in oxygen-handling systems are reviewed, with consideration of cleanliness specifications, film flammability, film migration and hazard mechanisms.

See Hydrocarbons, above

Organic analytical samples

1. Pack, D. E., *Chem. Eng. News*, 1966, **44**(2), 4
2. Devereaux, H. D., *Chem. Eng. News*, 1966, **44**(6), 4
3. Heysel, R. M. *et al.*, *Chem. Eng. News*, 1966, **44**(10), 4
4. MacDonald, A. M. G., *Chem. Eng. News*, 1966, **44**(19), 7
5. Jenkin, J. B., *Chem. & Ind.*, 1974, 585
6. Guthrie, R. D. *et al.*, *Chem. Brit.*, 1979, **15**, 614; *J. Chem. Educ.*, 1980, **57**(3), 226

Hazards involved in the use of the oxygen flask combustion technique are discussed and illustrated with several examples [1]. Some explosions occurred after completion of combustion, when the oxygen concentration in the flask is still some 75% [2]. Shielding of the flask [3], minimally with wire gauze [4] seems advisable. Explosion during a blank combustion (of a folded filter paper) was attributed possibly to thermal strains. Similar occurrences had been noted previously [5]. Fitting a simple pressure relief (Bunsen) valve to the flask improves safety aspects, and the oxygen flask method is also to be preferred (for qualitative work) to the more hazardous sodium fusion method [6].

See Sodium: Halocarbons, or: Organic analytical samples

Organic materials

Burnett, R. J. *et al.*, *J. Chem. Educ.*, 1985, **62**(5), A157, A159

Facilities and procedures necessary to encourage safety in laboratory work involving oxidation of organic materials with pressurised oxygen are reviewed.

Ozone, Rubber

See Ozone: Oxygen, Rubber

Pentaborane(9)

See Pentaborane(9): Oxygen

Perfluorocarbons

See FLUOROCARBONS: OXYGEN

Phosphine

See Phosphine: Oxygen

Phosphorus(III) oxide

See Tetraphosphorus hexaoxide: Oxygen

Phosphorus tribromide

Bailar, 1973, Vol. 2, 426

Oxidation of the bromide with gaseous oxygen is not easily controlled and becomes explosive.

Plastic cylinder caps

Anon., *Loss Prev. Bull.*, 1978, (020), 49

The plastic caps (or tape) which cover the outlets on refilled oxygen cylinders will burn readily in oxygen and must be completely removed and no bits trapped in the fittings attached to the cylinder outlet.

See Organic materials, above

Plastic tubes

1. Wolf, G. L. *et al.*, *Anaesthesiology*, 1987, **67**, 236–239

2. Davies, P., Personal communication, 1993

Following fires in which endotracheal tubes became ignited by surgical lasers or electrocautery in atmospheres enriched by oxygen and/or nitrous oxide, the flammability of PVC, silicone rubber and red rubber tubes in enriched atmospheres was studied [1]. Ozonised oxygen was reacted with hydrogen at low pressure to generate hydroxyl radicals. Pressure in the apparatus was maintained by a vacuum pump protected from ozone by a tube of heated silver foil. On two occasions there was an explosion in the plastic vent pipe from the vacuum pump. The vent gas should have been outside explosive limits and the exact cause is not clear; the editor suspects peroxide formation.

Polymers

1. *MCA Case History No. 395*
2. *MCA Case History No. 1111*
3. *491M*, 1975, 294

A foam rubber sample, being tested for oxidation resistance under oxygen at 34 bar at 90°C exploded with extreme violence after 4 days [1]. Use of a Neoprene-lined hose in a high-pressure oxygen manifold caused failure and ignition of the burst hose. Possible ignition sources include adiabatic compressive heating, and friction from vibration of metal reinforcing fibres in the high velocity stream of escaping oxygen [2]. Polytetrafluoroethylene (Teflon) ignited at 705°C in oxygen at 0.34 bar pressure when used as wire insulation, while polyolefine insulation ignited at 593°C under the same conditions [3].

Polytetrafluoroethylene, Stainless steel

Anon., *Petroleum Rev.*, 1979, 48

In two test methods (IP40, IP38) for the oxidative stability of hydrocarbons, samples are heated in a valve-sealed stainless bomb under an oxygen atmosphere. On several occasions minor explosions occurred when valves with Teflon sealing discs (rather than the older metal needle closures) were operated during oxygen purging or venting operations. (This was associated with presence of metal particles and traces of grease being embedded in the Teflon discs). Only stainless needle valves, and a controlled rate of pressure release, are recommended for these tests.

See PERFLUOROCARBONS

Propylene oxide

See Propylene oxide: Oxygen

Rhenium

Mellor, 1942, Vol. 12, 471

The metal ignites in oxygen at 300°C.

Rubberised fabric

Anon., *Loss Prev. Bull.*, 1978, (020), 52

Accidental connection of a cylinder of oxygen instead of medical air to an air breathing set led to ignition of the face-piece.

Tetrafluoroethylene

1. Pajaczkowski, A., *Chem. & Ind.*, 1964, 659
2. Ger. Pat. 773 900, 1971

Accidental admixture of oxygen gas with unstabilised liquid tetrafluoroethylene produced a polymeric peroxide which was powerfully explosive, and sensitive to heat, impact or friction [1]. Removal of oxygen by treatment with pyrophoric copper to prevent explosion of tetrafluoroethylene has been claimed [2].

See other POLYPEROXIDES

Tetramethyldisiloxane

1. Anon., *Jahresber.*, 1983, 71–72
2. Plewinsky, B. *et al.*, *Thermochim. Acta*, 1985, **94**, 33–42
3. Plewinsky, B., *Chem. Abs.*, 1990, **113**, 234231

A 4.9 g sample of the liquid siloxane in a glass dish was put into a bomb calorimeter (on an open bench) containing 5 ml of sodium hydroxide solution to absorb combustion gases. The electric igniter system consisted of a metal wire in contact with a cotton-wool wick which dipped into the siloxane sample. The bomb was sealed, pressured up to 39–44 bar with oxygen, and the igniter was fired. A violent explosion blew the lid off the bomb (rated at 190 bar working, 250 bar test), and examination of the deformed bomb indicated that a maximum detonation pressure of around 900 bar had been attained.

Detailed examination of the reaction showed a 2 stage mechanism. Hydrolysis of the volatile partially methylated siloxane by the alkaline solution liberated hydrogen which formed an explosive mixture with oxygen. When the igniter was fired, the hydrogen–oxygen explosion atomised the remaining liquid siloxane leading to a very violent secondary explosion. The wick also played a decisive separate role, as the siloxane-soaked cotton also underwent extremely rapid detonative decomposition on ignition. The rate of pressure rise was measured as 10 kbar/s, with a maximum pressure calculated for a 4.9 g sample of 800 bar. Hexamethyldisiloxane (which can't produce hydrogen by hydrolysis) burned smoothly under the same conditions. It is recommended that no more than 0.5 g samples should be used, with oxygen pressure limited to 20 bar and protective enclosure of the bomb equipment [1]. A more detailed investigation of the combustion of tetramethyldisiloxane in oxygen showed that, even in the absence of hydrogen, combustion was still extremely violent, the rate of pressure increase being inversely proportional to the pressure of oxygen, as expected for a homogeneous gas-phase combustion reaction. The unusually high rate of pressure rise (265 kbar/s, to max 113 bar for 10 bar oxygen pressure) is indicative of detonation. Under heterogeneous conditions, liquid tetramethyldisiloxane gave a maximum rate of pressure rise of 90 kbar/s to max 160 kbar with 30 bar oxygen pressure, with an abrupt change to the higher values between 15 and 20 bar oxygen pressure [2]. Further study showed that detonation-like transmission of combustion across the surface of either a soaked wick, or bulk liquid, could take place with a gas-phase composition below the lower flammability limit [3]. Pentamethyldisiloxane showed generally similar behaviour (maximum rate 38 kbar/s to max 205 bar for 30 bar oxygen pressure), with an abrupt increase between 25 and 30 bar pressure of oxygen [2].

See SILANES; OXYGEN BOMB CALORIMETRY
See entry PRESSURE INCREASE IN EXOTHERMIC DECOMPOSITION

Titanium

Anon., *ABCM Quart. Safety Summ.*, 1962, **33**, 24

Passage of oxygen through a titanium feed pipe into a titanium autoclave caused a titanium–oxygen fire and explosion at 44 bar. When the surface oxide film is damaged, titanium can ignite at 24 bar under static conditions and at 3.4 bar under dynamic conditions, with oxygen at ambient temperature.

See Aluminium–titanium alloys: Oxidants

Titanium trichloride, Vapours of low ignition temperatures

Huneker, K. H., *Chem. Abs.*, 1977, **87**, 15458

The Dräger oxygen-measuring tube contains titanium trichloride as the indicating substance, and a possible hazard in using these tubes to measure oxygen contents above 25 vol% in gas mixtures has been noted. The temperature in the tube can reach 120°C during oxidation of titanium trichloride to titanium dichloride oxide, so the test should not be used if compounds with ignition points below 135°C (e.g. carbon disulfide) are present.

Trirhenium nonachloride

1. Edwards, P. G. *et al.*, *J. Chem. Soc., Dalton Trans.*, 1980, 2467, 2472

2. Edwards, P. G., *Lab. Haz. Bull.*, 1984(4), 3, item 212

During preparation of perrhenyl chloride by combustion of rhenium nonachloride in an oxygen stream [1], a violent explosion, possibly involving chlorine oxides, occurred on heating the chloride to 250°C. A safe procedure involving heating the chloride to 250°C under nitrogen, and then introducing oxygen, is proposed [2].

See other NON-METALS, OXIDANTS

4832. Oxygen (Liquid)

[7782-44-7]

O₂

O₂

HCS 1980, 709

NFPA 50, Quincy (Ma), Natl. Fire Prot. Assoc., 1985

The recent US National Fire Code covers all aspects of equipment, installation and safe operational practices necessary for bulk oxygen storage at consumer sites.

Preparative hazard

Anon., *Sichere Chemiearbeit*, 1994, **46**(8), 94

During regeneration of an adsorber in the cold box of an air separation plant, regeneration gas exited via the adsorber drain pipe. This still contained some liquid oxygen. Very low regeneration-gas flows permitted desorption of acetylene and other hydrocarbons at above normal (ppm) concentrations, these condensed or crystallised into the liquid oxygen and the mixture finally detonated, causing

limited damage. It is recommended that regeneration gas should never be vented via the drain pipe.

See Hydrocarbons, below

Author's comments

Although the entries below refer to the specific hazards observed with a few organic or oxidisable materials, it is probable that most organic materials, inorganic reducing agents, and many unoxidised inorganic materials may be highly hazardous with liquid oxygen under appropriate conditions of contact and initiation. Liquid oxygen also implies a region of oxygen enriched atmosphere near the vent of a storage vessel, or during transfer; this has its own hazards.

See CRYOGENIC LIQUIDS, OXYGEN ENRICHMENT

Acetone

Blau, K., private comm., 1965

Accidental addition of liquid oxygen to vacuum jars containing acetone residues from trap-cooling use caused a violent explosion. Liquid nitrogen is less hazardous as a trap coolant, but only under controlled conditions.

See Nitrogen (Liquid)

Acetylene, Oil

Basurov, Z. B. *et al.*, *Chem. Abs.*, 1960, **54**, 12587a

The detonation capacity of mixtures of acetylene and liquid oxygen is increased by the presence of organic material (oils) in the oxygen. Hazards of accumulation of oil in air-liquefaction and -fractionation plants are emphasised.

See Hydrocarbons, below

Ammonia, Ammonium nitrate, Diphenyl carbonate

Buzard, J. *et al.*, *J. Amer. Chem. Soc.*, 1952, **74**, 2925–2926

During the synthesis of ¹⁵N-labelled urea by interaction of labelled ammonium nitrate, liquid ammonia and diphenyl carbonate in presence of copper powder, a series of explosions of the refrigerated sealed tubes was encountered. This was almost certainly caused by condensation of traces of oxygen in the tubes cooled to –196°C during condensation of ammonia before sealing the tubes. Cooling to –80°C would have been adequate and have avoided the hazard.

See Nitrogen (Liquid)

Asphalt

1. Weber, U., *Chim. et Ind. (Paris)*, 1963, **90**, 178

2. English, W. D. *et al.*, *Chem. Eng. News*, 1992, **70**(10), 18

3. Gayle, B., *Rev. Tech. Feu*, 1979, **20**(189), 62

Mechanical impact on a road surface on to which liquid oxygen had leaked caused a violent explosion [1]. Anecdotal evidence that there have been several like incidents is given [2]. Mixtures of asphalt and liquid oxygen were shown to be impact-sensitive on the small scale, but on the larger scale a detonator was necessary to initiate mild explosion of liquid oxygen on a layer of asphalt [3]. Oil,

rubber or other impurities may have been present on the road surface in the first incident.

Carbon

1. Kirshenbaum, 1965, 4
2. Stettbacher, A., *Spreng-u. Schiess-Stoffe*, 71, Zurich, Racher, 1948
3. Angot, A., *Ann. Mines [13]*, 1934, **5**, 46–51
4. Hempseed, J. W. *et al.*, *Plant/Oper. Progr.*, 1991, **10**(3), 184

Mixtures of carbon and liquid oxygen have been used as blasting explosives for some time [1,2], Mixtures with carbon black appear unusually sensitive to impact, and a blasting cartridge exploded when dropped [3]. Purification of helium containing only 1.4% oxygen and 0.7% nitrogen by passing through a carbon absorber at –195°C led to an explosion of 20 kg TNT equivalent. It was found that a liquid phase of 85% oxygen content was capable of condensing onto the carbon at the entry.

Carbon, Iron(II) oxide

Leleu, *Cahiers*, 1975, (78), 121

Carbon containing 3.5% of the oxide explodes on contact with liquid oxygen.

See other CATALYTIC IMPURITY INCIDENTS

Halocarbons

Anon., *Chem. Eng. News*, 1965, **43**(24), 41

Mixtures of liquid oxygen with dichloromethane, 1,1,1-trichloroethane, trichloroethylene and ‘chlorinated dye penetrants 1 and 2’ exploded violently when initiated with a blasting cap. Carbon tetrachloride exploded only mildly, and a partly fluorinated chloroalkane not at all. Trichloroethylene has been used for degreasing metallic parts before use with liquid oxygen, but is not safe.

See Oxygen (Gas): Halocarbons

Hexafluoropropene, Oxygen difluoride

See Oxygen difluoride: Hexafluoropropene, etc.

Hydrocarbons

1. Mellor, 1941, 1, 379
2. Kirshenbaum, 1956, 17, 22
3. *MCA Case History No. 865*
4. Wright, G. T., *Chem. Eng. Progr.*, 1961, **57**(4), 45–48
5. Rotzler, R. W., *et al.*, *Chem. Eng. Progr.*, 1960, **56**(6), 68–73
6. Matthews, L. G., *Chem. Eng. Progr.*, 1961, **57**(4), 48–49
7. Karwat., E., *Chem. Eng. Progr.*, 1958, **54**(10), 96–101
8. Streng, A. G. *et al.*, *J. Chem. Eng. Data*, 1959, **4**, 127–131
9. Pelman, M., *AD Rep. 768744/5GA*, Springfield (Va.), USNTIS, 1973
10. Gustov, V. F. *et al.*, *Chem. Abs.*, 1984, **100**, 12006

Mixtures of hydrocarbons with liquid oxygen are highly dangerous explosives, not always requiring external initiation. The earliest example must be that of Claudé, who accidentally knocked a lighted candle into a bucket of liquid oxygen

in 1903 and suffered from the violent explosion ensuing [1]. Mixtures with liquid methane or benzene are specifically described as explosive [2]. Traces of oil in a liquid oxygen transfer pump caused the explosion described in the Case History [3]. Mixtures with petroleum and absorbent charcoal have been used experimentally as blasting explosives [1]. Addition of a little aluminium powder to liquid methane–oxygen mixtures increases the explosive power [2]. An explosion in a liquid oxygen evaporator was attributed to the presence of acetylene, arising from unusual plant conditions and higher than usual hydrocarbon concentrations in the atmospheric air taken in for liquefaction [4]. Two similar explosions in main condensers were attributed to presence of acetylene [5] or unspecified hydrocarbons [6]. Suitable precautions are detailed in all cases. Experimental work appeared to implicate ozone as a major contributory factor [7].

The explosive properties of liquid methane–oxygen mixtures were determined [8]. During investigation of an explosion in a portable air liquefaction–separation plant, hydrocarbon oil was found in a silica filtration bed [9]. The mechanism of slow heterogeneous accumulation of hydrocarbons dissolved in trace amounts in liquid oxygen on the liquid evaporator surfaces is discussed. It was concluded that months of continuous evaporation would be required to attain explosion-hazardous levels in real evaporators [10].

Hydrogen (Liquid)

See Hydrogen (Liquid): Oxygen

Liquefied gases

1. Kirshenbaum, 1956, 17, 30, 34
2. Lester, D. H., *Rept. NASA-CR-148711*, Richmond (Va), USNTIS, 1976
Mixtures with liquid carbon monoxide, cyanogen (solidified), and methane are highly explosive [1]. Autoignition in liquid oxygen–hydrogen propellant systems has been reviewed [2].

Lithium hydride

Kirshenbaum, 1956, 44

Mixtures of lithium hydride powder and liquid oxygen are detonable explosives of greater power than TNT.

Metals

1. Anon., *Chem. Eng. News*, 1957, **35**(24), 90
2. Anon., *Chem. Eng. News*, 1957, **35**(31), 10
3. Austin, C. M. *et al.*, *J. Chem. Educ.*, 1959, **36**, 54, 308
4. *MCA Case History No. 824*
5. Kirshenbaum, 1956, 4, 16
6. *MCA Case History No. 988*
7. Strauss, W. A., *Amer. Inst. Aeronaut. Astronaut. J.*, 1968, **6**, 1753
8. Rozovskii, A. S. *et al.*, *Chem. Abs.*, 1979, **91**, 23447
9. Lewis, D. J., *Haz. Cargo. Bull.*, 1983, **4**(11), 28–29
10. Bodner, G. M., *J. Chem. Educ.*, 1985, **62**(12), 1106

A demonstration of combustion of aluminium powder in oxygen exploded violently, probably owing to presence of unevaporated liquid at ignition [1]. Stoichiometric mixtures of the two are explosives much more powerful than TNT [2]. Further details and comments were given later [3]. An aluminium filter in a high-capacity liquid oxygen transfer line exploded violently, possibly owing to friction or impact ignition of an aluminium component in contact with an abrasive particle, which would penetrate the protective oxide layer [4]. Powdered magnesium, titanium or zirconium mixed with liquid oxygen are detonable [5]. A nitrogen-pressurised liquid oxygen dispenser made of titanium alloy failed during nitrogen pressurising. The vessel failed because of reaction of the titanium alloy with liquid oxygen. Titanium is more reactive towards oxygen than either stainless steel or aluminium, and should therefore not be used for oxygen service, either gas or liquid [6]. Mixtures of liquid oxygen with 48–64 wt% of fine aluminium powder are detonable, and the parameters were investigated [7]. Detonation of mixtures with powdered aluminium, iron, titanium and chromium–nickel alloy powders was studied [8]. A new liquid oxygen road tanker constructed of aluminium ruptured violently during manoeuvring operations after a delivery, when the contents had been pressured up with gaseous oxygen (evaporated from the liquid) to expel the liquid. The rupture was traced to erosion and thinning of the tank wall adjacent to an internal baffle joint, where aluminium particles, oil and a halocarbon degreasing solvent were trapped in a cavity in the weld. These had reacted with the pressurising oxygen atmosphere, raising the wall temperature in the vapour space to that where direct reaction of the aluminium container with gaseous oxygen was possible. Some 70 kg of metal was missing in all [9]. Accounts of many further accidents in lecture demonstrations of rapid combustion of aluminium powder–liquid oxygen mixtures have been published [10].

See Aluminum: Oxidants, *also* Magnesium: Oxidants

Organic materials

1. Hauser, R. L. *et al.*, *Adv. Cryog. Eng.*, 1962, **8**, 242–250
2. Clippinger, D. E. *et al.*, *Aircraft. Eng.*, 1959, **32**(365), 204–205

Many common polymers, polymeric additives and lubricants oxidise so rapidly after impact in liquid oxygen that they are hazardous. Of those tested, only acrylonitrile–butadiene, poly(cyanoethylsiloxane), poly(dimethylsiloxane) and polystyrene exploded after impact of 6.8–95 J intensity (5–70 ft.lbf). All plasticisers (except dibutyl sebacate) and antioxidants examined were very reactive. A theoretical treatment of rates of energy absorption and transfer is included [1]. Previously, many resins and lubricants had been examined similarly, and 35 were found acceptable in liquid oxygen systems [2].

1,3,5-Trioxane

Kirshenbaum, 1956, 31

Mixtures with liquid oxygen are highly explosive.

Wood (+ charcoal)

1. Weber, U., *Chim. et Ind. (Paris)*, 1963, **90**, 178
2. Lang, A., *Chem. Eng. Progr.*, 1962, **58**(2), 71

Several major accidents during handling of tonnage amounts of liquid oxygen are described [1]. One involved a violent explosion caused by leakage of liquid oxygen on to metal-encased timber (which previously had been charred) and spark ignition from welding in the oxygen-enriched atmosphere of an air-separation plant [2].

See other CRYOGENIC LIQUIDS, NON-METALS, OXIDANTS

4833. Osmium(IV) oxide

[12036-02-1]

O₂Os

OsO₂

Preparative hazard

Sidgwick, 1950, 1493

The amorphous form, prepared by dehydration at low temperature, is a pyrophoric powder.

See Sodium chlorate: Osmium

See other METAL OXIDES, PYROPHORIC MATERIALS

4834. Lead(IV) oxide (Lead dioxide)

[1209-60-0]

O₂Pb

PbO₂

HCS 1980, 590

Carbon black, Chlorinated paraffin, Manganese(IV) oxide

1. Rasmussen, B., *Unwanted chemical reactions; Appendix B*, RisøI-279, RisøNational Laboratory, Roskilde (DK), 1987; *J. Haz. Mat.*, 1988, **19**, 279–288

2. Author's comments to Dr Rasmussen, 1988

During the established manufacture of an accelerator mixture for a joint-filling composition, carbon black, lead(IV) oxide and manganese(IV) oxide were dispersed in a mixture of viscous chlorinated paraffins in presence of a silicone surfactant by agitation in a 1200 l reactor. The mixture was subsequently milled in a continuously fed shear disk bead mill to reduce the particle size and increase the degree of dispersion to a predetermined value. The bulk temperature of the material was increased to 55°C by the milling operation, and it was returned to the reactor for storage before re-milling, as the particle size was too coarse. After 2 days, exothermic decomposition set in with evolution of hydrogen chloride and probably chlorine also, and the mixture in the reactor eventually ignited. The reactor was eventually quenched with water and was found to contain a solid residue resembling clinker.

Some variations from the normal routine had occurred in the course of preparing this batch, including:- addition of a major portion of the chlorinated paraffin to the reactor which still contained 100–150 kg of a previous batch of accelerator, followed by standing for 16 days over a winter holiday break; use of 20% more

manganese dioxide than usual; the mixture had not dispersed properly during the normal single milling pass, so had to be stored.

Subsequent investigation of the incident was undertaken both by DSC and by both 5 kg and 1 kg simulations of the incident in electrically heated glass or metal reaction vessels. DSC showed that the stabilised chlorinated paraffin began to decompose exothermally only above 300°C, with $\Delta H = 1.25$ kJ/g, while the reaction mixture showed an initial exotherm at 91°C, with the main exotherms above 200°C, and $\Delta H = 3.64$ kJ/g. The 5 kg process simulation experiment showed that non-acidic gas began to be evolved at about 150°C, and no hydrogen chloride was evolved until after the reaction mixture had been heated for several hours at temperatures up to about 200°C (during which period the inhibitor had probably been consumed). Further heating to 220°C initiated a thermal runaway reaction, with temperatures exceeding 300°C and violent evolution of white fumes, which continued (without external heating) for over an hour. The residue was a grey clinker, samples of which showed a relatively much lower lead content and higher manganese content than the original mixture. The addition of ferric chloride to the chlorinated paraffin caused considerable reduction of the thermal stability of the latter.

It was concluded that the reaction mixture was thermally unstable if subjected to thermal initiation, and that temperatures of over 300°C were capable of attainment, when direct reaction between the metal oxides and the chlorinated paraffins or their degradation products were possible. The origin of the heat energy necessary to increase the temperature of the milled batch from 55 to 91°C (the DSC-identified exotherm temperature) could not be identified [1].

However, a reasonable and coherent explanation seems possible if the following assumptions are made.

1. During the 16-day contact of the first 250 kg portion of chlorinated paraffin with the 150 kg residue of previously processed accelerator, the 1% inhibitor content of the former became depleted or exhausted by adsorption and/or reaction, reducing the thermal stability of the liquid phase.

2. During the 2 day storage (apparently without agitation) of the milled reaction mixture, initially at 55°C, the oxidants (mainly lead dioxide) began to react with the carbon black (effectively a very finely divided organic fuel), the 12.5 kg of which could release up to 373 MJ of heat if oxidised fully to carbon dioxide. The heat release may well have not been uniform throughout the batch, probably being concentrated in the bottom of the unstirred batch, where the not fully dispersed lead dioxide ($d = 9.4$) would settle out. This heat release may have been sufficient to increase the temperature, at least locally, to 91°C, when the total exotherm would then have been increased.

3. Once the thermal runaway had become established, the high temperature could allow a 'smelting reaction' (reduction of lead oxides by carbon to metallic lead) to occur, and the low m.p. and high density (327°C, 10.6, respectively) would cause the molten lead to concentrate at the base of the reactor, decreasing the lead content and raising the manganese content of the bulk of the clinker residue, as was observed [2].

See other AGITATION INCIDENTS, INDUCTION PERIOD INCIDENTS, REDOX REACTIONS

Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Hydrogen sulfides

MRH Hydrogen sulfide 0.56/82

Mellor, 1941, Vol. 7, 689; 1947, Vol. 10, 129, 159

Contact of hydrogen sulfide with dry or moist lead dioxide causes attainment of red heat and ignition. Contact of hydrogen trisulfide with the dioxide causes violent decomposition and ignition.

Metal acetylides or carbides

Mellor, 1946, Vol. 5, 850, 872

Interaction with aluminium carbide is incandescent, and with caesium acetylide at 350°C is explosive. Other related compounds may be expected to be oxidised violently if unmoderated.

Metals

MRH Magnesium 3.22/17

1. Mellor, 1963, Vol. 2, Suppl. 2.2, 1571
2. Mellor, 1940, Vol. 4, 272
3. Mellor, 1946, Vol. 5, 217
4. Mellor, 1941, Vol. 7, 658, 691
5. Collvin, P., *Chem. Abs.*, 1982, **96**, 87999

Warm potassium reacts explosively with lead dioxide, and sodium probably behaves similarly [1]. Magnesium reacts violently [2], and powdered aluminium probably does also, as it reacts violently with the monoxide [3]. Mixtures of powdered molybdenum or tungsten with the dioxide incandesce on heating [4]. The combination with zirconium is rated highest on a scale for sensitivity to deflagration, friction, ignition and static electrical initiation [5].

See other STATIC INITIATION INCIDENTS

Metal sulfides

Mellor, 1941, Vol. 3, 745

Calcium sulfide, strontium sulfide or barium sulfide react vigorously with lead dioxide on heating.

Nitroalkanes

See NITROALKANES: metal oxides

Nitrogen compounds

MRH Hydroxylamine 1.92/36

1. Mellor, 1941, Vol. 7, 637
2. Mellor, 1940, Vol. 8, 291

Hydroxylamine ignites in contact with lead dioxide [1], while phenylhydrazine immediately reacts vigorously [2].

Non-metal halides

MRH Phosphorus trichloride 0.67/31

1. Mellor, 1941, Vol. 7, 690
2. Mellor, 1947, Vol. 10, 676, 909

Warm phosphorus trichloride reacts with incandescence [1] and seleninyl chloride vigorously. Sulfonyl dichloride may react explosively [2].

Non-metals

1. Mellor, 1946, Vol. 5, 17
2. Mellor, 1941, Vol. 7, 689–690
3. Gibson, N., *Runaway Reactions*, 1981, Paper 3/R, 6

Boron or yellow phosphorus explode violently on grinding with lead dioxide, while red phosphorus ignites [1]. Mixtures with sulfur ignite on grinding or addition of sulfuric acid [2]. An initiating mixture of silicon and lead dioxide (2:1) attains a temperature around 1100°C after ignition by a small flame [3].

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

Potassium

See Potassium: Metal oxides

Sulfur dioxide

Mellor, 1941, Vol. 7, 689

Interaction is incandescent.

See other METAL OXIDES, OXIDANTS

4835. Palladium(IV) oxide

[12036-04-3]

PdO₂

O₂Pd

Bailar, 1973, Vol. 3, 1279

The hydrated oxide is a strong oxidant, and slowly evolves oxygen at ambient temperature.

See other METAL OXIDES, OXIDANTS *See related* PLATINUM COMPOUNDS

4836. Platinum(IV) oxide

[1314-15-4]

PtO₂

O₂Pt

Acetic acid, Hydrogen

Adams, R. *et al.*, *Org. Synth.*, 1941, Coll. Vol. 1, 66

Addition of fresh platinum oxide catalyst to a hydrogenation reaction in acetic acid caused immediate explosion. Several similar incidents, usually involving acetic acid as solvent, are known to the author.

See other METAL OXIDES, OXIDANTS, PLATINUM COMPOUNDS

4837. Sulfur dioxide
[7446-09-5]

SO₂

O₂S

(MCA SD-52, 1953); HCS 1980, 876 (cylinder)

Preparative hazard

See Sulfuric acid: Copper

Barium peroxide

See Barium peroxide: Non-metal oxides

Boranes

See BORANES: aluminium chloride, sulfur dioxide

Caesium azide

See Caesium azide: Sulfur dioxide

Diethylzinc

See Diethylzinc: Sulfur dioxide

Halogens, or Interhalogens

MRH values show % of interhalogen

See Bromine pentafluoride: Acids, etc.

MRH 1.72/53

Chlorine trifluoride: Metals, etc.

MRH 2.43/74

Fluorine: Non-metal oxides

MRH 5.10/64

Lithium acetylide–ammonia

See Monolithium acetylide–ammonia: Gases, etc.

Lithium nitrate,

MRH 1.00/30

Propene

MRH 2.18/30

See Propene: Lithium nitrate, etc.

Metal acetylides

Mellor, 1946, Vol. 5, 848–849

Monocaesium acetylide or monopotassium acetylide, and the ammoniate of monolithium acetylide all ignite and incandesce in unheated sulfur dioxide. The dimetal derivatives including sodium acetylide appear to be less reactive, needing heat before ignition occurs.

Metal oxides

MRH Iron(II) oxide 1.21/40

1. Mellor, 1941, Vol. 2, 487

2. Mellor, 1942, Vol. 13, 715

3. Mellor, 1941, Vol. 7, 388, 689

Caesium oxide [1], iron(II) oxide [2], tin oxide and lead(IV) oxide [3] all ignite and incandesce on heating in the gas.

Metals MRH Chromium 3.39/52, manganese 2.89/46, sodium 4.56/26

1. Mellor, 1943, Vol. 11, 161
2. Mellor, 1942, Vol. 12, 187
3. Gilbert, H. N., *Chem. Eng. News*, 1948, **26**, 2604
4. Mellor, 1961, Vol. 2, Suppl. 2.1, 468

Finely divided (pyrophoric) chromium incandesces in sulfur dioxide [1], while pyrophoric manganese burns brilliantly on heating in the gas [2]. Molten sodium reacts violently with the dry gas or liquid, while the moist gas reacts as vigorously as water with cold sodium [3].

Other reactants

Yoshida, 1980, 267

MRH values calculated for 12 combinations with various materials are given.

Peat

Byrne, P. J., *Chem. Abs.*, 1976, **104**, 21731

Treatment of peat stockpiles with sulfur dioxide surprisingly accelerated the self-heating process.

Polymeric tubing

MCA Case History No. 1044

Plastics tubing normally capable of withstanding an internal pressure of 7 bar failed below 2 bar when used to convey gaseous sulfur dioxide.

Potassium chlorate

MRH 1.80/44

See Potassium chlorate: Sulfur dioxide

Silver azide

See Silver azide: Sulfur dioxide

Sodium hydride

MRH 3.72/23

See Sodium hydride: Sulfur dioxide

See *other* NON-METAL OXIDES, REDUCANTS

4838. Selenium dioxide

[7446-08-4]

O₂Se

SeO₂

HCS 1980, 814

Preparative hazard

See Selenium: Oxygen, Organic matter

1,3-Bis(trichloromethyl)benzene

See 1,3-Bis(trichloromethyl)benzene: Oxidants

Phosphorus trichloride

Mellor, 1940, Vol. 8, 1005

A mixture of the cold components attains red-heat.
See related METAL OXIDES *See other* OXIDANTS

4839. Silicon dioxide (Silica)
[7631-86-9]

O₂Si

SiO₂

HCS 1980, 819

Hydrochloric acid

MCA Case History No. 1857

A glass drying trap on a hydrochloric acid storage tank was filled with silica gel instead of the calcium sulfate specified. The glass trap fractured, probably owing to thermal shock from the much higher heat of adsorption of water and hydrogen chloride on silica gel.

See other GLASS INCIDENTS

Metals

See Magnesium: Silicon dioxide

Sodium: Non-metal oxides

Oxygen difluoride

See Oxygen difluoride: Adsorbents

Ozone

See Ozone: Silica gel

Silicone liquid, Unstated salt

See SILICONE LIQUID: SILICA, ETC.

Steel

These two anciently served as a means of ignition. They still can. Although the sparking of flint and steel is strictly a mechanical phenomenon, any subsequent fires will not be. Other forms of silica still serve as igniters to this day, via the piezoelectric effect.

See GRAVEL

Vinyl acetate

See Vinyl acetate: Desiccants

Xenon hexafluoride

See Xenon hexafluoride: Silicon dioxide

See other NON-METAL OXIDES

4840. Tin(IV) oxide
[18282-10-5] **O₂Sn**

SnO₂

Chlorine trifluoride
See Chlorine trifluoride: Metals, etc.

Hydrogen trisulfide
See Hydrogen trisulfide: Metal oxides

Metals
See Aluminium: Metal oxides
Magnesium: Metal oxides
Potassium: Metal oxides
Sodium: Metal oxides
See other METAL OXIDES

4841. Strontium peroxide
[1314-18-7] **O₂Sr**

SrO₂

Organic materials
Haz. Chem. Data, 1975, 272
Mixtures of the peroxide with combustible organic materials very readily ignite on friction or contact with moisture.
See other METAL PEROXIDES

4842. Titanium(IV) oxide (Titanium dioxide)
[13463-67-7] **O₂Ti**

TiO₂

HCS 1980, 903

Metals
Mellor, 1941, Vol. 7, 10, 44; 1961, Vol. 2, Suppl. 2.1, 81
Reduction of the oxide by aluminium, calcium, magnesium, potassium, sodium or zinc is accompanied by more or less incandescence (lithium, magnesium and zinc especially).
See other METAL OXIDES

4843. Uranium(IV) oxide
[1344-57-6] **O₂U**

UO₂

Mellor, 1942, Vol. 12, 42

1865

The finely divided oxide prepared at low temperature is pyrophoric when heated in air, and burns brilliantly.

See other METAL OXIDES, PYROPHORIC MATERIALS

4844. Tungsten(IV) oxide
[12036-22-5]



O₂W

Chlorine

See Chlorine: Tungsten dioxide

See other METAL OXIDES

4845. Zinc peroxide
[1314-22-3]



O₂Zn

Alone, or Metals

1. Mellor, 1940, Vol. 4, 530

2. Sidgwick, 1950, 270

The hydrated peroxide (of indefinite composition) explodes at 212°C [1], and mixtures with aluminium or zinc powders burn brilliantly [2].

See other METAL PEROXIDES

4846. Ozone (Trioxygen)
[10028-15-6]



O₃

HCS 1980, 710 (cyl., dissolved gas); RSC Lab. Hazards Data Sheet No. 66, 1987

1. Sidgwick, 1950, 860

2. Adley, F. E., *Nucl. Sci. Abs.*, 1962, **17**, 18

3. Streng, A. G., *Explosivstoffe*, 1960, **8**, 225

4. Clough, P. N. *et al.*, *Chem. & Ind.*, 1966, 1971

5. Gatwood, G. T. *et al.*, *J. Chem. Educ.*, 1969, **46**, A103

6. Loomis, J. S. *et al.*, *Nucl. Instr. Methods*, 1962, **15**, 243

7. Brereton, S. J., *Fusion Technol.*, 1988, **15**(2 pt.23), 833

8. *Ozone Reactions with Organic Compounds*, ACS 112, Washington, ACS, 1972

9. Walker, K. G., *Chem. Brit.*, 1987, **23**, 839

Ozone is strongly endothermic (ΔH_f° (g) +142.2 kJ/mol, 2.96 kJ/g) and the pure solid or liquid materials are highly explosive. Evaporation of a solution of ozone in liquid oxygen causes ozone enrichment and ultimately explosion [1]. Organic liquids and oxidisable materials dropped into liquid ozone will also cause explosion of the ozone [2]. Ozone technology and hazards have been reviewed [3], a safe process to concentrate ozone by selective adsorption on silica gel at low

temperatures has been described [4], and safe techniques for laboratory generation and handling of ozone have been detailed [5]. Ozone has become available as a dissolved gas in cylinders. Explosive hazards involved in use of liquid nitrogen as coolant where ozone may be incidentally produced during operation of van der Graaff generators [6] and radiation sources [7], etc. have been discussed. The chemistry of interaction with organic compounds has been extensively reviewed [8]. Methods and equipment for generation of ozone (up to 13%) in oxygen and storage at high pressure and low temperature for extended periods are described [9].

See Silica gel, below

Acetylene

Grignard, 1935, Vol. 3, 166

Passage of ozone (endothermic oxidant) into acetylene (endothermic reductant) leads to a violent explosion when 50 mg/l of ozone is present.

See other REDOX REACTIONS

Alkenes

1. Sidgwick, 1950, 862

2. Davies, 1961, 97

3. Stull, 1977, 21

Interaction of alkenes with ozonised oxygen tends to give several types of products or their polymers, some of which show more pronounced explosive tendencies than others [1]. The cyclic *gem*-diperioxides are more explosive than the true ozonides [2]. It has been calculated that ozonisation of the endothermic *trans*-stilbene ($\Delta H_f^\circ +135.4$ kJ/mol, 0.78 kJ/g) would give, in the event of decomposition of the unstable ozonide, an exothermic release of 1.41 kJ/g which would attain an adiabatic decomposition temperature approaching 750°C with a 27-fold pressure increase in a closed vessel [3].

See CYCLIC PEROXIDES

Alkylmetals

Lee, H. U. *et al.*, *Combust. Flame*, 1975, **24**, 27–34

The spectra of flames from spontaneous interaction of ozone with dimethylzinc and diethylzinc in a flow system have been studied.

Ammoniacal vapours

1. Bond, 1991, 31

2. Anon., *Fire Prevention*, 1986, (191), 45

Vent air from an animal facility was deodourised by ozone treatment; an extensive fire was started by cutting into the polypropylene ducting carrying the treated air to atmosphere. This was attributed to formation of ammonium nitrate by ammonia oxidation, which mixed with organic dusts also present in the vent-line [1]. A more detailed account of the incident reveals that initial ignition was probably by sawing into metal. Ozone was also found to have degraded the polypropylene pipework, increasing flammability (peroxide formation?) [2].

See Ammonium nitrate

Aromatic compounds

Mellor, 1941, Vol. 1, 911

Benzene, aniline and other aromatic compounds give explosive gelatinous ozonides, among other products, on contact with ozonised oxygen.

Benzene, Oxygen, Rubber

Morrell, S. H., private comm., 1968

During ozonisation of rubber dissolved in benzene, an explosion occurred. This seems unlikely to have been owing to formation of benzene triozone (which separates as a gelatinous precipitate after prolonged ozonisation), since the solution remained clear. A rubber ozonide may have been involved, but the benzene–oxygen system itself has high potential for hazard.

Bromine

Lewis, B. *et al.*, *J. Amer. Chem. Soc.*, 1931, **53**, 2710

Interaction becomes explosive above 20°C and a minimum critical pressure.

See Hydrogen bromide, below, *also* Bromine trioxide

Charcoal, Potassium iodide

Anon., *Lab. Haz. Bull.*, 1983(9), 3, item 660

Carbon impregnated with potassium iodide was used as an ozone-scrubbing filter in a chemiluminescence NO_x analyser. When the level of iodide was inadvertently increased to the high level of 40%, the filter exploded violently during replacement after use. This was attributed to oxidation of iodide to iodate by ozone, and frictional initiation of the iodate–carbon mixture when the filter was dismantled.

Citronellic acid

Yost, Y., *Chem. Eng. News*, 1977, **55**(21), 4

During work-up of the products of ozonolysis of *R*- and *S*-citronellic acids, a substantial quantity of the highly explosive trimeric acetone peroxide (3,3,6,6,9,9-hexamethyl-1,2,4,5,7,8-hexoxonane) was unwittingly isolated by distillation at 105–135°C to give the solid m.p. 95°C. The peroxide appears to have been produced by ozonolysis of the isopropylidene group in citronellic acid, and presumably the same could occur when any isopropylidene group is ozonised. Appropriate care is advised.

Combustible gases

1. Streng, A. G., *Chem. Abs.*, 1964, **55**, 8862c

2. Nomura, Y. *et al.*, *Chem. Abs.*, 1952, **46**, 4234a

Carbon monoxide and ethylene admitted into contact with ozone via an aluminium tip ignited and burned smoothly, while normal contact of carbon monoxide, nitrogen oxide, ammonia or phosphine causes immediate explosion at 0° or –78°C [1]. At pressures below 10 mbar, contact with ethylene is explosive at –150°C [2]. See Nitrogen oxide, below

trans-2,3-Dichloro-2-butene

1. Griesbaum, K. *et al.*, *J. Amer. Chem. Soc.*, 1976, **98**, 2880

2. Gäb, S. *et al.*, *J. Org. Chem.*, 1984, **49**, 2711–2714

The crude products of ozonolysis at -30°C of the chloroalkene tended to decompose explosively on warming to ambient temperature, particularly in absence of solvents. The products included the individually explosive compounds acetyl 1,1-dichloroethyl peroxide, 3,6-dichloro-3,6-dimethyl-2,3,5,6-tetraoxane and diacetyl peroxide [1]. Ozonolysis in ethyl formate saturated with hydrogen chloride gives a high yield of 1,1-dichloroethyl hydroperoxide as a further unstable intermediate product [2].

See OZONIDES

Dicyanogen

Anon., *Chem. Eng. News*, 1957, **35**(21), 22

This hypergolic combination of two extremely endothermic compounds appears to have rocket-propellant capabilities.

See ROCKET PROPELLANTS

Dienes, Oxygen

Griesbaum, K. *et al.*, *Chem. Eng. News*, 1982, **60**(8), 63

Treatment of isoprene, 2,3-dimethylbutadiene or cyclopentadiene in pentane at -78°C with a high ratio ozone/oxygen stream led to immediate ignition and flames in both liquid and gas phases.

Diethyl ether

Mellor, 1941, Vol. 1, 911

Contact of ether with ozonised oxygen produces some of the explosive diethyl peroxide.

1,1-Difluoroethylene

1. Gillies, C. W. *et al.*, *J. Phys. Chem.*, 1979, **83**, 1545

2. Agopovich, J. W. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 5049

The product of ozonolysis of 1,1-difluoroethylene is a shock-sensitive peroxidic species [1], which reacts vigorously with aqueous potassium iodide [2].

Ethylene

Mellor, 1941, Vol. 1, 911

Interaction may be explosive, owing to the very low stability of ethylene ozonide.

See Combustible gases, above, *also* Ethylene ozonide

Ethylene, Formyl fluoride

Mazur, U. *et al.*, *J. Org. Chem.*, 1979, **44**, 3188

Two attempts to ozonise mixtures of ethylene and formyl fluoride in absence of solvent led to explosions.

Fluoroethylene

See Fluoroethylene ozonide

Hydrogen (liquid)

Schwab, G. M., *Umschau*, 1922, 538–539

Liquid hydrogen and solid ozone form very powerfully explosive mixtures.

Hydrogen, Oxygen difluoride

Boehm, R., *Chem. Abs.*, 1975, **83**, 82254

The use of liquid oxygen difluoride (40 or 90%) to stabilise liquid ozone as oxidant for gaseous hydrogen in a rocket motor was not entirely successful, explosions occurring at both concentrations.

See Oxygen fluorides, below

Hydrogen bromide

Lewis, B. *et al.*, *J. Amer. Chem. Soc.*, 1931, **53**, 3565

Interaction is rapid and becomes explosive above a total pressure of ~40 mbar, even at -104°C .

See Bromine, above

4-Hydroxy-4-methyl-1,6-heptadiene

See 4-Hydroxy-4-methyl-1,6-heptadiene: Ozone

3-Hydroxy-2,2,4-trimethyl-3-pentenoic acid lactone

Perrin, C. L. *et al.*, *J. Org. Chem.*, 1980, **45**, 1706

During work-up after ozonolysis of the lactone to give dimethylmaleic anhydride, removal of excess ozone by a stream of oxygen leads to the precipitation of peroxidic material at the concentration used in the preparation.

Isopropylidene compounds

See Citronellic acid, above

2-Methyl-1,3-butadiene

See Dienes, above, *also* 2-Methyl-1,3-butadiene: Ozone

Nitrogen

Strakhov, B. V. *et al.*, *Chem. Abs.*, 1963, **58**, 5069a

The explosive oxidation of nitrogen in admixture with ozone in metallic vessels has been studied.

Nitrogen oxide

Nomura, Y. *et al.*, *Sci. Reps. Tohoku Univ.*, 1950[A], **2**, 229–232

Violent explosions occurred in mixtures at -189°C .

See Combustible gases, above

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Oxygen, Rubber powder

Anon., *Jahresber.*, 1983, 72

In an attempt to introduce carboxyl groups to improve the adhesive properties of rubber, 6 l of powdered rubber (0.1–0.5 mm particle size) in an 8 l flask was treated with ozonised oxygen. The gas stream, containing 5% of ozone was led to the bottom of the flask via a dip tube at the rate of 4 l/min for 2 min, when

treatment was discontinued and the flask closed. After 5 min a violent explosion occurred. This was attributed to formation of ozonides, but exothermic interaction of the high surface area rubber with almost pure oxygen under virtually adiabatic conditions may also have been involved.

See Oxygen: Polymers

Oxygen fluorides

Streng, A. G., *Chem. Rev.*, 1963, **63**, 613, 619

Mixtures with dioxygen difluoride explode at -148°C , and with 'dioxygen trifluoride' (probably mixed dioxygen di- and tetra-fluorides) at -183°C .

Silica gel

Cohen, Z. *et al.*, *J. Org. Chem.*, 1975, **40**, 2142, footnote 6

Silica gel at -78°C adsorbs 4.5 wt% of ozone, and below this temperature the concentration increases rapidly. At below -112°C ozone liquefies and there is a potential explosion hazard at temperatures below -100°C if organic material is present.

See reference 3, main entry

Stibine

Stock, A. *et al.*, *Ber.*, 1905, **38**, 3837

Passage of oxygen containing 2% of ozone through stibine at -90°C caused an explosion. On standing, a suspension of solid stibine in a liquid oxygen solution of ozone eventually exploded, as oxygen evaporated increasing the concentration of ozone and the temperature.

Tetrafluorohydrazine

See Tetrafluorohydrazine: Ozone

Tetramethylammonium hydroxide

Solomon, I. J. *et al.*, *J. Amer. Chem. Soc.*, 1960, **82**, 5640

During interaction to form tetramethylammonium ozonide, constant agitation and slow ozonisation were necessary to prevent ignition.

Trifluoroethylene

Agopovich, J. W. *et al.*, *J. Amer. Chem. Soc.*, 1983, **105**, 5048

After ozonisation of trifluoroethylene in chlorotrifluoromethane solution at -95°C , the reactor must only be allowed to warm slightly (to -88°C) during subsequent cryogenic distillation. When allowed to warm towards ambient temperature the reactor exploded violently. The residue from distillation is peroxidic and probably sensitive to sudden temperature increases.

Unsaturated acetals

Schuster, L., *Ger. Offen.* 2 514 001, 1976

Ozonolysis of unsaturated acetals at -70 to 0°C to give glyoxal monoacetals is uneventful in organic solvents, but leads to explosions in aqueous solutions.

See other NON-METALS, OXIDANTS

4847. Phosphorus(III) oxide
[1314-24-5]



See Tetraphosphorus hexaoxide

4848. Palladium(III) oxide
[80680-07-5]



Sidgwick, 1950, 1573

If the hydrated oxide is heated to remove water, it incandescens or explodes, giving the monoxide.

See other METAL OXIDES See related PLATINUM COMPOUNDS

4849. Sulfur trioxide
[7446-11-9]



HCS 1980, 880

The metastable liquid form boils at 44°C, the more stable solid alpha form melts at 62°C and has 25–30 J/g heat of fusion. This permits pressure generation when proceeding in either direction through the phase change.

Acetonitrile, Sulfuric acid

See Acetonitrile: Sulfuric acid, etc.

Chlorinated solvents

Gilbert, E. E. *Chem. Eng. News* 1989 **67**(41), 2

Details of exothermic and gas-evolving reactions are given, storage of such mixtures is not recommended.

Cyclohexanone oxime, Sulfuric acid

See Cyclohexanone oxime: Oleum

Dimethylformamide

See Sulfur trioxide–dimethylformamide

Dimethyl sulfoxide

Seel, F., *Inorg. Synth.*, 1947, **2**, 174

Dissolution of sulfur trioxide in the sulfoxide is very exothermic and must be done slowly with cooling to avoid decomposition.

Dioxane

Sisler, H. H. *et al.*, *Inorg. Synth.*, 1947, **2**, 174

Since the 1:1 addition complex sometimes decomposes violently on storing at ambient temperature, it should only be prepared immediately before use (if refrigerated storage is not possible).

Dioxygen difluoride

See Dioxygen difluoride: Sulfur trioxide

Diphenylmercury

See Diphenylmercury: Chlorine monoxide, etc.

Ethylene oxide

See Ethylene oxide: Pyridine, Sulfur dioxide

Formamide, Iodine, Pyridine

See Formamide: Iodine, etc.

Metal oxides

1. Partington, 1967, 706
2. Mellor, 1941, Vol. 7, 654

The violent interaction with sulfur trioxide causes incandescence with barium oxide [1] and lead oxide [2].

Nitryl chloride

See Nitryl chloride: Inorganic materials

Organic materials, Water

Unremembered suppliers brochure, 1979

Organic materials, particularly if fibrous with adsorbed or absorbed moisture present, may char or ignite in contact with the stabilised liquid form because of the very high heat of hydration (2.1 kJ/g) and formation of hot oleum which then functions as an oxidant.

See ACID ANHYDRIDES

Other reactants

Yoshida, 1980, 128

MRH values calculated for 18 combinations with various materials are given.

Phosphorus

See Phosphorus: Non-metal oxides

Sulfuric acid

HCS 1980, 702 (65%), 703 (30%), 704 (20%)

Solutions of sulfur trioxide in conc. sulfuric acid (oleum) are available with various proportions of dissolved sulfur trioxide, and the 3 entries in *Handling Chemicals Safely 1980* deal with the properties and hazards of solutions with 65, 30 and 20% of sulfur trioxide, respectively. As expected, the reactivity and oxidising power increase with the content of sulfur trioxide.

Tetrafluoroethylene

1. Anderson, A. W., *Chem. Eng. News*, 1971, **49**(22), 3
2. Xu, B. *et al.*, *Chem. Abs.*, 1982, **97**, 182248
3. Aihara, R. *et al.*, *Chem. Abs.*, 1986, **101** 05, 99522

Reaction to give tetrafluorooxathietane 2,2-dioxide (tetrafluoroethane sultone) had been used industrially and uneventfully, but reaction with excess sulfur trioxide may cause explosive decomposition to carbonyl fluoride and sulfur dioxide [1]. An incident involving the same explosion hazard was reported 11 years later [2]. Use of inert gas to prevent explosion has been patented [3].

Water

1. Mellor, 1947, Vol. 10, 344
2. Willis, P. B., *Chem. Eng. News*, 1986, **64**(38), 2
3. Eccles, E. J., *Chem. Eng. News*, 1987, **65**(12), 2
4. Rosin, J., *Chem. Eng. News*, 1987, **65**(24), 2

Interaction is vigorously exothermic, sometimes explosive, with evolution of light and heat. A 4:1 mixture of oxide and water completely vaporises, with simultaneous emission of light. Unused portions of ampoules of sulfur trioxide were prepared for disposal by dripping a little water each day during several weeks into the ampoules. When inverted to empty the contents, one such ampoule exploded, scattering conc. sulfuric acid around [2]. This technique probably produces a layer of water above one of conc. sulfuric acid, and when the layers mixed the exotherm burst the narrow-necked ampoule. Moderate quantities of liquid sulfur trioxide (100–500 g) may be disposed of into a strong water stream discharging into an acid-proof drain, using personal protection [3]. It is, however, stressed that all disposal methods involving addition to water are inherently unsafe, because sulfur trioxide is insoluble in water and also highly hydrophobic, so the potential for sudden and violent reaction always exists. The safest method is to dissolve sulfur trioxide in a large volume of conc. sulfuric acid, and then dispose of the weak oleum by slow addition to a large and well stirred volume of water [4].

See other ACID ANHYDRIDES, NON-METAL OXIDES, OXIDANTS

4850. Sulfur trioxide–dimethylformamide

[29584-42-7]

$O_3S.C_3H_7NO$

$SO_3.Me_2NCO.H$

Koch-Light Laboratories Ltd., private comm., 1976

A bottle of the complex exploded in storage. No cause was established, but diffusive ingress of moisture to form sulfuric acid, and subsequent hydrolysis of the solvent with formation of carbon monoxide appear to be possible contributory factors.

See related ACID ANHYDRIDES, NON-METAL OXIDES

4851. Antimony(III) oxide
[1309-64-4]

O:SbOSb:O

O₃Sb₂

HCS 1980, 164

Mellor, 1939, Vol. 9, 425

The powdered oxide ignites on heating in air.

Bromine trifluoride

See Bromine trifluoride: Antimony trichloride oxide

Chlorinated rubber

See CHLORINATED RUBBER: metal oxides

See other METAL OXIDES

4852. Selenium trioxide
[13768-86-0]

SeO₃

O₃Se

Organic materials

Sorbe, 1968, 124

A powerful oxidant which forms explosive mixtures with organic materials.

See other NON-METAL OXIDES

4853. Tellurium trioxide
[13451-18-8]

TeO₃

O₃Te

Metals, or Non-metals

Bailar, 1973, Vol. 2, 971

The yellow α -form is a powerful oxidant, reacting violently when heated with a variety of metallic and non-metallic elements.

See other NON-METAL OXIDES

4854. Thallium(III) oxide
[1314-32-5]

O:TIOTI:O

O₃Tl₂

Sulfur, or Sulfur compounds

Mellor, 1946, Vol. 5, 434

Mixtures of the oxide with sulfur or antimony trisulfide explode on grinding in a mortar. Dry hydrogen sulfide ignites, and sometimes feebly explodes, over thallium oxide.

See other METAL OXIDES

4855. Vanadium(III) oxide
[1314-34-7]

O:VOV:O

O₃V₂

Sidgwick, 1950, 825
It ignites on heating in air.
See other METAL OXIDES

4856. Tungsten(VI) oxide
[1314-35-8]

WO₃

O₃W

Boron tribromide
See Boron tribromide: Tungsten trioxide

Bromine, Tungsten
See Bromine: Tungsten, etc.

Interhalogens
See Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Metals, etc.

Lithium
See Lithium: Metal oxides
See other METAL OXIDES

4857. Xenon trioxide
[13776-58-4]

XeO₃

O₃Xe

1. Malm, J. G. *et al.*, *Chem. Rev.*, 1965, **65**, 199
2. Chernick, C. L. *et al.*, *J. Chem. Educ.*, 1966, **43**, 619; *Inorg. Synth.*, 1968, **11**, 206, 209
3. Anon., *Chem. Eng. News*, 1963, **41**(13), 45
4. Holloway, J. H., *Talanta*, 1967, **14**, 871–873
5. Shackelford, S. A. *et al.*, *Inorg. Nucl. Chem. Lett.*, 1973, **9**, 609, footnote 9
6. Shackelford, S. A. *et al.*, *J. Org. Chem.*, 1975, **40**, 1869–1875
7. Foropoulos, J. *et al.*, *Inorg. Chem.*, 1982, **21**, 2503–2504

It is a powerful explosive [1] produced when xenon tetrafluoride or xenon hexafluoride are exposed to moist air and hydrolysed. Some tetrafluoride is usually present in xenon difluoride, so the latter is potentially dangerous. Although safe to handle in small amounts in aqueous solution, great care must be taken to avoid solutions drying out, e.g. around ground stoppers [2]. Full safety precautions have been discussed [2,3,4]. Precautions necessary for use of aqueous solutions of the trioxide as an epoxidation reagent are detailed [5,6]. A safe method of preparation,

so far free from explosions, from xenon hexafluoride and difluorophosphoric acid at 0°C has been described [7].

See other NON-METAL OXIDES, OXIDANTS, XENON COMPOUNDS

4858. Osmium(VIII) oxide

[20816-12-0]



O₄Os

Hydrogen peroxide

See Hydrogen peroxide: Metals, etc.

1-Methylimidazole

Nielsen, A. J., *J. Chem. Soc., Dalton Trans.*, 1979, 1084–1088


An explosion occurred when the oxidant was added to the imidazole.

See other METAL OXIDES, OXIDANTS

4859. Tetraphosphorus tetraoxide trisulfide

[]

O₄P₄S₃



Water

Mellor, 1971, Vol. 8, Suppl. 3, 435

It ignites if moistened with water.

See related NON-METAL OXIDES, NON-METAL SULFIDES

4860. Lead sulfate

[7446-14-2]



O₄PbS

Potassium

See Potassium: Oxidants

See other METAL OXONON-METALLATES

4861. Dilead(II)lead(IV) oxide

[1314-41-6]



O₄Pb₃

HCS 1980, 808

Dichloromethylsilane

See Dichloromethylsilane: Oxidants

Eosin lead salts

Anon., *Chem. & Ind.*, 1995, (22), 912

1877

The mixture of the lead salts of eosin and red lead formerly used as a pigment, proved to be a very easily ignited pyrotechnic mix.

Peroxyformic acid

See Peroxyformic acid: Metals, etc.

Seleninyl chloride

See Seleninyl chloride: Metal oxides

2,4,6-Trinitrotoluene

See 2,4,6-Trinitrotoluene: Added impurities

See *other* METAL OXIDES, OXIDANTS

4862. Ruthenium(VIII) oxide

[20427-56-9]

O₄Ru

RuO₄

1. Mellor, 1942, Vol. 15, 520
2. Brauer, 1965, Vol. 2, 1600
3. Bailar, 1973, Vol. 3, 1194
4. Ullmann, 1992, Vol. A21, 125/6

The liquid (which shows no true b.p.) is said to decompose explosively either above 106° or above 180°C [1,2,3]. The oxide should not be stored in bulk, either as liquid or solid [4].

Ammonia

Watt, J., *J. Inorg. Nucl. Chem.*, 1965, **27**, 262

Rapid interaction at -70°C/1 mbar causes ignition. Slow interaction produced a solid which exploded at 206°C, probably owing to formation of ruthenium amide oxide or ruthenium nitride.

Hydriodic acid

Mellor, 1942, Vol. 15, 520

Reaction with the acid to give ruthenium triiodide is explosively violent.

Organic materials

1. Mellor, 1942, Vol. 15, 517, 520
2. Brauer, 1965, Vol. 2, 1600

The solid oxide, or its concentrated solutions or vapour, tends to oxidise ethanol, cellulose fibres, etc., explosively [1,2]. Ethanol also gives an explosive by-product, probably an organoruthenium derivative [1].

Phosphorus tribromide

Mellor, 1971, Vol. 8, Suppl. 3, 521

Interaction is vigorously exothermic.

Sulfur

Mellor, 1942, Vol. 15, 520

The vapour explodes in contact with sulfur.

See other METAL OXIDES, OXIDANTS

4863. Xenon tetraoxide

[12340-14-6]



O_4Xe

Selig, H. *et al.*, *Science* (New York), 1964, **143**, 1322

Decomposition may be explosive, even at -40°C .

See other NON-METAL OXIDES, OXIDANTS, XENON COMPOUNDS

4864. Phosphorus(V) oxide

[1314-56-3]



O_5P_2

See Tetraphosphorus decaoxide

4865. Tantalum(V) oxide

[1314-61-0]



O_5Ta_2

Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

See other METAL OXIDES

4866. Vanadium(V) oxide

[1314-62-1]



O_5V_2

HCS 1980, 950

Calcium, Sulfur

491M, 1975, 81

A mixture of the metal and oxidant, contaminated with sulfur and some water, led to a serious fire.

Chlorine trifluoride

See Chlorine trifluoride: Metals, etc.

Lithium

See Lithium: Metal oxides

MRH 2.43/7

1879

Other reactants

Yoshida, 1980, 118

MRH values calculated for 9 combinations, largely with oxidisable materials, are given.

Peroxyformic acid

MRH 5.69/99+


See Peroxyformic acid: Metals, etc.

See other METAL OXIDES, OXIDANTS

4867. Tetraphosphorus hexaoxide (Phosphorus(III) oxide)

[10248-58-5]

O₆P₄



Ammonia

Thorpe, T. E. *et al.*, *J. Chem. Soc.*, 1891, **59**, 1019

Interaction of the molten oxide and ammonia under nitrogen is rather violent and the mixture ignites. Use of a solvent and cooling controls the reaction to produce diamidophosphinic acid.

Disulfur dichloride

Mellor, 1940, Vol. 8, 898

Interaction is very violent.

Halogens

1. Thorpe, T. E. *et al.*, *J. Chem. Soc.*, 1891, **59**, 1019

2. Mellor, 1940, Vol. 8, 897

The oxide ignites in contact with excess chlorine gas [1], and reacts violently, usually igniting, with liquid bromine [2].

Organic liquids

1. Thorpe, T. E. *et al.*, *J. Chem. Soc.*, 1890, **57**, 569–573

2. Mellor, 1971, Vol. 8, Suppl. 3, 382

The oxide ignites immediately with ethanol at ambient temperature [1]. The liquid oxide (above 24°C) reacts very violently with methanol, dimethylformamide, dimethyl sulfite or dimethyl sulfoxide (also with arsenic trifluoride) and charring may occur [2].

Oxygen

Mellor, 1940, Vol. 8, 897–898; 1971, Vol. 8, Suppl. 3, 380–381

Interaction with air or oxygen is rapid and, at slightly elevated temperatures in air or at high concentration of oxygen, ignition is very probable, particularly if the oxide is molten (above 22°C) or distributed as a thin layer. The solid in contact with oxygen at 50–60°C ignites and burns very brilliantly. During distillation of phosphorus(III) oxide, air must be rigorously excluded to avoid the possibility of

explosion. The later reference states that the carefully purified oxide does not ignite in oxygen, and that the earlier observations were on material containing traces of white phosphorus.

Phosphorus pentachloride

See Phosphorus pentachloride: Phosphorus(III) oxide

Sulfur

1. Thorpe, T. E. *et al.*, *J. Chem. Soc.*, 1891, **59**, 1019
2. Pernert, J. C. *et al.*, *Chem. Eng. News*, 1949, **27**, 2143

Interaction of a mixture of sulfur and the oxide under inert atmosphere above 160°C to form tetraphosphorus hexaoxide tetrasulfide is violent and dangerous on scales of working other than small [1]. A safer procedure involving distillation of phosphorus(V) oxide and phosphorus(V) sulfide is described [2].

Sulfuric acid

Mellor, 1940, Vol. 8, 898

Addition of sulfuric acid to the oxide causes violent oxidation, and ignition if more than 1–2 g is used.

Water


Mellor, 1940, Vol. 8, 897

Reaction with cold water is slow, but with hot water, violent, the evolved phosphine igniting. With more than 2 g of oxide, violent explosions occur.

See *other* NON-METAL OXIDES, REDUCANTS

4868. Tetraphosphorus hexaoxide–bis(borane)

[14940-94-4]



Water



Mellor, 1971, Vol. 8, Suppl. 3, 382

The compound ignites in contact with a little water.

See *related* BORANES, NON-METAL OXIDES

4869. Tetraphosphorus hexaoxide tetrasulfide

[15780-31-1]




Preparative hazard

See Tetraphosphorus hexaoxide: Sulfur

See *related* NON-METAL OXIDES, NON-METAL SULFIDES

4870. Disulfur heptaoxide

[12065-85-9]

O₇S₂Meyer, F. *et al.*, *Ber.*, 1922, **55**, 2923

The crystalline material ('peroxydisulfuric anhydride', of unknown structure) soon explodes on exposure to moist air.

See other ACID ANHYDRIDES, NON-METAL OXIDES**4871. Triuranium octaoxide**

[1344-59-8]

O₈U₃


Barium oxide

Mellor, 1942, Vol. 12, 49

Interaction below 300°C is vigorous and very exothermic.

See other METAL OXIDES**4872. Tetraphosphorus decaoxide (Phosphorus(V) oxide)**

[16752-60-6]

O₁₀P₄

(MCA SD-28, 1948); HCS 1980, 747

Barium sulfide

See Barium sulfide: Phosphorus(V) oxide

Formic acid

MRH 0.29/99+

See Formic acid: Phosphorus pentaoxide

Hydrogen fluoride

Gore, G., *J. Chem Soc.*, 1869, **22**, 368

Interaction is vigorous below 20°C.

Inorganic bases

1. Mellor, 1940, Vol. 8, 945

2. Van Wazer, 1958, Vol. 1, 279

Dry mixtures with calcium or sodium oxides do not react in the cold, but interact violently if warmed or moistened, evolving phosphorus(V) oxide vapour [1]. A

1882

mixture of the oxide and sodium carbonate may be initiated by strong local heating, when the whole mass will suddenly become hot [2].

Iodides

Pascal, 1960, Vol. 16.1, 538
Interaction is violent

Metals

MRH Aluminium 4.89/39, magnesium 5.48/45

Mellor, 1940, Vol. 8, 945
Interaction with warm sodium or potassium is incandescent and explosive with heated calcium.

Methyl hydroperoxide

See Methyl hydroperoxide: Alone, etc.

Other reactants

Yoshida, 1980, 119
MRH values calculated for 5 combinations with other materials are given.

Oxidants

See Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Metals, etc.
Hydrogen peroxide: Phosphorus(V) oxide
Oxygen difluoride: Phosphorus(V) oxide
Perchloric acid: Dehydrating agents

3-Propynol

See 3-Propynol: Phosphorus pentaoxide

Water

1. Mellor, 1940, Vol. 8, 944
2. *MCA SD-28*, 1948
Phosphorus(V) oxide is a very powerful desiccant and its interaction with liquid water is very energetic and highly exothermic [1]. The increase in temperature may be enough to ignite combustible materials if present and in contact [2].
See other ACID ANHYDRIDES, NON-METAL OXIDES

4873. Osmium

[7440-04-2]

Os

Os

Chlorine trifluoride

See Chlorine trifluoride: Metals

Fluorine

See Fluorine: Metals

Phosphorus

See Phosphorus: Metals

See other METALS

†4874. Phosphorus
[7723-14-0]

P



(MCA SD-16, 1947); FPA H19, 1973; HCS 1980, 748 (red), 750 (white); RSC Lab. Hazard Data Sheet No. 79, 1989 (white P)

1. Albright and Wilson, (Manufacturer's safety sheet)
2. Anon., *Chem. Trade J.*, 1936, **98**, 522
3. Anon., *CISHC Chem. Safety Summ.*, 1974–1975, **45–46**, 3
4. Anon., *Jahresber.*, 1979, 73
5. Anon., *Fire Prev.*, 1987, (197), 42–43
6. Anon., *Fire Prevention*, 1992, (249), 45

White phosphorus has an autoignition temperature only slightly above ambient, dispersed it will soon heat itself to that by the slow oxidation responsible for its glow. Red is not spontaneously combustible, however if it does catch fire white will be produced, so that the fire, once extinguished, may spontaneously re-ignite. Both can produce phosphine, among other products, by slow reaction with water. Sealed containers of damp phosphorus (white is often stored under water) may pressurise with highly toxic, pyrophoric, gas mixtures [1].

Red phosphorus powder being charged into a reaction vessel exploded when the lid of the tin fell in and was struck by the agitator [2]. A further case of ignition of red phosphorus by impact from its metal container is reported [3]. Impact of a rubber hammer on a nitrogen-purged stirred metal feed container of red phosphorus powder led to ignition and a jet of flame [4]. A rail tanker containing 44 kl of molten white phosphorus at about 50°C was damaged in a derailment, and the contents on exposure to ambient air immediately ignited, and burned for 5 days before being brought under control [5]. A very similar rail tankwagon fire occurred a few years later [6].

See Oxygen, below

See other FRICTIONAL INITIATION INCIDENTS

Alkalies

Mellor, 1940, Vol. 8, 802

Contact of phosphorus with boiling caustic alkalies or with hot calcium hydroxide evolves phosphine, which usually ignites in air.

Chlorosulfuric acid

Heumann, K. *et al.*, *Ber.*, 1882, **15**, 417

MRH 2.51/75

1884

P

White phosphorus begins to reduce the acid at 25–30°C and the vigorous reaction accelerates to explosion. Red phosphorus is similar at a higher initial temperature.

Cyanogen iodide

Mellor, 1940, Vol. 8, 791

Molten phosphorus reacts incandescently.

Halogen azides

See HALOGEN AZIDES: metals, or phosphorus

Halogen oxides

MRH values show % of oxidant

See Chlorine dioxide: Non-metals

MRH 7.87/73

Dichlorine oxide: Oxidisable materials

MRH 0.92/92

Oxygen difluoride: Non-metals

Trioxxygen difluoride: Various materials

Halogens, or Interhalogens

MRH Chlorine 1.92/85

1. Mellor, 1940, Vol. 8, 785; 1956, Vol. 2, Suppl. 1, 379

2. Christomanos, A. C., *Z. Anorg. Chem.*, 1904, **41**, 279

3. Kuhn, R. *et al.*, *Helv. Chim. Acta*, 1928, **11**, 107

4. Newkome, G. R. *et al.*, *J. Chem. Soc., Chem. Comm.*, 1975, 885–886

Both yellow and red phosphorus ignite on contact with fluorine and chlorine; red ignites in liquid bromine or in a heptane solution of chlorine at 0°C. Yellow phosphorus explodes in liquid bromine or chlorine, and ignites in contact with bromine vapour or solid iodine [1]. Interaction of bromine and white phosphorus in carbon disulfide gives a slimy by-product which explodes violently on heating [2]. Interaction of phosphorus and iodine in carbon disulfide is rather rapid [3]. A less hazardous preparation of diphosphorus tetraiodide from phosphorus trichloride and potassium iodide in ether is recommended [4].

See Bromine pentafluoride: Acids, etc.

Bromine trifluoride: Halogens, etc.

Chlorine trifluoride: Metals, etc.

Iodine pentafluoride: Metals, etc.

Iodine trichloride: Phosphorus

Hexalithium disilicide

Mellor, 1940, Vol. 6, 169

Interaction is incandescent.

Hydriodic acid

1. Blau, K., private comm., 1965

2. Agranat, I. *et al.*, *J. Chem. Soc., Perkin Trans. 1*, 1974, 1159

3. Villain, F. J. *et al.*, *J. Med. Chem.*, 1964, **7**, 457, footnote 9

During removal of free iodine from hydriodic acid by distillation from red phosphorus, phosphine was produced. When air was admixed by changing the receiver flask, an explosion occurred. Omission of distillation by boiling the reactants in inert atmosphere, and separating the phosphorus by filtration through a sintered

glass funnel containing solid carbon dioxide gave a colourless product [1]. Potential hazards incidental to the use of red phosphorus and hydriodic acid as a reducing agent for organic carbonyl compounds have been noted. Accumulation in a reflux condenser of phosphonium iodide which will react violently with water [2], and 2 unexplained explosions [3], possibly owing to the same cause, have been reported. *See* Phosphonium iodide

Hydrogen peroxide

MRH 7.78/73

See Hydrogen peroxide: Phosphorus

Magnesium perchlorate

1965 Summary of Serious Accidents, Washington, USAEC, 1966

Phosphorus and magnesium perchlorate exploded violently while being mixed. It seems likely that this was initiated by interaction of phosphorus and traces of perchloric acid to form magnesium phosphate. The acid may have been present as impurity in the magnesium perchlorate, or might have been formed from traces of phosphoric acid in the phosphorus. Once formed, the perchloric acid would be rendered anhydrous by the powerful dehydrating action of magnesium perchlorate.

Metal acetylides

Mellor, 1946, Vol. 5, 848–849

Monorubium acetylide and monocaesium acetylide incandesce with warm phosphorus. Lithium acetylide and sodium acetylide burn vigorously in phosphorus vapour, and the potassium, rubidium and caesium analogues should react with increasing violence.

Metal halogenates

MRH values below references

Mellor, 1941, Vol. 2, 310; 1940, Vol. 8, 785–786

MRH Barium chlorate 5.06/83, calcium chlorate 5.61/77, potassium chlorate 6.07/76, sodium bromate 4.98/80, sodium chlorate 7.32/75, zinc chlorate 6.11/76

Dry finely divided mixtures of red (or white) phosphorus with chlorates, bromates or iodates of barium, calcium, magnesium, potassium, sodium or zinc will readily explode on initiation by friction, impact or heat. Fires have been caused by accidental contact in the pocket between the red phosphorus in the friction strip on safety-match boxes and potassium chlorate tablets. Addition of a little water to a mixture of white or red phosphorus and potassium iodate causes a violent or explosive reaction. Addition of a little of a solution of phosphorus in carbon disulfide to potassium chlorate causes an explosion when the solvent evaporates. The extreme danger of mixtures of red phosphorus (or sulfur) with chlorates was recognised in the UK some 50 years ago when unlicensed preparation of such mixtures was prohibited by Orders in Council.

Metal halides

Mellor, 1939, Vol. 9, 467; 1943, Vol. 11, 234

Phosphorus ignites in contact with antimony pentachloride and explodes with chromyl chloride in presence of moisture at ambient temperature.

P

Metal oxides or peroxides MRH values below references
Mellor, 1941, Vol. 2, 490; Vol. 7, 690; 1940, Vol. 8, 792; 1943, Vol. 11, 234
MRH Chromium trioxide 3.68/84, lead peroxide 1.30/90, potassium peroxide
5.90/81, sodium peroxide 2.30/86

Red phosphorus reacts vigorously on heating with copper oxide or manganese dioxide; and on grinding with lead oxide, mercury oxide or silver oxide, ignition may occur. Red phosphorus ignites in contact with lead peroxide, potassium peroxide or sodium peroxide, while white phosphorus explodes, and also in contact with molten chromium trioxide at 200°C.

Metals

1. Mellor, 1941, Vol. 7, 115; 1940, Vol. 8, 842, 847, 853; 1971, Vol. 8, Suppl. 3, 228
 2. Van Wazer, 1958, Vol. 1, 159
 3. Mellor, 1942, Vol. 15, 696; 1937, Vol. 16, 160
 4. Iseler, G. W. *et al.*, *Int. Conf. Indium Phosphide Relat. Mater.*, 1992, 266
- Reaction of beryllium, copper, manganese, thorium or zirconium is incandescent when heated with phosphorus [1] and that of cerium, lanthanum, neodymium and praseodymium is violent above 400°C [2]. Osmium incandesces in phosphorus vapour, and platinum burns vividly below red-heat [3]. Red phosphorus shows very variable vapour pressure between batches (not surprising, it is an indeterminate material). This leads to explosions when preparing indium phosphide by reactions involving fusion with phosphorus in a sealed tube [4].
See Aluminium: Non-metals

Metal sulfates

Berger, E., *Compt. rend.*, 1920, **170**, 1492
Excess red phosphorus will burn admixed with barium or calcium sulfates if primed at a high temperature with potassium nitrate–calcium silicide mixture.
See Potassium nitrate: Calcium silicide

Nitrates MRH Ammonium nitrate 4.60/86, potassium nitrate 3.14/73,
silver nitrate 3.89/82, sodium nitrate 4.43/73

1. Mellor, 1941, Vol. 3, 470; 1940, Vol. 4, 987; Vol. 8, 788
 2. Anon., *Jahresber.*, 1975, 83–84
- Yellow phosphorus ignites in molten ammonium nitrate, and mixtures of phosphorus with ammonium nitrate, mercury(I) nitrate or silver nitrate explode on impact. Red phosphorus is oxidised vigorously when heated with potassium nitrate [1]. During development of new refining agents for aluminium manufacture, a mixture containing red phosphorus (16%) and sodium nitrate (35%) was being pressed into 400 g tablets. When the die pressure was increased to 70 bar, a violent explosion occurred [2].

Nitric acid MRH 7.66/67
See Nitric acid: Non-metals

Nitrogen halides

See Nitrogen tribromide hexaammoniate: Initiators
Nitrogen trichloride: Alone, or Non-metals MRH 3.35/87

Nitrosyl fluoride

See Nitrosyl fluoride: Metals, etc.

Nitryl fluoride

See Nitryl fluoride: Non-metals

Non-metal halides

1. Pascal, 1960, Vol. 13.2, 1162

2. Mellor, 1946, Vol. 5, 136; 1940, Vol. 8, 787; 1947, Vol. 10, 906

Either the white or red form incandescences with boron triiodide [1]. Red phosphorus incandescences in seleninyl chloride while white explodes. Red phosphorus reacts vigorously on warming with sulfuryl chloride or disulfuryl chloride, and violently with disulfur dibromide [2].

Non-metal oxides

MRH Dinitrogen tetraoxide 9.12/65

Mellor, 1940, Vol. 8, 786–787

Warm or molten white phosphorus burns vigorously in nitrogen oxide, dinitrogen tetraoxide or dinitrogen pentaoxide. White phosphorus ignites after some delay in contact with the vapour of sulfur trioxide, but immediately in contact with the liquid if a large portion is used.

Other reactants

Yoshida, 1980, 73–75

MRH values calculated for 36 combinations, largely with oxidants, are given.

Oxidants

Inoue, Y. *et al.*, *Chem. Abs.*, 1988, **108**, 134435

Impact sensitivities of mixtures of red phosphorus with various oxidants were determined in a direct drop-ball method, which indicated higher sensitivities than those determined with an indirect striker mechanism. Mixtures with silver chlorate were most sensitive, those with bromates, chlorates and chlorites were extremely sensitive, and mixtures with sodium peroxide and potassium superoxide were more sensitive than those with barium, calcium, magnesium, strontium or zinc peroxides. Mixtures with perchlorates or iodates had sensitivities comparable to those of unmixed explosives, such as lead azide, 3,5-dinitrobenzenediazonium-2-oxide etc.

Oxygen

1. Mellor, 1940, Vol. 8, 771

2. Van Wazer, 1958, 1, 97

3. Mellor, 1971, Vol. 8, Suppl. 3, 237–245

The reactivity of phosphorus with air or oxygen depends on the allotrope of phosphorus involved and the conditions of contact, white (yellow) phosphorus being by far the more reactive. White phosphorus readily ignites in air if warmed, finely divided (e.g. from an evaporating solution) or under conditions where the slow oxidative exotherm cannot be dissipated. Contact with finely divided charcoal or lampblack promotes ignition, probably by adsorbed oxygen [1]. Contact with amalgamated aluminium also promotes ignition [2]. Oxidation of phosphorus under both explosive and non-explosive conditions has been reviewed in detail [3].

Peroxyformic acid MRH 6.65/72
See Peroxyformic acid: Non-metals

Potassium nitride
Mellor, 1940, Vol. 8, 99
Potassium and other alkali-metal nitrides react on heating with phosphorus to give a highly flammable mixture which evolves ammonia and phosphine with water.

Potassium permanganate MRH 2.34/84
Mellor, 1942, Vol. 12, 322
Grinding mixtures of phosphorus and potassium permanganate causes explosion, more violent if also heated.

Selenium
See Selenium: Phosphorus

Sodium chlorite MRH 6.23/79
See Sodium chlorite: Phosphorus

Sulfur
See Sulfur: Non-metals

Sulfuric acid MRH 2.64/72
See Sulfuric acid: Phosphorus
See other NON-METALS, PYROPHORIC MATERIALS

4875. Phosphorus(V) sulfide **P₂S₅**
[1314-80-3]



See Tetraphosphorus decasulfide

4876. Zinc phosphide **P₂Zn₃**
[1314-85-7]




Perchloric acid
See Perchloric acid: Zinc phosphide

Water
Fehse, W., *Feuerschutz*, 1938, **18**, 17
Contact of one drop of water with a zinc phosphide rodenticide preparation caused ignition.
See other METAL NON-METALLIDES

4877. Tetraphosphorus trisulfide
[1314-85-8]

P₄S₃



Gibson, 1969, 118–119
It ignites in air above 100°C.

Other reactants


Yoshida, 1980, 131

MRH values calculated for 16 combinations, largely with oxidants, are given.

See other NON-METAL SULFIDES

4878. Tetraphosphorus decasulfide (Phosphorus(V) sulfide)
[15857-57-5]

P₄S₁₀



1. (MCA SD-71, 1958)
2. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74
3. Albright and Wilson, (Manufacturer's safety sheet)

The general reactivity of the sulfide depends markedly on the physical form, and differences of a factor of 10 may be involved. It is ignitable by friction, sparks or flames, and ignites in dry air if heated close to the m.p., 275–280°C. The dust (200 mesh) forms explosive mixtures in air above a concentration of 0.5% w/v [1], and maximum explosion pressures of 4.35 bar, with maximum rate of rise exceeding 680 bar/s have been determined [2]. The dust can acquire sufficient static electricity from movement for ignition to occur [3].

Alcohols

1. MCA Case History No. 227
2. Ashford, J. S., private comm., 1964

A mixture of the sulfide, ethylene glycol and hexane in a mantle-heated flask spontaneously overheated and exploded at an internal temperature of around 180°C. It had been intended to maintain the reaction temperature at 60°C, but since alcoholysis of the sulfide is exothermic, presence of the heating mantle prevented the dissipation of heat, and the reaction accelerated continuously until explosive decomposition occurred [1]. An incident in similar circumstances involving interaction of the sulfide with 4-methyl-2-pentanol also led to violent eruption of the flask contents.

Diethyl maleate, Methanol

Nolan, 1983, Case history 152

In manufacture of Malathion, the phosphorus sulfide dissolved in toluene is treated first with methanol, then diethyl maleate. Use of insufficient toluene to dissolve the sulfide led to a runaway exothermic reaction and explosion.

See Alcohols, above

Water

Haz. Chem. Data, 1975, 239

It heats and may ignite in contact with limited amounts of water, hydrogen sulfide also being evolved.

See other NON-METAL SULFIDES

4879. Tetraphosphorus decasulfide . pyridine complex
[71257-31-3]

P₄S₁₀·4C₅H₅N

P₄S₁₀·4Py


Riedel-de Haën catalogue, 1986

Although convenient to handle, the complex is classed as highly flammable.

See related NON-METAL SULFIDES

4880. Tetraphosphorus triselenide
[1314-86-9]

P₄Se₃



1. Mellor, 1947, Vol. 10, 790

2. Stoppioni, P. *et al.*, *Gazz. Chim. Ital.*, 1988, **118**, 581–582

It ignites when heated in air [1]. A new and safer synthesis from red phosphorus and grey selenium has been described [2].

See related NON-METAL SULFIDES

4881. Lead pentaphosphide
[]

P₅Pb

PbP₅

Bailar, 1973, Vol. 2, 117

It ignites in air

See other METAL NON-METALLIDES, PYROPHORIC MATERIALS

1891

4882. Lead

[7439-92-1]

Pb

Pb

HCS 1980, 534 (powder)

Mézáros, L., *Tetrahedron Lett.*, 1967, 4951

The finely divided lead produced by reduction of the oxide with furfural vapour at 290°C is pyrophoric and chemically reactive.

Hydrogen peroxide, Trioxane

See Hydrogen peroxide: Lead, Trioxane

Nitric acid, Rubber

See Nitric acid: Lead-containing rubber

Oxidants

See Ammonium nitrate: Metals

Chlorine trifluoride: Metals

Hydrogen peroxide: Metals

Sodium acetylide

See Sodium acetylide: Metals

See other METALS, PYROPHORIC METALS

4883. Lead–tin alloys

[12333-97-0] (1:1 alloy)

Pb–Sn

Pb–Sn

Palm oil, Steel sheets

Nedelcu, C. *et al.*, Rom. Pat. 68 602, 1979

The explosion hazards associated with use of palm oil when coating steel sheets with molten alloys (8–12% tin) is eliminated by use of molten sodium acetate in place of the oil.

See other ALLOYS

4884. Lead–zirconium alloys

[12412-94-1] (3:5 alloy)

Pb–Zr

Pb–Zr

Alexander, P. P., US Pat. 2 611 316. 1952

Alloys containing 10–70% of zirconium will pulverise and ignite on impact.

See other ALLOYS, PYROPHORIC ALLOYS

4885. Palladium
[7440-05-3]

Pd

Pd

Finely divided palladium (palladium-black) used as a hydrogenation catalyst is usually pyrophoric and needs appropriate handling precautions.

Aluminium

See Aluminium: Catalytic metals

Arsenic

See Arsenic: Metals

Carbon

1. Mazingo, R., *Org. Synth.*, 1955, Coll. Vol. 3, 687
2. Rachlin, A. I. *et al.*, *Org. Synth.*, 1971, **51**, 687
3. Fraser, R., private comm., 1973

Palladium-on-carbon catalysts prepared by formaldehyde reduction are less pyrophoric than those reduced with hydrogen [1]. Such catalysts become extremely pyrophoric on thorough vacuum drying [2]. Those prepared on high surface-area supports (up to 2000 m²/g) are highly active and readily will cause catalytic ignition of hydrogen-air or solvent-air mixtures. Methanol is noted for easy ignition because of its high volatility [3].

Formic acid

See Formic acid: Palladium-carbon catalyst

Hydrogen, Hydrogen peroxide

See Hydrogen peroxide: Hydrogen, Palladium

Hydrogen, Poly(tetrafluoroethylene)

See Poly(tetrafluoroethylene): Metal hydrides

Ozonides

See OZONIDES (reference 4)

Sodium tetrahydroborate

Augustine, 1968, 75

In preparation for the reduction of a nitro compound, the tetrahydroborate solution is added to an aqueous suspension of palladium-on-charcoal catalyst. The reversed addition of dry catalyst to the tetrahydroborate solution may cause ignition of liberated hydrogen.

Sulfur

See Sulfur: Metals

See other HYDROGENATION CATALYSTS, METALS, PYROPHORIC METALS

4886. Praseodymium
[7440-10-0]

Pr

Pr

Bromine

See Bromine: Metals,

See also LANTHANIDE METALS See other METALS

4887. Platinum
[7440-06-4]

Pt

Pt

1. van Campen, M. G., *Chem. Eng. News*, 1954, **32**, 4698

2. Tilford, C. H., private comm., 1965

Finely divided (catalytic) forms of platinum are hazardous to handle if allowed to dry. Used Adams' hydrogenation catalyst exploded while being sieved in air. It had been well washed, finally with water, and then air dried [1]. Manipulation of used catalyst on filter paper caused a violent explosion [2]. The explosive catalysed oxidation of adsorbed hydrogen is suspected here.

See Zinc: Catalytic metals

Aluminium

See Aluminium: Catalytic metals

Acetone, Nitrosyl chloride

See Nitrosyl chloride: Acetone, etc.

Arsenic

Wöhler, L., *Z. Anorg. Chem.*, 1930, **186**, 324

During interaction to give platinum diarsenide in a sealed tube at 270°C, the highly exothermic reaction may burst the tube.

Carbon, Methanol

Unpublished information, 1982

Ignition occurred when methanol was charged into an unpurged autoclave containing platinum-on-carbon catalyst.

Dioxygen difluoride

See Dioxygen difluoride: Various materials

Ethanol

Elkins, J. S., private comm., 1968

Addition of platinum-black catalyst to ethanol caused ignition. Pre-reduction with hydrogen and/or nitrogen purging of air prevented this.

Hydrazine

See Hydrazine: Metal catalysts

Hydrogen, Air

Coleman, J. W., private comm., 1965

A platinised alumina catalyst had been treated with flowing hydrogen at ambient temperature. Purging with air caused an explosion. Inert gas purging is essential after treating catalysts or adsorbent with hydrogen before admitting air.

Hydrogen peroxide

See Hydrogen peroxide: Metals

Lithium

Nash, C. P., *Chem. Eng. News*, 1961, **39**(5), 42

Interaction to form an intermetallic compound sets in violently at $540^{\circ}\text{C}\pm 20^{\circ}\text{C}$.

Methyl hydroperoxide

See Methyl hydroperoxide: Alone, etc.

Ozonides

See OZONIDES (reference 4)

Peroxomonosulfuric acid

See Peroxomonosulfuric acid: Catalysts

Phosphorus

See Phosphorus: Metals

Selenium, or Tellurium

Mellor, 1937, Vol. 16, 158

Finely divided or spongy platinum reacts incandescently when heated with selenium or tellurium.

Vanadium dichloride, Water

See Vanadium dichloride: Platinum, Water

See other HYDROGENATION CATALYSTS, METALS, PYROPHORIC METALS

4888. Plutonium

[7440-07-5]

Pu

Pu

1. Bailar, 1973, Vol. 5, 44–45

2. Williamson, G. K., *Chem. & Ind.*, 1960, 1384

The reactive metal is pyrophoric in air at elevated temperatures; lathe turnings have ignited at 265°C and 140-mesh powder at 135°C [1]. This most extremely toxic metal may be worked under nitrogen, argon, or helium (but not carbon dioxide) at temperatures above 600°C [2].

Carbon tetrachloride

Serious Accid. Series, No. 246, Washington, USAEC, 1965

While draining after degreasing in the solvent, plutonium chips ignited, and when accidentally dropped into the solvent, caused an explosion.

See other METAL–HALOCARBON INCIDENTS

Water

MCA Case History No. 1212

Plutonium components, normally kept under argon, were accidentally exposed to air and moisture, probably forming plutonium hydride and hydrated oxides. When the plastics containing bag was disturbed, ignition occurred, causing widespread radiation contamination.

See Plutonium(III) hydride

See other METALS, PYROPHORIC METALS

4889. Rubidium

[7440-17-7]

Rb

Rb

Mellor, 1941, Vol. 2, 468; 1963, Vol. 2, Suppl. 2.2, 2172

Rubidium is a typical but very reactive member of the series of alkali metals. It is appreciably more reactive than potassium, but less so than caesium, and so would be expected to react more violently with those materials that are hazardous with potassium or sodium. Rubidium ignites on exposure to air or dry oxygen, largely forming the oxide.

Halogens

Mellor, 1941, Vol. 2, 13; 1963, Vol. 2, Suppl. 2.2, 2174

Ignition occurs in fluorine, chlorine and bromine or iodine vapours. Contact with liquid bromine would be expected to cause a violent explosion.

Mercury

Mellor, 1941, Vol. 2, 469

Interaction is exothermic and may be violent.

Non-metals

Mellor, 1963, Vol. 2, Suppl. 2.2, 2174–2175

The molten metal ignites in sulfur vapour, and reacts with various forms of carbon exothermically, one product, the interstitial rubidium octacarbide, being pyrophoric.

Vanadium trichloride oxide

Mellor, 1963, Vol. 2, Suppl. 2.2, 2176

Interaction is violent at 60°C.

Water

1. Mellor, 1941, Vol. 2, 469

2. Markowitz, M. M., *J. Chem. Educ.*, 1963, **40**, 633–636

Contact with cold water is exothermic enough to ignite the hydrogen evolved [1]. Reactivity of rubidium and other alkali metals with water has been discussed in detail [2].

See ALKALI METALS *See other* METALS, PYROPHORIC METALS

4890. Rhenium

[7440-15-5]

Re

Re

Oxygen (Gas)

See Oxygen (Gas): Rhenium

See other METALS

4891. Rhenium(VII) sulfide

[12038-67-4]

Re₂S₇

Re₂S₇

Sidgwick, 1950, 1299

It is readily oxidised in air, sometimes igniting.

See other METAL SULFIDES

4892. Rhodium

[7440-16-6]

Rh

Rh

Sidgwick, 1950, 1513

Metallic rhodium prepared by heating its compounds in hydrogen must be allowed to cool in an inert atmosphere to prevent catalytic ignition of the sorbed hydrogen on exposure to air.

See Zinc: Catalytic metals

Interhalogens

See Bromine pentafluoride: Acids, etc.

Chlorine trifluoride: Metals

See other HYDROGENATION CATALYSTS, METALS, PYROPHORIC METALS

4893. Radon

[10043-92-2]

Rn

Rn

Water

John, G., *Health Phys.*, 1985, **49**, 977–979

A 22 year old glass ampoule containing an aqueous solution of radon was under considerable pressure when opened and ejected the top vigorously. This was attributed to formation of hydrogen by radiolysis of the solvent water.

See Americium trichloride
See other IRRADIATION DECOMPOSITION INCIDENTS

4894. Ruthenium
[7440-18-8]

Ru

Ru

1. Mellor, 1942, Vol. 15, 502
2. Ullmann, 1992, Vol. A21, 125/6

Dissolution of a zinc–ruthenium alloy in hydrochloric acid leaves an explosive residue of finely divided ruthenium [1]. More probably this is the hydride, which may decompose on slight stimulus, the evolved hydrogen probably igniting because of the catalytic activity of the metal. Ruthenium prepared from its compounds by borohydride reduction is especially dangerous in this respect [2].

Aqua regia, Potassium chlorate
See Potassium chlorate: Aqua regia, etc.
See other HYDROGENATION CATALYSTS, METALS

4895. Ruthenium(IV) sulfide
[12166-20-0]

RuS₂

RuS₂

Air, or Carbon dioxide
Mellor, 1942, Vol. 15, 541
The sulfide decomposes explosively if heated in air or air–nitrogen mixtures, and incandescens, then explodes if heated under carbon dioxide.
See other METAL SULFIDES

4896. Ruthenium salts
[22541-59-9] (2+)
[22541-88-4] (3+)


RuZ₂
RuZ₃

RuZ₂, etc.

Sodium tetrahydroborate
See Sodium tetrahydroborate: Ruthenium salts

†**4897. Sulfur**
[7704-34-9]

S



(MCA SD-74, 1959); NSC 612, 1979 (solid); NSC 592, 1978 (liquid); FPA H14, 1973; HCS 1980, 871 (liquid), 872 (solid)
 NFPA 655, Quincy (Ma), Natl. Fire Prot. Assoc., 1988

1. MCA SD-74, 1959
2. Anon., *Loss Prev. Bull.*, 1987, (078), 23–25
3. Ayers, P. G., *Loss. Prev. Bull.*, 1988, (094), 34
4. Poole, G. A., *Loss. Prev. Bull.*, 1988, (084), 35

The new US National Fire Code covers the precautions necessary to prevent fires or explosions during crushing of the solid, or during handling of the molten element.

It may become ignited through frictional heat or incendive sparks, particularly when suspended as dust in air, for which the approximate explosive limits are 35–1400 g/m³. Handling precautions for the solid and molten element are detailed. In the hazardous reactions listed below, the dual capacity of sulfur to oxidise or be oxidised is apparent [1]. Molten sulfur at '150–200°C' was being transported in a truck with two tank trailers each holding 12 t (57% of capacity). A traffic accident on a bridge caused both tanks to vault over the central barrier (thought to be caused by surging of the tank contents), and the impact caused tank rupture and ignition of the spilled hot sulfur. The fire led to dense clouds of sulfur dioxide and smoke which greatly hampered emergency services. The flash point of pure sulfur is 188°C, but impure sulfur may ignite at 168°C [2]. Later correspondents corrected the temperature limits for transportation of molten sulfur to 130–140° or 132–149°C [3,4]. Several values have been quoted in the literature for both flash point and autoignition temperature of molten sulfur, e.g. fl.p. 168, 188; a.i.t. 190, 232°C.

See entry HIGH SURFACE-AREA SOLIDS

Aluminium, Copper

See Aluminium: Copper, Sulfur

Aluminium, Starch

Biasutti, 1981, 138

Ignition of a dry mixture of the 3 powdered materials being mixed in a screw extruder led to a minor explosion, then a major dust explosion.

Calcium, Vanadium pentoxide

See Vanadium(V) oxide: Calcium, etc.

Diethyl ether

Taylor, H. F., *J. R. Inst. Chem.*, 1955, **79**, 43

Evaporation of an ethereal extract of sulfur caused an explosion of great violence. Experiment showed that evaporation of wet, peroxidised ether gave a mildly explosive residue, which became violently explosive on addition of sulfur.

Fibreglass, Iron

Anon., *Info. Circ. No. 8272(10)*, Washington, US Bur. Mines, 1964

Rubidium: Non-metals
Sodium: Non-metals
See Selenium: Metals

Non-metals

1. Mellor, 1946, Vol. 5, 15
2. von Schwartz, 1918, 328
3. Mellor, 1940, Vol. 8, 786
4. Phillips, R., *Org. Synth.*, 1943, Coll. Vol. 2, 579

Boron reacts with sulfur at 600°C becoming incandescent [1]. Mixtures of sulfur with lamp black or freshly calcined charcoal ignite spontaneously, probably owing to adsorbed oxygen on the catalytic surface [2]. Mixtures of yellow phosphorus and sulfur ignite and/or explode on heating [3]. Ignition of an intimate mixture of red phosphorus and sulfur causes a violent exothermic reaction [4].

Other reactants

Yoshida, 1980, 37–38

MRH values calculated for 32 combinations, mainly with oxidants, are given.

Oxidants

MRH values show % of oxidant

1. Ivanov, V. G. *et al.*, *Chem. Abs.*, 1980, **93**, 116708
2. Ivanov, V. G. *et al.*, *Chem. Abs.*, 1981, **94**, 86592

The self-ignition of sulfur with potassium chlorate or iodine(V) oxide at 145–160°C, and with potassium perchlorate at 385°C was studied using DTA [1], and combustion characteristics of the mixtures were determined [2].

See Halogen oxides, Interhalogens, Metal oxides, all above

Ammonium nitrate: Sulfur	MRH 3.01/88
Barium bromate: Sulfur	MRH 2.26/88
Chromyl chloride: Non-metals	
Fluorine: Non-metals	MRH 8.28/78
Heptasilver nitrate octaoxide: Alone, or Sulfides, etc.	
Lead(II) chlorite: Non-metals	
Lead chromate: Sulfur	
Phosphorus(III) oxide	
Potassium bromate: Non-metals	MRH 2.55/86
Potassium chlorate: Charcoal, etc., or: Non-metals	MRH 3.22/82
Potassium chlorite: Sulfur	
Potassium perchlorate: Sulfur	MRH 3.39/79
Potassium permanganate: Non-metals	MRH 1.63/90
Silver bromate: Sulfur compounds	MRH 2.00/88
Silver chlorite: Hydrochloric acid, etc.	
Silver nitrate: Non-metals	MRH 1.67/84
Sodium chlorate: Ammonium salts, etc.	MRH 4.27/80
Tetraphosphorus hexaoxide: Sulfur	

Potassium nitride

Mellor, 1940, Vol. 8, 99

S

Potassium nitride and other alkali metal nitrides react with sulfur to form a highly flammable mixture, which evolves ammonia and hydrogen sulfide in contact with water.

See Caesium nitride

Sodium hydride

Mellor, 1941, Vol. 2, 483

Interaction with sulfur vapour is vigorous.

Static discharges

Enstad, G., *A Reconsideration of the Concept of Minimum Ignition Energy*, paper at Euro. Fed. Chem. Engrs. Wkg. Party meeting, 16th March, 1975.

Sulfur dust has a very low minimum ignition energy (3 mJ), so particular care is necessary in formulating protective measures.

Tetraphenyllead

See Tetraphenyllead: Sulfur

See other NON-METALS, OXIDANTS

4898. Silicon monosulfide

[12504-41-5]

SiS

SSi

Water

Bailar, 1973, Vol. 1, 1355

It reacts rapidly with water and may ignite.

See other NON-METAL SULFIDES

4899. Samarium sulfide

[29678-92-0]

SmS

SSm

Gadolinium sulfide

Jayaraman, A. *et al.*, *Phys. Rev. Lett.*, 1973, **31**, 700

Cooling of samples of crystalline samarium sulfide containing 22 atom% of gadolinium in the lattice to below -153°C causes a near-explosive rearrangement to a powder form. This was attributed to changes in the lattice constants corresponding to a volume increase of 7.5%.

See other METAL SULFIDES

4900. Tin(II) sulfide

[1314-95-0]

SnS

SSn

See entry REFRACTORY POWDERS

1903

Oxidants

See Chloric acid: Metal sulfides

Dichlorine oxide: Oxidisable materials

See also METAL HALOGENATES: metals, etc.

See other METAL SULFIDES

4901. Strontium sulfide

[1314-96-1]

SrS

SSr

Lead dioxide

See Lead(IV) oxide: Metal sulfides

See other METAL SULFIDES

4902. Tin(IV) sulfide

[1315-01-1]

SnS₂

S₂Sn

Oxidants

See Chloric acid: Metal sulfides

Dichlorine oxide: Oxidisable materials

See METAL HALOGENATES: metals, etc.

See other METAL SULFIDES

4903. Tantalum(IV) sulfide

[12143-72-5]

TaS₂

S₂Ta

Preparative hazard

See Sulfur: Metals (reference 9)

See other METAL SULFIDES

4904. Titanium(IV) sulfide

[12039-13-3]

TiS₂

S₂Ti

Potassium nitrate

See Potassium nitrate: Metal sulfides

See other METAL SULFIDES

4905. Uranium(IV) sulfide
[12039-14-4]

US₂

S₂U

Nitric acid

See Nitric acid: Uranium disulfide

See other METAL SULFIDES

4906. Antimony trisulfide
[1345-04-6]

Sb₂S₃

S₃Sb₂

Other reactants

Yoshida, 1980, 130

MRH values calculated for 11 combinations with various materials are given.

Oxidants

MRH values show % of oxidant.

See Chloric acid: Metal sulfides

Dichlorine oxide: Oxidisable materials

MRH 3.72/49

Fluorine: Sulfides

MRH 6.40/56

Heptasilver nitrate octaoxide: Alone, or Sulfides

Lead(II) chlorite: Antimony sulfide

Potassium chlorate: Metal sulfides

MRH 2.84/61

Potassium nitrate: Metal sulfides

MRH 2.30/63

Silver(I) oxide: Metal sulfides

Thallium(III) oxide: Sulfur, etc.

See other METAL SULFIDES

4907. Antimony
[7440-36-0]

Sb

Sb

NSC 408, 1978 (metal and compounds)

Krebs, H. *et al.*, *Z. Anorg. Chem.*, 1955, **282**, 177

Electrolysis of acidified, stirred antimony halide solutions at low current density produces explosive antimony which contains substantial amounts of halogen.

Alkali nitrates

Mellor, 1947, Vol. 9, 382

Mixtures detonate on heating, forming antimonates.

Aluminium

See Aluminium: Antimony, etc.

Disulfur dibromide

See Disulfur dibromide: Metals

Halogens MRH Chlorine 1.67/47, fluorine 5.10/32
Mellor, 1947, Vol. 9, 379
Antimony ignites in fluorine, chlorine and bromine. With iodine, ignition or explosion may occur if quantities are large enough.

Other reactants
Yoshida, 1980, 34
MRH values calculated for 9 combinations with oxidants are given.

Oxidants	MRH values show % of oxidant
<i>See</i> Halogens, above	
Ammonium nitrate: Metals	MRH 2.30/62
Bromine pentafluoride: Acids, etc.	MRH 2.76/46
Bromine trifluoride: Halogens, etc.	MRH 2.34/53
Chloric acid: Metals, etc.	
Chlorine trifluoride: Metals, etc.	MRH 3.47/36
Dichlorine oxide: Oxidisable materials	MRH 2.64/41
Iodine pentafluoride: Metals	
Iodine: Metals	
Nitrosyl fluoride: Metals	
Perchloric acid: Antimony(III) compounds	
Potassium dioxide: Metals	
Potassium permanganate: Antimony, etc.	MRH 0.25/86
Seleninyl chloride: Antimony	
Sodium nitrate: Antimony	
<i>See other</i> METALS	

4908. Selenium
[7782-49-2]

Se

Se

NSC 578, 1982 (element and compounds)

Hexalithium disilicide
See Hexalithium disilicide: Non-metals

Metal acetylides or carbides
Mellor, 1946, Vol. 5, 862, 886
Metal acetylides incandesce on heating in selenium vapour; barium acetylide at 150°C, calcium acetylide and strontium acetylide at 500°C; and thorium carbide at an unstated temperature.
See Rubidium acetylide: Non-metals

Metal amides
Pascal, 1960, Vol. 16.2, 1724
Interaction of selenium with alkali metal amides and alkaline earth metal amides gives explosive products.

Metal chlorates

Mellor, 1956, Vol. 2, Suppl. 1, 583

A slightly moist mixture of selenium with any chlorate (except of alkali metals) becomes incandescent.

See METAL HALOGENATES: non-metals

Metals

1. Mellor, 1942, Vol. 15, 151
2. Mellor, 1947, Vol. 10, 766–777
3. Mellor, 1942, Vol. 12, 31
4. Mellor, 1940, Vol. 4, 480
5. Mellor, 1942, Vol. 16, 158
6. Reisman, A. *et al.*, *J. Phys. Chem.*, 1963, **67**, 22
7. Bailar, 1973, Vol. 2, 79

Nickel and selenium interact with incandescence on gentle heating [1], as do also sodium and potassium, the latter mildly explosively [2]. Uranium [3] and zinc [4] also incandesce when their mixtures with selenium are heated, and platinum sponge incandesces vividly [5]. The particle size of cadmium and selenium must be below a critical size to prevent explosions during synthesis of cadmium selenide by heating the elements together. Similar considerations also apply to interaction of cadmium or zinc with sulfur, selenium or tellurium [6]. Interaction of powdered tin and selenium at 350°C is extremely exothermic [7].

Nitrogen trichloride

See Nitrogen trichloride: Initiators

Oxidants

See Metal chlorates, above; Oxygen, etc., below

- Barium peroxide: Selenium
- Bromine pentafluoride: Acids, etc.
- Chlorine trifluoride: Metals, etc.
- Chromium trioxide: Selenium
- Fluorine: Non-metals
- Potassium bromate: Non-metals
- Silver(I) oxide: Non-metals
- Sodium peroxide: Non-metals

Oxygen, Organic matter

1. Astin, S. *et al.*, *J. Chem. Soc.*, 1933, 391
2. Watkins, C. R. *et al.*, *Chem. Rev.*, 1945, **36**, 235

During attempted conversion of recovered selenium metal to the dioxide by heating in oxygen, a vigorous explosion occurred. This was attributed to selenium-catalysed oxidation of traces of organic impurities in the selenium. Oxidation of recovered selenium with nitric acid is rendered vigorous by presence of organic impurities, but is a safe procedure [1]. The later reference incorrectly attributed the explosions to the nitric acid procedure [2].

Phosphorus

Pascal, 1956, Vol. 10, 724

Interaction warm attains incandescence.

See related METALS, NON-METALS

4909. Silicon

[7440-21-3]

Si

Si

1. Eckhoff, R. K. *et al.*, *J. Electrochem. Soc.*, 1986, **133**, 2631–2637

2. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74

3. See entry Silane (reference 4)

Finely powdered silicon can give significant dust explosion hazards. Relationships of sensitivity to spark ignition and of explosibility to particle size are studied [1]. Maximum explosion pressures of 6.4 bar, with maximum rate of rise of 884 bar/s have been determined [2]. Silicon dust is likely to result from processes using silanes in the gas phase [3].

Calcium

See Calcium: Silicon

Metal acetylides

See Caesium acetylide: Non-metals

Rubidium acetylide: Non-metals

Metal carbonates

Mellor, 1940, Vol. 6, 164

Amorphous or crystalline silicon both react exothermally when heated with alkali-metal carbonates, attaining incandescence and evolving carbon monoxide.

Metal hexafluorides

1. Paine, R. T. *et al.*, *Inorg. Chem.*, 1975, **14**, 1111

2. Paine, R. T. *et al.*, *Inorg. Synth.*, 1979, **19**, 138

During the reduction of iridium hexafluoride, osmium hexafluoride or rhenium hexafluoride with silicon to the pentafluorides, the hexafluorides must not be condensed directly onto undiluted silicon powder, or explosions may result [1]. The same is true for molybdenum hexafluoride and uranium hexafluoride [2].

Oxidants

See Metal hexafluorides, above

Bromine trifluoride: Halogens, etc.

Chlorine trifluoride: Metals, etc.

Chlorine: Non-metals

Cobalt trifluoride: Silicon

Fluorine: Non-metals

Iodine pentafluoride: Metals, etc.

Lead(II) oxide: Non-metals
Nitrosyl fluoride: Metals, etc.
Oxygen difluoride: Non-metals
Peroxyformic acid: Non-metals
Silver fluoride: Non-metals

Water

See Boron: Water

See related METALS, NON-METALS

4910. Silicon–zirconium alloys

[12138-26-0] (1:1)

Si–Zr

[12211-03-9] (1:2)

Si–Zr

See Zirconium (reference 6)

See other ALLOYS

4911. Samarium

[7440-31-5]

Sm

Sm

1,1,2-Trichlorotrifluoroethane

Anon., *NSC Newsletter, R & D Sect.*, 1970, (8)

An attempt to mill a slurry of metal in the solvent caused a violent explosion. Experiment showed violent interaction of the freshly ground metal surfaces with the halocarbon solvent.

See other METAL–HALOCARBON INCIDENTS *See other* METALS

4912. Tin

[7440-31-5]

Sn

Sn

HCS 1980, 901 (powder)

Carbon tetrachloride, Water

Pascal, 1963, Vol. 8.3, 309

In the presence of water vapour, interaction is violent

See other METAL–HALOCARBON INCIDENTS

Disulfur dichloride

See Disulfur dichloride: Tin

Other reactants

Yoshida, 1980, 217

MRH Values calculated for 9 combinations with oxidants are given.

Oxidants	MRH values show % of oxidant
<i>See</i> Ammonium nitrate: Metals	MRH 2.92/58
Chlorine: Metals	MRH 2.09/55
Chlorine trifluoride: Metals	
Copper(II) nitrate: Tin	MRH 2.68/38
Fluorine: Metals	
Iodine bromide: Metals	
Iodine heptafluoride: Metals	
Potassium dioxide: Metals	
Sodium peroxide: Metals	MRH 1.38/56
Sulfur: Metals	
Tellurium: Tin	

Water

1. Board, S. J. *et al.*, *Int. J. Heat Mass Transf.*, 1974, **17**, 331–339
2. Board, S. J. *et al.*, *Chem. Abs.*, 1975, **83**, 12950
3. Frost, D. L. *et al.*, *Progr. Astronaut. Aeronaut.*, 1988, **114** (Dynam. Explos.), 451–473

Experiments involving explosions of molten tin and water are described [1], and the mechanism of propagation of the thermal explosions was studied [2]. The dynamics of explosive interaction between multiple drops of molten tin and water has been examined experimentally [3].

See MOLTEN METAL EXPLOSIONS *See other* METALS

4913. Strontium

[7440-25-7]

Sr

Sr

Sorbe, 1968, 154

The finely divided metal may ignite in air.

Halogens

Pascal, 1958, Vol. 4, 579

At 300°C strontium incandescs in chlorine, and at 400°C it ignites in bromine vapour.

Water

Pascal, 1958, Vol. 4, 579

Interaction is more vigorous than with calcium.

See other METALS, PYROPHORIC METALS

4914. Tantalum

[7440-25-7]

Ta

Ta

Although very unreactive in massive forms, the finely divided metal may be pyrophoric.

See entry PYROPHORIC METALS (references 3,9)

Bromine trifluoride
See Bromine trifluoride: Halogens, etc.

Fluorine
See Fluorine: Metals

Lead chromate
See Lead chromate: Tantalum
See other METALS

4915. Technetium
[7440-26-8]

Tc

Tc

Bailar, 1973, Vol. 3, 881
In finely divided forms (sponge or powder) it readily burns in air to the heptaoxide.
See other METALS, PYROPHORIC METALS

4916. Tellurium
[13494-80-9]

Te

Te

Halogens, or Interhalogens
See Bromine pentafluoride: Acids, etc.
Chlorine: Non-metals
Chlorine fluoride: Tellurium
Chlorine trifluoride: Metals
Fluorine: Non-metals

Hexalithium disilicide
See Hexalithium disilicide: Non-metals

Metals
See Cadmium: Selenium, etc.
Platinum: Selenium, etc.
Potassium: Selenium, etc.
Sodium: Non-metals

Silver iodate
See Silver iodate: Tellurium

Tin
Bailar, 1973, Vol. 2, 80
Interaction hot to give tin telluride attains incandescence.
See Selenium: Metals
See related METALS, NON-METALS

4917. Thorium
[7440-29-1]

Th

Th

1. Mellor, 1941, Vol. 7, 207
2. Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64
3. *Fire Record Bull. FR58-3*, Boston, NFPA, 1958

The massive metal ignites below red-heat when heated in air, and the powdered metal ignites when rubbed or crushed [1], or on pouring as a stream in air, owing to intergranular friction. Storage and handling precautions are discussed [2]. During destruction by burning of radioactive thorium scrap, addition of a 4 cm lump of the metal waste to burning thorium wafers led to a series of violent explosions. This was attributed to presence of water or other impurities in the lump [3].

See THORIUM FURNACE RESIDUES

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, Nitrogen

Halogens

Mellor, 1941, Vol. 7, 207

Thorium incandescs in chlorine, bromine or iodine, and would be expected to ignite readily in fluorine.

Nitryl fluoride

See Nitryl fluoride: Metals

Oxygen

Mellor, 1941, Vol. 7, 207

The massive metal ignites on heating in oxygen.

Peroxyformic acid

See Peroxyformic acid: Metals

Phosphorus

See Phosphorus: Metals

Silver

See Silver–thorium alloy

Sulfur

See Sulfur: Metals

See other METALS, PYROPHORIC METALS

4918. Thorium salts
[16065-92-2]

Th⁴⁺

Th⁴⁺

‘Cupferron’

See Ammonium *N*-Nitrosophenylaminooxide: Thorium salts

4919. Titanium
[7440-32-6]

Ti

Ti

NSC 485, 1968; FPA H87, 1980

1. Mellor, 1941, Vol. 7, 19
2. Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64
3. *NSC Data Sheet 485*, 1968
4. Lowry, R. N. *et al.*, *Inorg. Synth.*, 1967, **10**, 3
5. *Standard for Production, Processing, Handling and Storage of Titanium*, NFPA 481, Quincy (Ma), 1987
6. Alekseev, A. G. *et al.*, *Chem. Abs.*, 1975, **83**, 83527
7. Popov, E. I. *et al.*, *Chem. Abs.*, 1975, **83**, 150922
8. Elrod, C. W., *Chem. Abs.*, 1982, **97**, 184846
9. Poyarkov, V. G. *et al.*, *Chem. Abs.*, 1983, **98**, 37077
10. Grossel, S. S., *J. Loss Prev. Proc. Ind.*, 1988, **1**, 62–74
11. Prine, B. A., *Plant/Oper. Progr.*, 1992, **11**(2), 113
12. Mahnken, G. E. *et al.*, *Process Safety Progress*, 1997, **16**(1), 54

While massive titanium ignites in air only at 1200°C [1], the finely divided metal is very pyrophoric and should be stored damp in metal containers [2]. Titanium fires are difficult to extinguish as it burns brilliantly in nitrogen above 800°C, in carbon dioxide above 550°C [3], and may react explosively with metal carbonates [1]. Residual titanium sponge after vapour phase reaction with iodine at 400–425°C is often pyrophoric [4]. All aspects of preventing hazards in use of titanium are covered in a US Standard [5]. Powdered metal made by calcium hydride reduction shows a greater explosion hazard than electrolytically reduced metal. The effect of inert gases was also examined [6]. The relationship between spontaneous ignition temperatures (250–600°C) and lower explosive limits (40–300 mg/l) of titanium powders was studied [7]. Incidents involving combustion of titanium components in aviation gas turbines are reviewed [8]. Ignition parameters for titanium powder of <100 μ particle size were determined. Ignition temperature decreases from 640 to 508°C with increase in specific surface area from 960 to 1520 cm²/g [9]. Maximum explosion pressures of 5.8 bar, with maximum rates of rise of 816 bar/s have been determined [10]. Titanium combustion as a result of welding operations appears to be potentiated by carbon steel, there are several recorded cases, mechanism not established [11]. Another fire when cutting into a steel-clad Ti heat-exchanger, with explosions when sprayed with water is reported [12].

Aluminium

See Aluminium–titanium alloys

Bromine trifluoride

See Bromine trifluoride: Halogens, etc.

Carbon black

Afanas'eva, L. F. *et al.*, *Chem. Abs.*, 1981, **95**, 64482

Fire hazards and pyrophoricity of mixed charges for preparing titanium carbide are discussed.

Carbon dioxide, or Metal carbonates

1. Stout, E. L., *Chem. Eng. News*, 1958, **36**(8), 64

2. Mellor, 1941, Vol. 7, 20

Titanium powder has ignited as a thin layer under carbon dioxide, and burns in the gas above 550°C [1]. Contact of titanium with fused alkali-metal carbonates causes incandescence [2].

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, or: Metals, Nitrogen

Halocarbons

Pot. Incid. Rep. 39, ASESB, Washington, 1968

Mixtures of powdered titanium and trichloroethylene or 1,1,2-trichlorotrifluoroethane flash or spark under heavy impact.

See other METAL–HALOCARBON INCIDENTS

Halogens

1. Mellor, 1941, Vol. 7, 192

2. Kirk-Othmer, 1969, Vol. 20, 372

3. Denes, G., *Chem. Eng. News*, 1987, **65**(7), 2

4. Fast, J. D., *Rec. Trav. Chim.*, 1939, **58**, 174

5. Cobb, C. M., *Chem. Eng. News*, 1987, **65**(14), 2

6. Sisler, H. H., *Chem. Abs.*, 1988, **108**, 15143

Reaction of heated titanium with the halogens usually causes ignition and incandescence (chlorine at 350°C, bromine at 360°C, iodine higher, but weak combustion) [1], especially in absence of moisture [2]. Attempts to prepare titanium tetraiodide (5 g) by heating titanium powder with iodine in sealed copper tubes heated in a water bath led to explosions when the bath temperature reached 90°C. Rupture of the tubes also occurred with 0.5 g preparations [3]. Attempts to prepare titanium diiodide according to an earlier high-vacuum procedure [4] also led to violent explosions after two days [5]. A procedure to prepare the tetraiodide by heating a mixture of titanium and excess iodine in a Pyrex glass apparatus is reported [6].

See other HALOGENATION INCIDENTS

See Chlorine: Metals (references 2,3), *also* Fluorine: Metals

Metal oxides MRH Chromium trioxide 4.85/73

Mellor, 1961, Vol. 7, 20

Interaction with copper(II) oxide or lead(II,IV) oxide is violent.

Metal oxosalts

Mellor, 1941, Vol. 7, 20

Contact of powdered titanium with molten potassium chlorate, alkali-metal carbonates or mixed potassium carbonate/nitrate causes vivid incandescence.

Nitric acid

MRH 8.79/51

1. Anon., *Occ. Health and Safety*, 1957, **7**, 214

2. Anon., *Chem. Eng. News*, 1953, **31**, 3320

Investigation showed that commercial titanium alloys in contact with acid containing less than 1.34% water and more than 6% of dinitrogen tetraoxide may become sensitive to impact, and react explosively with the acid. Possible causes are discussed [1]. The spongy residue formed by prolonged corrosion of titanium–manganese alloys by red fuming nitric acid will explode on exposure to friction or heat [2].

See other CORROSION INCIDENTS

Nitrogen

Bailar, 1973, Vol. 3, 359

Titanium is claimed to be the only element that will burn in nitrogen.

See Lithium, Nitrogen: Magnesium

Nitryl fluoride

See Nitryl fluoride: Metals

Other reactants

Yoshida, 1980, 231

MRH values for 13 combinations with oxidants are given.

Oxidants

Mellor, 1941, Vol. 7, 20

MRH Ammonium nitrate 5.48/77, hydrogen peroxide 8.74/59, potassium permanganate 3.97/63, sodium bromate 6.19/68, sodium chlorate 8.45/60, sodium chlorite 7.36/65, sodium nitrate 6.07/58, sodium perchlorate 8.37/57, sodium peroxide 3.51/76

Mixtures of the powdered metal with potassium chlorate, potassium nitrate or potassium permanganate react explosively on heating. Addition of the powdered metal to fused potassium chlorate at 370°C leads to incandescence.

See Metal oxides; Nitric acid, both above

Iodine: Metals

Potassium perchlorate: Metal powders (references 1,3)

Sodium chlorate, below

Oxygen

1. Kirk-Othmer, 1969, Vol. 20, 372

2. Komamiya, K., *Chem. Abs.*, 1982, **96**, 90906

3. Shenoy, V. R., *Chem. Eng. News*, 1992, **70**(40), 4

4. Anon., *Sichere Chemiearbeit*, 1994, **46**(6), 67

Ignition may occur in liquid or gaseous oxygen and in gas mixtures containing over 46% oxygen, under impact or abrasion. Mixtures of powdered titanium and liquid oxygen are detonable [1]. In the absence of a surface film of grease, etc. on a freshly abraded surface, titanium will not burn under high pressure oxygen at high temperature [2]. Two explosions were experienced when studying treatment of ppm levels of organics in waste under 3-4 bar of oxygen at around 200°C. In each case the problem seems to have started at joints where abrasion, flakes of metal, grease and/or polymeric seals might have been present [3]. An inaccessible valve of partly titanium construction in a line filled with oxygen at 140°C (pressure unspecified) ignited for no apparent reason and burnt out. Titanium is not a suitable material of construction for oxygen lines, being even less protected by its oxide coating than is aluminium [4].

See Oxygen (Gas): Titanium, *and* Oxygen (Liquid): Metals, *also* Oxygen (Gas): Polytetrafluoroethylene, Stainless steel

Platinum group metals

Ullmann, 1992, Vol. A21, 125/6

Alloy formation between the powdered metals may be violent

Silver fluoride

See Silver fluoride: Titanium

Silver nitrate

See Silver nitrate: titanium

Sodium chlorate, Water

MRH 8.45/60

Scribner, H. C., *Tappi J.*, 1986, **69**(12), 16

A steam-heated titanium heat exchanger was used to heat sodium chlorate solution passing through its tubes, some of which were blocked by polypropylene film and wood chip debris. This led to local boiling and separation of solid sodium chlorate, which ignited the organic material and heated the titanium tube to its ignition point (427°C). It then started to burn, reacting with steam to liberate hydrogen which led to a violent explosion.

See other GAS EVOLUTION INCIDENTS

Water

1. Mellor, 1941, Vol. 7, 19

2. Anon., *Loss Prev. Bull.*, 1987, (075), 22

Finely divided titanium reacts with steam at 700°C, and in presence of air the evolved hydrogen may ignite or explode [1]. A heat exchanger with titanium tubes was being dismantled with flame cutting equipment, which caused ignition of the hot titanium in air. Application of water to extinguish the violent fire led to evolution of hydrogen and then an explosion [2].

See other METALS, PYROPHORIC METALS

4920. Titanium salts
[16043-45-1] (4+)

TiZ₄

TiZ₄

‘Cupferron’

See Ammonium *N*-nitrosophenylaminoxide: Thorium salts

4921. Titanium–zirconium alloys
[50646-37-2] (1:1)

Ti–Zr

Ti–Zr

See Zirconium (reference 6)

See other ALLOYS

4922. Thallium
[7440-28-0]

Tl

Tl

Fluorine

See Fluorine: Metals

See other METALS

4923. Uranium
[7440-61-1]

U

U

1. *MCA Case History No. 1296*

2. Bailar, 1973, Vol. 5, 40

3. Rieke, R. D. *et al.*, *J. Org. Chem.*, 1979, **44**, 3446

4. Starks, D. F. *et al.*, *J. Amer. Chem. Soc.*, 1973, **95**, 3423

Storage of uranium foil in closed containers in presence of air and water may produce a pyrophoric surface [1]. Uranium must be machined in a fume hood because, apart from the radioactivity hazard, the swarf is easily ignited. The massive metal ignites at 600–700°C in air [2]. The finely divided reactive form of uranium produced by pyrolysis of the hydride is pyrophoric [3], while that produced as a slurry by reduction of uranium tetrachloride in dimethoxyethane by potassium–sodium alloy is not [4].

Ammonia

Mellor, 1942, Vol. 12, 31

At dull red heat the metal incandesces in ammonia.

Bromine trifluoride

See Bromine trifluoride: Uranium, etc.

Carbon dioxide

Bailar, 1973, Vol. 5, 42

At 750°C interaction is so rapid that ignition will occur with the finely divided metal, and at 800°C the massive metal will ignite.

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, Nitrogen

Carbon tetrachloride

491M, 1975, 435

Use of a carbon tetrachloride fire extinguisher on a small uranium fire led to an explosion.

See other METAL-HALOCARBON INCIDENTS

Halogens

Mellor, 1942, Vol. 12, 31

Uranium powder ignites in fluorine at ambient temperature, in chlorine at 150–180°, in bromine vapour at 210–240° and in iodine vapour at 260°C.

Nitric acid, Trichloroethylene

MCA Case History No. 1104

Uranium scrap containing rather fine turnings was being pickled, after degreasing with trichloroethylene, by treatment with 4–6 M nitric acid solution. During subsequent rinsing with cold water, an explosion ejected the metal and acid liquor from the beaker. It has been reported in the literature that uranium reacts with nitric acid or dinitrogen tetraoxide either explosively, or with the formation of an explosive surface coating or residue. This corrosive effect is inhibited by presence of fluoride ion during pickling. Preventive measures proposed include separate treatment of finely divided uranium scrap, thorough pre-mixing of the acid solution and addition of ammonium hydrogen fluoride. An alternative, or contributory, cause could have been the violent reaction between dinitrogen tetraoxide and residual trichloroethylene.

See Dinitrogen tetraoxide: Halocarbons

See other CORROSION INCIDENTS

Nitrogen oxide

See Nitrogen oxide: Metals

Nitryl fluoride

See Nitryl fluoride: Metals

Selenium, or Sulfur

Mellor, 1942, Vol. 12, 31

Uranium incandescences in sulfur vapour, and with selenium also.

Water

See Magnesium: Water, Uranium

See other METALS, PYROPHORIC METALS

4924. Vanadium
[7440-68-2]

V

V

Pascal, 1958, Vol. 12, 45

Some samples of vanadium prepared by magnesium reduction of vanadium dichloride or vanadium trichloride are pyrophoric.

Oxidants

See Bromine trifluoride: Halogens, etc.

Chlorine: Metals

Nitryl fluoride: Metals

See other METALS, PYROPHORIC METALS

4925. Tungsten
[7440-33-7]

W

W

Mellor, 1943, Vol. 11, 729

The finely divided metal may ignite on heating in air.

Oxidants

1. Mellor, 1943, Vol. 11, 729

2. Rose, J. E., *Chem. Abs.*, 1980, **92**, 8483

The finely divided metal may ignite in contact with a range of oxidants, usually on heating [1]. Mixtures of the metal powder with barium chromate and potassium chlorate are used as a pyrotechnic fuse [2].

See Bromine: Tungsten, etc.

Bromine pentafluoride: Acids, etc.

Chlorine trifluoride: Metals

Fluorine: Metals

Hydrogen sulfide: Metals

Iodine pentafluoride: Metals

Lead(IV) oxide: Metals

Nitryl fluoride: Metals

Oxygen difluoride: Metals

Potassium dichromate: Tungsten

Potassium perchlorate: Barium chromate, etc.

Sodium peroxide: Metals

See other METALS, PYROPHORIC METALS

4926. Xenon
[7440-63-3]

Xe

Xe

Sponsles, M. *et al.*, *J. Amer. Chem. Soc.*, 1989, **111**(17), 6841

1919

Liquid xenon, used as a solvent, will burst sealed apparatus if allowed to warm up. Pressure relief valves are recommended.

Fluorine

See Fluorine: Xenon

See other NON-METALS

4927. Zinc

[7440-66-6]

Zn

Zn

NSC 267, 1978; FPA H96, 1980; HCS 1980, 966 (dust); RSC Lab. Hazard Data Sheet No. 76, 1988 (zinc and compounds)

1. Anon., *Chem. Ztg.*, 1940, 289
2. Anon., *Ind. Eng. Chem.*, 1944, **20**, 330
3. Robson, S., *School Sci. Rev.*, 1934, **16**(62), 165
4. Anon., *Jahresber.*, 1978, 72

A cloud of zinc dust generated by sieving the hot dried material exploded violently, apparently after initiation by a spark from the percussive sieve-shaking mechanism [1]. Precautions recommended include use of cold zinc and total enclosure of such processes [2]. The possibility of explosions of zinc dust suspended in air is presented as a serious hazard [3]. A serious dust explosion in the air filter unit of a zinc grinding mill was initiated by a spark from an explosion relief panel [4].

See Zinc chloride, below

Acetic acid

Wehrli, S., *Chem. Fabrik*, 1940, 362–363

The residues from zinc dust–acetic acid reduction operations may ignite after a long delay if discarded into waste-bins with paper. Small amounts appear to ignite more rapidly than larger portions.

See other PYROPHORIC MATERIALS

Aluminium

See Aluminium: Zinc

Aluminium, Magnesium, Rusted steel

See Aluminium–magnesium–zinc alloy: Rusted steel

Ammonium nitrate,

MRH 3.22/5

Ammonium chloride, Water

1. Sorbe, 1968, 158
2. Hanson, R. M., *J. Chem. Educ.*, 1976, **53**, 578

A mixture of zinc dust with ammonium nitrate [1], or mixed nitrate and chloride [2] ignites when moistened.

See Ammonium nitrate: Metals

Ammonium sulfide

See Ammonium sulfide: Zinc

Arsenic alloy, Water

Carson, P. A. *et al.*, *Loss Prev. Bull.* 1991, **102**, 7

An arsenic alloy in a galvanised container with water generated the extremely toxic gas arsine.

See Aluminium: Arsenic trioxide, Sodium arsenate, Sodium hydroxide

See Water, below

Arsenic trioxide

Mellor, 1940, Vol. 4, 486

A mixture with excess zinc filings explodes on heating

Calcium chloride

See Calcium chloride: Zinc

Carbon disulfide

MRH 2.43/37

Mellor, 1940, Vol. 4, 4

Carbon disulfide and zinc powder interact with incandescence.

Carbon tetrachloride,

MRH 2.43/54

Methanol

See Methanol: Carbon tetrachloride, Metals

Chlorinated rubber

See Halocarbons (reference 2), below

Catalytic metals, Acids

Sidgwick, 1950, 1530, 2578

Alloys of zinc with iridium, platinum or rhodium, after extraction with acid, leave residues which explode on warming in air, owing to the presence of occluded hydrogen (or oxygen) in the catalytic metal powders so produced.

Electrolytes, Silver

See Silver: Electrolytes, etc.

Ethyl acetoacetate, Tris(bromomethyl)ethanol

See Ethyl acetoacetate: 2,2,2-Tris(bromomethyl)ethanol, Zinc

Halocarbons

MRH Carbon tetrachloride 2.43/54

1. Berger, E., *Compt. rend.*, 1920, **170**, 29

2. Anon., *ABCM Quart. Safety Summ.*, 1963, **34**, 12

3. Lamouroux, A. *et al.*, *Mém. Poudres*, 1957, **39**, 435–445

A paste of zinc powder and carbon tetrachloride (with kieselguhr as thickener) will readily burn after ignition by a high-temperature primer [1]. Intimate mixtures of

chlorinated rubber with powdered zinc (or its oxide) in presence or absence of hydrocarbon or halocarbon solvents react violently or explosively at about 216°C [2]. Powdered zinc initially reacts more violently with hexachloroethane in ethanol than does aluminium [3].
See Bromomethane: Metals *also* Chloromethane: Metals
See also CHLORINATED RUBBER: metal oxides
See other METAL-HALOCARBON INCIDENTS

Halogens, or Interhalogens MRH Chlorine 3.05/52, fluorine 7.45/47
Mellor, 1940, Vol. 4, 476
Warm zinc powder incandesces in fluorine, and zinc foil ignites in cold chlorine if traces of moisture are present.
See Bromine pentafluoride: Acids, etc.
Chlorine trifluoride: Metals
Iodine: Metals, Water

Lead azide MRH 1.63/99+
See Lead(II) azide: Copper, etc.

Manganese dichloride
Terreill, A., *Bull. Soc. Chim. Fr.* [2], 1874, **21**, 289
Zinc foil reacts explosively when heated with anhydrous manganese dichloride.

Metal oxides MRH Chromium trioxide 2.51/50, potassium peroxide 1.72/45
See Potassium dioxide: Metals
Titanium dioxide: Metals
Zinc peroxide: Alone, etc.

2-Nitroanisole, Sodium hydroxide
See 2-Nitroanisole: Sodium hydroxide, etc.

Nitrobenzene MRH 4.52/99+
Muir, G. D., private comm., 1968
Zinc residues from reduction of nitrobenzene to *N*-phenylhydroxylamine are often pyrophoric and must be kept wet during disposal.

Nitrogen compounds MRH values show % of nitrogen compound
See Hydrazinium nitrate: Alone, etc. MRH 1.67/50
Hydroxylamine: Metals MRH 3.85/99+

Non-metals MRH Sulfur 2.09/33
1. Mellor, 1940, Vol. 4, 480, 485
2. Read, C. W. W., *School Sci. Rev.*, 1940, **21**(83), 976
Arsenic, selenium and tellurium all react with incandescence on heating [1]. Interaction on heating powdered zinc and sulfur is considered to be too violent for use as a school experiment [2].

Other reactants

Yoshida, 1980, 1–2

MRH values calculated for 31 combinations, mainly with oxidants, are given.

Oxidants

MRH values show % of oxidant

See Halogens; Interhalogens; Metal oxides, all above

Ammonium nitrate: Metals

MRH 3.22/55

Nitric acid: Metals

MRH 3.59/28

Nitryl fluoride: Metals

Peroxyformic acid: Metals

MRH 5.69/99+

Potassium chlorate: Metals

MRH 3.63/38

Paint primer base

Muller, K. A., *Spontaneous Combustion of Zinc-rich Paint Mixtures*, Washington, API, 1969

Zinc residues from poorly mixed priming paint spontaneously inflamed after prolonged exposure to atmosphere. Aluminium residues may behave similarly.

Pentacarbonyliron, Transition metal halides

See Pentacarbonyliron: Transition metal halides, etc.

Seleninyl bromide

See Seleninyl bromide: Metals

Sodium hydroxide

1. Anon., *ABCM Quart. Safety Summ.*, 1963, **34**, 14

2. Jackson, H., *Spectrum*, 1969, **7**(2), 82

3. Summerlin, L. *unpublished letter to Chem. Eng. News*, 1988

Accidental contamination of a metal scoop with flake sodium hydroxide, prior to its use with zinc dust, caused ignition of the latter [1]. A stiff paste prepared from zinc dust and 10% sodium hydroxide solution attains a temperature above 100°C after exposure to air for 15 min [2]. The residue of zinc dust and sodium hydroxide solution from a lecture demonstration involving zinc plating a copper coin presents a high fire risk if discarded onto paper in a waste bin, ignition of the paper having occurred on many occasions. Dissolve the residue in dilute acid before flushing away with water [3].

See PAPER TOWELS

See Water, next below

Water

1. Anon., *Metallwirtschaft*, 1941, **20**, 475

2. Anon., *Chem. Fabrik*, 1940, 362

3. Mellor, 1940, Vol. 4, 474

4. Carson, P. J. *et al.*, *Loss Prev. Bull.*, 1992, **102**, 15

In contact with atmospheric oxygen and limited amounts of water, zinc dust will generate heat, and may become incandescent [1]. Presence of acetic acid and copper shortens the induction period. Store the metal dry, and keep residues

thoroughly wet until disposal [2]. Hydrogen is evolved, especially under acid or alkaline conditions [3]. An incident of a burst galvanised drum from hydrogen evolution is recorded [4].

See other CORROSION INCIDENTS, GAS EVOLUTION INCIDENTS, INDUCTION PERIOD INCIDENTS

Zinc chloride

Bylo, Z. *et al.*, *Chem. Abs.*, 1983, **98**, 146031

Flammability of some commercial zinc dusts or powders is attributed to presence of zinc chloride.

See other METALS, PYROPHORIC METALS, REDUCANTS

4928. Zirconium

[7440-67-7]

Zr

Zr

(MCA SD-92, 1966); NSC 382, 1968 (powder)

1. *Standard for Production, Processing, Handling and Storage of Zirconium*, NFPA 482M, 1974
2. *Zirconium Fire and Explosion Incidents*, TID-5365, Washington, USAEC, 1956
3. Gleason, R. P. *et al.*, *J. Amer. Soc. Safety Eng.*, 1964, **9**(3), 15
4. Bulmer, G. H. *et al.*, *Rept. AHSB(S) R59*, London, UKAEA, 1964
5. *MCA Case History No. 1234*
6. Ioffe, V. G., *Chem. Abs.*, 1967, **66**, 12614d
7. Karłowicz, P. *et al.*, *J. Electrochem. Soc.*, 1961, **108**, 659–663
8. Leuschke, G. *et al.*, *Chem. Abs.*, 1983, **98**, 221174
9. Baumgart, H., *Chem. Abs.*, 1985, **103**, 73412
10. Cooper, T D., *Chem. Abs.*, 1986, **104**, 65449
11. *See entry* DUST EXPLOSION INCIDENTS (reference 22)
12. Bond, 1991, 60

All aspects of hazards in the production, use and storage of zirconium are covered in a US Standard [1], and there is a compilation of 43 abstracts of unusual zirconium fire and explosion incidents, most of which involved the pyrophoric finely divided metal, moisture and friction, occasionally accompanied by the possibility of static sparks from polythene bags [2]. The increasing use of zirconium necessitated a review of methods of controlling the pronounced pyrophoric properties of zirconium powder. Five considerations for the safe use of zirconium include: exclusion of air or oxygen by use of inert gases; exclusion of water, its vapour and other contaminants or oxidants; control of particle size; limitation of amount of powder handled or exposed; and limitation of exposure of personnel [3]. Storage and disposal of irradiated metal wastes in various forms has also been discussed [4]. Ignition may not be immediate, however, as fine zirconium dust left uncleared from previous grinding operations was ignited violently by a spark from subsequent grinding operations [5]. The lower ignition limits of air suspensions of zirconium powder, of its alloys with silicon and titanium, and of zirconium–titanium hydride have been studied in relation to the history of exposure to air [6]. Treatment of

Zr

zirconium powder with 1% hydrogen fluoride solution considerably desensitises it to electrostatic ignition [7]. Particles of Zircaloy (98–99% Zr) blown from tubes by compressed air ignited by friction or impact, even at 2 bar blow-off pressure, and could initiate dust explosions [8]. Ignition and combustion properties of pyrotechnic zirconium and zirconium hydride powders (<10 μ) and safety measures are discussed [9]. In a review of pyrophoricity of zirconium and its alloys, it emerged that the maximum particle size for pyrophoric behaviour is 54 μ , and that smaller particles may also explode as well as ignite in air. Factors influencing ignition temperature were explored, and recommendations made for safe handling, transportation and storage [10]. In dust explosions, maximum pressures of 5.1 bar, with maximum rates of rise of 748 bar/s have been determined [11]. Flakes of zirconium came away when a manhole cover was opened on a zirconium-distillation column containing methanol. The zirconium particles ignited, setting fire to the methanol vapour [12].

See Lead–zirconium alloys

See METAL DUSTS, PYROPHORIC METALS

See Hafnium

Carbon dioxide, Nitrogen

See Carbon dioxide: Metals, or: Metals, Nitrogen

Carbon tetrachloride

1. *Zirconium Fire and Explosion Incidents*, TID-5365, Washington, USAEC, 1956

2. *491M*, 1975, 446

A mixture of powdered zirconium and carbon tetrachloride exploded violently while being heated [1], and immersion of zirconium sponge in the unheated solvent also caused explosion [2].

See other METAL–HALOCARBON INCIDENTS

Nitryl fluoride

See Nitryl fluoride: Metals

Oxygen-containing compounds

1. Mellor, 1941, Vol. 7, 116

2. Ellern, 1968, 249

The affinity of zirconium for oxygen is great, particularly when the metal is finely divided, and extends to many oxygen-containing compounds. The vigorous reactions with potassium chlorate, potassium nitrate, copper(II) oxide or lead oxide might be anticipated, but there are also explosive reactions with the combined oxygen when zirconium is heated with alkali-metal hydroxides or carbonates, or even zirconium hydroxide. Hydrated sodium tetraborate behaves similarly [1]. Several alkali-metal oxosalts (chromates, dichromates, molybdates, sulfates, tungstates) react violently or explosively when reduced to the parent metals by heating with zirconium [2].

See Copper(II) oxide: Metals

Lead(II) oxide: Metals

Lithium chromate: Zirconium

Potassium chlorate: Metals
Potassium nitrate : Metals

Phosphorus

See Phosphorus: Metals

Sulfuric acid, Potassium bisulfate

Ullmann, 1996, A28, 554,

Zirconium powder mixed with an approximately equal quantity of 17% potassium bisulfate in concentrated sulfuric acid exploded violently.

See Oxygen-containing compounds, above

Water

1. *MCA SD-92*, 1966

2. Muir, G. D., private comm., 1968

3. *Haz. Chem Data*, 1975, 305

4. Hall, A. N. *et al.*, *Loss Prev. Bull.*, 1987, (076), 22–29

Powder damp with 5–10% of water may ignite, and although 25% of water is regarded as a safe concentration [1], ignition of a 50% paste on breakage of a glass container has been observed [2]. Although water is used to prevent ignition, the powder, once ignited, will burn under water (88.8% oxygen) more violently than in air (21% oxygen) [3]. The relatively minor part played by steam and zircalloy fuel containment cans (ruptured by the steam explosion) to form hydrogen towards the end of the catastrophic sequence involved in the Chernobyl nuclear reactor accident is noted in a comprehensive account of the causes of the disaster [4].

See other METALS, PYROPHORIC METALS

4929. Zirconium salts

[15543-40-5]

Zr⁴⁺

Zr⁴⁺

‘Cupferron’

See Ammonium *N*-nitrosophenylaminoxide: Thorium salts

Source Title Abbreviations used in Handbook References

The abbreviations used in the references for titles of journals and periodicals are those used in BP publications practice and conform closely to the recommendations of the *Chemical Abstracts* system. Abbreviations which have been used to indicate textbook and reference book sources of information are set out below with the full titles and publication details.

<i>ABCM Quart. Safety Summ.</i> ,	<i>Quarterly Safety Summaries</i> , London, Association of British Chemical Manufacturers, 1930–1964
<i>ACS 54</i> , 1966	<i>Advanced Propellant Chemistry</i> , ACS 54, Washington, American Chemical Society, 1966
<i>ACS 88</i> , 1969	<i>Propellants Manufacture, Hazards and Testing</i> , ACS 88, Washington, American Chemical Society, 1969
Albright, Hanson, 1976	<i>Industrial and Laboratory Nitrations</i> , ACS Symposium Series 22, Albright, L. F., Hanson, C., (Eds.), Washington, American Chemical Society, 1976
<i>Alkali Metals</i> , 1957	<i>Alkali Metals</i> , ACS 19, Washington, American Chemical Society, 1957
<i>ASHRAE. J.</i>	<i>American Society of Heating, Refrigeration and Air-conditioning Engineers Journal</i> , New York.
Augustine, 1968	<i>Reduction, Techniques and Applications in Organic Chemistry</i> , Augustine, R. L., London, Edward Arnold, 1968

- Augustine, 1969, 1971 *Oxidation, Techniques and Applications in Organic Synthesis*, Augustine, R. L., (Ed.), New York, Marcel Dekker, Vol. 1, 1968, Vol. 2 (with Trecker, D. J.), 1971
- Bahme, 1972 *Fire Officer's Guide to Dangerous Chemicals*, FSP-36, Bahme, C. W., Boston, National Fire Protection Association, 1972
- Bailar, 1973 *Comprehensive Inorganic Chemistry*, Bailar, J. C., Emeléus, H. J., Nyholm, R. S., Trotman-Dickenson, A. F., (Eds.), Oxford, Pergamon, 5 vols., 1973
- BCISC Quart. Safety Summ.*, *Quarterly Safety Summaries*, London, British Chemical Industries Safety Council, 1965–1973
- Barton & Ollis, 1979 *Comprehensive Organic Chemistry*, Barton, D. H. R., Ollis, W. D., (Eds), Oxford, Pergamon, 1979.
- Biasutti, 1981 *History of Accidents in the Explosives Industry*, Biasutti, G. S. (Author and publisher), Vevey, 1981
- Bond, 1991 *Sources of Ignition*, Bond, J., Oxford, Butterworth-Heinemann, 1991
- Braker, 1981 *Matheson Gas Data Book*, Braker, W., Mossman, A. L., Lyndhurst (NJ), Matheson Div. Searle, 6th edn. 1981
- Brandsma, 1971 *Preparative Acetylenic Chemistry*, Brandsma, L., Barking, Elsevier, 1971
- Brauer, 1961, 1965 *Handbook of Preparative Inorganic Chemistry*, Brauer, G., (Translation Ed. Riley, R. F.), London, Academic Press, 2nd edn. Vol. 1, 1963; Vol. 2, 1965
- Bretherick, 1986 *Hazards in the Chemical Laboratory*, Bretherick, L. (Ed.), London, Royal Society of Chemistry, 4th edn., 1986
- Castrantas, 1965 *Fire and Explosion Hazards of Peroxy Compounds*, Special Publication No. 394, Castrantas, H. M., Banerjee, D. K., Noller, D. C., Philadelphia, ASTM, 1965
- Castrantas, 1970 *Laboratory Handling and Storage of Peroxy Compounds*, Special Publication No. 491, Castrantas, H. M., Banerjee, D. K., Philadelphia, ASTM, 1970

- CHAS Notes* *Chemical Health and Safety Notes*, ACS Division of Chemical Health and Safety, Washington, 1982 to date (4 issues/yr)
- Chemiarbeit* *Chemiearbeit, Schutzen und Helfen*, Dusseldorf, Berufsgenossenschaft der Chemische Industrie, 1949–1962 (supplement to *Chemische Industrie*)
- Chem. Hazards Ind.* *Chemical Hazards in Industry*, Royal Society of Chemistry Information Group, Cambridge, 1984 to date (monthly abstracts)
- CISHC Chem. Safety Summ.*, *Chemical Safety Summaries*, London, Chemical Industry Safety and Health Council of the Chemical Industries Association, 1974 to 1986 (terminated)
- CHETAH 1990* *The ASTM Chemical Thermodynamic and Energy Release Evaluation Programme*, Version 4.4, 2nd Edn (ASTM Data Series Publication 51A), Davies C. A. *et al.*, Philadelphia, ASTM, Appendix V. Upgrades continue to be issued (1998)
- Cloyd, 1965 *Handling Hazardous Materials*, Technical Survey No. SP-5032, Cloyd, D. R., Murphy, W. J., Washington, NASA, 1965
- Coates, 1960 *Organometallic Compounds*, Coates, G. E., London, Methuen, 1960
- Coates, 1967, 1968 *Organometallic Compounds*, Coates, G. E., Green, M. L. H., Wade, K., London, Methuen, Vol. 1, 1967, Vol. 2, 1968
- Compendium* 1985, 1987 *Compendium of Safety Data Sheets, for Research and Industrial Chemicals*, Keith, L. H., Walters, D. B., Deerfield Beach/Weinheim, VCH, vols. 1–3 of 867 sheets 1985, vols 4–6 of 723 sheets, 1987
- Dangerous Loads*, 1972 *Dangerous Loads*, Leicester, Institution of Fire Engineers, 1972
- Dangerous Substances*, 1972 *Dangerous Substances: Guidance on Fires and Spillages* (Section 1, 'Inflammable liquids'), London, HMSO, 1972
- Davies, 1961 *Organic Peroxides*, Davies, A. G., London, Butterworths, 1961
- Davis, 1943 *Chemistry of Powder and Explosives*, Davis, T. L., New York, Wiley, 1943

- DOC 5, 1982 *Dictionary of Organic Compounds*, Buckingham, J., (Ed.), London, Chapman & Hall, 5th edn. in 5 vols. + indices, 1982, annual supplementary vols. to date
- Dunlop, 1953 *The Furans*, ACS 119, Dunlop, A. P., Peters, F. N., New York, Reinhold, 1953
- Ellern, 1968 *Military and Civilian Pyrotechnics*, Ellern, H., New York, Chemical Publishing Co., 1968
- Emeléus, 1960 *Modern Aspects of Inorganic Chemistry*, Emeléus, H. J., Anderson, J. S., London, Rutledge and Kegan Paul, 1960
- Ephraim, 1939 *Inorganic Chemistry*, Ephraim, F., (3rd Engl. edn., Thorne, P. C. L., Ward, A. M.), London, Gurney & Jackson, 1939
- Federoff, 1960 *Encyclopaedia of Explosives and Related Compounds*, Federoff, B. T. (Ed.), Dover (NJ), Piccatinny Arsenal, Vol. 1, 1960; Vol. 10, 1984 (from USNTIS)
- Fieser, 1967–1986 *Reagents for Organic Synthesis*, Fieser, L. F. and Fieser, M., New York, Wiley, Vol. 1, 1967, The series continues after the death of the originator.
- Freifelder, 1971 *Practical Catalytic Hydrogenation*, Freifelder, M., New York, Wiley-Interscience, 1971
- Gallais, 1957 *Chimie Minérale Théorique et Expérimentale (Chimie Électronique)*, Gallais, F., Paris, Masson, 1957
- Gaylord, 1956 *Reductions with Complex Metal Hydrides*, Gaylord, N. G., New York, Interscience, 1956
- Gibson, 1969 *Handbook of Selected Properties of Air-and Water-Reactive Materials*, Gibson, H., Crane (In.), US Naval Ammunition Depot, 1969
- Goehring, 1957 *Ergebnisse und Probleme der Chemie der Schwefelstickstoff-verbinding*, *Scientia Chemica*, Vol. 9, Goehring, M., Berlin, Akademie Verlag, 1957
- Grignard, 1935–54 *Traité de Chimie Organique*, Grignard, V. and various volume Eds., Paris, Masson, Vol. 1, 1935, Vol. 23, 1954

- Guide for Safety*, 1972 *Guide for Safety in the Chemical Laboratory*, MCA, New York, Van Nostrand Reinhold, 2nd edn., 1972
- Harmon, 1974 *A Review of Violent Monomer Polymerisation*, Harmon, M., King, J., Rept. AD-017443, Springfield (Va.), NTIS, 1974
- Haz. Chem. Data*, 1975 *Hazardous Chemical Data*, NFPA 49, Boston, National Fire Protection Association, 1975
- HCS 1980 *Handling Chemicals Safely*, Dutch Assoc. of Safety Experts, Dutch Chemical Industry Assoc., Dutch Safety Inst., 1980 (Engl. transl. of 2nd Dutch edn.)
- Houben-Weyl, 1953 to date *Methoden der Organischen Chemie*, Müller, E. (Ed.), Stuttgart, Georg Thieme, 4th edn., published unsequentially from 1953.
- Inorg. Synth.*, 1939–86 *Inorganic Syntheses*, various Eds., Maidenhead, McGraw-Hill, Vol. 1, 1939, Vol. 24, 1986
- Jahresber. 1960 to date *Jahresberichte*, Berufsgenossenschaft der Chemischen Industrie, Heidelberg, annually since 1960
- JANAF, 1971 *JANAF Thermochemical Tables*, Stull, D. R., Prophet, H., National Bureau of Standards, 2nd edn., 1971
- Karrer, 1950 *Organic Chemistry*, Karrer, P., London, Elsevier, 1950
- Janz, 1976 *Eutectic Data, Table 2 (Safety Hazards, etc.)* Janz, G. J., Allen, C. B., Molten Salts Data Center, ERDA Rept. TID 27163-21, 1976; subsequently publ. in *J. Haz. Mat.*, 1980–81, **4** (1), 145–176
- Kharasch and Reinmuth, 1954 *Grignard Reactions of Non-Metallic Substances*, Kharasch, M. S. and Reinmuth, O., London, Constable, 1954
- King, 1990 *Safety in the Process Industries*, King, R., London, Butterworth-Heinemann, 1990
- Kirk-Othmer, 1963–71 *Encyclopaedia of Chemical Technology*, (3rd edn., 1978–83) Kirk, R. E., Othmer, D. F., London, Wiley-Interscience, Vol. 1, 1963, Vol. 22, 1970, Suppl. Vol. 1971. (4th edn., from 1991) 1993;

- Kirshenbaum, 1956 *Final Report on Fundamental Studies of New Explosive Reactions*, Kirshenbaum, A. D., Philadelphia, Research Institute of Temple University, 1956
- Kit and Evered, 1960 *Rocket Propellant Handbook*, Kit, B., Evered, D. S., London, MacMillan, 1960
- Lab. Hazards Bull.* *Laboratory Hazards Bulletin*, Royal Society of Chemistry Information Group, Cambridge, 1981 to date (monthly abstracts)
- Lawless, 1968 *High Energy Oxidisers*, Lawless, E. W. and Smith, I. C., New York, Marcel Dekker, 1968
- Leleu, Cahiers, 1972–80 *Les Réactions Chimique Dangereuse*, Leleu, J., published serially in *Cahiers de Notes Documentaires*, from 1972 (**68**), 319 to 1980, (**100**), 426
- Loss Prev. Bull.*, 1974 to date *Loss Prevention Bulletin*, Rugby, Institution of Chemical Engineers, 6 issues annually since 1974
- Mackay, 1966 *Hydrogen Compounds of the Metallic Elements*, Mackay, K. M., London, Spon, 1966
- Major Loss Prevention*, 1971 *Major Loss Prevention in the Process Industries, Symposium Series No. 34*, London, Institution of Chemical Engineers, 1971
- Martin, 1971, *Dimethylsulfoxid*, Martin, D., Hauthal, H. G., Berlin, Akademie Verlag, 1971
- Matheson Guide, 1982 *Guide to Safe Handling of Compressed Gases*, Matheson Div. Searle Medical Products, Lyndhurst (NJ), 1982
- MCA Case Histories*, 1950–78 *Case Histories of Accidents in the Chemical Industry*, Washington, MCA, (published monthly, republished as 4 indexed collected volumes 1953–75, discontinued 1978)
- (*MCA SD-Series*) 1947–76 *Safety Data Sheets*, MCA, Washington. The whole series of 101 sheets was withdrawn in 1980, but remains a useful source of collected information on basic industrial chemicals.
- Meidl, 1972 *Hazardous Materials Handbook*, Meidl, J. H., Beverley Hills, Glencoe Press, 1972

- Mellor, 1947–71 *Comprehensive Treatise on Inorganic and Theoretical Chemistry*, Mellor, J. W., London, Longmans Green, Vol. 1, repr. 1941, Vol. 16, publ. 1937; and isolated Supplementary vols. up to Vol. 8 Suppl. 3, 1971
- Mellor MIC, 1961 *Mellor's Modern Inorganic Chemistry*, Parkes, G. D. (Revision Ed.), London, Longmans Green, 1961
- Merck, 1976, 1983, 1989, 1996 *The Merck Index*, Chapman and Hall, London., 9th edn. 1976, 10th edn. 1983, 11th edn. 1989, 12th edn. 1996
- Merrifield, 1988 *Fire and Explosion Hazards associated with Storage and Handling of Hydrogen Peroxide*, Merrifield, R., Specialist Inspectors Report No. 19, Bootle, HSE. 1988
- Meyer, 1977 *Chemistry of Hazardous Materials*, Meyer, E., Englewood Cliffs (NJ), Prentice-Hall, 1977
- Miller, 1965–66 *Acetylene, its Properties, Manufacture and Uses*, Miller, S. A., London, Academic Press, Vol. 1, 1965, Vol. 2, 1966
- Moissan, 1900 *Le Fluor et ses Composés*, Moissan, H., Paris, Steinheil, 1900
- NFPA 491M, 1975 *Manual of Hazardous Chemical Reactions*, NFPA 491M, Quincy (MA), National Fire Protection Association, 5th edn., 1975 (larger format of same text, 1985)
- Nolan, 1983 *Case Histories of Runaway Reactions*, Nolan, P. F., London, Polytechnic of the South Bank, 1983. A summary of the conclusions on the causes was published in *J. Haz. Materials*, 1987, **14**, 233–239
- Org. Synth.*, 1941 to date *Organic Syntheses*, various Eds., New York, Wiley, Collective Vol. 1, 1944; Coll. Vol. 2, 1944; Coll. Vol. 3, 1955; Coll. Vol. 4, 1962; Coll. Vol. 5, 1973; Coll. Vol. 6, 1987; Coll. Vol. 7, 1990; Coll. Vol. 8, 1993; annual volumes (from 70) thereafter.
- Partington, 1967 *General and Inorganic Chemistry*, Partington, J. R., London, MacMillan, 4th edn., 1967

- Pascal, 1956–70 *Nouveau Traité de Chimie Minérale*, Pascal, P. (Ed.), Paris, Masson, Vol. 1, 1956; Vol. 20.3, 1964; Vol. 15.5, 1970
- Pieters, 1957 *Safety in the Chemical Laboratory*, Pieters, H. A. J., Creighton, J. W., London, Academic Press, 2nd edn., 1957
- Plant/Oper. Progress* *Plant Operations Progress* continues as *Process Safety Progress* 1993-
- Pot. Incid. Rept.* *ASESB Potential Incident Report*, Washington, Armed Services Explosives Safety Board
- Proc. nth Combust. Symp.* *Proceedings of Combustion Symposia*, various Eds. and Publishers, dated year following the event. 1st Symposium 1928; 19th Symposium, 1982
- Renfrew, 1981 *Safety in the Chemical Laboratory*, Collected reprinted articles from *J. Chem. Educ.*, Renfrew, M. M. (Ed.), Vol. IV, 1981
- Rodd, 1951–62 *The Chemistry of Carbon Compounds*, Rodd, E. H. (Ed.), London, Elsevier, Vol. 1A, 1951, Vol. 5, 1962; 2nd edn. began in 1979
- RSC Data Sheets*, 1982 to date *Laboratory Hazard Data Sheet Series*, Cambridge, Royal Society of Chemistry, No. 1, 1982; No. 89, 1989, issued monthly, ceased 1993
- RSC Safety Data Sheets*, 1989 *Chemical Safety Data Sheets*, Cambridge, Royal Society of Chemistry, 1989 Generally similar to the monthly Laboratory Hazards Safety Data Sheet series, but the information is presented under headings reflecting legislative requirements. Volume 1 deals with some 100 largely organic liquid solvents and reagents, while Volume 2 covers main group metals and their compounds, 122 items in all. Vol. 3; Corrosives and Irritants, 1990. Vol 4; Toxic Chemicals, Pt. I 1990, Pt. II 1991. Vol 5, Flammables, 1989. There is considerable overlap among the volumes
- Runaway Reactions*, 1981 *Runaway Reactions, Unstable Products and Combustible Powders*, Symposium Series No. 68 (Chester 1981), Rugby, Institution of Chemical Engineers, 1981

- Rüst, 1948 *Unfälle beim Chemischen Arbeiten*, Rüst, E., Ebert, A., Zurich, Rascher Verlag, 2nd edn., 1948
- Rutledge, 1968 *Acetylenic Compounds*, Rutledge, T. F., New York, Reinhold, 1968
- Schmidt, 1967 *Handling and Use of Fluorine and Fluorine-Oxygen Mixtures in Rocket Systems*, SP-3037, Schmidt, H. W., Harper, J. T., Washington, NASA, 1967
- Schumacher, 1960 *Perchlorates, their Properties, Manufacture and uses*, ACS 146, Schumacher, J. C., New York, Reinhold, 1960
- Schumb, 1955 *Hydrogen Peroxide*, ACS 128, Schumb, W. C., Satterfield, C. N., Wentworth, R. L., New York, Reinhold, 1955
- Shriver, 1986 *Manipulation of Air-Sensitive Compounds*, Shriver, D. F. and Drozdov, M. A. (Eds.), New York, Wiley, 2nd edn., 1986
- Sichere Chemiarb.* *Sichere Chemiarbeit*, Heidelberg, Berufsgenossenschaft der Chemischen Industrie, 1963 to date
- Sidgwick, 1950 *The Chemical Elements and their Compounds*, Sidgwick, N. V., Oxford, Oxford University Press, 1950
- Sittig, 1981 *Handbook of Toxic and Hazardous Chemicals*, Sittig, M., Park Ridge (NJ), Noyes Data Corp., 2nd ed., 1985
- Smith, 1965–6 *Open-Chain Nitrogen Compounds*, Smith, P. A. S., New York, Benjamin, Vol. 1, 1965; Vol. 2, 1966
- Sorbe, 1968 *Giftige und Explosive Substanzen*, Sorbe, G., Frankfurt, Umschau Verlag, 1968
- Steere, 1971 *Handbook of Laboratory Safety*, Steere, N. V. (Ed.), Cleveland, Chemical Rubber Publishing Co., 2nd edn., 1971. 3rd edn., Furr, A. K. (Ed.), 1989
- Steere, 1967, '71, '74 *Safety in the Chemical Laboratory*, N. V. Steere (Ed.), Collected reprinted articles from *J. Chem. Educ.*, Vol. I, 1967; Vol. II, 1971; Vol. III, 1974

- Stull, 1977 *Fundamentals of Fire and Explosion*, AIChE Monograph Series No. 10, Stull, D. R., New York, American Institute of Chemical Engineers, 1977
- Swern, 1970-52 *Organic Peroxides*, Swern, D. (Ed.), London, Wiley-Interscience, Vol. 1, 1970; Vol. 2, 1971; Vol. 3, 1972
- Ullmann, 1998 *Ullmann's Encyclopedia of Industrial Chemistry*, Weinheim, 6th edn. 1998 (CD update 1999)
- Urbanski, 1964-84 *Chemistry and Technology of Explosives*, Urbanski, T., London, MacMillan, Vol. 1, 1964; Vol. 2, 1965; Vol. 3, 1967; Oxford, Pergamon, Vol. 4, 1984
- Van Wazer, 1958-61 *Phosphorus and its Compounds*, Van Wazer, J. R., New York, Interscience, Vol. 1, 1958; Vol. 2, 1961
- Vervalin, 1973 *Fire Protection Manual for Hydrocarbon Processing Plants*, Vervalin, C. H. (Ed.), Houston, Gulf Publishing Co., 2nd edn., 1973
- Vogel, 1957 *A Textbook of Practical Organic Chemistry*, Vogel, A. I., London, Longmans Green, 3rd edn., 1957
- von Schwartz, 1918 *Fire and Explosion Risks*, von Schwartz, E., London, Griffin, 1918
- Weast, 1988 *Handbook of Chemistry and Physics*, Weast, R. C. (Ed.), Boca Raton (Fa.), CRC Press, 1988; continued under the editorship of Lide, D. R.
- Whitmore, 1921 *Organic Compounds of Mercury*, Whitmore, F. C., New York, Chemical Catalog Co. Inc., 1921
- Yoshida, 1980 *Handbook of Hazardous Reactions with Chemicals*, Yoshida, T., Tokyo, Tokyo Fire Department, 1980 (copy in British Library, Boston Spa)
- Zabetakis, 1965 *Flammability Characteristics of Combustible Gases and Vapours*, Zabetakis, M. G., Washington, US Bureau of Mines, 1965

Tabulated Fire-related Data

Compounds which are considered to be unusually hazardous in a fire context because of their low flash points (below 25°C) or auto-ignition temperatures (below 225°C) are included in the table. The names used are those titles in the text of Section 1 which are prefixed with a dagger. Synonyms may be found either in Section 1 or in the alphabetical index of chemical names and synonyms in Appendix 4. Boiling points are given for those compounds boiling below 50°C.

The figures for flash points are closed-cup values except where a suffix (o) indicates the (usually higher) open-cup value. The figures for explosive limits (or flammability limits) are % by volume in air at ambient temperature except where indicated otherwise. Where no figure has been found for the upper limit, a query has been inserted. Figures for auto-ignition temperatures are usually those determined in glass (without catalytic effects) except where stated.

Most of the values are those quoted in the references given in the topic entries:

AUTOIGNITION TEMPERATURES

FIRE

FLAMMABILITY

FLASH POINTS

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Arsine	AsH ₃	-62	—	flammable	—
Diborane	B ₂ H ₆	-93	-90	0.9-98	38-52
Pentaborane(9)	B ₅ H ₉	—	30	0.4-?	35
Bromosilane	BrH ₃ Si	2	< 0	—	ambient
Tribromosilane	Br ₃ HSi	112	—	flammable	ambient
Hydrogen cyanide	CHN	25.7	-18	6.0-41	—
Dichloromethane	CH ₂ Cl ₂	39.7	calc -9	12.0-19.0	605
Formaldehyde	CH ₂ O	-19	-19	7.0-73	—
Bromomethane	CH ₃ Br	3.5	none	see text	—
Chloromethane	CH ₃ Cl	-24	< 0	8.1-17	—
Methyltrichlorosilane	CH ₃ ClSi	—	8	7.6-?	—
Fluoromethane	CH ₃ F	-78	—	flammable	—
Methyl nitrite	CH ₃ NO ₂	-12	—	flammable	explodes
Methane	CH ₄	-161	-187	5.3-15	—
Dichloromethylsilane	CH ₄ Cl ₂ Si	—	-32	—	ambient
Methanol	CH ₄ O	—	10	6-36.5/60°	—
Methanethiol	CH ₄ S	6	-18	3.9-21.8	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Methylamine	CH ₅ N	-6	-18	4.5-21	—
35% w/v in water		—	7.5	—	—
Methylhydrazine	CH ₆ N ₂	—	18	2.5-97	194
Carbon monoxide	CO	-192	—	12.5-74	—
Carbonyl sulfide	COS	-50	—	6.5-28.2	—
Carbon disulfide	CS ₂	46	-30	0.6-50	125 (below 100 if rust present)
Bromotrifluoroethylene	C ₂ BrF ₃	-3	—	—	ambient
Chlorotrifluoroethylene	C ₂ ClF ₃	-28	—	8.4-38.7	—
Tetrafluoroethylene	C ₂ F ₄	-76	—	11-60	180
Trichloroethylene	C ₂ HCl ₃	—	32 (above 30°C)	12.5-90	410
Trifluoroethylene	C ₂ HF ₃	—	—	15.3-27	—
Acetylene	C ₂ H ₂	-83	-18	2.5-82	—
1,1-Dichloroethylene	C ₂ H ₂ Cl ₂	32	-15(o)	7.3-16	—
<i>cis</i> -1,2-Dichloroethylene	C ₂ H ₂ Cl ₂	—	6	3.3-15	—
<i>trans</i> -1,2-Dichloroethylene	C ₂ H ₂ Cl ₂	4	2	9.7-12.8	—
1,1-Difluoroethylene	C ₂ H ₂ F ₂	-86	—	5.5-21.3	—
Ethanedial	C ₂ H ₂ O ₂	50	—	—	—
Bromoethylene	C ₂ H ₃ Br	16	< -8	6-16	—
Chloroethylene	C ₂ H ₃ Cl	-14	-8	4-22 (or 33)	—
1-Chloro-1,1-difluoroethane	C ₂ H ₃ ClF ₂	-9	—	9-14.8	—
Acetyl chloride	C ₂ H ₃ ClO	—	4	5.0-?	—
Methyl chloroformate	C ₂ H ₃ ClO ₂	—	12	—	—
1,1,1-Trichloroethane	C ₂ H ₃ Cl ₃	—	none	8-10.5	500
Trichlorovinylsilane	C ₂ H ₃ Cl ₃ Si	—	-9	—	—
Fluoroethylene	C ₂ H ₃ F	-73	—	2.6-22	—
1,1,1-Trifluoroethane	C ₂ H ₃ F ₃	-47	—	9.2-18.4	—
Acetonitrile	C ₂ H ₃ N	—	6(o)	4-16	—
Methyl isocyanate	C ₂ H ₃ NO	39	-7	—	—
Ethylene	C ₂ H ₄	-104	-136	3.0-34	—
1,1-Dichloroethane	C ₂ H ₄ Cl ₂	—	-12	5.6-11.4	—
1,2-Dichloroethane	C ₂ H ₄ Cl ₂	—	13	6.2-15.9	—
Bis(chloromethyl) ether	C ₂ H ₄ Cl ₂ O	—	< 19	—	—
1,1-Difluoroethane	C ₂ H ₄ F ₂	-25	—	3.7-18	—
Acetaldehyde	C ₂ H ₄ O	20	-38	4.0-57	204 (140 by DIN 51794)
Ethylene oxide	C ₂ H ₄ O	11	-20	3.0-100	—
Thiolacetic acid	C ₂ H ₄ OS	—	11	—	—
Methyl formate	C ₂ H ₄ O ₂	32	-32	5.9-20	—
Thiiran	C ₂ H ₄ S	—	10	—	—
Bromoethane	C ₂ H ₅ Br	38	-20	6.7-11.3	—
Chloroethane	C ₂ H ₅ Cl	12	-50	3.8-15.4	—
Chloromethyl methyl ether	C ₂ H ₅ ClO	—	-8	—	—
Trichloroethylsilane	C ₂ H ₅ Cl ₃ Si	—	14	—	—
Fluoroethane	C ₂ H ₅ F	-38	—	—	—
Aziridine	C ₂ H ₅ N	—	-11	3.3-54.8	—
Acetaldehyde oxime	C ₂ H ₅ NO	—	< 22	—	—
Ethyl nitrite	C ₂ H ₅ NO ₂	17	35	3.1-> 50	expl. > 90
Ethyl nitrate	C ₂ H ₅ NO ₃	—	10	3.8-?	expl. 85
Ethane	C ₂ H ₆	-89	-93	3-12.5	—
Dichlorodimethylsilane	C ₂ H ₆ Cl ₂ Si	-9	—	3.4-9.5	—
Dichloroethylsilane	C ₂ H ₆ Cl ₂ Si	—	-1	—	—
Dimethyl ether	C ₂ H ₆ O	-24	-41	3.4-18 (27)	—
Ethanol	C ₂ H ₆ O	—	12	3.3-19/60°C	—
Dimethyl sulfoxide	C ₂ H ₆ OS	—	—	—	215
Dimethyl sulfate	C ₂ H ₆ O ₄ S	—	—	—	188
Dimethyl sulfide	C ₂ H ₆ S	37	-37	2.2-19.7	206
Ethanethiol	C ₂ H ₆ S	36	-17	2.8-18	—
Dimethyl disulfide	C ₂ H ₆ S ₂	—	7	1.1-?	—
Dimethylamine	C ₂ H ₇ N	7	-50	2.8-14.4	—
40% w/v in water		—	-16	—	—
25% w/v in water		—	6	—	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Ethylamine	C ₂ H ₇ N	17	-16	3.5-14	—
70% w/v in water	—	—	< 22	—	—
1,1-Dimethylhydrazine	C ₂ H ₈ N ₂	—	1	2.1-95	—
1,2-Dimethylhydrazine	C ₂ H ₈ N ₂	—	-17	—	—
1,1-Dimethyldiborane	C ₂ H ₁₀ B ₂	-3	< -10	—	—
1,2-Dimethyldiborane	C ₂ H ₁₀ B ₂	-49	< -55	—	—
Dicyanogen	C ₂ N ₂	-21	—	6-32	—
Hexafluoropropene	C ₃ F ₆	—	—	15-20	—
2-Chloroacrylonitrile	C ₃ H ₂ ClN	—	8	—	—
3-Bromopropyne	C ₃ H ₃ Br	—	10	3.0-?	—
3-Chloropropyne	C ₃ H ₃ Cl	—	-16	—	—
Acryloyl chloride	C ₃ H ₃ ClO	—	18	—	—
3,3,3-Trifluoropropene	C ₃ H ₃ F ₃	-18	< -30	4.7-23.5	—
1,1,1-Trifluoroacetone	C ₃ H ₃ F ₃ O	22	-30	—	—
Acrylonitrile	C ₃ H ₃ Ne	—	-1	3.0-17	—
5% w/v in water	—	—	< 9	—	—
Isoxazole	C ₃ H ₃ NO	—	8	—	—
Oxazole	C ₃ H ₃ NO	—	18	—	—
Propadiene	C ₃ H ₄	-34	—	1.7-12	—
Propyne	C ₃ H ₄	-23	—	2.15-12.5	—
1-Chloro-3,3,3-trifluoropropane	C ₃ H ₄ ClF ₃	—	—	6-14	—
2,3-Dichloropropene	C ₃ H ₄ Cl ₂	—	10	—	—
Acrylaldehyde	C ₃ H ₄ O	—	-26	2.8-31	—
Methoxyacetylene	C ₃ H ₄ O	23	< -20	—	—
Vinyl formate	C ₃ H ₄ O ₂	46	< 0	—	—
3-Bromo-1-propene	C ₃ H ₅ Br	—	-1	4.4-7.3	—
1-Chloro-1-propene	C ₃ H ₅ Cl	35	< -6	4.5-16	—
2-Chloropropene	C ₃ H ₅ Cl	22	-37	4.5-16	—
3-Chloropropene	C ₃ H ₅ Cl	45	-32	3.3-11.2	—
1-Chloro-2,3-epoxypropane	C ₃ H ₅ ClO	—	21	—	—
Propionyl chloride	C ₃ H ₅ ClO	—	12	—	—
Ethyl chloroformate	C ₃ H ₅ ClO ₂	—	16	—	—
1-Fluoro-2,3-epoxypropane	C ₃ H ₅ FO	—	4	—	—
3-Iodopropene	C ₃ H ₅ I	—	5	—	—
Propionitrile	C ₃ H ₅ N	—	6	3.1-?	—
Cyclopropane	C ₃ H ₆	-33	—	2.4-10.4	—
Propene	C ₃ H ₆	-48	-108	2.4-11.1	—
				(see text)	
1,1-Dichloropropane	C ₃ H ₆ Cl ₂	—	21	3.1-?	—
1,2-Dichloropropane	C ₃ H ₆ Cl ₂	—	15	3.4-14.5	—
Dichloromethylvinylsilane	C ₃ H ₆ Cl ₂ Si	—	-1	—	—
Acetone	C ₃ H ₆ O	—	-17	2.6-12.8	—
Methyl vinyl ether	C ₃ H ₆ O	5.5	-56(o)	2.6-39.0	—
Oxetane	C ₃ H ₆ O	50	-28	—	—
2-Propen-1-ol	C ₃ H ₆ O	—	21	3.0-18.0	—
Propionaldehyde	C ₃ H ₆ O	49	-9	2.9-17	207
Propylene oxide	C ₃ H ₆ O	34	-37	3.1-27.5	—
1,3-Dioxolane	C ₃ H ₆ O ₂	—	2	—	—
Ethyl formate	C ₃ H ₆ O ₂	—	-34	2.7-13.5	—
Methyl acetate	C ₃ H ₆ O ₂	—	-9	3.1-16	—
Dimethyl carbonate	C ₃ H ₆ O ₃	—	18	—	—
2-Propene-1-thiol	C ₃ H ₆ S	—	-10	—	—
1-Bromopropane	C ₃ H ₇ Br	—	-1	4.6-?	—
2-Bromopropane	C ₃ H ₇ Br	—	1	—	—
1-Chloropropane	C ₃ H ₇ Cl	47	< -18	2.6-11.1	—
2-Chloropropane	C ₃ H ₇ Cl	36	-32	2.8-10.7	—
Chloromethyl ethyl ether	C ₃ H ₇ ClO	—	15	—	—
2-Iodopropane	C ₃ H ₇ I	—	20	—	—
3-Aminopropene	C ₃ H ₇ N	—	-29	2.2-22	—
Azetidine	C ₃ H ₇ N	61	-20	—	—
Cyclopropylamine	C ₃ H ₇ N	—	1	—	—
2-Methylaziridine	C ₃ H ₇ N	—	-10	—	—
Isopropyl nitrite	C ₃ H ₇ NO ₂	40	< 10	—	—
Propyl nitrite	C ₃ H ₇ NO ₂	47	< 20	—	—
Isopropyl nitrate	C ₃ H ₇ NO ₃	—	11	?-100	175

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Propyl nitrate	C ₃ H ₇ NO ₃	—	20	2–100	—
Propane	C ₃ H ₈	7–45	–104	2.2–9.5	—
Ethyl methyl ether	C ₃ H ₈ O	11	–37	2–10.1	190
Propanol	C ₃ H ₈ O	—	15	2.5–13.5	—
2-Propanol	C ₃ H ₈ O	—	12	2.3–12.7	—
Dimethoxymethane	C ₃ H ₈ O ₂	44	–18(o)	—	—
Ethyl methyl sulfide	C ₃ H ₈ S	—	–15	—	—
Propanethiol	C ₃ H ₈ S	—	–20	—	—
2-Propanethiol	C ₃ H ₈ S	—	–35	—	—
Trimethyl borate	C ₃ H ₉ BO ₃	—	–8	—	—
Chlorotrimethylsilane	C ₃ H ₉ ClSi	—	–20	—	—
Iodotrimethylsilane	C ₃ H ₉ ISi	106	–31	—	—
Isopropylamine	C ₃ H ₉ N	32	–50	2.3–10.4	—
Propylamine	C ₃ H ₉ N	49	–37	2.0–10.4	—
Trimethylamine	C ₃ H ₉ N	3	–5	2.0–11.6	190
25% w/v in water	—	—	5	—	—
2-Methoxyethylamine	C ₃ H ₉ NO	—	9	—	—
Trimethylsilyl azide	C ₃ H ₉ N ₃ Si	—	23	—	—
1,2-Diaminopropane	C ₃ H ₁₀ N ₂	—	24	—	—
1,3-Diaminopropane	C ₃ H ₁₀ N ₂	—	24(o)	—	—
Propadiene-1,3-dione	C ₃ O ₂	7	—	6–30	—
1,3-Butadiyne	C ₄ H ₂	10	—	1.4–100	—
Buten-3-yne	C ₄ H ₄	5	< –5	1.8–40(-93% in oxygen)	—
Furan	C ₄ H ₄ O	32	–36	2.3–14.3	—
Diketene	C ₄ H ₄ O ₂	—	33	2–11.7	200
Methyl propiolate	C ₄ H ₄ O ₂	—	10	—	—
Thiophene	C ₄ H ₄ S	—	–6	—	—
2-Chloro-1,3-butadiene	C ₄ H ₅ Cl	—	–20	2.5–20	—
Ethyl trifluoroacetate	C ₄ H ₅ F ₃ O ₂	—	–17	—	—
1-Cyanopropene	C ₄ H ₅ N	—	16	—	—
3-Cyanopropene	C ₄ H ₅ N	—	19	—	—
1,2-Butadiene	C ₄ H ₆	19	< 0	—	—
1,3-Butadiene	C ₄ H ₆	–4	< –17	2–11.5	—
1-Butyne	C ₄ H ₆	8	< –7	—	—
2-Butyne	C ₄ H ₆	28	–31	1.4–?	—
Cyclobutene	C ₄ H ₆	2	< –10	—	—
1-Buten-3-one	C ₄ H ₆ O	—	–7	2.1–15.6	—
Crotonaldehyde	C ₄ H ₆ O	—	13	2.1–15.5/60°	207
2,3-Dihydrofuran	C ₄ H ₆ O	54	–24	—	—
Divinyl ether	C ₄ H ₆ O	29	< –30	1.7–27	—
3,4-Epoxybutene	C ₄ H ₆ O	—	< –50	—	—
Ethoxyacetylene	C ₄ H ₆ O	50	< –7	—	—
Methacrylaldehyde	C ₄ H ₆ O	—	–15	—	—
Allyl formate	C ₄ H ₆ O ₂	—	< 22	—	—
Butane-2,3-dione	C ₄ H ₆ O ₂	—	7	—	—
1,2:3,4-Diepoxybutane	C ₄ H ₆ O ₂	—	–10	—	—
Methyl acrylate	C ₄ H ₆ O ₂	—	–3(o)	2.8–25	—
Vinyl acetate	C ₄ H ₆ O ₂	—	–8	2.6–13.4	—
1-Bromo-2-butene	C ₄ H ₇ Br	—	13	—	—
4-Bromo-1-butene	C ₄ H ₇ Br	—	1	—	—
2-Chloro-2-butene	C ₄ H ₇ Cl	—	–25	2.3–9.3	—
3-Chloro-1-butene	C ₄ H ₇ Cl	—	–27	2.2–?	—
3-Chloro-2-methyl-1-propene	C ₄ H ₇ Cl	—	–19	2.3–9.3	—
Butyryl chloride	C ₄ H ₇ ClO	—	18	—	—
2-Chloroethyl vinyl ether	C ₄ H ₇ ClO	—	25	—	—
Isobutyryl chloride	C ₄ H ₇ ClO	—	8	—	—
Isopropyl chloroformate	C ₄ H ₇ ClO ₂	—	11	—	—
Butyronitrile	C ₄ H ₇ N	—	21	—	—
Isobutyronitrile	C ₄ H ₇ N	—	8	—	—
2-Methyl-2-oxazoline	C ₄ H ₇ NO	—	20	—	—
1-Butene	C ₄ H ₈	–6	–80	1.6–9.3	—
cis-2-Butene	C ₄ H ₈	4	–12	1.7–9.7	—
trans-2-Butene	C ₄ H ₈	3	< –6	1.8–9.7	—
Cyclobutane	C ₄ H ₈	13	< 10	1.8–?	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Methylcyclopropane	C ₄ H ₈	4	< 0	—	—
2-Methylpropene	C ₄ H ₈	-7	< -10	1.8-8.8	—
<i>mixo</i> -Dichlorobutane	C ₄ H ₈ Cl ₂	—	21	—	—
2-Butanone	C ₄ H ₈ O	—	-1	1.8-11.5	—
Butyraldehyde	C ₄ H ₈ O	—	-6	2.5-?	230
Cyclopropyl methyl ether	C ₄ H ₈ O	—	< 10	—	—
1,2-Epoxybutane	C ₄ H ₈ O	44.7	-15	1.5-18.3	—
Ethyl vinyl ether	C ₄ H ₈ O	36	-46	1.7-29	202
Isobutyraldehyde	C ₄ H ₈ O	—	-25	1.6-10.6	223
Tetrahydrofuran	C ₄ H ₈ O	—	-17	1.8-11.8	224
1,3-Dioxane	C ₄ H ₈ O ₂	—	2(o)	2-22	—
1,4-Dioxane	C ₄ H ₈ O ₂	—	12	2-22	180
Ethyl acetate	C ₄ H ₈ O ₂	—	-4	2.2-11.5	—
Isopropyl formate	C ₄ H ₈ O ₂	—	-6	—	—
Methyl propionate	C ₄ H ₈ O ₂	—	-2	2.5-13	—
Propyl formate	C ₄ H ₈ O ₂	—	-3	2.3-?	—
Tetrahydrothiophene	C ₄ H ₈ S	—	13	—	—
1-Bromobutane	C ₄ H ₉ Br	—	18	2.8-6.6/100	—
2-Bromobutane	C ₄ H ₉ Br	—	21	—	—
1-Bromo-2-methylpropane	C ₄ H ₉ Br	—	22	—	—
2-Bromo-2-methylpropane	C ₄ H ₉ Br	—	-18	—	—
2-Bromoethyl ethyl ether	C ₄ H ₉ BrO	—	21	—	—
1-Chlorobutane	C ₄ H ₉ Cl	—	-12	1.9-10.1	—
2-Chlorobutane	C ₄ H ₉ Cl	—	-10	—	—
1-Chloro-2-methylpropane	C ₄ H ₉ Cl	—	-6	2.0-8.7	—
2-Chloro-2-methylpropane	C ₄ H ₉ Cl	—	0	—	—
2-Iodobutane	C ₄ H ₉ I	—	-10	—	—
1-Iodo-2-methylpropane	C ₄ H ₉ I	—	0	—	—
2-Iodo-2-methylpropane	C ₄ H ₉ I	—	-10	—	—
Pyrrolidine	C ₄ H ₉ N	—	3	—	—
Butyl nitrite	C ₄ H ₉ NO ₂	—	10	—	—
<i>tert</i> -Butyl nitrite	C ₄ H ₉ NO ₂	—	-13	—	—
Cyanotrimethylsilane	C ₄ H ₉ NSi	114	1	—	—
Butane	C ₄ H ₁₀	1	-60	1.9-8.5	—
Isobutane	C ₄ H ₁₀	-12	-81	1.9-8.5	—
Dichlorodiethylsilane	C ₄ H ₁₀ Cl ₂ Si	—	24	—	—
2-Butanol	C ₄ H ₁₀ O	—	14	(1.7-9.8/100°)	—
<i>tert</i> -Butanol	C ₄ H ₁₀ O	—	10	2.4-8	—
Diethyl ether	C ₄ H ₁₀ O	36	-45	1.8-36.5	180
Methyl propyl ether	C ₄ H ₁₀ O	39	-20	2.0-14.8	—
1,1-Dimethoxyethane	C ₄ H ₁₀ O ₂	—	1(o)	—	—
1,2-Dimethoxyethane	C ₄ H ₁₀ O ₂	—	1	—	—
Trimethyl orthoformate	C ₄ H ₁₀ O ₃	—	15	—	—
1-Butanethiol	C ₄ H ₁₀ S	—	2	—	—
2-Butanethiol	C ₄ H ₁₀ S	—	-23	—	—
Diethyl sulfide	C ₄ H ₁₀ S	—	-10	—	—
2-Methylpropanethiol	C ₄ H ₁₀ S	—	-10	—	—
2-Methyl-2-propanethiol	C ₄ H ₁₀ S	< -29	—	—	—
Butylamine	C ₄ H ₁₁ N	—	-12	1.7-9.8	—
2-Butylamine	C ₄ H ₁₁ N	—	-9	—	—
<i>tert</i> -Butylamine	C ₄ H ₁₁ N	45	-7	(1.7-9.8/100°)	—
Diethylamine	C ₄ H ₁₁ N	—	-39	1.8-10.1	—
Ethyl dimethylamine	C ₄ H ₁₁ N	—	-36	—	—
Isobutylamine	C ₄ H ₁₁ N	—	-9	—	—
2-Dimethylaminoethylamine	C ₄ H ₁₂ N ₂	—	11	—	—
Tetramethyl orthosilicate	C ₄ H ₁₂ O ₄ Si	—	20	—	—
Tetramethylsilane	C ₄ H ₁₂ Si	26	< 0	—	—
Tetramethyltin	C ₄ H ₁₂ Sn	—	-12	1.9-?	—
Dicyanoacetylene	C ₄ N ₂	76	—	—	130
Tetracarbonylnickel	C ₄ NiO ₄	43	< -20	2.0-?	—
Pentacarbonyliron	C ₅ FeO ₅	—	-15	—	ambient
Pyridine	C ₅ H ₅ N	—	20	1.8-12.4	—
Cyclopentadiene	C ₅ H ₆	42	-3	—	—
2-Methyl-1-buten-3-yne	C ₅ H ₆	—	< -7(o)	—	—
2-Methylfuran	C ₅ H ₆ O	—	-30	—	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
2-Methylthiophene	C ₅ H ₆ S	—	8	—	—
1-Methylpyrrole	C ₅ H ₇ N	—	16	—	—
Cyclopentene	C ₅ H ₈	44	-29	—	—
3-Methyl-1,2-butadiene	C ₅ H ₈	40	-12	—	—
2-Methyl-1,3-butadiene	C ₅ H ₈	34	-53	2.0-8.9	220
3-Methyl-1-butyne	C ₅ H ₈	29.3	—	—	—
1,2-Pentadiene	C ₅ H ₈	44.9	—	—	—
1,3-Pentadiene	C ₅ H ₈	42	-43	2-8.3	—
1,4-Pentadiene	C ₅ H ₈	26	4	—	—
1-Pentyne	C ₅ H ₈	—	-20	—	—
2-Pentyne	C ₅ H ₈	56	—	—	—
Allyl vinyl ether	C ₅ H ₈ O	—	< 21(o)	—	—
Cyclopentene oxide	C ₅ H ₈ O	—	10	—	—
Cyclopropyl methyl ketone	C ₅ H ₈ O	—	13	—	—
2,3-Dihydropyran	C ₅ H ₈ O	—	-16	—	—
2-Methyl-3-butyne-2-ol	C ₅ H ₈ O	—	20	—	—
Methyl isopropenyl ketone	C ₅ H ₈ O	—	21	(1.8-9.0/50°)	—
Allyl acetate	C ₅ H ₈ O ₂	—	22	—	—
Ethyl acrylate	C ₅ H ₈ O ₂	—	9	1.8-?	—
Methyl crotonate	C ₅ H ₈ O ₂	—	-1	—	—
Methyl cyclopropanecarboxylate	C ₅ H ₈ O ₂	—	17	—	—
Methyl methacrylate	C ₅ H ₈ O ₂	—	8	2.1-12.5	—
Isopropenyl acetate	C ₅ H ₈ O ₂	—	16	1.9-?	—
Vinyl propionate	C ₅ H ₈ O ₂	—	1	—	—
Chlorocyclopentane	C ₅ H ₉ Cl	—	16	—	—
Pivaloyl chloride	C ₅ H ₉ ClO	—	8	—	—
Pivalonitrile	C ₅ H ₉ N	—	21	—	—
1,2,3,6-Tetrahydropyridine	C ₅ H ₉ N	—	16	—	—
Butyl isocyanate	C ₅ H ₉ NO	—	11	—	—
Cyclopentane	C ₅ H ₁₀	49	-25	—	—
1,1-Dimethylcyclopropane	C ₅ H ₁₀	20	—	—	—
Ethylcyclopropane	C ₅ H ₁₀	36	< 10	—	—
2-Methyl-1-butene	C ₅ H ₁₀	39	-20	—	—
2-Methyl-2-butene	C ₅ H ₁₀	38	-20	—	—
3-Methyl-1-butene	C ₅ H ₁₀	20	-7	1.5-9.1	—
Methylcyclobutane	C ₅ H ₁₀	41	< 10	—	—
1-Pentene	C ₅ H ₁₀	30	-18	1.5-8.7	—
2-Pentene	C ₅ H ₁₀	30	-18	—	—
Allyl ethyl ether	C ₅ H ₁₀ O	—	-20	—	—
Ethyl propenyl ether	C ₅ H ₁₀ O	—	< -5	—	—
Isopropyl vinyl ether	C ₅ H ₁₀ O	—	-32	—	—
Isovaleraldehyde	C ₅ H ₁₀ O	—	-5	—	—
3-Methyl-2-butanone	C ₅ H ₁₀ O	—	-3	—	—
2-Methyl-3-buten-2-ol	C ₅ H ₁₀ O	—	18	—	—
2-Methyltetrahydrofuran	C ₅ H ₁₀ O	—	-12	—	—
2-Pentanone	C ₅ H ₁₀ O	—	7	1.6-7.2	—
3-Pentanone	C ₅ H ₁₀ O	—	13	1.6-?	—
4-Penten-2-ol	C ₅ H ₁₀ O	—	25	—	—
Tetrahydropyran	C ₅ H ₁₀ O	—	-20	—	—
Valeraldehyde	C ₅ H ₁₀ O	—	12	—	—
Butyl formate	C ₅ H ₁₀ O ₂	—	18	1.7-8	—
3,3-Dimethoxypropene	C ₅ H ₁₀ O ₂	—	19	—	—
2,2-Dimethyl-1,3-dioxolane	C ₅ H ₁₀ O ₂	—	-1	—	—
Ethyl propionate	C ₅ H ₁₀ O ₂	—	12	1.9-11	—
Isobutyl formate	C ₅ H ₁₀ O ₂	—	5	2.0-8.9	—
Isopropyl acetate	C ₅ H ₁₀ O ₂	—	4	1.7-7.8	—
2-Methoxyethyl vinyl ether	C ₅ H ₁₀ O ₂	—	18(o)	—	—
Methyl butyrate	C ₅ H ₁₀ O ₂	—	14	—	—
4-Methyl-1,3-dioxane	C ₅ H ₁₀ O ₂	—	16	—	—
Methyl isobutyrate	C ₅ H ₁₀ O ₂	—	13(o)	—	—
Propyl acetate	C ₅ H ₁₀ O ₂	—	14	1.7-8	—
Diethyl carbonate	C ₅ H ₁₀ O ₃	—	25	—	—
1-Bromo-3-methylbutane	C ₅ H ₁₁ Br	—	21	—	—
2-Bromopentane	C ₅ H ₁₁ Br	—	20	—	—
1-Chloro-3-methylbutane	C ₅ H ₁₁ Cl	—	< 21	1.5-7.4	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
2-Chloro-2-methylbutane	C ₅ H ₁₁ Cl	—	3	1.5–7.4	—
1-Chloropentane	C ₅ H ₁₁ Cl	—	1	1.6–8.6	—
2-Iodopentane	C ₅ H ₁₁ I	—	< 23	—	—
Cyclopentylamine	C ₅ H ₁₁ N	—	13	—	—
1-Methylpyrrolidine	C ₅ H ₁₁ N	—	3	—	—
Piperidine	C ₅ H ₁₁ N	—	3	—	—
4-Methylmorpholine	C ₅ H ₁₁ NO	—	24	—	—
Isopentyl nitrite	C ₅ H ₁₁ NO ₂	—	–18	1–?(calc.)	209
Pentyl nitrite	C ₅ H ₁₁ NO ₂	—	< 23	—	—
2,2-Dimethylpropane	C ₅ H ₁₂	9	< –7	1.4–7.5	—
2-Methylbutane	C ₅ H ₁₂	28	–51	1.4–7.6	—
Pentane	C ₅ H ₁₂	36	–40	1.4–8.0	—
Butyl methyl ether	C ₅ H ₁₂ O	—	–10	—	—
<i>tert</i> -Butyl methyl ether	C ₅ H ₁₂ O	55	–28	2.5–15.1	—
Ethyl isopropyl ether	C ₅ H ₁₂ O	—	< –15	—	—
Ethyl propyl ether	C ₅ H ₁₂ O	—	< –20	1.7–9	—
<i>tert</i> -Pentanol	C ₅ H ₁₂ O	—	19	1.2–9	—
Diethoxymethane	C ₅ H ₁₂ O ₂	—	(55)	—	—
1,1-Dimethoxypropane	C ₅ H ₁₂ O ₂	—	< 10	—	—
2,2-Dimethoxypropane	C ₅ H ₁₂ O ₂	—	–7	—	—
Tetramethyl orthocarbonate	C ₅ H ₁₂ O ₄	—	6	—	—
2-Methyl-2-butanethiol	C ₅ H ₁₂ S	—	–1	—	—
3-Methylbutanethiol	C ₅ H ₁₂ S	—	18	—	—
Pentanethiol	C ₅ H ₁₂ S	—	18	—	—
1,1-Dimethylpropylamine	C ₅ H ₁₃ N	—	–1	—	—
1,2-Dimethylpropylamine	C ₅ H ₁₃ N	—	–20	—	—
2,2-Dimethylpropylamine	C ₅ H ₁₃ N	—	25	—	—
<i>N,N</i> -Dimethylpropylamine	C ₅ H ₁₃ N	—	–11	—	—
Isopentylamine	C ₅ H ₁₃ N	—	–1	2.3–22	—
<i>N</i> -Methylbutylamine	C ₅ H ₁₃ N	—	2	—	—
Pentylamine	C ₅ H ₁₃ N	—	–1	—	—
2-Dimethylamino- <i>N</i> -methyl-ethylamine	C ₅ H ₁₄ N ₂	—	14	—	—
Dimethylaminotrimethylsilane	C ₅ H ₁₅ NSi	—	–19	—	—
Hexafluorobenzene	C ₆ F ₆	—	10	—	—
1,2,4,5-Tetrafluorobenzene	C ₆ H ₂ F ₄	—	4(o)	—	—
1,2,4-Trifluorobenzene	C ₆ H ₃ F ₃	—	–5(o)	—	—
2-, 3-, or 4-Chlorofluorobenzene	C ₆ H ₄ ClF	—	all 18	—	—
1,3-Difluorobenzene	C ₆ H ₄ F ₂	—	< 0	—	—
1,4-Difluorobenzene	C ₆ H ₄ F ₂	—	–5(o)	—	—
Fluorobenzene	C ₆ H ₅ F	—	–15	—	—
Benzene	C ₆ H ₆	—	–11	1.4–8.0	—
1,5-Hexadien-3-yne	C ₆ H ₆	—	< –20	1.5–?	—
1,3-Cyclohexadiene	C ₆ H ₈	—	–6	—	—
1,4-Cyclohexadiene	C ₆ H ₈	—	–11	—	—
2,5-Dimethylfuran	C ₆ H ₈ O	—	16	—	—
Cyclohexene	C ₆ H ₁₀	—	–12	1.2–?/100°	310
1,3-Hexadiene	C ₆ H ₁₀	—	–3	—	—
1,4-Hexadiene	C ₆ H ₁₀	—	–21	2.0–6.1	—
1,5-Hexadiene	C ₆ H ₁₀	—	–46	—	—
<i>c</i> - or <i>t</i> -2- <i>trans</i> -4-Hexadiene	C ₆ H ₁₀	—	–7	—	—
1-, 2- or 3-Hexyne	C ₆ H ₁₀	—	–10	—	—
2-Methyl-1,3-pentadiene	C ₆ H ₁₀	—	–12	—	—
4-Methyl-1,3-pentadiene	C ₆ H ₁₀	—	–34	—	—
Diallyl ether	C ₆ H ₁₀ O	—	–6	—	—
3-Methyl-2-methylenebutanal	C ₆ H ₁₀ O	—	11	—	—
Ethyl crotonate	C ₆ H ₁₀ O ₂	—	2	—	—
Ethyl cyclopropanecarboxylate	C ₆ H ₁₀ O ₂	—	18	—	—
Ethyl methacrylate	C ₆ H ₁₀ O ₂	—	21	1.8–?	—
Vinyl butyrate	C ₆ H ₁₀ O ₂	—	21(o)	1.4–8.8	—
Diallylamine	C ₆ H ₁₁ N	—	15	—	—
Cyclohexane	C ₆ H ₁₂	—	–20	1.3–8.4	—
Ethylcyclobutane	C ₆ H ₁₂	—	–15	1.2–7.7	210
1-Hexene	C ₆ H ₁₂	—	–26	1.2–6.9	—
2-Hexene	C ₆ H ₁₂	—	–21	—	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Methylcyclopentane	C ₆ H ₁₂	—	-29	—	—
2-Methyl-1-pentene	C ₆ H ₁₂	—	-28	—	—
4-Methyl-1-pentene	C ₆ H ₁₂	—	-7	—	—
<i>cis</i> 8-4-Methyl-2-pentene	C ₆ H ₁₂	—	-32	—	—
<i>trans</i> 8-4-Methyl-2-pentene	C ₆ H ₁₂	—	-29	—	—
Butyl vinyl ether	C ₆ H ₁₂ O	—	-1	—	—
3,3-Dimethyl-2-butanone	C ₆ H ₁₂ O	—	12	—	—
2-Ethylbutyraldehyde	C ₆ H ₁₂ O	—	21	1.2-7.7	—
2-Hexanone	C ₆ H ₁₂ O	—	23	1.2-8.0	—
3-Hexanone	C ₆ H ₁₂ O	—	14	—	—
Isobutyl vinyl ether	C ₆ H ₁₂ O	—	-9	—	—
2-Methylpentanal	C ₆ H ₁₂ O	—	20	—	195
3-Methylpentanal	C ₆ H ₁₂ O	—	< 23	—	—
2-Methyl-3-pentanone	C ₆ H ₁₂ O	—	1	—	—
3-Methyl-2-pentanone	C ₆ H ₁₂ O	—	15	—	—
4-Methyl-2-pentanone	C ₆ H ₁₂ O	—	17	1.2-8	—
Butyl acetate	C ₆ H ₁₂ O ₂	—	23	1.4-7.5	—
2-Butyl acetate	C ₆ H ₁₂ O ₂	—	18	1.3-7.5	—
2,6-Dimethyl-1,4-dioxane	C ₆ H ₁₂ O ₂	—	24	—	—
Ethyl isobutyrate	C ₆ H ₁₂ O ₂	—	13	—	—
2-Ethyl-2-methyl-1,3-dioxolane	C ₆ H ₁₂ O ₂	—	23(o)	—	—
4-Hydroxy-4-methyl-2-pentanone	C ₆ H ₁₂ O ₂	—	9	—	—
Isobutyl acetate	C ₆ H ₁₂ O ₂	—	17	2.4-10.5	—
Isopentyl formate	C ₆ H ₁₂ O ₂	—	23	1.2-8.0	—
Isopropyl propionate	C ₆ H ₁₂ O ₂	—	21	—	—
Methyl isovalerate	C ₆ H ₁₂ O ₂	—	20	—	—
Methyl pivalate	C ₆ H ₁₂ O ₂	—	7	—	—
Methyl valerate	C ₆ H ₁₂ O ₂	—	< 25	—	—
Isobutyl peroxyacetate	C ₆ H ₁₂ O ₃	—	< 25	—	—
2,4,6-Trimethyltrioxane	C ₆ H ₁₂ O ₃	—	17	1.3-?	—
Cyclohexylamine	C ₆ H ₁₃ N	—	21	—	—
1-Methylpiperidine	C ₆ H ₁₃ N	—	3	—	—
2-Methylpiperidine	C ₆ H ₁₃ N	—	10	—	—
3-Methylpiperidine	C ₆ H ₁₃ N	—	8	—	—
4-Methylpiperidine	C ₆ H ₁₃ N	—	9	—	—
Perhydroazepine	C ₆ H ₁₃ N	—	22	—	—
2,2-Dimethylbutane	C ₆ H ₁₄	50	-48	(1.2-7.0/100°)	—
2,3-Dimethylbutane	C ₆ H ₁₄	—	-29	1.2-7.0	—
Hexane	C ₆ H ₁₄	—	-23	1.1-7.5	225
Isohexane	C ₆ H ₁₄	—	-7	1.2-7.0	—
3-Methylpentane	C ₆ H ₁₄	—	-7	1.2-7.0	—
Butyl ethyl ether	C ₆ H ₁₄ O	—	-1	—	—
<i>tert</i> -Butyl ethyl ether	C ₆ H ₁₄ O	—	-19	—	—
Diisopropyl ether	C ₆ H ₁₄ O	—	-36	1.4-21	—
Dipropyl ether	C ₆ H ₁₄ O	—	-21	—	215
2-Methyl-2-pentanol	C ₆ H ₁₄ O	—	21	—	—
1,1-Diethoxyethane	C ₆ H ₁₄ O ₂	—	-21	1.7-10.4	—
1,2-Diethoxyethane	C ₆ H ₁₄ O ₂	—	—	—	205
Triethylaluminium	C ₆ H ₁₅ Al	—	< -53	—	< -53
Triethyl borate	C ₆ H ₁₅ BO ₃	—	11	—	—
Butylethylamine	C ₆ H ₁₅ N	—	18(o)	—	—
Diisopropylamine	C ₆ H ₁₅ N	—	-7	—	—
1,3-Dimethylbutylamine	C ₆ H ₁₅ N	—	13	—	—
Dipropylamine	C ₆ H ₁₅ N	—	17	—	—
Triethylamine	C ₆ H ₁₅ N	—	-7	1.2-8.0	—
1,2-Bis(dimethylamino)ethane	C ₆ H ₁₆ N ₂	—	18	—	—
Diethoxydimethylsilane	C ₆ H ₁₆ O ₂ Si	—	11	—	—
Bis(trimethylsilyl) oxide	C ₆ H ₁₈ OSi ₂	—	-1	—	—
Hexamethyldisilazane	C ₆ H ₁₉ NSi ₂	—	14	—	—
Benzaldehyde	C ₇ H ₆ O	—	—	—	190
2-Fluorotoluene	C ₇ H ₇ F	—	8	—	—
3-Fluorotoluene	C ₇ H ₇ F	—	12	—	—
4-Fluorotoluene	C ₇ H ₇ F	—	10	—	—
Bicyclo[2.2.1]-2,5-heptadiene	C ₇ H ₈	—	-21	—	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
1,3,5-Cycloheptatriene	C ₇ H ₈	—	4	—	—
Toluene	C ₇ H ₈	—	4	1.3–7.0	—
Cycloheptene	C ₇ H ₁₂	—	–6	—	—
1-Heptyne	C ₇ H ₁₂	—	–2	—	—
4-Methylcyclohexene	C ₇ H ₁₂	—	–1(o)	—	—
Cycloheptane	C ₇ H ₁₄	—	6	—	—
Ethylcyclopentane	C ₇ H ₁₄	—	15	1.1–6.7	—
1-Heptene	C ₇ H ₁₄	—	–1	—	—
2-Heptene	C ₇ H ₁₄	—	< –1	—	—
3-Heptene	C ₇ H ₁₄	—	< –7	—	—
Methylcyclohexane	C ₇ H ₁₄	—	–4	1.2–6.7	—
2,3,3-Trimethylbutene	C ₇ H ₁₄	—	< 0	—	—
2,4-Dimethyl-3-pentanone	C ₇ H ₁₄ O	—	15	—	—
3,3-Diethoxypropene	C ₇ H ₁₄ O ₂	—	4	—	—
Ethyl isovalerate	C ₇ H ₁₄ O ₂	—	25	—	—
Isobutyl propionate	C ₇ H ₁₄ O ₂	—	18	—	—
Isopentyl acetate	C ₇ H ₁₄ O ₂	—	25	(1.1–7.0/100°)	—
Isopropyl butyrate	C ₇ H ₁₄ O ₂	—	24	—	—
Isopropyl isobutyrate	C ₇ H ₁₄ O ₂	—	20	—	—
Pentyl acetate	C ₇ H ₁₄ O ₂	—	25	1.1–7.5	—
2-Pentyl acetate	C ₇ H ₁₄ O ₂	—	23	1.1–7.5	—
2,6-Dimethylpiperidine	C ₇ H ₁₅ N	—	16	—	—
1-Ethylpiperidine	C ₇ H ₁₅ N	—	19	—	—
2-Ethylpiperidine	C ₇ H ₁₅ N	—	–12	—	—
2,3-Dimethylpentane	C ₇ H ₁₆	—	–56	1.1–6.7	—
2,4-Dimethylpentane	C ₇ H ₁₆	—	–12	—	—
Heptane	C ₇ H ₁₆	—	–4	1.2–6.7	223
2-Methylhexane	C ₇ H ₁₆	—	–1	1.0–6.0	—
3-Methylhexane	C ₇ H ₁₆	—	–1	1.0–6.0	—
2,2,3-Trimethylbutane	C ₇ H ₁₆	—	–20	1.0–?	—
Phenylglyoxal	C ₈ H ₆ O ₂	—	< 23	—	—
1,3,5,7-Cyclooctatetraene	C ₈ H ₈	—	21	—	—
Ethylbenzene	C ₈ H ₁₀	—	15	1.2–6.8	—
<i>o</i> -Xylene	C ₈ H ₁₀	—	17	1.0–6.0	—
<i>m</i> - or <i>p</i> -Xylene	C ₈ H ₁₀	—	25	1.1–7.0	—
4-Vinylcyclohexene	C ₈ H ₁₂	—	16	—	—
1,7-Octadiene	C ₈ H ₁₄	—	25	—	—
1-Octyne	C ₈ H ₁₄	—	16	—	—
2-, 3-, or 4-Octyne	C ₈ H ₁₄	—	all < 23	—	—
Vinylcyclohexane	C ₈ H ₁₄	—	16	—	—
<i>cis</i> -1,2-Dimethylcyclohexane	C ₈ H ₁₆	—	11	—	—
<i>trans</i> -1,2-Dimethylcyclohexane	C ₈ H ₁₆	—	7	—	—
1,3-Dimethylcyclohexane	C ₈ H ₁₆	—	6	—	—
<i>cis</i> -1,4-Dimethylcyclohexane	C ₈ H ₁₆	—	10	—	—
<i>trans</i> -1,4-Dimethylcyclohexane	C ₈ H ₁₆	—	16	—	—
1-Octene	C ₈ H ₁₆	—	8	—	—
2-Octene	C ₈ H ₁₆	—	14	—	—
2,3,4-Trimethyl-1-pentene	C ₈ H ₁₆	—	5	—	—
2,4,4-Trimethyl-1-pentene	C ₈ H ₁₆	—	–29	—	—
2,3,4-Trimethyl-2-pentene	C ₈ H ₁₆	—	1	—	—
2,4,4-Trimethyl-2-pentene	C ₈ H ₁₆	—	2	—	—
3,4,4-Trimethyl-2-pentene	C ₈ H ₁₆	—	< 21	—	—
2-Ethylhexanal	C ₈ H ₁₆ O	—	—	—	197
2,3-Dimethylhexane	C ₈ H ₁₈	—	7(o)	—	—
2,4-Dimethylhexane	C ₈ H ₁₈	—	10(o)	—	—
3-Ethyl-2-methylpentane	C ₈ H ₁₈	—	< 21	—	—
2-Methylheptane	C ₈ H ₁₈	—	4	—	—
3-Methylheptane	C ₈ H ₁₈	—	–23	—	—
Octane	C ₈ H ₁₈	—	13	1.0–4.7	220
2,2,3-Trimethylpentane	C ₈ H ₁₈	—	< 21	—	—
2,2,4-Trimethylpentane	C ₈ H ₁₈	—	–12	1.1–6.0	—
2,3,3-Trimethylpentane	C ₈ H ₁₈	—	<21	—	—
2,3,4-Trimethylpentane	C ₈ H ₁₈	—	4	—	—
Dibutyl ether	C ₈ H ₁₈ O	—	25	1.5–7.6	194
Di- <i>tert</i> -butyl peroxide	C ₈ H ₁₈ O ₂	—	12	—	—

Name	Formula	B.P./°C	Fl.P./°C	E.L./%	A.I.T./°C
Bis(2-ethoxyethyl) ether	C ₈ H ₁₈ O ₃	—	—	—	205
2-(2-Butoxyethoxy)ethanol	C ₈ H ₁₈ O ₃	—	—	—	225
Di-2-butylamine	C ₈ H ₁₉ N	—	24	—	—
Diisobutylamine	C ₈ H ₁₉ N	—	21	—	—
2,6-Dimethyl-3-heptene	C ₉ H ₁₈	—	21(o)	—	—
1,3,5-Trimethylcyclohexane	C ₉ H ₁₈	—	19	—	—
3,3-Diethylpentane	C ₉ H ₂₀	—	< 21	0.7–7.7	—
2,5-Dimethylheptane	C ₉ H ₂₀	—	23	—	—
3,5-Dimethylheptane	C ₉ H ₂₀	—	23	—	—
4,4-Dimethylheptane	C ₉ H ₂₀	—	21	—	—
3-Ethyl-2,3-dimethylpentane	C ₉ H ₂₀	—	8	—	—
3-Ethyl-4-methylhexane	C ₉ H ₂₀	—	24	—	—
4-Ethyl-2-methylhexane	C ₉ H ₂₀	—	< 21	—	—
2- or 3-Methyloctane	C ₉ H ₂₀	—	—	—	220
4-Methyloctane	C ₉ H ₂₀	—	—	—	225
Nonane	C ₉ H ₂₀	—	—	—	205
2,2,5-Trimethylhexane	C ₉ H ₂₀	—	13	—	—
2,2,3,3-Tetramethylpentane	C ₉ H ₂₀	—	< 21	0.8–4.9	—
2,2,3,4-Tetramethylpentane	C ₉ H ₂₀	—	< 21	—	—
Dicyclopentadiene	C ₁₀ H ₁₂	—	-7(o)	—	—
2-Ethylhexyl vinyl ether	C ₁₀ H ₂₀ O	—	—	—	202
Isopentyl isovalerate	C ₁₀ H ₂₀ O ₂	—	24	—	—
Decane	C ₁₀ H ₂₂	—	—	—	210
2-Methylnonane	C ₁₀ H ₂₂	—	—	—	210
Dipentyl ether	C ₁₀ H ₂₂ O	—	—	—	170
<i>tert</i> -Butyl peroxybenzoate	C ₁₁ H ₁₄ O ₃	—	19	—	—
Diethyl ether	C ₁₂ H ₂₆ O	—	—	—	185
Triisobutylaluminium	C ₁₂ H ₂₇ Al	—	< 0	—	< 4
Tributylphosphine	C ₁₂ H ₂₇ P	—	—	—	200
Tetradecane	C ₁₄ H ₂₀	—	—	—	200
Hexadecane	C ₁₆ H ₃₄	—	—	—	205
Dichlorosilane	Cl ₂ H ₂ Si	8.3	—	—	58 ± 5°
Trichlorosilane	Cl ₃ HSi	32	-50	—	104
Deuterium	D ₂	-249	—	5–75	—
Trifluorosilane	F ₃ HSi	-95	—	flammable	—
Tetrafluorohydrazine	F ₄ N ₂	-73	—	—	explosive
Germane	GeH ₄	-90	—	—	ambient
Digermane	Ge ₂ H ₆	29	—	—	—
Hydrogen	H ₂	-253	—	4.1–74.2	—
Hydrogen sulfide	H ₂ S	-62	—	4–44	—
Hydrogen disulfide	H ₂ S ₂	—	< 22	—	—
Hydrogen selenide	H ₂ Se	-42	—	—	—
Hydrogen telluride	H ₂ Te	-2	—	—	50°
Ammonia	H ₃ N	-33	—	15.8–25	—
Phosphine	H ₃ P	-88	—	1.79–?	see text
Stibine	H ₃ Sb	-17	—	—	—
Hydrazine	H ₄ N ₂	—	38	4.7–100	23 on rust 132 on iron 156 on st. steel
Diphosphane	H ₄ P ₂	—	—	—	ambient
Silane	H ₄ Si	-112	—	1.4–?	-162
Disilane	H ₆ Si ₂	-14.5	—	—	ambient
Trisilane	H ₈ Si ₃	< 0	< 0	—	ambient
Phosphorus (white) P	—	—	—	—	38.5
Sulfur (hot solid) S	—	—	see text	see text	see text
Sulfur (hot liquid) S	—	—	see text	—	see text

Appendix 3

Glossary of Abbreviations and Technical Terms

Aerobic	In presence of air
AIT	Autoignition temperature
Alloy	Mixture of 2 or more metals
Amalgam	Alloy of a metal with mercury
Ambient	Usual or surrounding
Anaerobic	In absence of air
Analogue	Compound of the same structural type
AO	Active oxygen content of peroxides
Aprotic	Without labile hydrogen atom(s)
Aqua regia	Mixture of nitric and hydrochloric acids
ARC	Accelerating rate calorimetry
ASTM	American Society for Testing and Materials
Autoxidation	Slow reaction with air
BAM	Bundes Anstalt fur Materialsprufung (similar to ASTM)
Basic	Fundamental, or, alkaline (acting as a base)
BLEVE	Boiling liquid expanding vapour explosion
Blowing agent	Material producing much gas on decomposition
b.p.	Boiling point
BSC	Bench scale calorimeter
Carbonaceous	Containing elemental carbon (as opposed to organic, containing combined carbon)
CHETAH	A computer program to predict energy release hazards
CIT	Critical ignition temperature
Class	Collection of related chemical groups or topics
COI	Critical oxygen index
conc.	concentrated
Congener	Compound with related but not identical structure
COT	Cyclooctatetraene
Critical diameter	Minimum diameter of an explosive charge capable of maintaining detonation
Cryogenic	At a very low (freezing) temperature
CVD	Chemical vapour deposition

Deflagration	Self sustaining internal combustion propagating by means of molecular heat transfer slower than the speed of sound (the explosion mechanism gunpowder and other 'low' explosives)
DH°f	Standard heat of formation
Desiccate	Dry intensively
Detonable	Capable of detonation
Detonation	A self sustaining decomposition reaction propagating faster than the speed of sound by means of a shock wave (the characteristic property of 'high' as opposed to 'low' explosives).
Diglyme	Diethyleneglycol dimethyl ether
Digol	Diethyleneglycol
DMF	Dimethylformamide
DMSO	Dimethyl sulfoxide
DSC	Differential scanning calorimetry
DTA	Differential thermal analysis
EL	Explosive limits (or Flammable limits), vol% in air
Endotherm	Absorption of heat
ESCA	Electron scanning chemical analysis
Exotherm.	Liberation of (reaction) heat
Freeze drying	Drying without heat by vacuum evaporation of frozen solvent
GLC	Gas-liquid chromatography
Glyme	Diethyleneglycol monomethyl ether
Halocarbon	Partially or fully halogenated hydrocarbon
HMPA	Hexamethylphosphoramide
HMSO	Her Majesty's Stationery Office
Homologue	Compound of the same (organic) series
Hypergolic	Ignites on contact
Ignition source	A source of energy which ignites flammables
IMS	Industrial methylated spirit (ethanol)
Induction period	Delay in reaction starting, caused by inhibitors
Inorganic	Not containing combined carbon, of mineral origin
Intermolecular	Between different molecules
Intramolecular	Within the same molecule
IR	Infrared spectroscopy
Initiation	Triggering of explosion or decomposition
LAH	Lithium tetrahydroaluminate
Lanthanide	Of the group of rare-earth metals
LRS	Laser Raman spectroscopy
LPG	Liquefied petroleum gas
MAPP	Methylacetylene/propadiene/propene mixture
Molecular sieve	A zeolite lattice with micropores of specific sizes, useful for molecular separations

m.p.	Melting point
MRH	Maximum reaction heat
NMR	Nuclear magnetic resonance spectroscopy
Nomenclature	System of naming chemicals
Off-spec.	Off-specification (low quality)
Oleum	Sulfur trioxide dissolved in sulfuric acid
Organic	A compound containing combined carbon
Organometallic	Containing carbon to metal bonding
Oxidant	Oxidising agent (electron sink)
PCB	Polychlorinated biphenyl
PTFE	Polytetrafluoroethylene
Propagation	Spread or transmission of decomposition, flame or explosion
Propellant	Energetic composition used in ballistics
PVC	Polyvinyl chloride
Pyrophoric	Igniting on contact with air (or frictional sparking)
Q	Heat of (exothermic) reaction or polymerisation
Quaternary salt	Tetra-substituted ammonium salt etc.
RCHD	Reactive Chemical Hazards Database, which you are using
Redox compound	Compound with reducing and oxidising features
Reducant	Reducing agent (electron source)
Refractory	Heat resisting
REITP2	A computer program to calculate MRH
RSC	Reaction Safety Calorimeter, or Royal Society of Chemistry
Runaway	Reaction out of control
Self-accelerating	Reaction catalysed by its own products
Self-heating	When substance generates heat faster than it is dissipated
Silicones	Organic derivatives of a polysiloxane chain
Slurry	Pourable mixture of solid and liquid
Smelt	Molten sodium sulfide and carbonate from evaporated sulfite liquor
Substance	Not a single chemical species, often of natural origin
Superheated	A liquid at a temperature above its boiling point
Tait24	Temperature of adiabatic storage which gives an induction time to exothermic decomposition of 24 hours
Thermite reaction	Reaction of aluminium powder and iron oxide producing molten iron (and analogous reactions)
THF	Tetrahydrofuran
TGA	Thermogravimetric analysis
Thermochemistry	Study of heat effects of chemical reactions
TLC	Thin layer chromatography
TNT	Trinitrotoluene
U	Heat of (exothermic) decomposition

Unit operation	A single operational stage of a chemical process sequence
Unit process	A single chemical reaction stage in a process sequence
USNTIS	US National Technical Information Service
UV	Ultraviolet spectroscopy

Index of Chemical Names and Synonyms used in Section 1

The majority of the names for chemicals in this alphabetically arranged index conform to one of the systematic series permitted under various sections of the IUPAC Definitive Rules for Nomenclature. Where there is a marked difference between these names and the alternative names recommended in the IUPAC-based BS2472:1983 or ASE 1985 nomenclature lists, or long established traditional names, these are given as synonyms in parentheses after the main title. These synonyms also have their own index entry, cross-referenced back to the IUPAC-based names used as bold titles in the text of Volume 1.

Those titles prefixed in the text with a dagger to show high flammability risks are prefixed similarly in this index. The four character numbers following the names or synonyms are the serial numbers of the entries, not page numbers.

It should be noted that italicised hyphenated prefixes which indicate structure, such as *cis-*, *o-*, *m-*, *tert-*, *mixo-*, *N-*, *O-*, etc., have been ignored during the alpha-sorting routine used on this index and the group-lists, while the roman character structural prefixes iso and neo, and roman multiplying prefixes such as di, tris, tetra and hexakis, have been included in the indexing procedure.

- † Acetaldehyde, 0828
- † Acetaldehyde oxime, 0865
- Acetic acid, 0833
- Acetic anhydride, 1534
- Acetoacet-4-phenetidide, 3529
- Acetohydrazide, 0912
- † Acetone, 1220
- Acetone oxime, 1258
- † Acetonitrile, 0758
- Acetonitrileimidazole-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II) perchlorate, 3828
- Acetonitrilethiol, *see* Mercaptoacetonitrile, 0767
- Acetoxymercurio(perchloratodimercurio)ethenone, 1413
- 1-Acetoxy-1-hydroperoxy-6-cyclodecanone, 3669
- 3-Acetoxy-4-iodo-3,7,7-trimethylbicyclo[4.1.0]heptane, 3544
- Acetoxymercurio(perchloratomercurio)ethenone, 1412
- 4-Acetoxy-3-methoxybenzaldehyde, 3290
- 4-Acetoxy-3-methoxy-2-nitrobenzaldehyde, 3266
- 1-Acetoxy-6-oxo-cyclodecyl hydroperoxide, 3547
- 3-Acetoxy-1,2,4-trioxolane, *see* Vinyl acetate ozonide, 1543
- 2-Acetylamino-3,5-dinitrothiophene, 2280
- Acetyl azide, 0771
- Acetyl bromide, 0728
- † Acetyl chloride, 0735
- Acetyl cyclohexanesulfonyl peroxide, 3033
- Acetyldimethylarsine, 1628
- † Acetylene, 0686
- Acetylenebis(triethyllead), 3672
- Acetylenebis(triethyltin), 3674
- Acetylenedicarboxaldehyde, 1403
- Acetylenedicarboxylic acid, 1405
- Acetyl hypobromite, 0729
- Acetyl hypofluorite, 0751
- 2-Acetyl-3-methyl-4,5-dihydrothiophen-4-one, 2807
- Acetyl nitrate, 0765
- Acetyl nitrite, 0762
- Acetyl nitro peroxide, *see* Peroxyacetyl nitrate, 0766
- O*-Acetylsalicylic acid, 3137
- 1-Acetyl-4-(4'-sulfophenyl)-3-tetrazene, 2982
- 4-(4-Acetyl-1-tetrazenyl)benzenesulfonic acid, *see* 1-Acetyl-4-(4'-sulfophenyl)-3-tetrazene, 2982
- Aconitic acid, *see* *E*-Propene-1,2,3-tricarboxylic acid, 2342
- † Acrylaldehyde, 1145
- † Acrylamide, 1180
- † Acrylic acid, 1148
- † Acrylonitrile, 1107

- † Acrylonitrile, 1107
- † Acryloyl chloride, 1093
- 1,2-Cyclohexanedione, 2385a
- Adipic acid, 2441
- Alane–*N,N*-dimethylethylamine, 0071
- † Allene, *see* Propadiene, 1124
- all isomers except 3,5-, *see* Dinitroanilines, 2276
- † ‘Alloxan’, *see* Pyrimidine-2,4,5,6-(1*H*,3*H*)-tetrone 1398
- † Allyl acetate, 1912
- † Allyl alcohol, *see* 2-Propen-1-ol, 1223
- † Allylamine, *see* 3-Aminopropene, 1254
- † Allyl azide, *see* 3-Azidopropene, 1188
- Allyl benzenesulfonate, 3155
- † Allyl bromide, *see* 3-Bromo-1-propene, 1153
- † Allyl chloride, *see* 3-Chloropropene, 1158
- Allyldimethylarsine, 1983
- † Allyl ethyl ether, 1955
- † Allyl formate, 1524
- Allyl glycidyl ether, 2-Propenyloxymethyloxirane, *see* 1-Allyloxy-2,3-epoxypropane, 2434
- Allyl hydroperoxide, 1226
- † Allyl iodide, *see* 3-Iodopropene, 1174
- Allyl isothiocyanate, 1471
- Allyllithium, 1177
- † Allyl mercaptan, *see* 2-Propene-1-thiol, 1239
- Allylmercury(II) iodide, 1173
- 1-Allyloxy-2,3-epoxypropane, 2434
- Allyl phosphorodichloridite, 1169
- N*-Allylthiourea, 1600
- Allyl 4-toluenesulfonate, 3315
- † Allyl vinyl ether, 1904
- Allylzinc bromide, 1155
- ‘Alumina’, *see* Aluminium oxide 0087
- Aluminium, 0048
- Aluminium abietate, 3917
- Aluminium amalgam, 0051
- Aluminium azide, 0082
- Aluminium bromide, 0060
- Aluminium carbide, 1031
- Aluminium chlorate, 0065
- Aluminium chloride, 0062
- Aluminium chloride–nitromethane, 0063
- Aluminium chloride–trimethylaluminium complex, *see* Trimethyldialuminium trichloride, 1292
- Aluminium–cobalt alloy, 0049
- Aluminium copper(I) sulfide, 0084

Aluminium–copper–zinc alloy, 0050
 Aluminium dichloride hydride diethyl etherate, 0061
 Aluminium formate, 1089
 Aluminium hydride, 0070
 Aluminium hydride–diethyl ether, 0073
 Aluminium hydride–trimethylamine, 0072
 Aluminium hydroxide, 0074
 Aluminium iodide, 0079
 Aluminium isopropoxide, 3216
 Aluminium–lanthanum–nickel alloy, 0080
 Aluminium–lithium alloy, 0052
 Aluminium–magnesium alloy, 0053
 Aluminium–magnesium–zinc alloy, 0054
 Aluminium–mercury alloy, *see* Aluminium amalgam, 0051
 Aluminium–nickel alloys, 0055
 Aluminium oxide, 0087
 Aluminium perchlorate, 0066
 Aluminium phosphide, 0083
 Aluminium phosphinate, 0078
 Aluminium stearate, 3914
 Aluminium tetraazidoborate, 0059
 Aluminium tetrahydroborate, 0058
 Aluminium–titanium alloys, 0056
 Americium trichloride, 0090
 Amidosulfuric acid, 4499
 Amidosulfuryl azide, *see* Sulfamoyl azide, 4472
 4-Aminobenzenediazonium perchlorate, 2304
 † 1-Aminobutane, *see* Butylamine, 1723
 † 2-Aminobutane, *see* 2-Butylamine, 1724
 2-Amino-4,6-dihydroxy-5-nitropyrimidine, 1436
 2-Amino-4,6-dinitrophenol, 2278
 2-Amino-3,5-dinitrothiophene, 1421
 3-Amino-2,5-dinitrotoluene, 2783
 2-(2-Aminoethyl)aminoethanol, *see* *N*-2-Hydroxyethyl-1,2-diaminoethane,
 1756
 2-(2-Aminoethylamino)-5-methoxynitrobenzene, 3172
 2-Aminoethylammonium perchlorate, 0958
 5-Amino-2-ethyl-2*H*-tetrazole, 1268
 Aminoguanidine, 0507
 Aminoguanidinium nitrate, 0512
 3-Aminoisoxazole, 1136
 5-Aminoisoxazole-3-carbonamide, 1476
 4-Amino-3-isoxazolidinone, 1137
 3-Amino-5-methylisoxazole, 1501
 5-Amino-3-methylisoxazole, 1502
 2-Amino-4-methyloxazole, 1500

- † 2-Amino-2-methylpropane, *see tert*-Butylamine, 1725
- 5-Amino-3-methylthio-1,2,4-oxadiazole, 1192
- 1-Amino-3-nitroguanidine, 0495
- 2-Amino-5-nitrophenol, 2316
- 3-Amino-6-nitro-1,2,4,5-tetrazine-2,4-dioxide, 0718
- 2-Amino-5-nitrothiazole, 1117
- 6-Aminopenicillanic acid *S*-oxide, 3010
- 4-Aminophenylazobenzene, 3487
- 5-Amino-3-phenyl-1,2,4-triazole, 2952
- 1-Amino-2-propanol, 1311
- † 3-Aminopropene, 1254
- 3-Aminopropionitrile, 1212
- 5-Aminotetrazole, 0461
- 5-Amino-1,2,3,4-thiatriazole, 0413
- 2-Aminothiazole, 1140
- 4-Amino-4*H*-1,2,4-triazole, 0812
- Amminebarium bis(nitrophenylide), 3518
- Amminedecahydroxydiplatinum, 4593
- Ammine-1,2-diaminoethanediperoxochromium(IV), 0968
- Amminepentahydroxyplatinum, 4571
- † Ammonia, 4497
- 4-(2-Ammonio-2-carboxyethylthio)-5,7-dinitro-4,5-dihydrobenzofurazanide
 N-oxide, 3146
- 2-Ammoniothiazole nitrate, 1194
- Ammonium amidoselenate, 4557
- Ammonium amidosulfate, 4556
- Ammonium azide, 4526
- Ammonium bromate, 0252
- Ammonium bromide, 0251
- Ammonium chlorate, 4003
- Ammonium chloride, 4001
- Ammonium decahydrodecaborate(2-), 0199
- Ammonium dichromate, 4246
- Ammonium dinitramide, 4528
- Ammonium 3,5-dinitro-1,2,4-triazolide, 0823
- Ammonium 1,2-ethylenebis(nitramide), 0972
- Ammonium fluoride, 4299
- Ammonium fluorochromate, 4224
- Ammonium hexachloroplatinate, 4186
- Ammonium hexacyanoferrate(II), 2577
- Ammonium hexanitrocerate, 3964
- Ammonium hexanitrocobaltate(3-), 4206
- Ammonium hydrogen sulfite, 4545
- Ammonium hydroxide, 4544
- Ammonium hydroxyoxodiperoxochromate(1-), 4230
- Ammonium iodate, 4513

Ammonium iodide, 4512
Ammonium iron(III) sulfate, 4393
Ammonium 3-methyl-2,4,6-trinitrophenoxide, 2803
Ammonium nitrate, 4522
Ammonium nitridoosmate, 4523
Ammonium nitrite, 4521
Ammonium *aci*-nitromethanide, 0504
Ammonium *N*-nitrosophenylaminoxide, 2399
Ammonium pentaperoxodichromate(2-), 4247
Ammonium perchlorate, 4004
Ammonium perchlorylamide, 4009
Ammonium periodate, 4514
Ammonium permanganate, 4518
Ammonium peroxoborate, 0146
Ammonium peroxodisulfate, 4576
Ammonium persulphate, *see* Ammonium peroxodisulfate, 4576
Ammonium phosphinate, 4554
Ammonium picrate, 2322
Ammonium sulfamate, *see* Ammonium amidosulfate, 4556
Ammonium sulfate, 4574
Ammonium sulfide, 4577
Ammonium tetrachromate(2-), 4253
Ammonium tetranitroplatinate(II), 4578
Ammonium tetraperochromate(3-), 4232
Ammonium thiocyanate, 0479
Ammonium thiosulfate, 4573
Ammonium trichromate(2-), 4252
Ammonium 2,4,5-trinitroimidazolid, 1144
Ammonium 2,4,6-tris(dioxoselena)hexahydrotriazine-1,3,5-triide, 4591
'Ammonium triselenimidate', *see* Ammonium 2,4,6-tris(dioxoselena)
hexahydrotriazine-1,3,5-triide, 4591
Aniline, 2354
Anilinium chloride, 2365
Anilinium nitrate, 2379
Anilinium perchlorate, 2367
Anisaldehyde, *see* 4-Methoxybenzaldehyde, 2956
m-Anisidine, *see* 3-Methoxyaniline, 2817
o-Anisidine, *see* 2-Methoxyaniline, 2816
p-Anisidine, *see* 4-Methoxyaniline, 2818
Anisoyl chloride, *see* 4-Methoxybenzoyl chloride, 2930
Anisyl chloride, *see* 4-Methoxybenzyl chloride, 2962
Anthracene, 3632
Antimony, 4907
Antimony(III) chloride oxide, 4041
Antimony(III) nitride, 4729
Antimony(III) oxide, 4851

Antimony(III) oxide perchlorate, 4046
Antimonyl chloride, *see* Antimony trichloride oxide, 4150
Antimony pentachloride, 4184
Antimony trichloride, 4157
Antimony trichloride oxide, 4150
Antimony trisulfide, 4906
Aqua-1,2-diaminoethanediperoxochromium(IV), 0965
Aqua-1,2-diaminopropanediperoxochromium(IV) dihydrate, 1336
trans-Aquadioxo(terpyridine)ruthenium(2+) diperchlorate, 3687
Aquafluorobis(1,10-phenanthroline)chromium(III) perchlorate, 3846
Aqua-glycinatonitratocopper, *see* Copper(II) glycinate nitrate, 0905
Argon, 0091
Arsenic, 0092
Arsenic pentafluoride, 0095
Arsenic pentaoxide, 0106
Arsenic trichloride, 0094
Arsenic trioxide, 0105
Arsenic trisulfide, 0108
Arsenous triazide, *see* Triazidoarsine 0103
† Arsine, 0100
Arsine–boron tribromide, 0101
Ascaridole, *see* 1,4-Epidioxy-2-*p*-menthene, 3347
2-Aza-1,3-dioxolanium perchlorate, 0898
2-Azatricyclo[2.2.1.0^{2,6}]hept-7-yl perchlorate, 2368
AZDN, *see* Azoisobutyronitrile, 3011
Azelaic acid, 3187
† Azetidine, 1255
Azidoacetaldehyde, 0772
Azidoacetic acid, 0774
† Azidoacetone, 1190
Azidoacetone oxime, 1215
Azidoacetonitrile, 0714
Azidoacetyl chloride, 0693
4-Azidobenzaldehyde, 2697
Azidobenzene, *see* Phenyl azide, 2271
7-Azidobicyclo[2.2.1]hepta-2,5-diene, 2777
2-Azido-1,3-butadiene, 1472
1-Azidobutane, 1666
Azido-2-butyne, 1473
N-Azidocarbonylazepine, 2728
Azidocarbonyl fluoride, 0339
Azidocarbonylguanidine, 0820
4-Azidocarbonyl-1,2,3-thiadiazole, 1069
2-*trans*-1-Azido-1,2-dihydroacenaphthyl nitrate, 3466
N-Azidodimethylamine, 0915
4-Azido-*N,N*-dimethylaniline, 3331

Azidodimethylborane, 0888
 2-Azido-3,5-dinitrofurane, 1384
 Azidodithioformic acid, 0386
 2-Azidoethanol, 0873
 Azidogermane, 4416
 6-Azidohexyl 6-azidohexanoate, 3551
 Azidoiodoiodonium hexafluoroantimonate, 4361
 2-Azidomethylbenzenediazonium tetrafluoroborate, 2708
 Azidomethyloxirane, *see* Glycidyl azide, 1191
 Azidopentafluorotungsten, 4358
 2-Azidoperfluoronaphthalene, 3234
 2-Azido-2-phenylpropane, 3159
 1-(2-Azido-2-phenylvinyl)phenyliodonium 4-toluenesulfonate, 3821
 † Azido-2-propanone, *see* Azidoacetone, 1190
 † 3-Azidopropene, 1188
 3-Azidopropyne, 1114
 Azidosilane, 4501
 5-Azidotetrazole, 0388
 3-Azido-1,2,4-triazole, 0717
 Azidotrifluoromethane, *see* Trifluoromethyl azide, 0347
 † Azidotrimethylsilane, *see* Trimethylsilyl azide, 1314
 Azinphosmethyl, *see* *O,O*-Dimethyl-*S*-[(4-oxo-1,2,3-benzotriazin-3[4*H*]-yl)methyl] phosphorodithioate, 3307
 † Aziridine, 0863
 Azobenzene, 3483
cis-Azobenzene, 3484
 2,2'-Azobis(2-amidiniopropane) chloride, 3089
 2,2'-Azobis(2-amidiniopropane) peroxodisulfate, 3091
 2,2'-Azobis(2,4-dimethylvaleronitrile), 3668
 Azocarbonitrile, *see* Dicyanodiazene, 1005
 Azo-*N*-chloroformamidine, 0792
 1,1'-Azo-1,2-dicarbadeborane(14), 1796
 Azodicarboxamide, *see* Azoformamide, 0816
 2,2'-Azo-3,5-dinitropyridine, 3238
 Azoformaldoxime, 0815
 Azoformamide, 0816
 Azoisobutyronitrile, 3011
 2,2'-Azoisovaleronitrile, 3345
 Azomethane, 0910
 1,1'-Azo-[2-methyl-1,2-dicarbadeborane(14)] 2624
 Azo-*N*-methylformamide, 1601
 3-Azoniabicyclo[3.2.2]nonane nitrate, 3048
 Azo-*N*-nitroformamidine, 0825
 3,3'-Azo-(1-nitro-1,2,4-triazole), 1401
 2-Azoxyanisole, 3653
 Azoxybenzene, 3485

Barium, 0200
 Barium acetate, 1486
 Barium acetylide, 0575
 Barium amidosulfate, 0209
 Barium azide, 0214
 Barium 5,5'-azotetrazolide, 0577
 Barium bromate, 0202
 Barium chlorate, 0205
 Barium chlorite, 0204
 Barium 1,3-di(5-tetrazolyl)triazene, 1424
 Barium hydride, 0207
 Barium hydroxide, 0208
 'Barium hypophosphite', *see* Barium phosphinate, 0210
 Barium iodate, 0211
 Barium methyl hydroperoxide, 0889
 Barium nitrate, 0212
 Barium nitride, 0219
 Barium nitridoosmate, 0213
 Barium oxide, 0215
 Barium perchlorate, 0206
 Barium perchlorylamide, 0203
 Barium *N*-perchlorylbzylamide, 3649
 Barium peroxide, 0216
 Barium phosphinate, 0210
 Barium sulfate, 0217
 Barium sulfide, 0218
 Barium tetrafluorobromate, 0201
 Barium thiocyanate, 0576
 Basic lead carbonate, *see* Lead carbonate–lead hydroxide, 0726
 'Basic stannous nitrate', *see* Tetrahydroxotritin(2+) nitrate, 4525
 Basic trihydrazinecadmium perchlorate, 3951
 † Benzaldehyde, 2731
 Benzaldehyde oxime, 2760
 Benzanthrone, 3736
 † Benzene, 2288
 Benzeneamine, *see* Aniline, 2354
 Benzene-1,4-bis(diazonium perchlorate), 2160
 Benzene-1,3-bis(sulfonyl azide), 2210
 η -Benzenecyclopentadienyliron(II) perchlorate, 3396
 Benzenediazonium-2-carboxylate, 2659
 Benzenediazonium-3-carboxylate, 2660
 Benzenediazonium-4-carboxylate, 2661
 Benzenediazonium chloride, 2230
 Benzenediazonium hydrogen sulfate, 2318
 Benzenediazonium iodide, 2250
 Benzenediazonium nitrate, 2275

Benzenediazonium-4-oxide, 2189
 Benzenediazonium perchlorate, 2232
 Benzenediazonium-2-sulfonate, 2192
 Benzenediazonium-4-sulfonate, 2193
 Benzenediazonium tetrachlorozincate, 3476
 Benzenediazonium tetrafluoroborate, 2223
 Benzenediazonium tetraphenylborate, 3876
 Benzenediazonium tribromide, 2227
 Benzenediazonium triiodide, 2254
 1,4-Benzenedicarbonyl chloride, *see* Terephthaloyl chloride, 2889
 1,2-Benzenediol, 2331
 1,3-Benzenediol, 2332
 1,4-Benzenediol, 2333
 1,4-Benzenediol–oxygen complex, 2334
 Benzeneethanol, 2983
 Benzeneperoxyseleninic acid, 2337
 Benzeneperoxysulfonic acid, 2341
 η^6 -Benzeneruthenium(II) dinitrate, 2320
 Benzeneseleninic acid, 2335
 Benzeneseleninic anhydride, 3495
 Benzeneselenonic acid, 2338
 (Benzenesulfinato-*S*)pentaamminecobalt(III) trichloro(perchlorato)stannate(II),
 2614
 Benzenesulfinyl azide, 2273
 Benzenesulfinyl chloride, 2234
 Benzenesulfonic anhydride, 3498
 Benzenesulfonyl azide, 2274
 Benzenesulfonyl chloride, 2235
 Benzenethiol, 2344
cis-Benzene triimine, *see* 3,6,9-Triazatetracyclo[6.1.0.0^{2,4}.0^{5,7}]nonane, 2398
 Benzene triozone, 2343
 Benzidine (4,4'-Diaminobiphenyl), 3516
 Benzimidazole, 2724
 Benzimidazolium 1-nitroimidate, 2704
 3-Benzocyclobutenylpotassium, 2904
 Benzodiazole, *see* Benzimidazole, 2724
 2,3-Benzodioxin-1,4-dione, *see* Phthaloyl peroxide, 2900
 1,3-Benzoditholium perchlorate, 2677
 Benzofurazan-*N*-oxide, 2190
 Benzohydrazide, 2798
 Benzoic acid, 2732
 1,4-Benzoquinone, 2214
 Benzoquinone 1,4-bis(chloroimine), 2159
 1,4-Benzoquinone-4-chloroimine, 2140
 1,4-Benzoquinone diimine, 2310
 1,4-Benzoquinone monoimine, 2260

1,2-Benzoquinone monoxime, *see* 2-Nitrosophenol, 2263
 1,4-Benzoquinone monoxime, *see* 4-Nitrosophenol, 2264
 Benzo-1,2,3-thiadiazole, 2199
 Benzo-1,2,3-thiadiazole 1,1-dioxide, 2191
 Benzotriazole, 2269
 Benzotriazolium 1-nitroimidate, 2281
 Benzotriazolium 2-nitroimidate, 2282
 Benzotri(furazan *N*-oxide), 2631
 Benzoyl azide, 2698
 5-Benzoylbenzenediazonium-2-oxide, 3597
 Benzoyl chloride, 2675
 Benzoyl nitrate, 2689
 1-Benzoyl-1-phenyldiazomethane, 3638
 Benzvalene, 2289
 Benzyl alcohol, 2806
 Benzylamine, 2812
 Benzyl azide, 2779
 Benzyl bromide, 2735
 Benzyl chloride, 2738
 Benzyl chloroformate, 2931
 Benzyl dimethylamine, 3168
 Benzyl fluoride, 2746
 2-Benzylideneamino-4,6-dinitrophenol, 3600
 Benzyl nitrate, 2765
 Benzyloxyacetylene, 3133
 Benzylsilane, 2832
 Benzylsodium, 2787
 1-Benzyl-3-(4-tolyl)triazene, 3657
 Benzyltriethylammonium permanganate, 3617
 Benzyltrimethylammonium permanganate, 3342
 Beryllium, 0220
 Beryllium chloride, 0221
 Beryllium fluoride, 0223
 Beryllium hydride, 0224
 Beryllium oxide, 0225
 Beryllium perchlorate, 0222
 Beryllium tetrahydroborate, 0157
 Beryllium tetrahydroborate–trimethylamine, 1337
 1,1'-Biaziridinyl, 1593
 † Bicyclo[2.2.1]hepta-2,5-diene, 2789
 Bicyclo[2.1.0]pent-2-ene, 1856
 2,2'-Bi-5,6-dihydro-1,3-dithiolo[4,5-b][1,4]dithiinylidene, 3263
 2,2'-Bi-1,3-dithiole, 2215
 1,1'-Biphenyl-4,4'-bisdiazonium perchlorate, *see* 4,4'-Biphenylenebis
 (diazonium) perchlorate, 3457
 2,2-Biphenyldicarbonyl peroxide, 3631

4,4'-Biphenylenebis(diazonium) perchlorate, 3457
 4,4'-Biphenylene-bis-sulfonylazide, 3468
 2,2'-Bipyridine *N,N'*-dioxide-dicarbonylrhodium(I) perchlorate, 3454
 4,4'-Bipyridyl-bis(pentaammineruthenium(III) perchlorate), 3385
 2,2'-Bipyridyldichloropalladium(IV) perchlorate, 3255
 2,2'-Bipyridyl 1-oxide, 3258
 1,2-Bis(acetoxymethoxy)ethane, 3355
 Bis(acrylonitrile)nickel(0), 2312
 Bis(2-aminoethyl)amine, 1777
 Bis(2-aminoethyl)aminocobalt(III) azide, 1775
 Bis(2-aminoethyl)aminediperoxochromium(IV), 1776
 Bis(2-aminoethyl)aminesilver nitrate, 1773
N,N'-Bis(2-aminoethyl)1,2-diaminoethane, 2599
N,N-Bis(3-aminopropyl)-1,4-diazacycloheptanenickel(II) perchlorate, 3415
 1,3-Bis(5-amino-1,3,4-triazol-2-yl)triazene, 1576
 Bis(2-azidobenzoyl) peroxide, 3628
cis-1,2-Bis(azidocarbonyl)cyclobutane, 2323
 1,2-Bis(azidocarbonyl)cyclopropane, 1835
 1,2-Bis(2-azidoethoxy)ethane, 2479
 Bis(2-azidoethoxymethyl)nitramine, 2481
N,N-Bis(azidomethyl)nitric amide, *see* 1,3-Diazido-2-nitroazapropane, 0824
 3,3-Bis(azidomethyl)oxetane, 1902
 Bis(azidothiocarbonyl) disulfide, 1016
 Bis(benzeneazo) oxide, 3490
 Bis(η -benzene)chromium(0), 3511
 Bis(benzene)chromium dichromate, 3851
 Bis(benzenediazo) sulfide, 3491
 Bis(η -benzene)iron(0), 3512
 Bis(benzeneiron)-fulvalenediyl complex, 3834
 Bis(η -benzene)molybdenum(0), 3514
 Bis(1-benzo[d]triazolyl) carbonate, 3598
 Bis(1-benzo[d]triazolyl) oxalate, 3629
 1,1-Bis(benzoylperoxy)cyclohexane, 3804
 Bis(benzyl 1-methylhydrazinocarbodithioate *N*²,*S'*)(perchlorato-*O,O'*)
 copper(1+) perchlorate, 3766
 Bis[1,5-bis(4-methylphenyl)-1,3-pentaazadienato-N3,N5]-(T-4) cobalt, 3868
 Bis(borane)-hydrazine, 0143
 Bis(bromobenzenesulfonyl) peroxide, 3453
N,N-Bis(bromomercurio)hydrazine, 0268
 1,2-Bis(*tert*-butylphosphino)ethane, 3378
 Bis(3-carboxypropionyl) peroxide, 2990
 1,4-Bis(chlorimido)-2,5-cyclohexadiene, *see* Benzoquinone
 1,4-bis(chloroimine), 2159
 Bis(4-chlorobenzenediazo) oxide, 3456
 Bis(4-chlorobenzenesulfonyl) peroxide, 3458
I,I-Bis(3-chlorobenzoylperoxy)-4-chlorophenyliodine, 3793

I,I-Bis(3-chlorobenzoylperoxy)phenyliodine, 3794
 1,4-Bis(2-chloroethyl)-1,4-bis(azonia)bicyclo[2.2.1]heptane periodate, 3192
 Bis(2-chloroethyl)sulfide, 1589
 Bis(1-chloroethylthallium chloride) oxide, 1591
N,N'-Bis(chloromercurio)hydrazine, 4064
 1,2-Bis(chloromethyl)benzene, 2946
 † Bis(chloromethyl) ether, 0793
 Bis(chloromethyl)thallium chloride, 0796
 2,5-Bis(chloromethyl)thiophene, 2305
 1,6-Bis(4-chlorophenyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane, 3743
 Bis(2-cyanoethyl)amine, 2397
 μ -1,2-Bis(cyanomercurio)ethanediylidenedimercury(II), 1798
 Bis(cyclooctatetraene)uranium(0), 3708
 Bis(η^8 -1,3,5,7-cyclooctatetraene)cerium, *see* Bis(cyclooctatetraenyl)cerium, 3706
 Bis(cyclooctatetraenyl)cerium, 3706
 Bis(cyclopentadienyl)bis(pentafluorophenyl)zirconium, 3830
 Bis(cyclopentadienyldinitrosylchromium), 3275
 Bis(cyclopentadienyl)hexafluoro-2-butynechromium, 3636
 Bis(cyclopentadienyl)lead, 3294
 Bis(cyclopentadienyl)magnesium, 3277
 Bis(cyclopentadienyl)manganese, 3278
 Bis(cyclopentadienyl)niobium tetrahydroborate, 3324
 Bis(cyclopentadienyl)pentafluorophenylzirconium hydroxide, 3703
 Bis(cyclopentadienyl)phenylvanadium, 3705
 Bis(cyclopentadienyl)titanium, 3295
 Bis(cyclopentadienyl)titanium selenate, 3293
 Bis(cyclopentadienyl)tungsten diazide oxide, 3285
 Bis(cyclopentadienyl)vanadium diazide, 3286
 Bis(cyclopentadienyl)zirconium, 3296
 Bis(1,2-diaminoethane)diaquacobalt(III) perchlorate, 1793
 Bis(1,2-diaminoethane)dichlorocobalt(III) chlorate, 1786
 Bis(1,2-diaminoethane)dichlorocobalt(III) perchlorate, 1787
cis-Bis(1,2-diaminoethane)dinitrocobalt(III) iodate, 1788
 Bis(1,2-diaminoethane)dinitrocobalt(III) perchlorate, 1784
 Bis(1,2-diaminoethane)hydroxoorhenium(V) perchlorate, 1791
 Bis(1,2-diaminopropane)-*cis*-dichlorochromium(III) perchlorate, 2613
 1,10-Bis(diazonio)decaboran(8)ate, 0197
 1,4-Bis(1,2-dibromoethyl)benzene, 3268
 Bis(dibutylborino)acetylene, 3775
 1,2-Bis(di-*tert*-butylphosphino)ethane, 3776
 Bis(dicarbonylcyclopentadienyliron)-bis(tetrahydrofuran)magnesium, 3835
 Bis[dicarbonyl(cyclopentadienyl)tributylphosphinemolybdenum]-
 tetrakis(tetrahydrofuran)magnesium, 3913
 Bis(3,4-dichlorobenzenesulfonyl) peroxide, 3445
 Bis(2,4-dichlorobenzoyl) peroxide, 3623

1,2-Bis(dichlorophosphino)ethane, 0797
N,N-Bis(diethylboryl)methylamine, 3223
 Bis(diethylenetriamine)cobalt(III) perchlorate, 3106
 1,2-Bis(diethylphosphino)ethane, 3379
 Bis(difluoroamino)difluoromethane, 0362
 1,2-Bis(difluoroamino)ethanol, 0802
 1,2-Bis(difluoroamino)ethyl vinyl ether, 1496
 4,4-Bis(difluoroamino)-3-fluoroimino-1-pentene, 1867
 1,2-Bis(difluoroamino)-*N*-nitroethylamine, 0803
 Bis(1,*n*-difluorobenzene)chromium(0) isomers, 3461
 Bis(difluoroboryl)methane, 0394
 Bis(*S,S*-difluoro-*N*-sulfimido)sulfur tetrafluoride, 4381
 Bis(dihydroxygold)imide, 0114
 Bis(dimethylaminoborane)aluminium tetrahydroborate, 1795
 Bis(dimethylamino)dimethylstannane, 2597
 † 1,2-Bis(dimethylamino)ethane, 2579
 Bis(2-dimethylaminopyridine *N*-oxide)copper(II) perchlorate, 3665
 Bis(dimethylamino) sulfoxide, 1757
 Bis(dimethylarsinyldiazomethyl)mercury, 2467
 Bis(dimethylarsinyl) oxide, 1737
 Bis(dimethylarsinyl) sulfide, 1738
 Bis(dimethylstibinyl) oxide, 1761
 1,1-Bis(dimethyl sulfoxide)telluracyclopentane diperchlorate, 1588
 Bis(dimethylthallium)acetylide, 2519
 Bis(2,4-dinitrophenyl) disulfide, 3449
 1,2-Bis(diphenylphosphino)ethanepalladium(II) perchlorate, 3864
 Bis(dipropylborino)acetylene, 3670
 1,2-Bis-(di-2-propylphosphino)ethane, 3673
 1,4-Bis(2,3-epoxypropoxy)butane, 3353
 2,2-Bis[4(2',3'-epoxypropoxy)phenyl]propane, 3825
 1,2-Bisethoxycarbonyldiazene, *see* Diethyl azoformate, 2425
 Bis(ethoxycarbonyldiazomethyl)mercury, 2978
 † Bis(2-ethoxyethyl) ether, 3076
 1,2-Bis(ethylammonio)ethane perchlorate, 2589
 Bis(fluorobenzene)chromium(0), 3477
 Bis(fluorobenzene)vanadium(0), 3478
 Bis(2-fluoro-2,2-dinitroethoxy)dimethylsilane, 2421
 Bis(2-fluoro-2,2-dinitroethyl)amine, 1459
 Bis(fluoroformyl) peroxide, 0625
 1,1-Bis(fluoroxy)hexafluoropropane, 1061
 2,2-Bis(fluoroxy)hexafluoropropane, 1062
 1,1-Bis(fluoroxy)tetrafluoroethane, 0641
 Bis-3-(2-furyl)acryloyl peroxide, 3640
 Bis(hexafluorobenzene)chromium(0), 3421
 Bis(hexafluorobenzene)cobalt(0), 3419
 Bis(hexafluorobenzene)iron(0), 3424

Bis(hexafluorobenzene)nickel(0), 3425
 Bis(hexafluorobenzene)titanium(0), 3426
 Bis(hexafluorobenzene)vanadium(0), 3427
 Bis(hydrazine)[butanedioato(2-)]cadmium, *see* Dihydrazinecadmium(II) succinate, 1741
 Bis(hydrazine)[butanedioato(2-)]nickel, *see* Dihydrazinenickel(II) succinate, 1760
 Bis(hydrazine)diperchloratochromium(III) perchlorate, 4128
 Bis(hydrazine)nickel perchlorate, 4069
 Bis(hydrazine)[propanedioato(2-)]cadmium, *see* Dihydrazinecadmium(II) malonate, 1324
 Bis(hydrazine)[propanedioato(2-)]nickel, *see* Dihydrazinenickel(II) malonate, 1328
 Bis(hydrazine)tin(II) chloride, 4070
 Bis(2-hydroperoxy-2-butyl) peroxide (Dioxybis(1-methylpropylhydroperoxide)), 3078
 Bis(1-hydroperoxycyclohexyl) peroxide, 3556
 Bis(2-hydroperoxy-4-methyl-2-pentyl) peroxide, 3566
 2,2-Bis(hydroperoxy)propane, 1287
 1,2-Bis(2-hydroperoxy-2-propyl)benzene, 3541
 1,4-Bis(2-hydroperoxy-2-propyl)benzene, 3542
 1,2-Bis(hydroxomercurio)-1,1,2,2-bis(oxydimercurio)ethane, 0708
 Bis(1-hydroxycyclohexyl) peroxide, 3553
 Bis(2-hydroxyethyl) ether, 1705
 Bis(2-hydroxyethyl)methylphosphine, 2035
 Bis(2-hydroxyethyl) terephthalate, 3526
 Bis(hydroxylamine)zinc chloride, 4067
 Bis(hydroxymethyl) peroxide, 0928
 2,2-Bis(4-hydroxyphenyl)propane, 3686
 Bis(4-hydroxyphenyl) sulfone, 3497
 Bis-*N*(imidosulfurdifluoridato)mercury, 4342
 Bis(iodomethyl)zinc, 0805
 1,3-Bis(isocyanomethyl)benzene, 3257
 1,4-Bis(isocyanomethyl)cyclohexane, 3330
 1,4-Bis(isothiuronio)-2-butene dichloride, 2536
 Bis-(2-methoxyethyl) ether, 2549
 Bis(2-methoxyethyl) peroxydicarbonate, 3037
 Bis(1-methylbenzotriazole)cobalt(II) nitrate, 3652
 Bis(methylcyclopentadienyl)peroxoniobium chloride, 3522
 Bis(μ_3 -methylidyne) triangulo tris(pentamethylcyclopentadienylrhodium), 3879
 Bis(methyl 1-methylhydrazinocarbodithioate *N*²,*S'*)(perchlorato-*O*,*O'*) copper(1+)perchlorate, 2576
 Bis(2-methyl-1,8-naphthyridine)tetracarbonyldirhodium(I) perchlorate, 3833
 Bis(2-methylpyridine)sodium, 3524
 Bismuth, 0226

Bismuth amide oxide, 0229
 'Bismuth chromate', *see* Dibismuth dichromium nonaoxide, 0232
 Bismuthic acid, 0228
 Bismuth nitride, 0230
 Bismuth oxide hydroxide, *see* Bismuthic acid, 0228
 Bismuth pentafluoride, 0227
 Bismuth trioxide, 0233
 Bismuth trisulfide, 0234
 Bis(1,8-naphthyridine)tetracarbonyldirrhodium(I) perchlorate, 3791
 Bis(2-nitratoethyl)nitric amide, 1602
 Bis(4-nitrobenzenediazo) sulfide, 3467
I,I-Bis(4-nitrobenzoylperoxy)-4-chlorophenyl iodine, 3789
 1,1-Bis(4-nitrobenzoylperoxy)cyclohexane, 3800
I,I-Bis(4-nitrobenzoylperoxy)-2-methoxyphenyl iodine, 3819
I,I-Bis(4-nitrobenzoylperoxy)phenyl iodine, 3795
 2,2-Bis[(nitrooxy)methyl]propane-1,3-diol dinitrate, 1901
 Bis(2-nitrophenyl) disulfide, 3465
 1,2-Bis(2-nitrophenyl)ethane, 3646
 Bis(pentafluorophenyl)aluminium bromide, 3418
 Bis(pentafluorophenyl)ytterbium, 3423
 Bis(2,4-pentanedionato)chromium, 3326
 Bis-*O,N*[(*N'*-pent-2-en-2-oxy-4-ylidene)-*N,S*-dimethyldithiocarbazate]
 copper(II) perchlorate, 3718
 Bis(perchloryloxy)xenon, *see* Xenon(II) perchlorate, 4110
 Bis(2-phenyl-1,2-dicarbadodecaborane(12)-1-yl)diazene, 3721
 1,3-Bis(phenyltriazeno)benzene, 3757
 2,6-Bis(picrylazo)-3,5-dinitropyridine, 3734
 2,6-Bis(2-picrylhydrazino)-3,5-dinitropyridine, 3735
 Bis(2-propyloxy)diazene, *see* Diisopropyl hyponitrite, 2538
 Bis(propyloxy)diazene, *see* Dipropyl hyponitrite, 2539
 Bis(*O*-salicylidenaminopropylaziridine)iron(III) perchlorate, 3853
 Bis(tetramethyldiphosphane disulfide)cadmium perchlorate, 3102
 1,6-Bis(5-tetrazolyl)hexaaza-1,5-diene, 0827
 3,4-Bis(1,2,3,4-thiaziazol-5-ylthio)maleimide, 2073
 Bis(1,2,3,4-thiaziazol-5-ylthio)methane, 1084
 Bis(toluenediazo) oxide, 3654
 Bis(trichloroacetyl) peroxide, 1361
 Bis(2,4,5-trichlorobenzenediazo) oxide, 3431
 1,3-Bis(trichloromethyl)benzene, 2891
 Bis(trichlorophosphoranylidene)sulfamide, 4188
 1,2-Bis(triethylsilyl)hydrazine, 3583
 1,2-Bis(triethylsilyl)trimethylsilylhydrazine, 3696
 Bis(triethyltin) peroxide, 3581
 [*I,I*-Bis(trifluoroacetoxy)iodo]benzene, 3239
 Bis(trifluoroacetoxy)xenon, 1365
 Bis(trifluoroacetyl) peroxide, 1367

1,3-Bis(trifluoromethyl)benzene, 2894
 Bis(trifluoromethyl)cadmium, 0586
 Bis(trifluoromethyl)cadmium-1,2-dimethoxyethane adduct, 0587
 Bis(trifluoromethyl)chlorophosphine, 0597
 Bis(trifluoromethyl)cyanophosphine, 1053
 Bis(trifluoromethyl)dioxirane, 1055
 Bis(trifluoromethyl) disulfide, 0644
 1,3-Bis(trifluoromethyl)-5-nitrobenzene, 2885
 Bis(trifluoromethyl) nitroxide, 0638
 3,6-Bis(trifluoromethyl)pentacyclo [6.2.0.0^{2,4}.0^{3,6}.0^{5,7}]dec-9-ene, 3462
 Bis(trifluoromethyl)phosphorus(III) azide, 0639
 Bis(trifluoromethyl) sulfide, 0643
 Bis(trifluoromethyl)sulfur difluoride, 0648
 Bis(trifluoromethyl)trioxide, 0642
 Bis(trimethylphosphine)di(3,5-dibromo-2,6-dimethoxyphenyl)nickel, 3837
 Bis(trimethylphosphine)nickel(0)–acetylene complex, 3093
N,N-Bis(trimethylsilyl)aminoborane, 2612
cis-Bis(trimethylsilylamino)tellurium tetrafluoride, 2615
 Bis(trimethylsilyl) chromate, 2591
 1,2-Bis(trimethylsilyl)hydrazine, 2616
 Bis(trimethylsilyl)mercury, 2593
 Bis(trimethylsilylmethyl)magnesium, 3099
 † Bis(trimethylsilyl) oxide, 2601
 Bis(trimethylsilyl) peroxomonosulfate, 2602
 Bis(trimethylsilyl) phosphonite, 2611
N,N'-Bis(2,2,2-trinitroethyl)urea, 1875
 Bis(triperchloratosilicon) oxide, 4190
 1,2-Bis(triphenylphosphonio)ethane permanganate, 3899
 1,2-Bis(tripropylsilyl)hydrazine, 3777
 Biuret, *see* Imidodicarbonic diamide, 0874
 Borane, 0135
 Borane–ammonia, 0141
 Borane–bis(2,2-dinitropropylhydrazine), 0137
 Borane–dimethylsulfide, 0136
 Borane–hydrazine, 0142
 Borane–phosphorus trifluoride, 0140
 Borane–pyridine, 0139
 Borane–tetrahydrofuran, 0138
 Borazine, 0176
 Boric acid, 0144
 Boron, 0119
 Boron azide dichloride, 0126
 Boron azide diiodide, 0148
 Boron bromide diiodide, 0120
 Boron diiodophosphide, 0149
 Boron nitride, 0152

Boron phosphide, 0156
Boron triazide, 0153
Boron tribromide, 0122
Boron trichloride, 0127
Boron trifluoride, 0129
Boron trifluoride diethyl etherate, 1674
Boron triiodide, 0150
Boron trioxide, 0170
Boron trisulfide, 0172
Bromamide, *see* Bromamine, 0249
Bromamine, 0249
Bromic acid, 0248
Bromine, 0261
Bromine azide, 0256
Bromine bromate, 0276
Bromine dioxide, 0258
Bromine fluoride, 0238
Bromine(I) trifluoromethanesulfonate, 0312
Bromine pentafluoride, 0243
Bromine perchlorate, 0235
Bromine trifluoride, 0241
Bromine trioxide, 0259
N-Bromoacetamide, 0784
Bromoacetone oxime, 1201
Bromoacetylene, 0651
4-Bromoaniline, 2296
trans-Bromoazido(meso-5,12-dimethyl-1,4,8,11-tetraazacyclotetradeca-4,11-diene)cobalt(III) perchlorate, 3559
1-Bromoaziridine, 0783
Bromobenzene, 2224
4-Bromobenzenediazonium salts, 2131
4(4'-Bromobenzoyl)acetanilide, 3681
4-Bromobenzoyl azide, 2644
† 1-Bromobutane, 1631
† 2-Bromobutane, 1632
† 1-Bromo-2-butene, 1548
† 4-Bromo-1-butene, 1549
Bromochloroacetylene, 0578
2-Bromocyclohexanone, 2392
4-Bromocyclopentene, 1884
1-Bromo-1,2-cyclotridecadien-4,8,10-triylne, 3599
4-Bromo-1,2-diaminobenzene, 2345
Bromodiborane, 0159
2-Bromo-3,5-dimethoxyaniline, 2974
4-Bromodimethylaniline, 2973
Bromodimethylborane, 0887

2-Bromo-4,6-dinitroaniline, 2132
 4-Bromo-2,6-dinitroaniline, 2133
 6-Bromo-2,4-dinitrobenzenediazonium hydrogen sulfate, 2097
 3-Bromo-2,7-dinitrobenzo[b]thiophene-5-diazonium-4-oxide, 2883
 † Bromoethane, 0846
 † Bromoethylene, 0727
 † 2-Bromoethyl ethyl ether, 1635
 Bromo(5-fluoro-2-methylphenyl)magnesium, *see* 5-Fluoro-2-methylphenyl-
 magnesium bromide, 2709
 Bromoform, 0368
 Bromogermane, 0246
 Bromo-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetra-
 decadieneiron(II) perchlorate, 3722
 † Bromomethane, 0429
 4-Bromomethylbenzoic acid, 2926
 † 1-Bromo-3-methylbutane, 1984
 3-(Bromomethyl)-1,3-dibromopropane, *see* 1,1,1-Tris(bromomethyl)methane,
 1550
 2-Bromomethylfuran, 1840
 2-Bromomethyl-5-methylfuran, 2346
 † 1-Bromo-2-methylpropane, 1633
 † 2-Bromo-2-methylpropane, 1634
 2-Bromo-4-methylpyridine *N*-oxide, 2297
 3-Bromo-3(4-nitrophenyl)-3*H*-diazirine, 2645
N-Bromo-3-nitrophthalimide, 2884
 1-Bromo-2-oximinopropane, *see* Bromoacetone oxime, 1201
 3-Bromo-2-oxotetrahydrothiophene, *see* α -Bromo- γ -thiobutyrolactone, 1450
 1-Bromopentaborane(9), 0186
 † 2-Bromopentane, 1985
 α -(4-Bromophenylazo)benzyl hydroperoxide, 3606
 α -(4-Bromophenylazo)phenylethyl α -hydroperoxide, 3648
 2-(4-Bromophenylazo)-2-propyl hydroperoxide, 3156
 3-Bromophenyllithium, 2129
 4-Bromophenyllithium, 2130
 2-(4-Bromophenyl)-2-propyl 1-(1,1-dimethyl-2-pentyn-4-enyl) peroxide, 3709
 † 1-Bromopropane, 1241
 † 2-Bromopropane, 1242
 † 3-Bromo-1-propene, 1153
 Bromo-2-propenylzinc, *see* Allylzinc bromide, 1155
 † 3-Bromopropyne, 1090
 3-Bromopyridine, 1827
 3-Bromopyridine *N*-oxide, 1828
 5-Bromo-4-pyrimidinone, 1410
 1-Bromo-2,5-pyrrolidinedione, *see* *N*-Bromosuccinimide, 1425
 † Bromosilane, 0250
N-Bromosuccinimide, 1425

- N*-Bromotetramethylguanidine, 2002
 α -Bromo- γ -thiobutyrolactone, 1450
 Bromotrichloromethane, 0310
 3-Bromo-1,1,1-trichloropropane, 1126
 † Bromotrifluoroethylene, 0579
 Bromotrifluoromethane, 0311
N-Bromotrimethylammonium bromide(?), 1300
 2-Bromo-2,5,5-trimethylcyclopentanone, 3018
 Bromotrimethyl-*N,N,N',N'*-tetramethylethanediamine-*N,N'*-palladium, *see*
N,N,N',N'-Tetramethylethane-1,2-diamine, trimethylpalladium(IV)
 bromide, 3225
 Bromyl fluoride, 0239
 Buckminsterfullerene, 3916
 † 1,2-Butadiene, 1479
 † 1,3-Butadiene, 1480
 † 1,3-Butadiyne, 1385
 4,4'-(Butadiyne-1,4-diyl)bis(2,2,6,6-tetramethyl-4-hydroxypiperidine-1-oxyl),
 3838
 † Butanal, *see* Butyraldehyde, 1607
 † Butane, 1668
 Butane-1,4-diisocyanate, 2376
 1,4-Butanedinitrile, *see* Succinodinitrile, 1433
 † Butane-2,3-dione, 1525
 Butane-2,3-dione dioxime, 1595
 2,3-Butanedione monoxime, 1570
 † 2-Butanethiol, 1713
 † Butanethiol, 1712
 Butanoic acid, *see* Butyric acid, 1614
 1-Butanol, 1694
 † 2-Butanol, 1695
 † *tert*-Butanol, 1696
 † 2-Butanone, 1606
 2-Butanone oxime, 1654
 2-Butanone oxime hydrochloride, 1678
 † Butanonitrile, *see* Butyronitrile, 1563
 † Butanoyl chloride, *see* Butyryl chloride, 1555
 † 2-Butenal, *see* Crotonaldehyde, 1516
 † 1-Butene, 1577
 † *cis*-2-Butene, 1578
 † *trans*-2-Butene, 1579
trans-2-Butene-1,4-dioic acid, *see* Fumaric acid, 1446
trans-2-Butene ozonide, 1623
 † 1-Buten-3-one, 1515
 † 2-Butenonitrile, *see* 1-Cyanopropene, 1464
 † 3-Butenonitrile, *see* 3-Cyanopropene, 1465
 2-Buten-1-yl benzenesulfonate, 3316

- But-2-en-2-ylboronic acid, 1630
 2-Buten-1-yl diazoacetate, 2377
- † Buten-3-yne, 1423
 3-Buten-1-ynyldiethylaluminium, 3017
 3-Buten-1-ynyldiisobutylaluminium, 3549
 3-Buten-1-nyltriethyllead, 3357
 Butoxyacetylene, 2428
- † 2-(2-Butoxyethoxy)ethanol, 3077
 1-Butoxyethyl 3-trimethylplumbylpropionate, 3552
- † 2-Butyl acetate, 2497
- † Butyl acetate, 2496
tert-Butyl 1-adamantaneperoxy-carboxylate, 3692
N-Butylamidosulfuryl azide, 1693
- † 2-Butylamine, 1724
- † Butylamine, 1723
- † *tert*-Butylamine, 1725
 Butyl azide, *see* 1-Azidobutane, 1666
N-Butyl-*N*-2-azidoethylnitramine, 2528
tert-Butyl azidoformate, 1936
 Butylbenzene, 3323
- † *n*-Butyl bromide, *see* 1-Bromobutane, 1631
- † *sec*-Butyl bromide, *see* 2-Bromobutane, 1632
- † *tert*-Butyl bromide, *see* 2-Bromo-2-methylpropane, 1634
- † *n*-Butyl chloride, *see* 1-Chlorobutane, 1637
- † *sec*-Butyl chloride, *see* 2-Chlorobutane, 1638
- † *tert*-Butyl chloride, *see* 2-Chloro-2-methylpropane, 1640
tert-Butyl chloroperoxyformate, 1926
tert-Butyl diazoacetate, 2423
tert-Butyl 2-diazoacetoacetate, 3009
 5-*tert*-Butyl-3-diazo-3*H*-pyrazole, 2831
tert-Butyldibromamine, *see* 1,1-Dimethylethyldibromamine, 1636
 Butyldichloroborane, 1629
tert-Butyldifluorophosphine, 1646
N-tert-Butyl-3,3-dinitroazetidine, *see* *N*-Dimethylethyl-3,3-dinitroazetidine,
 2848
tert-Butyl diperoxyoxalate, 3356
O-O-tert-Butyl diphenyl monoperoxophosphate, 3712
O-O-tert-Butyl di(4-tolyl) monoperoxophosphate, 3763
- † Butylethylamine, 2565
- † Butyl ethyl ether, 2540
- † *tert*-Butyl ethyl ether, 2541
N-tert-Butylformamide, 1994
- † Butyl formate, 1967
O-O-tert-Butyl hydrogen monoperoxymaleate, 3015
tert-Butyl hydroperoxide, 1698
tert-Butyl hypochlorite, 1642

- 3-*N*-*tert*-Butyliminioisobenzofuranone tetrafluoroborate, *see*
N-*tert*-Butylphthalisoimidium tetrafluoroborate, 3521
- † *sec*-Butyl iodide, *see* 2-Iodobutane, 1647
- † *tert*-Butyl iodide, *see* 2-Iodo-2-methylpropane, 1649
- 4-*tert*-Butyliodylbenzene, 3319
- (*tert*-Butyl isobutyl ether), *see* 1-(1,1-Dimethylethoxy)-2-methylpropane,
 3072
- † Butyl isocyanate, 1931
- O*-*O*-*tert*-Butyl isopropyl monoperoxycarbonate, 3051
- Butyllithium, 1651
- tert*-Butyllithium, 1652
- Butylmagnesium chloride, 1641
- Butyl 4-methylbenzenesulfonate, *see* Butyl toluenesulfonate, 3409
- † Butyl methyl ether, 2010
- † *tert*-Butyl methyl ether, 2011
- Butyl nitrate, 1661
- † Butyl nitrite, 1658
- † *tert*-Butyl nitrite, 1659
- tert*-Butylnitroacetylene, *see* 3,3-Dimethyl-1-nitro-1-butyne, 2395
- tert*-Butyl 4-nitroperoxybenzoate, 3400
- tert*-Butyl peroxophosphoryl dichloride, 1644
- tert*-Butyl peroxyacetate, 2509
- † *tert*-Butyl peroxybenzoate, 3401
- tert*-Butyl peroxynitrate, 1663
- 2-*tert*-Butyl-3-phenyloxaziridine, 3406
- N*-*tert*-Butylphthalisoimidium tetrafluoroborate, 3521
- Butylsodium, 1667
- Butyl toluenesulfonate, 3409
- N*-*tert*-Butyl-*N*-trimethylsilylaminoborane, 2878
- 5-*tert*-Butyl-2,4,6-trinitro-1,3-xylene, 3532
- † Butyl vinyl ether, 2484
- † 1-Butyne, 1481
- † 2-Butyne, 1482
- † 2-Butynedinitrile, *see* Dicyanoacetylene, 1801
- 2-Butyne-1,4-diol, 1526
- 1-Butyne-4-nitrile, *see* 3-Cyanopropyne, 1416
- 2-Butyne-1-thiol, 1546
- 3-Butyn-1-yl 4-toluenesulfonate, 3399
- † Butyraldehyde, 1607
- Butyraldehyde oxime, 1655
- Butyric acid, 1614
- Butyrolactone, 1527
- † Butyronitrile, 1563
- † Butyryl chloride, 1555
- Butyryl nitrate, 1573
- Butyryl peroxonitrate, 1574

Cadmium, 3949
Cadmium amide, 3954
Cadmium azide, 3957
Cadmium chlorate, 3952
Cadmium cyanide, 0588
Cadmium fulminate, 0589
Cadmium hydrazinium tris(hydrazinecarboxylate) trihydrate, 1338
Cadmium hydride, 3953
Cadmium nitride, 3960
Cadmium oxide, 3958
Cadmium propionate, 2418
Cadmium selenide, 3959
Caesium, 4254
Caesium acetylide, 0614
Caesium amide, 4260
Caesium azide, 4262
Caesium bromoxenate, 0237
Caesium chloroxenate, 3970
Caesium cyanotridecahydrodecaborate(2-), 0518
Caesium fluoride, 4255
Caesium fluoroxysulfate, 4256
Caesium graphite, 2882
Caesium hexafluorobromate, 0236
Caesium hexahydroaluminate(3-), 0067
Caesium hydride, 4258
Caesium hydrogen xenate, 4259
Caesium lithium tridecahydroonaborate, 0194
Caesium nitrate, 4261
Caesium nitride, 4266
Caesium oxide, 4264
Caesium ozonide, *see* Caesium trioxide, 4263
Caesium pentacarbonylvandate(3-), 1811
Caesium pentafluorotelluramide, 4257
Caesium selenide, 4265
Caesium tetrafluorochlorate(1-), 3969
Caesium tetraperchloratoiodate, 4161
Caesium trioxide, 4263
Calcium, 3922
Calcium abietate, 3900
Calcium acetylide, 0585
Calcium azide, 3936
Calcium bis(*O*-hydroxylamide), 3930
Calcium bromide, 0262
Calcium carbide, *see* Calcium acetylide, 0585
Calcium carbonate, 0317
Calcium chloride, 3923

Calcium chlorite, 3925
 Calcium chromate, 3926
 Calcium cyanamide, 0316
 Calcium disilicide, 3944
 Calcium hydrazide, 3932
 Calcium hydride, 3927
 Calcium hydroxide, 3928
 Calcium hydroxide *O*-hydroxylamide, 3929
 Calcium hypochlorite, 3924
 Calcium hypophosphite, *see* Calcium phosphinate, 3931
 Calcium 2-iodylbenzoate, 3625
 Calcium nitrate, 3935
 Calcium nitride, 3946
 Calcium oxide, 3937
 Calcium 2,4-pentanedionate, 3325
 Calcium permanganate, 3934
 Calcium peroxide, 3938
 Calcium peroxodisulfate, 3940
 Calcium phosphide, 3947
 Calcium phosphinate, 3931
 Calcium picrate, 3430
 Calcium polysulfide, 3942
 Calcium silicide, 3943
 Calcium stearate, 3896
 Calcium sulfate, 3939
 Calcium sulfide, 3941
 Calcium tetrahydroborate, 0158
 Calcium triperoxochromate, 3945
 Caprolactam, 2451
 Carbamide, *see* Urea, 0475
 † 2-Carbamoyl-2-nitroacetonitrile, 1116
N-Carboethoxyiminophosphoryl chloride, 1170
N-Carbomethoxyiminophosphoryl chloride, 0743
 Carbon, 0298
 Carbon dioxide, 0557
 † Carbon disulfide, 0560
 Carbonic diazide, 0550
 Carbonic dihydrazide, 0508
 † Carbon monoxide, 0555
 † Carbon oxide sulfide, *see* Carbonyl sulfide, 0556
 ‘10-Carbon oxime’ 3362
 † ‘Carbon suboxide’, *see* Propadiene-1,3-dione, 1349
 Carbon subsulfide, *see* Propadienedithione, 1350
 Carbon sulfide, 0559
 Carbon tetraboride, 0309
 Carbon tetrabromide, 0315

Carbon tetrachloride, 0332
 Carbon tetrafluoride, 0349
 Carbon tetraiodide, 0525
 Carbonyl azide, *see* Carbonic diazide, 0550
 Carbonyl-bis(triphenylphosphine)iridium–silver diperchlorate, 3898
 Carbonyl dichloride, 0329
 Carbonyl dicyanide, *see* Oxopropanedinitrile, 1341
 Carbonyl difluoride, 0343
 Carbonyl diisothiocyanate, 1342
 Carbonyl(pentasulfur pentanitrido)molybdenum, 0535
 † Carbonyl sulfide, 0556
 Carboxybenzenesulfonyl azide, 2699
 2-Carboxy-3,6-dimethylbenzenediazonium chloride, 3138
 Carboxymethyl carbamimioniothioate chloride, *see* S-Carboxymethyl-
 isothiuronium chloride, 1245
 S-Carboxymethylisothiuronium chloride, 1245
 ‘Carbyl sulfate’, *see* 2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide, 0839
 Δ3-Carene, 3336
 Ceric ammonium nitrate, Ammonium cerium(IV) nitrate, *see* Ammonium
 hexanitrocerate, 3964
 Cerium, 3961
 Cerium azide, 3966
 Cerium carbide, 0590
 Cerium dihydride, 3962
 Cerium(III) tetrahydroaluminate, 0089
 Cerium nitride, 3965
 Cerium trihydride, 3963
 Cerium trisulfide, 3967
 Cetyltrimethylammonium permanganate, *see* Hexadecyltrimethylammonium
 permanganate, 3785
 Chloral hydrate, *see* 2,2,2-Trichloro-1,1-ethanediol, 0745
 Chloramide, *see* Chloramine, 4000
 Chloramine, 4000
 ‘Chloramine B’, *see* Sodium N-chlorobenzenesulfonamide, 2229
 Chloratomercurio(formyl)methylenemercury(II), 0654
 Chloric acid, 3996
 Chlorimide, *see* Dichloramine, 4063
 3-Chlorimino-1-phenylpropene, *see* N-Chlorocinnamaldimin, 3126
 Chloriminovanadium trichloride, 4165
 Chlorine, 4047
 Chlorine azide, 4030
 Chlorine dioxide, 4042
 Chlorine dioxygen trifluoride, 3983
 Chlorine fluoride, 3971
 Chlorine fluorosulfate, 3975
 Chlorine(I) trifluoromethanesulfonate, 0320

- 'Chlorine nitrate', *see* Nitryl hypochlorite, 4026
- Chlorine pentafluoride, 3989
- Chlorine perchlorate, 4101
- Chlorine trifluoride, 3981
- Chlorine trifluoride oxide, 3982
- Chlorine trioxide, 4044
- Chloroacetaldehyde oxime, 0787
- N*-Chloroacetamide, 0789
- Chloroacetamide, 0788
- Chloroacetamide oxime, *see* 2-Chloro-*N*-hydroxyacetimidine, 0849
- † Chloroacetone, 1161
- 4-Chloroacetophenone, 2929
- Chloroacetylene, 0652
- † 2-Chloroacrylonitrile, 1074
- † *N*-Chloroallylamine, 1202
- 4-Chloro-2-aminophenol, 2303
- † *N*-Chloro-3-aminopropene, *see* *N*-Chloroallylamine, 1202
- N*-Chloro-3-aminopropyne, 1129
- 2-Chloroaniline, 2300
- 3-Chloroaniline, 2301
- 4-Chloroaniline, 2302
- 1-Chloroaziridine, 0786
- Chlorobenzene, 2228
- 2-Chloro-1,4-benzenediamine, 2349
- 4-Chloro-1,2-benzenediamine, 2347
- 4-Chloro-1,3-benzenediamine, 2348
- 2-Chlorobenzenediazonium salts, 2148
- 3-Chlorobenzenediazonium salts, 2149
- 4-Chlorobenzenediazonium triiodide, 2137
- 4-Chlorobenzenesulfonyl azide, 2152
- 1-Chlorobenzotriazole, 2150
- 4-Chlorobenzoyl azide, 2654
- 2-Chlorobenzylidenemalononitrile, 3333
- N*-Chloro-bis(2-chloroethyl)amine, 1590
- 2-Chloro-1,1-bis(fluoroxy)trifluoroethane, 0595
- N*-Chlorobis(trifluoromethanesulfonyl)imide, 0596
- † 2-Chloro-1,3-butadiene, 1451
- † 1-Chlorobutane, 1637
- † 2-Chlorobutane, 1638
- 1-Chloro-2-butanone, 1556
- 4-Chlorobutanonitrile, *see* 4-Chlorobutyronitrile, 1487
- † 2-Chloro-2-butene, 1551
- † 3-Chloro-1-butene, 1552
- 1-Chloro-1-buten-3-one, 1454
- 4-Chloro-2-butyne, 1455
- 4-Chlorobutyronitrile, 1487

(Chlorocarbonyl)imidodisulfur difluoride, 0318
N-(Chlorocarbonyloxy)trimethylurea, 1924
 2-Chloro-5-chloromethylthiophene, 1830
N-Chlorocinnamaldimin, 3126
 Chlorocycloacetylene, 1036
 2-Chloro-1-cyanoethanol, 1130
trans-2-Chlorocyclohexyl perchlorate, 2420
 Chloro(η^8 1,3,5,7-cyclooctatetraene)(tetrahydrofuran)lutetium, 3536
 † Chlorocyclopentane, 1923
 3-Chlorocyclopentene, 1885
 4-Chloro-2,6-diamino-*N*-methylaniline, 2822
 Chlorodiborane, 0160
 Chlorodibutylborane, 3065
 4-Chloro-2,5-diethoxynitrobenzene, 3301
 Chlorodiethylborane, 1673
 Chlorodifluoroacetyl hypochlorite, 0603
 † 1-Chloro-1,1-difluoroethane, 0731
 Chlorodifluoromethane, 0369
 1-Chloro-3,3-difluoro-2-methoxycyclopropene, 1411
 1-Chlorodifluoromethyl-1-trifluoromethyldioxirane, 1033
N-Chlorodimethylamine, 0895
B-Chlorodimethylaminodiborane, 0962
 1-Chloro-3-dimethylaminopropane, 2003
 Chlorodimethylarsine, 0884
 5-Chloro-1,3-dimethyl-4-nitro-1*H*-pyrazole, 1864
 Chlorodimethylphosphine, 0899
N-Chloro-4,5-dimethyltriazole, 1490
 2-Chloro-4,6-dinitroaniline, 2153
 4-Chloro-2,6-dinitroaniline, 2154
 1-Chloro-2,4-dinitrobenzene, 2098
 6-Chloro-2,4-dinitrobenzenediazonium hydrogen sulfate, 2102
 4-Chloro-2,5-dinitrobenzenediazonium 6-oxide, 2066
 2-Chloro-1,3-dinitro-5-trifluoromethylbenzene, 2637
 3-Chloro-1,3-diphenylcyclopropene, 3679
 Chlorodipropylborane, 2534
 1-Chloro-2,3-di(2-thienyl)cyclopropenium perchlorate, 3388
 † 1-Chloro-2,3-epoxypropane, 1162
 † Chloroethane, 0848
 † 2-Chloroethoxyethene, *see* 2-Chloroethyl vinyl ether, 1557
 2-Chloroethylamine, 0896
 2-Chloroethylamine hydrochloride, *see* 2-Chloroethylammonium chloride,
 0940
 2-Chloroethylammonium chloride, 0940
 Chloroethylbismuthine, 0892
 † Chloroethylene, 0730
N-(2-Chloroethyl)-*N*-nitrosocarbamoyl azide, 1131

- † 2-Chloroethyl vinyl ether, 1557
- † 2-Chlorofluorobenzene, 2134
- † 3-Chlorofluorobenzene, 2135
- † 4-Chlorofluorobenzene, 2136
- Chloroform, 0372
- Chloroformamidinium chloride, 0468
- Chloroformamidinium nitrate, 0467
- Chloroformamidinium perchlorate, 0469
- Chlorogermane, 3992
- Chloro-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetra-
decadieneiron(II) perchlorate, 3724
- 2-Chloro-*N*-hydroxyacetamide, 0849
- 2-Chloro-*N*-(2-hydroxyethyl)aniline, 2975
- 3-Chloro-2-hydroxypropyl perchlorate, 1207
- Chloroiodoacetylene, 0598
- Chloroiodoethyne, *see* Chloroiodoacetylene, 0598
- 3-Chloro-1-iodopropyne, 1073
- 3-Chlorolactonitrile, *see* 2-Chloro-1-cyanoethanol, 1130
- Chloromercuriodiphenylgold, 3473
- † Chloromethane, 0432
- 3-Chloro-4-methoxyaniline, 2795
- 3-Chloro-3-methoxydiazirine, 0734
- † Chloromethoxyethane, *see* Chloromethyl ethyl ether, 1246
- Chloro-(4-methoxyphenyl)diazirine, 2928
- 2-Chloro-5-methylaniline, 2793
- 4-Chloro-2-methylaniline, 2794
- 4-Chloro-2-methylbenzenediazonium salts, 2719
- † 1-Chloro-3-methylbutane, 1986
- † 2-Chloro-2-methylbutane, 1987
- 3-Chloro-3-methyldiazirine, 0733
- † Chloromethyl ethyl ether, 1246
- 2-Chloromethylfuran, 1841
- N*-Chloro-4-methyl-2-imidazolinone, 1554
- 4-Chloro-1-methylimidazolium nitrate, 1453
- † Chloromethyl methyl ether, 0850
- 2-Chloromethyl-5-methylfuran, 2352
- N*-Chloro-5-methyl-2-oxazolidinone, 1488
- † Chloromethyloxirane, Epichlorohydrin, *see* 1-Chloro-2,3-epoxypropane, 1162
- 4-Chloro-2-methylphenol, 2741
- 2-Chloro-5-methylphenylhydroxylamine, 2796
- Chloromethylphenylsilane, 2810
- † 1-Chloro-2-methylpropane, 1639
- † 2-Chloro-2-methylpropane, 1640
- † 3-Chloro-2-methyl-1-propene, 1553
- 2-Chloromethylthiophene, 1842
- N*-Chloro-3-morpholinone, 1489

N-Chloro-4-nitroaniline, 2231
 2-Chloro-5-nitrobenzaldehyde, 2649
 2-Chloro-6-nitrobenzaldehyde, 2650
 4-Chloro-3-nitrobenzaldehyde, 2651
 5-Chloro-2-nitrobenzaldehyde, 2652
 2-Chloronitrobenzene, 2141
 4-Chloronitrobenzene, 2142
 1-(4-Chloro-2-nitrobenzeneazo)-2-hydroxynaphthalene, 3697
 2-Chloro-5-nitrobenzenesulfonic acid, 2144
 4-Chloro-3-nitrobenzenesulfonic acid, 2145
 2-Chloro-5-nitrobenzyl alcohol, 2715
 4-Chloro-2-nitrobenzyl alcohol, 2716
 4-Chloro-3-nitrobenzyl alcohol, 2717
 5-Chloro-2-nitrobenzyl alcohol, 2718
 6-Chloro-2-nitrobenzyl bromide, 2671
 2-Chloro-4-nitrobenzyl chloride, 2678
 4-Chloro-2-nitrobenzyl chloride, 2679
 Chloronitromethane, 0396
 2-Chloro-1-nitro-4-nitroso-2-butene, 1452
 1-Chloro-4-(2-nitrophenyl)-2-butene, 3270
N-(4-Chloro-2-nitrophenyl)-1,2-diaminoethane, 2977
 1-Chloro-1-nitrosocyclohexane, 2419
 2-Chloro-1-nitroso-2-phenylpropane, 3147
 2-Chloro-4-nitrotoluene, 2711
 4-Chloro-3-nitrotoluene, 2712
 Chloronium perchlorate, 4065
 † 1-Chloropentane, 1988
 3-Chloroperoxybenzoic acid, 2676
 Chloroperoxytrifluoromethane, 0319
 2(3-Chlorophenoxy)ethylamine, 2976
 1-(4-Chlorophenyl)biguanidinium hydrogen dichromate, 3005
 3-(4-Chlorophenyl)butanoic acid, 3297
 2-(4-Chlorophenyl)-1,1-dimethylethyl hydroperoxide, 3018
 5-*p*-Chlorophenyl-2,2-dimethyl-3-hexanone, 3664
 4-Chlorophenyl isocyanate, 2648
 3-Chlorophenyllithium, 2138
 4-Chlorophenyllithium, 2139
 1-Chloro-3-phenylpent-1-en-4-yn-3-ol, 3402
 2-(4-Chlorophenyl)-2-propyl 1-(1,1-dimethyl-2-pentyn-4-enyl) peroxide, 3711
N-Chloro-5-phenyltetrazole, 2674
 'Chloropicrin', *see* Trichloronitromethane, 0331
N-Chloropiperidine, 1948
 † Chloroprene, *see* 2-Chloro-1,3-butadiene, 1451
 † 1-Chloropropane, 1243
 † 2-Chloropropane, 1244
 3-Chloro-1,2-propanediol, 1248

- † Chloro-2-propanone, *see* Chloroacetone, 1161
- N*-Chloropropargylamine, *see N*-Chloro-3-aminopropyne, 1129
- † 1-Chloro-1-propene, 1156
- † 2-Chloropropene, 1157
- † 3-Chloropropene, 1158
- 2-Chloro-2-propenyl trifluoromethanesulfonate, 1426
- 1-Chloro-2,3-propylene dinitrate, 1160
- 1-Chloro-2-propyl perchlorate, 1206
- † 3-Chloropropyne, 1092
- Chloropropyne nitrile, *see* Chlorocyanoacetylene, 1036
- 2-Chloro-3-pyridinediazonium tetrafluoroborate, 1816
- 2-Chloropyridine *N*-oxide, 1829
- N*-Chloropyrrolidine, 1584
- 1-Chloro-2,5-pyrrolidinedione, *see N*-Chlorosuccinimide, 1427
- N*-Chlorosuccinimide, 1427
- N*-Chlorosulfinylimide, 4024
- Chlorosulfonyl isocyanate, 0324
- Chlorosulfuric acid, 3997
- Chlorotetrafluorophosphorane, 3987
- N*-Chlorotetramethylguanidine, 2004
- 5-Chloro-1,2,3-thiadiazole, 0655
- 5-Chlorotoluene-2-diazonium tetrachlorozincate, 3644
- N*-Chlorotoluene-4-sulfonamide, 2797
- 3-Chloro-3-trichloromethyldiazirine, 0610
- † Chlorotrifluoroethylene, 0591
- 4-Chlorotrifluoromethylbenzene, 2646
- 2-Chloro-5-trifluoromethylbenzenediazonium hydrogen sulfate, 2647
- 3-Chloro-3-trifluoromethyldiazirine, 0593
- † 1-Chloro-3,3,3-trifluoropropane, 1127
- 2-Chloro-*N,N,N'*-trifluoropropionamidine, 1128
- † Chlorotrimethylsilane, 1304
- Chloro-1,2,4-triselenadiazolium chloride, 4089
- 3-Chloro-1,3,4-triselenadiazolium chloride, *see* Chloro-1,2,4-triselenadiazolium chloride, 4089
- 2-Chlorovinyltrimethyllead, 1989
- Chloryl hypofluorite, 3973
- Chloryl perchlorate, 4104
- Chromic acid, 4229
- Chromium, 4222
- Chromium(II) acetate, 1493
- Chromium(II) chloride, 4052
- Chromium(III) chloride, 4127
- Chromium(III) oxide, 4251
- Chromium(III) perchlorate . 6dimethyl sulfoxide, 4130
- Chromium(II) oxide, 4241
- Chromium(II) sulfate, 4244

Chromium(II) sulfide, 4245
Chromium nitride, 4237
Chromium pentafluoride, 4226
Chromium trioxide, 4242
Chromocene, *see* Dicyclopentadienylchromium, 3273
Chromyl acetate, 1494
Chromyl azide, 4239
Chromyl azide chloride, 3968
Chromyl chloride, 4054
'[(Chromyldioxy)iodo]benzene', *see* Phenyliodine(III) chromate, 2247
Chromyl fluorosulfate, 4225
Chromyl isocyanate, 0613
Chromyl isothiocyanate, 0612
Chromyl nitrate, 4238
Chromyl perchlorate, 4055
Cinnamaldehyde, 3134
Citral, *see* 3,7-Dimethyl-2,6-octadienal, 3346
Citric acid, 2389
Citronellic acid, 3352
Cobalt, 4199
Cobalt(II) acetate tetrahydrate, *see* Diacetatotetraquocobalt, 1780
Cobalt(II) azide, 4216
Cobalt(II) bromide, 0263
Cobalt(II) chelate of bi(1-hydroxy-3,5-diphenylpyrazol-4-yl *N*-oxide), 3872
Cobalt(II) chelate of 1,3-bis(*N*-nitrosohydroxylamino)benzene, 2164
Cobalt(II) chloride, 4048
Cobalt(III) amide, 4201
Cobalt(III) chloride, 4122
Cobalt(III) nitride, 4214
Cobalt(III) oxide, 4221
Cobalt(II) nitrate, 4215
Cobalt(II) oxide, 4217
Cobalt(II) perchlorate hydrates, 4051
Cobalt(II) picramate, 3460
Cobalt(II) sulfide, 4218
Cobalt trifluoride, 4200
Cobalt tris(dihydrogenphosphide), 4203
Columbium, 3948
Copper, 4267
Copper bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide), 3861
Copper chromate oxide, 4223
Copper diphosphide, 4284
Copper(I) azide, 4287
Copper(I) benzene-1,4-bis(ethynide), 3236
Copper(I) bromide, 0265
Copper(I) chloride, 4056

Copper(I) chloroacetylide, 1354
 Copper(I) cyanide, 0620
 Copper(I) hydride, 4286
 Copper(II) acetylide, 0615
 Copper(II) azide, 4280
 Copper(II) azide hydroxide, 4270
 Copper(II) cyanide, 0616
 Copper(II) 3,5-dinitroanthranilate, 3627
 Copper(II) 1,3-di(5-tetrazolyl)triazene, 1429
 Copper(II) fulminate, 0617
 Copper(II) glycinate nitrate, 0905
 Copper(II) nitrate, 4279
 Copper(II) oxide, 4281
 Copper(II) perchlorate, 4057
 Copper(II) phosphide, 4290
 Copper(II) phosphinate, 4272
 Copper(II) picrate, 3432
 Copper(II) sulfate, 4282
 Copper(II) sulfide, 4285
 Copper(II) thiocyanate, 0618
 Copper(I) nitride, 4289
 Copper(I) oxalate, 0622
 Copper(I) oxide, 4288
 Copper(I) perchlorate, 4058
 Copper iron(II) sulfide, 4269
 Copper(I) tetrahydroaluminate, 0068
 Copper monophosphide, 4283
 Copper 1,3,5-octatrien-7-ynide, 2933
 Copper 1,3,5-octatrien-7-ynide, 3704
 Copper–zinc alloys, 4268
 † Crotonaldehyde, 1516
 ‘18-Crown-6’, *see* 1,4,7,10,13,16-Hexaoxacyclooctadecane, 3561
 ‘Crown ethers’, *see* Cyclic poly(ethylene oxides), 0830
 ‘CS gas’, *see* 2-Chlorobenzylidenemalononitrile, 3333
 Cubane-1,2-dicarboxylic acid, *see* Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane-1,2-dicarboxylic acid, 3261
 ‘Cupferron’, *see* Ammonium *N*-nitrosophenylaminoxide, 2399
 Cyanamide, 0404
 Cyanoacetic acid, 1113
 Cyanoacetyl chloride, 1075
 Cyanoborane oligomer, 0393
N-Cyano-2-bromoethylbutylamine, 2846
N-Cyano-2-bromoethylcyclohexylamine, 3178
 Cyanocyclopropane, 1463
 Cyanodiazocetyl azide, 1346
 5-Cyano-4-diazo-4*H*-1,2,3-triazole, 1345

- Cyanodimethylarsine, 1199
- † 2-Cyanoethanol, 1181
- N*-(2-Cyanoethyl)ethanolamine, *see* 3-(2-Hydroxyethylamino)propionitrile, 1952
- N*-(2-Cyanoethyl)ethylhydroxylamine, 1951
- 3-Cyanoethylhydrazine, *see* 3-Hydrazinopropanenitrile, 1275
- Cyanoforn, 1383
- Cyanofornyl chloride, 0600
- Cyanogen azide, 0544
- Cyanogen bromide, 0313
- Cyanogen chloride, 0323
- Cyanoguanidine, 0813
- Cyanohydrazonoacetyl azide, 1083
- 5-Cyano-2-methyltetrazole, 1120
- Cyanonitrene, 0540
- 2-Cyano-4-nitrobenzenediazonium hydrogen sulfate, 2668
- 4-Cyano-3-nitrotoluene, 2917
- 3(3-Cyano-1,2,4-oxadiazol-5-yl)-4-cyanofurazan 2-(or 5-) oxide, 2630
- 2-Cyano-2-propanol, 1566
- † 1-Cyanopropene, 1464
- † 3-Cyanopropene, 1465
- 1-Cyano-2-propen-1-ol, 1467
- 2-Cyano-2-propyl nitrate, 1506
- 3-Cyanopropyne, 1416
- 2(5-Cyanotetrazole)pentaamminecobalt(III) perchlorate, 0974
- † Cyanotrimethylsilane, 1665
- 2-Cyano-1,2,3-tris(difluoroamino)propane, 1432
- † Cyanuric acid, *see* 2,4,6-Trihydroxy-1,3,5-triazine, 1118
- Cyanuric chloride, *see* 2,4,6-Trichloro-1,3,5-triazine, 1038
- Cyclic poly(ethylene oxides), 0830
- Cyclo-*oligo*-butadiynyl-2,3-dioxocycloutenylidene, *see* Oligo(octacarbon-dioxide), 3108
- † Cyclobutane, 1580
- 1,2-Cyclobutanedione, 1440
- † Cyclobutene, 1483
- Cyclobutyl 4-methylbenzenesulfonate, 3404
- Cyclobutylmethyl methanesulfonate, 2510
- cis*-Cyclododecene, 3351
- † Cycloheptane, 2849
- † 1,3,5-Cycloheptatriene, 2790
- † Cycloheptene, 2837
- † 1,3-Cyclohexadiene, 2361
- † 1,4-Cyclohexadiene, 2362
- Cyclohexadiene-1,4-dione, *see* 1,4-Benzoquinone, 2214
- 2,5-Cyclohexadienone-4-imine, *see* 1,4-Benzoquinone monoimine, 2260
- † Cyclohexane, 2457

- 1,2,3-Cyclohexanetrione trioxime, 2400
 1,3,5-Cyclohexanetrione trioxime, 2401
 Cyclohexanol, 2485
 Cyclohexanone, 2429
 Cyclohexanone oxime, 2452
 † Cyclohexene, 2406
 2-Cyclohexenyl hydroperoxide, 2435
 † Cyclohexylamine, 2521
 1,7,13-Cyclooctadecatriene-3,5,9,11,15,17-hexayne, 3740
 1,5-Cyclooctadiene-bis(4-chloropyridine *N*-oxide)rhodium(I) perchlorate,
 3761
 † 1,3,5,7-Cyclooctatetraene, 2943
 (η^8 1,3,5,7-Cyclooctatetraene)[2-(dimethylaminomethyl)phenyl-*C,N*](tetra-
 hydrofuran)lutetium, 3826
 1,3,5-Cyclooctatriene, 2966
 Cyclopentaazathienium chloride, 4034
 † Cyclopentadiene, 1857
 Cyclopentadiene–silver perchlorate, 1859
 Cyclopentadienyldimethylborane, 2833
 Cyclopentadienylgold(I), 1839
 μ -Cyclopentadienyl(methyl)-bis-(*N*-methyl-*N*-nitrosohydroxylamino)titanium,
 3028
 Cyclopentadienylsodium, 1855
 μ -Cyclopentadienyltrimethyltitanium, 3038
 † Cyclopentane, 1939
 4,5-Cyclopentanofurazan-*N*-oxide, 1871
 Cyclopentanone, 1905
 Cyclopentanone oxime, 1932
 † Cyclopentene, 1891
 † Cyclopentene oxide, 1906
 † Cyclopentylamine, 1991
 Cyclopentylmethyl methanesulfonate, 2867
 † Cyclopropane, 1197
 Cyclopropanecarbonitrile, *see* Cyanocyclopropane, 1463
 † Cyclopropylamine, 1256
 † Cyclopropyl methyl ether, 1608
 † Cyclopropyl methyl ketone, 1907
 1,7,13,19-Cyclotetracosatetraene-3,5,9,11,15,17,21,23-octayne, 3843
 Cyclotrimethylenetrinitramine, *see* 1,3,5-Trinitrohexahydro-1,3,5-triazine,
 1219
 Cyclotrisiloxane, *see* 2,4,6-Trisilatrioxane, 4566

 Decaborane(14), 0198
 Decacarbonyldirhenium, 3386
 Decachlorotetrasilane, 4197
 Decafluorobutyramidine, 1371

Decafluoro-2,5-diazahehexane 2,5-dioxyl, 1372

† Decane, 3365

Decaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]heptaconta-4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62,67,69-eicosayne, 3920

Decyl nitrite, 3364

Dehydrofurfural oxime, *see* 2,2'-Oxybis(iminomethylfuran) mono-*N*-oxide, 3260

† Deuterium oxide, 4291

2-Deuterobicyclo[2.2.1]hept-2-ene, 2811

Devarda's alloy, *see* Aluminium-copper-zinc alloy, 0050

9,10 Dewar anthracene, *see* 2,3:5,6-Dibenzobicyclo[3.3.0]hexane, 3633

Diacetatoplatinum(II) nitrate, 1510

Diacetatotetraaquocobalt, 1780

Diacetonitrile-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetra-decadieneiron(II) perchlorate, 3813

1,3-Diacetoxybenzene, 3291

N,N'-Diacetyl-*N,N'*-dinitro-1,2-diaminoethane, 2426

Diacetyl peroxide, 1537

† Diallylamine, 2450

† Diallyl ether, 2431

Diallyl peroxydicarbonate, 2989

Diallyl phosphite, 2456

Diallyl sulfate, 2443

Diallyl sulfide, 2447

Dialuminium octavanadium tridecasilicide, 0088

Diamidophosphorous acid, 4547

1,4-Diaminobenzene, 2371

4,4'-Diaminobiphenyl, *see* Benzidine (4,4'-Diaminobiphenyl), 3516

1,2-Diaminoethane, 0953

1,2-Diaminoethanebis(trimethylgold), 3105

Diaminoguanidinium nitrate, 0514

1,2-Diamino-2-methylpropaneaquadiperoxochromium(IV), 1781

1,2-Diamino-2-methylpropaneoxodiperoxochromium(VI), 1746

† 1,2-Diaminopropane, 1325

† 1,3-Diaminopropane, 1326

3,6-Diamino-1,2,4,5-tetrazine-1,4-dioxide, 0822

3,6-Diamino-1,2,4,5-tetrazine-1-oxide, 0821

1,11-Diamino-3,6,9-triazaundecane, 3101

Diammineboronium heptahydrotetraborate, 0190

Diammineboronium tetrahydroborate, 0167

Diamminedichloroaminotrichloroplatinum(IV), 4179

Diamminedihydroxyosmium, 4572

Diamminenitratocobalt(II) nitrate, 4202

Diamminepalladium(II) nitrate, 4562

Diamminepalladium(II) nitrite, 4559

cis-Diammineplatinum(II) nitrite, 4560
Diamminesilver permanganate, 0018
trans-Diammine(1,4,8,11-tetraazacyclotetradecane)chromium(III) perchlorate,
3383
1,2-Diammonioethane nitrate, 0966
Diammonium Aquabis(peroxotartatovanadate)(2-), 3079
1,4,3,6-Dianhydroglucitol 2-nitrate, 2396
1,4:3,6-Dianhydro-D-glucitol dinitrate, *see* Isosorbide dinitrate, 2380
Dianilineoxodiperoxochromium(VI), 3523
Dianilinium dichromate, 3537
o-Dianisidine, *see* 3,3'-Dimethoxy-4,4'-diaminobiphenyl, 3660
Diaquabis(cyclopentadienyl)titanium dichromate, 3327
2,3-Diazabicyclo[2.2.2]octa-2,5-diene *N*-oxide, 2374
1,4-Diazabicyclo[2.2.2]octane, 2473
1,4-Diazabicyclo[2.2.2]octane 1,4-bis(nitroimidate), 2480
1,4-Diazabicyclo[2.2.2]octane hydrogen peroxidate, 2474
1,6-Diaza-3,4,8,9,12,13-hexaoxabicyclo[4.4.4]tetradecane, 2476
Diazene, 4469
1,3-Diazidobenzene, 2207
1,4-Diazidobenzene, 2208
Diazido-bis(1,2-diaminoethane)ruthenium(II) hexafluorophosphate, 1789
2,3-Diazido-1,3-butadiene, 1437
2,2-Diazidobutane, 1603
1,2-Diazidocarbonylhydrazine, 0720
Diazido[(1,2,5,6- η)-1,5-Cyclooctadiene]platinum, 3012
2,5-Diazido-3,6-dichlorobenzoquinone, 2054
Diazidodichlorosilane, 4092
Diazidodicyanomethane, *see* Diazidomalnonitrile, 1347
2,4-Diazido-6-dimethylamino-1,3,5-triazine, 1876
Diazidodimethylsilane, 0918
1,12-Diazido-3,10-dioxa-5,8-dinitrazadodecane, 3049
Diazidodiphenylmethane, 3603
1,1-Diazidoethane, 0818
1,2-Diazidoethane, 0819
Diazidomalnonitrile, 1347
Diazidomethane, 0414
Diazidomethyleneazine, 1021
Diazidomethylenecyanamide, 1017
1,3-Diazido-2-nitroazapropane, 0824
1,1-Diazidophenylethane, 2953
1,2-Diazido-1-phenylethane, 2954
2,2-Diazidopropane, 1216
1,3-Diazidopropene, 1143
2,3-Diazidopropionitrile, 1121
2,6-Diazidopyrazine, 1400
Diazido(*N,N,N,N*-tetramethylethanediamine)palladium, 2581

Diazido(trifluoromethyl)arsine, 0307
Diazirine, 0405
Diazirine-3,3-dicarboxylic acid, 1080
Diazoacetaldehyde, 0710
Diazoacetone, 0675
Diazoacetyl azide, 0679
'Diazoaminomethane', *see* 1,3-Dimethyltriazene, 0945
2-Diazocyclohexanone, 2375
Diazocyclopentadiene, 1832
Diazodicyanoimidazole, *see* 2-Diazonio-4,5-dicyanoimidazolidine, 2050
Diazodicyanomethane, *see* Diazomalononitrile, 1344
2-Diazo-2,6-dihydro-6-oxo-1,3-azulenedicarbonitrile, *see* 1,3-Dicyano-2-diazo-2,6-azulene quinone, 3434
2-Diazo-2*H*-imidazole, 1082
1-Diazoindene, 3116
Diazomalonic acid, 1081
Diazomalononitrile, 1344
Diazomethane, 0406
5-(Diazomethylazo)tetrazole, 0719
Diazomethyldimethylarsine, 1240
5-(4-Diazoniobenzenesulfonamido)thiazole tetrafluoroborate, 3119
2-Diazonio-4,5-dicyanoimidazolidine, 2050
1-(4-Diazoniophenyl)-1,2-dihydropyridine-2-iminosulfinate, 3392
1(2'-Diazoniophenyl)2-methyl-4,6-diphenylpyridinium diperchlorate, 3847
3-Diazoniopyrazolidine-4-carboxamide, 1399
3-Diazoniopyrazolidine-4-carboxamide, 1422
5-Diazoniotetrazolidine, 0548
3-Diazo-5-phenyl-3*H*-pyrazole, 3118
3-Diazo-5-phenyl-3*H*-1,2,4-triazole, 2909
4-Diazo-5-phenyl-1,2,3-triazole, 2910
† 3-Diazopropene, 1135
Diazopropyne, 1070
 α -Diazo- γ -thiobutyrolactone, 1434
3-Diazo-3*H*-1,2,4-triazole, 0677
4-Diazo-1,2,3-triazole, 0678
Dibenzoylsulfonyl peroxide, 3499
2,3:5,6-Dibenzobicyclo[3.3.0]hexane, 3633
Dibenzocyclododeca-1,7-dienetetrayne, 3786
Dibenzo-d,f-1,2-dioxocin-3,8-dione, *see* 2,2-Biphenyldicarbonyl peroxide, 3631
1,5-Dibenzoylnaphthalene, 3845
Dibenzoyl peroxide, 3639
'Dibenzyl chlorophosphate', *see* Dibenzyl phosphorochloridate, 3650
Dibenzyl ether, 3655
Dibenzyl phosphite, 3658
Dibenzyl phosphorochloridate, 3650

1,3-Di[bis(cyclopentadienyl)iron]-2-propen-1-one, 3840
 Di-1,2-bis(difluoroaminoethyl) ether, 1497
 Dibismuth dichromium nonaoxide, 0232
 Di[bis(trifluoromethyl)phosphido]mercury, 1375
 † Diborane(6), 0166
 Diboron oxide, 0169
 Diboron tetrachloride, 0161
 Diboron tetrafluoride, 0162
 Dibromine pentoxide, 0277
 Dibromine trioxide, *see* Bromine bromate, 0276
 Dibromoacetylene, 0582
 2,4-Dibromoaniline, 2226
 2,6-Dibromobenzoquinone-4-chloroimide, 2076
 Dibromoborylphosphine, 0121
 3,5-Dibromo-7-bromomethylene-7,7a-dihydro-1,1-dimethyl-1*H*-
 azirino[1,2-*a*]indole, 3474
 1,4-Dibromo-1,3-butadiyne, 1351
 3,5-Dibromocyclopentene, 1860
 1,2-Dibromo-1,2-diisocyanatoethane polymers, 1387
 2,2-Dibromo-1,3-dimethylcyclopropanoic acid, 2364
 4,4-Dibromo-3,5-dimethylpyrazole, 1861
 2,3-Dibromo-5,6-epoxy-7,8-dioxabicyclo[2.2.2]octane, 2298
 1,2-Dibromoethane, 0785
 Dibromogermane, 0267
 Dibromomethane, 0395
N,N-Dibromomethylamine, 0430
 Dibromomethylborane, 0426
 1,2-Dibromopentaborane(9), 0187
N,N'-Dibromosulfurdiimide, 0272
 Di-2-butenylcadmium, 3027
 1,2-Di(3-buten-1-ynyl)cyclobutane, 3509
 Di-2-butoxyethyl ether, 3564
 † Di-2-butylamine, 3084
 Dibutylamine, 3083
 3-Dibutylaminopropylamine, 3416
 2,2'-Di-*tert*-butyl-3,3'-bioxaziridine, 3359
N,N'-Di-*tert*-butyl-*N,N'*-bis(trimethylsilyl)diaminophosphene, 3675
 Di-*tert*-butyl chromate, 3066
 2,6-Di-*tert*-butyl-4-cresol, 3691
 Di-*tert*-butyl diazomalonate, 3411
 Di-*tert*-butyl diperoxycarbonate, 3194
 Di-*tert*-butyl diperoxyphthalate, 3715
 † Dibutyl ether, 3071
 Di(*O-O-tert*-butyl) ethyl diperoxophosphate, 3374
 Di-*tert*-butylfluorophosphine, 3067
 Dibutyl hydrogen phosphite, 3086

Dibutyl hyponitrite, 3069
trans-Di-*tert*-butyl hyponitrite, 3070
 Dibutylmagnesium, 3068
 Dibutyl-3-methyl-3-buten-1-ynborane, 3618
 2,6-Di-*tert*-butyl-4-nitrophenol, 3666
 † Di-*tert*-butyl peroxide, 3074
 2,2-Di(*tert*-butylperoxy)butane, 3565
 Dibutyl phthalate, 3714
 2,4-Di-*tert*-butyl-2,2,4,4-tetrafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine, 3377
 Dibutylthallium isocyanate, 3193
 Dibutylzinc, 3080
 Di-3-camphoroyl peroxide, 3807
 Dicarbonyl- π -cycloheptatrienyltungsten azide, 3123
cis-Dicarbonyl(cyclopentadienyl)cyclooctenemanganese, 3689
 Dicarbonylmolybdenum diazide, 0995
 Dicarbonyl(phenanthroline *N*-oxide)rhodium(I) perchlorate, 3626
 Dicarbonylpyrazinerhodium(I) perchlorate, 2147
 Dicarbonyl- η -trichloropropenyldinickel chloride dimer, 3233
 Dicarbonyltungsten diazide, 1013
 Dichloramine, 4063
 Dichlorodifluoromethane, 0326
 Dichlorine heptaoxide, *see* Perchloryl perchlorate, 4107
 Dichlorine oxide, 4095
 Dichlorine trioxide, 4100
 Dichloroacetylene, 0602
 † *N,N*-Dichloro- β -alanine, 1168
 † 2-Dichloroaminopropanoic acid, *see N,N*-Dichloro- β -alanine, 1168
 1-Dichloroaminotetrazole, 0371
N,N-Dichloroaniline, 2242
 2,3-Dichloroaniline, 2237
 2,4-Dichloroaniline, 2238
 2,5-Dichloroaniline, 2239
 2,6-Dichloroaniline, 2240
 3,4-Dichloroaniline, 2241
 1,2-Dichlorobenzene, 2156
N,N-Dichlorobenzenesulfonamide, 2244
 2,6-Dichlorobenzoquinone-4-chloroimide, 2078
cis-Dichlorobis(2,2'-bipyridyl)cobalt(III) chloride, 3799
trans-Dichlorobis[1,2-phenylenebis(dimethylarsine)]palladium(IV) diperchlorate, 3808
N,N'-dichlorobis(2,4,6-trichlorophenyl)urea, 3596
 Dichloroborane, 0124
 1,4-Dichloro-1,3-butadiyne, 1353
 † *miso*-Dichlorobutane, 1587
 1,4-Dichloro-2-butyne, 1428

- Dichlorodi- μ -chlorobis(pentamethylcyclopentadienyl)dirhodium, 3806
- † Dichlorodiethylsilane, 1683
- 3,4-Dichloro-2,5-dilithiothiophene, 1357
- 2,2-Dichloro-3,3-dimethylbutane, 2469
- 1,3-Dichloro-5,5-dimethyl-2,4-imidazolidindione, 1865
- † Dichlorodimethylsilane, 0902
- 3,6-Dichloro-3,6-dimethyltetraoxane, 1492
- 1,5-Dichloro-2,4-dinitrobenzene, 2077
- Dichlorodinitromethane, 0328
- Dichlorodisilylaminoborane, 0125
- † 1,1-Dichloroethane, 0790
- † 1,2-Dichloroethane, 0791
- 2,2-Dichloroethylamine, 0853
- Dichloroethylborane, 0845
- † 1,1-Dichloroethylene, 0695
- † *cis*-1,2-Dichloroethylene, 0696
- † *trans*-1,2-Dichloroethylene, 0697
- 1,1-Dichloroethyl hydroperoxide, 0794
- 1,1-Dichloroethyl peroxyacetate, 1491
- † Dichloroethylsilane, 0903
- Dichloroethyne, *see* Dichloroacetylene, 0602
- Dichlorofluoramine, 4059
- 1,1-Dichloro-1-fluoroethane, 0738
- (2,2-Dichloro-1-fluorovinyl)ferrocene, 3470
- N,N*-Dichloroglycine, 0739
- 1,6-Dichloro-2,4-hexadiyne, 2157
- 4,5-Dichloro-3,3,4,5,6,6-hexafluoro-1,2-dioxane, 1355
- Dichloro-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(III) perchlorate, 3725
- Dichloromaleic anhydride, 1359
- Dichloromaleimide, 1378
- † Dichloromethane, 0397
- N,N*-Dichloromethylamine, 0437
- Dichloromethylarsine, 0424
- Dichloromethylenediethylammonium chloride, 1950
- Dichloromethylenedimethylammonium chloride, 1209
- Dichloromethylphosphine, 0438
- † Dichloromethylsilane, 0470
- † Dichloromethylvinylsilane, 1208
- 2,4,-Dichloronitrobenzene, 2104
- 3,4-Dichloronitrobenzene, 2105
- Dichloronitrobenzene, 2103
- 2,6-Dichloro-4-nitrobenzenediazonium hydrogen sulfate, 2106
- N,N*-Dichloropentafluorosulfanylamine, 4060
- Dichlorophenol mixed isomers, 2161
- 4-(2,4-Dichlorophenoxy)butyric acid, 3272

Dichlorophenylborane, 2222
 3,4-Dichlorophenylhydroxylamine, 2243
 † 1,1-Dichloropropane, 1203
 † 1,2-Dichloropropane, 1204
 2,2-Dichloropropane, 1205
 † 2,3-Dichloropropene, 1132
 3,4-Dichloro-2,5-pyrrolidinedione, *see* Dichloromaleimide, 1378
 † Dichlorosilane, 4066
 1,2-Dichlorotetrafluoroethane, 0604
 1,2-Dichlorotetramethyldisilane, 1743
trans-Dichlorotetrapyridinecobalt(III) chloride, 3802
 endo-2,5-Dichloro-7-thiabicyclo[2.2.1]heptane, 2369
 Dicobalt boride, 0128
 Dicopper(I) acetylide, 0619
 Dicopper(I) 1,5-hexadiynide, 2165
 Dicopper(I) ketenide, 0621
 Dicrotonoyl peroxide, 2986
 Dicumenechromium(0), 3767
 † Dicyanoacetylene, 1801
 1,4-Dicyano-2-butene, 2311
 'Dicyanodiamide', *see* Cyanoguanidine, 0813
 Dicyanodiazene, 1005
 1,3-Dicyano-2-diazo-2,6-azulene quinone, 3434
 Dicyanofurazan, 1803
 Dicyanofurazan *N*-oxide, 1804
 Dicyanofuroxan, *see* Dicyanofurazan *N*-oxide, 1804
 † Dicyanogen, 0996
 Dicyanogen *N,N'*-dioxide, 0998
 Dicyanoiodonium triflate, 1048
 Dicyclohexylcarbonyl peroxide, 3667
 † Dicyclopentadiene, 3299
 Dicyclopentadienefurazan *N*-oxide, *see* 4a,5,7a,8-Tetrahydro-4,8-methano-4*H*-indeno[5,6-*c*]-1,2,5-oxadiazole 1-oxide, 3281
 Dicyclopentadienefurazan *N*-oxide, *see* 4a,5,7a,8-Tetrahydro-4,8-methano-4*H*-indeno[5,6-*c*]-1,2,5-oxadiazole 3-oxide, 3282
 Dicyclopentadienylchromium, 3273
 Dicyclopentadienylperoxyniobium chloride, 3271
 Dicyclopropyldiazomethane, 2824
 Dideuterodiazomethane, 0336
 Didodecanoyl peroxide, 3857
 Didodecyl peroxydicarbonate, 3865
 † 1,2:3,4-Diepoxbutane, 1507
 † 1,2:3,4-Diepoxbutane, 1529
 2,3:5,6-Diepox-7,8-dioxabicyclo[2.2.2]octane, 2339
 Diethoxydiazene, *see* Diethyl hyponitrite, 1690
 † Diethoxydimethylsilane, 2582

- 1,1-Diethoxy-1,3-disiladioxetane, 1762
- † 1,1-Diethoxyethane, 2545
- † 1,2-Diethoxyethane, 2546
- † Diethoxymethane, 2016
- † 3,3-Diethoxypropene, 2858
- † Diethyl acetal, *see* 1,1-Diethoxyethane, 2545
- Diethyl acetylenedicarboxylate, 2987
- Diethylaluminium bromide, 1670
- Diethylaluminium chloride, 1671
- Diethylaluminium hydride, 1719
- Diethylaluminium 2,2,2-trifluoroethoxide, 2466
- † Diethylamine, 1726
- 1-Diethylamino-1-buten-3-yne, 3020
- S-Diethylamino(methylimino)sulfur(IV) fluoride, 2027
- 3-Diethylaminopropylamine, 2877
- Diethylaminosulfinyl chloride, 1679
- Diethylaminosulfur trifluoride, 1684
- 2-Diethylammonioethyl nitrate nitrate, 2571
- Diethylarsine, 1720
- Diethyl azoformate, 2425
- Diethylberyllium, 1675
- Diethylbismuth chloride, 1676
- Diethylcadmium, 1677
- † Diethyl carbonate, 1981
- Diethyl chlorophosphate, *see* Diethyl phosphorochloridate, 1681
- Diethyl diazomalonate, 2828
- Diethyl dicarbonate, 2444
- Diethyl-3-diethylaminopropylaluminium, 3414
- Diethyl 1,4-dihydro-2,6-dimethylpyridine-3,5-dicarboxylate, 3615
- 3,6-Diethyl-3,6-dimethyl-1,2,4,5-tetraoxane, 3052
- Di[*N,N'*-Ethylenebis(2-oxidoacetophenoneiminato)copper(II)]
oxovanadium(IV) diperchlorate, 3893
- Diethylene glycol, *see* Bis(2-hydroxyethyl) ether, 1705
- † Diethyleneglycol monobutyl ether, *see* 2-(2-Butoxyethoxy)ethanol, 3077
- Diethylenetriamine, *see* Bis(2-aminoethyl)amine, 1777
- Diethyl ethanephosponite, 2572
- † Diethyl ether, 1697
- Diethylethoxyaluminium, 2554
- Diethylgallium hydride, 1721
- Diethylgold bromide, 1672
- Diethylhydrogallium, *see* Diethylgallium hydride, 1721
- Diethylhydroxytin hydroperoxide, 1763
- Diethyl hyponitrite, 1690
- Diethylketene, 2430
- Diethyllead dinitrate, 1692
- Diethylmagnesium, 1687

Diethylmethylphosphine, 2036
 Diethyl 4-nitrophenyl phosphate, 3329
 Diethyl 4-nitrophenyl thionophosphate, 3328
 † 3,3-Diethylpentane, 3199
 Diethyl peroxide, 1699
 Diethyl peroxydicarbonate, 2446
 Diethylphosphine, 1734
N-(Diethylphosphinoyl)hydroxylamine, 1752
 Diethyl phosphite, 1733
 Diethyl phosphorochloridate, 1681
 Diethyl succinate, 3031
 Diethyl sulfate, 1710
 † Diethyl sulfide, 1714
 Diethyl sulfite, 1709
 Diethyl telluride, 1717
 Diethylthallium perchlorate, 1682
 Diethyl trifluoroacetosuccinate, 3320
 Diethylzinc, 1718
 1,4-Diethynylbenzene, 3241
cis-3,4-Diethynylcyclobut-3-ene-1,2-diol, 2921
 11,12-Diethynyl-9,10-dihydro-9,10-ethenoanthracene, 3792
 2,4-Diethynyl-5-methylphenol, 3393
 2,4-Diethynylphenol, 3244
 11,12-Diethynyl-1,4,7,10-tetraoxadispiro[4.0.4.2]dodec-11-ene, 3496
 4,4-Diferrocenylpentanoyl chloride, 3859
 Difluoramine, 4311
 Difluoroacetic acid, 0702
 Difluoroacetylene, *see* Difluoroethyne, 0623
 Difluoroacetyl hypochlorite, 0653
 (Difluoroamino)difluoroacetonitrile, 0630
 2-(Difluoroamino)-2,2-dinitroacetamide, 0701
 2-(Difluoroamino)-2,2-dinitroacetonitrile, 0624
 2-(Difluoroamino)-2,2-dinitroethane, 0753
 (Difluoroaminodinitromethyl)benzene, *N,N*-Difluoro- α,α -dinitrobenzene-
 methanamine, *see* α -(Difluoroamino)- α,α -dinitrotoluene, 2680
 α -(Difluoroamino)- α,α -dinitrotoluene, 2680
 3-Difluoroamino-1,2,3-trifluorodiaziridine, 0358
 Difluoroammonium hexafluoroantimonate, 4380
 Difluoroammonium hexafluoroarsenate, 0098
 † 1,3-Difluorobenzene, 2168
 † 1,4-Difluorobenzene, 2169
 Difluorodiazene, 4314
 Difluorodiazirine, 0342
 3,3-Difluoro-1,2-dimethoxycyclopropene, 1866
 1,5-Difluoro-2,4-dinitrobenzene, 2080
 † 1,1-Difluoroethane, 0801

- † 1,1-Difluoroethylene, 0700
- 1,1-Difluoroethylene ozonide, 0703
- Difluoroethyne, 0623
- Difluoro-*N*-fluoromethanimine, 0344
- Difluoromethylene dihypofluorite, 0355
- 2,4-Difluoronitrobenzene, 2109
- 3,4-Difluoro-2-nitrobenzenediazonium 6-oxide, 2067
- 3,6-Difluoro-2-nitrobenzenediazonium 4-oxide, 2068
- Difluoroperchloryl hexafluoroarsenate, 0093
- Difluoroperchloryl hexafluoroplatinate, 3991
- Difluoroperchloryl tetrafluoroborate, 0123
- N,N*-Difluorotrifluoromethylamine, 0357
- Difluorotrifluoromethylphosphine, 0361
- Difluorotrifluoromethylphosphine oxide, 0360
- 3,3-Difluoro-1,2,4-trioxolane, *see* 1,1-Difluoroethylene ozonide, 0703
- Difluorotrioxenon, 4325
- 1,1-Difluorourea, 0398
- Di-2-furoyl peroxide, 3245
- Digallane, 4410
- † Digermane, 4420
- Digold(I) ketenide, 0574
- Dihexanoyl peroxide, 3554
- † Dihexyl ether, 3563
- Dihydrazinecadmium(II) malonate, 1324
- Dihydrazinecadmium(II) succinate, 1741
- Dihydrazinecobalt(II) chlorate, 4049
- Dihydrazinemanganese(II) nitrate, 4570
- Dihydrazinenickel(II) malonate, 1328
- Dihydrazinenickel(II) succinate, 1760
- Dihydrazinesilver nitrate, 0019
- 6,6'-Dihydrazino-2,2'-bipyridylnickel(II) perchlorate, 3303
- 2,4-Dihydro-1*H*-1,2,6a-methanopentalene, *see* Tetracyclo[4.3.0^{3.5}.0^{4.6}]
nona-1,7-diene, 3125
- 1,4-Dihydrodicyclopropa[*b,g*]naphthalene, 3452
- 4,5-Dihydro-4,4-dimethyl-3,5,5-triphenyl-3*H*-pyrazol-3-yl hydroperoxide, *see*
3,3,5-Triphenyl-4,4-dimethyl-5-hydroperoxy-4,5-dihydro(3*H*)pyrazole,
3841
- † 2,3-Dihydrofuran, 1517
- 2,5-Dihydrofuran-2,5-dione, *see* Maleic anhydride, 1404
- 2(3*H*)-Dihydrofuranone, *see* Butyrolactone, 1527
- 3,6-Dihydro-1,2,2*H*-oxazine, 1567
- 5,8-Dihydro-2-methyl-5,8-ethano-1*H*-[1,2,4]triazolo[1,2*a*] pyridazine-
1,3(2*H*)dione, *see* 4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2.6}]undeca-
8-ene-3,5-dione, 3160
- 1,2-Dihydroperoxy-1,2-bis(benzeneazo)cyclohexane, 3762
- Di(hydroperoxy)trimethylantimony(V), 1333

- † 2,3-Dihydropyran, 1908
- 1,2-Dihydropyridazine-3,6-dione, *see* Maleic hydrazide, 1435
- 1,2-Dihydropyrido[2,1-e]tetrazole, 1834
- 2,3-Dihydroxybutanedioic acid, *see* Tartaric acid, 1545
- Dihydroxybutenedioic acid, *see* Dihydroxymaleic acid, 1447
- Dihydroxymaleic acid, 1447
- 1,8-Dihydroxy-2,4,5,7-tetranitroanthraquinone, 3621
- 1,4-Diimido-2,5-cyclohexadiene, *see* 1,4-Benzoquinone diimine, 2310
- Diiodamine, 4426
- Diiodoacetylene, 0985
- 1,2-Diiodobenzene, 2172
- 1,4-Diiodo-1,3-butadiyne, 1799
- Diiodoethyne, *see* Diiodoacetylene, 0985
- 1,6-Diiodo-2,4-hexadiyne, 2173
- Diiodomethane, 0400
- N,N'*-Diiodosulfurdiimide, 4626
- Di- μ -iodotetranitrosyldiiron, 4402
- Diisobutylaluminium chloride, 3064
- Diisobutylaluminium hydride, 3082
- † Diisobutylamine, 3085
- Diisobutylzinc, 3081
- Diisobutyryl peroxide, 3032
- 1,6-Diisocyanatohexane, 3008
- † Diisocyanatomethane, 1079
- 2,4-Diisocyanatotoluene, 3117
- 1,4-Diisocyanobutane, 2372
- 1,6-Diisocyanohexane, 3006
- Diisocyanomethane, 1077
- 1,8-Diisocyanooctane, 3343
- 1,5-Diisocyanopentane, 2825
- 1,3-Diisocyanopropane, 1869
- Diisopentylmercury, 3368
- Diisopentylzinc, 3371
- † Diisopropylamine, 2566
- Diisopropylberyllium, 2535
- † Diisopropyl ether, 2542
- Diisopropyl hyponitrite, 2538
- Diisopropyl malonate, 3188
- Diisopropyl peroxydicarbonate, 3034
- † Diketene, 1441
- Dilauroyl peroxide, *see* Didodecanoyl peroxide, 3857
- Dilead(II)lead(IV) oxide, 4861
- 1,3-Dilithiobenzene, 2178
- 1,4-Dilithiobenzene, 2179
- Dilithioferrocene, *see* $\mu,1,1'$ -Ferrocenediyldilithium, 3256
- Dilithium 1,1-bis(trimethylsilyl)hydrazide, 2595

Dilithium μ -cyclooctatetraene, 2948
 Dimercury dicyanide oxide, 0983
 1,10-Di(methanesulfonyloxy)deca-4,6-diyne, 3543
 Dimethanesulfonyl peroxide, 0931
 2,4-Dimethoxyaniline, 2998
 3,5-Dimethoxyaniline, 2999
 3,3'-Dimethoxy-4,4'-diaminobiphenyl, 3660
 Dimethoxydiazene, *see* Dimethyl hyponitrite, 0913
mixo-Dimethoxydinitroanthraquinone, 3701
 † 1,1-Dimethoxyethane, 1700
 † 1,2-Dimethoxyethane, 1701
 † Dimethoxymethane, 1281
 1,6-Di(4'-methoxyphenyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane, 3797
 † 1,1-Dimethoxypropane, 2017
 † 2,2-Dimethoxypropane, 2018
 † 3,3-Dimethoxypropene, 1968
N,N-Dimethylacetamide, 1656
 Dimethyl 4-acetamidophthalate, 3520
 Dimethyl 4-acetamino-5-nitrophthalate, 3515
 Dimethyl acetylenedicarboxylate, 2340
 Dimethylaluminium bromide, 0882
 Dimethylaluminium chloride, 0883
 Dimethylaluminium hydride, 0936
 † Dimethylamine, 0941
 4-Dimethylaminoacetophenone, 3321
 (Dimethylamino)acetylene, 1564
 5-(4-Dimethylaminobenzeneazo)tetrazole, 3161
 Dimethylaminobis(trifluoromethyl)borane, 1485
 Dimethylaminodiborane, 0967
 2-Dimethylaminoethanol, *see* *N*-2-Hydroxyethyl dimethylamine, 1729
 † 2-Dimethylaminoethylamine, 1755
 2-Dimethylamino-3-methylbenzoyl azide, 3308
 † 2-Dimethylamino-*N*-methylethylamine, 2038
 Dimethylaminomethylferrocene, 3613
 2-(Dimethylaminomethyl)fluoroferrrocene, 3612
 3-Dimethylaminopropylamine, 2039
 † Dimethylaminotrimethylsilane, 2043
 Dimethylammonium perchlorate, 0950
N,N-Dimethylaniline, 2993
 Dimethylantimony chloride, 0900
 Dimethylarsine, 0937
 Dimethyl azoformate, 1528
 3,5-Dimethylbenzenediazonium-2-carboxylate, 3128
 4,6-Dimethylbenzenediazonium-2-carboxylate, 3129
 2,4-Dimethylbenzenediazonium triiodide, 2963
 1,3-Dimethyl-2,1-benzisoxazolium perchlorate, 3148

3,5-Dimethylbenzoic acid, 3152
 Dimethylberyllium, 0890
 Dimethylberyllium-1,2-dimethoxyethane, 0891
 6,6-Dimethylbicyclo[3.1.1]heptan-2-one, 3175
 Dimethylbismuth chloride, 0893
 3,5-Dimethyl-4-[*I,I*-bis(trifluoroacetoxy)iodo]isoxazole, 3115
N,N-Dimethyl-1,1-bis(trifluoromethyl)boranamine, *see* Dimethylaminobis
 (trifluoromethyl)borane, 1485
 † 1,2-Dimethylborane, 0961
 2,3-Dimethyl-1,3-butadiene, 2407
 † 2,2-Dimethylbutane, 2529
 † 2,3-Dimethylbutane, 2530
 † 3,3-Dimethyl-2-butanone, 2486
 † 1,3-Dimethylbutylamine, 2567
 3,3-Dimethyl-1-butyne, 2408
 Dimethyl 2-butyndioate, *see* Dimethyl acetylenedicarboxylate, 2340
 Di-2-methylbutyryl peroxide, 3354
 Dimethylcadmium, 0894
 † Dimethyl carbonate, 1233
N,N-Dimethyl-2-chloroacetamide, 1585
 Dimethyl 2-chloro-4-nitrophenyl thionophosphate, 2960
 Dimethyl 3-chloro-4-nitrophenyl thionophosphate, 2961
 † Dimethylcyclohexanes, 3039
 † Dimethylcyclohexanes 5 isomers, *see* Dimethylcyclohexanes, 3039
 1,2-Dimethylcyclopentene ozonide, 2845
 † 1,1-Dimethylcyclopropane, 1940
 Dimethyldiazene, *see* Azomethane, 0910
 Dimethyl diazenedicarboxylate, *see* Dimethyl azoformate, 1528
 1,1-Dimethyldiazanium perchlorate, 0938
 † 1,1-Dimethyldiborane, 0960
 Dimethyl *N,N*-dichlorophosphoramidate, 0901
 2,5-Dimethyl-2,5-dihydrofuran-2,5-endoperoxide, *see* 1,4-Dimethyl-2,3,7-
 trioxabicyclo[2.2.1]hept-5-ene, 2388
 2,5-Dimethyl-2,5-dihydrothiophene-2,5-endoperoxide, 2387
 3,4-Dimethyl-4-(3,4-dimethyl-5-isoxazolylazo)isoxazolin-5-one, 3309
N,N'-Dimethyl-*N,N'*-dinitrooxamide, 1513
N,N'-Dimethyl-*N,N'*-dinitrosoxamide, 1512
N,N'-Dimethyl-*N,N'*-dinitrosoterephthalamide, 3284
 2,6-Dimethyl-1,3-dioxo-2,6-diphosphacyclooctane, 2548
 † 2,6-Dimethyl-1,4-dioxane, 2498
 3,3-Dimethyl-1,2-dioxetane, 1615
 Dimethyldioxirane, 1227
 † 2,2-Dimethyl-1,3-dioxolane, 1969
 1-(4,5-Dimethyl-1,3-diselenonylidene)piperidinium perchlorate, 3340
 † Dimethyl disulfide, 0934
 Di- μ -methylenebis(methylpentamethylcyclopentadienyl)dirhodium, 3855

- (*N,N*-Dimethylethanamine)trihydroaluminium, *see* Alane–*N,N*-dimethylethylamine, 0071
- Dimethyl ethanephosponite, 1732
- † Dimethyl ether, 0919
- (Dimethyl ether)oxodiperoxochromium(VI), 0904
- 1-(1,1-Dimethylethoxy)-2-methylpropane, 3072
- 1,1-Dimethylethyldibromamine, 1636
- N*-Dimethylethyl-3,3-dinitroazetidine, 2848
- 1,1-Dimethylethyl peroxyxynitrate, *see tert*-Butyl peroxyxynitrate, 1663
- 4-(Dimethylethyl)-2,2,6,6-tetramethyl-3-oxa-5-phospha-2,6-disilahept-4-ene, *see* 1-Trimethylsilyloxy-1-trimethylsilylphosphylidene-2,2-dimethylpropane, 3417
- N,N*-Dimethylethynamine, *see* (Dimethylamino)acetylene, 1564 (
- Dimethylfluoroarsine, 0885
- Dimethylformamide, 1259
- 6,6-Dimethylfulvene, 2967
- † 2,5-Dimethylfuran, 2382
- Dimethylgold selenocyanate, 1200
- 2,6-Dimethyl-2,5-heptadien-4-one diozonide, 3176
- † 2,5-Dimethylheptane, 3200
- † 3,5-Dimethylheptane, 3201
- † 4,4-Dimethylheptane, 3202
- † 2,6-Dimethyl-3-heptene, 3190
- 2,10-Dimethyl-1,2,3,4,5,10-hexahydroazepino[3-4,b]indole, 3662
- 1,3-Dimethylhexahydropyrimidone, 2475
- † 2,3-Dimethylhexane, 3054
- † 2,4-Dimethylhexane, 3055
- 3,5-Dimethyl-3-hexanol, 3073
- 3,7-Dimethyl-6-hexenoic acid, *see* Citronellic acid, 3352
- 2,5-Dimethyl-2,5-hexylenebis diperoxyoxalate, *see* 7,7,10,10-Tetramethyl-1,2,5,6-tetroxecane-3,4-dione, 3349
- 3,5-Dimethyl-3-hexyl hydroperoxide, 3075
- Dimethyl hydrazidophosphate, 0959
- † 1,1-Dimethylhydrazine, 0954
- † 1,2-Dimethylhydrazine, 0955
- Dimethyl hyponitrite, 0913
- Dimethyliodoarsine, 0886
- 3,5-Dimethylisoxazole, 1888
- Dimethylketene, 1518
- Dimethylmagnesium, 0908
- Dimethylmanganese, 0909
- Dimethylmercury, 0907
- † *N,N*-Dimethylmethanamine, *see* Trimethylamine, 1310
- O,O*-Dimethyl *S*-methylcarbamoylmethyl phosphorodithioate, 2009
- Dimethylmethyleneoxosulfanenickel(0) diethylene complex, 2856
- 3,3-Dimethyl-1-nitro-1-butyne, 2395

Dimethyl 4-nitrophenyl thionophosphate, 2979
N,N-Dimethyl-4-nitrosoaniline, 2980
 1,2-Dimethylnitrosohydrazine, 0946
 3,7-Dimethyl-2,6-octadienal, 3346
O,O-Dimethyl-*S*-[(4-oxo-1,2,3-benzotriazin-3[4*H*]-yl)methyl]
 phosphorodithioate, 3307
 † 2,3-Dimethylpentane, 2871
 † 2,4-Dimethylpentane, 2872
 † 2,4-Dimethyl-3-pentanone, 2857
S,S-Dimethylpentasulfur hexanitride, 0917
 Dimethyl peroxide, 0923
 Dimethyl peroxydicarbonate, 1544
 Dimethylphenylarsine, 2991
 Dimethyl(phenylethynyl)thallium, 3298
 Dimethylphenylphosphine, 3003
 Dimethylphenylphosphine oxide, 3002
 3,3-Dimethyl-1-phenyltriazene, 2996
 Dimethylphosphine, 0948
 Dimethyl phosphoramidate, 0952
 Dimethylphosphorohydrazine, *see* Dimethyl hydrazidophosphate, 0959
 † 2,6-Dimethylpiperidine, 2868
 † 2,2-Dimethylpropane, 1999
 † 1,1-Dimethylpropanol, *see tert*-Pentanol, 2015
 † *N,N*-Dimethylpropylamine, 2031
 † 1,1-Dimethylpropylamine, 2028
 † 1,2-Dimethylpropylamine, 2029
 † 2,2-Dimethylpropylamine, 2030
 Dimethylpropyleneurea, *see* 1,3-Dimethylhexahydropyrimidone, 2475
 Dimethyl-1-propynylthallium, 1938
 2,5-Dimethylpyrazine 1,4-dioxide, 2378
 2,6-Dimethylpyridine *N*-oxide, 2819
 3,3-Dimethyl-1-(3-quinolyl)triazene, 3398
 Dimethyl selenate, 0930
 (Dimethylsilylmethyl)trimethyllead, 2605
 † Dimethyl sulfate, 0929
 † Dimethyl sulfide, 0932
 Dimethyl sulfite, 0927
 † Dimethyl sulfoxide, 0921
 Dimethyl terephthalate, 3292
 3,6-Dimethyl-1,2,4,5-tetraoxane, 1625
 Dimethylthallium fulminate, 1211
 Dimethylthallium *N*-methylacetohydroxamate, 2008
 Dimethyltin dinitrate, 0914
 2,2-Dimethyltriazanium perchlorate, 0963
 1,3-Dimethyltriazene, 0945
 1,2-Dimethyl-2-trimethylsilylhydrazine, 2047

Dimethyltrimethylsilylphosphine, 2045
 1,4-Dimethyl-2,3,7-trioxabicyclo[2.2.1]hept-5-ene, 2388
 1,5-Dimethyl-6,7,8-trioxabicyclo[3.2.1]octane, *see* 1,2-Dimethylcyclopentene
 ozonide, 2845
 3,5-Dimethyl-1,2,4-trioxolane, *see trans*-2-Butene ozonide, 1623
 Dimethylzinc, 0935
 Di-1-naphthoyl peroxide, 3831
 Dinitramide, *see* Dinitramine, 4442
 Dinitramine, 4442
 Dinitroacetonitrile, 0676
 2,4-Dinitroaniline, 2277
 Dinitroanilines, 2276
 3,3-Dinitroazetidine, 1195
 3,3-Dinitroazetidinium trifluoromethanesulfonate, 1495
 1,2-Dinitrobenzene, 2194
 1,3-Dinitrobenzene, 2195
 1,4-Dinitrobenzene, 2196
 1-(2,4-Dinitrobenzeneazo)-2-hydroxynaphthalene, 3702
 2,4-Dinitrobenzenediazonium hydrogen sulfate, 2206
 3,5-Dinitrobenzenediazonium 2-oxide, 2087
 4,6-Dinitrobenzenediazonium 2-oxide, 2088
 4,6-Dinitro-1,3-benzenediol, 2198
 2,4-Dinitrobenzenesulfonyl chloride, 2099
 2,4-Dinitrobenzenesulfonyl chloride, 2100
 4,6-Dinitrobenzofurazan *N*-oxide, 2089
 4,6-Dinitrobenzofurazan *N*-oxide.cysteine complex, *see* 4-(2-Ammonio-2-
 carboxyethylthio)-5,7-dinitro-4,5-dihydrobenzofurazanide *N*-oxide,
 3146
 5,7-Dinitrobenzoxa-1,2,3-diazole, *see* 3,5-Dinitrobenzenediazonium 2-oxide,
 2087
 2,6-Dinitrobenzyl bromide, 2672
 2,2'-Dinitrobibenzyl, *see* 1,2-Bis(2-nitrophenyl)ethane, 3646
 1,1-Dinitro-3-butene, 1508
 2,3-Dinitro-2-butene, 1509
 4,6-Dinitro-2-*sec*-butylphenol, 3306
N,N'-Dinitro-1,2-diaminoethane, 0916
 Dinitrodiazomethane, 0545
 5,6-Dinitro-2-dimethylaminopyrimidinone, 2358
N'-dinitroethylenediamino)cyclotriphosphazene
 1,4,8,11,14,17-Hexanitro-1,4,6,8,11,12,14,17,18-nonaaza-5 λ ⁵,7 λ ⁵,13 λ ⁵-
 triphosphatatripiro[4.1.4.1.4.1]octadeca-5,7(12),13(18)-triene, *see*
 1,1,3,3,5,5-Tris-spiro(*N*,, 2483
 5(1,1-Dinitroethyl)-2-methylpyridine, 2965
 2,2-Dinitro-2-fluoroethoxycarbonyl chloride, 1072
 3,4-Dinitrofurazan-2-oxide, 1008

Dinitrogen oxide, 4744
 Dinitrogen pentaoxide, 4748
 Dinitrogen tetraoxide, 4747
 Dinitrogen tetraoxide, *see* Nitrogen dioxide, 4725
 Dinitrogen trioxide, 4746
 3,5-Dinitro-4-hydroxybenzenediazonium 2-oxide, 2090
 1,3-Dinitro-2-imidazolidinone, 1142
 Dinitromethane, 0410
N,N'-Dinitromethanediamine, lead(II) salt, *see* Lead methylenebis(nitramide),
 0412
 3,5-Dinitro-2-methylbenzenediazonium-4-oxide, 2666
 3,5-Dinitro-6-methylbenzenediazonium-2-oxide, 2667
 2,5-Dinitro-3-methylbenzoic acid, 2919
N,N'-Dinitro-*N*-methyl-1,2-diaminoethane, 1276
 1,5-Dinitronaphthalene, 3242
 1,4-Dinitropentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane, 2918
 2,4-Dinitropentane isomers, 1953
 2,6-Dinitro-4-perchlorylphenol, 2101
 2,4-Dinitrophenol, 2197
 2,4-Dinitrophenylacetyl chloride, 2902
 2,4-Dinitrophenylhydrazine, 2321
 2,4-Dinitrophenylhydrazinium perchlorate, 2351
O-(2,4-Dinitrophenyl)hydroxylamine, 2279
 3,8-Dinitro-6-phenylphenanthridine, 3780
 3,5-Dinitro-2-(picrylazo)pyridine, 3387
 4,6-Dinitro-1-picrylbenzotriazole, 3438
 5,7-Dinitro-1-picrylbenzotriazole, 3439
 2,2-Dinitropropylhydrazine, 1277
 1-(3,5-Dinitro-2-pyridyl)-2-picrylhydrazine, 3389
 4,6-Dinitroresorcinol, *see* 4,6-Dinitro-1,3-benzenediol, 2198
 3,5-Dinitrosalicylic acid, 2665
 3,7-Dinitroso-1,3,5,7-tetraazabicyclo[3.3.1]nonane, 1954
 Dinitrosylnickel, 4741
 3,5-Dinitro-2-toluamide, 2941
 2,4-Dinitrotoluene, 2726
 4,6-Dinitro-2-(2,4,6-trinitrophenyl)benzotriazole *N*-oxide, 3441
 7,8-Dioxabicyclo[4.2.2]-2,4,7-decatriene, 2955
 † 1,3-Dioxane, 1616
 † 1,4-Dioxane, 1617
 cis-1,4-Dioxenedioxetane, 1538
 1,4-Dioxobut-2-yne, *see* Acetylenedicarboxaldehyde, 1403
 3,6-Dioxo-1,2-dioxane, 1445
 † 1,3-Dioxolane, 1228
 † 1,3-Dioxol-4-en-2-one, 1087
 Dioxonium hexamanganato(VII)manganate, 4553

5,12-Dioxo-4,4,11,11-tetrahydroxy-14*H*-[1,2,5]oxadiazolo[3,4-*e*]
 [1,2,5]oxadiazolo[3',4':4,5]benzotriazolo[2,1-*a*]benzotriazolo-*b*-ium
 1,8-dioxide, 3440
 1,1'-Dioxybiscyclohexanol, *see* Bis(1-hydroxycyclohexyl) peroxide, 3553
 Dioxydicyclohexylidenebishydroperoxide, *see* Bis(1-hydroperoxycyclohexyl)
 peroxide, 3556
 Dioxygen difluoride, 4320
 Dioxygenyl tetrafluoroborate, 0132
 † Dipentyl ether, 3369
 2,6-Diperchloryl-4,4-diphenquinone, 3444
 Diperoxyazelaic acid, 3189
 Diperoxyterephthalic acid, 2925
 Diphenylacetylene, *see* Diphenylethyne, 3634
 Diphenylamine, 3504
 Diphenyl azidophosphate, 3489
 2,5-Diphenyl-3,4-benzofuran-2,5-endoperoxide (1,4-Epoxy-1,4-dihydro-
 1,4-diphenyl-2,3-benzodioxin), 3798
 4,4'-Diphenyl-2,2'-bi(1,3-dithiol)-2'-yl-2-ylum perchlorate, 3746
 1,6-Diphenyl-2,5-bis(diazo)-1,3,4,6-tetraoxohexane, 3744
 Diphenylcyclopropenylum perchlorate, 3680
 Diphenyldiazene, *see* Azobenzene, 3483
 Diphenyl diselenide, 3501
 Diphenyldistibene, 3500
 Diphenyl ether, 3492
 1,1-Diphenylethylene, 3642
trans-1,2-Diphenylethylene, 3643
 Diphenylethyne, 3634
 1,2-Diphenylhydrazine, 3517
 1,1-Diphenylhydrazinium chloride, 3519
 Diphenylmagnesium, 3482
 Diphenylmercury, 3480
 1,8-Diphenyloctatetrayne, 3788
 1,5-Diphenylpentaazadiene, 3507
N,N-Diphenyl-3-phenylpropenylidenimmonium perchlorate
 Diphenylphosphine, 3508
 Diphenylphosphorus(III) azide, 3488
 Diphenylselenone, 3494
 Diphenyl sulfoxide, 3493
 2,2-Diphenyl-1,3,4-thiadiazoline, 3647
 Diphenyltin, 3502
 1,3-Diphenyltriazene, 3506
 5,6-Diphenyl-1,2,4-triazine-3-diazonium tetrafluoroborate 2-oxide, 3677
 † Diphosphane, 4538
 1,2-Diphosphinoethane, 0957
 Diphosphoryl chloride, 4168
 Dipotassium μ -cyclooctatetraene, 2947

Dipotassium diazirine-3,3-dicarboxylate, 1340
 Dipotassium *aci*-nitroacetate, 0672
 Dipotassium phosphinate, 4431
 Dipotassium triimidotellurite, 4494
 Dipropargyl ether, *see* Di(2-propynyl) ether, 2327
 † Di-2-propenyl ether, *see* Diallyl ether, 2431
 Di-2-propenyl phosphonite, *see* Diallyl phosphite, 2456
 Dipropionyl peroxide, 2442
 Dipropylaluminium hydride, 2552
 † Dipropylamine, 2568
 † Dipropyl ether, 2543
 Dipropyl hyponitrite, 2539
 Dipropylmercury, 2537
 Dipropyl peroxide, 2547
 Dipropyl peroxydicarbonate, 3035
 Dipropylzinc, 2551
 Di(1-propynyl) ether, 2326
 Di(2-propynyl) ether, 2327
 Diprotium monoxide, 4475
 Dipyridinesilver(I) perchlorate, 3267
 Dipyridinesodium, 3280
 Dipyridinium dichromate, 3304
 2-(1,3-Diselena-4,5,6,7-tetrahydroindanylidene)piperidinium perchlorate, 3408
 Diselenium dichloride, 4115
 Diseleniumdisulfur tetranitride, 4769
 † Disilane, 4569
 Disilver cyanamide, 0305
 Disilver diazomethanide, 0306
 Disilver ketenide, 0570
 Disilver ketenide–silver nitrate, 0571
 Disilver pentatin undecaoxide, 0036
 Disilyl oxide, 4565
 Disilyl sulfide, 4568
 Disodium 5,5'-azotetrazolide, 1018
 Disodium 5,5'-azoxytetrazolide, 1019
 Disodium dicyanodiazene, 1006
 Disodium *N,N'*-dimethoxysulfonyldiamide, 0911
 Disodium 4-nitrophenylphosphate, 2185
 Disodium 5-tetrazolazocarboxylate, 1012
 Disodium tetrazole-5-diazoate, 0549
 Disodium tridecahydrononaborate(2–), 0195
 3,6-Di(spirocyclohexane)-1,2,4,5-tetraoxane (7,8,15,16-Tetraoxadispiro
 [5.2.5.2]hexadecane), 3546
 Disulfur dibromide, 0282
 Disulfur dichloride, 4114
 Disulfur dinitride, 4754

- Disulfur heptaoxide, 4870
 Disulfur thiocyanate, 1003
 Disulfuryl diazide, 4780
 Disulfuryl dichloride, 4103
 Disulfuryl difluoride, 4326
 1,3-Di(5-tetrazolyl)triazene, 0778
 2,4-Dithia-1,3-dioxane-2,2,4,4-tetraoxide, 0839
 1,4-Di(1,2,3,4-thiatriazol-5-yloxy)but-2-yne, 2211
 5,5'-Dithiobis[1H]tetrazole, *see* 5,5'-Tetrazolyl disulfide, 0721
 4,4'-Dithiodimorpholine, 3047
 † 'Dithiohydantoin', *see* Imidazoline-2,4-dithione, 1141
 1,3-Dithiolium perchlorate, 1094
 Di-4-toluenesulfonyl peroxide, 3656
 1,6-Di(4'-tolyl)-2,5-bis(diazo)-1,3,4,6-tetraoxohexane, 3796
 Di[tris(1,2-diaminoethanecromium(III)) triperoxodisulfate, 3593
 Di[tris-1,2-diaminoethanecobalt(III)] triperoxodisulfate, 3592
 Di(triselenadiazolium) dichloride, 4091
 Ditungsten carbide, 0564
 † Divinylacetylene, *see* 1,5-Hexadien-3-yne, 2291
 † Divinyl ether, 1519
 Divinyl ketone, 1877
 Divinylmagnesium, 1498
 Divinylzinc, 1547
 Dodecacarbonyltetracobalt, 3420
 Dodecacarbonyltriiron, 3428
 Dodecachloropentasilane, 4198
 Dodecahydro-4,8-dinitro-2,6-bis(nitroimino)diimidazo[4,5-*b*:4',5'-*e*]pyrazine,
 2381
 Dodecamethyltetraplatinum(IV) azide, 3590
 Dodecamethyltetraplatinum(IV) perchlorate, 3587
 Dodecanethiol, 3567
 Dodecaspino[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]tetraoctaconta-
 4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62,67,69,74,76,81,83-
 tetracosayne, 3921
 Dysprosium perchlorate, 4131

 Endrin, 3459
 Eosin, *see* 2',4',5',7'-Tetrabromo-3',6'-dihydroxyspiro[isobenzofuran-
 1(3*H*),9'-[9*H*]xanthen-3-one], 3787
 9,10-Epidioxyanthracene, 3630
 1,4-Epidioxy-1,4-dihydro-6,6-dimethylfulvene, 2984
 1,4-Epidioxy-2-*p*-menthene, 3347
 † 1,2-Epoxybutane, 1609
 † 3,4-Epoxybutene, 1520
 endo-2,3-Epoxy-7,8-dioxabicyclo[2.2.2]oct-5-ene, 2336
 2,3-Epoxypropanol, 1229

- 2,3-Epoxypropionaldehyde 2,4-dinitrophenylhydrazone, 3130
 2,3-Epoxypropionaldehyde oxime, 1182
 4(2,3-Epoxypropoxy)butanol, 2866
 † 2,3-Epoxypropyl nitrate, 1186
 3(2,3-Epoxypropyloxy)2,2-dinitropropyl azide, 2403
 Erbium perchlorate, 4132
 † Ethanal, *see* Acetaldehyde, 0828
 † Ethanamine, *see* Ethylamine, 0942
 † Ethane, 0881
 † Ethanedial, 0723
 † Ethanedinitrile, *see* Dicyanogen, 0996
 Ethanedioic acid, *see* Oxalic acid, 0725
 1,2-Ethanediol, *see* Ethylene glycol, 0924
 'Ethane hexamercarbide', *see* 1,2-Bis(hydroxomercurio)-1,1,2,2-bis(oxydimercurio)ethane, 0708
 † Ethanenitrile, *see* Acetonitrile, 0758
 Ethaneperoxoic acid, *see* Peroxyacetic acid, 0837
 † Ethanethiol, 0933
 Ethanoic acid, *see* Acetic acid, 0833
 † Ethanol, 0920
 Ethanolamine, *see* 2-Hydroxyethylamine, 0943
 Ethanolamine perchlorate, *see* 2-Hydroxyethylaminium perchlorate, 0951
 † Ethanoyl chloride, *see* Acetyl chloride, 0735
 Etheneamine, *see* Vinylamine, 0864
 † Ethene, *see* Ethylene, 0781
 Ethenone, *see* Ketene, 0722
 † 3-Ethenoxypropene, *see* Allyl vinyl ether, 1904
 Ethenylbenzene, *see* Styrene, 2945
 † Ethenyl ethanoate, *see* Vinyl acetate, 1532
 Ethenylmagnesium chloride, 0732
 † Ethenyl methanoate, *see* Vinyl formate, 1149
 † Ethoxyacetylene, 1521
 4-Ethoxybutyldiethylaluminium, 3372
 3-Ethoxycarbonyl-4,4,5,5-tetracyano-3-trimethylplumbyl-4,5-dihydro-3*H*-pyrazole, 3611
 Ethoxydiisobutylaluminium, 3373
 † Ethoxyethane, *see* Diethyl ether, 1697
 2-Ethoxyethanol, 1702
 † Ethoxyethene, *see* Ethyl vinyl ether, 1610
 2-Ethoxyethyl acetate, 2511
 † Ethoxyethyne, *see* Ethoxyacetylene, 1521
 2-Ethoxy-1-iodo-3-butene, 2449
 4-Ethoxy-2-methyl-3-butyn-2-ol, 2844
 3-Ethoxymethylene-2,4-pentanedione, 3014
 † 3-Ethoxypropene, *see* Allyl ethyl ether, 1955
 † 1-Ethoxypropene, *see* Ethyl propenyl ether, 1956

- 1-Ethoxy-2-propyne, 1909
- Ethoxytriethyldiphosphinyl oxide, 3094
- † Ethyl acetate, 1618
- Ethyl acetoacetate, 2440
- 2-Ethylacrylaldehyde oxime, 1933
- † Ethyl acrylate, 1913
- † Ethyl alcohol, *see* Ethanol, 0920
- Ethylaluminium bromide iodide, 0841
- Ethylaluminium dibromide, 0842
- Ethylaluminium dichloride, 0843
- Ethylaluminium diiodide, 0844
- † Ethylamine, 0942
- N*-Ethylaniline, 2994
- Ethyl azide, 0872
- Ethyl α -azido-*N*-cyanophenylacetimidate, 3397
- † Ethyl azidoformate, 1193
- Ethyl 2-azido-2-propenoate, 1890
- † Ethylbenzene, 2968
- † Ethyl bromide, *see* Bromoethane, 0846
- Ethyl 4-bromo-1,1,3-trioxoisothiazoleacetate, 2792
- † 2-Ethylbutanal, 2487
- 2-Ethyl-1-butene-1-one, *see* Diethylketene, 2430
- Ethyl carbamate, 1261
- Ethyl carbamioniothioate hydrogen sulfate, *see* *S*-Ethylisothiuronium hydrogen sulfate, 1327
- Ethyl carbonazidate, *see* Ethyl azidoformate
- † Ethyl carbonochloridate, *see* Ethyl chloroformate, 1164
- † Ethyl chloride, *see* Chloroethane, 0848
- Ethyl chloroacetate, 1559
- † Ethyl chloroformate, 1164
- Ethyl chloroglyoxylate, *see* Ethyl oxalyl chloride, 1456
- Ethyl *N*-chloro-*N*-sodiocarbamate, 1159
- † Ethyl crotonate, 2436
- Ethyl cyanoacetate, 1889
- Ethyl 2-cyano-2-(1-*H*-tetrazol-5-ylhydrazono)acetate, 2359
- † Ethylcyclobutane, 2458
- † Ethylcyclopentane, 2850
- † Ethylcyclopropane, 1941
- † Ethyl cyclopropanecarboxylate, 2437
- Ethyl 2,3-diazidopropionate, 1903
- Ethyl diazoacetate, 1503
- Ethyl 4-diazo-1,2,3-triazole-5-carboxylate, 1853
- Ethyl 3,4-dihydroxybenzenesulfonate, 2988
- † Ethyldimethylamine, 1727
- † 3-Ethyl-2,3-dimethylpentane, 3203
- Ethyldimethylphosphine, 1735

- † Ethylene, 0781
 - 1,2-Ethylenebis(ammonium) perchlorate, 0964
 - Ethylenebis(dimethylphosphine), 2585a
 - 2,2'-(1,2-Ethylenebis)3-phenyloxaziridine, 3707
 - Ethylenediamine, *see* 1,2-Diaminoethane, 0953
 - Ethylenediamine dinitrate, *see* 1,2-Diammonioethane nitrate, 0966
 - Ethylene dibromide, *see* 1,2-Dibromoethane, 0785
 - Ethylenedicaesium, 0798
- † Ethylene dichloride, *see* 1,2-Dichloroethane, 0791
 - Ethylenedioxyammonium perchlorate, *see* 2-Aza-1,3-dioxolanium perchlorate, 0898
 - Ethylene diperchlorate, 0795
 - Ethylene glycol, 0924
 - Ethylene glycol monomethyl ether, *see* 2-Methoxyethanol, 1284
- † Ethylene oxide, 0829
 - Ethylene ozonide, 0836
- † Ethylene sulfide, *see* Thiirane, 0840
- † Ethylenimine, *see* Aziridine, 0863
- † Ethyl ethanoate, *see* Ethyl acetate, 1618
 - Ethyl 6-ethoxycarbonyl-3,4-dihydro-1,1,3-trioxo-2-pyrazolo[3,4-d]isothiazole-2-acetate, 3403
- † Ethyl fluoride, *see* Fluoroethane, 0855
 - Ethyl fluorosulfate, 0856
- † Ethyl formate, 1230
 - Ethyl 2-formylpropionate oxime, 2454
 - 4-Ethyl-3-furazanone, *see* 3-Ethyl-4-hydroxy-1,2,5-oxadiazole, 1504
 - N*-Ethylheptafluorobutyramide, 2307
- † 2-Ethylhexanal, 3050
- † 2-Ethylhexyl vinyl ether, 3360
 - Ethyl hydroperoxide, 0925
 - 3-Ethyl-4-hydroxy-1,2,5-oxadiazole, 1504
 - Ethyl hypochlorite, 0851
 - Ethylidene dinitrate, 0811
 - Ethyl iminioformate chloride, 1273
 - Ethyl iodide, *see* Iodoethane, 0858
 - Ethyliodomethylarsine, 1272
- † Ethyl isobutyrate, 2499
- † Ethyl isocyanide, 1178
- † Ethyl isopropyl ether, 2012
 - S*-Ethylisothiuronium hydrogen sulfate, 1327
- † Ethyl isovalerate, 2859
 - Ethyllithium, 0862
 - Ethylmagnesium bromide, 0847
 - Ethylmagnesium iodide, 0859
- † Ethyl methacrylate, 2438
- † Ethyl methanoate, *see* Ethyl formate, 1230

Ethylmethylarsine, 1293
 † 2-Ethyl-2-methyl-1,3-dioxolane, 2500
 † Ethyl methyl ether, 1278
 † 3-Ethyl-4-methylhexane, 3204
 † 4-Ethyl-2-methylhexane, 3205
 † Ethyl methyl ketone, *see* 2-Butanone, 1606
 Ethyl *N*-methyl-*N*-nitrosocarbamate, 1597
 † 3-Ethyl-2-methylpentane, 3056
 Ethyl methyl peroxide, 1282
 3-Ethyl-1(4-methylphenyl)triazene, 3171
 5-Ethyl-2-methylpyridine, 2995
 † Ethyl methyl sulfide, 1288
 3-Ethyl-5-methyl-1,2,4-trioxolane, *see trans*-2-Pentene ozonide, 1982
 † Ethyl nitrate, 0870
 † Ethyl nitrite, 0867
 Ethyl 2-nitroethyl ether, 1662
 'Ethylnitrolic acid', *see* 1-Nitro-1-oximinoethane, 0810
 Ethyl oxalyl chloride, 1456
 † Ethyloxirane, *see* 1,2-Epoxybutane, 1609
 Ethyl 3-oxobutanoate, *see* Ethyl acetoacetate, 2440
 Ethylpentaborane(9), 0973
 Ethyl perchlorate, 0852
 Ethylphenylthallium(III) acetate perchlorate, 3302
 Ethylphosphine, 0949
 † 1-Ethylpiperidine, 2869
 † 2-Ethylpiperidine, 2870
 † Ethyl propanoate, *see* Ethyl propionate, 1970
 † Ethyl propenoate, *see* Ethyl acrylate, 1913
 † Ethyl propenyl ether, 1956
 † Ethyl propionate, 1970
N-Ethyl-*N*-propylcarbonyl chloride, 2468
 † Ethyl propyl ether, 2013
 2-Ethylpyridine-4-carbothioamide, 2981
 Ethyl pyrocarbonate, *see* Diethyl dicarbonate, 2444
 Ethyl sodioacetoacetate, 2404
 Ethylsodium, 0877
 1-Ethyl-1,1,3,3-tetramethyltetrazenium tetrafluoroborate, 2586
 2-Ethyltetrazole, 1213
 5-Ethyltetrazole, 1214
 † Ethyl trifluoroacetate, 1460
 Ethyl 2,2,3-trifluoropropionate, 1886
 † Ethyl vinyl ether, 1610
 † Ethyne, *see* Acetylene, 0686
 2-Ethynylfuran, 2212
 † Ethynyl methyl ether, *see* Methoxyacetylene, 1146
 Ethynyl vinyl selenide, 1449

Europium, 4292
Europium(II) sulfide, 4293

Ferrocene, 3276
Ferrocene-1,1'-dicarboxylic acid, 3479
 μ ,1,1'-Ferrocenediyldilithium, 3256
Ferrocenium perchlorate, 3269
1-Ferrocenyl-2-methylpropyl azide, 3661
Ferromanganese, 4389
Ferrosilicon, 4390
Ferrotitanium, 4391
Fluoramide, *see* Fluoramine, 4296
Fluoramine, 4296
Fluorimide, *see* Difluoramine, 4311
Fluorine, 4310
Fluorine azide, 4307
Fluorine fluorosulfate, 4324
'Fluorine nitrate', *see* Nitryl hypofluorite, 4304
Fluorine perchlorate, 3976
Fluoroacetylene, 0662
† Fluorobenzene, 2248
Fluorobis(trifluoromethyl)phosphine, 0646
Fluorobis(trifluoromethyl)phosphine oxide, 0645
Fluorocarbonyl azide, *see* Azidocarbonyl fluoride, 0339
Fluorocarbonylperoxonitrate, 0338
Fluorodiodomethane, 0373
1-Fluoro-2,4-dinitrobenzene, 2108
1-Fluoro-1,1-dinitrobutane, 1562
4-Fluoro-4,4-dinitrobutene, 1458
1-Fluoro-1,1-dinitroethane, 0748
2(?) -Fluoro-1,1-dinitroethane, 0749
2-Fluoro-2,2-dinitroethanol, 0750
2-Fluoro-2,2-dinitroethylamine, 0800
Fluorodinitromethane, 0374
Fluorodinitromethyl azide, 0341
1-Fluoro-1,1-dinitro-2-phenylethane, 2934
† 1-Fluoro-2,3-epoxypropane, 1171
† Fluoroethane, 0855
† Fluoroethylene, 0747
Fluoroethylene ozonide, 0752
Fluorohydrocyanokrypton(II) hexafluoroarsenate, 0367
1-Fluoroiminohexafluoropropane, 1058
2-Fluoroiminohexafluoropropane, 1059
N-Fluoroiminosulfur tetrafluoride, 4357
Fluorokrypton hexafluoroarsenate, 0096
† Fluoromethane, 0441

- † Fluoromethyloxirane, *see* 1-Fluoro-2,3-epoxypropane, 1171
- 5-Fluoro-2-methylphenylmagnesium bromide, 2709
- Fluoromethylsilane, 0492
- N*-Fluoro-*N*-nitrobutylamine, 1645
- 5-Fluoro-2-nitrophenol, 2167
- Fluoronium perchlorate, 3972
- 4-Fluorophenyllithium, 2166
- Fluorophosphoric acid, 4297
- 2-Fluoropyridine, 1831
- Fluoroselenic acid, 4295
- Fluorosilane, 4298
- Fluorosulfonyl hypofluorite, *see* Fluorine fluorosulfate, 4324
- Fluorothiophosphoryl diazide, 4308
- † 2-Fluorotoluene, 2747
- † 3-Fluorotoluene, 2748
- † 4-Fluorotoluene, 2749
- α -Fluorotoluene, *see* Benzyl fluoride, 2746
- 3-Fluoro-3-(trifluoromethyl)-3*H*-diazirine, 0631
- Fluorotrinitromethane, 0340
- 3-Fluoro-1,2,4-trioxolane, *see* Fluoroethylene ozonide, 0752
- Fluoroxymethane, *see* Methyl hypofluorite, 0442
- † Formaldehyde, 0416
- Formaldehyde oxide polymer (Poly(dioxymethylene)), 0419
- Formamide, 0453
- Formic acid (Methanoic acid), 0418
- 2-Formylamino-3,5-dinitrothiophene, 1823
- 2-Formylamino-1-phenyl-1,3-propanediol, 3322
- Formylhydrazine, 0473
- Formyloxymethyl hydroperoxide, *see* Hydroperoxymethyl formate, 0838
- Formyl(triisopropyl)silane, 3370
- Freon 22, *see* Chlorodifluoromethane, 0369
- Freon 12, *see* Dichlorodifluoromethane, 0326
- Freon 11, *see* Trichlorofluoromethane, 0330
- Fullerene, 3919
- Fulminic acid, 0381
- † 'Fulminuric acid', *see* 2-Carbamoyl-2-nitroacetonitrile 1116
- 6-Fulvenoselone, 2216
- Fumaric acid, 1446
- Fumarodinitrile, 1397
- 2-Furaldehyde, 1836
- 2-Furaldehyde azine, 3259
- † Furan, 1439
- Furan-2-amidoxime, 1872
- 2-Furanemethanol, *see* Furfuryl alcohol, 1880
- Furfural, *see* 2-Furaldehyde, 1836
- Furfuryl alcohol, 1880

2-Furoyl azide, 1821
Furoyl chloride, 1818
2-Furylchlorodiazirine, 1817

Gallium, 4406
Gallium azide, 4409
Gallium(I) oxide, 4411
Gallium perchlorate, 4135
† Germane, 4417
Germanium, 4412
Germanium(II) sulfide, 4419
Germanium imide, 4414
Germanium isocyanate, 1377
Germanium tetrachloride, 4162
Glucose, 2518
Glutarodinitrile, 1870
Glutaryl diazide, 1874
Glycerol, 1286
† Glyceryl trinitrate, 1196
Glycidyl azide, 1191
Glycidyl phenyl ether, *see* 3-Phenoxy-1,2-epoxypropane, 3150
Glycolonitrile, 0760
† Glyoxal, *see* Ethanedial, 0723
Glyoxylic acid, 0724
Gold, 0110
Gold(I) acetylde, 0573
Gold(I) cyanide, 0308
Gold(III) chloride, 0111
Gold(III) hydroxide–ammonia, 0112
Gold(III) nitride trihydrate, 0118
Gold(III) oxide, 0115
Gold(III) sulfide, 0116
Gold(I) nitride–ammonia, 0117
Graphite hexafluorogermanate, 3422
Guanidinium dichromate, 0971
Guanidinium nitrate, 0509
Guanidinium perchlorate, 0502

Hafnium, 4599
Hafnium carbide, 0521
Hafnium(IV) tetrahydroborate, 0182
Hafnium tetrachloride, 4163
Heptafluorobutyramide, 1392
Heptafluorobutyryl hypochlorite, 1352
Heptafluorobutyryl hypofluorite, 1369

Heptafluorobutyryl nitrite, 1368
 Heptafluoroisopropyl hypochlorite, 1035
 2-Heptafluoropropyl-1,3,4-dioxazolone, 1812
 Heptafluoropropyl hypofluorite, 1060
 Heptakis(dimethylamino)trialuminium triboron pentahydride, 3676
 † Heptane, 2873
 1,1,1,3,5,5,5-Heptanitropentane, 1854
 Heptasilver nitrate octaoxide, 0047
 Heptaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]nonatetraconta-
 4,6,11,13,18,20,25,27,32,34,39,41,46,48-tetradecayne, 3912
 Hepta-1,3,5-triyne, 2641
 † 1-Heptene, 2851
 † 2-Heptene, 2852
 † 3-Heptene, 2853
 1-Heptene-4,6-diyne, 2707
 † 1-Heptyne, 2838
 2-Heptyn-1-ol, 2841
 Hexaamminecalcium, 3933
 Hexaamminechromium(III) nitrate, 4234
 Hexaamminechromium(III) perchlorate, 4129
 Hexaamminecobalt(III) chlorate, 4125
 Hexaamminecobalt(III) hexanitrocobaltate(3-), 4219
 Hexaamminecobalt(III) iodate, 4208
 Hexaamminecobalt(III) nitrate, 4210
 Hexaamminecobalt(III) perchlorate, 4126
 Hexaamminecobalt(III) permanganate, 4209
 Hexaamminenickel chlorate, 4074
 Hexaamminenickel perchlorate, 4075
 1,1,3,3,5,5-Hexaazido-2,4,6-triaza-1,3,5-triphosphorine, 4795
 Hexaborane(10), 0191
 Hexaborane(12), 0192
 Hexabromoethane, 0584
 Hexacarbonylchromium, 2056
 Hexacarbonylmolybdenum, 2628
 Hexacarbonyltungsten, 2636
 Hexacarbonylvandium, 2635
 Hexachlorobenzene, 2055
 1,2,3,4,5,6-Hexachlorocyclohexane, 2306
 Hexachlorocyclopentadiene, 1808
 Hexachlorocyclotriphosphazine, 4189
 Hexachlorodisilane, 4191
 1,2,3,4,10,10-Hexachloro-6,7-epoxy-1,4,4a,5,6,7,8,8a-octahydro-
 endo,endo-1,4:5,8-dimethanonaphthalene), *see* Endrin, 3459
 Hexachloroethane, 0611
 Hexachloromelamine, *see* 2,4,6-Tris(dichloroamino)-1,3,5-triazine, 1043

- Hexacyclohexyldilead, 3894
- † Hexadecane, 3726
- Hexadecanethiol, 3727
- N*-Hexadecylpyridinium permanganate, 3827
- Hexadecyltrimethylammonium permanganate, 3785
- 2,4-Hexadienal, 2383
- † 1,3-Hexadiene, 2409
- † 1,4-Hexadiene, 2410
- † 1,5-Hexadiene, 2411
- † *cis*-2-*trans*-4-Hexadiene, 2412
- † *trans*-2-*trans*-4-Hexadiene, 2413
- 2,4-Hexadienoic acid, 2385b
- 1,3-Hexadien-5-yne, 2290
- † 1,5-Hexadien-3-yne, 2291
- 4,5-Hexadien-2-yn-1-ol, 2328
- 1,5-Hexadiyne, 2292
- 2,4-Hexadiyne, 2293
- 2,4-Hexadiyne-1,6-dioic acid, 2094
- 2,5-Hexadiyn-1-ol, 2329
- 1,5-Hexadiyn-3-one, 2213
- 2,4-Hexadiynylene chloroformate, 2890
- 2,4-Hexadiynylene chlorosulfite, 2162
- Hexaethylaluminium trithiocyanate, 3695
- † Hexafluorobenzene, 2060
- † 2,2,2,4,4,4-Hexafluoro-1,3-dimethyl-1,3,2,4-diazadiphosphetidine, 0906
- Hexafluoroglutaryl dihypochlorite, 1806
- † Hexafluoroisopropylideneamine, 1067
- Hexafluoroisopropylideneaminolithium, 1052
- Hexafluorophosphoric acid, *see* Hydrogen hexafluorophosphate, 4360
- † Hexafluoropropene, 1051
- Hexahydro[18]annulene, *see* 1,7,13-Cyclooctadecatriene-3,5,9,11,15,17-hexayne, 3740
- Hexahydropyrazine, *see* Piperazine, 1689
- Hexahydroxylaminecobalt(III) nitrate, 4211
- Hexakis(pyridine)iron(II) tridecacarbonyltetraferate(2-), 3906
- Hexalithium disilicide, 4689
- Hexamethyldiplatinum, 2606
- Hexamethyldisilane, 2608
- † Hexamethyldisilazane, 2610
- 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetracyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²cobalt(II) perchlorate, 3768
- 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetracyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²iron(II) perchlorate, 3769

- 5,6,14,15,20,21-Hexamethyl-1,3,4,7,8,10,12,13,16,17,19,22-dodecaazatetracyclo[8.8.4.1^{3,17}.1^{8,12}]tetracos-4,6,13,15,19,21-hexaene-N⁴,N⁷,N¹³,N¹⁶,N¹⁹,N²²nickel(II) perchlorate, 3770
- † Hexamethyleneimine, *see* Perhydroazepine, 2526
- Hexamethylenetetramine, 2477
- Hexamethylenetetramine tetraiodide, 2471
- Hexamethylenetetrammonium tetraperoxochromate(V)? 3779
- 'Hexamethylenetriperoxydiamine', *see* 1,6-Diaza-3,4,8,9,12,13-hexaoxabicyclo[4.4.4]tetradecane, 2476
- Hexamethylerbium-hexamethylethylenediaminelithium complex, 2592
- 2,4,6,8,9,10-Hexamethylhexaaza-1,3,5,7-tetraphosphaadamantane, 2600
- 3,3,6,6,9,9-Hexamethyl-1,2,4,5,7,8-hexoxonane, 3195
- 2,7,12,18,23,28-Hexamethyl-3,7,11,19,23,27,33,34-octaazatricyclo [27.3.1.1(13,17)]tetratriaconta-1(33),2,11,13,15,17(34), 18,27,29,31-decaenediaquadichlorodiiron(II) diperchlorate, 3880
- Hexamethylrhenium, 2607
- 5,7,7,12,14,14-Hexamethyl-1,4,8,11-tetraaza-4,11,-cyclotetradecadiene-1,10-phenanthrolineiron(II) perchlorate, 3869
- 1,1,1,3,3,3-Hexamethyl-2-(trimethylsilyl)trisilane, *see* Tris(trimethylsilyl)silane, 3231
- Hexamethyltrisiloxane, 2617
- 2,2,4,4,6,6-Hexamethyltrithiane, 3197
- Hexamethyltungsten, 2609
- † Hexane, 2531
- Hexane-1,6-dioic acid, *see* Adipic acid, 2441
- 1,6-HexanediyI perchlorate, 2470
- 6-Hexanelactam, *see* Caprolactam, 2451
- Hexanitrobenzene, 2632
- Hexanitroethane, 1014
- Hexanitrohexaazaisowurtzitane, 2325
- Hexanitrosobenzene, *see* Benzotri(furazan *N*-oxide), 2631
- † 2-Hexanone, 2488
- † 3-Hexanone, 2489
- 1,4,7,10,13,16-Hexaoxacyclooctadecane, 3561
- Hexaoxygen difluoride, 4327
- Hexaphenylhexaarsane, 3891
- Hexaspiro[2.4.2.4.2.4.2.4.2.4]dotetraconta-4,6,11,13,18,20,25, 27,32,34,39,41-dodecayne, 3905
- 1,3,5-Hexatriyne, 2074
- Hexaureachromium(III) nitrate, 2623
- Hexaureagallium(III) perchlorate, 2620
- 2-Hexenal, 2432
- † 1-Hexene, 2459
- † 2-Hexene, 2460
- trans*-2-Hexene ozonide, 2513
- Hexyl perchlorate, 2520

† 1-Hexyne, 2414
 † 3-Hexyne, 2415
 † Hydrazine, 4520
 Hydrazinium azide, 4550
 Hydrazinium chlorate, 4010
 Hydrazinium chloride, 4007
 Hydrazinium chlorite, 4008
 Hydrazinium dinitrate, 4561
 Hydrazinium diperchlorate, 4068
 Hydrazinium hydrogen selenate, 4558
 Hydrazinium nitrate, 4549
 Hydrazinium nitrite, 4548
 Hydrazinium nitroformate, *see* Hydrazinium trinitromethanide, 0496
 Hydrazinium perchlorate, 4011
 Hydrazinium salts, 4546
 Hydrazinium trinitromethanide, 0496
 4-[2-(4-Hydrazino-1-phthalazinyl)hydrazino]-4-methyl-2-pentanone
 (4-hydrazino-1-phthalazinyl)hydrazonedinickel(II) tetraperchlorate, 3836
 3-Hydrazinopropanenitrile, 1275
 5-Hydrazino-1,2,3,4-thiazotriazole, 0462
 Hydrazoic acid, *see* Hydrogen azide, 4441
 5,5'-Hydrazotetrazole, 0826
 Hydriodic acid, 4423
 Hydrochloric acid, *see* Hydrogen chloride, 3993
 Hydrogen azide, 4441
 Hydrogen bromide, 0247
 Hydrogen chloride, 3993
 † Hydrogen cyanide, 0380
 Hydrogen cyanide *N*-oxide, *see* Fulminic acid, 0381
 † Hydrogen disulfide, 4484
 Hydrogen fluoride, 4294
 Hydrogen (Gas), 4453
 Hydrogen hexaazidothallate(III), 4502
 Hydrogen hexafluorophosphate, 4360
 Hydrogen (Liquid), 4454
 Hydrogen peroxide, 4477
 † Hydrogen selenide, 4486
 † Hydrogen sulfide, 4483
 † Hydrogen telluride, 4488
 Hydrogen trisulfide, 4485
 1-Hydroperoxyphenylethane, 2985
 3-Hydroperoxy-1-butanol, *see* 1-Hydroxy-3-butyl hydroperoxide, 1706
 1-[(1-Hydroperoxycyclohexyl)dioxy]cyclohexanol, *see* 1-Hydroperoxy-
 1'-hydroxydicyclohexyl peroxide, 3555
 1-Hydroperoxy-1'-hydroxydicyclohexyl peroxide, 3555
 trans-4-Hydroperoxy-5-hydroxy-4-methylimidazolin-2-one, 1598

Hydroperoxymethyl formate, 0838
 2-Hydroperoxy-2-methylpropane, *see tert*-Butyl hydroperoxide, 1698
 2-Hydroperoxypropane, *see* Isopropyl hydroperoxide, 1283
 Hydroquinone, *see* 1,4-Benzenediol, 2333
 Hydroquinone–oxygen clathrate, *see* 1,4-Benzenediol–oxygen complex, 2334
 † Hydroximinoethane, *see* Acetaldehyde oxime, 0865
 Hydroxobis[5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II)] triperchlorate, 3881
 Hydroxyacetone, 1231
 Hydroxyacetonitrile, *see* Glycolonitrile, 0760
 Hydroxyaqua(oxo)diperxorheniumVII, 4507
N-Hydroxybenzeneamine, *see N*-Phenylhydroxylamine, 2356
 4-Hydroxybenzenediazonium-3-carboxylate, 2663
 Hydroxybenzene, *see* Phenol, 2330
 1-Hydroxy-1,2-benziodoxol-3-one-1-oxide, *see* 2-Iodylbenzoic acid, 2681
 1-Hydroxybenzotriazole, 2272
 2-Hydroxy-4,6-bis(nitroamino)-1,3,5-triazine, 1122
 2-Hydroxy-3-butenonitrile, *see* 1-Cyano-2-propen-1-ol, 1467
 1-Hydroxy-3-butyl hydroperoxide, 1706
 4-Hydroxy-*trans*-cinnamic acid, 3136
 Hydroxycopper(II) glyoximate, 0799
 4-Hydroxy-3,5-dimethyl-1,2,4-triazole, 1575
 4-Hydroxy-3,5-dinitrobenzenearsonic acid, 2220
 2-Hydroxy-3,5-dinitrobenzoic acid, *see* 3,5-Dinitrosalicylic acid, 2665
 2-Hydroxy-3,5-dinitropyridine, 1822
N-Hydroxydithiocarbamic acid, 0454
 2(2-Hydroxyethoxy)ethyl perchlorate, 1643
 2-Hydroxyethylamine, 0943
 2-Hydroxyethylaminium perchlorate, 0951
 3-(2-Hydroxyethylamino)propionitrile, 1952
N-2-Hydroxyethyl-1,2-diaminoethane, 1756
N-2-Hydroxyethyl-dimethylamine, 1729
O-(2-Hydroxyethyl)hydroxylamine, 0944
 1-Hydroxyethylidene-1,1-diphosphonic acid, 0956
 2-Hydroxyethylmercury(II) nitrate, 0857
 1-Hydroxyethyl peroxyacetate, 1626
 4(1-Hydroxyethyl)pyridine *N*-oxide
 ‘Hydroxyethyltriethylenetetramine’, *see* 3,6,9-Triaza-11-aminoundecanol, 3100
 2-Hydroxyethyltrimethylstibonium iodide, 2037
 1-Hydroxy-2-hydroxylamino-1,3,4-triazole, 0817
 1-Hydroxyimidazole-2-carboxaldoxime 3-oxide, 1477
 1-Hydroxyimidazole *N*-oxide, 1134
 1-Hydroxyiminobutane, *see* Butyraldehyde oxime, 1655
 2-Hydroxyiminopropane, *see* Acetone oxime, 1258
 Hydroxylamine, 4498

Hydroxylaminium chloride, 4002
 Hydroxylaminium nitrate, 4524
 Hydroxylaminium perchlorate, 4005
 Hydroxylaminium phosphinate, 4555
 Hydroxylaminium sulfate, 4575
 2-Hydroxyliminiobutane chloride, *see* 2-Butanone oxime hydrochloride, 1678
 4-Hydroxy-3-methoxybenzaldehyde, 2958
N-Hydroxymethylacrylamide, 1571
 2-Hydroxy-2-methylglutaric acid, 2445
 4-Hydroxy-4-methyl-1,6-heptadiene, 3029
 4-Hydroxy-4-methyl-1,6-heptadiene diozonide, 3036
 Hydroxymethyl hydroperoxide, 0487
 Hydroxymethyl methyl peroxide, 0926
 2-Hydroxymethyl-2-nitropropane-1,3-diol, *see* Tris(hydroxymethyl)
 nitromethane, 1664
 † 4-Hydroxy-4-methyl-2-pentanone, 2501
N-Hydroxymethyl-2-propenamide, *see N*-Hydroxymethylacrylamide, 1571
 3-Hydroxy-4-nitrobenzaldehyde, 2690
 4-Hydroxy-3-nitrobenzaldehyde, 2691
 5-Hydroxy-2-nitrobenzaldehyde, 2692
 4-Hydroxy-3-nitrobenzenesulfonyl chloride, 2146
 2-Hydroxy-6-nitro-1-naphthalenediazonium-4-sulfonate, 3240
 (Hydroxy)(oxo)(phenyl)- λ^3 -iodanium perchlorate, 2299
 † 3-Hydroxypropanenitrile, *see* 2-Cyanoethanol, 1181
 † 1-Hydroxypropane, *see* Propanol, 1279
 2-Hydroxypropane-1,2,3-tricarboxylic acid, *see* Citric acid, 2389
 2-Hydroxypropanoic acid, *see* Lactic acid, 1234
N-Hydroxysuccinimide, 1469
 Hydroxyurea, 0477
 Hypochlorous acid, 3995
 Hyponitrous acid, 4470
 'Hypophosphorous acid', *see* Phosphinic acid, 4503

 † Imidazoline-2,4-dithione, 1141
 Imidodicarbonic diamide, 0874
 Iminobisacetoneitrile, 1474
 3,3'-Iminobispropylamine, 2587
 † 2-Iminohexafluoropropane, *see* Hexafluoroisopropylideneamine, 1067
 2-Indanecarboxaldehyde, 3288
 Indium, 4640
 Indium bromide, 0289
 Indium(II) oxide, 4641
 Indium(I) perchlorate, 4014
 Indium phosphide, 4642
 Iodic acid, 4424
 Iodimide, *see* Diiodamine, 4426

Iodinated poly(sulfur nitride), 4622
 Iodine, 4625
 Iodine azide, 4621
 Iodine bromide, 0254
 Iodine chloride, 4013
 Iodine dioxide trifluoride, 4334
 Iodine heptafluoride, 4378
 Iodine(III) perchlorate, 4140
 Iodine isocyanate, 0524
 Iodine pentafluoride, 4355
 Iodine triacetate, 2394
 Iodine trichloride, 4139
 Iodine(VII) oxide, 4628
 Iodine(V) oxide, 4627
 Iodoacetylene, 0670
 Iodobenzene, 2249
 4-Iodobenzenediazonium-2-carboxylate, 2639
 Iodobis(2,6,6-trimethylbicyclo [3.1.1]hept-3-yl)borane, *see*
 B-Iododiisopinocampheylborane, 3812
 1-Iodo-1,3-butadiyne, 1381
 † 2-Iodobutane, 1647
 Iododiborane, 0165
B-Iododiisopinocampheylborane, 3812
 4-Iodo-3,5-dimethylisoxazole, 1868
 2-Iodo-3,5-dinitrobiphenyl, 3450
 Iododisilane, 4543
 Iodoethane, 0858
 2-Iodoethanol, 0860
 2(2-Iodoethyl)-1,3-dioxolane, 1928
 Iodoform, 0376
 1-Iodohexa-2,4-diene, 2393
 Iodo-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-
 4,11-cyclotetradecadieneiron(II) perchlorate, 3723
 Iodomethane, 0445
 3-Iodo-4-methoxy-4,7,7-trimethylbicyclo[4.1.0]heptane, 3412
 Iodomethylmagnesium, *see* Methylmagnesium iodide, 0446
 † 1-Iodo-2-methylpropane, 1648
 † 2-Iodo-2-methylpropane, 1649
 Iodomethylzinc, *see* Methylzinc iodide, 0447
 † 2-Iodopentane, 1990
 1-Iodo-3-penten-1-yne, 1845
 3-Iodo-1-phenylpropyne, 3120
 † 2-Iodopropane, 1252
 † 3-Iodopropene, 1174
 3-Iodopropyne, 1105
 2-Iodosovinyl chloride, 0691

Iodosylbenzene, 2252
Iodosylbenzene tetrafluoroborate, 2295
4-Iodosyltoluene, 2753
4-Iodotoluene, 2750
† Iodotrimethylsilane, 1306
4-Iodylanisole, 2755
Iodylbenzene, 2253
Iodylbenzene perchlorate, *see* (Hydroxy)(oxo)(phenyl)- λ^3 -iodanium perchlorate, 2299
2-Iodylbenzoic acid, 2681
4-Iodyltoluene, 2754
2-Iodylvinyl chloride, 0692
Iridium, 4643
Iridium hexafluoride, 4362
Iridium(IV) oxide, 4644
Iron, 4388
Iron carbide, 0365
Iron disulfide, 4401
Iron(II) bromide, 0266
Iron(II) chelate of bis-*N,N*-(2-pentanone-4-ylidene)-1,3-diamino-2-hydroxypropane, 3616
Iron(II) chloride, 4061
Iron(II) hydroxide, 4392
Iron(III) bromide, 0286
Iron(III) chloride, 4133
Iron(III) hexacyanoferrate(4-), 3739
Iron(II,III) oxide, 4405
Iron(III) maleate, 3447
Iron(III) nitrate, 4397
Iron(II) iodide, 4395
Iron(III) oxalate, 2065
Iron(III) oxide, 4403
Iron(III) perchlorate, 4134
Iron(III) phosphinate, 4394
Iron(III) phthalate, 3844
Iron(III) sulfide, 4404
Iron(II) maleate, 1393
Iron(II) oxide, 4398
Iron(II) perchlorate, 4062
Iron(II) sulfate, 4399
Iron(II) sulfide, 4400
Iron–manganese alloy, *see* Ferromanganese, 4389
† Iron pentacarbonyl, *see* Pentacarbonyliron, 1814
Iron pyrites, *see* Iron disulfide, 4401
Iron–silicon alloy, *see* Ferrosilicon, 4390
Iron–titanium alloy, *see* Ferrotitanium, 4391

- † Isobutane, 1669
- † Isobutene, *see* 2-Methylpropene, 1582
- † Isobutyl acetate, 2502
- † Isobutylamine, 1728
- † Isobutyl bromide, *see* 1-Bromo-2-methylpropane, 1633
- † Isobutyl chloride, *see* 1-Chloro-2-methylpropane, 1639
- † Isobutyl formate, 1971
- † Isobutyl iodide, *see* 1-Iodo-2-methylpropane, 1648
- † Isobutyl peroxyacetate, 2512
- † *p*-Isobutylphenyl- α -methylacetaldehyde oxime, *see* 2-Methyl-2-[4-(2-methylpropyl)phenyl]oximinoethane, 3614
- † Isobutyl propionate, 2860
- † Isobutyl vinyl ether, 2490
- † Isobutyraldehyde, 1611
- † Isobutyronitrile, 1565
- † Isobutyrophenone, 3311
- † Isobutyryl chloride, 1558
- † Isocyanoamide, 0407
- † Isocyanoethane, *see* Ethyl isocyanide
- † Isocyanoethene, *see* Vinyl isocyanide, 1109
- † 2-Isocyanoethyl benzenesulfonate, 3143
- † 'Isodiazomethane', *see* Isocyanoamide, 0407
- † Isohexane, 2532
- † Isonicotinamidopentaammineruthenium(II) perchlorate, 2618
- † Isopentane, *see* 2-Methylbutane, 2000
- † Isopentanol, 2014
- † Isopentyl acetate, 2861
- † Isopentylamine, 2032
- † Isopentyl formate, 2503
- † Isopentyl isovalerate, 3361
- † Isopentyl nitrite, 1996
- † Isophthaloyl chloride, 2888
- † Isoprene diozonide, 1920
- † Isoprene, *see* 2-Methyl-1,3-butadiene, 1893
- † Isopropanol, *see* 2-Propanol, 1280
- † Isopropenyl acetate, 1916
- † 4-Isopropenyl-1-methylcyclohexene, *see* Limonene, 3337
- † Isopropyl acetate, 1972
- † Isopropylacrylaldehyde, *see* 2-3-Methyl-2-methylenebutanal, 2433
- † 2-Isopropylacrylaldehyde oxime, 2453
- † Isopropylamine, 1308
- † Isopropyl bromide, *see* 2-Bromopropane, 1242
- † Isopropyl butyrate, 2862
- † Isopropyl chloride, *see* 2-Chloropropane, 1244
- † Isopropyl chloroformate, 1560
- † Isopropyldiazene, 1274

- † Isopropyl formate, 1619
- Isopropyl hydroperoxide, 1283
- Isopropyl hypochlorite, 1247
- 6-Isopropylideneethiacyclodeca-3,8-diyne, 3527
- † Isopropyl iodide, *see* 2-Iodopropane, 1252
- † Isopropyl isobutyrate, 2863
- Isopropylisocyanide dichloride, 1561
- 5-Isopropyl-2-methyl-1,3-cyclohexadiene, *see* 1,5-*p*-Menthadiene, 3338
- † Isopropyl nitrate, 1266
- † Isopropyl nitrite, 1262
- † Isopropyl propionate, 2504
- † Isopropyl vinyl ether, 1957
- Isosorbide dinitrate, 2380
- Isosorbide mononitrate, *see* 1,4,3,6-Dianhydroglucitol 2-nitrate, 2396
- 3-Isothiocyanatopropene, *see* Allyl isothiocyanate, 1471
- † Isovaleraldehyde, 1958
- † Isoxazole, 1110

Ketene, 0722

Krypton difluoride, 4313

Lactic acid, 1234

Lactose, 3557

Lanthanum, 4677

Lanthanum carbide, 0991

Lanthanum dihydride, 4461

Lanthanum hexaboride, 0193

Lanthanum hydride, 4495

Lanthanum–nickel alloy, 4678

Lanthanum 2-nitrobenzoate, 3815

Lanthanum oxide, 4679

Lanthanum pentanickel hexahydride, 4552

Lanthanum picrate, 3741

Lead, 4882

Lead abietate, 3902

Lead acetate–lead bromate, 1540

Lead 2-amino-4,6-dinitrophenoxide, 3469

Lead 5,5'-azotetrazolide, 1020

Lead bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide), 3862

Lead bromate, 0278

Lead carbonate, 0558

Lead carbonate–lead hydroxide, 0726

Lead chloride, 4112

Lead chromate, 4243

Lead dioxide, *see* Lead(IV) oxide, 4834

Lead hyponitrite, 4745
Lead(II) azide, 4782
Lead(II) chlorate, 4105
Lead(II) chlorite, 4102
Lead(II) cyanide, 0999
Lead(II) fluoride, 4329
Lead(II) hypochlorite, 4098
Lead(II) imide, 4439
Lead(II) nitrate, 4749
Lead(II) nitrate phosphinate, 4468
Lead(II) oxide, 4824
Lead(II) phosphinate, 4531
Lead(II) phosphite, 4535
Lead(II) picrate, 3436
Lead(II) thiocyanate, 1000
Lead(II) trinitrosobenzene-1,3,5-trioxide, 3594
Lead(IV) acetate azide, 2402
Lead(IV) azide, 4790
Lead(IV) oxide, 4834
Lead methylenebis(nitramide), 0412
Lead nitride, 4753
Lead oleate, 3895
Lead pentaphosphide, 4881
Lead perchlorate, 4108
'Lead styphnate', *see* Lead 2,4,6-trinitroresorcinoxide, 2071
Lead sulfate, 4860
Lead tetrachloride, 4172
Lead-tin alloys, 4883
Lead 2,4,6-trinitroresorcinoxide, 2071
Lead-zirconium alloys, 4884
Limonene, 3337
Linolenic acid, *see* 9,12,15-Octadecatrienoic acid, 3771
Lithium, 4680
Lithium acetylide, 0992
Lithium aluminium hydride, *see* Lithium tetrahydroaluminate, 0075
Lithium amide, 4462
Lithium azide, 4685
Lithium benzenehexoxide, 2627
Lithium bis(trimethylsilyl)amide, 2594
Lithium borohydride, *see* Lithium tetrahydroborate, 0145
Lithium bromoacetylide, 0580
Lithium carbonate, 0533
Lithium chlorite, 4020
Lithium chloroacetylide, 0599
Lithium chloroethynide, *see* Lithium chloroacetylide, 0599
Lithium chromate, 4236

Lithium diazomethanide, 0379
Lithium diethylamide, 1686
Lithium dihydrocuprate, 4271
Lithium 2,2-dimethyltrimethylsilylhydrazide, 2041
Lithium diphenylhydridotungstate(2-), 3503
Lithium dithionite, 4687
Lithium ethynediolate, 0993
Lithium heptapotassium di(tetrasilicide), 4676
Lithium 1-heptynide, 2834
Lithium hexaazidocuprate(4-), 4278
Lithium hexamethylchromate(3-), 2590
Lithium hexaphenyltungstate(2-), 3892
Lithium hydrazide, 4496
Lithium hydride, 4432
Lithium-magnesium alloy, 4681
Lithium nitrate, 4684
Lithium nitride, 4688
Lithium 4-nitrothiophenoxide, 2177
Lithium octacarbonyltrickelate, 3109
Lithium oxide, 4686
Lithium pentahydrocuprate(4-), 4273
Lithium pentamethyltitanate-bis(2,2'-bipyridine), 2042
Lithium perchlorate, 4021
Lithium sodium nitroxylate, 4683
Lithium tetraazidoaluminate, 0081
Lithium tetraazidoborate, 0151
Lithium tetradeuteroaluminate, 0069
Lithium 3-(1,1,2,2-tetrafluoroethoxy)propynide, 1820
Lithium tetrahydroaluminate, 0075
Lithium tetrahydroborate, 0145
Lithium tetrahydrogallate, 4407
Lithium tetramethylborate, 1740
Lithium tetramethylchromate(II), 1744
Lithium 1,1,2,2-tetranitroethanediide, 0994
Lithium thiocyanate, 0532
Lithium-tin alloys, 4682
Lithium triethylsilylamide, 2578
Lithium trifluoropropynide, 1046
Lithium tripotassium tetrasilicide, 4675

Magnesium, 4690
Magnesium azide, 4694
Magnesium boride, 0168
Magnesium carbonate hydroxide, 0534
Magnesium chlorate, 4083
Magnesium chloride, 4081

Magnesium hydride, 4463
Magnesium hypochlorite, 4082
Magnesium–nickel hydride, 4464
Magnesium nitrate, 4693
Magnesium nitride, 4698
Magnesium nitrite, 4692
Magnesium oxide, 4695
Magnesium perchlorate, 4084
Magnesium permanganate, 4691
Magnesium phosphide, 4699
Magnesium phosphinate, 4517
Magnesium silicide, 4697
Magnesium sulfate, 4696
Magnesium tetrahydroaluminate, 0085
'Magnetite', *see* Iron(II,III) oxide, 4405
Maleic anhydride, 1404
Maleic anhydride ozonide, 1406
Maleic hydrazide, 1435
Maleimide, 1418
Malonic acid, 1151
† Malononitrile, 1078
Manganese, 4700
Manganese abietate, 3901
Manganese chloride trioxide, 4022
Manganese diazide hydroxide, 4433
Manganese dichloride dioxide, 4086
Manganese dioxide, *see* Manganese(IV) oxide, 4705
Manganese fluoride trioxide, 4301
Manganese(II) bis(acetylide), 1396
Manganese(II) chlorate, 4087
Manganese(II) chloride, 4085
Manganese(II) *N,N*-diethyldithiocarbamate, 3358
Manganese(III) azide, 4702
Manganese(II) nitrate, 4701
Manganese(II) oxide, 4704
Manganese(II) perchlorate, 4088
Manganese(II) phosphinate, 4519
Manganese(II) sulfide, 4706
Manganese(II) telluride, 4708
Manganese(II) tetrahydroaluminate, 0086
Manganese(IV) oxide, 4705
Manganese(IV) sulfide, 4707
Manganese phosphide, 4711
Manganese picrate hydroxide, 3742
Manganese tetrafluoride, 4343
Manganese trichloride oxide, 4141

Manganese trifluoride, 4335
Manganese(VII) oxide, 4709
1,5-*p*-Menthadiene, 3338
Mercaptoacetonitrile, 0767
Mercury, 4600
Mercury 5,5'-azotetrazolide, 0980
Mercury bis(chloroacetylde), 1356
'Mercury diazocarbide', *see* Poly(diazomethylenemercury), 0522
Mercury(I) azide, 4612
Mercury(I) bromate, 0271
Mercury(I) chlorite, 4080
Mercury(I) cyanamide, 0523
Mercury(I) fluoride, 4312
Mercury(I) hypophosphate, 4617
Mercury(II) acetylde, 0975
Mercury(II) amide chloride, 3999
Mercury(II) azide, 4604
Mercury(II) bromate, 0270
Mercury(II) bromide, 0269
Mercury(II) chloride, 4076
Mercury(II) chlorite, 4077
Mercury(II) cyanate, 0977
Mercury(II) cyanide, 0976
Mercury(II) *aci*-dinitromethanide, 0707
Mercury(II) formohydroxamate, 0804
Mercury(II) fulminate, 0978
Mercury(II) iodide, 4602
Mercury(II) nitrate, 4603
Mercury(II) 5-nitrotetrazolide, 0981
Mercury(II) oxalate, 0982
Mercury(II) oxide, 4605
Mercury(II) oxycyanide, *see* Dimercury dicyanide oxide, 0983
Mercury(II) perchlorate, 4078
Mercury(II) perchlorate . 6 (or 4)dimethyl sulfoxide, 4079
Mercury(II) *N*-perchlorylbenzylamide, 3651
Mercury(II) peroxybenzoate, 3637
Mercury(II) picrate, 3433
Mercury(II) sulfide, 4607
Mercury(II) thiocyanate, 0979
Mercury(I) nitrate, 4609
'Mercury(I) oxide' 4613
Mercury(I) thionitrosylate, 4610
Mercury nitride, 4615
Mercury peroxide, 4606
Mesitylene, 3162
O-Mesitylenesulfonylhydroxylamine, 3170

- † Methacrylaldehyde, 1522
- Methacrylic acid, 1530
- Methacryloyl chloride, *see* 2-Methyl-2-propenoyl chloride, 1457
- † Methanamine, *see* Methylamine, 0493
- Methanaminium nitrate, *see* Methylammonium nitrate, 0506
- † Methane, 0466
- Methaneboronic anhydride–pyridine complex, 0427
- Methaneperoxoic acid, *see* Peroxyformic acid, 0420
- Methanesulfinyl chloride, 0435
- Methanesulfonic acid, 0488
- Methanesulfonyl azide, 0465
- 2-Methanesulfonyl-4-nitrobenzenediazonium hydrogen sulfate, 2784
- Methanetellurol, 0490
- † Methanethiol, 0489
- Methanimidamide monoperoxylate, *see* Chloroformamidinium perchlorate, 0469
- † Methanol, 0484
- ‘Methazonic acid’, *see* 2-Nitroacetaldehyde oxime, 0809
- † Methoxyacetyl chloride, 1165
- † Methoxyacetylene, 1146
- 2-Methoxyaniline, 2816
- 3-Methoxyaniline, 2817
- 4-Methoxyaniline, 2818
- 2-Methoxyanilinium nitrate, 2829
- 4-Methoxybenzaldehyde, 2956
- 2-Methoxybenzenediazonium triiodide, 2757
- 4-Methoxybenzenediazonium triiodide, 2758
- 4-Methoxybenzoyl chloride, 2930
- 3-Methoxybenzylamine, 2997
- 4-Methoxybenzyl chloride, 2962
- Methoxycarbonylmethyl perchlorate, 1166
- 3-Methoxycarbonylpropen-2-yl trifluoromethanesulfonate, 2353
- Methoxy-1,3,5,7-cyclooctatetraene, 3149
- † Methoxycyclopropane, *see* Cyclopropyl methyl ether, 1608
- † Methoxyethane, *see* Ethyl methyl ether, 1278
- 2-Methoxyethanol, 1284
- † 2-Methoxyethanoyl chloride, *see* Methoxyacetyl chloride, 1165
- † Methoxyethene, *see* Methyl vinyl ether, 1221
- Z-1-(2-Methoxyethenyl)-2,3-dioxabicyclo[2.2.2]octa-5,7-diene, *see*
 1-(*cis*-Methoxyvinyl)-1,4-endoperoxy-2,5-cyclohexadiene, 3153
- 2-(2-Methoxyethoxy)ethanol, 2019
- † 2-Methoxyethylamine, 1312
- 2-Methoxyethylbenzene, 3164
- † 2-Methoxyethyl vinyl ether, 1973
- 1-Methoxyimidazole *N*-oxide, 1505
- 3-Methoxy-3-methyl-5,6-benzo-6*H*-1,2-dioxin, 3313

- 3-Methoxy-4-methyl-5,6-benzo-6*H*-1,2-dioxin, 3314
N-Methoxy-*N*-methylbenzylidenimmonium methylsulfate, 3334
 [(1-Methoxy-2-methylbutenyl)oxy]trimethylsilane, *see* 1-Methoxy-2-methyl-1-(trimethylsilyloxy)-1-butene, 3213
 2-Methoxy-1-methylethyl hydroperoxide, *see* 2-Methoxyprop-2-yl hydroperoxide, 1707
 1-Methoxy-2-methyl-1-(trimethylsilyloxy)-1-butene, 3213
 2-Methoxy-5-nitroaniline, 2802
 3-Methoxy-2-nitrobenzaldehyde, 2937
 3-Methoxy-2-nitrobenzoyl chloride, 2915
 4-Methoxy-3-nitrobenzoyl chloride, 2916
 3-Methoxy-2-nitrobenzoyldiazomethane, 3124
 4-Methoxy-1,3-phenylenediamine, 2826
 1-(4-Methoxyphenyl)-3-methyltriazene, 3000
 † 1-Methoxypropane, *see* Methyl propyl ether, 1703
 3-Methoxypropylamine, 1730
 2-Methoxyprop-2-yl hydroperoxide, 1707
 3-Methoxypropyne (Methyl propargyl ether), 1523
 5-Methoxy-1,2,3,4-thiatriazole, 0773
 1-Methoxy-3,4,5-trimethylpyrazole *N*-oxide, 2840
 1-(*cis*-Methoxyvinyl)-1,4-endoperoxy-2,5-cyclohexadiene, 3153
 † Methyl acetate, 1232
 2-Methylacryaldehyde oxime, 1568
 † Methyl acrylate, 1531
 Methylaluminiumbis(pentamethylcyclopentadienyltrimethyliridium), 3866
 Methylaluminium diiodide, 0423
 † Methylamine, 0493
 4-Methylaminobenzene-1,3-bis(sulfonyl azide), 2785
 1-Methylamino-1-methylthio-2-nitroethene, 1596
 Methylammonium chlorite, 0500
 Methylammonium nitrate, 0506
 Methylammonium nitrite, 0505
 Methylammonium perchlorate, 0501
 Methyl azide, 0459
 Methyl 2-azidobenzoate, 2939
 † 2-Methylaziridine, 1257
 Methyl benzenediazoate, 2799
 † Methylbenzene, *see* Toluene, 2791
 4-Methylbenzyl chloride, 2959
 Methylbismuth oxide, 0428
 Methylbis(η^2 -peroxo)rhenium oxide hydrate, 0497
 Methylborylene, 0425
 † Methyl bromide, *see* Bromomethane, 0429
 Methyl 4-bromobenzenediazoate, 2737
 † 2-Methyl-1,3-butadiene, 1893
 † 3-Methyl-1,2-butadiene, 1892

- † 2-Methylbutane, 2000
- † 2-Methylbutane-2-thiol, 2023
- † 3-Methylbutanethiol, 2024
- 3-Methylbutanol, *see* Isopentanol, 2014
- † 3-Methyl-2-butanone, 1959
- † 2-Methyl-1-butene, 1942
- † 2-Methyl-2-butene, 1943
- † 3-Methyl-1-butene, 1944
- † Methyl 2-butenolate, *see* Methyl crotonate, 1914
- † 2-Methyl-3-buten-2-ol, 1960
- † 2-Methylbuten-3-one, *see* Methyl isopropenyl ketone, 1911
- † 2-Methyl-1-buten-3-yne, 1858
- 3-Methyl-3-buten-1-ynyltriethyllead, 3413
- † *N*-Methylbutylamine, 2033
- † 3-Methyl-1-butyne, 1894
- Methyl 2-butyrate, 1881
- † 2-Methyl-3-buten-2-ol, 1910
- † Methyl butyrate, 1974
- Methylcadmium azide, 0431
- Methyl carbamate, 0868
- † Methyl carbonochloridate, *see* Methyl chloroformate, 0736
- † Methyl chloride, *see* Chloromethane, 0432
- † Methyl chloroformate, 0736
- Methylcopper, 0440
- † Methyl crotonate, 1914
- † Methylcyclobutane, 1945
- † Methylcyclohexane, 2854
- 3-Methylcyclohexanone, 2842
- 4-Methylcyclohexanone, 2843
- † 4-Methylcyclohexene, 2839
- Methylcyclopentadiene, 2363
- † Methylcyclopentane, 2461
- † Methylcyclopropane, 1581
- † Methyl cyclopropanecarboxylate, 1917
- Methyldiazene, 0474
- Methyl 3,3-diazido-2-cyanoacrylate, 1824
- 3-Methyldiazirine, 0808
- Methyl diazoacetate, 1138
- Methyldiborane, 0513
- † 4-Methyl-1,3-dioxane, 1975
- 1-(4-Methyl-1,3-diselenonylidene)piperidinium perchlorate, 3174
- O*-Methyl-*N,N*-disilylhydroxylamine, 0516
- Methylenebis(nitramine), 0483
- Methylenebis(3-nitramino-4-methylfuran), 2805
- 2-Methylenebutanal oxime, *see* 2-Ethylacrylaldehyde oxime, 1933
- † Methylene diisocyanate, *see* Diisocyanatomethane, 1079

- Methylenedilithium, 0401
 Methylenemagnesium, 0402
 4-Methylene-2-oxetanone, *see* Diketene
 † Methyl ethanoate, *see* Methyl acetate, 1232
 7-(1-Methylethylidene)-2,3-dioxabicyclo[2.2.1]hept-5-ene, *see* 1,4-Epidioxy-
 1,4-dihydro-6,6-dimethylfulvene, 2984
 (3(1-Methylethyl)-4-methyldiphosphazole), *see* 5-Methyl-1(1-methylethyl)-
 1,2,3-azadiphosphole, 2455
 † Methyl fluoride, *see* Fluoromethane, 0441
N-Methylformamide, 0866
 † Methyl formate, 0834
 † 2-Methylfuran, 1878
 † 2-Methylheptane, 3057
 † 3-Methylheptane, 3058
 2-(1-Methylheptyl)-4,6-dinitrophenyl crotonate, 3764
 † 2-Methylhexane, 2874
 † 3-Methylhexane, 2875
 † Methylhydrazine, 0503
 Methylhydrazinium nitrate, 0511
 Methyl hydroperoxide, 0486
 Methyl hypochlorite, 0434
 Methyl hypofluorite, 0442
 1-Methylimidazole, 1499
 Methyl iminioformate chloride, 0897
 Methyl iodide, *see* Iodomethane, 0445
 † Methyl isobutyrate, 1976
 † Methyl isocyanate, 0761
 Methyl isocyanide, 0759
 Methyl isocyanoacetate, 1470
 † Methyl isopropenyl ketone, 1911
 † Methyl isovalerate, 2505
 5-Methylisoxazole, 1468
 Methyllithium, 0452
 Methylmagnesium iodide, 0446
 Methylmercury perchlorate, 0433
 † Methyl methacrylate, 1915
 † *N*-Methylmethanamine, *see* Dimethylamine, 0941
 † Methyl methanoate, *see* Methyl formate, 0834
 Methyl 3-methoxycarbonylazocrotonate, 2830
 † 2-3-Methyl-2-methylenebutanal, 2433
 5-Methyl-1(1-methylethyl)-1,2,3-azadiphosphole, 2455
 † Methyl 2-methylpropenoate, *see* Methyl methacrylate, 1915
 2-Methyl-2-[4-(2-methylpropyl)phenyl]oximinoethane, 3614
N-Methyl-1-(methylthio)-2-nitroethenamine, *see* 1-Methylamino-1-
 methylthio-2-nitroethene, 1596
 † 4-Methylmorpholine, 1995

- N*-Methylmorpholine oxide, 1997
 4-Methylmorpholine-4-oxide, *see N*-Methylmorpholine oxide, 1997
 Methyl nitrate, 0457
 2-Methyl-1-nitratodimercurio-2-nitratomercuriopropane, 1592
 † Methyl nitrite, 0455
 N-Methyl-4-nitroaniline, 2800
 2-Methyl-5-nitroaniline, 2801
 N-Methyl-*p*-nitroanilinium 2(*N*-methyl-*N*-*p*-nitrophenylaminosulfonyl) ethylsulfate, 3713
 3'-Methyl-2-nitrobenzanilide, 3645
 Methyl 2-nitrobenzenediazoate, 2782
 2-Methyl-5-nitrobenzenesulfonic acid, 2772
 4-Methyl-3-nitrobenzenesulfonic acid, 2773
 2-Methyl-5-nitrobenzimidazole, 2940
 3-Methyl-2-nitrobenzoyl chloride, 2913
 3-Methyl-4-nitro-1-buten-3-yl acetate, 2835
 3-Methyl-4-nitro-2-buten-1-yl acetate, 2836
 1-Methyl-3-nitroguanidinium nitrate, 0947
 1-Methyl-3-nitroguanidinium perchlorate, 0939
 2-Methyl-4-nitroimidazole, 1475
 'Methylnitrolic acid', *see* Nitrooximinomethane, 0409
 1-Methyl-3-nitro-1-nitrosoguanidine, 0876
 3-Methyl-4-nitrophenol, 2766
 4-Methyl-2-nitrophenol, 2767
 2-Methyl-2-nitropropane, *see tert*-Nitrobutane, 1660
 3-Methyl-4-nitropyridine *N*-oxide, 2317
 N-Methyl-*N*-nitrosoarea, 0875
 Methylnitrothiophene, 1850
 † 2-Methylnonane, 3366
 S-7-Methylnonylthiuronium picrate, 3737
 † 2-Methyloctane, 3206
 † 3-Methyloctane, 3207
 † 4-Methyloctane, 3208
 2-Methyl-3,5,7-octatriyn-2-ol, 3135
 † 2-Methyl-2-oxazoline, 1569
 † Methyloxirane, *see* Propylene oxide, 1225
 Methyloxobismuthine, *see* Methylbismuth oxide, 0428
 † 2-Methyl-1,3-pentadiene, 2416
 † 4-Methyl-1,3-pentadiene, 2417
 † 2-Methylpentanal, 2491
 † 3-Methylpentanal, 2492
 † 3-Methylpentane, 2533
 † 2-Methylpentane, *see* Isohexane, 2532
 † 2-Methyl-2-pentanol, 2544
 † 2-Methyl-3-pentanone, 2493
 † 3-Methyl-2-pentanone, 2494

† 4-Methyl-2-pentanone, 2495
 † 2-Methyl-1-pentene, 2462
 † 4-Methyl-1-pentene, 2463
 † *cis*-4-Methyl-2-pentene, 2464
 † *trans*-4-Methyl-2-pentene, 2465
 3-Methyl-2-penten-4-yn-1-ol, 2384
 Methyl perchlorate, 0436
 1-(1-Methyl-1-phenylethyl)-4-(2-propynyloxy)benzene, 3760
N-(3-Methylphenyl)-2-nitrobenzimidyl chloride, 3641
N-(Methylphenylphosphinoyl)hydroxylamine, 2823
 Methylphosphine, 0498
 † 1-Methylpiperidine, 2522
 † 2-Methylpiperidine, 2523
 † 3-Methylpiperidine, 2524
 † 4-Methylpiperidine, 2525
N-Methylpiperidine *N*-oxide, 2527
 † Methyl pivalate, 2506
 Methylpotassium, 0448
 † 2-Methylpropanal, *see* Isobutyraldehyde, 1611
 † 2-Methylpropane, *see* Isobutane, 1669
 † 2-Methyl-2-propanethiol, 1716
 † 2-Methylpropanethiol, 1715
 † 2-Methylpropanonitrile, *see* Isobutyronitrile, 1565
 † 2-Methylpropanoyl chloride, *see* Isobutyryl chloride, 1558
 † 2-Methylpropenal, *see* Methacrylaldehyde, 1522
 † 2-Methylpropene, 1582
 2-Methyl-1-propene-1-one, *see* Dimethylketene, 1518
 † Methyl propenoate, *see* Methyl acrylate, 1531
 2-Methylpropenoic acid, *see* Methacrylic acid, 1530
 2-Methyl-2-propenoyl chloride, 1457
 † Methyl propiolate, 1442
 † Methyl propionate, 1620
 † Methyl propyl ether, 1703
 3-Methyl-5-propyl-1,2,4-trioxolane, *see trans*-2-Hexene ozonide, 2513
 † Methyl propynoate, *see* Methyl propiolate, 1442
 2-Methylpyridine, 2355
 † 1-Methylpyrrole, 1887
 † *N*-Methylpyrrolidine, 1992
 1-Methylsilacyclopenta-2,4-diene, 1921
 Methylsilane, 0510
 1-Methylsilole, *see* 1-Methylsilacyclopenta-2,4-diene, 1921
 Methylsilver, 0421
 Methylsodium, 0463
 Methyl spiro[5-bromobicyclo[2.2.1]heptane 2,2' 1,3-dioxolane]-7-carboxylate,
 3405
 Methylstibine, 0499

- † 2-Methyltetrahydrofuran, 1961
- 2-Methyltetrazole, 0814
- S-Methylthioacetohydroximate, 1260
- † 2-Methylthiophene, 1882
- 4-Methyl-2,4,6-triazatricyclo[5.2.2.0^{2,6}]undeca-8-ene-3,5-dione, 3160
- 1-Methyl-1,2,3-triazole, 1189
- Methyl trichloroacetate, 1096
- † Methyltrichlorosilane, 0439
- Methyltrifluoromethyldioxirane, 1102
- Methyltrifluoromethyltrichlorophosphorane, 0742
- Methyl trifluorovinyl ether, 1101
- 3-Methyl-2,4,6-trinitrophenol, 2703
- 3-Methyl-1,2,4-trioxolane, 1235
- 3-(3-Methyl-1,2,4-trioxolan-3-yl)-1,2,4-trioxolane, *see* Isoprene diozonide, 1920
- Methyltriphenylphosphonium permanganate, 3784
- † Methyl valerate, 2507
- † Methyl vinyl ether, 1221
- † Methyl vinyl ketone, *see* 1-Buten-3-one, 1515
- 2-Methyl-5-vinyltetrazole, 1511
- Methylzinc iodide, 0447
- † 2-Methylpropylamine, *see* Isobutylamine, 1728
- 'Millon's base anhydride', *see* Poly(dimercuryimmonium hydroxide), 4422
- Molybdenum, 4712
- Molybdenum azide tribromide, 0288
- Molybdenum diazide tetrachloride, 4164
- Molybdenum hexafluoride, 4365
- Molybdenum hexamethoxide, 2596
- Molybdenum(IV) oxide, 4716
- Molybdenum(IV) sulfide, 4719
- Molybdenum nitride, 4714
- Molybdenum pentachloride, 4180
- Molybdenum(VI) oxide, 4717
- Monocaesium acetylide, 0661
- Monofluoroxonium hexafluoroarsenate, 0097
- Monolithium acetylide, 0673
- Monolithium acetylide-ammonia, 0674
- Monoperoxysuccinic acid, 1542
- Monopotassium acetylide, 0671
- Monopotassium perchlorylamide, 3994
- Monorubidium acetylide, 0685
- Monosilver acetylide, 0650
- Monosodium acetylide, 0684
- Monosodium cyanamide, 0382
- Morpholine, 1657

4-Morpholinesulfenyl chloride, 1586
Morpholinium perchlorate, 1680

Naphthalene, 3254

1-Naphthalenediazonium perchlorate, 3246
2-Naphthalenediazonium perchlorate, 3247
1-Naphthalenediazonium salts, 3251
2-Naphthalenediazonium salts, 3252
2-Naphthalenediazonium trichloromercurate, 3248

1-Naphthylamine, 3264

2-Naphthylamine, 3265

1-Naphthyl isocyanate, 3390

Naphthylsodium, 3253

1-(2-Naphthyl)-3-(5-tetrazolyl)triazene, 3394

(1-Naphthyl)-1,2,3,4-thiatriazole, 3391

Neodymium, 4819

Neodymium perchlorate . 2acetonitrile, 4148

† Neopentane, *see* 2,2-Dimethylpropane, 1999

† Neopentylamine, *see* 2,2-Dimethylpropylamine, 2030

Neptunium hexafluoride, 4366

Nickel, 4820

Nickel azide, 4777

Nickel chlorite, 4093

Nickel 2,4-dinitrophenoxide hydroxide, 3745

Nickel(II) cyanide, 0997

Nickel(III) oxide, 4823

Nickel(II) nitrate, 4743

Nickel(II) oxide, 4821

Nickel(IV) oxide, 4822

Nickel 2-nitrophenoxide, 3464

Nickelocene, 3287

Nickel perchlorate, 4094

Nickel picrate, 3435

† Nickel tetracarbonyl, *see* Tetracarbonylnickel, 1805

Niobium, 4817

Niobium(V) oxide, 4818

Nitric acid, 4436

Nitric amide (Nitramide), 4471

'Nitric oxide', *see* Nitrogen oxide, 4724

Nitrilotris(oxiranemethane), 3181

Nitritonitrosylnickel, 4742

Nitroacetaldehyde, 0763

2-Nitroacetaldehyde oxime, 0809

4-Nitroacetanilide, 2950

† Nitroacetone, 1184

Nitroacetonitrile, *see* 2-Nitroethanonitrile, 0711

2-Nitroacetophenone, 2936
1-(Nitroamino)benzotriazole, 2283
2-(*N*-Nitroamino)pyridine *N*-oxide, 1851
4-(*N*-Nitroamino)pyridine *N*-oxide, 1852
5-*N*-Nitroaminotetrazole, 0415
4-Nitroamino-1,2,4-triazole, 0777
2-Nitroaniline, 2313
3-Nitroaniline, 2314
4-Nitroaniline, 2315
4-Nitroaniline-2-sulfonic acid, 2319
4-Nitroanilinium perchlorate, 2350
2-Nitroanisole, 2768
1-Nitroanthraquinone, 3624
2-Nitrobenzaldehyde, 2686
3-Nitrobenzaldehyde, 2687
4-Nitrobenzaldehyde, 2688
Nitrobenzene, 2262
(Nitrobenzene)(cyclopentadienyl)iron(II) hexafluorophosphate, 3395
4-Nitrobenzenediazonium azide, 2209
3-Nitrobenzenediazonium chloride, 2151
4-Nitrobenzenediazonium nitrate, 2205
3-Nitrobenzenediazonium perchlorate, 2155
2-Nitrobenzenediazonium salts, 2201
4-Nitrobenzenediazonium salts, 2202
2-Nitrobenzenediazonium tetrachloroborate, 2128
4-Nitrobenzenesulfinyl azide, 2204
3-Nitrobenzenesulfonic acid, 2268
2-Nitrobenzoic acid, 2693
3-Nitrobenzoic acid, 2694
4-Nitrobenzoic acid, 2695
2-Nitrobenzonitrile, 2662
2-Nitrobenzoyl chloride, 2653
3-Nitrobenzoyl nitrate, 2664
2-Nitrobenzyl alcohol, 2769
3-Nitrobenzyl alcohol, 2770
4-Nitrobenzyl alcohol, 2771
2-Nitrobenzyl bromide, 2710
2-Nitrobenzyl chloride, 2713
4-Nitrobenzyl chloride, 2714
tert-Nitrobutane, 1660
4-Nitro-1-butene, 1572
1-Nitrocyclohepta-1,3,5-triene, 2761
1-Nitro-3-(2,4-dinitrophenyl)urea, 2705
4-Nitrodiphenyl ether, 3472
Nitroethanal, *see* Nitroacetaldehyde, 0763
Nitroethane, 0869

2-Nitroethanol, 0871
2-Nitroethanol, sodium salt, 0807
2-Nitroethanonitrile, 0711
2-Nitroethoxyethane, *see* Ethyl 2-nitroethyl ether, 1662
'Nitroform', *see* Trinitromethane, 0385
Nitrogen chloride difluoride, 3978
Nitrogen dioxide, 4725
Nitrogen dioxide, *see* Dinitrogen tetraoxide, 4747
Nitrogen (Gas), 4734
Nitrogen iodide, *see* Nitrogen triiodide, 4633
Nitrogen (Liquid), 4735
Nitrogen oxide, 4724
Nitrogen tribromide hexaammoniate, 0290
Nitrogen trichloride (Trichloramine), 4143
Nitrogen trifluoride, 4336
Nitrogen triiodide–ammonia, 4635
Nitrogen triiodide–silver amide, 4634
Nitrogen triiodide, 4633
Nitroguanidine, 0482
miso-Nitroindane, 3140
4-Nitroisopropylbenzene, 3157
3-Nitro-2-isoxazoline, 1139
Nitromesitylene, 3158
Nitromethane, 0456
N-Nitromethanimine, 0408
N-Nitromethylamine, 0478
1-Nitronaphthalene, 3249
5-Nitro-2-(nitroimino)hexahydro-1,3,5-triazine, 1218
Nitronium perchlorate, 4028
Nitronium tetrafluoroborate, 0131
1-Nitro-1-oximinoethane, 0810
Nitrooximinomethane, 0409
3-Nitroperchlorylbenzene, 2143
2-Nitrophenol, 2265
3-Nitrophenol, 2266
4-Nitrophenol, 2267
2-(4-Nitrophenoxyazo)benzoic acid, 3601
4-Nitrophenylacetic acid, 2938
2-Nitrophenylacetyl chloride, 2914
3-Nitrophenylacetylene, 2905
N-(2-Nitrophenyl)-1,2-diaminoethane, 3001
3-(2-Nitrophenyl)-2-hydroxyiminopropanoic acid, 3131
1-(2-Nitrophenyl)-5-phenyltetrazole, 3602
4-Nitrophenylphosphorodichloridate, 2158
4-Nitrophenylpropan-2-one, 3142
2-Nitrophenylpropionic acid, 3113

o-Nitrophenylpyruvic acid oxime, *see* 3-(2-Nitrophenyl)-2-hydroxyiminopropanoic acid, 3131
2-Nitrophenylsulfonyldiazomethane, 2700
3-Nitrophthalic acid, 2906
5-Nitro-2-picryltetrazole, 2638
4-Nitro-1-picryl-1,2,3-triazole, 2886
1-Nitropropane, 1263
2-Nitropropane, 1264
† 1-Nitro-2-propanone, *see* Nitroacetone, 1184
2-Nitropropene, 1183
3-Nitropropiofenone, 3141
N-Nitro-4-pyridinamine 1-oxide, *see* 4-(*N*-Nitroamino)pyridine *N*-oxide, 1852
4-Nitropyridine *N*-oxide, 1833
5-Nitrosalicylhydrazide, 2778
N-Nitrosoacetanilide, 2949
Nitrosobenzene, 2261
N-Nitrosodiphenylamine, 3486
N-Nitrosoethyl-2-hydroxyethylamine, 1691
Nitrosoguanidine, 0481
N-Nitroso-6-hexanelactam, 2424
1-Nitroso-2-naphthol, 3250
2-Nitrosophenol, 2263
4-Nitrosophenol, 2264
Nitrosotrifluoromethane, 0345
Nitrosyl azide, 4766
Nitrosyl chloride, 4023
Nitrosyl cyanide, 0541
Nitrosyl fluoride, 4302
Nitrosyl perchlorate, 4027
Nitrosylruthenium trichloride, 4144
Nitrosylsulfuric acid, 4438
Nitrosyl tetrafluorochlorate, 3985
Nitrosyl tribromide, *see* Tribromamine oxide, 0291
Nitroterephthalic acid, 2907
5-Nitrotetrazole, 0387
4-Nitrothiophene-2-sulfonyl chloride, 1388
2-Nitrotoluene, 2763
4-Nitrotoluene, 2764
miso-Nitrotoluene, 2762
3-Nitro-1,2,4-triazolone, 0716
Nitrourea, 0460
Nitrous acid, 4435
'Nitrous oxide', *see* Dinitrogen oxide, 4744
Nitryl chloride, 4025
Nitryl fluoride, 4303

- Nitryl hypochlorite, 4026
 Nitryl hypofluorite, 4304
 Nitryl nitrate, *see* Dinitrogen pentaoxide, 4748
 Nitryl perchlorate, 4029
 Nonacarbonyldiiron, 3111
 † Nonane, 3209
 Nonaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]trihexaconta-
 4,6,11,13,18,20,25,27,32,39,41,46,48,53,55,60,62-octadecayne, 3918
 2-Nonen-4,6,8-triyn-1-al, 3112
 Nopinone, *see* 6,6-Dimethylbicyclo[3.1.1]heptan-2-one, 3175
 Norspermidine, *see* 3,3'-Iminobispropylamine, 2587
- Octaammine- μ -hydroxy[μ -(peroxy-*O* : *O'*)]dirhodium(3+) perchlorate, 4138
 Octaammine- μ -hydroxy[μ -superoxido-*O*,*O'*)]dirhodium(4+) nitrate, 4597
 (Octacarbondioxide) pentamer, 3904
 Octacarbondioxide tetramer, 3884
 Octacarbonyldicobalt, 2881
 Octachlorotrisilane, 4194
 1,4-Octadecanolactone, 3773
 9,12,15-Octadecatrienoic acid, 3771
cis-9-Octadecenoic acid, 3774
cis-9-Octadecenoyl chloride, *see* Oleoyl chloride, 3772
 5,6,7,8,13,14,15,16-Octadehydridibenzo[*a,g*]cyclododecene, *see*
 Dibenzocyclododeca-1,7-dienetetrayne, 3786
 † 1,7-Octadiene, 3021
 Octafluoroadipamide, 2170
 Octahydro[24]annulene, *see* 1,7,13,19-Cyclotetracosatetraene-
 3,5,9,11,15,17,21,23-octayne, 3843
 Octahydro-2,5-bis(nitroimino)imidazo[4,5-*d*]imidazole, 1514
 Octahydro-1,3,4,7,8,10-hexanitro-5,2,6-(iminomethenimino-1*H*-
 imidazo[4,5-*b*]pyrazine), *see* Hexanitrohexaazaisowurtzitane, 2325
 Octakis(trifluorophosphine)dirhodium, 4387
 1,4,8,11,15,18,22,25-Octamethyl-29*H*,31*H*-tetrabenzo[*b.g.l.q*]
 porphinato(2-)-cobalt(II), 3910
 † Octane, 3059
 Octaspiro[2.4.2.4.2.4.2.4.2.4.2.4.2.4.2.4]hexapentaconta-
 4,6,11,13,18,20,25,27,32,34,39,41,46,48,53,55-hexadecayne, 3915
 Octatetrayne-1,8-dicarboxylic acid, 3235
 1,3,7-Octatrien-5-yne, 2944
 † 1-Octene, 3040
 † 2-Octene, 3041
 Octylsodium, 3053
 † 1-Octyne, 3022
 † 2-Octyne, 3023
 † 3-Octyne, 3024
 † 4-Octyne, 3025

Oleic acid, *see cis-9-Octadecenoic acid*, 3774
 Oleoyl chloride, 3772
 Oligo(octacarbondioxide), 3108
 1,5-(or 1,8-)Bis(dinitrophenoxy)-4,8-(or 4,5-)dinitroanthraquinone, 3860
 1(2', 3', or 4'-Diazoniophenyl)-2,4,6-triphenylpyridinium diperchlorate, 3870
 3- or 4-Methoxy-5,6-benzo-6*H*-1,2-dioxin, 3154
 Orthoperiodic acid, 4542
 Orthophosphoric acid, 4505
 2-, 3- or 4-Trifluoromethylphenylmagnesium bromide, 2643
 Osmium, 4873
 Osmium hexafluoride, 4370
 Osmium(IV) oxide, 4833
 Osmium(VIII) oxide, 4858
 † 6-Oxabicyclo[3.1.0]hexane, *see Cyclopentene oxide*, 1906
 Oxalic acid, 0725
 Oxalyl chloride, *see Oxalyl dichloride*, 0605
 Oxalyl dibromide, 0583
 Oxalyl dichloride, 0605
 1,4-Oxathiane, 1613
 † Oxazole, 1111
 Oxepin-3,6-endoperoxide, *see endo-2,3-Epoxy-7,8-dioxabicyclo*
 [2.2.2]oct-5-ene, 2336
 † Oxetane, 1222
 2-Oximino-butane, *see 2-Butanone oxime*, 1654
 4-Oximino-4,5,6,7-tetrahydrobenzofurazan *N*-oxide, 2357
 Oxiranecarboxaldehyde oxime, *see 2,3-Epoxypropionaldehyde oxime*, 1182
 † Oxirane, *see Ethylene oxide*, 0829
 Oxiranemethanol, *see 2,3-Epoxypropanol*, 1229
 † Oxiranemethanol nitrate, *see 2,3-Epoxypropyl nitrate*, 1186
 Oxoacetic acid, *see Glyoxylic acid*, 0724
 Oxobis[aqua(oxo)diperoxorhenium(VII)], 4537
 μ -Oxo-*I,I*-bis(trifluoroacetato-*O*)-*I,I*-diphenyldiodine(III), 3698
 Oxodiperoxodipiperidinechromium(VI), 3367
 Oxodiperoxodipyridinechromium(VI), 3274
 Oxodiperoxodi(pyridine *N*-oxide)molybdenum, 3279
 Oxodiperoxodi(pyridine *N*-oxide)tungsten, 3283
 Oxodiperoxodiquinolinechromium(VI), 3748
 Oxodiperoxomolybdenum-hexamethylphosphoramidate, 4718
 Oxodiperoxopyridinechromium *N*-oxide, 1844
 (Oxidiperoxy(1,3-dimethyl-2,4,5,6-tetrahydro-2-1*H*)-
 pyrimidinone)molybdenum, 2472
 Oxodiperoxy(pyridine)(1,3-dimethyl-2,4,5,6-tetrahydro-2-1*H*)-
 (pyrimidinone)molybdenum, 3410
 Oxodisilane, 4530
 Oxopropanedinitrile, 1341
 2-Oxopropanoic acid, *see Pyruvic acid*, 1150

Oxosilane, 4476
 1-Oxo-1,2,3,4-tetrahydronaphthalene, 3289
 Oxybis[aqua-5,7,7,12,14,14-hexamethyl-1,4,8,11-tetraaza-4,11-cyclotetradecadieneiron(II)] tetraperchlorate, 3882
 1,1'-Oxybis-2-azidoethane, 1604
 4,4'-Oxybis(benzenesulfonylhydrazide), 3525
 Oxybis[bis(cyclopentadienyl)titanium] 3803
 Oxybis(*N,N*-dimethylacetamidetriphenylstibonium) perchlorate, 3911
 [1,1'-Oxybis[ethane]]tris(pentafluorophenyl)aluminium, 3829
 [1,1'-Oxybis[ethane]]tris(trimethyl silyl)aluminium, *see* Tris(trimethylsilyl)aluminium etherate, 3620
 † 1,1'-Oxybisethene, *see* Divinyl ether, 1519
 2,2'-Oxybis(ethyl nitrate), 1599
 2,2'-Oxybis(iminomethylfuran) mono-*N*-oxide, 3260
 † Oxybismethane, *see* Dimethyl ether, 0919
 1,1-Oxybis(4-methylphenyldiazeno), *see* Bis(toluenediazo) oxide, 3654
 Oxybisphenyliodonium bistetrafluoroborate, 3475
 Oxygen difluoride, 4317
 Oxygen (Gas), 4831
 Oxygen (Liquid), 4832
 Oxynitrotriazole, *see* 3-Nitro-1,2,4-triazolone, 0716
 Ozobenzene, *see* Benzene triozone, 2343
 Ozone, 4846

 Palladium, 4885
 Palladium(II) acetate, 1541
 Palladium(II) azide, 4783
 Palladium(II) azidodithioformate, 1015
 Palladium(III) oxide, 4848
 Palladium(II) oxide, 4825
 Palladium(IV) oxide, 4835
 Palladium tetrafluoride, 4347
 Palladium trifluoride, 4341
 Paraformaldehyde, 0417
 Pentaammineaquacobalt(III) chlorate, 4124
 Pentaammineazidoruthenium(III) chloride, 4073
 Pentaamminechlorocobalt(III) perchlorate, 4123
 Pentaamminechlororuthenium chloride, 4137
 Pentaamminedinitrogenosmium(II) perchlorate, 4072
 Pentaamminedinitrogenruthenium(II) salts, 4596
 Pentaamminenitratocobalt(III) nitrate, 4207
 Pentaamminenitrochromium(III) nitrate, 4233
 Pentaamminephosphinatochromium(III) perchlorate, 4053
 Pentaamminephosphinatocobalt(III) perchlorate, 4050
 Pentaamminepyrazineruthenium(II) perchlorate, 1792
 Pentaamminepyridineruthenium(II) perchlorate, 2048

- Pentaamminethiocyanatocobalt(III) perchlorate, 0519
 Pentaamminethiocyanatoruthenium(III) perchlorate, 0520
 Pentaazidophosphorane, 4793
 Pentaborane(11), 0189
 † Pentaborane(9), 0188
 Pentaborane(9) diammoniate, *see* Diammineboronium heptahydrotetraborate, 0190
 † Pentacarbonyliron, 1814
 Pentachloroethane, 0660
 Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane, 2942
 Pentacyclo[4.2.0.0^{2,5}.0^{3,8}.0^{4,7}]octane-1,2-dicarboxylic acid, 3261
 † 1,2-Pentadiene, 1895
 † 1,3-Pentadiene, 1896
 † 1,4-Pentadiene, 1897
 1,4-Pentadien-3-one, *see* Divinyl ketone, 1877
 1,3-Pentadiyne, 1825
 1,3-Pentadiyn-1-ylcopper, 1819
 1,3-Pentadiyn-1-ylsilver, 1815
 Pentaerythritol, 2021
 1,2,3,-4,5-Pentafluorobicyclo[2.2.0]hexa-2,5-diene, 2069
 1,2,3,-5,6-Pentafluorobicyclo[2.2.0]hexa-2,5-diene, 2070
 Pentafluoroethylithium, 0636
 Pentafluoroguanidine, 0359
 Pentafluoroiodosylbenzene, 2058
 Pentafluoromethanamine, *see* *N,N*-Difluorotrifluoromethylamine, 0357
 Pentafluoroorthoselenic acid, 4354
 Pentafluorophenylaluminium dibromide, 2053
 Pentafluorophenyllithium, 2059
 Pentafluoropropionyl fluoride, 1054
 Pentafluoropropionyl hypochlorite, 1034
 Pentafluoropropionyl hypofluorite, 1056
 Pentafluoroselenium hypofluorite, 4368
 Pentafluorosulfur hypofluorite, 4367
 Pentafluorosulfur peroxyacetate, 0755
 Pentafluorosulfur peroxyhypochlorite, 3990
 Pentafluorotellurium hypofluorite, 4369
 Pentakis(dimethylamino)tantalum, 3384
 Pentamethylbismuth, 2040
 Pentamethylcyclopentadienyl-bis(thiolatothiatriazolyl)rhodium, 3534
 3,3-Pentamethylenediazirine, 2422
 Pentamethyltantalum, 2046
 † Pentanal, *see* Valeraldehyde, 1966
 † Pentane, 2001
 Pentanesulfonic acid, 2020
 † Pentanethiol, 2025
 Pentanitroaniline, 2093

† *tert*-Pentanol, 2015
 † 2-Pentanone, 1962
 † 3-Pentanone, 1963
 Pentasilver diamidophosphate, 0046
 Pentasilver diimidotriphosphate, 0045
 Pentasilver orthodiamidophosphate, 0044
 Pentaspiro[2.4.2.4.2.4.2.4.2.4]pentatriaconta-4,6,11,13,18,20,25,27,32,34-decayne, 3887
 Pentasulfur hexanitride, 4784
 Pentazole, 4443
 † 1-Pentene, 1946
 † 2-Pentene, 1947
trans-2-Pentene ozonide, 1982
 † 4-Penten-1-ol, 1964
 2-Penten-4-yn-3-ol, 1879
 † 2-Pentyl acetate, 2865
 † Pentyl acetate, 2864
 † Pentylamine, 2034
 α -Pentylcinnamaldehyde (3-Phenyl-2-pentylpropenal), 3663
 1-Pentyl methanesulfonate, 2550
 † Pentyl nitrite, 1998
 † 1-Pentyne, 1898
 † 2-Pentyne, 1899
 Perbromyl fluoride, 0240
 Perchloratotris(triethylphosphine)palladium(II) perchlorate, 3778
 Perchloric acid, 3998
 Perchlorobenzene, *see* Hexachlorobenzene, 2055
 Perchlorylbenzene, 2236
 Perchloryl fluoride, 3974
 Perchloryl hypofluorite, *see* Fluorine perchlorate, 3976
 Perchloryl perchlorate, 4107
N-Perchlorylpiperidine, 1949
 Perfluoro-1-aminomethylguanidine, 0647
C,N-Perfluoro-*N*-aminomethyltriaminomethane, 0649
 Perfluorobutadiene, 1364
 Perfluoro-*tert*-butanol, 1380
 Perfluoro-*tert*-butyl peroxyhypofluorite, 1373
 Perfluoro-*N*-cyanodiaminomethane, 0637
 Perfluoroethylithium, *see* Pentafluoroethylithium, 0636
 Perfluoroformamidine, 0350
 Perfluorohexyl iodide, 2062
 Perfluoro-*tert*-nitrosobutane, 1370
 Perfluorosuccinic acid, 1391
 † Perhydroazepine, 2526
 Perhydro-9b-boraphenalene, 3550
 Periodic acid, 4425

Permanganic acid, 4434
 μ -Peroxo-bis[ammine(2,2',2''-triaminotriethylamine)cobalt(III)](4+)
perchlorate, 3591
Peroxydisulfuric acid, 4482
Peroxydisulfuryl difluoride, 4328
Peroxymonophosphoric acid, 4506
Peroxymonosulfuric acid, 4481
Peroxonitric acid, 4437
Peroxyacetic acid, 0837
Peroxyacetyl nitrate, 0766
Peroxyacetyl perchlorate, 0737
Peroxybenzoic acid, 2733
Peroxy-2-butenic acid, *see* Peroxycrotonic acid, 1535
3-Peroxy-camphoric acid, 3348
Peroxycrotonic acid, 1535
Peroxyformic acid, 0420
Peroxyfuroic acid, 1837
Peroxyhexanoic acid, 2514
Peroxypropionic acid, 1236
Peroxypropionyl nitrate, 1187
Peroxypropionyl perchlorate, 1167
Peroxytrifluoroacetic acid, 0666
Perrhenyl chloride, *see* Rhenium chloride trioxide, 4045
Perylenium perchlorate, 3790
1,10-Phenanthroline-5,6-dione, 3578
Phenethyl alcohol, *see* Benzeneethanol, 2983
Phenol, 2330
Phenoxyacetylene, 2920
3-Phenoxy-1,2-epoxypropane, 3150
Phenylacetonitrile, 2935
Phenylacetylene, 2912
Phenylaminothia-2,3,4-triazole, *see* *N*-Phenyl-1,2,3,4-thiatriazolamine, 2729
Phenyl azide, 2271
 α -Phenylazobenzyl hydroperoxide, 3609
 α -Phenylazo-4-bromobenzyl hydroperoxide, 3607
1-Phenylazocyclohexyl hydroperoxide, 3539
 α -Phenylazo-4-fluorobenzyl hydroperoxide, 3608
N-Phenylazopiperidine, 3407
1-Phenylbiguanidinium hydrogen dichromate, 3019
1-Phenylboralane, 3317
1-Phenyl-3-*tert*-butyltriazene, 3335
Phenylchlorodiazirine, 2673
Phenyldiazidomethane, 2730
Phenyldiazomethane, 2725
2-Phenyl-1,1-dimethylethyl hydroperoxide, 3332
1-(4-Phenyl-1,3-diselenolydene)piperidinium perchlorate, 3659

2,2'-[1,4-Phenylenebis(azidomethylidene)]bis(propanedinitrile), 3622
p-Phenylenediamine, *see* 1,4-Diaminobenzene, 2371
 2-Phenylethyl isocyanate, 3139
 † Phenylglyoxal, 2922
 Phenylgold, 2221
 Phenylhydrazine, 2373
N-Phenylhydroxylamine, 2356
 Phenylhydroxylaminium chloride, 2366
 9-Phenyl-9-iodafluorene, 3747
 Phenyliodine(III) chromate, 2247
 Phenyliodine(III) nitrate, 2251
 Phenyl isocyanate, 2685
 3-Phenyl-5-isoxazolone, 3122
 Phenyllithium, 2259
 Phenylmagnesium bromide, 2225
 1-Phenyl-2-methoxydiazene, *see* Methyl benzenediazoate, 2799
 Phenyl, phenylethynyliodonium perchlorate, 3635
 Phenylphosphine, 2360
 Phenylphosphonic azide chloride, 2233
 Phenylphosphonic diazide, 2284
 Phenylphosphonyl dichloride, 2245
 3-Phenylpropanol, 3165
 3-Phenylpropenal, *see* Cinnamaldehyde, 3134
 3-Phenylpropionyl azide, 3144
 2-Phenyl-2-propyl hydroperoxide (Cumyl hydroperoxide), 3166
 Phenylsilver, 2217
 Phenylsodium, 2287
 5-Phenyltetrazole, 2727
 3-Phenyl-1-(5-tetrazolyl)-1-tetrazene, 2804
 Phenylthallium diazide, 2286
N-Phenyl-1,2,3,4-thiatriazolamine, 2729
 Phenylthiophosphonic diazide, 2285
 Phenylvanadium(V) dichloride oxide, 2246
 Phosgene, *see* Carbonyl dichloride, 0329
 Phosham, 4440
 † Phosphine, 4508
 Phosphinic acid, 4503
 Phosphonic acid, 4504
 Phosponium iodide, 4515
 Phosponium perchlorate, 4006
 Phosphorous acid, *see* Phosphonic acid, 4504
 † Phosphorus, 4874
 Phosphorus azide difluoride, 4315
 Phosphorus azide difluoride–borane, 4316
 Phosphorus chloride difluoride, 3979
 Phosphorus diiodide triselenide, 4629

Phosphorus(III) oxide, 4847
Phosphorus(III) oxide, *see* Tetraphosphorus hexaoxide, 4867
Phosphorus pentachloride, 4183
Phosphorus triazide, 4789
Phosphorus triazide oxide, 4788
Phosphorus tribromide, 0293
Phosphorus trichloride, 4153
Phosphorus tricyanide, 1343
Phosphorus trifluoride, 4339
Phosphorus triiodide, 4636
Phosphorus(V) oxide, 4864
Phosphorus(V) oxide, *see* Tetraphosphorus decaoxide, 4872
Phosphorus(V) sulfide, 4875
Phosphorus(V) sulfide, *see* Tetraphosphorus decasulfide, 4878
Phosphoryl chloride, 4149
Phosphoryl dichloride isocyanate, 0327
Phthalic acid, 2923
Phthalic anhydride, 2899
Phthaloyl diazide, 2898
Phthaloyl peroxide, 2900
2-Picoline, *see* 2-Methylpyridine, 2355
Picric acid, 2118
Picryl azide, 2092
1-Picryl-1,2,3-triazole, 2893
Pigment orange 5, *see* 1-(2,4-Dinitrobenzeneazo)-2-hydroxynaphthalene, 3702
2-Pinene, 3339
Piperazine, 1689
† Piperidine, 1993
2-Piperidone, 1934
† Pivalonitrile, 1929
Pivaloyl azide, 1935
† Pivaloyl chloride, 1925
Pivaloyloxydiethylborane, 3198
Platinum, 4887
Platinum diarsenide, 0107
Platinum hexafluoride, 4371
Platinum(IV) oxide, 4836
Platinum tetrafluoride, 4348
Plutonium, 4888
Plutonium bismuthide, 0231
Plutonium hexafluoride, 4372
Plutonium(III) hydride, 4509
Plutonium(IV) nitrate, 4768
Plutonium nitride, 4726
Poly(acetylene), 0687
Poly(acrylonitrile), 1108

Poly[bis(2,2,2-trifluoroethoxy)phosphazene] 1431
Poly([7,8-bis(trifluoromethyl)tetracyclo [4.2.0.0^{2,8}.0^{5,7}]octane-3,4-diyl)-
1,2-ethenediyl), 3463
Poly[borane(1)], 0134
cis-Poly(butadiene), 1484
Poly(1,3-butadiene peroxide), 1533
Poly(butadiyne), 1386
Poly(carbon monofluoride), 0337
Poly(chlorotrifluoroethylene), 0592
Poly(1,3-cyclohexadiene peroxide), 2386
Poly(cyclopentadienyltitanium dichloride), 1843
Poly(diazidophosphazene), 4787
Poly(diazomethylenemercury), 0522
Poly(dibromosilylene), 0283
Poly(difluorosilylene), 4330
Poly(dihydroxydioxodisilane), 4480
Poly(dimercurymmonium acetylide), 0669
Poly(dimercurymmonium azide), 4611
Poly(dimercurymmonium bromate), 0253
Poly(dimercurymmonium hydroxide), 4422
Poly(dimercurymmonium iodide hydrate), 4455
Poly(dimercurymmonium perchlorate), 4012
Poly(dimercurymmonium permanganate), 4608
Poly(dimethylketene peroxide), *see* Poly(peroxyisobutyrolactone), 1536
Poly(dimethylsiloxane), 0922
Poly(disilicon nitride), 4757
Poly(ethenyl nitrate), *see* Poly(vinyl nitrate), 0764
Poly(ethylene), 0782
Poly(ethylene terephthalate), 3262
Poly(ethylidene peroxide), 0835
Poly(furan-2,5-diyl), 1402
Poly(germanium dihydride), 4415
Poly(germanium monohydride), 4413
Poly(2,4-hexadiyne-1,6-ylyene carbonate), 2669
Poly(isobutene), 1583
Poly(methyl methacrylate peroxide), 1919
Poly(oxycarbonyl-2,4-hexadiyne-1,6-diyl), *see* Poly(2,4-hexadiyne-1,6-ylyene
carbonate), 2669
Poly[oxy(methyl)silylene], 0485
Poly(1-pentafluorothio-1,2-butadiyne), 1379
Poly(peroxyisobutyrolactone), 1536
Poly(pyrrole), 1417
Poly(selenium nitride), 4730
Poly(silylene), 4487
Poly(styrene peroxide), 2957
Poly(sulfur nitride), 4728

Poly(tetrafluoroethylene), 0629
 Poly(thiophene), 1407
 Poly(vinyl acetate peroxide), 1539
 Poly(vinyl alcohol), 0831
 Poly(vinyl butyral), 3030
 Poly(vinyl nitrate), 0764
 Potassium, 4645
 Potassium acetylene-1,2-dioxide, 0990
 Potassium acetylide, 0987
 Potassium amide, 4456
 Potassium amidosulfate, 4457
 Potassium antimonide, 4673
 Potassium azide, 4652
 Potassium azidodisulfate, 4655
 Potassium azidopentacyanocobaltate(3-), 1809
 Potassium azidosulfate, 4653
 Potassium azodisulfonate, 4663
 Potassium benzenehexoxide, 2626
 Potassium benzenesulfonylperoxosulfate, 2257
 Potassium *O-O*-benzoylmonoperoxosulfate, 2684
 Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato)
 cobaltate(1-), 3719
 Potassium bis(8-(4-nitrophenylthio)undecahydrodicarbaundecaborato)
 ferrate(1-), 3720
 Potassium bis(phenylethynyl)palladate(2-), 3699
 Potassium bis(phenylethynyl)platinate(2-), 3700
 Potassium bis(propynyl)palladate, 2308
 Potassium bis(propynyl)platinate, 2309
 Potassium bromate, 0255
 Potassium *tert*-butoxide, 1650
 Potassium carbonate, 0531
 Potassium chlorate, 4017
 Potassium chloride, 4015
 Potassium chlorite, 4016
 Potassium citrate tri(hydrogen peroxidate), 2258
 Potassium cyanate, 0528
 Potassium cyanide, 0526
 Potassium cyanide-potassium nitrite, 0527
 Potassium cyclohexanehexone 1,3,5-trioximate, 2625
 Potassium cyclopentadienide, 1846
 Potassium [(7,8,9,10,11- η)-1,2,3,4,5,6,7,8,9,11-decahydro-10-(4-nitrobenzenethiolato-*S*)-7,8-dicarbaundecaborato(2-)][(7,8,9,10,11- η)-undecahydro-7,8-dicarbadeborato(2-)]cobaltate(1-)], *see* Potassium [8-(4-nitrophenylthio)undecahydrodicarbaundecaborato]undecahydro-dicarbaundecaboratocobaltate(1-) 3380
 Potassium 1,3-dibromo-1,3,5-triazine-2,4-dione-6-oxide, 1032

Potassium dichromate, 4248
 Potassium diethylamide, 1685
 Potassium diethynylpalladate(2-), 1394
 Potassium diethynylplatinate(2-), 1395
 Potassium dihydrogenphosphide, 4460
 Potassium dinitramide, 4654
 Potassium dinitroacetamide, 0709
 Potassium 2,5-dinitrocyclopentanone, 1847
 Potassium dinitrogenbis(trimethylphosphine)cobaltate(1-), 3228
 Potassium dinitromethanide, 0377
 Potassium dinitrooxalatoplatinate(2-), 0988
 Potassium 1,1-dinitropropanide, 1175
 'Potassium dinitrosulfite', *see* Potassium *N*-nitrosohydroxylamine-
N-sulfonate, 4662
 Potassium 3,5-dinitro-2(1-tetrazenyl)phenoxide, 2256
 Potassium 2,6-dinitrotoluene, 2721
 Potassium dioxide, 4656
 Potassium diperoxomolybdate, 4659
 Potassium diperoxoorthovanadate, 4667
 Potassium *O,O*-diphenyl dithiophosphate, 3481
 Potassium dithioformate, 0378
 Potassium ethoxide, 0861
 † Potassium *O*-ethyl dithiocarbonate, 1176
 'Potassium ethyl xanthate', *see* Potassium *O*-ethyl dithiocarbonate
 Potassium ethynediolate, *see* Potassium acetylene-1,2-dioxide, 0990
 'Potassium ferricyanide', *see* Potassium hexacyanoferrate(III), 2063
 'Potassium ferrocyanide', *see* Potassium hexacyanoferrate(II), 2064
 Potassium fluoride hydrogen peroxidate, 4300
 Potassium graphite, 3107
 Potassium heptafluorotantalate(V), 4379
 Potassium hexaazidoplatinate(IV), 4665
 Potassium hexachloroplatinate, 4187
 Potassium hexacyanoferrate(II), 2064
 Potassium hexacyanoferrate(III), 2063
 Potassium hexaethynylcobaltate(4-), 3446
 Potassium hexaethynylmanganate(3-), 3448
 Potassium hexafluorobromate, 0244
 Potassium hexafluoromanganate(IV), 4363
 Potassium hexafluorosilicate(2-), 4364
 Potassium hexahydroaluminatate(3-), 0077
 Potassium hexanitrocobaltate(3-), 4213
 Potassium hexaooxoxenonate-xenon trioxide, 4674
 Potassium hydride, 4427
 Potassium hydrogen acetylenedicarboxylate, 1382
 Potassium hydrogen diazirine-3,3-dicarboxylate, 1068
 Potassium hydrogen peroxomonosulfate, 4430

Potassium hydrogen tartrate, 1462
 Potassium hydrogen xenate, 4429
 Potassium hydroxide, 4428
 Potassium 4-hydroxyamino-5,7-dinitro-4,5-dihydrobenzofurazanide
 3-oxide, 2176
 Potassium 4-hydroxy-5,7-dinitro-4,5-dihydrobenzofurazanide 3-oxide, 2114
 Potassium hydroxylamine-*O*-sulfonate, 4458
 Potassium hydroxyoxodiperoxochromate(1-), 4227
 Potassium hypoborate, 0163
 'Potassium hypophosphite', *see* Potassium phosphinate, 4459
 Potassium iodate, 4619
 Potassium iodide, 4618
 Potassium methanediazoate, 0449
 Potassium methoxide, 0450
 Potassium 4-methoxy-1-*aci*-nitro-3,5-dinitro-2,5-cyclohexadienonide, 2722
 Potassium methylamide, 0472
 Potassium methyldiazenoxide, *see* Potassium methanediazoate, 0449
 Potassium 3-methylfurazan-4-carboxylate 2-oxide, 1415
 Potassium methylselenide, 0451
 Potassium nitrate, 4650
 Potassium nitride, 4671
 Potassium nitridoosmate, 4651
 Potassium nitrite, 4649
 Potassium nitrobenzene, 2255
 Potassium 4-nitrobenzeneazosulfonate, 2175
 Potassium 6-*aci*-nitro-2,4-dinitro-2,4-cyclohexadieniminide, 2113
 Potassium 6-*aci*-nitro-2,4-dinitro-1-phenylimino-2,4-cyclohexadienide, 3451
 Potassium 1-nitroethane-1-oximate, 0756
 Potassium 1-nitroethoxide, 0806
 Potassium 4-nitrophenoxide, 2174
 Potassium [8-(4-nitrophenylthio)undecahydrodicarbaundecaborato]
 undecahydrodicarbaundecaboratocobaltate(1-) 3380
 Potassium nitrosodisulfate, 4661
 Potassium *N*-nitrosohydroxylamine-*N*-sulfonate, 4662
 Potassium octacyanodicobaltate(8-), 2880
 Potassium ozonide, *see* Potassium trioxide, 4657
 Potassium pentacarbonylvandate(3-), 2049
 Potassium pentacyanodiperoxochromate(5-), 1810
 Potassium pentaperoxodichromate, 4249
 Potassium perchlorate, 4018
 Potassium perchlorylamide, 4019
 Potassium periodate, 4620
 Potassium permanganate, 4647
 Potassium peroxide, 4666
 Potassium peroxodisulfate, 4668
 Potassium peroxyferrate(2-), 4396

Potassium phenyldinitromethanide, 2682
 Potassium 1-phenylethanediazoate, 2964
 Potassium phosphinate, 4459
 Potassium picrate, 2085
 Potassium *O*-propionohydroxamate, 1210
 Potassium silicide, 4658
 Potassium–sodium alloy, 4646
 Potassium sulfide, 4670
 Potassium sulfurdiimidate, 4664
 Potassium superoxide, *see* Potassium dioxide, 4656
 Potassium tetracyanomercurate(II), 1797
 Potassium tetracyanotitanate(IV), 1800
 Potassium tetraethylnickelate(2–), 2896
 Potassium tetraethylnickelate(4–), 2897
 Potassium tetrafluorochlorate(1–), 3984
 Potassium tetrahydrozincate, 4516
 Potassium tetrakis(propynyl)nickelate(4–), 3513
 Potassium 1,1,2,2-tetranitroethanediide, 0989
 Potassium tetraperoxochromate(3–), 4235
 Potassium tetraperoxomolybdate, 4660
 Potassium tetraperoxotungstate, 4669
 Potassium 1-tetrazolacetate, 1106
 Potassium thiazate, 4648
 Potassium thiocyanate, 0529
 Potassium triamidothallate ammoniate, 4551
 Potassium triazidocobaltate(1–), 4212
 Potassium tricarbonyltris(propynyl)molybdate(3–), 3471
 Potassium tricyanodiperoxochromate(3–), 1044
 Potassium 1,3,5-trinitrobenzene, 2112
 Potassium trinitromethanide, 0530
 Potassium 2,4,6-trinitrophenoxide, *see* Potassium picrate, 2085
 Potassium 2,4,6-trinitrotoluene, 2683
 Potassium trioxide, 4657
 Potassium 2,4,6-tris(dioxoselena)hexahydrotriazine-1,3,5-triide, 4672
 ‘Potassium triselenimidate’, *see* Potassium 2,4,6-tris(dioxoselena)-
 hexahydrotriazine-1,3,5-triide, 4672
 Praseodymium, 4886
 Prismane, 2294
 † Propadiene, 1124
 † Propadiene-1,3-dione, 1349
 Propadienedithione, 1350
 † Propanal, *see* Propionaldehyde, 1224
 † 2-Propanamine, *see* Isopropylamine, 1308
 † Propanamine, *see* Propylamine, 1309
 † Propane, 1271
 † 1,2-Propanediamine, *see* 1,2-Diaminopropane, 1325

- † 1,3-Propanediamine, *see* 1,3-Diaminopropane, 1326
- † Propanedinitrile, *see* Malononitrile, 1078
- Propanedioic acid, *see* Malonic acid, 1151
- 1,2-Propanediol, *see* Propylene glycol, 1285
- Propaneperoxoic acid, *see* Peroxypropionic acid, 1236
- † 2-Propanethiol, 1290
- † Propanethiol, 1289
- 1,2,3-Propanetriol, *see* Glycerol, 1286
- † 1,2,3-Propanetriyl nitrate, *see* Glyceryl trinitrate, 1196
- † 2-Propanol, 1280
- † Propanol, 1279
- Propan-1-ol-2-one, *see* Hydroxyacetone, 1231
- † 2-Propanone, *see* Acetone, 1220
- Propanonitrile, *see* Propiononitrile
- † Propionyl chloride, *see* Propionyl chloride, 1163
- † Propargyl alcohol, *see* 3-Propynol, 1147
- Propargyl azide, *see* 3-Azidopropyne, 1114
- † Propargyl bromide, *see* 3-Bromopropyne 1090
- † Propargyl chloride, *see* 3-Chloropropyne 1092
- † Propenal, *see* Acrylaldehyde, 1145
- Propenamide, *see* Acrylamide
- † Propene, 1198
- † Propenenitrile, 1107
- † Propenenitrile, *see* Acrylonitrile, 1107
- Propene ozonide, *see* 3-Methyl-1,2,4-trioxolane, 1235
- † 2-Propene-1-thiol, 1239
- E*-Propene-1,2,3-tricarboxylic acid, 2342
- † Propenoic acid, *see* Acrylic acid, 1148
- † 2-Propen-1-ol, 1223
- † 2-Propenoyl chloride, *see* Acryloyl chloride, 1093
- 2-Propenyl hydroperoxide, *see* Allyl hydroperoxide, 1226
- † 3-Propenyl methanoate, *see* Allyl formate, 1524
- Prop-2-enyl trifluoromethanesulfonate, 1461
- † Propionaldehyde, 1085
- Propiolic acid, 1086
- Propioloxy chloride, 1065
- † Propionaldehyde, 1224
- † Propiononitrile, 1179
- † Propionyl chloride, 1163
- Propionyl hypobromite, 1154
- Propionyl nitrite, 1185
- Propiophenone, 3151
- † Propyl acetate, 1977
- † Propylamine, 1309
- † Propyl bromide, *see* 1-Bromopropane, 1241

- † Propyl chloride, *see* 1-Chloropropane, 1243
- Propylcopper(I), 1251
- 3-Propyldiazirine, 1594
- Propylene glycol, 1285
- † Propyleneimine, *see* 2-Methylaziridine, 1257
- † Propylene oxide, 1225
- † Propylene, *see* Propene, 1198
- † Propyl formate, 1621
- Propyllithium, 1253
- † Propyl nitrate, 1267
- † Propyl nitrite, 1265
- 2-Propyl perchlorate, 1250
- Propyl perchlorate, 1249
- 2-Propylpyridine-4-carbothioamide, 3163
- Propylsilane, 1331
- Propylsodium, 1269
- 2-Propyl(thioisonicotinamide), *see* 2-Propylpyridine-4-carbothioamide, 3163
- † Propynal, *see* Propionaldehyde, 1085
- † Propyne, 1125
- 3-Propynethiol, 1152
- Propynoic acid, *see* Propiolic acid, 1086
- † 3-Propynol, 1147
- Propynoyl chloride, *see* Propioloyl chloride, 1065
- 1-Propynylcopper(I), 1098
- 5-(Prop-2-ynyloxy)-1,2,3,4-thiatriazole, 1420
- 2-Propynyl vinyl sulfide, 1883
- † Pyridine, 1848
- 3-Pyridinediazonium tetrafluoroborate, 1826
- 2,5-Pyridinedicarboxylic acid, 2696
- Pyridine *N*-oxide, 1849
- Pyridinium chlorochromate, 1862
- Pyridinium nitrate, 1873
- Pyridinium perchlorate, 1863
- † Pyrimidine-2,4,5,6-(1*H*,3*H*)-tetrone, 1398
- Pyrocatecholato(2-)(quinolin-8-olato-*N,O*)-trioxygenido(2-)-phosphorus, 3678
- Pyrocatechol, *see* 1,2-Benzenediol, 2331
- Pyrrole, 1466
- † Pyrrolidine, 1653
- 1-Pyrrolidinylcyclohexene, 3350
- Pyruvic acid, 1150

- Quinoline, 3121
- 6-Quinolinecarbonyl azide, 3243

Radon, 4893
Raney cobalt alloy, *see* Aluminium–cobalt alloy, 0049
Raney nickel alloys, *see* Aluminium–nickel alloys, 0055
Resorcinol, *see* 1,3-Benzenediol, 2332
Resorcinol diacetate, *see* 1,3-Diacetoxybenzene, 3291
Rhenium, 4890
Rhenium chloride trioxide, 4045
Rhenium hexafluoride, 4373
Rhenium hexamethoxide, 2603
Rhenium nitride tetrafluoride, 4344
Rhenium tetrachloride oxide, 4167
Rhenium(VII) sulfide, 4891
'Rhodanine', *see* 2-Thioxo-4-thiazolidinone, 1112
Rhodium, 4892
Rhodium(III) chloride, 4155
Rhodium tetrafluoride, 4349
Rubidium, 4889
Rubidium acetylide, 1025
Rubidium fluoroxysulfate, 4309
Rubidium graphite, 3110
Rubidium hexafluorobromate, 0245
Rubidium hydride, 4450
Rubidium hydrogen xenate, 4448
Rubidium nitride, 4727
Rubidium tetrafluorochlorate(1–), 3988
Ruthenium, 4894
Ruthenium(III) chloride, 4156
Ruthenium(IV) hydroxide, 4532
Ruthenium(IV) sulfide, 4895
Ruthenium salts, 4896
Ruthenium(VIII) oxide, 4862

Samarium, 4911
Samarium sulfide, 4899
Scandium 3-nitrobenzoate, 3816
Sebacoyl chloride, 3341
Seleninyl bis(dimethylamide), 1758
Seleninyl bromide, 0275
Seleninyl chloride, 4097
Selenium, 4908
Selenium difluoride dioxide, 4321
Selenium dioxide, 4838
Selenium tetrabromide, 0295
Selenium tetrafluoride, 4351
Selenium trioxide, 4852
† Silane, 4539

Silica, *see* Silicon dioxide, 4839
Silicon, 4909
Silicon dibromide sulfide, 0281
Silicon dioxide, 4839
Silicon monohydride (Silylidyne), 4451
Silicon monosulfide, 4898
Silicon oxide, 4828
Silicon tetraazide, 4791
Silicon tetrachloride, *see* Tetrachlorosilane, 4173
Silicon tetrafluoride, 4352
Silicon–zirconium alloys, 4910
Silver, 0001
Silver acetylide, 0568
Silver acetylide–silver nitrate, 0569
Silver–aluminium alloy, 0002
Silver amide, 0015
Silver 5-aminotetrazolide, 0392
Silver azide, 0023
Silver azide chloride, 0009
Silver 2-azido-4,6-dinitrophenoxide, 2075
Silver azidodithioformate, 0303
Silver 1-benzeneazothiocarbonyl-2-phenylhydrazide, 3605
Silver benzeneselenonate, 2219
Silver benzo-1,2,3-triazole-1-oxide, 2127
Silver bromate, 0007
Silver buten-3-ynide, 1408
Silver chlorate, 0011
Silver chloride, 0008
Silver chlorite, 0010
Silver chloroacetylide, 0566
Silver cyanate, 0300
Silver cyanide, 0299
Silver cyanodinitromethanide, 0567
Silver 3-cyano-1-phenyltriazene-3-ide, 2670
Silver cyclopropylacetylide, 1838
Silver difluoride, 0014
Silver dinitroacetamide, 0689
Silver 3,5-dinitroanthranilate, 2642
Silver 1,3-di(5-tetrazolyl)triazene, 0690
Silvered copper, 0003
Silver fluoride, 0013
Silver fulminate, 0301
Silver hexahydrohexaborate(2-), 0027
Silver hexanitrodiphenylamide, 3429
Silver 1,3,5-hexatriynide, 2052
Silver 3-hydroxypropynide, 1088

Silver hyponitrite, 0031
Silver(II) oxide, 0025
Silver imide, 0030
Silver iodate, 0020
Silver(I) oxide, 0032
Silver isophthalate, 2887
Silver malonate, 1071
Silver 3-methylisoxazolin-4,5-dione-4-oximate, 1409
Silver nitrate, 0022
Silver nitride, 0038
Silver nitroguanidide, 0422
Silver 4-nitrophenoxide, 2126
Silver *N*-nitrosulfuric diamidate, 0016
Silver nitroureide, 0391
Silver osmate, 0034
Silver oxalate, 0572
Silver perchlorate, 0012
Silver perchlorylamide, 0028
Silver *N*-perchlorylbenzylamide, 2734
Silver permanganate, 0021
Silver peroxide, 0033
Silver peroxochromate, 0037
Silver phenoxide, 2218
Silver phosphinate, 0017
Silver sulfide, 0026
Silver sulfide, 0035
Silver tetrafluoroborate, 0005
Silver tetrafluorobromate, 0006
Silver tetrazolide, 0366
Silver–thorium alloy, 0004
Silver 1,2,3-triazolide, 0688
Silver trichloromethanephosphonate, 0304
Silver trifluoropropynide, 1030
Silver trinitromethanide, 0302
Silver 2,4,6-tris(dioxoselena)perhydrotriazine-1,3,5-triide, 0039
'Silver triselenimidate', *see* Silver 2,4,6-tris(dioxoselena)perhydrotriazine-1,3,5-triide, 0039
Silver trisulfurpentanitridate, 0024
Silyl trifluoromethanesulfonate, 0444
3-Sodio-5-(5'-nitro-2'-furfurylideneamino)imidazolidin-2,4-dione, 2908
Sodium, 4796
Sodium abietate, 3805
Sodium acetate, 0779
Sodium acetylde, 1022
Sodium 2-allyloxy-6-nitrophenylpyruvate oxime, 3505
Sodium aluminium hydride, *see* Sodium tetrahydroaluminate, 0076

Sodium amide, 4465
Sodium amidosulfate, 4467
Sodium–antimony alloy, 4797
Sodium arachidonate, *see* Sodium 5,8,11,14-eicosatetraenoate, 3809
Sodium azide, 4758
Sodium azidosulfate, 4759
Sodium 5-azidotetrazolide, 0551
Sodium benzenehexoxide, 2634
Sodium 2-benzothiazolylthiolate, 2658
Sodium 1,4-bis(*aci*-nitro)-2,5-cyclohexadienide, 2188
Sodium 3,5-bis(*aci*-nitro)cyclohexene-4,6-diiminide, 2203
Sodium borate hydrogen peroxidate, 0154
Sodium borohydride, *see* Sodium tetrahydroborate, 0147
Sodium bromate, 0257
Sodium bromoacetylde, 0581
Sodium carbonate, 0552
Sodium carbonate hydrogen peroxidate, 0553
Sodium chlorate, 4039
Sodium chloride, 4036
Sodium chlorite, 4038
Sodium chloroacetate, 0694
Sodium 4-chloroacetophenone oximate, 2927
Sodium chloroacetylde, 0601
Sodium *N*-chlorobenzenesulfonamide, 2229
Sodium chloroethynide, *see* Sodium chloroacetylde, 0601
Sodium 4-chloro-2-methylphenoxide, 2720
Sodium *N*-chloro-4-toluenesulfonamide, 2739
Sodium 4-cresolate, *see* Sodium 4-methylphenoxide, 2788
Sodium cyanide, 0536
Sodium diazomethanide, 0383
Sodium 1,3-dichloro-1,3,5-triazine-2,4-dione-6-oxide, 1037
Sodium dichromate, 4250
Sodium diethylamide, 1688
Sodium diformylnitromethanide hydrate, 1076
Sodium dihydrobis(2-methoxyethoxy)aluminate, 2575
Sodium dihydrogen phosphide, 4474
Sodium 1,3-dihydroxy-1,3-bis(*aci*-nitromethyl)-2,2,4,4-tetramethylcyclo-
butandiide, 3344
Sodium 4,4-dimethoxy-1-*aci*-nitro-3,5-dinitro-2,5-cyclohexadienide, 2951
Sodium dimethylsulfinat, 0879
Sodium dinitroacetamide, 0712
Sodium dinitromethanide, 0384
Sodium 5-(dinitromethyl)tetrazolide, 0681
Sodium 2,4-dinitrophenoxide, 2115
Sodium dioxide, 4799
Sodium diphenylketyl, 3604

Sodium disulfide, 4812
Sodium disulfite, 4808
Sodium dithionite, 4807
Sodium 5,8,11,14-eicosatetraenoate, 3809
Sodium ethaneperoxoate, *see* Sodium peroxyacetate, 0780
Sodium ethoxide, 0878
Sodium ethoxyacetylde, 1478
Sodium ethylenediaminetetraacetate, 3305
Sodium ethynediolate, 1023
Sodium ethynide, *see* Sodium acetylde, 1022
Sodium fulminate, 0537
Sodium germanide, 4418
Sodium hexaazidophosphate, 4794
Sodium hexahydroxyplatinate(IV), 4564
Sodium hexakis(propynyl)ferrate(4-), 3759
Sodium hexaoxodinitrate, 4740
Sodium hydrazide, 4500
Sodium hydride, 4444
Sodium hydrogen carbonate, 0390
Sodium hydrogen sulfate, 4446
Sodium hydrogen xenate, 4447
'Sodium hydrosulfite', *see* Sodium dithionite, 4807
Sodium hydroxide, 4445
Sodium 2-hydroxyethoxide, 0880
Sodium *O*-hydroxylamide, 4466
Sodium 3-hydroxymercurio-2,6-dinitro-4-*aci*-nitro-2,5-cyclohexa-
dienonide, 2083
Sodium 2-hydroxymercurio-4-*aci*-nitro-2,5-cyclohexadienonide, 2171
Sodium 2-hydroxymercurio-6-nitro-4-*aci*-nitro-2,5-cyclohexadienonide, 2111
Sodium 5(5'-hydroxytetrazol-3'-ylazo)tetrazolide, 0682
Sodium hypoborate, 0164
Sodium hypochlorite, 4037
Sodium hyponitrite, 4736
'Sodium hypophosphite', *see* Sodium phosphinate, 4473
Sodium iodate, 4624
Sodium iodide, 4623
Sodium isopropoxide, 1270
'Sodium metabisulfite', *see* Sodium disulfite, 4808
Sodium metasilicate, 4805
Sodium methoxide, 0464
Sodium methoxyacetylde, 1123
Sodium 3-methylisoxazolin-4,5-dione-4-oximate, 1419
Sodium 4-methylphenoxide, 2788
Sodium molybdate, 4713
Sodium monoperoxycarbonate, 0554
Sodium nitrate, 4721

Sodium nitride, 4723
Sodium nitrite, 4720
Sodium 3-nitrobenzenesulfonate, 2184
Sodium *aci*-nitromethanide, 0403
Sodium 6-*aci*-nitro-4-nitro-2,4-cyclohexadieniminide, 2200
Sodium 4-nitrophenoxide, 2183
Sodium 4-nitrosophenoxide, 2181
Sodium 5-nitrotetrazolide, 0547
Sodium 2-nitrothiophenoxide, 2182
Sodium nitroxylate, 4722
Sodium octahydrotriborate, 0177
Sodium orthophosphate, 4814
Sodium oxide, 4802
Sodium pentacarbonylrhenate, 2051
Sodium pentacyanonitrosylferrate(2-), 1813
Sodium pentaoxidinitrate, 4739
'Sodium percarbonate', *see* Sodium carbonate hydrogen peroxidate, 0553
Sodium perchlorate, 4040
Sodium permanganate, 4703
Sodium peroxide, 4803
Sodium peroxoborate, 0155
Sodium peroxodisulfate, 4809
Sodium peroxyacetate, 0780
Sodium peroxydicarbonate, 1024
'Sodium perpyrophosphate' 4529
Sodium phenylacetylde, 2911
Sodium 3-phenylisoxazolin-4,5-dione-4-oximate, 3114
Sodium phosphide, 4815
Sodium phosphinate, 4473
Sodium picrate, 2086
Sodium polysulfide, 4813
Sodium pyrophosphate hydrogen peroxidate, 4816
Sodium silicide, 4801
Sodium sulfate, 4806
Sodium sulfide, 4811
Sodium superoxide, *see* Sodium dioxide, 4799
Sodium tetraborate, 0185
Sodium tetracarbonylferrate(2-), 1376
Sodium tetracyanatopalladate(II), 1802
Sodium tetradecahydranonaborate, 0196
Sodium tetrahydroaluminate, 0076
Sodium tetrahydroborate, 0147
Sodium tetrahydrogallate, 4408
Sodium 1,1,2,2-tetranitroethanediide, 1007
Sodium tetraoxidinitrate, 4738
Sodium tetraperochromate(3-), 4240

Sodium tetraperoxomolybdate, 4715
 Sodium tetraperoxotungstate, 4810
 Sodium tetrasulfur pentanitride, 4775
 Sodium 1-tetrazolacetate, 1119
 Sodium thiocyanate, 0538
 Sodium thiosulfate, 4804
 Sodium triamine, 4581
 Sodium triazidoaurate(?), 0113
 Sodium tricarbonylnitrosylferrate, 1064
 Sodium trichloroacetate, 0608
 Sodium 2,2,2-trifluoroethoxide, 0705
 Sodium 2,2,2-trinitroethanide, 0713
 Sodium 2,4,6-trinitrophenoxide, *see* Sodium picrate, 2086
 Sodium trioxide, 4800
 Sodium trioxodinitrate, 4737
 Sodium tris(*O,O'*-1-oximatonaphthalene-1,2-dione)ferrate (Pigment green B), 3871
 Sodium trisulfurtrinitride, 4760
 Sodium–zinc alloy, 4798
 ‘Solid Phosphorus hydride’ 4449
 Sorbic acid, *see* 2,4-Hexadienoic acid, 2385b
 Spiro[3*H*-diazirine-3,9′pentacyclo[4,3,0,0,2[·]5,0,3[·]8,0,4[·]7]nonane], *see* Spiro-(homocubane-9,9′-diazirine), 3127
 Spiro(homocubane-9,9′-diazirine), 3127
 † Stibine, 4510
 Stibobenzene, *see* Diphenyldistibene, 3500
 Stilbene, *see* *trans*-1,2-Diphenylethylene, 3643
 Strontium, 4913
 Strontium acetylide, 1026
 Strontium azide, 4785
 Strontium peroxide, 4841
 Strontium sulfide, 4901
 Styrene, 2945
 Succinic anhydride, 1443
 Succinodinitrile, 1433
 Succinoyl diazide, 1438
 Succinyl peroxide, *see* 3,6-Dioxo-1,2-dioxane, 1445
 Sucrose, 3558
 Sulfamic acid, *see* Amidosulfuric acid, 4499
 Sulfamoyl azide, 4472
 Sulfinyl azide, 4778
 † Sulfinylbismethane, *see* Dimethyl sulfoxide, 0921
 Sulfinyl bromide, 0274
 Sulfinyl chloride, 4096
 Sulfinylcyanamide, 0542
 Sulfinyl fluoride, 4318

Sulfinylsulfamoyl fluoride, *see* Sulfur oxide-(*N*-fluorosulfonyl)imide, 4305
 Sulfolane, *see* Tetrahydrothiophene-1,1-dioxide, 1622
 Sulfonyl chloride, 4099
 † Sulfur, 4897
 Sulfur dibromide, 0280
 Sulfur dichloride, 4113
 Sulfur dioxide, 4837
 Sulfur hexafluoride, 4374
 Sulfuric acid, 4479
 Sulfur oxide-(*N*-fluorosulfonyl)imide, 4305
 Sulfur tetrafluoride, 4350
 Sulfur thiocyanate, 1002
 Sulfur trioxide, 4849
 Sulfur trioxide–dimethylformamide, 4850
 Sulfuryl azide chloride, 4031
 Sulfuryl chloride, *see* Sulfonyl chloride, 4099
 Sulfuryl diazide, 4779

 Tantalum, 4914
 Tantalum(IV) sulfide, 4903
 Tantalum pentachloride, 4185
 Tantalum(V) oxide, 4865
 Tartaric acid, 1545
 Technetium, 4915
 Tellurane-1,1-dioxide, 1980
 Tellurium, 4916
 Tellurium tetrabromide, 0296
 Tellurium tetrachloride, 4175
 Tellurium trioxide, 4853
 Terephthalic acid, 2924
 Terephthaloyl chloride, 2889
 Tetraacrylonitrilecopper(I) perchlorate, 3510
 Tetraallyl-2-tetrazene, 3545
 Tetraallyluranium, 3548
 Tetra(3-aminopropanethiolato)trimercury perchlorate, 3582
 Tetraamminebis(dinitrogen)osmium(II) perchlorate, 4071
 Tetraamminebis(5-nitro-2*H*-tetrazolato)cobalt(1⁺) perchlorate, 0970
 Tetraammine-2,3-butanediimineruthenium(III) perchlorate, 1794
 Tetraamminecadmium permanganate, 3956
 Tetraamminecopper(II) azide, 4277
 Tetraamminecopper(II) bromate, 0264
 Tetraamminecopper(II) nitrate, 4276
 Tetraamminecopper(II) nitrite, 4275
 Tetraamminecopper(II) sulfate, 4274
trans Tetraamminediazidocobalt(III) *trans*-diamminetetraazido-
 cobaltate(1⁻), 4220

Tetraamminedinitratoplatinum, *see* Tetraammineplatinum(II) nitrate, 4589
 Tetraamminedinitratoplatinum, *see* Tetraammineplatinum(II) nitrate, 4590
 Tetraamminedithiocyanatocobalt(III) perchlorate, 0969
 Tetraamminehydroxynitratoplatinum(IV) nitrate, 4594
 Tetraamminelithium dihydrogenphosphide, 4595
 Tetraamminenickel(II) nitrate, 4587
 Tetraamminepalladium(II) nitrate, 4588
 Tetraammineplatinum(II) nitrate, 4589
 Tetraammineplatinum(II) nitrate, 4590
 Tetraamminezinc peroxodisulfate, 4586
 Tetraarsenic tetrasulfide, 0109
 1,4,8,11-Tetraazacyclotetradecanenickel(II) perchlorate, 3376
 4a,8a,9a,10a-Tetraaza-2,3,6,7-tetraoxaperhydroanthracene, 2478
 1,3,5,7-Tetraazatricyclo[3.3.1.1^{3,7}]decane, *see* Hexamethylenetetramine, 2477
 1,3,6,8-Tetraazatricyclo[6.2.1.1^{3,6}]dodecane tetranitrate, 3092
 Tetraazidoarsonium hexafluoroarsenate, 0104
 Tetraazido-1,4-benzoquinone, 2633
 1,2:5,6:11,12:15,16-Tetrabenzocycloconta-1,5,11,15-tetraene-3,7,9,13,17,19-hexayne, 3888
 1,2:5,6:11,12:15,16-Tetrabenzocycloconta-1,5,11,15-tetraene-3,7,9,13,17,19-hexahydro[20]annulene, *see* 1,2:5,6:11,12:15,16-Tetrabenzocycloconta-1,5,11,15-tetraene-3,7,9,13,17,19-hexayne, 3888
 Tetraborane(10), 0181
 Tetra(boron nitride) fluorosulfate, 0180
 Tetraboron tetrachloride, 0179
 2',4',5',7'-Tetrabromo-3',6'-dihydroxySpiro[isobenzofuran-1(3*H*),9'-[9*H*]xanthen-3-one], 3787
 Tetrabromomethane, *see* Carbon tetrabromide, 0315
 Tetrabutylammonium fluoride, 3728
 Tetrabutylammonium hydrogen monoperoxysulfate, 3732
 Tetrabutylammonium permanganate, 3729
 Tetracarbon monofluoride, 1362
 Tetracarbonylmolybdenum dichloride, 1358
 † Tetracarbonylnickel, 1805
N,N,N',N'-Tetrachloroadipamide, 2370
 1,2,4,5-Tetrachlorobenzene, 2079
 1,1,2,3-Tetrachloro-1,3-butadiene, 1389
 1,1,4,4-Tetrachlorobutatriene, 1360
 Tetrachlorodiazocyclopentadiene, 1807
 3,3,4,5-Tetrachloro-3,6-dihydro-1,2-dioxin, 1390
 1,2,3,4-Tetrachloro-7,7-dimethoxy-5-phenylbicyclo[2.2.1]hepta-2,5-diene, 3682
 Tetrachlorodiphosphane, 4171
 1,1,2,2-Tetrachloroethane, 0699
 Tetrachloroethylene, 0609
 Tetrachloroethylene carbonate, 1042

Tetrachloromethane, *see* Carbon tetrachloride, 0332
 Tetrachloropropadiene, 1041
 Tetrachlorosilane, 4173
 Tetrachlorotrifluoromethylphosphorane, 0333
 Tricyanoethylene, 2629
 Tetracyanooctaethyltetragold, 3814
 Tetracyclo[4.3.0^{3,5}.0^{4,6}]nona-1,7-diene, 3125
 Tetracyclo[20.2.0.0^{6,9}.0^{14,17}]tetracos-1(22),6(9),14(17)-triene-
 2,4,10,12,18,20-hexayne-7,8,15,16,23,24-hexone, 3858
 † Tetradecane, 3671
 2,4,6,8-Tetraethenyl-2,4,6,8-tetramethylcyclotetrasiloxane. 3560
 Tetraethylammonium perchlorate, 3088
 Tetraethylammonium periodate, 3090
 Tetraethyldiarsane, 3087
 'Tetraethyldiborane' 3098
 'Tetraethylenepentamine', *see* 1,11-Diamino-3,6,9-triazaundecane, 3101
 Tetraethyllead, 3095
 Tetraethyltin, 3097
 Tetraethynylgermanium, 2895
 Tetraethynyltin, 2901
 Tetrafluoroammonium hexafluoromanganate, 4384
 Tetrafluoroammonium hexafluoronickelate, 4385
 Tetrafluoroammonium octafluoroxenate, 4386
 Tetrafluoroammonium perbromate, 0242
 Tetrafluoroammonium perchlorate, 3986
 Tetrafluoroammonium tetrafluoroborate, 0133
 † 1,2,4,5-Tetrafluorobenzene, 2081
 Tetrafluorobenzene-1,4-diol, 2082
 Tetrafluoroboric acid, 0130
 Tetrafluorobutanedioic acid, *see* Perfluorosuccinic acid, 1391
 1,1,4,4-Tetrafluorobutatriene, 1363
 Tetrafluorodiaziridine, 0351
 1,1,2,2-Tetrafluoroethane, 0706
 Tetrafluoroethane sultone, *see* Tetrafluorooxathietane-2,2-dioxide, 0635
 † Tetrafluoroethylene, 0628
 Tetrafluoroethylene oxide, *see* Tetrafluorooxirane, 0632
 1,1,2,2-Tetrafluoroethyl hypofluorite, *see* 1,1,2,2-Tetrafluoro-
 1-(fluoroxy)ethane, 0668
 1,1,2,2-Tetrafluoro-1-(fluoroxy)ethane, 0668
 † Tetrafluorohydrazine, 4345
 Tetrafluoroiodosyl hypofluorite, 4356
 Tetrafluoromethane, *see* Carbon tetrafluoride, 0349
 Tetrafluorooxathietane-2,2-dioxide, 0635
 Tetrafluorooxirane, 0632
 2,2,3,3-Tetrafluoropropanol, 1133
 Tetrafluorosuccinamide, 1430

- Tetrafluorourea, 0352
- 1.3a,4,6a-Tetrahydro-*N,N'*-dinitroimidazo[4,5-*d*]imidazole-2,5-diamine, *see*
 Octahydro-2,5-bis(nitroimino)imidazo[4,5-*d*]imidazole, 1514
- 1,4,5,6-Tetrahydro-*N*,5-dinitrotriazin-2-amine, *see* 5-Nitro-2-
 (nitroimino)hexahydro-1,3,5-triazine, 1218
- † Tetrahydrofuran, 1612
- 2-Tetrahydrofuranylidene(dimethylphenylphosphine-
 trimethylphosphine)-2,4,6-trimethylphenylnickel perchlorate, 3854
- Tetrahydrofurfuryl alcohol, 1978
- 2-Tetrahydrofuryl hydroperoxide, 1624
- 4a,5,7a,8-Tetrahydro-4,8-methano-4*H*-indeno[5,6-*c*]-1,2,5-oxadiazole 1-oxide,
 3281
- 4a,5,7a,8-Tetrahydro-4,8-methano-4*H*-indeno[5,6-*c*]-1,2,5-oxadiazole 3-oxide,
 3282
- Tetrahydronaphthalene, 3300
- 1,2,3,4-Tetrahydro-1-naphthyl hydroperoxide, 3312
- Tetrahydro-1,4-oxazine, *see* Morpholine, 1657
- † Tetrahydropyran, 1965
- 2-Tetrahydropyranylidene-bis(dimethylphenylphosphine)-
 3,4,6-trimethylphenylnickel perchlorate, 3877
- † 1,2,3,6-Tetrahydropyridine, 1930
- 1,4,5,8-Tetrahydro-1,4,5,8-tetrathiafulvalene, *see* 2,2'-Bi-1,3-dithiole, 2215
- † Tetrahydrothiophene, 1627
- Tetrahydrothiophene-1,1-dioxide, 1622
- Tetrahydroxotritin(2+) nitrate, 4525
- Tetrahydroxydioxotrisilane, 4536
- Tetraimide (Tetraazetidine), 4527
- Tetraiodoaronium tetrachloroaluminate, 0057
- Tetraiododiphosphane, 4637
- Tetraiodoethylene, 0986
- Tetraiodomethane, *see* Carbon tetraiodide, 0525
- Tetraisocyanatogermane, *see* Germanium isocyanate, 1377
- Tetraisopropylchromium, 3577
- 1,1,6,6-Tetrakis(acetylperoxy)cyclododecane, 3810
- Tetrakis(μ_3 -2-amino-2-methylpropanolato)tetrakis(μ_2 -2-amino-
 2-methylpropanolato)hexacopper(II) perchlorate, 3883
- 3,3,6,6-Tetrakis(bromomethyl)-9,9-dimethyl-1,2,4,5,7,8-hexoxonane, 3173
- Tetrakis(butylthio)uranium, 3731
- Tetrakis(chloroethynyl)silane, 2879
- Tetrakis(*N,N*-dichloroaminomethyl)methane, 1900
- Tetrakis(diethylphosphino)silane, 3733
- Tetrakis(dimethylamino)titanium, 3104
- Tetrakis(ethylthio)uranium, 3096
- 1,1,4,4-Tetrakis(fluoroxy)hexafluorobutane, 1374
- Tetrakis(hydroxymercurio)methane, 0471
- Tetrakis(hydroxymethyl)phosphonium nitrate, 1754

Tetrakis(3-methylpyrazole)cadmium sulfate, 3716
 Tetrakis(3-methylpyrazole)manganese(II) sulfate, 3717
 Tetrakis(4-*N*-methylpyridinio)porphinecobalt(III)(5+) perchlorate, 3908
 Tetrakis(4-*N*-methylpyridinio)porphineiron(III)(5+) perchlorate, 3909
 1,3,4,6-Tetrakis(2-methyltetrazol-5-yl)-hexaaza-1,5-diene, 3013
 5,10,15,20-Tetrakis(2-nitrophenyl)porphine, 3907
 Tetrakis(pentafluorophenyl)titanium, 3842
 Tetrakis(pyrazole)manganese(II) sulfate, 3538
 Tetrakis(pyridine)bis(tetracarbonylcobalt)magnesium, 3867
 Tetrakis(thiourea)manganese(II) perchlorate, 1785
 Tetrakis(trimethylsilyl)diaminodiphosphene, 3588
 Tetrakis(2,2,2-trinitroethyl)orthocarbonate, 3132
 Tetramethoxyethylene, 2516
 † Tetramethoxymethane, *see* Tetramethyl orthocarbonate, 2022
 † Tetramethoxysilane, 1764
 Tetramethylammonium amide, 1782
 Tetramethylammonium azidocyanatoiodate(I), 2006
 Tetramethylammonium azidocyanatoiodate(I), 2005
 Tetramethylammonium azidoselenocyanatoiodate(I), 2007
 Tetramethylammonium chlorite, 1742
 Tetramethylammonium diazidoiodate(I), 1750
 Tetramethylammonium monoperchromate, 1745
 Tetramethylammonium ozonate, 1753
 Tetramethylammonium pentafluoroxenoxide, 1747
 Tetramethylammonium pentaperoxodichromate, 3103
 Tetramethylammonium periodate, 1749
 Tetramethylammonium superoxide, 1751
 Tetramethylbis(trimethylsilanoxy)digold, 3382
 2,4,6,8-Tetramethylcyclotetrasiloxane, 1790
 Tetramethyldialuminium dihydride, 1778
 Tetramethyldiarsane, 1736
 Tetramethyldiarsine, *see* Tetramethyldiarsane, 1736
 Tetramethyldiborane, 1779
 Tetramethyldigallane, 1748
 Tetramethyldigold diazide, 1739
 Tetramethyl-1,2-dioxetane, 2508
 Tetramethyldiphosphane, 1765
 Tetramethyldiphosphane disulfide, 1766
 Tetramethyldiphosphine, *see* Tetramethyldiphosphane, 1765
 Tetramethyldisiloxane, 1783
 Tetramethyldistibane, 1769
 Tetramethyldistibine, *see* Tetramethyldistibane, 1769
N,N,N',N'-Tetramethylethane-1,2-diamine, trimethylpalladium(IV) bromide, 3225
N,N,N,N-Tetramethylethane-1,2-diamine, trimethylpalladium(IV) iodide, 3226
N,N,N',N'-Tetramethylformamidinium perchlorate, 2026

1,3,4,7-Tetramethylisindole, 3528
 Tetramethyllead, 1767
 2,12,18,28-Tetramethyl-3,7,11,19,23,27,33,34-octaazatricyclo
 [27.3.1.1(13,17)]tetratriaconta-1(33),2,11,13,15,17(34)
 18,27,29,31-decaenetetraaquadiron(II) tetraperchlorate, 3878
 † Tetramethyl orthocarbonate, 2022
 † 2,2,3,3-Tetramethylpentane, 3211
 † 2,2,3,4-Tetramethylpentane, 3212
 Tetramethylplatinum, 1768
 Tetramethylplumbane, *see* Tetramethyllead, 1767
 2,3,4,6-Tetramethylpyrilium perchlorate, 3167
 † Tetramethylsilane, 1770
 † Tetramethyl silicate, *see* Tetramethoxysilane, 1764
 1,1,3,3-Tetramethylsiloxalane, 1774
 † Tetramethylstannane, *see* Tetramethyltin, 1771
 Tetramethylsuccinodinitrile, 3007
 Tetramethyltellurium(IV), 1772
 3,3,6,6-Tetramethyl-1,2,4,5-tetraoxane, 2517
 Tetramethyl-2-tetrazene, 1759
 7,7,10,10-Tetramethyl-1,2,5,6-tetroxecane-3,4-dione, 3349
 † Tetramethyltin, 1771
 1,3,5,7-Tetranitroadamantane, 3310
N,2,3,5-Tetranitroaniline, 2121
N,2,4,6-Tetranitroaniline, 2123
 2,3,4,6-Tetranitroaniline, 2122
 1,3,6,8-Tetranitrocarbazole, 3442
 Tetranitroethylene, 1010
 Tetranitromethane, 0546
N,2,4,6-Tetranitro-*N*-methylaniline, 2706
 1,3,6,8-Tetranitronaphthalene, 3237
 1,3,5,7-Tetranitroperhydro-1,3,5,7-tetrazocine, 1605
 2,3,4,6-Tetranitrophenol, 2091
trans-1,4,5,8-Tetranitro-1,4,5,8-tetraazadecahydronaphthalene, 2427
 2,5,7,8-Tetraoxa[4.2.0]bicyclooctane, *see cis*-1,4-Dioxenedioxetane, 1538
 4,5,11,12-Tetraoxo-14*H*-[1,2,5]oxadiazolo[3,4-*e*][1,2,5]oxadiazolo[3',4':4,5]
 benzotriazolo[2,1-*a*]benzotriazolo-*b*-ium 1,8-dioxide, 3595
 Tetraperchloratosilicon, 4169
 Tetraphenylarsonium permanganate, 3848
 3,3,6,6-Tetraphenylhexahydro-3,6-disilatetrazine, 3852
 Tetraphenyllead, 3850
 1,3,6,8-Tetraphenyloctaazatriene, 3849
 Tetraphosphoric acid, 4567
 Tetraphosphorus decaoxide, 4872
 Tetraphosphorus decasulfide, 4878
 Tetraphosphorus decasulfide . pyridine complex, 4879
 'Tetraphosphorus hexanitride' 4781

Tetraphosphorus hexaoxide, 4867
 Tetraphosphorus hexaoxide–bis(borane), 4868
 Tetraphosphorus hexaoxide tetrasulfide, 4869
 Tetraphosphorus tetraoxide trisulfide, 4859
 Tetraphosphorus triselenide, 4880
 Tetraphosphorus trisulfide, 4877
 Tetra-2-propenyl-2-tetrazene, *see* Tetraallyl-2-tetrazene, 3545
 ‘Tetrapropyldiborane’ 3580
 Tetrapyridinecobalt(II) chloride, 3801
 Tetraselenium dinitride, 4756
 Tetraselenium tetranitride, 4771
 Tetrasilane, 4584
 Tetrasilver diimidodioxosulfate, 0043
 Tetrasilver diimidotriphosphate, 0042
 Tetrasilver orthodiamidophosphate, 0041
 Tetrasilylhydrazine, 4585
 Tetraspiro[2.2.2.2.2.2.4]docosa-4,9,14,19,21pentyne, 3832
 Tetrasulfur dinitride, 4755
 Tetrasulfurtetraimide–silver perchlorate, 0029
 Tetrasulfur tetranitride, 4770
 Tetratellurium tetranitride, 4773
 Travinyllead, 3016
 1,2,4,5-Tetrazine, 0715
 Tetrazirconium tetraoxide hydrogen nonaperchlorate, 4195
 Tetrazole, 0411
 Tetrazole-5-diazonium chloride, 0370
 5,5'-Tetrazolyl disulfide, 0721
 Tetryl, *see* *N*,2,4,6-Tetranitro-*N*-methylaniline, 2706
 Thallium, 4922
 Thallium bromate, 0260
 Thallium fulminate, 0539
 Thallium hydroxyoxodiperoxochromate(1–), 4228
 Thallium(I) azide, 4765
 Thallium(I) azidodithiocarbonate, 0543
 Thallium(I) chlorite, 4043
 Thallium(I) dithiocarbonazidate, *see* Thallium(I) azidodithiocarbonate, 0543
 Thallium(III) nitrate, 4762
 Thallium(III) oxide, 4854
 Thallium(I) iodacetylde, 0984
 † Thallium(I) methanediazoate (Thallium *N*-nitrosomethylamide), 0458
 Thallium(I) nitride, 4731
 Thallium(I) 2-nitrophenoxide, 2186
 Thallium(I) 4-nitrophenoxide, 2187
 Thallium(I) peroxodiborate, 0171
 Thallium(I) tetraazidothallate, 4792
 Thallium *aci*-phenylnitromethanide, 2723

Thallium 2,4,6-tris (dioxoselena) hexahydrotriazine-1,3-5-triide, 4761
'Thallium triselenimidate', *see* Thallium 2,4,6-
tris(dioxoselena)hexahydrotriazine-1,3-5-triide, 4761
Thianthrenium perchlorate, 3455
† 3-Thiapentane, *see* Diethyl sulfide, 1714
Thiazyl chloride, *see* 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazine, 4145
Thiazyl fluoride, 4306
3-Thietanol-1,1-dioxide, 1238
† Thiirane, 0840
1,1'-Thiobis(2-chloroethane), *see* Bis(2-chloroethyl)sulfide, 1589
† Thiobismethane, *see* Dimethyl sulfide, 0932
Thiocarbonyl azide thiocyanate, 1011
3-Thiocresol, 2809
Thiocyanogen, 1001
† Thiolacetic acid, 0832
Thionyl bromide, *see* Sulfinyl bromide, 0274
Thionyl chloride, *see* Sulfinyl chloride, 4096
† Thiophene, 1448
Thiophenol, *see* Benzenethiol, 2344
Thiophosphoryl chloride, 4154
Thiophosphoryl chloride difluoride, 3980
Thiophosphoryl fluoride, 4340
Thiotriazyl chloride, 4033
Thiotriazyl nitrate, 4767
Thiotriazyl perchlorate, 4032
Thiourea, 0480
2-Thioxo-4-thiazolidinone, 1112
Thorium, 4917
Thorium dicarbide, 1027
Thorium dihydride, 4489
Thorium hydride, 4540
Thorium oxide sulfide, 4826
Thorium salts, 4918
Tin, 4912
Tin azide trichloride, 4146
Tin(II) chloride, 4116
Tin(II) fluoride, 4331
Tin(II) nitrate, 4750
Tin(II) oxide, 4829
Tin(II) perchlorate, 4109
Tin(II) sulfide, 4900
Tin(IV) chloride, 4174
Tin(IV) oxide, 4840
Tin(IV) sulfide, 4902
Titanium, 4919
Titanium azide trichloride, 4147

Titanium butoxide, 3730
 Titanium carbide, 0561
 Titanium diazide dibromide, 0273
 Titanium dibromide, 0284
 Titanium dihydride, 4490
 Titanium diiodide, 4630
 Titanium dioxide, *see* Titanium(IV) oxide, 4842
 Titanium(II) chloride, 4117
 Titanium(III) methoxide, 1317
 Titanium(IV) oxide, 4842
 Titanium(IV) sulfide, 4904
 Titanium salts, 4920
 Titanium tetrachloride, 4176
 Titanium tetraiodide, 4638
 Titanium tetraisopropoxide, 3579
 Titanium tetraperchlorate, 4170
 Titanium trichloride, 4158
 Titanium–zirconium alloys, 4921
 Titanium–zirconium hydride, 4491
 † Toluene, 2791
 2-Toluenediazonium bromide, 2736
 2-Toluenediazonium iodide, 2751
 3-Toluenediazonium iodide, 2752
 2-Toluenediazonium perchlorate, 2740
 2-Toluenediazonium salts, 2775
 3-Toluenediazonium salts, 2774
 4-Toluenediazonium salts, 2776
 4-Toluenediazonium triiodide, 2756
 4-Toluenesulfinyl azide, 2780
 4-Toluenesulfonic acid, 2808
 4-Toluenesulfonyl azide, 2781
 4-Toluenesulfonylhydrazide, 2827
 O-4-Toluenesulfonylhydroxylamine, 2821
 2-Toluidine, 2813
 3-Toluidine, 2814
 4-Toluidine, 2815
 4-Tolylbiguanidium hydrogen dichromate, 3182
 2-Tolylcopper, 2743
 3-Tolylcopper, 2744
 4-Tolylcopper, 2745
 1,1,1-Triacetoxy-1,2-benziodoxol-3-one, 3610
 Triacetyl borate, 2390
 1,3,5-Triacetylhexahydro-1,3,5-triazine, 3183
 2,4,6-Tri(2-acetylhydrazino)-1,3,5-trinitrobenzene, 3535
 1,3,5-Triacrylylhexahydro-*sym*-triazine, *see* 1,3,5-Tris(1-oxoprop-2-enyl)-hexahydro-1,3,5-triazine, 3531

Triallylborane, *see* Tri-2-propenylborane, 3177
 Triallylchromium, 3180
 'Triallylcyanurate', *see* 2,4,6-Triallyloxy-1,3,5-triazine, 3530
 2,4,6-Triallyloxy-1,3,5-triazine, 3530
 Triallyl phosphate, 3184
 Triaminoguanidinium nitrate, 0517
 Triaminoguanidinium perchlorate, 0515
 1,3,5-Triaminotrinitrobenzene, 2324
 Triamminediperoxochromium(IV), 4231
 Triamminenitratoplatinum(II) nitrate, 4582
 Triamminetrinitrocobalt(III), 4204
 3,6,9-Triaza-11-aminoundecanol, 3100
 1,4,7-Triazacyclononanetricarbonylmolybdenum hydride perchlorate, 3185
 1,4,7-Triazacyclononanetricarbonyltungsten hydride perchlorate, 3186
 3,6,9-Triazatetracyclo[6.1.0.0^{2,4}.0^{5,7}]nonane, 2398
 Triazidoarsine, 0103
 Triazidoborane, *see* Boron triazide, 0153
 Triazidocarbenium perchlorate, 0325
 Triazidochlorosilane, 4035
 1,1,1-Triazidodinitroethane, 0683
 Triazidogallane, *see* Gallium azide, 4409
 Triazidomethane, 0389
 Triazidomethylum hexachloroantimonate, 0335
 Triazidomethylum perchlorate, *see* Triazidocarbenium perchlorate, 0325
 2,4,6-Triazido-1,3,5-triazine, 1348
 1,3,5-Triazine, 1115
 1,2,3-Triazole, 0768
 1,2,4-Triazole, 0769
 1,2,4-Triazole-3-diazonium nitrate, 0680
 1,2,4-Triazolo[4,3-a]pyridine silver nitrate, 2270
 Tribenzylarsine, 3822
 Triboron pentafluoride, 0175
 Tribromamine hexaammoniate, *see* Nitrogen tribromide hexaammoniate, 0290
 Tribromamine oxide, 0291
 Tribromomethane, *see* Bromoform, 0368
 Tribromonitromethane, 0314
 † Tribromosilane, 0287
 Tributylbismuth, 3571
 Tri-2-butylborane, 3570
miso-Tributylborane, 3569
 Tributylgallium, 3572
 Tributylindium, 3573
 Tributyl(phenylethynyl)lead, 3811
 Tributyl phosphate, 3574
 † Tributylphosphine, 3575
 2-(Tricarbonylferra)hexaborane(10), 1297

Tricarbonyl(1-methylpyrrole)chromium(0), 2932
 Tricarbonyl[(1,2,3,4,5- η)-1-methyl-1*H*-pyrrole]chromium,
 see Tricarbonyl(1-methylpyrrole)chromium(0), 2932
 Trichloroacetaldehyde oxime, 0698
 Trichloroacetic acid, 0658
 Trichloroacetonitrile, 0607
 2,3,4-Trichloroaniline, 2163
B-1,3,5-Trichloroborazine, 0174
 † 1,1,1-Trichloroethane, 0740
 1,1,2-Trichloroethane, 0741
 2,2,2-Trichloro-1,1-ethanediol, 0745
 2,2,2-Trichloroethanol, 0744
 † Trichloroethylene, 0656
 † Trichloroethylsilane, 0854
 Trichlorofluoromethane, 0330
 ‘Trichloroisocyanuric acid’, *see* 1,3,5-Trichloro-1,3,5-triazinetriene, 1039
 † Trichloromelamine, *see* 2,4,6-Tris(chloroamino)-1,3,5-triazine 1095
 Trichloromethane, *see* Chloroform, 0372
 † Trichloromethyl carbonate, 1040
 Trichloromethyl perchlorate, 0334
 Trichloromethyl peroxyacetate, 1097
 5-Trichloromethyltetrazole, 0657
 5-Trichloromethyl-1-trimethylsilyltetrazole, 1927
 Trichloronitromethane, 0331
 Trichloroperoxyacetic acid, 0659
 2,4,5-Trichlorophenol, 2107
 † Trichlorosilane, 4136
 2,4,6-Trichloro-1,3,5-triazine, 1038
 1,3,5-Trichloro-1,3,5-triazinetriene, 1039
 1,3,5-Trichloro-2,4,6-trifluoroborazine, 0173
 1,1,2-Trichlorotrifluoroethane, 0606
 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazine, 4145
 1,3,5-Trichlorotrithiahexahydro-1,3,5-triazinemolybdenum, 4142
 † Trichlorovinylsilane, 0746
 Tricyanomethane, *see* Cyanoform, 1383
 Tricyclopentadienyluranium tetrahydroaluminate, 3688
 Tridecanal, 3619
 Triethoxydialuminium tribromide, 2555
 Triethoxysilane, 2584
 † Triethylaluminium, 2553
 † Triethylamine, 2569
 Triethylamine hydrogen peroxidate, 2570
 Triethylammonium nitrate, 2580
 Triethylantimony, 2574
 Triethylarsine, 2557
 Triethylbismuth, 2561

Triethylborane, 2559
 † Triethyl borate, 2560
 Triethyldialuminium trichloride, 2556
 Triethyldiborane, 2588
 Triethylene glycol diacetate, *see* 1,2-Bis(acetoxyethoxy)ethane, 3355
 Triethylenetetramine, *see* *N,N'*-Bis(2-aminoethyl)1,2-diaminoethane, 2599
 Triethylgallium, 2563
 3,6,9-Triethyl-1,2,4,5,7,8-hexoxonane, 3196
 Triethylindium, 2564
 Triethylphosphine, 2573
 Triethylphosphinegold nitrate, 2558
 Triethylsilane, 2585b
 Triethylsilyl-1,2-bis(trimethylsilyl)hydrazine, 3586
 Triethylsilyl perchlorate, 2562
 Triethyltin hydroperoxide, 2583
 3,6,9-Triethyl-3,6,9-trimethyl-1,2,4,5,7,8-hexoxonane, 3562
 Triethynylaluminium, 2095
 Triethynylantimony, 2125
 Triethynylarsine, 2096
 1,3,5-Triethynylbenzene, 3443
 Triethynylphosphine, 2124
 Triferrocenylcyclopropenium perchlorate, 3885
 'Triflic acid', *see* Trifluoromethanesulfonic acid, 0375
 Trifluoroacetamide, 0704
 Trifluoroacetic acid, 0665
 Trifluoroacetic anhydride, 1366
 † 1,1,1-Trifluoroacetone, 1100
N-Trifluoroacetoxy-2,4-dinitroaniline, 2892
 Trifluoroacetyl azide, 0627
 2-Trifluoroacetyl-1,3,4-dioxalane, 1047
O-Trifluoroacetyl-*S*-fluoroformyl thioperoxide, 1050
 Trifluoroacetyl hypochlorite, 0594
 Trifluoroacetyl hypofluorite, 0633
 Trifluoroacetyliminoiodobenzene, 2903
 Trifluoroacetyl nitrite, 0626
 Trifluoroacetyl trifluoromethanesulfonate, 1057
 Trifluoroacryloyl fluoride, 1049
 Trifluoroamine oxide, 4337
 † 1,2,4-Trifluorobenzene, 2110
 2,2,2-Trifluorodiazaoethane, 0664
 † 1,1,1-Trifluoroethane, 0754
 † Trifluoroethylene, 0663
 Trifluoroethylene ozonide, *see* 3,3,5-Trifluoro-1,2,4-trioxolane, 0667
N,N,N'-Trifluorohexanamidine, 2448
 Trifluoromethanesulfonyl chloride, 0322
 Trifluoromethanesulfinyl fluoride, 0354

- Trifluoromethanesulfonic acid, 0375
 1,3,3-Trifluoro-2-methoxycyclopropene, 1414
 Trifluoromethyl azide, 0347
 Trifluoromethyl 3-fluorocarbonylhexafluoroperoxybutyrate, 2061
 Trifluoromethyl hypofluorite, 0353
 Trifluoromethyl iodine(III) isocyanate, 1045
 Trifluoromethyl perchlorate, 0321
 Trifluoromethyl peroxonitrate, 0346
 Trifluoromethyl peroxyacetate, 1103
 2-Trifluoromethylphenyllithium, 2655
 3-Trifluoromethylphenyllithium, 2656
 4-Trifluoromethylphenyllithium, 2657
 Trifluoromethylphosphine, 0399
 † *N*-(Trifluoromethylsulfinyl)trifluoromethylimidodisulfinyl azide, 0640
 Trifluoromethylsulfonyl azide, 0348
 Trifluoromethylsulfur trifluoride, 0363
 † 3,3,3-Trifluoropropene, 1099
N,N,N'-Trifluoropropionamide, 1172
 3,3,3-Trifluoropropyne, 1066
 Trifluoroselenium hexafluoroarsenate, 0099
 † Trifluorosilane, 4333
 Trifluorosulfur nitride, 4338
 1,3,5-Trifluorotrinitrobenzene, 2057
 3,3,5-Trifluoro-1,2,4-trioxolane, 0667
 Trigermane, 4421
 Triglycidylamine, *see* Nitrilotris(oxiranemethane), 3181
 'Triglycidylisocyanurate', *see* 1,3,5-Tris(2,3-epoxypropyl)triazine-2,4,6-trione, 3533
 Trihydrazinealuminium perchlorate, 0064
 Trihydrazinecadmium chlorate, 3950
 Trihydrazinecadmium(II) nitrate, 3955
 Trihydrazinecobalt(II) nitrate, 4205
 Trihydrazinenickel(II) nitrate, 4592
 † 2,4,6-Trihydroxy-1,3,5-triazine, 1118
 3,4,5-Triiodobenzenediazonium nitrate, 2084
 Triiodocyclopropenium iodide, 1339
 Triiodomethane, *see* Iodoform, 0376
 † Triisobutylaluminium, 3568
 Triisopropylaluminium, 3214
 Triisopropylphosphine, 3220
N,N,4-Trilithioaniline, 2180
 Trimercury tetraphosphide, 4616
 † Trimethoxymethane, *see* Trimethyl orthoformate, 1708
 † Trimethylacetonitrile, *see* Pivalonitrile, 1929
 Trimethylacetyl azide, *see* Pivaloyl azide, 1935
 † Trimethylacetyl chloride, *see* Pivaloyl chloride, 1925

- Trimethylaluminium, 1291
- † Trimethylamine, 1310
- Trimethylamine oxide, 1313
- † Trimethylamine oxide perchlorate, *see* Trimethylhydroxylammonium perchlorate, 1323
- Trimethylammonium perchlorate, 1322
- 2,4,6-Trimethylaniline, 3169
- Trimethylantimony, 1320
- Trimethylarsine, 1294
- 1,3,5-Trimethylbenzene, *see* Mesitylene, 3162
- 3,7,7-Trimethylbicyclo[4.1.0]hept-3-ene, *see* Δ^3 -Carene, 3336
- 2,6,6-Trimethylbicyclo[3.1.1]hept-2-ene, *see* 2-Pinene, 3339
- Trimethylbismuthine, 1298
- Trimethylborane, 1295
- † Trimethyl borate, 1296
- 2,4,6-Trimethylborazine, 1335
- B*-Trimethylborazine, *see* 2,4,6-Trimethylborazine, 1335
- † 2,2,3-Trimethylbutane, 2876
- † 2,3,3-Trimethylbutene, 2855
- 1,2,2-Trimethyl-1-carboxycyclopentane-3-peroxycarboxylic acid, *see* 3-Peroxyamphoric acid, 3348
- † 1,3,5-Trimethylcyclohexane, 3191
- 2,2,4-Trimethyldecahydroquinolinium picrate, 3765
- Trimethylaluminium trichloride, 1292
- Trimethyldiborane, 1334
- Trimethyl-1,2-dioxetane, 1979
- Trimethylenefuroxan, *see* 4,5-Cyclopentanofurazan-*N*-oxide, 1871
- Trimethylgallium, 1305
- Trimethylgermylphosphine, 1332
- † 2,2,5-Trimethylhexane, 3210
- † Trimethylhydroxylammonium perchlorate, 1323
- Trimethylindium, 1307
- N,N,N*-Trimethylmethaminium periodate, *see* Tetramethylammonium periodate, 1749
- N,N,N*-Trimethylmethanaminium ozonide, *see* Tetramethylammonium ozonate, 1753
- † Trimethyl orthoformate, 1708
- † 2,2,3-Trimethylpentane, 3060
- † 2,2,4-Trimethylpentane, 3061
- † 2,3,3-Trimethylpentane, 3062
- † 2,3,4-Trimethylpentane, 3063
- † 2,3,4-Trimethyl-1-pentene, 3042
- † 2,3,4-Trimethyl-2-pentene, 3044
- † 2,4,4-Trimethyl-1-pentene, 3043
- † 2,4,4-Trimethyl-2-pentene, 3045
- † 3,4,4-Trimethyl-2-pentene, 3046

Tri-4-methylphenylsilyl perchlorate, 3823
 Trimethyl phosphate, 1318
 Trimethylphosphine, 1319
 Trimethyl phosphite, 1315
 Trimethyl phosphorothioate, *see* Trimethyl thiophosphate, 1316
 Trimethylplatinum hydroxide, 1329
 Trimethylplatinum(IV) azide tetramer, *see* Dodecamethyltetraplatinum(IV) azide, 3590
 Trimethylplatinum(IV) perchlorate tetramer, *see* Dodecamethyltetraplatinum(IV) perchlorate, 3587
 2,4,6-Trimethylpyrilium perchlorate, 2992
 † Trimethylsilyl azide, 1314
 Trimethylsilyl chlorochromate, 1301
 Trimethylsilyl hydroperoxide, 1330
 Trimethylsilylmethyl lithium, 1722
 1-Trimethylsilyloxy-1-trimethylsilylphosphylidene-2,2-dimethylpropane, 3417
 Trimethylsilyl perchlorate, 1302
 3-Trimethylsilylprop-2-enyl trifluoromethanesulfonate, 2847
N-Trimethylsilyl-*N*-trimethylsilyloxyacetamide, 3375
 Trimethylstibine, *see* Trimethylantimony, 1320
 Trimethylsulfonium chloride, 1303
 Trimethylsulfoxonium bromide, 1299
 Trimethylthallium, 1321
 Trimethyl thiophosphate, 1316
 † 2,4,6-Trimethyltrioxane, 2515
 Trinitroacetonitrile, 1009
 1,3,5-Trinitrobenzene, 2116
 2,3,5-Trinitrobenzenediazonium-4-oxide, 2072
 2,4,6-Trinitrobenzene-1,3-diol, *see* Trinitroresorcinol, 2119
 2,4,6-Trinitrobenzene-1,3,5-triol, *see* Trinitrophloroglucinol, 2120
 2,4,6-Trinitrobenzoic acid, 2640
 1,1,1-Trinitroethane, 0775
 2,2,2-Trinitroethanol, 0776
 1,3,5-Trinitrohexahydro-1,3,5-triazine, 1219
 Trinitromethane, 0385
 2,4,6-Trinitrophenol, *see* Picric acid, 2118
 2,4,6-Trinitrophenyl azide, *see* Picryl azide, 2092
 Trinitrophloroglucinol, 2120
 Trinitroresorcinol, 2119
 2,4,6-Trinitrosobenzene-1,3,5-triol, *see* Trinitrosophloroglucinol, 2117
 1,3,5-Trinitrosohexahydro-1,3,5-triazine, 1217
 Trinitrosophloroglucinol, 2117
 2,4,5-Trinitrotoluene, 2702
 2,4,6-Trinitrotoluene, 2701
 2,3,7-Trioxabicyclo[2,2,1]hept-5-ene, 1444
 1,3,5-Trioxane, 1237

1,2,4-Trioxolane, *see* Ethylene ozonide, 0836
 'Trixygen difluoride' 4323
 Trixygen, *see* Ozone, 4846
 Triphenylaluminium, 3749
 Triphenylchromium tetrahydrofuranate, 3751
 3,3,5-Triphenyl-4,4-dimethyl-5-hydroperoxy-4,5-dihydro(3*H*)pyrazole, 3841
 Triphenyllead nitrate, 3753
 Triphenylmethyl azide, 3783
 Triphenylmethyl nitrate, 3782
 Triphenylmethylpotassium, 3781
 1,3,5-Triphenyl-1,4-pentaazadiene, 3754
 Triphenylphosphine, 3756
 Triphenylphosphine oxide hydrogen peroxidate, 3755
 Triphenylphosphine oxide-oxodiperoxochromium(VI), 3752
 2,4,6-Triphenylpyrilium perchlorate, 3839
 Triphenylsilyl perchlorate, 3750
 Triphenyltin hydroperoxide, 3758
 † Triphosgene, *see* Trichloromethyl carbonate, 1040
 Triphosphorus pentanitride, 4776
 Tripotassium hexafluoroferrate(3-), 4359
 Tri-2-propenylborane, 3177
 Tripropylaluminium, 3215
 Tripropylantimony, 3221
 Tripropylborane, 3217
 Tripropylindium, 3219
 Tripropyllead fulminate, 3363
 Tripropylsilyl perchlorate, 3218
 Trirhenium nonachloride, 4196
 1,1,1-Tris(aminomethyl)ethane, 2044
 Tris(2-azidoethyl)amine, 2482
 1,1,1-Tris(azidomethyl)ethane, 1937
 1,3,5-Tris(4-azidosulfonylphenyl)-1,3,5-triazinetriene, 3818
 Tris(2,2'-bipyridine)chromium(0), 3875
 Tris(2,2'-bipyridine)chromium(II) perchlorate, 3874
 Tris(2,2'-bipyridine)silver(II) perchlorate, 3873
 Tris(bis-2-methoxyethyl ether)potassium hexacarbonylniobate(1-), 3856
 2,4,6-Tris(bromoamino)-1,3,5-triazine (Tribromomelamine), 1091
 2,2,2-Tris(bromomethyl)ethanol, 1922
 1,1,1-Tris(bromomethyl)methane, 1550
 † 2,4,6-Tris(chloroamino)-1,3,5-triazine, 1095
 Tris(cyclopentadienyl)cerium, 3683
 Tris(cyclopentadienyl)plutonium, 3684
 Tris(cyclopentadienyl)uranium, 3685
 Tris(2,3-diaminobutane)nickel(II) nitrate, 3589
 Tris(1,2-diaminoethane)chromium(III) perchlorate, 2619
 Tris(1,2-diaminoethane)cobalt(III) nitrate, 2622

Tris(1,2-diaminoethane)ruthenium(III) perchlorate, 2621
 2,4,6-Tris(dichloroamino)-1,3,5-triazine, 1043
 Tris(difluoroamino)fluoromethane, 0364
 Tris(dimethylamino)antimony, 2598
 2,4,6-Tris(dimethylaminomethyl)phenol, 3694
 Tris(2,2-dimethylethyl)triphosphirane, 3576
 Tris(dimethylfluorosilylmethyl)borane, 3224
 1,3,5-Tris(2,3-epoxypropyl)triazine-2,4,6-trione, 3533
 Tris(ethylthio)cyclopropenium perchlorate, 3179
 Tris(hydroxymethyl)methylamine, 1731
 Tris(hydroxymethyl)nitromethane, 1664
 † Trisilane, 4579
 2,4,6-Trisilatrioxane, 4566
 Trisilver tetranitride, 0040
 Trisilylamine, 4580
 Trisilylarsine, 0102
 Trisilylphosphine, 4583
 Tris(iodomercurio)phosphine, 4614
 Tris(4-methoxy-2,2''bipyridine)ruthenium(II) perchlorate, 3886
 Tris(3-methylpyrazole)zinc sulfate, 3540
 1,3,5-Tris(nitromethyl)benzene, 3145
 1,3,5-Tris(1-oxoprop-2-enyl)-hexahydro-1,3,5-triazine, 3531
 Tris(pentafluorophenyl)aluminium etherate, *see* [1,1'-Oxybis[ethane]]
 tris(pentafluorophenyl)aluminium, 3829
 Tris(pentafluorophenyl)boron, 3738
 Tris(2,4-pentanedionato)molybdenum(III), 3690
 Tris(1,10-phenanthroline)cobalt(III) perchlorate, 3890
 Tris(1,10-phenanthroline)ruthenium(II) perchlorate, 3889
 Tri(spirocyclopentane)1,1,4,4,7,7-hexoxonane, 3693
 Tris(2-propylthio)silane, 3222
 1,1,3,3,5,5-Tris-spiro(*N,N'*-dinitroethylenediamino)cyclotriphosphazene, 2483
 Tris(thionitrosyl)thallium, 4764
 Tris(trifluoromethyl)nitrosomethane, *see* Perfluoro-*tert*-nitrosobutane, 1370
 Tris(trifluoromethyl)phosphine, 1063
 Tris(trimethylsilyl)aluminium, 3227
 Tris(trimethylsilyl)aluminium etherate, 3620
N,N,N'-Tris(trimethylsilyl)diaminophosphine, 3232
 Tris(trimethylsilyl)hydrazine, 3230
 Tris(trimethylsilylmethyl)aluminium, 3584
 Tris(trimethylsilyl)methylcaesium, 3381
 Tris(trimethylsilylmethyl)indium, 3585
 Tris(trimethylsilyl)phosphine, 3229
 Tris(trimethylsilyl)silane, 3231
 Tris(2,2,2-trinitroethyl) orthoformate, 2786
 Tritellurium tetranitride, 4772
 1,4,7-Trithia[7]ferrocenophene–acetonitrilecopper(I) perchlorate, 3710

Trithorium tetranitride, 4774
Tri(4-tolyl)ammonium perchlorate, 3824
Triuranium octaoxide, 4871
Trivinylantimony, 2405
Trivinylbismuth, 2391
Tropylium perchlorate, 2742
Tungsten, 4925
Tungsten azide pentabromide, 0297
Tungsten azide pentachloride, 4182
Tungsten carbide, 0563
Tungsten dichloride, 4119
Tungsten diiodide, 4631
Tungsten hexachloride, 4193
Tungsten hexafluoride, 4376
Tungsten hexamethoxide, 2604
Tungsten(IV) oxide, 4844
Tungsten tetrabromide oxide, 0294
Tungsten(VI) oxide, 4856

UndecaamminetetRARuthenium dodecaoxide, 4598
'Unsaturated' Silicon hydride, 4452
Uranium, 4923
Uranium azide pentachloride, 4181
Uranium carbide, 0562
Uranium dicarbide, 1028
Uranium hexachloride, 4192
Uranium hexafluoride, 4375
Uranium(III) hydride, 4511
Uranium(III) nitride, 4732
Uranium(III) tetrahydroborate, 0178
Uranium(IV) hydride, 4541
Uranium(IV) oxide, 4843
Uranium(IV) sulfide, 4905
Uranium(IV) tetrahydroborate etherates, 0183
Uranyl nitrate, 4752
Uranyl perchlorate, 4111
Urea, 0475
Urea hydrogen peroxidate, 0476
Urea nitrate, *see* Uronium nitrate, 0494
Urea perchlorate, *see* Uronium perchlorate, 0491
Uronium nitrate, 0494
Uronium perchlorate, 0491

† Valeraldehyde, 1966
Vanadium, 4924

Vanadium azide tetrachloride, 4166
Vanadium dichloride, 4118
Vanadium(III) oxide, 4855
Vanadium tetrachloride, 4177
Vanadium tribromide oxide, 0292
Vanadium trichloride, 4159
Vanadium trichloride oxide, 4151
Vanadium trinitrate oxide, 4763
Vanadium(V) oxide, 4866
Vanadyl azide dichloride, 4090
Vanadyl chloride, *see* Vanadium trichloride oxide, 4151
Vanadyl perchlorate, 4152
Vanillin, *see* 4-Hydroxy-3-methoxybenzaldehyde, 2958
† Vinyl acetate, 1532
Vinyl acetate ozonide, 1543
† Vinylacetylene, *see* Buten-3-yne, 1423
Vinylamine, 0864
Vinyl azide, 0770
† Vinyl bromide, *see* Bromoethylene, 0727
† Vinyl butyrate, 2439
† Vinyl chloride, *see* Chloroethylene, 0730
† Vinylcyclohexane, 3026
† 4-Vinylcyclohexene, 3004
† Vinyl diazomethane, *see* 3-Diazopropene, 1135
† Vinylene carbonate, *see* 1,3-Dioxol-4-en-2-one, 1087
† Vinyl fluoride, *see* Fluoroethylene, 0747
† Vinyl formate, 1149
† Vinylidene chloride, *see* 1,1-Dichloroethylenem, 0695
† Vinylidene fluoride, *see* 1,1-Difluoroethylene, 0700
Vinyl isocyanide, 1109
Vinyl lithium, 0757
Vinylmagnesium chloride, Chloroethenylmagnesium, *see* Ethenylmagnesium chloride, 0732
† Vinyloxirane, *see* 3,4-Epoxybutene, 1520
† Vinyl propionate, 1918
Vinylpyridine, 2759

Water, *see* Diprotium monoxide, 4475

Xenon, 4926
Xenon difluoride, 4332
Xenon difluoride dioxide, 4322
Xenon difluoride oxide, 4319
Xenon hexafluoride, 4377
Xenon(II) fluoride methanesulfonate, 0443

Xenon(II) fluoride perchlorate, 3977
Xenon(II) fluoride trifluoroacetate, 0634
Xenon(II) fluoride trifluoromethanesulfonate, 0356
Xenon(II) pentafluoroorthoselenate, 4382
Xenon(II) pentafluoroorthotellurate, 4383
Xenon(II) perchlorate, 4110
Xenon(IV) hydroxide, 4533
Xenon tetrafluoride, 4353
Xenon tetrafluoride oxide, 4346
Xenon tetraoxide, 4863
Xenon trioxide, 4857
† *m*-Xylene, 2971
† *o*-Xylene, 2970
† *p*-Xylene, 2972
† *mixo*-Xylene, 2969

Yttrium 4-nitrobenzoate trihydrate, 3817

Zinc, 4927
Zinc abietate, 3903
Zinc amalgam, 4601
Zinc azide, 4786
Zinc bis(1-benzeneazothiocarbonyl-2-phenyl-2-hydrazide), 3863
Zinc bromate, 0279
Zinc chlorate, 4106
Zinc chloride, 4120
Zinc cyanide, 1004
Zinc dihydrazide, 4563
Zinc ethoxide, 1704
Zinc ethylsulfinate, 1711
Zinc hydride, 4492
Zinc hydroxide, 4478
Zinc iodide, 4632
Zinc nitrate, 4751
Zinc oxide, 4830
Zinc permanganate, 4710
Zinc peroxide, 4845
Zinc phosphide, 4876
Zinc picrate, 3437
Zinc stearate, 3897
Zirconium, 4928
Zirconium carbide, 0565
Zirconium dibromide, 0285
Zirconium dicarbide, 1029
'Zirconium hydride', 4493

Zirconium hydroxide, 4534
Zirconium(II) chloride, 4121
Zirconium(IV) tetrahydroborate, 0184
Zirconium nitride, 4733
Zirconium oxide sulfide, 4827
Zirconium salts 4929
Zirconium tetrachloride, 4178
Zirconium tetraiodide, 4639
Zirconium trichloride, 4160

Index of CAS Registry Numbers vs Serial Numbers in Section 1

The CAS Registry numbers are the key to the full structural and stereochemical information held within the CAS chemical identification system, available both in printed form and on-line via several hosts worldwide.

The Index is arranged in numerically increasing order of the registry numbers. The serial numbers of those compounds for which no registry number has been found are listed separately at the end of the Index.

[50-00-0], 0416	[62-23-7], 2695	[74-82-8], 0466	[75-09-2], 0397	[75-47-8], 0376
[50-71-5], 1398	[62-53-3], 2354	[74-83-9], 0429	[75-11-6], 0400	[75-50-3], 1310
[50-78-2], 3137	[62-56-6], 0480	[74-84-0], 0881	[75-12-7], 0453	[75-52-5], 0456
[50-99-7], 2518	[63-42-3], 3557	[74-85-1], 0781	[75-15-0], 0560	[75-54-7], 0470
[51-17-2], 2724	[64-17-5], 0920	[74-86-2], 0686	[75-18-3], 0932	[75-55-8], 1257
[51-28-5], 2197	[64-18-6], 0418	[74-87-3], 0432	[75-19-4], 1197	[75-56-9], 1225
[51-79-6], 1261	[64-19-7], 0833	[74-88-4], 0445	[75-20-7], 0585	[75-62-7], 0310
[54-87-5], 2908	[64-67-5], 1710	[74-89-5], 0493	[75-21-8], 0829	[75-63-8], 0311
[55-63-0], 1196	[65-85-0], 2732	[74-90-8], 0380	[75-24-1], 1291	[75-64-9], 1725
[56-18-8], 2587	[67-56-1], 0484	[74-93-1], 0489	[75-25-2], 0368	[75-65-0], 1696
[56-23-5], 0332	[67-63-0], 1280	[74-94-5], 0746	[75-26-3], 1242	[75-66-1], 1716
[56-81-5], 1286	[67-64-1], 1220	[74-95-3], 0395	[75-28-5], 1669	[75-68-3], 0731
[57-06-7], 1471	[67-66-3], 0372	[74-96-4], 0846	[75-29-6], 1244	[75-69-4], 0330
[57-13-6], 0475	[67-72-1], 0611	[74-98-6], 1271	[75-30-9], 1252	[75-71-8], 0326
[57-14-7], 0954	[68-12-2], 1259	[74-99-7], 1125	[75-31-0], 1308	[75-73-0], 0349
[57-50-1], 3558	[68-39-3], 1137	[75-00-3], 0848	[75-33-2], 1290	[75-74-1], 1767
[57-55-6], 1285	[69-68-5], 0921	[75-01-4], 0730	[75-34-3], 0790	[75-75-2], 0488
[57-71-6], 1570	[70-25-7], 0876	[75-02-5], 0747	[75-35-4], 0695	[75-76-3], 1770
[60-09-3], 3487	[70-34-8], 2108	[75-03-6], 0858	[75-36-5], 0735	[75-77-4], 1304
[60-12-8], 2983	[71-23-8], 1279	[75-04-7], 0942	[75-37-6], 0801	[75-78-5], 0902
[60-29-7], 1697	[71-43-2], 2288	[75-05-8], 0758	[75-38-7], 0700	[75-79-6], 0439
[60-34-4], 0503	[71-55-6], 0740	[75-07-0], 0828	[75-44-5], 0329	[75-83-2], 2529
[60-51-5], 2009	[72-20-8], 3459	[75-08-1], 0933	[75-45-6], 0369	[75-85-4], 2015

[75-86-5], 1566	[79-10-7], 1148	[91-22-5], 3121	[97-96-1], 2487	[100-99-2], 3568
[75-91-2], 1698	[79-15-2], 0784	[91-23-6], 2768	[97-99-4], 1978	[101-25-7], 1954
[75-97-8], 2486	[79-17-4], 0507	[91-43-0], 3301	[98-00-0], 1880	[101-38-2], 2078
[76-01-7], 0660	[79-20-9], 1232	[91-59-8], 3265	[98-01-1], 1836	[101-57-9], 1600
[76-03-9], 0658	[79-21-0], 0837	[92-87-5], 3516	[98-09-9], 2235	[101-84-8], 3492
[76-05-1], 0665	[79-22-1], 0736	[93-55-0], 3151	[98-56-6], 2646	[102-54-5], 3276
[76-06-2], 0331	[79-24-3], 0869	[93-59-4], 2733	[98-82-9], 2239	[102-67-0], 3215
[76-13-1], 0606	[79-29-8], 2530	[94-36-0], 3639	[98-88-4], 2675	[103-30-0], 3643
[76-14-2], 0604	[79-30-1], 1558	[94-82-6], 3272	[98-95-3], 2262	[103-33-3], 3483
[76-37-9], 1133	[79-34-5], 0699	[94-87-1], 2975	[99-09-2], 2314	[103-34-4], 3047
[77-47-4], 1808	[79-37-8], 0605	[95-14-7], 2269	[99-35-4], 2116	[103-44-6], 3360
[77-73-6], 3299	[79-38-9], 0591	[95-45-4], 1595	[99-54-7], 2105	[103-50-4], 3655
[77-76-9], 2018	[79-41-4], 1530	[95-47-6], 2970	[99-55-8], 2801	[103-69-5], 2994
[77-78-1], 0929	[79-46-9], 1264	[95-50-1], 2156	[99-59-2], 2802	[103-71-9], 2685
[77-86-1], 1731	[80-05-7], 3686	[95-51-2], 2300	[99-61-6], 2687	[104-03-0], 2938
[77-92-9], 2389	[80-09-1], 3497	[95-53-4], 2813	[99-63-8], 2888	[104-04-1], 2950
[78-00-2], 3095	[80-15-9], 3166	[95-69-2], 2794	[99-65-0], 2195	[104-12-1], 2648
[78-11-5], 1901	[80-51-3], 3525	[95-76-1], 2241	[99-83-2], 3338	[104-15-4], 2808
[78-18-2], 3555	[80-56-8], 3339	[95-81-8], 2793	[99-91-2], 2929	[104-51-8], 3323
[78-62-6], 2582	[80-62-6], 1915	[95-83-0], 2347	[99-99-0], 2764	[104-55-2], 3134
[78-67-1], 3011	[81-15-2], 3532	[95-85-2], 2303	[100-00-5], 2142	[104-82-5], 2959
[78-76-2], 1632	[82-05-3], 3736	[95-94-3], 2079	[100-00-5], 2142	[104-90-5], 2995
[78-77-3], 1633	[82-34-8], 3624	[95-95-4], 2107	[100-01-6], 2315	[104-91-6], 2264
[78-78-4], 2000	[82-71-3], 2119	[96-10-6], 1671	[100-02-7], 2267	[105-29-3], 2384
[78-79-5], 1893	[83-80-7], 3845	[96-14-0], 2533	[100-07-2], 2930	[105-37-3], 1970
[78-80-8], 1858	[84-74-2], 3714	[96-22-0], 1963	[100-14-1], 2714	[105-38-4], 1918
[78-81-9], 1728	[85-44-9], 2899	[96-24-2], 1248	[100-15-2], 2800	[105-39-5], 1559
[78-82-0], 1565	[86-30-6], 3486	[96-29-7], 1654	[100-20-9], 2889	[105-46-4], 2497
[78-83-1], 1694	[86-50-0], 3307	[96-33-3], 1531	[100-21-0], 2924	[105-56-6], 1889
[78-84-2], 1611	[86-57-7], 3249	[96-37-7], 2461	[100-25-4], 2196	[105-57-7], 2545
[78-85-3], 1522	[87-31-0], 2087	[96-40-2], 1885	[100-26-5], 2696	[105-58-8], 1981
[78-86-4], 1638	[87-33-2], 2380	[96-47-9], 1961	[100-34-5], 2230	[105-60-2], 2451
[78-87-5], 1204	[87-68-3], 1364	[96-48-0], 1527	[100-39-0], 2735	[105-64-6], 3034
[78-88-6], 1132	[87-69-4], 1545	[96-50-4], 1140	[100-40-3], 3004	[105-74-8], 3857
[78-90-0], 1325	[87-90-1], 1039	[96-54-8], 1887	[100-41-4], 2968	[106-40-1], 2296
[78-92-2], 1695	[88-05-1], 3169	[96-73-1], 2144	[100-42-5], 2945	[106-42-3], 2972
[78-93-3], 1606	[88-72-2], 2763	[96-75-3], 2319	[100-44-7], 2738	[106-47-8], 2302
[78-94-4], 1515	[88-73-3], 2141	[96-91-3], 2278	[100-46-9], 2812	[106-49-0], 2815
[78-95-5], 1161	[88-74-4], 2313	[97-00-7], 2098	[100-51-6], 2806	[106-50-3], 2371
[78-96-6], 1311	[88-75-5], 2265	[97-02-9], 2276	[100-52-7], 2731	[106-51-4], 2214
[78-99-9], 1203	[88-85-7], 3306	[97-02-9], 2277	[100-58-3], 2225	[106-86-1], 2224
[79-00-5], 0741	[88-89-1], 2118	[97-06-3], 2773	[100-63-0], 2373	[106-88-7], 1609
[79-01-6], 0656	[88-99-3], 2923	[97-62-1], 2499	[100-64-1], 2452	[106-89-8], 1162
[79-03-8], 1163	[90-04-0], 2816	[97-63-2], 2438	[100-65-2], 2356	[106-92-3], 2434
[79-06-1], 1180	[90-72-2], 3694	[97-93-8], 2553	[100-94-9], 2818	[106-93-4], 0785
[79-07-2], 0788	[91-20-3], 3254	[97-94-9], 2559	[100-97-0], 2477	[106-94-5], 1241

[106-95-6], 1153	[108-21-4], 1972	[109-86-4], 1284	[111-40-0], 1777	[121-03-9], 2772
[106-96-7], 1090	[108-23-6], 1560	[109-87-5], 1281	[111-41-1], 1756	[121-14-2], 2726
[106-97-8], 1668	[108-24-7], 1534	[109-89-7], 1726	[111-43-3], 2543	[121-33-5], 2958
[106-98-9], 1577	[108-30-5], 1443	[109-92-2], 1610	[111-46-6], 1705	[121-43-7], 1296
[106-99-0], 1480	[108-31-6], 1404	[109-93-3], 1519	[111-49-9], 2526	[121-44-8], 2569
[107-00-6], 1481	[108-38-3], 2971	[109-94-4], 1230	[111-65-9], 3059	[121-45-9], 1315
[107-02-8], 1145	[108-40-7], 2809	[109-95-5], 0867	[111-66-0], 3040	[121-46-0], 2789
[107-03-9], 1289	[108-42-9], 2301	[109-97-7], 1466	[111-67-1], 3041	[121-66-4], 1117
[107-05-1], 1158	[108-44-1], 2814	[109-99-9], 1612	[111-77-3], 2019	[121-69-7], 2993
[107-06-2], 0791	[108-46-3], 2332	[110-00-9], 1439	[111-84-2], 3209	[121-82-4], 1219
[107-10-8], 1309	[108-58-7], 3291	[110-01-0], 1627	[111-92-2], 3083	[121-86-8], 2711
[107-11-9], 1254	[108-64-5], 2859	[110-02-1], 1448	[111-94-4], 2397	[121-88-0], 2316
[107-12-0], 1179	[108-67-3], 3162	[110-05-4], 3074	[111-96-6], 2549	[121-92-6], 2694
[107-13-1], 1107	[108-72-5], 1916	[110-17-8], 1446	[112-24-3], 2599	[122-39-4], 3504
[107-15-3], 0953	[108-77-0], 1038	[110-18-9], 2579	[112-34-5], 3077	[122-40-7], 3663
[107-16-4], 0760	[108-80-5], 1118	[110-19-0], 2502	[112-36-7], 3076	[122-56-5], 3569
[107-18-6], 1223	[108-87-2], 2854	[110-22-2], 1537	[112-57-2], 3101	[122-60-1], 3150
[107-19-7], 1147	[108-88-3], 2791	[110-44-1], 2385b	[112-58-3], 3563	[122-82-7], 3529
[107-21-1], 0924	[108-90-7], 2228	[110-45-2], 2503	[112-73-2], 3564	[122-97-4], 3165
[107-22-2], 0723	[108-91-8], 2521	[110-46-3], 1996	[112-77-6], 3772	[123-05-7], 3050
[107-25-5], 1221	[108-93-0], 2485	[110-51-0], 0139	[112-80-1], 3774	[123-11-5], 2956
[107-29-9], 0865	[108-94-1], 2429	[110-54-3], 2531	[115-00-6], 3465	[123-15-9], 2491
[107-30-2], 0850	[108-95-2], 2330	[110-58-7], 2034	[115-07-1], 1198	[123-20-6], 2439
[107-31-3], 0834	[108-98-5], 2344	[110-61-2], 1433	[115-10-6], 0919	[123-23-9], 2990
[107-32-4], 0420	[109-02-4], 1995	[110-62-3], 1966	[115-11-7], 1582	[123-25-1], 3031
[107-39-1], 3043	[109-05-7], 2523	[110-65-6], 1526	[115-18-4], 1960	[123-31-9], 2333
[107-40-4], 3045	[109-06-8], 2355	[110-66-7], 2025	[115-19-5], 1910	[123-33-1], 1435
[107-46-0], 2601	[109-53-5], 2490	[110-68-9], 2033	[115-20-8], 0744	[123-38-6], 1224
[107-71-1], 2509	[109-54-6], 2003	[110-71-4], 1701	[115-21-9], 0854	[123-39-7], 0866
[107-81-3], 1985	[109-55-7], 2039	[110-74-7], 1621	[115-77-5], 2021	[123-42-2], 2501
[107-82-4], 1984	[109-60-4], 1977	[110-75-8], 1557	[116-09-6], 1231	[123-51-3], 2014
[107-83-5], 2532	[109-63-7], 1674	[110-80-5], 1702	[116-14-3], 0628	[123-63-7], 2515
[107-84-6], 1986	[109-65-9], 1631	[110-82-7], 2457	[116-15-4], 1051	[123-72-8], 1607
[107-85-7], 2032	[109-66-0], 2001	[110-83-8], 2406	[118-52-5], 1865	[123-75-1], 1653
[107-87-9], 1962	[109-69-3], 1637	[110-85-0], 1689	[118-74-1], 2055	[123-77-3], 0816
[107-92-6], 1614	[109-71-7], 1073	[110-86-1], 1848	[118-96-7], 2701	[123-86-4], 2496
[108-00-9], 1755	[109-72-8], 1651	[110-87-2], 1908	[119-26-6], 2321	[123-91-1], 1617
[108-01-0], 1729	[109-73-9], 1723	[110-88-3], 1237	[119-33-5], 2767	[123-92-2], 2861
[108-03-2], 1263	[109-74-0], 1563	[110-89-4], 1993	[119-64-2], 3300	[123-99-9], 3187
[108-05-4], 1532	[109-75-1], 1465	[110-91-8], 1657	[119-90-4], 3660	[124-02-7], 2450
[108-08-7], 2872	[109-76-1], 1946	[110-96-3], 3085	[120-12-7], 3632	[124-04-9], 2441
[108-09-8], 2567	[109-76-2], 1326	[111-15-9], 2511	[120-37-0], 3691	[124-18-5], 3365
[108-10-1], 2495	[109-77-3], 1078	[111-19-3], 3341	[120-61-6], 3292	[124-38-9], 0557
[108-18-9], 2566	[109-78-4], 1181	[111-21-7], 3355	[120-80-9], 2331	[124-40-3], 0941
[108-19-0], 0874	[109-79-5], 1712	[111-34-2], 2484	[120-92-3], 1905	[124-41-4], 0464
[108-20-3], 2542	[109-85-3], 1312	[111-36-4], 1931	[120-94-5], 1992	[124-43-6], 0476

[124-47-0], 0494	[142-84-7], 2568	[305-80-6], 2193	[402-26-6], 2643	[501-53-1], 2931
[124-70-9], 1208	[142-96-1], 3071	[311-45-5], 3329	[402-31-3], 2894	[501-65-5], 3634
[126-11-4], 1664	[143-33-9], 0536	[327-54-8], 2081	[402-51-7], 2643	[502-26-1], 3773
[126-33-0], 1622	[144-55-8], 0390	[327-92-4], 2080	[407-25-0], 1366	[502-47-6], 3352
[126-39-6], 2500	[144-62-7], 0725	[328-75-6], 2885	[420-04-2], 0404	[502-98-7], 0792
[126-73-8], 3574	[144-86-5], 2797	[333-20-0], 0529	[420-12-2], 0840	[503-09-3], 1171
[126-76-1], 3078	[148-01-6], 2941	[334-88-3], 0406	[420-23-5], 0885	[503-17-3], 1482
[126-99-8], 1451	[149-73-5], 1708	[334-99-6], 0345	[420-46-2], 0754	[503-28-6], 0910
[127-06-0], 1258	[150-46-9], 2560	[335-01-3], 0357	[420-52-0], 0399	[503-29-7], 1255
[127-07-1], 0477	[151-00-8], 1655	[338-66-9], 0344	[420-87-1], 0705	[503-30-0], 1222
[127-09-3], 0779	[151-18-8], 1212	[348-51-6], 2134	[421-17-0], 0322	[503-41-3], 0839
[127-17-3], 1150	[151-50-8], 0526	[352-32-9], 2749	[421-50-1], 1100	[503-49-1], 2445
[127-18-4], 0609	[151-56-4], 0863	[352-33-0], 2136	[422-61-7], 1054	[503-80-0], 1737
[127-19-5], 1656	[152-18-1], 1316	[352-70-5], 2748	[429-41-4], 3728	[504-03-0], 2868
[127-52-6], 2229	[156-59-2], 0697	[352-93-2], 1714	[431-03-8], 1525	[504-60-9], 1896
[127-65-1], 2739	[156-60-5], 0696	[353-07-1], 1275	[431-97-0], 1053	[504-64-3], 1349
[127-68-4], 2184	[156-62-7], 0316	[353-36-6], 0855	[432-04-2], 1063	[505-14-6], 1001
[128-08-5], 1425	[157-22-2], 0405	[353-50-4], 0343	[446-35-5], 2109	[505-22-6], 1616
[128-09-6], 1427	[183-84-6], 3546	[354-38-1], 0704	[446-36-6], 2167	[505-60-2], 1589
[129-66-8], 2640	[220-42-8], 3630	[354-93-8], 1370	[454-50-2], 1383	[505-71-5], 0916
[131-62-4], 2681	[262-38-4], 2478	[355-43-1], 2062	[460-12-8], 1385	[506-63-8], 0890
[131-74-8], 2322	[273-77-8], 2199	[355-66-8], 2170	[460-19-5], 0996	[506-64-9], 0299
[131-91-9], 3250	[277-10-1], 2942	[359-11-5], 0663	[460-35-5], 1127	[506-65-0], 0308
[133-14-2], 3623	[280-57-9], 2473	[359-46-6], 0633	[461-58-5], 0813	[506-68-3], 0313
[133-55-1], 3284	[283-66-9], 2476	[359-48-8], 0666	[462-06-6], 2248	[506-77-4], 0323
[134-32-7], 3264	[285-67-6], 1906	[367-23-7], 2110	[462-95-3], 2016	[506-82-1], 0894
[135-20-6], 2399	[287-23-0], 1580	[368-49-0], 2656	[463-04-7], 1998	[506-85-4], 0381
[136-35-6], 3506	[287-92-3], 1939	[369-57-3], 2223	[463-40-1], 3771	[506-96-7], 0728
[138-89-6], 2980	[288-14-2], 1110	[371-67-5], 0664	[463-49-0], 1124	[507-09-5], 0832
[140-29-4], 2935	[288-42-6], 1111	[371-69-7], 0856	[463-51-4], 0722	[507-16-4], 0274
[140-88-5], 1913	[288-88-0], 0769	[371-78-8], 0643	[463-58-1], 0556	[507-19-7], 1634
[140-89-6], 1176	[288-94-8], 0411	[372-09-8], 1113	[463-82-1], 1999	[507-20-0], 1640
[141-43-5], 0943	[289-14-5], 0836	[372-18-9], 2168	[464-06-2], 2876	[507-25-5], 0525
[141-52-6], 0878	[289-19-0], 4443	[372-48-5], 1831	[464-10-8], 0314	[507-40-4], 1642
[141-75-3], 1555	[290-87-9], 1115	[372-64-5], 0644	[471-25-0], 1086	[509-14-8], 0546
[141-78-6], 1618	[290-96-0], 0715	[373-91-1], 0353	[471-34-1], 0317	[512-35-6], 3498
[141-82-2], 1151	[291-50-9], 4566	[374-10-7], 0363	[471-35-2], 1736	[512-56-1], 1318
[141-84-4], 1112	[291-64-5], 2849	[375-31-3], 1541	[473-34-7], 2244	[512-85-6], 3347
[141-97-9], 2440	[294-93-9], 0830	[377-37-7], 1430	[475-08-1], 1116	[513-35-9], 1943
[142-04-1], 2365	[298-00-0], 2979	[377-38-8], 1391	[479-45-8], 2706	[513-36-0], 1639
[142-25-6], 2038	[298-12-4], 0724	[381-73-7], 0702	[480-96-6], 2190	[513-38-2], 1648
[142-29-0], 1891	[298-18-0], 1507	[383-63-1], 1460	[495-48-7], 3485	[513-44-0], 1715
[142-45-0], 1405	[298-18-0], 1529	[383-73-3], 1367	[497-18-7], 0508	[513-48-4], 1647
[142-68-7], 1965	[300-87-8], 1888	[392-56-3], 2060	[497-19-8], 0552	[513-53-1], 1713
[142-82-5], 2873	[302-01-2], 4520	[393-75-9], 2637	[499-06-9], 3152	[513-81-5], 2407
[142-83-6], 2383	[302-17-0], 0745	[395-47-1], 2643	[500-28-7], 2961	[513-92-8], 0986

[517-25-9], 0385	[544-92-3], 0616	[565-75-3], 3063	[592-55-2], 1635	[598-61-8], 1945
[526-84-1], 1447	[544-92-3], 0620	[565-76-4], 3042	[592-57-4], 2361	[598-63-0], 0558
[528-29-0], 2194	[544-97-8], 0935	[565-77-5], 3044	[592-76-7], 2851	[598-63-0], 0726
[528-76-7], 2099	[545-06-2], 0607	[565-80-0], 2857	[592-77-8], 2852	[598-73-2], 0579
[529-27-8], 3057	[546-68-9], 3579	[573-83-1], 2085	[592-78-9], 2853	[598-74-3], 2029
[529-34-0], 3289	[547-63-7], 1976	[576-44-6], 1468	[592-84-7], 1967	[598-82-3], 1234
[530-47-2], 3519	[552-16-9], 2693	[582-61-6], 2698	[592-85-8], 0979	[598-96-9], 3046
[530-48-3], 3642	[552-89-6], 2686	[584-08-7], 0531	[592-87-0], 1000	[598-98-1], 2506
[530-50-7], 3517	[554-00-7], 2238	[584-13-4], 0812	[592-88-1], 2447	[598-99-2], 1096
[530-85-8], 3113	[554-12-1], 1620	[584-84-9], 3117	[593-53-3], 0441	[600-26-0], 0810
[533-51-7], 0572	[554-13-2], 0533	[584-94-1], 3054	[593-57-7], 0937	[602-03-9], 2276
[534-15-6], 1700	[554-14-3], 1882	[586-77-6], 2973	[593-60-2], 0727	[602-99-3], 2703
[534-22-5], 1878	[554-70-1], 2573	[586-86-9], 2261	[593-61-3], 0651	[603-11-2], 2906
[536-33-4], 2981	[554-81-7], 2266	[586-92-5], 1826	[593-63-5], 0652	[603-11-2], 2907
[536-74-3], 2912	[555-16-5], 2688	[587-85-9], 3480	[593-67-9], 0864	[603-35-0], 3756
[536-80-1], 2252	[555-31-2], 3216	[589-34-4], 2875	[593-68-0], 0949	[603-71-4], 3158
[536-90-3], 2817	[555-55-4], 3482	[589-38-8], 2489	[593-74-8], 0907	[605-71-0], 3242
[537-45-1], 2076	[556-24-1], 2505	[589-43-5], 3055	[593-75-9], 0759	[606-22-4], 2276
[538-37-4], 3650	[556-52-5], 1229	[589-81-1], 3058	[593-78-2], 0434	[608-27-5], 2237
[540-36-3], 2169	[556-56-9], 1174	[589-92-4], 2843	[593-88-4], 1294	[608-31-1], 2240
[540-42-1], 2860	[556-65-0], 0532	[590-18-1], 1579	[593-89-5], 0424	[608-73-1], 2306
[540-47-6], 1608	[556-88-7], 0482	[590-19-2], 1479	[593-90-8], 1295	[609-26-7], 3056
[540-54-5], 1243	[556-89-8], 0460	[590-21-6], 1156	[593-91-9], 1298	[609-99-4], 2665
[540-67-0], 1278	[557-05-1], 3897	[590-28-3], 0528	[594-09-2], 1319	[610-14-0], 2653
[540-72-7], 0538	[557-17-5], 1703	[590-36-3], 2544	[594-10-5], 1320	[610-25-3], 2702
[540-73-8], 0955	[557-18-6], 1687	[590-86-3], 1958	[594-11-6], 1581	[610-41-3], 2276
[540-80-7], 1659	[557-19-7], 0997	[591-09-3], 0765	[594-19-4], 1652	[611-06-3], 2104
[540-84-1], 3061	[557-20-0], 1718	[591-10-6], 1738	[594-20-7], 1205	[611-70-1], 3311
[541-31-1], 2024	[557-21-1], 1004	[591-24-2], 2842	[594-27-4], 1771	[612-08-8], 3087
[541-41-3], 1164	[557-31-3], 1955	[591-47-9], 2839	[594-36-5], 1987	[612-12-4], 2946
[541-42-4], 1262	[557-40-4], 2431	[591-50-4], 2249	[594-39-8], 2028	[612-23-7], 2713
[541-59-3], 1418	[557-89-1], 0884	[591-51-5], 2259	[594-56-9], 2855	[612-24-8], 2662
[542-15-4], 2379	[557-98-2], 1157	[591-76-4], 2874	[594-70-7], 1660	[612-25-9], 2769
[542-55-2], 1971	[558-13-4], 0315	[591-78-6], 2488	[594-73-0], 0584	[612-31-7], 2192
[542-63-2], 1675	[558-17-8], 1649	[591-87-7], 1912	[594-84-3], 2469	[613-94-5], 2798
[542-83-6], 0588	[560-21-4], 3062	[591-89-9], 1797	[595-86-8], 0775	[614-21-1], 2936
[542-88-1], 0793	[563-43-9], 0843	[591-93-5], 1897	[595-89-1], 3850	[614-45-9], 3401
[542-92-7], 1857	[563-45-1], 1944	[591-95-7], 1895	[597-64-8], 3097	[615-05-4], 2826
[543-53-3], 1873	[563-46-2], 1942	[592-02-9], 1677	[598-23-2], 1894	[615-42-9], 2172
[543-59-9], 1988	[563-47-3], 1553	[592-04-1], 0976	[598-25-4], 1892	[615-53-2], 1597
[543-67-9], 1265	[563-52-0], 1552	[592-09-2], 0999	[598-26-5], 1518	[615-57-6], 2226
[543-80-6], 1486	[563-80-4], 1959	[592-41-6], 2459	[598-49-2], 0789	[615-66-7], 2349
[544-13-8], 1870	[564-02-3], 3060	[592-42-7], 2411	[598-55-0], 0868	[615-88-9], 1841
[544-16-1], 1658	[565-59-3], 2871	[592-43-8], 2460	[598-56-1], 1727	[616-27-3], 1556
[544-25-2], 2790	[565-61-7], 2494	[592-45-0], 2410	[598-57-2], 0478	[616-38-6], 1233
[544-76-3], 3726	[565-69-5], 2493	[592-48-3], 2409	[598-58-3], 0457	[616-42-2], 0927

[616-47-7], 1499	[626-56-2], 2524	[641-16-7], 2091	[693-02-7], 2414	[821-08-9], 2291
[616-74-0], 2198	[626-58-4], 2525	[646-04-8], 1947	[693-04-9], 1641	[821-09-0], 1964
[617-50-5], 2863	[626-67-5], 2522	[646-06-0], 1228	[693-21-0], 1599	[821-10-3], 1428
[617-75-4], 2557	[627-13-4], 1267	[650-42-0], 2294	[693-65-2], 3369	[821-13-6], 1188
[617-77-6], 2561	[627-19-0], 1898	[650-51-1], 0608	[693-85-6], 0342	[822-06-0], 3008
[617-85-6], 2574	[627-21-4], 1899	[650-52-2], 0597	[694-05-3], 1930	[822-35-3], 1483
[617-86-7], 2585b	[627-26-9], 1464	[659-70-1], 3361	[694-17-7], 0632	[822-85-5], 2392
[617-88-9], 1842	[627-34-9], 1350	[659-85-8], 2289	[694-59-7], 1849	[823-87-0], 2181
[619-18-1], 2276	[627-41-8], 1523	[659-86-9], 1105	[694-94-0], 2811	[824-72-6], 2245
[619-25-0], 2770	[627-49-6], 1734	[661-54-1], 1066	[695-12-5], 3026	[824-78-2], 2183
[619-37-8], 2771	[627-54-3], 1717	[662-50-0], 1392	[695-64-7], 2419	[824-94-2], 2962
[619-96-7], 2275	[628-16-0], 2292	[663-25-2], 1368	[696-23-1], 1475	[828-26-2], 3197
[622-22-7], 2401	[628-20-6], 1487	[667-29-8], 0626	[696-26-4], 2991	[829-85-6], 3508
[622-37-7], 2271	[628-28-4], 2010	[667-49-2], 1049	[696-33-3], 2253	[841-76-9], 3749
[622-79-7], 2779	[628-32-0], 2013	[670-54-2], 2629	[697-18-7], 0635	[865-33-8], 0450
[623-42-7], 1974	[628-33-1], 1909	[672-66-2], 3003	[704-13-2], 2690	[865-37-2], 0936
[623-43-8], 1914	[628-37-5], 1699	[674-81-7], 0481	[728-40-5], 3666	[865-44-1], 4139
[623-70-1], 2436	[628-41-1], 2362	[674-82-8], 1441	[760-19-0], 1670	[865-47-4], 1650
[623-73-4], 1503	[628-52-4], 1493	[675-20-7], 1934	[762-04-9], 1733	[866-87-5], 3016
[623-81-4], 1709	[628-63-7], 2864	[676-54-0], 0877	[762-21-0], 2987	[868-14-4], 1462
[624-24-8], 2507	[628-71-7], 2838	[676-59-5], 0948	[762-42-5], 2340	[870-23-5], 1239
[624-31-7], 2750	[628-81-9], 2540	[676-75-5], 0886	[763-29-1], 2462	[870-24-6], 0940
[624-61-3], 0582	[628-85-3], 2537	[676-83-5], 0438	[764-05-6], 0544	[870-30-4], 0327
[624-64-6], 1578	[628-86-4], 0978	[676-85-7], 0435	[765-30-0], 1256	[871-27-2], 1719
[624-65-7], 1092	[628-87-5], 1474	[677-21-4], 1099	[765-31-1], 0808	[871-31-8], 0872
[624-67-9], 1085	[628-91-1], 2551	[680-31-9], 4718	[765-43-5], 1907	[872-36-6], 1087
[624-74-8], 0985	[628-92-2], 2837	[681-84-5], 1764	[765-87-7], 2385a	[873-51-8], 2222
[624-76-0], 0860	[629-05-0], 3022	[683-45-4], 1199	[766-06-3], 2832	[877-66-7], 2827
[624-79-3], 1178	[629-13-0], 0819	[683-60-3], 1270	[766-09-6], 2869	[881-68-5], 3290
[624-83-9], 0761	[629-14-1], 2546	[684-93-5], 0875	[766-42-1], 1397	[881-99-2], 2891
[624-84-0], 0473	[629-20-9], 2943	[687-78-5], 2096	[766-91-6], 2725	[890-28-4], 1877
[624-85-1], 0851	[629-59-4], 3671	[687-80-9], 2124	[768-34-3], 1949	[890-60-1], 2712
[624-89-5], 1288	[630-08-0], 0555	[687-81-0], 2125	[771-29-9], 3312	[917-54-4], 0452
[624-90-8], 0459	[630-18-2], 1929	[688-61-9], 3177	[771-63-1], 2082	[917-57-7], 0757
[624-91-9], 0455	[630-72-8], 1009	[689-93-0], 1293	[775-09-9], 4039	[917-58-8], 0861
[624-92-0], 0934	[634-67-3], 2163	[689-97-4], 1423	[777-52-6], 2158	[917-64-6], 0446
[625-48-9], 0871	[637-12-7], 3914	[689-98-5], 0896	[778-28-9], 3409	[917-92-0], 2408
[625-49-0], 0409	[637-61-6], 2140	[689-99-6], 0623	[811-49-4], 0862	[918-37-6], 1014
[625-54-7], 2012	[637-62-7], 2263	[690-02-8], 0923	[811-62-1], 0899	[918-54-7], 0776
[625-55-8], 1619	[637-62-7], 2264	[691-35-0], 1983	[811-97-2], 0706	[918-99-0], 1875
[625-58-1], 0870	[637-70-7], 2159	[691-37-2], 2463	[812-12-4], 0354	[920-37-6], 1074
[625-76-3], 0410	[637-78-5], 2504	[691-38-3], 2464	[814-49-3], 1681	[920-46-7], 1457
[625-86-5], 2382	[637-92-3], 2541	[692-42-2], 1720	[814-68-6], 1093	[921-22-2], 0676
[625-98-9], 2135	[637-97-8], 1990	[692-45-5], 1149	[814-78-8], 1911	[922-67-8], 1442
[626-23-3], 3084	[638-11-9], 2862	[692-56-8], 2616	[816-43-3], 1686	[922-89-4], 2426
[626-55-1], 1827	[638-21-1], 2360	[692-74-0], 0625	[817-87-8], 1193	[923-34-2], 2564

[924-42-5], 1571	[1066-48-4], 0333	[1189-85-1], 3066	[1309-37-1], 4403	[1330-20-7], 2969
[925-90-6], 0847	[1067-20-5], 3199	[1189-93-1], 2617	[1309-48-4], 4695	[1330-43-4], 0185
[926-56-7], 2417	[1068-57-1], 0912	[1191-15-7], 3082	[1309-64-4], 4851	[1332-67-8], 3942
[926-63-6], 2031	[1069-12-1], 2516	[1191-47-5], 3068	[1310-58-3], 4428	[1332-69-9], 3942
[926-65-8], 1957	[1069-19-5], 3202	[1191-91-7], 1517	[1310-73-2], 4445	[1332-76-9], 4186
[926-82-9], 3201	[1070-19-5], 1936	[1191-96-4], 1941	[1312-73-8], 4670	[1333-74-0], 4453
[927-80-0], 1521	[1070-75-3], 0992	[1192-27-4], 1832	[1312-81-8], 4679	[1333-82-0], 4242
[928-45-0], 1661	[1071-98-3], 1801	[1192-28-5], 1932	[1313-13-9], 4705	[1335-31-5], 0983
[928-49-4], 2415	[1072-67-9], 1501	[1193-54-0], 1378	[1313-27-5], 4717	[1336-21-6], 4544
[928-55-2], 1956	[1073-23-0], 2819	[1209-60-0], 4834	[1313-59-3], 4802	[1337-81-1], 2759
[928-84-1], 1382	[1073-91-2], 2517	[1270-21-9], 2722	[1313-60-6], 4803	[1344-08-7], 4813
[929-25-9], 2372	[1074-12-0], 2922	[1271-24-5], 3273	[1313-82-2], 4811	[1344-28-1], 0087
[929-57-7], 3006	[1074-98-2], 2317	[1271-27-8], 3278	[1313-96-8], 4818	[1344-43-0], 4704
[930-22-3], 1520	[1075-17-8], 3317	[1271-28-9], 3287	[1313-99-1], 4821	[1344-48-5], 4607
[930-28-9], 1923	[1076-44-4], 2059	[1271-29-9], 3295	[1314-06-3], 4823	[1344-57-6], 4843
[930-82-5], 2422	[1109-15-5], 3738	[1271-54-1], 3511	[1314-08-5], 4825	[1344-59-8], 4871
[932-64-9], 0716	[1111-63-3], 0671	[1271-86-9], 3613	[1314-13-2], 4830	[1345-04-6], 4906
[932-90-1], 2760	[1111-64-4], 0673	[1284-72-6], 3277	[1314-15-4], 4836	[1345-07-9], 0234
[935-14-8], 3241	[1112-04-5], 0361	[1293-87-4], 3479	[1314-18-7], 4841	[1345-25-1], 4398
[937-14-4], 2676	[1113-78-6], 3570	[1294-74-2], 3294	[1314-22-3], 4845	[1423-11-6], 2057
[938-10-3], 2274	[1115-12-4], 1341	[1298-53-9], 3683	[1314-24-5], 4847	[1426-40-0], 0646
[938-71-6], 2679	[1115-99-7], 2563	[1299-86-1], 1031	[1314-32-5], 4854	[1445-79-0], 1305
[938-81-8], 2949	[1116-01-4], 1343	[1300-32-9], 1818	[1314-34-7], 4855	[1447-06-1], 2836
[940-71-6], 4189	[1116-39-8], 3569	[1301-96-8], 0025	[1314-35-8], 4856	[1450-14-2], 2608
[940-93-2], 2992	[1116-61-6], 3217	[1303-28-2], 0106	[1314-41-6], 4861	[1479-48-7], 0639
[941-55-9], 2781	[1117-94-8], 0619	[1303-33-9], 0108	[1314-56-3], 4864	[1484-80-6], 2870
[945-51-7], 3493	[1117-99-3], 0698	[1303-58-8], 0115	[1314-61-0], 4865	[1484-88-4], 3839
[946-32-7], 2778	[1118-58-7], 2416	[1303-61-3], 0116	[1314-62-1], 4866	[1493-01-2], 0373
[950-32-3], 3539	[1119-22-8], 1547	[1303-86-2], 0170	[1314-80-3], 4875	[1493-13-6], 0375
[952-52-3], 2747	[1119-69-3], 2311	[1304-28-5], 0215	[1314-85-7], 4876	[1493-23-8], 2166
[959-26-2], 3526	[1119-90-0], 3080	[1304-29-6], 0216	[1314-85-8], 4877	[1498-47-1], 1169
[959-52-4], 3531	[1120-46-3], 3895	[1304-56-9], 0225	[1314-86-9], 4880	[1510-31-2], 0398
[984-47-5], 2268	[1120-64-5], 1569	[1304-76-3], 0233	[1314-95-0], 4900	[1516-70-7], 0465
[992-94-9], 0510	[1120-97-4], 1975	[1305-62-0], 3928	[1314-96-1], 4901	[1517-05-1], 0873
[993-74-8], 2597	[1121-14-8], 1139	[1305-78-8], 3937	[1315-01-1], 4902	[1522-92-5], 1922
[994-99-0], 3674	[1121-53-5], 2787	[1305-79-9], 3938	[1317-33-5], 4719	[1528-27-4], 3781
[998-30-1], 2584	[1122-17-4], 1359	[1305-99-3], 3947	[1317-36-8], 4824	[1557-57-9], 1005
[998-40-3], 3575	[1124-31-8], 2174	[1306-19-0], 3958	[1317-37-9], 4400	[1570-64-5], 2741
[999-97-3], 2610	[1124-33-0], 1833	[1306-24-7], 3959	[1317-38-0], 4281	[1575-37-7], 2345
[1001-62-3], 0931	[1125-99-1], 3350	[1307-80-8], 4187	[1317-39-1], 4288	[1585-74-6], 0895
[1002-36-4], 2841	[1149-23-1], 3615	[1307-96-6], 4217	[1317-40-4], 4285	[1586-92-1], 2554
[1003-03-8], 1991	[1184-53-8], 0440	[1308-04-9], 4221	[1317-42-6], 4218	[1587-41-3], 0328
[1004-22-4], 2911	[1184-58-3], 0883	[1308-09-4], 4223	[1317-61-9], 4405	[1590-87-0], 4569
[1012-17-5], 3297	[1184-78-7], 1313	[1308-38-9], 4251	[1321-12-6], 2762	[1592-23-0], 3896
[1025-15-6], 3530	[1186-53-4], 3212	[1308-80-1], 4289	[1322-36-7], 3567	[1600-31-3], 2092
[1066-26-8], 0684	[1189-71-5], 0324	[1309-36-0], 4401	[1327-53-3], 0105	[1605-51-2], 1735

[1605-58-9], 2036	[1873-77-4], 3231	[2386-62-1], 0842	[2809-21-4], 0956	[3221-61-2], 3206
[1607-30-3], 3354	[1884-64-6], 0540	[2397-67-3], 3214	[2809-67-8], 3023	[3236-56-4], 3194
[1609-47-8], 2444	[1886-68-6], 3656	[2400-59-1], 3554	[2809-69-0], 2293	[3242-56-6], 2375
[1618-08-2], 1344	[1886-69-7], 3458	[2402-95-1], 1829	[2829-31-4], 3609	[3248-28-0], 2442
[1623-99-0], 2287	[1890-04-6], 1860	[2402-97-3], 1828	[2845-00-3], 1173	[3251-23-8], 4279
[1624-19-4], 3184	[1931-62-0], 3015	[2404-52-6], 4340	[2851-55-0], 0990	[3264-86-6], 2626
[1630-94-0], 1940	[1941-24-8], 0506	[2406-51-1], 0807	[2868-37-3], 1917	[3272-96-6], 0849
[1631-82-9], 2810	[1941-79-3], 3189	[2407-94-5], 3553	[2875-36-7], 3053	[3277-26-7], 1783
[1634-04-4], 2011	[1942-45-6], 3025	[2417-93-8], 1253	[2881-62-1], 1022	[3279-95-6], 0944
[1637-31-6], 1389	[1944-83-8], 3332	[2424-09-1], 1758	[2893-78-9], 1037	[3282-30-2], 1925
[1640-89-7], 2850	[1956-39-4], 2927	[2425-74-3], 1994	[2899-02-7], 3596	[3296-62-3], 0694
[1645-75-6], 1067	[1972-28-7], 2425	[2425-79-8], 3353	[2916-31-6], 1969	[3312-92-6], 1024
[1653-57-2], 3364	[1984-04-9], 3390	[2428-04-8], 1043	[2917-26-2], 3727	[3315-16-0], 0300
[1656-16-2], 2073	[2003-31-8], 1036	[2446-84-6], 1528	[2918-51-8], 2682	[3315-89-7], 0972
[1656-44-6], 2100	[2028-76-4], 2151	[2506-80-1], 2026	[2923-28-6], 0444	[3319-45-7], 3755
[1663-35-0], 1973	[2032-04-4], 1135	[2524-02-9], 3980	[2925-21-5], 0657	[3327-66-1], 0586
[1666-13-3], 3501	[2036-15-9], 2552	[2536-18-7], 1142	[2938-46-7], 0423	[3329-56-4], 2428
[1675-54-3], 3825	[2041-76-1], 2844	[2551-62-4], 4374	[2938-73-0], 0844	[3333-52-6], 3007
[1679-09-0], 2023	[2045-70-7], 1421	[2554-06-5], 3560	[2944-05-0], 0559	[3356-63-6], 1170
[1686-88-3], 1472	[2083-91-2], 2043	[2567-83-1], 3088	[2944-66-3], 2065	[3385-78-2], 1307
[1711-42-8], 2925	[2092-17-3], 0576	[2581-34-2], 2766	[2958-89-6], 1586	[3400-09-7], 4063
[1712-64-7], 1266	[2152-13-7], 3267	[2589-01-7], 3533	[2980-33-8], 1822	[3437-84-1], 3032
[1717-00-6], 0738	[2154-71-4], 0638	[2592-85-0], 2178	[2997-92-4], 3089	[3444-13-1], 0982
[1718-18-9], 0642	[2155-71-7], 3715	[2592-95-2], 2272	[2999-74-8], 0908	[3445-52-1], 1476
[1719-53-5], 1683	[2156-66-3], 0915	[2611-42-9], 1023	[3003-15-4], 1321	[3454-11-3], 0756
[1730-69-4], 3065	[2156-71-0], 1948	[2612-33-1], 1160	[3009-34-5], 2260	[3463-17-0], 3604
[1739-53-3], 0845	[2167-23-9], 3565	[2614-76-8], 1287	[3011-34-5], 2691	[3468-63-1], 3702
[1746-09-4], 2591	[2169-38-2], 1740	[2629-78-9], 1361	[3013-38-5], 2672	[3469-17-8], 3638
[1750-42-1], 1136	[2175-91-9], 2967	[2643-84-5], 2960	[3015-98-3], 3219	[3470-17-5], 2631
[1762-95-4], 0479	[2198-20-1], 3351	[2644-70-4], 4007	[3017-85-4], 0882	[3495-54-3], 0063
[1779-25-5], 3064	[2200-13-7], 3876	[2651-85-6], 2572	[3021-39-4], 0977	[3504-13-0], 1542
[1781-66-4], 1884	[2203-57-8], 1060	[2675-89-0], 1585	[3031-73-0], 0486	[3514-82-7], 3481
[1789-58-8], 0903	[2216-12-8], 3472	[2691-41-0], 1605	[3031-74-1], 0925	[3522-94-9], 3210
[1792-40-1], 2940	[2216-30-0], 3200	[2696-92-6], 4023	[3031-75-2], 1283	[3525-44-8], 1667
[1794-84-9], 0396	[2216-33-3], 3207	[2697-42-9], 0952	[3034-94-4], 2905	[3531-19-9], 2153
[1795-27-3], 3191	[2216-34-4], 3208	[2698-41-1], 3333	[3052-45-7], 1177	[3536-96-7], 0732
[1809-19-4], 3086	[2217-55-2], 3449	[2698-69-3], 3266	[3054-95-3], 2858	[3558-60-9], 3164
[1809-53-6], 3020	[2235-08-7], 1416	[2699-12-9], 3556	[3058-38-0], 2324	[3570-93-2], 2400
[1817-47-6], 3157	[2245-68-3], 1685	[2712-78-9], 3239	[3071-32-7], 2985	[3585-32-8], 0945
[1817-73-8], 2132	[2252-95-1], 1363	[2713-09-9], 0662	[3074-75-7], 3205	[3607-48-5], 0524
[1838-59-1], 1524	[2278-22-0], 0766	[2735-04-8], 2998	[3074-77-9], 3204	[3618-05-1], 4469
[1840-42-2], 0340	[2294-47-5], 2208	[2738-18-3], 3190	[3086-29-1], 1303	[3676-91-3], 1765
[1850-14-2], 2022	[2306-67-0], 0888	[2749-79-3], 2328	[3159-98-6], 3542	[3676-97-9], 1766
[1854-19-9], 3081	[2370-88-9], 1790	[2783-96-2], 0681	[3161-99-7], 2074	[3686-43-9], 1567
[1871-52-9], 2966	[2372-21-6], 3051	[2786-01-8], 2657	[3179-56-4], 3033	[3692-81-7], 3571
[1872-00-0], 1722	[2378-02-1], 1380	[2802-70-2], 1059	[3188-13-4], 1246	[3698-54-2], 2122

[3698-83-7], 2077	[4415-23-0], 1007	[4897-21-6], 1453	[5789-17-3], 1141	[6824-18-6], 1444
[3706-77-2], 1094	[4417-80-5], 2433	[4904-55-6], 3667	[5796-89-4], 1187	[6832-13-9], 0710
[3710-30-3], 3021	[4418-61-5], 0461	[4911-55-1], 1825	[5796-89-4], 1574	[6832-16-2], 1138
[3751-44-8], 1576	[4419-11-8], 3668	[4922-98-9], 2952	[5797-06-8], 1837	[6834-92-0], 4805
[3768-60-3], 1757	[4437-18-7], 1840	[4972-29-6], 2234	[5809-59-6], 1467	[6865-68-5], 2622
[3802-95-7], 0347	[4452-58-8], 0554	[4981-48-0], 1935	[5813-49-0], 0762	[6866-10-0], 3147
[3811-04-9], 4017	[4457-90-3], 2053	[4984-82-1], 1855	[5813-64-9], 2030	[6890-38-6], 2378
[3823-94-7], 1101	[4457-91-4], 3418	[5071-96-5], 2997	[5813-77-4], 1535	[6918-51-0], 0930
[3851-22-7], 1704	[4460-46-2], 2673	[5106-46-7], 2514	[5888-61-9], 3822	[6921-27-3], 2327
[3855-45-6], 0348	[4461-41-0], 1551	[5120-07-0], 3672	[5960-88-3], 0853	[6925-01-5], 2189
[3880-04-4], 0906	[4461-48-7], 2465	[5128-28-9], 2089	[5989-27-5], 3337	[6926-39-2], 2004
[3884-87-0], 0305	[4472-06-4], 0386	[5131-60-2], 2348	[6044-68-4], 1968	[6926-40-5], 2002
[3917-15-5], 1904	[4474-17-3], 0600	[5131-88-4], 0630	[6066-82-6], 1469	[6928-74-1], 1498
[3946-86-9], 1751	[4504-27-2], 1190	[5158-50-9], 0887	[6069-42-7], 1226	[6996-92-5], 2335
[3958-60-9], 2710	[4531-35-5], 2895	[5162-44-7], 1549	[6088-91-1], 1381	[7119-27-9], 1454
[3982-91-0], 4154	[4538-37-8], 2376	[5164-35-2], 1856	[6109-04-2], 3631	[7154-79-2], 3211
[3999-10-8], 2727	[4543-33-3], 3442	[5194-50-3], 2412	[6130-87-6], 1759	[7176-89-8], 3149
[4000-16-2], 0517	[4547-68-6], 2152	[5194-51-4], 2413	[6147-53-1], 1780	[7182-87-8], 0374
[4023-65-8], 2342	[4547-69-7], 2210	[5256-74-6], 2828	[6180-21-8], 0599	[7203-21-6], 3786
[4039-32-1], 2594	[4549-46-6], 1690	[5274-48-6], 2217	[6232-88-8], 2926	[7226-23-5], 2475
[4100-38-3], 3309	[4572-12-7], 3788	[5314-83-0], 1673	[6269-50-7], 2220	[7227-91-0], 2996
[4104-85-2], 0515	[4591-46-2], 2123	[5314-85-2], 1335	[6303-21-5], 4503	[7245-18-3], 1317
[4109-96-0], 4066	[4606-07-9], 2437	[5329-14-6], 4499	[6361-21-3], 2649	[7279-42-0], 2179
[4112-22-5], 1494	[4635-46-5], 4788	[5332-73-0], 1730	[6361-22-4], 2650	[7289-92-1], 2598
[4113-57-9], 0655	[4648-54-8], 1314	[5332-96-7], 3142	[6381-06-2], 3502	[7318-34-5], 3270
[4150-34-9], 3758	[4656-04-6], 2593	[5345-54-0], 2795	[6407-56-3], 3701	[7332-00-5], 1666
[4154-69-2], 1678	[4676-82-8], 1624	[5366-84-7], 3240	[6410-13-5], 3697	[7360-53-4], 1089
[4170-30-3], 1516	[4682-03-5], 2087	[5388-62-5], 2154	[6411-21-8], 3379	[7388-28-5], 0880
[4185-47-1], 1602	[4702-38-9], 0407	[5390-07-8], 2236	[6443-91-0], 1146	[7400-08-0], 3136
[4209-91-0], 3073	[4711-74-4], 2606	[5392-40-5], 3346	[6470-09-3], 1342	[7416-48-0], 1626
[4212-43-5], 1236	[4711-95-9], 2866	[5425-78-5], 1245	[6476-36-4], 3220	[7429-90-5], 0048
[4222-21-3], 0733	[4712-19-0], 3468	[5428-37-5], 3259	[6477-64-1], 3436	[7429-96-5], 4700
[4222-26-8], 2928	[4733-52-2], 2900	[5434-72-0], 3902	[6484-52-2], 4522	[7436-07-9], 3140
[4222-27-9], 0734	[4744-10-9], 2017	[5470-11-1], 4002	[6488-00-2], 2976	[7439-88-5], 4643
[4230-63-1], 3488	[4747-90-4], 1079	[5492-69-3], 3090	[6522-40-3], 2369	[7439-89-6], 4388
[4254-22-2], 0729	[4749-28-4], 1183	[5500-21-0], 1463	[6540-76-7], 2394	[7439-91-0], 4677
[4264-83-9], 2185	[4755-77-5], 1456	[5518-62-7], 0957	[6569-51-3], 0176	[7439-92-1], 4882
[4279-76-9], 2920	[4756-66-5], 1377	[5573-38-3], 3229	[6592-86-1], 2129	[7439-93-2], 4680
[4312-28-1], 3848	[4774-73-6], 0918	[5588-74-9], 2745	[6610-25-9], 3809	[7439-95-4], 4690
[4328-17-0], 2120	[4784-77-4], 1548	[5593-70-4], 3730	[6628-86-0], 2652	[7439-97-6], 4600
[4331-98-0], 0998	[4806-61-5], 2458	[5610-59-3], 0301	[6630-99-5], 0413	[7439-98-7], 4712
[4343-68-4], 0541	[4845-05-0], 2435	[5613-68-3], 2405	[6659-62-7], 1186	[7440-00-8], 4819
[4377-73-5], 2310	[4873-09-0], 3315	[5613-69-4], 3221	[6713-82-2], 3894	[7440-02-0], 4820
[4388-03-8], 1593	[4884-18-8], 3693	[5637-83-2], 1348	[6728-26-3], 2432	[7440-03-1], 3948
[4394-93-8], 0362	[4887-24-5], 2390	[5653-21-4], 0809	[6786-32-9], 2689	[7440-03-1], 4817
[4403-63-8], 3581	[4888-97-5], 3010	[5702-61-4], 3500	[6798-76-1], 3903	[7440-04-2], 4873

[7440-05-3], 4885	[7446-08-4], 4838	[7681-57-4], 4808	[7773-06-0], 4556	[7783-75-7], 4362
[7440-06-4], 4887	[7446-09-5], 4837	[7681-82-5], 4623	[7774-29-0], 4602	[7783-77-9], 4365
[7440-07-5], 4888	[7446-11-9], 4849	[7693-26-7], 4427	[7775-14-6], 4807	[7783-81-5], 4375
[7440-09-7], 4645	[7446-14-2], 4860	[7693-27-8], 4463	[7775-27-1], 4809	[7783-82-6], 4376
[7440-10-0], 4886	[7446-70-0], 0062	[7697-37-2], 4436	[7775-41-9], 0013	[7783-83-7], 4393
[7440-15-5], 4890	[7447-40-7], 4015	[7704-34-9], 4897	[7778-18-9], 3939	[7783-84-8], 4394
[7440-16-6], 4892	[7529-22-8], 1997	[7704-98-5], 4490	[7778-50-9], 4248	[7783-86-0], 4395
[7440-17-7], 4889	[7544-40-3], 3027	[7704-99-6], 4493	[7778-54-3], 3924	[7783-89-3], 0007
[7440-18-8], 4894	[7550-45-0], 4176	[7705-07-9], 4158	[7778-74-7], 4018	[7783-90-6], 0008
[7440-21-3], 4909	[7553-56-2], 4625	[7705-08-0], 4133	[7779-88-6], 4751	[7783-91-7], 0010
[7440-22-4], 0001	[7568-37-8], 0431	[7705-12-6], 1393	[7782-40-3], 0298	[7783-92-8], 0011
[7440-23-5], 4796	[7570-25-4], 0770	[7713-79-3], 3122	[7782-41-4], 4310	[7783-93-9], 0012
[7440-25-7], 4913	[7572-29-4], 0602	[7718-98-1], 4159	[7782-42-5], 0298	[7783-95-1], 0014
[7440-25-7], 4914	[7575-57-7], 3155	[7719-09-7], 4096	[7782-44-7], 4831	[7783-97-3], 0020
[7440-26-8], 4915	[7601-54-9], 4814	[7719-12-2], 4153	[7782-44-7], 4832	[7783-98-4], 0021
[7440-28-0], 4922	[7601-87-8], 1512	[7720-78-7], 4399	[7782-49-2], 4908	[7784-04-5], 0031
[7440-29-1], 4917	[7601-89-0], 4040	[7720-83-4], 4638	[7782-50-3], 4047	[7784-21-6], 0070
[7440-31-5], 4911	[7601-90-3], 3998	[7721-01-9], 4185	[7782-63-9], 3175	[7784-23-8], 0079
[7440-31-5], 4912	[7616-83-3], 4078	[7722-64-7], 4647	[7782-65-2], 4417	[7784-34-1], 0094
[7440-32-6], 4919	[7616-94-6], 3974	[7722-84-1], 4477	[7782-68-5], 4424	[7784-36-3], 0095
[7440-33-7], 4925	[7631-86-9], 4839	[7722-86-3], 4481	[7782-77-6], 4435	[7784-42-1], 0100
[7440-36-0], 4907	[7631-99-4], 4721	[7723-14-0], 4874	[7782-78-7], 4438	[7786-30-3], 4081
[7440-37-1], 0091	[7632-00-0], 4720	[7726-95-6], 0261	[7782-79-8], 4441	[7787-47-5], 0221
[7440-38-2], 0092	[7632-04-4], 0155	[7727-15-3], 0060	[7782-87-8], 4459	[7787-49-7], 0223
[7440-39-3], 0200	[7632-51-1], 4177	[7727-18-6], 4151	[7782-89-0], 4462	[7787-52-2], 0224
[7440-41-7], 0220	[7637-07-2], 0129	[7727-21-1], 4668	[7782-92-5], 4465	[7787-62-4], 0227
[7440-42-8], 0119	[7646-69-7], 4444	[7727-37-9], 4734	[7782-94-7], 4471	[7787-70-4], 0265
[7440-43-9], 3949	[7646-78-8], 4174	[7727-43-7], 0217	[7783-06-4], 4483	[7787-71-5], 0241
[7440-44-0], 0298	[7646-79-9], 4048	[7727-54-0], 4576	[7783-07-5], 4486	[7789-09-5], 4246
[7440-45-1], 3961	[7646-85-7], 4120	[7731-34-2], 3406	[7783-09-7], 4488	[7789-18-6], 4261
[7440-46-2], 4254	[7647-01-0], 3993	[7732-18-5], 4475	[7783-17-7], 4517	[7789-20-0], 4291
[7440-47-3], 4222	[7647-14-5], 4036	[7738-94-5], 4229	[7783-18-8], 4573	[7789-25-5], 4302
[7440-48-4], 4199	[7647-18-9], 4184	[7740-53-1], 4292	[7783-20-2], 4574	[7789-26-6], 4304
[7440-50-8], 4267	[7651-91-4], 0437	[7757-79-1], 4650	[7783-26-8], 4579	[7789-30-2], 0243
[7440-55-3], 4406	[7659-31-6], 0568	[7758-01-2], 0255	[7783-29-1], 4584	[7789-31-3], 0248
[7440-56-4], 4412	[7664-38-2], 4505	[7758-05-6], 4619	[7783-41-7], 4317	[7789-33-5], 0254
[7440-57-5], 0110	[7664-39-3], 4294	[7758-09-0], 4649	[7783-42-8], 4318	[7789-38-0], 0257
[7440-58-6], 4599	[7664-41-7], 4497	[7758-19-2], 4038	[7783-44-0], 4320	[7789-41-5], 0262
[7440-61-1], 4923	[7664-93-9], 4479	[7758-89-6], 4056	[7783-46-2], 4329	[7789-43-7], 0263
[7440-63-3], 4926	[7673-09-8], 1095	[7758-94-3], 4061	[7783-47-3], 4331	[7789-46-0], 0266
[7440-66-6], 4927	[7677-24-9], 1665	[7758-95-4], 4112	[7783-53-1], 4335	[7789-47-1], 0269
[7440-67-7], 4928	[7681-38-1], 4446	[7758-97-6], 4243	[7783-54-2], 4336	[7789-51-7], 0275
[7440-68-2], 4924	[7681-38-1], 4806	[7761-88-8], 0022	[7783-55-3], 4339	[7789-57-3], 0287
[7440-69-9], 0226	[7681-52-9], 4037	[7772-98-7], 4804	[7783-60-0], 4350	[7789-60-8], 0293
[7440-70-2], 3922	[7681-53-0], 4473	[7772-99-8], 4116	[7783-61-1], 4352	[7789-65-3], 0295
[7440-74-6], 4640	[7681-55-2], 4624	[7773-01-5], 4085	[7783-66-6], 4355	[7789-78-8], 3927

[7789-79-9], 3931	[10026-07-0], 4175	[10158-43-7], 3298	[10544-73-7], 4746	[12033-54-4], 4726
[7790-21-8], 4620	[10026-11-6], 4178	[10192-29-7], 4003	[10545-99-0], 4113	[12033-59-9], 4730
[7790-69-4], 4684	[10026-13-8], 4183	[10192-30-0], 4545	[10557-85-4], 1868	[12033-67-9], 4731
[7790-89-8], 3971	[10026-18-3], 4200	[10204-97-2], 4744	[10567-69-8], 0211	[12033-88-6], 4771
[7790-91-2], 3981	[10026-22-9], 4215	[10210-68-1], 2881	[10578-16-2], 4314	[12033-90-8], 4774
[7790-92-3], 3995	[10027-74-4], 1707	[10218-83-4], 0430	[10580-52-6], 4118	[12034-12-7], 4799
[7790-93-4], 3996	[10028-15-6], 4846	[10229-09-1], 1508	[10588-01-9], 4250	[12035-36-8], 4822
[7790-94-5], 3997	[10031-26-2], 0286	[10230-68-9], 1184	[10599-90-3], 4000	[12036-02-1], 4833
[7790-98-9], 4004	[10031-27-3], 0296	[10233-03-1], 4082	[11079-26-8], 3708	[12036-04-3], 4835
[7790-99-0], 4013	[10031-43-3], 4279	[10241-03-9], 4160	[11144-29-9], 0002	[12036-22-5], 4844
[7791-03-9], 4021	[10034-81-8], 4084	[10241-04-4], 4122	[11713-32-0], 0631	[12038-67-4], 4891
[7791-08-4], 4041	[10034-85-2], 4423	[10241-05-1], 4180	[11802-21-2], 3469	[12039-13-3], 4904
[7791-21-1], 4095	[10035-10-6], 0247	[10248-58-5], 4867	[12001-65-9], 2471	[12039-14-4], 4905
[7791-23-3], 4097	[10036-47-2], 4345	[10256-92-5], 0352	[12001-89-7], 3767	[12042-38-5], 0053
[7791-25-5], 4099	[10038-98-9], 4162	[10272-07-8], 2999	[12003-23-5], 0084	[12043-56-0], 0049
[7791-27-7], 4103	[10039-54-0], 4575	[10294-33-4], 0122	[12003-69-9], 0051	[12044-52-9], 0107
[7796-16-9], 0659	[10043-11-5], 0152	[10294-34-5], 0127	[12003-96-2], 0056	[12045-01-1], 0128
[7803-49-8], 4498	[10043-35-3], 0144	[10294-46-9], 4057	[12007-25-9], 0168	[12047-79-9], 0219
[7803-51-2], 4508	[10043-52-4], 3923	[10294-47-0], 4105	[12007-33-9], 0172	[12051-08-0], 4028
[7803-52-3], 4510	[10043-84-2], 4519	[10294-48-1], 4107	[12007-46-4], 0177	[12053-67-7], 4263
[7803-62-5], 4539	[10043-92-2], 4893	[10294-58-3], 4531	[12008-21-8], 0193	[12055-24-2], 4628
[7803-65-8], 4554	[10045-94-0], 4603	[10308-82-4], 0512	[12008-61-6], 0199	[12057-24-8], 4686
[7847-94-7], 4076	[10049-03-3], 3976	[10308-83-5], 0514	[12010-53-6], 0231	[12057-71-5], 4698
[7861-11-0], 4618	[10049-04-4], 4042	[10308-84-6], 0502	[12011-67-5], 0365	[12057-74-8], 4699
[7958-20-3], 2550	[10049-05-5], 4052	[10308-90-4], 1276	[12012-32-7], 0590	[12057-92-0], 4709
[8049-11-4], 0050	[10049-06-6], 4117	[10308-93-7], 3048	[12013-55-7], 3943	[12058-54-7], 4800
[9002-81-7], 0417	[10049-07-7], 4155	[10308-95-9], 3092	[12013-56-8], 3944	[12058-85-4], 4815
[9002-83-9], 0592	[10049-08-8], 4156	[10311-08-7], 3002	[12013-82-0], 3946	[12063-27-3], 4404
[9002-84-0], 0629	[10049-17-9], 4373	[10325-39-0], 0124	[12014-93-6], 3967	[12065-85-9], 4870
[9002-88-4], 0782	[10051-06-6], 0359	[10326-21-3], 4083	[12015-30-4], 4033	[12069-32-8], 0309
[9002-89-5], 0831	[10058-23-8], 4430	[10361-95-2], 4106	[12015-76-8], 4269	[12069-85-1], 0521
[9003-17-2], 1484	[10097-28-6], 4828	[10369-17-2], 2600	[12018-00-7], 4241	[12070-08-5], 0561
[9003-27-4], 1583	[10099-74-8], 4749	[10377-60-3], 4693	[12018-06-3], 4245	[12070-09-6], 0562
[9004-73-3], 0485	[10101-50-5], 4703	[10377-62-5], 4691	[12019-11-3], 4284	[12070-12-1], 0563
[9016-00-6], 0922	[10102-03-1], 4748	[10377-66-9], 4701	[12019-27-1], 4268	[12070-13-2], 0564
[9018-45-5], 3836	[10102-06-4], 4752	[10397-28-1], 2916	[12020-65-4], 4293	[12070-14-3], 0565
[10022-31-8], 0212	[10102-40-6], 4713	[10405-27-3], 4311	[12024-20-3], 4411	[12070-14-3], 1029
[10022-50-1], 4303	[10102-43-9], 4724	[10415-75-5], 4609	[12025-32-0], 4419	[12070-27-8], 0575
[10025-67-9], 4114	[10102-44-0], 4725	[10420-90-3], 2290	[12027-06-4], 4512	[12071-15-7], 0991
[10025-68-0], 4115	[10112-08-0], 0492	[10421-42-4], 4397	[12029-98-0], 4627	[12071-29-3], 1026
[10025-73-7], 4127	[10118-76-0], 3934	[10437-85-1], 3404	[12030-49-8], 4644	[12071-33-9], 1028
[10025-78-2], 4136	[10124-37-5], 3935	[10450-60-9], 4542	[12030-88-5], 4656	[12075-68-2], 2556
[10025-85-1], 4143	[10124-48-8], 3999	[10467-10-4], 0859	[12030-89-6], 4657	[12079-66-2], 2882
[10025-87-3], 4149	[10137-74-3], 3952	[10504-99-1], 3494	[12032-88-1], 4708	[12081-54-8], 3098
[10025-91-9], 4157	[10138-17-7], 2498	[10534-86-8], 4210	[12033-19-1], 4714	[12081-88-8], 3107
[10026-04-7], 4173	[10139-47-6], 4632	[10544-72-6], 4747	[12033-31-7], 4714	[12082-46-1], 3180

[12097-97-1], 3830	[12265-93-9], 4418	[13093-00-0], 4111	[13453-07-1], 0111	[13537-24-1], 4134
[12116-83-5], 3296	[12266-58-9], 2312	[13101-58-1], 0548	[13453-57-1], 4102	[13537-32-1], 4297
[12124-97-9], 0251	[12273-50-6], 4674	[13138-21-1], 0675	[13455-00-0], 4637	[13537-33-2], 4298
[12125-02-9], 4001	[12275-58-0], 2951	[13138-45-9], 4743	[13455-01-1], 4636	[13551-73-0], 1864
[12125-09-6], 4515	[12279-90-2], 0109	[13147-25-6], 1691	[13455-15-7], 4348	[13556-31-5], 3172
[12125-23-4], 4707	[12281-65-1], 0567	[13154-66-0], 1331	[13463-30-4], 4172	[13556-50-8], 2207
[12129-68-9], 3514	[12298-67-8], 4606	[13168-78-0], 2263	[13463-39-3], 1805	[13569-43-2], 0246
[12133-31-2], 3933	[12300-03-7], 4797	[13172-31-1], 0282	[13463-40-6], 1814	[13569-50-1], 3962
[12133-60-7], 3983	[12312-33-3], 3269	[13196-64-7], 3188	[13463-67-7], 4842	[13572-99-1], 4413
[12134-29-1], 4266	[12316-91-3], 4776	[13213-29-1], 3804	[13463-98-4], 3900	[13596-23-1], 4194
[12134-35-9], 4290	[12325-31-4], 4564	[13214-58-9], 0748	[13464-46-5], 0090	[13596-24-2], 4198
[12135-76-1], 4577	[12331-76-9], 4235	[13223-78-4], 1010	[13464-97-6], 4549	[13597-73-4], 4565
[12136-15-1], 4615	[12333-57-2], 4729	[13232-74-1], 2632	[13464-98-7], 4561	[13597-95-0], 0222
[12136-26-4], 4641	[12333-97-0], 4883	[13235-16-0], 3940	[13465-07-1], 4484	[13598-36-2], 4504
[12136-61-7], 4689	[12340-14-6], 4863	[13270-95-6], 2578	[13465-08-2], 4524	[13598-47-5], 4500
[12136-83-3], 4723	[12354-85-7], 3806	[13271-93-7], 2047	[13465-33-3], 0271	[13598-52-5], 4506
[12136-85-5], 4727	[12359-06-7], 4682	[13271-97-1], 3583	[13465-41-3], 4434	[13598-56-6], 4511
[12138-26-0], 4910	[12380-95-9], 3960	[13271-98-2], 3777	[13465-66-2], 4351	[13600-88-9], 4204
[12139-70-7], 4214	[12384-02-0], 4681	[13272-02-1], 3230	[13465-71-9], 4333	[13601-08-6], 4588
[12140-58-8], 4410	[12397-26-1], 4439	[13272-03-2], 3586	[13465-73-1], 0250	[13620-57-0], 3653
[12143-72-5], 4903	[12397-29-4], 4616	[13272-04-3], 3696	[13465-77-5], 4191	[13637-63-3], 3989
[12154-56-2], 3605	[12412-94-1], 4884	[13272-05-4], 3852	[13465-95-7], 0206	[13637-65-5], 3992
[12164-01-1], 4773	[12430-27-2], 0194	[13280-07-4], 1455	[13466-78-9], 3336	[13637-71-3], 4094
[12164-12-4], 4801	[12438-96-9], 4757	[13283-01-7], 4193	[13469-09-3], 0289	[13637-76-8], 4108
[12164-94-2], 4526	[12442-63-6], 4104	[13283-31-3], 0135	[13470-12-7], 4119	[13637-87-1], 3978
[12164-95-3], 4827	[12501-01-8], 4810	[13292-87-0], 0136	[13470-17-2], 4631	[13637-88-2], 3987
[12166-20-0], 4895	[12504-41-5], 4898	[13298-76-5], 3009	[13477-00-4], 0205	[13659-67-1], 4460
[12171-97-0], 3840	[12505-77-0], 0169	[13360-63-9], 2565	[13477-09-3], 0207	[13693-05-5], 4371
[12181-34-9], 4532	[12517-41-8], 4283	[13400-13-0], 4255	[13478-33-6], 4051	[13693-06-6], 4372
[12191-80-9], 4327	[12529-66-7], 4422	[13422-81-6], 1782	[13478-38-1], 4279	[13693-09-9], 4377
[12193-29-2], 3110	[12532-69-3], 4646	[13424-46-9], 4782	[13483-59-5], 0252	[13701-67-2], 0161
[12196-72-4], 4678	[12540-13-5], 0615	[13444-71-8], 4425	[13492-26-7], 4431	[13706-14-4], 3986
[12206-14-3], 4240	[12542-85-7], 1292	[13444-85-4], 4633	[13494-80-9], 4916	[13709-32-5], 4328
[12211-03-9], 4910	[12576-63-5], 3790	[13444-89-8], 0291	[13497-91-1], 4171	[13709-36-9], 4332
[12212-56-5], 3705	[12635-29-9], 0055	[13444-90-1], 4025	[13497-97-7], 0061	[13709-55-2], 4347
[12213-13-7], 3861	[12674-40-7], 1027	[13445-49-3], 4482	[13498-06-1], 4148	[13709-61-0], 4353
[12216-68-9], 3684	[12774-81-1], 1362	[13445-50-6], 4538	[13498-14-1], 4168	[13718-25-7], 4783
[12218-77-8], 4826	[12785-36-3], 0004	[13446-09-8], 4513	[13499-05-3], 4163	[13730-91-1], 0143
[12228-13-6], 0132	[13007-92-6], 2056	[13446-09-8], 4514	[13517-00-5], 4286	[13746-66-2], 2063
[12228-28-3], 0149	[13036-75-4], 4326	[13446-10-1], 4518	[13517-10-7], 0150	[13746-98-0], 4762
[12232-97-2], 0230	[13078-30-3], 2729	[13446-48-5], 4521	[13517-64-7], 4736	[13749-37-6], 1126
[12244-59-6], 2113	[13084-45-2], 1496	[13446-75-8], 4450	[13520-74-6], 0281	[13749-94-5], 1260
[12245-12-4], 3451	[13084-46-3], 1497	[13449-15-5], 4031	[13520-77-9], 0294	[13762-26-0], 4121
[12258-22-9], 0047	[13084-47-4], 0802	[13449-16-6], 4472	[13520-90-6], 0292	[13762-80-6], 4011
[12259-52-6], 4577	[13086-63-0], 0366	[13449-22-4], 1693	[13529-75-4], 2041	[13763-19-4], 4197
[12263-33-1], 4711	[13092-75-6], 0650	[13451-18-8], 4853	[13536-85-1], 4324	[13763-23-0], 4192

[13766-26-2], 0134	[13939-06-5], 2628	[14309-25-2], 3783	[14666-78-5], 2446	[15192-14-0], 4325
[13767-10-0], 4557	[13943-58-3], 2064	[14312-20-0], 0280	[14668-82-7], 1109	[15192-76-4], 0618
[13768-38-2], 4370	[13952-84-6], 1724	[14314-27-3], 4016	[14674-72-7], 3925	[15195-58-1], 4343
[13768-86-0], 4852	[13957-21-6], 2449	[14322-50-0], 4075	[14674-74-9], 0204	[15203-80-2], 4129
[13769-36-3], 0267	[13965-73-6], 0162	[14335-40-1], 3979	[14678-02-5], 1502	[15213-49-7], 0988
[13770-16-6], 4088	[13967-25-4], 4312	[14336-80-2], 4287	[14691-44-2], 4421	[15219-34-8], 0583
[13770-86-0], 0209	[13967-90-3], 0202	[14351-66-7], 3805	[14696-82-3], 4621	[15220-72-1], 3376
[13770-96-2], 0076	[13968-14-4], 4722	[14353-90-3], 2058	[14705-99-8], 1809	[15232-76-5], 3024
[13772-47-9], 4258	[13973-87-0], 0256	[14355-21-6], 0120	[14746-03-3], 2377	[15245-44-0], 2071
[13773-81-4], 4313	[13973-88-1], 4030	[14362-68-6], 0364	[14751-89-4], 3875	[15246-55-6], 2619
[13774-85-1], 4346	[13980-04-6], 1217	[14362-70-0], 0350	[14760-99-7], 1513	[15263-28-7], 4234
[13774-94-2], 4451	[13985-87-0], 3801	[14368-49-1], 2202	[14774-78-8], 2139	[15284-39-6], 1511
[13776-58-4], 4857	[13986-26-0], 4639	[14380-76-8], 4543	[14781-32-9], 1784	[15285-42-4], 2765
[13780-64-8], 4319	[13997-90-5], 3975	[14391-39-6], 0140	[14781-33-0], 1784	[15320-25-9], 4019
[13782-01-9], 4213	[14014-86-9], 4635	[14391-40-9], 0142	[14796-11-3], 2367	[15321-10-5], 3988
[13783-04-5], 0284	[14017-53-9], 4131	[14399-53-2], 0805	[14848-01-2], 2654	[15321-67-7], 3969
[13783-07-8], 4630	[14017-55-1], 4132	[14402-89-2], 1813	[14856-86-1], 1046	[15336-58-0], 0569
[13812-39-0], 4068	[14018-82-7], 4492	[14404-36-5], 4207	[14871-79-5], 0210	[15383-68-3], 1408
[13813-62-2], 4567	[14024-00-1], 2635	[14435-92-8], 0550	[14871-92-2], 3255	[15388-46-2], 3874
[13814-76-1], 4167	[14024-50-1], 3326	[14448-38-5], 4470	[14876-96-1], 4387	[15411-45-7], 1544
[13815-21-9], 3985	[14038-43-8], 3739	[14451-00-4], 3447	[14877-32-9], 0283	[15457-77-9], 4509
[13815-30-0], 4359	[14040-11-0], 2636	[14452-39-2], 0066	[14878-31-0], 1376	[15457-87-1], 4540
[13816-48-3], 3926	[14044-65-6], 0138	[14459-54-2], 4150	[14884-42-5], 4226	[15477-33-5], 0065
[13819-89-8], 4420	[14090-22-3], 1629	[14475-63-9], 4534	[14913-74-7], 0496	[15511-25-8], 3251
[13820-81-0], 4124	[14104-20-2], 0005	[14477-33-9], 1329	[14917-59-0], 2644	[15521-60-5], 4535
[13820-83-2], 4126	[14128-54-2], 0069	[14481-29-9], 2577	[14931-97-6], 2590	[15529-45-0], 3382
[13823-27-2], 4768	[14168-44-6], 0483	[14488-49-4], 0950	[14932-06-0], 1787	[15538-67-7], 0175
[13823-36-4], 4461	[14215-28-2], 4786	[14516-66-6], 3890	[14940-94-4], 4868	[15543-40-5], 4929
[13823-50-2], 4457	[14215-29-3], 3957	[14519-07-4], 0279	[14973-59-2], 4196	[15576-35-9], 1322
[13825-86-0], 4244	[14215-30-6], 4280	[14519-10-9], 0249	[14976-54-6], 3070	[15584-65-3], 3682
[13826-64-7], 4737	[14215-31-7], 4216	[14521-05-2], 4366	[14977-61-8], 4054	[15587-44-7], 4426
[13826-86-3], 0131	[14215-33-9], 4604	[14522-39-5], 3799	[14984-81-7], 4321	[15588-62-2], 4005
[13842-82-5], 4341	[14222-60-7], 3163	[14524-44-8], 3829	[14986-53-9], 4295	[15598-34-2], 1863
[13845-18-6], 4467	[14233-86-4], 1645	[14545-08-5], 0670	[14986-60-8], 4307	[15605-27-3], 4022
[13845-23-3], 4485	[14243-51-7], 2558	[14545-72-3], 4026	[14989-38-9], 4165	[15605-28-4], 4027
[13847-60-4], 4501	[14259-65-5], 4188	[14548-58-4], 3132	[14989-89-0], 1114	[15630-89-4], 0553
[13847-65-9], 4337	[14259-66-6], 4164	[14548-59-5], 2786	[15002-08-1], 4402	[15640-93-4], 0133
[13847-66-0], 4765	[14259-67-7], 3968	[14550-84-6], 0260	[15039-07-3], 4816	[15651-35-1], 4587
[13862-16-3], 4580	[14268-23-6], 0530	[14589-65-2], 4208	[15061-57-1], 4058	[15663-42-0], 0519
[13863-59-7], 0238	[14283-05-7], 4274	[14597-45-6], 2776	[15070-34-5], 4692	[15676-66-1], 3573
[13863-88-2], 0023	[14284-90-3], 3690	[14604-29-6], 2774	[15100-34-6], 0102	[15677-44-8], 3572
[13864-01-2], 4495	[14285-68-8], 3386	[14621-84-2], 0336	[15110-33-5], 4583	[15685-17-3], 3358
[13864-02-3], 3963	[14286-02-3], 4560	[14642-66-1], 2877	[15114-46-2], 1032	[15712-13-7], 1358
[13875-06-4], 4322	[14293-70-0], 4661	[14649-03-7], 3139	[15114-92-8], 2595	[15715-42-1], 1732
[13932-10-0], 4044	[14306-88-8], 3838	[14652-46-1], 4206	[15156-18-0], 4123	[15718-71-5], 0964
[13933-23-8], 4062	[14307-35-8], 4236	[14666-77-4], 3356	[15179-32-5], 4367	[15721-33-2], 2535

[15727-43-2], 3476	[16438-87-3], 0542	[17068-96-0], 0158	[17658-52-8], 3428	[18925-10-5], 1155
[15736-98-8], 0537	[16462-53-6], 3751	[17082-12-1], 3484	[17683-09-9], 3657	[18925-69-4], 3321
[15742-33-3], 4219	[16544-95-9], 4568	[17083-63-5], 0077	[17697-12-0], 3495	[18939-43-0], 4696
[15755-09-6], 0897	[16588-34-4], 2651	[17088-37-8], 3195	[17702-41-9], 0198	[18939-61-2], 4282
[15760-45-9], 3019	[16607-77-5], 2944	[17088-73-2], 0928	[17761-23-8], 1070	[19155-52-3], 0773
[15760-46-0], 3182	[16664-33-8], 3550	[17097-12-0], 0146	[17814-73-2], 0448	[19162-21-1], 3223
[15769-72-9], 3373	[16668-69-2], 3740	[17117-12-3], 2297	[17814-74-3], 3443	[19162-94-8], 0360
[15780-31-1], 4869	[16681-65-5], 1189	[17122-97-3], 2730	[17835-81-3], 4135	[19195-69-8], 3984
[15790-54-2], 1269	[16681-78-0], 0814	[17156-85-3], 0179	[17836-90-7], 4407	[19212-11-4], 3420
[15829-53-5], 4613	[16689-88-6], 4489	[17156-88-6], 0961	[17891-55-3], 3222	[19232-39-4], 2404
[15842-89-4], 3005	[16694-46-5], 1273	[17168-82-0], 0965	[17927-57-0], 0160	[19267-68-6], 1791
[15857-57-5], 4878	[16714-23-1], 2939	[17168-82-0], 0968	[17933-16-2], 0426	[19273-47-3], 1562
[15861-05-9], 4296	[16747-33-4], 3203	[17168-83-1], 1781	[18139-02-1], 0800	[19273-49-5], 1458
[15865-57-3], 2915	[16752-60-6], 4872	[17168-85-3], 4231	[18139-03-2], 1459	[19287-45-7], 0166
[15873-50-4], 2350	[16774-21-3], 3964	[17176-77-1], 3658	[18178-05-7], 4529	[19372-47-2], 3325
[15875-44-2], 0500	[16789-24-5], 4658	[17185-68-1], 1336	[18204-79-0], 1302	[19416-93-4], 0449
[15876-39-8], 3787	[16823-94-2], 4673	[17194-00-2], 0208	[18230-75-6], 1330	[19419-98-8], 0672
[15877-57-3], 2492	[16824-78-5], 3430	[17206-00-7], 2527	[18244-91-2], 2562	[19423-89-3], 4745
[15930-75-3], 4338	[16824-81-0], 3437	[17224-08-7], 0358	[18264-75-0], 0495	[19465-88-8], 3936
[15932-89-5], 0487	[16825-72-2], 2754	[17224-09-8], 0351	[18273-30-8], 0803	[19465-89-5], 4785
[15980-15-1], 1613	[16825-74-4], 2755	[17242-52-3], 4456	[18282-10-5], 4840	[19482-31-6], 2048
[15995-42-3], 2044	[16829-28-0], 4323	[17292-62-5], 0382	[18283-93-7], 0181	[19504-40-6], 4596
[16007-16-2], 3173	[16829-30-4], 4305	[17300-62-8], 0085	[18356-02-0], 0463	[19521-84-7], 2227
[16016-20-0], 2396	[16853-85-3], 0075	[17333-74-3], 2775	[18365-12-3], 1251	[19528-48-4], 3433
[16017-37-1], 4763	[16871-90-2], 4364	[17333-82-2], 2131	[18380-68-2], 0900	[19597-69-4], 4685
[16017-38-2], 4238	[16872-11-0], 0130	[17333-83-4], 2148	[18428-81-4], 4145	[19624-22-7], 0188
[16029-98-4], 1306	[16875-44-2], 0501	[17333-84-5], 2149	[18433-84-6], 0189	[19708-47-5], 0335
[16043-45-1], 4920	[16903-32-5], 0164	[17333-86-7], 2659	[18501-44-5], 4211	[19733-68-7], 1584
[16065-92-2], 4918	[16921-96-3], 4378	[17333-87-8], 2660	[18523-48-3], 0774	[19792-18-8], 1360
[16066-38-9], 3035	[16924-00-8], 4379	[17333-88-9], 2661	[18532-87-1], 4137	[19808-30-1], 1410
[16099-75-5], 4299	[16924-32-6], 0960	[17408-16-1], 3141	[18558-16-4], 0415	[19816-89-8], 0783
[16102-24-2], 2824	[16940-66-2], 0147	[17417-38-8], 4059	[18608-30-5], 1041	[19899-80-0], 1122
[16130-58-8], 1075	[16940-81-1], 4360	[17421-82-8], 3603	[18624-44-7], 4392	[19932-64-0], 0679
[16143-80-9], 3871	[16949-15-8], 0145	[17440-85-6], 0157	[18649-64-4], 2212	[19987-14-5], 2845
[16166-61-3], 3400	[16962-07-5], 0058	[17455-13-9], 0830	[18727-07-6], 0491	[20039-37-6], 3304
[16176-02-6], 1432	[16962-31-5], 4363	[17455-13-9], 3561	[18810-58-7], 0214	[20205-91-8], 0156
[16187-03-4], 1982	[16968-06-2], 1345	[17495-81-7], 4029	[18815-16-2], 4179	[20281-00-9], 4264
[16187-15-8], 1623	[16968-19-7], 3646	[17496-59-2], 4100	[18815-73-1], 0447	[20394-65-4], 4009
[16260-59-6], 2157	[16971-92-9], 3994	[17498-10-1], 1692	[18820-29-6], 4706	[20421-00-5], 3679
[16282-67-0], 0355	[16978-76-0], 3407	[17508-17-7], 2279	[18820-63-8], 4306	[20427-56-9], 4862
[16329-92-3], 0641	[16986-83-7], 2418	[17510-52-0], 1159	[18864-05-6], 3549	[20427-58-1], 4478
[16329-93-4], 1062	[17003-75-7], 0750	[17524-18-4], 4582	[18868-43-4], 4716	[20436-27-5], 0165
[16408-92-7], 0649	[17003-82-6], 0341	[17530-57-3], 3599	[18880-14-3], 1601	[20442-99-3], 3628
[16408-93-8], 0647	[17013-07-9], 0072	[17548-36-6], 3695	[18902-42-6], 4144	[20443-67-8], 3316
[16408-94-9], 0637	[17014-71-0], 4666	[17596-45-1], 0141	[18918-17-7], 3637	[20446-69-9], 0963
[16413-88-0], 2901	[17043-56-0], 0436	[17607-20-4], 1902	[18922-71-9], 2893	[20471-57-2], 2334

[20519-92-0], 1332	[21651-19-4], 4829	[22750-67-0], 4547	[23834-96-0], 0159	[25474-92-4], 4754
[20539-85-9], 2978	[21774-03-8], 4651	[22750-69-2], 1143	[23840-95-1], 0184	[25480-76-6], 1488
[20548-54-3], 3941	[21790-54-5], 3863	[22750-93-2], 0852	[23936-60-9], 2585a	[25483-10-7], 0242
[20611-53-4], 4072	[21844-15-1], 0153	[22751-18-4], 2705	[24094-93-7], 4237	[25496-08-6], 2746
[20634-12-2], 4589	[21892-31-9], 2069	[22751-23-1], 2914	[24115-83-1], 3124	[25523-79-9], 4110
[20667-12-3], 0032	[21908-53-2], 4605	[22751-24-2], 2155	[24156-53-4], 0771	[25546-98-9], 4563
[20720-71-2], 0685	[21916-66-5], 1883	[22754-96-7], 0987	[24167-76-8], 4474	[25573-59-5], 4585
[20731-44-6], 2143	[21958-87-5], 2093	[22754-97-8], 1025	[24173-36-2], 2697	[25573-61-9], 0125
[20737-02-4], 0038	[21961-22-8], 1365	[22755-00-6], 4530	[24264-08-2], 2430	[25582-86-9], 3977
[20738-31-2], 3891	[21981-95-3], 3552	[22755-01-7], 4476	[24308-92-7], 1711	[25596-24-1], 1299
[20762-60-1], 4652	[22086-53-9], 2534	[22755-07-3], 3491	[24345-74-2], 1604	[25604-70-0], 0578
[20762-98-5], 1821	[22113-51-5], 2788	[22755-09-5], 2366	[24356-04-5], 3865	[25604-71-1], 0598
[20816-12-0], 4858	[22143-20-0], 4301	[22755-14-2], 1249	[24423-68-5], 3368	[25639-45-6], 3245
[20829-66-7], 0966	[22205-57-8], 4260	[22755-22-2], 4466	[24516-71-0], 3462	[25658-42-8], 4733
[20854-03-9], 2743	[22281-49-8], 3616	[22755-25-5], 2182	[24573-95-3], 3331	[25658-43-9], 4732
[20854-05-1], 2744	[22289-82-3], 4578	[22755-27-7], 4750	[24621-17-8], 0285	[25682-07-9], 0958
[20859-73-8], 0083	[22295-11-0], 1768	[22755-34-6], 1091	[24704-64-1], 0078	[25710-89-8], 0634
[20914-03-0], 3712	[22295-99-4], 4795	[22755-36-8], 1323	[24748-23-0], 3562	[25737-28-4], 3328
[20919-99-7], 1854	[22388-72-3], 4209	[22784-01-6], 3580	[24748-25-2], 1754	[25764-08-3], 3965
[20944-65-4], 3528	[22392-07-0], 1682	[22826-61-5], 2633	[24869-88-3], 1564	[25854-38-0], 0403
[20972-64-9], 1450	[22398-80-7], 4642	[22831-39-6], 4697	[24903-95-5], 3175	[25854-41-5], 0384
[20982-74-5], 3111	[22410-18-0], 1374	[22834-73-8], 3907	[25005-56-5], 4383	[25869-93-6], 0182
[20991-79-1], 0539	[22432-68-4], 1042	[22868-13-9], 4812	[25014-41-9], 1108	[25870-02-4], 0028
[21041-85-0], 0717	[22471-42-7], 2623	[22887-15-6], 4270	[25021-40-5], 3029	[25875-18-7], 1064
[21050-95-3], 2150	[22480-64-4], 2130	[22978-89-8], 3640	[25038-59-9], 3262	[25910-37-6], 2201
[21107-27-7], 1334	[22493-01-2], 4523	[22996-18-5], 2716	[25067-58-7], 0687	[25987-94-4], 0302
[21109-95-5], 0218	[22524-35-2], 1238	[23097-77-0], 4141	[25077-87-6], 2138	[26028-46-6], 3183
[21138-22-7], 4416	[22524-45-4], 2510	[23115-33-5], 2602	[25081-01-0], 4024	[26134-62-3], 4688
[21230-20-6], 2273	[22541-59-9], 4896	[23129-50-2], 3906	[25136-85-0], 0337	[26156-56-9], 4125
[21245-43-2], 3692	[22541-88-4], 4896	[23143-88-6], 0339	[25147-05-1], 0743	[26157-96-0], 2054
[21249-40-0], 3811	[22575-95-7], 3037	[23273-02-1], 0967	[25167-31-1], 0786	[26163-27-9], 0689
[21251-20-7], 1403	[22585-64-4], 0239	[23292-52-6], 0627	[25167-81-1], 2161	[26163-27-9], 0712
[21254-73-9], 1839	[22620-90-2], 0617	[23303-78-8], 4342	[25196-24-9], 3378	[26222-92-4], 0236
[21255-83-4], 0258	[22650-46-0], 4060	[23326-27-4], 1881	[25230-72-2], 2742	[26241-10-1], 4662
[21261-07-4], 3371	[22653-19-6], 1739	[23362-09-6], 0499	[25233-34-5], 1407	[26249-01-4], 2658
[21299-20-7], 3455	[22675-68-9], 1035	[23380-38-3], 3799	[25238-02-2], 1128	[26257-00-1], 4414
[21372-60-1], 1172	[22688-10-6], 1979	[23411-89-4], 3587	[25240-93-1], 0545	[26283-13-6], 3970
[21380-82-5], 1628	[22691-91-4], 2395	[23414-72-4], 4710	[25251-03-0], 0240	[26299-14-9], 1862
[21473-42-1], 0401	[22692-30-4], 2934	[23418-85-1], 3399	[25253-54-7], 4109	[26351-01-9], 0073
[21482-59-7], 1779	[22710-73-2], 2954	[23679-20-1], 2456	[25284-83-7], 0490	[26386-83-9], 3489
[21483-09-8], 3753	[22722-03-8], 1327	[23705-25-1], 2880	[25382-52-9], 0402	[26388-78-3], 1786
[21519-20-0], 3862	[22722-98-1], 2575	[23753-67-5], 0186	[25393-98-0], 3268	[26445-82-9], 0174
[21548-73-2], 0026	[22750-47-6], 0228	[23777-55-1], 0513	[25395-83-9], 0213	[26464-99-3], 2045
[21548-73-2], 0035	[22750-53-4], 3954	[23777-63-1], 0167	[25398-08-7], 3253	[26493-63-0], 1775
[21572-61-2], 1807	[22750-56-7], 0614	[23777-80-2], 0191	[25402-50-0], 4334	[26519-91-5], 2363
[21645-51-2], 0074	[22750-57-8], 4262	[23784-96-5], 1830	[25455-73-6], 0033	[26522-91-8], 0270

[26604-66-0], 4437	[28781-86-4], 1886	[30962-16-4], 0378	[32707-10-1], 0235	[34489-60-6], 3474
[26628-22-8], 4758	[28815-10-3], 0203	[30994-24-2], 1846	[32714-73-5], 3747	[34521-47-6], 3302
[26645-10-3], 1672	[28853-06-7], 0973	[31044-86-7], 1937	[32750-98-4], 2061	[34529-29-8], 1504
[26717-79-3], 1847	[28872-17-5], 3106	[31052-46-7], 4265	[32755-26-3], 0319	[34641-74-2], 4635
[26761-81-9], 1587	[28950-34-7], 4770	[31058-64-7], 4276	[32817-38-2], 3429	[34701-14-9], 3140
[26830-95-5], 2917	[28965-70-0], 0190	[31122-35-7], 3723	[32970-52-8], 3198	[34729-49-2], 3824
[26981-93-1], 0474	[28995-89-3], 3237	[31122-42-6], 3724	[33017-08-2], 1103	[34876-30-7], 1679
[26981-95-3], 1274	[29014-32-2], 3541	[31122-43-7], 3722	[33021-02-2], 3072	[34992-80-8], 1688
[27063-40-7], 2443	[29119-58-2], 1561	[31123-28-1], 2886	[33100-27-5], 0830	[35038-45-0], 0551
[27070-49-1], 0768	[29128-41-4], 0913	[31247-72-0], 0019	[33175-34-7], 2243	[35038-46-1], 0388
[27096-31-7], 2580	[29146-24-5], 3067	[31255-94-4], 4233	[33196-65-5], 1778	[35059-50-8], 2423
[27165-08-8], 2719	[29149-32-4], 1646	[31266-90-1], 2258	[33212-68-9], 0909	[35175-75-8], 1680
[27195-67-1], 3039	[29149-89-1], 1015	[31279-70-6], 2618	[33272-09-2], 3256	[35207-75-1], 3411
[27218-12-8], 4368	[29202-04-8], 1357	[31328-33-3], 3818	[33327-66-1], 0587	[35295-56-5], 0610
[27218-16-2], 4101	[29285-24-3], 4671	[31330-22-0], 2448	[33362-98-0], 4202	[35321-46-1], 1134
[27275-90-7], 0370	[29342-58-3], 3453	[31332-72-6], 2620	[33364-51-1], 1867	[35356-48-3], 1033
[27278-01-4], 0570	[29342-61-8], 3499	[31340-36-0], 1052	[33393-70-3], 3460	[35357-46-1], 1055
[27318-90-7], 3578	[29362-18-3], 2647	[31366-25-3], 2215	[33421-43-1], 3258	[35398-20-0], 1843
[27505-49-3], 4020	[29584-42-7], 4850	[31724-50-2], 0522	[33497-88-0], 4225	[35452-30-3], 2020
[27610-45-2], 4811	[29671-92-9], 0468	[31841-79-9], 1072	[33506-43-3], 2932	[35551-00-7], 3349
[27661-44-5], 3764	[29674-96-2], 0511	[31890-52-5], 1662	[33529-09-8], 3432	[35629-70-0], 1500
[27662-11-9], 2164	[29678-92-0], 4899	[31921-03-6], 3348	[33543-69-0], 3402	[35638-41-6], 3729
[27662-34-6], 3910	[29727-86-4], 0901	[32006-07-8], 0273	[33634-75-2], 2051	[35784-01-1], 2424
[27698-99-3], 0971	[29903-04-6], 3831	[32028-94-7], 4415	[33689-28-0], 1440	[35847-40-6], 3116
[27778-27-4], 3889	[29906-35-2], 2090	[32061-49-7], 1429	[33691-99-3], 3828	[35856-82-7], 2508
[27846-30-6], 1152	[29914-92-9], 2547	[32062-14-4], 0259	[33744-03-5], 1489	[35869-33-1], 2604
[27878-36-0], 3131	[30030-30-9], 3795	[32078-95-8], 4487	[33759-44-3], 1952	[35869-74-0], 1575
[27883-34-7], 3802	[30047-26-8], 3843	[32106-51-7], 4408	[33774-96-8], 4067	[35941-38-9], 3167
[27890-58-0], 4791	[30148-50-6], 2611	[32175-44-3], 4300	[33817-92-4], 3052	[35967-34-3], 1477
[27961-70-2], 1050	[30180-52-0], 0661	[32229-04-2], 3393	[33827-55-3], 4051	[35967-89-6], 0006
[28016-59-3], 3676	[30242-75-2], 3794	[32283-21-9], 2677	[33842-02-3], 1950	[35967-90-9], 0201
[28051-68-5], 1568	[30341-38-9], 0648	[32312-22-4], 0244	[33843-02-3], 1209	[36016-40-7], 3170
[28068-05-5], 4559	[30426-58-5], 0693	[32312-23-5], 0245	[33849-97-1], 4440	[36097-38-8], 3252
[28103-68-6], 1509	[30427-53-3], 3244	[32315-10-9], 1040	[33873-53-9], 2957	[36133-73-0], 2609
[28128-14-5], 1185	[30459-28-0], 1789	[32315-88-1], 1615	[33884-41-2], 3014	[36211-73-1], 2318
[28212-50-2], 1431	[30478-96-7], 3357	[32349-29-4], 1572	[33965-80-9], 1130	[36294-69-6], 0969
[28223-30-5], 4073	[30533-63-2], 1175	[32366-26-0], 3159	[34005-83-9], 0645	[36333-41-2], 1351
[28240-69-9], 0797	[30582-57-1], 4330	[32390-97-1], 4252	[34018-28-5], 0278	[36336-08-0], 0442
[28255-99-4], 2329	[30604-81-0], 1417	[32434-37-5], 3955	[34037-79-1], 2989	[36525-64-1], 1372
[28262-13-7], 3168	[30616-12-7], 3754	[32454-63-0], 3917	[34090-17-0], 4562	[36609-92-4], 3991
[28375-94-2], 0192	[30641-57-7], 0121	[32607-15-1], 4755	[34277-48-0], 2820	[36691-97-1], 3725
[28569-48-4], 2305	[30645-13-7], 1098	[32607-31-1], 0989	[34324-17-9], 4516	[36732-77-1], 3869
[28597-01-5], 2246	[30708-80-6], 3982	[32617-22-4], 0377	[34410-88-3], 3956	[36794-93-1], 3138
[28655-95-0], 1533	[30734-83-9], 3416	[32673-60-2], 0841	[34461-00-2], 1076	[36841-23-3], 3882
[28663-71-0], 2070	[30911-12-7], 4629	[32707-02-1], 2186	[34461-68-2], 4272	[36913-76-5], 3881
[28711-64-0], 3548	[30957-47-2], 1104	[32707-03-2], 2187	[34464-38-5], 3366	[37035-60-2], 3911

[37067-42-8], 4496	[39422-73-6], 4798	[43135-83-7], 3363	[51593-99-8], 4672	[53474-95-6], 4718
[37150-27-9], 2191	[39422-94-1], 3461	[44023-01-6], 4595	[51655-36-8], 1636	[53477-43-3], 3000
[37205-27-9], 3706	[39437-99-5], 1743	[44397-85-5], 0763	[51680-55-8], 4541	[53490-67-8], 2035
[37218-25-0], 0003	[39597-90-5], 2336	[44750-17-6], 2173	[51688-25-6], 3397	[53496-15-4], 2865
[37235-82-8], 0232	[39687-95-1], 1470	[45101-14-2], 1760	[51819-69-3], 4086	[53504-62-4], 3461
[37245-81-1], 4714	[39729-81-2], 4275	[45722-89-2], 2388	[51821-32-0], 0857	[53504-66-8], 3461
[37271-59-3], 0049	[39733-13-6], 4586	[48114-68-7], 2822	[51832-56-5], 3520	[53504-80-6], 3285
[37279-26-8], 3424	[39753-42-9], 2840	[49558-46-5], 2700	[51877-12-4], 1387	[53534-20-6], 0938
[37297-87-3], 0975	[39957-12-5], 4570	[49559-20-8], 4458	[52003-45-9], 0321	[53535-47-0], 3434
[37298-41-2], 3324	[40089-12-1], 3133	[49571-31-5], 2655	[52096-16-9], 0777	[53578-07-7], 1247
[37346-96-6], 4669	[40237-34-1], 0914	[49716-47-4], 3689	[52096-18-1], 2283	[53654-97-0], 0318
[37388-50-4], 4315	[40330-52-7], 4227	[50274-95-8], 2678	[52096-20-5], 2282	[53808-72-3], 3174
[37414-44-1], 3288	[40330-52-7], 4228	[50277-65-1], 1065	[52096-22-7], 2704	[53808-74-5], 3340
[37436-17-2], 2374	[40330-52-7], 4230	[50311-48-3], 0346	[52106-86-2], 2720	[53808-76-7], 3659
[37480-75-4], 3591	[40358-04-1], 1388	[50327-56-5], 3813	[52118-12-4], 3844	[53847-48-6], 2723
[37609-69-1], 2948	[40428-75-9], 1438	[50424-93-6], 2913	[52157-57-0], 2352	[53852-44-1], 2204
[37647-36-2], 3680	[40560-76-7], 2780	[50577-64-5], 0392	[52324-65-9], 4212	[53939-55-2], 3701
[37759-21-0], 3842	[40561-27-1], 1506	[50623-64-8], 3645	[52412-56-3], 3515	[53966-09-9], 3478
[37895-51-5], 1415	[40575-26-6], 3235	[50645-52-8], 4390	[52470-25-4], 0509	[54007-00-0], 3685
[38005-27-5], 4308	[40575-26-6], 3236	[50646-37-2], 4921	[52716-12-8], 1924	[54065-28-0], 4553
[38005-31-1], 4140	[40661-97-0], 0433	[50650-92-5], 0709	[52732-14-6], 3590	[54086-40-7], 0306
[38078-09-0], 1684	[40898-91-7], 4043	[50662-24-3], 2042	[52762-90-0], 3611	[54086-41-8], 0574
[38092-76-1], 0951	[40953-35-3], 2050	[50707-40-9], 3171	[52781-75-6], 3452	[54153-83-2], 4253
[38115-19-4], 4316	[40985-41-9], 0469	[50744-08-6], 3179	[52795-23-0], 4381	[54196-34-8], 0451
[38139-15-0], 0520	[41029-45-2], 1461	[50764-78-8], 1214	[52851-26-0], 2398	[54217-98-0], 3464
[38232-63-2], 4612	[41084-90-6], 0621	[50831-29-3], 1785	[52913-14-1], 2821	[54217-99-1], 3464
[38293-27-5], 1844	[41334-40-1], 3119	[50846-98-5], 1082	[52936-24-0], 2520	[54222-33-2], 3359
[38344-58-0], 4382	[41337-62-6], 3798	[50859-36-2], 0096	[52936-25-1], 0795	[54222-34-3], 3707
[38361-85-2], 3100	[41409-50-1], 1371	[50892-86-9], 3438	[52936-53-1], 1250	[54448-39-4], 0472
[38386-55-9], 3038	[41422-43-9], 1769	[50892-90-5], 3439	[52999-19-6], 3545	[54514-12-4], 2736
[38551-29-0], 4550	[41463-64-3], 1018	[50892-99-4], 1872	[53004-03-8], 3161	[54514-15-7], 2751
[38682-34-7], 0123	[41481-90-7], 1792	[50906-29-1], 2898	[53055-05-3], 2937	[54514-15-7], 2752
[38711-69-2], 3577	[41570-79-0], 3742	[51006-26-9], 3867	[53078-10-7], 4161	[54524-31-1], 0767
[38787-96-1], 1235	[41909-62-0], 4665	[51027-90-8], 2984	[53088-06-5], 3600	[54567-24-7], 0820
[38870-89-2], 1165	[41924-27-0], 3584	[51104-87-1], 1353	[53144-64-2], 0825	[54573-23-8], 1871
[38892-09-0], 2115	[42017-07-2], 2834	[51138-16-0], 3001	[53149-77-2], 3908	[54675-76-2], 3901
[38910-25-8], 2474	[42081-47-0], 4257	[51184-04-4], 4553	[53151-88-5], 3566	[54689-17-7], 4220
[38989-47-8], 4354	[42087-90-1], 3477	[51249-08-2], 2280	[53182-16-4], 3178	[54747-08-9], 3612
[39003-82-2], 0093	[42238-29-9], 2205	[51269-39-7], 3803	[53201-99-3], 4271	[54847-21-1], 4552
[39047-21-7], 1795	[42246-25-3], 4045	[51286-83-0], 1106	[53213-77-7], 3746	[55011-46-6], 0387
[39108-12-8], 4694	[42294-95-6], 0589	[51286-84-1], 1119	[53213-78-8], 2589	[55042-15-4], 4181
[39108-14-0], 0082	[42404-59-1], 3543	[51317-78-3], 3322	[53214-97-4], 1799	[55044-04-7], 0811
[39254-48-3], 2338	[42454-06-8], 2692	[51325-04-3], 1402	[53346-21-7], 3303	[55058-76-9], 3664
[39254-49-4], 2219	[42489-15-6], 4715	[51329-17-0], 3099	[53378-72-6], 2046	[55060-79-2], 3717
[39274-39-0], 0356	[42494-76-1], 1200	[51451-05-9], 0787	[53421-36-6], 0622	[55060-80-5], 3540
[39409-82-0], 0534	[43008-25-9], 2160	[51470-74-7], 2480	[53436-80-9], 0067	[55060-82-7], 3716

[55106-91-7], 3238	[57508-69-7], 3115	[59598-02-6], 2613	[63104-54-1], 3232	[65343-66-0], 3227
[55106-92-8], 3389	[57530-25-3], 1194	[59643-77-5], 3625	[63105-40-8], 4385	[65381-02-4], 3473
[55106-93-9], 3387	[57664-97-8], 2887	[59675-70-6], 4130	[63128-08-5], 3770	[65521-60-0], 1044
[55106-95-1], 3735	[57670-85-6], 1002	[59744-77-3], 4687	[63148-65-2], 3030	[65579-06-8], 0089
[55106-96-2], 3734	[57678-14-5], 3143	[59795-18-5], 4739	[63160-33-8], 2830	[65579-07-9], 0183
[55124-14-6], 0394	[57707-64-9], 0714	[59865-91-7], 4777	[63181-09-9], 3228	[65597-24-2], 0320
[55153-36-1], 2512	[57846-03-4], 2346	[59871-26-0], 1240	[63246-75-3], 3335	[65597-25-3], 0594
[55341-15-6], 1554	[57891-85-7], 2114	[60241-33-0], 4775	[63338-74-9], 4502	[65651-91-4], 2586
[55358-23-1], 3870	[57965-32-9], 2250	[60345-95-1], 0981	[63354-77-8], 2669	[65664-23-5], 2896
[55358-25-3], 3847	[58249-49-3], 3990	[60380-67-8], 4432	[63382-64-9], 2467	[65776-39-8], 0086
[55358-29-7], 3870	[58281-24-6], 3470	[60489-76-1], 3808	[63609-41-6], 3148	[65901-39-5], 4053
[55358-33-3], 3870	[58302-41-3], 3627	[60553-18-6], 0752	[63618-84-8], 3636	[66080-22-6], 3423
[55374-98-6], 4070	[58302-42-4], 2642	[60580-20-3], 4170	[63643-78-7], 0799	[66217-76-3], 2799
[55644-07-0], 1803	[58412-05-8], 4659	[60617-65-4], 4349	[63672-69-3], 4152	[66326-45-2], 4008
[55644-07-0], 1804	[58426-27-0], 1206	[60633-76-3], 0393	[63706-85-4], 0163	[66326-46-3], 4010
[55761-75-8], 3602	[58430-57-2], 0879	[60672-60-8], 0755	[63839-60-1], 1144	[66427-01-8], 0454
[55800-08-3], 3835	[58501-09-0], 4224	[60784-40-9], 1131	[63915-60-6], 1590	[66459-02-7], 0080
[55864-39-6], 1192	[58572-21-7], 4753	[60955-41-1], 4794	[64010-65-7], 4273	[66486-68-8], 2641
[55870-36-5], 1793	[58609-72-6], 3614	[61014-23-1], 3181	[64057-57-4], 0780	[66599-67-5], 3589
[55912-20-4], 2717	[58656-11-4], 3377	[61033-15-6], 2359	[64297-36-5], 3233	[66649-16-9], 3275
[55999-69-4], 1749	[58659-15-7], 3028	[61204-16-8], 2095	[64297-64-9], 2681	[66737-76-7], 2213
[56090-02-9], 2607	[58674-00-3], 4527	[61224-59-7], 3392	[64297-75-2], 3319	[66793-67-7], 1373
[56092-91-2], 2630	[58675-02-8], 1452	[61447-07-2], 2835	[64365-00-0], 3909	[66862-11-1], 2592
[56094-15-6], 4660	[58696-86-9], 2803	[61514-68-9], 2218	[64445-49-4], 2206	[66946-48-3], 3263
[56139-33-4], 1926	[58776-08-2], 2251	[61565-16-0], 1386	[64624-44-8], 1874	[66955-43-9], 0737
[56280-76-3], 4789	[58816-51-8], 0959	[61695-12-3], 3576	[64771-59-1], 1356	[66955-44-0], 1167
[56295-87-5], 4793	[58911-30-1], 0593	[61832-41-5], 1596	[64781-77-7], 2910	[66986-75-2], 3699
[56370-81-1], 1794	[58941-14-3], 0739	[62072-11-1], 2831	[64781-78-8], 0677	[67016-28-8], 4464
[56386-22-2], 3859	[58941-15-4], 1168	[62072-18-8], 3118	[64825-48-5], 2614	[67165-30-4], 4232
[56391-49-2], 3789	[59034-32-1], 1414	[62116-25-0], 1201	[64867-41-0], 0029	[67187-50-2], 2621
[56411-66-6], 2298	[59034-33-2], 1866	[62126-20-9], 0068	[65036-47-7], 2097	[67312-43-0], 0547
[56411-67-7], 2339	[59034-34-3], 1411	[62127-48-4], 1550	[65037-43-6], 2674	[67362-62-3], 0824
[56422-03-8], 4728	[59104-93-7], 0680	[62133-36-2], 2588	[65039-20-5], 2796	[67400-85-2], 0408
[56433-01-3], 2671	[59183-17-4], 1492	[62316-46-5], 4766	[65082-64-6], 3412	[67401-50-7], 3305
[56522-42-0], 0683	[59183-18-5], 1491	[62375-91-1], 2782	[65107-36-0], 4760	[67405-47-4], 2536
[56533-30-3], 0098	[59261-17-5], 1538	[62497-91-0], 3224	[65232-69-1], 2555	[67517-99-7], 3665
[56533-31-4], 4380	[59320-16-0], 2977	[62554-90-9], 2133	[65235-78-1], 3372	[67536-44-1], 0708
[56581-58-9], 3885	[59327-98-9], 2708	[62597-99-3], 4055	[65235-79-2], 0015	[67711-62-0], 2387
[56743-33-0], 2699	[59348-62-8], 1081	[62763-56-8], 4014	[65313-31-7], 1272	[67722-96-7], 1182
[56892-92-5], 0154	[59391-85-4], 2947	[62785-41-5], 0425	[65313-32-8], 0101	[67772-28-5], 1536
[56929-36-3], 0778	[59419-71-5], 1776	[62861-56-7], 1426	[65313-33-9], 0892	[67849-01-8], 0059
[56943-26-1], 0173	[59420-01-8], 3421	[62861-57-8], 2353	[65313-34-0], 1676	[67849-02-9], 0081
[57284-58-7], 2121	[59483-61-3], 2231	[62865-97-6], 2337	[65313-35-1], 2391	[67870-97-7], 2576
[57304-94-6], 3856	[59544-89-7], 0099	[62905-84-4], 2847	[65313-36-2], 0798	[67870-99-9], 3766
[57421-56-4], 1071	[59559-04-7], 2479	[62909-81-3], 4079	[65313-37-3], 1748	[67880-11-9], 0772
[57425-12-4], 4491	[59589-13-8], 3123	[63085-06-3], 3435	[65321-68-8], 2783	[67880-13-1], 0009

[67880-14-2], 4655	[68979-48-6], 1835	[71178-86-4], 3534	[73452-31-0], 2612	[76695-67-5], 3796
[67880-15-3], 4759	[68993-02-2], 4784	[71231-59-9], 3102	[73452-32-1], 2878	[76695-68-6], 3743
[67880-17-5], 0720	[69042-75-7], 3385	[71250-00-5], 0518	[73469-45-1], 2624	[76710-75-3], 3342
[67880-19-7], 4092	[69101-54-8], 4592	[71257-31-3], 4879	[73506-23-7], 4780	[76710-76-4], 3899
[67880-20-0], 0818	[69167-99-3], 3018	[71356-36-0], 3018	[73506-39-5], 1478	[76710-77-5], 3827
[67880-21-1], 1347	[69180-59-2], 2753	[71359-61-0], 1034	[73513-15-2], 3120	[76828-34-7], 2403
[67880-22-2], 1017	[69338-55-2], 3521	[71359-62-1], 1352	[73513-16-3], 4790	[77486-94-3], 3395
[67880-25-5], 4035	[69517-30-2], 0742	[71359-63-2], 0653	[73513-17-4], 4077	[77625-37-7], 3522
[67880-26-6], 2737	[69587-03-5], 3247	[71359-64-3], 1806	[73526-98-4], 2421	[77628-04-7], 3192
[67880-27-7], 0379	[69597-04-6], 2740	[71369-69-2], 2343	[73566-71-5], 3414	[77664-36-9], 3820
[67880-28-8], 0383	[69681-97-6], 3154	[71404-13-2], 3762	[73593-33-6], 4525	[77695-66-4], 3797
[67922-18-3], 1742	[69681-98-7], 3154	[71799-32-1], 2145	[73680-23-6], 3279	[77812-44-3], 3846
[68206-59-5], 3466	[69681-99-8], 3313	[71901-54-7], 0917	[73730-53-7], 0049	[78114-26-8], 3417
[68348-85-6], 0995	[69682-00-4], 3313	[71939-10-1], 1210	[73771-13-8], 2086	[78141-00-1], 3243
[68379-32-8], 1013	[69682-01-5], 3314	[72040-09-6], 1490	[73954-59-3], 4051	[78343-32-5], 1058
[68436-99-7], 0180	[69720-86-1], 3864	[72044-13-4], 0523	[74087-85-7], 1227	[78350-94-4], 2177
[68448-47-5], 3248	[69833-15-4], 3585	[72151-31-6], 4356	[74093-43-9], 0303	[78398-48-8], 3834
[68499-65-0], 3394	[69932-17-8], 0703	[72172-64-6], 3953	[74167-05-8], 3425	[78657-29-1], 1753
[68533-38-0], 0148	[70142-16-4], 0312	[72248-72-7], 4093	[74190-87-7], 3559	[78790-57-5], 3160
[68560-64-3], 2209	[70187-32-5], 1997	[72250-07-8], 4779	[74273-75-9], 1400	[78925-61-8], 3535
[68574-13-0], 1750	[70212-12-3], 3415	[72385-44-5], 1927	[74311-45-7], 4128	[78937-12-9], 1811
[68574-15-2], 2006	[70247-32-4], 0974	[72399-85-0], 1869	[74386-96-2], 0391	[78937-14-1], 2049
[68574-17-4], 2005	[70247-49-3], 3114	[72437-42-4], 3606	[74415-60-4], 3234	[78948-09-1], 0751
[68594-17-2], 0371	[70247-50-6], 1419	[72443-10-8], 2216	[74449-37-9], 4384	[79151-63-6], 3853
[68594-19-4], 0827	[70247-51-7], 1409	[72447-41-7], 3156	[74542-20-4], 4357	[79251-56-2], 2548
[68594-24-1], 1012	[70277-99-5], 2974	[72644-04-3], 3768	[74637-34-6], 3193	[79251-57-3], 2548
[68596-88-3], 3128	[70278-00-1], 2242	[72780-86-0], 4601	[74936-21-3], 0815	[79263-47-1], 3146
[68596-89-4], 2663	[70299-48-8], 1282	[72848-18-1], 0721	[74999-19-2], 0826	[79403-75-1], 3633
[68596-91-8], 2758	[70324-20-8], 2670	[72957-64-3], 2364	[75027-71-3], 3872	[79472-87-0], 1434
[68596-92-9], 2757	[70324-23-1], 3398	[72985-54-7], 1061	[75038-71-0], 1800	[79723-21-0], 0427
[68596-93-0], 2137	[70324-26-4], 3757	[72985-56-9], 0595	[75109-11-4], 3271	[79796-14-8], 4664
[68596-94-1], 2756	[70324-27-5], 2982	[73017-74-2], 2867	[75142-81-3], 3662	[79796-40-0], 4767
[68596-95-2], 2254	[70324-35-5], 2256	[73033-58-6], 2718	[75441-10-0], 3620	[79999-60-3], 3647
[68596-99-6], 2084	[70343-15-6], 2919	[73132-51-5], 1921	[75476-36-7], 3310	[80005-73-8], 3761
[68597-02-4], 3246	[70343-16-7], 2666	[73202-12-7], 4389	[75516-37-9], 3769	[80044-09-3], 1191
[68597-05-7], 2102	[70343-16-7], 2667	[73257-07-5], 3785	[75524-40-2], 0467	[80044-10-6], 1191
[68597-06-8], 2106	[70348-66-2], 1594	[73335-41-8], 3784	[75900-58-2], 4358	[80410-45-3], 3855
[68597-08-0], 2784	[70473-76-6], 2307	[73347-64-5], 4309	[75952-58-8], 1297	[80466-56-4], 1436
[68597-10-4], 2668	[70664-49-2], 1384	[73378-53-7], 4447	[75993-65-6], 0421	[80605-27-2], 3509
[68602-57-3], 1057	[70682-07-4], 1816	[73378-54-8], 4429	[76002-04-5], 0817	[80605-36-3], 3509
[68674-44-2], 0603	[70806-67-6], 4256	[73378-55-9], 4448	[76321-00-0], 3422	[80670-36-6], 2909
[68795-10-8], 0749	[70843-45-7], 3345	[73378-56-0], 4259	[76371-73-8], 3408	[80680-07-5], 4848
[68825-98-9], 0288	[70865-40-6], 0196	[73399-68-5], 1412	[76429-97-5], 1340	[80841-08-3], 3505
[68844-25-7], 3617	[70950-00-4], 0573	[73399-71-0], 1413	[76429-98-6], 1080	[80866-80-4], 2715
[68868-87-1], 3512	[70992-03-9], 4277	[73399-75-4], 3721	[76556-13-3], 0823	[80913-65-1], 1997
[68897-47-2], 0505	[71155-12-9], 3405	[73400-02-9], 4648	[76695-64-2], 3744	[80913-66-2], 1997

[81181-41-5], 3109	[85620-95-7], 1216	[90584-32-0], 0794	[101672-04-2], 1215	[113634-30-3], 3868
[81341-46-0], 0640	[85807-68-7], 0678	[90635-49-7], 3791	[101672-05-3], 1546	[113728-41-9], 4138
[81852-50-8], 1890	[85807-69-8], 1853	[90635-53-3], 3833	[101672-06-4], 1573	[113753-12-1], 4597
[81924-93-8], 2037	[85836-55-1], 3286	[91364-94-2], 3648	[101672-07-5], 1207	[113925-83-0], 0458
[81971-84-8], 2420	[85861-34-3], 3125	[91410-67-2], 3518	[101672-08-6], 3973	[114328-95-9], 2232
[82140-87-2], 1824	[85923-32-6], 3675	[91511-39-6], 1120	[101672-09-7], 3537	[114515-72-9], 3815
[82148-84-3], 3913	[85962-45-4], 0064	[91686-86-1], 1437	[101672-10-0], 0946	[114752-52-2], 1835
[82177-75-1], 2638	[86013-87-8], 0667	[91742-17-5], 0596	[101672-11-1], 1449	[115045-20-4], 4442
[82177-80-8], 2075	[86045-52-5], 2615	[91879-79-7], 3698	[101672-14-4], 2299	[115057-40-4], 1346
[82198-80-9], 1154	[86202-43-9], 0412	[91935-83-0], 0636	[101672-16-6], 2627	[115057-44-8], 1083
[82434-14-8], 3678	[86255-25-6], 2126	[93228-65-0], 3752	[101672-17-7], 1406	[115127-49-6], 2645
[82434-33-1], 4205	[86341-95-9], 2176	[93481-56-2], 1721	[101672-18-8], 4080	[115383-22-7], 0298
[82522-47-2], 2538	[86425-12-9], 2165	[93506-63-9], 2986	[101672-19-9], 2664	[115464-59-0], 1102
[82647-51-6], 3877	[86886-16-0], 2538	[93693-72-2], 2127	[101672-20-2], 1879	[115609-87-5], 1819
[82865-33-6], 1202	[86886-17-1], 3069	[93895-68-2], 3893	[101672-21-3], 4006	[115627-20-8], 3817
[82906-06-7], 1820	[87050-94-0], 1047	[94089-34-6], 1097	[101672-22-4], 2634	[116005-09-5], 3812
[82921-86-6], 3780	[87050-95-1], 1812	[94193-96-3], 2823	[101672-23-5], 1543	[116778-64-4], 2466
[82944-28-3], 1213	[87272-16-0], 2953	[94286-09-6], 2903	[101697-88-5], 3280	[117412-28-9], 0970
[82963-15-3], 4386	[87318-59-0], 3375	[94288-47-8], 3778	[101944-90-5], 1121	[118112-40-6], 1920
[83195-98-6], 3013	[87413-09-0], 3610	[94484-09-0], 3383	[102210-64-0], 4675	[118112-41-7], 1920
[83297-43-2], 1876	[87425-02-3], 2785	[94633-25-7], 3320	[102210-65-1], 4676	[118171-56-5], 1625
[83314-21-0], 4369	[87478-73-7], 0577	[94643-57-9], 3878	[102488-24-4], 1301	[118235-47-5], 3821
[83350-11-2], 3879	[87489-52-9], 1020	[94643-57-9], 3880	[102502-32-9], 1625	[119391-53-6], 0195
[83459-96-5], 3837	[87490-22-0], 4391	[94791-14-7], 0905	[102613-59-2], 1399	[119701-48-3], 3475
[83665-55-8], 1928	[87532-69-2], 3673	[95112-14-4], 1268	[102613-59-2], 1422	[120312-56-3], 2295
[83673-81-8], 2427	[87552-19-0], 0980	[95247-08-1], 3582	[102868-70-2], 3463	[121832-04-0], 0753
[83844-91-1], 3608	[87604-86-2], 3441	[95322-62-6], 3391	[103698-31-3], 1129	[122530-01-2], 3787
[83844-92-2], 3607	[87612-48-4], 3293	[95407-64-0], 2470	[103760-93-6], 0052	[123002-86-8], 3108
[83931-97-8], 2955	[87612-49-5], 3327	[95407-67-3], 1166	[104172-76-1], 2777	[123002-86-8], 3858
[84159-29-5], 4344	[87830-30-6], 2918	[95650-61-6], 1997	[104364-15-0], 1401	[123002-87-9], 3108
[84393-12-4], 3213	[88253-43-4], 3186	[96504-50-6], 3536	[104503-71-1], 2351	[123002-87-9], 3884
[84487-84-3], 3551	[88433-73-2], 4034	[96570-67-1], 3462	[104548-30-3], 3732	[123002-93-7], 3496
[84521-55-1], 3588	[88487-87-6], 2481	[97850-58-3], 2825	[104813-96-9], 4087	[123147-84-2], 3226
[84578-88-1], 3454	[88487-88-7], 3049	[97953-19-0], 4714	[105017-95-6], 3661	[123147-95-5], 3225
[84578-90-5], 3626	[88495-99-2], 4182	[98316-02-0], 3093	[105224-90-6], 1485	[123311-08-0], 1772
[84582-81-0], 3826	[88519-63-5], 3709	[98456-61-2], 2468	[105643-77-4], 1473	[123668-66-6], 3652
[84691-20-3], 1861	[88544-01-8], 3598	[98566-27-9], 2368	[107585-03-5], 0414	[124330-23-0], 0071
[84864-30-2], 1379	[88906-07-4], 0993	[98571-91-6], 3677	[107633-87-4], 3688	[124687-11-2], 0057
[84928-98-3], 2528	[89028-37-5], 3629	[99319-30-9], 0462	[107783-62-0], 3776	[125261-72-5], 1630
[84928-99-4], 2482	[89452-69-0], 3644	[99685-96-8], 0298	[107981-97-5], 0419	[125261-73-6], 1630
[84937-00-8], 4581	[89521-42-6], 3649	[99685-96-8], 3916	[109745-04-2], 1953	[125358-28-3], 2921
[85060-25-9], 1851	[89521-43-7], 2734	[99705-27-8], 1933	[109745-05-3], 1953	[125735-38-8], 2848
[85060-27-1], 1852	[89521-44-8], 3651	[99796-78-8], 1339	[109799-62-4], 2902	[126208-50-2], 3635
[85489-61-8], 1997	[90052-22-5], 1505	[100113-85-7], 3883	[110725-79-6], 3710	[126330-30-1], 2455
[85576-52-9], 1598	[90238-04-3], 3654	[100242-83-9], 1588	[111016-97-8], 3816	[126487-10-3], 0298
[85590-62-1], 1903	[90329-44-5], 1603	[101652-99-7], 3750	[111422-41-4], 3782	[126487-13-6], 3792

[126623-64-1], 2792 [133610-01-2], 3841 [153757-93-8], 0822 [159148-53-5], 3832 [168848-75-7], 3920
[126623-77-6], 3403 [134927-04-1], 2381 [153757-95-0], 0718 [159793-62-1], 1328 [168848-76-8], 3921
[127510-65-0], 1045 [135285-90-4], 2325 [155147-41-4], 3720 [159793-64-3], 1324 [171565-26-7], 0104
[127526-90-3], 2892 [135840-44-7], 2596 [155147-43-6], 3380 [159793-67-6], 1741 [174753-06-1], 4537
[128568-96-7], 2472 [139394-51-7], 1514 [155147-44-7], 3719 [160693-88-2], 2381 [174872-99-2], 3256
[128575-71-3], 3410 [140456-78-6], 4528 [155793-91-2], 2483 [161535-67-6], 1747 [180570-35-8], 0624
[129103-51-1], 3261 [140456-79-7], 4654 [155797-52-7], 3905 [161616-67-7], 3308 [180570-38-1], 0701
[129215-84-5], 3127 [144077-02-2], 3802 [155797-58-3], 3918 [164077-35-4], 3887 [187585-04-6], 0389
[129660-17-9], 1195 [144673-73-4], 3012 [156545-33-4], 2761 [164077-36-5], 3915
[130400-13-4], 1218 [148739-67-7], 2040 [157066-32-5], 2680 [165610-13-9], 3381
[132724-39-1], 4756 [153498-61-4], 1008 [157951-76-3], 0307 [167771-41-7], 0103
[133163-34-5], 4590 [153757-93-8], 0821 [158668-93-0], 2581 [168848-73-5], 3912

Serial numbers with no Registry number

0016	0334	0904	1591	2094	2539	3129	3507	3807
0017	0338	0911	1592	2101	2570	3130	3510	3810
0018	0367	0926	1643	2103	2571	3135	3513	3814
0024	0422	0939	1644	2111	2583	3144	3523	3819
0027	0428	0947	1663	2112	2603	3145	3524	3823
0030	0443	0962	1706	2117	2605	3153	3527	3849
0034	0471	0984	1744	2128	2625	3176	3538	3851
0036	0497	0994	1745	2146	2639	3185	3544	3854
0037	0504	1003	1746	2147	2683	3196	3547	3860
0039	0516	1006	1752	2162	2684	3218	3592	3866
0040	0527	1011	1761	2171	2707	3257	3593	3873
0041	0535	1016	1762	2175	2709	3260	3594	3886
0042	0543	1019	1763	2180	2721	3274	3595	3888
0043	0549	1021	1773	2188	2728	3281	3597	3892
0044	0566	1030	1774	2200	2804	3282	3601	3898
0045	0571	1048	1788	2203	2805	3283	3618	3904
0046	0580	1056	1796	2211	2807	3330	3619	3919
0054	0581	1068	1798	2221	2829	3334	3621	3929
0088	0601	1069	1802	2233	2833	3343	3622	3930
0097	0612	1077	1810	2247	2846	3344	3641	3932
0112	0613	1084	1815	2255	2856	3362	3669	3945
0113	0654	1088	1817	2257	2879	3367	3670	3950
0114	0668	1123	1823	2270	2883	3370	3681	3951
0117	0669	1211	1834	2281	2884	3374	3687	3966
0118	0674	1277	1838	2284	2890	3384	3700	3972
0126	0682	1300	1845	2285	2897	3388	3703	4012
0137	0688	1333	1850	2286	2904	3396	3704	4032
0151	0690	1337	1859	2304	2933	3413	3711	4046
0171	0691	1338	1900	2308	2963	3419	3713	4049
0178	0692	1354	1919	2309	2964	3426	3718	4050
0187	0707	1355	1938	2320	2965	3427	3731	4064
0197	0711	1369	1951	2323	2988	3431	3733	4065
0229	0713	1375	1980	2326	3017	3440	3737	4069
0237	0719	1390	1989	2341	3036	3444	3741	4071
0253	0764	1394	2007	2357	3075	3445	3745	4074
0264	0796	1395	2008	2358	3079	3446	3748	4089
0268	0804	1396	2027	2370	3091	3448	3759	4090
0272	0806	1420	2052	2386	3094	3450	3760	4091
0276	0835	1424	2066	2393	3096	3456	3763	4098
0277	0838	1445	2067	2402	3103	3457	3765	4142
0290	0889	1495	2068	2453	3104	3467	3775	4146
0297	0891	1510	2072	2454	3105	3471	3779	4147
0304	0893	1539	2083	2513	3112	3490	3793	4166
0325	0898	1540	2088	2519	3126	3503	3800	4169

4190	4361	4468	4551	4598	4634	4740	4781
4195	4396	4480	4555	4608	4653	4741	4787
4201	4409	4494	4558	4610	4663	4742	4792
4203	4433	4507	4571	4611	4667	4761	4859
4239	4449	4533	4572	4614	4683	4764	4881
4247	4452	4536	4591	4617	4702	4769	
4249	4454	4546	4593	4622	4735	4772	
4278	4455	4548	4594	4626	4738	4778	