

Astrophysics and Space Science Library 391

Boris V. Somov

Plasma Astrophysics, Part I

Fundamentals and Practice

Second Edition

AS
SL



Springer

Plasma Astrophysics, Part I

Astrophysics and Space Science Library

EDITORIAL BOARD

Chairman

W. B. BURTON, *National Radio Astronomy Observatory, Charlottesville, Virginia, U.S.A.*
(bburton@nrao.edu); *University of Leiden, The Netherlands*
(burton@strw.leidenuniv.nl)

F. BERTOLA, *University of Padua, Italy*

J. P. CASSINELLI, *University of Wisconsin, Madison, U.S.A.*

C. J. CESARSKY, *Commission for Atomic Energy, Saclay, France*

P. EHRENFREUND, *Leiden University, The Netherlands*

O. ENGVOLD, *University of Oslo, Norway*

A. HECK, *Strasbourg Astronomical Observatory, France*

E. P. J. VAN DEN HEUVEL, *University of Amsterdam, The Netherlands*

V. M. KASPI, *McGill University, Montreal, Canada*

J. M. E. KUIJPERS, *University of Nijmegen, The Netherlands*

H. VAN DER LAAN, *University of Utrecht, The Netherlands*

P. G. MURDIN, *Institute of Astronomy, Cambridge, UK*

F. PACINI, *Istituto Astronomia Arcetri, Firenze, Italy*

V. RADHAKRISHNAN, *Raman Research Institute, Bangalore, India*

B. V. SOMOV, *Astronomical Institute, Moscow State University, Russia*

R. A. SUNYAEV, *Space Research Institute, Moscow, Russia*

For further volumes:

<http://www.springer.com/series/5664>

Boris V. Somov

Plasma Astrophysics, Part I

Fundamentals and Practice

Second Edition



Springer

Boris V. Somov
Astronomical Institute and Faculty
of Physics
M.V. Lomonosov Moscow State University
Universitetskij Prospekt 13
119991 Moskva, Russia

ISSN 0067-0057

ISBN 978-1-4614-4282-0

ISBN 978-1-4614-4283-7 (eBook)

DOI 10.1007/978-1-4614-4283-7

Springer New York Heidelberg Dordrecht London

Library of Congress Control Number: 2012940742

© Springer Science+Business Media New York 2013

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed. Exempted from this legal reservation are brief excerpts in connection with reviews or scholarly analysis or material supplied specifically for the purpose of being entered and executed on a computer system, for exclusive use by the purchaser of the work. Duplication of this publication or parts thereof is permitted only under the provisions of the Copyright Law of the Publisher's location, in its current version, and permission for use must always be obtained from Springer. Permissions for use may be obtained through RightsLink at the Copyright Clearance Center. Violations are liable to prosecution under the respective Copyright Law.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

While the advice and information in this book are believed to be true and accurate at the date of publication, neither the authors nor the editors nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Cover illustration: Interaction of plasma with magnetic fields and light of stars creates many beautiful views of the night sky, like this one - a part of the famous nebula IC434 located about 1600 light-years away from Earth and observed by the National Science Foundation's 0.9-meter telescope on Kitt Peak.

Cover design: Photograph reproduced with kind permission from T.A. Rector (NOAO/AURA/NSF) and Hubble Heritage Team (STScI/AURA/NASA)

Printed on acid-free paper

Springer is part of Springer Science+Business Media (www.springer.com)



Sergey Ivanovich Syrovatskii (1925–1979)

*Dedicated to the memory of my teacher,
Professor Sergey Ivanovich Syrovatskii,
from whom I learnt much of what is in
this book and life.*

Preface to the Second Edition

The first edition, published in 2006, was addressed mainly to graduate students in astrophysics, space sciences and geophysics as well as to advanced students in applied physics and mathematics seeking a unified view of plasma physics and fluid mechanics. Lectures and seminars, given regularly since that time in the Faculty of Physics at the Moscow State University, have been found satisfactory by students, although new scientific topics, on which the students wanted to gain a view, have grown and multiplied. So, a considerable amount of new material, related to new space experiments that studied the Sun, interplanetary space, magnetosphere and distant astrophysical objects, should be added, at least briefly, to the lectures.

Solar physics gave wonderful examples of such news. Active regions at the Sun have been examined in extraordinary detail by space missions *Hinode*, *STEREO* and *Solar Dynamic Observatory (SDO)*. Much attention to solar plasma is conditioned by the possibility of the all-round observational test of theoretical models. Moreover, new observations have provided the important data that needed to answer many new questions being not answered by previous observations. One of them, having not only scientific but also a practical relevance, is how to predict large solar flares. They strongly influence the near-Earth space by large-amplitude shock waves and flows of particles accelerated to high energies.

It is well known that solar active regions can accumulate a huge energy before a large flare in the form of interacting magnetic fluxes in the solar corona. That is why the topological features of magnetic fields play a fundamental role and have to be studied. The text of the first edition was much complemented in the sections on the possibility to investigate the topological properties of the large-scale magnetic fields in active regions in order to recognize a pre-flare situation at the Sun. In particular, such an analysis of the field topology shows that the so-called “topological trigger” effect should be taken into account when we model large eruptive flares and try to predict them.

Another trend in modern solar physics was initiated by the *RHESSI* observations of hard-X-ray sources in the solar corona. Such emission interpreted as the bremsstrahlung of fast electrons was not predicted by theory because of very low density of coronal plasma. However its appearance during solar flares

clearly indicates generation of the so-called *super-hot* (with electron temperature $T_e \gtrsim 10$ keV) plasma and an additional acceleration of electrons in the vicinity of reconnecting current layers, i.e. the main source of flare energy.

In the second edition, a reader will find a summary and illustrations of the most principal observational features of the coronal HXR sources. Then it is considered the model in which the fast electrons are accelerated in collapsing magnetic traps created by magnetic reconnection in super-hot turbulent-current layers in the corona. The comprehensive analytical consideration is taken for joint action of the Fermi-type first-order mechanism and betatron acceleration of fast electrons. Efficient trapping and continuous acceleration of the fastest electrons in the collapsing traps can also produce a large flux of microwaves in solar flares.

Thus, new space-borne observations of the Sun from *Hinode*, *RHESSI*, *SDO* and *STEREO* have produced stunning results, invigorated solar research and challenged existing theoretical models.

Significant progress in the theory of magnetic reconnection has been achieved in the last years. Based on the results of numerical and laboratory experiments on magnetic reconnection, new analytical models have been suggested. They have generalized the famous classical models by Petschek and Syrovatskii. The magnetic field in these models was assumed to be potential in the exterior of a current configuration that included Syrovatskii's thin current layer and four MHD discontinuous flows attached to the edges of the current layer. The flow pattern near the current layer was not prescribed but determined from a self-consistent solution of the MHD problem in the approximation of a strong magnetic field. These new models allow to study the global structure of magnetic field near the reconnection region and its local properties in the vicinity of the current layer and attached discontinuities.

Interesting results have been obtained in the classical theory of the small-amplitude MHD waves propagating in a compressible optically thin medium whose radiative cooling depends on the temperature in the form of one or more sharp maxima. In astrophysical plasmas, these maxima are produced by the radiation of a small admixture of heavy ions. In the solar corona, the damping of MHD oscillations due to the radiative energy losses is present and turns out to be strong for slow magnetoacoustic waves. Where the rate of energy losses at a maximum (i.e. at $T \sim 10^5$ K), the brightness of the oscillating loops is at a maximum (i.e. in EUV) and, as a consequence, the MHD oscillations are damped more rapidly than in other places. That is why the rapidly damped oscillations are best seen in a small group of coronal loops observed in EUV radiation by satellites *TRACE* and *Hinode*.

New materials were added in almost every chapter of the second edition. For example, considering macroscopic description of astrophysical plasma, it was very useful to add the thirteen moment equations by Grad. They play a significant role in breaching the gap between hydrodynamics and kinetic theory in the problem of non-stationary powerful heat fluxes. In particular, it has appeared that the heat flux with consideration for the so-called "collisional relaxation" describes the heat transfer in solar flares much better than the classical Fourier law or anomalous heat fluxes.

References have been well updated but there will be no doubt that some important and relevant publications have been overlooked. This was not intentional. Plasma astrophysics is quickly developing in many astrophysical and related directions. So it is impossible to be aware of all the excellent papers published in these fields. I sincerely apologize for possible omissions in citations.

Moscow, Russia

Boris V. Somov

About This Book

If you want to learn the most fundamental things about plasma astrophysics with the least amount of time and effort—and who doesn't?—this text is for you. The book is addressed to young people, mainly to students, without a background in plasma physics; it grew from the lectures given many times in the Faculty of General and Applied Physics at the Moscow Institute of Physics and Technics (the well-known “fiz-tekh”) since 1977. A similar full-year course was also offered to the students of the Astronomical Division in the Faculty of Physics at the Moscow State University over the years after 1990. A considerable amount of new material, related to modern astrophysics, has been added to the lectures. So the contents of this book can hardly be presented during a one-year lecture course, without additional seminars.

In fact, just the seminars with the topics *how to make a cake* were especially pleasant for the author and useful for students. In part, the text of the book retains the imprint of the seminar form, implying a more lively dialogue with the reader and more visual representation of individual notions and statements. At the same time, the author's desire was that these digressions from the academic language of the monograph will not harm the rigour of presentation of this textbook's subject—the physical and mathematical introduction to plasma astrophysics.

There is no unique simple model of a plasma, which encompasses all situations in space. We have to familiarize ourselves with many different models applied to different situations. We need clear guidelines when a model works and when it does not work. Hence *the best strategy* is to develop an intuition about plasma physics, but how to develop it?

The idea of the book is not typical for the majority of textbooks on plasma astrophysics. Its idea is

the consecutive consideration of physical principles, starting from the most general ones, and of simplifying assumptions which give us a simpler description of plasma under cosmic conditions.

Thus I would recommend the students to read the book straight through each chapter to see the central line of the plasma astrophysics, its *classic fundamentals*. In so doing, the boundaries of the domain of applicability of the approximation at

hand will be outlined from the viewpoint of physics rather than of many possible astronomical applications. After that, as an aid to detailed understanding, please return with pencil and paper to work out the missing steps (if any) in the formal mathematics.

On the basis of such an approach the student interested in modern astrophysics, its *current practice*, will find the answers to two key questions:

1. What approximation is the best one (the simplest but sufficient) for description of a phenomenon in astrophysical plasma?
2. How to build an adequate model for the phenomenon, for example, a solar flare or a flare in the corona of an accretion disk?

Practice is really important for the theory of astrophysical plasma. Related exercises (problems and answers supplemented to each chapter) to improve skill do not thwart the theory but serve to better understanding of plasma astrophysics.

As for the applications, preference evidently is given to physical processes in the solar plasma. Why?—Much attention to solar plasma physics is conditioned by the possibility of the all-round observational test of theoretical models. This statement primarily relates to the processes in the solar atmosphere. For instance, flares on the Sun, in contrast to those on other stars as well as a lot of other analogous phenomena in the Universe, *can be seen* in their development, i.e. we can obtain a sequence of images during the flare's evolution, not only in the optical and radio ranges but also in the ultraviolet, soft and hard X-ray, gamma-ray ranges.

This book is mainly intended for students who have mastered a course of general physics and have some initial knowledge of theoretical physics. For beginning students, who may not know in which subfields of astrophysics they wish to specialize,

it is better to cover a lot of fundamental theories thoroughly than to dig deeply into any particular astrophysical subject or object,

even a very interesting one, for example black holes. Astronomers and astrophysicists of the future will need tools that allow them to explore in many different directions. Moreover astronomy of the future will be, more than hitherto, *precise science* similar to mathematics and physics.

The beginning graduate students are usually confronted with a confusing amount of work on plasma astrophysics published in a widely dispersed scientific literature. Knowing this difficulty, the author has tried as far as possible to represent the material in a self-contained form which does not require the reading of additional literature. However there is an extensive bibliography at the end of the book, allowing one to find the original works. In many cases, particularly where a paper in Russian is involved, the author has aimed to give the full bibliographic description of the work, including its title, etc.

Furthermore the book contains recommendations as to an introductory (unavoidable) reading needed to refresh the memory about a particular fact, as well as to additional (further) reading to refine one's understanding of the subject. Separate *remarks of an historical character* are included in many places. It is sometimes

simpler to explain the interrelation of discoveries by representing the subject in its development. It is the author's opinion that the outstanding discoveries in plasma astrophysics are by no means governed by chance. With the same thought in mind, the author gives preference to original papers on a topic under consideration; it happens in science, as in art, that an original is much better than nice-looking modernizations. Anyway,

knowledge of the history of science and especially of natural science is of great significance for its understanding and development.

The majority of the book's chapters begin from an "elementary account" and illustrative simple examples but finish with the most modern results of scientific importance. New problems determine the most interesting perspectives of plasma astrophysics as a new developing science. The author hopes, in this context, that professionals in the field of plasma astrophysics and adjacent sciences will enjoy reading this book too. Open issues are the focus of our attention in many places where they are. In this way, *perspectives of plasma astrophysics* with its many applications will be also of interest for readers. The book can be used as a textbook but has higher potential of modern scientific monograph.

The first volume of the book is unique in covering the basic principles and main practical tools required for understanding and work in plasma astrophysics. The second volume *Plasma Astrophysics. Part II. Reconnection and Flares* (Somov 2012, in the text as Part II) represents the basic physics of the magnetic reconnection phenomenon and the flares of electromagnetic origin in space plasmas in the solar system, relativistic objects, accretion disks, their coronae.

Acknowledgements

This textbook has grown out of lectures; and, first of all, I want to thank my students. During many years of my teaching at the Moscow Institute of Physics and Technics (the “fiz-tekh”) and later on at the Moscow State University, they were never silent if something was not or seemed to be not crystal clear immediately. The students forced me to write down this book; they read the chapters (included in a program of exams) as well as they suggested critical comments and corrections.

I am grateful to my colleagues: Ester Antonucci, Sergei I. Bezrodnykh, Sergei A. Bogachev, Cornelis de Jager, O.G. Den, Sergei V. Diakonov, Andre Duijveman, Pavel A. Gritsyk, Jean-Claude Hénoux, Hugh S. Hudson, Irina A. Kovalenko, Victor A. Kovalev, Takeo Kosugi, Pankaj Kumar, Leonid S. Ledentsov, Yuri E. Litvinenko, Marcos E. Machado, Satoshi Masuda, Sergei A. Markovskii, Elena Yu. Merenkova, Anna V. Oreshina, Inna V. Oreshina, Alexandr I. Podgornii, Olga V. Ptitsina, David M. Rust, Taro Sakao, Yuri I. Skrynnikov, Andrei R. Spektor, Abhishek Srivastava, Zdeněk Švestka, Igor P. Tindo, Vyacheslav S. Titov, Alexandr I. Verneta and Vladimir I. Vlasov for the pleasure of working together, for generous help and valuable remarks. I am also happy to acknowledge helpful discussions with many of my colleagues and friends in the world. There is no way to adequately express how much I owed to my wife, Irina Konstantinovna Somova, for her love and patience.

This work was supported by the Russian Foundation for Fundamental Research, grants Nos. 04-02-16125-a, 08-02-01033-a and 11-02-00843-a.

Moscow, Russia

Boris V. Somov

Plasma Astrophysics

History and Neighbours

Plasma astrophysics studies electromagnetic processes and phenomena in space, mainly the role of forces of an electromagnetic nature in the dynamics of cosmic matter. Two factors are specific to the latter: its gaseous state and high conductivity. Such a combination is unlikely to be found under natural conditions on Earth; the matter is either a non-conducting gas (the case of gas dynamics or hydrodynamics) or a liquid or a solid conductor. By contrast, *plasma is the main state of cosmic matter*. It is precisely the poor knowledge of cosmic phenomena and cosmic plasma properties that explains the retarded development of plasma astrophysics. It has been distinguished as an independent branch of physics in the pioneering works of Alfvén (see [Alfvén 1950](#)).

Soon after that, the problem of thermonuclear reactions initiated a great advance in plasma research ([Simon 1959](#); [Glasstone and Loveberg 1960](#); [Leontovich 1960](#)); see review of a current state in [Freidberg \(2007\)](#). This branch has been developing rather independently, although being partly “fed” by astrophysical ideas. They contributed to the growth of plasma physics, for example, the idea of stellarators. Presently, the reverse influence of laboratory plasma physics on astrophysics is also important.

From the physical viewpoint,

plasma astrophysics is a part of plasma theory related in the first place to the dynamics of a low-resistivity plasma in space.

However it is this part that is the most poorly studied one under laboratory conditions. During the 1930s, scientists began to realize that the Sun and other stars are powered by nuclear fusion and they began to think of re-creating the process in the laboratory. The ideas of astro- and geophysics dominate here, as before. At present time, they mainly come from many space experiments and fine astronomical ground-based observations. From this viewpoint, plasma astrophysics belongs to experimental science.

Electric currents and, therefore, magnetic fields are easily generated in the astrophysical plasma owing to its low resistivity. The energy of magnetic fields is accumulated in plasma, and the sudden release of this energy—an original

electrodynamical “burst” or “explosion”—takes place under definite but quite general conditions. It is accompanied by fast directed plasma ejections (jets), powerful flows of heat and radiation and impulsive acceleration of particles to high energies.

This phenomenon is quite a widespread one. It can be observed in flares on the Sun and other stars, in the Earth’s magnetosphere as magnetic storms and substorms, in coronae of accretion disks of cosmic X-ray sources, in nuclei of active galaxies and quasars. The second volume of this book, Part II, is devoted to the physics of *magnetic reconnection and flares* generated by reconnection in plasma in the solar system, single and double stars, relativistic objects, and other astrophysical objects.

The subject of the first volume of present book, Part I, is the systematic description of the most important topics of plasma astrophysics. However the aim of the book is not the strict substantiation of the main principals and basic equations of plasma physics; this can be found in many wonderful monographs ([Klimontovich 1986](#); [Schram 1991](#); [Liboff 2003](#); [Diamond et al. 2010](#)). There are also many nice textbooks ([Goldston and Rutherford 1995](#); [Choudhuri 1998](#); [Parks 2004](#)) to learn general plasma physics without or with some astrophysical applications.

The primary aim of the book in your hands is rather the solution of a much more modest but still important problem, namely to help the students of astrophysics to understand the interrelation and limits of applicability of different approximations which are used in plasma astrophysics. If, on his/her way, the reader will continuously try, following the author, to reproduce all mathematical transformation, he/she will soon find the pleasant feeling of real knowledge of the subject and the real desire for constructive work in plasma astrophysics.

The book will help the young reader to master the modern methods of plasma astrophysics and will teach the application of these methods while solving concrete problems in the physics of the Sun and many other astronomical objects. A good working knowledge of plasma astrophysics is essential for the modern astrophysicist.

Contents

Preface to the Second Edition	vii
About This Book	xi
Acknowledgements	xv
Plasma Astrophysics: History and Neighbours	xvii
1 Particles and Fields: Exact Self-consistent Description	1
1.1 Interacting Particles, Liouville's Theorem	1
1.1.1 Continuity in Phase Space	1
1.1.2 The Character of Particle Interactions	3
1.1.3 The Lorentz Force, Gravity	5
1.1.4 Collisional Friction in Plasma	6
1.1.5 The Exact Distribution Function	8
1.2 Charged Particles in the Electromagnetic Field	10
1.2.1 General Formulation of the Problem	10
1.2.2 The Continuity Equation for Electric Charge	11
1.2.3 Initial Equations and Initial Conditions	12
1.2.4 Astrophysical Plasma Applications	13
1.3 Gravitational Systems	13
1.4 Practice: Exercises and Answers	14
2 Statistical Description of Interacting Particle Systems	19
2.1 The Averaging of Liouville's Equation	19
2.1.1 Averaging Over Phase Space	19
2.1.2 Two Statistical Postulates	20
2.1.3 A Statistical Mechanism of Mixing in Phase Space	22
2.1.4 The Derivation of a General Kinetic Equation	25
2.2 A Collisional Integral and Correlation Functions	27
2.2.1 Binary Interactions	27
2.2.2 Binary Correlation	29
2.2.3 The Collisional Integral and Binary Correlation	30

2.3	Equations for Correlation Functions	32
2.4	Practice: Exercises and Answers	35
3	Weakly-Coupled Systems with Binary Collisions	37
3.1	Approximations for Binary Collisions	37
3.1.1	The Small Parameter of Kinetic Theory	37
3.1.2	The Vlasov Kinetic Equation	39
3.1.3	The Landau Collisional Integral	40
3.1.4	The Fokker-Planck Equation	42
3.2	Correlations and Debye-Hückel Shielding	44
3.2.1	The Maxwellian Distribution Function	44
3.2.2	The Averaged Force and Electric Neutrality	45
3.2.3	Pair Correlations and the Debye-Hückel Radius	46
3.2.4	Plasma Parameter and Collective Phenomena.....	49
3.3	Gravitational Systems	50
3.3.1	Vlasov-Type Equation	50
3.3.2	Collisional Integral and Other Models.....	51
3.4	Comments on Numerical Simulations.....	52
3.5	Practice: Exercises and Answers	53
4	Propagation of Fast Particles in Plasma	59
4.1	Derivation of the Basic Kinetic Equation	59
4.1.1	Basic Approximations	59
4.1.2	Dimensionless Kinetic Equation in Energy Space	61
4.2	A Kinetic Equation at High Speeds	63
4.3	The Classical Thick-Target Model	65
4.3.1	General Solution for Fast Electrons	65
4.3.2	The Injection Spectrum.....	66
4.3.3	Fast Electrons Inside a Target	67
4.3.4	Hard X-Ray Bremsstrahlung	68
4.4	The Role of Angular Diffusion	69
4.4.1	An Approximate Account of Scattering	69
4.4.2	The Thick-Target Model	70
4.5	The Reverse-Current Electric-Field Effect	72
4.5.1	The Necessity for a Beam-Neutralizing Current	72
4.5.2	Formulation of a Realistic Kinetic Problem	74
4.5.3	Dimensionless Parameters of the Problem	77
4.5.4	Coulomb Losses of Energy	78
4.5.5	New Physical Results.....	81
4.5.6	Plasma Heating by a Reverse Current	82
4.5.7	To the Future Models	83
4.6	Practice: Exercises and Answers	84
5	Motion of a Charged Particle in Given Fields	87
5.1	A Particle in Constant Homogeneous Fields	87
5.1.1	Relativistic Equation of Motion.....	87

5.1.2	Constant Homogeneous Non-magnetic Forces	88
5.1.3	Constant Homogeneous Magnetic Fields	89
5.1.4	Non-magnetic Force in a Magnetic Field	91
5.1.5	Electric and Gravitational Drifts	93
5.2	Weakly Inhomogeneous Slowly Changing Fields	94
5.2.1	Small Parameters in the Motion Equation	94
5.2.2	Expansion in Powers of m/e	96
5.2.3	The Averaging Over Gyromotion	99
5.2.4	Spiral Motion of the Guiding Center	101
5.3	Gradient and Inertial Drifts	102
5.3.1	Basic Formulae	102
5.3.2	The Centrifugal Drift	103
5.3.3	The Curvature-Dependent Drift	104
5.3.4	The Curvature-Independent Gradient Drift	105
5.4	Practice: Exercises and Answers	107
6	Adiabatic Invariants in Astrophysical Plasma	115
6.1	General Definitions	115
6.2	Two Main Invariants	116
6.2.1	Motion in the Larmor Plane	116
6.2.2	Magnetic Mirrors and Traps	118
6.2.3	Bounce Motion	120
6.2.4	The Fermi Acceleration	121
6.3	The Flux Invariant	123
6.4	Approximation Accuracy. Exact Solutions	124
6.5	Practice: Exercises and Answers	125
7	Wave-Particle Interaction in Astrophysical Plasma	129
7.1	The Basis of Kinetic Theory	129
7.1.1	The Linearized Vlasov Equation	129
7.1.2	The Landau Resonance and Landau Damping	131
7.1.3	Gyroresonance	134
7.2	Stochastic Acceleration of Particles by Waves	136
7.2.1	The Principles of Particle Acceleration by Waves	136
7.2.2	The Kolmogorov Theory of Turbulence	139
7.2.3	MHD Turbulent Cascading	141
7.3	The Relativistic Electron-Positron Plasma	142
7.4	Practice: Exercises and Answers	143
8	Coulomb Collisions in Astrophysical Plasma	147
8.1	Close and Distant Collisions	147
8.1.1	The Collision Parameters	147
8.1.2	The Rutherford Formula	149
8.1.3	The Test Particle Concept	149
8.1.4	Particles in a Magnetic Trap	150
8.1.5	The Role of Distant Collisions	151

8.2	Debye-Hückel Shielding	153
8.2.1	Simple Illustrations of the Shielding Effect	153
8.2.2	The Real Shielding Effect	156
8.2.3	Charge Neutrality and Oscillations in Plasma	156
8.3	Collisional Relaxations in Plasma	158
8.3.1	Some Exact Solutions	158
8.3.2	Plasma Heating by Accelerated Electrons	159
8.3.3	Plasma Heating by Magnetic Reconnection	164
8.3.4	An Adiabatic Model for Two-Temperature Plasma	166
8.3.5	Two-Temperature Accretion Flows	168
8.4	Dynamic Friction in Astrophysical Plasma	169
8.4.1	The Collisional Drag Force and Energy Losses	169
8.4.2	Electric Runaway	174
8.4.3	Thermal Runaway in Astrophysical Plasma	175
8.5	Practice: Exercises and Answers	177
9	Macroscopic Description of Astrophysical Plasma	181
9.1	Summary of Microscopic Description	181
9.2	Definition of Macroscopic Quantities	182
9.3	Macroscopic Transfer Equations	184
9.3.1	Equation for the Zeroth Moment	184
9.3.2	The Momentum Conservation Law	185
9.4	The Energy Conservation Law	188
9.4.1	The Second Moment Equation	188
9.4.2	The Case of Thermodynamic Equilibrium	190
9.4.3	The General Case of Non-equilibrium Plasma	191
9.4.4	The Sources and Sinks of Energy	192
9.5	General Properties of Transfer Equations	194
9.5.1	Divergent and Hydrodynamic Forms	194
9.5.2	Status of the Conservation Laws	195
9.6	Equation of State and Transfer Coefficients	197
9.6.1	General Approach to the Problem	197
9.6.2	The Classical Heat Conductive Flux	198
9.7	The Higher-Order Moment Equations	199
9.7.1	The 13 Moment Equations by Grad	199
9.7.2	The Powerful Heat Flux Problem	201
9.7.3	Large-Amplitude Heat Waves in Solar Flares	204
9.8	Gravitational Systems	206
9.9	Practice: Exercises and Answers	207
10	Multi-Fluid Models of Astrophysical Plasma	211
10.1	Multi-fluid Models in Astrophysics	211
10.2	Langmuir Waves	212
10.2.1	Langmuir Waves in a Cold Plasma	212
10.2.2	Langmuir Waves in a Warm Plasma	214
10.2.3	Ion Effects in Langmuir Waves	216

10.3	Electromagnetic Waves in Plasma	217
10.4	What Do We Miss?	219
10.5	Practice: Exercises and Answers	220
11	The Generalized Ohm's Law in Plasma	223
11.1	The Classic Ohm's Law	223
11.2	Derivation of Basic Equations	224
11.3	The General Solution	227
11.4	The Conductivity of Magnetized Plasma	228
11.4.1	Two Limiting Cases	228
11.4.2	The Physical Interpretation	229
11.5	Currents and Charges in Plasma	231
11.5.1	Collisional and Collisionless Plasmas	231
11.5.2	Volume Charge and Quasi-Neutrality	233
11.6	Practice: Exercises and Answers	235
12	Single-Fluid Models for Astrophysical Plasma	237
12.1	Derivation of the Single-Fluid Equations	237
12.1.1	The Continuity Equation	237
12.1.2	The Momentum Conservation Law in Plasma	238
12.1.3	The Energy Conservation Law	240
12.2	Basic Assumptions and the MHD Equations	242
12.2.1	Old Simplifying Assumptions	242
12.2.2	New Simplifying Assumptions	244
12.2.3	Non-relativistic Magnetohydrodynamics	247
12.2.4	The Energy Conservation Law	248
12.2.5	Relativistic Magnetohydrodynamics	250
12.3	Magnetic Flux Conservation. Ideal MHD	251
12.3.1	Integral and Differential Forms of the Law	251
12.3.2	The Equations of Ideal MHD	253
12.3.3	The "Frozen Field" Theorem	255
12.4	Magnetic Reconnection	257
12.4.1	General Properties	257
12.4.2	Reconnection in the Solar Wind	257
12.5	Practice: Exercises and Answers	260
13	Magnetohydrodynamics in Astrophysics	263
13.1	The Main Approximations in Ideal MHD	263
13.1.1	Dimensionless Equations	263
13.1.2	Weak Magnetic Fields in Astrophysical Plasma	265
13.1.3	Strong Magnetic Fields in Plasma	267
13.2	Accretion Disks of Stars	270
13.2.1	Angular Momentum Transfer in Binary Stars	270
13.2.2	Magnetic Accretion in Cataclysmic Variables	271
13.2.3	Accretion Disks Near Black Holes	272
13.2.4	Flares in Accretion Disk Coronae	274

13.3	Astrophysical Jets	275
13.3.1	Magnetic and Gravitomagnetic Forces	275
13.3.2	The Space Experiment “Gravity Probe B”	275
13.3.3	Relativistic Jets Near Black Holes	277
13.3.4	Relativistic Jets from Disk Coronae	278
13.4	Practice: Exercises and Answers	279
14	Plasma Flows in a Strong Magnetic Field	285
14.1	The General Formulation of a Problem	285
14.2	The Formalism of Two-Dimensional Problems	287
14.2.1	The First Type of Problems	288
14.2.2	The Second Type of MHD Problems	289
14.3	On the Existence of Continuous Flows	295
14.4	Flows in a Time-Dependent Dipole Field	296
14.4.1	Plane Magnetic Dipole Fields	296
14.4.2	Axial-Symmetric Dipole Fields in Plasma	299
14.5	Practice: Exercises and Answers	301
15	MHD Waves in Astrophysical Plasma	307
15.1	The Dispersion Equation in Ideal MHD	307
15.2	Small-Amplitude Waves in Ideal MHD	310
15.2.1	Entropy Waves	310
15.2.2	Alfvén Waves	311
15.2.3	Magnetoacoustic Waves	313
15.2.4	The Phase Velocity Diagram	314
15.3	Dissipation of Alfvén Waves	316
15.3.1	Small Damping of Alfvén Waves	316
15.3.2	Slightly Damped MHD Waves	318
15.4	Stability of Plasma-Compressing Waves	319
15.4.1	Derivation of the Dispersion Equation	319
15.4.2	Two Types of MHD Waves	322
15.4.3	Waves in Strong Magnetic Field	324
15.4.4	The Instability of Entropy Waves	325
15.4.5	The Damping of Magnetoacoustic Waves	328
15.5	MHD Oscillations in the Solar Corona	329
15.6	Practice: Exercises and Answers	331
16	Discontinuous Flows in a MHD Medium	333
16.1	Discontinuity Surfaces in Hydrodynamics	333
16.1.1	The Origin of Shocks in Ordinary Hydrodynamics	333
16.1.2	Boundary Conditions and Classification	334
16.1.3	Dissipative Processes and Entropy	336
16.2	Magnetohydrodynamic Discontinuities	337
16.2.1	Boundary Conditions at a Discontinuity Surface	337
16.2.2	Discontinuities Without Plasma Flows Across Them	340
16.2.3	Perpendicular Shock Wave	342

16.2.4	Oblique Shock Waves	345
16.2.5	Trans-Alfvénic Shock Waves	350
16.2.6	Peculiar Shock Waves	351
16.2.7	The Alfvén Discontinuity	352
16.3	Transitions Between Discontinuities	354
16.4	Shock Waves in Collisionless Plasma	355
16.5	Practice: Exercises and Answers	356
17	Evolutionarity of MHD Discontinuities	363
17.1	Conditions for Evolutionarity	363
17.1.1	The Physical Meaning and Definition	363
17.1.2	Linearized Boundary Conditions	365
17.1.3	The Number of Small-Amplitude Waves	367
17.1.4	Domains of Evolutionarity	370
17.2	Consequences of Evolutionarity Conditions	372
17.2.1	The Order of Wave Propagation	372
17.2.2	Continuous Transitions Between Discontinuities	373
17.3	Dissipative Effects in Evolutionarity	374
17.3.1	Dissipative and Non-dissipative Waves	374
17.3.2	Dissipation and Evolutionarity	375
17.4	The Structure of Discontinuity and Its Evolutionarity	377
17.4.1	Perpendicular Shock Waves	377
17.4.2	Discontinuities with Penetrating Magnetic Field	381
17.5	Practice: Exercises and Answers	384
18	Particle Acceleration by Shock Waves	387
18.1	Two Basic Mechanisms	387
18.2	Shock Diffusive Acceleration	388
18.2.1	The Canonical Model of Diffusive Mechanism	388
18.2.2	Some Properties of Diffusive Mechanism	391
18.2.3	Nonlinear Effects in Diffusive Acceleration	392
18.3	Shock Drift Acceleration	393
18.3.1	Perpendicular Shock Waves	393
18.3.2	Quasi-Perpendicular Shock Waves	395
18.3.3	Oblique Shock Waves	399
18.4	Practice: Exercises and Answers	400
19	Plasma Equilibrium in Magnetic Field	403
19.1	The Virial Theorem in MHD	403
19.1.1	A Brief Pre-history	403
19.1.2	Deduction of the Scalar Virial Theorem	404
19.1.3	Some Astrophysical Applications	407
19.2	Force-Free Fields and Shafranov's Theorem	410
19.2.1	The Simplest Examples of Force-Free Fields	410
19.2.2	The Energy of a Force-Free Field	412

19.3	Properties of Equilibrium Configurations	414
19.3.1	Magnetic Surfaces	414
19.3.2	The Specific Volume of a Magnetic Tube	415
19.3.3	The Flute or Convective Instability	418
19.3.4	Stability of an Equilibrium Configuration	419
19.4	The Archimedean Force in MHD	420
19.4.1	A General Formulation of the Problem	420
19.4.2	A Simplified Consideration of the Effect	421
19.4.3	MHD Equilibrium in the Solar Atmosphere	423
19.5	Grad-Shafranov Equation	424
19.6	Practice: Exercises and Answers	425
20	Stationary Flows in a Magnetic Field	429
20.1	Ideal Plasma Flows	429
20.1.1	Incompressible Medium	430
20.1.2	Compressible Medium	431
20.1.3	Astrophysical Collimated Streams (Jets)	431
20.1.4	MHD Waves of Arbitrary Amplitude	432
20.1.5	Differential Rotation and Isorotation	433
20.2	Flows at Small Magnetic Reynolds Numbers	436
20.2.1	Stationary Flows Inside a Duct	436
20.2.2	The MHD Generator or Pump	439
20.2.3	Weakly-Ionized Plasma in Astrophysics	440
20.3	The σ -Dependent Force and Vortex Flows	441
20.3.1	Simplifications and Problem Formulation	441
20.3.2	The Solution for a Spherical Ball	443
20.3.3	Forces and Flows Near a Spherical Ball	444
20.4	Large Magnetic Reynolds Numbers	448
20.4.1	The General Formula for the σ -Dependent Force	449
20.4.2	The σ -Dependent Force in Solar Prominences	452
20.5	Practice: Exercises and Answers	453
Appendix A	Notation	455
Appendix B	Useful Expressions	461
Appendix C	Constants	465
References		467
Index		487

Chapter 1

Particles and Fields: Exact Self-consistent Description

Abstract There exist two different ways to describe *exactly* the behavior of a system of charged particles in electromagnetic and gravitational fields. The first description, the Newton set of motion equations, is convenient for a small number of interacting particles. For systems of large numbers of particles, it is more advantageous to deal with the single Liouville equation for an *exact* distribution function.

1.1 Interacting Particles, Liouville's Theorem

1.1.1 Continuity in Phase Space

Let us consider a system of N interacting particles. Without much justification (which will be given in Chap. 2), let us introduce the distribution function

$$f = f(\mathbf{r}, \mathbf{v}, t) \quad (1.1)$$

for particles as follows. We consider the six-dimensional (6D) space called *phase space* $X = \{\mathbf{r}, \mathbf{v}\}$. The number of particles present in a small volume $dX = d^3\mathbf{r} d^3\mathbf{v}$ at a point X (see Fig. 1.1) at a moment of time t is defined to be

$$dN(X, t) = f(X, t) dX. \quad (1.2)$$

Accordingly, the total number of the particles at this moment is

$$N(t) = \int f(X, t) dX \equiv \iint f(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{r} d^3\mathbf{v}. \quad (1.3)$$

If, for definiteness, we use the Cartesian coordinates, then

$$X = \{x, y, z, v_x, v_y, v_z\}$$

Fig. 1.1 The 6D phase space X . A small volume dX at a point X





Fig. 1.2 The 6D phase space X . The volume U is enclosed by the surface S . $\mathbf{J}$ is the particle flux density



is a point of the phase space (Fig. 1.2) and

$$\dot{X} = \{v_x, v_y, v_z, \dot{v}_x, \dot{v}_y, \dot{v}_z\} \quad (1.4)$$

denotes the velocity of this point in the phase space. If we watch a particle at a point X then $\{\dot{v}_x, \dot{v}_y, \dot{v}_z\}$ represents the acceleration of its motion under action on this particle from all other particles.

Let us suppose, for a while, that the coordinates and velocities of the particles are changing *continuously*—“from point to point”, i.e. the particles move smoothly at all times. So the distribution function $f(X, t)$ is differentiable. Moreover we shall assume that this continuous motion of the particles in phase space can be expressed by the *continuity equation*

$$\boxed{\frac{\partial f}{\partial t} + \text{div}_X f \dot{X} = 0} \quad (1.5)$$

or

$$\frac{\partial f}{\partial t} + \operatorname{div}_{\mathbf{r}} f \mathbf{v} + \operatorname{div}_{\mathbf{v}} f \dot{\mathbf{v}} = 0.$$

Equation (1.5) expresses the *conservation law* in a divergent form for the particles, since the integration of (1.5) over a volume U enclosed by the surface S in Fig. 1.2 gives

$$\frac{\partial}{\partial t} \int_U f dX + \int_U \operatorname{div}_X f \dot{X} dX =$$

by virtue of definition (1.2) and the Ostrogradskii-Gauss theorem

$$= \frac{\partial}{\partial t} N(t) \Big|_U + \int_S f \dot{X} dS = \frac{\partial}{\partial t} N(t) \Big|_U + \int_S \mathbf{J} \cdot d\mathbf{S} = 0.$$

Here a surface element $d\mathbf{S}$, normal to the boundary S , is oriented towards its outside, so that imports are counted as negative (e.g., [Smirnov 1965](#), Sect. 126).

$$\mathbf{J} = f \dot{X} \tag{1.6}$$

is the *particle flux density* in phase space. So

$$\frac{\partial}{\partial t} N(t) \Big|_U = - \int_S \mathbf{J} \cdot d\mathbf{S}.$$

(1.7)

Thus

a change of the particle number in a given volume U of the phase space X is defined by the particle flux through the boundary surface S only.

The reason is clear. There are no sources or sinks for the particles inside the volume. Otherwise the source and sink terms must be added to the right-hand side of (1.5).

1.1.2 The Character of Particle Interactions

Let us rewrite the continuity equation (1.5) in another form in order to understand the meaning of divergent terms. The first of them is

$$\operatorname{div}_{\mathbf{r}} f \mathbf{v} = f \operatorname{div}_{\mathbf{r}} \mathbf{v} + (\mathbf{v} \cdot \nabla_{\mathbf{r}}) f = 0 + (\mathbf{v} \cdot \nabla_{\mathbf{r}}) f,$$

since $\mathbf{r}$ and $\mathbf{v}$ are independent variables in the phase space X . The second divergent term is

$$\operatorname{div}_{\mathbf{v}} f \dot{\mathbf{v}} = f \operatorname{div}_{\mathbf{v}} \dot{\mathbf{v}} + \dot{\mathbf{v}} \cdot \nabla_{\mathbf{v}} f .$$

So far no assumption has been made as to the character of particle interactions. It is worth doing here. Let us restrict our consideration to the interactions with

$$\boxed{\operatorname{div}_{\mathbf{v}} \dot{\mathbf{v}} = 0 ,} \quad (1.8)$$

then (1.5) can be rewritten in the following form:

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} f + \frac{\mathbf{F}}{m} \cdot \nabla_{\mathbf{v}} f = 0$$

or

$$\frac{\partial f}{\partial t} + \dot{X} \cdot \nabla_X f = 0 , \quad (1.9)$$

where

$$\dot{X} = \left\{ v_x, v_y, v_z, \frac{F_x}{m}, \frac{F_y}{m}, \frac{F_z}{m} \right\} . \quad (1.10)$$

Having written that, we “trace” the particle *phase trajectories* when the particles move under action of a force field $\mathbf{F}(\mathbf{r}, \mathbf{v}, t)$.

Thus we have found Liouville’s theorem in the following formulation:

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} f + \frac{\mathbf{F}}{m} \cdot \nabla_{\mathbf{v}} f = 0 . \quad (1.11)$$

| Liouville’s theorem: *The distribution function remains constant on the particle phase trajectories* if condition (1.8) is satisfied.

We shall call (1.11) the *Liouville equation*. The first term in the Liouville equation (1.11), the *partial* time derivative $\partial f / \partial t$, characterizes a change of the distribution function $f(t, X)$ at a given point X in the phase space with time t .

Let us define also the Liouville operator

$$\frac{D}{Dt} \equiv \frac{\partial}{\partial t} + \dot{X} \cdot \frac{\partial}{\partial X} \equiv \frac{\partial}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} + \frac{\mathbf{F}}{m} \cdot \nabla_{\mathbf{v}} . \quad (1.12)$$

This operator is just the *total* time derivative following a particle motion in the phase space X . By using definition (1.12), we rewrite Liouville’s theorem as follows:




Fig. 1.3 Action of the two different terms of the Liouville operator in the 6D phase space X . (a) A simple kinematic effect. (b) Particle acceleration

$$\boxed{\frac{Df}{Dt} = 0.} \quad (1.13)$$

What factors do lead to the changes in the distribution function?—Let dX be a small volume in the phase space X . The second term in (1.11), $\mathbf{v} \cdot \nabla_r f$, means that the particles come into and go out of the phase volume element considered, because their velocities are not zero (Fig. 1.3a). So this term describes a simple kinematic effect. If the distribution function f has a non-zero gradient over $\mathbf{r}$, then a number of particles inside the volume dX changes because they move with velocity $\mathbf{v}$.

The third term, $(\mathbf{F}/m) \cdot \nabla_v f$, means that the particles escape from the phase volume element dX or come into this element due to their *acceleration* (see Fig. 1.3b) or deceleration (as illustrated in Sect. 1.1.4) under the influence of the force field $\mathbf{F}$.

Some important properties of the Liouville equation are considered in Exercises 1.1–1.4.

1.1.3 The Lorentz Force, Gravity

Let us recall that, in order that the Liouville theorem can be valid, the force field $\mathbf{F}$ has to satisfy condition (1.8). We rewrite it as follows:

$$\frac{\partial \dot{v}_\alpha}{\partial v_\alpha} = \frac{1}{m} \frac{\partial F_\alpha}{\partial v_\alpha} = 0$$

or

$$\frac{\partial F_\alpha}{\partial v_\alpha} = 0. \quad (1.14)$$

Here the index $\alpha = 1, 2, 3$. In particular, this condition holds if

the component F_α of the force vector $\mathbf{F}$ does not depend upon the velocity component v_α .

This is a sufficient condition, of course.

The classical Lorentz force

$$F_\alpha = e \left[E_\alpha + \frac{1}{c} (\mathbf{v} \times \mathbf{B})_\alpha \right] \quad (1.15)$$

obviously has that property. The gravitational force in the classical approximation is entirely independent of velocity.

Other forces may be considered, depending on the situation, for example the force resulting from the emission of radiation (the *radiation reaction*) and/or absorption of radiation by astrophysical plasma, which is electromagnetic in nature, though maybe quantum. These forces when they are important should be considered with account of their relative significance, conservative or *dissipative* character (see Sect. 1.1.4), and other physical properties taken.

1.1.4 Collisional Friction in Plasma

As a contrary example we consider the friction force (cf. formula (8.71) for the collisional drag force which acts on a particle moving with velocity $\mathbf{v}$ in plasma):

$$\mathbf{F} = -k \mathbf{v}, \quad (1.16)$$

where the constant $k > 0$. In this case the right-hand side of Liouville's equation (1.11) is not zero:

$$-f \operatorname{div}_{\mathbf{v}} \dot{\mathbf{v}} = -f \operatorname{div}_{\mathbf{v}} \frac{\mathbf{F}}{m} = \frac{3k}{m} f,$$


because

$$\frac{\partial v_\alpha}{\partial v_\alpha} = \delta_{\alpha\alpha} = 3.$$

Instead of Liouville's equation we have

$$\frac{Df}{Dt} = \frac{3k}{m} f > 0. \quad (1.17)$$

Fig. 1.4 Particle density increases in the phase space in the vicinity of the $\mathbf{r}$ -axis as a result of the action of the friction force $\mathbf{F}$



Therefore the distribution function (that is the particle density) does not remain constant on particle trajectories but increases as the time elapses. Along the phase trajectories, it increases exponentially:

$$f(t, \mathbf{r}, \mathbf{v}) \sim f(0, \mathbf{r}, \mathbf{v}) \exp\left(\frac{3k}{m} t\right). \quad (1.18)$$

The physical sense of this phenomenon is obvious. As the particles are decelerated by the friction force, they go down in the schematic Fig. 1.4. By so doing, they are concentrated in a diminishing region of the phase space situated in the vicinity of the axis $\mathbf{v} = 0$.


There is a viewpoint that the Liouville theorem is valid for the forces that *do not disperse* particle velocities (Shkarofsky et al. 1966, Chap. 2). Why?—It is usually implied that particle *collisions* enlarge such a dispersion: $\text{div}_{\mathbf{v}} \dot{\mathbf{v}} > 0$. So

$$\frac{Df}{Dt} = \left(\frac{\partial f}{\partial t}\right)_c = -f \text{div}_{\mathbf{v}} \dot{\mathbf{v}} < 0. \quad (1.19)$$

In this case the right-hand side of (1.19) is called the *collisional* integral (see Sects. 2.1 and 2.2). In contrast to the right-hand side of (1.17), that of (1.19) is usually negative.

The above example of the friction force is instructive in that it shows how the forces that are diminishing the velocity dispersion ($\text{div}_{\mathbf{v}} \dot{\mathbf{v}} < 0$) lead to the violation of Liouville's theorem; in other words, how they lead to a change of the distribution function along the particle trajectories. For validity of Liouville's theorem only the condition (1.8) is important; in the velocity space, the divergence of the forces has to equal zero. The sign of this divergence is unimportant.

Fig. 1.5 The one-dimensional analogy of the exact distribution function



1.1.5 The Exact Distribution Function

Let us consider another property of the Liouville theorem. We introduce the N -particle distribution function of the form

$$\hat{f}(t, \mathbf{r}, \mathbf{v}) = \sum_{i=1}^N \delta(\mathbf{r} - \mathbf{r}_i(t)) \delta(\mathbf{v} - \mathbf{v}_i(t)). \quad (1.20)$$

The delta function of the vector-argument is defined as usually (see [Vladimirov 1971](#)):

$$\delta(\mathbf{r} - \mathbf{r}_i(t)) = \prod_{\alpha=1}^3 \delta_{\alpha} = \delta(r_x - r_x^i(t)) \delta(r_y - r_y^i(t)) \delta(r_z - r_z^i(t)). \quad (1.21)$$

We shall call function (1.20) the *exact* distribution function. It is illustrated by schematic Fig. 1.5.

Let us substitute the exact distribution function in the Liouville equation (1.11). The resulting three terms are

$$\begin{aligned} \frac{\partial \hat{f}}{\partial t} &= \sum_i \delta'_{\alpha}(\mathbf{r} - \mathbf{r}_i(t)) (-1) \dot{r}_{\alpha}^i \delta(\mathbf{v} - \mathbf{v}_i(t)) + \\ &+ \sum_i \delta(\mathbf{r} - \mathbf{r}_i(t)) \delta'_{\alpha}(\mathbf{v} - \mathbf{v}_i(t)) (-1) \dot{v}_{\alpha}^i, \end{aligned} \quad (1.22)$$

$$\mathbf{v} \cdot \nabla_{\mathbf{r}} \hat{f} \equiv v_{\alpha} \frac{\partial \hat{f}}{\partial r_{\alpha}} = \sum_i v_{\alpha} \delta'_{\alpha}(\mathbf{r} - \mathbf{r}_i(t)) \delta(\mathbf{v} - \mathbf{v}_i(t)), \quad (1.23)$$

$$\frac{\mathbf{F}}{m} \cdot \nabla_{\mathbf{v}} \hat{f} \equiv \frac{F_{\alpha}}{m} \frac{\partial \hat{f}}{\partial v_{\alpha}} = \sum_i \frac{F_{\alpha}}{m_i} \delta(\mathbf{r} - \mathbf{r}_i(t)) \delta'_{\alpha}(\mathbf{v} - \mathbf{v}_i(t)). \quad (1.24)$$

Here the index $\alpha = 1, 2, 3$ or (x, y, z) . The prime denotes the derivative with respect to the argument of a function; for the delta function, see definition of the derivative

in [Vladimirov \(1971\)](#), Chap. 2. The overdot denotes differentiation with respect to time t . Summation over the repeated index α (contraction) is implied:

$$\delta'_\alpha \dot{r}_\alpha^i = \delta'_x \dot{r}_x^i + \delta'_y \dot{r}_y^i + \delta'_z \dot{r}_z^i.$$

The sum of terms (1.22)–(1.24) equals zero. Let us rewrite it as follows

$$\begin{aligned} 0 = & \sum_i (-\dot{r}_\alpha^i + v_\alpha^i) \delta'_\alpha (\mathbf{r} - \mathbf{r}_i(t)) \delta (\mathbf{v} - \mathbf{v}_i(t)) + \\ & + \sum_i \left(-\dot{v}_\alpha^i + \frac{F_\alpha}{m_i} \right) \delta (\mathbf{r} - \mathbf{r}_i(t)) \delta'_\alpha (\mathbf{v} - \mathbf{v}_i(t)). \end{aligned}$$

This can occur just then that all the coefficients of different combinations of delta functions with their derivatives equal zero as well. Therefore we find

$$\frac{dr_\alpha^i}{dt} = v_\alpha^i(t), \quad \frac{dv_\alpha^i}{dt} = \frac{1}{m_i} F_\alpha (\mathbf{r}_i(t), \mathbf{v}_i(t)). \quad (1.25)$$

Thus

the Liouville equation for an exact distribution function is *equivalent* to the Newton set of equations for a particle motion.

Both methods describe a purely dynamic behavior of the particles. It is natural since this distribution function is exact. No statistical averaging has been done so far. It is for this reason that both descriptions—namely, the Newton set and the Liouville theorem for the exact distribution function—are *dynamic* (as well as *reversible*, of course) and equivalent. Statistics will appear in the next chapter when, instead of the exact description of a system, we begin to use some mean characteristics such as temperature, density etc. This is the statistical description that is valid for systems containing a large number of particles.

We have shown that finding a solution of the Liouville equation for an exact distribution function

$$\boxed{\frac{D\hat{f}}{Dt} = 0} \quad (1.26)$$

is the same as the integration of the motion equations. However

for systems of a large number of interacting particles, it is much more advantageous to deal with the single Liouville equation for the exact distribution function which describes the entire system.

Recommended Reading: [Landau and Lifshitz \(1976\)](#), Chaps. 2 and 7; [Landau and Lifshitz \(1959b\)](#), Chap. 1, Sects. 1–3.

1.2 Charged Particles in the Electromagnetic Field

1.2.1 General Formulation of the Problem

Let us start from recalling the basic physics notations and establishing a common basis. Maxwell's equations for the electric field $\mathbf{E}$ and magnetic field $\mathbf{B}$ are well known to have the form (see [Landau and Lifshitz 1975](#), Chap. 4, Sect. 26):

$$\text{rot } \mathbf{B} = \frac{4\pi}{c} \mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t}, \quad (1.27)$$

$$\text{rot } \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}, \quad (1.28)$$

$$\text{div } \mathbf{B} = 0, \quad (1.29)$$

$$\text{div } \mathbf{E} = 4\pi\rho^q. \quad (1.30)$$

Here $\mathbf{B}$ and $\mathbf{E}$ are the exact magnetic and electric fields. The fields are completely determined by electric charges and electric currents.

Note that, in general, Maxwell's equations imply the continuity equation for electric charge (see Exercise 1.5) as well as the conservation law for electromagnetic field energy (Exercise 1.6).

Let there be N particles with charges $e_1, e_2, \dots, e_i, \dots, e_N$, coordinates $\mathbf{r}_i(t)$ and velocities $\mathbf{v}_i(t)$, see Fig. 1.6. By definition, the electric charge density

$$\rho^q(\mathbf{r}, t) = \sum_{i=1}^N e_i \delta(\mathbf{r} - \mathbf{r}_i(t)) \quad (1.31)$$

and the density of electric current

$$\mathbf{j}(\mathbf{r}, t) = \sum_{i=1}^N e_i \mathbf{v}_i(t) \delta(\mathbf{r} - \mathbf{r}_i(t)). \quad (1.32)$$




Fig. 1.6 A system of N charged particles

The coordinates and velocities of particles can be found by integrating the equations of motion—the Newton equations:

$$\dot{\mathbf{r}}_i \equiv \frac{d\mathbf{r}_i}{dt} = \mathbf{v}_i(t), \quad (1.33)$$

$$\dot{\mathbf{v}}_i \equiv \frac{d\mathbf{v}_i}{dt} = \frac{1}{m_i} e_i \left[\mathbf{E}(\mathbf{r}_i(t)) + \frac{1}{c} \mathbf{v}_i \times \mathbf{B}(\mathbf{r}_i(t)) \right]. \quad (1.34)$$

Let us count the number of unknown quantities: the vectors $\mathbf{B}$, $\mathbf{E}$, $\mathbf{r}_i$, and $\mathbf{v}_i$. We obtain: $3 + 3 + 3N + 3N = 6(N + 1)$. The number of equations is equal to $8 + 6N = 6(N + 1) + 2$. Therefore two equations seem to be unnecessary. Why is this so?

1.2.2 The Continuity Equation for Electric Charge

At first let us make sure that the definitions (1.31) and (1.32) conform to the conservation law for electric charge. Differentiating (1.31) with respect to time gives (see Exercise 1.7)

$$\frac{\partial \rho^q}{\partial t} = \sum_i e_i \delta'_\alpha (-1) \dot{r}_\alpha^i. \quad (1.35)$$

Here the index $\alpha = 1, 2, 3$. The prime denotes the derivative with respect to the argument of the delta function, see Vladimirov (1971). The overdot denotes differentiation with respect to time t .

For the electric current density (1.32) we have the divergence

$$\text{div } \mathbf{j} = \frac{\partial}{\partial r_\alpha} j_\alpha = \sum_i e_i v_\alpha^i \delta'_\alpha. \quad (1.36)$$

Comparing formula (1.35) with (1.36) we see that

$$\frac{\partial \rho^q}{\partial t} + \text{div } \mathbf{j} = 0.$$

(1.37)

Therefore the definitions for ρ^q and $\mathbf{j}$ conform to the *continuity equation*.

As we shall see it in Exercise 1.5, conservation of electric charge follows also directly from the Maxwell equations (1.27) and (1.30). The difference is that above we have not used (1.30). This is essential for the next section.

1.2.3 Initial Equations and Initial Conditions

Operating with the divergence on (1.27) and using the continuity equation (1.37), we obtain

$$0 = \frac{4\pi}{c} \left(-\frac{\partial \rho^q}{\partial t} \right) + \frac{1}{c} \frac{\partial}{\partial t} \operatorname{div} \mathbf{E}.$$

Thus, by postulating the definitions (1.31) and (1.32), by virtue of the continuity equation (1.37) and without using the Maxwell equation (1.30), we find that

$$\frac{\partial}{\partial t} (\operatorname{div} \mathbf{E} - 4\pi \rho^q) = 0. \quad (1.38)$$

Hence (1.30) will be valid at any moment of time, provided it is true at the initial moment.

Let us operate with the divergence on (1.28):

$$\frac{\partial}{\partial t} \operatorname{div} \mathbf{B} = 0. \quad (1.39)$$

Equation (1.29) implies the absence of magnetic charges or, which is the same, the *solenoidal* character of the magnetic field.

We come to the conclusion that the scalar equations (1.29) and (1.30) play the role of *initial conditions* for the time-dependent vector equations

$$\frac{\partial}{\partial t} \mathbf{B} = -c \operatorname{rot} \mathbf{E} \quad (1.40)$$

and

$$\frac{\partial}{\partial t} \mathbf{E} = +c \operatorname{rot} \mathbf{B} - 4\pi \mathbf{j}. \quad (1.41)$$

Thus, in order to describe the gas consisting of N charged particles, we consider the time-dependent problem of N bodies with a given interaction law.

The electromagnetic part of the interaction is described by Maxwell's equations, the time-independent scalar equations playing the role of initial conditions for the time-dependent problem.

Therefore the set consisting of eight Maxwell's equations and $6N$ Newton's equations is neither over- nor under-determined. It is *closed* with respect to the time-dependent problem, i.e. it consists of $6(N + 1)$ equations for $6(N + 1)$ unknown variables, once the initial and boundary conditions are given.

1.2.4 Astrophysical Plasma Applications

The set of equations described above can be treated analytically in just three cases:

1. $N = 1$, the motion of a charged particle in a given electromagnetic field, for example, drift motions and the so-called adiabatic invariants, wave-particle interaction and the problem of particle acceleration in astrophysical plasma; e.g., Chaps. 7 and 8.
2. $N = 2$, Coulomb collisions of two charged particles, i.e. binary collisions. This is important for the kinetic description of physical processes, for example, the kinetic effects under propagation of accelerated particles in plasma, collisional heating of plasma by a beam of fast electrons or/and ions, see Chaps. 4 and 8.
3. $N \rightarrow \infty$, a very large number of particles. This case is the frequently considered one in plasma astrophysics, because it allows us to introduce macroscopic descriptions of plasma, the widely-used magnetohydrodynamic (MHD) approximation; Chaps. 9 and 12.

Numerical integration of (1.27)–(1.34) in the case of large but finite N , like $N \approx 3 \times 10^6$, is possible by using powerful modern computers. Such computations called *particle simulations* have proved to be increasingly useful for understanding many properties of astrophysical plasma and for demonstration of them. One important example of a simulation is *magnetic reconnection* in a collisionless plasma (Horiuchi and Sato 1994; Cai and Lee 1997). This process often leads to fast energy conversion from field energy to particle energy, flares in astrophysical plasma (see Part II).

Note also that the set of equations described above can be well generalized to include consideration of neutral particles. This is necessary, for instance, in the study of the generalized Ohm's law (Chap. 9) which can be applied in the investigation of physical processes in *weakly-ionized* plasmas, for example in the solar photosphere and prominences (see Part II).

Dusty and *self-gravitational* plasmas in space are interesting in view of the diverse and often surprising facts about planetary rings and comet environments, interstellar dark space (Bliokh et al. 1995; Kikuchi 2001; Fernández 2005). Two effects are often of basic importance, gravitational and electric, since charged or polarized *dust grains* involved in such environments are much heavier than electrons and ions. So a variety of electric rather than magnetic phenomena are taking place predominantly; and gravitational forces acting on dust particles can become appreciable.

1.3 Gravitational Systems

Gravity plays a central role in the dynamics of many astrophysical systems—from stars to the Universe as a whole (Lahav et al. 1996; Rose 1998; Bertin 1999; Dadhich

and Kembhavi 2000). It is important for many astrophysical applications that a *gravitational* force acts on the particles as follows:

$$m_i \dot{\mathbf{v}}_i = -m_i \nabla \phi. \quad (1.42)$$

Here the gravitational potential

$$\phi(t, \mathbf{r}) = - \sum_{n=1}^N \frac{G m_n}{|\mathbf{r}_n(t) - \mathbf{r}|}, \quad n \neq i, \quad (1.43)$$

G is the gravitational constant. We shall return to this subject many times, for example, while studying the *virial theorem* in MHD (Chap. 19). This theorem is widely used in astrophysics.

At first sight, it may seem that a gravitational system, like stars in a galaxy, will be easier to study than a plasma, because there is gravitational charge (i.e. mass) of only one sign compared to the electric charges of two opposite signs. However the reality is the other way round. Though the potential (1.43) of the gravitational interaction looks similar to the Coulomb potential of charged particles (see formula (8.1)),

physical properties of gravitational systems differ so much from properties of astrophysical plasma.

We shall see this fundamental difference, for example, in Sect. 3.3 and many times in what follows. A deep unifying theme which underlies many astrophysical results is that self-gravity is incompatible with thermodynamic equilibrium (see Sect. 9.8).

1.4 Practice: Exercises and Answers

Exercise 1.1 (Sect. 1.1.2). Show that

any distribution function that is a function of the constants of motion—the *invariants* of motion or *integrals* of motion—satisfies Liouville's equation (1.13).

Answer. A general solution of the equations of motion (1.25) depends on $6N$ constants C_i where index $i = 1, 2, \dots, 6N$ (e.g., Landau and Lifshitz 1976, Chap. 2, Sect. 6). If we assume that the distribution function is a function of these constants of the motion

$$f = f(C_1, \dots, C_i, \dots, C_{6N}), \quad (1.44)$$

we can rewrite the left-hand side of (1.13) as

$$\frac{Df}{Dt} = \sum_{i=1}^{6N} \left(\frac{DC_i}{Dt} \right) \left(\frac{\partial f}{\partial C_i} \right). \quad (1.45)$$

Because C_i are constants of the motion, $DC_i/Dt = 0$. Therefore the right-hand side of (1.45) is also zero, and the distribution function (1.44) satisfies the Liouville equation. This is the so-called *Jeans theorem*. It will be used, for example, in the theory of wave-particle interaction in astrophysical plasma (Sect. 7.1).

Exercise 1.2 (Sect. 1.1.2). Rewrite the Liouville theorem by using the Hamilton equations instead of the Newton equations.

Answer. Rewrite the Newton set of the motion equations (1.25) in the Hamilton form (see Landau and Lifshitz 1976, Chap. 7, Sect. 40):

$$\dot{q}_s = \frac{\partial H}{\partial P_s}, \quad \dot{P}_\alpha = -\frac{\partial H}{\partial q_s}. \quad (1.46)$$

Here index $s = 1, 2, 3, \dots, 3N$, $H(P, q)$ is the Hamiltonian of the system under consideration, q_α and P_α are the *generalized* coordinates and momenta, respectively.

Let us substitute the variables $\mathbf{r}$ and $\mathbf{v}$ in the Liouville equation (1.11) by the generalized variables $\mathbf{q}$ and $\mathbf{P}$. By doing so and using (1.46), we obtain the following form of the Liouville equation

$$\frac{\partial f}{\partial t} + \nabla_{\mathbf{P}} H \cdot \nabla_{\mathbf{q}} f - \nabla_{\mathbf{q}} H \cdot \nabla_{\mathbf{P}} f = 0. \quad (1.47)$$

Therefore, from a mathematical point of view,

the Liouville equation represents the linear differential equation in partial derivatives of the first order, for which the *characteristic equations* coincide with the Hamilton equations.

Because of symmetry of (1.47), it is convenient here to use the *Poisson brackets* (see Landau and Lifshitz 1976, Chap. 7, Sect. 42). Recall that the Poisson brackets for arbitrary quantities A and B are defined to be

$$[A, B] = \sum_{s=1}^{3N} \left(\frac{\partial A}{\partial q_s} \frac{\partial B}{\partial P_s} - \frac{\partial A}{\partial P_s} \frac{\partial B}{\partial q_s} \right). \quad (1.48)$$

Applying definition (1.48) to (1.47), we find the final form of the Liouville theorem

$$\frac{\partial f}{\partial t} + [f, H] = 0.$$

(1.49)

Q.e.d. Note that for a system in equilibrium

$$[f, H] = 0. \quad (1.50)$$

Exercise 1.3 (Sect. 1.1.2). Discuss what to do with the Liouville theorem, if it is impossible to disregard *quantum indeterminacy* and assume that the classical description of a system is justified. Consider the case of dense fluids inside stars, for example, white dwarfs.

Comment. Inside a *white dwarf* star the temperature $T \sim 10^5$ K, but the density is very high: $n \sim 10^{28} - 10^{30} \text{ cm}^{-3}$ (e.g., [de Martino et al. 2003](#)). The electrons cannot be regarded as classical particles. We have to consider them as a quantum system with a Fermi-Dirac distribution (see Sect. 57 in [Landau and Lifshitz 1959b](#); [Kittel 1995](#)).

Exercise 1.4 (Sect. 1.1.2). Recall the Liouville theorem in a course of mechanics—the phase volume of a system is independent of t , i.e. it is the invariant of motion (e.g., [Landau and Lifshitz 1976](#), Chap. 7, Sect. 46). Show that this formulation is equivalent to (1.13).

Exercise 1.5 (Sect. 1.2.1). Show that Maxwell's equations imply the continuity equation for the electric charge.

Answer. Operating with the divergence on (1.27), we have

$$0 = \frac{4\pi}{c} \operatorname{div} \mathbf{j} + \frac{1}{c} \frac{\partial}{\partial t} \operatorname{div} \mathbf{E}.$$

Substituting (1.30) in this equation gives us the continuity equation for the electric charge

$$\frac{\partial}{\partial t} \rho^q + \operatorname{div} \mathbf{j} = 0. \quad (1.51)$$

Thus Maxwell's equations conform to the charge continuity equation.

Exercise 1.6 (Sect. 1.2.1). Starting from Maxwell's equations, derive the energy conservation law for an electromagnetic field.

Answer. Let us multiply (1.27) by the electric field vector $\mathbf{E}$ and add it to (1.28) multiplied by the magnetic field vector $\mathbf{B}$. The result is

$$\frac{1}{c} \mathbf{E} \frac{\partial \mathbf{E}}{\partial t} + \frac{1}{c} \mathbf{B} \frac{\partial \mathbf{B}}{\partial t} = -\frac{4\pi}{c} \mathbf{j} \mathbf{E} - (\mathbf{B} \operatorname{rot} \mathbf{E} - \mathbf{E} \operatorname{rot} \mathbf{B}).$$

By using the known formula from vector analysis

$$\operatorname{div} [\mathbf{a} \times \mathbf{b}] = \mathbf{b} \operatorname{rot} \mathbf{a} - \mathbf{a} \operatorname{rot} \mathbf{b},$$

we rewrite the last equation as follows

$$\frac{1}{2c} \frac{\partial}{\partial t} (E^2 + B^2) = -\frac{4\pi}{c} \mathbf{j} \mathbf{E} - \operatorname{div} [\mathbf{E} \times \mathbf{B}]$$

or

$$\boxed{\frac{\partial}{\partial t} W = -\operatorname{div} \mathbf{G} - \mathbf{j} \cdot \mathbf{E}.} \quad (1.52)$$

Here

$$W = \frac{E^2 + B^2}{8\pi} \quad (1.53)$$

is the energy of electromagnetic field in a unit volume of space;

$$\mathbf{G} = \frac{c}{4\pi} [\mathbf{E} \times \mathbf{B}] \quad (1.54)$$

is the flux of electromagnetic field energy through a unit surface in space, i.e. the energy flux density for electromagnetic field. This is called the Poynting vector.

The second term on the right-hand side of (1.52) is the power of work done by the electric field on all the charged particles in the unit volume of space. In the simplest approximation

$$e \mathbf{v} \cdot \mathbf{E} = \frac{d}{dt} \mathcal{E}, \quad (1.55)$$

where $\mathcal{E}$ is the particle kinetic energy (see (5.6)). Hence instead of (1.52) we write the divergent form of the energy conservation law:

$$\frac{\partial}{\partial t} \left(\frac{E^2 + B^2}{8\pi} + \frac{\rho v^2}{2} \right) + \operatorname{div} \left(\frac{c}{4\pi} [\mathbf{E} \times \mathbf{B}] \right) = 0. \quad (1.56)$$

Compare this simple approach to the energy conservation law for charged particles and an electromagnetic field with the more general situation considered in Sect. 12.1.3.

Exercise 1.7 (Sect. 1.2.2). Clarify the meaning of the right-hand side of (1.35).

Answer. Substitute definition (1.21) of the delta-function in definition (1.31) of the electric charge density and differentiate the result over time t :

$$\begin{aligned} \frac{\partial \rho^q}{\partial t} &= \sum_{i=1}^N e_i \sum_{\alpha=1}^3 \left[\frac{\partial}{\partial (r_\alpha - r_\alpha^i(t))} \prod_{\beta=1}^3 \delta(r_\beta - r_\beta^i(t)) \right] \frac{\partial}{\partial t} (r_\alpha - r_\alpha^i(t)) = \\ &= - \sum_{i=1}^N e_i \sum_{\alpha=1}^3 \left[\frac{\partial}{\partial (r_\alpha - r_\alpha^i(t))} \prod_{\beta=1}^3 \delta(r_\beta - r_\beta^i(t)) \right] \frac{dr_\alpha^i(t)}{dt}. \end{aligned} \quad (1.57)$$

This is the right-hand side of (1.35).

Chapter 2

Statistical Description of Interacting Particle Systems

Abstract In a system which consists of many interacting particles, the statistical mechanism of “mixing” in phase space works and makes description of the system’s behavior *on average* more simple. A state and evolution of the system is described by a *statistically smoothed* distribution function and the averaged Liouville equation, i.e. the kinetic equation, with attached chain of the equations for correlation functions.

2.1 The Averaging of Liouville’s Equation

2.1.1 Averaging Over Phase Space

As was shown in the first chapter, the exact state of a system consisting of N interacting particles can be given by the *exact* distribution function (see definition (1.20)) in the six-dimensional (6D) phase space $X = \{\mathbf{r}, \mathbf{v}\}$. This function is defined as the sum of δ -functions at N points of the phase space:


$$\hat{f}(\mathbf{r}, \mathbf{v}, t) = \sum_{i=1}^N \delta(\mathbf{r} - \mathbf{r}_i(t)) \delta(\mathbf{v} - \mathbf{v}_i(t)). \quad (2.1)$$

Instead of the equations of motion, we use Liouville’s equation to describe the change of the system state (Sect. 1.1.5):

$$\frac{\partial \hat{f}}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} \hat{f} + \frac{\hat{\mathbf{F}}}{m} \cdot \nabla_{\mathbf{v}} \hat{f} = 0. \quad (2.2)$$

Once the exact initial state of all the particles is known, it can be represented by N points in the phase space X (Fig. 2.1). The motion of these points is described by Liouville’s equation or by the $6N$ equations of motion (1.25).

Fig. 2.1 Particle trajectories in the 6D phase space X



In fact we usually know only some average characteristics of the system's state, such as the temperature, density, etc. Moreover the behavior of each single particle is in general of no interest. For this reason, instead of the exact distribution function (2.1), let us introduce the distribution function *averaged* over a small volume ΔX of phase space, i.e. over a small interval of coordinates $\Delta \mathbf{r}$ and velocities $\Delta \mathbf{v}$ centered at the point $(\mathbf{r}, \mathbf{v})$, at a moment of time t :

$$\begin{aligned} \langle \hat{f}(\mathbf{r}, \mathbf{v}, t) \rangle_X &= \frac{1}{\Delta X} \int_{\Delta X} \hat{f}(X, t) dX = \\ &= \frac{1}{\Delta \mathbf{r} \Delta \mathbf{v}} \int_{\Delta \mathbf{r} \Delta \mathbf{v}} \hat{f}(\mathbf{r}, \mathbf{v}, t) d^3 \mathbf{r} d^3 \mathbf{v}. \end{aligned} \quad (2.3)$$

Here $d^3 \mathbf{r} = dx dy dz$ and $d^3 \mathbf{v} = dv_x dv_y dv_z$, if use is made of Cartesian coordinates.

To put the same in another way, the *mean* number of particles that present at a moment of time t in an element of phase volume ΔX is

$$\langle \hat{f}(\mathbf{r}, \mathbf{v}, t) \rangle_X \cdot \Delta X = \int_{\Delta X} \hat{f}(\mathbf{r}, \mathbf{v}, t) dX. \quad (2.4)$$


The total number N of particles in the system is the integral over the whole phase space X .

Obviously the distribution function averaged over phase volume differs from the exact one as shown in Fig. 2.2.

2.1.2 Two Statistical Postulates

Let us average the same exact distribution function (2.1) over a small time interval Δt centered at a moment of time t :

Fig. 2.2 The one-dimensional analogy of the distribution function averaging over phase space X : (a) the exact distribution function (2.1), (b) the averaged function (2.3)



$$\langle \hat{f}(\mathbf{r}, \mathbf{v}, t) \rangle_t = \frac{1}{\Delta t} \int_{\Delta t} \hat{f}(\mathbf{r}, \mathbf{v}, t) dt. \quad (2.5)$$

Here Δt is small in comparison with the characteristic time of the system's evolution:

$$\Delta t \ll \tau_{ev}. \quad (2.6)$$

We assume that the following *two statistical postulates* concerning systems containing a large number of particles are applicable to the system considered.

The first postulate:

The mean values $\langle \hat{f} \rangle_X$ and $\langle \hat{f} \rangle_t$ exist for sufficiently small ΔX and Δt and are *independent* of the averaging scales ΔX and Δt .


Clearly the first postulate implies that the number of particles should be large. For a small number of particles the mean value depends upon the averaging scale: if, for instance, $N = 1$ then the exact distribution function (2.1) is simply a δ -function, and the average over the variable X is $\langle \hat{f} \rangle_X = 1/\Delta X$. For illustration, the case $(\Delta X)_1 > \Delta X$ is shown in Fig. 2.3.

The second postulate is

$$\langle \hat{f}(X, t) \rangle_X = \langle \hat{f}(X, t) \rangle_t = f(X, t). \quad (2.7)$$

In other words, the averaging of the distribution function over phase space is *equivalent* to the averaging over time.

Fig. 2.3 Averaging of the exact distribution function $\hat{f}$ which is equal to a δ -function. ΔX is a small volume of the phase space X



While speaking of the small ΔX and Δt , we assume that they are not too small: ΔX must contain a reasonably large number of particles while Δt must be large in comparison with the duration of drastic changes of the exact distribution function, such as the duration of the particle Coulomb collisions: $\Delta t \gg \tau_c$. Thus we assume that

$$\tau_c \ll \Delta t \ll \tau_{ev} \quad (2.8)$$

and

$$\langle l \rangle \ll \Delta X \ll L. \quad (2.9)$$

Here $\langle l \rangle \approx n^{-1/3}$ is a mean distance between the particles, n is a number of the particles in a unit volume; L is a distance over which the macroscopic quantities of the system change considerably. In this case the statistical mechanism of particle “mixing” in phase space works, and

the averaging of the exact distribution function over the time Δt is equivalent to the averaging over the phase volume ΔX .

2.1.3 A Statistical Mechanism of Mixing in Phase Space

Let us understand qualitatively how the mixing mechanism works in phase space. We start from the dynamical description of the N -particle system in $6N$ -dimensional phase space in which

$$\Gamma = \{ \mathbf{r}_i, \mathbf{v}_i \}, \quad i = 1, 2, \dots, N, \quad (2.10)$$

a point is determined ($t = 0$ in Fig. 2.4) by the initial conditions of all the particles. The motion of this point, that is the dynamical evolution of the system, can be




Fig. 2.4 The dynamical trajectory of a system of N interacting particles in the $6N$ -dimensional phase space Γ




Fig. 2.5 The dynamical trajectories of two systems never cross each other

described by Liouville's equation or equations of motion. The point moves along a complicated *dynamical trajectory* because the interactions in a many-particle system are extremely intricate and complicated.

The dynamical trajectory has a remarkable property which we shall illustrate by the following example. Imagine a glass vessel containing a gas consisting of a large number N of particles (e.g., 10^{23} molecules or charged particles). The state of this gas at any moment of time is depicted by a single point in the phase space Γ .

Let us imagine another vessel which is identical to the first one, with one exception, being that at any moment of time the gas state in the second vessel is different from that in the first one. These states are depicted by two different points in the space Γ . For example, at $t = 0$, they are points 1 and 2 in Fig. 2.5.

With the passage of time, the gas states in both vessels change, whereas the two points in the space Γ draw two *different* dynamical trajectories (Fig. 2.5). These trajectories do *not* intersect. If they had intersected at just one point (see point A in Fig. 2.6a), then the state of the first gas, determined by $6N$ numbers $(\mathbf{r}_i, \mathbf{v}_i)$, would have coincided with the state of the second gas. These numbers could have




Fig. 2.6 Two types of unbelievable dynamical trajectories




Fig. 2.7 Two types of statistical averaging

been taken as the initial conditions which, in turn, would have uniquely determined the motion. The two trajectories would have merged into one. For the same reason the trajectory of a system cannot intersect itself (Fig. 2.6b). Thus we come to the conclusion that

only one dynamical trajectory of a many particle system passes through each point of the phase space Γ .

Since the trajectories differ in initial conditions, we can introduce an infinite *ensemble of systems* (glass vessels) corresponding to the different initial conditions. In a finite time the ensemble of dynamical trajectories will *closely* fill the phase space Γ , without intersections. By averaging over the ensemble we can answer the question of what the probability is that, at a moment of time t , the system will be found in an element $\Delta\Gamma = \Delta r_i \Delta v_i$ of the phase space Γ (see Fig. 2.7a):

$$dw = \langle \hat{f}(\mathbf{r}_i, \mathbf{v}_i) \rangle_{\Gamma} d\Gamma. \quad (2.11)$$

Here $\langle \hat{f}(\mathbf{r}_i, \mathbf{v}_i) \rangle_\Gamma$ is a function of all the coordinates and velocities. It plays the role of the *probability distribution density* in the phase space Γ and is called the statistical distribution function or simply the distribution function. It is obtained by way of statistical averaging over the ensemble and evidently corresponds to definition (2.3).

* * *

It is rather obvious that the same *probability density* can be obtained in another way—through the averaging over time. The dynamical trajectory of a system, given a sufficient time Δt , will closely cover the phase space Γ . There will be no self-intersections; but since the trajectory is very intricate it will repeatedly pass through the phase space element $\Delta \Gamma$ (Fig. 2.7b).

Let $(\Delta t)_\Gamma$ be the time during which the system locates in $\Delta \Gamma$. For a sufficiently large Δt , which is formally restricted by the characteristic time of a relatively slow evolution of the system as a whole, the ratio $(\Delta t)_\Gamma / \Delta t$ tends to the limit

$$\lim_{\Delta t \rightarrow \infty} \frac{(\Delta t)_\Gamma}{\Delta t} = \frac{dw}{d\Gamma} = \langle \hat{f}(\mathbf{r}_i, \mathbf{v}_i, t) \rangle_t. \quad (2.12)$$

By virtue of the role of the probability density, it is clear that

the statistical averaging over the ensemble (2.11) is equivalent to the averaging over time (2.12) as well as to the definition (2.5).

2.1.4 The Derivation of a General Kinetic Equation

Now we have everything what we need to average the exact Liouville equation (2.2). Since the equation contains the derivatives with respect to time t and phase-space coordinates $(\mathbf{r}, \mathbf{v})$, the procedure of averaging over the small interval $\Delta X \Delta t$ is defined as follows:

$$f(X, t) = \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \hat{f}(X, t) dX dt. \quad (2.13)$$

Averaging the first term of the Liouville equation gives

$$\begin{aligned} \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{\partial \hat{f}}{\partial t} dX dt &= \frac{1}{\Delta t} \int_{\Delta t} \frac{\partial}{\partial t} \left[\frac{1}{\Delta X} \int_{\Delta X} \hat{f} dX \right] dt = \\ &= \frac{1}{\Delta t} \int_{\Delta t} \frac{\partial}{\partial t} f dt = \frac{\partial f}{\partial t}. \end{aligned} \quad (2.14)$$

In the last equality the use is made of the fact that, by virtue of the second postulate of statistics (2.7), the averaging of the smooth averaged function does not change it.

Let us average the second term in (2.2):

$$\begin{aligned} \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} v_\alpha \frac{\partial \hat{f}}{\partial r_\alpha} dX dt &= \frac{1}{\Delta X} \int_{\Delta X} v_\alpha \frac{\partial}{\partial r_\alpha} \left[\frac{1}{\Delta t} \int_{\Delta t} \hat{f} dt \right] dX = \\ &= \frac{1}{\Delta X} \int_{\Delta X} v_\alpha \frac{\partial}{\partial r_\alpha} f dX = v_\alpha \frac{\partial f}{\partial r_\alpha}. \end{aligned} \quad (2.15)$$

Here the index $\alpha = 1, 2, 3$.

In order to average the term containing the force $\hat{\mathbf{F}}$, let us represent this exact force as a sum of a *mean force* $\langle \mathbf{F} \rangle$ and the force due to the difference of the real force field from the mean (statistically smoothed) one:

$$\hat{\mathbf{F}} = \langle \mathbf{F} \rangle + \mathbf{F}'. \quad (2.16)$$

Substituting definition (2.16) in the third term in (2.2) and averaging this term, we have

$$\begin{aligned} \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{\hat{F}_\alpha}{m} \frac{\partial \hat{f}}{\partial v_\alpha} dX dt &= \\ &= \frac{\langle F_\alpha \rangle}{m} \frac{1}{\Delta X} \int_{\Delta X} \frac{\partial}{\partial v_\alpha} \left[\frac{1}{\Delta t} \int_{\Delta t} \hat{f} dt \right] dX + \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{F'_\alpha}{m} \frac{\partial \hat{f}}{\partial v_\alpha} dX dt = \\ &= \frac{\langle F_\alpha \rangle}{m} \frac{\partial f}{\partial v_\alpha} + \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{F'_\alpha}{m} \frac{\partial \hat{f}}{\partial v_\alpha} dX dt. \end{aligned} \quad (2.17)$$

Gathering all three terms together, we write the averaged Liouville equation in the form

$$\boxed{\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} f + \frac{\langle \mathbf{F} \rangle}{m} \cdot \nabla_{\mathbf{v}} f = \left(\frac{\partial \hat{f}}{\partial t} \right)_c}, \quad (2.18)$$

where

$$\boxed{\left(\frac{\partial \hat{f}}{\partial t} \right)_c = - \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{F'_\alpha}{m} \frac{\partial \hat{f}}{\partial v_\alpha} dX dt}. \quad (2.19)$$

Equation (2.18) and its right-hand side (2.19) are called the *kinetic equation* and the *collisional integral* (cf. definition (1.19)), respectively.

Therefore we have found the *most general* form of the kinetic equation with a collisional integral, which is nice but cannot be directly used in plasma astrophysics, without making some additional simplifying assumptions. The main assumption, the binary character of collisions, will be taken into account in the next section, see also Sect. 3.3.

2.2 A Collisional Integral and Correlation Functions

2.2.1 Binary Interactions

The statistical mechanism of mixing in phase space makes particles have no individuality. However we have to distinguish *different kinds* of particles, for example, electrons and protons, because their behaviors differ. Let $\hat{f}_k(\mathbf{r}, \mathbf{v}, t)$ be the exact distribution function (2.1) of particles of the *kind* k , i.e.

$$\hat{f}_k(\mathbf{r}, \mathbf{v}, t) = \sum_{i=1}^{N_k} \delta(\mathbf{r} - \mathbf{r}_{ki}(t)) \delta(\mathbf{v} - \mathbf{v}_{ki}(t)), \quad (2.20)$$

the index i denoting the i th particle of kind k , N_k being the number of particles of kind k . The Liouville equation (2.2) for the particles of kind k takes a view

$$\frac{\partial \hat{f}_k}{\partial t} + \mathbf{v} \cdot \nabla_{\mathbf{r}} \hat{f}_k + \frac{\hat{\mathbf{F}}_k}{m_k} \cdot \nabla_{\mathbf{v}} \hat{f}_k = 0, \quad (2.21)$$

m_k is the mass of a particle of kind k .

The force acting on a particle of kind k at a point $(\mathbf{r}, \mathbf{v})$ of the phase space X at a moment of time t , $\hat{F}_{k,\alpha}(\mathbf{r}, \mathbf{v}, t)$, is the sum of forces acting on this particle from all other particles (see Fig. 2.8):


$$\hat{F}_{k,\alpha}(\mathbf{r}, \mathbf{v}, t) = \sum_l \sum_{i=1}^{N_l} \hat{F}_{kl,\alpha}^{(i)}(\mathbf{r}, \mathbf{v}, \mathbf{r}_{li}(t), \mathbf{v}_{li}(t)). \quad (2.22)$$

So the total force $\hat{F}_{k,\alpha}(\mathbf{r}, \mathbf{v}, t)$ depends upon the instant positions and velocities of all the particles generally with the time delay taken into account (see Landau and Lifshitz 1975, Chap. 8, Sect. 63).

With the help of the exact distribution function, we can rewrite formula (2.22) as follows:

$$\hat{F}_{k,\alpha}(\mathbf{r}, \mathbf{v}, t) = \sum_l \int_{X_1} F_{kl,\alpha}(X, X_1) \hat{f}_l(X_1, t) dX_1. \quad (2.23)$$

Fig. 2.8 An action of a particle e_{li} located at the point $\mathbf{r}_{li}$ on a particle of kind k at a point $\mathbf{r}$ at a moment of time t



Here we assume that an interaction law $F_{kl,\alpha}(X, X_1)$ is explicitly independent of time t ;

$$\hat{f}_l(X, t) = \sum_{i=1}^{N_l} \delta(X - X_{li}(t))$$

is the exact distribution function of particles of kind l , the variable of integration is designated as $X_1 = \{\mathbf{r}_1, \mathbf{v}_1\}$ and $dX_1 = d^3\mathbf{r}_1 d^3\mathbf{v}_1$.

Formula (2.23) takes into account that the forces considered are *binary* ones, i.e. they can be represented as a sum of interactions between *two* particles.

Making use of the representation (2.23), let us average the force term in the Liouville equation (2.2), as this has been done in formula (2.17). We have

$$\begin{aligned} & \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{1}{m_k} \hat{F}_{k,\alpha}(\mathbf{r}, \mathbf{v}, t) \frac{\partial \hat{f}_k}{\partial v_\alpha} dX dt = \\ & = \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) \hat{f}_l(X_1, t) \frac{\partial}{\partial v_\alpha} \hat{f}_k(X, t) dX dX_1 dt = \\ & = \frac{1}{\Delta X} \int_{\Delta X} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) \times \\ & \quad \times \frac{\partial}{\partial v_\alpha} \left[\frac{1}{\Delta t} \int_{\Delta t} \hat{f}_k(X, t) \hat{f}_l(X_1, t) dt \right] dX dX_1. \end{aligned} \quad (2.24)$$

Here we have taken into account that the exact distribution function $\hat{f}_l(X_1, t)$ is independent of the velocity $\mathbf{v}$, which is a part of the variable $X = \{\mathbf{r}, \mathbf{v}\}$ related to the particles of the kind k .

Formula (2.24) contains the *pair products* of the exact distribution functions of different particle kinds, as is natural for the case of *binary interactions*.

2.2.2 Binary Correlation

Let us represent the exact distribution function $\hat{f}_k$ as

$$\hat{f}_k(X, t) = f_k(X, t) + \hat{\varphi}_k(X, t), \quad (2.25)$$

where $f_k(X, t)$ is the *statistically averaged* distribution function, $\hat{\varphi}_k(X, t)$ is the deviation of the exact distribution function from the averaged one. In general the deviation is not small, of course. It is obvious that, according to definition (2.25),

$$\hat{\varphi}_k(X, t) = \hat{f}_k(X, t) - f_k(X, t);$$

hence

$$\langle \hat{\varphi}_k(X, t) \rangle = 0. \quad (2.26)$$

Let us consider the integrals of pair products, appearing in the averaged force term (2.24). In view of definition (2.25), they can be rewritten as

$$\frac{1}{\Delta t} \int_{\Delta t} \hat{f}_k(X, t) \hat{f}_l(X_1, t) dt = f_k(X, t) f_l(X_1, t) + f_{kl}(X, X_1, t), \quad (2.27)$$

where

$$f_{kl}(X, X_1, t) = \frac{1}{\Delta t} \int_{\Delta t} \hat{\varphi}_k(X, t) \hat{\varphi}_l(X_1, t) dt. \quad (2.28)$$

The function f_{kl} is referred to as the *correlation function* or, more exactly, the *binary correlation function*.

The physical meaning of the correlation function is clear from (2.27). The left-hand side of (2.27) means the probability to find a particle of kind k at a point X of the phase space at a moment of time t *under condition* that a particle of kind l places at a point X_1 at the same time. By definition this is a *conditional probability* (e.g., Gnedenko 1965, Sects. 23 and 52). In the right-hand side of (2.27) the distribution function $f_k(X, t)$ characterizes the probability that a particle of kind k stays at a point X at a moment of time t . The function $f_l(X_1, t)$ plays the analogous role for the particles of kind l .

If the particles of kind k did not interact with those of kind l , then their distributions would be independent, i.e. probability densities would simply multiply:

$$\langle \hat{f}_k(X, t) \hat{f}_l(X_1, t) \rangle = f_k(X, t) f_l(X_1, t). \quad (2.29)$$

So in the right-hand side of (2.27) there should be

$$f_{kl}(X, X_1, t) = 0. \quad (2.30)$$

In other words, there would be no correlation in the particle distribution.

We consider a system of interacting particles. With the proviso that the parameter characterizing the binary interaction, namely Coulomb collision considered below,

$$\zeta_i \approx \frac{e^2}{\langle l \rangle} \left/ \left\langle \frac{mv^2}{2} \right\rangle \right., \quad (2.31)$$

is small under conditions in a wide range, the correlation function must be *relatively small*:

if the interaction is weak, the second term in the right-hand side of (2.27) must be small in comparison with the first one.

We shall come back to the discussion of this property in Sect. 3.1. This fundamental property allows us to construct a theory of plasma in many cases of astrophysical interest.

2.2.3 The Collisional Integral and Binary Correlation

Now let us substitute (2.27) in formula (2.24) for the statistically averaged force term in the kinetic equation:

$$\begin{aligned} \frac{1}{\Delta X \Delta t} \int_{\Delta X} \int_{\Delta t} \frac{1}{m_k} \hat{F}_{k,\alpha}(X, t) \frac{\partial \hat{f}_k}{\partial v_\alpha} dX dt = \\ = \frac{1}{\Delta X} \int_{\Delta X} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) \frac{\partial}{\partial v_\alpha} [f_k(X, t) f_l(X_1, t) + \\ + f_{kl}(X, X_1, t)] dX dX_1 = \end{aligned}$$

since: (a) the averaging of smooth functions does not change them, (b) the function $f_k(X, t)$ does not depend of X_1 , and (c) the function $f_l(X_1, t)$ does not depend of X , we can proceed as follows

$$\begin{aligned}
&= \frac{1}{m_k} \left[\frac{\partial}{\partial v_\alpha} f_k(X, t) \right] \left\{ \sum_l \int_{X_1} F_{kl, \alpha}(X, X_1) f_l(X_1, t) dX_1 \right\} + \\
&+ \sum_l \int_{X_1} \frac{1}{m_k} F_{kl, \alpha}(X, X_1) \frac{\partial}{\partial v_\alpha} f_{kl}(X, X_1, t) dX_1 = \\
&= \frac{1}{m_k} F_{k, \alpha}(X, t) \frac{\partial f_k(X, t)}{\partial v_\alpha} + \\
&+ \sum_l \int_{X_1} \frac{1}{m_k} F_{kl, \alpha}(X, X_1) \frac{\partial f_{kl}(X, X_1, t)}{\partial v_\alpha} dX_1. \tag{2.32}
\end{aligned}$$

Here we have taken into account the following definition of the *averaged force*

$$F_{k, \alpha}(X, t) = \sum_l \int_{X_1} F_{kl, \alpha}(X, X_1) f_l(X_1, t) dX_1. \tag{2.33}$$

This definition follows from averaging the definition (2.23) of the exact force $\hat{\mathbf{F}}_k$ and coincides with the previous definition (2.17) of the *mean* (smoothed or averaged) force, since

all the deviations of the real force $\hat{\mathbf{F}}_k$ from the mean (smooth) force $\mathbf{F}_k$ are taken care of in the deviations $\hat{\varphi}_k$ and $\hat{\varphi}_l$ of the real distribution functions $\hat{f}_k$ and $\hat{f}_l$ from their mean values f_k and f_l .

Thus the collisional integral can be represented in the form

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \sum_l \int_{X_1} \frac{1}{m_k} F_{kl, \alpha}(X, X_1) \frac{\partial f_{kl}(X, X_1, t)}{\partial v_\alpha} dX_1. \tag{2.34}$$

Moreover, if in the last term of (2.32) the binary interactions can be represented by smooth functions of the type $e_k e_l (|\mathbf{r}_k - \mathbf{r}_l|)^{-2}$ with or without account of the Debye-Hückel shielding (see Sects. 3.2.3 and 8.2.1), then formally the velocity dependence may be neglected.

Let us recall an important particular case considered in Sect. 1.1. For the Lorentz force (1.15) as well as for the gravitational one (1.43), the condition (1.8) is satisfied. Let us require that in formula (2.34)

$$\frac{\partial}{\partial v_\alpha} F_{kl, \alpha}(X, X_1) = 0. \tag{2.35}$$

In fact this condition was tacitly assumed from the early beginning, from (2.2). Anyway, in the case (2.35), we obtain from formula (2.34) the following expression

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \frac{\partial}{\partial v_\alpha} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) f_{kl}(X, X_1, t) dX_1. \quad (2.36)$$

Hence

the collisional integral, at least, for the Coulomb and gravity forces can be written in the *divergent* form in the *velocity* space $\mathbf{v}$:

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \frac{\partial}{\partial v_\alpha} J_{k,\alpha}, \quad (2.37)$$

where the flux of particles of kind k in the velocity space (cf. Fig. 1.3b) is

$$J_{k,\alpha}(X, t) = \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) f_{kl}(X, X_1, t) dX_1. \quad (2.38)$$

Therefore we arrive to conclusion that the averaged Liouville equation or *the kinetic equation for particles of kind k*

$$\begin{aligned} \frac{\partial f_k(X, t)}{\partial t} + v_\alpha \frac{\partial f_k(X, t)}{\partial r_\alpha} + \frac{F_{k,\alpha}(X, t)}{m_k} \frac{\partial f_k(X, t)}{\partial v_\alpha} = \\ = - \frac{\partial}{\partial v_\alpha} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) f_{kl}(X, X_1, t) dX_1 \end{aligned} \quad (2.39)$$

contains the *unknown* function f_{kl} . Hence the kinetic equation (2.39) for distribution function f_k is not closed. We have to find the equation for the correlation function f_{kl} . This will be done in the next section.

2.3 Equations for Correlation Functions

To derive the equations for correlation functions (in the first place for the function of pair correlations f_{kl}), it is not necessary to introduce any new postulates or develop new formalisms. All the necessary equations and averaging procedures are at hand.

Looking at definition (2.28), we see that we need an equation which will describe the deviation of distribution function from its mean value, i.e. the function

$\hat{\varphi}_k = \hat{f}_k - f_k$. In order to derive such equation, we simply have to subtract the averaged representation (2.39) from the exact Liouville equation (2.2). The result is

$$\begin{aligned} \frac{\partial \hat{\varphi}_k(X, t)}{\partial t} + v_\alpha \frac{\partial \hat{\varphi}_k(X, t)}{\partial r_\alpha} + \frac{\hat{F}_{k,\alpha}}{m_k} \frac{\partial \hat{f}_k}{\partial v_\alpha} - \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} = \\ = \frac{\partial}{\partial v_\alpha} \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) f_{kl}(X, X_1) dX_1. \end{aligned} \quad (2.40)$$

Here

$$\hat{F}_{k,\alpha}(X, t) = \sum_l \int_{X_1} F_{kl,\alpha}(X, X_1) \hat{f}_l(X_1, t) dX_1 \quad (2.41)$$

is the *exact* force (2.23) acting on a particle of the kind k at the point X of phase space, and

$$F_{k,\alpha}(X, t) = \sum_l \int_{X_1} F_{kl,\alpha}(X, X_1) f_l(X_1, t) dX_1 \quad (2.42)$$

is the statistically *averaged* force (2.33).

Thus the difference between the exact force and the averaged one is

$$\hat{F}_{k,\alpha} - F_{k,\alpha} = \sum_l \int_{X_1} F_{kl,\alpha}(X, X_1) \hat{\varphi}_l(X_1, t) dX_1. \quad (2.43)$$

We substitute it in (2.40) where, at first, the difference of force terms can be rewritten as follows:

$$\begin{aligned} \frac{\hat{F}_{k,\alpha}}{m_k} \frac{\partial \hat{f}_k}{\partial v_\alpha} - \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} &= \frac{\hat{F}_{k,\alpha}}{m_k} \frac{\partial}{\partial v_\alpha} (f_k + \hat{\varphi}_k) - \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} = \\ &= \frac{\hat{F}_{k,\alpha} - F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} + \frac{\hat{F}_{k,\alpha}}{m_k} \frac{\partial \hat{\varphi}_k}{\partial v_\alpha}. \end{aligned}$$

The result of the substitution is

$$\begin{aligned} \frac{\hat{F}_{k,\alpha}}{m_k} \frac{\partial \hat{f}_k}{\partial v_\alpha} - \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} &= \\ &= \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) \hat{\varphi}_l(X_1, t) dX_1 \frac{\partial f_k}{\partial v_\alpha} + \frac{F_{k,\alpha}}{m_k} \frac{\partial \hat{\varphi}_k}{\partial v_\alpha} + \\ &+ \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) \hat{\varphi}_l(X_1, t) dX_1 \frac{\partial \hat{\varphi}_k}{\partial v_\alpha}. \end{aligned} \quad (2.44)$$

On substituting (2.44) in (2.40) we have the equation for the deviation $\hat{\varphi}_k$ of the exact distribution function $\hat{f}_k$ from its mean value f_k :

$$\frac{\partial \hat{\varphi}_k(X, t)}{\partial t} + v_\alpha \frac{\partial \hat{\varphi}_k(X, t)}{\partial r_\alpha} + \dots = 0. \quad (2.45)$$

Considering that we have to derive the equation for the pair correlation function

$$f_{kl}(X_1, X_2, t) = \langle \hat{\varphi}_k(X_1, t) \hat{\varphi}_l(X_2, t) \rangle,$$

let us take two equations:

one for $\hat{\varphi}_k(X_1, t)$

$$\frac{\partial \hat{\varphi}_k(X_1, t)}{\partial t} + v_{1,\alpha} \frac{\partial \hat{\varphi}_k(X_1, t)}{\partial r_{1,\alpha}} + \frac{F_{k,\alpha}}{m_k} \frac{\partial \hat{\varphi}_k(X_1, t)}{\partial v_{1,\alpha}} + \dots = 0 \quad (2.46)$$

and another for $\hat{\varphi}_l(X_2, t)$

$$\frac{\partial \hat{\varphi}_l(X_2, t)}{\partial t} + v_{2,\alpha} \frac{\partial \hat{\varphi}_l(X_2, t)}{\partial r_{2,\alpha}} + \frac{F_{l,\alpha}}{m_l} \frac{\partial \hat{\varphi}_l(X_2, t)}{\partial v_{2,\alpha}} + \dots = 0. \quad (2.47)$$

Now we add the equations resulting from (2.46) multiplied by $\hat{\varphi}_l$ and (2.47) multiplied by $\hat{\varphi}_k$. We obtain

$$\hat{\varphi}_l \frac{\partial \hat{\varphi}_k}{\partial t} + \hat{\varphi}_k \frac{\partial \hat{\varphi}_l}{\partial t} + v_{1,\alpha} \frac{\partial \hat{\varphi}_k}{\partial r_{1,\alpha}} \hat{\varphi}_l + v_{2,\alpha} \frac{\partial \hat{\varphi}_l}{\partial r_{2,\alpha}} \hat{\varphi}_k + \dots = 0$$

or

$$\frac{\partial (\hat{\varphi}_k \hat{\varphi}_l)}{\partial t} + v_{1,\alpha} \frac{\partial (\hat{\varphi}_k \hat{\varphi}_l)}{\partial r_{1,\alpha}} + v_{2,\alpha} \frac{\partial (\hat{\varphi}_k \hat{\varphi}_l)}{\partial r_{2,\alpha}} + \dots = 0. \quad (2.48)$$

On averaging (2.48) we finally have the equation for the *pair correlation* function in the following form:

$$\begin{aligned} & \frac{\partial f_{kl}(X_1, X_2, t)}{\partial t} + v_{1,\alpha} \frac{\partial f_{kl}(X_1, X_2, t)}{\partial r_{1,\alpha}} + v_{2,\alpha} \frac{\partial f_{kl}(X_1, X_2, t)}{\partial r_{2,\alpha}} + \\ & + \frac{F_{k,\alpha}(X_1, t)}{m_k} \frac{\partial f_{kl}(X_1, X_2, t)}{\partial v_{1,\alpha}} + \frac{F_{l,\alpha}(X_2, t)}{m_l} \frac{\partial f_{kl}(X_1, X_2, t)}{\partial v_{2,\alpha}} + \\ & + \frac{\partial f_k(X_1, t)}{\partial v_{1,\alpha}} \sum_n \int_{X_3} \frac{1}{m_k} F_{kn,\alpha}(X_1, X_3) f_{nl}(X_3, X_2, t) dX_3 + \\ & + \frac{\partial f_l(X_2, t)}{\partial v_{2,\alpha}} \sum_n \int_{X_3} \frac{1}{m_l} F_{ln,\alpha}(X_2, X_3) f_{nk}(X_3, X_1, t) dX_3 = \end{aligned}$$

$$\begin{aligned}
&= -\frac{\partial}{\partial v_{1,\alpha}} \sum_n \int_{X_3} \frac{1}{m_k} F_{kn,\alpha}(X_1, X_3) f_{kln}(X_1, X_2, X_3, t) dX_3 - \\
&\quad -\frac{\partial}{\partial v_{2,\alpha}} \sum_n \int_{X_3} \frac{1}{m_l} F_{ln,\alpha}(X_2, X_3) f_{kln}(X_1, X_2, X_3, t) dX_3. \quad (2.49)
\end{aligned}$$

Here

$$f_{kln}(X_1, X_2, X_3, t) = \frac{1}{\Delta t} \int_{\Delta t} \hat{\varphi}_k(X_1, t) \hat{\varphi}_l(X_2, t) \hat{\varphi}_n(X_3, t) dt \quad (2.50)$$

is the function of *triple correlations* (see also Exercise 2.51).

Thus (2.49) for the pair correlation function contains the *unknown* function of triple correlations. In general,

the chain of equations for correlation functions can be shown to be *unclosed*: the equation for the correlation function of s th order contains the function of the order $(s + 1)$.

2.4 Practice: Exercises and Answers

Exercise 2.1 (Sect. 2.3). By analogy with formula (2.27), show that

$$\begin{aligned}
&\langle \hat{f}_k(X_1, t) \hat{f}_l(X_2, t) \hat{f}_n(X_3, t) \rangle = \\
&= f_k(X_1, t) f_l(X_2, t) f_n(X_3, t) + \\
&\quad + f_k(X_1, t) f_{ln}(X_2, X_3, t) + f_l(X_2, t) f_{kn}(X_1, X_3, t) + \\
&\quad + f_n(X_3, t) f_{kl}(X_1, X_2, t) + f_{kln}(X_1, X_2, X_3, t). \quad (2.51)
\end{aligned}$$

Exercise 2.2. Discuss a similarity and difference between the kinetic theory presented in this chapter and the famous BBGKY hierarchy theory developed by Bogoliubov (1946), Born and Green (1949), Kirkwood (1946), and Yvon (1935).

Hint. Show that essential to both derivations is the weak-coupling assumption, according to which

grazing encounters, involving small fractional energy and momentum exchange between colliding particles, dominate the evolution of the velocity distribution function.

The *weak-coupling* assumption provides justification of the widely appreciated practice (see Chap. 3) which leads to a very significant simplification of the original collisional integral; for more detail see [Klimontovich \(1975, 1986\)](#).

Chapter 3

Weakly-Coupled Systems with Binary Collisions

Abstract In a system which consists of many interacting particles, the weak-coupling assumption allows us to introduce a well controlled approximation to consider the chain of the equations for correlation functions. This leads to a very significant simplification of the original collisional integral to describe collisional relaxation and transport in astrophysical plasmas but not in cosmic self-gravitating systems.

3.1 Approximations for Binary Collisions

3.1.1 *The Small Parameter of Kinetic Theory*

The infinite chain of equations for the distribution function and correlation functions does not contain more information in itself than the initial Liouville equation for the exact distribution function. Moreover, the statistical mixing in phase space with subsequent statistical smoothing the distribution function over the physically infinitesimal volume allows to lose “*useless information*”—the information about the exact motion of particles. Just for this reason, description of the system’s behavior becomes irreversible.

The value of the chain is also that the chain allows a direct introduction of new physical assumptions which make it possible to break the chain off at some term (Fig. 3.1) and to estimate the resulting error. We call this procedure a *well controlled approximation* because it looks, in a sense, similar to the Taylor expansion series.

There is no universal way of breaking the chain off. It is intimately related, in particular, to the physical state of a plasma. Different states (as well as different aims) require different approximations. In general, the physical state of a plasma can be characterized, at least partially, by the ratio of the mean energy of two particle interaction to their mean kinetic energy (parameter (2.31)). If the last one can be reasonably characterized by some effective temperature T (see Sect. 9.4.1), then this ratio




Fig. 3.1 How to break the infinite chain of the equations for correlation functions? *LT* is the Liouville theorem (1.13) or (2.21) for an exact distribution function $\hat{f}_k$. *KE* and *BC* are the kinetic equation (2.39) for f_k and (2.49) for the binary correlation function f_{kl}

$$\zeta_i \approx \frac{e^2}{\langle l \rangle} \bigg/ k_B T. \quad (3.1)$$

As a mean distance between the particles we take $\langle l \rangle \approx n^{-1/3}$. Hence the ratio

$$\zeta_i = \frac{e^2}{n^{-1/3}} \bigg/ k_B T = \frac{e^2}{k_B} \times \frac{n^{1/3}}{T} \quad (3.2)$$

is termed the *interaction parameter*. It is small for a sufficiently *hot* and *rarefied* plasma.

In many astrophysical plasmas, for example in the solar corona (see Exercise 3.2), the interaction parameter is really very small. So the thermal kinetic energy of plasma particles is much larger than their interaction energy.

The particles are almost free or moving on definite trajectories in the external fields if the later are present.

We shall call this case the approximation of *weak* Coulomb interaction. An existence of the small parameter allows us to have a complete description of this interaction by using the perturbation procedure. Moreover such a description is the simplest and the most exact one.

While constructing the kinetic theory, it is natural to use the perturbation theory with respect to the small parameter ζ_i . This means that

the distribution function f_k must be taken to be of order unity, the pair correlation function f_{kl} of order ζ_i , the triple correlation function f_{klm} of order ζ_i^2 , etc.

We shall see in Sect. 3.2.3 that this principle has a deep physical sense in kinetic theory. Such plasmas are said to be “*weakly coupled*”.

An opposite case, when the interaction parameter takes values larger than unity, is very dense, relatively cold plasmas, for example in the interiors of white dwarf stars (Exercise 3.3). These plasmas are “*strongly coupled*”. The major part of matter in the Universe finds itself in the state of high-pressure plasmas. Estimations show that about 95% of matter (without taking a *dark matter* into account; see, however, Bradt 2008) are the plasmas of stars, pulsars, black holes, and giant planets. Plasma nonideality defines the behavior of matter in a wide range of the phase diagram, from solid and liquid to neutral gas, the phase boundaries of melting and boiling, and the

metal-dielectric transition region. Investigation of strongly compressed Coulomb systems is now one of the hottest and most intensively developed fundamental branches of physics (Fortov et al. 2006).

3.1.2 The Vlasov Kinetic Equation

In the zeroth order with respect to the small parameter ζ_i , we obtain the Vlasov equation with the self-consistent electromagnetic field (Vlasov 1938, 1945):

$$\begin{aligned} \frac{\partial f_k(X, t)}{\partial t} + v_\alpha \frac{\partial f_k(X, t)}{\partial r_\alpha} + \\ + \frac{e_k}{m_k} \left(\mathbf{E} + \frac{1}{c} \mathbf{v} \times \mathbf{B} \right)_\alpha \frac{\partial f_k(X, t)}{\partial v_\alpha} = 0. \end{aligned} \quad (3.3)$$

Here $\mathbf{E}$ and $\mathbf{B}$ are the *statistically averaged* electric and magnetic fields obeying Maxwell's equations:

$$\begin{aligned} \text{rot } \mathbf{E} &= -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}, \quad \text{div } \mathbf{E} = 4\pi (\rho^0 + \rho^q), \\ \text{rot } \mathbf{B} &= \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} + \frac{4\pi}{c} (\mathbf{j}^0 + \mathbf{j}^q), \quad \text{div } \mathbf{B} = 0. \end{aligned} \quad (3.4)$$

ρ^0 and $\mathbf{j}^0$ are the densities of external charges and currents; they describe the large-scale external fields, for example, the uniform magnetic field $\mathbf{B}_0$. ρ^q and $\mathbf{j}^q$ are the *statistically smoothed* charge and current densities due to the plasma particles themselves:

$$\rho^q(\mathbf{r}, t) = \sum_k e_k \int_{\mathbf{v}} f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}, \quad (3.5)$$

$$\mathbf{j}^q(\mathbf{r}, t) = \sum_k e_k \int_{\mathbf{v}} \mathbf{v} f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (3.6)$$

Therefore the electric and magnetic fields are also statistically smoothed.

In other words, if we are considering the physical processes which occur on a time scale much shorter than the time scale of collisions,

$$\tau_{ev} \ll \tau_c, \quad (3.7)$$

we may use the description which includes the *averaged* electric and magnetic fields arising from the *averaged* charge density and current density in plasma, but

neglects the microfields responsible for binary collisions. This means that $\mathbf{F}' = 0$ in formula (2.16) for the mean force. Therefore the collisional integral (2.19) is also equal to zero.

The Vlasov kinetic equation (3.3) together with the definitions (3.5) and (3.6), and with Maxwell's equations (3.4) is a *nonlinear integro-differential* equation. It serves as a classic basis for the theory of oscillations and waves in a plasma (e.g., Silin 1971; Schmidt 1979; Benz 2002) with the small parameter ζ_i and small correlational effects of higher orders.

The Vlasov equation is also a proper basis for kinetic theory of wave-particle interactions in turbulent plasmas (see Diamond et al. 2010), in astrophysical plasmas (Chap. 7), collisionless *shock waves* (Sect. 16.4) and collisionless *reconnecting current layers* (Harris 1962; Coppi et al. 1966; see review of current state in Part II). The Vlasov equation was strongly criticized by Ginzburg et al. (1946).

One of the natural limitations of the Vlasov equation is that it will not make a plasma relax to a Maxwellian distribution (see Sect. 9.6), since we effectively neglect collisions by neglecting the binary correlation function.

Vlasov was the first to recognize that

the electromagnetic interaction among plasma particles is qualitatively different from the interaction in an ordinary gas.

3.1.3 The Landau Collisional Integral

Using the perturbation theory with respect to the small interaction parameter ζ_i in the first order, and, therefore, neglecting the *close* Coulomb collisions (this will be justified in Sect. 8.1.5), we can find the kinetic equation with the collisional integral given by Landau (1937)

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \frac{\partial}{\partial v_\alpha} J_{k,\alpha}, \quad (3.8)$$

where the flux of particles of kind k in the velocity space (cf. formula (2.38)) is

$$\begin{aligned} J_{k,\alpha} = & \frac{\pi e_k^2 \ln \Lambda}{m_k} \sum_l e_l^2 \int_{\mathbf{v}_l} \left\{ f_k \frac{\partial f_l}{m_l \partial v_{l,\beta}} - f_l \frac{\partial f_k}{m_k \partial v_{k,\beta}} \right\} \times \\ & \times \frac{(u^2 \delta_{\alpha\beta} - u_\alpha u_\beta)}{u^3} d^3 \mathbf{v}_l. \end{aligned} \quad (3.9)$$

Here $\mathbf{u} = \mathbf{v} - \mathbf{v}_l$ denotes the relative velocity, $d^3 \mathbf{v}_l$ corresponds to the integration over the whole velocity space of “*field*” particles l . $\ln \Lambda$ is the Coulomb logarithm which takes into account divergence of the Coulomb-collision cross-section

(see Sect. 8.1.5). The full kinetic equation (2.18) with the Landau collisional integral is a nonlinear integro-differential equation for the distribution function $f_k(\mathbf{r}, \mathbf{v}, t)$ of particles of the kind k .

The date of publication of the Landau (1937) paper may be considered as the date of birth of the kinetic theory of *collisional* fully-ionized plasma. The theory of *collisionless* plasma begins with the classical paper of Vlasov (1938). In fact, these two approaches correspond to *different limiting cases*.

The Landau integral takes into account the part of the particle interaction which determines dissipation while the Vlasov equation allows for the averaged field, and is thus reversible.

For example, in the Vlasov theory the question of the role of collisions in the neighbourhood of resonances remains open. The famous paper by Landau (1946) was devoted to this problem. Landau used the reversible Vlasov equation as the basis to study the dynamics of a small perturbation of the Maxwell distribution function, $f^{(1)}(\mathbf{r}, \mathbf{v}, t)$. In order to solve the linearized Vlasov equation (Sect. 7.1.1), he made use of the Laplace transformation, and defined the rule to avoid a pole at $\omega = k_{\parallel} v_{\parallel}$ in the divergent integral (see formula (7.11) in Sect. 7.1.2) by the replacement $\omega \rightarrow \omega + i0$.

This technique for avoiding singularities may be formally replaced by a different procedure. Namely it is possible to add a small dissipative term $-\nu f^{(1)}(\mathbf{r}, \mathbf{v}, t)$ to the right-hand side of the linearized Vlasov equation. In this way, the Fourier transform of the kinetic equation involves the complex frequency $\omega = \omega' + i\nu$, leading with $\nu \rightarrow 0$ to the *same* expression for the *Landau damping* (see formula (7.13) in Sect. 7.1.2) Note, however, that

the Landau damping is not by randomizing Coulomb collisions but by a *collisionless transfer* of wave field energy into oscillations of resonant particles

(see Sect. 7.1.2 for more detail).

Thus there are two different approaches to the description of plasma oscillation damping. The first is based on mathematical regularization of the Cauchy integral (e.g., Lavrent'ev and Shabat 1973, Chap. 3, Sect. 3) divergence. In this approach the physical nature of the damping seems to be not considered since the initial equation remains reversible. However the Landau method is really a beautiful example of complex analysis leading to an important new physical result.

The second approach reduces the reversible Vlasov equation to an irreversible one. Although the dissipation is assumed to be negligibly small, one cannot take the limit $\nu \rightarrow 0$ directly in the master equations: this can be done only in the final formulae. This second method of introducing the collisional damping is more natural. It shows that

even very rare collisions play the principal role in the physics of collisionless plasma.

It is this approach that has been adopted in [Klimontovich \(1986\)](#). A more comprehensive solution of this principal question, however, can only be obtained on the basis of the dissipative kinetic equation.

The example of the Landau resonance and Landau damping demonstrates that some fundamental problems still remain unsolved in the kinetic theory of plasma. They arise from inconsistent descriptions of the transition from the reversible equations of the mechanics of charge particles and fields to the irreversible kinetic equations for statistically averaged distribution functions ([Klimontovich 1998](#)).

In the first approximation with respect to the small interaction parameter ζ_i we shall find the Maxwellian distribution function and the effect of Debye shielding. This will be the subject of the Sect. 3.2.

3.1.4 The Fokker-Planck Equation

The smallness of the interaction parameter ζ_i signifies that, in the Landau collisional integral, the sufficiently *distant* Coulomb collisions are taken care of as the interactions with a *small momentum and energy transfer* (see Sect. 8.1). For this reason, it comes as no surprise that the Landau integral can be considered as a particular case of a different approach which is the Fokker-Planck equation ([Fokker 1914](#); [Planck 1917](#)). The latter generally describes systems of many particles that move under action of *stochastic* forces producing *small* changes in particle velocities (for a review see [Chandrasekhar 1943a](#)).

Let us consider a distribution function independent of space so that $f = f(\mathbf{v}, t)$. The Fokker-Planck equation describes the distribution function evolution due to *nonstop overlapping weak collisions* resulting in particle diffusion in velocity space:

$$\frac{\partial f}{\partial t} = \left(\frac{\partial \hat{f}}{\partial t} \right)_c = - \frac{\partial}{\partial v_\alpha} [a_\alpha f] + \frac{\partial^2}{\partial v_\alpha \partial v_\beta} [b_{\alpha\beta} f]. \quad (3.10)$$

The Fokker-Planck equation formally coincides with the diffusion-type equation (which is irreversible of course) for some admixture with concentration f , for example, *Brownian particles*¹ (or *test* particles) in a gas, on which stochastic forces are exerted by the molecules of the gas. The coefficient $b_{\alpha\beta}$ plays the role of the diffusion coefficient and is expressed in terms of the averaged velocity change $\langle \delta v_\alpha \rangle$ in an elementary act—a collision:

$$b_{\alpha\beta} = \frac{1}{2} \langle \delta v_\alpha \delta v_\beta \rangle. \quad (3.11)$$

¹In 1827, the botanist Robert Brown described the chaotic motion of a grain of pollen suspended on the water and repeatedly hit by water molecules. In 1900, Louis Bachelier and independently Albert Einstein studied the *Brown motion* from the mathematical point of view.

The other coefficient is

$$a_\alpha = \langle \delta v_\alpha \rangle. \quad (3.12)$$

It is known as the Fokker-Planck coefficient of *dynamic friction*. For example, a Brownian particle moving with velocity $\mathbf{v}$ through the gas experiences a drag opposing the motion (see Fig. 1.4). We easily expect such friction to be present in *collisional* plasma (Sect. 8.4.1(c)). It is more difficult to imagine this effect in system like a star cluster where the particles interact through long-range forces. Chandrasekhar (1943b) derived the famous result that such a friction term in the Fokker-Planck equation should arise in a *stellar system* (see Exercise 3.5).

In order to find the mean values appearing in the Fokker-Planck equation, we have to make clear the physical and mathematical sense of expressions (3.11) and (3.12), see Exercise 3.4.

The mean values of velocity changes are in fact statistically averaged and determined by the forces acting between a test particle and scatterers (*field* particles or waves).

Because of this, these averaged quantities have to be expressed by the collisional integral with the corresponding cross-sections (Exercises 3.5 and 3.6). The “standard” derivation of the Fokker-Planck equation from the Boltzmann integral, with discussion of its particular features, can be found, for example, in Shoub (1987); however see Sects. 11.5 and 11.6 in Balescu (1975) for a detailed analysis.

For the test particles interacting with the thermal electrons and ions in a plasma, such calculations can be made and give us the Landau collisional integral (Sect. 3.1.3); see, for example, Sect. 11.8 in Balescu (1975). Thus one did not anticipate any major problems in rewriting the Landau integral in the Fokker-Planck form (e.g., Somov 1982). The kinetic equation found in this way will allow us to study the Coulomb interaction of accelerated particle beams with astrophysical plasma (Chap. 4). As was indicated above, the first term in the Fokker-Planck equation is a *friction*.

Collisional friction in plasma slows down the fast particles of the beam and moves them toward the zero velocity in the velocity space


as illustrated by Fig. 3.2. The second term represents the three-dimensional *diffusion* of the beam particles in the velocity space. Diffusion disperses the distribution of beam particles.

During the motion of a beam of accelerated particles in a plasma a *reverse* current of thermal electrons is generated, which tends to compensate the electric current of accelerated particles—the *direct* current (Sect. 4.5.1).

The electric field driving the reverse current in astrophysical plasmas makes a great impact on the particle beam kinetics.

That is why, in order to solve the problem of accelerated particle propagation in, for example, the solar atmosphere, we inevitably have to apply a *combined approach*,

Fig. 3.2 A beam of fast particles in plasma can generate the Langmuir waves due to the *bump-on tail* instability which will be shown in Chap. 7. Here we illustrate only the simplest effects of Coulomb collisions, that will be considered in Chap. 4



which takes into account both the electric field influence on the accelerated particles (as in the Vlasov equation; Sect. 3.1.2) and their scattering from the thermal particles of a plasma (as in the Landau integral; see Sect. 4.5.2).

The Landau collisional integral is effectively used in many problems of plasma astrophysics. It permits a considerable simplification of the calculations of many quantities determined by collisions of charged particles, such as the viscosity coefficient, thermal conductivity, electric conductivity, etc. (Sect. 9.6).

The Landau collisional integral does not take into account the *close* collisions since they are responsible for large exchange of the particle momentum (see Sect. 8.1). So the interaction parameter is not small, and the perturbation theory is not applicable (Exercise 3.6). However the close Coulomb collisions of charged particles can play an important role in *collective* plasma phenomena (Klimontovich 1986).


3.2 Correlations and Debye-Hückel Shielding

We are going to understand the most fundamental property of the binary correlation function. With this aim in mind, we shall solve the second equation in the chain illustrated by Fig. 3.1. In order to determine and to solve this equation we have to know two functions for each kind of interacting particles: the distribution function f_k from the first link *KE* in the chain and the triple correlation function f_{kln} from the third link (see Fig. 3.3).

3.2.1 The Maxwellian Distribution Function

Let us consider the stationary ($\partial/\partial t = 0$) solution to the equations for correlation functions, assuming the interaction parameter ζ_i to be small and using the *method of successive approximations* in the following iterative form.

Fig. 3.3 Formulation of the problem on the binary correlation function f_{kl}



First, we set $f_{kl} = 0$ in the averaged Liouville equation (2.39) for the distribution function f_k . Second, we assume that the triple correlation function f_{kln} is zero in (2.49) for the correlation function f_{kl} etc.

Let us start from the first iteration. The plasma is supposed to be stationary, uniform and in the *thermodynamic equilibrium* state, i.e. the particle velocity distribution is assumed to be a Maxwellian function

$$f_k(X) = f_k(v^2) = c_k \exp\left(-\frac{m_k v^2}{2k_B T_k}\right). \quad (3.13)$$

The constant c_k is determined by the normalizing condition and equals

$$c_k = n_k \left(\frac{m_k}{2\pi k_B T_k}\right)^{3/2}. \quad (3.14)$$

Under assumptions made above, the Maxwellian function satisfies the kinetic equation

$$\underbrace{\frac{\partial f_k(v^2)}{\partial t}}_{=0} + v_\alpha \underbrace{\frac{\partial f_k(v^2)}{\partial r_\alpha}}_{=0} + F_{k,\alpha}(X) \underbrace{\frac{1}{m_k}}_{\neq 0} \underbrace{\frac{\partial f_k(v^2)}{\partial v_\alpha}}_{\neq 0} = 0,$$

if the averaged force is equal to zero:

$$F_{k,\alpha}(X) = 0. \quad (3.15)$$

We shall justify this assumption in the next section.

3.2.2 The Averaged Force and Electric Neutrality

Let us substitute the Maxwellian distribution function (3.13) in the kinetic equation (2.39), neglecting all the interactions except the Coulomb ones. For the latter, in circumstances where the averaged distribution functions for the components are *uniform*, we obtain the following expression for the averaged force (2.33):

$$\begin{aligned}
F_{k,\alpha}(X_1) &= \sum_l \int_{X_2} F_{kl,\alpha}(X_1, X_2) f_l(X_2) dX_2 = \\
&\text{since plasma is uniform, the function } f_l \text{ does not depend of } \mathbf{r}_2 \\
&= \sum_l \int_{\mathbf{r}_2} F_{kl,\alpha}(\mathbf{r}_1, \mathbf{r}_2) d^3\mathbf{r}_2 \cdot \int_{\mathbf{v}_2} f_l(\mathbf{v}_2) d^3\mathbf{v}_2 = \\
&= - \int_{\mathbf{r}_2} \sum_l \frac{\partial}{\partial r_{1,\alpha}} \left(\frac{e_k e_l}{|\mathbf{r}_1 - \mathbf{r}_2|} \right) d^3\mathbf{r}_2 \times n_l = \\
&= - \int_{\mathbf{r}_2} \frac{\partial}{\partial r_{1,\alpha}} \left(\frac{e_k}{|\mathbf{r}_1 - \mathbf{r}_2|} \right) d^3\mathbf{r}_2 \times \sum_l n_l e_l. \tag{3.16}
\end{aligned}$$

Therefore

$$F_{k,\alpha} = 0, \tag{3.17}$$

if the plasma is assumed to be *electrically neutral*:

$$\boxed{\sum_l n_l e_l = 0,} \tag{3.18}$$

or *quasi-neutral* (see Sect. 8.2). *Balanced charges of ions and electrons* determine the name *plasma* according [Langmuir \(1928\)](#).

So the averaged (statistically smoothed) force (2.33) is equal to zero in the electrically neutral plasma but is not equal to zero in a system of charged particles of the same charge sign: positive or negative, it does not matter. Such a system tends to expand.

There is no neutrality in *gravitational systems* like the stellar clusters.

■ The large-scale gravitational field makes an *overall* thermodynamic equilibrium in a gravitational system impossible

(see Sect. 9.8). Moreover, on the contrary to plasma, the gravitational systems tend to contract and collapse.

3.2.3 Pair Correlations and the Debye-Hückel Radius

As a first approximation, on putting the triple correlation function $f_{kln} = 0$, we obtain from (2.49), in view of condition (3.17), the following equation for the binary or pair correlation function f_{kl} :

$$\begin{aligned}
& v_{1,\alpha} \frac{\partial f_{kl}}{\partial r_{1,\alpha}} + v_{2,\alpha} \frac{\partial f_{kl}}{\partial r_{2,\alpha}} = \\
& = - \sum_n \int_{X_3} \left\{ \frac{1}{m_k} F_{kn,\alpha}(X_1, X_3) f_{nl}(X_3, X_2) \frac{\partial f_k}{\partial v_{1,\alpha}} + \right. \\
& \quad \left. + \frac{1}{m_l} F_{ln,\alpha}(X_2, X_3) f_{nk}(X_3, X_1) \frac{\partial f_l}{\partial v_{2,\alpha}} \right\} dX_3. \quad (3.19)
\end{aligned}$$

Let us consider the particles of two kinds: electrons and ions, assuming the ions to be motionless and homogeneously distributed. Then the ions do not take part in any kinetic processes; hence $\hat{\varphi}_i \equiv 0$ for ions, and all the correlation functions associated with $\hat{\varphi}_i$ equal zero as well:

$$f_{ii} = 0, \quad f_{ei} = 0 \quad \text{etc.} \quad (3.20)$$

Among the pair correlation functions, only $f_{ee}(X_1, X_2)$ has a non-zero magnitude. Taking into account (3.20) and (3.13), rewrite (3.19) as follows

$$\begin{aligned}
& \mathbf{v}_1 \frac{\partial f_{ee}}{\partial \mathbf{r}_1} + \mathbf{v}_2 \frac{\partial f_{ee}}{\partial \mathbf{r}_2} = \\
& = \frac{1}{k_B T} \int_{X_3} [\mathbf{v}_1 \cdot \mathbf{F}(X_1, X_3) f_{ee}(X_3, X_2) f_e(\mathbf{v}_1) + \\
& \quad + \mathbf{v}_2 \cdot \mathbf{F}(X_2, X_3) f_{ee}(X_1, X_3) f_e(\mathbf{v}_2)] dX_3. \quad (3.21)
\end{aligned}$$

Since $\mathbf{v}_1$ and $\mathbf{v}_2$ are arbitrary and refer to the same kind of particles (electrons), (3.21) takes the form

$$\frac{\partial f_{ee}}{\partial \mathbf{r}_1} = \frac{1}{k_B T} \int_{X_3} \mathbf{F}(X_1, X_3) f_{ee}(X_3, X_2) f_e(\mathbf{v}_1) dX_3. \quad (3.22)$$

Taking into account the character of Coulomb force in the same approximation as in formula (3.17) and assuming the correlation to exist only between the positions of the particles in space (rather than between velocities), let us integrate both sides of (3.22) over $d^3\mathbf{v}_1 d^3\mathbf{v}_2$. The result is

$$\frac{\partial g(\mathbf{r}_1, \mathbf{r}_2)}{\partial \mathbf{r}_1} = -\frac{ne^2}{k_B T} \int_{\mathbf{r}_3} \nabla_{\mathbf{r}_1} \frac{1}{|\mathbf{r}_1 - \mathbf{r}_3|} g(\mathbf{r}_2, \mathbf{r}_3) d^3\mathbf{r}_3. \quad (3.23)$$

Here the function

$$g(\mathbf{r}_1, \mathbf{r}_2) = \int_{\mathbf{v}_1} \int_{\mathbf{v}_2} f_{ee}(X_1, X_2) d^3\mathbf{v}_1 d^3\mathbf{v}_2. \quad (3.24)$$

We integrate (3.23) over $\mathbf{r}_1$ and designate the function

$$g(\mathbf{r}_1, \mathbf{r}_2) = g(r_{12}),$$

where $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$. So we obtain the equation

$$g(r_{12}) = -\frac{ne^2}{k_B T} \int_{\mathbf{r}_3} \frac{g(r_{23})}{r_{13}} d^3\mathbf{r}_3. \quad (3.25)$$

Its solution is

$$g(r) = \frac{c_0}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right), \quad (3.26)$$

where

$$r_{\text{DH}} = \left(\frac{k_B T}{4\pi n e^2}\right)^{1/2} \quad (3.27)$$

is the *Debye-Hückel radius* or, more exactly, the electron Debye-Hückel radius. It will be defined in just this form (see formula (8.33)) for the case when the Debye-Hückel shielding is due to the particles of one kind—due to electrons. A more general formula for the Debye-Hückel radius will be derived in Sect. 8.2.1.

The constant of integration

$$c_0 = -\frac{1}{4\pi r_{\text{DH}}^2 n} \quad (3.28)$$

(see Exercise 3.8). Substituting (3.28) in solution (3.26) gives the sought-after pair correlation function, more exactly the velocity-integrated pair-correlation function

$$g(r) = -\frac{1}{k_B T} \frac{e^2}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right). \quad (3.29)$$

Formula (3.29) shows that

the Debye-Hückel radius is a characteristic length scale of the pair correlations in a fully-ionized equilibrium plasma.

This result proves to be fair in the context of Sect. 8.2.1 where the Debye-Hückel shielding will be considered in another approach. Comparison of formula (3.29) with (8.32) shows that, as one might have anticipated,

the binary correlation function reproduces the shape of the actual potential of interaction, i.e. the *shielded* Coulomb potential:

$$g(r) \sim \varphi(r) \sim \frac{1}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right).$$

(3.30)

Note that the probability to find an electron near another one is

$$dw = g(r) 4\pi r^2 dr \sim r \exp\left(-\frac{r}{r_{\text{DH}}}\right) dr \rightarrow 0 \quad (3.31)$$

when $r \rightarrow 0$. Naturally this is because the repulsive force between the electrons.

Note also that, for $r = \langle l \rangle$, the coefficient before exponent in formula (3.29) coincides with the interaction parameter (3.1). This is well consistent with our assumption that, if we prescribe to distribution function f_k the order of unity, then the pair correlation function f_{kl} is of order the interaction parameter ζ_i . That is why

we do have the well-controlled approximation considering the chain of equations for correlation functions in weakly-coupled plasmas.

3.2.4 Plasma Parameter and Collective Phenomena

It is known that astrophysical plasma can exhibit *collective phenomena* arising out of mutual interactions of many charged particles. Since the Debye-Hückel radius r_{DH} is a characteristic length scale of pair correlations, the number $n r_{\text{DH}}^3$ gives us a measure of the number of particles which can interact simultaneously. The inverse of this number is the so-called *plasma parameter*

$$\zeta_p = (n r_{\text{DH}}^3)^{-1}. \quad (3.32)$$

This is a small quantity as well as it can be expressed in terms of the small interaction parameter ζ_i (see Exercise 3.1). The fact that $\zeta_p \ll 1$ implies *a large number of plasma particles* in a volume enclosed by the sphere of the Debye-Hückel radius. In many astrophysical applications, the plasma parameter (3.32) is really small. Therefore the number of particles inside the Debye-Hückel sphere is very large (e.g., Exercise 3.2). So

the collective phenomena can be really important in astrophysical plasmas in many places where a plasma is weakly coupled.

The electrostatic oscillations are the simplest example of such collective behavior. Each charged particle moves due to the electrostatic field produced by all the particles around it. However different charged particles moving in this way give rise to coherent oscillations, *Langmuir waves*; see Sect. 10.2.

3.3 Gravitational Systems

3.3.1 Vlasov-Type Equation

There is a fundamental difference between the astrophysical plasmas and the gravitational systems with the interaction potential (1.43), for example, the stars in a galaxy. This difference lies in the nature of the gravitational force: there is *no shielding* to vitiate this long-range $1/r^2$ force. The collisional integral formally equals infinity because the binary correlation function $g(r) \sim 1/r$.

The conventional wisdom of such system dynamics (see [Binney and Tremaine 1987](#)) asserts that the structure and evolution of a system of N self-gravitating point masses can be described by the collisionless kinetic equation, the gravitational analog of the Vlasov equation (see also Exercise 16.7). The basic assumption underlying the Vlasov equation is that the gravitational N -body system can be described probabilistically in terms of a statistically smooth distribution function $f(X, t)$. The Vlasov equation manifests the idea that this function will stream freely in the self-consistent gravitational potential $\phi(\mathbf{r}, t)$ (cf. (1.43)) associated with $f(X, t)$, so that

$$\frac{\partial f(X, t)}{\partial t} + v_\alpha \frac{\partial f(X, t)}{\partial r_\alpha} - \frac{\partial \phi}{\partial r_\alpha} \frac{\partial f(X, t)}{\partial v_\alpha} = 0. \quad (3.33)$$

Here

$$\Delta \phi = -4\pi G \rho(\mathbf{r}, t) \quad (3.34)$$

and

$$\rho(\mathbf{r}, t) = \int f(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (3.35)$$

Note that, in the context of the *mean field* theory, a distribution of particles over their masses has no effect.

Applying for example to the system of stars in a galaxy, (3.33) implies that

the net gravitational force acting on a star is assumed to be determined by the large-scale structure of the galaxy rather than by whether the star happens to lie close to some other star.

The force acting on any star does not vary rapidly, and each star is supposed to accelerate smoothly through the force field generated by the galaxy as a whole.

In fact, encounters between stars may cause the acceleration $\dot{\mathbf{v}}$ to differ from the smoothed gravitational force $-\nabla\phi$ and therefore invalidate (3.33). *Gravitational encounters are not screened*, they can be thought of as leading to an additional collisional term on the right side of the equation—a collisional integral. However very little is known mathematically about such possibility.

On the basis of what we have seen in Sect. 3.2.3,

the collisionless approach in gravitational systems, i.e. the entire neglect of particle pair correlations, constitutes an *uncontrolled approximation*, even if the interaction parameter is small.

Unlike the case of astrophysical plasma, we *cannot* derive the next order correction to the collisionless kinetic equation in the context of a systematic perturbation expansion.

3.3.2 Collisional Integral and Other Models

The $1/r$ potential yields an infinite cross-section. So, when evaluating the effects of collisions in the usual way (see Sect. 8.1.5) for a gravitational system, we encounter logarithmic divergences in the limit of large impact parameter (see formula (8.18)). We may hope to circumvent this difficulty, the problematic Coulomb logarithm of gravitational dynamics, by first identifying the bulk mean field force $\langle \mathbf{F} \rangle$ in definition (2.16), acting at any given point in space. Then we treat fluctuations $\mathbf{F}'$ away from the mean field force. This splitting into a mean field plus fluctuations can be introduced formally (Kandrup 1998) and allows one to write down the collisional integral of the type (2.19). However this is difficult to implement concretely because of the apparent absence of a clean separation of time scales.

For the N -body problem with $N \gg 1$ we might expect that these fluctuations are small, so that their effects do in fact constitute a small perturbation. So it is assumed that, on long time scales, one must allow for discreteness effects, described by the Fokker-Plank equation (3.10) or the kinetic equation with the Landau collisional integral (3.8) in the following form (e.g., Lancellotti and Kiessling 2001):

$$\left(\frac{\partial \hat{f}}{\partial t} \right)_c = \sigma \frac{\partial}{\partial \mathbf{v}} \int_{\mathbf{v}'} \frac{\partial^2 |\mathbf{v} - \mathbf{v}'|}{\partial \mathbf{v} \partial \mathbf{v}'} \cdot \left(\frac{\partial}{\partial \mathbf{v}} - \frac{\partial}{\partial \mathbf{v}'} \right) [f(\mathbf{r}, \mathbf{v}, t) f(\mathbf{r}, \mathbf{v}', t)] d^3 \mathbf{v}'. \quad (3.36)$$

Here σ is a constant determined by the effective collision rate.

Given that theoretical analyses have as yet proven inconclusive, one might instead seek resource to numerical experiments. This, however, is also difficult for gravitational systems not characterized by a high degree of symmetry. There is in fact only one concrete setting where detailed computations have been done, namely the *toy model* of one-dimensional gravity.

In summary, even though a *mean* gravitational field theory based on the Vlasov-type equation may seem well motivated physically, there is as yet no rigorous proof of its validity. In particular, no rigorous estimate as to the time scale on which it might be expected to fail. On the other hand, the collisional or even hydrodynamic descriptions of gravitational systems have a difficulty of the same origin. The gravitational attraction cannot be screened (see also Sect. 9.8).

3.4 Comments on Numerical Simulations

At present, the astrophysical plasma processes are typically investigated in well developed and distinct approaches. One approach, described by the Vlasov equation, is the collisionless limit used when collective effects dominate. In cases where the plasma dynamics is determined by collisional processes in external fields and where the self-consistent fields can be neglected, the Fokker-Planck approach is used. At the same time, it is known that

both collective kinetic effects and Coulomb collisions can play an essential role in a great variety of astrophysical phenomena

starting from the most simple one—propagation of fast particles in plasma (Chap. 4). Besides, as was mentioned in Sect. 3.1.3, *collisions play the principal role in the physics of collisionless plasma*. Taking collisions into account may lead not only to quantitative but also qualitative changes in the plasma behavior, even if the collision frequency ν is much less than the electron plasma frequency.

It is known that, even in the collisionless limit, the kinetic equation is still too difficult for numerical simulations, and the “*macroparticle*” methods are the most widely used algorithms. In these methods, instead of direct numerical solution of the kinetic equation, a set of ordinary differential equations for every macroparticle is solved. These equations are the characteristics of the Vlasov equation.

In the case of a collisional plasma, the position of a macroparticle satisfies the usual equation of the collisionless case

$$\dot{\mathbf{r}} \equiv \frac{d\mathbf{r}}{dt} = \mathbf{v}(t), \quad (3.37)$$

but the momentum equation is modified owing to the Coulomb collisions. They are described by the Fokker-Planck operator (3.10) which introduces a *friction* (the coefficient a_α) and *diffusion* (the coefficient $b_{\alpha\beta}$) in velocity space. Thus it is necessary to find the effective collisional force $\mathbf{F}_c$ which acts on the macroparticles:

$$\dot{\mathbf{v}} \equiv \frac{d\mathbf{v}}{dt} = \frac{1}{m} (\mathbf{F}_L + \mathbf{F}_c). \quad (3.38)$$

The collisional force can be introduced phenomenologically (see Jones et al. 1996) but a more mathematically correct approach can be constructed using the stochastic equivalence of the Fokker-Planck and Langevin equations (see Cadjan and Ivanov 1999). So *stochastic differential equations* can be regarded as an alternative to the description of astrophysical plasma in terms of distribution function.

The Langevin approach allows one to overcome some difficulties related to the Fokker-Planck equation and to simulate actual plasma processes, taking account of both collective effects and Coulomb collisions.

A broad treatment of the basic ideas and algorithms associated with the so-called sampling-based methods, often also referred to as *stochastic simulation*, can be found in [Asmussen and Glynn \(2007\)](#). The reach of these ideas is well illustrated there, in Chap. 10, by discussing a wide range of different applications, in particular, related to stochastic differential equations (see also [Iacus 2008](#)).

Generally, if we want to construct an effective method for the simulation of complex nonlinear processes in astrophysical plasma, we have to satisfy the following obvious but conflicting conditions.

First, the method should be adequate for the task in hand. For a number of problems the application of simplified models of the collisional integral can provide a correct description and ensure good accuracy. The constructed model should describe collisional effects with the desired accuracy.

Second, the method should be computationally efficient. The algorithm should not be extremely time-consuming. In practice, some compromise between accuracy and complexity of the method should be achieved. Otherwise, we restrict ourselves either to a relatively simple setup of the problem or to a too-rough description of the phenomena.

A “recipe”:

the choice of a particular collisional model (or a model of the collisional integral) is determined by the importance and particular features of the collisional processes in a given astrophysical problem.

3.5 Practice: Exercises and Answers

Exercise 3.1 (Sect. 3.1.1). Show that the interaction parameter ζ_i is related to the plasma parameter ζ_p as follows:

$$\zeta_i = \frac{1}{4\pi} \zeta_p^{2/3}, \quad (3.39)$$

if the Debye-Hückel radius is given by formula (3.27). Discuss the difference between ζ_i and ζ_p .

Exercise 3.2 (Sect. 3.1.1). How many particles are inside the Debye-Hückel sphere in plasma of the solar corona?

Answer. From formula (8.31) for the Debye-Hückel radius in a two-component equilibrium plasma (see also formula (8.82) in Exercise 3.3) it follows that, for an electron-proton plasma with $T \approx 2 \times 10^6$ K and $n \approx 2 \times 10^8$ cm⁻³, the Debye-Hückel radius

$$r_{\text{DH}} = \left(\frac{kT}{8\pi e^2 n} \right)^{1/2} \approx 4.9 \left(\frac{T}{n} \right)^{1/2} \approx 0.5 \text{ cm}. \quad (3.40)$$

The number of particles inside the Debye-Hückel sphere

$$N_{\text{DH}} = n \frac{4}{3} \pi r_{\text{DH}}^3 \sim 10^8. \quad (3.41)$$

Hence the typical value of plasma parameter (3.32) in the corona is really small: $\zeta_p \sim 10^{-8}$. The interaction parameter (3.2) is also small:

$$\zeta_i \sim 10^{-6}$$

(see formula (3.39)).

Exercise 3.3 (Sect. 3.1.1). Estimate the interaction parameter (3.2) in the interior of *white dwarf* stars (de Martino et al. 2003; see also Exercise 3.3).

Comment. It may seem at first sight that the mutual interactions between electrons would be very important inside a white dwarf star. However, in a system of fermions with most states filled up to the Fermi energy,

collisions among nearby electrons are suppressed due to the fact that the electrons may not have free state available for occupation after the collision

(see Kittel 1995). Hence electrons inside a white dwarf star are often approximated as a perfect gas made up of non-interacting fermions (see Sect. 57 in Landau and Lifshitz 1959b). For this reason, some results of plasma astrophysics are applicable to the electron gas inside white dwarfs.

Exercise 3.4 (Sect. 3.1.4). Let $w = w(\mathbf{v}, \delta\mathbf{v})$ be the probability that a test particle changes its velocity $\mathbf{v}$ to $\mathbf{v} + \delta\mathbf{v}$ in the time interval δt . The velocity distribution at the time t can be written as

$$f(\mathbf{v}, t) = \int f(\mathbf{v} - \delta\mathbf{v}, t - \delta t) w(\mathbf{v} - \delta\mathbf{v}, \delta\mathbf{v}) d^3\delta\mathbf{v}. \quad (3.42)$$

Bearing in mind that the interaction parameter (3.1) is small and, therefore, $|\delta\mathbf{v}| \ll |\mathbf{v}|$, expand the product $f w$ under the integral into a Taylor series.

Take the first three terms in the series and show that, in formulae (3.12) and (3.11), the average velocity change per time interval δt :

$$\langle \delta v_\alpha \rangle = \int \delta v_\alpha w d^3\delta\mathbf{v}, \quad (3.43)$$

$$\langle \delta v_\alpha \delta v_\beta \rangle = \int \delta v_\alpha \delta v_\beta w d^3\delta\mathbf{v}. \quad (3.44)$$

Show that

the Fokker-Planck equation (3.10) follows from the Taylor series expansion of the function $f(\mathbf{v}, t)$ given by formula (3.42).

Exercise 3.5 (Sect. 3.1.4). Express the collisional integral in terms of the differential cross-sections of interaction between particles (Smirnov 1981).

Discussion. Boltzmann (1872) considered a dilute neutral gas. Since the particles in a neutral gas do not have long-range interactions like the charged particles in a plasma, they are assumed to interact only when they collide, i.e. when the separation between two particles is not much larger than $2a$, where a is the “radius of a particle”. A particle moves freely in a straight line between two collisions.

In a binary collision, let $\mathbf{v}_k$ and $\mathbf{v}_l$ be the velocities of particles k and l before the collision, $\mathbf{v}'_k$ and $\mathbf{v}'_l$ be the velocities of the same particles after the collision. There are two types of collisions: (a) one that increases the density of the particles at a given point of phase space by bringing in particles from other phase space locations, (b) the other that reduces the density of particles by taking particles away from this point to other phase space locations; these are the collisions $\mathbf{v}_k + \mathbf{v}_l \rightarrow \mathbf{v}'_k + \mathbf{v}'_l$.

By using notations taken into account that k and l can be different kinds of particles, we write the Boltzmann collisional integral in the form (cf. Boltzmann 1956):

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = \sum_l \int \int_{\Omega} (f'_k f'_l - f_k f_l) v_{kl} d\sigma_{kl} d^3 \mathbf{v}_l. \quad (3.45)$$

Here $\mathbf{v}_{kl} = \mathbf{v}_k - \mathbf{v}_l$ is the relative velocity, $d^3 \mathbf{v}_l$ corresponds to the integration over the whole velocity space of “field” particles l . $f_k = f_k(t, \mathbf{r}, \mathbf{v}_k)$ is the distribution function of particles of the kind k , $f'_k = f_k(t, \mathbf{r}, \mathbf{v}'_k)$. The product $f'_k f'_l$ corresponds to the collisions $\mathbf{v}'_k + \mathbf{v}'_l \rightarrow \mathbf{v}_k + \mathbf{v}_l$ which enhance the particle density.

The precollision velocities $\mathbf{v}_k$ and $\mathbf{v}_l$ are related to the postcollision velocities $\mathbf{v}'_k$ and $\mathbf{v}'_l$ through the conservation laws of momentum and energy. These relations give us four scalar equations. However we need six equations to find two vectors $\mathbf{v}'_k$ and $\mathbf{v}'_l$.

A fifth condition comes from the fact the vectors $\mathbf{v}'_k$ and $\mathbf{v}'_l$ will have to lie in the plane of the vectors $\mathbf{v}_k$ and $\mathbf{v}_l$. This follows from the momentum conservation law and means that collisions are coplanar if the force of interaction between two particles is radial.

We need one more condition. We do not expect, of course, that the outcome of a collision is independent of the nature of interaction. If the impact parameter of the collision is given, we can calculate the deflection produced by the collision from the interaction potential. The case of the Coulomb potential is considered in Chap. 8.

Since we are interested here in a statistical treatment, it is enough for us to know the probability of deflection in different direction or a differential scattering cross-section

$$d\sigma_{kl} = \frac{d\sigma_{kl}(v_{kl}, \chi)}{d\Omega} d\Omega, \quad (3.46)$$

where $d\Omega = 2\pi \sin \chi d\chi$ is a solid angle. If the particles are modeled as hard spheres undergoing two-body elastic collisions, the differential scattering cross-section is a function of the scattering angle χ alone.

The Boltzmann gas model can be used for low-density neutral particles as well as for interactions of charged particles with neutral particles.

In plasma astrophysics, the Rutherford formula (8.8) is used to characterize the Coulomb collisions of charged particles. A general case is considered, for example, in Kogan (1967), Silin (1971), Lifshitz and Pitaevskii (1981).

Exercise 3.6 (Sect. 3.1.4). Show that

the Fokker-Planck collisional model can be derived from the Boltzmann collisional integral (3.45)

under the assumption that the change in the velocity of a particle due to a collision is rather small.

Exercise 3.7 (Sect. 3.1.4). The Landau collisional integral is generally thought to approximate the Boltzmann integral (3.45) for the $1/r$ potentials to a “dominant order”, i.e. to within terms of order $1/\ln\Lambda$, where $\ln\Lambda$ is the Coulomb logarithm (see formula (8.23)). However this is not the whole truth. Show that

the Landau integral approximates the Boltzmann integral to the dominant order only in parts of the velocity space.

Hint. This can be established by carrying the Taylor series expansion of the Boltzmann integral to the fourth order. The first term in the series will be the familiar Landau-type collisional integral. The conclusion, drawn from the higher-order terms (Shoub 1987), is that the large-angle scattering processes can play a more significant role in the evolution of the distribution function than currently believed. The normally “non-dominant” part of the diffusion tensor can make a contribution to the collisional term that decays more slowly with increasing velocity than do terms that are retained. In general, the approximations made are not uniformly valid in the velocity space, if the particle distribution functions are not sufficiently close to equilibrium distributions (Cercignani 1969).

Exercise 3.8 (Sect. 3.2.3). Find the constant of integration c_0 in formula (3.26).

Answer. Let us solve the Poisson equation for the shielded potential φ (more justification will be given in Sect. 8.2.1):

$$\begin{aligned}\Delta\varphi &= -4\pi en \left\{ 1 - \left[1 + \frac{c_0}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right) \right] \right\} = \\ &= n \frac{4\pi e c_0}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right).\end{aligned}\tag{3.47}$$

Here it is taken into account that

$$\int_{\mathbf{v}_1} \int_{\mathbf{v}_2} \langle \hat{f}_k(X_1) \hat{f}_l(X_2) \rangle d^3\mathbf{v}_1 d^3\mathbf{v}_2 = n_k(\mathbf{r}_1) n_l(\mathbf{r}_2) + g_{kl}(\mathbf{r}_1, \mathbf{r}_2).$$

The general solution of (3.47) in the spherically symmetric case, i.e. the solution of equation

$$\frac{1}{r} \frac{d^2}{dr^2} (r\varphi) = \frac{4\pi e c_0}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right) n,$$

is of the form

$$\varphi(r) = n \frac{4\pi e r_{\text{DH}}^2 c_0}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right) + c_1 + \frac{c_2}{r}.$$

Since, as $r \rightarrow 0$, the potential φ takes the form $(-e)/r$, $c_1 = c_2 = 0$, and the only non-zero constant is

$$c_0 = -\frac{1}{4\pi r_{\text{DH}}^2 n}. \quad (3.48)$$

Chapter 4

Propagation of Fast Particles in Plasma

Abstract Among a variety of kinetic phenomena related to fast particles in astrophysical plasma, the simplest effect is Coulomb collisions under propagation of the particles in a plasma. The so-called thick-target model is considered. An important role of the reverse-current electric field in this model is demonstrated. In particular, we inevitably have to take into account the reverse current effects in order to study the problem of acceleration of electrons in solar flares. However this problem is of general astrophysical significance how to interpret the observed cosmic hard X-rays quantitatively correct.

4.1 Derivation of the Basic Kinetic Equation

4.1.1 Basic Approximations

We shall call a test particle of kind k a *fast* one if it moves in plasma with velocity larger than the mean thermal velocity of the field particles of the kind k . Among a rich variety of kinetic phenomena related to accelerated fast electrons and ions in astrophysical plasma (Kivelson and Russell 1995) let us consider the simplest effect—Coulomb collisions under propagation of a *fast particle beam in a fully-ionized thermal plasma*. We shall assume that there exists some external (background) magnetic field $\mathbf{B}_0$ which determines a way of particle propagation. This field can be considered locally as a uniform one.

Electric and magnetic fields, $\mathbf{E}$ and $\mathbf{B}$, related to a beam of fast particles will be discussed in Sect. 4.5. Heating of plasma will be considered, for example, in Sect. 8.3. So, until this will be necessary,

fast particles will be considered as “test” particles that do not influence the background plasma and magnetic field $\mathbf{B}_0$.

Let $f = f(t, \mathbf{r}, \mathbf{v})$ be an unknown distribution function of test particles. In what follows, $q = Ze$ and $m = Am_p$ are electric charge and mass of a test particle, respectively.

We restrict a problem by consideration of fast but non-relativistic particles interacting with background plasma which consists of thermal electrons ($m_1 = m_e$ and $e_1 = -e$) and thermal protons ($m_2 = m_p$ and $e_2 = +e$). Both components of a plasma are in thermodynamic equilibrium. Using the kinetic equation with the Landau collisional integral (3.8) we obtain

$$\frac{\partial f}{\partial t} + v_\alpha \frac{\partial f}{\partial r_\alpha} + \frac{q}{m} \left\{ E_\alpha + \frac{1}{c} [\mathbf{v} \times (\mathbf{B} + \mathbf{B}_0)]_\alpha \right\} \frac{\partial f}{\partial v_\alpha} = - \frac{\partial}{\partial v_\alpha} J_\alpha, \quad (4.1)$$

with $\mathbf{E} = 0$ and $\mathbf{B} = 0$,

$$J_\alpha = \frac{\pi q^2 \ln \Lambda}{m} \sum_{l=1}^2 e_l^2 \int_{\mathbf{v}_l} \left\{ f \frac{\partial f_l}{m_l \partial v_{l,\beta}} - f_l \frac{\partial f}{m \partial v_\beta} \right\} \times \\ \times \frac{(u^2 \delta_{\alpha\beta} - u_\alpha u_\beta)}{u^3} d^3 \mathbf{v}_l. \quad (4.2)$$

Here $\mathbf{u} = \mathbf{v} - \mathbf{v}_l$ is the relative velocity, $d^3 \mathbf{v}_l$ corresponds to the integration over the whole velocity space of the plasma particles $l = 1, 2$. They are distributed by the Maxwellian function (3.13):

$$f_e(\mathbf{v}) = n_e \left(\frac{m_e}{2\pi k_B T_e} \right)^{3/2} \exp \left(- \frac{m_e v^2}{2k_B T_e} \right) \quad (4.3)$$

and

$$f_p(\mathbf{v}) = n_p \left(\frac{m_p}{2\pi k_B T_p} \right)^{3/2} \exp \left(- \frac{m_p v^2}{2k_B T_p} \right). \quad (4.4)$$

For the sake of simplicity we assume $T_e = T_p = T$ (see, however, Sect. 8.3.2) as well as $n_e = n_p = n$. Also for the sake of simplicity we shall consider the stationary situation ($\partial/\partial t = 0$).

Moreover we shall assume that the distribution function f depends on one spatial variable—the coordinate z measured along the field $\mathbf{B}_0$, on the value of velocity v and the angle θ between the velocity vector $\mathbf{v}$ and the axis z . Therefore

$$f = f(z, v, \theta). \quad (4.5)$$

In this case of the axial symmetry, the term containing the Lorentz force, related to the external field $\mathbf{B}_0$, in (4.1) is equal to zero because the vector $\mathbf{v} \times \mathbf{B}_0$ is perpendicular to the plane $(\mathbf{v}, \mathbf{B}_0)$ but the vector $\partial f / \partial \mathbf{v}$ is placed in this plane.

Under assumptions made above, (4.1) takes the following form:

$$v \cos \theta \frac{\partial f}{\partial z} = -\frac{1}{v^2} \frac{\partial}{\partial v} (v^2 J_v) - \frac{1}{v} \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} (\sin \theta J_\theta). \quad (4.6)$$

The distribution function f is *not* an isotropic one. So the angular component J_θ of the particle flux is not equal to zero.

4.1.2 Dimensionless Kinetic Equation in Energy Space

Let us introduce the dimensionless non-relativistic energy of the fast particles

$$x = \frac{mv^2}{2k_B T} \left(\frac{m_e}{m} \right) \quad (4.7)$$

and the dimensionless column depth along the magnetic field

$$\zeta = \xi / \tilde{\xi}. \quad (4.8)$$

Here

$$\xi = \int_0^z n(z) dz, \quad \text{cm}^{-2}, \quad (4.9)$$

is the dimensional column depth passed by the fast particles along the z axis; the unit of its measurement is

$$\tilde{\xi} = \frac{k_B^2 T^2}{\pi e^2 q^2 \ln \Lambda} \left(\frac{m}{m_e} \right)^2, \quad \text{cm}^{-2}. \quad (4.10)$$

Equation (4.6) in the dimensionless variables (4.7) and (4.8) takes the following form (Somov 1982):

$$\sqrt{x} \cos \theta \frac{\partial f}{\partial \zeta} = \frac{1}{\sqrt{x}} \frac{\partial}{\partial x} \left\{ \sqrt{x} D_\gamma(x) \left[\frac{\partial f}{\partial x} + \left(\frac{m}{m_e} \right) f \right] \right\} + D_\theta(x) \Delta_\theta f. \quad (4.11)$$

Here

$$\begin{aligned} D_\gamma(x) = & \left[\frac{\text{erf}(\sqrt{x})}{\sqrt{x}} - \frac{2}{\sqrt{\pi}} \exp(-x) \right] + \\ & + \left(\frac{m_e}{m_p} \right)^{1/2} \left[\frac{\text{erf}(\sqrt{\mathcal{X}})}{\sqrt{\mathcal{X}}} - \frac{2}{\sqrt{\pi}} \exp(-\mathcal{X}) \right] \end{aligned} \quad (4.12)$$

with

$$\mathcal{X} = \frac{m_p}{m_e} x$$

and

$$\operatorname{erf}(w) = \frac{2}{\sqrt{\pi}} \int_0^w \exp(-t^2) dt ,$$

which is the error function. The diffusion coefficient over the angle θ

$$D_\theta(x) = \frac{1}{8x^2} \left\{ \left[\frac{\operatorname{erf}(\sqrt{x})}{\sqrt{x}} (2x - 1) + \frac{2}{\sqrt{\pi}} \exp(-x) \right] + \left(\frac{m_e}{m_p} \right)^{1/2} \left[\frac{\operatorname{erf}(\sqrt{\mathcal{X}})}{\sqrt{\mathcal{X}}} (2\mathcal{X} - 1) + \frac{2}{\sqrt{\pi}} \exp(-\mathcal{X}) \right] \right\} , \quad (4.13)$$

and

$$\Delta_\theta = \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right)$$

is the θ -dependent part of the Laplace operator.

To point out the similarity of the equation obtained with the Fokker-Planck equation (3.10), let us rewrite (4.11) as follows:

$$\sqrt{x} \cos \theta \frac{\partial f}{\partial \xi} = -\frac{\partial}{\partial x} [F(x)f] + \frac{\partial^2}{\partial x^2} [D(x)f] + D_\theta(x) \Delta_\theta f. \quad (4.14)$$

Here the first coefficient

$$F(x) = \frac{dD_\gamma}{dx} - \left(\frac{m}{m_e} + \frac{1}{2x} \right) D_\gamma(x) \quad (4.15)$$

characterized the *regular losses* of energy when accelerated particles pass through the plasma. The second coefficient

$$D(x) = D_\gamma(x) \quad (4.16)$$

describes the *energy diffusion*. The third coefficient $D_\theta(x)$ corresponds to the fast particle diffusion over the pitch angle θ .

Kudriavtsev (1958) derived the time-dependent equation which has the right-hand side similar to the one in our (4.11) but for the isotropic distribution function $f = f(t, x)$ for fast ions in a thermal plasma. By using the Laplace transformation, Kudriavtsev solved the problem of maxwellization of fast ions that initially had the mono-energetic distribution $f(0, x) \sim \delta(x - x_0)$. The same problem has been solved numerically by MacDonald et al. (1957). (Note that in formula (8) by Kudriavtsev for the “radial” component j_v of the fast ion flow in the velocity space, the factor $\sqrt{\pi}$ must be in the nominator but not in the denominator.) Both solutions (analytical and numerical) show, of course, that the higher the ion energy, the longer the maxwellization process.

In the particular case when all the particles are the same ($m = m_e = m_p$), the right-hand side of (4.11) can be found, for example, by using the formulae for the Fokker-Planck coefficients (3.12) and (3.11) from Balescu (1963).

4.2 A Kinetic Equation at High Speeds

Bearing in mind the particles accelerated to high speeds in astrophysical plasma, let us consider some approximations and some solutions of the kinetic equation (4.11) that correspond to these approximations. First of all, we shall assume that the dimensionless energy (4.7) of the fast particles

$$x \gg 1. \quad (4.17)$$

This means that speeds of the particles are much higher than the mean thermal velocity of plasma electrons (8.15). However, for the sake of simplicity, we restrict the problem by consideration of the fast but non-relativistic particles.

Under condition (4.17), we obtain from (4.12) and (4.13) the following simple formulae for the coefficients in the kinetic equation (4.11):

$$D_\gamma(x) = \frac{1}{\sqrt{x}} \left(1 + \frac{m_e}{m_p} \right), \quad (4.18)$$

$$D_\theta(x) = \frac{1}{2x\sqrt{x}}. \quad (4.19)$$

It is *not* taken into account here yet that $m_e \ll m_p$. The first term on the right-hand side of formula for D_γ (see the unit inside the brackets) is a contribution of collisions with the thermal electrons of a plasma, the second term (see the ratio m_e/m_p) comes from collisions with the thermal protons. However the electrons and protons give equal contributions to the *angular diffusion* coefficient D_θ . This is important to see when we derive formula (4.19) from (4.13).

Under the same assumption, the Fokker-Planck type equation (4.14) has the following coefficients:

$$D(x) = \frac{1}{\sqrt{x}} \left(1 + \frac{m_e}{m_p} \right), \quad (4.20)$$

$$F(x) = -\frac{m}{m_e} \frac{1}{\sqrt{x}} \left(1 + \frac{m_e}{m} \frac{1}{x} \right), \quad (4.21)$$

and the same coefficient of angular diffusion $D_\theta(x)$ of course.

Formulae (4.18) and (4.20) demonstrate that

energy diffusion due to collisions with thermal electrons is faster in m_p/m_e times than that due to collisions with thermal protons.

However the angular diffusion rate is equally determined by both electrons and protons in a plasma.

The second term on the right-hand side of the formula for $F(x)$ describes the regular losses of fast particle energy by collisions with thermal protons of plasma.

Since $x \gg 1$ and $m \geq m_e$, this term is always smaller than the first one. Taking into account that $m_e \ll m_p$ we also neglect the second term in formula for $D(x)$. Hence, in approximation under consideration,

$$F(x) = -\frac{m}{m_e} \frac{1}{\sqrt{x}}, \quad D(x) = \frac{1}{\sqrt{x}}, \quad D_\theta(x) = \frac{1}{2x\sqrt{x}}. \quad (4.22)$$

Let us estimate a relative role of the first and second terms on the right-hand side of (4.14). Dividing the former by the last with account of (4.22) taken gives the ratio

$$\frac{x F(x)}{D(x)} = \frac{m}{m_e} x, \quad (4.23)$$

which is always much greater than unity. So, for fast particles with speeds much greater than the thermal velocity of plasma electrons,

the regular losses of energy due to Coulomb collisions always dominate the energy diffusion.

However the energy diffusion may appear significant near the lower energy boundary $\mathcal{E}_1$ of the fast particle spectrum if $\mathcal{E}_1 \approx k_B T$. This seems to be the case of electron acceleration in the so-called *super-hot turbulent-current* layers in *solar flares* (see Part II, Sects. 6.3 and 7.1). This simply means that, near the lower energy $\mathcal{E}_1 \approx 10$ keV, the initial assumption (4.17) becomes invalid. Instead of (4.17), $x \rightarrow 1$; so we have to solve exactly (4.11).

Let us compare the first and third terms on the right-hand side of (4.14). Dividing the former by the last with account of (4.22) taken gives the ratio

$$\frac{F(x)}{x D_\theta(x)} = 2 \frac{m}{m_e}. \quad (4.24)$$

For fast protons and heavier ions, we can neglect angular scattering in comparison with the regular losses of energy.

Formula (4.24) shows, however, that

for fast electrons, it is impossible to neglect the angular diffusion in comparison with the regular losses of energy.

Since the case of fast electrons will be considered later on in more detail, let us rewrite the non-relativistic kinetic equation in the high-speed approximation as follows:

$$\cos \theta \frac{\partial f}{\partial \zeta} = \frac{1}{x} \frac{\partial f}{\partial x} + \frac{1}{2x^2} \Delta_\theta f.$$

(4.25)

Recall that the energy diffusion is neglected in (4.25) according to (4.23).

4.3 The Classical Thick-Target Model

4.3.1 General Solution for Fast Electrons

We have seen that, in the fast electron kinetic equation (4.25), it is not reasonable to neglect the angular diffusion. Let us, however, consider the well-known and widely-used model of a *thick target*. From (4.25), by *neglecting* the angular diffusion, we obtain the following simple equation

$$\cos \theta \frac{\partial f}{\partial \xi} = \frac{1}{x} \frac{\partial f}{\partial x}. \quad (4.26)$$

With a new variable $y = \xi/\mu$, where $\mu = \cos \theta$, this equation becomes especially simple:

$$\frac{1}{x} \frac{\partial f}{\partial x} - \frac{\partial f}{\partial y} = 0. \quad (4.27)$$

General solution of this equation can be written as

$$f(x, y) = \mathcal{F}\left(\frac{x^2}{2} + y\right), \quad (4.28)$$

where $\mathcal{F}$ is an arbitrary function of its argument. Recall that $\mu = \text{const}$, because we have neglected the angular diffusion; so the fast electrons move along straight lines $\theta = \text{const}$ without any scattering.

Let us consider the initial ($y = 0$) energy distribution of fast electrons—the *injection spectrum*—as a power law:

$$f(x, 0) = c_0 x^{-\gamma_0} \Theta(x - x_1) \Theta(x_2 - x) p_0(\mu). \quad (4.29)$$

Here $\Theta(x)$ is the theta-function; $p_0(\mu)$ is the angular distribution of fast electrons, for example, for a beam of electrons injected parallel to the z axis

$$p_0(\mu) = \frac{1}{(1 - \mu^2)^{1/2}} \delta(\mu - 1). \quad (4.30)$$

According to (4.28) the general solution of the kinetic equation for the fast electrons at the column depth y has the following form:

$$f(x, y) = c_0 2^{-\gamma_0/2} \left(\frac{x^2}{2} + y\right)^{-\gamma_0/2} \Theta(x - x'_1) \Theta(x'_2 - x) p_0(\mu), \quad (4.31)$$

where

$$x'_{1,2} = \text{Re} (x_{1,2}^2 - 2y)^{1/2}.$$

4.3.2 The Injection Spectrum

Let us consider the normalization condition for the distribution function, first, in the dimensional variables z , v , and θ (see definition (4.5)). If $n_b(z)$ is the *density of electrons in the beam* at distance z from the injection plane $z = 0$, then

$$n_b(z) = \int_0^\infty \int_0^\pi f(z, v, \theta) v^2 dv 2\pi \sin \theta d\theta, \text{ cm}^{-3}. \quad (4.32)$$

It is taken into account here that we consider the case of a beam with the axial symmetry in velocity space (see Sect. 4.1.1).

Now we rewrite the same normalization condition in the dimensionless variable ζ , x , and μ :

$$n_b(\zeta) = \pi \left(\frac{2k_B T}{m_e} \right)^{3/2} \int_0^\infty \int_{-1}^1 f(\zeta, x, \mu) \sqrt{x} dx d\mu, \text{ cm}^{-3}. \quad (4.33)$$

For initial energy distribution (4.29) and initial angular distribution (4.30), formula (4.33) gives

$$n_b(0) = \pi \left(\frac{2k_B T}{m_e} \right)^{3/2} c_0 \int_{x_1}^{x_2} x^{-\gamma_0+1/2} dx \equiv \int_{x_1}^{x_2} N(0, x) dx, \text{ cm}^{-3}. \quad (4.34)$$

Here

$$N(0, x) = \pi \left(\frac{2k_B T}{m_e} \right)^{3/2} c_0 x^{-\gamma_0+1/2} \Theta(x - x_1) \Theta(x_2 - x) \quad (4.35)$$

is the differential spectrum of the fast electron density at the boundary $\zeta = 0$ where they are injected.

Let $\mathcal{E}$ be the kinetic energy of a fast electron measured in keV. Then we rewrite (4.35) as

$$N(0, \mathcal{E}) = K \mathcal{E}^{-(\gamma+1/2)} \Theta(\mathcal{E} - \mathcal{E}_1) \Theta(\mathcal{E}_2 - \mathcal{E}), \text{ cm}^{-3} \text{ keV}^{-1}, \quad (4.36)$$

where the coefficient

$$K = \pi \left(\frac{2k_B T}{m_e} \right)^{3/2} c_0 \left(\frac{k_B T}{\text{keV}} \right)^{\gamma+1/2}, \quad \text{cm}^{-3} \text{keV}^{\gamma-1/2}, \quad (4.37)$$

and the spectral index

$$\boxed{\gamma = \gamma_0 - 1.} \quad (4.38)$$

Hence the *injection* spectrum of fast electrons is determined by parameters (4.37) and (4.38).

4.3.3 Fast Electrons Inside a Target

Substituting c_0 and γ_0 from (4.37) and (4.38) in (4.31) allows us to obtain the differential spectrum of the number density of fast electrons passed the column depth ξ measured in cm^{-2} according to definition (4.9); see Syrovatskii and Shmeleva (1972):

$$N(\xi, \mathcal{E}) = K \mathcal{E}^{1/2} (\mathcal{E}^2 + \mathcal{E}_0^2)^{-(\gamma+1)/2} \times \\ \times \Theta(\mathcal{E} - \mathcal{E}'_1) \Theta(\mathcal{E}'_2 - \mathcal{E}), \quad \text{cm}^{-3} \text{keV}^{-1}. \quad (4.39)$$

Here

$$\mathcal{E}_0 = (2a_0\xi)^{1/2} \quad (4.40)$$

is the minimal energy of electrons that can pass the depth ξ , the “constant” a_0 (a slow function of energy $\mathcal{E}$) originates from the Coulomb logarithm and equals

$$a_0 = 2\pi e^4 \ln \Lambda \approx \\ \approx 1.3 \times 10^{-19} \times \left[\ln \left(\frac{\mathcal{E}}{mc^2} \right) - \frac{1}{2} \ln n + 38.7 \right], \quad \text{keV}^2 \text{cm}^2. \quad (4.41)$$

In formula (4.39)

$$\mathcal{E}'_{1,2}(\xi) = (\mathcal{E}_{1,2}^2 - \mathcal{E}_0^2(\xi))^{1/2} \quad (4.42)$$


are the new boundaries of energetic spectrum, when the fast electrons have passed the column depth ξ .

Solution (4.39) shows that

the regular losses of energy due to Coulomb collisions shift the spectrum of fast electrons to lower energies and make it harder

as illustrated by schematic Fig. 4.1; see also Exercises 4.1 and 4.2. Both effects follow from the fact that, in (4.26), we have taken into account *only the regular losses* of energy (4.22). For non-relativistic electrons $F(x) = -1/\sqrt{x}$.

Fig. 4.1 An injection spectrum ($\xi = 0$) and the spectrum of fast electrons that have passed the column depth ξ inside a target



4.3.4 Hard X-Ray Bremsstrahlung

In the solar system, the Sun is the most energetic particle accelerator, producing electrons of up to tens of MeV and ions of up tens of GeV. The accelerated 20–100 keV electrons appear to contain a significant part of the total energy of a large solar flare (Lin and Hudson 1971; Syrovatskii and Shmeleva 1972), indicating that the particle acceleration and energy release processes are intimately linked. Flare accelerated electrons colliding with the ambient solar atmosphere produce the *bremsstrahlung* hard X-ray (HXR) emission.

The question of the spectrum of the electrons accelerated in a flare in connection with the solar HXR bursts was posed at first by Peterson and Winckler (1959). Syrovatskii and Shmeleva (1972) used the solution (4.39) to calculate the HXR bremsstrahlung which arises during inelastic collisions of accelerated electrons with thermal ions in the solar atmosphere during flares (e.g., Strong et al. 1999). Brown (1971), in a less accurate approximation and using a different method, has found a similar formula for HXR intensity but with the different numerical coefficient by factor π in his Sect. 4.5 (see formulae (14) and (15)). Anyway, since that time,

the simplest thick-target model is widely accepted as a likely mechanism and an approximate tool to explain and describe the HXR emission observed during flares

on the Sun and other stars or generally in astrophysical plasma (see, however, Sects. 4.4 and 4.5). In the classical formulation of the thick-target model, beams of accelerated electrons stream along the magnetic field lines and lose their energy by Coulomb collisions in denser layers of the solar atmosphere, mainly in the chromosphere.

4.4 The Role of Angular Diffusion

4.4.1 An Approximate Account of Scattering

As we have seen in Sect. 4.2, for fast electrons, we cannot neglect the angular scattering in comparison with the regular losses of energy in kinetic equation (4.14). Hence, in the classical thick-target model, we have to take the angular scattering into account at least approximately.

If, for example, the beam of fast electrons penetrates a plane-parallel stratified plasma such as the solar chromosphere should be according to a classical definition, the scattering of an *average* beam of electrons may conveniently be described by the Chandrasekhar-Spitzer formulae (8.56) and (8.57) in terms of a coordinate z normal to the atmospheric strata and directed into the plasma. Then the *mean* electron energy $\mathcal{E}$ may be expressed as a function of z while the scattering is measured in terms of the angle $\theta(z)$ which the *mean* electron velocity $\mathbf{v}$ makes with the z axis at that point. So

$$v_{\parallel} \equiv v_z = v\mu, \quad \text{where} \quad \mu = \cos \theta. \quad (4.43)$$

The dimensional column depth passed by electrons along the z axis is

$$\xi = \int_0^z n(z) dz, \quad \text{cm}^{-2}. \quad (4.44)$$

In terms of ξ , the Chandrasekhar-Spitzer formulae (8.56) and (8.57) are:

$$\frac{d\mathcal{E}}{d\xi} = -\frac{a_0}{\mathcal{E}} \frac{v}{v_z} \quad (4.45)$$

and

$$\frac{dv_z}{d\xi} = -\frac{3}{2} \frac{a_0}{\mathcal{E}^2} v, \quad (4.46)$$

where $a_0 = 2\pi e^4 \ln \Lambda$ (see definition (4.41)). Thus we have an ordinary differential equation

$$\frac{3}{2} \frac{1}{\mathcal{E}} \frac{d\mathcal{E}}{d\xi} = \frac{1}{v_z} \frac{dv_z}{d\xi}$$

with solution

$$\left(\frac{\mathcal{E}}{\mathcal{E}_0} \right)^{3/2} = \frac{v_z}{v_{z0}}, \quad (4.47)$$

where the suffix 0 refers to values at $\xi = 0$. Since $v_z/\mu = v$ and $v^2/v_0^2 = \mathcal{E}/\mathcal{E}_0$, we find that

$$\frac{v_z}{v_{z0}} = \frac{\mu}{\mu_0} \left(\frac{\mathcal{E}}{\mathcal{E}_0} \right)^{1/2}.$$

Therefore it follows from (4.47) that

$$\boxed{\frac{\mu}{\mu_0} = \frac{\mathcal{E}}{\mathcal{E}_0}}. \quad (4.48)$$

This nice formula (Brown 1972) shows that on average when an electron has suffered a 60° deflection its energy has been reduced by 50%.

Substituting (4.48) in (4.45) and (4.46) gives the solutions for μ and $\mathcal{E}$:

$$\frac{\mu}{\mu_0} = \frac{\mathcal{E}}{\mathcal{E}_0} = \left(1 - \frac{3a_0\xi}{\mu_0\mathcal{E}_0^2} \right)^{1/3}. \quad (4.49)$$

For small depth ξ

$$\frac{\mu}{\mu_0} = \frac{\mathcal{E}}{\mathcal{E}_0} \approx 1 - \frac{a_0}{\mu_0\mathcal{E}_0^2} \xi. \quad (4.50)$$

Let us compare these results with the general solution (4.28) obtained without account taken of scattering in the classical thick-target model.

4.4.2 The Thick-Target Model

According to (4.28)

$$\frac{x^2}{2} + y = \frac{x_0^2}{2}, \quad (4.51)$$

where x_0 is an initial energy of an electron. Hence

$$\frac{x}{x_0} = \left(1 - \frac{2y}{x_0^2} \right)^{1/2}, \quad (4.52)$$


where $y = \zeta/\mu$ and $\mu = \text{const} = \mu_0$. Therefore for electrons with initial energy $\mathcal{E}_0$ solution (4.28) gives us:

$$\frac{\mathcal{E}}{\mathcal{E}_0} = \left(1 - \frac{2a_0}{\mu_0\mathcal{E}_0^2} \xi \right)^{1/2}. \quad (4.53)$$

If

$$\xi \ll \xi_0 = \frac{\mathcal{E}_0^2}{2a_0},$$

Fig. 4.2 The mean energy $\mathcal{E}$ of the fast electrons that have passed the column depth ξ (from Somov 1982)



then

$$\frac{\mathcal{E}}{\mathcal{E}_0} \approx 1 - \frac{a_0}{\mu_0 \mathcal{E}_0^2} \xi. \quad (4.54)$$

Formula (4.54) coincides with (4.50). The fast electrons in the thick-target model have the same behavior at small depth ξ as that one predicted by the approximate Chandrasekhar-Spitzer formulae.

However, with increase of the column depth ξ , the approximate formula (4.49) predicts much faster losses of energy in comparison with the classical thick-target model which does not take collisional scattering into account.

In Fig. 4.2, the dashed straight line (a) corresponds to the asymptotic formula (4.50) which is valid for small column depth ξ . Moreover here $\mu_0 = 0$, so

$$\frac{\mathcal{E}}{\mathcal{E}_0} \approx 1 - \frac{1}{2} \frac{\xi}{\xi_0}. \quad (4.55)$$

The solid curve (b) represents the *classical thick-target model*; it takes only the collisional losses of energy into account. With $\mu_0 = 0$, formula (4.53) is

$$\frac{\mathcal{E}}{\mathcal{E}_0} = \left(1 - \frac{\xi}{\xi_0}\right)^{1/2}. \quad (4.56)$$

An approximate scattering model described above is presented by the curve (c) which corresponds to formula (4.49) with $\mu_0 = 0$, so

$$\frac{\mathcal{E}}{\mathcal{E}_0} = \left(1 - \frac{3}{2} \frac{\xi}{\xi_0}\right)^{1/3}. \quad (4.57)$$

Figure 4.2 shows that

the collisional scattering and energy losses become very great in comparison with the classical thick-target model if the column depth ξ is not very small.

Brown (1972) used the approximate formula (4.49) to develop an approximate thick-target model in which accelerated electrons penetrate downward into the solar chromosphere during a flare. Here the electron distribution is greatly modified by collisions—not only by energy losses but also by scattering. *Directivity and polarization* of the hard X-ray bremsstrahlung emission have been calculated in such oversimplified thick-target model in which the guiding field $\mathbf{B}_0$ is vertical. The model predicted that the degree of polarization should rise from zero to around 30% near the solar limb.

Unfortunately the accuracy of the model decreases when the collisional scattering and energy losses become not small. The reason is that the mean rates (4.45) and (4.46) represent well the modification of the electron velocity distribution only at small depth ξ . A more accurate formulation of the kinetic problem will be given in the next section with account taken of the collisional scattering and one more mechanism of the electron beam anisotropization. Generally, it seems true that the total absorption of the accelerated electrons in a thick target might result in *negligible* directivity and polarization of the hard X-ray emission.

4.5 The Reverse-Current Electric-Field Effect

4.5.1 The Necessity for a Beam-Neutralizing Current

We assume that some external magnetic field $\mathbf{B}_0$ channels a fast particle propagation and can be locally considered as uniform. The electric and magnetic fields $\mathbf{E}$ and $\mathbf{B}$ related to a *beam* of fast electrons are superposed on this field. In this way, the beam will be considered as a real electric current $\mathbf{J}$ which influences the background plasma and magnetic field $\mathbf{B}_0$. In order not to obscure the essential physical points related to the electromagnetic field of the beam, we shall neglect all other processes like the radiative and hydrodynamic response of the background plasma to a fast heating by the electron beam (Sect. 8.3.2).

In the classical thick-target model for hard X-ray bremsstrahlung emission during solar flares, if the fast *accelerated electrons* are supposed to have about the parallel velocities, then the number of injected beam particles per unit time has to be very large—in the order of $\gtrsim 10^{36}$ electrons s^{-1} above 25 keV during the *impulsive phase* of a flare (Hoyng et al. 1976). Given the large electron fluxes implied by the hard X-ray observations, various authors realized that the beam electric current must be enormously large: $J \gtrsim 10^{17}$ A.

This huge value would imply the magnetic field of the beam $B \gtrsim 10^5$ G. So the magnetic energy contents of a flare volume in the solar corona should be more than

six orders of magnitude larger than the pre-flare contents for an average coronal field $B_0 \approx 100$ G. Such situation is not likely to occur because the electron beams are thought to be created by conversion of the magnetic energy available in the corona into kinetic energy.

Apart from this energy problem there is another difficulty related to beams of $\sim 10^{36}$ electrons s^{-1} ; they create an enormous charge displacement. For a typical coronal volume of 10^{28} cm^3 and an electron density 10^9 cm^{-3} , the total number of electrons is 10^{37} . A stream of 10^{36} electrons s^{-1} would evacuate all the electrons out of the volume in about 10 s. As a result an enormous charge difference between the corona and the chromosphere would be build up.

In reality the above mentioned problems will not occur, because the beam propagates in a background *well-conducting plasma*. The charge displacement by the beam will quickly create an electric field $\mathbf{E}_1$ which causes the plasma electrons to redistribute in such a way as to *neutralize* the local charge built:

$$\text{div } \mathbf{E}_1 = 4\pi\rho^q. \quad (4.58)$$

Because this electric field is caused by charge separation, it is frequently referred to as an *electrostatic* field. So the electrostatic effect allows the plasma in the solar corona to “absorb” the excess charge imposed by the beam of fast electrons accelerated in solar flare.

The second effect is related to the inductive properties of a well-conducting plasma. In such a plasma the magnetic field will not vary considerably on a timescale shorter than the *magnetic diffusion* time. For beams with radii comparable to the radii of coronal flaring loops this scale is much longer than the duration of the impulsive phase. When the current varies in magnitude, an *inductive* electric field $\mathbf{E}_2$ will be immediately created. This field drives a current $\mathbf{j}_2$ of plasma electrons in such a way to prevent magnetic field variations on a time scale shorter than the magnetic diffusion time. As a result the magnetic field will not vary much during the impulsive phase:

$$\text{rot } \mathbf{B} \approx \text{const} \approx 0 \approx \frac{4\pi}{c} \mathbf{j}_2 + \frac{1}{c} \frac{\partial}{\partial t} \mathbf{E}_2. \quad (4.59)$$

So the inductive effect tends to prevent the magnetic field in the solar corona from changing faster than the allowed diffusion time.

Both the electrostatic and the inductive electric field will effectively result in an electron plasma current which is in opposite direction of the beam current $\mathbf{J}$.

This electron plasma current is commonly referred to as the *reverse* or *return* current $\mathbf{J}_{rc}$.

van den Oord (1990) has analyzed the electrostatic and inductive response of a plasma to a prescribed electron beam. By using the Maxwell equations together with the time-dependent Ohm’s law (see Sect. 11.2) and with the equation of motion for

the plasma electrons in the hydrodynamic approximation (Sect. 9.5), he has shown that the non-linear terms are responsible for a coupling between the electrostatic (irrotational) and inductive (solenoidal) vector fields generated by the beam in a plasma. In order to obtain analytical solutions, van den Oord has decoupled the electrostatic and inductive fields, by ignoring the non-linear terms in the equation of motion, and has found solutions for a mono-energetic *blunt* beam.

An application of the van den Oord model in conditions of the solar corona leads to the following results. Charge neutralization is accompanied by plasma oscillations (see formula (8.35)), that are present behind the beam front, and occurs on a time-scale of a few electron-ion collision times. This is also the time scale on which the *plasma waves* damp out. The net current in the system quickly becomes too low and therefore also the resulting magnetic field strength remains low ($B \ll B_0$).

Although the electric field near the beam front is locally strong, the oscillatory character prevents strong acceleration of the plasma electrons. According to the van den Oord model, all the beam energy is used initially to accelerate the plasma electrons from rest and later on to drive the reverse current against collisional losses. In what follows, we shall use these results and shall formulate an opposite problem in the kinetic approximation. We shall not consider the beam as prescribed. On the contrary, we shall consider *an influence of the electric field*, which drives the reverse current, *on the distribution function of fast electrons* in the thick-target plasma.

4.5.2 Formulation of a Realistic Kinetic Problem

The *direct* electric current carried by the fast electrons is equal to

$$j_{dc}(z) = e \int_{\mathbf{v}} f(v, \theta, z) v \cos \theta d^3 \mathbf{v}. \quad (4.60)$$

In order to make the problem easier to solve, we shall consider this current to be fully balanced by the *reverse* current, j_{rc} , of the thermal electrons in the ambient plasma,

$$|j_{dc}(z)| = j_{rc}(z) \equiv j(z). \quad (4.61)$$

This simplification means that here we do not consider a very fast process of the reverse current generation. The time-dependent process of current neutralization, with account of both electrostatic and inductive effects taken (Sect. 4.5.1), has been investigated in linear approximation by van den Oord (1990) in the frame of his model. Instead of that we shall construct a self-consistent approach for solving the pure kinetic problem with a steady electric field $E = E(z)$ which drives the reverse current.

So, using Ohm's law, we determine the reverse-current electric field to be equal to

$$E(z) = \frac{j(z)}{\sigma}. \quad (4.62)$$

Here σ is conductivity of the plasma; we can assume that the conductivity is determined by, for example, Coulomb collisions (Sect. 11.1). This is the case of a cold dense astrophysical plasma, for example, solar chromosphere.

On the other hand, the plasma turbulence effects are also important, for example, in the heat conductive front between the high-temperature source of energy and cold plasma of the thick-target. Anyway, even though we expect the wave-particle interactions to have some effects on the fast electrons (Chap. 7), it is unlikely that such effects can change significantly the distribution function of fast electrons with energies far exceeding the energies of the particles in a background cold plasma.

What is really important is the *reverse-current electric field*, it results in an essential change of the fast electron behaviour in the plasma. That is why, to solve the thick-target problem, we develop a combined approach which takes into account the electric field (4.62) as in the Vlasov equation and Coulomb collisions as in the Landau equation. So the distribution function for the fast electrons in the target is described by the following equation (Diakonov and Somov 1988):

$$v \cos \theta \frac{\partial f}{\partial z} - \frac{eE(z)}{m_e} \cos \theta \frac{\partial f}{\partial v} - \frac{eE(z)}{m_e v} \sin^2 \theta \frac{\partial f}{\partial \cos \theta} = \left(\frac{\partial f}{\partial t} \right)_c. \quad (4.63)$$

Here the second and the third terms are the expression of the term

$$\frac{e_e}{m_e} \mathbf{E}(\mathbf{r}) \frac{\partial f}{\partial \mathbf{v}}$$

in the dimensional variables v and θ . On the right-hand side of (4.63)


$$\begin{aligned} \left(\frac{\partial f}{\partial t} \right)_c &= \frac{1}{v^2} \frac{\partial}{\partial v} \left[v^2 \nu(v) \left(\frac{k_B T_e}{m_e} \frac{\partial f}{\partial v} + \nu f \right) \right] + \\ &+ \nu(v) \frac{\partial}{\partial \cos \theta} \left(\sin^2 \theta \frac{\partial f}{\partial \cos \theta} \right) \end{aligned} \quad (4.64)$$

is the *linearized* collisional integral; $\nu(v)$ is the collisional rate for fast electrons in the cold plasma.

To set the mathematical problem in the simplest form (see Fig. 4.3), we assume that “super-hot” ($T_{e,0} = T_0 \gtrsim 10^8$ K) and “cold” ($T_{e,1} = T_1 \sim 10^4 - 10^6$ K $\ll T_0$) plasmas occupy the two half-spaces separated by the plane turbulent front ($z = 0$). The super-hot region represents the source of energy, for example, the high-temperature reconnecting current layer (RCL) in a solar flare. Let

$$f_s = f_s(v, \theta) \quad (4.65)$$

Fig. 4.3 The fast electron propagation in a thick-target cold plasma. TF is the turbulent front between the super-hot source of fast electrons and the cold plasma



be the electron distribution function in the *source*. f_s is, for example, the Maxwellian function for the case of thermal electron runaway (Diakonov and Somov 1988) or a superposition of thermal and non-thermal functions in the general case. To study the effect of the reverse-current electric field in the classical thick-target model, Litvinenko and Somov (1991b) considered only accelerated electrons with an energetic power-law spectrum. Anyway, the function f_s is normalized to the electron number density n_0 in the source:

$$\int f_s(v, \theta) d^3\mathbf{v} = n_0. \quad (4.66)$$

Because the *electron runaway* in a turbulent plasma (Gurevich and Zhivlyuk 1966) is similar to the ordinary collisional runaway effect (Sect. 8.4.3), the electrons with velocities

$$v_e > v_{cr}, \quad (4.67)$$

where v_{cr} is some critical velocity, can freely penetrate through the turbulent front into the cold plasma. Electrons with lower velocities remain *trapped* in the source. In this section, we are going to consider the distribution function for the fast electrons escaping into the cold plasma and propagating there. The boundary condition for the forward-flying (the suffix *ff*) fast electrons may be taken as

$$f_{ff}(v, \theta, 0) = f_s(v, \theta) \Theta(v - v_{cr}), \quad 0 \leq \theta \leq \pi/2, \quad (4.68)$$

where Θ is the theta-function.

The distribution function for the *back-flying* electrons is determined from the solution of (4.63) everywhere, including the boundary $z = 0$. Therefore the problem has been formulated. Note the obvious but important thing; (4.63) contains *two unknown functions*: the fast electron distribution function $f(v, \theta, z)$ and the electric field $E(z)$. So the kinetic equation (4.63) must be solved together with (4.60)–(4.62). This is the complete set of equations to be solved self-consistently.

4.5.3 Dimensionless Parameters of the Problem

In the dimensionless variables (4.7), (4.8) and $\mu = \cos \theta$, (4.63) takes the form

$$\mu x^2 \frac{\partial f}{\partial \xi} - 2\varepsilon \mu x^2 \frac{\partial f}{\partial x} - \varepsilon x (1 - \mu^2) \frac{\partial f}{\partial \mu} = x \frac{\partial f}{\partial x} + \tau x \frac{\partial^2 f}{\partial x^2} + \frac{1}{2} \Delta_\mu f. \quad (4.69)$$

Here the dimensionless electron energy

$$x = \frac{m_e v^2}{2k_B T_0} \quad (4.70)$$

is normalized with the temperature T_0 of the super-hot plasma; for example, $T_0 = T_{e,eff} \approx 100$ MK is an effective electron temperature of the *super-hot turbulent-current* layer (SHTCL, see Part 2, Sect. 6.3). The ratio of the cold-to-super-hot plasma temperature

$$\tau = \frac{T_1}{T_0} \approx 10^{-4}, \quad (4.71)$$

if we consider as example the injection of fast electrons into the solar chromosphere. The dimensionless column depth ξ (see definition (4.8)) equals the dimensional column depth passed by fast electrons

$$\xi = \int_0^z n(z) dz, \text{ cm}^{-2}, \quad (4.72)$$

divided by the unit of its measurement

$$\tilde{\xi} = \frac{k_B^2 T_0^2}{\pi e^4 \ln \Lambda}, \text{ cm}^{-2}. \quad (4.73)$$

The dimensionless electric field

$$\varepsilon = \frac{E}{E_{D,1}} \frac{2}{\tau}, \quad (4.74)$$

where

$$E_{D,1} = \frac{4\pi e^3 \ln \Lambda}{k_B} \frac{n_1}{T_1} \quad (4.75)$$

is the *Dreicer field* in the cold plasma of the target (cf. definition (8.75)).

The parameter ε can be found from the self-consistent solution of the complete set of equations and the boundary conditions as described in Sect. 4.5.2.

The parameter ε is not small in a general case and, in particular, in the solar flare problem $\varepsilon \approx 2 - 20$ (see Fig. 4 in [Diakonov and Somov 1988](#)). Therefore, from (4.74)

$$E = \varepsilon \frac{\tau}{2} E_{D,1} \approx (10^{-4} - 10^{-3}) E_{D,1}, \quad (4.76)$$

so Ohm's law (4.62) is well applicable in this case. However

the physical effects related to a reverse-current electric field in the solar chromosphere are not small

(see Exercise 4.3).

Let us set the specific form of the boundary distribution function (4.68). The processes of *electron acceleration* in astrophysical plasma and their heating are always closely related. However, for the sake of contrast of them to each other, we consider separately two different functions.

- (a) We shall suppose that the electron distribution in the super-hot plasma is near to the Maxwellian one. So the distribution function

$$f_s(x, \mu) = n_0 c_0 \exp(-x) h(\mu), \quad \mu \geq 0, \quad (4.77)$$

with the constant

$$c_0 = \left(\frac{m_e}{2\pi k_B T_0} \right)^{3/2}.$$

- (b) For accelerated electrons we shall use the power-law spectrum as the boundary distribution function for the forward-flying electrons

$$f_{ff}(x, \mu) = f_s(x, \mu) \Theta(\mu - 1) = n_0 c_0 x^{-\gamma} h(\mu), \quad \mu \geq 0, \quad (4.78)$$

with another normalization constant c_0 . In principle, the function $h(\mu)$ is indefinite but should satisfy some additional conditions; at least the function $h(\mu)$ should be maximally smooth ([Diakonov and Somov 1988](#)).

4.5.4 Coulomb Losses of Energy

4.5.4.1 Electric Current in the Thick Target

In (4.69), the term $\tau x (\partial^2 f / \partial x^2)$ describes the *energy diffusion*. As we know from Sect. 4.2, for fast electrons with velocities much greater than the thermal velocity of plasma electrons, the *regular losses* of energy due to Coulomb collisions always dominate the energy diffusion. So we neglect this term in comparison with the term $x (\partial f / \partial x)$.

However, as we also know from Sect. 4.2, we cannot neglect the term with the μ -dependent part $\Delta_\mu f$ of the differential operator Laplacian Δ . This term is responsible for the *angular diffusion* of electrons and is not small in comparison to the regular losses term $x (\partial f / \partial x)$.

Therefore we can ignore only the term with small parameter τ in (4.69). After that we have

$$\mu x^2 \frac{\partial f}{\partial \zeta} = 2\varepsilon \mu x^2 \frac{\partial f}{\partial x} + \varepsilon x (1 - \mu^2) \frac{\partial f}{\partial \mu} + x \frac{\partial f}{\partial x} + \frac{1}{2} \frac{\partial}{\partial \mu} \left[(1 - \mu^2) \frac{\partial f}{\partial \mu} \right]. \quad (4.79)$$

By using this equation, we would like to obtain the equation which determines the behavior of the *direct electric current* carried by fast electrons in the target. It follows from definition (4.60) that

$$j_{dc}(\zeta) = 2\pi e \left(\frac{2k_B T_0}{m_e} \right)^2 \int_0^\infty \int_{-1}^{+1} f(x, \mu, \zeta) x \mu dx d\mu. \quad (4.80)$$

So we have to divide (4.79) by x and to integrate it as in formula (4.80).

All terms on the right-hand side of (4.79), except one, give zero contributions. The only term $x (\partial f / \partial x)$, describing the regular energy losses due to Coulomb collisions, determines the changes of electric current

$$j(\zeta) = j_{dc}(\zeta) = j_{rc}(\zeta) \quad (4.81)$$

along the column depth ζ into the target. It gives the right-hand side of the equation:

$$\frac{dj}{d\zeta} = -c_j \int_{-1}^{+1} f(x, \mu, \zeta) d\mu \quad (4.82)$$

with constant

$$c_j = \pi e \left(\frac{2k_B T_0}{m_e} \right)^2. \quad (4.83)$$

The physical meaning of (4.82) is that

fast electrons lose their energy and mix with thermal particles of the ambient cold plasma just due to Coulomb collisions.

Thus the self-consistent reverse-current problem demands to consider the term $x (\partial f / \partial x)$, describing the Coulomb energy losses; see also Exercise 4.4.

4.5.4.2 2D Versus 1D Models for the Thick Target

Equation (4.82) shows that the electric current $j(\zeta)$ decreases along the column depth ζ into the target because of the “falling out” of “completely” stopped ($x = 0$) electrons from the distribution function owing to collisional losses of energy. From the electric current continuity equation it follows that a current change is possible only when there are electron “sources” and/or “sinks” in the thick target.

In the energy region where (4.69) is valid ($x \gg \tau$), the collisional *friction force* (Sect. 8.4.1) is inversely proportional to x . For this reason, the electrons with low energies quickly slow down to energies of the order of τ and thus mix with the thermal electrons in the ambient plasma. Since in (4.79) formally $\tau = 0$, the “falling out” takes place under $x = 0$ according to formula (4.82).

The models under consideration in this chapter, except the classical thick-target model in Sect. 4.3, are *two-dimensional* (2D) *in the velocity space* (see definition (4.5)). This fact has an important consequence.

Some electrons after injection into the thick target make a curve trajectory and cross the boundary in the reverse direction without significant losses of energy.

These electrons come back to the source (the place of acceleration) without being stopped in the target; they determine the boundary distribution function for *back-flying* electrons and constitute a *significant* part (possibly the bulk) of all injected electrons.

Such a process is impossible in one-dimensional (1D) models, like the classical thick-target model, because an electron cannot change the initial direction to the opposite one without being stopped to zeroth velocity and accelerated by the *reverse-current electric field* from the zeroth velocity in the reverse direction. So collisional losses of energy are involved twice in the 1D dynamics of all fast electrons stopped in the target. In general,

the 1D (in velocity space) kinetic models taking Coulomb collisions into account are non-physical approximations.

The other group of injected electrons considered in 2D models is composed of the fast electrons which, after moving in the target under electrostatic and friction forces, do not come back in the particle source. With suitable values of energy x and angle θ , they lose a lot of their initial energy and stop their motion in the target not far from the boundary. There seem to be small amounts of such particles. They determine the electric current change. Thus the current $j(\zeta)$ and, hence, the electric field $E(\zeta)$ can change slowly near the boundary.

Among the particles that determine the current, we may choose a small subgroup of fast electrons which penetrate to such a depth into the target where the electric field is very small ($\varepsilon \ll 1$) and further on they are moving affected only by collisions. Even for this small subgroup the 2D models are certainly more realistic in comparison with the 1D models which do not take into account the collisional scattering (Sect. 4.4).

4.5.5 *New Physical Results*

Usually, in order to solve the 2D (in velocity space) kinetic equation, one have to develop a complicated numerical method. [Diakonov and Somov \(1988\)](#) have developed a new technique to obtain an *approximate analytical* solution of (4.63) taking the Coulomb collisions and the reverse-current electric field into account. They have applied this technique to the case of thermal *runaway* electrons in solar flares. It appears that the reverse-current field leads to a *significant reduction of the convective heat flux* carried by fast electrons escaping from the high-temperature plasma to the cold one ([Diakonov and Somov 1990](#)).

It is not justified to exclude the reverse-current electric-field effects in studies of convective heat transport by fast thermal electrons in astrophysical plasma, for example, in solar flares.

[Litvinenko and Somov \(1991b\)](#) have used the same technique to study the behavior of the electrons accelerated inside a reconnecting current layer (RCL) in the solar atmosphere during flares. They have shown that the reverse-current electric field results in an essential change of the fast electron behavior in the thick target.

The reverse-current electric field leads to a *quicker* decrease of the distribution function with the column depth in comparison with the classical thick-target model.

It is worth mentioning here that both models (thermal and non-thermal) lead to practically the same value of the field near the boundary, ε_0 , and this value is large: $\varepsilon_0 \gg 1$. So the effects of the reverse-current field are not small.

The distribution function appears to be an *almost isotropic* one. The main part of the injected electrons returns into the source. As a result, the hard X-ray polarization appears much smaller than that calculated in the collisional thick-target model without taking account of the reverse current. In calculations by [Litvinenko and Somov \(1991b\)](#), the maximum polarization was found to be of about 4% only; see also [Gritsyk and Somov \(2011\)](#), [Somov and Gritsyk \(2012\)](#). The computed degree of polarization is one order of magnitude less than that obtained in the simplified thick-target models and appreciably less than the degree of polarization obtained in the 2D (in velocity space) collisional models without the reverse current.

So a major conclusion of this section is that

in order to have a more precise insight into the problem of electron acceleration in solar flares, we inevitably have to take into account the reverse-current electric-field effects.

They make the accelerated electron distribution to be almost isotropic and leads to a significant decrease of expected hard X-ray bremsstrahlung polarization ([Somov and Tindo 1978](#)).

4.5.6 Plasma Heating by a Reverse Current

The reverse current can play the significant role in the heating of the solar atmosphere by accelerated electrons. [Sermulyan'sh and Somov \(1982, 1983\)](#) compared the Coulomb energy losses of an electron beam with the Joule heating of the solar plasma by the reverse current dissipation.

The heating by reverse current depends on the temperature and density distribution in the target plasma.

Therefore we have to obtain a self-consistent solution of the hydrodynamic problem taking account of both heating mechanisms as well as the kinetic problem considered above.

It appears that the Joule heating of the chromosphere by a reverse current can be larger than the direct heating by the beam of accelerated electrons in the upper chromosphere. The reverse current makes the accelerated electrons lose more energy in the uppermost chromospheric layers (for more detail see Figs. 2.1.23 and 2.1.24 in [Somov 1992](#)). The energy losses are larger (as compared to the Coulomb losses alone) the larger is the beam energy flux and the steeper is the beam spectrum. The reverse current heating can also become considerable in the low-temperature regions since plasma resistivity is large there.

Considering the problem of solar atmosphere heating by a reverse current, we should keep in mind the possibility that

plasma turbulence and anomalous resistivity can develop in low-density regions in the corona and upper chromosphere,

which would increase the effective resistance of the target plasma to the reverse current and thus increase the plasma heating ([Cromwell et al. 1988](#)).

Whether the resistivity is normal or anomalous, the reverse-current electric field hinders penetration of fast electrons into the low chromosphere and increases energy losses in its upper layers. It is important that the heat conduction mechanism of energy transfer is active here and that the kinetic energy of a beam is transformed to the heat flow energy in a collisionless process. Thus it is not strange that the non-thermal hard X-ray (HXR) flux observed in solar flares by the *RHESSI* (the Reuven Ramaty High Energy Solar Spectroscopic Imager) telescope asymptotically approaches some limiting value with increasing flare intensity. [Alexander and Daou \(2007\)](#) argue that such saturation of the bremsstrahlung HXR production sets in as a result of the transition from Coulomb collisions to the reverse-current dissipation dominating the energy losses as the flare magnitudes increase.

Recall however that, even in a situation with the dominated heating of a hot plasma by a reverse current, we cannot neglect the rare Coulomb collisions of fast electrons of a beam with thermal electrons of the plasma if we want to obtain the physically correct results (see Sect. 4.5.4.1). Once again,

both the collective kinetic effects and Coulomb collisions play the essential role in the physics of plasma heating by electron beams.

The *RHESSI* observations presented by Alexander and Daou (2007) shows that, for the most part of the beam fluxes, significant HXR fluxes are detected in the chromosphere, say in the thick target. The balance of thermal to non-thermal HXR emission in solar flares, its spatial distribution and its evolution in time may provide important diagnostic information on the development and evolution of beam systems in flares.

4.5.7 To the Future Models

The spectrum, polarization and directivity of hard X-ray radiation are the most delicate observational parameters that should be calculated very accurately to judge the validity of the theoretical models accounting for acceleration of electrons and their radiation. After all said above in this chapter, it is rather surprising to conclude that the most of the above mentioned 2D (in velocity space) models, which have been developed after the *classical* thick-target model (Sect. 4.3), are however not used to obtain a more realistic quantitative information on the fast electrons in solar and stellar flares. *The simplest classical thick-target model is still very popular* especially in astrophysical applications. Up to now we do not have a realistic time-dependent self-consistent thick-target model (which must be simple enough to be easily used) to interpret and analyze the hard X-ray emission so frequently detected in space.

Future models will incorporate such fine effects like a nonuniform initial ionization of chromospheric plasma in the thick-target (Brown et al. 1998a, 2003), the time-of-flight effect (Aschwanden et al. 1998; Brown et al. 1998b; Aschwanden 2002), with account taken of the effect of the *reverse-current electric field* as an effect of primary importance. Otherwise the accuracy of a model is lower than the accuracy of modern hard X-ray data obtained by *RHESSI* (Lin et al. 2002, 2003).

* * *

Now let us clarify our plans. Before transition to the hydrodynamic-type descriptions of astrophysical plasma, that are valid for systems containing a large number of colliding particles, we have to study two particular but interesting cases.

First, $N = 1$, a particle in a given force field. This simplest approximation gives us clear approach to several fundamental issues of collisionless plasma. In particular, it is necessary to outline the basis of kinetic theory for wave-particle interactions in astrophysical plasma (Chap. 7).

Second, $N = 2$, binary collisions of particles with the Coulomb potential of interaction. They are typical for collisional plasma. We have to know the Coulomb collisions well to justify the hydrodynamic description of astrophysical plasma (Chap. 9).

In the next chapter we start from the former.

4.6 Practice: Exercises and Answers

Exercise 4.1 (Sect. 4.3). How deep can the accelerated electrons with the initial energy $\mathcal{E}_0 \approx 10 \text{ keV}$ penetrate from the solar corona into the chromosphere?

Answer. From formula (4.40) we find the simplest estimation for the column depth

$$\xi = \frac{\mathcal{E}_0^2}{2a_0}, \text{ cm}^{-2}. \quad (4.84)$$

Substituting $\mathcal{E}_0 \approx 10 \text{ keV}$ and $n \approx 10^{12} \text{ cm}^{-3}$ in formula (4.41) gives $a_0 \approx 3 \times 10^{-18} \text{ keV}^2 \text{ cm}^2$. With this value a_0 we find $\xi \approx 10^{19} \text{ cm}^{-2}$. At such depth in the chromosphere, the density of the plasma $n \approx 10^{12} \text{ cm}^{-3}$ indeed.

Accelerated electrons with energies $\mathcal{E} > 10 \text{ keV}$ penetrate deeper and contribute significantly to impulsive heating of the optical part of a solar flare (see a temperature enhancement at $\xi \approx 10^{20} \text{ cm}^{-2}$ in Fig. 8.4).

Exercise 4.2 (Sect. 4.3). By using the thick-target model, find the energy spectrum of fast electrons, passing in high-temperature plasma the column depth $\xi = 1.5 \times 10^{19} \text{ cm}^{-2}$. The power-law injection spectrum ($\xi = 0$) has the boundaries $\mathcal{E}_1 = 10 \text{ keV}$ and $\mathcal{E}_2 = 100 \text{ keV}$, the spectral index $\gamma = 3$; the energy flux of fast electrons $F_0 = 10^{11} \text{ erg cm}^{-2} \text{ s}^{-1}$.

Answer. In order to solve the problem, we use formulae (4.39)–(4.42). At first, we have to find the coefficient K . With this aim, let us write expression for the energy flux

$$\begin{aligned} F_0 &= \int_{\mathcal{E}_1}^{\mathcal{E}_2} \mathcal{E} v N(\mathcal{E}, 0) d\mathcal{E} = \\ &= K \left(\frac{2}{m} \right)^{1/2} \times \begin{cases} (\mathcal{E}_2^{2-\gamma} - \mathcal{E}_1^{2-\gamma}) / (2 - \gamma), & \text{if } \gamma \neq 2, \\ \ln(\mathcal{E}_2 / \mathcal{E}_1), & \text{if } \gamma = 2. \end{cases} \end{aligned} \quad (4.85)$$

Thus

$$K = F_0 \left(\frac{m}{2} \right)^{1/2} \times \begin{cases} (2 - \gamma) / (\mathcal{E}_2^{2-\gamma} - \mathcal{E}_1^{2-\gamma}), & \text{if } \gamma \neq 2, \\ [\ln(\mathcal{E}_2 / \mathcal{E}_1)]^{-1}, & \text{if } \gamma = 2. \end{cases} \quad (4.86)$$

From here, with $\gamma = 3$, we calculate coefficient

$$K = 1.48 \times 10^7 \text{ cm}^{-3} \text{ keV}^{2.5}.$$

The energy spectrum is presented on Fig. 4.4.




Fig. 4.4 The energy spectrum of fast electrons in a high-temperature plasma on the column depth $\xi = 0$ (the *right straight line*) and on the column depth $\xi = 1.5 \times 10^{19} \text{ cm}^{-2}$ (*left curve*). Here $\xi'_1 = 3.41 \text{ keV}$ and $\xi'_2 = 99.56 \text{ keV}$ according to (4.42)

Exercise 4.3 (Sect. 4.5.3). How strong is the reverse-current electric field in the chromosphere during a solar flare?

Answer. According to (4.76), the electric field

$$E = \varepsilon \frac{\tau}{2} E_{D,1} \approx (10^{-4} - 10^{-3}) E_{D,1}. \quad (4.87)$$

In the chromosphere (Exercise 8.4), the Dreicer field $E_D > 0.1 \text{ V cm}^{-1}$. So, under injection of accelerated electrons into the chromosphere during the impulsive phase of a flare, the reverse-current field $E > 10^{-5} - 10^{-4} \text{ V cm}^{-1}$. With the length scale $l \sim 10^3 \text{ km}$, this electric field gives rise to a potential $\phi \approx E l \gtrsim 1 - 10 \text{ keV}$.

From this rough estimate, it is clear that

the physical effects related to a reverse-current electric field in the solar chromosphere should appear in the X-ray range

of flare observations by the *RHESSI* telescope. This seems to be well confirmed by Alexander and Daou (2007); see Sect. 4.5.6.

Exercise 4.4 (Sect. 4.5.4). Discuss expected properties of a solution of (4.79) without the collisional energy losses term $x (\partial f / \partial x)$.

Chapter 5

Motion of a Charged Particle in Given Fields

Abstract Astrophysical plasma is often an extremely tenuous hot gas of charged particles, without net charge on average. If there are very few encounters between particles, we need only to consider the responses of a particle to the force fields in which it moves. The simplest situation, a single particle in given fields, allows us to understand the drift motions of different origin and electric currents in such collisionless plasma.

5.1 A Particle in Constant Homogeneous Fields

5.1.1 Relativistic Equation of Motion

In order to study the motion of a charged particle, let us consider the following basic equation:

$$\frac{d\mathbf{p}}{dt} = e\mathbf{E} + \frac{e}{c}\mathbf{v} \times \mathbf{B} + m\mathbf{g}. \quad (5.1)$$

In relativistic mechanics (see [Landau and Lifshitz 1975](#), Chap. 2, Sect. 9) the particle momentum and energy are

$$\mathbf{p} = \frac{m\mathbf{v}}{\sqrt{1-v^2/c^2}} \quad \text{and} \quad \mathcal{E} = \frac{mc^2}{\sqrt{1-v^2/c^2}}, \quad (5.2)$$


respectively. By using the Lorentz factor

$$\gamma_L = \frac{1}{\sqrt{1-v^2/c^2}}, \quad (5.3)$$

we rewrite formulae (5.2) as

$$\mathbf{p} = \gamma_L m \mathbf{v} \quad \text{and} \quad \mathcal{E} = \gamma_L mc^2. \quad (5.4)$$

Fig. 5.1 The trajectory of particle motion under the action of a constant homogeneous non-magnetic force



Hence

$$\mathbf{p} = \frac{\mathcal{E}}{c^2} \mathbf{v}. \quad (5.5)$$

By taking the scalar product of (5.1) with the velocity vector $\mathbf{v}$ we obtain

$$\frac{d\mathcal{E}}{dt} = \mathbf{F} \cdot \mathbf{v}, \quad (5.6)$$

where

$$\mathbf{F} = e \mathbf{E} + m \mathbf{g}$$

is a *non-magnetic* force. The particle kinetic energy change during the time dt is $d\mathcal{E} = \mathbf{v} \cdot d\mathbf{p}$. Therefore, according to (5.6), *the work on a particle is done by the non-magnetic force only*. In what follows we shall remember that magnetic fields are “lazy” and do not work.

Let us consider the particle motion in *constant homogeneous* fields.

5.1.2 Constant Homogeneous Non-magnetic Forces

Now let a non-magnetic force be parallel to the y axis, $\mathbf{F} = F \mathbf{e}_y$, and let the initial momentum of the particle be parallel to the x axis, $\mathbf{p}_0 = p_0 \mathbf{e}_x$.

Then we integrate (5.1) to find that the particle moves along the *catenary* shown in Fig. 5.1:

$$y - y_0 = \frac{\mathcal{E}_0}{F} \left\{ \cosh \left[\frac{F}{p_0 c} (x - x_0) \right] - 1 \right\}. \quad (5.7)$$

Here $\mathcal{E}_0$ is an initial energy of the particle.

Formula (5.7) in the non-relativistic limit is that of a parabola:

$$y - y_0 = \frac{F}{2mv_0^2} (x - x_0)^2.$$

5.1.3 Constant Homogeneous Magnetic Fields

Let the non-magnetic force $\mathbf{F} = 0$. The magnetic force in a constant and homogeneous field results in particle motions. Let us show that. From (5.1) we have

$$\frac{d\mathbf{p}}{dt} = \frac{e}{c} \mathbf{v} \times \mathbf{B}. \quad (5.8)$$

We know by virtue of (5.6) that the particle kinetic energy $\mathcal{E} = \text{const}$. Therefore $|\mathbf{v}| = \text{const}$, and from (5.8)

$$\dot{\mathbf{v}} = \omega_B \mathbf{v} \times \mathbf{n}. \quad (5.9)$$

Here the overdot denotes the derivative with respect to time t , $\mathbf{n}$ is the unit vector along the field $\mathbf{B} = B \mathbf{n}$, and the constant

$$\omega_B = \frac{ecB}{\mathcal{E}} \quad (5.10)$$

is the *gyrofrequency* or *cyclotron frequency*. We use sometimes, in what follows, the name Larmor frequency. The last is a slightly confusing terminology in view of the fact that there is the frequency of the Larmor precession (see Sect. 45 in [Landau and Lifshitz 1975](#)), ω_L , which turns out to be half of the gyrofrequency ω_B .

In the non-relativistic limit, the gyrofrequency

$$\omega_B = \frac{eB}{mc}.$$

(5.11)

By integrating (5.9) we find the linear differential equation

$$\dot{\mathbf{r}} = \omega_B \mathbf{r} \times \mathbf{n} + \mathbf{C}, \quad (5.12)$$

where vector $\mathbf{C} = \text{const}$.

By taking the scalar product of (5.12) with the unit vector $\mathbf{n}$ we have

$$\mathbf{n} \cdot \dot{\mathbf{r}} = C_{\parallel} \equiv v_{\parallel} (t = 0).$$

The constant $\mathbf{C}_\perp$ can be removed from consideration by an appropriate choice of the moving reference system. $\mathbf{C}_\perp = 0$ in the reference system where $\mathbf{F} = 0$ (Sect. 5.1.4), and this choice is consistent with the initial equation (5.8). Therefore

$$\dot{\mathbf{r}}_\perp = \omega_B \mathbf{r}_\perp \times \mathbf{n}. \quad (5.13)$$

The vector $\mathbf{r}_\perp$ is changing with the velocity $\mathbf{v}_\perp$ which is perpendicular to $\mathbf{r}_\perp$ itself. Hence the change of vector $\mathbf{r}_\perp$ is a *rotation with the constant frequency* $\omega = \omega_B \mathbf{n}$. Thus we have

$$v_\perp = \omega_B r_\perp = \text{const} = v_\perp(0),$$

and

$$r_\perp = \frac{v_\perp(0)}{\omega_B} = \frac{\mathcal{E} v_\perp(0)}{ecB} = \frac{c p_\perp}{eB},$$

since it follows from formula (5.5) that

$$\mathcal{E} v_\perp = c^2 p_\perp.$$

We have obtained the expression for the *gyroradius* or the Larmor radius

$$\boxed{r_L = \frac{c p_\perp}{eB}}. \quad (5.14)$$

The term “*rigidity*” is introduced in cosmic physics:

$$\mathcal{R} = \frac{c \mathbf{p}}{e}. \quad (5.15)$$

The rigidity of a particle is measured in Volts:

$$[\mathcal{R}] = \frac{[cp]}{[e]} = \frac{eV}{e} = V.$$

Rigidity is usually used together with the term “*pitch-angle*”

$$\theta = \left(\widehat{\mathbf{v}_0, \mathbf{B}} \right). \quad (5.16)$$

From (5.14) and (5.15) it follows that the particle’s gyroradius or Larmor radius is

$$r_L = \frac{\mathcal{R}_\perp}{B}. \quad (5.17)$$

That is why

the particles with the same rigidity and pitch-angle move along the same trajectories in a magnetic field.

This fact is used in the physics of the magnetospheres of the Earth and other planets, as well as in general physics of “*cosmic rays*” (Ginzburg and Syrovatskii 1964; Schlickeiser 2002; Dorman 2009).

The cosmic rays, high-energy (from 10^9 eV to somewhat above 10^{20} eV) particles of cosmological origin, were discovered almost a century ago. However they are one of the very few means available to an Earth-based observer to study astrophysical or cosmological phenomena. The knowledge of their incoming direction and their energy spectrum are the bits and pieces of a complex puzzle which can give us information on the mechanism that produced them at the origin, unfortunately distorted by many effects they undergo during their journey over huge distances.

5.1.4 Non-magnetic Force in a Magnetic Field

Let us consider the case when a non-magnetic force $\mathbf{F}$ is perpendicular to the homogeneous magnetic field $\mathbf{B}$ as shown in Fig. 5.2. For the sake of simplicity, we shall consider the non-relativistic equation of motion:

$$m \dot{\mathbf{v}} = \mathbf{F} + \frac{e}{c} \mathbf{v} \times \mathbf{B}. \quad (5.18)$$

Let us try to find the solution of this equation in the form

$$\mathbf{v} = \mathbf{v}_d + \mathbf{u}. \quad (5.19)$$

Here $\mathbf{v}_d$ is some constant velocity, so that substituting (5.19) in (5.18) gives

$$m \dot{\mathbf{u}} + \mathbf{0} = \frac{e}{c} \mathbf{u} \times \mathbf{B} + \mathbf{F} + \frac{e}{c} \mathbf{v}_d \times \mathbf{B}.$$

We choose $\mathbf{v}_d$ in such a way that the two last terms vanish:

$$\mathbf{F} + \frac{e}{c} \mathbf{v}_d \times \mathbf{B} = \mathbf{0}.$$

This is the case if the following expression is chosen:

$$\mathbf{v}_d = \frac{c}{e} \frac{\mathbf{F} \times \mathbf{B}}{B^2}.$$

(5.20)




Fig. 5.2 The trajectory of motion a positively charged particle in a homogeneous magnetic field under action of a non-magnetic force. Slow (a) and fast (b) drifts

Actually, by using the known vector identity

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = \mathbf{b}(\mathbf{a} \cdot \mathbf{c}) - \mathbf{c}(\mathbf{a} \cdot \mathbf{b}),$$

we infer

$$\frac{e}{c} \mathbf{v}_d \times \mathbf{B} = \mathbf{n}(\mathbf{n} \cdot \mathbf{F}) - \mathbf{F} = -\mathbf{F},$$

since $\mathbf{F} \perp \mathbf{n} = \mathbf{B}/B$. So formula (5.20) is correct.

Thus if a non-magnetic force $\mathbf{F}$ is perpendicular to the field $\mathbf{B}$, the particle motion is a sum of the *drift* with the velocity (5.20) called *drift velocity*, which is perpendicular to both $\mathbf{F}$ and $\mathbf{B}$, and the *spiral* motion round the magnetic field lines—the *gyromotion*:

$$m \dot{\mathbf{u}} = \frac{e}{c} \mathbf{u} \times \mathbf{B}. \quad (5.21)$$

Depending on a relative speed of these two motions, we distinguish *slow* ($v_d < u$) and *fast* ($v_d > u$) drifts, see (a) and (b) in Fig. 5.2.




Fig. 5.3 Electric drift. The kinetic energy $\mathcal{E}$ of a positively charged particle p^+ is a maximum at the upper point 1, hence the curvature radius r_L of the trajectory is a maximum at this point

To understand the motion, let us think first about how the particle would move if only the magnetic field were present. It would gyrate in a circle, and the direction of motion around the circle would depend on the sign of the particle's charge. The radius of the circle, r_L , would vary with the particle's mass and would therefore much larger for an ion than for an electron if their velocities were the same (see formula (5.14)).

The non-magnetic force $\mathbf{F}$ accelerates the particle during part of each orbit (see $1 \rightarrow 2$ in Fig. 5.2a) and decelerates it during the remaining part of the orbit (see $2 \rightarrow 3$ in Fig. 5.2a). The result is that the orbit is a distorted circle with a larger-than-average radius of curvature during half of the orbit and a smaller-than-average radius of curvature during the remaining half of the orbit. A net displacement is perpendicular to the force $\mathbf{F}$ and the magnetic field $\mathbf{B}$.

5.1.5 Electric and Gravitational Drifts

As we saw above, any force $\mathbf{F}$, that is capable of accelerating or decelerating charged particles as they gyrate about the magnetic field $\mathbf{B}$ in *collisionless plasma*, will result in a drift perpendicular to both the field and the force.

(a) If $\mathbf{F} = e \mathbf{E}$, then the drift is called *electric drift*, its velocity

$$\mathbf{v}_d = c \frac{\mathbf{E} \times \mathbf{B}}{B^2} \quad (5.22)$$

being independent of the particle charge and mass (Fig. 5.3).

Since the drift velocity depends upon neither the charge nor the mass of the particle,

the electric drift generates the motion of collisionless plasma as a whole with the velocity $\mathbf{v} = \mathbf{v}_d$ relative to a magnetic field.




Fig. 5.4 Gravitational drift. Initiation of an electric current by the action of the gravity force in a collisionless plasma with magnetic field

Being involved in the electric drift, the collisionless plasma tends: (a) to flow similar to a fluid, and (b) to be “squeezed out” from direct action of the electric field $\mathbf{E}$ applied in a direction which is perpendicular to the magnetic field $\mathbf{B}$.

Formula (5.22) says that the drift velocity is perpendicular to both the electric and magnetic fields. This is sometimes referred to as an “ E -cross- B drift”, but its magnitude is inversely proportional to the magnitude of $\mathbf{B}$. In fact, formula (5.22) would formally result in $v_d \geq c$ for $E \geq B$. However we should not forget that formula (5.22) was obtained in the non-relativistic limit.

- (b) For the gravitational force $\mathbf{F} = m \mathbf{g}$ formula (5.20) gives the drift velocity

$$\mathbf{v}_d = \frac{mc}{e} \frac{\mathbf{g} \times \mathbf{B}}{B^2}. \quad (5.23)$$

The *gravitational* drift velocity is seen to depend upon the particle mass and charge. Positively charged particles drift in the direction coinciding with that of the product $\mathbf{g} \times \mathbf{B}$, while negatively charged particles drift in the opposite direction as shown in Fig. 5.4. Therefore

■ a gravitational field is capable of generating an electric current in a magnetized collisionless plasma.

This may often happen in astrophysical conditions.

5.2 Weakly Inhomogeneous Slowly Changing Fields

5.2.1 Small Parameters in the Motion Equation

Let us take the non-relativistic equation (5.18) for the motion of a charged particle and rewrite it as follows:

$$\frac{m}{e} (\ddot{\mathbf{r}} - \mathbf{g}) = \mathbf{E} + \frac{1}{c} \dot{\mathbf{r}} \times \mathbf{B}. \quad (5.24)$$

On making this expression non-dimensional

$$\mathbf{r}^* = \frac{\mathbf{r}}{L}, \quad t^* = \frac{t}{\tau}, \quad \mathbf{v}^* = \frac{\mathbf{v}}{v_0}, \quad \mathbf{g}^* = \frac{\mathbf{g}}{g}, \quad \mathbf{B}^* = \frac{\mathbf{B}}{B_0}, \quad \mathbf{E}^* = \frac{\mathbf{E}}{E_0},$$

we have the following equation

$$\frac{m}{e} \frac{L}{\tau^2} \left(\ddot{\mathbf{r}}^* - \frac{g\tau^2}{L} \mathbf{g}^* \right) = E_0 \mathbf{E}^* + \frac{L}{c\tau} B_0 \dot{\mathbf{r}}^* \times \mathbf{B}^*.$$

Let us normalize this equation with respect to the last term (the Lorentz force) by dividing the equation by $LB_0/c\tau$:

$$\frac{m}{e} \frac{c}{B_0} \frac{1}{\tau} (\ddot{\mathbf{r}}^* - \alpha_g \mathbf{g}^*) = \frac{E_0}{B_0} \frac{c\tau}{L} \mathbf{E}^* + \dot{\mathbf{r}}^* \times \mathbf{B}^*.$$

Introduce the dimensionless parameter

$$\alpha_B = \frac{m}{e} \frac{c}{B_0} \frac{1}{\tau}.$$

Two situations are conceivable.

- (a) Spatially homogeneous magnetic and electric fields are slowly changing in time. The characteristic time $\tau = 1/\omega$, where ω is a characteristic field change frequency. Therefore the dimensionless parameter α_B is equal to

$$\alpha_B = \frac{\omega}{\omega_B}, \quad (a)$$

where

$$\omega_B = \frac{eB}{mc}$$

is the Larmor frequency.

- (b) For the fields that are constant in time but weakly inhomogeneous, the characteristic time is to be defined as $\tau = L/v_0$, L and v_0 being the characteristic values of the field dimensions and the particle velocity (more exactly we assume $v_\perp \sim v_0$), respectively. In this case

$$\alpha_B = \frac{r_L}{L}, \quad (b)$$

where

$$r_L = \frac{c p_\perp}{eB} = \frac{cm v_\perp}{eB}$$

is the Larmor radius.

Generally, a superposition of these two cases takes place. The field is called *weakly inhomogeneous slowly changing* field, if

$$\alpha_B \approx \frac{\omega}{\omega_B} \approx \frac{r_L}{L} \ll 1. \quad (5.25)$$

The second parameter of the problem,

$$\alpha_E = \frac{E_0}{B_0} \frac{c\tau}{L},$$

characterizes the relative role of the electric field. We assume $\alpha_E = 1$, because, if this parameter is small or large, this can be easily taken into account in the final result.

The third dimensionless parameter $\alpha_g = g\tau^2/L$ is not important for our consideration in this section; so we put $\alpha_g = 1$ too.

Thus we have

$$\alpha_B (\ddot{\mathbf{r}}^* - \mathbf{g}^*) = \mathbf{E}^* + \dot{\mathbf{r}}^* \times \mathbf{B}^*, \quad (5.26)$$

the equation formally coinciding with the initial dimensional one. That is why it is possible to work with (5.24), using as a *small parameter* the dimensional quantity m/e . The corresponding expansion in the Taylor series is termed the expansion in powers of m/e . This method is widely used in plasma physics, starting presumably from [Chew et al. \(1956\)](#). We shall find such a solution of (5.24).


5.2.2 Expansion in Powers of m/e

Now let us represent the solution of (5.24) as a sum of two terms,

$$\mathbf{r}(t) = \mathbf{R}(t) + \mathbf{r}_L(t). \quad (5.27)$$

The first term $\mathbf{R}(t)$ describes the motion of the *guiding center* of the Larmor circle, the second term $\mathbf{r}_L(t)$ corresponds to the rotational motion or gyromotion of the particle. The case of an electron e^- is shown in Fig. 5.5.

Fig. 5.5 The Larmor motion of a negatively charged particle (an electron) in a weakly inhomogeneous slowly changing field



Recall that for the constant homogeneous magnetic field (see (5.14))

$$r_L = \frac{c p_{\perp}}{eB} = \frac{m}{e} \frac{c v_{\perp}}{B},$$

i.e., the Larmor radius is proportional to the parameter m/e . It is natural to suppose that the dependence is the same for the weakly inhomogeneous slowly changing field, i.e.

$$|r_L| \sim \frac{m}{e}.$$

For example, if the magnetic field does not change in time and does not change much within the gyroradius, then the particle moves through a nearly uniform magnetic field while making a circular round. However the non-uniformities make the guiding center move in a way different from a simple translatory motion. We are going to find the equation describing the guiding center motion.

Let us substitute (5.27) in (5.24) and expand the fields $\mathbf{g}$, $\mathbf{E}$, and $\mathbf{B}$ in the Taylor series about the point $\mathbf{r} = \mathbf{R}$:

$$\begin{aligned} \mathbf{g}(\mathbf{r}) &= \mathbf{g}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{g}(\mathbf{R}) + \dots, \\ \mathbf{E}(\mathbf{r}) &= \mathbf{E}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{E}(\mathbf{R}) + \dots, \\ \mathbf{B}(\mathbf{r}) &= \mathbf{B}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R}) + \dots. \end{aligned} \quad (5.28)$$

From (5.24) we have

$$\ddot{\mathbf{r}} = \mathbf{g} + \left(\frac{m}{e}\right)^{-1} \left[\mathbf{E}(\mathbf{r}) + \frac{1}{c} \dot{\mathbf{r}} \times \mathbf{B}(\mathbf{r}) \right].$$

Hence the basic equation contains the small parameter m/e to the power (-1) . By substituting (5.27) and (5.28) in this equation we obtain

$$\begin{aligned}
\ddot{\mathbf{R}} + \ddot{\mathbf{r}}_L &= \mathbf{g}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{g}(\mathbf{R}) + \\
&+ \left(\frac{m}{e}\right)^{-1} \{ \mathbf{E}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{E}(\mathbf{R}) \} + \\
&+ \left(\frac{m}{e}\right)^{-1} \left\{ \frac{1}{c} (\dot{\mathbf{R}} + \dot{\mathbf{r}}_L) \times [\mathbf{B}(\mathbf{R}) + (\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})] \right\} + \dots \quad (5.29)
\end{aligned}$$

Note that we have to think carefully about smallness of different terms in (5.29). For example, the magnitude of $\dot{\mathbf{r}}_L$ is not small:

$$|\dot{\mathbf{r}}_L| \sim \frac{|\mathbf{r}_L|}{\tau} \sim r_L \omega_B \sim \frac{m}{e} \times \left(\frac{m}{e}\right)^{-1} \sim 1.$$

since the Larmor frequency

$$\omega_B = \frac{eB}{mc} \sim \left(\frac{m}{e}\right)^{-1}.$$

Therefore, *the particle velocity is not small, although the Larmor radius is small.* That is the physical reason for the term

$$\left(\frac{m}{e}\right)^{-1} \frac{1}{c} [\dot{\mathbf{r}}_L \times (\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})]$$

having zero order with respect to the small parameter m/e .

The acceleration term $\ddot{\mathbf{r}}_L$ is not small either:


$$|\ddot{\mathbf{r}}_L| \sim \frac{|\mathbf{r}_L|}{\tau^2} \sim r_L \omega_B^2 \sim \left(\frac{m}{e}\right)^{-1}.$$

In the expansion (5.29) let us retain only the terms with the order of smallness less than one, that is

$$\begin{aligned}
\underbrace{\ddot{\mathbf{R}}}_{(0)} &= \underbrace{-\ddot{\mathbf{r}}_L}_{(-1)} + \underbrace{\mathbf{g}(\mathbf{R})}_{(0)} + \underbrace{\left(\frac{m}{e}\right)^{-1} \left[\mathbf{E}(\mathbf{R}) + \frac{1}{c} \dot{\mathbf{R}} \times \mathbf{B}(\mathbf{R}) \right]}_{(-1)} + \\
&+ \underbrace{\left(\frac{m}{e}\right)^{-1} (\mathbf{r}_L \cdot \nabla) \mathbf{E}(\mathbf{R})}_{(0)} + \underbrace{\left(\frac{m}{e}\right)^{-1} \frac{1}{c} \dot{\mathbf{R}} \times [(\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})]}_{(0)} + \\
&+ \underbrace{\left(\frac{m}{e}\right)^{-1} \frac{1}{c} \dot{\mathbf{r}}_L \times [(\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})]}_{(0)} + O\left(\frac{m}{e}\right). \quad (5.30)
\end{aligned}$$

Here the orders of smallness of the corresponding terms are given in brackets under the braces.

Fig. 5.6 The motion of a negatively charged particle on the Larmor orbit and its magnetic moment. The moment is antiparallel to the magnetic field



5.2.3 The Averaging Over Gyromotion

In order to obtain the equation for *guiding center* motion let us average (5.30) over a small period of the Larmor rotation,

$$T_B = \frac{2\pi}{\omega_B} = 2\pi \frac{mc}{eB} \sim \frac{m}{e}.$$

Since $\langle \mathbf{r}_L \rangle = \langle \dot{\mathbf{r}}_L \rangle = \langle \ddot{\mathbf{r}}_L \rangle = 0$, we infer the following equation

$$\begin{aligned} \ddot{\mathbf{R}} = & \mathbf{g}(\mathbf{R}) + \frac{e}{m} \left[\mathbf{E}(\mathbf{R}) + \frac{1}{c} \dot{\mathbf{R}} \times \mathbf{B}(\mathbf{R}) \right] + O\left(\frac{m}{e}\right) + \\ & + \frac{e}{mc} \langle \dot{\mathbf{r}}_L \times [(\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})] \rangle. \end{aligned} \quad (5.31)$$

Let us consider the last term which also has to be averaged. Here we may put

$$\dot{\mathbf{r}}_L = \omega_B \mathbf{r}_L \times \mathbf{n}.$$

On rearrangement (see Exercise 5.9), we obtain

$$\frac{e}{mc} \langle \dot{\mathbf{r}}_L \times [(\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})] \rangle = -\frac{\mathcal{M}}{m} \nabla B. \quad (5.32)$$

Here


$$\mathcal{M} = \frac{1}{c} \frac{e \omega_B}{2\pi} (\pi r_L^2) = \frac{1}{c} JS \quad (5.33)$$

is the *magnetic moment* of a particle on the Larmor orbit (Fig. 5.6). The case of electron e^- is shown here.

We interpret $-e(\omega_B/2\pi)$ as the current $+J$ associated with the gyrating electron. That is why we call $\mathcal{M}$ a *dipole* magnetic moment as the name usually refers to a property of a *current loop* defined as the current J flowing through the loop times the area S of the loop (see [Sivukhin 1965](#)). Hence it is clear from (5.33) that $\mathcal{M}$ is the magnetic moment of the gyrating particle.

So a single gyrating charge generates a magnetic dipole. Note that, for any charge of a particle, positive or negative,

Fig. 5.7 The *diamagnetic force* acts on the guiding center moving along the symmetry axis of a magnetic mirror configuration



the direction of the dipole magnetic moment is opposite to the direction of the magnetic field.

Therefore the *diamagnetic effect* has to occur.

Substituting the non-relativistic formula $\omega_B = eB/mc$ in (5.33) gives

$$\mathcal{M} = \frac{1}{2\pi} \frac{e^2}{mc^2} B \pi r_L^2. \quad (5.34)$$

Therefore

the magnetic moment is proportional to the magnetic field flux through the surface covering the particle's Larmor orbit.

It is also obvious from (5.32) that we can use the following formula for the force acting on the magnetic moment:

$$\mathbf{F} = -\mathcal{M} \nabla B.$$

(5.35)

Let the field strength increase along the field direction. For the sake of simplicity, we consider a magnetic configuration symmetric around the central field line as shown in Fig. 5.7. The strength of the magnetic field increases when the guiding center (not a particle) of a particle moves along the central line in the direction of the axis z .

The diamagnetic force (5.35) is exerted along the magnetic field and *away from the direction of increase of the field*.

As a consequence, the parallel component of the guiding center velocity $\dot{\mathbf{R}}$ decreases to zero at some maximum strength of the magnetic field and then changes sign. We say that the particle experiences a *mirror force*, and we shall call the place where it turns around a *magnetic mirror*. Note that a charged particle moving along the symmetry axis z (without gyration) is unaffected by magnetic force of course.

Finally, from (5.31), we obtain the equation of the guiding center motion:

$$\ddot{\mathbf{R}} = \mathbf{g}(\mathbf{R}) + \frac{e}{m} \left[\mathbf{E}(\mathbf{R}) + \frac{1}{c} \dot{\mathbf{R}} \times \mathbf{B}(\mathbf{R}) \right] - \frac{\mathcal{M}}{m} \nabla B(\mathbf{R}) + O\left(\frac{m}{e}\right). \quad (5.36)$$

The guiding center calculations involve considerably less amount of numerical work and produce trajectories in good agreement with detailed calculations if the non-uniformities of the magnetic and other fields are really small over the region through which the particle is making the circular motion. Moreover

the guiding center theory helps us to develop an intuition about the motions of charged particles in magnetic field.

And this intuition turns out to be useful in solving many practical problems of plasma astrophysics, for example, in physics of the Earth magnetosphere.

In the problem of magnetic reconnection, the guiding center approximation allows us to investigate particle motion in a *reconnecting current layer* with a three-component magnetic field (Oreshina and Somov 2009). Given the electric field related to reconnection, the analytical solution describes the stable and unstable three-dimensional orbits of particles. Charge separation is shown to take place in the current layer: oppositely charged particles are localized mostly in different regions of the layer.

5.2.4 Spiral Motion of the Guiding Center

Even without regarding the terms $O(m/e)$, (5.36) is more difficult in comparison with the motion (5.24). The term $\mathbf{g}(\mathbf{R})$, the term with electric field $\mathbf{E}(\mathbf{R})$, and the two last terms in (5.36) apart, it is seen that

$$\ddot{\mathbf{R}} = \frac{e}{mc} \dot{\mathbf{R}} \times \mathbf{B}. \quad (5.37)$$

Therefore the guiding center *spirals*, as does the particle (cf. (5.8)).

By analogy with formula (5.14), the guiding center spiral radius can be found

$$R_{\perp} = \frac{mc \dot{R}_{\perp}}{eB}. \quad (5.38)$$

So it is a small quantity of order

$$\frac{R_{\perp}}{r_L} = \frac{\dot{R}_{\perp}}{v_{\perp}} \sim \frac{r_L}{L}$$

as compared with the particle Larmor radius (5.14).

The radius of the guiding center spiral is of the order of m/e as compared with the particle Larmor radius. Consequently, this spiral has a higher order with respect to the small parameter m/e and can be neglected in the approximation under study.

5.3 Gradient and Inertial Drifts

5.3.1 Basic Formulae

Let us neglect the term $O(m/e)$ in (5.36) and take the vector product of (5.36) with the unit vector $\mathbf{n} = \mathbf{B}/B$:

$$\ddot{\mathbf{R}} \times \mathbf{n} = \mathbf{g} \times \mathbf{n} + \frac{e}{m} \mathbf{E} \times \mathbf{n} + \frac{eB}{mc} (\dot{\mathbf{R}} \times \mathbf{n}) \times \mathbf{n} + \frac{\mathcal{M}}{m} \mathbf{n} \times \nabla B.$$

From the third term on the right-hand side of this equation, we find the *drift velocity* across the magnetic field

$$\begin{aligned} \dot{\mathbf{R}}_{\perp} \equiv \mathbf{n} \times (\dot{\mathbf{R}} \times \mathbf{n}) &= c \frac{\mathbf{E} \times \mathbf{n}}{B} + \frac{mc}{eB} \mathbf{g} \times \mathbf{n} + \\ &+ \frac{\mathcal{M}c}{eB} \mathbf{n} \times \nabla B - \frac{mc}{eB} \ddot{\mathbf{R}} \times \mathbf{n}. \end{aligned} \quad (5.39)$$

The first term on the right-hand side of (5.39) corresponds to the *electric drift* (5.22), the second one presents the *gravitational drift* (5.23). The third term is new for us in this chapter; it describes the *gradient drift* arising due to the magnetic field inhomogeneity. The gradient drift velocity

$$\mathbf{v}_d = \frac{\mathcal{M}c}{eB} \mathbf{n} \times \nabla B.$$

(5.40)

The same formula follows of course from (5.20) after substituting in it the formula (5.35) for the force acting on the magnetic moment $\mathcal{M}$ in the weakly inhomogeneous field.

So, if a particle gyrates in a magnetic field whose strength changes from one side of its gyration orbit to the other, the instantaneous radius of the curvature of the orbit will become alternately smaller and larger. Averaged over several gyrations,

the particle drifts in a direction perpendicular to both the magnetic field and the direction in which the strength of the field changes.

The fourth term on the right-hand side of (5.39) corresponds to the *inertial* drift:

$$\mathbf{v}_d = -\frac{mc}{eB} \ddot{\mathbf{R}} \times \mathbf{n}. \quad (5.41)$$

Let us consider it in some detail. For calculating the inertial drift velocity (5.41), we have to know the guiding center *acceleration* $\ddot{\mathbf{R}}$. It will suffice for the calculation of $\ddot{\mathbf{R}}$ to consider (5.39) in the zeroth order, since the last term of (5.39) contains the small parameter m/e . In this order with respect to m/e , we have

$$\dot{\mathbf{R}}_{\perp} = c \frac{\mathbf{E} \times \mathbf{n}}{B}.$$

Hence the guiding center acceleration

$$\ddot{\mathbf{R}} = \frac{d}{dt} \dot{\mathbf{R}} = \frac{d}{dt} (\dot{\mathbf{R}}_{\parallel} + \dot{\mathbf{R}}_{\perp}) = \frac{d}{dt} \left(v_{\parallel} \mathbf{n} + c \frac{\mathbf{E} \times \mathbf{n}}{B} \right). \quad (5.42)$$

Because this aspect of particle motion is important in accounting for the special properties of a collisionless astrophysical plasma, it is good to understand it not only mathematically but also in an intuitive manner.

5.3.2 The Centrifugal Drift

At first, we consider the particular case assuming the electric field $\mathbf{E} = 0$ in formula (5.42), the magnetic field $\mathbf{B}$ being *time-independent* but *weakly inhomogeneous*. Under these conditions

$$\ddot{\mathbf{R}} = \frac{d}{dt} (v_{\parallel} \mathbf{n}) = \mathbf{n} \frac{dv_{\parallel}}{dt} + v_{\parallel} \frac{d\mathbf{n}}{dt}.$$


The first term on the right-hand side does not contribute to the drift velocity since $\mathbf{n} \times \mathbf{n} = \mathbf{0}$. Rewrite the second term as follows:

$$v_{\parallel} \frac{d\mathbf{n}}{dt} = v_{\parallel} \left(\frac{\partial \mathbf{n}}{\partial t} + v_{\parallel} (\mathbf{n} \cdot \nabla) \mathbf{n} \right). \quad (5.43)$$

In this formula, the first term on the right equals zero for the time-independent field. The second one is equal to

$$v_{\parallel}^2 (\mathbf{n} \cdot \nabla) \mathbf{n} = -v_{\parallel}^2 \left(\frac{\mathbf{e}_c}{\mathcal{R}_c} \right). \quad (5.44)$$

Fig. 5.8 The frame of reference for derivation of the formula for the inertial drift in weakly inhomogeneous magnetic field



Here $\mathcal{R}_c$ is a radius of *curvature* for the field line at a given point $\mathbf{R}$. At this point the unit vector $\mathbf{e}_c$ is directed from the curvature center O_c as shown in Fig. 5.8.

Thus the dependence of the inertial drift velocity on the curvature of the weakly inhomogeneous magnetic field is found

$$\dot{\mathbf{R}}_{\perp}|_c = \frac{1}{\mathcal{R}_c \omega_B} v_{\parallel}^2 \mathbf{e}_c \times \mathbf{n}. \quad (5.45)$$

This is the drift of a particle under action of the *centrifugal* force

$$\mathbf{F}_c = \frac{mv_{\parallel}^2}{\mathcal{R}_c} \mathbf{e}_c. \quad (5.46)$$

In formula (5.45), the centrifugal force produced by motion of a particle along the magnetic field appears explicitly. Therefore the *centrifugal* drift velocity can be seen to be a special case of the expression (5.20) obtained for drift produced by an arbitrary non-magnetic force $\mathbf{F}$.

5.3.3 The Curvature-Dependent Drift

Let us come back to the gradient drift and consider a time-independent weakly-inhomogeneous magnetic field. Its gradient

$$\nabla B = \frac{1}{2B} \nabla (\mathbf{B} \cdot \mathbf{B}) = \frac{1}{B} [(\mathbf{B} \cdot \nabla) \mathbf{B} + \mathbf{B} \times \text{rot } \mathbf{B}].$$

In a current-free region $\text{rot } \mathbf{B} = 0$, and hence

$$\begin{aligned} \nabla B &= \frac{1}{B} (\mathbf{B} \cdot \nabla) \mathbf{B} = (\mathbf{n} \cdot \nabla) \mathbf{B} = (\mathbf{n} \cdot \nabla) B \mathbf{n} = B (\mathbf{n} \cdot \nabla) \mathbf{n} + \\ &+ \mathbf{n} (\mathbf{n} \cdot \nabla B) = -B \left(\frac{\mathbf{e}_c}{\mathcal{R}_c} \right) + \mathbf{n} (\mathbf{n} \cdot \nabla B). \end{aligned}$$

The last term does not contribute to the gradient drift velocity (5.40). The contribution of the first term to the drift velocity is

$$\dot{\mathbf{R}}_{\perp} = \frac{\mathcal{M}c}{eB} \mathbf{n} \times \left((-B) \frac{\mathbf{e}_c}{\mathcal{R}_c} \right) = -\frac{\mathcal{M}}{e \mathcal{R}_c} \mathbf{n} \times \mathbf{e}_c = \frac{\mathcal{M}}{e \mathcal{R}_c} \mathbf{e}_c \times \mathbf{n}. \quad (5.47)$$

Here, according to definition (5.33) and formula (5.14), the *magnetic moment*

$$\mathcal{M} = \frac{1}{c} JS = \frac{e \omega_B r_L^2}{2c} = \frac{e v_{\perp}^2}{2c \omega_B}. \quad (5.48)$$

On substituting formula (5.48) into (5.47) we see that the gradient drift in a time-independent weakly-inhomogeneous magnetic field has a structure analogous to the centrifugal drift (5.45):

$$\dot{\mathbf{R}}_{\perp}|_{\text{gr}} = \frac{1}{\mathcal{R}_c \omega_B} \frac{1}{2} v_{\perp}^2 \mathbf{e}_c \times \mathbf{n}. \quad (5.49)$$

Therefore we can add the curvature-dependent part of the gradient drift (5.49) to the centrifugal drift (5.45):

$$\dot{\mathbf{R}}_{\perp} = \frac{1}{\mathcal{R}_c \omega_B} \left(v_{\parallel}^2 + \frac{1}{2} v_{\perp}^2 \right) \mathbf{e}_c \times \mathbf{n}. \quad (5.50)$$

■ This formula unites the two drifts that depend on the field line curvature of a weakly inhomogeneous magnetic field.

In a curved magnetic field, the gradient drift is present in combination with the centrifugal drift.

5.3.4 The Curvature-Independent Gradient Drift

It is worth considering the part of the gradient drift, that is independent of the field line curvature. Let the field lines be straight ($\mathcal{R}_c \rightarrow \infty$), their density increasing unidirectionally as shown in Fig. 5.9. The field strength B_2 at a point 2 is greater than that one at a point 1. So, according to (5.17), the Larmor radius

$$r_L|_2 < r_L|_1.$$

The particle moves in the manner indicated in Fig. 5.9.

For comparison purposes, it is worth remembering another illustration. This is related to, on the contrary, the non-magnetic force $\mathbf{F}$ (Sect. 5.1.4). Under action of the force, the particle velocity at a point 1 in Fig. 5.10, v_1 , is greater than at a point 2. Hence the Larmor radius $r_L = cp_{\perp}/eB$ is greater at a point 1 than at a point 2 as well.




Fig. 5.9 The simplest interpretation of the gradient drift. A gradient in the field strength, ∇B , in the direction perpendicular to $\mathbf{B}$ will produce a drift motion of ions and electrons




Fig. 5.10 The physical nature of the drift under the action of a non-magnetic force $\mathbf{F}$ which is perpendicular to the uniform magnetic field $\mathbf{B}$

In other words, when the particle is at the point 2 at the top of its trajectory, the force $\mathbf{F}$ and the Lorentz force $(e/c) \mathbf{v} \times \mathbf{B}$ both act in the downward direction in Fig. 5.10. This enhanced normal acceleration makes the trajectory more sharply bent than it would have been in the absence of the force $\mathbf{F}$. On the other hand, when the particle is at the bottom point 1, the Lorentz force is diluted by $\mathbf{F}$, thereby causing the trajectory to be less sharply bent. As a result, there is a drift of the guiding center in a direction perpendicular to both $\mathbf{B}$ and $\mathbf{F}$.

Figures 5.9 and 5.10 also demonstrate the validity of formula (5.35).

The drifts with velocity which depends on the particle charge and mass, like the gradient drift, can give rise to a current by making the electrons and ions drift in opposite directions. Such drifts can also be important for the problem of element abundances or element fractionation (see Part II, Sects. 13.4 and 13.5).

Recommended Reading: Sivukhin (1965), Morozov and Solov'ev (1966b).

5.4 Practice: Exercises and Answers

Exercise 5.1 (Sect. 5.1). Evaluate the gyrofrequency for thermal electrons and protons in the *solar corona* above a sunspot.

Answer. At typical temperature in the corona, $T \approx 2 \times 10^6$ K, from the non-relativistic formula (5.11), it follows that: the electron gyrofrequency

$$\omega_B^{(e)} = 1.76 \times 10^7 B \text{ (G)}, \text{ rad s}^{-1}; \quad (5.51)$$

the proton gyrofrequency

$$\omega_B^{(p)} = 9.58 \times 10^3 B \text{ (G)}, \text{ rad s}^{-1}. \quad (5.52)$$

The gyrofrequency of electrons is $m_p / m_e \approx 1.84 \times 10^3$ times larger than that one of protons. Just above a sunspot the field strength can be as high as $B \approx 3,000$ G. Here $\omega_B^{(e)} \approx 5 \times 10^{10} \text{ rad s}^{-1}$. The emission of thermal electrons at this height in the corona can be observed at wavelength $\lambda \approx 4$ cm.

Exercise 5.2 (Sect. 5.1). Under conditions of the solar corona (see Exercise 5.1), evaluate the *mean thermal velocity* and the Larmor radius of thermal electrons and protons.

Answer. The thermal velocity of particles with mass m_i and temperature T_i is

$$V_{Ti} = \left(\frac{3k_B T_i}{m_i} \right)^{1/2}. \quad (5.53)$$

Respectively, for electrons and protons:

$$V_{Te} = 6.74 \times 10^5 \sqrt{T_e \text{ (K)}}, \text{ cm s}^{-1}, \quad (5.54)$$

and

$$V_{Tp} = 1.57 \times 10^4 \sqrt{T_p \text{ (K)}}, \text{ cm s}^{-1}. \quad (5.55)$$

At the coronal temperature $V_{Te} \approx 9.5 \times 10^3 \text{ km s}^{-1} \sim 10^9 \text{ cm s}^{-1}$ and $V_{Tp} \approx 220 \text{ km s}^{-1}$.

From (5.14) we find the following formulae for the Larmor radius:

$$r_L^{(e)} = \frac{V_{Te}}{\omega_B^{(e)}} = 3.83 \times 10^{-2} \frac{\sqrt{T_e \text{ (K)}}}{B \text{ (G)}}, \text{ cm}, \quad (5.56)$$

and

$$r_L^{(p)} = \frac{V_{Tp}}{\omega_B^{(p)}} = 1.64 \frac{\sqrt{T_p \text{ (K)}}}{B \text{ (G)}}, \text{ cm.} \quad (5.57)$$

At $T \approx 2 \times 10^6 \text{ K}$ and $B = 3,000 \text{ G}$ we find $r_L^{(e)} \approx 0.2 \text{ mm}$ and $r_L^{(p)} \approx 1 \text{ cm}$.

Exercise 5.3 (Sect. 5.1). During *solar flares* many electrons are accelerated to energies higher than 20–30 keV. These electrons produce the hard X-ray bremsstrahlung emission. The lower boundary of the spectrum of accelerated electrons is not well known because the thermal X-ray emission of the high-temperature (super-hot) plasma masks the lower boundary of the non-thermal X-ray spectrum. Assuming that the lower energy of accelerated electrons $\mathcal{K} \approx 30 \text{ keV}$, find their velocity and the Larmor radius in the corona.

Answer. The kinetic energy of a particle

$$\mathcal{K} = \mathcal{E} - mc^2, \quad (5.58)$$

where $\mathcal{E}$ is the total energy (5.2), $mc^2 = 511 \text{ keV}$ for an electron. Since $\mathcal{K}/mc^2 \ll 1$, formula (5.58) can be used in the non-relativistic limit: $\mathcal{K} = mv^2/2$. From here the velocity of a 30 keV electron $v \approx 10^{10} \text{ cm s}^{-1} \approx 0.3 c$.

The Larmor radius of a non-relativistic electron according to (5.14)

$$r_L^{(e)} = 5.69 \times 10^{-8} \frac{v_{\perp} \text{ (cm s}^{-1}\text{)}}{B \text{ (G)}}. \quad (5.59)$$

For a 30 keV electron

$$r_L^{(e)} \approx 5.6 \times 10^2 \frac{1}{B \text{ (G)}}. \quad (5.60)$$

Above a sunspot with $B \approx 3,000 \text{ G}$ the Larmor radius $r_L^{(e)} \approx 2 \text{ mm}$. Inside a coronal *magnetic trap* with a field $B \approx 100 \text{ G}$ the electrons with kinetic energy $\mathcal{K} \approx 30 \text{ keV}$ have the Larmor radius $r_L^{(e)} \approx 6 \text{ cm}$.

Exercise 5.4 (Sect. 5.1). Under conditions of the previous Exercise estimate the Larmor radius of a proton moving with the same velocity as a 30 keV electron.

Answer. For a non-relativistic proton it follows from formula (5.14) that

$$r_L^{(p)} = 1.04 \times 10^{-4} \frac{v_{\perp} \text{ (cm s}^{-1}\text{)}}{B \text{ (G)}}, \text{ cm.} \quad (5.61)$$

Above a sunspot a proton with velocity $\approx 0.3 c$ has the Larmor radius $\approx 3 \text{ m}$. Inside a coronal trap with magnetic field $\approx 100 \text{ G}$ the Larmor radius $\approx 10^4 \text{ cm}$. So

non-relativistic protons (and other ions) can be well trapped in coronal magnetic traps including collapsing ones

(see Part II, Chap. 7). This is important for the problem of ion acceleration in solar flares.

Exercise 5.5 (Sect. 5.1). The stronger magnetic field, the smaller is the Larmor radius r_L of an electron. Find the condition when r_L is so small as the *de Broglie* wavelength of the electron

$$\lambda_B = \frac{h}{m_e v} = 1.22 \times 10^{-7} \frac{1}{\sqrt{\mathcal{K}(\text{eV})}}. \quad (5.62)$$

Here h is Planck's constant, $\mathcal{K}$ is the kinetic energy (5.58) of the electron. If $\mathcal{K} = 1 \text{ eV}$, the de Broglie wavelength $\lambda_B \approx 10^{-7} \text{ cm} \approx 10 \text{ \AA}$.

Answer. In the non-relativistic limit, the electron with kinetic energy $\mathcal{K}$ has the Larmor radius

$$r_L = 3.37 \frac{\sqrt{\mathcal{K}(\text{eV})}}{B(\text{G})}, \text{ cm}. \quad (5.63)$$

When the energy of the electron is 1 eV and the field has a strength of 1 G, the Larmor radius $r_L \approx 3 \text{ cm}$. However for a field of $3 \times 10^7 \text{ G}$, the Larmor radius is diminished to the de Broglie wavelength $\approx 10^{-7} \text{ cm}$. So for *white dwarfs* which have $B > 10^7 \text{ G}$, and especially for *neutron stars* (see Becker 2009), we have to take into account that

the *quantization* effect of the magnetic field: the Larmor radius is no longer arbitrary but can take only certain definite values.

We call a magnetic field the *superstrong* one, if $r_L < \lambda_B$. Substituting (5.63) and (5.62) into this condition, we rewrite it as follows

$$B > 3 \times 10^7 \mathcal{K}(\text{eV}), \text{ G}. \quad (5.64)$$

In superstrong fields the classic theory of particle motion, developed above, is no longer valid and certain *quantum effects* appear.

The energy difference between the levels of a non-relativistic electron in a superstrong field is

$$\delta \mathcal{E}_B \approx \frac{eB}{mc} \frac{h}{2\pi} \sim 10^{-8} B, \text{ eV}. \quad (5.65)$$

On the other hand, the difference between energy levels in an atom, for example a hydrogen atom, is of about 10 eV; this is comparable with $\delta \mathcal{E}_B$ in a superstrong field $B > 10^8 - 10^9 \text{ G}$. In ordinary conditions B is not so large and does not affect the internal structure of atoms.

Inside and near neutron stars $B > 10^{11} - 10^{12}$ G. In such fields a lot of abnormal phenomena come into existence due to the profound influence of the external field on the interior of atoms. For example, the electron orbits around nuclei become very oblate. Two heavy atoms, e.g. iron atoms, combine into a molecule (Fe_2) and, moreover, these molecules form polymolecular substances, which are constituents of the hard surface of neutron stars (Ruderman 1971; Rose 1998).

Exceedingly superstrong (ultrastrong) fields, $\gtrsim 10^{14}$ G, are suggested in the so-called *magnetars*, the highly-magnetized, newly-born neutron stars (see Sect. 19.1.3).

Exercise 5.6 (Sect. 5.1). Is it justified to neglect the *radiation reaction* in the motion equation (5.8) while considering the gyromotion of electrons in astrophysical plasmas?

Answer. In the non-relativistic limit $v^2 \ll c^2$, the total energy radiated per unit time by a charge e moving with acceleration $\ddot{\mathbf{r}}$ can be calculated in the dipolar approximation (see Landau and Lifshitz 1975, Chap. 9, Sect. 67):

$$I = \frac{2}{3c^3} \ddot{\mathbf{d}}^2. \quad (5.66)$$

Here $\mathbf{d} = e \mathbf{r}$ and $\ddot{\mathbf{d}} = e \ddot{\mathbf{r}}$.

In a uniform magnetic field $\mathbf{B}$, an electron moves in a helical trajectory. For the transversal motion in the Larmor orbit $r = r_L$, the total power radiated by the electron

$$I = \frac{2}{3c^3} e^2 r_L^2 \omega_B^4 = \frac{2e^2}{3c^3} v^2 \omega_B^2. \quad (5.67)$$

Here $v = \omega_B r_L$ is the velocity of the electron in the Larmor orbit.

Let us estimate the strength of the magnetic field such that an electron with kinetic energy $\mathcal{K} = mv^2/2$ would radiate an appreciable amount of energy during one period of gyration, $\tau_B = 2\pi/\omega_B$. Consider a ratio

$$\gamma_r = \frac{\tau_B}{\tau_r} = \frac{1}{\mathcal{K}} \frac{d\mathcal{K}}{dt} \frac{2\pi}{\omega_B}. \quad (5.68)$$

Substituting (5.67) in (5.68) gives

$$\gamma_r = \frac{8\pi}{3} \frac{e^3}{(mc^2)^2} B \approx 1.4 \times 10^{-15} B \text{ (G)}. \quad (5.69)$$

Therefore, while considering the gyromotion of non-relativistic electrons in astrophysical plasmas, the radiation reaction could be important in the motion (5.8) only in ultrastrong magnetic fields with

$$B \gtrsim \frac{3}{8\pi} \frac{(mc^2)^2}{e^3} \approx 7 \times 10^{14} \text{ G}. \quad (5.70)$$

However other physical processes already dominate under such conditions; see discussion in Exercise 5.5.

Recall that formula (5.67) is not valid for a relativistic electron moving in the Larmor orbit; see next Exercise.

Exercise 5.7 (Sect. 5.1). For a relativistic electron moving in the Larmor orbit with a speed $v = \beta c$, the total power of radiation is given by formula (see Landau and Lifshitz 1975, Chap. 9, Sect. 74):

$$I = \frac{2}{3c^3} \frac{e^4}{m^2} \frac{\beta^2}{1 - \beta^2} B^2. \quad (5.71)$$

Therefore, in contrast to the non-relativistic formula (5.67), $I \rightarrow \infty$ when $\beta \rightarrow 1$. Find the rate of energy loss for such an electron.

Answer. According to (5.4), for a relativistic particle

$$\mathcal{E}^2 = (pc)^2 + (mc^2)^2. \quad (5.72)$$

By using this expression we rewrite formula (5.71) as follows

$$\frac{d\mathcal{E}}{dt} = -I = \frac{2e^4 B^2}{3m^4 c^7} ((mc^2)^2 - \mathcal{E}^2). \quad (5.73)$$

From here we find

$$\frac{\mathcal{E}}{mc^2} = \text{cth} \left(\frac{2e^4 B^2}{3m^3 c^5} t + \text{const} \right). \quad (5.74)$$

With an increase of time t , the particle's energy monotonously decreases to the value $\mathcal{E} = mc^2$ with the characteristic time

$$\tau_r = \frac{3m^3 c^5}{2e^4 B^2}. \quad (5.75)$$

Comparing between this time and $2\pi \omega_B^{-1}$ gives us the characteristic value of magnetic field

$$B = \frac{3m^2 c^4}{4\pi e^3} (1 - \beta^2)^{1/2}. \quad (5.76)$$

We see that $B \rightarrow 0$ when $\beta \rightarrow 1$. So, for relativistic electrons, there is no need in strong magnetic fields to radiate efficiently unless they become non-relativistic particles (see Exercise 5.6). This means that

for relativistic electrons, the radiative losses of energy can be important even in relatively weak magnetic fields.

That is why the *synchrotron radiation* is very widespread in astrophysical conditions (e.g., [Ginzburg and Syrovatskii 1965](#)). It was the first radio-astronomical radiation mechanism which had been successfully used by classical astrophysics to interpret the continuum spectrum of the *Crab nebula*. The synchrotron mechanism of radio emission works in any source which contains relativistic electrons in a magnetic field: in the solar corona during flares, in the Jovian magnetosphere, interstellar medium, supernova remnants etc.

Exercise 5.8 (Sect. 5.2.3). Consider an actual force acting on a particle gyrating around the central field line in the magnetic mirror configuration shown in Fig. 5.7.

Answer. Let us use the cylindrical coordinates (r, φ, z) with the axis z along the central field line as shown in Fig. 5.7. In the weakly inhomogeneous magnetic field, the predominant component is B_z but there is a small component B_r which produces the z component of the Lorentz force:

$$F_z = -\frac{q}{c} v_\varphi B_r. \quad (5.77)$$

Here the φ -component is the gyromotion velocity $\mathbf{v}_\perp$, for a positively charged particle $q > 0$ but $v_\varphi < 0$ (cf. Fig. 5.6).

The component B_r of the magnetic field can be found from condition $\text{div } \mathbf{B} = 0$ as follows:

$$B_r = -\frac{1}{2} r \frac{\partial B_z}{\partial z}. \quad (5.78)$$

Substituting (5.78) into (5.77) gives

$$F_z = -\mathcal{M} \frac{\partial B_z}{\partial z}, \quad (5.79)$$

where $\mathcal{M}$ is the magnetic moment (5.33) of the gyrating particle.

Exercise 5.9 (Sect. 5.2.3). Derive formula (5.32) for the last term in the averaged (5.31).

Answer. We have to write down the following expression explicitly

$$(\mathbf{r}_L \times \mathbf{n}) \times [(\mathbf{r}_L \cdot \nabla) \mathbf{B}(\mathbf{R})]$$

and then to average it. It is a matter to do that, once we make use of the following tensor identities:

$$(\mathbf{a} \times \mathbf{b})_\alpha = e_{\alpha\beta\gamma} a_\beta b_\gamma.$$

Here $e_{\alpha\beta\gamma}$ is the unit antisymmetric tensor, and

$$e_{\alpha\beta\gamma} e_{\mu\nu\gamma} = \delta_{\alpha\mu} \delta_{\beta\nu} - \delta_{\alpha\nu} \delta_{\beta\mu}.$$

On rearrangement, we average the last term in (5.31) to obtain

$$\frac{e}{mc} \langle \dot{\mathbf{r}}_{\text{L}} \times [(\mathbf{r}_{\text{L}} \cdot \nabla) \mathbf{B}(\mathbf{R})] \rangle = -\frac{\mathcal{M}}{m} \nabla B, \quad (5.80)$$

where

$$\mathcal{M} = \frac{1}{c} \frac{e \omega_{\text{B}}}{2\pi} (\pi r_{\text{L}}^2) = \frac{1}{c} JS \quad (5.81)$$

is the *magnetic moment* of a particle on the Larmor orbit (see Fig. 5.6).

Chapter 6

Adiabatic Invariants in Astrophysical Plasma

Abstract Adiabatic invariants are useful to understand many interesting properties of collisionless plasma in cosmic magnetic fields: trapping and acceleration of charged particles in collapsing magnetic traps, the Fermi acceleration, “cosmic rays” origin.

6.1 General Definitions

As is known from mechanics (see Landau and Lifshitz 1976, Chap. 7, Sect. 49), the so-called *adiabatic invariants* remain constant under changing conditions of motion, if these changes are slow. Recall that the system executing a *finite one-dimensional* motion is assumed to be characterized by a parameter λ that is slowly—adiabatically—changing with time:

$$\lambda / \dot{\lambda} \gg T. \quad (6.1)$$

Here T is a characteristic time for the system (e.g., a particle in given fields) motion.

More precisely, if the parameter λ did not change, the system would be closed and would execute a strictly periodic motion with the period T like a simple pendulum in gravitational field. In this case the energy of the system, $\mathcal{E}$, would be invariant.

Under the slowly changing parameter λ , if $\dot{\mathcal{E}} \sim \dot{\lambda}$, then the integral

$$I = \oint P dq, \quad (6.2)$$

rather than the energy $\mathcal{E}$, is conserved. Here P and q are the *generalized* momentum and coordinate, respectively. The integral is taken along the trajectory of motion under given $\mathcal{E}$ and λ . The integral I is referred to as the adiabatic invariant.

6.2 Two Main Invariants

6.2.1 Motion in the Larmor Plane

The motion of a charged particle in *slowly changing weakly inhomogeneous fields* has been considered in the previous chapter. Several types of periodic motion have been found. In particular, the particle's motion in the plane perpendicular to the magnetic field—the Larmor motion—is periodic. Let $\mathbf{P}$ be the generalized momentum. According to definition (6.2) for such a motion the adiabatic invariants are the integrals

$$I_1 = \oint P_1 dq_1 = \text{const} \quad \text{and} \quad I_2 = \oint P_2 dq_2 = \text{const},$$

taken over a period of the motion of coordinates q_1 and q_2 in the plane of the Larmor orbit.

It is convenient to combine these integrals, that is simply to add them together:

$$I = \oint \mathbf{P}_\perp \cdot d\mathbf{q} = \text{const}. \quad (6.3)$$

Here

$$\mathbf{P}_\perp = \mathbf{p}_\perp + \frac{e}{c} \mathbf{A}$$

is the generalized momentum (see Landau and Lifshitz 1975, Chap. 3, Sect. 16) projection onto the plane mentioned above. In this plane the generalized coordinate vector $\mathbf{q} = \mathbf{r}_\perp$, $d\mathbf{q} = d\mathbf{r}_\perp = (r_\perp d\varphi) \mathbf{e}_\varphi$ with $0 \leq \varphi \leq 2\pi$. The vector potential $\mathbf{A}$ is perpendicular to the vector $\mathbf{B}$ since $\mathbf{B} = \text{rot} \mathbf{A}$, and $\mathbf{p}$ is the ordinary kinetic momentum of a particle, directed also along the Larmor orbit.

Now perform the integration in formula (6.3)

$$\begin{aligned} I &= \oint \mathbf{P}_\perp \cdot d\mathbf{r}_\perp = \oint \mathbf{p}_\perp \cdot d\mathbf{r}_\perp + \frac{e}{c} \oint \mathbf{A} \cdot d\mathbf{r}_\perp = \\ &= 2\pi r_\perp p_\perp - \frac{e}{c} \int_S \text{rot} \mathbf{A} \cdot d\mathbf{S} = \end{aligned}$$

by virtue of the Stokes theorem

$$= 2\pi r_\perp p_\perp - \frac{e}{c} \int_S \mathbf{B} \cdot d\mathbf{S} = 2\pi r_\perp p_\perp - \frac{e}{c} B \pi r_\perp^2. \quad (6.4)$$

Substituting $r_\perp = c p_\perp / eB$ (cf. formula (5.17)) into (6.4) gives

$$I = \frac{\pi c}{e} \frac{p_{\perp}^2}{B} = \text{const.}$$

Thus we come to the conclusion that the conserving quantity is

$$\boxed{\frac{p_{\perp}^2}{B} = \text{const.}} \quad (6.5)$$

This quantity is called the *first* or *transversal* adiabatic invariant.

According to definition (5.33), the particle magnetic moment for the Larmor orbit is

$$\mathcal{M} = \frac{1}{c} JS = \frac{p_{\perp}^2}{2mB} = \frac{\mathcal{K}_{\perp}}{B}. \quad (6.6)$$

Here use is made of the non-relativistic formula for the Larmor frequency (5.11) and the non-relativistic kinetic energy of the particle transversal motion is designated as

$$\mathcal{K}_{\perp} = \frac{p_{\perp}^2}{2m}.$$

When formula (6.5) is compared with (6.6), it is apparent that

the *magnetic moment* of a particle at the Larmor orbit is conserved in the non-relativistic approximation.

In the relativistic limit the particle magnetic moment (6.6) does not remain constant; however the first adiabatic invariant can be interpreted to represent the magnetic field flux through the surface covering the particle Larmor orbit,

$$\Phi = B \pi r_{\perp}^2 = \frac{\pi c^2}{e^2} \frac{p_{\perp}^2}{B} = \text{const.} \quad (6.7)$$

This also follows directly from (6.4), when we substitute the relativistic formula

$$p_{\perp} = r_{\perp} \frac{eB}{c} \quad (6.8)$$


into the first term on the right-hand side of formula (6.4). We obtain

$$I = \frac{e}{c} (B \pi r_{\perp}^2) = \frac{e}{c} \Phi. \quad (6.9)$$

Therefore

in the *relativistic* case, the *magnetic field flux* Φ through the surface S covering the particle Larmor orbit is conserved.

Fig. 6.1 A converging flux of field lines forms a magnetic “mirror”. At the point M_1 , the parallel component of momentum reverses under action of the diamagnetic force (5.35)



6.2.2 Magnetic Mirrors and Traps

Let us imagine the time-independent magnetic field, the field lines forming the convergent flux. As a rule, the field takes such a form in the vicinity of its sources, for instance, a sunspot S in the photosphere Ph in Fig. 6.1.

The particle transversal momentum is

$$p_{\perp} = p \sin \theta, \quad (6.10)$$

it being known that $p = \text{const}$, since by virtue of (5.6) we have $\mathcal{E} = \text{const}$. Substituting (6.10) into (6.5) gives

$$\frac{\sin^2 \theta}{B} = \text{const} = \frac{\sin^2 \theta_0}{B_0}$$

or

$$\sin^2 \theta = \frac{B}{B_0} \sin^2 \theta_0. \quad (6.11)$$

This formula shows that, for the increasing B , a point M_1 must appear in which $\sin^2 \theta_1 = 1$. The corresponding value of the field is equal to

$$B_1 = B_0 / \sin^2 \theta_0. \quad (6.12)$$


At this point the particle “reflection” takes place:

$$p_{\parallel} = p \cos \theta_1 = 0.$$

The regions of convergent field lines are frequently referred to as magnetic “mirrors”.

So, if there is a field-aligned gradient of the magnetic-field strength, the component of velocity parallel to the field decreases as the particle moves into a region

Fig. 6.2 A coronal magnetic tube as a trap for particles accelerated in a solar flare. $\theta < \theta_0$ is the loss cone. Motion between the mirror points M_1 and M_2 is called bounce motion



of increasing field magnitude, although the total velocity is conserved. Eventually, under action of the diamagnetic force (5.35), the parallel velocity reverses (see the point M_1 in Fig. 6.1). Such reflections constitute the principle of a *magnetic trap*. For example, magnetic fields create traps for fast particles in the solar atmosphere as shown in Fig. 6.2. The particles are injected into the coronal magnetic tubes called flaring loops, during a solar flare. Let us suppose that this injection occurs at the loop apex.

Let us also suppose that, having hit the chromosphere Ch , the particles “die” because of collisions. The particles do not return to the coronal part of the trap, their energy being transferred to the chromospheric plasma, leading to its heating. Such particles are termed *precipitating* ones. Their pitch-angles have to be less than θ_0 :

$$\theta < \theta_0 \quad (6.13)$$


with

$$\theta_0 = \arcsin \sqrt{B_0 / B_1} \quad (6.14)$$

in accordance with (15.72). Here B_0 is the magnetic field at the trap apex, B_1 is the field at the upper chromosphere level at the mirror points M_1 and M_2 as shown in Fig. 6.2. The quantity B_1 / B_0 is called the *cork ratio*.

The angle region (6.13) is termed the *loss cone*. The particles with the initial momenta inside the loss cone precipitate from the trap. By contrast, the particles with $\theta > \theta_0$ at the loop apex experience reflection and do not reach the chromosphere. Such particles are termed *trapped* ones.

Fig. 6.3 An idealized model of a long trap with a short moving cork. Unless a charged particle has its velocity vector very close to the axis of the trap, it is reflected back and forth between the mirrors, thereby remaining trapped



An interesting situation arises if the diffusion of the trapped particles into the loss cone is slower than their precipitation from the trap into the chromosphere. Then the distribution function becomes anisotropic, since the loss cone is “eaten away”, and non-equilibrium. The situation is quite analogous to the case of the distribution function formation with the positive derivative in some velocity region, like the *bump-on-tail* distribution (Fig. 7.2). As a result, some *kinetic instabilities* (e.g., [Silin 1971](#); [Schram 1991](#); [Shu 1992](#)) can be excited which lead to such plasma processes as wave excitation, anomalous particle transfer owing to the particles scattering off the waves, and *anomalous diffusion* into the loss cone (see also Chap. 7).

6.2.3 Bounce Motion

Let us consider another example of a particle motion in a magnetic trap, namely that of a motion between two magnetic corks, the transversal drift being small during the period of longitudinal motion. In other words, the conditions of periodic longitudinal motion are changing adiabatically slowly. Then the *second* adiabatic invariant, referred to as the *longitudinal* one, is conserved:

$$I = \oint P_{\parallel} dl = p \oint \sqrt{1 - \sin^2 \theta} dl = p \oint \sqrt{1 - \frac{B}{B_1}} dl. \quad (6.15)$$

Here account is taken of the facts that the vector $\mathbf{A}$ is perpendicular to the vector $\mathbf{B}$ and

$$p = |\mathbf{p}| = \text{const}$$

since $\mathcal{E} = \text{const}$. The formula (6.11) for the first adiabatic invariant is used in the last equality in (6.15).

Let us apply formula (6.15) to the case of a long trap with short corks: $l \ll L$ in Fig. 6.3. The longitudinal invariant for such a trap is

$$I = \oint p_{\parallel} dl \approx 2 p_{\parallel} L = \text{const}.$$

Therefore the second adiabatic invariant is associated with the cyclical *bounce* motion between two mirrors or corks and is equal to

$$p_{\parallel} L = \text{const.}$$

(6.16)

Let us suppose now that the distance between the corks is changing, that is the trap length $L = L(t)$. Then from (6.16) it follows that

$$p_{\parallel}(t) = p_{\parallel}(0) \frac{L(0)}{L(t)}. \quad (6.17)$$

It is evident from (6.17) that:

- (a) Increasing the distance between the corks decreases the longitudinal momentum and, consequently, the particle energy, and
- (b) Particle acceleration takes place in the trap if two magnetic corks are approaching each other as is shown by vector $\mathbf{v}$ in Fig. 6.3.

The former case can describe the so-called “adiabatic cooling” of accelerated particles, for example, in a magnetic trap which is captured by the *solar wind* and is expanding into the interplanetary space. The latter case is more interesting. It corresponds to the Fermi mechanism considered in the next section.

6.2.4 The Fermi Acceleration

The famous theory of Fermi (1949) discussed the so-called interstellar “clouds” that carry magnetic fields and could reflect charged particles. The same role could be played for instance by magnetic inhomogeneities in the solar wind or interplanetary medium. Fermi visualized that charged particles can be accelerated by being repeatedly hit by the moving magnetic clouds.

The energy of a particle, $\mathcal{E}$, will increase or decrease according to whether a cloud (an inhomogeneity of magnetic field) that causes the reflection moves toward the particle (*head-on* collision) or away from it (*overtaken* collision). The particle gains energy in a head-on collision but there can be also *trailing* collisions in which energy is lost. It was shown by Fermi (1949, 1954) that

on the average, the energy increases because the head-on collisions are more probable than the overtaking collisions

(see a non-relativistic treatment of the problem in Exercise 6.1). Through this *stochastic* mechanism

the energy of the particle increases at a rate that, for relativistic particles, is proportional to their energy

(Exercise 6.2):

$$\frac{d\mathcal{E}}{dt} \propto \mathcal{E}. \quad (6.18)$$

The higher the energy $\mathcal{E}$, the faster acceleration. This is the most important feature of the Fermi mechanism. We shall call this mechanism the *stochastic* Fermi acceleration. There is a parameter (a relative velocity of magnetic clouds) which characterizes the coefficient of proportionality in the problem under consideration (see Exercise 6.2).

From formula (6.18) follows that the energy $\mathcal{E}$ increases exponentially with time:

$$\mathcal{E}(t) = \mathcal{E}_0 \exp \frac{t}{t_a}, \quad (6.19)$$

where $\mathcal{E}_0$ is the initial energy, t_a is the acceleration time scale.


Large-scale MHD turbulence is generally considered as a source of magnetic inhomogeneities accelerating particles in astrophysical plasma. Acceleration of particles by MHD turbulence has long been recognized as a possible mechanism for solar and galactic cosmic rays (Davis 1956).

Though the Fermi acceleration has been popular, it appears to be neither efficient nor selective. A mirror reflects particles on a *nonselective* basis: thermal particles may be reflected as well as suprathermal ones. Therefore one is faced with the conclusion (Eichler 1979) that *most of the energy in the MHD turbulence goes into bulk heating* of the plasma rather than the selective acceleration of only a minority of particles. We shall come back to this question in Chap. 7.

If we somehow arrange that only head-on collisions take place, then the acceleration process will be much more efficient. We should call the acceleration resulting from such a situation the *regular* Fermi acceleration. More often, however, this mechanism is called the first-order (in the small parameter v_m/c , where v_m is the velocity of the moving magnetic clouds; see Exercise 6.1). The simplest example of this type mechanism is a pair of converging shock waves (Wentzel 1964). In this case, there is no deceleration by trailing collisions (see formula (6.22) in Exercise 6.1) that reduce the net efficiency to the second order in the parameter v_m/c (Exercise 6.2).

One of several well-known examples of this type of the Fermi acceleration is the impulsive (with high rate of energy gain) acceleration between two approaching shocks S_{up} in the model of a flaring loop as shown in Fig. 6.4. To explain the hard X-ray and gamma-ray time profiles in solar flares, Bai et al. (1983) assumed that pre-accelerated electrons penetrate into the flare loop and heat the upper chromosphere to high-temperatures rapidly. As a consequence of the fast expansion of a high-temperature plasma into the corona—the process of chromospheric “evaporation”, two shock waves S_{up} move upward from both footpoints.

Fig. 6.4 The flare-heated chromospheric plasma P rapidly expands into the corona. Particle acceleration of the first order Fermi type may occur in a magnetic loop between two converging shock waves S_{up}



Energetic particles are to be reflected only by colliding with the shock fronts. In such a way, the regular Fermi acceleration of particles between two shocks was suggested as a mechanism for the second-step acceleration of protons and electrons in flares. A similar example of the regular Fermi-type acceleration also related to a *collapsing* ($L(t) \rightarrow 0$) magnetic trap in solar flares (Somov and Kosugi 1997) is considered in Part II, Chap. 7.

The cosmic rays (see Sect. 5.1.3) were assumed to be accelerated by crossing shock fronts generated in explosive phenomena such as supernovae. However a very simple dimensional argument shows the kind of difficulties encountered even by the most violent phenomena in the Universe.

The more energetic are the particles, the larger are their Larmor radius and/or the higher are the magnetic fields B necessary to confine them within the limits of a cosmic accelerator.


The size of an accelerator R must be larger than the Larmor radius of a particle. The product BR large enough to suit the 10^{20} eV energy range exists in no known standard astrophysical object.

6.3 The Flux Invariant

Let us consider the axisymmetric trap which is modelled on, for example, the Earth's magnetic field. Three types of the particle's motion are shown in Fig. 6.5.

First, on the time scales of Larmor period, the particle spirals about a field line. Second, since there is a field-aligned gradient of the field strength, the particle oscillates between two mirrors M_1 and M_2 . Third, if the guiding center does not lie on the trap's symmetry axis then *the radial gradient of field* (cf. Fig. 5.9) *causes the drift* around this axis. This drift (formula (5.40)) is superimposed on the particle's oscillatory motion of rotation.

Fig. 6.5 Particle drift in a trap, due to the radial gradient of field



As the particle bounces between the mirrors and also drifts from one field line to another one, it traces some magnetic surface S_d . The latter is called the *drift shell*. Let T_s be the period of particle motion on this surface.

If the magnetic field $\mathbf{B} = \mathbf{B}(t)$ is changing so slowly that

$$B / \dot{B} \gg T_s,$$

then a *third* adiabatic invariant, referred to as a flux one, is conserved:

$$\Phi = \int_S \mathbf{B} \cdot d\mathbf{S} = \text{const}. \quad (6.20)$$

Thus the first adiabatic invariant implies conservation of the magnetic flux through the Larmor orbit, $B \pi r_L^2$, whereas

the flux invariant implies conservation of the magnetic flux through the closed orbit of guiding center motion,

that is the flux through the shaded surface S in Fig. 6.5.

6.4 Approximation Accuracy. Exact Solutions

Adiabatic invariants have been obtained in the approximation of weakly inhomogeneous slowly changing magnetic fields. The invariants are *approximate* integrals of motion, widely used in plasma astrophysics. However we should not forget two important facts. First, the adiabatic theory has a limited, though *exponential*, accuracy. Second, this theory has a limited, though wide, area of applicability. The second volume of this book (Part II) will be devoted to the effect of magnetic reconnection and will present a situation when the adiabatic theory a priori does not apply.

Exact solutions to the equations of charged particle motion usually require numerical integration. The motion in the field of a magnetic dipole is a simple case that, nevertheless, is of practical significance. The reason for that is the possibility to approximate the Earth's magnetic field at moderately large distances by the dipole field. It was [Störmer \(1955\)](#) who contributed significantly to the solution of this problem.

Two types of trajectories are considered.

- (a) The first type is the trajectories coming from infinity and returning there. These have been calculated in order to find out whether a particle can reach a given point on the Earth along a given direction from cosmic space. An answer to this question is important for cosmic ray theories. For each point on the Earth and for each direction the so-called “*threshold rigidity*” has been calculated. If a rigidity is greater than the threshold one, then the particle can reach the point. The vertical threshold rigidity is the most universally used one. This characterizes particle arrival in the direction of the smallest column depth of the Earth atmosphere.
- (b) The second type is the orbits of *trapped* particles. Two radiation belts of the Earth, the inner and the outer, have been shown to exist. The mechanisms which generate trapped particles are not yet fully understood. They are presumably related to geomagnetic storms ([Tverskoy 1969](#); [Walt 1994](#)).

Both gradient drift and curvature drift cause the positive particles in the radiation belt to drift westward in the Earth dipole magnetic field. Thus the radiation belt forms a ring of westward current circulating the Earth. This current tends to decrease the strength of the basic northward magnetic field observed at low latitudes on the Earth surface. There is a simple theoretical relationship between the depression of the magnetic field at the surface of the Earth and the total energy in the trapped particles. This relation allows us to use the observed change of the magnetic field as an indication of the amount of the energy in ring-current particles.

Recommended Reading: [Northrop \(1963\)](#), [Kivelson and Russell \(1995\)](#).

6.5 Practice: Exercises and Answers

Exercise 6.1 (Sect. 6.2.4). Show that a non-relativistic particle on average gains energy in collisions with moving magnetic clouds.

Answer. Let us consider the simplest model of one-dimensional motions of clouds: half of the clouds are moving in one direction and the other half moving in the opposite direction with the same velocity v_m .

Let a particle of initial velocity V_0 undergo a *head-on* collision. The initial velocity seen from the rest frame of the cloud is $V_0 + v_m$. If the collision is elastic, the particle bounces back in the opposite direction with the same magnitude of velocity

$V_0 + v_m$ in this rest frame. In the observer's frame, the reflected velocity appears to be $V_0 + 2v_m$. Hence the gain of kinetic energy $\mathcal{K}$ according to the observer equals

$$\delta\mathcal{K}_+ = \frac{1}{2} m (V_0 + 2v_m)^2 - \frac{1}{2} m V_0^2 = 2m v_m (V_0 + v_m). \quad (6.21)$$

Similarly, the energy loss in a *trailing* collision

$$\delta\mathcal{K}_- = -2m v_m (V_0 - v_m). \quad (6.22)$$

The probability of head-on collisions is proportional to the relative velocity $V_0 + v_m$, whereas the probability of trailing collisions is proportional to $V_0 - v_m$. Therefore the average gain of kinetic energy is equal to

$$\delta\mathcal{K}_{av} = \delta\mathcal{K}_+ \frac{V_0 + v_m}{2v_m} + \delta\mathcal{K}_- \frac{V_0 - v_m}{2v_m} = 4m v_m^2. \quad (6.23)$$

So a particle is accelerated.

Exercise 6.2 (Sect. 6.2.4). Prove the Fermi formula (6.18) for a *relativistic* particle.

Answer. Make the same procedure as that one in Exercise 6.1 by using the corresponding expressions in special relativity to see that the average energy gain

$$\delta\mathcal{E}_{av} = 4 \left(\frac{v_m}{c} \right)^2 \mathcal{E}. \quad (6.24)$$

Formula (6.24) obviously reduces to (6.23) in the non-relativistic limit on putting $\mathcal{E} = mc^2$.

So the average energy gain is proportional to the energy. Therefore the energy of a relativistic particle suffering repeated collisions with moving magnetic clouds increases according to formula

$$\frac{d\mathcal{E}}{dt} = \alpha_f \mathcal{E}, \quad (6.25)$$

where α_f is a constant. Q.e.d.

Note also that the average energy gain (6.24) is proportional to the dimensionless parameter $(v_m/c)^2$. Since actual interstellar clouds are moving at non-relativistic velocities, this parameter should be a very small number. Hence the acceleration process is quite inefficient. Because of this quadratic dependence on v_m , this process is often referred as the *second-order* Fermi acceleration.

If only head-on collisions take place, then the acceleration is much more efficient. It follows from formula (6.21) that, for $V_0 \gg v_m$, the energy gain will depend linearly on v_m . So the acceleration resulting from such conditions is called the *first-order* Fermi acceleration. Such conditions are well possible, for example, in collapsing magnetic traps created by the magnetic reconnection process in solar flares (see Part II, Chap. 7).

Powerful shock waves in astrophysical plasma with magnetic field (like the solar wind) may well provide sites for the first-order Fermi acceleration. Magnetic inhomogeneities are expected on both sides of the shock front. It is possible that a charged particle is trapped near the front and repeatedly reflected from magnetic inhomogeneities on both sides. Such collisions may lead to more efficient acceleration (see Chap. 18) compared to original Fermi's acceleration by moving interstellar clouds.

Chapter 7

Wave-Particle Interaction in Astrophysical Plasma

Abstract The growth or damping of the waves, the emission of radiation, the scattering and acceleration of particles—all these phenomena may result from wave-particle interaction, a process in which a wave exchanges energy with the particles in astrophysical plasma.

7.1 The Basis of Kinetic Theory

7.1.1 The Linearized Vlasov Equation

In this chapter we shall only outline the physics and main methods used to describe the wave-particle interaction in collisionless astrophysical plasmas as well as in Maxwellian plasmas where fast particles interact with electromagnetic waves. In the simplest—*linear*—approach, the idea is in the following.

We assume the unperturbed plasma to be uniform and characterized by the distribution functions $f_k^{(0)}$ of its components k : electrons and ions. The unperturbed plasma is also assumed to be steady. So

$$f_k^{(0)} = f_k^{(0)}(\mathbf{v}). \quad (7.1)$$

Let $\mathbf{B}^{(0)}$ be the unperturbed uniform magnetic field inside the plasma. We further assume that the only zero-order force is the Lorentz force with $\mathbf{E}^{(0)} = 0$.

The dynamics of individual particles is determined by the first-order forces related to the wave electric field $\mathbf{E}^{(1)}$ and wave magnetic field $\mathbf{B}^{(1)}$. To describe these particles we shall use the perturbation function $f_k^{(1)}$, which is linear in $\mathbf{E}^{(1)}$ and $\mathbf{B}^{(1)}$. Under the assumptions made, we see that the Vlasov equation (Sect. 3.1.2) can be a proper basis for the kinetic theory of wave-particle interaction. For this reason we shall realize the following procedure.

- (a) We linearize the Vlasov equation (3.3) together with the Maxwell equations (3.4) for the self-consistent wave field. Equation (3.3) becomes

$$\begin{aligned} & \frac{\partial f_k^{(1)}(X, t)}{\partial t} + v_\alpha \frac{\partial f_k^{(1)}(X, t)}{\partial r_\alpha} \\ & + \frac{e_k}{m_k} \left(\frac{1}{c} \mathbf{v} \times \mathbf{B}^{(0)} \right)_\alpha \frac{\partial f_k^{(1)}(X, t)}{\partial v_\alpha} \\ & - \frac{e_k}{m_k} \left(\mathbf{E}^{(1)} + \frac{1}{c} \mathbf{v} \times \mathbf{B}^{(1)} \right)_\alpha \frac{\partial f_k^{(0)}(\mathbf{v})}{\partial v_\alpha} . \end{aligned} \quad (7.2)$$

The left-hand side of the linear equation (7.2) is the Liouville operator (1.12) acting on the first-order distribution function for particles following the *unperturbed* trajectories in phase space $X = \{\mathbf{r}, \mathbf{v}\}$:

$$\frac{D}{Dt} f_k^{(1)} = - \frac{F_{k,\alpha}^{(1)}}{m_k} \frac{\partial f_k^{(0)}}{\partial v_\alpha} . \quad (7.3)$$

This fact (together with the linear Lorentz force in the right-hand side of (7.3) and the linearized Maxwell equations) can be used to find the general solution of the problem. We are not going to do this here (see Exercise 7.1). Instead, we shall make several simplifying assumptions to demonstrate the most important features of kinetic theory on the basis of the linearized Vlasov equation (7.3).

- (b) Let us consider a small harmonic perturbation varying as

$$f_k^{(1)}(t, \mathbf{r}, \mathbf{v}) = \tilde{f}_k(\mathbf{v}) \exp[-i(\omega t - \mathbf{k} \cdot \mathbf{r})] . \quad (7.4)$$

Substituting the plane wave expression (7.4) with a similar presentation of the perturbed electromagnetic field in (7.2) gives us the following linear equation:

$$\begin{aligned} & i(\omega - \mathbf{k} \cdot \mathbf{v}) \tilde{f}_k(\mathbf{v}) - \frac{e_k}{m_k} \left(\frac{1}{c} \mathbf{v} \times \mathbf{B}^{(0)} \right)_\alpha \frac{\partial \tilde{f}_k(\mathbf{v})}{\partial v_\alpha} \\ & = \frac{e_k}{m_k} \left[\tilde{\mathbf{E}} \left(1 - \frac{\mathbf{k} \cdot \mathbf{v}}{\omega} \right) + \mathbf{k} \left(\frac{\mathbf{v} \cdot \tilde{\mathbf{E}}}{\omega} \right) \right]_\alpha \frac{\partial f_k^{(0)}(\mathbf{v})}{\partial v_\alpha} . \end{aligned} \quad (7.5)$$

Here the Faraday law (1.28) has been used to substitute for the wave magnetic field.

- (c) We shall assume that the waves propagate parallel to the ambient field $\mathbf{B}^{(0)}$ which defines the z axis of a Cartesian system. From Sect. 5.1 it follows that in a uniform magnetic field there exist two constants of a particle's motion, i.e. two integrals of motion or two invariants of motion: the *parallel* velocity

$$\mathbf{v}_\parallel = v_z$$

and the magnitude of the *perpendicular* velocity

$$v_{\perp} = |\mathbf{v}_{\perp}| = \left(v_x^2 + v_y^2 \right)^{1/2}.$$

Hence the unperturbed distribution function

$$f_k^{(0)} = f_k^{(0)}(v_{\parallel}, v_{\perp}), \quad (7.6)$$

as required by Jeans's theorem (Exercise 7.1). Therefore in what follows we can consider two cases of resonance, corresponding two variables in the distribution function (7.6).

7.1.2 The Landau Resonance and Landau Damping

Let us consider the so-called *electrostatic* waves which have only a parallel electric field $\mathbf{E}^{(1)} = \mathbf{E}_{\parallel}$ under the assumption of parallel propagation:

$$\mathbf{k} \times \mathbf{B}^{(0)} = 0. \quad (7.7)$$

In this case the linearized Vlasov equation (7.5) reduces to

$$i(\omega - k_{\parallel} v_{\parallel}) \tilde{f}_k - \frac{e_k}{m_k} \left(\frac{1}{c} \mathbf{v} \times \mathbf{B}^{(0)} \right)_{\alpha} \frac{\partial \tilde{f}_k}{\partial v_{\alpha}} = \frac{e_k}{m_k} \tilde{E}_{\parallel} \frac{\partial f_k^{(0)}}{\partial v_{\alpha}}. \quad (7.8)$$

Now let us find the perturbation of charge density according to definition (3.5):

$$\rho^{q(1)}(\mathbf{r}, t) = \sum_k e_k \int_{\mathbf{v}} f_k^{(1)}(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (7.9)$$

Hence the amplitude

$$\tilde{\rho}^q = \sum_k e_k \int_{\mathbf{v}} \tilde{f}_k(\mathbf{v}) d^3\mathbf{v}. \quad (7.10)$$

When we calculate the charge density by using (7.8), the second term on the left-hand side of this equation vanishes on integration over perpendicular velocity.

Therefore, for parallel propagating electrostatic waves, the harmonic perturbation of charge density is given by

$$\tilde{\rho}^q = -i \tilde{\mathbf{E}}_{\parallel} \sum_k \frac{e_k^2}{m_k} \int_{v_{\parallel}} \frac{1}{(\omega - k_{\parallel} v_{\parallel})} \frac{\partial f_k^{(0)}}{\partial v_{\parallel}} dv_{\parallel}. \quad (7.11)$$




Fig. 7.1 The Landau damping. **(a)** The initial distribution function of thermal electrons with some narrow region centered at the resonance with the wave. **(b)** The distribution function after an evolution due to interaction of the electrons with the wave

Formula (7.11) shows that there is a *resonance* which occurs when

$$\omega - k_{\parallel} v_{\parallel} = 0$$

(7.12)

or when the particle velocity equals the parallel phase velocity of the wave, $\omega/k_{\parallel}$. This is the *Landau resonance*.

A physical picture of Landau resonance is simple.

When the resonance condition (7.12) is satisfied the particle “sees” the electric field of the wave as a *static* electric field in the particle’s rest system

(see Exercise 7.3).


Particles in resonance moving slightly faster than the wave will lose energy, while those moving slightly slower will gain energy. Since the Maxwellian distribution is decreasing with velocity,

in a Maxwellian plasma, near the Landau resonance, there are more particles at lower velocities than at higher velocities. That is why *the plasma gains energy at the expense of the wave*.

This effect, illustrated by Fig. 7.1 (see also Exercise 7.6), is called the *Landau damping* (Landau 1946) or collisionless damping of the electrostatic-type waves. The effect is described by the last term in the following dispersion equation

$$\omega^2 = \left(\omega_{pl}^{(e)}\right)^2 + 3 \frac{k_{\parallel} T}{m_e} \frac{k_{\parallel}^2}{\omega^2} \left(\omega_{pl}^{(e)}\right)^2 + i \pi \frac{\omega^2}{k_{\parallel}^2} \left(\omega_{pl}^{(e)}\right)^2 \frac{\partial f^{(0)}}{\partial v_{\parallel}} \Big|_{v_{\parallel}=\omega/k_{\parallel}}. \quad (7.13)$$

Fig. 7.2 The bump-on-tail distribution function with the resonance condition in the region of a positive slope



Here

$$\omega_{pl}^{(e)} = \pm \left(\frac{4\pi e^2 n_e}{m_e} \right)^{1/2} \quad (7.14)$$

is the *electron plasma frequency* (see Sect. 10.2.1).

Normally we think of damping as a dissipative process and hence expect it to be present only in systems where collisions can convert a part of the wave energy into thermal energy. At first sight, damping in a collisionless system seems mystifying since we ask the question where could the energy have gone. For a negative slope of the distribution function at the phase velocity ω/k , there are more particles which are accelerated than which are decelerated. For this reason the wave puts a net amount of energy in the particles so that there is a loss of wave energy. Therefore the Landau damping is not by randomizing collisions but by a transfer of wave field energy into oscillations of resonant particles.

Landau damping is often the dominant *damping mechanism for waves*, such as ion-acoustic waves and Langmuir waves, in thermal plasma without a magnetic field.

■ The absorption of longitudinal waves in plasma in the thermal equilibrium is often determined by the Landau collisionless damping (e.g., [Zheleznyakov 1996](#)).

On the other hand, if a distribution function has more particles at higher velocities than at lower velocities in some region of phase space as shown in Fig. 7.2, this distribution will be unstable to waves that are in resonance with the particles. This is the known “bump-on-tail” instability. Due to this type of instability, a beam of fast electrons (with velocities much higher than the thermal speed of electrons in the plasma) causes Langmuir waves to grow. Langmuir waves generated through the bump-on-tail instability play an essential role, for example, in solar radio bursts.

There are many examples in plasma astrophysics in which one species (e.g., electrons) moves relative to another. Solar flares produce a significant flux of fast electrons moving through the plasma in the interplanetary space. Fast electrons move away from a planetary shock through the solar wind. Aurorae are produced by fast electrons moving along Earth’s magnetic-field lines. If we consider a stream of

plasma with an average velocity impinging on another plasma at rest, we have just the same situation. The system has an instability such that

the kinetic energy of the relative motion between the plasma streams is fed into a plasma wave of the appropriate phase velocity.

So all the two-stream instabilities have, in fact, the same origin.

The above derivation emphasizes the close relation of the Landau damping with the *Cherenkov effect* (see Exercises 7.2–7.5). It has been definitely pointed out by [Ginzburg and Zheleznyakov \(1958\)](#) that

the Landau damping and the Cherenkov absorption of plasma waves, the inverse Cherenkov effect, are the same phenomenon

initially described in two different ways.

The discussion hitherto has focused on the linear Landau damping, i.e. the behaviour of a small perturbation which satisfies the linearized Vlasov equation. However this picture can be extended to *finite amplitude* perturbations ([Kadomtsev 1976](#), Chap. 4). In the context of plasma astrophysics, this means considering *nonlinear* Landau damping, which generalized the linear theory by incorporating the possibility of mode-mode couplings that allow energy transfer between different modes.

In fact, the linear theory illuminates only a narrow window out of the wealth of all effects related to wave-particle interactions. Mathematically, the *linear theory uses a well-developed algorithm*. Few analytical methods are known to treat the much wider field of nonlinear effects, and most of these methods rely on approximations and lowest-order perturbation theory. The theory of *weak* wave-particle interaction or *weak turbulence* as well as the *quasi-linear* theory for different types of waves are still today the most important parts in astrophysical applications (e.g., [Treumann and Baumjohann 1997](#); [Benz 2002](#)).

7.1.3 Gyroresonance

As for the Landau resonance, we shall use the linear equation (7.5) as a basis, assuming that a wave is propagating parallel to the ambient field $\mathbf{B}^{(0)}$. However, this time, we shall further assume that the wave electric field $\mathbf{E}^{(1)}$ and hence the wave magnetic field $\mathbf{B}^{(1)}$ are perpendicular to the ambient magnetic field.

Under the assumption of a harmonic perturbation (7.4) we shall make use of the so-called *polarized coordinates*:

$$\tilde{E}_l = \frac{\tilde{E}_x + i\tilde{E}_y}{\sqrt{2}}, \quad \tilde{E}_r = \frac{\tilde{E}_x - i\tilde{E}_y}{\sqrt{2}}. \quad (7.15)$$

Subscripts l and r correspond to the waves with left- and right-hand circular polarizations, respectively.

By definition, the wave is right-hand circular polarized if $\tilde{E}_x$ leads $\tilde{E}_y$ by a quarter of a wave period. If, for such a wave, we multiply (7.5) by velocity

$$\mathbf{v}_r = \frac{v_x - i v_y}{\sqrt{2}} \quad (7.16)$$

and integrate over velocity space, making use of (7.6) and the fact that the unperturbed distribution function $f^{(0)}$ is a symmetric function of $v_\perp$, we find the equation which determines (see definition (3.6)) the current density in the harmonic perturbation:

$$\begin{aligned} \tilde{\mathbf{j}}_r^q = -i \sum_k \frac{e_k^2}{m_k} \tilde{E}_r \times \\ \times \int_{\mathbf{v}} \frac{1}{\left(\omega - k_\parallel v_\parallel - s \omega_B^{(k)}\right)} \left[\left(1 - \frac{k_\parallel v_\parallel}{\omega}\right) \frac{\partial f_k^{(0)}}{\partial v_\perp} + \frac{k_\parallel v_\perp}{\omega} \frac{\partial f_k^{(0)}}{\partial v_\parallel} \right] v_\perp d^3 \mathbf{v}. \end{aligned} \quad (7.17)$$

Here $\omega_B^{(k)}$ is the Larmor frequency of a particle of a kind k , the integer s can be positive or negative. The resonance condition in formula (7.17) for current density is the *gyroresonance*:

$$\omega - k_\parallel v_\parallel - s \omega_B^{(k)} = 0.$$

(7.18)

We see that a gyroresonant interaction occurs when the Doppler-shifted wave frequency

$$\omega_D = \omega - k_\parallel v_\parallel, \quad (7.19)$$

as observed by a particle moving with the parallel velocity $v_\parallel$, is an integer multiple s of the Larmor frequency in the guiding center frame, i.e.

$$\omega_D = s \omega_B^{(k)}. \quad (7.20)$$

Depending upon the initial relative phase of the wave and particle, the particle will corotate with either an accelerating or decelerating electric field over a significant portion of its Larmor motion, resulting in an appreciable gain or loss of energy, respectively.

If the particle and transversal electric field rotate in the same sense, the integer $s > 0$, whereas an opposite sense of rotation requires $s < 0$. However the strongest interaction usually occurs when the Doppler-shifted frequency exactly matches the particle Larmor frequency.

The gyroresonance is important for generating waves such as the *whistlers*, the short bursts of audio-frequency radio noise in the ionosphere (Ferencz et al. 2001). Whistlers are polarized predominantly perpendicular to the Earth magnetic field.

For a wave to grow from gyroresonance, there should be a net decrease in particle energy as the particle diffuses down the phase-space density gradient defined by the numerator in formula (7.17),

i.e. by the expression enclosed in large square brackets under the integral in formula (7.17).

For the parallel propagation of a wave in astrophysical plasma, the Landau resonance is associated with parallel electric fields. For perpendicular electric fields, particles and fields can be in gyroresonance. It is clear that the Landau resonance diffuses particles parallel to the ambient magnetic field, whereas *gyroresonance causes diffusion in the pitch angle*. This can be seen in the wave frame, i.e. the frame in which the parallel phase velocity of the wave is zero. If we transform the expression enclosed in large square brackets in formula (7.17) to the wave frame, we find that in this frame the gradient in velocity space is gradient with respect to pitch angle θ . Hence

the main effect of gyroresonance is to cause particles to change pitch angle in the wave frame.

This is contrasted with the Landau resonance, where the diffusion is in the parallel velocity $v_{\parallel}$ due to the term $\partial f^{(0)} / \partial v_{\parallel}$ and therefore mainly in energy, rather than pitch angle.

As such, then the Landau-resonant instabilities are often driven by bump-on-tail distributions of particles, whereas gyroresonant instabilities are driven by *pitch-angle* anisotropy. Thus the gyroresonance-type instabilities can appear as soon as a “tail” or beam is formed in the direction parallel to the background field $\mathbf{B}^{(0)}$. They excite waves that scatter the particles back to a nearly isotropic state.

7.2 Stochastic Acceleration of Particles by Waves

7.2.1 The Principles of Particle Acceleration by Waves

In Sect. 7.1 we considered the resonant interaction between particles and one wave propagating parallel to the uniform magnetic field $\mathbf{B}^{(0)}$ in a uniform plasma without an external electric field $\mathbf{E}^{(0)} = 0$. The dynamics of individual particles was determined by the first-order forces related to the wave electric field $\mathbf{E}^{(1)}$ and wave magnetic field $\mathbf{B}^{(1)}$. We described behavior of these particles by the linearized Vlasov equation (7.2) for the perturbation function $f_k^{(1)}$, which is linear in $\mathbf{E}^{(1)}$ and $\mathbf{B}^{(1)}$.

Under simplifying assumptions made, we saw that, in addition to the Landau resonance (7.12):

$$\omega_D = 0, \quad (7.21)$$

other resonances (7.18) arise in wave-particle interaction. These are the gyroresonances which occur when the Doppler-shifted frequency

$$\omega_D = \omega - k_{\parallel} v_{\parallel} \quad (7.22)$$

(as observed by a particle moving with parallel velocity $v_{\parallel}$) is some integer multiple s of the particle Larmor frequency $s \omega_B^{(k)}$:

$$\omega_D = s \omega_B^{(k)}.$$

(7.23)

If a wave is, in general, oblique, its electric field has components transversal and parallel to $\mathbf{B}^{(0)}$, whereas if the wave is parallel, its electric field is transversal. Since the transversal field typically consists of left- and right-hand polarized components, the integer s may be either positive or negative. Anyway

the energy gain is severely limited due to the particle losing resonance with the wave.

Large gains of energy are possible, in principle, if a spectrum of waves is present. In this case, the resonant interaction of a particle with one wave can result in an energy change that brings this particle into resonance with a neighboring wave, which then changes the energy so as to allow the particle to resonate with another wave, and so on. Such an energy change can be diffusive, but over long time scales there is a net gain of energy, resulting in *stochastic* acceleration.

A traditional problem of the process under discussion is the so-called *injection* energy. The problem arises since for many waves in plasma their phase velocity along the ambient magnetic field, $\omega/k_{\parallel}$, is much greater than the mean thermal velocity of particles. Let us rewrite the gyroresonance condition (7.23) as

$$\gamma_L \left(\frac{\omega}{k_{\parallel}} - v_{\parallel} \right) = \frac{s \omega_B^{(k)}}{k_{\parallel}}. \quad (7.24)$$

Here the relativistic Lorentz factor γ_L has been taken into account (see Exercise 7.3). Consider two opposite cases.

- (a) For low thermal velocities we can neglect $v_{\parallel}$ in (7.24) and see that, in order to resonate with a thermal particle, the waves must have very high frequencies $\omega \approx \omega_B^{(k)}$ or very small $k_{\parallel}$.

For the case of thermal electrons and protons in the solar corona, their Larmor frequencies are very high (Exercise 5.1). If we try to choose a minimal value of $k_{\parallel}$, we are strongly restricted by a maximal value of wavelength λ ,

which must be certainly smaller than the maximal size of an acceleration region, L . These difficulties naturally lead to much doubt about the viability of stochastic acceleration and to a search for *preacceleration* mechanisms.

- (b) On the other hand, high energy particles need, according to the resonance condition (7.24), waves with very low frequencies: $\omega \ll \omega_B^{(k)}$. Therefore

a very *broad-band* spectrum of waves (extending from very high $\approx \omega_B^{(k)}$ to very low frequencies) is necessary to accelerate particles from thermal to relativistic energies in astrophysical plasma.

In principle, the so-called *wave cascading* from low to high frequencies can be a way of producing the necessary broad-band spectrum. The idea comes from the theory of hydrodynamic turbulence (Kolmogorov 1941). Here the evolution of turbulence can be described by the *Kolmogorov dimensional analysis* or by a *diffusion of energy in wavenumber space*. The last idea was subsequently introduced to MHD by Zhou and Matthaeus (1990). They presented a general transfer equation for the wave spectral density. In Sect. 7.2.2, we shall discuss briefly both approaches and their applications; see also Goldreich and Sridhar (1997).

The stochastic acceleration of particles by waves is essentially the resonant form of Fermi acceleration (see Sect. 6.2(c)). An important feature of stochastic acceleration is an isotropization process because

the pitch-angle scattering increases the volume of wave phase space that can be sampled by the resonant particles (7.24).

In general, if isotropization exists and keeps the distribution isotropic during an acceleration time, it increases the acceleration efficiency in astrophysical plasma. For example, Alfvén (1949) considered the betatron acceleration in an uniform magnetic field $\mathbf{B}^{(0)}(t)$ which changes periodically in time and has local nonuniformities $\mathbf{B}^{(1)}$ characterized by significant variations at distances smaller than the Larmor radius of accelerated particles.


When a particle passes through such nonuniformities its motion becomes random, with the momenta tending to be uniformly distributed between the three degrees of freedom. For this reason, when the field $\mathbf{B}^{(0)}(t)$ contracts, a fraction of the energy acquired due to betatron acceleration is transferred to the parallel component of the particle motion. As a consequence, the decrease in the energy of the transverse motion with decreasing magnetic field is smaller than its increase in the growth time. Thus the particle acquires an additional energy on completion of the full cycle. Therefore

the total particle energy can systematically increase even if the fluctuating magnetic field does not grow.

This phenomenon is known as the *Alfvén pumping*.

Tverskoy (1969, 1968) showed that in a turbulent astrophysical plasma, the Fermi acceleration related to the reflection from long strong waves is efficient only in the presence of fast particle scattering by short waves whose length is comparable to the particle Larmor radius.

Fig. 7.3 The energy per unit wave number in Kolmogorov's turbulence is plotted as a function of wavenumber in the inertial range *B* between the source *A* at small *k* and the sink *C* at large *k*



7.2.2 The Kolmogorov Theory of Turbulence

In general terms, a hydrodynamic flow tends to become turbulent if the ratio of inertial to viscous terms in the equation of motion, as described by the Reynolds number (see Chap. 12), is sufficiently large. In order not to obscure the essential physical point made in this section, we assume that a turbulence is *isotropic* and *homogeneous*. So we can define a one-dimensional spectral density $W(k)$, which is the wave energy density per unit volume in the wave vector space $\mathbf{k}$.

First, we remind the Kolmogorov (1941) treatment of *stationary* turbulence of *incompressible* fluid. The steady state assumption implies that the energy flux F through a sphere of radius k is independent of time. In the *inertial* range of wave numbers, for which supply and dissipation of energy are neglected, the flux F is also independent of the wave vector k . If $\mathcal{P}$ denotes the total rate of energy dissipation at the short wave ($k = k_{\max}$) edge of the inertial range, which equals the rate of energy supply at the long wave ($k = k_{\min}$) edge, then $F = \mathcal{P}$ and $dF/dk = 0$ in the inertial range in Fig. 7.3.

Kolmogorov's theory adopts the hypothesis that with the above assumptions the flux F through a sphere of radius k in the *inertial* range depends only upon the energy in that sphere and upon the wave number. Thus by dimensional analysis we arrive at

$$F = \mathcal{P} \sim W^{3/2} k^{5/2}. \quad (7.25)$$

From here it follows that the one-dimensional spectral density

$$W(k) = C_k \mathcal{P}^{2/3} k^{-5/3}.$$

(7.26)

This is the famous *Kolmogorov spectrum* for the hydrodynamic isotropic turbulence, involving the Kolmogorov constant C_k .

The turbulent velocity field in fluid can be thought of as being made of many eddies of different sizes. The input energy is usually fed into the system in a way to produce the largest eddies. Kolmogorov had realized that these large eddies can feed energy to the smaller eddies and these in turn feed the still smaller eddies, resulting in a cascade of energy from the larger eddies to the smaller ones.

If we anticipate the viscosity ν (see Sect. 12.2.3) to be not important for this process, we neglect dissipation of energy. However we cannot have eddies of indefinitely small size. For sufficiently small eddies of size $l_{\min}$ and velocity $v_{\min}$, the Reynolds number is of order unity, i.e.

$$l_{\min} v_{\min} \sim \nu. \quad (7.27)$$

So the energy in these small eddies is dissipated by viscosity.

Let the energy be fed into the turbulence at some rate $\mathcal{P}$ per unit mass per unit time at the largest eddies of size $l_{\max}$ and velocity $v_{\max}$, for which the Reynolds number

$$\text{Re} = \frac{l_{\max} v_{\max}}{\nu} \gg 1. \quad (7.28)$$

Then this energy cascades to smaller and smaller eddies until it reaches the smallest eddies satisfying condition (7.27).

The intermediate eddies merely transmit the energy to the smaller eddies. Let characterize these intermediate eddies only by their size l and velocity v . Since they are able to transmit the energy at the required rate $\mathcal{P}$, Kolmogorov postulated that it must be possible to express $\mathcal{P}$ in terms of l and v . On dimensional grounds, there is only one way of writing $\mathcal{P}$ in terms of l and v :

$$\mathcal{P} \sim \frac{v^3}{l}. \quad (7.29)$$

From here

$$v \sim (\mathcal{P}l)^{1/3}. \quad (7.30)$$

So

the velocity associated with the turbulent eddies of a particular size is proportional to the cube root of this size.

This result is known as the *Kolmogorov scaling law*. The scaling law (7.30) expresses the same thing as the Kolmogorov spectrum (7.26). This is shown in Exercise 7.10.

The Kolmogorov scaling law (7.30) was verified by doing experiments on a turbulent fluid with a sufficiently large inertial range. In laboratory it is very difficult to reach high enough Reynolds numbers to produce a sufficiently broad inertial range. One of the first confirmations of it was reported by Grant et al. (1962) by conducting experiments in a tidal channel between Vancouver Island and mainland Canada (see also Stewart and Grant 1969).

The Kolmogorov power spectrum (7.26) is observed in the turbulent *boundary layer* on the ground and in some other turbulent flows in astrophysical plasma (for example, in the *solar wind*), in spite of the fact that, in all these cases, the original assumptions of incompressibility and isotropy are not fulfilled.

7.2.3 MHD Turbulent Cascading

The Kolmogorov concept of independence of widely separated wave numbers in the inertial range of hydrodynamic turbulence was modified for the MHD case by Iroshnikov (1964) and Kraichnan (1965). When the magnetic energy in subinertial wave numbers exceeds the total energy in the inertial range, the predicted inertial range spectrum is proportional to $k^{-3/2}$, instead of $k^{-5/3}$. Note that the Kolmogorov spectrum is steeper than the Kraichnan spectrum ($5/3 > 3/2$).

Leith (1967) introduced a *diffusion* approximation for spectral transfer of energy in isotropic hydrodynamic turbulence. This approach may be viewed as an alternative to the straight-forward dimensional analysis discussed above. However it is a natural extension since this approach approximates the spectral transfer as a *local* process in wave number space, i.e. in accordance with the spirit of the Kolmogorov hypotheses that the total energy is conserved with respect to couplings between waves. Therefore

just diffusion of the wave energy density $W(\mathbf{k})$ in the wave number space $\mathbf{k}$ is a physically appealing framework for the simplest model to describe this kind of *local conservative* transfer.

If some waves, propagating parallel to the uniform field $\mathbf{B}^{(0)}$, are injected at the longest wavelength $\lambda = \lambda_{\max}$ and if a Kolmogorov-like nonlinear cascade transfers the wave energy to smaller scales, then the diffusion equation in wave number space

$$\frac{\partial W}{\partial t} = \frac{\partial}{\partial k_{\parallel}} \left(D_{\parallel\parallel} \frac{\partial W}{\partial k_{\parallel}} \right) - \gamma(k_{\parallel}) W + S \quad (7.31)$$

can describe injection, cascading, and damping of the waves. Here $D_{\parallel\parallel}$ is a diffusion coefficient that depends on W and can be determined for Kolmogorov-type cascading. $\gamma(k_{\parallel})$ is the damping rate usually due to particle acceleration in high-temperature low-density astrophysical plasma. The wave energy is dissipated by accelerating particles in smallest scales $\lambda \sim \lambda_{\min}$.

The source term S in (7.31) is proportional to the injection rate Q of the wave energy. A mechanism by which the waves are generated is typically unknown but easily postulated. For example, the MHD waves can be formed by a large-scale restructuring of the magnetic field in astrophysical plasma, which presumably occurs in non-stationary phenomena with flare-like energy releases due to *magnetic reconnection* (see Part II).

In summary,

the wave cascading and particle acceleration are described by one wave-diffusion equation, in which the damping depends on the accelerating particle spectra, and by diffusion equations (one for each kind k of particles: electrons and ions) for accelerating particles.

The system is therefore highly coupled and generally nonlinear or quasilinear in the case of small-amplitude waves.

7.3 The Relativistic Electron-Positron Plasma

According to present views, in a number of astrophysical objects there is a relativistic plasma that mainly consists of electrons and positrons. Among these objects are pulsar magnetospheres (Ruderman and Sutherland 1975; Michel 1991; recent successes in this field are reviewed by Becker 2009), accretion disks in close binary systems (Takahara and Kusunose 1985; Rose 1998), relativistic jets from active galactic nuclei (Begelman et al. 1984; Peacock 1999), and magnetospheres of rotating black holes in active galactic nuclei (Hirotani and Okamoto 1998).

Because of synchrotron losses, the relativistic collisionless plasma in a strong magnetic field should be strongly *anisotropic*: its particle momenta should have a virtually one-dimensional distribution distended along the field. The transversal (with respect to the field) momentum of a particle is small compared with the longitudinal momentum. In accordance with Ruderman and Sutherland (1975), such a particle distribution is formed near the pulsar surface under the action of a strong longitudinal electric field and synchrotron radiation. What equations can be used as starting ones for a description of the electron-positron plasma?—The answer depends upon a property of the plasma, which we would like to describe.

It is known that the anisotropy can result in various types of instabilities, for example, the *fire-hose instability* of the relativistic electron-positron plasma (Mikhailovskii 1979). Behavior of Alfvén waves in the isotropic and anisotropic plasmas can be essentially different (Mikhailovskii et al. 1985).

We suppose that the anisotropic relativistic approach of a type of the CGL approximation (Sect. 11.5) can be used to consider the problem of Alfvén waves of finite amplitude. However the dispersion effects are important for such waves and are not taken into account in the CGL approximation. The problem can be analyzed on the basis of the standard kinetic approach with use of the Vlasov equation (Sect. 3.1.2). As we saw above, such a procedure is sufficiently effective in the case of linear problems but is complicated in study of nonlinear processes when one must deal with parts of the distribution function square and cubic to the wave amplitude.

More effective kinetic approaches are demonstrated in Mikhailovskii et al. (1985). One of them is based on expansion in the series of the inverse power of the background magnetic field (Sect. 5.2) and allowance for the cyclotron effects as a small corrections. Using this approach, Mikhailovskii et al. consider the nonlinear

Alfvén waves both in the case of an almost one-dimensional momentum particle distribution (the case of a pulsar plasma) and in the case of an isotropic plasma. The later case is interesting, in particular, for the reason that it has been also analyzed by means of the MHD equations (Sect. 20.1.4). Two types of Alfvén solitons (the moving-wave type and the nonlinear wave-packet type) can exist in relativistic collisionless electron-positron plasma.

Magnetic reconnection in a collisionless relativistic electron-positron plasma is considered as a mechanism of electron and lepton acceleration in large-scale extragalactic jets, pulsar outflows like the Crab nebula and core regions of active galactic nuclei (AGN) as the respective jet origin (see Larrabee et al. 2003).

In the electron-ion plasma, mass asymmetry produces at least two inertial scales. Magnetic reconnection proceeds on ion scales, and provides the framework with respect to flow conditions for fast reconnection on the electron demagnetization scale. In the relativistic electron-positron plasma, the reconnection rates observed in numerical simulations (e.g., Zenitani and Hoshino 2001, 2007; Jaroschek et al. 2004), correspond to rather vacuum type reconnection which is only limited by electron inertia.

Recommended Reading: Lifshitz and Pitaevskii (1981), Chaps. 3 and 6.

7.4 Practice: Exercises and Answers

Exercise 7.1 (Sect. 7.1.1). Write the general solution of the linear equation (7.2).

Answer. Since the left-hand side of (7.2) is the time derivative (more exactly, the Liouville operator (1.12) acting on the first-order distribution function for particles following the *unperturbed* trajectories), the solution of (7.2) is formally the integral over time

$$f_k^{(1)}(\mathbf{r}, \mathbf{v}, t) = -\frac{e_k}{m_k} \int_{-\infty}^t \left(\mathbf{E}^{(1)} + \frac{1}{c} \mathbf{v} \times \mathbf{B}^{(1)} \right)_{\alpha} \frac{\partial f_k^{(0)}(\mathbf{r}, \mathbf{v}, \tau)}{\partial v_{\alpha}} d\tau. \quad (7.32)$$

Here the integration follows an unperturbed-particle trajectory to the point $(\mathbf{r}, \mathbf{v})$ in phase space X .


In principle, substitution of (7.32) into the Poisson law for electrostatic waves gives a perturbation of electric charge density (3.5). Similarly, one can determine a perturbation of current density (3.6) by substitution of (7.32) into the Ampère law in the case of electromagnetic waves. In practice, however, solving (7.32) is fairly complicated.

Exercise 7.2 (Sect. 7.1.2). Show that, for a particle with velocity $\mathbf{v}$ in a plasma without magnetic field, the resonance condition corresponds to:

$$\omega - \mathbf{k} \cdot \mathbf{v} = 0. \quad (7.33)$$

This is usually called the *Cherenkov condition*.

Fig. 7.4 The wave-vector cone of the Cherenkov emission



Exercise 7.3 (Sects. 7.1.2, 7.2.1). Consider a wave that has frequency ω and wave vector $\mathbf{k}$ in the laboratory frame. Show that in the rest frame of a relativistic particle the frequency of the wave is

$$\omega_0 = \gamma_L (\omega - \mathbf{k} \cdot \mathbf{v}), \quad (7.34)$$

where

$$\gamma_L = \left(1 - \frac{v^2}{c^2}\right)^{-1/2} \quad (7.35)$$

is the Lorentz factor of the particle. Therefore the Cherenkov resonance condition (7.33) corresponds to $\omega_0 = 0$, which means that the fields appear static in the rest frame of the particle.

Answer. Apply the Lorentz transformation to the four-vector $\{\mathbf{k}, i\omega/c\}$ (see Landau and Lifshitz (1975), Chap. 6, Sect. 48).

Exercise 7.4 (Sect. 7.1.2). In a transparent medium with a refraction index n , greater than unity, the Cherenkov condition (7.33) can be satisfied for fast particles with

$$\beta = \frac{v}{c} \geq \frac{1}{n}. \quad (7.36)$$

Let χ be the angle between the particle's velocity $\mathbf{v}$ and the wave vector $\mathbf{k}$ of appearing emission which is called *Cherenkov emission* (Cherenkov 1934, 1937).


As we know, a charged particle must move non-uniformly to radiate in vacuum. As an example we may recall the formula (5.66) for dipole emission. In a medium, however, condition (7.36) allows the uniformly moving particle to radiate.

Show that Cherenkov emission is confined to the surface of a cone with the cone half-angle (as shown in Fig. 7.4)

$$\chi = \arccos \frac{1}{n}. \quad (7.37)$$

Radiation with wave vectors along the conic surface (7.37) is generated as a result of the Cherenkov emission. Discuss an analogy between the Cherenkov emission pattern and the bow wave of a ship or a supersonic aircraft.

Fig. 7.5 (a) The electric field potential in a Langmuir wave of a small but finite amplitude. (b) The phase trajectories of an electron in the wave



Exercise 7.5 (Sect. 7.1.2). Consider the one-dimensional motion of an electron in the electric field of a Langmuir wave of a small but finite amplitude.

Answer. Let the electric field potential of the wave be of the form

$$\varphi = \varphi_0 \cos(\omega_{pl}^{(e)} t - kx). \quad (7.38)$$

In the reference frame moving with the wave (see Sect. 10.2.2), the field is static:

$$\varphi = \varphi_0 \cos kx. \quad (7.39)$$

This potential is shown in Fig. 7.5a.

For an electron having a small velocity near $x = 0$, we have the following equation of motion:

$$m_e \ddot{x} = e \frac{\partial \varphi}{\partial x} = -e\varphi_0 k \sin kx \approx -e\varphi_0 k^2 x. \quad (7.40)$$

So such a trapped electron is oscillating with frequency

$$\omega_{tr}^{(e)} = k \left(\frac{e\varphi_0}{m_e} \right)^{1/2}. \quad (7.41)$$

This is illustrated by particle trajectories in the two-dimensional phase space (Fig. 7.5b).

The potential energy $-e\varphi$ of the trapped electron has maximum at the minimum of the potential φ , at points M which determine the separatrix S .

Exercise 7.6 (Sect. 7.1.2). Consider the Landau resonance for electrons in a Maxwellian plasma. It is clear that electrons moving much slower or much faster than the wave tend to see the electric field that averages to zero. So we have to consider only the particles in some small part of velocity space close to the phase velocity as shown in Fig. 7.1.

Since the slope of the initial distribution function is negative, there are more electrons at lower velocity than at higher velocity near the resonance (7.12). Estimate a difference.

Exercise 7.7 (Sect. 7.1.2). Show that the Landau damping prevents plasma waves from escaping the region where $\omega = \omega_{pl}^{(e)}$ (see definition (8.83)) into rarefied plasma, for example, from the solar corona to the interplanetary space.

Hint. Consider the dispersion equation for the Langmuir waves in a homogeneous equilibrium plasma without a magnetic field (Sect. 10.2.2). Show that the phase velocity V_{ph} decreases with decreasing the electron plasma frequency $\omega_{pl}^{(e)}$ (see formula (10.10)). Therefore the Langmuir waves are absorbed in the regions where V_{ph} falls to the values comparable to the mean thermal velocity (e.g., Zheleznyakov 1996).

Exercise 7.8 (Sect. 7.1.2). In the fire-hose instability, the driving force is the beam pressure parallel to the magnetic field. Show that this pressure increases the amplitude of an electromagnetic transverse wave in a way analogous to that of a water flowing through a hose.

Hint. Consider low-frequency transverse waves in a homogeneous equilibrium plasma with a magnetic field. Such waves are called the *kinetic* Alfvén waves. They extend to frequencies higher than that are valid for magnetohydrodynamics (MHD). Let a beam of protons or electrons travel parallel to the magnetic field. An analysis of linear disturbances similar to the MHD waves will introduce an additional term into the dispersion equation of the Alfvén wave. Note that an instability occurs for beams of protons or electrons. Consider the threshold condition in both cases.

Exercise 7.9 (Sect. 7.1.3). Show that fast ions can generate *whistler* waves when the resonant particles are travelling faster than the wave. Show that, in this case, the effect of Doppler shift is to change the sense of rotation of the wave electric field in the resonant-particle frame from right-handed to left-handed.

Exercise 7.10 (Sect. 7.2.2). Show that the Kolmogorov spectrum formula (7.26) follows from the Kolmogorov scaling law (7.30).

Answer. The kinetic energy density associated with some wave number k is $W(k) dk$, which can be roughly written as

$$W(k) k \sim v^2. \quad (7.42)$$

Substituting for v from formula (7.30) with $l \sim 1/k$, we have

$$W(k) k \sim \mathcal{P}^{2/3} k^{-2/3}. \quad (7.43)$$

From here the Kolmogorov spectrum (7.26) readily follows.

Chapter 8

Coulomb Collisions in Astrophysical Plasma

Abstract Binary collisions of particles with the Coulomb potential of interaction are typical for physics of collisional plasmas in space and especially for gravitational systems. Coulomb collisions of fast particles with plasma particles determine momentum and energy losses of fast particles, the relaxation processes in astrophysical plasma.

8.1 Close and Distant Collisions

8.1.1 The Collision Parameters

Binary interactions of particles, described by the Coulomb potential

$$\varphi(r) = \frac{e}{r}, \quad (8.1)$$

have been studied in mechanics (see [Landau and Lifshitz \(1976\)](#), Chap. 4, Sect. 19). Considering binary interactions as *collisions*, we are interested only in their final result, the duration of the interaction and the actual form of particle trajectories being neglected. Thus in the centre-of-mass system, each particle is deflected through an angle χ defined by the relation

$$\tan \frac{\chi}{2} = \frac{e_1 e_2}{m v^2 l} \quad (8.2)$$

or

$$l(\chi) = \frac{e_1 e_2}{m v^2} \cot \frac{\chi}{2}. \quad (8.3)$$

Here

$$m = \frac{m_1 m_2}{m_1 + m_2} \quad (8.4)$$

Fig. 8.1 The trajectory of a light particle with mass m_1 near a heavy particle with mass m_2





Fig. 8.2 Close (a) and distant (b) collisions of particles in the momentum space in the centre-of-mass system

is the reduced mass, v is the relative particle velocity at infinity, l is the “*impact parameter*”. The last is the closest distance of the particle’s approach, were it not for their interaction as shown in Fig. 8.1.

For particles deflected through a right angle

$$l\left(\frac{\pi}{2}\right) \equiv l_{\perp} = \frac{e_1 e_2}{mv^2}, \quad (8.5)$$

so the initial formula (8.2) is conveniently rewritten as

$$\tan \frac{\chi}{2} = \frac{l_{\perp}}{l}.$$

(8.6)

The collisions are called *close* if

$$\pi/2 \leq \chi \leq \pi, \quad \text{i.e.} \quad 0 \leq l \leq l_{\perp}. \quad (8.7)$$

Correspondingly, for *distant* collisions $l > l_{\perp}$ and $0 \leq \chi < \pi/2$. Both cases are shown in Fig. 8.2.

8.1.2 The Rutherford Formula

The average characteristics of the Coulomb collisions are obtained with the aid of the formula for the *differential* cross-section. It is called the Rutherford formula and is derived from (8.3) as follows (see Fig. 8.1):

$$\begin{aligned} d\sigma &= 2\pi l(\chi) dl = 2\pi l(\chi) \left| \frac{dl}{d\chi} \right| d\chi = \\ &= \frac{\pi e_1^2 e_2^2}{m^2 v^4} \frac{\cos(\chi/2)}{\sin^3(\chi/2)} d\chi = \left(\frac{e_1 e_2}{2mv^2} \right)^2 \frac{d\Omega}{\sin^4(\chi/2)}. \end{aligned} \quad (8.8)$$

Here the modulus bars indicate the absolute value of the derivative $dl/d\chi$ because it has a negative sign: with increase of the impact parameter l , the scattering angle χ decreases; the solid angle $d\Omega = 2\pi \sin \chi d\chi$.

By integrating (8.8) over the back hemisphere (8.7), we find the *total* cross-section of close collisions

$$\sigma_{cl} = \frac{\pi e_1^2 e_2^2}{m^2 v^4} = \pi l_{\perp}^2. \quad (8.9)$$

This formula follows directly from definition (8.5), of course, without integrating the differential cross-section (8.8).

8.1.3 The Test Particle Concept

By analogy with the usual gas, the concept of a “test” particle is introduced to analyze the collisions in plasma. For instance the frequency of test particle (m_1, e_1) collisions with “field” particles (m_2, e_2) is introduced:

$$\nu_{cl} = n_2 v_1 \sigma_{cl} = \frac{\pi e_1^2 e_2^2 n_2}{m_1^2 v_1^3}. \quad (8.10)$$

Here, for simplicity’s sake, it is assumed that $m_2 \gg m_1 \approx m$ (see formula (8.4)) and $v_2 \ll v_1$. So this is, for example, the case of an electron colliding with “cold” ions.

The length of *mean free path* λ of a test particle in a gas consisting of field particles is, by definition, the distance along which the particle suffers one collision,

$$\lambda = v_1 \nu^{-1}. \quad (8.11)$$

From (8.10) and (8.11) it follows for *close* collisions that

$$\lambda_{cl} = \frac{1}{n_2 \sigma_{cl}} \sim \frac{1}{n_2} v_1^4. \quad (8.12)$$

Hence the time between two consecutive collisions is

$$\tau_{cl} = \frac{\lambda_{cl}}{v_1} = \frac{m_1^2 v_1^3}{\pi e_1^2 e_2^2 n_2} \sim \frac{1}{n_2} v_1^3, \quad (8.13)$$

or the frequency of close collisions

$$\nu_{cl} = \frac{1}{\tau_{cl}} = \frac{\pi e_1^2 e_2^2 n_2}{m_1^2 v_1^3} \sim n_2 \frac{1}{v_1^3}, \quad (8.14)$$

which is the same as formula (8.10) of course. So, it is natural that

the higher the velocity v_1 of a test particle, the longer are its mean free path and collisional time

at least under the assumptions made above.

8.1.4 Particles in a Magnetic Trap

Formulae (8.10) and (8.13) are frequently used in order to find out what approximation we have to use to consider the astrophysical plasma under definite physical conditions. For example, if the length of mean free path λ of a test particles inside a magnetic trap (Sect. 6.2) is greater than the trap's size, then such particles can be considered in the collisionless approximation. Here *charge separation* may be found to be essential, as well as the electric field resulting from it (Alfvén and Fälthammar 1963; Persson 1963).

While the magnetic mirror is the primary trapping mechanism, the electrostatic potential also traps electrons

with energies low to overcome the electrostatic potential.

In the solar atmosphere, the electrostatic potential produced, in solar-flare magnetic traps, has an energy equivalent of the average energy of accelerated electrons. The number and energy fluxes of the electrons that escape from the trap can be reduced by as much as ~ 50 or more depending on the magnetic mirror ratio of the flare loop and the ratio of the ion and electron anisotropy factors (Spicer and Emslie 1988).

Some other effects due to non-collisional particles in the so-called *collapsing* magnetic traps are mentioned in Sect. 6.2; they will be considered in Sect. 18.3 and Part II, Chap. 9. For example,

the electric potential mentioned above increases the efficiency of confinement and acceleration of electrons in solar flares

(Kovalev and Somov 2002).

On the other hand, if the length of the mean free path of the test particles is much less than the trap's size, the collisions play an important role. As a rule they maxwellise the plasma (the gas of test particles), making it an equilibrium

one. In such a plasma the notion of *temperature* is meaningful, as we shall see in Chap. 9. For example, while considering thermal electrons (having the density n_e and the temperature T_e) in the trap, an electron with the *mean thermal velocity* (see definition (5.53))

$$V_{Te} = \sqrt{\frac{3k_B T_e}{m_e}} \quad (8.15)$$

should be taken as the test particle. Then we obtain the well-known “ T to the 3/2 power” law for the time of the Coulomb collisions (8.13):

$$\tau \sim \frac{T_e^{3/2}}{n_e}. \quad (8.16)$$

■ The hotter the astrophysical plasma is, the more collisionless is it with respect to some physical phenomenon or another.

In general the characteristic time τ of the Coulomb collisions has to be compared with the characteristic times of other physical processes: the time of particle motion between magnetic corks in the trap, the period of the Larmor rotation, the time of heating or cooling, etc. Then we can conclude what is the most important physical effect and what is the best approximation to describe this effect in astrophysical plasma. As for the collapsing magnetic traps in solar flares, the intermediate case of *rare-collisional plasma* can be easily and often released (Bogachev and Somov 2009).

8.1.5 The Role of Distant Collisions

Because for small angles χ the differential cross-section (8.8) is

$$d\sigma \sim \frac{d\chi}{\chi^3}, \quad (8.17)$$

the total cross-section diverges.

■ Such divergence of the collisional cross-section always occurs, once the interaction potential has no restricting factor,

or, to put the same in another way, if the interaction forces do not break off at some distance, as in the case of hard balls. This fact is of fundamental importance, for example, in *stellar dynamics* (Jeans 1929; Chandrasekhar 1943a) or, more exactly, in any astrophysical system governed by gravitational force (say a gravitational system), see Sects. 3.3 and 9.8.

Although each distant collision causes only a small deflection of the test particle trajectory, they are present in such large numbers that their total action upon the particle is *greater* or much greater than that of relatively rare close collisions. Let us convince ourselves that this is true.

Each distant collision causes a small change in momentum perpendicular to the initial direction of the particle's motion:

$$\delta p_{\perp} = p \sin \chi = m_1 v_1 \frac{2 \tan(\chi/2)}{1 + \tan^2(\chi/2)} = \frac{2 m_1 v_1 (l_{\perp}/l)}{1 + (l_{\perp}/l)^2} = 2 m_1 v_1 \frac{x}{1 + x^2}.$$

Here $x = l_{\perp}/l$, and $0 \leq x \leq 1$.

Since distant collisions occur chaotically, we are usually interested in the mean rate of change in the quantity $p_{\perp}^2$:

$$\begin{aligned} \frac{d}{dt} p_{\perp}^2 &= \int_{\chi=\pi/2}^{\chi=0} (\delta p_{\perp})^2 n_2 v_1 d\sigma = \\ &= 8\pi n_2 m_1^2 v_1^3 l_{\perp}^2 \int_1^0 \frac{dx}{(1+x^2)^2 x} \sim \ln x \Big|_1^0. \end{aligned} \quad (8.18)$$

The integral diverges logarithmically on the upper limit. Let us restrict it to some maximal value of the impact parameter

$$\Lambda = l_{\max}/l_{\perp}. \quad (8.19)$$

Then the integral is approximately equal to

$$\frac{d}{dt} p_{\perp}^2 = 8\pi n_2 m_1^2 v_1^3 l_{\perp}^2 \ln \Lambda = \pi e_1^2 e_2^2 \frac{n_2}{v_1} 8 \ln \Lambda. \quad (8.20)$$

The factor $\ln \Lambda$ is referred to as the Coulomb logarithm.

Introduce the characteristic time $\tau_{\perp}$ during which the perpendicular component of the momentum acquires a value equal to the initial momentum $m_1 v_1$:

$$\tau_{\perp} = (m_1 v_1)^2 \left(\frac{d}{dt} p_{\perp}^2 \right)^{-1} = \frac{m_1^2 v_1^3}{\pi e_1^2 e_2^2 n_2 (8 \ln \Lambda)}. \quad (8.21)$$

In other words, the mean resulting deflection becomes comparable with the quantity $\pi/2$ in a time $\tau_{\perp}$. Recall that this deflection through a large angle is a result of many distant collisions.

The effective frequency of distant collisions that corresponds to the time $\tau_{\perp}$ is

$$\nu_{\perp} = \frac{1}{\tau_{\perp}} = \frac{\pi e_1^2 e_2^2 n_2}{m_1^2 v_1^3} 8 \ln \Lambda, \quad (8.22)$$

which is $8 \ln \Lambda$ larger than the close collisions frequency (8.14):

$$\nu_{\perp} = 8 \ln \Lambda \cdot \nu_{cl}.$$

(8.23)

The factor $8 \ln \Lambda$ is usually much greater than unity; its typical value is $\gtrsim 10^2$ under physical definition of $\ln \Lambda$ given in Sect. 8.2.

The influence of the close Coulomb collisions on kinetic processes in astrophysical plasma is, as a rule, negligibly small in comparison to the action of distant collisions.

For example, the distant collisions determine an evolution of the distribution function of fast electrons injected into the thermal plasma in the solar atmosphere during solar flares. However *this does not mean that the close collisions do never play any role* in plasma astrophysics. Just in the same example, the close collisions of fast electrons with thermal ions create the collisional *bremsstrahlung* emission in the energy range $\gtrsim 10$ keV, because the close collisions are responsible for large exchange of the particle momentum. For typical flare parameters ($h\nu \approx 20$ keV, $\ln \Lambda \approx 20$) the efficiency of the *bremsstrahlung* process is $\sim 3 \times 10^{-6}$ (Brown 1971; Korchak 1971).

8.2 Debye-Hückel Shielding

8.2.1 Simple Illustrations of the Shielding Effect

While considering the distant collisions, we have removed the divergence of the integral (8.18) which describes the mean rate of change of the test particle transversal momentum, purely formally—by artificially restricting the radius of action of the Coulomb forces at some maximal distance $l_{\max}$. Meanwhile this maximal distance may be chosen quite justifiably, based on the following reasoning. In a plasma,

each charged particle attracts oppositely charged particles and, at the same time, repels the particles of the same charge.

As a consequence, the oppositely charged particles tend to gather around a test unitary charge, thus weakening its Coulomb field. As a result of such “*shielding*” the action of the electric field extends over a distance no greater than some length r_{DH} called the *Debye-Hückel radius*.

The concept of *Debye-Hückel shielding* (Debye and Hückel 1923) has a clear meaning. Let us assume that a plasma contains an immovable charge which then creates the electrostatic field in its vicinity. After a sufficiently long time, electrons

and ions rearrange themselves in response to the forces acting on them. For example, if the test charge is positive, the electron density near it increases. As a final result of shielding interactions, some new *equilibrium* distribution of *two components*: positive and negative plasma particles is established in this field. Its electrostatic potential φ is related to the densities of ions n_i and electrons n_e via the Poisson equation

$$\Delta\varphi = -4\pi e (Zn_i - n_e), \quad (8.24)$$

where Ze is the ion charge.

In the thermodynamic equilibrium state the ion and electron densities in the electrostatic field with potential $\varphi(r)$ are to be distributed according to Boltzmann's law

$$n_i = n_i^0 \exp\left(-\frac{Ze\varphi}{k_B T_i}\right), \quad n_e = n_e^0 \exp\left(\frac{e\varphi}{k_B T_e}\right). \quad (8.25)$$

The constant coefficients are set equal to the mean densities n_i^0 and n_e^0 of plasma particles, since $\varphi \rightarrow 0$ far from the test particle considered.

Supposing that the Coulomb interaction is so weak that

$$Ze\varphi \ll k_B T_i \quad \text{and} \quad e\varphi \ll k_B T_e, \quad (8.26)$$

or restricting our consideration to the approximate solutions applicable at large distances from the shielded charge, we expand both exponents (8.25) in a series and substitute in (8.24). We obtain the following linearized equation:

$$\begin{aligned} \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\varphi}{dr} \right) &= -4\pi e \left[Zn_i^0 \left(1 - \frac{Ze\varphi}{k_B T_i} \right) - n_e^0 \left(1 + \frac{e\varphi}{k_B T_e} \right) \right] = \\ &= 4\pi e \left[(n_e^0 - Zn_i^0) + \frac{e}{k_B} \left((Zn_i^0) \frac{Z}{T_i} + (n_e^0) \frac{1}{T_e} \right) \varphi \right]. \end{aligned} \quad (8.27)$$

As usual the actual plasma is *quasi-neutral on average* (see Sect. 8.2.3); instead of this let us assume here (like in Sects. 3.2.2 and 3.2.3) that the plasma is *ideally neutral*:

$$Zn_i^0 = n_e^0. \quad (8.28)$$

Thus we have an equation

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\varphi}{dr} \right) = \frac{4\pi e^2 n_e^0}{k_B} \left(\frac{Z}{T_i} + \frac{1}{T_e} \right) \varphi = \frac{\varphi}{r_{DH}^2}. \quad (8.29)$$

On the right-hand side of (8.29) we have two terms for a two-component plasma. We divide them by φ , then

$$\frac{1}{r_{DH}^2} = \frac{1}{r_{DH}^{(i)2}} + \frac{1}{r_{DH}^{(e)2}} = \frac{4\pi e^2 n_e^0}{k_B T_e} \left(1 + Z \frac{T_e}{T_i} \right). \quad (8.30)$$

Therefore

$$r_{\text{DH}} = \left(\frac{k_{\text{B}}}{4\pi e^2 n_e^0} \frac{T_e T_i}{Z T_e + T_i} \right)^{1/2} \quad (8.31)$$

is known as the Debye-Hückel (D-H) radius, being first derived by [Debye and Hückel \(1923\)](#) in the theory of electrolytes.

The solution of (8.29) corresponding to the test charge Ze situated at the origin of the coordinates, $r = 0$, is the potential

$$\varphi = \frac{Ze}{r} \exp\left(-\frac{r}{r_{\text{DH}}}\right). \quad (8.32)$$

At distances greater than r_{DH} , the electrostatic interaction is exponentially small.

The D-H length is an effective range for collisions, the potential between charged particles being the *shielded* Coulomb potential (8.32) rather than the Coulomb one (8.1) which would apply in a vacuum.

That is why:

- (a) The binary correlation function (3.30) reproduces the shape of the shielded Coulomb potential (8.32),
- (b) The D-H radius r_{D} is substituted in the Coulomb logarithm (8.20) in place of l_{max} .

A formula that is simpler than (8.31) is frequently used for the D-H radius, namely

$$r_{\text{DH}}^{(\text{e})} = \left(\frac{k_{\text{B}} T}{4\pi e^2 n_e} \right)^{1/2}. \quad (8.33)$$

This variant of the formula for the D-H radius implies that the shielding is due to just the particles of one sign, more exactly, *electrons*, i.e. in the formulae (8.25) we have $T_i = 0$ (the approximation of *cold ions*) and $T_e = T$. This is the *electron* D-H radius. The corresponding formula for the Coulomb logarithm is

$$\ln \Lambda = \ln \frac{3}{2e^3} \left(\frac{k_{\text{B}}^3 T^3}{\pi n_e} \right)^{1/2}. \quad (8.34)$$

Its values typical of the solar atmosphere are around 20 (Exercise 8.1).

Formula (8.33) shows that

the electron D-H radius increases with an increase of temperature, since electrons with higher kinetic energy can withstand the attraction of the positive ion charge Ze up to larger distances.

The electron D-H radius decreases with an increase of plasma density n_0 , since a larger number of electrons and ions can be accommodated in shorter distances to screen the electric field of charge Ze .

8.2.2 *The Real Shielding Effect*

In the standard approach to the Debye-Hückel (D-H) shielding effect, reproduced in Sect. 8.2.1, two equations are needed: one coming from electromagnetism (the Poisson equation (8.24)), the other from statistical mechanics (the Boltzmann law (8.25)). After linearization of the Poisson equation, the electrostatic screened potential (8.32) has been found.

The assumptions made to derive the famous D-H relation are, among other things, that: (a) the induced perturbation is slow; only the electrostatic fields are present, while induced electromagnetic fields are negligible; (b) a temperature can always be defined; (c) the number of particles inside the D-H sphere is large (see Sect. 3.2.3) and therefore fluctuations are small; (d) although ions and electrons have random thermal motion, perturbations induced around the equilibrium, which are responsible for small spatial variations in the electrostatic potential, can be neglected.

Of course, the real situation in astrophysical plasmas can differ from that required by one or more of these assumptions. In stellar cores, ions and nuclei are embedded in an electronic environment made by mainly free electrons. Nuclear fusion reaction *cross-section and rates* are sensitive to the screening effect of the electron cloud around the reacting nuclei. Thus the use of the standard D-H potential may produce errors in the evaluation of nuclear fusion rates, for example, in the solar core and in other dense astrophysical plasmas.

By choosing the inverse of the D-H length, $1/r_{\text{DH}}$, as a fluctuating parameter, Quarati and Scarfone (2007) have modified the shielded electrostatic potential with respect to the standard D-H potential. They have obtained the results that differ from those calculated with the standard potential and that are useful in the interpretation of experimental results on atomic and nuclear rates in several astrophysical systems and processes.

8.2.3 *Charge Neutrality and Oscillations in Plasma*

The Debye-Hückel shielding length is fundamental to the nature of a plasma. That is why this important characteristic appears again and again in plasma astrophysics, starting from the binary correlation function (3.30).

The first point to note is that a plasma maintains *approximate charge neutrality* called *quasi-neutrality* (see Sects. 3.2.2 and 11.5.2). The reason for this is simply that

any significant imbalance of positive and negative charge could only be maintained by a huge electric field.

The movement of electrons to neutralize a charge inhomogeneity would be followed by an oscillatory motion (e.g., [Alfvén and Fälthammar 1963](#), see Chap. 4).

This brings us to a second characteristic of plasmas called the *plasma frequency* or, more exactly, the electron plasma frequency:

$$\omega_{pl}^{(e)} = \left(\frac{4\pi e^2 n_e}{m_e} \right)^{1/2}. \quad (8.35)$$

A charge density disturbance oscillates with this frequency (see Sect. 10.2.1). These oscillations are called *Langmuir waves* or *plasma waves*. Therefore, under most circumstances,

an actual plasma cannot sustain electric fields for lengths in excess of the Debye-Hückel radius r_{DH} or times in excess of a plasma period $T_{pl}^{(e)} = 2\pi/\omega_{pl}^{(e)}$.

However one cannot talk of plasma oscillations unless a large number of thermal particles are involved in the motion. It is the Debye-Hückel shielding length which determines the spatial range of the electric-field set up by the charge inequality:

$$r_{DH} = \frac{1}{\sqrt{3}} \frac{V_{Te}}{\omega_{pl}^{(e)}}. \quad (8.36)$$

Here V_{Te} is the mean thermal velocity of electrons. Therefore the Debye-Hückel length

$$r_{DH} \approx \frac{V_{Te}}{\omega_{pl}^{(e)}}. \quad (8.37)$$

So a fully-ionized plasma in the thermodynamic equilibrium is a quasi-neutral medium. The *space* and *time* scales of charge separation in such plasma are the Debye-Hückel radius and the inverse plasma frequency. Therefore the plasma oscillations are a typical example of *collective phenomena* (Sect. 3.2.3).

The Coulomb collisions, of course, damp the amplitude of the plasma oscillations with the rate which is proportional to the frequency ν_{ei} of electron-ion collisions (see Exercise 8.3).

8.3 Collisional Relaxations in Plasma

8.3.1 Some Exact Solutions

It was shown in Sect. 8.1 that, as a result of the Coulomb collisions, a particle deflects through an angle comparable with $\pi/2$ in a characteristic time given by formula (8.21). More exact calculations of the Coulomb collisions times, that take into account the thermal motion of field particles, have been carried out by Spitzer (1940) and Chandrasekhar (1943a,b,c). These calculations are cumbersome, so we give only their final results.

Let us consider the electron component of a plasma. Suppose that the test particles likewise are electrons moving with mean thermal velocity. Then the exact calculation gives instead of the formula (8.21) the time

$$\tau_{ee} = \frac{m_e^2 (3k_B T_e / m_e)^{3/2}}{\pi e_e^4 n_e (8 \ln \Lambda)} \cdot \frac{1}{0.714}. \quad (8.38)$$

This is called the time of mutual electron collisions or simply the *electron collisional time*. Comparison of formula (8.38) with (8.21) shows that the difference (the last factor in (8.38)) is not large. So the consideration of binary collisions in the approximation used in Sect. 8.1 is accurate enough, at least for astrophysical applications.

The analogous time of mutual collisions for ions, having mass m_i , charge e_i , temperature T_i and density n_i , is equal to

$$\tau_{ii} = \frac{m_i^2 (3k_B T_i / m_i)^{3/2}}{\pi e_i^4 n_i (8 \ln \Lambda)} \cdot \frac{1}{0.714}. \quad (8.39)$$

If a plasma is quasi-neutral: $e_i n_i \approx -e_e n_e = en$, where $e_i = -Ze_e$, and if $T_e \approx T_i$, then the ratio

$$\boxed{\frac{\tau_{ii}}{\tau_{ee}} \approx \left(\frac{m_i}{m_e} \right)^{1/2} \frac{1}{Z^3}}. \quad (8.40)$$

Coulomb collisions between thermal ions in plasma occur much more rarely than those between thermal electrons.

However it is not the time of collisions between ions τ_{ii} —the *ion collisional time*, but rather the time of electron-ion collisions that is the greatest. This characterizes, in particular, the process of temperature equalizing between the electron and ion components in a plasma. The rate of temperature equalizing can be determined from the equation

$$\frac{dT_e}{dt} = \frac{T_i - T_e}{\tau_{ei}(\mathcal{E})}, \quad (8.41)$$

where $\tau_{ei}(\mathcal{E})$ is the time of equilibrium establishment between the electron and ion plasma components. It characterizes the rate of exchange of energy $\mathcal{E}$ between the components and equals (Spitzer 1940, 1962; see also Sivukhin 1966, Sects. 9 and 17; cf. formulae (42.5) in Lifshitz and Pitaevskii 1981, Sect. 42)

$$\tau_{ei}(\mathcal{E}) = \frac{m_e m_i [3k_B (T_e/m_e + T_i/m_i)]^{3/2}}{e_e^2 e_i^2 (6\pi)^{1/2} (8 \ln \Lambda)}. \quad (8.42)$$

For comparison with formula (8.40) let us put $T_i = T_e$. Then

$$\tau_{ei}(\mathcal{E}) = 0.517 \frac{e_i^2}{e_e^2} \left(\frac{m_i}{m_e} \right)^{1/2} \tau_{ii}. \quad (8.43)$$

Thus the time of energy exchange between electrons and ions is much greater than the time of mutual ion collisions.

In a plasma consisting of electrons and protons with equal temperatures, we have

$$\tau_{ep}(\mathcal{E}) \approx 22 \tau_{pp} \approx 950 \tau_{ee}. \quad (8.44)$$

The energy exchange between electron and ion components occurs so slowly that for each component a distribution may be set up which is close to Maxwellian with the proper temperature.

That is the reason why we often deal with a *two-temperature* plasma. Moreover the so-called adiabatic model for two-temperature plasma (Sect. 8.3.4) is often used in astrophysics. However, at first, we illustrate the situation, discussed above, by two more accurate examples from the physics of solar flares.

8.3.2 Plasma Heating by Accelerated Electrons

Electrons and ions are rapidly accelerated to high energies during the so-called impulsive phase of solar flares. The electromagnetic radiation of non-relativistic electrons is routinely measured by hard X-ray (HXR) detectors in space and is interpreted in terms of the thick-target model as bremsstrahlung radiation (see Sect. 4.3.4). It appears that the accelerated 20–100 keV electrons contain a significant part of the total energy of a solar flare (Lin and Hudson 1971; Syrovatskii and Shmeleva 1972), indicating that they play a significant role in energetics of a flare.

In this section, we shall consider an impulsive heating of the solar atmosphere by a beam of accelerated electrons. The beam impinges on the chromosphere from the coronal part of a flare along the magnetic field tubes. Initially, the maximal energy flux was estimated as $F_{\max} \gtrsim 10^{11} \text{ erg cm}^{-2} \text{ s}^{-1}$. The time profile with the maximum at $t \lesssim 5 \text{ s}$ of the energy flux at the upper boundary of the chromosphere was typically

used for numerical solution of the two-temperature dissipative hydrodynamic equations (see Chap. 2 in [Somov 1992](#)), more exactly, in the *single-fluid two-temperature* approximation.

The electrons and ions are assumed to move with the same velocity implying no charge separation, i.e. *charge neutrality* (see Sect. 3.2.2). However

the electrons and ions of an impulsively heated plasma may each have their own Maxwellian distribution about different temperatures, necessitated by the slow temperature *relaxation time* of the electrons and ions as compared with self-relaxation.

In the solar corona and upper chromosphere, a strong magnetic field channels all the basic physical processes (except emission of radiation, of course) namely the propagation of accelerated electrons, heating by them, heating and cooling by the heat-conductive flows of energy, and other hydrodynamic processes. Thus the numerical code must solve the one-fluid two-temperature equations of hydrodynamics in *one dimensional* geometry. By using the Lagrangian coordinates, we can start with the following set of equations ([Somov et al. 1977, 1979, 1981](#)):

the continuity equation

$$\frac{\partial n}{\partial t} + n^2 \frac{\partial v}{\partial \xi} = 0, \quad (8.45)$$

the equation of motion

$$\frac{\partial v}{\partial t} + \frac{1}{m_H} \frac{\partial}{\partial \xi} [n k_B (T_i + x T_e)] = \frac{4}{3} \frac{1}{m_H} \frac{\partial}{\partial \xi} \left(\eta_i n \frac{\partial v}{\partial \xi} \right), \quad (8.46)$$

the electron temperature equation

$$\begin{aligned} \frac{n k_B}{(\gamma_g - 1)} \frac{\partial}{\partial t} (x T_e) + n \chi \frac{\partial x}{\partial \xi} - x k_B T_e \frac{\partial n}{\partial t} &= n \frac{\partial}{\partial \xi} \left(\kappa n \frac{\partial T_e}{\partial \xi} \right) + \\ &+ \mathcal{P}_e(n, \xi) - \mathcal{L}(n, T_e) - \mathcal{Q}(n, T_e, T_i), \end{aligned} \quad (8.47)$$

and the ion temperature equation

$$\frac{n k_B}{(\gamma_g - 1)} \frac{\partial T_i}{\partial t} - k_B T_i \frac{\partial n}{\partial t} = \frac{4}{3} \eta_i n^2 \left(\frac{\partial v}{\partial \xi} \right)^2 + \mathcal{Q}(n, T_e, T_i). \quad (8.48)$$

Here n , v , T_e and T_i are the number density, bulk velocity, electron and ion temperatures, respectively. We assume that neutral atoms have the same temperature as the ions. m_H is the atomic mass of Hydrogen, the most abundant element in the solar plasma. The presence of other elements is important in the radiative energy losses $\mathcal{L}$, see formula (12.23). $\gamma_g = c_p/c_v$ is the specific heat ratio. The symbols x and χ denote the degree of ionization of Hydrogen and its ionization potential. $\mathcal{P}_e$

is the rate of plasma heating by accelerated electrons. The rate of energy exchange between the electron and ion components of plasma (Braginskii 1965)

$$\mathcal{Q}(n, T_e, T_i) = \frac{4e^4(2\pi m_e)^{1/2}}{m_i k_B^{1/2}} \ln \Lambda n^2 \frac{(T_e - T_i)}{T_e^{3/2}}. \quad (8.49)$$

κ and η_i are the heat conductivity and ion viscosity, respectively. The Lagrange variable

$$\xi = - \int_{z_{\max}}^z n(z) dz + \xi_{\min}, \text{ cm}^{-2}, \quad (8.50)$$

z is the height above the photosphere. When we consider an impulsive heating of the solar chromosphere, $z_{\max}$ corresponds to the transition layer between the chromosphere and corona before the impulsive heating. Therefore, in this case, ξ is the column depth—the number of atoms and ions in a column (of the unit cross-section) measured down into the chromosphere from its upper boundary, the transition layer.

* * *

Before we shall illustrate the basic physics in the model under consideration, let us mention some results of space observations related to the accelerated electrons in solar flares. At first, the *Yohkoh* satellite (Ogawara et al. 1991) observations, made using three of the instruments on board—the Hard X-ray Telescope (HXT), the Soft X-ray Telescope (SXT), and the Bragg Crystal Spectrometer (BCS)—show that the non-thermal electron energy flux can be huge: even larger than that, for example, in the flare of 16 December 1991 (see Fig. 6a in McDonald et al. 1999), the maximal energy flux is

$$F_{\max} \approx 2.5 \times 10^{29} \text{ erg s}^{-1} / 2 \times 10^{17} \text{ cm}^2 \sim 10^{12} \text{ erg cm}^{-2} \text{ s}^{-1}.$$

Weak beams do not produce a significant response of the chromosphere (see Fig. 6b in McDonald et al. 1999), of course, just hard X-ray bremsstrahlung. So the most interesting are the most powerful electron beams.

In our days, it appears that the higher space-and-time resolution of optical and HXR observations, the higher the energy flux density of accelerated electron beams. One of the well-studied example is the X6.5 class white-light flare of 2006 December 6 (Krucker et al. 2011). The imaging observations were made at 430 nm (the Fraunhofer G band) by the Solar Optical Telescope (SOT; Tsuneta et al. 2008) on board *Hinode* (Kosugi et al. 2007) and in HXR by the *Ramaty High Energy Solar Spectroscopic Imager* (RHESSI). These two sets of data show closely similar ribbon structures, suggesting that the flare emissions in white light and in HXR have physically linked emission mechanisms. The G-band observations resolved the width of the ribbon with values between ~ 0.5 and ~ 1.8 arcsec.




Fig. 8.3 A scheme of the energy exchange in the two-temperature model of a single-fluid hydrodynamic response of the solar atmosphere to impulsive heating by an electron beam

The unresolved HXR observations revealed an even narrow ribbon in HXR. Applying the simplest thick-target model (Sect. 4.3.4), the derived energy deposition rate is estimated as

$$F_{\max} > 5 \times 10^{12} \text{ erg cm}^{-2} \text{ s}^{-1}.$$

This poses serious questions to the simplified thick-target models.

* * *

Anyway, if beam electrons lose their energy by mainly Coulomb collisions in the chromosphere according to this model,


the fastest process is the primary one, namely that of energy transfer from the beam electrons to the thermal electrons

of chromospheric plasma (Fig. 8.3).

As a result, plasma electrons are rapidly heated to high temperatures: in a matter of seconds the electron temperature reaches values of the order of ten million degrees. At the same time, the ion temperature lags considerably, by one order of magnitude, behind the electron temperature (Fig. 8.4).

The column depth $\xi_{\min} = n_{\text{cor}} l_{\text{rec}}$ is the number of ions inside a flaring loop which is the coronal part of a reconnected magnetic-field-line tube (see

Fig. 8.4 The distribution of the electron and ion temperatures over the column depth ξ into the chromosphere for $F_{\max} = 3 \times 10^{11} \text{ erg cm}^{-2} \text{ s}^{-1}$, the electron spectrum exponent $\gamma = 3$, and $\mathcal{E}_1 = 10 \text{ keV}$



Part II, Sect. 3.2.1); l_{rec} is the length of the reconnected field line, n_{cor} is the number density of plasma inside the tube above the transition layer between the chromosphere and corona before an impulsive heating. Let us assume, for simplicity, that

$$\xi_{\min} \ll \xi_1 = \frac{\mathcal{E}_1^2}{2a_1}, \text{ cm}^{-2}. \quad (8.51)$$

Here ξ_1 is the column thickness that the accelerated electrons with the minimal energy $\mathcal{E}_1$ measured in keV can pass in a plasma before they stop (see formula (4.40)). The assumption (8.51) means that we neglect the energy losses of the electrons in the coronal part of the loop. In this way, we consider direct impulsive heating of the chromosphere by an electron beam. Accelerated electrons penetrate into the chromosphere to significant depth. For this reason a significant fraction of the beam energy is lost as the radiation in optical and EUV lines. The column depth of evaporated plasma $\xi \approx 2 \times 10^{19} \text{ cm}^{-2}$ but its temperature does not exceed $T_{\max} \approx 10^7 \text{ K}$ in the example presented in Fig. 8.4, which is not the case of an extremely large energy deposition rate (Krucker et al. 2011).

The difference between the electron and ion temperatures is essential, at first, for the dynamics of high-temperature plasma

which absorbs the main part ($\geq 90\%$) of the beam energy flux. Let us imagine that only the electrons are heated, while the ion heating can be neglected. In this case the electron temperature is twice as large as it would be in the case of equal heating of the electrons and ions,

$$(T_e)_1 \simeq 2(T_e)_2.$$

The rate of high-temperature plasma cooling is mainly determined by heat fluxes into colder plasma. These can be evaluated by the formula for the classical heat flux

$$F_c = -\kappa_e \nabla T_e \quad (8.52)$$

under conditions when this formula is applicable, of course (see Somov et al. 1981). Here $\kappa_e = \kappa_0 T_e^{5/2}$ is the classical heat conductivity due to the Coulomb collisions

of plasma electrons. From formula (8.52) we see that the heat flux is proportional to $T_e^{7/2}$. Therefore the real heat flux

$$F_c(T_e)_1 \simeq 2^{7/2} F_c(T_e)_2 \quad (8.53)$$

can be an order of magnitude ($2^{7/2} \sim 10$) larger than the flux calculated in one-temperature ($T_e = T_i$) models. Because of this, the one-temperature models are much less dynamic than one would expect. We shall call them “*lazy*” models (e.g., McClymont and Canfield 1983; Allred et al. 2005); however they take into account the radiative transfer in a low-temperature plasma of solar flares.

The effect becomes even more important if the accelerated electrons heat a preliminary (before a flare) evaporated “*hot*” plasma. This formally means that, in formula (8.50), the column depth $\xi_{\min} = n_c l_r$ is not small in comparison with ξ_1 . So we have to take into account the direct impulsive heating of the plasma inside the coronal part of the flaring loop. Such process (Duijveman et al. 1983; MacNeice et al. 1984) can very efficiently produce a “*super-hot*” plasma which has an electron temperature T_e much higher than the maximal temperature in the case of chromospheric heating considered above.

In conclusion, we recall that heat conduction and radiative losses are strongly affected by only the electron temperature T_e . However *viscous heating* of plasma and *viscous force* are associated with ions and, hence, are determined by the ion temperature T_i ; see Exercise 8.1. Since a *dynamic viscosity* coefficient $\eta \sim T_i^{5/2}$ (10.31),

the higher ion temperature, the larger viscosity of high-temperature plasma heated by an electron beam as well as the stronger the viscous force;

see the positive feedback in Fig. 8.3. Note that the ion temperature is higher in one-temperature models than the ion temperature in two-temperature models,

$$(T_i)_1 > (T_i)_2,$$

in the region where $T_e > 10^6$ K in Fig. 8.4, i.e. above the flare *transition layer*. So, the “*lazy*” models are also much more viscous and, therefore, even much more lazy than the actual two-temperature models of hydrodynamic response of the solar atmosphere to impulsive heating by accelerated electrons in solar flares (Somov et al. 1977, 1979, 1981). It is clear that the physical features discussed above can be even more pronounced in the solar-type flares on other stars or generally in astrophysical plasmas.

8.3.3 Plasma Heating by Magnetic Reconnection

The difference between the electron and ion temperatures is known to be critical for a wide variety of kinetic effects, in particular for the generation of some turbulence (for example, ion-acoustic or ion-cyclotron) in the impulsively heated plasma. The




Fig. 8.5 Magnetic reconnection changes the topology of magnetic field lines and creates the fast plasma outflows. v_1 and $-v_1$ are the high-temperature plasma velocities near a reconnecting current layer

turbulence, in its turn, has a great impact on the efficiency of heating and particle acceleration in a plasma.

The second example, when the electron component of a plasma has a temperature that is considerably different from the ion temperature, is supplied by the physics of high-temperature (or even *super-hot*) *turbulent-current layers* (Somov 1981, 1986; Somov and Titov 1983) in the regions of magnetic reconnection. Since the layer thickness $2a$ is small in comparison with its width $2b$ (see Fig. 8.5), the plasma inflows quickly enter the region of the Joule dissipation of reconnecting magnetic field components. Here the impulsively fast heating of the electrons and ions takes place, resulting in considerably different temperatures.

The conditions in a *reconnecting current layer* (RCL) in the solar corona, especially, in flares (Part II, Sect. 6.3) are such that

the Coulomb exchange of energy between the impulsively heated electrons and ions inside the RCL can be entirely neglected.

One of distinctive features of fast reconnection in RCLs, proposed as the primary energy source in solar flares, is the presence of fast plasma outflows, or jets, whose velocities are nearly equal to the Alfvén velocity, see definition (15.27). Outflows can give origin to plasma velocity distributions with equal and opposite components along the x axis in Fig. 8.5 and, as a consequence, along the line-of-sight (l.o.s.) to an observer. Therefore, in this way, they can create a *symmetric supra-thermal broadening* in the soft X-ray and EUV lines observed during solar flares. The broadening mainly depends on the electron and ion temperatures inside the RCL (Antonucci and Somov 1992).

A comparison of the supra-thermal profiles of the Fe XXV emission lines observed at flare onset with the predictions of the high-temperature turbulent-current layer model suggests that the observed supra-thermal broadenings are consistent with the presence in the flare region of several small-scale or one (a few) curved large-scale RCLs (Antonucci et al. 1996).

The energy release by the reconnection effect has been invoked to explain both large-scale events, such as solar flares and coronal mass ejections (CMEs), and

small-scale phenomena, such as the coronal and chromospheric microflares that probably heat the corona (Part II, Sect. 12.4) and accelerate the solar wind.

The ultraviolet observations of the so-called explosive events in the solar chromosphere by SUMER (the Solar Ultraviolet Measurements of Emitted Radiation instrument) on the spacecraft *SOHO* (the Solar and Heliospheric Observatory) reveal the presence of *bi-directional plasma jets* ejected from small sites above the solar surface (Innes et al. 1997; cf. Antonucci and Somov 1992). The structure of these jets evolves in the manner predicted by theoretical models of reconnection (see Fig. 1 in Somov and Syrovatskii 1976a), thereby leading the first strong support to the view that reconnection is the fundamental process for accelerating plasma on the Sun.

In conclusion note that an impulsive heating of the solar atmosphere by a very powerful heat fluxes from super-hot turbulent-current layers can be considered in the *single-fluid two-temperature* approximation (Somov et al. 1982), by solving (8.45)–(8.48).

8.3.4 An Adiabatic Model for Two-Temperature Plasma

As we saw in Sect. 8.3.1, equilibrium in an electron-proton plasma is achieved in three stages. First, the electrons reach a Maxwellian distribution with temperature T_e on a time τ_{ee} . Then, on a longer time,

$$\tau_{pp} \approx (m_p/m_e)^{1/2} \tau_{ee},$$

the protons reach a Maxwellian distribution with temperature T_p . Finally, the two temperatures equalize on the longest time of order

$$\tau_{ep} \sim (m_p/m_e) \tau_{ee}.$$

Let us suppose that a two-temperature plasma is created by a strong shock wave in an electron-proton plasma. *The shock primarily heats ions* because the kinetic energy of a particle is proportional to the particle mass. In the post-shock region, the protons reach thermal equilibrium on a time τ_{pp} after they are heated through the shock (Zel'dovich and Raizer 1966, 2002). Within this time the proton temperature is significantly higher than the electron one. Subsequently the protons share their thermal energy with the electrons through Coulomb collisions.

In astrophysical plasma, sometimes, a difference between electron and ion temperatures can be observed at huge linear scales.

For example, the so-called *X-ray clusters*, or clusters of galaxies, with the X-ray temperatures $(4 - 10) \times 10^7$ K show noticeable differences between their electron and ion temperatures at radii greater than 2 Mpc $\approx 6 \times 10^{24}$ cm.

The clusters of galaxies are the largest objects in the Universe, containing galaxies and dark matter, collisionless particles and a diffuse gas component.

The last one is called the *intracluster medium* and has a temperature of about 10^8 K, thus emitting hard X-rays (HXR) mainly through the thermal bremsstrahlung of the electrons. In the outer parts of the clusters, the free-free cooling time is much longer than the Hubble time. So we neglect radiative cooling in such plasma which is supposed to be heated by the shock in the accretion flow (Takizawa 1998).

If we could also neglect heat conduction (for example, by assuming that the thermal conductivity of the intracluster medium is strongly reduced by a temperature gradient-driven kinetic instability, see Hattori and Umetsu 2000), then the electrons would be considered as an adiabatic gas. It would be very convenient to calculate the electron and ion temperature profiles by using the *adiabatic model* of a two-temperature plasma by Fox and Loeb (1997). This is also the case if tangled magnetic fields, for example of turbulent origin, can suppress heat conduction in high-temperature plasma. So one may assume that

there exists a chaotic magnetic field that is sufficiently strong to suppress heat conduction in high-temperature astrophysical plasma, *yet small enough* to have negligible dynamical and dissipative effects including Joule heating.

These conditions seem to be approximately satisfied in cluster environments; for more detail see Fox and Loeb (1997).

The general case of a strong shock in a fully ionized plasma with heat conduction is complicated by the fact that the electron thermal speed exceeds the shock speed, allowing the electrons to preheat the plasma ahead of the shock (Zel'dovich and Raizer 1966, Chap. 7, Sect. 12, Figs. 7.19 and 7.20).

Usually *heat conduction determines internal scales of the problem* being in competition with the thermal instability driven by radiative cooling (Field 1965; see also Somov and Syrovatskii 1976a). Radiation emitted by the high-temperature plasma behind the shock also may heat a pre-shock region. Fast particles, escaping from the high-temperature (super-hot) plasma (see Sect. 8.4.3), may contribute the pre-shock heating too. So we have to be very careful when we apply the adiabatic model of two-temperature plasma to astrophysical conditions.

If come back to the HXR tails observed in the X-ray spectra of some clusters, one suggestion is that all or part of this emission might be non-thermal bremsstrahlung from supra-thermal electrons with energies of $\sim 10\text{--}100$ keV. This non-thermal electrons would form a population in excess of the normal thermal gas, which is the bulk of the intra-cluster medium. The most natural explanation of this supra-thermal population would be that they are particles currently being accelerated to high energies by turbulence in the intra-cluster medium. Sarazin and Kempner (2000) have calculated models for the non-thermal HXR bremsstrahlung in the clusters of galaxies.

The high-Mach-number shocks in young supernova remnants (SNRs) do not produce electron-ion temperature equilibration either. The heating process in these collisionless shocks is not well understood, but the Coulomb collisions times are too long to provide the required heating. Presumably the plasma collective processes

should be responsible for the heating; see discussion and references in Sect. 16.4. This raises the question of whether the heating process leads to temperature equilibration or not. It appears that the observed electron temperature ($T_e \sim 1$ keV) remains very low compared to the observed ion temperature ($T_i \sim 500$ keV for ions O VII) behind the shock.

8.3.5 Two-Temperature Accretion Flows

Magnetized accretion disks have become the most convincing physical paradigm to explain a low emission from the central engines of *active galactic nuclei* (AGN) and X-ray binary sources (see also Sect. 13.2). The observed radiation comes from the energy dissipation required to maintain steady accretion of plasma on to the central object. In the standard model of the optically-thin accretion disk, the heat energy released by viscous dissipation is radiated almost immediately by the accreting plasma. So

the net luminosity must be equal to ($\approx$ one-half) the gravitational energy released as the mass falls onto the central object.

In a few of binary stellar systems, the mass of the primary star has been measured and found to be consistent with the mass of a neutron star, $\sim 1.4 M_\odot$. In several other systems, however, the mass of the primary is found to be greater than $3 M_\odot$, which makes these stars too massive to be neutron stars. These are considered as black hole candidates.

Although neutron stars and black holes have been distinguished on the basis of their masses, the real physical distinction between the two is that black holes must have a horizon (a surface through which the matter and energy fall in but from which nothing escapes) while neutron stars are normal stars with surfaces. This basic difference provides an opportunity to test the reality of black holes (Narayan et al. 1997).

Two-temperature advection-dominated accretion flows (ADAFs) have received much attention in an effort to explain low-luminosity stellar and galactic accreting sources (Blackman 1999; Wiita 1999; Manmoto 2000). Here the ions are assumed to receive the energy dissipated by the steady accretion *without having enough time to transfer their energy to the cooler electrons* before falling on to the central object.

While the electrons can almost always radiate efficiently, the protons will not, as long as Coulomb processes are the only thing that share energy between electrons and protons. So some or most of the dissipated energy is *advected* (Sect. 13.2.3), not radiated, as it would have been if the electrons received all of the dissipated energy. In the ADAF model, the heat generated via viscosity is advected inward rather than radiated away locally like a standard accretion disk (Novikov and Thorne 1973, Shakura and Sunyaev 1973).

When the central object is a black hole, the advected energy is lost forever rather than reradiated as it would be for a neutron star.

Precisely such observed differences between corresponding X-ray binary systems have been purported to provide evidence for black hole horizons (Narayan et al. 1997; see also Chakrabarti 1999); see, however, discussion of the ADAF model in Sect. 9.4.1.

8.4 Dynamic Friction in Astrophysical Plasma

8.4.1 The Collisional Drag Force and Energy Losses

8.4.1(a) Chandrasekhar-Spitzer's Formulae

As in Sects. 8.1 and 8.3, we use the concept of a test particle to illustrate the effects of the collisional drag force in astrophysical plasma. A test particle of mass m_1 and charge e_1 is incident with velocity $\mathbf{v}$ in a gas containing field particles of mass m_2 , charge e_2 and density n_2 . In what follows, $v_{\parallel}$ will be the component of the test particle velocity parallel to the original direction of its motion.

First, for the sake of simplicity, let us consider the field particles *at rest*. As in Sect. 8.1.5, integration over all possible values of the impact parameter up to the upper cut-off at $l = l_{\max}$ yields the following formulae describing the *mean rates* of energy losses and of scattering for the incident particle (Spitzer 1962):

$$\frac{d\mathcal{E}}{dt} = -\frac{2\pi e_1^2 e_2^2 \ln \Lambda}{\mathcal{E}} \frac{m_1}{m_2} n_2 v \quad (8.54)$$

and

$$\frac{d}{dt} v_{\parallel} = -\frac{\pi e_1^2 e_2^2 \ln \Lambda}{\mathcal{E}^2} \left(1 + \frac{m_1}{m_2}\right) n_2 v^2. \quad (8.55)$$

Here $\mathcal{E}$ is the energy of the incident particle (see definition (5.2)).

If we consider a beam of accelerated electrons in astrophysical ionized plasma, the most important are interactions with electrons and protons. So

$$\frac{d\mathcal{E}}{dt} = -\frac{2\pi e^4 \ln \Lambda}{\mathcal{E}} \left(1 + \frac{m_e}{m_p}\right) n_e v \quad (8.56)$$

and

$$\frac{d}{dt} v_{\parallel} = -\frac{\pi e^4 \ln \Lambda}{\mathcal{E}^2} \left(3 + \frac{m_e}{m_p}\right) n_e v^2. \quad (8.57)$$

Thus

both ambient electrons and protons produce scattering of the incident electrons but *only ambient electrons contribute significantly to the energy losses*;

the contribution of protons in the rate of energy losses (8.56) is proportional to the small ratio m_e/m_p . This is consistent, of course, with what we have concluded in Sect. 4.2 for fast particles propagating in thermal plasma.

We neglect collective effects due to interaction of the plasma and the electron beam as a whole without any justification here. It must be emphasized also at this point that formulae (8.56) and (8.57) describe the *mean* rates of change of $\mathcal{E}$ and $v_{\parallel}$ for the electrons of an incident beam but neglect the dispersions about these means. The accuracy of such procedure decreases as the scattering and energy losses become not small. These restrictions have been discussed in Sect. 4.4. Now we recall that we have neglected the proper motions of the plasma particles. Let us take them into account.

8.4.1(b) Energy Losses in Plasma

The most general non-relativistic formula for Coulomb losses in the many-component thermal plasma is given, for example, in Trubnikov (1965), Sivukhin (1966) and can be expressed as follows:

$$P \equiv \frac{d\mathcal{E}}{dt} = \sum_k \left(\frac{d\mathcal{E}}{dt} \right)_k = - \sum_k \frac{4\pi e^4 \ln \Lambda}{m_k} \frac{Z^2 Z_k^2 n_k}{v_k} \mathcal{P}_k \left(\frac{v}{v_k}, \frac{m_k}{M} \right). \quad (8.58)$$

Here Z_k , m_k , n_k and v_k are the charge, mass, density and thermal velocity of the plasma particles of the kind k ; they have a temperature T_k . Z , $M = Am_p$ and v are the charge, mass and velocity of the incident particles; their kinetic energy $\mathcal{E} = Mv^2/2$. Contrary to definition (8.15) of the mean thermal velocity, in formula (8.58) the thermal velocity is equal to the *most probable* velocity of thermal particles (Sivukhin 1966):

$$v_k = \left(\frac{2k_B T_k}{m_k} \right)^{1/2}. \quad (8.59)$$

It is convenient to determine the dimensionless variable

$$x_k = \frac{v}{v_k} = \left(\frac{m_k}{M} \frac{\mathcal{E}}{k_B T_k} \right)^{1/2} \quad (8.60)$$

and to rewrite the dimensionless function $\mathcal{P}_k$ as follows


$$\mathcal{P}_k \left(x_k, \frac{m_k}{M} \right) = \frac{1}{x_k} \operatorname{erf}(x_k) - \left(1 + \frac{m_k}{M} \right) \frac{2}{\sqrt{\pi}} \exp(-x_k^2). \quad (8.61)$$

Here

$$\operatorname{erf}(x_k) = \frac{2}{\sqrt{\pi}} \int_0^{x_k} \exp(-t^2) dt \quad (8.62)$$

is the probability integral.

Fig. 8.6 The Coulomb losses (with the sign *minus* in formula (8.58)) of energy of a test particle as a function of its velocity measured in the most probable velocity of the field thermal particles of the kind k



Let us consider the low-energy limit. Note that

$$\mathcal{P}_k \left(x_k, \frac{m_k}{M} \right) \approx \frac{2}{\sqrt{\pi}} \left[-\frac{m_k}{M} + \frac{2}{3} \left(1 + \frac{m_k}{M} \right) x_k^2 \right] \quad \text{if } x_k \ll 1. \quad (8.63)$$

Hence the dimensionless function

$$\mathcal{P}_k \left(0, \frac{m_k}{M} \right) = -\frac{2}{\sqrt{\pi}} \frac{m_k}{M} < 0 \quad (8.64)$$

and, according to formula (8.58), the energy losses rate

$$P_k \equiv \left(\frac{d\mathcal{E}}{dt} \right)_k = \frac{8\sqrt{\pi}e^4 \ln \Lambda}{M} \frac{Z^2 Z_k^2 n_k}{v_k} > 0. \quad (8.65)$$

This means that

■ a test particle with zeroth (or very small) velocity takes energy from the field particles having the temperature T_k .

The hot field particles heat a cold test particle.

Consider an opposite limiting case. If $x_k \gg 1$, then, being positive, the function

$$\mathcal{P}_k \left(x_k, \frac{m_k}{M} \right) \sim \frac{1}{x_k} \rightarrow 0 \quad \text{when } x_k \gg 1. \quad (8.66)$$

So the higher the energy of a test particle, the smaller are the Coulomb losses.

The maximum of the dimensionless function $\mathcal{P}_k$ is reached at $x_{k,\max} \approx 1.52$, see schematic Fig. 8.6.

Astrophysical plasma consists of many components. To obtain the total losses it is necessary to sum over all of them in formula (8.58). However two components—electrons and protons—give the largest contribution. In a plasma consisting of

electrons and protons with $n_e = n_p = n$ and temperatures T_e and T_p we have (Korchak 1980):

$$P = -c_\varepsilon \frac{Z^2}{A} \frac{n \ln \Lambda}{\sqrt{k_B T_e}} \left[\mathcal{P}_e \left(x_e, \frac{m_e}{M} \right) + \left(\frac{m_e T_e}{m_p T_p} \right)^{1/2} \mathcal{P}_p \left(x_p, \frac{m_p}{M} \right) \right], \quad (8.67)$$

where the constant $c_\varepsilon \approx 1.6 \times 10^{-23}$.

The location of both maxima of the function (8.67) is determined by conditions:

$$x_1 = x_p \approx 1.52 \quad \text{and} \quad x_2 = x_e \approx 1.52. \quad (8.68)$$

As follows from formula (8.67), the ratio of losses in the maxima

$$\frac{P_{\max, p}}{P_{\max, e}} = \left(\frac{m_e}{m_p} \frac{T_e}{T_p} \right)^{1/2} \approx \frac{1}{43} \left(\frac{T_e}{T_p} \right)^{1/2}. \quad (8.69)$$

■ The maximum of the electron Coulomb losses is the main energy threshold of the particle acceleration from low energies.

The proton barrier is considerably lower than the electron one.


The energy loss contribution of the proton component of astrophysical plasma does not seem to be important. This is not always true, however. First of all, formula (8.69) shows that the Coulomb losses on thermal protons increase with the growth of the ratio T_e/T_p . This may be an important case if particles of low energies are accelerated in *super-hot turbulent-current layers* (SHTCLs, see Part II, Sect. 6.3). The second argument comes from a consideration of very low energies of accelerated particles. In this region, the efficiency of acceleration is low for the majority of accelerating mechanisms. However, just in this region of low energies,

■ the Coulomb losses can strongly influence the nuclear composition and the charge-state of accelerated particles in astrophysical plasma

(Korchak 1980; see also Holman 1995; Bodmer and Bochsler 2000; Bykov et al. 2000).

When particular acceleration mechanisms in a astrophysical plasma are considered, the role of Coulomb collisions often reduces to the energy losses of the accelerated particles and, in particular, to the presence of the loss barrier at low velocities. As a result, Coulomb collisions decrease the efficiency of any acceleration mechanism. Contrary to this statement, we shall see that in many cases Coulomb collisions can play a much less trivial and not so passive role (e.g., Part II, Sect. 12.3.1). This makes plasma astrophysics more interesting.

Fig. 8.7 The collisional drag force F_k (with the sign *minus* in formula (8.70)) on a test particle as a function of its velocity $v_{\parallel}$ measured in the most probable velocity v_k of the field particles of the kind k



8.4.1(c) Dynamic Friction in Plasma

The collisional drag force F_f acts on a test particle (mass M , charge Ze) moving through the many-component plasma with the Maxwellian distribution of field particles:

$$M \frac{d}{dt} v_{\parallel} = -F_f = -\sum_k F_k(v_{\parallel}). \quad (8.70)$$

Here the velocity component $v_{\parallel}$ is parallel to the vector of the initial velocity of an incident test particle.

For a test particle with a velocity v much below the thermal velocity (8.59) of the field particles with the mass m_k , temperature T_k , and number density n_k ,

$$F_f \approx \sum_k \frac{4\pi e^4 \ln \Lambda}{k_B} \frac{Z^2 Z_k^2 n_k}{T_k} \left(1 + \frac{m_k}{M}\right) \frac{2}{3\sqrt{\pi}} \frac{v_{\parallel}}{v_k} \sim v_{\parallel}. \quad (8.71)$$

Therefore at small velocities the collisional drag force is proportional to the component $v_{\parallel}$ (cf. formula (1.16)).

When the test particle velocity exceeds the thermal velocity of the field particles, the drag force decreases with $v_{\parallel}$ as follows:

$$F_f = \sum_k F_k \approx \sum_k \frac{2\pi e^4 \ln \Lambda}{k_B} \frac{Z^2 Z_k^2 n_k}{T_k} \left(1 + \frac{m_k}{M}\right) \left(\frac{v_{\parallel}}{v_k}\right)^{-2} \sim v_{\parallel}^{-2}. \quad (8.72)$$

The general formula for collisional drag force is given, for example, in [Sivukhin \(1966\)](#) and is illustrated by schematic Fig. 8.7; here the dimensionless variable $x_{\parallel k} = v_{\parallel}/v_k$. The drag force vanishes when $x_{\parallel k} = 0$; it linearly increases with increasing $x_{\parallel k}$, becoming a maximum when

$$x_{\parallel k} = x_{\parallel k, \max} \approx 0.97, \quad (8.73)$$

and then falls off, approaching zero asymptotically as $x_{\parallel k} \rightarrow \infty$. This behavior of the drag force has important consequences discussed below.

8.4.2 Electric Runaway

It has been assumed above that the plasma is characterized by the Maxwellian distribution and that there are no external fields. Let us now assume that a uniform electric field $\mathbf{E}$ is switched on at some instant of time, the velocity distribution being assumed to be Maxwellian at this time. At least, at the beginning of the process when the velocity distribution has not yet changed appreciably, the time variation of the test-particle momentum $M\mathbf{v}$ due to Coulomb collisions with plasma particles will still be given by formulae (8.71) and (8.72) supplemented by the electric force $Ze\mathbf{E}$ in (8.70).

Thus, considering the component $v_{\parallel}$ as a component of the test-particle velocity $\mathbf{v}$ which is parallel to the electric field $\mathbf{E}$, we rewrite (8.70) as follows:

$$M \frac{d}{dt} v_{\parallel} = -F_f + ZeE = - \sum_k F_k + ZeE. \quad (8.74)$$

If the test-particle velocity is not small in comparison with the thermal velocity v_k , then the collisional drag force on a test particle falls off with increasing velocity v , according to formula (8.72), while the electric force is velocity independent. Therefore

for all particles with high enough velocities, the electric force exceeds the collisional drag force, and the particles are able to *run away* from the thermal distribution.

Equating the electric and collisional drag forces allows us to see the critical velocity v_{cr} above which runaway will occur for a given electric field strength E , see point *B* in Fig. 8.7. Runaway in astrophysical plasma can occur as long as there is a component of the electric field along the magnetic field. Before the acceleration of the heavy ions becomes significant, the acceleration of the light electrons gives rise to the *electron runaway* effect which was first predicted by Giovanelli (1949). He has shown that

- As the electric field applied to a highly ionized gas is increased, *the current*, which is initially limited by elastic collisions between electrons and positive ions, *increases* rapidly as the field strength reaches a critical value;
- This is due to a reduction in the cross-section of positive ions for scattering of electrons with increasing electron velocity.

In a strong electric field (or in a plasma of sufficiently low density and high temperature), all the electrons are accelerated by the field, i.e. become the runaway electrons. The Dreicer field (Dreicer 1959):

$$E_{\text{Dr}} = \frac{4\pi e^3 \ln \Lambda}{k_B} \frac{n_e}{T_e} \quad (8.75)$$

approximately corresponds to the electric field strength for which $v_{\text{cr}} = v_e$. Here v_e is the most probable velocity of thermal electrons (8.59).

In a weak field, only very fast electrons will run away, i.e. those velocity $v_{\parallel} \gg v_{\text{cr}}$. The velocity v_{cr} depends in an essential manner on the magnitude of electric field. In a weak field, the velocity v_{cr} is naturally much larger than the thermal velocity of electrons in the plasma. Therefore the number of runaway electrons should be very small if their distribution would remain Maxwellian for velocities $v_{\parallel} \lesssim v_{\text{cr}}$. This is not true however.

In order to determine the flux of runaway electrons we must know the way in which the density of electrons having a velocity $v_{\parallel} \sim v_{\text{cr}}$ varies under action of the runaway effect. This means that we must know the velocity distribution for the electrons for $v_{\parallel} \sim v_{\text{cr}}$. To consider this problem self-consistently it is necessary to solve the kinetic equation taking both collisions and the electric field into account (Sect. 4.5). It appears that

Coulomb collisions create a power-law tail distribution between a region of thermal velocities and the region where $v_{\parallel} \approx v_{\text{cr}}$ with a constant flux of electrons directed from low to high velocities.

By so doing, *Coulomb collisions increase the flux of runaway electrons* (Gurevich 1961).

To have an idea of the magnitude of the Dreicer field (see Exercise 8.4), let us substitute the definition of the Debye-Hückel radius (8.31) in formula (8.75) and assume that $T_e = T_p = T$ and $n_e = n_p = n$. We find

$$E_{\text{Dr}} = \frac{e}{r_{\text{DH}}^2} \frac{\ln \Lambda}{2} \sim \frac{e}{r_{\text{DH}}^2}. \quad (8.76)$$

So the Dreicer field is approximately equal to the electric field of a positive charge at a distance slightly smaller than the Debye-Hückel radius.

8.4.3 Thermal Runaway in Astrophysical Plasma

Let us consider a plasma with a non-uniform distribution of electron temperature T_e . Let l_T be the characteristic length of the temperature profile and λ_e be the mean free path of thermal electrons. For the classical heat conductivity to be applicable, it is necessary to satisfy a condition (Sect. 9.6):

$$\lambda_e \ll l_T \equiv \frac{T_e}{|\nabla T_e|}. \quad (8.77)$$

The mean free path of a particle increases with its velocity. This can be seen from formula (8.13) which gives us the mean free path

$$\lambda = \tau v_1 \sim v_1^4. \quad (8.78)$$

That is why

■ a number of fast electrons can penetrate from a hot plasma into cold one even if the gradient of temperature is very small.

In such a way, the hot plasma can lose some part of its thermal energy transferred by fast thermal escaping electrons. In addition to the usual heat flux (8.52), which is determined *locally* by the Coulomb collisions of plasma electrons, there appears a *non-local* energy flux carried by the fast electrons practically without collisions. A classical diffusive heat transfer and a convective one, determined by *thermal runaway* electrons, are always present in plasma.

It is interesting for astrophysical applications that, at not too small temperature gradients, the convective transfer of thermal energy can play a principal role. Gurevich and Istomin (1979) have examined the case of a *small temperature gradient*. By using a perturbation analysis for the high-speed kinetic equation (Sect. 4.2), they have shown that the fast growth of the mean free path with increasing velocity gives an abrupt growth of the number of fast electrons in the cold plasma.

The opposite case of a *large temperature gradient* in the narrow transition layer between a high-temperature plasma and a cold one was investigated by many authors with applications to the problem of energy transfer in the solar atmosphere. For example, Shoub (1983) has solved numerically the boundary-value problem for the Fokker-Planck equation in the model of the transition layer between the corona and the chromosphere in quiet conditions. An excess of fast electrons has been found in the low transition layer region. As for solar flares, the prevailing view is that

■ the high-temperature plasma can lose energy efficiently by the convective heat transfer by the thermal runaway electrons

(see Somov 1992, Sect. 2.2).

In both cases, however, it is important to take into account that the *fast runaway electrons*, similar to any beam of fast particles, *generate the electric field which drives the reverse current* of thermal electrons. Diakonov and Somov (1988) have found an analytical solution to the self-consistent kinetic problem on the beam of escaping thermal electrons and its associated reverse current (Sect. 4.5). They have shown that the reverse-current electric field in solar flares leads to a significant reduction of the convective heat flux carried by fast electrons escaping from the high-temperature plasma to the cold one.

Recommended Reading: Sivukhin (1966), Somov (1992).

8.5 Practice: Exercises and Answers

Exercise 8.1 (Sect. 8.1). For an electron, which moves in the solar corona with a mean thermal velocity (Exercise 8.2), evaluate the characteristic time of *close* and *distant* collisions with thermal protons.

Answer. Characteristic time of close electron-proton collisions follows from formula (8.13) and is equal to

$$\tau_{cl, ep} = \frac{m_e^2}{\pi e^4} \frac{V_{Te}^3}{n_p} \approx 4.96 \times 10^{-18} \frac{V_{Te}^3}{n_p}, \text{ s.} \quad (8.79)$$

At typical temperature of electrons in the corona $T_e \approx 2 \times 10^6$ K, their thermal velocity (5.54) $V_{Te} \approx 9.5 \times 10^8 \text{ cm s}^{-1}$. Substituting this value in (8.79) and assuming $n_p \approx n_e \approx 2 \times 10^8 \text{ cm}^{-3}$, we find that $\tau_{cl, ep} \approx 22$ s.

According to (8.21) the characteristic time of distant collisions is $8 \ln \Lambda$ shorter than the close collision time (8.79). Hence, first, we have to find the value of the Coulomb logarithm (8.34):

$$\ln \Lambda = \ln \left[\left(\frac{3k_B^{3/2}}{2\pi^{1/2} e^3} \right) \left(\frac{T_e^3}{n_e} \right)^{1/2} \right] \approx \ln \left[1.25 \times 10^4 \left(\frac{T_e^3}{n_e} \right)^{1/2} \right]. \quad (8.80)$$

At typical coronal temperature and density, formula (8.80) gives

$$\ln \Lambda \approx 22.$$

With this value of $\ln \Lambda$ formula (8.21) gives

$$\tau_{\perp, ep} = \frac{m_e^2}{\pi e^4} \frac{1}{8 \ln \Lambda} \frac{V_{Te}^3}{n_p} \approx 2.87 \times 10^{-20} \frac{V_{Te}^3}{n_p}, \text{ s.} \quad (8.81)$$

In the solar corona $\tau_{\perp, ep} \approx 0.1$ s. Therefore the *distant* collisions of thermal electrons with thermal protons in the corona are really much more frequent in comparison with close collisions.

Exercise 8.2 (Sect. 8.2). Evaluate the Debye-Hückel radius and the plasma frequency in the solar corona.

Answer. From (8.31) it follows that for electron-proton plasma with $T_e = T_p = T$ and $n_e = n_p = n$ the Debye-Hückel radius

$$r_{DH} = \left(\frac{k_B T}{8\pi e^2 n} \right)^{1/2} \approx 4.9 \left(\frac{T}{n} \right)^{1/2}, \text{ cm.} \quad (8.82)$$

Under conditions in the solar corona $r_{DH} \approx 0.5$ cm.

The electron plasma frequency (8.35)

$$\omega_{pl}^{(e)} = \left(\frac{4\pi e^2 n_e}{m_e} \right)^{1/2} \approx 5.64 \times 10^4 \sqrt{n_e}, \text{ rad s}^{-1}, \quad (8.83)$$

or

$$\nu_{pl}^{(e)} = \omega_{pl}^{(e)} / 2\pi \approx 10^4 \sqrt{n_e}, \text{ Hz}. \quad (8.84)$$

In the solar corona $\omega_{pl}^{(e)} \sim 10^9 \text{ rad s}^{-1}$.

Exercise 8.3 (Sect. 8.3). Under conditions of Exercise 8.1 evaluate the *exact* (determined by formulae (8.38) and (8.39)) *collisional times* between thermal electrons and between thermal protons, respectively. Compare these times with the characteristic time of energy exchange between electrons and protons in the coronal plasma and in a *transition layer* between the solar corona and chromosphere.

Answer. By substituting $\ln \Lambda$ in (8.38), we have the following expression for the thermal electron collisional time

$$\tau_{ee} = \frac{m_e^2}{0.714 e^4 8\pi \ln \Lambda} \frac{V_{Te}^3}{n_e} \approx 4.04 \times 10^{-20} \frac{V_{Te}^3}{n_e}, \text{ s}. \quad (8.85)$$

In the solar corona $\tau_{ee} \approx 0.2 \text{ s}$. For thermal protons formula (8.39) gives

$$\tau_{pp} = \frac{m_p^2}{0.714 e^4 8\pi \ln \Lambda} \frac{V_{Tp}^3}{n_p} \approx 1.36 \times 10^{-13} \frac{V_{Tp}^3}{n_p}, \text{ s}. \quad (8.86)$$

Assuming $T_p = T_e$ and $n_p = n_e$, we find the proton collisional time in the solar corona $\tau_{pp} \approx 7 \text{ s}$; this is in a good agreement with formula (8.40), of course.

Let us find the time of energy exchange between electrons and protons. By using formula (8.44), we have

$$\tau_{ep}(\mathcal{E}) \approx 22 \tau_{pp} \approx 164 \text{ s}. \quad (8.87)$$

So the energy exchange between electron and proton components in the coronal plasma is the slowest process determined by Coulomb collisions.

It is convenient to rewrite formula (8.85) for the collisional time of thermal electrons in the following form

$$\tau_{ee} \approx 1.24 \times 10^{-2} \frac{T_e^{3/2}}{n_e}, \text{ s}. \quad (8.88)$$

In order to estimate this time in a transition layer between the solar corona and chromosphere we put $T_e \approx 10^5 \text{ K}$ and $n_e \approx 4 \times 10^9 \text{ cm}^{-3}$. Thus we find $\tau_{ee} \sim 10^{-4} \text{ s}$. For thermal protons formula (8.86) gives

$$\tau_{pp} \approx 5.26 \times 10^{-1} \frac{T_p^{3/2}}{n_p}, \text{ s}. \quad (8.89)$$

So in the transition layer $\tau_{ee} \sim 4 \times 10^{-3}$ s. Finally the energy exchange time between electrons and protons in the transition layer is also small: $\tau_{ep} \sim 0.1$ s. This is why the approximation $T_e = T_p = T$ is sufficient for description of many phenomena in the solar transition region ([Shmeleva and Syrovatskii 1973](#)).

Exercise 8.4 (Sect. 8.4). Evaluate and compare Dreicer's electric fields in the solar corona and in the chromosphere.

Answer. From (8.75) it follows that

$$E_{\text{Dr}} = \frac{4\pi e^3 \ln \Lambda}{k_B} \frac{n_e}{T_e} \approx 7.54 \times 10^{-8} \frac{n_e (\text{cm}^{-3})}{T_e (\text{K})}, \text{ V cm}^{-1}. \quad (8.90)$$

Here it was taken $\ln \Lambda \approx 21.6$ according to Exercise 8.1.

At typical temperature and number density of electrons in the solar corona $T_e \approx 2 \times 10^6$ K and $n_e \approx 2 \times 10^8 \text{ cm}^{-3}$, we find that the Dreicer electric field

$$E_{\text{Dr}} \approx 7 \times 10^{-6} \text{ V cm}^{-1} \sim 10^{-5} \text{ V cm}^{-1}.$$

The same value follows, of course, from formula (8.76) with $r_{\text{DH}} \approx 0.5$ cm (see Exercise 8.3).

In the solar chromosphere $n_e > 2 \times 10^{10} \text{ cm}^{-3}$ and $T_e < 10^4$ K. According to formula (8.90), the Dreicer electric field

$$E_{\text{Dr}} > 0.1 \text{ V cm}^{-1}$$


in the chromosphere is, at least, 10^4 times stronger than the coronal one.

Exercise 8.5. Define the dynamic friction by gravitational force as momentum loss by a massive moving object, for example a star in a galaxy, due to its gravitational interaction with its own gravitationally induced wake. Discuss two possibilities: (a) the background medium consists of collisionless matter (other stars in the galaxy), (b) the medium is entirely gaseous (e.g., [Ostriker 1999](#)). The first case, the *gravitational drag* in collisionless systems ([Chandrasekhar 1943b](#)), has widespread theoretical application in modern astrophysics.

Hint. At first, let us qualitatively understand why a friction force should arise in a collisionless gravitational system. Suppose a star has moved from a point *A* to a point *B* as shown in Fig. 8.8.

While passing from *A* to *B*, the star attracted the surrounding stars towards itself. Hence the number density of stars around *AB* should be slightly larger than that ahead of *B*. Therefore the star at the point *B* experiences a net gravitational attraction in the backward direction, i.e. in the direction opposite to the direction of the star velocity vector *v*.

Fig. 8.8 An illustration of the origin of dynamic friction in a collisionless gravitational system



The variety of consequences of the gravitational drag force in collisionless astronomical systems includes the mass segregation in star clusters, sinking satellites in dark matter galaxy halo, orbital decay of binary supermassive black holes after galaxy mergers, etc. (Binney and Tremaine 1987).

Exercise 8.6. Discuss why the rate of escape of stars from a galactic cluster, evaluated ignoring dynamic friction, is too rapid to be compatible with a life for the cluster (Chandrasekhar 1943c). Show that the escape rate is drastically reduced when dynamic friction is allowed for.

Chapter 9

Macroscopic Description of Astrophysical Plasma

Abstract In this chapter we treat individual kinds of particles as continuous media, mutually penetrating charged gases which interact between themselves and with an electromagnetic field. This approach gives us the multi-fluid model which is useful to consider many properties of astrophysical plasmas, for example, the solar wind.

9.1 Summary of Microscopic Description

The averaged Liouville equation or kinetic equation gives us a *microscopic* (though averaged in a statistical sense) description of the plasma state's evolution. Let us consider the way of transition to a less comprehensive *macroscopic* description of a plasma. We start from the kinetic equation for particles of kind k , in the form derived in Sect. 2.2.3:

$$\frac{\partial f_k(X, t)}{\partial t} + v_\alpha \frac{\partial f_k(X, t)}{\partial r_\alpha} + \frac{F_{k,\alpha}(X, t)}{m_k} \frac{\partial f_k(X, t)}{\partial v_\alpha} = \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c \quad (9.1)$$

Here the *statistically averaged* force is

$$F_{k,\alpha}(X, t) = \sum_l \int_{X_1} F_{kl,\alpha}(X, X_1) f_l(X_1, t) dX_1 \quad (9.2)$$

and the collisional integral

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \frac{\partial}{\partial v_\alpha} J_{k,\alpha}(X, t), \quad (9.3)$$

where the flux of particles of kind k

$$J_{k,\alpha}(X, t) = \sum_l \int_{X_1} \frac{1}{m_k} F_{kl,\alpha}(X, X_1) f_{kl}(X, X_1, t) dX_1 \quad (9.4)$$

in the six-dimensional phase space $X = \{\mathbf{r}, \mathbf{v}\}$.

9.2 Definition of Macroscopic Quantities

Before turning our attention to the deduction of equations for the macroscopic quantities or macroscopic *transfer* equations, let us define the following *moments* of the distribution function.

- (a) *The zeroth moment* (without multiplying the distribution function f_k by the velocity $\mathbf{v}$)

$$\int_{\mathbf{v}} f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v} = n_k(\mathbf{r}, t) \quad (9.5)$$

is obviously the number of particles of kind k in a unit volume, i.e. the *number density* of particles of kind k . It is related to the *mass density* of particles of kind k in a natural way:

$$\rho_k(\mathbf{r}, t) = m_k n_k(\mathbf{r}, t).$$

The plasma mass density is accordingly

$$\rho(\mathbf{r}, t) = \sum_k \rho_k(\mathbf{r}, t) = \sum_k m_k n_k(\mathbf{r}, t). \quad (9.6)$$

- (b) *The first moment* of the distribution function, i.e. the integral of the product of the velocity $\mathbf{v}$ to the first power and the distribution function f_k ,

$$\int_{\mathbf{v}} v_\alpha f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v} = n_k u_{k,\alpha} \quad (9.7)$$

is the particle flux density, i.e. product of the number density of particles of kind k by their *mean velocity*

$$u_{k,\alpha}(\mathbf{r}, t) = \frac{1}{n_k} \int_{\mathbf{v}} v_\alpha f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (9.8)$$

Consequently, the *mean momentum* of particles of kind k in a unit volume is expressed in terms of the first moment of the distribution function as follows

$$m_k n_k u_{k,\alpha} = m_k \int_{\mathbf{v}} v_\alpha f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (9.9)$$

(c) *The second moment* of the distribution function is defined to be

$$\Pi_{\alpha\beta}^{(k)}(\mathbf{r}, t) = m_k \int_{\mathbf{v}} v_\alpha v_\beta f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v} = m_k n_k u_{k,\alpha} u_{k,\beta} + p_{\alpha\beta}^{(k)}. \quad (9.10)$$

Here we have introduced

$$v_\alpha = u_{k,\alpha} + v'_\alpha,$$

thus

$$v'_\alpha = v_\alpha - u_{k,\alpha}$$

is the deviation of the particle velocity from its mean value

$$u_{k,\alpha} = \langle v_{k,\alpha} \rangle_v$$

in the sense of the definition (9.8). Therefore

$$\langle v'_\alpha \rangle_v = 0;$$

and

$$p_{\alpha\beta}^{(k)} = m_k \int_{\mathbf{v}} v'_\alpha v'_\beta f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}, \quad (9.11)$$

is termed the *pressure tensor*.

$\Pi_{\alpha\beta}^{(k)}$ is the *tensor of momentum flux density* for particles of kind k . Its component $\Pi_{\alpha\beta}^{(k)}$ is the α th component of the momentum transported by the particles of kind k , in a unit time, across the unit area perpendicular to the axis r_β .

Once we know the distribution function $f_k(\mathbf{r}, \mathbf{v}, t)$, which contains all the statistically averaged information on the particles of kind k at the *microscopic* level, we can derive all *macroscopic* quantities related to these particles. So, higher moments of the distribution function will be introduced as needed in what follows.

Actually one of the related problems is that we do not know the distribution function f_k but can observe or measure several moments of it in an astrophysical phenomenon or laboratory experiment. Another problem is complexity of the equations that describes these moments.

9.3 Macroscopic Transfer Equations

Note that the deduction of macroscopic equations is nothing but just the derivation of the equations for the distribution function moments.

9.3.1 Equation for the Zeroth Moment

Let us calculate the *zeroth* moment of the kinetic equation (9.1):

$$\int_{\mathbf{v}} \frac{\partial f_k}{\partial t} d^3\mathbf{v} + \int_{\mathbf{v}} v_\alpha \frac{\partial f_k}{\partial r_\alpha} d^3\mathbf{v} + \int_{\mathbf{v}} \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} d^3\mathbf{v} = \int_{\mathbf{v}} \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v}. \quad (9.12)$$

We interchange the order of integration over velocities and the differentiation with respect to time t in the first term and with respect to coordinates r_α in the second one. Under the second integral

$$v_\alpha \frac{\partial f_k}{\partial r_\alpha} = \frac{\partial}{\partial r_\alpha} (v_\alpha f_k) - f_k \frac{\partial v_\alpha}{\partial r_\alpha} = \frac{\partial}{\partial r_\alpha} (v_\alpha f_k) - 0,$$

since $\mathbf{r}$ and $\mathbf{v}$ are independent variables in the phase space X .

Taking into account that the distribution function quickly approaches zero as $v \rightarrow \infty$, the integral of the third term is taken by parts and is equal to zero (Exercise 9.1).

Finally, the integral of the right-hand side of (9.12) describes the change in the number of particles of kind k in a unit volume, in a unit time, as a result of collisions with particles of other kinds. If the processes of transformation, during which the particle kind can be changed (such as ionization, recombination, charge exchange, dissociation etc., see Exercise 9.2), are not allowed for, then the last integral is zero as well:

$$\int_{\mathbf{v}} \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v} = 0. \quad (9.13)$$

Thus, by integration of (9.12), the following equation is found to result from (9.1)

$$\frac{\partial n_k}{\partial t} + \frac{\partial}{\partial r_\alpha} n_k u_{k,\alpha} = 0.$$

(9.14)

This is the usual *continuity equation* expressing the conservation of particles of kind k or (that is the same, of course) conservation of their mass:




Fig. 9.1 From the microscopic to the macroscopic view of a plasma. LT is the Liouville theorem and the Liouville equation (1.26) for an exact distribution function $\hat{f}_k$. KE and BC are the kinetic equation (2.39) and the equation for the binary correlation function. m_0 is the equation for the zeroth moment of the distribution function f_k , the number density n_k of the particles of kind k . This equation is unclosed

$$\frac{\partial \rho_k}{\partial t} + \frac{\partial}{\partial r_\alpha} \rho_k u_{k,\alpha} = 0. \quad (9.15)$$

Here

$$\rho_k(\mathbf{r}, t) = m_k n_k(\mathbf{r}, t)$$

is the mass density of particles of kind k .

Equation (9.14) for the zeroth moment n_k depends on the unknown first moment $u_{k,\alpha}$. This is illustrated by Fig. 9.1.

9.3.2 The Momentum Conservation Law

Now let us calculate the *first* moment of the kinetic equation (9.1) multiplied by the mass m_k :

$$\begin{aligned} m_k \int_{\mathbf{v}} \frac{\partial f_k}{\partial t} v_\alpha d^3 \mathbf{v} + m_k \int_{\mathbf{v}} v_\alpha v_\beta \frac{\partial f_k}{\partial r_\beta} d^3 \mathbf{v} + \int_{\mathbf{v}} v_\alpha F_{k,\beta} \frac{\partial f_k}{\partial v_\beta} d^3 \mathbf{v} = \\ = m_k \int_{\mathbf{v}} v_\alpha \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3 \mathbf{v}. \end{aligned} \quad (9.16)$$

With allowance made for the definitions (9.7) and (9.10), we obtain the *momentum conservation law*

$$\begin{aligned} \frac{\partial}{\partial t} (m_k n_k u_{k,\alpha}) + \frac{\partial}{\partial r_\beta} \left(m_k n_k u_{k,\alpha} u_{k,\beta} + p_{\alpha\beta}^{(k)} \right) - \\ - \langle F_{k,\alpha}(\mathbf{r}, t) \rangle_v = \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v. \end{aligned} \quad (9.17)$$

Here $p_{\alpha\beta}^{(k)}$ is the pressure tensor (9.11), i.e. a part of the unknown second moment which is the tensor of momentum flux density (9.10).

The *mean force* acting on the particles of kind k in a unit volume (the mean force *per unit volume*) is (see Exercise 9.3):

$$\langle F_{k,\alpha}(\mathbf{r}, t) \rangle_v = \int_{\mathbf{v}} F_{k,\alpha}(\mathbf{r}, \mathbf{v}, t) f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v}. \quad (9.18)$$

This should not be confused with the *statistical mean force* $F_{k,\alpha}(\mathbf{r}, \mathbf{v}, t)$ acting on a single particle (see definition (9.2)). The statistically averaged force (9.2) is under the integral in formula (9.18).

In the particular case of the Lorentz force, we rewrite the mean force per unit volume as follows:

$$\langle F_{k,\alpha}(\mathbf{r}, t) \rangle_v = n_k e_k \left[E_\alpha + \frac{1}{c} (\mathbf{u}_k \times \mathbf{B})_\alpha \right]$$

or

$$\langle F_{k,\alpha}(\mathbf{r}, t) \rangle_v = \rho_k^q E_\alpha + \frac{1}{c} (\mathbf{j}_k^q \times \mathbf{B})_\alpha. \quad (9.19)$$

Here ρ_k^q and $\mathbf{j}_k^q$ are the mean densities of electric charge and current, produced by the particles of kind k . However note that

the mean electromagnetic force couples all the charged components of astrophysical plasma together

because the electric and magnetic fields, $\mathbf{E}$ and $\mathbf{B}$, act on all charged components and, at the same time, all charged components contribute to the electric and magnetic fields according to Maxwell's equations.

The right-hand side of (9.17) contains the mean force resulting from collisions, i.e. the *mean collisional force* (see Exercise 9.4):

$$\langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = m_k \int_{\mathbf{v}} v_\alpha \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v}. \quad (9.20)$$

Substituting (9.3) in definition (9.20) gives us the following formula

$$\langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = -m_k \int_{\mathbf{v}} v_\alpha \frac{\partial}{\partial v_\beta} J_{k,\beta} d^3\mathbf{v}. \quad (9.21)$$

Let us integrate (9.21) by parts. For this purpose, at first, we find the derivative

$$\frac{\partial}{\partial v_\beta} (v_\alpha J_{k,\beta}) = J_{k,\beta} \frac{\partial v_\alpha}{\partial v_\beta} + v_\alpha \frac{\partial}{\partial v_\beta} J_{k,\beta}.$$

From this it follows that

$$\begin{aligned} v_\alpha \frac{\partial}{\partial v_\beta} J_{k,\beta} &= -J_{k,\beta} \delta_{\alpha\beta} + \frac{\partial}{\partial v_\beta} (v_\alpha J_{k,\beta}) = \\ &= -J_{k,\alpha} + \frac{\partial}{\partial v_\beta} (v_\alpha J_{k,\beta}). \end{aligned} \quad (9.22)$$

On substituting (9.22) and (9.4) in (9.20) and integrating, we obtain the most general formula for the mean collisional force

$$\begin{aligned} \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v &= m_k \int_{\mathbf{v}} J_{k,\alpha}(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v} = \\ &= \sum_{l \neq k} \int_{\mathbf{v}} \int_{\mathbf{v}_1} \int_{\mathbf{r}_1} F_{kl,\alpha}(\mathbf{r}, \mathbf{v}, \mathbf{r}_1, \mathbf{v}_1) f_{kl}(\mathbf{r}, \mathbf{v}, \mathbf{r}_1, \mathbf{v}_1, t) d^3\mathbf{r}_1 d^3\mathbf{v}_1 d^3\mathbf{v}. \end{aligned} \quad (9.23)$$

Note that

for the particles of the same kind, the *elastic* collisions cannot change the total particle momentum per unit volume.

That is why $l \neq k$ in the sum (9.23).

Formula (9.23) contains the unknown binary correlation function f_{kl} . The last should be found from the correlation function (2.49) indicated as the second link *BC* in Fig. 9.2. Thus

the equation for the first moment of the distribution function is as much unclosed as the initial kinetic equation,


which is the first equation of the chain for correlation functions (see the first link *KE* in Fig. 9.2). Therefore the equation for the first moment is unclosed in *two* directions.

If there are several kinds of particles, and if each of them is in the state of thermodynamic equilibrium (as will be discussed in Sect. 9.4.2), then the mean collisional force can be expressed in terms of the *mean momentum loss* during the collisions of a particle of kind k with the particles of other kinds:

$$\langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = - \sum_{l \neq k} \frac{m_k n_k (u_{k,\alpha} - u_{l,\alpha})}{\tau_{kl}}.$$

(9.24)

Fig. 9.2 KE and BC are the kinetic equation and the equation for the binary correlation function. m_0, m_1 are the equations for the first two moments of the distribution function f_k . The link m_1 is unclosed in two directions



Here $\tau_{kl}^{-1} = \nu_{kl}$ is the mean frequency of collisions between the particles of kinds k and l . Formula (9.24) has the status of a good approximation in plasma astrophysics. General theoretical derivation of this very useful formula is given, for example, in [Golant et al. \(1977\)](#), see Sect. 6.3, formula (6.59).

If $u_{k,\alpha} > u_{l,\alpha}$ then the mean collisional force is negative:

the fast particles of kind k slow down by dint of collisions with the slowly moving particles of other kinds.

The force is zero, once the particles of all kinds have identical mean velocities. Therefore the mean collisional force, as well as the mean electromagnetic force, tends to make astrophysical plasma be a single hydrodynamic medium (see Sect. 12.1).

9.4 The Energy Conservation Law

9.4.1 The Second Moment Equation

The second moment (9.10) of a distribution function f_k is the tensor of momentum flux density $\Pi_{\alpha\beta}^{(k)}$. In general, in order to find an equation for this tensor, we should multiply the kinetic equation (9.1) by the factor $m_k v_\alpha v_\beta$ and integrate over velocity space $\mathbf{v}$. In this way, we could arrive to a *matrix equation* in partial derivatives; see also Sect. 9.7 where the classical Grad moment method is applied to obtain similar equations. If we take the trace of this equation we could obtain the partial differential *scalar equation* for energy density of the particles under consideration

(e.g., [Shkarofsky et al. 1966](#); Sect. 9.2). This is the correct self-consistent way which is the basis of the moment method. For our aims, a simpler direct procedure is sufficient and correct in some applications to astrophysical problems.

In order to derive the *energy conservation law*, we multiply (9.1) by the particle's kinetic energy $m_k v_\alpha^2/2$ and integrate over velocities, taking into account that

$$v_\alpha = u_{k,\alpha} + v'_\alpha, \quad \langle v'_\alpha \rangle_v = 0,$$

and

$$v_\alpha^2 = u_{k,\alpha}^2 + (v'_\alpha)^2 + 2 u_{k,\alpha} v'_\alpha.$$

A straightforward integration yields

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\rho_k u_k^2}{2} + \rho_k \varepsilon_k \right) + \frac{\partial}{\partial r_\alpha} \left[\rho_k u_{k,\alpha} \left(\frac{u_k^2}{2} + \varepsilon_k \right) + p_{\alpha\beta}^{(k)} u_{k,\beta} + q_{k,\alpha} \right] = \\ = \rho_k^q (\mathbf{E} \cdot \mathbf{u}_k) + (\mathbf{F}_k^{(c)} \cdot \mathbf{u}_k) + Q_k^{(c)}(\mathbf{r}, t) + \mathcal{L}_k^{(rad)}(\mathbf{r}, t). \end{aligned} \quad (9.25)$$

Here

$$\begin{aligned} m_k \varepsilon_k(\mathbf{r}, t) &= \frac{1}{n_k} \int_{\mathbf{v}} \frac{m_k (v'_\alpha)^2}{2} f_k(\mathbf{r}, \mathbf{v}, t) d^3 \mathbf{v} = \\ &= \frac{m_k}{2n_k} \int_{\mathbf{v}} (v'_\alpha)^2 f_k(\mathbf{r}, \mathbf{v}, t) d^3 \mathbf{v} \end{aligned} \quad (9.26)$$

is the *mean kinetic energy* of *chaotic* (non-directed) motion per single particle of kind k . Thus the first term on the left-hand side of (9.25) represents the time derivative of the energy of the particles of kind k in a unit volume, which is the sum of kinetic energy of a *regular* motion with the mean velocity $\mathbf{u}_k$ and the so-called *internal* energy.

As every tensor, the pressure tensor can be written as

$$p_{\alpha\beta}^{(k)} = p_k \delta_{\alpha\beta} + \pi_{\alpha\beta}^{(k)}. \quad (9.27)$$

Thus, on rearrangement, we obtain the following general equation

$$\begin{aligned} \frac{\partial}{\partial t} \left(\frac{\rho_k u_k^2}{2} + \rho_k \varepsilon_k \right) + \frac{\partial}{\partial r_\alpha} \left[\rho_k u_{k,\alpha} \left(\frac{u_k^2}{2} + w_k \right) + \pi_{\alpha\beta}^{(k)} u_{k,\beta} + q_{k,\alpha} \right] = \\ = \rho_k^q (\mathbf{E} \cdot \mathbf{u}_k) + (\mathbf{F}_k^{(c)} \cdot \mathbf{u}_k) + Q_k^{(c)}(\mathbf{r}, t) + \mathcal{L}_k^{(rad)}(\mathbf{r}, t). \end{aligned} \quad (9.28)$$

Here

$$w_k = \varepsilon_k + \frac{p_k}{\rho_k} \quad (9.29)$$

is the *heat function* per unit mass. Therefore the second term on the left-hand side contains the energy flux

$$\rho_k u_{k,\alpha} \left(\frac{u_k^2}{2} + w_k \right),$$

which can be called the “*advective*” flux of kinetic energy.

Let us mention the well known astrophysical application of this term. In principle,

the *advective cooling* of ions heated by viscosity might dominate the cooling by the electron-ion collisions,

for example, in a low-density high-temperature plasma flow near a rotating black hole. In such an *advection-dominated accretion flow* (ADAF), the heat generated via viscosity is transferred inward the black hole rather than radiated away locally like in a standard accretion disk model (see Sects. 8.3.5 and 13.2).

On the other hand, discussing the ADAF model as a solution for the important astrophysical problem should be treated with reasonable cautions. Looking at (9.25) for electrons and ions separately, we see that

too many assumptions have to be made to arrive to the ADAF approximation.

For example, this is not realistic to assume that plasma electrons are heated only due to Coulomb collisions with plasma ions and, for this reason, the electrons are much cooler than the ions because the electrons do not have enough time to be heated to a high temperature of ions. However it is not necessary to heat electrons to high temperatures (say $T_e > 10^6$ K) when collisions become unimportant. If electrons reach $T_e \sim 10^5$ K they quickly lose a lot of energy via the optically thin emission of EUV radiation. So the ions also become cool.

Therefore the suggestions underlying the ADAF approximation ignore several physical effects. Moreover, in general, they ignore *reconnection* and dissipation of magnetic fields (regular and random) in astrophysical plasma. This makes a physical basis of the ADAF model very uncertain.

9.4.2 The Case of Thermodynamic Equilibrium

In order to clarify the physical meaning of the definitions given above, let us, for a while, come back to the general principles of plasma physics. If the particles of the k th kind are in *thermodynamic equilibrium* state, then f_k is the Maxwellian function with the *temperature* T_k . In the case of local thermodynamic equilibrium (LTE), frequent collisions keep the distribution function f_k in a local region at a point $\mathbf{r}$ close to the Maxwellian

$$f_k^{(0)}(\mathbf{r}, \mathbf{v}) = n_k(\mathbf{r}) \left[\frac{m_k}{2\pi k_B T_k(\mathbf{r})} \right]^{3/2} \exp \left\{ -\frac{m_k |\mathbf{v} - \mathbf{u}_k(\mathbf{r})|^2}{2 k_B T_k(\mathbf{r})} \right\}, \quad (9.30)$$

see Sect. 9.6. In this case, according to formula (9.26), the mean kinetic energy of chaotic motion per single particle of kind k

$$m_k \varepsilon_k = \frac{3}{2} k_B T_k. \quad (9.31)$$

The pressure tensor (9.11) is isotropic, i.e. diagonal and constant:

$$p_{\alpha\beta}^{(k)} = p_k \delta_{\alpha\beta}, \quad (9.32)$$

where the scalar

$$p_k = n_k k_B T_k \quad (9.33)$$

is the gas pressure of the particles of kind k . This is also the equation of state for the *ideal* gas. Thus we have found that the pressure tensor is diagonal. This implies the absence of *viscosity* for the ideal gas:

$$\pi_{\alpha\beta}^{(k)} = 0. \quad (9.34)$$

The *heat function* per unit mass or, more exactly, the *specific enthalpy* is

$$w_k = \varepsilon_k + \frac{p_k}{\rho_k} = \frac{5}{2} \frac{k_B T_k}{m_k}. \quad (9.35)$$

This is a particular case of the thermodynamic equilibrium state; it will be discussed in Sect. 9.6.

9.4.3 The General Case of Non-equilibrium Plasma

In general, we do not expect that the system of the particles of kind k has reached thermodynamic equilibrium. Nevertheless we often use the mean kinetic energy (9.26) to define the *effective* or *kinetic* temperature

$$T_{\text{ef},k} = \frac{2}{3} \frac{m_k}{k_B} \varepsilon_k \quad (9.36)$$

according to definition (9.31).

■ A kinetic temperature is just a measure for the spread of the particle distribution in velocity space.

The kinetic temperatures of different components (electrons and ions) in astrophysical plasma may differ from each other:

$$T_{\text{ef},e} \neq T_{\text{ef},i} . \quad (9.37)$$

Moreover, in an anisotropic plasma, the kinetic temperatures parallel and perpendicular to the magnetic field are different:

$$T_{\text{ef},k \parallel} \neq T_{\text{ef},k \perp} . \quad (9.38)$$

Therefore, two heat-flux vectors $q_{k \parallel}$ and $q_{k \perp}$ are introduced to describe separately the transfer of longitudinal and transverse thermal energies.

Without supposing thermodynamic equilibrium, in an *anisotropic* plasma, the part associated with the deviation of the distribution function from the isotropic one (which does not need to be a Maxwellian function in general, of course) is distinguished in the pressure tensor:

$$p_{\alpha\beta}^{(k)} - p_k \delta_{\alpha\beta} = \pi_{\alpha\beta}^{(k)} \neq 0 . \quad (9.39)$$

Here $\pi_{\alpha\beta}^{(k)}$ is the *viscous stress tensor*. Recall that we did not derived an equation for this traceless symmetric tensor.

The term $\pi_{\alpha\beta}^{(k)} u_{k,\beta}$ in the energy-conservation (9.25) represents the flux of energy released by the *viscous force* in the particles of kind k . The non-diagonality of tensor $\pi_{\alpha\beta}^{(k)}$ implies that this force is related to the transport of momentum (see (9.17)) from a fast-moving layer of plasma to the adjoining slow-moving layer.

The last term on the left-hand side of the energy-conservation (9.25), the vector

$$q_{k,\alpha} = \int_{\mathbf{v}} \frac{m_k (v')^2}{2} v'_\alpha f_k(\mathbf{r}, \mathbf{v}, t) d^3\mathbf{v} \quad (9.40)$$

is the *heat flux density* due to the particles of kind k in a system of coordinates, in which the gas of these particles is immovable at a given point $\mathbf{r}$ of space. Formula (9.40) shows that a *third-order-moment* term appears in the second order moment of the kinetic equation.

9.4.4 The Sources and Sinks of Energy

The left-hand side of the energy conservation law (9.25) has a typical divergent form. The right-hand side of (9.25) contains the following four terms that play the role of sources and sinks of energy.

(a) The first term

$$\rho_k^q(\mathbf{E} \cdot \mathbf{u}_k) = n_k e_k E_\alpha u_{k,\alpha} \quad (9.41)$$

is the work done by the Lorentz force (without the magnetic field, of course) in unit time on unit volume.

(b) The second term

$$\left(\mathbf{F}_k^{(c)} \cdot \mathbf{u}_k \right) = u_{k,\alpha} \int_{\mathbf{v}} m_k v'_\alpha \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3 \mathbf{v} \quad (9.42)$$

is the work done by the collisional force of friction of the particles of kind k with all other particles in unit time on unit volume. This means that

the work of friction force results from the mean momentum change of particles of kind k (*moving with the mean velocity* $\mathbf{u}_k$) owing to collisions with all other particles.

This work equals zero if $\mathbf{u}_k = 0$.

(c) The third term

$$\mathcal{Q}_k^{(c)}(\mathbf{r}, t) = \int_{\mathbf{v}} \frac{m_k (v')^2}{2} \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3 \mathbf{v} \quad (9.43)$$

is the rate of thermal energy release (heating or cooling) in a gas of the particles of kind k due to collisions with other particles. Recall that the collisional integral depends on the binary correlation function f_{kl} .

(d) The last term $\mathcal{L}_k^{(rad)}(\mathbf{r}, t)$ takes into account that a plasma component k can gain energy by absorbing radiations of different kinds and can lose energy by emitting radiations. Various processes of emission by charged particles contribute to the emissivity of astrophysical plasmas. The major contributions in a fully-ionized *super-hot* plasma come from collisional *bremsstrahlung* (Sect. 8.1.5), Cherenkov emission (Sects. 7.1.2 and 7.4), and magnetic *bremsstrahlung*.

These processes do not cover, of course, the whole variety of those realized in astrophysical conditions. For example, a lot of spectral lines are registered in the spectra of cosmic sources. They originate due to electron transitions between discrete energy levels of ions, atoms, and molecules. In optically thin plasma in the thermodynamical equilibrium state, with a cosmic abundance of elements, the power of radiative cooling is characterized by the presence of several maxima (see Sect. 12.1.3). For this reason, the astrophysical plasmas are often subject to thermal instability (Field 1965).

9.5 General Properties of Transfer Equations

9.5.1 Divergent and Hydrodynamic Forms

Equations (9.14), (9.17), and (9.25) are referred to as the equations of particle, momentum and energy *transfer*, respectively; and the approximation in which they have been obtained is called the model of *mutually penetrating* charged gases. These gases are not assumed to be in the thermodynamic equilibrium. However the definition of the temperature (9.31) may be generally considered as formally coinciding with the corresponding definition pertaining to the gas of particles of kind k in thermodynamic equilibrium.

The equations of mass, momentum and energy transfer are written in the *divergent* form. This essentially states the conservation laws and turns out to be convenient in numerical work, to construct the conservative schemes for computations. Sometimes, other forms are more convenient. For instance, the equation of momentum transfer or simply the equation of motion (9.17) can be brought into the frequently used form (with the aid of the continuity (9.14) to remove the derivative $\partial \rho_k / \partial t$):

$$\rho_k \left(\frac{\partial u_{k,\alpha}}{\partial t} + u_{k,\beta} \frac{\partial u_{k,\alpha}}{\partial r_\beta} \right) = - \frac{\partial}{\partial r_\beta} p_{\alpha\beta}^{(k)} + \langle F_{k,\alpha}(\mathbf{r}, t) \rangle_v + \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v. \quad (9.44)$$

The so-called *substantial derivative* appears on the left-hand side of this equation:

$$\frac{d^{(k)}}{dt} = \frac{\partial}{\partial t} + u_{k,\beta} \frac{\partial}{\partial r_\beta} = \frac{\partial}{\partial t} + \mathbf{u}_k \cdot \nabla_{\mathbf{r}}.$$

(9.45)


This substantial or *advective* derivative—the total time derivative following a *fluid element* of kind k —is typical of hydrodynamic-type equations, to which the equation of motion (9.44) belongs.

In the frame of reference, in which the fluid element of the particles of kind k is not moving, the mean velocity $\mathbf{u}_k = 0$ but the time partial derivative $\partial/\partial t$ does not vanish of course.

The total time derivative with respect to the mean velocity $\mathbf{u}_k$ of the particles of kind k is different for each kind k . In Chap. 12 on the one-fluid MHD theory, we shall introduce the substantial derivative with respect to the average velocity of the plasma as a whole.

For the case of the Lorentz force (9.19), the equation of motion of the particles of kind k can be rewritten as follows:

Fig. 9.3 LT is the Liouville theorem and the Liouville equation (1.26) for an exact distribution function $\hat{f}_k$. KE and BC are the kinetic equation and the equation for the binary correlation function. m_0, m_1 etc. are the chain of the equation for the moments of the distribution function f_k



$$\rho_k \frac{d^{(k)} u_{k,\alpha}}{dt} = -\frac{\partial}{\partial r_\beta} p_{\alpha\beta}^{(k)} + \rho_k^q E_\alpha + \frac{1}{c} (\mathbf{j}_k^q \times \mathbf{B})_\alpha + \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v. \quad (9.46)$$

Here the right-hand side represents the density of the forces acting on the fluid element of kind k , in particular, the last term is the mean collisional force (9.20) or, more specifically, (9.24). The left-hand side of (9.46) is the net change of the momentum density of this fluid element.

9.5.2 Status of the Conservation Laws

As we saw in Sect. 9.3, when we treat a plasma as several continuous media (the mutually penetrating charged gases), for each of them,

the main three average properties (density, velocity, and a quantity like temperature or pressure) are governed by the *basic conservation laws* for mass, momentum, and energy in the media.

These conservation equations are useful, of course, except they contain more unknowns than the number of equations. The transfer equations for local macroscopic quantities are as much unclosed as the initial kinetic equation (9.1) which is the first equation of the chain for correlation functions (see KE in Fig. 9.3). For example, formula (9.23) for the mean collisional force contains the unknown binary correlation function f_{kl} . The last should be found from the correlation function (2.49) indicated as the second link BC in Fig. 9.3. The terms (9.42) and (9.43) in the energy conservation (9.25) also depend on the unknown binary correlation function f_{kl} .

It is also important that the transfer equations are unclosed in “orthogonal” direction: (9.14) for the zeroth moment (see the link m_0 in Fig. 9.3), density n_k , depends on the unknown first moment, the mean velocity $\mathbf{u}_k$, and so on. This process of generating equations for the higher moments could be extended indefinitely depending solely on how many primary variables (n_k , $\mathbf{u}_k$, ε_k , ...) one is prepared to introduce. However, if at any level the distribution function is known, or can be approximated to, in terms of the primary variables for which the equations have already been generated, then this set of equation should be closed. We will come back to this critical point in the next section. Anyway we know now that

three basic conservation laws for mass, momentum, and energy in the components of astrophysical plasma represent the main transfer equations that are *the first three links* in the chain of the equations for the distribution function moments.

It certainly would not be easy if possible to arrive to this fundamental conclusion and would be difficult to derive the conservation laws in the form of the transfer equations (9.14), (9.17), and (9.25) in the way which is typical for the majority of textbooks (even very nice ones, like [Clarke and Carswell 2007](#); [Bradt 2008](#)): from simple specific knowledge to more general ones. Such generalization means that we go from well-known things to more complicated ones, for example, from the Newton equation of motion of a particle to the ordinary hydrodynamic equation of fluid motion. Though this way makes a text easier to read, it does not give a reader the complete knowledge of a subject. That is why we selected the opposite way: from general to specific knowledge.

The consecutive consideration of physical principles, starting from the most general ones, and of simplifying assumptions, which give us a simpler description of plasma under astrophysical conditions, allows us to find the answers to *two key questions*:

1. What approximation is the best one (the simplest but sufficient) for description of a phenomenon in astrophysical plasma;
2. How to build an adequate model for the phenomenon, for example, a solar flare or a flare in the corona of an accretion disk.

From a mathematical point of view, an elegant treatment of the particle transfer in plasma can be also well based on the use of non-canonical conjugate variables (for example, $\mathbf{r}$ and $\mathbf{p}$ are not canonically conjugate for a system of charged particles moving under the Lorentz force) and the associated Lie algebra (see [Balescu 1988](#)).

9.6 Equation of State and Transfer Coefficients

9.6.1 General Approach to the Problem

The first three transfer equations for a plasma component k would be closed with respect to the three unknown variables ρ_k , $\mathbf{u}_k$, and ε_k , if it were possible to express the other *unknown* quantities p_k , $\pi_{\alpha\beta}^{(k)}$, and $q_\alpha^{(k)}$ in terms of these three variables, or the variables ρ_k , $\mathbf{u}_k$ and the formally defined temperature T_k . For this purpose, we have to know the *equation of state* and the so-called *transfer coefficients*. How can we find them?

Formally, we should write equations for higher (than second) moments of the distribution function, e.g., (9.58) and (9.59). However these equations will not be closed either. So, how shall we proceed?

According to the general principles of statistical physics (Maxwell 1860, see Sects. 72 and 74 in [Sivukhin 1990](#)),

by virtue of collisions in a closed system of particles, any distribution function tends to assume the Maxwellian form.

In this case the equation of state is that of the ideal gas.

The *Maxwellian distribution* is the kinetic equation solution for a stationary homogeneous gas in the absence of any mean force in the thermal equilibrium state, i.e. for a gas in *thermodynamic equilibrium* ([Boltzmann 1872](#); see also [Boltzmann 1956](#); [Kubbinga 2006](#)). Then spatial gradients and derivatives with respect to time are zero. In fact they are always nonzero. For this reason, the assumption of full thermodynamic equilibrium is replaced with the *local* thermodynamic equilibrium (LTE). Moreover

if the gradients and derivatives are *small*, then the real distribution function differs *little* from the local Maxwellian one, the difference being *proportional* to the small gradients or derivatives.

Thus, in plasma, if we are interested in the processes occurring in a time t , which is much greater than the characteristic collision time τ , and at a distance L , which is much larger than the particle mean free path λ ,

$$t \gg \tau, \quad L \gg \lambda, \quad (9.47)$$

then the particle distribution function $f_k(\mathbf{r}, \mathbf{v}, t)$ can be thought of as a sum of the local Maxwellian distribution

$$f_k^{(0)}(\mathbf{r}, \mathbf{v}, t) = n_k(\mathbf{r}, t) \left[\frac{m_k}{2\pi k_B T_k(\mathbf{r}, t)} \right]^{3/2} \times \exp \left\{ -\frac{m_k |\mathbf{v} - \mathbf{u}_k(\mathbf{r}, t)|^2}{2 k_B T_k(\mathbf{r}, t)} \right\} \quad (9.48)$$

and some *small* additional term $f_k^{(1)}(\mathbf{r}, \mathbf{v}, t)$. Therefore

$$f_k(\mathbf{r}, \mathbf{v}, t) = f_k^{(0)}(\mathbf{r}, \mathbf{v}, t) + f_k^{(1)}(\mathbf{r}, \mathbf{v}, t), \quad \left| f_k^{(1)} \right| < f_k^{(0)}. \quad (9.49)$$

According to (9.48), the function $f_k^{(0)}$ depends on t and $\mathbf{r}$ through the functions $n_k(\mathbf{r}, t)$, $T_k(\mathbf{r}, t)$ and $\mathbf{u}_k(\mathbf{r}, t)$. Therefore we can find the derivatives $\partial f_k^{(0)} / \partial t$ and $\partial f_k^{(0)} / \partial r_\alpha$. Now, by using these derivatives, we substitute formula (9.49) in the kinetic equation (9.1) and linearly approximate the collisional integral (9.3) by using one or another of the models introduced in Chap. 3; alternatively, see Exercise 9.5 as a specific example. Then we seek the additional term $f_k^{(1)}$ in the *linear* approximation with respect to the factors disturbing the Maxwellian distribution, such as gradients of physical parameters, electric fields etc. The quantities $q_\alpha^{(k)}$, $\pi_{\alpha\beta}^{(k)}$ etc., which in their turn are proportional to the same factors, can be expressed in terms of $f_k^{(1)}$. The proportionality coefficients are the sought-after *transfer coefficients*.

9.6.2 The Classical Heat Conductive Flux

As example, let us consider the case of the collisional heat flux q_α , both the additional term $f_k^{(1)}$ and the flux q_α are chosen to be proportional to the temperature gradient. Thus, in a fully ionized plasma in the limit of a vanishing magnetic field, we find the heat flux in the electron component of plasma:

$$q_e = -\kappa_e \nabla T_e, \quad (9.50)$$

where

$$\kappa_e \approx \frac{1.84 \times 10^{-5}}{\ln \Lambda} T_e^{5/2} \quad (9.51)$$

is the coefficient of electron *thermal conductivity* (Spitzer 1962).

In the presence of strong magnetic field in astrophysical plasma, all the transfer coefficients become highly anisotropic. Since the Maxwellian function (9.48) and its derivatives are uniquely determined by the parameters n_k , $\mathbf{u}_k$, and T_k , the transfer coefficients are expressed in terms of the same quantities and magnetic field B , of course.

■ This procedure makes it possible to close the set of transfer equations for astrophysical plasma under the conditions (9.47).

The first step is to calculate the departure $f_k^{(1)}$ from the Maxwellian distribution function by using some method of handling collisions. This allows us to disconnect the third link m_2 in the vertical chain of the moment equations in Fig. 9.3 from the horizontal chain of equations related to correlation functions. Several models have

been suggested on different grounds to account for collisions in plasma (Shkarofsky et al. 1966; Krall and Trivelpiece 1973). The simplest of them is considered in Exercise 9.5.

The first three moment equations were extensively used in astrophysics, for example, in investigations of the *solar wind*. They led to a significant understanding of phenomena such as escape, acceleration, and cooling. However, as more detailed solar wind observations become available, it appeared that the simplified, collisionally dominated models are not adequate for most of the interplanetary range and for most of the times, i.e. most physical states of the solar wind.

A higher order, closed set of equations for the six moments have been derived for multi-fluid, *moderately non-Maxwellian plasma* of the solar wind (Cuperman and Dryer 1985). On the basis of these equations, for example, the generalized expression for heat flux relates the flux to the temperature gradients, relative streaming velocity, thermal anisotropy, temperature differences of the components.

By investigating the breakdown of classical hydrodynamics, for example, in describing electron thermal energy transport away from a strong heat source (e.g., in a solar flare plasma heated by reconnecting super-hot turbulent-current layers, SHTCLs) we conclude that

significant problems rest in the time-dependent treatment of high-temperature plasma heating in the solar atmosphere

(Oreshina and Somov 2011a). Classical formula (9.50), called the Fourier law, does not adequately treat this very rapid process; see Sect. 9.7.3. However the Grad thirteen moment equations, introduced in the next section, do provide rapid time response through a time dependent differential equation for the heat flux in addition to the standard hydrodynamic equations (Oreshina and Somov 2011b).

Recommended Reading: Braginskii (1965), Lifshitz and Pitaevskii (1981), Hollweg (1986).

9.7 The Higher-Order Moment Equations

9.7.1 The 13 Moment Equations by Grad

In the classical method of Grad's moments (Grad 1949), the distribution function, which is written in this section for a single species for convenience (cf. Sect. 12.1), is expanded in velocity moments about a local Maxwellian distribution $f^{(0)}$, i.e.

$$f(\mathbf{r}, \mathbf{v}, t) = f^{(0)}(\mathbf{r}, \mathbf{v}, t) \sum_{n=0}^{\infty} \frac{1}{n!} a_{[\alpha]}^{(n)}(\mathbf{r}, t) H_{[\alpha]}^{(n)}(\mathbf{v}). \quad (9.52)$$

Here the index $\alpha = 1, 2, 3$ or (x, y, z) ; $H_{[\alpha]}^{(n)}$ is a Chebishev–Hermite polynomial of n th degree (see [Lavrent'ev and Shabat 1973](#), Chap. 7, Sect. 92–94); $H_{[\alpha]}^{(n)}$ and $a_{[\alpha]}^{(n)}$ are tensors of n th rank. Due to the orthogonality properties of the Chebishev–Hermite polynomials, the inversion formula for the coefficients is

$$a_{[\alpha]}^{(n)} = \frac{1}{n(\mathbf{r}, t)} \int f(\mathbf{r}, \mathbf{v}, t) H_{[\alpha]}^{(n)} d^3 \mathbf{v}. \quad (9.53)$$

The first few coefficients $a_{[\alpha]}^{(n)}$ can be related to the familiar definitions for density, fluid velocity, temperature, pressure tensor, and heat flux.

When the expansion (9.52) is substituted into the kinetic equation (9.1) and integrated over $d^3 \mathbf{v}$, an infinite set of first order partial differential equations for the $a_{[\alpha]}^{(n)}$ is obtained. If the equations up to $a_{\alpha\beta\gamma}^{(3)}$ are retained, there are a total of twenty coupled equations. In fact, the equations never decouple for any order $a^{(n)}$ because in general $a_{[\alpha]}^{(n+1)}$ will appear in the equation for $\partial a_{\alpha\beta\gamma}^{(n)} / \partial t$. To close the set, the $a_{[i]}$ term is evaluated formally using an ansatz for the distribution function that is complete up to the n th term in the infinite series expansion.

This system of equations is typically reduced to thirteen by contracting the third rank tensor ($a_{\alpha\beta\gamma}^{(3)} \rightarrow a_{\alpha\beta\beta} \rightarrow q_\alpha$) to a vector that can be related to the thermal heat flux. The thirteen coefficients (i.e., variables in the moment equations) are then n , $u_1, u_2, u_3, p_{11}, p_{22}, p_{33}, p_{12} = p_{21}, p_{23} = p_{32}, p_{31} = p_{13}, q_1, q_2$, and q_3 , while the corresponding ansatz for the distribution function takes the form (e.g., [Golant et al. 1977](#), Sect. 6.2; [Moses and Duderstadt 1977](#)):

$$f(\mathbf{r}, \mathbf{v}, t) = f^{(0)} \left[1 + \pi_{\alpha\beta} \frac{v'_\alpha v'_\beta}{p V_T^2} + 4 q_\alpha v'_\alpha \frac{(v'_\beta v'_\beta - (5/2) V_T^2)}{5 p V_T^4} \right]. \quad (9.54)$$

Here the pressure tensor

$$p_{\alpha\beta} = p \delta_{\alpha\beta} + \pi_{\alpha\beta}, \quad p = (1/3) p_{\alpha\alpha},$$

$\pi_{\alpha\beta}$ is the viscous stress tensor which allows for the transport of momentum from one layer of a plasma flow to other layers; the deviation of the particle velocity from its mean value u_α , i.e. $v'_\alpha = v_\alpha - u_\alpha$; and the mean thermal velocity $V_T = (3k_B T/m)^{1/2}$.

The resulting set of thirteen moment equations is then comprised of the conventional *hydrodynamic equations* (the general consideration of multi-fluid plasma in electromagnetic field is given in [Golant et al. 1977](#), Sects. 6.3, 6.4) of mass, momentum, and energy conservation:

$$\frac{\partial \rho}{\partial t} + u_\alpha \frac{\partial \rho}{\partial r_\alpha} + \rho \frac{\partial u_\alpha}{\partial r_\alpha} = 0, \quad (9.55)$$

$$\rho \left(\frac{\partial u_\alpha}{\partial t} + u_\beta \frac{\partial u_\alpha}{\partial r_\beta} \right) = - \frac{\partial p_{\alpha\beta}}{\partial r_\beta}, \quad (9.56)$$

$$\frac{3}{2} n k_B \left(\frac{\partial T}{\partial t} + u_\alpha \frac{\partial T}{\partial r_\alpha} \right) = - \frac{\partial q_\alpha}{\partial r_\alpha} - p_{\alpha\beta} \frac{\partial u_\beta}{\partial r_\alpha}, \quad (9.57)$$

augmented by two additional equations which represent generalizations of the traditional transport laws:

$$\frac{\partial p_{\beta\gamma}}{\partial t} + u_\alpha \frac{\partial p_{\beta\gamma}}{\partial r_\alpha} + p_{\alpha\gamma} \frac{\partial u_\beta}{\partial r_\alpha} + p_{\alpha\beta} \frac{\partial u_\gamma}{\partial r_\alpha} + p_{\beta\gamma} \frac{\partial u_\alpha}{\partial r_\alpha} + \frac{\partial p_{\alpha\beta\gamma}}{\partial r_\alpha} = R(p_{\beta\gamma}), \quad (9.58)$$

$$\begin{aligned} & \frac{\partial q_\beta}{\partial t} + u_\alpha \frac{\partial q_\beta}{\partial r_\alpha} + \frac{7}{5} q_\alpha \frac{\partial u_\beta}{\partial r_\alpha} + \frac{7}{5} q_\beta \frac{\partial u_\alpha}{\partial r_\alpha} + \frac{2}{5} q_\gamma \frac{\partial u_\gamma}{\partial r_\beta} + \\ & + p \frac{\partial}{\partial r_\alpha} \left(\frac{\pi_{\alpha\beta}}{\rho} \right) + \frac{5}{2} \frac{\pi_{\alpha\beta}}{\rho} \frac{\partial k_B T}{\partial r_\alpha} + \frac{5}{2} \frac{p}{m} \frac{\partial k_B T}{\partial r_\beta} - \\ & - \frac{\pi_{\beta\gamma}}{\rho} \frac{\partial \pi_{\alpha\gamma}}{\partial r_\alpha} = R(q_\beta). \end{aligned} \quad (9.59)$$

Here

$$\begin{aligned} p_{\alpha\beta\gamma} &= \int f v'_\alpha v'_\beta v'_\gamma d^3\mathbf{v}, \\ R(q_\beta) &= \int \left(\frac{\partial \hat{f}}{\partial t} \right)_c (f) \frac{m}{2} v'_\alpha v'_\alpha v'_\beta d^3\mathbf{v}, \end{aligned}$$

and

$$R(p_{\beta\gamma}) = \int \left(\frac{\partial \hat{f}}{\partial t} \right)_c (f) m v'_\beta v'_\gamma d^3\mathbf{v},$$

while $(\partial \hat{f} / \partial t)_c(f)$ is the collisional integral, e.g., Fokker-Planck (3.10) or Boltzmann (3.45). So, the equations for second moments of the distribution function f are not closed of course.

9.7.2 The Powerful Heat Flux Problem

Since our concern in Sect. 9.6 was with heat transfer processes in solar flares, we focus our attention on (9.59). We suggest that the unphysical infinite heat flux predicted by Fourier's law (see Oreshina and Somov 2011a) can be corrected by including higher order conservation equations, as given by Grad's method, in

the description of the dynamic behavior of the solar plasma. This constitutes a systematic application of a classical technique for solving the kinetic equation. Since the conventional hydrodynamic equations (9.55)–(9.57) are the result of the same technique, we are only suggesting that within the same framework, we carry the application to the next higher order equations. We suspect that the functional dependence $f(\mathbf{r}, \mathbf{v}, |n, v'_\alpha, T, p_{\alpha\beta}, q_\alpha)$ is a better representation of the distribution function than is $f(\mathbf{r}, \mathbf{v}, |n, v'_\alpha, T)$.

The equation governing the time dependent heat flux as it is written in (9.59) is unsuitable for further detailed analysis of the impulsive heating plasma in solar flares. Therefore some approximations and additions must be made. To model the plasma we chose the one-fluid two-temperature description, see Sect. 8.3.2. Hence the single fluid velocity will be that of the massive ions, i.e. protons. Within this context, all convective terms in the electron heat flux equation will be regarded as small in this analysis. This can be justified by noting that the electron conduction characteristic velocity, V_{T_e} , is the electron thermal velocity whereas the convection characteristic velocity is the ion thermal velocity, V_{T_i} . Thus,

$$V_{T_i}/V_{T_e} \sim (m_e/m_i)^{1/2}.$$

We also recognize that the electron viscosity is generally much smaller than the ion viscosity. So since we are treating the electrons in this analysis, we shall also assume that viscosity terms are small. We anticipate that the dominant feature in our analysis will be the time dependent heat flux. These assumptions greatly simplify the heat flux equation

$$\frac{\partial q_\alpha}{\partial t} + \frac{5}{2} \frac{p}{m} \frac{\partial k_B T}{\partial r_\alpha} = R(q_\alpha). \quad (9.60)$$

To render this equation suitable for a further treatment, we use a simple BGK-like approximation (see Exercise 9.5) for the collision integral. Then

$$R(q_\alpha) = -q_\alpha/\tau,$$

where τ is some characteristic relaxation time such as the electron-electron mean collisional time (for more detail see derivation of formula (6.85) in Golant et al. 1977):

$$\tau \approx \frac{1}{2} \frac{m_e^{1/2} (k_B T_e)^{3/2}}{\pi^{1/2} e^4 n_e \ln \Lambda},$$

cf. (8.38). Some times, especially if the actual physical conditions are not very clear and should be clarified on the basis of numerical experiments, it is convenient to consider the characteristic time τ as an adjustable parameter.

Anyway the simplified flux equation is

$$\frac{\partial q_\alpha}{\partial t} + \frac{5}{2} \frac{p k_B}{m} \frac{\partial T}{\partial r_\alpha} + \frac{q_\alpha}{\tau} = 0. \quad (9.61)$$

Several remarks can be made about (9.61).

- (a) In the absence of macroscopic spatial gradients, the heat flux decays from an initial condition to zero with a decay rate of $1/\tau$. This is plausible because

█ Coulomb collisions isotropize the electron distribution function and relax it to a local Maxwellian distribution

on this time scale, and the heat flux is exponentially decreasing to zero. This also indicates a principal difference between (1) the equations for density, fluid velocity, and temperature and (2) the equations for the pressure tensor and heat flux. Since the first three equations are the result of moments using collisional invariants, no collision term explicitly appears in them (see also Sect. 12.1.3). For this reason, their time response is governed by macroscopic gradients and time scales. The electron heat flux equation contains a collision term, however. So the time dependent heat flux is described by this equation on time scales comparable to an electron mean free time. Hence,

█ by including the time dependent heat flux in the functional dependence of the temperature distribution we include the short time scale response of the electron distribution function.

- (b) For steady state, the heat flux equation reduces to Fourier's law

$$q_\alpha(\mathbf{r}, t) = -\frac{5}{2} \frac{p k_B \tau}{m} \frac{\partial T}{\partial r_\alpha}. \quad (9.62)$$

So, a thermal conductivity can be defined as

$$\kappa = \frac{5}{2} P k_B \tau / m, \quad (9.63)$$


which is the same as (9.51) of course.

* * *

Our contention is that the thirteen moment equations can be used, in their fully time dependent form, to better describe the time evolution of the plasma through their additional state variables.

█ The thirteen moment equations play a significant role in breaching the gap between hydrodynamics and kinetic theory in the problem of non-stationary powerful heat fluxes

Fig. 9.4 Temperature distributions over the column depth ξ along a reconnected magnetic-flux tube at consecutive times calculated for a heat flux with (*black curves*) and without (*gray curves*) account of the collisional relaxation effect. The numbers near the curves indicate the time in sec; the heating time $t_h = 7$ s



in non-equilibrium astrophysical plasmas, for example, in solar flares. In particular, it has appeared that the heat flux with consideration for collisional relaxation describes the heat transfer in solar flares much better than the classical Fourier law. This will be discussed in the next section.

However the question of accurate evaluating the thermal conductivity still remains in this analysis; and more elaborate methods than the simple BGK model have been used (starting presumably from [Salat 1975](#)) to approximate the thermal conductivity.

9.7.3 Large-Amplitude Heat Waves in Solar Flares

During a solar flare, a *reconnecting current layer* (RCL) in the solar corona becomes turbulent and super-hot. The electron temperature in it reaches huge values: $T_e \gtrsim 10^8$ K ([Somov 1981](#)). We call such a RCL the *super-hot turbulent-current layer* (SHTCL, see Part II, Chap. 8).

Let us imagine that two thin magnetic-flux tubes reconnect at the center of the SHTCL and move inside it toward its edges, while the plasma electrons in the tubes are heated to its temperature. Let t_h be this heating time. Having emerged from the SHTCL, the tubes of reconnected magnetic-field lines continue to move in the corona, being already disconnected from the heating source. However, at the same time, the redistribution of thermal energy continues in the form of a large-amplitude heat wave traveling along the magnetic-flux tubes. They are illustrated by Fig. 9.4.

In the case of classical heat conduction, the heat flux due to Coulomb collisions between electrons is described by the classical Fourier law (9.62). We rewrite (9.57) with condition $\mathbf{u} = 0$ in the following simple form:

$$\frac{\partial T}{\partial t} = A_{cl} \frac{\partial}{\partial l} \left(T^\alpha \frac{\partial T}{\partial l} \right), \quad (9.64)$$

where

$$A_{\text{cl}} = \frac{2\kappa_0}{3n_e k_B} = \text{const}, \quad \kappa_0 = \frac{1.84 \times 10^{-5}}{\ln \Lambda}, \quad \alpha = 5/2.$$

Since the plasma temperature T_0 in a magnetic-flux tube before its contact with the CHTCL is negligible compared to the CHTCL temperature T_s , we put $T_0 = 0$. As a consequence, the solution of a self-similar problem can be reduced to integrating ordinary differential equations (Oreshina and Somov 2011a).

However, in this approximation, the heat wave front moves too fast at the initial times; its mean velocity is greater than the speed of light. This unacceptable property follows from the self-similar dependence: the front velocity is formally proportional to $t^{-1/2}$; see Chap. 10 in Zel'dovich and Raizer (1966, 2002). Therefore it tends to infinity as $t \rightarrow 0$.

In fact, the contradiction mentioned above stems from the fact that the well-known classical formulae (9.62) and (9.63) are applicable for a plasma in a state *very close* to the state of local thermodynamic equilibrium. The latter suggests that the characteristic time scale t_T exceeds greatly the electron collision time τ_{ee} , and the characteristic length scale l_T , on which the temperature changes by a factor e , is much larger than the mean free path for thermal electrons λ_{ee} :

$$t_T = \frac{T}{|dT/dt|} \gg \tau_{ee}, \quad l_T = \frac{T}{|dT/dl|} \gg \lambda_{ee}. \quad (9.65)$$

However the temperature profiles (gray in Fig. 9.4) formally calculated by using Fourier's collisional law do not satisfy conditions (9.65).

Recall that, apart from conditions (9.65), the Fourier law implies that the classical heat flux q_{cl} must be less than the anomalous heat flux q_{an} and the saturated heat flux q_{sat} :

$$q_{\text{cl}} < \min(q_{\text{an}}, q_{\text{sat}}); \quad (9.66)$$

see, e.g., Somov (1992), Sect. 2.2.2. Here

$$q_{\text{an}} = \frac{n_e (k_B T_e)^{3/2}}{m_i^{1/2}}, \quad (9.67)$$

if a plasma turbulence is attributed to the ion-acoustic waves generated by a reverse electric current, and if the ratio $T_e/T_i \rightarrow \infty$. The saturated heat flux is the maximum possible heat flux because the diffusive heat transport cannot be greater than the convective one with a characteristic velocity V_{cr} :

$$q_{\text{sat}} = \frac{3}{2} n_e k_B T_e V_{\text{cr}}, \quad (9.68)$$

where

$$V_{\text{cr}} = V_{T_e} = \left(\frac{3k_B T_e}{m_e} \right)^{1/2}.$$

It appears (see [Oreshina and Somov 2011a](#)) that the heat flux calculated in terms of the self-similar model using the classical Fourier law is greater than the anomalous flux and even the saturated one. Therefore, all the necessary conditions for applicability of the classical heat conduction in the vicinity of SHTCLs in solar flares are violated.

Fortunately, allowance for the effect of collisional flux relaxation (as discussed in Sect. 9.7.2) removes both contradictions in the theory of solar flares. The heat wave shape (see Fig. 9.4) changes significantly. The wave front moves along a magnetic-flux tube more slowly; and the heat flux no longer exceeds the saturated flux. Thus we may conclude that

the heat flux with collisional relaxation describes the thermal energy transport in solar flares much better than the classical and anomalous heat fluxes.

However the heat flux still remains larger than the anomalous one calculated by using the asymptotic formula (9.67). This implies that, in general, one might expect the generation of particular waves, primarily at the heat wave front, for the temperature distributions similar to that shown in Fig. 9.4 (black), i.e. the distributions calculated by taking into account the collisional relaxation. In principle, such waves can turbulize the front but they cannot completely suppress the *thermal runaway* of fast electrons from a super-hot plasma and the so-called *nonlocal heat conduction* related to it ([Diakonov and Somov 1988](#)).

9.8 Gravitational Systems

There is a big difference between astrophysical plasmas and astrophysical gravitational systems (Sect. 3.3). The gravitational attraction cannot be screened. A large-scale gravitational field always exists over a system. This follows from the formula (3.17) which shows that the *averaged* gravitational force cannot be equal to zero because the neutrality condition (3.18) cannot be satisfied if all the particles have the same charge sign.

The large-scale gravitational field makes an overall thermodynamic equilibrium impossible. On the contrary, the electric force in a plasma is well screened beyond the Debye-Hückel radius and does not come in the way of the plasma having a proper thermodynamic equilibrium. Therefore, as one might have anticipated,

those results of plasma astrophysics which explicitly depend upon the plasma being in thermodynamic equilibrium do not hold for gravitational systems.

For gravitational systems, like the stars in a galaxy, we may hope that the observed distribution function reflects something about the *initial conditions* rather than just reflecting the relaxation mechanism. The random motions of the stars

may be not only non-Maxwellian but even direction dependent within the system. So galaxies may be providing us with clues on how they were formed (Palmer 1994; Bertin 1999; Peacock 1999).

If we assume that the stars form a collisionless system (see, however, Sect. 3.3), they do not exert pressure. Such a pressureless gravitating system is unstable (Jean's instability). Presumably a real galaxy should possess something akin to pressure to withstand the collapsing action of its gravity. This “pressure” is associated with the random motion of stars. So the role of sound speed is assumed to be played by the root mean velocity of the stars.

Another justification for treating a galaxy in the *hydrodynamic* approximation is that we consider processes on a spatial scale which is large enough to contain a large number of stars—one of the main requirements of the continuum mechanics. Anyway, several aspects of the structure of a galaxy can be understood by assuming that it is made up of a continuum medium (see, e.g., Alfaro et al. 2004). Often

hydrodynamics provides a first level description of an astrophysical phenomenon governed predominantly by the gravitational force.

Magnetic fields are usually included later on in order to address additional issues. For example, the early stages of star formation during which an *interstellar cloud* of low density collapses under the action of its own gravity can be modeled in the hydrodynamic approximation. However, when we want to explain the difference between the angular momentum of the cloud and that of the born star, we have to include the effect of a magnetic field.

9.9 Practice: Exercises and Answers

Exercise 9.1 (Sect. 9.3.1). Show that the third integral in (9.12) equals zero.

Answer. Let us find the derivative

$$\frac{\partial}{\partial v_\alpha} \left(\frac{F_{k,\alpha}}{m_k} f_k \right) = \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} + \frac{f_k}{m_k} \frac{\partial F_{k,\alpha}}{\partial v_\alpha} = \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha}.$$

The condition (1.8) has been used on the right-hand side as the condition

$$\frac{\partial F_{k,\alpha}}{\partial v_\alpha} = 0. \quad (9.69)$$

Hence

$$\int_v \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} d^3 \mathbf{v} = \frac{F_k}{m_k} f_k(\mathbf{r}, \mathbf{v}, t) \Big|_{\mathbf{v} \rightarrow -\infty}^{\mathbf{v} \rightarrow +\infty} = 0,$$

if the distribution function f_k quickly approaches zero as $v \rightarrow \infty$; q.e.d.

Exercise 9.2 (Sect. 9.3.1). Write the continuity equation with account of ionization and recombination.

Answer. The continuity equation including the source/sink terms related to ionization/recombination or charge exchange reads

$$\frac{\partial n_k}{\partial t} + \frac{\partial}{\partial r_\alpha} n_k u_{k,\alpha} = \sum_l (\gamma_{lk} n_l - \gamma_{kl} n_k). \quad (9.70)$$

Here n_k denotes the particle density of species k , either neutral or ionized. The right-hand side of the equation is the change of n_k due to collisions. The coefficients γ_{kl} and γ_{lk} denote the rate of transformation of species k into species l and vice versa. These rates must obey the relation

$$\sum_k \sum_l (\gamma_{lk} n_l - \gamma_{kl} n_k) = 0, \quad (9.71)$$

which ensures the total particle number density conservation.

Exercise 9.3 (Sect. 9.3.2). Consider the third integral in the first moment (9.16).

Answer. Let us find the derivative

$$\begin{aligned} \frac{\partial}{\partial v_\beta} (v_\alpha F_{k,\beta} f_k) &= v_\alpha F_{k,\beta} \frac{\partial f_k}{\partial v_\beta} + v_\alpha \frac{\partial F_{k,\beta}}{\partial v_\beta} f_k + F_{k,\beta} f_k \frac{\partial v_\alpha}{\partial v_\beta} = \\ &= v_\alpha F_{k,\beta} \frac{\partial f_k}{\partial v_\beta} + 0 + F_{k,\beta} f_k \delta_{\alpha\beta}. \end{aligned} \quad (9.72)$$

The condition (1.8) has been used on the right-hand side as the condition

$$\frac{\partial F_{k,\beta}}{\partial v_\beta} = 0. \quad (9.73)$$

It follows from (9.72) that

$$v_\alpha F_{k,\beta} \frac{\partial f_k}{\partial v_\beta} = \frac{\partial}{\partial v_\beta} (v_\alpha F_{k,\beta} f_k) - F_{k,\alpha} f_k.$$

Thus

$$\int_{\mathbf{v}} v_\alpha F_{k,\beta} \frac{\partial f_k}{\partial v_\beta} d^3\mathbf{v} = v_\alpha F_k f_k \Big|_{\mathbf{v} \rightarrow -\infty}^{\mathbf{v} \rightarrow +\infty} - \int_{\mathbf{v}} F_{k,\alpha} f_k d^3\mathbf{v}. \quad (9.74)$$

The first term on the right-hand side equals zero, if the distribution function f_k quickly approaches zero as $v \rightarrow \infty$. Therefore, for the mean force acting on the particles of kind k in a unit volume, formula (9.18) has finally arrived.

Exercise 9.4 (Sect. 9.3.2). Find a condition under which the mean collisional force (9.20) is determined only by random motions of the particles of kind k .

Answer. In definition (9.20), let us take into account that

$$v_\alpha = u_{k,\alpha} + v'_\alpha.$$

Thus we obtain

$$\langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = m_k u_{k,\alpha} \int_{\mathbf{v}} \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v} + m_k \int_{\mathbf{v}} v'_\alpha \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v}. \quad (9.75)$$

The first integral on the right-hand side equals zero if condition (9.13) is satisfied. The remaining part

$$\langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = m_k \int_{\mathbf{v}} v'_\alpha \left(\frac{\partial \hat{f}_k}{\partial t} \right)_c d^3\mathbf{v}. \quad (9.76)$$

Thus the average transfer of momentum from the particles of kind k to the particles of other kinds is solely due to the random motions of the particles of kind k if the processes of transformation, during which the particle kind can be changed, are not allowed for.

Exercise 9.5 (Sect. 9.6). Let us approximate the collisional integral (9.3) by the following simple form (Bhatnagar et al. (1954)):

$$\left(\frac{\partial \hat{f}_k}{\partial t} \right)_c = - \frac{f_k(\mathbf{r}, \mathbf{v}, t) - f_k^{(0)}(\mathbf{r}, \mathbf{v}, t)}{\tau_c}, \quad (9.77)$$

where an arbitrary distribution function $f_k(\mathbf{r}, \mathbf{v}, t)$ relaxes to the Maxwellian distribution function $f_k^{(0)}(\mathbf{r}, \mathbf{v}, t)$, as discussed in Sect. 9.6, in a collisional time τ_c . Discuss why this simple approximation illuminates much of the basic physics of transfer phenomena in a relatively less-painful way for neutral gases but is not very reliable for plasmas, especially in the presence of magnetic fields.

Comment. The departure of the distribution function from the pure Maxwellian one, the function

$$f_k^{(1)}(\mathbf{r}, \mathbf{v}, t) = f_k(\mathbf{r}, \mathbf{v}, t) - f_k^{(0)}(\mathbf{r}, \mathbf{v}, t) \quad (9.78)$$

satisfies the following equation:

$$\frac{\partial f_k}{\partial t} + v_\alpha \frac{\partial f_k}{\partial r_\alpha} + \frac{F_{k,\alpha}}{m_k} \frac{\partial f_k}{\partial v_\alpha} = - \frac{f_k^{(1)}}{\tau_c}, \quad (9.79)$$

which is called the BGK (Bhatnagar, Gross and Krook) equation.

If a gradient in space, $\partial/\partial r_\alpha$, gives rise to the departure from the Maxwellian distribution, then in order to have a rough estimate of the effect, we may balance the second term on the left-hand side of (9.79) with its right-hand side:

$$\frac{|v_\alpha| f_k^{(0)}}{L} \approx \frac{|f_k^{(1)}|}{\tau_c}. \quad (9.80)$$

Here $|v_\alpha|$ is the typical velocity of the particles of kind k , L is the typical length scale over which properties of the system change appreciably. From (9.80) it follows that

$$\frac{|f_k^{(1)}|}{f_k^{(0)}} \approx \frac{\lambda_c}{L}. \quad (9.81)$$

Thus the departure from the Maxwellian distribution will be small if the mean free path λ_c is small compared to the typical length scale. This is consistent with the second condition of (9.47).

Chapter 10

Multi-Fluid Models of Astrophysical Plasma

Abstract The multi-fluid models of plasma in electric and magnetic fields allow us to consider many important properties of astrophysical plasma, in particular the Langmuir and electromagnetic waves, as well as many other interesting effects and applications.

10.1 Multi-fluid Models in Astrophysics

The transfer equations (9.14), (9.17), and (9.25) give us the hydrodynamic-type description of multi-component astrophysical plasma in electric and magnetic fields. The problem is that, if we would like to solve the equations for one of the plasma components, we could not escape solving the transfer equations for all of the components since they depend on each other and on the electric and magnetic fields. For this reason, we should minimize the number of plasma components under consideration.

The “*two-fluid*” hydrodynamic-type equations are often used to describe the flow of the electrons and protons of a fully-ionized astrophysical plasma under the action of an electric and magnetic fields. Such treatment yields, for example, the generalized Ohm’s law in astrophysical plasma (Chap. 11) as well as a *dynamic friction* force which maximizes when the relative drift velocity is equal to the sum of the most probable random velocities of the electrons and ions. For relative drift velocities in excess of this value, the friction force decreases rapidly. The electron and ion currents flowing parallel to the existing magnetic fields increase steadily in time, i.e. runaway (Dreicer 1959; see also Sect. 8.4).

The “*multi-fluid*” models are useful, for example, to explore properties of the solar wind (e.g., Bodmer and Bochsler 2000). The electrons, protons, and alpha particles in the solar wind constitute the main three components, while the less abundant elements and isotopes are treated as test species. To model the main gases, we have to study solutions for the conservation-law equations of the three components. The behaviour of minor ions depends in a complicated manner on

their mass and on their charge, structured by the interplay of acceleration, gravity, pressure gradient, electromagnetic fields, Coulomb friction force, and thermal diffusion. Such models allow one to explore the efficiency of isotope fractionation processes in the solar corona.

10.2 Langmuir Waves

Because a plasma consists of at least two components (electrons and ions), the number of possible waves is larger than in a normal gas, where *acoustic* and *entropy* waves are the two possible types of waves. In this section we shall discuss the *simplest* waves in astrophysical plasma, whose properties can be deduced from the hydrodynamic-type equations for two mutually penetrating charged gases (Sect. 9.5).

Although astrophysical plasma is almost always magnetized, we can quite often neglect the magnetic field in discussing small-amplitude plasma waves; the condition will become clear later. The reduced complexity of the governing equations can be further simplified by approximations.

10.2.1 Langmuir Waves in a Cold Plasma

Let us assume that the ions do not move at all (they are infinitely massive) and they are uniformly distributed in space. So the ions have a fixed number density n_0 . This is a cold ion approximation.

Let us also neglect all magnetic fields. We shall assume that any variations of electron density n_e , electron velocity u_e , and related electric field $\mathbf{E}$ occur only in one dimension—the x axis. Then we are left with a set of three equations:

- (a) The continuity equation (9.14) for electrons

$$\frac{\partial n_e}{\partial t} + \frac{\partial}{\partial x} n_e u_e = 0, \quad (10.1)$$

- (b) The motion equation (9.44)

$$m_e n_e \left(\frac{\partial u_e}{\partial t} + u_e \frac{\partial u_e}{\partial x} \right) = - \frac{\partial p_e}{\partial x} - e n_e E_x, \quad (10.2)$$

- (c) The electric field equation

$$\frac{\partial E_x}{\partial x} = 4\pi e (n_0 - n_e). \quad (10.3)$$

In general, we cannot solve these nonlinear equations exactly, except for very special cases. One of them is trivial:

$$n_e = n_0 = \text{const}, \quad u_e = 0, \quad p_e = p_0 = \text{const}, \quad E_x = 0. \quad (10.4)$$

This solution corresponds to a *stationary* electron gas of uniform density and temperature.

Let us linearize (10.1)–(10.3) with respect to the state (10.4). This yields the following set of *linear* equations:

$$\frac{\partial n_1}{\partial t} + n_0 \frac{\partial u_1}{\partial x} = 0, \quad (10.5)$$

$$m_e n_0 \frac{\partial u_1}{\partial t} = - \frac{\partial p_1}{\partial x} - e n_e E_1, \quad (10.6)$$

$$\frac{\partial E_1}{\partial x} = 4\pi e n_1. \quad (10.7)$$

Let us consider the special case of *cold* electrons: $T_e = 0$ and, therefore,

$$p_e = 0. \quad (10.8)$$

Now we eliminate u_1 and E_1 from the set of equation by taking the time derivative of (10.5) to obtain the oscillator equation

$$\frac{\partial^2 n_1}{\partial t^2} + \left(\frac{4\pi e^2 n_0}{m_e} \right) n_1 = 0. \quad (10.9)$$

If we displace some electrons to produce an initial perturbation, we create a positive-charge density at the position where they started. This positive-charge perturbation attracts the electrons, which will tend to move back to their original position, but will overshoot it. They come back again, overshoot it, and so on. Without any damping, the energy put into the plasma to create the perturbation will remain in the plasma. So the oscillation will continue forever with the frequency

$$\omega_{pl}^{(e)} = \pm \left(\frac{4\pi e^2 n_e}{m_e} \right)^{1/2}$$

(10.10)

called the *electron plasma frequency*.

Therefore, in a two-component cold plasma, there exist the *oscillations of charge density*—*Langmuir waves* which frequency is independent of the wave vector $\mathbf{k}$; so the group velocity, $\mathbf{V}_{gr} = d\omega/d\mathbf{k}$, is zero. Thus

in a cold plasma, Langmuir waves are spatially localized oscillations of electric charge density which do not propagate at all.

Note that there is no equivalent to these oscillations in gasdynamics or gravitational dynamics, for which there is no charge separation and related electric-type force.

10.2.2 Langmuir Waves in a Warm Plasma

What happens with behavior of a Langmuir wave, if the electron temperature is not equal to zero?—Let us drop the assumption (10.8) of zero pressure in the linear equations (10.5)–(10.7). We then must include the perturbation of electron pressure

$$\frac{\partial p_1}{\partial x} = n_0 k_B \frac{\partial T_1}{\partial x} + k_B T_0 \frac{\partial n_1}{\partial x} \quad (10.11)$$

in (10.6).

Now we must relate n_1 to T_1 and vice versa. For example, we could argue that for long-wavelength waves the compression is the one-dimensional ($N = 1$) adiabatic process with the index $\gamma = (N + 2)/N = 3$. In this case, the perturbation of electron pressure becomes

$$\frac{\partial p_1}{\partial x} = 3k_B T_0 \frac{\partial n_1}{\partial x}. \quad (10.12)$$

Naturally we expect now an initial perturbation to propagate through the plasma as a wave. Thus a plane-wave solution of the form

$$f_1(x, t) = \tilde{f}_1 \exp[-i(\omega t - kx)] \quad (10.13)$$


should satisfy the linear differential equations. The quantities with tildes are the complex amplitudes. They obey three linear algebraic equations:

$$\begin{aligned} -i\omega \tilde{n}_1 + ik n_0 \tilde{u}_1 &= 0, \\ -i\omega m_e n_0 \tilde{u}_1 + ik 3k_B T_0 \tilde{n}_1 + e n_0 \tilde{E}_1 &= 0, \\ ik \tilde{E}_1 + 4\pi e \tilde{n}_1 &= 0. \end{aligned}$$

To have a nontrivial solution, the determinant must be zero:

$$\det \begin{vmatrix} -i\omega & ik n_0 & 0 \\ ik 3k_B T_0 (m_e n_0)^{-1} & -i\omega e m_e^{-1} & \\ 4\pi e & 0 & ik \end{vmatrix} = 0.$$

Fig. 10.1 A dispersion diagram (*solid curves*) for Langmuir waves in a warm plasma. The ions do not move at all. *Dashed straight lines* are drawn for Langmuir waves in a cold plasma



Its solution is

$$\omega = \pm \omega_{pl}^{(e)} (1 + 3 r_{DH}^2 k^2)^{1/2}, \quad (10.14)$$

where

$$r_{DH} = \frac{1}{\sqrt{3}} \frac{V_{Te}}{\omega_{pl}^{(e)}} \quad (10.15)$$

is the electron Debye-Hückel radius, V_{Te} is the mean thermal velocity (8.15) of electrons in a plasma.

The dispersion equation (10.14) can be also derived from the Vlasov equation, of course (see formula (49) in Vlasov 1938). It is similar to the well-known relation for the propagation of transverse electromagnetic waves in a vacuum, except that the role of the light velocity c is here played by the thermal velocity V_{Te} . This dispersion relation is shown in Fig. 10.1.

Therefore the frequency ω of Langmuir waves in a plasma with warm electrons depends on the wave vector $\mathbf{k}$ which is parallel to the x -axis. So


the group velocity, $\partial \omega / \partial k$, of Langmuir waves in a warm plasma without magnetic field is not equal to zero.

They oscillate at the electron plasma frequency $\omega_{pl}^{(e)}$ and propagate in a warm plasma. It follows from (10.14) and (10.15) that the group velocity is

$$V_{gr} = \frac{\partial \omega}{\partial k} = V_{Te}^2 \frac{k}{\omega} = \frac{3k_B T}{m_e} \frac{k}{\omega}. \quad (10.16)$$

Therefore the plasma waves are propagating as long as the electron temperature is non-zero. Moreover, due to the small mass of the electrons, the group velocity (10.16) is always relatively large.

Fig. 10.2 A dispersion diagram for ion-acoustic waves (part *a*) and for ion plasma waves (part *b*) in a warm plasma without magnetic field



10.2.3 Ion Effects in Langmuir Waves

Let us show that, when the ions are allowed to move, ion contributions are important only for slow variations or *low-frequency* waves because the ions cannot react quickly enough.

We are still dealing with linear waves which involve only the first-order electric field $\mathbf{E}^{(1)}$ directed along the wave vector $\mathbf{k}$ which is parallel to the x -axis. Linearizing the continuity equations for electrons and ions, the motion equations for electrons and ions, as well as the electric field equation, let us assume that the electrons and ions both obey the adiabatic equation (10.12). Then we again use the wave solution (10.13) to reduce the linearized differential equations to algebraic ones and to obtain the determinant. Because $m_i/m_e \gg 1$, we neglect the term $m_e \omega^2$ in this determinant as compared with the term $m_i \omega^2$. By so doing, we obtain the relation

$$\omega^2 = k^2 \left(\frac{\gamma_i k_B T_i}{m_i} + \frac{\gamma_e k_B T_e}{m_i} \frac{1}{1 + \gamma_e k^2 r_D^2} \right). \quad (10.17)$$

This dispersion relation is shown in Fig. 10.2.

In the limit of small kr_D

$$\omega^2 = k^2 \left(\frac{\gamma_i k_B T_i}{m_i} + \frac{\gamma_e k_B T_e}{m_i} \right) = k^2 V_{ia}^2. \quad (10.18)$$

This is the so-called *ion-acoustic waves*. They are shown by a curve part (*a*) in Fig. 10.2. The group velocity of the wave is independent of k :

$$V_{gr} = \frac{\partial \omega}{\partial k} = V_{ia} = \left(\frac{\gamma_i k_B T_i + \gamma_e k_B T_e}{m_i} \right)^{1/2}. \quad (10.19)$$

An opposite limit is obtained for cold ions. If ion temperature $T_i \rightarrow 0$, then $kr_D \gg 1$, i.e., short wavelengths are under consideration. In this case, shown by the curve part (*b*) in Fig. 10.2, the cold ions oscillate with a frequency

$$\omega_{pl}^{(i)} = \pm \left(\frac{4\pi e^2 n_e}{m_i} \right)^{1/2} \quad (10.20)$$

called the *ion* plasma frequency.

Ion-acoustic waves are observed in many cases. They were registered, for example, by the spacecraft *Voyager 1* in the upstream side of the Jovian bow shock. Ion-acoustic waves presumably play an important role in solar flares, for example, in super-hot turbulent-current layers (see Part II, Sect. 6.3).

10.3 Electromagnetic Waves in Plasma

In this section we still assume that the unperturbed plasma has no magnetic field: $\mathbf{B}^{(0)} = 0$. However we shall discuss the waves that carry not only an electric field $\mathbf{E}^{(1)}$ but also a magnetic field $\mathbf{B}^{(1)}$.

Let us consider transversal waves, so that $\mathbf{k} \cdot \mathbf{E}^{(1)} = 0$ and $\mathbf{k} \cdot \mathbf{B}^{(1)} = 0$. The last equality is imposed by (1.29) and is always true. We do not need (1.30) in this case either. We shall neglect the ion motion, which is justified for high-frequency waves. So the remaining equations in their linearized form are

$$\frac{\partial \mathbf{u}_e^{(1)}}{\partial t} = -\nabla p_e^{(1)} - e n_e^{(0)} \mathbf{E}^{(1)}, \quad (10.21)$$

$$\frac{\partial n_e^{(1)}}{\partial t} + n_e^{(0)} \operatorname{div} \mathbf{u}_e^{(1)} = 0, \quad (10.22)$$

$$\operatorname{curl} \mathbf{B}^{(1)} = \frac{4\pi}{c} \mathbf{j}^{(1)} + \frac{1}{c} \frac{\partial \mathbf{E}^{(1)}}{\partial t}, \quad (10.23)$$

$$\operatorname{curl} \mathbf{E}^{(1)} = -\frac{1}{c} \frac{\partial \mathbf{B}^{(1)}}{\partial t}, \quad (10.24)$$

$$\mathbf{j}^{(1)} = e n_e^{(0)} \mathbf{u}_e^{(1)}. \quad (10.25)$$


The Lorentz force does not appear in the electron-motion equation (10.21) because it is of the second-order small value proportional to $\mathbf{u}_e^{(1)} \times \mathbf{B}^{(1)}$. Furthermore vectors $\mathbf{E}^{(1)}$ and $\mathbf{u}_e^{(1)}$ are perpendicular to the wave vector $\mathbf{k}$, and thus

$$n_e^{(1)} = 0 \quad \text{and} \quad p_e^{(1)} = 0. \quad (10.26)$$

After assuming the exponential plane-wave form (10.13) and using usual algebra, we find the dispersion equation for electromagnetic waves:

$$\omega^2 = \omega_{pl}^{(e)2} + k^2 c^2. \quad (10.27)$$

Fig. 10.3 The dispersion diagram for electromagnetic waves in a cold plasma without magnetic field. For large values of k (*short wavelengths*), the group velocity (the slope of the *solid curve*) and phase velocity approach the speed of the light (*dashed line*). For small values of k (*long wavelengths*), the group velocity goes to zero



Here c is the speed of light in a vacuum. This dispersion relation is shown in Fig. 10.3.

If the wave frequency ω is much larger the electron plasma frequency $\omega_{pl}^{(e)}$, the wave becomes a free-space light wave with $\omega = kc$.

If $\omega \rightarrow \omega_{pl}^{(e)}$, a wave would decay in space and not propagate. In this case, the index of refraction

$$n_r = \frac{c}{V_{ph}} = \frac{ck}{\omega} = \left(1 - \frac{\omega_{pl}^{(e)2}}{\omega^2}\right)^{1/2} \quad (10.28)$$

goes to zero. If $\omega < \omega_{pl}^{(e)}$, the refraction index becomes imaginary.

Moving through astrophysical plasma of changing $\omega_{pl}^{(e)}$, a wave is reflected when $\omega = \omega_{pl}^{(e)}$ and, therefore, $n_r = 0$. This allows one to measure remotely the plasma density, for example, in the Earth ionosphere.

Another application is in ionospheric heating. At the height where $\omega_{pl}^{(e)}$ is equal to the wave frequency, the group velocity

$$V_{gr} = \frac{\partial \omega}{\partial k} = \frac{kc^2}{\omega} = n_r c \quad (10.29)$$

also goes to zero.

The wave amplitude becomes large near a reflecting point because its flux of energy cannot propagate at this point.

The large electric field of the wave can accelerate electrons and drive currents in the ionospheric plasma. In this way, the wave can heat and modify the plasma. If the power from a transmitter on the ground emitting a radiation at a frequency ω is large enough, the heating is quite significant.

10.4 What Do We Miss?

We have considered two basic types of waves in a two-fluid plasma. The Langmuir wave or plasma wave (as well as the ion plasma wave) does not have a *wave magnetic field* $\mathbf{B}^{(1)}$. The electromagnetic wave does have a magnetic field $\mathbf{B}^{(1)}$ but can propagate only if its frequency is above the plasma frequency:

$$\omega > \omega_{pl}^{(e)}. \quad (10.30)$$

We should see that, when there is a *stationary magnetic field* $\mathbf{B}^{(0)}$ in the plasma, the wave properties become more complex and more interesting (e.g., [Stix 1992](#); [Zheleznyakov 1996](#)).

In particular, we could find that

an electromagnetic wave with its frequency below the plasma frequency can propagate through a magnetized plasma.

For low-frequency waves this effect will be demonstrated in the magnetohydrodynamic (MHD) approximation in Chap. 15. What else has been lost in the above consideration?

The advantage of the hydrodynamic approach used in this section to study the basic properties of waves in a two-fluid plasma is the relative simplicity. The hydrodynamic-type equations have three spatial dimensions and time, rather than the seven-dimensional phase space of the Vlasov kinetic theory (Sect. 3.1.2).

The obvious disadvantage is that some *fine effects*, such as Landau damping (Sect. 7.1.2) which is caused by a resonance with particles moving at the phase velocity of a wave, cannot be obtained from the hydrodynamic-type equations. We have to use a kinetic treatment to specify how a distribution of particles responds to a wave. In this case we use the Vlasov equation to specify how the distribution functions of electrons and ions are affected by the wave fields (e.g., Chap. 7).

In order to calculate the collisional damping of plasma waves simply, the simplest hydrodynamic model is useful (Exercise 10.3).

The hydrodynamic-type models work well only when a finite number of the low-order moments are sufficient to provide all the essential information about the system.

If the distribution function has some unusual features, then a few low-order moments may not carry all the necessary information,

and we may lose important physics by restricting ourselves to the quasi-hydrodynamic description of astrophysical plasma.

10.5 Practice: Exercises and Answers

Exercise 10.1. Show that in the solar corona a *dynamic viscosity* coefficient can be given by a simple formula (Hollweg 1986):

$$\eta \approx 10^{-16} T_p^{5/2}, \quad \text{g cm}^{-1} \text{ s}^{-1}, \quad (10.31)$$

where T_p is the proton temperature, and the Coulomb logarithm has been taken to be 22. So, with $T_p \approx 2 \times 10^6$ K, the viscosity coefficient in the corona

$$\eta \sim 1 \text{ g cm}^{-1} \text{ s}^{-1}.$$

Why does the viscosity grow with the proton temperature? Why is it so large and does it grow with temperature so quickly?

Hint. Consider a fully-ionized hydrogen plasma in a magnetic field. Let τ_{pp} represent the typical Coulomb collisional time (8.39) for thermal protons. Let $\omega_B^{(p)}$ denote the proton cyclotron frequency (5.52).

Write the viscous stress tensor (9.39) for the protons. This tensor involves five coefficients of viscosity, denoted $\eta_0, \eta_1, \dots, \eta_4$ by Braginskii (1965). Show that the coefficient η_0 is by far the largest one (10.31). The coefficients η_3 and η_4 are smaller by factors $\sim (\omega_B^{(p)} \tau_{pp})^{-1}$, while η_1 and η_2 are smaller than η_0 by factors $\sim (\omega_B^{(p)} \tau_{pp})^{-2}$.

Thus the parts of the viscous stress tensor involving the off-diagonal terms can often be neglected. The part involving η_0 can be dynamically and thermodynamically important.

Exercise 10.2. Discuss a famous puzzle of plasma astrophysics. Solar flares generate electron beams that move through the solar corona and the interplanetary space at velocities $\sim 0.3c$ (see Exercise 10.3). These fast beams should lose their energy quickly to plasma waves. In fact, they do generate waves called the solar type III radio bursts. However the solar fast electrons are still seen far beyond the orbit of the Earth. Why?

Hint. The link between the electron beams and the waves observed in space near the Earth or even on the ground is a little more complex than it seems. It involves the transformation of the electrostatic plasma oscillations with frequency near $\omega_{pl}^{(e)}$ into electromagnetic waves at the same frequency. In any realistic situation, the electrons in the beam are not cold but have a thermal spread. They cause a plasma wave to grow. But as the electric field in the wave grows, the electrons are heated.

The spreading and slowing of the beam in the velocity space cannot be described by fluid equations. This process is often referred to as *quasi-linear diffusion*. We can expect that the electron beam has slowed and spread in the velocity space to such a degree that waves do not grow anymore. A stable situation can occur, and a warm electron beam can propagate through the plasma without losing energy.

Exercise 10.3. Show that Coulomb collisions damp the Langmuir plasma waves with the rate

$$\operatorname{Im} \omega = -2 \nu_{ei} . \quad (10.32)$$

Hint. Following formula (9.24), add to the right-hand side of the electron motion equation (10.2) the collisional friction term

$$+ m_e n_e \nu_{ei} (u_{i,\alpha} - u_{e,\alpha}) .$$

Chapter 11

The Generalized Ohm's Law in Plasma

Abstract The multi-fluid models of the astrophysical plasma in magnetic field allow us to derive the generalized Ohm's law and to consider different physical approximations, including the collisional, rarely-collisional and collisionless models of plasma, as well as many interesting applications.

11.1 The Classic Ohm's Law

The usual Ohm's law, $\mathbf{j} = \sigma \mathbf{E}$, relates the current density $\mathbf{j}$ to the electric field $\mathbf{E}$ in a *solid conductor in rest*. The coefficient σ is electric conductivity. As we know, the electric field in every equation of motion in any approximation determines acceleration of charged particles rather than their velocity. That is why, generally, such a simple relation as the classic Ohm's law does not exist. Moreover, while considering astrophysical plasmas, it is necessary to take into account the presence of a magnetic field and the motion of a plasma as a whole or as a medium consisting of several moving components, their compressibility and viscosity, presence of gradients of temperature and density.

However, at first, let us recall the way of deriving the classic Ohm's law in a conductor or in a plasma without magnetic field. The electric current is determined by the relative motion of electrons and ions. Considering the processes in which all quantities vary only slightly in a time between the electron-ion collisions, electron inertia can be neglected. An equilibrium is set up between the electric field action and electrons-on-ions *friction* (see point *A* in Fig. 8.7). Let us assume that the ions do not move. Then the condition for this equilibrium with respect to the electron gas

$$0 = -e n_e E_\alpha + m_e n_e v_{ei} (0 - u_{e,\alpha})$$

results in Ohm's law

$$j_\alpha = -e n_e u_{e,\alpha} = + \frac{e^2 n_e}{m_e v_{ei}} E_\alpha = \sigma E_\alpha . \quad (11.1)$$

Here

$$\sigma = \frac{e^2 n_e}{m_e \nu_{ei}} \quad (11.2)$$

is the *electric conductivity*.

In order to deduce the generalized Ohm's law for a plasma with magnetic field, we have to consider at least two equations of motion—for the electron and ion components. Therefore a theory of conductivity in a fully-ionized plasma, considered in Sects. 11.2–11.5, can be given in terms of a *two-fluid* approximation.

A more general case, with motion of neutrals taken into account, has been considered by Schlüter (1951); Alfvén and Fälthammar (1963), Sect. 5.2.3.2. Different applications of the generalized Ohm's law in the *three-component* astrophysical plasma were considered by many authors (Schabansky 1971; Kunkel 1984; Hénoux and Somov 1991 and 1997; Alperovich and Fedorov 2007).

11.2 Derivation of Basic Equations

Let us write the momentum-transfer equations (9.17) for the electrons and ions, taking account of the pressure tensor (9.11), the Lorentz force (9.19) and the friction force (9.24). We have two following equations:

$$m_e \frac{\partial}{\partial t} (n_e u_{e,\alpha}) = - \frac{\partial \Pi_{\alpha\beta}^{(e)}}{\partial r_\beta} - e n_e \left[\mathbf{E} + \frac{1}{c} (\mathbf{u}_e \times \mathbf{B}) \right]_\alpha + m_e n_e \nu_{ei} (u_{i,\alpha} - u_{e,\alpha}), \quad (11.3)$$

$$m_i \frac{\partial}{\partial t} (n_i u_{i,\alpha}) = - \frac{\partial \Pi_{\alpha\beta}^{(i)}}{\partial r_\beta} + Z_i e n_i \left[\mathbf{E} + \frac{1}{c} (\mathbf{u}_i \times \mathbf{B}) \right]_\alpha + m_e n_e \nu_{ei} (u_{e,\alpha} - u_{i,\alpha}). \quad (11.4)$$

Here the tensor of momentum flux density for electrons and ions is given by formula (9.10):

$$\Pi_{\alpha\beta}^{(e)}(\mathbf{r}, t) = m_e n_e u_{e,\alpha} u_{e,\beta} + p_{\alpha\beta}^{(e)} \quad (11.5)$$

and

$$\Pi_{\alpha\beta}^{(i)}(\mathbf{r}, t) = m_i n_i u_{i,\alpha} u_{i,\beta} + p_{\alpha\beta}^{(i)}. \quad (11.6)$$

The last term in (11.3) represents the mean momentum transferred, because of collisions (see formula (9.24)), between electrons and ions. It is equal, with opposite sign, to the last term in (11.4). It is assumed that there are just two kinds of particles, their total momentum remaining constant under the action of elastic collisions.

Now let us suppose that the ions are protons ($Z_i = 1$) and electrical *neutrality* is observed:

$$n_i = n_e = n. \quad (11.7)$$

Let us multiply (11.3) by $-e/m_e$ and add it to (11.4) multiplied by e/m_i . The result is

$$\begin{aligned} \frac{\partial}{\partial t} [en(u_{i,\alpha} - u_{e,\alpha})] &= \left[\frac{e}{m_i} F_{i,\alpha} - \frac{e}{m_e} F_{e,\alpha} \right] + \\ &+ e^2 n \left(\frac{1}{m_e} + \frac{1}{m_i} \right) E_\alpha \\ &+ \frac{e^2 n}{c} \left[\left(\frac{\mathbf{u}_e}{m_e} \times \mathbf{B} \right)_\alpha + \left(\frac{\mathbf{u}_i}{m_i} \times \mathbf{B} \right)_\alpha \right] - \\ &- v_{ei} en \left[(u_{i,\alpha} - u_{e,\alpha}) + \frac{m_e}{m_i} (u_{i,\alpha} - u_{e,\alpha}) \right]. \end{aligned} \quad (11.8)$$

Here

$$F_{e,\alpha} = -\frac{\partial \Pi_{\alpha\beta}^{(e)}}{\partial r_\beta} \quad \text{and} \quad F_{i,\alpha} = -\frac{\partial \Pi_{\alpha\beta}^{(i)}}{\partial r_\beta}. \quad (11.9)$$

Let us introduce the velocity of the centre-of-mass system

$$\mathbf{u} = \frac{m_i \mathbf{u}_i + m_e \mathbf{u}_e}{m_i + m_e}. \quad (11.10)$$

Since $m_i \gg m_e$,

$$\mathbf{u} = \mathbf{u}_i + \frac{m_e}{m_i} \mathbf{u}_e \approx \mathbf{u}_i. \quad (11.11)$$

On treating (11.8), we neglect the small terms of the order of the ratio m_e/m_i . On rearrangement, we obtain the equation for the current density

$$\mathbf{j} = en(\mathbf{u}_i - \mathbf{u}_e) \quad (11.12)$$

in the system of coordinates (11.11). This equation is

$$\begin{aligned} \frac{\partial \mathbf{j}'}{\partial t} &= \frac{e^2 n}{m_e} \left[\mathbf{E} + \frac{1}{c} (\mathbf{u} \times \mathbf{B}) \right] - \frac{e}{m_e c} (\mathbf{j}' \times \mathbf{B}) - \\ &- v_{ei} \mathbf{j}' + \frac{e}{m_i} \mathbf{F}_i - \frac{e}{m_e} \mathbf{F}_e. \end{aligned} \quad (11.13)$$

The prime designates the electric current in the system of moving plasma, i.e. in the rest-frame of the plasma. Let $\mathbf{E}_u$ denote the electric field in this frame of reference, i.e.

$$\mathbf{E}_u = \mathbf{E} + \frac{1}{c} \mathbf{u} \times \mathbf{B}. \quad (11.14)$$

Now we divide (11.13) by ν_{ei} and represent it in the form

$$\mathbf{j}' = \frac{e^2 n}{m_e \nu_{ei}} \mathbf{E}_u - \frac{\omega_B^{(e)}}{\nu_{ei}} \mathbf{j}' \times \mathbf{n} - \frac{1}{\nu_{ei}} \frac{\partial \mathbf{j}'}{\partial t} + \frac{1}{\nu_{ei}} \left(\frac{e}{m_i} \mathbf{F}_i - \frac{e}{m_e} \mathbf{F}_e \right). \quad (11.15)$$

Here the unit vector $\mathbf{n} = \mathbf{B}/B$ and

$$\omega_B^{(e)} = \frac{eB}{m_e c} \quad (11.16)$$

is the electron gyro-frequency (Sect. 5.1.3).

Thus we have derived a differential equation for the current density $\mathbf{j}'$. This equation is interesting as it is (see Exercise 11.1). However we shall use it in another form.

The third and the fourth terms on the right in (11.15) do not depend of magnetic field. Let us replace them by some *effective* electric field such that

$$\sigma \mathbf{E}_{\text{eff}} = -\frac{1}{\nu_{ei}} \frac{\partial \mathbf{j}'}{\partial t} + \frac{e}{\nu_{ei}} \left(\frac{1}{m_i} \mathbf{F}_i - \frac{1}{m_e} \mathbf{F}_e \right), \quad (11.17)$$

where

$$\boxed{\sigma = \frac{e^2 n}{m_e \nu_{ei}}} \quad (11.18)$$

is the *plasma conductivity* in the absence of magnetic field. Combine the fields (11.14) and (11.17),

$$\mathbf{E}' = \mathbf{E}_u + \mathbf{E}_{\text{eff}}, \quad (11.19)$$

in order to rewrite (11.15) in the most compact form

$$\mathbf{j}' = \sigma \mathbf{E}' - \frac{\omega_B^{(e)}}{\nu_{ei}} \mathbf{j}' \times \mathbf{n}. \quad (11.20)$$

We shall consider (11.20) as an *algebraic* equation in $\mathbf{j}'$, neglecting the $\partial \mathbf{j}'/\partial t$ dependence of the field (11.17). Note, however, that

the term $\partial \mathbf{j}'/\partial t$ is by no means small in the problem of the particle acceleration by a strong electric field in astrophysical plasma.

Collisionless reconnection is a phenomenon in which **particle inertia** (usually combined with *anomalous resistivity*, σ_{an}^{-1} , see Part II, i.e. Somov 2012, Sect. 8.4) of the current density replaces *classical resistivity*, σ^{-1} , in allowing fast reconnection to occur (Drake and Kleva 1991; Horiuchi and Sato 1994; Tsiklauri and Haruki 2007).

11.3 The General Solution

Let us find the solution to (11.20) as a sum of three currents

$$\mathbf{j}' = \sigma_{\parallel} \mathbf{E}'_{\parallel} + \sigma_{\perp} \mathbf{E}'_{\perp} + \sigma_H \mathbf{n} \times \mathbf{E}'_{\perp}. \quad (11.21)$$

Substituting formula (11.21) in (11.20) gives

$$\sigma_{\parallel} = \sigma = \frac{e^2 n}{m_e \nu_{ei}}, \quad (11.22)$$

$$\sigma_{\perp} = \sigma \frac{1}{1 + \left(\omega_B^{(e)} \tau_{ei} \right)^2}, \quad \tau_{ei} = \frac{1}{\nu_{ei}}; \quad (11.23)$$

$$\sigma_H = \sigma_{\perp} \left(\omega_B^{(e)} \tau_{ei} \right) = \sigma \frac{\omega_B^{(e)} \tau_{ei}}{1 + \left(\omega_B^{(e)} \tau_{ei} \right)^2}. \quad (11.24)$$

Formula (11.21) is called the *generalized Ohm's law*. It shows that the presence of a magnetic field in a plasma not only changes the magnitude of the conductivity but changes the form of Ohm's law as well: generally, the electric field and the resulting current are not parallel, since $\sigma_{\perp} \neq \sigma_{\parallel}$. Therefore the electric conductivity of a plasma in a magnetic field is *anisotropic*. Moreover the current component $\mathbf{j}'_H$ which is perpendicular to both the magnetic and electric fields, appears in the plasma. This *out-of-plane* component is the so-called Hall current (Fig. 11.1).




Fig. 11.1 The generalized Ohm's law in a magnetized plasma: the direct ($\mathbf{j}'_{\parallel}$ and $\mathbf{j}'_{\perp}$) and Hall's ($\mathbf{j}'_H$) currents in a plasma with electric ($\mathbf{E}'$) and magnetic ($\mathbf{B}$) fields

11.4 The Conductivity of Magnetized Plasma

11.4.1 Two Limiting Cases

The magnetic-field influence on the conductivity $\sigma_{\perp}$ of the “direct” current $\mathbf{j}'_{\perp}$ across the magnetic field $\mathbf{B}$ and on the Hall-current conductivity σ_H is determined by the parameter $\omega_B^{(e)}\tau_{ei}$ which is the *turning angle* of an electron on the Larmor circle in the intercollisional time. This parameter characterizes the competition between the magnetic field and collisions for control of the electron motion. More general cases are considered in many books, e.g., [Alperovich and Fedorov 2007](#), Sect. 1.4 in context of study the gas-plasma shells of the Earth, i.e. the ionosphere and magnetosphere.

In the simplest case under consideration, we distinguish *two limiting cases*.

(a) Let the turning angle be small:

$$\omega_B^{(e)}\tau_{ei} \ll 1. \quad (11.25)$$

Obviously this inequality corresponds to the *weak* magnetic field or *dense cool* plasma, so that the electric current is scarcely affected by the magnetic field:

$$\sigma_{\perp} \approx \sigma_{\parallel} = \sigma, \quad \frac{\sigma_H}{\sigma} \approx \omega_B^{(e)}\tau_{ei} \ll 1. \quad (11.26)$$

Thus in a frame of reference associated with the plasma, the usual Ohm's law with *isotropic* conductivity holds.

(b) The opposite case, when the electrons *spiral freely* between rare collisions of electrons with ions:

$$\omega_B^{(e)}\tau_{ei} \gg 1, \quad (11.27)$$

corresponds to the *strong* magnetic field and hot rarefied plasma. This plasma is termed the *magnetized* one. It is frequently encountered under astrophysical conditions. In this case

$$\sigma_{\parallel} = \sigma, \quad \sigma_{\perp} \approx \sigma (\omega_B^{(e)}\tau_{ei})^{-2}, \quad \sigma_H \approx \sigma (\omega_B^{(e)}\tau_{ei})^{-1}, \quad (11.28)$$

or

$$\sigma_{\parallel} \approx (\omega_B^{(e)}\tau_{ei}) \sigma_H \approx (\omega_B^{(e)}\tau_{ei})^2 \sigma_{\perp}. \quad (11.29)$$

Hence in the magnetized plasma, for example in the solar corona (see Exercises [11.2](#) and [11.3](#)),

$$\sigma_{\parallel} \gg \sigma_H \gg \sigma_{\perp}.$$

$$(11.30)$$




Fig. 11.2 Initiation of the current in the direction of the perpendicular field $\mathbf{E}'_{\perp}$ as the result of rare collisions (1, 2, 3, ...) against a background of the electric drift. Only collisions of electrons are shown

In other words, the impact of the magnetic field on the direct current is especially strong for the component resulting from the electric field $\mathbf{E}'_{\perp}$. The current in the $\mathbf{E}'_{\perp}$ direction is considerably weaker than it would be in the absence of a magnetic field. Why is this so?

11.4.2 The Physical Interpretation

The physical mechanism of the perpendicular current $\mathbf{j}'_{\perp}$ is illustrated by Fig. 11.2.

The primary effect of the electric field $\mathbf{E}'_{\perp}$ in the presence of the magnetic field $\mathbf{B}$ is not the current in the direction $\mathbf{E}'_{\perp}$, but rather the electric *drift* in the direction perpendicular to both $\mathbf{B}$ and $\mathbf{E}'_{\perp}$.

The electric drift velocity (5.22) is independent of the particle's mass and charge. The electric drift of electrons and ions generates the motion of the plasma as a whole with the velocity

$$\mathbf{v} = \mathbf{v}_d = c \frac{\mathbf{E} \times \mathbf{B}}{B^2}. \quad (11.31)$$

This would be the case if there were no collisions at all (Fig. 5.3).

Collisions, even the rare ones, result in a disturbance of the particle's Larmor motion, leading to a displacement of the ions (not shown in Fig. 11.2) along the field $\mathbf{E}'_{\perp}$, and the electrons in the opposite direction as shown in Fig. 11.2. The small electric current $\mathbf{j}'_{\perp}$ (a factor of $\omega_B^{(e)} \tau_{ei}$ smaller than the drift one) appears in the direction $\mathbf{E}'_{\perp}$.

In order to ensure the current across the magnetic field, the so-called *Hall electric field* is necessary, that is the electric field component perpendicular to both the current $\mathbf{j}'_{\perp}$ and the field $\mathbf{B}$ (e.g., [Braginskii 1965](#); [Sivukhin 1996](#), Chap. 7, Sect. 98). This is the secondary effect but it is not small in a strong magnetic field.

The Hall electric field balances the Lorentz force acting on the carriers of the perpendicular electric current in a rarely collisional plasma due to the presence of a magnetic field,

i.e. the force

$$\begin{aligned} \mathbf{F}(\mathbf{j}'_{\perp}) &= \frac{en}{c} \mathbf{u}_{i\perp} \times \mathbf{B} - \frac{en}{c} \mathbf{u}_{e\perp} \times \mathbf{B} = \\ &= \frac{1}{c} en (\mathbf{u}_{i\perp} - \mathbf{u}_{e\perp}) \times \mathbf{B}. \end{aligned} \quad (11.32)$$

Hence the magnitude of the Hall electric field is

$$\mathbf{E}'_H = \frac{1}{en c} \mathbf{j}'_{\perp} \times \mathbf{B}. \quad (11.33)$$

The current $\mathbf{j}'_{\perp}$ is small but magnetic field $\mathbf{B}$ is strong according to (11.27). That is why the Hall electric field and Hall electric current can play significant role in plasma astrophysics.

The Hall electric field in astrophysical plasma is frequently set up automatically, as a consequence of small charge separation within the limits of quasi-neutrality. In this case the “external” field, which has to be applied to the plasma, is determined by the expressions

$$E'_{\parallel} = j'_{\parallel} / \sigma_{\parallel} \quad \text{and} \quad E'_{\perp} = j'_{\perp} / \sigma_{\perp}. \quad (11.34)$$

We shall not discuss here the dissipation process under the conditions of anisotropic conductivity. In general, considering the media with complicated properties, the symmetric highest component of the *conductivity tensor* can play the most important role in Joule heating (see [Landau et al. 1984](#), Chap. 3).

In the particular case of a fully-ionized rarely-collisional plasma, in the process of fundamental significance for the flare energy release problem, the tendency for a particle to spiral round the magnetic field lines insures the great reduction in the transversal conductivity (11.23). However, since the dissipation of the energy of the electric current into Joule heat, $\mathbf{j}' \cdot \mathbf{E}'$, is due solely to collisions between particles (if the particle acceleration can be neglected), the reduced conductivity does not lead to significantly increased dissipation in rarely-collisional plasma (Exercise 11.4).

On the other hand, the Hall electric field and Hall current can significantly modify conditions of magnetic reconnection ([Bhattacharjee 2004](#); [Cassak et al. 2007](#)). For example,

compared with ordinary resistive magnetohydrodynamic (MHD), the Hall MHD reconnection is distinguished by qualitatively different magnetic field distributions, electron and ion signatures

in reconnecting current layers (RCLs); see Part II, Sect. 2.4.4.

Although the Hall effect itself is nondissipative,

$$\mathbf{j}'_{\text{H}} \mathbf{E}' = 0, \quad (11.35)$$

it can lead to dissipation through a turbulent “*Hall cascade*”, magnetic energy cascading from large to small scales, where it dissipates by ohmic decay (Goldreich and Reisenegger 1992). The Hall effect can dominate ohmic decay of currents in the crust of neutron stars and therefore can determine evolution of their magnetic field. In an initial poloidal dipole field, the toroidal currents initially “twist” the field. The resulting poloidal currents then generate a quadrupole poloidal field via a “pinch” (see Fig. 10 in Cumming et al. 2004).

11.5 Currents and Charges in Plasma

11.5.1 Collisional and Collisionless Plasmas

Let us point out another property of the generalized Ohm’s law. Under laboratory conditions, as a rule, one cannot neglect the gradient forces (11.9). On the contrary, these forces usually play no part in astrophysical plasma. We shall often ignore them. However this simplification may be not well justified in such important applications as reconnecting current layers (RCLs), shock waves and other discontinuities.

Moreover let us also restrict our consideration to very *slow* (say *hydrodynamic*) motions of plasma. These motions are supposed to be so slow that the following *three conditions* are fulfilled.

First, it is supposed that

$$\omega = \frac{1}{\tau} \ll \nu_{\text{ei}} \quad \text{or} \quad \nu_{\text{ei}} \tau \gg 1, \quad (11.36)$$

where τ is a characteristic time of the plasma motions. Recall that

$$\nu_{\text{ei}} \ll \nu_{\text{ii}} \ll \nu_{\text{ee}}.$$

Therefore the frequency ω is lower than the lowest collisional frequency.

As a consequence,

the departures of actual distribution functions for electrons and ions from the Maxwellian distribution are small.

This allows us to handle the transfer phenomena in linear approximation.

Moreover, if a single-fluid model is to make physical sense, the electrons and ions could have comparable temperatures, ideally, the same temperature T which is the temperature of the plasma as a whole:

$$T_e = T_p = T .$$

Second, we neglect the electron inertia in comparison with that of the ions and make use of (11.11). This condition is usually written in the form

$$\omega \ll \omega_B^{(i)} = \frac{eB}{m_i c} . \quad (11.37)$$

Recall that the gyro-frequency of ions $\omega_B^{(i)} \ll \omega_B^{(e)}$.

Thus

the plasma motions have to be so slow that their frequency is smaller than the lowest gyro-frequency of the particles.

The third condition

$$\omega_B^{(e)} \tau_{ei} \ll 1 . \quad (11.38)$$

Hence the hydrodynamic approximation can be used, the conductivity σ being isotropic.

With all these three conditions together, the generalized Ohm's law assumes the following form which is specific to the classic or ordinary *magnetohydrodynamics* (MHD):

$$\mathbf{j}' = \sigma \left(\mathbf{E} + \frac{1}{c} \mathbf{u} \times \mathbf{B} \right) . \quad (11.39)$$

The MHD approximation is the subject of the next chapter. Numerous applications of MHD to various phenomena in astrophysical plasma will be considered in many places in the remainder of the book.

In the opposite case (11.27), when the parameter $\omega_B^{(e)} \tau_{ei}$ is large, charged particles revolve around magnetic field lines, and a typical particle may spend a considerable time in a region of a size of the order of the gyroradius (5.14). Hence, if the length scale of a phenomenon is much larger than the gyroradius, we may expect the hydrodynamic-type models to work.

It appears that, even when the parameter $\omega_B^{(e)} \tau_{ei}$ tends to infinity (like in the solar corona, see Exercise 11.3) and collisions are negligible, the *quasi-hydrodynamic* description of plasma, the Chew-Goldberger-Low (CGL) approximation (Chew et al. 1956) is possible (especially if the actual electric field $\mathbf{E}$ in a collisionless plasma is perpendicular to a sufficiently strong magnetic field $\mathbf{B}$) and quite useful. This is because

the strong magnetic field makes the plasma, even a collisionless one, more “interconnected”, so to speak, more hydrodynamic in the directions perpendicular to the magnetic field.

That allows one to write down a well-justified set of two-dimensional MHD equations for the collisionless plasma in a magnetic field (see [Volkov 1966](#), (42)–(45)). As for the motion of collisionless particles along the magnetic field, some important kinetic features and physical restrictions still are significant ([Klimontovich et al. 1961](#); [Shkarofsky et al. 1966](#), Chap. 10). [Chew et al. \(1956\)](#) emphasized that “a strictly hydrodynamic approach to the problem is appropriate only when some special circumstance suppresses the effects of pressure transport along the magnetic lines”.

There is ample experimental evidence that

strong magnetic fields do make astrophysical plasmas behave like hydrodynamic charged fluids.

This does not mean, of course, that there are no pure kinetic phenomena in such plasmas. There are many of them indeed. The most interesting of them is magnetic reconnection, for example, in the solar corona and solar wind (e.g., [Sheeley et al. 2007](#); [Gosling et al. 2007a](#)).

11.5.2 Volume Charge and Quasi-Neutrality

One more remark concerning the generalized Ohm’s law is important for the following. While deriving the law in Sect. 11.2, the *exact* charge neutrality of plasma or the exact electric neutrality was assumed:

$$\sum_i Z_i n_i = n_e, \quad (11.40)$$

i.e. the absolute absence of the *volume charge* in plasma: $\rho^q = 0$. However there is no need for such a strong restriction. It is sufficient to require *quasi-neutrality*, i.e. the numbers of ions (with account of their charge taken) and electrons per unit volume are very nearly equal:

$$\left(\sum_i Z_i n_i - n_e \right) n_e^{-1} \ll 1. \quad (11.41)$$

So

the volume charge density has to be small in comparison to the plasma density.

Once the volume charge density

$$\rho^q = e \left(\sum_i Z_i n_i - n_e \right) \neq 0, \quad (11.42)$$

yet another term must be taken into account in the generalized Ohm's law:

$$\mathbf{j}_u^q = \rho^q \mathbf{u}. \quad (11.43)$$

This is the so-called *convective* current. It is caused by the volume charge transfer and must be added to the *conductive* current (11.21).

The volume charge, the associated electric force $\rho^q \mathbf{E}$ and the convective current $\rho^q \mathbf{u}$ are of great importance in electrodynamics of relativistic objects such as *black holes* (Novikov and Frolov 1989) and *pulsars* (Michel 1991). *Charge-separated plasmas* originate in magnetospheres of pulsars and rotating black holes, for example, a super-massive black hole in active galactic nuclei (AGN).

The shortage of charge leads to the emergence of a strong electric field along the magnetic field lines.

The parallel electric field accelerates migratory electrons and/or positrons to ultra-relativistic energies (e.g., Hirotani and Okamoto 1998).

Charge density oscillations in a plasma, the Langmuir waves, are considered in Sect. 10.2.

The volume charge density can be evaluated in the following manner. On the one hand, from Maxwell's equation $\text{div } \mathbf{E} = 4\pi\rho^q$ we estimate

$$\rho^q \approx \frac{E}{4\pi L}. \quad (11.44)$$

On the other hand, the non-relativistic equation of plasma motion yields

$$en_e E \approx \frac{p}{L} \approx \frac{n_e k_B T}{L},$$

so that

$$E \approx \frac{k_B T}{eL}. \quad (11.45)$$

On substituting (11.45) in (11.44), we find the following estimate

$$\frac{\rho^q}{en_e} \approx \frac{k_B T}{eL} \frac{1}{4\pi L} \frac{1}{en_e} = \frac{1}{L^2} \left(\frac{k_B T}{4\pi e^2 n_e} \right)$$

or

$$\frac{\rho^q}{en_e} \approx \frac{r_{\text{DH}}^2}{L^2}.$$

(11.46)

Here

$$r_{\text{DH}} = r_{\text{DH}}^{(\text{e})} = \left(\frac{k_{\text{B}} T}{4\pi e^2 n_{\text{e}}} \right)^{1/2}. \quad (11.47)$$

is the electron Debye-Hückel radius (see Sect. 8.2.1).

Since the usual *concept of plasma* implies that the Debye-Hückel radius

$$r_{\text{DH}} \ll L, \quad (11.48)$$

the volume charge density is small in comparison with the plasma density. When we consider astrophysical phenomena with a length scale L much larger than the Debye-Hückel radius r_{DH} and a time scale τ much larger than the inverse the plasma frequency $\omega_{\text{pl}}^{(\text{e})}$ (see Sect. 10.2.2, formula (10.15)), the charge separation in the plasma can be neglected.

On the contrary, in laboratory and technical devices, we often deal with non-neutral charged plasmas (see, e.g., Davidson 1974).

11.6 Practice: Exercises and Answers

Exercise 11.1 (Sect. 11.2). Consider a plasma system with given distributions of magnetic and velocity fields. Is it possible to use (11.15) in order to estimate the growth rate of electric current and, as a consequence, of magnetic field in such a system, for example, a protostar?

Exercise 11.2 (Sect. 11.4.1). Evaluate the characteristic value of the parallel conductivity (11.22) in the *solar corona*.

Answer. It follows from formula (11.22) that

$$\sigma_{\parallel} = \frac{e^2 n}{m_{\text{e}}} \tau_{\text{ei}} = 2.53 \times 10^8 n \tau_{\text{ei}} \sim 10^{16} - 10^{17}, \text{ s}^{-1}, \quad (11.49)$$

if we take $\tau_{\text{ep}} \sim 0.2 - 2.0 \text{ s}$ (Exercise 11.1).

Exercise 11.3 (Sect. 11.4.1). Estimate the parameter $\omega_{\text{B}}^{(\text{e})} \tau_{\text{ei}}$ in the solar corona above a sunspot.

Answer. Just above a large sunspot the field strength can be as high as $B \approx 3,000 \text{ G}$. Here the electron Larmor frequency $\omega_{\text{B}}^{(\text{e})} \approx 5 \times 10^{10} \text{ rad s}^{-1}$ (Exercise 11.1). Characteristic time of *close* electron-proton collisions $\tau_{\text{ep}}(cl) \approx 22 \text{ s}$ (see Exercise 11.1). Therefore $\omega_{\text{B}}^{(\text{e})} \tau_{\text{ei}}(cl) \sim 10^{12} \text{ rad} \gg 1$.

Distant collisions are much more frequent (Exercise 11.1). However, even with $\tau_{\text{ep}} \approx 0.1 \text{ s}$, we obtain

$$\omega_{\text{B}}^{(\text{e})} \tau_{\text{ei}} \sim 10^{10} \text{ rad} \gg 1.$$

So, for anisotropic conductivity in the solar corona, the approximate formulae (11.28) can be well used.

Exercise 11.4 (Sect. 11.4.2). Consider the generalized Ohm's law in the case when the electric field is perpendicular to the magnetic field $\mathbf{B} = B \mathbf{n}$. So

$$\mathbf{j}' = \sigma_{\perp} \mathbf{E}'_{\perp} + \sigma_H \mathbf{n} \times \mathbf{E}'_{\perp}, \quad (11.50)$$

where

$$\sigma_{\perp} = \sigma \frac{1}{1 + \left(\omega_B^{(e)} \tau_{ei}\right)^2} \quad \text{and} \quad \sigma_H = \sigma \frac{\omega_B^{(e)} \tau_{ei}}{1 + \left(\omega_B^{(e)} \tau_{ei}\right)^2}. \quad (11.51)$$

This indicates that the current $\mathbf{j}'_{\perp}$ in the direction of $\mathbf{E}'_{\perp}$ is reduced in the ratio

$$1 / \left(1 + \left(\omega_B^{(e)} \tau_{ei}\right)^2\right) \approx \left(\omega_B^{(e)} \tau_{ei}\right)^{-2}, \quad \text{if } \omega_B^{(e)} \tau_{ei} \gg 1,$$

by the magnetic field. In addition, the other current $\left(\omega_B^{(e)} \tau_{ei}\right)^2$ times as large flows in the direction perpendicular to both $\mathbf{B}$ and $\mathbf{E}'_{\perp}$; this is the *Hall current* $\mathbf{j}'_H$.

Show that

the reduction in the “perpendicular” conductivity (Fig. 11.1) does not increase the rate of dissipation of current energy

(see Cowling 1976, Sect. 6.2).

Chapter 12

Single-Fluid Models for Astrophysical Plasma

Abstract Single-fluid models are the simplest but sufficient approximation to describe many large-scale low-frequency phenomena in astrophysical plasma: regular and turbulent dynamo, plasma motions driven by strong magnetic fields, accretion disks, and relativistic jets.

12.1 Derivation of the Single-Fluid Equations

12.1.1 The Continuity Equation

In order to consider astrophysical plasma as a *single* hydrodynamic medium, we have to sum each of the three transfer equations over all kinds of particles. Let us start from the continuity equation (9.14). With allowance for the definition of the plasma mass density (9.6), we have

$$\frac{\partial \rho}{\partial t} + \operatorname{div} \left(\sum_k \rho_k \mathbf{u}_k \right) = 0. \quad (12.1)$$

The mean velocities of motion for all kinds of particles are supposed to be equal to the plasma hydrodynamic velocity:

$$\mathbf{u}_1(\mathbf{r}, t) = \mathbf{u}_2(\mathbf{r}, t) = \cdots = \mathbf{u}(\mathbf{r}, t), \quad (12.2)$$

as a result of action of the mean collisional force (9.24). However this is not a general case.

In general, the mean velocities are not the same, but a frame of reference can be chosen in which

$$\rho \mathbf{u} = \sum_k \rho_k \mathbf{u}_k. \quad (12.3)$$

Then from (12.1) and (12.3) we obtain the usual *continuity equation*

$$\boxed{\frac{\partial \rho}{\partial t} + \operatorname{div} \rho \mathbf{u} = 0.} \quad (12.4)$$

In what follows we shall consider both cases mentioned above.

12.1.2 The Momentum Conservation Law in Plasma

In much the same way as in previous section, we handle the momentum transfer equation (9.46). On summing over all kinds of particles, we obtain the following equation:

$$\rho \frac{d u_\alpha}{dt} = - \frac{\partial}{\partial r_\beta} p_{\alpha\beta} + \rho^q E_\alpha + \frac{1}{c} (\mathbf{j} \times \mathbf{B})_\alpha + \sum_k \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v. \quad (12.5)$$

Here the pressure tensor

$$p_{\alpha\beta} = \sum_k p_{\alpha\beta}^{(k)},$$

the *volume charge* density in plasma is

$$\rho^q = \sum_k n_k e_k = \frac{1}{4\pi} \operatorname{div} \mathbf{E}, \quad (12.6)$$

and the electric current density is

$$\mathbf{j} = \sum_k n_k e_k \mathbf{u}_k = \frac{c}{4\pi} \operatorname{rot} \mathbf{B} - \frac{1}{4\pi} \frac{\partial \mathbf{E}}{\partial t}. \quad (12.7)$$

The electric and magnetic fields, $\mathbf{E}$ and $\mathbf{B}$, involved in this description of astrophysical plasma are the averaged fields associated with the total electric charge density ρ^q and the total current density $\mathbf{j}$. They satisfy the macroscopic Maxwell equations. In astrophysical plasma, the magnetic *permeability* and the electric *permittivity* can almost always be replaced by their vacuum values. For this reason, the macroscopic Maxwell equations have the same structure as (1.30) and (1.27) that have been used on the right-hand side of formulae (12.6) and (12.7).

Since elastic collisions do not change the total momentum, we have

$$\sum_k \langle F_{k,\alpha}^{(c)}(\mathbf{r}, t) \rangle_v = 0. \quad (12.8)$$

On substituting (12.6)–(12.8) in (12.5), the latter can be rearranged to give the *momentum conservation law* for plasma

$$\rho \frac{d u_\alpha}{dt} = - \frac{\partial}{\partial r_\beta} p_{\alpha\beta} + F_\alpha(\mathbf{E}, \mathbf{B}). \quad (12.9)$$

Here the electromagnetic force is written in terms of the electric and magnetic field vectors:

$$F_\alpha(\mathbf{E}, \mathbf{B}) = - \frac{\partial}{\partial t} \frac{(\mathbf{E} \times \mathbf{B})_\alpha}{4\pi c} - \frac{\partial}{\partial r_\beta} M_{\alpha\beta} + \frac{1}{4\pi} E_\alpha \operatorname{div} \mathbf{E}. \quad (12.10)$$

The tensor

$$M_{\alpha\beta} = \frac{1}{4\pi} \left[-E_\alpha E_\beta - B_\alpha B_\beta + \frac{1}{2} \delta_{\alpha\beta} (E^2 + B^2) \right] \quad (12.11)$$

is called the *Maxwellian tensor* of stresses. The last term in (12.10) represents the electric force acting on the volume charge ρ^q .

The divergent form of the *momentum conservation law* is

$$\frac{\partial}{\partial t} \left[\rho u_\alpha + \frac{(\mathbf{E} \times \mathbf{B})_\alpha}{4\pi c} \right] + \frac{\partial}{\partial r_\beta} (\Pi_{\alpha\beta} + M_{\alpha\beta}) = 0. \quad (12.12)$$

The operator $\partial/\partial t$ acts on two terms that correspond to momentum density: $\rho \mathbf{u}$ is the momentum of the motion of the plasma as a whole in a unit volume, $\mathbf{E} \times \mathbf{B}/4\pi c$ is the momentum density of the electromagnetic field.

The divergency operator $\partial/\partial r_\alpha$ acts on

$$\Pi_{\alpha\beta} = p_{\alpha\beta} + \rho u_\alpha u_\beta \quad (12.13)$$

which is the *momentum flux density* tensor

$$\Pi_{\alpha\beta} = \sum_k \Pi_{\alpha\beta}^{(k)}, \quad (12.14)$$

see definition (9.10). Therefore the *pressure tensor*

$$p_{\alpha\beta} = p \delta_{\alpha\beta} + \pi_{\alpha\beta}, \quad (12.15)$$

where

$$p = \frac{1}{3} \delta_{\alpha\alpha} = \sum_k p_k \quad (12.16)$$

is the total plasma pressure, the sum of *partial pressures*, and

$$\pi_{\alpha\beta} = \sum_k \pi_{\alpha\beta}^{(k)} \quad (12.17)$$

is the *viscous stress* tensor (see definition (9.39)), which allows for the transport of momentum from one layer of the plasma flow to the other layers so that relative motions inside the plasma are damped out. If we accept condition (12.2) then the random velocities are now defined with respect to the macroscopic velocity $\mathbf{u}$ of the plasma as a whole.

The momentum conservation law in the form (12.9) or (12.12) is applied for a wide range of conditions in astrophysical plasmas like *fluid relativistic flows*, for example, astrophysical jets (Sect. 13.3). The assumption that the astrophysical plasma behaves as a continuum medium, which is essential if these forms of the momentum conservation law are to be applied, is excellent in the cases in which we are often interested:

the Debye-Hückel length and the particle Larmor radii are much smaller than the plasma flow scales.

On the other hand, going from the multi-fluid description to a single-fluid model is a serious damage because we loose an information not only on the small-scale dynamics of the electrons and ions but also on the high-frequency processes in plasma.

The single-fluid equations describe well the *low-frequency large-scale* behaviour of plasma in astrophysical conditions.

12.1.3 The Energy Conservation Law

In a similar manner as above, the energy conservation law is derived. We sum the general equation (9.25) over k and then substitute in the resulting equation the total electric charge density (12.6) and the total electric current density (12.7) expressed in terms of the electric field $\mathbf{E}$ and magnetic field $\mathbf{B}$. On rearrangement, the following divergent form of the energy conservation law (cf. the simplified equation (1.56) for electromagnetic field energy and kinetic energy of charged particles is obtained:

$$\begin{aligned} & \frac{\partial}{\partial t} \left(\frac{\rho u^2}{2} + \rho \varepsilon + \frac{E^2 + B^2}{8\pi} \right) + \\ & + \frac{\partial}{\partial r_\alpha} \left[\rho u_\alpha \left(\frac{u^2}{2} + w \right) + \frac{c}{4\pi} (\mathbf{E} \times \mathbf{B})_\alpha + \pi_{\alpha\beta} u_\beta + q_\alpha \right] = \\ & = (u_\alpha F_\alpha^{(c)})_{ff} + \mathcal{L}^{(rad)}(\mathbf{r}, t). \end{aligned} \quad (12.18)$$

On the left-hand side of this equation, an additional term has appeared: the operator $\partial/\partial t$ acts on the energy density of the electromagnetic field

$$W = \frac{E^2 + B^2}{8\pi}. \quad (12.19)$$

The divergency operator $\partial/\partial r_\alpha$ acts on the *Poynting vector*, the electromagnetic energy flux through a unit surface in space:

$$\mathbf{G} = \frac{c}{4\pi} [\mathbf{E} \times \mathbf{B}]. \quad (12.20)$$

The right-hand side of (12.18) contains the total work of *friction forces* (9.42) in unit time on unit volume

$$(u_\alpha F_\alpha^{(c)})_{ff} = \sum_k (F_{k,\alpha}^{(c)} u_{k,\alpha}) = \sum_k u_{k,\alpha} \int_{\mathbf{v}} m_k v'_\alpha \left(\frac{\partial f_k}{\partial t} \right)_c d^3 \mathbf{v}. \quad (12.21)$$

This work related to the relative motion of the plasma components is not zero. Recall that we consider general case (12.3).

By contrast, the total heat release under elastic collisions between particles of different kinds (see definition (9.43)) is

$$\sum_k Q_k^{(c)}(\mathbf{r}, t) = \sum_k \int_{\mathbf{v}} \frac{m_k (v')^2}{2} \left(\frac{\partial f_k}{\partial t} \right)_c d^3 \mathbf{v} = 0. \quad (12.22)$$

█ Elastic collisions in a plasma conserve both the total momentum (see (12.8)) and the total energy (see (12.22)).

If we accept condition (12.2) then, with account of formula (9.24), the collisional heating (12.21) by friction force is also equal to zero. In this limit, there is not any term which contains the collisional integral. *Elastic collisions have done a good job.*


Inelastic collisions are still important in radiative cooling and heating. In optically thin plasma in the thermodynamical equilibrium state, with collisional excitations of ions, the power of radiation from a unit volume of plasma is proportional to the square of plasma density n (cm^{-3}):

$$\mathcal{L}^{(rad)} \simeq -n^2 q(T). \quad (12.23)$$

The function $q(T)$ measured in $\text{erg cm}^{+3} \text{s}^{-1}$ is called the *radiative losses function*. It depends strongly on the temperature T as illustrated by Fig. 12.1 but weakly depends on the plasma density n (Cox and Tucker 1969).

The radiative energy losses in astrophysical plasma often create the thermal non-equilibrium processes which govern the plasma evolution. Under action of magnetic reconnection, for example, in the solar corona, plasma can be raked

Fig. 12.1 Radiative energy loss function q vs. temperature T at fixed value of density $n = 10^9 \text{ cm}^{-3}$. (From Somov et al. 2007; reproduced with permission © Pleiades Publishing, Ltd)



and compressed by magnetic field (see Chap. 3 in Somov and Syrovatskii 1976b). Radiation can become dominant as the energy loss mechanism and lead to rapid cooling and compression of plasma by different modes of *thermal instability* (Field 1965); see also Somov and Syrovatskii (1976a). This instability results in the formation of cold dense loops in the corona.

Luna et al. (2012) have found that the condensations in the corona can be divided into populations: threads and blobs. The threads are the principal contributors to the total mass of *solar prominences*, whereas the blob contribution is small. Luna et al. have generated synthetic images of prominences, which look similar to the images observed by the *Solar Dynamics Observatory* (SDO).

Note in conclusion that, in the frame of our simplified treatment of the second-order moments (Sect. 9.4.1), we still do not have any equations for the anisotropic part of the pressure tensor, which is the viscous stress tensor $\pi_{\alpha\beta}$, and for the flux q_α of heat due to random motions of particles. This is not unexpected, of course, but inherent at the method of the moments as discussed in Sect. 9.5. In order to be correct we have to find these transfer coefficients by using the procedure similar to that described in Sect. 9.6, see also Sect. 9.7, where the Grad moment method has been used.

12.2 Basic Assumptions and the MHD Equations

12.2.1 Old Simplifying Assumptions

As we saw in Chap. 9, the set of transfer equations for local macroscopic quantities determines the behavior of different kinds of particles, such as electrons and ions in astrophysical plasma, once two main conditions are complied with:

- (a) Many collisions occur in a characteristic time τ of the process or phenomenon under consideration:

$$\tau \gg \tau_c, \quad (12.24)$$

- (b) The particle's path between two collisions—the particle's free path—is significantly smaller than the distance L , over which macroscopic quantities change considerably:

$$L \gg \lambda_c. \quad (12.25)$$

Here τ_c and λ_c are the *collisional characteristic time* and the collisional *mean free path*, respectively. Once these conditions are satisfied, we can close the set of *transfer* equations, as has been discussed in Sect. 9.6.

While considering the generalized Ohm's law in Chap. 11, three other assumptions have been made, that are complementary to the restriction (12.24) on the characteristic time τ of the process.

The first condition can be written in the form

$$\tau \gg \tau_{ei} = \nu_{ei}^{-1}, \quad (12.26)$$

where τ_{ei} is the electron-ion collisional time, the longest collisional relaxation time. Thus departures from the Maxwellian distribution are small. Moreover the electrons and ions should have comparable temperatures, ideally, the same temperature T being the temperature of the plasma as a whole.

Second, we neglect the electron inertia in comparison with that of the ions. This condition is usually written as

$$\tau \gg (\omega_B^{(i)})^{-1}, \quad \text{where} \quad \omega_B^{(i)} = \frac{eB}{m_i c}. \quad (12.27)$$

Thus the plasma motions have to be so slow that their frequency $\omega = 1/\tau$ is smaller than the lowest gyro-frequency of the particles. Recall that the gyro-frequency of ions $\omega_B^{(i)} \ll \omega_B^{(e)}$.

The third condition,

$$\omega_B^{(e)} \tau_{ei} \ll 1, \quad (12.28)$$

is necessary to write down Ohm's law in the form

$$\mathbf{j} = \sigma \left(\mathbf{E} + \frac{1}{c} \mathbf{v} \times \mathbf{B} \right) + \rho^q \mathbf{v}. \quad (12.29)$$

Here $\mathbf{v}$ is the macroscopic velocity of plasma considered as a continuous medium, $\mathbf{E}$ and $\mathbf{B}$ are the electric and magnetic fields in the “laboratory” system of coordinates, where we measure the velocity $\mathbf{v}$. Accordingly, the field

$$\mathbf{E}_v = \mathbf{E} + \frac{1}{c} \mathbf{v} \times \mathbf{B} \quad (12.30)$$

is the electric field in a frame of reference related to the plasma. The isotropic conductivity is (see formula (11.18)):

$$\sigma = \frac{e^2 n}{m_e v_{ei}} . \quad (12.31)$$

Complementary to the restriction (12.25) on the characteristic length L of the phenomenon under consideration, we have to add the condition

$$L \gg r_{\text{DH}} , \quad (12.32)$$

where r_{DH} is the Debye-Hückel radius. Then the volume charge density ρ^q is small in comparison with the plasma density ρ .

Under the conditions listed above, we use the *general* hydrodynamic-type equations which are the conservation laws for mass (12.4), momentum (12.5) and energy (12.18).

The general hydrodynamic-type equations have a much wider area of applicability in astrophysics than the equations of *ordinary* magnetohydrodynamics (MHD) derived below.

The latter will be much simpler than the equations derived in Sect. 12.1. Therefore *additional simplifying assumptions* are necessary. Let us introduce them. There are two.

12.2.2 New Simplifying Assumptions

First assumption: the plasma conductivity σ is assumed to be large, the electromagnetic processes being not very fast. Then, in Maxwell's equation (1.27)

$$\text{rot } \mathbf{B} = \frac{4\pi}{c} \mathbf{j} + \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} ,$$

we ignore the *displacement* current in comparison to the *conductive* one. The corresponding condition is found by evaluating the currents as follows

$$\frac{1}{c} \frac{E}{\tau} \ll \frac{4\pi}{c} j \quad \text{or} \quad \omega E \ll 4\pi \sigma E .$$

Thus we suppose that

$$\omega \ll 4\pi \sigma .$$

(12.33)

In the same order with reference to the small parameter ω/σ (more exactly, $\omega/4\pi\sigma$), we can neglect the *convective* current (see formula (11.43) and its discussion in Sect. 11.5.2) in comparison with the *conductive* current in Ohm's law (12.29). Actually,

$$\rho^q v \approx v \operatorname{div} \mathbf{E} \frac{1}{4\pi} \approx \frac{L}{\tau} \frac{E}{L} \frac{1}{4\pi} \approx \frac{\omega}{4\pi} E \ll \sigma E,$$

once the condition (12.33) is satisfied.

The conductivity of astrophysical plasma, which is often treated in the MHD approximation, is very high (e.g., Exercise 11.1). This is the reason why condition (12.33) is satisfied up to frequencies close to optical ones.

Neglecting the displacement current and the convective current, Maxwell's equations and Ohm's law result in the following relations:

$$\mathbf{j} = \frac{c}{4\pi} \operatorname{rot} \mathbf{B}, \quad (12.34)$$

$$\mathbf{E} = -\frac{1}{c} \mathbf{v} \times \mathbf{B} + \frac{c}{4\pi\sigma} \operatorname{rot} \mathbf{B}, \quad (12.35)$$

$$\rho^q = -\frac{1}{4\pi c} \operatorname{div} (\mathbf{v} \times \mathbf{B}), \quad (12.36)$$

$$\operatorname{div} \mathbf{B} = 0, \quad (12.37)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \operatorname{rot} (\mathbf{v} \times \mathbf{B}) + \frac{c^2}{4\pi\sigma} \Delta \mathbf{B}. \quad (12.38)$$

Once two vectors, $\mathbf{B}$ and $\mathbf{v}$, are given, the current density $\mathbf{j}$, the electric field $\mathbf{E}$, and the volume charge density ρ^q are completely determined by formulae (12.34)–(12.36). Thus

the problem is reduced to finding the interaction of the magnetic field $\mathbf{B}$ and the hydrodynamic velocity field $\mathbf{v}$.

As a consequence, the approach under discussion has come to be known as *magnetohydrodynamics* (Alfvén 1950; Syrovatskii 1957).

The corresponding equation of plasma motion is obtained by substitution of formulae (12.34)–(12.36) in the equation of momentum transfer (12.5). With due regard for the manner in which *viscous force* is usually written in ordinary hydrodynamics, we have

$$\begin{aligned} \rho \frac{d\mathbf{v}}{dt} = & -\nabla p + \rho^q \mathbf{E} - \frac{1}{4\pi} \mathbf{B} \times \operatorname{rot} \mathbf{B} + \\ & + \eta \Delta \mathbf{v} + \left(\zeta + \frac{\eta}{3} \right) \nabla \operatorname{div} \mathbf{v}. \end{aligned} \quad (12.39)$$

Here η is the *first viscosity* coefficient, ζ is the *second viscosity* coefficient (see [Landau and Lifshitz 1959a](#), Chap. 2, Sect. 15). Formulae for these coefficients as well as for the viscous force should be derived from the moment equation for the pressure tensor, which we were not inclined to write down in Sect. 9.3 being busy in the way to the energy conservation law.

* * *

The *second additional assumption* has to be introduced now. Treating (12.39), the electric force $\rho^q \mathbf{E}$ can be ignored in comparison to the magnetic one if

$$v^2 \ll c^2, \quad (12.40)$$

that is in the non-relativistic approximation. To make certain that this is true, evaluate the electric force using (12.36) and (12.35):

$$\rho^q E \approx \frac{1}{4\pi c} \frac{vB}{L} \frac{vB}{c} \approx \frac{B^2}{4\pi} \frac{1}{L} \frac{v^2}{c^2}, \quad (12.41)$$

the magnetic force being proportional to

$$\frac{1}{4\pi} |\mathbf{B} \times \text{rot} \mathbf{B}| \approx \frac{B^2}{4\pi} \frac{1}{L}. \quad (12.42)$$

Comparing (12.41) with (12.42), we see that the electric force is a factor of v^2/c^2 short of the magnetic one.

In a great number of astrophysical applications of MHD, the plasma velocities fall far short of the speed of light. The Sun is a good case in point. Here the largest velocities observed, for example, in coronal transients and coronal mass ejections (CMEs) do not exceed several thousands of km/s, i.e. $\lesssim 3 \times 10^8$ cm/s. Under these conditions, *we neglect the electric force acting upon the volume charge* in comparison with the magnetic force.

However the relativistic objects such as accretion disks near rotating black holes (see Chap. 7 in [Novikov and Frolov 1989](#)), pulsar magnetospheres ([Michel 1991](#); [Rose 1998](#); [Becker 2009](#)), and relativistic objects in compact binaries ([Colpi et al. 2009](#)) are at the other extreme.

The electric force acting on the volume charge plays a crucial role in the electrodynamics of relativistic objects.

For example, the fast rotation of a magnetised neutron star in vacuum induces an enormous electric field. This strong field tears particles from the stellar surface and accelerates them up to high energies. So a vacuum rotator has to form a charged magnetosphere as charged particles move from the surface to short out $\mathbf{E} \cdot \mathbf{B}$. The charge density is expected to be close to the density required to screen the induced electric field.

The primary beams of accelerated particles initiate an *electron-positron cascade* which, according to conventional wisdom, populates the entire magnetosphere with plasma. In a simplified picture of an aligned rotator, the charges in the a fully *charge-separated plasma* flowing parallel to the poloidal magnetic field, generate a poloidal convective current and, therefore, a toroidal magnetic field.

12.2.3 Non-relativistic Magnetohydrodynamics

With the assumptions made above, the considerable simplifications have been obtained; and now we write the following set of equations of non-relativistic MHD:

$$\frac{\partial}{\partial t} \rho v_\alpha = -\frac{\partial}{\partial r_\beta} \Pi_{\alpha\beta}^*, \quad (12.43)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \text{rot}(\mathbf{v} \times \mathbf{B}) + \nu_m \Delta \mathbf{B}, \quad (12.44)$$

$$\text{div} \mathbf{B} = 0, \quad (12.45)$$

$$\frac{\partial \rho}{\partial t} + \text{div} \rho \mathbf{v} = 0, \quad (12.46)$$

$$\frac{\partial}{\partial t} \left(\frac{\rho v^2}{2} + \rho \varepsilon + \frac{B^2}{8\pi} \right) = -\text{div} \mathbf{G}, \quad (12.47)$$

$$p = p(\rho, T). \quad (12.48)$$

In contrast to (12.12), the momentum of electromagnetic field does not appear on the left-hand side of the non-relativistic equation (12.43). It is negligibly small in comparison to the plasma motion momentum ρv_α . This fact is a consequence of neglecting the displacement current in Maxwell's equations.

On the right-hand side of (12.43), the asterisk refers to the total (unlike (12.13)) momentum flux density tensor $\Pi_{\alpha\beta}^*$ which is now equal to

$$\Pi_{\alpha\beta}^* = \rho v_\alpha v_\beta + \left(p \delta_{\alpha\beta} - \sigma_{\alpha\beta}^v \right) + \frac{1}{4\pi} \left(\frac{B^2}{2} \delta_{\alpha\beta} - B_\alpha B_\beta \right). \quad (12.49)$$

There is no electric field in this formula either.

In (12.44)

$$\nu_m = \frac{c^2}{4\pi\sigma} \quad (12.50)$$

is the *magnetic diffusivity* (or magnetic viscosity). It plays the same role in (12.44) as the kinematic viscosity $\nu = \eta/\rho$ in the equation of plasma motion (12.43).

The vector $\mathbf{G}$ is defined as the energy flux density (cf. (12.18))

$$\begin{aligned} G_\alpha = & \rho v_\alpha \left(\frac{v^2}{2} + w \right) + \frac{1}{4\pi} [\mathbf{B} \times (\mathbf{v} \times \mathbf{B})]_\alpha - \\ & - \frac{v_m}{4\pi} (\mathbf{B} \times \text{rot } \mathbf{B})_\alpha - \sigma_{\alpha\beta}^v v_\beta - \kappa \nabla_\alpha T, \end{aligned} \quad (12.51)$$

where the *specific enthalpy* is

$$w = \varepsilon + \frac{p}{\rho} \quad (12.52)$$

(see definition (9.35)).

The Poynting vector appearing as a part in expression (12.51) is rewritten using formula (12.35):

$$\mathbf{G}_p = \frac{c}{4\pi} \mathbf{E} \times \mathbf{B} = \frac{1}{4\pi} \mathbf{B} \times (\mathbf{v} \times \mathbf{B}) - \frac{v_m}{4\pi} \mathbf{B} \times \text{rot } \mathbf{B}. \quad (12.53)$$

As usually in electrodynamics, the flux of electromagnetic energy equals zero when electric field $\mathbf{E}$ is parallel to magnetic field $\mathbf{B}$.

The energy flux density due to friction processes is written as the contraction of the velocity vector $\mathbf{v}$ and the viscous stress tensor

$$\sigma_{\alpha\beta}^v = \eta \left(\frac{\partial v_\alpha}{\partial r_\beta} + \frac{\partial v_\beta}{\partial r_\alpha} - \frac{2}{3} \delta_{\alpha\beta} \frac{\partial v_\gamma}{\partial r_\gamma} \right) + \zeta \delta_{\alpha\beta} \frac{\partial v_\gamma}{\partial r_\gamma} \quad (12.54)$$

(see Landau and Lifshitz 1959a, Chap. 2, Sect. 15). How should we find formula (12.54) and formulae for coefficients η and ζ ?—In order to find an equation for the second moment (9.10), we should multiply the kinetic equation (9.1) by the factor $m_k v_\alpha v_\beta$ and integrate over velocity space $\mathbf{v}$. In this way, we could derive the equations for the anisotropic part of the pressure tensor and for the flux of heat due to random motions of particles (Shkarofsky et al. 1966; Sect. 9.2). We restrict ourself just by recalling the expressions for the viscous stress tensor (12.54) and heat flux density $-\kappa \nabla T$, where κ is the plasma thermal conductivity.

12.2.4 The Energy Conservation Law

The non-relativistic MHD equations are frequently used to model solar flares, coronal mass ejections (CMEs), eruptive prominences. A key goal of such studies is to deduce how energy of magnetic fields is stored and then suddenly released to drive these phenomena. However, because most MHD models use a relatively simple energy equation (one or another), the discussion often centers on the over-simplified interpretation or simple comparison of magnetic field structure and evolution in the models with corresponding features observed in emission.

With new capabilities to study X-ray and EUV emission from *Hinode*, as well as complementary observations from *SOHO*, *RHESSI*, *STEREO*, *SDO*, and other satellites,

it now becomes imperative that new models advance to more quantitative comparisons with the emission measurements in space.

Thus we have to develop the MHD models that include radiative losses and other dissipative processes, the energy transport by anisotropic heat conduction in the solar corona and transition region between the corona and chromosphere. Such more accurate representation of energy equation is necessary for example to compute simulated EUV and X-ray emission and to compare it directly with observations.

The equation of state (12.48) can be rewritten in other thermodynamic variables. In order to do this, we have to make use of (12.43)–(12.46) and the thermodynamic identities

$$d\varepsilon = T ds + \frac{p}{\rho^2} d\rho \quad \text{and} \quad dw = T ds + \frac{1}{\rho} dp.$$

Here s is the entropy per unit mass.

At the same time, it is convenient to transform the energy conservation law (12.47) from the divergent form to the hydrodynamic one containing the substantial derivative (9.45). On rearrangement, (12.47) results in the *heat transfer equation*

$$\rho T \frac{ds}{dt} = \frac{\nu_m}{4\pi} (\text{rot } \mathbf{B})^2 + \sigma_{\alpha\beta}^v \frac{\partial v_\alpha}{\partial r_\beta} + \text{div } \kappa \nabla T + \mathcal{L}^{(rad)}(\mathbf{r}, t). \quad (12.55)$$

It shows that

the heat abundance change $dQ = \rho T ds$ in a moving element of unit volume is a sum of the Joule and viscous heating, conductive heat redistribution to neighbourhood elements, radiative cooling,

see also Exercise 12.1.

The momentum conservation law (12.43) can be also recast into the equation of plasma motion in the hydrodynamic form:

$$\frac{d\mathbf{v}}{dt} = -\frac{\nabla p}{\rho} - \frac{1}{4\pi\rho} \mathbf{B} \times \text{rot } \mathbf{B} + \frac{\eta}{\rho} \Delta \mathbf{v} + \frac{1}{\rho} \left(\zeta + \frac{\eta}{3} \right) \nabla \text{div } \mathbf{v}. \quad (12.56)$$

Once again, we see that the momentum of electromagnetic field does not appear in the non-relativistic equation of plasma motion.

12.2.5 Relativistic Magnetohydrodynamics

Relativistic MHD models are of considerable interest in several areas of modern astrophysics. The theory of gravitational collapse and models of *supernova* explosions are based on relativistic hydrodynamic models for a star. In most models a key feature is the occurrence of a relativistic shock, for example, to expel the bulk of the star. The effects of deviations from spherical symmetry due to an initial angular momentum and magnetic field require the use of relativistic MHD models.

In the theories of galaxy formation, relativistic fluid models have been used, for example, in order to describe the evolution of perturbations of the baryon and radiation components of the cosmic medium. Theories of relativistic stars are also based on *relativistic fluid* model (Zel'dovich and Novikov 1971; [Rose 1998](#)). Relativistic hydrodynamics is presumably applied to the so-called *quark-gluon plasma*, a hot soup of quarks and gluons, which is the primordial state of hadronic matter in the Universe (see Sect. 6.2 in [Letessier and Rafelski 2004](#)).

When the medium interacts electromagnetically and is highly conducting, the simplest description is in terms of relativistic MHD. From the mathematical viewpoint, the relativistic MHD was mainly treated in the framework of *general relativity*. This means that the MHD equations were studied in conjunction with Einstein's equations. [Lichnerowicz \(1967\)](#) has made a thorough and deep investigation of the initial value problem. Gravito-hydromagnetics describes one of the most fascinating phenomena in the outer space (e.g., [Punsly 2001](#)).

In many applications, however, one neglects the gravitational field generated by the conducting medium in comparison with the background gravitational field as well as in many cases one simply uses *special relativity*. Mathematically this amounts to taking into account only the *conservation laws for matter and the electromagnetic field*, neglecting Einstein's equations. Such relativistic MHD theory is much simpler than the full general relativistic theory. So more detailed results can be obtained ([Anile 1989](#); [Novikov and Frolov 1989](#); [Koide et al. 1999](#)).

Astrophysical understanding the pulsar electrodynamics comes from using modern observations and relativistic MHD theory to find and follow the flow of energy, mass and momentum. For pulsars, this has been a challenge. They are quintessential “dogs that don't bark in the night”, with only a small fraction of the energy flux the pulsars broadcast into the Universe appearing in directly observable forms. Recent successes are nicely reviewed by [Arons \(2009\)](#).

According to Arons, the unsteady *magnetic reconnection* in the current layer separating the closed from the open zones of the *pulsar magnetosphere* is responsible for the torque fluctuations observed in some pulsars, as well as for departures of the braking index from the canonical value of 3.1. The role of reconnection in the transfer of open to closed magnetic flux has begun to be assessed, and is full of promise as a part to a physical theory of pulsars.

12.3 Magnetic Flux Conservation. Ideal MHD

12.3.1 Integral and Differential Forms of the Law

Equations (12.46), (12.43), and (12.47) are the conservation laws for mass, momentum, and energy, respectively. Let us show that, with the proviso that $\nu_m = 0$, (12.44) is the magnetic flux conservation law.

Let us consider the vector $\mathbf{B}$ flux through a surface S moving with the plasma (Fig. 12.2). We are interested to know the time derivative of this flux.

According to the known formula of vector analysis (see Smirnov 1965), we have

$$\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = \int_S \left(\frac{\partial \mathbf{B}}{\partial t} + \mathbf{v} \operatorname{div} \mathbf{B} + \operatorname{rot} (\mathbf{B} \times \mathbf{v}) \right) \cdot d\mathbf{S}. \quad (12.57)$$

By virtue of (12.45), formula (12.57) is rewritten as follows

$$\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = \int_S \left(\frac{\partial \mathbf{B}}{\partial t} - \operatorname{rot} (\mathbf{v} \times \mathbf{B}) \right) \cdot d\mathbf{S},$$


or, making use of (12.44),

$$\boxed{\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = \nu_m \int_S \Delta \mathbf{B} \cdot d\mathbf{S}.} \quad (12.58)$$

Thus, if we cannot neglect magnetic diffusivity $\nu_m = c^2/4\pi\sigma$, then

the change rate of magnetic flux through a surface moving together with a conducting plasma is proportional to the magnetic diffusivity of the plasma.

Fig. 12.2 The magnetic field $\mathbf{B}$ flux through the surface S moving with a plasma with velocity $\mathbf{v}$. L is the “liquid contour” bounding the surface S



The right-hand side of formula (12.58) can be rewritten with the help of the Stokes theorem:

$$\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = -v_m \oint_L \text{rot } \mathbf{B} \cdot d\mathbf{l}. \quad (12.59)$$

Here L is the “liquid contour” bounding the surface S . We have also used here that

$$\Delta \mathbf{B} = -\text{rot rot } \mathbf{B},$$

since

$$\text{rot rot } \mathbf{B} = \nabla \text{div } \mathbf{B} - \Delta \mathbf{B} = -\Delta \mathbf{B}.$$

By using (12.34) we have

$$\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = -\frac{c}{\sigma} \oint_L \mathbf{j} \cdot d\mathbf{l}.$$

(12.60)

The change rate of flux through a surface connected with the moving plasma is proportional to the electric resistivity σ^{-1} of the plasma.

Equation (12.60) is equivalent to the differential equation (12.44) and presents an *integral* form of the magnetic flux conservation law.

The magnetic flux through any surface moving with the plasma is conserved, once the electric resistivity σ^{-1} can be ignored.

Let us clarify the conditions when it is possible to neglect electric resistivity of plasma. The relative role of a dissipation process in the *differential* equation (12.44) can be evaluated by proceeding as follows. In a spirit similar to that of Sect. 5.2, we pass on to the dimensionless variables

$$\mathbf{r}^* = \frac{\mathbf{r}}{L}, \quad t^* = \frac{t}{\tau}, \quad \mathbf{v}^* = \frac{\mathbf{v}}{v}, \quad \mathbf{B}^* = \frac{\mathbf{B}}{B_0}.$$

Therefore

$$\mathbf{r} = L \cdot \mathbf{r}^*, \quad t = \tau \cdot t^*, \quad \mathbf{v} = L \cdot \mathbf{v}^*, \dots \quad (12.61)$$

On substituting definition (12.61) into (12.44) we obtain

$$\frac{B_0}{\tau} \frac{\partial \mathbf{B}^*}{\partial t^*} = \frac{v B_0}{L} \text{rot}^* (\mathbf{v}^* \times \mathbf{B}^*) + v_m \frac{B_0}{L^2} \Delta^* \mathbf{B}^*.$$

Now we normalize this equation with respect to its left-hand side, i.e.

$$\frac{\partial \mathbf{B}^*}{\partial t^*} = \frac{v\tau}{L} \text{rot}^* (\mathbf{v}^* \times \mathbf{B}^*) + \frac{v_m \tau}{L^2} \Delta^* \mathbf{B}^*. \quad (12.62)$$

The dimensionless equation (12.62) contains two dimensionless parameters. The first one,

$$\delta = \frac{\nu\tau}{L}, \quad (12.63)$$

will be discussed later on, in Chap. 13. Here, for simplicity, we assume that $\delta = 1$.

The second parameter,

$$\text{Re}_m = \frac{L^2}{\nu_m \tau} = \frac{\nu L}{\nu_m}, \quad (12.64)$$

is termed the *magnetic* Reynolds number, by analogy with the *hydrodynamic* Reynolds number $\text{Re} = \nu L/\nu$, $\nu = \eta/\rho$ is the kinematic viscosity.

The magnetic Reynolds number characterizes the ratio of the first term on the right-hand side of (12.62) to the second one. Omitting the asterisk, we rewrite (12.62) in the *dimensionless* form

$$\frac{\partial \mathbf{B}}{\partial t} = \text{rot}(\mathbf{v} \times \mathbf{B}) + \frac{1}{\text{Re}_m} \Delta \mathbf{B}. \quad (12.65)$$

The larger the magnetic Reynolds number, the smaller the role played by magnetic diffusivity.

So the magnetic Reynolds number is the measure of a relative importance of resistivity. If $\text{Re}_m \gg 1$, we neglect the plasma resistivity and, as a consequence, not only the magnetic field *diffusion* but also magnetic field *dissipation*. This is just the same as one neglects viscosity effects under large Reynolds numbers in ordinary hydrodynamic equations: in the equation of motion and in the energy equation.

In laboratory experiments, for example in devices for studying the processes of current layer formation and rupture during magnetic reconnection (e.g., Altyntsev et al. 1977; Bogdanov et al. 1986, 2000), because of a small value L^2 , the magnetic Reynolds number is usually not large: $\text{Re}_m \sim 1 - 3$. In this case the electric resistivity has a dominant role, and *Joule dissipation* is important.

12.3.2 The Equations of Ideal MHD

Under astrophysical conditions, owing to the *low resistivity* of plasma and the enormously large length scales usually considered, the magnetic Reynolds number is usually huge: $\text{Re}_m > 10^{10}$ (for example, in the solar corona; see Exercise 12.2).

Therefore, in a great number of problems of plasma astrophysics, it is sufficient to consider plasma as a medium with *infinite conductivity*:

$$\text{Re}_m \gg 1. \quad (12.66)$$

Furthermore the usual Reynolds number can be also large (see, however, Exercise 12.3):

$$\text{Re} \gg 1. \quad (12.67)$$

Let us also assume (without much justification) the heat-conductive exchange of energy (see Exercise 12.1) to be of minor importance. This assumption is not universally true either. Sometimes the thermal conductivity (due to thermal electrons or radiation) is so effective that the astrophysical plasma behavior must be considered as isothermal, rather than adiabatic. However, conventionally,

while treating the “ideal medium”, all dissipative transfer coefficients as well as the thermal conductivity are set equal to zero in the non-relativistic MHD equations (12.43)–(12.50):

$$\nu_m = 0, \quad \eta = \zeta = 0, \quad \kappa = 0.$$

(12.68)

The complete set of the MHD equations for the ideal medium has two different (but equivalent, of course) forms. The first one (with the energy equation (12.55)) is the form of *transfer* equations:

$$\begin{aligned} \frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} &= -\frac{\nabla p}{\rho} - \frac{1}{4\pi\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \\ \frac{\partial \mathbf{B}}{\partial t} &= \text{rot } (\mathbf{v} \times \mathbf{B}), \quad \text{div } \mathbf{B} = 0, \\ \frac{\partial \rho}{\partial t} + \text{div } \rho \mathbf{v} &= 0, \quad \frac{\partial s}{\partial t} + (\mathbf{v} \cdot \nabla) s = 0, \quad p = p(\rho, s). \end{aligned} \quad (12.69)$$

The other form of ideal MHD equations is the *divergent* form which also corresponds to the *conservation laws* for energy, momentum, mass and magnetic flux:

$$\frac{\partial}{\partial t} \left(\frac{\rho v^2}{2} + \rho \varepsilon + \frac{B^2}{8\pi} \right) = -\text{div } \mathbf{G}, \quad (12.70)$$

$$\frac{\partial}{\partial t} \rho v_\alpha = -\frac{\partial}{\partial r_\beta} \Pi_{\alpha\beta}^*, \quad (12.71)$$

$$\frac{\partial \rho}{\partial t} = -\operatorname{div} \rho \mathbf{v}, \quad (12.72)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \operatorname{rot} (\mathbf{v} \times \mathbf{B}), \quad (12.73)$$

$$\operatorname{div} \mathbf{B} = 0, \quad (12.74)$$

$$p = p(\rho, s). \quad (12.75)$$

Here the energy flux density and the momentum flux density tensor are, respectively, equal to (cf. (12.51) and (12.49))

$$\mathbf{G} = \rho \mathbf{v} \left(\frac{v^2}{2} + w \right) + \frac{1}{4\pi} (B^2 \mathbf{v} - (\mathbf{B} \cdot \mathbf{v}) \mathbf{B}) \quad (12.76)$$

and

$$\Pi_{\alpha\beta}^* = p \delta_{\alpha\beta} + \rho v_\alpha v_\beta + \frac{1}{4\pi} \left(\frac{B^2}{2} \delta_{\alpha\beta} - B_\alpha B_\beta \right). \quad (12.77)$$

12.3.3 The “Frozen Field” Theorem

The magnetic flux conservation law (12.73) written in the integral form

$$\frac{d}{dt} \int_S \mathbf{B} \cdot d\mathbf{S} = 0, \quad (12.78)$$


where the integral is taken over an arbitrary surface moving together with the plasma, is quite characteristic of ideal MHD. It allows us to clearly represent the magnetic field as a set of field lines attached to the medium, as if they were “frozen into” it. For this reason, (12.73) is frequently referred to as the “freezing-in” equation.

The freezing-in property converts the notion of magnetic field line from the purely geometric to the material sphere. The MHD “frozen field” theorem can be formulated as follows.

In the ideally conducting medium, the magnetic field lines move together with the plasma. A medium motion conserves not only the magnetic flux but each of the field lines as well.

To convince ourselves that this is the case, we have to imagine a thin tube of magnetic field lines (Fig. 12.3). There is no magnetic flux through any part of the surface formed by the collection of the boundary field lines that intersect the closed curve L . The flux tube evolves in time, moving with plasma. Because of flux conservation, the “fluid particles” that are initially in the flux tube must remain in the same flux tube.

Fig. 12.3 The magnetic-field flux tube through the surface $d\mathbf{S}$ moves with a plasma with velocity $\mathbf{v}$. L is the “liquid contour” bounding the surface $d\mathbf{S}$. The “fluid particles” (FP) that are initially in this flux tube remain in the same tube



In ideal MHD flows, magnetic field lines inside the thin flux tube accompany the plasma. They are therefore materialized and are unbreakable because the flux tube links the same fluid particles. As a result its *topology cannot change*. Fluid particles which are not initially on a common field line cannot become linked by one later on.

This general topological constraint restricts the ideal MHD motions, forbidding a lot of motions that would otherwise appear.

The individual flux tubes must move exactly together with the fluid, since any relative motion between the flux tube and fluid will result in the violation of (12.73) or (12.78).

Conversely, the constraint that the thin flux tube follows the fluid particle motion, whatever its complexity, may create situations where the magnetic field structure becomes itself very complex (see Part II, Chap. 14).

In general, the field intensity $\mathbf{B}$ is a *local* quantity. However the magnetic field lines (even in vacuum) are *integral* characteristics of the field. Their analysis becomes more complicated. Nonetheless, a large number of actual fields have been studied because the general features of the morphology—an investigation of *non-local* structures—of magnetic fields are fairly important in plasma astrophysics.

The *geometry* of the field lines appears in different ways in the equilibrium criteria for astrophysical plasma. For example, much depends on whether the field lines are concave or convex, on the value of the gradient of the so-called *specific volume* of magnetic flux tubes (Chap. 19). However even more depends on the presence of X-type points (Sect. 14.3) as well as on a number of other *topological* characteristics, for example, the global *magnetic helicity* (see Part II, Chap. 12).

12.4 Magnetic Reconnection

12.4.1 General Properties

Reconnection of magnetic field lines is the fundamental physical process which involves a breakdown of the MHD “frozen field” theorem (Sect. 12.3.3). The effects of electric resistivity, normally negligible in the large, *become locally dominant* with dramatic consequences in the nature of the large-scale plasma flows and magnetic field configuration which could not be achieved otherwise (see Fig. 8.5). *Reconnection changes topology* of magnetic field. Such a process is mathematically singular and in most astrophysical situations highly non-linear. Its implications will be easy to grasp conceptually in Part II.

The origin of the concept of magnetic reconnection lies in an attempt by Giovanelli (1946, 1947) to explain *solar flares* in terms of a fundamental feature of plasma. This is an interplay of energy forms that can occur within a plasma, between the energy of fields on one hand and the kinetic energy (thermal and non-thermal) of particles on the other.

Reconnection is the means by which the magnetic topologies are determined in astrophysical plasma and by which energy stored in magnetic fields is released rapidly

to produce such phenomena as solar flares and magnetospheric substorms. Furthermore, reconnection has been found to play important roles in many areas of plasma astrophysics (see Part II).

12.4.2 Reconnection in the Solar Wind

Depending on complexity of magnetic fields and boundary conditions, the reconnection process can occur over an extended region in space or can be *patchy* and “unpredictable in nature” (Nishida 1989). For example, in the Earth’s magnetosphere the *reconnecting current layers* (RCLs) are formed by the interaction between the solar wind and the geomagnetic field (see Part II, Sect. 10.2). Such RCLs have finite extents, and their boundary conditions (determined mainly by the solar wind magnetic field) often change rapidly.

On the contrary,

in the solar wind, the magnetic field orientations on the two sides of the interplanetary current layers are usually well defined, and the boundary conditions seem to be relatively stable.

Phan et al. (2006) report the three-spacecraft observations of accelerated plasma flow associated with large-scale reconnection in a RCL embedded in the solar wind and extended at least 390 Earth radii R_E (or 2.5×10^{11} cm) as illustrated by Fig. 12.4.




Fig. 12.4 Reconnection in the solar wind. Diagram encounters the three spacecraft: *Wind*, *ACE* and *Cluster* on 2 February 2002. The spacecraft positions are shown in units of Earth radius (R_E) and in geocentric solar ecliptic (GSE) coordinates with the x -axis pointing from Earth to Sun, the y -axis pointing towards dusk and the z -axis is perpendicular to the ecliptic plane (in yellow). (Reproduced with permission © NATURE Publishing Group)

Reconnection in the RCL (in blue) occurs at the X-line between magnetic field lines with the large-scale anti-parallel components $B_{L,1}$ and $B_{L,2}$; $V_{in,1}$ and $V_{in,2}$ are the inflow velocities. The resulting bi-directional plasma jets (confined to the reconnection “exhausts”) are observed far from the X-line. Also shown is the LMN current layer coordinate system, with N along the layer normal, M along the X-line direction and L along the anti-parallel magnetic field direction. The thick solid red line is the portion of the X-line whose effect is observed by the three spacecraft. The solid orange lines denote the spacecraft trajectory relative to the solar wind. The angle of the diverging exhausts is exaggerated for illustration; the actual angle is $\sim 4^\circ$. B_M is the magnetic field component along the X-line.

Observations of this and 27 similar events imply that reconnection is fundamentally a large-scale physical process (Phan et al. 2006). Patchy reconnection observed in the Earth’s magnetosphere is therefore likely to be a geophysical effect associated with rapidly changing boundary conditions, rather than a fundamental property of reconnection.




Fig. 12.5 A model of a wavy reconnecting current layer (RCL). Traditional view of the solar wind origins suggests that the fast wind (FW) blows along the open magnetic field lines, whereas the slow wind (SW) emanates from the coronal streamers

In the most astrophysical situations the magnetic reconnection process is presumably predictable in nature and occurs in an internal scale of a phenomenon, which is responsible to the initial and boundary conditions

(see Part II). In the solar wind the scale of a reconnecting current layer (RCL) around the Sun can be very large (Fig. 12.5).

This model explains the sudden changes of the interplanetary magnetic fields (**B** in Fig. 12.5) observed in the solar wind. The reconnecting current layer (RCL) separates the magnetic fields of nearly opposite directions. The electric currents **j** flow azimuthally in this layer. The average plane of the layer is approximately the plane of the equator of the Sun's average magnetic dipole field. Thus the magnetic equator (**M** in Fig. 12.5) is tilted with respect to its equator of rotation. The polarity change occurs when an observer (a spacecraft) crosses the heliospheric current layer; e.g., [Smith et al. \(1978\)](#).

[Gosling et al. \(2007b\)](#) presented another convincing evidence for prolonged (at least 5 h) magnetic reconnection at a continuous X-line in the solar wind, observed by five spacecrafts of a broad (1.85×10^6 km) reconnection exhaust within the heliospheric current layer. At least 1.2×10^{24} ergs of magnetic energy were extracted from the CL in this event and converted to kinetic and thermal energy of the exhaust plasma. The reconnection produced the field lines disconnected from the Sun, thereby slightly reducing the amount of open magnetic flux present in the heliosphere.

On the other hand, the high-speed solar wind that originates in coronal holes is permeated by evolved Alfvénic-type fluctuations associated with MHD turbulence. The spacecraft *Wind* at 1 AU allows us to study reconnection in this turbulent flow

(Gosling 2007). The *Wind* observations demonstrate that reconnection is one way in which the solar wind turbulence is dissipated and the high-speed wind is heated far from the Sun.

In the solar wind, the kinetic and thermal energies of plasma exceed the magnetic energy density. It means that, in the first consideration, we can neglect the magnetic force as compared to the inertia force of moving plasma and its pressure gradient. This is valid everywhere except the current layers discussed above. Here all three forces can become approximately equal, and magnetic reconnection can operate in a quasi-stationary manner. This assumption seems to be well consistent with spacecraft observations (Phan et al. 2006). We shall call such a process as reconnection in a *weak magnetic field*. A particular example of this phenomenon is the photospheric reconnection (see Part II, Sect. 15.2).

Reconnection in a *strong magnetic field* is a fundamental feature of astrophysical plasmas like a plasma of the solar corona. Such reconnection explains an accumulation of magnetic energy and a sudden release of this energy, an original *electrodynamical explosion* or *flare*. This phenomenon is accompanied by fast ejections of plasma, powerful flows of heat and hard electromagnetic radiation as well as by acceleration of charged particles to high energies (see Part II).

12.5 Practice: Exercises and Answers

Exercise 12.1 (Sect. 12.2.4). Show that

heat conduction redistributes thermal energy conserving its total amount in a plasma but increases the total entropy.

Answer. Let us consider (12.55) taking into account only the heat-conductive term on the right-hand side in the one-dimensional approximation:

$$\rho T \frac{ds}{dt} = \frac{\partial}{\partial x} \left(\kappa \frac{\partial T}{\partial x} \right). \quad (12.79)$$

Now we integrate this equation from point x_1 to point x_2 that restrict a volume V occupied by a plasma. We find the time derivative of the total thermal energy of the plasma inside the volume V

$$\frac{dQ}{dt} = \int_{x_1}^{x_2} \rho T \frac{ds}{dt} dx = \kappa \frac{\partial T}{\partial x} \Big|_{x_1}^{x_2} = 0, \quad (12.80)$$

if the plasma is thermally isolated at the boundaries x_1 and x_2 . Thus heat conduction does redistribute the thermal energy inside the plasma but conserves the total thermal energy.

In order to consider the change of total entropy let us divide (12.79) by temperature T and integrate it from point x_1 to point x_2 by parts:

$$\int_{x_1}^{x_2} \rho \frac{ds}{dt} dx = \frac{1}{T} \kappa \frac{\partial T}{\partial x} \Big|_{x_1}^{x_2} + \int_{x_1}^{x_2} \frac{\kappa}{T^2} \left(\frac{\partial T}{\partial x} \right)^2 dx. \quad (12.81)$$

The first term on the right-hand side equals zero but the second one is always positive. Therefore

the heat-conductive redistribution of thermal energy inside a plasma increases its total entropy.

The increase in entropy indicates that irreversible process occurs in a plasma.

Exercise 12.2 (Sect. 12.3.2). Estimate the magnetic diffusivity and the magnetic Reynolds number under typical conditions in the *solar corona*.

Answer. Let us take characteristic values of the parallel conductivity as they were estimated in Exercise 11.1:

$$\sigma_{\parallel} = \sigma \sim 10^{16} - 10^{17} \text{ s}^{-1}.$$

Substituting these values in formula (12.50) we obtain

$$\nu_m = \frac{c^2}{4\pi\sigma} \approx 7.2 \times 10^{19} \frac{1}{\sigma} \sim 10^3 - 10^4 \text{ cm}^2 \text{ s}^{-1}. \quad (12.82)$$

According to definition (12.64) the magnetic Reynolds number

$$\text{Re}_m = \frac{\nu L}{\nu_m} \sim 10^{11} - 10^{12}, \quad (12.83)$$

if the characteristic values of length and velocity, $L \sim 10^4 \text{ km} \sim 10^9 \text{ cm}$ and $\nu \sim 10 \text{ km s}^{-1} \sim 10^6 \text{ cm s}^{-1}$, are taken for the corona. Thus the ideal MHD approximation can be well used to consider, for example, magnetic field diffusion in coronal linear scales.

Exercise 12.3 (Sect. 12.3.2). Show that

in the solar corona, usual viscosity of plasma can be a much more important dissipative mechanism than its electric resistivity.

Answer. By using the formula (10.31) for viscosity, let us estimate the value of *kinematic viscosity* in the solar corona:

$$\nu = \frac{\eta}{\rho} \approx 3 \times 10^{15} \text{ cm}^2 \text{ s}^{-1}. \quad (12.84)$$

Here $T_p \approx 2 \times 10^6$ K and $n_p \approx n_e \approx 2 \times 10^8 \text{ cm}^{-3}$ have been taken as the typical proton temperature and density.

If the characteristic values of length and velocity, $L \sim 10^9 \text{ cm}$ and $v \sim 10^6 \text{ cm s}^{-1}$, are taken (see Exercise 12.1), then the hydrodynamic Reynolds number

$$\text{Re} = \frac{vL}{\nu} \sim 0.3. \quad (12.85)$$

This value demonstrates the potential importance of usual viscosity in the solar corona. A comparison between (12.85) and (12.83) shows that

$$\text{Re}_m \gg \text{Re}.$$

Clearly, the viscous effects can dominate the effects of electric resistivity in coronal plasma.

Chapter 13

Magnetohydrodynamics in Astrophysics

Abstract Magnetohydrodynamics (MHD) is appropriate for many phenomena in astrophysical plasma, that take place on a relatively large scale. In the non-relativistic limit, MHD is applied to dynamo theory, flows in the solar atmosphere, prominences, flares, coronal heating, solar and stellar winds. Relativistic MHD describes well accretion disks near relativistic objects, and relativistic jets.

13.1 The Main Approximations in Ideal MHD

13.1.1 Dimensionless Equations

The equations of MHD, even the ideal MHD, constitute a set of nonlinear differential equations in partial derivatives. The order of the set is rather high, and its structure is complicated. To formulate a problem in the context of MHD, we have to know the initial and boundary conditions admissible by this set of equations. To do this, in turn, we have to know the type of these equations, in the sense adopted in mathematical physics (see [Vladimirov 1971](#)).

To formulate a problem, we usually use one or another approximation, which makes it possible to point up and isolate the main effect—the essence of a phenomenon. For instance, if the magnetic Reynolds number is small, then the plasma moves comparatively easily with respect to the magnetic field. This is the case in MHD generators and other laboratory and technical devices ([Sutton and Sherman 1965](#), Sect. 1.3; [Shercliff 1965](#), Sect. 6.5).

The opposite approximation is that of large magnetic Reynolds numbers, when magnetic field “freezing in” takes place in the plasma (see Sect. [12.3.2](#)). Obviously, the transversal (with respect to the magnetic field) plasma flows are implied. For any flow along the field, [\(12.73\)](#) holds. This approximation is quite characteristic of the astrophysical plasma dynamics.

How can we isolate the main effect in a physical phenomenon and correctly formulate the problem?—From the above examples concerning the magnetic Reynolds number, the following rule suggests itself:

take the dimensional parameters characterizing the phenomenon at hand, combine them into dimensionless combinations and then, on calculating their numerical values, make use of the corresponding approximation in the set of *dimensionless* equations.

Such an approach is effective in hydrodynamics (Sedov 1973, vol. 1). However the following procedure is more efficient in MHD.

Let us start with the set of the ideal MHD equations (12.69):

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla p}{\rho} - \frac{1}{4\pi\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \quad (13.1)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \text{rot} (\mathbf{v} \times \mathbf{B}), \quad (13.2)$$

$$\frac{\partial \rho}{\partial t} + \text{div } \rho \mathbf{v} = 0, \quad (13.3)$$

$$\frac{\partial s}{\partial t} + (\mathbf{v} \cdot \nabla) s = 0, \quad (13.4)$$

$$\text{div } \mathbf{B} = 0, \quad (13.5)$$

$$p = p(\rho, s). \quad (13.6)$$

Let the quantities L , τ , v , ρ_0 , p_0 , s_0 , and B_0 be the characteristic values of length, time, velocity, density, pressure, entropy and field strength, respectively. Rewrite (13.1)–(13.6) in the dimensionless variables

$$\mathbf{r}^* = \frac{\mathbf{r}}{L}, \quad t^* = \frac{t}{\tau}, \quad \mathbf{v}^* = \frac{\mathbf{v}}{v}, \dots \quad \mathbf{B}^* = \frac{\mathbf{B}}{B_0}.$$

Therefore, in (13.1)–(13.6)

$$\mathbf{r} = L \cdot \mathbf{r}^*, \quad t = \tau \cdot t^*, \quad \mathbf{v} = L \cdot \mathbf{v}^*, \dots \quad (13.7)$$

After omitting the asterisk, we obtain the following set of equations in dimensionless variables (Somov and Syrovatskii 1976b):

$$\varepsilon^2 \left\{ \frac{1}{\delta} \frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right\} = -\gamma^2 \frac{\nabla p}{\rho} - \frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \quad (13.8)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \delta \text{rot} (\mathbf{v} \times \mathbf{B}), \quad (13.9)$$

$$\frac{\partial \rho}{\partial t} + \delta \operatorname{div} \rho \mathbf{v} = 0, \quad (13.10)$$

$$\frac{\partial s}{\partial t} + \delta (\mathbf{v} \cdot \nabla) s = 0, \quad (13.11)$$

$$\operatorname{div} \mathbf{B} = 0, \quad (13.12)$$

$$p = p(\rho, s). \quad (13.13)$$

Here

$$\delta = \frac{v\tau}{L}, \quad \varepsilon^2 = \frac{v^2}{V_A^2}, \quad \gamma^2 = \frac{p_0}{\rho_0 V_A^2} \quad (13.14)$$

are three dimensionless parameters characterizing the problem;

$$V_A = \frac{B_0}{\sqrt{4\pi\rho_0}} \quad (13.15)$$

is the characteristic value of the Alfvén speed (see Exercise 13.1).

If the gravitational force were taken into account in the equation of motion (13.1), then (13.8) would contain another dimensionless parameter, gL/V_A^2 , where g is the gravitational acceleration. The analysis of these parameters allows us to separate the approximations which are possible in the ideal MHD.

13.1.2 Weak Magnetic Fields in Astrophysical Plasma

We begin with the assumption that

$$\varepsilon^2 \gg 1 \quad \text{and} \quad \gamma^2 \gg 1. \quad (13.16)$$

As is seen from (13.8), in the zero-order approximation relative to the small parameters ε^{-2} and γ^{-2} , we can neglect the magnetic force as compared to the inertia force and the gas pressure gradient. In subsequent approximations, the magnetic effects are treated as small corrections to the hydrodynamic ones.

A lot of problems of plasma astrophysics are solved in this approximation, termed the *weak* magnetic field approximation. Among the simplest of them are the ones concerning the weak field's influence on hydrostatic equilibrium. An example is the problem of the influence of poloidal and toroidal magnetic fields on the equilibrium of a self-gravitating plasma ball (a star, the magnetoid of quasar's kernel etc., see examples in Sect. 19.1.3).

Some other problems are in fact analogous to the previously mentioned ones. They are called *kinematic* problems, since

they treat the influence of a given plasma flow on the magnetic field; the reverse influence is considered to be negligible.

Such problems are reduced to finding the magnetic field distribution resulting from the known velocity field. An example is the problem of magnetic field amplification and support by stationary plasma flows (the *magnetic dynamo* problem) or turbulent amplification. The simplest example is the problem of magnetic field amplification by plasma *differential rotation* (Elsasser 1956; Moffatt 1978).

For example, a neutron star undergoes vigorous *convection* soon after its formation. When coupled with rapid rotation, this makes the star a likely site for dynamo action. The neutron stars with unusually strong magnetic fields (up to $\sim 10^{15}$ G) called *magnetars* can form when conditions for efficient helical dynamo are met during the first few seconds after gravitational collapse (Duncan and Thompson 1992). Such neutron stars initially rotate with short periods (~ 1 ms) but quickly lose most of their rotational energy via magnetic braking, giving a large energy boost to the associated *supernova* explosion. The main observational signature of magnetars is *gamma-ray bursts* powered by their vast reservoirs of magnetic energy.

A leading candidate to explain the origin of large-scale magnetic fields in astrophysical plasma is the mean-field *turbulent magnetic dynamo* theory (Moffatt 1978; Parker 1979; Zel'dovich et al. 1983; Rüdiger and Rekowski 1998). The theory appeals to a combination of helical turbulence (leading to the so-called α effect), differential rotation (the Ω effect) and turbulent diffusion to exponentiate an initial seed mean magnetic field. The total magnetic field is split into a mean component and a fluctuating component, and the rate of growth of the mean field is sought.

The mean field grows on a length scale much larger than the outer scale of the turbulent velocity, with a growth time much larger than the eddy turnover time at the outer scale. A combination of kinetic and magnetic helicities provides a statistical correlation of small-scale loops favorable to exponential growth. Turbulent diffusion is needed to redistribute the amplified mean field rapidly to ensure a net mean flux gain inside the system of interest (a star or galaxy). Rapid growth of the fluctuating field necessarily accompanies the mean-field dynamo. Its impact upon the growth of the mean field, and the impact of the mean field itself on its own growth are controversial and depends crucially on the boundary conditions (see Blackman and Field 2000).

More recent models of large-scale dynamos include also such phenomena as the large-scale velocity shear (which creates anisotropy of turbulence), randomly forced shearing waves (that drive a large-scale dynamo), kinetic helicity fluctuations, etc. (e.g., Elperin et al., 2007; Rogachevskii and Kleeorin 2007; Kleeorin et al. 2009; Heinemann et al. 2011). Corresponding dynamo models for particular astrophysical bodies (including the Earth, the Sun, the Milky Way, etc.) result in magnetic field

configurations that are compatible with available phenomenology at least in a crude approximation. On the other hand, some important features in astrophysical dynamos still remain unclear.

13.1.3 Strong Magnetic Fields in Plasma

The opposite approximation—that of the *strong* magnetic field—has been less well studied because it is more complicated. It reflects the specificity of MHD to a greater extent than the weak field approximation. The strong field approximation is valid when the *magnetic force*

$$\mathbf{F}_m = -\frac{1}{4\pi} \mathbf{B} \times \text{rot } \mathbf{B} \quad (13.17)$$

dominates all the others (inertia force, pressure gradient, gravity, etc.). Within the framework of (13.8), the magnetic field is referred to as a strong one if in some region under consideration

$$\varepsilon^2 \ll 1 \quad \text{and} \quad \gamma^2 \ll 1, \quad (13.18)$$

i.e. if the magnetic energy density greatly exceeds that of the kinetic and thermal energies:

$$\frac{B_0^2}{8\pi} \gg \frac{\rho_0 v^2}{2} \quad \text{and} \quad \frac{B_0^2}{8\pi} \gg 2n_0 k_B T_0.$$

From (13.8) it follows that, in the zeroth order with respect to the small parameters (13.18), the magnetic field is *force-free*, i.e. it obeys the equation

$$\mathbf{B} \times \text{rot } \mathbf{B} = 0. \quad (13.19)$$

This conclusion is quite natural:

if the magnetic force dominates all the others, then the magnetic field must balance itself in the region under consideration.

Condition (13.19) obviously means that electric currents flow parallel to magnetic field lines. If, in addition, electric currents are absent in some region (in the zeroth approximation relative to the small parameters ε^2 and γ^2), then the strong field is simply *potential*:

$$\text{rot } \mathbf{B} = 0, \quad \mathbf{B} = \nabla \psi, \quad \Delta \psi = 0. \quad (13.20)$$

In principle, the magnetic field can be force-free or even potential for another reason: due to the equilibrium of non-magnetic forces. However this does not happen frequently.

Let us consider the *first order* in the small parameters (13.18). If they are not equally significant, i.e. $\varepsilon^2 \neq \gamma^2$, then there are two possibilities.

(a) We suppose, at first, that

$$\varepsilon^2 \ll \gamma^2 \ll 1. \quad (13.21)$$

Then we neglect the inertia force in (13.8) as compared to the gas pressure gradient. Decomposing the magnetic force into a *magnetic tension* force and a *magnetic pressure* gradient force (see Exercises 13.2 and 13.3),

$$\mathbf{F}_m = -\frac{1}{4\pi} \mathbf{B} \times \text{rot } \mathbf{B} = \frac{1}{4\pi} (\mathbf{B} \cdot \nabla) \mathbf{B} - \nabla \frac{B^2}{8\pi}, \quad (13.22)$$

we obtain the following dimensionless equation:

$$(\mathbf{B} \cdot \nabla) \mathbf{B} = \nabla \left(\frac{B^2}{2} + \gamma^2 p \right). \quad (13.23)$$

Owing to the presence of the gas pressure gradient, the magnetic field differs from the force-free one at any moment of time:

the magnetic tension force $(\mathbf{B} \cdot \nabla) \mathbf{B}/4\pi$ must balance not only the magnetic pressure gradient but that of the gas pressure as well.

Obviously the effect is proportional to the small parameter γ^2 .

This approximation can be naturally called the *magnetostatics* since $\mathbf{v} = 0$. It effectively works in regions of a strong magnetic field where the gas pressure gradients are large, for example, in coronal loops and *reconnecting current layers* (RCLs) in the solar corona (Exercise 13.4). The most fundamental properties of magnetostatic equilibrium are considered in Sects. 19.3 and 19.5.

(b) The *inertia force* also causes the magnetic field to deviate from the force-free one:

$$\varepsilon^2 \left\{ \frac{1}{\delta} \frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right\} = -\frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}. \quad (13.24)$$

Here we ignored (in the first order) the gas pressure gradient as compared with the inertia force. Thus it is not the relation (13.21) between the small parameters (13.18), but rather its converse, that should be obeyed, i.e.

$$\gamma^2 \ll \varepsilon^2 \ll 1. \quad (13.25)$$

The problems on plasma flows in a strong magnetic field are of great interest in plasma astrophysics. To solve them, inequalities (13.25) can be assumed to hold. Then we can use (13.24) as the MHD equation of motion. The approximation corresponding to inequalities (13.25) is naturally termed the approximation of *strong field* and *cold* plasma.

The main applications of the strong-field-cold-plasma approximation are concerned with the solar atmosphere (see Part II, Chaps. 2 and 8) and the Earth's magnetosphere. Both astrophysical objects are well studied from the observational viewpoint. So we can proceed with confidence from qualitative interpretation to the

construction of quantitative models. The presence of a sufficiently strong magnetic field and a comparatively rarefied plasma is common for both phenomena. This justifies the applicability of the approximation at hand.

■ A sufficiently strong magnetic field easily moves a comparatively rarefied plasma in many non-stationary phenomena in space.

Analogous conditions are reproduced under laboratory modeling of these phenomena (e.g., [Hoshino et al. 2001](#)). Some other astrophysical applications of the strong-field-cold-plasma approximation will be discussed in the following two sections.

* * *

In closing, let us consider the dimensionless parameter $\delta = \nu\tau/L$. As is seen from (13.24), it characterizes the relative role of the local $\partial/\partial t$ and transport $(\mathbf{v} \cdot \nabla)$ terms in the substantial derivative d/dt .

If $\delta \gg 1$ then, in the zeroth approximation relative to the small parameter δ^{-1} , the plasma flow can be considered to be *stationary*

$$\varepsilon^2 (\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}. \quad (13.26)$$

If $\delta \ll 1$, i.e. plasma displacement is small during the magnetic field change, then the transport term $(\mathbf{v} \cdot \nabla)$ can be ignored in the substantial derivative and the equation of motion in the strong-field-cold-plasma approximation takes the form

$$\varepsilon^2 \frac{\partial \mathbf{v}}{\partial t} = -\frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \quad (13.27)$$

other equations becoming linear. This case corresponds to small plasma displacements from the equilibrium state, i.e. small perturbations. (If need be, the right-hand side of (13.27) can be linearized in the usual way.)

Generally the parameter $\delta \approx 1$ and the set of MHD equations in the approximation of strong field and cold plasma for ideal medium assumes the following dimensionless form:

$$\varepsilon^2 \frac{d\mathbf{v}}{dt} = -\frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \quad (13.28)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \text{rot}(\mathbf{v} \times \mathbf{B}), \quad (13.29)$$

$$\frac{\partial \rho}{\partial t} + \text{div } \rho \mathbf{v} = 0. \quad (13.30)$$

In the next chapter we shall consider some continuous plasma flows in a strong magnetic field, which are described by (13.28)–(13.30).

13.2 Accretion Disks of Stars

13.2.1 Angular Momentum Transfer in Binary Stars

Magnetic fields were discussed as a possible means of angular transport in the development of *accretion disk* theory in the early seventies (Shakura and Sunyaev 1973; Novikov and Thorne 1973). Interest in the role of magnetic fields in binary stars steadily increased after the discovery of the nature of AM Herculis (see Sect. 13.2.2). It appeared that the optical counterpart of the soft X-ray source has linear and circular polarization in the V and I spectral bands, of a strength an order of magnitude larger than previously observed in any object. This suggested the presence of a very strong (as thought at that time) field, with $B \sim 10^8$ G, assuming the fundamental cyclotron frequency to be observed.


Other similar systems were soon discovered. Evidence for strong magnetic fields was subsequently found in the X-ray binary pulsars and the *intermediate polar binaries* (see Sect. 13.2.2), both believed to include accretion disks. A magnetically channeled wind from the main sequence star has been invoked to explain the higher rates of mass transfer observed in binaries above the period gap, and in an explanation of the gap. The winds from accretion disks have been suggested as contributing to the inflow by removing angular momentum.

Magnetohydrodynamics in binary stars is now an area of central importance in stellar astrophysics (Campbell 1997; Rose 1998). Magnetic fields are believed to play a role even in apparently non-magnetic binaries. They provide the most viable means, through the so-called *shear-type instabilities*, of generating the MHD turbulence in an accretion disk necessary to drive the plasma inflow via the resulting magnetic and viscous stresses.

The fundamental problem is the role of magnetic fields in redistributing angular momentum in binary stars. The disk is fed by the plasma stream originated in the L_1 region (Fig. 13.1) of the secondary star. In a steady state,

plasma is transported through the disk at the rate it is supplied by the stream and the angular momentum will be advected outwards.

Fig. 13.1 The standard model of a binary stellar system with an accretion disk. The tidally and rotationally distorted secondary star SS loses plasma from the unstable L_1 point. The resulting plasma stream feeds an accretion disk D , centered on the primary star



Angular momentum advection requires coupling between rings of rotating plasma; the ordinary hydrodynamic viscosity is too weak to provide this. Hence some form of *anomalous viscosity* must be invoked to explain the plasma flow through the disk.

The key point is the recognition that a simple linear instability, which we refer to here as the standard *magnetorotational* instability (Hawley et al. 1995), generates MHD turbulence. This turbulence transports angular momentum outward through the disk, allowing accretion to proceed. Although turbulence seems like a natural and straightforward transport mechanism, it turns out that the *magnetic fields are essential*. Purely hydrodynamic turbulence is not self-sustaining and does not produce sustained outward transport of angular momentum (Hawley and Balbus 1999). MHD turbulence greatly enhances angular momentum transport associated with the so-called α -disks (Balbus and Papaloizou 1999).

It is most probable that the accretion disks have turbulent motions generated by the shear instabilities. The turbulence and strong radial shear lead to the generation and maintenance of a large-scale magnetic field.

Turbulent viscous and magnetic stresses cause radial advection of the angular momentum via the azimuthal forces.

Provided these forces oppose the large-scale azimuthal motion, plasma will spiral in through the disk as angular momentum flows outwards. Presumably, the approximation of a weak field (Sect. 13.1.2) can be used inside the disk to model these effects. Most models to date involve a vertically averaged structure. The future aim is to find 3D solutions which self-consistently incorporate the magnetic shear instabilities and vertical structure.

The stellar spin dynamics and stability are also important, of course. For example, in spin evolution calculations, a compact white dwarf, or neutron star, is usually treated as a rigid body. This is valid provided the dynamic time-scale for adjustments in the stellar structure is short compared to the spin evolution time scale. In general, however, a strongly-magnetic primary star may experience significant distortions from spherical symmetry due to non-radial internal magnetic forces. This fact can be demonstrated by the tensor virial theorem in MHD (Sect. 19.1.3).

13.2.2 Magnetic Accretion in Cataclysmic Variables

Cataclysmic variables (CVs) are the binary systems composed of a white dwarf (primary star) and a late-type, main-sequence companion (secondary star). The secondary star fills its Roche lobe, and plasma is transferred to the compact object through the inner Lagrangian point (see point L_1 in Fig. 13.1). The way this plasma falls towards the primary depends on the intensity of a magnetic field of the white dwarf.

If the magnetic field is weak, the mass transfer occurs through an optically thick accretion disk. Such CVs are classified as non-magnetic ones.

The strong magnetic field ($B \gtrsim 10^7$ G) may entirely dominate the geometry of the accretion flow. The magnetic field is strong enough to synchronize the white dwarf rotation (spin) with the orbital period. Synchronization occurs when the magnetic torque between primary and secondary overcomes the accretion torque, and no disk is formed. Instead, the magnetic field channels accretion towards its polar regions. Such synchronous systems are known as AM Herculis binaries or *polars*.

The *intermediate* ($B \sim 2 - 8 \times 10^6$ G) magnetic field primary stars harbor magnetically truncated accretion disks which can extend until magnetic pressure begins to dominate. A shock should appear when the plasma streams against the white dwarf's magnetosphere. The shock should occur close to the corotation radius (the distance from the primary at which the Keplerian and white dwarf angular velocities match), inside and above the disk plane. Presumably the plasma is finally accreted onto the magnetic poles of the white dwarf. The asynchronous systems are known as DQ Herculis binaries or *intermediate polars* (IPs).

General properties of plasma flows driven by a strong magnetic field will be discussed in Chap. 14.

The accretion geometry strongly influences the emission properties at all wavelengths and its variability. The knowledge of the behavior in all energy domains can allow one to locate the different accreting regions (Bianchini et al. 1995). Reid et al. (2001) discovered the first magnetic white dwarf of the spectral type DZ, which shows lines of heavy elements like Ca, Mg, Na, and Fe. The cool white dwarf LHS 2534 offers the first empirical data in an astrophysical setting of the Zeeman effect on neutral Na, Mg, and both ionized and neutral Ca. The Na I splittings result in a mean surface field strength estimate of 1.92×10^6 G. In fact, there are direct laboratory measurements of the Na I D lines that overlap this field strength.

13.2.3 Accretion Disks Near Black Holes

In the binary stars discussed above, there is an abundance of evidence for the presence of accretion disks: (a) double-peaked emission lines are observed, (b) eclipses of an extended light source centered on the primary occur, and (c) in some cases eclipses of the secondary star by the disk are also detected. The case for the presence of accretion disks in *active galactic nuclei* (AGN) is less clear. Nonetheless the disk-fed accretion onto a super-massive black hole is the commonly accepted model for these astronomical objects (see Fig. 7.7 in Bradt 2008). In fact, AGN also exhibit the classical double-peaked, broad emission lines which are considered to be characteristic for a rotating disk.

As the plasma accretes in the gravitational field of the central mass, magnetic field lines are convected inwards, amplified and finally deposited on *the horizon of the black hole* (Sect. 8.3.5). As long as a magnetic field is confined by the disk, a

differential rotation causes the field to wrap up tightly (see Sect. 20.1.5), becoming highly sheared and predominantly azimuthal in orientation. A dynamo in the disk may be responsible for the maintenance and amplification of the magnetic field.

In the standard model of an accretion disk (Shakura and Sunyaev 1973; Novikov and Thorne 1973), the gravitational energy is locally radiated from the optically thin disk, and the plasma keeps its Keplerian rotation. However *the expected power far exceeds the observed luminosity*.

There are two possible explanations for the low luminosities of nearby black holes: (a) the accretion occurs at extremely low rates, or (b) the accretion occurs at low radiative efficiency. *Advection* has come to be thought of as an important process and results in a structure different from the standard model. The advection process physically means that

the energy generated via viscous dissipation is restored as entropy of the accreting plasma flow rather than being radiated.

The advection effect can be important if the radiation efficiency decreases under these circumstances (Sect. 8.3.5). An optically thin advection-dominated accretion flow (ADAF) seemed to be a hydrodynamic model that can reproduce the observed hard spectra of black hole systems such as AGN and Galactic black hole candidates (e.g., Manmoto 2000).

This situation is perhaps best illustrated by the case of nearby elliptical galaxy nuclei (Di Matteo et al. 2000). Assuming that the accretion occurs primarily from the hot, quasi-spherical interstellar medium (ISM), the Bondi (1952) theory can be used to estimate the accretion rates onto the supermassive black holes. Such estimates, however, require accurate measurements of both the density and the temperature of the ISM at the Bondi accretion radius, i.e., the radius at which the gravitational force of the black hole begins to dominate the dynamics of the hot plasma.

In order to determine unambiguously whether or not the low luminosities of nearby black holes are due to a low radiative efficiency in the accreting plasma, it is also necessary to measure the nuclear power. When combined with the estimated accretion rates, this gives us a direct measurement of the radiative efficiency η_r .

Thanks to its high spatial resolution and sensitivity, the *Chandra X-ray Observatory* is able, for the first time, to detect nuclear X-ray point sources in nearby galaxies and provide us with direct measurements of their luminosities. *Chandra* also allows us to measure the central temperatures and densities of the ISM close to accretion radii of the central black holes and therefore to determine the Bondi accretion rates in these systems to much greater accuracy than before.

Di Matteo et al. (2001) explored the implications of *Chandra* observations of the giant elliptical galaxy NGC 6166. They show that, if the central black hole of $\sim 10^9 M_\odot$ is fed at the estimated Bondi rate, the inferred efficiency $\eta_r \lesssim 10^{-5}$. At the given accretion rate, ADAF models can explain the observed nuclear luminosity. However the presence of *fast outflows* in the accretion flow is also consistent with the present constraints. The power output from the jets in NGC 6166 is also important to the energetics of the system.

13.2.4 Flares in Accretion Disk Coronae

Following the launch of several X-ray satellites, astrophysicists have tried to observe and analyze the violent variations of high energy flux from black hole candidates (e.g., [Negoro et al. 1995](#); see also review in [Di Matteo et al. 1999](#)). So far, similar solar and astrophysical statistical studies have been done almost independently of each other. [UeNo \(1998\)](#) first compared X-ray light curves from the solar corona and from the accretion disk in Cyg X-1, a famous black hole candidate. He analyzed also the power spectral densities, the peak interval distributions (the interval of time between two consecutive flares), and the peak intensity distributions.

It has appeared that

there are many relationships between flares in the solar corona and “X-ray shots” in accretion disks.

(Of course, there are many differences and unexplained features.) For example, the peak interval distribution of Cyg X-1 shows that the occurrence frequency of large X-ray shots is reduced. A second large shot does not occur soon after a previous large shot. This suggests the existence of energy-accumulation structures, such as non-potential magnetic fields in the solar corona.

It is likely that accretion disks have a corona. [Galeev et al. \(1979\)](#) suggested that the corona is confined in strong magnetic loops which have buoyantly emerged from the disk. Buoyancy constitutes a mechanism able to channel a part of the energy released in the accretion process directly into the corona outside the disk.

Magnetic reconnection of buoyant fields in the lower density surface regions may supply the energy source for a hot corona.

On the other hand, the coronal magnetic field can penetrate the disk and is stressed by its motions. The existence of a disk corona with a *strong* field (Sect. 13.1.3) raises the possibility of a wind flow similar to the solar wind. In principle, this would result in angular momentum transport away from the disk, which could have some influence on the inflow. Another feature related to the accretion disk corona is the possibility of a flare energy release similar to solar flares (see Part II, Sect. 8.3).

When a plasma in the disk corona is optically thin and has a dominant magnetic pressure, the circumstances are likely to be similar to the solar corona. Therefore

it is possible to imagine some similarity between the mechanisms of solar flares and X-ray shots in accretion disk coronae.

Besides the effect of heating the disk corona, reconnection is able to accelerate particles to high energies ([Lesch and Pohl 1992](#); [Bednarek and Protheroe 1999](#)). Some geometrical and physical properties of the flares in accretion disk coronae can be inferred almost directly from soft- and hard X-ray observations of Galactic black hole candidates ([Beloborodov 1999](#); [Di Matteo et al. 1999](#)).

13.3 Astrophysical Jets

13.3.1 Magnetic and Gravitomagnetic Forces

Jet-like phenomena, including relativistic jets (Begelman et al. 1984; Birkinshaw 1997), are observed on a wide range of scales in accretion disk systems. Active galactic nuclei (AGN) show extremely energetic outflows extending even to scales beyond the outer edge of a galaxy in the form of strongly collimated radio jets. The luminosities of the radio jets give an appreciable fraction of the luminosity of the underlying central object. There is substantial evidence that the ordinary *magnetic forces are involved in the driving mechanism* and that the magnetic fields also provide the collimation of relativistic flows (see also Sect. 20.1.3). So numerical simulations must incorporate relativistic MHD in a four-dimensional space-time (Nishikawa et al. 1999; Koide et al. 1999).

Rotating black holes are thought to be the prime-mover behind the activity detected in centers of galaxies. The gravitational field of rotating black holes is more complex than that of non-rotating ones. In addition to the ordinary gravitational force, $m\mathbf{g}$, the rotation generates the so-called *gravitomagnetic force* which is just an analogy of the Lorentz force. In fact, the full weak-gravity (far from the hole) low-velocity (replacing the relativistic unified space-time with an equivalent Galilean “absolute-space-plus-universal-time”) coordinate acceleration of uncharged particle (Macdonald et al. 1986; see also Chap. 4 in Novikov and Frolov 1989)

$$\frac{d^2\mathbf{r}}{dt^2} = \mathbf{g} + \frac{d\mathbf{r}}{dt} \times \mathbf{H}_{gr}. \quad (13.31)$$

This looks like the Lorentz force with the electric field $\mathbf{E}$ replaced by $\mathbf{g}$, the magnetic field $\mathbf{B}$ replaced by the vector

$$\mathbf{H}_{gr} = \text{rot } \mathbf{A}_{gr},$$

and the electric charge e replaced by the particle mass m . These analogies lie behind the use of the words “*gravitoelectric*” and “*gravitomagnetic*” to describe the gravitational acceleration field $\mathbf{g}$ and to describe the “*shift function*” $\mathbf{A}_{gr}$ and its derivatives (Exercise 13.6).

13.3.2 The Space Experiment “Gravity Probe B”

It is interesting that the idea that a rotating body may exert not only the static gravitational force of Newtonian type but also an additional “dragging force” on test particles deflecting them in the direction of the rotation was formulated very

many years ago by [Mach \(1872, 1883\)](#). In our days, we understand that a massive body warps space-time, and a rotating massive body drags space-time around with it as predicted by the Einstein theory of general relativity. However

detection of gravitomagnetic components as predicted by Einstein's theory of general relativity is one of the primary goals of experimental gravitational physics

(see, e.g., Chap. 4 in [Ciufolini and Matzner 2010](#)). However the *frame-dragging effect*, the amount by which the rotating massive body drags its local space-time with it, has never been measured directly before the *Gravity Probe B* (GP-B), a NASA space experiment launched in 2004 ([Everitt et al. 2011](#)). This experiment used four cryogenic spherical gyroscopes and a star-tracking telescope on board a satellite orbiting above the Earth. The four were set in line (two spinning clockwise and two counter-clockwise) with axes initially aligned with the sight of the telescope.

In astrophysical plasma, the frame-dragging effect has been involved as a mechanism to drive the relativistic jets from galactic nuclei ([Thorne 1988](#)). Moreover the non-linear frame-dragging effect is of particular interest for astrophysicists studying black hole because, like the air in a hurling tornado in the Earth atmosphere, the hurling space in the vicinity of a massive rotating black hole presumably has enormous destructive potential.

The analogy with electromagnetism remains strong so long as all velocities are small compared with that of light and gravity is weak enough to be linear. Thus, far from the horizon, the gravitational acceleration

$$\mathbf{g} = -\frac{M}{r^2} \mathbf{e}_r \quad (13.32)$$

is the radial Newtonian acceleration, and the gravitomagnetic field

$$\mathbf{H}_{gr} = 2 \frac{\mathbf{J} - 3(\mathbf{J} \cdot \mathbf{e}_r) \mathbf{e}_r}{r^3} \quad (13.33)$$

is a dipole field with the role of dipole moment played by the hole's angular momentum

$$\mathbf{J} = \int (\mathbf{r} \times \rho_m \mathbf{v}) dV. \quad (13.34)$$

The results of the GP-B experiment ([Everitt et al. 2011](#)) enables us to verify that the frame-dragging effect does exist, and that it is indeed proportional to the angular momentum of our rotating Earth. So we are able to extrapolate these frame-dragging measurements around the Earth to much more massive celestial objects such as black holes ([Thorne 1988](#)).




Fig. 13.2 An accretion disk D around a rotating black hole is driven into the hole's equatorial plane at small radii by a combination of gravitomagnetic forces (action of the gravitomagnetic field $\mathbf{H}_{gr}$ on orbiting plasma) and viscous forces

13.3.3 Relativistic Jets Near Black Holes

A physical manifestation of the gravitomagnetic field (13.33) is the precession that is induced in gyroscopes far from the black hole. The electromagnetic analogy suggests that not only should the gravitomagnetic field exert a torque on a gyroscope outside a black hole, it should also exert a force.

The gravitomagnetic force drives an accretion disk into the hole's equatorial plane and holds it there

indefinitely regardless of how the disk's angular momentum may change (see Fig. 13.2).

Consequently, at radii where the bulk of the disk's gravitational energy is released and where the hole-disk interactions are strong,

there is only one geometrically preferred direction along which a jet might emerge: the normal to the disk plane, which coincides with the rotation axis of the black hole.

In some cases the jet might be produced by winds off the disk, in other cases by electrodynamic acceleration of the disk, and in others by currents in the hole's magnetosphere (see Begelman et al. 1984). However whatever the mechanism, the jet presumably is locked to the hole's rotation axis.

The black hole acts as a gyroscope to keep the jet aligned. The fact that it is very difficult to torque a black hole accounts for the constancy of the observed jet directions over length scales as great as millions of light years and thus over time scales of millions of years or longer.

A black hole by itself is powerless to produce the observed jets. It does so only with the aid of surrounding plasma and magnetic fields. A supermassive hole in

a galactic nucleus can acquire surrounding matter either by gravitationally pulling interstellar gas into its vicinity, or by tidally disrupting passing stars and smearing their matter out around itself. In either case the gas is likely to have so much angular momentum that, instead of being swallowed directly and radially into the hole, it forms an orbiting disk around the hole. The orientation of the disk at large radii is determined by the direction of the angular momentum of the recently acquired gas, see an external part of the accretion disk in Fig. 13.2.

In the highly-conducting plasma, the gravitomagnetic force couples with electromagnetic fields over Maxwell's equations.

This effect has interesting consequences for the magnetic fields advected from the interstellar matter towards the black hole (Camenzind 1995). It leads to a *gravitomagnetic dynamo* which amplifies any seed field near a rotating compact object. This process builds up the dipolar magnetic structures which may be behind the bipolar outflows seen as relativistic jets (for comparison with a non-relativistic process see Sect. 14.4).

Magnetic fields also influence the accretion towards the rotating black hole. For rapidly rotating holes, the accretion can carry negative angular momentum inwards, spinning down the black hole.

13.3.4 Relativistic Jets from Disk Coronae

Relativistic jets are produced perpendicular to the accretion disk plane (see Fig. 13.2) around a super-massive black hole in the central part of an AGN. The shock of the jets on intergalactic media, at a distance of several hundreds of kpc from the central engine, is considered as being able to accelerate particles up to the highest energies, say 10^{20} eV for cosmic rays. This hypothesis needs, however, to be completed by some further and necessary ingredients since such powerful galaxies are rare objects.

Subramanian et al. (1999) consider the possibility that the relativistic jets observed in many AGN may be powered by the Fermi acceleration of protons in a tenuous corona above a two-temperature accretion disk (Sect. 8.3.5). The acceleration arises, in this scenario, as a consequence of the shearing motion of the magnetic field lines in the corona, that are anchored in the underlying Keplerian disk. The protons in the corona have a power-law distribution because the density there is too low for proton-proton collisions (formula (8.39)) to thermalize the energy supplied via Fermi acceleration.

The same mechanism also operates in the disk itself. However there the density is high enough for thermalization to occur and consequently the disk protons have the Maxwellian distribution. Particle acceleration in the corona leads to the development of a pressure-driven wind that passes through a critical point and subsequently transforms into a relativistic jet at large distances from the black hole.

13.4 Practice: Exercises and Answers

Exercise 13.1 (Sect. 13.1.1). Evaluate the characteristic value of Alfvén speed in the solar corona above a large sunspot.

Answer. From definition (13.15) we find the following formula for Alfvén speed

$$V_A \approx 2.18 \times 10^{11} \frac{B}{\sqrt{n}} \text{ , cm s}^{-1} \text{ .} \quad (13.35)$$

In this formula, in the coefficient, we have neglected a small contribution of the ions that are heavier than protons into the plasma density ρ . Another thing is much more important however.

Above a sunspot the field strength can be as high as $B \approx 3,000 \text{ G}$. Plasma density in the low corona $n \approx 2 \times 10^8 \text{ cm}^{-3}$. For these values formula (13.35) gives unacceptably high values of the Alfvén speed: $V_A \approx 5 \times 10^{10} \text{ cm s}^{-1} > c$. This means that

in a strong magnetic field and low density plasma, the Alfvén waves propagate with velocities approaching the light speed c .

So the non-relativistic formula (13.35) has to be corrected by a *relativistic* factor which takes this fact into account.

Alfvén (1950) pointed out that the “magnetohydrodynamic waves” are just an extreme case of electromagnetic waves (Sect. 15.2.2 and Exercise 13.3). Alfvén has shown that the transition between electromagnetic and Alfvén waves can be surveyed by the help of the following formula for the speed of propagation along the magnetic field:

$$V_A^{rel} = \frac{B}{\sqrt{4\pi\rho}} \frac{1}{\sqrt{1 + B^2/4\pi\rho c^2}} \text{ ,} \quad (13.36)$$

which agrees with (13.15) when $B^2 \ll 4\pi\rho c^2$. Therefore the relativistic Alfvén wave speed is always smaller than the light speed:

$$V_A^{rel} = \frac{c}{\sqrt{1 + 4\pi\rho c^2/B^2}} \leq c \text{ .} \quad (13.37)$$

For values of the magnetic field and plasma density mentioned above, this formula gives $V_A^{rel} \approx 2 \times 10^{10} \text{ cm s}^{-1} < c$ of course. However formula (13.37) shows that, in low density astrophysical plasmas, the Alfvén speed can easily approach the light speed c .

Exercise 13.2 (13.1.3). Discuss properties of the Lorentz force (13.17) in terms of the Maxwellian stress tensor (12.11).

Answer. In non-relativistic MHD, the Maxwellian stress tensor has only magnetic field components (see formula (12.49)):

$$M_{\alpha\beta} = \frac{1}{4\pi} \left(\frac{B^2}{2} \delta_{\alpha\beta} - B_\alpha B_\beta \right). \quad (13.38)$$

Let us write down these components in the reference system which has the z coordinate in the direction of the magnetic field at a given point. In its neighbourhood, formula (13.38) implies

$$M_{\alpha\beta} = \begin{vmatrix} B^2/8\pi & 0 & 0 \\ 0 & B^2/8\pi & 0 \\ 0 & 0 & -B^2/8\pi \end{vmatrix}. \quad (13.39)$$

According to definition (13.38) the zz component of the tensor has two parts:

$$M_{zz} = \frac{B^2}{8\pi} - \frac{B^2}{4\pi}. \quad (13.40)$$

The first part, $B^2/8\pi$, combines with the M_{xx} and M_{yy} components to give an isotropic magnetic pressure. The remaining part, $-B^2/4\pi$, corresponds to excess “negative pressure” or *tension* in the z direction. Thus

■ a magnetic field has a tension along the field lines in addition to having the isotropic pressure, $B^2/8\pi$.

The second term on the right-hand side of the Maxwellian stress tensor (13.38) describes the *magnetic tension* along field lines.

Recall that the diagonal components of the ordinary pressure tensor (12.15), in exactly the same way, correspond to isotropic gas pressure and the off-diagonal components to viscous shear.

Exercise 13.3 (Sect. 13.1.3). Show that the *magnetic tension force* is directed to the local centre of curvature.

Answer. The Lorentz force is

$$\mathbf{F}_m = -\frac{1}{4\pi} \mathbf{B} \times \text{rot } \mathbf{B} = \frac{1}{4\pi} (\mathbf{B} \cdot \nabla) \mathbf{B} - \nabla \frac{B^2}{8\pi}. \quad (13.41)$$

Here $(\mathbf{B} \cdot \nabla)$ is the directional derivative along a magnetic field line. Hence we can use formulae that are similar to (5.43) and (5.44) to rewrite the first term on the right-hand side of (13.41) as follows

$$\frac{1}{4\pi} (\mathbf{B} \cdot \nabla) \mathbf{B} = -\frac{B^2}{4\pi} \frac{\mathbf{e}_c}{\mathcal{R}_c} + \frac{\partial}{\partial l} \frac{B^2}{8\pi} \mathbf{n}. \quad (13.42)$$

Here $\mathbf{n} = \mathbf{B}/B$ is the unit vector along the magnetic field, l is the distance along the field line, $\mathcal{R}_c$ is a radius of *curvature* for the field line at a given point $\mathbf{R}$. At this point the unit vector $\mathbf{e}_c$ is directed from the curvature center 0_c as shown in Fig. 5.8.

Let us decompose the second term on the right-hand side of (13.41) as

$$-\nabla \frac{B^2}{8\pi} = -\nabla_{\perp} \frac{B^2}{8\pi} - \frac{\partial}{\partial l} \frac{B^2}{8\pi} \mathbf{n}, \quad (13.43)$$

where the operator $\nabla_{\perp}$ operates in the planes normal to the magnetic field lines.

Now we combine formulae (13.42) and (13.43) to write the Lorentz force as

$$\mathbf{F}_m = -\nabla_{\perp} \frac{B^2}{8\pi} - \frac{B^2}{4\pi} \frac{\mathbf{e}_c}{\mathcal{R}_c}. \quad (13.44)$$

The first term in the Lorentz force is the magnetic pressure force which is isotropic in the planes normal to the magnetic field lines. It is directed from high magnetic pressure (strong magnetic field) to low magnetic pressure (low field strength) in the same way as the gas pressure. Therefore

the magnetic pressure force acts when the strength of the magnetic field is not a constant in space.

The second term on the right-hand side of (13.44), the magnetic tension force, is directed to the local center of curvature (see point 0_c is Fig. 5.8). It is inversely proportional to the curvature radius $\mathcal{R}_c$. Thus the more a field line is curved, the stronger the tension force is.

The magnetic tension force behaves in an identical way as the tension force in an elastic string.

It is present for magnetic fields with curved field lines and tends to make curved field lines straight, for example, in an Alfvén wave (see Fig. 15.1).

The sum of both terms, the Lorentz force, has no component along the magnetic field. We already knew this since the vector product $\mathbf{B} \times \text{rot} \mathbf{B}$ is perpendicular to the vector $\mathbf{B}$.

Exercise 13.4 (Sect. 13.1.1). For the conditions in the low corona, used in Exercise 13.1, estimate the parameter γ^2 .

Answer. Substitute $p_0 = 2n_0k_B T_0$ in definition (13.14):

$$\gamma^2 = \frac{n_0 k_B T_0}{B_0^2 / 8\pi} \approx 3.47 \times 10^{-15} \frac{n_0 T_0}{B_0^2}. \quad (13.45)$$

Let us take as the characteristic values of temperature $T_0 \approx 2 \times 10^6$ K and magnetic field $B_0 \approx 3,000$ G. For these values formula (13.45) gives the dimensionless parameter

$$\gamma^2 \sim 10^{-7}.$$

Hence, in the solar corona above sunspots, the conditions (13.25) of a strong magnetic field can be satisfied well for a wide range of plasma parameters.

Exercise 13.5 (Sect. 13.3.2). By using general formula (12.76) for the energy flux in ideal MHD, find the magnetic energy influx into a reconnecting current layer (RCL).

Answer. Let us consider a current layer as a neutral one (Fig. 8.5). In this simplest approximation, near the layer, the magnetic field $\mathbf{B} \perp \mathbf{v}$. Therefore in formula (12.76) the scalar product $\mathbf{B} \cdot \mathbf{v} = 0$ and the energy flux density

$$\mathbf{G} = \rho \mathbf{v} \left(\frac{v^2}{2} + w \right) + \frac{B^2}{4\pi} \mathbf{v}. \quad (13.46)$$

If the approximation of a strong field is satisfied, the last term in (13.46) is dominating, and we find the magnetic energy flux density or the Poynting vector (cf. general definition (12.53)) directed into the current layer

$$\mathbf{G}_p = \frac{B^2}{4\pi} \mathbf{v}. \quad (13.47)$$

For a quarter of the current layer assumed to be symmetrical and for a unit length along the current, the total flux of magnetic energy

$$\mathcal{E}_{mag}^{in} = \frac{B_0^2}{4\pi} v_0 b. \quad (13.48)$$

Here b is half-width of the layer (see Fig. 8.5), B_0 is the field strength on the inflow sides of the reconnecting current layer, v_0 is the inflow velocity.

Exercise 13.6 (Sect. 13.3.1). Consider a weakly gravitating, slowly rotating body such as the Earth or the Sun, with all nonlinear gravitational effects neglected. Compute the *gravitational* force and *gravitomagnetic* force (as in Sect. 13.3.1) from the linearized Einstein equations (see Landau and Lifshitz 1975, Chap. 10, Sect. 100). Show that, for a time-independent body, these equations are identical to the Maxwell equations (1.27)–(1.30):

$$\text{rot } \mathbf{g} = 0, \quad \text{div } \mathbf{g} = -4\pi G \rho_m, \quad (13.49)$$

$$\text{rot } \mathbf{H}_{gr} = -16\pi G \rho_m \mathbf{v}, \quad \text{div } \mathbf{H}_{gr} = 0. \quad (13.50)$$

Here the differences are: (a) two minus signs due to gravity being attractive rather than repulsive, (b) the factor 4 in the $\text{rot } \mathbf{H}_{gr}$ equation, (c) the presence of the gravitational constant G , (d) the replacement of charge density ρ^q by mass density ρ_m , and (e) the replacement of electric current density $\mathbf{j}$ by the density of mass flow $\rho_m \mathbf{v}$ with $\mathbf{v}$ the velocity of the mass.

Exercise 13.7 (Sect. 13.3.1). Schiff (1960) showed that a gyroscope in an orbit around the Earth will undergo two relativistic precessions. The first is a geodetic drift in the orbit plane due to motion through the space-time curved by the Earth mass. The second is the frame-dragging due to Earth's rotation. Discuss how to distinguish these two effects by measuring the displacement angles of the spin axes of an ideal gyroscope placed in a polar-orbiting satellite.

Hint. The two effects are at right angles. The warping of space-time exerts a torque on the gyroscope so its axis precesses in the plane of the satellite's orbit (see Fig. 1 in Will 2011). However the Earth rotation exerts the precession which is perpendicular to the orbital plane.

Chapter 14

Plasma Flows in a Strong Magnetic Field

Abstract A sufficiently strong magnetic field easily moves a comparatively rarefied plasma in many non-stationary phenomena in the astrophysical environment. The best studied example is the solar flares accompanied by coronal mass ejections (CMEs) which strongly influence the interplanetary and terrestrial space.

14.1 The General Formulation of a Problem

As was shown in Sect. 13.1.3, the set of MHD equations for an ideal medium in the approximation of strong field and cold plasma is characterized only by the small parameter $\varepsilon^2 = v^2/V_A^2$:

$$\varepsilon^2 \frac{d\mathbf{v}}{dt} = -\frac{1}{\rho} \mathbf{B} \times \text{rot } \mathbf{B}, \quad (14.1)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \text{rot}(\mathbf{v} \times \mathbf{B}), \quad (14.2)$$

$$\frac{\partial \rho}{\partial t} + \text{div } \rho \mathbf{v} = 0. \quad (14.3)$$

Let us try to find the solution to this set as a power series in the parameter ε^2 , i.e. representing all the unknown quantities in the form


$$f(\mathbf{r}, t) = f^{(0)}(\mathbf{r}, t) + \varepsilon^2 f^{(1)}(\mathbf{r}, t) + \dots. \quad (14.4)$$

Then we try to find the solution in three consequent steps.

(a) To zeroth order with respect to ε^2 , the magnetic field is determined by the equation

$$\mathbf{B}^{(0)} \times \text{rot } \mathbf{B}^{(0)} = 0. \quad (14.5)$$

Fig. 14.1 The boundary and initial conditions for the ideal MHD problems



This must be supplemented with a boundary condition, which generally depends on time:

$$\mathbf{B}^{(0)}(\mathbf{r}, t) \big|_S = \mathbf{f}_1(\mathbf{r}, t). \quad (14.6)$$

Here S is the boundary of the region G (Fig. 14.1), in the interior of which the force-free-field equation (14.5) applies.

- The strong force-free magnetic field, changing in time according to the boundary condition (14.6), sets the plasma in motion.
- (b) The kinematics of this motion is uniquely determined by two conditions. The first one follows from the equation of motion and signifies the orthogonality of acceleration to the magnetic field lines

$$\mathbf{B}^{(0)} \cdot \frac{d\mathbf{v}^{(0)}}{dt} = 0. \quad (14.7)$$

This equation is obtained by taking the scalar product of (14.1) and the vector $\mathbf{B}^{(0)}$.

The second condition is a consequence of the freezing-in equation (14.2)

$$\frac{\partial \mathbf{B}^{(0)}}{\partial t} = \text{rot}(\mathbf{v}^{(0)} \times \mathbf{B}^{(0)}). \quad (14.8)$$

Equations (14.7) and (14.8) determine the velocity field $\mathbf{v}^{(0)}(\mathbf{r}, t)$, if the initial condition inside the region G is given:

$$\mathbf{v}_{\parallel}^{(0)}(\mathbf{r}, 0) \big|_G = \mathbf{f}_2(\mathbf{r}). \quad (14.9)$$

Here $\mathbf{v}_{\parallel}^{(0)}$ is the velocity component along the magnetic-field lines.

The velocity component across the field lines, $\mathbf{v}_{\perp}^{(0)}$, is uniquely defined, once the field $\mathbf{B}^{(0)}(\mathbf{r}, t)$ is known, by the freezing-in equation (14.8) at any moment, including the initial one. Therefore we do not need the initial condition for $\mathbf{v}_{\perp}^{(0)}$.

- (c) Since we know the velocity field $\mathbf{v}^{(0)}(\mathbf{r}, t)$, the continuity equation

$$\frac{\partial \rho^{(0)}}{\partial t} + \text{div} \rho^{(0)} \mathbf{v}^{(0)} = 0 \quad (14.10)$$

allows us to find the plasma density distribution $\rho^{(0)}(\mathbf{r}, t)$, if we know its initial distribution

$$\rho^{(0)}(\mathbf{r}, 0) \big|_G = f_3(\mathbf{r}). \quad (14.11)$$

Therefore (14.5), (14.7) and (14.8), together with the continuity equation (14.10), completely determine the unknown zero-order quantities $\mathbf{B}^{(0)}(\mathbf{r}, t)$, $\mathbf{v}^{(0)}(\mathbf{r}, t)$ and $\rho^{(0)}(\mathbf{r}, t)$, once the boundary condition (14.6) at the boundary S is given, and the initial conditions (14.9) and (14.11) inside the region G are given (Somov and Syrovatskii 1976b).

At any moment of time, the field $\mathbf{B}^{(0)}(\mathbf{r}, t)$ is found from (14.5) and the boundary condition (14.6). Thereupon the velocity $\mathbf{v}^{(0)}(\mathbf{r}, t)$ is determined from (14.7) and (14.8) and the initial condition (14.9). Finally the continuity equation (14.10) and the initial condition (14.11) give the plasma density distribution $\rho^{(0)}(\mathbf{r}, t)$.

From here on we restrict our attention to the consideration of the zeroth order relative to the parameter ε^2 , neglecting the magnetic field deviation from a force-free state. However the consecutive application of the expansion (14.4) to the set of equations (14.1)–(14.3) allows us to obtain *a closed set of equations* for determination of ideal MHD quantities *in any order* relative to the small parameter ε^2 .

An important point, however, is that, during the solution of the problem in the zeroth order relative to ε^2 , regions can appear, where the gas pressure gradient cannot be ignored. Here effects proportional to the small parameter γ^2 must be taken into account (Sect. 13.1.3). This fact usually imposes a limitation on the applicability of the strong-field-cold-plasma approximation.

The question of the existence of general solutions to the ideal MHD equations in this approximation will be considered in Sect. 14.3, using two-dimensional problems as an example.

14.2 The Formalism of Two-Dimensional Problems


While being relatively simple from the mathematical viewpoint, two-dimensional (2D) ideal MHD problems allow us to gain some general knowledge concerning the actual flows of plasma with the frozen-in strong magnetic field. Moreover the 2D problems are sometimes a close approximation of the real three-dimensional (3D) flows and can be used to compare the theory with experiments and observations, both qualitatively and quantitatively.

There are *two types of problems* (Somov 1994a) treating the plane flows of plasma, i.e. the flows with the velocity field of the form

$$\mathbf{v} = \{v_x(x, y, t), v_y(x, y, t), 0\}. \quad (14.12)$$

All the quantities are dependent on the variables x , y and t .

Fig. 14.2 Structure of the magnetic field near a separator. A longitudinal field $\mathbf{B}_{\parallel}$ parallel to the z axis is superimposed on the 2D hyperbolic field in the plane (x, y)



14.2.1 The First Type of Problems

The first type incorporates the problems with a magnetic field which is everywhere parallel to the z axis of a Cartesian system of coordinates:

$$\mathbf{B} = \{0, 0, B(x, y, t)\}. \quad (14.13)$$

Thus the corresponding electric current is parallel to the (x, y) plane:

$$\mathbf{j} = \{j_x(x, y, t), j_y(x, y, t), 0\}. \quad (14.14)$$

As an example of a problem of the first type, let us discuss the effect of a *longitudinal* magnetic field in a reconnecting current layer (RCL). Under real conditions, reconnection does occur not at the zeroth lines, but rather at the “limiting lines” of the magnetic field or “*separators*” (see Part II, Sect. 3.2). The latter differ from the zeroth lines only in that the separators contain the longitudinal component of the field as shown in Fig. 14.2.


With the appearance of the longitudinal field, the force balance in the RCL that is formed at the separator is changed. The field and plasma pressure outside the layer must balance not only the gas pressure but also that of the longitudinal field inside the layer (Fig. 14.3)

$$\mathbf{B}_{\parallel} = \{0, 0, B_{\parallel}(x, y, t)\}. \quad (14.15)$$

This effect is well known, for example, in the so-called theta-pinch. In axially symmetric geometry, in cylindrical coordinates r, θ, z , an azimuthal current density j_{θ} crossed with an axial field B_z can support a radial pressure gradient.

If the longitudinal field accumulated in the layer during reconnection, the field pressure $B_{\parallel}^2/8\pi$ would considerably limit the layer compression as well as the reconnection rate. However the solution of the problem of the first type with respect

Fig. 14.3 A model of a reconnecting current layer with a longitudinal component $\mathbf{B}_{\parallel}$ of a magnetic field



to $\mathbf{B}_{\parallel}$ (see Part II, Sect. 6.2.2) shows that another effect is of importance in the real plasma with finite conductivity.

The effect, in essence, is this: the *longitudinal field compression* in the RCL produces a gradient of this field and a corresponding electric current circulating in the transversal (relative to the main current j_z in the layer) plane (x, y) . This current circulation is of the type (14.14); it is represented schematically in Fig. 14.3.

The circulating current plays just the same role as the j_{θ} -current in the theta-pinch, a one-dimensional equilibrium in a cylindric geometry with an axial field $B_z(r)$. *Ohmic dissipation of the circulating current* under conditions of finite conductivity leads to longitudinal field diffusion outwards from the layer. More exactly, because of dissipation of the circulating current, plasma moves into the RCL relatively free with respect to the longitudinal component of magnetic field, thus limiting its accumulation in the RCL.

14.2.2 The Second Type of MHD Problems

14.2.2(a) Magnetic Field and Its Vector Potential

From this point on we shall be mainly interested in the 2D problems of the second type. They treat the plane plasma flows (14.12) associated with the plane magnetic field

$$\mathbf{B} = \{ B_x(x, y, t), B_y(x, y, t), 0 \}. \quad (14.16)$$


The electric currents corresponding to this field are parallel to the z axis

$$\mathbf{j} = \{ 0, 0, j(x, y, t) \}. \quad (14.17)$$

The vector-potential $\mathbf{A}$ of such a field has an its only non-zero component:

$$\mathbf{A} = \{ 0, 0, A(x, y, t) \}.$$

Fig. 14.4 The curve L connects the points 1 and 2 situated in different field lines



The magnetic field $\mathbf{B}$ is defined by the z -component of the vector-potential:

$$\mathbf{B} = \left\{ \frac{\partial A}{\partial y}, -\frac{\partial A}{\partial x}, 0 \right\}. \quad (14.18)$$

The scalar function $A(x, y, t)$ is often termed the *vector potential*. This function is quite useful, owing to its properties.

Property 14.1. Substitute (14.18) in the differential equations describing the magnetic field lines

$$\frac{dx}{B_x} = \frac{dy}{B_y} = \frac{dz}{B_z}. \quad (14.19)$$

Equations (14.19) imply parallelism of the vector $d\mathbf{l} = \{dx, dy, dz\}$ to the vector $\mathbf{B} = \{B_x, B_y, B_z\}$. In the case under study $B_z = 0$, $dz = 0$, and

$$\frac{dx}{\partial A / \partial y} = -\frac{dy}{\partial A / \partial x}$$

or

$$\frac{\partial A}{\partial x} dx + \frac{\partial A}{\partial y} dy = 0.$$

On integrating the last, we come to the conclusion that the relation

$$A(x, y, t) = \text{const} \quad \text{for} \quad t = \text{const}$$

(14.20)

is the equation for a family of magnetic field lines in the plane $z = \text{const}$ at the moment t .

Property 14.2. Let L be some curve in the plane (x, y) and $d\mathbf{l}$ an arc element along the curve in Fig. 14.4.

Let us calculate the magnetic flux $d\Phi$ through the arc element $d\mathbf{l}$. By definition,

$$\begin{aligned}
 d\Phi &= \mathbf{B} \cdot d\mathbf{S} = \mathbf{B} \cdot (\mathbf{e}_z \times d\mathbf{l}) = \mathbf{B} \cdot \begin{vmatrix} \mathbf{e}_x & \mathbf{e}_y & \mathbf{e}_z \\ 0 & 0 & 1 \\ dx & dy & 0 \end{vmatrix} \\
 &= \mathbf{B} \cdot \{(-dy)\mathbf{e}_x + dx\mathbf{e}_y\} = -B_x dy + B_y dx. \tag{14.21}
 \end{aligned}$$

On substituting definition (14.18) in formula (14.21) we find that

$$d\Phi = -\frac{\partial A}{\partial y} dy - \frac{\partial A}{\partial x} dx = -dA. \tag{14.22}$$

On integrating (14.22) along the curve L from point 1 to point 2 we obtain the magnetic flux

$$\Phi = A_2 - A_1. \tag{14.23}$$

Thus the fixed value of the vector potential A is not only the field line “tag” determined by formula (14.20);

the difference of values of the vector potential A on two field lines is equal to the magnetic flux between them.

From this, in particular, the following simple rule holds: we have to plot the magnetic field lines corresponding to equidistant values of A .

Property 14.3. Let us substitute definition (14.18) in the freezing-in equation (14.2). We obtain the following general equation

$$\text{rot} \frac{d\mathbf{A}}{dt} = 0. \tag{14.24}$$

Disregarding a gradient of an arbitrary function, which can be eliminated by a gauge transformation, and considering the second type of 2D problems, we have

$$\frac{dA}{dt} \equiv \frac{\partial A}{\partial t} + (\mathbf{v} \cdot \nabla)A = 0. \tag{14.25}$$

This equation means that, in the plane (x, y) , the lines

$$A(x, y, t) = \text{const} \tag{14.26}$$

are *Lagrangian* lines, i.e. they move together with the plasma. According to (14.20) they are composed of the field lines, hence (14.25) expresses the magnetic field freezing in plasma.

Thus (formally it follows from (14.25) on passing to the Lagrangian variables) we have one of the integrals of motion

$$A(x, y, t) = A(x_0, y_0, 0) \equiv A_0$$

(14.27)

at an arbitrary t . Here x_0, y_0 are the coordinates of some “*fluid particle*” at the initial moment of time $t = 0$; x, y are the coordinates of the same particle at a moment of time t or (by virtue of (14.27)) the coordinates of any other particle situated on the same field line A_0 at the moment t .

Property 14.4. Equation of motion (14.1) rewritten in terms of the vector potential $A(x, y, t)$ is of the form

$$\varepsilon^2 \frac{d\mathbf{v}}{dt} = -\frac{1}{\rho} \Delta A \nabla A. \quad (14.28)$$

In the zeroth order relative to the small parameter ε^2 , outside the zeroth points (where $\nabla A = 0$) and the magnetic field sources (where $\Delta A \neq 0$) we have:

$$\Delta A = 0.$$

(14.29)

So the vector potential is a *harmonic* function of variables x and y . Hence, while considering the (x, y) plane as a complex plane $z = x + iy$, it is convenient to relate an *analytic* function F to the vector potential A in the region under consideration:

$$F(z, t) = A(x, y, t) + iA^+(x, y, t). \quad (14.30)$$

Here $A^+(x, y, t)$ is a conjugate harmonic function connected with $A(x, y, t)$ by the Cauchy-Riemann conditions (see [Lavrent'ev and Shabat 1973](#), Sect. 2):

$$\frac{\partial A}{\partial x} = \frac{\partial A^+}{\partial y}, \quad \frac{\partial A}{\partial y} = -\frac{\partial A^+}{\partial x}.$$

Thus

$$\begin{aligned} A^+(x, y, t) &= \int \left(-\frac{\partial A}{\partial y} dx + \frac{\partial A}{\partial x} dy \right) + A^+(t) = \\ &= -\int \mathbf{B} \cdot d\mathbf{l} + A^+(t), \end{aligned} \quad (14.31)$$

where $A^+(t)$ is a quantity independent of the coordinates x and y . So the function $A^+(x, y, t)$ can be used as a scalar potential of magnetic field at a moment t , i.e.

$$\mathbf{B} = \left\{ -\frac{\partial A^+}{\partial x}, -\frac{\partial A^+}{\partial y}, 0 \right\} = -\nabla A^+. \quad (14.32)$$

The function $F(z, t)$ is termed the *complex potential*. The magnetic field vector, according to (14.18) and (14.30), is

$$\mathbf{B} = B_x + i B_y = -i \left(\frac{dF}{dz} \right)^*. \quad (14.33)$$

Here the asterisk denotes the complex conjugation. After the introduction of the complex potential, we can widely apply the methods of the complex variable function theory, in particular the method of conform transformations or *conform mapping*, to determine the magnetic field in zeroth order in the small parameter ε^2 (e.g., Exercise 14.4).

This has been done successfully many times in order to determine the structure of the magnetic field: in vicinity of *reconnecting current layer* (RCL; Syrovatskii 1971), in solar coronal streamers (Somov and Syrovatskii 1972b) and the Earth's magnetosphere (Unti and Atkinson 1968; Atkinson and Unti 1969; Obertz 1973), the accretion disk magnetosphere (Somov et al. 2003; see also Part II, Sect. 8.3).

Markovskii and Somov (1989) suggested a generalization of the Syrovatskii model by attaching four shock MHD waves at the endpoints (edges) of the RCL. Such model reduces exactly to the Riemann-Hilbert problem (in an analytical form on the basis of the Christoffel-Schwarz integral) in order to analyze the structure of magnetic field in vicinity of reconnection region and its variations with model parameters (Bezrodnykh et al. 2007, 2011).

14.2.2(b) Motion of the Plasma and Its Density

In the strong field approximation, the plasma motion kinematics due to changes in a potential field is uniquely determined by two conditions:

- (i) The freezing-in condition (14.25) or its solution (14.27) and
- (ii) The acceleration orthogonality with respect to the field lines

$$\frac{d\mathbf{v}^{(0)}}{dt} \times \nabla A^{(0)} = 0 \quad (14.34)$$

(cf. (14.7)). A point to be noted is that (14.34) is a result of eliminating the unknown $\Delta A^{(1)}$, which has a first order in ε^2 , from two components of the vector equation

$$\frac{d\mathbf{v}^{(0)}}{dt} = -\frac{1}{\rho^{(0)}} \Delta A^{(1)} \nabla A^{(0)}. \quad (14.35)$$

If $x(t)$ and $y(t)$ are the coordinates of a fluid particle, (14.34) and (14.25) are reduced to the set of *ordinary differential equations* (see [Somov and Syrovatskii 1976b](#), Chap. III, Sect. 1).

Once the kinematic part of the problem is solved, the trajectories of fluid particles are known:

$$x = x(x_0, y_0, t), \quad y = y(x_0, y_0, t). \quad (14.36)$$

In this case the continuity equation (14.3) solution presents no problem. In fact, the fluid particle density change on moving along the found trajectory is determined by the continuity equation (14.3), rewritten in the Lagrangian form, and is equal to

$$\frac{\rho(x, y, t)}{\rho_0(x_0, y_0)} = \frac{dU_0}{dU} = \frac{\mathcal{D}(x_0, y_0)}{\mathcal{D}(x, y)}. \quad (14.37)$$

Here dU_0 is the initial volume of a particle, dU is the volume of the same particle at a moment of time t ;

$$\frac{\mathcal{D}(x_0, y_0)}{\mathcal{D}(x, y)} = \frac{\partial x_0}{\partial x} \frac{\partial y_0}{\partial y} - \frac{\partial x_0}{\partial y} \frac{\partial y_0}{\partial x} \quad (14.38)$$

is the Jacobian of the transformation that is inverse to the transformation (14.36) of coordinates at a fixed value of time t .

The 2D equations of the MHD strong-field-cold-plasma approximation ([Somov and Syrovatskii 1976b](#)) in the problem of the second type are relatively simple but rather useful for applications to astrophysical plasmas. In particular, they enable us to study the fast plasma flows in the solar atmosphere ([Syrovatskii and Somov 1980](#)) and to understand some aspects of the reconnection process.

In spite of their numerous applications, the list of exact solutions to them is rather poor. Still, we can enrich it significantly,

relying on many astrophysical objects, for example in the accretion disk coronae (see Part II, Sect. 8.3), and some mathematical ideas.

[Titov and Priest \(1993\)](#) have shown that the equations of zeroth order can be reduced to a set of Cauchy-Riemann and ordinary differential equations, by using a conformal system of coordinates in which the positions of particles are fixed by magnitudes of two conjugate functions. These are the flux function and the potential of magnetic field. The set obtained has a special class of solutions. First, in such

flows the conjugate potential is frozen into the moving medium as well as the vector potential $A(x, y, t)$. Second, each flow is realized as a continuous sequence of conformal mappings. A linear diffusion-like equation describes such flows. The equation was solved analytically for examples describing the *magnetic collapse* (cf. Part II, Chap. 2) in the neighbourhood of the X-point.

14.3 On the Existence of Continuous Flows

Thus, in the strong-field-cold-plasma approximation, the MHD equations for a plane two-dimensional flow of ideally conducting plasma (for the second-type problems) are reduced, in the zeroth order in the small parameter ε^2 , to the following closed set of equations:

$$\Delta A = 0, \quad (14.39)$$

$$\frac{d\mathbf{v}}{dt} \times \nabla A = 0, \quad (14.40)$$

$$\frac{dA}{dt} = 0, \quad (14.41)$$

$$\frac{\partial \rho}{\partial t} + \operatorname{div} \rho \mathbf{v} = 0. \quad (14.42)$$

Seemingly, the solution of this set is completely defined inside some region G (Fig. 14.5) on the plane (x, y) , once the boundary condition is given at the boundary S

$$A(x, y, t) \big|_S = f_1(x, y, t) \quad (14.43)$$

together with the initial conditions inside the region G

$$\mathbf{v}_{\parallel}(x, y, 0) \big|_G = \mathbf{f}_2(x, y), \quad (14.44)$$

$$\rho(x, y, 0) \big|_G = f_3(x, y). \quad (14.45)$$




Fig. 14.5 The boundary and initial conditions for the second-type MHD problems

Here $\mathbf{v}_{\parallel}$ is the velocity component along field lines. Once the potential $A(x, y, t)$ is known, the transversal velocity component is uniquely determined by the freezing-in equation (14.41) and is equal, at any moment including the initial one, to

$$\mathbf{v}_{\perp}(x, y, t) = (\mathbf{v} \cdot \nabla A) \frac{\nabla A}{|\nabla A|^2} = -\frac{\partial A}{\partial t} \frac{\nabla A}{|\nabla A|^2}. \quad (14.46)$$

From (14.39) and boundary condition (14.43) we find the vector potential $A(x, y, t)$ at any moment of time. Next, from (14.40) and (14.41) and the initial condition (14.44), the velocity $\mathbf{v}(x, y, t)$ is determined; the density $\rho(x, y, t)$ is found from the continuity equation (14.42) and the initial density distribution (14.45). The next section is devoted to the consideration of an example which may have interesting applications.

14.4 Flows in a Time-Dependent Dipole Field

14.4.1 Plane Magnetic Dipole Fields

Two straight parallel currents, equal in magnitude but opposite in direction, engender the magnetic field which far enough from the currents can be described by a complex potential

$$F(z) = \frac{i\mathbf{m}}{z}, \quad \mathbf{m} = m e^{i\psi} \quad (14.47)$$

and is called the plane *dipole* field. The quantity $m = 2Il/c$ has the meaning of the *dipole moment*, I is the current magnitude, l is the distance between the currents. Formula (14.47) corresponds to the plane dipole situated at the origin of coordinates in the plane (x, y) and directed at an angle of ψ to the x axis. The currents are parallel to the z axis of the Cartesian system of coordinates.

Let us consider the plasma flow caused by the change with time of the strong magnetic field of the plane dipole. Let $\psi = \pi/2$ and $m = m(t)$, $m(0) = m_0$.

- (a) Let us find the first integral of motion. According to (14.30) and (14.47), the complex potential


$$F(z, t) = \frac{im(t)e^{i\pi/2}}{x + iy} = \frac{-m(t)x + im(t)y}{x^2 + y^2}. \quad (14.48)$$

So, according to (14.20), the field lines constitute a family of circles

$$A(x, y, t) = -\frac{m(t)x}{x^2 + y^2} = \text{const} \quad \text{for} \quad t = \text{const}. \quad (14.49)$$

They have centres on the axis x and the common point $x = 0$, $y = 0$ in Fig. 14.6.

Fig. 14.6 The field lines of a plane magnetic dipole



Therefore the freezing-in condition (14.27) results in a first integral of motion

$$\frac{m x}{x^2 + y^2} = \frac{m_0 x_0}{x_0^2 + y_0^2}. \quad (14.50)$$

Here x_0, y_0 are the coordinates of some fluid particle at the initial moment of time $t = 0$; Lagrangian variables x and y are the coordinates of the same particle at a moment t .

- (b) The second integral is easily found in the limit of small changes of the dipole moment $m(t)$ and respectively *small* plasma displacements. Assuming the parameter $\delta = \nu\tau/L$ to be small, (13.27), which is *linear in velocity*, takes the place of (14.34). The integration over time (with zero initial values for the velocity) allows us to reduce (13.27) to the form

$$\frac{\partial x}{\partial t} = K(x, y, t) \frac{\partial A}{\partial x}, \quad \frac{\partial y}{\partial t} = K(x, y, t) \frac{\partial A}{\partial y}. \quad (14.51)$$

Here $K(x, y, t)$ is some function of coordinates and time. Eliminating it from two equations (14.51), we arrive at

$$\frac{\partial y}{\partial x} = \frac{\partial A}{\partial y} \bigg/ \frac{\partial A}{\partial x}. \quad (14.52)$$

Thus, in the approximation of small displacements, not only the acceleration but also the plasma displacements are normal to the field lines.


On substituting (14.49) in (14.52), we obtain an ordinary differential equation

$$\frac{dy}{dx} = \frac{2xy}{x^2 - y^2}.$$

Its integral

$$\frac{y}{x^2 + y^2} = \text{const}$$

Fig. 14.7 A trajectory of a fluid particle driven by a changing magnetic field of a plane dipole



describes a family of circles, orthogonal to the field lines, and presents fluid particle trajectories. In particular, the trajectory of a particle, situated at a point (x_0, y_0) at the initial moment of time $t = 0$, is an arc of the circle

$$\frac{y}{x^2 + y^2} = \frac{y_0}{x_0^2 + y_0^2} \quad (14.53)$$

from the point (x_0, y_0) to the point (x, y) on the field line (14.50) as shown in Fig. 14.7.

Thus the integrals of motion (14.50) and (14.53) completely determine the plasma flow in terms of the Lagrangian coordinates

$$x = x(x_0, y_0, t), \quad y = y(x_0, y_0, t). \quad (14.54)$$

This flow has a simple form: the particles are connected with the magnetic field lines and move together with them in a transversal direction. Such simple kinematics is a result of considering small plasma displacements (from the state having zero initial velocity) under the action of the force perpendicular to the field lines.

- (c) The plasma density change is defined by (14.37). On calculating the Jacobian for the transformation implicitly given by formulae (14.50) and (14.53), we obtain (for the case of a homogeneous initial density distribution ρ_0) the formula

$$\frac{\rho(x, y, t)}{\rho_0} = \left(\frac{m}{m_0}\right) \frac{m_0^4}{(m^2 x^2 + m_0^2 y^2)^4} \left\{ \left[m^2 x^4 + m_0^2 y^4 + x^2 y^2 (3m^2 - m_0^2) \right]^2 - \left[2x^2 y^2 (m_0^2 - m^2) \right]^2 \right\}. \quad (14.55)$$

In particular, on the dipole axis ($x = 0$)

$$\boxed{\frac{\rho(0, y, t)}{\rho_0} = \frac{m}{m_0}}, \quad (14.56)$$

whereas in the “equatorial plane” ($y = 0$)

$$\frac{\rho(x, 0, t)}{\rho_0} = \left(\frac{m_0}{m}\right)^3. \quad (14.57)$$

With increasing dipole moment m , the plasma density on the dipole axis grows proportionally to the moment,

whereas that at the equatorial plane falls in inverse proportion to the third power of the moment. The opposite process takes place as the moment decreases.

The result pertains to the case of small changes in the dipole moment and can demonstrate just the tendency of plasma behavior in the strong magnetic field of a plane dipole. The exception is formula (14.56). It applies to any changes of the dipole moment. The reason is in the following. In the approximation of a strong field and cold plasma, the acceleration of plasma is perpendicular to the field lines and is zero at the dipole axis. Hence, if the plasma is motionless at the initial moment, arbitrary changes of the dipole moment do not cause a plasma motion on the dipole axis ($\mathbf{v} = 0$). Plasma displacements in the vicinity of the dipole axis always remain small ($\delta \ll 1$) and the solution obtained applies.

(d) In the general case of arbitrarily large dipole moment changes,

the *inertial effects* resulting in plasma flows along the magnetic field lines are of considerable importance

(Somov and Syrovatskii 1972a). In this case, the solution of the problem requires the integration of the *ordinary differential equations* that follows from (14.34) or (14.35) together with the freezing-in equation (14.25).

One can obtain exact analytical solutions for a linearly changing magnetic moment using the “*frozen-in coordinates*” technique (Gorbachev and Kel’ner 1988). These coordinates can be quite useful while solving non-stationary MHD problems. One introduces a set which is *doubly Lagrangian*: in the parameter s_1 along a stream line (along the velocity field $\mathbf{v}$) and in the parameter s_2 along a magnetic field line.

14.4.2 Axial-Symmetric Dipole Fields in Plasma

Two-dimensional axial-symmetric problems can better suit the astrophysical applications of the second-type problem considered. The ideal MHD equations are written, using the approximation of a strong field and cold plasma, in spherical

coordinates with due regard for axial symmetry. The role of the vector potential is fulfilled by the so-called *stream function*

$$\Phi(r, \theta, t) = r \sin \theta A_\varphi(r, \theta, t). \quad (14.58)$$

Here A_φ is the only non-zero φ -component of the vector-potential $\mathbf{A}$.

In terms of the stream functions, the equations take the form

$$\frac{d\mathbf{v}}{dt} = \varepsilon^{-2} K(r, \theta, t) \nabla \Phi, \quad \frac{d\Phi}{dt} = 0, \quad \frac{d\rho}{dt} = -\rho \operatorname{div} \mathbf{v}, \quad (14.59)$$

where

$$K(r, \theta, t) = \frac{j_\varphi(r, \theta, t)}{\rho r \sin \theta} \quad (14.60)$$

(Somov and Syrovatskii 1976b). The equations formally coincide with the corresponding equations (14.28), (14.25) and (14.3) describing the plane flows in terms of the vector potential.

As a zeroth approximation in the small parameter ε^2 , we may take, for example, the dipole field. In this case the stream function is of the form

$$\Phi^{(0)}(r, \theta, t) = m(t) \frac{\sin^2 \theta}{r}, \quad (14.61)$$

where $m(t)$ is a time-varying moment.

Let us imagine a homogeneous magnetized ball of radius $R(t)$ with the frozen field $\mathbf{B}_{int}(t)$. The dipole moment of such a ball (a star or its envelope) is

$$m(t) = \frac{1}{2} B_{int}(t) R^3(t) = \frac{1}{2\pi} (B_0 \pi R_0^2) R(t), \quad (14.62)$$

where B_0 and R_0 are the values of $B_{int}(t)$ and $R(t)$ at the initial moment of time $t = 0$. The second equality takes account of the magnetic field freezing-in as conservation of the flux $B_{int}(t) \pi R^2(t)$ through the ball. Formula (14.62) shows that the dipole moment of the ball is thereby proportional to its radius $R(t)$.

The solution to the problem (Somov and Syrovatskii 1972a) shows that as the dipole moment grows (when the ball expands)

the magnetic field rakes the plasma up to the dipole axis, compresses it and simultaneously accelerates it along the field lines.

A distinguishing characteristic of the solution is that the density at the axis grows in proportion to the dipole moment, just as in the 2D plane case (see formula (14.56)).

Envelopes of *nova* and *supernova* stars (Rose 1998; Padmanabhan 2006) present a wide variety of different shapes. We can hardly find the ideally round envelopes, even among the ones of regular shape. It is more common to find either flattened or stretched envelopes. As a rule, their surface brightness is maximal at the ends of the main axes of an oval image. This phenomenon can sometimes be interpreted as a gaseous ring observed almost from an edge. However, if there is no luminous belt

between the brightness maxima, which would be characteristic of the ring, then the remaining possibility is that single gaseous compressions—condensations—exist in the envelope.

At the early stages of the expansion during the explosion of a nova, the condensations reach such brightness that they give the impression that the nova “*bifurcates*”. Consider one of the models in which a magnetic field plays a decisive role. Suppose that the star’s magnetic field was a dipole one before the explosion. At the moment of the explosion a massive envelope with the frozen-in field separated from the star and began to expand. According to (14.62), the expansion results in the growth of the dipole moment. According to the solution of the problem considered above, the field will rake the interstellar plasma surrounding the envelope, as well as external layers of the envelope, up in the direction of the dipole axis.

The process of polar condensate formation can be conventionally divided into *two stages* (Somov and Syrovatskii 1976b, Chap. 2). At the first one, the plasma is raked up by the magnetic field into the polar regions, a corresponding growth in density and pressure at the dipole axis taking place. At the second stage, the increased pressure hinders the growth of the density at the axis, thus stopping compression, but the plasma raking-up still continues. At the same time, the gas pressure gradient, arising ahead of the envelope, gives rise to the motion along the axis. As a result, by the time the magnetic force action stops, all the plasma is raked up into two compact condensates.

The plasma raking-up by the strong magnetic field seems to be capable of explaining some types of chromospheric ejections on the Sun (Somov and Syrovatskii 1976b, Chap. 2, Sect. 4).

If a magnetized ball compresses, plasma flows from the poles to the equatorial plane, thus forming a *dense disk* or *ring*. This case is the old problem of astrophysics concerning the compression of a gravitating cloud with the frozen-in field. The process of magnetic raking-up of plasma into dense disks can effectively work in the atmospheres of collapsing stars.

14.5 Practice: Exercises and Answers

Exercise 14.1. For a three-dimensional (3D) magnetic field $\mathbf{B}$, consider general properties of the vector-potential $\mathbf{A}$ which is determined in terms of two scalar functions α and β :

$$\mathbf{A} = \alpha \nabla \beta + \nabla \psi. \quad (14.63)$$

Here ψ is an arbitrary scalar function.

Answer. Formula (14.63) permits $\mathbf{B}$ to be written as

$$\mathbf{B} = \text{rot } \mathbf{A} = \nabla \alpha \times \nabla \beta, \quad (14.64)$$

where the last step follows from the fact that the rot of a gradient vanishes.

This representation of $\mathbf{B}$ provides another way to obtain information about the magnetic field in 3D problems of plasma astrophysics. According to (14.64)

$$\mathbf{B} \cdot \nabla \alpha = 0 \quad \text{and} \quad \mathbf{B} \cdot \nabla \beta = 0. \quad (14.65)$$

Thus $\nabla \alpha$ and $\nabla \beta$ are perpendicular to the vector $\mathbf{B}$, and functions α and β are constant along $\mathbf{B}$. The surfaces $\alpha = \text{const}$ and $\beta = \text{const}$ are orthogonal to their gradients and tangent to $\mathbf{B}$. Hence

| a magnetic field line can be conveniently defined in terms of a pair of values: α and β .

A particular set of α and β labels a field line.

The functions α and β are referred to as *Euler potentials* or *Clebsch variables*. Depending on a problem to be examined, one form may have an advantage over another. The variables α and β , while in general not easily obtained, are available for some axial-symmetric geometries.

Another advantage of these variables appears in the study of field line motions in the context of the *ideal* MHD theory (see Sect. 5.7 in Parks 2004). Since the time evolution of the magnetic field is governed by the induction equation (14.2), the functions $\alpha(\mathbf{r}, t)$ and $\beta(\mathbf{r}, t)$ satisfy the equations:

$$\frac{\partial \alpha}{\partial t} + (\mathbf{v} \cdot \nabla) \alpha = 0 \quad \text{and} \quad \frac{\partial \beta}{\partial t} + (\mathbf{v} \cdot \nabla) \beta = 0. \quad (14.66)$$

That is, the functions $\alpha(\mathbf{r}, t)$ and $\beta(\mathbf{r}, t)$ take constant values for a point that moves with the plasma.

Exercise 14.2 (Sect. 14.4.2). Evaluate the typical value of the magnetic dipole moment for a *neutron star*.

Answer. Typical neutron stars have $B \sim 10^{12}$ G. With the star radius $R \sim 10$ km, it follows from formula (14.62) that $m \sim 10^{30}$ G cm³. Some of neutron stars, related to the so-called “Soft Gamma-ray Repeaters” (SGRs), are the spinning super-magnetized neutron stars created by *supernova* explosions. The rotation of such stars called *magnetars* is slowing down so rapidly that a super-strong field of the unprecedented strength, $B \sim 10^{15}$ G, could provide so fast braking (see Sect. 19.1.3). For a magnetar the dipole moment $m \sim 10^{33}$ G cm³.

Note that the same value of magnetic dipole moment is typical for white dwarfs. Even with a relatively weak magnetic field $B \sim 10^6$ G (see, e.g., Sect. 13.2.2) they have the star radius $R \approx 0.013 R_{\odot} \sim 10^9$ cm which is small in comparison with the solar radius $R_{\odot}$ but by three orders of magnitude larger than the radius of a neutron star.

Exercise 14.3. Show that, prior to the onset of a solar flare, the magnetic energy density in the corona is of about three orders of magnitude greater than any of the other types. So the flares occur in a plasma environment well dominated by the magnetic field.

Hint. Take the coronal field of about 100 G, and the coronal plasma velocity of order of 1 km s^{-1} .

Exercise 14.4 (Sect. 14.2.2(a)). By using the method of *conform mapping*, determine the shape of a magnetic cavity, *magnetosphere*, created by a plane dipole inside a perfectly conducting uniform plasma with a gas pressure p_0 . Determine the magnetic field inside the cavity.

Answer. The conditions to be satisfied along the boundary S of the magnetic cavity G are equality of magnetic and gas pressure,

$$\left. \frac{B^2}{8\pi} \right|_S = p_0 = \text{const}, \quad (14.67)$$

and tangency of the magnetic field,

$$\mathbf{B} \cdot \mathbf{n} \Big|_S = 0. \quad (14.68)$$

Condition (14.68) means that

$$\text{Re } F(z) = A(x, y) = \text{const}, \quad (14.69)$$

where a complex potential $F(z)$ is an analytic function (14.30) within the region G in the complex plane z except at the point $z = 0$ of the dipole $\mathbf{m}$.

Let us assume that a conform transformation $w = w(z)$ maps the region G onto the circle $|w| \leq 1$ in an auxiliary complex plane $w = u + iv$ so that the point $z = 0$ goes into the center of the circle without rotation of the dipole as shown in Fig. 14.8.

The boundary $|w| = 1$ is the field line S' of the solution in the plane w , which we easily construct:

$$F(w) = \left(w - \frac{1}{w} \right). \quad (14.70)$$

Note that we have used only the boundary condition (14.68).

The other boundary condition (14.67) will allow us to find an unknown conform transformation $w = w(z)$. With account of definition (14.33) taken, condition (14.67) gives us the following relation

$$B^2 = \left| \frac{dF}{dw} \right|^2 \left| \frac{dz}{dw} \right|^{-2} = 8\pi p_0$$




Fig. 14.8 The field lines of a plane dipole $\mathbf{m}$ inside: (a) the unit circle in the plane w , (b) the cavity in a plasma of constant pressure

or

$$\left| \frac{dz}{dw} \right|^2 = \frac{1}{8\pi p_0} \left| \frac{dF}{dw} \right|^2. \quad (14.71)$$

At the boundary $|w| = 1$, this condition reduces to an ordinary differential equation relative to the real part, $x = x(u)$, of an unknown function $z = z(w)$:

$$\left(\frac{dx}{du} \right)^2 = M^2 u^4, \quad \text{where} \quad M^2 = \frac{1}{2\pi}. \quad (14.72)$$

By integrating this equation we find

$$x = \pm M \frac{u^3}{3} + c_1 = \pm \frac{M}{3} \cos^3 \varphi + c_1, \quad (14.73)$$

here φ is an argument of the complex number w , and c_1 is a constant of integration.

Since we know the real part $x = x(\varphi)$ on the circle boundary, we find the complex function $z = z(w)$ in the entire region $|w| \leq 1$, for example, by expanding the function $x = x(\varphi)$ in the Fourier series

$$x(\varphi) = c_1 + \frac{M}{4} \cos \varphi + \frac{M}{12} \cos 3\varphi. \quad (14.74)$$

So, inside the circle, the power series has only three terms:

$$x(r, \varphi) = c_1 + \frac{M}{4} r \cos \varphi + \frac{M}{12} r^3 \cos 3\varphi, \quad (14.75)$$

$$y(r, \varphi) = c_2 + \frac{M}{4} r \sin \varphi + \frac{M}{12} r^3 \sin 3\varphi. \quad (14.76)$$




Fig. 14.9 A model of the star magnetosphere with an accretion disk; Γ_l and Γ_r are the cross sections of the disk. The plane z corresponds to the complex variable $z = x + iy$. S_u and S_d together with Γ_l and Γ_r constitute the boundary of the singly connected domain G in the plane z

Moreover $c_1 = c_2 = 0$ because $z(0) = 0$. Therefore

$$z(w) = \frac{M}{4} \left(w + \frac{w^3}{3} \right). \quad (14.77)$$

The conform mapping (14.77) and the potential (14.70) determine the general solution of the problem, the complex potential (Oreshina and Somov 1999):

$$F(z) = B_0 L^{2/3} \frac{K^4 - 3L^{2/3}K^2 + L^{4/3}}{K(K^2 - L^{2/3})}. \quad (14.78)$$

Here $B_0 = p_0^{1/2}$ is the unit of magnetic field strength, the function

$$K(z) = \left(6\sqrt{2\pi} \cdot z + \sqrt{L^2 + 72\pi \cdot z^2} \right)^{1/3}, \quad (14.79)$$

and $L = m^{1/3} p_0^{-1/6}$ is the unit of length; it shows that, when the dipole moment m increases, the size of the magnetic cavity also increases. This is consistent with what we discussed in Sect. 14.4.

The field lines corresponding solution (14.78) are shown in Fig. 14.8b. Therefore, in addition to the shape of the boundary (Cole and Huth 1959), we have found an analytic solution for the magnetic field inside the static dipole cavity. This solution

can be used in the zero-order approximation, described in Sect. 14.1, to analyze properties of plasma flows near collapsing or exploding astrophysical objects with strong magnetic fields.

Exercise 14.5 (Sect. 14.2.2(a)). To estimate characteristic values of the large-scale magnetic field in the corona of an accretion disk (see Part II, Sect. 8.3.1), we have to find the structure of the field inside an open magnetosphere created by a dipole field of a star and a regular field generated by the disk (Somov et al. 2003).

Consider a simplified 2D problem, demonstrated by Fig. 14.9, on the shape of a magnetic cavity and the shape of the accretion disk under assumption that this cavity, i.e. the magnetosphere, is surrounded by a perfectly conducting uniform plasma with a gas pressure p_0 . Discuss a way to solve the problem by using the method of conform mapping (Part II, Sect. 8.3.2).

Chapter 15

MHD Waves in Astrophysical Plasma

Abstract There are four different modes of magnetohydrodynamic (MHD) waves in an ideal plasma with magnetic field. They can create turbulence, nonlinearly cascade in a wide range of wave-numbers, accelerate particles and produce a lot of interesting effects under astrophysical conditions. New interesting results have been obtained in the classical theory of the small-amplitude MHD waves propagating in a compressible optically-thin medium whose radiative cooling depends on the temperature in the form of one or more sharp maxima. In astrophysical plasmas, these maxima are produced by the radiation of a small admixture of heavy ions.

15.1 The Dispersion Equation in Ideal MHD

Small disturbances in a conducting medium with a magnetic field propagate as waves, their properties being different from those of the usual sound waves in a gas or electromagnetic waves in a vacuum. First, the conducting medium with a magnetic field has a characteristic anisotropy: the wave propagation velocity depends upon the direction of propagation relative to the magnetic field. Second, as a result of the interplay of electromagnetic and hydrodynamic phenomena, the waves in MHD are generally neither longitudinal nor transversal.

The study of the behavior of small-amplitude waves, apart from being interesting in itself, has a direct bearing on the analysis of large-amplitude waves, in particular shock waves and other discontinuous flows in MHD, including reconnecting current layers.

Initially we shall study the possible types of small-amplitude waves, restricting ourselves to the *ideal* MHD equations (12.69). Let us suppose a plasma in the initial stationary state is subjected to a small perturbation, so that velocity $\mathbf{v}_0$, magnetic field $\mathbf{B}_0$, density ρ_0 , pressure p_0 and entropy s_0 acquire some small deviations $\mathbf{v}'$, $\mathbf{B}'$, ρ' , p' and s' :

$$\begin{aligned} \mathbf{v} &= \mathbf{v}_0 + \mathbf{v}', & \mathbf{B} &= \mathbf{B}_0 + \mathbf{B}', \\ \rho &= \rho_0 + \rho', & p &= p_0 + p', & s &= s_0 + s'. \end{aligned} \quad (15.1)$$

The initial state is assumed to be a uniform flow of an homogeneous medium in a constant magnetic field:

$$\begin{aligned} \mathbf{v}_0 &= \text{const}, & \mathbf{B}_0 &= \text{const}, \\ \rho_0 &= \text{const}, & p_0 &= \text{const}, & s_0 &= \text{const}. \end{aligned} \quad (15.2)$$

Needless to say, the latter simplification can be ignored, i.e. we may study waves in inhomogeneous media, the coefficients in linearized equations being dependent upon the coordinates. For the sake of simplicity we restrict our consideration to the case (15.2).

It is convenient to introduce the following designations:

$$\mathbf{u} = \frac{\mathbf{B}_0}{\sqrt{4\pi\rho_0}}, \quad \mathbf{u}' = \frac{\mathbf{B}'}{\sqrt{4\pi\rho_0}}. \quad (15.3)$$

Let us linearize the set of MHD equations for an ideal medium. We substitute definitions (15.1)–(15.3) in the set of equations (12.69), neglecting the products of small quantities. Hereafter the subscript “0” for undisturbed quantities will be omitted. We shall get the following set of *linear differential* equations for the primed quantities characterizing small perturbations:

$$\begin{aligned} \partial \mathbf{u}' / \partial t + (\mathbf{v} \cdot \nabla) \mathbf{u}' &= (\mathbf{u} \cdot \nabla) \mathbf{v}' - \mathbf{u} \operatorname{div} \mathbf{v}', & \operatorname{div} \mathbf{u}' &= 0, \\ \partial \mathbf{v}' / \partial t + (\mathbf{v} \cdot \nabla) \mathbf{v}' &= -\rho^{-1} \nabla (p' + \rho \mathbf{u} \cdot \mathbf{u}') + (\mathbf{u} \cdot \nabla) \mathbf{u}', \\ \partial \rho' / \partial t + (\mathbf{v} \cdot \nabla) \rho' &= -\rho \operatorname{div} \mathbf{v}', \\ \partial s' / \partial t + (\mathbf{v} \cdot \nabla) s' &= 0, & p' &= (\partial p / \partial \rho)_s \rho' + (\partial p / \partial s)_\rho s'. \end{aligned} \quad (15.4)$$

The latter equation is the linearized equation of state. We rewrite it as follows:

$$p' = V_s^2 \rho' + b s'. \quad (15.5)$$

Here

$$V_s = (\partial p / \partial \rho)_s^{1/2} \quad (15.6)$$

is the velocity of *sound* in a medium without a magnetic field (Exercise 15.1), the coefficient $b = (\partial p / \partial s)_\rho$.

By virtue of (15.2), the set of equations (15.4) is that of linear differential equations with *constant coefficients*. That is why we may seek a solution in the form of a superposition of plane waves with a dependence on coordinates and time of the type

$$f'(\mathbf{r}, t) \sim \exp[i(\mathbf{k} \cdot \mathbf{r} - \omega t)], \quad (15.7)$$

where ω is the wave frequency and $\mathbf{k}$ is the wave vector. An arbitrary disturbance can be expanded into such waves by means of a Fourier transform. As this takes place, the set of equations (15.4) is reduced to the following set of *linear algebraic* equations:

$$\begin{aligned} (\omega - \mathbf{k} \cdot \mathbf{v}) \mathbf{u}' + (\mathbf{k} \cdot \mathbf{u}) \mathbf{v}' - \mathbf{u} (\mathbf{k} \cdot \mathbf{v}') &= 0, & \mathbf{k} \cdot \mathbf{u}' &= 0, \\ (\omega - \mathbf{k} \cdot \mathbf{v}) \mathbf{v}' + (\mathbf{k} \cdot \mathbf{u}) \mathbf{u}' - \rho^{-1} (p' + \rho \mathbf{u} \cdot \mathbf{u}') \mathbf{k} &= 0, \\ (\omega - \mathbf{k} \cdot \mathbf{v}) \rho' - \rho (\mathbf{k} \cdot \mathbf{v}') &= 0, \\ (\omega - \mathbf{k} \cdot \mathbf{v}) s' &= 0, & p' - V_s^2 \rho' - b s' &= 0. \end{aligned} \quad (15.8)$$

The quantities $\mathbf{k}$ and ω appearing in this set are assumed to be known from the initial conditions. The unknown terms are the primed ones. With respect to these the set of equations (15.8) is closed, linear and *homogeneous* (the right-hand sides equal zero). For this set to have non-trivial solutions, its determinant must be equal to zero.

The determinant can be conveniently calculated in a frame of reference with one of the axes directed along the wave vector $\mathbf{k}$. In addition, it is also convenient to use the frequency

$$\omega_0 = \omega - \mathbf{k} \cdot \mathbf{v}, \quad (15.9)$$

i.e. the frequency in a frame of reference moving with the plasma.

Setting the determinant equal to zero, we get the following equation

$$\begin{aligned} \omega_0^2 [\omega_0^2 - (\mathbf{k} \cdot \mathbf{u})^2] \times \\ \times [\omega_0^4 - k^2 (V_s^2 + u^2) \omega_0^2 + k^2 V_s^2 (\mathbf{k} \cdot \mathbf{u})^2] &= 0. \end{aligned} \quad (15.10)$$

This equation is called the *dispersion* equation. It defines four values of ω_0^2 . Since they differ in absolute magnitude, *four different modes* of waves are defined, each of them having its own velocity of propagation with respect to the plasma

$$\mathbf{V}_{\text{ph}} = \frac{\omega_0}{\mathbf{k}}. \quad (15.11)$$

Clearly this is the *phase* velocity of a wave. It should be distinguished from the *group* velocity

$$\mathbf{V}_{\text{gr}} = \frac{d \omega_0}{d \mathbf{k}}. \quad (15.12)$$

Let us consider the properties of the waves defined by the dispersion equation (15.10) in greater detail.

15.2 Small-Amplitude Waves in Ideal MHD

15.2.1 Entropy Waves

The first root of the dispersion equation (15.10)

$$\omega_0 = \omega - \mathbf{k} \cdot \mathbf{v} = 0 \quad (15.13)$$

corresponds to the small perturbation which is immobile with respect to the medium:

$$\mathbf{V}_{\text{ph}} = 0.$$

If the medium is moving, the disturbance is carried with it.

Substituting (15.13) in (15.8), we obtain the following equations:

$$(\mathbf{k} \cdot \mathbf{u}) \mathbf{v}' - \mathbf{u} (\mathbf{k} \cdot \mathbf{v}') = 0, \quad (15.14)$$

$$\mathbf{k} \cdot \mathbf{u}' = 0, \quad (15.15)$$

$$(\mathbf{k} \cdot \mathbf{u}) \mathbf{u}' - \rho^{-1} (p' + \rho \mathbf{u} \cdot \mathbf{u}') \mathbf{k} = 0, \quad (15.16)$$

$$\mathbf{k} \cdot \mathbf{v}' = 0, \quad (15.17)$$

$$p' - V_s^2 \rho' - b s' = 0. \quad (15.18)$$

Let us make use of (15.17) in (15.14). Then we take the scalar product of (15.16) with the vector $\mathbf{k}$ and make allowance for (15.15). We write

$$(\mathbf{k} \cdot \mathbf{u}) \mathbf{v}' = 0, \quad (15.19)$$

$$\mathbf{k} \cdot \mathbf{u}' = 0, \quad (15.20)$$

$$p' + \rho \mathbf{u} \cdot \mathbf{u}' = 0, \quad (15.21)$$

$$\mathbf{k} \cdot \mathbf{v}' = 0, \quad (15.22)$$

$$p' - V_s^2 \rho' - b s' = 0. \quad (15.23)$$

Substitution of (15.21) in (15.16) gives us the following set of equations:

$$(\mathbf{k} \cdot \mathbf{u}) \mathbf{u}' = 0, \quad (\mathbf{k} \cdot \mathbf{u}) \mathbf{v}' = 0,$$

$$p' + \rho \mathbf{u} \cdot \mathbf{u}' = 0, \quad p' - V_s^2 \rho' - b s' = 0.$$

Since generally $\mathbf{k} \cdot \mathbf{u} \neq 0$, the velocity, magnetic field and gas pressure are undisturbed in the wave under discussion:

$$\mathbf{v}' = 0, \quad \mathbf{u}' = 0, \quad p' = 0. \quad (15.24)$$

The only disturbed quantities are the density and entropy related by the condition

$$\rho' = -\frac{b}{V_s^2} s'.$$

(15.25)

This is the reason why these disturbances are called the *entropy* waves. They are well known in hydrodynamics (Exercise 15.2). The meaning of an entropy wave is that regions containing hotter but more rarefied plasma can exist in a plasma flow.

The entropy waves are only arbitrarily termed *waves*, since their velocity of propagation with respect to the medium is zero. Nevertheless the entropy waves must be taken into account together with the real waves in such cases as the study of shock waves behavior under small perturbations. Blokhintsev (1945) has considered the passage of small perturbations through a shock in ordinary hydrodynamics. He came to the conclusion that

the entropy wave must be taken into account in order to match the linearized solutions at the shock front

(see Exercise 17.1). In MHD, the entropy waves are important in the problem of evolutionarity of the MHD discontinuities (Chap. 17) and reconnecting current layers (see Part II, Chap. 10). The entropy waves can be principally essential in astrophysical plasma where plasma motions are not slow, for example in *helioseismology* of the chromosphere and corona of the Sun or another star.

Meanwhile the entropy waves are generally (i.e., with the inclusion of particular dissipative processes) believed to be damped ones. For this reason it is commonly assumed that the entropy waves may be ignored, for example, in the problem of solar coronal heating (see Kumar et al. 2006). However Somov et al. (2007) have solved the stability problem for small MHD perturbations in an optically thin, perfectly conducting plasma with a cosmic abundance of elements. It appears that the entropy waves can grow exponentially in a wide range of physical conditions closest to the conditions in stellar coronae with the proper allowance for radiative losses of energy (see Sect. 15.4.4).

15.2.2 Alfvén Waves

The second root of the dispersion equation (15.10),

$$\omega_0^2 = (\mathbf{k} \cdot \mathbf{u})^2 \quad \text{or} \quad \omega_0 = \pm \mathbf{k} \cdot \mathbf{u}, \quad (15.26)$$




Fig. 15.1 The transversal displacements of plasma and magnetic field in the Alfvén wave

corresponds to waves with the phase velocity

$$V_A = \pm \frac{B}{\sqrt{4\pi\rho}} \cos \theta. \quad (15.27)$$

Here θ is the angle between the direction of wave propagation $\mathbf{k}/k$ and the ambient field vector $\mathbf{B}_0$ (Fig. 15.1). In formula (15.27) the value $B = |\mathbf{B}_0|$ and $\rho = \rho_0$. These are the *Alfvén* waves.

By substituting (15.26) in the algebraic equations (15.8) we check that the thermodynamic characteristics of the medium remain unchanged

$$\rho' = 0, \quad p' = 0, \quad s' = 0, \quad (15.28)$$

while the perturbations of the velocity and magnetic field are subject to the conditions

$$\mathbf{v}' = \mp \mathbf{u}', \quad \mathbf{u} \cdot \mathbf{u}' = 0, \quad \mathbf{k} \cdot \mathbf{u}' = 0. \quad (15.29)$$

Thus the Alfvén waves are the displacements of plasma together with the magnetic field frozen into it. They are transversal with respect to both the field direction and the wave vector as shown in Fig. 15.1.

The Alfvén waves have no analogue in hydrodynamics. They are specific to MHD and were called the *magnetohydrodynamic* waves. This term emphasized that, according to (15.28), they do not change the density of a medium. The fact that the Alfvén waves are transversal signifies that

■ a conducting plasma in a magnetic field has a characteristic elasticity resembling that of stretched strings under tension.

The *magnetic tension force* is one of the characteristics of MHD (see Exercise 13.3). According to (15.29), the perturbed quantities are related by an energy equipartition:

$$\frac{1}{2} \rho (v')^2 = \frac{1}{8\pi} (B')^2. \quad (15.30)$$

Let us note also that

the energy of Alfvén waves, much like the energy of oscillations in a stretched string, propagates along the field lines only.

Unlike the phase velocity, the group velocity of the Alfvén waves (15.12)

$$\mathbf{V}_{\text{gr}} = \pm \frac{\mathbf{B}}{\sqrt{4\pi\rho}} \quad (15.31)$$

is directed strictly along the magnetic field; here $\mathbf{B} = \mathbf{B}_0$ of course.

In low density astrophysical plasmas with a strong field, like the solar corona, the Alfvén speed V_A can approach the light speed c (Exercise 15.3). The discovery of Alfvén waves was a major stage in the development of plasma astrophysics (Alfvén 1950).

15.2.3 Magnetoacoustic Waves

The dispersion equation (15.10) has two other branches—two types of waves defined by a bi-square equation

$$\omega_0^4 - k^2 (u^2 + V_s^2) \omega_0^2 + k^2 V_s^2 (\mathbf{k} \cdot \mathbf{u})^2 = 0. \quad (15.32)$$

Its solutions are two values of ω_0^2 , which differ in absolute magnitude, corresponding to two different waves with the phase velocities V_+ and V_- which are equal to

$$V_{\pm}^2 = \frac{1}{2} \left[u^2 + V_s^2 \pm \sqrt{(u^2 + V_s^2)^2 - 4u^2 V_s^2 \cos^2 \theta} \right]. \quad (15.33)$$

These waves are called the *fast* (+) and the *slow* (−) *magnetoacoustic* waves, respectively (van de Hulst 1951). The point is that the entropy of the medium, as follows from (15.8) under condition (15.32), does not change in such waves

$$s' = 0, \quad (15.34)$$

as is also the case in an usual sound wave. Perturbations of the other quantities can be expressed in terms of the density perturbation

$$p' = V_s^2 \rho', \quad (15.35)$$

$$\mathbf{v}' = -\frac{\omega_0}{\rho k^2} \left(\frac{k^2 (\mathbf{k} \cdot \mathbf{u}) \mathbf{u} - \omega_0^2 \mathbf{k}}{\omega_0^2 - (\mathbf{k} \cdot \mathbf{u})^2} \right) \rho', \quad (15.36)$$

$$\mathbf{u}' = \frac{\omega_0^2}{\rho k^2} \left(\frac{k^2 \mathbf{u} - (\mathbf{k} \cdot \mathbf{u}) \mathbf{k}}{\omega_0^2 - (\mathbf{k} \cdot \mathbf{u})^2} \right) \rho'. \quad (15.37)$$

Formulae (15.36) and (15.37) show that the magnetoacoustic waves are *neither* longitudinal *nor* transversal. Perturbations of the velocity and magnetic field intensity, $\mathbf{v}'$ and $\mathbf{u}'$, as differentiated from the Alfvén wave, lie in the $(\mathbf{k}, \mathbf{B}_0)$ plane in Fig. 15.1. They have components both in the direction of the wave propagation $\mathbf{k}/k$ and in the perpendicular direction. That is why

the magnetoacoustic waves generally have a linearly polarized electric field $\mathbf{E}'$ normal to both $\mathbf{B}_0$ and $\mathbf{k}$.

The perturbation of magnetic pressure $B^2/8\pi$ may be written in the form (see definition (15.3))

$$p'_m = \rho \mathbf{u} \cdot \mathbf{u}'$$

or

$$p'_m = \left(\frac{V_{\pm}^2}{V_s^2} - 1 \right) p'. \quad (15.38)$$

Therefore for the fast wave, by virtue of that $V_+^2 > V_s^2$, the perturbation of magnetic pressure p'_m is of the same sign as that of gas pressure p' .

The magnetic pressure and the gas pressure are added in the fast magnetoacoustic wave. The wave propagates faster, since the effective elasticity of the plasma is greater.

A different situation arises with the slow magnetoacoustic wave. In this case $V_-^2 < V_s^2$ and p'_m is *opposite* in sign to p' . Magnetic and gas pressure deviations partially compensate each other. That is why such a slow wave propagates slowly.

15.2.4 The Phase Velocity Diagram

The dependence of the wave velocities on the angle θ between the undisturbed field $\mathbf{B}_0$ and the wave vector $\mathbf{k}$ is clearly demonstrated in a polar diagram—the phase velocity diagram. In Fig. 15.2, the radius-vector length from the origin of the coordinates to a curve is proportional to the corresponding phase velocity (15.11). The horizontal axis corresponds to the direction of the undisturbed magnetic field.

As the angle $\theta \rightarrow 0$, the fast magnetoacoustic wave V_+ transforms to the usual sound one V_s if

$$V_s > V_{A\parallel} = \frac{B}{\sqrt{4\pi\rho}} \equiv u_A \quad (15.39)$$

in Fig. 15.2a or to the Alfvén wave if $V_s < u_A$ in Fig. 15.2b.

For the angle $\theta \rightarrow \pi/2$, the propagation velocities of the Alfvén and slow waves approach zero. As this takes place, both waves convert to the weak tangential





Fig. 15.2 The phase velocities of MHD waves versus the angle θ for the two cases: (a) $u_A < V_s$ and (b) $u_A > V_s$

Fig. 15.3 The phase velocity diagram for the small-amplitude MHD waves in an ideal plasma with a strong magnetic field



discontinuity in which disturbances of velocity and magnetic field are parallel to the front plane. As $\theta \rightarrow \pi/2$, the fast magnetoacoustic wave velocity tends to

$$V_{\perp} = \sqrt{V_{A\parallel}^2 + V_s^2} = \sqrt{u_A^2 + V_s^2}. \quad (15.40)$$

In the *strong* field limit ($V_{A\parallel}^2 \gg V_s^2$) the diagram for the fast magnetoacoustic wave becomes practically isotropic as shown in Fig. 15.3.

Such a wave may be called the “*magnetic sound*” wave since its phase velocity $V_+ \approx V_{A\parallel} \equiv u_A$ is almost independent of the angle θ .

Generally the sound speed is the minimum velocity of disturbance propagation in ordinary hydrodynamics. By contrast, there is *no minimum velocity* in magnetohydrodynamics.

This property is of fundamental importance for what follows in Chaps. 16 and 17—in study of the principal questions related to discontinuous flows of astrophysical plasma. The first of these questions is what kinds of discontinuities can really exist?

MHD waves produce a lot of effects in astrophysical plasma. The fast magneto-acoustic wave *turbulence* can presumably accelerate electrons in solar flares (see Part II, Sect. 12.3.1). The heavy ions observed in interplanetary space after impulsive flares can result from *stochastic acceleration* by the cascading Alfvén wave turbulence (Part II, Sect. 12.3.2).

15.3 Dissipation of Alfvén Waves

15.3.1 Small Damping of Alfvén Waves

We shall start by treating a plane Alfvén wave propagating along a uniform field $\mathbf{B}_0$; so the angle $\theta = 0$ in Fig. 15.1. Perturbations of the magnetic field and the velocity are small and parallel to the z axis:

$$\mathbf{B}' = \{0, 0, b(t, y)\}, \quad \mathbf{v}' = \{0, 0, v(t, y)\}.$$

In general, the damping effects for such a wave are determined by viscosity and conductivity. Let us consider, first, only the uniform finite conductivity σ . In this case we obtain the extended equation of the wave type with a *dissipative term*:

$$\frac{\partial^2 b}{\partial t^2} = u_A^2 \frac{\partial^2 b}{\partial y^2} + \nu_m \frac{\partial^3 b}{\partial^2 y \partial t}. \quad (15.41)$$

Here $u_A = V_{A\parallel}$ and ν_m is the magnetic diffusivity (12.50). In the case of infinite conductivity, (15.41) is reduced to the wave equation and represents an Alfvén wave with velocity u_A .

Let us suppose that the conductivity is finite. We suppose further that the small perturbations are functions of t and y only:

$$b(t, y) = b_0 \exp(i\omega t + \alpha y), \quad v(t, y) = v_0 \exp(i\omega t + \alpha y). \quad (15.42)$$

Here ω , α , b_0 , and v_0 are constants, all of which except ω may be complex numbers. Substituting (15.42) in (15.41) gives us the dispersion equation:

$$\omega^2 + (u_A^2 + i\nu_m\omega)\alpha^2 = 0 \quad (15.43)$$

or

$$\alpha = \pm i \frac{\omega}{u_A} \left(1 + i \frac{\nu_m \omega}{u_A^2} \right)^{-1/2}. \quad (15.44)$$

For small damping

$$\alpha = \pm \left(\frac{v_m \omega^2}{2 u_A^3} + i \frac{\omega}{u_A} \right). \quad (15.45)$$

The distance l_d in which the amplitude of the wave is reduced to $1/e$ is the inverse value of the real part of α . Thus we have

$$l_d = \frac{2 u_A^3}{v_m \omega^2} = \frac{8 \pi \sigma u_A^3}{\omega^2 c^2} = \frac{2 \sigma u_A}{\pi c^2} \lambda^2, \quad (15.46)$$

where $\lambda = 2 \pi u_A / \omega$ is the wave length. The *short waves suffer more damping* than do the long waves.

Since we treat the dissipative effects as small, the expression (15.46) is valid if $\lambda \ll l_d$. Thus we write

$$b(t, y) = b_0 \exp\left(-\frac{y}{l_d}\right) \exp\left[i \omega \left(t - \frac{y}{u_A}\right)\right], \quad (15.47)$$

$$v(t, y) = v_0 \exp\left(-\frac{y}{l_d}\right) \exp\left[i \omega \left(t - \frac{y}{u_A}\right)\right] \quad (15.48)$$

with

$$v_0 = u_A \frac{b_0}{B_0} \left(1 - i \frac{v_m \omega}{2 u_A^2}\right). \quad (15.49)$$

The imaginary part indicates the phase shift of the velocity v in relation to the magnetic perturbation field b . Therefore

$$v(t, y) = u_A \frac{b_0}{B_0} \exp\left(-\frac{y}{l_d}\right) \exp\left\{i \left[\omega \left(t - \frac{y}{u_A}\right) - \varphi \right]\right\}, \quad (15.50)$$

where

$$\varphi = \frac{v_m \omega}{2 u_A^2} = \frac{\omega c^2}{8 \pi \sigma u_A^2} = \frac{\omega c^2 \rho}{2 \sigma B_0^2}.$$

So the existence of Alfvén waves requires an external field B_0 enclosed between two limits.

The magnetic field should be strong enough to make the damping effects small ($l_d \gg \lambda$) but yet weak enough to keep the Alfvén speed well below the velocity of light,

because otherwise the wave becomes an ordinary electromagnetic wave (see Exercise 13.1). In optical and radio frequencies it is not possible to satisfy both conditions. However longer periods often observed in astrophysical plasma leave a wide range between both limits so that Alfvén waves may easily exist.

One of favorable sites for excitation of MHD waves is the solar atmosphere. The chromosphere and corona are highly inhomogeneous media supporting a variety of filamentary structures in the form of arches and loops. The foot points of these structures are anchored in the poles of the photospheric magnetic fields. They undergo a continuous twisting and turning due to convective motions in the subphotospheric layers. This twisting and turning excite MHD waves. The waves then dissipate and heat the corona (see Part II, Sect. 12.5). Presumably this energy is enough to explain coronal heating, but the unambiguous detection of the MHD waves heating the corona is still awaited.

15.3.2 Slightly Damped MHD Waves

The damping effects due to a finite conductivity σ and due to a kinematic viscosity $\nu = \eta/\rho$ (Sect. 12.2.3) can be included in a general treatment of MHD waves of small amplitudes (van de Hulst 1951). Well developed waves are the waves that travel at least a few wave lengths before they lose a considerable fraction of their energy if the two dimensionless parameters

$$p_\nu = \frac{\omega \nu}{c^2} \quad \text{and} \quad p_{\nu_m} = \frac{\omega \nu_m}{c^2}, \quad (15.51)$$

that characterize *two dissipative processes*, are much smaller than the two small dimensionless parameters

$$p_s = \frac{V_s^2}{c^2} \quad \text{and} \quad p_A = \frac{u_A^2}{c^2}, \quad (15.52)$$

that characterize the propagation speeds of undamped waves.

Let us postulate the form

$$X \equiv c^2 / V_{ph}^2 = X_0 (1 - i q) \quad (15.53)$$

for a general solution of the linearized equations of dissipative MHD. Here

$$X_0 = c^2 / V_{ph,0}^2$$

represents any solution for an undamped wave.

We shall not review all special cases here but shall mention only one, the same case as in previous section. For Alfvén wave we find the following solution

$$X = X_m \equiv c^2 / u_A^2, \quad q = (p_\nu + p_{\nu_m}) X_m. \quad (15.54)$$

This shows that, if dissipative effects are small,

the relative importance of resistivity and viscosity as damping effects in Alfvén wave is independent of frequency ω .

The *damping length*, i.e., the distance l_d , in which the amplitude of a wave decreases by a factor $1/e$, and the *damping time* τ_d , in which this distance is covered by the wave, can be found:

$$l_d = \frac{1}{kq} = \frac{u_A}{q\omega} = \frac{u_A^3}{\omega^2(\nu + \nu_m)}, \quad (15.55)$$

$$\tau_d = \frac{l_d}{u_A} = \frac{1}{q\omega} = \frac{u_A^2}{\omega^2(\nu + \nu_m)}. \quad (15.56)$$

So the high frequency waves have a short damping length and time.

The magnetoacoustic waves (Sect. 15.2.3), being compressional, have an additional contribution to their damping rate from compressibility of the plasma. If dissipative effects are not small, they result in additional waves propagating in a homogeneous medium (see Sect. 17.3).

15.4 Stability of Plasma-Compressing Waves

Let us consider the stability problem for small MHD perturbations in an optically thin, perfectly conducting, uniform plasma with a cosmic abundance of elements. In Sect. 15.4.1, we formulate the problem in the linear approximation and derive a general dispersion equation. In Sect. 15.4.2, we consider two special cases: (a) small perturbations that do not change the plasma density and (b) perturbations accompanied by density variations. Subsequently, in Sect. 15.4.3, the dispersion equation for magnetoacoustic and entropy waves is written in the strong-field approximation. We show in Sects. 15.4.4 and 15.4.5 that the entropy waves can be unstable, they can grow exponentially, while the magnetoacoustic waves are always damped.

15.4.1 Derivation of the Dispersion Equation

Disregarding the damping effects of ordinary and magnetic viscosities, considered in Sect. 15.3, let us write the set of MHD equations as

$$\begin{aligned} \frac{d\rho}{dt} + \rho \operatorname{div} \mathbf{v} &= 0, \\ \rho \frac{d\mathbf{v}}{dt} + \nabla \left(p + \frac{B^2}{8\pi} \right) - (\mathbf{B} \cdot \nabla) \mathbf{B} \frac{1}{4\pi} &= 0, \end{aligned}$$

$$\begin{aligned}\rho T \frac{ds}{dt} &= \frac{1}{\gamma - 1} \frac{dp}{dt} - \frac{\gamma}{\gamma - 1} \frac{p}{\rho} \frac{d\rho}{dt} = \\ &= \nabla_{\parallel} \cdot (\kappa_{\parallel} \nabla_{\parallel} T) + \nabla_{\perp} \cdot (\kappa_{\perp} \nabla_{\perp} T) - \mathcal{L}(\rho, T) + \mathcal{H},\end{aligned}\quad (15.57)$$

$$\frac{d\mathbf{B}}{dt} + \mathbf{B} \operatorname{div} \mathbf{v} - (\mathbf{B} \cdot \nabla) \mathbf{v} = \mathbf{0},$$

$$\operatorname{div} \mathbf{B} = 0,$$

$$p = \frac{R}{\mu} \rho T.$$

Here s is the entropy per unit mass, $\gamma = c_p/c_v$ is the ratio of the specific heats at constant pressure and constant volume, and $\mathcal{L}$ is the rate of energy losses through the optically thin plasma radiation per unit volume measured in $\text{erg s}^{-1} \text{cm}^{-3}$ (see formula (12.23) and Fig. 12.1).

The *heat conduction* plays a major role in a high-temperature plasma, for example, in the solar corona. It is anisotropic in the presence of a strong magnetic field. The heat conduction along ($\kappa_{\parallel}$) and across ($\kappa_{\perp}$) the field is determined, respectively, by electrons and, for a completely ionized hydrogen plasma, by protons (Spitzer 1962). The contribution from heavier ions to the transverse heat conduction can be ignored in view of their low abundance. Meanwhile, precisely these ions determine the huge rate of energy losses through EUV and soft X-ray radiation (Cox and Tucker 1969).

We will be interested in the behavior of small perturbations against the background of a uniform equilibrium plasma with a frozen-in uniform constant magnetic field $\mathbf{B}_0$. The existence of a steady state in such a plasma implies the presence of a steady uniform heating source whose power $\mathcal{H}$ is equal to the rate of *radiative energy losses*. Since the approximation of an ideal conductivity is often applicable in astrophysical plasma, we can disregard the Joule heating and the related *overheating instability* (see Kadomtsev 1966). Instead, various heating mechanisms are widely discussed in astrophysical literature. For example, the heating by particular waves, X-ray radiation, microflares, etc. is taken into account when the stellar atmospheres are investigated. Unfortunately, nothing is known about the physical properties of this heating from observations. Therefore, not the Joule heating, but a background source of heat independent of plasma parameters is commonly assumed; i.e., a constant $\mathcal{H}$ is added to the right-hand side of (15.57).

Let us represent all of the physical variables as $f(\mathbf{r}, t) = f_0 + f'(\mathbf{r}, t)$. In particular, the gas pressure is $p(\mathbf{r}, t) = p_0 + p'(\mathbf{r}, t)$, where $p_0 = (R/\mu) \rho_0 T_0$, the constants ρ_0 and T_0 are the unperturbed plasma density and temperature. In addition, since the medium and the field are assumed to be uniform, we seek a solution in the form of plane waves

$$f'(\mathbf{r}, t) = f_1 \exp i(\mathbf{k} \cdot \mathbf{r} - \omega t). \quad (15.58)$$

Instead of the initial set of partial differential equations, we obtain a set of linear algebraic equations with constant coefficients:

$$\omega \rho_1 - \rho_0 (\mathbf{k} \cdot \mathbf{v}_1) = 0, \quad (15.59)$$

$$\omega \rho_0 \mathbf{v}_1 - \mathbf{k} \left(p_1 + \frac{\mathbf{B}_0 \cdot \mathbf{B}_1}{4\pi} \right) + (\mathbf{k} \cdot \mathbf{B}_0) \frac{\mathbf{B}_1}{4\pi} = 0, \quad (15.60)$$

$$\frac{(-i)\omega}{\gamma - 1} (p_1 - c_s^2 \rho_1) + Q_\rho \rho_1 + Q_T T_1 + \mathcal{K} T_1 = 0, \quad (15.61)$$

$$\omega \mathbf{B}_1 - \mathbf{B}_0 (\mathbf{k} \cdot \mathbf{v}_1) + (\mathbf{k} \cdot \mathbf{B}_0) \mathbf{v}_1 = 0, \quad (15.62)$$

$$(\mathbf{k} \cdot \mathbf{B}_1) = 0, \quad (15.63)$$

$$\frac{p_1}{\rho_0} = \frac{\rho_1}{\rho_0} + \frac{T_1}{T_0}. \quad (15.64)$$

Here $c_s^2 = \gamma p_0 / \rho_0$ is the sound speed squared; the coefficients

$$Q_\rho = \left. \frac{\partial \mathcal{L}}{\partial \rho} \right|_T, \quad Q_T = \left. \frac{\partial \mathcal{L}}{\partial T} \right|_\rho \quad (15.65)$$

and

$$\mathcal{K} = \kappa_{\parallel} k_{\parallel}^2 + \kappa_{\perp} k_{\perp}^2 \quad (15.66)$$

characterize the plasma emissivity and anisotropic heat conductivity. The constant $\mathcal{H}$ vanishes after the differentiation; there are no derivatives of it in (15.61).

Let us eliminate ρ_1 and p_1 from the set of equations (15.59)–(15.64) using the relations

$$\omega \rho_1 = \rho_0 (\mathbf{k} \cdot \mathbf{v}_1), \quad p_1 = -\Sigma (\mathbf{k} \cdot \mathbf{v}_1), \quad (15.67)$$

where

$$\Sigma = \frac{(-i)\omega \rho_0 c_s^2 - (\gamma - 1) \rho_0^2 Q_\rho + (\gamma - 1) T_0 (Q_T + \mathcal{K})}{\omega [i \omega \rho_0 - \mu (\gamma - 1) (Q_T + \mathcal{K}) / R]} \rho_0. \quad (15.68)$$

As a result, we obtain two equations,

$$\omega \rho_0 \mathbf{v}_1 - \mathbf{k} \left[-\Sigma (\mathbf{k} \cdot \mathbf{v}_1) + \frac{\mathbf{B}_0 \cdot \mathbf{B}_1}{4\pi} \right] + (\mathbf{k} \cdot \mathbf{B}_0) \frac{\mathbf{B}_1}{4\pi} = 0 \quad (15.69)$$

and

$$\omega \mathbf{B}_1 - \mathbf{B}_0 (\mathbf{k} \cdot \mathbf{v}_1) + (\mathbf{k} \cdot \mathbf{B}_0) \mathbf{v}_1 = 0, \quad (15.70)$$

for two unknowns, $\mathbf{v}_1$ and $\mathbf{B}_1$, as well as the standard condition of magnetic field divergence:

$$(\mathbf{k} \cdot \mathbf{B}_1) = 0. \quad (15.71)$$

Multiplying (15.69) scalarwise by the wave vector $\mathbf{k}$ and using condition (15.71), we obtain

$$(\mathbf{B}_0 \cdot \mathbf{B}_1) = \frac{4\pi}{k^2} (\Sigma k^2 + \omega \rho_0) (\mathbf{k} \cdot \mathbf{v}_1). \quad (15.72)$$

The scalar product of (15.69) by $\mathbf{B}_0$ yields

$$(\mathbf{B}_0 \cdot \mathbf{v}_1) = -\frac{\Sigma}{\omega \rho_0} (\mathbf{B}_0 \cdot \mathbf{k}) (\mathbf{k} \cdot \mathbf{v}_1). \quad (15.73)$$

Multiplying (15.70) scalarwise by $\mathbf{B}_0$, we find

$$\omega (\mathbf{B}_0 \cdot \mathbf{B}_1) - B_0^2 (\mathbf{k} \cdot \mathbf{v}_1) + (\mathbf{k} \cdot \mathbf{B}_0) (\mathbf{B}_0 \cdot \mathbf{v}_1) = 0. \quad (15.74)$$

Substituting formulae (15.72) and (15.73) into (15.74) yields the sought-for dispersion equation

$$(\mathbf{k} \cdot \mathbf{v}_1) \left[-\omega \frac{4\pi}{k^2} (\Sigma k^2 + \omega \rho_0) + B_0^2 + \frac{\Sigma}{\omega \rho_0} (\mathbf{k} \cdot \mathbf{B}_0)^2 \right] = 0. \quad (15.75)$$

The function $\Sigma = \Sigma(\omega)$ in this equation is defined by formula (15.68).

15.4.2 Two Types of MHD Waves

15.4.2.1 Plasma-Noncompressing Waves

Let in (15.75) $(\mathbf{k} \cdot \mathbf{v}_1) = 0$. It then follows from (15.72) and (15.73) that

$$(\mathbf{B}_0 \cdot \mathbf{B}_1) = 0, \quad (\mathbf{B}_0 \cdot \mathbf{v}_1) = 0,$$

and (15.69) and (15.70) take the form

$$4\pi \omega \rho_0 \mathbf{v}_1 = -(\mathbf{k} \cdot \mathbf{B}_0) \mathbf{B}_1, \quad (15.76)$$

$$\omega \mathbf{B}_1 = -(\mathbf{k} \cdot \mathbf{B}_0) \mathbf{v}_1. \quad (15.77)$$

If $\mathbf{v}_1 \neq 0$, then $\mathbf{B}_1 \neq 0$.

Multiplying (15.76) by (15.77), we find the first solution to the dispersion equation (15.75),

$$\omega^2 = \frac{(\mathbf{k} \cdot \mathbf{B}_0)^2}{4\pi \rho_0} = k^2 u_A^2 \cos^2 \theta,$$

which corresponds to *Alfvén waves* (see Sect. 15.2.2). The constant $u_A^2 = B_0^2 / 4\pi \rho_0$ is the Alfvén speed squared. Being transverse electromagnetic perturbations by

their nature, the Alfvén waves do not affect the thermodynamic parameters of the medium. Therefore they are not involved in the radiative cooling and the conductive transfer of energy under consideration.

15.4.2.2 Plasma-Compressing Waves

Let now $(\mathbf{k} \cdot \mathbf{v}_1) \neq 0$. It then follows from (15.75) that

$$-\omega \frac{4\pi}{k^2} (\Sigma k^2 + \omega \rho_0) + B_0^2 + \frac{\Sigma}{\omega \rho_0} (\mathbf{k} \cdot \mathbf{B}_0)^2 = 0. \quad (15.78)$$

Let us introduce the notation

$$a_\rho = \frac{\mu(\gamma - 1) Q_\rho}{R V_s T_0}, \quad a_T = \frac{\mu(\gamma - 1) Q_T}{R V_s \rho_0}, \quad a'_T = \frac{\mu(\gamma - 1)}{R V_s \rho_0} (Q_T + \mathcal{K}).$$

In this notation (15.78) takes the form

$$\begin{aligned} \omega^5 + i V_s a'_T \omega^4 - k^2 (V_s^2 + u_A^2) \omega^3 - i k^2 \left[V_s^3 (a'_T - a_\rho) \frac{1}{\gamma} + V_s u_A^2 a'_T \right] \omega^2 + \\ + V_s^2 u_A^2 k^4 \cos^2 \theta \cdot \omega + \frac{i}{\gamma} (a'_T - a_\rho) V_s^3 u_A^2 k^4 \cos^2 \theta = 0. \end{aligned} \quad (15.79)$$

In the approximation of an ideal medium,

$$a_\rho = a_T = \mathcal{K} = 0,$$

and the dispersion equation (15.79) is simplified to

$$\omega \left[\omega^4 - k^2 (V_s^2 + u_A^2) \omega^2 + V_s^2 u_A^2 k^4 \cos^2 \theta \right] = 0. \quad (15.80)$$

This form is typical for the description of *entropy* and *magnetoacoustic* waves (e.g., Syrovatskii 1957; see Sects. 15.2.1 and 15.2.3).

Returning to the general case and taking into account the fact that $R/\mu = c_v(\gamma - 1)$, let us introduce the characteristic frequencies

$$\omega_T = V_s a'_T = \frac{1}{c_v \rho_0} (Q_T + \mathcal{K}) \quad (15.81)$$

and

$$\omega_\rho = V_s a_\rho = \frac{Q_\rho}{c_v T_0}. \quad (15.82)$$

The dispersion equation (15.79) for small-amplitude waves that change the plasma density with allowance made for radiative energy losses and the anisotropic heat conduction in a magnetic field can then be rewritten as

$$\begin{aligned} \omega^5 + i \omega_T \omega^4 - k^2 u_A^2 (1 + \beta) \omega^3 - i k^2 u_A^2 \left[(\omega_T - \omega_\rho) \frac{\beta}{\gamma} + \omega_T \right] \omega^2 + \\ + \beta u_A^4 k^4 \cos^2 \theta \cdot \omega + \frac{i}{\gamma} (\omega_T - \omega_\rho) \beta u_A^4 k^4 \cos^2 \theta = 0. \end{aligned} \quad (15.83)$$

Here the dimensionless parameter $\beta = V_s^2 / u_A^2$ is a quantity proportional to the ratio of the gas pressure in the plasma to the magnetic pressure.

15.4.3 Waves in Strong Magnetic Field

Before we turn to quantitative calculations that include the peculiar features of the radiative cooling, we must pass to the following dimensionless variables in the dispersion equation (15.83):

$$\Omega = i \frac{\omega}{k u_A} = i \frac{v_{ph}}{u_A}, \quad \Omega_T = \frac{\omega_T}{k u_A}, \quad \Omega_\rho = \frac{\omega_\rho}{k u_A}.$$

As a result, we obtain the dimensionless equation

$$\begin{aligned} \Omega^5 - \Omega_T \Omega^4 + (1 + \beta) \Omega^3 - \left[(\Omega_T - \Omega_\rho) \frac{\beta}{\gamma} + \Omega_T \right] \Omega^2 + \\ + \beta \cos^2 \theta \cdot \Omega - \frac{\beta}{\gamma} (\Omega_T - \Omega_\rho) \cos^2 \theta = 0. \end{aligned} \quad (15.84)$$

It is applicable for any values of the parameters β , both small and large ones.

In strong magnetic fields $u_A^2 \gg V_s^2$. Therefore, let us pass to the limit $\beta \rightarrow 0$ in (15.84). We then obtain

$$\Omega^2 (\Omega^2 + 1) (\Omega - \Omega_T) \approx 0. \quad (15.85)$$

The first root of the dispersion equation (15.85)

$$\Omega^2 \approx 0 \quad (15.86)$$

corresponds to *slow* magnetoacoustic waves (Sect. 15.2.3). If $u_A > V_s$, then their phase velocity v_{ph} along the magnetic field is equal to the speed of sound V_s and, hence, tends to zero as $\beta \rightarrow 0$. The sign “ $\approx$ ” in formula (15.86) emphasizes that Ω^2 is a small quantity proportional to the small parameter β that is not identically equal zero.

The second root of (15.85)

$$\omega = \pm k u_A \quad (15.87)$$

corresponds to *fast* magnetoacoustic waves (Sect. 15.2.3). In the strong-field limit ($u_A^2 \gg V_s^2$), the phase velocity diagram for these waves is almost isotropic (Fig. 15.3).

The third root of the dispersion equation (15.85)

$$\omega = -i \omega_T \quad (15.88)$$

corresponds to *entropy waves* (Sect. 15.2.1).

15.4.4 The Instability of Entropy Waves

Entropy waves are stationary relative to the medium. Indeed, substituting (15.88) into (15.58), we see that

$$f'(\mathbf{r}, t) = f_1 \exp i(\mathbf{k} \cdot \mathbf{r}) \cdot \exp(-\omega_T t). \quad (15.89)$$

According to (15.89), the entropy waves grow exponentially if $\omega_T < 0$. In view of definition (15.81), the instability condition can be rewritten as

$$Q_T + \mathcal{K} < 0. \quad (15.90)$$

Since the coefficient $\mathcal{K}$ is always greater than zero by definition (15.66), the heat conduction has a stabilizing effect on the entropy waves.

Given definition (15.65), it thus follows that the instability can manifest itself only at negative values of the derivative of the rate of radiative energy losses,

$$\mathcal{L} \simeq -n^2 q(T), \text{ erg s}^{-1} \text{ cm}^{-3},$$

(see the cooling function $q(T)$ in Fig. 12.1) with respect to the temperature. The logarithmic derivative of the cooling function,

$$\alpha(T) = d \ln q(T) / d \ln T,$$

is shown in Fig. 15.4.

Thus, in the strong-field approximation, the general dispersion equation (15.84) is reduced to the simple equation (15.85) which allowed us to obtain the explicit condition (15.90) for the instability of entropy waves. Now we return to the general case, to (15.84) with arbitrary values of the parameter β .

Fig. 15.4 Logarithmic derivative of the radiative cooling function q with respect to the temperature for an optically thin plasma with a cosmic abundance of elements. (From Somov et al. 2007; reproduced with permission © Pleiades Publishing, Ltd)





Fig. 15.5 The growth time for entropy wave instability vs. temperature at fixed plasma densities. The weak dependence on the magnetic field manifests itself as a slight stratification of the curves increasing with density. (From Somov et al. 2007; reproduced with permission © Pleiades Publishing, Ltd)




The exact solutions of this equation show that there are temperature ranges in which the entropy waves grow exponentially (Somov et al. 2007). The characteristic growth time τ is plotted against the temperature T in Fig. 15.5 for three fixed plasma densities: $n_1 = 10^9$, $n_2 = 10^{10}$, $n_3 = 10^{11} \text{ cm}^{-3}$.

A weak dependence on the magnetic field enters into the calculations with formulae for the heat conductivity (Spitzer 1962):

$$\kappa_{\parallel} \simeq 9 \times 10^{-7} T^{5/2}, \quad \frac{\kappa_{\perp}}{\kappa_{\parallel}} \simeq 2 \times 10^{-11} \frac{n^2}{T^3 B^2}, \quad (15.91)$$

where the heat conductivity is measured in CGS units and the magnetic field B is measured in Gauss. The calculations were performed for a wide range of field strengths, $B = (10^{-1} - 10^4) \text{ G}$. This manifested itself in Fig. 15.5 as a slight “stratification” of the curves for the instability growth time τ at the plasma

Fig. 15.6 The growth time for entropy wave instability as a function of temperature and the wavelength λ , which is indicated as a number near the curves; the unit of length is taken to be 10^8 cm. (From [Somov et al. 2007](#); reproduced with permission © Pleiades Publishing, Ltd)



densities $n_2 = 10^{10}$ and $n_3 = 10^{11} \text{ cm}^{-3}$. Here we took the angle θ between the wave vector $\mathbf{k}$ and the magnetic field vector $\mathbf{B}_0$ to be $\pi/4$ and chose the wavelength to be $\lambda = 10^8$ cm.

The wavelength dependence of the instability growth time τ is shown in Fig. 15.6. The magnetic field strength is $B_0 = 1,000$ G.

The growth time for the entropy waves in stellar coronae can vary over a wide range: from tenths of a second to tens minutes.

The higher the plasma density, the faster the instability growth. It is significant that two “main” minima (regions of strong instability) are present at temperatures $T_1 \approx 2 \times 10^4$ K and $T_2 \approx 3 \times 10^5$ K.


The instability mechanism is simple. In the temperature regions of a rapid decrease in the radiative loss function with temperature, a small decrease in temperature causes a large increase in the rate of radiative energy losses. Conversely a small increase in temperature is accompanied by a decrease in the rate of radiative plasma cooling. As a result, small perturbations grow rapidly.

Clearly, in general,

the entropy waves can be unstable in any compressible optically thin medium whose radiative cooling depends on the temperature in the form of one or more sharp maxima.

In astrophysical and laboratory plasmas, these maxima are produced by the radiation of a small admixture of heavy ions. However, in contrast to the astrophysical plasma, the approximation of an ideal conductivity is inapplicable in the laboratory plasma. The Joule heating becomes the dominant heating mechanism. The dispersion equation includes the derivative with respect to the temperature not only the bulk radiation intensity but also of the plasma conductivity. This leads to instabilities and complex dynamics of plasma, for example, in emitting Z-pinch (Imshennik and Bobrova 1997).

Fig. 15.7 Damping decrement of fast magnetoacoustic waves as a function of the plasma temperature and density for the same parameters as those in Fig. 15.6. (From Somov et al. 2007; reproduced with permission © Pleiades Publishing, Ltd)



15.4.5 The Damping of Magnetoacoustic Waves

In contrast to the entropy waves, the magnetoacoustic waves are damped. At the assumed physical model parameters consistent with our views of the solar and stellar coronae, the damping decrement is

$$d = \frac{\text{Im } \omega}{\text{Re } \omega} < 0.$$


For *fast* magnetoacoustic waves, Fig. 15.7 presents the solutions to the dispersion equation at the same plasma density, angle θ and the magnetic field strength as those in Fig. 15.6. The damping decrement d in the temperature range $10^4 \leq T \leq 2 \times 10^6$ K is very small: $-d < 10^{-6}$.

For *slow* magnetoacoustic waves, the damping decrement is several orders of magnitude larger (Fig. 15.8). At $T_2 \approx 3 \times 10^5$ K, the damping decrement lies within the range $\sim 0.05 < -d < 0.16$. This corresponds to a decrease in the wave amplitude by a factor of e in a time τ equal to ~ 3 –1 wave periods $\tau_\omega = 2\pi / \text{Re } \omega$. As the temperature decreases to $T \approx T_1 \approx 2 \times 10^4$ K, the damping decrement increases to values of the order of several seconds. Thus

the damping decrement for slow magnetoacoustic waves in stellar coronae is 4–6 orders of magnitude larger than that for fast waves.

What is the cause of such a large difference?—In a fast magnetoacoustic wave, the sign of the change in magnetic pressure is the same as that of the change in gas pressure (see formula (15.38)), i.e. the magnetic and gas pressures add up. Where the plasma density increases, the magnetic pressure also increases, preventing the plasma compression. The effective plasma elasticity for a fast wave is larger. The picture is different in a slow wave. In this case, the changes in magnetic and gas pressures have opposite signs.

Fig. 15.8 Damping decrement for slow magnetoacoustic waves. Parameters are the same as those in Fig. 15.7. (From Somov et al. 2007; reproduced with permission © Pleiades Publishing, Ltd)



Where the plasma density increases, the magnetic field decreases, without preventing the plasma from compressing. As a result, the radiative energy losses in a slow magnetoacoustic wave grow faster than that ones in a fast wave.

For this reason, the radiative damping of slow waves is much stronger than that of fast waves.

One would think that slow waves resemble more closely the ordinary sound and that the excitation of thermal instability might be expected for them (Field 1965). Instead, the calculations demonstrate only rapid damping of slow waves. However there is no contradiction in this fact. The point is that the magnetoacoustic waves are neither longitudinal nor transverse; see formulae (15.36) and (15.37). The perturbations in velocity $\mathbf{v}_1$ and magnetic field $\mathbf{B}_1$ lie in the plane $(\mathbf{k}, \mathbf{B}_0)$ and have components both in the direction of wave propagation $\mathbf{k}/k$, i.e., partly in the direction transverse to the field $\mathbf{B}_0$. Moreover, according to formulae (15.36) and (15.37), the longitudinal components are not independent of the transverse ones. For this reason, the stabilizing effect of a strong magnetic field (see (49) in Field 1965) dominates over the excitation of oscillations by *thermal instability* in the wide range of parameters under consideration. Of course, the damping of oscillations due to the radiative energy losses is present and turns out to be strong for slow waves.

The viscous and ohmic damping of magnetoacoustic waves was described, for example, by Alfvén and Fälthamar (1963), Akhiezer et al. (1975).

15.5 MHD Oscillations in the Solar Corona

In the solar atmosphere, particularly in the corona, the low-frequency MHD oscillations can be studied comprehensively. They are observed almost at all wavelengths (see Aschwanden 2004). Most of these oscillations are commonly interpreted as standing oscillations of various types in coronal magnetic loops. Meanwhile the oscillations of coronal loops observed from *TRACE* satellite in EUV are, as a

rule, damped rapidly. According to [Aschwanden et al. \(2003\)](#), the ratio of the characteristic damping time τ_d to the oscillation period τ_ω is $\tau_d/\tau_\omega = 1.8 \pm 0.8$ in the range of periods $\tau_\omega = 317 \pm 114$ s. Such *rapid damping* of the MHD oscillations is difficult to explain.

The current models of damping mechanisms consider the following factors: (a) nonideality of the solar plasma as an MHD medium, i.e. the presence of electric resistivity, viscosity, and heat conduction and radiative losses of energy in it; (b) the escape of waves from the magnetic loop through its side boundaries into corona and through the loop footpoints into the chromosphere; (c) phase mixing in regions where the magnetic field and the plasma are nonuniform; (d) resonant absorption. The first two factors are believed to be negligible and the last two factors are believed to be the main ones (see [Aschwanden 2004](#)).

Since the coronal oscillations are damped rapidly, the question of their excitation mechanism also arises. If the oscillations grew rapidly, there would be no problem: low-amplitude perturbations would suffice. However the formal number of oscillations is, in average,

$$N_{obs} = \frac{\tau_d}{\tau_\omega} \approx 2. \quad (15.92)$$

The current models just postulate instantaneous excitation of large-amplitude MHD oscillations in an isolated magnetic flux tube.

Why rapidly damped oscillations are seen best in a small group of loops precisely in EUV radiation is probably a key question. Contrary to popular belief, the answer may be simple.

Where the rate of energy losses via optically thin plasma radiation is at a maximum (i.e. at $T \sim 10^5$ K), the brightness of the oscillating loops is at a maximum (i.e. in EUV) and, as a consequence, the MHD oscillations are damped more rapidly than in other places.

The calculations presented in Sect. 15.4.5 show that the predicted number of oscillations for *slow* magnetoacoustic waves is

$$N_{th} = \frac{\tau}{\tau_\omega} \approx 1 - 3 \quad (15.93)$$

in the range of temperatures corresponding to the second main maximum, i.e. $T_2 \approx 3 \times 10^5$ K.

Moreover, at the plasma density $n_1 = 10^9 \text{ cm}^{-3}$, the damping decrement remains in the same range as the temperature decreases to $T \approx T_1 \approx 2 \times 10^4$ K. For the 11 events with oscillating coronal loops that were studied in detail by [Aschwanden et al. \(2003\)](#), the plasma density inside the loops was, on average, $n_{int} = (1.4 \pm 0.7) \times 10^9 \text{ cm}^{-3}$. Consequently $n_{int} \approx n_1$ and $N_{th} \approx N_{obs}$. Thus we have reason to suggest that

the observed oscillations are a manifestation of the rapid damping of slow waves due to the radiative energy losses in the solar corona.

The significant advantage of slow magnetoacoustic waves over *fast* ones is that the regions of reduced magnetic field in the former coincide with the regions of enhanced plasma density. It is here that the rapid radiative losses manifest themselves. Meanwhile, as calculations show, fast magnetoacoustic waves radiate little and are damped too slowly (see Fig. 15.7).

The excitation mechanism of magnetoacoustic waves should be studied further. The rapid growth of *entropy waves* calculated in Sect. 15.4.4 must probably be taken into account to explain this phenomenon. Clearly, we cannot ignore the fast plasma flows (particularly those from reconnecting current layers) and, hence, the growing entropy waves transferred by these flows in the solar corona. Magnetic and plasma nonuniformities probably create conditions for a nonlinear interaction and transformation of part of the energy of entropy waves into magnetoacoustic waves.

15.6 Practice: Exercises and Answers

Exercise 15.1. Evaluate the sound speed in the solar corona.

Answer. For an ideal gas with constant specific heats c_p and c_v , the sound speed (15.6) is

$$V_s = \left(\gamma_g \frac{p}{\rho} \right)^{1/2}, \quad (15.94)$$

where $\gamma_g = c_p/c_v$. Let us consider the coronal plasma as a “monatomic gas” ($\gamma_g = 5/3$) of electrons and protons with $T_e = T_p = T \approx 2 \times 10^6$ K and $n_e = n_p = n$. So $p = 2nk_B T$ and $\rho = nm_p$. Hence

$$V_s = \left(\frac{10}{3} \frac{k_B}{m_p} \right)_s^{1/2} \sqrt{T} = 1.66 \times 10^4 \sqrt{T(\text{K})}, \text{ cm s}^{-1}. \quad (15.95)$$

In the solar corona $V_s \approx 230 \text{ km s}^{-1}$.

Exercise 15.2. Consider entropy waves in ordinary hydrodynamics.

Answer. Let us take the linear algebraic equations (15.19)–(15.23). In the absence of a magnetic field we put $\mathbf{u} = 0$ and $\mathbf{u}' = 0$. It follows that the perturbation of the velocity $\mathbf{v}'$ can be an arbitrary value except the gas pressure must be undisturbed. Instead of (15.24), we write

$$\mathbf{v}' \neq 0, \quad p' = 0. \quad (15.96)$$

Perturbations of the density and entropy remain to be related by condition (15.25). So the velocity perturbation is independent of the entropy perturbation and, according to (15.13) and (15.22), satisfies the equation

$$\mathbf{k} \cdot \mathbf{v}' = \frac{\omega}{v} v'_x + k_y v'_y = 0. \quad (15.97)$$

This is in the reference frame in which $v = v_x$.

Note that for such velocity perturbation (see [Landau and Lifshitz 1959a](#), Chap. 9):

$$\text{rot } \mathbf{v}' \neq 0. \quad (15.98)$$

That is why the wave is called the *entropy-vortex* wave.

I In the presence of a magnetic field in plasma, it is impossible to create a vortex without a perturbation of the magnetic field.

For this reason, in a MHD entropy wave, the only disturbed quantities are the entropy and the density (see (15.25)).

Exercise 15.3. Show that the inclusion of the displacement current modifies the dispersion relation for the Alfvén waves (15.26) to the following equation

$$\omega_0^2 = \frac{(\mathbf{k} \cdot \mathbf{u})^2}{1 + u^2/c^2} \quad \text{or} \quad \omega_0 = \pm \frac{\mathbf{k} \cdot \mathbf{u}}{\sqrt{1 + u^2/c^2}}. \quad (15.99)$$

So the phase velocity of the relativistic Alfvén waves

$$V_A = \pm \frac{B}{\sqrt{4\pi\rho}} \cos \theta \frac{1}{\sqrt{1 + B^2/4\pi\rho c^2}}, \quad (15.100)$$

which coincides with the Alfvén formula (13.36).

Exercise 15.4. Discuss the following situation. A star of the mass M moves along a uniform magnetic field $\mathbf{B}_0$ at a constant velocity $\mathbf{v}_0$ which exceeds the phase velocity of a fast magnetoacoustic wave ([Dokuchaev 1964](#)).

Hint. The moving star emits magnetoacoustic waves by the Cherenkov radiation (see Exercises 7.2–7.5).

Chapter 16

Discontinuous Flows in a MHD Medium

Abstract The phenomena related to shock waves and other discontinuous flows in astrophysical plasma are so numerous that the study of MHD discontinuities on their own is of independent great interest for space sciences.

16.1 Discontinuity Surfaces in Hydrodynamics

16.1.1 The Origin of Shocks in Ordinary Hydrodynamics

First of all, let us recall the way the shock waves are formed in ordinary hydrodynamic media without a magnetic field. Imagine a piston moving into a tube occupied by a gas. Let the piston velocity increase from zero by small jumps δv . As soon as the piston starts moving, it begins to rake the gas up and compress it. The front edge of the compression region thereby travels down the undisturbed gas inside the tube with the velocity of sound


$$V_s = \left(\frac{\partial p}{\partial \rho} \right)_s^{1/2}. \quad (16.1)$$

Each following impulse of compression $\delta \rho$ will propagate in a denser medium and hence with greater velocity. Actually, the derivative of the sound speed with respect to density

$$\frac{\partial V_s}{\partial \rho} = \frac{1}{2} \left(\frac{\partial^2 p}{\partial \rho^2} \right)_s \left(\frac{\partial p}{\partial \rho} \right)_s^{-1/2} \approx \sqrt{\gamma_g} (\gamma_g - 1) \rho^{(\gamma_g - 3)/2} > 0,$$

since for all real substances $\gamma_g > 1$ in the adiabatic process $p \sim \rho^{\gamma_g}$. Therefore $\delta V_s > 0$.

Fig. 16.1 The behavior of small perturbations in front of a moving piston



As a consequence of this fact, successive compression impulses will catch up with each other as shown in Fig. 16.1a. As a result, the compression region front steepens (Fig. 16.1b). The gradients of the gas parameters become so large that the description of the gas as a hydrodynamic medium (Sect. 12.2) is no longer valid. The density, pressure and velocity of the gas change abruptly over a distance comparable to a particle's mean free path λ .

The physical processes inside such a jump, called a *shock wave*, are determined by the kinetic phenomena in the gas. As far as the hydrodynamic approximation is concerned,

the surface, at which the continuity of the hydrodynamic parameters of a medium is violated, represents some discontinuity surface—a discontinuous solution of the hydrodynamic equations.

It stands to reason that some definite boundary conditions must hold at the discontinuity surface. What are they?

16.1.2 Boundary Conditions and Classification

Let us choose a frame of reference connected with a discontinuity surface. The frame is supposed to move with a constant velocity with respect to the medium. Generally, if the gas flow is non-stationary in the vicinity of the discontinuity, we could consider the discontinuity surface over a small period of time, so that the changes of velocity and other hydrodynamic quantities in time could be neglected.

In order to formulate the boundary conditions, let us consider an element of the discontinuity surface. Let the axis x be directed normally to it.

The flux of mass through such a surface element must conserve:

$$\rho_1 v_{x1} = \rho_2 v_{x2}. \quad (16.2)$$

Here the indices 1 and 2 refer to the two sides of the discontinuity surface.

In this chapter, the difference in a quantity across the discontinuity surface will be designated by curly brackets, e.g.,

$$\{ \rho v_x \} = \rho_1 v_{x1} - \rho_2 v_{x2}.$$

Then (16.2) is rewritten as

$$\{ \rho v_x \} = 0. \quad (16.3)$$

The energy flux must also be continuous at the discontinuity surface. For a hydrodynamic medium without a magnetic field (cf. formula (12.76)) we obtain the following condition for the energy flux conservation:

$$\left\{ \rho v_x \left(\frac{v^2}{2} + w \right) \right\} = 0. \quad (16.4)$$

Here w is the specific enthalpy (9.35).

The momentum flux must be also continuous (cf. (12.77)):

$$\Pi_{\alpha\beta} = p \delta_{\alpha\beta} + \rho v_\alpha v_\beta, \quad \alpha = x.$$

The continuity of the x -component of the momentum flux means that

$$\{ p + \rho v_x^2 \} = 0,$$

while the continuity of y - and z -components gives the two conditions

$$\{ \rho v_x v_y \} = 0, \quad \{ \rho v_x v_z \} = 0.$$

Taking care of condition (16.3), let us rewrite the full set of boundary conditions at the discontinuity surface as follows:

$$\begin{aligned} \{ \rho v_x \} &= 0, & \rho v_x \{ \mathbf{v}_\tau \} &= 0, \\ \rho v_x \left\{ \frac{v^2}{2} + w \right\} &= 0, & \{ p + \rho v_x^2 \} &= 0. \end{aligned} \quad (16.5)$$

Here the index τ identifies the tangential components of the velocity.

Obviously the set of equations (16.5) falls into two *mutually exclusive* groups, depending on whether the matter flux across the discontinuity surface is zero or not. Consider these groups.

(a) If

$$v_x = 0$$

then the gas pressure is continuous at the discontinuity surface,

$$\{p\} = 0, \quad (16.6)$$

while the tangential velocity component $\mathbf{v}_\tau$ as well as the density may experience an arbitrary jump:

$$\{\mathbf{v}_\tau\} \neq 0, \quad \{\rho\} \neq 0, \quad \left\{ \frac{v^2}{2} + w \right\} \neq 0.$$

Such discontinuities are called *tangential* (see Landau and Lifshitz 1959a, Chap. 9, Sect. 84).

(b) By contrast, if

$$v_x \neq 0$$

then

$$\{\rho v_x\} = 0, \quad \{\mathbf{v}_\tau\} = 0, \quad \{p + \rho v_x^2\} = 0, \quad \left\{ \frac{v^2}{2} + w \right\} = 0. \quad (16.7)$$

Discontinuities of this type are termed *shock waves*. Their properties are also well known in hydrodynamics (see Exercises 16.1–16.6; see also Landau and Lifshitz 1959a, Chap. 9, Sect. 84).

Therefore

the equations of ideal hydrodynamics allow just two *mutually exclusive* types of discontinuities to exist: the shock wave and the tangential discontinuity.

16.1.3 Dissipative Processes and Entropy

The equations of *ideal* hydrodynamics, as a specific case ($\mathbf{B} = 0$) of the ideal MHD equations (12.70)–(12.75), do not take into account either viscosity or thermal conductivity:

$$\eta = \zeta = 0, \quad \kappa = 0. \quad (16.8)$$

For this reason the ideal hydrodynamics equations describe three conservation laws: conservation of mass, momentum, and entropy. The last one,

$$\frac{\partial s}{\partial t} + (\mathbf{v} \cdot \nabla) s = 0, \quad (16.9)$$

is the specific form of the energy conservation law (see (12.55)) under assumption that the process under consideration is adiabatic. In Sect. 16.1.2 to obtain the boundary conditions at the discontinuity surface we used *conservation of mass, momentum, and energy*, but not entropy. The entropy increases across a shock (Exercise 16.6).

The increase in entropy indicates that *irreversible* dissipative processes (which can be traced to the presence of viscosity and heat conduction in a medium) occur in the shock wave. The model which does not take into account these processes (Sect. 16.1.2) admits the existence of discontinuities but is not capable of describing the continuous transition from the initial to the final state. The ideal hydrodynamics cannot describe either the mechanism of shock compression or the structure of the very thin but finite layer where the plasma undergoes a transition from the initial to the final state.

The entropy increase across the shock is entirely independent of the dissipative mechanism and is defined exclusively by the conservation laws of mass, momentum, and energy

(see Exercise 16.6). Only the thickness of the discontinuity depends upon the rate of the irreversible heating of the plasma compressed by the shock. The following analogy in everyday life is interesting. A glass of hot water will invariably cool from a given temperature (the initial state) to a room temperature (the final state), independently of the mechanism of heat exchange with the surrounding air; the mechanism determines only the rate of cooling.

Recommended Reading: Zel'dovich and Raizer 1966; 2002, v. 1, Chap. 2.


16.2 Magnetohydrodynamic Discontinuities

16.2.1 Boundary Conditions at a Discontinuity Surface

Much like ordinary hydrodynamics, the equations of MHD for an ideal medium (Sect. 12.3) allow discontinuous solutions. de Hoffmann and Teller (1950) were the first to consider shock waves in MHD, based on the relativistic energy-momentum tensor for an ideal medium and the electromagnetic field.

Syrovatskii (1953) has given a general formulation of the problem of the possible types of discontinuity surfaces in a conducting medium with a magnetic field. He has formulated a closed set of equations of ideal MHD and, using this, has written the *boundary conditions* at the discontinuity. We shall briefly reproduce the derivation of the boundary conditions.

Fig. 16.2 The contour L for the derivation of the boundary condition on electric field tangential component



We start from the equations of ideal MHD (12.70)–(12.75). Rewrite them (the equation of state (12.75) is omitted for brevity) as follows:

$$\begin{aligned} \operatorname{div} \mathbf{B} &= 0, & \frac{\partial \mathbf{B}}{\partial t} &= \operatorname{rot}(\mathbf{v} \times \mathbf{B}), & \frac{\partial \rho}{\partial t} &= -\operatorname{div} \rho \mathbf{v}, \\ \frac{\partial}{\partial t} \left(\frac{\rho v^2}{2} + \rho \varepsilon + \frac{B^2}{8\pi} \right) &= -\operatorname{div} \mathbf{G}, & \frac{\partial}{\partial t} (\rho v_\alpha) &= -\frac{\partial}{\partial r_\beta} \Pi_{\alpha\beta}^*. \end{aligned} \quad (16.10)$$

In a frame of reference moving with the discontinuity surface, all the conditions are stationary ($\partial/\partial t = 0$). Hence

$$\operatorname{div} \mathbf{B} = 0, \quad (16.11)$$

$$\operatorname{rot}(\mathbf{v} \times \mathbf{B}) = 0, \quad (16.12)$$

$$\operatorname{div} \rho \mathbf{v} = 0, \quad \operatorname{div} \mathbf{G} = 0, \quad \frac{\partial}{\partial r_\beta} \Pi_{\alpha\beta}^* = 0. \quad (16.13)$$

Four of these conditions have the divergent form and are therefore reduced in the integral form to the conservation of fluxes of vectors appearing at the divergence. Thus the following quantities must conserve at the discontinuity: the perpendicular (to the discontinuity surface) component of the magnetic field vector B_n , the mass flux ρv_n , the energy flux G_n , and the momentum flux $\Pi_{\alpha n}^*$.

The exception is condition (16.12). It is written as the rot of $\mathbf{v} \times \mathbf{B}$. Integration of (16.12) over the area S enclosed by the contour L shown in Fig. 16.2 gives, by virtue of the Stokes theorem,

$$\int_S \operatorname{rot}(\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{S} = \oint_L (\mathbf{v} \times \mathbf{B}) \cdot d\mathbf{l} = 0.$$

Thus condition (16.12) demonstrates the continuity of the tangential component of the vector $(\mathbf{v} \times \mathbf{B})_\tau$, i.e. the electric field $\mathbf{E}_\tau$ in the discontinuity surface.

As in the previous section, the jump of a quantity on crossing the discontinuity surface is designated by curly brackets. The full set of boundary conditions at the surface is written as follows:

$$\{ B_n \} = 0, \quad (16.14)$$

$$\{ (\mathbf{v} \times \mathbf{B})_\tau \} = 0, \quad (16.15)$$

$$\{ \rho v_n \} = 0, \quad (16.16)$$

$$\{ G_n \} = 0, \quad (16.17)$$

$$\{ \Pi_{an}^* \} = 0. \quad (16.18)$$

The physical meaning of the boundary conditions obtained is obvious. The first two are the usual electrodynamic continuity conditions for the normal component of the magnetic field and the tangential component of the electric field. The last three equations represent the continuity of fluxes of mass, energy and momentum, respectively.

As distinct from that in ordinary hydrodynamics (see (16.5)),

in general, the set of the ideal MHD boundary conditions does *not* fall into mutually exclusive groups of equations.

This means that, with a few exceptions, any discontinuity, once accepted by these equations, can, generally speaking, transform to any other discontinuity under continuous change of the conditions of the motion (Syrovatskii 1956).

Hence the classification of discontinuities in MHD seems to be a matter of convention. Any classification is based on the external properties of the flow near the surface, such as the absence or presence of normal components of the velocity v_n and magnetic field B_n , continuity or jump in density. The classification given below is due to Syrovatskii (1953). It is quite convenient for investigating MHD discontinuities.

Before turning our attention to the discussion of the classification mentioned above, let us rewrite the boundary conditions obtained, using (12.76) and (12.77) for the densities of the energy and momentum fluxes and substituting (16.14) in (16.15) and (16.16) in (16.18). We have

$$\{ B_n \} = 0, \quad (16.19)$$

$$\{ v_n \mathbf{B}_\tau \} = B_n \{ \mathbf{v}_\tau \}, \quad (16.20)$$

$$\{ \rho v_n \} = 0, \quad (16.21)$$

$$\left\{ \rho v_n \left(\frac{v^2}{2} + w \right) + \frac{1}{4\pi} (B^2 v_n - (\mathbf{v} \cdot \mathbf{B}) B_n) \right\} = 0, \quad (16.22)$$

$$\left\{ p + \rho v_n^2 + \frac{B^2}{8\pi} \right\} = 0, \quad (16.23)$$

$$\rho v_n \{ \mathbf{v}_\tau \} = \frac{B_n}{4\pi} \{ \mathbf{B}_\tau \}. \quad (16.24)$$

For later use, we write down the boundary conditions in the Cartesian frame of reference, the x axis being perpendicular to the discontinuity surface:

$$\{B_x\} = 0, \quad (16.25)$$

$$\{v_x B_y - v_y B_x\} = 0, \quad (16.26)$$

$$\{v_x B_z - v_z B_x\} = 0, \quad (16.27)$$

$$\{\rho v_x\} = 0, \quad (16.28)$$

$$\left\{ \rho v_x \left(\frac{v^2}{2} + w \right) + \frac{1}{4\pi} (B^2 v_x - (\mathbf{v} \cdot \mathbf{B}) B_x) \right\} = 0, \quad (16.29)$$

$$\left\{ p + \rho v_x^2 + \frac{B^2}{8\pi} \right\} = 0, \quad (16.30)$$

$$\left\{ \rho v_x v_y - \frac{1}{4\pi} B_x B_y \right\} = 0, \quad (16.31)$$

$$\left\{ \rho v_x v_z - \frac{1}{4\pi} B_x B_z \right\} = 0. \quad (16.32)$$

The set consists of *eight* boundary conditions. For $\mathbf{B} = 0$ it converts to the set of four equations (16.5).

Let us consider the classification of discontinuity surfaces in MHD, which stems from the boundary conditions (16.19)–(16.24).

16.2.2 Discontinuities Without Plasma Flows Across Them

Let us suppose the plasma flow through the discontinuity surface is absent:

$$v_n = 0. \quad (16.33)$$

The discontinuity type depends on whether the magnetic field penetrates through the surface or not. Consider both possibilities.


(a) If the perpendicular component of the magnetic field

$$B_n \neq 0, \quad (16.34)$$

then the set of equations (16.19)–(16.24) becomes

$$\begin{aligned} \{B_n\} = 0, \quad B_n \{v_\tau\} = 0, \quad B_n \{\mathbf{B}_\tau\} = 0, \\ \left\{ p + \frac{B^2}{8\pi} \right\} = 0, \quad \{\rho\} \neq 0. \end{aligned} \quad (16.35)$$

Fig. 16.3 Discontinuity surfaces without a plasma flow across them: **(a)** contact discontinuity, **(b)** tangential discontinuity



The velocity, magnetic field strength and (by virtue of the fourth equation) gas pressure are continuous at the surface. The density jump does not have to be zero; otherwise, all values change continuously.

The discontinuity type considered is called the *contact* discontinuity and constitutes just a *boundary between two media*, which moves together with them. It is schematically depicted in Fig. 16.3a.

(b) On the other hand, if

$$B_n = 0 \quad (16.36)$$

then the velocity and magnetic field are parallel to the discontinuity surface (plane $x = 0$). In this case all the boundary conditions (16.19)–(16.24) are satisfied identically, with the exception of one. The remaining equation is

$$\left\{ p + \frac{B^2}{8\pi} \right\} = 0.$$

In other words, the velocity and magnetic field are parallel to the discontinuity surface and may experience arbitrary jumps in magnitude and direction, the only requirement being that the total pressure, that is the sum of the usual gas pressure and the magnetic one, remains continuous at the discontinuity surface:

$$p^* = p + \frac{B^2}{8\pi}. \quad (16.37)$$

Such a discontinuity is called a *tangential* discontinuity (Fig. 16.3b). As treated in MHD, it has a remarkable property. The tangential discontinuity in ordinary hydrodynamics is always unstable (Syrovatskii 1954; see also Landau and Lifshitz 1959a, Chap. 9, Sect. 84, Problem 1). The velocity jump engenders vortices, thus resulting in a turbulence near the discontinuity. Another situation occurs in MHD.




Fig. 16.4 (a) The simplest type of the MHD tangential discontinuities. (b) Formation of a turbulent vortex gives rise to the magnetic field growth

Syrovatskii (1953) has shown that the magnetic field exerts a stabilizing influence on the tangential discontinuity. In particular, if the density ρ_0 and magnetic field B_0 are continuous, the only discontinuous quantity being the tangential velocity component, $v_2 - v_1 = v_0 \neq 0$, then the condition for the tangential discontinuity stability is especially simple:

$$\frac{B_0^2}{8\pi} \geq \frac{1}{4} \frac{\rho_0 v_0^2}{2}.$$

(16.38)

To put it another way, such a discontinuity (Fig. 16.4a) becomes stable with respect to small perturbations (of the general rather than a particular type) once the magnetic energy density reaches one quarter of the kinetic energy density.

The general conclusion concerning the influence of the magnetic field on the stability of hydrodynamic motions of a conducting fluid is as follows:

the magnetic field can only increase the stability of a given velocity distribution as compared to the stability of the same distribution in the absence of a magnetic field.

The point is that any flow instability and turbulence give rise, in view of the freezing-in of the field, to an increase of the magnetic energy (Fig. 16.4b), which is always disadvantageous from the standpoint of the energetic principle of stability.

16.2.3 Perpendicular Shock Wave

Now let

$$v_n \neq 0 \quad \text{and} \quad B_n = 0, \quad (16.39)$$

i.e. a flow through the discontinuity surface is present whereas the magnetic field does not penetrate through the surface. Under these conditions, the following two statements result from (16.19)–(16.24).

(a) From (16.24) the continuity of the tangential velocity component follows:

$$\{\mathbf{v}_\tau\} = 0. \quad (16.40)$$

This makes it possible to transform to such a frame of reference in which the tangential velocity component is absent on either side of the discontinuity:

$$\mathbf{v}_{\tau 1} = \mathbf{v}_{\tau 2} = 0.$$

(b) The tangential electric field continuity (16.20) results in

$$\{v_n \mathbf{B}_\tau\} = 0. \quad (16.41)$$

If the frame of reference is rotated with respect to the x axis in such a way that $B_z = 0$ on one side of the surface, then the same is true on the other side (for clarity see (16.27)). Thus a frame of reference exists in which, in view of (a),

$$\mathbf{v} = (v_n, 0, 0) = (v, 0, 0)$$

and in addition, by virtue of (b),

$$\mathbf{B} = (0, B_\tau, 0) = (0, B, 0).$$

In this frame of reference, the other boundary conditions take the form:

$$\{\rho v\} = 0, \quad (16.42)$$

$$\{B/\rho\} = 0, \quad (16.43)$$

$$\left\{ \rho v^2 + p + \frac{B^2}{8\pi} \right\} = 0, \quad (16.44)$$


$$\left\{ \frac{v^2}{2} + w + \frac{B^2}{4\pi\rho} \right\} = 0. \quad (16.45)$$

Such a discontinuity is called the *perpendicular* shock wave, since it constitutes the compression shock (see (16.7)) propagating perpendicular to the magnetic field as shown Fig. 16.5.

Condition (16.43) reflects the fact of the magnetic field “freezing-in” into the plasma. The role of pressure in such a wave is played by the total pressure

$$p^* = p + \frac{B^2}{8\pi}, \quad (16.46)$$

Fig. 16.5 The character of the plasma motion and magnetic field compression ($B_2 > B_1$) in a perpendicular shock wave



whereas the role of the specific enthalpy is fulfilled by

$$w^* = w + \frac{B^2}{4\pi\rho}. \quad (16.47)$$

Therefore the role of the internal energy density is played by the total internal energy

$$\varepsilon^* = w^* - \frac{p^*}{\rho} = \varepsilon + \frac{B^2}{8\pi\rho} \quad (16.48)$$

(cf. corresponding terms in (12.70) and (12.71)).

For $\mathbf{B} = 0$, the perpendicular shock degenerates to the usual compression shock wave (16.7).

For $\mathbf{B} \neq 0$, the propagation velocity of the perpendicular shock depends on the magnetic field strength.

■ A magnetic field decreases the compressibility of plasma while increasing its elasticity.

This is seen from (16.46) and the freezing-in condition (16.43). Accordingly, the magnetic field increases the shock wave propagation velocity.

If the intensity of a perpendicular shock is diminished, it converts to a fast magnetoacoustic wave propagating across the magnetic field (the angle $\theta = \pi/2$ in Fig. 15.2) with the speed (15.40), i.e.

$$V_{\perp} = \sqrt{V_s^2 + u_A^2}. \quad (16.49)$$

16.2.4 Oblique Shock Waves

The types of discontinuity surfaces treated above are the limiting cases of a more general discontinuity type for which

$$v_n \neq 0 \quad \text{and} \quad B_n \neq 0. \quad (16.50)$$

16.2.4(a) The de Hoffmann-Teller Frame of Reference

In investigating the discontinuities (16.50), a frame of reference would be convenient in which $\mathbf{v}_1$ and $\mathbf{B}_1$ are parallel to each other. Such a frame does exist. It moves with respect to the laboratory one with the velocity

$$\mathbf{U}_{\text{dHT}} = \mathbf{v}_1 - \frac{v_{x1}}{B_{x1}} \mathbf{B}_1 \quad (16.51)$$

parallel to the discontinuity surface. Actually, the scalar product

$$\mathbf{U}_{\text{dHT}} \cdot \mathbf{e}_x = v_{x1} - \frac{v_{x1}}{B_{x1}} B_{x1} = 0.$$

In this frame

$$\mathbf{v}_1(\mathbf{U}_{\text{dHT}}) = \mathbf{v}_1 - \mathbf{U}_{\text{dHT}} = \frac{v_{x1}}{B_{x1}} \mathbf{B}_1$$

and hence

$$\mathbf{v}_1 \times \mathbf{B}_1 = 0. \quad (16.52)$$

Therefore the velocity vector $\mathbf{v}_1$ is parallel to the vector of magnetic field $\mathbf{B}_1$, q.e.d.

Then condition (16.15),

$$\{(\mathbf{v} \times \mathbf{B})_\tau\} = 0,$$

in its coordinate form (16.26) and (16.27) can be used to obtain two equations valid to the right of the discontinuity, i.e. downstream of the shock:

$$v_{x2}B_{y2} - v_{y2}B_{x2} = 0, \quad v_{x2}B_{z2} - v_{z2}B_{x2} = 0.$$

On rewriting these conditions as

$$\frac{v_{x2}}{v_{y2}} = \frac{B_{x2}}{B_{y2}} \quad \text{and} \quad \frac{v_{x2}}{v_{z2}} = \frac{B_{x2}}{B_{z2}},$$

we ensure that

the magnetic field is parallel to the velocity field (in the chosen reference frame) to the right of the discontinuity too.

In such frame of reference, called the *de Hoffmann-Teller frame* (de Hoffmann and Teller 1950), the *electric field does not appear* according to (16.52). More exactly, the tangential electric field equals zero in the de Hoffmann-Teller reference frame.

This fact does not mean, of course, that the local *cross-shock* electric fields do not appear inside the shock transition layer, i.e. inside the discontinuity. The quasi-static electric and magnetic fields may determine the dynamics of particles in the shock front especially if Coulomb collisions play only a minor role. In collisionless shock waves, this dynamics depends on the particular mechanism of the energy redistribution among the perpendicular (with respect to the local magnetic field) and parallel degrees of freedom (Sect. 16.4).

16.2.4(b) Two Types of Shock Waves

Thus $\mathbf{v}$ is parallel to $\mathbf{B}$ on either side of the discontinuity. As a consequence, of the eight boundary conditions (16.25)–(16.32)) initially considered, the two conditions, namely (16.26) and (16.27), have been satisfied by choice of the de Hoffmann-Teller frame of reference. Therefore there remain *six* equations:

$$\{ B_x \} = 0, \quad (16.53)$$

$$\{ \rho v_x \} = 0, \quad (16.54)$$

$$\left\{ \frac{v^2}{2} + w \right\} = 0, \quad (16.55)$$

$$\left\{ p + \rho v_x^2 + \frac{B^2}{8\pi} \right\} = 0, \quad (16.56)$$

$$\left\{ \rho v_x v_y - \frac{B_x B_y}{4\pi} \right\} = 0, \quad (16.57)$$

$$\left\{ \rho v_x v_z - \frac{B_x B_z}{4\pi} \right\} = 0. \quad (16.58)$$

Let us again take account of the parallelism of vectors $\mathbf{v}$ and $\mathbf{B}$ in the chosen reference frame:

$$\mathbf{v}_1 = q_1 \mathbf{B}_1, \quad \mathbf{v}_2 = q_2 \mathbf{B}_2, \quad (16.59)$$

where q_1 and q_2 are some proportionality coefficients. On substituting (16.59) in (16.53)–(16.58) we obtain from three equations (16.54), (16.57) and (16.58) the following three conditions:

$$\{ \rho q \} = 0, \quad (16.60)$$

$$\left\{ \left(1 - \frac{1}{4\pi\rho q^2} \right) v_y \right\} = 0, \quad (16.61)$$

$$\left\{ \left(1 - \frac{1}{4\pi\rho q^2} \right) v_z \right\} = 0. \quad (16.62)$$

These equations admit two essentially different discontinuity types, depending on whether the density of the plasma is continuous or experiences a jump.

First we consider the discontinuity accompanied by a *density jump*:

$$\{\rho\} \neq 0.$$

(16.63)

Discontinuities of this type are called *oblique* shock waves. A discontinuity without a density jump will be considered in Sect. 16.2.7.

Let us rotate the reference frame with respect to the x axis in such a way that

$$v_{z1} = 0.$$

Then from (16.62) the following equation follows:

$$\left(1 - \frac{1}{4\pi\rho_2 q_2^2} \right) v_{z2} = 0. \quad (16.64)$$

This suggests two possibilities: either

(Case I)

$$v_{z2} = 0, \quad (16.65)$$

i.e.

the motion is planar: the velocity and magnetic field are in the plane (x, y) on either side of the discontinuity,

or

(Case II)

$$v_{z2} \neq 0 \quad \text{but} \quad q_2^2 = \frac{1}{4\pi\rho_2}. \quad (16.66)$$

Note that in this case (it will be considered in Sect. 16.2.6)

$$q_1^2 \neq \frac{1}{4\pi\rho_1} \quad (16.67)$$

since concurrently valid equations

$$q_2^2 = \frac{1}{4\pi\rho_2} \quad \text{and} \quad q_1^2 = \frac{1}{4\pi\rho_1}$$

would imply that

$$\rho_2 q_2 = \frac{1}{4\pi q_2} \quad \text{and} \quad \rho_1 q_1 = \frac{1}{4\pi q_1},$$

thus obviously contradicting (16.60) and (16.63). Therefore condition (16.67) must be valid.

We shall consider, of course, both cases indicated above.

Note that we did not use (16.55) and (16.56). These equations determine the relationship between the thermodynamic parameters p and w of a plasma ahead and behind of the shock front.

16.2.4(c) Fast and Slow Shock Waves

Let us consider first the **Case I**. On the strength of (16.65), the boundary conditions (16.53)–(16.58) take the form

$$\begin{aligned} \{B_x\} = 0, \quad \{\rho v_x\} = 0, \quad \left\{\frac{v^2}{2} + w\right\} = 0, \\ \left\{p + \rho v_x^2 + \frac{B_y^2}{8\pi}\right\} = 0, \quad \left\{\rho v_x v_y - \frac{1}{4\pi} B_x B_y\right\} = 0. \end{aligned} \quad (16.68)$$


The oblique shock waves interact with the magnetic field in an intricate way. The relationship between the parameters determining the state of a plasma before and after the wave passage is the topic of a large body of research (see reviews: Syrovatskii 1957; Polovin 1961; monographs: Anderson 1963, Chap. 5; Priest 1982, Chap. 5).

Boundary conditions (16.68) can be rewritten in such a way as to represent the *Rankine–Hugoniot* relation (see Exercises 16.2 and 16.3 for an ordinary shock wave) for shocks in MHD (see Landau et al. 1984, Chap. 8). Moreover the *Zemplen theorem* on the increase of density and pressure in a shock wave can be proved in MHD (Iordanskii 1958; Liubarskii and Polovin 1958; Polovin and Liubarskii 1958; see also Zank 1991). The *fast* and the *slow* oblique shock waves are distinguished.

In the fast shock wave, the magnetic field increases across the shock and is bent towards the shock front surface $x = 0$ as shown in Fig. 16.6. So the magnetic pressure increases as well as the gas pressure:

$$\delta p_m > 0, \quad \delta p > 0. \quad (16.69)$$

Fig. 16.6 The magnetic field change ($B_2 > B_1$), velocity field and plasma density at the front of the fast shock wave



In other words, and this seems to be a natural behavior,

compression of the plasma in a fast MHD shock wave is accompanied by compression of the magnetic field.

In the limiting case of small intensity, the fast shock converts to the fast magnetoacoustic wave (see (16.46)). The speed of the fast shock wave with respect to the medium equals v_{x1} . It is greater than or equal to the speed of the fast magnetoacoustic wave:

$$v_{x1} \geq V_+. \quad (16.70)$$

No small perturbation running in front of the shock can exist upstream of the fast shock wave.

In the slow shock wave, the magnetic field decreases across the shock and is bent towards the shock normal as illustrated by Fig. 16.7. Therefore

$$\delta p_m < 0, \quad \delta p > 0. \quad (16.71)$$

Compression of the plasma is accompanied by a *decrease* of the magnetic field strength in the slow MHD shock wave.


As the amplitude decreases, the slow shock wave will transform to the slow magnetoacoustic wave. The speed of the slow shock propagation is

$$V_- \leq v_{x1} \leq V_A. \quad (16.72)$$

In the particular case

$$B_y = 0 \quad (16.73)$$

Fig. 16.7 The magnetic field change ($B_2 < B_1$), velocity field and plasma density at the front of the slow shock wave (SSW). For comparison, the trans-Alfvénic shock wave (TASW) is shown by vector of magnetic field $\mathbf{B}_2$ (TASW) the tangential component of that changes sign at the shock front



the set of boundary conditions (16.68) results in the set (16.5). This means that the oblique shock wave converts to the parallel (longitudinal) shock wave propagating along the magnetic field, mutual interaction being absent.

16.2.5 Trans-Alfvénic Shock Waves

The set of boundary conditions (16.53)–(16.58) formally admits four other types of discontinuous solutions (Sect. 17.4.2), apart from those indicated above. These four discontinuities are shown in Fig. 17.10 as transitions $\text{II} \rightarrow \text{III}$, $\text{I} \rightarrow \text{III}$, $\text{II} \rightarrow \text{IV}$ and $\text{I} \rightarrow \text{IV}$. They are the so called *trans-Alfvénic* shock waves (TASWs; e.g., Shercliff 1965, Chap. 7) because their velocity is larger than the Alfvén velocity. These shocks are also called *intermediate* shocks.


The distinctive feature of TASWs is that they reverse the magnetic field; see vector $\mathbf{B}_2$ (TASW) in Fig. 16.7. More exactly, the tangential component of the magnetic field changes sign at the shock front. That is why it is tempting to suppose that TASWs are present in the magnetic configurations with the *field reversal* that occur, for example, in the solar wind interaction with planets (Steinolfson and Cable 1993) or during *coronal mass ejections* (CMEs) from the Sun (Steinolfson and Hundhausen 1990).

The peculiarity of these discontinuous solutions is that they have no counterpart among the small amplitude waves or simple waves.

This is the reason why the TASWs are not included in the classification of discontinuities under consideration. What is more important is that the TASWs are *non-evolutionary* (see Sects. 17.1 and 17.4.2).

The **Case II** mentioned above (see conditions (16.66)) shall be considered in the next section.

Fig. 16.8 A switch-on shock: $\mathbf{B}_{\tau 1} = 0$, but $\mathbf{B}_{\tau 2} \neq 0$



16.2.6 Peculiar Shock Waves

We return to the consideration of the oblique shock waves in particular case (16.66) and (16.67):

$$v_{z2} \neq 0, \quad q_1^2 \neq \frac{1}{4\pi\rho_1}, \quad q_2^2 = \frac{1}{4\pi\rho_2}. \quad (16.74)$$

On the strength of (16.61) and (16.62), the following conditions must be satisfied at such a discontinuity:

$$\left(1 - \frac{1}{4\pi\rho_1 q_1^2}\right) v_{y1} = 0, \quad \left(1 - \frac{1}{4\pi\rho_1 q_1^2}\right) v_{z1} = 0.$$

Because the expression in the parentheses is not zero, we get

$$v_{y1} = v_{z1} = 0, \quad (16.75)$$

i.e. in front of such a discontinuity the tangential velocity component $\mathbf{v}_{\tau 1}$ is absent. The tangential field component $\mathbf{B}_{\tau 1}$ is also zero in front of the discontinuity, i.e. the motion follows the pattern seen in the parallel shock wave. However arbitrary tangential components of the velocity and magnetic field are permissible downstream of the shock, the only condition being that

$$\mathbf{v}_2 = \frac{\mathbf{B}_2}{\sqrt{4\pi\rho_2}}. \quad (16.76)$$


Such a discontinuity is called the *switch-on* shock wave. The character of motion of this wave is shown in Fig. 16.8.

The switch-on shock exists in the interval

$$1 < \frac{v_{x1}^2}{V_{Ax1}^2} < \frac{4v_{x1}^2}{v_{x1}^2 + V_{s1}^2}$$

(e.g., Liberman 1978).

Fig. 16.9 A switch-off wave:
 $\mathbf{B}_{\tau 2} = 0$ but $\mathbf{B}_{\tau 1} \neq 0$



Assuming the tangential magnetic field component to be zero to the rear of the peculiar shock wave,

$$\mathbf{B}_{\tau 2} = 0, \quad (16.77)$$

the fluid velocity in front of the discontinuity is the Alfvén one:

$$\mathbf{v}_1 = \frac{\mathbf{B}_1}{\sqrt{4\pi\rho_1}}. \quad (16.78)$$

Such a peculiar shock wave is called the *switch-off* shock wave (Fig. 16.9).

16.2.7 The Alfvén Discontinuity

Returning to the general set of equations (16.53)–(16.58), consider the discontinuity at which the density is constant:

$$\{\rho\} = 0. \quad (16.79)$$

On substituting this condition in (16.54), we see that the normal component of the velocity is continuous at the discontinuity:

$$\{v_x\} = 0.$$


Furthermore, in view of (16.60), the quantity q does not change at the discontinuity:

$$\{q\} = 0.$$

Hence the quantity

$$\left(1 - \frac{1}{4\pi\rho q^2}\right)$$

Fig. 16.10 An Alfvén or rotational discontinuity



is also continuous and may be factored out in (16.61) and (16.62). Rewrite them as follows:

$$\left(1 - \frac{1}{4\pi\rho q^2}\right) \{v_\tau\} = 0. \quad (16.80)$$

If the expression in the parentheses is not zero then the tangential velocity component is continuous and all other quantities are easily checked to be continuous solutions. So, to consider the discontinuous solutions, we put

$$q = \pm \frac{1}{\sqrt{4\pi\rho}}.$$

Thus the velocity vector is connected with the magnetic field strength through the relations

$$\mathbf{v}_1 = \pm \frac{\mathbf{B}_1}{\sqrt{4\pi\rho}}, \quad \mathbf{v}_2 = \pm \frac{\mathbf{B}_2}{\sqrt{4\pi\rho}}. \quad (16.81)$$

The following relations also hold at the discontinuity surface

$$\{p\} = 0, \quad \{\mathbf{B}_\tau^2\} = 0. \quad (16.82)$$

Therefore the normal components and the absolute values of the tangential components of the magnetic field and velocity as well as all thermodynamical parameters conserve at the discontinuity. For given values of $\mathbf{B}_1$ and $\mathbf{v}_1$, possible values of $\mathbf{B}_2$ and $\mathbf{v}_2$ lie on a conical surface, the cone angle being equal to that between the normal to the discontinuity surface and the vector $\mathbf{B}_1$ (Fig. 16.10). A discontinuity of this type is called *Alfvén* or *rotational*.

Its peculiarity is reflected in the second name. On passing the discontinuity surface, a medium can acquire a directionally arbitrary tangential momentum, so that the flow is not generally planar.

The speed of the discontinuity propagation relative to the plasma

$$v_{x1} = \mp \frac{B_{x1}}{\sqrt{4\pi\rho}}. \quad (16.83)$$

In the limiting case of small intensity, the *Alfvén* or *rotational* discontinuity converts to the Alfvén wave (see (15.26)).

16.3 Transitions Between Discontinuities

As was shown by Syrovatskii (1956), *continuous* transitions can occur between *discontinuous* MHD solutions of different types. This statement is easily verified on passing from the discontinuities (Sect. 16.2) to the limit of small-amplitude waves (Sect. 15.1). In this limit the fast and slow magnetoacoustic waves correspond to the oblique shocks, whereas the Alfvén wave corresponds to the Alfvén or rotational discontinuity.

The phase velocity diagrams for the small-amplitude waves are shown in Fig. 15.2. Reasoning from it, the following scheme of continuous transitions between discontinuous solutions in ideal MHD can be suggested (Somov 1993, 1994a).

Let us recall that θ is the angle between the wave vector $\mathbf{k}$ and the magnetic field direction $\mathbf{B}_0/B_0$, i.e. axis x in Fig. 15.2. If $\theta \rightarrow \pi/2$ then the fast magnetoacoustic wave (V_+) converts to the perpendicular wave propagating across the field with the velocity $V_\perp$ (15.40). In the limit of large-amplitude waves this corresponds to the transition from the fast shock (S_+) to the perpendicular one ($S_\perp$) as shown in Fig. 16.11.

If $\theta \rightarrow 0$, the fast magnetoacoustic wave (V_+) converts to the usual sound one (V_s) if $V_s > V_A$ or to the Alfvén wave (V_A) if $V_A > V_s$. Therefore the fast shock (S_+) must convert, when $\theta \rightarrow 0$, either to the longitudinal shock ($S_\parallel$) if $V_{s1} > V_{A1}$ or to the Alfvén discontinuity (A) if $V_{A1} > V_{s1}$.

In much the same way, we conclude, reasoning from Fig. 15.2, that the slow shock (S_-) converts either to the longitudinal shock ($S_\parallel$) for $V_{s1} < V_{A1}$ or to the Alfvén discontinuity (A) for $V_{s1} > V_{A1}$. This transition takes place as $\theta \rightarrow 0$.




Fig. 16.11 A scheme of the continuous transitions between discontinuous solutions in MHD, following from comparison of the properties of the discontinuities and small-amplitude waves on the phase velocity diagram

For $\theta \rightarrow \pi/2$, both the slow shock wave (S_-) and Alfvén discontinuity (A) transform to the tangential discontinuity (T) as demonstrated by the fact that the corresponding phase velocities of the slow magnetoacoustic (V_-) and Alfvén (V_A) waves tend to zero for $\theta \rightarrow \pi/2$.

* * *

How are such transitions realized? —They are effected through some discontinuities which are called *transitional* since they conform to boundary conditions for both types of discontinuities and may be classified as either of the two. The existence of transitional discontinuities means that the discontinuity of one type can convert to the discontinuity of another type under a *continuous* change of parameters (Syrovatskii 1956).

The absence of transitional discontinuities in MHD, manifested as the absence of transitions between small-amplitude waves in the phase velocity diagram (Fig. 15.2), signifies the impossibility similar to that one in ordinary hydrodynamics because there exists a minimal velocity of shock propagation—the sound velocity V_s . That is why small perturbations in hydrodynamics cannot convert the shock wave (S) into the tangential discontinuity (T).

For the same reason the continuous transition between fast (S_+) and slow (S_-) shocks is impossible in MHD. The fast shock (S_+) cannot continuously convert to the perpendicular one ($S_\perp$). These and other restrictions on continuous transitions between discontinuities in MHD will be explained in Chap. 17 from the viewpoint of evolutionarity conditions.

The classical theory of the MHD discontinuous flows is of great utility in analyzing the results of numerical calculations, for example time dependent numerical solutions of the dissipative MHD equations, in order to determine whether the numerical solutions are physically correct (e.g., Falle and Komissarov 2001).

16.4 Shock Waves in Collisionless Plasma

In ordinary collision-dominated gases or plasmas the density rise across a shock wave occurs in a distance of the order of a few collision mean free paths. The velocity distributions on both sides of the front are constrained by collisions to be Maxwellian and, if there is more than one kind of particles (for example, ions and electrons), the temperature of the various constituents of the plasma reach equality. Moreover, as we saw in Sects. 16.1 and 16.2, the conditions (density, pressure, and flow velocity) on one side of the front are rigidly determined in terms of those on the other side by requirement that the flux of mass, momentum, and energy through the front be conserved. For weak shocks the front structure itself can be determined relatively simple, by taking into account collisional transfer coefficients representing viscosity, resistivity, and so on (Sirotna and Syrovatskii 1960; Zel'dovich and Raizer 1966, 2002; see also Sect. 17.4).

In a collisionless plasma the mechanisms by which the plasma state is changed by the passage of the shock front are more complex. Energy and momentum can be transferred from the plasma flow into electric and magnetic field oscillations for example by some *kinetic instabilities*. The energy of these collective motions must be taken into account when conservation laws are applied to relate the pre-shock state to the post-shock state. The ions and electrons are affected differently by instabilities. So there is no reason for their temperatures to remain equal. Since kinetic instabilities are seldom isotropic, it is unlikely that the temperatures will remain isotropic. These anisotropies further change the jump conditions.

The change in state derives from the collective interactions between particles and electric and magnetic fields. In general these fields are of two types. They can be: (a) *constant in time*, more exactly, quasi-static fields produced by charge separation, currents (e.g., [Gedalin and Griv 1999](#)), or (b) *fluctuating in time*, produced by kinetic instabilities. The first situation is usually termed laminar, the second one turbulent. The fields often are turbulent. So the scattering of particles by turbulence can play the role of dissipation in the collisionless shock structure. This turbulence can be either a small-scale one generated by plasma instabilities inside a laminar shock front, or a large-scale turbulence associated with the dominant mode of the shock interaction itself (see [Tidman and Krall 1971](#)).

Since we are discussing the kinetic processes which occur on a time scale much shorter than the time scale of Coulomb collisions, we may efficiently use the Vlasov equation (3.3) or the fluid-type descriptions derived from it ([Chew et al. 1956](#); [Klimontovich et al. 1961](#); [Volkov 1966](#)) to study the properties of shock waves in collisionless plasma.

The high Mach number collisionless shocks are well observed in some astrophysical objects, for example in young supernova remnants (SNRs). It has been suspected for many years that such shocks do not produce the electron-ion temperature equilibration. A clear hint for non-equilibration is the low electron temperature in young SNRs, which in no object seems to exceed 5 keV, whereas a typical shock velocity of $4,000 \text{ km s}^{-1}$ should give rise to a mean plasma temperature of about 20 keV. X-ray observations usually allow only the electron temperature to be determined.

The reflective grating spectrometer on board *XMM-Newton* allowed a direct measurement of an oxygen (O VII) temperature $T_i \approx 500 \text{ keV}$ in SN 1006 ([Vink et al. 2003](#)). Combined with the observed electron temperature $T_e \sim 1.5 \text{ keV}$, this measurement confirms, with a high statistical confidence, that shock heating process resulted in only a small degree ($\sim 3\%$) of electron-ion equilibration at the shock front and that the subsequent equilibration process is slow.

16.5 Practice: Exercises and Answers

Exercise 16.1 (Sect. 16.1.2). Relate the flow variables ρ , v , and p at the surface of an ordinary shock wave.

Answer. From formula (16.7) with $v_\tau = 0$ and $v_x = v$, we find

$$\rho_1 v_1 = \rho_2 v_2, \quad (16.84)$$

$$p_1 + \rho_1 v_1^2 = p_2 + \rho_2 v_2^2, \quad (16.85)$$

$$\frac{v_1^2}{2} + w_1 = \frac{v_2^2}{2} + w_2. \quad (16.86)$$

Here

$$w = \varepsilon + \frac{p}{\rho} \quad (16.87)$$

is the specific enthalpy; the thermodynamic relationship for the specific internal energy $\varepsilon(p, \rho)$ is assumed to be known.

Exercise 16.2 (Sect. 16.1.2). Assuming that the value of a parameter describing the strength of the shock in Exercise 16.1 is known (for example, the relative velocity $\delta v = v_1 - v_2$ which plays the role of the “piston” velocity), find the general relationships that follow from the conservation laws (16.84)–(16.86).

Answer. Instead of the density let us introduce the specific volume $U = 1/\rho$. From (16.84) we obtain

$$\frac{U_2}{U_1} = \frac{v_2}{v_1}. \quad (16.88)$$

Eliminating the velocities v_1 and v_2 from (16.85) and (16.86), we find

$$v_1^2 = U_1^2 \frac{p_2 - p_1}{U_1 - U_2}, \quad (16.89)$$

$$v_2^2 = U_2^2 \frac{p_2 - p_1}{U_1 - U_2}. \quad (16.90)$$

The velocity of the compressed plasma with respect to the undisturbed one

$$\delta v = v_1 - v_2 = [(p_2 - p_1)(U_1 - U_2)]^{1/2}. \quad (16.91)$$


Substituting (16.89) and (16.90) in the energy equation (16.86), we obtain

$$\delta w = w_2 - w_1 = \frac{1}{2} (p_2 - p_1)(U_1 + U_2). \quad (16.92)$$

This is the most general form of the Rankine–Hugoniot relation.

Exercise 16.3 (Sect. 16.1.2). Consider the Rankine–Hugoniot relation for an ideal gas.

Fig. 16.12 The Rankine–Hugoniot curve



Answer. For an ideal gas with constant specific heats c_p and c_v , the specific enthalpy

$$w(p, U) = c_p T = \frac{\gamma_g}{\gamma_g - 1} p U, \quad (16.93)$$

where $\gamma_g = c_p/c_v$ is the specific heat ratio.

If we substitute (16.93) in (16.92), we obtain the Rankine–Hugoniot relation in the explicit form

$$\frac{p_2}{p_1} = \frac{(\gamma_g + 1) U_1 - (\gamma_g - 1) U_2}{(\gamma_g + 1) U_2 - (\gamma_g - 1) U_1}. \quad (16.94)$$

From here, the density ratio

$$r = \frac{\rho_2}{\rho_1} = \frac{U_1}{U_2} = \frac{(\gamma_g + 1) p_2 + (\gamma_g - 1) p_1}{(\gamma_g - 1) p_2 + (\gamma_g + 1) p_1}. \quad (16.95)$$

It is evident from (16.95) that *the density ratio* across a very strong shock, where the pressure p_2 behind the wave front is much higher than the initial pressure p_1 , *does not increase infinitely* with increasing strength p_2/p_1 , but approaches a certain finite value. This limiting density ratio is a function of the specific heat ratio γ_g only, and is equal to

$$r_\infty = \frac{\rho_2}{\rho_1} = \frac{\gamma_g + 1}{\gamma_g - 1}. \quad (16.96)$$

For a monatomic gas with $\gamma_g = 5/3$ the limiting compression ratio $r_\infty = 4$.

A curve on the diagram (p, U) passing through the initial state (p_1, U_1) according to (16.94) is called the Rankine–Hugoniot curve; it is shown in Fig. 16.12.

Exercise 16.4 (Sect. 16.1.2). What is the value of the limiting density ratio r in relativistic shock waves?

Answer. Note that (16.84) is valid only for non-relativistic flows. In relativistic shock waves, the Lorentz factor (5.3) for the upstream and downstream flows must be included, and we have (de Hoffmann and Teller 1950):

$$\gamma_{L,1} \rho_1 v_1 = \gamma_{L,2} \rho_2 v_2. \quad (16.97)$$

The density ratio

$$r = \frac{\rho_2}{\rho_1} = \frac{v_1}{v_2} \frac{\gamma_{L,1}}{\gamma_{L,2}}. \quad (16.98)$$

In highly relativistic shock waves, the ratio v_1/v_2 remains finite, while the density ratio $r \rightarrow \infty$.

This is important fact for particle acceleration by shock waves (see Chap. 18).

Exercise 16.5 (Sect. 16.1.2). Write the density ratio r as a function of the upstream Mach number.

Answer. Let us use the definition of the sound speed (16.1) in an ideal gas with constant specific heats

$$V_s = \left(\gamma_g \frac{p}{\rho} \right)^{1/2} = (\gamma_g p U)^{1/2}. \quad (16.99)$$

The upstream Mach number (to the second power)

$$M_1^2 = \frac{v_1^2}{V_{s1}^2} = \frac{U_1}{\gamma_g p_1} \frac{p_2 - p_1}{V_1 - V_2}. \quad (16.100)$$

Here the solution (16.89) has been taken into account.

Substituting (16.100) in (16.95) gives us the compression ratio as a function of the upstream Mach number

$$r = \frac{(\gamma_g + 1) M_1^2}{(\gamma_g - 1) M_1^2 + 2}. \quad (16.101)$$

When $M_1 \rightarrow \infty$, the density ratio

$$r \rightarrow (\gamma_g + 1)/(\gamma_g - 1)$$

of course. This is the limiting case of a *strong* but non-relativistic shock wave.

When $M_1 \rightarrow 1$, which is the limiting case of a *weak* shock wave, the density ratio $r \rightarrow 1$ too. By using formula (16.94), we see that the pressures on both sides of a weak shock wave are close to each other: $p_1 \approx p_2$ and $(p_2 - p_1)/p_1 \ll 1$. Thus a weak shock wave is practically the same as an acoustic compression wave.

For $M_1 < 1$ we could formally have an expansion shock wave with $r < 1$ and $p_2 < p_1$. However it can be shown (see the next Exercise) that such a transition would involve a *decrease* of entropy rather than an increase. So such transitions are ruled out by the second law of thermodynamics.

Exercise 16.6 (Sect. 16.1.3). Show that

the entropy jump of a gas compressed by a shock increases with the strength of the shock wave but is entirely independent of the dissipative mechanism.

Answer. To within an arbitrary constant the entropy of an ideal gas with constant specific heats is given by formula (see Landau and Lifshitz 1959b, Chap. 4):

$$s = c_v \ln p U^{\gamma_g}. \quad (16.102)$$

The difference between the entropy on each side of the shock front, as derived from (16.95), is

$$s_2 - s_1 = c_v \ln \left\{ \left(\frac{p_2}{p_1} \right) \left[\frac{(\gamma_g - 1)(p_2/p_1) + (\gamma_g + 1)}{(\gamma_g + 1)(p_2/p_1) + (\gamma_g - 1)} \right]^{\gamma_g} \right\}. \quad (16.103)$$

In the limiting case of a weak wave ($p_2 \approx p_1$) the expression in braces is close to unity. Therefore $s_2 \approx s_1$ and $s_2 > s_1$ if $p_2 > p_1$.

As the strength of the wave increases, that is, as the ratio p_2/p_1 increases beyond unity, the expression in braces increases monotonically and approaches infinity as $p_2/p_1 \rightarrow \infty$. Thus the entropy jump is positive and does increase with the strength of the shock wave.

The increase in entropy indicates that irreversible dissipative processes occur in the shock front.

This can be traced to the presence of viscosity and heat conduction in the gas or plasma (see the discussion in Sect. 16.1.3).


Exercise 16.7 (Sect. 16.1.3). Consider and discuss Fig. 16.13 which illustrates character of a solution of hydrodynamic problem on the structure of a strong shock wave with electron heat conduction in a fully ionized hydrogen plasma.

Comment. Consider a stationary one-dimensional flow of a plasma in the *single-fluid two-temperature* approximation (see Sect. 8.3.2) taking account of ion viscosity in the ion-entropy equation and electron heat conduction in the electron-entropy equation, as well as the collisional exchange of energy between electrons and ions in both equations. Thus we obtain the following set of ordinary differential equations (Zel'dovich and Raizer 1966, 2002, Chap. 7):

$$\frac{d}{dx} \rho v = 0, \quad (16.104)$$

$$\rho v \frac{dv}{dx} = -\frac{dp}{dx} + \frac{d}{dx} \left(\frac{4}{3} \eta_p \frac{dv}{dx} \right), \quad (16.105)$$

Fig. 16.13 The behavior of:
(a) temperatures of electrons
 and ions **(b)** plasma density in
 a strong shock wave in an
 ionized hydrogen plasma



$$\rho v T_p \frac{ds_p}{dx} = \frac{4}{3} \eta_p \left(\frac{dv}{dx} \right)^2 + Q, \quad (16.106)$$

$$\rho v T_e \frac{ds_e}{dx} = -\frac{d}{dx} q_e - Q. \quad (16.107)$$

Here T_p and T_e are the proton and electron temperatures respectively, s_p and s_e are the specific entropies of proton and electron components in plasma, η_p is the proton viscosity (see Exercise 10.1), q_e is the heat flux in the electron component in plasma (9.50), Q is the rate of thermal energy exchange between protons and electrons due to Coulomb collisions (9.43).

Exercise 16.8. Consider a collisionless gravitational system described by the gravitational analog of the Vlasov equation (Sect. 3.3.1). Explain qualitatively why the Vlasov equation (3.33) does not predict the existence of a shock wave. In other words, unlike the case of gas or plasma, an evolution governed by the set of equations (3.33)–(3.35) never leads to caustics or shocks.

Hint. By analogy with the discussion of the shock origin in ordinary hydrodynamics (Sect. 16.1.1), it is necessary to show that

given sufficiently smooth initial data, the distribution function of a collisionless gravitational system will never diverge.

So the gravitational analog of the Vlasov equation manifests the so-called “global existence” (Pfaffelmoser 1992).

Chapter 17

Evolutionarity of MHD Discontinuities

Abstract A discontinuity cannot exist in astrophysical plasma with magnetic field if small perturbations disintegrate it into other discontinuities or transform it to a more general non-steady flow. In this chapter we consider the so-called evolutionarity or structural stability of the classic discontinuous solution of the MHD equations.

17.1 Conditions for Evolutionarity

17.1.1 The Physical Meaning and Definition

Of concern to us is the issue of the stability of MHD discontinuities with respect to their decomposition into more than one discontinuity. To answer this question small perturbations must be imposed on the discontinuity surface. If they do not instantaneously lead to large changes of the discontinuity, then the discontinuity is termed *evolutionary*.


Obviously the property of evolutionarity does not coincide with stability in the ordinary sense. The usual instability means exponential ($e^{\gamma t}$, $\gamma > 0$) growth of the disturbance, it remains small for some time ($t \leq \gamma^{-1}$). The discontinuity gradually evolves. By contrast,

┌ a disturbance *instantaneously* becomes large in a non-evolutionary discontinuity.

By way of illustration, the decomposition of a density jump $\rho(x)$ is shown in Fig. 17.1. The disturbance $\delta\rho$ is not small, though it occupies an interval δx which is small for small t , when the two discontinuities have not become widely separated.

The problem of disintegration of discontinuities has a long history. Kotchine (1926) considered the disintegration of an arbitrary discontinuity into a set of other discontinuities and rarefaction waves in the frame of ordinary hydrodynamics.

Fig. 17.1 Disintegration of a density jump into two successive jumps



Bethe (1942) studied the disintegration of a shock wave. The mathematical idea of evolutionarity was expressed for the first time in the context of the study of discontinuities in hydrodynamics (Courant and Friedrichs 1985; see also Gel'fand 1959).

With respect to evolutionary discontinuities, the usual problem of linear stability can be formulated,

i.e. we find solutions to the linearized equations giving rise to small amplitudes which grow or decay in time.

The *evolutionarity criterion* may be obtained by counting the number of equations supplied by linearized boundary conditions at the discontinuity surface under consideration, and the number of independent parameters determining an arbitrary, initially small disturbance of the discontinuity. If the numbers are equal, then the boundary conditions uniquely define further development—evolution—of the disturbance which remains small for small $t > 0$. Such a discontinuity is *evolutionary*. By contrast, if the number of parameters is greater or less than the number of independent equations, then the problem of a small perturbation of the discontinuity has an infinitely large number of solutions or no solutions at all. Thus

the initial assumption of the smallness of the disturbance for small t is incorrect, hence the discontinuity is *non-evolutionary*.

Such a discontinuity cannot exist as a stationary configuration because a small perturbation leads to a finite variation of the initial flow. This variation is the disintegration of the discontinuity into other discontinuities that move away from the place of their formation (Fig. 17.1), or a transition to a more general non-steady flow.

Let us count the number of equations which must be satisfied by an arbitrary small perturbation at an *ideal MHD discontinuity*. Thus we take as the initial conditions the set of eight boundary conditions (16.25)–(16.32). It is to be linearized.

Now we consider perturbations of the discontinuity, which generate plane waves propagating along the x axis. Then the quantity B_x remains constant on either side of the discontinuity, and condition (16.25) (both exact and linearized) is satisfied

identically. Hence, on either side of the discontinuity, seven quantities are perturbed: three velocity components (v_x, v_y, v_z), two field components (B_y, B_z), density ρ and pressure p . Small perturbations of these quantities,

$$\delta v_x, \delta v_y, \delta v_z, \delta B_y, \delta B_z, \delta \rho, \delta p,$$

on either side of the discontinuity surface are characterized by the coordinate and time dependence

$$\delta f(t, x) \sim \exp[i(kx - \omega t)]$$

typical of the plane wave.

If the number of waves leaving the discontinuity is equal to the number of boundary conditions, then the problem of small perturbations has only one solution and the discontinuity is evolutionary.

This form of evolutionarity conditions has been obtained for the first time by Lax (1957, 2006). The small perturbations must obey the linearized boundary conditions, i.e. linear algebraic equations following from (16.26)–(16.32). In addition to the seven quantities mentioned above, the velocity of propagation of the discontinuity surface is disturbed. It acquires a small increment δu_x relative to the chosen frame of reference in which the undisturbed discontinuity is at rest.

17.1.2 Linearized Boundary Conditions

Let us write down the linearized boundary conditions in a reference frame rotated with respect to the x axis in such a way that the undisturbed values $B_z = 0$ and $v_z = 0$. Thus we restrict our consideration to those discontinuity surfaces in which the undisturbed fields $\mathbf{B}_1, \mathbf{B}_2$ and the velocities $\mathbf{v}_1, \mathbf{v}_2$ lie in the plane (x, y) .

From the boundary conditions (16.25)–(16.32) we find a set of linear equations which falls into *two groups* describing different perturbations:

(a) Alfvén perturbations ($\delta v_z, \delta B_z$)

$$\left\{ \rho v_x \delta v_z - \frac{1}{4\pi} B_x \delta B_z \right\} = 0, \quad (17.1)$$

$$\{ v_x \delta B_z - B_x \delta v_z \} = 0; \quad (17.2)$$

(b) magnetoacoustic and entropy perturbations ($\delta v_x, \delta v_y, \delta B_y, \delta \rho, \delta p$)

$$\{ \rho (\delta v_x - \delta u_x) + v_x \delta \rho \} = 0, \quad (17.3)$$

$$\left\{ \rho v_x \delta v_y + v_y [\rho (\delta v_x - \delta u_x) + v_x \delta \rho] - \frac{1}{4\pi} B_x \delta B_y \right\} = 0, \quad (17.4)$$

$$\left\{ \delta p + v_x^2 \delta \rho + 2\rho v_x (\delta v_x - \delta u_x) + \frac{1}{4\pi} B_y \delta B_y \right\} = 0, \quad (17.5)$$

$$\{ B_x \delta v_y - B_y (\delta v_x - \delta u_x) - v_x \delta B_y \} = 0, \quad (17.6)$$

$$\begin{aligned} & \left\{ \rho v_x [v_x (\delta v_x - \delta u_x) + v_y \delta v_y + \delta w] + \right. \\ & + \left(\frac{v_x^2 + v_y^2}{2} + w \right) [\rho (\delta v_x - \delta u_x) + v_x \delta \rho] + \\ & + \frac{B_y}{4\pi} [B_y (\delta v_x - \delta u_x) + v_x \delta B_y - B_x \delta v_y] + \\ & \left. + \frac{1}{4\pi} (v_x B_y - v_y B_x) \delta B_y \right\} = 0. \end{aligned} \quad (17.7)$$

Condition (17.3) allows us to express the disturbance of the propagation velocity of the discontinuity surface δu_x in terms of perturbations of ρ and v_x :

$$\delta u_x \{ \rho \} = \{ \rho \delta v_x + v_x \delta \rho \}. \quad (17.8)$$

On substituting (17.8) in (17.4)–(17.7) there remain four independent equations in the second group of boundary conditions, since the disturbance of the velocity of the discontinuity surface δu_x can be eliminated from the set.

Therefore the MHD boundary conditions for perturbations of the discontinuity, which generate waves propagating perpendicular to the discontinuity surface, fall into two *isolated* groups. As this takes place,

the conditions of evolutionarity (the number of waves leaving the MHD discontinuity is equal to the number of independent linearized boundary conditions) must hold not only for the variables in total but also for *each* isolated group

(Syrovatskii 1959). The number of Alfvén waves leaving the discontinuity must be two, whereas there must be four magnetoacoustic and entropy waves. This makes the evolutionary requirement more stringent.

Whether or not a discontinuity is evolutionary is clearly a *purely kinematic* problem. We have to count the number of small-amplitude waves leaving the discontinuity on either side. Concerning the boundary conditions the following comment should be made. As distinct from the unperturbed MHD equations, the perturbed ones are not stationary. Therefore the arguments used to derive (16.19)–(16.24) from (16.10) are not always valid.

To derive boundary conditions at a disturbed discontinuity we have to transform them to the reference frame connected with the surface. For example, for a perturbation (see Exercise 17.2)

$$\xi_x(y, t) = \xi_0 \exp [i (k_y y - \omega t)],$$

where ξ_x is a displacement of the surface, this is equivalent to the following substitution in the linearized MHD equations

$$\frac{\partial}{\partial t} \delta \rightarrow -i\omega \left(\delta - \xi_0 \frac{\partial}{\partial y} \right), \quad \frac{\partial}{\partial y} \delta \rightarrow ik_y \left(\delta - \xi_0 \frac{\partial}{\partial y} \right),$$

where $-i\omega \xi_0 = \delta u_x$ is the amplitude of the time derivative of ξ . Consider, for example, the linearized continuity equation which after the integration over the discontinuity thickness takes the form

$$\begin{aligned} i \int_{-a}^{+a} (\omega - k_y v_y) \delta \rho dx - ik_y \int_{-a}^{+a} \rho \delta v_y dx &= \\ &= \{ v_x \delta \rho + \rho [\delta v_x + i(\omega - k_y v_y) \xi_0] \}. \end{aligned} \quad (17.9)$$

If the integrals on the left-hand side of (17.9) are equal to zero in the limit $a \rightarrow 0$ then, for $k_y = 0$, formula (17.9) transforms to (17.3). However this possibility is based on the supposition that $\delta \rho$ and δv_y inside the discontinuity do not increase in the limit $a \rightarrow 0$. We shall see in Part II, Chap. 10 that this supposition is not valid at least for more complicated, two-dimensional, configurations such as a *reconnecting current layer*.

17.1.3 The Number of Small-Amplitude Waves

If the discontinuity is immovable with respect to the plasma (no flow across the discontinuity), then *on either side* of the surface of discontinuity there exist three waves leaving it as shown in Fig. 17.2:

$$-V_{+x1}, -V_{Ax1}, -V_{-x1}, V_{-x2}, V_{Ax2}, V_{+x2}. \quad (17.10)$$

Let the discontinuity move with a velocity v_{x1} relative to the plasma (see Fig. 17.3). The positive direction of the axis x is chosen to coincide with the direction of the plasma motion at the discontinuity surface. The index “1” refers to the region in front of the surface ($x < 0$) whereas the index “2” refers to the region behind the discontinuity ($x > 0$), i.e. downstream of the flow. Then there exist fourteen different phase velocities of propagation of small-amplitude waves:

$$\begin{aligned} v_{x1} \pm V_{+x1}, \quad v_{x1} \pm V_{Ax1}, \quad v_{x1} \pm V_{-x1}, \quad v_{x1}, \\ v_{x2} \pm V_{-x2}, \quad v_{x2} \pm V_{Ax2}, \quad v_{x2} \pm V_{+x2}, \quad v_{x2}. \end{aligned}$$

Waves leaving the discontinuity have negative phase velocities in the region 1 and positive phase velocities in the region 2.





Fig. 17.2 Six small-amplitude waves leaving an immovable discontinuity surface ($x = 0$) being perturbed

Fig. 17.3 Small-amplitude waves in a plasma moving through the MHD discontinuity



In the region 1, four velocities, corresponding to the waves moving toward the discontinuity surface, can be immediately discarded:

$$v_{x1} + V_{+x1}, \quad v_{x1} + V_{Ax1}, \quad v_{x1} + V_{-x1}, \quad v_{x1}.$$

The remaining three waves ($7 - 4$) can leave the discontinuity or propagate toward it, depending on the plasma flow velocity towards the discontinuity v_{x1} .

In the region 2, four waves always have positive phase velocities:

$$v_{x2} + V_{+x2}, \quad v_{x2} + V_{Ax2}, \quad v_{x2} + V_{-x2}, \quad v_{x2}. \quad (17.11)$$

These waves leave the discontinuity. Other waves will be converging or diverging, depending on relations between the quantities

$$v_{x2}, \quad V_{+x2}, \quad V_{Ax2}, \quad V_{-x2}.$$

v_{x2}	$3 + 7 = 10$	$2 + 7 = 9$	$1 + 7 = 8$	$0 + 7 = 7$
V_{+x2}	$3 + 6 = 9$	$2 + 6 = 8$	$1 + 6 = 7$	$0 + 6 = 6$
V_{Ax2}	$3 + 5 = 8$	$2 + 5 = 7$	$1 + 5 = 6$	$0 + 5 = 5$
V_{-x2}	$3 + 4 = 7$	$2 + 4 = 6$	$1 + 4 = 5$	$0 + 4 = 4$
	0	V_{-x1}	V_{Ax1}	V_{+x1}
				v_{x1}

Fig. 17.4 The number of small-amplitude waves leaving a discontinuity surface

Let

$$0 < v_{x1} < V_{-x1} . \quad (17.12)$$

Then there are three waves leaving the discontinuity in the region 1:

$$v_{x1} - V_{-x1}, \quad v_{x1} - V_{Ax1}, \quad v_{x1} - V_{-x1}.$$

If

$$0 < v_{x2} < V_{-x2} , \quad (17.13)$$

then four waves (17.11) propagate downstream of the discontinuity since the waves

$$v_{x2} - V_{-x2}, \quad v_{x2} - V_{Ax2}, \quad v_{x2} - V_{+x2}$$

converge to the discontinuity.

We shall write down the number of diverging waves to the left (in front of) and to the right (behind) the discontinuity as their sum (e.g. $3 + 4 = 7$ in the case considered) in the corresponding rectangle in the plane (v_{x1}, v_{x2}) presented in Fig. 17.4. This rectangle is the lower left one. In the rectangle situated to the right of this one, two rather than three waves are diverging in the region 1:

$$v_{x1} - V_{Ax1}, \quad v_{x1} - V_{+x1} .$$

The wave $v_{x1} - V_{-x1}$ is carried by the flow to the discontinuity since

$$V_{-x1} < v_{x1} < V_{Ax1} .$$

(17.14)

Thus we write $2 + 4 = 6$ in this rectangle. The whole table is filled up in a similar manner.




Fig. 17.5 The evolutionarity domains for the fast (S_+) and slow (S_-) shocks and the Alfvén discontinuity (the point A)

17.1.4 Domains of Evolutionarity

If one considers the total number of boundary conditions (six), without allowance being made for their falling into two groups, then just three rectangles in Fig. 17.4 should be inspected for a possible evolutionarity. The boundaries of these rectangles are shown by solid lines.

However, as indicated above, the equality of the total number of independent boundary conditions to the number of outgoing waves is insufficient for the existence and uniqueness of the solutions in the class of small perturbations (Syrovatskii 1959). Take into account that

the linearized boundary conditions fall into two groups with two equations in the first group; for this reason the number of Alfvén waves must equal two and that of outgoing magnetoacoustic and entropy waves must equal four.

Then one of the three rectangles (that one for which the number of outgoing waves $1 + 5 = 6$) becomes the point A in Fig. 17.5.

The figure also shows that there exist only two domains of evolutionarity of shock waves:

(a) *fast shock waves* (S_+) for which

$$v_{x1} > V_{+x1}, \quad V_{Ax2} < v_{x2} < V_{+x2}, \quad (17.15)$$

(b) *slow shock waves* (S_-) for which

$$V_{-x1} < v_{x1} < V_{Ax1}, \quad v_{x2} < V_{-x2}. \quad (17.16)$$

Therefore,

in the ideal MHD approximation, the fast and slow oblique shock waves are evolutionary.

Satisfaction of the boundary conditions at such a shock uniquely determines the amplitudes of the outgoing waves, for example, in terms of the amplitudes of the incoming waves if they are a cause of perturbations of the shock front.

In contrast with this, for the *trans-Alfvénic shock waves* (TASWs) introduced in Sect. 16.2.5 (d), the boundary conditions overdetermine the amplitudes of outgoing waves in the remaining three rectangles: $1 + 4 = 5$, $0 + 5 = 5$ and $0 + 4 = 4$. Thus there is no solution at all. Therefore,

the trans-Alfvénic shock waves (TASWs) are non-evolutionary in the ideal MHD approximation.

The last fact will be also justified from a viewpoint of structure of such discontinuities (see Sect. 17.4.2).

Recall that our treatment of the Alfvén discontinuity was not quite satisfactory. It was treated as a flow in the plane (x, y) . Generally this is not the case (Fig. 16.10). The result of the above analysis is also not quite satisfactory: the evolutionarity of the Alfvén discontinuity, as well as the *switch-on* and *switch-off* shocks, is more complicated. While investigating the evolutionarity of these discontinuities, dissipative effects must be allowed for (Sect. 17.3).

Although *dissipative* waves quickly damp as they propagate away from the discontinuity surface, they play an important role in the system of small-amplitude waves leaving the discontinuity. Thus only one solution exists for the switch-off shock, i.e. it is evolutionary. By contrast,

the switch-on shock wave, as well as the Alfvén or rotational discontinuity, are non-evolutionary

in the linear approximation.

Roikhvarger and Syrovatskii (1974) have shown that attention to dissipation in the dispersion equation for magnetoacoustic and entropy waves leads to the appearance of *dissipative waves* and, as a consequence, to the non-evolutionarity of tangential and contact discontinuities (Sect. 17.3).

Recall that in an ideal medium the disintegration of a discontinuity is instantaneous in the sense that the secondary discontinuities become separated in the beginning of the disintegration process (Fig. 17.1). In a dissipative medium the spatial profiles of the MHD discontinuities are continuous. Nevertheless the principal result remains the same. The steady flow is rearranged toward a non-steady state, and after a large enough period of time the disintegration manifests itself (Sect. 17.4).

17.2 Consequences of Evolutionarity Conditions

17.2.1 The Order of Wave Propagation

Some interesting inferences concerning the order of shock propagation result from the evolutionarity conditions (17.15) and (17.16).

If a shock wave follows another one of the same type (fast or slow), the back shock will catch up with the front one (Akhiezer et al. 1959). Let us consider, as an example, two slow shock waves, S_-^A and S_-^B , propagating in the direction of the x axis as shown in Fig. 17.6.

In a reference frame connected with the front of the first shock S_-^A , we get, by virtue of the evolutionary condition (17.16),

$$V_{-x1}^A < v_{x1}^A < V_{Ax1}^A, \quad v_{x2}^A < V_{-x2}^A. \quad (17.17)$$

In a reference frame connected with the front of the second shock S_-^B , analogous conditions hold:

$$V_{-x1}^B < v_{x1}^B < V_{Ax1}^B, \quad v_{x2}^B < V_{-x2}^B. \quad (17.18)$$

Since the velocities of slow magnetoacoustic waves of small amplitude V_{-x2}^A and V_{-x1}^B refer to the same region (between the shocks), they are equal

$$V_{-x2}^A = V_{-x1}^B. \quad (17.19)$$

Substituting (17.19) in the second part of (17.17) and in the first part of (17.18) gives the inequality




Fig. 17.6 Plasma flow velocities relative to: (a) shock wave fronts, (b) the plasma between the shock waves

$$v_{x2}^A < v_{x1}^B. \quad (17.20)$$

Hence, relative to the plasma between the shocks (Fig. 17.6b), the shock S_-^B catches up with the shock S_-^A , which was to be proved.

As for different types of waves, the following inferences can be drawn: the Alfvén discontinuity will catch up with the slow shock, whereas the fast shock will catch up with all possible types of discontinuities. If shock waves are generated by a single source (for example, a flare in the solar atmosphere), then no more than three shocks can move in the same direction: the fast shock is followed by the Alfvén discontinuity, the slow shock being to the rear of the Alfvén discontinuity.

17.2.2 Continuous Transitions Between Discontinuities

Reasoning from the polar diagram for phase velocities of small-amplitude waves, in Sect. 16.3 we have treated the possibility of continuous transitions between different types of discontinuous solutions in MHD. However the evolutionarity conditions have not been taken into account. They are known to impose limitations on possible continuous transitions between the discontinuities under changes of external parameters (magnetic field, flow velocity, etc.).

Continuous transition is impossible between the fast and slow shock waves. This stems from the fact that the evolutionarity domains for fast (S_+) and slow (S_-) shocks have no common points (Fig. 17.5). Similarly, the lines of phase velocities V_+ and V_- in polar diagrams (Figs. 15.2 and 15.3) are out of contact. That was the basis for banning transitions between the fast and slow shocks in Fig. 16.11.

The fast shock (S_+) cannot continuously convert to the tangential discontinuity (T) since that would go against the evolutionarity condition $v_{x1} > V_{Ax1}$. The same ban stems from the consideration of the phase velocity diagram. The perpendicular shock ($S_\perp$) is the limiting case of the fast shock. That is why the continuous transition of the perpendicular shock to the tangential discontinuity is forbidden, as shown in Fig. 16.11.

As was indicated in the previous section, the issue of evolutionarity of the Alfvén discontinuity has no satisfactory solution in the framework of ideal MHD. The established viewpoint is that the continuous transition of shock waves (S_- and S_+) to the Alfvén discontinuity (A) is impossible, as is predicted by the phase velocity diagram with $\theta \rightarrow 0$. Transitions between the Alfvén (A) and tangential (T) discontinuities, between the tangential discontinuity and the slow shock (S_-), between the tangential and contact (C) discontinuities are assumed to be possible. These discontinuities convert to the tangential discontinuity in the limiting case $B_x \rightarrow 0$ (Polovin 1961; Akhiezer et al. 1975).

We shall consider the evolutionarity conditions and their consequences for reconnecting current layers (RCLs) as a MHD discontinuity in Part II, Chap. 10.

17.3 Dissipative Effects in Evolutionarity

17.3.1 Dissipative and Non-dissipative Waves

Roikhvarger and Syrovatskii (1974) have taken into account the effect of dissipation on evolutionarity of the peculiar shocks. In this case the dispersion relation of the Alfvén waves has the following general form:

$$k^2 V_{Ax}^2 - (\omega - k v_x - i k^2 \nu_m) (\omega - k v_x - i k^2 \nu) = 0. \quad (17.21)$$

Here the vector $\mathbf{k}$ is directed along the x axis, ν_m is the magnetic diffusivity, and $\nu = \eta/\rho$ is the kinematic viscosity.

After expansion of the solutions of this equation in powers of a small ω (the conditions under which ω is small will be discussed below) the expression for k reads as follows:

(a) for $v_x = V_{Ax}$

$$k^d = \pm \sqrt{\frac{\omega}{\nu_m + \nu}} (1 - i), \quad (17.22)$$

$$k^A = \frac{\omega}{2v_x} - i \frac{(\nu_m + \nu) \omega^2}{16 v_x^3}, \quad (17.23)$$

$$k^* = -\frac{\omega (\nu_m^2 + \nu^2)}{v_x (\nu_m + \nu)^2} + i \frac{v_x (\nu_m + \nu)}{\nu_m \nu}; \quad (17.24)$$

(b) for $v_x \neq V_{Ax}$

$$k^A = \frac{\omega}{v_x \pm V_{Ax}} - i \frac{(\nu_m + \nu) \omega^2}{2 (v_x \pm V_{Ax})^3}, \quad (17.25)$$

$$k^* = -\frac{\omega [(\nu_m - \nu)^2 v_x \pm (\nu_m + \nu) K]}{4 V_{Ax}^2 \nu_m \nu + v_x^2 (\nu_m - \nu)^2 \pm v_x (\nu_m + \nu) K} + i \frac{v_x (\nu_m + \nu) \pm K}{2 \nu_m \nu}, \quad (17.26)$$

where

$$K = \left[v_x^2 (\nu_m - \nu)^2 + 4 V_{Ax}^2 \nu_m \nu \right]^2.$$

Thus

the dissipative effects result in additional small-amplitude waves propagating in a homogeneous MHD medium.

The width of an MHD shock (at least of small amplitude) is proportional, in order of magnitude, to the dissipative transport coefficients and inversely proportional to the shock intensity (Sirotnina and Syrovatskii 1960). The intensity is determined by the difference $v_x - V_{Ax}$ on the side of the discontinuity on which it is not zero. Since the *switch-off* shock, as a slow one, has a finite intensity, and the *switch-on* shock exists in the interval (see Sect. 16.2.6)

$$1 < \frac{v_{x1}^2}{V_{Ax1}^2} < \frac{4v_{x1}^2}{v_{x1}^2 + V_{s1}^2},$$

the width of the peculiar shock can be estimated as

$$l \sim \frac{v_m + v}{|v_x - V_{Ax}|} \quad (17.27)$$

(Roikhvarger and Syrovatskii 1974). It is just this distance within which the perturbations k^* from (17.24) and (17.26) damp considerably. Therefore outside the shock front these waves are absent, and their amplitudes do not enter into the boundary conditions which relate perturbations outside the shock front.

The situation is different for the remaining perturbations, in particular, for the *purely dissipative* waves k^d from (17.22). For small enough ω their wave numbers are much larger than the thickness l of the shock. This is true under the condition

$$\omega \ll \frac{(v_x - V_{Ax})^2}{v_m + v}, \quad (17.28)$$

which coincides with that used to derive formulae (17.22)–(17.26). Since the characteristic length scale of such perturbations is much larger than the shock thickness l , their amplitudes satisfy the boundary conditions at the discontinuity surface (17.1) and (17.2) obtained for an ideal medium.

17.3.2 Dissipation and Evolutionarity

The classification of dissipative perturbations on incoming and outgoing waves should be made according to the sign of the imaginary part of the wave vector, because in a stable medium such waves damp in the direction of the propagation (Sect. 15.3). Consequently, there are two outgoing perturbations leaving the peculiar shock, one of them being the dissipative wave. Much like the case of non-peculiar shocks, both waves propagate downstream away from the (fast) switch-on shock, while there is one outgoing wave on each side of the (slow) switch-off shock (Fig. 17.7).




Fig. 17.7 The direction of the wave propagation in the case of a switch-on shock (a) and a switch-off shock (b)

With the precision adopted when deriving formulae (17.22)–(17.26), the perturbations δv_z and δB_z in the dissipative wave k^* from (17.22) are related by the formula

$$\delta v_z^d = \left(1 \pm \frac{v_m - v}{v_x} \sqrt{\frac{i\omega}{2(v_m + v)}} \right) \frac{\delta B_z^d}{\sqrt{4\pi\rho}}. \quad (17.29)$$

From here and boundary conditions (17.1) and (17.2), it follows that if an Alfvén wave is incident onto the switch-off shock from upstream or downstream then the amplitude of the dissipative wave equals respectively

$$\delta B_{z1}^d = -\frac{2v_{x1}}{v_m - v} \sqrt{\frac{2(v_m + v)}{i\omega}} \delta B_{z1}^\downarrow, \quad (17.30)$$

or

$$\delta B_{z1}^d = -\frac{2v_{x1}}{v_m - v} \sqrt{\frac{2(v_m + v)}{i\omega}} \delta B_{z2}^\uparrow. \quad (17.31)$$

The amplitude $\delta B_{z2}^\downarrow$ of the traveling (non-dissipative) wave equals zero in the first case and $-\delta B_{z2}^\uparrow$ in the second case. Thus only one solution exists for the switch-off shock. Consequently, the *switch-off shock is evolutionary*.

On the contrary, the switch-on shock is non-evolutionary. Indeed, (17.1) and (17.2), with regard for the relation at the switch-on shock

$$v_{x1} v_{x2} = V_{Ax1}^2 \quad \text{and} \quad \frac{\rho_2}{\rho_1} = \frac{v_{x1}^2}{V_{Ax1}^2}, \quad (17.32)$$

can be rewritten as

$$v_{x1} \left(\delta v_{z2} - \frac{\delta B_{z2}}{\sqrt{4\pi\rho}} \right) = v_{x1} \delta v_{z1} - V_{Ax1} \frac{\delta B_{z1}}{\sqrt{4\pi\rho}}, \quad (17.33)$$

$$V_{Ax1} \left(\delta v_{z2} - \frac{\delta B_{z2}}{\sqrt{4\pi\rho}} \right) = V_{Ax1} \delta v_{z1} - v_{x1} \frac{\delta B_{z1}}{\sqrt{4\pi\rho}}. \quad (17.34)$$

The set of equations (17.33) and (17.34) is incompatible with a non-zero amplitude of the incident wave, i.e. when δv_{z1} and δB_{z1} are not equal to zero. Note that if the incident wave is absent, this set has an infinite number of solutions. Hence the *switch-on shock is non-evolutionary*.

Finally it should be mentioned that the additional dissipative waves appear only for $v_x = V_{Ax}$. This means that

the dissipative effects do not alter the evolutionarity conditions for non-peculiar (*fast* and *slow*) MHD shock waves.

An increase or decrease of dissipation coefficients results only in a larger or smaller thickness of fast and slow shocks.

As for the Alfvén discontinuity, it becomes non-evolutionary with respect to dissipative Alfvén waves. This is well consistent with the fact that, in the presence of dissipation, the Alfvén discontinuity cannot have a stationary thickness and smooths out with time (see Landau et al. 1984).

It was also pointed out by Roikhvarger and Syrovatskii (1974) that the inclusion of dissipation into the dispersion relation for magnetoacoustic and entropy waves results in the appearance of dissipative waves, and, as a consequence, in non-evolutionarity of *tangential*, *contact*, and *weak* discontinuities (discontinuities of the derivatives of the MHD properties).

As we saw in Sect. 17.1.4, the *trans-Alfvénic shock waves* (TASWs) in the ideal MHD medium are non-evolutionary. Markovskii (1999) has shown that the dissipative processes do not alter this conclusion (for more detail see Sect. 17.4.2).

17.4 The Structure of Discontinuity and Its Evolutionarity


17.4.1 Perpendicular Shock Waves

It is natural to assume that

the stationary problem of the structure of an evolutionary MHD discontinuity has a unique solution, while for the non-evolutionary one this problem does not have such a solution.

To illustrate this assumption let us obtain the structure of the perpendicular shock (Sect. 16.2.3). With this aim the one-dimensional dissipative MHD equations should

Fig. 17.8 The structure of the perpendicular shock (*bold arrow*) connecting the states 1 and 2



be integrated over x . After that the conservation laws of mass, momentum, and energy, and Maxwell equations take the form (Polovin and Demutskii 1990):

$$\rho v = J, \quad (17.35)$$

$$Jv + p + \frac{B^2}{8\pi} - \mu \frac{dv}{dx} = S, \quad (17.36)$$

$$J \left[\frac{v^2}{2} + \frac{p}{\rho(\gamma_g - 1)} \right] + p v + \frac{vB^2}{4\pi} - \mu v \frac{dv}{dz} - \frac{v_m}{4\pi} B \frac{dB}{dx} = Q, \quad (17.37)$$

$$vB - v_m \frac{dB}{dx} = cE. \quad (17.38)$$

Here the thermal conductivity of the medium is assumed to be zero. J , S , and Q are constants of integration, γ_g is the adiabatic index, $\mu = (4/3)\eta + \zeta$, and ζ is a bulk viscosity (the indexes x and y at the quantities v_x and B_y are omitted).

From (17.35)–(17.38) we obtain the set of ordinary differential equations which describes the structure of the perpendicular shock:

$$\mu \frac{dv}{dx} = f(v, B), \quad (17.39)$$

$$v_m \frac{dB}{dx} = g(v, B), \quad (17.40)$$

where the right-hand-side functions

$$f(v, B) = \frac{\gamma_g + 1}{2} Jv - \gamma_g \left(S - \frac{B^2}{2\pi} \right) + \frac{\gamma_g - 1}{v} \left(Q - \frac{cE}{4\pi} B \right), \quad (17.41)$$

$$g(v, B) = vB - cE. \quad (17.42)$$

The curves $f(v, B) = 0$ and $g(v, B) = 0$ on the plane (v, B) are shown schematically in Fig. 17.8. At the points 1 and 2 of intersection of these curves the derivatives dv/dx and dB/dx equal zero simultaneously.

The points (B_1, v_1) and (B_2, v_2) correspond to the states ahead of the shock ($x \rightarrow -\infty$) and behind the shock ($x \rightarrow +\infty$). These are *stationary points* of the set of differential equations (17.39) and (17.40). The structure of the shock

$$v = v(x), \quad B = B(x) \quad (17.43)$$

is a solution to the set (17.39), (17.40) which leaves the initial point 1 and enters into the final point 2.

To consider the behavior of the integral curves in the vicinity of the stationary points 1 and 2 (Fig. 17.8) the quantities J , S , and Q should be expressed in terms of the MHD properties v_i and B_i ahead of the shock ($i = 1$) and behind the shock ($i = 2$). Then, by virtue of the fact that the derivatives dv/dx and dB/dx tend to zero for $x \rightarrow \pm \infty$, (17.35)–(17.37) yield

$$J = \rho_i v_i, \quad (17.44)$$

$$S = J v_i + p_i + \frac{B_i^2}{8\pi}, \quad (17.45)$$

$$Q = J \left(\frac{v_i^2}{2} + \frac{\gamma_g}{\gamma_g - 1} \frac{p_i}{\rho_i} \right) + \frac{v_i B_i^2}{4\pi}, \quad (17.46)$$

where $i = 1, 2$.

Let us now represent the quantities B and v in the form

$$B = B_i + \delta B_i, \quad v = v_i + \delta v_i, \quad (17.47)$$

with δ being a small perturbation. Substituting this together with (17.44)–(17.46) in (17.41) and (17.42), and expanding the result in powers of δB_i and δv_i , we find to the first order

$$\mu \frac{d\delta v_i}{dx} = \frac{\rho_i}{v_i} (v_i^2 - V_{si}^2) \delta v_i + \frac{B_i}{4\pi} \delta B_i, \quad (17.48)$$

$$v_m \frac{d\delta B_i}{dx} = B_i \delta v_i + v_i \delta B_i. \quad (17.49)$$

As is known (e.g., Fedoryuk 1985), a stationary point $\delta v_i = 0$, $\delta B_i = 0$ of the set of autonomous differential equations

$$\frac{d\delta v_i}{dx} = a_{11} \delta v_i + a_{12} \delta B_i, \quad (17.50)$$

$$\frac{d\delta B_i}{dx} = a_{21} \delta v_i + a_{22} \delta B_i \quad (17.51)$$


is a *saddle* if the roots of characteristic equation

$$(a_{11} - \lambda)(a_{22} - \lambda) - a_{12} a_{21} = 0 \quad (17.52)$$

are real numbers and have opposite signs, i.e. if

$$(a_{11} - a_{22})^2 + 4a_{12} a_{21} > 0, \quad a_{11} a_{22} - a_{12} a_{21} < 0. \quad (17.53)$$

Fig. 17.9 Stationary points of the set of autonomous differential equations. (a) Saddle. (b) Unstable node



In this case only two integral curves enter the stationary point $\delta v_i = 0$, $\delta B_i = 0$ from the opposite directions (Fig. 17.9a). And in the orthogonal way only two curves leave the stationary point.

In the case when the roots of characteristic equation (17.52) are real numbers and have the same sign, i.e. if

$$(a_{11} - a_{22})^2 + 4a_{12}a_{21} > 0, \quad a_{11}a_{22} - a_{12}a_{21} > 0, \quad (17.54)$$

then the stationary point is a *node*. If in addition

$$a_{11} + a_{22} > 0 \quad (17.55)$$

then the node is unstable, and all the integral curves leave the stationary point (Fig. 17.9b).

In a perpendicular MHD shock

$$a_{11}a_{22} - a_{12}a_{21} = \frac{\rho_i (v_i^2 - V_{\perp i}^2)}{\mu v_m}, \quad (17.56)$$

as follows from (17.48) and (17.49). Here

$$V_{\perp} = \sqrt{V_{A\parallel}^2 + V_s^2} = \sqrt{u_A^2 + V_s^2}. \quad (17.57)$$

(Sect. 15.2.4). So the second inequality (17.54) is always valid. As for the quantity $a_{11} + a_{22}$, it equals

$$a_{11} + a_{22} = \frac{\rho_i (v_i^2 - V_{si}^2)}{\mu v_i} + \frac{v_i}{v_m}. \quad (17.58)$$

It follows from (17.56) and (17.58) that in the case of the perpendicular shock the stationary points of (17.48), (17.49) can be only of two types: either a saddle or an unstable node (recall that v_i is assumed to be positive).

Let us consider at first the case when

$$v_1 > V_{\perp 1}, \quad v_2 < V_{\perp 2}. \quad (17.59)$$

Then point 2 is a saddle, while point 1 is an unstable node. The only integral curve enters into point 2 in Fig. 17.8 from the side of larger values of v . If the quantities v and B vary along this curve in the opposite direction, i.e. upstream of the shock, then they will inevitably reach the values (v_1, B_1) , i.e. point 1, because all integral curves leave point 1 (unstable node in the case under consideration). This curve describes a unique structure of the perpendicular shock. The inequalities (17.59) coincide with the conditions of evolutionarity of the perpendicular shock (see (17.15)), because $V_+ = V_{\perp}$ for perpendicular propagation. Therefore

the conditions that the perpendicular shock wave has the unique structure coincide with the conditions of its evolutionarity.

Now we consider the structure of a non-evolutionary perpendicular shock wave. If

$$v_2 > V_{\perp 2}, \quad (17.60)$$

then point 2 is an unstable node. Neither integral curve enters this point, i.e. the problem of structure of the shock does not have a solution.

If

$$v_1 < V_{\perp 1}, \quad v_2 < V_{\perp 2}, \quad (17.61)$$

then both stationary points 1 and 2 are saddles. In this case one of two integral curves, leaving point 1, may coincide with one of two curves entering point 2. However this can take place only for the definite exclusive values of the parameters ahead of the shock front. An infinitesimal perturbation of the state upstream of the shock destroys its structure. In other words, the integral curve cannot connect the states 1 and 2 in a general case under conditions (17.61).

17.4.2 Discontinuities with Penetrating Magnetic Field

Let us turn to the discontinuity type for which

$$v_x \neq 0 \quad \text{and} \quad B_x \neq 0 \quad (17.62)$$

(Sects. 16.2.4 and 16.2.6). Consider at first the discontinuity accompanied by a density jump:

$$\{\rho\} \neq 0. \quad (17.63)$$

(the oblique shock waves). In this case the boundary conditions (16.68) can be rewritten in such a way as to represent the Rankine–Hugoniot relation for shock




Fig. 17.10 Transitions with increasing entropy. Evolutionarity domains (*bold rectangles*) for the fast ($I \rightarrow II$) and slow ($III \rightarrow IV$) shock waves

waves in MHD. [Germain \(1960\)](#) has shown that the boundary conditions allow four states (see also [Shercliff 1965](#)):

$$\begin{aligned}
 I: \quad & v_x > V_+, \\
 II: \quad & V_+ > v_x > V_{Ax}, \\
 III: \quad & V_{Ax} > v_x > V_-, \\
 IV: \quad & V_- > v_x.
 \end{aligned} \tag{17.64}$$

The states are arranged in order of increasing entropy. The second law of thermodynamics requires that a shock transition is possible only from a lower state of entropy to an upper one. There are thus six transitions shown in Fig. 17.10.

The evolutionarity of an oblique shock wave is related to its structure in the following way ([Germain 1960](#); [Kulikovskii and Liubimov 1961](#); [Anderson 1963](#)). *The evolutionary fast and slow shocks always have a unique structure.*

The *trans-Alfvénic shock* (TASW) transition $II \rightarrow III$ may have a unique structure only for the definite relationship between the dissipative transport coefficients. If these coefficients fall into the certain intervals, the $I \rightarrow III$ and $II \rightarrow IV$ shocks also may have a unique structure, while the $I \rightarrow IV$ transition may be connected by an infinite number of integral curves.

[Markovskii \(1999\)](#) has considered the trans-Alfvénic shocks (TASWs) in the two-fluid approximation. He has shown that

the dissipative and dispersive processes that form the shock structure do not alter the conclusions based on the principle of evolutionarity.

An Alfvén discontinuity becomes non-evolutionary in the presence of arbitrary small but nonzero dissipation. The TASW disintegrates or transforms to some other nonsteady configuration. The effects of electrical resistivity in these processes




Fig. 17.11 (a) 2D schematic of the bow shock (BS) with the primary trans-Alfvénic shock wave (TASW) segment (ab); vector $\mathbf{n}$ is a normal to the shock front. The magnetopause (MP) is shown for reference. (b) Disintegration of the bow shock: TASW is a secondary trans-Alfvénic shock; S_+ and S_- are fast and slow shocks, respectively

were considered at first by [Todd \(1964\)](#). Such configuration includes an Alfvén discontinuity or another TASW which takes the *magnetic reversal* given at the initial shock (see Fig. 16.7).

According to [Markovskii \(1999\)](#), because of this reversal, the time evolution of a non-evolutionary shock can take the form of *oscillatory disintegration*, i.e. reversible transformation of one non-evolutionary discontinuity to the other. Therefore, most of the time of its evolution, such flow can be in the intermediate state between these two, and it cannot be identified as one of known types of discontinuities.

TASWs are widely used to model various phenomena in astrophysical plasmas. For example, one can imagine the following scenario for evolution of planetary *bow shocks*. As illustrated by Fig. 17.11a, depending on an angle between the *solar wind* (SW) and direction of *interplanetary magnetic field* (IMF), there appears some segment (ab) of the bow shock (BS) where the tangential component of magnetic field changes sign at the shock front. This segment represents the non-evolutionary TASW. Here disintegration of the TASW can appear according to [Markovskii and Skorokhodov \(2000\)](#).

After the disintegration, at least for some time, the solar wind is decelerated at the planet through several discontinuous structures (Fig. 17.11b) instead of one shock. In the space observations this behavior may manifest itself as an unusual position of the bow shock even under the condition that the upstream parameters do not differ much from their typical values.

As shown by [Liberman \(1978\)](#) with the help of the method discussed in Sect. 17.4.1, the *switch-on shock*, which is not evolutionary with respect to

dissipative waves, has a unique structure. The possible reason is that the peculiarity of the switch-on and switch-off shocks is related to the absence of B_τ on one side of the discontinuity surface. The small asymmetry, that is assumed when studying the stationary points, removes the degeneration, and thus can make the shock evolutionary.

17.5 Practice: Exercises and Answers

Exercise 17.1 (Sect. 17.1.1). Show that an ordinary shock wave is evolutionary.

Answer. From (16.7) it follows that there exist three boundary conditions at the surface of a shock wave in ordinary hydrodynamics:

$$\{\rho v_x\} = 0, \quad \{p + \rho v_x^2\} = 0, \quad \left\{\frac{v^2}{2} + w\right\} = 0. \quad (17.65)$$

The boundary condition

$$\{v_\tau\} = 0 \quad (17.66)$$

makes it possible to transform to such a frame of reference in which the tangential velocity component is absent on either side of the discontinuity:

$$\mathbf{v}_{\tau 1} = \mathbf{v}_{\tau 2} = 0.$$

So we obtain three linearized conditions for small perturbations. Since the disturbance of the velocity of the shock front surface δu_x can be eliminated from the set of boundary conditions, there remain two independent equations in the set.

Let us count the number of outgoing small-amplitude waves. There are no such waves upstream the shock because of the condition

$$v_{x1} > V_{s1} = 0, \quad (17.67)$$


where V_{s1} is the upstream sound velocity. At the downstream side of the shock there are two waves: the sound wave propagating with velocity $v_{x2} + V_{s2}$ and the entropy-vortex wave (Exercise 15.2) propagating with velocity v_{x2} as shown in Fig. 17.12. Therefore the number of waves leaving the shock is equal to the number of independent linearized boundary conditions; q.e.d.

Exercise 17.2 (Sect. 17.1.1). Since an ordinary shock wave is evolutionary, consider the linear problem of its stability in the ordinary sense of small perturbations.

Answer. Suppose that the surface of a shock is perturbed in the following way:

$$\xi = \xi_0 \exp [i (k_y y - \omega t)], \quad (17.68)$$

Fig. 17.12 Small-amplitude waves in a plasma moving through a shock wave without a magnetic field



where ξ is a displacement of the surface. The shock front thus becomes corrugated. The corrugation causes a perturbation of the flow. An arbitrary hydrodynamic perturbation is represented as a sum of the entropy-vortex wave and the sound wave. Since the flow is stationary and homogeneous in the y direction, all perturbations have the same frequency ω and tangential component of the wave vector k_y .

Since the flow velocity ahead of the shock $v_1 > V_{s1}$, only the downstream flow is perturbed. The usual condition of compatibility of the linear equation set is that the determinant of the coefficients at unknown quantities is zero, which yields the dispersion equation

$$\frac{\omega v_2}{v_1} \left(k_y^2 + \frac{\omega^2}{v_2^2} \right) - \left(\frac{\omega^2}{v_1 v_2} + k_x^2 \right) (\omega - k_y v_2) \left[1 + J^2 \left(\frac{\partial U_2}{\partial p_2} \right)_{\text{RH}} \right] = 0. \quad (17.69)$$

Here $U = 1/\rho$ is a specific volume, and

$$J = \rho_1 v_1 = \rho_2 v_2.$$

The subscript RH means that the derivative is taken along the Rankine–Hugoniot curve (see Fig. 16.12).

The shock front as a discontinuity is unstable if

$$\text{Im } \omega > 0, \quad \text{Im } k_x > 0. \quad (17.70)$$

The second condition (17.70) means that the perturbation is excited by the shock itself, but not by some external source. As shown by D'yakov (1954), (17.69) has solution which satisfies the condition (17.70), when

$$J^2 \left(\frac{\partial U_2}{\partial p_2} \right)_{\text{RH}} < -1 \quad (17.71)$$

or

$$J^2 \left(\frac{\partial U_2}{\partial p_2} \right)_{\text{RH}} > 1 + 2 \frac{v_2}{V_{s2}}. \quad (17.72)$$

If the parameters of the flow fall into the interval (17.71) or (17.72) then the small perturbation of the shock grows exponentially with time. This is the so-called *corrugational* instability of shock waves in ordinary hydrodynamics.

Along with this there is a possibility that (17.69) has solutions with real ω and k_x which correspond to *non-damping waves* outgoing from the discontinuity (D'yakov 1954). In this case

the shock spontaneously radiates sound and entropy-vortex waves, with the energy being supplied from the whole moving medium.

Apparently this instability is the reason of the flow inhomogeneities observed, for example, in laboratory experiments when a strong shock propagates in a gas (see Markovskii and Somov 1996).

Exercise 17.3 (Sect. 17.1.1). Show that an ordinary tangential discontinuity introduced in Sect. 16.1.2 is non-evolutionary.

Answer. From (16.6) it follows that there exists only one boundary condition at the surface of a tangential discontinuity in ordinary hydrodynamics. However two sound waves can propagate from the discontinuity at its both sides. Therefore the number of small-amplitude waves is greater than the number of linearized boundary equations at a tangential discontinuity.

Chapter 18

Particle Acceleration by Shock Waves

Abstract Sir Charles Darwin (Nature **164**, 1112–1114 (1949)) presumably thought that shock waves are responsible for accelerating cosmic rays. Nowadays shocks are widely recognized as a key to understanding high-energy particle acceleration in a variety of astrophysical environments.

18.1 Two Basic Mechanisms

Astrophysical plasma, being tenuous, differs from laboratory plasma in many ways. One of them is the following. In most environments where accelerated particles are observed, typical sound speeds are considerably less than easily obtainable bulk flow velocities, and shock waves are expected to develop. In fact, shocks are associated with most energetic particle populations seen in space.

In the heliosphere, *collisionless shocks* are directly observable with spacecrafts and they have been well studied. In every case where direct observations have been made, shocks are seen to accelerate particles, often to power-law distributions. Investigations of heliospheric shocks, along with a great deal of theoretical work, also show that collective field-particle interactions control the shock dissipation and structure. The physics of shock dissipation and particle acceleration seem to be intimately related.

In this chapter, we introduce only the most important aspects of the shock acceleration theory including two fundamental mechanisms of particle acceleration by a shock wave. Analytical models and numerical simulations (Jones and Ellison (1991); Blandford (1994); Giacalone and Ellison (2000); Parks (2004)) illustrate the possible high efficiency of *diffusive* and *drift* accelerations to high energies.

18.2 Shock Diffusive Acceleration

18.2.1 The Canonical Model of Diffusive Mechanism

Axford et al. (1977) and Krymskii (1977) considered the idealized problem of the particle acceleration by a shock wave of plane geometry propagating in a medium containing small-scale inhomogeneities of a magnetic field which scatter fast particles. The origin of these scatterers will be discussed later on. This may be, for example, the case of parallel or nearly parallel MHD shocks. In shocks of this kind (see case (16.73)) the average magnetic field plays essentially no role since it is homogeneous, while fluctuations in the average field play a secondary role producing particle scattering. Assuming this, we consider a shock wave as an ordinary hydrodynamic shock with scatterers.

If the medium is homogeneous, and if the propagation of the shock is stationary, then the front of the shock separates the two half-spaces: $x < 0$ and $x > 0$, and the velocity of the medium is given by the following formula:

$$v(x) = \begin{cases} v_1 & \text{for } x < 0, \\ v_2 = r^{-1}v_1 & \text{for } x > 0. \end{cases} \quad (18.1)$$

Here

$$r = \frac{\rho_2}{\rho_1} = \frac{v_1}{v_2} \quad (18.2)$$

is the compression ratio. It follows from formula (16.95) that, in a very strong (but non-relativistic) shock wave, the ratio

$$r \rightarrow r_\infty = \frac{\gamma_g + 1}{\gamma_g - 1}$$

and

$$v_2 = \frac{v_1}{r_\infty} = \frac{\gamma_g - 1}{\gamma_g + 1} v_1. \quad (18.3)$$

The adiabatic index γ_g is considered constant on both sides of the shock front $x = 0$.

Following Axford et al. (1977) and Krymskii (1977), let us assume that the distribution function in space and the scalar momentum of the accelerated particles, $f(\mathbf{r}, p)$, is isotropic (see generalization in Gieseler et al. 1999; Ruffolo 1999). This means that $f(\mathbf{r}, p)$ is the same in all reference frames to first order in the small parameter v/v_p , where v_p and p are the individual particle velocity and momentum measured in the local plasma frame.

As long as scattering is strong enough to insure the isotropy assumption, the kinetic equation (2.18) describing the transport of particles with $v_p \gg v$ in space and velocity *diffusion-convection* equation (see Krymskii (1977) and references therein):

$$\frac{\partial f}{\partial t} = \nabla_{\mathbf{r}} (D \nabla_{\mathbf{r}} f) - \nabla_{\mathbf{r}} (f \mathbf{v}) + \frac{1}{3} \frac{\partial(f p)}{\partial p} \operatorname{div} \mathbf{v}. \quad (18.4)$$

Here $D = D(\mathbf{r}, p)$ is the coefficient of diffusion of fast particles.

For our problem under consideration, with one-dimensional geometry, we have in the *stationary* case

$$\frac{\partial}{\partial x} \left[v f(x, p) - D(x, p) \frac{\partial f(x, p)}{\partial x} \right] = \frac{1}{3} \frac{\partial v}{\partial x} \frac{\partial}{\partial p} [p f(x, p)]. \quad (18.5)$$

Let us integrate (18.5) over x from $x = -\infty$ to $x = +\infty$. By employing the boundary conditions

$$f(x = -\infty, p) = f_1(p) \quad \text{and} \quad f(x = +\infty, p) = f_2(p), \quad (18.6)$$

where $f_2(p)$ is an unknown spectrum of accelerated particles, we obtain the following differential equation in p

$$v_2 f_2(p) - v_1 f_1(p) - 0 + 0 = \frac{1}{3} (v_2 - v_1) \frac{d}{dp} [p f_2(p)]. \quad (18.7)$$

Using the definition of the compression ratio (18.2), we obtain an ordinary differential equation for the downstream distribution function $f_2(p)$ in the form

$$p \frac{d}{dp} f_2(p) + \frac{r+2}{r-1} f_2(p) = \frac{3r}{r-1} f_1(p); \quad (18.8)$$

recall that $r > 1$.

The general solution of this equation is

$$f_2(p) = \frac{3}{r-1} p^{-\gamma_p} \int_{p_0}^p f_1(p') (p')^{-\gamma_p} dp' + c_1 p^{-\gamma_p}. \quad (18.9)$$


Here

$$\gamma_p = \frac{r+2}{r-1}$$

(18.10)

plays the role of the *spectral index* of the accelerated particles, c_1 is an arbitrary constant of integration which multiplies the homogeneous term, the distribution function $f_1(p)$ is the far upstream spectrum of ambient particles that are accelerated by the shock, and p_0 is large enough so that the assumption $v_p \gg v$ holds.

Fig. 18.1 The compression ratio r and spectral index γ_p versus the Mach number M



So the solution of the diffusion-convection equation does show that a planar shock, propagating through a region in which fast particles are diffusing, produces a superthermal population of particles with the power-law momentum distribution

$$f_2(p) \sim p^{-\gamma_p}. \quad (18.11)$$

The property which gave the diffusive acceleration process a wide appeal is the fact that, with the simplest assumptions made above,

the spectral index (18.10) of the accelerated particles depends only on the compression ratio r of the shock wave.

Most astrophysical shocks, since they are strong, have compression ratios constrained to a rather narrow range of values near $r_\infty = 4$ assuming $\gamma_g = 5/3$. For a shock with Mach number M (see Exercise 16.5) greater than 3 say, as we see in Fig. 18.1, the compression ratio $3 < r < 4$ and the spectral index $2 < \gamma_p < 2.5$.

A spectral index of $\gamma_p \approx 2$ is characteristic of energy particle spectra observed in a wide range of astrophysical environments (Jones and Ellison 1991; Blandford 1994). For example, $\gamma_p \approx 2$ closely fits the inferred source spectrum of *galactic cosmic rays* for high energies below approximately 10^{15} eV (e.g., Gombosi 1999). In the cosmic rays observed at Earth, the spectrum of cosmic-ray ions is an unbroken power law from 10^9 to 10^{15} eV. The supernova shocks are one of the few mechanisms known to be capable of providing adequate energy to supply the pool of Galactic cosmic rays. Supernova remnants (SNRs) have long been suspected as the primary site of Galactic cosmic-ray acceleration.

The earliest evidence of non-thermal X-ray emission in a SNR came from featureless observed spectra interpreted as the extrapolation of a radio synchrotron spectrum. However early data were poor and the models were simplistic. New observations and theoretical results (e.g., Dyer et al. 2001) indicate that joint thermal and non-thermal fitting, using sophisticated models, will be required for analysis of

most supernova-remnant X-ray data in the future to answer two questions: (a) Do SNRs accelerate ions? (b) Are they capable of accelerating particles to energies of 10^{15} eV?

In the *solar wind* the shock-associated low-energy-proton events seem to be well studied. The most intensive of them have a power-low energy spectrum, suggesting that protons are accelerated by the diffusive-shock acceleration mechanism (e.g., [Rodrigues-Pacheco et al. 1998](#)). Nevertheless the correlation between the spectral exponent γ with the solar wind velocity compression ratio is found to be linear. This result differs from that presented above. The discrepancy of the spectral-exponent dependence on the shock-wave parameters could lie on the event selection criterion or on the account of nonlinear effects (Sect. 18.2.3) or on another mechanism of acceleration.

18.2.2 Some Properties of Diffusive Mechanism

As we saw above, the spectral index γ_p of energetic particles produced by diffusive shock acceleration does not depend on the diffusion coefficient D . However the diffusion coefficient D , together with the characteristic flow velocity $v \sim v_1$, determines the overall length scale of the acceleration region

$$l_D \sim D(p)/v \quad (18.12)$$

and acceleration time

$$t_D \sim D(p)/v^2. \quad (18.13)$$

The first-order Fermi or diffusive shock acceleration is a statistical process in which particles undergo spatial diffusion and are accelerated as they scatter back and forth across the shock, thereby being compressed between scattering centers fixed in the converging upstream and downstream flows.

Particle energies are derived just from the relative motion, the converging flow with velocity $v_1 - v_2$, between scatterers (waves) on either side of a shock front.

This is a main advantage of the diffusive mechanism. Its disadvantage is that particles can achieve very high energies by diffusion acceleration, but

since particles spend most of their time random walking in the upstream or downstream plasma, the acceleration time can become excessively large

compared with, for example, the shock's life time.

Another disadvantage in applying it to some astrophysical phenomena, for example solar flares, consists of the lack of actual knowledge about the assumed scattering waves. However diffusion determines only the length scale (18.12) and characteristic time (18.13) of the acceleration process. In this context, let us recall

once more (Sect. 16.1.3) the following analogy from everyday life. A glass of hot water with a temperature T_1 will invariably cool to a given room temperature T_2 , independently of the mechanism of heat exchange with the surrounding medium, while the mechanism determines only the time of cooling.

In the presence of a magnetic field in plasma, the diffusive acceleration requires that the particles are able to traverse the shock front in both directions either along the field or by scattering across the field, in order that they may couple to the shock compression by pitch-angle scattering both upstream and downstream of the shock. At *quasi-parallel* shocks this condition on particle mobility is easily met. For sufficiently fast shocks, downstream shock-heated particles can be kinematically able to return to the shock along the downstream magnetic field to initiate the process of diffusive shock acceleration. At *quasi-perpendicular* shocks (Sect. 18.3.2), however, this condition is stringent. Although the diffusive mechanism is rapid since particles are confined closer to the shock front, there is a *high threshold speed*, significantly exceeding v_1 , in order that diffusive acceleration can occur (Webb et al. 1995).

18.2.3 Nonlinear Effects in Diffusive Acceleration

The test particle (i.e., *linear*) model demonstrated above yields the most important result: the power law (18.11) with the spectral index (18.10) is the natural product of the diffusive acceleration in shock waves. The equally important question of the actual efficiency of the process can only be adequately addressed to a fully *nonlinear* (and more complex) theory.

Using observations of the Earth bow shock and interplanetary observations, numerical modeling of different shocks shows that the inherent efficiency of shock acceleration implies that

the hydrodynamic feedback effects between the accelerated particles and the shock structure are important

and therefore essential to any complete description of the process. This has turned out to be a formidable task because of the wide range of spatial and energy scales that must be self-consistently included in numerical simulations.

On the one hand, the plasma micro-processes of the shock dissipation control injection from the thermal population. On the other hand, the highest energy particles (extending to 10^{14} – 10^{15} eV in the case of galactic cosmic rays) with extremely long diffusion lengths (18.12) are dynamically significant in strong shock waves and feed back on the shock structure. Ranges of interacting scales of many orders of magnitudes must be described self-consistently (for review, see Parks 2004).

18.3 Shock Drift Acceleration

The principal process whereby a particle gains energy upon crossing a shock wave with a magnetic field may be the so-called shock drift acceleration (Hudson 1965). The drift mechanism, in contrast to the diffusive one, neglects any shock-front associated turbulence. So many not-well-justified assumptions concerning the physics of scatterers have not to be made in applying the drift acceleration model to an astrophysical phenomenon.

If the fast particle Larmor radius

$$r_L = \frac{c p_\perp}{eB} \gg l_f, \quad (18.14)$$

where l_f is the front thickness, we can replace the shock by a simple discontinuity (the shock surface) and can approximate the particle motion as scatter-free on both sides of the shock. Let us begin by considering an interaction of individual particles with such a discontinuity. We shall consider very fast particles:

$$v_p \gg v_1 > v_2. \quad (18.15)$$

These assumptions are basic for further considerations that we start from the simplest case—a perpendicular shock (Sect. 16.2.3).

18.3.1 Perpendicular Shock Waves

As shown in Fig. 16.5, the magnetic fields $\mathbf{B}_1$ and $\mathbf{B}_2$ are parallel to the shock front, the plane $x = 0$; and plasma moves perpendicularly to the front. According to (16.41), there exists an identical electric field on both sides of the shock:

$$\mathbf{E} = -\frac{1}{c} \mathbf{v}_1 \times \mathbf{B}_1 = -\frac{1}{c} \mathbf{v}_2 \times \mathbf{B}_2. \quad (18.16)$$

The fast particles rotate on the magnetic field lines and move together with the field lines with the plasma speed across the front as shown in Fig. 18.2.

Nothing will happen before the Larmor ring touches the front; a particle simply drifts to the front. For what follows it is important that the particle will make many rotations (Fig. 18.3) during the motion of the Larmor ring across the front because of the condition (18.15). A “single encounter” consists of many individual penetrations by the particle through the shock surface as the particle follows its nearly helical trajectory. Because of the difference between the Larmor radius ahead of and behind the front, a drift parallel to the front will appear, accompanying the drift across the front.




Fig. 18.2 The Larmor ring moves together with the plasma and the magnetic field across the perpendicular shock front




Fig. 18.3 The trajectory of a negatively charged particle (an electron) multiply crossing the perpendicular shock front

During each rotation, the electric field $\mathbf{E}$ accelerates a particle on the upstream side ($x < 0$) of the shock and decelerates it on the downstream side ($x > 0$). However the work of the field $\mathbf{E}$ on a larger circle exceeds the work on the smaller circle:

$$\delta A_1 = +eE \times AC > -\delta A_2 = eE \times BC, \quad (18.17)$$

since the length AC is larger than the length AB . Therefore, during each rotation, the particle is slightly accelerated. How much energy does the particle take during the motion of its Larmor ring across the shock front?

Since we consider the shock as a discontinuity, the adiabatic approximation is formally not suitable. However it appears that the *first* or *transversal* invariant (Sect. 6.2) conserves:

$$\frac{p_{\perp}^2}{B} = \text{const} \quad (18.18)$$

(Hudson 1965; Alekseyev and Kropotkin 1970). From (18.18) it follows that

$$p_{\perp 2}^2 = p_{\perp 1}^2 \times \frac{B_2}{B_1}.$$

Therefore the transversal kinetic energy of a non-relativistic particle

$$\frac{\mathcal{K}_{\perp 2}}{\mathcal{K}_{\perp 1}} = \frac{p_{\perp 2}^2}{p_{\perp 1}^2} \propto \frac{B_2}{B_1} = r. \quad (18.19)$$

■ An increase of transversal energy (18.19) is relatively small when the Larmor ring of a particle crosses the front only once.

Multiple interactions of a particle with the shock is a necessary condition for a considerable increase of energy.

Drift acceleration typically involves several shock crossings and results from a net displacement δz of an ion (electron) guiding center parallel (anti-parallel) to the convection electric field $\mathbf{E}$. The energy gain is proportional to this displacement, which in general depends upon the plasma and shock parameters, the particle species and velocity, and the intensity of possible electromagnetic fluctuations in the vicinity of the shock as well as within the shock front itself.

It is popular to discuss the displacement δz as the consequence of a gradient drift (see formula (5.14) in Jones and Ellison 1991). Such a treatment is not reasonable when we consider the shock as a discontinuity; so formally $\nabla B \rightarrow \infty$. A wonderful thing is that the adiabatic approximation is not applicable for such a situation but the first adiabatic invariant (18.18) conserves.

18.3.2 Quasi-Perpendicular Shock Waves

18.3.2(a) Classical Model of Acceleration

The basic aspects of drift acceleration of fast particles by an almost perpendicular shock wave, as a discontinuity, emerge from a simple model which is valid for a certain range of incident pitch angles and which allows us to derive analytical expressions for the reflection and transmission coefficients, the energy and the angular distributions (Toptyghin 1980; Decker 1983).

By definition, in a *quasi-perpendicular shock*, the angle Ψ_1 (Fig. 18.4) between the shock normal $\mathbf{n}$ and the upstream magnetic field vector $\mathbf{B}_1$ is greater than about 80° . Hence the field lines form small angles α_1 and α_2 with the shock front plane $x = 0$. Under this condition, as well as for the perpendicular shock case considered above, *the first adiabatic invariant is conserved* (Hudson 1965; see also Sect. 4 in Wentzel 1964). This enables analytical calculations of the energy increase on the front of a quasi-perpendicular shock as well as the reflection and transmission of fast particles (Sarris and Van Allen 1974).




Fig. 18.4 A quasi-perpendicular shock wave in the frame of reference S where $\mathbf{v}_1 \parallel \mathbf{n}$

Since the particles conserve the first adiabatic invariant (Sect. 6.2.1), all particles with pitch angles

$$\theta > \theta_0 \quad (18.20)$$

will be reflected. To find the critical pitch angle θ_0 , consider two frames of reference: S and S' .

In the frame of reference S , where the shock front is in the plane (y, z) and the shock normal $\mathbf{n}$ is along the x axis, there is an electric field

$$\mathbf{E} = -\frac{1}{c} \mathbf{v}_1 \times \mathbf{B}_1 = -\frac{1}{c} \mathbf{v}_2 \times \mathbf{B}_2. \quad (18.21)$$

In the frame of reference S' , where $\mathbf{B}_1 \parallel \mathbf{v}_1$ and $\mathbf{B}_2 \parallel \mathbf{v}_2$ (Sect. 16.2.4), there is *no electric field*. The system S' moves along the y axis (perpendicular to the vector $\mathbf{E}$) with velocity

$$\mathbf{v}_y = c \frac{\mathbf{E} \times \mathbf{B}_n}{B_n^2}, \quad (18.22)$$

where $\mathbf{B}_n$ is the normal component of the magnetic field. Since $\mathbf{E}' = 0$, there is no change in the energy of a fast particle after reflection from the front: $\delta\mathcal{E}' = 0$.

We shall assume that B_n is very small but $v_y < c$. Using the relativistic Lorentz transformation for the energy-momentum 4D-vector with condition $\delta\mathcal{E}' = 0$, we obtain the relative energy increment of the reflected fast particles (see Exercise 18.1):

$$\frac{\delta\mathcal{K}}{\mathcal{K}} \approx \frac{4v_1^2}{v_p^2} \left[\frac{v_p \cos \theta}{v_1} + \operatorname{tg} \Psi_1 \right] \operatorname{tg} \Psi_1. \quad (18.23)$$

Here $\mathcal{K} = mv_p^2/2$ is the kinetic energy of a particle in the shock wave frame of reference S , v_p is the particle velocity in the same frame, and θ is the pitch angle also in the frame S . The connection between θ' and θ is given by

$$\cos \theta' = \frac{v_p \cos \theta + v_1 \operatorname{tg} \Psi_1}{\left[v_p^2 + (v_1 \operatorname{tg} \Psi_1)^2 + 2v_p (v_1 \operatorname{tg} \Psi_1) \cos \theta \right]^{1/2}}. \quad (18.24)$$

In the S' frame of reference, where the electric field is zero, the first adiabatic invariant can be written as (see definition (6.11)):

$$\frac{\sin^2 \theta'}{B} = \text{const}. \quad (18.25)$$

So the critical pitch angle θ'_0 satisfies equation

$$\sin^2 \theta'_0 = \frac{B_1}{B_2}. \quad (18.26)$$

This allows us to calculate the critical pitch angle θ_0 in the shock-front frame S . For example, if a non-relativistic proton has an initial energy $\mathcal{K} = 0.3 \text{ MeV}$ and if a shock wave has an upstream velocity $v_1 = 150 \text{ km/s}$, the ratio $B_1/B_2 = 1/3$, and the angle $\Psi_1 = 88^\circ$ and 89° , then we find, correspondingly, $\theta_0 = 55^\circ$ and 77° . As the angle Ψ_1 increases toward 90° , most of the particles are really transmitted into the downstream side. At $\Psi_1 = 90^\circ$, which is the perpendicular shock case, there are no reflecting particles.

Formula (18.23) shows that

the relative increment of kinetic energy of a fast particle increases when the angle Ψ_1 increases toward 90° .

The model under consideration predicts the *high field-aligned anisotropy* of accelerated particles for a large Ψ_1 because of conservation of first adiabatic invariant and the large energy gains.

It is widely believed that the slow *thermal particles* inside the shock front can also be considered as adiabatic, at least, in thick collisionless shocks: the electron magnetic moment is conserved throughout the shock and $v_\perp^2/B = \text{const}$ (Feldman et al. 1982). In very thin collisionless shock (with a large cross-shock potential) the adiabaticity may break down, so that electrons become demagnetized. It means that the magnetic moment is no longer conserved, and a more substantial part of the energy may be transferred into the perpendicular degree of freedom (Balikhin et al. 1993; Gedalin and Griv 1999).

18.3.2(b) Some Astrophysical Applications

Observations of interplanetary shocks (e.g., Balogh and Erdős 1991) show that the intensive acceleration of protons occurs when the upstream magnetic field is almost parallel to the shock front. Energetic particles entering the shock front stay with it, crossing it many times and being accelerated by the electric field of the front.




Fig. 18.5 A collapsing magnetic trap on the upstream side of a quasi-perpendicular shock wave

After the direction of the interplanetary magnetic field changes again away from the parallel to the front, the intensive acceleration ceases.

Owing to interplanetary magnetic field fluctuations the upstream field vector $\mathbf{B}_1$, if it is found to be parallel to the shock front, stays as such for only a short time (a few minutes, in general). This time is enough for the low-energy protons ($\mathcal{K}_p < 1$ MeV) to be accelerated to about 2–3 times their original energy but not enough for the high-energy protons ($\mathcal{K}_p < 10$ MeV) to be noticeably affected by the shock wave.

Single scatter-free shock drift interactions at quasi-perpendicular shocks can accelerate particles to at most a few times the shock compression ratio. Weak scattering during single drift interactions can increase this upper limit for a small fraction of an incident particle distribution, but the energy spectra will be still rather steep. One anticipates large energy gains and flatter spectra that extend to high energies if some particles can return to the shock for many drift interactions.

This is suggestive of the classical case of a *collapsing magnetic trap* (Sect. 6.2), and is the basis of the model of proton trapping and acceleration due to multiple drift interactions along magnetic loops that convect through a planar quasi-perpendicular shock (Wentzel 1963, 1964; Gisler and Lemons 1990). Figure 18.5 represents a quasi-perpendicular shock, with a small perturbation of the magnetic field superimposed on the un-shocked homogeneous field $\mathbf{B}_1$. The heavy line displays a particular field line which intersects the shock front plane $x = 0$ two times, forming a magnetic loop in the upstream region.

Upstream particles bounce back and forth along a loop and gain parallel energy at each reflection until they fall within the loss cone and transmit downstream.

Simple analytic models and detailed numerical study have shown that the collapse of the trap by the convection of the loop field lines through the shock is accompanied by a considerable increase of the accelerated proton flux, which may be responsible for the “shock spike” events observed near fast mode interplanetary shock waves (Decker 1993; Erdős and Balogh 1994).

In general, if the magnetic field contains fluctuations with wavelengths that are considerably larger than the gyro-radii of the fast particles, a fraction of particles is accelerated by a quasi-perpendicular shock to energy well above the thermal energy (Giacalone and Ellison 2000).

18.3.3 Oblique Shock Waves

If values of the angle α between the magnetic field and the shock front plane are arbitrary, then the phase-averaged coefficients of reflection and transmission are complicated and can be found, in principle, by numerical calculations. When

$$\frac{v_1}{v_p} \leq \alpha_1 \leq \frac{\pi}{2} \quad (18.27)$$

and the pitch angle θ is arbitrary, the order of magnitude of the energy increase

$$\delta\mathcal{K} \approx \frac{p v_1}{\alpha_1} \ll \mathcal{K} = \mathcal{E} - mc^2 \quad (18.28)$$

is small in comparison with the initial kinetic energy. In a general case, the increase of particle energy is small when the Larmor ring of a particle crosses the front once. Multiple interactions of a particle with the shock front is the necessary condition for a considerable increase of energy.

One possibility for multiple interactions of a particle with the shock is a *strong* MHD turbulence. More exactly, it is assumed that in a sufficiently large region of space there exists an ensemble of MHD shocks which interact successively with the particles. The investigation of particle acceleration by a random shock wave ensemble is of certain interest in astrophysical applications but the conditions of such an acceleration mechanism are not totally clear yet.

Another possibility is the propagation of one shock in a turbulent medium or of an oblique collisionless shock when magnetic turbulence exists in the regions upstream and downstream of the shock (Decker and Vlahos 1986). It is important, however, that the particle acceleration near the shock front in a turbulent medium, i.e. the *diffusive* mechanism (Sect. 18.2) will take place in the absence of a regular electric field. No terms should be added to the basic *diffusion-convection* equation (18.4) to take account of the drift mechanism in an oblique shock. The process is already included in the energy change which is proportional to the $\text{div } \mathbf{v}$ term. This, of course, assumes that there is sufficient scattering and that other assumptions used in deriving the diffusion-convection equation are also valid. That is not trivial.

The interesting possibility discussed in Sect. 18.3.2 is a combination of a magnetic trap with an oblique shock wave. In Part II, Sect. 7.3, this idea is applied to the particle acceleration in solar flares.

18.4 Practice: Exercises and Answers

Exercise 18.1. Derive formula (18.23) in Sect. 18.3.2.

Answer. According to the geometry shown in Fig. 18.4, the frame of reference S' moves with respect to the shock wave frame of reference S with velocity (18.22):

$$\mathbf{v}_y = -\frac{cE}{B_n} \mathbf{e}_y. \quad (18.29)$$

In the frame S' there is no electric field; therefore there is no change in the energy of a particle reflecting at the shock front, that is $\delta\mathcal{E}' = 0$, where $\mathcal{E}'$ is the energy of the particle in S' . Let us transform this condition back to S by using the Lorentz transformation of the energy-momentum 4D-vector (Landau and Lifshitz 1975, Chap. 2, Sect. 9):

$$\begin{aligned} p_x &= p'_x, & p_y &= \gamma_L \left(p'_y + \frac{v_y}{c^2} \mathcal{E}' \right), & p_z &= p'_z, \\ \mathcal{E} &= \gamma_L \left(\mathcal{E}' + v_y p'_y \right). \end{aligned} \quad (18.30)$$

Since $\delta\mathcal{E}' = 0$, it follows from (18.30) that

$$\delta\mathcal{E} = \gamma_L v_y \delta p'_y. \quad (18.31)$$

The change in the y component of momentum of the reflected particle in the frame of reference S' is

$$\delta \mathbf{p}'_y = -2\mathbf{p}'_y = 2\gamma_L \left(\mathbf{p}_y - \frac{v_y}{c^2} \mathcal{E} \mathbf{e}_y \right). \quad (18.32)$$

Note that vectors $\mathbf{p}_y$ and $\mathbf{v}_y$ point in opposite directions.

Substituting (18.32) into (18.31) gives us

$$\delta\mathcal{E} = \frac{2v_y}{1 - v_y^2/c^2} \left[p_y + \frac{v_y}{c^2} (\mathcal{K} + mc^2) \right], \quad (18.33)$$

where $\mathcal{K} = mv_p^2/2$ is kinetic energy of a particle. Assuming $\mathcal{K} \ll mc^2$ and using (18.29), we obtain

$$\delta\mathcal{E} = \delta\mathcal{K} = \frac{2E}{B_n^2 - E^2} \left(\frac{v_{p,y}}{c} B_n + E \right) mc^2, \quad (18.34)$$

where $v_{p,y}$ is the y component of the particle velocity.

According to (18.21) the electric field

$$E = \frac{1}{c} v_1 B_{y1} , \quad (18.35)$$

where B_{y1} is the y component of the vector $\mathbf{B}_1$. So we rewrite formula (18.34) as follows

$$\delta\mathcal{K} = 2mv_1^2 \frac{(v_{p,y}/v_1)(B_n/B_{y1}) + 1}{(B_n/B_{y1})^2 - (v_1/c)^2} . \quad (18.36)$$

The condition $v_y < c$ can equivalently be written as

$$\frac{B_n}{B_{y1}} > \frac{v_1}{c} \quad \text{or} \quad \text{tg } \Psi_1 < \frac{v_1}{c} . \quad (18.37)$$

If we further assume that

$$\frac{B_n}{B_{y1}} \gg \frac{v_1}{c} , \quad (18.38)$$

we obtain from (18.36) the following formula

$$\delta\mathcal{K} = 2mv_1^2 \left(\frac{v_p \cos \theta}{v_1} \text{tg } \Psi_1 + \text{tg}^2 \Psi_1 \right) , \quad (18.39)$$

where θ is the pitch angle in the shock-front frame of reference S . Dividing (18.39) by $\mathcal{K}$, we obtain formula (18.23).

Chapter 19

Plasma Equilibrium in Magnetic Field

Abstract The concept of equilibrium is fundamental to any discussion of the energy contained in an astrophysical object or phenomenon. The MHD non-equilibrium is often related to the onset of dynamic phenomena in astrophysical plasma, for example, in the solar corona. In this chapter, we derive the virial theorem and the famous Shafranov theorem as well as we consider some general properties of equilibrium configurations.

19.1 The Virial Theorem in MHD

19.1.1 A Brief Pre-history

An integral equality relating different kinds of energy (kinetic, thermal, gravitational, etc.) of some region with a volume V and a surface S , is commonly referred to as the *virial theorem*. It has been proved for mechanical systems for the first time by [Clausius \(1870\)](#). The derivation of the virial theorem for a mechanical system executing a motion in some finite region of space, velocities also being finite, can be found, for example, in [Landau and Lifshitz \(1976, Chap. 2, Sect. 10\)](#). Its relativistic form is presented in [Landau and Lifshitz \(1975, Chap. 4, Sect. 34\)](#).

The generalization of the virial theorem to include the magnetic energy in the context of MHD was achieved by [Chandrasekhar and Fermi \(1953\)](#) when addressing the question of the gravitational stability of infinitely conductive masses of cosmic dimensions in the presence of a magnetic field. Although “most students of physics will recognize the name of the virial theorem, few can state it correctly and even fewer appreciate its power” ([Collins 1978](#)).

19.1.2 Deduction of the Scalar Virial Theorem

The virial theorem is deduced from the momentum conservation law (see the ideal MHD motion equation (12.71) or (13.1)) rather than the energy conservation law. We have

$$\rho \frac{dv_\alpha}{dt} \equiv \rho \left(\frac{\partial v_\alpha}{\partial t} + v_\beta \frac{\partial v_\alpha}{\partial r_\beta} \right) = - \frac{\partial p}{\partial r_\alpha} - \frac{\partial M_{\alpha\beta}}{\partial r_\beta} - \rho \frac{\partial \phi}{\partial r_\alpha}. \quad (19.1)$$

Here

$$M_{\alpha\beta} = \frac{1}{4\pi} \left(\frac{B^2}{2} \delta_{\alpha\beta} - B_\alpha B_\beta \right) \quad (19.2)$$

is the Maxwellian stress tensor. So we consider an ideal MHD plasma distributed within a limited region V of space. The gravitational potential at a point $\mathbf{r}$ is

$$\phi(\mathbf{r}) = -G \int \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d^3\mathbf{r}', \quad (19.3)$$

where G is the gravitational constant (Appendix 3), $d^3\mathbf{r}' = dx' dy' dz'$.

The partial differential equations (19.1) are often very difficult to solve. Moreover, in astrophysics, we may have such incomplete knowledge of a system that it may not be worthwhile to work out an elaborate solution. In many situations, it is possible to make important conclusions if we know some global relationships among the different forms of energy in the system.

Let us multiply the plasma motion equation (19.1) by r_α and integrate it over the volume V . We observe in passing that multiplication of (19.1) by r_γ rather than r_α would result, on integrating, in the *tensor* virial theorem and not in the *scalar* one (Chandrasekhar 1981; see also Strittmatter 1966; Choudhuri 1998).

First let us integrate the left-hand side of (19.1) multiplied by r_α . We get

$$\int \rho r_\alpha \frac{dv_\alpha}{dt} dV = \int r_\alpha \frac{d^2 r_\alpha}{dt^2} \rho dV = \int r_\alpha \frac{d^2 r_\alpha}{dt^2} dm. \quad (19.4)$$

Here we have passed from the integration over volume to integration over mass: $dm = \rho dV$. We rearrange formula (19.4) as follows

$$\begin{aligned} r_\alpha \frac{d^2 r_\alpha}{dt^2} &= \frac{d}{dt} \left(r_\alpha \frac{dr_\alpha}{dt} \right) - \left(\frac{dr_\alpha}{dt} \right)^2 = \\ &= \frac{d}{dt} \left(\frac{1}{2} \frac{dr_\alpha^2}{dt} \right) - \left(\frac{dr_\alpha}{dt} \right)^2 = \frac{1}{2} \frac{d^2}{dt^2} r_\alpha^2 - v_\alpha^2. \end{aligned}$$

On substituting this into (19.4), we obtain

$$\int \rho r_\alpha \frac{dv_\alpha}{dt} dV = \frac{1}{2} \frac{d^2}{dt^2} \int r^2 dm - \int v^2 dm = \frac{1}{2} \frac{d^2 I}{dt^2} - 2U_{kin}. \quad (19.5)$$

Here

$$I = \int r^2 dm \quad (19.6)$$

is the *moment of inertia* in the reference frame related to the mass center of the system. When the system expands, its moment of inertia I increases.

$$U_{kin} = \int \frac{v^2}{2} dm \quad (19.7)$$

is *kinetic energy* or (to be more specific) the kinetic energy of macroscopic motions inside the system.

Let us multiply the first term on the right-hand side of (19.1) by r_α and integrate it over volume:

$$-\int_V r_\alpha \frac{\partial p}{\partial r_\alpha} dV = -\oint_S p r_\alpha dS_\alpha + 3 \int_V p dV, \quad (19.8)$$

since

$$\frac{\partial}{\partial r_\alpha} (p r_\alpha) = r_\alpha \frac{\partial p}{\partial r_\alpha} + p \frac{\partial r_\alpha}{\partial r_\alpha} = r_\alpha \frac{\partial p}{\partial r_\alpha} + 3p.$$

The Ostrogradskii-Gauss theorem [Smirnov 1965](#), Sect. 126 was used to integrate the divergence over the volume V in formula (19.8).

If U_{th} is the *thermal energy* of the plasma, γ_g is the ratio of specific heats at constant pressure and at constant volume, then

$$\int_V p dV = (\gamma_g - 1) U_{th}. \quad (19.9)$$

Therefore

$$-\int_V r_\alpha \frac{\partial p}{\partial r_\alpha} dV = -\oint_S p (\mathbf{r} \cdot d\mathbf{S}) + 3(\gamma_g - 1) U_{th}. \quad (19.10)$$

Similarly we calculate the integral

$$-\int_V r_\alpha \frac{\partial M_{\alpha\beta}}{\partial r_\beta} dV = -\int_S M_{\alpha\beta} r_\alpha dS_\beta + \int_V M_{\alpha\beta} \delta_{\alpha\beta} dV \quad (19.11)$$

since

$$\frac{\partial}{\partial r_\beta} (r_\alpha M_{\alpha\beta}) = r_\alpha \frac{\partial M_{\alpha\beta}}{\partial r_\beta} + M_{\alpha\beta} \delta_{\alpha\beta}.$$

On rearranging, we find from (19.11) and (19.2)

$$-\int_V r_\alpha \frac{\partial M_{\alpha\beta}}{\partial r_\beta} dV = \mathcal{M} - \int_S \left[\frac{B^2}{8\pi} (\mathbf{r} \cdot d\mathbf{S}) - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{r}) (\mathbf{B} \cdot d\mathbf{S}) \right], \quad (19.12)$$

where

$$\mathcal{M} = \int_V \frac{B^2}{8\pi} dV \quad (19.13)$$

is the *magnetic energy* of the system.

The third term on the right-hand side of (19.1) gives

$$\begin{aligned} -\int_V r_\alpha \frac{\partial \phi}{\partial r_\alpha} \rho dV &= \int_V \rho r_\alpha \frac{\partial}{\partial r_\alpha} \int_{V'} \frac{G \rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} dV' dV = \\ &= G \int_V \int_{V'} \rho \rho' r_\alpha \frac{\partial}{\partial r_\alpha} \frac{1}{\sqrt{(r_\beta - r'_\beta)^2}} dV dV'. \end{aligned} \quad (19.14)$$

We rewrite the expression as follows. Let the distance

$$R = \sqrt{(r_\beta - r'_\beta)^2}.$$

Then

$$r_\alpha \frac{\partial}{\partial r_\alpha} \frac{1}{R} = \frac{1}{2} \left(r_\alpha \frac{\partial}{\partial r_\alpha} \frac{1}{R} + r'_\alpha \frac{\partial}{\partial r'_\alpha} \frac{1}{R} \right) = -\frac{1}{R}$$

and

$$-\int_V r_\alpha \frac{\partial \phi}{\partial r_\alpha} \rho dV = \Omega, \quad (19.15)$$

where

$$\Omega = -\frac{G}{2} \int_V \int_{V'} \frac{\rho \rho'}{R} dV dV', \quad (19.16)$$

is the *gravitational energy* of the system. Obviously, the energy is negative.

Combining (19.5), (19.10), (19.12), and (19.15) into a single equation, we finally obtain

$$\begin{aligned} \frac{1}{2} \frac{d^2 I}{dt^2} = & 2U_{kin} + 3(\gamma_g - 1) U_{th} + \mathcal{M} + \Omega - \oint_S p (\mathbf{r} \cdot d\mathbf{S}) - \\ & - \oint_S \left[\frac{B^2}{8\pi} (\mathbf{r} \cdot d\mathbf{S}) - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{r}) (\mathbf{B} \cdot d\mathbf{S}) \right]. \end{aligned} \quad (19.17)$$

Formula (19.17) is called the *virial theorem* or, more exactly, the *scalar virial theorem*. It has repeatedly been used in astrophysics when discussing the question of the stability of equilibrium systems of various types. More exactly, this integral force balance relation is nothing more than a *necessary condition for equilibrium*. So it may be well used as a *non-existence theorem* for the equilibrium problem to find circumstances when non-equilibrium may occur.

19.1.3 Some Astrophysical Applications

The positive terms on the right-hand side of (19.17), in the virial theorem, lead to an increase in the moment of inertia I of an astrophysical system under consideration. It is no wonder that the kinetic energy U_{kin} or the thermal energy U_{th} tends to expand the system. The effect of magnetic field is more subtle. The magnetic field has tension along field lines and magnetic pressure. So we expect the overall average effect to be expansive. On the other hand, a negative term on the right-hand side, which is the gravitational energy Ω , tries to make the system more compact. Gravity is the only force which introduces a confining tendency in the system.

By way of illustration, let us consider some consequences of the virial theorem for the case of a *steady* system, i.e. when gravity balances the expansive forces so that

$$\frac{d^2 I}{dt^2} = 0. \quad (19.18)$$

Moreover let the kinetic energy of macroscopic motions be equal to zero

$$U_{kin} = 0, \quad (19.19)$$

i.e. the system is in *static* equilibrium. Both assumptions must be justified carefully, if they are applied to astrophysical plasma.

Let us suppose also that the system is finite and the surface S , over which the integration in formulae (19.10) and (19.12) is performed, can be moved sufficiently far away (formally speaking, to infinity), so that

$$\oint_S p (\mathbf{r} \cdot d\mathbf{S}) = 0 \quad (19.20)$$

and

$$\oint_S \left[\frac{B^2}{8\pi} (\mathbf{r} \cdot d\mathbf{S}) - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{r}) (\mathbf{B} \cdot d\mathbf{S}) \right] = 0. \quad (19.21)$$

Then from the virial theorem (19.17) it follows that

$$3(\gamma_g - 1) U_{th} + \mathcal{M} + \Omega = 0. \quad (19.22)$$

Introduce the “total” (without what has been neglected) energy of the system

$$\mathcal{E} = U_{th} + \mathcal{M} + \Omega. \quad (19.23)$$

Eliminating the thermal energy U_{th} from (19.22) and (19.23), the total energy is expressed as follows

$$\mathcal{E} = -\frac{(3\gamma_g - 4)}{3(\gamma_g - 1)} (|\Omega| - \mathcal{M}). \quad (19.24)$$

In a sense, the equilibrium is stable if $\mathcal{E} < 0$, i.e.

$$\frac{(3\gamma_g - 4)}{3(\gamma_g - 1)} (|\Omega| - \mathcal{M}) > 0, \quad (19.25)$$

which is equivalent, once $\gamma_g > 4/3$, to

$$|\Omega| > \mathcal{M}. \quad (19.26)$$

It is self-evident that inequality (19.26) is just a *necessary* condition for the dynamical *global* stability of a system. The condition is by no means sufficient. It can be used to show a non-existence of equilibrium of the system.

Let us consider *two particular cases of astrophysical interest*.

- (a) If $\mathcal{M} = 0$ then the system can be stable only for $\gamma_g > 4/3$. This condition is easy to understand. The pressure inside the system under adiabatic compression ($p \sim \rho^{\gamma_g}$) must grow faster than the gravitational pressure $p_g \sim \rho^{4/3}$. It is in this case that the system, for instance a star, can be sufficiently resilient to resist the gravitational collapse. That is why a star consisting of a monatomic gas (with $\gamma_g = 5/3$) can be dynamically stable.
- (b) Let $\mathcal{M} > 0$. Generally, the necessary condition for stability (19.25) can be, in principle, violated. What this means is that the field diminishes the stability of a star. Given a sufficiently strong field, gravitational attraction forces cannot balance the magnetic repulsion of the constituents of the system. However, such a situation is difficult to conceive.

In actuality, *gravitational compression* cannot result in $\mathcal{M} > |\Omega|$ since, given the freezing-in condition and isotropic compression, $p_{\text{mag}} \sim \rho^{4/3}$ in common with $p_g \sim \rho^{4/3}$. It is also impossible to obtain $\mathcal{M} > |\Omega|$ by dint of magnetic field amplification owing to *differential rotation*, since $|\Omega| > 2U_{\text{kin}}$ in a gravitationally bound system. On the other hand, the energy of a magnetic field generated by differential rotation must remain less than the kinetic energy U_{kin} of the rotation motion, i.e. $\mathcal{M} < U_{\text{kin}}$. Hence $\mathcal{M} < |\Omega|$.

At most, the condition $\mathcal{M} \sim |\Omega|$ can be realized. This situation is probably realized in stars of the *cold giant* type with a large radius. Perhaps such stars are at the limit of stability, which reveals itself as non-steady behavior.

In general, condition (19.26) allows us to evaluate the upper limit of the mean intensity of a magnetic field inside a star or other equilibrium configuration. Substitute the gravitational energy of a uniform ball,

$$\Omega = -\frac{3}{5} \frac{GM^2}{R}, \quad (19.27)$$

in (19.26). The result is (Syrovatskii 1957)

$$B < B_{cr} = 2 \times 10^8 \left(\frac{M}{M_\odot} \right) \left(\frac{R}{R_\odot} \right)^{-2}. \quad (19.28)$$

For the Sun, magnetic field B must be less than 2×10^8 G. For the most *magnetic stars* of the spectral class A, which are observed to have fields $\sim 10^4$ G, the condition $B < 3 \times 10^7$ G must hold. Hence these magnetic stars called the Ap stars, because they possess some peculiar properties (e.g., Hubrig et al., 2000), still are very far from the stability limit. As is seen from the Syrovatskii condition (19.28), the cold giants with large radii could be closer to such a limit.

Given a uniform field inside a star, on approaching the limit established by (19.28), the form of the star increasingly deviates from a sphere:

the magnetic field resists gravitational compression of a collapsing star in the direction perpendicular to the field, whereas the plasma may freely flow along the field lines.

As a result, the equilibrium configuration is represented by a rotation ellipsoid compressed in the field direction. The virial theorem can be written (e.g., Nakano 1998) for an axially-symmetric oblate magnetic cloud of mass M and semi-major axes $a_\perp$ and $a_\parallel$, respectively, embedded in a medium of pressure p_s . This is typical for the problem of star formation in magnetic clouds.

The action of a magnetic field is analogous to rotation (Strittmatter 1966). Furthermore, both the *strong field* and *fast rotation* are typical of *pulsars*, especially of the magnetars (see Exercise 14.2). So both these factors determine the real flattening of a neutron star. The flattening can be calculated using the *tensor* virial theorem. Note, however, that for a neutron star with $M \sim M_\odot$ and $R \sim 10$ km the critical magnetic field (19.28) is still unprecedentedly high: $B_{cr} \sim 10^{18}$ G. We call such fields *ultrastrong*.

Magnetars, or “magnetically powered neutron stars”, could form via a magnetic dynamo action in hot, nascent neutron stars if they are born spinning rapidly enough. Magnetism may be strong enough within these stars to evolve diffusively, driving internal heat dissipation that would keep the neutron stars hot and X-ray bright. Above a field strength of $\sim 10^{14}$ G, the evolving field inevitably induces stresses in the solid crust. Observations (e.g., [Feroci et al. 2001](#)) indicate that *giant flares*, involved a relativistic outflow of pairs and hard gamma rays, can plausibly be triggered by a large fracture in the crust of a neutron star with a field exceeding 10^{14} G. So the observed giant flares are presumably due to *local* magnetic instabilities in magnetars.

On the other hand, numerical studies ([Bocquet et al. 1995](#)) have confirmed that neutron stars with the ultrastrong internal magnetic fields are *globally* stable up to the order of 10^{18} G. They also have found that, for such values, the maximum mass of neutron stars increases by 13–29% relative to the maximum mass of non-magnetized neutron stars.

If ultrastrong fields exist in the interior of neutron stars, such fields will primarily affect the behavior of the residual charged particle. Moreover, contributions from the anomalous magnetic moment of the particles in a magnetic field should also be significant ([Broderick et al. 2000](#)). In particular, in a ultrastrong field, complete spin polarization of the neutrons occurs as a result of the interaction of the neutron magnetic moment with the magnetic field. The presence of a sufficiently strong field changes the ratio of protons to neutrons as well as the neutron drip density ([Suh and Mathews 2001](#)).

The virial theorem was applied in solar physics, for example, while studying active regions (see Sect. 19.4.3). It allows us to evaluate the energy of equilibrium electric currents and show that the energy can be large enough to explain the flaring activity ([Litvinenko and Somov 1991a](#)); see also discussion of the problem of the global MHD equilibria and filament eruptions in the solar corona ([Litvinenko and Somov 2001](#)).

19.2 Force-Free Fields and Shafranov’s Theorem

19.2.1 The Simplest Examples of Force-Free Fields

A particular case of equilibrium configurations of astrophysical plasma in a magnetic field is the *force-free field*, i.e. the field which does not require external forces. As was noted in Sect. 13.1, force-free fields naturally occur when the magnetic force dominates all the others, and hence the magnetic field must balance itself

$$\mathbf{B} \times \text{rot } \mathbf{B} = 0. \quad (19.29)$$

Let us consider several examples of such equilibrium configurations.

19.2.1.1 The Syrovatskii Force-Free Field

Let the magnetic field vector be situated in the plane parallel to the plane (x, y) , but depends only on z :

$$\mathbf{B} = \{ B_x(z), B_y(z), 0 \}. \quad (19.30)$$

Substitute (19.30) in (19.29):

$$\text{rot } \mathbf{B} = \left\{ -\frac{\partial B_y}{\partial z}, \frac{\partial B_x}{\partial z}, 0 \right\}, \quad (19.31)$$

$$\mathbf{B} \times \text{rot } \mathbf{B} = \left\{ 0, 0, B_x \frac{\partial B_x}{\partial z} + B_y \frac{\partial B_y}{\partial z} \right\} = 0. \quad (19.32)$$

The resulting equation is

$$\frac{\partial}{\partial z} (B_x^2 + B_y^2) = 0, \quad (19.33)$$

with the solution

$$B^2 = B_x^2 + B_y^2 = \text{const}. \quad (19.34)$$

This is the simplest example of a force-free field. The magnitude of the field vector is independent of z . A one-dimensional force-free field of this type may be considered to be a *local* approximation of an arbitrary force-free field in a region of the magnetic “*shear*” in the solar atmosphere. As a particular example, suitable for formal analysis, one may adopt the force-free field of the type

$$\mathbf{B} = \{ B_0 \cos kz, B_0 \sin kz, 0 \} \quad (19.35)$$

(Bobrova and Syrovatskii 1979). The field lines, and hence the electric current, lie in the plane (x, y) . The direction of the lines rotates with increasing z .


19.2.1.2 The Lundquist Force-Free Field

The magnetic field of a direct current flowing along the z axis tends to compress the plasma to the axis, owing to the tension of the field lines (see Sect. 13.1.3). By contrast,

■ a bundle of parallel field lines tends to expand by the action of the magnetic pressure gradient.

Given the superposition of these fields for a certain relationship between them, the total magnetic force can be zero. Field lines for such a force-free field have the shape of spirals shown in Fig. 19.1.

Fig. 19.1 A helical magnetic field in the form of a spiral of constant slope on a cylindrical surface $r_{\perp} = \text{const}$



The corresponding *axially symmetric* solution to (19.29) in the cylindrical coordinates $r_{\perp}$, ϕ , z is of the form (Lundquist 1951):

$$B_z = A J_0(\alpha r_{\perp}), \quad B_{\phi} = A J_1(\alpha r_{\perp}), \quad B_r = 0. \quad (19.36)$$

Here J_0 and J_1 are the Bessel functions, A and α are constants.

A distinguishing feature of the field is that $B^2 \sim r_{\perp}^{-1}$ for large $r_{\perp}$ since Bessel functions $J_n \sim r_{\perp}^{-1/2}$ as $r_{\perp} \rightarrow \infty$ ($n = 0, 1$). The magnetic energy

$$\mathcal{M} = \int \frac{B^2}{8\pi} dV \sim r_{\perp}^{-1} r_{\perp}^2 \sim r_{\perp} \quad (19.37)$$

diverges for large $r_{\perp}$. Such a *divergence of magnetic energy* is known to be typical of force-free fields and will be explained below.

19.2.2 The Energy of a Force-Free Field

Let us retain only magnetic terms in the virial theorem; we have

$$\mathcal{M} - \oint_S \left[\frac{B^2}{8\pi} (\mathbf{r} \cdot d\mathbf{S}) - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{r}) (\mathbf{B} \cdot d\mathbf{S}) \right] = 0. \quad (19.38)$$

Provided the electric currents occupy a *finite* region, the value of the magnetic field is proportional to r^{-3} (or higher degrees of r^{-1}). Once the surface of integration S is expanded to infinity, the surface integral tends to zero. Equality (19.38) becomes impossible.

Therefore any finite magnetic field cannot contain itself. There must be *external* forces to balance the outwardly directed pressure due to the *total magnetic* energy $\mathcal{M}$.

The same statement may be formulated as follows. *The force-free field cannot be created in the whole space.* This is the so-called *Shafranov theorem* (Shafranov 1966). While stresses may be eliminated in a given region V , they cannot be canceled everywhere. In general

■ a force-free configuration requires the forces needed to balance the outward pressure of the magnetic field to be reduced in magnitude by spreading them out over the bounding surface S .

In this way, the virial theorem sets limits on the space volume V that can be force-free.

The Shafranov theorem is the counterpart of the known Irnshaw theorem (see Sivukhin 1996 Chap. 1, Sect. 9) concerning the equilibrium configuration of a system of electric charges. Such a configuration also can be stable only in the case that some external forces, other than the electric ones, act in the system.

In fact, Shafranov (1966) has proved a stronger statement than the above theorem on the force-free field. He has taken into account not only the terms corresponding to the magnetic force in (19.17) but the gas pressure as well:

$$\begin{aligned} \int_V \left(3p + \frac{B^2}{8\pi} \right) dV &= \\ &= \oint_S \left[\left(p + \frac{B^2}{8\pi} \right) (\mathbf{r} \cdot d\mathbf{S}) - \frac{1}{4\pi} (\mathbf{B} \cdot \mathbf{r}) (\mathbf{B} \cdot d\mathbf{S}) \right]. \end{aligned} \quad (19.39)$$

If the plasma occupies some finite volume V , the pressure outside of this volume being zero, and if electric currents occupy a finite region, then the surface integral tends to zero, once the surface of integration, S , is expanded to infinity. On the other hand, the expression under the integral sign on the left-hand side is always positive. Hence the integral is positive. Thus the equality (19.39) turns out to be impossible. Therefore

■ any finite equilibrium configuration of a plasma with a magnetic field can exist only in the presence of external forces which, *apart from the gas pressure*, serve to fix the electric currents.

In a laboratory, fixed current conductors must be present. In this case the right-hand side of (19.39) is reduced to the integral over the surface of the conductors.

Under astrophysical conditions, the role of the external force is frequently played by the gravitational force or by an external magnetic field having its sources outside the volume under investigation. However these sources must be kept and driven by non-magnetic forces.

A typical example of such a situation is the magnetic field of an active region on the Sun, above the photosphere. This is the sum of the proper field created by currents flowing inside the active region, and the external field with the sources

situated (and fixed) below the photosphere (Litvinenko and Somov 1991a). In this case the formula for the magnetic energy of the equilibrium system contains a term due to the interaction of internal currents (in particular, current layers in the regions of reconnection, i.e. the *reconnecting current layers*) with the external magnetic field.

19.3 Properties of Equilibrium Configurations

19.3.1 Magnetic Surfaces

Let us consider the case of *magnetostatic* equilibrium

$$-\nabla p + \frac{1}{4\pi} \operatorname{rot} \mathbf{B} \times \mathbf{B} - \rho \nabla \phi = 0. \quad (19.40)$$

The gravitational force is supposed to be negligible:

$$\rho \nabla \phi = 0. \quad (19.41)$$

On dropping the third term in (19.40) and taking the scalar product with vector $\mathbf{B}$ we obtain

$\mathbf{B} \cdot \nabla p = 0,$

(19.42)

i.e. magnetic field lines in an equilibrium configuration are situated on the surface $p = \text{const}$. Therefore

in order to contain a plasma by the magnetic field, the field lines are forbidden to leave the volume occupied by the plasma.

There is a common viewpoint that, by virtue of the condition

$$\operatorname{div} \mathbf{B} = 0, \quad (19.43)$$

field lines may either close or go to infinity. However the other variant is possible, when a field line fills up an entire surface—*magnetic surface*.

Let us consider the field of two electric currents—a line current J_1 flowing along the vertical z axis (Fig. 19.2a) and a plane current ring J_2 (see Tamm 1989 Chap. 4, Sect. 53). If there were only the current J_1 , the field lines of this current $\mathbf{B}_1$ would constitute circumferences centred at the z axis. The field lines $\mathbf{B}_2$ of the ring current J_2 lie in meridional planes. The total field $\mathbf{B} = \mathbf{B}_1 + \mathbf{B}_2$ forms a helical line




Fig. 19.2 (a) A line current J_1 and a ring current J_2 . (b) The field lines of the total field $B_1 + B_2$ form a toroidal surface S

on a toroidal surface S . The course of this spiral depends on the ratio B_1/B_2 . Once this is a rational number, the spiral will close. However, in general, it does not close but continuously fills up the entire toroidal surface S (Fig. 19.2b).

By virtue of condition (19.42), the plasma pressure at such a surface (called the magnetic one) is constant. Such a magnetic field can serve as a trap for the plasma. This fact constitutes the basis for constructing laboratory devices for plasma containment in stellarators, suggested by Spitzer.

Take the scalar product of (19.40), without the gravitational force, with the electric current vector

$$\mathbf{j} = \frac{c}{4\pi} \text{rot } \mathbf{B}. \quad (19.44)$$

The result is

$$\mathbf{j} \cdot \nabla p = 0$$

(19.45)

which, in combination with (19.42), signifies that,

in a pressure-balanced magnetic field configuration, the electric current also flows on magnetic surfaces (Fig. 19.2b).

In general, magnetic fields do not form magnetic surfaces. Such surfaces arise in magnetohydrostatic equilibria and for some highly symmetric field configurations. In the case of the latter, (14.19) for the magnetic field lines admit an exact integral which is the equation for the magnetic surface.

19.3.2 The Specific Volume of a Magnetic Tube

Let us consider two closed magnetic surfaces: $p = \text{const}$ and $p + dp = \text{const}$. Construct a system of non-crossing partitions between them (Fig. 19.3). Let $d\mathbf{l}_1$ be the line element directed normally to the surface $p = \text{const}$:




Fig. 19.3 The calculation of the electric current between two magnetic surfaces

$$d\mathbf{l}_1 = \frac{\nabla p}{|\nabla p|^2} dp. \quad (19.46)$$

The vectors $d\mathbf{l}_2$ and $d\mathbf{l}_3$ are directed along the two independent contours l_2 and l_3 which may be drawn on a toroidal surface: for example the curve l_2 is directed along a large circle of the toroid while l_3 lies along the small one. The surface element of this partition is

$$d\mathbf{S}_3 = d\mathbf{l}_1 \times d\mathbf{l}_2. \quad (19.47)$$

The total current dJ_3 flowing through the partition situated on the contour l_2 is

$$dJ_3 = \oint_{l_2} \mathbf{j} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2). \quad (19.48)$$

According to (19.45), the total current flowing through the system of non-crossing partitions between the two magnetic surfaces is constant. In other words, dJ_3 is independent of the choice of the integration contour. We are concerned with the physical consequences of this fact.

In order to find the expression for the current density $\mathbf{j}$ in an equilibrium configuration, take the vector product of (19.40) with the magnetic field $\mathbf{B}$. The result is

$$\mathbf{B} \times \nabla p = \frac{1}{c} \mathbf{B} \times (\mathbf{j} \times \mathbf{B}),$$

which, on applying the formula for a double vector product to the right-hand side, becomes

$$c \mathbf{B} \times \nabla p = \mathbf{j} B^2 - \mathbf{B} (\mathbf{j} \times \mathbf{B}).$$

Thus we have

$$\mathbf{j} = c \frac{\mathbf{B} \times \nabla p}{B^2} + f \mathbf{B}, \quad (19.49)$$

where $f = f(\mathbf{r})$ is an arbitrary function. If need be, it can be found from the condition $\text{div } \mathbf{j} = 0$.

Substitute (19.49) in the integral (19.48). The last takes the following form (see Exercise 19.4):

$$dJ_3 = -c \, dp \oint_{l_2} \frac{\mathbf{B} \cdot d\mathbf{l}_2}{B^2} + \oint_{l_2} f(\mathbf{r}) \mathbf{B} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2). \quad (19.50)$$

Provided the contour l_2 coincides with a closed field line, the vector

$$d\mathbf{l}_2 = \frac{\mathbf{B}}{B} dl,$$

and, therefore, the second term on the right-hand side of Formula (19.50) vanishes.

Once a magnetic field line closes on making one circuit of the toroid, the expression

$$dJ_n = -c \, dp \oint \frac{dl}{B} \quad (19.51)$$

defines the total current flowing between neighbouring magnetic surfaces normal (the subscript n) to the field line. Since the magnitude of this current is independent of the choice of contour, for each field line on a magnetic surface the integral

$$\boxed{U = \oint \frac{dl}{B}} \quad (19.52)$$

is constant. The condition of constancy of U can be generalized to include the surface with unclosed field lines (Shafranov 1966). Thus (Kadomtsev 1966),

under the condition of magnetostatic equilibrium, the magnetic surface consists of the field lines with the same value of U .

Let us introduce the notion of the *specific volume* of a magnetic tube (Rosenbluth and Longmire 1957) or simply the *specific magnetic volume* as the ratio of its geometric volume dV to the magnetic flux $d\Phi$ through the tube. If dS_n is the cross-sectional surface of the tube, its geometric volume is

$$dV = \oint dS_n dl$$

whereas the magnetic flux

$$d\Phi = B dS_n.$$

On the basis of the magnetic flux constancy inside the tube of field lines, i.e. $d\Phi = \text{const}$, we deduce that

$$\frac{dV}{d\Phi} = \oint \frac{dS_n}{B dS_n} dl = \oint \frac{dl}{B} = U. \quad (19.53)$$

The stability of an equilibrium MHD configuration can be judged by the condition (19.52). This property will be discussed in the next section.

19.3.3 The Flute or Convective Instability

Much like any gas with a finite temperature, the plasma in a magnetic field tends to expand. However, given a high conductivity, it cannot move independently of the magnetic field. The plasma moves together with the field lines in such a way that it travels to a region of the field characterized by a greater specific volume.

In order for an equilibrium configuration to be stable with respect to a given perturbation type—deformation of a tube of magnetic field lines—the following condition is necessary (Rosenbluth and Longmire 1957):

$$\delta U = \delta \oint \frac{dl}{B} < 0. \quad (19.54)$$

To put it another way,

the magnetostatic equilibrium is stable once the given type of deformation does not facilitate the plasma spreading,

i.e. increasing its specific volume.

As an example, let us consider the plasma in the magnetic field of a linear current J :

$$B_\varphi = \frac{2J}{cr}, \quad (19.55)$$

here r, z, φ are the cylindrical coordinates. In such a field there exists an equilibrium plasma configuration in the form of an infinite hollow cylinder C as shown in Fig. 19.4a.

Let us calculate the specific volume for such a configuration. The geometric volume of the tube of field lines is

$$dV = 2\pi r dr dz,$$

whereas the magnetic flux

$$d\Phi = B_\varphi dr dz = \frac{2J}{cr} dr dz.$$




Fig. 19.4 (a) An equilibrium plasma configuration. (b) Unstable perturbations of the outer boundary

Hence the specific volume

$$U = \frac{dV}{d\Phi} = \frac{\pi c}{J} r^2. \quad (19.56)$$

It is seen from (19.56) that the specific volume grows with the radius. In particular, for small perturbations δr of the external surface S of the plasma cylinder C

$$\delta U = \frac{2\pi c}{J} r \delta r > 0$$

once $\delta r > 0$. It is sufficient to have a small perturbation of the external boundary of the plasma to obtain ring flutes which will rapidly grow towards the wall W of the chamber as shown in Fig. 19.4b.

19.3.4 Stability of an Equilibrium Configuration

The problems of plasma *equilibrium and stability* are of great value for plasma astrophysics as a whole (Zel'dovich and Novikov 1971; Chandrasekhar 1981), and especially for solar physics (Parker 1979; Priest 1982).

■ The Sun seems to maintain stability of solar prominences and coronal loops with great ease

(Tandberg-Hanssen 1995; Acton 1996) in contrast to the immense difficulty of containing plasmas in a laboratory.

Therefore, sometimes, we need to explain how an equilibrium can remain stable for a very long time. This is, for example, the case of *reconnecting current layers* (RCLs) in the solar atmosphere and the *geomagnetic tail* (see Part II, Sects. 8.2 and 11.6.3). At other times, we want to understand

why magnetic structures on the Sun suddenly become unstable and produce dynamic events

of great beauty such as eruptive prominences, solar flares and coronal mass ejections (CMEs).

The methods employed to investigate the stability of an equilibrium MHD system are natural generalizations of those for studying a particle in one-dimensional motion. One approach is to seek normal mode solutions as we did it in Chap. 15.

An alternative approach for tackling stability is to consider the change in potential energy due to a displacement from equilibrium. The main property of a stable equilibrium is that it is at the minimum of the potential energy. So any perturbations around the equilibrium ought to increase the total potential energy. Hence, in order to determine if an equilibrium is stable, one finds out if *all types* of perturbations increase the potential energy of the system (Bernstein et al. 1958).

Recommended Reading: Morozov and Solov'ev (1966a), Kadomtsev (1960, 1966), Shu (1992).

19.4 The Archimedean Force in MHD

19.4.1 A General Formulation of the Problem

Now we return to the equation of magnetostatic equilibrium (19.40). Let us rewrite it as follows:

$$\nabla p = \rho \mathbf{g} + \mathbf{f} \quad (19.57)$$

where

$$\mathbf{f} = \frac{1}{c} \mathbf{j} \times \mathbf{B} \quad (19.58)$$


is the Lorentz force, $\mathbf{g} = -g \mathbf{e}_z$ is the gravity acceleration.

We begin by considering an incompressible conducting fluid situated in an uniform magnetic field $\mathbf{B}_0$ and electric field $\mathbf{E}_0$ as illustrated by Fig. 19.5. Provided the current $\mathbf{j}_0$ flowing in the fluid is *uniform*, the Lorentz force created is uniform as well:

$$\mathbf{f}_0 = \frac{1}{c} \mathbf{j}_0 \times \mathbf{B}_0 \quad (19.59)$$

By virtue of (19.57), *the Lorentz force makes the fluid heavier or lighter*. In both cases the uniform volume force is potential and, much like the gravity force, will be

Fig. 19.5 Formulation of the problem concerning the Archimedean force in magnetohydrodynamics (see Somov 1994b)



balanced by an additional pressure gradient appearing in the fluid. As will be shown later, that allows the creation of a regulated *expulsion* force (Fig. 19.5) analogous to the Archimedean force in ordinary hydrodynamics.

A body plunged into the fluid is acted upon by the force

$$\mathbf{F} = \int_V (\rho_1 \mathbf{g} + \mathbf{f}_1) dV + \oint_S p_0 \mathbf{n} dS. \quad (19.60)$$

Here ρ_1 is the density of the submerged body, which is generally not equal to that of the fluid ρ_0 ;

$$\mathbf{f}_1 = \frac{1}{c} \mathbf{j}_1 \times \mathbf{B}_0 \quad (19.61)$$

is the volume Lorentz force, $\mathbf{j}_1$ is the current inside the body, $\mathbf{n}$ is the inward normal to the surface S , and p_0 is the pressure on the body from the fluid, resulted from (19.57):

$$\nabla p_0 = \rho_0 \mathbf{g} + \mathbf{f}_0. \quad (19.62)$$

19.4.2 A Simplified Consideration of the Effect

If the current $\mathbf{j}_0$ was uniform, the right-hand side of (19.62) would be a uniform force, and formula (19.60) could be rewritten as

$$\mathbf{F} = \int_V (\rho_1 \mathbf{g} + \mathbf{f}_1) dV - \int_V \nabla p_0 dV \quad (19.63)$$




Fig. 19.6 Opposite orientation of the σ -dependent force in two opposite cases: (a) $\sigma_1 > \sigma_0$ and (b) $\sigma_1 < \sigma_0$. Appearance of a non-uniform distribution of electric current is shown

or

$$\mathbf{F} = \int_V (\rho_1 - \rho_0) \mathbf{g} dV + \frac{1}{c} \int_V (\mathbf{j}_1 - \mathbf{j}_0) \times \mathbf{B}_0 dV. \quad (19.64)$$

The first term in (19.64) corresponds to the usual Archimedean force in hydrodynamics. It equals zero once $\rho_1 = \rho_0$. When $\rho_1 > \rho_0$, the direction of this force coincides with the gravitational acceleration $\mathbf{g}$. The second term describes the *magnetic expulsion* force. It vanishes once $\mathbf{j}_1 = \mathbf{j}_0$, i.e. $\sigma_1 = \sigma_0$.

The second term in formula (19.64) shows that the magnetic expulsion force, different from the known Parker's magnetic buoyancy force (see Chap. 8 in [Parker 1979](#)) by its origin, appears provided $\sigma_1 \neq \sigma_0$. This fact has been used to construct, for example, MHD devices for the separation of mechanical mixtures. In what follows we shall call the second term in (19.64) the magnetic σ -dependent force:

$$\mathbf{F}_\sigma = \frac{1}{c} \int_V (\sigma_1 - \sigma_0) \mathbf{E}_0 \times \mathbf{B}_0 dV. \quad (19.65)$$

Note, however, that the simplest formula (19.64) is of purely illustrative value since *the electric field and current density are not uniform in the presence of a body with conductivity σ_1 which is not equal to that of the fluid σ_0* (Fig. 19.6). In this case, the appearing σ -dependent force is generally not potential. Hence it cannot be balanced by potential forces. That is the reason why

the magnetic σ -dependent force generates MHD vortex flows of the conducting fluid.

The general analysis of the corresponding MHD problem was made by [Andres et al. \(1963\)](#). The stationary solutions for a ball and a cylinder were obtained by [Syrovatskii and Chesalin \(1963\)](#) for the specific case when both the magnetic and usual Reynolds numbers are small; similar stationary solutions for a cylinder see also in [Marty and Alemany \(1983\)](#), [Gerbeth et al. \(1990\)](#). The character of the MHD vortex flows and the forces acting on submerged bodies will be analyzed in Sects. 20.3 and 20.4.

19.4.3 MHD Equilibrium in the Solar Atmosphere

The magnetic configuration in an active region in the solar atmosphere is, in general, very complex and modeling of dynamical processes in these regions requires a high degree of idealization. First, as regards the most powerful and fascinating of these processes, the *two-ribbon flare*, the typical pre-flare magnetic field distribution seems to conform to a certain standard picture: a magnetic arcade including a more or less pronounced plage filament, prominence. Second, instead of dynamics, models often deal with a static or steady-state equilibrium in order to understand the causes of a flare or another transient activity in the solar atmosphere as a result of some MHD instability or lack of equilibrium.

So it is assumed that initially the configuration of prominence and overlying arcade is in equilibrium but later the eruption takes place.

Either the MHD equilibrium of solar plasma has become unstable or the equilibrium has been lost.

One limiting possibility is that the magnetic field around the prominence evolves into an unstable or non-equilibrium configuration and then drives the overlying magnetic arcade. However observations imply that this is unlikely. An alternative is that the overlying arcade evolves until it is no longer in stable equilibrium and then its eruption stimulates the prominence to erupt by removing stabilizing field lines. Presumably this is the case of a coronal loop transient and coronal mass ejection (CME).

The idealized models used in theoretical and numerical studies of this problem usually consider two-dimensional *force-free* arcade configurations with foot points anchored in the photosphere which are energized, for example, by photospheric shear flows in the direction along the arcade (see [Biskamp and Welter 1989](#)). Some other models take into account the gas pressure gradient and the gravitational force ([Webb 1986](#)).

However it is important to investigate more general circumstances when equilibrium and non-equilibrium may occur. The electromagnetic expulsion force—a MHD analogue of the usual Archimedean force—can play an important part in the dynamics of coronal plasma with a non-uniform distribution of temperature and, hence, electric conductivity. More exactly,

the condensation mode of the radiatively-driven *thermal instability* in an active region may result in the formation of cold dense loops or filaments surrounded by hot rarified plasma

(see [Somov 1992](#)). This well-known effect results in a great difference of electric conductivities outside and inside the filaments. Therefore the conductivity-dependent force can generate vortex flows (see Sect. 20.4) inside and in the vicinity of the filaments as well as initiate the non-equilibrium responsible for transient activities: flares, CMEs etc.

The virial theorem confirms this possibility and clarifies the role of pre-flare reconnecting current layers in MHD equilibrium and non-equilibrium of an active

region. Correct use of the virial theorem confirms the applicability of reconnection in current layers for explaining the energetics of solar flares (Litvinenko and Somov 1991a, 2001) and other non-steady phenomena in the solar atmosphere.

19.5 Grad-Shafranov Equation

The famous Grad-Shafranov equation was formulated in connection with the classical problem of controlled thermonuclear fusion in the late 1950s (Grad and Rubin 1958; Shafranov 1958; Grad 1960). The equation was originally related to the static equilibrium of ideal MHD configurations (Sect. 13.1.3). Thus we start from two equations:

$$-\nabla p + \frac{1}{4\pi} \text{rot } \mathbf{B} \times \mathbf{B} = 0 \quad (19.66)$$

and

$$\text{div } \mathbf{B} = 0 \quad (19.67)$$

Both of them are the equations for the magnetic-field vector $\mathbf{B}$ but, in addition, the former is a relationship between the magnetic field and unknown pressure p of plasma. We would like to see what follows from this relationship.

At first, let us rewrite (19.66) as

$$\nabla p = \frac{1}{c} \mathbf{j} \times \mathbf{B}, \quad (19.68)$$

where the electric current

$$\mathbf{j} = \frac{c}{4\pi} \text{rot } \mathbf{B}. \quad (19.69)$$

By taking the scalar product of equation (19.68) with vector $\mathbf{B}$ and with vector $\mathbf{j}$ we find two conditions:

$$\mathbf{B} \cdot \nabla p = 0 \quad \text{and} \quad \mathbf{j} \cdot \nabla p = 0, \quad (19.70)$$

respectively. Thus the magnetic field lines and the electric current lines lay on surfaces

$$p(\mathbf{r}) = \text{const}, \quad (19.71)$$

that are called the *magnetic surfaces*. Recall that we already derived this property (*Property I*) from the same equations in Sect. 19.3.1.

Now let us illustrate another property (*Property II*) of a pressure-balanced magnetic field by a simple example. We assume that a system is the 2D one with z as the invariant axis: all variables are independent of z . Therefore $\partial/\partial z = 0$. Then the magnetic field can be written in cartesian coordinates as

$$\mathbf{B}(x, y) = \text{rot } \mathbf{A} + B_z(x, y) \mathbf{e}_z, \quad (19.72)$$

where the vector potential

$$A(x, y) = \{0, 0, A(x, y)\} . \quad (19.73)$$

Thus the magnetic field

$$\mathbf{B}(x, y) = \left\{ \frac{\partial A}{\partial y}, -\frac{\partial A}{\partial x}, B_z(x, y) \right\} . \quad (19.74)$$

With this definition and Property I it follows from (19.66) that

$$p + \frac{B_z^2}{8\pi} = f(A) , \quad (19.75)$$

where $f(A)$ is an arbitrary function of scalar potential A (see, e.g., [Parker 1979](#); Sect. 6.2).

If $\nabla A \neq 0$ then it also follows from (19.66) that

$$\frac{\partial^2 A}{\partial x^2} + \frac{\partial^2 A}{\partial y^2} + 4\pi \frac{d f}{d A} = 0 . \quad (19.76)$$

This is well known as the Grad-Shafranov equation. More exactly, the Grad-Shafranov method was mainly addressed to the ideal MHD 2D equilibrium, mainly the axisymmetric toroidal plasma in a tokamak. However it is also often used in plasma astrophysics.

19.6 Practice: Exercises and Answers

Exercise 19.1 (Sect. 19.2.1). Show that, apart from the trivial case of a potential field, the magnetic fields for which

$$\text{rot } \mathbf{B} = \alpha \mathbf{B} \quad (19.77)$$

will be force-free. In the most general case, α will be spatially dependent.

Answer. Just substitute formula (19.77) in (19.29).

Exercise 19.2 (Sect. 19.2.2). Show that the force-free fields with $\alpha = \text{const}$ represent the state of *minimal* magnetic energy in a closed system ([Woltjer 1958](#)).

Hint. First, let us assume perfect conductivity and rewrite the freezing-in equation (12.73) by using $\mathbf{B} = \text{rot } \mathbf{A}$ as follows

$$\frac{\partial \mathbf{A}}{\partial t} = \mathbf{v} \times (\nabla \times \mathbf{A}) . \quad (19.78)$$

Here $\mathbf{A}$ is the vector potential. Using (19.78), show that

$$\mathcal{H} = \int_V \mathbf{A} \cdot (\nabla \times \mathbf{A}) dV = \text{const}$$

(19.79)

for all $\mathbf{A}$ which are constant on the boundary S of the region V . The integral $\mathcal{H}$ is called the *global magnetic helicity* of the closed system under consideration (for more detail see Part II, Sect. 12.1.1).

Second, examine the stationary values of the magnetic energy

$$\mathcal{M} = \int_V \frac{B^2}{8\pi} dV = \int_V \frac{1}{8\pi} (\text{rot } \mathbf{A})^2 dV. \quad (19.80)$$

Introduce a Lagrangian multiplier $\alpha/8\pi$ and obtain the following condition for stationary values

$$\delta \int_V \left[(\text{rot } \mathbf{A})^2 - \alpha \mathbf{A} \cdot \text{rot } \mathbf{A} \right] dV = 0. \quad (19.81)$$

Performing the variation, (19.77) follows with $\alpha = \text{const}$. Such fields are called *linear force-free fields*.

Exercise 19.3 (Sect. 19.4.3). The highly-conductive plasma in the solar corona can support an electric field $\mathbf{E}_{\parallel}$ if $E_{\parallel} \ll E_{\text{Dr}}$ where E_{Dr} is the Dreicer field (8.75). In the corona $E_{\text{Dr}} \approx 7 \times 10^{-6} \text{ V cm}^{-1}$ (Exercise 8.4). Evaluate the characteristic values of the magnetic field B and the velocity v of plasma motions in the corona which allow us to consider an equilibrium of moving plasma in the corona as a *force-free* one.

Answer. Let us evaluate an electric field as the electric field related to a motion of magnetic field lines in the corona

$$E_{\parallel} \approx E \approx \frac{1}{c} v B \approx 10^{-8} v (\text{cm s}^{-1}) B (\text{G}), \quad \text{V cm}^{-1}. \quad (19.82)$$

From the condition that this field must be much smaller than the Dreicer field we find that

$$v (\text{cm s}^{-1}) B (\text{G}) \ll 10^8 E_{\text{Dr}} \approx 7 \times 10^2. \quad (19.83)$$

So, with the magnetic field in the corona $B \sim 100 \text{ G}$, the plasma motion velocity must be very small: $v \ll 10 \text{ cm s}^{-1}$. Hence, if the electric fields that are parallel to the magnetic field lines have the same order of magnitude as the perpendicular electric fields, the solar corona hardly can remain force-free with ordinary collisional conductivity because of the motion of magnetic field lines.

The electric runaway effects (Sect. 8.4.2) can become important even at very slow motions of the field lines in the corona. The *minimum current corona* (see Part II, Sects. 3.3.1 and 3.4.3) seems to be a more realistic approximation everywhere except the strongly-twisted magnetic-flux tubes.

Exercise 19.4 (Sect. 19.3.2). Derive formula (19.50) for the total electric current flowing through the system of non-crossing partitions between two magnetic surfaces.

Answer. Substitute the electric current density (19.49) in the integral (19.48):

$$\mathbf{j} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2) = \frac{c}{B^2} (\mathbf{B} \times \nabla p) \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2) + f \mathbf{B} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2). \quad (19.84)$$

Let us rearrange the first item, using the well-known *Lagrange identity* in vector analysis (e.g., Smirnov 1965):

$$(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{b} \cdot \mathbf{c})(\mathbf{a} \cdot \mathbf{d}).$$

We get

$$(\mathbf{B} \times \nabla p) \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2) = (\mathbf{B} \cdot d\mathbf{l}_1)(\nabla p \cdot d\mathbf{l}_2) - (\mathbf{B} \cdot d\mathbf{l}_2)(\nabla p \cdot d\mathbf{l}_1).$$

By virtue of (19.42) and (19.46),

$$(\mathbf{B} \cdot d\mathbf{l}_1) = (\mathbf{B} \cdot \nabla p) \frac{dp}{|\nabla p|^2} = 0, \quad (\nabla p \cdot d\mathbf{l}_1) = dp.$$

Hence

$$(\mathbf{B} \times \nabla p) \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2) = -(\mathbf{B} \cdot d\mathbf{l}_2) dp. \quad (19.85)$$

Substitute (19.85) in (19.84):

$$\mathbf{j} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2) = -c \frac{dp}{B^2} (\mathbf{B} \cdot d\mathbf{l}_2) + f(\mathbf{r}) \mathbf{B} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2).$$

Thus the expression (19.48) for current dJ_3 takes the form

$$dJ_3 = -c dp \oint_{l_2} \frac{\mathbf{B} \cdot d\mathbf{l}_2}{B^2} + \oint_{l_2} f(\mathbf{r}) \mathbf{B} \cdot (d\mathbf{l}_1 \times d\mathbf{l}_2), \quad (19.86)$$

q.e.d.

Chapter 20

Stationary Flows in a Magnetic Field

Abstract There exist two different sorts of stationary MHD flows depending on whether or not a plasma can be considered as ideal or non-ideal medium. Both cases have interesting applications in modern astrophysics.

20.1 Ideal Plasma Flows

Stationary motions of an ideal conducting medium in a magnetic field are subject to the following set of MHD equations (cf. (12.69)):

$$(\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{1}{\rho} \nabla \left(p + \frac{B^2}{8\pi} \right) + \frac{1}{4\pi\rho} (\mathbf{B} \cdot \nabla) \mathbf{B}, \quad (20.1)$$

$$\text{rot} (\mathbf{v} \times \mathbf{B}) = 0, \quad (20.2)$$

$$\text{div } \rho \mathbf{v} = 0, \quad (20.3)$$

$$\text{div } \mathbf{B} = 0, \quad (20.4)$$

$$(\mathbf{v} \cdot \nabla) s = 0, \quad (20.5)$$

$$p = p(\rho, s). \quad (20.6)$$

The induction equation (20.2) is satisfied identically, provided the motion of the medium occurs along the magnetic field lines, i.e.

$\mathbf{v} \parallel \mathbf{B}.$

(20.7)

20.1.1 Incompressible Medium

In the case of an incompressible fluid ($\rho = \text{const}$) (20.1)–(20.6) have the general solution Syrovatskii (1956, 1957):

$$\mathbf{v} = \pm \frac{\mathbf{B}}{\sqrt{4\pi\rho}}, \quad (20.8)$$

$$\nabla \left(p + \frac{B^2}{8\pi} \right) = 0. \quad (20.9)$$

Here $\mathbf{B}$ is an arbitrary magnetic field: the form of the field lines is unimportant, once condition (20.4) holds. A conducting fluid flows parallel or anti-parallel to the magnetic field. We shall learn more about such equilibrium flows later on.

It follows from (20.8) that

$$\frac{\rho v^2}{2} = \frac{B^2}{8\pi}, \quad (20.10)$$

while (20.9) gives

$$p + \frac{B^2}{8\pi} = \text{const}. \quad (20.11)$$

For the considered class of plasma motions along the field lines, the equipartition of energy between that of the magnetic field and the kinetic energy of the medium takes place, whereas the sum of the gas pressure and the magnetic pressure is everywhere constant.

The existence of the indicated solution means that

an arbitrary magnetic field and an ideal incompressible medium in motion are in MHD equilibrium, provided the motion of the medium occurs with the Alfvén speed along magnetic field lines.


Stationary flows of this type can be continuous in the whole space as well as discontinuous at some surfaces. For example, the solution (20.10) and (20.11) can be realized as a stream or non-relativistic jet of an arbitrary form, flowing in an immovable medium without a magnetic field.

Note that the tangential discontinuity at the boundary of such a jet is *stable*, since, by virtue of (20.10), the condition (16.38) by Syrovatskii is valid:

$$\frac{B^2}{8\pi} > \frac{1}{4} \frac{\rho v^2}{2}. \quad (20.12)$$

Such stable stationary jets of an incompressible fluid can close in *rings* and *loops* of an arbitrary type.

Fig. 20.1 Rotational (a) and helical (b) stationary flows of a compressible plasma



20.1.2 Compressible Medium

In a compressible plasma ($\rho \neq \text{const}$) the stationary solution (20.8) is still possible, once the density of the plasma does not change along the field lines:

$$\mathbf{B} \cdot \nabla \rho = 0. \quad (20.13)$$

Obviously, this condition is necessary but not sufficient. On substituting the solution (20.8) in (20.3), we have

$$\text{div } \rho \mathbf{v} = \pm \frac{1}{\sqrt{4\pi}} \left[\frac{1}{\sqrt{\rho}} \text{div } \mathbf{B} - \frac{1}{2} \rho^{-3/2} \mathbf{B} \cdot \nabla \rho \right] = 0$$


by virtue of (20.4) and (20.13). Thus the condition (20.13) is enough for (20.3) to be satisfied identically. However, in order to ensure the fulfilment of condition (20.9), we must require constancy of the gas and the magnetic pressure or the absolute value of the magnetic field intensity. The latter means that each magnetic flux tube must have a *constant cross-section*. Hence, by virtue of (20.8), the flow velocity along the flux tube will be constant as well.

Therefore stationary flows corresponding to the solutions (20.8) and (20.9), which are flows with a *constant velocity* in magnetic tubes of a constant cross-section, are possible in a compressible medium. An example of such a flow is the plasma rotation in a ring tube (Fig. 20.1a). We can envisage spiral motions of the plasma, belonging to the same type of stationary solutions in MHD (Fig. 20.1b). This may be, for example, the case of an astrophysical jet when plasma presumably moves along a spiral trajectory.

20.1.3 Astrophysical Collimated Streams (Jets)

Powerful extragalactic radio sources comprise two extended regions containing magnetic field and synchrotron-emitting relativistic electrons, each linked by a jet to a central compact radio source located in the nucleus of the associated active galaxy

Fig. 20.2 A MHD wave of arbitrary amplitude



Begelman et al. (1984). These jets are well *collimated streams of plasma* that emerge from the nucleus in opposite directions, along which flow mass, momentum, energy, and magnetic flux. The oscillations of jets about their mean directions are observed. The origin of the jet is crucial to understanding all active nuclei (Sect. 13.3).

The *microquasars* recently discovered in our Galaxy offer a unique opportunity for a deep insight into the physical processes in relativistic jets observed in different source populations Mirabel and Rodriguez (1998); Atoyan and Aharonian (1999). Microquasars are stellar-mass black holes in our Galaxy that mimic, on a small scale, many of the phenomena seen in *quasars*. Their discovery opens the way to study the connection between the accretion of plasma onto the black holes and the origin of the relativistic jets observed in remote quasars (Sect. 13.3).

In spite of the vast differences in luminosity and the sizes of microquasars in our Galaxy and those in active galaxies, both phenomena are believed to be powered by gravitational energy released during the accretion of plasmas onto black holes. Since the accreting plasmas have non-zero angular momentum, they form *accretion disks* orbiting around black holes. If the accreting plasmas have non-zero poloidal magnetic field, the magnetic flux accumulates in the inner region of the disk to form a global poloidal field penetrating the disk. Such poloidal fields could also be generated by dynamo action inside the accretion disk.

In either case, poloidal fields are twisted by the rotating disk toward the azimuthal direction. Moreover this process extracts angular momentum from the disk, enabling efficient accretion of disk plasmas onto black holes. In addition, magnetic twist generated during this process accelerates plasmas in the surface layer of the disk toward the polar direction by the Lorentz force to form bi-directional relativistic jets which are also collimated by the magnetic force (Lovell, 1976).

20.1.4 MHD Waves of Arbitrary Amplitude

Let us return to the case of an incompressible medium. Consider a steady flow of the type (20.8) and (20.9) in the magnetic field shown in Fig. 20.2.

In the region 1, transformed to the frame of reference, the wave front of an arbitrary amplitude $\mathbf{h} = \mathbf{h}(x)$ runs against the immovable plasma in the uniform magnetic field $\mathbf{B}_0$, the front velocity being the Alfvén one:

$$\mathbf{v}_0 = \frac{\mathbf{B}_0}{\sqrt{4\pi\rho}}. \quad (20.14)$$

On the strength of condition (20.11), in such a wave

$$p + \frac{(\mathbf{B}_0 + \mathbf{h})^2}{8\pi} = p_0 + \frac{B_0^2}{8\pi} = \text{const}, \quad (20.15)$$

i.e. the gas pressure is balanced everywhere by the magnetic pressure.

The non-compensated magnetic tension, $(\mathbf{B} \cdot \nabla) \mathbf{B}/4\pi$, provides the wave motion of arbitrary amplitude

(cf. Sect. 15.2.2). In this sense, the MHD waves are analogous to elastic waves in a string. MHD waves of an arbitrary amplitude were found for the first time by Alfvén (1950) as non-stationary solutions of the MHD equations for an incompressible medium (see also Alfvén 1981).

The Alfvén or rotational discontinuity considered in Sect. 16.2 is a particular case of the solutions (20.8) and (20.9), corresponding to a discontinuous velocity profile. Behavior of Alfvén waves in the isotropic and anisotropic astrophysical plasmas can be essentially different (see Sect. 7.3).

20.1.5 Differential Rotation and Isorotation

Now we consider another exact solution to the stationary equations of ideal MHD, (20.1)–(20.6). Let us suppose that an equilibrium configuration (for example, a star) rigidly rotates about the symmetry axis of the cylindrically symmetric ($\partial/\partial\varphi = 0$) magnetic field. The angular velocity $\boldsymbol{\omega}$ is a constant vector. Then

$$\mathbf{v} = \mathbf{r} \times \boldsymbol{\omega} = \{0, 0, v_\varphi\}, \quad (20.16)$$

where

$$v_\varphi = \omega r.$$

The induction equation (20.2) is satisfied identically in this case.

Now we relax the assumption that $\boldsymbol{\omega}$ is a constant. Consider the case of the so-called *differential rotation*. Let the vector $\boldsymbol{\omega}$ be everywhere parallel to the z axis, i.e. the symmetry axis of the field $\mathbf{B}$, but the quantity $|\boldsymbol{\omega}| = \omega$ be dependent on the coordinates r and z , where r is the cylindrical radius:

$$\boldsymbol{\omega} = \omega(r, z).$$




Fig. 20.3 Differential rotation creates the toroidal (B_ϕ) component of a magnetic field inside a star

Hence

$$v_\phi = \omega(r, z) r. \quad (20.17)$$

Substitution of (20.17) in the induction equation (20.2), with allowance being made for $\partial/\partial\phi = 0$ and (20.4), gives

$$\text{rot}(\mathbf{v} \times \mathbf{B}) = \mathbf{e}_\phi r (\mathbf{B} \cdot \nabla \omega) = 0.$$

Therefore

$$\mathbf{B} \cdot \nabla \omega = 0,$$

(20.18)

i.e. the magnetic field lines are situated at $\omega = \text{const}$ surfaces. When treated in astrophysics, this case is called *isorotation*.

As a consequence of cylindrical symmetry, the $\omega = \text{const}$ surfaces are those of rotation, hence *isorotation does not change the magnetic field*.

On the other hand, if the condition for isorotation (20.18) is not valid, *differential rotation twists the field lines*, for example as shown in Fig. 20.3, creating a *toroidal field* B_ϕ . The magnetic field is amplified.

Rigid rotation and isorotation are widely discussed, when applied to stellar physics, because

rotation is an inherent property of the majority of the stars having strong magnetic fields

Schrijver and Zwaan (1999). What is the actual motion of the plasma in the interior of stars?

Suppose there is no tangential stress at the surface of a star. The rigid rotation must be gradually established owing to viscosity in the star. However the observed motion of the Sun, as a well studied example, is by no means rigid: *the equator rotates faster than the poles*. This effect cannot be explained by surface rotation.

Fig. 20.4 Schematic summary of a radial gradient in rotation that have been inferred from helioseismic measurements





Fig. 20.5 Schematic of the flow of angular momentum in the convective zone, tachocline, and photosphere, that may be responsible for the rotation gradients summarized in previous figure



Deep layers of the Sun and fast-rotating solar-type stars participate in complex motions: differential rotation, convection, and meridional circulation (see [Rüdiger and Rekowski 1998](#)). Such motions ensure mixing of deep solar layers down to the solar core. The circumstantial evidence for this comes from observations of the solar neutrino flux as well as helioseismological data. The latter show, in particular, that *the solar core rotates faster than the surface*. The results of the *SOHO* helioseismology enable us to know the structure of the solar internal differential rotation [Schou et al. \(1998\)](#).

Roughly speaking, in the convective zone (see CZ in Fig. 20.4) the angular velocity ω is independent of radius r . The radiative interior (RI) appears to rotate almost uniformly, and is separated from the differentially rotating convective zone by a thin shear layer called the *tachocline* (shown by T in Fig. 20.4). The last is, in fact, too thin to be convincingly resolved by the *SOHO* data.

Numerical simulations are still rather far from producing a radius-independent differential rotation in the convective zone. A qualitative perspective, which probably will define a context for progress in the future, invokes the concepts of angular momentum balance and transport, and angular momentum cycles in the Sun. With this perspective, it is possible to consider all the angular velocity domains in the outer part of the Sun in a unified way [Gilman \(2000\)](#). Figure 20.5 illustrates how angular momentum could be continually cycling in the convective zone and adjacent layers.

If we accept that some process dominates in the cycle by transporting angular momentum from high latitudes to low in the bulk of the convective zone, then everything else follows. All that is required is that some of this momentum “leaks” into the tachocline below and the granulation and supergranulation layers above. Then, to complete the cycle, there is transport of angular momentum back toward the pole in both layers. There the momentum reenters the bulk of the convective zone to be recycled again.

Recommended Reading: [Elsasser \(1956\)](#), [Parker \(1979\)](#), [Moreau \(1990\)](#).

20.2 Flows at Small Magnetic Reynolds Numbers

While investigating MHD flows in a laboratory, the finite conductivity being significant, one has to account for the magnetic field dissipation. Furthermore one has to take account of the fact that the freezing-in condition breaks down owing to the smallness of the magnetic Reynolds number ([12.64](#)):

$$\text{Re}_m = \frac{\nu L}{\nu_m} \ll 1. \quad (20.19)$$

The analogous situation takes place, for example, in deep layers of the solar atmosphere near the temperature minimum. The conductivity is small here, since the number of neutral atoms is relatively large (e.g., [Hénoux and Somov 1987, 1991](#)).

Stationary flows are possible in the case of finite conductivity. However they differ greatly from the ideal medium flows considered in the previous section. The difference manifests itself in the fact that, given dissipative processes, steady flows are realized only under action of some external force, a pressure gradient, for instance. A second difference is that *the plasma of finite conductivity can flow across the magnetic field lines*.


20.2.1 Stationary Flows Inside a Duct

We shall examine a flow which has been well studied for reasons of practical importance. Let us consider the steady flow of a viscous conducting fluid along a duct with a transversal magnetic field. Let the x axis of the Cartesian system (Fig. [20.6](#)) be chosen in the flow direction, the external uniform field $\mathbf{B}_0$ coinciding with the z axis:

$$\mathbf{v} = \{v(z), 0, 0\}, \quad \mathbf{B}_0 = \{0, 0, B_0\}. \quad (20.20)$$

Let the width of the duct be $2l$.

Fig. 20.6 Formulation of the problem on the finite conductivity plasma flow in a duct



We start from the set of equations (12.43)–(12.48) for a steady flow of an incompressible medium:

$$\rho = \text{const.} \quad (20.21)$$

Consider two equations:

$$\text{rot}(\mathbf{v} \times \mathbf{B}) + \nu_m \Delta \mathbf{B} = 0, \quad (20.22)$$

$$(\mathbf{v} \cdot \nabla) \mathbf{v} = -\frac{\nabla p}{\rho} - \frac{\mathbf{B} \times \text{rot} \mathbf{B}}{4\pi\rho} + \nu \Delta \mathbf{v}. \quad (20.23)$$

The pressure gradient $\partial p / \partial x$ along the x axis, which is independent of x , is assumed to be the cause of the motion. Supposing the flow to be relatively slow, neglect the term on the left-hand side of (20.23).

Let $b = b(z)$ be the magnetic field component along the velocity. In the coordinate form, (20.22) and (20.23) are reduced to the following three equations:

$$B_0 \frac{\partial v}{\partial z} + \nu_m \frac{\partial^2 b}{\partial z^2} = 0, \quad (20.24)$$

$$\rho \nu \frac{\partial^2 v}{\partial z^2} + \frac{B_0}{4\pi} \frac{\partial b}{\partial z} - \frac{\partial p}{\partial x} = 0, \quad (20.25)$$

$$\frac{\partial}{\partial z} \left(p + \frac{b^2}{8\pi} \right) = 0. \quad (20.26)$$


Differentiating (20.26) with respect to x gives

$$\frac{\partial^2 p}{\partial x \partial z} = 0. \quad (20.27)$$

Differentiating (20.25) with respect to z , with care taken of (20.27), gives

$$\rho \nu \frac{\partial^3 v}{\partial z^3} + \frac{B_0}{4\pi} \frac{\partial^2 b}{\partial z^2} = 0. \quad (20.28)$$

Fig. 20.7 Usual parabolic ($Ha = 0$) and Hartmann profiles of the viscous flow velocity in a duct with a transverse magnetic field



Eliminate $\partial^2 b / \partial z^2$ between (20.24) and (20.28). The result is

$$\frac{d^3 v}{dz^3} - \frac{B_0^2}{4\pi\rho\nu v_m} \frac{dv}{dz} = 0. \quad (20.29)$$

This equation is completed by the boundary conditions on the duct walls

$$v(l) = v(-l) = 0. \quad (20.30)$$

The corresponding solution is of the form

$$v(z) = v_0 \frac{\cosh Ha - \cosh(Ha z/l)}{\cosh Ha - 1}. \quad (20.31)$$

Here $v_0 = v(0)$ is the flow velocity at the center of the duct, the dimensionless parameter characterizing the flow is

$$Ha = \frac{l B_0}{\sqrt{4\pi\rho\nu v_m}}.$$

(20.32)

It is called the *Hartmann number*, the flow (20.31) being the Hartmann flow. As $Ha \rightarrow 0$, formula (20.31) converts to the usual parabolic velocity profile which is typical of viscous flows in a duct without a magnetic field:

$$v(z) = v_0 \left(1 - \frac{z^2}{l^2}\right). \quad (20.33)$$

The influence of a transversal magnetic field shows itself as the appearance of *an additional drag to the plasma flow* and the change of the velocity profile which becomes *flatter* in the central part of the duct (Fig. 20.7).

In the limit $Ha \rightarrow \infty$, the Hartmann formula (20.31) gives

$$v(z) = v_0 \left\{ 1 - \exp \left[-Ha \left(1 - \frac{z}{l} \right) \right] \right\}. \quad (20.34)$$

Such a velocity profile is flat, $v(z) \approx v_0$, the exception being a thin layer near the walls, the *boundary layer* of the thickness l/Ha .

20.2.2 The MHD Generator or Pump

What factors determine the value of velocity v_0 at the center of the duct? To find them let us calculate the electric current density in the duct

$$\begin{aligned} j_y &= \frac{c}{4\pi} \frac{\partial b}{\partial z} = \frac{c}{4\pi} \left(\frac{4\pi}{B_0} \frac{\partial p}{\partial x} - \rho v \frac{4\pi}{B_0} \frac{\partial^2 v}{\partial z^2} \right) = \\ &= \frac{c}{B_0} \left(\frac{\partial p}{\partial x} - \rho v \frac{\partial^2 v}{\partial z^2} \right). \end{aligned} \quad (20.35)$$

Here the use is made of formula (20.25) to find the derivative $\partial b/\partial z$. Let us substitute in (20.35) an expression for velocity of the type (20.31), i.e.

$$v(z) = A \left(\cosh Ha - \cosh \frac{Ha z}{l} \right). \quad (20.36)$$

We get the following equation

$$\frac{j_y B_0}{c} = \frac{\partial p}{\partial x} - \rho v A \left(\frac{Ha}{l} \right)^2 \cosh \frac{Ha z}{l}. \quad (20.37)$$

Let us integrate (20.37) over z from $-l$ to $+l$. The result is


$$\frac{IB_0}{c} = 2l \frac{\partial p}{\partial x} - A 2\rho v \left(\frac{Ha}{l} \right) \sinh Ha, \quad (20.38)$$

where

$$I = \int_{-l}^l j_y dz \quad (20.39)$$

is the total current per unit length of the duct. We shall assume that there is an electrical circuit for this current to flow outside the duct. The opposite case is considered in Landau and Lifshitz 1959a, Chap. 8, Sect. 67.

Fig. 20.8 Utilization of the MHD duct as the generator of the current I ; R is an external load



Finally it follows from (20.38) that the sought-after coefficient in formula (20.36) is

$$A = \frac{\partial p / \partial x - (1/2lc) IB_0}{(\rho v / l^2) \text{Ha} \sinh \text{Ha}}. \quad (20.40)$$

Thus

the velocity of the plasma flow in the duct is proportional to the gas pressure gradient and the magnetic Lorentz force.

This is why two different operational regimes are possible for the duct.

If the flow in the duct is realized under the action of an external pressure gradient, the duct operates as the MHD generator shown in Fig. 20.8. The same principle explains the action of flowmeters (for more detail see Shercliff 1965, Sect. 6.5; Sutton and Sherman 1965, Sect. 10.2) which are important, for example, in controlling the flow of the metallic heat conductor in reactors.

The second operating mode of the duct occurs when an external electromagnetic force (instead of a passive load R in Fig. 20.8) creates the electric current I between the walls of the duct. Interaction of the current with the external magnetic field $\mathbf{B}_0$ gives rise to the Lorentz force that makes the plasma move along the duct, i.e. in the direction of the x axis. Hence the duct operates as the MHD pump, and this is also used in some technical applications.

20.2.3 Weakly-Ionized Plasma in Astrophysics

Under astrophysical conditions, both operating modes of the MHD duct are realized, once the plasma resistivity is high due, for instance, to its low temperature. In the solar atmosphere, in the minimum temperature region, neutral atoms move in the directions of convective flows and collide with ions, thus setting them in motion.

At the same time, electrons remain “frozen” in the magnetic field. This effect (termed the *photospheric dynamo*) can generate electric currents and amplify the magnetic field in the photosphere and the low chromosphere (see Part II, Sect. 12.4).

A violent outflow of high-velocity weakly-ionized plasma is one of the first manifestations of the formation of a new stars (Bachiller 1996; Bontemps et al. 1996). Such outflows emerge bipolarly from the young object and involve amounts of energy similar to those involved in accretion processes. The youngest protostellar low-mass objects known to date (the class 0 protostars) present a particularly efficient outflow activity, indicating that outflow and infall motions happen simultaneously and are closely linked since the very first stages of the star formation processes.

The idea of a new star forming from relatively simple hydrodynamic infall of weakly-ionized plasma is giving place to a picture in which magnetic fields play a crucial role and stars are born through the formation of complex engines of accretion/ejection. It seems inevitable that future theories of star formation will have to take into account, together with the structure of the protostar and its surrounding accretion disk, the processes related to *multi-fluid hydrodynamics of weakly-ionized plasma*. These are the effects similar to the photospheric dynamo and magnetic reconnection in weakly-ionized plasma (Part II, Sect. 12.3).

Recommended Reading: Sutton and Sherman (1965), Ramos and Winowich (1986).

20.3 The σ -Dependent Force and Vortex Flows

20.3.1 Simplifications and Problem Formulation

As was shown in Sect. 19.4, a body plunged into a conducting fluid with magnetic and electric fields is acted upon by an *expulsion* force or, more exactly, by the magnetic σ -dependent force. As this takes place, the electric field $\mathbf{E}$ and current density $\mathbf{j}$ are non-uniform, and the volume *Lorentz force inside the fluid is non-potential*. The force generates vortex flows of the fluid in the vicinity of the body.

- (a) Let us consider the stationary problem for an incompressible fluid having uniform constant viscosity ν and magnetic diffusivity ν_m Syrovatskii and Chesalin (1963); Marty and Alemany (1983); Gerbeth et al. (1990). Let, at first, both the usual and magnetic Reynolds numbers be small:

$$\text{Re} = \frac{\nu L}{\nu} \ll 1, \quad (20.41)$$

$$\text{Re}_m = \frac{\nu L}{\nu_m} \ll 1. \quad (20.42)$$

The freezing-in condition (12.65) can be rewritten in the form

$$\Delta \mathbf{B} + \text{Re}_m \text{curl}(\mathbf{v} \times \mathbf{B}) = 0, \quad (20.43)$$

where, in view of (20.42), Re_m is a small parameter. In a zeroth approximation in this parameter, the magnetic field is potential:

$$\Delta \mathbf{B} = 0.$$

Moreover the magnetic field will be assumed to be uniform, in accordance with the formulation of the problem discussed in Sect. 19.4. Strictly speaking, the assumption of a *uniform magnetic field* implies the inequality

$$B \gg \frac{4\pi}{c} Lj. \quad (20.44)$$

Its applicability will be discussed later on, in connection with the simplified form of Ohm's law to be used while solving the problem.

- (b) Assuming the stationary flows occurring in the fluid to be slow, the inertial force (proportional to v^2) will be ignored in the equation of motion (20.23) as compared to the other forces: *pressure gradient*, *Lorentz force*, *viscous force*. The term describing the gravity force will be dropped, since its effect has already been studied in Sect. 19.4. Finally, on multiplying the equation

$$0 = -\frac{\nabla p}{\rho} - \frac{\mathbf{B} \times \text{rot} \mathbf{B}}{4\pi\rho} + \nu \Delta \mathbf{v}$$

by the fluid density $\rho = \rho_0$, it is rewritten in the form

$$\eta \Delta \mathbf{v} = \nabla p - \mathbf{f}. \quad (20.45)$$

Here $\eta = \rho_0 \nu$ is the *dynamic viscosity coefficient*, and

$$\mathbf{f} = \frac{1}{c} \mathbf{j} \times \mathbf{B}_0 \quad (20.46)$$

is the Lorentz force in the same approximation.

Recall that, in view of the assumed incompressibility of the fluid, the velocity field obeys the equation


$$\text{div} \mathbf{v} = 0, \quad (20.47)$$

which formally follows from the continuity (20.3).

- (c) The electric field $\mathbf{E}$ is assumed to be uniform at infinity

$$\mathbf{E} \rightarrow \mathbf{E}_0, \quad r \rightarrow \infty. \quad (20.48)$$

Fig. 20.9 An uniform conducting ball of radius a , submerged in a conducting fluid with electric and magnetic fields



Given the conductivities of the fluid σ_0 and of the submerged body σ_1 , we can find the current $\mathbf{j}$ in the whole space using the following conditions:

$$\operatorname{div} \mathbf{j} = 0, \quad (20.49)$$

$$\mathbf{j} = \sigma \mathbf{E}, \quad (20.50)$$

$$\operatorname{rot} \mathbf{E} = 0. \quad (20.51)$$

The current $(\sigma/c) \mathbf{v} \times \mathbf{B}$ has been ignored in Ohm's law (20.50). This may be done, once the velocity of engendered vortex flows is much less than the drift velocity, i.e. once the inequality

$$v \ll v_d = c \frac{E}{B} \quad (20.52)$$

holds. Note that substituting (20.44) in (20.52) results in the inequality

$$\frac{vL}{(c^2/4\pi\sigma)} \ll 1, \quad (20.53)$$

which coincides with the initial assumption (20.42).

20.3.2 The Solution for a Spherical Ball

Let us solve the problem for a ball of radius a . We choose the Cartesian frame of reference, in which the direction of the x axis is parallel to $\mathbf{E}_0$, and the origin of coordinates coincides with the center of the ball as shown in Fig. 20.9.

By virtue of Ohm's law (20.50), the electric current at infinity

$$\mathbf{j}_0 = \sigma_0 \mathbf{E}_0 \quad (20.54)$$

is also parallel to the x axis.

It follows from (20.51) that the current can be represented in the form

$$\mathbf{j} = \nabla \psi. \quad (20.55)$$

Here a scalar function ψ , in view of (20.49), satisfies the Laplace equation

$$\Delta\psi = 0. \quad (20.56)$$

Let us try to find the solution to the problem in the form of uniform and dipole components:

$$\psi = \mathbf{j}_0 \cdot \mathbf{r} + c_0 \mathbf{j}_0 \cdot \nabla \frac{1}{r}, \quad r \geq a, \quad (20.57)$$

and

$$\psi = \mathbf{j}_1 \cdot \mathbf{r}, \quad r < a. \quad (20.58)$$

Here c_0 is an unknown constant, $\mathbf{j}_1 = \{j_1, 0, 0\}$ is an unknown current density inside the ball. Both unknowns are to be found from the matching conditions at the surface of the ball:

$$\{j_r\} = 0 \quad \text{and} \quad \{\mathbf{E}_\tau\} = 0.$$

These conditions can be rewritten as follows

$$\frac{\mathbf{j} \cdot \mathbf{r}}{r} = \frac{\mathbf{j}_1 \cdot \mathbf{r}}{r} \quad \text{at } r = a, \quad (20.59)$$

and

$$\frac{\mathbf{j}_\tau}{\sigma_0} = \frac{\mathbf{j}_{\tau 1}}{\sigma_1} \quad \text{at } r = a. \quad (20.60)$$

On substituting (20.57) and (20.58) in (20.59) and (20.60), the constants c_0 and j_1 are found. The result is

$$\psi = \left[1 + \beta \left(\frac{a}{r} \right)^3 \right] \mathbf{j}_0 \cdot \mathbf{r} \quad \text{for } r \geq a, \quad (20.61)$$

and

$$\psi = (1 - 2\beta) \mathbf{j}_0 \cdot \mathbf{r} \quad \text{for } r < a. \quad (20.62)$$

Here the constant

$$\beta = \frac{\sigma_0 - \sigma_1}{2\sigma_0 + \sigma_1}. \quad (20.63)$$

Specifically, inside the ball

$$\mathbf{j}_1 = (1 - 2\beta) \mathbf{j}_0, \quad (20.64)$$

and $\mathbf{j}_1 = \mathbf{j}_0$, once $\sigma_1 = \sigma_0$.

20.3.3 Forces and Flows Near a Spherical Ball

Knowing the current in the whole space, we can find the Lorentz force (20.46)

$$\mathbf{f} = \frac{1}{c} \nabla \psi \times \mathbf{B}_0 = \text{rot} \frac{\psi \mathbf{B}_0}{c}. \quad (20.65)$$

In the case at hand,

the volume Lorentz force has a rotational character and hence generates vortex flows in the conducting fluid.

Let us operate with rot rot on (20.45). Using the known vector identity

$$\text{rot rot } \mathbf{a} = \nabla (\nabla \mathbf{a}) - \Delta \mathbf{a}$$

and taking account of relations (20.49)–(20.51), a biharmonic equation for the velocity field is obtained

$$\Delta \Delta \mathbf{v} = 0. \quad (20.66)$$

Operating with divergence on (20.45) and taking account of (20.49)–(20.51), we get

$$\Delta p = 0. \quad (20.67)$$

Equations (20.66) and (20.67) are to be solved together with (20.45) and (20.47). For bodies with spherical or cylindrical symmetry, it is convenient to make use of the identity

$$\mathbf{r} \cdot \Delta \mathbf{q} = \Delta (\mathbf{q} \cdot \mathbf{r}), \quad (20.68)$$

where $\mathbf{q}$ is any vector satisfying the condition $\text{div } \mathbf{q} = 0$. Then from (20.66) subject to the condition (20.47) we find

$$\Delta \Delta (v_r r) = 0. \quad (20.69)$$

The boundary conditions are taken to be

$$v|_S = 0, \quad v|_\infty = 0. \quad (20.70)$$

Here S is the surface of the submerged body which is assumed to be a ball of radius a (cf. Fig. 20.9). At its surface $r = a = \text{const}$, (20.47) and the first of conditions (20.70) give

$$\left. \frac{\partial v_r}{\partial r} \right|_S = 0. \quad (20.71)$$

The solution of (20.69), satisfying the boundary condition (20.71) and the second of conditions (20.70), is clearly seen to be

$$v_r \equiv 0. \quad (20.72)$$

Thus

in the case of a spherical ball, the flow lines of a conducting incompressible fluid are situated at $r = \text{const}$ surfaces.

Next an equation for the pressure is found using (20.45) and taking into account that, by virtue of (20.68),

$$\mathbf{r} \cdot \Delta \mathbf{v} = \Delta (v_r r) = 0.$$

The resulting equation is

$$\frac{\partial p}{\partial r} = f_r. \quad (20.73)$$

The function f_r occurring on the right-hand side is the radial component of the above mentioned Lorentz force (20.65).

Once the plasma pressure has been found by integrating (20.73) and (20.67), the velocity is determined from (20.45) with the known right-hand side.

Choose the Cartesian frame of reference in which

$$\mathbf{B}_0 = \{ B_{0x}, 0, B_{0z} \},$$

$B_{0x} = B_{\parallel}$ and $B_{0z} = B_{\perp}$ being the magnetic field components parallel and perpendicular to $\mathbf{j}_0$, respectively (see Fig. 20.9). The current in the conducting fluid (cf. formula (20.61)) is

$$\mathbf{j} = \nabla \psi, \quad \psi = j_0 x + j_0 \frac{\beta a^3 x}{r^3}, \quad (20.74)$$

the current inside the ball being defined by formula (20.64). The pressure in the fluid

$$p = \frac{1}{c} j_0 B_{\perp} y \left(\frac{\beta a^3}{2r^3} - 1 \right) + \text{const}. \quad (20.75)$$

It is convenient to rewrite the velocity distribution in spherical coordinates

$$\mathbf{v} = \{ v_r, v_{\theta}, v_{\varphi} \} \quad (20.76)$$

(cf. Syrovatskii and Chesalin, 1963):


$$\begin{aligned} v_r &= 0, \\ v_{\theta} &= \frac{\beta j_0 a^2}{4c\eta} \frac{a}{r} \left(1 - \frac{a^2}{r^2} \right) (-B_{\perp} \cos \theta \sin \varphi + B_{\parallel} \sin \theta \sin 2\varphi), \\ v_{\varphi} &= \frac{\beta j_0 a^2}{4c\eta} \frac{a}{r} \left(1 - \frac{a^2}{r^2} \right) (B_{\perp} \cos 2\theta \cos \varphi + B_{\parallel} \sin 2\theta \cos^2 \varphi). \end{aligned}$$

This velocity field pattern is shown in Fig. 20.10.

The force acting on the body is defined to be (cf. formula (19.60))

$$\mathbf{F} = \frac{1}{c} \int_V \mathbf{j} \times \mathbf{B}_0 dV + \oint_S p \mathbf{n} dS - \oint_S \sigma'_n dS, \quad (20.77)$$

Fig. 20.10 Vortex flows near the conducting ball submerged in a conducting fluid with electric and magnetic fields



where $\mathbf{n}$ is the inward normal to the sphere;

$$\sigma'_n = (\sigma'_{\alpha\beta} n_\beta)_n, \quad (20.78)$$

$\sigma'_{\alpha\beta}$ being the viscous stress tensor, see definition (12.54).

On substituting the velocity distribution (20.76) in the viscous force formula (20.78) and integrating (20.77) over the surface S of the ball,

the sum of the viscous forces is concluded to be zero. The moment of the viscous forces acting on the ball is also zero.

The remaining force determined by (20.77) is directed along the y axis and is equal to

$$F = \frac{4\pi a^3}{3} \frac{j_0 B_\perp}{c} \left\{ -(1 - 2\beta) + \left(1 - \frac{\beta}{2}\right) \right\}. \quad (20.79)$$

The constant β is defined by formula (20.63):

$$\beta = \frac{\sigma_0 - \sigma_1}{2\sigma_0 + \sigma_1}.$$

The first term in the roty brackets corresponds to the force $\mathbf{j}_1 \times \mathbf{B}_0 / c$ which immediately acts on the current $\mathbf{j}_1$ inside the ball. Note that


$$1 - 2\beta = \frac{3\sigma_1}{2\sigma_0 + \sigma_1} > 0,$$

in agreement with the direction of the vector product $\mathbf{j}_1 \times \mathbf{B}_0$ or $\mathbf{j}_0 \times \mathbf{B}_0$ (Fig. 20.9). Moreover, provided $\sigma_1 = 0$, the term $(1 - 2\beta) = 0$ as it should be the case for a non-conducting ball, since there is no current inside it.

The second term in the roty brackets of formula (20.79) expresses the sum of the forces of the pressure on the surface of the ball. The coefficient

$$1 - \frac{\beta}{2} = \frac{3(\sigma_0 + \sigma_1)}{2(2\sigma_0 + \sigma_1)} > 0,$$

Fig. 20.11 The expulsion force $\mathbf{F}$ acting on the non-conducting ball submerged in a conducting fluid with electric and magnetic fields



signifying that

the actual σ -dependent force is always somewhat less than the force owing to the interaction of the current $\mathbf{j}_1$ and the magnetic field $\mathbf{B}_0$. Moreover the total force can be opposite in sign.

In the particular case $\sigma_1 = 0$, when the current $j_1 = 0$

$$1 - \frac{\beta}{2} = \frac{3}{4}.$$

Hence $F > 0$. The non-conducting ball is expelled in the direction opposite to that of the vector product $\mathbf{j}_0 \times \mathbf{B}_0$ (Fig. 20.11).

The above properties of the magnetic σ -dependent force are used, for example, in technical MHD. They constitute the principle of action for magnetic separators which are intended for dividing mechanical mixtures having different conductivities.

Having the physical sense of the two terms determining the magnetic σ -dependent force (20.79), let us combine them in the following descriptive formula:

$$\mathbf{F} = -\mathbf{f}_0 V \times \frac{3}{2} \beta.$$

(20.80)

Here $V = 4\pi a^3/3$ is the volume of the ball, $\mathbf{f}_0 = \mathbf{j}_0 \times \mathbf{B}_0 / c$ is the Lorentz force in the conducting fluid with uniform magnetic $\mathbf{B}_0$ and electric $\mathbf{E}_0$ fields (cf. (19.59)), the coefficient β being determined by formula (20.63).

20.4 Large Magnetic Reynolds Numbers

In the previous section we have considered the solution to the MHD problem concerning the magnetic σ -dependent force in the limit of small (usual and magnetic) Reynolds numbers. [Leenov and Kolin \(1954\)](#) were the first to obtain similar solutions in connection with the problem of electromagnetophoresis.

As a rule the opposite limiting case is applicable for astrophysical use. In this case, the problem of the magnetic σ -dependent force is difficult and can hardly be solved completely, especially given

$$\text{Re} \ll 1, \quad \text{Re}_m \gg 1. \quad (20.81)$$

A situation of this kind occurs, for example, in solar prominences (Sect. 20.4.2). In what follows we will show (Litvinenko and Somov 1994; Somov 1994b) that an expression for the magnetic σ -dependent force can be found for large magnetic Reynolds numbers, without rigorous calculations of the characteristics of the plasma flow near a body.

20.4.1 The General Formula for the σ -Dependent Force

The equations of stationary MHD for flows of an incompressible fluid with density ρ_0 and dynamic viscosity $\eta = \rho_0 \nu$ are of the form:

$$\begin{aligned} \rho (\mathbf{v} \cdot \nabla) \mathbf{v} &= -\nabla p + \frac{1}{c} \mathbf{j} \times \mathbf{B} + \eta \Delta \mathbf{v}, \\ \text{rot} (\mathbf{v} \times \mathbf{B}) + \nu_m \Delta \mathbf{B} &= 0, \\ \text{rot} \mathbf{B} &= \frac{4\pi}{c} \mathbf{j}, \quad \text{div} \mathbf{v} = 0, \quad \text{div} \mathbf{B} = 0. \end{aligned} \quad (20.82)$$

Let us find the σ -dependent force density $\mathbf{f}$ on the basis of *similarity* considerations. The given set of equations implies that five quantities are the determining parameters of the problem: ν , ν_m , a , ρ_0 , and $\mathbf{f}_0$. By way of example, velocity v_0 depends on these parameters. Hence v_0 rather than ρ_0 may be treated as a determining parameter. The standard procedure of *dimensional analysis*, described by Bridgman (1931), gives us the formula

$$\mathbf{f} = -\mathbf{f}_0 \Phi(\text{Re}, \text{Re}_m). \quad (20.83)$$

In the limit $\text{Re}_m = 0$ it reproduces (in a slightly different notation) the result presented in the theoretical part of the paper by Andres et al. (1963). Experimental data, which are stated in the same paper for $\text{Re} < 10^2$, allow one to conclude that, with an accuracy which is completely sufficient for astrophysical applications,

$$\Phi(\text{Re}, \text{Re}_m) \approx \Phi_1(\text{Re}_m), \quad (20.84)$$

where $\Phi_1(0) \approx 1$.

Generally, the behaviour of the magnetic field lines near the body for $\text{Re}_m \neq 0$ can become nonregular and intricate, as a consequence of the electric current redistribution and vortex flow generation. For example, if $\text{Re}_m < 1$, then the value of

the nonregular field component $\delta B \approx \text{Re}_m B_0$. The effective magnitude of the field and the magnetic σ -dependent force decrease as compared to the case $\text{Re}_m = 0$.

The form of the decreasing function Φ_1 for $\text{Re}_m \gg 1$ can be determined as follows. Far from the body, at infinity, the electromagnetic energy flux is equal to

$$\mathbf{G}_0 = \frac{c}{4\pi} \mathbf{E}_0 \times \mathbf{B}_0. \quad (20.85)$$

In close proximity to the body, the magnitude of the Poynting vector must diminish once the disordered behaviour of lines of force is assumed. The difference $(G_0 - G)$ is equal to the power of engendered vortex flows, hence generally we get

$$fa^3 v_0 \leq G_0 a^2. \quad (20.86)$$

The equality (20.86) is achieved in the limit $\text{Re}_m \rightarrow \infty$. Here the characteristic velocity v_0 is determined from the equation of motion in the set (20.4.1):

$$v_0 = fa^2/\eta \quad \text{for } \text{Re} \ll 1, \quad (20.87)$$

$$v_0 = (fa/\rho_0)^{1/2} \quad \text{for } \text{Re} \gg 1. \quad (20.88)$$

When $\text{Re}_m \rightarrow \infty$, relations (20.84)–(20.88) allow us to obtain the sought-after function appearing in formula (20.83):

$$\Phi(\text{Re}, \text{Re}_m) = \begin{cases} 1 & \text{for } \text{Re}_m < 1, \\ \text{Re}_m^{-1} & \text{for } \text{Re}_m > 1. \end{cases} \quad (20.89)$$

The case $\text{Re}_m < 1$ was treated by [Leenov and Kolin \(1954\)](#).

Strictly speaking, we could take also into account the dependence of the function Φ on the usual Reynolds number Re . We could obtain

$$\Phi(\text{Re}, \text{Re}_m) = \frac{1}{\text{Re}_m} \Phi_2(\text{Re}), \quad (20.90)$$

where the function $\Phi_2(\text{Re})$ is practically constant.

Note that formula (20.90) can be interpreted as a manifestation of an *incomplete self-similarity* of the function Φ relative to the similarity parameter Re_m ([Barenblatt 1979](#)). The point is that, from the viewpoint of a “naive” analysis, the function Φ does not depend on a dimensionless parameter whose magnitude is much greater (or less) than unity. This statement is true only if there exists a final non-zero limit of the function Φ as the parameter at hand tends to infinity (or zero). However, in general, this is not the case, as is clearly demonstrated by (20.90). In fact, $\Phi \rightarrow 0$ when $\text{Re}_m \rightarrow \infty$. At the same time the function Φ is a power-law one in Re_m ; that allows us to write down an expression for the force density $\mathbf{f}$ in a self-similar form. As this takes place, the exact form of dimensionless combinations cannot be determined from the formal dimensional analysis alone.

Therefore an order-of magnitude expression is obtained for the density of the magnetic σ -dependent force acting on a body submerged into a conducting fluid or plasma (Litvinenko and Somov 1994; Somov 1994b):

$$\mathbf{f} = -\frac{c}{4\pi v_0 a} \mathbf{E}_0 \times \mathbf{B}_0. \quad (20.91)$$

The expression (20.91) is valid in the limit of large magnetic Reynolds numbers. For a body with a non-zero conductivity σ_1 , the electric current flowing inside the body must be taken care of in formula (20.91). The corresponding treatment was presented in Sect. 20.3.

The physical sense of formula (20.91) is obvious. Comparison of (20.91) with formula (19.59) for the σ -dependent force, which then holds a uniform current flow in the plasma, shows that for $\text{Re}_m \rightarrow \infty$ ($\sigma \rightarrow \infty$) the plasma in the vicinity of the body possesses, as it were, an *effective* conductivity

$$\sigma_{\text{ef}} \approx \frac{c^2}{v_0 a}. \quad (20.92)$$

This finite conductivity of a plasma is a result of the electromagnetic energy losses to generation of *macroscopic* vortex flows.

This mechanism of conductivity of a plasma is different from the usual microscopic one, in which energy losses result from Coulomb collisions of current-carrying electrons with thermal electrons and ions of the plasma. It is no accident that an expression for conductivity, which is equivalent to (20.92), has emerged in quite another problem—while calculating the electrical resistivity of necks in Z-pinches appearing in a highly conductive plasma (Chernov and Yan'kov 1982).

Note in this context that the σ -dependent force, as well as the characteristic velocity of the plasma flow, depends in a *non-linear* way on the quantity $E_0 B_0$. Using (20.87), (20.88) and (20.91), we see that

$$f \sim \begin{cases} (E_0 B_0)^{1/2}, & \text{Re} \ll 1, \\ (E_0 B_0)^{2/3}, & \text{Re} \gg 1. \end{cases} \quad (20.93)$$

Litvinenko and Somov (1994) have supposed that

the magnetic σ -dependent force may play an important part in the dynamics of astrophysical plasma with a non-uniform distribution of temperature and, hence, electric conductivity.

It is this force that can generate large-scale vortex flows of plasma in space. This possibility is illustrated in the next section.

20.4.2 The σ -Dependent Force in Solar Prominences

The solar corona is a natural “plasma physics laboratory” where formula (20.91), which is applicable at large magnetic Reynolds numbers, can be tested. Recall several of characteristics of the solar corona: low density $\rho_0 \approx 10^{-16} \text{ g cm}^{-3}$, high temperature $T_0 \approx 10^6 \text{ K}$, dynamic viscosity $\eta \approx 1 \text{ g cm}^{-1} \text{ s}^{-1}$, magnetic field $B_0 \approx 10 - 100 \text{ G}$, electric field $E_0 \approx 10^{-5} \text{ CGSE units}$.

On the other hand, according to observational data (Tandberg-Hanssen, 1995), prominences consist of numerous fine threads—*cold dense* formations having a transversal scale $a \approx 10^7 \text{ cm}$ and temperature $T_1 \approx 10^4 \text{ K}$. Hence the ratio

$$\sigma_1/\sigma_0 \approx 10^{-3} \ll 1,$$

as applied to prominences in the corona. In the vicinity of the threads, as well as near a prominence as a whole, rather fast plasma flows are actually observed.

According to the model under discussion, these flows can be generated by the vortex component of the magnetic σ -dependent force. For $\text{Re} \ll 1$, their maximum velocity, as follows from relations (20.87) and (20.91), is determined by the expression

$$v_0 \approx \left(\frac{c E_0 B_0 a}{4\pi \eta} \right)^{1/2} \approx 10 - 30 \text{ km s}^{-1}, \quad (20.94)$$

that, generally speaking, corresponds to the characteristic values of observed velocities. However the spatial resolution of modern optical, EUV and soft X-ray observations is smaller than is necessary for the model to be confirmed or refuted. Let us consider another possibility.

The symmetric distribution of velocities on the line-of-sight projection (i.e., in the direction towards the observer) is a distinguishing feature of the model since it predicts the presence of *a large number of vortex flows of plasma* inside the prominence. Such a distribution can be observed as a symmetric broadening of spectral lines, which it will be necessary to study if one wishes to study the effect quantitatively. A similar observational effect can be related to the existence of reconnecting current layers in the same region (Antonucci and Somov 1992; Antonucci et al. 1996).

The gravity force acting on the prominences is supposed to be balanced by the σ -dependent expulsion. The equilibrium condition makes it possible to evaluate the characteristic value of the plasma density related to the fine threads forming the prominence

$$(\rho_1 - \rho_0) g_{\odot} \approx f. \quad (20.95)$$

Here the specific gravity of the Sun $g_{\odot} \approx 3 \times 10^4 \text{ cm s}^{-2}$. Formulae (20.91), (20.92), and (20.95) result in

$$\rho_1 \approx \left(\frac{c E_0 B_0 \eta}{4\pi g_{\odot}^2 a^3} \right)^{1/2} \approx 3 \times 10^{-13} \text{ g cm}^{-3}, \quad (20.96)$$

in accordance with observational data.

Even faster flows with characteristic velocities $10^2 - 10^3 \text{ km s}^{-1}$ in so-called *eruptive* prominences are probably a consequence of the fact that the coronal fields $\mathbf{E}_0$ and $\mathbf{B}_0$ can change (in magnitude or direction) during the course of evolution. As this takes place, the equilibrium described by (20.95) can be violated.

Observations with high spectral resolution in EUV and soft X-ray ranges are necessary to study the effect of the magnetic force stimulated by the presence of plasma regions with considerably different conductivity in the solar atmosphere.

20.5 Practice: Exercises and Answers

Exercise 20.1. Discuss a possible behavior of electrically conducting spheres in an insulating bounded fluid placed in a vertical traveling magnetic field.

Hint. The spheres move in response to the induced electromagnetic forces, the motion being influenced by gravity, viscous drag, vessel boundary reaction, and collisions. The range of possible behaviors, stable, unstable, and chaotic, is very wide. The term “electromagnetic billiards” seems appropriate to describe this phenomenon Bolcato et al. (1993).

Appendix A

Notation

Latin Alphabet

<i>Symbol</i>	<i>Description</i>	<i>Introduced in Section (Formula)</i>
a	Current layer half-thickness	8.3
$\mathbf{A}$	Vector potential of a magnetic field	6.2
b	Half-width of a reconnecting current layer (RCL)	8.3
$\mathbf{b}$	Perturbation of a magnetic field	20.2.1
$\mathbf{B}$	Magnetic field	1.2
$\mathbf{B}_\tau$	Tangential magnetic field	16.2
e, e_a	Electric charge	1.2
$\mathbf{e}_c$	Unit vector from the curvature centre	5.2
$\mathcal{E}$	Energy of a particle	5.1
$\mathbf{E}$	Electric field	1.2
$\mathbf{E}_u$	Electric field in the plasma rest-frame	11.1
f_k	Averaged distribution function for particles of kind k	1.1
f_{kl}	Binary correlation function	2.2
f_{kln}	Triple correlation function	2.3
$\hat{f}_k$	Exact distribution function for particles of kind k	2.2
F	Complex potential	14.2
$\mathbf{F}, \mathbf{F}_k$	Force	1.1
$\langle \mathbf{F}_k \rangle_v$	Mean force per unit volume	9.1
$\mathbf{F}_{kl}$	Force density in the phase space	2.2
$\mathbf{F}'$	Fluctuating force	2.1
g	Velocity-integrated correlation function	3.2
G	Gravitational constant	1.2

G	Energy flux density	(1.54)
h	Magnetic field at a wave front	20.1
Ha	Hartmann number	20.2
j	Electric current density	1.2
j'	Current density in the plasma rest-frame	11.1
j_k^q	Current density due to particles of kind <i>k</i>	9.1
j_k	Particle flux density in the phase space	3.1
J	Electric current	19.3
k	Friction coefficient	1.1
k	Wave vector	15.1
K	Kinetic energy of a particle	(5.58)
m	Magnetic dipole moment	14.4
m, m_a	Particle mass	1.2
M	Mass of star	19.1
M	Magnetic moment of a particle	5.2
	Magnetic energy of a system	19.1
n, n_k	Number density	8.1
n	Unit vector along a magnetic field	5.1
N_k	Number of particles of kind <i>k</i>	1.1
p_k	Gas pressure of particles of kind <i>k</i>	9.1
p_m	Magnetic pressure	15.1
p_{αβ}	Pressure tensor	9.1
p	Particle momentum	5.1
P	Generalized momentum	6.2
q	Generalized coordinate	6.2
q	Heat flux density	12.1
q_k	Heat flux density due to particles of kind <i>k</i>	9.1
Q_k	Rate of energy release in a gas of particles of kind <i>k</i>	9.1
r_{DH}	Debye-Hückel radius	8.2
r_L	Larmor radius	5.1
r_a	Coordinates of <i>a</i> th particle	1.2
R	Radius of star	14.4
R_⊥	Guiding centre spiral radius	5.2
R	Rigidity of a particle	5.1
R	Guiding centre vector	5.2
Re	Reynolds number	12.3
Re_m	Magnetic Reynolds number	12.3
s	Entropy per unit mass	12.2
T	Temperature	12.2
T_B	Period of the Larmor rotation	5.2
T_{αβ}	Maxwellian stress tensor	12.1

$\mathbf{u}$	Relative velocity	5.1
	Velocity of the centre-of-mass system	11.1
$\mathbf{u}_e$	Mean electron velocity	11.1
$\mathbf{u}_i$	Mean ion velocity	11.1
$\mathbf{u}_k$	Mean velocity of particles of kind k	9.1
U	Interaction potential	8.1
	Volume of a fluid particle	14.2
	Specific volume of a magnetic tube	19.3
U_{th}	Thermal energy	19.1
$\mathbf{U}$	Velocity of the moving reference frame	16.2
	Shock speed	17.1
$\mathbf{v}$	Macroscopic velocity of a plasma	12.2
$\mathbf{v}, \mathbf{v}_a$	Particle velocity	1.2
$\mathbf{v}_d$	Drift velocity	5.1
v_n	Normal component of the velocity	16.2
v_x	Velocity orthogonal to a discontinuity surface	16.1
$\mathbf{v}'$	Deviation of particle velocity from its mean value	9.1
$\mathbf{v}_\tau$	Tangential velocity	16.1
$v_\parallel$	Velocity component along the magnetic field lines	5.1
$v_\perp$	Transversal velocity	5.1
V_A	Alfvén speed	13.1
$\mathbf{V}_{gr}$	Group velocity of a wave	15.1
$\mathbf{V}_{ph}$	Phase velocity of a wave	15.1
V_s	Sound speed velocity	15.1
V_{Te}	Mean thermal velocity of electrons	(5.54)
V_{Ti}	Mean thermal velocity of ions	(5.53)
V_{Tp}	Mean thermal velocity of protons	(5.55)
$V_\pm$	Speed of a fast (slow) magnetoacoustic wave	15.1
w	Probability density	2.1
w, w_k	Heat function per unit mass	9.1
W	Energy density of an electromagnetic field	(1.53)
X	Phase space	1.1
Z	Ion charge number	8.2

Greek Alphabet

<i>Symbol</i>	<i>Description</i>	<i>Introduced in Section (Formula)</i>
α_B	Parameter of the magnetic field inhomogeneity	5.2
α_E	Parameter of the electric field inhomogeneity	5.2
β	Coefficient in an expulsion force	20.3
γ	Dimensionless parameter of ideal MHD	13.1
γ_g	Ratio of specific heats	16.1
Γ	$6N$ -dimensional phase space	2.1
δ	Dimensionless parameter of ideal MHD	12.3
ε	Mean kinetic energy of a chaotic motion	12.1
	Dimensionless parameter of ideal MHD	13.1
ζ	Second viscosity coefficient	12.2
ζ_i	Interaction parameter	3.1
ζ_p	Plasma parameter	3.1
η	First viscosity coefficient (dynamic viscosity)	12.2
θ	Pitch-angle	5.1
	Angle between a wave vector and the magnetic field	15.1
κ_e	Classical electron conductivity	8.3
λ	Mean free path	8.1
$\ln A$	Coulomb logarithm	8.1
ν	Collisional frequency	8.1
ν	Kinematic viscosity	12.2
ν_{ei}	Electron-ion mean collisional frequency	11.1
ν_{kl}	Mean collisional frequency	9.1
ν_m	Magnetic diffusivity	12.2
ξ	Column depth	8.3
$\pi_{\alpha\beta}^{(k)}$	Viscous stress tensor	9.1
$\Pi_{\alpha\beta}^*$	Total momentum flux density tensor	12.2
ρ	Plasma mass density	9.1
ρ_k	Mass density for particles of kind k	9.1
ρ^q	Electric charge density	1.2
ρ_k^q	Charge density due to particles of kind k	9.1
$\boldsymbol{\rho}$	Rotational motion vector	5.2
σ	Isotropic electric conductivity	11.1
σ_H	Hall conductivity	11.1
$\sigma_{\parallel}$	Conductivity parallel to the magnetic field	11.1

$\sigma_{\perp}$	Conductivity perpendicular to the magnetic field	11.1
$\sigma_{\alpha\beta}^v$	Viscous stress tensor	12.2
τ	Characteristic time scale	5.2
τ_{ee}	Electron collisional time	8.3
τ_{ei}	Electron-ion collisional time	8.3
τ_{ii}	Ion collisional time	8.3
ϕ	Gravitational potential	1.2
φ	Electrostatic potential	8.2
φ	Angle in the spherical frame	14.4
ϕ, φ	Angle in the cylindrical frame	19.2
$\hat{\phi}_k$	Deviation of the exact distribution function	
	From an averaged distribution function	2.2
Φ	Magnetic flux	14.2
	Stream function	14.4
χ	Deflection angle	8.1
ψ	Angle to the x axis	14.4
	Potential of an electric current	20.3
Ψ	Potential of a current-free magnetic field	13.1
ω	Wave frequency	15.1
ω_0	Wave frequency in a moving frame of reference	15.1
ω_B	Cyclotron or Larmor frequency	5.1
ω_{pl}	Electron plasma frequency	8.2
Ω	Gravitational energy	19.1
ω	Vector of angular velocity	20.1

Appendix B

Useful Expressions

Source formulae

Larmor frequency of a non-relativistic electron (5.11), (5.51)

$$\omega_{\text{B}}^{(\text{e})} = \frac{eB}{m_{\text{e}}c} \approx 1.76 \times 10^7 B \text{ (G)}, \text{ rad s}^{-1}.$$

Larmor frequency of a non-relativistic proton (5.52)

$$\omega_{\text{B}}^{(\text{p})} = \frac{eB}{m_{\text{p}}c} \approx 9.58 \times 10^3 B \text{ (G)}, \text{ rad s}^{-1}.$$

Larmor radius of a non-relativistic electron (5.14), (5.59)

$$r_{\text{L}}^{(\text{e})} = \frac{m_{\text{e}}c v_{\perp}}{eB} \approx 5.69 \times 10^{-8} \frac{v_{\perp} \text{ (cm s}^{-1}\text{)}}{B \text{ (G)}}, \text{ cm}.$$

Larmor radius of a non-relativistic proton (5.14), (5.61)

$$r_{\text{L}}^{(\text{p})} \approx 1.04 \times 10^{-4} \frac{v_{\perp} \text{ (cm s}^{-1}\text{)}}{B \text{ (G)}}, \text{ cm}.$$

Mean thermal velocity of electrons in plasma (5.54)

$$V_{\text{Te}} = \left(\frac{3k_{\text{B}} T_{\text{e}}}{m_{\text{e}}} \right)^{1/2} \approx 6.74 \times 10^5 \sqrt{T_{\text{e}} \text{ (K)}}, \text{ cm s}^{-1}.$$

Mean thermal velocity of protons in plasma (5.55)

$$V_{\text{Tp}} \approx 1.57 \times 10^4 \sqrt{T_{\text{p}} \text{ (K)}}, \text{ cm s}^{-1}.$$

Larmor radius of non-relativistic *thermal* electrons (5.56)

$$r_L^{(e)} = \frac{V_{Te}}{\omega_B^{(e)}} \approx 3.83 \times 10^{-2} \frac{\sqrt{T_e(\text{K})}}{B(\text{G})}, \text{ cm}.$$

Larmor radius of non-relativistic *thermal* protons (5.57)

$$r_L^{(p)} = \frac{V_{Tp}}{\omega_B^{(p)}} \approx 1.64 \frac{\sqrt{T_p(\text{K})}}{B(\text{G})}, \text{ cm}.$$

Drift velocity (5.20)

$$\mathbf{v}_d = \frac{c}{e} \frac{\mathbf{F} \times \mathbf{B}}{B^2}.$$

Magnetic moment of a particle on the Larmor orbit (6.6)

$$\mathcal{M} = \frac{1}{c} JS = \frac{e \omega_B r_L^2}{2c} = \frac{p_\perp^2}{2mB} = \frac{\mathcal{E}_\perp}{B}.$$

Debye-Hückel radius ($T_e = T$, $T_i = 0$ or $T_e \gg T_i$) (8.33)

$$r_{\text{DH}} = \left(\frac{k_B T}{4\pi n e^2} \right)^{1/2}.$$

Debye-Hückel radius in electron-proton thermal plasma ($T_e = T_p = T$) (8.82)

$$r_{\text{DH}} = \left(\frac{k_B T}{8\pi e^2 n} \right)^{1/2} \approx 4.9 \left(\frac{T}{n} \right)^{1/2}, \text{ cm}.$$

Coulomb logarithm (8.80)

$$\ln \Lambda = \ln \left[\left(\frac{3k_B^{3/2}}{2\pi^{1/2} e^3} \right) \left(\frac{T_e^3}{n_e} \right)^{1/2} \right] \approx \ln \left[1.25 \times 10^4 \left(\frac{T_e^3}{n_e} \right)^{1/2} \right].$$

Electron plasma frequency (8.83)

$$\omega_{pl}^{(e)} = \left(\frac{4\pi e^2 n_e}{m_e} \right)^{1/2} \approx 5.64 \times 10^4 \sqrt{n_e}, \text{ rad s}^{-1}.$$

Thermal electron collisional time (8.85)

$$\tau_{ee} = \frac{m_e^2}{0.714 e^4 8\pi \ln \Lambda} \frac{V_{Te}^3}{n_e} \approx 4.04 \times 10^{-20} \frac{V_{Te}^3}{n_e}, \text{ s},$$

$$\tau_{ee} \approx 1.24 \times 10^{-2} \frac{T_e^{3/2}}{n_e}, \text{ s.}$$

Thermal proton collisional time (8.86)

$$\tau_{pp} = \frac{m_p^2}{0.714 e^4 8\pi \ln \Lambda} \frac{V_{Tp}^3}{n_p} \approx 1.36 \times 10^{-13} \frac{V_{Tp}^3}{n_p}, \text{ s.},$$

$$\tau_{pp} \approx 5.26 \times 10^{-1} \frac{T_p^{3/2}}{n_p}, \text{ s.}$$

Electron-ion collision (energy exchange) time Sect. 8.3

$$\tau_{ei}(\mathcal{E}) = \frac{m_e m_i [3k_B (T_e/m_e + T_i/m_i)]^{3/2}}{e_e^2 e_i^2 (6\pi)^{1/2} 8 \ln \Lambda}.$$

Time of energy exchange between electrons and protons (8.44)

$$\tau_{ep}(\mathcal{E}) \approx 22 \tau_{pp} \approx 950 \tau_{ee}.$$

Rate of energy exchange between electrons and ions (8.49)

$$\mathcal{Q}_{ei}(n, T_e, T_i) = \frac{4e^4 (2\pi m_e)^{1/2}}{m_i k_B^{1/2}} \ln \Lambda n^2 \frac{(T_e - T_i)}{T_e^{3/2}}.$$

Rate of energy exchange between electrons and protons (8.49)

$$\mathcal{Q}_{ep}(n, T_e, T_p) \approx 8.2 \times 10^{-19} \ln \Lambda n^2 \frac{(T_e - T_p)}{T_e^{3/2}}, \text{ erg cm}^{-3} \text{ s}^{-1}.$$

Dreicer field (8.90)

$$E_{Dr} = \frac{4\pi e^3 \ln \Lambda}{k_B} \frac{n_e}{T_e} \approx 6.54 \times 10^{-8} \frac{n_e}{T_e}, \text{ V cm}^{-1}.$$

Conductivity of magnetized plasma Sect. 11.3

$$\sigma_{\parallel} = \sigma = \frac{e^2 n}{m_e} \tau_{ei} \approx 2.53 \times 10^8 n (\text{cm}^{-3}) \tau_{ei} (\text{s}), \text{ s}^{-1},$$

$$\sigma_{\perp} = \sigma \frac{1}{1 + \left(\omega_B^{(e)} \tau_{ei}\right)^2}, \quad \sigma_H = \sigma \frac{\omega_B^{(e)} \tau_{ei}}{1 + \left(\omega_B^{(e)} \tau_{ei}\right)^2}.$$

Magnetic diffusivity (or viscosity) (12.50)

$$\nu_m = \frac{c^2}{4\pi\sigma} \approx 7.2 \times 10^{19} \frac{1}{\sigma}, \text{ cm}^2 \text{ s}^{-1}.$$

Magnetic Reynolds number (12.64)

$$\text{Re}_m = \frac{L^2}{\nu_m \tau} = \frac{\nu L}{\nu_m}.$$

Alfvén speed (13.15), (13.35)

$$V_A = \frac{B}{\sqrt{4\pi\rho}} \approx 2.18 \times 10^{11} \frac{B}{\sqrt{n}}, \text{ cm s}^{-1}.$$

Sound speed in electron-proton plasma (16.99)

$$V_s = \left(\gamma_g \frac{p}{\rho} \right)^{1/2} \approx 1.66 \times 10^4 \sqrt{T(\text{K})}, \text{ cm s}^{-1}.$$

Electric field in magnetized plasma (19.82)

$$E \approx \frac{1}{c} \nu B \approx 10^{-8} \nu (\text{cm s}^{-1}) B (\text{G}), \text{ V cm}^{-1}.$$

Appendix C

Constants

Fundamental Physical Constants

Speed of light	c	$2.998 \times 10^{10} \text{ cm s}^{-1}$
Electron charge	e	$4.802 \times 10^{-10} \text{ CGSE}$
Electron mass	m_e	$9.109 \times 10^{-28} \text{ g}$
Proton mass	m_p	$1.673 \times 10^{-24} \text{ g}$
Boltzmann constant	k_B	$1.381 \times 10^{-16} \text{ erg K}^{-1}$
Gravitational constant	G	$6.673 \times 10^{-8} \text{ dyne cm}^2 \text{ g}^{-2}$
Planck's constant	h	$6.625 \times 10^{-27} \text{ erg s}$

Some Useful Constants and Units

Ampere (current)	A	$3 \times 10^9 \text{ CGSE}$
Angström (length)	Å	10^{-8} cm
Electron volt (energy)	eV	$1.602 \times 10^{-12} \text{ erg}$
	eV	11,605 K
Gauss (magnetic induction)	G	$3 \times 10^{10} \text{ CGSE}$
Henry (inductance)	H	$1.111 \times 10^{-12} \text{ s}^2 \text{ cm}^{-1}$
Ionization potential of hydrogen		13.60 eV
Joule (energy)	J	10^7 erg
Maxwell (magnetic flux)	M	$3 \times 10^{10} \text{ CGSE}$
Ohm (resistance)	Ω	$1.111 \times 10^{-12} \text{ s cm}^{-1}$
Tesla (magnetic induction)		10^4 G
Volt (potential)	V	$3.333 \times 10^{-3} \text{ CGSE}$
Watt (power)	W	10^7 erg s^{-1}
Weber (magnetic flux)	Wb	10^8 Maxwell

Some Astrophysical Constants

Astronomical unit	AU	$1.496 \times 10^{13} \text{ cm}$
Mass of the Sun	$M_{\odot}$	$1.989 \times 10^{33} \text{ g}$
Mass of the Earth	M_{E}	$5.976 \times 10^{27} \text{ g}$
Parsec	pc	$3.086 \times 10^{18} \text{ cm}$
Radius of the Earth	R_{E}	$6.371 \times 10^8 \text{ cm}$
Solar radius	$R_{\odot}$	$6.960 \times 10^{10} \text{ cm}$
Solar surface gravity	$g_{\odot}$	$2.740 \times 10^4 \text{ cm s}^{-2}$
Solar luminosity	$L_{\odot}$	$3.827 \times 10^{33} \text{ erg s}^{-1}$
Mass loss rate	$\dot{M}_{\odot}$	10^{12} g s^{-1}
Rotation period of the Sun	$T_{\odot}$	26 days (at equator)

References

- Acton, L.: Coronal structures, local and global. In: Uchida, Y., Kosugi, T., Hudson, H. (eds.) *Magnetohydrodynamic Phenomena in the Solar Atmosphere: Prototypes of Stellar Magnetic Activity*, pp. 3–11. Kluwer Academic, Dordrecht (1996) [Sect. [19.3.4](#)]
- Akhiezer, A.I., Lyubarskii, G.Ya., Polovin, R.V.: On the stability of shock waves in MHD. *Sov. Phys. JETP* **8**(3), 507–512 (1959) [Sect. [17.2.1](#)]
- Akhiezer, A.I., Akhiezer, I.A., Polovin, R.V., et al.: *Plasma Electrodynamics*. Oxford, Pergamon (1975) [Sects. [15.4.5](#), [17.2.2](#)]
- Alekseyev, I.I., Kropotkin, A.P.: Passage of energetic particles through a MHD discontinuity. *Geomagn. Aeron.* **10**(6), 755–758 (1970) [Sect. [18.3.1](#)]
- Alexander, D., Daou, A.G.: Saturation of nonthermal hard X-ray emission in solar flares. *Astrophys. J.* **666**(2), 1268–1276 (2007) [Sects. [4.5.6](#), [4.6](#)]
- Alfaro, E.J., Pérez, E., Franco, J. (eds.): *How does the Galaxy work? A galactic tertulia with Don Cox and Ron Reynolds*. Kluwer Academic, Dordrecht (2004) [Sect. [9.8](#)]
- Alfvén, H.: On the solar origin of cosmic radiation. *Phys. Rev.* **75**(11), 1732–1735 (1949) [Sect. [7.2](#)]
- Alfvén, H.: *Cosmic Electrodynamics*, p. 228. Clarendon Press, Oxford (1950) [Intr., Sects. [12.2.2](#), [13.4](#), [15.2.2](#), [20.1.4](#)]
- Alfvén, H.: *Cosmic Plasma*, p. 164. D. Reidel Publishers, Dordrecht (1981) [Sect. [20.1.4](#)]
- Alfvén, H., Fälthammar, C.-G.: *Cosmic Electrodynamics*, p. 228. Clarendon Press, Oxford (1963) [Sects. [8.1.4](#), [8.2.3](#), [11.1](#), [15.4.5](#)]
- Allred, J.C.; Hawley, S.L., Abbott, W.P., Carlsson, M.: Radiative hydrodynamic models of the optical and ultraviolet emission from solar flares. *Astrophys. J.* **630**(1), 573–586 (2005) [Sect. [8.3.2](#)]
- Alperovich, L.S., Fedorov, E.N.: *Hydromagnetic Waves in the Magnetosphere and Ionosphere*, p. 426. Springer, Berlin (2007) [Sects. [11.1](#), [11.4.1](#)]
- Altynets, A.T., Krasov, V.I., Tomozov V.M.: Magnetic field dissipation in neutral current sheets. *Solar Phys.* **55**(1), 69–81 (1977) [Sect. [12.3.1](#)]
- Anderson, J.E.: *Magnetohydrodynamic Shock Waves*, p. 226. MIT, Cambridge (1963) [Sects. [16.2.4\(c\)](#), [17.4.2](#)]
- Andres, U.T., Polak, L.S., Syrovatskii, S.I.: Electromagnetic expulsion of spherical bodies from a conductive fluid. *Soviet Phys. Tech. Phys.* **8**(3), 193–196 (1963) [Sects. [19.4.2](#), [20.4](#)]
- Anile, A.M.: *Relativistic Fluids and Magneto-Fluids*, p. 336. Cambridge University Press, Cambridge (1989) [Sect. [12.2](#)]
- Antonucci, E., Somov, B.V.: A diagnostic method for reconnecting magnetic fields in the solar corona. In: *Coronal Streamers, Coronal Loops, and Coronal and Solar Wind Composition, Proceedings of First SOHO Workshop, ESA SP-348*, pp. 293–294 (1992) [Sects. [8.3.3](#), [20.4](#)]

- Antonucci, E., Benna, C., Somov, B.V.: Interpretation of the observed plasma ‘turbulent’ velocities as a result of reconnection in solar flares. *Astrophys. J.* **456**(2), 833–839 (1996) [Sects. 8.3.3, 20.4]
- Arons, J.: Pulsar emission: Where to go? In: Becker, W. (ed.) *Neutron Stars and Pulsars*, pp. 373–420. Springer-Verlag, Berlin, Heidelberg (2009) [Sect. 12.2.5]
- Aschwanden, M.J.: Particle Acceleration and Kinematics in Solar Flares: A Synthesis of Recent Observations and Theoretical Concepts, p. 227. Kluwer Academic, Dordrecht (2002) [Sect. 4.5.7]
- Aschwanden, M.J.: Physics of the Solar Corona: An Introduction, p. 227. Springer, Berlin (2004) [Sect. 15.5]
- Aschwanden, M.J., Kliem, B., Schwarz, U., et al.: Wavelet analysis of solar flare hard X-rays. *Astrophys. J.* **505**(2), 941–956 (1998) [Sect. 4.5.7]
- Aschwanden, M.J., Nightingale, R.W., Andries, J., et al.: Observational tests of damping by resonant absorption in coronal loop oscillations. *Astrophys. J.* **598**, 1375–1386 (2003) [Sect. 15.5]
- Asmussen, S., Glynn, P.W.: *Stochastic Simulations: Algorithms and Analysis*, p. 476. Springer, New York (2007) [Sect. 3.4]
- Atkinson, G., Unti, T.: Two-dimensional Chapman-Ferraro problem with neutral sheet. 1. The interior field. *J. Geophys. Res. Space Phys.* **74**(14), 3713–3716 (1969) [Sect. 14.2.2(a)]
- Atoyan, A.M., Aharonian, F.A.: Modeling of the non-thermal flares in the Galactic microquasar GRS 1915+105. *Mon. Not. Roy. Astron. Soc.* **302**(1), 253–276 (1999) [Sect. 20.1.3]
- Axford, W.I., Leer, E., Skadron, G.: The acceleration of cosmic rays by shock waves. In: *Proc. 15th Int. Cosmic Ray Conf.* (Plovdiv, August 13–26, 1977), Bulgarian Acad. Sci., Sofia, vol. 11, pp. 132–137 (1977) [Sect. 18.2.1]
- Bachiller, R.: Bipolar molecular outflows from young stars and protostars. *Ann. Rev. Astron. Astrophys.* **34**, 111–154 (1996) [Sect. 20.2]
- Bai, T., Hudson, H.S., Pelling, R.M., et al.: First-order Fermi acceleration in solar flares as a mechanism for the second-step acceleration of protons and electrons. *Astrophys. J.* **267**(1), 433–441 (1983) [Sect. 6.2.4]
- Balbus, S.A., Papaloizou, J.C.B.: On the dynamical foundations of α disks. *Astrophys. J.* **521**(2), 650–658 (1999) [Sect. 13.2.1]
- Balescu, R.: *Statistical Mechanics of Charged Particles*, p. 477. Wiley, London (1963) [Sect. 4.1.2]
- Balescu, R.: *Equilibrium and Nonequilibrium Statistical Mechanics*. Wiley, London (1975) [Sect. 3.1.4]
- Balescu, R.: *Transport Processes in Plasmas*. North-Holland, Amsterdam (1988) [Sect. 9.5]
- Balikhin, M., Gedalin, M., Petrukovich, A.: New mechanism for electron heating in shocks. *Phys. Rev. Lett.* **70**, 1259–1262 (1993) [Sect. 18.3.2(a)]
- Balogh, A., Erdös, G.: Fast acceleration of ions at quasi-perpendicular shocks. *J. Geophys. Res.* **96**(A9), 15853–15862 (1991) [Sect. 18.3.2(b)]
- Barenblatt, G.I.: *Similarity, Self-Similarity, and Intermediate Asymptotics*. Plenum, New York (1979) [Sect. 20.4]
- Becker, W. (ed.): *Neutron Stars and Pulsars*, p. 997. Springer, Berlin (2009) [Sects. 5.4, 7.3, 12.2.2]
- Bednarek, W., Protheroe, R.J.: Gamma-ray and neutrino flares produced by protons accelerated on an accretion disc surface in active galactic nuclei. *Mon. Not. Royal Astron. Soc.* **302**, 373–380 (1999) [Sect. 13.2.4]
- Begelman, M.C., Blandford, R.D., Rees, M.J.: Theory of extragalactic radio sources. *Rev. Mod. Phys.* **56**(2), 255–351 (1984) [Sects. 7.3, 13.3.1, 13.3.3, 20.1.3]
- Beloborodov, A.M.: Plasma ejection from magnetic flares and the X-ray spectrum of Cygnus X-1. *Astrophys. J.* **510**, L123–L126 (1999) [Sect. 13.2.4]
- Benz, A.: *Plasma Astrophysics: Kinetic Processes in Solar and Stellar Coronae*, 2nd edn., p. 299. Kluwer Academic, Dordrecht (2002) [Sects. 3.1.2, 7.1]
- Bernstein, I.B., Frieman, E.A., Kruskal, M.D., et al.: An energy principle for hydromagnetic stability problems. *Proc. Roy. Soc.* **244**(A1), 17–40 (1958) [Sect. 19.3.4]
- Bertin, G.: *The Dynamics of Galaxies*, p. 448. Cambridge University Press, Cambridge (1999). [Sects. 1.3, 9.8]

- Bethe, H.A.: Office of Scientific Research and Development, Rep. No. 445 (1942) [Sect. 17.1.1]
- Bezrodnykh, S.I., Vlasov, V.I., Somov, B.V.: Analytical model of magnetic reconnection in the presence of shock waves attached to a current sheet. *Astron. Lett.* **33**(2), 130–136 (2007) [Sect. 14.2.2(a)]
- Bezrodnykh, S.I., Vlasov, V.I., Somov, B.V.: Generalized analytical models of Syrovatskii's current sheet. *Astron. Lett.* **37**(2), 113–130 (2011) [Sect. 14.2.2(a)]
- Bhatnagar, P.L., Gross, E.P., Krook, M.: A model for collision processes in gases. 1. Small amplitude processes in charged and neutral one-component systems. *Phys. Rev.* **94**(3), 511–525 (1954) [Sect. 9.9]
- Bhattacharjee, A.: Impulsive magnetic reconnection in the Earth's magnetotail and the solar corona. *Ann. Rev. Astron. Astrophys.* **42**, 365–384 (2004) [Sect. 11.4.2]
- Bianchini, A., Della Valle, M., Orio, M. (eds.): *Cataclysmic Variables*, p. 540. Kluwer Academic, Dordrecht (1995) [Sect. 13.2.2]
- Binney, J., Tremaine, S.: *Galactic Dynamics*. Princeton University Press, New Jersey (1987) [Sects. 3.3.1, 8.5]
- Birkinkshaw, M.: Instabilities in astrophysical jets. In: de Gouveia Dal Pino, E.M., et al. (eds.) *Advanced Topics on Astrophysical and Space Plasmas*, pp. 17–91. Kluwer Academic, Dordrecht (1997) [Sect. 13.3.1]
- Biskamp, D., Welter, H.: Magnetic arcade evolution and instability. *Solar Phys.* **120**(1), 49–77 (1989) [Sect. 19.4.3]
- Blackman, E.G.: On particle energization in accretion flow. *Mon. Not. Roy. Astron. Soc.* **302**(4), 723–730 (1999) [Sect. 8.3.5]
- Blackman, E.G., Field, G.B.: Constraints on the magnitude of α in dynamo theory. *Astrophys. J.* **534**(2), 984–988 (2000) [Sect. 13.1.2]
- Blandford, R.D.: Particle acceleration mechanisms. *Astrophys. J. Suppl.* **90**(2), 515–520 (1994) [Sects. 18.1, 18.2.1]
- Bliokh, P., Sinitsin, V., Yaroshenko, V.: *Dusty and Self-Gravitational Plasmas in Space*, p. 250. Kluwer Academic, Dordrecht (1995) [Sect. 1.2.4]
- Blokhintsev, D.I.: Moving receiver of sound. *Doklady Akademii Nauk SSSR (Soviet Physics Doklady)*, **47**(1), 22–25 (in Russian) (1945) [Sect. 15.2.1]
- Bobrova, N.A., Syrovatskii, S.I.: Singular lines of 1D force-free field. *Solar Phys.* **61**(2), 379–387 (1979) [Sect. 19.2.1.1]
- Bocquet, M., Bonazzola, S.,ourgoulhon, E., et al.: Rotating neutron star models with a magnetic field. *Astron. Astrophys.* **301**(3), 757–775 (1995) [Sect. 19.1.3]
- Bodmer, R., Bochsler, P.: Influence of Coulomb collisions on isotopic and elemental fractionation in the solar wind. *J. Geophys. Res.* **105**(A1), 47–60 (2000) [Sects. 8.4.1(b), 10.1]
- Bogachev, S.A., Somov, B.V.: Effect of Coulomb collisions on the particle acceleration in collapsing magnetic traps. *Astron. Lett.* **35**(1), 57–69 (2009) [Sect. 8.1.4]
- Bogdanov, S.Yu., Frank, A.G., Kyrei, N.P., et al.: Magnetic reconnection, generation of plasma fluxes and accelerated particles in laboratory experiments. *Plasma Astrophys. ESA SP-251*, 177–183 (1986) [Sect. 12.3.1]
- Bogdanov, S.Yu., Kyrei, N.P., Markov, V.S., et al.: Current sheets in magnetic configurations with singular X-lines. *JETP Lett.* **71**(2), 78–84 (2000) [Sect. 12.3.1]
- Bogoliubov, N.N.: *Problems of a Dynamical Theory in Statistical Physics*. State Technical Press, Moscow (in Russian) (1946) [Sect. 2.4]
- Bolcato, R., Etay, J., Fautrelle, Y., et al.: Electromagnetic billiards. *Phys. Fluids* **5**(A7), 1852–1853 (1993) [Sect. 20.5]
- Boltzmann, L.: *Sitzungsber. Kaiserl. Akad. Wiss. Wien.* **66**, 275–284 (1872) [Sects. 3.5, 9.6.1]
- Boltzmann, L.: *Lectures on the Theory of Gases*. Gostehizdat, Moscow (in Russian) (1956) [Sects. 3.5, 9.6.1]
- Bondi, H.: On spherical symmetrical accretion. *Mon. Not. Roy. Astron. Soc.* **112**(1), 195–204 (1952) [Sect. 13.2.3]
- Bontemps, S., André, P., Terebey, S., et al.: Evolution of outflow activity around low-mass embedded young stellar objects. *Astron. Astrophys.* **311**, 858–875 (1996) [Sect. 20.2]

- Born, M., Green, H.S.: A General Kinetic Theory of Liquids. Cambridge University Press, Cambridge (1949) [Sect. 2.4]
- Bradt, H.: Astrophysics Processes, p. 504. Cambridge University Press, Cambridge (2008) [Sects. 3.1.1, 9.5.2, 13.2.3]
- Braginskii, S.I.: Transport processes in plasma. In: Leontovich, M. (ed.) Reviews of Plasma Physics, vol. 1, pp. 205–311. Consultants Bureau, New York (1965) [Sects. 8.3.2, 9.6, 10.5, 11.4.2]
- Bridgman, P.W.: Dimensional Analysis, p. 113. Yale University Press, New Haven (1931) [Sect. 20.4]
- Broderick, A., Prakash, M., Lattimer, J.M.: The equation of state of neutron star matter in strong magnetic fields. *Astrophys. J.* **537**(1), 351–367 (2000) [Sect. 19.1.3]
- Brown, J.C.: The deduction of energy spectra of non-thermal electrons in flares from the observed dynamic spectra of hard X-ray bursts. *Solar Phys.* **18**(2), 489–502 (1971) [Sects. 4.3.4, 8.1.5]
- Brown, J.C.: The directivity and polarization of thick target X-ray bremsstrahlung from flares. *Solar Phys.* **26**(2), 441–459 (1972) [Sects. 4.4.1, 4.4.2]
- Brown, J.C., McArthur, G.K., Barrett, R.K., et al.: Inversion of the thick-target bremsstrahlung spectra from non-uniformly ionized plasmas. *Solar Phys.* **179**(2), 379–404 (1998a) [Sect. 4.5.7]
- Brown, J.C., Conway, A.J., Aschwanden, M.J.: The electron injection function and energy-dependent delays in thick-target hard X-rays. *Astrophys. J.* **509**(2), 911–917 (1998b) [Sect. 4.5.7]
- Brown, J.C., Emslie, A.G., Kontar, E.P.: The determination and use of mean electron flux spectra in solar flares. *Astrophys. J.* **595**(2), L115–L117 (2003) [Sect. 4.5.7]
- Bykov, A.M., Chevalier, R.A., Ellison, D.C., et al.: Non-thermal emission from a supernova remnant in a molecular cloud. *Astrophys. J.* **538**(1), 203–216 (2000) [Sect. 8.4.1(b)]
- Cadjan, M.G., Ivanov, M.F.: Langevin approach to plasma kinetics with collisions. *J. Plasma Phys.* **61**(1), 89–106 (1999) [Sect. 3.4]
- Cai, H.J., Lee, L.C.: The generalized Ohm’s law in collisionless reconnection. *Phys. Plasmas* **4**(3), 509–520 (1997) [Sect. 1.2.4]
- Camenzind, M.: Magnetic fields and the physics of active galactic nuclei. *Rev. Mod. Astron.* **8**, 201–233 (1995) [Sect. 13.3.3]
- Campbell, C.G.: Magnetohydrodynamics of Binary Stars, p. 306. Kluwer Academic, Dordrecht (1997) [Sect. 13.2.1]
- Cassak, P.A., Drake, J.F., Shay, M.A., et al.: Onset of fast magnetic reconnection. *Phys. Rev. Lett.* **98**(21), id. 215001 (2007) [Sect. 11.4.2]
- Cercignani, C.: Mathematical Methods in Kinetic Theory. MacMillan, London (1969) [Sect. 3.5]
- Chakrabarti, S.K. (ed.): Observational Evidence for Black Holes in the Universe, p. 399. Kluwer Academic, Dordrecht (1999) [Sect. 8.3.5]
- Chandrasekhar, S.: Stochastic problems in physics and astronomy. *Rev. Mod. Phys.* **15**(1), 1–89 (1943a) [Sects. 3.1.4, 8.1.5, 8.3.1]
- Chandrasekhar, S.: Dynamical friction. 1. General considerations. *Astrophys. J.* **97**(1), 255–262 (1943b) [Sects. 3.1.4, 8.3.1, 8.5]
- Chandrasekhar, S.: Dynamical friction. 2. The rate of escape of stars from clusters and the evidence for the operation of dynamic friction. *Astrophys. J.* **97**(1), 263–273 (1943c) [Sects. 8.3.1, 8.5]
- Chandrasekhar, S.: Hydrodynamic and Hydromagnetic Stability, p. 654. Dover, New York (1981) [Sects. 19.1.2, 19.3.4]
- Chandrasekhar, S., Fermi, E.: Problems of gravitational stability in the presence of a magnetic field. *Astrophys. J.* **118**(1), 116–141 (1953) [Sect. 19.1.1]
- Cherenkov, P.A.: C. R. Acad. Sci. U.S.S.R. **8**, 451 (in Russian) (1934) [Sect. 7.4]
- Cherenkov, P.A.: Visible radiation produced by electrons moving in a medium with velocities exceeding that of light. *Phys. Rev.* **52**, 378–379 (1937) [Sect. 7.4]
- Chernov, A.A., Yan’kov, V.V.: Electron flow in low-density pinches. *Soviet J. Plasma Phys.* **8**(5), 522–528 (1982) [Sect. 20.4]

- Chew, G.F., Goldberger, M.L., Low, F.E.: The Boltzmann equation and the one-fluid hydromagnetic equations in the absence of particle collisions. *Proc. Roy. Soc. Lond.* **A236**(1), 112–118 (1956) [Sects. 5.2.1, 11.5.1, 16.4]
- Choudhuri, A.R.: *The Physics of Fluids and Plasmas: An Introduction for Astrophysicists*, p. 427. Cambridge University Press, Cambridge (1998) [Intr., Sect. 19.1.2]
- Ciufolini, I., Matzner, R.A. (eds.): *General Relativity and John Archibald Wheeler*, p. 545. Springer Science+Business Media B.V., Dordrecht (2010) [Sect. 13.3.2]
- Clarke, C., Carswell, P.: *Principles of Astrophysical Fluid Dynamics*, p. 226. Cambridge University Press, Cambridge (2007) [Sect. 9.5.2]
- Clausius, R.: On a mechanical theorem applicable to heat. *Phil. Mag. (Series 4)* **40**(1), 122–127 (1870) [Sect. 19.1.1]
- Cole, J.D., Huth, J.H.: Some interior problems of hydromagnetics. *Phys. Fluids* **2**(6), 624–626 (1959) [Sect. 14.5]
- Collins, G.W.: *The Virial Theorem in Stellar Astrophysics*. Pachart, Tucson (1978) [Sect. 19.1.1]
- Colpi, M., Casella, P., Gorini, V., et al. (eds.): *Physics of Relativistic Objects in Compact Binaries: From Birth to Coalescence*, Springer, Dordrecht (2009) [Sect. 12.2.2]
- Coppi, B., Laval, G., Pellat, R.: Dynamics of the geomagnetic tail. *Phys. Rev. Lett.* **6**(26), 1207–1210 (1966) [Sect. 3.1.2]
- Courant, R., Friedrichs, K.O.: *Supersonic Flow and Shock Waves*, p. 464. Springer, New York (1985) [Sect. 17.1.1]
- Cowling, T.G.: *Magnetohydrodynamics*, p. 135. Adam Hilger, Bristol (1976) [Sect. 11.6]
- Cox, D.P., Tucker, W.H.: Ionization equilibrium and radiative cooling of a low-density plasma. *Astrophys. J.* **157**(3), 1157–1167 (1969) [Sects. 12.1.3, 15.4.1]
- Cromwell, D., McQuillan, P., Brown, J.C.: Beam-driven return current instability and anomalous plasma heating in solar flares. *Solar Phys.* **115**(2), 289–312 (1988) [Sect. 4.5.6]
- Crooker, N., Joselyn, J.A., Feynman, J. (eds.): *Coronal Mass Ejections*, p. 299. American Geophysical Union, Washington (1997) [Intr.]
- Cumming, A., Arras, P., Zweibel, E.: Magnetic field evolution in neutron star crusts due to the Hall effect and ohmic decay. *Astrophys. J.* **609**, 999–1017 (2004) [Sect. 11.4.2]
- Cuperman, S., Dryer, M.: On the heat conduction in multicomponent, non-Maxwellian spherically symmetric solar wind plasmas. *Astrophys. J.* **298**, 414–420 (1985) [Sect. 9.6.2]
- Dadhich, N., Kembhavi, A. (eds.): *The Universe: Visions and Perspectives*, p. 346. Kluwer Academic, Dordrecht (2000) [Sect. 1.3]
- Darwin, C.: Source of the cosmic rays. *Nature* **164**, 1112–1114 (1949) [Sect. 18.1]
- Davidson, R.C.: *Theory of Nonneutral Plasmas*. W.A. Benjamin, London (1974) [Sect. 11.5.2]
- Davis, L.Jr.: Modified Fermi mechanism for the acceleration of cosmic rays. *Phys. Rev.* **101**, 351–358 (1956) [Sect. 6.2.4]
- de Hoffmann, F., Teller, E.: Magnetohydrodynamic shocks. *Phys. Rev.* **80**(4), 692–703 (1950) [Sects. 16.2.1, 16.2.4(a), 16.5]
- de Martino, D., Silvotti, R., Solheim, J.-E., et al. (eds.): *White Dwarfs*, p. 429. Kluwer Academic, Dordrecht (2003) [Sects. 1.4, 3.5]
- Debye, P., Hückel, E.: *Phys. Z.* **24**, 185 (1923) [Sect. 8.2.1]
- Decker, R.B.: Formation of shock-spike events in quasi-perpendicular shocks. *J. Geophys. Res.* **88**(A12), 9959–9973 (1983) [Sect. 18.3.2(a) (a)]
- Decker, R.B.: The role of magnetic loops in particle acceleration at nearly perpendicular shocks. *J. Geophys. Res.* **98**(A1), 33–46 (1993) [Sect. 18.3.2(b) (b)]
- Decker, R.B., Vlahos, L.: Numerical studies of particle acceleration at turbulent, oblique shocks with an application to prompt ion acceleration during solar flares. *Astrophys. J.* **306**(2), 710–729 (1986) [Sect. 18.3.3]
- Diakonov, S.V., Somov, B.V.: Thermal electrons runaway from a hot plasma during a flare in the reverse-current model and their X-ray bremsstrahlung. *Solar Phys.* **116**(1), 119–139 (1988) [Sects. 4.5.2, 4.5.3, 4.5.5, 8.4.3, 9.7.3]

- Diakonov, S.V., Somov, B.V.: A thermal model with return current for source of hard X-ray radiation and microwave radiation of solar flare. *Kinematics Phys. Celes. Bodies* (Allerton Press, Inc.) **6**(1), 47–53 (1990) [Sect. 4.5.5]
- Diamond, P.H., Itoh, S.I., Itoh, K.: *Modern Plasma Physics*. Vol. 1: Physical Kinetics of Turbulent Plasmas, p. 417. Cambridge University Press, Cambridge (2010) [Intr., Sect. 3.1.2]
- Di Matteo, T., Celotti, A., Fabian, A.C.: Magnetic flares in accretion disc coronae and the spectral states of black hole candidates: The case of GX339-4. *Mon. Not. Roy. Astron. Soc.* **304**, 809–820 (1999) [Sect. 13.2.4]
- Di Matteo, T., Quataert, E., Allen, S.W., et al.: Low-radiative-efficiency accretion in the nuclei of elliptical galaxies. *Mon. Not. Roy. Astron. Soc.* **311**(3), 507–521 (2000) [Sect. 13.2.3]
- Di Matteo, T., Johnstone, R.M., Allen, S.W., et al.: Accretion onto nearby supermassive black holes: *Chandra* constraints on the dominant cluster galaxy NGC 6166. *Astrophys. J.* **550**(1), L19–L23 (2001) [Sect. 13.2.3]
- Dokuchaev, V.P.: Emission of magnetoacoustic waves in the motion of stars in cosmic space. *Sov. Astron. AJ* **8**(1), 23–31 (1964) [Sect. 15.6]
- Dorman, L.: *Cosmic Rays in Magnetospheres of the Earth and other Planets*, p. 770. Springer Science+Business Media B.V., Dordrecht (2009) [Sect. 5.1.3]
- Drake, J.F., Kleva R.G.: Collisionless reconnection and the sawtooth crash. *Phys. Rev. Lett.* **66**(11), 1458–1461 (1991) [Sect. 11.2]
- Dreicer, H.: Electron and ion runaway in a fully ionized gas *Phys. Rev.* **115**(2), 238–249 (1959) [Sects. 8.4.2, 10.1]
- Duijveman, A., Somov, B.V., Spektor, A.R.: Evolution of a flaring loop after injection of fast electrons. *Solar Phys.* **88**(1), 257–273 (1983) [Sect. 8.3.2]
- Duncan, R.C., Thompson, C.: Formation of very strongly magnetized neutron stars: Implications for gamma-ray bursts. *Astrophys. J.* **392**(1), L9–L13 (1992) [Sect. 13.1.2]
- D'yakov, S.P.: *Zhurnal Exper. Teor. Fiz.* **27**, 288–297 (in Russian) (1954) [Sect. 17.5]
- Dyer, K.K., Reynolds, S.R., Borkowski, K.J., et al.: Separating thermal and non-thermal X-rays in supernova remnants. I. Total fits to SN 1006 AD. *Astrophys. J.* **551**(1), 439–453 (2001) [Sect. 18.2.1]
- Eichler, D.: Particle acceleration in solar flares by cyclotron damping of cascading turbulence. *Astrophys. J.* **229**(1), 413–418 (1979) [Sect. 6.2.4]
- Elperin, T., Golubev, I., Kleeorin, N., et al.: Large-scale instability in a sheared nonhelical turbulence: Formation of vortical structures. *Phys. Rev. E* **76**(6), id. 066310 (2007) [Sect. 13.1.2]
- Elsasser, W.M.: Hydromagnetic dynamo theory. *Rev. Mod. Phys.* **28**(2), 135–163 (1956) [Sects. 13.1.2, 20.1.5]
- Erdős, G., Balogh, A.: Drift acceleration at interplanetary shocks. *Astrophys. J. Suppl.* **90**(2), 553–559 (1994) [Sect. 18.3.2(b)]
- Everitt, C.W.F., DeBra, D.B., Parkinson, B.W., et al.: Gravity Probe B: Final results of a space experiment to test general relativity. *Phys. Rev. Lett.* **106**, 221101–221105 (2011) [Sect. 13.3.2]
- Falle, S.A., Komissarov, S.S.: On the inadmissibility of non-evolutionary shocks. *J. Plasma Phys.* **65**(1), 29–58 (2001) [Sect. 16.3]
- Fedoruk, V.M.: *Ordinary Differential Equations*. Nauka, Moscow (in Russian) (1985) [Sect. 17.4.1]
- Feldman, W.C., Bame, S.J., Gary, S.P., et al.: Electron heating within the Earth's bow shock. *Phys. Rev. Lett.* **49**, 199–202 (1982) [Sect. 18.3.2(a)]
- Ferencz, C., Ferencz, O.E., Hamar, D., et al.: *Whistler Phenomena*, p. 260. Kluwer Academic, Dordrecht (2001) [Sect. 7.1.3]
- Fermi, E.: On the origin of cosmic radiation. *Phys. Rev.* **75**, 1169–1174 (1949) [Sect. 6.2.4]
- Fermi, E.: Galactic magnetic fields and the origin of cosmic radiation. *Astrophys. J.* **119**(1), 1–6 (1954) [Sect. 6.2.4]
- Fernández, J.A.: *Comets: Nature, Dynamics, Origin, and their Cosmogonical Relevance*, p. 383. Springer, Dordrecht (2005) [Sect. 1.2.4]

- Feroci, M., Hurley, K., Duncan, R.C., et al.: The giant flare of 1998 August 27 from SGR 1900+14. I. An interpretive study of *BeppoSax* and *Ulysses* observations. *Astrophys. J.* **549**, 1021–1038 (2001) [Sect. 19.1.3]
- Field, G.B.: Thermal instability. *Astrophys. J.* **142**(2), 531–567 (1965) [Sects. 8.3.4, 9.4.3, 12.1.3, 15.4.5]
- Fokker, A.D.: Die mittlere Energie rotieren der elektrischer Dipole im Strahlungsfeld. *Ann. der Physik* **43**(5), 810–820 (1914) [Sect. 3.1.4]
- Fortov, V.E., Iakubov, I.T., Khrapak, A.G.: *Physics of Strongly Coupled Plasma*, p.376. Clarendon Press, Oxford (2006) [Sect. 3.1.1]
- Fox, D.C., Loeb, A.: Do the electrons and ions in X-ray clusters share the same temperature? *Astrophys. J.* **491**(2), 459–466 (1997) [Sect. 8.3.4]
- Freidberg, J.P.: *Plasma Physics and Fusion Energy*, p. 671. Cambridge University Press, Cambridge (2007) [Intr.]
- Galeev, A.A., Rosner, R., Vaiana, G.S.: Structured coronae of accretion discs. *Astrophys. J.* **229**(1), 318–326 (1979) [Sect. 13.2.4]
- Gedalin, M., Griv, E.: Collisionless electrons in a thin high Mach number shocks: Dependence on angle and β . *Ann. Geophysicae* **17**(10), 1251–1259 (1999) [Sects. 16.4, 18.3.2(a)]
- Gel'fand, I.M.: Some problems of the theory of quasilinear equations. *Usp. Mat. Nauk* **14**(2), 87–158 (in Russian) (1959) [Sect. 17.1.1]
- Gerbeth, G., Thess, A., Marty, P.: Theoretical study of the MHD flow around a cylinder in crossed electric and magnetic fields. *Eur. J. Mech. B/Fluids* **9**(3), 239–257 (1990) [Sects. 19.4.2, 20.3]
- Germain, P.: Shock waves and shock-wave structure in magneto-fluid dynamics. *Rev. Mod. Phys.* **32**(4), 951–958 (1960) [Sect. 17.4.2]
- Giacalone, J., Ellison, D.C.: Three-dimensional numerical simulations of particle injection and acceleration at quasi-perpendicular shocks. *J. Geophys. Res.* **105**(A6), 12541–12556 (2000) [Sects. 18.1, 18.3.2(b)]
- Gieseler, U.D.J., Kirk, J.G., Gallant, Y.A., et al.: Particle acceleration at oblique shocks and discontinuities of the density profile. *Astron. Astrophys.* **435**(1), 298–306 (1999) [Sect. 18.2.1]
- Gilman, P.A.: Fluid dynamics and MHD of the solar convection zone and tachocline. *Solar Phys.* **192**(1), 27–48 (2000) [Sect. 20.1.5]
- Ginzburg, V.L., Syrovatskii, S.I.: *The Origin of Cosmic Rays*. Pergamon Press, Oxford (1964) [Sect. 5.1.3]
- Ginzburg, V.L., Syrovatskii, S.I.: Cosmic magneto-bremsstrahlung (synchrotron) radiation. *Annu. Rev. Astron. Astrophys.* **3**, 297–350 (1965) [Sect. 5.4]
- Ginzburg, V.L., Zheleznyakov, V.V.: On the possible mechanisms of sporadic solar radio emission. *Sov. Astron. AJ* **2**(5), 653–668 (1958) [Sect. 7.1]
- Ginzburg, V., Landau, L., Leontovich, M., et al.: On the insolvency of the A.A. Vlasov works on general theory of plasma and solid-state matter. *Zhur. Eksp. Teor. Fiz.* **16**(3), 246–252 (in Russian) (1946) [Sect. 3.1.2]
- Giovannelli, R.G.: A theory of chromospheric flares. *Nature* **158**(4003), 81–82 (1946) [Sect. 12.4.1]
- Giovannelli, R.G.: Magnetic and electric phenomena in the Sun's atmosphere associated with sunspots. *Mon. Not. Roy. Astron. Soc.* **107**(4), 338–355 (1947) [Sect. 12.4.1]
- Giovannelli, R.G.: Electron energies resulting from an electric field in a highly ionized gas. *Phil. Mag. Seventh Series* **40**(301), 206–214 (1949) [Sect. 8.4.2]
- Gisler, G., Lemons, D.: Electron Fermi acceleration in collapsing magnetic traps: Computational and analytical models. *J. Geophys. Res.* **95**(A9), 14925–14938 (1990) [Sect. 18.3.2(b)]
- Glasstone, S., Loveberg, R.H.: *Controlled Thermonuclear Reactions*, p. 523. Van Nostrand, Princeton (1960) [Intr.]
- Gnedenko, B.V.: *A Course of Probability Theory*, 4th edn. Nauka, Moscow (in Russian) (1965) [Sect. 2.2.2]
- Golant, V.E., Zhilinskii, A.P., Sakharov, I.E.: *The Basis of Plasma Physics*. Atomizdat, Moscow (in Russian) (1977) [Sects. 9.3.2, 9.7.1, 9.7.2]
- Goldreich, P., Reisenegger, A.: Magnetic field decay in isolated neutron stars. *Astrophys. J.* **395**(1), 250–258 (1992) [Sect. 11.4.2]

- Goldreich, P., Sridhar, S.: Magnetohydrodynamic turbulence revisited. *Astrophys. J.* **485**(2), 680–688 (1997) [Sect. 7.2]
- Goldston, R.J., Rutherford, P.H.: *Introduction to Plasma Physics*, p. 492. Institute of Physics Publishing, Bristol (1995) [Intr.]
- Gombosi, T.I.: *Physics of the Space Environment*, p. 339. Cambridge University Press, Cambridge (1999) [Sect. 18.2.1]
- Gorbachev, V.S., Kel’ner, S.R.: Formation of plasma condensations in fluctuating strong magnetic field. *Sov. Phys. JETP* **67**(9), 1785–1790 (1988) [Sect. 14.4.1]
- Gosling, J.T.: Observations of magnetic reconnection in the turbulent high-speed solar wind. *Astrophys. J.* **671**(1), L73–L76 (2007) [Sect. 12.4.2]
- Gosling, J.T., Eriksson, S., McComas, D.J., et al.: Multiple magnetic reconnection sites associated with a coronal mass ejection in the solar wind. *Geophys. Res.* **112**(A8), CiteID A08106 (2007a) [Sect. 11.5.1]
- Gosling, J.T., Eriksson, S., Phan, T.D., et al.: Direct evidence for prolonged magnetic reconnection at a continuous X-line within the heliospheric current sheet. *Geophys. Res. Lett.* **34**(6), CiteID L06102 (2007b) [Sect. 12.4.2]
- Grad, H.: Note on N-dimensional Hermite polynomials. *Commun. Pure Appl. Math.* **2**(4), 325–330 (1949) [Sect. 9.7.1]
- Grad, H.: Reducible problems in magneto-fluid dynamic steady flows. *Rev. Mod. Phys.* **32**(4), 830–847 (1960) [Sect. 19.5]
- Grad, H., Rubin, H.: Hydromagnetic equilibria and force-free fields. *Proc. Second Int. Conf. on Peaceful Uses of Atomic Energy* **31**, 190–197 (1958) [Sect. 19.5]
- Grant, H.L., Stewart, R.W., Moilliet, A.: Turbulence spectra from a tidal channel. *J. Fluid Mech.* **12**, 241–248 (1962) [Sect. 7.2]
- Gritsyk, P.A., Somov, B.V.: The kinetic description of the accelerated-electron flux in solar flares. *Moscow Univ. Phys. Bull.* **66**(5), 466–472 (2011) [Sect. 4.5.5]
- Gurevich, A.V.: On the theory of runaway electrons. *Sov. Phys. JETP* **12**(5), 904–912 (1961) [Sect. 8.4.2]
- Gurevich, A.V., Istomin, Y.N.: Thermal runaway and convective heat transport by fast electrons in a plasma. *Sov. Phys. JETP* **50**(3), 470–475 (1979) [Sect. 8.4.3]
- Gurevich, A.V., Zhivlyuk, Y.N.: Runaway electrons in a non-equilibrium plasma. *Sov. Phys. JETP* **22**(1), 153–159 (1966) [Sect. 4.5.2]
- Harris, E.G.: On a plasma sheath separating regions of oppositely directed magnetic field. *Nuovo Cimento* **23**(1), 115–121 (1962) [Sect. 3.1.2]
- Hattori, M., Umetsu, K.: A possible route to spontaneous reduction of the heat conductivity by a temperature gradient-driven instability in electron-ion plasmas. *Astrophys. J.* **533**(1), 84–94 (2000) [Sect. 8.3.4]
- Hawley, J.F., Balbus, S.A.: Instability and turbulence in accretion discs. In: Miyama, S.M., et al. (eds.) *Numerical Astrophysics*, pp. 187–194. Kluwer Academic, Dordrecht (1999) [Sect. 13.2.1]
- Hawley, J.F., Gammie, C.F., Balbus, S.A.: Local three-dimensional magnetohydrodynamic simulations of accretion disks. *Astrophys. J.* **440**(2), 742–763 (1995) [Sect. 13.2.1]
- Heinemann, T., McWilliams, J.C., Schekochihin, A.A.: Large-scale magnetic field generation by randomly forced shearing waves. *Phys. Rev. Lett.*, **107**(25), 255004 (2011) [Sect. 13.1.2]
- Hénoux, J.-C., Somov, B.V.: Generation and structure of the electric currents in a flaring activity complex. *Astron. Astrophys.* **185**(1), 306–314 (1987) [Sect. 20.2]
- Hénoux, J.-C., Somov, B.V.: The photospheric dynamo. 1. Magnetic flux-tube generation. *Astron. Astrophys.* **241**(2), 613–617 (1991) [Sects. 11.1, 20.2]
- Hénoux, J.-C., Somov, B.V.: The photospheric dynamo. 2. Physics of thin magnetic flux tubes. *Astron. Astrophys.* **318**(3), 947–956 (1997) [Sect. 11.1]
- Hirovani, K., Okamoto, I.: Pair plasma production in a force-free magnetosphere around a supermassive black hole. *Astrophys. J.* **497**(2), 563–572 (1998) [Sects. 7.3, 11.5.2]
- Hollweg, J.V.: Viscosity and the Chew-Goldberger-Low equations in the solar corona. *Astrophys. J.* **306**(2), 730–739 (1986) [Sects. 9.6, 10.5]

- Holman, G.D.: DC electric field acceleration of ions in solar flares. *Astrophys. J.* **452**(2), 451–456 (1995) [Sect. 8.4.1(b)]
- Horiuchi, R., Sato, T.: Particle simulation study of driven reconnection in a collisionless plasma. *Phys. Plasmas* **1**(11), 3587–3597 (1994) [Sects. 1.2.4, 11.2]
- Hoshino, M., Stenzel, R.L., Shibata, K. (eds.): *Magnetic Reconnection in Space and Laboratory Plasmas*, p. 693. Terra Scientific Publ. Co., Tokyo (2001) [Sect. 13.1.3]
- Hoyng, P., Brown, J.C., van Beek, H.F.: High time resolution analysis of solar hard X-ray flares observed on board the ESRO TD-1A satellite. *Solar Phys.* **48**(2), 197–254 (1976) [Sect. 4.5.1]
- Hubrig, S., North, P., Mathys, G.: Magnetic Ap stars in the Hertzsprung-Russell diagram. *Astrophys. J.* **539**(1), 352–363 (2000) [Sect. 19.1.3]
- Hudson, P.D.: Reflection of charged particles by plasma shocks. *Mon. Not. Roy. Astron. Soc.* **131**(1), 23–50 (1965) [Sects. 18.3, 18.3.1, 18.3.2(a)]
- Iacus, S.M.: *Simulation and Inference for Stochastic Differential Equations*, p. 284. Springer Science+Business Media, LLC, New York (2008) [Sect. 3.4]
- Imshennik, V.S., Bobrova, N.A.: *Dynamics of Collisional Plasma*. Energoatomizdat, Moscow (in Russian) (1997) [Sect. 15.4.4]
- Innes, D.E., Inhester, B., Axford, W.I., et al.: Bi-directional jets produced by reconnection on the Sun. *Nature* **386**, 811–813 (1997) [Sect. 8.3.3]
- Iordanskii, S.V.: On compression waves in magnetohydrodynamics. *Sov. Phys. Doklady* **3**(4), 736–738 (1958) [Sect. 16.2.4(c)]
- Iroshnikov, P.S.: Turbulence of a conducting fluid in a strong magnetic field. *Sov. Astron. AJ.* **7**(4), 566–571 (1964) [Sect. 7.2.3]
- Jaroschek, C.H., Treumann, R.A., Lesch, H., et al.: Fast reconnection in relativistic pair plasmas: Analysis of particle acceleration in self-consistent full particle simulations. *Phys. Plasm.* **11**(3), 1151–1163 (2004) [Sect. 7.3]
- Jeans, J.: *Astronomy and Cosmogony*. Cambridge University Press, Cambridge (1929) [Sect. 8.1.5]
- Jones, F.C., Ellison D.C.: The plasma physics of shock acceleration. *Space Sci. Rev.* **58**(3), 259–346 (1991) [Sects. 18.1, 18.2.1, 18.3.1]
- Jones, M.E., Lemons, D.S., Mason, R.J., et al.: A grid-based Coulomb collision model for PIC codes. *J. Comput. Phys.* **123**(1), 169–181 (1996) [Sect. 3.4]
- Kadomtsev, B.B.: Convective instability of a plasma. In: Leontovich, M.A. (ed.) *Plasma Physics and the Problem of Controlled Thermonuclear Reactions*, vol. 4, pp. 450–453. Pergamon Press, Oxford (1960) [Sect. 19.3.4]
- Kadomtsev, B.B.: Hydrodynamic stability of a plasma. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 2, pp. 153–198. Consultants Bureau, New York (1966) [Sects. 15.4.1, 19.3.2, 19.3.4]
- Kadomtsev, B.B.: *Collective Phenomena in Plasma*, p. 238. Nauka, Moscow (in Russian) (1976) [Sect. 7.1]
- Kandrup, H.E.: Collisionless relaxation in galactic dynamics and the evolution of long-range order. *Ann. New York Acad. Sci.* **848**, 28–47 (1998) [Sect. 3.3.2]
- Kikuchi, H.: *Electrohydrodynamics in Dusty and Dirty Plasmas*, p. 207. Kluwer Academic, Dordrecht (2001) [Sect. 1.2.4]
- Kirkwood, J.G.: The statistical mechanical theory of transport processes. I. General theory. *J. Chem. Phys.* **14**, 180–201 (1946) [Sect. 2.4]
- Kittel, C.: *Introduction to Solid State Physics*, 7th edn. Wiley, New York (1995) [Sects. 1.4, 3.5]
- Kivelson, M.G., Russell, C.T. (eds.): *Introduction to Space Physics*, p. 568. Cambridge University Press, Cambridge (1995) [Sects. 4.1.1, 6.2.4]
- Kleeorin, N., Rogachevskii, I., Sokoloff, D., et al.: Mean-field dynamos in random Arnold-Beltrami-Childress and Roberts flows. *Phys. Rev. E* **79**(4), 046302 (2009) [Sect. 13.1.2]
- Klimontovich, Yu.L.: *Kinetic Theory of Non-ideal Gas and Non-ideal Plasma*, p. 352. Nauka, Moscow (in Russian) (1975) [Sect. 2.4]
- Klimontovich, Yu.L.: *Statistical Physics*. Harwood Academic, New York (1986) [Intr., Sects. 2.4, 3.1.3, 3.1.4]

- Klimontovich, Yu.L.: Two alternative approaches in the kinetic theory of a fully ionized plasma. *J. Plasma Phys.* **59**(4), 647–656 (1998) [Sect. 3.1.3]
- Klimontovich, Yu.L., Silin, V.P.: On magnetic hydrodynamics for a non-isothermal plasma without collisions. *Sov. Phys. JETP* **40**, 1213–1223 (1961) [Sects. 11.5.1, 16.4]
- Kogan, M.N.: Dynamics of a Dilute Gas. Nauka, Moscow (in Russian) (1967) [Sect. 3.5]
- Koide, S., Shibata, K., Kudoh, T.: Relativistic jet formation from black hole magnetized accretion discs. *Astrophys. J.* **522**, 727–752 (1999) [Sects. 12.2, 13.3.1]
- Kolmogorov, A.N.: The local structure of turbulence in incompressible viscous fluid for very large Reynolds numbers. *C.R. Acad. Sci. USSR*, **30**, 201–206 (1941) [Sect. 7.2]
- Korchak, A.A.: On the origin of solar flare X-rays. *Solar Phys.* **18**(2), 284–304 (1971) [Sect. 8.1.5]
- Korchak, A.A.: Coulomb losses and the nuclear composition of the solar flare accelerated particles. *Solar Phys.* **66**(1), 149–158 (1980) [Sect. 8.4.1(b)]
- Kosugi, T., Matsuzaki, K., Sakao, T., et al.: The Hinode (Solar-B) mission: An overview. *Solar Phys.* **243**(1), 3–17 (2007) [Sect. 8.3.2]
- Kotchine, N.E.: *Rendiconti del Circolo Matematico di Palermo* **50**, 305–314 (1926) [Sect. 17.1.1]
- Kovalev, V.A., Somov, B.V.: On the acceleration of solar-flare charged particles in a collapsing magnetic trap with an electric potential. *Astron. Lett.* **28**(7), 488–493 (2002) [Sect. 8.1.4]
- Kraichnan, R.H.: Inertial-range spectrum of hydromagnetic turbulence. *Phys. Fluids* **8**(7), 1385–1389 (1965) [Sect. 7.2]
- Krall, N.A., Trivelpiece, A.W.: *Principles of Plasma Physics*. McGraw-Hill Book Co., New York (1973) [Sect. 9.6.2]
- Krucker, S., Hudson, H.S., Jeffrey, N.L.S., et al.: High-resolution imaging of solar flare ribbons and its implication on the thick-target beam model. *Astrophys. J.* **739**(2), 96 (7pp) (2011) [Sect. 8.3.2]
- Krymskii, G.F.: A regular mechanism for the acceleration of charged particles on the front of a shock wave. *Sov. Phys. Dokl.* **22**(6), 327–328 (1977) [Sect. 18.2.1]
- Kubbinga, H.: A tribute to Boltzmann. *Europhysicsnews* **37**(6), 28–29 (2006) [Sect. 9.6.1]
- Kudriavtsev, V.S.: Energetic diffusion of fast ions in equilibrium plasma. *Sov. Phys. JETP* **7**(6), 1075–1079 (1958) [Sect. 4.1.2]
- Kulikovskii, A.G., Liubimov, G.A.: On the structure of an inclined MHD shock wave. *Appl. Math. Mech.* **25**(1), 171–179 (1961) [Sect. 17.4.2]
- Kumar, N., Kumar, P., Singh, S.: Coronal heating by MHD waves. *Astron. Astrophys.* **453**(2), 1067–1078 (2006) [Sect. 15.2.1]
- Kunkel, W.B.: Generalized Ohm's law for plasma including neutral particles. *Phys. Fluids* **27**(9), 2369–2371 (1984) [Sect. 11.1]
- Lahav, O., Terlevich, E., Terlevich, R.J. (eds.): *Gravitational Dynamics*, p. 270. Cambridge University Press, Cambridge (1996) [Sect. 1.3]
- Lancellotti, C., Kiessling, M.: Self-similar gravitational collapse in stellar dynamics. *Astrophys. J.* **549**, L93–L96 (2001) [Sect. 3.3.2]
- Landau, L.D.: Kinetic equation in the case of Coulomb interaction. *Zhur. Exper. Teor. Fiz.* **7**(1), 203–212 (in Russian) (1937) [Sect. 3.1.3]
- Landau, L.D.: On the vibrations of the electron plasma. *J. Phys. USSR* **10**(1), 25–30 (1946) [Sects. 3.1.3, 7.1]
- Landau, L.D., Lifshitz, E.M.: *Fluid Mechanics*, p. 536. Oxford, London (1959a) [Sects. 12.2.2, 12.2.3, 15.6, 16.1.2, 16.2.2, 20.2]
- Landau, L.D., Lifshitz, E.M.: *Statistical Physics*, p. 478. Pergamon Press, London (1959b) [Sects. 1.1.5, 1.4, 3.5, 16.5]
- Landau, L.D., Lifshitz, E.M.: *Classical Theory of Field*, 4th edn., p. 374. Oxford, New York (1975) [Sects. 1.2.1, 2.2.1, 5.1.1, 5.1.3, 5.4, 6.2.1, 7.4, 13.4, 18.4, 19.1.1]
- Landau, L.D., Lifshitz, E.M.: *Mechanics*, 3rd edn., p. 165. Oxford, London (1976) [Sects. 1.1.5, 1.4, 6.1, 8.1.1, 19.1.1]
- Landau, L.D., Lifshitz, E.M., Pitaevskii, L.P.: *Electrodynamics of Continuous Media*, p. 460. Pergamon Press, Oxford (1984) [Sects. 11.4.2, 16.2.4(c), 17.3.2]
- Langmuir, I.: *Proc. Nat. Acad. Sci. U.S.A.* **14**, 627 (1928) [Sect. 3.2.2]

- Larrabee, D.A., Lovelace, R.V.E., Romanova, M.M.: Lepton acceleration by relativistic collisionless magnetic reconnection. *Astrophys. J.* **586**(1), 72–78 (2003) [Sect. 7.3]
- Lavrent'ev, M.A., Shabat, B.V.: *Methods of the Theory of Complex Variable Functions*, p. 736. Nauka, Moscow (in Russian) (1973) [Sects. 3.1.3, 9.7.1, 14.2.2(a)]
- Lax, P.: Hyperbolic systems of conservation laws. *Comm. Pure Appl. Math.* **10**(4), 537–566 (1957) [Sect. 17.1.1]
- Lax, P.: *Hyperbolic Partial Differential Equations*, AMS, Courant Inst. of Math. Sci. (2006) [Sect. 17.1.1]
- Leenov, D., Kolin, A.: Theory of electromagnetophoresis. 1. MHD forces experienced by spherical and cylindrical particles. *J. Chem. Phys.* **22**(4), 683–688 (1954) [Sect. 20.4]
- Leith, C.E.: Diffusion approximation to inertial energy transfer in isotropic turbulence. *Phys. Fluids* **10**(7), 1409–1416 (1967) [Sect. 7.2]
- Leontovich, M.A. (ed.): *Plasma Physics and the Problem of Controlled Thermonuclear Reactions*, vols. 1–4. Pergamon Press, London (1960) [Intr.]
- Lesch, H., Pohl, M.: A possible explanation for intraday variability in active galactic nuclei. *Astron. Astrophys.* **254**(1), 29–38 (1992) [Sect. 13.2.4]
- Letessier, J., Rafelski, J.: *Hadrons and Quark-gluon Plasma*, p. 397. Cambridge University Press, Cambridge (2004) [Sect. 12.2.5]
- Lieberman, M.A.: On actuating shock waves in a completely ionized plasma. *Sov. Phys. JETP* **48**(5), 832–840 (1978) [Sects. 16.2.6, 17.4.2]
- Liboff, R.: *Kinetic Theory: Classical, Quantum, and Relativistic Descriptions*, p. 571. Springer, Heidelberg (2003) [Intr.]
- Lichnerowicz, A.: *Relativistic Hydrodynamics and Magnetohydrodynamics*, p. 196. Benjamin, New York (1967) [Sect. 12.2]
- Lifshitz, E.M., Pitaevskii, L.P.: *Physical Kinetics*, p. 452. Pergamon Press, Oxford (1981) [Sects. 3.5, 7.3, 8.3.1, 9.6]
- Lin, R.P., Hudson, H.S.: 10–100 keV electron acceleration and emission from solar flares. *Solar Phys.* **17**(2), 412–435 (1971) [Sects. 4.3.4, 8.3.2]
- Lin, R.P., Dennis, B.R., Hurford, G.J., et al.: The Reuven Ramaty High-Energy Solar Spectroscopic Imager (RHESSI). *Solar Phys.* **210**(1), 3–32 (2002) [Sect. 4.5.7]
- Lin, R.P., Krucker, S., Hurford, G.J., et al.: *RHESSI* observations of particle acceleration and energy release in an intense solar gamma-ray line flare. *Astrophys. J.* **595**(2), L69–L76 (2003) [Sect. 4.5.7]
- Litvinenko, Y.E., Somov, B.V.: Solar flares and virial theorem. *Sov. Astron. AJ* **35**(2), 183–188 (1991a) [Sects. 19.1.3, 19.2.2, 19.4.3]
- Litvinenko, Y.E., Somov, B.V.: Nonthermal electrons in the thick-target reverse-current model for hard X-ray bremsstrahlung. *Solar Phys.* **131**(2), 319–336 (1991b) [Sects. 4.5.2, 4.5.5]
- Litvinenko, Y.E., Somov, B.V.: Electromagnetic expulsion force in cosmic plasma. *Astron. Astrophys.* **287**(1), L37–L40 (1994) [Sect. 20.4]
- Litvinenko, Y.E., Somov, B.V.: Aspects of the global MHD equilibria and filament eruptions in the solar corona. *Space Sci. Rev.* **95**(1), 67–77 (2001) [Sects. 19.1.3, 19.4.3]
- Liubarskii, G.Ya., Polovin, R.V.: Simple magnetoacoustic waves. *Sov. Phys. JETP* **8**(2), 351 (1958) [Sect. 16.2.4(c)]
- Lovelace, R.V.E.: Dynamo model of double radio sources. *Nature* **262**, 649–652 (1976) [Sect. 20.1.3]
- Luna, M., Karpen, J.T., DeVore, C.R.: Formation and evolution of a multi-threaded solar prominence. *Astrophys. J.* **746**(1), article id. 30 (2012) [Sect. 12.1.3]
- Lundquist, S.: Magneto-hydrostatic fields. *Ark. Fys.* **2**(35), 361–365 (1951) [Sect. 19.2.1.2]
- Macdonald, D.A., Thorne, K.S., Price, R.H., et al.: Astrophysical applications of black-hole electrodynamics. In: Thorne, K.S., Price, R.H., Macdonald, D.A. (eds.) *Black Holes: The Membrane Paradigm*, pp. 121–137. Yale University Press, New Haven (1986) [Sect. 13.3.1]
- MacDonald, W.M., Rothenbluth, M.N., Chuck, W.: Relaxation of a system of particles with Coulomb interactions. *Phys. Rev.* **107**(2), 350–353 (1957) [Sect. 4.1.2]

- Mach, E.: Die Geschichte und die Wurzel des Satzes von der Erhaltung der Arbeit. Calve, Prag (1872) [Sect. 13.3.2]
- Mach, E.: Die Mechanik in ihrer Entwicklung. Historisch-kritisch Dargestellt. Brockhaus, Leipzig (1883) [Sect. 13.3.2]
- MacNeice, P., McWhirter, R.W.P., Spicer, D.S., et al.: A numerical model of a solar flare based on electron beam heating of the chromosphere. *Solar Phys.* **90**(2), 357–353 (1984) [Sect. 8.3.2]
- Manmoto, T.: Advection-dominated accretion flow around a Kerr black hole. *Astrophys. J.* **534**(2), 734–746 (2000) [Sects. 8.3.5, 13.2.3]
- Markovskii, S.A.: Nonevolutionarity of trans-Alfvénic shocks in a magnetized plasma. *J. Geophys. Res.* **104**(A3), 4427–4436 (1999) [Sects. 17.3.2, 17.4.2]
- Markovskii, S.A., Skorokhodov, S.L.: Disintegration of trans-Alfvénic shocks due to variable viscosity and resistivity. *J. Geophys. Res.* **105**(A6), 12702–12711 (2000) [Sect. 17.4.2]
- Markovskii, S.A., Somov, B.V.: A model of magnetic reconnection in a current sheet with shock waves. *Fizika Solnechnoi Plazmy (Physics of Solar Plasma)*, pp. 456–472. Nauka, Moscow (in Russian) (1989) [Sect. 14.2.2(a)]
- Markovskii, S.A., Somov, B.V.: MHD discontinuities in space plasmas: Interrelation between stability and structure. *Space Sci. Rev.* **78**(3–4), 443–506 (1996) [Sect. 17.5]
- Marty, P., Alemany, A.: Écoulement dû à des champs magnétique et électrique croisés autour d'un cylindre de conductivité quelconque. *Journal de Mécanique Théorique et Appliquée* **2**(2), 227–243 (1983) [Sects. 19.4.2, 20.3]
- Maxwell, J.C.: Illustrations of the dynamical theory of gases. *Phil. Mag. Ser.* **4**(19), 19–24 (1860) [Sect. 9.6.1]
- McClymont, A.N., Canfield, R.C.: Flare loop radiative hydrodynamics. I – Basic methods. *Astrophys. J.* **265**, 483–506 (1983) [Sect. 8.3.2]
- McDonald, L., Harra-Murnion, L.K., Culhane, J.L.: Non-thermal electron energy deposition in the chromosphere and the accompanying soft X-ray flare emission. *Solar Phys.* **185**(2), 323–350 (1999) [Sect. 8.3.2]
- Michel, F.C.: *Theory of Neutron Star Magnetospheres*, p. 456. Chicago University Press, Chicago (1991) [Sects. 7.3, 11.5.2, 12.2.2]
- Mikhailovskii, A.B.: Nonlinear excitation of electromagnetic waves in a relativistic electron-positron plasma. *Sov. J. Plasma Phys.* **6**(3), 336–340 (1979) [Sect. 7.3]
- Mikhailovskii, A.B., Onishchenko, O.G., Tatarinov, E.G.: Alfvén solitons in a relativistic electron-positron plasma. *Plasma Phys. Contr. Fusion* **27**(5), 539–556 (1985) [Sect. 7.3]
- Mirabel, I.F., Rodriguez, L.F.: Microquasars in our Galaxy. *Nature* **392**, 673–676 (1998) [Sect. 20.1.3]
- Moffatt, H.K.: *Magnetic Field Generation in Electrically Conducting Fluids*, p. 343. Cambridge University Press, London (1978) [Sect. 13.1.2]
- Moreau, R.: *Magnetohydrodynamics*, p. 328. Kluwer Academic, Dordrecht (1990) [Sect. 20.1.5]
- Morozov, A.I., Solov'ev, L.S.: The structure of magnetic fields. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 2, pp. 1–101. Consultants Bureau, New York (1966a) [Sect. 19.3.4]
- Morozov, A.I., Solov'ev, L.S.: Motion of particles in electromagnetic fields. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 2, pp. 201–297. Consultants Bureau, New York (1966b) [Sect. 5.3.4]
- Moses, G.A., Duderstadt, J.J.: Improved treatment of electron thermal conduction in plasma hydrodynamics calculations. *Phys. Fluids* **20**(5), 762–770 (1977) [Sect. 9.7.1]
- Nakano, T.: Star formation in magnetic clouds. *Astrophys. J.* **494**(2), 587–604 (1998) [Sect. 19.1.3]
- Narayan, R., Garcia, M.R., McClintock, J.E.: Advection-dominated accretion and black hole horizons. *Astrophys. J.* **478**(2), L79–L82 (1997) [Sect. 8.3.5]
- Negoro, H., Kitamoto, S., Takeuchi, M., et al.: Statistics of X-ray fluctuations from Cygnus X-1: Reservoirs in the disk? *Astrophys. J.* **452**(1), L49–L52 (1995) [Sect. 13.2.4]
- Nishida, A.: Can random reconnection on the magnetopause produce the low latitude boundary layer? *Geophys. Res. Lett.* **16**, 227–230 (1989) [Sect. 12.4.2]

- Nishikawa, K.I., Frank, J., Christodoulou, D.M., et al.: 3D relativistic MHD simulations of extragalactic jets. In: Miyama, S.M., et al. (eds.) *Numerical Astrophysics*, pp. 217–218. Kluwer Academic, Dordrecht (1999) [Sect. 13.3.1]
- Northrop, T.G.: *The Adiabatic Motion of Charged Particles*. Wiley, New York (1963) [Sect. 6.4]
- Novikov, I.D., Frolov, V.P.: *Physics of Black Holes*, p. 341. Kluwer Academic, Dordrecht (1989) [Sects. 11.5.2, 12.2.2, 13.3.1]
- Novikov, I.D., Thorne, K.S.: In: Dewitt, C.D., Dewitt, B. (eds.) *Black Holes*, pp. 345–354. Gordon and Breach, New York (1973) [Sects. 8.3.5, 13.2.1, 13.2.3]
- Obertz, P.: Two-dimensional problem of the shape of the magnetosphere. *Geomagn. Aeron.* **13**(5), 758–766 (1973) [Sect. 14.2.2(a)]
- Ogawara, Y., Takano, T., Kato, T., et al.: The Solar-A mission: An overview. *Solar Phys.* **136**(1), 1–16 (1991) [Sect. 8.3.2]
- Oreshina, A.V., Somov, B.V.: Analytical description of charged particle motion in a reconnecting current layer *Astron. Lett.* **35**(3), 195–206 (2009) [Sect. 5.2.3]
- Oreshina, A.V., Somov, B.V.: On the heat-transfer mechanisms in solar flares. 1. Classical and anomalous heat conduction. *Moscow Univ. Phys. Bull.* **66**(3), 286–291 (2011a) [Sects. 9.6.2, 9.7.3]
- Oreshina, A.V., Somov, B.V.: On the heat-transfer mechanisms in solar flares. 1. Account of heat-flux relaxation. *Moscow Univ. Phys. Bull.* **66**(3), 292–297 (2011b) [Sects. 9.6.2, 9.7.3]
- Oreshina, I.V., Somov, B.V.: Conformal mapping for solving problems of space electrodynamics. *Bull. Russ. Acad. Sci. Phys.* **63**(8), 1209–1212 (1999) [Sect. 14.5]
- Ostriker, E.C.: Dynamical friction in a gaseous medium. *Astrophys. J.* **513**(1), 252–258 (1999) [Sect. 8.5]
- Padmanabhan, T.: *An Invitation to Astrophysics*. World Scientific Publ. Co., New Jersey (2006) [Sect. 14.4.2]
- Palmer, P.L.: *Stability of Collisionless Stellar Systems*, p. 349. Kluwer Academic, Dordrecht (1994) [Sect. 9.8]
- Parker, E.N.: *Cosmic Magnetic Fields. Their Origin and Their Activity*, p. 841. Clarendon Press, Oxford (1979) [Sects. 13.1.2, 19.3.4, 19.4.2, 20.1.5]
- Parks, G.K.: *Physics of Space Plasmas, An Introduction*, 2nd edn., p. 597. Westview Press, Oxford (2004) [Intr., Sects. 14.5, 18.1, 18.2.3]
- Peacock, J.A.: *Cosmological Physics*, p. 682. Cambridge University Press, Cambridge (1999) [Sects. 7.3, 9.8]
- Persson, H.: Electric field along a magnetic line of force in a low-density plasma. *Phys. Fluids* **6**(12), 1756–1759 (1963) [Sect. 8.1.4]
- Peterson, L.E., Winckler, J.B.: Gamma-ray burst from a solar flare. *J. Geophys. Res.* **64**(7), 697–707 (1959) [Sect. 4.3.4]
- Pfaffelmoser, K.: Global classic solutions of the Vlasov-Poisson system in three dimensions for general initial data. *J. Diff. Equat.* **95**, 281–303 (1992) [Sect. 16.5]
- Phan, T.D., Gosling, J.T., Davis, M.S., et al.: A magnetic reconnection X-line extending more than 390 Earth radii in the solar wind. *Nature* **439**(04393), 175–178 (2006) [Sect. 12.4.2]
- Planck, M.: Über einen Satz der Statistischen Dynamik und seine Erweiterung in der Quantentheorie. *Sitzber Preuss. Akad. Wiss., Phys-Math. Klasse* 324 (1917) [Sect. 3.1.4]
- Polovin, R.V.: Shock waves in MHD. *Soviet Phys. Usp.* **3**(5), 677–688 (1961) [Sects. 16.2.4(c), 17.2.2]
- Polovin, R.V., Demutskii, V.P.: *Fundamentals of Magnetohydrodynamics*. Consultants Bureau, New York (1990) [Sect. 17.4.1]
- Polovin, R.V., Liubarskii, G.Ya.: Impossibility of rarefaction shock waves in MHD. *Sov. Phys. JETP* **8**(2), 351–352 (1958) [Sect. 16.2.4(c)]
- Priest, E.R.: *Solar Magnetohydrodynamics*, p. 472. D. Reidel Publ. Co., Dordrecht (1982) [Sects. 16.2.4(c), 19.3.4]
- Punsly, B.: *Black Hole Gravito-hydro-magnetics*, p. 400. Springer, New York (2001) [Sect. 12.2.5]
- Quarati, P., Scarfone, A.M.: Modified Debye-Hückel electron shielding and penetration factor. *Astrophys. J.* **666**(2), 1303–1310 (2007) [Sect. 8.2.2]

- Ramos, J.I., Winowich, N.S.: Magnetohydrodynamic channel flow study. *Phys. Fluids* **29**(4), 992–997 (1986) [Sect. 20.2]
- Reid, I.N., Liebert, J., Schmidt, G.D.: Discovery of a magnetic DZ white dwarf with Zeeman-split lines of heavy elements. *Astrophys. J.* **550**(1), L61–L63 (2001) [Sect. 13.2.2]
- Rodriguez-Pacheco, J., Sequeiros, J., del Peral, L., et al.: Diffusive-shock-accelerated interplanetary ions at several energies during the solar cycle 21 maximum. *Solar Phys.* **181**(1), 185–200 (1998) [Sect. 18.2.1]
- Rogachevskii, I., Kleeorin, N.: Shear-current effect in a turbulent convection with a large-scale shear. *Phys. Rev. E* **75**(4), 046305 (2007) [Sect. 13.1.2]
- Roikhvarger, Z.B., Syrovatskii, S.I.: Evolutionarity of MHD discontinuities with allowance for dissipative waves. *Sov. Phys. JETP* **39**(4), 654–656 (1974) [Sects. 17.1.4, 17.3.1, 17.3.2]
- Rose, W.K.: *Advanced Stellar Astrophysics*, p. 494. Cambridge University Press, Cambridge (1998) [Sects. 1.3, 5.4, 7.3, 12.2.2, 13.2.1, 14.4.2]
- Rosenbluth, M., Longmire, C.: Stability of plasmas confined by magnetic fields. *Ann. Phys.* **1**(1), 120–140 (1957) [Sects. 19.3.2, 19.3.3]
- Ruderman, M.: Matter in superstrong magnetic fields: The surface of a neutron star. *Phys. Rev. Lett.* **27**(19), 1306–1308 (1971) [Sect. 5.4]
- Ruderman, M.A., Sutherland, P.G.: Theory of pulsars: Polar gaps, sparks, and coherent radiation. *Astrophys. J.* **196**(1), 51–72 (1975) [Sect. 7.3]
- Rüdiger, G., von Rekowski, B.: Differential rotation and meridional flow for fast-rotating solar-type stars. *Astrophys. J.* **494**(2), 691–699 (1998) [Sects. 13.1.2, 20.1.5]
- Ruffolo, D.: Transport and acceleration of energetic particles near an oblique shock. *Astrophys. J.* **515**(2), 787–800 (1999) [Sect. 18.2.1]
- Salat, A.: Non-linear plasma transport equations for high flow velocity. *Plasma Phys. J.* **17**, 589–607 (1975) [Sect. 9.7.2]
- Sarazin, C.L., Kempner, J.C.: Nonthermal bremsstrahlung and hard X-ray emission from clusters of galaxies. *Astrophys. J.* **533**(1), 73–83 (2000) [Sect. 8.3.4]
- Sarris, E.T., Van Allen, J.A.: Effects of interplanetary shocks on energetic particles. *J. Geophys. Res.* **79**(28), 4157–4173 (1974) [Sect. 18.3.2(a)]
- Schabansky, V.P.: Some processes in the magnetosphere. *Space Sci. Rev.* **12**(3), 299–418 (1971) [Sect. 11.1]
- Schiff, L.I.: Possible new experimental test of general relativity theory. *Phys. Rev. Lett.* **4**(5), 215–217 (1960) [Sect. 13.4]
- Schlickeiser, R.: *Cosmic Ray Astrophysics*, p. 519. Springer, New York (2002) [Sect. 5.1.3]
- Schlüter, A.: Dynamic des Plasmas. *Zeitschrift für Naturforschung* **6A**(2), 73–78 (1951) [Sect. 11.1]
- Schmidt, G.: *Physics of High Temperature Plasmas*, p. 408. Academic, New York (1979) [Sect. 3.1.2]
- Schou, J., Antia, H.M., Basu, S., et al.: Helioseismic studies of differential rotation in the solar envelope by the solar oscillations investigation using the Michelson Doppler Imager. *Astrophys. J.* **505**(1), 390–417 (1998) [Sect. 20.1.5]
- Schram, P.P.J.: *Kinetic Theory of Gases and Plasmas*, p. 426. Kluwer Academic, Dordrecht (1991) [Intr., Sect. 6.2.2]
- Schrijver, C.J., Zwaan, C.: *Solar and Stellar Magnetic Activity*, p. 400. Cambridge University Press, Cambridge (1999) [Sect. 20.1.5]
- Sedov, L.I.: *Mechanics of Continuous Medium*, vol. 1, p. 536, vol. 2, p. 584. Nauka, Moscow (in Russian) (1973) [Sect. 13.1.1]
- Sermulyan'sh, B.A., Somov, B.V.: The problem of reverse current under heating of the solar atmosphere by accelerated electrons. In: *Proc. 12th Leningrad Seminar on Space Physics: Complex Study of the Sun*, pp. 90–95. LIYaF, Leningrad (in Russian) (1982) [Sect. 4.5.6]
- Sermulyan'sh, B.A., Somov, B.V.: On the influence of reverse current on the chromospheric heating by accelerated electrons. *Investig. Sun Red Stars* **18**, 86–92 (in Russian) (1983) [Sect. 4.5.6]
- Shafranov, V.D.: On MHD equilibrium configurations. *Sov. Phys. JETP* **6**, 545–551 (1958) [Sect. 19.5]

- Shafranov, V.D.: Plasma equilibrium in a magnetic field. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 2, pp. 103–151. Consultants Bureau, New York (1966) [Sects. [19.2.2](#), [19.3.2](#)]
- Shakura, N.I., Sunyaev, R.A.: Black holes in binary systems, Observational appearance. *Astron. Astrophys.* **24**(2), 337–355 (1973) [Sects. [8.3.5](#), [13.2.1](#), [13.2.3](#)]
- Sheeley, N.R., Jr., Warren, H.P., Wang, Y.-M.: A streamer ejection with reconnection close to the Sun. *Astrophys. J.* **671**(1), 926–935 (2007) [Sect. [11.5.1](#)]
- Shercliff, A.J.: *A Textbook of Magnetohydrodynamics*, p. 265. Pergamon Press, Oxford (1965) [Sects. [13.1.1](#), [16.2.4\(c\)](#), [17.4.2](#), [20.2.2](#)]
- Shkarofsky, I.P., Johnston, T.W., Bachynski, M.P.: *The Particle Kinetics of Plasma*, p. 518. Addison-Wesley, Reading (1966) [Sects. [1.1.4](#), [9.4.1](#), [9.6.2](#), [11.5.1](#), [12.2.3](#)]
- Shmeleva, O.P., Syrovatskii, S.I.: Distribution of temperature and emission measure in a steadily heated solar atmosphere. *Solar Phys.* **33**(2), 341–362 (1973) [Sect. [8.5](#)]
- Shoub, E.C.: Invalidity of local thermodynamic equilibrium for electrons in solar transition region. *Astrophys. J.* **266**(1), 339–369 (1983) [Sect. [8.4.3](#)]
- Shoub, E.C.: Failure of the Fokker-Planck approximation to the Boltzmann integral for (1/r) potentials. *Phys. Fluids* **30**(5), 1340–1352 (1987) [Sects. [3.1.4](#), [3.5](#)]
- Shu, F.H.: *The Physics of Astrophysics*, vol. 2. Gas Dynamics, p. 476. California Univ. Science Books, Mill Valley (1992) [Sects. [6.2.2](#), [19.3.4](#)]
- Silin, V.P.: *Introduction to the Kinetic Theory of Gases*, p. 332. Nauka, Moscow (in Russian) (1971) [Sects. [3.1.2](#), [3.5](#), [6.2.2](#)]
- Simon, A.L.: *An Introduction to Thermonuclear Research*, p. 182. Pergamon Press, London (1959) [Intr.]
- Sirotina, E.P., Syrovatskii, S.I.: Structure of low intensity shock waves in MHD. *Sov. Phys. JETP* **12**(3), 521–526 (1960) [Sects. [16.4](#), [17.3.1](#)]
- Sivukhin, D.V.: Motion of charged particles in electromagnetic fields in the drift approximation. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 1, pp. 1–104. Consultants Bureau, New York (1965) [Sects. [5.2.3](#), [5.3.4](#)]
- Sivukhin, D.V.: Coulomb collisions in a fully ionized plasma. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 4, pp. 93–341. Consultants Bureau, New York (1966) [Sects. [8.3.1](#), [8.4.1\(b\)](#), [8.4.3](#)]
- Sivukhin, D.V.: *A Course of General Physics. Vol. II, Thermodynamics and Molecular Physics*, 3rd edn. Nauka, Moscow (in Russian) (1990) [Sect. [9.6.1](#)]
- Sivukhin, D.V.: *A Course of General Physics. Vol. III, Electricity*, 3rd edn. Nauka, Moscow (in Russian) (1996) [Sects. [11.4.2](#), [19.2.2](#)]
- Smirnov, B.M.: *Physics of Weakly Ionized Gases: Problems and Solutions*, p. 432. Mir Publ., Moscow (1981) [Sect. [3.5](#)]
- Smirnov, V.I.: *A Course of Higher Mathematics*, vol. 2. Pergamon Press, Oxford (1965) [Sects. [1.1.1](#), [12.3.1](#), [19.1.2](#), [19.6](#)]
- Smith, E.J., Tsurutani, B.T., Rosenberg, R.L.: Observations of the interplanetary sector structure up to heliographic latitudes of 16°: Pioneer 11. *J. Geophys. Res.* **83**, 717–724 (1978) [Sect. [12.4.2](#)]
- Somov, B.V.: Fast reconnection and transient phenomena with particle acceleration in the solar corona. *Bull. Acad. Sci. USSR, Phys. Ser.* **45**(4), 114–116 (1981) [Sects. [8.3.3](#), [9.7.2](#)]
- Somov, B.V.: Accumulation and release of flare energy. In: *Proc. 12th Leningrad Seminar on Space Physics: Complex Study of the Sun*, pp. 6–49. LIYaF, Leningrad (in Russian) (1982) [Sects. [3.1.4](#), [4.1.2](#), [4.4](#)]
- Somov, B.V.: Non-neutral current sheets and solar flare energetics. *Astron. Astrophys.* **163**(1), 210–218 (1986) [Sect. [8.3.3](#)]
- Somov, B.V.: *Physical Processes in Solar Flares*, p. 248. Kluwer Academic, Dordrecht (1992) [Sects. [4.5.6](#), [8.3.2](#), [8.4.3](#), [9.7.3](#), [19.4.3](#)]
- Somov, B.V.: *Cosmic Electrodynamics and Solar Physics*, p. 288. Moscow State Univ. Publ., Moscow (in Russian) (1993) [Sect. [16.3](#)]
- Somov, B.V.: *Fundamentals of Cosmic Electrodynamics*, p. 364. Kluwer Academic, Dordrecht (1994a) [Sects. [14.2](#), [16.3](#)]

- Somov, B.V.: Features of mass supply and flows related with reconnection in the solar corona. *Space Sci. Rev.* **70**(1), 161–166 (1994b) [Sects. [19.4.1](#), [20.4](#)]
- Somov, B.V.: *Plasma Astrophysics, Part II, Reconnection and Flares*, p. 504. Springer Science + Business Media, New York (2012) [Intr.]
- Somov, B.V., Gritsyk, P.A.: Bremsstrahlung radiation of accelerated electrons in solar flares. *Moscow Univ. Phys. Bull.* **67**(1), 110–116 (2012) [Sect. [4.5.5](#)]
- Somov, B.V., Kosugi, T.: Collisionless reconnection and high-energy particle acceleration in solar flares. *Astrophys. J.* **485**(2), 859–868 (1997) [Sect. [6.2.4](#)]
- Somov, B.V., Syrovatskii, S.I.: Plasma motion in an increasing strong dipolar field. *Sov. Phys. JETP* **34**(2), 332–335 (1972a) [Sects. [14.4.1](#), [14.4.2](#)]
- Somov, B.V., Syrovatskii, S.I.: Appearance of a current sheet in a plasma moving in the field of a two-dimensional magnetic dipole. *Sov. Phys. JETP* **34**(5), 992–997 (1972b) [Sect. [14.2.2\(a\)](#)]
- Somov, B.V., Syrovatskii, S.I.: Physical processes in the solar atmosphere associated with flares. *Sov. Phys. Usp.* **19**(10), 813–835 (1976a) [Sects. [8.3.3](#), [8.3.4](#), [12.1.3](#)]
- Somov, B.V., Syrovatskii, S.I.: Hydrodynamic plasma flows in a strong magnetic field. In: Basov, N.G. (ed.) *Neutral Current Sheets in Plasma*, Proc. P.N. Lebedev Phys. Inst., vol. 74, pp. 13–71. Consultants Bureau, New York (1976b) [Sects. [12.1.3](#), [13.1.1](#), [14.1](#), [14.2.2\(b\)](#), [14.4.2](#)]
- Somov, B.V., Tindo, I.P.: Polarization of hard X-rays from solar flares. *Cosmic Res.* **16**(5), 555–564 (1978) [Sect. [4.5.5](#)]
- Somov, B.V., Titov, V.S.: Magnetic reconnection as a mechanism for heating the coronal loops. *Sov. Astron. Lett.* **9**(1), 26–28 (1983) [Sect. [8.3.3](#)]
- Somov, B.V., Spektor, A.R., Syrovatskii, S.I.: Gas dynamics of a flare plasma. *Bull. Acad. Sci. USSR Phys. Ser.* **41**(2), 32–43 (1977) [Sect. [8.3.2](#)]
- Somov, B.V., Spektor, A.R., Syrovatskii, S.I.: Hydrodynamics of an optically transparent plasma with a distributed heating source. In: Basov, N.G. (ed.) *Flare Processes in Plasmas*, Proc. P.N. Lebedev Phys. Inst., vol. 110, pp. 73–94. Nauka, Moscow (in Russian) (1979) [Sect. [8.3.2](#)]
- Somov, B.V., Syrovatskii, S.I., Spektor, A.R.: Hydrodynamic response of the solar chromosphere to elementary flare burst. 1. Heating by accelerated electrons. *Solar Phys.* **73**(1), 145–155 (1981) [Sect. [8.3.2](#)]
- Somov, B.V., Sermulina, B.J., Spektor, A.R.: Hydrodynamic response of the solar chromosphere to elementary flare burst. 1. Thermal model. *Solar Phys.* **81**(1), 281–292 (1982) [Sect. [8.3.3](#)]
- Somov, B.V., Oreshina, A.V., Oreshina, I.V., et al.: Flares in accretion disk coronae. *Adv. Space Res.* **32**(6), 1087–1096 (2003) [Sects. [14.2.2\(a\)](#), [14.5](#)]
- Somov, B.V., Dzhalilov, N.S., Staudte, J.: Peculiarities of entropy and magnetosonic waves in optically thin cosmic plasma. *Astron. Lett.* **33**(5), 309–318 (2007) [Sects. [12.1.3](#), [15.2.1](#), [15.4.4](#)]
- Spicer, D.S., Emslie, A.G.: A new quasi-thermal trap model for solar hard X-ray bursts: An electrostatic trap model. *Astrophys. J.* **330**(2), 997–1007 (1988) [Sect. [8.1.4](#)]
- Spitzer, L.: The stability of isolated clusters. *Mon. Not. Roy. Astron. Soc.* **100**(5), 396–413 (1940) [Sect. [8.3.1](#)]
- Spitzer, L.: *Physics of Fully Ionized Gases*, p. 170. Wiley Interscience, New York (1962) [Sects. [8.3.1](#), [8.4.1\(a\)](#), [9.6.2](#), [15.4.1](#), [15.4.4](#)]
- Steinolfson, R.S., Cable, S.: Venus bow shock at unusually large distances from the planet. *Geophys. Res. Lett.* **20**, 755–758 (1993) [Sect. [16.2.5](#)]
- Steinolfson, R.S., Hundhausen, A.J.: MHD intermediate shocks in coronal mass ejections. *J. Geophys. Res.* **95**, 6389–6401 (1990) [Sect. [16.2.5](#)]
- Stewart, R.W., Grant, H.L.: Determination of the rate of dissipation of turbulent energy near the sea surface in the presence of waves. *J. Geophys. Res.* **67**, 3177–3184 (1969) [Sect. [7.2.2](#)]
- Stix, T.H.: *Waves in Plasmas*. American Institute of Physics, New York (1992) [Sect. [10.4](#)]
- Störmer, C.: *The Polar Aurora*. Clarendon Press, Oxford (1955) [Sect. [6.4](#)]
- Strittmatter, P.A.: Gravitational collapse in the presence of a magnetic field. *Monthly Not. Roy. Astron. Soc.* **132**(3), 359–378 (1966) [Sects. [19.1.2](#), [19.1.3](#)]
- Strong, K.T., Saba, J.L.R., Haisch, B.M., et al. (eds.): *The Many Faces of the Sun*, p. 610. Springer, New York (1999) [Sect. [4.3.4](#)]

- Subramanian, P., Becker, P.A., Kazanas, D.: Formation of relativistic outflows in shearing black hole accretion coronae. *Astrophys. J.* **523**(1), 203–222 (1999) [Sect. 13.3.4]
- Suh, I.S., Mathews, G.J.: Cold ideal equation of state for strongly magnetized neutron star matter: Effects on muon production and pion condensation. *Astrophys. J.* **546**(3), 1126–1136 (2001) [Sect. 19.1.3]
- Sutton, G.W., Sherman, A.: *Engineering Magnetohydrodynamics*, p. 548. McGraw-Hill Book Co., New York (1965) [Sects. 13.1.1, 20.2]
- Syrovatskii, S.I.: On the stability of tangential discontinuities in MHD medium. *Zhur. Exper. Teor. Fiz.* **24**(6), 622–630 (in Russian) (1953) [Sects. 16.2.1, 16.2.2]
- Syrovatskii, S.I.: Instability of tangential discontinuities in a compressive medium. *Zhur. Exper. Teor. Fiz.* **27**(1), 121–123 (in Russian) (1954) [Sect. 16.2.2]
- Syrovatskii, S.I.: Some properties of discontinuity surfaces in MHD. *Proc. P.N. Lebedev Phys. Inst.* **8**, 13–64 (in Russian) (1956) [Sects. 16.2.1, 16.3, 20.1.1]
- Syrovatskii, S.I.: *Magnetohydrodynamics*. *Uspehi Fiz. Nauk* **62**(3), 247–303 (in Russian) (1957) [Sects. 12.2.2, 15.4.2, 16.2.4(c), 19.1.3, 20.1.1]
- Syrovatskii, S.I.: The stability of shock waves in MHD. *Sov. Phys. JETP* **8**(6), 1024–1028 (1959) [Sects. 17.1.2, 17.1.4]
- Syrovatskii, S.I.: Formation of current sheets in a plasma with a frozen-in strong field. *Sov. Phys. JETP* **33**(5), 933–940 (1971) [Sect. 14.2.2(a)]
- Syrovatskii, S.I., Chesalin, L.S.: Electromagnetic generation of conductive fluid flows near bodies and expulsive force. *Questions of Magnetohydrodynamics*, pp. 17–22. Zinatne, Riga (in Russian) (1963) [Sects. 19.4.2, 20.3]
- Syrovatskii, S.I., Shmeleva, O.P.: Heating of plasma by high-energy electrons, and the non-thermal X-ray emission in solar flares. *Sov. Astron. AJ* **16**(2), 273–283 (1972) [Sects. 4.3.3, 4.3.4, 8.3.2]
- Syrovatskii, S.I., Somov, B.V.: Physical driving forces and models of coronal responses. In: Dryer, M., Tandberg-Hanssen, E. (eds.) *Solar and Interplanetary Dynamics*, IAU Symp. vol. 91, pp. 425–441. Reidel, Dordrecht (1980) [Sect. 14.2.2(b)]
- Takahara, F., Kusunose, M.: Electron-positron pair production in a hot accretion plasma around a massive black hole. *Progr. Theor. Phys.* **73**(6), 1390–1400 (1985) [Sect. 7.3]
- Takizawa, M.: A two-temperature model of the intracluster medium. *Astrophys. J.* **509**(2), 579–584 (1998) [Sect. 8.3.4]
- Tamm, I.E.: *Basic Theory of Electricity*, 10th edn., p. 504. Nauka, Moscow (in Russian) (1989) [Sect. 19.3.1]
- Tandberg-Hanssen, E.: *The Nature of Solar Prominences*, p. 308. Kluwer Academic, Dordrecht (1995) [Sects. 19.3.4, 20.4]
- Thorne, K.: Gravitomagnetism, Jets in Quasars, and the Stanford Gyroscope Experiment. In: Fairbank, J.D., et al. (eds.) *Near Zero: New Frontiers of Physics*, pp. 573–586. W.H. Freeman and Co., New York (1988) [Sect. 13.3.2]
- Tidman, D.A., Krall, N.A.: *Shock Waves in Collisionless Plasma*, p. 175. Wiley-Interscience, New York (1971) [Sect. 16.4]
- Titov, V.S., Priest, E.R.: The collapse of an X-type neutral point to form a reconnecting current sheet. *Geophys. Astrophys. Fluid Dyn.* **72**, 249–276 (1993) [Sect. 14.2.2(b)]
- Todd, L.: Evolution of the trans-Alfvénic normal shock in a gas of finite electrical conductivity. *J. Fluid Mech.* **18**, 321–336 (1964) [Sect. 17.4.2]
- Toptyghin, I.N.: Acceleration of particles by shocks in a cosmic plasma. *Space Sci. Rev.* **26**(1), 157–213 (1980) [Sect. 18.3.2(a)]
- Treumann, R.A., Baumjohann, W.: *Advanced Space Plasma Physics*, p. 381. Imperial College Press, London (1997) [Sect. 7.1]
- Trubnikov, B.A.: Particle interactions in a fully ionized plasma. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 1, pp. 105–204. Consultants Bureau, New York (1965) [Sect. 8.4.1(b)]
- Tsiklauri, D., Haruki, T.: Magnetic reconnection during collisionless, stressed, X-point collapse using particle-in-cell simulation. *Phys. Plasma* **14**(11), 112905–112905-10 (2007) [Sect. 11.2]
- Tsuneta, S., Ichimoto, K., Katsukawa, Y., et al.: The Solar Optical Telescope for the Hinode mission: An overview. *Solar Phys.* **249**(2), 167–196 (2008) [Sect. 8.3.2]

- Tverskoy, B.A.: Contribution to the theory of Fermi statistical acceleration. *Soviet Phys. JETP*. **25**(2), 317–325 (1967) [Sect. 7.2]
- Tverskoy, B.A.: Theory of turbulent acceleration of charged particles in a plasma. *Soviet Phys. JETP*. **26**(4), 821–828 (1968) [Sect. 7.2]
- Tverskoy, B.A.: Main mechanisms in the formation of the Earth's radiation belts. *Rev. Geophys.* **7**(1), 219–231 (1969) [Sect. 6.4]
- UeNo, S.: Comparison between statistical features of X-ray fluctuations from the solar corona and accretion disks. In: Watanabe, T., Kosugi, T., Sterling, A.C. (eds.) *Observational Plasma Astrophysics: Five Years of Yohkoh and Beyond*, pp. 45–50. Kluwer Academic, Dordrecht (1998) [Sect. 13.2.4]
- Unti, T., Atkinson, G.: Two-dimensional Chapman-Ferraro problem with neutral sheet. 1. The boundary. *J. Geophys. Res. Space Phys.* **73**(23), 7319–7327 (1968) [Sect. 14.2.2(a)]
- van de Hulst, H.C.: Interstellar polarization and MHD waves. In: Burgers, J.M., van de Hulst, H.C. (eds.) *Problems of Cosmical Aerodynamics*, pp. 45–57, Central Air Documents Office, Dayton, Ohio (1951) [Sects. 15.2.3, 15.3.2]
- van den Oord, G.H.J.: The electrodynamics of beam/return current systems in the solar corona. *Astron. Astrophys.* **234**(2), 496–518 (1990) [Sects. 4.5.1, 4.5.2]
- Vink, J., Laming, J.M., Gu, M.F., et al.: The slow temperature equilibration behind the shock front of SN 1006. *Astrophys. J.* **587**(1), L31–L34 (2003) [Sect. 16.4]
- Vladimirov, V.S.: *Equations of Mathematical Physics*, p. 418. M. Dekker, New York (1971) [Sects. 1.1.5, 1.2.2, 13.1.1]
- Vlasov, A.A.: On the oscillation properties of an electron gas. *Zhur. Eksp. Teor. Fiz.* **8**(1), 29–33 (in Russian). English translation: 1968, The vibrational properties of an electron gas. *Sov. Phys. Uspekhi* **10**(4), 721–733, see also *Sov. Phys. Uspekhi* **19**(6), 545–546 (1938) [Sects. 3.1.2, 3.1.3, 10.2.2]
- Vlasov, A.A.: On the kinetic theory of an ensemble of particles with collective interactions. *Soviet J. Phys.* **9**(1), 25–28 (1945) [Sect. 3.1.2]
- Volkov, T.F.: Hydrodynamic description of a collisionless plasma. In: Leontovich, M.A. (ed.) *Reviews of Plasma Physics*, vol. 4, pp. 1–21. Consultant Bureau, New York (1966) [Sects. 11.5.1, 16.4]
- Walt, M.: *Introduction to Geomagnetically Trapped Radiation*, p. 188. Cambridge University Press, Cambridge (1994) [Sect. 6.4]
- Webb, G.M.: Similarity considerations and conservation laws for magnetostatic atmospheres. *Solar Phys.* **106**(2), 287–313 (1986) [Sect. 19.4.3]
- Webb, G.M., Zank, G.P., Ko, C.M., et al.: Multi-dimensional Green's functions and the statistics of diffusive shock acceleration. *Astrophys. J.* **453**(1), 178–189 (1995) [Sect. 18.2.2]
- Wentzel, D.G.: Fermi acceleration of charged particles. *Astrophys. J.* **137**(1), 135–146 (1963) [Sect. 18.3.2(b)]
- Wentzel, D.G.: Motion across magnetic discontinuities and Fermi acceleration of charged particles. *Astrophys. J.* **140**(3), 1013–1024 (1964) [Sects. 6.2.4, 18.3.2(b)]
- Wiita, P.J.: Accretion disks around black holes. In: Iyer, B.R., Bhawal, B. (eds.) *Black Holes, Gravitational Radiation and the Universe*, pp. 249–263. Kluwer Academic, Dordrecht (1999) [Sect. 8.3.5]
- Will, C.M.: Finally, results from Gravity Probe B. *Physics* **4**, 43 (2011) [Sect. 13.4]
- Woltjer, L.: A theorem on force-free magnetic fields. *Proc. Nat. Acad. Sci. USA* **44**(6), 489–491 (1958) [Sect. 19.6]
- Yvon, J.: *La Theorie des Fluids et l'Equation d'Etat*. Hermann et Cie, Paris (1935) [Sect. 2.4]
- Zank, G.P.: Weyl's theorem for MHD. *J. Plasma Phys.* **46**(1), 11–14 (1991) [Sect. 16.2.4(c)]
- Zel'dovich, Ya.B., Novikov, I.D.: *Relativistic Astrophysics*. Vol. 1, *Stars and Relativity*. University of Chicago Press, Chicago (1971) [Sects. 12.2, 19.3.4]
- Zel'dovich, Ya.B., Raizer, Yu.P.: *Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena*, vol. 1, p. 464, vol. 2, p. 452. Academic, New York (1966) [Sects. 8.3.4, 9.7.3, 16.1.3, 16.4, 16.5]

- Zel'dovich, Ya.B., Raizer, Yu.P.: *Physics of Shock Waves and High-Temperature Hydrodynamic Phenomena*. Dover, Mineola (2002) [Sects. 8.3.4, 9.7.3, 16.1.3, 16.4, 16.5]
- Zel'dovich, Ya.B., Ruzmaikin, A.A., Sokolov, D.D.: *Magnetic Fields in Astrophysics*. Gordon and Breach, New York (1983) [Sect. 13.1.2]
- Zenitani, S., Hoshino, M.: The generation of nonthermal particles in the relativistic magnetic reconnection of pair plasmas. *Astrophys. J.* **562**(1), L63–L66 (2001) [Sect. 7.3]
- Zenitani, S., Hoshino, M.: Particle acceleration and magnetic dissipation in relativistic current sheet of pair plasmas. *Astrophys. J.* **670**(1), 702–726 (2007) [Sect. 7.3]
- Zheleznyakov, V.V.: *Radiation in Astrophysical Plasmas*, p. 462. Kluwer Academic, Dordrecht (1996) [Sects. 7.1, 7.4, 10.4]
- Zhou, Y., Matthaeus, W.H.: Models of inertial range spectra of MHD turbulence. *J. Geophys. Res.* **95**(A9), 14881–14892 (1990) [Sect. 7.2]

Index

A

- abundance
 - elements, 106
- acceleration
 - Alfvén pumping, 138
 - by electric field, 226
 - by magnetic inhomogeneities, 121, 388
 - by MHD turbulence, 122
 - by shock waves, 122, 387
 - diffusive, 388
 - drift, 387, 393
 - by waves, 136, 316
 - electrons, 68, 72, 78, 81, 108
 - Fermi, 121, 138, 278
 - guiding center, 103
 - in solar flares, 64, 108, 122, 316
 - ions, 68, 109, 316
 - particle, 5, 13, 121, 165
 - protons, 123, 278, 397
 - stochastic, 121, 137, 316
- accretion
 - magnetic, 271
- accretion disk, 142, 168, 190, 270, 441
 - black hole, 272, 275, 432
 - white dwarf, 271
- acoustic wave, 212
- active galaxy, 142, 168, 234, 272, 275, 278, 432
- active region, 413
- adiabatic cooling, 121
- adiabatic invariant, 13, 115
 - first *or* transversal, 117, 394, 395
 - second *or* longitudinal, 120
 - third *or* flux invariant, 124
- adiabatic process, 333
- adiabatic theory
 - accuracy, 124
- advection, 168, 190, 273
 - advection-dominated accretion flow, 168, 190
 - advective derivative, 194
 - Alfvén, xvii, 279, 313, 433
 - Alfvén discontinuity, 353, 433
 - non-evolutionarity, 377
 - propagation order, 373
 - Alfvén pumping, 138
 - Alfvén soliton, 143
 - Alfvén wave, 142, 312, 322, 353, 365, 370, 433
 - dissipation, 316, 318, 374
 - kinetic, 146
 - algebra of Lie, 196
 - angular diffusion, 63, 69, 79, 120
 - Ap star, 409
 - approximation
 - binary collisions, 37, 147
 - CGL, 142, 232
 - cold ions, 149, 155, 158, 212
 - collisionless, 150
 - diffusion, 141
 - hydrodynamic, 74, 207, 219
 - isotropic conductivity, 232
 - kinetic, 74, 219
 - large mag. Reynolds number, 263, 449
 - magnetostatic, 268, 414, 424
 - multi-fluid, 211
 - non-relativistic, 91, 117, 246
 - single-fluid two-temperature, 160, 166, 360
 - small displacement, 269, 297
 - small mag. Reynolds number, 263, 436, 441
 - stationary, 269, 338, 429
 - strong magnetic field, 228, 267, 315
 - strong-field-cold-plasma, 268, 285, 294, 295
 - two-dimensional, 287
 - two-fluid, 211, 224

- approximation (*cont.*)
 - two-temperature, 278
 - weak Coulomb interaction, 154
 - weak magnetic field, 228, 265, 271
- atmosphere
 - solar, 43, 150, 153, 155, 268, 318, 373, 411, 436
- Aurora, 133
- averaged force, 31, 45, 181

- B**
- BBGKY hierarchy, 35
- beam
 - blunt, 74
 - electrons, 72
- betatron acceleration, 138
- binary interaction, 29, 147
- binary stars, 168, 270
- black hole, 38, 168, 432
 - rotating, 142, 190, 234, 246, 275
 - supermassive, 180, 272, 277
- Boltzmann integral, 43, 55
- Boltzmann law, 154
- Bondi accretion, 273
- bounce motion, 121
- boundary conditions, 12, 355
 - ideal MHD, 337
 - isolated groups, 335, 366
 - linearized, 365
- boundary layer, 141, 439
- bow shock, 383
- bremsstrahlung, 68, 159, 161, 167
 - collisional, 153, 193
 - magnetic, 193
- Brown motion, 42
- bump-on-tail instability, 133

- C**
- catenary, 88
- centrifugal force, 104
- Chandra, 273
- characteristic equations, 15
- characteristic time
 - diffusive shock acceleration, 391
- charge neutrality, 157, 160, 233
- charge-separated plasma, 247
- Cherenkov condition, 143, 332
- Cherenkov effect, 134
- chromospheric evaporation, 122
- Clebsch variables, 302
- cluster of galaxies, 166
- collapse
 - gravitational, 46, 408
 - star, 301, 408
- collapsing magnetic trap, 109, 123, 126, 150, 398
- collective phenomena, 44, 49, 157
- collision
 - characteristic time, 151, 188, 197, 243
 - close, 44, 148, 177, 235
 - Coulomb, 13, 41, 59, 147
 - cross-section, 149
 - distant, 42, 148, 177, 235
 - elastic, 224
 - frequency, 152
 - head-on, 121, 125
 - mean free path, 149
 - overtaken, 121
 - trailing, 121, 126
- collisional integral, 7, 27, 31, 51, 181
 - Boltzmann, 55
 - Landau, 40, 44, 56, 60
 - linearized, 75
- collisional time
 - between electrons, 158, 178
 - between electrons and ions, 159, 243
 - between ions, 158, 278
- collisionless plasma, 93, 103, 129, 355
- collisionless shock, 356, 387
- complex potential, 293
- conditional probability, 29
- conditions
 - boundary, 12, 337, 355
 - isolated groups, 366
 - linearized, 365
 - electrodynamic continuity, 339
 - evolutionarity, 365, 372
 - initial, 12, 286, 295
- conductive current, 234, 245
- conductivity
 - anisotropic, 235
 - electric, 223
 - Hall, 227
 - isotropic, 228, 232, 244
 - parallel, 235
 - perpendicular, 227
 - thermal, 167, 248
- conform mapping, 293, 303
- conservation law
 - energy, 16, 240, 335, 338
 - magnetic flux, 251, 255
 - mass, 184, 335, 338
 - momentum, 239, 249, 335, 338
 - particles, 3
- contact discontinuity, 341
 - non-evolutionarity, 371, 377

- continuity equation, 286
 - electric charge, 11
 - for particles of kind k , 184
 - for plasma, 238
 - in phase space, 2
 - Lagrangian form, 294
 - convective current, 234, 245
 - cooling
 - adiabatic, 121
 - by heat flux, 163
 - radiative, 162
 - coordinates
 - doubly Lagrangian, 299
 - generalized, 15, 115
 - Lagrangian, 298
 - polarized, 134
 - cork ratio, 119
 - coronal heating, 318
 - coronal mass ejection, 165, 246, 350, 420, 423
 - coronal transient, 246, 423
 - correlation function
 - binary, 29, 32
 - triple, 35, 46
 - corrugational instability, 386
 - cosmic rays, 91, 123, 125, 278
 - galactic, 390, 392
 - Coulomb collision, 13, 22, 59, 147
 - Coulomb force, 32
 - Coulomb logarithm, 40, 152, 155, 177
 - Crab nebula, 112, 143
 - current
 - conductive, 234, 244
 - convective, 234, 245
 - direct, 43, 74, 79
 - displacement, 244
 - Hall, 236
 - reverse, 43, 73, 176
 - current layer
 - collisionless, 40
 - reconnecting, 199, 204, 257, 367, 414, 420, 423
 - super-hot turbulent-current, 64, 77, 165, 172, 199, 204
 - curvature
 - magnetic field line, 104, 281
 - cyclotron *or* gyrofrequency, 107
 - Cyg X-1, 274
- D**
- damping
 - Alfvén wave, 316, 318, 319
 - collisional, 157
 - Landau, 41, 132
 - length, 319
 - time, 319
 - dark matter, 38
 - de Broglie wavelength, 109
 - de Hoffmann-Teller frame, 346
 - Debye-Hückel radius, 153, 175, 206, 235, 244
 - electron, 48, 215
 - density
 - charge, 233, 238
 - in MHD, 245
 - current in MHD, 245
 - energy flux, 17, 248, 255
 - friction, 248
 - heat, 248
 - magnetic field energy, 267, 282, 342
 - mass, 182
 - momentum flux, 247
 - number, 182
 - particle flux in phase space, 3
 - plasma, 298
 - probability distribution, 25
 - spectral, 138
 - description
 - exact, 9
 - kinetic *or* microscopic, 181
 - kinetic *or* microscopic, 13
 - macroscopic, 13, 181
 - statistical, 9
 - diamagnetic effect, 100
 - differential rotation, 273, 409, 433
 - diffusion
 - angular, 63, 79, 120
 - energy, 62, 78
 - magnetic, 73
 - pitch angle, 62, 136
 - quasi-linear, 220
 - turbulent, 266
 - diffusivity
 - magnetic, 251
 - dimensionless parameters, 95, 265
 - dipole moment, 296
 - direct current, 43, 74, 79
 - discontinuity
 - Alfvén *or* rotational, 353, 354
 - evolutionarity, 371
 - boundary conditions, 355
 - hydrodynamics, 334
 - ideal MHD, 337
 - linearized, 365
 - classification, 337
 - contact, 341, 377
 - evolutionary, 363
 - non-evolutionary, 363
 - shock wave, 334, 336, 343

- discontinuity (*cont.*)
 - small perturbations, 363
 - switch-off shock
 - evolutionarity, 371
 - switch-on shock
 - evolutionarity, 371
 - tangential, 315, 336, 341, 377, 386, 430
 - transitional, 355
 - weak, 377
 - discontinuous flow, 307
 - discontinuous solutions
 - continuous transitions, 354, 373
 - dispersion equation, 309, 385
 - displacement current, 244
 - dissipation
 - Joule, 165, 230, 253
 - dissipative wave, 371, 375
 - distribution function, 1
 - averaged, 20, 29
 - bump-on-tail, 120, 133
 - exact, 8, 19, 27
 - Maxwellian, 197
 - non-equilibrium, 120
 - Dreicer field, 77, 85, 174, 179, 426
 - drift
 - centrifugal, 103, 105
 - curvature-dependent, 105, 125
 - electric, 93, 102
 - fast, 92
 - gradient, 102, 125
 - gravitational, 94, 102
 - inertial, 103
 - non-magnetic force, 92
 - slow, 92
 - drift shell, 124
 - dust, 13
 - dynamic friction, 43, 169, 211
 - gravitational, 179
 - dynamical trajectory, 23
 - dynamo
 - gravitomagnetic, 278
 - magnetic, 266, 273, 410
 - photospheric, 441
 - turbulent, 266
- E**
- Earth
 - magnetic field, 91, 123, 125, 136
 - electric conductivity
 - anisotropic, 227
 - isotropic, 223
 - electric drift, 93, 102
 - electric field, 10
 - cross-shock, 346
 - Dreicer, 77, 85, 174, 426
 - electrostatic, 73
 - Hall, 230
 - in MHD, 245
 - in moving plasma, 244
 - inductive, 73
 - reverse current, 74, 80, 85, 176
 - electric neutrality, 46, 225, 233
 - electric resistivity, 252
 - electromagnetic force, 239
 - electron-positron cascade, 247
 - electrons
 - back-flying, 76, 80
 - runaway, 76, 81
 - trapped, 76
 - energy
 - internal, 344
 - energy conservation law, 189, 240, 335
 - energy diffusion, 62, 78
 - energy flux density, 17, 248, 255, 282
 - energy losses
 - radiative, 320
 - regular, 62, 78
 - enthalpy
 - specific, 191, 248, 344, 357
 - entropy, 249
 - entropy wave, 212, 311, 323, 325, 331, 366, 370, 377
 - equation
 - biharmonic, 445
 - continuity, 2, 11, 160, 184, 212, 238, 286
 - correlation function, 32, 34
 - diffusion, 42
 - diffusion-convection, 388, 399
 - dispersion, 132, 309, 385
 - Fokker-Planck, 42, 51
 - freezing-in, 255, 286
 - Grad-Shafranov, 424
 - guiding center motion, 99, 101
 - heat transfer, 249
 - kinetic, 27, 32, 181
 - Langevin, 52
 - linear oscillator, 213
 - Liouville, 4, 19
 - motion, 87, 212, 245, 249, 292
 - Poisson, 56, 154
 - state, 249
 - ideal gas, 191, 197
 - linearized, 308
 - Vlasov, 39, 129, 142, 219, 356
 - equations
 - autonomous, 379

- characteristic, 15, 52
- Einstein, 250, 282
- Hamilton, 15
- ideal MHD, 254, 264, 338
 - linearized, 308
- magnetic field line, 290
- Maxwell, 10, 39, 130, 245
- MHD, 242
- Newton, 11
- particle motion, 11
- transfer, 138, 182, 194, 254
- equilibrium
 - magnetostatic, 414, 420
 - MHD, 403, 423, 430
 - non-existence, 408, 423
 - static, 407
 - thermodynamic, 45, 46, 154, 190
- Euler potentials, 302
- evolutionarity
 - Alfvén discontinuity, 371
 - conditions, 365, 372
 - consequences, 372
 - contact discontinuity, 371
 - continuous transitions, 373
 - definitions, 363
 - entropy wave, 311
 - fast shock wave, 370
 - slow shock wave, 370
 - switch-off shock, 371
 - switch-on shock, 371
 - tangential discontinuity, 371
- exact distribution function, 8, 19, 27
- expulsion force, 423, 441

- F**
- Fermi acceleration, 138, 278
 - first-order, 126
 - regular, 122
 - second-order, 126
 - stochastic, 122
- field
 - constant homogeneous, 88
 - force-free, 410
 - slowly changing weakly inhomogeneous, 96, 116
 - weakly inhomogeneous, 95
- field reversal, 350, 383
- fire-hose instability, 142
- flare
 - accretion disk, 274
 - giant, 410
 - solar, 64, 68, 84, 108, 112, 119, 122, 133, 150, 153, 159, 176, 217, 220, 274, 373, 420
- fluid particle, 292, 297
- Fokker-Planck equation, 42, 51
- force
 - Archimedean, 421, 423, 441
 - averaged, 31, 45
 - binary, 28
 - centrifugal, 104
 - collisional drag, 169, 223
 - Coulomb, 32
 - diamagnetic, 100
 - electric
 - in MHD, 246
 - electromagnetic, 239
 - expulsion, 441
 - friction, 6, 80, 193, 212, 223, 241
 - gravitational, 6, 14, 31, 50, 179, 206, 265, 282
 - gravitomagnetic, 275, 282
 - inertia, 268
 - Lorentz, 6, 31, 186, 420
 - magnetic, 89, 246
 - magnetic σ -dependent, 422, 441
 - magnetic buoyancy, 422
 - mean, 26, 51, 186
 - mean collisional, 186
 - non-magnetic, 88, 91, 267
 - statistically averaged, 181, 186
 - stochastic, 42
 - viscous, 164, 192, 245, 447
- force-free field, 267, 286, 410, 423
 - helicity, 426
 - linear, 426
- Fourier law, 199
- fractionation
 - elements, 106, 212
- frame-dragging effect, 276
- freezing-in equation, 286
- frequency
 - collision, 152, 157, 188
 - Doppler-shifted, 135
 - electron plasma, 133, 157, 213
 - gyrofrequency *or* cyclotron, 89, 107
 - ion plasma, 217
- friction force, 6, 80, 179, 211, 223
 - work, 193, 241
- function
 - correlation, 29
 - delta, 8
 - distribution, 1
 - error, 62

function (*cont.*)

- heat, 191
- Maxwellian, 45, 190
- radiative losses, 241
- stream, 300
- theta, 65

G

- galaxy
 - active, 272, 275
 - elliptical, 273
- gamma-ray burst, 266
- gas
 - ideal, 191, 197
- general relativity, 250
- generalized momentum, 115
- geomagnetic storm, 125
- geomagnetic tail, 420
- giant flare, 410
- Giovanelli, 174
- Grad moment method, 188, 199, 242
- Grad-Shafranov equation, 424
- gradient drift, 102
- gravitational drag, 179
- gravitational drift, 94, 102
- gravitational energy, 406
- gravitational force, 6, 14, 31, 50, 179, 265, 361
 - averaged, 206
- gravitational pressure, 408
- gravitational system, 14, 46, 50, 151, 206
 - thermodynamic equilibrium, 46
- Gravity Probe B, 276
- group velocity, 213, 309
- guiding center, 96, 123, 135
- guiding center acceleration, 103
- guiding center motion
 - flux invariant, 124
- guiding center spiral, 101
- gyromotion, 96
- gyroradius *or* Larmor radius, 90
- gyroresonance, 135

H

- Hall cascade, 231
- Hall current, 227, 236
- Hamilton equations, 15
- Hamiltonian
 - usual, 15
- hard X-ray bremsstrahlung, 68, 153
- Hartmann number, 438
- heat conduction
 - anisotropic, 320

- nonlocal, 206
- heat flux density, 192
- heat function, 191
- heating
 - by electron beam, 159
 - chromospheric, 119
 - coronal, 318
 - Joule, 249
 - viscous, 249
- helioseismology, 435
- Hinode, viii, 161, 249
- horizon
 - black hole, 168, 272
- hydrodynamic velocity, 237
- hydrodynamics
 - ideal, 336

I

- ideal gas, 191, 197
- ideal hydrodynamics, 336
- ideal MHD, 254
- impact parameter, 148
- inertial drift, 103
- initial conditions, 12, 286, 295
- injection energy, 137
- injection spectrum, 65
- instability
 - bump-on-tail, 133, 136
 - corrugational, 386
 - fire-hose, 142, 146
 - gyroresonance-type, 136
 - Jean, 207
 - kinetic, 120, 356
 - magnetorotational, 271
 - MHD, 423
 - overheating, 320
 - shear, 270
 - thermal, 329, 423
 - two-stream, 134
- integral
 - Boltzmann, 43, 55
 - Cauchy, 41
 - collisional, 7, 27, 31, 51, 181
 - Landau, 40, 51, 56
 - motion, 14, 124, 130, 292, 297
- interaction
 - binary, 29
 - Coulomb, 43, 147
 - weak, 38, 154
 - electromagnetic, 12
 - particles, 4
 - wave-particle, 13, 15, 129
- interaction parameter, 30, 38

interplanetary magnetic field, 383
 interplanetary space, 121
 interstellar cloud, 121, 207
 interstellar medium, 273
 intra-cluster medium, 167
 invariant
 adiabatic, 13, 115
 motion, 14
 ion-acoustic wave, 133, 216
 ionosphere, 136, 218, 228
 Irnshaw theorem, 413
 isorotation, 434

J

Jean's instability, 207
 Jeans, 151
 Jeans theorem, 15, 131
 jet
 astrophysical, 431
 bi-directional, 166, 258, 432
 disk corona, 278
 non-relativistic, 430
 relativistic, 142, 240, 275, 432
 Joule dissipation, 165
 Joule heating, 167, 249

K

kinematic problems, 266
 kinematic viscosity, 253, 374
 kinetic energy, 189, 405
 kinetic equation, 27, 32, 181
 kinetic instability, 356
 Kolmogorov
 scaling law, 140
 spectrum, 139

L

Lagrange identity, 427
 Lagrangian coordinates, 298
 Lagrangian lines, 291
 Lagrangian variables, 292, 297
 Landau
 collisional integral, 40, 56, 60
 gravitational analog, 51
 Landau damping, 41, 132, 219
 nonlinear, 134
 Landau resonance, 42, 132
 Langevin equation, 52
 Langmuir wave, 49, 133, 157, 212, 213, 234
 finite amplitude, 145
 Larmor radius, 97, 107, 393

law

T to the $3/2$ power, 151
 Boltzmann, 154
 conservation, 189, 239, 255, 338
 Ohm's, 13, 223

layer

boundary, 141, 439
 reconnecting, 452

Lichnerowicz, 250

Lie algebra, 196

Liouville equation, 4, 19

Liouville operator, 4, 130, 143

Liouville theorem, 4

liquid contour, 252

loop

flaring, 119, 122

Lorentz factor, 87, 137, 144

Lorentz force, 6, 31, 95, 186, 432

loss cone

anomalous diffusion, 120
 magnetic trap, 119

Lundquist force free field, 412

M

macroparticle method, 52
 magnetar, 110, 266, 302, 409
 magnetic collapse, 295, 306
 magnetic diffusivity, 73, 247, 251, 374, 441
 magnetic dynamo, 266, 273
 magnetic energy, 406, 426
 magnetic field
 force free, 267, 286, 410
 helical, 411
 interplanetary, 398
 limiting line, 288
 longitudinal, 288
 plane dipole, 296
 potential *or* current free, 267
 shear, 411
 superstrong, 109, 302
 toroidal, 434
 transversal, 436
 ultrastrong, 110, 409
 zeroth point *or* line, 288
 magnetic field line
 equations, 290
 meaning, 255
 separator, 288
 magnetic flux, 290
 magnetic flux conservation, 251, 255
 magnetic flux tube
 coronal, 119
 specific volume, 417

- magnetic force, 89, 246
 - magnetic helicity
 - global, 256, 426
 - magnetic mirror, 100, 118
 - magnetic moment, 99, 105, 113, 117, 300
 - magnetic pressure, 268, 341, 348, 411, 430
 - perturbation, 314
 - magnetic reconnection, 141, 165, 190, 250, 257, 274, 288, 294
 - collisionless, 13, 143
 - magnetic Reynolds number, 253
 - magnetic separator, 448
 - magnetic sound, 315
 - magnetic stresses, 270
 - magnetic surface, 124, 414, 424
 - magnetic tension, 268, 280, 312, 411, 433
 - magnetic trap, 109, 119, 150, 398
 - collapsing, 123, 126
 - magnetoacoustic wave, 323, 366, 370
 - damping, 328
 - fast, 325, 328, 331, 344, 349
 - slow, 324, 328, 330, 349
 - magnetohydrodynamics, 13, 232, 242, 245
 - magnetosphere
 - black hole, 142, 234
 - Earth, 91, 228, 257, 268
 - Jovian, 112
 - magnetic dipole, 303
 - Ohm's law, 228
 - pulsar, 142, 234, 246, 250
 - substorm, 257
 - white dwarf, 272
 - Maxwell equations, 10, 39, 130, 245
 - Maxwellian function, 45, 190, 197
 - Maxwellian stress tensor, 239, 404
 - mean collisional force, 186, 208
 - mean field, 50, 266
 - mean force, 26, 51, 186
 - mean free path, 149, 175, 197, 243
 - mean kinetic energy, 189
 - mean momentum, 183
 - mean thermal velocity, 151
 - electron, 107, 215
 - proton, 107
 - mean velocity, 182
 - MHD
 - ideal, 254, 264, 285
 - non-relativistic, 247
 - relativistic, 250, 275
 - two-fluid, 382
 - MHD assumptions, 244
 - MHD pump, 440
 - MHD turbulence, 122, 139, 141
 - microquasar, 432
 - minimum current corona, 427
 - mixing mechanism, 22
 - model
 - Boltzmann gas, 56
 - toy, 51
 - moment
 - inertia, 405
 - magnetic, 99, 113, 117, 300
 - of distribution function, 182
 - viscous force, 447
 - momentum
 - angular, 270, 435
 - conservation, 335
 - electromagnetic field, 247
 - generalized, 15, 115
 - longitudinal, 142
 - mean, 183
 - plasma motion, 247
 - transversal, 118, 142
 - momentum flux density tensor, 183, 224, 239, 247, 255
 - motion
 - bounce, 121
 - guiding center, 99, 124
 - random, 207, 208
 - spiral, 92
- N**
- neutron star, 109, 168, 231, 302, 409
 - Newton equations, 11
- O**
- Ohm's law
 - generalized, 13, 227
 - in MHD, 232, 243
 - usual, 13, 75, 223
 - operator
 - Liouville, 4, 130, 143
 - oscillatory disintegration, 383
- P**
- parameter
 - m/e , 96
 - interaction, 30, 38
 - plasma, 49
 - Parker buoyancy force, 422
 - particle
 - accelerated, 43, 121
 - field, 40, 149
 - fluid, 292
 - precipitating, 119

- test, 42, 149
- trapped, 119, 125
- particle flux density, 3
- particle interaction, 4
- particle simulation, 13
- permeability, 238
- permittivity, 238
- perpendicular shock, 343
 - structure, 377
- Petschek, viii
- phase space, 1, 19
- phase trajectory, 4
- phase velocity, 309
- phase velocity diagram, 314, 354
- pinch effect, 288
- pitch-angle, 90, 119, 136
- plasma, 46
 - anisotropic, 142, 192
 - charge-separated, 150, 234
 - collisional, 41, 43, 129
 - collisionless, 41, 93, 103, 129, 142, 233
 - dusty, 13
 - electron-positron, 142
 - fully-ionized, 41, 59, 211
 - hot, 164
 - quark-gluon, 250
 - quasi-neutrality, 46, 157
 - rare-collisional, 151
 - self-gravitational, 13
 - strongly-coupled, 38
 - super-hot, 108, 164, 193
 - three-component, 224
 - two-temperature, 159
 - weakly-coupled, 38
 - weakly-ionized, 13, 440
- plasma frequency, 157
 - ion, 217
- plasma parameter, 49
- plasma wave, 74, 157
- Poisson brackets, 15
- Poisson equation, 56, 154
- polarized coordinates, 134
- postulates of statistics, 21
- potential
 - complex, 293
 - conjugate harmonic, 295
 - Coulomb, 147
 - Euler, 302
 - gravitational, 404
 - magnetic field, 267
 - shielded, 48
 - vector, 290
- Poynting vector, 17, 241, 248, 282, 450

- pressure
 - partial, 240
 - total, 240
- pressure tensor, 183, 189, 192, 224, 239
- probability density, 25
- prominence, 13, 419, 452
- protostar, 441
- pulsar
 - magnetosphere, 142, 234, 250
- purely dissipative wave, 375

Q

- quantum effects, 109
- quantum indeterminacy, 16
- quark-gluon plasma, 250
- quasar, 432

R

- radiation
 - synchrotron, 112, 142
- radiation belts, 125
- radiation reaction, 6, 110
- radio source
 - extragalactic, 431
- Rankine–Hugoniot relation, 348, 357, 381, 385
- reconnecting current layer, 101, 165, 204, 268
- reconnection
 - collisionless, 13, 40, 143, 226
 - large-scale, 258
 - magnetic, 141, 190, 274, 452
 - patchy, 257
 - strong magnetic field, 260
 - weak magnetic field, 260
- reduced mass, 148
- refraction index, 144, 218
- resistivity
 - anomalous, 226
 - classical, 226
- resonance
 - Landau, 42, 132
- reverse current, 43, 59, 73, 176
- Reynolds number
 - hydrodynamic, 140, 253
 - magnetic, 253
- RHESSI, viii, 82, 83, 85, 161, 249
- rigidity, 90
 - threshold, 125
- ring current, 125
- rotation
 - differential, 266, 433

runaway

- collisional, 76
- electric, 174, 211
- thermal, 176, 206

Rutherford formula, 56, 149

S

scaling law

Kolmogorov, 140

SDO, vii, 242, 249

separation

charge, 150

MHD, 422

separator, 288

Shafranov theorem, 413

shear, 270

shock wave

bow, 383

collisionless, 40, 346, 355, 387, 397

discontinuity surface, 336

fast

evolutionarity, 370, 377

high Mach number, 167, 356

intermediate *or* trans-Alfvénic, 350, 371, 377, 382

interplanetary, 397

longitudinal, 354

oblique, 347

fast, 348, 354

slow, 348, 354

ordinary, 388

perpendicular, 343, 354, 393

propagation order, 372

quasi-parallel, 392

quasi-perpendicular, 392, 395

Rankine–Hugoniot relation, 348, 382

slow

evolutionarity, 370, 377

switch-off, 352

evolutionary, 376

switch-on, 351

non-evolutionary, 377

soft gamma-ray repeater, 302

SOHO, 166, 249, 435

solar atmosphere, 43, 68, 150, 155, 373

solar corona, 84, 107, 112, 137, 146, 177, 212, 220, 228, 235, 249, 261, 318, 331, 452

solar flare, 64, 68, 84, 108, 119, 122, 133, 159, 220, 248, 257, 420

impulsive phase, 72, 85

two-ribbon, 423

solar photosphere, 13, 118

solar prominence, 242

solar wind, 121, 141, 199, 211, 257, 383, 391

sound velocity, 308, 333

space

interplanetary, 121, 133, 146

phase, 1, 19

wave number, 141

special relativity, 126, 250

specific enthalpy, 191, 248, 344, 357

specific magnetic volume, 256, 417

specific volume, 357

spectrum

injection, 65

Kolmogorov, 139, 146

Störmer solutions, 125

star

AM Herculis, 272

binary, 168, 270

cataclysmic variable, 271

class A, 409

cold giant, 409

collapse, 301, 408

DQ Herculis, 272

formation, 441

in galaxy, 50, 179, 206

intermediate polars, 272

magnetar, 110, 302, 409

magnetic, 409

neutron, 168, 302, 409

nova, 300

polars, 270, 272

rotation, 434

Sun, 246, 409

supernova, 112, 123, 167, 250, 266, 300, 302, 356, 390

white dwarf, 16, 38, 54, 109, 271, 302

statistical averaging, 25

stellar system, 43

STEREO, vii, 249

stochastic acceleration, 136

stochastic force, 42

stochastic simulation, 53

stream function, 300

substantial derivative, 194, 269

Sun

active region, 413

chromosphere, 68, 75, 84, 119, 162, 249, 441

corona, 53, 72, 107, 112, 122, 137, 220, 228, 235, 331, 452

photosphere, 118, 441

prominence, 248, 419

rotation, 434

- sunspot, 118, 235
 - transition layer, 178
 - superstrong magnetic field, 109
 - supernova remnant, 356
 - switch-off shock wave, 352, 371, 375, 384
 - switch-on shock wave, 351, 371, 375, 384
 - synchrotron radiation, 112
 - Syrovatskii, viii, 337, 366, 411, 430
- T**
- tachocline, 435
 - tangential discontinuity, 355
 - hydrodynamics, 336, 386
 - ideal MHD, 341, 430
 - non-evolutionarity, 371, 377, 386
 - stability, 342
 - weak, 315
 - temperature, 151
 - effective *or* kinetic, 191
 - equilibrium, 190
 - tensor
 - conductivity, 230
 - Maxwellian stress, 404
 - momentum flux density, 183, 239, 247, 255
 - pressure, 183, 189, 192
 - unit antisymmetric, 112
 - viscous stress, 192, 220, 248, 447
 - test particle, 42
 - theorem
 - Irish, 413
 - Jeans, 15, 131
 - Liouville, 4
 - MHD “frozen field”, 255
 - non-existence, 407
 - Ostrogradskii-Gauss, 3, 405
 - Shafranov, 413
 - virial, 14, 403
 - Zemplen, 348
 - thermal conductivity, 248
 - thermal energy, 405
 - thermal instability, 242, 329
 - thermal runaway, 206
 - thermodynamic equilibrium
 - overall, 46
 - theta-pinch, 289
 - thick target, 65
 - threshold rigidity, 125
 - toy model, 51
 - TRACE, viii
 - transfer coefficients, 197, 355
 - transfer equations, 138, 182, 194
 - transition layer, 164
 - trapped particle, 109, 119, 125, 398
 - triple correlation function, 35, 46
 - turbulence
 - Hall cascade, 231
 - helical, 266
 - hydrodynamic, 138, 139
 - MHD, 122, 141, 266, 271
 - strong, 399
 - MHD wave, 316
 - plasma, 40, 164
 - weak, 134
 - two-dimensional problem
 - axial-symmetric, 299
 - first type, 288
 - second type, 289
 - two-temperature plasma, 159
- V**
- vector potential, 290
 - velocity
 - drift, 92
 - group, 213, 309
 - hydrodynamic, 237
 - mean thermal, 151
 - most probable, 170
 - phase, 132, 309
 - sound *or* acoustic, 308, 333
 - virial theorem, 14
 - scalar, 403, 407
 - tensor, 404, 409
 - viscosity
 - dynamic, 164, 220, 442
 - kinematic, 140, 247, 261
 - viscosity coefficient, 246
 - viscous force, 164, 192, 245
 - viscous heating, 164
 - viscous stress tensor, 192, 220, 240, 248
 - Vlasov equation, 39, 75, 129, 142, 219, 356
 - gravitational analog, 50, 361
 - volume charge, 233, 238, 246
 - vortex flow, 422
 - Voyager, 217
- W**
- wave
 - acoustic, 212
 - Alfvén, 312, 322, 353, 354, 370, 433
 - kinetic, 146
 - nonlinear, 143
 - relativistic, 142
 - de Broglie, 109
 - dissipative, 371, 375
 - electromagnetic, 217

wave (*cont.*)

- entropy, 212, 311, 323, 325, 331, 366, 370
 - entropy-vortex, 332
 - ion-acoustic, 133, 216
 - Langmuir, 49, 133, 145, 157, 213
 - large-amplitude, 307, 354
 - low-frequency, 216
 - magnetoacoustic, 319, 323, 370
 - fast, 313, 325, 328, 331, 344, 349, 354
 - slow, 313, 324, 328, 330, 349
 - plane, 308, 365
 - shock, 40, 307, 334, 336
 - small-amplitude, 307, 354
 - sound *or* acoustic, 354
 - whistler, 136, 146
- wave cascading, 138
- wave spectral density, 138
- weak discontinuity, 377
- white dwarf, 16, 38, 54, 109, 271

X

- X-ray binary system, 168
- X-ray cluster, 166
- X-ray emission
 - bremsstrahlung, 72, 108, 161, 167
 - hard, 68, 122, 153, 161, 167
 - polarization, 72, 81
 - soft, 165
- X-type zeroth point, 256, 288, 295
- XMM-Newton, 356

Y

- Yohkoh, 161

Z

- Zeeman effect, 272
- Zemlen theorem, 348